

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number  
WO 03/000656 A2

(51) International Patent Classification<sup>7</sup>:

C07D

(21) International Application Number:

PCT/US02/19665

(22) International Filing Date:

19 June 2002 (19.06.2002)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09/888,361 21 June 2001 (21.06.2001) US

(71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MURRAY, Susan, F.** [US/US]; 12348 Ann-O-Reno Lane, Poway, CA 92064 (US). **WYATT, Jacqueline, R.** [US/US]; 1065 Hymettus Avenue, Encinitas, CA 92024 (US).

(74) Agents: **LICATA, Jane, Massey** et al.; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/000656 A2

(54) Title: ANTISENSE MODULATION OF TRANSFORMING GROWTH FACTOR BETA RECEPTOR II EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of Transforming growth factor beta receptor II. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding Transforming growth factor beta receptor II. Methods of using these compounds for modulation of Transforming growth factor beta receptor II expression and for treatment of diseases associated with expression of Transforming growth factor beta receptor II are provided.

**ANTISENSE MODULATION OF TRANSFORMING GROWTH FACTOR BETA RECEPTOR  
II EXPRESSION**

**FIELD OF THE INVENTION**

The present invention provides compositions and methods for modulating the expression of Transforming growth factor beta receptor II. In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding Transforming growth factor beta receptor II. Such compounds have been shown to modulate the expression of Transforming growth factor beta receptor II.

**BACKGROUND OF THE INVENTION**

The transforming growth factor beta (TGF- $\beta$ ) superfamily of cytokines regulates a diverse array of physiologic functions including cell proliferation and growth, cell migration, differentiation, development, production of extracellular matrix, and the immune response. Each subgroup of this superfamily initiates a unique intracellular signaling cascade stimulated by ligand-induced formation and activation of specific serine/threonine kinase receptor complexes. The TGF- $\beta$  subfamily signal transduction occurs through a requisite interaction between transforming growth factor beta type I and II receptors. Mutant function or overactivity of TGF- $\beta$  signaling components is implicated in cancers of the colon, esophagus, pancreas, lung, and breast, as well as in hyperproliferative disorders of the kidney, atherosclerosis, and rheumatoid

-2-

arthritis (Imai et al., *J. Nephrol.*, 1998, 11, 16-19; Markowitz, *J. Clin. Invest.*, 1997, 100, 2143-2145; Pasche, *J. Cell Physiol.*, 2001, 186, 153-168; Piek et al., *Faseb J.*, 1999, 13, 2105-2124).

Transforming growth factor beta receptor II (also known as TGF-beta receptor II, TGFBR2, Tgfbr2, TGF- $\beta$  RII, T $\beta$ R-II, TbetaRII) has high affinity for the TGF- $\beta$ 1 ligand but low affinity for the TGF- $\beta$ 2 ligand. Thus, in some cell types, a third receptor (TGF- $\beta$  receptor type III), is required to facilitate TGF- $\beta$ 2 binding and presentation of the ligand to transforming growth factor beta receptor II (Rotzer et al., *Embo J.*, 2001, 20, 480-490). Once the growth factor/type II receptor complex is formed, the kinase domain of transforming growth factor beta receptor II activates the type I receptor by transphosphorylation, and the activated type I receptor then associates with cytoplasmic effectors, the Smad proteins, to propagate the kinase signal (Piek et al., *Faseb J.*, 1999, 13, 2105-2124).

Transforming growth factor beta receptor II was cloned from LLC-PK<sub>1</sub> porcine renal epithelial cells (Lin et al., *Cell*, 1992, 68, 775-785). Disclosed and claimed in US Patent 6,008,011 are nucleic acid sequences encoding full length and soluble polypeptides encoding transforming growth factor beta receptor II, complementary fragments of such nucleic acids useful as probes, as well as the corresponding vectors, host cells, and their use in the production of recombinant receptors (Lin et al., 1999). The human transforming growth factor beta receptor II gene was mapped to the 3p22 locus, a region implicated in lung and breast carcinomas (Mathew et al., *Genomics*, 1994, 20, 114-115), and the mouse Tgfbr2 gene was mapped to distal mouse chromosome 9 within a region of synteny with the human chromosome at 3p22-p21 (Bonyadi et al., *Genomics*, 1996, 33, 328-329).

Alternate transcripts of transforming growth factor beta

-3-

receptor II genes in mouse and human have been identified. One human isoform, T $\beta$ RII-B, has acquired a high affinity binding site for TGF- $\beta$ 2 in its extracellular domain and can bind the ligand without assistance from the type III receptor. The sites of predominant expression of the T $\beta$ RII-B type II receptor isoform in osteoblasts and mesenchymal precursor cells also correlate with expression of the TGF- $\beta$  2 ligand in chondrocytes and osteocytes (Rotzer et al., *Embo J.*, 2001, 20, 480-490).

Transforming growth factor beta receptor II propagates the potent antiproliferative effect of TGF- $\beta$  during the normal process of apoptosis that occurs during development in the kidney and after acute vascular injury such as angioplasty. Disruptions of transforming growth factor beta receptor II signaling have been shown to be involved in glomerulosclerosis, a progressive disease characterized by an accumulation of extracellular matrix (ECM) following proliferation of glomerular cells in the kidney (Imai et al., *J. Nephrol.*, 1998, 11, 16-19). In another fibroproliferative vascular disease, lesion-derived human vascular cells exhibit resistance to the antiproliferative effects of TGF- $\beta$  resulting from an acquired mutation in transforming growth factor beta receptor II (Markowitz, *J. Clin. Invest.*, 1997, 100, 2143-2145; McCaffrey et al., *J. Mol. Cell Cardiol.*, 1999, 31, 1627-1642).

Tumor-specific transforming growth factor beta receptor II mutations have been reported in gastrointestinal malignancies (including gastric, esophageal, liver, biliary and pancreatic, and colorectal), in brain tumors, adenocarcinoma, and in cervical, endometrial, breast, head and neck, and small-cell lung cancers (Pasche, *J. Cell Physiol.*, 2001, 186, 153-168). In particular, the vast majority of hereditary nonpolyposis colorectal cancer cell lines with microsatellite instability bear mutations in transforming growth factor beta receptor II and these mutations are associated with the absence of receptors from the cell surface and a loss of responsiveness to TGF- $\beta$ .

-4-

(Markowitz et al., *Science*, 1995, 268, 1336-1338). Mutations in transforming growth factor beta receptor II have also been identified in cell lines derived from recurrent human breast tumors, in subpopulations of human pancreatic adenocarcinomas, and in glioma cancers (Hata, *Exp. Cell Res.*, 2001, 264, 111-116; Lucke et al., *Cancer Res.*, 2001, 61, 482-485; Pasche, *J. Cell Physiol.*, 2001, 186, 153-168; Venkatasubbarao et al., *Genes Chromosomes Cancer*, 1998, 22, 138-144). Furthermore, in patients with Ewing's sarcoma (EWS) and related peripheral primitive neuroectodermal tumors, the EWS/Fli-1 fusion protein resulting from chromosomal translocation has been shown to repress the expression of transforming growth factor beta receptor II (Hahm et al., *Nat Genet*, 1999, 23, 222-227.). Once generated, these mutations lead to an escape from negative growth regulation by TGF- $\beta$  and a proliferative advantage of cells lacking functional type II receptors, which is believed to drive tumor progression and account for a predisposition to cancer.

The modulation of transforming growth factor beta receptor II activity and/or expression is an ideal target for therapeutic intervention aimed at modulating the TGF- $\beta$  signaling pathway in the prevention and treatment of many cancers and fibroproliferative diseases.

For example, many tumor cells have a cytokine mediated immunosuppressive defense mechanism involving the secretion of TGF- $\beta$ , which downregulates the tumoricidal capabilities of antigen-specific T-cells. This immune evasion has been referred to as the tumor "firewall." By inhibiting the function of transforming growth factor beta receptor II in a subpopulation of leukocytes, it should be possible to provide a means of circumventing the tumor firewall by making these immune cells insensitive to the immunosuppressive effects of TGF- $\beta$  (Shah and Lee, *Prostate*, 2000, 45, 167-172).

A transforming growth factor beta receptor II knockout

-5-

mouse has been generated and heterozygous mice are developmentally normal, whereas the homozygous mutation causes defects in yolk sac hematopoiesis and vasculogenesis, resulting in embryonic lethality around 10.5 days of gestation (Oshima et al., *Dev. Biol.*, 1996, 179, 297-302).

Investigative strategies aimed at modulating transforming growth factor beta receptor II function have involved the use of antibodies directed against an N-terminal peptide of transforming growth factor beta receptor II to perturb ligand-receptor binding and functionally block signaling, and the use of antisense oligonucleotides.

In two studies of the role of this receptor in TGF- $\beta$  signaling, a phosphorothioate antisense oligodeoxynucleotide, 16 nucleotides in length, surrounding the translation start site of the murine transforming growth factor beta receptor II was used to show that when the function of murine transforming growth factor beta receptor II is abrogated, lung branching morphogenesis in cultured murine embryonic lung cells is stimulated (Zhao et al., *Dev. Biol.*, 1996, 180, 242-257) and tooth formation in early morphogenesis of mouse mandibular explants is enhanced (Chai et al., *Mech. Dev.*, 1999, 86, 63-74).

Two phosphorothioate antisense oligodeoxynucleotides, 16 nucleotides and 18 nucleotides in length and spanning the translation start site, have been used to inhibit rat transforming growth factor beta receptor II function and show its involvement in branching of embryonic lung explants (Liu et al., *Dev. Dyn.*, 2000, 217, 343-360) and in the TGF- $\beta$ 1 mediated inhibition of MAP kinase activation in cultured rat glomerular epithelial cells, respectively (Higashiyama et al., *Hypertens. Res.*, 1999, 22, 173-180).

Transient transfection of MDA-MB-435 human breast cancer cells with a phosphorothioate antisense oligodeoxynucleotide, 21 nucleotides in length and targeting an unspecified region of transforming growth factor beta receptor II, was found to block

-6-

TGF- $\beta$  -induced apoptosis (Yu et al., *Nutr. Cancer*, 1997, 27, 267-278).

Currently, there are no known therapeutic agents that effectively inhibit the synthesis of transforming growth factor beta receptor II. Consequently, there remains a long felt need for additional agents capable of effectively inhibiting transforming growth factor beta receptor II function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and therefore may prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of transforming growth factor beta receptor II expression.

The present invention provides compositions and methods for modulating transforming growth factor beta receptor II expression, including modulation of the truncated mutants and alternatively spliced forms of transforming growth factor beta receptor II.

#### **SUMMARY OF THE INVENTION**

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding Transforming growth factor beta receptor II, and which modulate the expression of Transforming growth factor beta receptor II. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of Transforming growth factor beta receptor II in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of Transforming growth factor beta receptor II by administering a therapeutically or

prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

#### **DETAILED DESCRIPTION OF THE INVENTION**

The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding Transforming growth factor beta receptor II, ultimately modulating the amount of Transforming growth factor beta receptor II produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding Transforming growth factor beta receptor II. As used herein, the terms "target nucleic acid" and "nucleic acid encoding Transforming growth factor beta receptor II" encompass DNA encoding Transforming growth factor beta receptor II, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of Transforming growth factor beta receptor II. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred

form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding Transforming growth factor beta receptor II. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or

-9-

tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding Transforming growth factor beta receptor II, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-

-10-

most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be

-11-

complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the

-12-

function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with other antisense compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

Expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds which affect expression patterns.

Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, **2000**, 480, 17-24; Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., *Drug Discov. Today*, **2000**, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, **1999**, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U. S. A.*, **2000**, 97, 1976-81), protein arrays and proteomics (Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16; Jungblut, et al., *Electrophoresis*, **1999**, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16; Larsson, et al., *J. Biotechnol.*, **2000**, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., *Anal. Biochem.*, **2000**,

-13-

286, 91-98; Larson, et al., *Cytometry*, **2000**, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, **2000**, 3, 316-21), comparative genomic hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*, **1998**, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, *Eur. J. Cancer*, **1999**, 35, 1895-904) and mass spectrometry methods (reviewed in (To, *Comb. Chem. High Throughput Screen*, **2000**, 3, 235-41).

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The

-14-

antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the

backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphoro-dithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkyl-phosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed

-16-

by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone.

Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylimino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- [wherein the native phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having

similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, also described in examples hereinbelow.

A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methelyne (-CH<sub>2</sub>-)<sub>n</sub> group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 2'-allyl (2'-CH<sub>2</sub>-CH=CH<sub>2</sub>), 2'-O-allyl (2'-O-CH<sub>2</sub>-CH=CH<sub>2</sub>) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920,

-19-

certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl (-C≡C-CH<sub>3</sub>) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example

-20-

7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include sterols, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, *86*, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1994**, *4*, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, **1992**, *660*, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1993**, *3*, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, **1992**, *20*, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, **1991**, *10*, 1111-1118; Kabanov et al., *FEBS Lett.*, **1990**, *259*,

327-330; Svinarchuk et al., *Biochimie*, **1993**, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, **1995**, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, **1990**, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, **1995**, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, **1995**, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, **1995**, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, **1996**, 277, 923-937. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667;

-23-

5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed

-24-

as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized *in vitro* and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the *in vivo* synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721;

-25-

4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine,

ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid,

-27-

ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of Transforming growth factor beta receptor II is treated by administering antisense compounds in accordance with this

-28-

invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding Transforming growth factor beta receptor II, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding Transforming growth factor beta receptor II can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of Transforming growth factor beta receptor II in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at

least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1</sub>-10 alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

Compositions and formulations for oral administration

-30-

include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucholic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodieshydrofusidate,. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and

starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in detail in United States applications 08/886,829 (filed July 1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed February 23, 1999), 09/082,624 (filed May 21, 1998) and 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of

bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p.

-33-

199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase

-34-

and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin

-35-

and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-

-36-

hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to

-37-

form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, *in: Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, *in Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, *in Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, *in Pharmaceutical Dosage Forms*, Lieberman, Rieger and Bunker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (S0750), decaglycerol deaoleate

(DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, **1994**, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, **1993**, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, **1994**, 11, 1385; Ho et al., *J. Pharm. Sci.*, **1996**, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating

thermolabile drugs, peptides or oligonucleotides.

Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

-40-

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic

-41-

absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang *et al.*, *Biochem. Biophys. Res. Commun.*, **1987**, *147*, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou *et al.*, *Journal of Controlled Release*, **1992**, *19*, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from

-42-

dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to

liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G<sub>M1</sub>, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen *et al.*, *FEBS Letters*, **1987**, 223, 42; Wu *et al.*, *Cancer Research*, **1993**, 53, 3765).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos *et al.* (*Ann. N.Y. Acad. Sci.*, **1987**, 507, 64) reported the ability of monosialoganglioside G<sub>M1</sub>, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon *et al.* (*Proc. Natl. Acad. Sci. U.S.A.*, **1988**, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen *et al.*, disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G<sub>M1</sub> or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb *et al.*) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim *et al.*).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto *et al.* (*Bull. Chem. Soc. Jpn.*, **1980**, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum *et al.* (*FEBS Lett.*, **1984**, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycals results in significantly enhanced blood half-lives. Synthetic

-44-

phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov *et al.* (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume *et al.* (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle *et al.* (U.S. Patent No's. 5,013,556 and 5,356,633) and Martin *et al.* (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin *et al.*) and in WO 94/20073 (Zalipsky *et al.*) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi *et al.*). U.S. Patent Nos. 5,540,935 (Miyazaki *et al.*) and 5,556,948 (Tagawa *et al.*) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry *et al.* discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa *et al.* discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman *et al.* describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787

-45-

to Love *et al.* discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, *e.g.* they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and

-46-

ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or

lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

**Surfactants:** In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi *et al.*, *J. Pharm. Pharmacol.*, 1988, 40, 252).

**Fatty acids:** Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (*n*-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (*e.g.*, methyl, isopropyl and *t*-butyl), and mono- and di-glycerides thereof (*i.e.*, oleate,

-48-

laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodieshydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other

-50-

compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

#### Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'isothiocyanostilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

### Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases.

-52-

The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium

carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUDR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual*

-54-

of *Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC<sub>50</sub>s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging

-55-

from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

**EXAMPLES****Example 1****Nucleoside Phosphoramidites for Oligonucleotide Synthesis****Deoxy and 2'-alkoxy amidites**

2'-Deoxy and 2'-methoxy beta-cyanoethyldiisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

**2'-Fluoro amidites****2'-Fluorodeoxyadenosine amidites**

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP

and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

#### **2'-Fluorodeoxyguanosine**

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropyldisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

#### **2'-Fluorouridine**

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-O-(2-Methoxyethyl) modified amidites**

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of

Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

**2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenyl-carbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

**2'-O-Methoxyethyl-5-methyluridine**

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate

-59-

evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL). The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride

-60-

(24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C; and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The

residue was triturated with EtOAc to give the title compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-**

**cytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-cytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra(isopropyl)-phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

**2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites****2'-(Dimethylaminooxyethoxy) nucleoside amidites**

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

**5'-O-tert-Butyldiphenylsilyl- $O^2$ -2'-anhydro-5-methyluridine**  
 **$O^2$ -2'-anhydro-5-methyluridine** (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. *tert*-Butyldiphenylchlorosilane (125.8g,

-63-

119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (Rf 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to

-10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl- $O^2$ -2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone,

-64-

the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine**

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxypthalimide (7.24g, 44.36mmol). It was then dried over  $P_2O_5$  under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

**5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine**

-65-

2'-O-([2-phthalimidoxy)ethyl]-5'-*t*-butyldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH<sub>2</sub>Cl<sub>2</sub> and the combined organic phase was washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solution was concentrated to get 2'-O-(aminoxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%).

**5'-O-*tert*-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine**

5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the

reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

**2'-O-(dimethylaminoxyethyl)-5-methyluridine**

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

**5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine**

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to

-67-

get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

**5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

The 2'-O-aminoxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinoss, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxypthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-[(2-phthalmidoxy)ethyl]-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2-</sup>, 2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and

heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)-ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction

-70-

mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

**Example 2**

**Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as

-71-

described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### **Example 3**

#### **Oligonucleoside Synthesis**

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethyl-hydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

### **Example 4**

**PNA Synthesis**

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

**Example 5****Synthesis of Chimeric Oligonucleotides**

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

**[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and

-73-

the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

**[2'-O-(2-Methoxyethyl)]-[2'-deoxy]-[2'-O-(Methoxyethyl)]**

**Chimeric Phosphorothioate Oligonucleotides**

[2'-O-(2-methoxyethyl)]-[2'-deoxy]-[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

**[2'-O-(2-Methoxyethyl) Phosphodiester]-[2'-deoxy Phosphorothioate]-[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides**

[2'-O-(2-methoxyethyl) phosphodiester]-[2'-deoxy phosphorothioate]-[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3, H-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

-74-

Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

**Example 6**

**Oligonucleotide Isolation**

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* **1991**, *266*, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

**Example 7**

**Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-1,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethylisopropyl phosphoramidites were purchased from

-75-

commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyldiisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated NH<sub>4</sub>OH at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

#### **Example 8**

##### **Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

#### **Example 9**

##### **Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example,

-76-

PCR or Northern blot analysis. The following 6 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

**T-24 cells:**

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

**A549 cells:**

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

**NHDF cells:**

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

**HEK cells:**

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

**HepG2 cells:**

The human hepatoblastoma cell line HepG2 was obtained from the American Type Culture Collection (Manassas, VA). HepG2 cells were routinely cultured in Eagle's MEM supplemented with 10% fetal calf serum, non-essential amino acids, and 1 mM sodium pyruvate (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

**b.END cells:**

The mouse brain endothelial cell line b.END was obtained

-78-

from Dr. Werner Risau at the Max Plank Institute (Bad Nauheim, Germany). b.END cells were routinely cultured in DMEM, high glucose (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 3000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM<sup>TM</sup>-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM<sup>TM</sup>-1 containing 3.75  $\mu$ g/mL LIPOFECTIN<sup>TM</sup> (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCAAGGA**, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold)

with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

**Example 10****Analysis of oligonucleotide inhibition of Transforming growth factor beta receptor II expression**

Antisense modulation of Transforming growth factor beta receptor II expression can be assayed in a variety of ways known in the art. For example, Transforming growth factor beta receptor II mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied

-80-

Biosystems, Foster City, CA and used according to manufacturer's instructions.

Protein levels of Transforming growth factor beta receptor II can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to Transforming growth factor beta receptor II can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

#### **Example 11**

##### **Poly(A)+ mRNA isolation**

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al.,

-81-

*Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

#### **Example 12**

##### **Total RNA Isolation**

Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 100  $\mu$ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection

-82-

tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 µL water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 µL water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

#### **Example 13**

#### **Real-time Quantitative PCR Analysis of Transforming growth factor beta receptor II mRNA Levels**

Quantitation of Transforming growth factor beta receptor II mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in

real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated

-84-

cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN<sup>TM</sup> buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD<sup>TM</sup>, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25  $\mu$ L total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD<sup>TM</sup>, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreen<sup>TM</sup> (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen<sup>TM</sup> RNA quantification reagent from Molecular Probes. Methods of RNA quantification by RiboGreen<sup>TM</sup> are taught in Jones, L.J., et al, *Analytical Biochemistry*, 1998, 265, 368-374.

-85-

In this assay, 175  $\mu$ L of RiboGreen<sup>TM</sup> working reagent (RiboGreen<sup>TM</sup> reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25uL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

Probes and primers to human Transforming growth factor beta receptor II were designed to hybridize to a human Transforming growth factor beta receptor II sequence, using published sequence information (GenBank accession number D50683, incorporated herein as SEQ ID NO:3). For human Transforming growth factor beta receptor II the PCR primers were:  
forward primer: AGAGATCGAGGGCGACCAG (SEQ ID NO: 4)  
reverse primer: TCAACGTCTCACACACCATCTG (SEQ ID NO: 5) and the PCR probe was: FAM-TCCCAGCTTCTGGCTAACCAACCA-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For human GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 7)  
reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 8) and the PCR probe was: 5' JOE-CAAGCTTCCGTTCTCAGCC- TAMRA 3' (SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

Probes and primers to mouse Transforming growth factor beta receptor II were designed to hybridize to a mouse Transforming growth factor beta receptor II sequence, using published sequence information (GenBank accession number D32072, incorporated herein as SEQ ID NO:10). For mouse Transforming growth factor beta receptor II the PCR primers were:

forward primer: CGACCGCTCCGACATCA (SEQ ID NO:11)  
reverse primer: TCGATGGGCAGCAGCTC (SEQ ID NO: 12) and the PCR

-86-

probe was: FAM-CTCCACGTGCGCCAACAAACATCA-TAMRA (SEQ ID NO: 13) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For mouse GAPDH the PCR primers were:

forward primer: GGCAAATTCAACGGCACAGT (SEQ ID NO: 14)

reverse primer: GGGTCTCGCTCCTGGAAGAT (SEQ ID NO: 15) and the PCR probe was: 5' JOE-AAGGCCGAGAATGGGAAGCTTGTACATC- TAMRA 3' (SEQ ID NO: 16) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

#### **Example 14**

#### **Northern blot analysis of Transforming growth factor beta receptor II mRNA levels**

Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA) and then robed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions.

To detect human Transforming growth factor beta receptor II, a human Transforming growth factor beta receptor II specific probe was prepared by PCR using the forward primer AGAGATCGAGGGCGACCAAG (SEQ ID NO: 4) and the reverse primer TCAACGTCTCACACACCATCTG (SEQ ID NO: 5). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

To detect mouse Transforming growth factor beta receptor II, a mouse Transforming growth factor beta receptor II specific probe was prepared by PCR using the forward primer CGACCGCTCCGACATCA (SEQ ID NO: 11) and the reverse primer TCGATGGGCAGCAGCTC (SEQ ID NO: 12). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for mouse glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

#### **Example 15**

#### **Antisense inhibition of human Transforming growth factor beta receptor II expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human Transforming growth factor beta receptor II RNA, using published sequences (GenBank accession number D50683, incorporated herein as SEQ ID NO: 3, GenBank accession number NM\_003242, incorporated herein as SEQ ID NO: 17, GenBank accession number D50682, incorporated herein as SEQ ID NO: 18,

-88-

and GenBank accession number AW020512, incorporated herein as SEQ ID NO: 19). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human Transforming growth factor beta receptor II mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1  
Inhibition of human Transforming growth factor beta receptor II mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
123751	Start Codon	3	1568	cagcccccgaccatggcag	0	20
123760	Coding	3	1768	ctgcagttgctcatgcagga	34	21
123772	Coding	3	2266	ttgtggttgatgttgtggc	61	22
123777	Coding	3	2350	ttctgcttcagctggcctt	0	23
123786	Coding	3	2593	aggtcctcccagctgtatgac	0	24
123788	Coding	3	2674	ggcatcttgggcctccaca	27	25
123793	Coding	3	2787	catccacagacagacttaggg	26	26
123794	Coding	3	2794	gccaggtcatccacagacag	41	27
123797	Coding	3	2908	agagccatggagtagacatc	48	28
123799	Coding	3	2992	tcccgcaccttggAACAAA	0	29
123803	Coding	3	3073	tggttgagccagaagctggg	8	30
123804	Coding	3	3083	gatgccctggtggttagcc	17	31
128885	5'UTR	17	34	ctcaactcaactcaactcag	27	32
128886	5'UTR	17	89	cctgtcccggcgggtgcac	40	33
128887	5'UTR	17	121	gcggccgaggaaagctgcac	38	34

-89-

128888	5'UTR	3	445	atatgaaagggttattacaa	0	35
128889	5'UTR	3	591	aaaaaactgcattcaacat	0	36
128890	5'UTR	3	618	tctacacacttagttaagga	23	37
128891	5'UTR	3	688	cttagggttccatgtttcc	18	38
128892	5'UTR	3	1065	ataaaagactgtaatcgtat	0	39
128893	5'UTR	3	1144	gtttacaagaacagtccaaa	10	40
128894	5'UTR	3	1159	aagatgcatacacctgttta	40	41
128895	5'UTR	3	1249	catcccagtgaggctttct	60	42
128896	5'UTR	3	1324	gagtctagacatattggtgg	0	43
128897	5'UTR	3	1335	tgttagcctctgagtctaga	44	44
128898	5'UTR	3	1346	actagcagtcatgttagcct	16	45
128899	5'UTR	3	1424	agcttgtgactttaaagg	15	46
128900	5'UTR	3	1551	cagaccccgctgcgtcat	51	47
128901	Start Codon	3	1567	agcccccgaccatggcaga	19	48
128902	Coding	3	1658	gtcggttattaaccgacttct	40	49
128903	Coding	3	1739	gtcacaggtggaaaatctca	66	50
128904	Coding	3	2412	aaggaggcatactcctcatag	26	51
128905	Coding	3	2447	attgatgtctgagaagatgt	0	52
128906	Coding	3	2756	ggaaagcccaaagtcacaca	47	53
128907	Coding	3	2873	ctcagcatttccaaattca	1	54
128908	Coding	3	2938	ttacagcgagatgtcatttc	0	55
128909	Coding	3	3045	gtcgccctcgatctctcaac	48	56
128910	Coding	3	3236	gccgtcttcaggaatcttct	50	57
128911	Stop Codon	3	3265	cagaagagctatttggtagt	24	58
128912	3'UTR	3	3469	agcttatccatgacaatgt	0	59
128913	3'UTR	3	3499	caatctcatttcctgaggaa	0	60
128914	3'UTR	3	3623	aggtaggcataatataatata	34	61
128915	3'UTR	3	3682	agttctccaataaaaccaat	18	62
128916	3'UTR	3	3746	ctgcattgtgttgcataaa	47	63
128917	3'UTR	3	3840	tttgcaaaagcaagtgcata	21	64
128918	3'UTR	3	3938	taagggcacacaggaacccc	35	65
128919	3'UTR	3	3947	caggagaaataagggcacac	47	66
128920	3'UTR	3	4027	gttaggtttctaaacttaggt	58	67
128921	3'UTR	3	4083	tgcaacccatgaaggtaaaa	50	68
128922	3'UTR	3	4208	aaaggatgaaggctggagc	2	69
128923	3'UTR	3	4304	ccattgggtttgtatagaa	6	70
128924	3'UTR	3	4326	caggagcccagaaagatgga	20	71
128925	3'UTR	3	4333	gagcaatcaggagcccagaa	76	72
128926	3'UTR	3	4462	aaagtggcttcacttttga	38	73
128927	3'UTR	3	4540	ggactggagataactgaaaa	36	74
128928	3'UTR	3	4569	gtctccacacccatcacatt	10	75
128929	3'UTR	3	4628	ccaggttaggcagtggaaaga	42	76
128930	3'UTR	3	4789	cttaaaggagttcccttta	29	77
128931	3'UTR	3	4931	ttctgcaaagggttggagaagg	38	78
128932	3'UTR	3	4999	tttgtggacacaaatttctta	0	79
128933	3'UTR	3	5062	acacttcttgcatttgcata	31	80
128934	3'UTR	3	5172	ccatccaaacagagctgata	0	81
128935	3'UTR	3	5412	ttgaatatctcatgaatgga	28	82
128936	3'UTR	3	5413	cttgaatatctcatgaatgg	0	83
128937	3'UTR	3	5692	taaaaataagtgcgttggagac	20	84
128938	3'UTR	3	5731	aaacattttatttgcataaa	0	85

-90-

128939	5'UTR	18	312	actatgccactttacttcat	0	86
128940	5'UTR	18	344	gggttatgcagtgcggcccttt	30	87
128941	5'UTR	18	414	gtggctgtacagacaccttcc	0	88
128942	5'UTR	18	491	attagtccccagaatcctgt	0	89
128943	5'UTR	18	497	acatgaatttagtccccagaa	0	90
128944	5'UTR	18	519	atatgtcttgtaaattcagt	62	91
128945	5'UTR	18	578	ccttcctactgtatgagattt	0	92
128946	5'UTR	18	665	ggtcattatgttaaaaccatt	42	93
128947	5'UTR	18	738	tctcaaattccatggctcaa	37	94
128948	5'UTR	18	854	acctcagcctcccaagtagc	5	95
128949	3'UTR	19	286	aaatggatttagctactagg	23	96
128950	3'UTR	19	287	caaatggatttagctactag	22	97

As shown in Table 1, SEQ ID NOS 21, 22, 27, 28, 33, 34, 41, 42, 44, 47, 49, 50, 53, 56, 57, 61, 63, 65, 66, 67, 68, 72, 73, 74, 76, 78, 80, 87, 91, 93 and 94 demonstrated at least 30% inhibition of human Transforming growth factor beta receptor II expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

#### Example 16

##### Antisense inhibition of mouse Transforming growth factor beta receptor II expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap.

In accordance with the present invention, a second series of oligonucleotides were designed to target different regions of the mouse Transforming growth factor beta receptor II RNA, using published sequences (GenBank accession number D32072, incorporated herein as SEQ ID NO: 10). The oligonucleotides are shown in Table 2. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides,

-91-

which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on mouse Transforming growth factor beta receptor II mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 2

Inhibition of mouse Transforming growth factor beta receptor II mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
123751	Start Codon	10	151	cagccccgacccatggcag	78	20
123760	Coding	10	426	ctgcagttgctcatgcagga	13	21
123772	Coding	10	924	tttgtggttgatgttgtggc	63	22
123777	Coding	10	1008	ttctgcttcagcttggcctt	65	23
123786	Coding	10	1251	aggtcctcccagctgatgac	71	24
123788	Coding	10	1332	ggcatcttgggcctccaca	61	25
123793	Coding	10	1445	catccacagacagactagg	29	26
123794	Coding	10	1452	gccaggtcatccacagacag	80	27
123797	Coding	10	1566	agagccatggagtagacatc	69	28
123799	Coding	10	1650	tcccgcaccttggAACCAA	67	29
123803	Coding	10	1731	tggttgagccagaaggctggg	70	30
123804	Coding	10	1741	gatgccctgggtggtagcc	62	31
123745	5'UTR	10	7	ctcgaggcccggaggaggc	38	98
123746	5'UTR	10	18	gatccccggagctggaggc	63	99
123747	5'UTR	10	35	ggccagatgtggccggcgat	44	100
123748	5'UTR	10	78	ggaccccggctgcgcctt	78	101
123749	5'UTR	10	105	cctgggtgcgccacgaaccga	70	102
123750	5'UTR	10	132	gcccccgtcgctcgatcgat	77	103
123752	Start Codon	10	156	cggagcagccccgacccat	82	104
123753	Coding	10	166	ccacaggccccggagcagcc	80	105
123754	Coding	10	196	gatgcgcgtccacaggacga	58	106
123755	Coding	10	206	tcgtgctggcgatgcgcgtc	47	107
123756	Coding	10	216	tgcggcgggatcgtgctggc	63	108
123757	Coding	10	290	ggatggccattacagctt	61	109

-92-

123758	Coding	10	368	acttgcacagctgtggaagc	69	110
123759	Coding	10	416	tcatgcaggacttctggttg	60	111
123761	Coding	10	553	ttccagagtgaagccgtggt	55	112
123762	Coding	10	598	cgcctttctttccttca	56	113
123763	Coding	10	606	gtctcgcccgcctttctt	81	114
123764	Coding	10	684	ggactgctgggtgttattc	74	115
123765	Coding	10	738	agcggaggcaggaggctgac	68	116
123766	Coding	10	753	gctatggcaatccccagcgg	88	117
123767	Coding	10	758	tgacagctatggcaatcccc	65	118
123768	Coding	10	804	ctcagcttctgctgcccgtg	44	119
123769	Coding	10	881	ggtcgtcctccaggatgatg	32	120
123770	Coding	10	905	cgcacgtggagctgatgtcg	78	121
123771	Coding	10	918	ttgatgttgttggcgcacgt	67	122
123773	Coding	10	929	ccgtgttgtgggtatgttg	76	123
123774	Coding	10	944	cgatgggcagcagctccgtg	72	124
123775	Coding	10	951	tccagctcgatggcagcag	74	125
123776	Coding	10	993	gcctttagacacctcgccgaa	0	126
123778	Coding	10	1040	tgacagccacggtctcaaac	81	127
123779	Coding	10	1134	aggaactgcaggatgtctc	49	128
123780	Coding	10	1144	ctcggccgtcaggaactgca	36	129
123781	Coding	10	1150	ccgctcctcgccgtcagga	45	130
123782	Coding	10	1185	gtgatcagccagtaactgctt	63	131
123783	Coding	10	1208	ggttgcccttcgcgtgaaac	78	132
123784	Coding	10	1218	tactcctgcagggtgcctt	74	133
123785	Coding	10	1239	ctgatgacatgcctcgtag	79	134
123787	Coding	10	1276	ggccaggagctgcccagct	86	135
123789	Coding	10	1350	aggccctgtgaacaatggg	83	136
123790	Coding	10	1364	tgttagagctttaggtcc	63	137
123791	Coding	10	1380	tcgttcttcaactaggatgtt	78	138
123792	Coding	10	1424	ccaggcgcaaggacagcccg	64	139
123795	Coding	10	1515	ttcatcctggattctagaac	29	140
123796	Coding	10	1558	ggagtagacatccgtctgct	28	141
123798	Coding	10	1593	cagcgggacgtcatttccca	30	142
123800	Coding	10	1674	ttcatgctctccacacaggg	66	143
123801	Coding	10	1709	tttccggccgcctcggtct	55	144
123802	Coding	10	1718	agctggaaatttccggccgc	56	145
123805	Coding	10	1752	cacacgatctggatgcctg	79	146
123806	Coding	10	1779	tggcccagcactcggtcaa	70	147
123807	Coding	10	1879	cttctcctgggagcagctcc	31	148
123808	Stop Codon	10	1923	agaaaaagctattggtagt	47	149
123809	3'UTR	10	1935	cccagcctgcccagaaaaag	77	150
123810	3'UTR	10	2034	ggaagaaggagcaagtcct	0	151
123811	3'UTR	10	2086	ctatggttgttgcgtctgct	59	152
123812	3'UTR	10	2151	tgcctatgacagtctccaagg	77	153
123813	3'UTR	10	2259	tataaaacatagctattcgt	73	154
123814	3'UTR	10	2422	taggaaggagagtgtttagt	40	155
123815	3'UTR	10	2527	ttgtttctgagccagacagt	71	156
123816	3'UTR	10	2785	cagttcaactcccttttaatg	22	157
123817	3'UTR	10	2870	acagaacctcacacctggcg	63	158
123818	3'UTR	10	2977	agagagagaaaggcaagctg	36	159
123819	3'UTR	10	3033	cttaaaccacatcagaaatc	66	160
123820	3'UTR	10	3110	tattttagtgtctttctgggt	65	161

-93-

123821	3'UTR	10	3452	gattgttcattcagatgaga	10	162
123822	3'UTR	10	3804	gtgtggtattattcaggatc	19	163

As shown in Table 2, SEQ ID NOS 20, 22, 23, 24, 25, 27, 28, 29, 30, 31, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 121, 122, 123, 124, 125, 127, 128, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 143, 144, 145, 146, 147, 149, 150, 152, 153, 154, 155, 156, 158, 160 and 161 demonstrated at least 40% inhibition of mouse Transforming growth factor beta receptor II expression in this experiment and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

#### **Example 17**

##### **Western blot analysis of Transforming growth factor beta receptor II protein levels**

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to Transforming growth factor beta receptor II is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

**What is claimed is:**

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding Transforming growth factor beta receptor II, wherein said compound specifically hybridizes with said nucleic acid molecule encoding Transforming growth factor beta receptor II and inhibits the expression of Transforming growth factor beta receptor II.

2. The compound of claim 1 which is an antisense oligonucleotide.

3. The compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 21, 22, 27, 28, 33, 34, 41, 42, 44, 47, 49, 50, 53, 56, 57, 61, 63, 65, 66, 67, 68, 72, 73, 74, 76, 78, 80, 87, 91, 93, 94, 20, 23, 24, 25, 29, 30, 31, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 121, 122, 123, 124, 125, 127, 128, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 143, 144, 145, 146, 147, 149, 150, 152, 153, 154, 155, 156, 158, 160 or 161.

4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.

6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.

10. The compound of claim 2 wherein the antisense

oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding Transforming growth factor beta receptor II.

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of Transforming growth factor beta receptor II in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of Transforming growth factor beta receptor II is inhibited.

16. A method of treating an animal having a disease or condition associated with Transforming growth factor beta receptor II comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of Transforming growth factor beta receptor II is inhibited.

17. The method of claim 16 wherein the disease or condition is a hyperproliferative disorder.

18. The method of claim 17 wherein the hyperproliferative disorder is cancer.

19. The method of claim 18 wherein the cancer is lung, liver, bone, breast, cervical, colon, gastric, pancreatic, esophageal, or hematopoetic cancer.

20. The method of claim 16 wherein the disease or condition involves activation of the immune system.

## SEQUENCE LISTING

<110> Susan Murray  
Jacqueline R. Wyatt  
Isis Pharmaceuticals, Inc.

<120> ANTISENSE MODULATION OF TRANSFORMING GROWTH  
FACTOR BETA RECEPTOR II EXPRESSION

<130> RTSP-0383

<150> 09/888,361  
<151> 2001-06-21

<160> 163

<210> 1  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1  
tccgtcatcg ctcctcaggg 20

<210> 2  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2  
atgcattctg cccccaagga 20

<210> 3  
<211> 5759  
<212> DNA  
<213> Homo sapiens

<220>

<221> CDS  
<222> (1573) ... (3276)

<400> 3  
ataccctgtg gtgctagtgt aaaatcaaat ataatctacc tcttagctt ttctgttatg 60  
taattggaat gaaaaaatag tgctttttt ttttttcat tttttagatt gagttaccat 120  
ggagggaaag aacatgtgtt ttaccagtgg aacgcattct ttgtgttatac tcttcgattt 180  
ttaacatttt atgtttccag tttataaact ttgcctccag ttgtatgaag aaaatttagc 240  
tattagaaag tatttaaaat ctaaaaagta tgacatttta aaatgttagc aggtttaaag 300

ataacttagt ttttcaactc ttaacccaaa attatacttg tatactataa tttgtgttga	360
attacttggt attatttatac atcctagtaa gatctatttt atataatttg attcaacttt	420
tatattgaat tatgttaatt gttttgtaa tacacccccc atatccccc aaagaaaaact	480
agtgttgggt tattgctaattttgttgcgt ggtttggtaa ttacatataaaaattat	540
tgtttaaaaa attgcaaaagt gatcatttt tgacacatca gcaagaattt atgttggaaat	600
gcaagttttt tccccatcc ttaactaagt gtgttagatat ctttggggttt tttttttat	660
agatagcaaa caataaaagat gcattgagga agacatggaa ccctaagttt acattgagaa	720
gtcactttga tggcatccga gcccttgctt tccatcccat tgagcctgtt ttgataacag	780
catcagagga tcacacatataaaaatgtgaa atttacagaa aacagccccca gccaaaaagt	840
gagaatattc tacttttaca ttatttgaat attttaagta acattttcag cagaaaaacta	900
catattttat gatttctgtt aaatttccca aacatttctt ctagttctt cttccatgtc	960
atgttaatttta ttgtgtttt ccatttatac aacctaattgc taggttagaa aactgtatgc	1020
taggaagata tggattttatc ttttgggttc accagggttt gtgaataactg attacagtct	1080
ttatacataa cagtattttt aaatggctgc tatatttggg gagcccccttgc tctatactga	1140
agggttggac tggttttgta aacaggtata tgcatcttta aaataatcaa atgtccaccc	1200
ccaccccttgc ccatttgccttta ctagatttag acttattgaga tggagcagag aaaagccctca	1260
ctggatgggg tggagggtct tggctttgtc ttttggccctt gagacttttag gtactttcca	1320
tacccaccaa tatgtctaga ctcagaggct aacatgactg ctagtgcctca cttggccct	1380
ttagttcctg tctaaggcatg gatagacatg agtttagata aaccctttaa aagtcacaaa	1440
gctaaatctg gttctacttg cttgggtctt tattggagct ccactccctt gggtcacatg	1500
tcaaaaaaaaaa aaaaaaaaaaag gcgcgcgtccg ctgcgcgtggg ggctcggtct atgacgagca	1560
gcgggggtctg cc atg ggt cgg ggg ctg ctc agg ggc ctg tgg ccg ctg cac	1611
Met Gly Arg Gly Leu Leu Arg Gly Leu Trp Pro Leu His	
1 5 10	
atc gtc ctg tgg acg cgt atc gcc agc acg atc cca ccg cac gtt cag	1659
Ile Val Leu Trp Thr Arg Ile Ala Ser Thr Ile Pro Pro His Val Gln	
15 20 25	
aag tcg gtt aat aac gac atg ata gtc act gac aac aac ggt gca gtc	1707
Lys Ser Val Asn Asn Asp Met Ile Val Thr Asp Asn Asn Gly Ala Val	
30 35 40 45	
aag ttt cca caa ctg tgt aaa ttt tgt gat gtg aga ttt tcc acc tgt	1755
Lys Phe Pro Gln Ile Cys Lys Phe Cys Asp Val Arg Phe Ser Thr Cys	
50 55 60	
gac aac cag aaa tcc tgc atg agc aac tgc agc atc acc tcc atc tgt	1803
Asp Asn Gln Lys Ser Cys Met Ser Asn Cys Ser Ile Thr Ser Ile Cys	
65 70 75	
gag aag cca cag gaa gtc tgt gtg gct gta tgg aga aag aat gac gag	1851
Glu Lys Pro Gln Glu Val Cys Val Ala Val Trp Arg Lys Asn Asp Glu	

80

85

90

aac ata aca cta gag aca gtt tgc cat gac ccc aag ctc ccc tac cat	1899
Asn Ile Thr Leu Glu Thr Val Cys His Asp Pro Lys Leu Pro Tyr His	
95 100 105	
gac ttt att ctg gaa gat gct gct tct cca aag tgc att atg aag gaa	1947
Asp Phe Ile Leu Glu Asp Ala Ala Ser Pro Lys Cys Ile Met Lys Glu	
110 115 120 125	
aaa aaa aag cct ggt gag act ttc ttc atg tgt tcc tgt agc tct gat	1995
Lys Lys Lys Pro Gly Glu Thr Phe Phe Met Cys Ser Cys Ser Ser Asp	
130 135 140	
gag tgc aat gac aac atc atc ttc tca gaa gaa tat aac acc agc aat	2043
Glu Cys Asn Asp Asn Ile Ile Phe Ser Glu Glu Tyr Asn Thr Ser Asn	
145 150 155	
cct gac ttg ttg cta gtc ata ttt caa gtg aca ggc atc agc ctc ctg	2091
Pro Asp Leu Leu Leu Val Ile Phe Gln Val Thr Gly Ile Ser Leu Leu	
160 165 170	
cca cca ctg gga gtt gcc ata tct gtc atc atc atc ttc tac tgc tac	2139
Pro Pro Leu Gly Val Ala Ile Ser Val Ile Ile Ile Phe Tyr Cys Tyr	
175 180 185	
cgc gtt aac cgg cag cag aag ctg agt tca acc tgg gaa acc ggc aag	2187
Arg Val Asn Arg Gln Gln Lys Leu Ser Ser Thr Trp Glu Thr Gly Lys	
190 195 200 205	
acg cgg aag ctc atg gag ttc agc gag cac tgt gcc atc atc ctg gaa	2235
Thr Arg Lys Leu Met Glu Phe Ser Glu His Cys Ala Ile Ile Leu Glu	
210 215 220	
gat gac cgc tct gac atc agc tcc acg tgt gcc aac aac atc aac cac	2283
Asp Asp Arg Ser Asp Ile Ser Ser Thr Cys Ala Asn Asn Ile Asn His	
225 230 235	
aac aca gag ctg ctg ccc att gag ctg gac acc ctg gtg ggg aaa ggt	2331
Asn Thr Glu Leu Leu Pro Ile Glu Leu Asp Thr Leu Val Gly Lys Gly	
240 245 250	
cgc ttt gct gag gtc tat aag gcc aag ctg aag cag aac act tca gag	2379
Arg Phe Ala Glu Val Tyr Lys Ala Lys Leu Lys Gln Asn Thr Ser Glu	
255 260 265	
cag ttt gag aca gtc gca gtc aag atc ttt ccc tat gag gag tat gcc	2427
Gln Phe Glu Thr Val Ala Val Lys Ile Phe Pro Tyr Glu Glu Tyr Ala	
270 275 280 285	
tct tgg aag aca gag aag gac atc ttc tca gac atc aat ctg aag cat	2475
Ser Trp Lys Thr Glu Lys Asp Ile Phe Ser Asp Ile Asn Leu Lys His	
290 295 300	
gag aac ata ctc cag ttc ctg acg gct gag gag cgg aag acg gag ttg	2523
Glu Asn Ile Leu Gln Phe Leu Thr Ala Glu Glu Arg Lys Thr Glu Leu	
305 310 315	
ggg aaa caa tac tgg ctg atc acc gcc ttc cac gcc aag ggc aac cta	2571
Gly Lys Gln Tyr Trp Leu Ile Thr Ala Phe His Ala Lys Gly Asn Leu	
320 325 330	
cag gag tac ctg acg cgg cat gtc atc agc tgg gag gac ctg cgc aag	2619
Gln Glu Tyr Leu Thr Arg His Val Ile Ser Trp Glu Asp Leu Arg Lys	

335	340	345	
ctg ggc agc tcc ctc gcc cg <sup>g</sup> ggg att gct cac ctc cac agt gat cac Leu Gly Ser Ser Leu Ala Arg Gly Ile Ala His Leu His Ser Asp His 350 355 360 365			2667
act cca tgt ggg agg ccc aag atg ccc atc gtg cac agg gac ctc aac Thr Pro Cys Gly Arg Pro Lys Met Pro Ile Val His Arg Asp Leu Asn 370 375 380			2715
agc tcc aat atc ctc gtg aag aac gac cta acc tgc tgc ctg tgt gac Ser Ser Asn Ile Leu Val Lys Asn Asp Leu Thr Cys Cys Leu Cys Asp 385 390 395			2763
ttt ggg ctt tcc ctg cgt ctg gac cct act ctg tct gtg gat gac ctg Phe Gly Leu Ser Leu Arg Leu Asp Pro Thr Leu Ser Val Asp Asp Leu 400 405 410			2811
gct aac agt ggg cag gtg gga act gca aga tac atg gct cca gaa gtc Ala Asn Ser Gly Gln Val Gly Thr Ala Arg Tyr Met Ala Pro Glu Val 415 420 425			2859
cta gaa tcc agg atg aat ttg gag aat gct gag tcc ttc aag cag acc Leu Glu Ser Arg Met Asn Leu Glu Asn Ala Glu Ser Phe Lys Gln Thr 430 435 440 445			2907
gat gtc tac tcc atg gct ctg gtg ctc tgg gaa atg aca tct cgc tgt Asp Val Tyr Ser Met Ala Leu Val Leu Trp Glu Met Thr Ser Arg Cys 450 455 460			2955
aat gca gtg gga gaa gta aaa gat tat gag cct cca ttt ggt tcc aag Asn Ala Val Gly Glu Val Lys Asp Tyr Glu Pro Pro Phe Gly Ser Lys 465 470 475			3003
gtg cgg gag cac ccc tgt gtc gaa agc atg aag gac aac gtg ttg aga Val Arg Glu His Pro Cys Val Glu Ser Met Lys Asp Asn Val Leu Arg 480 485 490			3051
gat cga ggg cga cca gaa att ccc agc ttc tgg ctc aac cac cag ggc Asp Arg Gly Arg Pro Glu Ile Pro Ser Phe Trp Leu Asn His Gln Gly 495 500 505			3099
atc cag atg gtg tgt gag acg ttg act gag tgc tgg gac cac gac cca Ile Gln Met Val Cys Glu Thr Leu Thr Glu Cys Trp Asp His Asp Pro 510 515 520 525			3147
gag gcc cgt ctc aca gcc cag tgt gtg gca gaa cgc ttc agt gag ctg Glu Ala Arg Leu Thr Ala Gln Cys Val Ala Glu Arg Phe Ser Glu Leu 530 535 540			3195
gag cat ctg gac agg ctc tcg ggg agg agc tgc tcg gag gag aag att Glu His Leu Asp Arg Leu Ser Gly Arg Ser Cys Ser Glu Glu Lys Ile 545 550 555			3243
cct gaa gac ggc tcc cta aac act acc aaa tag ctcttctggg gcaggctggg Pro Glu Asp Gly Ser Leu Asn Thr Thr Lys 560 565			3296
ccatgtccaa agaggctgcc cctctcacca aagaacagag gcagcaggaa gctgcccctg aactgatgct tcctggaaaa ccaagggggt cactccctc cctgttaagct gtggggataa gcagaaacaa cagcagcagg gagtgggtga catagagcat tctatgcctt ttacattgtc			3356 3416 3476

ataggataag ctgtgttagc acttcctcag gaaatgagat tgattttaca atagccaata 3536  
acatttgcac ttttattaatg cctgtatata aatatgaata gctatgtttt atatatatat 3596  
atatatatct atatatgtct atagctctat atatatagcc ataccttcaa aagagacaag 3656  
gaaaaacatc aaatattccc agggaaattgg ttttattgga gaactccaga accaagcaga 3716  
gaaggaaggg acccatgaca gcattagcat ttgacaatca cacatgcagt ggttctctga 3776  
ctgtaaaaca gtgaactttg catgaggaaa gaggctccat gtctcacagc cagctatgac 3836  
cacattgcac ttgctttgc aaaataatca ttccctgcct agcacttctc ttctggccat 3896  
ggaactaagt acagtggcac tgttttagga ccagtgtcc cggggttcct gtgtgcctt 3956  
atttctcctg gacttttcat ttaagctcca agccccaaat ctggggggct agtttagaaa 4016  
ctctccctca acctagtttta gaaactctac cccatcttta ataccttcaa tgtttgaac 4076  
cccactttt accttcatgg gttcagaaa aatcagaaca gatgtccccca tccatgcgt 4136  
tgccccacca tctactaatg aaaaattgtt cttttttca tctttccct gcacttatgt 4196  
tactattctc tgctcccagc cttcatcctt ttctaaaaag gagcaaattc tcactctagg 4256  
ctttatctg tttactttt cattacactt gacttgattt tctagtttc tatacaaaca 4316  
ccaatgggtt ccacatcttctt gggctcctga ttgctcaagc acagtttggc ctgatgaaga 4376  
ggatttcaac tacacaatac tatcattgtc aggactatga cctcaggcac tctaaacata 4436  
tgtttgttt ggtcagcaca gcgtttcaaa aagtgaagcc actttataaa tatttggaga 4496  
ttttgcagga aaatctggat cccaggtaa ggatagcaga tggtttcag ttatctccag 4556  
tccacgttca caaaatgtga aggtgtggag acacttacaa agctgcctca cttctcactg 4616  
taaacattag ctctttcac tgcttacctg gacccagtc taggaattaa atctgcaccc 4676  
aaccaggcgc ctttgtaaga aatgtccatt caagcagtca ttctctgggt atataatatg 4736  
atttgacta ccttatctgg tgttaagatt tgaagtggc cttttattgg actaaagggg 4796  
aactccctta agggtctcag ttggccaaag tttctttgc ttatatgtta atagtttac 4856  
cctctgcatt ggagagagga gtgtttact ccaagaagct ttctcatgg ttaccgttct 4916  
ctccatcatg ccagccttctt caacccttgc agaaattact agagaggatt tgaatgtggg 4976  
acacaaagggt cccatggca gttagaaaat ttgtgtccac aaggacaaga acaaagtatg 5036  
agctttaaaa ctccatagga aacttgttaa tcaacaaaga agtgttaatg ctgcaagtaa 5096  
tctctttttt aaaactttttt gaagctactt attttcagcc aaataggaat attagagagg 5156  
gactggtagt gagaatatca gctctgtttg gatggtgaa ggtctcattt tattgagatt 5216  
tttaagatac atgcaaagggt ttggaaatag aacctctagg caccctcctc agtgtgggtg 5276  
ggctgagagt taaagacagt gtggctgcag tagcatagag ggcgcctagaa attccacttg 5336  
caccgttaggg catgctgata ccatcccaat agctgttgcc cattgacctc tagtggtgag 5396

tttctagaat actggtccat tcatgagata ttcaagattc aagagtattc tcacttctgg 5456  
gttatcagca taaaactggaa tggactgtca gaggatactg tggcttgttt tgtttatgtt 5516  
ttttttctt attcaagaaa aaagaccaag gaataacatt ctgtagttcc taaaaaatact 5576  
gacttttttc actactatac ataaaggaa agttttattc ttttatggaa cacttcagct 5636  
gtactcatgt attaaaatag gaatgtgaat gctatatact ctttttatata caaaagtctc 5696  
aagcacttat ttttattcta tgcattgttt gtctttaca taaataaaaat gtttattaga 5756  
ttg 5759

<210> 4  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 4  
agagatcgag ggcgaccag

19

<210> 5  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 5  
tcaacgtctc acacaccatc tg

22

<210> 6  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 6  
tcccagcttc tggctcaacc acca

24

<210> 7  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 7

gaaggtaag gtcggagtc 19  
 <210> 8  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> PCR Primer  
 <400> 8  
 gaagatggtg atgggatttc 20  
 <210> 9  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> PCR Probe  
 <400> 9  
 caagcttccc gttctcagcc 20  
 <210> 10  
 <211> 3839  
 <212> DNA  
 <213> Mus musculus  
 <220>  
 <221> CDS  
 <222> (156) ... (1934)  
 <400> 10  
 ccgggggcct cctccgggcc tccgagctcc ggggatcgcc ggccacatct ggcccgcatc 60  
 ctgagagggc gaggagtaaa ggccgcagccc ggggtccccg aggctcggtt cgtggcgcac  
 caggggcccgg tctatgacga gcgcacggggg ctgcc atg ggt cgg ggg ctg ctc 120  
 Met Gly Arg Gly Leu Leu  
 1 5  
 cg ggc ctg tgg ccg ctg cat atc gtc ctg tgg acg cgc atc gcc agc 221  
 Arg Gly Leu Trp Pro Leu His Ile Val Leu Trp Thr Arg Ile Ala Ser  
 10 15 20  
 acg atc ccg ccg cac gtt ccc aag tcg gat gtg gaa atg gaa gcc cag 269  
 Thr Ile Pro Pro His Val Pro Lys Ser Asp Val Glu Met Glu Ala Gln  
 25 30 35  
 aaa gat gca tcc atc cac cta agc tgt aat agg acc atc cat cca ctg 317  
 Lys Asp Ala Ser Ile His Leu Ser Cys Asn Arg Thr Ile His Pro Leu  
 40 45 50  
 aaa cat ttt aac agt gat gtc atg gcc agc aat ggc ggt ggc gtc 365  
 Lys His Phe Asn Ser Asp Val Met Ala Ser Asp Asn Gly Gly Ala Val  
 55 60 65 70  
 aag ctt cca caq ctq aq ttt tqc qat qtq aqa ctq tcc act tqc 413

Lys Leu Pro Gln Leu Cys Lys Phe Cys Asp Val Arg Leu Ser Thr Cys	75	80	85	
gac aac cag aag tcc tgc atg agc aac tgc agc atc acg gcc atc tgt				461
Asp Asn Gln Lys Ser Cys Met Ser Asn Cys Ser Ile Thr Ala Ile Cys				
90	95	100		
gag aag ccg cat gaa gtc tgc gtg gcc gtg tgg agg aag aac gac aag				509
Glu Lys Pro His Glu Val Cys Val Ala Val Trp Arg Lys Asn Asp Lys				
105	110	115		
aac att act ctg gag acg gtt tgc cac gac ccc aag ctc acc tac cac				557
Asn Ile Thr Leu Glu Thr Val Cys His Asp Pro Lys Leu Thr Tyr His				
120	125	130		
ggc ttc act ctg gaa gat gcc gct tct ccc aag tgt gtc atg aag gaa				605
Gly Phe Thr Leu Glu Asp Ala Ala Ser Pro Lys Cys Val Met Lys Glu				
135	140	145	150	
aag aaa agg gcg ggc gag act ttc ttc atg tgt gcc tgt aac atg gaa				653
Lys Lys Arg Ala Gly Glu Thr Phe Phe Met Cys Ala Cys Asn Met Glu				
155	160	165		
gag tgc aac gat tac atc atc ttt tcg gaa gaa tac acc acc agc agt				701
Glu Cys Asn Asp Tyr Ile Ile Phe Ser Glu Glu Tyr Thr Ser Ser				
170	175	180		
ccc gac ctg ttg ttg gtc att atc caa gtg acg ggt gtc agc ctc ctg				749
Pro Asp Leu Leu Val Ile Ile Gln Val Thr Gly Val Ser Leu Leu				
185	190	195		
cct ccg ctg ggg att gcc ata gct gtc atc atc ttc tac tgc tac				797
Pro Pro Leu Gly Ile Ala Ile Ala Val Ile Ile Ile Phe Tyr Cys Tyr				
200	205	210		
cgt gtc cac cgg cag cag aag ctg agc ccg tcc tgg gag agc agc aag				845
Arg Val His Arg Gln Gln Lys Leu Ser Pro Ser Trp Glu Ser Ser Lys				
215	220	225	230	
ccc cgg aaa ctg atg gat ttc agt gac aat tgt gcc atc atc ctg gag				893
Pro Arg Lys Leu Met Asp Phe Ser Asp Asn Cys Ala Ile Ile Leu Glu				
235	240	245		
gac gac cgc tcc gac atc agc tcc acg tgc gcc aac aac atc aac cac				941
Asp Asp Arg Ser Asp Ile Ser Ser Thr Cys Ala Asn Asn Ile Asn His				
250	255	260		
aac acg gag ctg ctg ccc atc gag ctg gac acg ctg gtg ggg aag ggc				989
Asn Thr Glu Leu Leu Pro Ile Glu Leu Asp Thr Leu Val Gly Lys Gly				
265	270	275		
cgc ttc gcc gag gtc tac aag gcc aag ctg aag cag aac acc tca gag				1037
Arg Phe Ala Glu Val Tyr Lys Ala Lys Leu Lys Gln Asn Thr Ser Glu				
280	285	290		
cag ttt gag acc gtg gct gtc aag atc ttc ccc tac gag gag tac tcc				1085
Gln Phe Glu Thr Val Ala Val Lys Ile Phe Pro Tyr Glu Glu Tyr Ser				
295	300	305	310	
tcg tgg aaa aca gag aag gac atc ttc tcc gat atc aac ctg aag cat				1133
Ser Trp Lys Thr Glu Lys Asp Ile Phe Ser Asp Ile Asn Leu Lys His				
315	320	325		
gag aac atc ctg cag ttc ctg acg gcc gag gag cggt aag aca gag ctg				1181

Glu Asn Ile Leu Gln Phe Leu Thr Ala Glu Glu Arg Lys Thr Glu Leu			
330	335	340	
ggc aag cag tac tgg ctg atc acg gcg ttc cac gcg aag ggc aac ctg			1229
Gly Lys Gln Tyr Trp Leu Ile Thr Ala Phe His Ala Lys Gly Asn Leu			
345	350	355	
cag gag tac ctc acg agg cat gtc atc agc tgg gag gac ctg agg aag			1277
Gln Glu Tyr Leu Thr Arg His Val Ile Ser Trp Glu Asp Leu Arg Lys			
360	365	370	
ctg ggc agc tcc ctg gcc cgg ggc atc gct cat ctc cac agt gac cac			1325
Leu Gly Ser Ser Leu Ala Arg Gly Ile Ala His Leu His Ser Asp His			
375	380	385	390
act cct tgt ggg agg ccc aag atg ccc att gtt cac agg gac ctc aag			1373
Thr Pro Cys Gly Arg Pro Lys Met Pro Ile Val His Arg Asp Leu Lys			
395	400	405	
agc tct aac atc cta gtg aag aac gac ttg acc tgt tgc ctg tgt gac			1421
Ser Ser Asn Ile Leu Val Lys Asn Asp Leu Thr Cys Cys Leu Cys Asp			
410	415	420	
ttc ggg ctg tcc ttg cgc ctg gac cct act ctg tct gtg gat gac ctg			1469
Phe Gly Leu Ser Leu Arg Leu Asp Pro Thr Leu Ser Val Asp Asp Leu			
425	430	435	
gcc aac agc ggg cag gtg gga acg gca aga tac atg gcc ccg gaa gtt			1517
Ala Asn Ser Gly Gln Val Gly Thr Ala Arg Tyr Met Ala Pro Glu Val			
440	445	450	
cta gaa tcc agg atg aat ctg gaa aac gtg gag tcg ttc aag cag acg			1565
Leu Glu Ser Arg Met Asn Leu Glu Asn Val Glu Ser Phe Lys Gln Thr			
455	460	465	470
gat gtc tac tcc atg gct ctg gta ctc tgg gaa atg acg tcc cgc tgc			1613
Asp Val Tyr Ser Met Ala Leu Val Leu Trp Glu Met Thr Ser Arg Cys			
475	480	485	
aat gct gtg gga gaa gtg aag gat tac gag ccc cca ttt ggt tcc aag			1661
Asn Ala Val Gly Glu Val Lys Asp Tyr Glu Pro Pro Phe Gly Ser Lys			
490	495	500	
gtg cgg gag cac ccc tgt gtg gag agc atg aaa gac agt gtg ctg aga			1709
Val Arg Glu His Pro Cys Val Glu Ser Met Lys Asp Ser Val Leu Arg			
505	510	515	
gac cga ggg cgg ccg gaa att ccc agc ttc tgg ctc aac cac cag ggc			1757
Asp Arg Gly Arg Pro Glu Ile Pro Ser Phe Trp Leu Asn His Gln Gly			
520	525	530	
atc cag atc gtg tgt gag act ttg acc gag tgc tgg gac cat gac ccc			1805
Ile Gln Ile Val Cys Glu Thr Leu Thr Glu Cys Trp Asp His Asp Pro			
535	540	545	550
gaa gcc cgt ctc aca gca cag tgt gtg gca gag cgc ttc agt gag ctg			1853
Glu Ala Arg Leu Thr Ala Gln Cys Val Ala Glu Arg Phe Ser Glu Leu			
555	560	565	
gag cat ccg gag aga ctc tct ggg agg agc tgc tcc cag gag aag att			1901
Glu His Pro Glu Arg Leu Ser Gly Arg Ser Cys Ser Gln Glu Lys Ile			
570	575	580	
cca gaa gat ggc tgc ctg aac act acc aaa tag cttttctgg gcaggctggg			1954

Pro Glu Asp Gly Ser Leu Asn Thr Thr Lys  
 585 590

ccaagcctcc agaagccgtc ctctagccaa agaccagagg cagcaggatt ctctcctgac 2014  
 ttagtgcgttctt ggaaaaccaa ggacttgctc ccttcttccc caggagctgc cccgtgttta 2074  
 gaagcggcag cagcagcagc aacaaccata gcggcggtgg cagcggcggg ggtatgagtga 2134  
 cagagagcgt cctatgcctt ggagactgtc atggcataag ctgtgcttagc acctcctcag 2194  
 gaaatgagat tgatttttac aacagccaaat aacgtttgca ctttattaaat gctgtgtgt 2254  
 aaatacgaat agctatgttt tatataatatac tatataatcta tatgtctata tctctctata 2314  
 tatagccata ctctgcaagg agaccgggtgt tttcttttg gtaacttgaa cgcataggga 2374  
 gcttcttcc gggctctctgg attgtgtgtt tctaccccttag ggtatctcat caacactctc 2434  
 cttcctaaca gttgctaagg agtacttgaa gtttgcgtcgt ggacacactt ggttctcttt 2494  
 ggagccatgg ctgctctgtt gctagtcctat tcactgtctg gctcagaaac aatactgaca 2554  
 agatgaatga attataatca cccccccttag tttattttaa ttttatctta ttttccttca 2614  
 caaccgtatt ctcagctttt gtctgtttta acctgataaa ttattctgaa ctgaaatcca 2674  
 agagacactt tgctgggctc caagagagac tgcacggaaa ggactaagag aaaactctta 2734  
 ttactttcaa actgttatgc gagggataga tgccgttcc tccatctcca cattaaaagg 2794  
 gagtgaactg cccgtggcca tagaaccata catgcagtgg tctctggact cctacaaagc 2854  
 aaactccttc tatcccgcca ggtgtgaggt tctgtgctta agcaccagac acgcccataac 2914  
 cacaagaact tgttctcctg ctgtaaacat gatcttctct ttcttccttc aggctccctc 2974  
 ttcagcttgc ctttctctct ctctctctgc cagcttgcctc tgccccccaca aagactctga 3034  
 tttctgtatgt ggttaagct gctgttttag attctggctg caaccactta tatctaccaa 3094  
 tgtggctcct ctctcaccca gaaagacact aaatacaaag gctgtgtttt ttatttcctt 3154  
 agacatcctt cacagaccac agtgtggct ataagctttt tccattatgc aaggttctta 3214  
 ttttcaaga gcaactatgc ttcaagtcct acctgctgtt ttgtatattt taaatctcct 3274  
 ttgattgtta agtgttgaag gcatgcacaa ctttgcattgg agcagtaatt gtgcatacat 3334  
 agtatcagct gaaagtgaca tggaaatttc cagaatagca ttcacagctg tgagagttga 3394  
 agcctgtggg gctatatgac cagagacaga gaattagaaa atcttggttt atactgtct 3454  
 catctgaatg aacaatctta aaccaatggc tcagagagtt gagctgccaa gctcaattca 3514  
 gagagatact atctttttt tttcccttct tcttctgttc aaacttcgac aagctgtaga 3574  
 caatctgcta atcttttagtc atatagaaac ataaatctct catgtgagaa gaataaaata 3634  
 cgagaacaat tcaaattaga tccatgtgtt aaatccctct ggcccactta cataggcaac 3694  
 ttcccttaca cattttgggg aacagtctgt tagaatattt actcaactga agtctatttc 3754  
 tagtataatt tctctatata tcccaacctg gggagatacg caagagtctg atcctgaata 3814

11

ataccacaca gaaaaaggcg gccgc

3839

<210> 11  
<211> 17  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; PCR Primer

<400> 11  
cgaccgctcc gacatca

17

<210> 12  
<211> 17  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; PCR Primer

<400> 12  
tcgatggca gcagctc

17

<210> 13  
<211> 23  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; PCR Probe

<400> 13  
ctccacgtgc gccaacaaca tca

23

<210> 14  
<211> 20  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; PCR Primer

<400> 14  
ggcaaattca acggcacagt

20

<210> 15  
<211> 20  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; PCR Primer

<400> 15  
 gggtctcgct cctggaagat 20

<210> 16  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Probe

<400> 16  
 aaggccgaga atgggaagct tgtcatc 27

<210> 17  
 <211> 2090  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (336) ... (2039)

<400> 17  
 gttggcgagg agtttccgtt ttcccccga gcgctgagtt gaagttgagt gagtcactcg 60  
 cgcgcacgga gcgacgacac ccccgcggt gcacccgctc gggacaggag ccggactcct 120  
 gtgcagcttc cctcggccgc cgggggcctc cccgcgcctc gccggcctcc agggccctcc 180  
 tggctggcga gcgggggcca catctggccc gcacatctgc gctgccggcc cggcgcgcccc 240  
 tccggagagg gcgcggcgcg gagcgagcc aggggtccgg gaaggcgccg tccgtgcgt 300  
 gggggctcgg tctatgacga gcagcgggtt ctgcc atg ggt cgg ggg ctg ctc 353  
 Met Gly Arg Gly Leu Leu  
 1 5

agg ggc ctg tgg ccg ctg cac atc gtc ctg tgg acg cgt atc gcc agc 401  
 Arg Gly Leu Trp Pro Leu His Ile Val Leu Trp Thr Arg Ile Ala Ser  
 10 15 20

acg atc cca ccg cac gtt cag aag tcg gtt aat aac gac atg ata gtc 449  
 Thr Ile Pro Pro His Val Gln Lys Ser Val Asn Asn Asp Met Ile Val  
 25 30 35

act gac aac aac ggt gca gtc aag ttt cca caa ctg tgt aaa ttt tgt 497  
 Thr Asp Asn Asn Gly Ala Val Lys Phe Pro Gln Leu Cys Lys Phe Cys  
 40 45 50

gat gtg aga ttt tcc acc tgt gac aac cag aaa tcc tgc atg agc aac 545  
 Asp Val Arg Phe Ser Thr Cys Asp Asn Gln Lys Ser Cys Met Ser Asn  
 55 60 65 70

tgc agc atc acc tcc atc tgt gag aag cca cag gaa gtc tgt gtg gct 593  
 Cys Ser Ile Thr Ser Ile Cys Glu Lys Pro Gln Glu Val Cys Val Ala  
 75 80 85

gta tgg aga aag aat gac gag aac ata aca cta gag aca gtt tgc cat 641  
 Val Trp Arg Lys Asn Asp Glu Asn Ile Thr Leu Glu Thr Val Cys His  
 90 95 100

gac ccc aag ctc ccc tac cat gac ttt att ctg gaa gat got gct tct	689		
Asp Pro Lys Leu Pro Tyr His Asp Phe Ile Leu Glu Asp Ala Ala Ser			
105	110	115	
cca aag tgc att atg aag gaa aaa aaa aag cct ggt gag act ttc ttc	737		
Pro Lys Cys Ile Met Lys Glu Lys Lys Pro Gly Glu Thr Phe Phe			
120	125	130	
atg tgt tcc tgt agc tct gat gag tgc aat gac aac atc atc ttc tca	785		
Met Cys Ser Cys Ser Asp Glu Cys Asn Asp Asn Ile Ile Phe Ser			
135	140	145	150
gaa gaa tat aac acc agc aat cct gac ttg ttg cta gtc ata ttt caa	833		
Glu Glu Tyr Asn Thr Ser Asn Pro Asp Leu Leu Val Ile Phe Gln			
155	160	165	
gtg aca ggc atc agc ctc ctg cca cca ctg gga gtt gcc ata tct gtc	881		
Val Thr Gly Ile Ser Leu Leu Pro Pro Leu Gly Val Ala Ile Ser Val			
170	175	180	
atc atc atc ttc tac tgc tac cgc gtt aac cgg cag cag aag ctg agt	929		
Ile Ile Ile Phe Tyr Cys Tyr Arg Val Asn Arg Gln Gln Lys Leu Ser			
185	190	195	
tca acc tgg gaa acc ggc aag acg cgg aag ctc atg gag ttc agc gag	977		
Ser Thr Trp Glu Thr Gly Lys Thr Arg Lys Leu Met Glu Phe Ser Glu			
200	205	210	
cac tgt gcc atc atc ctg gaa gat gac cgc tct gac atc agc tcc acg	1025		
His Cys Ala Ile Ile Leu Glu Asp Asp Arg Ser Asp Ile Ser Ser Thr			
215	220	225	230
tgt gcc aac aac atc aac cac aca gag ctg ctg ccc att gag ctg	1073		
Cys Ala Asn Asn Ile Asn His Asn Thr Glu Leu Leu Pro Ile Glu Leu			
235	240	245	
gac acc ctg gtg ggg aaa ggt cgc ttt gct gag gtc tat aag gcc aag	1121		
Asp Thr Leu Val Gly Lys Gly Arg Phe Ala Glu Val Tyr Lys Ala Lys			
250	255	260	
ctg aag cag aac act tca gag cag ttt gag aca gtg gca gtc aag atc	1169		
Leu Lys Gln Asn Thr Ser Glu Gln Phe Glu Thr Val Ala Val Lys Ile			
265	270	275	
ttt ccc tat gag gag tat gcc tct tgg aag aca gag aag gac atc ttc	1217		
Phe Pro Tyr Glu Glu Tyr Ala Ser Trp Lys Thr Glu Lys Asp Ile Phe			
280	285	290	
tca gac atc aat ctg aag cat gag aac ata ctc cag ttc ctg acg gct	1265		
Ser Asp Ile Asn Leu Lys His Glu Asn Ile Leu Gln Phe Leu Thr Ala			
295	300	305	310
gag gag cgg aag acg gag ttg ggg aaa caa tac tgg ctg atc acc gcc	1313		
Glu Glu Arg Lys Thr Glu Leu Gly Lys Gln Tyr Trp Leu Ile Thr Ala			
315	320	325	
ttc cac gcc aag ggc aac cta cag gag tac ctg acg cgg cat gtc atc	1361		
Phe His Ala Lys Gly Asn Leu Gln Glu Tyr Leu Thr Arg His Val Ile			
330	335	340	
agc tgg gag gac ctg cgc aag ctg ggc agc tcc ctc gcc cgg ggg att	1409		
Ser Trp Glu Asp Leu Arg Lys Leu Gly Ser Ser Leu Ala Arg Gly Ile			
345	350	355	

gct cac ctc cac agt gat cac act cca tgt ggg agg ccc aag atg ccc Ala His Leu His Ser Asp His Thr Pro Cys Gly Arg Pro Lys Met Pro 360 365 370	1457
atc gtg cac agg gac ctc aag agc tcc aat atc ctc gtg aag aac gac Ile Val His Arg Asp Leu Lys Ser Ser Asn Ile Leu Val Lys Asn Asp 375 380 385 390	1505
cta acc tgc tgc ctg tgt gac ttt ggg ctt tcc ctg cgt ctg gac cct Leu Thr Cys Cys Leu Cys Asp Phe Gly Leu Ser Leu Arg Leu Asp Pro 395 400 405	1553
act ctg tct gtg gat gac ctg gct aac agt ggg cag gtg gga act gca Thr Leu Ser Val Asp Asp Leu Ala Asn Ser Gly Gln Val Gly Thr Ala 410 415 420	1601
aga tac atg gct cca gaa gtc cta gaa tcc agg atg aat ttg gag aat Arg Tyr Met Ala Pro Glu Val Leu Glu Ser Arg Met Asn Leu Glu Asn 425 430 435	1649
gct gag tcc ttc aag cag acc gat gtc tac tcc atg gct ctg gtg ctc Ala Glu Ser Phe Lys Gln Thr Asp Val Tyr Ser Met Ala Leu Val Leu 440 445 450	1697
tgg gaa atg aca tct cgc tgt aat gca gtg gga gaa gta aaa gat tat Trp Glu Met Thr Ser Arg Cys Asn Ala Val Gly Glu Val Lys Asp Tyr 455 460 465 470	1745
gag cct cca ttt ggt tcc aag gtg cg <sup>g</sup> gag cac ccc tgt gtc gaa agc Glu Pro Pro Phe Gly Ser Lys Val Arg Glu His Pro Cys Val Glu Ser 475 480 485	1793
atg aag gac aac gtg ttg aga gat cga ggg cga cca gaa att ccc agc Met Lys Asp Asn Val Leu Arg Asp Arg Gly Arg Pro Glu Ile Pro Ser 490 495 500	1841
ttc tgg ctc aac cac cag ggc atc cag atg gtg tgt gag acg ttg act Phe Trp Leu Asn His Gln Gly Ile Gln Met Val Cys Glu Thr Leu Thr 505 510 515	1889
gag tgc tgg gac cac gac cca gag gcc cgt ctc aca gcc cag tgt gtg Glu Cys Trp Asp His Asp Pro Glu Ala Arg Leu Thr Ala Gln Cys Val 520 525 530	1937
gca gaa cgc ttc agt gag ctg gag cat ctg gac agg ctc tcg ggg agg Ala Glu Arg Phe Ser Glu Leu Glu His Leu Asp Arg Leu Ser Gly Arg 535 540 545 550	1985
agc tgc tcg gag gag aag att cct gaa gac ggc tcc cta aac act acc Ser Cys Ser Glu Glu Lys Ile Pro Glu Asp Gly Ser Leu Asn Thr Thr 555 560 565	2033
aaa tag ctcttatggg gcaggctggg catgtccaaa gaggctgccc ctctcaccaa Lys	2089
a	2090

<210> 18  
 <211> 1225  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (917) ... (1225)

<400> 18 aactacaaat tcacatctcc tgtgtgagcc atgggtgagt tgagtagaaat ggtggtagaa ttatatgggt tacattaact acttagactt tgtcctattt ttcctatatg aaaatagttg aaattctcat aaaaatgcat gtagatgtg aaactgaagt ataatgtaca gtcagataat tttgaatata ttttgtgaaa tagatgtta tgatttataa cagcagtatt ttaatttcta tgcagcttaa attgatgctt ggatagggaa aaaggatgaa gaagaaactc aatggatgt taactgagag aatgaagtaa agtggcatag taacgacagt tataaaaggg gcactgcata ccccatgtgtcaaggat tgtaaacttc cagactggc taaacgaatg ctggaaagg tctgtacagc cacaacaagt ccttagactt aggacttaga agcttattt cagtagttat ttttaggat acaggattct gggactaat tcatgttaac tgaatttaca agacatataat atattttagtt gagacattta attaccaaga taactagaaa tctcatcagt aggaaggct ccttggctgt tttgttgcta tgtttatgaa gatggttca tgaacatgga gggagaagcc gtaaaatggt tttaactaat gaccaggcat gatagcttat tccttggaa agccaaggcg agaggatccc ttgagcctt agcctaggaa tttgagacca gcctggcaa cataataaga ccccatctgt acaaaacatt aaaagttacc tggcataac tttcctataat atcctataat tcctataatc ccagctactt gggaggctga ggtgggagga ttgcttgagc ctggaggtc gaggatgcag tgagcg atg ata cta cca ctg cat aac ctg ggc aat gga gtg Met Ile Leu Pro Leu His Asn Leu Gly Asn Gly Val 1 5 10	60 120 180 240 300 360 420 480 540 600 660 720 780 840 900 952 1000 1048 1096 1144 1192 1225
agg tcc cat aac ttt tct ttt ctt tac ttt att ctg gaa gat gct gct Arg Ser His Asn Phe Ser Phe Leu Tyr Phe Ile Leu Glu Asp Ala Ala 15 20 25	1000
tct cca aag tgc att atg aag gaa aaa aaa aag cct ggt gag act ttc Ser Pro Lys Cys Ile Met Lys Glu Lys Lys Lys Pro Gly Glu Thr Phe 30 35 40	1048
ttc atg tgt tcc tgt agc tct gat gag tgc aat gac aac atc atc ttc Phe Met Cys Ser Cys Ser Asp Glu Cys Asn Asp Asn Ile Ile Phe 45 50 55 60	1096
tca gaa gaa tat aac acc agc aat cct gac ttg ttg cta gtc ata ttt Ser Glu Glu Tyr Asn Thr Ser Asn Pro Asp Leu Leu Leu Val Ile Phe 65 70 75	1144
caa gtg aca ggc atc agc ctc ctg cca cca ctg gga gtt gcc ata tct Gln Val Thr Gly Ile Ser Leu Leu Pro Pro Leu Gly Val Ala Ile Ser	1192
80 85 90	
gtc atc atc atc ttc tac tgc tac cgc gtt aac Val Ile Ile Ile Phe Tyr Cys Tyr Arg Val Asn 95 100	1225

<210> 19  
<211> 425  
<212> DNA  
<213> Homo sapiens

<220>

<221> unsure  
<222> 6  
<223> unknown

<221> unsure  
<222> 29  
<223> unknown

<221> unsure  
<222> 42  
<223> unknown

<221> unsure  
<222> 89  
<223> unknown

<221> unsure  
<222> 204  
<223> unknown

<223>

<400> 19  
tcggcnccga aggaaaaaaag accaaggant aacattctgt antgcctaaa aatactgact 60  
tttttcacta ctatacatga aggaaaagnt ttattctttt atggaacact tcagctgtac 120  
tcatgtatta aaataggaat gtgaatgcta tatactcttt ttatatcaaa agtctcaagc 180  
acttattttt attctatgca ttgnttgcct 360  
tttacataaaa taaaatgttt attagattga  
ataaaagcaaa atactcaggt gagcatcctg cctcctgttc ccattcctag tagctaaatc  
cattgcctc atttggttat tttgcaattt atgcagaaaa cgtcaccaag taaataatct  
ccaacttttt catactgtct ttctcaactg acagtctcac agtccttaaa aaaaaaaaaa 420  
aaaaaa 425

<210> 20  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 20  
cagccccccga cccatggcag 20

<210> 21  
<211> 20

17

<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 21  
ctgcagttgc tcatgcagga 20

<210> 22  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 22  
ttgtgggtga tgttgtggc 20

<210> 23  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 23  
ttctgcttca gcttggcctt 20

<210> 24  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 24  
aggcctcccc agctgatgac 20

<210> 25  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 25  
ggcatcttgg gcctccacaca 20

<210> 26

<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 26  
catccacaga cagagtaggg 20

<210> 27  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 27  
gccaggtcat ccacagacag 20

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 28  
agagccatgg agtagacatc 20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 29  
tcccgcacct tggaaccaaa 20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 30  
tggttgagcc agaagctggg 20

19

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 31  
gatgccctgg tggttgagcc 20  
  
<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 32  
ctcactcaac ttcaactcag 20  
  
<210> 33  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 33  
cctgtccccga gcgggtgcac 20  
  
<210> 34  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 34  
gcggccgagg gaagctgcac 20  
  
<210> 35  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 35  
atatgaaagg tgtattacaa 20

20

<210> 36  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 36  
aaaaaacttgc atttcaacat

20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 37  
tctacacact tagttaagga

20

<210> 38  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 38  
cttagggttc catgtcttcc

20

<210> 39  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 39  
ataaaagactg taatcagtgat

20

<210> 40  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 40  
gtttacaaga acagtccaaa

20

21

<210> 41  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 41  
aagatgcata tacctgttta 20

<210> 42  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 42  
catcccaagtg aggctttct 20

<210> 43  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 43  
gagtcctagac atattggtgg 20

<210> 44  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 44  
tgttagcctc tgagtctaga 20

<210> 45  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 45  
actagcagtc atgttagcct 20

<210> 46  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 46  
agctttgtga cttttaaagg 20

<210> 47  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 47  
cagaccccgac tgctcgatcat 20

<210> 48  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 48  
agccccccgac ccatggcaga 20

<210> 49  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 49  
gtcgttatta accgacttct 20

<210> 50  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 50  
gtcacaggtg gaaaatctca 20

<210> 51  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 51  
aagaggcata ctcctcatag 20

<210> 52  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 52  
attgatgtct gagaagatgt 20

<210> 53  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 53  
ggaaagccca aagtcacaca 20

<210> 54  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 54  
ctcagcattc tccaaattca 20

<210> 55  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 55  
ttacagcgag atgtcatttc 20

<210> 56  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 56  
gtcgccctcg atctctcaac 20

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 57  
gccgtcttca ggaatcttct 20

<210> 58  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 58  
cagaagagct atttggttagt 20

<210> 59  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 59  
agcttatacct atgacaatgt 20

<210> 60  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 60  
caatctcatt tcctgaggaa 20

25

<210> 61  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 61  
aggtatggct atatatata 20  
tag

<210> 62  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 62  
agttctccaa taaaaccaat 20  
at

<210> 63  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 63  
ctgcatgtgt gattgtcaaa 20  
ttt

<210> 64  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 64  
tttgcaaaag caagtgc 20  
aat

<210> 65  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 65  
taagggcaca caggaacccc 20  
cc

<210> 66  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 66  
caggagaaaat aagggcacac 20

<210> 67  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 67  
gtagagtttc taaaacttaggt 20

<210> 68  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 68  
tgcaaccat gaaggtaaaa 20

<210> 69  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 69  
aaaggatgaa ggctgggagc 20

<210> 70  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 70  
ccattgggtgt ttgtatagaa 20

<210> 71  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 71  
caggagccca gaaagatgga 20

<210> 72  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 72  
gagcaatcag gagcccgagaa 20

<210> 73  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 73  
aaagtggctt cacttttga 20

<210> 74  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 74  
ggactggaga taactgaaaa 20

<210> 75  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 75  
gtctccacac cttcacat 20

<210> 76  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 76  
ccaggttaggc agtggaaaga

20

<210> 77  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 77  
cttaaaggag ttcccccttta

20

<210> 78  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 78  
ttctgcaaag gttgagaagg

20

<210> 79  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 79  
ttgtggacac aaattttctta

20

<210> 80  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 80  
acacttcttt gttgatttaac

20

<210> 81  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 81  
ccatccaaac agagctgata

20

<210> 82  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 82  
ttgaatatct catgaatgga

20

<210> 83  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 83  
cttgaatatac tcatgaatgg

20

<210> 84  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 84  
taaaaataaag tgcttgagac

20

<210> 85  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 85  
aaacatttta tttatgtaaa

20

30

<210> 86  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 86  
actatgccac tttacttcat

20

<210> 87  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 87  
gggtatgcag tgcccccttt

20

<210> 88  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 88  
gtggctgtac agacctttcc

20

<210> 89  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 89  
attagtcccc agaattcctgt

20

<210> 90  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 90  
acatgaattt gtcccccagaa

20

<210> 91  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 91  
atatgtcttg taaaattcagt 20

<210> 92  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 92  
ccttcctact gatgagattt 20

<210> 93  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 93  
ggtcattagt taaaaccatt 20

<210> 94  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 94  
tctcaaattc ctaggctcaa 20

<210> 95  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 95  
acctcagcct cccaaatgc 20

<210> 96  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 96  
aaatggattt agctactagg

20

<210> 97  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 97  
caaatggattt tagctactag

20

<210> 98  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 98  
ctcgagggcc cgaggaggc

20

<210> 99  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 99  
gatccccgga gctcggaggc

20

<210> 100  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 100  
ggccagatgt ggccggcgat

20

<210> 101  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 101  
ggaccccgaaa ctgcgccttt 20

<210> 102  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 102  
cctgggtgcgc cacgaaccga 20

<210> 103  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 103  
gccccccgtcg ctcgtcatag 20

<210> 104  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 104  
cgagcagcc cccgaccat 20

<210> 105  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 105  
ccacaggccc cgagcagcc 20

<210> 106  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 106  
gatgcgcgtc cacaggacga 20

<210> 107  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 107  
tcgtgctggc gatgcgcgtc 20

<210> 108  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 108  
tgccgcggga tcgtgctggc 20

<210> 109  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 109  
ggatggtcct attacagctt 20

<210> 110  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 110  
acttgcacag ctgtggaagc 20

<210> 111  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 111  
tcatgcagga cttctggttg 20

<210> 112  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
,

<220>

<223> Antisense Oligonucleotide

<400> 112  
ttccagagtg aagccgtggt 20

<210> 113  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 113  
cgcccttttc ttttccttca 20

<210> 114  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 114  
gtctcgcccg ccctttctt 20

<210> 115  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 115  
ggactgctgg tggtgtattc 20

<210> 116  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 116  
agcggaggca ggaggctgac 20

<210> 117  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 117  
gctatggcaa tccccagcgg 20

<210> 118  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 118  
tgacagctat ggcaatcccc 20

<210> 119  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 119  
ctcagttct gctgccggtg 20

<210> 120  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 120  
ggtcgtcctc caggatgatg 20

<210> 121  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 121  
cgcacgtgga gctgatgtcg 20

<210> 122  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 122  
ttgatgttgt tggcgacgt 20

<210> 123  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 123  
ccgtgttgtg gttgatgttg 20

<210> 124  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 124  
cgatggcag cagctccgtg 20

<210> 125  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 125  
tccagctcga tggcagcag 20

<210> 126  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 126  
gccttgtaga cctcggcgaa 20

<210> 127  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 127  
tgacagccac ggtctcaaac 20

<210> 128  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 128  
aggaactgca ggtatgttctc 20

<210> 129  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 129  
ctcgccgtc aggaactgca 20

<210> 130  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 130  
ccgctcctcg gccgtcagga 20

<210> 131  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 131  
gtgatcagcc agtactgctt 20

<210> 132  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 132  
ggttgccctt cgcgtgaaac 20

<210> 133  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 133  
tactcctgca ggttgccctt 20

<210> 134  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 134  
ctgatgacat gcctcgtgag 20

<210> 135  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 135  
ggccagggag ctgcccagct 20

40

<210> 136  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 136  
aggtccctgt gaacaatggg

20

<210> 137  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 137  
tgttagagct cttgaggtcc

20

<210> 138  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 138  
tcgttttca ctaggatgtt

20

<210> 139  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 139  
ccaggcgcaa ggacagcccg

20

<210> 140  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 140  
ttcatcctgg attctagaac

20

41

<210> 141  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 141  
ggagtagaca tccgtatgct 20

<210> 142  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 142  
cagcgggacg tcatttccca 20

<210> 143  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 143  
ttcatgctct ccacacaggg 20

<210> 144  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 144  
tttccggcccg ccctcggtct 20

<210> 145  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 145  
agctggaaat ttccggccgc 20

42

<210> 146  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 146  
cacacgatct ggatgccctg

20

<210> 147  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 147  
tggtcccagc actcggtcaa

20

<210> 148  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 148  
cttctcctgg gagcagctcc

20

<210> 149  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 149  
agaaaaagct atttggtagt

20

<210> 150  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 150  
cccagcctgc ccagaaaaag

20

<210> 151  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 151  
ggaagaagggg agcaagtccct 20

<210> 152  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 152  
ctatggttgt tgctgctgct 20

<210> 153  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 153  
tgccatgaca gtctccaagg 20

<210> 154  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 154  
tataaaacat agctattcgt 20

<210> 155  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 155  
taggaaggag agtgttgatg 20

44

<210> 156  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 156  
ttgtttctga gccagacagt 20

<210> 157  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 157  
cagttcactc ccttttaatg 20

<210> 158  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 158  
acagaacctc acacctggcgt 20

<210> 159  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 159  
agagagagaa aggcaagctg 20

<210> 160  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 160  
cttaaaccac atcagaaatc 20

45

<210> 161  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 161  
tatttagtgt ctttctgggt

20

<210> 162  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 162  
gattgttcat tcagatgaga

20

<210> 163  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
  
<220>  
  
<223> Antisense Oligonucleotide  
  
<400> 163  
gtgtggatt attcaggatc

20