



(22) Date de dépôt/Filing Date: 1993/01/14

(41) Mise à la disp. pub./Open to Public Insp.: 1993/07/18

(45) Date de délivrance/Issue Date: 2003/09/09

(30) Priorité/Priority: 1992/01/17 (MI 92 A 000082) IT

(51) Cl.Int.⁵/Int.Cl.⁵ C07J 31/00, A61K 31/575


(72) Inventeurs/Inventors:
GIULIANI, GIANGERMANO, IT;
FRIGERIO, GIULIANO, IT;
PELLICCIARI, ROBERTO, IT

(73) Propriétaire/Owner:
GIULIANI S.P.A., IT

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : DERIVES SULFURES D'UN ACIDE BILIAIRE

(54) Title: BILE ACID SULFURATED DERIVATIVES




(57) Abrégé/Abstract:

Compounds of formula I: (see formula I) in which the various R groups have the meanings defined in the disclosure, are useful as medicaments for the treatment of hepatobiliary diseases.



AbstractBILE ACID SULFURATED DERIVATIVES

5 Compounds of formula I:




15 in which the various R groups have the meanings defined
in the disclosure, are useful as medicaments for the
treatment of hepatobiliary diseases.

BILE ACID SULFURATED DERIVATIVES

The present invention relates to bile acid derivatives, a process for the preparation thereof and pharmaceutical compositions containing them.

5 The derivatives of the invention have the following general formula I:



wherein:

R is OH or SH in α - or β - position;

10 R^I and R^{II} are hydrogen, or one is hydrogen and the other is OH or SH;

R^{III} and R^{IV} are hydrogen or one is hydrogen and the other is OH or SH;

R^V is hydrogen or methyl;

15 with the proviso that at least one of R, R^I , R^{II} , R^{III} and R^{IV} is a SH group,

the glyco- or tauro-conjugated derivatives thereof and the pharmaceutically acceptable salts thereof with the exception of the compounds 3 β -mercapto-5 β -cholan-24-oic acid and 12 α -hydroxy-3 β -mercapto-5 β -cholan-24-oic acid.

20 Compounds I can be considered as the thio-derivatives of the natural bile acids ursodeoxycholic (UDCA) (3 α , 7 β OH),ursocholic (3 α , 7 β , 12 α OH), chenodeoxycholic (3 α , 7 α OH), lithocholic (3 α OH), deoxycholic

(3 α , 12 α OH) and cholic (3 α , 7 α , 12 α OH) and of the corresponding 23-methyl derivatives described in EP 135782.

Particularly preferred are the compounds I in which only one of the R, R^I, R^{II}, R^{III} and R^{IV} groups is SH.

A further preferred group includes compounds wherein either R or R_I is SH and the other groups are different from SH.

The compounds of formula I are useful in therapy as anticholestatic, choleretic, antidyslipemic, hepatocyte-protecting agents as well as for the up to now known traditional uses of the bile acids (treatment of cholelithiasis, bile desaturation, cholesterol metabolic, etc.). Compared with the natural bile acids and the up to now known derivatives thereof, the compounds of the invention show an increased metabolic stability as well as antioxidant and radical scavenging activities, the latter being particularly desirable for the treatment of hepatobiliary pathologies in view of the cyto-protecting effect from exogenous or physiological events (cell aging, toxic metabolites).

The compounds of the invention, for the envisaged therapeutical uses, are administered in form of pharmaceutical compositions prepared according to conventional techniques and excipients, as described for example in "Remington's Pharmaceutical Sciences Handbook", Mack Pub., N.Y. USA.

The preferred administration route is the oral one, whereas the doses, which will vary depending on the pathology to treat and the patient's conditions, in

principle will range from 0.5 to 5 g, one or more times daily.

Examples of suitable pharmaceutical forms comprise capsules, tablets, sugar-coated pills, syrups, granulates, solutions, vials. The compounds of the invention
5 can also be used for local perfusion before or after of surgery, in form of solutions or dispersible powders.

The compounds of formula I can be prepared starting from the natural bile acids or the derivatives described in EP 135782, by selective substitution of a OH
10 group with a SH group, according to conventional procedures. For example, the OH group to be replaced will be transformed into a good leaving group, such as a mesyl or tosyl group and then reacted with a sulhydrylation
15 agent such as thiourea, KSH or equivalent reagents, whereas the remaining OH groups will be protected by means of inert groups, under the used reaction conditions.

The substitution reaction can occur with inversion
20 or retention of the α or β stereochemical configuration.

The known difference in chemical reactivity of the OH groups at the 3, 7 and 12 -positions of the bile acids allows to prepare the desired derivatives of formula I by means of suitable schemes of protection/deprotection known to those skilled in the art.
25

The following example further illustrates the invention.

EXAMPLE

30 a) methyl 3 α -7 β -diacetoxy-12 α -hydroxy-5 β -cholanoate

A solution of methyl 3 α ,7 β ,12 α -triidroxy-5 β -chola-

noate (10 g, 23.66 mmol) in benzene (50 ml), pyridine (12.5 ml) and acetic anhydride (12.5 ml) was kept under magnetic stirring in the dark and at room temperature, for 24 hours. The reaction mixture was washed with water (4 x 40 ml), then with 10% hydrochloric acid to acid pH, then again with water (2 x 40 ml). The organic phase was evaporated and the residue (10.5 g) was subjected to flash chromatography (SiO_2 , $h = 18$ cm, $\phi = 1.7$ cm). Eluting first with petroleum ether/ethyl ether 97/3 v/v, then with petroleum ether/ethyl ether 94/6 v/v, the acetylated product is obtained (9.30 g, 77% yield).

NMR (CDCl_3): $\delta = 0.71$ (3H, s, C-18 Me); 0.96 (3H, s, C-19 Me); 1.98 (3H, s, C-3 OCOCH_3); 2.01 (3H, s, C-7 OCOCH_3); 3.66 (3H, s, COOMe); 3.90 - 4.06 (1H, m, C-12 CHOH); 4.43 - 4.96 (2H, m, C-3 and C-7 CHOAc).

b) methyl 3 α ,7 β -diacetoxy-12 α -mesyloxy-5 β -cholanoate

A solution of methyl 3 α ,7 β -diacetoxy-12 α -hydroxy-5 β -cholanoate (1.54 g, 2.96 mmol) in anhydrous pyridine (4,40 ml), kept under magnetic stirring in nitrogen atmosphere at room temperature, was added with methanesulfonyl chloride (0.32 ml, 4.11 mmol). After 24 hours the reaction mixture was poured into ice-water (50 ml) and extracted with ether (3 x 40 ml). The combined ether phases were dried over magnesium sulfate, evaporated and washed repeatedly with hexane to remove pyridine. 1.75 g of the mesyl derivative are obtained (quantitative yield) (m.p. 88°-94°C).

IR (CHCl_3): $\nu = 2950$, 2870, (CH_2 , CH_3), 1725 (CO), 1335 (SO_2), 905 (SO) cm^{-1} .

NMR (CDCl_3): $\delta = 0.81$ (3H, s, C-18 Me); 0.96 (3H, s, C-

19 Me); 1.98 (3H, s, C-3 OCOCH₃); 2.01 (3H, s, C-7 OCOCH₃); 3.08 (3H, s, SO₂ Me); 3.65 (3H, s, COOMe); 4.41 - 4.93 (2H, m, C-3 and C-7 CHOAc); 4.98 - 5.15 (1H, m, C-12 CHO Ms).

5 c) 3 α ,7 α -dihydroxy-12 β -mercapto-5 β -cholan-24-oic acid

A mixture of 3 α ,7 α -diacetox-12 α -mesyloxy-5 β -cho-
lan-24-oic acid (1.90 g, 3.50 mmoles) and thiourea
(1.14 g, 3.50 mmoles) in 250 ml of isopropanol was re-
fluxed for 3 hours. Then the reaction mixture was added
10 with a sodium hydroxide solution (30 g) in 300 ml of
water and the mixture was refluxed for 2 hours, then it
was acidified with diluted sulfuric acid and extracted
with several ethyl acetate portions. The combined orga-
nic extracts were washed with water and dried over
15 anhydrous sodium sulfate. Then the solvent was evapora-
ted off to obtain a product (65% yield) which, on the
basis of the analytical and spectroscopic data, is
identified as the 3 α ,7 α -dihydroxy-12 β -mercapto-5 β -cho-
lan-24-oic acid.

20 In a similar way, the following compounds I have
also been prepared:

- 3 β -mercapto-5 β -lithocholan-24-oic acid (Formula I,
R = 3 β SH, R^I = R^{II} = R^{III} = R^{IV} = R^V = H),
obtained from methyl, 3 α -hydroxy-5 β -litho-24-
25 cholanoate through the following steps:

- tosylation of the 3-OH group;
- substitution of the 3 α -tosyloxy group with
potassium thioacetate in DMF and hydrolysis.

M.P. 140-145°C.

30 ¹H-NMR (CDCl₃ + CD₃OD): δ 0.7 (s, 3H, 18-CH₃);
0.8-1.0 (m, 6H, 19-CH₃, 21CH₃); 3.55 (m, 1H, 3-

CH).

^{13}C -NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$): δ 37.1 (3-CH); 176 (24-COOH).

MS-GC: 406 [M-73(Me_3Si -)]

- 5 - 3 α -mercapto-7 α -hydroxy-5 β -chenodesoxycholan-24-oic acid (Formula I $\text{R} = 3\alpha \text{ SH}$, $\text{R}^{\text{I}} = \text{R}^{\text{III}} = \text{R}^{\text{IV}} = \text{R}^{\text{V}} = \text{H}$, $\text{R}^{\text{II}} = \text{OH}$), obtained from methyl 3 α , 7 α -dihydroxy-5 β -chenodesoxy-24-cholanoate through the following steps:
 - 10 - acetylation of the OH groups;
 - selective hydrolysis of the 3 α acetoxy group in a 9% NaOH methanol solution at room temperature for 30 minutes;
 - mesylation of the 3 α OH group;
 - 15 - substitution of the 3 α mesyl group with 3 β OH group with concomitant hydrolysis of the 7 α acetoxy group;
 - acetylation of the 3 β , 7 α hydroxy groups;
 - selective hydrolysis of the 3 β -hydroxy group in a 9% NaOH methanol solution at room temperature for 4 hours;
 - 20 - tosylation of the 3 β OH group;
 - substitution of the 3 β -tosyl group with potassium thioacetate in DMF and hydrolysis.
 - 25 ^1H -NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$): δ 0.7 (1, 3H, 18- CH_3); 0.8-1.0 (m, 6H, 19- CH_3 , 21- CH_3); (m, 1H, 3-CH); 3.9 (m, 1H, 7-CH).
 - ^{13}C -NMR ($\text{CDCl}_3 + \text{CD}_3\text{OD}$): δ (3-CH-SH); 67.6 (7-CH-OH); 177.0 (24-COOH).
 - 30 - 3 β -mercapto-7 α -hydroxy-5 β -chenodesoxycholan-24-oic acid, obtained from methyl 3 α , 7 α -dihydroxy-5 β -

chenodesoxy-24-cholanoate through the following steps:


- acetylation of the OH groups;
- selective hydrolysis of the 3 α acetoxy group as disclosed above;
- tosylation of the 3 α OH group;
- substitution of the 3 α tosyloxy group with potassium thioacetate and hydrolysis.

¹H-NMR (CDCl₃ + CD₃OD): δ 0.7 (1, 3H, 18-CH₃);
0.8-1.0 (m, 6H, 19-CH₃, 21-CH₃); 2.5 (m, 1H, 3-CH); 3.9 (m, 1H, 7-CH).

MS-GC: 461.50 (M-S-Si-CH₃).

CLAIMS

1. Compounds of formula I:



wherein:

R is OH or SH in α - or β - position;

5 R^I and R^{II} are hydrogen, or one is hydrogen and the other is OH or SH;

R^{III} and R^{IV} are hydrogen or one is hydrogen and the other is OH or SH;

R^V is hydrogen or methyl;

10 with the proviso that at least one of R, R^I, R^{II}, R^{III} and R^{IV} is a SH group,

the glyco- or tauro-conjugated derivatives thereof and the pharmaceutically acceptable salts thereof with the exception of the compounds 3 β -mercapto-5 β -cholan-24-oic acid and 12 α -hydroxy-3 β -mercapto-5 β -cholan-24-oic acid.

2. Compounds according to claim 1, in which only one of the R, R^I, R^{II}, R^{III} and R^{IV} groups is SH.

3. A compound according to claim 1, selected from:

3 α , 7 α -dihydroxy-12 β -mercapto-5 β -cholan-24-oic acid;

20 3 α -mercapto-7 α -hydroxy-5 β -chenodesoxycholan-24-oic acid;
and

3 β -mercapto-7 α -hydroxy-5 β -chenodesoxycholan-24-oic acid.

4. Pharmaceutical compositions containing as the active ingredient a compound of claim 1, 2 or 3 in admixture with a pharmaceutically acceptable carrier or diluent.

5. The use of the compounds of claim 1, 2 or 3 and

5 3 β -mercapto-5 β -cholan-24-oic acid and 12 α -hydroxy-3 β -mercapto-5 β -cholan-24-oic acid for the preparation of a medicament for the treatment of hepatobiliary diseases.

6. The use of the compounds of claim 1, 2 or 3 and

3 β -mercapto-5 β -cholan-24-oic acid and 12 α -hydroxy-3 β -
10 mercapto-5 β -cholan-24-oic acid for the treatment of hepatobiliary diseases.

