




(86) Date de dépôt PCT/PCT Filing Date: 2011/09/15
(87) Date publication PCT/PCT Publication Date: 2012/03/22
(45) Date de délivrance/Issue Date: 2020/07/21
(85) Entrée phase nationale/National Entry: 2013/03/11
(86) N° demande PCT/PCT Application No.: US 2011/051827
(87) N° publication PCT/PCT Publication No.: 2012/037397
(30) Priorité/Priority: 2010/09/15 (US61/383,208)

(51) Cl.Int./Int.Cl. *C07K 14/47* (2006.01),
A61K 38/17 (2006.01), *A61P 25/00* (2006.01),
C07K 19/00 (2006.01)
(72) Inventeur/Inventor:
PAUL, SUROJIT, US
(73) Propriétaire/Owner:
STC.UNM, US
(74) Agent: SMART & BIGGAR LLP

(54) Titre : PEPTIDE DERIVE DE LA PROTEINE STEP UTILISE DANS LE TRAITEMENT DES LESIONS CEREBRALES
(54) Title: STEP-DERIVED PEPTIDE FOR BRAIN INJURY TREATMENT



(57) **Abrégé/Abstract:**

A novel peptide sequence that is a modified derivative of a neuron-specific tyrosine phosphatase is shown and described. Specifically, the novel peptide sequence is a modified derivative of striatal-enriched tyrosine phosphatase (STEP). The peptide sequence has been modified so as to be able to ameliorate and treat brain injury resulting from excessive glutamate release and / or oxidative stress. Examples of the types of brain injury which the presently disclosed peptide sequence is useful for treating includes acute brain injury resulting from stroke or traumatic brain injury and chronic disorders such as Huntington's chorea and schizophrenia. Furthermore, the presently described peptide sequence may further be useful in the treatment and amelioration of disorders associated with fear memory such as post-traumatic stress disorder.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2012/037397 A3

(43) International Publication Date
22 March 2012 (22.03.2012)

(51) International Patent Classification:

C07K 14/47 (2006.01) *A61K 38/17* (2006.01)
C07K 19/00 (2006.01) *A61P 25/00* (2006.01)

(21) International Application Number:

PCT/US2011/051827

(22) International Filing Date:

15 September 2011 (15.09.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/383,208 15 September 2010 (15.09.2010) US

(71) Applicant (for all designated States except US):

STC.UNM [US/US]; MSC 04 2750, 801 University Blvd.
SE, Suite 101, Albuquerque, NM 87106 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): PAUL, Surojit
[IN/US]; 8910 Greenarbor Road, NE, Albuquerque, NM
87122 (US).

(74) Agent: GONZALES, Ellen; Gonzales Patent Services,
4605 Congress Ave NW, Albuquerque, NM 87114 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:


— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

28 June 2012

(54) Title: STEP-DERIVED PEPTIDE FOR BRAIN INJURY TREATMENT



(57) Abstract: A novel peptide sequence that is a modified derivative of a neuron-specific tyrosine phosphatase is shown and described. Specifically, the novel peptide sequence is a modified derivative of striatal-enriched tyrosine phosphatase (STEP). The peptide sequence has been modified so as to be able to ameliorate and treat brain injury resulting from excessive glutamate release and / or oxidative stress. Examples of the types of brain injury which the presently disclosed peptide sequence is useful for treating includes acute brain injury resulting from stroke or traumatic brain injury and chronic disorders such as Huntington's chorea and schizophrenia. Furthermore, the presently described peptide sequence may further be useful in the treatment and amelioration of disorders associated with fear memory such as post-traumatic stress disorder.

STEP-Derived Peptide for Brain Injury Treatment

Statement Regarding Government Sponsored Research

[001] This invention was made with Government support under 5R01NS059962-04 awarded by NINDS. The U.S. Government has certain rights in this invention.

Cross-reference to Related Applications

[002] The following application claims benefit of U.S. Provisional Application No. 61/383,208, filed September 15, 2010.

[003]

Background

[004] Brain injury resulting from ischemic insults such as stroke, post-traumatic lesions and a variety of neurological disorders frequently causes significant neuro-degeneration resulting in mild to severe impairment and even death to those suffering from these injuries. Current treatment methods are limited and are typically related to invasive surgical procedures and post-injury rehabilitative treatments rather than limiting or reversing the damage caused by the initial injury to the brain.

[005] Various studies implicate excitotoxic glutamate receptor activation in neuro-degeneration following ischemic insults, post-traumatic lesions and a range of other neurological disorders (Lipton and Rosenberg 1994; Arundine and Tymianski 2004). In pathological conditions excessive release of the neurotransmitter glutamate leads to activation of NMDA-subtype of glutamate receptors that triggers a deleterious cascade of intracellular events including Ca^{2+} overload and activation of effector molecules like p38 mitogen activated protein kinase (p38 MAPK) that is known to be involved in excitotoxic cell death (Barone et al. 2001a; Barone et al. 2001b). Recent findings suggest the existence of two functionally distinct pools of NMDA receptors (NMDAR) defined by the presence of NR2A (NR2A-NMDAR) or NR2B (NR2B-NMDAR) containing subunits (Luo et al. 1997; Tovar and Westbrook 1999; Kohr 2006). NR2B-NMDARs are generally thought to be involved in triggering cell death and subsequent brain damage following an excitotoxic insult (Liu et al.

2007; Tu et al. 2010). However pharmaceutical application of NR2B-NMDAR antagonists have been unsuccessful in clinical trials, since they interfere with physiological functions of these receptors, in addition to blocking excitotoxic cascades (Rodrigues et al. 2001; Zhao et al. 2005; Walker and Davis 2008).

[006] Accordingly, targeting components of the intracellular signaling cascades downstream of NMDAR signaling may provide novel compounds for therapeutic approaches to interventions in excitotoxicity.

[007] The brain-enriched tyrosine phosphatase STEP (also known as STriatal Enriched Phosphatase or PTPN5) may be one such target molecule that is activated following stimulation of NMDARs and is emerging as an important regulator of neuronal survival and death. STEP is expressed specifically in neurons of the striatum, neo-cortex and hippocampus (Boulanger et al. 1995). STEP₆₁ and STEP₄₆, the two STEP isoforms (Bult et al. 1997) contain a highly conserved substrate-binding domain termed as the kinase interacting motif or KIM domain (Pulido et al. 1998). Phosphorylation of a critical serine residue within the KIM domain is mediated through dopamine/D1 receptor dependent activation of the Protein Kinase A (PKA) pathway (Paul et al. 2000). Dephosphorylation of this residue by Ca²⁺-dependent phosphatase calcineurin, following glutamate/NMDA receptor stimulation, renders STEP active in terms of its ability to bind to its substrates (Paul et al. 2003). Active STEP, in turn can bind to and modulate the activity of its substrate through tyrosine dephosphorylation of a regulatory tyrosine residue. Known substrates of STEP include ERK (extracellular regulated kinase 1/2) and p38 MAPKs (Paul et al. 2007, Poddar et al., 2010, Xu et al., 2009), Src family tyrosine kinases and NMDAR subunits (Nguyen et al., 2002; Braithwaite et al., 2006), all of which are involved in neuronal survival and death.

Brief Description of the Drawings

[008] Fig. 1 is a schematic diagram of STEP₆₁.

[009] Fig. 2 is a schematic diagram of STEP₄₆.

[010] Fig. 3 is a schematic diagram of a STEP-derived peptide according to an embodiment of the present disclosure, wherein the PKA phosphorylation site * has been modified.

[011] Fig. 4 is a schematic diagram of a STEP-derived peptide according to another embodiment of the present disclosure, wherein both the PKA phosphorylation site * and the KIS-domain phosphorylation sites have been modified.

[012] Fig. 5 is a schematic diagram of a STEP-derived peptide according to still another embodiment of the present disclosure, wherein the PKA phosphorylation site * has been modified and further including a TAT sequence at the N-terminal of the peptide.

[013] Fig. 6 is a schematic diagram of a STEP-derived peptide according to another embodiment of the present disclosure, wherein both the PKA phosphorylation site * and the KIS-domain phosphorylation sites have been modified and further including a TAT sequence at the N-terminal of the peptide.

[014] Fig. 7 is a schematic diagram of a STEP-derived peptide according to yet another embodiment of the present disclosure, wherein the PKA phosphorylation site * has been modified and further including a TAT sequence at the N-terminal of the peptide and a myc-tag at the C-terminal of the KIM domain.

[015] Fig. 8 is a schematic diagram of a STEP-derived peptide according to another embodiment of the present disclosure, wherein both the PKA phosphorylation site * and the KIS-domain phosphorylation sites have been modified and further including a TAT sequence at the N-terminal of the peptide and a myc-tag at the C-terminal of the KIM domain.

[016] Fig. 9 is a schematic illustration showing the injection site of the TAT-STEP peptide.

[017] Fig. 10 shows the immunohistochemistry of coronal section of striatum with anti-myc antibody to detect exogenous TAT-STEP.

[018] Fig. 11 shows the immunohistochemistry of cortex with anti-myc antibody to detect exogenous TAT-STEP.

[019] Fig. 12 shows representative TTC (triphenyltetrazolium chloride) stained sections from saline and TAT-STEP-myc treatment groups where treatment was administered immediately before an ischemic event.

[020] Fig. 13 is a quantitative analysis of the infarct volume. Infarct volumes are expressed as a percentage of the contralateral (control) hemisphere and the data are mean $\pm$ SEM ($p < 0.01$).

[021] Fig. 14 shows representative TTC (triphenyltetrazolium chloride) stained sections from saline and TAT-STEP-myc treatment groups where treatment was administered 1.5 hours after an ischemic event.

[022] Fig. 15 is a quantitative analysis of the infarct volume. Infarct volumes are expressed as a percentage of the contralateral (control) hemisphere and the data are mean $\pm$ SEM ($p < 0.01$).

Detailed Description

[023] According to various embodiments, the present disclosure provides a plurality of novel peptide sequences that are modified derivatives of a neuron-specific tyrosine phosphatase. Specifically, the novel peptide sequences are modified derivatives of striatal-enriched tyrosine phosphatase (STEP). In each of the peptide sequences, STEP has been modified so as to be able to ameliorate and treat brain injury resulting from glutamate toxicity, hypoxia and/or transient ischemic insult. Examples of the types of brain injury which the presently disclosed peptide sequences are useful for treating includes acute brain injury resulting from stroke and traumatic brain injury, and chronic disorders such as Huntington's chorea and schizophrenia. Furthermore, the presently described peptide sequences may further be useful in the treatment and amelioration of disorders associated with fear memory such as post-traumatic stress disorder.

[024] Glutamate, the most common excitatory neurotransmitter in the brain, is a major target for clinical intervention in ischemic stroke. Sustained elevation of glutamate has been observed for at least 24 hours in patients with acute ischemic stroke (Bullock et al., 1995; Davalos et al., 1997). This maintained glutamate excitotoxicity is presumably related to the progression of ischemic brain damage over a period of hours to days after the primary event. The events leading up to glutamate toxicity begin as oxygen and glucose deprivation from ischemia result in the excessive release of glutamate followed by over activation of N-methyl-D-aspartate (NMDA)-subtype of glutamate receptors that constitutes about 80% of all the excitatory receptors in the brain. Glutamate-mediated NMDAR stimulation induces large increases in intracellular Ca^{2+} , leading to the activation of multiple signaling pathways that are involved in the propagation and maintenance of inflammatory response in the injured brain (Mehta et al., 2007). Activation of p38 MAPK signaling pathway in particular has been implicated in enhancing ischemic brain injury and has been linked to the production of cyclooxygenase-2, a key enzyme involved in inflammatory response and generation of reactive oxygen species (Nito et al., 2008; Iadecola et al., 2001, Barone et al., 2001a and b). However a significant concern in the pharmaceutical application of NMDA receptor antagonists is that they not only block its pathological effects but also interferes with physiological functions of NMDAR (Chazot et al., 2004; Dirnagl et al., 1999; Kemp et al., 2002; Lipton et al., 2004) that includes excitatory synaptic transmission and plasticity. Therefore, identifying specific molecular targets that are activated following NMDAR stimulation and will interfere selectively with the pathological effects of NMDAR may provide novel compounds for the attenuation of ischemic brain damage.

[025] STEP is an intracellular protein tyrosine phosphatase (PTP) that is exclusively expressed in the central nervous system. STEP is preferentially expressed in neurons of the basal ganglia, hippocampus, cortex and related structures (Lombroso et al. 1993, Boulanger et al. 1995). The STEP-family of PTPases includes both membrane associated (STEP₆₁) and cytosolic (STEP₄₆) variants that are formed by alternative splicing of a single gene (Bult et al. 1997). STEP₆₁ differs from STEP₄₆ by the presence of an additional 172 amino acids at its N-terminus (Bult et al. 1996). For the purposes of this disclosure, specific amino acids residues within the STEP protein are frequently referred to using the numbering from the STEP₄₆ variant. The 107 amino acid sequence of the STEP protein discussed herein is highly conserved between animal species with 95% sequence homology between rat and human. (Compare SEQ IDs 1 and 2.) The present disclosure provides for a number of possible mutations of the STEP protein in order to achieve the useful peptides of the disclosed herein. Where the described mutation sites are conserved between species, it will be well understood that the various STEP-based peptides described herein may be based on or derived from the STEP proteins from a variety of species and that such peptides may or may not be derived from proteins that are endogenous to the specific species being treated and/or studied. Accordingly, the peptides disclosed herein should not be construed as being limited to the specific peptides sequences included in the sequence listing.

[026] STEP along with two other PTPs, PTPRR and HePTP belongs to a family of PTPs that contains a highly conserved 16-amino acid substrate-binding domain termed the kinase interacting motif (KIM domain) (Pulido et al. 1998). A regulatory serine residue, ser 49 (ser 221 in STEP₆₁) lies in the middle of the KIM domain and dephosphorylation of this residue renders STEP active in terms of its ability to bind to its substrates. Phosphorylation of ser 49 is mediated by dopamine/D1 receptor dependent activation of the cAMP/PKA pathway (Paul et al. 2000) while dephosphorylation is mediated by glutamate/NMDA receptor induced activation of the Ca²⁺ dependent phosphatase, calcineurin (Paul et al. 2003).

[027] A second conserved domain carboxy-terminal to the KIM domain is present in STEP, PTP-SL and HePTP. As described in further detail below, we have determined that this domain, termed the kinase specificity sequence (KIS domain) includes two phosphorylation sites, Thr 59 (Thr 231 in STEP₆₁) and Ser 72 (Ser 244 in STEP₆₁), which are important in regulating the stability of the STEP protein.

[028] Figs. 1 and 2 are schematic diagrams of STEP₆₁ and STEP₄₆, respectively, indicating the positions of the phosphatase domain, putative proteolytic sites (PEST), transmembrane domain (TM), polyproline rich domain (PP), kinase interacting motif (KIM), kinase

specificity sequence (KIS) and the above-mentioned phosphorylation sites involved in the activation and subsequent degradation of STEP. Additionally we have indicated the position of a cysteine residue (Cys 23 in STEP₄₆/Cys 195 in STEP₆₁) that has been shown to be involved in intermolecular dimerization and a threonine residue (Thr 18 in STEP₄₆/Thr 190 in STEP₆₁) that is known to be phosphorylatable by both ERK and p38 MAPKs.

[029] In its active form STEP can modulate synaptic plasticity by regulating the activity of extracellular regulated kinase 1/2 (ERK1/2), a key protein involved in memory formation (Paul et al. 2007 and Thomas and Huganir 2004). Active STEP can also modulate NMDA receptor-dependent long term potentiation by interfering with NMDA receptor trafficking to synaptic membrane, possibly through regulation of the upstream kinase Fyn (Nguyen et al. 2002) and tyrosine dephosphorylation of NR2B-NMDA receptor subunits (Snyder et al. 2005). Several studies also indicate a role of active STEP in neuroprotection through its regulation of p38 MAPK (Poddar et al., 2010, Xu et al., 2009).

[030] Because the STEP protein is known to interfere with NMDAR, a constitutively active peptide based on the STEP protein is a likely candidate for treatment, amelioration and/or prevention of ischemic brain damage. Fig. 3 is a diagram of a constitutively active STEP-derived peptide according to an embodiment of the present disclosure. The peptide contains the first 107 amino acids of STEP₄₆ including the KIM and KIS domains. Furthermore, as shown, the serine residue (Ser 49) which acts as a PKA phosphorylation site * has been modified. (See SEQ. ID. No. 3.) In the depicted embodiment, Ser 49 has been converted, using standard point mutation techniques, to alanine, which is non-phosphorylatable, resulting in a constitutively active peptide. Modification of the PKA phosphorylation site addresses the problem of inactivation of the STEP-derived peptide due to phosphorylation.

[031] Additional studies, as described in Mukherjee et al. 2011, shows that the active STEP protein is more susceptible to degradation. The study highlights the role of the KIS domain in regulation of the level of active STEP. We identified two SP/TP sites (Fig. 2, Thr 59 and Ser 72) in the KIS domain whose endogenous phosphorylation is mediated primarily through the basal activity of ERK and p38 MAPKs. Dephosphorylation of these two sites selectively results in ubiquitin-mediated proteasomal degradation of the active form of STEP. These findings imply that ubiquitin-mediated proteasomal degradation of active STEP may also lead to secondary activation of p38 MAPK.

[032] Accordingly, it may be desirable to produce an active STEP-derived peptide that remains stable and is not susceptible to ubiquitin-mediated proteasomal degradation in vivo.

[033] Figure 4 is a schematic illustration of a STEP-derived peptide according to an embodiment of the present disclosure, wherein Ser49 has been converted to alanine, which is non-phosphorylatable and Thr 59 and Ser 72 are converted, to glutamic acid to mimic the phosphorylatable form. (See SEQ. ID. No. 4.) However, it will be appreciated that other phospho-mimics may be used including, for example, Aspartic acid.

[034] More recent studies as described in Deb et al., 2011, show that enzymatic activity of both STEP₆₁ and STEP₄₆ are also regulated by intermolecular dimerization. Dimerization of STEP involves intermolecular disulfide bond formation involving several cysteine residues and oxidative stress leads to increase in dimerization of STEP resulting in reduced activity. One such cysteine residue (Cys 23 in STEP₄₆/Cys 195 in STEP₆₁) that is involved in intermolecular dimerization is present in the STEP derived peptide described herein. It is possible that oxidative stress during an ischemic insult and / or reperfusion can lead to dimerization, at least in part, of the STEP derived peptide, thereby reducing its therapeutic efficacy. Accordingly, the present disclosure further provides a STEP peptide including a mutated Cys 23 residue (cysteine to alanine) (SEQ. ID. No. 5).

[035] We have also identified another phosphorylation site in the STEP-derived peptide reported herein (Thr 18 in STEP₄₆/Thr 190 in STEP₆₁). *In vitro* studies show that this site is phosphorylatable by both ERK and p38 MAPKs. Based on the functional significance of phosphorylation, the present disclosure further provides a STEP peptide including a non-phosphorylatable or mimic phosphorylatable form of Thr 18 (See SEQ. ID No. 6, Thr18 mutated to Glutamic Acid and SEQ ID. No. 7, Thr18 mutated to Alanine).

[036] SEQ ID. No. 8 provides the amino acid sequence of the human STEP Peptide including all the mutations points discussed above.

[037] According to some embodiments, the present disclosure provides for the prevention, treatment, or amelioration of brain injury caused by excessive glutamate release and oxidative stress. Accordingly, the present disclosure also provides STEP-derived peptides which can be delivered to the brain of a patient who is or has suffered from an incident, such as a stroke, resulting in this type of injury. Figs. 5 and 6 are schematic illustrations of the previously described constitutively active STEP-derived peptides including a TAT sequence at the N-terminal of the peptide. In Fig. 5 only the phosphorylation site in the KIM domain has been altered. (See SEQ. ID No. 9) In Fig. 6, the phosphorylation sites in both the KIM and KIS domains have been altered. (See SEQ ID. No. 10). TAT is an 11 amino acid peptide that renders peptide sequences cell permeable and enables these peptides to cross the blood brain barrier. The ability of the STEP-derived peptide to cross the blood brain barrier

enables the peptide to be delivered to a patient's brain via the significantly less invasive mechanism of intravenous injection, for example via the femoral vein, rather than previous treatment mechanisms that require direct surgical access to the brain. Of course, those of skill in the art will be familiar with other suitable delivery mechanisms that could be employed including, for example, known targeted and viral-based delivery systems.

[038] For the purposes of experimentation, it is frequently useful to include an observable tag on the molecule being studied. Accordingly, the present disclosure further provides STEP-derived peptides including an observable tag. Figs. 7 and 8 are schematic illustrations of the previously described constitutively active STEP-peptides including a TAT sequence at the N-terminus and wherein the C-terminus of the peptide has been modified to include a myc tag. In Fig. 7 only the phosphorylation site in the KIM domain has been altered. In Fig. 8, the phosphorylation sites in both the KIM and KIS domains have been altered. Of course, those of skill in the art will be familiar with various observable tags and mechanisms for modifying peptides to include these tags.

[039] Using a cell culture model of glutamate toxicity we first investigated the functional significance of STEP's activation following excitotoxic NMDA receptor stimulation. As described in Paul et al., 2010, we have demonstrated that following an excitotoxic insult, NR2B-NMDA receptor mediated increases in Ca^{2+} leads to dephosphorylation and subsequent activation of STEP. As described in Poddar et al., 2010, using a substrate-trapping mutant we established that p38 MAPK is a substrate of STEP and the KIM domain of STEP is involved in STEP-p38 interaction. NR2B-NMDAR mediated activation of STEP promotes cell survival by down-regulating p38 MAPK signaling pathway. However sustained NMDAR stimulation results in degradation of active STEP leading to secondary activation of p38 MAPK that triggers cell death pathways. Based on these findings we generated a constitutively active form of a STEP peptide that can bind irreversibly to p38 MAPK. It was rendered cell permeable by fusion to the protein transduction domain of the human immunodeficiency virus-type I, TAT. Transduction of the purified TAT-STEP-myc peptide in cultured neurons prior to or immediately after treatment with an excitatory dose of glutamate (100 μM , 30 min) blocked neuronal cell death as assessed by Hoechst DNA staining 24 hr later. Additional studies in an animal model of glutamate toxicity showed that the TAT-STEP peptide, when injected stereotactically into the striatum prior to glutamate, could attenuate glutamate-induced excitotoxic cell death *in vivo*.

[040] In subsequent experiments we used oxygen-glucose deprivation (OGD) as an *in vitro* model of cerebral ischemia to evaluate the role of TAT-STEP peptide in neuroprotection

following an ischemic insult. Cultured neurons were submerged in glucose free buffer in a hypoxic chamber (oxygen-glucose deprivation or OGD) for 2 hr in the presence or absence of the peptide and cell viability was tested by Hoechst DNA staining 24 hr later. Our findings show that pre-incubation with the TAT-STEP peptide resulted in a significant reduction in OGD-induced cell death (manuscript under preparation).

[041] The specific methods and compositions described herein are representative of preferred embodiments and are exemplary and not intended as limitations on the scope of the invention. Other objects, aspects, and embodiments will occur to those skilled in the art upon consideration of this specification, and are encompassed within the spirit of the invention as defined by the scope of the claims. It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention. The invention illustratively described herein suitably may be practiced in the absence of any element or elements, or limitation or limitations, which is not specifically disclosed herein as essential. The methods and processes illustratively described herein suitably may be practiced in differing orders of steps, and that they are not necessarily restricted to the orders of steps indicated herein or in the claims. As used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality (for example, a culture or population) of such host cells, and so forth.

[042] Under no circumstances may the patent be interpreted to be limited to the specific examples or embodiments or methods specifically disclosed herein. Under no circumstances may the patent be interpreted to be limited by any statement made by any Examiner or any other official or employee of the Patent and Trademark Office unless such statement is specifically and without qualification or reservation expressly adopted in a responsive writing by Applicants.

[043] The terms and expressions that have been employed are used as terms of description and not of limitation, and there is no intent in the use of such terms and expressions to exclude any equivalent of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention as claimed. Thus, it will be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such

modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

[044] All patents and publications referenced below and/or mentioned herein are indicative of the levels of skill of those skilled in the art to which the invention pertains.

[045] References:

[046] Lipton S. A. and Rosenberg P. A. (1994) Excitatory amino acids as a final common pathway for neurologic disorders. *The New England journal of medicine* 330, 613-622.

[047] Arundine M. and Tymianski M. (2004) Molecular mechanisms of glutamate-dependent neurodegeneration in ischemia and traumatic brain injury. *Cell Mol Life Sci* 61, 657-668.

[048] Barone F. C., Irving E. A., Ray A. M., Lee J. C., Kassis S., Kumar S., Badger A. M., Legos J. J., Erhardt J. A., Ohlstein E. H., Hunter A. J., Harrison D. C., Philpott K., Smith B. R., Adams J. L. and Parsons A. A. (2001a) Inhibition of p38 mitogen-activated protein kinase provides neuroprotection in cerebral focal ischemia. *Medicinal research reviews* 21, 129-145.

[049] Barone F. C., Irving E. A., Ray A. M., Lee J. C., Kassis S., Kumar S., Badger A. M., White R. F., McVey M. J., Legos J. J., Erhardt J. A., Nelson A. H., Ohlstein E. H., Hunter A. J., Ward K., Smith B. R., Adams J. L. and Parsons A. A. (2001b) SB 239063, a second-generation p38 mitogen-activated protein kinase inhibitor, reduces brain injury and neurological deficits in cerebral focal ischemia. *The Journal of pharmacology and experimental therapeutics* 296, 312-321.

[050] Luo J., Wang Y., Yasuda R. P., Dunah A. W. and Wolfe B. B. (1997) The majority of N-methyl-D-aspartate receptor complexes in adult rat cerebral cortex contain at least three different subunits (NR1/NR2A/NR2B). *Molecular pharmacology* 51, 79-86.

[051] Tovar K. R. and Westbrook G. L. (1999) The incorporation of NMDA receptors with a distinct subunit composition at nascent hippocampal synapses in vitro. *J Neurosci* 19, 4180-4188.

[052] Kohr G. (2006) NMDA receptor function: subunit composition versus spatial distribution. *Cell and tissue research* 326, 439-446.

[053] Liu Y., Wong T. P., Aarts M., Rooyakkers A., Liu L., Lai T. W., Wu D. C., Lu J., Tymianski M., Craig A. M. and Wang Y. T. (2007) NMDA receptor subunits have

differential roles in mediating excitotoxic neuronal death both in vitro and in vivo. *J Neurosci* 27, 2846-2857.

[054] Tu W., Xu X., Peng L., Zhong X., Zhang W., Soundarapandian M. M., Babel C., Wang M., Jia N., Zhang W., Lew F., Chan S. L., Chen Y. and Lu Y. (2010) DAPK1 interaction with NMDA receptor NR2B subunits mediates brain damage in stroke. *Cell* 140, 222-234.

[055] Rodrigues S. M., Schafe G. E. and LeDoux J. E. (2001) Intra-amygdala blockade of the NR2B subunit of the NMDA receptor disrupts the acquisition but not the expression of fear conditioning. *J Neurosci* 21, 6889-6896.

[056] Zhao M. G., Toyoda H., Lee Y. S., Wu L. J., Ko S. W., Zhang X. H., Jia Y., Shum F., Xu H., Li B. M., Kaang B. K. and Zhuo M. (2005) Roles of NMDA NR2B subtype receptor in prefrontal long-term potentiation and contextual fear memory. *Neuron* 47, 859-872.

[057] Walker D. L. and Davis M. (2008) Amygdala infusions of an NR2B-selective or an NR2A-preferring NMDA receptor antagonist differentially influence fear conditioning and expression in the fear-potentiated startle test. *Learning & memory* (Cold Spring Harbor, N.Y. 15, 67-74.

[058] Boulanger L. M., Lombroso P. J., Raghunathan A., During M. J., Wahle P. and Naegle J. R. (1995) Cellular and molecular characterization of a brain-enriched protein tyrosine phosphatase. *J Neurosci* 15, 1532-1544.

[059] Bult A., Zhao F., Dirks R., Jr., Raghunathan A., Solimena M. and Lombroso P. J. (1997) STEP: a family of brain-enriched PTPs. Alternative splicing produces transmembrane, cytosolic and truncated isoforms. *European journal of cell biology* 72, 337-344.

[060] Pulido R., Zuniga A. and Ullrich A. (1998) PTP-SL and STEP protein tyrosine phosphatases regulate the activation of the extracellular signal-regulated kinases ERK1 and ERK2 by association through a kinase interaction motif. *The EMBO journal* 17, 7337-7350.

[061] Paul S., Snyder G. L., Yokakura H., Picciotto M. R., Nairn A. C. and Lombroso P. J. (2000) The Dopamine/D1 receptor mediates the phosphorylation and inactivation of the protein tyrosine phosphatase STEP via a PKA-dependent pathway. *J Neurosci* 20, 5630-5638.

[062] Paul S., Olausson P., Venkitaramani D. V., Ruchkina I., Moran T. D., Tronson N., Mills E., Hakim S., Salter M. W., Taylor J. R. and Lombroso P. J. (2007) The striatal-enriched protein tyrosine phosphatase gates long-term potentiation and fear memory in the lateral amygdala. *Biological psychiatry* 61, 1049-1061.

- [063] Poddar R. and Paul S. (2009) Homocysteine-NMDA receptor-mediated activation of extracellular signal-regulated kinase leads to neuronal cell death. *Journal of neurochemistry* 110, 1095-1106.
- [064] Xu J., Kurup P., Zhang Y., Goebel-Goody S. M., Wu P. H., Hawasli A. H., Baum M. L., Bibb J. A. and Lombroso P. J. (2009) Extrasynaptic NMDA receptors couple preferentially to excitotoxicity via calpain-mediated cleavage of STEP. *J Neurosci* 29, 9330-9343.
- [065] Nguyen T. H., Liu J. and Lombroso P. J. (2002) Striatal enriched phosphatase 61 dephosphorylates Fyn at phosphotyrosine 420. *The Journal of biological chemistry* 277, 24274-24279.
- [066] Braithwaite S. P., Xu J., Leung J., Urfer R., Nikolic K., Oksenberg D., Lombroso P. J. and Shamloo M. (2008) Expression and function of striatal enriched protein tyrosine phosphatase is profoundly altered in cerebral ischemia. *The European journal of neuroscience* 27, 2444-2452.
- [067] Bullock R, Zauner A, Woodward J, Young HF. Massive persistent release of excitatory amino acids following human occlusive stroke. *Stroke*. 1995;26:2187-2189
- [068] Davalos A, Castillo J, Serena J, Noya M. Duration of glutamate release after acute ischemic stroke. *Stroke*. 1997;28:708-710
- [069] Mehta SL, Manhas N, Raghubir R. Molecular targets in cerebral ischemia for developing novel therapeutics. *Brain Res Rev*. 2007;54:34-66
- [070] Chazot PL. The nmda receptor nr2b subunit: A valid therapeutic target for multiple cns pathologies. *Curr Med Chem*. 2004;11:389-396
- [071] Dirnagl U, Iadecola C, Moskowitz MA. Pathobiology of ischaemic stroke: An integrated view. *Trends Neurosci*. 1999;22:391-397
- [072] Kemp JA, McKernan RM. Nmda receptor pathways as drug targets. *Nat Neurosci*. 2002;5 Suppl:1039-1042
- [073] Lipton SA. Failures and successes of nmda receptor antagonists: Molecular basis for the use of open-channel blockers like memantine in the treatment of acute and chronic neurologic insults. *NeuroRx*. 2004;1:101-110
- [074] Nito C, Kamada H, Endo H, Niizuma K, Myer DJ, Chan PH. Role of the p38 mitogen-activated protein kinase/cytosolic phospholipase a2 signaling pathway in blood-brain barrier disruption after focal cerebral ischemia and reperfusion. *J Cereb Blood Flow Metab*. 2008;28:1686-1696

- [075] Iadecola C, Niwa K, Nogawa S, Zhao X, Nagayama M, Araki E, Morham S, Ross ME. Reduced susceptibility to ischemic brain injury and n-methyl-d-aspartate-mediated neurotoxicity in cyclooxygenase-2-deficient mice. *Proc Natl Acad Sci U S A*. 2001;98:1294-1299.
- [076] Deb I, Poddar R, Paul S. Oxidative stress-induced oligomerization inhibits the activity of the non-receptor tyrosine phosphatase STEP61. *J Neurochem*. 2011;116:1097-1111.
- [077] Braithwaite SP, Adkisson M, Leung J, Nava A, Masterson B, Urfer R, Oksenberg D, Nikolic K. Regulation of NMDA receptor trafficking and function by striatal-enriched tyrosine phosphatase (STEP). *Eur J Neurosci*. 2006;23:2847-2856.
- [078] Candelario-Jalil E, González-Falcón A, García-Cabrera M, León OS, Fiebich BL. Wide therapeutic time window for nimesulide neuroprotection in a model of transient focal cerebral ischemia in the rat. *Brain Res*. 2004 ;1007:98-108.
- [079] Lombroso, P. J., Naegele, J. R., Sharma, E., and Lerner, M. (1993) *J Neurosci* 13(7), 3064-3074
- [080] Bult, A., Zhao, F., Dirks, R., Jr., Sharma, E., Lukacs, E., Solimena, M., Naegele, J. R., and Lombroso, P. J. (1996) *J Neurosci* 16(24), 7821-7831
- [081] Paul, S., Nairn, A. C., Wang, P., and Lombroso, P. J. (2003) *Nature neuroscience* 6(1), 34-42
- [082] Thomas, G. M., and Huganir, R. L. (2004) *Nature reviews* 5(3), 173-183
- [083] Snyder, E. M., Nong, Y., Almeida, C. G., Paul, S., Moran, T., Choi, E. Y., Nairn, A. C., Salter, M. W., Lombroso, P. J., Gouras, G. K., and Greengard, P. (2005) *Nature neuroscience* 8(8), 1051-1058
- [084] Mukherjee, S.; Poddar, R.; Deb, I., Paul, S., (2011) "Dephosphorylation of Specific Sites in the KIS domain leads to Ubiquitin-Mediated Degradation of the tyrosine Phosphatase" (In preparation)
- [085] Deb, I.; Poddar, R.; Paul, S.; (2011) "Oxidative stress induced dimerization inhibits the activity of the non-receptor tyrosine phosphatase STEP61" *Journal of Neurochemistry* (In publication)

Experiments – Prevention, treatment and amelioration of brain damage with TAT-STP peptide.

- [086] Using an animal model of cerebral ischemia (middle cerebral artery occlusion for 90 min to generate embolic stroke followed by reperfusion for 24 hr), we evaluated whether

degradation of active STEP contributes to neuronal cell death and subsequent brain damage and if the modified STEP peptide can mitigate ischemic brain damage. At first we examined the temporal profile of STEP's activation and subsequent degradation following cerebral ischemia and reperfusion and also evaluated its correlation with p38 MAPK activation. The findings show a rapid but transient increase in phosphorylation of p38 MAPK within 15 min of ischemic insult. This was followed by a time-dependent decrease in p38 phosphorylation from 30 min, which returned to basal levels by 90 min. The ischemic insult not only led to rapid activation of p38 MAPK but also resulted in dephosphorylation and subsequent activation of STEP, which begins at 30 min and leads to complete dephosphorylation by 90 min. As a consequence, active STEP limits the duration of activation of p38 MAPK. Progressive down regulation of active STEP through degradation is evident from ~6 hours after reperfusion resulting in secondary activation of p38 MAPK. The secondary but not the primary activation of p38 MAPK pathway correlate with the expression of cyclooxygenase-2 (COX-2), a key enzyme that is linked to p38 MAPK activation and is involved in inflammatory response and reactive oxygen species generation (manuscript in preparation). Consistent with our hypothesis, the above findings indicate that STEP is a negative regulator of cell death pathways and that degradation of active STEP trigger signaling cascades that leads to ischemic brain injury.

[087] We next investigated whether a modified TAT-STEP peptide delivered *in vivo* can reduce ischemic brain damage. A major technical challenge here was to successfully deliver the peptide in the brain in a non-invasive manner. Figs. 9-15 show that following a single intravenous injection of a TAT-STEP peptide through the femoral vein of an adult rat, the peptide was detectable both in the cortex (Fig. 10) and striatum (Fig. 11). In these experiments, Male Wistar rats received a single intravenous injection of 1 ml (1 mg/ml) of the TAT-STEP peptide of Fig. 8 through the femoral vein (Fig. 9). Detection took place 1 hour after injection.

[088] To determine the efficacy of the peptide in attenuating ischemic brain injury, the TAT-STEP peptide of SEQ ID. No. 11 was administered intravenously immediately before or 1.5 hr after the onset of stroke. The findings demonstrate that intravenous injection of the cell-permeable STEP peptide either before or after an acute ischemic stroke can provide significant protection against ischemic brain injury. In these experiments, to generate acute ischemic stroke, male Wistar rats (260-280 gm) were subjected to middle cerebral artery occlusion (MCAO) for 90 minutes using the intraluminal suture method as described in Cadelario-Jalil et al., 2004. TAT-STEP peptide or saline was injected in the rats immediately

prior to (Figs. 12-13) or 1.5 hours after (Figs. 14-15) MCAO followed by reperfusion for 24 hr. The fact that the peptide when injected even 1.5 hr after the onset of stroke can mitigate ischemic brain injury suggests that the secondary activation of p38 MAPK following the degradation of endogenous STEP is possibly responsible for triggering intracellular death pathways. Taken together the findings further imply that the modified STEP-derived peptide may be an alternative therapeutic target, downstream of NMDAR, for intervention in diseases that involve excitotoxicity.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 52789-5 Seq 16-MAY-13 vl.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> STC.UNM

<120> STEP-Derived Peptide for Brain Injury Treatment

<130> 52789-5

<140> CA 2,811,086

<141> 2011-09-15

<150> US 61/383,208

<151> 2010-09-15

<160> 11

<170> PatentIn version 3.5

<210> 1

<211> 107

<212> PRT

<213> Rat

<400> 1

Met	Glu	Glu	Lys	Val	Glu	Asp	Asp	Phe	Leu	Asp	Leu	Asp	Ala	Val	Pro
1			5						10					15	
Glu	Thr	Pro	Val	Phe	Asp	Cys	Val	Met	Asp	Ile	Lys	Pro	Glu	Thr	Asp
			20					25					30		
Pro	Ala	Ser	Leu	Thr	Val	Lys	Ser	Met	Gly	Leu	Gln	Glu	Arg	Arg	Gly
			35					40					45		
Ser	Asn	Val	Ser	Leu	Thr	Leu	Asp	Met	Cys	Thr	Pro	Gly	Cys	Asn	Glu
			50				55				60				
Glu	Gly	Phe	Gly	Tyr	Leu	Val	Ser	Pro	Arg	Glu	Glu	Ser	Ala	His	Glu
65					70				75					80	
Tyr	Leu	Leu	Ser	Ala	Ser	Arg	Val	Leu	Arg	Ala	Glu	Glu	Leu	His	Glu
				85					90					95	
Lys	Ala	Leu	Asp	Pro	Phe	Leu	Leu	Gln	Ala	Glu					
				100					105						

<210> 2
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
 1 5 10 15
 Glu Thr Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp
 20 25 30
 Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
 35 40 45
 Ser Asn Val Ser Leu Thr Leu Asp Met Cys Thr Pro Gly Cys Asn Glu
 50 55 60
 Glu Gly Phe Gly Tyr Leu Met Ser Pro Arg Glu Glu Ser Ala Arg Glu
 65 70 75 80
 Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
 85 90 95
 Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
 100 105

<210> 3
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 3
 Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
 1 5 10 15
 Glu Thr Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp
 20 25 30
 Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
 35 40 45
 Ala Asn Val Ser Leu Thr Leu Asp Met Cys Thr Pro Gly Cys Asn Glu
 50 55 60
 Glu Gly Phe Gly Tyr Leu Met Ser Pro Arg Glu Glu Ser Ala Arg Glu
 65 70 75 80
 Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
 85 90 95
 Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
 100 105

<210> 4
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
 1 5 10 15
 Glu Thr Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp
 20 25 30
 Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
 35 40 45

Ala Asn Val Ser Leu Thr Leu Asp Met Cys Gly Pro Gly Cys Asn Glu
 50 55 60
 Glu Gly Phe Gly Tyr Leu Met Gly Pro Arg Glu Glu Ser Ala Arg Glu
 65 70 75 80
 Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
 85 90 95
 Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
 100 105

<210> 5
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
 1 5 10 15
 Glu Thr Pro Val Phe Asp Ala Val Met Asp Ile Lys Pro Glu Ala Asp
 20 25 30
 Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
 35 40 45
 Ser Asn Val Ser Leu Thr Leu Asp Met Cys Thr Pro Gly Cys Asn Glu
 50 55 60
 Glu Gly Phe Gly Tyr Leu Met Ser Pro Arg Glu Glu Ser Ala Arg Glu
 65 70 75 80
 Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
 85 90 95
 Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
 100 105

<210> 6
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 6
 Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
 1 5 10 15
 Glu Gly Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp
 20 25 30
 Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
 35 40 45
 Ser Asn Val Ser Leu Thr Leu Asp Met Cys Thr Pro Gly Cys Asn Glu
 50 55 60
 Glu Gly Phe Gly Tyr Leu Met Ser Pro Arg Glu Glu Ser Ala Arg Glu
 65 70 75 80
 Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
 85 90 95
 Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
 100 105

<210> 7
 <211> 107

<212> PRT

<213> Homo sapiens

<400> 7

```

Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
1          5          10          15
Glu Ala Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp
          20          25          30
Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
          35          40          45
Ser Asn Val Ser Leu Thr Leu Asp Met Cys Thr Pro Gly Cys Asn Glu
          50          55          60
Glu Gly Phe Gly Tyr Leu Met Ser Pro Arg Glu Glu Ser Ala Arg Glu
65          70          75          80
Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
          85          90          95
Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
          100          105

```

<210> 8

<211> 107

<212> PRT

<213> Homo sapiens

<400> 8

```

Met Glu Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro
1          5          10          15
Glu Glu Pro Val Phe Asp Ala Val Met Asp Ile Lys Pro Glu Ala Asp
          20          25          30
Pro Thr Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly
          35          40          45
Ala Asn Val Ser Leu Thr Leu Asp Met Cys Glu Pro Gly Cys Asn Glu
          50          55          60
Glu Gly Phe Gly Tyr Leu Met Ala Pro Arg Glu Glu Ser Ala Arg Glu
65          70          75          80
Tyr Leu Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu
          85          90          95
Lys Ala Leu Asp Pro Phe Leu Leu Gln Ala Glu
          100          105

```

<210> 9

<211> 121

<212> PRT

<213> Homo sapiens

<400> 9

```

Met Ala Leu Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg Gly Glu
1          5          10          15
Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro Glu Thr
          20          25          30
Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp Pro Thr
          35          40          45
Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly Ala Asn
          50          55          60

```

```

Val Ser Leu Thr Leu Asp Met Cys Thr Pro Gly Cys Asn Glu Glu Gly
65          70          75          80
Phe Gly Tyr Leu Met Ser Pro Arg Glu Glu Ser Ala Arg Glu Tyr Leu
          85          90          95
Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu Lys Ala
          100          105          110
Leu Asp Pro Phe Leu Leu Gln Ala Glu
          115          120

```

```

<210> 10
<211> 121
<212> PRT
<213> Homo sapiens

```

```

<400> 10
Met Ala Leu Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg Gly Glu
1          5          10          15
Glu Lys Ile Glu Asp Asp Phe Leu Asp Leu Asp Pro Val Pro Glu Thr
          20          25          30
Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Ala Asp Pro Thr
          35          40          45
Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly Ala Asn
          50          55          60
Val Ser Leu Thr Leu Asp Met Cys Gly Pro Gly Cys Asn Glu Glu Gly
65          70          75          80
Phe Gly Tyr Leu Met Gly Pro Arg Glu Glu Ser Ala Arg Glu Tyr Leu
          85          90          95
Leu Ser Ala Ser Arg Val Leu Gln Ala Glu Glu Leu His Glu Lys Ala
          100          105          110
Leu Asp Pro Phe Leu Leu Gln Ala Glu
          115          120

```

```

<210> 11
<211> 121
<212> PRT
<213> Rat

```

```

<400> 11
Met Ala Leu Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg Gly Glu
1          5          10          15
Glu Lys Val Glu Asp Asp Phe Leu Asp Leu Asp Ala Val Pro Glu Thr
          20          25          30
Pro Val Phe Asp Cys Val Met Asp Ile Lys Pro Glu Thr Asp Pro Ala
          35          40          45
Ser Leu Thr Val Lys Ser Met Gly Leu Gln Glu Arg Arg Gly Ala Asn
          50          55          60
Val Ser Leu Thr Leu Asp Met Cys Gly Pro Gly Cys Asn Glu Glu Gly
65          70          75          80
Phe Gly Tyr Leu Val Gly Pro Arg Glu Glu Ser Ala His Glu Tyr Leu
          85          90          95
Leu Ser Ala Ser Arg Val Leu Arg Ala Glu Glu Leu His Glu Lys Ala
          100          105          110
Leu Asp Pro Phe Leu Leu Gln Ala Glu
          115          120

```

CLAIMS:

1. An isolated peptide comprising a kinase interacting motif (KIM) domain and a kinase specificity sequence (KIS) domain of a striatal-enriched tyrosine phosphatase (STEP) protein, wherein a serine in the KIM domain is converted so as to prevent phosphorylation at that site and at least one of a threonine and a serine in the KIS domain is converted to a phospho-mimic so as to prevent phosphorylation at that site.
2. The isolated peptide of claim 1, wherein the serine residue of the KIM domain is converted to an alanine, resulting in a constitutively active peptide.
3. The isolated peptide of claim 2, wherein the serine residue is at position 49 using the numbering from SEQ ID NO: 1.
4. The isolated peptide of claim 1, wherein the threonine residue of the KIS domain is converted to an aspartic acid residue or a glutamic acid residue.
5. The isolated peptide of claim 1, wherein the serine residue of the KIS domain is converted to an aspartic acid residue or a glutamic acid residue.
6. The isolated peptide of claim 1, wherein the threonine residue and the serine residues of the KIS domain are converted to a phospho-mimic amino acid residue.
7. The isolated peptide of claim 5, wherein the serine residue of the KIS domain is at position 72 using the numbering from SEQ ID NO: 1.
8. The isolated peptide of claim 1, further comprising a region configured to render the peptide cell permeable.
9. The isolated peptide of claim 8, wherein the region configured to render the peptide cell permeable is a TAT domain.
10. The isolated peptide of claim 1, further comprising a region configured to act as an observable marker.

11. The isolated peptide of claim 10, wherein the region configured to act as an observable marker is a myc region.
12. The isolated peptide of claim 4, wherein the threonine residue of the KIS domain is at position 59 using the numbering from SEQ ID NO: 1.
13. The isolated peptide of claim 6, wherein the threonine residue of the KIS domain is replaced with an aspartic acid or a glutamic acid.
14. The isolated peptide of claim 1, wherein the isolated peptide is SEQ ID NO: 4.
15. The isolated peptide of claim 1, wherein the isolated peptide is SEQ ID NO: 8.
16. The isolated peptide of claim 1, wherein the isolated peptide is SEQ ID NO: 10.
17. The isolated peptide of claim 1, wherein the isolated peptide is SEQ ID NO: 11.
18. Use of the isolated peptide of any one of claims 1 to 17 for treating or ameliorating ischemia in a patient.

1/4

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7





Fig. 8



2/4

Fig. 9




Fig. 10





Fig. 11



TAT-STEP (anti-myc)

Fig. 12





Fig. 13



3/4

Fig. 14





Fig. 15



4/4

