

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
16 May 2002 (16.05.2002)

PCT

(10) International Publication Number  
**WO 02/38742 A2**

(51) International Patent Classification<sup>7</sup>: **C12N 9/00**

(74) Agent: **MUETING, Ann, M.**; Mueting, Raasch, Gebhardt, P.A., P.O. Box 581415, Minneapolis, MN 55458-1415 (US).

(21) International Application Number: PCT/US01/46782

(22) International Filing Date:  
8 November 2001 (08.11.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/246,827 8 November 2000 (08.11.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **THE UNIVERSITY OF GEORGIA RESEARCH FOUNDATION, INC.** [US/US]; 632 Boyd Graduate Studies Research Center, Athens, GA 30602-7411 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **TRAVIS, James** [US/US]; 825 Riverbend Parkway, Athens, GA 30605 (US). **POTEMPA, Jan, S.** [PL/US]; 170 Barrington Drive, Apt. #102, Athens, GA 30605 (US). **BANBULA, Agnieszka** [PL/US]; 6204 Drew Hill Lane, Chapel Hill, NC 27514 (US). **BUGNO, Marcin** [PL/US]; 6204 Drew Hill Lane, Chapel Hill, North Carolina 27514 (US).

Published:

— *without international search report and to be republished upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: DIPEPTIDYLPEPTIDASES AND METHODS OF USE

(57) Abstract: The present invention provides isolated polypeptides, dipeptidylpeptidases, active analogs, active fragments, or active modifications thereof, having amidolytic activity for cleavage of a peptide bond between the second and third amino acids from the N-terminal end of a target polypeptide, wherein the target polypeptide has an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the peptide. Isolated nucleic acids encoding dipeptidylpeptidases are also provided, as are methods of reducing growth of a bacterium by inhibiting a dipeptidylpeptidase.



WO 02/38742 A2

## DIPEPTIDYLPEPTIDASES AND METHODS OF USE

5

### RELATED APPLICATIONS

This application claims the benefit of the U.S. Provisional Application No. 60/246,827, filed November 8, 2000, which is incorporated by reference in its entirety.

10

### GOVERNMENT FUNDING

The present invention was made with government support under Grant No. DE 09761, awarded by the National Institutes of Health. The Government may have certain rights in this invention.

15

### BACKGROUND

*Porphyromonas gingivalis* (*P. gingivalis*), an oral anaerobic bacterium, has been implicated as a causative agent of adult type periodontitis. As an asaccharolytic organism, *P. gingivalis* is totally dependent on external sources of peptides that are necessary for its growth and proliferation. In order to fulfill such a fastidious nutritional requirement this bacterium evolved a complex system of proteolytic enzymes which are now recognized as important virulence factors in the development of periodontal disease (Travis et al., *J. Adv. Exp. Med. Biol.*, 477:455-65 (2000)). The best known and well characterized enzymes of this system are gingipains R and K, arginine and lysine specific, cysteine proteinases (Curtis et al., *J. Periodontal Res.*, 34:464-72 (1999)). Working in concert with the proteinases periodontain (Nelson et al., *J. Biol. Chem.*, 274:12245-51 (1999)), collagenases/gelatinases (Birkedal-Hansen et al., *J. Periodontal Res.*, 23:258-64 (1988); Lawson et al., *Infect. Immun.*, 60:1524-29 (1992); Kato et al., *J. Bacteriol.*, 174:3889-95 (1992), prtT (Otogoto et al., *Infect. Immun.*, 61:117-23 (1993)), and Tpr (Bourgeau et al., *Infect. Immun.*, 60:3186-92 (1992)) as well as host proteinases, this array of enzymes has the

potential to degrade proteins from both the periodontal ligamentum and surrounding tissues. Their concerted action leads to the formation of a large pool of oligopeptides, which can be further utilized by *P. gingivalis* and other oral bacteria. However, *P. gingivalis* cannot transport poly- and oligo- peptides into the cell, even though it has the ability to thrive on dipeptides as a sole source of carbon. This has led to an interest in studying a specialized group of *P. gingivalis* peptidases capable of hydrolyzing oligopeptides to di- and tripeptides, which can be subsequently metabolized by this periodontopathogen. The purification, characterization and cloning of prolyl tripeptidylpeptidase A (PtpA), an enzyme which liberates tripeptides from the N-terminal regions of substrates containing proline residues in the third position has been previously reported (Banbula et al., *J. Biol. Chem.*, 274:9246-52 (1999)).

Dipeptidylpeptidase-IV (DPP-IV), an enzyme with similar specificity, but only dipeptidylpeptidase activity, has also been cloned (Kiyama et al., *Biochim. Biophys. Acta*, 1396:39-46 (1998)), purified, and characterized (Kumagai et al., *Infect. Immun.*, 68:716-24 (2000); Banbula et al., *Infect. Immun.*, 68:1176-82 (2000)). Together with a recently described angiotensinogen-converting enzyme analogue (Awano et al., *FEBS Lett.*, 460:139-44 (1999)) all of these proteases can hydrolyze peptide bonds containing proline residues. In addition, the *P. gingivalis* genome contains three further putative coding sequences encoding proteinases homologous with dipeptidylpeptidase-IV, although their activities have not yet been identified (Banbula et al., *J. Biol. Chem.*, 274:9246-52 (1999)).

## SUMMARY OF THE INVENTION

In one aspect, the present invention provides an isolated dipeptidylpeptidase, active analog, active fragment, or active modification thereof having amidolytic activity for cleavage of a peptide bond between the second and third amino acids from the N-terminal end of a target polypeptide, wherein the target polypeptide has an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the polypeptide. Preferably, the dipeptidylpeptidase is isolated from *Porphyromonas gingivalis*. Preferably, the dipeptidylpeptidase is a serine

protease. Preferably, the dipeptidylpeptidase includes an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, and SEQ ID NO:26. Preferably the  
5 dipeptidylpeptidase is encoded by a nucleic acid including a nucleotide sequence SEQ ID NO:1.

In another aspect, the present invention provides an isolated polypeptide including an amino acid sequence having a percentage amino acid identity greater than about 40% with SEQ ID NO:2.

10 In another aspect, the present invention provides an isolated nucleic acid including a coding sequence encoding a dipeptidylpeptidase, active analog, active fragment, or active modification thereof having amidolytic activity for cleavage of a peptide bond between the second and third amino acids from the N-terminal end of a target polypeptide, wherein the target polypeptide has an  
15 aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the polypeptide. Preferably the nucleic acid includes a nucleotide sequence SEQ ID NO:1. Alternatively, the complement of the nucleic acid preferably hybridizes to SEQ ID NO:1 under hybridization conditions of 0.5 M phosphate buffer, pH 7.2, 7% SDS, 10 mM  
20 EDTA, at 68°C, followed by three for 20 minutes washes in 2x SSC, and 0.1% SDS, at 65°C, wherein at least about 20 nucleotides of the complement hybridize.

In another aspect, the present invention provides an isolated nucleic acid encoding a polypeptide, wherein the polypeptide includes an amino acid  
25 sequence having a percentage amino acid identity greater than about 40% with SEQ ID NO:2.

In another aspect, the present invention provides a method of identifying an inhibitor of a dipeptidylpeptidase, active analog, active fragment, or active modification thereof. The method includes identifying a compound that inhibits  
30 the amidolytic activity of the dipeptidylpeptidase by incubating the dipeptidylpeptidase with the compound under conditions that promote amidolytic activity of the dipeptidylpeptidase and determining if the amidolytic

activity of the dipeptidylpeptidase is inhibited relative to the amidolytic activity in the absence of the compound.

In another aspect, the present invention provides a method of reducing growth of a bacterium including inhibiting a dipeptidylpeptidase, active analog,  
5 active fragment, or active modification thereof, by contacting the dipeptidylpeptidase with an inhibitor of the dipeptidylpeptidase. Preferably the dipeptidylpeptidase is a serine protease.

In another aspect, the present invention provides a method for protecting an animal from a periodontal disease caused by *Porphyromonas gingivalis*  
10 including administering to the animal an inhibitor of dipeptidylpeptidase, wherein the disease is selected from the group consisting of gingivitis and periodontitis. Preferably the inhibitor is administered by a method selected from the group consisting of subgingival application and controlled release delivery.

In another aspect, the present invention provides an immunogenic  
15 composition including an isolated dipeptidylpeptidase, an antigenic analog, an antigenic fragment, or an antigenic modification thereof having amidolytic activity for cleavage of a peptide bond present in a target polypeptide, the peptide bond being located between the second and third amino acids from the N-terminal end of the target polypeptide, wherein the second amino acid from  
20 the N-terminal end has an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom. Preferably the dipeptidylpeptidase is a serine protease. Preferably the second amino acid is selected from the group consisting of alanine, phenylalanine, isoleucine, and leucine. The immunogenic composition may optionally include an adjuvant.

25 In another aspect, the present invention provides a composition including an inhibitor of an isolated dipeptidylpeptidase and a pharmaceutically acceptable carrier.

#### Definitions

"Polypeptide" as used herein refers to a polymer of amino acids and does  
30 not refer to a specific length of a polymer of amino acids. Thus, for example, the terms peptide, oligopeptide, protein, and enzyme are included within the definition of polypeptide. This term also includes post-expression modifications

of the polypeptide, for example, glycosylations, acetylations, phosphorylations, and the like. A polypeptide can be produced by an organism, or produced using recombinant techniques, or chemically or enzymatically synthesized.

"Polynucleotide" and "nucleic acid" are used herein interchangeably and refer to a linear polymeric form of nucleotides of any length, either  
5 ribonucleotides or deoxynucleotides, and include both double- and single-stranded DNA and RNA. A nucleic acid may include both coding and non-coding regions that can be obtained directly from a natural source (e.g., a microorganism), or can be prepared with the aid of recombinant or synthetic  
10 techniques. A nucleic acid may be equivalent to this nucleic acid or it can include, in addition, one or more other polynucleotides. For example, the nucleic acid of the invention can be a vector, such as an expression of a coding sequence.

"Peptidase," "proteinase," and "protease" all refer to enzymes that  
15 catalyze the hydrolysis of peptide bonds in a polypeptide. A "peptide bond" or "amide bond" is a covalent bond between the alpha-amino group of one amino acid and the carboxyl group of another amino acid. "Peptidase inhibitor," "proteinase inhibitor," "protease inhibitor," and "inhibitor" all refer to compounds that inhibit a peptidase that catalyzes the hydrolysis of peptide bonds  
20 in a polypeptide.

"Serine protease" refers to an enzyme that uses the hydroxy-functional side chain of serine as a nucleophile in a catalytic reaction.

"Amidolytic activity" refers to the ability of a polypeptide to catalyze the hydrolysis of at least one peptide bond in a polypeptide. The term "cleavage" can  
25 also be used to refer to the hydrolysis of a peptide bond in a polypeptide. A "dipeptidylpeptidase" is able to hydrolyze the peptide bond between the second and third amino acids from the N-terminal end of a target polypeptide including the general formula H-Xaa-Yaa-Xaa-, wherein Xaa is a natural or modified amino acid, and Yaa is an amino acid including an aliphatic or an aromatic  
30 residue as a substituent on the  $\alpha$ -carbon atom. Preferred amino acids in the Yaa position include alanine, phenylalanine, isoleucine, and leucine.

A "target polypeptide" is a polypeptide that is the potential substrate of the amidolytic activity of a dipeptidylpeptidase. A "dipeptidylpeptidase" does not have to cleave all members of the target polypeptide. The term "natural amino acid" refers to the 20 amino acids typically produced by a cell. The term  
5 "modified amino acid" refers to, for instance, acetylation, hydroxylation, methylation, amidation, or the attachment of carbohydrate or lipid moieties, cofactors, and the like.

As used herein, the term "isolated" means that a polypeptide or a polynucleotide has been either removed from its natural environment, produced  
10 using recombinant techniques, or chemically or enzymatically synthesized. Preferably, the polypeptide or polynucleotide is purified, i.e., essentially free from any other polypeptides, polynucleotides, and associated cellular products or other impurities.

An active analog, active fragment, or active modification of a  
15 polypeptide of the invention is one that has amidolytic activity by hydrolysis of a peptide bond present in the target polypeptide as described herein. Active analogs, active fragments, and active modifications are described in greater detail herein.

An antigenic analog, antigenic fragment, or antigenic modification of a  
20 polypeptide of the invention is one that has amidolytic activity by hydrolysis of a peptide bond present in the target polypeptide as described herein. Antigenic analogs, antigenic fragments, and antigenic modifications are described in greater detail herein.

"Percentage amino acid identity" refers to a comparison of the amino  
25 acids of two polypeptides as described herein.

As used herein, "aliphatic residue" means an organic radical having carbon atoms linked in open chains.

As used herein, "aromatic residue" means an organic radical that includes an aromatic ring (e.g., an aromatic group, an alkaryl group, or an aralkyl group).

30 As used herein, the "P1" position of a polypeptide is the amino acid on the N-terminal end of the scissile bond that is being cleaved. For dipeptidylpeptidases that cleave the peptide bond between the second and third

amino acids from the N-terminal end of a target polypeptide, the P1 position is the second amino acid from the N-terminal end of the target polypeptide (i.e., the penultimate position).

## 5 BRIEF DESCRIPTION OF THE FIGURES

Figure 1 depicts a plot of the absorbance at 280 nm ( $\Delta$ ) and amidolytic activity against Ala-Phe-pNA ( $\bullet$ ) for the purification of *P. gingivalis* dipeptidylpeptidase (DPP-7) from the acetone precipitate of the *P. gingivalis* cell extract. The straight solid lines indicate gradients in the eluting composition.

- 10 Figure 1(a) illustrates the separation of DPP-7 on hydroxyapatite (100 ml) equilibrated with 20 mM potassium phosphate buffer, pH 7.0, and using a potassium phosphate gradient from 20 mM to 300 mM. Figure 1(b) illustrates the separation of DPP-7 obtained from the previous step on Phenyl-Sepharose HP (25 ml) equilibrated with 50 mM potassium phosphate, 1M ammonium
- 15 sulfate, pH 7.0, at a flow rate of 30 ml/hour, and using an ammonium sulfate gradient from 0.4M to 0M. Figure 1(c) illustrates the separation of DPP-7 on a MonoS FPLC column using a sodium chloride gradient from 0M to 0.3M then from 0.3M to 1M.

- Figure 2 is a depiction of the SDS-PAGE of fractions obtained during the purification of *P. gingivalis* DPP-7 with *Lane A* representing molecular mass
- 20 markers (phosphorylase B, 97 kDa; bovine serum albumin, 68 kDa; ovalbumin, 43 kDa; carbonic anhydrase, 30 kDa; soybean trypsin inhibitor, 20 kDa;  $\alpha$ -lactalbumin, 14 kDa); *Lane B* representing acetone precipitate from Triton X-100 extract of *P. gingivalis*; *Lane C* representing hydroxyapatite column eluate;
- 25 *Lane D* representing Phenyl-Sepharose column eluate; and *Lane E* representing MonoS column eluate.

- Figure 3 depicts a plot of the DPP-7 activity against Ala-Phe-pNA vs. pH. Enzyme activity was tested on Ala-Phe-pNA substrate in different buffers including: HEPES ( $\bullet$ ); PIPES ( $\square$ ); potassium phosphate ( $\blacksquare$ ); Tris ( $\circ$ ); and MES
- 30 ( $\blacktriangle$ ).

Figure 4 depicts the coding sequence (SEQ ID NO:1) encoding *P. gingivalis* DPP-7 (SEQ ID NO:2). Sequences obtained from the Edman



degradation of the trypsin fragmented DPP-7 polypeptide chain are underlined. The putative active site serine residue is marked by the black background.

Figure 5 is a listing of sequences comparing the C-terminal regions of the *P. gingivalis* DPP-7 (residues 664-695; SEQ ID NO:3) and *S. aureus* V8 endopeptidase (residues 704-863; SEQ ID NO:4). Common residues are indicated by the single letter amino acid in the line between the two sequences. The "+" symbol in the line between the two sequences indicates similar residues.

Figure 6 depicts a multiple sequence alignment of *P. gingivalis* DPP-7 and its putative homologues. Sequences of DPP-7 related proteinases were obtained from the conceptual translation of the following ORFs retrieved from unfinished and finished genomes databases (available at [www.tigr.org](http://www.tigr.org)): S1- *Shewanella putrefaciens* gnl | TIGR\_24 | sputre 6401 (SEQ ID NO:5); S2- *Shewanella putrefaciens* gnl | TIGR\_24 | sputre 6410 (SEQ ID NO:6); X- *Xylella fastidiosa* gb | AE004008.1 | (SEQ ID NO:7); P1- *Porphyromonas* *gingivalis* gnl | TIGR | *P. gingivalis*\_CPG.con (SEQ ID NO:8); P2- *P. gingivalis* DPP-7 gnl | TIGR | *P. gingivalis*\_CPG.con (SEQ ID NO:9). The sequences were subsequently aligned using the ClustalW multiple sequence alignment tool.

## 20 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides isolated polypeptides, preferably isolated dipeptidylpeptidases, that have amidolytic activity by hydrolysis of a peptide bond present in a target polypeptide, wherein the bond is between the second and third amino acid from the N-terminus of the peptide. The dipeptidylpeptidase has amidolytic activity by hydrolysis of a peptide bond present in a target polypeptide including the fragment H-Xaa-Yaa-Xaa-, wherein Xaa is a natural or modified amino acid, Yaa is an amino acid including an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom, and the peptide bond of the target polypeptide that is hydrolyzed is the bond between the second and third amino acids from the N-terminus of the peptide. In increasing order of preference,

isolated polypeptides can cleave a target polypeptide that is at least about 5 amino acids or at least about 400 Da, at least about 10 amino acids or at least about 750 Da, at least about 20 amino acids or at least about 1,500 Da, or at least about 30 amino acids or at least about 3,000 Da. Preferably, the

- 5 dipeptidylpeptidases cleave peptides including a sequence of H-Xaa-Yaa-Xaa-, wherein Yaa is alanine, phenylalanine, isoleucine, or leucine. More preferably, the dipeptidylpeptidases cleave peptides including a sequence of SEQ ID NO:10, SEQ ID NO:11; SEQ ID NO:12; or SEQ ID NO:13 as shown in Table 3.

- The polypeptides disclosed in the present application are preferably
- 10 dipeptidylpeptidases. Preferably, the dipeptidylpeptidase is isolated from *Porphyromonas gingivalis*. Preferably, the dipeptidylpeptidase is a serine protease. Most preferably, the dipeptidylpeptidase is *P. gingivalis* dipeptidylpeptidase-7 (DPP-7). The polypeptides can be used as a source of antibodies for inhibiting the amidolytic activity and thereby possibly reducing
- 15 periodontitis, loss of tooth attachment and periodontal pocket formation. Antibodies to dipeptidylpeptidases can also be used to identify and/or isolate additional dipeptidylpeptidases. Knowledge of dipeptidylpeptidases can also be used to make inhibitors of dipeptidylpeptidases and to make immunogenic compositions that could be used to elicit the production of antibodies to
- 20 dipeptidylpeptidases and thereby possibly reduce gingivitis, periodontitis, loss of tooth attachment, and/or periodontal pocket formation.

- Dipeptidylpeptidase-7, either alone or in a mixture with other dipeptidylpeptidases, can be used to generate a pool of dipeptides from polypeptides. Dipeptides may be preferably imported by cells. Thus, pools of
- 25 dipeptides might be useful substrates for transport.

- Dipeptidylpeptidase-7 (DPP-7) was purified from the membrane fraction of *Porphyromonas gingivalis*. This enzyme, preferably having an apparent molecular mass of about 76 kDa, has specificity for polypeptides having either an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the
- 30 second amino acid from the N-terminal end of the polypeptide. Although it belongs to the serine class of peptidases, it does not resemble other known dipeptidylpeptidases. Interestingly, the amino acid sequence around the putative

active site serine residue shows significant homology to the C-terminal region of the *Staphylococcus aureus* V-8 endopeptidase. In *P. gingivalis*, DPP-7 probably serves nutritional functions by providing dipeptides to this assaccharolytic bacterium.

5           Several studies indicate that the outer membrane of *P. gingivalis* contains a complex, proteolytic machinery which serves multiple physiological functions. The present application discloses the identification of a novel proteinase localized on the bacterial surface.

10           The purified enzyme migrated as a single band of about 76 kDa on SDS-PAGE and its amino-terminal sequence was located within the primary structure of the translated product of the *dpp-7* coding sequence. Apparently, the enzyme is truncated at the amino terminus (i.e., amino acid 24 of SEQ ID NO:2 is the first amino acid of the truncated form) due to the action of a lysine specific proteinase, most likely gingipain K. Taking into account that the N-terminus of  
15   DPP-7 contains membrane anchorage domains it is likely that the N-terminal truncation noted here occurred during the isolation procedure and may not represent its true membrane form.

          The dipeptidylpeptidases of the present invention are preferably serine proteases that are inhibited by serine protease inhibitors. The  
20   dipeptidylpeptidases of the present invention are preferably inhibited by serine protease inhibitors including, for example, diisopropylfluorophosphate (DFP), 4-(2-aminoethyl)-benzenesulfonyl fluoride hydrochloride (PEFABLOCK), and phenylmethanesulfonyl fluoride (PMSF). In addition, the dipeptidylpeptidases of the present invention are resistant to sulfhydryl group blocking reagents and  
25   chelating agents, which is also consistent with the enzyme being a serine protease. However, the *P. gingivalis* DPP-7 does not belong to any of the six previously described types of dipeptidylpeptidases (Barrett et al., *Handbook of Proteolytic Enzymes*, Academic Press, London (1998)). DPP-I is a member of a cysteine class of peptidases and possesses a broad specificity, but has an  
30   exclusion for basic amino acid and proline residues in the P1 site of the scissile peptide bond (McGuire et al., *Arch. Biochem. Biophys.*, 295:280-88 (1992)). DPP-VI is another representative of the cysteine proteinase family with

dipeptidylpeptidase activity towards broad spectrum of substrates (Vacheron et al., *Eur. J. Biochem.*, 100:189-96 (1979)). DPP-II, DPP-IV and DPP-V belong to the S9 family of the serine proteases (Barrett et al., *Handbook of Proteolytic Enzymes*, Academic Press, London (1998)). Both DPP-II and DPP-IV share  
5 similar specificity directed against Pro and Ala residues in the penultimate position whereas DPP-V is an enzyme secreted by *Aspergillus fumigatus* with a unique substrate specificity limited to X-Ala, His-Ser, and Ser-Tyr dipeptides (Beauvais et al., *J. Biol. Chem.*, 272:6238-44 (1997)). DPP-III is also classified as a serine peptidase, with its action being restricted to Arg residue in the P1  
10 position (Ellis et al., *J. Biol. Chem.*, 242:4623-29 (1967)). In terms of biochemical features, DPP-7 resembles a dipeptidyl aminopeptidase (DAP-BII), which was isolated from *Pseudomonas sp.* strain WO24, but the coding sequence of that enzyme remains unknown and does not allow a sequence comparison of these proteins (Ogasawara et al., *J. Bacteriol.*, 178:6288-95  
15 (1996)). Because *P. gingivalis* dipeptidylpeptidase does not exhibit any significant homology to any of the dipeptidylpeptidases described so far, this enzyme has been designated DPP-7.

Interestingly the *P. gingivalis* DPP-7 displays the consensus sequence characteristic for the catalytic site of the V-8 like proteases, a group of  
20 endopeptidases cleaving after glutamic or aspartic acid residues (Carmona et al., *Nucleic Acids Res.*, 15:6757 (1987)). This region of homology is specifically located only at the C-terminal region of both proteases and includes the putative active site serine residue. Interestingly, more coding sequences encoding putative, DPP-7 related proteases in *P. gingivalis*, *Xylella fastidiosa* and  
25 *Shewanella putrefaciens* were identified. Based on the enzymological and coding sequence data presented above, the *P. gingivalis* DPP-7 does not belong to any of the peptidase families previously reported and should, therefore, be regarded as a prototype enzyme that defines a new family of dipeptidylpeptidases.

30 The invention further includes a polypeptide, preferably a dipeptidylpeptidase, that shares a significant level of primary structure (referred to as "percent identity") with SEQ ID NO:2. The level of identity is determined

by aligning the two amino acid sequences (i.e., the amino acid sequence of the polypeptide and the sequence SEQ ID NO:2) such that the residues that make up the putative active site sequence (e.g., about amino acid 644 to about 653, preferably about amino acid 644 to about 658) are in register, then further  
5 aligned to maximize the number of amino acids that they have in common along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to place the residues that make up the putative active site sequence (e.g., about amino acid 644 to about 653, preferably about amino acid 644 to about 658) in register and to maximize the number of shared  
10 amino acids, although the amino acids in each sequence must nonetheless remain in their proper order. Preferably, two amino acid sequences are compared using the blastp program of the BLAST search algorithm, which is described by Altshul et al., (*Nucl. Acids Res.*, 25, 3389-3402 (1997)), and available at the National Center for Biotechnology Information (e.g.,  
15 [www.ncbi.nlm.nih.gov/Microb\\_blast/unfinishedgenome.html](http://www.ncbi.nlm.nih.gov/Microb_blast/unfinishedgenome.html) or [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). Preferably, the default values for all BLAST search parameters are used. In the comparison of two amino acid sequences using the BLAST search algorithm, structural similarity is referred to as "identities." Preferably, a dipeptidylpeptidase has, in increasing order of  
20 preference, at least about 40% identity, at least about 50% identity, at least about 60% identity, at least about 70% identity, at least about 80% identity, and most preferably at least about 90% identity with SEQ ID NO:2. Preferably, about amino acid 543 to about 699 of SEQ ID NO:2 are used, more preferably about amino acid 71 to about 712 of SEQ ID NO:2 are used. Preferably the invention  
25 includes an isolated polypeptide including an amino acid sequence having a percentage amino acid identity of greater than about 40% with SEQ ID NO:2.

In general, the amidolytic activity of the polypeptides of the invention, preferably dipeptidylpeptidases, can be measured by assay of the cleavage of a target polypeptide in the presence of dipeptidylpeptidase and a buffer.  
30 Preferably, the ratio of dipeptidylpeptidase to target polypeptide is at least about 1:1; more preferably at least about 1:100; even more preferably at least about 1:1,000; and most preferably at least about 1:10,000. Preferably, the ratio of

dipeptidylpeptidase to target polypeptide is at most about 1:10,000,000; more preferably at most about 1:1,000,000; and most preferably at most about 1:100,000. Buffers in which a dipeptidylpeptidase is active are suitable for the assay. Preferably, the buffer is at most about 200 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid), more preferably at most about 50 mM HEPES, and most preferably at most about 20 mM HEPES. Preferably, the pH of the buffer is at least about pH 6.0. Preferably, the pH of the buffer is at most about pH 8.0 and more preferably at most about pH 7.5. Preferably, the temperature of the assay is about 37°C. The assay can be carried out for at least about 1 minute to at most about 24 hours. Preferably, the amidolytic activity of the dipeptidylpeptidases are measured at a dipeptidylpeptidase:target polypeptide ratio of at least about 1:100 and at most about 1:1,000,000 in about 200 mM HEPES, about pH 7.5 at about 37°C for at least about 3 hours. In general, the time of the assay can vary depending on the substrate and enzyme:substrate ratio. Typically, target polypeptides are stable under these conditions, and typically it is difficult to detect background levels of hydrolysis in the absence of a dipeptidylpeptidase. Preferably, the assay is allowed to continue until at least about 1% of the target polypeptide is hydrolyzed.

Dipeptidylpeptidases of the present invention are preferably inhibited by compounds including, for example, 4-(2-aminoethyl)-benzenesulfonyl fluoride hydrochloride (PEFABLOCK); diisopropylfluorophosphate (DFP); phenylmethanesulfonyl fluoride (PMSF); 3,4-dichlorisocoumarin; and combinations thereof. The peptidases of the present invention are preferably not inhibited by a compounds including, for example, specific inhibitors of metallo peptidases, cysteine peptidases, and aspartic peptidases.

An active analog, active fragment, or active modification of a polypeptide including the amino acid sequence SEQ ID NO:2 is one that has amidolytic activity by hydrolysis of the target polypeptide described above. Active analogs of a polypeptide including the amino acid sequence SEQ ID NO:2 include dipeptidylpeptidases having amino acid substitutions that do not eliminate hydrolysis of the target polypeptide at the peptide bond between the second and third amino acids. Substitutes for an amino acid may be selected

from other members of the class to which the amino acid belongs. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, and tryptophan. Polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine and glutamine. The

5 positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Examples of preferred conservative substitutions include Lys for Arg and *vice versa* to maintain a positive charge; Glu for Asp and *vice versa* to maintain a negative charge; Ser for Thr so that a free -OH is maintained; and Gln for Asn

10 to maintain a free NH<sub>2</sub>.

Active fragments of a dipeptidylpeptidase of the invention include dipeptidylpeptidases containing deletions or additions of one or more contiguous or noncontiguous amino acids such that the resulting polypeptide will hydrolyze the target polypeptide at the bond between the second and third amino acids.

15 Modified dipeptidylpeptidases include dipeptidylpeptidases that are chemically and enzymatically derivatized at one or more constituent amino acid, including side chain modifications, backbone modifications, and N- and C- terminal modifications including acetylation, hydroxylation, methylation, amidation, and the attachment of carbohydrate or lipid moieties, cofactors, and the like.

20 Modified dipeptidylpeptidases will hydrolyze the target polypeptide at the peptide bond between the second and third amino acids.

Preferably, a dipeptidylpeptidase includes the sequence TGGNSGSPV (SEQ ID NO:26), and more preferably includes the consensus sequence for the active-site serine residue of serine type proteases, TGGNSGSPVF (SEQ ID

25 NO:25), where T is Threonine, G is glycine, N is Asparagine, P is Proline, V is valine, F is Phenylalanine, and S is serine, with the putative active site serine being underlined. The active site serine can be identified by, for instance, labeling with diisopropylfluorophosphate as described herein. Preferably, the catalytic domain of the dipeptidylpeptidases of the invention begins at about

30 residue 543 of SEQ ID NO:2 and includes the remaining 169 amino acids, more preferably begins at about residue 540 of SEQ ID NO:2 and includes the

remaining 172 amino acids, and most preferably begins at about residue 522 of SEQ ID NO:2 and includes the remaining carboxy-terminal amino acids.

Dipeptidylpeptidases can be obtained by several methods. Isolation of a dipeptidylpeptidase present on the surface of a cell producing the peptidase typically requires lysis of the cell followed by purification methods that are well known in the art. Alternatively, cells can be treated with a detergent, for instance Triton X-100, to remove the peptidase from the cell surface. The following are nonlimiting examples of suitable protein purification procedures: fractionation on immunoaffinity, ion-exchange, hydroxyapatite, Phenyl-Sephadex HP, MonoQ HR 5/5, or MonoP columns; ethanol precipitation; reverse phase HPLC; chromatography on silica or on an ion-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75. Preferably, isolation of a dipeptidylpeptidase from *P. gingivalis* is accomplished using a combination of hydroxyapatite, Phenyl-Sephadex HP, MonoS HR 5/5 and MonoP column chromatography steps as described herein.

Dipeptidylpeptidases can also be isolated from organisms other than *P. gingivalis*. Other organisms can express a dipeptidylpeptidase that is encoded by a coding region having similarity to the coding region encoding SEQ ID NO:2. A "coding region," a "coding sequence," or an "open reading frame" (ORF) is a linear form of nucleotides that encodes a polypeptide, usually via mRNA, when placed under the control of appropriate regulatory sequences. The boundaries of a coding region are generally determined by a translation start codon at its 5' end and a translation stop codon at its 3' end. "Regulatory region" refers to a nucleic acid that regulates expression of a coding region to which a regulatory region is operably linked. Non limiting examples of regulatory regions include promoters, transcription initiation sites, translation start sites, translation stop sites, and terminators. "Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A regulatory element is "operably linked" to a coding region when it is joined in such a way that expression of the coding region is achieved under conditions compatible with the regulatory region. Alternatively,



other organisms can express a dipeptidylpeptidase from a recombinant coding region encoding the peptidase. The identification of similar coding regions in other organisms can be accomplished as described herein. A dipeptidylpeptidase can be isolated using purification methods that are well known in the art.

- 5 Alternatively, the peptidase can be chemically synthesized using methods that are well known in the art including, for instance, solid phase synthesis. Examples of, for instance, coding and regulatory regions are described herein.

- The expression of a dipeptidylpeptidase by an organism other than *P. gingivalis* can be detected using specific substrates of the general formula
- 10 Xaa-Xaa-LG, wherein Xaa represents any natural amino acid and LG is a leaving group. The leaving group can be a chromogenic or fluorogenic group known to the art. The expression of a dipeptidylpeptidase by an organism and subsequent cleavage of a specific substrate results in a free amino acid or a free leaving group, each of which can be assayed using techniques known to those of
- 15 skill in the art. Other methods can be based on immunogenic properties of DPP-7, for instance immunoassays and histochemistry, the detection of mRNA, and PCR related methods, all of which are known to one of skill in the art.

- In one aspect, the present invention is directed to a nucleic acid encoding a polypeptide, particularly a dipeptidylpeptidase, active analog, active fragment,
- 20 or active modification thereof. The nucleic acid can have a nucleotide sequence as shown in SEQ ID NO:1. Alternatively, nucleic acids of the invention include those whose complement hybridize to SEQ ID NO:1 under standard hybridization conditions as described herein. During hybridization the entire nucleotide sequence of the complement can hybridize with SEQ ID NO:1.
- 25 Preferably, at least about 20 nucleotides of the complement hybridize with SEQ ID NO:1, more preferably at least about 50 nucleotides, most preferably at least about 100 nucleotides.

- The identification of similar coding regions in other organisms can be accomplished by screening individual wild-type microorganisms for the
- 30 presence of nucleotide sequences that are similar to the coding region of DPP-7, which is shown in SEQ ID NO:1. Screening methods include, for instance, hybridization of a detectably labeled probe with a nucleic acid.

Standard hybridizing conditions are a modification of the conditions used by Church et al. ((1984) *Proc. Natl. Acad. Sci. USA* 81, 1991): 0.5 M phosphate buffer, pH 7.2, 7% SDS, 10 mM EDTA, at 68°C, and three washes, each for 20 minutes in 2x SSC (preferably 0.1 SSC), 0.1% SDS, at 65°C. Preferably, a probe will hybridize to the nucleotide sequence set forth in SEQ ID NO:1 under standard hybridizing conditions. Generally the probe does not have to be complementary to all the nucleotides of the nucleic acid as long as there is hybridization under the above-stated conditions.

“Complement” and “complementary” refer to the ability of two single stranded nucleic acids to base pair with each other, where an adenine on one nucleic acid will base pair to a thymine on a second nucleic acid and a cytosine on one nucleic acid will base pair to a guanine on a second nucleic acid. Two nucleic acids are complementary to each other when a nucleotide sequence in one nucleic acid can base pair with a nucleotide sequence in a second nucleic acid. For instance, 5'-ATGC and 5'-GCAT are complementary. The term complement and complementary also encompasses two nucleic acids where one nucleic acid contains at least one nucleotide that will not base pair to at least one nucleotide present on a second nucleic acid. For instance the third nucleotide of each of the two nucleic acids 5'-ATTGC and 5'-GCTAT will not base pair, but these two nucleic acids are complementary as defined herein. Typically two nucleic acids are complementary if they hybridize under the standard conditions referred to herein.

Preferred probes are nucleic acids complementary to a coding region or another nucleotide sequence that encodes a dipeptidylpeptidase. For instance, a probe can include a consecutive series of nucleotides complementary to a portion of SEQ ID NO:1. Preferably a probe is at least about 18 bases, more preferably at least about 21 bases, and most preferably at least about 24 bases in length. One of skill in the art could select useful probes as desired. Methods of detectably labeling a probe are well known to the art.

The nucleic acid that is identified by the probe is further analyzed to determine if it encodes a polypeptide with amidolytic activity of the peptide bond between the second and third amino acids from the N-terminus on a target polypeptide of the general formula H-Xaa-Yaa-Xaa, wherein Xaa is a natural or modified amino acid and Yaa is an amino acid including an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom. Another method for screening individual microorganisms for the presence of nucleotide sequences that are similar to the coding regions of the present invention is the polymerase chain reaction (PCR).

Individual wild-type microorganisms containing nucleic acids encoding a dipeptidylpeptidase can also be identified using antibody. Preferably the antibody is directed to DPP-7. The production of antibodies to a particular polypeptide is known to a person of skill in the art, and is further detailed herein.

The use of hybridization of a probe to a coding region present in individual wild-type microorganisms can be used as a method to identify a coding region identical or similar to a coding region present in SEQ ID NO:1. The coding region can then be isolated and ligated into a vector as described below.

The present invention is also directed to coding regions sharing a significant level of primary structure with the coding region present at SEQ ID NO:1. The level of identity is determined by aligning the two nucleotide sequences (i.e., the nucleotide sequence of the polynucleotide and the sequence SEQ ID NO:1) such that the residues that encode the putative active site of the encoded protein (e.g., about nucleotide 1929 to about 1974) are in register, then further aligned to maximize the number of nucleotides that they have in common along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to place the residues that encode the putative active site of the encoded protein (e.g., about nucleotide 1929 to about 1974) in register and to maximize the number of shared nucleotides, although the nucleotides in each sequence must nonetheless remain in their proper order. Preferably, two nucleotide sequences are compared using the blastn program of the BLAST search algorithm, which is described by Altshul et al., (*Nucl. Acids*

Res., 25, 3389-3402 (1997)), and available at the National Center for Biotechnology Information (e.g., [www.ncbi.nlm.nih.gov/Microb\\_blast/unfinishedgenome.html](http://www.ncbi.nlm.nih.gov/Microb_blast/unfinishedgenome.html) or [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). Preferably, the default values for all BLAST search parameters are used. In the comparison of two nucleotide sequences using the BLAST search algorithm, structural similarity is referred to as "identities." Preferably, two nucleotide acid sequences have, in increasing order of preference, preferably at least about 70%, at least about 80%, at least about 90%, at least about 95%, and most preferably at least about 95% identity.

As mentioned above, a nucleic acid of the invention can be inserted in a vector. Construction of vectors containing a nucleic acid of the invention employs standard ligation techniques known in the art. See, e.g., Sambrook et al, *Molecular Cloning: A Laboratory Manual.*, Cold Spring Harbor Laboratory Press (1989) or Ausubel, R.M., ed. *Current Protocols in Molecular Biology* (1994). A vector can provide for further cloning (amplification of the nucleic acid), i.e., a cloning vector, or for expression of the polypeptide encoded by the coding region, i.e., an expression vector. The term vector includes, but is not limited to, plasmid vectors, viral vectors, cosmid vectors, or artificial chromosome vectors. Typically, a vector is capable of replication in a bacterial host, for instance *E. coli*. Preferably the vector is a plasmid.

Selection of a vector depends upon a variety of desired characteristics in the resulting construct, such as a selection marker, vector replication rate, and the like. Suitable plasmids for expression in *E. coli*, for example, include pUC(X), pKK223-3, pKK233-2, pTrc99A, and pET-(X) wherein (X) denotes a vector family in which numerous constructs are available. pUC(X) vectors can be obtained from Pharmacia Biotech (Piscataway, NH) or Sigma Chemical Co. (St. Louis, MO). pKK223-3, pKK233-2 and pTrc99A can be obtained from Pharmacia Biotech. pET-(X) vectors can be obtained from Promega (Madison, WI) Stratagene (La Jolla, CA) and Novagen (Madison, WI). To facilitate replication inside a host cell, the vector preferably includes an origin of replication (known as an "ori") or replicon. For example, ColE1 and P15A replicons are commonly used in plasmids that are to be propagated in *E. coli*.

An expression vector optionally includes regulatory regions operably linked to the coding region. The invention is not limited by the use of any particular promoter, and a wide variety are known. Promoters act as regulatory signals that bind RNA polymerase in a cell to initiate transcription of a downstream (3' direction) coding region. The promoter used in the invention can be a constitutive or an inducible promoter. It can be, but need not be, heterologous with respect to the host cell. Preferred promoters for bacterial transformation include *lac*, *lacUV5*, *tac*, *trc*, T7, SP6 and *ara*.

An expression vector can optionally include a Shine Dalgarno site (e.g., a ribosome binding site), and a start site (e.g., the codon ATG) to initiate translation of the transcribed message to produce the enzyme. It can also include a termination sequence to end translation. A termination sequence is typically a codon for which there exists no corresponding aminoacyl-tRNA, thus ending polypeptide synthesis. The nucleic acid used to transform the host cell can optionally further include a transcription termination sequence. The *rrnB* terminators, which is a stretch of DNA that contains two terminators, T1 and T2, is an often used terminator that is incorporated into bacterial expression systems (J. Brosius et al., (1981) *J. Mol. Biol.* 148 107-127).

The nucleic acid used to transform the host cell optionally includes one or more marker sequences, which typically encode a polypeptide that inactivates or otherwise detects or is detected by a compound in the growth medium. For example, the inclusion of a marker sequence can render the transformed cell resistant to an antibiotic, or it can confer compound-specific metabolism on the transformed cell. Examples of a marker sequence are sequences that confer resistance to kanamycin, ampicillin, chloramphenicol, and tetracycline.

Antibodies to a polypeptide including the sequence SEQ ID NO:2 can be produced. Alternatively, antibodies to an antigenic analog, antigenic fragment, or antigenic modification of a polypeptide including the sequence SEQ ID NO:2 can be made. An antigenic analog, antigenic fragment, or antigenic modification of a polypeptide including the amino acid sequence SEQ ID NO:2 is one that generates an immune response in an animal. Preferably, an antigenic analog, antigenic fragment, or antigenic modification has amidolytic activity. Antigenic

analogs of a polypeptide including the amino acid sequence SEQ ID NO:2 include dipeptidylpeptidases having amino acid substitutions that do not eliminate peptide antigenicity in an animal. Substitutes for an amino acid may be selected from other members of the class to which the amino acid belongs, as  
5 described herein. Fragments of a dipeptidylpeptidase of the invention include dipeptidylpeptidases containing deletions or additions of one or more contiguous or noncontiguous amino acids such that the resulting polypeptide will generate an immune response in an animal. Modified dipeptidylpeptidases include dipeptidylpeptidases that are chemically and enzymatically derivatized at one or  
10 more constituent amino acids, including side chain modifications, backbone modifications, and N- and C- terminal modifications including acetylation, hydroxylation, methylation, amidation, and the attachment of carbohydrate or lipid moieties, cofactors, and the like.

Accordingly, an aspect of the invention is an immunogenic composition  
15 including an isolated dipeptidylpeptidase, an antigenic analog, antigenic fragment, or antigenic modification thereof. The dipeptidylpeptidase preferably has amidolytic activity for cleavage of the target polypeptide described herein.

The immunogenic composition can further include excipients or diluents that are pharmaceutically acceptable as carriers and compatible with the  
20 immunogenic composition. The term "pharmaceutically acceptable carrier" refers to a carrier(s) that is "acceptable" in the sense of being compatible with the other ingredients of a composition and not deleterious to the recipient thereof. Suitable excipients include, for example, water, saline, dextrose, glycerol, ethanol, and combinations thereof. In addition, if desired, the  
25 immunogenic composition may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the immune-stimulating composition.

The immunogenic composition can be used in a method for protecting an animal from a disease caused by *P. gingivalis*. This method includes  
30 administering the immunogenic composition and eliciting antibodies to a dipeptidylpeptidase, antigenic analog, antigenic fragment, or antigenic modification. The diseases that can be treated in this manner include

periodontal diseases, which include gingivitis and periodontitis. Clinical hallmarks of periodontitis include loss of tooth attachment and periodontal pocket formation.

Alternatively and preferably, periodontal diseases can be treated by the use of inhibitors of a dipeptidylpeptidase. An inhibitor of a dipeptidylpeptidase can be present in a composition that preferably contains a pharmaceutically acceptable carrier. For instance, inhibitors can be applied systemically, subgingivally (e.g., subgingival irrigation), and/or by controlled release delivery directly into the periodontal pocket using methods well known in the art (see, e.g., Kornman, *J. Periodontol.* 64:782-91 (1993). Preferably, an inhibitor is applied subgingivally or by controlled release delivery.

The dipeptidylpeptidases, active analogs, active fragments, and active modifications thereof can be used in a method of reducing growth of bacteria *in vitro* or *in vivo*. Preferably, the bacteria is a periodontal pathogen, i.e., a bacterial pathogen that causes periodontal disease, more preferably the bacteria is *P. gingivalis*. The inability of asaccharolytic *P. gingivalis* to utilize free amino acids makes the bacterium entirely dependant on an external peptide supply. The action of the polypeptides of the invention may be required for bacterial growth, and inhibition of the polypeptides of the invention may inhibit the *in vivo* growth of organisms, including *P. gingivalis*. The method includes decreasing the amount of dipeptides (e.g., the result of cleavage of the target polypeptide by a dipeptidylpeptidase) and the amount of free amino acids that result from further cleavage of the dipeptides present by inhibiting a dipeptidylpeptidase, active analog, active fragment, or active modification thereof, such that the amount of dipeptides generated by the polypeptides is decreased. The amount of dipeptides is decreased relative to the amount of dipeptides present in the absence of the inhibitor. Preferably, the amount of dipeptides generated is decreased by an inhibitor, a monoclonal antibody that inhibits the dipeptidylpeptidase, or polyclonal antibodies that inhibit the dipeptidylpeptidase, more preferably, the amount of dipeptides generated is decreased by an inhibitor. Preferably, an inhibitor acts to inhibit a polypeptide of the invention, preferably a dipeptidylpeptidase, by blocking the active site of the

polypeptide. The polypeptide can be present on the surface of the bacteria or secreted into the environment, preferably the polypeptide is present in the surface of the bacteria.

The present invention is also directed to a method of developing an inhibitor of a dipeptidylpeptidase, active analog, active fragment, or active modification thereof, preferably a dipeptidylpeptidase. The method includes identifying a compound that inhibits the amidolytic activity of the dipeptidylpeptidase. Such compounds include, for example, polypeptides, organic compounds, inorganic compounds, metals, non-ribosomal polypeptides, polyketides, and peptidomimetics. The identification of compounds can be accomplished by, for instance, incubating the dipeptidylpeptidase with a candidate compound under conditions that promote amidolytic activity of the dipeptidylpeptidase and determining if the amidolytic activity of the dipeptidylpeptidase is decreased relative to the amidolytic activity in the absence of the compound. The amidolytic activity can be measured by cleavage of the peptide bond between the second and third amino acids of the target polypeptide as described herein. One method of developing an inhibitor includes using the target polypeptide and replacing the Xaa residues with modified amino acids. It is expected that some modified amino acids will cause the target polypeptide to act as an inhibitor.

The present invention is illustrated by the following examples. It is to be understood that the particular examples, materials, amounts, and procedures are to be interpreted broadly in accordance with the scope and spirit of the invention as set forth herein.

## EXAMPLES

### *Materials*

Diisopropylfluorophosphate (DFP), leupeptin and 3,4-dichloroisocoumarin, were purchased from Calbiochem (La Jolla, CA). Other peptides used in this study were synthesized at the Molecular Genetic Instrumental Facility (University of Georgia, Athens, GA) using Fmoc protocol



with an advanced ChemTech MPS350 automated synthesizer. The peptides H-Xaa-Xaa-pNA and Z-Xaa-Xaa-pNA, where pNA is p- Nitroanilide; Z is benzyloxycarbonyl; and H is hydrogen and denotes an unblocked amino-terminal group were obtained from Bachem (King of Prussia, PA).

5

### *Methods*

#### EXAMPLE 1

*Source and Cultivation of Bacteria* - *P. gingivalis* DPP-7 was purified from strain HG66, a kind gift of Dr. Roland Arnold (University of North Carolina, Chapel Hill, NC). The cells were grown as described previously (Chen et al., *J. Biol. Chem.*, 267:18896-901 (1992)).

*Protein Determination* - Protein concentration was determined with the BCA reagent kit (Sigma), using bovine serum albumin as a standard.

*Localization of Dipeptidylpeptidase Activity* - The localization of active enzyme was checked in bacterial cells that had been subjected to a previously described fractionation procedure (Banbula et al., *Infect. Immun.*, 68:1176-82 (2000)). All fractions, as well as the full culture, culture medium, and full culture after sonication, were assayed for amidolytic activity against H-Ala-Phe-pNA.

*Enzyme Purification* - All purification steps were performed at 4°C except for FPLC separations, which were carried out at room temperature. The cells were collected by centrifugation (6,000 x g, 30 minutes) and resuspended in 50 mM potassium phosphate buffer, pH 7.4. The outer membrane proteins were solubilized with 0.05% Triton X-100. After 2 hours of gentle stirring, unbroken cells were removed by centrifugation (28,000-x g, 60 minutes). Proteins from the supernatant were precipitated with cold acetone (60% final concentration), collected by centrifugation, and redissolved in 50 mM potassium phosphate buffer, pH 7.0. After extensive dialysis against the same buffer the sample was loaded onto a hydroxyapatite column (BioRad) previously equilibrated with 20 mM potassium phosphate, pH 7.0, at a flow rate of 20 ml/hour. The column was then washed until the  $A_{280}$  fell to zero. Bound proteins were eluted with a potassium phosphate gradient (20-300 mM) and fractions (7 ml) were analyzed

for amidolytic activity against H-Ala-Phe-pNA. The active fractions were saturated with 1 M ammonium sulfate and loaded onto a Phenyl-Sepharose HP column (Pharmacia) equilibrated with 50 mM potassium phosphate, pH 7.0, containing 1M ammonium sulfate. The column was washed with two volumes  
5 of the equilibration buffer, followed by a wash with buffer containing 0.4 M ammonium sulfate, and developed with a descending gradient of ammonium sulfate from 0.4 to 0 M. Active fractions were pooled, extensively dialyzed against 20 mM MES, pH 6.6, and applied onto a MonoS HR 5/5 FPLC (Pharmacia) column equilibrated with the same buffer. Bound proteins were  
10 eluted with a 0-300 mM NaCl gradient. This allowed us to obtain a homogenous preparation of active proteinase.

*Electrophoretic Techniques* - The SDS-PAGE system of Schagger and von Jagow (Schagger et al., *Anal. Biochem.*, 166:368-79 (1987)), was used to monitor enzyme purification and estimate the enzyme molecular mass. For  
15 amino-terminal sequence analysis, proteins resolved in SDS-PAGE were electroblotted onto polyvinylidene difluoride membranes using 10 mM CAPS, pH 11, 10% methanol (Matsudaira et al., *J. Biol. Chem.*, 262:10035-38 (1987)). After staining with Coomassie Blue G250 the blot was air dried, and protein bands cut out and subjected to amino-terminal sequence analysis with an  
20 Applied Biosystems 491 Protein Sequencer using the program designed by the manufacturers.

*Kinetic Analysis* - Routinely, the dipeptidylpeptidase amidolytic activity was measured with H-Ala-Phe-pNA (1mM) in 0.2 M HEPES, pH 7.8 at 37°C. The reaction was followed for specific time intervals in a thermostated ELISA  
25 reader (SpectraMax, Applied Biosystem) and the release of p-nitroaniline was monitored at 405 nm. Other p-nitroanilide substrates were used in the same manner. For inhibition studies, the enzyme was first preincubated with an inhibitor for 15 minutes at 37°C, substrate added, and residual activity recorded. The initial steady-state velocity ( $v_0$ ) was determined by continuous assay for the  
30 range of substrate concentrations (100 nM to 1 mM).  $K_m$  and  $V_{max}$  were determined by hyperbolic regression of the kinetic data using the software

package Hyper Version 1.02 obtained from Dr. J. S. Easterby (University of Liverpool, UK).

*Enzyme Fragmentation* - The purified dipeptidylpeptidase was subjected to in-gel tryptic digestion (Rosenfeld et al., *Anal. Biochem.*, 203:173-79 (1992)).

5 Peptides were extracted and separated by microbore reverse-phase HPLC. Fractions absorbing at 210 nm were manually collected, and their masses were determined by reflectron MALDI-TOF mass spectrometry using a Bruker Daltonics ProFlex instrument as described previously (Pohl et al., *Lett. Peptide Sci.*, 1:291-97 (1995)). Selected peptides were subjected to Edman degradation  
10 in a model Procise-cLS sequencer (PE Biosystems, CA).

*Identification of the DPP-7 Coding Sequence* - An unfinished *P. gingivalis* W83 genome database, available from the Institute for Genomic Research, was searched for the presence of nucleotide sequences corresponding to the amino-terminal and the internal DPP-7 amino acid sequences using the  
15 TBLASTN algorithm (Altschul et al., *Nucleic Acids Res.*, 25:3389-402 (1997)). An identified contig gnl | TIGR | *P. gingivalis*\_1208 was retrieved from the Institute for Genomic Research database. The position of the DPP-7 coding sequence was localized using the National Center for Biotechnology Information (NCBI) open reading frame (ORF) finder and the amino acid sequence, obtained  
20 by conceptual translation of the entire ORF, was further used for homology screening by use of the NCBI BLAST search tool.

*Enzyme Specificity* - The determination of substrate specificity was based on the separation of the products of peptide hydrolysis by reverse-phase chromatography. Peptides were first incubated with 1 microgram of DPP-7 at an  
25 enzyme: substrate molar ratio of 1:100 for 3 hours or 24 hours in 50 microliters of 200 mM HEPES, 100 mM NaCl pH 8.0, at 37°C, and the reaction stopped by acidification with trifluoroacetic acid. The samples were then subjected to reverse-phase high pressure liquid chromatography using a Supelcosil LC 18 column (Supelco) with an acetonitrile gradient 0-60% in 0.075% trifluoroacetic  
30 acid in 50 minutes. Each peak, detected at 210 nm, was collected, lyophilized, re-dissolved in 50% (v/v) methanol, 0.1% acetic acid and subjected to analysis by mass spectrometry.

## EXAMPLE 2

A 76 kDa dipeptidylpeptidase associated with *P. gingivalis* membranes was solubilized by mild detergent treatment. This procedure released more than 5 90% of the amidolytic activity against H-Ala-Phe-pNA into the medium. After acetone precipitation and subsequent chromatography steps including the use of hydroxyapatite, Phenyl-Sepharose and MonoS columns (Fig. 1) a pure enzyme preparation was obtained. The homogeneity of the preparation and molecular mass of the protein were checked both by SDS PAGE (Fig. 2) and gel filtration 10 on a TSK G3000 SW column.

## EXAMPLE 3

*Inhibition Profile* - Based on the inhibition studies (Table I), DPP-7 was classified as a serine protease. DPP-7 was inactivated by 15 diisopropylfluorophosphate, PEFABLOCK and 3,4-dichloroisocoumarin, but not by typical cysteine class inhibitors such as E-64 or iodoacetic acid. Metal chelators including EDTA and 1,10-orthophenanthroline, as well as reducing agents did not influence its activity. The enzyme was not sensitive to inactivation by either detergents (0.5% SDS, 1% Triton X-100) or heavy metal 20 ions including  $Zn^{2+}$ ,  $Co^{2+}$  and  $Ni^{2+}$ . Human plasma inhibitors, such as  $\alpha_1$ -proteinase inhibitor,  $\alpha_1$ -antichymotrypsin, and  $\alpha_2$ -macroglobulin, did not effect enzyme activity nor were they cleaved by DPP-7.

TABLE I: Effect of different compounds on *P. gingivalis* DPP-7 activity.

Inhibitor	Concentration	% of residual activity
Diisopropylfluorophosphate	10 mM	34
PEFABLOCK	4 mg/ml	1
3,4-dichloroisocoumarin	2 mM	0
E-64	1 micromolar	96
Iodoacetic acid	0.1 mM	102
EDTA	10 mM	90
1,10-orthophenanthroline	1 mM	98

Leupeptin	0.1 mM	107
Aprotinin	0.5 mg/ml	128
Pepstatin	0.5 mg/ml	127
Cysteine	10 mM	90
Gly-Ala	100 mM	102
Arg-Phe	100 mM	69
Ala-Gly	100 mM	96
Arg-Gly	10 mM	84
Lys-Gly	10 mM	96
Ni <sup>++</sup>	1 mM	95
Zn <sup>++</sup>	1 mM	95
Co <sup>++</sup>	1 mM	116
SDS	0.5%	65
SDS	1%	0
Triton X-100	0.1%	144
Triton X-100	0.5%	103
Triton X-100	1%	94

#### EXAMPLE 4

- pH Optimum and Stability* - Purified DPP-7 was active against H-Ala-Phe-pNA over a broad pH range, from neutral to basic pH (6.5-9.0) (Fig. 3). This activity also changed with the ionic strength of the buffer, reaching 200% at 0.5 M NaCl concentration in 100 mM HEPES, pH 8.0. DPP-7 was stable in 0.2 M HEPES, pH 8.0, for one week at 4°C. The protease showed no appreciable loss of activity when kept frozen at -80°C for one month. After 3 hours incubation at either room temperature or 37°C, activity was reduced to 62% and 20%, respectively. The optimum temperature for the hydrolysis of H-Ala-Phe-pNA was determined to be 43°C.

## EXAMPLE 5

*Substrate Specificity* - Among several chromogenic substrates tested, only those with an aliphatic or an aromatic side chain residues in the second, penultimate position were rapidly hydrolyzed by DPP-7 (Table 2).

5

TABLE 2: Kinetic analysis for paranitroanalides cleavage by DPP-7.

Substrate	K <sub>m</sub> [mM]	V <sub>max</sub>
H-Ala-Ala-pNA	0.313	129.65
H-Ala-Phe-pNA	0.441	170.06
H-Gly-Phe-pNA	0.256	54.54

Several other substrates including H-Ala-Pro-pNA, H-Ala-pNA, H-Gly-pNA, H-Ile-pNA, H-Leu-pNA, H-Lys-pNA, H-Phe-pNA, H-Gly-Arg-pNA, H-Gly-Glu-pNA, H-Gly-Lys-pNA, H-Ala-Gly-pNA, H-Gly-Gly-pNA, H-Ala-Ala-Phe-pNA, H-Ala-Gly-Arg-pNA, H-Leu-Thr-Arg-pNA, H-Ala-Phe-Pro-pNA, N $\alpha$ -benzoyl-DL-arginine-pNA, N-met-Ala-Pro-Val-pNA, N-suc-Ala-Ala-pNA, N-suc-Ala-Ala-Pro-Glu-pNA, N-suc-Ala-Ala-Pro-Leu-pNA, N-suc-Ala-Ala-Val-Ala-pNA, Z-Ala-Ala-pNA, Z-Lys-pNA, Z-Arg-pNA, Z-Glu-Glu-pNA, Z-Leu-Leu-Glu-pNA, Z-Lys-Arg-pNA, Z-Phe-Arg-pNA, Z-Phe-Val-Arg-pNA, Z-Tyr-Lys-Arg-pNA were tested, but none of these was hydrolysed by DPP-7.

To further confirm specificity, several synthetic peptides were also tested as substrates for this enzyme. Again, only those polypeptides having an amino acid with an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the polypeptide were cleaved (Table 3), with glycine, proline, or charged amino acids not being acceptable as the second amino acid from the N-terminal end of the polypeptide. The protease did not show any endopeptidase activity on gelatin, insulin  $\beta$  chain, carboxymethylated lysosyme, azocazein or type I collagen. Purified DPP-7 was

devoid of any aminopeptidase activity and did not cleave model substrates with blocked amino-termini.

TABLE 3: Specificity of *P. gingivalis* DPP-7 on synthetic peptides.

Peptides cleaved	Peptides not cleaved
Trp-Ala-↓-Gly-Gly-Asp-Ala-Ser-Gly-Glu (SEQ ID NO:10)	Trp-His-Trp-Leu-Glu-Leu-Lys-Pro-Gly- Glu-Pro-Met-Tyr (SEQ ID NO:14)
Ile-Ala-↓-Arg-Arg-His-Pro-Tyr-Phe-Leu (SEQ ID NO:11)	Ser-Pro-Tyr-Ser-Ser-Glu-Thr-Thr (SEQ ID NO:15)
Lys-Ile-↓-Ala-Gly-Tyr-His-Leu-Glu-Leu (SEQ ID NO:12)	Ala-Pro-Val-Arg-Ser-Leu (SEQ ID NO:16)
Phe-Leu-↓-Arg-Glu-Pro-Val-Ile-Phe-Leu (SEQ ID NO:13)	Gln-Lys-Gln-Met-Ser-Asp-Arg-Arg-Glu (SEQ ID NO:17)

5      *An arrow indicates cleavage site*

#### EXAMPLE 6

*DPP-7 Sequence Analysis* - Purified DPP-7 was resolved on SDS-PAGE and electroblotted onto a PVDF membrane. It had an amino-terminal sequence  
 10    ADKGMWLLNELNQENLDRMRELGFT (SEQ ID NO:18). After proteolytic in-gel digestion of the enzyme additional internal sequences were obtained, including: DNKPYK (SEQ ID NO:19), EMTYL (SEQ ID NO:20), FAQFAN (SEQ ID NO:21), VLPAML (SEQ ID NO:22), SVVPY (SEQ ID NO:23), LFFAGL (SEQ ID NO:24). All of this sequence data allowed us to  
 15    identify the *P. gingivalis* genomic contig gln | TIGR | *P. gingivalis*\_ in the Unfinished Microbial Genomes database, TIGR. An ORF corresponding to the DPP-7 amino acid sequence (SEQ ID NO:1) was found, as indicated by the fact, that all sequences of the DPP-7 derived peptides obtained by the enzyme polypeptide fragmentation by trypsin were present in the protein primary  
 20    structure inferred from the nucleotide sequence of the ORF as shown in Fig. 4. Including a signal peptide (residues 1-24), the entire ORF corresponds to a 712 amino acid polypeptide (see Fig. 4). Interestingly, the DPP-7 ORF contains the

consensus sequence for the active-site serine residue of serine type proteases, TGGNSGSPVF (SEQ ID NO:25). As indicated in Fig. 5 the DPP-7 carboxy-terminus (SEQ ID NO:3) exhibits high degree of identity to that of the V8 serine protease (SEQ ID NO:4), particularly around the putative active site serine residue. This is surprising since the *P. gingivalis* DPP-7 is a dipeptidylpeptidase specific for substrates having an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the substrate, whereas *Staphylococcus aureus* V8 endopeptidase is specific towards substrates including glutamic acid or aspartic acid as the second amino acid from the N-terminal end of the substrate. The similarity search performed using the NCBI TBLASTN tool against GenBank, EMBL, DDBJ and PDB databases showed no significant similarity of DPP-7 to any other known dipeptidylpeptidases, indicating that this enzyme could be regarded as a member of a new family of proteases. Additional searches against databases containing unfinished and finished microbial genomes allowed us to identify more coding sequences encoding similar proteases with consensus active site sequence TGGNSGSPV (Fig. 6; SEQ ID NO:26). A coding sequence of related protein has been found in *P. gingivalis* W83 unfinished portion of complete genome between positions 1360759 and 1362718. This putative proteinase reveals significant similarity to DPP-7 (267/691 identities). Another organism *Shewanella putrefaciens* possesses two related coding sequences (gnl |TIGR\_24|sputre 6401 and gnl |TIGR\_24|sputre 6410) while a plant pathogen *Xylella fastidiosa* contains one coding sequence encoding similar proteinase (gb |AE004008.1|). In addition, the computer assisted search for sequential motifs characteristic for transmembrane domains revealed the presence of two such putative regions within the amino-terminal sequence of DPP-7, with residues 7 to 24 and 62 to 78 most likely folded into hydrophobic  $\alpha$ -helices responsible for membrane anchoring of this enzyme.

The complete disclosure of all patents, patent applications, and publications, and electronically available material (e.g., GenBank amino acid and nucleotide sequence submissions) cited herein are incorporated by reference.



What is claimed is:

1. An isolated dipeptidylpeptidase, active analog, active fragment, or active modification thereof having amidolytic activity for cleavage of a peptide bond between the second and third amino acids from the N-terminal end of a target polypeptide, wherein the target polypeptide has an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the polypeptide.
2. The dipeptidylpeptidase of claim 1 wherein the dipeptidylpeptidase is isolated from *Porphyromonas gingivalis*.
3. The dipeptidylpeptidase of claim 1 wherein the dipeptidylpeptidase is a serine protease.
4. The dipeptidylpeptidase of claim 1 comprising an amino acid sequence TGGNSGSPV (SEQ ID NO:26).
5. The dipeptidylpeptidase of claim 1 comprising an amino acid sequence TGGNSGSPVF (SEQ ID NO:25).
6. The dipeptidylpeptidase of claim 1 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, and SEQ ID NO:26.
7. The dipeptidylpeptidase of claim 1 comprising an amino acid sequence SEQ ID NO:2.
8. The dipeptidylpeptidase of claim 1 wherein the dipeptidylpeptidase is encoded by a nucleic acid comprising a nucleotide sequence SEQ ID NO:1.

9. An isolated polypeptide comprising an amino acid sequence having a percentage amino acid identity greater than about 40% with SEQ ID NO:2.
10. An isolated nucleic acid comprising a coding sequence encoding a dipeptidylpeptidase, active analog, active fragment, or active modification thereof having amidolytic activity for cleavage of a peptide bond between the second and third amino acids from the N-terminal end of a target polypeptide, wherein the target polypeptide has an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom of the second amino acid from the N-terminal end of the polypeptide.
11. The nucleic acid of claim 10 wherein the nucleic acid comprises a nucleotide sequence SEQ ID NO:1.
12. The nucleic acid of claim 10 wherein a complement of the nucleic acid hybridizes to SEQ ID NO:1 under hybridization conditions of 0.5 M phosphate buffer, pH 7.2, 7% SDS, 10 mM EDTA, at 68°C, followed by three for 20 minutes washes in 2x SSC, and 0.1% SDS, at 65°C, wherein at least about 20 nucleotides of the complement hybridize.
13. An isolated nucleic acid encoding a polypeptide, wherein the polypeptide comprises an amino acid sequence having a percentage amino acid identity greater than about 40% with SEQ ID NO:2.
14. A method of identifying an inhibitor of a dipeptidylpeptidase, active analog, active fragment, or active modification thereof, comprising identifying a compound that inhibits the amidolytic activity of the dipeptidylpeptidase by incubating the dipeptidylpeptidase with the compound under conditions that promote amidolytic activity of the dipeptidylpeptidase and determining if the amidolytic activity of the


dipeptidylpeptidase is inhibited relative to the amidolytic activity in the absence of the compound.

15. A method of reducing growth of a bacterium comprising inhibiting a dipeptidylpeptidase, active analog, active fragment, or active modification thereof, by contacting the dipeptidylpeptidase with an inhibitor of the dipeptidylpeptidase.
16. The method of claim 15 wherein the dipeptidylpeptidase is a serine protease.
17. A method for protecting an animal from a periodontal disease caused by *Porphyromonas gingivalis* comprising administering to the animal an inhibitor of dipeptidylpeptidase, wherein the disease is selected from the group consisting of gingivitis and periodontitis.
18. The method of claim 17 wherein the inhibitor is administered by a method selected from the group consisting of subgingival application and controlled release delivery.
19. An immunogenic composition comprising an isolated dipeptidylpeptidase, an antigenic analog, an antigenic fragment, or an antigenic modification thereof having amidolytic activity for cleavage of a peptide bond present in a target polypeptide, the peptide bond being located between the second and third amino acids from the N-terminal end of the target polypeptide, wherein the second amino acid from the N-terminal end of the polypeptide has an aliphatic or an aromatic residue as a substituent on the  $\alpha$ -carbon atom.
20. The composition of claim 19 wherein the dipeptidylpeptidase is a serine protease.

21. The composition of claim 19 wherein the second amino acid is selected from the group consisting of alanine, phenylalanine, isoleucine, and leucine.
22. The immunogenic composition of claim 19 further comprising an adjuvant.
23. A composition comprising an inhibitor of an isolated dipeptidylpeptidase and a pharmaceutically acceptable carrier.


1/7

Fig. 1




2/7

Fig. 2



3/7

Fig. 3



atgccaatgaaatataaaagtattcttcctcgagcagccctgctgttgggtgcttcagggttagccaaagccgacaaggcatgtggctc	90
M Q M K L K S I L L G A A L L L G A S G V A K A D K G M W L	30
ctcaacgaactcaatcaggagaatctggatcgatgcgtgagctcggtcttacgctcccggtggatctcagcttttcgacaagccg	180
L N E L N Q E N L D R M R E L G F T L P L D S L Y S F D K P	60
tccattgccaatgcgctgggttatcttcggtggcggtatgtaccggtatcacagtgctccgatcaggcgctgatctttaccaaccaccactgc	270
S I A N A V V I F G G G C T G I T V S D Q G L I F T N H H C	90
ggatacgggtgctatccagagccaaagcacggtggatcacgactatctgcgcgatggtttcgtttctcgacgatgggtgaggagcttcgg	360
G Y G A I Q S Q S T V D H D Y L R D G F V S R T M G E E L P	120
attccgggtctttccgctaagatctctgcgcaagatcgtgaaggttaacggacaaggtagaaggacagctcaaggggtatcactgacgagatg	450
I P G L S V K Y L R K I V K V T D K V E G Q L K G I T D E M	150
gagcgtctgcgcaagactcaggaggtatgccaaagactggccaaaaaagaaaatcgacagcagaaccaactctgcactcgtagagcctttc	540
E R L R K A Q E V C Q E L A K K E N A D E N Q L C I V E P F	180
tattccaacaacgaataacttctctcgtctacgatgtattcaaggacggttcgtatggattttgctctcccgactctgttaggtaagttc	630
Y S N N E Y F L I V Y D V F K D V R M V F A P P S S V G K F	210
ggaggcgatacggacaactggatgtggcgcgctcacacggcgacttcagcgtattcccgctgtatgccggtgccgacaaccggcgccg	720
G G D T D N W M W P R H T G D F S V F R V Y A G A D N R P A	240
gaatacagcaaggacaataaacctataagcccgctttacttcgctgcggtatccatgcaaggctacaaggctgacgactatgccatgacc	810
E Y S K D N K P Y K P V Y F A A V S M Q G Y K A D D Y A M T	270
atcggtttccgggagcgtacggatcgctacctcacttcttgggtgtggaagatcgatcgaaaacgagacaacatcctcgatcgaaagt	900
I G F P G S T D R Y L T S W G V E D R I E N E N N P R I E V	300
cgcggtatcaagcaaggcatctggaaggaagccatgagcgcagatcaggctacccggtatcaaatatgccagcaagtatgctcagagtgt	990
R G I K Q G I W K E A M S A D Q A T R I K Y A S K Y A Q S A	330
aactattggaagaattcgatcggtatgaaccgcggtctcgtcgtcttgacgtgataggtcgtaagcgtgccgaggaaagagcattcgca	1080
N Y W K N S I G M N R G L A R L D V I G R K R A E E R A F A	360
gactggatcgtaagaacggcaagagtgtgtctatggcgatgtattgtcttctctcgaaaaggcttataaggaaggagccaaggccaac	1170
D W I R K N G K S A V Y G D V L S S L E K A Y K E G A K A N	390
cgtagatgacttatttgagcagacgctcttcgggtgtaccgaggtgggttcggtttgcacagtttgccaacgcattggctacaaaatcct	1260
R E M T Y L S E T L F G G T E V V R F A Q F A N A L A T N P	420
gatgctcatgcgggtatcctcaaatcgcttgacgacaagtcacaagactacctccctcgtctcgaccgtaaggtgctgccgcgatgctc	1350
D A H A G I L K S L D D K Y K D Y L P S L D R K V L P A M L	450
gatattgtacgcggcggtatccctgccgacaagctcccgatatattcaagaatgtaatcgacaagaattcaaaaggcagacagaagaag	1440
D I V R R R I P A D K L P D I F K N V I D K K F K G D T K K	480
tatgcagacttcgtatttcgacaagagtgtggttctctatagcgacaagttccatgccatgctcaagtcattggacaaggaaaagttgcc	1530
Y A D F V F D K S V V P Y S D K F H A M L K S M D K E K F A	510
aaggctatcgagaaagatccggcagtagagctttccaagagcgtaatatgctgctgctgcgctattcaggccgatgcgatggccaatgcc	1620
K A I E K D P A V E L S K S V I A A R A I Q A D A M A N A	540
tatgccattgagaaggcgcaagcgtcttttcttgcgggtttgctgagatgtaccccgagcgtgctctgcgagcagatgccaaacttcacc	1710
Y A I E K G K R L F F A G L R E M Y P G R A L P S D A N F T	570
atgcgtatgagctacggctccatcaagggaatatgaaccgcaggacgggtgcctggtacaactatcatacgacaggcaaggcggtattggag	1800
M R M S Y G S I K G Y E P Q D G A W Y N Y H T T G K G V L E	600
aagcaggatcctaagagcagtaggtttgcggtacaggagaatatctcgcacctcttcgccacaaaaaactatggctcgctatgccgagaaac	1890
K Q D P K S D E F A V Q E N I L D L F R T K N Y G A E N	630
ggtcagctccatcatcgctttctctatcgaaacacacatcacggcggttaactccggtagccccgatttcgataagaacggcggtctgatc	1980
G Q L H I A F L S N N D I T G G N S G S P V F D K N G R L I	660
ggctctgtcttcgatggcaactgggaagctatgagtggtgacatcgagttcgaaaccgatctgcagcgcaacaatcagcgtggacatccgc	2070
G L A F D G N W E A M S G D I E F E P D L Q R T I S V D I R	690
tacgtttctcttcgatgattgacaaatggggctcagtgcccccgctctcatccaagagctgaagttgatctaa	2139
Y V L F M I D K W G Q C P R L I Q E L K L I *	711



Fig. 5

SEQ ID NO. 3	DPP-7	644	TGGNSGSPVFDKNGRLIGLAFDGNWEAMSGDIEFFPDLQRTISVDIRYVLFM	695
			TGGNSGSPVF++ +IG+ + G +G + +++ + +I + F	
SEQ ID NO. 4	V-8	704	TGGNSGSPVFNEKNEVIGIHWGGVPNEFFNGAVFINENVRNFLKQNIEDIAHFA	863

6/7

Fig. 6

SEQ ID NO: 5  
 SEQ ID NO: 6  
 SEQ ID NO: 7  
 SEQ ID NO: 8  
 SEQ ID NO: 9

1 MASQALGFLHONGLNTMKKWLTSVAFAAFASHADEGMWPHOLFANADVYKAKGIEIDAKSHISKLTEFPMN--AVIISL  
 1 -----MFAIAVAINLTCGLATADEGQOPYOP-SHADKLSARGDIPADKLADLTSYPMN--AVVGL  
 1 -----MRENTISISVLAITITVDSTHAGECMVPOOLP-EHAGFLKQAGLOHSPQLSNLTGDEMG--AVVSL  
 1 -----MOMKLKSLHGAALHIGASGVAKADKGMWFLNENQENLDRNELGFTPLDLSLSYFDKPSHANAVVIFG  
 1 -----MKKRLHLEFAMLCLOIAHADEGMWPMQOLG-RKYAQMKERKKEKYEIDLYNPNGTSLKDAVWIFD  
 1 -----

78 GGCTASFSVPKGLVVTNHHCAYSIQNSTPEKNLLQDGFHAKTEDELPAAPGSRVAVTEDVTVNTERVKALENKIG-  
 S1  
 62 CYCTASFSVPQGLVVTNHHCAIKALQVNTKKHNYLEQGFHATSMDKSPSAGPNERVITEAVTDVTSVTKOLSODP--  
 S2  
 66 GNCITASTVSPEGLVVTNHHCAICAIQVNSTPKNLTKEGNAITQADEVSAGPNARVAVHETLTDVTAQAKAALAAAGND  
 X  
 71 GGCTGTTVSDQGLFETNHHCGYCAIQSOSTVDHDIYRDGFVSRIMG-BELPTEGLSVKYRKYVVDKVEGOLKGITD-  
 P1  
 67 GGCTGEVVSDRGLVVTNHHCGYDMIOAHSTLEHNYLENCEWAMREA-BELPNKDI SVVFEDKIEDVDVDYVKKOLKAIKDP  
 P2

157 -----REFYQGVENQEKALVAECEEKDEGYRCQVYSEHGGLEYMLVKOLEITDVRVAVNPAGSVGKYGGVDNMMWPRHT  
 S1  
 140 -----LKRVEEHNHSHKALTKSCAADNRYRCNVRSEHNGLEYMLVKOLEITDVRVAVNPAGSVGKYGGVDNMMWPRHS  
 S2  
 146 -----PFKRTTAEFTFSKQELAKCEEQGYRCQVYSEHNGLEYMLVKOLEITDVRVAVNPAGSVGKYGGVDNMMWPRHT  
 X  
 149 -----EMERLRKAEVCOELAKKENADENQICIVEPHYSNVEYELLVIDVEKDVRYVAVNPAGSVGKYGGVDNMMWPRHT  
 P1  
 146 NSMDYLSPKYLOKLADKAKGNFSANKPGLSVEIKAFYCGNLYLMTFKKTYTIDVRVAVNPAGSVGKYGGVDNMMWPRHT  
 P2

231 GDMSEFRAVVSKNCKPAEFESADNPVPEPKSELKVSAGVSEGEVMVAGYPCPTNNRYTATEVONEFEWAYPEGKQIRER  
 S1  
 214 GDEAFLRAVVGKDKPAEFESADNPVTPKSYLAKVNADGVKACDGVFVAGYPCPTNNRYNLTSELKFASDWLYPTQAKRYOL  
 S2  
 221 GDFSFRVAVVGKDKPASPSENIPYRPHKWLKESDQPLGDEVMVAGYPCPTNNRYALVAEFETAHTWTPYVIGQHFKN  
 X  
 224 GDFSVERVACADNRPAEKSKDNKPKPVYFAAVSMQYKADYAMTIGFPGSDPRYLTSWGVDRDIENENNPRIEVARGI  
 P1  
 226 GDFSIERHIVADKNGNPAPYSEDNVPLKPKRFNFNLSLGGVQENDYAMIMGCPCHIEYFETASEVDEWKSIDNDIRIRMRDI  
 P2

311 FIETIKATAPEGSDERIKVEQTAGHANYAKNFTSMIEFYCKSTMLADREKALBAKTAELAKDSS--REAKYGKTLAEL  
 S1  
 294 QIDTIEAMGQKDADIAIKVAGNMAWMANRMKLNGLLAGFKATDVGIKQQRNDFHAWTKNPN-----LNONLISEL  
 S2  
 301 LIJALIEAASKQNPDIQVAVASTLAGHNTSKNFDQDGFERRINAIHQKQSEETAVHAWKQQGI--RGHEALAAHOTL  
 X  
 304 KQGIWKEAMSAQOATRIKYASKYAQSANYWKNISGMNRLARLDVIGERKEAFADWIRKNGKSAV---YGDVLSLSE  
 P1  
 306 RQGVMLREMLADPOLKIMYSAKYAASQNAKYKRAIGANWAIKTRGIRONKQANQDRHIANWAKQCTP-----RYEEAVHE  
 P2

388 DALIAKSKAHOERDMATSYISSHTMPTANNIYRIAHEKQLPDMQREPCQDRDNTREKASMERIDRRYAAASVDKAVLFD  
 S1  
 368 EVLLAEQOLQTOINYYFTNAQSSMLTAANNIYRLAKEKQKSDAEREIGQOERDFAMFSRPHKIDSSESDVKVDKTLWLQ  
 S2  
 378 VDLTEQYKANODEVILGOENGSGVIGAVNIYRLAIERTKSDAOREACQOERDIPTEIGNIKQMERHILPEMDROMQQY  
 X  
 381 KAYKEGAKANREMTVLSFTLFGGEVVRFAQFANALATNPDAHAGILKSLDPKYDYLPSLDKVLPAMLDIVRRIPAD  
 P1  
 380 IDATVAKRADLRRRWMIIEGIRGHEFARSPITEDETKALQENDA SRKEADIKIRTRYSKFANKQYSAEVDKKVAVA  
 P2

468 MKRYAALPEAQRIPAMDKAFGIDNKVNEAKLAKTIDKMYAKTELGNKDYRLANWMEKSVDDDEKASKDBFIQEAVAMYDTN  
 S1  
 448 DIIN--AYLSQPNRVAALDNNMLNDKN--VSLAAKIDGLYSITLITDQACRLANWMAKAKABETSSDPFIIRLAVAILDNTN  
 S2

458 WLTENKIPVQORVANDVWEDGIPAT-----L-KRLGDTKLSSEERIKWENADRAAFESSQDPAIRVAVAMPAL  
 P1 461 KLPDLEKNVIDKGEKCDTKKYADEVFD--KS-----VVPYSDKPHAMKSMQKEKFAKAIKDPDAVEISKSVIAAA  
 P2 460 MLTEYLKEIPYENPLHLRLVKDRFAG----DVQAYVDDIFARSYFGSEAQDFAFAPVPSVEKLAEDPMVLEASSVEDEY  
  
 S1 548 MSEERKEKELDCGLMKVRPOTMDATHAYNLEQCKPVIYADANSSLRVTVCHVKGYSPKDCIVAVPFRLEGTIVQKDTG-ID  
 S2 524 MAQEKAEKILAGKLTSTARPAYAAVADYYKANNWPVYPDANGCHLRISYGMVDGYQSRDAIKKQPFTRLDGIVAKHTG-VE  
 X 530 LETERONKIRTCGLLKKARPIYLCAIADYNKSHCKFVAPDANSSLRITRCHVKGYSPKDCGVEYTPPFTTLQGVMAKNTG-VE  
 P1 530 RALQADAMANAYAEKGRLEFAGIREMYP--GRALPSDANFTMRMSYCSHKGYEPQDCAWNYHTTGKGVLEKQDPKSD  
 P2 536 RKQYNELEPYDDPILRAQRTYTAGILEMDG--DQDQEPDANLTLRFTYQOVKGYSPRDNVARYGHQTTLLDGVLEKEDPDNW  
  
 S1 627 PEDAPKQOLEHTTKQVCGDEYMKSHDS-----VPVNELSLHDTTCGNSGSPVTLNGRAELVGLLE  
 S2 603 PYNAPRKLLDAISVQRECDHLVKSYYQDPRGWCRLFSCLDKPEETNSVPVNELSSVDTCGNSGSPVNGKGLVGLNE  
 X 609 PRDSPKSLUNALIKAKSYANLADORTGT-----VPVNELSDLEDTTCGNSGSPVLDAGCKLVGLAR  
 P1 608 ERAVQENHLDLFRKNTYGRVAENGOLH-----IAF--LSNNDLTTCGNSGSPVFDKNCRLTGLAR  
 P2 614 EFVVDPKKAYIERKDEGRKADRSGRMP-----VAECATHTTTCGNSGSPVMNANGELTGLNE  
  
 S1 686 DGVYESHICGMEDNEINRSIHVDSRMYLWVMKYLHDHADNLTAEHEHVN-  
 S2 683 DSTMEALTKDWFENETLTRAVEVDIRYLLMMMDVDDHADNLTKEPDIVRN  
 X 668 DGNWESVSSNWVFDPPVMTRTIIAVDSRYVQWIMTEVAPAPHKKEINLYR-  
 P1 665 DGNWEAMSGDIEFEPDLQRTISVDIRYVLEMDKW-----  
 P2 672 DRNWEGVGEDIQMLADYQRSIIVDIRYVILVIDKVGCCQRIIDEMANIMP-

The foregoing detailed description and examples have been given for clarity of understanding only. No unnecessary limitations are to be understood therefrom. The invention is not limited to the exact details shown and described, for variations obvious to one skilled in the art will be included within the invention defined by the claims.

### SEQUENCE LISTING FREE TEXT

	SEQ ID NO:1	Coding sequence encoding <i>Porphyromonas gingivalis</i> DPP-7
10	SEQ ID NO:2	<i>Porphyromonas gingivalis</i> DPP-7
	SEQ ID NO:3	C-terminal region of <i>Porphyromonas gingivalis</i> DPP-7
	SEQ ID NO:4	C-terminal region of <i>Staphylococcus aureus</i> V8 endopeptidase
15	SEQ ID NO:5	Coding sequence for <i>Shewanella putrefaciens</i> gnl   TIGR_24   sputre 6401
	SEQ ID NO:6	Coding sequence for <i>Shewanella putrefaciens</i> gnl   TIGR_24   sputre 6410
	SEQ ID NO:7	Coding sequence for <i>Xylella fastidiosa</i> gb   AE004008.1
20	SEQ ID NO:8	Coding sequence for <i>Porphyromonas gingivalis</i> gnl   TIGR   <i>P. gingivalis</i> _CPG.con
	SEQ ID NO:9	Coding sequence for <i>Porphyromonas gingivalis</i> DPP-7 gnl   TIGR   <i>P. gingivalis</i> _CPG.con
	SEQ ID NO:10-17	Synthetic peptides
	SEQ ID NO:18	N-terminal region of <i>Porphyromonas gingivalis</i> DPP-7
25	SEQ ID NO:19-24	Internal sequences of <i>Porphyromonas gingivalis</i> DPP-7
	SEQ ID NO:25-26	Consensus sequences for active sites for serine type proteases

## SEQUENCE LISTING

<110> THE UNIVERSITY OF GEORGIA RESEARCH FOUNDATION, INC.

<120> Dipeptidylpeptidases And Methods Of Use

<130> 235.00440201

<150> US 60/246,827

<151> 2000-11-08

<160> 26

<170> PatentIn version 3.0

<210> 1

<211> 2139

<212> DNA

<213> Porphyromonas gingivalis

```

<400> 1
atgcaaatga aattaaag tattcttctc ggagcagccc tgctgttggg tgcttcaggg      60
gtagccaaag ccgacaaagg catgtggctc ctcaacgaac tcaatcagga gaatctggat      120
cgaatgcgtg agctcggctt tacgctcccg ttggattcgc tctacagttt cgacaagccg      180
tccattgcca atgccgtggt tatcttcggt ggcggatgta ccggtatcac agtgtccgat      240
cagggcctga tctttacca ccaccactgc ggatacgggtg ctatccagag ccaaagcacg      300
gtggatcacg actatctgcg cgatgggttc gtttctcgca cgatgggtga ggagcttccg      360
attccgggtc tttccgtgaa gtatctgcgc aagatcgtga aggtaacgga caaggtagaa      420
ggacagctca agggatcac tgacgagatg gagcgtctgc gcaaagctca ggaggtatgc      480
caagaactgg ccaaaaaaga aaatgcagac gagaaccaac tctgcatcgt agagcctttc      540
tattccaaca acgaatactt cctcatcgtc tacgatgtat tcaaggacgt tcgtatggta      600
tttgctcctc ccagctctgt aggttaagttc ggaggcgata cggacaactg gatgtggccg      660
cgtcacacgg gcgacttcag cgtattccgc gtgtatacca atgccgacaa ccggccggcc      720

```

gaatacagca aggacaataa accctataag cccgtttact tcgctgccgt atccatgcaa 780  
 ggctacaagg ctgacgacta tgccatgacc atcggtttcc cgggcagtac ggatcgctac 840  
 ctcaatttctt ggggtgtgga agatcgatc gaaaacgaga acaatcctcg tatcgaagtt 900  
 cgcggtatca agcaaggcat ctggaaggaa gccatgagcg cagatcaggc taccggtatc 960  
 aaatatgcca gcaagtatgc tcagagtgc aactattgga agaattcgat cgggtatgaac 1020  
 cgcggtctcg ctcgctttga cgtgataggt cgtaagcgtg ccgaggaaag agcattcgca 1080  
 gactggatcc gtaagaacgg caagagtgc gtctatggcg atgtattgtc ttctctcgaa 1140  
 aaggcttata aggaaggagc caaggccaac cgtgagatga cttatttgag cgagacgctc 1200  
 ttcggtggtg ccgagggtgt tcgttttgca cagtttgcca acgcattggc tacaatcct 1260  
 gatgctcatg ccggtatcct caaatcgctt gacgacaagt acaaagacta cctcccctcg 1320  
 ctcgaccgta aggtgctgcc cgccatgctc gatattgtac gccggcgat ccctgccgac 1380  
 aagctccccg atatattcaa gaatgtaatc gacaagaaat tcaaaggcga cacgaagaag 1440  
 tatgcagact tcgtattcga caagagtgtg gttccttata gcgacaagtt ccatgccatg 1500  
 ctcaagtcca tggacaagga aaagtttgcc aaggctatcg agaaagatcc ggcagtagag 1560  
 ctttccaaga gcgtaatagc tgctgctcgc gctattcagg ccgatgcat ggccaatgcc 1620  
 tatgccattg agaagggcaa gcgtctttt tttgccggtt tgcgtgagat gtaccccgga 1680  
 cgtgctctgc cgagcgatgc caacttcacc atgcgtatga gctacggctc catcaaggga 1740  
 tatgaaccgc aggacggtgc ctggtacaac tatcatcga caggcaaggg cgtattggag 1800  
 aagcaggatc ctaagagcga tgagtttgcc gtacaggaga atatcctcga cctcttcgcg 1860  
 accaaaaact atggtcgcta tgccgagaac ggtcagctcc atatcgcttt cctatcgaac 1920  
 aacgacatca cgggcggtaa ctccggtagc cccgtattcg ataagaacgg ccgtctgatc 1980  
 ggtcttgctt tcgatggcaa ctgggaagct atgagtgtg acatcgagtt cgaacccgat 2040  
 ctgcagcgca caatcagcgt ggacatccgc tacgttctct tcatgattga caaatggggt 2100  
 cagtgccccc gtctcatcca agagctgaag ttgatctaa 2139

<210> 2

<211> 712

<212> PRT

<213> Porphyromonas gingivalis

<400> 2

Met Gln Met Lys Leu Lys Ser Ile Leu Leu Gly Ala Ala Leu Leu Leu  
 1 5 10 15

Gly Ala Ser Gly Val Ala Lys Ala Asp Lys Gly Met Trp Leu Leu Asn  
 20 25 30  
 Glu Leu Asn Gln Glu Asn Leu Asp Arg Met Arg Glu Leu Gly Phe Thr  
 35 40 45  
 Leu Pro Leu Asp Ser Leu Tyr Ser Phe Asp Lys Pro Ser Ile Ala Asn  
 50 55 60  
 Ala Val Val Ile Phe Gly Gly Gly Cys Thr Gly Ile Thr Val Ser Asp  
 65 70 75 80  
 Gln Gly Leu Ile Phe Thr Asn His His Cys Gly Tyr Gly Ala Ile Gln  
 85 90 95  
 Ser Gln Ser Thr Val Asp His Asp Tyr Leu Arg Asp Gly Phe Val Ser  
 100 105 110  
 Arg Thr Met Gly Glu Glu Leu Pro Ile Pro Gly Leu Ser Val Lys Tyr  
 115 120 125  
 Leu Arg Lys Ile Val Lys Val Thr Asp Lys Val Glu Gly Gln Leu Lys  
 130 135 140  
 Gly Ile Thr Asp Glu Met Glu Arg Leu Arg Lys Ala Gln Glu Val Cys  
 145 150 155 160  
 Gln Glu Leu Ala Lys Lys Glu Asn Ala Asp Glu Asn Gln Leu Cys Ile  
 165 170 175  
 Val Glu Pro Phe Tyr Ser Asn Asn Glu Tyr Phe Leu Ile Val Tyr Asp  
 180 185 190  
 Val Phe Lys Asp Val Arg Met Val Phe Ala Pro Pro Ser Ser Val Gly  
 195 200 205  
 Lys Phe Gly Gly Asp Thr Asp Asn Trp Met Trp Pro Arg His Thr Gly  
 210 215 220  
 Asp Phe Ser Val Phe Arg Val Tyr Ala Gly Ala Asp Asn Arg Pro Ala  
 225 230 235 240  
 Glu Tyr Ser Lys Asp Asn Lys Pro Tyr Lys Pro Val Tyr Phe Ala Ala  
 245 250 255  
 Val Ser Met Gln Gly Tyr Lys Ala Asp Asp Tyr Ala Met Thr Ile Gly  
 260 265 270  
 Phe Pro Gly Ser Thr Asp Arg Tyr Leu Thr Ser Trp Gly Val Glu Asp  
 275 280 285  
 Arg Ile Glu Asn Glu Asn Asn Pro Arg Ile Glu Val Arg Gly Ile Lys  
 290 295 300  
 Gln Gly Ile Trp Lys Glu Ala Met Ser Ala Asp Gln Ala Thr Arg Ile  
 305 310 315 320  
 Lys Tyr Ala Ser Lys Tyr Ala Gln Ser Ala Asn Tyr Trp Lys Asn Ser  
 325 330 335  
 Ile Gly Met Asn Arg Gly Leu Ala Arg Leu Asp Val Ile Gly Arg Lys  
 340 345 350

Arg Ala Glu Glu Arg Ala Phe Ala Asp Trp Ile Arg Lys Asn Gly Lys  
 355 360 365  
 Ser Ala Val Tyr Gly Asp Val Leu Ser Ser Leu Glu Lys Ala Tyr Lys  
 370 375 380  
 Glu Gly Ala Lys Ala Asn Arg Glu Met Thr Tyr Leu Ser Glu Thr Leu  
 385 390 395 400  
 Phe Gly Gly Thr Glu Val Val Arg Phe Ala Gln Phe Ala Asn Ala Leu  
 405 410 415  
 Ala Thr Asn Pro Asp Ala His Ala Gly Ile Leu Lys Ser Leu Asp Asp  
 420 425 430  
 Lys Tyr Lys Asp Tyr Leu Pro Ser Leu Asp Arg Lys Val Leu Pro Ala  
 435 440 445  
 Met Leu Asp Ile Val Arg Arg Arg Ile Pro Ala Asp Lys Leu Pro Asp  
 450 455 460  
 Ile Phe Lys Asn Val Ile Asp Lys Lys Phe Lys Gly Asp Thr Lys Lys  
 465 470 475 480  
 Tyr Ala Asp Phe Val Phe Asp Lys Ser Val Val Pro Tyr Ser Asp Lys  
 485 490 495  
 Phe His Ala Met Leu Lys Ser Met Asp Lys Glu Lys Phe Ala Lys Ala  
 500 505 510  
 Ile Glu Lys Asp Pro Ala Val Glu Leu Ser Lys Ser Val Ile Ala Ala  
 515 520 525  
 Ala Arg Ala Ile Gln Ala Asp Ala Met Ala Asn Ala Tyr Ala Ile Glu  
 530 535 540  
 Lys Gly Lys Arg Leu Phe Phe Ala Gly Leu Arg Glu Met Tyr Pro Gly  
 545 550 555 560  
 Arg Ala Leu Pro Ser Asp Ala Asn Phe Thr Met Arg Met Ser Tyr Gly  
 565 570 575  
 Ser Ile Lys Gly Tyr Glu Pro Gln Asp Gly Ala Trp Tyr Asn Tyr His  
 580 585 590  
 Thr Thr Gly Lys Gly Val Leu Glu Lys Gln Asp Pro Lys Ser Asp Glu  
 595 600 605  
 Phe Ala Val Gln Glu Asn Ile Leu Asp Leu Phe Arg Thr Lys Asn Tyr  
 610 615 620  
 Gly Arg Tyr Ala Glu Asn Gly Gln Leu His Ile Ala Phe Leu Ser Asn  
 625 630 635 640  
 Asn Asp Ile Thr Gly Gly Asn Ser Gly Ser Pro Val Phe Asp Lys Asn  
 645 650 655  
 Gly Arg Leu Ile Gly Leu Ala Phe Asp Gly Asn Trp Glu Ala Met Ser  
 660 665 670  
 Gly Asp Ile Glu Phe Glu Pro Asp Leu Gln Arg Thr Ile Ser Val Asp  
 675 680 685  
 Ile Arg Tyr Val Leu Phe Met Ile Asp Lys Trp Gly Gln Cys Pro Arg



690                                      695                                      700

Leu Ile Gln Glu Leu Lys Leu Ile  
 705                                      710

<210> 3

<211> 52

<212> PRT

<213> Porphyromonas gingivalis

<400> 3

Thr Gly Gly Asn Ser Gly Ser Pro Val Phe Asp Lys Asn Gly Arg Leu  
 1                                      5                                      10                                      15

Ile Gly Leu Ala Phe Asp Gly Asn Trp Glu Ala Met Ser Gly Asp Ile  
                                     20                                      25                                      30

Glu Phe Glu Pro Asp Leu Gln Arg Thr Ile Ser Val Asp Ile Arg Tyr  
                                     35                                      40                                      45

Val Leu Phe Met  
                                     50

<210> 4

<211> 52

<212> PRT

<213> staphylococcus aureus

<400> 4

Thr Gly Gly Asn Ser Gly Ser Pro Val Phe Asn Glu Lys Asn Glu Val  
 1                                      5                                      10                                      15

Ile Gly Ile His Trp Gly Gly Val Pro Asn Glu Phe Asn Gly Ala Val  
                                     20                                      25                                      30

Phe Ile Asn Glu Asn Val Arg Asn Phe Leu Lys Gln Asn Ile Glu Asp  
                                     35                                      40                                      45

Ile His Phe Ala  
                                     50

<210> 5

<211> 734

<212> PRT

<213> Shewanella putrefaciens

<400> 5

Met Ala Ser Gln Ala Leu Gly Phe Leu His Gln Asn Gly Leu Asn Thr  
 1 5 10 15  
 Met Lys Lys Trp Leu Leu Ser Val Ala Val Ala Ala Ser Phe Ala Ser  
 20 25 30  
 His Ala Asp Glu Gly Met Trp Gln Pro His Gln Leu Pro Ala Met Ala  
 35 40 45  
 Asp Val Leu Lys Ala Lys Gly Leu Glu Ile Asp Ala Lys Ser Ile Ser  
 50 55 60  
 Lys Leu Thr Glu Phe Pro Met Asn Ala Val Ile Ser Leu Gly Gly Cys  
 65 70 75 80  
 Thr Ala Ser Phe Val Ser Pro Lys Gly Leu Val Val Thr Asn His His  
 85 90 95  
 Cys Ala Tyr Gly Ser Ile Gln Tyr Asn Ser Thr Pro Glu Lys Asn Leu  
 100 105 110  
 Leu Gln Asp Gly Phe Leu Ala Lys Thr Phe Ala Asp Glu Leu Pro Ala  
 115 120 125  
 Ala Pro Gly Ser Arg Val Tyr Val Thr Glu Asp Val Thr Asn Val Thr  
 130 135 140  
 Glu Arg Val Lys Ala Gly Leu Glu Asn Lys Thr Gly Arg Glu Phe Tyr  
 145 150 155 160  
 Gln Gly Val Glu Asn Gln Glu Lys Ala Leu Val Ala Glu Cys Glu Lys  
 165 170 175  
 Asp Glu Gly Tyr Arg Cys Gln Val Tyr Ser Phe His Gly Gly Leu Glu  
 180 185 190  
 Tyr Tyr Leu Val Lys Gln Leu Glu Ile Arg Asp Val Arg Leu Val Tyr  
 195 200 205  
 Asn Pro Ala Gly Ser Val Gly Lys Tyr Gly Gly Asp Val Asp Asn Trp  
 210 215 220  
 Met Trp Pro Arg His Thr Gly Asp Tyr Ser Phe Tyr Arg Ala Tyr Val  
 225 230 235 240  
 Ser Lys Asn Gly Lys Pro Ala Glu Phe Ser Ala Asp Asn Val Pro Tyr  
 245 250 255  
 Glu Pro Lys Ser Phe Leu Lys Val Ser Ala Lys Gly Val Ser Glu Gly  
 260 265 270  
 Asp Phe Val Met Val Ala Gly Tyr Pro Gly Arg Thr Asn Arg Tyr Arg  
 275 280 285  
 Thr Ala Thr Glu Val Gln Asn Glu Phe Glu Trp Ala Tyr Pro Glu Gly  
 290 295 300  
 Lys Met Leu Arg Glu Arg Phe Ile Glu Ile Ile Lys Ala Thr Ala Pro  
 305 310 315 320  
 Ile Gly Ser Asp Glu Arg Ile Lys Tyr Glu Ser Gln Ile Ala Gly Leu  
 325 330 335

Ala Asn Tyr Ala Lys Asn Phe Thr Ser Met Ile Glu Phe Tyr Gly Lys  
 340 345 350  
 Ser Thr Met Leu Ala Asp Arg Lys Ala Leu Glu Ala Lys Leu Ala Glu  
 355 360 365  
 Trp Ile Ala Lys Asp Ser Ser Arg Glu Ala Lys Tyr Gly Lys Thr Leu  
 370 375 380  
 Ala Glu Leu Asp Ala Leu Ile Ala Lys Ser Lys Ala His Gln Glu Arg  
 385 390 395 400  
 Asp Met Ile Leu Ser Tyr Ile Ser Ser Thr Thr Met Leu Pro Thr Ala  
 405 410 415  
 Asn Asn Leu Tyr Arg Leu Ala His Glu Lys Gln Leu Pro Asp Met Gln  
 420 425 430  
 Arg Glu Pro Gly Phe Gln Asp Arg Asp Met Thr Arg Phe Lys Ala Ser  
 435 440 445  
 Met Glu Arg Ile Asp Arg Arg Tyr Ala Ala Ser Val Asp Lys Ala Val  
 450 455 460  
 Leu Phe Asp Met Leu Lys Arg Tyr Ala Ala Leu Pro Glu Ala Gln Arg  
 465 470 475 480  
 Leu Pro Ala Met Asp Lys Ala Phe Gly Ile Asp Asn Lys Val Asn Glu  
 485 490 495  
 Ala Lys Leu Ala Lys Thr Leu Asp Lys Met Tyr Ala Lys Thr Glu Leu  
 500 505 510  
 Gly Asn Lys Asp Val Arg Leu Ala Trp Met Glu Lys Ser Val Asp Asp  
 515 520 525  
 Phe Lys Ala Ser Lys Asp Pro Phe Ile Gln Phe Ala Val Ala Met Tyr  
 530 535 540  
 Asp Thr Asn Met Ser Glu Glu Lys Lys Glu Lys Glu Leu Asp Gly Glu  
 545 550 555 560  
 Leu Met Lys Val Arg Pro Gln Tyr Met Asp Ala Ile Ile Ala Tyr Asn  
 565 570 575  
 Leu Glu Gln Gly Lys Pro Val Tyr Ala Asp Ala Asn Ser Ser Leu Arg  
 580 585 590  
 Val Thr Val Gly His Val Lys Gly Tyr Ser Pro Lys Asp Gly Leu Val  
 595 600 605  
 Ala Val Pro Phe Thr Arg Leu Glu Gly Ile Val Gln Lys Asp Thr Gly  
 610 615 620  
 Ile Asp Pro Phe Asp Ala Pro Lys Gln Gln Leu Glu Leu Ile Lys Gln  
 625 630 635 640  
 Lys Gln Tyr Gly Asp Phe Tyr Met Lys Ser Ile Asp Ser Val Pro Val  
 645 650 655  
 Asn Phe Leu Ser Thr Leu Asp Thr Thr Gly Gly Asn Ser Gly Ser Pro  
 660 665 670  
 Thr Leu Asn Gly Arg Ala Glu Leu Val Gly Leu Leu Phe Asp Gly Val

675                      680                      685  
 Tyr Glu Ser Ile Ile Gly Gly Trp Ala Phe Asp Asn Glu Ile Asn Arg  
     690                      695                      700  
 Ser Ile His Val Asp Ser Arg Tyr Met Leu Trp Val Met Lys Tyr Leu  
     705                      710                      715                      720  
 Asp His Ala Asp Asn Leu Leu Ala Glu Met Glu Ile Val Asn  
                     725                      730

<210> 6

<211> 732

<212> PRT

<213> Shewanella putrefaciens

<400> 6

Met Arg Ile Ala Leu Val Ala Ala Leu Val Leu Thr Cys Gly Ile Ala  
 1                      5                      10  
 Thr Ala Asp Glu Gly Gln Trp Gln Pro Tyr Gln Met Pro Ser Ile Ala  
                     20                      25                      30  
 Asp Lys Leu Ser Ala Arg Gly Ile Asp Ile Pro Ala Asp Lys Leu Ala  
                     35                      40                      45  
 Asp Leu Thr Ser Tyr Pro Met Asn Ala Val Val Gly Leu Gly Tyr Cys  
                     50                      55                      60  
 Thr Ala Ser Phe Val Ser Pro Gln Gly Leu Val Val Thr Asn His His  
 65                      70                      75                      80  
 Cys Ala Tyr Lys Ala Ile Gln Tyr Asn Thr Lys Lys Glu His Asn Tyr  
                     85                      90                      95  
 Leu Glu Gln Gly Phe Leu Ala Thr Ser Met Asp Lys Glu Pro Ser Ala  
                     100                      105                      110  
 Gly Pro Asn Glu Arg Leu Tyr Ile Thr Glu Ala Val Thr Asp Val Thr  
                     115                      120                      125  
 Ser Asp Val Thr Lys Asp Leu Ser Gln Asp Pro Leu Lys Arg Tyr Glu  
                     130                      135                      140  
 Glu Ile Glu Asn His Ser Lys Ala Leu Ile Lys Ser Cys Glu Ala Asp  
 145                      150                      155                      160  
 Asp Asn Tyr Arg Cys Asn Val Arg Ser Phe His Asn Gly Leu Glu Tyr  
                     165                      170                      175  
 Tyr Leu Ile Lys Gln Leu Met Ile Arg Asp Val Arg Leu Val Tyr Ala  
                     180                      185                      190  
 Pro Pro Glu Ser Val Gly Gly Tyr Gly Gly Asp Ile Asp Asn Tyr Glu  
                     195                      200                      205  
 Tyr Pro Arg His Ser Gly Asp Phe Ala Phe Leu Arg Ala Tyr Val Gly  
                     210                      215                      220

Lys Asp Gly Lys Pro Ala Ala Phe Ser Glu Asp Asn Ile Pro Tyr Thr  
 225 230 235 240  
 Pro Lys Ser Tyr Leu Lys Val Asn Ala Asp Gly Val Lys Ala Gly Asp  
 245 250 255  
 Gly Val Phe Val Ala Gly Tyr Pro Gly Thr Thr Asn Arg Tyr Asn Leu  
 260 265 270  
 Thr Ser Glu Leu Lys Phe Ala Ser Asp Trp Leu Tyr Pro Thr Gln Ala  
 275 280 285  
 Lys Arg Tyr Gln Leu Gln Ile Asp Thr Ile Glu Ala Met Gly Gln Lys  
 290 295 300  
 Asp Ala Asp Ile Ala Ile Lys Tyr Ala Gly Asn Met Ala Ser Met Ala  
 305 310 315 320  
 Asn Arg Met Lys Lys Leu Asn Gly Leu Leu Ala Gly Phe Lys Ala Thr  
 325 330 335  
 Asp Ile Val Gly Ile Lys Gln Gln Arg Glu Asn Asp Phe Leu Ala Trp  
 340 345 350  
 Leu Thr Lys Asn Pro Asn Leu Asn Gln Asn Leu Ile Ser Glu Leu Glu  
 355 360 365  
 Val Leu Leu Ala Glu Gln Gln Leu Gln Thr Gln Thr Asn Tyr Tyr Phe  
 370 375 380  
 Thr Asn Ala Gln Ser Ser Thr Leu Leu Thr Ala Ala Asn Asn Leu Tyr  
 385 390 395 400  
 Arg Leu Ala Lys Glu Lys Gln Lys Ser Asp Ala Glu Arg Glu Ile Gly  
 405 410 415  
 Tyr Gln Glu Arg Asp Leu Ala Met Phe Ser Ser Arg Leu Lys Arg Ile  
 420 425 430  
 Asp Ser Ser Phe Asp Val Lys Val Asp Lys Thr Leu Trp Leu Gln Asp  
 435 440 445  
 Leu Asn Ala Tyr Leu Ser Gln Pro Asn Arg Val Ala Ala Leu Asp Asn  
 450 455 460  
 Met Leu Asn Leu Asn Asp Lys Asn Val Ser Leu Ala Ala Lys Leu Asp  
 465 470 475 480  
 Gly Leu Tyr Ser Leu Thr Thr Leu Thr Asp Gln Ala Gln Arg Leu Ala  
 485 490 495  
 Trp Met Glu Ala Asp Ala Lys Ala Phe Glu Thr Ser Ser Asp Pro Phe  
 500 505 510  
 Ile Arg Leu Ala Val Ala Leu Tyr Asp Thr Asn Met Ala Gln Glu Lys  
 515 520 525  
 Ala Glu Lys Ile Leu Ala Gly Lys Leu Ser Thr Ala Arg Pro Ala Tyr  
 530 535 540  
 Met Ala Ala Val Ile Asp Tyr Tyr Lys Ala Asn Asn Trp Pro Val Tyr  
 545 550 555 560



12/22

Glu Met Asp Arg Gln Met Gln Gln Tyr Trp Leu Thr Glu Tyr Asn Lys  
 450 455 460  
 Leu Pro Val Lys Gln Arg Val Ala Ala Ile Asp Val Trp Leu Gly Asp  
 465 470 475 480  
 Gly Ile Pro Ala Thr Leu Lys Arg Leu Gly Asp Thr Lys Leu Ser Ser  
 485 490 495  
 Ser Glu Glu Arg Leu Lys Trp Phe Asn Ala Asp Arg Ala Ala Phe Glu  
 500 505 510  
 Ser Ser Gln Asp Pro Ala Ile Arg Tyr Ala Val Ala Ile Met Pro Ala  
 515 520 525  
 Leu Leu Glu Ile Glu Arg Gln Asn Lys Ile Arg Thr Gly Glu Leu Leu  
 530 535 540  
 Lys Ala Arg Pro Ile Tyr Leu Gln Ala Leu Ala Asp Tyr Asn Lys Ser  
 545 550 555 560  
 His Gly Lys Phe Val Tyr Pro Asp Ala Asn Ser Ser Leu Arg Ile Thr  
 565 570 575  
 Phe Gly His Val Lys Gly Tyr Ser Pro Lys Asp Gly Val Glu Tyr Thr  
 580 585 590  
 Pro Phe Thr Thr Leu Gln Gly Val Met Ala Lys Asn Thr Gly Val Glu  
 595 600 605  
 Pro Phe Asp Ser Pro Lys Ser Leu Ile Asn Ala Ile Lys Ala Lys Ser  
 610 615 620  
 Tyr Ala Asn Leu Ala Asp Gln Arg Ile Gly Thr Val Pro Val Asn Phe  
 625 630 635 640  
 Leu Ser Asp Leu Asp Ile Thr Gly Gly Asn Ser Gly Ser Pro Val Leu  
 645 650 655  
 Asp Ala His Gly Lys Leu Val Gly Leu Ala Phe Asp Gly Asn Trp Glu  
 660 665 670  
 Ser Val Ser Ser Asn Trp Val Phe Asp Pro Val Met Thr Arg Thr Ile  
 675 680 685  
 Ala Val Asp Ser Arg Tyr Val Gln Trp Ile Met Thr Glu Val Ala Pro  
 690 695 700  
 Ala Pro His Leu Leu Lys Glu Leu Asn Leu Tyr Arg  
 705 710 715

&lt;210&gt; 8

&lt;211&gt; 699

&lt;212&gt; PRT

&lt;213&gt; Porphyromonas gingivalis

&lt;400&gt; 8



Met Gln Met Lys Leu Lys Ser Ile Leu Leu Gly Ala Ala Leu Leu Leu  
 1 5 10 15  
 Gly Ala Ser Gly Val Ala Lys Ala Asp Lys Gly Met Trp Leu Leu Asn  
 20 25 30  
 Glu Leu Asn Gln Glu Asn Leu Asp Arg Met Arg Glu Leu Gly Phe Thr  
 35 40 45  
 Leu Pro Leu Asp Ser Leu Tyr Ser Phe Asp Lys Pro Ser Ile Ala Asn  
 50 55 60  
 Ala Val Val Ile Phe Gly Gly Gly Cys Thr Gly Ile Thr Val Ser Asp  
 65 70 75 80  
 Gln Gly Leu Ile Phe Thr Asn His His Cys Gly Tyr Gly Ala Ile Gln  
 85 90 95  
 Ser Gln Ser Thr Val Asp His Asp Tyr Leu Arg Asp Gly Phe Val Ser  
 100 105 110  
 Arg Thr Met Gly Glu Glu Leu Pro Ile Pro Gly Leu Ser Val Lys Tyr  
 115 120 125  
 Leu Arg Lys Ile Val Lys Val Thr Asp Lys Val Glu Gly Gln Leu Lys  
 130 135 140  
 Gly Ile Thr Asp Glu Met Glu Arg Leu Arg Lys Ala Gln Glu Val Cys  
 145 150 155 160  
 Gln Glu Leu Ala Lys Lys Glu Asn Ala Asp Glu Asn Gln Leu Cys Ile  
 165 170 175  
 Val Glu Pro Phe Tyr Ser Asn Asn Glu Tyr Phe Leu Ile Val Tyr Asp  
 180 185 190  
 Val Phe Lys Asp Val Arg Met Val Phe Ala Pro Pro Ser Ser Val Gly  
 195 200 205  
 Lys Phe Gly Gly Asp Thr Asp Asn Trp Met Trp Pro Arg His Thr Gly  
 210 215 220  
 Asp Phe Ser Val Phe Arg Val Tyr Ala Gly Ala Asp Asn Arg Pro Ala  
 225 230 235 240  
 Glu Tyr Ser Lys Asp Asn Lys Pro Tyr Lys Pro Val Tyr Phe Ala Ala  
 245 250 255  
 Val Ser Met Gln Gly Tyr Lys Ala Asp Asp Tyr Ala Met Thr Ile Gly  
 260 265 270  
 Phe Pro Gly Ser Thr Asp Arg Tyr Leu Thr Ser Trp Gly Val Glu Asp  
 275 280 285  
 Arg Ile Glu Asn Glu Asn Asn Pro Arg Ile Glu Val Arg Gly Ile Lys  
 290 295 300  
 Gln Gly Ile Trp Lys Glu Ala Met Ser Ala Asp Gln Ala Thr Arg Ile  
 305 310 315 320  
 Lys Tyr Ala Ser Lys Tyr Ala Gln Ser Ala Asn Tyr Trp Lys Asn Ser  
 325 330 335  
 Ile Gly Met Asn Arg Gly Leu Ala Arg Leu Asp Val Ile Gly Arg Lys

15/22

Ile Arg Tyr Val Leu Phe Met Ile Asp Lys Trp  
 690 695

<210> 9

<211> 720

<212> PRT

<213> Porphyromonas gingivalis

<400> 9

Met Lys Lys Arg Leu Leu Leu Pro Leu Phe Ala Val Leu Cys Leu Cys  
 1 5 10 15  
 Gln Ile Ala His Ala Asp Glu Gly Met Trp Leu Met Gln Gln Leu Gly  
 20 25 30  
 Arg Lys Tyr Ala Gln Met Lys Glu Arg Gly Leu Lys Met Lys Glu Tyr  
 35 40 45  
 Asp Leu Tyr Asn Pro Asn Gly Thr Ser Leu Lys Asp Ala Val Val Leu  
 50 55 60  
 Phe Asp Gly Gly Cys Thr Gly Glu Val Val Ser Asp Arg Gly Leu Val  
 65 70 75 80  
 Leu Thr Asn His His Cys Gly Tyr Asp Met Ile Gln Ala His Ser Thr  
 85 90 95  
 Leu Glu His Asn Tyr Leu Glu Asn Gly Phe Trp Ala Met Arg Glu Ala  
 100 105 110  
 Asp Glu Leu Pro Asn Lys Asp Ile Ser Val Val Phe Ile Asp Lys Ile  
 115 120 125  
 Glu Asp Val Thr Asp Tyr Val Lys Lys Asp Leu Lys Ala Ile Lys Asp  
 130 135 140  
 Pro Asn Ser Met Asp Tyr Leu Ser Pro Lys Tyr Leu Gln Lys Leu Ala  
 145 150 155 160  
 Asp Lys Lys Ala Gly Lys Asn Phe Ser Ala Lys Asn Pro Gly Leu Ser  
 165 170 175  
 Val Glu Ile Lys Ala Phe Tyr Gly Gly Asn Leu Tyr Leu Met Phe Thr  
 180 185 190  
 Lys Lys Thr Tyr Thr Asp Val Arg Leu Val Gly Ala Pro Pro Thr Ser  
 195 200 205  
 Ile Gly Lys Phe Gly Ala Asp Thr Asp Asn Trp Ile Trp Pro Arg His  
 210 215 220  
 Thr Gly Asp Phe Ser Ile Phe Arg Ile Tyr Ala Asp Lys Asn Gly Asn  
 225 230 235 240  
 Pro Ala Pro Tyr Ser Glu Asp Asn Val Pro Leu Lys Pro Lys Arg Phe  
 245 250 255

Phe Asn Ile Ser<sub>260</sub> Leu Gly Gly Val<sub>265</sub> Gln Glu Asn Asp Tyr Ala<sub>270</sub> Met Ile  
 Met Gly Phe<sub>275</sub> Pro Gly Thr Thr His<sub>280</sub> Arg Tyr Phe Thr Ala<sub>285</sub> Ser Glu Val  
 Asp Glu<sub>290</sub> Trp Lys Ser Ile Asp<sub>295</sub> Asn Asp Ile Arg Ile<sub>300</sub> Arg Met Arg Asp  
 Ile Arg Gln Gly Val<sub>310</sub> Met Leu Arg Glu Met Leu<sub>315</sub> Ala Asp Pro Gln Ile<sub>320</sub>  
 Lys Ile Met Tyr Ser<sub>325</sub> Ala Lys Tyr Ala<sub>330</sub> Ala Ser Gln Asn Ala Tyr<sub>335</sub> Lys  
 Arg Ala Ile Gly<sub>340</sub> Ala Asn Trp Ala Ile<sub>345</sub> Lys Thr Arg Gly Leu<sub>350</sub> Arg Gln  
 Asn Lys Gln<sub>355</sub> Ala Met Gln Asp Arg<sub>360</sub> Leu Ile Ala Trp Gly<sub>365</sub> Ala Lys Gln  
 Gly Thr<sub>370</sub> Pro Arg Tyr Glu Glu<sub>375</sub> Ala Val His Glu Ile<sub>380</sub> Asp Ala Thr Val  
 Ala Lys Arg Ala Asp<sub>390</sub> Leu Arg Arg Arg Tyr Trp<sub>395</sub> Met Ile Glu Glu Gly<sub>400</sub>  
 Ile Ile Arg Gly Ile<sub>405</sub> Glu Phe Ala Arg Ser<sub>410</sub> Pro Ile Pro Thr Glu Asp<sub>415</sub>  
 Glu Thr Lys Ala<sub>420</sub> Leu Gln Gly Asn Asp<sub>425</sub> Ala Ser Ala Arg Lys Glu Ala  
 Ile Asp Lys<sub>435</sub> Ile Arg Thr Arg Tyr<sub>440</sub> Ser Lys Phe Ala Asn<sub>445</sub> Lys Asp Tyr  
 Ser Ala Glu Val Asp Lys Lys<sub>455</sub> Val Ala Val Ala Met<sub>460</sub> Leu Thr Glu Tyr  
 Leu Lys Glu Ile Pro Tyr<sub>470</sub> Glu Asn Leu Pro Leu<sub>475</sub> His Leu Arg Leu Val<sub>480</sub>  
 Lys Asp Arg Phe Ala<sub>485</sub> Gly Asp Val Gln Ala Tyr Val Asp Asp Ile Phe<sub>495</sub>  
 Ala Arg Ser Val<sub>500</sub> Phe Gly Ser Glu Ala<sub>505</sub> Gln Phe Asp Ala Phe Ala Ala<sub>510</sub>  
 Val Pro Ser<sub>515</sub> Val Glu Lys Leu Ala<sub>520</sub> Glu Asp Pro Met Val<sub>525</sub> Leu Phe Ala  
 Ser Ser<sub>530</sub> Val Phe Asp Glu Tyr<sub>535</sub> Arg Lys Leu Tyr Asn Glu Leu Arg Pro  
 Tyr Asp Asp Pro Ile Leu<sub>550</sub> Arg Ala Gln Arg Thr<sub>555</sub> Tyr Ile Ala Gly Leu<sub>560</sub>  
 Leu Glu Met Asp Gly<sub>565</sub> Asp Gln Asp Gln Phe Pro Asp Ala Asn Leu Thr<sub>575</sub>  
 Leu Arg Phe Thr<sub>580</sub> Tyr Gly Gln Val Lys<sub>585</sub> Gly Tyr Ser Pro Arg Asp Asn  
 Val Tyr Tyr Gly His Gln Thr Thr Leu Asp Gly Val Met Glu Lys Glu

595                      600                      605  
 Asp Pro Asp Asn Trp Glu Phe Val Val Asp Pro Lys Leu Lys Ala Val  
 610                      615                      620  
 Tyr Glu Arg Lys Asp Phe Gly Arg Tyr Ala Asp Arg Ser Gly Arg Met  
 625                      630                      635                      640  
 Pro Val Ala Phe Cys Ala Thr Thr His Thr Thr Gly Gly Asn Ser Gly  
 645                      650  
 Ser Pro Val Met Asn Ala Asn Gly Glu Leu Ile Gly Leu Asn Phe Asp  
 660                      665                      670  
 Arg Asn Trp Glu Gly Val Gly Gly Asp Ile Gln Tyr Leu Ala Asp Tyr  
 675                      680                      685  
 Gln Arg Ser Ile Ile Val Asp Ile Arg Tyr Val Leu Leu Val Ile Asp  
 690                      695                      700  
 Lys Val Gly Gly Cys Gln Arg Leu Leu Asp Glu Met Asn Ile Val Pro  
 705                      710                      715                      720

&lt;210&gt; 10

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Synthetic peptide

&lt;400&gt; 10

Trp Ala Gly Gly Asp Ala Ser Gly Glu  
 1                      5

&lt;210&gt; 11

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Synthetic peptide

&lt;400&gt; 11

Ile Ala Arg Arg His Pro Tyr Phe Leu  
 1                      5

&lt;210&gt; 12

&lt;211&gt; 9

<212> PRT

<213> Artificial

<220>

<223> Synthetic peptide

<400> 12

Lys Ile Ala Gly Tyr His Leu Glu Leu  
1 5

<210> 13

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Synthetic peptide

<400> 13

Phe Leu Arg Glu Pro Val Ile Phe Leu  
1 5

<210> 14

<211> 13

<212> PRT

<213> Artificial

<220>

<223> Synthetic peptide

<400> 14

Trp His Trp Leu Glu Leu Lys Pro Gly Glu Pro Met Tyr  
1 5 10

<210> 15

<211> 8

<212> PRT

<213> Artificial

&lt;220&gt;

&lt;223&gt; Synthetic peptide

&lt;400&gt; 15

Ser	Pro	Tyr	Ser	Ser	Glu	Thr	Thr
1				5			

&lt;210&gt; 16

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Synthetic peptide

&lt;400&gt; 16

Ala	Pro	Val	Arg	Ser	Leu
1				5	

&lt;210&gt; 17

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; Synthetic peptide

&lt;400&gt; 17

Gln	Lys	Gln	Met	Ser	Asp	Arg	Arg	Glu
1				5				

&lt;210&gt; 18

&lt;211&gt; 26

&lt;212&gt; PRT

&lt;213&gt; Porphyromonas gingivalis

&lt;400&gt; 18

Ala	Asp	Lys	Gly	Met	Met	Trp	Leu	Leu	Asn	Glu	Leu	Asn	Gln	Glu	Asn
1				5					10					15	

Leu	Asp	Arg	Met	Arg	Glu	Leu	Gly	Phe	Thr
			20					25	

<210> 19  
<211> 6  
<212> PRT  
<213> Porphyromonas gingivalis

<400> 19  
Asp Asn Lys Pro Tyr Lys  
1 5

<210> 20  
<211> 5  
<212> PRT  
<213> Porphyromonas gingivalis

<400> 20  
Glu Met Thr Tyr Leu  
1 5

<210> 21  
<211> 6  
<212> PRT  
<213> Porphyromonas gingivalis

<400> 21  
Phe Ala Gln Phe Ala Asn  
1 5

<210> 22  
<211> 6  
<212> PRT  
<213> Porphyromonas gingivalis

<400> 22  
Val Leu Pro Ala Met Leu  
1 5

<210> 23  
<211> 5



<212> PRT

<213> Porphyromonas gingivalis

<400> 23

Ser Val Val Pro Tyr  
1 5

<210> 24

<211> 6

<212> PRT

<213> Porphyromonas gingivalis

<400> 24

Leu Phe Phe Ala Gly Leu  
1 5

<210> 25

<211> 10

<212> PRT

<213> Artificial

<220>

<223> Consensus sequence

<400> 25

Thr Gly Gly Asn Ser Gly Ser Pro Val Phe  
1 5 10

<210> 26

<211> 9

<212> PRT

<213> Artificial

<220>

<223> Consensus sequence

<400> 26

Thr Gly Gly Asn Ser Gly Ser Pro Val  
1 5