

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11)** Application No. **AU 199743839 B2**  
**(10)** Patent No. **730992**

(54) Title  
**Liquid compositions comprising stability enhancing surfactants and a method of enhancing low temperature stability thereof**

(51)<sup>6</sup> International Patent Classification(s)  
**A61K 007/50**

(21) Application No: **199743839** (22) Application Date: **1997.09.04**

(87) WIPO No: **WO98/13022**

(30) Priority Data

(31) Number	(32) Date	(33) Country
<b>60/026652</b>	<b>1996.09.24</b>	<b>US</b>
<b>60/026650</b>	<b>1996.09.24</b>	<b>US</b>

(43) Publication Date : **1998.05.29**

(43) Publication Journal Date : **1998.07.23**

(44) Accepted Journal Date : **2001.03.22**

(71) Applicant(s)  
**Unilever PLC**

(72) Inventor(s)  
**Sudhakar Puvvada; Richard Kolodziej; May Shana'a**

(74) Agent/Attorney  
**DAVIES COLLISON CAVE,GPO Box 3876,SYDNEY NSW 2001**

(56) Related Art  
**US 5409640**  
**AU 67373/96**  
**US 5478490**

OPI DATE 29/05/98 APPLN. ID 43839/97  
AOJP DATE 23/07/98 PCT NUMBER PCT/EP97/04924



AU9743839

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 6:</b> <b>A61K 7/50</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/13022</b> <b>(43) International Publication Date:</b> 2 April 1998 (02.04.98)
<b>(21) International Application Number:</b> PCT/EP97/04924 <b>(22) International Filing Date:</b> 4 September 1997 (04.09.97) <b>(30) Priority Data:</b> 60/026,652 24 September 1996 (24.09.96) US 60/026,650 24 September 1996 (24.09.96) US <b>(71) Applicant (for AU BB CA GB GH IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW only):</b> UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). <b>(71) Applicant (for all designated States except AU BB CA GB GH IE IL KE LC LK LS MN MW NZ SD SG SL SZ TT UG ZW):</b> UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). <b>(72) Inventors:</b> PUVVADA, Sudhakar; Apartment 5A, 130 Orient Way, Rutherford, NJ 07070 (US). KOLODZIEJ, Richard; Apartment 405, 102, rue de Genève, F-74240 Gaillard (FR). SHANA'A, May; 200 Old Palisade Road, Fort Lee, NJ 07024 (US). <b>(74) Agent:</b> MOLE, Peter, Geoffrey; Unilever PLC, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> LIQUID COMPOSITIONS COMPRISING STABILITY ENHANCING SURFACTANTS AND A METHOD OF ENHANCING LOW TEMPERATURE STABILITY THEREOF		
<b>(57) Abstract</b> <p>The present invention relates to lamellar structured liquid cleansing compositions comprising 5 % to 50 % of a surfactant system comprising (a) an anionic or mixture of anionics and (b) an amphoteric and/or zwitterionic surfactant in mixture, wherein alkalimetal alkylamphoacetate comprises 25 % to 90 % of component (b). A method of enhancing low temperature stability of such lamellar structured liquid cleansing compositions by the selection of an alkalimetal alkylamphoacetate in an amount of 25 % to 90 % of the component (b) is provided. Excellent low temperature stability is achieved.</p>		

LIQUID COMPOSITIONS COMPRISING STABILITY  
ENHANCING SURFACTANTS AND A METHOD OF ENHANCING  
LOW TEMPERATURE STABILITY THEREOF

5

The present invention relates to lamellar structured liquid cleansing compositions such as those described, for example, in applicant's co-pending US Serial No 08/512,010 filed August 7, 1995. These compositions are generally used in skin cleansing or shower gel compositions. In particular, the invention relates to such compositions in which the surfactant system is carefully selected to ensure good stability even at low temperature storage (e.g. -6.7°C (20°F) and below down to -17.8°C (0°F) and to a method of enhancing such low temperature stability by careful selection of the surfactant system.

15

Typically, lamellar structured liquid cleansing composition (e.g. shower gel compositions) comprise a mixture of anionic surfactants (for cleansing and foaming attributes) and mild surfactant. In a typical shower formulation the mild surfactant may be an amphoteric and/or zwitterionic surfactant such as those described in US Serial No 08/512,010 mentioned above, hereby incorporated by reference into the subject application.

In such lamellar structured compositions, however it has been found that there is considerable thinning of product as the product is cooled down to temperatures of -6.7-17.8°C (20° to 0°F). This loss of viscosity is not a desirable property.

Unexpectedly, applicants have found that when alkalimetal alkyl amphotacetate is used as 25% to 90%, preferably 30% to 90% and more preferably about 40% to 90% of the amphoteric and/or zwitterionic



AMENDED SHEET

component, in the surfactant system in lamellar structured compositions, there is a significant increase in product stability. The applicants have also found a method of enhancing low temperature stability of a lamellar  
5 structured liquid cleansing composition by selecting an alkali metal alkylamphoacetate as 25-90% of the amphoteric and/or zwitterionic surfactant in a composition further comprising anionic surfactant.

US Serial No 08/512,010 shows one example (Example IX at page  
10 23) where sodium cocoamphoacetate is used. However, in neither that example or in the other eight examples are there ever taught blends of other amphoteric (e.g. betaine) and amphoacetate. Nor is there any teaching or suggestion in that application that blends of amphoteric will ameliorate low temperature instability in such compositions. Indeed, until  
15 the problem of low temperature instability was even appreciated, it could not have been known that the specifically selected surfactant system of the invention could ameliorate the problem.

WO96/214266 describes mild antibacterial cleansing compositions  
20 containing anionic surfactant, amphoteric surfactant, skin moisturizer antibacterial agent, and water, wherein the anionic and amphoteric surfactants are present at specified ratios and levels of the composition.

WO94/18292 describes personal cleansing compositions comprising  
25 5-50% anionic and/or amphoteric surfactants, 0.1-20% soluble or dispersible ethoxylated anionic surfactant, 1.5-10% water soluble citrate salt, up to 3% perfume or cosmetic oil and water.

The present invention relates to lamellar structured liquid cleansing  
30 compositions comprising 5% to 50% of a surfactant system wherein said



AMENDED SHEET

2a

surfactant system comprises (a) an anionic or mixture of anionics and (b) a blend of amphoteric and/or zwitterionic surfactants wherein said blend comprises alkalimetal alkylamphoacetate and said alkalimetal  
5 alkylamphoacetate comprises 25% to 90%, preferably 30% to 90% and more preferably 40% to 90% of the blend

The present invention further relates to a method of enhancing stability of low temperature compositions (i.e. temperatures of from about  
10 -6.7 to 17.8°C (20°F to about 0°F) in lamellar structured liquid cleansing compositions comprising about 5%



AMENDED SHEET

to about 50% of a surfactant system which surfactant system in turn comprises:

- (a) anionic or mixture of anionic surfactant; and
- 5 (b) an amphoteric and/or zwitterionic surfactant or mixture thereof,

wherein said method comprises selecting component (b) such that the alkalimetal alkyl amphoacetate comprises greater than 25% to 90%, preferably about 30% to 90%, more preferably about 40% to 90% of said  
10 component (b).

Unexpectedly, the applicants have found that when the alkalimetal alkylamphoacetate is used as at least a minimum amount of amphoteric/zwitterionic blend, the component significantly enhances cold  
15 temperature stability of the lamellar structured composition relative to compositions where the alkalimetal alkylamphoacetate does not comprise a portion or comprise less than 25% of the amphoteric/zwitterionic blend.

The present invention is directed towards improving the low  
20 temperature stability of lamellar structured liquid compositions comprising an amphoteric and/or zwitterionic surfactant or mixture thereof.

According to one aspect the present invention provides lamellar structured liquid cleansing compositions comprising about 5% to about  
25 50% of a surfactant system which surfactant system in turn comprises:

- (a) anionic or mixture of anionic surfactant; and
- (b) an amphoteric and/or zwitterionic surfactant or mixture thereof;



AMENDED SHEET

3a

wherein alkalimetal alkyl amphoacetate comprises greater than 25% to 90%, preferably about 30% to 90%, more preferably about 40% to 90% of component (b); and

- 5 (c) a structurant selected from saturated or unsaturated, branched or unbranched C<sub>8</sub>-C<sub>24</sub> liquid fatty acids, or polyalkylene glycol fatty acid esters, capric or caprylic acid, or mixtures thereof.



AMENDED SHEET

According to a further aspect the present invention provides a method of enhancing low temperature stability of a lamellar structured liquid cleansing compositions comprising 5% to 50% of a surfactant system which surfactant  
5 system comprises:

- (a) anionic or mixture of anionic surfactant; and
- (b) an amphoteric and/or zwitterionic surfactant or mixture thereof;

wherein said method comprises selecting the  
10 amphoteric and/or zwitterionic surfactant component (b) such that alkalimetal alkyl amphotoacetate comprises greater than 25% to 90% of said component (b).

According to a further aspect the present invention  
15 provides a lamellar structured liquid cleansing composition comprising 5% to 50% of a surfactant system comprising:

- a) anionic or mixture of anionic surfactants; and
- b) an amphoteric and/or zwitterionic surfactant or mixture thereof;

wherein alkalimetal alkylamphotoacetate comprises 25% to  
20 90% of component (b); and

- c) a structurant selected from liquid fatty acids comprising oleic acid, isostearic acid, linoleic acid, ricinoleic acid, elaidic acid, arichodonic acid, myristoleic acid, palmitoleic acid and  
25 mixtures thereof, or

prolyalkylene glycol fatty acid esters comprising propylene glycol isostearate, propylene glycol oleate, glyceryl isostearate, glyceryl oleate, polyglyceryl diisostearate, and mixtures thereof.


30 The anionic surfactant may be, for example, an aliphatic sulfonate, such as a primary alkane (e.g.,  $C_8-C_{22}$ )





sulfonate, primary alkane (e.g.,  $C_8-C_{22}$ ) disulfonate,  $C_8-C_{22}$  alkene sulfonate,  $C_8-C_{22}$  hydroxyalkane sulfonate or alkyl glyceryl ether sulfonate (AGS); or an aromatic sulfonate such as alkyl benzene sulfonate.

The anionic surfactant may also be an alkyl sulfate (e.g.,  $C_{12}-C_{18}$  alkyl sulfate) or alkyl ether sulfate (including alkyl glyceryl ether sulfates). Among the alkyl ether sulfates are those having the formula:



wherein R is an alkyl or alkenyl having 8 to 18 carbons, preferably 12 to 18 carbons, n has an average value of greater than 1.0, preferably between 2 and 3; and M is a solubilizing cation such as sodium, potassium, ammonium or substituted ammonium. Ammonium and sodium lauryl ether sulfates are preferred.

55


55




- 5 -

The anionic surfactant may also be alkyl  
 sulfosuccinates (including mono- and dialkyl, e.g., C<sub>6</sub>-C<sub>22</sub>  
 sulfosuccinates); alkyl and acyl taurates, alkyl and acyl  
 sarcosinates, sulfoacetates, C<sub>8</sub>-C<sub>22</sub> alkyl phosphates and  
 5 phosphates, alkyl phosphate esters and alkoxyalkyl  
 phosphate esters, acyl lactates, C<sub>8</sub>-C<sub>22</sub> monoalkyl succinates  
 and maleates, sulphoacetates, and acyl isethionates.

Sulfosuccinates may be monoalkyl sulfosuccinates having  
 10 the formula:




amido-MEA sulfosuccinates of the formula  
 15




wherein R<sup>4</sup> ranges from C<sub>8</sub>-C<sub>22</sub> alkyl and M is a  
 solubilizing cation;  
 20

amido-MIPA sulfosuccinates of formula



25 where M is as defined above.

Also included are the alkoxyated citrate  
 sulfosuccinates; and alkoxyated sulfosuccinates such as the  
 following:



wherein n = 1 to 20; and M is as defined above.


- 6 -

Sarcosinates are generally indicated by the formula  $RCON(CH_3)CH_2CO_2M$ , wherein R ranges from  $C_8$  to  $C_{20}$  alkyl and M is a solubilizing cation.

5 Taurates are generally identified by formula  $R^2CONR^1CH_2CH_2SO_3M$

wherein  $R^2$  ranges from  $C_8$ - $C_{20}$  alkyl,  $R^1$  ranges from  $C_1$ - $C_4$  alkyl and M is a solubilizing cation.

10 Another class of anionic surfactants are carboxylates such as follows:



15 wherein R is  $C_8$  to  $C_{20}$  alkyl; n is 0 to 20; and M is as defined above.

Another carboxylate which can be used is amido alkyl polypeptide carboxylates such as, for example, Monteine


20 LCQ<sup>(R)</sup> by Seppic.

Another anionic surfactant which may be used are the  $C_8$ - $C_{18}$  acyl isethionates. These esters are prepared by reaction between alkali metal isethionate with mixed aliphatic fatty acids having from 6 to 18 carbon atoms and an iodine value of less than 20. At least 75% of the mixed fatty acids have from 12 to 18 carbon atoms and up to 25% have from 6 to 10 carbon atoms.

30 Acyl isethionates, when present, will generally range from about 0.5-15% by weight of the total composition. Preferably, this component is present in an amount from 1 to 10%.

- 7 -

The acyl isethionate may be an alkoxyated isethionate such as is described in Ilardi et al., U.S. Patent No. 5,393,466, hereby incorporated by reference into the subject application. This compound has the general formula:




wherein R is an alkyl group having 8 to 18 carbons, m is an integer from 1 to 4, X and Y are hydrogen or an alkyl group having 1 to 4 carbons and M<sup>+</sup> is a monovalent cation such as, for example, sodium, potassium or ammonium.

In general the anionic surfactant component will comprise from 1 to 20% by weight of the composition, preferably 2 to 15%, most preferably 5 to 12% by weight of the composition.

The anionic surfactant mixture may comprise alkyl sulphates, acylisethionates and mixtures thereof.

The compositions of the present invention further comprise zwitterionic surfactants. Zwitterionic surfactants are exemplified by those which can be broadly described as derivatives of aliphatic quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight or branched chain, and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. A general formula for these compounds is:




wherein  $R^2$  contains an alkyl, alkenyl, or hydroxy alkyl radical of from about 8 to about 18 carbon atoms, from 0 to about 10 ethylene oxide moieties and from 0 to about 1 glyceryl moiety; Y is selected from the group consisting of nitrogen, phosphorus, and sulfur atoms;  $R^3$  is an alkyl or monohydroxyalkyl group containing about 1 to about 3 carbon atoms; X is 1 when Y is a sulfur atom, and 2 when Y is a nitrogen or phosphorus atom;  $R^4$  is an alkylene or hydroxyalkylene of from about 1 to about 4 carbon atoms and Z is a radical selected from the group consisting of carboxylate, sulfonate, sulfate, phosphonate, and phosphate groups.

Examples of such surfactants include:

- 15 4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate;
- 5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxypentane-1-sulfate;
- 3-[P,P-diethyl-P-3,6,9-trioxatetradecylphosphonio]-2-hydroxypropane-1-phosphate;
- 20 3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropylammonio]-propane-1-phosphonate;
- 3-(N,N-dimethyl-N-hexadecylammonio)propane-1-sulfonate;
- 3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxypropane-1-sulfonate;
- 25 4-[N,N-di(2-hydroxyethyl)-N-(2-hydroxydodecyl)ammonio]-butane-1-carboxylate;
- 3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-propane-1-phosphate;
- 30 3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-phosphonate; and
- 5-[N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-hydroxy-pentane-1-sulfate.

- 9 -

Amphoteric detergents which may be used in this invention include at least one acid group. This may be a carboxylic or a sulphonic acid group. They include quaternary nitrogen and therefore are quaternary amido acids. They should generally include an alkyl or alkenyl group of 7 to 18 carbon atoms. They will usually comply with an overall structural formula:



where R<sup>1</sup> is alkyl or alkenyl of 7 to 18 carbon atoms;

R<sup>2</sup> and R<sup>3</sup> are each independently alkyl, hydroxyalkyl or carboxyalkyl of 1 to 3 carbon atoms;


n is 2 to 4;

m is 0 to 1;

X is alkylene of 1 to 3 carbon atoms optionally substituted with hydroxyl, and


Y is -CO<sub>2</sub>- or -SO<sub>3</sub>-

Suitable amphoteric detergents within the above general formula include simple betaines of formula:



and amido betaines of formula:

- 10 -




where m is 2 or 3.


10 Betaines are typically present in the compositions of the present inventions in amounts of 0 to 25% by weight, for example, 0.1 to 25% by weight.

In both formulae  $R^1$ ,  $R^2$  and  $R^3$  are as defined previously.  $R^1$  may in particular be a mixture of  $C_{12}$  and  $C_{14}$  15 alkyl groups derived from coconut so that at least half, preferably at least three quarters of the groups  $R^1$  have 10 to 14 carbon atoms.  $R^2$  and  $R^3$  are preferably methyl.


20 A further possibility is that the amphoteric detergent is a sulphobetaine of formula



or



35 where m is 2 or 3, or variants of these in which  $(\text{CH}_2)_3\text{SO}_3^-$  is replaced by



- 11 -

In these formulae  $R^1$ ,  $R^2$  and  $R^3$  are as discussed previously.

The amphoteric/zwitterionic surfactant generally  
5 comprises 0.1 to 20% by weight, preferably 5% to 15% of the composition.

A critical aspect of this invention is that the  
zwitterionic/amphoteric compounds must be used in blends of  
10 zwitterionic/amphoteric wherein one component of the blend is an alkalimetal alkylamphoacetate. Further, the alkali metal alkylamphoacetate must comprise 25% to 90%, preferably 30% to 90%, more preferably 40% to 90% of the blend. Suitably the amount of alkalimetal alkylamphoacetate may be  
15 within the range 30 to 75%, for example 40 to 60%, of the blend.

Examples of alkalimetal alkyl amphoacetate compounds include, but are not limited to, sodium or potassium lauro  
20 or cocoamphoacetate.

The total amount of amphoteric/zwitterionic including the amphoacetate, preferably should be no greater than 20%, more preferably no greater than 15%. The total  
25 amphoteric/zwitterionic should comprise at least 5% of the composition.

In addition to one or more anionic and amphoteric and/or zwitterionic, the surfactant system may optionally  
30 comprise a nonionic surfactant.

The nonionic which may be used includes in particular the reaction products of compounds having a hydrophobic group and a reactive hydrogen atom, for example aliphatic  
35 alcohols, acids, amides or alkyl phenols with alkylene




- 12 -

oxides, especially ethylene oxide either alone or with propylene oxide. Specific nonionic detergent compounds are alkyl ( $C_6-C_{22}$ ) phenols-ethylene oxide condensates, the condensation products of aliphatic ( $C_6-C_{18}$ ) primary or  
 5 secondary linear or branched alcohols with ethylene oxide, and products made by condensation of ethylene oxide with the reaction products of propylene oxide and ethylenediamine. Other so-called nonionic detergent compounds include long chain tertiary amine oxides, long chain tertiary phosphine  
 10 oxides and dialkyl sulphoxides.

The nonionic may also be a sugar amide, such as a polysaccharide amide. Specifically, the surfactant may be one of the lactobionamides described in U.S. Patent No.  
 15 5,389,279 to Au et al. which is hereby incorporated by reference or it may be one of the sugar amides described in Patent No. 5,009,814 to Kelkenberg, hereby incorporated into the subject application by reference.

20 Other surfactants which may be used are described in U.S. Patent No. 3,723,325 to Parran Jr. and alkyl polysaccharide nonionic surfactants as disclosed in U.S. Patent No. 4,565,647 to Llenado, both of which are also  
 25 incorporated into the subject application by reference.

Preferred alkyl polysaccharides are alkylpolyglycosides of the formula



30 wherein  $R^2$  is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 0 to 3, preferably 2; t is from 0 to about 10,

- 13 -

preferably 0; and x is from 1.3 to about 10, preferably from 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted  
5 with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4- and/or 6-position, preferably predominantly the 2-position.

10

Nonionic comprises 0 to 10% by wt. of the composition.

In general, the compositions of the invention are soap-free compositions.

15

The present invention provides compositions utilizing typically 0.1% to 15% by wt., preferably 1 to 10% by wt. of a structuring agent which works in the compositions to form a lamellar phase. Such lamellar phase is preferred because  
20 it enables the compositions to suspend particles more readily (e.g., emollient particles) while still maintaining good shear thinning properties. The lamellar phase also provides consumers with desired rheology ("heaping").

25

More particularly, where the composition is not lamellar structured and enhanced particle suspension/enhancing is desired, it is usually necessary to add external structurants such as carbomers (e.g., cross-linked polyacrylate such as Carbopol<sup>(R)</sup>) and clays. However,  
30 these external structurants have poorer shear thinning properties that significantly reduce consumer acceptability.

The structurant is generally an unsaturated and/or branched long chain (C<sub>8</sub>-C<sub>24</sub>) liquid fatty acid or ester  
35 derivative thereof; and/or unsaturated and/or branched long

chain liquid alcohol or ether derivatives thereof. It may also be a short chain saturated fatty acid such as capric acid or caprylic acid. While not wishing to be bound by theory, it is believed that the unsaturated part of the fatty acid of alcohol or the branched part of the fatty acid or alcohol acts to "disorder" the surfactant hydrophobic chains and induce formation of lamellar phase.

Examples of liquid fatty acids which may be used are oleic acid, isostearic acid, linoleic acid, linolenic acid, ricinoleic acid, elaidic acid, arachidonic acid, myristoleic acid and palmitoleic acid. Ester derivatives include propylene glycol isostearate, propylene glycol oleate, glyceryl isostearate, glyceryl oleate and polyglyceryl diisostearate.

Examples of alcohols include oleyl alcohol and isostearyl alcohol. Examples of ether derivatives include isosteareth or oleth carboxylic acid; or isosteareth or oleth alcohol.

The structuring agent may be defined as having melting point below about 25 C centigrade.

One of the principle benefits of the invention is the ability to suspend oil/emollient particles in a lamellar phase composition.

Various classes of oils are set forth below.

Vegetable oils: Arachis oil, castor oil, cocoa butter, coconut oil, corn oil, cotton seed oil, olive oil, palm kernel oil, rapeseed oil, safflower seed oil, sesame seed oil and soybean oil.

- 15 -

Esters: Butyl myristate, cetyl palmitate, decyloleate, glyceryl laurate, glyceryl ricinoleate, glyceryl stearate, glyceryl isostearate, hexyl laurate, isobutyl palmitate, isocetyl stearate, isopropyl isostearate, isopropyl laurate,  
5 isopropyl linoleate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, propylene glycol monolaurate, propylene glycol ricinoleate, propylene glycol stearate, and propylene glycol isostearate.

10 Animal Fats: Acetylated lanolin alcohols, lanolin, lard, mink oil and tallow.

Fatty acids and alcohols: Behenic acid, palmitic acid, stearic acid, behenyl alcohol, cetyl alcohol, eicosanyl  
15 alcohol and isocetyl alcohol.

Other examples of oil/emollients include mineral oil, petrolatum, silicone oil such as dimethyl polysiloxane, lauryl and myristyl lactate.

20 It should be understood that where the emollient may also function as a structurant, it should not be doubly included such that, for example, if the structurant is 15% oleyl alcohol, no more than 5% oleyl alcohol as "emollient"  
25 would be added since the emollient (whether functioning as emollient or structurant) never comprises more than 20%, preferably no more than 15% of the composition.

The emollient/oil is generally used in an amount of 1  
30 to 20%, preferably 1 to 15% by wt. of the composition. Generally, it should comprise no more than 20% of the composition.

In addition, the compositions of the invention may  
35 include optional ingredients as follows:

- 16 -

Organic solvents, such as ethanol; auxiliary thickeners, such as carboxymethylcellulose, magnesium aluminum silicate, hydroxyethylcellulose, methylcellulose, carbopols, glucamides, or Antil<sup>(R)</sup> from Rhone Poulenc; perfumes; sequestering agents, such as tetrasodium ethylenediaminetetraacetate (EDTA), EHDP or mixtures in an amount of 0.01 to 1%, preferably 0.01 to 0.05%; and coloring agents, opacifiers and pearlizers such as zinc stearate, magnesium stearate, TiO<sub>2</sub>, EGMS (ethylene glycol monostearate) or Lytron 621 (Styrene/Acrylate copolymer); all of which are useful in enhancing the appearance or cosmetic properties of the product.

The compositions may further comprise antimicrobials such as 2-hydroxy-4,2'4' trichlorodiphenylether (DP300); preservatives such as dimethyloldimethylhydantoin (Glydant XL1000), parabens, sorbic acid etc.

The compositions may also comprise coconut acyl mono- or diethanol amides as suds boosters, and strongly ionizing salts such as sodium chloride and sodium sulfate may also be used to advantage.

Antioxidants such as, for example, butylated hydroxytoluene (BHT) may be used advantageously in amounts of 0.01% or higher if appropriate.

Cationic conditioners which may be used include Quatrisoft LM-200 Polyquaternium-24, Merquat Plus 3330 - Polyquaternium 39; and Jaguar<sup>(R)</sup> type conditioners.

Polyethylene glycols which may be used include:

Polyox	WSR-205	PEG 14M,
Polyox	WSR-N-60K	PEG 45M, or

- 17 -

Polyox WSR-N-750 PEG 7M.

Thickeners which may be used include Amerchol Polymer HM 1500 (Nonoxynyl Hydroethyl Cellulose); Glucam DOE 120  
5 (PEG 120 Methyl Glucose Dioleate); Rewoderm<sup>®</sup> (PEG modified glyceryl cocoate, palmate or tallowate) from Rewo Chemicals; Antil<sup>®</sup> 141 (from Goldschmidt). A particularly preferred thickener is xanthan gum. Indeed, xanthan gum, particularly  
10 when used with the surfactant system of the invention, also helps ameliorate cold storage instability.

Another optional ingredient which may be added are the defloculating polymers such as are taught in U.S. Patent No. 5,147,576 to Montague, hereby incorporated by reference.  
15

Another ingredient which may be included are exfoliants such as polyoxyethylene beads, walnut sheets and apricot seeds.

20 The compositions of the present invention are typically personal products but are not to be construed as strictly limited thereto.

The invention will be described in greater detail by  
25 way of the following non-limiting examples. The examples are for illustrative purposes only and not intended to limit invention in any way. Further modifications within the scope of the present invention will be obvious to the skilled man.

30 Figure 1 shows cold temperature stability of various lamellar structured liquid cleansers at 15 and 0°F. As seen, when alkali metal amphoteric comprises 25% of all amphoteric (for example, amphoteric plus betaine),  
35 stability is dramatically increased.

All percentages in the specification and examples are by weight unless stated otherwise.

EXAMPLES

5

The following compositions are used in the examples:

Ingredients	I	II	III
Cocoamido Propyl Betaine	12	9	6
Sodium Lauroamphoacetate	0	3	6
Sodium Cocoyl Isethionate	6.5	6.5	6.5
Sodium Laureth Sulfate	6.5	6.5	6.5
Thickener/Polymer (e.g., Cationic, Guar or Xanthan Gum)	0.1 to 1%	0.1 to 1%	0.1 to 1%
Emollient	1 to 7%	1 to 7%	1 to 7%
Structurant	3 to 10%	3 to 10%	3 to 10%
Titanium Dioxide	0.2	0.2	0.2
DMDM Hydantoin	0.2	0.2	0.2
Fragrance	1.0	1.0	1.0
BHT	0.0075	0.0075	0.0075
Water	to 100.0	to 100.0	to 100.0

10

EXAMPLES 1-3

The compositions shown above in I-III were stored in plastic cups at 15°F and 0°F for a period of 1 day and then equilibrated back to room temperature. Care was taken not to disturb the sample since viscosity increase when these

15



products are shaken. The viscosity of the sample is then measured using a Brookfield RV Viscometer attached to a helipath accessory and using T-Bar Spindle A.

5

The results are set forth in Table I below:

Example	% Amphoacetate in Betaine/ Amphoacetate Blend	T-Bar Viscosity		
		Room Temperature	After 1 Day at -9°C (15°F)	After 1 Day at -17.8°C (0°F)
1	0	88400	22800	22400
2	25	91200	26000	33200
3	50	87200	84000	83200

As seen from the Table and from Figure 1, (Examples 2 and 3),  
10 when amphoacetate comprises about 25% and greater, preferably about 30% to 90% and most preferably about 40% to 90% of blend of amphoteric (betaine/amphoacetate blend), viscosity at low temperature - 9°C (15°F), -17.8°C (0°F) remains much higher. Thus, clearly, low  
temperature viscosity/phase stability is much superior relative to  
15 compositions in which amphoacetate is not used or comprises less than 25% of the blend (e.g. Example 1).



AMENDED SHEET



Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a  
5 stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference to any prior art in this specification is  
10 not, and should not be taken as, an acknowledgment or any form of suggestion that that prior art forms part of the common general knowledge in Australia.

3  
5  
7  
9



The Claims defining the invention are as follows:

1. A lamellar structured liquid cleansing composition comprising 5% to 50% of a surfactant system comprising:
  - 5 a) anionic or mixture of anionic surfactants; and
  - b) an amphoteric and/or zwitterionic surfactant or mixture thereof;wherein alkalimetal alkylamphoacetate comprises 25% to 90% of component (b); and
  - 10 c) a structurant selected from liquid fatty acids comprising oleic acid, isostearic acid, linoleic acid, ricinoleic acid, elaidic acid, arichodonic acid, myristoleic acid, palmitoleic acid and mixtures thereof, or
  - 15 polyalkylene glycol fatty acid esters comprising propylene glycol isostearate, propylene glycol oleate, glyceryl isostearate, glyceryl oleate, polyglyceryl diisostearate, and mixtures thereof.
- 20 2. A composition according to Claim 1 wherein alkalimetal alkylamphoacetate comprises 30 to 90% of component (b).
3. A composition according to Claim 2, wherein alkalimetal alkylamphoacetate comprises 40% to 90% of components (b).
- 25 4. A composition according to Claim 1, wherein the anionic surfactant is selected from the group consisting of alkyl sulfates, acylisethionates and mixtures thereof.
- 30 5. A composition according to Claim 1, wherein component (b) comprises 0.1% to 25% betaine.



6. A composition according to Claim 1, wherein the composition additionally comprises 0% to 10% of nonionic surfactant.
- 5 7. A method of enhancing low temperature stability of a lamellar structured cleansing composition as claimed in Claim 1 wherein said method comprises selecting the amphoteric and/or zwitterionic surfactant component (b) such that alkalimetal alkyl amphotoacetate comprises 25% to 90% of  
10 component (b).
8. A method according to Claim 7, wherein alkalimetal alkylamphoacetate comprises 30% to 90% of component (b).
- 15 9. A method according to Claim 8, wherein alkali metal alkylamphoacetate comprises 40% to 90% of component (b).
10. A method according to any one of Claims 7 to 9 wherein the lamellar structured liquid cleansing composition  
20 comprises one or more of the components of Claims 4 to 6.
11. A composition as hereinbefore described with reference to the examples.

DATED THIS 22nd day of November, 2000.

UNILEVER PLC

By Its Patent Attorneys

DAVIES COLLISON CAVE



Fig.1.

