



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A01N 33/12, 33/08, 25/30 // (A01N 33/08, 61:00, 59:24, 59:06, 59:00, 47:28, 25:30) (A01N 33/12, 61:00, 59:24, 59:06, 59:00, 47 :28, 25 :30)</p>	A1	<p>(11) International Publication Number: WO 96/01049</p> <p>(43) International Publication Date: 18 January 1996 (18.01.96)</p>
<p>(21) International Application Number: PCT/EP95/02575</p> <p>(22) International Filing Date: 3 July 1995 (03.07.95)</p> <p>(30) Priority Data: 08/270,857 5 July 1994 (05.07.94) US</p> <p>(71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): PARR, William, John, Ernest [GB/NL]; Emmastraat 33, NL-7411 EK Deventer (NL). BUTSELAAR, Robert, Jan [NL/NL]; Zwinglilaan 40, NL-1216 ME Hilversum (NL). NORTH, Michael, Shaun [ZA/NL]; Velperweg 76, NL-6824 BM Arnhem (NL).</p> <p>(74) Agent: SCHALKWIJK, Pieter, Cornelis; Akzo Nobel N.V., Patent Dept. (Dept. APTA), P.O. Box 9300, NL-6800 SB Arnhem (NL).</p>	<p>(81) Designated States: AU, BR, MX, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: ACTIVITY PROMOTING ADDITIVES FOR REST-BREAKING AGENTS</p>		
<p>(57) Abstract</p> <p>Activity promoting additives for rest-breaking agents are disclosed. Further, rest-breaking compositions comprising these additives and a rest-breaking agent are also disclosed. Finally, a process for breaking the rest of bushes, shrubs, vines, nuts, berries and non-deciduous fruit trees is disclosed. The activity promoting additives enhance the activity of rest-breaking agents thereby leading to improvements in the yields and quality of fruit, nuts and berries in regions which have mild winter weather conditions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Activity Promoting Additives for Rest-Breaking Agents

The present invention relates to the use of particular compounds as activity promoting additives for rest-breaking agents, to rest-breaking compositions comprising these additives and to the use of these compositions for rest-breaking of bushes, shrubs, vines, nuts, berries and non-deciduous fruit trees.

Background of the invention

Fruit and nut trees, as well as bushes, shrubs, vines and berries, require winter chilling to grow normally. The amount of chilling required depends upon the kind of fruit and the cultivar. If winter chilling is insufficient, then growth abnormalities such as delayed and uneven blossoming, poor leaf cover, insufficient fruit-set and reduced fruit size can occur. These symptoms are generally referred to as delayed foliation according to the article, Blommaert, K.L.J., "Winter Dormancy and Delayed Foliation," The Deciduous Fruit Grower, (1956).

Measures to reduce the symptoms of delayed foliation include treatment with high volumes of chemical rest-breaking agents during later winter and various physical manipulations such as pruning.

In South Africa, for example, most apple trees receive insufficient winter chilling to break rest completely and thus annual application of a chemical rest-breaking agent is standard practice. The most widely used rest-breaking agent in commercial apple orchards is dinitro-ortho-cresol (DNOC) in combination with winter-oil emulsion (a lipophilic agent) (Erez, A. et al., "Improved Methods of Breaking Rest in the Peach and Other Deciduous Fruit Species," J. Amer. Soc. Hort. Sci., 96, pp. 519-522 (1971)). This article also mentions the use of the potassium salt of gibberellic acid, kinetin, indoleacetic acid and thiourea as potential rest-breaking agents.

5 DNOC is a non-systemic stomach poison and contact insecticide, ovicidal to the eggs of certain insects. It is strongly phytotoxic and its permissible use as an insecticide is limited to dormant sprays or on waste ground. Further, DNOC is known to act as a cumulative
10 poison in man. Although DNOC is relatively inexpensive and effective as a rest-breaking agent, it is toxic enough to limit its continued use and it is currently on the European Red List of agricultural chemicals which will be prohibited in the future. DNOC is also banned in the United States for use as an agricultural chemical.

15 One of the most effective alternatives to DNOC is hydrogen cyanamide according to the publications, Snir, I., "Chemical Dormancy Breaking of Red Raspberry," HortScience, 18, pp. 719-713 (1983); North, M.S., "Effects of Cyanamide and DNOC/oil on Budbreak, Yield and Fruit Size of Golden Delicious Apples," S. Afr. J. Plant Soil, 6(3), pp. 176-178 (1989); Stadler, J.D., North, M.S. and Lutze, G.F.A., "Artificial Rest-Breaking of Apricot and Plum Cultivars Using Hydrogen Cyanamide,"
20 J.S.Afr.Soc.Hort.Sci., 1(1), pp. 9-11, (1991); Nee, C.C. and Fuchigami, L.H., "Overcoming Rest at Different Growth with Hydrogen Cyanamide," Scientia Horticulturae, 50, pp. 107-113 (1992); and North, M.S., "Alternative Rest-Breaking Agents to DNOC/oil for Apples," S. Afr. J. Plant Soil, 9(1), pp. 39-40 (1992).

25 Hydrogen cyanamide is a skin and eye irritant and is especially acute when used in combination with the consumption of alcohol. The toxicity and relatively high price limit its market acceptance and hydrogen cyanamide has also been placed on the European Red List.

30 Thus, there exists an immediate need for new, milder rest-breaking agents, not only in South Africa, but also in countries which are currently developing agricultural industries but do not have an ideal climate therefor, such as Brazil. Further, there is a need for such new rest-breaking agents in countries with existing agricultural

industries that, up to now, did not appreciate the extent of the problem of delayed foliation.

5 The response of a plant to a rest-breaking agent is dependent upon the chemical composition of the agent, its application rate and timing and on the nutritional status of the plant according to Terblanche, J.H. and Strydom, D.K., "Effects of Autumnal Nitrogen Nutrition, Urea Sprays and a Winter Rest-Breaking Spray on Budbreak and Blossoming of Young 'Golden Delicious' Trees Grown in Sand Culture," Deciduous Fruit
10 Grower, 23 pp. 8-14 (1973).

Many chemicals have been found to have rest-breaking ability. A summary of some of these chemicals can be found in Saure, M.C., "Dormancy Release in Deciduous Fruit Trees," Janick, J. (Ed.),
15 Horticultural Reviews, 7, pp. 239-287, AVI Publishing Co. Inc., Westport, Connecticut (1985). The efficacy of these various rest-breaking agents appears to be directly related to their physiological harshness.

20 One of the milder rest-breaking agents is potassium nitrate which has been shown to have a positive effect on peaches. Generally, fruit trees having a lower chill requirement, such as peaches, also require lower concentrations of rest-breaking agents, whereas fruits with a higher chill requirement, such as apples, require excessive
25 concentrations of a mild rest-breaking agent. Articles demonstrating the effects of thiourea, potassium nitrate and combinations thereof include, Wolak, R.J. and Couvillon, G.A., "Time of Thiourea-KNO₃ Application on the Rest Requirement and Bud Development in 'Loring' Peach," HortScience, 11(4), pp. 400-402, (1976) and Fernandez-Escobar, R. and Martin, R., "Chemical Treatments for Breaking Rest in Peach in
30 Relation to Accumulated Chilling," J. Hort. Sci., 62(4), pp. 457-461, (1987).


For the foregoing reasons, there exists a need in the art for improved rest-breaking agents which are, affordable, effective, less toxic than the present rest-breaking agents and which can be employed in reasonable concentrations.

5

Summary of the Invention

The present invention relates to a process for enhancing the rest-breaking in bushes, shrubs, vines, nuts, berries and non-deciduous fruit trees which comprises the step of applying to at least one of these plants, before blossom, an effective amount of at least one rest-breaking agent and a compound selected from the group consisting of alkoxyated amines represented by the following general formula:


15



20

wherein n is an integer from 1 to 50, A represents an alkylene group and when n>1, each A may be the same or different alkylene groups, R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms and groups represented by the formula:


25




30

wherein m is an integer from 1-50, A₁ represents an alkylene group and when n>1, each A₁ may be the same or different alkylene groups, x is an integer from 1-6, and R₂ is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms, R₃ is selected from hydrogen, 1-8 carbon atom straight or branched chain alkyl and alkenyl groups and aryl groups having up to 8 carbon atoms;


and R_1 is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-22 carbon atoms, a group represented by the formula:



wherein n' is an integer from 1 to 50 and A_2 represents an alkylene group and when $n' > 1$ each A_2 may be the same or different alkylene groups; and alkoxyated quaternary ammonium compounds represented by the following general formula:



wherein n , A , R_1 and R_3 are as defined above, X is an anion, R_5 is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 4 carbon atoms, and benzyl, or R_5X^- is carboxymethyl as in betaines or oxygen as in amine oxides; R_4 is selected from straight or branched chain alkyl and alkenyl groups having 8-22 carbon atoms and groups represented by the formula:



wherein A_1 , m , R_2 , R_3 , R_5 and x are as defined above.

30

More particularly, X may be halides such as Cl^- , Br^- , $CH_3SO_4^-$, and $C_2H_5SO_4^-$, among others. The anion associated with these quaternary

ammonium compounds is not critical to the process of the present invention.

5 These compounds are known from European patent application publication number 0 463 241 where it is disclosed that these compounds are useful as blossom thinning agents for stone fruits. In addition, a small group of the foregoing compounds are known from European patent 0 257 686 which discloses several alkoxyated amines and their use as activity promoting additives for herbicides and fungicides. Also 10 disclosed therein is a method for making these compounds, which method is hereby incorporated by reference.

As a result of extensive research it has now been found that the above-identified compounds can be used as adjuvants for chemical rest-breaking agents for plants. These compounds exhibit a substantial 15 effect when combined with rest-breaking agents, and human toxicity and phytotoxicity studies have shown that these compounds have acceptably low levels of toxicity to humans and other plants. Further, the compounds do not cause significant harm to useful insect populations. Finally, the activity-enhancing effect of these compounds allows the 20 use of milder rest-breaking agents which are less toxic, as well as the use of lower concentrations of rest-breaking agents.

The amino compounds of the present invention may be prepared by reacting an amine selected from the group consisting of $R-NH_2$, $RRNH$, and $R-NH-((CH_2)_x-NH)_n-R'$ wherein R and R' are aliphatic hydrocarbon groups having 8-22 carbon atoms, $n=1-5$ and x is an integer from 1-6; 25 with at least one alkylene oxide.

30 The preferred alkylene oxides for use in the present invention are ethylene oxide, propylene oxide, isobutylene oxide and butylene oxide. The compounds of the present invention are made in such a way as to introduce varying numbers of alkylene oxide units onto the amino

nitrogen. Thus, these alkylene oxide groups may be all the same, such as, for example, one or more ethylene oxide units, or the groups may be different to form, for example, block copolymer chains of ethylene oxide and propylene oxide units, random copolymer chains consisting of several units of each of two or more different alkylene oxides, or alternating units of two or more alkylene oxides. Any conceivable combination of alkylene oxide units up to 50 units long may be employed at each location on the amino nitrogen which is to contain such units. In addition, a single amino nitrogen may contain two different alkylene oxide chains attached thereto or two chains which are the same.

In the most preferred embodiments of the present invention, block copolymer chains of ethylene oxide and one or more of propylene oxide or butylene oxide are employed. Preferably, the molar weight of the compounds used in the present invention is less than 8000 though higher molecular weight compounds can be employed in some circumstances.

The amino compounds can also be quaternized by known quaternization methods to produce quaternary ammonium compounds which are also useful in the process of the present invention. The fourth substituent added to the amino nitrogen by quaternization may be an alkyl, aryl or alkenyl group having 1 to 4 carbon atoms. The anion associated with such quaternary ammonium compounds is not critical to the process of the present invention.

Typical compounds suitable for use in the process of the present invention include, but are not limited to, cocobis (2-hydroxyethyl)methylammonium chloride, polyoxyethylene (15) cocomethylammonium chloride, oleylbis (2-hydroxyethyl)methylammonium chloride, polyoxyethylene (15) stearyl methylammonium chloride, cocobis (2-hydroxyethyl)amine, polyoxyethylene(5)cocoamine,

polyoxyethylene(15)cocoamine, tallowbis (2-hydroxyethyl)amine,
polyoxyethylene(5)tallowamine, polyoxyethylene(15)tallowamine,
tallow/oleylbis(2-hydroxyethyl)amine, oleylbis(2-hydroxyethyl)amine,
polyoxyethylene(5)oleylamine, polyoxyethylene(15)oleylamine,
5 hydrogenated tallowbis(2-hydroxyethyl)amine,
hydrogenated polyoxyethylene(5)tallowamine,
hydrogenated polyoxyethylene(15)tallowamine,
hydrogenated polyoxyethylene(50)tallowamine,
N,N',N'-tris(2-hydroxyethyl)-N-tallow-1,3-diaminopropane,
10 N,N',N'-polyoxyethylene(10)-N-tallow-1,3-diaminopropane, and
N,N',N'-polyoxyethylene(15)-N-tallow-1,3-diaminopropane.

The process of the present invention is useful for rest-breaking of
bushes, shrubs, vines, nuts, berries and non-deciduous fruit trees to
15 produce improvements advancing the time of bloom, budbreak and/or leaf
cover and fruit set. The present process also enables the use of
either milder rest-breaking agents or smaller quantities of harsher
rest-breaking agents thus providing considerable advantages in
toxicity to man and insect populations as well as safety advantages in
20 the application of these agents.

Examples of the types of plants to which the present invention is
applicable are grape vines, olive trees, raspberries, strawberries,
cranberries, blackberries, loganberries, blackcurrants, redcurrants
25 and any non-deciduous fruit trees. Nuts include, but are not limited
to, almonds, walnuts and chestnuts.

When reference is made in this patent application to enhancement of
rest-breaking, what is meant is that the time of bloom is advanced or
one or more budbreak and/or leaf cover or fruit set are improved over
30 a control tree.

5 The process of the present invention is to be applied to the respective plants, prior to blossom. The process is particularly useful in growing areas where the winter chilling of the plants is insufficient to provide good budbreak and fruit set for the particular cultivar. Winter chilling can be measured and is often represented by the unit, "Richardson Chill Units."

10 The optimum time to break rest for a particular plant will depend upon several factors including the type of fruit, the cultivar, the climatic conditions and the type and amount of rest-breaking agent being applied. For some fruit or cultivar species, the best rest-breaking effects are accomplished by early application of the rest-breaking agent to the plants whereas for others it is best to wait until just before blossom. In general, the rest-breaking agent will be applied at some point between the time when winter has peaked and the time when blossoming begins.

15 One of the effects of the rest-breaking agents of the present invention is to accelerate the blossoming of the plants. Accordingly, in the application of these rest-breaking agents consideration should be given to this effect and the agents should not be applied too early such that blooming occurs when there is still a risk of harsh weather.

25 The process of the present invention is carried out by the application of an effective amount of at least one rest-breaking agent in combination with an amount of at least one compound as defined herein to enhance the rest-breaking activity of said rest-breaking agent. The rest-breaking compositions are preferably applied in the form of an aqueous solution in a concentration of 0.25 to 30% and more preferably from 0.5 to 10%. The lower limit is generally determined by the upper limit on application volume for the particular application equipment being employed, as well as by the type of fruit, the cultivar and the particular rest-breaking composition.

30

The upper concentration limit will generally be dictated by phytotoxicity considerations since higher concentrations of certain compounds have a localized phytotoxic effect on the trees. Thus, a concentration should be selected which provides adequate rest-breaking without unwanted phytotoxic effects on the remainder of the plant. Such concentrations can be selected by routine experimentation with the particular species of plant.

The activity promoting additive of the present invention is generally employed in a concentration of 0.1-10.0% in the aqueous solution and more preferably, 0.2-5.0%. Again, the amount of activity promoting additive required will depend on the fruit, the cultivar and the particular rest-breaking agent, as well as the quantity of rest-breaking agent to be employed.

Rest-breaking agents which may be used in the process and composition of the invention include any known rest-breaking agent. Examples include hydrogen cyanamide, calcium ammonium nitrate, urea ammonium nitrate, potassium nitrate, the potassium salt of gibberellic acid, kinetin, indoleacetic acid, thiourea, and combinations thereof. The most preferred rest-breaking agents for use in the present invention are calcium ammonium nitrate and urea ammonium nitrate, especially when used on grapes. Also preferred is potassium nitrate. These nitrates are milder agents for which the activity can be enhanced to the desired level by the adjuvants of the present invention.

The composition is preferably applied in a manner similar to the manner in which commercial insecticides are applied. More particularly, conventional equipment such as knapsack sprayers, hand held spray guns, mist blowers, and aerial spraying equipment among others may be used. The composition is applied the same way as in pesticide application. The compositions may also be applied directly to the plant by hand, if desired.

The process of the present invention has the significant advantages that it breaks rest to the extent that the use of known, highly toxic rest-breaking agents can be eliminated or considerably reduced, it can be done in a manner which is safe for the crops and the treatment has no long term phytotoxic effect on the plants, if carried out correctly. Further, the rest-breaking process will cause significantly less harm to beneficial insects when applied within the normal application volume, and the process appears to be environmentally acceptable, non-hazardous to operators of the application equipment, and non-corrosive to the equipment.

The present invention will be further illustrated by the examples appended hereto.

Examples 1-12 and Comparative Examples A-C

Effect of Armobreak™ adjuvant used with Dormex™ Rest-Breaking Agent on the Perlette Table Grape

In Examples 1-12, various combinations of Dormex™ (hydrogen cyanamide rest breaking agent) and Armobreak™ (activity promoting additive) in aqueous spray solutions were employed as rest-breaking agents in Perlette grapes. These tests were carried out in the Coachella Valley of California with, as the purpose, to overcome the effects of inadequate winter chilling.


These examples were conducted in a commercial Perlette table grape vineyard located near Thermal, California. The own-rooted vines were planted 12 feet apart between rows and 6 feet apart between vines and were cordon trained and spur pruned. The vines were drip irrigated. The vines were pruned to 40, two-bud spurs on 15 December. The experimental treatments of the present examples were applied the following day.

Each treatment was replicated eight times using single vine plots. The treatments were arranged in a randomized complete block, split-plot design. A single vine served as a buffer between each plot.

5 The treatments of all the field trials in the examples have all been applied by a hand held spray wand at the rate of 100 gallons of spray solution per acre. The vines were monitored two to three times per week to determine the rate of bud emergence. The date of shoot emergence was recorded for the buds of ten randomly selected spurs on
10 each vine.

Armobreak™ which is a tallowamine containing five moles of ethylene oxide derived units and a total of 12 moles of propylene oxide units, can be represented by the general formula:

15



wherein $(p1+p2)=12$, $(q1+q2)=3$ and R is a hydrocarbon group derived from tallow fatty acid.

25

Comparative Examples A-C are the control examples where only Dormex™ was applied.

Results:

30

The following Table I gives the results obtained in these examples. All amounts are volume percent, based on the total volume of the aqueous solution.

The results for budbreak represent the number of days required to reach 70% of the total budbreak for a given grapevine.

TABLE 1

5	Example	Dormex™ (%v/v)	Armobreak™ (%v/v)	Days Until 70% Budbreak
	A	1.0	0.0	58
	B	2.0	0.0	47
	C	4.0	0.0	42
10	1	1.0	0.5	48
	2	2.0	0.5	37
	3	4.0	0.5	35
	4	1.0	1.0	42
	5	2.0	1.0	34
	6	4.0	1.0	36
	7	1.0	2.0	38
	8	2.0	2.0	34
15	9	4.0	2.0	36
	10	1.0	3.0	36
	11	2.0	3.0	33
	12	4.0	3.0	33

20 These results demonstrate that using 1.0% Dormex™ in combination with 1.0% Armobreak™ gives the same result (42 days) as is obtained using 4.0% Dormex™ without Armobreak™. Thus, the amount of rest-breaking agent can be reduced by up to 75% when employing the Armobreak™ adjuvant of the present invention without compromising the effects obtained with the rest-breaking agent.

25

Examples 13-18 and Comparative Examples D-L

Effect of Armobreak™ adjuvant used with Dormex™ Rest-Breaking Agent on different varieties of Table Grapes

30

This example illustrates that the ability of Armobreak to act effectively as an adjuvant for the rest breaking agent Dormex™ is

general across different varieties of vines. The results are shown in Table 2.

TABLE 2

5	Example	Variety	Dormex™ %(v/v)	Armobreak™ %(v/v)	% Budbreak 55 days after application
	D	Thompson	1	0	12
	E	Seedless	2	0	28
	F		4	0	43
	13		1	2	33
10	14		2	2	44
	G	Black	1	0	13
	H	Seedless	2	0	29
	I		4	0	36
	15		1	2	37
	16		2	2	31
	J	Red Globe	1	0	17
15	K		2	0	23
	L		4	0	34
	17		1	2	38
	18		2	2	39

20 The results show that the addition of 2% Armobreak to Dormex™ rest breaking agent allows equal or better advance of budbreak to be attained at a given time as when using two to four times the concentration of Dormex™ alone, and that this effect is common across different varieties of table grapevines.

25 Examples 19-28 and Comparative Examples M-N

Effect of Armobreak™ adjuvant used with Nitrate Rest-Breaking Agents on Perlette Table Grapes

30 The following examples illustrate the utility of Armobreak™ when used with a variety of nitrate rest breaking agents applied to Perlette variety grapevines. The results are shown in Table 3.

TABLE 3

Example	Rest breaking agent Name	Rest breaking agent %	Armobreak™ %(v/v)	Days to reach budbreak	
				70%	90%
5	M	-	-	50	53
	N	Dormex™	4	33	40
	19	NH ₄ NO ₃	25	52	55
	20	NH ₄ NO ₃	25	48	51
	21	KNO ₃	15	53	56
	22	KNO ₃	15	43	47
	23	CaNO ₃	15	52	57
	24	CaNO ₃	15	43	47
10	25	UAN32*	25	52	58
	26	UAN32*	25	37	46
	27	CAN17*	25	50	55
	28	CAN17*	25	33	38

*UAN32 is a commercially available solution of urea ammonium nitrate;
 15 CAN17 is a commercially available solution of calcium ammonium
 nitrate.

The use of nitrates alone, at the concentrations employed, causes
 little advance in budbreak relative to control (Example M). The
 20 addition of 2% Armobreak causes an improvement in the advance of
 budbreak with all nitrates tested. However, particular nitrates
 combined with 2% Armobreak can provide especially large improvements
 in advance of budbreak. Thus Calcium ammonium nitrate/Armobreak
 (Example 28) can be seen to provide equal advance of budbreak to the
 25 use of 4% Dormex alone (Example N). The combination of Armobreak with
 urea ammonium nitrate (Example 26) also effects a large advance of
 budbreak, and the combination of Armobreak with potassium or calcium
 nitrates also provides some improvement in advance of budbreak
 relative to the control. The CAN17/Armobreak combination is
 30 particularly preferred for use on grapevines.

Examples 29-32 and Comparative Example M (control)Effect of varying the concentration of Calcium Ammonium Nitrate as a Rest-Breaking Agent on Perlette Table Grapes

5 The following examples demonstrate the effect of varying the concentration of calcium ammonium nitrate used as a rest breaking agent in an experiment conducted on Perlette variety grapevines. The results are shown in Table 4.

10

TABLE 4

Example	CAN17 %(v/v)	Armobreak™ %(v/v)	Days to reach Budbreak	
			70%	90%
M	-	-	50	53
29	10	-	50	53
27	25	-	50	55
30	50	-	46	50
31	10	2	37	46
28	25	2	33	38
20 32	50	2	42	51

15

It can be seen that in the absence of Armobreak very high concentrations of CAN17 are required to produce effects upon advance of budbreak. In contrast, when 2% Armobreak is added lower concentrations of CAN17 produce considerably larger advances in budbreak. The optimum advance of budbreak produced by CAN17/Armobreak combination can be seen to depend upon the CAN17 concentration.

25

30

Examples 33-34 and Comparative Example 0 (control)Effect of Armobreak™ adjuvant used with Potassium Nitrate Rest-Breaking agent on Kiwi Fruit

5 The following table shows the effect of Armobreak™, used with potassium nitrate dormancy breaking agent, upon budbreak of kiwi fruit vines. The data are from an experiment in which the treatment solutions were applied to dormant vines about 6 weeks prior to
10 anticipated budbreak, at a solution spray volume of 1600 l/ha. In kiwi fruit it is desirable that any budbreaking treatment also reduce the number of side flowers so as to avoid excessive hand labour. The results are shown in Table 5.

TABLE 5

15

Example	Rest Breaking Name	Breaking %	Armobreak™	Budbreak Total%	75 days Floral%	Side flowers
0	-	-	-	36.2	26.4	13.3
33	KNO ₃	4	-	33.8	26.1	10.2
20 34	KNO ₃	4	2	38.5	31.6	7.4


It can be seen that Armobreak enhances the effect of potassium nitrate upon budbreak, and upon the reduction of undesirable side flowers.

25 The foregoing examples have been presented for purposes of illustration and description only and are not to be construed as limiting the scope of the invention in any manner. Accordingly, the scope of the invention is to be determined by the claims appended hereto.


30

What is claimed is:


1. A process for enhancing the rest-breaking of plants which comprises the step of applying to at least one plant selected from bushes, shrubs, vines, nuts, berries and non-deciduous fruit trees before blossom, an effective amount of at least one rest-breaking agent and a compound selected from alkoxyated amines represented by the following general formula:



wherein n is an integer from 1 to 50, A represents an alkylene group and when n>1, each A may be the same or different alkylene groups, R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms and groups represented by the formula:




wherein m is an integer from 1-50, A₁ represents an alkylene group and when m>1, each A₁ may be the same or different alkylene groups, x is an integer from 1-6, and R₂ is selected from straight or branched chain alkyl groups having 8 to 22 carbon atoms, R₃ is selected from hydrogen, 1-8 carbon atom straight or branched chain alkyl and alkenyl groups and aryl groups having up to 8 carbon atoms; and R₁ is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-22 carbon atoms, a group represented by the formula:



wherein n' is an integer from 1 to 50 and A₂ represents an alkylene group and when n' > 1 each A₂ may be the same or different alkylene groups; and alkoxyated quaternary ammonium compounds represented by the following general formula:


5



10

wherein n, A, R₁ and R₃ are as defined above, X is an anion, R₅ is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 4 carbon atoms, and benzyl, or R₅X⁻ is carboxymethyl or oxygen; R₄ is selected from straight or branched chain alkyl and alkenyl groups having 8-22 carbon atoms and groups represented by the formula:

15



20

wherein A₁, m, R₂, R₃, R₅ and x are as defined above.


25

2. The process of claim 1 wherein said rest-breaking agent is applied in the form of an aqueous solution having a concentration of from 0.1 to 10% of said compound and from 0.5-30% of said rest-breaking agent.

30

3. The process of claim 2 wherein said compound has a molecular weight of less than 8000 grams/mole.


4. The process of claim 3 wherein said compound is an alkoxyated quaternary ammonium compound represented by the general formula:



5

wherein n is an integer from 1 to 50, X⁻ is an anion, A represents an alkylene group and when n>1, each A may be the same or different alkylene groups, R₅ is selected from hydrogen, straight or branched chain alkyl and alkenyl groups having 1-4 carbon atoms, and benzyl, or R₅X⁻ is carboxymethyl or oxygen; R₄ is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms or a group represented by the formula:


15



20

wherein m is an integer from 1 to 50, R₅ is as defined above, A₁ represents an alkylene group and when n>1, each A₁ may be the same or different alkylene groups, x is an integer from 1 to 6, R₂ is selected from straight or branched chain alkyl or alkenyl groups having 8-22 carbon atoms; and R₁ is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 22 carbon atoms, a group represented by the formula:


25



30

wherein n' is an integer from 1 to 50 and A₂ represents an alkylene group; when n'>1 each A₂ may be the same or different alkylene groups, and R₃ is selected from hydrogen, straight or branched chain alkyl or alkenyl groups having 1 to 8 carbon atoms and aryl groups having up to 8 carbon atoms.


5. The process of claim 3 wherein said compound is an alkoxyated amine represented by the following general formula:




10 wherein n and n' are integers from 1 to 50, A represents an alkylene group and when n or n' are greater than 1, each A may be the same or different alkylene groups, and R is selected from straight or branched chain alkyl or alkenyl groups having 8 to 22 carbon atoms.

- 15 6. The process of claim 5 wherein n>1 and A includes both ethoxy groups and propoxy groups.

7. The process of claim 3 wherein said compound is an alkoxyated amine represented by the following general formula:



25 wherein n and n' are integers from 1 to 50, A represents an alkylene group and when n or n' are greater than 1, each A may be the same or different alkylene groups, R is a group represented by the formula:




wherein m is an integer from 1 to 50, A₁ represents an alkylene group and when m>1, each A₁ may be the same or different alkylene groups; x is an integer from 1 to 6, and R₂ is selected from straight or branched chain alkyl or alkenyl groups having 8-22 carbon atoms.

5

8. The process of claim 3 wherein R is selected from alkyl groups having 12-22 carbon atoms.

10

9. The process of claim 3 wherein said compound is an amine oxide represented by the following general formula:



15

wherein R, A, n and m are as defined in claim 1.

20

10. The process of claim 1 wherein said plant is selected from the group consisting of grape vines, olive trees, raspberries, strawberries, cranberries, blackberries, loganberries, blackcurrants, redcurrants, almonds, walnuts and chestnuts.

25

11. The process of any one of claims 1-10 wherein said rest-breaking compound is selected from the group consisting of hydrogen cyanamide, calcium ammonium nitrate, urea ammonium nitrate, potassium nitrate, the potassium salt of gibberellic acid, kinetin, indoleacetic acid, thiourea, and combinations thereof.

30

INTERNATIONAL SEARCH REPORT

International Application No
PC./EP 95/02575

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 A01N33/12 A01N33/08 A01N25/30 //(A01N33/08, 61:00, 59:24, 59:06, 59:00, 47:28, 25:30), (A01N33/12, 61:00, 59:24, 59:06, 59:00, 47:28, 25:30) According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 A01N Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Week 9418 Derwent Publications Ltd., London, GB; AN 94-151677 & ZA,A,9 303 611 (AKZO) , 23 February 1994 see abstract & WO,A,94 23574 (...) 27 October 1994 ---	1-11
P,X	EP,A,0 257 686 (AKZO) 2 March 1988 cited in the application see the whole document ---	1-11
Y	DE,A,31 50 404 (SKW TROSTBERG) 30 June 1983 see page 4, paragraph 5 see page 6 - page 7 see examples 1-4 see claim 3 ---	1-11
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 6 November 1995		Date of mailing of the international search report 23.11.95
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016		Authorized officer Lamers, W

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/02575

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 117, no. 1, 6 July 1992 Columbus, Ohio, US; abstract no. 2721, C.C.NEE 'Breaking dormancy in kiwifruit with chilling or chemical treatments' see abstract & N.Z.J.CROP HORTIC.SCI., vol. 19, no. 4, 1991 pages 419-421, ---	1-11
Y	GB,A,1 604 321 (R.C.BARBA) 9 December 1981 see page 1, line 26 - line 47 see page 1, line 69 - line 85 see page 2, line 16 - line 37 ---	1-11
Y	CHEMICAL ABSTRACTS, vol. 118, no. 15, 12 April 1993 Columbus, Ohio, US; abstract no. 141787, T.MOCHIZUKI ET AL. 'Studies on hastening of sprouting of grape buds.V. Nitrogen metabolism in the grape buds treated with ammonium nitrate and calcium cyanamide' see abstract & NIPPON DOJO HIRYOGAKU ZASSHI, vol. 64, no. 1, 1993 pages 62-70, ---	1-11
Y	GB,A,2 257 044 (IMPERIAL CHEMICAL INDUSTRIES) 6 January 1993 see page 1, paragraph 3 see page 1, paragraph 5 see page 1, line 24 see page 11; table IV see claims 1,2,21 ---	1-11
A	US,A,4 556 410 (P.M.RONNING ET AL.) 3 December 1985 see column 1, line 6 - column 4, line 48 ---	1-11
A	US,A,4 525 200 (M.KIMPARA ET AL.) 25 June 1985 see column 1, line 6 - column 3, line 3 ---	1-11
A	EP,A,0 272 542 (BASF) 29 June 1988 see page 3, line 1 - page 4, line 20 ---	1-11
A	EP,A,0 463 241 (AKZO) 2 January 1992 cited in the application see the whole document ---	1-11
A	EP,A,0 232 755 (MITSUBISHI GAS CHEMICAL) 19 August 1987 -----	1-11

INTERNATIONAL SEARCH REPORT

International Application No
PL./EP 95/02575

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0257686	02-03-88	CA-A- 1329538	17-05-94
		US-A- 5226943	13-07-93
		ZA-A- 8706184	22-02-88
DE-A-3150404	30-06-83	JP-B- 2062524	26-12-90
		JP-A- 58110511	01-07-83
		US-A- 4487625	11-12-84
GB-A-1604321	09-12-81	NONE	
GB-A-2257044	06-01-93	NONE	
US-A-4556410	03-12-85	NONE	
US-A-4525200	25-06-85	JP-C- 1783174	31-08-93
		JP-B- 4072801	19-11-92
		JP-A- 59157002	06-09-84
EP-A-0272542	29-06-88	DE-A- 3643246	30-06-88
		AU-B- 600059	02-08-90
		AU-B- 8264887	23-06-88
		DE-A- 3784658	15-04-93
		ES-T- 2044903	16-01-94
		US-A- 4936899	26-06-90
EP-A-0463241	02-01-92	AU-B- 635836	01-04-93
		AU-B- 7630991	02-01-92
		CA-A- 2045352	27-12-91
		US-A- 5118340	02-06-92
EP-A-0232755	19-08-87	JP-C- 1863243	08-08-94
		JP-A- 62167704	24-07-87
		US-A- 4799950	24-01-89