



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/04/12
(87) Date publication PCT/PCT Publication Date: 2019/11/21
(85) Entrée phase nationale/National Entry: 2020/10/19
(86) N° demande PCT/PCT Application No.: EP 2019/059522
(87) N° publication PCT/PCT Publication No.: 2019/219312
(30) Priorités/Priorities: 2018/05/16 (US62/672,411);
2018/05/16 (DE10 2018 111 819.8)

(51) Cl.Int./Int.Cl. *A61K 39/00* (2006.01),
C07K 14/47 (2006.01), *C07K 14/74* (2006.01)

(71) Demandeur/Applicant:
IMMATICS BIOTECHNOLOGIES GMBH, DE

(72) Inventeurs/Inventors:
SONG, COLETTE, DE;
HOFFGAARD, FRANZISKA, DE;
KOWALEWSKI, DANIEL, DE;
SCHOOR, OLIVER, DE;
FRITSCHKE, JENS, DE;
WEINSCHENK, TONI, DE;
SINGH, HARPREET, DE

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : PEPTIDES DESTINES A ETRE UTILISES EN IMMUNOTHERAPIE CONTRE DES CANCERS
(54) Title: PEPTIDES FOR USE IN IMMUNOTHERAPY AGAINST CANCERS

(57) **Abrégé/Abstract:**

The present invention relates to peptides, proteins, nucleic acids and cells for use in immunotherapeutic methods. In particular, the present invention relates to the immunotherapy of cancer. The present invention furthermore relates to tumor-associated T-cell peptide epitopes, alone or in combination with other tumor-associated peptides that can for example serve as active pharmaceutical ingredients of vaccine compositions that stimulate anti-tumor immune responses, or to stimulate T cells ex vivo and transfer into patients. Peptides bound to molecules of the major histocompatibility complex (MHC), or peptides as such, can also be targets of antibodies, soluble T-cell receptors, and other binding molecules.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

21 November 2019 (21.11.2019)



(10) International Publication Number

WO 2019/219312 A1

(51) International Patent Classification:

A61K 39/00 (2006.01) C07K 14/47 (2006.01)

C07K 14/74 (2006.01)

EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/EP2019/059522

(22) International Filing Date:

12 April 2019 (12.04.2019)

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10 2018 111 819.8

62/672,411 16 May 2018 (16.05.2018) DE

62/672,411 16 May 2018 (16.05.2018) US

(71) Applicant: IMMATICS BIOTECHNOLOGIES GMBH

[DE/DE]; Paul-Ehrlich-Straße 15, 72076 Tübingen (DE).

(72) Inventors: SONG, Colette; Willi-Baumeister-Weg 21,

73760 Ostfildern (DE). HOFFGAARD, Franziska;

Schwärzlocher Strasse 64, 72070 Tübingen (DE).

KOWALEWSKI, Daniel; Denzenbergerstrasse 29, 72074

Tübingen (DE). SCHOOR, Oliver; Eichhaldenstrasse 19,

72074 Tübingen (DE). FRITSCHE, Jens; Lärchenweg 11,

72144 Dusslingen (DE). WEINSCHENK, Toni; Im Mor-

genrain 15, 73773 Aichwald (DE). SINGH, Harpreet;

Heckscherstraße 25a, 80804 München (DE).

(74) Agent: KRAUß, Jan; BOEHMERT & BOEHMERT, Pet-

tenkoflerstraße 20-22, 80336 München (DE).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,

HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,

KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,

MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,

OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,

TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,

GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,

(54) Title: PEPTIDES FOR USE IN IMMUNOTHERAPY AGAINST CANCERS

(57) **Abstract:** The present invention relates to peptides, proteins, nucleic acids and cells for use in immunotherapeutic methods. In particular, the present invention relates to the immunotherapy of cancer. The present invention furthermore relates to tumor-associated T-cell peptide epitopes, alone or in combination with other tumor-associated peptides that can for example serve as active pharmaceutical ingredients of vaccine compositions that stimulate anti-tumor immune responses, or to stimulate T cells ex vivo and transfer into patients. Peptides bound to molecules of the major histocompatibility complex (MHC), or peptides as such, can also be targets of antibodies, soluble T-cell receptors, and other binding molecules.



WO 2019/219312 A1

Peptides for use in immunotherapy against cancers

The present invention relates to peptides, proteins, nucleic acids and cells for use in immunotherapeutic methods. In particular, the present invention relates to the immunotherapy of cancer. The present invention furthermore relates to tumor-associated T-cell peptide epitopes, alone or in combination with other tumor-associated peptides that can for example serve as active pharmaceutical ingredients of vaccine compositions that stimulate anti-tumor immune responses, or to stimulate T cells ex vivo and transfer into patients. Peptides bound to molecules of the major histocompatibility complex (MHC), or peptides as such, can also be targets of antibodies, soluble T-cell receptors, and other binding molecules.

The present invention relates to several novel peptide sequences and their variants derived from HLA class I molecules of human tumor cells that can be used in vaccine compositions for eliciting anti-tumor immune responses, or as targets for the development of pharmaceutically / immunologically active compounds and cells.

BACKGROUND OF THE INVENTION

According to the World Health Organization (WHO), cancer ranged among the four major non-communicable deadly diseases worldwide in 2012. For the same year, colorectal cancer, breast cancer and respiratory tract cancers were listed within the top 10 causes of death in high income countries.

Considering the severe side-effects and expenses associated with common strategies for treating cancer, there is a need to identify factors that can be used in the treatment of cancer in general and acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell

- 2 -

lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer in particular.

There is also a need to identify factors representing biomarkers for cancer in general and acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer in particular, leading to better diagnosis of cancer, assessment of prognosis, and prediction of treatment success.

Immunotherapy of cancer represents an option of specific targeting of cancer cells while minimizing side effects. Cancer immunotherapy makes use of the existence of tumor associated antigens.

The current classification of tumor associated antigens (TAAs) comprises the following major groups:

a) Cancer-testis antigens: The first TAAs ever identified that can be recognized by T cells belong to this class, which was originally called cancer-testis (CT) antigens because of the expression of its members in histologically different human tumors and, among normal tissues, only in spermatocytes/spermatogonia of testis and, occasionally, in placenta. Since the cells of testis do not express class I and II HLA molecules, these antigens cannot be recognized by T cells in normal tissues and can therefore be considered as immunologically tumor-specific. Well-known examples for CT antigens are the MAGE family members and NY-ESO-1.

b) Differentiation antigens: These TAAs are shared between tumors and the normal tissue from which the tumor arose. Most of the known differentiation antigens are found

- 3 -

in melanomas and normal melanocytes. Many of these melanocyte lineage-related proteins are involved in biosynthesis of melanin and are therefore not tumor specific but nevertheless are widely used for cancer immunotherapy. Examples include, but are not limited to, tyrosinase and Melan-A/MART-1 for melanoma or PSA for prostate cancer.

c) Over-expressed TAAs: Genes encoding widely expressed TAAs have been detected in histologically different types of tumors as well as in many normal tissues, generally with lower expression levels. It is possible that many of the epitopes processed and potentially presented by normal tissues are below the threshold level for T-cell recognition, while their over-expression in tumor cells can trigger an anticancer response by breaking previously established tolerance. Prominent examples for this class of TAAs are Her-2/neu, survivin, telomerase, or WT1.

d) Tumor-specific antigens: These unique TAAs arise from mutations of normal genes (such as β -catenin, CDK4, etc.). Some of these molecular changes are associated with neoplastic transformation and/or progression. Tumor-specific antigens are generally able to induce strong immune responses without bearing the risk for autoimmune reactions against normal tissues. On the other hand, these TAAs are in most cases only relevant to the exact tumor on which they were identified and are usually not shared between many individual tumors. Tumor-specificity (or -association) of a peptide may also arise if the peptide originates from a tumor- (-associated) exon in case of proteins with tumor-specific (-associated) isoforms.

e) TAAs arising from abnormal post-translational modifications: Such TAAs may arise from proteins which are neither specific nor overexpressed in tumors but nevertheless become tumor associated by posttranslational processes primarily active in tumors. Examples for this class arise from altered glycosylation patterns leading to novel epitopes in tumors as for MUC1 or events like protein splicing during degradation which may or may not be tumor specific.

f) Oncoviral proteins: These TAAs are viral proteins that may play a critical role in the oncogenic process and, because they are foreign (not of human origin), they can evoke a T-cell response. Examples of such proteins are the human papilloma type 16 virus proteins, E6 and E7, which are expressed in cervical carcinoma.

- 4 -

T-cell based immunotherapy targets peptide epitopes derived from tumor-associated or tumor-specific proteins, which are presented by molecules of the major histocompatibility complex (MHC). The antigens that are recognized by the tumor specific T lymphocytes, that is, the epitopes thereof, can be molecules derived from all protein classes, such as enzymes, receptors, transcription factors, etc. which are expressed and, as compared to unaltered cells of the same origin, usually up-regulated in cells of the respective tumor.

There are two classes of MHC-molecules, MHC class I and MHC class II. MHC class I molecules are composed of an alpha heavy chain and beta-2-microglobulin, MHC class II molecules of an alpha and a beta chain. Their three-dimensional conformation results in a binding groove, which is used for non-covalent interaction with peptides.

MHC class I molecules can be found on most nucleated cells. They present peptides that result from proteolytic cleavage of predominantly endogenous proteins, defective ribosomal products (DRIPs) and larger peptides. However, peptides derived from endosomal compartments or exogenous sources are also frequently found on MHC class I molecules. This non-classical way of class I presentation is referred to as cross-presentation in the literature (Brossart and Bevan, 1997; Rock et al., 1990). MHC class II molecules can be found predominantly on professional antigen presenting cells (APCs), and primarily present peptides of exogenous or transmembrane proteins that are taken up by APCs e.g. during endocytosis and are subsequently processed.

Complexes of peptide and MHC class I are recognized by CD8-positive T cells bearing the appropriate T-cell receptor (TCR), whereas complexes of peptide and MHC class II molecules are recognized by CD4-positive-helper-T cells bearing the appropriate TCR. It is well known that the TCR, the peptide and the MHC are thereby present in a stoichiometric amount of 1:1:1.

CD4-positive helper T cells play an important role in inducing and sustaining effective responses by CD8-positive cytotoxic T cells. The identification of CD4-positive T-cell

- 5 -

epitopes derived from tumor associated antigens (TAA) is of great importance for the development of pharmaceutical products for triggering anti-tumor immune responses (Gnjatic et al., 2003). At the tumor site, T helper cells, support a cytotoxic T cell- (CTL-) friendly cytokine milieu (Mortara et al., 2006) and attract effector cells, e.g. CTLs, natural killer (NK) cells, macrophages, and granulocytes (Hwang et al., 2007).

In the absence of inflammation, expression of MHC class II molecules is mainly restricted to cells of the immune system, especially professional antigen-presenting cells (APC), e.g., monocytes, monocyte-derived cells, macrophages, dendritic cells. In cancer patients, cells of the tumor have been found to express MHC class II molecules (Dengjel et al., 2006).

Longer (elongated) peptides of the invention can act as MHC class II active epitopes.

T-helper cells, activated by MHC class II epitopes, play an important role in orchestrating the effector function of CTLs in anti-tumor immunity. T-helper cell epitopes that trigger a T-helper cell response of the TH1 type support effector functions of CD8-positive killer T cells, which include cytotoxic functions directed against tumor cells displaying tumor-associated peptide/MHC complexes on their cell surfaces. In this way tumor-associated T-helper cell peptide epitopes, alone or in combination with other tumor-associated peptides, can serve as active pharmaceutical ingredients of vaccine compositions that stimulate anti-tumor immune responses.

It was shown in mammalian animal models, e.g., mice, that even in the absence of CD8-positive T lymphocytes, CD4-positive T cells are sufficient for inhibiting manifestation of tumors via inhibition of angiogenesis by secretion of interferon-gamma (IFN γ) (Beatty and Paterson, 2001; Mumberg et al., 1999). There is evidence for CD4 T cells as direct anti-tumor effectors (Braumuller et al., 2013; Tran et al., 2014).

Since the constitutive expression of HLA class II molecules is usually limited to immune cells, the possibility of isolating class II peptides directly from primary tumors was

- 6 -

previously not considered possible. However, Dengjel et al. were successful in identifying a number of MHC Class II epitopes directly from tumors (WO 2007/028574, EP 1 760 088 B1).

Since both types of response, CD8 and CD4 dependent, contribute jointly and synergistically to the anti-tumor effect, the identification and characterization of tumor-associated antigens recognized by either CD8+ T cells (ligand: MHC class I molecule + peptide epitope) or by CD4-positive T-helper cells (ligand: MHC class II molecule + peptide epitope) is important in the development of tumor vaccines.

For an MHC class I peptide to trigger (elicit) a cellular immune response, it also must bind to an MHC-molecule. This process is dependent on the allele of the MHC-molecule and specific polymorphisms of the amino acid sequence of the peptide. MHC-class-I-binding peptides are usually 8-12 amino acid residues in length and usually contain two conserved residues ("anchors") in their sequence that interact with the corresponding binding groove of the MHC-molecule. In this way each MHC allele has a "binding motif" determining which peptides can bind specifically to the binding groove.

In the MHC class I dependent immune reaction, peptides not only have to be able to bind to certain MHC class I molecules expressed by tumor cells, they subsequently also have to be recognized by T cells bearing specific T cell receptors (TCR).

For proteins to be recognized by T-lymphocytes as tumor-specific or -associated antigens, and to be used in a therapy, particular prerequisites must be fulfilled. The antigen should be expressed mainly by tumor cells and not, or in comparably small amounts, by normal healthy tissues. In a preferred embodiment, the peptide should be over-presented by tumor cells as compared to normal healthy tissues. It is furthermore desirable that the respective antigen is not only present in a type of tumor, but also in high concentrations (i.e. copy numbers of the respective peptide per cell). Tumor-specific and tumor-associated antigens are often derived from proteins directly involved in transformation of a normal cell to a tumor cell due to their function, e.g. in cell cycle

- 7 -

control or suppression of apoptosis. Additionally, downstream targets of the proteins directly causative for a transformation may be up-regulated and thus may be indirectly tumor-associated. Such indirect tumor-associated antigens may also be targets of a vaccination approach (Singh-Jasuja et al., 2004). It is essential that epitopes are present in the amino acid sequence of the antigen, in order to ensure that such a peptide ("immunogenic peptide"), being derived from a tumor associated antigen, leads to an *in vitro* or *in vivo* T-cell-response.

Basically, any peptide able to bind an MHC molecule may function as a T-cell epitope. A prerequisite for the induction of an *in vitro* or *in vivo* T-cell-response is the presence of a T cell having a corresponding TCR and the absence of immunological tolerance for this particular epitope.

Therefore, TAAs are a starting point for the development of a T cell based therapy including but not limited to tumor vaccines. The methods for identifying and characterizing the TAAs are usually based on the use of T-cells that can be isolated from patients or healthy subjects, or they are based on the generation of differential transcription profiles or differential peptide expression patterns between tumors and normal tissues. However, the identification of genes over-expressed in tumor tissues or human tumor cell lines, or selectively expressed in such tissues or cell lines, does not provide precise information as to the use of the antigens being transcribed from these genes in an immune therapy. This is because only an individual subpopulation of epitopes of these antigens are suitable for such an application since a T cell with a corresponding TCR has to be present and the immunological tolerance for this particular epitope needs to be absent or minimal. In a very preferred embodiment of the invention it is therefore important to select only those over- or selectively presented peptides against which a functional and/or a proliferating T cell can be found. Such a functional T cell is defined as a T cell, which upon stimulation with a specific antigen can be clonally expanded and is able to execute effector functions ("effector T cell").

- 8 -

In case of targeting peptide-MHC by specific TCRs (e.g. soluble TCRs) and antibodies or other binding molecules (scaffolds) according to the invention, the immunogenicity of the underlying peptides is secondary. In these cases, the presentation is the determining factor.

In a first aspect of the present invention, the present invention relates to a peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464 or a variant sequence thereof which is at least 77%, preferably at least 88%, homologous (preferably at least 77% or at least 88% identical) to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464, wherein said variant binds to MHC and/or induces T cells cross-reacting with said peptide, or a pharmaceutical acceptable salt thereof, wherein said peptide is not the underlying full-length polypeptide.

The present invention also relates to a peptide of the present invention comprising a sequence that is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464 or a variant thereof, which is at least 77%, preferably at least 88%, homologous (preferably at least 77% or at least 88% identical) to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464, wherein said peptide or variant thereof has an overall length of between 8 and 100, preferably between 8 and 30, and most preferred of between 8 and 14 amino acids.

The following tables show the peptides according to the present invention, their respective SEQ ID NOs, and the prospective source (underlying) genes for these peptides. In Table 1a, peptides with SEQ ID NO: 1 to SEQ ID NO: 382 and in Table 1b, peptides with SEQ ID NO: 463 to SEQ ID NO: 464 bind to HLA-A*24. The peptides in Table 2 have been disclosed before in large listings as results of high-throughput screenings with high error rates or calculated using algorithms but have not been associated with cancer at all before. In Table 2, peptides with SEQ ID NO: 383 to SEQ ID NO: 387 bind to HLA-A*24. The peptides in Table 3 are additional peptides that may

be useful in combination with the other peptides of the invention. In Table 3, peptides with SEQ ID NO: 388 to SEQ ID NO: 460 bind to HLA-A*24.

Table 1a: Peptides according to the present invention.

Seq ID No	Sequence	Gene(s)	HLA allotype
1	IFPKTGLLII	MAGEA4	A*24
2	LYAPTILLW	AFP	A*24
3	KFLTHDVLTELF	TRPM8	A*24
4	MVLQPQPQLF	POTEG, POTEH	A*24
5	LQPQPQLFFSF	POTEG, POTEH	A*24
6	IVTFMNKTLGTF	ADAM29	A*24
7	GYPLRGSSI	ALPP, ALPPL2	A*24
8	IMKPLDQDF	ADAM2	A*24
9	TYINSLAIL	TGM4	A*24
10	QYPEFSIEL	TGM4	A*24
11	RAMCAMMSF	TGM4	A*24
12	KYMSRVLFVY	CHRNA9	A*24
13	KYYIATMAL	CHRNA9	A*24
14	YYIATMALI	CHRNA9	A*24
15	FMVIAGMPLF	SLC6A3	A*24
16	GYFLAQYLM	TRPM8	A*24
17	IYPEAIATL	SLC6A3	A*24
18	KYVDINTFRL	MMP12	A*24
19	ILLCMSLLLF	CYP4Z1, CYP4Z2P	A*24
20	ELMAHPFLL	CYP4Z1, CYP4Z2P	A*24
21	LYMRFVNTHF	SPINK2	A*24
22	VYSSFVFNL	NLRP4	A*24
23	VYSSFVFNLF	NLRP4	A*24
24	KMLPEASLLI	NLRP4	A*24
25	MLPEASLLI	NLRP4	A*24
26	TYFFVDNQYW	MMP12	A*24
27	LSCTATPLF	KHDC1L	A*24
28	FWFDSREISF	OR51E2	A*24
29	IYLLPPIVI	OR51E2	A*24
30	RQAYSVYAF	SLC45A3	A*24
31	KMQQEFFGL	MMP1	A*24
32	FYPEVELNF	MMP1	A*24
33	FYQPDLKYLFS	NLRP4	A*24
34	LIFALALAAF	GAST	A*24
35	FSSTLVSLF	MAGEC1	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
36	VYLASVAAF	SLC45A3	A*24
37	ISFSDTVNVW	ITIH6	A*24
38	RYAHTLVTSVLF	ITIH6	A*24
39	KTYLPTFETTI	ENPP3, OR2A4	A*24
40	NYPEGAAAYEF	ESR1	A*24
41	IYFATQVVF	SLC45A3	A*24
42	VYDSIWCNM	SCGB2A1	A*24
43	KYKDHFEI	MAGEB1	A*24
44	FYHEDMPLW	FCRL5	A*24
45	YGQSKPWTF	PAX3	A*24
46	IYPSIQEL	LOC645382, LOC645399, PRAMEF10	A*24
47	SYLWTDNLQEF	DNAH17	A*24
48	AWSPPATLFLF	LOXL4	A*24
49	QYLSIAERAEF	MSX1, MSX2	A*24
50	RYFDENIQKF	HEPHL1	A*24
51	YFDENIQKF	HEPHL1	A*24
52	SWHKATFLF	COL24A1	A*24
53	LFQRVSSVSF	HMCN1	A*24
54	SYQEAIQQL	NEFH	A*24
55	AVLRHLETF	CDK6	A*24
56	FYKLIQNGF	FLT3	A*24
57	RYLQVLLY	NPSR1	A*24
58	IYYSHENLI	F5	A*24
59	VFPLVTPLL	PTPRZ1	A*24
60	RYSPVKDAW	KLHDC7B	A*24
61	RIFTARLYF	AICDA	A*24
62	VYIVPVIVL	OXTR	A*24
63	LYIDKGQYL	HMCN1	A*24
64	QFSHVPLNNF	ALX1	A*24
65	EYLLMIFKLV	HRNR, RPTN	A*24
66	IYKDYYRYNF	PLA2G2D	A*24
67	SYVLQIVAI	PTPRZ1	A*24
68	VYKEDLPQL	EML5, EML6	A*24
69	KWFDSHIPRW	ERVV-1, ERVV-2	A*24
70	RYTGQWSEW	IL9R	A*24
71	RYLPNPSLNAF	CYP1A1	A*24
72	RWLDGSPVTL	CLEC17A	A*24
73	YFCSTKGQLF	FCRL2	A*24
74	NYVLVPTMF	CAPN6	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
75	VYEHNHVSL	BTBD16	A*24
76	IYIYPFAHW	NPFFR2	A*24
77	LYGFFFKI	BTBD16	A*24
78	TYSKTIALYGF	BTBD16	A*24
79	FYIVTRPLAF	SUCNR1	A*24
80	SYATPVDLW	CDK6	A*24
81	AYLKLLPMF	SLC5A4	A*24
82	SYLENSASW	DLX5	A*24
83	VLQGEFFLF	KBTBD8	A*24
84	YTIERYFTL	GABRP	A*24
85	KYLSIPTVFF	UGT1A3	A*24
86	SYLPTAERL	SYT12	A*24
87	NYTRLVLQF	GABRP	A*24
88	TYVPSTFLV	GABRP	A*24
89	TYVPSTFLVVL	GABRP	A*24
90	TDLVQFLLF	MAGEA10	A*24
91	KQQVVKFLI	LOC100288966, POTE _B , POTE _C , POTE _D , POTE _E , POTE _F , POTE _G , POTE _H , POTE _I , POTE _J , POTE _{KP} , POTE _M	A*24
92	RALTETIMF	ALPP, ALPPL2	A*24; B*15
93	TDWSPPPVEF	FAM178B	A*24
94	THSGGTNLF	MMP12	A*24; B*38
95	IGLSVVHRF	OR51E2	A*24
96	SHIGVVLAFL	OR51E2	A*24; B*15
97	TQMFFIHAL	OR51E2	A*24; B*39
98	LQIPVSPSF	MAGEC1	A*24; B*15
99	ASAALTGFTF	SLC45A3	A*24; A*32
100	KVWSDVTPLTF	MMP11	A*24; A*32
101	VYAVSSDRF	DCX	A*24
102	VLASAHILQF	BTBD16	A*24
103	EMFFSPQVF	ACTL8	A*24
104	GYGLTRVQPF	ACTL8	A*24
105	ITPATALLL	LOC100996407, LOC101060778, LOC441242	A*24
106	LYAFLGSHF	KISS1R	A*24

- 12 -

Seq ID No	Sequence	Gene(s)	HLA allotype
107	FFFKAGFVWR	MMP11	A*24
108	WFFQGAQYW	MMP11	A*24
109	AQHSLTQLF	GPC2	A*24; B*15
110	VYSNPDLFW	TRDV3	A*24
111	IRPDYSFQF	TRDV3	A*24
112	LYPDSVFGRLF	SMC1B	A*24
113	ALMSAFYTF	MMP11	A*24
114	KALMSAFYTF	MMP11	A*24
115	IMQGFIRAF	PAEP	A*24
116	TYFFVANKY	MMP1	A*24
117	RSMEHPGKLLF	ESR1	A*24; B*15
118	IFLPPFIVF	ADAM18	A*24
119	VWSCEGCKAF	ESR1, ESR2	A*24
120	VYAFMNENF	QRFPR	A*24
121	RRYFGEKVAL	ANO7	A*24; B*27
122	YFLRGRLYW	MMP11	A*24
123	FFLQESPVF	ABCC11	A*24
124	EYNVFPRTL	MMP13	A*24
125	LYYGSILYI	LOC100996718, OR9G1, OR9G9	A*24
126	YSLLDPAQF	SOX14	A*24
127	FLPRAYYRW	ANO7	A*24
128	AFQNVISF	NMUR2	A*24
129	IYVSLAHVL	ANO7	A*24
130	RPEKVFVF	COL11A1	A*24
131	MHRTWRETF	ANO7	A*24; A*32
132	TFEGATVTL	FCRL5	A*24
133	FFYVTETTF	TERT	A*24
134	IYSSQLPSF	TFEC	A*24
135	KYKQHFPEI	MAGEB17	A*24
136	YLKSVQLF	RFX8	A*24
137	ALFAVCWAPF	QRFPR	A*24
138	MMVTVVALF	QRFPR	A*24
139	AYAPRGSIIYKF	HHIPL2	A*24
140	IFQHFCEEI	SMC1B	A*24
141	QYAAAITNGL	SALL3	A*24
142	PYWWNANMVF	NOTUM	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
143	KTKRWLWDF	COL11A1	A*24
144	LFDHGGTVFF	ANO7	A*24
145	MYTIVTPML	OR1N1	A*24
146	NYFLDPVTI	TRIM51, TRIM51HP	A*24
147	FPYPSSILSV	DLL3	A*24; B*51
148	MLPQIPFLLL	COL10A1	A*24; A*02
149	TQFFIPYTI	COL10A1	A*24; A*32
150	FIPVAWLIF	MRGPRX4	A*24
151	RRLWAYVTI	ITIH6	A*24
152	MHPGVLA AFLF	MMP13	A*24; B*15
153	AWSPATLF	LOXL4	A*24
154	DYSKQALSL	LAMC2	A*24
155	PYSIYPHGVT F	F5	A*24
156	IYPHGVT FSP	F5	A*24
157	SIYPHGVT F	F5	A*24; B*15
158	SYLKDPMIV	DDX53	A*24
159	VFQPNPLF	WISP3	A*24
160	YIANLISCF	GLYATL3	A*24
161	ILQAPLSVF	FCRL5	A*24; B*15
162	YYIGIVEEY	HEPHL1	A*24
163	YYIGIVEEYW	HEPHL1	A*24
164	MFQEMLQRL	TRIML2	A*24
165	KDQPQVPCVF	NAT1	A*24; B*15
166	MMALWSLLHL	ZACN	A*24
167	LQPPWTTVF	FCRL5	A*24; B*15
168	LSSPVHLDF	FCRL5	A*24; B*58
169	MYDLHHLYL	EPYC	A*24
170	IFIPATILL	ACSM1	A*24
171	LYTVPFNLI	SLC45A2	A*24
172	RYFIAAEKILW	HEPHL1	A*24
173	RYLSVCERL	NKX1-1, NKX1-2	A*24
174	TYGEEMPEEI	DNAH17	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
175	SYFEYRRLL	LAMC2	A*24
176	TQAGEYLLF	FLT3	A*24
177	KYLITTFSL	NLRP2	A*24
178	AYPQIRCTW	FLT3	A*24
179	MYNMVPPF	DCT	A*24
180	IYNKTKMAF	SLCO6A1	A*24
181	IHGKIFHYF	NMUR2	A*24
182	AQGSGTVTF	FCRL3	A*24; B*15
183	YQVAKGMEF	FLT3	A*24
184	VYVRPRVF	HMCN1	A*24
185	LYICKVELM	CTLA4	A*24
186	RRVTWNVLF	BTBD17	A*24
187	KWFNVRMGFGF	LIN28A, LIN28B	A*24
188	SLPGSFIYVF	HMCN1	A*24
189	FYPDEDDFYF	MYCN	A*24
190	IYIIMQSCW	FLT3	A*24
191	MSYSCGLPSL	KRT33A	A*24
192	CYSFIHLSF	NLRP2, NLRP7	A*24
193	KYKPVALQCIA	HMCN1	A*24
194	EYFLPSLEII	SLC24A5	A*24
195	IYNEHGIQQI	COL11A1	A*24
196	VGRSPVFLF	COL11A1	A*24
197	YYHSGENLY	PSG1, PSG3, PSG7	A*24
198	VLAPVSGQF	FCRL5	A*24; B*15
199	MFQFEHIKW	FBXW10	A*24
200	LYMSVEDFI	STK31	A*24
201	VFPSVDVSF	PTPRZ1	A*24
202	VYDTMIEKFA	PTPRZ1	A*24
203	VYPSESTVM	PTPRZ1	A*24
204	WQNVTPLTF	MMP16	A*24
205	ISWEVVHTVF	HMCN1	A*24
206	EVVHTVFLF	HMCN1	A*24; A*26
207	IYKFIMDRF	FOXB1	A*24
208	QYLQQQAKL	FOXB1	A*24
209	DIYVTGGHLF	KLHDC7B	A*24
210	EAYSYPPATI	HMCN1	A*24
211	MLYFAPDLIL	PGR	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
212	VYFVQYKIM	IL22RA2	A*24
213	FYNRLTKLF	OFCC1	A*24
214	YIPMSVMLF	HTR7	A*24
215	KASKITFHW	PTPRZ1	A*24
216	RHYHSIEVF	LOXL4	A*24; C*12
217	QRYGFSSVGF	RHBG	A*24; B*27
218	FYFYNCSSL	ERVV-1, ERVV-2	A*24
219	KVVSIFYI	CCR8	A*24
220	TYATHVTEI	CCR8	A*24
221	VFYCLLFVF	CCR8	A*24
222	HYHAESFLF	HEPHL1	A*24
223	KLRALSILF	PTPRZ1	A*24
224	AYLQFLSVL	GREB1	A*24
225	ISMSATEFLL	CYP1A1	A*24
226	TYSTNRTMI	FLT3	A*24
227	YLPNPSLNAF	CYP1A1	A*24
228	VYLRIGGF	WNT7A	A*24
229	CAMPVAMEF	KBTBD8	A*24
230	RWLSKPSLL	KBTBD8	A*24
231	KYSVAFYSLD	LAMA1	A*24
232	IWPGFTTSI	PIWIL1	A*24
233	LYSRRGVRTL	DPPA3, DPPA3P2, LOC101060236	A*24
234	RYKMLIPF	NLRP2	A*24
235	VYISDVSVY	CLECL1	A*24
236	LHLYCLNTF	PGR	A*24
237	RQGLTVLTW	DNAH8	A*24
238	YTCSRASLFL	OTOG	A*24
239	IYTFSNVTF	BTN1A1	A*24
240	RVHANPLLI	APOB	A*24; B*15
241	QKYYITGEAEGF	ESR1	A*24
242	SYTPLLSYI	C1orf94	A*24
243	ALFPMGPLTF	LILRA4	A*24; B*15
244	TYIDTRTVFL	CAPN6	A*24
245	VLPLHFLPF	HBG2	A*24
246	KIYTTVLFANI	NPFFR2	A*24
247	VHSYLGSPF	MPL	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
248	CWGPHCFEM	SEMA5B	A*24
249	HQYGGAYNRV	DLX5	A*24
250	VYSDRQIYLL	ABCC11	A*24
251	DYLLSWLLF	CNR2	A*24
252	RYLIKYPF	SUCNR1	A*24
253	QYYCLLLIF	KLRF2	A*24
254	KQHAWLPLTI	TCL1A	A*24
255	VYLDEKQHAW	TCL1A	A*24
256	QHAWLPLTI	TCL1A	A*24; B*39
257	MLILFFSTI	OR56A3	A*24
258	VCWNPFNNTF	RNF183	A*24
259	FFLFIPFF	ADAM2	A*24
260	FLFIPFFIIF	ADAM2	A*24
261	IMFCLKNFVW	TBC1D7	A*24
262	YIMFCLKNF	TBC1D7	A*24
263	AYVTEFVSL	SCN3A	A*24
264	AYAIPSASLSW	HMCN1	A*24
265	LYQQSDTWSL	KIAA1407	A*24
266	TQIITFESF	CSF2	A*24; B*15
267	QHMLPFWTDL	NLRP2	A*24
268	YQFGWSPNF	CFHR5	A*24
269	FSFSTSMNEF	CAPN6	A*24
270	GTGKLFWVF	BTLA	A*24
271	INGDLVFSF	CAPN6	A*24
272	IYFNHRCF	SFMBT1	A*24
273	VTMYLPLLL	GPR143	A*24
274	EYSLPVLTF	PTPRZ1	A*24
275	PEYSLPVLTF	PTPRZ1	A*24
276	KFLGSKCSF	HAS3	A*24
277	MSAIWISAF	SLC24A5	A*24
278	TYESVVTGFF	HAS3	A*24
279	KYKNPYGF	MMP20	A*24
280	TIYSLEMKMSF	GLB1L3	A*24
281	MDQNQVWTF	ROS1	A*24
282	ASYQQSTSSFF	FAM82A1	A*24
283	SYIVDGKII	PSG9	A*24
284	QFYSTLPNTI	ROS1	A*24
285	YFLPGPHYF	SOX30	A*24

- 17 -

Seq ID No	Sequence	Gene(s)	HLA allotype
286	HHTQLIFVF	ELP4, EXOSC7, KCNG2, TM4SF19, TOP2A	A*24
287	LVQPQAVLF	PAX5	A*24
288	MGKGSISFLF	PCSK1	A*24
289	RTLNEIYHW	FOXP3	A*24
290	VTPKMLISF	OR5H8P	A*24
291	YTRLVLQF	GABRP	A*24
292	KMFPKDFRF	TTLL6	A*24
293	MYAYAGWFY	SLC7A11	A*24
294	KMGRIVDYF	GABRP	A*24
295	KYNRQSMTL	APOB	A*24
296	YQRPDLLLF	GEN1	A*24; B*15
297	LKSPRLFTF	BTBD16	A*24
298	TYETVMTFF	BTBD16	A*24
299	FLPALYSLL	CXCR3	A*24
300	LFALPDFIF	CXCR3	A*24
301	RTALSSTDTF	CXCR3	A*24
302	YQGSLEVLV	MROH2A	A*24
303	RFLDRGWGF	ADAMTS12	A*24
304	YFGNPQKF	LAMA3	A*24
305	RNAFSIYIL	MRGPRX1	A*24
306	RYILEPFFI	SLC7A11	A*24
307	RILTEFELL	TRIM31	A*24
308	AAFISVPLLI	TAS2R38	A*24
309	AFISVPLLI	TAS2R38	A*24
310	EFINGWYVL	MCOLN2	A*24
311	IQNAILHLF	OR51B5	A*24
312	YLCMLYALF	KCNK18	A*24
313	IFMENAFEL	APOB	A*24
314	SQHFNLATF	DNMT3B	A*24; B*15
315	VYDYIPLLL	MROH2A	A*24
316	IWAERIMF	TDRD1	A*24
317	DWIWRILFLV	IGHV1-58	A*24
318	VQADAKLLF	FERMT1	A*24; B*15
319	ATATLHLIF	PCDHGB1, PCDHGB2	A*24; B*15
320	EYQKILKF	PASD1	A*24
321	VYTVGHNLI	KLB	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
322	SFISPRYSWLF	SPNS3	A*24
323	NYSPVTGKF	OTOL1	A*24
324	RYFVSNIYL	PRSS21	A*24
325	IFMGAVPTL	LPAR3	A*24
326	VHMKDFFYF	DYRK4	A*24
327	KWKPSPLLF	GPR126	A*24
328	IYLVGGYSW	KLHL14	A*24
329	YLGKNWSF	SPNS3	A*24
330	DYIQMIPEL	RTL1	A*24
331	EYIDEFQSL	RTL1	A*24
332	VYCSLDKSQF	RTL1	A*24
333	RYADLLIYTY	MYO3B	A*24
334	KVFGSFLTL	AGTR2	A*24
335	RYQSVIYPF	AGTR2	A*24
336	VYSDLHAFY	MANEAL	A*24; A*29
337	SHSDHEFLF	ARSH	A*24; B*38
338	VYLTWLPGL	IFNLR1	A*24
339	KQVIGIHTF	SFMBT1	A*24; B*15
340	FPPTPPLF	BCL11A	A*24
341	RYENVSILF	ADCY8	A*24
342	MYGIVIRTI	NPSR1	A*24
343	EYQQYHPSL	CLEC4C	A*24
344	YAYATVLTF	ABCC4	A*24; B*46
345	RYLEEHTEF	MROH2A	A*24
346	TYIDFVPYI	TEX15	A*24
347	AWLIVLLFL	CTCFL	A*24
348	RSWENIPVTF	C18orf54	A*24
349	IYMTTGVLL	TDRD9	A*24
350	VYKWTEEFK	TSPEAR	A*24
351	GYFGTASLF	SLC16A14	A*24
352	NAFEAPLTF	BRCA2	A*24; C*07
353	AAFPGAFSF	CRB2	A*24; C*07
354	QYIPTFHVY	SLC44A5	A*24
355	VYNNNSSRF	MYO10	A*24
356	YSLEHLTQF	ZCCHC16	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
357	RALLPSPLF	SPATA31D1	A*24
358	IYANVTEMLL	CYP27C1	A*24
359	TQLPAPLRI	GPR45	A*24
360	LYITKVTTI	FSTL4	A*24
361	KQPANFIVL	LOC100124692	A*24
362	NYMDTDNLMF	LOC100124692	A*24
363	QYGFNYNKF	PNLDC1	A*24
364	KQSQVVFVL	HMCN1, HMCN2, LOC101060175	A*24; B*48
365	KDLMKAYLF	TXNDC16	A*24; B*37
366	RLGEFVLLF	TGM6	A*24
367	HWSHITHLF	DPY19L1	A*24
368	AYFVAMHLF	TENM4	A*24
369	NFYLFPTTF	PNLDC1	A*24
370	TQMDVKLVF	GEN1	A*24; B*15
371	FRSWAVQTF	NOS2	A*24; C*06
372	LYHNWRHAF	PDE11A	A*24
373	IWDALERTF	ABCC11	A*24
374	MIFAVVLF	CCR4	A*24
375	YYAADQWVF	CCR4	A*24
376	KYVGEVFNI	DMXL1	A*24
377	SLWREVVTF	CEP250	A*24
378	VYAVISNIL	TNR	A*24
379	KLPTEWNVL	AKAP13	A*24
380	FYIRRLPMF	CHRNA6	A*24
381	IYTDITYSF	CHRNA6	A*24
382	SYPKELMKF	MROH2A	A*24

Table 1b: Peptides according to the present invention.

Seq ID No	Sequence	Gene(s)	HLA allotype
463	VGGNVTSNF	CT45A4, CT45A5	A*24
464	VGGNVTSSF	CT45A1, CT45A2, CT45A3, CT45A4, CT45A6, LOC101060208, LOC101060210, LOC101060211	A*24

- 20 -

Table 2: Additional peptides according to the present invention with no prior known cancer association.

Seq ID No	Sequence	Gene(s)	HLA allotype
383	PYFSPSASF	SPERT	A*24
384	RTRGWVQTL	C6orf223	A*24
385	GYFGNPQKF	LAMA3	A*24
386	YQSRDYYNF	AR	A*24
387	THAGVRLYF	NUP155	A*24; B*38

Table 3: Peptides according to the present invention useful for e.g. personalized cancer therapies.

Seq ID No	Sequence	Gene(s)	HLA allotype
388	LFHPEDTGQVF	KLK3	A*24
389	AYSEKVTEF	KLK2	A*24
390	KYKDYFPVI	LOC392555 ,MAGEC2	A*24
391	VYGEPRELL	LOC392555, MAGEC2	A*24
392	SYEKVINYL	MAGEA9, MAGEA9B	A*24
393	SYNDALLTF	TRPM8	A*24
394	VYLPKIPSW	KCNU1	A*24
395	NYEDHFPLL	MAGEA10	A*24
396	SYVKVLHHL	LOC101060230, MAGEA12	A*24
397	RMPTVLQCV	KLK4	A*24
398	GYLQGLVSF	KLK4	A*24
399	VWSNVTPLKF	MMP12	A*24
400	RYLEKFYGL	MMP12	A*24
401	YLEKFYGL	MMP12	A*24
402	TYKYVDINTF	MMP12	A*24
403	LYFEKGEYF	ACPP	A*24
404	SYLKAVFNL	CYP4Z1, CYP4Z2P	A*24
405	NYPKSIHSF	MMP12	A*24
406	KYLEKYYNL	MMP1	A*24
407	RILRFPWQL	MMP11	A*24
408	VWSDVTPLTF	MMP11	A*24
409	VYTFLSSTL	ESR1	A*24
410	IYISNSIYF	CXorf48	A*24
411	VYPPYLNLYL	PGR	A*24
412	STIRGELFFF	MMP11	A*24
413	RYMKKDYL I	SLC35D3	A*24
414	TDSIHAWTF	SLC35D3	A*24
415	KYEKIFEML	CT45A1, CT45A2, CT45A3, CT45A4, CT45A5, CT45A6, LOC101060208,	A*24

- 21 -

Seq ID No	Sequence	Gene(s)	HLA allotype
		LOC101060210, LOC101060211	
416	VFMKDGFFYF	MMP1	A*24
417	GYIDKVRQL	NEFH	A*24
418	VHFEDTGKTLF	MMP13	A*24
419	RYVFPLPYL	SOX14	A*24
420	VYEKNGYIYF	MMP13	A*24
421	RYILENHDF	RFPL4B	A*24
422	VWSDVTPLNF	MMP13	A*24
423	IYPDVTYAF	CHRNA2	A*24
424	VQQWSVAVF	PTHLH	A*24
425	GYIDNVTLI	LAMC2	A*24
426	SVHKITSTF	LAMC2	A*24
427	VYFVAPAKF	LAMC2	A*24
428	WYVNGVNYF	TRPM8	A*24
429	YYSKSVGFMQW	FAM111B	A*24
430	VYIAELEKI	SMC1B	A*24
431	KTPTNYLFL	NMUR2	A*24
432	TRTGLFLRF	NLRP2,NLRP7	A*24
433	NYTSLLVTW	PTPRZ1	A*24
434	VYDTMIEKF	PTPRZ1	A*24
435	IYVTGGHLF	KLHDC7B	A*24
436	KYLQVVGMF	OXTR	A*24
437	VFKASKITF	PTPRZ1	A*24
438	SYSSCYSF	KRT13,KRT17	A*24
439	VYALKVRTI	CCR8	A*24
440	NYGVLHVTF	NLRP11	A*24
441	NYLVDPVTI	TRIM43,TRIM43B	A*24
442	KYLNSVQYI	RALGPS2	A*24
443	VFIHHL PQF	ACSM1	A*24
444	IYLSDLTYI	RALGPS2	A*24
445	TYIDTRTVF	CAPN6	A*24
446	IYGFFNENF	NPFFR2	A*24
447	EYIRALQQL	ASCL1	A*24
448	PFLPPAACFF	ASCL1	A*24
449	QYIEELQKF	RALGPS2	A*24
450	QYDPTPLTW	ADAMTS12	A*24
451	FGLARIYSF	CDK6	A*24
452	VYKDSIYYI	KBTBD8	A*24
453	YYTVRNFTL	PTPRZ1	A*24

Seq ID No	Sequence	Gene(s)	HLA allotype
454	TLPNTIYRF	ROS1	A*24
455	KYLSIPTVF	UGT1A3	A*24
456	PYDPALGSPSRLF	SP5	A*24
457	LIFMLANVF	GABRP	A*24
458	DYLNEWGSRF	CDH3	A*24
459	SYEVRSTF	TAS2R38	A*24
460	VYPWLGALL	CYP2W1	A*24

The present invention furthermore generally relates to the peptides according to the present invention for use in the treatment of proliferative diseases, such as cancer, such as, for example, acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

Particularly preferred are the peptides – alone or in combination - according to the present invention selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID: 464. More preferred are the peptides – alone or in combination - selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 89 and SEQ ID NO: 463 to SEQ ID: 464 (see Table 1a and Table 1b), and their uses in the immunotherapy of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, and preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric

- 23 -

cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

Another aspect of the present invention relates to the use of the peptides according to the present invention for the - preferably combined - treatment of a proliferative disease selected from the group of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

The present invention furthermore relates to peptides according to the present invention that have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or - in an elongated form, such as a length-variant - MHC class -II.

The present invention further relates to the peptides according to the present invention wherein said peptides (each) consist or consist essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID: 464.

The present invention further relates to the peptides according to the present invention, wherein said peptide is modified and/or includes non-peptide bonds.

The present invention further relates to the peptides according to the present invention, wherein said peptide is part of a fusion protein, in particular fused to the N-terminal amino acids of the HLA-DR antigen-associated invariant chain (Ii) or fused to (or into

- 24 -

the sequence of) an antibody, such as, for example, an antibody that is specific for dendritic cells.

The present invention further relates to a nucleic acid, encoding the peptides according to the present invention. The present invention further relates to the nucleic acid according to the present invention that is DNA, cDNA, PNA, RNA or combinations thereof.

The present invention further relates to an expression vector capable of expressing and/or expressing a nucleic acid according to the present invention.

The present invention further relates to a peptide according to the present invention, a nucleic acid according to the present invention or an expression vector according to the present invention for use in the treatment of diseases and in medicine, in particular in the treatment of cancer.

The present invention further relates to antibodies that are specific against the peptides according to the present invention or complexes of said peptides according to the present invention with MHC, and methods of making these.

The present invention further relates to T-cell receptors (TCRs), in particular soluble TCR (sTCRs) and cloned TCRs engineered into autologous or allogeneic T cells, and methods of making these, as well as NK cells or other cells bearing said TCR or cross-reacting with said TCRs.

The antibodies and TCRs are additional embodiments of the immunotherapeutic use of the peptides according to the invention at hand.

The present invention further relates to a host cell comprising a nucleic acid according to the present invention or an expression vector as described before. The present

- 25 -

invention further relates to the host cell according to the present invention that is an antigen presenting cell, and preferably is a dendritic cell.

The present invention further relates to a method for producing a peptide according to the present invention, said method comprising culturing the host cell according to the present invention, and isolating the peptide from said host cell or its culture medium.

The present invention further relates to said method according to the present invention, wherein the antigen is loaded onto class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell or artificial antigen-presenting cell by contacting a sufficient amount of the antigen with an antigen-presenting cell.

The present invention further relates to the method according to the present invention, wherein the antigen-presenting cell comprises an expression vector capable of expressing or expressing said peptide containing SEQ ID No. 1 to SEQ ID No.: 387 and SEQ ID NO: 463 to SEQ ID: 464, preferably containing SEQ ID No. 1 to SEQ ID No. 89 and SEQ ID NO: 463 to SEQ ID: 464, or a variant amino acid sequence.

The present invention further relates to activated T cells, produced by the method according to the present invention, wherein said T cell selectively recognizes a cell which expresses a polypeptide comprising an amino acid sequence according to the present invention.

The present invention further relates to a method of killing target cells in a patient which target cells aberrantly express a polypeptide comprising any amino acid sequence according to the present invention, the method comprising administering to the patient an effective number of T cells as produced according to the present invention.

The present invention further relates to the use of any peptide as described, the nucleic acid according to the present invention, the expression vector according to the present invention, the cell according to the present invention, the activated T lymphocyte, the T

- 26 -

cell receptor or the antibody or other peptide- and/or peptide-MHC-binding molecules according to the present invention as a medicament or in the manufacture of a medicament. Preferably, said medicament is active against cancer.

Preferably, said medicament is a cellular therapy, a vaccine or a protein based on a soluble TCR or antibody.

The present invention further relates to a use according to the present invention, wherein said cancer cells are acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, and preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells.

The present invention further relates to biomarkers based on the peptides according to the present invention, herein called "targets" that can be used in the diagnosis of cancer, preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell

- 27 -

carcinoma, urinary bladder carcinoma, uterine and endometrial cancer. The marker can be over-presentation of the peptide(s) themselves, or over-expression of the corresponding gene(s). The markers may also be used to predict the probability of success of a treatment, preferably an immunotherapy, and most preferred an immunotherapy targeting the same target that is identified by the biomarker. For example, an antibody or soluble TCR can be used to stain sections of the tumor to detect the presence of a peptide of interest in complex with MHC.

Optionally the antibody carries a further effector function such as an immune stimulating domain or toxin.

The present invention also relates to the use of these novel targets in the context of cancer treatment.

Stimulation of an immune response is dependent upon the presence of antigens recognized as foreign by the host immune system. The discovery of the existence of tumor associated antigens has raised the possibility of using a host's immune system to intervene in tumor growth. Various mechanisms of harnessing both the humoral and cellular arms of the immune system are currently being explored for cancer immunotherapy.

Specific elements of the cellular immune response are capable of specifically recognizing and destroying tumor cells. The isolation of T-cells from tumor-infiltrating cell populations or from peripheral blood suggests that such cells play an important role in natural immune defense against cancer. CD8-positive T-cells in particular, which recognize class I molecules of the major histocompatibility complex (MHC)-bearing peptides of usually 8 to 10 amino acid residues derived from proteins or defect ribosomal products (DRIPS) located in the cytosol, play an important role in this response. The MHC-molecules of the human are also designated as human leukocyte-antigens (HLA).

As used herein and except as noted otherwise all terms are defined as given below.

The term "T-cell response" means the specific proliferation and activation of effector functions induced by a peptide *in vitro* or *in vivo*. For MHC class I restricted cytotoxic T cells, effector functions may be lysis of peptide-pulsed, peptide-precursor pulsed or naturally peptide-presenting target cells, secretion of cytokines, preferably Interferon-gamma, TNF-alpha, or IL-2 induced by peptide, secretion of effector molecules, preferably granzymes or perforins induced by peptide, or degranulation.

The term "peptide" is used herein to designate a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. The peptides are preferably 9 amino acids in length, but can be as short as 8 amino acids in length, and as long as 10, 11, or 12 or longer, and in case of MHC class II peptides (elongated variants of the peptides of the invention) they can be as long as 13, 14, 15, 16, 17, 18, 19 or 20 or more amino acids in length.

Furthermore, the term "peptide" shall include salts of a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. Preferably, the salts are pharmaceutical acceptable salts of the peptides, such as, for example, the chloride or acetate (trifluoroacetate) salts. It has to be noted that the salts of the peptides according to the present invention differ substantially from the peptides in their state(s) *in vivo*, as the peptides are not salts *in vivo*.

The term "peptide" shall also include "oligopeptide". The term "oligopeptide" is used herein to designate a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. The length of the oligopeptide is not critical to the invention, as long as the correct epitope or epitopes are maintained therein. The oligopeptides are typically less

than about 30 amino acid residues in length, and greater than about 15 amino acids in length.

The term “polypeptide” designates a series of amino acid residues, connected one to the other typically by peptide bonds between the alpha-amino and carbonyl groups of the adjacent amino acids. The length of the polypeptide is not critical to the invention as long as the correct epitopes are maintained. In contrast to the terms peptide or oligopeptide, the term polypeptide is meant to refer to molecules containing more than about 30 amino acid residues.

A peptide, oligopeptide, protein or polynucleotide coding for such a molecule is “immunogenic” (and thus is an “immunogen” within the present invention), if it is capable of inducing an immune response. In the case of the present invention, immunogenicity is more specifically defined as the ability to induce a T-cell response. Thus, an “immunogen” would be a molecule that is capable of inducing an immune response, and in the case of the present invention, a molecule capable of inducing a T-cell response. In another aspect, the immunogen can be the peptide, the complex of the peptide with MHC, oligopeptide, and/or protein that is used to raise specific antibodies or TCRs against it.

A class I T cell “epitope” requires a short peptide that is bound to a class I MHC receptor, forming a ternary complex (MHC class I alpha chain, beta-2-microglobulin, and peptide) that can be recognized by a T cell bearing a matching T-cell receptor binding to the MHC/peptide complex with appropriate affinity. Peptides binding to MHC class I molecules are typically 8-14 amino acids in length, and most typically 9 amino acids in length.

In humans there are three different genetic loci that encode MHC class I molecules (the MHC-molecules of the human are also designated human leukocyte antigens (HLA)): HLA-A, HLA-B, and HLA-C. HLA-A*01, HLA-A*02, and HLA-B*07 are examples of different MHC class I alleles that can be expressed from these loci.

Table 4: Expression frequencies F of HLA-A*02, HLA-A*01, HLA-A*03, HLA-A*24, HLA-B*07, HLA-B*08 and HLA-B*44 serotypes. Haplotype frequencies Gf are derived from a study which used HLA-typing data from a registry of more than 6.5 million volunteer donors in the U.S. (Gragert et al., 2013). The haplotype frequency is the frequency of a distinct allele on an individual chromosome. Due to the diploid set of chromosomes within mammalian cells, the frequency of genotypic occurrence of this allele is higher and can be calculated employing the Hardy-Weinberg principle ($F = 1 - (1-Gf)^2$).

Allele	Population	Calculated phenotype from allele frequency (F)
A*02	African (N=28557)	32.3%
	European Caucasian (N=1242890)	49.3%
	Japanese (N=24582)	42.7%
	Hispanic, S + Cent Amer. (N=146714)	46.1%
	Southeast Asian (N=27978)	30.4%
A*01	African (N=28557)	10.2%
	European Caucasian (N=1242890)	30.2%
	Japanese (N=24582)	1.8%
	Hispanic, S + Cent Amer. (N=146714)	14.0%
	Southeast Asian (N=27978)	21.0%
A*03	African (N=28557)	14.8%
	European Caucasian (N=1242890)	26.4%
	Japanese (N=24582)	1.8%
	Hispanic, S + Cent Amer. (N=146714)	14.4%
	Southeast Asian (N=27978)	10.6%
A*24	African (N=28557)	2.0%
	European Caucasian (N=1242890)	8.6%
	Japanese (N=24582)	35.5%
	Hispanic, S + Cent Amer. (N=146714)	13.6%
	Southeast Asian (N=27978)	16.9%
B*07	African (N=28557)	14.7%

Allele	Population	Calculated phenotype from allele frequency (F)
	European Caucasian (N=1242890)	25.0%
	Japanese (N=24582)	11.4%
	Hispanic, S + Cent Amer. (N=146714)	12.2%
	Southeast Asian (N=27978)	10.4%
B*08	African (N=28557)	6.0%
	European Caucasian (N=1242890)	21.6%
	Japanese (N=24582)	1.0%
	Hispanic, S + Cent Amer. (N=146714)	7.6%
	Southeast Asian (N=27978)	6.2%
B*44	African (N=28557)	10.6%
	European Caucasian (N=1242890)	26.9%
	Japanese (N=24582)	13.0%
	Hispanic, S + Cent Amer. (N=146714)	18.2%
	Southeast Asian (N=27978)	13.1%

The peptides of the invention, preferably when included into a vaccine of the invention as described herein bind to A*24. A vaccine may also include pan-binding MHC class II peptides. Therefore, the vaccine of the invention can be used to treat cancer in patients that are A*24-positive, whereas no selection for MHC class II allotypes is necessary due to the pan-binding nature of these peptides.

If A*24 peptides of the invention are combined with peptides binding to another allele, for example A*02, a higher percentage of any patient population can be treated compared with addressing either MHC class I allele alone. While in most populations less than 50% of patients could be addressed by either allele alone, a vaccine comprising HLA-A*24 and HLA-A*02 epitopes can treat at least 60% of patients in any relevant population. Specifically, the following percentages of patients will be positive for at least one of these alleles in various regions: USA 61%, Western Europe 62%, China 75%, South Korea 77%, Japan 86% (calculated from www.allelefrequencies.net).

Table 5: HLA alleles coverage in European Caucasian population (calculated from (Gragert et al., 2013)).

	coverage (at least one A-allele)	combined with B*07	combined with B*44	combined with B*07 and B*44
A*02 / A*01	70%	78%	78%	84%
A*02 / A*03	68%	76%	76%	83%
A*02 / A*24	61%	71%	71%	80%
A*01 / A*03	52%	64%	65%	75%
A*01 / A*24	44%	58%	59%	71%
A*03 / A*24	40%	55%	56%	69%
A*02 / A*01 / A*03	84%	88%	88%	91%
A*02 / A*01 / A*24	79%	84%	84%	89%
A*02 / A*03 / A*24	77%	82%	83%	88%
A*01 / A*03 / A*24	63%	72%	73%	81%
A*02 / A*01 / A*03 / A*24	90%	92%	93%	95%

In a preferred embodiment, the term “nucleotide sequence” refers to a heteropolymer of deoxyribonucleotides.

The nucleotide sequence coding for a particular peptide, oligopeptide, or polypeptide may be naturally occurring or they may be synthetically constructed. Generally, DNA segments encoding the peptides, polypeptides, and proteins of this invention are assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene that is capable of being expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial or viral operon.

As used herein the term “a nucleotide coding for (or encoding) a peptide” refers to a nucleotide sequence coding for the peptide including artificial (man-made) start and stop codons compatible for the biological system the sequence is to be expressed by, for example, a dendritic cell or another cell system useful for the production of TCRs.

As used herein, reference to a nucleic acid sequence includes both single stranded and double stranded nucleic acid. Thus, for example for DNA, the specific sequence, unless the context indicates otherwise, refers to the single strand DNA of such sequence, the duplex of such sequence with its complement (double stranded DNA) and the complement of such sequence.

The term “coding region” refers to that portion of a gene which either naturally or normally codes for the expression product of that gene in its natural genomic environment, i.e., the region coding *in vivo* for the native expression product of the gene.

The coding region can be derived from a non-mutated (“normal”), mutated or altered gene, or can even be derived from a DNA sequence, or gene, wholly synthesized in the laboratory using methods well known to those of skill in the art of DNA synthesis.

The term “expression product” means the polypeptide or protein that is the natural translation product of the gene and any nucleic acid sequence coding equivalents resulting from genetic code degeneracy and thus coding for the same amino acid(s).

The term “fragment”, when referring to a coding sequence, means a portion of DNA comprising less than the complete coding region, whose expression product retains essentially the same biological function or activity as the expression product of the complete coding region.

The term “DNA segment” refers to a DNA polymer, in the form of a separate fragment or as a component of a larger DNA construct, which has been derived from DNA isolated at least once in substantially pure form, i.e., free of contaminating endogenous materials and in a quantity or concentration enabling identification, manipulation, and recovery of the segment and its component nucleotide sequences by standard biochemical methods, for example, by using a cloning vector. Such segments are provided in the form of an open reading frame uninterrupted by internal non-translated sequences, or

- 34 -

introns, which are typically present in eukaryotic genes. Sequences of non-translated DNA may be present downstream from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

The term “primer” means a short nucleic acid sequence that can be paired with one strand of DNA and provides a free 3'-OH end at which a DNA polymerase starts synthesis of a deoxyribonucleotide chain.

The term “promoter” means a region of DNA involved in binding of RNA polymerase to initiate transcription.

The term “isolated” means that the material is removed from its original environment (e.g., the natural environment, if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The polynucleotides, and recombinant or immunogenic polypeptides, disclosed in accordance with the present invention may also be in “purified” form. The term “purified” does not require absolute purity; rather, it is intended as a relative definition, and can include preparations that are highly purified or preparations that are only partially purified, as those terms are understood by those of skill in the relevant art. For example, individual clones isolated from a cDNA library have been conventionally purified to electrophoretic homogeneity. Purification of starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated. Furthermore, a claimed polypeptide which has a purity of preferably 99.999%, or at least 99.99% or 99.9%; and even desirably 99% by weight or greater is expressly encompassed.

- 35 -

The nucleic acids and polypeptide expression products disclosed according to the present invention, as well as expression vectors containing such nucleic acids and/or such polypeptides, may be in "enriched form". As used herein, the term "enriched" means that the concentration of the material is at least about 2, 5, 10, 100, or 1000 times its natural concentration (for example), advantageously 0.01%, by weight, preferably at least about 0.1% by weight. Enriched preparations of about 0.5%, 1%, 5%, 10%, and 20% by weight are also contemplated. The sequences, constructs, vectors, clones, and other materials comprising the present invention can advantageously be in enriched or isolated form. The term "active fragment" means a fragment, usually of a peptide, polypeptide or nucleic acid sequence, that generates an immune response (i.e., has immunogenic activity) when administered, alone or optionally with a suitable adjuvant or in a vector, to an animal, such as a mammal, for example, a rabbit or a mouse, and also including a human, such immune response taking the form of stimulating a T-cell response within the recipient animal, such as a human. Alternatively, the "active fragment" may also be used to induce a T-cell response *in vitro*.

As used herein, the terms "portion", "segment" and "fragment", when used in relation to polypeptides, refer to a continuous sequence of residues, such as amino acid residues, which sequence forms a subset of a larger sequence. For example, if a polypeptide were subjected to treatment with any of the common endopeptidases, such as trypsin or chymotrypsin, the oligopeptides resulting from such treatment would represent portions, segments or fragments of the starting polypeptide. When used in relation to polynucleotides, these terms refer to the products produced by treatment of said polynucleotides with any of the endonucleases.

In accordance with the present invention, the term "percent identity" or "percent identical", when referring to a sequence, means that a sequence is compared to a claimed or described sequence after alignment of the sequence to be compared (the "Compared Sequence") with the described or claimed sequence (the "Reference Sequence"). The percent identity is then determined according to the following formula:
percent identity = $100 [1 - (C/R)]$

- 36 -

wherein C is the number of differences between the Reference Sequence and the Compared Sequence over the length of alignment between the Reference Sequence and the Compared Sequence, wherein

(i) each base or amino acid in the Reference Sequence that does not have a corresponding aligned base or amino acid in the Compared Sequence and

(ii) each gap in the Reference Sequence and

(iii) each aligned base or amino acid in the Reference Sequence that is different from an aligned base or amino acid in the Compared Sequence, constitutes a difference and

(iiii) the alignment has to start at position 1 of the aligned sequences;

and R is the number of bases or amino acids in the Reference Sequence over the length of the alignment with the Compared Sequence with any gap created in the Reference Sequence also being counted as a base or amino acid.

If an alignment exists between the Compared Sequence and the Reference Sequence for which the percent identity as calculated above is about equal to or greater than a specified minimum Percent Identity then the Compared Sequence has the specified minimum percent identity to the Reference Sequence even though alignments may exist in which the herein above calculated percent identity is less than the specified percent identity.

As mentioned above, the present invention thus provides a peptide comprising a sequence that is selected from the group of consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464 or a variant thereof which is 88% homologous to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464, or a variant thereof that will induce T cells cross-reacting with said peptide. The peptides of the invention have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or elongated versions of said peptides to class II.

In the present invention, the term "homologous" shall refer to the degree of identity (see percent identity above) between sequences of two amino acid sequences, i.e. peptide

or polypeptide sequences. The aforementioned "homology" is determined by comparing two sequences aligned under optimal conditions over the sequences to be compared. Such a sequence homology can be calculated by creating an alignment using, for example, the ClustalW algorithm. Commonly available sequence analysis software, more specifically, Vector NTI, GENETYX or other tools are provided by public databases.

A person skilled in the art will be able to assess, whether T cells induced by a variant of a specific peptide will be able to cross-react with the peptide itself (Appay et al., 2006; Colombetti et al., 2006; Fong et al., 2001; Zaremba et al., 1997).

By a "variant" of the given amino acid sequence the inventors mean that the side chains of, for example, one or two of the amino acid residues are altered (for example by replacing them with the side chain of another naturally occurring amino acid residue or some other side chain) such that the peptide is still able to bind to an HLA molecule in substantially the same way as a peptide consisting of the given amino acid sequence in consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464. For example, a peptide may be modified so that it at least maintains, if not improves, the ability to interact with and bind to the binding groove of a suitable MHC molecule, such as HLA-A*02 or -DR, and in that way it at least maintains, if not improves, the ability to bind to the TCR of activated T cells.

These T cells can subsequently cross-react with cells and kill cells that express a polypeptide that contains the natural amino acid sequence of the cognate peptide as defined in the aspects of the invention. As can be derived from the scientific literature and databases (Godkin et al., 1997; Rammensee et al., 1999), certain positions of HLA binding peptides are typically anchor residues forming a core sequence fitting to the binding motif of the HLA receptor, which is defined by polar, electrophysical, hydrophobic and spatial properties of the polypeptide chains constituting the binding groove. Thus, one skilled in the art would be able to modify the amino acid sequences set forth in SEQ ID NO: 1 to SEQ ID NO 387 and SEQ ID NO: 463 to SEQ ID NO: 464,

- 38 -

by maintaining the known anchor residues, and would be able to determine whether such variants maintain the ability to bind MHC class I or II molecules. The variants of the present invention retain the ability to bind to the TCR of activated T cells, which can subsequently cross-react with and kill cells that express a polypeptide containing the natural amino acid sequence of the cognate peptide as defined in the aspects of the invention.

The original (unmodified) peptides as disclosed herein can be modified by the substitution of one or more residues at different, possibly selective, sites within the peptide chain, if not otherwise stated. Preferably those substitutions are located at the end of the amino acid chain. Such substitutions may be of a conservative nature, for example, where one amino acid is replaced by an amino acid of similar structure and characteristics, such as where a hydrophobic amino acid is replaced by another hydrophobic amino acid. Even more conservative would be replacement of amino acids of the same or similar size and chemical nature, such as where leucine is replaced by isoleucine. In studies of sequence variations in families of naturally occurring homologous proteins, certain amino acid substitutions are more often tolerated than others, and these are often show correlation with similarities in size, charge, polarity, and hydrophobicity between the original amino acid and its replacement, and such is the basis for defining "conservative substitutions."

Conservative substitutions are herein defined as exchanges within one of the following five groups: Group 1-small aliphatic, nonpolar or slightly polar residues (Ala, Ser, Thr, Pro, Gly); Group 2-polar, negatively charged residues and their amides (Asp, Asn, Glu, Gln); Group 3-polar, positively charged residues (His, Arg, Lys); Group 4-large, aliphatic, nonpolar residues (Met, Leu, Ile, Val, Cys); and Group 5-large, aromatic residues (Phe, Tyr, Trp).

Less conservative substitutions might involve the replacement of one amino acid by another that has similar characteristics but is somewhat different in size, such as replacement of an alanine by an isoleucine residue. Highly non-conservative

- 39 -

replacements might involve substituting an acidic amino acid for one that is polar, or even for one that is basic in character. Such “radical” substitutions cannot, however, be dismissed as potentially ineffective since chemical effects are not totally predictable and radical substitutions might well give rise to serendipitous effects not otherwise predictable from simple chemical principles.

Of course, such substitutions may involve structures other than the common L-amino acids. Thus, D-amino acids might be substituted for the L-amino acids commonly found in the antigenic peptides of the invention and yet still be encompassed by the disclosure herein. In addition, non-standard amino acids (i.e., other than the common naturally occurring proteinogenic amino acids) may also be used for substitution purposes to produce immunogens and immunogenic polypeptides according to the present invention.

If substitutions at more than one position are found to result in a peptide with substantially equivalent or greater antigenic activity as defined below, then combinations of those substitutions will be tested to determine if the combined substitutions result in additive or synergistic effects on the antigenicity of the peptide. At most, no more than 4 positions within the peptide would be simultaneously substituted.

A peptide consisting essentially of the amino acid sequence as indicated herein can have one or two non-anchor amino acids (see below regarding the anchor motif) exchanged without that the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or –II is substantially changed or is negatively affected, when compared to the non-modified peptide. In another embodiment, in a peptide consisting essentially of the amino acid sequence as indicated herein, one or two amino acids can be exchanged with their conservative exchange partners (see herein below) without that the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or –II is substantially changed, or is negatively affected, when compared to the non-modified peptide.

- 41 -

		F								L
		F								

Longer (elongated) peptides may also be suitable. It is possible that MHC class I epitopes, although usually between 8 and 11 amino acids long, are generated by peptide processing from longer peptides or proteins that include the actual epitope. It is preferred that the residues that flank the actual epitope are residues that do not substantially affect proteolytic cleavage necessary to expose the actual epitope during processing.

The peptides of the invention can be elongated by up to four amino acids, that is 1, 2, 3 or 4 amino acids can be added to either end in any combination between 4:0 and 0:4. Combinations of the elongations according to the invention can be found in Table 7.

Table 7: Combinations of the elongations of peptides of the invention

C-terminus	N-terminus
4	0
3	0 or 1
2	0 or 1 or 2
1	0 or 1 or 2 or 3
0	0 or 1 or 2 or 3 or 4
N-terminus	C-terminus
4	0
3	0 or 1
2	0 or 1 or 2
1	0 or 1 or 2 or 3
0	0 or 1 or 2 or 3 or 4

The amino acids for the elongation/extension can be the peptides of the original sequence of the protein or any other amino acid(s). The elongation can be used to enhance the stability or solubility of the peptides.

Thus, the epitopes of the present invention may be identical to naturally occurring tumor-associated or tumor-specific epitopes or may include epitopes that differ by no more than four residues from the reference peptide, as long as they have substantially identical antigenic activity.

- 42 -

In an alternative embodiment, the peptide is elongated on either or both sides by more than 4 amino acids, preferably to a total length of up to 30 amino acids. This may lead to MHC class II binding peptides. Binding to MHC class II can be tested by methods known in the art.

Accordingly, the present invention provides peptides and variants of MHC class I epitopes, wherein the peptide or variant has an overall length of between 8 and 100, preferably between 8 and 30, and most preferred between 8 and 14, namely 8, 9, 10, 11, 12, 13, 14 amino acids, in case of the elongated class II binding peptides the length can also be 15, 16, 17, 18, 19, 20, 21 or 22 amino acids.

Of course, the peptide or variant according to the present invention will have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class I or II. Binding of a peptide or a variant to a MHC complex may be tested by methods known in the art.

Preferably, when the T cells specific for a peptide according to the present invention are tested against the substituted peptides, the peptide concentration at which the substituted peptides achieve half the maximal increase in lysis relative to background is no more than about 1 mM, preferably no more than about 1 μ M, more preferably no more than about 1 nM, and still more preferably no more than about 100 pM, and most preferably no more than about 10 pM. It is also preferred that the substituted peptide be recognized by T cells from more than one individual, at least two, and more preferably three individuals.

In a particularly preferred embodiment of the invention the peptide consists or consists essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464.

- 43 -

“Consisting essentially of” shall mean that a peptide according to the present invention, in addition to the sequence according to any one of SEQ ID NO: 1 to SEQ ID NO 387 and SEQ ID NO: 463 to SEQ ID NO: 464 or a variant thereof contains additional N- and/or C-terminally located stretches of amino acids that are not necessarily forming part of the peptide that functions as an epitope for MHC molecules epitope.

Nevertheless, these stretches can be important to provide an efficient introduction of the peptide according to the present invention into the cells. In one embodiment of the present invention, the peptide is part of a fusion protein which comprises, for example, the 80 N-terminal amino acids of the HLA-DR antigen-associated invariant chain (p33, in the following “Ii”) as derived from the NCBI, GenBank Accession number X00497. In other fusions, the peptides of the present invention can be fused to an antibody as described herein, or a functional part thereof, in particular into a sequence of an antibody, so as to be specifically targeted by said antibody, or, for example, to or into an antibody that is specific for dendritic cells as described herein.

In addition, the peptide or variant may be modified further to improve stability and/or binding to MHC molecules in order to elicit a stronger immune response. Methods for such an optimization of a peptide sequence are well known in the art and include, for example, the introduction of reverse peptide bonds or non-peptide bonds.

In a reverse peptide bond amino acid residues are not joined by peptide (-CO-NH-) linkages but the peptide bond is reversed. Such retro-inverso peptidomimetics may be made using methods known in the art, for example such as those described in Meziere et al (1997) (Meziere et al., 1997), incorporated herein by reference. This approach involves making pseudopeptides containing changes involving the backbone, and not the orientation of side chains. Meziere et al. (Meziere et al., 1997) show that for MHC binding and T helper cell responses, these pseudopeptides are useful. Retro-inverse peptides, which contain NH-CO bonds instead of CO-NH peptide bonds, are much more resistant to proteolysis.

- 44 -

A non-peptide bond is, for example, $-\text{CH}_2\text{-NH}$, $-\text{CH}_2\text{S-}$, $-\text{CH}_2\text{CH}_2\text{-}$, $-\text{CH}=\text{CH-}$, $-\text{COCH}_2\text{-}$, $-\text{CH}(\text{OH})\text{CH}_2\text{-}$, and $-\text{CH}_2\text{SO-}$. US 4,897,445 provides a method for the solid phase synthesis of non-peptide bonds ($-\text{CH}_2\text{-NH}$) in polypeptide chains which involves polypeptides synthesized by standard procedures and the non-peptide bond synthesized by reacting an amino aldehyde and an amino acid in the presence of NaCNBH_3 .

Peptides comprising the sequences described above may be synthesized with additional chemical groups present at their amino and/or carboxy termini, to enhance the stability, bioavailability, and/or affinity of the peptides. For example, hydrophobic groups such as carbobenzoxy, dansyl, or t-butyloxycarbonyl groups may be added to the peptides' amino termini. Likewise, an acetyl group or a 9-fluorenylmethoxy-carbonyl group may be placed at the peptides' amino termini. Additionally, the hydrophobic group, t-butyloxycarbonyl, or an amido group may be added to the peptides' carboxy termini.

Further, the peptides of the invention may be synthesized to alter their steric configuration. For example, the D-isomer of one or more of the amino acid residues of the peptide may be used, rather than the usual L-isomer. Still further, at least one of the amino acid residues of the peptides of the invention may be substituted by one of the well-known non-naturally occurring amino acid residues. Alterations such as these may serve to increase the stability, bioavailability and/or binding action of the peptides of the invention.

Similarly, a peptide or variant of the invention may be modified chemically by reacting specific amino acids either before or after synthesis of the peptide. Examples for such modifications are well known in the art and are summarized e.g. in R. Lundblad, *Chemical Reagents for Protein Modification*, 3rd ed. CRC Press, 2004 (Lundblad, 2004), which is incorporated herein by reference. Chemical modification of amino acids includes but is not limited to, modification by acylation, amidination, pyridoxylation of lysine, reductive alkylation, trinitrobenzylation of amino groups with 2,4,6-trinitrobenzene

- 45 -

sulphonic acid (TNBS), amide modification of carboxyl groups and sulphydryl modification by performic acid oxidation of cysteine to cysteic acid, formation of mercurial derivatives, formation of mixed disulphides with other thiol compounds, reaction with maleimide, carboxymethylation with iodoacetic acid or iodoacetamide and carbamoylation with cyanate at alkaline pH, although without limitation thereto. In this regard, the skilled person is referred to Chapter 15 of Current Protocols In Protein Science, Eds. Coligan et al. (John Wiley and Sons NY 1995-2000) (Coligan et al., 1995) for more extensive methodology relating to chemical modification of proteins.

Briefly, modification of e.g. arginyl residues in proteins is often based on the reaction of vicinal dicarbonyl compounds such as phenylglyoxal, 2,3-butanedione, and 1,2-cyclohexanedione to form an adduct. Another example is the reaction of methylglyoxal with arginine residues. Cysteine can be modified without concomitant modification of other nucleophilic sites such as lysine and histidine. As a result, a large number of reagents are available for the modification of cysteine. The websites of companies such as Sigma-Aldrich (<http://www.sigma-aldrich.com>) provide information on specific reagents.

Selective reduction of disulfide bonds in proteins is also common. Disulfide bonds can be formed and oxidized during the heat treatment of biopharmaceuticals. Woodward's Reagent K may be used to modify specific glutamic acid residues. N-(3-(dimethylamino)propyl)-N'-ethylcarbodiimide can be used to form intra-molecular crosslinks between a lysine residue and a glutamic acid residue. For example, diethylpyrocarbonate is a reagent for the modification of histidyl residues in proteins. Histidine can also be modified using 4-hydroxy-2-nonenal. The reaction of lysine residues and other α -amino groups is, for example, useful in binding of peptides to surfaces or the cross-linking of proteins/peptides. Lysine is the site of attachment of poly(ethylene)glycol and the major site of modification in the glycosylation of proteins. Methionine residues in proteins can be modified with e.g. iodoacetamide, bromoethylamine, and chloramine T.

- 46 -

Tetranitromethane and N-acetylimidazole can be used for the modification of tyrosyl residues. Cross-linking via the formation of dityrosine can be accomplished with hydrogen peroxide/copper ions.

Recent studies on the modification of tryptophan have used N-bromosuccinimide, 2-hydroxy-5-nitrobenzyl bromide or 3-bromo-3-methyl-2-(2-nitrophenylmercapto)-3H-indole (BPNS-skatole).

Successful modification of therapeutic proteins and peptides with PEG is often associated with an extension of circulatory half-life while cross-linking of proteins with glutaraldehyde, polyethylene glycol diacrylate and formaldehyde is used for the preparation of hydrogels. Chemical modification of allergens for immunotherapy is often achieved by carbamylation with potassium cyanate.

A peptide or variant, wherein the peptide is modified or includes non-peptide bonds is a preferred embodiment of the invention.

Another embodiment of the present invention relates to a non-naturally occurring peptide wherein said peptide consists or consists essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID NO: 463 to SEQ ID NO: 464 and has been synthetically produced (e.g. synthesized) as a pharmaceutically acceptable salt. Methods to synthetically produce peptides are well known in the art. The salts of the peptides according to the present invention differ substantially from the peptides in their state(s) *in vivo*, as the peptides as generated *in vivo* are no salts. The non-natural salt form of the peptide mediates the solubility of the peptide, in particular in the context of pharmaceutical compositions comprising the peptides, e.g. the peptide vaccines as disclosed herein. A sufficient and at least substantial solubility of the peptide(s) is required in order to efficiently provide the peptides to the subject to be treated. Preferably, the salts are pharmaceutically acceptable salts of the peptides. These salts according to the invention include alkaline and earth alkaline salts such as salts of the Hofmeister series comprising as anions PO_4^{3-} , SO_4^{2-} , CH_3COO^- , Cl^- , Br^- ,

- 47 -

NO₃⁻, ClO₄⁻, I⁻, SCN⁻ and as cations NH₄⁺, Rb⁺, K⁺, Na⁺, Cs⁺, Li⁺, Zn²⁺, Mg²⁺, Ca²⁺, Mn²⁺, Cu²⁺ and Ba²⁺. Particularly salts are selected from (NH₄)₃PO₄, (NH₄)₂HPO₄, (NH₄)H₂PO₄, (NH₄)₂SO₄, NH₄CH₃COO, NH₄Cl, NH₄Br, NH₄NO₃, NH₄ClO₄, NH₄I, NH₄SCN, Rb₃PO₄, Rb₂HPO₄, RbH₂PO₄, Rb₂SO₄, Rb₄CH₃COO, Rb₄Cl, Rb₄Br, Rb₄NO₃, Rb₄ClO₄, Rb₄I, Rb₄SCN, K₃PO₄, K₂HPO₄, KH₂PO₄, K₂SO₄, KCH₃COO, KCl, KBr, KNO₃, KClO₄, KI, KSCN, Na₃PO₄, Na₂HPO₄, NaH₂PO₄, Na₂SO₄, NaCH₃COO, NaCl, NaBr, NaNO₃, NaClO₄, NaI, NaSCN, ZnCl₂, Cs₃PO₄, Cs₂HPO₄, CsH₂PO₄, Cs₂SO₄, CsCH₃COO, CsCl, CsBr, CsNO₃, CsClO₄, CsI, CsSCN, Li₃PO₄, Li₂HPO₄, LiH₂PO₄, Li₂SO₄, LiCH₃COO, LiCl, LiBr, LiNO₃, LiClO₄, LiI, LiSCN, Cu₂SO₄, Mg₃(PO₄)₂, Mg₂HPO₄, Mg(H₂PO₄)₂, Mg₂SO₄, Mg(CH₃COO)₂, MgCl₂, MgBr₂, Mg(NO₃)₂, Mg(ClO₄)₂, MgI₂, Mg(SCN)₂, MnCl₂, Ca₃(PO₄)₂, Ca₂HPO₄, Ca(H₂PO₄)₂, CaSO₄, Ca(CH₃COO)₂, CaCl₂, CaBr₂, Ca(NO₃)₂, Ca(ClO₄)₂, CaI₂, Ca(SCN)₂, Ba₃(PO₄)₂, Ba₂HPO₄, Ba(H₂PO₄)₂, BaSO₄, Ba(CH₃COO)₂, BaCl₂, BaBr₂, Ba(NO₃)₂, Ba(ClO₄)₂, BaI₂, and Ba(SCN)₂. Particularly preferred are NH acetate, MgCl₂, KH₂PO₄, Na₂SO₄, KCl, NaCl, and CaCl₂, such as, for example, the chloride or acetate (trifluoroacetate) salts.

Generally, peptides and variants (at least those containing peptide linkages between amino acid residues) may be synthesized by the Fmoc-polyamide mode of solid-phase peptide synthesis as disclosed by Lukas et al. (Lukas et al., 1981) and by references as cited therein. Temporary N-amino group protection is afforded by the 9-fluorenylmethyloxycarbonyl (Fmoc) group. Repetitive cleavage of this highly base-labile protecting group is done using 20% piperidine in N, N-dimethylformamide. Side-chain functionalities may be protected as their butyl ethers (in the case of serine threonine and tyrosine), butyl esters (in the case of glutamic acid and aspartic acid), butyloxycarbonyl derivative (in the case of lysine and histidine), trityl derivative (in the case of cysteine) and 4-methoxy-2,3,6-trimethylbenzenesulphonyl derivative (in the case of arginine). Where glutamine or asparagine are C-terminal residues, use is made of the 4,4'-dimethoxybenzhydryl group for protection of the side chain amido functionalities. The solid-phase support is based on a polydimethyl-acrylamide polymer constituted from the three monomers dimethylacrylamide (backbone-monomer), bisacryloylethylene diamine (cross linker) and acryloylsarcosine methyl ester

- 48 -

(functionalizing agent). The peptide-to-resin cleavable linked agent used is the acid-labile 4-hydroxymethyl-phenoxyacetic acid derivative. All amino acid derivatives are added as their preformed symmetrical anhydride derivatives with the exception of asparagine and glutamine, which are added using a reversed N, N-dicyclohexylcarbodiimide/1hydroxybenzotriazole mediated coupling procedure. All coupling and deprotection reactions are monitored using ninhydrin, trinitrobenzene sulphonic acid or isotin test procedures. Upon completion of synthesis, peptides are cleaved from the resin support with concomitant removal of side-chain protecting groups by treatment with 95% trifluoroacetic acid containing a 50 % scavenger mix. Scavengers commonly used include ethanedithiol, phenol, anisole and water, the exact choice depending on the constituent amino acids of the peptide being synthesized. Also a combination of solid phase and solution phase methodologies for the synthesis of peptides is possible (see, for example, (Bruckdorfer et al., 2004), and the references as cited therein).

Trifluoroacetic acid is removed by evaporation in vacuo, with subsequent trituration with diethyl ether affording the crude peptide. Any scavengers present are removed by a simple extraction procedure which on lyophilization of the aqueous phase affords the crude peptide free of scavengers. Reagents for peptide synthesis are generally available from e.g. Calbiochem-Novabiochem (Nottingham, UK).

Purification may be performed by any one, or a combination of, techniques such as re-crystallization, size exclusion chromatography, ion-exchange chromatography, hydrophobic interaction chromatography and (usually) reverse-phase high performance liquid chromatography using e.g. acetonitrile/water gradient separation.

Analysis of peptides may be carried out using thin layer chromatography, electrophoresis, in particular capillary electrophoresis, solid phase extraction (CSPE), reverse-phase high performance liquid chromatography, amino-acid analysis after acid hydrolysis and by fast atom bombardment (FAB) mass spectrometric analysis, as well as MALDI and ESI-Q-TOF mass spectrometric analysis.

In order to select over-presented peptides, a presentation profile is calculated showing the median sample presentation as well as replicate variation. The profile juxtaposes samples of the tumor entity of interest to a baseline of normal tissue samples. Each of these profiles can then be consolidated into an over-presentation score by calculating the p-value of a Linear Mixed-Effects Model (Pineiro et al., 2015) adjusting for multiple testing by False Discovery Rate (Benjamini and Hochberg, 1995) (cf. Example 1, Figure 1).

For the identification and relative quantitation of HLA ligands by mass spectrometry, HLA molecules from shock-frozen tissue samples were purified and HLA-associated peptides were isolated. The isolated peptides were separated, and sequences were identified by online nano-electrospray-ionization (nanoESI) liquid chromatography-mass spectrometry (LC-MS) experiments. The resulting peptide sequences were verified by comparison of the fragmentation pattern of natural tumor-associated peptides (TUMAPs) recorded from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples (N = 263 samples) with the fragmentation patterns of corresponding synthetic reference peptides of identical sequences. Since the peptides were directly identified as ligands of HLA molecules of primary tumors, these results provide direct evidence for the natural processing and presentation of the identified peptides on primary cancer tissue obtained from 263 acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer,

renal cell carcinoma, urinary bladder carcinoma, and uterine and endometrial cancer patients.

The discovery pipeline XPRESIDENT® v2.1 (see, for example, US 2013-0096016, which is hereby incorporated by reference in its entirety) allows the identification and selection of relevant over-presented peptide vaccine candidates based on direct relative quantitation of HLA-restricted peptide levels on cancer tissues in comparison to several different non-cancerous tissues and organs. This was achieved by the development of label-free differential quantitation using the acquired LC-MS data processed by a proprietary data analysis pipeline, combining algorithms for sequence identification, spectral clustering, ion counting, retention time alignment, charge state deconvolution and normalization.

Presentation levels including error estimates for each peptide and sample were established. Peptides exclusively presented on tumor tissue and peptides over-presented in tumor versus non-cancerous tissues and organs have been identified.

HLA-peptide complexes from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer tissue samples were purified and HLA-associated peptides were isolated and analyzed by LC-MS (see example 1). All TUMAPs contained in the present application were identified with this approach on acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell

- 51 -

lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples confirming their presentation on acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

TUMAPs identified on multiple acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer and normal tissues were quantified using ion-counting of label-free LC-MS data. The method assumes that LC-MS signal areas of a peptide correlate with its abundance in the sample. All quantitative signals of a peptide in various LC-MS experiments were normalized based on central tendency, averaged per sample and merged into a bar plot, called presentation profile. The presentation profile consolidates different analysis methods like protein database search, spectral clustering, charge state deconvolution (decharging) and retention time alignment and normalization.

Besides over-presentation of the peptide, mRNA expression of the underlying gene was tested. mRNA data were obtained via RNASeq analyses of normal tissues and cancer tissues (cf. Example 2, Figure 2). Peptides which are derived from proteins whose coding mRNA is highly expressed in cancer tissue, but very low or absent in vital normal tissues, were preferably included in the present invention.

The present invention provides peptides that are useful in treating cancers/tumors, preferably acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer that over- or exclusively present the peptides of the invention. These peptides were shown by mass spectrometry to be naturally presented by HLA molecules on primary human acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples.

Many of the source gene/proteins (also designated “full-length proteins” or “underlying proteins”) from which the peptides are derived were shown to be highly over-expressed in cancer compared with normal tissues – “normal tissues” in relation to this invention shall mean either healthy blood cells, blood vessels, brain, heart, liver, lung, adipose tissue, adrenal gland, bile duct, bladder, bone marrow, esophagus, eye, gallbladder, head & neck, large intestine, small intestine, kidney, lymph node, peripheral nerve, pancreas, parathyroid gland, peritoneum, pituitary, pleura, skeletal muscle, skin, spleen, stomach, thyroid, trachea, ureter cells or other normal tissue cells, demonstrating a high degree of tumor association of the source genes (see Example 2). Moreover, the peptides themselves are strongly over-presented on tumor tissue – “tumor tissue” in relation to this invention shall mean a sample from a patient suffering from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic

leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, but not on normal tissues (see Example 1).

HLA-bound peptides can be recognized by the immune system, specifically T lymphocytes. T cells can destroy the cells presenting the recognized HLA/peptide complex, e.g. acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells presenting the derived peptides.

The peptides of the present invention have been shown to be capable of stimulating T cell responses and/or are over-presented and thus can be used for the production of antibodies and/or TCRs, such as soluble TCRs, according to the present invention (see Example 3, Example 4). Furthermore, the peptides when complexed with the respective MHC can be used for the production of antibodies and/or TCRs, in particular sTCRs, according to the present invention, as well. Respective methods are well known to the person of skill, and can be found in the respective literature as well (see also below). Thus, the peptides of the present invention are useful for generating an immune response in a patient by which tumor cells can be destroyed. An immune response in a patient can be induced by direct administration of the described peptides or suitable precursor substances (e.g. elongated peptides, proteins, or nucleic acids encoding these peptides) to the patient, ideally in combination with an agent enhancing the

- 54 -

immunogenicity (i.e. an adjuvant). The immune response originating from such a therapeutic vaccination can be expected to be highly specific against tumor cells because the target peptides of the present invention are not presented on normal tissues in comparable copy numbers, preventing the risk of undesired autoimmune reactions against normal cells in the patient.

The present description further relates to T-cell receptors (TCRs) comprising an alpha chain and a beta chain ("alpha/beta TCRs"). Also provided are peptides according to the invention capable of binding to TCRs and antibodies when presented by an MHC molecule.

The present description also relates to fragments of the TCRs according to the invention that are capable of binding to a peptide antigen according to the present invention when presented by an HLA molecule. The term particularly relates to soluble TCR fragments, for example TCRs missing the transmembrane parts and/or constant regions, single chain TCRs, and fusions thereof to, for example, with Ig.

The present description also relates to nucleic acids, vectors and host cells for expressing TCRs and peptides of the present description; and methods of using the same.

The term "T-cell receptor" (abbreviated TCR) refers to a heterodimeric molecule comprising an alpha polypeptide chain (alpha chain) and a beta polypeptide chain (beta chain), wherein the heterodimeric receptor is capable of binding to a peptide antigen presented by an HLA molecule. The term also includes so-called gamma/delta TCRs.

In one embodiment the description provides a method of producing a TCR as described herein, the method comprising culturing a host cell capable of expressing the TCR under conditions suitable to promote expression of the TCR.

- 55 -

The description in another aspect relates to methods according to the description, wherein the antigen is loaded onto class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell or artificial antigen-presenting cell by contacting a sufficient amount of the antigen with an antigen-presenting cell or the antigen is loaded onto class I or II MHC tetramers by tetramerizing the antigen/class I or II MHC complex monomers.

The alpha and beta chains of alpha/beta TCR's, and the gamma and delta chains of gamma/delta TCRs, are generally regarded as each having two "domains", namely variable and constant domains. The variable domain consists of a concatenation of variable region (V) and joining region (J). The variable domain may also include a leader region (L). Beta and delta chains may also include a diversity region (D). The alpha and beta constant domains may also include C-terminal transmembrane (TM) domains that anchor the alpha and beta chains to the cell membrane.

With respect to gamma/delta TCRs, the term "TCR gamma variable domain" as used herein refers to the concatenation of the TCR gamma V (TRGV) region without leader region (L), and the TCR gamma J (TRGJ) region, and the term TCR gamma constant domain refers to the extracellular TRGC region, or to a C-terminal truncated TRGC sequence. Likewise, the term "TCR delta variable domain" refers to the concatenation of the TCR delta V (TRDV) region without leader region (L) and the TCR delta D/J (TRDD/TRDJ) region, and the term "TCR delta constant domain" refers to the extracellular TRDC region, or to a C-terminal truncated TRDC sequence.

TCRs of the present description preferably bind to a peptide-HLA molecule complex with a binding affinity (KD) of about 100 μ M or less, about 50 μ M or less, about 25 μ M or less, or about 10 μ M or less. More preferred are high affinity TCRs having binding affinities of about 1 μ M or less, about 100 nM or less, about 50 nM or less, about 25 nM or less. Non-limiting examples of preferred binding affinity ranges for TCRs of the present invention include about 1 nM to about 10 nM; about 10 nM to about 20 nM; about 20 nM to about 30 nM; about 30 nM to about 40 nM; about 40 nM to about 50 nM;

- 56 -

about 50 nM to about 60 nM; about 60 nM to about 70 nM; about 70 nM to about 80 nM; about 80 nM to about 90 nM; and about 90 nM to about 100 nM.

As used herein in connect with TCRs of the present description, "specific binding" and grammatical variants thereof are used to mean a TCR having a binding affinity (KD) for a peptide-HLA molecule complex of 100 μ M or less.

Alpha/beta heterodimeric TCRs of the present description may have an introduced disulfide bond between their constant domains. Preferred TCRs of this type include those which have a TRAC constant domain sequence and a TRBC1 or TRBC2 constant domain sequence except that Thr 48 of TRAC and Ser 57 of TRBC1 or TRBC2 are replaced by cysteine residues, the said cysteines forming a disulfide bond between the TRAC constant domain sequence and the TRBC1 or TRBC2 constant domain sequence of the TCR.

With or without the introduced inter-chain bond mentioned above, alpha/beta heterodimeric TCRs of the present description may have a TRAC constant domain sequence and a TRBC1 or TRBC2 constant domain sequence, and the TRAC constant domain sequence and the TRBC1 or TRBC2 constant domain sequence of the TCR may be linked by the native disulfide bond between Cys4 of exon 2 of TRAC and Cys2 of exon 2 of TRBC1 or TRBC2.

TCRs of the present description may comprise a detectable label selected from the group consisting of a radionuclide, a fluorophore and biotin. TCRs of the present description may be conjugated to a therapeutically active agent, such as a radionuclide, a chemotherapeutic agent, or a toxin.

In an embodiment, a TCR of the present description having at least one mutation in the alpha chain and/or having at least one mutation in the beta chain has modified glycosylation compared to the unmutated TCR.

- 57 -

In an embodiment, a TCR comprising at least one mutation in the TCR alpha chain and/or TCR beta chain has a binding affinity for, and/or a binding half-life for, a peptide-HLA molecule complex, which is at least double that of a TCR comprising the unmutated TCR alpha chain and/or unmutated TCR beta chain. Affinity-enhancement of tumor-specific TCRs, and its exploitation, relies on the existence of a window for optimal TCR affinities. The existence of such a window is based on observations that TCRs specific for HLA-A-restricted pathogens have KD values that are generally about 10-fold lower when compared to TCRs specific for HLA-A-restricted tumor-associated self-antigens. It is now known, although tumor antigens have the potential to be immunogenic, because tumors arise from the individual's own cells only mutated proteins or proteins with altered translational processing will be seen as foreign by the immune system. Antigens that are upregulated or overexpressed (so called self-antigens) will not necessarily induce a functional immune response against the tumor: T-cells expressing TCRs that are highly reactive to these antigens will have been negatively selected within the thymus in a process known as central tolerance, meaning that only T-cells with low-affinity TCRs for self-antigens remain. Therefore, affinity of TCRs or variants of the present description to peptides can be enhanced by methods well known in the art.

The present description further relates to a method of identifying and isolating a TCR according to the present description, said method comprising incubating PBMCs from HLA-A*024-negative healthy donors with A24/peptide monomers, incubating the PBMCs with tetramer-phycoerythrin (PE) and isolating the high avidity T-cells by fluorescence activated cell sorting (FACS)–Calibur analysis.

The present description further relates to a method of identifying and isolating a TCR according to the present description, said method comprising obtaining a transgenic mouse with the entire human TCR $\alpha\beta$ gene loci (1.1 and 0.7 Mb), whose T-cells express a diverse human TCR repertoire that compensates for mouse TCR deficiency, immunizing the mouse with a peptide, incubating PBMCs obtained from the transgenic

mice with tetramer-phycoerythrin (PE), and isolating the high avidity T-cells by fluorescence activated cell sorting (FACS)—Calibur analysis.

In one aspect, in order to obtain T-cells expressing TCRs of the present description, nucleic acids encoding TCR-alpha and/or TCR-beta chains of the present description are cloned into expression vectors, such as gamma retrovirus or lentivirus. The recombinant viruses are generated and then tested for functionality, such as antigen specificity and functional avidity. An aliquot of the final product is then used to transduce the target T-cell population (generally purified from patient PBMCs), which is expanded before infusion into the patient.

In another aspect, to obtain T-cells expressing TCRs of the present description, TCR RNAs are synthesized by techniques known in the art, e.g., in vitro transcription systems. The in vitro-synthesized TCR RNAs are then introduced into primary CD8+ T-cells obtained from healthy donors by electroporation to re-express tumor specific TCR-alpha and/or TCR-beta chains.

To increase the expression, nucleic acids encoding TCRs of the present description may be operably linked to strong promoters, such as retroviral long terminal repeats (LTRs), cytomegalovirus (CMV), murine stem cell virus (MSCV) U3, phosphoglycerate kinase (PGK), β -actin, ubiquitin, and a simian virus 40 (SV40)/CD43 composite promoter, elongation factor (EF)-1a and the spleen focus-forming virus (SFFV) promoter. In a preferred embodiment, the promoter is heterologous to the nucleic acid being expressed.

In addition to strong promoters, TCR expression cassettes of the present description may contain additional elements that can enhance transgene expression, including a central polypurine tract (cPPT), which promotes the nuclear translocation of lentiviral constructs (Follenzi et al., 2000), and the woodchuck hepatitis virus posttranscriptional regulatory element (wPRE), which increases the level of transgene expression by increasing RNA stability (Zufferey et al., 1999).

The alpha and beta chains of a TCR of the present invention may be encoded by nucleic acids located in separate vectors, or may be encoded by polynucleotides located in the same vector.

Achieving high-level TCR surface expression requires that both the TCR-alpha and TCR-beta chains of the introduced TCR be transcribed at high levels. To do so, the TCR-alpha and TCR-beta chains of the present description may be cloned into bicistronic constructs in a single vector, which has been shown to be capable of overcoming this obstacle. The use of a viral intraribosomal entry site (IRES) between the TCR-alpha and TCR-beta chains results in the coordinated expression of both chains, because the TCR-alpha and TCR-beta chains are generated from a single transcript that is broken into two proteins during translation, ensuring that an equal molar ratio of TCR-alpha and TCR-beta chains are produced (Schmitt et al., 2009).

Nucleic acids encoding TCRs of the present description may be codon optimized to increase expression from a host cell. Redundancy in the genetic code allows some amino acids to be encoded by more than one codon, but certain codons are less "optimal" than others because of the relative availability of matching tRNAs as well as other factors (Gustafsson et al., 2004). Modifying the TCR-alpha and TCR-beta gene sequences such that each amino acid is encoded by the optimal codon for mammalian gene expression, as well as eliminating mRNA instability motifs or cryptic splice sites, has been shown to significantly enhance TCR-alpha and TCR-beta gene expression (Scholten et al., 2006).

Furthermore, mispairing between the introduced and endogenous TCR chains may result in the acquisition of specificities that pose a significant risk for autoimmunity. For example, the formation of mixed TCR dimers may reduce the number of CD3 molecules available to form properly paired TCR complexes, and therefore can significantly decrease the functional avidity of the cells expressing the introduced TCR (Kuball et al., 2007).

To reduce mispairing, the C-terminus domain of the introduced TCR chains of the present description may be modified in order to promote interchain affinity, while decreasing the ability of the introduced chains to pair with the endogenous TCR. These strategies may include replacing the human TCR-alpha and TCR-beta C-terminus domains with their murine counterparts (murinized C-terminus domain); generating a second interchain disulfide bond in the C-terminus domain by introducing a second cysteine residue into both the TCR-alpha and TCR-beta chains of the introduced TCR (cysteine modification); swapping interacting residues in the TCR-alpha and TCR-beta chain C-terminus domains ("knob-in-hole"); and fusing the variable domains of the TCR-alpha and TCR-beta chains directly to CD3 ζ (CD3 ζ fusion) (Schmitt et al., 2009).

In an embodiment, a host cell is engineered to express a TCR of the present description. In preferred embodiments, the host cell is a human T-cell or T-cell progenitor. In some embodiments the T-cell or T-cell progenitor is obtained from a cancer patient. In other embodiments the T-cell or T-cell progenitor is obtained from a healthy donor. Host cells of the present description can be allogeneic or autologous with respect to a patient to be treated. In one embodiment, the host is a gamma/delta T-cell transformed to express an alpha/beta TCR.

A "pharmaceutical composition" is a composition suitable for administration to a human being in a medical setting. Preferably, a pharmaceutical composition is sterile and produced according to GMP guidelines.

The pharmaceutical compositions comprise the peptides either in the free form or in the form of a pharmaceutically acceptable salt (see also above). As used herein, "a pharmaceutically acceptable salt" refers to a derivative of the disclosed peptides wherein the peptide is modified by making acid or base salts of the agent. For example, acid salts are prepared from the free base (typically wherein the neutral form of the drug has a neutral -NH₂ group) involving reaction with a suitable acid. Suitable acids for preparing acid salts include both organic acids, e.g., acetic acid, propionic acid, glycolic

- 61 -

acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methane sulfonic acid, ethane sulfonic acid, p-toluene sulfonic acid, salicylic acid, and the like, as well as inorganic acids, e.g., hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid phosphoric acid and the like. Conversely, preparation of basic salts of acid moieties which may be present on a peptide are prepared using a pharmaceutically acceptable base such as sodium hydroxide, potassium hydroxide, ammonium hydroxide, calcium hydroxide, trimethylamine or the like.

In an especially preferred embodiment, the pharmaceutical compositions comprise the peptides as salts of acetic acid (acetates), trifluoro acetates or hydrochloric acid (chlorides).

Preferably, the medicament of the present invention is an immunotherapeutic such as a vaccine. It may be administered directly into the patient, into the affected organ or systemically i.d., i.m., s.c., i.p. and i.v., or applied *ex vivo* to cells derived from the patient or a human cell line which are subsequently administered to the patient or used *in vitro* to select a subpopulation of immune cells derived from the patient, which are then re-administered to the patient. If the nucleic acid is administered to cells *in vitro*, it may be useful for the cells to be transfected so as to co-express immune-stimulating cytokines, such as interleukin-2. The peptide may be substantially pure or combined with an immune-stimulating adjuvant (see below) or used in combination with immune-stimulatory cytokines, or be administered with a suitable delivery system, for example liposomes. The peptide may also be conjugated to a suitable carrier such as keyhole limpet haemocyanin (KLH) or mannan (see WO 95/18145 and (Longenecker et al., 1993)). The peptide may also be tagged, may be a fusion protein, or may be a hybrid molecule. The peptides whose sequence is given in the present invention are expected to stimulate CD4 or CD8 T cells. However, stimulation of CD8 T cells is more efficient in the presence of help provided by CD4 T-helper cells. Thus, for MHC Class I epitopes that stimulate CD8 T cells the fusion partner or sections of a hybrid molecule suitably

- 62 -

provide epitopes which stimulate CD4-positive T cells. CD4- and CD8-stimulating epitopes are well known in the art and include those identified in the present invention.

In one aspect, the vaccine comprises at least one peptide having the amino acid sequence set forth SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464, and at least one additional peptide, preferably two to 50, more preferably two to 25, even more preferably two to 20 and most preferably two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen or eighteen peptides. The peptide(s) may be derived from one or more specific TAAs and may bind to MHC class I molecules.

A further aspect of the invention provides a nucleic acid (for example a polynucleotide) encoding a peptide or peptide variant of the invention. The polynucleotide may be, for example, DNA, cDNA, PNA, RNA or combinations thereof, either single- and/or double-stranded, or native or stabilized forms of polynucleotides, such as, for example, polynucleotides with a phosphorothioate backbone and it may or may not contain introns so long as it codes for the peptide. Of course, only peptides that contain naturally occurring amino acid residues joined by naturally occurring peptide bonds are encodable by a polynucleotide. A still further aspect of the invention provides an expression vector capable of expressing a polypeptide according to the invention.

A variety of methods have been developed to link polynucleotides, especially DNA, to vectors for example via complementary cohesive termini. For instance, complementary homopolymer tracts can be added to the DNA segment to be inserted to the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA molecules.

Synthetic linkers containing one or more restriction sites provide an alternative method of joining the DNA segment to vectors. Synthetic linkers containing a variety of restriction endonuclease sites are commercially available from a number of sources including International Biotechnologies Inc. New Haven, CN, USA.

- 63 -

A desirable method of modifying the DNA encoding the polypeptide of the invention employs the polymerase chain reaction as disclosed by Saiki RK, et al. (Saiki et al., 1988). This method may be used for introducing the DNA into a suitable vector, for example by engineering in suitable restriction sites, or it may be used to modify the DNA in other useful ways as is known in the art. If viral vectors are used, pox- or adenovirus vectors are preferred.

The DNA (or in the case of retroviral vectors, RNA) may then be expressed in a suitable host to produce a polypeptide comprising the peptide or variant of the invention. Thus, the DNA encoding the peptide or variant of the invention may be used in accordance with known techniques, appropriately modified in view of the teachings contained herein, to construct an expression vector, which is then used to transform an appropriate host cell for the expression and production of the polypeptide of the invention. Such techniques include those disclosed, for example, in US 4,440,859, 4,530,901, 4,582,800, 4,677,063, 4,678,751, 4,704,362, 4,710,463, 4,757,006, 4,766,075, and 4,810,648.

The DNA (or in the case of retroviral vectors, RNA) encoding the polypeptide constituting the compound of the invention may be joined to a wide variety of other DNA sequences for introduction into an appropriate host. The companion DNA will depend upon the nature of the host, the manner of the introduction of the DNA into the host, and whether episomal maintenance or integration is desired.

Generally, the DNA is inserted into an expression vector, such as a plasmid, in proper orientation and correct reading frame for expression. If necessary, the DNA may be linked to the appropriate transcriptional and translational regulatory control nucleotide sequences recognized by the desired host, although such controls are generally available in the expression vector. The vector is then introduced into the host through standard techniques. Generally, not all of the hosts will be transformed by the vector. Therefore, it will be necessary to select for transformed host cells. One selection

- 64 -

technique involves incorporating into the expression vector a DNA sequence, with any necessary control elements, that codes for a selectable trait in the transformed cell, such as antibiotic resistance.

Alternatively, the gene for such selectable trait can be on another vector, which is used to co-transform the desired host cell.

Host cells that have been transformed by the recombinant DNA of the invention are then cultured for a sufficient time and under appropriate conditions known to those skilled in the art in view of the teachings disclosed herein to permit the expression of the polypeptide, which can then be recovered.

Many expression systems are known, including bacteria (for example *E. coli* and *Bacillus subtilis*), yeasts (for example *Saccharomyces cerevisiae*), filamentous fungi (for example *Aspergillus spec.*), plant cells, animal cells and insect cells. Preferably, the system can be mammalian cells such as CHO cells available from the ATCC Cell Biology Collection.

A typical mammalian cell vector plasmid for constitutive expression comprises the CMV or SV40 promoter with a suitable poly A tail and a resistance marker, such as neomycin. One example is pSVL available from Pharmacia, Piscataway, NJ, USA. An example of an inducible mammalian expression vector is pMSG, also available from Pharmacia. Useful yeast plasmid vectors are pRS403-406 and pRS413-416 and are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Plasmids pRS403, pRS404, pRS405 and pRS406 are Yeast Integrating plasmids (YIps) and incorporate the yeast selectable markers HIS3, TRP1, LEU2 and URA3. Plasmids pRS413-416 are Yeast Centromere plasmids (Ycps). CMV promoter-based vectors (for example from Sigma-Aldrich) provide transient or stable expression, cytoplasmic expression or secretion, and N-terminal or C-terminal tagging in various combinations of FLAG, 3xFLAG, c-myc or MAT. These fusion proteins allow for detection, purification and analysis of recombinant protein. Dual-tagged fusions provide flexibility in detection.

The strong human cytomegalovirus (CMV) promoter regulatory region drives constitutive protein expression levels as high as 1 mg/L in COS cells. For less potent cell lines, protein levels are typically ~0.1 mg/L. The presence of the SV40 replication origin will result in high levels of DNA replication in SV40 replication permissive COS cells. CMV vectors, for example, can contain the pMB1 (derivative of pBR322) origin for replication in bacterial cells, the b-lactamase gene for ampicillin resistance selection in bacteria, hGH polyA, and the f1 origin. Vectors containing the pre-pro-trypsin leader (PPT) sequence can direct the secretion of FLAG fusion proteins into the culture medium for purification using ANTI-FLAG antibodies, resins, and plates. Other vectors and expression systems are well known in the art for use with a variety of host cells.

In another embodiment two or more peptides or peptide variants of the invention are encoded and thus expressed in a successive order (similar to “beads on a string” constructs). In doing so, the peptides or peptide variants may be linked or fused together by stretches of linker amino acids, such as for example LLLLLL, or may be linked without any additional peptide(s) between them. These constructs can also be used for cancer therapy and may induce immune responses both involving MHC I and MHC II.

The present invention also relates to a host cell transformed with a polynucleotide vector construct of the present invention. The host cell can be either prokaryotic or eukaryotic. Bacterial cells may be preferred prokaryotic host cells in some circumstances and typically are a strain of *E. coli* such as, for example, the *E. coli* strains DH5 available from Bethesda Research Laboratories Inc., Bethesda, MD, USA, and RR1 available from the American Type Culture Collection (ATCC) of Rockville, MD, USA (No ATCC 31343). Preferred eukaryotic host cells include yeast, insect and mammalian cells, preferably vertebrate cells such as those from a mouse, rat, monkey or human fibroblastic and colon cell lines. Yeast host cells include YPH499, YPH500 and YPH501, which are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Preferred mammalian host cells include Chinese hamster ovary (CHO)

- 66 -

cells available from the ATCC as CCL61, NIH Swiss mouse embryo cells NIH/3T3 available from the ATCC as CRL 1658, monkey kidney-derived COS-1 cells available from the ATCC as CRL 1650 and 293 cells which are human embryonic kidney cells. Preferred insect cells are Sf9 cells which can be transfected with baculovirus expression vectors. An overview regarding the choice of suitable host cells for expression can be found in, for example, the textbook of Paulina Balbás and Argelia Lorence "Methods in Molecular Biology Recombinant Gene Expression, Reviews and Protocols," Part One, Second Edition, ISBN 978-1-58829-262-9, and other literature known to the person of skill.

Transformation of appropriate cell hosts with a DNA construct of the present invention is accomplished by well-known methods that typically depend on the type of vector used. With regard to transformation of prokaryotic host cells, see, for example, Cohen et al. (Cohen et al., 1972) and (Green and Sambrook, 2012) . Transformation of yeast cells is described in Sherman et al. (Sherman et al., 1986) . The method of Beggs (Beggs, 1978) is also useful. Regarding vertebrate cells, reagents useful in transfecting such cells, for example calcium phosphate and DEAE-dextran or liposome formulations, are available from Stratagene Cloning Systems, or Life Technologies Inc., Gaithersburg, MD 20877, USA. Electroporation is also useful for transforming and/or transfecting cells and is well known in the art for transforming yeast cell, bacterial cells, insect cells and vertebrate cells.

Successfully transformed cells, i.e. cells that contain a DNA construct of the present invention, can be identified by well-known techniques such as PCR. Alternatively, the presence of the protein in the supernatant can be detected using antibodies.

It will be appreciated that certain host cells of the invention are useful in the preparation of the peptides of the invention, for example bacterial, yeast and insect cells. However, other host cells may be useful in certain therapeutic methods. For example, antigen-presenting cells, such as dendritic cells, may usefully be used to express the peptides of the invention such that they may be loaded into appropriate MHC molecules. Thus, the

current invention provides a host cell comprising a nucleic acid or an expression vector according to the invention.

In a preferred embodiment the host cell is an antigen presenting cell, in particular a dendritic cell or antigen presenting cell. APCs loaded with a recombinant fusion protein containing prostatic acid phosphatase (PAP) were approved by the U.S. Food and Drug Administration (FDA) on April 29, 2010, to treat asymptomatic or minimally symptomatic metastatic HRPC (Sipuleucel-T) (Rini et al., 2006; Small et al., 2006).

A further aspect of the invention provides a method of producing a peptide or its variant, the method comprising culturing a host cell and isolating the peptide from the host cell or its culture medium.

In another embodiment, the peptide, the nucleic acid or the expression vector of the invention are used in medicine. For example, the peptide or its variant may be prepared for intravenous (i.v.) injection, sub-cutaneous (s.c.) injection, intradermal (i.d.) injection, intraperitoneal (i.p.) injection, intramuscular (i.m.) injection. Preferred methods of peptide injection include s.c., i.d., i.p., i.m., and i.v. Preferred methods of DNA injection include i.d., i.m., s.c., i.p. and i.v. Doses of e.g. between 50 µg and 1.5 mg, preferably 125 µg to 500 µg, of peptide or DNA may be given and will depend on the respective peptide or DNA. Dosages of this range were successfully used in previous trials (Walter et al., 2012).

The polynucleotide used for active vaccination may be substantially pure or contained in a suitable vector or delivery system. The nucleic acid may be DNA, cDNA, PNA, RNA or a combination thereof. Methods for designing and introducing such a nucleic acid are well known in the art. An overview is provided by e.g. Teufel et al. (Teufel et al., 2005). Polynucleotide vaccines are easy to prepare, but the mode of action of these vectors in inducing an immune response is not fully understood. Suitable vectors and delivery systems include viral DNA and/or RNA, such as systems based on adenovirus, vaccinia virus, retroviruses, herpes virus, adeno-associated virus or hybrids containing elements

- 68 -

of more than one virus. Non-viral delivery systems include cationic lipids and cationic polymers and are well known in the art of DNA delivery. Physical delivery, such as via a "gene-gun" may also be used. The peptide or peptides encoded by the nucleic acid may be a fusion protein, for example with an epitope that stimulates T cells for the respective opposite CDR as noted above.

The medicament of the invention may also include one or more adjuvants. Adjuvants are substances that non-specifically enhance or potentiate the immune response (e.g., immune responses mediated by CD8-positive T cells and helper-T (TH) cells to an antigen and would thus be considered useful in the medicament of the present invention. Suitable adjuvants include, but are not limited to, 1018 ISS, aluminum salts, AMPLIVAX®, AS15, BCG, CP-870,893, CpG7909, CyaA, dSLIM, flagellin or TLR5 ligands derived from flagellin, FLT3 ligand, GM-CSF, IC30, IC31, Imiquimod (ALDARA®), resiquimod, ImuFact IMP321, Interleukins as IL-2, IL-13, IL-21, Interferon-alpha or -beta, or pegylated derivatives thereof, IS Patch, ISS, ISCOMATRIX, ISCOMs, JuvImmune®, LipoVac, MALP2, MF59, monophosphoryl lipid A, Montanide IMS 1312, Montanide ISA 206, Montanide ISA 50V, Montanide ISA-51, water-in-oil and oil-in-water emulsions, OK-432, OM-174, OM-197-MP-EC, ONTAK, OspA, PepTel® vector system, poly(lactid co-glycolid) [PLG]-based and dextran microparticles, talactoferrin SRL172, Virosomes and other Virus-like particles, YF-17D, VEGF trap, R848, beta-glucan, Pam3Cys, Aquila's QS21 stimulon, which is derived from saponin, mycobacterial extracts and synthetic bacterial cell wall mimics, and other proprietary adjuvants such as Ribi's Detox, Quil, or Superfos. Adjuvants such as Freund's or GM-CSF are preferred. Several immunological adjuvants (e.g., MF59) specific for dendritic cells and their preparation have been described previously (Allison and Krummel, 1995). Also cytokines may be used. Several cytokines have been directly linked to influencing dendritic cell migration to lymphoid tissues (e.g., TNF-), accelerating the maturation of dendritic cells into efficient antigen-presenting cells for T-lymphocytes (e.g., GM-CSF, IL-1 and IL-4) (U.S. Pat. No. 5,849,589, specifically incorporated herein by reference in its entirety) and acting as immunoadjuvants (e.g., IL-12, IL-15, IL-23, IL-7, IFN-alpha, IFN-beta) (Gabrilovich et al., 1996).

CpG immunostimulatory oligonucleotides have also been reported to enhance the effects of adjuvants in a vaccine setting. Without being bound by theory, CpG oligonucleotides act by activating the innate (non-adaptive) immune system via Toll-like receptors (TLR), mainly TLR9. CpG triggered TLR9 activation enhances antigen-specific humoral and cellular responses to a wide variety of antigens, including peptide or protein antigens, live or killed viruses, dendritic cell vaccines, autologous cellular vaccines and polysaccharide conjugates in both prophylactic and therapeutic vaccines. More importantly it enhances dendritic cell maturation and differentiation, resulting in enhanced activation of TH1 cells and strong cytotoxic T-lymphocyte (CTL) generation, even in the absence of CD4 T cell help. The TH1 bias induced by TLR9 stimulation is maintained even in the presence of vaccine adjuvants such as alum or incomplete Freund's adjuvant (IFA) that normally promote a TH2 bias. CpG oligonucleotides show even greater adjuvant activity when formulated or co-administered with other adjuvants or in formulations such as microparticles, nanoparticles, lipid emulsions or similar formulations, which are especially necessary for inducing a strong response when the antigen is relatively weak. They also accelerate the immune response and enable the antigen doses to be reduced by approximately two orders of magnitude, with comparable antibody responses to the full-dose vaccine without CpG in some experiments (Krieg, 2006). US 6,406,705 B1 describes the combined use of CpG oligonucleotides, non-nucleic acid adjuvants and an antigen to induce an antigen-specific immune response. A CpG TLR9 antagonist is dSLIM (double Stem Loop Immunomodulator) by Mologen (Berlin, Germany) which is a preferred component of the pharmaceutical composition of the present invention. Other TLR binding molecules such as RNA binding TLR 7, TLR 8 and/or TLR 9 may also be used.

Other examples for useful adjuvants include, but are not limited to chemically modified CpGs (e.g. CpR, Idera), dsRNA analogues such as Poly(I:C) and derivatives thereof (e.g. AmpliGen®, Hiltonol®, poly-(ICLC), poly(IC-R), poly(I:C12U), non-CpG bacterial DNA or RNA as well as immunoactive small molecules and antibodies such as cyclophosphamide, sunitinib, Bevacizumab®, celebrex, NCX-4016, sildenafil, tadalafil,

- 70 -

varafenil, sorafenib, temozolomide, temsirolimus, XL-999, CP-547632, pazopanib, VEGF Trap, ZD2171, AZD2171, anti-CTLA4, other antibodies targeting key structures of the immune system (e.g. anti-CD40, anti-TGFbeta, anti-TNFalpha receptor) and SC58175, which may act therapeutically and/or as an adjuvant. The amounts and concentrations of adjuvants and additives useful in the context of the present invention can readily be determined by the skilled artisan without undue experimentation.

Preferred adjuvants are anti-CD40, imiquimod, resiquimod, GM-CSF, cyclophosphamide, sunitinib, bevacizumab, interferon-alpha, CpG oligonucleotides and derivates, poly-(I:C) and derivates, RNA, sildenafil, and particulate formulations with PLG or virosomes.

In a preferred embodiment, the pharmaceutical composition according to the invention the adjuvant is selected from the group consisting of colony-stimulating factors, such as Granulocyte Macrophage Colony Stimulating Factor (GM-CSF, sargramostim), cyclophosphamide, imiquimod, resiquimod, and interferon-alpha.

In a preferred embodiment, the pharmaceutical composition according to the invention the adjuvant is selected from the group consisting of colony-stimulating factors, such as Granulocyte Macrophage Colony Stimulating Factor (GM-CSF, sargramostim), cyclophosphamide, imiquimod and resiquimod. In a preferred embodiment of the pharmaceutical composition according to the invention, the adjuvant is cyclophosphamide, imiquimod or resiquimod. Even more preferred adjuvants are Montanide IMS 1312, Montanide ISA 206, Montanide ISA 50V, Montanide ISA-51, poly-ICLC (Hiltonol®) and anti-CD40 mAB, or combinations thereof.

This composition is used for parenteral administration, such as subcutaneous, intradermal, intramuscular or oral administration. For this, the peptides and optionally other molecules are dissolved or suspended in a pharmaceutically acceptable, preferably aqueous carrier. In addition, the composition can contain excipients, such as buffers, binding agents, blasting agents, diluents, flavors, lubricants, etc. The peptides

- 71 -

can also be administered together with immune stimulating substances, such as cytokines. An extensive listing of excipients that can be used in such a composition, can be, for example, taken from A. Kibbe, Handbook of Pharmaceutical Excipients (Kibbe, 2000). The composition can be used for a prevention, prophylaxis and/or therapy of adenomatous or cancerous diseases. Exemplary formulations can be found in, for example, EP2112253.

It is important to realize that the immune response triggered by the vaccine according to the invention attacks the cancer in different cell-stages and different stages of development. Furthermore different cancer associated signaling pathways are attacked. This is an advantage over vaccines that address only one or few targets, which may cause the tumor to easily adapt to the attack (tumor escape). Furthermore, not all individual tumors express the same pattern of antigens. Therefore, a combination of several tumor-associated peptides ensures that every single tumor bears at least some of the targets. The composition is designed in such a way that each tumor is expected to express several of the antigens and cover several independent pathways necessary for tumor growth and maintenance. Thus, the vaccine can easily be used "off-the-shelf" for a larger patient population. This means that a pre-selection of patients to be treated with the vaccine can be restricted to HLA typing, does not require any additional biomarker assessments for antigen expression, but it is still ensured that several targets are simultaneously attacked by the induced immune response, which is important for efficacy (Banchereau et al., 2001; Walter et al., 2012).

As used herein, the term "scaffold" refers to a molecule that specifically binds to an (e.g. antigenic) determinant. In one embodiment, a scaffold is able to direct the entity to which it is attached (e.g. a (second) antigen binding moiety) to a target site, for example to a specific type of tumor cell or tumor stroma bearing the antigenic determinant (e.g. the complex of a peptide with MHC, according to the application at hand). In another embodiment a scaffold is able to activate signaling through its target antigen, for example a T cell receptor complex antigen. Scaffolds include but are not limited to antibodies and fragments thereof, antigen binding domains of an antibody, comprising an antibody heavy chain variable region and an antibody light chain variable region,

- 72 -

binding proteins comprising at least one ankyrin repeat motif and single domain antigen binding (SDAB) molecules, aptamers, (soluble) TCRs and (modified) cells such as allogenic or autologous T cells. To assess whether a molecule is a scaffold binding to a target, binding assays can be performed.

“Specific” binding means that the scaffold binds the peptide-MHC-complex of interest better than other naturally occurring peptide-MHC-complexes, to an extent that a scaffold armed with an active molecule that is able to kill a cell bearing the specific target is not able to kill another cell without the specific target but presenting other peptide-MHC complex(es). Binding to other peptide-MHC complexes is irrelevant if the peptide of the cross-reactive peptide-MHC is not naturally occurring, i.e. not derived from the human HLA-peptidome. Tests to assess target cell killing are well known in the art. They should be performed using target cells (primary cells or cell lines) with unaltered peptide-MHC presentation, or cells loaded with peptides such that naturally occurring peptide-MHC levels are reached.

Each scaffold can comprise a labelling which provides that the bound scaffold can be detected by determining the presence or absence of a signal provided by the label. For example, the scaffold can be labelled with a fluorescent dye or any other applicable cellular marker molecule. Such marker molecules are well known in the art. For example a fluorescence-labelling, for example provided by a fluorescence dye, can provide a visualization of the bound aptamer by fluorescence or laser scanning microscopy or flow cytometry.

Each scaffold can be conjugated with a second active molecule such as for example IL-21, anti-CD3, and anti-CD28.

For further information on polypeptide scaffolds see for example the background section of WO 2014/071978A1 and the references cited therein.

- 73 -

The present invention further relates to aptamers. Aptamers (see for example WO 2014/191359 and the literature as cited therein) are short single-stranded nucleic acid molecules, which can fold into defined three-dimensional structures and recognize specific target structures. They have appeared to be suitable alternatives for developing targeted therapies. Aptamers have been shown to selectively bind to a variety of complex targets with high affinity and specificity.

Aptamers recognizing cell surface located molecules have been identified within the past decade and provide means for developing diagnostic and therapeutic approaches. Since aptamers have been shown to possess almost no toxicity and immunogenicity they are promising candidates for biomedical applications. Indeed aptamers, for example prostate-specific membrane-antigen recognizing aptamers, have been successfully employed for targeted therapies and shown to be functional in xenograft *in vivo* models. Furthermore, aptamers recognizing specific tumor cell lines have been identified.

DNA aptamers can be selected to reveal broad-spectrum recognition properties for various cancer cells, and particularly those derived from solid tumors, while non-tumorigenic and primary healthy cells are not recognized. If the identified aptamers recognize not only a specific tumor sub-type but rather interact with a series of tumors, this renders the aptamers applicable as so-called broad-spectrum diagnostics and therapeutics.

Further, investigation of cell-binding behavior with flow cytometry showed that the aptamers revealed very good apparent affinities that are within the nanomolar range.

Aptamers are useful for diagnostic and therapeutic purposes. Further, it could be shown that some of the aptamers are taken up by tumor cells and thus can function as molecular vehicles for the targeted delivery of anti-cancer agents such as siRNA into tumor cells.

- 74 -

Aptamers can be selected against complex targets such as cells and tissues and complexes of the peptides comprising, preferably consisting of, a sequence according to any one of SEQ ID NO 1 to SEQ ID NO 387 and SEQ ID No. 463 to SEQ ID No. 464, according to the invention at hand with the MHC molecule, using the cell-SELEX (Systematic Evolution of Ligands by Exponential enrichment) technique.

The peptides of the present invention can be used to generate and develop specific antibodies against MHC/peptide complexes. These can be used for therapy, targeting toxins or radioactive substances to the diseased tissue. Another use of these antibodies can be targeting radionuclides to the diseased tissue for imaging purposes such as PET. This use can help to detect small metastases or to determine the size and precise localization of diseased tissues.

Therefore, it is a further aspect of the invention to provide a method for producing a recombinant antibody specifically binding to a human major histocompatibility complex (MHC) class I or II being complexed with a HLA-restricted antigen (preferably a peptide according to the present invention), the method comprising: immunizing a genetically engineered non-human mammal comprising cells expressing said human major histocompatibility complex (MHC) class I or II with a soluble form of a MHC class I or II molecule being complexed with said HLA-restricted antigen; isolating mRNA molecules from antibody producing cells of said non-human mammal; producing a phage display library displaying protein molecules encoded by said mRNA molecules; and isolating at least one phage from said phage display library, said at least one phage displaying said antibody specifically binding to said human major histocompatibility complex (MHC) class I or II being complexed with said HLA-restricted antigen.

It is thus a further aspect of the invention to provide an antibody that specifically binds to a human major histocompatibility complex (MHC) class I or II being complexed with an HLA-restricted antigen, wherein the antibody preferably is a polyclonal antibody, monoclonal antibody, bi-specific antibody and/or a chimeric antibody.

- 75 -

Respective methods for producing such antibodies and single chain class I major histocompatibility complexes, as well as other tools for the production of these antibodies are disclosed in WO 03/068201, WO 2004/084798, WO 01/72768, WO 03/070752, and in publications (Cohen et al., 2003a; Cohen et al., 2003b; Denkberg et al., 2003), which for the purposes of the present invention are all explicitly incorporated by reference in their entireties.

Preferably, the antibody is binding with a binding affinity of below 20 nanomolar, preferably of below 10 nanomolar, to the complex, which is also regarded as "specific" in the context of the present invention.

The present invention relates to a peptide comprising a sequence that is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464, or a variant thereof which is at least 88% homologous (preferably identical) to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464 or a variant thereof that induces T cells cross-reacting with said peptide, wherein said peptide is not the underlying full-length polypeptide.

The present invention further relates to a peptide comprising a sequence that is selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464 or a variant thereof which is at least 88% homologous (preferably identical) to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464, wherein said peptide or variant has an overall length of between 8 and 100, preferably between 8 and 30, and most preferred between 8 and 14 amino acids.

The present invention further relates to the peptides according to the invention that have the ability to bind to a molecule of the human major histocompatibility complex (MHC) class-I or -II.

- 76 -

The present invention further relates to the peptides according to the invention wherein the peptide consists or consists essentially of an amino acid sequence according to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464.

The present invention further relates to the peptides according to the invention, wherein the peptide is (chemically) modified and/or includes non-peptide bonds.

The present invention further relates to the peptides according to the invention, wherein the peptide is part of a fusion protein, in particular comprising N-terminal amino acids of the HLA-DR antigen-associated invariant chain (Ii), or wherein the peptide is fused to (or into) an antibody, such as, for example, an antibody that is specific for dendritic cells.

The present invention further relates to a nucleic acid, encoding the peptides according to the invention, provided that the peptide is not the complete (full) human protein.

The present invention further relates to the nucleic acid according to the invention that is DNA, cDNA, PNA, RNA or combinations thereof.

The present invention further relates to an expression vector capable of expressing a nucleic acid according to the present invention.

The present invention further relates to a peptide according to the present invention, a nucleic acid according to the present invention or an expression vector according to the present invention for use in medicine, in particular in the treatment of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer,

- 77 -

esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

The present invention further relates to a host cell comprising a nucleic acid according to the invention or an expression vector according to the invention.

The present invention further relates to the host cell according to the present invention that is an antigen presenting cell, and preferably a dendritic cell.

The present invention further relates to a method of producing a peptide according to the present invention, said method comprising culturing the host cell according to the present invention, and isolating the peptide from said host cell or its culture medium.

The present invention further relates to the method according to the present invention, where-in the antigen is loaded onto class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell by contacting a sufficient amount of the antigen with an antigen-presenting cell.

The present invention further relates to the method according to the invention, wherein the antigen-presenting cell comprises an expression vector capable of expressing said peptide containing SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464 or said variant amino acid sequence.

The present invention further relates to activated T cells, produced by the method according to the present invention, wherein said T cells selectively recognizes a cell which aberrantly expresses a polypeptide comprising an amino acid sequence according to the present invention.

The present invention further relates to a method of killing target cells in a patient which target cells aberrantly express a polypeptide comprising any amino acid sequence

- 78 -

according to the present invention, the method comprising administering to the patient an effective number of T cells as according to the present invention.

The present invention further relates to the use of any peptide described, a nucleic acid according to the present invention, an expression vector according to the present invention, a cell according to the present invention, or an activated cytotoxic T lymphocyte according to the present invention as a medicament or in the manufacture of a medicament. The present invention further relates to a use according to the present invention, wherein the medicament is active against cancer.

The present invention further relates to a use according to the invention, wherein the medicament is a vaccine. The present invention further relates to a use according to the invention, wherein the medicament is active against cancer.

The present invention further relates to a use according to the invention, wherein said cancer cells are acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells or other solid or hematological tumor cells such as acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

- 79 -

The present invention further relates to particular marker proteins and biomarkers based on the peptides according to the present invention, herein called "targets" that can be used in the diagnosis and/or prognosis of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer. The present invention also relates to the use of these novel targets for cancer treatment.

The term "antibody" or "antibodies" is used herein in a broad sense and includes both polyclonal and monoclonal antibodies. In addition to intact or "full" immunoglobulin molecules, also included in the term "antibodies" are fragments (e.g. CDRs, Fv, Fab and Fc fragments) or polymers of those immunoglobulin molecules and humanized versions of immunoglobulin molecules, as long as they exhibit any of the desired properties (e.g., specific binding of an acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker (poly)peptide, delivery of a toxin to an acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and

- 80 -

endometrial cancer cell expressing a cancer marker gene at an increased level, and/or inhibiting the activity of an acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker polypeptide) according to the invention.

Whenever possible, the antibodies of the invention may be purchased from commercial sources. The antibodies of the invention may also be generated using well-known methods. The skilled artisan will understand that either full length acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker polypeptides or fragments thereof may be used to generate the antibodies of the invention. A polypeptide to be used for generating an antibody of the invention may be partially or fully purified from a natural source or may be produced using recombinant DNA techniques.

For example, a cDNA encoding a peptide according to the present invention, such as a peptide according to SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464 polypeptide, or a variant or fragment thereof, can be expressed in prokaryotic cells (e.g., bacteria) or eukaryotic cells (e.g., yeast, insect, or mammalian cells), after which the recombinant protein can be purified and used to generate a monoclonal or polyclonal antibody preparation that specifically bind the acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer,

- 81 -

gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer marker polypeptide used to generate the antibody according to the invention.

One of skill in the art will realize that the generation of two or more different sets of monoclonal or polyclonal antibodies maximizes the likelihood of obtaining an antibody with the specificity and affinity required for its intended use (e.g., ELISA, immunohistochemistry, *in vivo* imaging, immunotoxin therapy). The antibodies are tested for their desired activity by known methods, in accordance with the purpose for which the antibodies are to be used (e.g., ELISA, immunohistochemistry, immunotherapy, etc.; for further guidance on the generation and testing of antibodies, see, e.g., Greenfield, 2014 (Greenfield, 2014)). For example, the antibodies may be tested in ELISA assays or, Western blots, immunohistochemical staining of formalin-fixed cancers or frozen tissue sections. After their initial *in vitro* characterization, antibodies intended for therapeutic or *in vivo* diagnostic use are tested according to known clinical testing methods.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a substantially homogeneous population of antibodies, i.e.; the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. The monoclonal antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or

- 82 -

subclass, as well as fragments of such antibodies, so long as they exhibit the desired antagonistic activity (US 4,816,567, which is hereby incorporated in its entirety).

Monoclonal antibodies of the invention may be prepared using hybridoma methods. In a hybridoma method, a mouse or other appropriate host animal is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in US 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies).

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly Fab fragments, can be accomplished using routine techniques known in the art. For instance, digestion can be performed using papain. Examples of papain digestion are described in WO 94/29348 and US 4,342,566. Papain digestion of antibodies typically produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual Fc fragment. Pepsin treatment yields a F(ab')₂ fragment and a pFc' fragment.

The antibody fragments, whether attached to other sequences or not, can also include insertions, deletions, substitutions, or other selected modifications of particular regions or specific amino acids residues, provided the activity of the fragment is not significantly altered or impaired compared to the non-modified antibody or antibody fragment. These modifications can provide for some additional property, such as to remove/add amino acids capable of disulfide bonding, to increase its bio-longevity, to alter its secretory characteristics, etc. In any case, the antibody fragment must possess a bioactive

- 83 -

property, such as binding activity, regulation of binding at the binding domain, etc. Functional or active regions of the antibody may be identified by mutagenesis of a specific region of the protein, followed by expression and testing of the expressed polypeptide. Such methods are readily apparent to a skilled practitioner in the art and can include site-specific mutagenesis of the nucleic acid encoding the antibody fragment.

The antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab' or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed by substituting rodent CDRs or CDR

- 84 -

sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (US 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production can be employed. For example, it has been described that the homozygous deletion of the antibody heavy chain joining region gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. Human antibodies can also be produced in phage display libraries.

Antibodies of the invention are preferably administered to a subject in a pharmaceutically acceptable carrier. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the formulation to render the formulation isotonic. Examples of the pharmaceutically-acceptable carrier include saline, Ringer's solution and dextrose solution. The pH of the solution is preferably from about 5 to about 8, and more preferably from about 7 to about 7.5. Further carriers include sustained release preparations such as semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, liposomes or microparticles. It will be apparent to those persons skilled in the art that certain carriers may be more preferable depending upon, for instance, the route of administration and concentration of antibody being administered.

The antibodies can be administered to the subject, patient, or cell by injection (e.g., intravenous, intraperitoneal, subcutaneous, intramuscular), or by other methods such as infusion that ensure its delivery to the bloodstream in an effective form. The antibodies

- 85 -

may also be administered by intratumoral or peritumoral routes, to exert local as well as systemic therapeutic effects. Local or intravenous injection is preferred.

Effective dosages and schedules for administering the antibodies may be determined empirically, and making such determinations is within the skill in the art. Those skilled in the art will understand that the dosage of antibodies that must be administered will vary depending on, for example, the subject that will receive the antibody, the route of administration, the particular type of antibody used, and other drugs being administered. A typical daily dosage of the antibody used alone might range from about 1 ($\mu\text{g}/\text{kg}$) to up to 100 mg/kg of body weight or more per day, depending on the factors mentioned above. Following administration of an antibody, preferably for treating acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, the efficacy of the therapeutic antibody can be assessed in various ways well known to the skilled practitioner. For instance, the size, number, and/or distribution of cancer in a subject receiving treatment may be monitored using standard tumor imaging techniques. A therapeutically-administered antibody that arrests tumor growth, results in tumor shrinkage, and/or prevents the development of new tumors, compared to the disease course that would occur in the absence of antibody administration, is an efficacious antibody for treatment of cancer.

It is a further aspect of the invention to provide a method for producing a soluble T-cell receptor (sTCR) recognizing a specific peptide-MHC complex. Such soluble T-cell receptors can be generated from specific T-cell clones, and their affinity can be increased by mutagenesis targeting the complementarity-determining regions. For the purpose of T-cell receptor selection, phage display can be used (US 2010/0113300,

- 86 -

(Liddy et al., 2012)). For the purpose of stabilization of T-cell receptors during phage display and in case of practical use as drug, alpha and beta chain can be linked e.g. by non-native disulfide bonds, other covalent bonds (single-chain T-cell receptor), or by dimerization domains (Boulter et al., 2003; Card et al., 2004; Willcox et al., 1999). The T-cell receptor can be linked to toxins, drugs, cytokines (see, for example, US 2013/0115191), and domains recruiting effector cells such as an anti-CD3 domain, etc., in order to execute particular functions on target cells. Moreover, it could be expressed in T cells used for adoptive transfer. Further information can be found in WO 2004/033685A1 and WO 2004/074322A1. A combination of sTCRs is described in WO 2012/056407A1. Further methods for the production are disclosed in WO 2013/057586A1.

In addition, the peptides and/or the TCRs or antibodies or other binding molecules of the present invention can be used to verify a pathologist's diagnosis of a cancer based on a biopsied sample.

The antibodies or TCRs may also be used for *in vivo* diagnostic assays. Generally, the antibody is labeled with a radionucleotide (such as ^{111}In , ^{99}Tc , ^{14}C , ^{131}I , ^3H , ^{32}P or ^{35}S) so that the tumor can be localized using immunoscintigraphy. In one embodiment, antibodies or fragments thereof bind to the extracellular domains of two or more targets of a protein selected from the group consisting of the above-mentioned proteins, and the affinity value (Kd) is less than $1 \times 10\mu\text{M}$.

Antibodies for diagnostic use may be labeled with probes suitable for detection by various imaging methods. Methods for detection of probes include, but are not limited to, fluorescence, light, confocal and electron microscopy; magnetic resonance imaging and spectroscopy; fluoroscopy, computed tomography and positron emission tomography. Suitable probes include, but are not limited to, fluorescein, rhodamine, eosin and other fluorophores, radioisotopes, gold, gadolinium and other lanthanides, paramagnetic iron, fluorine-18 and other positron-emitting radionuclides. Additionally, probes may be bi- or multi-functional and be detectable by more than one of the

- 87 -

methods listed. These antibodies may be directly or indirectly labeled with said probes. Attachment of probes to the antibodies includes covalent attachment of the probe, incorporation of the probe into the antibody, and the covalent attachment of a chelating compound for binding of probe, amongst others well recognized in the art. For immunohistochemistry, the disease tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin. The fixed or embedded section contains the sample are contacted with a labeled primary antibody and secondary antibody, wherein the antibody is used to detect the expression of the proteins *in situ*.

Another aspect of the present invention includes an *in vitro* method for producing activated T cells, the method comprising contacting *in vitro* T cells with antigen loaded human MHC molecules expressed on the surface of a suitable antigen-presenting cell for a period of time sufficient to activate the T cell in an antigen specific manner, wherein the antigen is a peptide according to the invention. Preferably a sufficient amount of the antigen is used with an antigen-presenting cell.

Preferably the mammalian cell lacks or has a reduced level or function of the TAP peptide transporter. Suitable cells that lack the TAP peptide transporter include T2, RMA-S and Drosophila cells. TAP is the transporter associated with antigen processing.

The human peptide loading deficient cell line T2 is available from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, USA under Catalogue No CRL 1992; the Drosophila cell line Schneider line 2 is available from the ATCC under Catalogue No CRL 19863; the mouse RMA-S cell line is described in Ljunggren et al. (Ljunggren and Karre, 1985).

Preferably, before transfection the host cell expresses substantially no MHC class I molecules. It is also preferred that the stimulator cell expresses a molecule important for providing a co-stimulatory signal for T-cells such as any of B7.1, B7.2, ICAM-1 and LFA

3. The nucleic acid sequences of numerous MHC class I molecules and of the co-stimulator molecules are publicly available from the GenBank and EMBL databases.

In case of an MHC class I epitope being used as an antigen; the T cells are CD8-positive T cells.

If an antigen-presenting cell is transfected to express such an epitope, preferably the cell comprises an expression vector capable of expressing a peptide containing SEQ ID NO: 1 to SEQ ID NO: 387 and SEQ ID No. 463 to SEQ ID No. 464, or a variant amino acid sequence thereof.

A number of other methods may be used for generating T cells in vitro. For example, autologous tumor-infiltrating lymphocytes can be used in the generation of CTL. Plebanski et al. (Plebanski et al., 1995) made use of autologous peripheral blood lymphocytes (PLBs) in the preparation of T cells. Furthermore, the production of autologous T cells by pulsing dendritic cells with peptide or polypeptide, or via infection with recombinant virus is possible. Also, B cells can be used in the production of autologous T cells. In addition, macrophages pulsed with peptide or polypeptide, or infected with recombinant virus, may be used in the preparation of autologous T cells. S. Walter et al. (Walter et al., 2003) describe the in vitro priming of T cells by using artificial antigen presenting cells (aAPCs), which is also a suitable way for generating T cells against the peptide of choice. In the present invention, aAPCs were generated by the coupling of preformed MHC:peptide complexes to the surface of polystyrene particles (microbeads) by biotin:streptavidin biochemistry. This system permits the exact control of the MHC density on aAPCs, which allows to selectively elicit high- or low-avidity antigen-specific T cell responses with high efficiency from blood samples. Apart from MHC:peptide complexes, aAPCs should carry other proteins with co-stimulatory activity like anti-CD28 antibodies coupled to their surface. Furthermore such aAPC-based systems often require the addition of appropriate soluble factors, e. g. cytokines, like interleukin-12.

- 89 -

Allogeneic cells may also be used in the preparation of T cells and a method is described in detail in WO 97/26328, incorporated herein by reference. For example, in addition to *Drosophila* cells and T2 cells, other cells may be used to present antigens such as CHO cells, baculovirus-infected insect cells, bacteria, yeast, and vaccinia-infected target cells. In addition plant viruses may be used (see, for example, Porta et al. (Porta et al., 1994) which describes the development of cowpea mosaic virus as a high-yielding system for the presentation of foreign peptides.

The activated T cells that are directed against the peptides of the invention are useful in therapy. Thus, a further aspect of the invention provides activated T cells obtainable by the foregoing methods of the invention.

Activated T cells, which are produced by the above method, will selectively recognize a cell that aberrantly expresses a polypeptide that comprises an amino acid sequence of SEQ ID NO: 1 to SEQ ID NO 387 and SEQ ID No. 463 to SEQ ID No. 464.

Preferably, the T cell recognizes the cell by interacting through its TCR with the HLA/peptide-complex (for example, binding). The T cells are useful in a method of killing target cells in a patient whose target cells aberrantly express a polypeptide comprising an amino acid sequence of the invention wherein the patient is administered an effective number of the activated T cells. The T cells that are administered to the patient may be derived from the patient and activated as described above (i.e. they are autologous T cells). Alternatively, the T cells are not from the patient but are from another individual. Of course, it is preferred if the individual is a healthy individual. By "healthy individual" the inventors mean that the individual is generally in good health, preferably has a competent immune system and, more preferably, is not suffering from any disease that can be readily tested for and detected.

In vivo, the target cells for the CD8-positive T cells according to the present invention can be cells of the tumor (which sometimes express MHC class II) and/or stromal cells

- 90 -

surrounding the tumor (tumor cells) (which sometimes also express MHC class II; (Dengjel et al., 2006)).

The T cells of the present invention may be used as active ingredients of a therapeutic composition. Thus, the invention also provides a method of killing target cells in a patient whose target cells aberrantly express a polypeptide comprising an amino acid sequence of the invention, the method comprising administering to the patient an effective number of T cells as defined above.

By "aberrantly expressed" the inventors also mean that the polypeptide is over-expressed compared to levels of expression in normal tissues or that the gene is silent in the tissue from which the tumor is derived but in the tumor it is expressed. By "over-expressed" the inventors mean that the polypeptide is present at a level at least 1.2-fold of that present in normal tissue; preferably at least 2-fold, and more preferably at least 5-fold or 10-fold the level present in normal tissue.

T cells may be obtained by methods known in the art, e.g. those described above.

Protocols for this so-called adoptive transfer of T cells are well known in the art. Reviews can be found in: Gattioni et al. and Morgan et al. (Gattioni et al., 2006; Morgan et al., 2006).

Another aspect of the present invention includes the use of the peptides complexed with MHC to generate a T-cell receptor whose nucleic acid is cloned and is introduced into a host cell, preferably a T cell. This engineered T cell can then be transferred to a patient for therapy of cancer.

Any molecule of the invention, i.e. the peptide, nucleic acid, antibody, expression vector, cell, activated T cell, T-cell receptor or the nucleic acid encoding it, is useful for the treatment of disorders, characterized by cells escaping an immune response. Therefore any molecule of the present invention may be used as medicament or in the

manufacture of a medicament. The molecule may be used by itself or combined with other molecule(s) of the invention or (a) known molecule(s).

The present invention is further directed at a kit comprising:

- (a) a container containing a pharmaceutical composition as described above, in solution or in lyophilized form;
- (b) optionally a second container containing a diluent or reconstituting solution for the lyophilized formulation; and
- (c) optionally, instructions for (i) use of the solution or (ii) reconstitution and/or use of the lyophilized formulation.

The kit may further comprise one or more of (iii) a buffer, (iv) a diluent, (v) a filter, (vi) a needle, or (v) a syringe. The container is preferably a bottle, a vial, a syringe or test tube; and it may be a multi-use container. The pharmaceutical composition is preferably lyophilized.

Kits of the present invention preferably comprise a lyophilized formulation of the present invention in a suitable container and instructions for its reconstitution and/or use. Suitable containers include, for example, bottles, vials (e.g. dual chamber vials), syringes (such as dual chamber syringes) and test tubes. The container may be formed from a variety of materials such as glass or plastic. Preferably the kit and/or container contain/s instructions on or associated with the container that indicates directions for reconstitution and/or use. For example, the label may indicate that the lyophilized formulation is to be reconstituted to peptide concentrations as described above. The label may further indicate that the formulation is useful or intended for subcutaneous administration.

The container holding the formulation may be a multi-use vial, which allows for repeat administrations (e.g., from 2-6 administrations) of the reconstituted formulation. The kit may further comprise a second container comprising a suitable diluent (e.g., sodium bicarbonate solution).

- 92 -

Upon mixing of the diluent and the lyophilized formulation, the final peptide concentration in the reconstituted formulation is preferably at least 0.15 mg/mL/peptide (=75 µg) and preferably not more than 3 mg/mL/peptide (=1500 µg). The kit may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

Kits of the present invention may have a single container that contains the formulation of the pharmaceutical compositions according to the present invention with or without other components (e.g., other compounds or pharmaceutical compositions of these other compounds) or may have distinct container for each component.

Preferably, kits of the invention include a formulation of the invention packaged for use in combination with the co-administration of a second compound (such as adjuvants (e.g. GM-CSF), a chemotherapeutic agent, a natural product, a hormone or antagonist, an anti-angiogenesis agent or inhibitor, an apoptosis-inducing agent or a chelator) or a pharmaceutical composition thereof. The components of the kit may be pre-complexed, or each component may be in a separate distinct container prior to administration to a patient. The components of the kit may be provided in one or more liquid solutions, preferably, an aqueous solution, more preferably, a sterile aqueous solution. The components of the kit may also be provided as solids, which may be converted into liquids by addition of suitable solvents, which are preferably provided in another distinct container.

The container of a therapeutic kit may be a vial, test tube, flask, bottle, syringe, or any other means of enclosing a solid or liquid. Usually, when there is more than one component, the kit will contain a second vial or other container, which allows for separate dosing. The kit may also contain another container for a pharmaceutically acceptable liquid. Preferably, a therapeutic kit will contain an apparatus (e.g., one or

- 93 -

more needles, syringes, eye droppers, pipette, etc.), which enables administration of the agents of the invention that are components of the present kit.

The present formulation is one that is suitable for administration of the peptides by any acceptable route such as oral (enteral), nasal, ophthalmic, subcutaneous, intradermal, intramuscular, intravenous or transdermal. Preferably, the administration is s.c., and most preferably i.d. administration may be by infusion pump.

Since the peptides of the invention were isolated from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer, the medicament of the invention is preferably used to treat acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer.

The present invention further relates to a method for producing a personalized pharmaceutical for an individual patient comprising manufacturing a pharmaceutical composition comprising at least one peptide selected from a warehouse of pre-screened TUMAPs, wherein the at least one peptide used in the pharmaceutical composition is selected for suitability in the individual patient. In one embodiment, the pharmaceutical composition is a vaccine. The method could also be adapted to produce

- 94 -

T cell clones for down-stream applications, such as TCR isolations, or soluble antibodies, and other treatment options.

A “personalized pharmaceutical” shall mean specifically tailored therapies for one individual patient that will only be used for therapy in such individual patient, including actively personalized cancer vaccines and adoptive cellular therapies using autologous patient tissue.

As used herein, the term “warehouse” shall refer to a group or set of peptides that have been pre-screened for immunogenicity and/or over-presentation in a particular tumor type. The term “warehouse” is not intended to imply that the particular peptides included in the vaccine have been pre-manufactured and stored in a physical facility, although that possibility is contemplated. It is expressly contemplated that the peptides may be manufactured *de novo* for each individualized vaccine produced or may be pre-manufactured and stored. The warehouse (e.g. in the form of a database) is composed of tumor-associated peptides which were highly overexpressed in the tumor tissue of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer patients with various HLA-A HLA-B and HLA-C alleles. It may contain MHC class I and MHC class II peptides or elongated MHC class I peptides. In addition to the tumor associated peptides collected from several acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic

- 95 -

cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer tissues, the warehouse may contain HLA-A*02, HLA-A*01, HLA-A*03, HLA-A*24, HLA-B*07, HLA-B*08 and HLA-B*44 marker peptides. These peptides allow comparison of the magnitude of T-cell immunity induced by TUMAPS in a quantitative manner and hence allow important conclusion to be drawn on the capacity of the vaccine to elicit anti-tumor responses. Secondly, they function as important positive control peptides derived from a “non-self” antigen in the case that any vaccine-induced T-cell responses to TUMAPs derived from “self” antigens in a patient are not observed. And thirdly, it may allow conclusions to be drawn, regarding the status of immunocompetence of the patient.

TUMAPs for the warehouse are identified by using an integrated functional genomics approach combining gene expression analysis, mass spectrometry, and T-cell immunology (XPresident®). The approach assures that only TUMAPs truly present on a high percentage of tumors but not or only minimally expressed on normal tissue, are chosen for further analysis. For initial peptide selection, acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples from patients and blood from healthy donors were analyzed in a stepwise approach:

1. HLA ligands from the malignant material were identified by mass spectrometry
2. Genome-wide messenger ribonucleic acid (mRNA) expression analysis was used to identify genes over-expressed in the malignant tissue acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell

- 96 -

lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer) compared with a range of normal organs and tissues

3. Identified HLA ligands were compared to gene expression data. Peptides over-presented or selectively presented on tumor tissue, preferably encoded by selectively expressed or over-expressed genes as detected in step 2 were considered suitable TUMAP candidates for a multi-peptide vaccine.

4. Literature research was performed in order to identify additional evidence supporting the relevance of the identified peptides as TUMAPs

5. The relevance of over-expression at the mRNA level was confirmed by redetection of selected TUMAPs from step 3 on tumor tissue and lack of (or infrequent) detection on healthy tissues.

6. In order to assess, whether an induction of *in vivo* T-cell responses by the selected peptides may be feasible, *in vitro* immunogenicity assays were performed using human T cells from healthy donors as well as from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer patients.

In an aspect, the peptides are pre-screened for immunogenicity before being included in the warehouse. By way of example, and not limitation, the immunogenicity of the peptides included in the warehouse is determined by a method comprising *in vitro* T-cell priming through repeated stimulations of CD8+ T cells from healthy donors with artificial antigen presenting cells loaded with peptide/MHC complexes and anti-CD28 antibody.

This method is preferred for rare cancers and patients with a rare expression profile. In contrast to multi-peptide cocktails with a fixed composition as currently developed, the

- 97 -

warehouse allows a significantly higher matching of the actual expression of antigens in the tumor with the vaccine. Selected single or combinations of several “off-the-shelf” peptides will be used for each patient in a multitarget approach. In theory an approach based on selection of e.g. 5 different antigenic peptides from a library of 50 would already lead to approximately 17 million possible drug product (DP) compositions.

In an aspect, the peptides are selected for inclusion in the vaccine based on their suitability for the individual patient based on the method according to the present invention as described herein, or as below.

The HLA phenotype, transcriptomic and peptidomic data is gathered from the patient's tumor material, and blood samples to identify the most suitable peptides for each patient containing “warehouse” and patient-unique (i.e. mutated) TUMAPs. Those peptides will be chosen, which are selectively or over-expressed in the patients' tumor and, where possible, show strong *in vitro* immunogenicity if tested with the patients' individual PBMCs.

Preferably, the peptides included in the vaccine are identified by a method comprising: (a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from the individual patient; (b) comparing the peptides identified in (a) with a warehouse (database) of peptides as described above; and (c) selecting at least one peptide from the warehouse (database) that correlates with a tumor-associated peptide identified in the patient. For example, the TUMAPs presented by the tumor sample are identified by: (a1) comparing expression data from the tumor sample to expression data from a sample of normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and (a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor. Preferably, the sequences of MHC ligands are identified by eluting bound peptides from MHC

molecules isolated from the tumor sample and sequencing the eluted ligands. Preferably, the tumor sample and the normal tissue are obtained from the same patient.

In addition to, or as an alternative to, selecting peptides using a warehousing (database) model, TUMAPs may be identified in the patient *de novo*, and then included in the vaccine. As one example, candidate TUMAPs may be identified in the patient by (a1) comparing expression data from the tumor sample to expression data from a sample of normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and (a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor. As another example, proteins may be identified containing mutations that are unique to the tumor sample relative to normal corresponding tissue from the individual patient, and TUMAPs can be identified that specifically target the mutation. For example, the genome of the tumor and of corresponding normal tissue can be sequenced by whole genome sequencing: For discovery of non-synonymous mutations in the protein-coding regions of genes, genomic DNA and RNA are extracted from tumor tissues and normal non-mutated genomic germline DNA is extracted from peripheral blood mononuclear cells (PBMCs). The applied NGS approach is confined to the re-sequencing of protein coding regions (exome re-sequencing). For this purpose, exonic DNA from human samples is captured using vendor-supplied target enrichment kits, followed by sequencing with e.g. a HiSeq2000 (Illumina). Additionally, tumor mRNA is sequenced for direct quantification of gene expression and validation that mutated genes are expressed in the patients' tumors. The resultant millions of sequence reads are processed through software algorithms. The output list contains mutations and gene expression. Tumor-specific somatic mutations are determined by comparison with the PBMC-derived germline variations and prioritized. The *de novo* identified peptides can then be tested for immunogenicity as described above for the warehouse, and candidate TUMAPs possessing suitable immunogenicity are selected for inclusion in the vaccine.

- 99 -

In one exemplary embodiment, the peptides included in the vaccine are identified by: (a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from the individual patient by the method as described above; (b) comparing the peptides identified in a) with a warehouse of peptides that have been prescreened for immunogenicity and overpresentation in tumors as compared to corresponding normal tissue; (c) selecting at least one peptide from the warehouse that correlates with a tumor-associated peptide identified in the patient; and (d) optionally, selecting at least one peptide identified de novo in (a) confirming its immunogenicity.

In one exemplary embodiment, the peptides included in the vaccine are identified by: (a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from the individual patient; and (b) selecting at least one peptide identified de novo in (a) and confirming its immunogenicity.

Once the peptides for a personalized peptide-based vaccine are selected, the vaccine is produced. The vaccine preferably is a liquid formulation consisting of the individual peptides dissolved in between 20-40% DMSO, preferably about 30-35% DMSO, such as about 33% DMSO.

Each peptide to be included into a product is dissolved in DMSO. The concentration of the single peptide solutions has to be chosen depending on the number of peptides to be included into the product. The single peptide-DMSO solutions are mixed in equal parts to achieve a solution containing all peptides to be included in the product with a concentration of ~2.5 mg/ml per peptide. The mixed solution is then diluted 1:3 with water for injection to achieve a concentration of 0.826 mg/ml per peptide in 33% DMSO. The diluted solution is filtered through a 0.22 µm sterile filter. The final bulk solution is obtained.

Final bulk solution is filled into vials and stored at -20°C until use. One vial contains 700 µL solution, containing 0.578 mg of each peptide. Of this, 500 µL (approx. 400 µg per peptide) will be applied for intradermal injection.

- 100 -

In addition to being useful for treating cancer, the peptides of the present invention are also useful as diagnostics. Since the peptides were generated from acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer cells and since it was determined that these peptides are not or at lower levels present in normal tissues, these peptides can be used to diagnose the presence of a cancer.

The presence of claimed peptides on tissue biopsies in blood samples can assist a pathologist in diagnosis of cancer. Detection of certain peptides by means of antibodies, mass spectrometry or other methods known in the art can tell the pathologist that the tissue sample is malignant or inflamed or generally diseased, or can be used as a biomarker for acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer. Presence of groups of peptides can enable classification or sub-classification of diseased tissues.

The detection of peptides on diseased tissue specimen can enable the decision about the benefit of therapies involving the immune system, especially if T-lymphocytes are known or expected to be involved in the mechanism of action. Loss of MHC expression is a well described mechanism by which infected or malignant cells escape immuno-

- 101 -

surveillance. Thus, presence of peptides shows that this mechanism is not exploited by the analyzed cells.

The peptides of the present invention might be used to analyze lymphocyte responses against those peptides such as T cell responses or antibody responses against the peptide or the peptide complexed to MHC molecules. These lymphocyte responses can be used as prognostic markers for decision on further therapy steps. These responses can also be used as surrogate response markers in immunotherapy approaches aiming to induce lymphocyte responses by different means, e.g. vaccination of protein, nucleic acids, autologous materials, adoptive transfer of lymphocytes. In gene therapy settings, lymphocyte responses against peptides can be considered in the assessment of side effects. Monitoring of lymphocyte responses might also be a valuable tool for follow-up examinations of transplantation therapies, e.g. for the detection of graft versus host and host versus graft diseases.

The present invention will now be described in the following examples which describe preferred embodiments thereof, and with reference to the accompanying figures, nevertheless, without being limited thereto. For the purposes of the present invention, all references as cited herein are incorporated by reference in their entireties.

FIGURES

Figures 1A through 1J show the over-presentation of various peptides in different cancer tissues (black dots). Upper part: Median MS signal intensities from technical replicate measurements are plotted as dots for single HLA-A*24 positive normal (grey dots, left part of figure) and tumor samples (black dots, right part of figure) on which the peptide was detected. Boxes display median, 25th and 75th percentile of normalized signal intensities, while whiskers extend to the lowest data point still within 1.5 interquartile range (IQR) of the lower quartile, and the highest data point still within 1.5 IQR of the upper quartile. Normal organs are ordered according to risk categories (blood cells, blood vessels, brain, liver, lung: high risk, grey dots; reproductive organs, breast, prostate: low risk, grey dots; all other organs: medium risk; grey dots). Lower

- 102 -

part: The relative peptide detection frequency in every organ is shown as spine plot. Numbers below the panel indicate number of samples on which the peptide was detected out of the total number of samples analyzed for each organ (N = 75 for normal samples, N = 263 for tumor samples). If the peptide has been detected on a sample but could not be quantified for technical reasons, the sample is included in this representation of detection frequency, but no dot is shown in the upper part of the figure. Tissues (from left to right): Normal samples: blood cells; bloodvess (blood vessels); brain; heart; liver; lung; adrenal gl (adrenal gland); bile duct; gall bl (gallbladder); intest. la (large intestine); intest. sm (small intestine); kidney; nerve perith (peripheral nerve); pancreas; pituit (pituitary); skin; spinal cord; spleen; stomach; thyroid. Tumor samples: AML (acute myeloid leukemia); BRCA (breast cancer); CCC (cholangiocellular carcinoma); CLL (chronic lymphocytic leukemia); CRC (colorectal cancer); GBC (gallbladder cancer); GBM (glioblastoma); GC (gastric cancer); HCC (hepatocellular carcinoma); HNSCC (head and neck squamous cell carcinoma); MEL (melanoma); NHL (non-Hodgkin lymphoma); NSCLCadeno (non-small cell lung cancer adenocarcinoma); NSCLCother (NSCLC samples that could not unambiguously be assigned to NSCLCadeno or NSCLCsquam); NSCLCsquam (squamous cell non-small cell lung cancer); OC (ovarian cancer); OSCAR (esophageal cancer); PACA (pancreatic cancer); PRCA (prostate cancer); RCC (renal cell carcinoma); SCLC (small cell lung cancer); UBC (urinary bladder carcinoma); UEC (uterine and endometrial cancer). Figure 1A) Gene symbol: SLC6A3, Peptide: FMVIAGMPLF (SEQ ID NO.: 15), Figure 1B) Gene symbol: KLHDC7B, Peptide: RYSPVKDAW (SEQ ID NO.: 60), Figure 1C) Gene symbol: CAPN6, Peptide: NYVLVPTMF (SEQ ID NO.: 74), Figure 1D) Gene symbol: SYT12, Peptide: SYLPTAERL (SEQ ID NO.: 86), Figure 1E) Gene symbol: PTPRZ1, Peptide: VYDTMIEKFA (SEQ ID NO.: 202), Figure 1F) Gene symbol: PTPRZ1, Peptide: EYSLPVLTF (SEQ ID NO.: 274), Figure 1G) Gene symbol: LOC100124692, Peptide: NYMDTDNLMF (SEQ ID NO.: 362), Figure 1H) Gene symbol: AR, Peptide: YQSRDYYNF (SEQ ID NO.: 386), Figure 1I) Gene symbol: CT45A4, CT45A5, Peptide: VGGNVTSNF (SEQ ID NO.: 463), and Figure 1J) Gene symbol: CT45A1, CT45A2, CT45A3, CT45A4, CT45A6, LOC101060208, LOC101060210, LOC101060211, Peptide: VGGNVTSNF (SEQ ID NO.: 464).

Figures 2A through 2Q show exemplary expression profile of source genes of the present invention that are over-expressed in different cancer samples. Tumor (black dots) and normal (grey dots) samples are grouped according to organ of origin. Box-and-whisker plots represent median FPKM value, 25th and 75th percentile (box) plus whiskers that extend to the lowest data point still within 1.5 interquartile range (IQR) of the lower quartile and the highest data point still within 1.5 IQR of the upper quartile. Normal organs are ordered according to risk categories. FPKM: fragments per kilobase per million mapped reads. Tissues (from left to right): Normal samples: blood cells; bloodvess (blood vessels); brain; heart; liver; lung; adipose (adipose tissue); adrenal gl (adrenal gland); bile duct; bladder; bone marrow; esoph (esophagus); eye; gall bl (gallbladder); head&neck; intest. la (large intestine); intest. sm (small intestine); kidney; lymph node; nerve perith (peripheral nerve); pancreas; parathyr (parathyroid gland); perit (peritoneum); pituit (pituitary); pleura; skel. mus (skeletal muscle); skin; spleen; stomach; thyroid; trachea; ureter; breast; ovary; placenta; prostate; testis; thymus; uterus. Tumor samples: AML (acute myeloid leukemia); BRCA (breast cancer); CCC (cholangiocellular carcinoma); CLL (chronic lymphocytic leukemia); CRC (colorectal cancer); GBC (gallbladder cancer); GBM (glioblastoma); GC (gastric cancer); HCC (hepatocellular carcinoma); HNSCC (head and neck squamous cell carcinoma); MEL (melanoma); NHL (non-Hodgkin lymphoma); NSCLCadeno (non-small cell lung cancer adenocarcinoma); NSCLCother (NSCLC samples that could not unambiguously be assigned to NSCLCadeno or NSCLCsquam); NSCLCsquam (squamous cell non-small cell lung cancer); OC (ovarian cancer); OSCAR (esophageal cancer); PACA (pancreatic cancer); PRCA (prostate cancer); RCC (renal cell carcinoma); SCLC (small cell lung cancer); UBC (urinary bladder carcinoma); UEC (uterine and endometrial cancer). Figure 2A) Gene symbol: MAGEA4, Peptide: IFPKTGILLI (SEQ ID No.: 1), Figure 2B) Gene symbol: TRPM8, Peptide: KFLTHDVLTELF (SEQ ID No.: 3), Figure 2C) Gene symbol: CHRNA9, Peptide: KYIATMAL (SEQ ID No.: 13), Figure 2D) Gene symbol: MMP12, Peptide: KYVDINTFRL (SEQ ID No.: 18), Figure 2E) Gene symbol: SPINK2, Peptide: LYMRVNTHF (SEQ ID No.: 21), Figure 2F) Gene symbol: OR51E2, Peptide: FWFDSREISF (SEQ ID No.: 28), Figure 2G) Gene symbol: MMP1, Peptide:

- 104 -

KQMQEFFFGL (SEQ ID No.: 31), Figure 2H) Gene symbol: MAGEC1, Peptide: FSSTLVSLF (SEQ ID No.: 35), Figure 2I) Gene symbol: ENPP3, Peptide: KTYLPTFETTI (SEQ ID No.: 39), Figure 2J) Gene symbol: POTEH, Peptide: MVLQPQPQLF (SEQ ID No.: 4), Figure 2K) Gene symbol: OR51E2, Peptide: TQMFFIHAL (SEQ ID No.: 97), Figure 2L) Gene symbol: MMP11, Peptide: FFFKAGFVWR (SEQ ID No.: 107), Figure 2M) Gene symbol: MMP11, Peptide: YFLRGRLYW (SEQ ID No.: 122), Figure 2N) Gene symbol: SLC24A5, Peptide: EYFLPSLEII (SEQ ID No.: 194), Figure 2O) Gene symbol: SLC24A5, Peptide: MSAIWISAF (SEQ ID No.: 277), Figure 2P) Gene symbol: ELP4, EXOSC7, KCNG2, TM4SF19, TOP2A, Peptide: HHTQLIFVF (SEQ ID No.: 286), Figure 2Q) Gene symbol: LAMA3, Peptide: YFGNPQKF (SEQ ID No.: 304).

Figure 3A through 3G show exemplary results of peptide-specific in vitro CD8+ T cell responses of a healthy HLA-A*24+ donor. CD8+ T cells were primed using artificial APCs coated with anti-CD28 mAb and HLA-A*24 in complex with SEQ ID NO: 420 peptide (VYEKNGYIYF, Seq ID NO: 420) (A, left panel), SEQ ID NO: 411 peptide (VYPPYLNLYL, Seq ID NO: 411) (B, left panel), SEQ ID NO: 5 peptide (LQPQPQLFFSF, Seq ID NO: 5) (C, left panel), SEQ ID NO: 77 peptide (LYGFFFKI, Seq ID NO: 77) (D, left panel), SEQ ID NO: 76 peptide (IYIYPPFAHW, Seq ID NO: 76) (E, left panel), SEQ ID NO: 32 peptide (FYPEVELNLF, Seq ID NO: 32) (F, left panel), SEQ ID NO: 23 peptide (VYSSFVFNLF, Seq ID NO: 23) (G, left panel), respectively. After three cycles of stimulation, the detection of peptide-reactive cells was performed by 2D multimer staining with A*24/SEQ ID NO: 420 (A), A*24/SEQ ID NO: 411 (B), A*24/SEQ ID NO: 5 (C), A*24/SEQ ID NO: 77 (D), A*24/SEQ ID NO: 76 (E), A*24/SEQ ID NO: 32 (F), A*24/SEQ ID NO: 23 (G), respectively. Right panels (A) show control staining of cells stimulated with irrelevant A*24/peptide complexes. Viable singlet cells were gated for CD8+ lymphocytes. Boolean gates helped excluding false-positive events detected with multimers specific for different peptides. Frequencies of specific multimer+ cells among CD8+ lymphocytes are indicated.

EXAMPLES

EXAMPLE 1Identification and quantitation of tumor associated peptides presented on the cell surfaceTissue samples

Patients' tumor tissues were obtained from: Asterand Bioscience (Detroit USA & Royston, Herts, UK), BioServe (Beltsville, MD, USA), Conversant Bio (Huntsville, AL, USA), Geneticist Inc. (Glendale, CA, USA), Heidelberg University Hospital (dept. of Neurosurgery, Heidelberg, Germany), Heidelberg University Hospital (General, Visceral and Transplantation Surgery, Heidelberg, Germany), Heidelberg University Hospital (Thoraxklinik, Heidelberg, Germany), Istituto Nazionale Tumori "Pascale" (Molecular Biology and Viral Oncology Unit, Naples, Italy), Kyoto Prefectural University of Medicine (Kyoto, Japan), Leiden University Medical Center (LUMC) (Leiden, Netherlands), Osaka City University (Osaka, Japan), ProteoGenex Inc. (Culver City, CA, USA), Saint Savvas Hospital (Athens, Greece), Tissue Solutions (Glasgow, UK), University Hospital Bonn (dept. of Internal Medicine III, Hematology and Oncology, Bonn, Germany), University Hospital Tübingen (Dept. of General, Visceral and Transplant Surgery, Tübingen, Germany), University Hospital Tübingen (Dept. of Immunology, Tübingen, Germany), University Hospital Tübingen (Dept. of Urology, Tübingen, Germany), University of Geneva (Division of Oncology, Geneva, Switzerland)

Normal tissues were obtained from: Asterand Bioscience (Detroit USA & Royston, Herts, UK), BioServe (Beltsville, MD, USA), Capital BioScience Inc. (Rockville, MD, USA), Centre for Clinical Transfusion Medicine Tuebingen (Tübingen, Germany), Geneticist Inc. (Glendale, CA, USA), Heidelberg University Hospital (Thoraxklinik, Heidelberg, Germany), Kyoto Prefectural University of Medicine (Kyoto, Japan), ProteoGenex Inc. (Culver City, CA, USA), University Hospital Tübingen (Dept. of General, Visceral and Transplant Surgery, Tübingen, Germany), University Hospital Tübingen (Dept. of Urology, Tübingen, Germany)

- 106 -

Written informed consents of all patients had been given before surgery or autopsy. Tissues were shock-frozen immediately after excision and stored until isolation of TUMAPs at -70°C or below.

Isolation of HLA peptides from tissue samples

HLA peptide pools from shock-frozen tissue samples were obtained by immune precipitation from solid tissues according to a slightly modified protocol (Falk et al., 1991; Seeger et al., 1999) using the HLA-A*02-specific antibody BB7.2, the HLA-A, -B, C-specific antibody W6/32, the HLA-DR specific antibody L243 and the HLA DP specific antibody B7/21, CNBr-activated sepharose, acid treatment, and ultrafiltration.

Mass spectrometry analyses

The HLA peptide pools as obtained were separated according to their hydrophobicity by reversed-phase chromatography (nanoAcquity UPLC system, Waters) and the eluting peptides were analyzed in LTQ-velos and fusion hybrid mass spectrometers (ThermoElectron) equipped with an ESI source. Peptide pools were loaded directly onto the analytical fused-silica micro-capillary column (75 μm i.d. x 250 mm) packed with 1.7 μm C18 reversed-phase material (Waters) applying a flow rate of 400 nL per minute. Subsequently, the peptides were separated using a two-step 180 minute-binary gradient from 10% to 33% B at a flow rate of 300 nL per minute. The gradient was composed of Solvent A (0.1% formic acid in water) and solvent B (0.1% formic acid in acetonitrile). A gold coated glass capillary (PicoTip, New Objective) was used for introduction into the nanoESI source. The LTQ-Orbitrap mass spectrometers were operated in the data-dependent mode using a TOP5 strategy. In brief, a scan cycle was initiated with a full scan of high mass accuracy in the orbitrap ($R = 30\,000$), which was followed by MS/MS scans also in the orbitrap ($R = 7500$) on the 5 most abundant precursor ions with dynamic exclusion of previously selected ions. Tandem mass spectra were interpreted by SEQUEST at a fixed false discovery rate ($q \leq 0.05$) and additional manual control. In cases where the identified peptide sequence was uncertain it was additionally validated by comparison of the generated natural peptide fragmentation pattern with the fragmentation pattern of a synthetic sequence-identical reference peptide.

Label-free relative LC-MS quantitation was performed by ion counting i.e. by extraction and analysis of LC-MS features (Mueller et al., 2007). The method assumes that the peptide's LC-MS signal area correlates with its abundance in the sample. Extracted features were further processed by charge state deconvolution and retention time alignment (Mueller et al., 2008; Sturm et al., 2008). Finally, all LC-MS features were cross-referenced with the sequence identification results to combine quantitative data of different samples and tissues to peptide presentation profiles. The quantitative data were normalized in a two-tier fashion according to central tendency to account for variation within technical and biological replicates. Thus each identified peptide can be associated with quantitative data allowing relative quantification between samples and tissues. In addition, all quantitative data acquired for peptide candidates was inspected manually to assure data consistency and to verify the accuracy of the automated analysis. For each peptide a presentation profile was calculated showing the mean sample presentation as well as replicate variations. The profiles juxtapose acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer samples to a baseline of normal tissue samples. Presentation profiles of exemplary over-presented peptides are shown in Figure 1.

Table 8a and Table 8b show the presentation on various cancer entities for selected peptides, and thus the particular relevance of the peptides as mentioned for the diagnosis and/or treatment of the cancers as indicated (e.g. peptide SEQ ID No. 3 for prostate cancer (PRCA), peptide SEQ ID No. 20 for ovarian cancer (OC) and pancreatic cancer (PACA)).

Table 8a: Overview of presentation of selected tumor-associated peptides of the present invention across entities.

AML: acute myeloid leukemia; BRCA: breast cancer; CCC: cholangiocellular carcinoma; CLL: chronic lymphocytic leukemia; CRC: colorectal cancer; GBC: gallbladder cancer; GBM: glioblastoma; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCadeno: non-small cell lung cancer adenocarcinoma; NSCLCother: NSCLC samples that could not unambiguously be assigned to NSCLCadeno or NSCLCsquam; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer; PACA: pancreatic cancer; PRCA: prostate cancer; RCC: renal cell carcinoma; SCLC: small cell lung cancer; UBC: urinary bladder carcinoma; UEC: uterine and endometrial cancer.

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
1	IFPKTGLLII	NSCLCadeno, NSCLCsquam, OSCAR
2	LYAPTILLW	CRC, GC, NSCLCadeno, NSCLCsquam, UEC
3	KFLTHDVLTELF	PRCA
4	MVLQPQPQLF	NSCLCadeno
5	LQPQPQLFFSF	PRCA
6	IVTFMNKTLGTF	NSCLCadeno
7	GYPLRGSSI	GC
9	TYINSLAIL	NSCLCadeno, PRCA, RCC, UBC
10	QYPEFSIEL	PRCA
11	RAMCAMMSF	PACA
12	KYMSRVLFVY	CCC
13	KYYIATMAL	GC
14	YYIATMALI	PRCA
15	FMVIAGMPLF	AML, CLL, GC, HCC, NHL, NSCLCother
16	GYFLAQYLM	RCC
17	IYPEAIATL	BRCA, NSCLCadeno, NSCLCother, RCC
18	KYVDINTFRL	NSCLCsquam
19	ILLCMSLLLF	NSCLCadeno, NSCLCsquam, PRCA
20	ELMAHPFLL	OC, PACA
21	LYMRFVNTHF	AML
22	VYSSFVFNL	NHL
23	VYSSFVFNLF	NHL
24	KMLPEASLLI	NHL
25	MLPEASLLI	NHL

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
26	TYFFVDNQYW	NSCLCsquam, OSCAR
27	LSCTATPLF	NSCLCadenos
28	FWFDSREISF	PRCA
29	IYLLLPPVI	NSCLCadenos, PRCA
30	RQAYSVYAF	PRCA
31	KQMQUEFFGL	GC, NSCLCsquam
32	FYPEVELNF	CCC, UBC
33	FYQPDLKYLSF	NHL
34	LIFALALAAF	UEC
35	FSSTLVSLF	OC
36	VYLASVAAF	PRCA
37	ISFSDTVNVW	RCC
38	RYAHTLVTSVLF	BRCA, MEL
39	KTYLPTFETTI	UEC
40	NYPEGAAYEF	NSCLCadenos, OC
41	IYFATQVWF	PRCA
42	VYDSIWCNM	UEC
43	KYKDHFTFI	BRCA, GBC, NHL, NSCLCadenos, NSCLCsquam
44	FYHEDMPLW	NHL
45	YGQSKPWTF	HCC, HNSCC, NSCLCadenos, NSCLCsquam, OSCAR, RCC
46	IYPDSIQEL	HCC
47	SYLWTDNLQEF	OSCAR
48	AWSPATLFLF	GC, MEL, NSCLCsquam, OC, RCC
49	QYLSIAERAEF	GC, UBC
50	RYFDENIQKF	OSCAR
51	YFDENIQKF	NSCLCsquam, OSCAR
52	SWHKATFLF	AML
53	LFQRVSSVSF	HCC
54	SYQEAIQQL	PRCA
55	AVLRHLETF	CCC
56	FYKLIQNGF	AML
57	RYLQVLLY	GC
58	IYYSHENLI	CCC, CRC, GC, HCC, NSCLCadenos, NSCLCsquam, OC, PRCA
59	VFPLVTPLL	GBM
60	RYSVKDAW	CCC, GBC, NHL, NSCLCadenos, NSCLCother, NSCLCsquam, OC, OSCAR
61	RIFTARLYF	NHL
62	VYIVPVIVL	GBM, NSCLCsquam
63	LYIDKGQYL	GBM, HNSCC, MEL, NSCLCadenos, NSCLCsquam
64	QFSHVPLNNF	NSCLCsquam

- 110 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
65	EYLLMIFKLV	NSCLCadeno, NSCLCsquam, PRCA
66	IYKDYYRYNF	NHL
67	SYVLQIVAI	GBM, MEL
68	VYKEDLPQL	CLL, NSCLCsquam
69	KWFDSHIPRW	GBC
70	RYTGQWSEW	UBC
71	RYLPNP SLNAF	HCC, NSCLCadeno
72	RWLDGSPVTL	NHL
73	YFCSTKGQLF	CLL, NHL
74	NYVLVPTMF	CCC, CRC, GBC, GC, HCC, NSCLCadeno, NSCLCsquam, OC, UEC
75	VYEHNHVSL	UBC
76	IYIY PFAHW	HCC, NHL
77	LYGFFFKI	GC, HCC, UBC
78	TYSKTIALYGF	UBC
79	FYIVTRPLAF	NHL, NSCLCadeno
80	SYATPVDLW	AML, CCC, MEL, NSCLCsquam
81	AYLKLLPMF	BRCA, HNSCC, MEL
82	SYLENSASW	BRCA, HNSCC, NSCLCadeno, OSCAR, SCLC, UEC
83	VLQGEFFLF	NHL
84	YTIERYFTL	GC, NSCLCsquam, UEC
85	KYLSIPTVFF	CCC, GBC, GC, HCC, NSCLCadeno, PRCA, RCC, UBC
86	SYLPTAERL	GBC, GBM, GC, HCC, NSCLCadeno, NSCLCother, NSCLCsquam, PACA, PRCA, UEC
87	NYTRLVLQF	CCC, GBC, GC, NSCLCsquam, OSCAR, UEC
88	TYVPSTFLV	CCC, GBM, GC, NSCLCsquam, OSCAR, PACA
89	TYVPSTFLVVL	GC
90	TDLVQFLLF	RCC
91	KQQVVKFLI	PRCA
92	RALTETIMF	UEC
93	TDWSPPPVEF	NSCLCsquam
94	THSGGTNLF	NSCLCadeno
95	IGLSVVHRF	PRCA
96	SHIGVVLAF	PRCA
97	TQMFFIHAL	PRCA
98	LQIPVSPSF	MEL
99	ASAALTGFTF	OC, PRCA
100	KVWSDVTPLTF	CCC, NSCLCsquam, PACA
101	VYAVSSDRF	GBM
102	VLASAHILQF	NSCLCadeno
103	EMFFSPQVF	CRC, HCC, NSCLCadeno

- 111 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
104	GYGLTRVQPF	NSCLCadenos
105	ITPATALLL	NHL
106	LYAFLGSHF	RCC
107	FFFKAGFVWR	NSCLCsquam, UEC
108	WFFQGAQYW	HCC
109	AQHSLTQLF	GBM
110	VYSNPDLFW	NSCLCadenos
111	IRPDYSFQF	NSCLCadenos
112	LYPDSVFGRLF	OC
113	ALMSAFYTF	NSCLCsquam
114	KALMSAFYTF	CCC
115	IMQGFIRAF	NSCLCadenos
116	TYFFVANKY	GC
117	RSMEHPGKLLF	NSCLCadenos, UEC
118	IFLPFFIVF	GC
119	VWSCEGCKAF	UBC
120	VYAFMNENF	RCC
121	RRYFGEKVAL	PRCA
122	YFLRGRLYW	NSCLCsquam, OC
123	FFLQESPVF	HCC
124	EYNVFPRTL	CRC, GC, NSCLCadenos, NSCLCsquam
125	LYYGSILYI	MEL
126	YSLLDPAQF	CRC, NSCLCadenos, PACA, PRCA, UEC
127	FLPRAYYRW	PRCA
128	AFQNVISF	GC
129	IYVSLAHVL	GC, NSCLCother, NSCLCsquam, PRCA
130	RPEKVFVF	NSCLCadenos
131	MHRTWRETF	PRCA
132	TFEGATVTL	CRC, MEL, NHL
133	FFYVTETTF	NHL
134	IYSSQLPSF	CLL
135	KYKQHFPEI	HCC
136	YLKSVQLF	NSCLCsquam
137	ALFAVCWAPF	NSCLCsquam
138	MMVTVVALF	NSCLCsquam, RCC
139	AYAPRGSYKF	GBC, NSCLCadenos
140	IFQHFCEEI	CLL
141	QYAAAITNGL	NSCLCsquam
142	PYWWNANMVF	NHL
143	KTKRWLWDF	NSCLCadenos, NSCLCsquam
144	LFDHGGTVFF	GBM

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
145	MYTIVTPML	GC
146	NYFLDPVTI	BRCA, MEL, NSCLCadeno, NSCLCsquam, OC
147	FPYPSSILSV	GBM
148	MLPQIPFLLL	OC
149	TQFFIPYTI	CCC, GC, NSCLCsquam, OC
150	FIPVAWLIF	OC
151	RRLWAYVTI	MEL
152	MHPGVLA AFLF	NSCLCadeno
153	AWSPPATLF	MEL
154	DYSKQALSL	NSCLCadeno
155	PYSIYPHGVT F	HCC
156	IYPHGVT FSP	GC
157	SIYPHGVT F	PRCA
158	SYLKDPMIV	CLL
159	VFQPNPLF	UEC
160	YIANLISCF	GBC
161	ILQAPLSVF	CLL, GC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, OSCAR, PACA, RCC, SCLC
162	YYIGIVEEY	HCC, NSCLCsquam
163	YYIGIVEEYW	GC
164	MFQEMLQRL	MEL
165	KDQPQVPCVF	GC
166	MMALWSLLHL	NSCLCsquam
167	LQPPWTTVF	GC, MEL, NSCLCadeno, NSCLCsquam
168	LSSPVHLDF	CRC, HCC, NHL, NSCLCadeno, OSCAR, RCC, UEC
169	MYDLHHLYL	NSCLCsquam
170	IFIPATILL	HCC
171	LYTVPFNLI	NSCLCsquam
172	RYFIAAEKILW	HNSCC
173	RYLSVCERL	NSCLCsquam
174	TYGEEMPEEI	PRCA
175	SYFEYRRL	NSCLCadeno
176	TQAGEYLLF	AML
177	KYLITTFSL	NSCLCadeno
178	AYPQIRCTW	AML
179	MYNMVPPF	NSCLCadeno
180	IYNKTKMAF	SCLC
181	IHGKIFHYF	GC
182	AQGSQTVTF	CLL
183	YQVAKGMEF	NSCLCadeno
184	VYVRPRVF	NSCLCadeno

- 113 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
185	LYICKVELM	GC
186	RRVTWNVLF	NSCLCsquam
187	KWFNVRMGFG F	NHL
188	SLPGSFIYVF	NSCLCadenoc, RCC
189	FYPDEDDFYF	NSCLCother
190	IYIIMQSCW	AML
191	MSYSCGLPSL	GC, HCC
192	CYSFIHLSF	HCC, NHL
193	KYKPVALQCIA	NSCLCadenoc
194	EYFLPSLEII	CLL
195	IYNEHGIQQI	GBM, NSCLCadenoc, NSCLCsquam
196	VGRSPVFLF	OC
197	YYHSGENLY	GC, NSCLCadenoc
198	VLAPVSGQF	CLL, GBC, GC, MEL, NHL, NSCLCadenoc, NSCLCsquam, SCLC
199	MFQFEHIKW	NSCLCadenoc
200	LYMSVEDFI	NSCLCother
201	VFPSVDVSF	CRC, GBM
202	VYDTMIEKFA	GBM, NSCLCadenoc, NSCLCother, NSCLCsquam
203	VYPSESTVM	GBM
204	WQNVTPLETF	NSCLCsquam
205	ISWEVVHTVF	PACA
206	EVVHTVFLF	GBC, MEL, NSCLCadenoc, NSCLCsquam, PRCA, SCLC
207	IYKFIMDRF	AML, GBC, GC, HCC, NSCLCadenoc, NSCLCother, NSCLCsquam, OSCAR, PRCA, UBC
208	QYLQQQAKL	NSCLCadenoc
209	DIYVTGGHLF	PRCA
210	EAYSYPATI	NSCLCadenoc, NSCLCsquam
211	MLYFAPDLIL	GC
212	VYFVQYKIM	UBC
213	FYNRLTKLF	NSCLCadenoc
214	YIPMSVMLF	NSCLCadenoc
215	KASKITFHW	GBM
216	RHYHSIEVF	HCC
217	QRYGFSSVGF	HCC
218	FYFYNCSSL	UEC
219	KVVSGFYI	GC
220	TYATHVTEI	GC
221	VFYCLLFVF	CCC, NSCLCsquam
222	HYHAESFLF	OSCAR

- 114 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
223	KLRALSILF	PRCA
224	AYLQFLSVL	NSCLCother
225	ISMSATEFLL	OSCAR
226	TYSTNRMTI	CCC
227	YLPNPSLNAF	CLL, GC, NSCLCadeno, NSCLCsquam
228	VYLRIGGF	MEL, NSCLCadeno, NSCLCsquam, OSCAR, PRCA, RCC, SCLC, UBC
229	CAMPVAMEF	NHL
230	RWLSKPSLL	NHL
231	KYSVAFYSLD	GC
232	IWPGFTTSI	GC
233	LYSRRGVRTL	OC
234	RYKMLIPF	PRCA
235	VYISDVSVY	NSCLCadeno
236	LHLYCLNTF	CCC
237	RQGLTVLW	NSCLCadeno
238	YTCSRAVSLF	NSCLCadeno
239	IYTFSNVTF	GC
240	RVHANPLLI	HCC
241	QKYYITGEAEGF	OSCAR
242	SYTPLLSYI	MEL
243	ALFPMGPLTF	CLL
244	TYIDTRTVFL	HCC
245	VLPLHFLPF	RCC
246	KIYTTVLFANI	HCC
247	VHSYLGSPF	RCC
248	CWGPHCFEM	NSCLCsquam
249	HQYGGAYNRV	OC
250	VYSDRQIYLL	HCC
251	DYLLSWLLF	GC
252	RYLIIKYPF	GBC
253	QYYCLLLIF	NSCLCadeno, NSCLCsquam
254	KQHAWLPLTI	NHL
255	VYLDEKQHAW	CLL, NHL
256	QHAWLPLTI	NHL
257	MLILFFSTI	MEL
258	VCWNPFNNTF	CRC
259	FFLFIPFF	NSCLCadeno
260	FLFIPFFIF	GC
261	IMFCLKNFWW	NSCLCadeno, OC
263	AYVTEFVSL	BRCA, GBC

- 115 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
264	AYAIPSASLSW	HCC
265	LYQQSDTWSL	HCC, NSCLCadeno
266	TQIITFESF	MEL
267	QHMLPFWTDL	OSCAR
268	YQFGWSPNF	GC
269	FSFSTSMNEF	PRCA
270	GTGKLFWVF	PACA
271	INGDLVFSF	CLL
272	IYFNHRCF	NSCLCadeno
273	VTMYLPLLL	GC, NSCLCadeno, NSCLCsquam, OC, RCC
274	EYSLPVLTF	AML, CCC, CLL, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, OC, OSCAR, RCC, SCLC, UBC, UEC
275	PEYSLPVLTF	NHL
277	MSAIWISAF	OSCAR
278	TYESVVTGFF	SCLC
279	KYKNPYGF	HCC
280	TIYSLEMKMSF	NSCLCother, NSCLCsquam
281	MDQNQVWTF	NSCLCadeno
282	ASYQQSTSSFF	NSCLCadeno
283	SYIVDGKII	PRCA
284	QFYSTLPNTI	HCC
285	YFLPGPHYF	RCC
286	HHTQLIFVF	GBC, PRCA
287	LVQPQAVLF	GBM, GC, NSCLCadeno, SCLC, UEC
288	MGKGSISFLF	NSCLCadeno
289	RTLNEIYHW	NSCLCadeno, NSCLCsquam
290	VTPKMLISF	CCC, NSCLCother
291	YTRLVLQF	AML, BRCA, CCC, CLL, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NHL, NSCLCadeno, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
292	KMFPKDFRF	CRC
293	MYAYAGWFY	CRC
294	KMGRIVDYF	CCC, GBC, GC
295	KYNRQSM TL	HCC
296	YQRPDLLLF	CLL, MEL
297	LKSPRLFTF	GBC
298	TYETVM TFF	GBC, GC, HCC, NHL, UBC
299	FLPALYSLL	NSCLCadeno
300	LFALPDFIF	NSCLCadeno

- 116 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
301	RTALSSTDTF	NHL
302	YQGSLEVLV	NSCLCadenos
303	RFLDRGWGF	CRC
304	YFGNPQKF	GC, PACA
305	RNAFSIYIL	CRC, NSCLCadenos
306	RYILEPFFI	BRCA, CCC, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NHL, NSCLCadenos, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
307	RILTEFELL	GC
308	AAFISVPLLI	HCC
309	AFISVPLLI	NSCLCadenos, NSCLCother
310	EFINGWYVL	HCC
311	IQNAILHLF	SCLC
312	YLCMLYALF	CCC, GBC, GC, NHL
313	IFMENAFEL	HCC
315	VYDYIPLLL	OC
316	IWAERIMF	NSCLCsquam
317	DWIWRILFLV	GBM
318	VQADAKLLF	GC, NSCLCadenos, UBC
319	ATATLHLIF	UEC
320	EVYQKIILKF	GBC
321	VYTVGHNLI	HCC
322	SFISPRYSWLF	AML, CRC, NSCLCadenos, NSCLCsquam, UBC
323	NYSPVTGKF	OSCAR
324	RYFVSNIYL	GC, NSCLCadenos, NSCLCsquam, UEC
325	IFMGAVPTL	HNSCC, NSCLCadenos, OC, OSCAR
326	VHMKDFFYF	CLL, MEL, NHL, NSCLCadenos, NSCLCsquam, OC, PACA
328	IYLVGGYSW	CLL
329	YLGKNWSF	AML, CCC
330	DYIQMIPEL	BRCA, GBC, GC, SCLC
332	VYCSLDKSQF	BRCA, CCC, GBC, GC, MEL
333	RYADLLIYTY	UEC
334	KVFGSFLTL	UEC
335	RYQSVIYPF	UEC
336	VYSDLHAFY	CCC, GBC, HCC, NSCLCadenos, PACA
337	SHSDHEFLF	UEC
338	VYLTWLPGL	CLL, NHL
339	KQVIGIHTF	CLL, GBC, GBM, GC, RCC
340	FPPTPPLF	AML
341	RYENVSILF	GBM, OC
342	MYGIVIRTI	GC

- 117 -

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
343	EYQQYHPSL	NHL
344	YAYATVLTf	GBC, GC, NHL, NSCLCaden, UBC
345	RYLEEHTeF	GBC, NSCLCaden, OC, UEC
346	TYIDFVPYI	GBC, GC, NSCLCaden, OC, RCC, UEC
347	AWLIVLLFL	NHL, UEC
348	RSWENIPVTF	CCC
349	IYMTTGvLL	GC, NSCLCaden
350	VYKWTEEKf	BRCA, NSCLCother, OSCAR
351	GYFGTASLf	NSCLCaden, NSCLCother
352	NAFEAPLTf	OC
353	AAFPGAfSF	OC
354	QYIPTFHvY	BRCA, GBM, HCC, HNSCC, NSCLCsquam, OSCAR
355	VYNNSSRF	CCC, CRC, GBC, GC, HCC, HNSCC, MEL, NHL, NSCLCaden, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UEC
356	YSLEHLTQf	OC, UEC
357	RALLPSPLf	MEL, NHL
358	IYANVTEMLL	OSCAR
359	TQLPAPLRI	UEC
360	LYITKvTTI	NSCLCaden
361	KQPANFIVL	GBC
362	NYMDTDNLmf	CCC, GBC, GC, HCC, OC, OSCAR, SCLC
363	QYGFNYNKF	GC, NSCLCsquam
364	KQSQVVfVL	NSCLCsquam, OC, PRCA
365	KDLMKAYLf	GC, SCLC
366	RLGEFvLLF	GC
367	HWSHITHLf	CCC, CRC, GBM, GC, MEL
368	AYFVAMHLf	GBC, GBM, HNSCC, MEL, NHL, NSCLCaden, NSCLCother, NSCLCsquam, OC, SCLC, UEC
369	NFYLFPTTf	GC, NSCLCsquam
370	TQMDVKLVf	GBC, GC, HCC, NHL, NSCLCaden, NSCLCsquam, OSCAR, UBC, UEC
371	FRSWAVQTF	AML, CRC, GC
372	LYHNWRHAF	BRCA, OC, RCC
373	IWDALERTf	CLL, CRC, GBC, GBM, GC, HCC, HNSCC, MEL, NHL, NSCLCaden, NSCLCsquam, OC, OSCAR, PRCA
374	MIFAVVvLF	GBC, GBM, GC, MEL, NHL, NSCLCaden, OC, OSCAR, PRCA
375	YYAADQWvF	AML, GBC, GC, HCC, NHL, NSCLCaden, NSCLCother, NSCLCsquam, OSCAR, PACA
376	KYVGEVfNI	AML, CLL, HCC, NHL, NSCLCaden, NSCLCsquam, RCC
377	SLWREVvTF	OC, RCC

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
378	VYAVISNIL	GBM, NHL, NSCLCadenoc, SCLC
379	KLPTEWNVL	CCC, CLL, CRC, GC, HCC, NHL, NSCLCadenoc, NSCLCother, OSCAR, RCC, UEC
380	FYIRRLPMF	NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, RCC
381	IYTDITYSF	AML, CCC, GBC, HCC, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, RCC
382	SYPKELMKF	GBC, GC, HNSCC, NSCLCadenoc, OC, UEC
383	PYFSPSASF	NSCLCadenoc, NSCLCother, UBC
385	GYFGNPQKF	BRCA, CCC, CRC, GBC, GC, HCC, HNSCC, MEL, NSCLCadenoc, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, PRCA, RCC, SCLC, UBC, UEC
386	YQSRDYYNF	CRC, GBC, GBM, GC, HCC, HNSCC, NSCLCadenoc, NSCLCsquam, OC, PRCA, UBC, UEC
387	THAGVRLYF	CRC, GBC, GC, HCC, NHL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PRCA, UEC

Table 8b: Overview of presentation of selected tumor-associated peptides of the present invention across entities.

AML: acute myeloid leukemia; BRCA: breast cancer; GBC: gallbladder cancer; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; NSCLCadenoc: non-small cell lung cancer adenocarcinoma; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer;.

SEQ ID No.	Sequence	Peptide Presentation on cancer entities
463	VGGNVTSNF	AML, BRCA, GBC, GC, HCC, HNSCC, NSCLCsquam, OC
464	VGGNVTSNF	BRCA, GBC, GC, HNSCC, NSCLCadenoc, NSCLCsquam, OC

EXAMPLE 2

Expression profiling of genes encoding the peptides of the invention

Over-presentation or specific presentation of a peptide on tumor cells compared to normal cells is sufficient for its usefulness in immunotherapy, and some peptides are tumor-specific despite their source protein occurring also in normal tissues. Still, mRNA expression profiling adds an additional level of safety in selection of peptide targets for immunotherapies. Especially for therapeutic options with high safety risks, such as

- 119 -

affinity-matured TCRs, the ideal target peptide will be derived from a protein that is unique to the tumor and not found on normal tissues.

RNA sources and preparation

Surgically removed tissue specimens were provided as indicated above (see Example 1) after written informed consent had been obtained from each patient. Tumor tissue specimens were snap-frozen immediately after surgery and later homogenized with mortar and pestle under liquid nitrogen. Total RNA was prepared from these samples using TRI Reagent (Ambion, Darmstadt, Germany) followed by a cleanup with RNeasy (QIAGEN, Hilden, Germany); both methods were performed according to the manufacturer's protocol.

Total RNA from healthy human tissues for RNASeq experiments was obtained from: Asterand (Detroit, MI, USA & Royston, Herts, UK); Bio-Options Inc. (Brea, CA, USA); Geneticist Inc. (Glendale, CA, USA); ProteoGenex Inc. (Culver City, CA, USA); Tissue Solutions Ltd (Glasgow, UK). Total RNA from tumor tissues for RNASeq experiments was obtained from: Asterand (Detroit, MI, USA & Royston, Herts, UK); BioCat GmbH (Heidelberg, Germany); BioServe (Beltsville, MD, USA); Geneticist Inc. (Glendale, CA, USA); Istituto Nazionale Tumori "Pascale" (Naples, Italy); ProteoGenex Inc. (Culver City, CA, USA); University Hospital Heidelberg (Heidelberg, Germany).

Quality and quantity of all RNA samples were assessed on an Agilent 2100 Bioanalyzer (Agilent, Waldbronn, Germany) using the RNA 6000 Pico LabChip Kit (Agilent).

RNAseq experiments

Gene expression analysis of - tumor and normal tissue RNA samples was performed by next generation sequencing (RNAseq) by CeGaT (Tübingen, Germany). Briefly, sequencing libraries are prepared using the Illumina HiSeq v4 reagent kit according to the provider's protocol (Illumina Inc., San Diego, CA, USA), which includes RNA fragmentation, cDNA conversion and addition of sequencing adaptors. Libraries derived from multiple samples are mixed equimolar and sequenced on the Illumina HiSeq 2500

- 120 -

sequencer according to the manufacturer's instructions, generating 50 bp single end reads. Processed reads are mapped to the human genome (GRCh38) using the STAR software. Expression data are provided on transcript level as RPKM (Reads Per Kilobase per Million mapped reads, generated by the software Cufflinks) and on exon level (total reads, generated by the software Bedtools), based on annotations of the ensembl sequence database (Ensembl77). Exon reads are normalized for exon length and alignment size to obtain RPKM values.

Exemplary expression profiles of source genes of the present invention that are highly over-expressed or exclusively expressed in acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer are shown in Figure 2. Expression scores for further exemplary genes are shown in Table 9a and Table 9b.

Table 9a: Expression scores. The table lists for each peptide the tumor types for which the exon is very highly over-expressed in tumors compared to a panel of normal tissues (+++), highly over-expressed in tumors compared to a panel of normal tissues (++) or over-expressed in tumors compared to a panel of normal tissues (+). The baseline for this score was calculated from measurements of the following relevant normal tissues: blood cells, blood vessels, brain, heart, liver, lung, adipose tissue, adrenal gland, bile duct, bone marrow, esophagus, eye, gallbladder, head-and-neck, kidney, large intestine, lymph node, pancreas, parathyroid gland, peripheral nerve, peritoneum, pituitary, pleura, skeletal muscle, skin, small intestine, spleen, stomach, thyroid gland, trachea, ureter, urinary bladder. In case expression data for several samples of the same tissue type were available, the arithmetic mean of all respective samples was used for the calculation. AML: acute myeloid leukemia; BRCA: breast cancer; CCC:

cholangiocellular carcinoma; CLL: chronic lymphocytic leukemia; CRC: colorectal cancer; GBC: gallbladder cancer; GBM: glioblastoma; GC: gastric cancer; HCC: hepatocellular carcinoma; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCadeno: non-small cell lung cancer adenocarcinoma; NSCLCother: NSCLC samples that could not unambiguously be assigned to NSCLCadeno or NSCLCsquam; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer; PACA: pancreatic cancer; PRCA: prostate cancer; RCC: renal cell carcinoma; SCLC: small cell lung cancer; UBC: urinary bladder carcinoma; UEC: uterine and endometrial cancer.

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
1	IFPKTGLLII	BRCA, CRC, GC, UEC	HCC, NSCLCadeno, UBC	GBC, HNSCC, MEL, NSCLCsquam, OC, OSCAR, SCLC
2	LYAPTILLW	GBC		CCC, HCC
3	KFLTHDVLTELF	MEL	GBM, SCLC	PRCA
4	MVLQPQPQLF	GBC, HCC, NHL, OC, UBC		PRCA
5	LQPQPQLFFSF	GBC, HCC, NHL, OC, UBC		PRCA
6	IVTFMNKTLGTF	NHL		CLL
7	GYPLRGSSI		OC	UEC
8	IMKPLDQDF	BRCA, CRC, GBC, GC, NSCLCsquam, PACA, SCLC, UBC, UEC	NSCLCadeno	PRCA
9	TYINSLAIL			PRCA
10	QYPEFSIEL			PRCA
11	RAMCAMMSF			PRCA
12	KYMSRVLFVY	MEL, NSCLCadeno, NSCLCsquam, OC, OSCAR, UBC, UEC	GBM, HNSCC, SCLC	NSCLCother
13	KYYIATMAL	MEL, NSCLCadeno, NSCLCsquam, OC, OSCAR, UBC, UEC	GBM, HNSCC, SCLC	NSCLCother
14	YYIATMALI	MEL, NSCLCadeno, NSCLCsquam, OC, OSCAR, UBC, UEC	GBM, HNSCC, SCLC	NSCLCother
15	FMVIAGMPLF	NSCLCadeno,		RCC

- 122 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		NSCLCsquam		
16	GYFLAQYLM	GBM, MEL		PRCA
17	IYPEAIATL	NSCLCadenoc		RCC
18	KYVDINTFRL	HCC, RCC	CCC, GC, MEL, NHL, NSCLCadenoc, OC, PACA, UEC	CRC, GBC, HNSCC, NSCLCother, NSCLCsquam, OSCAR, SCLC, UBC
19	ILLCMSLLLF	OC		BRCA
20	ELMAHPFLL	OC		BRCA
21	LYMRFVNTHF			AML
22	VYSSFVFNL	GBC, GBM, MEL, OC, UBC	CLL	NHL
23	VYSSFVFNLF	GBC, GBM, MEL, OC, UBC	CLL	NHL
24	KMLPEASLLI	GBC, GBM, MEL, OC, UBC	CLL	NHL
25	MLPEASLLI	GBC, GBM, MEL, OC, UBC	CLL	NHL
26	TYFFVDNQYW	HCC, NHL, RCC	CCC, CRC, GBC, GC, HNSCC, MEL, NSCLCadenoc, OC, OSCAR, PACA, SCLC, UBC, UEC	NSCLCother, NSCLCsquam
27	LSCTATPLF	NSCLCother, NSCLCsquam, SCLC	OSCAR	HNSCC
28	FWFDSREISF			PRCA
29	IYLLPPVI			PRCA
30	RQAYSVYAF			PRCA
31	KQMQUEFFGL	CRC, MEL, PACA	GC, NSCLCadenoc, NSCLCsquam, OSCAR, UBC	HNSCC
32	FYPEVELNF	CRC, GC, MEL, PACA	HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, UBC	
33	FYQPDLKYLSF	CLL, GBC	NHL	
34	LIFALALAAF	GBC	GC, HNSCC, NSCLCsquam, OSCAR, PACA	
35	FSSTLVSLF	GBC, SCLC	HCC, MEL	
36	VYLASVAAF		PRCA	
37	ISFSDTVNVW	BRCA, PRCA	MEL	
38	RYAHTLVTSVLF		MEL	

- 123 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
39	KTYLPTFETTI	UEC	RCC	
40	NYPEGAAAYEF		BRCA, OC, UEC	
41	IYFATQVVF		PRCA	
42	VYDSIWCM		OC, UEC	
43	KYKDHFEI	BRCA, MEL, NSCLCadeno, NSCLCsquam, SCLC	GBC, HCC, NHL	
44	FYHEDMPLW	NHL	CLL	
45	YGQSKPWTF		MEL	
46	IYPDSIQEL		HCC	
47	SYLWTDNLQEF	GBC, SCLC	HNSCC, NSCLCsquam, OSCAR	
48	AWSPATLFLF	CCC	MEL	
49	QYLSIAERAEF		UEC	
50	RYFDENIQKF	NSCLCother, OSCAR	HNSCC	
51	YFDENIQKF	NSCLCother, OSCAR	HNSCC	
52	SWHKATFLF		AML	
53	LFQRVSSVSF		MEL	
54	SYQEAIQQL		PRCA	
55	AVLRHLETF	GBM, GC, OSCAR	AML	
56	FYKLIQNGF		AML	
57	RYLQVLLY	GBM, PACA	CRC, GBC, GC	
58	IYYSHENLI	CCC	HCC	
59	VFPLVTPLL		GBM	
60	RYSVKDAW	BRCA, NSCLCsquam, OSCAR, UBC	HNSCC	
61	RIFTARLYF		NHL	
62	VYIVPVIVL	GBM, OC	CCC	
63	LYIDKGQYL		MEL	
64	QFSHVPLNNF		MEL	
65	EYLLMIFKLV	NHL	HNSCC, MEL	
66	IYKDYYRYNF	MEL	NHL	
67	SYVLQIVAI		GBM	
68	VYKEDLPQL		NHL	
69	KWFDSHIPRW	BRCA, CLL, NHL, RCC	UBC	
70	RYTGQWSEW	AML	UBC	

- 124 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
71	RYLPNPSTLNAF		NSCLCother	
72	RWLDGSPVTL		CLL, NHL	
73	YFCSTKGQLF	NHL	CLL	
74	NYVLVPTMF	GC	UEC	
75	VYEHNHVSL		UBC	
76	IYIYPFAHW	GBC	HCC	
77	LYGFFFKI		UBC	
78	TYSKTIALYGF		UBC	
79	FYIVTRPLAF	AML	CCC	
80	SYATPVDLW	GBM, GC	AML	
81	AYLKLLPMF		MEL	
82	SYLENSASW		UEC	
83	VLQGEFFLF	NHL	CLL	
84	YTIERYFTL	GBC, GC, UEC	PACA	
85	KYLSIPTVFF		HCC	
86	SYLPTAERL		CCC	
87	NYTRLVLQF	GBC, GC, UEC	PACA	
88	TYVPSTFLV	GBC, GC, UEC	PACA	
89	TYVPSTFLVVL	GBC, GC, UEC	PACA	
90	TDLVQFLLF	HCC, NSCLCadenoc	GBC, GC, HNSCC, NSCLCsquam, OC, OSCAR, SCLC, UBC	MEL
91	KQQVVKFLI	GC, HNSCC, NHL, NSCLCsquam, OSCAR, PACA, SCLC, UBC, UEC	GBC, OC	BRCA, PRCA
92	RALTETIMF		OC	UEC
93	TDWSPPPVEF	GC, NSCLCsquam, PACA, SCLC, UEC	GBC, HCC, HNSCC, OSCAR	MEL
94	THSGGTNLF	HCC, RCC	CCC, CRC, GC, MEL, NHL, NSCLCadenoc, OC, OSCAR, PACA, SCLC, UBC, UEC	GBC, HNSCC, NSCLCother, NSCLCsquam
95	IGLSVVHRF			PRCA
96	SHIGVVLAF			PRCA
97	TQMFFIHAL			PRCA
98	LQIPVSPSF	GBC, SCLC	HCC, MEL	
99	ASAALTGFTF		PRCA	
100	KVWSDVTPLTF	HCC, RCC, SCLC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadenoc,	

- 125 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
			NSCLCsquam, OC, OSCAR, PACA, UBC, UEC	
101	VYAVSSDRF		GBM, SCLC	
102	VLASAHILQF		UBC	
103	EMFFSPQVF	GC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, RCC, SCLC	BRCA, CCC, CRC, GBC, HCC, OC, UBC	
104	GYGLTRVQPF	GC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, RCC, SCLC	BRCA, CCC, CRC, GBC, HCC, OC, UBC	
105	ITPATALLL	NHL	CLL	
106	LYAFLGSHF	BRCA, NSCLCadenoc, UEC	RCC	
107	FFFKAGFVWR	HCC, OC, SCLC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, UBC, UEC	
108	WFFQGAQYW	HCC, OC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, UBC, UEC	
109	AQHSLTQLF		GBM, SCLC	
110	VYSNPDLFW		CLL	
111	IRPDYSFQF		CLL	
112	LYPDSVFGRLF	AML, CCC, GC, HNSCC, NSCLCother, NSCLCsquam, OSCAR, UEC	BRCA, GBC, HCC, MEL, NSCLCadenoc, OC, SCLC, UBC	
113	ALMSAFYTF	HCC, OC, RCC, SCLC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, PACA, UBC, UEC	
114	KALMSAFYTF	HCC, OC, RCC, SCLC	BRCA, CCC, CRC, GBC, GC, HNSCC,	

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
			NSCLCadeno, NSCLCsquam, OSCAR, PACA, UBC, UEC	
115	IMQGFIRAF	CCC, HCC, OC	NSCLCadeno	
116	TYFFVANKY	CRC, GC, NSCLCadeno, PACA, UBC	HNSCC, NSCLCsquam, OSCAR	
117	RSMEHPGKLLF		BRCA, OC, UEC	
118	IFLPFFIVF	BRCA, GBC, HCC, OSCAR, PRCA, UBC, UEC	CCC, RCC	
119	VWSCEGCKAF		BRCA, OC, UEC	
120	VYAFMNENF	NSCLCsquam, OSCAR	RCC	
121	RRYFGEKVAL		PRCA	
122	YFLRGRLYW	HCC, OC	BRCA, CCC, CRC, GBC, GC, HNSCC, NSCLCadeno, NSCLCsquam, OSCAR, PACA, UBC, UEC	
123	FFLQESPVF	HCC, MEL, PRCA	BRCA	
124	EYNVFPRTL	NSCLCadeno, UBC	HNSCC, NSCLCsquam, OSCAR	
125	LYYGSILYI		MEL	
126	YSLLDPAQF	GBC, HCC, HNSCC, NSCLCadeno, NSCLCothers, OSCAR, SCLC, UBC, UEC	CRC, GC, PRCA	
127	FLPRAYRW		PRCA	
128	AFQNVISF	NSCLCsquam, UEC	GC, PACA	
129	IYVSLAHVL		PRCA	
130	RPEKVFVF	CCC, CRC, GBM, NSCLCothers, NSCLCsquam, OC, SCLC	BRCA, GBC, GC, HNSCC, MEL, NSCLCadeno, OSCAR, PACA	
131	MHRTWRETF		PRCA	
132	TFEGATVTL	NHL	CLL	
133	FFYVTETTF	AML, GC, OC, OSCAR, SCLC,	HCC, NHL	

- 127 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		UEC		
134	IYSSQLPSF	NHL	CLL	
135	KYKQHFPEI		HCC	
136	YLKSVQLF	BRCA, MEL	AML	
137	ALFAVCWAPF	NSCLCsquam, OSCAR	RCC	
138	MMVTVVALF	NSCLCsquam, OSCAR	RCC	
139	AYAPRGSIIYKF	BRCA, NSCLCadenoc, OC	HCC	
140	IFQHFCEEI	AML, BRCA, GBC, HCC, MEL, NSCLCadenoc	OC, SCLC	
141	QYAAAITNGL	SCLC	GBM	
142	PYWWNANMVF		HCC	
143	KTKRWLWDF	CCC, CRC, GBM, GC, NSCLCother, NSCLCsquam, OC	BRCA, GBC, HNSCC, MEL, NSCLCadenoc, OSCAR, PACA	
144	LFDHGGTVFF		PRCA	
145	MYTIVTPML	GBM	OC	
146	NYFLDPVTI		MEL	
147	FPYPSSILSV	MEL	GBM, SCLC	
148	MLPQIPFLLL	CRC, HNSCC, NSCLCsquam, OC, UBC, UEC	BRCA, CCC, GBC, GC, NSCLCadenoc, OSCAR, PACA	
149	TQFFIPYTI	CRC, HNSCC, NSCLCsquam, OC, UBC, UEC	BRCA, CCC, GBC, GC, NSCLCadenoc, OSCAR, PACA	
150	FIPVAWLIF	HNSCC, UBC	MEL	
151	RRLWAYVTI		MEL	
152	MHPGVLAAFLF	NSCLCadenoc, UBC	HNSCC, NSCLCsquam, OSCAR	
153	AWSPATLF	CCC	MEL	
154	DYSKQALSL	CCC, GC, NSCLCadenoc, NSCLCother, PACA, UBC, UEC	HNSCC, NSCLCsquam, OSCAR	
155	PYSIYPHGVTF	CCC	HCC	
156	IYPHGVTFSP	CCC	HCC	
157	SIYPHGVTF	CCC	HCC	
158	SYLKDPMIV	GBC, HNSCC, MEL,	HCC	

- 128 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		NSCLCadenoc, SCLC		
159	VFQPNPLF	GC, NSCLCsquam, PACA	CRC, HNSCC, MEL, OC, OSCAR, UBC	
160	YIANLISCF		SCLC, UEC	
161	ILQAPLSVF	NHL	CLL	
162	YYIGIVEEY	NSCLCother, OSCAR	HNSCC	
163	YYIGIVEEYW	NSCLCother, OSCAR	HNSCC	
164	MFQEMLQRL	BRCA, GBC, NSCLCsquam, OC, UBC	HNSCC, MEL, OSCAR, PACA	
165	KDQPQVPCVF		BRCA	
166	MMALWSLLHL		SCLC	
167	LQPPWTTVF	NHL	CLL	
168	LSSPVHLDF	NHL	CLL	
169	MYDLHHLYL	CRC, GBC, HNSCC, NHL, NSCLCsquam, UEC	BRCA, GC, MEL, NSCLCadenoc, OC, OSCAR, PACA	
170	IFIPATILL	PRCA	HCC	
171	LYTVPFNLI	HCC	MEL	
172	RYFIAAEKILW	OSCAR	HNSCC	
173	RYLSVCERL	HNSCC, NSCLCadenoc, NSCLCsquam, OSCAR, SCLC, UEC	NSCLCother	
174	TYGEEMPEEI	NSCLCother, NSCLCsquam	HNSCC, OSCAR	
175	SYFEYRRL	GC, NSCLCadenoc, NSCLCother, PACA, UBC	CCC, HNSCC, NSCLCsquam, OSCAR	
176	TQAGEYLLF		AML	
177	KYLITTFSL	BRCA, GC, HNSCC, NHL, NSCLCadenoc, OSCAR, PACA, PRCA, SCLC	CCC, GBC, NSCLCsquam, OC, UBC, UEC	
178	AYPQIRCTW		AML	
179	MYNMVPPF		MEL	
180	IYNKTKMAF	HNSCC, MEL, OC, OSCAR, SCLC	HCC, UBC	

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
181	IHGKIFHYF	PACA, UEC	GC	
182	AQGSGTVTF		CLL, NHL	
183	YQVAKGMEF		AML	
184	VYVRPRVF		MEL	
185	LYICKVELM	NHL	CLL	
186	RRVTWNVLF	SCLC	GBM	
187	KWFNVRMGFGF	GBC, MEL, OC, UBC, UEC	HCC, SCLC	
188	SLPGSFIYVF		MEL	
189	FYPDEDDFYF	GBM, MEL, OC, SCLC, UBC	AML, UEC	
190	IYIIMQSCW		AML	
191	MSYSCGLPSL	HNSCC	MEL	
192	CYSFIHLSF	BRCA, CCC, GC, HNSCC, NHL, NSCLCadenoc, OSCAR, PACA, PRCA, SCLC	GBC, NSCLCsquam, OC, UBC, UEC	
193	KYKPVALQCIA		MEL	
194	EYFLPSLEII		MEL	
195	IYNEHGIQQI	CCC, CRC, GBC, GBM, GC, HNSCC, NSCLCother, NSCLCsquam, OC, OSCAR	BRCA, MEL, NSCLCadenoc, PACA	
196	VGRSPVFLF	CCC, CRC, GBC, GBM, GC, HNSCC, NSCLCother, NSCLCsquam, OC, OSCAR	BRCA, MEL, NSCLCadenoc, PACA	
197	YYHSGENLY	MEL	CCC	
198	VLAPVSGQF	NHL	CLL	
199	MFQFEHIKW		HCC	
200	LYMSVEDFI	GC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, UBC	CRC, GBC	
201	VFPSVDVSF		GBM	
202	VYDTMIEKFA		GBM	
203	VYPSESTVM		GBM	
204	WQNVTPLTF	BRCA	GBM, MEL	
205	ISWEVVHTVF		MEL	

- 130 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
206	EVVHTVFLF		MEL	
207	IYKFIMDRF	NHL, NSCLCadeno, NSCLCsquam, OC	SCLC, UEC	
208	QYLQQQAKL	NHL, NSCLCadeno, NSCLCsquam, OC	SCLC, UEC	
209	DIYVTGGHLF	BRCA, NSCLCsquam, OSCAR, UBC	HNSCC	
210	EAYSYPATI		MEL	
211	MLYFAPDLIL	BRCA	UEC	
212	VYFVQYKIM	BRCA, HNSCC, NSCLCother, NSCLCsquam, OC, OSCAR, PACA, UBC, UEC	CCC, NHL, NSCLCadeno	
213	FYNRLTKLF	CRC, GBC, HNSCC, MEL, NHL, NSCLCadeno, NSCLCsquam, OC, PACA, UEC	OSCAR, UBC	
214	YIPMSVMLF	AML, NSCLCsquam, OSCAR	HNSCC, UBC	
215	KASKITFHW		GBM	
216	RHYHSIEVF		MEL	
217	QRYGFSSVGF		HCC	
218	FYFYNCSSL	BRCA, CLL, NHL, RCC	UBC	
219	KVVSGFYI	BRCA, CCC, HNSCC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, OSCAR, UBC	NHL	
220	TYATHVTEI	BRCA, CCC, HNSCC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, OSCAR, UBC	NHL	
221	VFYCLLFVF	BRCA, CCC, HNSCC, MEL, NSCLCadeno, NSCLCother, NSCLCsquam, OSCAR, UBC	NHL	

- 131 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
222	HYHAESFLF	OSCAR	HNSCC	
223	KLRALSILF		GBM	
224	AYLQFLSVL	BRCA, HCC, MEL, OC, SCLC, UEC	PRCA	
225	ISMSATEFLL		NSCLCother	
226	TYSTNRTMI		AML	
227	YLPNPSLNAF		NSCLCother	
228	VYLRIGGF	CCC, HNSCC, PACA, UBC, UEC	OC	
229	CAMPVAMEF		CLL, NHL	
230	RWLSKPSLL		CLL, NHL	
231	KYSVAFYSLD	MEL, NSCLCsquam, RCC, SCLC	OC, UEC	
232	IWPGFTTSI	GBC, GC, PACA, SCLC, UEC	CRC	
233	LYSRRGVRTL	CCC, OC	PACA	
234	RYKMLIPF	BRCA, CCC, GC, HNSCC, NHL, NSCLCadenocarcinoma, NSCLCsquam, OSCAR, PACA, SCLC, UEC	GBC, OC, UBC	
235	VYISDVSVY	AML, NHL	CLL	
236	LHLYCLNTF	BRCA	UEC	
237	RQGLTVLTW	AML, BRCA, CRC, GC, MEL, NSCLCadenocarcinoma, OC, OSCAR, PACA, PRCA, SCLC, UEC	NHL	
238	YTCSRAVSLF	GBC, HCC, NSCLCother, OC, RCC	SCLC	
239	IYTFSNVTF		NHL	
240	RVHANPLLI		HCC	
241	QKYYITGEAEGF	OC	BRCA, UEC	
242	SYTPLLSYI	SCLC	GBM	
243	ALFPMGPLTF	NHL	CLL	
244	TYIDTRTVFL	GC	UEC	
245	VLPLHFLPF	HCC, HNSCC, NSCLCsquam, UBC	GBC, MEL, NHL	
246	KIYTTVLFANI	GBC	HCC	
247	VHSYLGSPF		AML	

- 132 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
248	CWGPFCFEM	GBM	RCC	
249	HQYGGAYNRV		UEC	
250	VYSDRQIYLL		BRCA	
251	DYLLSWLLF	NHL	CLL	
252	RYLIIKYPF	AML	CCC	
253	QYYCLLLIF	CCC	AML	
254	KQHAWLPLTI		CLL, NHL	
255	VYLDEKQHAW		CLL, NHL	
256	QHAWLPLTI		CLL, NHL	
257	MLILFFSTI	GBC, HCC, HNSCC, MEL, NSCLCaden, NSCLCoth, NSCLCsquam, OC, OSCAR, SCLC, UBC, UEC	BRCA	
258	VCWNPFNNTF	NSCLCoth, OC, RCC, SCLC	UEC	
259	FFLFIPFF	NSCLCaden, PRCA	OC	
260	FLFIPFFIIF	NSCLCaden, PRCA	OC	
261	IMFCLKNFWW		MEL	
262	YIMFCLKNF		MEL	
263	AYVTEFVSL		SCLC	
264	AYAIPSASLSW		MEL	
265	LYQQSDTWSL		CLL	
266	TQIITFESF	NSCLCaden, NSCLCsquam, OSCAR, UBC	HNSCC	
267	QHMLPFWTDL	BRCA, CCC, NSCLCaden, NSCLCsquam, OC, OSCAR, UBC, UEC	GBC	
268	YQFGWSPNF		HCC	
269	FSFSTSMNEF		UEC	
270	GTGKLFVVF	NHL	CLL	
271	INGDLVFSF		UEC	
272	IYFNHRCF		CLL	
273	VTMYLPLLL		MEL	
274	EYSLPVLTF		GBM	
275	PEYSLPVLTF		GBM	
276	KFLGSKCSF	HNSCC,	UBC	

- 133 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
		NSCLCsquam, OSCAR		
277	MSAIWISAF		MEL	
278	TYESVVTGFF	HNSCC, NSCLCsquam, OSCAR	UBC	
279	KYKNPYGF	BRCA, CRC, GC, HNSCC, NHL, NSCLCother, OC, OSCAR, SCLC, UEC	NSCLCsquam	
280	TIYSLEMKMSF	NSCLCadenoc	PRCA	
281	MDQNQVVWTF	NSCLCother, NSCLCsquam	NSCLCadenoc	
282	ASYQQSTSSFF		CLL	
283	SYIVDGKII	HNSCC, MEL, NSCLCsquam, PACA	CCC	
284	QFYSTLPNTI	NSCLCother, NSCLCsquam	NSCLCadenoc	
285	YFLPGPHYF	CLL, CRC, GBM, GC, HNSCC, MEL, NSCLCadenoc, NSCLCsquam, OC, OSCAR, PACA, PRCA, SCLC	NHL	
286	HHTQLIFVF	MEL, NSCLCsquam, OSCAR	HNSCC	
287	LVQPQAVLF	NHL	CLL	
288	MGKGSISFLF	GBM, PACA	SCLC	
289	RTLNEIYHW		NHL	
290	VTPKMLISF		HCC	
291	YTRLVLQF	GBC, GC, UEC	PACA	
292	KMFPKDFRF	PACA	RCC	
293	MYAYAGWFY	HNSCC, NSCLCadenoc, OSCAR	NSCLCsquam	
294	KMGRIVDYF	GBC, GC, UEC	PACA	
295	KYNRQSMTL		HCC	
296	YQRPDLLLF	NHL	CLL	
297	LKSPRLFTF		UBC	
298	TYETVMTFF		UBC	
299	FLPALYSLL		CLL	

- 134 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
300	LFALPDFIF		CLL	
301	RTALSSTDTF		CLL	
302	YQGSLEVLFF	UEC	UBC	
303	RFLDRGWGF	BRCA, CRC, GBC, GC, HNSCC, NSCLCadeno, NSCLCother, NSCLCsquam, OSCAR, PACA, UBC	CCC	
304	YFGNPQKF	CCC, GC, HNSCC, NSCLCother, NSCLCsquam, PACA, UBC	OSCAR	
305	RNAFSIYIL	HNSCC	MEL	
306	RYILEPFFI	HNSCC, NSCLCadeno, OSCAR	NSCLCsquam	
307	RILTEFELL	GBC, GC	CRC	
308	AAFISVPLLI	GBC, GC, PACA, UBC	CRC	
309	AFISVPLLI	GBC, GC, PACA, UBC	CRC	
310	EFINGWYVL	NHL	CLL	
311	IQNAILHLF	GBC, HCC, HNSCC, MEL, NSCLCadeno, NSCLCother, PACA, SCLC	NHL	
312	YLCMLYALF	GBM	OC	
313	IFMENAFEL		HCC	
314	SQHFNLATF		AML	
315	VYDYIPLLL	UEC	UBC	
316	IWAERIMF	BRCA	PRCA	
317	DWIWRILFLV	BRCA, MEL, NSCLCsquam	GBC	
318	VQADAKLLF	GC, HNSCC, NSCLCsquam, OSCAR, UBC	CRC	
319	ATATLHLIF	GBM		
320	EVYQKIILKF	GBC, MEL, OC, SCLC		
321	VYTVGHNLI	HCC		
322	SFISPRYSWLF	AML		

- 135 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
323	NYSPVTGKF	MEL		
324	RYFVSNIYL	AML, GBC, MEL, OC		
325	IFMGAVPTL	HNSCC, OC, OSCAR, PRCA		
326	VHMKDFFYF	CLL, NHL		
327	KWKPSPLLF	CCC		
328	IYLVGGYSW	CLL, OC		
329	YLGKNWSF	AML		
330	DYIQMIPEL	SCLC		
331	EYIDEFQSL	SCLC		
332	VYCSLDKSQF	SCLC		
333	RYADLLIYTY	HNSCC, OC, UBC, UEC		
334	KVFGSFLTTL	NSCLCother, UEC		
335	RYQSVIYPF	NSCLCother, UEC		
336	VYSDLHAFY	SCLC		
337	SHSDHEFLF	BRCA, GBC, MEL, NSCLCadeno, NSCLCother, OC, PRCA, UBC, UEC		
338	VYLTWLPGL	CLL, NHL		
339	KQVIGIHTF	CLL		
340	FPPTPPLF	AML, CLL, NHL		
341	RYENVSILF	GBM		
342	MYGIVIRTI	CRC, GBC, GC, PACA		
343	EYQQYHPSL	CLL, NHL		
344	YAYATVLTF	PRCA		
345	RYLEEHTEF	UBC		
346	TYIDFVPYI	MEL, OC, RCC, SCLC		
347	AWLIVLLFL	GBC, NHL, OC		
348	RSWENIPVTF	BRCA, MEL, NHL, SCLC		
349	IYMTTGVL	MEL		
350	VYKWTEEF	CRC, PACA, SCLC		
351	GYFGTASLF	NSCLCother		
352	NAFEAPLTF	AML, CLL, CRC, HNSCC, NHL, NSCLCsquam, SCLC		
353	AAFPGAFSF	GBM		

- 136 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
354	QYIPTFHVY	GBM, HCC, HNSCC, NSCLCadeno, OC, OSCAR, SCLC, UBC		
355	VYNNNSSRF	MEL		
356	YSLEHLTQF	HCC		
357	RALLPSPLF	HCC		
358	IYANVTEMLL	HNSCC, OSCAR, UBC		
359	TQLPAPLRI	GBM		
360	LYITKVTTI	UBC		
361	KQPANFIVL	GBC, GC, PACA		
362	NYMDTDNLMF	GBC, GC, PACA		
363	QYGFNYNKF	BRCA, HNSCC, MEL, OSCAR		
364	KQSQVVFVL	MEL		
365	KDLMKAYLF	PRCA, SCLC		
366	RLGEFVLLF	HNSCC, NSCLCsquam		
367	HWSHITHLF	CCC, GBC, GBM, MEL		
368	AYFVAMHLF	GBM, OC, SCLC		
369	NFYLFPTTF	HNSCC, MEL, OSCAR		
370	TQMDVKLVF	CLL		
371	FRSWAVQTF	CRC		
372	LYHNWRHAF	PRCA		
373	IWDALERTF	BRCA		
374	MIFAVVLF	NHL		
375	YYAADQWVF	NHL		
376	KYVGEVFN	PRCA		
377	SLWREVTF	SCLC		
378	VYAVISNIL	GBM		
379	KLPTEWNVL	CLL		
380	FYIRRLPMF	MEL		
381	IYTDITYSF	MEL		
382	SYPKELMKF	UBC		
383	PYFSPSASF	CRC, GBC, HNSCC, NSCLCadeno, NSCLCsquam, OC, OSCAR, SCLC	UBC	

- 137 -

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
384	RTRGWVQTL	HCC, NSCLCadeno, NSCLCsquam, PACA, SCLC	CRC, GBC, GC, RCC	
385	GYFGNPQKF	CCC, GC, HNSCC, NSCLCother, NSCLCsquam, PACA, UBC	OSCAR	
386	YQSRDYYNF	BRCA, HCC, PRCA		
387	THAGVRLYF	NSCLCother, NSCLCsquam, SCLC		

Table 9b: Expression scores. The table lists for each peptide the tumor types for which the exon is very highly over-expressed in tumors compared to a panel of normal tissues (+++), highly over-expressed in tumors compared to a panel of normal tissues (++) or over-expressed in tumors compared to a panel of normal tissues (+). The baseline for this score was calculated from measurements of the following relevant normal tissues: blood cells, blood vessels, brain, heart, liver, lung, adipose tissue, adrenal gland, bile duct, bone marrow, esophagus, eye, gallbladder, head-and-neck, kidney, large intestine, lymph node, pancreas, parathyroid gland, peripheral nerve, peritoneum, pituitary, pleura, skeletal muscle, skin, small intestine, spleen, stomach, thyroid gland, trachea, ureter, urinary bladder. In case expression data for several samples of the same tissue type were available, the arithmetic mean of all respective samples was used for the calculation. AML: acute myeloid leukemia; GBC: gallbladder cancer; HNSCC: head and neck squamous cell carcinoma; MEL: melanoma; NHL: non-Hodgkin lymphoma; NSCLCsquam: squamous cell non-small cell lung cancer; OC: ovarian cancer; OSCAR: esophageal cancer.

Seq ID No	Sequence	Exon Expression in tumor types vs normal tissue panel		
		over-expressed (+)	highly over-expressed (++)	very highly over-expressed (+++)
463	VGGNVTSNF	AML, HNSCC, MEL, NHL, OSCAR	GBC, NSCLCsquam, OC	
464	VGGNVTSNF	HNSCC, NHL	GBC, NSCLCsquam, OC	

EXAMPLE 3

In vitro immunogenicity for MHC class I presented peptides

In order to obtain information regarding the immunogenicity of the TUMAPs of the present invention, the inventors performed investigations using an *in vitro* T-cell priming assay based on repeated stimulations of CD8⁺ T cells with artificial antigen presenting cells (aAPCs) loaded with peptide/MHC complexes and anti-CD28 antibody. This way the inventors could show immunogenicity for HLA-A*24:02 restricted TUMAPs of the invention, demonstrating that these peptides are T-cell epitopes against which CD8⁺ precursor T cells exist in humans (Table 10).

In vitro priming of CD8⁺ T cells

In order to perform *in vitro* stimulations by artificial antigen presenting cells loaded with peptide-MHC complex (pMHC) and anti-CD28 antibody, the inventors first isolated CD8⁺ T cells from fresh HLA-A*24 leukapheresis products via positive selection using CD8 microbeads (Miltenyi Biotec, Bergisch-Gladbach, Germany) of healthy donors obtained from the University clinics Mannheim, Germany, after informed consent.

PBMCs and isolated CD8⁺ lymphocytes were incubated in T-cell medium (TCM) until use consisting of RPMI-Glutamax (Invitrogen, Karlsruhe, Germany) supplemented with 10% heat inactivated human AB serum (PAN-Biotech, Aidenbach, Germany), 100 U/ml Penicillin/100 µg/ml Streptomycin (Cambrex, Cologne, Germany), 1 mM sodium pyruvate (CC Pro, Oberdorla, Germany), 20 µg/ml Gentamycin (Cambrex). 2.5 ng/ml IL-7 (PromoCell, Heidelberg, Germany) and 10 U/ml IL-2 (Novartis Pharma, Nürnberg, Germany) were also added to the TCM at this step.

Generation of pMHC/anti-CD28 coated beads, T-cell stimulations and readout was performed in a highly defined *in vitro* system using four different pMHC molecules per stimulation condition and 8 different pMHC molecules per readout condition.

The purified co-stimulatory mouse IgG2a anti human CD28 Ab 9.3 (Jung et al., 1987) was chemically biotinylated using Sulfo-N-hydroxysuccinimidobiotin as recommended

- 139 -

by the manufacturer (Perbio, Bonn, Germany). Beads used were 5.6 μm diameter streptavidin coated polystyrene particles (Bangs Laboratories, Illinois, USA).

pMHC used for positive and negative control stimulations were A*0201/MLA-001 (peptide ELAGIGILTV (SEQ ID NO. 461) from modified Melan-A/MART-1) and A*0201/DDX5-001 (YLLPAIVHI from DDX5, SEQ ID NO. 462), respectively.

800.000 beads / 200 μl were coated in 96-well plates in the presence of 4 x 12.5 ng different biotin-pMHC, washed and 600 ng biotin anti-CD28 were added subsequently in a volume of 200 μl . Stimulations were initiated in 96-well plates by co-incubating 1×10^6 CD8+ T cells with 2×10^5 washed coated beads in 200 μl TCM supplemented with 5 ng/ml IL-12 (PromoCell) for 3 days at 37°C. Half of the medium was then exchanged by fresh TCM supplemented with 80 U/ml IL-2 and incubating was continued for 4 days at 37°C. This stimulation cycle was performed for a total of three times. For the pMHC multimer readout using 8 different pMHC molecules per condition, a two-dimensional combinatorial coding approach was used as previously described (Andersen et al., 2012) with minor modifications encompassing coupling to 5 different fluorochromes. Finally, multimeric analyses were performed by staining the cells with Live/dead near IR dye (Invitrogen, Karlsruhe, Germany), CD8-FITC antibody clone SK1 (BD, Heidelberg, Germany) and fluorescent pMHC multimers. For analysis, a BD LSR II SORP cytometer equipped with appropriate lasers and filters was used. Peptide specific cells were calculated as percentage of total CD8+ cells. Evaluation of multimeric analysis was done using the FlowJo software (Tree Star, Oregon, USA). *In vitro* priming of specific multimer+ CD8+ lymphocytes was detected by comparing to negative control stimulations. Immunogenicity for a given antigen was detected if at least one evaluable *in vitro* stimulated well of one healthy donor was found to contain a specific CD8+ T-cell line after *in vitro* stimulation (i.e. this well contained at least 1% of specific multimer+ among CD8+ T-cells and the percentage of specific multimer+ cells was at least 10x the median of the negative control stimulations).

- 140 -

In vitro immunogenicity for acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-Hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer peptides

For tested HLA class I peptides, *in vitro* immunogenicity could be demonstrated by generation of peptide specific T-cell lines. Exemplary flow cytometry results after TUMAP-specific multimer staining for 2 peptides of the invention are shown in Figure 3 together with corresponding negative controls. Results for 47 peptides from the invention are summarized in Table 10a and Table 10b.

Table 10a: *in vitro* immunogenicity of HLA class I peptides of the invention

Exemplary results of *in vitro* immunogenicity experiments conducted by the applicant for the peptides of the invention.

<20 % = +; 20 % - 49 % = ++; 50 % - 69 % = +++; >= 70 % = ++++

Seq ID No	Sequence	Peptide Code	Wells positive [%]
390	KYKDYFPVI	MAGEC2-003	+
392	SYEKVINYL	MAGEA9-001	+
393	SYNDALLTF	TRPM8-004	+++
395	NYEDHFPLL	MAGEA10-002	++
398	GYLQGLVSF	KLK4-001	++
399	VWSNVTPLKF	MMP12-014	+
400	RYLEKFYGL	MMP12-006	+
402	TYKYVDINTF	MMP12-004	+
406	KYLEKYYNL	MMP1-001	++
408	VWSDVTPLTF	MMP11-001	+
409	VYTFLSSTL	ESR1-006	+
411	VYPPYLNLYL	PGR-002	++++
415	KYEKIFEML	CT45-002	+
416	VFMKDGFFYF	MMP1-002	+
420	VYEKNGYIYF	MMP13-001	++++

- 141 -

Seq ID No	Sequence	Peptide Code	Wells positive [%]
422	VWSDVTPLNF	MMP13-002	+
429	YYSKSVGFMQW	FAM111B-008	++
433	NYTSLLVTW	PTP-018	+
434	VYDTMIEKF	PTP-016	+
436	KYLQVVGMF	OXTR-001	+
440	NYGVLHVTF	NLRP11-002	+
447	EYIRALQQL	ASCL1-001	+
448	PFLPPAACFF	ASCL1-002	+
450	QYDPTPLTW	ADAMTS12-002	+
453	YYTVRNFTL	PTP-014	+
455	KYLSIPTVF	UGT1A3-001	+

Table 10b: *in vitro* immunogenicity of HLA class I peptides of the invention

Exemplary results of *in vitro* immunogenicity experiments conducted by the applicant for the peptides of the invention.

<20 % = +; 20 % - 49 % = ++; 50 % - 69 % = +++; >= 70 % = ++++

Seq ID No	Sequence	Peptide Code	Wells positive [%]
1	IFPKTGLLII	MAGEA4-008	++
5	LQPQPQLFFSF	POT-002	++
22	VYSSFVFNLF	NLRP4-006	+
23	VYSSFVFNLF	NLRP4-007	++
26	TYFFVDNQYW	MMP12-020	+
32	FYPEVELNF	MMP1-009	+++
38	RYAHTLVTSVLF	ITIH6-003	++
44	FYHEDMPLW	FCRL5-004	+
47	SYLWTDNLQEF	DNAH17-003	+
52	SWHKATFLF	COL24-002	+
56	FYKLIQNGF	FLT3-010	+
58	IYYSHENLI	F5-005	++
63	LYIDKGQYL	HMCN1-011	+
76	IYIYPFAHW	NPFFR2-002	+++
77	LYGFFFKI	BTBD16-002	++++
78	TYSKTIALYGF	BTBD16-004	++
79	FYIVTRPLAF	SUCN-002	+
81	AYLKLLPMF	SLC5A4-001	+
86	SYLPTAERL	SYT12-001	+

- 142 -

87	NYTRLVLQF	GABRP-004	+
88	TYVPSTFLV	GABRP-005	+

EXAMPLE 4

Synthesis of peptides

All peptides were synthesized using standard and well-established solid phase peptide synthesis using the Fmoc-strategy. Identity and purity of each individual peptide have been determined by mass spectrometry and analytical RP-HPLC. The peptides were obtained as white to off-white lyophilizes (trifluoro acetate salt) in purities of >50%. All TUMAPs are preferably administered as trifluoro-acetate salts or acetate salts, other salt-forms are also possible.

EXAMPLE 5

MHC Binding Assays

Candidate peptides for T cell based therapies according to the present invention were further tested for their MHC binding capacity (affinity). The individual peptide-MHC complexes were produced by UV-ligand exchange, where a UV-sensitive peptide is cleaved upon UV-irradiation, and exchanged with the peptide of interest as analyzed. Only peptide candidates that can effectively bind and stabilize the peptide-receptive MHC molecules prevent dissociation of the MHC complexes. To determine the yield of the exchange reaction, an ELISA was performed based on the detection of the light chain ($\beta 2m$) of stabilized MHC complexes. The assay was performed as generally described in Rodenko et al. (Rodenko et al., 2006)

96 well MAXISorp plates (NUNC) were coated over night with 2ug/ml streptavidin in PBS at room temperature, washed 4x and blocked for 1h at 37°C in 2% BSA containing blocking buffer. Refolded HLA-A*02:01/MLA-001 monomers served as standards, covering the range of 15-500 ng/ml. Peptide-MHC monomers of the UV-exchange reaction were diluted 100-fold in blocking buffer. Samples were incubated for 1h at 37°C, washed four times, incubated with 2ug/ml HRP conjugated anti- $\beta 2m$ for 1h at 37°C, washed again and detected with TMB solution that is stopped with NH_2SO_4 . Absorption was measured at 450nm. Candidate peptides that show a high exchange

- 143 -

yield (preferably higher than 50%, most preferred higher than 75%) are generally preferred for a generation and production of antibodies or fragments thereof, and/or T cell receptors or fragments thereof, as they show sufficient avidity to the MHC molecules and prevent dissociation of the MHC complexes.

Table 11: MHC class I binding scores. Binding of HLA-class I restricted peptides to HLA-A*24 was ranged by peptide exchange yield: >10% = +; >20% = ++; >50 = +++; >75% = ++++

Seq ID No	Sequence	Peptide Code	Peptide exchange
1	IFPKTGLLII	MAGEA4-008	++++
2	LYAPTILLW	AFP-001	++++
3	KFLTHDVLTELF	TRPM8-009	++++
5	LQPQPQLFFSF	POT-002	++++
6	IVTFMNKTLGTF	ADAM29-001	+
9	TYINSLAIL	TGM4-003	++++
10	QYPEFSIEL	TGM4-001	+++
11	RAMCAMMSF	TGM4-002	++
12	KYMSRVLFVY	CHRNA9-002	++
13	KYYIATMAL	CHRNA9-003	+++
14	YYIATMALI	CHRNA9-004	++++
15	FMVIAGMPLF	SLC6A3-001	++
16	GYFLAQYLM	TRPM8-008	+++
17	IYPEAIATL	SLC6A3-002	++++
18	KYVDINTFRL	MMP12-018	++++
20	ELMAHPFLL	CYP4Z-002	++
21	LYMRFVNTHF	SPINK2-002	++
22	VYSSFVFNL	NLRP4-006	+++
23	VYSSFVFNLF	NLRP4-007	+++
24	KMLPEASLLI	NLRP4-004	++
25	MLPEASLLI	NLRP4-005	++++
26	TYFFVDNQYW	MMP12-020	++++
27	LSCTATPLF	KHDC1L-001	+++
28	FWFDSREISF	OR51E2-002	++++
29	IYLLPPIVI	OR51E2-004	++++
30	RQAYSVYAF	SLC45A3-005	++
31	KQMQEFGGL	MMP1-010	+++
32	FYPEVELNF	MMP1-009	++++
33	FYQPDLKYLFSF	NLRP4-003	++++
34	LIFALALAAF	GAST-001	+

- 144 -

35	FSSTLVSLF	MAGEC1-003	++
36	VYLASVAAF	SLC45A3-006	++++
37	ISFSDTVNVW	ITIH6-001	+
38	RYAHTLVTSVLF	ITIH6-003	++++
39	KTYLPTFETTI	ENP-002	++
40	NYPEGAAYEF	ESR1-008	++++
41	IYFATQVVF	SLC45A3-004	++++
42	VYDSIWCNM	SCGB2A1-001	++++
43	KYKDHFEI	MAGEB1-001	++++
44	FYHEDMPLW	FCRL5-004	++++
45	YGQSKPWTF	PAX3-001	++
46	IYPDSIQEL	LOC-019	+++
47	SYLWTDNLQEF	DNAH17-003	++++
48	AWSPPATLFLF	LOXL4-003	++++
49	QYLSIAERAEF	MSX-001	++++
50	RYFDENIQKF	HEPHL-004	++++
51	YFDENIQKF	HEPHL-006	+++
52	SWHKATFLF	COL24-002	++++
53	LFQRVSSVSF	HMCN1-010	+++
54	SYQEAIQQL	NEFH-003	+++
55	AVLRHLETF	CDK6-006	++
56	FYKLIQNGF	FLT3-010	++++
58	IYSHENLI	F5-005	++++
59	VFPLVTPLL	PTP-042	++++
60	RYSVPKDAW	KLHDC7B-006	++++
61	RIFTARLYF	AICD-001	+++
62	VYIVPVIVL	OXTR-002	++++
63	LYIDKGQYL	HMCN1-011	++++
64	QFSHVPLNNF	ALX1-001	++
66	IYKDYRYNF	PLA2G2D-001	++++
67	SYVLQIVAI	PTP-041	+++
68	VYKEDLPQL	EML-002	++++
69	KWFDSHIPRW	ERV-002	++++
70	RYTGQWSEW	IL9R-001	++++
71	RYLPNPSLNAF	CYP1A1-002	++++
72	RWLDGSPVTL	CLEC17-005	++++
73	YFCSTKGQLF	FCRL2-004	++++
74	NYVLVPTMF	CAPN6-003	++++
75	VYEHNHVSL	BTBD16-006	+++
76	IYIYPFAHW	NPFFR2-002	++++
77	LYGFFFKI	BTBD16-002	++++

- 145 -

78	TYSKTIALYGF	BTBD16-004	++++
79	FYIVTRPLAF	SUCN-002	+++
80	SYATPVDLW	CDK6-007	++++
81	AYLKLLPMF	SLC5A4-001	++++
82	SYLENSASW	DLX5-003	++++
83	VLQGEFFLF	KBTBD8-005	++++
84	YTIERYFTL	GABRP-007	++++
85	KYLSIPTVFF	UGT1A3-002	++++
86	SYLPTAERL	SYT12-001	++++
87	NYTRLVLQF	GABRP-004	++++
88	TYVPSTFLV	GABRP-005	++++
89	TYVPSTFLVVL	GABRP-006	++++
90	TDLVQFLLF	MAGEA10-003	++
92	RALTETIMF	ALP-011	++
93	TDWSPPPVEF	FAM178B-001	+++
94	THSGGTNLF	MMP12-019	++
95	IGLSVVHRF	OR51E2-003	++++
96	SHIGVVLAFLAF	OR51E2-005	++
98	LQIPVSPSF	MAGEC1-004	+
99	ASAALTGFTF	SLC45A3-003	++
100	KVWSDVTPLTF	MMP11-023	+++
101	VYAVSSDRF	DCX-001	++++
102	VLASAHILQF	BTBD16-005	++
103	EMFFSPQVF	ACTL8-002	++
104	GYGLTRVQPF	ACTL8-003	++
106	LYAFLGSHF	KISS1R-003	++++
108	WFFQGAQYW	MMP11-024	++
109	AQHSLTQLF	GPC2-002	++
110	VYSNPDLFW	TRDV3-002	++++
111	IRPDYSFQF	TRDV3-001	++
112	LYPDSVFGRLF	SMC1B-003	++++
113	ALMSAFYTF	MMP11-020	++++
114	KALMSAFYTF	MMP11-022	+++
115	IMQGFIRAF	PAE-002	++
116	TYFFVANKY	MMP1-011	+
117	RSMEHPGKLLF	ESR1-010	+++
118	IFLPPFIVF	ADAM18-001	++++
119	VWSCEGCKAF	ESR-001	++
120	VYAFMNENF	QRFPR-004	++++
121	RRYFGKVAL	ANO7-005	++
123	FFLQESPVF	ABCC11-004	++++

- 146 -

124	EYNVFPRTL	MMP13-004	++
125	LYYGSILYI	OR9-001	++++
126	YSLLDPAQF	SOX14-002	+++
127	FLPRAYYRW	ANO7-001	++
128	AFQNVISF	NMUR2-003	++++
129	IYVSLAHVL	ANO7-002	++++
130	RPEKVFVF	COL11A1-005	++
131	MHRTWRETF	ANO7-004	++
133	FFYVTETTF	TERT-003	++++
134	IYSSQLPSF	TFEC-004	++++
135	KYKQHFPEI	MAGEB17-001	+++
136	YLKSVQLF	RFX8-001	++
137	ALFAVCWAPF	QRFPR-002	++
138	MMVTVVALF	QRFPR-003	+++
139	AYAPRGSYKF	HHIPL2-001	++++
140	IFQHFCEEI	SMC1B-002	++
141	QYAAAITNGL	SALL3-002	+++
142	PYWWNANMVF	NOTU-001	++++
143	KTKRWLWDF	COL11A1-004	+++
144	LFDHGGTVFF	ANO7-003	+++
145	MYTIVPML	OR1N1-001	++++
146	NYFLDPVTI	TRI-005	+++
148	MLPQIPFLLL	COL10-001	++
149	TQFFIPYTI	COL10-002	++
150	FIPVAWLIF	MRGPRX4-001	+++
151	RRLWAYVTI	ITIH6-002	+
152	MHPGVLAALF	MMP13-005	+++
153	AWSPPATLF	LOXL4-002	++++
154	DYSKQALSL	LAMC2-018	++
155	PYSIYPHGVTF	F5-006	++++
156	IYPHGVTFSP	F5-004	+
157	SIYPHGVTF	F5-007	++
158	SYLKDPMIV	DDX53-001	+++
159	VFQPNPLF	WISP3-002	++
160	YIANLISCF	GLYATL3-001	++
161	ILQAPLSVF	FCRL5-005	++
162	YYIGIVEEY	HEPHL-007	+++
163	YYIGIVEEYW	HEPHL-008	++++
164	MFQEMLQRL	TRIML2-001	+
165	KDQPQVPCVF	NAT1-001	+
166	MMALWSLLHL	ZAC-001	+

- 147 -

167	LQPPWTTVF	FCRL5-006	++
168	LSSPVHLDF	FCRL5-007	++++
169	MYDLHHLYL	EPY-001	++++
170	IFIPATILL	ACSM1-001	++++
171	LYTVPFNLI	SLC45A2-006	++++
172	RYFIAAEKILW	HEPHL-005	++++
173	RYLSVCERL	NKX-001	++++
174	TYGEEMPEEI	DNAH17-004	++
175	SYFEYRLL	LAMC2-019	++
176	TQAGEYLLF	FLT3-012	++++
177	KYLITTFSL	NLRP2-008	++++
178	AYPQIRCTW	FLT3-009	++++
179	MYNMVPPF	DCT-002	+++
180	IYNKTKMAF	SLCO6-001	++++
181	IHGKIFHYF	NMUR2-004	+++
182	AQSGSTVTF	FCRL3-004	++
183	YQVAKGMEF	FLT3-014	+
184	VYVRPRVF	HMCN1-013	++
185	LYICKVELM	CTL-001	++++
186	RRVTWNVLF	BTBD17-003	++
187	KWFNVRMGFGF	LIN-001	++
188	SLPGSFIYVF	HMCN1-012	++
189	FYPDEDDFYF	MYCN-002	+++
190	IYIIMQSCW	FLT3-011	+++
191	MSYSCGLPSL	KRT33A-001	+++
192	CYSFIHLSF	NLR-006	++++
193	KYKPVALQCIA	HMCN1-009	++
195	IYNEHQIQI	COL11A1-003	++++
196	VGRSPVFLF	COL11A1-006	++
198	VLAPVSGQF	FCRL5-009	+++
199	MFQFEHIKW	FBXW10-001	++
200	LYMSVEDFI	STK31-001	++++
201	VFPSVDVSF	PTP-043	++++
202	VYDTMIEKFA	PTP-044	++
203	VYPSESTVM	PTP-045	++
204	WQNVTPLEF	MMP16-002	++
205	ISWEVVHTVF	HMCN1-008	++++
206	EVVHTVFLF	HMCN1-007	++
207	IYKFIMDRF	FOXB1-001	++++
208	QYLQQQAKL	FOXB1-002	++++
209	DIYVTGGHLF	KLHDC7B-005	+++

- 148 -

210	EAYSYPPTI	HMCN1-006	+++
211	MLYFAPDLIL	PGR-004	++
212	VYFVQYKIM	IL22RA2-001	+
213	FYNRLTKLF	OFCC-001	++++
214	YIPMSVMLF	HTR7-002	+++
215	KASKITFHW	PTP-038	++
216	RHYHSIEVF	LOXL4-004	++
217	QRYGFSSVGF	RHBG-001	++++
218	FYFYNCSSL	ERV-001	++++
219	KVVSIFYI	CCR8-003	+
220	TYATHVTEI	CCR8-004	++++
222	HYHAESFLF	HEPHL-003	++++
223	KLRALSILF	PTP-039	+
224	AYLQFLSVL	GREB-002	++++
225	ISMSATEFLL	CYP1A1-001	+++
226	TYSTNRTMI	FLT3-013	++++
227	YLPNPSLNAF	CYP1A1-003	++
228	VYLRIGGF	WNT7A-002	++
229	CAMPVAMEF	KBTBD8-003	++
230	RWLSKPSLL	KBTBD8-004	++++
231	KYSVAFYSLD	LAMA1-008	++
232	IWPGFTTSI	PIWIL1-003	++++
234	RYKMLIPF	NLRP2-010	+++
235	VYISDVSVY	CLECL-003	++++
236	LHLYCLNTF	PGR-003	+++
238	YTCSRAVSLF	OTOG-001	++
239	IYTFSNVTF	BTN1-003	++++
240	RVHANPLLI	APOB-081	+
241	QKYIITGEAEGF	ESR1-009	++++
242	SYTPLLSYI	C1orf94-002	++++
243	ALFPMGPLTF	LILRA4-003	++
244	TYIDTRTVFL	CAPN6-005	++++
245	VLPLHFLPF	HBG2-001	++++
246	KIYTTVLFANI	NPFFR2-003	++++
247	VHSYLGSPF	MPL-001	++
249	HQYGGAYNRV	DLX5-002	++
250	VYSDRQIYLL	ABCC11-006	++++
251	DYLLSWLLF	CNR2-003	++++
252	RYLIIKYPF	SUCN-003	+++
254	KQHAWLPLTI	TCL1A-003	+
255	VYLDEKQHAW	TCL1A-005	++++

- 149 -

256	QHAWLPLTI	TCL1A-004	+
258	VCWNPFNNTF	RNF183-001	+++
259	FFLFIPFF	ADAM2-001	++
263	AYVTEFVSL	SCN3A-001	++++
264	AYAIPSASLSW	HMCN1-005	++
265	LYQQSDTWSL	KIAA140-001	++
266	TQITFESF	CSF2-001	++
267	QHMLPFWTDL	NLRP2-009	+++
269	FSFSTSMNEF	CAPN6-001	++
270	GTGKLFWVF	BTL-003	+
271	INGDLVFSF	CAPN6-002	++
272	IYFNHRCF	SFMBT1-001	+++
273	VTMYLPLLL	GPR143-001	++
274	EYSLPVLTF	PTP-037	+++
275	PEYSLPVLTF	PTP-040	++++
276	KFLGSKCSF	HAS3-003	++++
278	TYESVVTGFF	HAS3-004	++
279	KYKNPYGF	MMP20-001	+
280	TIYSLEMKMSF	GLB1L3-001	++
281	MDQNQVWTF	ROS-008	+++
282	ASYQQSTSSFF	FAM82A1-001	++
283	SYIVDGKII	PSG9-001	+
284	QFYSTLPNTI	ROS-009	+
285	YFLPGPHYF	SOX30-001	++++
288	MGKGSISFLF	PCSK1-001	++
289	RTLNEIYHW	FOXP3-001	++
290	VTPKMLISF	OR5H8P-001	+
291	YTRLVLQF	GABRP-008	++
292	KMFPKDFRF	TTLL6-001	++
294	KMGRIVDYF	GABRP-003	++++
295	KYNRQSMTL	APOB-080	++++
296	YQRPDLLLLF	GEN-004	++
297	LKSPRLFTF	BTBD16-001	++
298	TYETVMTFF	BTBD16-003	++++
299	FLPALYSLL	CXCR3-001	+++
300	LFALPDFIF	CXCR3-002	++
302	YQGSLEVLFF	MROH2A-006	++
303	RFLDRGWGF	ADAMTS12-005	++++
306	RYILEPFFI	SLC7A11-007	++++
307	RILTEFELL	TRIM31-001	+++
308	AAFISVPLLI	TAS2R38-002	++++

- 150 -

309	AFISVPLLI	TAS2R38-003	++++
310	EFINGWYVL	MCOLN2-002	++
311	IQNAILHLF	OR51B5-001	++++
312	YLCMLYALF	KCNK18-001	++
313	IFMENAFEL	APOB-079	++++
314	SQHFNLATF	DNMT3B-003	++
315	VYDYIPLLL	MROH2A-005	++++
316	IWAERIMF	TDRD1-001	+++
318	VQADAKLLF	C20-002	+
319	ATATLHLIF	PCD-007	+
320	EVYQKIILKF	PASD-001	++++
321	VYTVGHNLI	KLB-003	++++
322	SFISPRYSWLF	SPNS3-005	++++
323	NYSPVTGKF	OTOL-001	++++
325	IFMGAVPTL	LPAR3-001	++++
326	VHMKDFFYF	DYRK4-001	++++
327	KWKPSPLLF	GPR126-002	++++
328	IYLVGGYSW	KLHL14-007	++++
329	YLGNWSF	SPNS3-006	++
330	DYIQMPEL	RTL-002	++++
331	EYIDFQSL	RTL-003	++++
332	VYCSLDKSQF	RTL-004	++++
333	RYADLLIYTY	MYO3B-005	++++
334	KVFGSFLTL	AGTR2-001	++
335	RYQSVIYPF	AGTR2-002	++++
336	VYSDLHAFY	MANEAL-004	++++
337	SHSDHEFLF	ARSH-001	+
338	VYLTWLPGL	IFNLR-001	++++
339	KQVIGIHTF	SFMBT1-002	+
340	FPPTPPLF	BCL11A-002	+
341	RYENVSILF	ADCY8-001	++++
342	MYGIVIRTI	NPSR-001	++++
343	EYQQYHPSL	CLEC4C-001	++++
344	YAYATVLTf	ABCC4-002	++
345	RYLEEHTEF	MROH2A-003	++++
346	TYIDFVPYI	TEX15-001	++++
348	RSWENIPVTF	C18orf54-001	++
349	IYMTTGvLL	TDRD9-002	++++
350	VYKWTEEFK	TSPE-001	++++
351	GYFGTASLF	SLC16A14-001	++++
352	NAFEAPLTF	BRCA2-004	+++

353	AAFPGAFSF	CRB2-002	++
354	QYIPTFHVY	SLC44A5-003	++++
355	VYNNSSRF	MYO10-003	++++
356	YSLEHLTQF	ZCCHC16-001	++
357	RALLPSPLF	SPATA31D1-001	++
358	IYANVTEMLL	CYP27C-001	++++
359	TQLPAPLRI	GPR45-001	++
360	LYITKVTTI	FSTL4-002	++++
361	KQPANFIVL	LOC1001246-001	++
362	NYMDTDNLMF	LOC1001246-002	++++
363	QYGFNYNKF	PNLD-002	++++
365	KDLMKAYLF	TXNDC16-006	+++
366	RLGEFVLLF	TGM6-001	+++
367	HWSHITHLF	DPY19L1-003	++++
368	AYFVAMHLF	TENM4-007	++++
369	NFYLFPTTF	PNLD-001	++++
370	TQMDVKLVF	GEN-003	++
371	FRSWAVQTF	NOS2-002	++
372	LYHNWRHAF	PDE11-002	++++
373	IWDALERTF	ABCC11-005	++
375	YYAADQWVF	CCR4-004	++++
376	KYVGEVFNI	DMXL1-001	++++
377	SLWREVVTF	CEP250-004	+++
378	VYAVISNIL	TNR-003	++++
379	KLPTEWNVL	AKAP13-005	++++
380	FYIRRLPMF	CHRNA6-001	++++
381	IYTDITYSF	CHRNA6-002	++++
382	SYPKELMKF	MROH2A-004	++++
383	PYFSPSASF	SPER-001	++++
385	GYFGNPQKF	LAMA3-002	++++
386	YQSRDYYNF	AR-002	++++
387	THAGVRLYF	NUP155-012	++
463	VGGNVTSNF	CT45-004	+++
464	VGGNVTSSF	CT45-005	+++

Reference List

Allison, J. P. et al., Science. **270** (1995): 932-933

- 152 -

- Andersen, R. S. et al., Nat Protoc. **7** (2012): 891-902
- Appay, V. et al., Eur J Immunol. **36** (2006): 1805-1814
- Banchereau, J. et al., Cell. **106** (2001): 271-274
- Beatty, G. et al., J Immunol. **166** (2001): 2276-2282
- Beggs, J. D., Nature. **275** (1978): 104-109
- Benjamini, Y. et al., Journal of the Royal Statistical Society Series B (Methodological),. **Vol.57** (1995): 289-300
- Boulter, J. M. et al., Protein Eng. **16** (2003): 707-711
- Braumuller, H. et al., Nature.(2013)
- Brossart, P. et al., Blood. **90** (1997): 1594-1599
- Bruckdorfer, T. et al., Curr Pharm Biotechnol. **5** (2004): 29-43
- Card, K. F. et al., Cancer Immunol Immunother. **53** (2004): 345-357
- Cohen, C. J. et al., J Mol Recognit. **16** (2003a): 324-332
- Cohen, C. J. et al., J Immunol. **170** (2003b): 4349-4361
- Cohen, S. N. et al., Proc Natl Acad Sci U S A. **69** (1972): 2110-2114
- Coligan, J. E. et al., Current Protocols in Protein Science.(1995)
- Colombetti, S. et al., J Immunol. **176** (2006): 2730-2738
- Dengjel, J. et al., Clin Cancer Res. **12** (2006): 4163-4170
- Denkberg, G. et al., J Immunol. **171** (2003): 2197-2207
- Falk, K. et al., Nature. **351** (1991): 290-296
- Follenzi, A. et al., Nat Genet. **25** (2000): 217-222
- Fong, L. et al., Proc Natl Acad Sci U S A. **98** (2001): 8809-8814
- Gabrilovich, D. I. et al., Nat Med. **2** (1996): 1096-1103
- Gattinoni, L. et al., Nat Rev Immunol. **6** (2006): 383-393
- Gnjatic, S. et al., Proc Natl Acad Sci U S A. **100** (2003): 8862-8867
- Godkin, A. et al., Int Immunol. **9** (1997): 905-911
- Gragert, L. et al., Hum Immunol. **74** (2013): 1313-1320
- Green, M. R. et al., Molecular Cloning, A Laboratory Manual. **4th** (2012)
- Greenfield, E. A., Antibodies: A Laboratory Manual. **2nd** (2014)
- Gustafsson, C. et al., Trends Biotechnol. **22** (2004): 346-353
- Hwang, M. L. et al., J Immunol. **179** (2007): 5829-5838

- 153 -

- Jung, G. et al., Proc Natl Acad Sci U S A. **84** (1987): 4611-4615
- Kibbe, A. H., Handbook of Pharmaceutical Excipients. **rd** (2000)
- Krieg, A. M., Nat Rev Drug Discov. **5** (2006): 471-484
- Kuball, J. et al., Blood. **109** (2007): 2331-2338
- Liddy, N. et al., Nat Med. **18** (2012): 980-987
- Ljunggren, H. G. et al., J Exp Med. **162** (1985): 1745-1759
- Longenecker, B. M. et al., Ann N Y Acad Sci. **690** (1993): 276-291
- Lukas, T. J. et al., Proc Natl Acad Sci U S A. **78** (1981): 2791-2795
- Lundblad, R. L., Chemical Reagents for Protein Modification. **3rd** (2004)
- Meziere, C. et al., J Immunol. **159** (1997): 3230-3237
- Morgan, R. A. et al., Science. **314** (2006): 126-129
- Mortara, L. et al., Clin Cancer Res. **12** (2006): 3435-3443
- Mueller, L. N. et al., J Proteome Res. **7** (2008): 51-61
- Mueller, L. N. et al., Proteomics. **7** (2007): 3470-3480
- Mumberg, D. et al., Proc Natl Acad Sci U S A. **96** (1999): 8633-8638
- Pinheiro, J. et al., nlme: Linear and Nonlinear Mixed Effects Models (<http://CRANR-project.org/package=nlme>) (2015)
- Plebanski, M. et al., Eur J Immunol. **25** (1995): 1783-1787
- Porta, C. et al., Virology. **202** (1994): 949-955
- Rammensee, H. et al., Immunogenetics. **50** (1999): 213-219
- Rini, B. I. et al., Cancer. **107** (2006): 67-74
- Rock, K. L. et al., Science. **249** (1990): 918-921
- Rodenko, B. et al., Nat Protoc. **1** (2006): 1120-1132
- Saiki, R. K. et al., Science. **239** (1988): 487-491
- Schmitt, T. M. et al., Hum Gene Ther. **20** (2009): 1240-1248
- Scholten, K. B. et al., Clin Immunol. **119** (2006): 135-145
- Seeger, F. H. et al., Immunogenetics. **49** (1999): 571-576
- Sherman, F. et al., Laboratory Course Manual for Methods in Yeast Genetics.(1986)
- Singh-Jasuja, H. et al., Cancer Immunol Immunother. **53** (2004): 187-195
- Small, E. J. et al., J Clin Oncol. **24** (2006): 3089-3094
- Sturm, M. et al., BMC Bioinformatics. **9** (2008): 163

- 154 -

- Teufel, R. et al., Cell Mol Life Sci. **62** (2005): 1755-1762
Tran, T. T. et al., Photochem Photobiol. **90** (2014): 1136-1143
Walter, S. et al., J Immunol. **171** (2003): 4974-4978
Walter, S. et al., Nat Med. **18** (2012): 1254-1261
Willcox, B. E. et al., Protein Sci. **8** (1999): 2418-2423
Zaremba, S. et al., Cancer Res. **57** (1997): 4570-4577
Zufferey, R. et al., J Virol. **73** (1999): 2886-2892

- 155 -

I33190WO

Claims

1. A peptide comprising an amino acid sequence selected from the group consisting of SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464, and variant sequences thereof which are at least 88% homologous to SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464, and wherein said variant binds to molecule(s) of the major histocompatibility complex (MHC) and/or induces T cells cross-reacting with said variant peptide; and a pharmaceutical acceptable salt thereof, wherein said peptide is not a full-length polypeptide.
2. The peptide according to claim 1, wherein said peptide has the ability to bind to an MHC class-I or –II molecule, and wherein said peptide, when bound to said MHC, is capable of being recognized by CD4 and/or CD8 T cells.
3. The peptide or variant thereof according to claim 1 or 2, wherein the amino acid sequence thereof comprises a continuous stretch of amino acids according to any one of SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464.
4. The peptide or variant thereof according to any of claims 1 to 3, wherein said peptide or variant thereof has an overall length of from 8 to 100, preferably from 8 to 30, and more preferred from 8 to 16 amino acids, and most preferred wherein the peptide consists or consists essentially of an amino acid sequence according to any one of SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464.
5. The peptide or variant thereof according to any one of Claims 1 to 4, wherein said peptide is modified and/or includes non-peptide bonds.
6. The peptide or variant thereof according to any one of Claims 1 to 5, wherein said peptide is part of a fusion protein, in particular comprising N-terminal amino acids of the HLA-DR antigen-associated invariant chain (Ii).

- 156 -

7. An antibody, in particular a soluble or membrane-bound antibody, preferably a monoclonal antibody or fragment thereof, that specifically recognizes the peptide or variant thereof according to any one of claims 1 to 5, preferably the peptide or variant thereof according to any one of claims 1 to 5 when bound to an MHC molecule.

8. A T-cell receptor, preferably soluble or membrane-bound, or a fragment thereof, that is reactive with an HLA ligand, wherein said ligand is the peptide or variant thereof according to any one of claims 1 to 5, preferably the peptide or variant thereof according to any one of claims 1 to 5 when bound to an MHC molecule.

9. The T-cell receptor according to claim 8, wherein said ligand amino acid sequence is at least 88% identical to any one of SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464, or wherein said ligand amino acid sequence consists of any one of SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464.

10. The T-cell receptor according to claim 8 or 9, wherein said T-cell receptor is provided as a soluble molecule and optionally carries a further effector function such as an immune stimulating domain or toxin.

11. An aptamer that specifically recognizes the peptide or variant thereof according to any one of claims 1 to 5, preferably the peptide or variant thereof according to any one of claims 1 to 5 that is bound to an MHC molecule.

12. A nucleic acid, encoding for a peptide or variant thereof according to any one of claims 1 to 5, an antibody or fragment thereof according to claim 7, a T-cell receptor or fragment thereof according to claim 8 or 9, optionally linked to a heterologous promoter sequence, or an expression vector expressing said nucleic acid.

13. A recombinant host cell comprising the peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9 or the nucleic acid or the expression vector according to claim 12, wherein said host cell preferably is selected from an antigen presenting cell, such as a dendritic cell, a T cell or an NK cell.

- 157 -

14. An *in vitro* method for producing activated T lymphocytes, the method comprising contacting *in vitro* T cells with antigen loaded human class I or II MHC molecules expressed on the surface of a suitable antigen-presenting cell or an artificial construct mimicking an antigen-presenting cell for a period of time sufficient to activate said T cells in an antigen specific manner, wherein said antigen is a peptide according to any one of claims 1 to 4.

15. An activated T lymphocyte, produced by the method according to claim 14, that selectively recognizes a cell which presents a polypeptide comprising an amino acid sequence given in any one of claims 1 to 4.

16. A pharmaceutical composition comprising at least one active ingredient selected from the group consisting of the peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector according to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15, or a conjugated or labelled active ingredient, and a pharmaceutically acceptable carrier, and optionally, pharmaceutically acceptable excipients and/or stabilizers.

17. A method for producing the peptide or variant thereof according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, or the T-cell receptor or fragment thereof according to claim 8 or 9, the method comprising culturing the host cell according to claim 13, and isolating the peptide or variant thereof, the antibody or fragment thereof or the T cell receptor or fragment thereof from said host cell and/or its culture medium.

18. The peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector according

- 158 -

to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15 for use in medicine.

19. A method for killing target cells in a patient which target cells present a polypeptide comprising an amino acid sequence given in any one of claims 1 to 4, the method comprising administering to the patient an effective number of activated T cells as defined in claim 15.

20. The peptide according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector according to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15 for use in diagnosis and/or treatment of cancer, or for use in the manufacture of a medicament against cancer.

21. The use according to claim 20, wherein said cancer is selected from the group of acute myeloid leukemia, breast cancer, cholangiocellular carcinoma, chronic lymphocytic leukemia, colorectal cancer, gallbladder cancer, glioblastoma, gastric cancer, hepatocellular carcinoma, head and neck squamous cell carcinoma, melanoma, non-hodgkin lymphoma, lung cancer (including non-small cell lung cancer adenocarcinoma, squamous cell non-small cell lung cancer, and small cell lung cancer), ovarian cancer, esophageal cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, urinary bladder carcinoma, uterine and endometrial cancer and other tumors that show an overexpression of a protein from which a peptide SEQ ID No. 1 to SEQ ID No. 387 and SEQ ID No. 463 to SEQ ID No. 464 is derived from.

22. A kit comprising:

(a) a container comprising a pharmaceutical composition containing the peptide(s) or the variant according to any one of claims 1 to 6, the antibody or fragment thereof according to claim 7, the T-cell receptor or fragment thereof according to claim 8 or 9, the aptamer according to claim 11, the nucleic acid or the expression vector according

- 159 -

to claim 12, the host cell according to claim 13, or the activated T lymphocyte according to claim 15, in solution or in lyophilized form;

(b) optionally, a second container containing a diluent or reconstituting solution for the lyophilized formulation;

(c) optionally, at least one more peptide selected from the group consisting of SEQ ID No. 1 to SEQ ID No. 460 and SEQ ID No. 463 to SEQ ID No. 464, and

(d) optionally, instructions for (i) use of the solution or (ii) reconstitution and/or use of the lyophilized formulation.

23. The kit according to claim 22, further comprising one or more of (iii) a buffer, (iv) a diluent, (v) a filter, (vi) a needle, or (v) a syringe.

24. A method for producing a personalized anti-cancer vaccine or a compound-based and/or cellular therapy for an individual patient, said method comprising:

a) identifying tumor-associated peptides (TUMAPs) presented by a tumor sample from said individual patient;

b) comparing the peptides as identified in a) with a warehouse of peptides that have been pre-screened for immunogenicity and/or over-presentation in tumors as compared to normal tissues;

c) selecting at least one peptide from the warehouse that matches a TUMAP identified in the patient; and

d) manufacturing and/or formulating the personalized vaccine or compound-based or cellular therapy based on step c).

25. The method according to claim 24, wherein said TUMAPs are identified by:

a1) comparing expression data from the tumor sample to expression data from a sample of normal tissue corresponding to the tissue type of the tumor sample to identify proteins that are over-expressed or aberrantly expressed in the tumor sample; and

a2) correlating the expression data with sequences of MHC ligands bound to MHC class I and/or class II molecules in the tumor sample to identify MHC ligands derived from proteins over-expressed or aberrantly expressed by the tumor.

- 160 -

26. The method according to claim 24 or 25, wherein the sequences of MHC ligands are identified by eluting bound peptides from MHC molecules isolated from the tumor sample, and sequencing the eluted ligands.

27. The method according to any one of claims 24 to 26, wherein the normal tissue corresponding to the tissue type of the tumor sample is obtained from the same patient.

28. The method according to any one of claims 24 to 27, wherein the peptides included in the warehouse are identified based on the following steps:

aa. Performing genome-wide messenger ribonucleic acid (mRNA) expression analysis by highly parallel methods, such as microarrays or sequencing-based expression profiling, comprising identify genes that over-expressed in a malignant tissue, compared with a normal tissue or tissues;

ab. Selecting peptides encoded by selectively expressed or over-expressed genes as detected in step aa, and

ac. Determining an induction of *in vivo* T-cell responses by the peptides as selected comprising *in vitro* immunogenicity assays using human T cells from healthy donors or said patient; or

ba. Identifying HLA ligands from said tumor sample using mass spectrometry;

bb. Performing genome-wide messenger ribonucleic acid (mRNA) expression analysis by highly parallel methods, such as microarrays or sequencing-based expression profiling, comprising identify genes that over-expressed in a malignant tissue, compared with a normal tissue or tissues;

bc. Comparing the identified HLA ligands to said gene expression data;

bd. Selecting peptides encoded by selectively expressed or over-expressed genes as detected in step bc;

be. Re-detecting of selected TUMAPs from step bd on tumor tissue and lack of or infrequent detection on healthy tissues and confirming the relevance of over-expression at the mRNA level; and

- 161 -

bf. Determining an induction of *in vivo* T-cell responses by the peptides as selected comprising *in vitro* immunogenicity assays using human T cells from healthy donors or said patient.

29. The method according to any one of claims 24 to 28, wherein the immunogenicity of the peptides included in the warehouse is determined by a method comprising *in vitro* immunogenicity assays, patient immunomonitoring for individual HLA binding, MHC multimer staining, ELISPOT assays and/or intracellular cytokine staining.

30. The method according to any one of claims 24 to 29, wherein said warehouse comprises a plurality of peptides selected from the group consisting of SEQ ID No. 1 to SEQ ID No. 460 and SEQ ID No. 463 to SEQ ID No. 464.

31. The method according to any one of claims 24 to 30, further comprising identifying at least one mutation that is unique to the tumor sample relative to normal corresponding tissue from the individual patient, and selecting a peptide that correlates with the mutation for inclusion in the vaccine or for the generation of cellular therapies.

32. The method according to claim 31, wherein said at least one mutation is identified by whole genome sequencing.

Figure 1A

Peptide: FMVIAGMPLF (A*24)
SEQ ID NO: 15




Figure 1B
 Peptide: RYSPVKDAW (A*24)
 SEQ ID NO: 60




Figure 1C

Peptide: NYVLVPTMF (A*24)
 SEQ ID NO: 74




Figure 1D

Peptide: SYLPTAERL (A*24)
SEQ ID NO: 86






Figure 1E

Peptide: VYDTMIEKFA (A*24)
 SEQ ID NO: 202



6/31

Figure 1F
 Peptide: EYSLPVLTF (A*24)
 SEQ ID NO: 274



7/31

Figure 1G

Peptide: NYMDTDNLMF (A*24)
SEQ ID NO: 362






Figure 1H
 Peptide: YQSRDYNF (A*24)
 SEQ ID NO: 386



9/31


Figure 11

Peptide: VGGNVTSNF (A*24)
 SEQ ID NO: 463



10/31


Figure 1J
 Peptide: VGGNVTSSF (A*24)
 SEQ ID NO: 464



11/31

Figure 2A

Gene: MAGEA4
Peptide: IFPKTGLLI
SEQ ID NO: 1



12/31

Figure 2B

Gene: TRPM8
Peptide: KFLTHDVLTELF
SEQ ID NO: 3





Figure 2C


Gene: CHRNA9
Peptide: KYIATMAL
SEQ ID NO: 13



14/31

Figure 2D


Gene: MMP12
Peptide: KYVDINTFRL
SEQ ID NO: 18



15/31

Figure 2E

Gene: SPINK2
Peptide: LYMRFVNTHF
SEQ ID NO: 21



16/31

Figure 2F

Gene: OR51E2
 Peptide: FWFDSREISF
 SEQ ID NO: 28




Figure 2G

Gene: MMP1
 Peptide: KQMQEFFGL
 SEQ ID NO: 31





Figure 2H


Gene: MAGEC1
Peptide: FSSTLVSLF
SEQ ID NO: 35



19/31

Figure 2I

Gene: ENPP3
Peptide: KTYLPTFFETI
SEQ ID NO: 39




bioRxiv preprint doi: <https://doi.org/10.1101/2020.10.19.351111>; this version posted October 19, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

20/31

Figure 2J

Gene: POTEH, POTEH
Peptide: MVLQPQPQLF
SEQ ID NO: 4



21/31

Figure 2K

Gene: OR51E2
Peptide: TQMFFIHAL
SEQ ID NO: 97




Figure 2L

Gene: MMP11
Peptide: FFFKAGFVWR
SEQ ID NO: 107





Figure 2M

Gene: MMP11
Peptide: YFLRGRLYW
SEQ ID NO: 122



24/31

Figure 2N

Gene: SLC24A5

Peptide: EYFLPSLEII

SEQ ID NO: 194





Figure 20

Gene: SLC24A5
 Peptide: MSAIWISAF
 SEQ ID NO: 277



27/31

Figure 2Q

Gene: LAMA3
Peptide: YFGNPQKF
SEQ ID NO: 304




Figure 3




Figure 3 (continued)




Figure 3 (continued)





Figure 3 (continued)

