

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 5/00, 5/02	A1	(11) International Publication Number: WO 96/24660 (43) International Publication Date: 15 August 1996 (15.08.96)
(21) International Application Number: PCT/US96/01744 (22) International Filing Date: 8 February 1996 (08.02.96) (30) Priority Data: 08/385,772 9 February 1995 (09.02.95) US (71) Applicant: NOVAFLORA, INC. [US/US]; Suite 350, 3401 Market Street, Philadelphia, PA 19104 (US). (72) Inventors: DOBRES, Michael, S.; 1905 Rodman Street, Philadelphia, PA 19146 (US). WILLIAMS LOKAJ, Gail, R.; 8 North Lincoln Avenue, Wenonah, NJ 08090 (US). (74) Agents: MILLER, Suzanne, E. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KG, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: MICROPROPAGATION OF ROSE PLANTS (57) Abstract The present invention is directed to, <i>inter alia</i> , a method of micropropagation of a rose plant comprising culturing a stem bearing a node in a first enclosed vessel containing a first culture medium comprising about 0.5 to about 3.0 mg/l of a cytokine, 0 to about 1.0 mg/l of an auxin, and 0 to about 1.0 mg/l of a gibberellin until at least one shoot emerges from said node; excising said shoot from said stem and propagating said shoot in a second enclosed vessel containing a second culture medium comprising about 0.5 to about 3.0 mg/l of a cytokine, 0 to about 1.0 mg/l of an auxin, and 0 to about 1.0 mg/l of a gibberellin to produce a flowering rose plant capable of being transferred to soil. Among other things, the present invention is also directed to rose plants and culture media for the micropropagation of roses.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

MICROPROPAGATION OF ROSE PLANTS

BACKGROUND OF THE INVENTION

Micropropagation, also known as axillary bud propagation or in vitro asexual propagation, is one of a number of methods known in the art for the propagation of desirable species or varieties of plants. Other traditional methods of asexual propagation include greenhouse and field propagation of plant cuttings. There are many advantages, however, which favor micropropagation as a propagative method of choice. One advantage is that a plant which has a known and desirable phenotype can be chosen as the source of cells, and, in accordance with micropropagation techniques, these cells can be rapidly cultured into many genetically uniform plantlets. The resulting plantlets may then be cultivated into entire plants possessing roots and shoots. Thus, in accordance with this technique, plants with the same desirable phenotype as the parent can be mass produced, potentially at costs comparable to and often more quickly than standard greenhouse or field methods of asexual propagation.

The propagation of plants under sterile aseptic conditions has been used as cost effective means for the

- 2 -

mass propagation of certain plants. An important consideration of this technology is the ability to add value to plants being propagated. In many cases value is added by assuring uniformity of product, and products that are free
5 of bacterial, fungal, viral, or insect pests.

Commercial propagation of roses is normally carried out by budding or grafting of a specific shoot cutting, or scion, onto an established wild rose root stock. In recent years numerous publications have dealt with the
10 propagation of roses through tissue culture. Examination of the literature reveals that different varieties and types display greatly varying nutrient and hormone requirements. Indeed, the process of producing plants from explants or plant tissues, however, requires the development of a
15 culture medium and protocol suitable for the particular type of plant tissue. Various media and procedures have been attempted, but only some have been met with success. For this reason, most commercial growers of roses still propagate rose plants by traditional means.

For example, Hasegawa, *J. Amer. Soc. Hort. Sci.*
20 1980 105, 216-220, the disclosures of which are hereby incorporated by reference in their entirety, reported that media developed for the climbing rose Improved Blaze allowed significant growth of three miniature rose varieties but not
25 for all hybrid tea roses. Khosh-Khui and Sink, *J. Hort. Sci.* 1982 57, 315-319 and Khosh-Khui and Sink, *Scientia Horticulturae* 1982 17, 371-376, the disclosures of which are hereby incorporated by reference in their entirety, found that plant growth regulator requirements for multiplying the
30 *R. hybrida* cultivars, Tropicana and Bridal Pink differed considerably from those of two old world species *R. damascena* and *R. canina*.

To optimize growth conditions most studies have focused on variations in the concentrations of certain
35 growth regulators, Hasegawa, *supra*. A few studies have investigated the effect of certain other inorganic or organic components. For example, Khosh-Khui and Sink,

- 3 -

supra. investigated the effect of explant orientation, explant size and lighting on the growth of new and old world species. Bressan et al., *J. Amer. Soc. Hort. Sci.* 1982 107, 979-990, the disclosures of which are hereby incorporated by
5 reference in their entirety, investigated the effect of bud position, temperature, hours of light per day, Murashige and Skoog ingredient concentrations, and weeks in culture between transfers.

Despite some success in developing suitable
10 culture media and culturing processes, new and/or better media and methods, particularly for commercially significant roses, are needed. Toward this end, the present invention is directed to the development of media useful for the propagation of roses and for induction of flowering in
15 vitro.

SUMMARY OF THE INVENTION

The present invention is directed to a method of tissue culture micropropagation of a rose plant comprising culturing a stem bearing a node in a first enclosed vessel
20 containing a first culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid until at least one shoot is produced from said node; excising said shoot from said stem and propagating said shoot in a second
25 enclosed vessel containing a second culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid to produce a rose plant in an enclosed vessel. In accordance with this method, the rose plant may
30 be a flowering rose and the rose plant may be capable of growth in soil.

The present invention is also directed to a method of tissue culture micropropagation of a rose plant comprising culturing a stem bearing a node in an enclosed
35 vessel containing culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole

- 4 -

acetic acid, and 0 to about 1.0 mg/l gibberellic acid. The stem may be cultured until at least one shoot is produced from the node or until a rose plant is produced. A method of tissue culture micropropagation of a rose plant
5 comprising culturing a shoot in an enclosed vessel containing culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid is also an embodiment of the present invention. The shoot may be
10 cultured until a rose plant is produced within the enclosed vessel. In each of these methods, the rose plant may be a flowering rose, and the rose plant may be capable of growth in soil.

A culture medium for tissue culture
15 micropropagation of a rose plant is also an embodiment of the present invention and comprises about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid. The culture medium may further comprise a nutritive medium. Preferably
20 the culture medium contains 3.0 mg/l benzyladenine, more preferably about 2.0 mg/l benzyladenine, and even more preferably about 1.0 mg/l benzyladenine.

The rose plant may also be subject to growth in rooting medium.

25 A rose plant, including a flowering rose, produced by the method is also within the scope of the present invention. The rose is preferably *Rose hybrida*, and most preferably a cultivar selected from the group consisting of Meirutral, Fairy, and Pink Miniwonder. The rose plant may
30 be contained within the enclosed vessel. The rose plant may also be capable of growth in soil.

The present invention is also directed to a rose tissue culture comprising a rose stem bearing a node contained in an enclosed vessel containing culture medium
35 comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid and 0 to about 1.0 mg/l gibberellic acid. The rose stem may comprise at least one

- 5 -

shoot, the rose tissue culture capable of producing a rose, for example, a flowering rose. A rose explant is also within the scope of the present invention. The rose explant may be selected from a rose stem bearing a node, a rose stem
5 bearing a node and at least one shoot, and a rose shoot. The rose explant may be contained in an enclosed vessel comprising culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid and 0 to about 1.0 mg/l gibberellic acid, and may be
10 cultured until a rose plant is produced, a flowering rose plant, for example.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts full view (Figure 1A) and close up view (Figure 1B) of *Rosa hybrida* cv. Meirutral at an
15 early stage of flower opening *in vitro*.

Figure 2 shows full view (Figure 2A) and close up view (Figure 2B) of *Rosa hybrida* cv. Meirutral showing completely opened flower *in vitro*.

Figure 3 depicts three stages of flower
20 development of *Rosa hybrida* cv. Pink Miniwonder *in vitro*. Figure 3A shows floral bud with sepals enclosing petals. Figure 3B exhibits floral bud with retracted petals revealing pink colored petals. Figure 3C shows a fully opened flower. All plants have a well developed root system
25 allowing for transplanting to soil for indoor or outdoor growth.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed, *inter alia*, to a flowering rose plant which may remain in culture without
30 feeding or watering for a period of time which permits the plant to continue to be sustained. Once the rose flowers in the enclosed vessel, it typically may be maintained for at least about three (3) weeks, more preferably for at least about four (4), more preferably for at least about six (6),
35 even more preferably for at least about eight (8) weeks, and

- 6 -

most preferably for at least about twelve (12) weeks.

Alternatively, the flowering rose plant may be removed from the enclosed vessel and be transferred to soil for use as an indoor or outdoor plant.

5 In accordance with the present invention, micropropagation is defined as replicating a parent plant such that the resulting plants are identical to the parent plant. Micropropagation is carried out with rose plant tissue (a rose explant) obtained from a rose plant. In
10 accordance with the present invention, a rose plant is defined herein as a shoot having no roots (wherein said shoot may be induced to root), and a shoot having a root system. Each type of rose plant, with or without root
15 system may bear flowers in accordance with the methods of the present invention. The explant may be mature or immature, preferably immature. The explant may be at least one cell from a rose plant, a rose stem, preferably a rose stem bearing at least one node, a rose stem bearing a node and at least one shoot, a rose shoot, and a shoot having
20 roots. A rose stem is a cutting of a rose plant from the stem of the plant. A rose stem may be cut from any where along the length of the rose plant stem and generally is of a length of about one quarter of an inch ($\frac{1}{4}$) to about three (3) inches, preferably about one inch to about one and one
25 half inches. It has been found that variation in explant source may effect the performance of the explant in the enclosed vessel, such that rose stems obtained from the area of the stem closer to the shoot tip tends to perform more favorably in the methods of the present invention. A node
30 is an area on a stem which develops into a shoot. Similarly, a rose shoot generally is of a length of about one quarter of an inch ($\frac{1}{4}$) to about six (6) inches, preferably about one inch to about one and one half inches.

 The present invention is directed to a method of
35 tissue culture micropropagation of a rose plant comprising culturing a stem bearing a node in a first enclosed vessel containing a first culture medium comprising about 0.5 to

- 7 -

about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid and 0 to about 1.0 mg/l gibberellic acid until at least one shoot is produced from the node; excising the shoot from the stem and propagating the shoot in a second enclosed vessel containing a second culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid and 0 to about 1.0 mg/l gibberellic acid to produce a flowering rose plant in an enclosed vessel capable of being further propagated *in vitro* and/or transferred to soil.

The time needed to culture the stem to a flowering rose plant *in vitro*, is a time effective to produce a flowering rose plant in accordance with the methods of the present invention, typically about three (3) to about six (6) weeks, preferably about four (4) weeks. The second enclosed vessel may be the same as the first vessel, or a different vessel.

The present invention is also directed to each individual step of the two steps of the method of tissue culture micropropagation set forth above. A rose plant is propagated by culturing a stem bearing at least one node in an enclosed vessel comprising the aforementioned culture medium. The stem may be cultured until at least one shoot is produced from the node. Alternatively the stem may be cultured for a time effective to produce a rose plant, typically about three (3) to about six (6) weeks, preferably about four (4) weeks. The shoot produced from the node of the above- identified method, or a shoot obtained as an explant from a rose plant, may be used in a method of tissue culture micropropagation comprising culturing the shoot in an enclosed vessel comprising the aforementioned culture medium. The shoot may be cultured for a time effective to produce a rose plant, typically about three (3) to about six (6) weeks, preferably about four (4) weeks. The rose plant may be produced within the enclosed vessel and may be a flowering rose. Further the rose plant may be capable of

- 8 -

growth in soil such that the rose plant may be potted for indoor or outside use.

Preferably, the method present invention is carried out a medium comprising 3 mg/l benzyladenine, more preferably 2.0 mg/l benzyladenine, and even more preferably 1.0 mg/l benzyladenine while the concentration of indole acetic acid is about 0.3 mg/l and the concentration of gibberellic acid is about 0.01 to about 0.5 mg/l, more preferably about 0.1 mg/l. The concentration of benzyladenine is at least about 0.5 mg/l, while indole acetic acid and gibberellic acid may not be included, however, are preferably included at 0 to about 1.0 mg/l each.

Benzyladenine, a cytokine also known as 6-benzylaminopurine, may be substituted with another natural or synthetic cytokine selected from the group consisting of 6-benzylaminopurine riboside; 1-phenyl-3-(1,2,3-thiadiazol-5-yl) urea; 6-(γ - γ -dimethylallylamino)purine; DL-Dihydrozeatin; t-zeatin riboside; zeatin; N-(2-Chloro-4-pyridyl)-N'-phenylurea; N-benzyl-9-(2-tetrahydropyranyl)-adenine; kinetin; kinetin riboside; and isopentyladenosine, wherein the cytokine may be used alone or in combination with one or more cytokine. Indole acetic acid, an auxin, may be substituted with another natural or synthetic auxin selected from the group consisting of naphthaleneacetic acid (NAA), indole butyric acid (IBA), picloram, dicamba, and 2,4-dichlorophenoxy acetic acid (2,4-D), at a concentration of about 0.03 to about 0.3 mg/l, wherein the auxin may be used alone or in combination with one or more auxin. Gibberellins include various forms of gibberellic acid, such as and not limited to GA3, GA4, GA7, and iso-gibberellin A7.

As those skilled in the art would recognize, the general nutritional and growth requirements of tissue cultured plant cells must be satisfied. The general nutritional, hormonal, and growth requirements of plant cells are well known, and a number of conventional culture media and growth protocols have been developed which satisfy

- 9 -

these needs (hereinafter referred to herein as nutritive medium). See, e.g., MSG medium (Becwar, M.R., et al., "Development and Characterization of In Vitro Embryogenic Systems in Conifers", in Somatic Cell Genetics of Woody Plants, ed. Ahuja, M.R., Kluwer Academic Publishing, Dordrecht, The Netherlands (1988)), Shenk-Hildebrandt (SH) culture medium (Shenk et al., Can. J. Bot., Vol. 50, pp. 199-204 (1972)), Murashige-Skoog (M-S) basal media (Murashige et al., Physiol. Plant., Vol. 15, pp. 473-97 (1962)), and White's medium (White, The Cultivation of Animal and Plant Cells, 2nd ed., Ronald Press Co., New York (1963)). A comprehensive list of plant culture media and culture protocol are found in Huang et al., Plant Tissue Culture Media, TCA Manual, Vol. 3, pp. 539-48, Tissue Culture Association, Rockville, MD (1977). The disclosures of each of these references are incorporated herein by reference in their entirety. The foregoing culture media and culture protocol, as well as others known to those skilled in the art, can be employed in conjunction with the methods and medium of the present invention. In accordance with the invention, however, sucrose, or other carbon sources including and not limited to glucose, sorbitol, mannitol, fructose, and galactose, must necessarily be present, and indoleacetic acid preferably present, each in the amounts set forth herein, for the efficient and effective production of flowering rose plants from the rose explant.

Within this general framework, and in addition to the aforementioned components benzyladenine, indole acetic acid and gibberellic acid, the culture medium of the present invention may further comprise a nutritive medium. Preferably the medium is Murashige and Skoog (MS) medium. MS medium comprises salts such as and not limited to at least about 400 mg/l, preferably about 1650 mg/l ammonium nitrate, at least about 1 mg/l, preferably about 6.2 mg/l boric acid, at least about 75 mg/l, preferably about 332 mg/l, more preferably 332.2 mg/l calcium chloride

- 10 -

(anhydrous), at least about 0.005 mg/l, preferably about 0.025 mg/l cobalt chloride.6H₂O, at least about 0.005, preferably about 0.025 mg/l cupric sulfate.5H₂O, at least about 3 mg/l, preferably about 37 mg/l, more preferably about 37.26 mg/l sodium ethylene diamine tetraacetic acid (Na₂.EDTA), at least about 6 mg/l, preferably about 27-28 mg/l, more preferably about 27.8 mg/l ferrous sulfate.7H₂O, at least about 45 mg/l, preferably about 180 mg/l, more preferably about 180.7 mg/l magnesium sulfate, at least about 4 mg/l, preferably about 16-17 mg/l, more preferably about 16.9 mg/l manganese sulfate.H₂O, at least about 0.05 mg/l, preferably about 0.25 mg/l molybdic acid (sodium salt).2H₂O, at least about 0.2 mg/l, preferably about 0.83 mg/l potassium iodide, at least about 500 mg/l, preferably about 1900 mg/l potassium nitrate, about 170 mg/l potassium phosphate (monobasic), and at least about 2 mg/l, preferably about 8.6 mg/l zinc sulfate.7H₂O. The culture medium may further comprise Gamborg's B5 vitamins comprising about 10 to about 200 mg/l, preferably about 100 mg/l myo-inositol, at least about 0.5 to about 5 mg/l, preferably about 1.0 mg/l nicotinic acid, at least about 0.5 to about 5.0 mg/l, preferably about 1.0 mg/l pyridoxine.HCl, and at least about 5 to about 20 mg/l, preferably about 10 mg/l thiamine.HCl. These vitamins may also be used singularly or in various combinations of two or more vitamins. Further, other vitamins and vitamin combinations may be substituted therefor.

The preferred composition of the culture medium of the present invention comprises MS medium containing Gamborg's B5 vitamins, at least about 0.5 mg/l, preferably 2 mg/l glycine, at least about 10 mg/l, preferably 50 mg/l ascorbic acid, at least about 10 mg/l, preferably 50 mg/l citric acid, at least about 10 g/l, preferably about 30 g/l sucrose, and at least about 1.5 g/l, preferably about 3 g/l PHYTAGEL™. This medium is added to about 0.5 to about 3.0 mg/l benzyladenine, preferably 3 mg/l, more preferably about 2.0 mg/l, and even more preferably about 1.0 mg/l; 0 to

- 11 -

about 1.0 mg/l, preferably 0.3 mg/l indole acetic acid and 0 to about 1.0 mg/l, preferably 0.1 mg/l gibberellic acid.

The minimal requirements for shoot growth include the aforementioned basic salts, such as those of MS medium, 5 about 0.1 to about 0.5 mg/l cytokine, such as and not limited to benzyladenine, and about 5 to about 10 g/l of a carbon source such as and not limited to sucrose.

The media may be modified such that hormonal constituents are provided for specific varieties of plants. 10 Nutrient levels may be modified in the basal media to induce flowering. For example, Murashige and Skoog (MS) medium, which provides the general nutritional and growth requirements of tissue culture plant cells, may be substituted with other conventional culture or growth medium 15 such as and not limited to those disclosed by Becwar, M.R., et al., "Development and Characterization of In Vitro Embryogenic Systems in Conifers", in Somatic Cell Genetics of Woody Plants, ed. Ahuja, M.R., Kluwer Academic Publishing, Dordrecht, The Netherlands (1988), Shenk- 20 Hildebrandt (SH) culture medium Shenk et al., Can. J. Bot., Vol. 50, pp. 199-204 (1972), White, The Cultivation of Animal and Plant Cells, 2nd ed., Ronald Press Co., New York (1963) and Gamborg et al., Exp. Cell Res. 1968, 50, 151-158, the disclosure of each reference cited herein is 25 incorporated herein by reference in its entirety.

Within these general parameters, the precise amount of each component to be employed will depend upon various factors, such as the type of rose to be cultured, as will be readily apparent to those skilled in the art. 30 Typically, the compounds are initially employed at the lower levels, and the amounts increased as necessary to achieve the desired effect. For example, a preferred composition of MS medium containing Gamborg's B5 vitamins, about 0.5 to about 5.0 mg/l glycine, about 10 to about 100 mg/l ascorbic 35 acid, about 10 to about 100 mg/l citric acid, about 10 to about 30 g/l, preferably 30 g/l sucrose, and about 1.5 to

- 12 -

about 3.0 g/l, preferably 3 g/l Phytigel™ may be used in the present invention.

Glycine may be used at concentrations of about 0.5 to about 5.0 mg/l, and may be substituted by other amino acids, such as and not limited to alanine and proline; mixture of naturally occurring amino acids, at a similar concentration range; crude mixtures such as casein hydrolysate at concentration of about 0.5 to about 10 g/l; coconut water at a concentration of between about 1 to about 20 g/l. Ascorbic acid and citric acid may be used in the range of about 10 to about 100 mg/l. Gibberellic acid may be used in the range of about 0.01 to about 0.5 mg/l. Any suitable gelling agent such as and not limited to PHYTAGEL™ comprising gellan gum, GEL-RITE™ comprising gellan gum; AGARGEL™ comprising gellan gum; agar such as and not limited to agar types A, E, and M, High Gel Strength, Purified, Bacteriological Flake, and Washed; AGARGEL™ a blend of agar and PHYTAGEL™; agarose, such as and not limited to Type VII; alginic acid; carrageenan; transfergel or hydroxyethylcellulose; and the like. GEL-RITE™ is used at half the concentration recommended by the present invention for the other gelling agents. One or more of the ingredients of the culture medium may be omitted, without substitution therefor, including and not limited to gibberellic acid, glycine, ascorbic acid, citric acid, and/or indole acetic acid.

In addition to alternative medium compositions, success of the micropropagation technique may be enhanced by additional conditions as well. Such conditions will be readily apparent to those skilled in the art and include choosing donor tissues for culture initiation which come from plants of appropriate genotype as well as physiological and development states, variation of explant source and the plant variety, and physical environment in which the cultures are grown. For example, gaseous conditions may be modified within the enclosed vessel to allow ethylene and air to be exchanged. Temperature and light conditions may

- 13 -

be varied to provide for an improved rose plant contained in an enclosed vessel.

Further to the methods and culture medium set forth above, the present invention is also directed to propagating a shoot in a third enclosed vessel containing a rooting medium comprising nutritive medium. The nutritive medium contained in the rooting medium is preferably MS medium. However, any of the aforementioned nutritive media may be useful in the rooting medium as well. In addition to nutritive medium, the rooting medium also comprises about 1 to about 5 mg/l, preferably about 2.0 mg/l glycine, about 50 to about 150 mg/l, preferably about 100 mg/l myo-inositol, about 0.1 to about 1.0 mg/l, preferably about 0.5 mg/l nicotinic acid, about 0.1 to about 1.0 mg/l, preferably about 0.5 mg/l pyridoxine.HCl, about 0.01 to about 0.2 mg/l, preferably about 0.1 mg/l thiamine.HCl, about 10 to about 50 g/l, preferably about 30 g/l sucrose, and optionally containing about 1 to about 5 g/l, preferably about 3 g/l gelling agent, about 0.05 to about 1 mg/l, preferably about 1 mg/l, more preferably about 0.05 mg/l indole butyric acid. The gelling agent may be any of the aforementioned gelling agents, as well as water, sterilized soil, cellulose foam or fiber rooting materials such as and not limited to Sorbarod™. The rooting medium may even further comprise about 0.01 to about 0.2 mg/l, preferably about 0.1 mg/l naphthaleneacetic acid. At a minimum, rooting medium may simply comprise 1/4X MS salts. In addition about 0.1 mg/l of an auxin such as and not limited to indole acetic acid may be added to the 1/4X MS salts. Preferred rooting media of the present invention include and are not limited to the following compositions: 0.5 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose and 3.0 g/l PHYTAGEL™, and 1 mg/l IBA; 0.25 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, 3.0 g/l PHYTAGEL™, and 1 mg/l IBA; 0.5 X MS salts, 2 mg/l

- 14 -

glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, 3.0 g/l PHYTAGEL, 0.05 mg/l IBA, and 0.1 mg/l NAA; 0.25 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l
5 nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, 3.0 g/l PHYTAGEL™, 0.05 mg/l IBA and 0.1 mg/l NAA; 0.25 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, and
10 3.0 g/l PHYTAGEL™; and 0.5 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose and 3.0 g/l PHYTAGEL™.

The rose explant, *per se*, is an embodiment of the
15 present invention. The rose explant may be contained in an enclosed vessel comprising the aforementioned culture medium, and may be cultured until a rose plant is produced, a flowering rose plant, for example. The explants may be obtained from a plant, such as and not limited to the genus
20 *Rosa*, including and not limited to the hybrid rose *Rosa hybrida*, such as and not limited to the cultivars Meirutral (Red Sun Blaze™), Fairy, Pink Miniwonder, Mr. Lincoln, Royalty, Chrysler Imperial, Frisco, Sonia, Peach and the like; *Rosa damascena*, *Rosa multiflora*, and *Rosa gallica*, and
25 the like.

The tissue culture techniques of the present invention comprise culturing a rose explant in an enclosed vessel such as and not limited to a petri dish, a test tube, a Magenta™ box, an eppendorf tube, a baby food jar, a
30 canning jar, and any other enclosed container capable of supporting the growth of the rose plant of the present invention in accordance with the methods of the present invention. The enclosed vessel comprises a culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to
35 about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid.

- 15 -

A rose, including a flowering rose, produced by the methods of the present invention, is also within the scope of the present invention. The rose may be *Rose hybrida*, and a cultivar selected from, and not limited to, the group consisting of Meirutral (Red Sun Blaze™), Fairy, and Pink Miniwonder. The rose plant may be contained within the enclosed vessel.

Rose tissue culture, *per se*, is within the scope of the present invention. A rose tissue culture comprising an explant, such as and not limited to a rose stem bearing at least one node, a rose stem bearing a node and at least one shoot, or a rose shoot, contained in an enclosed vessel containing culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid, is also an embodiment of the present invention. The rose tissue culture is capable of producing a rose, such as a flowering rose, and may be transferred to soil, for indoor or outside use.

The rose in an enclosed vessel may be used to screen new cultivars for flower color and form. For example, plants may be subjected to mutational treatment during tissue culture of rose callus, explants or axillary bud. Mutated rose plants may then be grown in the culture of the present invention and screened for those with altered flower color, form, disease resistance, growth vigor, cold and heat sensitivity, and any other valuable trait. In addition, the present invention also permits performing cross pollination of rose plants in culture. For example, since flowers contain visibly developed male and female organs, such flowers may be pollinated with pollen derived from the same flower or from a plant of any other source.

In the preferred method of the invention, rose stems are selected and excised in sections less than about 6 inches in length. Terminal buds and all leaves are removed from stems and each stem is cut into segments containing at

- 16 -

least one node. Excised stem segments are then washed with warm soap and water and rinsed with tap water. Stem segments are then surface sterilized which may be carried out in a laminar flow hood. Stem segments are placed in
5 ethanol and rinsed in sterile, distilled water and then swirled in sterile, distilled water. Stem segments may then be placed in a shaking bleach solution. Finally, stem segments are rinsed three times with sterile, distilled water, for about 3 to about 5 minutes/rinse. Stem segments
10 are then cultured by placing 2 to 3 segments in a sterile, disposable petri dish. The bleached ends of stem pieces are removed with a scalpel and the basal end (the end closest to the root in the parent plant) of one stem segment is placed, with node, approximately ¼ inch into culture medium, into a
15 test tube containing about 5 to about 50 ml, preferably about 20 ml of such medium. Cultures are grown under lights (about 16 hours daylength) on growth rack in culture room at about 24-25 °C.

After about 25 days, nodes typically give rise to
20 multiple new shoots. Each shoot is separately excised and transferred to fresh media. Approximately 17 days after this transfer, about 10% of the new shoots may bear a single floral bud about 3-5 mm in length. These buds then proceed to open.

25 The present invention is further described in the following examples. These examples are not to be construed as limiting the scope of the appended claims.

- 17 -

EXAMPLES**EXAMPLE 1**

A plant designated *Rosa hybrida* cv. Meirutral (U.S. Plant Patent 7,021, the disclosures of which are hereby incorporated by reference in their entirety) was obtained from the greenhouses of Conard-Pyle Co., West Chester, PA. About one month later, young, healthy stems were selected and excised as sections approximately 4-5 inches in length. Terminal buds and all leaves were removed from stems and each stem was cut into segments containing at least one node, 1 inch to 1½ inches in length. Excised stem segments were then washed with warm soap and water and rinsed well with tap water.

Stem segments were surface sterilized in a laminar flow hood by the following procedure. Stem segments were placed in 70% ethanol for about 2 minutes then rinsed in sterile, distilled water followed by swirling in sterile, distilled water for about 3-5 minutes. Stem segments were then placed in a 20% bleach solution (Chlorox, Proctor & Gamble, Cincinnati, OH) containing one drop (ca. 100 µl) of detergent for 20 minutes with shaking. Finally, stem segments were rinsed three times with sterile, distilled water (3-5 minutes/rinse.)

Stem segments were cultured by the following procedure. Segments (2-3 at a time) were placed in a sterile, disposable petri dish. The bleached ends of stem pieces were removed with a scalpel. The basal end of one stem segment was placed, with node, approximately ¼ inch into culture medium, into a test tube containing about 20 ml of such medium. Each culture was labelled with experiment number, stem segment number and date. Cultures were grown under lights (16 hours daylength) on growth rack in a culture room (24-25 °C). The effects of three different media were tested. MS basal media (Murashige 1962, the disclosures of which hereby incorporated by reference in their entirety) containing Gamborg's B5 vitamins was supplemented with glycine 2.0 mg/l, 50 mg/l ascorbic acid

- 18 -

and 50 mg/l citric acid, sucrose (30.0 g/l) PHYTAGEL™ (3.0 g/l) and adjusted to pH 5.7. The three variations tested further contained the growth regulators, benzyladenine (BA), indole acetic acid (IAA), and gibberellic acid (GA3). The
5 three media (BIG-4, BIG-5, and BIG-6) tested varied in their concentration of BA. BIG-4 contained 3 mg/l BA, 0.3 mg/l IAA and 0.1 mg/l GA3; BIG-5 contained 2 mg/l BA, 0.3 mg/l IAA, 0.1 mg/l GA3; BIG-6 contained 1 mg/l BA, 0.3 mg/l IAA and 0.1 mg/l GA3.

10 After 25 days, most nodes had given rise to multiple (2-4) new shoots. Each shoot was separately excised and transferred to fresh media (BIG 4, 5 or 6 respectively), see Table 1, row 2. Approximately 17 days after this transfer, approximately 10% of the new shoots
15 were seen to bear a single floral bud about 3-5 mm in length, see Table 1, row 5. In most cases, these buds proceeded to open. The first floral buds appeared on the largest shoots and produced flowers with small but otherwise normal looking petals, sepals, stamens and ovaries.

20 Individual plants were subsequently divided by cutting between nodes and subcultured on the same medium as the previous culture. At 47 days (from the time of the time the original culture was started) flowers were observed on 2 out of 38, 5 out of 33, and 6 out of 30 plants grown on BIG-
25 4, BIG-5, and BIG-6, respectively.

On the 76th day, plants were again divided by cutting between nodes and subcultured on the same medium as the previous culture. At day 88, 4 out of 46, 3 out of 26, and 6 out of 40 plants bore flowers in BIG-4, BIG-5, and
30 BIG-6 respectively.

Since the percentage of flower buds increased with decreasing BA concentration, it appears that lower concentrations of BA induce optimal flower formation.

In a similar experiment, *R. hybrida* cv. The Fairy,
35 formed flower buds which have opened. The ability to induce flower formation in culture appears to rely on a specific combination of media and genotype. In this case, *R. hybrida*

- 19 -

cv. Meirutral is particularly suited to the formation of flowers *in vitro* using the media tested.

TABLE 1

			BIG-4	BIG-5	BIG-6
5	Culture 1	NODES 102-0 10 days	17	22	14
	Culture 2	SHOOTS 102-1 25 days	23	25	17
	Culture 3	SHOOTS 102-2 45 days	38	33	30
	Culture 4	SHOOTS 102-3 76 DAYS	46	26	40
10	Culture 2	FLOWERS 42 days	2/23	3/25	2/17
	Culture 3	FLOWERS 47 days	2/38	5/33	6/30
	Culture 4	FLOWERS 88 days	4/46	3/26	6/40
	Culture 2	% FLOWERS 42 days	8.7	12.0	11.8
	Culture 3	% FLOWERS 47 days	5.3	15.2	20.0
	Culture 4	% FLOWERS 88 days	8.7	11.54	15

- 20 -

EXAMPLE 2

The following procedure was used to produce a plant designated *Rosa Hybrida* cv. Pink Miniwonder that flowered *in vitro*.

5 Stem segments were surface sterilized in a laminar flow hood by the following procedure. Stem segments were placed in 70% ethanol for 2 minutes then rinsed in sterile, distilled water followed by swirling in sterile, distilled water for 3-5 minutes. Stem segments were then placed in a
10 20% chlorox solution containing one drop (ca. 100 μ l) of detergent for 20 minutes with shaking. Finally, stem segments were rinsed 3 times with sterile, distilled water (3-5 minutes/rinse).

Stem segments were cultured by the following
15 procedure. Segments (2-3 at a time) were placed in a sterile, disposable petri dish. The bleached ends of stem pieces were removed with a scalpel. The basal end of one stem segment was placed, with node, approximately 1/4-inch into culture medium, into a test-tube containing 20 ml of
20 such medium. Each culture was labelled with experiment number, stem segment number and date. Cultures under lights (16 hour day length) on growth rack in culture room (24-25 °C).

The effects of two different media were tested.
25 The first media termed BIG-3 contained MS basal media (Murashige and Skoog, *Physiologia Plantarum* 1962, 15, 473-497, the disclosures of which are incorporated herein by reference in their entirety) containing Gamborg's B5 vitamins, Gamborg et al., *Exp. Cell Res.* 1968, 50, 151-158,
30 the disclosures of which are incorporated herein by reference in their entirety. The media was supplemented with glycine 2.0 mg/l, sucrose (30.0 g/l), PHYTAGEL™ (3.0 g/l), 3 mg/l BA, 0.3 mg/l IAA, and 0.1 mg/l GA3, and adjusted to pH 5.7. The second media termed BIG-4 in
35 addition to the above contained 50 mg/l ascorbic acid and 50 mg/l citric acid.

- 21 -

Plants were transferred every 2 to 4 weeks into new medium, identical to the medium from which the plant was transferred, for a total of 9 transfers for about 4 months before being transferred to four different kinds of rooting media.

Rooting I: 0.5 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose and 3.0 g/l PHYTAGEL™, and 1 mg/l IBA.

Rooting IA: 0.25 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, 3.0 g/l PHYTAGEL™, and 1 mg/l IBA.

Rooting II: 0.5 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, 3.0 g/l PHYTAGEL, 0.05 mg/l IBA, and 0.1 mg/l NAA.

Rooting IIA: 0.25 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, 3.0 g/l PHYTAGEL™, 0.05 mg/l IBA and 0.1 mg/l NAA.

Rooting 4: 0.25 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose, and 3.0 g/l PHYTAGEL™.

Rooting 5: 0.5 X MS salts, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.5 mg/l pyridoxine.HCl, 0.1 mg/l thiamine.HCl, 30 g/l sucrose and 3.0 g/l PHYTAGEL™.

After about three weeks, roots had developed in each type of rooting media. In addition, in rooting media I and IIA about 20% of the plants had developed flower buds. The induction of flowering in this case is due to the absence of BA in rooting media and/or the presence of auxins that might act to enhance ethylene biosynthesis and thereby stimulate flower production. In addition, the tendency of plants to flower may also be enhanced by the reduced MS salts concentrations. That is, flowering of rose plants in

- 22 -

vivo is known to be enhanced by reduced nutrients (fertilizers).

Various modifications of the invention in addition to those shown and described herein will be apparent to
5 those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

- 23 -

WHAT IS CLAIMED IS:

1. A method of tissue culture micropropagation of a rose plant comprising culturing a stem bearing a node in a first enclosed vessel containing a first culture medium
5 comprising about 0.5 to about 3.0 mg/l of a cytokine, 0 to about 1.0 mg/l of an auxin, and 0 to about 1.0 mg/l of a gibberellin until at least one shoot is produced from said node; excising said shoot from said stem and propagating said shoot in a second enclosed vessel containing a second
10 culture medium comprising about 0.5 to about 3.0 mg/l of a cytokine, 0 to about 1.0 mg/l of an auxin, and 0 to about 1.0 mg/l gibberellic acid to produce a rose plant in an enclosed vessel.

2. The method of claim 1 wherein said cytokine
15 is benzyladenine, said auxin is indole acetic acid, and said gibberellin is gibberellic acid.

3. The method of claim 2 wherein said first and said second culture media comprise about 1.0 mg/l benzyladenine, about 0.3 mg/l indole acetic acid, and about
20 0.1 mg/l gibberellic acid.

4. The method of claim 2 wherein said first and said second culture media comprise about 2.0 mg/l benzyladenine, about 0.3 mg/l indole acetic acid, and about 0.1 mg/l gibberellic acid.

25 5. The method of claim 2 wherein said first and said second culture media comprise about 3.0 mg/l benzyladenine, about 0.3 mg/l indole acetic acid, and about 0.1 mg/l gibberellic acid.

6. The method of claim 2 wherein said first and
30 said second culture media further comprises a nutritive medium.

- 24 -

7. The method of claim 6 wherein said nutritive medium comprises MS medium in combination with Gamborg's B5 vitamins.

8. The method of 7 wherein said culture media
5 comprises about 0.5 to about 3.0 mg/l of benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

9. The method of 7 wherein said culture media comprises about 1.0 mg/l benzyladenine.

10 10. The method of 7 wherein said culture media comprises about 2.0 mg/l benzyladenine.

11. The method of 7 wherein said culture media comprises about 3.0 mg/l benzyladenine.

12. The method of claim 2 wherein said rose plant
15 is *Rosa hybrida*.

13. The method of claim 12 wherein said *Rose hybrida* is selected from the cultivars consisting of Meirutral, Fairy, and Pink Miniwonder.

14. The method of claim 2 wherein said rose plant
20 is capable of being transferred to soil.

15. The method of claim 2 wherein said rose plant is a rose plant bearing a flower.

16. The method of claim 12 wherein said rose plant is a rose plant bearing a flower.

25 17. The method of claim 13 wherein said rose plant is a rose plant bearing a flower.

- 25 -

18. The method of claim 2 further comprising removing said rose plant from an enclosed vessel.

19. The method of claim 18 further comprising planting said rose plant in soil.

5 20. A rose plant produced by the method of claim 1.

 21. A rose plant produced by the method of claim 16.

10 22. A rose plant produced by the method of claim 17.

 23. A rose plant produced by the method of claim 18.

 24. A rose plant produced by the method of claim 19.

15 25. A method of tissue culture micropropagation of a rose plant comprising culturing a stem bearing a node in an enclosed vessel containing culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic
20 acid.

 26. The method of claim 25 wherein said stem is cultured until at least one shoot is produced from said node or until a rose plant is produced in said enclosed vessel.

25 27. The method of claim 26 wherein said rose plant is a rose plant bearing a flower.

 28. The method of claim 26 wherein said rose plant is capable of being transferred to soil.

- 26 -

29. The method of claim 25 further comprising removing said rose plant from an enclosed vessel.

30. The method of claim 29 further comprising planting said rose plant in soil.

5 31. A method of tissue culture micropropagation of a rose plant comprising culturing a shoot in an enclosed vessel containing culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid.

10 32. The method of claim 31 wherein said shoot is cultured until a rose plant is produced in said enclosed vessel.

33. The method of claim 32 wherein said rose plant is a rose plant bearing a flower.

15 34. The method of claim 32 wherein said rose plant is capable of being transferred to soil.

35. The method of claim 34 further comprising removing said rose plant from an enclosed vessel.

20 36. The method of claim 35 further comprising planting said rose plant in soil.

37. A culture medium for tissue culture micropropagation of a rose plant comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid.

25 38. The culture medium of claim 37 further comprising a nutritive medium.

- 27 -

39. The culture medium of claim 38 wherein said nutritative medium comprises MS medium in combination with Gamborg's B5 vitamins.

40. The culture medium of 39 comprising about 0.5 to about 3.0 mg/l of benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

41. The culture medium of 39 comprising about 1.0 mg/l benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

42. The culture medium of 39 wherein comprising about 2.0 mg/l benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

43. The culture medium of 39 comprising about 3.0 mg/l benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

44. The culture medium of claim 37 wherein said rose plant is *Rosa hybrida*.

45. The culture medium of claim 44 wherein said *Rosa hybrida* is selected from the cultivar consisting of Meirutral, Fairy, and Pink Miniwonder.

46. A rose culture comprising a rose stem bearing a node contained in an enclosed vessel containing culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid.

47. The rose culture of 46 comprising about 0.5 to about 3.0 mg/l of benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

- 28 -

48. The rose culture of 46 comprising about 1.0 mg/l benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

49. The rose culture of 46 comprising about 2.0 mg/l benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

50. The rose culture of 46 comprising about 3.0 mg/l benzyladenine, about 0.3 mg/l of indole acetic acid, and about 0.1 mg/l gibberellic acid.

51. The rose culture of claim 46 capable of generating a rose plant.

52. The rose culture of claim 46 wherein said stem bearing a node comprises at least one shoot.

53. The rose culture of claim 46 wherein said rose plant is *Rosa hybrida*.

54. The rose culture of claim 53 wherein said *Rosa hybrida* is selected from the cultivar consisting of Meirutral, Fairy, and Pink Miniwonder.

55. A rose plant produced from the rose culture of claim 46.

56. The rose plant of claim 55 wherein said rose is a rose bearing a flower.

57. The rose plant of claim 55 wherein said rose is *Rosa hybrida*.

58. The rose plant of claim 57 wherein said *Rosa hybrida* is selected from the cultivar consisting of Meirutral, Fairy, and Pink Miniwonder.

- 29 -

59. A rose explant selected from the group consisting of a rose stem bearing a node, a rose stem bearing a node and at least one shoot, and a rose shoot, said rose explant contained in an enclosed vessel comprising
5 a culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid.

60. The rose explant of claim 59 capable of producing a rose plant.

10 61. The rose explant of claim 60 wherein said rose plant is capable of transfer to soil.

62. The rose explant of claim 59 wherein said rose explant is *Rose hybrida*.

15 63. The rose explant of claim 60 wherein *Rosa hybrida* is selected from the cultivar consisting of Meirutral, Fairy, and Pink Miniwonder.

64. The method of claim 2 wherein the step of excising said shoot from said stem and propagating said shoot in a second enclosed vessel containing a second
20 culture medium comprising about 0.5 to about 3.0 mg/l benzyladenine, 0 to about 1.0 mg/l indole acetic acid, and 0 to about 1.0 mg/l gibberellic acid to produce a rose plant in an enclosed vessel is repeated more than one time.

65. The method of claim 2 further comprising
25 propagating said shoot in a third enclosed vessel containing a rooting medium.

66. The method of claim 65 wherein said rooting medium comprises MS medium, about 1 to about 5 mg/l glycine, about 50 to about 150 mg/l myo-inositol, about 0.1 to about
30 1.0 mg/l nicotinic acid, about 0.1 to about 1.0 mg/l

- 30 -

pyridoxine.HCl, about 0.01 to about 0.2 mg/l thiamine.HCl, about 10 to about 50 g/l sucrose, about 1 to about 5 g/l gelling agent, and optionally containing about 0.05 to about 1 mg/l indole butyric acid.

5 67. The method of claim 66 further comprising about 0.01 to about 0.2 mg/l naphthaleneacetic acid.

68. The method of claim 25 further comprising propagating said shoot in a third enclosed vessel containing a rooting medium.

10 69. The method of claim 68 wherein said rooting medium comprises MS medium, about 1 to about 5 mg/l glycine, about 50 to about 150 mg/l myo-inositol, about 0.1 to about 1.0 mg/l nicotinic acid, about 0.1 to about 1.0 mg/l pyridoxine.HCl, about 0.01 to about 0.2 mg/l thiamine.HCl,
15 about 10 to about 50 g/l sucrose, about 1 to about 5 g/l gelling agent, and optionally containing about 0.05 to about 1 mg/l indole butyric acid.

70. The method of claim 69 further comprising about 0.01 to about 0.2 mg/l naphthaleneacetic acid.

20 71. A rooting medium for the rooting of rose plants comprising MS medium and about 1 to about 5 mg/l glycine, about 50 to about 150 mg/l myo-inositol, about 0.1 to about 1.0 mg/l nicotinic acid, about 0.1 to about 1.0 mg/l pyridoxine.HCl, about 0.01 to about 0.2 mg/l
25 thiamine.HCl, about 10 to about 50 g/l sucrose, and optionally containing about 1 to about 5 g/l gelling agent, about 0.05 to about 1 mg/l indole butyric acid.

72. The rooting medium of claim 71 further comprising about 0.01 to about 0.2 mg/l naphthaleneacetic
30 acid.

1/3



FIG. 1A



FIG. 1B
SUBSTITUTE SHEET (RULE 26)

2/3



FIG. 2A



FIG. 2B
SUBSTITUTE SHEET (RULE 26)

3/3



FIG. 3A



FIG. 3B



FIG. 3C
SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/01744

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 5/00, 5/02

US CL : 435/240.4, 240.45, 240.48, 240.49, 240.54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/240.4, 240.45, 240.48, 240.49, 240.54

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CA, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Journal of American Horticultural Science, Volume 105, No. 2, issued 1980, P. Hasegawa, "Factors Affecting Shoot and Root Initiation from Cultured Rose Shoot Tips", pages 216-220, see entire document.	1-72

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 MAY 1996

Date of mailing of the international search report

12 JUN 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

L. BLAINE LANKFORD

Telephone No. (703) 308-0196