



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2690334 C 2017/02/14

(11)(21) **2 690 334**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2008/06/04
(87) Date publication PCT/PCT Publication Date: 2008/12/18
(45) Date de délivrance/Issue Date: 2017/02/14
(85) Entrée phase nationale/National Entry: 2009/12/01
(86) N° demande PCT/PCT Application No.: US 2008/065766
(87) N° publication PCT/PCT Publication No.: 2008/154249
(30) Priorité/Priority: 2007/06/08 (US60/942,906)

(51) Cl.Int./Int.Cl. *C40B 40/06* (2006.01),
C12Q 1/68 (2006.01), *C40B 30/04* (2006.01)
(72) Inventeurs/Inventors:
LEE-HOEFLICH, SI TUEN, US;
STERN, HOWARD, US
(73) Propriétaire/Owner:
GENENTECH, INC., US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : MARQUEURS D'EXPRESSION DE GENE DE RESISTANCE TUMORALE A UN TRAITEMENT PAR
INHIBITEUR HER2
(54) Title: GENE EXPRESSION MARKERS OF TUMOR RESISTANCE TO HER2 INHIBITOR TREATMENT

(57) **Abrégé/Abstract:**

The present invention concerns markers of resistance of HER2 expressing tumors to treatment with HER2 inhibitors, such as HER2 antibodies, including trastuzumab.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 December 2008 (18.12.2008)

PCT

(10) International Publication Number
WO 2008/154249 A9

(51) International Patent Classification:

G01N 33/50 (2006.01) **C12Q 1/68** (2006.01)

(21) International Application Number:

PCT/US2008/065766

(22) International Filing Date: 4 June 2008 (04.06.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/942,906 8 June 2007 (08.06.2007) US

(71) Applicant (for all designated States except US): **GENEN-TECH, INC.** [US/US]; 1 DNA Way, South San Francisco, CA 90480 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LEE-HOEFLICH, Si Tuen** [CA/US]; 371 Ashton Avenue, Millbrae, CA 94030 (US). **STERN, Howard** [US/US]; 864 Bauer Drive, San Carlos, CA 94070 (US).(74) Agents: **DREGER, Ginger R.** et al.; Goodwin Procter LLP, 135 Commonwealth Drive, Menlo Park, CA 94025 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- with information concerning authorization of rectification of an obvious mistake under Rule 91.1

(48) Date of publication of this corrected version:

30 April 2009

(15) Information about Correction:

see Notice of 30 April 2009

(54) Title: GENE EXPRESSION MARKERS OF TUMOR RESISTANCE TO HER2 INHIBITOR TREATMENT

(57) Abstract: The present invention concerns markers of resistance of HER2 expressing tumors to treatment with HER2 inhibitors, such as HER2 antibodies, including trastuzumab.



WO 2008/154249 A9

**GENE EXPRESSION MARKERS OF TUMOR RESISTANCE TO HER2 INHIBITOR
TREATMENT**

Field of the Invention

5 The present invention concerns markers of resistance of HER2 expressing tumors to treatment with HER2 inhibitors, such as HER2 antibodies, including trastuzumab.

Description of the Related Art

HER receptors and HER antibodies

10 The HER family of receptor tyrosine kinases are important mediators of cell growth, differentiation and survival. The receptor family includes four distinct members including epidermal growth factor receptor (EGFR, ErbB1, or HER1), HER2 (ErbB2 or p185^{neu}), HER3 (ErbB3) and HER4 (ErbB4 or tyro2).

15 The second member of the HER family, p185^{neu}, was originally identified as the product of the transforming gene from neuroblastomas of chemically treated rats. The activated form of the *neu* proto-oncogene results from a point mutation (valine to glutamic acid) in the transmembrane region of the encoded protein. Amplification of the human homolog of *neu* is observed in breast and ovarian cancers and correlates with a poor prognosis (Slamon *et al.*, *Science*, 235:177-182 (1987); Slamon *et al.*, *Science*, 244:707-712 (1989); and US Pat No. 4,968,603). To date, no point mutation analogous
20 to that in the *neu* proto-oncogene has been reported for human tumors. Overexpression of HER2 (frequently but not uniformly due to gene amplification) has also been observed in other carcinomas including carcinomas of the stomach, endometrium, salivary gland, lung, kidney, colon, thyroid, pancreas and bladder. See, among others, King *et al.*, *Science*, 229:974 (1985); Yokota *et al.*, *Lancet*: 1:765-767 (1986); Fukushige *et al.*, *Mol Cell Biol.*, 6:955-958 (1986); Guerin *et al.*, *Oncogene Res.*,
25 3:21-31 (1988); Cohen *et al.*, *Oncogene*, 4:81-88 (1989); Yonemura *et al.*, *Cancer Res.*, 51:1034 (1991); Borst *et al.*, *Gynecol. Oncol.*, 38:364 (1990); Weiner *et al.*, *Cancer Res.*, 50:421-425 (1990); Kern *et al.*, *Cancer Res.*, 50:5184 (1990); Park *et al.*, *Cancer Res.*, 49:6605 (1989); Zhau *et al.*, *Mol. Carcinog.*, 3:254-257 (1990); Aasland *et al. Br. J. Cancer* 57:358-363 (1988); Williams *et al. Pathobiology* 59:46-52 (1991); and McCann *et al., Cancer*, 65:88-92 (1990). HER2 may be
30 overexpressed in prostate cancer (Gu *et al. Cancer Lett.* 99:185-9 (1996); Ross *et al. Hum. Pathol.* 28:827-33 (1997); Ross *et al. Cancer* 79:2162-70 (1997); and Sadasivan *et al. J. Urol.* 150:126-31 (1993)).

 Antibodies directed against the rat p185^{neu} and human HER2 protein products have been described.

35 Drebin and colleagues have raised antibodies against the rat *neu* gene product, p185^{neu}. See, for example, Drebin *et al., Cell* 41:695-706 (1985); Myers *et al., Meth. Enzym.* 198:277-290 (1991); and WO94/22478. Drebin *et al. Oncogene* 2:273-277 (1988) report that mixtures of antibodies

reactive with two distinct regions of p185^{neu} result in synergistic anti-tumor effects on *neu*-transformed NIH-3T3 cells implanted into nude mice. See also U.S. Patent 5,824,311 issued October 20, 1998.

Hudziak *et al.*, *Mol. Cell. Biol.* 9(3):1165-1172 (1989) describe the generation of a panel of HER2 antibodies which were characterized using the human breast tumor cell line SK-BR-3. Relative cell proliferation of the SK-BR-3 cells following exposure to the antibodies was determined by crystal violet staining of the monolayers after 72 hours. Using this assay, maximum inhibition was obtained with the antibody called 4D5 which inhibited cellular proliferation by 56%. Other antibodies in the panel reduced cellular proliferation to a lesser extent in this assay. The antibody 4D5 was further found to sensitize HER2-overexpressing breast tumor cell lines to the cytotoxic effects of TNF- α . See also U.S. Patent No. 5,677,171 issued October 14, 1997. The HER2 antibodies discussed in Hudziak *et al.* are further characterized in Fendly *et al. Cancer Research* 50:1550-1558 (1990); Kotts *et al. In Vitro* 26(3):59A (1990); Sarup *et al. Growth Regulation* 1:72-82 (1991); Shepard *et al. J. Clin. Immunol.* 11(3):117-127 (1991); Kumar *et al. Mol. Cell. Biol.* 11(2):979-986 (1991); Lewis *et al. Cancer Immunol. Immunother.* 37:255-263 (1993); Pietras *et al. Oncogene* 9:1829-1838 (1994); Vitetta *et al. Cancer Research* 54:5301-5309 (1994); Sliwkowski *et al. J. Biol. Chem.* 269(20):14661-14665 (1994); Scott *et al. J. Biol. Chem.* 266:14300-5 (1991); D'souza *et al. Proc. Natl. Acad. Sci.* 91:7202-7206 (1994); Lewis *et al. Cancer Research* 56:1457-1465 (1996); and Schaefer *et al. Oncogene* 15:1385-1394 (1997).

A recombinant humanized version of the murine HER2 antibody 4D5 (huMAb4D5-8, rhuMAb HER2, trastuzumab or HERCEPTIN[®]; U.S. Patent No. 5,821,337) is clinically active in patients with HER2-overexpressing metastatic breast cancers that have received extensive prior anti-cancer therapy (Baselga *et al.*, *J. Clin. Oncol.* 14:737-744 (1996)). Trastuzumab received marketing approval from the Food and Drug Administration September 25, 1998 for the treatment of patients with metastatic breast cancer whose tumors overexpress the HER2 protein. In November 2006, the FDA approved HERCEPTIN[®] (trastuzumab) as part of a treatment regimen containing doxorubicin, cyclophosphamide and paclitaxel, for the adjuvant treatment of patients with HER2-positive, node-positive breast cancer. See also, Press *et al.*, *Cancer Res.* 53:4960-4970 (1993); Baselga *et al.*, *Cancer Res.* 58:2825-2831 (1998); Pegram *et al.*, *Proc. Am. Assoc. Cancer* 38:602 (1997), Abstract 4044; Slamon *et al.*, *N. Engl. J. Med.* 344:783-792 (2001); Lee *et al.*, *Nature* 378:394-396 (1995); Romond *et al.*, *N. Engl. J. Med.* 353:1673-1684 (2005); Ta-Chiu *et al.*, *J. Clin. Oncol.* 7811-7819 (2005).

Other HER2 antibodies with various properties have been described in Tagliabue *et al. Int. J. Cancer* 47:933-937 (1991); McKenzie *et al. Oncogene* 4:543-548 (1989); Maier *et al. Cancer Res.* 51:5361-5369 (1991); Bacus *et al. Molecular Carcinogenesis* 3:350-362 (1990); Stancovski *et al. PNAS (USA)* 88:8691-8695 (1991); Bacus *et al. Cancer Research* 52:2580-2589 (1992); Xu *et al. Int. J. Cancer* 53:401-408 (1993); WO94/00136; Kasprzyk *et al. Cancer Research* 52:2771-2776 (1992); Hancock *et al. Cancer Res.* 51:4575-4580 (1991); Shawver *et al. Cancer Res.* 54:1367-1373 (1994);

WO 2008/154249**PCT/US2008/065766**

Arteaga *et al.* *Cancer Res.* 54:3758-3765 (1994); Harwerth *et al.* *J. Biol. Chem.* 267:15160-15167 (1992); U.S. Patent No. 5,783,186; and Klapper *et al.* *Oncogene* 14:2099-2109 (1997).

Additional patent publications related to HER antibodies include: US 5,677,171, US

5,720,937, US 5,720,954, US 5,725,856, US 5,770,195, US 5,772,997, US 6,165,464, US 6,387,371,

5 US 6,399,063, US2002/0192211A1, US 6,015,567, US 6,333,169, US 4,968,603, US 5,821,337, US

6,054,297, US 6,407,213, US 6,719,971, US 6,800,738, US2004/0236078A1, US 5,648,237, US

6,267,958, US 6,685,940, US 6,821,515, WO98/17797, US 6,127,526, US 6,333,398, US 6,797,814,

US 6,339,142, US 6,417,335, US 6,489,447, WO99/31140, US2003/0147884A1,

US2003/0170234A1, US2005/0002928A1, US 6,573,043, US2003/0152987A1, WO99/48527,

10 US2002/0141993A1, WO01/00245, US2003/0086924, US2004/0013667A1, WO00/69460,

WO01/00238, WO01/15730, US 6,627,196B1, US6,632,979B1, WO01/00244, US2002/0090662A1,

WO01/89566, US2002/0064785, US2003/0134344, WO 04/24866, US2004/0082047,

US2003/0175845A1, WO03/087131, US2003/0228663, WO2004/008099A2, US2004/0106161,

WO2004/048525, US2004/0258685A1, US 5,985,553, US 5,747,261, US 4,935,341, US 5,401,638,

15 US 5,604,107, WO 87/07646, WO 89/10412, WO 91/05264, EP 412,116 B1, EP 494,135 B1, US

5,824,311, EP 444,181 B1, EP 1,006,194 A2, US 2002/0155527A1, WO 91/02062, US 5,571,894, US

5,939,531, EP 502,812 B1, WO 93/03741, EP 554,441 B1, EP 656,367 A1, US 5,288,477, US

5,514,554, US 5,587,458, WO 93/12220, WO 93/16185, US 5,877,305, WO 93/21319, WO 93/21232,

US 5,856,089, WO 94/22478, US 5,910,486, US 6,028,059, WO 96/07321, US 5,804,396, US

20 5,846,749, EP 711,565, WO 96/16673, US 5,783,404, US 5,977,322, US 6,512,097, WO 97/00271,

US 6,270,765, US 6,395,272, US 5,837,243, WO 96/40789, US 5,783,186, US 6,458,356, WO

97/20858, WO 97/38731, US 6,214,388, US 5,925,519, WO 98/02463, US 5,922,845, WO 98/18489,

WO 98/33914, US 5,994,071, WO 98/45479, US 6,358,682 B1, US 2003/0059790, WO 99/55367,

WO 01/20033, US 2002/0076695 A1, WO 00/78347, WO 01/09187, WO 01/21192, WO 01/32155,

25 WO 01/53354, WO 01/56604, WO 01/76630, WO02/05791, WO 02/11677, US 6,582,919,

US2002/0192652A1, US 2003/0211530A1, WO 02/44413, US 2002/0142328, US 6,602,670 B2, WO

02/45653, WO 02/055106, US 2003/0152572, US 2003/0165840, WO 02/087619, WO 03/006509,

WO03/012072, WO 03/028638, US 2003/0068318, WO 03/041736, EP 1,357,132, US

2003/0202973, US 2004/0138160, US 5,705,157, US 6,123,939, EP 616,812 B1, US 2003/0103973,

30 US 2003/0108545, US 6,403,630 B1, WO 00/61145, WO 00/61185, US 6,333,348 B1, WO 01/05425,

WO 01/64246, US 2003/0022918, US 2002/0051785 A1, US 6,767,541, WO 01/76586, US

2003/0144252, WO 01/87336, US 2002/0031515 A1, WO 01/87334, WO 02/05791, WO 02/09754,

US 2003/0157097, US 2002/0076408, WO 02/055106, WO 02/070008, WO 02/089842 and WO

03/86467.

35 U.S. Application Publication No. 2005010 (published May 12, 2005) and its PCT counterpart, WO 20054432, concern method for treating cancer, including lung cancer, bone cancer and ovarian cancer, with a combination of an ErbB2 ligand and an ErbB antibody.

U.S. Application Publication No. 20050119288 (published June 2, 2005) and its PCT counterpart, WO 200516347, are directed to a method for treating overexpression of the erbB2 receptor by administering a therapeutically effective amount of a first inhibitor of the erbB2 receptor; and subsequently, after an interval comprising less than 24 hours, from one to six therapeutically effective amounts of a second inhibitor of the erbB2 receptor.

WO 2006026313, published March 9, 2006, concerns method for treating cancer by administering 4-quinazolinamines, which are dual inhibitors of EGFR and ErbB2, in combination with at least one other ErbB family inhibitor.

HERCEPTIN[®] (trastuzumab) provides clinical benefit to a large percentage of patients diagnosed with HER2 positive breast cancer, both alone and in the adjuvant setting, in combination with chemotherapy. However, a significant number of HER2 positive patients exhibits either primary resistance or acquired resistance to treatment with trastuzumab. It is, therefore, a great need for identifying genes that might be involved in resistance to treatment with trastuzumab and other HER2 antibodies.

Pertuzumab (also known as recombinant human monoclonal antibody 2C4; OMNITARG[™], Genentech, Inc, South San Francisco) represents the first in a new class of agents known as HER dimerization inhibitors (HDI) and functions to inhibit the ability of HER2 to form active heterodimers with other HER receptors (such as EGFR/HER1, HER3 and HER4) and is active irrespective of HER2 expression levels. See, for example, Harari and Yarden *Oncogene* 19:6102-14 (2000); Yarden and Sliwkowski. *Nat Rev Mol Cell Biol* 2:127-37 (2001); Sliwkowski *Nat Struct Biol* 10:158-9 (2003); Cho *et al. Nature* 421:756-60 (2003); and Malik *et al. Pro Am Soc Cancer Res* 44:176-7 (2003).

Pertuzumab blockade of the formation of HER2-HER3 heterodimers in tumor cells has been demonstrated to inhibit critical cell signaling, which results in reduced tumor proliferation and survival (Agus *et al. Cancer Cell* 2:127-37 (2002)).

Pertuzumab has undergone testing as a single agent in the clinic with a phase Ia trial in patients with advanced cancers and phase II trials in patients with ovarian cancer and breast cancer as well as lung and prostate cancer. In a Phase I study, patients with incurable, locally advanced, recurrent or metastatic solid tumors that had progressed during or after standard therapy were treated with pertuzumab given intravenously every 3 weeks. Pertuzumab was generally well tolerated. Tumor regression was achieved in 3 of 20 patients evaluable for response. Two patients had confirmed partial responses. Stable disease lasting for more than 2.5 months was observed in 6 of 21 patients (Agus *et al. Pro Am Soc Clin Oncol* 22:192 (2003)). At doses of 2.0-15 mg/kg, the pharmacokinetics of pertuzumab was linear, and mean clearance ranged from 2.69 to 3.74 mL/day/kg and the mean terminal elimination half-life ranged from 15.3 to 27.6 days. Antibodies to pertuzumab were not detected (Allison *et al. Pro Am Soc Clin Oncol* 22:197 (2003)).

Diagnostics

Patients treated with the HER2 antibody trastuzumab are selected for therapy based on HER2

WO 2008/154249

PCT/US2008/065766

overexpression/amplification. See, for example, WO99/31140 (Paton *et al.*), US2003/0170234A1 (Hellmann, S.), and US2003/0147884 (Paton *et al.*); as well as WO01/89566, US2002/0064785, and US2003/0134344 (Mass *et al.*). See, also, US Patent No. 6,573,043, US Patent No. 6,905,830, and US2003/0152987, Cohen *et al.*, concerning immunohistochemistry (IHC) and fluorescence *in situ* hybridization (FISH) for detecting HER2 overexpression and amplification.

WO2004/053497 and US2004/024815A1 (Bacus *et al.*), as well as US 2003/0190689 (Crosby and Smith), refer to determining or predicting response to trastuzumab therapy. US2004/013297A1 (Bacus *et al.*) concerns determining or predicting response to ABX0303 EGFR antibody therapy.

WO2004/000094 (Bacus *et al.*) is directed to determining response to GW572016, a small molecule,

EGFR-HER2 tyrosine kinase inhibitor. WO2004/063709, Amler *et al.*, refers to biomarkers and methods for determining sensitivity to EGFR inhibitor, erlotinib HCl. US2004/0209290 and

WO04/065583, Cobleigh *et al.*, concern gene expression markers for breast cancer prognosis. See, also, WO03/078662 (Baker *et al.*), and WO03/040404 (Bevilacqua *et al.*). WO02/44413 (Danenber, K.) refers to determining EGFR and HER2 gene expression for determining a chemotherapeutic regimen.

Patients treated with pertuzumab can be selected for therapy based on HER activation or dimerization. Patent publications concerning pertuzumab and selection of patients for therapy therewith include: US Patent No. 6,949,245, WO01/00245, US2005/0208043, US2005/0238640, US2006/0034842, and US2006/0073143 (Adams *et al.*); US2003/0086924 (Sliwkowski, M.); US2004/0013667A1 (Sliwkowski, M.); as well as WO2004/008099A2, and US2004/0106161 (Bossenmaier *et al.*).

Cronin *et al. Am. J. Path.* 164(1): 35-42 (2004) describes measurement of gene expression in archival paraffin-embedded tissues. Ma *et al. Cancer Cell* 5:607-616 (2004) describes gene profiling by gene oligonucleotide microarray using isolated RNA from tumor-tissue sections taken from archived primary biopsies.

Summary of the Invention

In one aspect, the invention concerns a method of predicting the likelihood of response of a mammalian subject diagnosed with or at risk of developing a HER2 expressing tumor to treatment with a HER2 inhibitor, comprising

determining, in a biological sample obtained from said subject, the expression level of RNA transcripts or their expression products of one or more genes selected from the group consisting of CDK11, DYRK1A, LATS2, STK10, Wee1, DUSP4, DUSP6, HIPK3, JNK, MAP4K4, PTPN11, Socs5, PPM1H, DKFZP586B16, DGKI, FLJ35107, FLT1, HK2, ITK, MOAP1, KIAA0685, KIAA1639, LIM/PDLIM5, PANK1, P14K2B, PPP2R1A, PRKWNK3, RYK, SPEC2, STK22C, STYK1, and TXND3,

wherein a lower level of expression relative to one or more positive and/or negative controls indicates that the subject is likely to be resistant to treatment with the HER2 inhibitor.

The mammalian subject preferably is a human patient, such as a human cancer patient diagnosed with or at risk of developing a HER2 expressing cancer.

In various embodiments, the diagnosis includes quantification of the HER2 expression level, such as by immunohistochemistry (IHC) and/or fluorescence *in situ* hybridization (FISH).

5 In other embodiments, the cancer expresses HER2 at least at a 1+ level, or at least at a 2+ level, or at a 3+ level.

In another embodiment, the cancer is selected from the group consisting of breast cancer, squamous cell cancer, small-cell lung cancer (SCLC), non-small cell lung cancer (NSCLC), adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum,
 10 hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, colon cancer, rectal cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, testicular cancer, esophageal cancer, tumors of the biliary tract, and head and neck cancer.

15 In yet another embodiment, the cancer is selected from the group consisting of Overexpression of HER2 (frequently but not uniformly due to gene amplification) has also been observed in other carcinomas including carcinomas of the stomach, endometrium, salivary gland, lung, kidney, colon, thyroid, pancreas and bladder, and prostate cancer.

In still another embodiment, the cancer is breast cancer, such as metastatic breast cancer.

20 In various embodiments, the resistance to a HER2 inhibitor is determined by using one or more genes are selected from the group consisting of DYRK1A, HK2, Socs5, STK10, KIAA1639, and MAP4K4, and/or the group consisting of PTPN11, KIAA0685, and PPM1H.

The HER2 inhibitor may be an agent which interferes with HER2 activation or function.

25 HER 2 inhibitors include, without limitation, HER antibodies and antibody fragments, small molecule HER2 antagonists, HER2 tyrosine kinases inhibitors, and antisense molecules.

In one embodiment, the HER2 inhibitor is a HER2 antibody or antibody fragment, or a small molecule which binds to and inhibits the HER2 receptor.

In various embodiments, the HER2 antibody may inhibits HER2 ectodomain cleavage, may block ligand activation of a HER receptor, or may inhibit HER2 dimerization.

30 In another embodiment, the HER2 antibody binds to the heterodimeric binding site of HER2.

In yet another embodiment, the HER2 antibody or antibody fragment binds to the 4D5 epitope, and may, for example, be selected from the group consisting of humanized antibodies huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6, huMAb4D5-7 and trastuzumab, and fragments thereof.

35 In a preferred embodiment, the HER2 antibody is trastuzumab or a fragment thereof.

In a further embodiment, the HER2 antibody blocks ligand activation of a HER2 receptor more effectively than trastuzumab.

In a different embodiment, the HER2 antibody binds the 2C4 epitope, and may, for example, be pertuzumab or a fragment thereof.

In various embodiments, the biological sample is a tumor sample, such as a sample is from a fixed, wax-embedded cancer tissue specimen of a patient.

5 In another embodiment, the tumor sample is a core biopsy tissue.

In yet another embodiment, the biological sample is biological fluid, such as, for example, blood, urine, saliva, ascites fluid, blood serum or blood plasma.

In another aspect, the invention concerns an array comprising polynucleotides hybridizing to two or more, or at least 3, or at least 5 of the following genes: CDK11, DYRK1A, LATS2, STK10,
10 Wee1, DUSP4, DUSP6, HIPK3, JNK, MAP4K4, PTPN11, Socs5, PPM1H, DKFZP586B16, DGKI, FLJ35107, FLT1, HK2, ITK, MOAP1, KIAA0685, KIAA1639, LIM/PDLIM5, PANK1, P14K2B, PPP2R1A, PRKWNK3, RYK, SPEC2, STK22C, STYK1, and TXND3.

In one embodiment, the array comprises polynucleotides hybridizing to all of the following genes: CDK11, DYRK1A, LATS2, STK10, Wee1, DUSP4, DUSP6, HIPK3, JNK, MAP4K4,
15 PTPN11, Socs5, PPM1H, DKFZP586B16, DGKI, FLJ35107, FLT1, HK2, ITK, MOAP1, KIAA0685, KIAA1639, LIM/PDLIM5, PANK1, P14K2B, PPP2R1A, PRKWNK3, RYK, SPEC2, STK22C, STYK1, and TXND3.

In another embodiment, the array comprises polynucleotides hybridizing to the following genes: DYRK1A, HK2, Socs5, STK10, KIAA1639, and MAP4K4.

20 In yet another embodiment, the array comprises polynucleotides hybridizing to the following genes: PTPN11, KIAA0685, and PPM1H.

Brief Description of the Drawings

Figure 1. Measurement of trastuzumab response of HER2 amplified cell line BT474 by 3H-
25 thymidine incorporation assay.

Figure 2. Further HTP screen refinement by pilot automation experiments. NTC = non-targeting (negative) control.

Figure 3. Optimization of the screening window coefficient –Z factor.

Figure 4. Overview of the trastuzumab-resistance screen.

30 Figure 5. Statistical analysis.

Figure 6. Data analysis by plotting raw values of the screen showed p27 is a 4-oligo hit.

Figure 7. Combined analysis of kinase library hits.

Figure 8. Results from the kinase library screen.

Figure 9. Development of the secondary screen.

35 Figure 10. Combined analysis of the screens.

Figure 11. Summary of the phosphatase library screen.

Figure 12. Genologic expression data.

Figure 13. Top hits based on strongest phenotype and >2 oligo hit.

Figure 14. 3H-Thymidine uptake assay after 72 hours of trastuzumab treatment in BT474 cell line, with and without the knockdown of candidate genes.

Figure 15. 3H-Thymidine uptake assay after 72 hours of trastuzumab treatment in BT474M1 cell line.

Figure 16. 3H-Thymidine uptake assay of BT474M cell line after 72 hours of trastuzumab treatment and cell titer glow assays after 7 days of trastuzumab treatment.

Figure 17. 3H-Thymidine uptake assay of multiple HER2-amplified breast cancer cell lines by a dose range of Lapatinib treatment for 72 hours.

Figure 18. Western hybridization to examine both phosphorylation level and total level of HER3 in BT474 after trastuzumab treatment over time (top). Phospho-Akt ELISA and total-Akt ELISA to measure Akt1 in BT474 cell line after treatment with trastuzumab over time (bottom).

Figure 19. Phospho-Akt ELISA and total-Akt ELISA to measure Akt1 in BT474 cells after trastuzumab treatment over time,

Figure 20. Cladogram – PPM1 family members. The relative amino acid sequence similarity between other PP2C-like family members and PPM1H. By aligning amino acid sequence of the family and analyzed by computer program cluster W.

Figure 21. 3H-Thymidine uptake assay after 72 hours of trastuzumab treatment in BT474 cell line with and without the knockdown of closely related PP2C family members PPM1H, PPM1J, PPM1M.

Figure 22. 3H-Thymidine uptake assay of an HER2-amplified breast cancer cell line, HCC1419, by a dose range of Lapatinib treatment for 72 hours with and without the knockdown of closely related PP2C family members PPM1H and PPM1M.

Table 1. List of trastuzumab resistance markers identified.

Table 2. Summary of expression data in basal-like cell lines and tumors.

Table 3. Accession numbers of markers identified herein.+

Detailed Description of the Preferred Embodiment

Definitions

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton *et al.*, Dictionary of Microbiology and Molecular Biology 2nd ed., J. Wiley & Sons (New York, NY 1994), and March, Advanced Organic Chemistry Reactions, Mechanisms and Structure 4th ed., John Wiley & Sons (New York, NY 1992), provide one skilled in the art with a general guide to many of the terms used in the present application.

One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. Indeed, the

present invention is in no way limited to the methods and materials described. For purposes of the present invention, the following terms are defined below.

A “HER receptor” or “HER” is a receptor protein tyrosine kinase which belongs to the HER receptor family and includes EGFR (ErbB1, HER1), HER2 (ErbB2), HER3 (ErbB3) and HER4 (ErbB4) receptors. The HER receptor will generally comprise an extracellular domain, which may bind an HER ligand and/or dimerize with another HER receptor molecule; a lipophilic transmembrane domain; a conserved intracellular tyrosine kinase domain; and a carboxyl-terminal signaling domain harboring several tyrosine residues which can be phosphorylated. The HER receptor may be a “native sequence” HER receptor or an “amino acid sequence variant” thereof. Preferably the HER receptor is native sequence human HER receptor. Thus, the term “HER”, as used herein, will encompass HER1, HER2, HER3, and HER4.

The terms “ErbB1”, “HER1”, “epidermal growth factor receptor” and “EGFR” are used interchangeably herein and refer to EGFR as disclosed, for example, in Carpenter *et al. Ann. Rev. Biochem.* 56:881-914 (1987), including naturally occurring mutant forms thereof (*e.g.* a deletion mutant EGFR as in Ullrich *et al. Nature* (1984) 309:418425 and Humphrey *et al. PNAS (USA)* 87:4207-4211 (1990)), as well as variants thereof, such as EGFRvIII. Variants of EGFR also include deletional, substitutional and insertional variants, for example those described in Lynch *et al. (New England Journal of Medicine)* 2004, 350:2129, Paez *et al. (Science)* 2004, 304:1497, and Pao *et al. (PNAS)* 2004, 101:13306).

The expressions “ErbB2” and “HER2” are used interchangeably herein and refer to human HER2 protein described, for example, in Semba *et al., PNAS (USA)* 82:6497-6501 (1985) and Yamamoto *et al. Nature* 319:230-234 (1986) (GenBank accession number X03363). The term “*erbB2*” refers to the gene encoding human HER2 and “*neu*” refers to the gene encoding rat p185^{neu}. Preferred HER2 is native sequence human HER2.

Herein, “HER2 extracellular domain” or “HER2 ECD” refers to a domain of HER2 that is outside of a cell, either anchored to a cell membrane, or in circulation, including fragments thereof. In one embodiment, the extracellular domain of HER2 may comprise four domains: “Domain I” (amino acid residues from about 1-195, “Domain II” (amino acid residues from about 196-319), “Domain III” (amino acid residues from about 320-488), and “Domain IV” (amino acid residues from about 489-630) (residue numbering without signal peptide). See Garrett *et al. Mol. Cell.* 11: 495-505 (2003), Cho *et al. Nature* 421: 756-760 (2003), Franklin *et al. Cancer Cell* 5:317-328 (2004), and Plowman *et al. Proc. Natl. Acad. Sci.* 90:1746-1750 (1993), as well as Fig. 1 herein.

“ErbB3” and “HER3” refer to the receptor polypeptide as disclosed, for example, in US Pat. Nos. 5,183,884 and 5,480,968 as well as Kraus *et al. PNAS (USA)* 86:9193-9197 (1989).

The terms “ErbB4” and “HER4” herein refer to the receptor polypeptide as disclosed, for example, in EP Pat. Appln. No. 599,274; Plowman *et al., Proc. Natl. Acad. Sci. USA*, 90:1746-1750

(1993); and Plowman *et al.*, *Nature*, 366:473-475 (1993), including isoforms thereof, *e.g.*, as disclosed in WO99/19488, published April 22, 1999.

By “HER ligand” is meant a polypeptide which binds to and/or activates a HER receptor. The HER ligand of particular interest herein is a native sequence human HER ligand such as epidermal growth factor (EGF) (Savage *et al.*, *J. Biol. Chem.* 247:7612-7621 (1972)); transforming growth factor alpha (TGF- α) (Marquardt *et al.*, *Science* 223:1079-1082 (1984)); amphiregulin also known as schwanoma or keratinocyte autocrine growth factor (Shoyab *et al.* *Science* 243:1074-1076 (1989); Kimura *et al.* *Nature* 348:257-260 (1990); and Cook *et al.* *Mol. Cell. Biol.* 11:2547-2557 (1991)); betacellulin (Shing *et al.*, *Science* 259:1604-1607 (1993); and Sasada *et al.* *Biochem. Biophys. Res. Commun.* 190:1173 (1993)); heparin-binding epidermal growth factor (HB-EGF) (Higashiyama *et al.*, *Science* 251:936-939 (1991)); epiregulin (Toyoda *et al.*, *J. Biol. Chem.* 270:7495-7500 (1995); and Komurasaki *et al.* *Oncogene* 15:2841-2848 (1997)); a heregulin (see below); neuregulin-2 (NRG-2) (Carraway *et al.*, *Nature* 387:512-516 (1997)); neuregulin-3 (NRG-3) (Zhang *et al.*, *Proc. Natl. Acad. Sci.* 94:9562-9567 (1997)); neuregulin-4 (NRG-4) (Harari *et al.* *Oncogene* 18:2681-89 (1999)); and cripto (CR-1) (Kannan *et al.* *J. Biol. Chem.* 272(6):3330-3335 (1997)). HER ligands which bind EGFR include EGF, TGF- α , amphiregulin, betacellulin, HB-EGF and epiregulin. HER ligands which bind HER3 include heregulins. HER ligands capable of binding HER4 include betacellulin, epiregulin, HB-EGF, NRG-2, NRG-3, NRG-4, and heregulins.

“Heregulin” (HRG) when used herein refers to a polypeptide encoded by the heregulin gene product as disclosed in U.S. Patent No. 5,641,869, or Marchionni *et al.*, *Nature*, 362:312-318 (1993). Examples of heregulins include heregulin- α , heregulin- β 1, heregulin- β 2 and heregulin- β 3 (Holmes *et al.*, *Science*, 256:1205-1210 (1992); and U.S. Patent No. 5,641,869); *neu* differentiation factor (NDF) (Peles *et al.* *Cell* 69: 205-216 (1992)); acetylcholine receptor-inducing activity (ARIA) (Falls *et al.* *Cell* 72:801-815 (1993)); glial growth factors (GGFs) (Marchionni *et al.*, *Nature*, 362:312-318 (1993)); sensory and motor neuron derived factor (SMDF) (Ho *et al.* *J. Biol. Chem.* 270:14523-14532 (1995)); γ -heregulin (Schaefer *et al.* *Oncogene* 15:1385-1394 (1997)).

A “HER dimer” herein is a noncovalently associated dimer comprising at least two HER receptors. Such complexes may form when a cell expressing two or more HER receptors is exposed to an HER ligand and can be isolated by immunoprecipitation and analyzed by SDS-PAGE as described in Sliwkowski *et al.*, *J. Biol. Chem.*, 269(20):14661-14665 (1994), for example. Other proteins, such as a cytokine receptor subunit (*e.g.* gp130) may be associated with the dimer. Preferably, the HER dimer comprises HER2.

A “HER heterodimer” herein is a noncovalently associated heterodimer comprising at least two different HER receptors, such as EGFR-HER2, HER2-HER3 or HER2-HER4 heterodimers.

A “HER inhibitor” is an agent which interferes with HER activation or function. Examples of HER inhibitors include HER antibodies (*e.g.* EGFR, HER2, HER3, or HER4 antibodies); EGFR-targeted drugs; small molecule HER antagonists; HER tyrosine kinase inhibitors; HER2 and EGFR

dual tyrosine kinase inhibitors such as lapatinib/GW572016; antisense molecules (see, for example, WO2004/87207); and/or agents that bind to, or interfere with function of, downstream signaling molecules, such as MAPK or Akt. Preferably, the HER inhibitor is an antibody or small molecule which binds to a HER receptor. The term “HER inhibitor” specifically includes HER1, HER2, HER3 and HER4 inhibitors. Thus, for example, a HER2 inhibitor is an agent which interferes with HER2 activation or function, including antibodies, small molecule HER2 antagonists, HER2 tyrosine kinase inhibitors, HER2 and EGFR dual tyrosine kinase inhibitors, antisense molecules, and the like.

A “HER dimerization inhibitor” or “HDI” is an agent which inhibits formation of a HER homodimer or HER heterodimer. Preferably, the HER dimerization inhibitor is an antibody, for example an antibody which binds to HER2 at the heterodimeric binding site thereof. However, HER dimerization inhibitors also include peptide and non-peptide small molecules, and other chemical

entities which inhibit the formation of HER homo- or heterodimers. The most preferred HER dimerization inhibitor herein is pertuzumab or MAb 2C4. Binding of 2C4 to the heterodimeric binding site of HER2 is illustrated in Fig. 4. Other examples of HER dimerization inhibitors include antibodies which bind to EGFR and inhibit dimerization thereof with one or more other HER receptors (for example EGFR monoclonal antibody 806, MAb 806, which binds to activated or “untethered” EGFR; see Johns *et al.*, *J. Biol. Chem.* 279(29):30375-30384 (2004)); antibodies which bind to HER3 and inhibit dimerization thereof with one or more other HER receptors; antibodies which bind to HER4 and inhibit dimerization thereof with one or more other HER receptors; peptide dimerization inhibitors (US Patent No. 6,417,168); antisense dimerization inhibitors; etc.

A “HER2 dimerization inhibitor” herein is a HER2 antibody or other HER2 antagonist, such as a peptide or on-peptide small molecule, which bind to HER2 and interferes with the formation of HER2-containing oligomers, including HER2 homo- and heterodimers, such as one or more of HER2-HER2, HER2-EGFR, HER2-HER3, and HER2-HER4 heterodimers. Preferably, the HER2 dimerization inhibitor is a molecule, such as an HER2 antibody or a peptide or non-peptide small molecule, that blocks the formation of all of HER2-HER2, HER2-EGFR and HER2-HER3 heterodimers, for example by binding to HER2 at a location required for heterodimerization, such as the heterodimeric binding site shown in Figure 4. A typical representative of such HER2 dimerization inhibitors is pertuzumab, which was also listed as a “HER dimerization inhibitor” in a broader sense.

A “HER antibody” or “HER antibody” is an antibody that binds to a HER receptor. Optionally, the HER antibody further interferes with HER activation or function. Preferably, the HER antibody binds to the HER2 receptor. A HER2 antibody of particular interest herein is trastuzumab. Another example of a HER2 antibody is pertuzumab.

“HER activation” refers to activation, or phosphorylation, of any one or more HER receptors. Generally, HER activation results in signal transduction (*e.g.* that caused by an intracellular kinase domain of a HER receptor phosphorylating tyrosine residues in the HER receptor or a substrate polypeptide). HER activation may be mediated by HER ligand binding to a HER dimer comprising

the HER receptor of interest. HER ligand binding to a HER dimer may activate a kinase domain of one or more of the HER receptors in the dimer and thereby results in phosphorylation of tyrosine residues in one or more of the HER receptors and/or phosphorylation of tyrosine residues in additional substrate polypeptides(s), such as Akt or MAPK intracellular kinases.

5 “Phosphorylation” refers to the addition of one or more phosphate group(s) to a protein, such as a HER receptor, or substrate thereof.

An antibody which “inhibits HER dimerization” is an antibody which inhibits, or interferes with, formation of a HER dimer, regardless of the underlying mechanism. Preferably, such an antibody binds to HER2 at the heterodimeric binding site thereof. The most preferred dimerization
10 inhibiting antibody herein is pertuzumab or MAb 2C4. Binding of 2C4 to the heterodimeric binding site of HER2 is illustrated in Fig. 4. Other examples of antibodies which inhibit HER dimerization include antibodies which bind to EGFR and inhibit dimerization thereof with one or more other HER receptors (for example EGFR monoclonal antibody 806, MAb 806, which binds to activated or “untethered” EGFR; see Johns *et al.*, *J. Biol. Chem.* 279(29):30375-30384 (2004)); antibodies which
15 bind to HER3 and inhibit dimerization thereof with one or more other HER receptors; and antibodies which bind to HER4 and inhibit dimerization thereof with one or more other HER receptors.

An antibody which “blocks ligand activation of a HER receptor more effectively than trastuzumab” is one which reduces or eliminates HER ligand activation of HER receptor(s) or HER dimer(s) more effectively (for example at least about 2-fold more effectively) than trastuzumab.
20 Preferably, such an antibody blocks HER ligand activation of a HER receptor at least about as effectively as murine monoclonal antibody 4D5 or a Fab fragment thereof, or as trastuzumab or a Fab fragment thereof. One can evaluate the ability of an antibody to block ligand activation of a HER receptor by studying HER dimers directly, or by evaluating HER activation, or downstream signaling, which results from HER dimerization, and/or by evaluating the antibody-HER2 binding site, etc.
25 Assays for screening for antibodies with the ability to inhibit ligand activation of a HER receptor more effectively than trastuzumab are described in Agus *et al. Cancer Cell* 2: 127-137 (2002) and US Patent No. 6,949,245 (Adams *et al.*). By way of example only, one may assay for: inhibition of HER dimer formation (see, *e.g.*, Fig. 1A-B of Agus *et al. Cancer Cell* 2: 127-137 (2002); and US Patent No. 6,949,245); reduction in HER ligand activation of cells which express HER dimers (US Patent
30 No. 6,949,245 and Fig. 2A-B of Agus *et al. Cancer Cell* 2: 127-137 (2002), for example); blocking of HER ligand binding to cells which express HER dimers (US Patent No. 6,949,245, and Fig. 2E of Agus *et al. Cancer Cell* 2: 127-137 (2002), for example); cell growth inhibition of cancer cells (*e.g.* MCF7, MDA-MD-134, ZR-75-1, MD-MB-175, T-47D cells) which express HER dimers in the presence (or absence) of HER ligand (US Patent No. 6,949,245 and Figs. 3A-D of Agus *et al. Cancer Cell* 2: 127-137 (2002), for instance); inhibition of downstream signaling (for instance, inhibition of HRG-dependent AKT phosphorylation or inhibition of HRG- or TGF α - dependent MAPK
35 phosphorylation) (see, US Patent No. 6,949,245, and Fig. 2C-D of Agus *et al. Cancer Cell* 2: 127-137

(2002), for example). One may also assess whether the antibody inhibits HER dimerization by studying the antibody-HER2 binding site, for instance, by evaluating a structure or model, such as a crystal structure, of the antibody bound to HER2 (See, for example, Franklin *et al. Cancer Cell* 5:317-328 (2004)).

5 A “heterodimeric binding site” on HER2, refers to a region in the extracellular domain of HER2 that contacts, or interfaces with, a region in the extracellular domain of EGFR, HER3 or HER4 upon formation of a dimer therewith. The region is found in Domain II of HER2. Franklin *et al. Cancer Cell* 5:317-328 (2004).

The HER2 antibody may “inhibit HRG-dependent AKT phosphorylation” and/or inhibit
10 “HRG- or TGF α -dependent MAPK phosphorylation” more effectively (for instance at least 2-fold more effectively) than trastuzumab (see Agus *et al. Cancer Cell* 2: 127-137 (2002) and WO01/00245, by way of example).

The HER2 antibody may be one which, like pertuzumab, does “not inhibit HER2 ectodomain cleavage” (Molina *et al. Cancer Res.* 61:4744-4749(2001)). Trastuzumab, on the other hand, can
15 inhibit HER2 ectodomain cleavage. Thus, the HER2 antibody may be one which, like trastuzumab, inhibits HER2 ectodomain cleavage.

A HER2 antibody that “binds to a heterodimeric binding site” of HER2, binds to residues in domain II (and optionally also binds to residues in other of the domains of the HER2 extracellular domain, such as domains I and III), and can sterically hinder, at least to some extent, formation of a
20 HER2-EGFR, HER2-HER3, or HER2-HER4 heterodimer. Franklin *et al. Cancer Cell* 5:317-328 (2004) characterize the HER2-pertuzumab crystal structure, deposited with the RCSB Protein Data Bank (ID Code IS78), illustrating an exemplary antibody that binds to the heterodimeric binding site of HER2.

An antibody that “binds to domain II” of HER2 binds to residues in domain II and optionally
25 residues in other domain(s) of HER2, such as domains I and III. Preferably the antibody that binds to domain II binds to the junction between domains I, II and III of HER2.

Herein “time to disease progression” or “TTP” refer to the time, generally measured in weeks or months, from the time of initial treatment until the cancer progresses or worsens. Such progression can be evaluated by the skilled clinician.

30 By “extending TTP” is meant increasing the time to disease progression in a treated patient relative to an untreated patient.

“Survival” refers to the patient remaining alive, and includes overall survival as well as progression free survival.

“Overall survival” refers to the patient remaining alive for a defined period of time, such as 1
35 year, 5 years, etc from the time of diagnosis or treatment.

“Progression free survival” refers to the patient remaining alive, without disease progression.

By “extending survival” is meant increasing overall or progression free survival in a treated patient relative to an untreated patient, or relative to a patient treated with an approved anti-tumor agent for the treatment of the cancer in question.

5 An “objective response” refers to a measurable response, including complete response (CR) or partial response (PR).

By “complete response” or “CR” is intended the disappearance of all signs of cancer in response to treatment. This does not always mean the cancer has been cured.

“Partial response” or “PR” refers to a decrease in the size of one or more tumors or lesions, or in the extent of cancer in the body, in response to treatment.

10 The term “refractory tumor” or “refractory cancer” is used to refer to tumors that fail to respond to or are resistant to a certain treatment, such as treatment with a HER2 inhibitor, such as a HER2 antibody, e.g. trastuzumab, when administered alone or in combination with other cancer treatments. For the purposes of this specification, refractory tumors also encompass tumors that appear to be inhibited by such treatment(s) but recur within 12 months from the completion of such
15 treatment.

A tumor which “responds poorly” to a certain treatment, such as treatment with a HER2 inhibitor, such as a HER2 antibody, e.g. trastuzumab, does not show statistically significant improvement in response to such treatment when compared to no treatment or treatment with placebo in a recognized animal model or a human clinical trial, or which responds to initial treatment but
20 grows as treatment is continued.

The term “standard of care” is used to refer to a treatment process that an ordinary skilled prudent physician uses to treat a certain disease, such as cancer. The standard of care varies depending on the type and stage of cancer, the patient’s condition and treatment history, and the like, and will be apparent to those skilled in the art.

25 Protein “expression” refers to conversion of the information encoded in a gene into messenger RNA (mRNA) and then to the protein.

Herein, a sample or cell that “expresses” a protein of interest (such as a HER receptor or HER ligand) is one in which mRNA encoding the protein, or the protein, including fragments thereof, is determined to be present in the sample or cell.

30 The phrase “gene amplification” refers to a process by which multiple copies of a gene or gene fragment are formed in a particular cell or cell line. The duplicated region (a stretch of amplified DNA) is often referred to as “amplicon.” Usually, the amount of the messenger RNA (mRNA) produced also increases in the proportion of the number of copies made of the particular gene expressed.

35 The term “modulate” is used herein to mean that the expression of the gene, or level of RNA molecule or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity

of one or more proteins or protein subunits is up regulated or down regulated, such that expression, level, or activity is greater than or less than that observed in the absence of the modulator.

The terms “inhibit”, “down-regulate”, and “reduce” are used interchangeably and mean that the expression of a gene, or level of RNA molecules or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits, is reduced relative to one or more controls, such as, for example, one or more positive and/or negative controls.

The term “up-regulate” is used to mean that the expression of a gene, or level of RNA molecules or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits, is elevated relative to one or more controls, such as, for example, one or more positive and/or negative controls.

An “interfering RNA” or “small interfering RNA (siRNA)” is a double stranded RNA molecule usually less than about 30 nucleotides in length that reduces expression of a target gene. Interfering RNAs may be identified and synthesized using known methods (Shi Y., *Trends in Genetics* 19(1):9-12 (2003), WO/2003056012 and WO2003064621), and siRNA libraries are commercially available, for example from Dharmacon, Lafayette, Colorado.

A “native sequence” polypeptide is one which has the same amino acid sequence as a polypeptide (*e.g.*, HER receptor or HER ligand) derived from nature, including naturally occurring or allelic variants. Such native sequence polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. Thus, a native sequence polypeptide can have the amino acid sequence of naturally occurring human polypeptide, murine polypeptide, or polypeptide from any other mammalian species.

The term “antibody” herein is used in the broadest sense and specifically covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.* bispecific antibodies), and antibody fragments, so long as they exhibit the desired biological activity.

The term “monoclonal antibody” as used herein refers to an antibody from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical and/or bind the same epitope(s), except for possible variants that may arise during production of the monoclonal antibody, such variants generally being present in minor amounts. Such monoclonal antibody typically includes an antibody comprising a polypeptide sequence that binds a target, wherein the target-binding polypeptide sequence was obtained by a process that includes the selection of a single target binding polypeptide sequence from a plurality of polypeptide sequences. For example, the selection process can be the selection of a unique clone from a plurality of clones, such as a pool of hybridoma clones, phage clones or recombinant DNA clones. It should be understood that the selected target binding sequence can be further altered, for example, to improve affinity for the target, to humanize the target binding sequence, to improve its production in cell culture, to reduce its immunogenicity *in vivo*, to create a multispecific antibody, *etc.*, and that an antibody comprising the altered target binding sequence is also a monoclonal antibody of this

invention. In contrast to polyclonal antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. In addition to their specificity, the monoclonal antibody preparations are advantageous in that they are typically

5 uncontaminated by other immunoglobulins. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including, for example, the hybridoma method (*e.g.*, Kohler *et al.*, *Nature*, 256:495

10 (1975); Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling *et al.*, in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681, (Elsevier,

N.Y., 1981)), recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567), phage display technologies (see, *e.g.*, Clackson *et al.*, *Nature*, 352:624-628 (1991); Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991); Sidhu *et al.*, *J. Mol. Biol.* 338(2):299-310 (2004); Lee *et al.*,

15 *J.Mol.Biol.*340(5):1073-1093 (2004); Fellouse, *Proc. Nat. Acad. Sci. USA* 101(34):12467-12472 (2004); and Lee *et al. J. Immunol. Methods* 284(1-2):119-132 (2004), and technologies for producing human or human-like antibodies in animals that have parts or all of the human immunoglobulin loci or

genes encoding human immunoglobulin sequences (see, *e.g.*, WO 1998/24893; WO 1996/34096; WO 1996/33735; WO 1991/10741; Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993);

20 Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggemann *et al.*, *Year in Immuno.*, 7:33 (1993); U.S. Patent Nos. 5,545,806; 5,569,825; 5,591,669 (all of GenPharm); U.S. Patent No. 5,545,807; WO 1997/17852; U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and 5,661,016; Marks *et al.*, *Bio/Technology*, 10: 779-783 (1992); Lonberg *et al.*, *Nature*, 368: 856-859 (1994);

Morrison, *Nature*, 368: 812-813 (1994); Fishwild *et al.*, *Nature Biotechnology*, 14: 845-851 (1996);
25 Neuberger, *Nature Biotechnology*, 14: 826 (1996); and Lonberg and Huszar, *Intern. Rev. Immunol.*, 13: 65-93 (1995)).

The monoclonal antibodies herein specifically include “chimeric” antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass,
30 while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include “primatized” antibodies comprising variable domain antigen-
35 binding sequences derived from a non-human primate (*e.g.* Old World Monkey, Ape etc) and human constant region sequences, as well as “humanized” antibodies.

“Humanized” forms of non-human (*e.g.*, rodent) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

Humanized HER2 antibodies include huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6, huMAb4D5-7 and huMAb4D5-8 or trastuzumab as described in Table 3 of U.S. Patent 5,821,337 expressly incorporated herein by reference; humanized 520C9 (WO93/21319); and humanized 2C4 antibodies such as pertuzumab as described herein.

An “intact antibody” herein is one which comprises two antigen binding regions, and an Fc region. Preferably, the intact antibody has a functional Fc region.

“Antibody fragments” comprise a portion of an intact antibody, preferably comprising the antigen binding region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragment(s).

“Native antibodies” are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains.

The term “variable” refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called

hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC).

The term “hypervariable region” when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region generally comprises amino acid residues from a “complementarity determining region” or “CDR” (*e.g.* residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a “hypervariable loop” (*e.g.* residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). “Framework Region” or “FR” residues are those variable domain residues other than the hypervariable region residues as herein defined.

Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, each with a single antigen-binding site, and a residual “Fc” fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

“Fv” is the minimum antibody fragment which contains a complete antigen-recognition and antigen-binding site. This region consists of a dimer of one heavy chain and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three hypervariable regions of each variable domain interact to define an antigen-binding site on the surface of the V_H - V_L dimer. Collectively, the six hypervariable regions confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three hypervariable regions specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab=fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

The term "Fc region" herein is used to define a C-terminal region of an immunoglobulin heavy chain, including native sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue.

Unless indicated otherwise, herein the numbering of the residues in an immunoglobulin heavy chain is that of the EU index as in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991), expressly incorporated herein by reference. The "EU index as in Kabat" refers to the residue numbering of the human IgG1 EU antibody.

A "functional Fc region" possesses an "effector function" of a native sequence Fc region. Exemplary "effector functions" include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (*e.g.* B cell receptor; BCR), etc. Such effector functions generally require the Fc region to be combined with a binding domain (*e.g.* an antibody variable domain) and can be assessed using various assays as herein disclosed, for example.

A "native sequence Fc region" comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof.

A “variant Fc region” comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification, preferably one or more amino acid substitution(s). Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, *e.g.* from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% homology therewith, more preferably at least about 95% homology therewith.

Depending on the amino acid sequence of the constant domain of their heavy chains, intact antibodies can be assigned to different “classes”. There are five major classes of intact antibodies:

IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into “subclasses” (isotypes), *e.g.*, IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

“Antibody-dependent cell-mediated cytotoxicity” and “ADCC” refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (*e.g.* Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express Fc γ RIII only, whereas monocytes express Fc γ RI, Fc γ RII and Fc γ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, *e.g.*, in a animal model such as that disclosed in Clynes *et al. PNAS (USA)* 95:652-656 (1998).

“Human effector cells” are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least Fc γ RIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, *e.g.* from blood or PBMCs as described herein.

The terms “Fc receptor” or “FcR” are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc γ RII receptors include Fc γ RIIA (an “activating receptor”) and Fc γ RIIB (an “inhibiting receptor”), which

have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof.

Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain (see review M. in Daëron, *Annu. Rev. Immunol.*

15 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991); Capel *et al.*, *Immunomethods* 4:25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, *J. Immunol.* 117:587 (1976) and Kim *et al.*, *J. Immunol.*

10 24:249 (1994)), and regulates homeostasis of immunoglobulins.

“Complement dependent cytotoxicity” or “CDC” refers to the ability of a molecule to lyse a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (*e.g.* an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, *e.g.* as described in Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996), may be performed.

15 “Single-chain Fv” or “scFv” antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Plückthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994). HER2 antibody scFv fragments are described in WO93/16185; U.S. Patent No. 5,571,894; and U.S. Patent No. 5,587,458.

20 The term “diabodies” refers to small antibody fragments with two antigen-binding sites, which fragments comprise a variable heavy domain (V_H) connected to a variable light domain (V_L) in the same polypeptide chain (V_H - V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

30 A “naked antibody” is an antibody that is not conjugated to a heterologous molecule, such as a cytotoxic moiety or radiolabel.

An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at

least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

An "affinity matured" antibody is one with one or more alterations in one or more hypervariable regions thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen.

Affinity matured antibodies are produced by procedures known in the art. Marks *et al. Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas *et al. Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier *et al. Gene* 169:147-155 (1995); Yelton *et al. J. Immunol.* 155:1994-2004 (1995); Jackson *et al., J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al, J. Mol. Biol.* 226:889-896 (1992).

The term "main species antibody" herein refers to the antibody structure in a composition which is the quantitatively predominant antibody molecule in the composition. In one embodiment, the main species antibody is a HER2 antibody, such as an antibody that binds to Domain II of HER2, antibody that inhibits HER dimerization more effectively than trastuzumab, and/or an antibody which binds to a heterodimeric binding site of HER2.

An "amino acid sequence variant" antibody herein is an antibody with an amino acid sequence which differs from a main species antibody. Ordinarily, amino acid sequence variants will possess at least about 70% homology with the main species antibody, and preferably, they will be at least about 80%, more preferably at least about 90% homologous with the main species antibody. The amino acid sequence variants possess substitutions, deletions, and/or additions at certain positions within or adjacent to the amino acid sequence of the main species antibody. Examples of amino acid sequence variants herein include an acidic variant (*e.g.* deamidated antibody variant), a basic variant, an antibody with an amino-terminal leader extension (*e.g.* VHS-) on one or two light chains thereof, an antibody with a C-terminal lysine residue on one or two heavy chains thereof, etc., and includes combinations of variations to the amino acid sequences of heavy and/or light chains. The antibody variant of particular interest herein is the antibody comprising an amino-terminal leader extension on one or two light chains thereof, optionally further comprising other amino acid sequence and/or glycosylation differences relative to the main species antibody.

A “glycosylation variant” antibody herein is an antibody with one or more carbohydrate moieties attached thereto which differ from one or more carbohydrate moieties attached to a main species antibody. Examples of glycosylation variants herein include antibody with a G1 or G2 oligosaccharide structure, instead a G0 oligosaccharide structure, attached to an Fc region thereof, antibody with one or two carbohydrate moieties attached to one or two light chains thereof, antibody with no carbohydrate attached to one or two heavy chains of the antibody, etc., and combinations of glycosylation alterations.

Where the antibody has an Fc region, an oligosaccharide structure may be attached to one or two heavy chains of the antibody, *e.g.* at residue 299 (298, Eu numbering of residues). For pertuzumab, G0 was the predominant oligosaccharide structure, with other oligosaccharide structures such as G0-F, G-1, Man5, Man6, G1-1, G1(1-6), G1(1-3) and G2 being found in lesser amounts in the pertuzumab composition.

Unless indicated otherwise, a “G1 oligosaccharide structure” herein includes G-1, G1-1, G1(1-6) and G1(1-3) structures.

An “amino-terminal leader extension” herein refers to one or more amino acid residues of the amino-terminal leader sequence that are present at the amino-terminus of any one or more heavy or light chains of an antibody. An exemplary amino-terminal leader extension comprises or consists of three amino acid residues, VHS, present on one or both light chains of an antibody variant.

A “deamidated” antibody is one in which one or more asparagine residues thereof has been derivatized, *e.g.* to an aspartic acid, a succinimide, or an iso-aspartic acid.

The term “tumor,” as used herein, refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues.

The terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma (including medulloblastoma and retinoblastoma), sarcoma (including liposarcoma and synovial cell sarcoma), neuroendocrine tumors (including carcinoid tumors, gastrinoma, and islet cell cancer), mesothelioma, schwannoma (including acoustic neuroma), meningioma, adenocarcinoma, melanoma, and leukemia or lymphoid malignancies. More particular examples of such cancers include squamous cell cancer (*e.g.* epithelial squamous cell cancer), lung cancer including small-cell lung cancer (SCLC), non-small cell lung cancer (NSCLC), adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer (including metastatic breast cancer), colon cancer, rectal cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, testicular cancer, esophageal cancer, tumors of the biliary tract, as well as head and neck cancer.

An “advanced” cancer is one which has spread outside the site or organ of origin, either by local invasion or metastasis.

A “recurrent” cancer is one which has regrown, either at the initial site or at a distant site, after a response to initial therapy.

5 The “pathology” of cancer includes all phenomena that compromise the well-being of the patient. This includes, without limitation, abnormal or uncontrollable cell growth, metastasis, interference with the normal functioning of neighboring cells, release of cytokines or other secretory products at abnormal levels, suppression or aggravation of inflammatory or immunological response, neoplasia, premalignancy, malignancy, invasion of surrounding or distant tissues or organs, such as
10 lymph nodes, etc.

Herein, a “subject” includes a mammalian and a human subject. The subject may be a “tumor subject” or a “cancer subject,” *i.e.* one who is suffering or at risk for suffering from one or more symptoms of tumor, such as cancer.

A “tumor sample” herein is a sample derived from, or comprising tumor cells from, a patient's
15 tumor. Examples of tumor samples herein include, but are not limited to, tumor biopsies, circulating tumor cells, circulating plasma proteins, ascitic fluid, primary cell cultures or cell lines derived from tumors or exhibiting tumor-like properties, as well as preserved tumor samples, such as formalin-fixed, paraffin-embedded tumor samples or frozen tumor samples.

A “fixed” tumor sample is one which has been histologically preserved using a fixative.

20 A “formalin-fixed” tumor sample is one which has been preserved using formaldehyde as the fixative.

An “embedded” tumor sample is one surrounded by a firm and generally hard medium such as paraffin, wax, celloidin, or a resin. Embedding makes possible the cutting of thin sections for microscopic examination or for generation of tissue microarrays (TMAs).

25 A “paraffin-embedded” tumor sample is one surrounded by a purified mixture of solid hydrocarbons derived from petroleum.

Herein, a “frozen” tumor sample refers to a tumor sample which is, or has been, frozen.

A cancer or biological sample which “displays HER expression, amplification, or activation”
30 is one which, in a diagnostic test, expresses (including overexpresses) a HER receptor, has amplified HER gene, and/or otherwise demonstrates activation or phosphorylation of a HER receptor.

A cancer or biological sample which “displays HER activation” is one which, in a diagnostic test, demonstrates activation or phosphorylation of a HER receptor. Such activation can be determined directly (*e.g.* by measuring HER phosphorylation by ELISA) or indirectly (*e.g.* by gene expression profiling or by detecting HER heterodimers, as described in U.S. patent application
35 publication No. 2004/0106161, published June 3, 2004).

Herein, “gene expression profiling” refers to an evaluation of expression of one or more genes as a surrogate for determining HER phosphorylation directly.

A “phospho-ELISA assay” herein is an assay in which phosphorylation of one or more HER receptors, especially HER2, is evaluated in an enzyme-linked immunosorbent assay (ELISA) using a reagent, usually an antibody, to detect phosphorylated HER receptor, substrate, or downstream signaling molecule. Preferably, an antibody which detects phosphorylated HER2 is used. The assay
5 may be performed on cell lysates, preferably from fresh or frozen biological samples.

A cancer cell with “HER receptor overexpression or amplification” is one which has significantly higher levels of a HER receptor protein or gene compared to a noncancerous cell of the same tissue type. Such overexpression may be caused by gene amplification or by increased transcription or translation. HER receptor overexpression or amplification may be determined in a
10 diagnostic or prognostic assay by evaluating increased levels of the HER protein present on the surface of a cell (*e.g.* via an immunohistochemistry assay; IHC). Alternatively, or additionally, one may measure levels of HER-encoding nucleic acid in the cell, *e.g.* via fluorescent *in situ* hybridization (FISH; see WO98/45479 published October, 1998), southern blotting, or polymerase chain reaction (PCR) techniques, such as quantitative real time PCR (qRT-PCR). One may also study HER receptor
15 overexpression or amplification by measuring shed antigen (*e.g.*, HER extracellular domain) in a biological fluid such as serum (see, *e.g.*, U.S. Patent No. 4,933,294 issued June 12, 1990; WO91/05264 published April 18, 1991; U.S. Patent 5,401,638 issued March 28, 1995; and Sias *et al.* *J. Immunol. Methods* 132: 73-80 (1990)). Aside from the above assays, various *in vivo* assays are available to the skilled practitioner. For example, one may expose cells within the body of the patient
20 to an antibody which is optionally labeled with a detectable label, *e.g.* a radioactive isotope, and binding of the antibody to cells in the patient can be evaluated, *e.g.* by external scanning for radioactivity or by analyzing a biopsy taken from a patient previously exposed to the antibody.

Herein, an “anti-tumor agent” refers to a drug used to treat cancer. Non-limiting examples of anti-tumor agents herein include chemotherapeutic agents, HER dimerization inhibitors, HER
25 antibodies, antibodies directed against tumor associated antigens, anti-hormonal compounds, cytokines, EGFR-targeted drugs, anti-angiogenic agents, tyrosine kinase inhibitors, growth inhibitory agents and antibodies, cytotoxic agents, antibodies that induce apoptosis, COX inhibitors, farnesyl transferase inhibitors, antibodies that binds oncofetal protein CA 125, HER2 vaccines, Raf or ras inhibitors, liposomal doxorubicin, topotecan, taxane, dual tyrosine kinase inhibitors, TLK286, EMD-
30 7200, pertuzumab, trastuzumab, erlotinib, and bevacizumab.

An “approved anti-tumor agent” is a drug used to treat cancer which has been accorded marketing approval by a regulatory authority such as the Food and Drug Administration (FDA) or foreign equivalent thereof.

Where an anti-tumor agent is administered as a “single anti-tumor agent” it is the only anti-
35 tumor agent administered to treat the cancer, *i.e.* it is not administered in combination with another anti-tumor agent, such as chemotherapy.

A “growth inhibitory agent” when used herein refers to a compound or composition which inhibits growth of a cell, especially a HER expressing cancer cell either *in vitro* or *in vivo*. Thus, the growth inhibitory agent may be one which significantly reduces the percentage of HER expressing cells in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxanes, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled “Cell cycle regulation, oncogenes, and antineoplastic drugs” by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13.

Examples of “growth inhibitory” antibodies are those which bind to HER2 and inhibit the growth of cancer cells overexpressing HER2. Preferred growth inhibitory HER2 antibodies inhibit growth of SK-BR-3 breast tumor cells in cell culture by greater than 20%, and preferably greater than 50% (*e.g.* from about 50% to about 100%) at an antibody concentration of about 0.5 to 30 µg/ml, where the growth inhibition is determined six days after exposure of the SK-BR-3 cells to the antibody (see U.S. Patent No. 5,677,171 issued October 14, 1997). The SK-BR-3 cell growth inhibition assay is described in more detail in that patent and hereinbelow. The preferred growth inhibitory antibody is a humanized variant of murine monoclonal antibody 4D5, *e.g.*, trastuzumab.

An antibody which “induces apoptosis” is one which induces programmed cell death as determined by binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies). The cell is usually one which overexpresses the HER2 receptor. Preferably the cell is a tumor cell, *e.g.* a breast, ovarian, stomach, endometrial, salivary gland, lung, kidney, colon, thyroid, pancreatic or bladder cell. *In vitro*, the cell may be a SK-BR-3, BT474, Calu 3 cell, MDA-MB-453, MDA-MB-361 or SKOV3 cell. Various methods are available for evaluating the cellular events associated with apoptosis. For example, phosphatidyl serine (PS) translocation can be measured by annexin binding; DNA fragmentation can be evaluated through DNA laddering; and nuclear/chromatin condensation along with DNA fragmentation can be evaluated by any increase in hypodiploid cells. Preferably, the antibody which induces apoptosis is one which results in about 2 to 50 fold, preferably about 5 to 50 fold, and most preferably about 10 to 50 fold, induction of annexin binding relative to untreated cell in an annexin binding assay using BT474 cells (see below). Examples of HER2 antibodies that induce apoptosis are 7C2 and 7F3.

The “epitope 4D5” is the region in the extracellular domain of HER2 to which the antibody 4D5 (ATCC CRL 10463) and trastuzumab bind. This epitope is close to the transmembrane domain of HER2, and within Domain IV of HER2. To screen for antibodies which bind to the 4D5 epitope, a

routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed to assess whether the antibody binds to the 4D5 epitope of HER2 (*e.g.* any one or more residues in the region from about residue 529 to about residue 625, inclusive of the HER2 ECD, residue numbering including signal peptide).

The “epitope 2C4” is the region in the extracellular domain of HER2 to which the antibody 2C4 binds. In order to screen for antibodies which bind to the 2C4 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Preferably the antibody blocks 2C4's binding to HER2 by about 50% or more. Alternatively, epitope mapping can be performed to assess whether the antibody binds to the 2C4 epitope of HER2. Epitope 2C4 comprises residues from Domain II in the extracellular domain of HER2. 2C4 and pertuzumab binds to the extracellular domain of HER2 at the junction of domains I, II and III. Franklin *et al.* *Cancer Cell* 5:317-328 (2004).

The “epitope 7C2/7F3” is the region at the N terminus, within Domain I, of the extracellular domain of HER2 to which the 7C2 and/or 7F3 antibodies (each deposited with the ATCC, see below) bind. To screen for antibodies which bind to the 7C2/7F3 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed to establish whether the antibody binds to the 7C2/7F3 epitope on HER2 (*e.g.* any one or more of residues in the region from about residue 22 to about residue 53 of the HER2 ECD, residue numbering including signal peptide).

“Treatment” refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with cancer as well as those in which cancer is to be prevented. Hence, the patient to be treated herein may have been diagnosed as having cancer or may be predisposed or susceptible to cancer.

The term “effective amount” refers to an amount of a drug effective to treat cancer in the patient. The effective amount of the drug may reduce the number of cancer cells; reduce the tumor size; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. The effective amount may extend progression free survival (*e.g.* as measured by Response Evaluation Criteria for Solid Tumors, RECIST, or CA-125 changes), result in an objective response (including a partial response, PR, or complete response, CR), increase overall survival time, and/or improve one or more symptoms of cancer (*e.g.* as assessed by FOSI).

The term “cytotoxic agent” as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive

isotopes (*e.g.* At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu), chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer.

- 5 Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN®); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and
- 10 bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scoplectin, and 9-aminocamptothecin); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins
- 15 (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine,
- 20 chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (*e. g.*, calicheamicin, especially calicheamicin gammaII and calicheamicin omegaII (see, *e.g.*, Agnew, *Chem Intl. Ed. Engl.*, 33: 183-186 (1994)); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin,
- 25 carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (including ADRIAMYCIN®, morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin, doxorubicin HCl liposome injection (DOXIL®) and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin,
- 30 puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate, gemcitabine (GEMZAR®), tegafur (UFTORAL®), capecitabine (XELODA®), an epothilone, and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; anti-adrenals
- 35 such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestabucil;

bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside (“Ara-C”); thiotepa; taxoid, *e.g.*, paclitaxel (TAXOL®), albumin-engineered nanoparticle formulation of paclitaxel (ABRAXANE™), and docetaxel (TAXOTERE®); chloranbucil; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); oxaliplatin; leucovovin; vinorelbine (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), etidronate (DIDROCAL®), NE-58095, zoledronic acid/zoledronate (ZOMETA®), alendronate (FOSAMAX®), pamidronate (AREDIA®), tiludronate (SKELID®), or risedronate (ACTONEL®); troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as THERATOPE® vaccine and gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; topoisomerase 1 inhibitor (*e.g.*, LURTOTECAN®); rmRH (*e.g.*, ABARELIX®); sorafenib (Bayer); SU-11248 (Pfizer); and pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovovin.

An “anti-hormonal agent” or “endocrine therapeutic” is an agent that acts to regulate, reduce, block, or inhibit the effects of hormones that can promote the growth of cancer. They may be hormones themselves. Examples include: anti-estrogens with mixed agonist/antagonist profile, including, tamoxifen (NOLVADEX®), 4-hydroxytamoxifen, toremifene (FARESTON®), idoxifene, droloxifene, raloxifene (EVISTA®), trioxifene, keoxifene, and selective estrogen receptor modulators (SERMs) such as SERM3; pure antiestrogens without agonist properties, such as EM800 (such agents may block estrogen receptor (ER) dimerization, inhibit DNA binding, increase ER turnover, and/or suppress ER levels); aromatase inhibitors, including steroidal aromatase inhibitors such as formestane and exemestane (AROMASIN®), and nonsteroidal aromatase inhibitors such as anastrozole (ARIMIDEX®), letrozole (FEMARA®) and aminoglutethimide, and other aromatase inhibitors

include vorozole (RIVISOR®), megestrol acetate (MEGASE®), fadrozole, and 4(5)-imidazoles; lutenizing hormone-releasing hormone agonists, including leuprolide (LUPRON® and ELIGARD®), goserelin, buserelin, and triptorelin; sex steroids, including progestins such as megestrol acetate and medroxyprogesterone acetate, estrogens such as diethylstilbestrol and premarin, and androgens/retinoids such as fluoxymesterone, all transretinoic acid and fenretinide; onapristone; anti-progesterones; estrogen receptor down-regulators (ERDs); anti-androgens such as flutamide, nilutamide and bicalutamide.

An “antimetabolite chemotherapeutic agent” is an agent which is structurally similar to a metabolite, but can not be used by the body in a productive manner. Many antimetabolite chemotherapeutic agents interfere with the production of the nucleic acids, RNA and DNA. Examples of antimetabolite chemotherapeutic agents include gemcitabine (GEMZAR®), 5-fluorouracil (5-FU), capecitabine (XELODAJ), 6-mercaptopurine, methotrexate, 6-thioguanine, pemetrexed, raltitrexed, arabinosylcytosine ARA-C cytarabine (CYTOSAR-U®), dacarbazine (DTIC-DOME®), azocytosine, deoxycytosine, pyrimidine, fludarabine (FLUDARA®), cladribine, 2-deoxy-D-glucose etc. The preferred antimetabolite chemotherapeutic agent is gemcitabine.

“Gemcitabine” or “2'-deoxy-2', 2'-difluorocytidine monohydrochloride (b-isomer)” is a nucleoside analogue that exhibits antitumor activity. The empirical formula for gemcitabine HCl is C₉H₁₁F₂N₃O₄ · HCl. Gemcitabine HCl is sold by Eli Lilly under the trademark GEMZAR®.

A “platinum-based chemotherapeutic agent” comprises an organic compound which contains platinum as an integral part of the molecule. Examples of platinum-based chemotherapeutic agents include carboplatin, cisplatin, and oxaliplatin.

By “platinum-based chemotherapy” is intended therapy with one or more platinum-based chemotherapeutic agents, optionally in combination with one or more other chemotherapeutic agents.

By “chemotherapy-resistant” cancer is meant that the cancer patient has progressed while receiving a chemotherapy regimen (*i.e.* the patient is “chemotherapy refractory”), or the patient has progressed within 12 months (for instance, within 6 months) after completing a chemotherapy regimen.

By “platinum-resistant” cancer is meant that the cancer patient has progressed while receiving platinum-based chemotherapy (*i.e.* the patient is “platinum refractory”), or the patient has progressed within 12 months (for instance, within 6 months) after completing a platinum-based chemotherapy regimen.

An “anti-angiogenic agent” refers to a compound which blocks, or interferes with to some degree, the development of blood vessels. The anti-angiogenic factor may, for instance, be a small molecule or antibody that binds to a growth factor or growth factor receptor involved in promoting angiogenesis. The preferred anti-angiogenic factor herein is an antibody that binds to vascular endothelial growth factor (VEGF), such as bevacizumab (AVASTIN®) (see U.S. Patent No. 6,884,879B1).

The term “cytokine” is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF- β ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1 α , IL-2 (e.g. PROLEUKIN[®]), IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12; a tumor necrosis factor such as TNF- α or TNF- β ; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

As used herein, the term “EGFR-targeted drug” refers to a therapeutic agent that binds to EGFR and, optionally, inhibits EGFR activation. Examples of such agents include antibodies and small molecules that bind to EGFR. Examples of antibodies which bind to EGFR include MAb 579 (ATCC CRL HB 8506), MAb 455 (ATCC CRL HB8507), MAb 225 (ATCC CRL 8508), MAb 528 (ATCC CRL 8509) (see, US Patent No. 4,943, 533, Mendelsohn *et al.*) and variants thereof, such as chimerized 225 (C225 or Cetuximab; ERBUTIX[®]) and reshaped human 225 (H225) (see, WO 96/40210, Imclone Systems Inc.); IMC-11F8, a fully human, EGFR-targeted antibody (Imclone); antibodies that bind type II mutant EGFR (US Patent No. 5,212,290); humanized and chimeric antibodies that bind EGFR as described in US Patent No. 5,891,996; and human antibodies that bind EGFR, such as ABX-EGF (see WO98/50433, Abgenix); EMD 55900 (Stragliotto *et al. Eur. J. Cancer* 32A:636-640 (1996)); EMD7200 (matuzumab) a humanized EGFR antibody directed against EGFR that competes with both EGF and TGF- α for EGFR binding; and mAb 806 or humanized mAb 806 (Johns *et al., J. Biol. Chem.* 279(29):30375-30384 (2004)). The anti-EGFR antibody may be conjugated with a cytotoxic agent, thus generating an immunoconjugate (see, *e.g.*, EP659,439A2, Merck Patent GmbH). Examples of small molecules that bind to EGFR include ZD1839 or Gefitinib (IRESSA; Astra Zeneca); CP-358774 or Erlotinib (TARCEVA[™]; Genentech/OSI); and AG1478, AG1571 (SU 5271; Sugen); EMD-7200.

A “tyrosine kinase inhibitor” is a molecule which inhibits tyrosine kinase activity of a tyrosine kinase such as a HER receptor. Examples of such inhibitors include the EGFR-targeted drugs noted in the preceding paragraph; small molecule HER2 tyrosine kinase inhibitor such as TAK165 available

from Takeda; CP-724,714, an oral selective inhibitor of the HER2 receptor tyrosine kinase (Pfizer and OSI); dual-HER inhibitors such as EKB-569 (available from Wyeth) which preferentially binds EGFR but inhibits both HER2 and EGFR-overexpressing cells; GW572016 (available from Glaxo) an oral HER2 and EGFR tyrosine kinase inhibitor; PKI-166 (available from Novartis); pan-HER inhibitors
 5 such as canertinib (CI-1033; Pharmacia); Raf-1 inhibitors such as antisense agent ISIS-5132 available from ISIS Pharmaceuticals which inhibits Raf-1 signaling; non-HER targeted TK inhibitors such as Imatinib mesylate (Gleevec[®]) available from Glaxo; MAPK extracellular regulated kinase I inhibitor CI-1040 (available from Pharmacia); quinazolines, such as PD 153035, 4-(3-chloroanilino)quinazoline; pyridopyrimidines; pyrimidopyrimidines; pyrrolopyrimidines, such as CGP 59326, CGP
 10 60261 and CGP 62706; pyrazolopyrimidines, 4-(phenylamino)-7H-pyrrolo[2,3-d] pyrimidines; curcumin (diferuloyl methane, 4,5-bis (4-fluoroanilino)phthalimide); tyrphostines containing nitrothiophene moieties; PD-0183805 (Warner-Lambert); antisense molecules (*e.g.* those that bind to HER-encoding nucleic acid); quinoxalines (US Patent No. 5,804,396); tryphostins (US Patent No. 5,804,396); ZD6474 (Astra Zeneca); PTK-787 (Novartis/Schering AG); pan-HER inhibitors such as
 15 CI-1033 (Pfizer); Affinitac (ISIS 3521; Isis/Lilly); Imatinib mesylate (Gleevec; Novartis); PKI 166 (Novartis); GW2016 (Glaxo SmithKline); CI-1033 (Pfizer); EKB-569 (Wyeth); Semaxinib (Sugen); ZD6474 (AstraZeneca); PTK-787 (Novartis/Schering AG); INC-1C11 (Imclone); or as described in any of the following patent publications: US Patent No. 5,804,396; WO99/09016 (American Cyanamid); WO98/43960 (American Cyanamid); WO97/38983 (Warner Lambert); WO99/06378
 20 (Warner Lambert); WO99/06396 (Warner Lambert); WO96/30347 (Pfizer, Inc); WO96/33978 (Zeneca); WO96/3397 (Zeneca); and WO96/33980 (Zeneca).

A “fixed” or “flat” dose of a therapeutic agent herein refers to a dose that is administered to a human patient without regard for the weight (WT) or body surface area (BSA) of the patient. The fixed or flat dose is therefore not provided as a mg/kg dose or a mg/m² dose, but rather as an absolute
 25 amount of the therapeutic agent.

A “loading” dose herein generally comprises an initial dose of a therapeutic agent administered to a patient, and is followed by one or more maintenance dose(s) thereof. Generally, a single loading dose is administered, but multiple loading doses are contemplated herein. Usually, the amount of loading dose(s) administered exceeds the amount of the maintenance dose(s) administered
 30 and/or the loading dose(s) are administered more frequently than the maintenance dose(s), so as to achieve the desired steady-state concentration of the therapeutic agent earlier than can be achieved with the maintenance dose(s).

A “maintenance” dose herein refers to one or more doses of a therapeutic agent administered to the patient over a treatment period. Usually, the maintenance doses are administered at spaced
 35 treatment intervals, such as approximately every week, approximately every 2 weeks, approximately every 3 weeks, or approximately every 4 weeks.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, typically: (1)

employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" may be identified as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

In the context of the present invention, reference to "at least one," "at least two," "at least three," "at least four," "at least five," etc. of the genes listed in any particular gene set means any one or any and all combinations of the genes listed.

Detailed Description

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, and

biochemistry, which are within the skill of the art. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", 2nd edition (Sambrook et al., 1989); "Oligonucleotide Synthesis" (M.J. Gait, ed., 1984); "Animal Cell Culture" (R.I. Freshney, ed., 1987); "Methods in Enzymology" (Academic Press, Inc.); "Handbook of Experimental Immunology", 4th edition (D.M. Weir & C.C. Blackwell, eds., Blackwell Science Inc., 1987); "Gene Transfer Vectors for Mammalian Cells" (J.M. Miller & M.P. Calos, eds., 1987); "Current Protocols in Molecular Biology" (F.M. Ausubel et al., eds., 1987); and "PCR: The Polymerase Chain Reaction", (Mullis et al., eds., 1994).

Identification of diagnostic markers of resistance to treatment with HER2 inhibitors

As discussed above, trastuzumab is used in clinical practice both in the adjuvant and the metastatic setting to treat breast cancer in patients whose tumor overexpresses the HER2 oncogene.

Currently, HER2 expression levels are typically measured by two main types of assay, immunohistochemistry (IHC) and fluorescent in situ hybridization (FISH). Thus, HER2 overexpression may be analyzed by IHC, *e.g.* using the HERCEPTEST® (Dako). Paraffin embedded tissue sections from a tumor biopsy may be subjected to the IHC assay and accorded HER2 protein staining intensity criteria as follows:

Score 0 - no staining is observed or membrane staining is observed in less than 10% of tumor cells.

Score 1+ a faint/barely perceptible membrane staining is detected in more than 10% of the tumor cells. The cells are only stained in part of their membrane.

Score 2+ a weak to moderate complete membrane staining is observed in more than 10% of the tumor cells.

Score 3+ a moderate to strong complete membrane staining is observed in more than 10% of the tumor cells.

Those tumors with 0 or 1+ scores for HER2 overexpression assessment may be characterized as not overexpressing HER2, whereas those tumors with 2+ or 3+ scores may be characterized as overexpressing HER2.

Tumors overexpressing HER2 may be rated by immunohistochemical scores corresponding to the number of copies of HER2 molecules expressed per cell, and can be determined biochemically:

0 = 0-10,000 copies/cell,

1+ = at least about 200,000 copies/cell,

2+ = at least about 500,000 copies/cell,

3+ = at least about 2,000,000 copies/cell.

Overexpression of HER2 at the 3+ level, which leads to ligand-independent activation of the tyrosine kinase (Hudziak *et al.*, *Proc. Natl. Acad. Sci. USA*, 84:7159-7163 (1987)), occurs in approximately 30% of breast cancers, and in these patients, relapse-free survival and overall survival

are diminished (Slamon *et al.*, *Science*, 244:707-712 (1989); Slamon *et al.*, *Science*, 235:177-182 (1987)).

Alternatively, or additionally, FISH assays such as the INFORM™ (sold by Ventana, Arizona) or PATHVISION™ (Vysis, Illinois) may be carried out on formalin-fixed, paraffin-
5 embedded tumor tissue to determine the extent (if any) of HER2 amplification in the tumor.

For review, see Winston *et al.*, *Am J. Pathol* 121(Suppl. 1):S33-49 (2004).

In patients with metastatic breast cancer, approximately 30% of patients who test positive for HER2 either by IHC or FISH (i.e. patients with HER2-expressing tumors) exhibit an objective response to trastuzumab alone, and about 50% to trastuzumab plus chemotherapy. Some of the
10 remaining patients may still derive clinical benefit without an objective response, but there still remains a proportion of patients that exhibit primary resistance to trastuzumab. Furthermore, many patients that do benefit initially in the metastatic setting eventually progress while on trastuzumab treatment (acquired resistance). Patients with primary or acquired resistance to treatment with trastuzumab and collectively referred to as “refractory” or “resistant” to such treatment. In the
15 adjuvant setting, the addition of trastuzumab to chemotherapy results in a significant improvement in disease-free survival. Nevertheless, there is still a group of patients whose tumor recurs after treatment.

The present invention is based on the identification of genes that are associated with trastuzumab resistance. Accordingly, the expression levels of such genes can serve as diagnostic
20 markers to identify patients with HER2 expressing tumors who are less likely to respond to current therapies with trastuzumab or other HER2 inhibitors, and might benefit from novel combination treatments including trastuzumab or other HER2 inhibitors in combination with other anti-cancer agents and/or other treatment modalities.

It is well known that kinases and phosphatases control the reversible process of
25 phosphorylation and are dysregulated in a variety of diseases, including cancer. Accordingly, a large-scale RNAi approach was elected to identify kinases and phosphatases that are associated with resistance to treatment with trastuzumab. In particular, performing a large-scale siRNA screen on HER2 positive cell lines that are sensitive to trastuzumab treatment *in vitro*, a group of kinases and phosphatases has been identified whose loss of function turned the cell lines resistant to treatment
30 with trastuzumab. The results were validated by re-assaying the siRNA and by confirming the results in two different cell lines (BT474 and SKBR3). Details of this screen are provided in the Example below.

Thus, according to the present invention, the following genes have been identified as being associated with resistance to treatment with HER2 inhibitors: CDK11, DYRK1A, LATS2, STK10,
35 Wee1, DUSP4, DUSP6, HIPK3, JNK, MAP4K4, PTPN11, Socs5, PPM1H, DKFZP586B16, DGKI, FLJ35107, FLT1, HK2, ITK, MOAP1, KIAA0685, KIAA1639, LIM/PDLIM5, PANK1, P14K2B, PPP2R1A, PRKWINK3, RYK, SPEC2, STK22C, STYK1, TXND3. These genes are also listed in

Table 1, along with their NCBI GenBank accession numbers. Reduced expression or activity of one or more of these genes, or the corresponding RNA molecules or encoded proteins in a biological sample obtained from the patient, relative to control, indicates that the patient's tumor is likely to show resistance to treatment with a HER2 inhibitor.

5 The control can, for example, be a gene, present in the same cell, which is known to be down-regulated in patients showing resistance to HER2 inhibitor treatment (positive control), such as, for example, p27 or PTEN. Alternatively, or in addition, the control can be the expression level of the same gene in a normal cell of the same cell type (negative control). Expression levels can also be normalized, for example, to the expression levels of housekeeping genes, such as glyceraldehyde-3-
 10 phosphate-dehydrogenase (GAPDH) and/or β -actin, or to the expression levels of all genes in the sample tested. In one embodiment, expression of one or more of the above noted genes is deemed positive expression if it is at the median or above, e.g. compared to other samples of the same tumor-type. The median expression level can be determined essentially contemporaneously with measuring gene expression, or may have been determined previously. These and other methods are well known
 15 in the art, and are apparent to those skilled in the art.

Although the present invention identifies specific markers of tumor resistance to treatment with a HER2 inhibitor, surrogate markers the expression of which positively or negatively coordinately regulated with the expression of a gene specifically disclosed herein, are also suitable as resistance markers. Thus, surrogate markers include genes that are positive regulators of the same
 20 pathway as the pathway positively regulated by a gene specifically identified herein, or a downstream pathway. The lower expression (inactivation or inhibition) of such genes will be a predictor of resistance of HER2 expressing tumors to treatment with HER2 inhibitors. Included within this group are genes which show a similar expression pattern to a gene specifically disclosed herein, where the similar expression pattern may, for example, result from involvement of both genes in a particular
 25 biological process and/or being under common regulatory control in tumor cells. Surrogate markers also include genes the expression of which inversely correlates with the expression of a gene specifically identified herein, i.e. genes the expression of which is negatively coordinately regulated with a specifically disclosed gene. Included in this group of surrogate markers are genes which are negative regulators of the same pathway as a pathway positively regulated by a gene specifically
 30 identified herein, or a downstream pathway. The higher expression (activation or upregulation) of such genes will be a predictor of resistance of HER2 expressing tumors to treatment with HER2 inhibitors.

Diagnostic Methods

35 Methods for identifying patients for treatment with HER2 inhibitors, such as HER2 antibodies have been discussed above. Of this patient population, patients who are likely to be resistant or not repond well to such treatment can be identified by determining the expression level of one or more of the genes, the corresponding RNA molecules or encoded proteins in a biological sample comprising

tumor cells obtained from the patient. The biological sample can, for example, be a fresh or frozen or archived paraffin-embedded and fixed (e.g. formalin-fixed) tissue sample, routinely prepared and preserved in everyday clinical practice. The biological sample can also be a different sample obtained from the patient, such as a biological fluid, including, without limitation, blood, urine, saliva, ascites fluid, or derivatives such as blood serum and blood plasma, and the like.

Various methods for determining expression of mRNA or protein include, but are not limited to, gene expression profiling, polymerase chain reaction (PCR) including quantitative real time PCR (qRT-PCR), microarray analysis that can be performed by commercially available equipment, following manufacturer's protocols, such as by using the Affymetrix GenChip technology, serial analysis of gene expression (SAGE) (Velculescu *et al.*, *Science* 270:484-487 (1995); and Velculescu *et al.*, *Cell* 88:243-51 (1997)), MassARRAY, Gene Expression Analysis by Massively Parallel Signature Sequencing (MPSS) (Brenner *et al.*, *Nature Biotechnology* 18:630-634 (2000)), proteomics, immunohistochemistry (IHC), etc. Preferably mRNA is quantified. Such mRNA analysis is preferably performed using the technique of polymerase chain reaction (PCR), or by microarray analysis. Where PCR is employed, a preferred form of PCR is quantitative real time PCR (qRT-PCR).

The steps of a representative protocol for profiling gene expression using fixed, paraffin-embedded tissues as the RNA source, including mRNA isolation, purification, primer extension and amplification are given in various published journal articles (for example: Godfrey *et al.* *J. Molec. Diagnostics* 2: 84-91 (2000); Specht *et al.*, *Am. J. Pathol.* 158: 419-29 (2001)). Briefly, a representative process starts with cutting about 10 microgram thick sections of paraffin-embedded tumor tissue samples. The mRNA is then extracted, and protein and DNA are removed. General methods for mRNA extraction are well known in the art and are disclosed in standard textbooks of molecular biology, including Ausubel *et al.*, *Current Protocols of Molecular Biology*, John Wiley and Sons (1997). Methods for RNA extraction from paraffin embedded tissues are disclosed, for example, in Rupp and Locker, *Lab Invest.* 56:A67 (1987), and De Andrés *et al.*, *BioTechniques* 18:42044 (1995). In particular, RNA isolation can be performed using purification kit, buffer set and protease from commercial manufacturers, such as Qiagen, according to the manufacturer's instructions. For example, total RNA from cells in culture can be isolated using Qiagen RNeasy mini-columns. Other commercially available RNA isolation kits include MasterPure™ Complete DNA and RNA Purification Kit (EPICENTRE®, Madison, WI), and Paraffin Block RNA Isolation Kit (Ambion, Inc.). Total RNA from tissue samples can be isolated using RNA Stat-60 (Tel-Test). RNA prepared from tumor can be isolated, for example, by cesium chloride density gradient centrifugation. After analysis of the RNA concentration, RNA repair and/or amplification steps may be included, if necessary, and RNA is reverse transcribed using gene specific promoters followed by PCR. Preferably, real time PCR is used, which is compatible both with quantitative competitive PCR, where internal competitor for each target sequence is used for normalization, and with quantitative

comparative PCR using a normalization gene contained within the sample, or a housekeeping gene for RT-PCR. For further details see, e.g. "PCR: The Polymerase Chain Reaction", Mullis et al., eds., 1994; and Held *et al.*, *Genome Research* 6:986-994 (1996). Finally, the data are analyzed to identify the best treatment option(s) available to the patient on the basis of the characteristic gene expression pattern identified in the tumor sample examined.

Expression levels can also be determined at the protein level, for example, using various types of immunoassays or proteomics techniques.

In immunoassays, the target diagnostic protein marker is detected by using an antibody specifically binding to the marker. The antibody typically will be labeled with a detectable moiety.

Numerous labels are available which can be generally grouped into the following categories:

Radioisotopes, such as ^{35}S , ^{14}C , ^{125}I , ^3H , and ^{131}I . The antibody can be labeled with the radioisotope using the techniques described in *Current Protocols in Immunology*, Volumes 1 and 2, Coligan et al. (1991) Ed. Wiley-Interscience, New York, New York, Pubs. for example and radioactivity can be measured using scintillation counting.

Fluorescent labels such as rare earth chelates (europium chelates) or fluorescein and its derivatives, rhodamine and its derivatives, dansyl, Lissamine, phycoerythrin and Texas Red are available. The fluorescent labels can be conjugated to the antibody using the techniques disclosed in *Current Protocols in Immunology*, *supra*, for example. Fluorescence can be quantified using a fluorimeter.

Various enzyme-substrate labels are available and U.S. Patent No. 4,275,149 provides a review of some of these. The enzyme generally catalyzes a chemical alteration of the chromogenic substrate which can be measured using various techniques. For example, the enzyme may catalyze a color change in a substrate, which can be measured spectrophotometrically. Alternatively, the enzyme may alter the fluorescence or chemiluminescence of the substrate. Techniques for quantifying a change in fluorescence are described above. The chemiluminescent substrate becomes electronically excited by a chemical reaction and may then emit light which can be measured (using a chemiluminometer, for example) or donates energy to a fluorescent acceptor. Examples of enzymatic labels include luciferases (*e.g.*, firefly luciferase and bacterial luciferase; U.S. Patent No. 4,737,456), luciferin, 2,3-dihydrophthalazinediones, malate dehydrogenase, urease, peroxidase such as horseradish peroxidase (HRPO), alkaline phosphatase, β -galactosidase, glucoamylase, lysozyme, saccharide oxidases (*e.g.*, glucose oxidase, galactose oxidase, and glucose-6-phosphate dehydrogenase), heterocyclic oxidases (such as uricase and xanthine oxidase), lactoperoxidase, microperoxidase, and the like. Techniques for conjugating enzymes to antibodies are described in O'Sullivan et al. (1981) *Methods for the Preparation of Enzyme-Antibody Conjugates for use in Enzyme Immunoassay*, in *Methods in Enzym.* (ed J. Langone & H. Van Vunakis), Academic press, New York 73:147-166.

Examples of enzyme-substrate combinations include, for example:

Horseradish peroxidase (HRPO) with hydrogen peroxidase as a substrate, wherein the hydrogen peroxidase oxidizes a dye precursor (*e.g.*, orthophenylene diamine (OPD) or 3,3',5,5'-tetramethyl benzidine hydrochloride (TMB));

alkaline phosphatase (AP) with para-Nitrophenyl phosphate as chromogenic substrate; and
5 β -D-galactosidase (β -D-Gal) with a chromogenic substrate (*e.g.*, p-nitrophenyl- β -D-galactosidase) or fluorogenic substrate 4-methylumbelliferyl- β -D-galactosidase.

Numerous other enzyme-substrate combinations are available to those skilled in the art. For a general review of these, see U.S. Patent Nos. 4,275,149 and 4,318,980.

Sometimes, the label is indirectly conjugated with the antibody. The skilled artisan will be
10 aware of various techniques for achieving this. For example, the antibody can be conjugated with biotin and any of the three broad categories of labels mentioned above can be conjugated with avidin, or *vice versa*. Biotin binds selectively to avidin and thus, the label can be conjugated with the antibody in this indirect manner. Alternatively, to achieve indirect conjugation of the label with the antibody, the antibody is conjugated with a small hapten (*e.g.*, digoxin) and one of the different types of labels
15 mentioned above is conjugated with an anti-hapten antibody (*e.g.*, anti-digoxin antibody). Thus, indirect conjugation of the label with the antibody can be achieved.

In other versions of immunoassay techniques, the antibody need not be labeled, and the presence thereof can be detected using a labeled antibody which binds to the antibody.

Thus, the diagnostic immunoassays herein may be in any assay format, including, for
20 example, competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp.147-158 (CRC Press, Inc. 1987).

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyze for binding with a limited amount of antibody. The amount of antigen in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To
25 facilitate determining the amount of standard that becomes bound, the antibodies generally are insolubilized before or after the competition, so that the standard and analyze that are bound to the antibodies may conveniently be separated from the standard and analyze which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample
30 analyze is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyze, thus forming an insoluble three-part complex. See, *e.g.*, U.S. Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in
35 which case the detectable moiety is an enzyme.

Protein levels can also be detected using proteomics techniques. The term "proteome" is defined as the totality of the proteins present in a sample (*e.g.* tissue, organism, or cell culture) at a

certain point of time. Proteomics includes, among other things, study of the global changes of protein expression in a sample (also referred to as “expression proteomics”). Proteomics typically includes the following steps: (1) separation of individual proteins in a sample by 2-D gel electrophoresis (2-D PAGE); (2) identification of the individual proteins recovered from the gel, e.g. by mass spectrometry or N-terminal sequencing, and (3) analysis of the data using bioinformatics. Proteomics methods are valuable alternatives or supplements to other methods of gene expression profiling, and can be used, alone or in combination with other methods, to detect the products of the tumor resistance markers of the present invention.

Preferred markers of the present invention, identified by the kinase library screen, include DYRK1A, HK2, Socs5, STK10, Kila1639, and MAP4K4. A particularly preferred group of kinase markers includes DYRK1A, HK2, Socs5, and STK10. Members of these groups, as single markers or in any combination, are preferred for use in the diagnostic assays of the present invention.

Preferred markers, identified by the phosphatase library screen, include PTPN11, KIAA0685, and PPM1H. These markers, as single markers or any any combination, are preferred for use in the diagnostic assays of the present invention.

Measurement of biomarker expression levels may be performed by using a software program executed by a suitable processor. Suitable software and processors are well known in the art and are commercially available. The program may be embodied in software stored on a tangible medium such as CD-ROM, a floppy disk, a hard drive, a DVD, or a memory associated with the processor, but persons of ordinary skill in the art will readily appreciate that the entire program or parts thereof could alternatively be executed by a device other than a processor, and/or embodied in firmware and/or dedicated hardware in a well known manner.

Following the measurement of the expression levels of the genes identified herein, or their expression products, and the determination that a subject is likely or not likely to respond to treatment with a HER2 inhibitor, the assay results, findings, diagnoses, predictions and/or treatment recommendations are typically recorded and communicated to technicians, physicians and/or patients, for example. In certain embodiments, computers will be used to communicate such information to interested parties, such as, patients and/or the attending physicians. In some embodiments, the assays will be performed or the assay results analyzed in a country or jurisdiction which differs from the country or jurisdiction to which the results or diagnoses are communicated.

In a preferred embodiment, a diagnosis, prediction and/or treatment recommendation based on the expression level in a test subject of one or more of the biomarkers herein is communicated to the subject as soon as possible after the assay is completed and the diagnosis and/or prediction is generated. The results and/or related information may be communicated to the subject by the subject's treating physician. Alternatively, the results may be communicated directly to a test subject by any means of communication, including writing, electronic forms of communication, such as email, or telephone. Communication may be facilitated by use of a computer, such as in case of email

communications. In certain embodiments, the communication containing results of a diagnostic test and/or conclusions drawn from and/or treatment recommendations based on the test, may be generated and delivered automatically to the subject using a combination of computer hardware and software which will be familiar to artisans skilled in telecommunications. One example of a healthcare-oriented communications system is described in U.S. Pat. No. 6,283,761; however, the present invention is not limited to methods which utilize this particular communications system. In certain embodiments of the methods of the invention, all or some of the method steps, including the assaying of samples, diagnosing of diseases, and communicating of assay results or diagnoses, may be carried out in diverse (e.g., foreign) jurisdictions.

Identification of HER2 inhibitors

The first step in identifying inhibitors of a HER2 polypeptide, is typically *in vitro* screening to identify compounds that selectively bind HER2. The binding affinity of the candidate compounds can be tested by direct binding (see, e.g. Schoemaker et al., J. Pharmacol. Exp. Ther., 285:61-69 (1983)) or by indirect, e.g. competitive, binding. In competitive binding experiments, the concentration of a compound necessary to displace 50% of another compound bound to the target polypeptide (IC₅₀) is usually used as a measure of binding affinity. If the test compound binds HER2 selectively and with high affinity, displacing another compound bound to HER2, such as a HER2 antibody, it is identified as HER2 inhibitor. Cell based assays can be used in a similar manner.

A preferred group of HER2 inhibitors includes antibodies specifically binding to HER2. Antibody "binding affinity" may be determined by equilibrium methods (e.g. enzyme-linked immunoabsorbent assay (ELISA) or radioimmunoassay (RIA)), or kinetics (e.g. BIACORE™ analysis), for example. Also, the antibody may be subjected to other "biological activity assays", e.g., in order to evaluate its "potency" or pharmacological activity and potential efficacy as a therapeutic agent. Such assays are known in the art and depend on the target antigen and intended use for the antibody.

Other HER2 inhibitors include peptide and non-peptide small molecules, and antisense, ribozyme and triple helix molecules.

Non-antibody HER2 inhibitors, such as peptide and non-peptide small molecule inhibitors of HER2, can be identified by binding or interaction assays, well known in the art.

All binding assays for inhibitors are common in that they call for contacting the candidate inhibitor with a HER2 polypeptide under conditions and for a time sufficient to allow these two components to interact. In binding assays, the interaction is binding, and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, either the HER2 or the candidate inhibitor is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the HER2 polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the HER2 polypeptide to be immobilized can be used to

anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the
5 originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

If the candidate compound is a polypeptide which interacts with but does not bind to HER2,
10 the interaction of HER2 with the respective polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, Nature (London), 340:245-246 (1989); Chien et
15 al., Proc. Natl. Acad. Sci. USA, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, Proc. Natl. Acad. Sci. USA, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this, and
20 employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit
25 (MATCHMAKERTM) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

It is emphasized that the screening assays specifically discussed herein are for illustration
30 only. A variety of other assays, which can be selected depending on the type of the antagonist candidates screened (e.g. polypeptides, peptides, non-peptide small organic molecules, nucleic acid, etc.) are well known to those skilled in the art and are equally suitable for the purposes of the present invention.

The assays described herein may be used to screen libraries of compounds, including, without
35 limitation, chemical libraries, natural product libraries (e.g. collections of microorganisms, animals, plants, etc.), and combinatorial libraries comprised of random peptides, oligonucleotides or small organic molecules. In a particular embodiment, the assays herein are used to screen antibody libraries,

including, without limitation, naïve human, recombinant, synthetic and semi-synthetic antibody libraries. The antibody library can, for example, be a phage display library, including monovalent libraries, displaying on average one single-chain antibody or antibody fragment per phage particle, and multi-valent libraries, displaying, on average, two or more antibodies or antibody fragments per viral particle. However, the antibody libraries to be screened in accordance with the present invention are not limited to phage display libraries. Other display techniques include, for example, ribosome or mRNA display (Mattheakis et al., Proc. Natl. Acad. Sci. USA 91:9022-9026 (1994); Hanes and Pluckthun, Proc. Natl. Acad. Sci. USA 94:4937-4942 (1997)), microbial cell display, such as bacterial display (Georgiou et al., Nature Biotech. 15:29-34 (1997)), or yeast cell display (Kieckhefer et al., Protein Eng. 10:1303-1310 (1997)), display on mammalian cells, spore display, viral display, such as retroviral display (Urban et al., Nucleic Acids Res. 33:e35 (2005)), display based on protein-DNA linkage (Odegrip et al., Proc. Acad. Natl. Sci. USA 101:2806-2810 (2004); Reiersen et al., Nucleic Acids Res. 33:e10 (2005)), and microbead display (Sepp et al., FEBS Lett. 532:455-458 (2002)). Libraries of other molecules, such as combinatorial libraries of synthetic small molecules can also be screened in a similar manner.

HER2 inhibitors can also be designed to reduce the level of endogenous HER2 gene expression, for example, by using well-known antisense or ribozyme approaches to inhibit or prevent translation of HER2 mRNA or triple helix approaches to inhibit transcription of HER2 genes. Such antisense, ribozyme, and triple helix antagonists may be designed to reduce or inhibit either unimpaired, or if appropriate, mutant HER2 gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art.

Antisense RNA and DNA molecules can act to directly block the translation of mRNA by hybridizing to targeted endogenous mRNA thereby preventing translation. Alternatively, antisense RNA or DNA can inhibit or prevent transcription of the target gene. The antisense approach involves designing oligonucleotides (either DNA or RNA) that are complementary to a HER2 mRNA, or complementary to a portion of the target gene, such as a regulatory element that controls transcription of the gene. Typically, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length.

Production of antibodies

Since, in the preferred embodiment, the HER2 inhibitor is an antibody, a description follows as to exemplary techniques for the production of HER antibodies used in accordance with the present invention. The HER antigen to be used for production of antibodies may be, *e.g.*, a soluble form of the extracellular domain of a HER receptor or a portion thereof, containing the desired epitope. Alternatively, cells expressing HER at their cell surface (*e.g.* NIH-3T3 cells transformed to overexpress HER2; or a carcinoma cell line such as SK-BR-3 cells, see Stancovski *et al.* PNAS (USA) 88:8691-8695 (1991)) can be used to generate antibodies. Other forms of HER receptor useful for generating antibodies will be apparent to those skilled in the art.

Polyclonal antibodies

Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfo succinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are different alkyl groups.

Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, *e.g.*, 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

Monoclonal antibodies

Various methods for making monoclonal antibodies herein are available in the art. For example, the monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived

from MOPC-21 and MPC-11 mouse tumors available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); and Brodeur *et al.*, *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp.59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional antibody purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce antibody protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Plückthun, *Immunol. Revs.*, 130:151-188 (1992).

In a further embodiment, monoclonal antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage

libraries (Waterhouse *et al.*, *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

The DNA also may be modified, for example, by substituting the coding sequence for human heavy chain and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; and Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

Humanized antibodies

Methods for humanizing non-human antibodies have been described in the art. Preferably, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, *Nature*, 321:522-525 (1986); Riechmann *et al.*, *Nature*, 332:323-327 (1988); Verhoeven *et al.*, *Science*, 239:1534-1536 (1988)), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework region (FR) for the humanized antibody (Sims *et al.*, *J. Immunol.*, 151:2296 (1993); Chothia *et al.*, *J. Mol. Biol.*, 196:901 (1987)). Another method uses a particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred

method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

US Patent No. 6,949,245 describes production of exemplary humanized HER2 antibodies which bind HER2 and block ligand activation of a HER receptor. The humanized antibody of particular interest herein blocks EGF, TGF- α and/or HRG mediated activation of MAPK essentially as effectively as murine monoclonal antibody 2C4 (or a Fab fragment thereof) and/or binds HER2 essentially as effectively as murine monoclonal antibody 2C4 (or a Fab fragment thereof). The humanized antibody herein may, for example, comprise nonhuman hypervariable region residues incorporated into a human variable heavy domain and may further comprise a framework region (FR) substitution at a position selected from the group consisting of 69H, 71H and 73H utilizing the variable domain numbering system set forth in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991). In one embodiment, the humanized antibody comprises FR substitutions at two or all of positions 69H, 71H and 73H.

An exemplary humanized antibody of interest herein comprises variable heavy domain complementarity determining residues GFTFTDYTMX, where X is preferably D or S (SEQ ID NO:1); DVNPNSGGSIYNQRFKG (SEQ ID NO:2); and/or NLGPSFYFDY (SEQ ID NO:3), optionally comprising amino acid modifications of those CDR residues, *e.g.* where the modifications essentially maintain or improve affinity of the antibody. For example, the antibody variant of interest may have from about one to about seven or about five amino acid substitutions in the above variable heavy CDR sequences. Such antibody variants may be prepared by affinity maturation, *e.g.*, as described below.

The humanized antibody may comprise variable light domain complementarity determining residues KASQDVSIGVA (SEQ ID NO:4); SASYX¹X²X³, where X¹ is preferably R or L, X² is preferably Y or E, and X³ is preferably T or S (SEQ ID NO:5); and/or QQYYIYPYT (SEQ ID NO:6), *e.g.* in addition to those variable heavy domain CDR residues in the preceding

paragraph. Such humanized antibodies optionally comprise amino acid modifications of the above CDR residues, *e.g.* where the modifications essentially maintain or improve affinity of the antibody. For example, the antibody variant of interest may have from about one to about seven or about five amino acid substitutions in the above variable light CDR sequences. Such antibody variants may be prepared by affinity maturation, *e.g.*, as described below.

The present application also contemplates affinity matured antibodies which bind HER2 and block ligand activation of a HER receptor. The affinity matured antibody preferably binds to HER2 receptor with an affinity superior to that of murine 2C4 or pertuzumab (*e.g.* from about two or about four fold, to about 100 fold or about 1000 fold improved affinity, *e.g.* as assessed using a HER2-extracellular domain (ECD) ELISA). Exemplary variable heavy CDR residues for substitution include H28, H30, H34, H35, H64, H96, H99, or combinations of two or more (*e.g.* two, three, four, five, six, or seven of these residues). Examples of variable light CDR residues for alteration include L28, L50, L53, L56, L91, L92, L93, L94, L96, L97 or combinations of two or more (*e.g.* two to three, four, five or up to about ten of these residues).

Various forms of the humanized antibody or affinity matured antibody are contemplated. For example, the humanized antibody or affinity matured antibody may be an antibody fragment, such as a Fab, which is optionally conjugated with one or more cytotoxic agent(s) in order to generate an immunoconjugate. Alternatively, the humanized antibody or affinity matured antibody may be an intact antibody, such as an intact IgG1 antibody.

Human antibodies

As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits *et al.*, *Nature*, 362:255-258 (1993); Bruggermann *et al.*, *Year in Immuno.*, 7:33 (1993); and U.S. Patent Nos. 5,591,669, 5,589,369 and 5,545,807. Alternatively, phage display technology (McCafferty *et al.*, *Nature* 348:552-553 (1990)) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody

fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B-cell. Phage display can be performed in a variety of formats; for their review see, *e.g.*, Johnson, Kevin S. and Chiswell, David J., *Current Opinion in Structural Biology* 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson *et al.*, *Nature*, 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks *et al.*, *J. Mol. Biol.* 222:581-597 (1991), or Griffith *et al.*, *EMBO J.* 12:725-734 (1993). See, also, U.S. Patent Nos. 5,565,332 and 5,573,905.

As discussed above, human antibodies may also be generated by *in vitro* activated B cells (see U.S. Patents 5,567,610 and 5,229,275).

Human HER2 antibodies are described in U.S. Patent No. 5,772,997 issued June 30, 1998 and WO 97/00271 published January 3, 1997.

Antibody fragments

Various techniques have been developed for the production of antibody fragments comprising one or more antigen binding regions. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, *e.g.*, Morimoto *et al.*, *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992); and Brennan *et al.*, *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, *Bio/Technology* 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; U.S. Patent No. 5,571,894; and U.S. Patent No. 5,587,458. The antibody fragment may also be a Alinear antibody@, *e.g.*, as described in U.S. Patent 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

Bispecific antibodies

Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the HER2 protein. Other such antibodies may combine a HER2 binding site with binding site(s) for EGFR, HER3 and/or HER4. Alternatively, a HER2 arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (*e.g.* CD2 or CD3), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular

defense mechanisms to the HER2-expressing cell. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express HER2. These antibodies possess a HER2-binding arm and an arm which binds the cytotoxic agent (*e.g.* saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length
5 antibodies or antibody fragments (*e.g.* F(ab')₂ bispecific antibodies).

WO 96/16673 describes a bispecific HER2/Fc γ RIII antibody and U.S. Patent No. 5,837,234 discloses a bispecific HER2/Fc γ RI antibody IDM1 (Osidem). A bispecific HER2/Fc α antibody is shown in WO98/02463. U.S. Patent No. 5,821,337 teaches a bispecific HER2/CD3 antibody. MDX-210 is a bispecific HER2-Fc γ RIII Ab.

10 Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein *et al.*, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which
15 only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain
20 sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into
25 a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular
30 significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific
35 compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific

antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

According to another approach described in U.S. Patent No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the C_H3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (*e.g.* tyrosine or tryptophan). Compensatory “cavities” of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (*e.g.* alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or “heteroconjugate” antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, *Science*, 229: 81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby *et al.*, *J. Exp. Med.*, 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the HER2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, *J. Immunol.*, 148(5):1547-1553 (1992). The leucine

zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*,
 5 *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming
 10 two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber *et al.*, *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.* *J. Immunol.* 147: 60 (1991).

Other amino acid sequence modifications

15 Amino acid sequence modification(s) of the antibodies described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of the antibody are prepared by introducing appropriate nucleotide changes into the antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the
 20 amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter post-translational processes of the antibody, such as changing the number or position of glycosylation sites.

A useful method for identification of certain residues or regions of the antibody that are
 25 preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells *Science*, 244:1081-1085 (1989). Here, a residue or group of target residues are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with antigen. Those amino acid locations demonstrating functional sensitivity to the
 30 substitutions then are refined by introducing further or other variants at, or for, the sites of substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed antibody variants are screened for the desired activity.

35 Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions

include antibody with an N-terminal methionyl residue or the antibody fused to a cytotoxic polypeptide. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.* for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

5 Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the antibody molecule replaced by a different residue. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 1 under the heading of “preferred substitutions”. If such substitutions result in a change in biological activity, then more substantial
10 changes, denominated “exemplary substitutions” in the following table, or as further described below in reference to amino acid classes, may be introduced and the products screened.

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in the biological properties of the antibody are accomplished by
15 selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the

polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Amino acids may be grouped according to similarities in the properties of their side chains (in A. L. Lehninger, in *Biochemistry*, second ed., pp. 73-75, Worth Publishers, New York (1975)):

- 5 non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M)
 uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q)
 acidic: Asp (D), Glu (E)
 basic: Lys (K), Arg (R), His(H)

Alternatively, naturally occurring residues may be divided into groups based on common
 10 side-chain properties:

-
- hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;
 neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
 acidic: Asp, Glu;
 basic: His, Lys, Arg;
 15 residues that influence chain orientation: Gly, Pro;
 aromatic: Trp, Tyr, Phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

Any cysteine residue not involved in maintaining the proper conformation of the antibody
 20 also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (*e.g.* a humanized or human antibody). Generally,
 25 the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (*e.g.* 6-7 sites) are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage
 30 particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (*e.g.* binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of
 35 the antigen-antibody complex to identify contact points between the antibody and human HER2. Such contact residues and neighboring residues are candidates for substitution according to the techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to

screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

Another type of amino acid variant of the antibody alters the original glycosylation pattern of the antibody. By altering is meant deleting one or more carbohydrate moieties found in the antibody, and/or adding one or more glycosylation sites that are not present in the antibody.

Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylglucosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. For example, antibodies with a mature carbohydrate structure that lacks fucose attached to an Fc region of the antibody are described in US Pat. Appl. No. US 2003/0157108 A1, Presta, L. See also US 2004/0093621 A1 (Kyowa Hakko Kogyo Co., Ltd). Antibodies with a bisecting N-acetylglucosamine (GlcNAc) in the carbohydrate attached to an Fc region of the antibody are referenced in WO03/011878, Jean-Mairet *et al.* and US Patent No. 6,602,684, Umana *et al.* Antibodies with at least one galactose residue in the oligosaccharide attached to an Fc region of the antibody are reported in WO97/30087, Patel *et al.* See, also, WO98/58964 (Raju, S.) and WO99/22764 (Raju, S.) concerning antibodies with altered carbohydrate attached to the Fc region thereof.

It may be desirable to modify the antibody of the invention with respect to effector function, *e.g.* so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antibody. This may be achieved by introducing one or more amino acid substitutions in an Fc region of the antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp Med.* 176:1191-1195 (1992) and Shopes, B. *J. Immunol.* 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using

heterobifunctional cross-linkers as described in Wolff *et al. Cancer Research* 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al. Anti-Cancer Drug Design* 3:219-230 (1989).

5 WO00/42072 (Presta, L.) describes antibodies with improved ADCC function in the presence of human effector cells, where the antibodies comprise amino acid substitutions in the Fc region thereof. Preferably, the antibody with improved ADCC comprises substitutions at positions 298, 333, and/or 334 of the Fc region (Eu numbering of residues). Preferably the altered Fc region is a human IgG1 Fc region comprising or consisting of substitutions at one, two or three of these positions. Such

10 substitutions are optionally combined with substitution(s) which increase C1q binding and/or CDC.

Antibodies with altered C1q binding and/or complement dependent cytotoxicity (CDC) are described in WO99/51642, US Patent No. 6,194,551B1, US Patent No. 6,242,195B1, US Patent No. 6,528,624B1 and US Patent No. 6,538,124 (Idusogie *et al.*). The antibodies comprise an amino acid substitution at one or more of amino acid positions 270, 322, 326, 327, 329, 313, 333 and/or 334 of

15 the Fc region thereof (Eu numbering of residues).

To increase the serum half life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in US Patent 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for

20 increasing the *in vivo* serum half-life of the IgG molecule.

Antibodies with improved binding to the neonatal Fc receptor (FcRn), and increased half-lives, are described in WO00/42072 (Presta, L.) and US2005/0014934A1 (Hinton *et al.*). These antibodies comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. For example, the Fc region may have substitutions at one or more of positions

25 238, 250, 256, 265, 272, 286, 303, 305, 307, 311, 312, 314, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424, 428 or 434 (Eu numbering of residues). The preferred Fc region-comprising antibody variant with improved FcRn binding comprises amino acid substitutions at one, two or three of positions 307, 380 and 434 of the Fc region thereof (Eu numbering of residues).

Engineered antibodies with three or more (preferably four) functional antigen binding sites are

30 also contemplated (US Appln. No. US2002/0004587 A1, Miller *et al.*).

Nucleic acid molecules encoding amino acid sequence variants of the antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette

35 mutagenesis of an earlier prepared variant or a non-variant version of the antibody.

Screening for antibodies with the desired properties

Techniques for generating antibodies have been described above. One may further select

antibodies with certain biological characteristics, as desired.

To identify an antibody which blocks ligand activation of a HER receptor, the ability of the antibody to block HER ligand binding to cells expressing the HER receptor (*e.g.* in conjugation with another HER receptor with which the HER receptor of interest forms a HER hetero-oligomer) may be determined. For example, cells naturally expressing, or transfected to express, HER receptors of the HER hetero-oligomer may be incubated with the antibody and then exposed to labeled HER ligand. The ability of the antibody to block ligand binding to the HER receptor in the HER hetero-oligomer may then be evaluated.

For example, inhibition of HRG binding to MCF7 breast tumor cell lines by HER2 antibodies may be performed using monolayer MCF7 cultures on ice in a 24-well-plate format essentially as described in US Patent No. 6,949,245. HER2 monoclonal antibodies may be added to each well and incubated for 30 minutes. 125I-labeled rHRG β 1177-224 (25 pm) may then be added, and the incubation may be continued for 4 to 16 hours. Dose response curves may be prepared and an IC₅₀ value may be calculated for the antibody of interest. In one embodiment, the antibody which blocks ligand activation of a HER receptor will have an IC₅₀ for inhibiting HRG binding to MCF7 cells in this assay of about 50nM or less, more preferably 10nM or less. Where the antibody is an antibody fragment such as a Fab fragment, the IC₅₀ for inhibiting HRG binding to MCF7 cells in this assay may, for example, be about 100nM or less, more preferably 50nM or less.

Alternatively, or additionally, the ability of an antibody to block HER ligand-stimulated tyrosine phosphorylation of a HER receptor present in a HER hetero-oligomer may be assessed. For example, cells endogenously expressing the HER receptors or transfected to expressed them may be incubated with the antibody and then assayed for HER ligand-dependent tyrosine phosphorylation activity using an anti-phosphotyrosine monoclonal (which is optionally conjugated with a detectable label). The kinase receptor activation assay described in U.S. Patent No. 5,766,863 is also available for determining HER receptor activation and blocking of that activity by an antibody.

In one embodiment, one may screen for an antibody which inhibits HRG stimulation of p180 tyrosine phosphorylation in MCF7 cells essentially as described in US Patent No. 6,949,245. For example, the MCF7 cells may be plated in 24-well plates and monoclonal antibodies to HER2 may be added to each well and incubated for 30 minutes at room temperature; then rHRG β 1₁₇₇₋₂₄₄ may be added to each well to a final concentration of 0.2 nM, and the incubation may be continued for 8 minutes. Media may be aspirated from each well, and reactions may be stopped by the addition of 100 μ l of SDS sample buffer (5% SDS, 25 mM DTT, and 25 mM Tris-HCl, pH 6.8). Each sample (25 μ l) may be electrophoresed on a 4-12% gradient gel (Novex) and then electrophoretically transferred to polyvinylidene difluoride membrane. Antiphosphotyrosine (at 1 μ g/ml) immunoblots may be developed, and the intensity of the predominant reactive band at M_r –180,000 may be quantified by reflectance densitometry. The antibody selected will preferably significantly inhibit HRG stimulation of p180 tyrosine phosphorylation to about 0-35% of control in this assay. A dose-response curve for

inhibition of HRG stimulation of p180 tyrosine phosphorylation as determined by reflectance densitometry may be prepared and an IC_{50} for the antibody of interest may be calculated. In one embodiment, the antibody which blocks ligand activation of a HER receptor will have an IC_{50} for inhibiting HRG stimulation of p180 tyrosine phosphorylation in this assay of about 50nM or less, more preferably 10nM or less. Where the antibody is an antibody fragment such as a Fab fragment, the IC_{50} for inhibiting HRG stimulation of p180 tyrosine phosphorylation in this assay may, for example, be about 100nM or less, more preferably 50nM or less.

One may also assess the growth inhibitory effects of the antibody on MDA-MB-175 cells, e.g., essentially as described in Schaefer *et al. Oncogene* 15:1385-1394 (1997). According to this assay, MDA-MB-175 cells may be treated with a HER2 monoclonal antibody (10 μ g/mL) for 4 days and stained with crystal violet. Incubation with a HER2 antibody may show a growth inhibitory effect on this cell line similar to that displayed by monoclonal antibody 2C4. In a further embodiment, exogenous HRG will not significantly reverse this inhibition. Preferably, the antibody will be able to inhibit cell proliferation of MDA-MB-175 cells to a greater extent than monoclonal antibody 4D5 (and optionally to a greater extent than monoclonal antibody 7F3), both in the presence and absence of exogenous HRG.

In one embodiment, the HER2 antibody of interest may block heregulin dependent association of HER2 with HER3 in both MCF7 and SK-BR-3 cells as determined in a co-immunoprecipitation experiment such as that described in US Patent No. 6,949,245 substantially more effectively than monoclonal antibody 4D5, and preferably substantially more effectively than monoclonal antibody 7F3.

To identify growth inhibitory HER2 antibodies, one may screen for antibodies which inhibit the growth of cancer cells which overexpress HER2. In one embodiment, the growth inhibitory antibody of choice is able to inhibit growth of SK-BR-3 cells in cell culture by about 20-100% and preferably by about 50-100% at an antibody concentration of about 0.5 to 30 μ g/ml. To identify such antibodies, the SK-BR-3 assay described in U.S. Patent No. 5,677,171 can be performed. According to this assay, SK-BR-3 cells are grown in a 1:1 mixture of F12 and DMEM medium supplemented with 10% fetal bovine serum, glutamine and penicillin streptomycin. The SK-BR-3 cells are plated at 20,000 cells in a 35mm cell culture dish (2mls/35mm dish). 0.5 to 30 μ g/ml of the HER2 antibody is added per dish. After six days, the number of cells, compared to untreated cells are counted using an electronic COULTER™ cell counter. Those antibodies which inhibit growth of the SK-BR-3 cells by about 20-100% or about 50-100% may be selected as growth inhibitory antibodies. See US Pat No. 5,677,171 for assays for screening for growth inhibitory antibodies, such as 4D5 and 3E8.

In order to select for antibodies which induce apoptosis, an annexin binding assay using BT474 cells is available. The BT474 cells are cultured and seeded in dishes as discussed in the preceding paragraph. The medium is then removed and replaced with fresh medium alone or medium containing 10 μ g/ml of the monoclonal antibody. Following a three day incubation period, monolayers

are washed with PBS and detached by trypsinization. Cells are then centrifuged, resuspended in Ca^{2+} binding buffer and aliquoted into tubes as discussed above for the cell death assay. Tubes then receive labeled annexin (*e.g.* annexin V-FTIC) (1 $\mu\text{g}/\text{ml}$). Samples may be analyzed using a FACSCAN™ flow cytometer and FACSCONVERT™ CellQuest software (Becton Dickinson). Those antibodies

5 which induce statistically significant levels of annexin binding relative to control are selected as apoptosis-inducing antibodies. In addition to the annexin binding assay, a DNA staining assay using BT474 cells is available. In order to perform this assay, BT474 cells which have been treated with the antibody of interest as described in the preceding two paragraphs are incubated with 9 $\mu\text{g}/\text{ml}$ HOECHST 33342™ for 2 hr at 37°C, then analyzed on an EPICS ELITE™ flow cytometer (Coulter

10 Corporation) using MODFIT LT™ software (Verity Software House). Antibodies which induce a change in the percentage of apoptotic cells which is 2 fold or greater (and preferably 3 fold or greater) than untreated cells (up to 100% apoptotic cells) may be selected as pro-apoptotic antibodies using this assay. See WO98/17797 for assays for screening for antibodies which induce apoptosis, such as 7C2 and 7F3.

15 To screen for antibodies which bind to an epitope on HER2 bound by an antibody of interest, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed to assess whether the antibody cross-blocks binding of an antibody, such as 2C4 or pertuzumab, to HER2. Alternatively, or additionally, epitope mapping can be performed by methods known in the art and/or one can study the

20 antibody-HER2 structure (Franklin *et al.* *Cancer Cell* 5:317-328 (2004)) to see what domain(s) of HER2 is/are bound by the antibody.

Methods of cancer treatment

The patients identified in accordance with the present invention as likely to be resistant to treatment with HER2 inhibitors, are likely to benefit from combination treatments.

25 Combination treatments may include chemotherapy in conjunction with use of a HER2 inhibitor, such as a HER2 antibody, *e.g.* trastuzumab or pertuzumab.

The purpose of chemotherapeutic treatment of cancer is to cure the patient or, at least, slow down disease progression, increase survival, reduce the likelihood of cancer recurrence, control symptoms and/or maintain or improve quality of life. Chemotherapy varies depending on the type of

30 cancer, and, in case of solid tumors, can be performed before and/or after surgical removal of primary tumor. For some cancers, there are a few universally accepted standard therapies, while the treatment of others is not yet standardized.

Exemplary chemotherapeutic agents have been listed before, and generally can be classified according to their mechanism of action. Some chemotherapeutic agents directly damage DNA and

35 RNA. By disrupting replication of the DNA such chemotherapeutics either completely halt replication, or result in the production of nonsense DNA or RNA. This category includes, for example, cisplatin (Platinol®), daunorubicin (Cerubidine®), doxorubicin (Adriamycin®), and etoposide

(VePesid®). Another group of cancer chemotherapeutic agents interfere with the formation of nucleotides or deoxyribonucleotides, so that RNA synthesis and cell replication is blocked. Examples of drugs in this class include methotrexate (Abitrexate®), mercaptopurine (Purinethol®), fluorouracil (Adrucil®), and hydroxyurea (Hydrea®). A third class of chemotherapeutic agents effects the

5 synthesis or breakdown of mitotic spindles, and, as a result, interrupt cell division. Examples of drugs in this class include vinblastine (Velban®), vincristine (Oncovin®) and taxenes, such as, paclitaxel (Taxol®), and docetaxel (Taxotere®). Other classifications, for example, based on the chemical structure of the chemotherapeutic agents, are also possible.

For breast cancer, doxorubicin (Adriamycin®) is considered by most the most effective single

10 chemotherapeutic agent. In addition, 5-FU has been in clinical use for several decades, and is the cornerstone of many combination therapies for breast cancer. Other chemotherapeutic agents commonly used for the treatment of breast cancer include, for example, anthracyclines, taxane derivatives, and various combinations therapies, such as CMF (cyclophosphamide-methotrexate-fluorouracil) chemotherapy. Most patients receive chemotherapy immediately following surgical

15 removal of tumor. This approach is commonly referred to as adjuvant therapy. However, chemotherapy can be administered also before surgery, as so called neoadjuvant treatment. Although the use of neo-adjuvant chemotherapy originates from the treatment of advanced and inoperable breast cancer, it has gained acceptance in the treatment of other types of cancers as well. The efficacy of neoadjuvant chemotherapy has been tested in several clinical trials. In the multi-center National

20 Surgical Adjuvant Breast and Bowel Project B-18 (NSAB B-18) trial (Fisher et al., J. Clin. Oncology 15:2002-2004 (1997); Fisher et al., J. Clin. Oncology 16:2672-2685 (1998)) neoadjuvant therapy was performed with a combination of adriamycin and cyclophosphamide ("AC regimen"). In another clinical trial, neoadjuvant therapy was administered using a combination of 5-fluorouracil (5-FU), epirubicin and cyclophosphamide ("FEC regimen") (van Der Hage et al., J. Clin. Oncol. 19:4224-

25 4237 (2001)). Other clinical trials have also used taxane-containing neoadjuvant treatment regimens. See, e.g. Holmes et al., J. Natl. Cancer Inst. 83:1797-1805 (1991) and Moliterni et al., Seminars in Oncology, 24:S17-10-S-17-14 (1999). For further information about neoadjuvant chemotherapy for breast cancer see, Cleator et al., Endocrine-Related Cancer 9:183-195 (2002).

5-FU, CPT-11 (irinotecan), and oxaliplatin, administered alone or in combination, have

30 proven effective in the treatment of advanced colorectal cancer (CRC) (see, e.g. Grothey et al. (2004) J. Clin. Oncol. 22:1209-15).

Non-small-cell lung cancer (NSCLC) has been shown to respond well to combination therapy with vinorelbine, cisplatin and optionally paclitaxel (see, e.g. Rodriguez et al. (2004) Am. J. Clin. Oncol. 27:299-303).

35 Chemotherapeutic regimens for the treatment of other types of cancer are also well known to those skilled in the art.

Further details of the invention will be described in the following non-limiting Example

ExampleIdentifying markers of trastuzumab resistance in HER2+ breast cancer

HER2 is overexpressed by gene amplification in about 20% of breast cancers. It is known that
 5 HER2 gene amplification leads to significantly higher level of HER2 receptor expression compared to
 normal cells: e.g., IHC3+ = 1×10^6 receptors/cell (normal cells = 2×10^4). It is also known that HER2
 amplification is associated with higher tumor grade, lymph node positivity and poor prognosis, and
 HER2 status in metastases is highly correlated with HER2 status in the primary tumor.

While trastuzumab has been highly successful in the treatment of HER2-positive tumors, such
 10 as HER2-positive breast cancer, certain tumors are non-responsive, or show or develop resistance to
 trastuzumab treatment.

Using a cell line that is known to be sensitive to trastuzumab *in vitro* (BT474), an siRNA
 screen was performed to identify genes that when knocked down (or inactivated) lead to induction of
 trastuzumab resistance *in vitro*. Validated hits from the screen are candidate diagnostic markers of
 15 trastuzumab resistance *in vivo*.

Methods:

Cell line and assay: The BT474 cell line was used, which is HER2-positive and sensitive to
 trastuzumab *in vitro*. Based on information in the literature, PTEN and p27 were used as positive
 controls to develop an assay for screening. Knockdown of both PTEN and p27 has been reported to
 20 reduce the ability of trastuzumab to slow cell proliferation *in vitro*. This effect has been observed in
 the present study as well and used these positive controls to optimize the assay. The most effective
 method of determining trastuzumab response *in vitro* was found to be measuring cell proliferation via
 a ^3H -thymidine uptake assay. Briefly, the siRNA and lipofectamine were distributed onto 96-well
 plates. Cells were then plated onto the aliquoted siRNA and at 24 hours, trastuzumab was added at a
 25 concentration of 25 $\mu\text{g}/\text{ml}$. At 72 hours, ^3H -thymidine was added to the cultures. The amount of
 incorporated ^3H was measured using a 96-well plate cell harvested on day 4 (outline on Figure 1).

siRNA screen: The screen was optimized for automated screening using a 96-well plate
 format using either luciferase or non-targeting controls as the negative control and PTEN and p27 as
 the positive control (Figures 2 and 3). The finalized screen format is depicted in Figure 4. Using this
 30 method, the Dharmacon kinase and phosphatase libraries, which covered 979 genes (779 genes from a
 kinase library and 200 genes from a phosphatase library), were screened and analyzed with 4
 individually screened siRNA's against each gene.

Data analysis: Data were analyzed in several ways. One method was to normalize data to
 various controls including the negative controls, the positive controls or to the plate average (Figure
 35 5). A gene was considered a hit if at least 2 of the 4 siRNA oligos were above a z-score threshold of
 1.5. The data were also analyzed manually by plotting the data and identifying spots that were greater

than 1.5 standard deviations above the mean for the non-targeting control, again with a minimum of 2 of the 4 siRNA oligos scoring positive to be considered a hit (Figure 6).

Results:

Primary screen data: From the analysis of the screen data, there were 25 genes that were identified as hits from the kinase library by all the data analysis methods that were used (Figure 7). An additional 5 genes that were identified manually were found to be very close to the threshold in the biostatistics analysis and were included in further follow-up. Both of the positive controls, PTEN and p27, were on the plates initially screened (kinase library and one plate from phosphatase library) and were identified as hits, suggesting that the screen performed well to detect the type of hits of interest.

The hits fell into several categories of potential interest including cell cycle regulators, major players in downstream-receptor tyrosine kinase signaling, and several other categories (Figure 8).

Hit validation: For further validation, we focused on 28 genes from this initial screen. This includes the 30 noted above minus the two positive controls PTEN and p27 which have already been validated in other studies. Two methods were used for validation. First, the siRNA's were re-screened in BT474 cells to determine whether the observation would repeat in the same system. The genes were then also screened in a different cell line (SKBR3) which is also HER2-positive and trastuzumab sensitive. Examples of how the positive controls PTEN and p27 performed in the validation screens is illustrated in Figure 9. The 28 hits (other than PTEN and p27) from the primary screen are listed in Figure 10 along with the results from the repeat screen and the screen in SKBR3 cells. Some of the most promising candidates considering the performance in validation screens are shaded.

In a smaller subsequent screen, the remaining phosphatase library plates were screened (other than the one plate containing PTEN which was screened with the kinase library). The results from the analysis of all phosphatase plates are shown in Figure 11. PTEN was identified as a 3 oligo hit by two methods. There were an additional 3 genes that were at least 2 oligo hits by all normalization methods and are shaded in the list of hits on Figure 11.

Another method of validation was to examine GeneLogic data to determine if any of the genes exhibited evidence of decreased expression in HER2-positive breast cancer compared to normal breast tissue or to HER2-negative breast cancer. Four genes did exhibit such a pattern – SOCS5, LATS2, PTPN11, and DYRK1A and are thus worth further follow-up even if the validation screen data is not as strong (e.g. LATS2) (Figure 12).

The top hits based on the strongest phenotype and >2 oligo hits (PPM1H, DYRK1A, STK10, and PTPN11) are shown in Figure 13.

Figures 14 and 15 show examples of the top hits augment cell proliferation in BT474 cells and BT474-M1 cells treated with trastuzumab.

Figure 16 shows that results of 3H-thymidine uptake and cell titre glow assays, and demonstrate that increased proliferation at 3 days (a) is associated with increased cell number at 7 days (b).

Figure 17 shows that knockdown of the candidate genes also attenuates lapatinib response in multiple cell lines (PPM1H in particular).

Figure 18 shows that PPM1H and PTPN11 negatively regulate the HER3/PI3K signaling axis.

The data set forth in Figure 19 show that knockdown of all four candidate genes (PPM1H, PTPN11, DYK1A and STK10) may increase Akt phosphorylation.

Based on these experimental data, PPM1H appears to be a particularly useful and reliable indicator of trastuzumab resistance. This molecule belongs to the protein phosphatase 2C family, and is known to play a role in other cell types, such as neurite outgrowth and putative oncogenic role in

colon adenocarcinoma. Other family members identified herein have been linked to diverse pathways, e.g., PP2C α and β binds CDK2/CDK6; ILKAP is linked to integrin/GSK-beta signaling; PHLPP I a pAkt phosphatase, and mouse PP2C γ ?FIN13 has been shown to negatively regulate growth.

Figure 20 is a cladogram showing PPM1 family members.

Figure 21 shows that PPM1M and PPM1J also attenuate trastuzumab response in vitro albeit weaker than PPM1H.

The results shown in Figure 22 show that PPM1M knockdown also decreased Lapatinib response in vitro.

From these data it appears that PPM1H has similar functions to other PP2C family members. In particular, without being bound by any theory, it is believed that PPM1H may function as pAkt phosphatase similar to another family member, PHLPP to dephosphorylate P-Akt1 (S473).

However, PPM1H is also different, and may have a novel function, distinct from other PP2C family members. This new function is the modulation of signaling upstream of HER3. Thus, PP2C may function like another PP2C family member, ILKAP, to indirectly regulate GSK3/cyclin D1 signaling and thereby modulate the PI3K/Akt signaling axis downstream of pAkt.

It has been found that several genes (PTEN, CDKN1B, PPM1H, PTPN11, PPM1A, PPM1J) genes identified by the screens described in the present invention exhibit decreased expression in basal-like cell lines and tumors (see Table 2). This is of great significance, since basal-like expression has been negatively associated with poor outcome in HER2-negative patients.

Although in the foregoing description the invention is illustrated with reference to certain embodiments, it is not so limited. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

SEQUENCE LISTING IN ELECTRONIC FORM

This description contains a sequence listing in electronic form in ASCII text format (file no. 84261-169_ca_seqlist_v1_2009Nov27.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following Table.

SEQUENCE TABLE

<110> GENENTECH, INC.

<120> GENE EXPRESSION MARKERS OF TUMOR RESISTANCE TO HER2
INHIBITOR TREATMENT

<130> 84261-169

<140> PCT/US2008/065766

<141> 2008-06-04

<150> 60/942,906

<151> 2007-06-08

<160> 6

<170> PatentIn Ver. 3.5

<210> 1

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<220>

<221> VARIANT

<222> (10)

<223> /replace="Ser"

<220>

<221> misc_feature

<222> (10)

<223> /note="Residue given in the sequence have no preference
with respect to those in the annotation for said position"

<400> 1

Gly Phe Thr Phe Thr Asp Tyr Thr Met Asp
1 5 10

<210> 2
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 2
 Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr Asn Gln Arg Phe Lys
 1 5 10 15

Gly

<210> 3
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 3
 Asn Leu Gly Pro Ser Phe Tyr Phe Asp Tyr
 1 5 10

<210> 4
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 4
 Lys Ala Ser Gln Asp Val Ser Ile Gly Val Ala
 1 5 10

<210> 5
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic peptide

<220>
 <221> VARIANT
 <222> (5)
 <223> /replace="Leu"

<220>
<221> VARIANT
<222> (6)
<223> /replace="Glu"

<220>
<221> VARIANT
<222> (7)
<223> /replace="Ser"

<220>
<221> misc_feature
<222> (5)..(7)
<223> /note="Residues given in the sequence have no preference
with respect to those in the annotations for said positions"

<400> 5
Ser Ala Ser Tyr Arg Tyr Thr
1 5

<210> 6
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 6
Gln Gln Tyr Tyr Ile Tyr Pro Tyr Thr
1 5

CLAIMS:


1. A method of predicting the likelihood of response of a human breast cancer patient diagnosed with a HER2 expressing tumor of the breast to treatment with a HER2 inhibitor, comprising:

determining, in a breast tumor sample obtained from said subject, the expression level of RNA transcripts or their expression products of the PPM1H gene,

wherein a lower level of expression relative to the expression level of the PPM1H gene in a normal cell of the same cell type indicates that the subject is likely to be resistant to treatment with the HER2 inhibitor.
2. The method of claim 1, wherein the diagnosis includes quantification of HER2 expression level.
3. The method of claim 2, wherein the HER2 expression level is quantified by immunohistochemistry (IHC) and/or fluorescence in situ hybridization (FISH).
4. The method of claim 2, wherein the HER2 expression level is quantified by immunohistochemistry (IHC) and/or fluorescence in situ hybridization (FISH) and wherein the cancer expresses HER2 at least at a 1 + level.
5. The method of claim 4, wherein the cancer expresses HER2 at least at a 2+ level.
6. The method of claim 4, wherein the cancer expresses HER2 at least at a 3+ level.
7. The method of any one of claims 1 to 6, wherein the breast cancer is metastatic breast cancer.
8. The method of any one of claims 1 to 7, wherein the level of expression in the tumor sample of one or more of PTPN11, DYRK1A and STK10 genes is additionally determined.
9. The method of claim 8, wherein the expression level of the RNA transcript or the expression product of one or more of said PTPN11, DYRK1A or STK10 genes is determined.

10. The method of any one of claims 1 to 9, wherein the HER2 inhibitor is an agent which interferes with HER2 activation or function.
11. The method of any one of claims 1 to 9, wherein the HER2 inhibitor is a HER antibody or antibody fragment, a small molecule HER2 antagonist, a HER2 tyrosine kinases inhibitor, or an antisense molecule.
12. The method of claim 11, wherein the HER2 inhibitor is a HER2 antibody or antibody fragment, or a small molecule which binds to and inhibits the HER2 receptor.
13. The method of claim 12, wherein the HER2 antibody inhibits HER2 ectodomain cleavage, blocks ligand activation of a HER receptor, inhibits HER2 dimerization, binds to the heterodimeric binding site of HER2, or binds to the 405 epitope.
14. The method of claim 12, wherein the HER2 antibody or antibody fragment binds to the 405 epitope and is one or more of trastuzumab, humanized antibodies huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6 and huMAb4D5-7, and fragments thereof.
15. The method of claim 12, wherein the HER2 antibody or antibody fragment blocks ligand activation of a HER2 receptor more effectively than trastuzumab.
16. The method of claim 12, wherein the HER2 antibody or antibody fragment inhibits HER2 dimerization and binds the 2C4 epitope.
17. The method of claim 16, wherein the HER2 antibody or antibody fragment is pertuzumab or a fragment thereof.
18. The method of any one of claims 1 to 17, wherein the tumor sample is from a fixed, wax-embedded cancer tissue specimen of said patient or is a core biopsy tissue.

Measurement of Trastuzumab Response of HER2 Amplified Cell Line BT474 by 3H-Thymidine Incorporation Assay



Maximum dynamic range (highest sensitivity)




FIG. 1

2/28




FIG. 2

Optimization of the Screening Window Coefficient -Z factor




FIG. 3B





FIG. 3C



Summary of the Phosphatase Library Screen

normalization	Li analysis z>1.5			Raw values analysis	
	ave	pos	neg	Nil	
PTEN	3	1	1	3	
PTPN11	2	2	2	3	
PPP2R1A	1*	2	3	3	
KIAA0685	2	2	2	2	
PPM1H	2	2	2	2	
SSH3	1	2	2	2	
MTMR3	1	2	2	1	
PPP2R2B	0	2	1	1	
PTENP1	2	1	1	2	
KIF1B	2	0	0	2	
MPHOSPH1	2	0	0	2	

2 or 3 oligo hits (disregard of mix results) = 11 (5.5% of the library)

2 oligo hits (in all analysis) = 3 (1.5% of the library)

* Average of plate is obscured due to one high data point

FIG. 4A





FIG. 4B

≥ 3 oligo hits or 2 oligos + pool



6/28

Data analysis by plotting raw values of the screen showed P27 is a 4 oligo hit




FIG. 6A




FIG. 6B

7/28





FIG. 6C

Combine Analysis of Kinase Library Hits



* 2 oligo hits in some but not all normalized analysis

FIG. 7

8/28

Result from the Kinase Library Screen

Known/putative functions	examples	hits
Cell cycle regulation	CDKs, RhoGTPases/ cytoskeleton	6 (2 oligo hits) 1 (3 oligo hits)
Key downstream Signaling pathways	MAPKs, PI 3/4 kinase	7 (2 oligo hits) 1 (3 oligo hits)
Less Characterized pathways	Metabolic pathways, PKC, KAK-STAT, Wnt	5 (2 oligo hits)
Others	Novel genes, unrelated pathways	7 (2 oligo hits) 1 (3 oligo hits)

FIG. 8

9/28

Development of the Secondary Screen

FIG. 9A




FIG. 9B






FIG. 9C

*P27=oligo2*

10/28



Automation

FIG. 9D

Combined analysis of the screens

Gene	1st screen # oligo hits	(oligos)	Validation screen # oligo hits	(oligos)	SKBR3 # oligo hits	(oligos)
CDK11	2	(2,3)	2	(2,3)	1	(2)
DKFZP586B16	2	(2,3)	2+P	(2,3)	0	0
DGKI	2	(2,3)	2+P	(2,3)	1	(2)
DUSP4	2	(1,4)	2	(1,4)	1+P	(1)
DUSP6	2	(2,4)	2	(2,4)	1	(1)
DYRK1A	2	(1,4)	4+P	(1,2,3,4)	4+P	(1,2,3,4)
FLJ35107	2	(1,2)	2	(1,2)	1	(2)
FLT1	3	(1,2,3)	4+P	(1,2,3,4)	2	(2,4)
HIPK3	2	(3,4)	1	(3)	0	0
HK2	2	(1,4)	3	(1,2,4)	3	(1,2,4)
ITK	2	(1,2)	2	(1,2)	1	(2)
MOAP1/ITPK1	2	(3,4)	1	(4)	1	(3)
KIAA1639	3 (4)	(1,2,3, 4?)	3+P	(1,3,4)	2	(1,3)
LATS2	2(3)	(1,3, 4?)	3+P	(1,3,4)	0	0
LIM/PDLIM5	2	(1,2)	2+P	(1,2)	0	0
MAPK8/JNK	2	(1,3)	1	(1)	0	0
MAP4K4	2	(1,2)	2	(1,2)	2	(1,2)
PANK1	2	(1,3)	3	(2,3,4)	1+P	(1)
PI4K2B	2 (3)	(1,2 4?)	2	(2,4)	1	(2)
PRKWNK3	2	(2,4)	2+P	(2,4)	0	0
RYK	2	(1,4)	2	(1,4)	0	0
SPEC2	2	(3,4)	2	(3,4)	2+P	(2,4)
Socs5	2	(1,3)	2	(1,3)	4+P	(1,2,3 4)
STK10	2	(1,2)	4+P	(1,2,3 4)	3	(2,3 4)
STK22C	2	(1,4)	3+P	(1,3,4)	2+P	(3,4)
STYK1/NOV*	2	(1,2)	3+P	(1,2,4)	0	0
TXND3	2	(1,3)	2	(1,3)	2	(1,2)
Wee1	2	(1,2)	2	(1,2)	0	0

FIG. 10

12/28

Summary of the Phosphatase Library Screen

normalization	Li analysis z>1.5			Raw values analysis
	ave	pos	neg	Nil
PTEN	3	1	1	3
PTPN11	2	2	2	3
PPP2R1A	1*	2	3	3
KIAA0685	2	2	2	2
PPM1H	2	2	2	2
SSH3	1	2	2	2
MTMR3	1	2	2	1
PPP2R2B	0	2	1	1
PTENP1	2	1	1	2
KIF1B	2	0	0	2
MPHOSPH1	2	0	0	2

2 or 3 oligo hits (disregard of mix results) = 11 (5.5% of the library)

2 oligo hits (in all analysis) = 3 (1.5% of the library)

*Average of plate is obscured due to one high data point

FIG. 11

13/28

GeneLogic Expression Data

FIG. 12A




FIG. 12B




FIG. 12C





FIG. 12D



14/28

Top Hits Based on Strongest Phenotype and >2 Oligo Hit

1	PPM1H	3+P strong	UNQ11299
2	DYRK1A	4+P strong	UNQ3880
3	STK10	4+P strong	UNQ4026
4	PTPN11*	2+P	UNQ7397

*PTPN11 was included despite being only a 2 oligo hit because of a known link to HER2-PI3K/Akt signaling

FIG. 13

15/28

Examples of how top hits augment cell proliferation
in BT474 cells treated with Herceptin




FIG. 14A




FIG. 14B

16/28

Examples of how top hits augment cell proliferation in
BT474-M1 cells treated with Herceptin




FIG. 15A




FIG. 15B

17/28

Increased proliferation at 3 days (a) is associated with increased cell number at 7 days (b)




FIG. 16A

18/28




FIG. 16B

19/28

Knockdown of candidate genes also attenuates
lapatinib response in mutiple cell lines
(PPM1H in particular)




FIG. 17A





FIG. 17B

20/28



Summary	Cell lines		
Genes	M1	HCC1419	SKBR3
PTEN	+	-	+/-
P27	+	+	-
PPM1H	+	+	+
PTPN11	+	-	N/D
STK10	-/+	+	N/D
DYRK1A	-	-/+	N/D

+ significant decreased of Lapatinib response

+/- moderate decreased of Lapatinib response

FIG. 17D

PPM1H and PTPN11 negatively regulates HER3/PI3K signaling axis

Knockdown augments pHER3





FIG. 18

22/28

Knockdown of all 4 candidate genes may increase Akt phosphorylation




FIG. 19

PPM1 Family Members




FIG. 20

24/28

PPM1M and PPM1J also attenuate Herceptin response in vitro,
albeit weaker than PPM1H




FIG. 21

25/28

PPM1M knockdown also decreased Lapatinib response in vitro




FIG. 22

26/28

TABLE 1

gene	# oligos worked		SKBR3	ccession number
	Primary	Validation		
CDK11	2	2		NM_015076
DYRK1A	2	4+P	*	NM_001396
LATS2	2-3	3+P		NM_014572
STK10	2	4+P	*	NM_005990
Wee1	2	2		NM_003390
DUSP4	2	2		NM_001394
DUSP6	2	2		NM_001946
HIPK3	2	1		NM_005734
JNK	2	1		NM_002752
MAP4K4	2	2		NM_004834
PTPN11	2-3	3+P	*	NM_002834
Socs5	2	2	**	NM_014011
PPM1H	2-3	3+P	*	NM_051093
DKFZP586B16	2	2+P		NM_015533
DGKI	2	2+P		NM_004717
FLJ35107	2	2		NM_496631
FLT1	3	4+P		NM_002019
HK2	2	3	*	NM_000189
ITK	2	2		NM_005546
MOAP1	2	1		NM_014216
KIAA0685	2	1+P		NM_014678
KIAA1639	3-4	3+P	*	NM_290923
LIM/PDLIM5	2	2+P		NM_006457
PANK1	2	3		NM_138316
PI4K2B	2-3	2		NM_018323
PPP2R1A	2-3	2+P		NM_014225
PRKWNK3	2	2+P		NM_020922
RYK	2	2		NM_002958
SPEC2	2	2		NM_020240
STK22C	2	3+P	*	NM_052841
STYK1	2	3+P		NM_018423
TXND3	2	2	*	NM_016616

* 2 or more oligo hits in SKBR3 cell line functions

Blue: cell cycle

Green: Erk/MARK pathway

Red: Post-translation modification

27/28

TABLE 2

Summary

Gene	Cell lines	Tissue
	Lower expression in Basal	Lower expression in Basal
PTEN	Y	Y
CDKN1B	Y	Y
PPM1H	Y	Y
PTPN11	Y	Y
DYRK1A	N	N
STK10	N	N
PPM1A	Y	Y
PPM1B	N	N
ILKAP	N	N
PPM1M	N	N
PPM1J	Y	Y

28/28
TABLE 3

	Refseq
CDKN1B	NM_004064
DYRK1A	NM_001396
TNNI3K	NM_015978
DUSP4	NM_001394
ATR	NM_001184
BCR	NM_004327
BMPR1A	NM_004329
CDK11	NM_015076
DKFZP586B1	NM_015533
CDC2L2	NM_024011
DUSP6	NM_001946
STYK1	NM_018423
DGKI	NM_004717
KIAA1639	NM_290923
FLJ35107	NM_496631
LIM	NM_006457
MARK8	NM_002750
FLT1	NM_002019
ITPK1	NM_014216
HK2	NM_000189
ITK	NM_005546
MAP4K4	NM_004834
LATS2	NM_014572
SPEC2	NM_020240
EXOSC10	NM_002685
PRKWNK3	NM_020922
PI4K2B	NM_018323
STK22C	NM_052841
STK10	NM_005990
PANK1	NM_138316
SOCS5	NM_014011
RYK	NM_002958
PDK3	NM_005391
PRKD2	NM_016457
TXNDC3	NM_016616
WEE1	NM_003390
PPM1H	NM_051093
PPP2R1A	NM_014225
PPP2R2B	NM_004576
MTMR3	NM_021090
PTPN11	NM_002834
SSH3	NM_017857