



(86) Date de dépôt PCT/PCT Filing Date: 1995/05/15
(87) Date publication PCT/PCT Publication Date: 1996/01/18
(45) Date de délivrance/Issue Date: 2006/07/18
(85) Entrée phase nationale/National Entry: 1996/12/30
(86) N° demande PCT/PCT Application No.: IT 1995/000073
(87) N° publication PCT/PCT Publication No.: 1996/001321
(30) Priorité/Priority: 1994/07/01 (ITTO94A000543)

(51) Cl.Int./Int.Cl. *C12N 15/62* (2006.01),
C07K 19/00 (2006.01), *C07K 14/045* (2006.01),
C12N 15/38 (2006.01), *G01N 33/569* (2006.01)

(72) Inventeurs/Inventors:
MAINE, GREGORY T., US;
FLANDERS, RICHARD T., US;
LANDINI, MARIA PAOLA, IT;
RIPALTI, ALESSANDRO, IT

(73) Propriétaires/Owners:
LANDINI, MARIA PAOLA, IT;
RIPALTI, ALESSANDRO, IT;
ABBOTT LABORATORIES, US

(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : MONO- ET POLY-ANTIGENES RECOMBINES POUR LA DETECTION DE L'IgM SPECIFIQUE DU
CYTOMEGALOVIRUS DANS LES SERUMS HUMAINS PAR DOSAGE IMMUNO-ENZYMATIQUE

(54) Title: RECOMBINANT MONO AND POLY ANTIGENS TO DETECT CYTOMEGALOVIRUS-SPECIFIC IgM IN
HUMAN SERA BY ENZYME IMMUNOASSAY

(57) **Abrégé/Abstract:**

A mixture of recombinant mono- and poly-epitope proteic materials able to fully replace the viral antigens when used in an enzyme immunoassay (EIA) is disclosed; the mixture includes a poly-epitope fusion protein having a first region formed by an amino acid sequence (H10) corresponding to that of the last 233 amino acids of the COOH terminus of the vital protein p52 or to a part thereof, a second region formed by an amino acid sequence (F3) corresponding to that of the last 43 amino acids of the COOH terminus of viral protein pp150 or to a part thereof, and a third region formed by an amino acid sequence (AIC2) corresponding to that taken from aa 595 to aa 614, proceeding in direction 5'→3', of the same viral protein pp150; and, in combination, a second fusion protein including a sequence of amino acids corresponding to that taken, proceeding in direction 5'→3' from aa 297 to aa 510 of the viral major matrix protein pp65 encoded by the viral gene UL83 and a third fusion protein including a sequence of amino acids corresponding to that taken, proceeding in direction 5'→3', from aa 117 to aa 373 of the vital assembly protein pp38 encoded by the viral gene UL80a. These three fusion proteins may be used combined together for the preparation of an ELISA test kit for detection of Cytomegalovirus-specific IgM in human sera.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/38, 15/62, G01N 33/569, C07K 16/08	A1	(11) International Publication Number: WO 96/01321 (43) International Publication Date: 18 January 1996 (18.01.96)
(21) International Application Number: PCT/IT95/00073 (22) International Filing Date: 15 May 1995 (15.05.95) (30) Priority Data: TO94A000543 1 July 1994 (01.07.94) IT (71) Applicant (for all designated States except US): ABBOTT LABORATORIES [US/US]; 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US). (71)(72) Applicants and Inventors: LANDINI, Maria, Paola [IT/IT]; Via S. Apollonia, 2, I-40127 Bologna (IT). RIPALTI, Alessandro [IT/IT]; Via Petrarca, 25, I-40127 Bologna (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): MAINE, Gregory, T. [US/US]; 5743 Constitution Avenue, Gurnee, IL 60031 (US). FLANDERS, Richard, T. [US/US]; 358 Woodland Drive, Grayslake, IL 60030 (US). (74) Agents: BONGIOVANNI, Guido et al.; Studio Torta, Via Viotti, 9, I-10121 Torino (IT).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). 2194135 Published With international search report.
(54) Title: RECOMBINANT MONO AND POLY ANTIGENS TO DETECT CYTOMEGALOVIRUS-SPECIFIC IgM IN HUMAN SERA BY ENZYME IMMUNOASSAY (57) Abstract A mixture of recombinant mono- and poly-epitope proteic materials able to fully replace the viral antigens when used in an enzyme immunoassay (EIA) is disclosed; the mixture includes a poly-epitope fusion protein having a first region formed by an amino acid sequence (H10) corresponding to that of the last 233 amino acids of the COOH terminus of the viral protein p52 or to a part thereof, a second region formed by an amino acid sequence (F3) corresponding to that of the last 43 amino acids of the COOH terminus of viral protein pp150 or to a part thereof, and a third region formed by an amino acid sequence (A1C2) corresponding to that taken from aa 595 to aa 614, proceeding in direction 5'→3', of the same viral protein pp150; and, in combination, a second fusion protein including a sequence of amino acids corresponding to that taken, proceeding in direction 5'→3' from aa 297 to aa 510 of the viral major matrix protein pp65 encoded by the viral gene UL83 and a third fusion protein including a sequence of amino acids corresponding to that taken, proceeding in direction 5'→3', from aa 117 to aa 373 of the viral assembly protein pp38 encoded by the viral gene UL80a. These three fusion proteins may be used combined together for the preparation of an ELISA test kit for detection of Cytomegalovirus-specific IgM in human sera.		

5

Recombinant mono and poly antigens to detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay.

*** **

10 Technical Field:

The present invention relates to recombinant proteic materials useful as mono- and poly-epitope artificial antigens able to detect HCMV-specific immunoglobulines in Human sera, especially IgM, in place of purified virions. In particular, the invention relates to the joint
15 use of three different fusion proteins obtained by recombinant methods, a first one including high immunogenic epitopes both from the non structural, DNA-binding viral protein of 52 kD (pp52) and from the structural phosphoprotein of 150 kD (pp150), a second one including an immunogenic epitope from the major matrix protein of the Human
20 Cytomegalovirus (HCMV), namely the viral protein of 65 kD (pp65) encoded by the viral gene UL83, and a third one including an immunogenic epitope from the assembly protein of 38 kD (pp38) encoded by the viral gene UL80a. The invention further relates to a

diagnostic kit and to diagnostic reagents derived therefrom and to plasmids expressing said recombinant antigens.

Background Art:

Human Cytomegalovirus (HCMV) is a ubiquitous herpesvirus in man. It is rarely pathogenic in healthy adults but is associated with several diseases in immunocompromised individuals (such as HIV-infected people and transplant recipients). Furthermore, HCMV is the most common cause of congenital infection in humans. Intrauterine primary infections are second only to Down's syndrome as a known cause of mental retardation. Less severe complications are the result of secondary infections. As infections are either asymptomatic or accompanied by symptoms that are not specific of HCMV (such as fever and leukopenia), laboratory techniques are the sole means of diagnosing acute HCMV infection. Diagnosis of HCMV infection can be obtained by direct demonstration of the virus or virus components in pathological materials or indirectly through serology [1]. Diagnosis of primary HCMV infection is exclusively accomplished by serological methods, i.e. demonstration of the appearance of antibodies to HCMV in a previously seronegative subject. HCMV-specific IgM is a sensitive and specific indicator of primary HCMV infection in immunocompetent subjects while it is very often produced during active viral reactivation in transplant recipients [2-4]. However its detection varies widely and a very poor agreement has been found among the results obtained with

different commercial kits [5].

From EP 262531 there are known immunogenic portions of HCMV structural phosphoprotein of 150 kD, encoded by the gene localized in the Hind III-Y/N fragment of the viral genome; according to said European patent, such immunogenic portions of pp150 are encoded, in particular, by an EcoRI-PstI fragment of approximately 1.5 kb, localized inside the region of EcoRI-Y fragment from HCMV genome of AD169 strain. Subsequent and more exhaustive studies have however shown the afore deductions to be incorrect, in that (Fig. 1), protein pp150 is shown to be encoded by UL32, much longer than 1.5 kb and, anyway, quite outside any such EcoRI-PstI fragment, as may be defined within the EcoRI-Y fragment of the AD169 strain. Furthermore, the EcoRI-Y fragment is wholly outside (in particular, to the side, towards the NH₂-terminus) of Hind III-Y/N fragment. According to the Applicants, therefore, there is a correlation between the immunogenic properties shown by the proteic material referred to in the afore European patent (which, however, providing an incorrect identification of the polypeptide, causes it to remain substantially undetermined) and the inclusion into the peptide of epitope A1C2, encoded by UL32 nucleotides 1783 to 1842, running 5'→3', i.e., of the region corresponding to aminoacids 595 to 614 of ppUL32, whose identification is posterior (Novak J.P. et al. 1991. Mapping of serologically relevant regions of human cytomegalovirus

phosphoprotein pp150 using synthetic peptides. J. Gen. Virol. 72; 1409-1413).

Antigenic materials composed of single well characterized viral proteins, or portions of them, produced via molecular biology or peptide chemistry have proven to be promising tools in improving serological diagnosis [6-14]. The analysis of the humoral immune response elicited during natural infection has repeatedly shown that the basic phosphoprotein of 150 kD encoded by UL32 (ppUL32) [15] and localized in the viral tegument is highly immunogenic and is recognized by sera from nearly 100% of the HCMV-seropositive subjects tested [6, 16]. In this molecule at least two epitopes have been identified and shown to react efficiently with human immunoglobulins. In particular, the analysis of several ppUL32 fusion proteins showed that a region localized in the sequence between aa 1006 and aa 1048, inclusive, read in 5'→3' direction, of the molecule (aa 1006-1048) reacts with more than 80% of IgM-positive sera [9,11]. When chemically synthesized, another region localized between aa 595 and 614 gave a positive reaction with almost 100% of IgG-positive sera [17]. When expressed as recombinant protein this region has proven to react very efficiently also with HCMV-specific IgM [12]. The two coding regions were recently fused together and shown to produce a double epitope fusion protein which can replace the entire pp150 molecule in its IgM-binding ability [12].

- 5 -

Another HCMV protein which reacts very well with IgM is the non structural DNA-binding phosphoprotein of 52 kD [18] encoded by viral gene UL44 (ppUL44) [7, 11]. The COOH part of the molecule shows efficient IgM binding and does not contain relevant aminoacid sequences cross-reacting with the homologous protein of other members of the *Herpesviridae* family (that are mainly present in the NH₂ half) [19]. The COOH half of this protein was expressed and tested with human sera. The DNA sequence expressing this region has been in fact inserted into plasmid p-ROS (which carries a truncated sequence of LacZ) at site SmaI and cloned into E.coli, thus obtaining a fusion protein H10, which has yielded good results [21].

Besides ppUL32 and ppUL44, two other HCMV proteins are well known: the major matrix phosphoprotein of 65 kD encoded by viral gene UL83 (ppUL83) and the assembly phosphoprotein of 38 kD encoded by the viral gene UL80a (ppUL80a). The first is known to induce a strong IgM response and antibodies reacting exclusively to this protein were described during primary infection [7]. Likewise, the IgM to this second protein are very often the first marker to be detected in HCMV seropositive transplanted patients undergoing a viral reactivation [Kraat et al. 1995, manuscript submitted]. Moreover, some of the Applicants observed a strong IgM reactivity to this protein in congenitally infected newborns (Lazzarotto and Landini, unpublished observation).

However, the above studies have not been successful, so far, in producing such recombinant proteic material, as are actually capable of replacing the virus or virus components (purified virions) in the diagnostic kits currently in use.

5 Disclosure of the Invention:

A primary object of the present invention is to provide a recombinant proteic material, in particular, a fusion protein, capable of binding by immunoreaction with antibodies produced against human Cytomegalovirus, in particular, both IgM and IgG, capable of being used
10 as an antigen for detecting such antibodies in the relevant serologic tests, with substantially the same efficacy as the virus and/or of infected cell lyses.

A further object of the present invention is to furnish diagnostic reagents and kits derived from said proteic material.

15 A further object of the present invention is to provide a plasmid, capable of being inserted into a prokaryote and/or eukaryote host organism so as to express said recombinant proteic material as a fusion protein.

It is also an object of the present invention to provide
20 recombinant antigens, in particular fusion proteins, to detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay.

A further object of the present invention is to furnish diagnostic reagents and kits derived from said proteic materials.

Finally, a further object of the present invention is to provide plasmids, capable of being inserted into a prokaryotes and/or eukaryotes host organism so as to express said recombinant proteic material as a fusion protein, in particular fused with CPT: CMP-3-deoxy-D-manno-
5 octulosonate cyidylyl-transferase or CMP-KDO transferase (CKS) protein.

According to the present invention, there is provided a recombinant proteic material, capable of binding with antibodies against human Cytomegalovirus (HCMV), characterized in that it consists of a fusion protein comprising a first region, carrying at least part of the
10 sequence between aa 202 and aa 434, inclusive, read in 5'→3' direction, of protein pp52, a second region, carrying at least a part of the sequence between aa 1006 and aa 1048, inclusive, read in 5'→3' direction, of pp150, and a third region, carrying at least a part of the sequence between aa 595 and 614 inclusive, in 5'→3' direction, of
15 pp150; said regions capable of being variably arranged with respect to one another within the fusion protein.

Said regions preferably comprise said sequences of pp52 and pp150 in full, and, according to a preferred, non-limiting, embodiment, a fusion protein according to the invention comprises in a sequential
20 manner, running 5'→3': said first region, immediately downstream wherefrom there is placed said third region, and then said second region, downstream from said third region, a bridge sequence being inserted between said third and second regions.

Said bridge sequence consists, running 5'→3', of the following

series of aminoacids: lys leu gln glu phe.

The present invention further provides for a reagent for diagnosing HCMV infection by serological methods, characterized in that it comprises a fusion protein as defined hereinabove.

5 In particular, the diagnostic reagent comprises a fusion protein, whereof at least part of the amino acid is encoded by the nucleotide sequence shown in the sequence listing as SEQ ID NO: 1, read from nucleotide 001 to nucleotide 900.

The present invention also relates to diagnostic reagents for
10 direct detection of HCMV through serology, comprising at least one monospecific polyclonal serum or monoclonal antibodies (MaB), directed against said fusion protein.

Finally, the present invention relates to a plasmid, capable of being inserted into a prokaryote or eukaryote host organism,
15 characterized in that it comprises, combined together, a first DNA sequence encoding at least part of the sequence between aa 202 to aa 434, inclusive, read in 5'→3' direction, of protein pp52 of HCMV, a second DNA sequence encoding at least part of the sequence between aa 1006 and aa 1048, inclusive, read in 5'→3' direction, of protein
20 pp150, and a third DNA sequence encoding the amino acid sequence, running 5'→3', between aa 595 and aa 614, included, of protein pp150.

In particular, said plasmid comprises either the nucleotide

.2194135

sequences shown in the sequence listing as SEQ ID No: 1, from nucleotide 002 to nucleotide 907, or that shown as SEQ ID No: 3.

Moreover, according to an improved aspect of the present invention, there is provided a mixture of recombinant antigens to
5 detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay, the mixture being characterized in that it includes, in combination:

(i)- a first fusion protein comprising: a first region, carrying an amino acid sequence (H10) corresponding to at least part of the sequence
10 between aa 202 and aa 434, inclusive, read in 5'→3' direction, of viral protein pp52; a second region, carrying an amino acid sequence (F3) corresponding to at least part of the sequence between aa 1006 to aa 1048, inclusive, read in 5'→3' direction, of viral protein pp150; and a third region, carrying an amino acid sequence (A1C2) corresponding to
15 at least part of the sequence between aa 595 and aa 614 inclusive, read in 5'→3' direction, of the same viral protein pp150; said regions capable of being variably arranged with respect to one another within the first fusion protein;

(ii)- a second fusion protein comprising an immunogenic region
20 consisting in a sequence of amino acids corresponding to at least part of the sequence between aa 297 to aa 510, inclusive, read in 5'→3' direction, of the major matrix protein pp65 encoded by the viral gene UL83; and

- 10 - 2194135

(iii)- a third fusion protein comprising an immunogen region consisting in a sequence of amino acids corresponding to at least part of the sequence between aa 117 to aa 373, read in 5'→3' direction, of the viral assembly protein pp38 encoded by the viral gene UL80a.

5 In particular, said regions include said amino acid sequences of pp52, pp150, pp65 and pp38 in full, fused together with protein CKS or at least a part thereof .

The invention relates, furthermore, to a plasmid, capable of being inserted into a prokaryote or eukaryote host organism, characterized
10 in that it comprises a DNA sequence encoding at least part of the sequence between aa 297 to aa 510, read in 5'→3' direction, of the HCMV major matrix protein pp65 encoded by the viral gene UL83, said sequence being linked at the 3' end of a second sequence corresponding to the sequence from amino acid 1 to amino acid 240
15 inclusive encoding for protein CKS.

The invention relates also to a further plasmid, capable of being inserted into a prokaryotes or eukaryotes host organism, characterized in that it comprises a DNA sequence encoding at least part of the sequence between aa 117 to aa 373, read in 5'→3' direction, of the
20 HCMV assembly protein pp38 encoded by the viral gene UL80a, said sequence being linked at the 3' end of a second sequence corresponding to the sequence from amino acid 1 to amino acid 240 inclusive encoding for protein CKS.

Finally, the invention relates to the use of a mixture of mono- and poly- epitope fusion proteins to detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay, the mixture being characterized in that it includes, in combination:

- 5 (i)- a first fusion protein comprising: a first region, carrying an amino acid sequence (H10) corresponding to at least part of the sequence between aa 202 and aa 434, inclusive, read in 5'→3' direction, of viral protein pp52; a second region, carrying an amino acid sequence (F3) corresponding to at least part of the sequence between aa 1006 to aa
10 1048, inclusive, read in 5'→3' direction, of viral pp150; and a third region, carrying an amino acid sequence (A1C2) corresponding to at least part of the sequence between aa 595 and 614 inclusive, read in 5'→3' direction, of the same viral protein pp150; said regions capable of being variably arranged with respect to one another within the first
15 fusion protein;
- (ii)- a second fusion protein comprising an immunogenic region consisting in a sequence of amino acids corresponding to at least part of the sequence between aa 297 to aa 510, inclusive, read in 5'→3' direction, of the major matrix protein pp65 encoded by the viral gene
20 UL83 ; and
- (iii)- a third fusion protein comprising an immunogenic region consisting in a sequence of amino acids corresponding to at least part of the sequence between aa 117 to aa 373, read in 5'→3' direction, of the

viral assembly protein pp38 encoded by the viral gene UL80a.

The invention further relates to the use of the above mixture of fusion proteins to produce a diagnostic kit to detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay.

5 According to the first aspect of the present invention, starting from the studies on polyepitope fusion proteins of some of the Applicants [12],

 a series of clones have been prepared, operating with plasmid p-ROS, as according to the well known techniques described
10 by Maniatis et al. (1982. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory. Cold Spring Harbor, N.Y.); these clones, once inserted into E. coli and activated with IPTG, are capable of producing a whole range of fusion proteins comprising a portion of β -galactosidase (β -Gal), which were subsequently comparatively tested
15 using the known techniques (Western blott, Immunoblott, ELISA, etc.) with HCMV-positive sera for HCMV-specific IgM and IgG from transplant recipients in order to ascertain their different detection levels.

 In particular, there have been prepared a first plasmid, pMB28,
20 capable of expressing aminoacid sequence A1C2; a second plasmid, pGT1, capable of expressing a new immunogenic sequence F3 of ppUL32, comprising the last 44 aminoacids of COOH terminus of viral protein pp150 and encoded by UL32 nucleotides from 3013 to 3144,

running 5'→3'; and a third plasmid, pMB34, capable of expressing A1C2 and F3 as a single fusion protein with β-Gal; said first plasmid having been obtained by inserting a modified and PCR amplified DNA fragment between sites Sall and HindIII of the polyclonation site of prokaryotes expression vector p-ROS according to Platchter et al. (1992. "Detection of cytomegalovirus..."; J. Clin. Microbiol. 30; 201-206); the second plasmid was obtained by inserting into site SmaI of p-ROS a DNA EcoRI fragment from a LambdaGT11 clone, suitably altered by mutagenesis by means of vector p-Alter, so as to replace a STOP codon with a corresponding EcoRI site. The third plasmid was obtained from plasmid pMB28, inserting therein the same 78 bp (base pairs) EcoRI fragment encoding F3 used for pGT1 at site StuI, thus allowing, between said two DNA sequences encoding A1C2 and F3, for a bridge sequence consisting of five aminoacid : lys leu gln glu phe (K, L, Q, E and F, according to IUPAC). Moreover, the known, highly immunogenic epitope (H10) of viral protein pp52, consisting of the last 233 aminoacid of COOH terminus of the protein itself, has also been included in the comparison.

To this end, the nucleotide sequence H10, contained in the genome of phage Lambda GT11 between two closely arranged EcoRI restriction sites, immediately upstream from lacZ, is obtained according to the well known techniques described by Berg et al. (Proc. Nat. Acad. Sci, 1999, 69: 2904-2909) and Kaiser et al. (1973 J. Mol. Biol 78: 453-47) by digestion with EcoRI; subsequently, the cohesive ends are caused to

become blunt by a filling-in operation in the presence of DNA polymerase, wherein the adenine/thymine bases (A/T), unpaired, are paired with the complementary nucleotides. In the meantime, plasmid p-ROS is digested with SmaI, which causes its opening and linearization; then, EcoRI H10 fragment of Lambda GT11 and the open plasmid are incubated together in the presence of DNA lygase, thus causing fragment H10 to be inserted at the appropriate site SmaI provided for in p-ROS and a plasmid pGH10 to be formed. The latter is in turn inserted in E.coli by electrophoration and cloned.

10 Finally, a last "mixed" plasmid, comprising both DNA sequences encoding A1C2 and F3 and the one encoding H10 was prepared from the previous plasmids. To this end, a pair of oligonucleotides were synthesized for pGH10 and one for pMB34, respectively, their sequences allowing for them to be used for chain amplification with
15 PCR of sequences H10 and A1C2-F3, respectively; said oligonucleotide pairs, containing sequences, capable of introducing suitable recognition sites for given restriction enzymes (in the case in point, EcoRI) at 5' ends, and sequences in common between 5' and 3' of H10 and A1C2-F3, respectively at opposite ends, were used for amplifying a single
20 DNA fragment containing said three epitopes, already in the desired reading frame and ready for cloning at the target sites of the expression vector, in the case in point, at the SmaI site of pROS for the epitopes to be expressed as fusion protein with β -Gal; in the last analysis, the

- 15 -

resulting plasmid, pMB40, therefore comprises, at site SmaI, DNA sequence SEQ ID NO: 1, corresponding to nucleotides 002 to 907, encoding a new fusion protein comprising, at the same time, both the most immunogenic known epitope of pp150 (A1C2) and the most
5 immunogenic epitope of pp52 (H10), and the new immunogenic epitope of pp150 (F3), deriving from the addition of further 19 aminoacids to the known epitope of pp150, consisting of the last 25 aminoacids (sequence D1) at the COOH terminus of the said protein.

Surprisingly, said new fusion protein, H10/A1C2/F3, was found
10 to have a much higher IgG and IgM reactivity than that obtainable by properly combining in one single test kit the proteins containing the three distinct epitopes. A synergic effect, therefore, takes place, probably owing to the fact that, arranging such epitopes, as are normally far off, close to one another on the same viral protein (A1C2
15 and F3 of ppUL32, for instance) or even on different proteins (such as H10 of ppUL44), causes conformational epitopes, likely to be present in the original complete viral proteins and/or on the virions, to be "mimicked".

The new fusion protein, comprising the said sequences, has
20 indeed proven to be as active as the purified virions in binding with IgG and IgM of infected subjects, thus finally allowing for a standardized antigen, both as to quality and as to quantity, capable of actually replacing purified virions and/or infected cells in the preparation of

diagnostic kits for HCMV infection serodiagnosis. Obviously, the previously described epitopes may be included into the new fusion protein as according to the present invention in the order of sequence SEQ ID NO: 2, currently preferred, or else in any other sequence, also
5 allowing for bridge sequences to be inserted between them, like that corresponding to nucleosides 757 to 771 of SEQ ID NO: 1.

Plasmid pMB40, expressing such a protein, can be inserted into a prokaryote host organism, such as E.coli. Besides, the specific method described for obtaining the "mixed" fragment further allows for
10 its cloning in a different vector, in particular, after digestion with EcoRI, at the corresponding site of vector pHIL-S1, thus obtaining plasmid pHIL-D1, which allows for the recombinant protein to be expressed in Pichia pastoris, fused with a short peptide representing a secretion signal outside the yeast cell, in the culture medium. Such a manner of
15 expression allows, among other things, for fusion protein production to be induced by the addition of methanol, instead of IPTG, as used for the expression in E.coli., thus representing a far less expensive method. Consequently, production of the poly-epitope fusion protein as according to the present invention can be large-scale, simple and
20 economical.

According to a further aspect of the present invention, and starting from the results described above, the poly-epitope protein according to the first aspect of the invention has been further studied

arranging the said regions in a different way and combining said recombinant material, expressed fused together with protein CKS, with immunogenic regions of other HCMV proteins, so obtaining a mixture of fusion proteins high effective against IgM.

5 The fusion proteins according to the present invention can be used alone or, according to the main aspect of the present invention, mixed together, for direct preparation of diagnostic reagents for serological detection of HCMV-specific antibodies, in that their ability of efficiently binding anti-HCMV IgM, in particular those produced during a
10 primary asymptomatic phase of the infection, is substantially 100%. Such reagents may comprise the mixture of fusion proteins according to the invention, eventually in combination with other HCMV antigens, or else more complex fusion proteins, including the immunogenic sequences repeated many times and anyway in combination with one
15 another.

 The proteic materials represented by the fusion proteins according to the present invention can be further used for the preparation, by means of known techniques, of diagnostic reagents for serological detection of human Cytomegalovirus and/or antigens for the
20 detection thereof represented by specific polyclonal sera, or by monoclonal antibodies (MaB) directed against the proteins of the invention and obtained via a number of different methods.

Said diagnostic reagents, containing the proteins according to the

present invention, can be used, in turn, for the production of diagnostic kits for the demonstration, by serological methods, of the appearance of antibodies to human Cytomegalovirus (HCMV), wherein the reagent is adsorbed on nitrocellulose paper (blotting). Finally, the proteins according to the present invention can be used, after being purified, in ELISA assays, latex agglutination tests, RIA, as well as in all known serological screening methods for the presence of HCMV-specific IgG and IgM, both manual and partially or completely automatized.

Brief Description of the Drawings:

- 10 - Fig. 1 illustrates schematically the position of the DNA fragments encoding two of the portions characterizing one of the fusion proteins according to the invention, in gene UL32 contained in EcoRI-Y fragment of the virus strain AD169;
- Fig. 2 schematically illustrates a method for preparing plasmids used in the invention;
- 15 - Figs. 3 and 4 are schematic representations of the construction of plasmids pMB28: lacZ-A1C2 and pMB34: lacZ-A1C2F3;
- Fig. 5 schematically illustrates a method for preparing plasmids used in the invention;
- 20 - Figs. 6 and 7 show two tables comparing experimental data relating to the detection of IgG and IgM from HCMV-infected subjects, by ELISA assay, using the entire virus or different recombinant fusion proteins, including one according to a first aspect of the present invention;
- Fig. 8 is a schematic representation of the construction of plasmids

pJO210 and pJO215;

- Fig. 9 is a schematic representation of the construction of plasmid pMC200;

5 - Fig. 10 is a schematic representation of the construction of plasmid pJO200;

- Fig. 11 is a schematic representation of the construction of plasmid pMB38: lacZpp38(106-373aa);

- Fig. 12 is a schematic representation of the construction of plasmid pCMV-1A: CKS-A1C2F3-CKS;

10 - Fig. 13A shows preparation of PCR fragments containing the A1C2F3 and H10 DNA sequences;

- Fig. 13B shows preparation of PCR fragments containing the pp65(297-510aa) and pp38(117-373aa) DNA sequences;

15 - Fig. 13C is a schematic representation of the construction of: plasmid pCMV-3B:CKS-H10, plasmid pCMV-4:CKS-A1C2F3-H10, plasmid pCMV-9:CKS-pp65(297-510aa), and plasmid pCMV-26:CKS-pp38(117-373aa);

- Fig. 14A is a schematic representation of the construction of plasmid pJO200- Δ Mlul;

20 - Figs. 14B, C, D show respectively: the nucleotide sequence of the plasmid pJO200, including the intended modification site at plasmid residues 151 to 180 (5'-3'); the double-stranded structure of the mutagenic oligonucleotide, 5'CGCGACGT3', synthesized for ligation

into plasmid pJO200/MluI/CIAP; and the nucleotide sequence of the plasmid pJO200 Δ MluI, including the modified residues 151 to 180 (5'-3');

- Fig. 15 is the schematic representation of the construction of plasmid pEE1;

- Fig. 16A shows the preparation of PCR fragments containing the A1C2F3-H10, pp65(297-510aa), and pp38(117-373aa) DNA sequences;

- Fig. 16B is a schematic representation of the construction of: plasmid pCMV-27:CKS-A1C2F3-H10-CKS, plasmid pCMV-28:CKS-pp65(297-510aa)-CKS, and plasmid pCMV-29:CKS- pp38(117-373aa)-CKS;

- Figs. 17A and 17B shows the follow up in six transplanted patients undergoing primary and, respectively, secondary, HCMV infection, wherein graphs A1 and A2 relate to kidney transplant recipients and graphs A3, B1, B2 and B3 relate to heart transplant recipients; and

- Figs. 18 to 25 show tables comparing experimental data relating to the detection of IgM by EIA.

Best mode of carrying out the Invention:

The present invention will be better described by way of non-limiting examples thereof, with reference to the accompanying drawings.

EXAMPLES

REAGENT, ENZYMES AND CULTURE MEDIUM - Restriction enzymes,

- 21 -

T4 DNA ligase, calf Intestinal alkaline phosphatase (CIAP), polynucleotide kinase, and the Klenow fragment of DNA Polymerase I were purchased from New England Biolabs, Inc., Beverly, Massachusetts or from Boehringer Mannheim Corp., Indianapolis, Indiana. DNA and Protein molecular weight standards, Dalichi pre-cast gradient polyacrylamide gels, and Semi-Dry Blotting System* with buffers were obtained from Integrated Separation Systems, Inc., Natick, Massachusetts. Isopropyl- β -D-thiogalactoside (IPTG), acrylamide, N-N'-methylene-bis-acrylamide. N, N,N',N'-tetramethylethylenediamine (TEMED), 4-chloro-1-naphthol, and sodium dodecylsulfate (SDS) were purchased from BioRad Laboratories, Richmond, California. Horseradish peroxidase (HRPO)- labeled antibodies were purchased from Kirkegeard & Perry Laboratories Inc. Gaithersburg, Maryland. "EPICUREAN COLI XL-1 Blue"*(recA1, endA1, gyrA96, thi-l, hsdR17, supE44, relA1, lac [F'proAB laI^qZdM15 Tn10 (Tet')]) Supercompetent E. coli cells, DNA isolation kit, RNA isolation kit, and ZAP-cDNA* Synthesis kit were obtained from Stratagene Cloning Systems Inc. La Jolla, California. GeneAmp*reagent kit and AmpliTaq*DNA Polymerase were purchased from Perkin-Elmer Cetus, Norwalk, CT. Nucleotide kit for DNA sequencing with Sequenase*and 7-deaza-dGTP and Sequenase version 2.0 DNA Polymerase were obtained from U.S. Biochemical Corp., Cleveland, Ohio. PolyA+ mRNA purification kit was purchased from Pharmacia LKB Biotechnology Inc.,

* Trade-mark

Piscataway, New Jersey. Luria Broth plates with ampicillin (LiSamp plates) were purchased from Micro Diagnostics, Inc., Lombard, Illinois. "OPTI-MEM"* Medium, fetal calf serum, phosphate-buffered saline (PBS), competent E.coli DH5 α (F' ϕ 80dlacZdM15 d(lacZYA-argF)U169
5 deoR recA1 endA1 phoA hsdR17 supE44 λ thi-1 gyrA96 relA1), and "ULTRAPURE"* agarose were purchased from GIBCO BRL, Inc., Grand Island, New York. Bacto-Tryptone, Bacto-Yeast Extract, and Bacto-Agar were obtained from Difco Laboratories, Detroit, Michigan. "NZY" Broth was purchased from Becton Dickinson Microbiology Systems,
10 Cockeysville, Maryland. Salmon Sperm DNA, lysozyme, ampicillin, N-lauroyl sarcosine, thimerosal, buffers, casein acid hydrolysate, urea, surfactants, like "TWEEN 20",* diethylpyrocarbonate (DEPC), and inorganic salts were purchased from Sigma Chemical Co., St. Louis, Missouri. "SUPERBROTH II"* contained 11.25 g/L tryptone, 22.5 g/L
15 yeast extract, 11.4 g/L potassium phosphate dibasic, 1.7 g/L potassium phosphate monobasic, 10 ml/L glycerol, adjust pH to 7.2 with sodium hydroxide. Tris-buffered saline (TBS) consisted of 20 mM Tris, 150 mM NaCl, pH 7.5. Tris-buffered saline "TWEEN 20" consisted of TBS + 0.05% "TWEEN 20". Membrane blocking solution consisted of 1%
20 Bovine Serum Albumin, 1% Casein acid hydrolysate, 0.05% "TWEEN 20" in TBS. Rubazyme specimen dilution buffer (Rubazyme SDB) consisted of 100 mM Tris, pH 7.5, 135 mM NaCl, 10 mM EDTA, 0.2 % "TWEEN 20", 0.01% thimerosal, and 4% bovine calf serum. Rubazyme
* Trade-mark

conjugate diluent dilution buffer consisted of 100 mM Tris, pH 7.5, 135 mM NaCl, 0.01% thimerosal, and 10% bovine calf serum. "HRP" color development solution consisted of 0.06% 4-chloro-1-naphthol, 0.02% H₂O₂, and 0.2% methanol in TBS. SDS-PAGE loading buffer consisted
5 of 62 mM Tris, pH 6.8, 2% SDS (Sodium Dodecyl Sulphate), 10% glycerol, 5% β-mercaptoethanol, and 0.1% bromophenol blue. TE buffer consisted of 10 mM Tris, 1 mM EDTA, pH 8.0.

VIRUS PROPAGATION AND PREPARATION OF cDNA - CMV strain AD169 and Towne were propagated in human fibroblasts grown in
10 "OPTI-MEM" Medium supplemented with 5% fetal calf serum. After 6 days post-infection, the infected cells were harvested by centrifugation, washed with PBS, and homogenized with a glass-PTFE ("TEFLON")* homogenizer. Total viral DNA was isolated as described in Mocarski, E.S. et al. Proc.Nat.Acad.Sci 82:1266, 1985. Total RNA was isolated
15 from the homogenized cells using the RNA Isolation Kit (Stratagene Cloning Systems, La Jolla, CA) and polyA+ RNA was isolated using a mRNA Isolation Kit (Pharmacia Biotech, Piscataway, NJ). HCMV cDNA was synthesized from the purified viral mRNA using a "ZAP-cDNA" Synthesis Kit (Stratagene Cloning Systems, La Jolla, CA).

20 GENERAL METHODS - All enzyme digestions of DNA were performed according to suppliers' instructions. At least 5 units of enzyme were used per microgram of DNA, and sufficient incubation time was allowed for complete digestion of DNA. Supplier protocols were followed for the

* Trade-mark

- 24 - 2194135

various kits used in manipulation of DNA and RNA, and also for PCR and sequence of DNA. Standard procedures were used for (1) small scale preparation and large scale preparation of plasmid DNA from E. coli, (2) preparation of phage lysate DNA from E. coli cells infected with phage λ , (3) preparation of phage lysates for the absorption of anti-E.coli antibodies, (4) extraction with phenol-chloroform, (5) precipitation of DNA with ethanol, (6) restriction analysis of DNA on agarose gels, (7) purification of DNA fragments from agarose and polyacrylamide gels, (8) filling the recessed 3' termini created by digestion of DNA with restriction enzymes using the Klenow fragment of DNA Polymerase I, (9) ligation of DNA fragments with T4 DNA ligase, and (10) preparation of competent E.coli TB1 cells (F^- ara d(lac-proAB) rpsL ϕ 80dlacZdM15 hsdR17) as described in Sambrock, Fritsch, and Maniatis, Molecular Cloning: A Laboratory Manual, 2nd ed., New York: Cold Spring Harbor Laboratory Press 1989, the content of which is hereby incorporated by reference. The DNA fragments for cloning into plasmids that were generated by PCR amplification were extracted with phenol-chloroform and precipitated with ethanol prior to restriction enzyme digestion of the PCR reaction mixture. Oligonucleotides for PCR and DNA sequencing were synthesized on an Applied Biosystems Oligonucleotide Synthesizer, model 380B or 394, per manufacturer's protocol.

EXAMPLE 1

- 25 -

Construction of lacZ-H10 expression Vector pROSH10

Operating with recombinant DNA standard methods, as described by Maniatis et al. (1982. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory. Cold Spring Harbor, N.Y.) and from plasmids
5 pROS supplied by the authors thereof (cf. Ellinger et al.), the genome of a clone, obtained from early gene expressing phage Lambda Gt11, which produces a fusion protein with β -galactosidase, highly reactive with CMV-positive sera, was purified. The genoma was then digested with restriction enzyme EcoRI to extract the DNA fragment expressed
10 by the phage. The fragment was then cloned into vector pUC18 and sequenced. The sequence represented a fragment of 699 base pairs expressing 233 aminoacids at COOH terminus of pp52 (UL44). The fragment was then cloned at site SmaI of expression vector pROS, after filling in the EcoRI ends of the fragment, as illustrated schematically in
15 Fig. 2. The plasmid so obtained was named pROSH10.

EXAMPLE 2Construction of lacZ-F3 expression Vector pGT1

Operating with the methods described in Example 1, the genome of a clone, obtained from early gene expressing phage Lambda Gt11, which
20 produces a fusion protein with β -galactosidase, highly reactive with CMV-positive sera, was purified. The genoma was then digested with restriction enzyme EcoRI to extract the DNA fragment expressed by the phage. The fragment was then cloned into vector pUC18 and

- 26 - 2194135

sequenced. The sequence represented a fragment of 132 base pairs expressing 44 aminoacids at COOH terminus of p150 (UL32). The fragment was then cloned at site SmaI of expression vector pROS, after filling in the EcoRI ends of the fragment, as illustrated schematically in Fig. 2 referred to Example 1. The plasmid so obtained was named pGT1.

EXAMPLE 3

Construction of lacZ-A1C2F3 expression Vector pMB34

The plasmid pMB34 is a derivative of the lacZ expression vector pROS described in Ellinger et al., J. Clin. Micro. 27: 971, 1989. This vector contains a truncated form of the lacZ gene (1-375 amino acids) with a polylinker cloning site located downstream of the lacZ gene. The pMB34 plasmid was constructed in two steps by joining two regions of ppUL32, which encodes the basic phosphoprotein of 150 kD of HCMV, to the 3' end of the lacZ gene in pROS.

A. Construction of pMB28: lacZ-A1C2

The plasmid pMB28 is a derivative of plasmid pROS as shown in Fig. 3. This plasmid was constructed by cloning a DNA fragment containing HCMV-A1C2, obtained by PCR amplification of Human Cytomegalovirus (HCMV) genomic DNA from the region of ppUL32 encoding amino acids 595-614 of pp150 (nucleotides 1763-1842), into the polylinker region of pROS. Large scale plasmid DNA (pROS) was isolated from DH5α cells operating as described in general methods. Plasmid pROS was

- 27 -

digested with Sall and HindIII and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide 1783 of ppUL32 containing a Sall site and an antisense primer starting at nucleotide 1842 of ppUL32 containing a HindIII site were synthesized and added
5 to a PCR reaction mixture containing genomic HCMV DNA. After PCR amplification, the reaction mixture was digested with Sall and HindIII, and the 60 base pair fragment containing A1C2 (nucleotides 1733-1842) was purified on an agarose gel. This purified fragment was then ligated to purified pROS/Sall/HindIII overnight at 16°C. The next day
10 the ligation mixture was transformed into competent DH5α cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the 60 base pair fragment in pROS at the end of the lacZ gene. Plasmid pMB28, which contains the A1C2 fragment, was isolated. The DNA sequence of A1C2 in pMB28 was
15 confirmed and the A1C2 coding region was in frame with the lacZ coding sequence.

B. Construction of pMB34: lacZ-A1C2F3

The plasmid pMB34 is a derivative of plasmid pMB28, as shown in Fig. 4. The plasmid pMB34 was constructed by cloning a DNA fragment,
20 containing HCMV-F3 obtained from a λgt11 subclone of ppUL32 encoding amino acids 1006-1043 of ppUL32 (nucleotides 3016-3144) derived from the λgt11 library of Mocarski, E.S. et al. Proc. Nat. Acad. Sci 82: 1266, 1985, into the polylinker region of pMB28 just

- 28 - 2194135

downstream of the A1C2 DNA sequence. Large scale plasmid DNA (pMB28) was isolated from DH5 α cells as described in general methods. Phage lysate DNA was prepared from phage λ gt11 clone λ -F3 as described in general methods. Plasmid pMB28 was digested with

5 Stul and the vector backbone with blunt-ends was purified on an agarose gel. Phage λ -F3 DNA was digested with EcoRI and the recessed 3' termini were filled in with the Klenow fragment of DNA Polymerase I, leaving blunt-ends. The blunt-ended 129 base pair λ -F3 fragment was purified on an agarose gel and then blunt-end ligated to

10 pMB28/Stul overnight at 16°C. The next day the ligation mixture was transformed into competent DH5 α cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the 129 base pair fragment in pMB28 at the end of the lacZ gene in the correct orientation. Plasmid pMB34, which contains the F3 fragment in

15 the correct orientation, was isolated. The DNA sequence of F3 in plasmid pMB34 was confirmed and the F3 coding region was in-frame with the lacZ-A1C2 coding sequence. The coding region of the lacZ-A1C2F3 construct in pMB34 contains a bridge of 5 amino acids between A1C2 and F3 as shown below using the International

20 Standard one-letter Codification for amino acids (according to which: L is Leu, Q is Gln, K is Lys, E is Glu, F is Phe):

(1) lacZ(1-375aa)-A1C2(595-614aa,pp150)-**K-L-Q-E-F**-F3(1006-1048aa,pp150)

EXAMPLE 4

- 29 -

Construction of lacZ-H10A1C2F3 Expression Vector pMB40

Starting from the previously described plasmids, and operating with such methods, as have been described in Examples 1, 2 and 3 hereinabove and as illustrated in Fig. 5, two pairs of oligonucleotides
 5 were synthesized, the sequences whereof allow for their use for chain amplification, with PCR, of H10 and A1C2-F3 sequences, respectively, localized in plasmids pGH10 and pMB34. The oligonucleotide sequences composing said primers are as follows:

EcoH10: GAA TTC ACA GCC AAT AAC CGC GTC AGT TTC;

10 **H10_{A1C2}: AGG CGT CGG CGT GCC GCA CTT TTG CTT CTT GGT GTT**

to amplify H10 from pGH10, introduce a site EcoRI in 5' in the amplified sequence and a sequence of 12 base pairs, homologous of A1C2, in 3' in the amplified fragment; and

H10A1C2: CAA AAG TGC GGC ACG CCG ACG CCT GTC AAT CCT TCC;

15 **F3_{Eco}: GAA TTC CTA TTC CTC CGT GTT CTT AAT CTT**

to amplify A1C2-F3 fragment from plasmide pMB34, introducing in the amplified product, at 5', a sequence, homologous of H10, and an EcoRI site in 3'.

The two fragments, resulting from said two independent amplifications
 20 (**EcoH10_{A1C2}** and **H10A1C2-F3_{Eco}**) were then mixed, together with external primers **EcoH10** and **F3_{Eco}**, and subjected to additional amplification. Owing to sequences **3'_{Eco}H10_{A1C2}** and **5'_{H10}A1C2-F3_{Eco}** being complementary, the amplified product contains all three epitopes and has an EcoRI site at each end.

The amplified sequence was then cloned into pROS and the plasmid derived therefrom, named pMB40, includes, in connection with corresponding site SmaI, DNA sequence SEQ ID No: 1, corresponding to nucleotides 002 to 907.

5

EXAMPLE 5

The plasmids described in Examples 1 to 4 hereinabove, were inserted in E.coli by electroporation and cloned; synthesis of the desired fusion protein was then induced by the addition of IPTG and the proteins so obtained were made competent by lysis of bacterial cells and
10 denaturation of the cell lysates with SDS and β -mercaptoethanol; the proteins were then run in 9%-12% acrylamide gel with SDS-PAGE and then transferred to nitrocellulose, where immunoreactions took place; a preliminary screening of a number of serum samples was carried out by Miniblotter* (Immunetics, Cambridge-Mass. USA); both IgG purified
15 ("ENDOBU LIN*", of Immuno AG, Wien AT) and a group of IgM-highly positive sera were used at a dilution of 1:100. Individual human serum samples were diluted at 1:80 for IgG and IgM detection; a peroxidase-conjugated anti-g or anti- μ chain was used as a second antibody.

Two groups of human serum samples were used: the first group of sera
20 consisted of seventeen HCMV-positive samples, eight whereof with a high IgG titre to HCMV and seven a medium/low level of HCMV-specific IgG, as detected by ELISA; the presence of HCMV-specific IgG was confirmed by immunoblotting. The second group of sera consisted of

* Trade-mark

nineteen HCMV-positive samples, ten whereof with a high IgM titre to HCMV and nine a medium/low level of HCMV-specific IgM, as detected by ELISA; also in this case, the presence of HCMV-specific IgM was confirmed by immunoblotting.

- 5 The evaluation of anti-HCMV IgG was carried out with a CMV kit of M.A. Bioproducts (Walkersville, Md. USA); plates were read on a "microELISA" automatic reader (Dynatech Products, Alexandria, Va. USA). To perform linear regression analysis and to standardize the test run, every plate included three serum calibrators. The evaluation of
- 10 anti-HCMV IgM was carried out with a CMV IgM "ELA"* kit of Technogenetics (Hamburg, Germany). The results were interpreted as suggested by the manufacturers.

In order to avoid false positive results, all samples were also subjected to a test run for the detection of Rheumatoid factor by latex

15 agglutination (Rheuma-Wellco test of Wellcome, Dartford, Great Britain) and only the negative samples were used for comparison. Results are shown in Tables 1 and 2 of Figs. 6 (IgG) and 7 (IgM).

As it is clear from the data presented, not only have the results obtained with H10/A1C2/F distinctly improved, as compared to those

20 obtained with the other fusion proteins, both to IgG and to IgM, but they are fully superimposable to the results obtained with antigens composed of the entire virus.

EXAMPLE 6

* Trade-mark

Construction of CKS Expression Vector pJO200

The CKS expression vector pJO200 allows the fusion of recombinant proteins to the CMP-KDO synthetase (CKS) protein. The DNA sequence for the structural gene encoding CKS (the kdsB gene) is published in Goldman et al. , J. Biol. Chem. 261:15831, 1986. The amino acid sequence, 248 total, for CKS derived from the DNA sequence, is described in the same article. This pJO200 vector is constructed by a three-step procedure starting with the plasmid pTB201 (Figure 8) described in Bolling and Mandecki, Biotechniques 8: 468, 1990. The construction plan for plasmid pJO200 involves the removal of two restriction enzyme sites and the addition of a multi-cloning site at the 3' end of the CKS gene. This was done to facilitate the cloning of HCMV (Human Cytomegalovirus) genes encoding protein antigens at the 3' end of CKS. The completed vector contains the coding sequence for 240 amino acids of the original kdsB gene plus an additional 20 amino acids at the end of the CKS gene contributed by the polylinker DNA sequence, for a total of 260 amino acids.

A. Construction of pJO210

The plasmid, pJO210, is a derivative of the CKS expression vector, pTB201. This plasmid was constructed by removing a single EcoRI site present in pTB201 located upstream from the promoter for the CKS gene. Large scale plasmid DNA (pTB201) was isolated from TB1 cells using the techniques described in the section above named

- 33 -

"General Methods". The DNA was digested with EcoRI to completion and purified on a polyacrylamide gel. The purified pTB201/EcoRI fragment was then treated with the Klenow fragment of DNA Polymerase I in the presence of deoxynucleotide triphosphates. This
5 enzyme fills in the recessed 3' termini produced after digestion with EcoRI, leaving blunt ends. After treatment with Klenow fragment, the DNA was phenol/chloroform extracted, ethanol precipitated, and resuspended in T4 DNA ligase buffer and ligated at room temperature with T4 DNA ligase for 4 hours. The ligation mixture was transformed
10 into competent TB1 cells. Small scale preparation DNA was prepared from the transformants and screened for the loss of the EcoRI site. Plasmid pJO210, which has lost the EcoRI site, was isolated.

B. Construction of pJO215

The plasmid pJO215 is a derivative of the plasmid pJO210 (Figure
15 8). This plasmid was constructed by removing a single BamH-I site located in the promoter for the CKS gene using bridge mutagenesis (Mandecki, W., Proc. Nat. Acad. Sci. 837177,1986). Large scale plasmid DNA (pJO210) was isolated from TB1 cells as described in general methods. The DNA was digested with BamH-I to completion
20 and purified on a acrylamide gel. The purified pJO210/BamH-I fragment was then mixed with the following mutagenic oligonucleotide:

(2) 5' GGATAACAAT TGGGCATCCA GTAAGGAGGT 3'

which is complementary to one of the DNA strands of pJO210 in the

region of the plasmid spanning the BamH-I site. This oligonucleotide contains a single base change, underlined, which will destroy the BamH-I site when incorporated into plasmid pJO210. After mixing the mutagenic oligonucleotide with pJO210/BamH-I, the mixture was boiled
5 for 3 minutes, cooled to room temperature for 5 minutes and then transformed into competent TB1 cells. Small scale preparation DNA was prepared from the transformants and screened for the loss of the BamH-I site. Plasmid pJO215, which has lost the BamH-I site, was isolated.

10 C. Construction of pMC200

The plasmid pMC200 is a derivative of plasmid pMW507 described in Mandecki and Bolling, Gene 68: 101, 1988 (Figure 9). This plasmid was constructed by cloning a synthetic oligonucleotide containing a multi-cloning site into pMW507 using the FokI method of
15 gene synthesis described by Mandecki and Bolling, 1988. Large scale plasmid DNA (pMW507) was isolated from TB1 cells as described in general methods. The DNA was digested to completion with SmaI and then mixed with the following oligonucleotide:

5' AGGATGCGGA TCCCCGATCT CGACCCGTCG ACGAATTCGA
20 (3) GCTCGGTACC CGGGGATCCT CTAGACTGCA GGCATGCTAA
GTAAGTAGAT CGGGAATTCA CATCCG 3'

which contains FokI arms at the end and several restriction enzyme sites, as follows:

- 35 -

5' FokI arm-BglII sticky end-SalI/AccI/HincII-EcoRI-SstI-KpnI-SmaI/XmaI-

(4) BamHI-XbaI-PstI-SphI- stop codons -BglII sticky end-FokI arm 3'

After mixing the oligonucleotide with pMW507/SmaI, the mixture was boiled for 3 minutes, cooled to room temperature for 5 minutes, and then transformed into competent TB1 cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the multi-cloning site. Plasmid pMC200, which contains the multi-cloning site, was isolated. The DNA sequence of the multi-cloning site was confirmed.

10 D. Construction of pJO200

The plasmid pJO200 is a derivative of plasmid pJO215 (Figure 10). This plasmid was constructed by removing the multi-cloning site from pMC200 and cloning this site at the 3' end of the CKS gene in pJO215. Large scale plasmid DNA (pMC200 and pJO215) was isolated from TB1 cells as described in general methods. Plasmid pJO215 was digested to completion with BglII and then treated with calf intestinal alkaline phosphatase (CIAP) to prevent recircularization of the plasmid during the ligation reaction. Plasmid pMC200 was digested with FokI and then the pJO215/BglII/CIAP DNA and the pMC200/FokI DNA containing the multi-cloning site (106 base pairs) were purified on a polyacrylamide gel. Digestion of plasmid pMC200 with FokI releases the multi-cloning site DNA from the plasmid. This DNA contains BglII sticky ends which will readily ligate into pJO215 DNA digested with BglII.

- 36 - 2194135

These purified DNA fragments were mixed and ligated at 16°C with T4 DNA ligase overnight. The next day the ligation mixture was transformed into competent TB1 cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the multi-cloning site in the correct orientation at the BglII site. Plasmid pJO200, which contains the multi-cloning site in the correct orientation, was isolated. The DNA sequence of the multi-cloning site in pJO200 at the BglII site was confirmed.

EXAMPLE 7

10 Construction of lacZ-pp38(106-373aa) Expression Vector pMB38

The plasmid pMB38 is a derivative of the lacZ expression vector pROS (Figure 11). This plasmid was constructed by cloning a DNA fragment containing HCMV-pp38(106-373aa), obtained by PCR amplification of genomic HCMV DNA from the region of ppUL80a encoding amino acids 106-373aa of pp38 (nucleotides 316-1119), into the polylinker region pROS. The ppUL80a region encodes a phosphoprotein of 38 kD of HCMV. Plasmid pROS was digested with Sall and HindIII and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide 316 of ppUL80a containing a Sall site and an antisense primer starting at nucleotide 1119 of ppUL80a containing a HindIII site were synthesized and added to a PCR reaction mixture containing genomic HCMV DNA. After PCR amplification, the reaction mixture was digested with Sall and HindIII and the 804 base pair fragment

- 37 -

containing pp38(106-373aa) was purified on an agarose gel. This purified fragment was then ligated to purified pROS/Sall/HindIII overnight at 16°C. The next day the ligation mixture was transformed into competent DH5α cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the 804 base pair pp38 (106-373aa) fragment in pROS at the end of the lacZ gene. Plasmid pMB38, which contains the pp38(106-373aa) fragment, was isolated. The DNA sequence of pp38 (106-373aa) in pMB38 was confirmed and the pp38(106-373aa) coding region was in-frame with the lacZ coding sequence.

EXAMPLE 8

Construction of CKS-CMV Expression Vectors Based on pJO200

The CKS expression vector pJO200 is the building block for a series of five CKS-CMV gene fusion constructs. Two types of CKS-CMV gene fusion plasmids were constructed. One type of CKS-CMV gene fusion plasmid was constructed in which the CMV gene sequence is embedded within the CKS gene at nucleotide 638 of pJO200 (amino acid 171 of CKS) as shown below:

(5) CKS(1 -171 aa)-**CMV**-CKS(171 -260aa)

This method of CKS-CMV gene fusion construction is called epitope-embedding. Fusion proteins expressed in E.coli from this type of construct contain the epitopes of the antigen embedded entirely within the CKS amino acid sequence. Plasmid pCMV-1A was constructed in

Another type of CKS-CMV gene fusion plasmid was constructed in which the CMV gene DNA sequence is linked to the 3' end of the CKS gene at amino acid 248 as shown below:

Plasmids pCMV-3B, pCMV-4, pCMV-9, and pCMV-26 were constructed in this manner. Large scale plasmid DNA (pJO200) was isolated using general methods for the constructs described below.

pCMV-1A is a derivative of plasmid pJO200 (Figure 12). This plasmid was constructed by cloning a DNA fragment containing HCMV-A1C2F3, obtained by PCR amplification of A1C2F3 DNA contained in plasmid pMB34, into the *Stu*I site of pJO200. Large scale plasmid DNA (pMB34) was isolated by general methods. Plasmid pJO200 was digested with *Stu*I and *Bam*H-I and the vector backbone was purified on an agarose gel. This digest removes a portion of the 3' end of the CKS gene which will be restored in the ligation reaction. Into this vector backbone two PCR-derived DNA fragments will be cloned in a three-way ligation reaction. A1C2F3 will be cloned as a *Stu*I/*Mlu*I DNA fragment and the remaining 3' portion of the CKS gene will be cloned as a *Mlu*I/*Bam*H-I DNA fragment, restoring the complete CKS gene. A sense primer starting at nucleotide 1783 of ppUL32 containing a *Stu*I site and an antisense primer starting at nucleotide 3144 of ppUL32

containing an MluI site were synthesized and added to a PCR reaction mixture containing plasmid pMB34. After PCR amplification, the reaction mixture was digested with StuI and MluI, and the 204 base pair fragment containing A1C2F3 was purified on an agarose gel. A
5 sense primer starting at nucleotide 640 of pJO200 containing an MluI site and an antisense primer starting at nucleotide 905 of pJO200 containing a BamH-I site were synthesized and added to a PCR reaction mixture containing plasmid pJO200. After PCR amplification, the reaction mixture was digested with MluI and BamH-I, and the 266 base
10 pair fragment containing the 3' portion of the CKS gene was gel purified. These purified PCR-derived DNA fragments were then ligated to pJO200/StuI/BamH-I overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and screened for
15 the presence of A1C2F3 inserted at the StuI site of pJO200. Plasmid pCMV-1A, which contains A1C2F3 inserted at the StuI site, was isolated. The DNA sequence of A1C2F3 and the 3' end of the CKS gene was confirmed. The coding region of the CKS-A1C2F3-CKS construct in pCMV-1A contains a bridge of 2 amino acids (threonine
20 and arginine) contributed from the MluI site between A1C2F3 and the 3' end of CKS. In addition, amino acid 171 of CKS is duplicated in the construct as shown below:

(7) CKS(1-171aa)-A1C2-K-L-Q-E-F-F3-T-R-CKS(171-260aa)

- 40 - 2194135

B. Construction of pCMV-3B: CKS-H10 - The plasmid pCMV-3B is a derivative of plasmid pJO200 (Figures 13A and 13C). This plasmid was constructed by cloning a DNA fragment containing HCMV-H10 from plasmid pROSH10, described in Example 1 [see also: Ripalti et al. J. Virological Methods 46: 39, 1994], into pJO200. The H10 DNA sequence is derived from ppUL44 which encodes the phosphoprotein of 52 kD of HCMV. The H10 portion of ppUL44 in pROSH10 contains nucleotides 604-1299 (amino acids 202-434). H10 encodes the C-terminal half of phosphoprotein pp52. Plasmid pCMV-3B was constructed by cloning the H10 DNA fragment from pROSH10, obtained by PCR amplification of the H10 DNA sequence, into the SacI/BamH-I sites of pJO200. Plasmid pJO200 was digested with SacI and BamH-I and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide 604 of ppUL44 containing a SacI site and an antisense primer starting at nucleotide 1299 of ppUL44 containing a stop codon at the end of the H10 coding sequence and a BamH-I site were synthesized and added to a PCR reaction mixture containing plasmid pROSH10. After PCR amplification, the reaction mixture was digested with SacI and BamH-I, and the 696 base pair fragment containing H10 was purified on an agarose gel and then ligated to pJO200/SacI/BamH-I overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and

- 41 -

screened for the presence of the 696 base pair fragment in pJO200 at the SacI/BamH-I sites. Plasmid pCMV-3B, which contains the H10 fragment fused in-frame with the CKS gene, was isolated. The DNA sequence of H10 and the 3' end of the CKS gene was confirmed. This

5 CKS-CMV fusion construct is diagrammed below:

(8) CKS(1-248aa)-H10

C. Construction of pCMV-4: CKS-A1C2F3-H10 - The plasmid pCMV-4 is a derivative of pJO200 (Figures 13A and 13C). This plasmid was constructed by cloning PCR amplified DNA fragments, containing
10 HCMV-A1C2-K-L-Q-E-F-F3 (briefly: A1C2F3) and HCMV-H10, derived from pMB34 and pROSH10, respectively, into pJO200. Plasmid pJO200 was digested with SacI and BamH-I and the vector backbone was purified on an agarose gel. Into this vector backbone the two PCR-derived DNA fragments will be cloned in a three-way ligation reaction.
15 A1C2-bridge-F3 will be cloned as a SacI/PstI DNA fragment and H10 will be cloned as a PstI/BamH-I DNA fragment. A sense primer starting at nucleotide 1783 of ppUL32 containing a SacI site and an antisense primer starting at nucleotide 3144 of ppUL32 containing an PstI site were synthesized and added to a PCR reaction mixture containing
20 plasmid pMB34. After PCR amplification, the reaction mixture was digested with SacI and PstI, and the 204 base pair fragment containing A1C2F3 was purified on an agarose gel. A sense primer starting at nucleotide 604 of ppUL44 containing a PstI site and an antisense

primer starting at nucleotide 1299 of ppUL44 containing a stop codon at the end of the H10 coding sequence and a BamH-I site were synthesized and added to a PCR reaction mixture containing plasmid pROSH10. After PCR amplification, the reaction mixture was digested
5 with PstI and BamH-I, and the 696 base pair fragment containing H10 was purified on an agarose gel. These purified PCR-derived DNA fragments were then ligated to pJO200/SacI/BamH-I overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the
10 transformants and screened for the presence of A1C2F3 and H10 inserted at the SacI/BamH-I sites of pJO200. Plasmid pCMV-4, which contains A1C2F3 and H10 at the end of the CKS gene in pJO200, was isolated. The DNA sequence of A1C2-K-L-Q-E-F-F3 and H10 was confirmed. The coding region of the CKS-A1C2F3-H10 construct in
15 pCMV-4 contains a bridge of 2 amino acids contributed from the PstI site between A1C2F3 and H10. This CKS-CMV fusion construct is diagrammed below:

(9) CKS(1 -248aa)-A1C2-K-L-Q-E-F-F3-L-Q-H10

D. Construction of pCMV-9: CKS-pp65(297-510aa) - The plasmid
20 pCMV-9 is a derivative of pJO200 (Figures 13B and 13C). This plasmid was constructed by cloning a DNA fragment containing HCMV-pp65(297-510aa), obtained by PCR amplification of HCMV cDNA from the region of ppUL83 encoding amino acids 297-510 of pp65

- 43 -

(nucleotides 889-1530), into pJO200. Plasmid pJO200 was digested with SacI and BamH-I and the vector backbone was purified on an agarose gel. A two-stage nested PCR reaction was used to generate the HCMV-pp65(297-510aa) DNA fragment using HCMV cDNA as
5 template, as follows. For the outer nest PCR amplification reaction, a sense primer starting at nucleotide 807 of ppUL83 and an antisense primer starting at nucleotide 1572 of ppUL83 were synthesized and added to a PCR reaction mixture containing HCMV cDNA. After PCR amplification, the outer nest PCR reaction mixture was used as
10 template DNA for the inner nest PCR amplification reaction. For the inner nest PCR amplification reaction, a sense primer starting at nucleotide 889 of ppUL83 containing a SacI site and an antisense primer starting at nucleotide 1530 of ppUL83 containing a stop codon at the end of the pp65(297-510aa) coding sequence and a BamH-I site,
15 were synthesized and added to a PCR reaction mixture containing outer nest amplified DNA. After PCR amplification, the reaction mixture was digested with SacI and BamH-I, and the 642 base pair fragment containing pp65(297-510aa) was purified on an agarose gel. This purified DNA fragment was then ligated to pJO200/SacI/BamH-I
20 overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of pp65(297-510aa) inserted at the SacI/BamH-I sites of pJO200. Plasmid

- 44 -

2194135

pCMV-9, which contains pp65(297-510aa) at the end of the CKS gene in pJO200, was isolated. The DNA sequence of pp65(297-510aa) was confirmed. This CKS-CMV fusion construct is diagrammed below:

(10) CKS(1-248 aa)-pp65(297-510 aa)

- 5 E. Construction of pCMV-26: CKS-pp38(117-373aa) - The plasmid pCMV-26 is a derivative of pJO200 (Figures 13B and 13C). This plasmid was constructed by cloning a DNA fragment containing HCMV-pp38(117-373aa), obtained by PCR amplification of pp38 DNA from the region of ppUL80a encoding amino acids 117-373 of pp38 (nucleotides
- 10 349-1119) derived from pMB38, into pJO200. Large scale plasmid DNA (pMB34) was isolated as described in general methods. Plasmid pJO200 was digested with SacI and BamH-I and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide
- 15 at nucleotide 1119 of ppUL80a containing a BamH-I site were synthesized and added to a PCR reaction mixture containing pMB38 DNA. After PCR amplification, the reaction mixture was digested with SacI and BamH-I, and the 771 base pair fragment containing pp38(117-373aa) was purified on an agarose gel. This purified DNA fragment was
- 20 then ligated to pJO200/SacI/BamH-I overnight at 16°C. The next day the ligation mixture was transformed into the competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of pp38(117-373aa)

inserted at the SacI/BamH-I sites of pJO200. Plasmid pCMV-26, which contains pp38(117-373aa) at the end of the CKS gene in pJO200, was isolated. The DNA sequence of pp38(117-373aa) was confirmed. This CKS-CMV fusion construct is diagrammed below:

5 (11) CKS(1-248aa)-pp38(117-373aa)

EXAMPLE 9

Construction of CKS Epitope-Embedding Expression Vector pEE1

The CKS epitope-embedding expression vector pEE1 allows the embedding of recombinant proteins containing epitopes within the CKS
10 protein as diagrammed below:


(12) CKS(1-171aa)- Recombinant Protein -T-R-CKS(171-260aa)

This pEE1 vector was constructed in two steps starting with the CKS expression vector pJO200. In the first step a mutagenic oligonucleotide is cloned into a pair of adjacent MluI sites located near
15 the 5' end of the CKS gene in pJO200, removing both MluI sites and a Sall site. This modification to pJO200 allows the use of a unique MluI cloning site to be introduced further downstream in the CKS gene in the next step. In the second step a fragment of DNA from pCMV-1A is cloned into this modified pJO200 vector, thus permitting the
20 embedding of genes as StuI/MluI fragments into the CKS gene.

A. Construction of pJO200-ΔMluI - The plasmid pJO200-ΔMluI is a derivative of the CKS expression vector pJO200 (Figure 14A). This plasmid was constructed by removing a pair of adjacent MluI sites and

- 46 - 2194135


a Sall site located at nucleotides 161-174 (11-15 amino acids) in the pJO200 DNA sequence shown below using a mutagenic oligonucleotide:



Plasmid pJO200 was digested with MluI and then ethanol precipitated
 10 and resuspended in alkaline phosphatase buffer. Plasmid pJO200/MluI
 was then treated with calf intestinal alkaline phosphatase (CIAP) to
 remove the 5' phosphate groups to prevent self-ligation. After
 treatment with CIAP the DNA was extracted with phenol-chloroform,
 ethanol precipitated, resuspended in TE buffer, and then the vector
 15 backbone was purified on an agarose gel. The following mutagenic
 oligonucleotide was synthesized in preparation for ligation to
 pJO200/MluI/CIAP:



This oligonucleotide is self-complementary at its 3' end and will form
 20 the following double-stranded structure after a heat denaturation step
 followed by an annealing step:



This oligonucleotide contains MluI sticky-ends permitting ligation into
 25 MluI digested pJO200 DNA. The sequence of this oligonucleotide
 differs from the native pJO200 DNA sequence in that the "T" and "A"

- 47 -

residues underlined in the pJO200 sequence (13) above have been reversed in the mutagenic oligonucleotide as underlined. Thus, the mutagenic oligonucleotide when cloned into the MluI site of pJO200 will destroy both MluI sites and the Sall site. After oligonucleotide synthesis, the synthetic oligonucleotide was phosphorylated at its 5' end using polynucleotide kinase. After treatment with this enzyme, the phosphorylated oligonucleotide was heated to 65°C for 20 minutes to inactivate the kinase. After cooling to room temperature, the phosphorylated oligonucleotide was mixed with the pJO200/MluI/CIAP, heated at 65°C for 5 minutes and then cooled to room temperature gradually to permit annealing of the phosphorylated oligonucleotide to itself. Ligation buffer and T4 DNA ligase were then added to the mixture and incubated overnight at range of temperatures from 20°C to 4°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and screened for the loss of the MluI and Sall sites. Plasmid pJO200ΔMluI, which has lost these restriction enzyme sites, was isolated. The DNA sequence of the 5' end of the CKS gene was confirmed. In addition to removing the MluI and Sall sites, the mutagenic oligonucleotide changes the amino acids coded by nucleotides 166-171 from *Ser-Thr* to *Thr-Ser* as shown below:

(13') ¹⁵¹CCC GCG CGC TAC GCG ACG TCG CGT CTG CCC¹⁸⁰
 pro Ala Arg Tyr Ala Thr Ser Arg Leu Pro
 pJO200ΔMluI DNA sequence nucleotides 151-180 taken in direction 5→3'

B. Construction of pEE1 - The plasmid pEE1 is a derivative of the plasmid pJO200ΔMlul (Figure 15). This plasmid was constructed by cloning a Stul/BamH-I fragment from pCMV-1A, which contains HCMV-A1C2F3 embedded within the CKS gene, into the Stul/BamH-I sites of

5 pJO200ΔMlul. By substituting the Stul/BamH-I DNA fragment within the CKS coding region present in pJO200ΔMlul with the Stul/BamH-I fragment from pCMV-1A, the resulting plasmid pEE1 contains HCMV-A1C2F3 embedded within the CKS gene. Plasmid pEE1 differs from plasmid pCMV-1A in that pEE1 does not contain the upstream Mlul

10 sites present in the 5' end of the CKS gene. Hence, digestion of pEE1 with Stul and Mlul will release the HCMV-A1C2F3 DNA fragment and provide a vector backbone, after purification on an agarose gel, capable of accepting other genes for embedding into the CKS gene as blunt/Mlul compatible sticky-end DNA fragments. Large scale plasmid

15 DNAs (pJO200ΔMlul and pCMV-1A) were isolated as described in general methods. Plasmid pCMV-1A was digested with Stul and BamH-I and the 461 base pair fragment, containing A1C2F3 and the 3' end of the CKS gene, was purified on an agarose gel. Plasmid pJO200ΔMlul was digested with Stul and BamH-I and the vector backbone was

20 purified on an agarose gel. These purified DNA fragments were then mixed together and ligated overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and screened for

- 49 -

the presence of the 461 base pair A1C2F3 DNA fragment in pJO200ΔMlul. Plasmid pEE1, which contains the A1C2F3 DNA fragment and no Mlul sites in the 5' end of the CKS gene, was isolated. The DNA sequence of the 3' end of the CKS gene and the A1C2F3 fragment was confirmed. Digestion of plasmid pEE1 with Stul and BamH-I followed by purification of the vector backbone on an agarose gel removes the A1C2F3 DNA fragment completely in preparation for ligation with other DNA fragments. This purified vector backbone can accept DNA fragments for embedding into the CKS gene in the correct reading frame in the following format:

(16) 5' X- **Gene of interest** -Y 3'

where X is a blunt end and Y is an Mlul compatible sticky-end, for example Mlul or BssHII.

EXAMPLE 10

15 Construction of CKS-epitope-embedding Expression vectors based on pEE1

The CKS expression vector pEE1 is the building block for a series of three CKS-CMV-CKS gene fusion constructs. For each construct plasmid pEE1 was digested with Stul and Mlul and the vector backbone was purified. This backbone is ready to accept CMV gene fragments generated by PCR which have a Stul site at their 5' end and a Mlul site at their 3' end. After digestion with Stul and Mlul, the PCR fragments are cloned in-frame into the pEE1/Stul/Mlul backbone. The CKS-CMV-

2194135

- 50 -

CKS fusion proteins expressed from these vectors are diagrammed below where T and R are the threonine and arginine residues, respectively, encoded by the synthetic MluI site introduced into the vector:

5 (17) CKS(1 -171 aa)-CMV-T-R-CKS(171 -260)

A. Construction of pCMV-27: CKS-A1C2F3-H10-CKS - The plasmid pCMV-27 is a derivative of pEE1 (Figures 16A and 16B). This plasmid was constructed by cloning a PCR amplified DNA fragment, containing HCMV-A1C2F3-H10 derived from pCMV-4 (therefore containing the
10 bridge KLQEF between A1C2 and F3), into pEE1. Large scale plasmid DNA (pEE1 and pCMV-4) was isolated operating as described in general methods. Plasmid pEE1 was digested with StuI and MluI and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide 1783 of ppUL32 containing a StuI site and an antisense
15 primer starting at nucleotide 1299 of ppUL44 containing an MluI site were synthesized and added to a PCR reaction mixture containing plasmid pCMV-4. After PCR amplification, the reaction mixture was digested with StuI and MluI, and the 906 base pair fragment containing A1C2F3-H10 was purified on an agarose gel. This purified DNA
20 fragment was ligated to pEE1/StuI/MluI overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the A1C2F3-H10 DNA

- 51 -

fragment inserted at the *Stu*I/*Mlu*I sites of pEE1. Plasmid pCMV-27, which contains A1C2F3-H10 embedded at the *Stu*I/*Mlu*I sites of pEE1, was isolated. The DNA sequence of A1C2F3-H10 and the adjacent DNA sequence of CKS was confirmed. This CKS-CMV-CKS fusion construct is diagrammed below, where L, Q and T, R are the leucine, glutamine and threonine, arginine residues, respectively, encoded by the synthetic *Pst*I and *Mlu*I sites introduced into the vector:

(18) CKS(1-171 aa)-A1C2-K-L-Q-E-F-F3-L-Q-H10-T-R-CKS(171-260 aa)

B. Construction of pCMV-28: CKS-pp65(297-510aa)-CKS - The

plasmid pCMV-28 is a derivative of pEE1 (Figures 16A and 16B). This plasmid was constructed by cloning a PCR amplified DNA fragment, containing HCMV-pp65(297-510 aa) derived from pCMV-9, into pEE1. Large scale plasmid DNAs (pEE1 and pCMV-9) were isolated by general methods. Plasmid pEE1 was digested with *Stu*I and *Mlu*I and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide 889 of ppUL83 containing a *Stu*I site and an antisense primer starting at nucleotide 1530 of ppUL83 containing an *Mlu*I site were synthesized and added to a PCR reaction mixture containing plasmid pCMV-9. After PCR amplification, the reaction mixture was digested with *Stu*I and *Mlu*I, and the 642 base pair fragment containing pp65(297-510aa) was purified on an agarose gel. This purified DNA fragment was ligated to pEE1/*Stu*I/*Mlu*I overnight at 16°C. The next day the ligation mixture was transformed into competent XL-1 Blue

cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the pp65(297-510aa) DNA fragment inserted at the *Stu*I/*Mlu*I sites of pEE1. Plasmid pCMV-28, which contains pp65(297-510aa) embedded at the *Stu*I/*Mlu*I sites of pEE1, was isolated. The DNA sequence of pp65(297-510aa) and the adjacent DNA sequence of CKS was confirmed. This CKS-CMV-CKS fusion construct is diagrammed below where T and R are the threonine and arginine residues, respectively, encoded by the synthetic *Mlu*I sites introduced into the vector:

10 (19) CKS(1-171aa)-pp65(297-510aa)-T-R-CKS(171-260aa)

C. Construction of pCMV-29: CKS-pp38(117-373aa)-CKS - The plasmid pCMV-29 is a derivative of pEE1 (Figures 16A and 16B). This plasmid was constructed by cloning a PCR amplified DNA fragment, containing HCMV-pp38(117-373aa) derived from pCMV-26, into pEE1.

15 Large scale plasmid DNA (pEE1 and pCMV-26) was isolated as described in general methods. Plasmid pEE1 was digested with *Stu*I and *Mlu*I and the vector backbone was purified on an agarose gel. A sense primer starting at nucleotide 349 of ppUL80a containing a *Stu*I site and an antisense primer starting at nucleotide 1119 of ppUL80a containing

20 an *Mlu*I site were synthesized and added to a PCR reaction mixture containing plasmid pCMV-26. After PCR amplification, the reaction mixture was digested with *Stu*I and *Mlu*I, and the 771 base pair fragment containing pp38(117-373aa) was purified on an agarose gel.

- 53 -

This purified DNA fragment was ligated to pEE1/Stul/Mlul overnight at 16°C. The next day the ligation mixture was transformed into competent XL-I Blue cells. Small scale preparation DNA was prepared from the transformants and screened for the presence of the pp38(117-373aa) DNA fragment inserted at the Stul/Mlul sites of pEE1. Plasmid pCMV-29, which contains pp38(117-373aa) embedded at the Stul/Mlul sites of pEE1, was isolated. The DNA sequence of pp38(117-373aa) and the adjacent DNA sequence of CKS was confirmed. This CKS-CMV-CKS fusion construct is diagrammed below where T and R are the threonine and arginine residues, respectively, encoded by the synthetic Mlul sites introduced into the vector:

(20) CKS(1-171 aa)-pp38(117-373aa)-T-R-CKS(171-260 aa)

EXAMPLE 11

A. Expression of HCMV genes.

Bacterial clones expressing HCMV fusion proteins of Example 10, the control bacterial strain expressing unfused CKS, and bacterial clones expressing HCMV fusion proteins of Example 8, were grown in "SUPERBROTH II" media containing 100 µg/ml ampicillin to log phase and the synthesis of the CKS-HCMV fusion proteins was induced by the addition of IPTG as previously described [22]. After 4 hours post-induction, the cells were harvested and the cell pellets were stored at -80°C until protein purification. The relevant data of all the expressed proteins are given in Table 3.

2194135

B. Purification of unfused CKS and recombinant CKS-HCMV fusion proteins.

Insoluble CKS-HCMV fusion proteins (rpCMV-1A, rpCMV-3B, rpCMV-4, rpCMV-9, rpCMV-27, rpCMV-28, rpCMV-29) were purified after lysis by a combination of detergent washes that were followed by solubilization in 8M urea [22]. After solubilization in 8M urea, the fusion proteins were purified by Q-Sepharose* chromatography (Pharmacia Biotech, Piscataway, New Jersey). Fusion protein 4 (rpCMV-4) and 9 (rpCMV-9) were subjected to additional purification by Sephacryl S-200 * chromatography (Pharmacia Biotech, Piscataway, New Jersey). Soluble CKS-HCMV fusion protein 26 (rpCMV-26) was purified after cell lysis on preparative SDS-PAGE using a BIO-Rad Prep Cell* (Bio-Rad, Richmond, CA). Soluble CKS protein was purified after cell lysis by ammonium sulfate precipitation followed by DEAE chromatography.

15

EXAMPLE 12

Automated HCMV IgM Immunoassay

The use of recombinant polypeptides, which contain epitopes within the ppUL32, ppUL44, ppUL83, and ppULB0a regions of the HCMV genome, provide immunological assays that have increased sensitivity and that may be more specific than HCMV IgM immunoassays using epitopes from the virus.

In the automated HCMV IgM immunoassay, three E.Coli expressed recombinant proteins, CKS-A1C2F3-H10-CKS (rpCMV-27), CKS-pp65-

* Trade-mark

- 55 -

CKS (rpCMV-28), and CKS-pp38-CKS (rpCMV-29), representing four distinct regions of the HCMV genome were used. Each of these recombinant polypeptides were prepared according to Example 10 (A,B and C) and 11. In the automated immunoassay, each of these three
5 recombinant antigens were coated separately onto polystyrene microparticles, one antigen per microparticle. The automated immunoassay can then be run in either a sequential mode of operation or in a combination mode of operation. In a modification of the automated immunoassay, all of these three recombinant antigens may
10 be coated together onto the same polystyrene microparticle, three antigens per microparticle. The modified automated immunoassay can then be run in a combination mode of operation.

The polystyrene microparticles (2.65 μm size) in distilled water were introduced into a solution containing 50 mM Tris, 137 mM sodium
15 chloride, 0.1% sodium azide, pH 7.5. The polystyrene microparticles were then dispensed into three separate bottles designated rpCMV-27, rpCMV-28, and rpCMV-29. Recombinant rpCMV-27 in 50 mM Tris, pH 8.5 was added to the bottle containing rpCMV-27 to give a final concentration of 25-100 $\mu\text{g/ml}$. Recombinant rpCMV-28 in 50 mM
20 Tris, pH 8.5 was added to the bottle containing rpCMV-28 to give a final concentration of 25-100 $\mu\text{g/ml}$. Recombinant rpCMV-29 in 50 mM Tris, pH 8.5 was added to the bottle containing rpCMV-29 to give a final concentration of 25-100 $\mu\text{g/ml}$. The final concentration of the

polystyrene microparticles in each bottle was 1-3% solids. The bottles were rotated end over end (30-50 rpm) for two hours at room temperature (19-22°C) and were then centrifuged at 10,000-15,000 xg for 10 minutes at room temperature. The microparticles, which were in
5 the form of pellets, were then resuspended in a mixture containing 90 mM Tris, 135 mM sodium chloride, 100 mM disodium EDTA, 6% sucrose, 0.18% "TWEEN-20", 50% fetal calf serum (CMV antibody free), 35 µg/ml CKS protein, pH 7.5 to give a final concentration of 0.1-0.25% solids. The microparticles were then loaded into plastic
10 bottles.

The polystyrene microparticles coated with rpCMV-27 rpCMV-28, and rpCMV-29 were used in an antibody capture format in an automated immunoassay performed on the ABBOTT "IMx"® instrument (Abbott Laboratories, Abbott Park, Illinois). The aforementioned coated
15 microparticles can also be used in an antibody capture format in an automated immunoassay performed on the Abbott "AxSYM"® instrument (Abbott Laboratories, Abbott Park, Illinois).

These systems employ pipetting devices that dispense serum samples and reagents automatically. These instruments employ an optical reader
20 that can measure the fluorescence emission reflectance at 448 nm from the sample matrix. In the sequential mode of operation, the serum sample is incubated separately with each of the three coated microparticles. In the combination mode of operation, an equal volume

- 57 -

of each of the three coated microparticles are mixed together prior to incubation with the serum sample. In either mode the appropriate stock microparticles are dispensed into plastic bottles and then are loaded onto the instrument.

- 5 The preferred conjugate is goat anti-human IgM alkaline phosphatase conjugate. The conjugate is titered to determine a working concentration of 1-5 µg/ml. The conjugate diluent includes 90 mM Tris, 135 mM sodium chloride, 5% fetal calf serum, 0.1% sodium azide, pH 7.4. The conjugate is sterile filtered, filled in plastic bottles, and loaded
10 onto the instrument. The preferred substrate for the conjugate is 4-methylumbelliferyl phosphate.

Anti-HCMV positive index calibrator is prepared from plasma units positive for antibodies to HCMV. The pool of units includes plasma with antibodies reactive to rpCMV-27, rpCMV-28, and rpCMV-29. The
15 pooled units are recalcified, aliquoted, and stored at -20°C or at 2-8°C. For each lot of positive Index calibrator, the stock solution is diluted with negative control containing 0.1% sodium azide as a preservative. The final material is sterile filtered and filled in plastic bottles.

Anti-HCMV negative control is prepared from recalcified human plasma
20 negative for antibodies to rpCMV-27, rpCMV-28, and rpCMV-29 proteins of HCMV. The plasma is also negative for antibodies to human immunodeficiency virus (HIV) and negative for hepatitis B surface antigen (HBsAg). The units are pooled, and 0.1% sodium azide is added

- 58 - 2194135

as a preservative. The final material is sterile filtered and filled in plastic bottles.

The serum samples to be tested are dispensed into the instrument's reaction vessels. The appropriate code to run the automated CMV IgM immunoassay is entered and then the assay is begun. The instrument dilutes the serum sample in line diluent (0.1 M sodium phosphate, 0.1% sodium azide, pH 7.5 or 0.3 M sodium chloride, 0.1 M Tris, 0.1% sodium azide, pH 7.5) and then adds the microparticles to the diluted serum sample. After incubation for 5-10 minutes, the mixture is then transferred to the reaction vessel matrix. The matrix is washed with line diluent and then the conjugate is added to the reaction vessel matrix. After incubation for 8-10 minutes, the matrix is washed with line diluent and then the substrate for the conjugate is added. The fluorescence emission reflectance at 448 nm from the sample matrix is read immediately and the instrument reports both a rate count value and an index value for each sample. The index value is equal to the rate count value for the sample divided by the rate count value for the positive index calibrator.

In order to maintain acceptable assay specificity, the cut-off value for the assay should be at least 3-4 standard deviations above the mean of the index values for the negative samples. The cut-off value for the automated HCMV IgM immunoassay was set at an index value of 0.6. Any samples with an index value greater than or equal to 0.5 had to be

- 59 -

treated with rheumatoid factor neutralization reagent and then rerun again on the instrument. Samples with an index value equal to or greater than 0.6 were considered positive for HCMV IgM antibody. Samples with an index value of 0.500-0.599 were considered
5 equivocal, and samples with an index value less than 0.500 were considered negative for HCMV IgM antibody. The characteristics of the fusion proteins used in the assay are referred to in Table 3.

Automated HCMV IgM Immunoassay Performance

The performance of the automated HCMV IgM immunoassay was
10 evaluated by testing seven HCMV IgM-negative serum samples and eleven HCMV IgM-positive serum samples. The titer of HCMV IgM antibody for these samples was initially determined using the HCMV IgM microtiter recombinant EIA described in Example 13. All HCMV IgM positive samples were confirmed to be positive by Western Blot using
15 purified viral antigen (see Table 5). All HCMV IgM positive samples were treated with rheumatoid factor neutralization reagent prior to running them in the immunoassays described below in (a) and (b).

(a) - Sequential Mode of Operation

In the sequential mode of operation of the automated
20 immunoassay, each serum/plasma sample is tested separately using the rpCMV-27 coated microparticles, rpCMV-28 coated microparticles, and rpCMV-29 coated microparticles.

A sample is considered positive for HCMV IgM antibody in the

2194135

sequential mode of operation when the index value for the sample is equal to or greater than 0.6 with any one of rpCMV-27 coated microparticles or rpCMV-28 coated microparticles or rpCMV-29 coated microparticles.

- 5 A sample is considered equivocal for HCMV IgM antibody in the sequential mode of operation when the index value for the sample is 0.500-0.599 with rpCMV-27 coated microparticles or rpCMV-28 coated microparticles or rpCMV-29 coated microparticles.

- A sample is considered negative for HCMV IgM antibody in the
10 sequential mode of operation when the index value for the sample is less than 0.500 with rpCMV-27, rpCMV-28, and rpCMV-29 coated microparticles. As shown in Tables 4 and 5, all negative samples were determined to be negative for HCMV IgM antibody and all positive samples were determined to be positive for HCMV IgM antibody in the
15 sequential mode of operation of the HCMV IgM automated immunoassay.

(b) - Combination Mode of Operation

- In the combination mode of operation of the automated immunoassay, each serum/plasma sample is tested using an equal
20 volume mixture of rpCMV-27, rpCMV-28, and rpCMV-29 coated microparticles (pooled microparticles). A sample is considered positive for HCMV IgM antibody in the combination mode of operation when the index value for the sample is equal to or greater than 0.6 with the

- 61 -

2194135

pooled microparticles. A sample is considered equivocal for HCMV IgM antibody in the combination mode of operation when the index value for the sample is 0.500-0.599 with pooled microparticles. A sample is considered negative for HCMV IgM antibody in the combination mode of operation when the index value for the sample is less than 0.500 with pooled microparticles. As shown in Tables 4 and 5, all negative samples were determined to be negative for HCMV IgM antibody and all positive samples were determined to be positive for HCMV IgM antibody in the combination mode of operation of the HCMV IgM automated immunoassay.

EXAMPLE 13

Manual HCMV IgG/IgM Immunoassay

1. HCMV serology.

1.1- Conventional EIA (conv-EIA) - The evaluation of anti-HCMV IgG was carried out with a commercial kit ("ENZYGNOST"* anti-HCMV/IgG EIA alpha method, Behring AG, Marburg, Germany). Plates were read on a microEIA automatic reader (Behring AG- Marburg, Germany). The evaluation of anti-HCMV IgM was performed using the "ENZYGNOST" Anti-HCMV/IgM kit (Behring AG, Germany). Both kits were used and the results interpreted as suggested by the manufacturers.

1.2 - Western blotting (WB). Protein extracts from purified viral particles (Towne strain) were run in 9% acrylamide gel and electrophoretically separated polypeptides were then transferred to

* Trade-mark

B

- 62 - 2194135

nitrocellulose paper. Infection of cells, virus purification, protein extraction, blotting and immune reaction with sera were done as previously described in detail [16].

1.3 - Recombinant EIA (rec-EIA). EIA was performed following the
5 procedure and using the reagents of the commercial kit Enrygnost anti-HCMV/IgG and IgM (Behring, Marburg, Germany). Briefly 0.05 µg of purified recombinant protein per well in bicarbonate buffer (pH 9.6) was used to coat EIA plastic plates (A/S NUNC, Roskilde, Denmark). After an overnight incubation at 4°C, plates were rinsed three times with
10 PBS-"TWEEN 20" (0.05%) and incubated with BSA (1%) in bicarbonate buffer (pH 9.6). After a 1 hour incubation period at room temperature, wells were rinsed three times with PBS-"TWEEN 20" (0.05%) and incubated with human serum samples at a dilution of 1:100 and 1:40 in PBS for IgG and IgM respectively (50 µl final volume). The incubation
15 was carried out at 37°C for 2 hours. After three washes with PBS-"TWEEN 20", peroxidase-conjugated goat anti-human IgG or IgM antibodies were deposited into wells, and plates were incubated at 37°C for 1.5 hours. After three washes with PBS-"TWEEN 20", immunoreactions were evidenced by addition of chromogen - TMB
20 (Tetramethylene benzidine dihydrochloride). The reaction was stopped after one hour by the addition of sulfuric acid (0.5 N), and the results read on a "MICROEIA" automatic reader. For each sample, the immunoreaction level was considered to be the difference between the

- 63 -

measured absorbances, (A rec.antigen- A control antigen), obtained. The cut-off value for each individual fusion protein was determined by testing 210 sera that did not contain IgM to HCMV (determined using both conv-EIA and WB). Cut off values for each fusion protein are reported in Table 3. For the determination of IgM antibodies, "The RF absorbent" (Behring, Marburg, Germany) was used to remove Rheumatoid factor from serum samples. To perform linear regression analysis and to standardize the test run, every plate included three serum calibrators. Reproducibility was controlled by including in each EIA plate six standard sera whose reactivity had previously been assayed in four independent experiments. Test runs were considered acceptable when the value of the internal control sera were within the interval of two standard deviations from the mean value previously established.

2. Human Serum samples.

Many groups of human serum samples (sera) were used in this work. The first group of samples, which was used for the determination of the cut-off values, consisted of 210 sera from blood donors (150) and healthy adults (60). While sixty-eight sera were IgG/IgM-, 142 were IgG + /IgM- as judged by both conv-EIA and Western blotting. The second group of sera consisted of 150 HCMV-positive samples from immunocompetent subjects (50 with a high IgM titre to HCMV as detected by conv-EIA, 50 with a medium IgM level and 50 with a low

- 64 - 2194135

level of HCMV-specific IgM). The presence of HCMV-specific IgM in this group of sera was determined by conv-EIA and confirmed by Western blotting. A third group of sera consisted of 51 sera from pregnant women (18 were from HCMV-uninfected women, 26 from HCMV infected women who did not transmit the infection and 7 from HCMV-infected pregnant women who transmitted the infection). All these sera were obtained between 22 and 24 weeks of gestation. A group of 35 sera from 35 newborns was also tested: 6 serum samples were obtained during the first week of life from 6 congenitally infected newborns, 19 sera were from newborns (1-8 months of life) in which the virus was repeatedly isolated from urine. Twelve newborns had different symptoms (maculopapular rash, encephalopathy, growth delay), the others were asymptomatic shedders. Finally ten sera were from HCMV-uninfected newborns during the first 8 months of life. Another group of samples consisted of 104 sera from 23 transplanted patients (20 heart and 3 kidney transplant recipients) who underwent a HCMV infection during the first 3 months after transplantation. Eight patients had a primary infection, while 15 underwent a viral reactivation. The last group consisted of 200 serum samples randomly selected among sera from renal transplant recipients (150) and pregnant women (150), who came to diagnostic laboratory for HCMV monitoring after transplantation and during pregnancy.

3. Determination of a HCMV infection in a pregnant woman and

- 65 -

newborns.

HCMV infection in pregnant women was determined by one or more of the following parameters: virus isolation from urine, saliva, blood, seroconversion for anti-HCMV antibodies. A congenital HCMV
5 infection in a newborn was determined by HCMV isolation from urine during the first week of life. A HCMV infection in newborns was determined by virus isolation from urine.

4. Determination of a HCMV infection in transplanted patients.

HCMV infection in transplanted patients was determined by the
10 presence of pp65-positive PMNL (antigenemia) in peripheral blood as determined by indirect immunofluorescence and the presence of the HCMV genome in the same cells as determined by PCR.

5. Virological detection of HCMV.

HCMV isolation. The "shell vial" procedure [23] was used for HCMV
15 isolation from urine and saliva. The inoculated cells were fixed 24-48 hours after inoculation and stained in an indirect immunofluorescence (IIF) assay using a monoclonal antibody reacting with HCMV-IE1/IE2 gene product (EI3 from Bioline, Paris, France).

Antigenemia. The presence of HCMV-pp65 (ppUL83) in PMNL was
20 determined as originally described by Schirm et al. [24] and modified by Revello et al [25] using a HCMV-pp65 specific pool of two monoclonal antibodies (Biotest, Frankfurt, Germany) in IIF tests.

Determination of the presence of HCMV genome by PCR. Aliquots of

5X10⁵ PMNL were resuspended in 100 µl of PCR buffer (KCl 50 mM, Tris-HCl 10 mM, pH 8.3, MgCl₂ 2 mM, gelatin 0.01%), with non-ionic detergents and Proteinase K. Samples were incubated at 60°C for one hour and then at 95°C for 20 minutes to inactivate the proteinase.

5 Thirty microliters of each sample were added to 20 µl of reaction buffer containing 50 mM KCl, 10 mM Tris-HCl pH 8.3, 2 mM MgCl₂, 0.2 mM each of the four deoxynucleotide triphosphates, 50 picomoles of each primer and 1.25 units of native Taq DNA-polymerase (Perkin Elmer Cetus, Norwalk, CT, USA). HCMV specific primers were synthesized by

10 Genset (Paris, France). Sequences are from the 4th exon of HCMV immediate early gene (EcoRI J fragment of the AD169 strain) and correspond to nucleotides 1767 to 1786 and from nucleotide 1894 to 1913. Using these primers, a fragment of 147 base pairs was amplified. A third oligonucleotide consisting of nucleotides 1807-1847

15 was complementary to the antisense DNA strand in the region between the binding sites of the other oligonucleotides and was used for hybridization [26]. The amplification reaction was carried out in a DNA thermal cycler (Perkin-Elmer/Cetus, Norwalk, CT, USA) for 35 cycles. Each sample was initially subjected to analysis with oligonucleotide

20 primers GH26 and GH27 (Perkin Elmer Cetus) (20 pmol each), with flank a conserved region of the HLA-DQa locus, to determine the capacity of DNA to be amplified. Fifteen microliters of the reaction mixture were subjected to electrophoresis on 1.2% "NUSIEVE" GTG-

- 67 -

agarose and 0.4% Seakem LE-agarose (FMC Bioproducts, Rockland, ME, USA) gel. The DNA was visualized by UV fluorescence after ethidium bromide staining. Nucleic acids were denatured and neutralized using standard procedures and then transferred to nylon
5 membranes (Mybond N, Amersham, UK) using a trans-blott apparatus (Hoefer Scientific Instruments, San Francisco, California). Hybridization with a terminally labeled oligonucleotide was carried out by a commercial kit (3'-oligolabelling and detection system, ECL, Amersham, UK) as suggested by the manufacturers. The sensitivity and specificity
10 of the PCR reaction were determined as described in a previous work [27].

6. Results.

As summarized in Table 3, five different fusion proteins have been comparatively tested.

15 The first fusion protein (rpCMV-1A) carries the two ppUL32 major antigenic regions fused together. The second fusion protein (rpCMV-3B) carries the COOH half of ppUL44. The third fusion protein (rpCMV-4) carries three regions (two regions of ppUL32 and one of ppUL44) fused together. Fusion proteins rpCMV-9 and rpCMV-26 carry
20 large fragments of ppUL83 and ppUL80a respectively. The cut-off values for the manual assay, for each recombinant protein, are reported in Table 3 and correspond to Optical Density values higher than the maximum value obtained from the 210 serum samples (they

- 68 - 2194135

approximately correspond to the mean values plus 4-5 standard deviations). In order to evaluate the sensitivity of the rec-EIA, 150 serum samples were selected from several hundred sera because they definitely contained IgM-to HCMV as they gave an IgM-positive result with both conv-EIA and WB. Sera were divided into three groups on the basis of the conv-EIA titre for HCMV-specific IgM. In particular, sera with OD values between 210 and 400 were considered to have a low IgM titre, those with OD ($\times 10^3$) from 401 to 800 were considered to have a medium titre, and those with OD ($\times 10^3$) higher than 800 were considered to have a high IgM titre. As summarized in Table 6, all the sera with medium and high IgM titres to HCMV reacted with one or more fusion proteins. Among the sera with a low IgM titre only two were not detected by any fusion protein. The % of reactivity is therefore higher than 98%. The highest reactivity obtained was against fusion proteins rpCMV-3B and rpCMV-4 and the combination of proteins rpCMV-4, rpCMV-9 and rpCMV-26 gave a reactivity of 98%. Only one serum did not react with this combination and was found to be reactive with fusion protein rpCMV-3B alone.

Having determined the high sensitivity of the recombinant EIA, several sera obtained from different groups of subjects were tested. Table 7 compares the results obtained by recombinant EIA with fifty-one sera obtained from pregnant women with the results obtained by a conv-EIA with the same sera. Among eighteen HCMV uninfected

- 69 -

pregnant women, none was found IgM-positive by either conventional or recombinant EIA. On the contrary, among twenty-six HCMV-infected pregnant women who did not transmit the infection to their offspring, recombinant EIA detected IgM in twenty cases, while conv-EIA
5 detected IgM in only twelve cases. Among a group of eight HCMV-infected pregnant women who transmitted the infection, all eight were found IgM-positive by recombinant and six by conv-EIA. The highest reactivity was always obtained with the triple antigen fusion protein (fusion protein rpCMV-4). The combination of proteins rpCMV-4,
10 rpCMV-9 and rpCMV-26 gave the maximum reactivity. As shown in Table 7, a significant difference in IgM titre to fusion protein rpCMV-4 was observed between HCMV-infected pregnant women who did not transmit the infection (lower titres) and those who transmitted the infection (higher titres). Although less significant, also the titre against
15 rpCMV-26 was different in the two groups (table 7). The differences in titres to fusion proteins rpCMV-3B and rpCMV-9 were not significant.

A group of thirty-five sera from newborns was also tested (Table 8). Among ten HCMV-uninfected newborns, none was found IgM-positive by either conventional or recombinant EIA. On the contrary,
20 among six sera obtained from six congenitally infected newborns during the first week of life, two gave a positive reaction by rec-EIA and none with the conventional test. The maximum IgM reactivity observed was specific for fusion proteins rpCMV-9 and rpCMV-26. In another group

of nineteen sera from newborns persistently excreting HCMV in urine during the first year of life (these newborns were not checked for HCMV at birth) rec-EIA detected IgM in ten cases while conv-EIA detected IgM in only two cases. The maximum IgM reactivity obtained
5 was against fusion proteins rpCMV-4 and rpCMV-26 and the combination of rpCMV-4, rpCMV-9 and rpCMV-26 gave the maximum reactivity.

Representative examples of the virological and serological monitoring of HCMV infection in transplant recipients are shown in
10 Figures 17A and 17B. In Figure 17A three primary and in Figure 17B three cases of secondary HCMV infections are shown. A1 and A2 are follow up of kidney, A3, B1,2 and B3 of heart transplant recipients. S+T means Symptoms and gancyclovir treatment; PCR+ means positive detection of HCMV genome in polymorphonuclear cells.

15 From the data presented, fusion protein rpCMV-4 was the one giving the highest reactivity with IgM antibodies presented in sera from healthy adults, pregnant women, infected newborns, and transplant recipients. However, fusion protein rpCMV-4 alone could not represent the entire complex of viral antigen reacting with anti-HCMV IgM. On
20 the contrary, the combination of fusion proteins rpCMV-4, rpCMV-9, rpCMV-26 can efficiently replace the virus in IgM detection. In fact, as shown in Tables 9 and 10, 99.6% of IgM-positive sera gave a positive IgM reaction with one or more of these fusion proteins. When the rec-

- 71 -

EIA with the three clones was compared with conv-EIA and WB, rec-EIA and WB gave the same results, and both procedures were more sensitive than conv-EIA. This means that the sensitivity and specificity of the rec-EIA according to the present invention is very similar to that of WB, which is a very sensitive and specific procedure for HCMV-IgM detection [30,31].

List of the References

1. Landini M. P. New approaches and perspectives in cytomegalovirus diagnosis. Prog. Med Microbiol 1993; 4, 157-177.
- 10 2. Nielsen St, Sorensen I, Andersen ELK. Kinetics of specific immunoglobulins M,E,A and G in congenital, primary, and secondary cytomegalovirus infection studied by antibody-capture enzyme-linked immunosorbent assay. J clin Microbiol 1988; 26: 654-661.
3. Basson J, Tardy JC, Aymard M. Pattern of anti-cytomegalovirus IgM antibodies determined by immunoblotting. A study of kidney graft recipients developing a primary or recurrent CMV infection. Arch Virol 15 1989; 108: 259-270.
4. Drew WL. Nonpulmonary manifestations of cytomegalovirus infection in immuno-compromized patients. Clin Microbiol Rev 1992; 20 5: 204-210.
5. Lazzarotto T, Dalla Casa B, Campisi B, Landini NIP. Enzyme-linked immunoadsorbent assay for the detection of cytomegalovirus-IgM: Comparison between eight commercial kits, immunofluorescence and

- 72 - 2194135

- immunoblotting. J Clin Lab Anal 1992; 6: 216-218.
6. Jahn O, Scholl BC, Traupe B, Fleckenstein B. The two major phosphoproteins (pp65 and pp150) of human cytomegalovirus and their antigenic properties. J Gen Virol 1987; 68, 1327-1337.
- 5 7. Landini MP., Lazzarotto T, Ripalti A, Guan MX, La Placa M. Antibody response to recombinant Lambda gt11 fusion proteins in Cytomegalovirus infection. J Clin Microbiol 1989; 27: 2324-2327.
8. Landini MP, Guan MX, Jahn G, Lindenmeier W, Mach M, Ripalti A, Necker A, Lazzarotto T, Plachter B. Large scale screening of human
10 sera with Cytomegalovirus recombinant antigens. J Clin Microbiology 1990; 28:1375-1379.
9. Landini MP, Ripalti A, Sra K , Pouletty P. Human cytomegalovirus structural proteins: immune reaction against pp150 synthetic peptides. J Clin Microbiol 1991; 29: 1868-1872.
- 15 10. Plachter B, Wieczorek L, Scholl B-C, Ziegelmaler R and Jahn G. Detection of Cytomegalovirus antibodies by an enzyme linked immunosorbent assay using recombinant polypeptides of the large phosphorylated tegument protein pp150. J Clin Microbiol 1992; 30:201-206.
- 20 11. Ripalti A, Landini MP, Mocarski ES and La Placa M. identification and preliminary use of recombinant lambda gt11 fusion proteins in Cytomegalovirus diagnosis. J Gen Virol 1989; 70:1247-1251.
12. Ripalti A, Ruan Q, Boccuni MC, Campanini F, Bergamini G and

- 73 -

Landini MP. Construction of a polyepitope fusion antigens of human cytomegalovirus ppUL32: reactivity with human antibodies J Clin Microbiol 1994 ; 32: 358-363.

13. Scholl BC, Von Hintzestein B, Borisch B, Traupe B, Broker M, Jahn G. Procaryotic expression of immunogenic polypeptides of the large phosphoprotein (pp150) of human cytomegalovirus J Gen Virol 1988; 69:1195-1204.

14. Vornhagen R, Plachter B, Hinderer W, The TH, Van Zanten J, Matter L, Schmidt CA, Sonneborn HH, Jahn G. Early serodiagnosis of acute cytomegalovirus infection by Enzyme-linked immunosorbent assay using recombinant antigens. J Clin Microbiol 1994; 32: 981-986.

15. Jahn G, Kouzarides T, Mach M, Scholl BC, Plachter B, Traupe B, Preddie E, Satchwell SC, Fleckenstein B, Barrell BO. Map position and nucleotide sequence of the gene for the large structural phosphoprotein of human cytomegalovirus . J Virol 1987; 61:1358-1367.

16. Landini MP, Mirolo G, Baldassarri B, La Placa M. Human immune response to Cytomegalovirus structural polypeptides studied by immunoblotting. J Med Virol 1985; 17:303-311.

17. Novak I, Sova P, Krchnak V, Hamsikova E, Zavadova H E Mapping of serologically relevant regions of human cytomegalovirus phosphoprotein pp150 using synthetic peptides. J Gen Virol 1991; 72:1409-1413.

18. Mocarski ES, Pereira L, Michael N. Precise localisation of genes on

- 74 - 2194135

large animal genomes: use of lambda gt11 and monoclonal antibodies to map the gene for a Cytomegalovirus protein family. Proceedings of the National Academy of Science (USA) 1985; 82:1266-1270

19. Karlin 5, Mocarski ES, Schachtel GA. Molecular evolution of herpes
5 viruses: genomic and protein sequence comparisons. Journal of Virology 1994; 68: 1886-1902.

20. Bolling TY, Mandecki M. An Escherichia coli expression vector for high-level production of heterologous proteins in fusion with CVP-KDO synthetase. Bio/Techniques 1990; 8: 488-490.

10 21. Ripalti A, Dal Monte P, Boccuni, MC, Campanini, F, Lazzarotto T, Campisi B, Ruan Q, Landini MP. Prokaryotic expression of a large fragment of the most antigenic cytomegalovirus DNA-binding protein (ppUL44) and its reactivity with human antibodies. J Virol Methods 1994;46: 39-50.

15 22. Robinson IM, Pilot-Matias TI, Pratt SD, Patel CB, Bevirt TS, Hunt IC. Analysis of the humoral response to the flagellin protein of Borrelia burgdorferi: cloning of regions capable of differentiating lyme disease from syphilis. J Clin Microbiol 1993; 31: 629-635.

23. Gleaves CA, Smith TF, Shuster EA, Pearson & R. Rapid detection of
20 cytomegalovirus in MRC5 cells inoculated with urine specimens by use of low speed centrifugation and monoclonal antibody to an early antigen. J Clin Microbiol 1984; 19: 917-919.

24. Van der Bijl, Torensma R, Son WJ, Schirm J, Tegzess AM, The

- TH. Rapid immunodiagnosis of active cytomegalovirus infection by monoclonal antibody staining of blood leukocytes. J Med Virol 1988; 25:179-188.
25. Revello MG, Zavattoni M, Percivalle E, Grossi P, Gerna G.
5 Correlation between immunofluorescent detection of human cytomegalovirus immediate early antigens in polymorphonuclear leukocytes and viremia. J Inf Dis 1989; 160:159-160.
27. Lazzarotto T, Furlini G, Re MC, Ramazzotti E, Campisi B, Landini MP. Human cytomegalovirus replication correlates with differentiation
10 in a hematopoietic progenitor cell line and can be modulated by HtV. Arch Virol 1994; 135:13-28.
30. Gold D, Ashley R, Handsfield HH, Verdon M, Leach L, Mills I, Drew L, Corey L. Immunoblot analysis of the humoral immune response in primary cytomegalovirus infection. J Inf Dis 1988; 157: 319-325.
- 15 31. Re MC, Landini MP. IgM to human cytomegalovirus: Comparison of two enzyme immunoassays and IgM reactivity to viral polypeptides detected by immunoblotting. J Clin Lab Anal 1989; 3:169-173.

- 76 - 2194135

SEQUENCE LISTING

(iii) NUMBER OF SEQUENCES: 4

5

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

10

(1) SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 908 base pairs

15

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

20

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..900

25

SEQ ID NO: 1:

30

GAA	TTC	ACA	GCC	AAT	AAC	CGC	GTC	AGT	TTC	CAC	GGC	GTA	AAA	AAC	ATG	48
Glu	Phe	Thr	Ala	Asn	Asn	Arg	Val	Ser	Phe	His	Gly	Val	Lys	Asn	Met	
1				5				10					15			

35

CGT	ATC	AAC	GTG	CAG	CTG	AAG	AAC	TTC	TAC	CAG	ACG	CTG	CTC	AAT	TGC	96
Arg	Ile	Asn	Val	Gln	Leu	Lys	Asn	Phe	Tyr	Gln	Thr	Leu	Leu	Asn	Cys	
		20				25						30				

40

GCC	GTC	ACC	AAA	CTA	CCG	TGC	ACG	CTG	CGT	ATA	GTT	ACG	GAG	CAC	GAC	144
Ala	Val	Thr	Lys	Leu	Pro	Cys	Thr	Leu	Arg	Ile	Val	Thr	Glu	His	Asp	
		35				40						45				

45

ACG	CTG	TTG	TAC	GTG	GCC	AGC	CGC	AAC	GGT	CTG	TTC	GCC	GTG	GAG	AAC	192
Thr	Leu	Leu	Tyr	Val	Ala	Ser	Arg	Asn	Gly	Leu	Phe	Ala	Val	Glu	Asn	
	50					55				60						

50

TTT	CTC	ACC	GAG	GAA	CCT	TTC	CAG	CGT	GGC	GAT	CCC	TTC	GAC	AAA	AAT	240
Phe	Leu	Thr	Glu	Glu	Pro	Phe	Gln	Arg	Gly	Asp	Pro	Phe	Asp	Lys	Asn	
65				70				75						80		

TAC	GTC	GGG	AAC	AGC	GGC	AAG	TCG	CGT	GGC	GGC	GGC	GGT	GGT	GGC	GGC	288
Tyr	Val	Gly	Asn	Ser	Gly	Lys	Ser	Arg	Gly	Gly	Gly	Gly	Gly	Gly	Gly	
			85					90						95		

55

AGC	CTC	TCT	TCG	CTG	GCC	AAT	GCC	GGC	GGT	CTG	CAC	GAC	GAC	GGC	CCG	336
Ser	Leu	Ser	Ser	Leu	Ala	Asn	Ala	Gly	Gly	Leu	His	Asp	Asp	Gly	Pro	
			100					105						110		

GGT	CTG	GAT	AAC	GAT	CTC	ATG	AAC	GAG	CCC	ATG	GGT	CTC	GGC	GGT	CTG	384
Gly	Leu	Asp	Asn	Asp	Leu	Met	Asn	Glu	Pro	Met	Gly	Leu	Gly	Gly	Leu	
		115				120						125				

60

GGA	GGA	GGT	GGC	GGC	GGT	GGC	GGC	AAG	AAG	CAC	GAC	CGC	GGT	GGC	GGC	432
Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Lys	Lys	His	Asp	Arg	Gly	Gly	Gly	
		130				135						140				

- 77 -

5 GGT GGT TCC GGT ACG CGG AAA ATG AGC AGC GGT GGC GGC GGC GGT GAT 480
 Gly Gly Ser Gly Thr Arg Lys Met Ser Ser Gly Gly Gly Gly Gly Asp
 145 150 155 160

10 CAC GAT CAC GGT CTT TCC TCC AAG GAA AAA TAC GAG CAG CAC AAG ATC 528
 His Asp His Gly Leu Ser Ser Lys Glu Lys Tyr Glu Gln His Lys Ile
 165 170 175

15 ACC AGT TAC CTG ACG TCC AAA GGT GGA TCG GGC GGC GGC GGA GGA GGA 576
 Thr Ser Tyr Leu Thr Ser Lys Gly Gly Ser Gly Gly Gly Gly Gly Gly
 180 185 190

20 GGA GGC GGC GGT TTG GAT CGC AAC TCC GGC AAT TAC TTC AAC GAC GCG 624
 Gly Gly Gly Gly Leu Asp Arg Asn Ser Gly Asn Tyr Phe Asn Asp Ala
 195 200 205

25 AAA GAG GAG AGC GAC AGC GAG GAT TCT GTA ACG TTC GAG TTC GTC CCT 672
 Lys Glu Glu Ser Asp Ser Glu Asp Ser Val Thr Phe Glu Phe Val Pro
 210 215 220

30 AAC ACC AAG AAG CAA AAG TGC GGC ACG CCG ACG CCT GTC AAT CCT TCC 720
 Asn Thr Lys Lys Gln Lys Cys Gly Thr Pro Thr Pro Val Asn Pro Ser
 225 230 235 240

35 ACG GCC CCC GCT CCG GCC CCG ACA CCT ACC TTC GCG AAG CTT CAG GAA 768
 Thr Ala Pro Ala Pro Ala Pro Thr Pro Thr Phe Ala Lys Leu Gln Glu
 245 250 255

40 TTC GGA TCG TCG CCC CAG AAG AGC GGT ACG GGG CCG CAA CCG GGT TCT 816
 Phe Gly Ser Ser Pro Gln Lys Ser Gly Thr Gly Pro Gln Pro Gly Ser
 260 265 270

45 GCC GGC ATG GGG GGC GCC AAA ACG CCG TCG GAC GCC GTG CAG AAC ATC 864
 Ala Gly Met Gly Gly Ala Lys Thr Pro Ser Asp Ala Val Gln Asn Ile
 275 280 285

50 CTC CAA AAG ATC GAG AAG ATT AAG AAC ACG GAG GAA TAGAATTC 908
 Leu Gln Lys Ile Glu Lys Ile Lys Asn Thr Glu Glu
 290 295 300

(2) SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 300 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

50

SEQ ID NO: 2:

55 Glu Phe Thr Ala Asn Asn Arg Val Ser Phe His Gly Val Lys Asn Met
 1 5 10 15

Arg Ile Asn Val Gln Leu Lys Asn Phe Tyr Gln Thr Leu Leu Asn Cys
 20 25 30

60 Ala Val Thr Lys Leu Pro Cys Thr Leu Arg Ile Val Thr Glu His Asp
 35 40 45

Thr Leu Leu Tyr Val Ala Ser Arg Asn Gly Leu Phe Ala Val Glu Asn

- 78 - 2194135

50 55 60

5 Phe Leu Thr Glu Glu Pro Phe Gln Arg Gly Asp Pro Phe Asp Lys Asn
65 70 75 80

Tyr Val Gly Asn Ser Gly Lys Ser Arg Gly Gly Gly Gly Gly Gly Gly
85 90 95

10 Ser Leu Ser Ser Leu Ala Asn Ala Gly Gly Leu His Asp Asp Gly Pro
100 105 110

Gly Leu Asp Asn Asp Leu Met Asn Glu Pro Met Gly Leu Gly Gly Leu
115 120 125

15 Gly Gly Gly Gly Gly Gly Gly Gly Lys Lys His Asp Arg Gly Gly Gly
130 135 140

Gly Gly Ser Gly Thr Arg Lys Met Ser Ser Gly Gly Gly Gly Asp
145 150 155 160

20 His Asp His Gly Leu Ser Ser Lys Glu Lys Tyr Glu Gln His Lys Ile
165 170 175

25 Thr Ser Tyr Leu Thr Ser Lys Gly Gly Ser Gly Gly Gly Gly Gly Gly
180 185 190

Gly Gly Gly Gly Leu Asp Arg Asn Ser Gly Asn Tyr Phe Asn Asp Ala
195 200 205

30 Lys Glu Glu Ser Asp Ser Glu Asp Ser Val Thr Phe Glu Phe Val Pro
210 215 220

35 Asn Thr Lys Lys Gln Lys Cys Gly Thr Pro Thr Pro Val Asn Pro Ser
225 230 235 240

Thr Ala Pro Ala Pro Ala Pro Thr Pro Thr Phe Ala Lys Leu Gln Glu
245 250 255

40 Phe Gly Ser Ser Pro Gln Lys Ser Gly Thr Gly Pro Gln Pro Gly Ser
260 265 270

Ala Gly Met Gly Gly Ala Lys Thr Pro Ser Asp Ala Val Gln Asn Ile
275 280 285

45 Leu Gln Lys Ile Glu Lys Ile Lys Asn Thr Glu Glu
290 295 300

50 (3) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 906 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

60 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- 79 -

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Human Cytomegalovirus (HCMV)

5

(vii) IMMEDIATE SOURCE:

(B) CLONE: pCMV-27

(ix) FEATURE:

10

(A) NAME/KEY: CDS

(B) LOCATION:1..906

(D) OTHER INFORMATION:/function= "HCMV antigen"

/product= "fusion protein alc2f3h10"

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

20	ACG CCG ACG CCT GTC AAT CCT TCC ACG GCC CCC GCT CCG GCC CCG ACA	48
	Thr Pro Thr Pro Val Asn Pro Ser Thr Ala Pro Ala Pro Ala Pro Thr	
	1 5 10 15	
	CCT ACC TTC GCG AAG CTT CAG GAA TTC GGA TCG TCG CCC CAG AAG AGC	96
	Pro Thr Phe Ala Lys Leu Gln Glu Phe Gly Ser Ser Pro Gln Lys Ser	
	20 25 30	
25	GGT ACG GGG CCG CAA CCG GGT TCT GCC GGC ATG GGG GGC GCC AAA ACG	144
	Gly Thr Gly Pro Gln Pro Gly Ser Ala Gly Met Gly Gly Ala Lys Thr	
	35 40 45	
30	CCG TCG GAC GCC GTG CAG AAC ATC CTC CAA AAG ATC GAG AAG ATT AAG	192
	Pro Ser Asp Ala Val Gln Asn Ile Leu Gln Lys Ile Glu Lys Ile Lys	
	50 55 60	
35	AAC ACG GAG GAA CTT CAG GAA TTC ACA GCC AAT AAC CGC GTC AGT TTC	240
	Asn Thr Glu Glu Leu Gln Glu Phe Thr Ala Asn Asn Arg Val Ser Phe	
	65 70 75 80	
40	CAC GGC GTA AAA AAC ATG CGT ATC AAC GTG CAG CTG AAG AAC TTC TAC	288
	His Gly Val Lys Asn Met Arg Ile Asn Val Gln Leu Lys Asn Phe Tyr	
	85 90 95	
45	CAG ACG CTG CTC AAT TGC GCC GTC ACC AAA CTA CCG TGC ACG CTG CGT	336
	Gln Thr Leu Leu Asn Cys Ala Val Thr Lys Leu Pro Cys Thr Leu Arg	
	100 105 110	
	ATA GTT ACG GAG CAC GAC ACG CTG TTG TAC GTG GCC AGC CGC AAC GGT	384
	Ile Val Thr Glu His Asp Thr Leu Leu Tyr Val Ala Ser Arg Asn Gly	
	115 120 125	
50	CTG TTC GCC GTG GAG AAC TTT CTC ACC GAG GAA CCT TTC CAG CGT GGC	432
	Leu Phe Ala Val Glu Asn Phe Leu Thr Glu Glu Pro Phe Gln Arg Gly	
	130 135 140	
55	GAT CCC TTC GAC AAA AAT TAC GTC GGG AAC AGC GGC AAG TCG CGT GGC	480
	Asp Pro Phe Asp Lys Asn Tyr Val Gly Asn Ser Gly Lys Ser Arg Gly	
	145 150 155 160	
60	GGC GGC GGT GGT GGC GGC AGC CTC TCT TCG CTG GCC AAT GCC GGC GGT	528
	Gly Gly Gly Gly Gly Gly Ser Leu Ser Ser Leu Ala Asn Ala Gly Gly	
	165 170 175	
	CTG CAC GAC GAC GGC CCG GGT CTG GAT AAC GAT CTC ATG AAC GAG CCC	576

- 80 -

2194135

	Leu	His	Asp	Asp	Gly	Pro	Gly	Leu	Asp	Asn	Asp	Leu	Met	Asn	Glu	Pro	
				180					185					190			
5	ATG	GGT	CTC	GGC	GGT	CTG	GGA	GGA	GGT	GGC	GGC	GGT	GGC	GGC	AAG	AAG	624
	Met	Gly	Leu	Gly	Gly	Leu	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Lys	Lys	
			195				200					205					
10	CAC	GAC	CGC	GGT	GGC	GGC	GGT	GGT	TCC	GGT	ACG	CGG	AAA	ATG	AGC	AGC	672
	His	Asp	Arg	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Thr	Arg	Lys	Met	Ser	Ser	
		210					215					220					
15	GGT	GGC	GGC	GGC	GGT	GAT	CAC	GAT	CAC	GGT	CTT	TCC	TCC	AAG	GAA	AAA	720
	Gly	Gly	Gly	Gly	Gly	Asp	His	Asp	His	Gly	Leu	Ser	Ser	Lys	Glu	Lys	
	225					230				235					240		
	TAC	GAG	CAG	CAC	AAG	ATC	ACC	AGT	TAC	CTG	ACG	TCC	AAA	GGT	GGA	TCG	768
	Tyr	Glu	Gln	His	Lys	Ile	Thr	Ser	Tyr	Leu	Thr	Ser	Lys	Gly	Gly	Ser	
				245					250						255		
20	GGC	GGC	GGC	GGA	GGA	GGA	GGA	GGC	GGC	GGT	TTG	GAT	CGC	AAC	TCC	GGC	816
	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Leu	Asp	Arg	Asn	Ser	Gly	
				260					265					270			
25	AAT	TAC	TTC	AAC	GAC	GCG	AAA	GAG	GAG	AGC	GAC	AGC	GAG	GAT	TCT	GTA	864
	Asn	Tyr	Phe	Asn	Asp	Ala	Lys	Glu	Glu	Ser	Asp	Ser	Glu	Asp	Ser	Val	
			275				280						285				
30	ACG	TTC	GAG	TTC	GTC	CCT	AAC	ACC	AAG	AAG	CAA	AAG	TGC	GGC			906
	Thr	Phe	Glu	Phe	Val	Pro	Asn	Thr	Lys	Lys	Gln	Lys	Cys	Gly			
		290					295					300					

(4) INFORMATION FOR SEQ ID NO: 4:

35

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 302 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

40

- (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

45	Thr	Pro	Thr	Pro	Val	Asn	Pro	Ser	Thr	Ala	Pro	Ala	Pro	Ala	Pro	Thr	
	1				5					10					15		
	Pro	Thr	Phe	Ala	Lys	Leu	Gln	Glu	Phe	Gly	Ser	Ser	Pro	Gln	Lys	Ser	
			20						25					30			
50	Gly	Thr	Gly	Pro	Gln	Pro	Gly	Ser	Ala	Gly	Met	Gly	Gly	Ala	Lys	Thr	
			35				40						45				
	Pro	Ser	Asp	Ala	Val	Gln	Asn	Ile	Leu	Gln	Lys	Ile	Glu	Lys	Ile	Lys	
		50					55					60					
55	Asn	Thr	Glu	Glu	Leu	Gln	Glu	Phe	Thr	Ala	Asn	Asn	Arg	Val	Ser	Phe	
	65					70					75					80	
	His	Gly	Val	Lys	Asn	Met	Arg	Ile	Asn	Val	Gln	Leu	Lys	Asn	Phe	Tyr	
				85					90						95		
60	Gln	Thr	Leu	Leu	Asn	Cys	Ala	Val	Thr	Lys	Leu	Pro	Cys	Thr	Leu	Arg	
				100					105					110			

- 81 -

	Ile	Val	Thr	Glu	His	Asp	Thr	Leu	Leu	Tyr	Val	Ala	Ser	Arg	Asn	Gly	
				115				120					125				
5	Leu	Phe	Ala	Val	Glu	Asn	Phe	Leu	Thr	Glu	Glu	Pro	Phe	Gln	Arg	Gly	
		130					135					140					
	Asp	Pro	Phe	Asp	Lys	Asn	Tyr	Val	Gly	Asn	Ser	Gly	Lys	Ser	Arg	Gly	
	145					150					155					160	
10	Gly	Gly	Gly	Gly	Gly	Gly	Ser	Leu	Ser	Ser	Leu	Ala	Asn	Ala	Gly	Gly	
					165					170					175		
	Leu	His	Asp	Asp	Gly	Pro	Gly	Leu	Asp	Asn	Asp	Leu	Met	Asn	Glu	Pro	
15				180					185					190			
	Met	Gly	Leu	Gly	Gly	Leu	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Lys	Lys	
		195					200						205				
20	His	Asp	Arg	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Thr	Arg	Lys	Met	Ser	Ser	
		210					215					220					
	Gly	Gly	Gly	Gly	Gly	Asp	His	Asp	His	Gly	Leu	Ser	Ser	Lys	Glu	Lys	
	225					230					235				240		
25	Tyr	Glu	Gln	His	Lys	Ile	Thr	Ser	Tyr	Leu	Thr	Ser	Lys	Gly	Gly	Ser	
				245						250					255		
	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Leu	Asp	Arg	Asn	Ser	Gly	
30				260					265					270			
	Asn	Tyr	Phe	Asn	Asp	Ala	Lys	Glu	Glu	Ser	Asp	Ser	Glu	Asp	Ser	Val	
		275						280					285				
35	Thr	Phe	Glu	Phe	Val	Pro	Asn	Thr	Lys	Lys	Gln	Lys	Cys	Gly			
		290					295					300					

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A fusion protein capable of binding with antibodies against human Cytomegalovirus (HCMV), said fusion protein comprising a first region, carrying at least one epitope binding HCMV-specific IgM between aa 202 and aa 434 inclusive, of the entire amino acid sequence of a protein pp52 encoded by a gene UL44, of said HCMV; a second region, carrying at least one epitope binding HCMV-specific IgM between aa 1006 and aa 1048 inclusive, of the entire amino acid sequence of a protein pp150 encoded by a gene UL32 of said HCMV; and a third region, carrying at least one epitope binding HCMV-specific IgM between aa 595 and aa 614 inclusive, of said pp150 protein amino acid sequence, said fusion protein comprising in a sequential manner, running in amino to carboxy terminal direction: said first region, immediately downstream wherefrom is placed said third region, and then said second region is placed downstream from said third region, a bridge sequence being inserted between said third region and said second region, said amino acids being encoded by a nucleotide sequence as set forth in SEQ ID NO:1.

2. A fusion protein as defined in claim 1, wherein said bridge sequence consists, running in amino to carboxy terminal direction of the following series of amino acids: lys leu gln glu phe.

3. A fusion protein as defined in any one of claims 1 to 2 comprising at least a portion of β -Galactosidase.

4. A fusion protein capable of binding with antibodies against human Cytomegalovirus (HCMV), comprising a fusion protein, at least part of the amino acids thereof being amino acids encoded either by the nucleotide sequence SEQ ID NO:1, read from nucleotides 001 to 900, or by the nucleotide sequence SEQ ID NO:3, read from nucleotides 001 to 906.

5. A mixture of recombinant antigens to detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay, said mixture comprising, in combination:

(i) a first fusion protein comprising: a first region, carrying an amino acid sequence (H10) corresponding to at least one epitope that binds HCMV-specific IgM between aa 202 and aa 434, inclusive, of an amino acid sequence of viral protein pp52; a second region, carrying an amino acid sequence (F3) corresponding to at least one epitope that binds HCMV-specific IgM between aa 1006 to aa 1048, inclusive, of an amino acid sequence of viral protein pp150; and a third region, carrying an amino acid sequence (A1C2) corresponding to at least one epitope that binds HCMV-specific IgM between aa 595 and aa 614, inclusive, of said viral protein pp150 amino acid sequence, said regions capable of being variably arranged with respect to one another within said first fusion protein; and said amino acids being encoded by a nucleotide sequence as set forth in SEQ ID NO:1 or SEQ ID NO:3;

(ii) a second fusion protein comprising an immunogenic region comprising a sequence of amino acids corresponding to at least one epitope that binds HCMV-specific IgM between aa 297 to aa 510, inclusive,

of the viral major matrix protein pp65 encoded by the viral gene UL83; and

(iii) a third fusion protein comprising an immunogen region comprising a sequence of amino acids corresponding to at least one epitope that binds HCMV-specific IgM between aa 117 to aa 373, of the viral assembly protein pp38 encoded by the viral gene UL80a.

6. A mixture of recombinant antigens as claimed in claim 5, wherein said regions include the entire amino acid sequence for each of said first region H10, said second region F3, said third region A1C2, said protein pp65 and said protein pp38.

7. A mixture of recombinant antigens as defined in claim 6, wherein wherein said fusion protein comprises in a sequential manner, running in amino to carboxy terminal direction, said third region (A1C2), immediately downstream wherefrom is placed said second region (F3), and then said first region (H10) is placed downstream from said second region, a first bridge sequence being inserted between said first region (H10) and said second region (F3); and downstream from said third region (A1C2), a second bridge sequence being inserted between said third region (A1C2) and second region (F3), wherein said bridge sequences consist, running in amino to carboxy terminal direction, of the following series of amino acids:

leu gln	for said first bridge; and
lys leu gln glu phe	for said second bridge.

8. A mixture of recombinant antigens as claimed in any one of the above claims 5 to 7, wherein said fusion protein comprises at least a portion of protein

CTP: CMP-3-deoxy-D-manno-octulosonate cytidylyl-transferase (CKS).

9. A plasmid, capable of being inserted into a prokaryote or eukaryote host microorganism, comprising, combined together, a first DNA sequence encoding at least one epitope that binds HCMV-specific IgM between aa 202 to aa 434, inclusive, of protein pp52 of HCMV, a second DNA sequence encoding one epitope that binds HCMV-specific IgM between aa 1006 and aa 1048, inclusive, of protein pp150, and a third DNA sequence encoding the amino acid sequence between aa 595 and aa 614, inclusive, of said protein pp150, wherein DNA sequence SEQ ID NO:3 or SEQ ID NO:1 is included in full.

10. A plasmid, capable of being inserted into either prokaryote or eukaryote host microorganism, comprising an expression vector pROS, wherein DNA sequence SEQ ID NO:1 has been inserted at site SmaI, from nucleotide 002 to 907.

11. A plasmid, capable of being inserted into either prokaryote or eukaryote host microorganism, comprising an expression vector pHIL-D1, wherein DNA sequence SEQ ID NO:1 has been inserted at site EcoRI, from nucleotide 002 to 907.

12. A plasmid as defined in claim 9, wherein DNA sequence SEQ ID NO:3 is included in full.

13. A plasmid, capable of being inserted into either a prokaryote or a eukaryote host microorganism, comprising an expression vector pJ0200, wherein DNA

sequence SEQ ID NO:3 has been inserted between the sites SacI and BamH-I, from nucleotide 001 to 906.

14. A recombinant host microorganism comprising a plasmid as claimed in claim 10, capable of producing a fusion protein comprising a first region, carrying an amino acid sequence (H10) corresponding to at least one epitope that binds HCMV-specific IgM between aa 202 and aa 434, inclusive, of viral protein pp52; a second region, carrying an amino acid sequence (F3) corresponding to at least one epitope that binds HCMV-specific IgM between aa 1006 to aa 1048, inclusive, of viral protein pp150; and a third region, carrying an amino acid sequence (A1C2) corresponding to at least one epitope that binds HCMV-specific IgM between aa 595 and aa 614, inclusive, of the same viral protein pp150.

15. A mixture of fusion proteins capable of binding with human Cytomegalovirus (HCMV)-specific antibodies, comprising fusion proteins expressed in host microorganism wherein there has been inserted at least one of the following plasmids: a plasmid as claimed in claim 10; a plasmid capable of being inserted into a prokaryote or eukaryote host organism, comprising a DNA sequence encoding at least one epitope that binds HCMV-specific IgM between aa 297 to aa 510, inclusive, of the HCMV major matrix protein pp65 encoded by the viral gene UL83, said sequence being linked at the 3' end of a second sequence corresponding to the sequence from amino acid 1 to amino acid 240, inclusive, encoding for protein CKS; and a plasmid capable of being inserted into a prokaryote or eukaryote host organism, comprising a DNA sequence encoding at least one epitope that binds HCMV-specific IgM between aa

117 to aa 373, inclusive, of the HCMV assembly protein pp38 encoded by the viral gene UL80a, said sequence being linked at the 3' end of a second sequence corresponding to the sequence from amino acid 1 to amino acid 240, inclusive, encoding for protein CKS.

16. A diagnostic reagent for diagnosing HCMV infection by serological methods, comprising a fusion protein as defined in any one of claims 1 to 4.

17. A diagnostic reagent for diagnosing HCMV infection by serological methods, comprising at least one polyclonal antibody specific for a fusion protein as defined in claim 4.

18. A diagnostic reagent for diagnosing HCMV infection by serological methods, said diagnostic reagent comprising a fusion protein, comprising sequence SEQ ID NO:2.

19. A diagnostic reagent for detecting Cytomegalovirus-specific IgM in human sera by enzyme immunoassay, said reagent comprising a mixture of fusion proteins as defined in any one of claims 5 to 8 and 15.

20. A diagnostic reagent suitable for detecting Cytomegalovirus-specific IgM in human sera by enzyme immunoassay comprising a first fusion protein as defined in claim 5.

21. A diagnostic reagent for diagnosing HCMV infection by serological methods, characterized in that it comprises a fusion protein, comprising a fusion protein comprising sequence SEQ ID NO:4.

22. A diagnostic kit for detecting, by serological methods, the presence of antibodies to human Cytomegalovirus (HCMV), comprising a diagnostic reagent as claimed in claims 16 to 21, wherein said diagnostic reagent is adsorbed on a solid medium.

23. A diagnostic kit for detecting Cytomegalovirus-specific IgM in Human sera by enzyme immunoassay, comprising a plurality of microparticles, each (of said) microparticle(s) being coated with at least one of the fusion proteins of the mixture as claimed in claim 5.

24. A diagnostic kit to detect Cytomegalovirus-specific IgM in Human sera by enzyme immunoassay, comprising a plurality of microparticles, each microparticle being coated with the mixture of the three fusion proteins, as defined in claims 5 to 8 and 15.

25. A method for detecting Cytomegalovirus-specific IgM in human sera by enzyme immunoassay said method comprising the steps of: providing a mixture of fusion proteins, the mixture comprising, in combination:

(i) a first fusion protein comprising: a first region, carrying an amino acid sequence (H10) corresponding to at least one epitope that binds HCMV-specific IgM between aa 202 and aa 434, inclusive, of viral protein pp52; a second region, carrying an amino acid sequence (F3) corresponding to at least one epitope that binds HCMV-specific IgM between aa 1006 to aa 1048, inclusive, of viral protein pp150; and a third region, carrying an amino acid sequence (A1C2) corresponding to at least one epitope that binds HCMV-

specific IgM between aa 595 and aa 614, inclusive, of the same viral protein pp150; at least part of the amino acids thereof being encoded either by the nucleotide sequence SEQ ID NO:1, read from nucleotides 001 to 900, or by the nucleotide sequence SEQ ID NO:3, read from nucleotides 001 to 906;

(ii) a second fusion protein comprising an immunogenic region consisting in a sequence of amino acids corresponding to at least one epitope that binds HCMV-specific IgM between aa 297 to aa 510, inclusive, of the viral major matrix protein pp65 encoded by the viral gene UL83; and

(iii) a third fusion protein comprising an immunogenic region consisting in a sequence of amino acids corresponding to at least one epitope that binds HCMV-specific IgM between aa 117 to aa 373, of the viral assembly protein pp38 encoded by the viral gene UL80a, contacting a sample of human serum with said mixture

to form a reaction mixture,
contacting said reaction mixture with a conjugate,
said conjugate comprising an antibody and an
enzyme,

adding a substrate for the enzyme, and
monitoring the reaction between the conjugate and the
substrate to determine the concentrate of
Cytomegalovirus-specific IgM in the serum.

26. A method for preparing a solid medium comprising a mixture of fusion proteins to prepare a diagnostic kit to detect Cytomegalovirus-specific IgM in human sera by enzyme immunoassay, said method comprising: producing a mixture by combining:

(i) a first fusion protein comprising: a first region, carrying an amino acid sequence (H10) corresponding to at least one epitope that binds HCMV-

specific IgM, between aa 202 and aa 434, inclusive, of viral protein pp52; a second region, carrying an amino acid sequence (F3) corresponding to at least one epitope that binds HCMV-specific IgM between aa 1006 to aa 1048, inclusive, of viral protein pp150; and a third region, carrying an amino acid sequence (A1C2) corresponding to at least one epitope that binds HCMV-specific IgM, between aa 595 and aa 614, inclusive, of the same viral protein pp150; at least part of the amino acids thereof being encoded either by the nucleotide sequence SEQ ID NO:1, read from nucleotides 001 to 900, or by the nucleotide sequence SEQ ID NO:3, read from nucleotides 001 to 906;

(ii) a second fusion protein comprising an immunogen region consisting in a sequence of amino acids corresponding to at least one epitope that binds HCMV-specific IgM, between aa 297 to aa 510, inclusive, of the viral major matrix protein pp65 encoded by the viral gene UL83; and

(iii) a third fusion protein comprising an immunogen region consisting in a sequence of amino acids corresponding to at least one epitope that binds HCMV-specific IgM, between aa 117 to aa 373, of the viral assembly protein pp38 encoded by the viral gene UL 80a; and allowing said mixture to be adsorbed onto at least one solid medium.

27. A plasmid according to claim 10, wherein said host microorganism consists of *E. coli*.

28. A plasmid according to claim 11, wherein said host microorganism consists of *Pichia pastoris*.

29. A fusion protein capable of binding with antibodies against human Cytomegalovirus (HCMV), said

fusion protein comprising a first region, carrying at least one epitope binding HCMV-specific IgM between aa 202 and aa 434 inclusive, of the entire amino acid sequence of a protein pp52 encoded by a gene UL44, of said HCMV; a second region, carrying at least one epitope binding HCMV-specific IgM between aa 1006 and aa 1048 inclusive, of the entire amino acid sequence of a protein pp150 encoded by a gene UL32 of said HCMV; and a third region, carrying at least one epitope binding HCMV-specific IgM between aa 595 and aa 614 inclusive, of said pp150 protein amino acid sequence, as set forth in SEQ ID NO:3.

1/30




Fig. 1

2/30





Fig. 2

3/30

HCMV Genomic DNA

Sall Sense Primer n. 1783
HindIII Antisense Primer n. 1842
ppUL32 (pp150)
PCR Amplification
Sall/HindIII Digestion
Gel Purify
60 bp Fragment

Sall-A1C2-HindIII



Sall/HindIII Digestion
Gel Purify Vector Backbone

Ligate




Fig. 3

4/30




Fig. 4

5/30




Fig. 5

TABLE 1

Comparison between IgG detection by ELISA with whole CMV or by recombinant ELISA with individual fusion proteins

ELISA (IgG) with whole CMV		ELISA (IgG) with the following fusion proteins:														
		F3			A1C2			A1C2/F3			H10			A1C2/F3/H10		
		N.	Mean Titre (Range)	Positives	N.	Mean Titre (Range)	Positives	N.	Mean Titre (Range)	Positives	N.	Mean Titre (Range)	Positives	N.	Mean Titre (Range)	Positives
28	291 (242-330)	19	75 (17-180)	10	41 (17-43)	20	172 (40-490)	19	214 (168-501)	27	176 (140-641)					
40	576 (434-705)	40	434 (53-2079)	22	110 (12-440)	40	825 (230-2209)	39	611 (206-1038)	40	880 (1202160)					
40	1161 (690-2138)	38	796 (50-2461)	22	114 (25-275)	38	805 (50-2351)	40	758 (300-2120)	40	940 (120-2405)					
108	TOTAL	98 (90.7%)		54 (50%)		97 (89.8%)		98 (90.7%)		107 (99.1%)						
14	0.18 (0.2-5)	0	—	0	—	0	—	0	—	0	—			0	—	

Fig. 6

TABLE 2
Comparison between IgM detection by ELISA with whole CMV or by recombinant ELISA with individual fusion proteins

ELISA (IgM) with whole CMV		ELISA (IgM) with the following fusion proteins:									
		F3		A1C2		A1C2/F3		H10		A1C2/F3/H10	
N.	Mean Titre (Range)	N. Positives	Mean Titre (Range)	N. Positives	Mean Titre (Range)	N. Positives	Mean Titre (Range)	N. Positives	Mean Titre (Range)	N. Positives	Mean Titre (Range)
20	294 (242-389)	10	521 (232-1171)	10	317 (85-612)	14	797 (210-1417)	19	578 (432-724)	20	517 (386-815)
24	516 (210-2297)	18	1090 (210-2297)	16	467 (143-1141)	22	1278 (370-2561)	23	887 (209-2270)	24	767 (298-1973)
20	1341 (881-2000)	16	1163 (451-2339)	14	1021 (202-2260)	16	1707 (521-2249)	20	1065 (579-2471)	20	1121 (421-2243)
64	TOTAL positives	44 (68,75%)		40 (62,5%)		52 (81,25%)		62 (96,87%)		64 (100%)	
5	27 (1-56)	0	—	0	—	0	—	0	—	0	—

Fig. 7

8/30




Fig. 8

9/30




Fig. 9

10/30




Fig. 10

11/30




Fig. 11

12/30




Fig. 12

13/30




Fig. 13A

14/30





Fig. 13B

15/30




Fig. 13C

16/30




Fig. 14A

17/30




Fig. 14B



Fig. 14C




Fig. 14D

18/30




Fig. 15

19/30




Fig. 16A

20/30




Fig. 16B

21/30




Fig. 17A

22/30




Fig. 17B

TABLE 3

Viral Protein	Approx. MW x10 ³ kD	Fusion Protein		Protein Data			Assay Cut-off	
		automated test	manual test	Nucleotides	Amino Acids	Manual test OD x 10 ³	Assay test	Autom test
ppUL32+ppUL44	58		rpCMV-4	1783-1842+3016-3144(ppUL32) + 604-1299(ppUL44)	595-616 + 1006-1048(ppUL32) + 202-434(ppUL44)	180		
ppUL80a	54		rpCMV-26	349-1119	117-373	180		
ppUL32	36		rpCMV-1A	1783-1842+3016-3144	595-614 + 1006-1048	130		
ppUL44	51		rpCMV-3B	604-1299	202-434	170		
ppUL83	51		rpCMV-9	889-1530	297-510	110		
ppUL32+ppUL44	60	rpCMV-27		1783-1842+3016-3144(ppUL32) + 604-1299(ppUL44)	595-616 + 1006-1048(ppUL32) + 202-434(ppUL44)			0.6
ppUL83	53	rpCMV-28		889-1530	297-510			0.6
ppUL80a	55	rpCMV-29		349-1119	117-373			0.6

Fig. 18

Table 4: Automated HCMV IgM immunoassay results obtained by testing 7 HCMV IgM-negative sera using microparticles coated separately and then run sequentially and in combination

Sample	Sample Index Value with Coated Microparticles				Sample Result (Pos or Neg)	
	27	28	29	27+28+29	Sequential	Combination
BT10578	0.076	0.071	0.106	0.075	Neg	Neg
BS18323	0.099	0.044	0.090	0.104	Neg	Neg
BT12170	0.202	0.070	0.140	0.167	Neg	Neg
BT12155	0.044	0.059	0.123	0.065	Neg	Neg
BT12173	0.204	0.111	0.316	0.255	Neg	Neg
BT12162	0.436	0.251	0.200	0.476	Neg	Neg
BT12267	0.198	0.141	0.496	0.296	Neg	Neg

Fig. 19

Table 5: Automated HCMV IgM immunoassay results obtained by testing 11 HCMV IgM-positive sera using microparticles coated separately and then run sequentially and in combination

Sample	Viral WB	Sample Index Value with Coated Microparticles			Sample Result (Pos or Neg)	
		27	28	29	27+28+29	Sequential Combination
495	150,65,55,38	3.860	0.894	4.049	4.550	Pos
530	150,65,55,38	1.437	1.416	2.395	1.647	Pos
705	150,65,55,38	1.752	0.120	0.560	2.091	Pos
525	150	0.784	0.214	0.438	0.833	Pos
587	150,38	0.824	1.144	4.780	2.011	Pos
316	150,38	0.629	0.047	0.724	0.653	Pos
498	150,38	0.616	0.084	1.502	0.787	Pos
317	150,65	0.036	1.041	0.109	0.973	Pos
472	150,65	0.960	1.988	0.572	0.632	Pos
457	150,65	2.768	0.744	0.717	3.127	Pos
301	150,65	2.888	0.157	0.658	2.214	Pos

Fig. 20

Table 6: Recombinant EIA results obtained by testing 150 IgM-positive sera

N. of sera	IgG	HCMV serology* IgM	N of sera positive for:					all fusion proteins
			1A	3B	4	9	26	4+9+26
50	+	high	42	49	49	44	39	50
50	+	medium	31	47	44	42	37	50
50	+	low	29	40	42	31	32	47
tot. 150			102	136	135	117	108	147 (98%)
								148 (98.7%)

* Sera were divided in three groups on the basis of EIA-IgM values. All the sera were IgM-confirmed by Western blotting. They were all IgG positive as shown by EIA.

Fig. 21

Table 7: Comparison between conventional and recombinant EIA in pregnant women. Determination of IgM titres to individual fusion proteins

Groups of subjects	N	N of positive samples detected by						IgM titres (ODx10 ³) detected by			
		Rec- EIA with fusion proteins		26 4+9+26		all		Rec-EIA 3B	with fusion 4	proteins 9	26
		1A		9	10	20	20	461 (1/211)	838 (1/124)	164 (1/59)	487 (1/220)
HCMV-infected pregnant women who did not transmit the infection*	25	1	15	19					p= 0.012		p=0.023
HCMV-infected pregnant women who transmitted the infection*	8	1	6	8	6	8	6	724 (1/344)	1450 (1/379)	191 (1/32)	1064 (1/490)
HCMV-uninfected pregnant women*	18	0	0	0	0	0	0	nd	nd	nd	nd

* Samples were obtained at 22-24 weeks of gestation

Fig. 22

Table 8: Comparison between conventional and recombinant EIA in newborns

Groups of subjects	N.	N. of positive samples detected by						Conventional EIA
		Rec - IA	EIA with 3B	fusion 4	proteins: 9	26	4+9+26 all	
Congenitally infected newborns*	6	0	1	1	2	2	2	0
Newborn excreting HCMV during the first year of life**	19	0	4	7	1	7	10	2
HCMV uninfected newborns**	10	0	0	0	0	0	0	0

* Sera were obtained 1-7 days after birth
** Sera were obtained between 1 and 12 months after birth.

Fig. 23

Table 9: Cumulative IgM reactions to the five fusion proteins obtained in 253 IgM-positive sera from different groups of subjects

Recombinant proteins	Number of positive		samples	among:		Infected newborns ^{oo} (n10)	Infected transplant recipients (n65) GROUPS §						Tot n (%)
	IgM+sera (n 148)	Infected pregnant women* (n20)		Infected pregnant women** (n8)	Infected newborns ^o (n2)		I	II	III	IV	V	VI	
							1	15	17	11	15	5	
1A	102	1	1	0		0	1	5	7	4	7	0	128 (50.6)
3B	136	15	6	1		4	1	7	13	9	10	2	204 (80.6)
4	135	19	8	1		7	1	12	17	11	14	3	228 (90.1)
9	117	9	3	0		1	2	8	7	8	7	2	164 (64.8)
26	108	10	6	2		7	1	11	15	6	10	3	179 (70.7)
4+9	144	20	8	1		7	2	14	17	11	14	3	241 (95.2)
4+26	142	19	8	2		10	1	13	17	11	15	5	243 (96.0)
4+9+26	147	20	8	2		10	2	15	17	11	15	5	252 (99.6)

* HCMV infected pregnant women who did not transmit the infection
** HCMV-infected pregnant women who transmitted the infection
§ Groups are referred to Table 5
° Sera were obtained during the first week of life
oo Sera were obtained from 1 to 12 months after birth

Fig. 24

Table 10: Comparison between conventional and recombinant EIA (fusion protein 4, 9 and 26) and Western blotting in 200 sera randomly selected from renal transplant recipients and pregnant women.

		Recombinant EIA		Western Blotting	
		positive	negative	positive	negative
Conventional EIA	positive	88	2	86	2
	negative	112	62	51	61
	totals	200	64	137	63

Fig. 25