

(43) International Publication Date
11 November 2010 (11.11.2010)(10) International Publication Number
WO 2010/129672 A1(51) International Patent Classification:
C07H 21/00 (2006.01)(21) International Application Number:
PCT/US2010/033729(22) International Filing Date:
5 May 2010 (05.05.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/175,690 5 May 2009 (05.05.2009) US(71) Applicant (for all designated States except US): **MIRAGEN THERAPEUTICS** [US/US]; 6200 Lookout Road, Boulder, Colorado 80301 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MARSHALL, William S.** [US/US]; c/o MIRAGEN THERAPEUTICS, 6200 Lookout Road, Boulder, Colorado 80301 (US). **VAGLE, Kurt** [US/US]; c/o MIRAGEN THERAPEUTICS, 6200 Lookout Road, Boulder, Colorado 80301 (US).(74) Agents: **MCLEOD, Bonnie Weiss** et al.; COOLEY LLP, 777 6th Street, N.W., Suite 1100, Washington, DC 20001 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,




AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: LIPOPHILIC POLYNUCLEOTIDE CONJUGATES

Natural Cholesterol**Current State of the Art Chol Conjugates****Novel Chol Conjugate with enhanced Lipophilicity**

(57) **Abstract:** Disclosed are lipophilic polynucleotide conjugates including polynucleotide-cholesterol conjugates and methods of delivering therapeutic polynucleotides to a mammalian cell or patient in need of treatment using said conjugates. The disclosure further provides methods of synthesizing the lipophilic polynucleotide conjugates. The conjugates are designed to mimic or target cellular miRNAs. The lipophilic moiety, such as cholesterol or cholesterol derivative, is spaced from the polynucleotide by a substantially linear hydrocarbon group. Due to an absence of significantly polar groups and/or exchangeable protons in the vicinity of the lipophilic moiety, the interaction between the lipophilic moiety and cell membranes is enhanced to provide for efficient entry into cells.

LIPOPHILIC POLYNUCLEOTIDE CONJUGATES

CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application claims the benefit of priority of U.S. Provisional Application No. 61/175,690 filed May 5, 2009, which is herein incorporated by reference in its entirety.

DESCRIPTION OF THE TEXT FILE SUBMITTED ELECTRONICALLY

[002] The contents of the text file submitted electronically herewith are incorporated herein by reference in their entirety: A computer readable format copy of the Sequence Listing (filename: MIRG_014_00WO_SeqList_ST25.txt, date recorded: May 4, 2010, file size 10 kilobytes)

FIELD OF THE INVENTION

[003] The present invention relates to lipophilic conjugates of nucleic acid for the delivery of polynucleotides to cells for therapeutic, research and/or diagnostic purposes. The invention relates in certain embodiments to the delivery of polynucleotides that target endogenous microRNA function.

BACKGROUND

[004] Oligonucleotide therapeutic strategies rely on delivering the hydrophilic, highly charged and anionic nucleic acid molecule into the cytoplasm by crossing the hydrophobic phospholipid bilayer that constitutes the cell membrane. Hydrophobic conjugates of oligonucleotides have been investigated for their ability to enhance cellular uptake. For example, cholesterol conjugates may be capable of binding to lipoproteins, and more specifically IDL and LDL particles. The particles contain ApoE and ApoB-100 lipoproteins, which target LDL cell surface receptors and, therefore, can be internalized via lipoprotein mediated endocytosis. These receptors are found mainly in the liver, but also in adipose tissue, heart and skeletal muscle. Additionally, pendant lipophilic groups may aid in "anchoring" their conjugates to the cell membrane by intercalating into the phospholipid bilayer and therefore aiding passive internalization of the oligonucleotide.

[005] Such lipophilic molecules or substituents must be formulated with, or attached to, the polynucleotide so as to preserve the functional and structural integrity of the polynucleotide for its desired therapeutic action. For example, therapeutic polynucleotides may function through interactions with large and/or intricate ribonucleoprotein (RNP) complex(es), and/or may target RNA components of these complexes. Lipophilic

modifications should therefore not impair the accessibility of the polynucleotide for interactions with cellular factors and molecular targets.

[006] MicroRNAs (miRs) have been implicated in a number of biological processes including regulation of developmental timing, apoptosis, fat metabolism, and hematopoietic cell differentiation among others, and therefore represent a relatively new class of therapeutic targets. miRs are small, non-protein coding RNAs of about 18 to about 25 nucleotides in length, and act as repressors of target mRNAs by promoting their degradation, when their sequences are perfectly complementary, or by inhibiting translation, when their sequences contain mismatches. The mature miRNA strand is incorporated into the RNA-induced silencing complex (RISC), where it associates with its target RNAs by base-pair complementarity.

[007] Compositions and methods for efficiently delivering therapeutic polynucleotides, including miRs and miR inhibitors, to cells without substantially or significantly impairing interactions with the molecular target(s) are needed.

SUMMARY OF THE INVENTION

[008] The present invention provides lipophilic polynucleotide conjugates, and methods of delivering therapeutic polynucleotides to a mammalian cell or a mammalian patient in need of treatment. The invention further provides methods of synthesizing the lipophilic polynucleotide conjugates. In accordance with the invention, the conjugates include conjugates of cholesterol and polynucleotides, designed to mimic or target cellular miRNAs. The lipophilic moiety, such as cholesterol or cholesterol derivative, is spaced from the polynucleotide by a hydrocarbon group. Due to an absence of significantly polar groups and/or exchangeable protons in the vicinity of the lipophilic moiety, the invention enhances the interaction between the lipophilic moiety and cell membranes to provide for efficient entry into cells. The conjugate does not substantially or significantly disrupt the ability of the polynucleotide to deliver its therapeutic effect.

[009] In one aspect, the invention provides a lipophilic polynucleotide conjugate as well as pharmaceutical compositions and formulations comprising the same. In accordance with this aspect, the lipophilic moiety may be a cholesterol or similar lipophilic structure as described herein, or other group that associates with membrane phospholipids. The polynucleotide may be DNA- or RNA-based, and/or may embody one or more nucleic acid modifications, for example, such as a modified polynucleotide backbone or one or more

modified nucleoside units. For example, the modified backbone may be a full or partial phosphorothioate modified backbone. The sequence of the polynucleotide may be designed so as to mimic a miRNA or target a miRNA by antisense inhibition, for example. The polynucleotide and lipophilic moiety are spaced by, and conjugated through, a substantially linear hydrocarbon moiety (e.g., linker). The hydrocarbon linker may be conjugated to the 5' and/or 3' end of the polynucleotide, for example.

[010] The conjugate of the invention has an absence of significantly polar groups, such as groups having an exchangeable proton, within the vicinity of the lipophilic moiety. For example, the hydrocarbon may be conjugated to the lipophilic moiety (e.g., cholesterol) through a relatively non-polar group such as an ether or thioether linkage, and the hydrophobic linker may comprise from about 3 to about 15 carbon atoms. The absence of a significantly polar group, e.g., having an exchangeable proton, in the vicinity of the lipophilic moiety allows the moiety to better intercalate with membrane phospholipids upon delivery to a cell or patient.

[011] In a second aspect, the invention provides a method for delivering polynucleotides to a mammalian cell, and methods for treating or preventing the progression of a condition in a mammalian patient. In this aspect, the invention comprises administering the polynucleotide as a conjugate with cholesterol or other lipophilic moiety as described herein. The polynucleotide may be a miRNA or a miRNA inhibitor (e.g., having a nucleotide sequence designed to inhibit expression or activity of a miRNA). In such embodiments, the patient has a condition associated with RNA expression, such as miRNA expression. Such conditions include, for example, cardiac hypertrophy, myocardial infarction, heart failure (e.g., congestive heart failure), vascular damage, restenosis, and/or pathologic cardiac fibrosis, among others.

[012] In a third aspect, the invention provides a method for synthesizing the polynucleotide conjugates. The invention provides methods of synthesizing polynucleotides having lipophilic moieties conjugated at the 5' and/or 3' end, or along the polynucleotide backbone, through a hydrocarbon moiety or linker. In certain embodiments where the lipophilic moiety is an alcohol, such as a sterol, the lipophilic moiety is etherized with a hydrocarbon having a suitable terminal functional group. This ether-linked intermediate may be further reacted to prepare suitable reagents compatible with solid phase nucleic acid synthesis. For example, the ether-linked intermediate may be further reacted to prepare a terminal phosphoramidite (e.g., through an ether linkage) for converting to a phosphate ester (e.g., coupling to a nucleotide 5' hydroxyl during polynucleotide synthesis). Alternatively, or

in addition, the ether-linked intermediate may be further reacted to prepare a suitable terminal group for coupling (e.g. via amidation) to a suitable support for nucleic acid synthesis.

[013] Alternatively, the invention provides a method for synthesizing cholesterol conjugates with polynucleotides, after synthesis of the polynucleotide is complete, using for example, carbonyl addition-elimination/reductive amination, amidation, maleimide-thiol coupling, aqueous Diels-Alder and "Click" chemistries.

DESCRIPTION OF THE FIGURES

[014] **Figure 1** illustrates an exemplary lipophilic (cholesterol) conjugate of the invention as compared to conventional conjugates. In accordance with the invention, polar groups such as those having exchangeable protons are removed from the vicinity of the cholestene ring structure.

[015] **Figure 2** illustrates the synthesis of exemplary conjugates in accordance with the invention, including the preparation of phosphoramidite intermediates for coupling to a nucleotide at the 5' position (**6**), and succinate intermediates for coupling to a solid support (**7**).

[016] **Figure 3** illustrates the synthesis of an exemplary 3'-cholesterol conjugate (1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3 β -oxy)dodecan-3-ol Functionalized CPG) with the cholesterol-linked intermediate anchored to a solid support.

[017] **Figure 4** illustrates that Cholest-5-en-3-oxyalkanol (**A**) can be modified to give functional groups amenable to common bioconjugation techniques such as amidation (**B**), Diels-Alder bioconjugation (**C**) and "Click" bioconjugation (**D**) when the oligonucleotide is functionalized with the complementary functional group.

[018] **Figure 5** is a Northern analysis of miR-208 in response to IV injection with the chemistries shown in Table 2.

[019] **Figure 6** shows realtime quantification of miR-208 knockdown in response to IV injection with the indicated chemistries compared to saline injected animals.

DETAILED DESCRIPTION OF THE INVENTION

[020] Active and/or passive cellular transport mechanisms may benefit from pendant lipophilic groups, including those that are native to the phospholipid bilayer and which should efficiently pack in the bilayer. Thus, the invention provides conjugates that contain a sufficiently hydrophobic linker to efficiently pack into a phospholipid bilayer, to thereby enhance the affinity of the polynucleotide conjugates for cell membranes, or alternatively liposomes or micelles that may act as delivery vehicles. Oligonucleotide-cholesterol type conjugates have generally employed cholesteryl carbamate linkages that result in a polar group containing an exchangeable proton immediately adjacent to the steroid ring structure. These conjugates do not have optimized lipophilic properties and the polar linkage point may impede efficient phospholipid bilayer packing.


[021] The present invention provides lipophilic polynucleotide conjugates, and methods of delivering therapeutic polynucleotides to a mammalian cell or a mammalian patient in need of treatment. The invention further provides methods of synthesizing the lipophilic polynucleotide conjugates. In accordance with the invention, the conjugates may be designed to target cellular RNAs, including miRNAs. For example, the conjugates may target cellular miRNAs by antisense inhibition, or may be designed to mimic a miRNA. The lipophilic moiety, such as cholesterol or other lipophilic group, is spaced from the polynucleotide by a hydrocarbon linker. Due to an absence of significantly polar groups and/or exchangeable protons in the vicinity of the lipophilic moiety, the invention promotes interaction of the lipophilic moiety with membrane phospholipids, thereby providing efficient entry into cells. The conjugate does not substantially or significantly disrupt the ability of the polynucleotide to deliver its therapeutic effect.

Lipophilic Moiety and Hydrocarbon Spacer


[022] The lipophilic moiety is a fat soluble moiety capable of intercalating with a phospholipid bi-layer, such as a cell membrane, or delivery vehicle such as a liposome or micelle as disclosed herein. The lipophilic moiety may be conjugated to the hydrocarbon spacer (described below) through an ether or thioether linkage. Exemplary lipophilic moieties include cholesterol and cholesterol derivatives (including cholestenes, cholestanes, and cholestadienes), bile acids (such as cholic acid, deoxycholic acid and dehydrocholic acid), sterols, steroids, or other fat soluble alcohol or thiol. In certain embodiments, the lipophilic moiety is cholesterol or a cholesterol derivative, or cholic acid or cholic acid derivative as described in US Patent 7,202,227, which is hereby incorporated by reference.

Such derivatives include C1-C4 alkyl (e.g., methyl) substituted cholesterol or cholic acid structures.


[023] For example, in certain embodiments, the cholesterol or derivative may have the following structure:



; or



In such structures, each of R¹ through R⁸ is a substantially non-polar group. For example, each of R¹ through R⁸ may be independently selected from hydrogen, alkyl (e.g., C1-C6 alkyl), substituted alkyl, alkoxy (e.g., C1-C6 alkoxy) which may be substituted, and alkoxyalkyl which may be substituted. In certain embodiments, the lipophilic moiety has the following formula, where R is alkyl (e.g., C1-C10 alkyl):



[024] A hydrocarbon moiety or spacer separates the lipophilic moiety (described above) from non-lipophilic, e.g., polar, groups, so as to allow the lipophilic moiety to better interact with, *inter alia*, membrane phospholipids upon delivery to a patient or upon contact with mammalian cells. Thus, the conjugate generally does not have exchangeable protons or other polar groups (e.g., hydroxyl or amide) within the vicinity of the lipophilic moiety. Generally, there are no significantly polar groups and exchangeable protons within about 3 to about 15 atoms from the lipophilic moiety. In certain embodiments, the first significantly polar group or exchangeable proton is at least 6, 7, 8 or 9 atoms from the lipophilic moiety.


[025] The hydrocarbon spacer may be conjugated to the lipophilic moiety through a relatively non-polar bond, such as an ether or thioether linkage. For example, where the lipophilic moiety is cholesterol or other sterol, the hydrocarbon spacer may be ether-linked with the sterol hydroxyl group.

[026] In certain embodiments, the hydrocarbon spacer comprises a C2 to C15 saturated or unsaturated hydrocarbon chain (e.g., alkylene or alkenylene), and may be substituted by relatively non-polar substituents. Such substituents generally do not contain exchangeable protons or polar groups within the vicinity of the lipophilic moiety as described. For example, the substituents may be independently selected from alkyl (e.g., C1-C6 alkyl), alkoxy (e.g., C1-C6 alkoxy), and alkoxyalkyl. Where polar substituents are present, they are spaced at least about 6, 7, 8, or 9 atoms from the lipophilic moiety. Where unsaturated, the hydrocarbon spacer may comprise 1, 2, or 3 double bonds. In certain embodiments, the hydrocarbon spacer comprises a C2 to C10, or a C4 to C8 saturated hydrocarbon chain. For example, the hydrocarbon spacer may comprise a C4 or C6 saturated hydrocarbon.

[027] In certain embodiments, the hydrocarbon chain contains one or more heteroatoms (e.g., from 1 to 5), such as heteroatoms independently selected from O, S, and/or N. In certain other embodiments, the hydrocarbon chain does not have any heteroatoms, except in some embodiments as desired or resulting from the attachment of the polynucleotide to the terminus by an ether or thioether linkage.


[028] The hydrocarbon spacer is conjugated on the end opposite the lipophilic moiety to the polynucleotide, e.g., through a nucleotide 5' or 3' position, or at other location along the polynucleotide backbone. The hydrocarbon spacer may be conjugated to the polynucleotide, e.g., to the nucleotide 5' or 3' position, via any suitable functional group, including phosphodiester, ether, thioether, ester, thioester, or amide.





[029] An exemplary conjugate of the invention may have a structure selected from structures (I), (II), and (III), respectively, below:




where n is 1 or 2, and where R may be a nucleotide or polynucleotide linked through the 3' or 5' position, optionally through a linking group.



[030] Additional exemplary conjugates include structures (IV)-(VI), where R and R' are each H or phosphate polynucleotide, and structures VII-IX, where R and R' are each H, OH, or a phosphate polynucleotide.







[031] In certain embodiments, the conjugate is a 5' conjugate having one of the following structures, where R is the polynucleotide chain:



[032] In certain embodiments, the conjugate is a 3' conjugate having one of the following structures where R is the polynucleotide chain:

Chemical structure (XII) is a steroid derivative. It features a four-ring steroid nucleus. At the C3 position, there is a phosphate group: $\text{HO}-\text{CH}_2-\text{CH}_2-\text{O}-\text{P}(\text{OR})(\text{O}^-)-\text{O}-$. At the C17 position, there is a long alkyl chain: $-(\text{CH}_2)_4-\text{CH}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$. The stereochemistry is indicated with wedges and dashes at various positions, including C13, C14, C15, and C17.

(XIII)

Polynucleotides

[033] The compound, compositions and methods of the invention may employ various polynucleotides (including oligonucleotides) and derivatives thereof. The polynucleotide may be DNA- or RNA-based, and/or may embody one or more nucleic acid modifications, for example, such as a modified polynucleotide backbone or one or more modified nucleoside units. The polynucleotide or derivative may have one or more single stranded and/or one or more double stranded regions. The polynucleotide may be an antisense oligonucleotide, short interfering RNA (siRNA), double stranded RNA (dsRNA), single stranded RNA (ssRNA), microRNA (miRNA), short hairpin RNA (shRNA), or ribozyme.

[034] In certain embodiments, the compound, composition, or method employs a conjugated miRNA inhibitor (e.g., antisense inhibitor). Such inhibitors are described for example, in WO 2008/016924, WO 2009/062169, WO 2009/058818, WO 2009/018492, WO 2009/018493, WO 2009/012468, and WO 2007/070483, which are hereby incorporated by reference in their entireties.

[035] Thus, the polynucleotide may be an antisense oligonucleotide. In some embodiments, the antisense oligonucleotide contains at least one chemical modification (e.g. sugar or backbone modification). For instance, antisense oligonucleotide may comprise one or more modified nucleotides, such as 2'-O-methyl-sugar modifications. In some embodiments, antisense oligonucleotides comprise only modified nucleotides. In certain embodiments, antisense oligonucleotides may also comprise one or more phosphorothioate linkages resulting in a partial or full phosphorothioate backbone. Antisense oligonucleotides suitable for inhibiting miRNAs may be about 5 to about 50 nucleotides in length, such as about 10 to about 30 nucleotides in length, or about 20 to about 25 nucleotides in length. In certain embodiments, antisense oligonucleotides are about 8 to about 18 nucleotides in length, and in other embodiments about 12 to about 16 nucleotides in length. The antisense oligonucleotide may comprise a sequence that is at least about 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% complementary to a mature miRNA sequence (see Table 1 below). In other embodiments, the antisense oligonucleotides comprise a sequence that is 100% complementary to the mature miRNA sequence. As used herein "complementary" or "base pair" includes or refers only to classic Watson-Crick nucleotide base-pairing. In one embodiment, the antisense oligonucleotide is an antagomir. An "antagomir" is a single-stranded, optionally chemically-modified, ribonucleotide that is at least partially complementary to a miRNA sequence.

[036] In certain embodiments, the polynucleotide has a sequence designed to mimic a cellular miRNA, such as a miRNA listed in Table 1. The polynucleotide in these embodiments may also comprise one or more modified nucleotides, such as 2'-O-methyl-sugar modifications, and one or more phosphorothioate linkages. Polynucleotides suitable for mimicking miRNAs may be about 15 to about 50 nucleotides in length, such as about 18 to about 30 nucleotides in length, or about 20 to about 25 nucleotides in length. The synthetic miRNA may comprise a sequence that is at least about 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to a mature miRNA sequence (see Table 1 below). In other embodiments, the polynucleotide comprises a sequence that is 100% identical to the mature miRNA sequence.

[037] The polynucleotide may be composed of predominately ribonucleotide units or predominately deoxyribonucleotide units, and may have one or more chemical modification(s). For instance, the polynucleotides may be comprised of one or more "conformationally constrained" or bicyclic sugar nucleoside modifications (BSN) that confer enhanced thermal stability to complexes formed between the oligonucleotide containing BSN and their complementary microRNA target strand. In certain embodiments, the

polynucleotide includes, for example, from about 1 to 10 locked nucleic acids. "Locked nucleic acids" (LNAs) contain the 2'-O, 4'-C-methylene ribonucleoside wherein the ribose sugar moiety is in a "locked" conformation. Alternatively or in addition, the polynucleotide may comprise peptide nucleic acids (PNAs), which contain a peptide-based backbone rather than a sugar-phosphate backbone. In these or other embodiments, the polynucleotide contains one or more sugar modifications and/or one or more backbone modifications. Exemplary sugar modifications include 2' and 4' modifications, such as 2'-O-alkyl (e.g. 2'-O-methyl, 2'-O-methoxyethyl), 2'-fluoro, and 4' thio modifications. Exemplary backbone modifications include phosphorothioate, morpholino, or phosphonocarboxylate linkages (see, for example, U.S. Patent Nos. 6,693,187 and 7,067,641, which are herein incorporated by reference in their entireties).

[038] In some embodiments, the polynucleotide is a 2'-O-methoxyethyl gapmer. A "gapmer" contains 2'-O-methoxy ethyl-modified ribonucleotides on both 5' and 3' ends with at least ten deoxyribonucleotides in the center. These "gapmers" are capable of triggering RNase-independent degradation mechanisms of RNA targets.

[039] Other modifications of polynucleotides, for example, to enhance stability and/or improve antisense efficacy, are known, and may be employed in connection with the invention. Exemplary modifications are described in U.S. Patent No. 6,838,283, which is hereby incorporated by reference in its entirety.

[040] The polynucleotide may have a nucleotide sequence designed to mimic or target a miRNA, such as a miRNA listed in Table 1 below. In certain embodiments, the polynucleotide designed to inhibit a miRNA may have a sequence containing from 1 to 5 (e.g., 2, 3, or 4) mismatches relative to the fully complementary miRNA sequence (shown in Table 1 below). In other embodiments, the polynucleotide designed to mimic a miRNA may have a sequence containing from 1 to 5 (e.g., 2, 3, or 4) nucleotide substitutions relative to the mature miRNA sequence (shown in Table 1 below). Such antisense and sense sequences may be incorporated into shRNAs or other RNA structures containing stem and loop portions, for example. Such sequences are useful for, among other things, mimicking or targeting miRNA function for treatment or amelioration of cardiac hypertrophy, myocardial infarction, heart failure (e.g., congestive heart failure), vascular damage, and/or pathologic cardiac fibrosis, among others. Exemplary miRNA therapeutic utilities are disclosed in the US and PCT patent references listed in Table 1 below, each of which is hereby incorporated by reference in its entirety.

Table 1

<u>miRNA</u>	<u>miRNA Sequence</u>	<u>SEQ ID NO:</u>	<u>Reference</u>
1	UGGAAUGUAAAGAAGUAUGUAU	1	WO 2009/012468
100	AACCCGUAGAUCCGAACUUGUG	2	WO 2009/012468
10b	UACCCUGUAGAACCGAAUUUGUG	3	WO 2009/012468
125b	UCCCUGAGACCCUAACUUGUGA	4	WO 2009/012468
128	UCACAGUGAACCGGUCUCUUU	5	WO 2007/070483
133a	UUUGGUCCCCUUAACCAGCUG	6	WO 2009/012468
133b	UUUGGUCCCCUUAACCAGCUA	7	WO 2009/012468
139	UCUACAGUGCACGUGUCUCCAG	8	WO 2009/012468
143	UGAGAUGAAGCACUGUAGCUC	9	WO 2007/070483
145	GUCCAGUUUUCCCAGGAAUCCCU	10	WO 2007/070483
150	UCUCCCAACCCUUGUACCAGUG	11	WO 2009/012468
15a	UAGCAGCACAUAAUGGUUUGUG	12	WO 2009/062169
15b	UAGCAGCACAUCAUGGUUUACA	13	WO 2009/062169
16	UAGCAGCACGUAAAUAUUGGCG	14	WO 2009/062169
181b	AACAUUCAUUGCUGUCGGUGGGU	15	WO 2009/012468
195	UAGCAGCACAGAAAUAUUGGC	16	WO 2009/012468
197	UUCACCACCUUCUCCACCCAGC	17	WO 2009/012468
199a	CCCAGUGUUCAGACUACCUGUUC	18	WO 2009/012468
199b	miR-199b-5p CCCAGUGUUUAGACUAUCUGUUC	19	US 61/047,005
	miR-199b-3p ACAGUAGUCUGCACAUUGGUUA	20	
206	UGGAAUGUAAGGAAGUGUGUGG	21	WO 2007/070483
208a	AUAAGACGAGCAAAAAGCUUGU	22	WO 2008/016924
208b	AUAAGACGAACAAAAGGUUUGU	23	WO 2009/018492

<u>miRNA</u>	<u>miRNA Sequence</u>	<u>SEQ ID NO:</u>	<u>Reference</u>
20a	UAAAGUGCUUAUAGUGCAGGUAG	24	US 60/950,565
21	UAGCUUAUCAGACUGAUGUUGA	25	WO 2009/058818
214	ACAGCAGGCACAGACAGGCAGU	26	US 61/047,005
22	AAGCUGCCAGUUGAAGAACUGU	27	WO 2009/012468
221	AGCUACAUUGUCUGCUGGGUUUC	28	WO 2009/012468
222	AGCUACAUCUGGCUACUGGGU	29	WO 2009/012468
224	CAAGUCACUAGUGGUUCCGUU	30	WO 2009/012468
23a	AUCACAUUGCCAGGGAUUUCC	31	WO 2009/012468
26a	UUCAAGUAAUCCAGGAUAGGCU	32	WO 2007/070483
26b	UUCAAGUAAUUCAGGAUAGGU	33	WO 2009/012468
28	AAGGAGCUCACAGUCUAUUGAG	34	WO 2009/012468
29a	UAGCACCAUCUGAAAUCGGUUA	35	WO 2009/018493
29b	UAGCACCAUUUGAAAUCAGUGUU	36	WO 2009/018493
29c	UAGCACCAUUUGAAAUCGGUUA	37	WO 2009/018493
30a	UGUAAACAUCCUCGACUGGAAG	38	PCT/US2010/031147
30b	UGUAAACAUCCUACACUCAGCU	39	PCT/US2010/031147
30c	UGUAAACAUCCUACACUCUCAGC	40	WO 2009/012468
30d	UGUAAACAUCCCCGACUGGAAG	41	PCT/US2010/031147
30e	UGUAAACAUCCUUGACUGGAAG	42	PCT/US2010/031147
342-3p	UCUCACACAGAAUCGCACCCGU	43	WO 2009/012468
382	GAAGUUGUUCGUGGUGGAUUCG	44	WO 2009/012468
422a	ACUGGACUUAGGGUCAGAAGGC	45	US 12/391,028
378	ACUGGACUUGGAGUCAGAAGG	46	WO 2009/012468
424	CAGCAGCAAUUCAUGUUUUGAA	47	WO 2009/062169
483-3p	UCACUCCUCUCCUCCCGUCUU	48	WO 2009/012468
484	UCAGGCUCAGUCCCCUCCCGAU	49	WO 2009/012468

<u>miRNA</u>	<u>miRNA Sequence</u>	<u>SEQ ID NO:</u>	<u>Reference</u>
486-5p	UCCUGUACUGAGCUGCCCCGAG	50	WO 2009/012468
497	CAGCAGCACACUGUGGUUUGU	51	WO 2009/062169
499	UUAAGACUUGCAGUGAUGUUU	52	WO 2009/018492
542-5p	UCGGGGAUCAUCAUGUCACGAGA	53	WO 2009/012468
92a	UAUUGCACUUGUCCCGGCCUGU	54	WO 2009/012468
92b	UAUUGCACUCGUCCCGGCCUCC	55	WO 2009/012468
let-7a	UGAGGUAGUAGGUUGUAUAGUU	56	WO 2009/012468
let-7b	UGAGGUAGUAGGUUGUGUGGUU	57	WO 2009/012468
let-7c	UGAGGUAGUAGGUUGUAUGGUU	58	WO 2009/012468
let-7d	AGAGGUAGUAGGUUGCAUAGUU	59	WO 2009/012468
let-7e	UGAGGUAGGAGGUUGUAUAGUU	60	WO 2009/012468
let-7f	UGAGGUAGUAGAUUGUAUAGUU	61	WO 2009/012468
let-7g	UGAGGUAGUAGUUUGUACAGUU	62	WO 2009/012468
451	AAACCGUUACCAUACUGAGUU	63	US 61/286,546

[041] The polynucleotide may carry from one to five lipophilic groups conjugated as described herein. The lipophilic groups may be conjugated through a polynucleotide 5' or 3' position, and/or conjugated to the polynucleotide backbone via formation of, for example, a phosphate ester. In certain embodiments, the polynucleotide carries one or two lipophilic groups at the 5' and/or 3' end, conjugated as described herein.

Pharmaceutical Compositions and Delivery

[042] The conjugates of the invention may be formulated as a variety of pharmaceutical compositions. Pharmaceutical compositions will be prepared in a form appropriate for the intended application. Generally, this will entail preparing compositions that are essentially free of pyrogens, as well as other impurities that could be harmful to humans or animals. Exemplary delivery/formulation systems include colloidal dispersion systems, macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Commercially

available fat emulsions that are suitable for delivering the nucleic acids of the invention to cardiac and skeletal muscle tissues include Intralipid®, Liposyn®, Liposyn® II, Liposyn® III, Nutrilipid, and other similar lipid emulsions. A preferred colloidal system for use as a delivery vehicle in vivo is a liposome (i.e., an artificial membrane vesicle). The preparation and use of such systems is well known in the art. Exemplary formulations are also disclosed in US 5,981,505; US 6,217,900; US 6,383,512; US 5,783,565; US 7,202,227; US 6,379,965; US 6,127,170; US 5,837,533; US 6,747,014; and WO03/093449, which are hereby incorporated by reference in their entireties.

[043] The pharmaceutical compositions and formulations may employ appropriate salts and buffers to render delivery vehicles stable and allow for uptake by target cells. Aqueous compositions of the present invention comprise an effective amount of the delivery vehicle comprising the inhibitor polynucleotides or miRNA polynucleotide sequences (e.g. liposomes or other complexes), dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. The phrases "pharmaceutically acceptable" or "pharmacologically acceptable" refers to molecular entities and compositions that do not produce adverse, allergic, or other untoward reactions when administered to an animal or a human. As used herein, "pharmaceutically acceptable carrier" may include one or more solvents, buffers, solutions, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like acceptable for use in formulating pharmaceuticals, such as pharmaceuticals suitable for administration to humans. The use of such media and agents for pharmaceutically active substances is well known in the art. Supplementary active ingredients also can be incorporated into the compositions.

[044] Administration or delivery of the pharmaceutical compositions according to the present invention may be via any route so long as the target tissue is available via that route. For example, administration may be by intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection, or by direct injection into target tissue (e.g., cardiac tissue). Pharmaceutical compositions comprising miRNA inhibitors or expression constructs comprising miRNA sequences may also be administered by catheter systems or systems that isolate coronary circulation for delivering therapeutic agents to the heart. Various catheter systems for delivering therapeutic agents to the heart and coronary vasculature are known in the art. Some non-limiting examples of catheter-based delivery methods or coronary isolation methods suitable for use in the present invention are disclosed in U.S. Patent No. 6,416,510; U.S. Patent No. 6,716,196; U.S. Patent No. 6,953,466, WO 2005/082440, WO 2006/089340, U.S. Patent Publication No. 2007/0203445, U.S. Patent

Publication No. 2006/0148742, and U.S. Patent Publication No. 2007/0060907, which are all hereby incorporated by reference in their entireties.

[045] The compositions or formulations may also be administered parenterally or intraperitoneally. By way of illustration, solutions of the conjugates as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations generally contain a preservative to prevent the growth of microorganisms.

[046] The pharmaceutical forms suitable for injectable use or catheter delivery include, for example, sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. Generally, these preparations are sterile and fluid to the extent that easy injectability exists. Preparations should be stable under the conditions of manufacture and storage and should be preserved against the contaminating action of microorganisms, such as bacteria and fungi. Appropriate solvents or dispersion media may contain, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[047] Sterile injectable solutions may be prepared by incorporating the conjugates in an appropriate amount into a solvent along with any other ingredients (for example as enumerated above) as desired. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the desired other ingredients, e.g., as enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation include vacuum-drying and freeze-drying techniques which yield a powder of the

active ingredient(s) plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[048] Upon formulation, solutions are preferably administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations may easily be administered in a variety of dosage forms such as injectable solutions, drug release capsules and the like. For parenteral administration in an aqueous solution, for example, the solution generally is suitably buffered and the liquid diluent first rendered isotonic for example with sufficient saline or glucose. Such aqueous solutions may be used, for example, for intravenous, intramuscular, subcutaneous and intraperitoneal administration. Preferably, sterile aqueous media are employed as is known to those of skill in the art, particularly in light of the present disclosure. By way of illustration, a single dose may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

Methods of Treatment

[049] The invention provides a method for delivering polynucleotides to a mammalian cell, and methods for treating, ameliorating, or preventing the progression of a condition in a mammalian patient. The method generally comprises administering the polynucleotide as a conjugate with cholesterol or other lipophilic moiety as described herein. The polynucleotide may be a miRNA or a miRNA inhibitor (e.g., having a nucleotide sequence designed to inhibit expression or activity of a miRNA). Thus, the patient may have a condition associated with RNA expression, such as miRNA expression. Such conditions include, for example, cardiac hypertrophy, myocardial infarction, heart failure (e.g., congestive heart failure), vascular damage, restenosis, or pathologic cardiac fibrosis.

[050] miRNAs involved in conditions such as cardiac hypertrophy, myocardial infarction, heart failure (e.g., congestive heart failure), vascular damage, restenosis, and/or pathologic cardiac fibrosis, as well as sequences for targeting miRNA function are described in WO 2008/016924, WO 2009/062169, WO 2009/058818, WO 2009/018492, WO 2009/018493, WO 2009/012468, and WO 2007/070483, which are hereby incorporated by

reference in their entireties. Such miRNAs and sequences are further listed in Table 1 and described herein.

[051] The lipophilic polynucleotide conjugate may be delivered to the patient as a pharmaceutical composition or formulation as also described herein.

[052] In certain embodiments, the patient has one or more risk factors including, for example, long standing uncontrolled hypertension, uncorrected valvular disease, chronic angina, recent myocardial infarction, congenital predisposition to heart disease and pathological hypertrophy. Alternatively or in addition, the patient may have been diagnosed as having a genetic predisposition to, for example, cardiac hypertrophy, or may have a familial history of, for example, cardiac hypertrophy.

[053] In this aspect, the present invention may provide for an improved exercise tolerance, reduced hospitalization, better quality of life, decreased morbidity, and/or decreased mortality in a patient with heart failure or cardiac hypertrophy.

Methods of Synthesizing Conjugates

[054] The invention further provides methods of synthesizing polynucleotides having lipophilic moieties conjugated at the 5' and/or 3' end, or conjugated to the polynucleotide backbone, through a hydrocarbon linker. In this aspect, the invention involves coupling the lipophilic moiety with a linear or cyclic hydrocarbon having terminal functional groups to prepare a first intermediate, and preparing a second intermediate having functional groups suitable for incorporation into a polynucleotide chain during solid phase polynucleotide synthesis. Alternatively, the method involves synthesizing cholesterol conjugates with polynucleotides, after synthesis of the polynucleotide is complete, using for example, carbonyl addition-elimination/reductive amination, amidation, maleimide-thiol coupling, aqueous Diels-Alder and "Click" chemistries.

[055] In certain embodiments where the lipophilic moiety is an alcohol, such as a sterol, the method comprises conjugating the lipophilic moiety to a hydrocarbon (as described herein) that has terminal functional groups. This step will prepare a first intermediate, preferably an ether-linked intermediate. For example, in certain embodiments a sterol is etherized (through the sterol hydroxyl) to a C3-C15 hydrocarbon. The hydrocarbon in certain embodiments is a substantially linear C4-C8 (e.g., C4 or C6) hydrocarbon, which may be substituted, and which may have one, two, or three heteroatoms in the backbone independently selected from O, N and/or S. Alternatively, the hydrocarbon comprises a C5

to C8 cyclic hydrocarbon, e.g., cyclohexane, or a 5 to 8 membered cyclic group having one or two heteroatoms.

[056] The first or ether-linked intermediate is further reacted to prepare a suitable second intermediate that is compatible with solid phase nucleic acid synthesis. For example, the ether-linked intermediate may be further reacted through one or more steps to prepare a phosphoramidite intermediate (e.g., through an ether linkage) for converting to a phosphate ester during polynucleotide synthesis. Alternatively, or in addition, the ether-linked intermediate may be further reacted through one or more steps to prepare a suitable terminal group for coupling (e.g. via amidation) to a suitable support to form the 3' end of a polynucleotide chain. Exemplary reactions for preparing 5' and 3' conjugates are illustrated in **Figure 2**.

[057] In certain embodiments involving the preparation of a phosphoramidite intermediate (shown as **6** in **Fig. 2**), the terminal alcohol of the ether-linked intermediate (**2**) is reacted to prepare the phosphoramidite by reaction with a phosphitylating agent, such as 2-cyanoethyl N,N-diisopropylchlorophosphoramidite. These embodiments may be employed particularly where there is no need for trityl monitoring or additional amidite couplings.

[058] Alternatively, other intermediates having a suitable hydroxyl for phosphoramidite preparation may be synthesized from (**2**). For example, compound (**2**) may be reacted to form two differentiated hydroxyl groups separated by at least one carbon, but preferably 2 or more carbons. An exemplary series of reactions for converting compound (**2**) to a compound having differentiated hydroxyls (**5**) is shown in **Fig. 2**. Cholesterol tosylate (**1**), formed by reacting cholesterol with tosyl chloride, undergoes an S_N1 reaction with 1,6-hexanediol in refluxing 1,4-dioxane. Neighboring group participation by the alkene at the 5-position provides the 3 β -isomer exclusively. The resultant (Cholest-5-en-3 β -oxy)hexan-6-ol (**2**) is converted to the mesylate (**3**) which is subsequently used to alkylate solketal. The crude solketal derivative is deprotected by treatment with trifluoroacetic acid in wet THF to give 3-(6-(Cholest-5-en-3 β -oxy)hexyloxy)propane-1,2-diol (**4**). The primary alcohol of the diol is selectively protected as the 4,4'-Dimethoxytrityl ether (**5**) using standard methods. The remaining secondary alcohol can be converted to the phosphoramidite (**6**) by reacting with a phosphitylating agent, such as 2-Cyanoethyl N,N-diisopropylchlorophosphoramidite, and subsequently used as a standard amidite monomer in the oligonucleotide synthesis cycle. Alternatively, the secondary alcohol can be converted to the succinate ester (**7**) by treatment with succinic anhydride in the presence of TEA. The resultant carboxylate can then be coupled, via HBTU, to excess amine functionalized controlled pore glass (CPG).

The excess amines and any free surface hydroxides of the CPG are rendered unreactive by "capping" with acetic anhydride. The cholesterol functionalized CPG is suitable for 3'-cholesterol modified oligonucleotide synthesis. The CPG can be used just as any standard nucleoside loaded CPG is used in automated oligonucleotide synthesis.

[059] Thus, in these embodiments the second intermediate is prepared from a compound having at least one 2°-alcohol and at least one 1°-alcohol. In accordance with these embodiments, during polynucleotide synthesis, one hydroxyl group forms a phosphodiester linkage with the nucleotide on the 5'-end of the chain. In the case of a 5' conjugate, two hydroxyl groups would allow for trityl monitoring of the reaction on an automated synthesizer and/or further chain elongation past the cholesterol conjugate, if desired.

[060] In certain embodiments involving the preparation of a second intermediate for coupling to a solid support (shown as **7** in **Fig. 2**), the alcohol of the ether-linked intermediate (**2**) is reacted in several steps to prepare a succinate intermediate (or other intermediate cleavable from a solid support). Intermediates for preparing 3' conjugates would, as above, preferably have two hydroxyl groups, preferably differentiated. For example, the intermediate for coupling to a solid support may be prepared from a compound having at least one 2° hydroxyl and at least one 1° hydroxyl. An exemplary series of reactions for converting compound **2** to a compound having differentiated hydroxyls (**5**) is shown in **Fig. 2**. In these embodiments, during polynucleotide synthesis the nucleotide chain will grow off the DMTr-protected 1°-alcohol and the conjugate moiety will be anchored to a solid support via some type of cleavable linkage on the 2°-hydroxyl group (e.g., succinate).

[061] Alternative embodiments of this invention includes the addition of a derivative of ether-linked intermediate (**2**) after oligonucleotide synthesis is complete. See **Fig. 4**. This can be accomplished, for example, via reactions as carbonyl addition-elimination/reductive amination, amidation, maleimide-thiol coupling, aqueous Diels-Alder and "Click" chemistries. These methods provide equivalent cholesterol ether constructs without the need to synthesize cholesterol-amidites. For example, a cholest-5-en-3 β -oxyalkanol, such as compound (**2**), can be readily modified by one of skill in the art to give a terminal aldehyde, carboxylate, amine, maleimide, thiol, diene, azide, alkyne or other reactive moiety. These compounds can be further reacted with an oligonucleotide suitably modified with a complimentary functional group that forms a covalent bond with the reactive moiety of the terminally functionalized cholest-5-en-3 β -oxyalkane. The product is a cholesterol conjugate that still retains a relatively non-polar, substantially linear alkyloxy linkage to the cholesterol

moiety, and maintains a 3 or greater atom distance between the lipophilic moiety and any group containing an exchangeable proton or other substantially polar functional group. These so-called "post-synthetic bioconjugations" can be performed while the protected synthetic oligonucleotide chain is still attached to the solid phase, or after deprotection when the synthetic oligonucleotide is in solution.

[062] Techniques for bioconjugation are well known, and are described, for example, in Hermanson GT, *Bioconjugate Techniques*, Second Edition, Academic Press; 2nd edition (April 1, 2008).

EXAMPLES

[063] Nucleic acid (NA)-Chol conjugates have been derived from amines reacted with readily available cholesterol chloroformate to give a carbamate linkage at the 3 β position of the cholestene ring structure. This system places a polar group with an exchangeable hydrogen very close to the lipophilic ring structure. This can limit the lipophilicity of the conjugate and the ability of conjugate to fully intercalate into any lipid bilayer or associate with other lipophilic moieties. See **Figure 1**. The conjugates as described herein avoid the use of linker groups that place polar groups with exchangeable protons next to the cholestene ring structure (**Fig. 1**).

Example 1: Synthesis of Exemplary Conjugates

[064] An ether linked cholesterol, such as Chol-O-HEX-OH (shown as **2** in **Figure 2**) may be used to create NA conjugates. **Figure 2** depicts the synthesis of a linker amidite, **6**, and succinate, **7** (herein referred to as "C6-Chol"). They may be used, respectively, as a 5'-modification (Compound **6**) on a common, chemically synthesized oligonucleotide made with normally available synthesis reagents and as a 3'-modification (Compound **7**) when conjugated via amidation to any commercially available amino support used for oligonucleotide synthesis. An amidite may alternatively be prepared from an alcohol such as **2** for 5'-terminally conjugated oligonucleotides, for example, when there is no need for trityl monitoring or additional amidite couplings.

[065] Generally, the hydrocarbon spacer contains a hydroxyl group at or near the terminus to form a phosphate ester. Preferably, there are two hydroxyl groups that would be separated by at least one carbon, but preferably 2 or more carbons. For example, the hydroxyls may be differentiated where one is a 2°-alcohol and the other is a 1°-alcohol. At least one hydroxyl group is necessary to form a phosphodiester linkage with the nucleotide on the 5'-end of the chain. In the case of a 5'-cholesterol conjugate, two hydroxyl groups

would allow for trityl monitoring of the reaction on an automated synthesizer and/or further chain elongation past the cholesterol conjugate. The 4,4'-dimethoxytrityl (DMTr) would be placed, for example, on one hydroxyl and the phosphoramidite (i.e. the phosphodiester precursor) would be on the remaining hydroxyl, which provides for more efficient phosphoramidite coupling.

[066] 3'-Cholesterol conjugates would require two hydroxyl groups, preferably differentiated (i.e. one is 2° and one is 1°) where the nucleotide chain would grow off the DMTr-protected 1°-alcohol and the where the conjugate moiety would be anchored to a solid support via some type of cleavable linkage on the 2°-hydroxyl group. The 5'-conjugate would have the phosphate on the 2° position.

[067] An exemplary synthesis of Chol-O-Hex intermediates for conjugation during solid phase polynucleotide synthesis are described below.

Cholest-5-en-3β-tosylate

[068] Cholesterol (25.0g, 64.7 mmol) is weighed into a 500 mL round bottomed flask and dissolved in pyridine (200 mL) to give a colorless solution that is cooled to 0 °C. Tosyl chloride (24.65g, 129 mmol) is weighed into a 100 mL round bottomed flask and dissolved in pyridine (40 mL) to give a slightly yellow solution. The tosyl chloride solution is then added, at once, to the stirring cholesterol solution and the resultant reaction mixture is allowed to stir overnight while coming to room temperature. The reaction mixture is concentrated on a rotary evaporator to give a white solid that is taken up in minimum amount of chloroform (40 mL) and precipitated via addition of methanol (500 mL). The resultant white solid is filtered and washed with methanol (500 mL) and acetonitrile (200 mL). The solid is transferred to a 1L round bottomed flask and dried overnight under high vacuum to give 34.2g (98%) of a white powder. ¹H (400 MHz, CDCl₃): δ0.67 (3H, s); 0.80-1.75 (33H, m); 1.76-2.03 (5H, m); 2.20-2.30 (1H, m); 2.44 (4H, m); 4.25-4.35 (1H, m); 5.29 (1H, d, *J* = 5 Hz); 7.32 (2H, d, *J* = 8 Hz); 7.79 (2H, d, *J* = 8 Hz).

Cholest-5-en-3β-oxyhexan-6-ol

[069] Cholest-5-en-3β-tosylate (15.0g, 27.7 mmol) and 1,6-hexanediol (65.5 g, 555 mmol) is weighed into a 500 mL round bottomed flask and dissolved in 1,4-dioxane (300 mL) to give a colorless suspension. The flask is fitted with an efficient reflux condenser and the reaction mixture is heated to reflux in a 110 °C oil bath for 17 hr. The reaction mixture is concentrated in vacuo and the residue is suspended in DI water (500 mL). The aqueous phase is extracted with EtOAc (2 x 150 mL) and the combined organic phases are washed

with water (2 x 300 mL) and brine (1 x 150 mL). The organic phase is dried over sodium sulfate, filtered, concentrated and dried overnight under high vacuum to give 12.85g (95%) of a light yellow wax. ^1H (400 MHz, CDCl_3): δ 0.67 (3H, s); 0.80-1.65 (42H, m); 1.75-2.05 (5H, m); 2.12-2.25 (1H, m); 2.30-2.40 (1H, m); 3.05-3.17 (1H, m); 3.45 (2H, dt, J = 6.5, 1.6 Hz); 3.57 (2H, t, J = 6.6 Hz); 5.34 (1H, d, J = 5.3 Hz).

Cholest-5-en-3 β -oxyhexan-6-mesylate

[070] Cholest-5-en-3 β -oxyhexan-6-ol (12.84g, 26.4 mmol) is weighed into a 500 mL round bottomed flask. The flask is charged with dichloromethane (100 mL) and triethylamine (7.35 mL, 52.7 mmol), flushed with argon and cooled to 0 °C. Mesyl chloride (2.55 mL, 32.96 mmol) is added dropwise over 5 minutes to the stirring solution and the mixture is allowed to stir for an additional 1h at 0 °C. Saturated sodium bicarbonate solution (150 mL) was added to the mixture which was then transferred to a separatory funnel. The aqueous phase was discarded and the organic phase washed with fresh saturated sodium bicarbonate solution (1 x 150 mL) and brine (1 x 50 mL). The organic phase is dried over Na_2SO_4 , filtered and concentrated to dryness to give 14.8g (99%) of a yellow wax. ^1H (400 MHz, CDCl_3): δ 0.67 (3H, s); 0.80-1.65 (41H, m); 1.70-2.05 (5H, m); 2.12-2.23 (1H, m); 2.30-2.37 (1H, m); 3.00 (3H, s); 3.05-3.17 (1H, m); 3.45 (2H, dt, J = 6.5, 1.4 Hz); 4.22 (2H, t, J = 6.6 Hz); 5.34 (1H, d, J = 5.2 Hz).

3-(6-(Cholest-5-en-3 β -oxy)hexyloxy)propane-1,2-diol

[071] Solketal (2.92g, 27.5 mmol) was weighed into a 200 mL round bottomed flask. The flask was flushed with argon and charged with toluene (40 mL). Sodium hydride, 60% dispersion in oil (3.14 g, 131 mmol) was added, at once, to the stirring solution and the mixture was allowed to stir for 30 minutes at room temperature. Cholest-5-en-3 β -oxyhexan-6-mesylate (14.8g, 26.2 mmol) was dissolved in toluene (40 mL) and the solution was added slowly to the alkoxide solution. The flask was fitted with a reflux condenser and the apparatus flushed with argon. The reaction mixture was heated to reflux and allowed to stir for 17 hrs. The reaction mixture was cooled to room temperature and quenched with a solution of ethanol in ethyl acetate added dropwise to the vigorously stirred reaction mixture over 15 minutes. The reaction mixture was further diluted with ethyl acetate and washed with 10% aqueous sodium carbonate solution (2 x 150 mL). The organic phase was dried over Na_2SO_4 , filtered and concentrated.

[072] The residue was re-dissolved in a 4:4:1 ratio of tetrahydrofuran:trifluoroacetic:water (18mL), then set to stir for 2 hours at room temperature.

TLC (15% acetone/hexanes) of the reaction mixture revealed that the deprotection of the desired compound in the crude mixture was complete. The reaction mixture was evaporated to dryness, applied to a 100g Biotage SNAP silica column and eluted with an ethyl acetate/hexanes gradient (0-35% over 2 L @ ~50 mL/min) to give 4.90 g (33% over two steps) of a colorless wax. ¹H (400 MHz, CDCl₃): δ 0.67 (3H, s); 0.80-1.65 (41H, m); 1.75-2.05 (5H, m); 2.12-2.40 (4H, m); 3.07-3.17 (1H, m); 3.40-3.57 (6H, m); 3.63 (1H, dd, J=11.4, 5.2); 3.71 (1H, dd, J=11.4, 3.9); 3.81-3.89 (1H, m); 5.34 (1H, d, J=5.2 Hz).

1-(4,4'-Dimethoxytrityloxy)-3-(6-(Cholest-5-en-3β-oxy)hexyloxy)propan-2-ol

[073] 3-(6-(Cholest-5-en-3β-oxy)hexyloxy)propane-1,2-diol (4.75 g, 8.47 mmol) was weighed into a 200 mL round bottomed flask with a stir bar. The flask was flushed with argon, charged with pyridine (30 mL) and cooled to 0 °C with stirring. DMTr-Cl (3.01 g, 8.89 mmol) was weighed into a 20 mL scintillation vial and dissolved in pyridine (17 mL). The DMTr-Cl solution was added, dropwise, to the stirring diol solution over 20 minutes. The reaction was allowed to stir overnight while coming to room temperature. The reaction was quenched by adding anhydrous methanol (1 mL) and allowing the reaction to stir for 30 minutes. Saturated sodium bicarbonate solution (50 mL) was added to the reaction mixture, which was stirred until CO₂ evolution ceased. The mixture was concentrated to dryness and partitioned with water (100 mL) and ethyl acetate (100 mL). The organic phase was dried over Na₂SO₄, filtered and concentrated. The residue was applied to a 100g Biotage SNAP column and eluted with an ethyl acetate/hexanes gradient (10-20% over 1.5 L @ ~50 mL/min) to give 3.50g (47.9%) of a colorless, sticky oil. ¹H (400 MHz, CDCl₃): δ 0.67 (3H, s); 0.87 (6H, dd, J=6.6, 1.8 Hz); 0.91 (3H, d, J=6.6 Hz); 0.93-1.60 (32H, m); 1.75-2.05 (5H, m); 2.15-2.25 (1H, m); 2.32-2.41 (1H, m); 2.42 (1H, d, J=4.6 Hz); 3.07-3.22 (2H, m); 3.39-3.57 (6H, m); 3.79 (6H, s); 3.80 (s, 1H); 3.90-3.98 (1H, m); 5.34 (1H, d, J=5.3 Hz); 6.78-6.85 (m, 4H); 7.14-7.25 (m, 1H); 7.26-7.34 (m, 6H); 7.40-7.44 (m, 2H).

4-(4,4'-Dimethoxytrityloxy)-3-(6-(Cholest-5-en-3β-oxy)hexyloxy)propan-2-yloxy)-4-oxobutanoic acid, TEA salt

[074] 1-(4,4'-Dimethoxytrityloxy)-3-(6-(Cholest-5-en-3β-oxy)hexyloxy)propan-2-ol (1.75 g, 2.02 mmol), succinic anhydride (0.812 g, 8.10 mmol) and 4-(dimethylamino)pyridine (25 mg, 0.20 mmol) was added to a 100 mL round bottomed flask with a stir bar. The flask was charged with pyridine (10 mL). The reaction mixture was stirred for 8 hours at 80 °C and 16 h at 50 °C. The reaction mixture was concentrated in vacuo and co-evaporated with EtOH (3 x 20 mL) to remove residual pyridine. The residue was applied to a 50 g Biotage SNAP column and eluted @ ~40 mL/min with 1L of 60/30/10 EtOAc/Hex/MeOH with 3% TEA

followed by 500 mL of 90/10 DCM/MeOH with 3% TEA to give 1.31g (61%) of a colorless wax. ^1H (400 MHz, CDCl_3): δ 0.67 (3H, s); 0.86 (6H, dd, $J=6.6$, 1.8 Hz); 0.91 (3H, d, $J=6.6$ Hz); 0.93-1.60 (41H, m); 1.75-2.05 (5H, m); 2.15-2.25 (1H, m); 2.32-2.41 (1H, m); 2.51-2.72 (4H, m); 2.97 (6H, q, $J=7.3$ Hz); 3.07-3.25 (2H, m); 3.31-3.48 (4H, m); 3.52-3.61 (2H, m); 3.78 (6H, s); 5.19 (1H, quintet, $J=5.1$ Hz); 5.35 (1H, d, $J=5.4$ Hz); 6.78-6.85 (m, 4H); 7.14-7.25 (m, 1H); 7.26-7.34 (m, 6H); 7.40-7.44 (m, 2H).

"Chol-O-Hex-LCAA-CPG"

[075] 4-(4,4'-Dimethoxytrityloxy)-3-(6-(Cholest-5-en-3 β -oxy)hexyloxy)propan-2-yloxy)-4-oxobutanoic acid, TEA salt (1.00 g, 0.952 mmol) is weighed into a 20 mL scintillation vial. The vial is charged with 6 mL of 2:1 ACN:DCM. HBTU (0.343 g, 0.904 mmol) is added to the succinate followed by triethylamine (0.25 mL). In a 200 mL round bottomed flask is weighed LCAA-CPG (12.56 g) with a loading of 132 $\mu\text{mol/g}$ amine. The CPG is suspended in 75 mL of 2:1 ACN:DCM. The succinate/HBTU mixture is added, at once, to the LCAA-CPG suspension, the flask is sealed and the combined mixture is agitated on an incubated orbital shaker at 25 $^\circ\text{C}$ @ 250 rpm for 4 hours. The CPG is filtered and washed with 200 mL aliquots of ACN, DCM, DMF, water, methanol, ACN and DCM. The CPG is air dried overnight. The CPG is then transferred to a clean 200 mL round bottomed flask. The flask is charged with 40 mL of CAP A solution and 40 mL of CAP B solution, septum sealed and the contents agitated at 25 $^\circ\text{C}$ @ 250 rpm for 3 hours. The CPG is then filtered and washed thoroughly with 200 mL aliquots of ACN, DCM, MeOH, H₂O, MeOH, DMF, ACN, DCM. The CPG is air dried overnight.

[076] The CPG loading is measured by a modification of the method described by Pon, R.T., *Current Protocols in Nucleic Acid Chemistry*. **2000** 3.2.1-3.2.23. CPG (4.24 mg) was weighed into a 20 mL scintillation vial. 5% Trifluoroacetic acid in dichloroethane was added and the vial was capped and shaken for 1 minute. The absorbance of the solution was measured at 503 nm via a ThermoSpectronic Genesys10 UV/Vis spectrophotometer to be 2.114. The loading was calculated by the formula:

[077]
$$[(\text{volume, mL} \times \text{absorbance}) / 76 \text{ mLcm}^{-1}\mu\text{mol}^{-1}] \times [1000 / \text{mass, mg}] = \text{loading in } \mu\text{mol/g}$$
 to give: $[(10 \text{ mL} \times 2.114) / 76 \text{ mLcm}^{-1}\mu\text{mol}^{-1}] \times [1000 / 4.24 \text{ mg}] = 66 \mu\text{mol/g}$.

"Chol-O-Hex-DMTr Amidite"

[078] 1-(4,4'-Dimethoxytrityloxy)-3-(6-(Cholest-5-en-3 β -oxy)hexyloxy)propan-2-ol (1.68 g, 1.95 mmol) is weighed into a 200 mL round bottomed flask with a stir bar. The flask is flushed with argon, septum sealed and charged with DCM (10 mL) and N,N-

diisopropylethylamine (0.68 mL, 3.89 mmol). 2-Cyanoethyl N,N-diisopropylchlorophosphoramidite (0.48 g, 2.0 mmol) is added to the stirring reaction dropwise and the reaction is allowed to stir for 17 hrs. The reaction mixture is diluted with DCM 50 mL and washed with saturated bicarbonate solution (2 x 50 mL) and brine (1 x 50 mL). The organic phase is dried over Na₂SO₄, filtered and concentrated. The residue is applied to a 25 g Biotage SNAP column and eluted with an ethyl acetate/hexanes gradient (5-10% over 750 mL @ 25 mL/min) to give 1.70g (82.1%) of a white foam. ¹H (400 MHz, DMSO-d₆): δ 0.62 (3H, s); 0.80-1.57 (53H, m); 1.70-1.95 (5H, m); 1.96-2.02 (1H, m); 2.21-2.26 (1H, m); 2.57-2.72 (2H, d, J=4.6 Hz); 2.97-3.80 (m, 20H); 3.97 (1H, quintuplet, J=5.3 Hz); 5.26 (1H, d, J=4.6 Hz); 6.78-6.86 (m, 4H); 7.13-7.29 (m, 7H); 7.34-7.39 (m, 2H).

Example 2: Synthesis of 1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3-oxy)dodecan-3-ol Functionalized CPG

[079] Figure 3 depicts the synthesis of a linker structure (III) described herein that can be used as a polynucleotide 3'-modification. The synthesis steps are described in more detail below.

10-(Cholest-5-en-3β-oxy)decan-1-ol

[080] In a 500 mL round-bottomed flask was cholesteryl tosylate (10.0 g, 18.5 mmol) and 1,10-Decanediol (48.3 g, 277 mmol) suspended in Dioxane (200 ml) to give a white suspension. The flask was fitted with a temperature sensor and reflux condenser. The mixture was heated to reflux and allowed to stir at reflux overnight. The reaction mixture was cooled to room temperature and the solids were filtered off. The solution was concentrated *in vacuo*. The mixture was re-dissolved in methanol (250 mL) and water (ca. 40 mL) was added to precipitate out the crude product. The precipitate was filtered and dried *in vacuo*. The residue was further purified via column chromatography on a 100g Biotage SNAP column eluting over 2L of a 0-25% EtOAc in Hexanes gradient. Pure fractions were combined and concentrated to give 7.85g (82%) of a white solid. ¹H NMR (300 MHz, CDCl₃) δ 0.69 (3H, s); 0.85-1.66 (50H, m); 1.75-2.09 (5H, m); 2.14-2.29 (1H, m); 2.32-2.43 (1H, m); 3.14 (1H, tt, J=11.2, 4.4Hz); 3.46 (2H, dt, J=6.9, 0.4 Hz); 3.65 (2H, t, J=6.6 Hz); 5.36 (1H, d, J=5.3Hz).

10-(Cholest-5-en-3β-oxy)decanal

[081] In a 100 mL round-bottomed flask was solid supported 2,2,6,6-Tetramethyl Piperidinyloxy, Free Radical (0.5 g, 1.1 mmol/g, 0.553 mmol) and 10-(Cholest-5-en-3β-oxy)decan-1-ol (3.00 g, 5.53 mmol) suspended in dichloromethane (20 ml) to give a

colorless suspension. Iodobenzene diacetate (1.958 g, 6.08 mmol) was added to the mixture, which was allowed to stir overnight at room temperature. The reaction mixture was filtered and the solid support washed with DCM. The combined filtrate was washed with saturated sodium thiosulfate (2 x 20 mL) and brine (1 x 20 mL). The organic phase was dried over Na₂SO₄, filtered and concentrated. The resultant oil was purified via column chromatography with 100% DCM as mobile phase to give 2.70g (90%) of a colorless wax. ¹H NMR (300MHz, CDCl₃) δ 0.69 (3H, s); 0.84-1.74 (47 H, m); 1.76-2.09 (5H, m); 2.12-2.26 (1H, m); 2.32-2.40 (1H, m); 2.44 (2H, dt, J=7.4, 1.9 Hz); 3.14 (1H, tt, J=11.2, 4.4 Hz); 3.46 (2H, t, J=6.9 Hz); 5.35 (1H, d, J=5.3Hz); 9.78 (1H, t, J=1.9Hz).

12-(Cholest-5-en-3β-oxy)-3-hydroxydodecanal

[082] In a dry 200 mL round-bottomed flask was 10-(Cholest-5-en-3β-oxy)decanal (2.65 g, 4.90 mmol) dissolved in THF (10 ml), under argon to give a colorless solution. A 0.5 M THF solution of ((1,3-dioxolan-2-yl)methyl)magnesium bromide (9.80 ml, 4.90 mmol) was added dropwise, and the resulting solution was allowed to stir for 4h at room temperature. The resultant solution was quenched by slowly adding 100mL of NaHCO₃ solution and extracting with EtOAc (3 x 100 mL). The combined organic phase was washed with brine (1 x 100mL), dried over sodium sulfate, filtered and concentrated in a 500 mL round bottomed flask. Trimethyl orthoformate (10.83 ml, 98 mmol), PPTS (0.246 g, 0.980 mmol) and methanol (10 ml) were added to the flask and the contents brought to reflux for 2 h. The solution was cooled to room temperature and evaporated to dryness. The residue was then taken up in Acetone (20 ml) and 1.0N HYDROCHLORIC ACID (4.90 ml, 4.90 mmol) and allowed to stir overnight. The reaction mixture was diluted to 200 mL with water and extracted with ethyl acetate (3 x 100 mL). The organic phase was combined, washed with brine (1 x 50 mL), dried over Na₂SO₄, filtered and concentrated. The resultant wax was purified via column chromatography on a 40g Biotage SNAP column eluting with a 0-50% gradient over 1L. The pure fractions were combined and evaporated to dryness to give 12-(Cholest-5-en-3β-oxy)-3-hydroxydodecanal (1.97 g, 3.37 mmol, 68.7 % yield) as a colorless wax. ¹H NMR (300MHz, CDCl₃) δ 0.69 (3H, s); 0.85-1.74 (49H, m); 1.75-2.13 (7H, m); 2.14-2.28 (1H, m); 2.33-2.44 (1H, m); 3.14 (1H, tt, J=11.2, 4.4Hz); 3.46 (2H, t, J=6.8Hz); 3.93-4.10 (1H, m); 5.36 (1H, d, J=5.2 Hz); 5.42-5.55 (1H, m); 9.78 (1H, s).

12-(Cholest-5-en-3β-oxy)dodecan-1,3-diol

[083] In a 100 mL round-bottomed flask was 12-(Cholest-5-en-3β-oxy)-3-hydroxydodecanal (1.90 g, 3.25 mmol) dissolved in 1,4-Dioxane (10 ml) to give a colorless

solution. MeOH (5 ml) was added slowly and only enough was added to keep the aldehyde in solution. SODIUM BOROHYDRIDE (0.184 g, 4.87 mmol) was added to the stirring solution which was then allowed to stir for 2 hours at room temperature. The reaction mixture was quenched by addition of 1N HCl (2 mL). The mixture was diluted to 100 mL with water and extracted with ethyl acetate (3 x 50 mL). The organic phase was combined, washed with brine (1 x 50 mL), dried over Na₂SO₄, filtered and concentrated to give 12-(Cholest-5-en-3 β -oxy)dodecan-1,3-diol (1.85 g, 3.15 mmol, 97 % yield) as a colorless wax. ¹H NMR (300 MHz, CDCl₃) δ 0.69 (3H, s); 0.83-2.07 (58H, m); 2.13-2.28 (1H, m); 2.32-2.43 (1H, m); 3.14 (1H, tt, *J*=11.2, 4.4 Hz); 3.46 (2H, t, *J*=6.8 Hz); 3.60-4.19 (3H, m); 5.36 (1H, d, *J*=5.2 Hz).

1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3 β -oxy)dodecan-3-ol

[084] In a 100 mL round-bottomed flask was 12-(Cholest-5-en-3 β -oxy)dodecan-1,3-diol (1.80 g, 3.07 mmol) dissolved in Pyridine (20 ml) to give a colorless solution. The mixture was cooled to 0 °C, under argon with stirring. DMTr-Cl (1.091 g, 3.22 mmol) was weighed into a septum sealed scintillation vial and dissolved in Pyridine (10 ml) to give a yellow solution. The DMTr-Cl solution was added dropwise to the diol. The reaction mixture was stirred overnight while coming to room temperature. Methanol (2 mL) was added and the mixture allowed to stir an additional 15 minutes. The reaction mixture was diluted to 200 mL with saturated NaHCO₃ solution and extracted with ethyl acetate (3 x 50 mL). The organic phase was combined, washed with brine (1 x 50 mL), dried over Na₂SO₄, filtered and concentrated. The resultant oil was purified via a 40g Biotage SNAP column eluted with a 0-50% EtOAc in Hexanes gradient over 1L. The pure fractions were combined and evaporated to give 1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3 β -oxy)dodecan-3-ol (2.60 g, 2.92 mmol, 95 % yield) as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 0.69 (3H, s); 0.83-2.09 (57H, m); 2.14-2.29 (1H, m); 2.33-2.44 (1H, m); 3.07-3.22 (3H, m); 3.47 (2H, t, *J*=6.9Hz); 3.53-3.66 (1H, m); 3.81 (6H, s); 5.37 (1H, d, *J*=5.2 Hz); 6.80-6.88 (4H, m); 7.20-7.35 (7H, m); 7.40-7.49 (2H, m).

4-((1-(bis(4-methoxyphenyl)(phenyl)methoxy)-12-(Cholest-5-en-3 β -oxy)dodecan-3-yl)oxy)-4-oxobutanoic acid

[085] In a 20 mL scintillation vial was 1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3 β -oxy)dodecan-3-ol (0.5 g, 0.562 mmol) and succinic anhydride (0.113 g, 1.124 mmol) in DCM (10 ml) to give a colorless solution. TEA (0.313 ml, 2.249 mmol) was added and the mixture set to stir for 17 hours. The reaction was judged complete by TLC

(25% EtOAc/Hex, PMA visualization). The reaction mixture was washed with 5% NaHPO₄ solution (3 x 15 mL) and brine (1 x 15 mL). The organic phase was dried over sodium sulfate, filtered and concentrated to dryness for use without further purification. ¹H NMR (300 MHz, CDCl₃) δ 0.70 (3H, s); 0.83-1.74 (49H, m); 1.77-2.12 (5H, m); 2.14-2.29 (1H, m); 2.33-2.44 (1H, m); 2.53-2.72 (4H, m); 3.00-3.09 (2H, m); 3.16 (1H, tt, *J*=11.2, 4.4 Hz); 3.48 (2H, t, *J*=6.8 Hz); 3.80 (6H, s); 4.85-5.00 (1H, m); 5.36 (1H, d, *J*=5.1 Hz); 6.78-6.90 (4H, m), 7.15-7.38 (7H, m); 7.40-7.49 (2H, m); 9.29 (1H, bs).

1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3β-oxy)dodecan-3-ol

Functionalized CPG

[086] 4-((1-(bis(4-methoxyphenyl)(phenyl)methoxy)-12-(Cholest-5-en-3β-oxy)dodecan-3-yl)oxy)-4-oxobutanoic acid (0.35 g, 0.36 mmol) is weighed into a 20 mL scintillation vial. The vial is charged with 6 mL of 2:1 ACN:DCM. HBTU (0.137 g, 0.35 mmol) is added to the succinate followed by triethylamine (0.25 mL). In a 100 mL round bottomed flask is weighed LCAA-CPG (5.00 g) with a loading of 132 μmol/g amine. The CPG is suspended in 30 mL of 2:1 ACN:DCM. The succinate/HBTU mixture is added, at once, to the LCAA-CPG suspension, the flask is sealed and the combined mixture is agitated on an incubated orbital shaker at 25 °C @ 250 rpm for 4 hours. The CPG is filtered and washed with 200 mL aliquots of ACN, DCM, DMF, water, methanol, ACN and DCM. The CPG is air dried overnight. The CPG is then transferred to a clean 200 mL round bottomed flask. The flask is charged with 40 mL of CAP A solution and 40 mL of CAP B solution, septum sealed and the contents agitated at 25 °C @ 250 rpm for 3 hours. The CPG is then filtered and washed thoroughly with 200 mL aliquots of ACN, DCM, MeOH, H₂O, MeOH, DMF, ACN, DCM. The CPG is air dried overnight.

[087] The CPG loading is measured by a modification of the method described by Pon, R.T., *Current Protocols in Nucleic Acid Chemistry*. **2000** 3.2.1-3.2.23. CPG (4.24 mg) was weighed into a 20 mL scintillation vial. 5% Trifluoroacetic acid in dichloroethane was added and the vial was capped and shaken for 1 minute. The absorbance of the solution was measured at 503 nm via a ThermoSpectronic Genesys10 UV/Vis spectrophotometer to be 2.114. The loading was calculated by the formula:

$$[(\text{volume, mL} \times \text{absorbance}) / 76 \text{ mLcm}^{-1}\mu\text{mol}^{-1}] \times [1000 / \text{mass, mg}] = \text{loading in } \mu\text{mol/g}$$

to give:

$$[(10 \text{ mL} \times 2.260) / 76 \text{ mLcm}^{-1}\mu\text{mol}^{-1}] \times [1000 / 5.12 \text{ mg}] = 58 \mu\text{mol/g}$$


Example 3: Post-synthetic addition of cholesterol

[088] The post synthetic addition of cholesterol via such reactions as carbonyl addition-elimination/reductive amination, amidation, maleimide-thiol coupling, aqueous Diels-Alder and "Click" would also provide equivalent cholesterol ether constructs without the need to synthesize cholesterol-amidites. For example, cholesterol tosylate may be alkylated to give a Cholest-5-en-3 β -oxyalkanol that can be further modified to give a terminal aldehyde, carboxylate, amine, maleimide, thiol, diene, azide, alkyne or other reactive moiety that could be further reacted with a suitably modified oligonucleotide to give a cholesterol conjugate that still retains a substantially non-polar, substantially linear alkyloxy linkage to the cholesterol moiety, and that maintains a 3 or greater atom distance between the lipophilic moiety and any group containing an exchangeable proton or other substantially polar functional group. See **Fig. 4**.

[089] Cholest-5-en-3-oxyalkanol (**A**) can be modified to give functional groups amenable to common bioconjugation techniques such as amidation (**B**), Diels-Alder bioconjugation (**C**) and "Click" bioconjugation (**D**) when the oligonucleotide is functionalized with the complementary functional group.

Example 4: In vivo cardiac knockdown of miR-208

[090] For all chemistries indicated in Table 2, 4 adult male C57Bl6 mice were injected via their tail vein with 2 x 80 mg/kg of either: the anti-miR sequence or the mismatch ("mm" in Figs. 5 and 6) sequence using the same chemistry dissolved in saline, and using saline alone ("sal" in Figs. 5 and 6) as a control (Table 2). Linker 1 in Table 2 corresponds to linker structures (XI) and (XIII) described herein. Linker 2 in Table 2 corresponds to linker structures (X) and (XII) described herein. For comparison, the linker of structure (XIV) was used as a control.



[091] Further, phosphorothioate (PS) refers to a substitution pattern in which the last two internucleotide bonds on the 5'-end are phosphorothioate linkages, and the first 5 internucleotide phosphate bonds on the 3'-end are phosphorothioate linkages (including the phosphate linkage between the cholesterol linker and oligonucleotide).

Table 2

Polynucleotide	Phosphorothioate (Ps)	Cholesterol	Side of conjugation	Linker
antagomiR	Yes	Yes	3'	Structure (XIV)
No PS	No	No	-	-
PS	Yes	No	-	-
3' chol linker #1, no PS	No	Yes	3'	linker #1
3' chol linker #2, no PS	No	Yes	3'	linker #2
5' chol linker #1	Yes	Yes	5'	linker #1
3' chol linker #1	Yes	Yes	3'	linker #1
5' chol linker #2	Yes	Yes	5'	linker #2
3' chol linker #2	Yes	Yes	3'	linker #2

[092] Three days after the last injection the animals were sacrificed and heart, liver, lungs and kidneys were collected. RNA was extracted from cardiac tissue to determine the effect on cardiac expression of miR-208 by Northern blot analysis (**Fig. 5**) and quantified using realtime PCR for miR-208 (**Fig. 6**). These data re-confirm efficient cardiac knockdown using the antagomiR design. However, the absence of either cholesterol or phosphorothioate prohibits the chemistry from establishing significant cardiac knockdown of miR-208 *in vivo*. Using either the C4 or C6 linker for cholesterol linkage to either the 5' or 3' end of the oligo, indicate that both linkers are suitable for the linkage of cholesterol. Using C6 at the 3' end of the oligo induces an even more efficient knockdown of miR-208 than the original antagomiR against miR-208 (**Fig. 5** and **6**). These data indicate that, unlike the *in*

vitro system, a lipid moiety such as cholesterol and phosphorothioate is desirable for cardiomyocyte knockdown, and both linkers, either 5' or 3', are sufficient in cholesterol linkage to the oligo and are able to establish a comparable if not enhanced knockdown of miR-208 compared to the original antagomiR design. See **Fig. 5**.

Experimental protocols

Northern blotting for miRNAs

To assess knockdown of miR-208 in response to the different chemistries and conjugates, Northern blot analysis using a probe directed against miR-208 served to detect the presence of miR-208 in the heart. Using Trizol reagent total RNA is collected from the heart, of which 10 ug is used for Northern blotting. To this end RNA samples are run on 20% acrylamide denaturing gels and transferred to Zeta-probe GT genomic blotting membranes (Bio-Rad) by electrophoresis. After transfer, blots are cross-linked and baked at 80° C for 1 hr. To maximize the sensitivity of miRNA detection, oligonucleotide probes are labeled with the Starfire Oligos Kit (IDT, Coralville, IA) and α -32P dATP (Amersham). Probes are hybridized to the membranes overnight at 39° C in Rapid-hyb buffer (Amersham), after which they are washed twice for 10 minutes at 39° C with 0.5x SSC containing 0.1% SDS. The blots are exposed and a U6 probe will serve as a loading control (U6 forward: 5-GTGCTCGCTTCGGCAGC-3 (SEQ ID NO: 64), U6 reverse: 5-AAAATATGGAACGCTTCACGAATTTGCG-3 (SEQ ID NO: 65)). The intensity of the radioactive signal will be used to quantify the fold change in expression using a phosphorimager and ImageQuant (Bio-Rad).

Real-time PCR analysis

[093] In addition to Northern blot analysis we performed miRNA specific realtime PCR analysis to both verify and quantify the level of miR-208 knockdown. Two μ g RNA from each tissue sample was used to generate cDNA using Super Script II reverse transcriptase per manufacturer's specifications (Invitrogen Life Technologies Inc., Burlington, Ontario, Canada). Real time PCR is cycled between 95 °C/30 s and 60 °C/30 for 40 cycles, following an initial denaturation step at 95°C for 3 min using Taqman probes purchased from ABI. Amplification products are routinely checked using dissociation curve software (Biorad), and transcript quantities are compared using the relative Ct method, where the amount of target normalized to the amount of endogenous SnoRNAs and relative to the control sample is given by $2^{-\Delta\Delta C_t}$.

REFERENCES

- Krieg, A. M., Tonkinson, J., Matson, S., Zhao, Q., Saxon, M., Zhang, L. M., Bhanja, U., Yakubov, L., and Stein, C. A. 1993. Modification of antisense phosphodiester oligodeoxynucleotides by a 5'-cholesteryl moiety increases cellular association and improves efficacy. *Proc Natl Acad Sci USA* 90:1048+–1052.
- Bijsterbosch, M. K., Manoharan, M., Dorland, R., Waarlo, I. H. E., Biessen, E. A. L., and van Berkel, T. J. C. 2001. Delivery of cholesteryl-conjugated phosphorothioate oligodeoxynucleotides to Kupffer cells by lactosylated low-density lipoprotein. *Biochem Pharmacol* 62:627–633.
- Manoharan, M., Tivel, K. L., Condon, T. P., Andrade, L. K., Barber-Peoch, I., Inamati, G., Shah, S., Mohan, V., Graham, M. J., Bennett, C. F., Crooke, S. T., and Cook, P. D. 1997. Conjugated antisense oligonucleotides. *Nucleosides Nucleotides* 16:1129–1138.
- Ghosh, YK, Visweswariah, SS, and Bhattacharya, S. 2002. Advantage of the Ether Linkage between the Positive Charge and the Cholesteryl Skeleton in Cholesterol-Based Amphiphiles as Vectors for Gene Delivery. *Bioconjugate Chem*, 13:378-384.
- Ghosh, YK, Visweswariah, SS, and Bhattacharya, S. 2000. Nature of linkage between the cationic headgroup and cholesteryl skeleton controls gene transfection efficiency. *FEBS Letters*, 473:341-344.
- Song, YK, Liu, F, Chu, S, and Liu, D. 1997. Characterization of Cationic Liposome-Mediated Gene Transfer In Vivo by Intravenous Administration. *Hum Gene Ther*, 8:1585-1592.

CLAIMS:

1. A lipophilic polynucleotide conjugate comprising a polynucleotide and a lipophilic moiety,

wherein the lipophilic moiety is spaced from the polynucleotide by a hydrocarbon linker, such that there is an absence of significantly polar groups and/or exchangeable protons in the vicinity of the lipophilic moiety.

2. The conjugate of claim 1, wherein the lipophilic moiety is capable of intercalating with a phospholipid bi-layer.

3. The conjugate of claim 1 or 2, wherein the lipophilic moiety is cholesteryl ether or derivative thereof.

4. The conjugate of claim 1 or 2, wherein the lipophilic moiety is a cholestene, cholestane, or cholestadiene.

5. The conjugate of claim 1 or 2, wherein the lipophilic moiety is a bile acid.

6. The conjugate of claim 5, wherein the bile acid is cholic acid, deoxycholic acid, and dehydrocholic acid.

7. The conjugate of claim 1 or 2, wherein the lipophilic moiety is a sterol or steroid.

8. The conjugate of claim 1 or 2, wherein the lipophilic moiety is a fat soluble alcohol or thiol.

9. The conjugate of any one of claims 1 to 8, wherein the hydrocarbon linker separates significantly polar groups and/or exchangeable protons from the lipophilic moiety by from 3 to about 15 atoms.
10. The conjugate of claim 9, wherein the hydrocarbon linker separates significantly polar groups and/or exchangeable protons from the lipophilic moiety by 6, 7, 8 or 9 atoms.
11. The conjugate of claim 10, wherein the hydrocarbon linker comprises a cyclic hydrocarbon.
12. The conjugate of any one of claims 1 to 11, wherein the hydrocarbon linker is conjugated to the lipophilic moiety through an ether or thioether linkage.
13. The conjugate of any one of claims 1 to 12, wherein the hydrocarbon linker comprises a C2 to C15 alkylene or alkenylene, which is optionally substituted.
14. The conjugate of claim 13, wherein the hydrocarbon linker comprises a C4 to C10, or a C4 to C8 saturated hydrocarbon.
15. The conjugate of any one of claims 1 to 14, wherein the hydrocarbon linker contains one or more heteroatoms in the carbon chain.
16. The conjugate of 15, wherein the hydrocarbon linker is conjugated on the end opposite the lipophilic moiety to the polynucleotide.

17. The conjugate of claim 16, wherein the hydrocarbon linker is conjugated to the polynucleotide through a phosphate ester or ether linkage.
18. The conjugate of claim 17, wherein the hydrocarbon linker is conjugated through the polynucleotide 5' position.
19. The conjugate of claim 17, wherein the hydrocarbon linker is conjugated through the polynucleotide 3' position.
20. The conjugate of any one of claims 1 to 19, wherein the conjugate has a structure selected from Structures (I) to (XIII).
21. The conjugate of any one of claims 1 to 20, wherein the polynucleotide is DNA-based.
22. The conjugate of any one of claims 1 to 20, wherein the polynucleotide is RNA-based.
23. The conjugate of any one of claims 1 to 22, wherein the polynucleotide has a modified polynucleotide backbone.
24. The conjugate of claim 23, wherein the polynucleotide contains one or more phosphorothioate modifications.
25. The conjugate of any one of claims 1 to 24, wherein the polynucleotide has one or more modified nucleoside units.

26. The conjugate of any one of claims 1 to 25, wherein the polynucleotide has one or more single stranded and one or more double stranded regions.

27. The conjugate of claim 26, wherein the polynucleotide is an antisense oligonucleotide, short interfering RNA (siRNA), double stranded RNA (dsRNA), single stranded RNA (ssRNA), microRNA (miRNA), short hairpin RNA (shRNA), or ribozyme.

28. The conjugate of claim 27, wherein the polynucleotide is an antagomir.

29. The conjugate of claim 27 or 28, wherein the polynucleotide is about 5 to about 50 nucleotides in length.

30. The conjugate of claim 27 or 28, wherein the polynucleotide is about 18 to about 30 nucleotides in length.

31. The conjugate of claim 27 or 28, wherein the polynucleotide comprises a sequence that is at least about 75% complementary to a mature miRNA.

32. The conjugate of claim 27 or 28, wherein the polynucleotide comprises a sequence that is at least about 75% complementary to a mature miRNA listed in Table 1.

33. The conjugate of claim 27 or 28, wherein the polynucleotide comprises a sequence that is 100% complementary to a mature miRNA.

34. The conjugate of claim 27 or 28, wherein the polynucleotide comprises a sequence that is 100% complementary to a mature miRNA listed in Table 1.
35. The conjugate of claim 27 or 28, wherein the polynucleotide has a sequence designed to mimic a cellular miRNA.
36. The conjugate of claim 27 or 28, wherein the polynucleotide has a sequence designed to mimic a miRNA listed in Table 1.
37. The conjugate of any one of claims 1 to 36, wherein the polynucleotide comprises one or more of a locked nucleic acid, peptide nucleic acid backbone, and/or a sugar modification at the 2' or 4' position.
38. The conjugate of any one of claims 1 to 37, wherein the conjugate has from one to five lipophilic moieties.
39. A pharmaceutical composition, comprising the conjugate of any one of claims 1 to 38 formulated for delivery to a patient.
40. The pharmaceutical composition of claim 39, wherein the conjugate is formulated as a colloidal dispersion system, macromolecular complex, nanocapsule, microsphere, bead, oil-in-water emulsion, micelle, mixed micelle, or liposome.
41. The pharmaceutical composition of claim 40, wherein the lipophilic moieties anchor the polynucleotide conjugate in a liposome.

42. The pharmaceutical composition of any one of claims 39 to claim 41, wherein the composition is formulated for intradermal delivery, subcutaneous delivery, intramuscular delivery, intraperitoneal or intravenous delivery.

43. The pharmaceutical composition of claim 42, wherein the composition is formulated for administration by a cardiac catheter system.

44. A method for delivering a polynucleotide to a mammalian cell, comprising administering the polynucleotide as a conjugate in accordance with any one of claims 1 to 38.

45. A method for delivering a polynucleotide to a mammalian patient, comprising administering the polynucleotide as a conjugate in accordance with any one of claims 1 to 38.

46. The method of claim 44 or 45, wherein the polynucleotide is a miRNA or a miRNA inhibitor.

47. The method of claim 46, wherein the patient has a condition associated with miRNA expression.

48. The method of claim 47, wherein the condition is one or more of cardiac hypertrophy, myocardial infarction, heart failure, vascular damage, and pathologic cardiac fibrosis.

49. A method for synthesizing a lipophilic polynucleotide conjugate, comprising:

preparing a lipophilic moiety with a substantially linear hydrocarbon chain having terminal functional groups to thereby prepare a first intermediate, wherein the lipophilic

moiety is spaced from the terminal functional group by a hydrocarbon linker, such that there is an absence of significantly polar groups and/or exchangeable protons in the vicinity of the lipophilic moiety,

preparing from said first intermediate, a second intermediate that has functional groups suitable for incorporation into a polynucleotide chain,

incorporating the second intermediate into a polynucleotide chain during solid phase synthesis.

50. The method of claim 49, where the lipophilic moiety is a sterol, and the first intermediate is an ether-linked intermediate.

51. The method of claims 49 or 50, wherein the substantially linear hydrocarbon linker is a C3-C15 alkyl, alkenyl, alkynyl, and wherein the hydrocarbon is optionally substituted.

52. The method of claim 51, wherein the substantially linear hydrocarbon linker is a C4-C8 alkyl, alkenyl, alkynyl, and wherein the hydrocarbon is optionally substituted.

53. The method of any one of claims 49 to 52, wherein the first intermediate is reacted to prepare a phosphoramidite as the second intermediate.

54. The method of claim 53, wherein the phosphoramidite is converted to a phosphate ester during solid phase polynucleotide synthesis.

55. The method of any one of claims 49 to 54, wherein the first intermediate is reacted to prepare a suitable terminal group for coupling to a support, thereby forming the 3' end of a polynucleotide chain.

56. The method of any one of claims 49 to 55, wherein the second intermediate is prepared from a compound having two differentiated hydroxyl groups separated by at least one carbon.

57. A method for synthesizing a polynucleotide conjugate, comprising:


preparing a lipophilic moiety with a hydrocarbon chain having terminal functional groups to thereby prepare an intermediate, wherein the lipophilic moiety is spaced from the terminal functional group by a hydrocarbon linker, such that there is an absence of significantly polar groups and/or exchangeable protons in the vicinity of the lipophilic moiety,

coupling the intermediate to a polynucleotide.


58. The method of claim 57, wherein the intermediate is coupled to the polynucleotide by a reaction or series of reactions selected from carbonyl addition-elimination/reductive amination, amidation, maleimide-thiol coupling, or aqueous Diels-Alder.

FIGURE 1

Natural Cholesterol



Current State of the Art Chol Conjugates



Novel Chol Conjugate with enhanced Lipophilicity




FIGURE 2




FIGURE 3

Synthesis of 1-((bis(4-methoxyphenyl)(phenyl)methoxy))-12-(Cholest-5-en-3-oxyl)dodecan-3-ol Functionalized CPG




FIGURE 4





FIGURE 5






FIGURE 6

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/33729

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C07H 21/00 (2010.01)

USPC - 536/23.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC - 536/23.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST -- PGPB,USPT,USOC,EPAB,JPAB; Dialog Classic Files -- 654, 652, 351, 349, 6, 35, 65, 155; Google Scholar; USPTO Web Page; Search terms -- lipophilic conjugate, nucleic acid, hydrocarbon linker, cholesterol, cholestene, cholesteryl ether, bile salts, fatty alcohol

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	US 2005/0107325 A1 (MANOHARAN et al.) 19 May 2005 (19.05.2005) para [0002], [0125], [0136], [0137], [0441], [0544], [0962], [0964], [1019]	1, 2, 5-7 ----- 3, 4, 8
Y	US 2003/0134420 A1 (LOLLO et al.) 17 July 2003 (17.07.2003) para [0066], [0170], [0253], Fig 5	3
Y	US 2005/0130911 A1 (UHLMANN et al.) 16 June 2005 (16.06.2005) abstract, para [0008]	4
Y	US 2008/0199960 A1 (JULIANO et al.) 21 August 2008 (21.08.2008) para [0002], [0028], [0034], [0035], [0169]	8

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

“A” document defining the general state of the art which is not considered to be of particular relevance

“E” earlier application or patent but published on or after the international filing date

“L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

“O” document referring to an oral disclosure, use, exhibition or other means

“P” document published prior to the international filing date but later than the priority date claimed

“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

“&” document member of the same patent family

Date of the actual completion of the international search

07 September 2010 (07.09.2010)

Date of mailing of the international search report

17 SEP 2010

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/33729

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 9-48 and 53-56
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I: Claims 1-8, directed to a lipophilic polynucleotide conjugate, wherein the lipophilic moiety is space from the polynucleotide by a hydrocarbon linker, such that there is an absence of polar groups or exchangeable protons in the vicinity of the lipophilic moiety.

Group II: claims 49-52, 57 and 58 directed to methods for synthesizing a lipophilic polynucleotide conjugate, comprising a lipophilic moiety with a substantially linear hydrocarbon chain having terminal functional groups, wherein the lipophilic moiety is spaced from the terminal functional group of the hydrocarbon linker, such that there is an absence of polar groups or exchangeable protons in the vicinity of the lipophilic moiety.

- Please see extra sheet for continuation -

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-8

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/33729

Continuation of Box III: Unity of Invention

The inventions listed as Groups I - II do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The special technical feature of the Group I claims is a lipophilic polynucleotide conjugate, wherein the lipophilic moiety is space from the polynucleotide by a hydrocarbon linker, such that there is an absence of polar groups or exchangeable protons in the vicinity of the lipophilic moiety. The special technical feature of the Group II claims is methods for synthesizing a lipophilic polynucleotide conjugate, comprising a lipophilic moiety with a substantially linear hydrocarbon chain having terminal functional groups, wherein the lipophilic moiety is spaced from the terminal functional group of the hydrocarbon linker, such that there is an absence of polar groups or exchangeable protons in the vicinity of the lipophilic moiety.

The common technical element shared by the above groups is a lipophilic polynucleotide conjugate, wherein the lipophilic moiety is space from the polynucleotide by a hydrocarbon linker, such that there is an absence of polar groups or exchangeable protons in the vicinity of the lipophilic moiety. This common technical element does not represent an improvement over the prior art of US 2006/0058266 A1 to Manoharan et al. (para [0005]; an oligonucleotide agent coupled to a ligand...In a preferred embodiment, the coupling is by a tether or a linker -- para [0009]; One of the most preferred moieties is a moiety which promotes entry into a cell, e.g., a lipophilic moiety -- para [0350]; Tethers are connected to the carrier at a tethering attachment point (TAP) and may include any C.sub.1-C.sub.100 carbon-containing moiety, (e.g. C.sub.1-C.sub.75, C.sub.1-C.sub.50, C.sub.1-C.sub.20, C.sub.1-C.sub.10; C.sub.1, C.sub.2, C.sub.3, C.sub.4, C.sub.5, C.sub.6, C.sub.7, C.sub.8, C.sub.9, or C.sub.10)...Preferred tethered ligands may include...TAP-(CH.sub.2).sub.n(LIGAND) -- para [0351] - In some embodiments, the tether can terminate with...an olefin (e.g., CH.dbd.CH.sub.2)...TAP-(CH.sub.2).sub.n-(CH.dbd.CH.sub.2), or TAP-C(O)(CH.sub.2).sub.n(CH.dbd.CH.sub.2)). Therefore, the inventions of Group I and Group II lack unity of invention under PCT Rule 13 because they do not share a same or corresponding special technical feature.