



(11)

EP 2 854 849 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

28.02.2018 Bulletin 2018/09

(51) Int Cl.:

A61K 39/395 ^(2006.01)

(21) Application number: **13730702.1**

(86) International application number:

PCT/US2013/043516

(22) Date of filing: **31.05.2013**

(87) International publication number:

WO 2013/181486 (05.12.2013 Gazette 2013/49)

(54) STABILIZED FORMULATIONS CONTAINING ANTI-DLL4 ANTIBODIES

STABILISIERTE FORMULIERUNGEN MIT ANTI-DLL4-ANTIKÖRPERN

FORMULATIONS STABILISÉES CONTENANT DES ANTICORPS ANTI-DLL4

(84) Designated Contracting States:

AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

Designated Extension States:

BA ME

(30) Priority: **31.05.2012 US 201261653478 P**

(43) Date of publication of application:

08.04.2015 Bulletin 2015/15

(73) Proprietor: **Regeneron Pharmaceuticals, Inc.**

Tarrytown, NY 10591 (US)

(72) Inventors:

- **WALSH, Scott**
Tarrytown, NY 10591 (US)
- **DIX, Daniel**
LaGrangeville, NY 12540 (US)

(74) Representative: **J A Kemp**

14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(56) References cited:

WO-A2-2008/076379 WO-A2-2011/109298

- **ANN L DAUGHERTY AND RANDALL J MRSNY ED - STEVEN J SHIRE ET AL: "Formulation and Delivery Issues for Monoclonal Antibody Therapeutics", CURRENT TRENDS IN MONOCLONAL ANTIBODY DEVELOPMENT AND MANUFACTURING, SPRINGER, US, 1 January 2010 (2010-01-01), pages 103-129, XP009133774, ISBN: 978-0-387-76642-3 [retrieved on 2010-11-11]**
- **FENG HE ET AL: "High-throughput assessment of thermal and colloidal stability parameters for monoclonal antibody formulations", JOURNAL OF PHARMACEUTICAL SCIENCES, vol. 100, no. 12, 24 December 2011 (2011-12-24), pages 5126-5141, XP055076385, ISSN: 0022-3549, DOI: 10.1002/jps.22712**
- **WANG W ET AL: "ANTIBODY STRUCTURE, INSTABILITY, AND FORMULATION", JOURNAL OF PHARMACEUTICAL SCIENCES, AMERICAN PHARMACEUTICAL ASSOCIATION, WASHINGTON, US, vol. 96, no. 1, 1 January 2007 (2007-01-01), pages 1-26, XP009084505, ISSN: 0022-3549, DOI: 10.1002/JPS.20727**
- **FENG HE ET AL: "High Throughput Thermostability Screening of Monoclonal Antibody Formulations", JOURNAL OF PHARMACEUTICAL SCIENCES, AMERICAN PHARMACEUTICAL ASSOCIATION, WASHINGTON, US, vol. 99, no. 4, 1 April 2010 (2010-04-01), pages 1707-1720, XP007913147, ISSN: 0022-3549, DOI: 10.1002/JPS.21955 [retrieved on 2009-09-24]**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 854 849 B1

Description

[0001] This application is being filed on 31 May 2013, as a PCT International patent application, and claims priority to U.S. Provisional Patent Application No. 61/653,478, filed May 31, 2012.

FIELD

[0002] The present invention relates to the field of therapeutic antibody formulations. More specifically, the present invention relates to the field of pharmaceutical formulations comprising an antibody that specifically binds to human delta-like ligand 4 (DII4).

SEQUENCE LISTING

[0003] An ST.25 compliant computer readable text file of a sequence listing is filed concurrently with the present specification according to 37 C.F.R. § 1.821(e). The contents of the text file are herein incorporated by reference. A paper copy of the sequence listing, which pursuant to 37 C.F.R. § 1.821(f) is identical in content to the ST.25 compliant computer readable text file, is included as part of the present specification and is herein incorporated by reference.

BACKGROUND

[0004] The Notch-signaling pathway is a system for cell-to-cell communication used by a wide range of eukaryotes for many biological processes, such as differentiation, proliferation, and homeostasis. Delta like 4 (DII4) or delta-like ligand 4 (DII4) (hereinafter "DII4") is a member of the Delta family of Notch ligands which exhibits highly selective expression by vascular endothelium (Shutter et al. (2000) Genes Develop. 14:1313-1318). DII4 is a ligand for Notch receptors, including Notch1 and Notch 4. The amino acid sequence for human DII4 is depicted in SEQ ID NO:9. Given the important role that the DII4-Notch signaling pathway plays in the regulation of angiogenesis and its implication in human health and medicine, it is desired to develop and deploy DII4-based medicaments. One such DII4-based medicament is a therapeutic antibody specific to DII4.

[0005] Methods to produce antibodies useful as human therapeutics include generation of chimeric antibodies and humanized antibodies (see, for example, U.S. 6,949,245). See, for example, WO 94/02602 (Abgenix) and U.S. 6,596,541 (Regeneron Pharmaceuticals), describing methods of generating nonhuman transgenic mice capable of producing human antibodies. U.S. Patents No. 7,488,806, 7,534,868, and 7,919,593, and U.S. Patent Application No. U.S. 2011-0150905A1 disclose antibodies to human DII4.

[0006] Therapeutic antibodies must be formulated in a manner that not only makes the antibodies suitable for administration to patients, but also in a manner that maintains their stability during storage and subsequent use. For example, therapeutic antibodies in liquid solution are prone to fragmentation, precipitation, aggregation, and undesired chemical modifications unless the solution is formulated properly. The stability of an antibody in liquid formulation depends not only on the kinds of excipients used in the formulation, but also on the amounts and proportions of the excipients relative to one another. Furthermore, other considerations aside from stability must be taken into account when preparing a liquid antibody formulation. Examples of such additional considerations include the viscosity of the solution and the concentration of antibody that can be accommodated by a given formulation, and the visual quality or appeal of the formulation. Thus, when formulating a therapeutic antibody, great care must be taken to arrive at a formulation that remains stable, contains an adequate concentration of antibody, and possesses a suitable viscosity as well as other properties which enable the formulation to be conveniently administered to patients.

[0007] Antibodies to DII4 are one example of a therapeutically relevant macromolecule that requires proper formulation. Although some anti-DII4 antibodies are known, there nonetheless remains a need in the art for novel pharmaceutical formulations comprising anti-DII4 antibodies that are sufficiently stable and suitable for administration to patients.

SUMMARY

[0008] The present invention satisfies the aforementioned need by providing pharmaceutical formulations comprising a human antibody that specifically binds to human delta-like ligand 4 (DII4). The scope of the present invention is defined by the claims. A liquid pharmaceutical formulation is provided, comprising: (i) an antibody that specifically binds to DII4; (ii) a buffer; (iii) an organic cosolvent; and (iv) thermal stabilizers. An antibody is provided at a concentration from about 20 ± 3 mg/mL to about 75 ± 11.25 mg/mL. The antibody is provided at a concentration of about 25 mg/mL ± 3.75 mg/mL. An antibody is provided at a concentration of about 50 mg/mL ± 7.5 mg/mL.

[0009] In one embodiment, exemplary anti-DII4 antibodies of the invention comprise HCDR1, HCDR2, HCDR3, LCDR1, LCDR2 and LCDR3 domains having the respective amino acid sequences of SEQ ID NO: 2, 3, 4, 6, 7 and 8 (e.g.,

REGN421). In one embodiment, the antibody comprises the CDR sequences contained within an HCVR having an amino acid sequence of SEQ ID NO:1, and an LCVR having an amino acid of SEQ ID NO:5.

[0010] In one embodiment, the pH of the liquid formulation is about pH 5.0 to about pH 6.6; pH 6.0 ± 0.5 , pH 6.0 ± 0.4 , pH 6.0 ± 0.3 , pH 6.0 ± 0.2 , pH 6.0 ± 0.1 , pH 6.0 ± 0.05 , pH 6.0 ± 0.01 , or pH 6.0. In the formulation of the invention the pH of the liquid formulation is about pH 6.0 ± 0.5 . In one embodiment, the buffer has an effective buffering range of about pH 5.8 to about pH 8.0; or about pH 6.0 to about 7.5. In one embodiment, the buffer has a pKa of about 6.0 to about 7.5. In one embodiment, the buffer has a pKa of 6.0. In one embodiment, the buffer has a pKa of 6.4. In one embodiment, the buffer has a pKa of 6.5. In one embodiment, the buffer has a pKa of 7.2.

[0011] In the formulation of the invention the buffer is a phosphate buffer. In one embodiment, the phosphate is at a concentration of $5 \text{ mM} \pm 0.75 \text{ mM}$ to $50 \text{ mM} \pm 7.5 \text{ mM}$. In one embodiment, the phosphate is at a concentration of $5 \text{ mM} \pm 0.75 \text{ mM}$ or about 5 mM. In the formulation of the invention the phosphate is at a concentration of $10 \text{ mM} \pm 1.5 \text{ mM}$ or about 10 mM. In one embodiment, the phosphate is at a concentration of $15 \text{ mM} \pm 2.25 \text{ mM}$ or about 15 mM. In one embodiment, the phosphate is at a concentration of $20 \text{ mM} \pm 3 \text{ mM}$ or about 20 mM. In one embodiment, the phosphate is at a concentration of $25 \text{ mM} \pm 3.75 \text{ mM}$ or about 25 mM. In one embodiment, the phosphate is at a concentration of $30 \text{ mM} \pm 4.5 \text{ mM}$ or about 30 mM. In one embodiment, the phosphate is at a concentration of $35 \text{ mM} \pm 5.25 \text{ mM}$ or about 35 mM. In one embodiment, the phosphate is at a concentration of $40 \text{ mM} \pm 6 \text{ mM}$ or about 40 mM. In one embodiment, the phosphate is at a concentration of $45 \text{ mM} \pm 6.75 \text{ mM}$ or about 45 mM. In one embodiment, the phosphate is at a concentration of $50 \text{ mM} \pm 7.5 \text{ mM}$ or about 50 mM.

[0012] In one embodiment, the organic cosolvent is a nonionic polymer containing a polyoxyethylene moiety. In some embodiments, the organic cosolvent is any one or more of polysorbate 20, polysorbate 80, poloxamer 188, and polyethylene glycol 3350. In a specific embodiment, the organic cosolvent is polysorbate 20.

[0013] In one embodiment, the organic cosolvent is at a concentration of from about $0.005\% \pm 0.00075\%$ to about $1\% \pm 0.15\%$ "weight to volume" or "w/v", wherein, e.g., $0.1 \text{ g/ml} = 10\%$ and $0.01 \text{ g/ml} = 1\%$. In the formulation of the invention the organic cosolvent is polysorbate 20, which is at a concentration of $0.2\% \pm 0.03\% \text{ w/v}$, or about $0.2\% \text{ w/v}$. In another embodiment, the organic cosolvent is polysorbate 20, which is at a concentration of $0.01\% \pm 0.0015\% \text{ w/v}$ or about $0.01\% \text{ w/v}$.

[0014] In one embodiment, the stabilizer is a sugar, a salt, an amino acid, or a combination of any one or more thereof. In one embodiment, the sugar is selected from the group consisting of sucrose, sorbitol, mannitol, glycerol and trehalose; the salt is sodium chloride; and the amino acid is glycine. In one embodiment, the stabilizer comprises a combination of sucrose and sodium chloride.

[0015] In one embodiment, the sucrose is at a concentration of from about 5% to about 40%, w/v; and the sodium chloride is from about 50 mM to about 250 mM. The inventive formulation comprises $10\% \pm 1.5\%$, about 10%, or 10% sucrose; and $150 \text{ mM} \pm 22.5 \text{ mM}$, about 150 mM, or 150 mM sodium chloride. In another embodiment, the formulation comprises $20\% \pm 3\%$, about 20%, or 20% sucrose.

[0016] In one embodiment, the viscosity of the formulation is about 1 cPoise to about 5 cPoise. In one embodiment, the viscosity of the formulation is $1.5 \text{ cPoise} \pm 0.23 \text{ cPoise}$, about 1.5 cPoise, or 1.5 cPoise.

[0017] In one embodiment, the osmolality of the formulation is close to within a physiological range. In one embodiment, the formulation has an osmolality of about 300 milli-Osmoles per kilogram (mOsm) to about 700 mOsm. In one embodiment, the osmolality of the formulation is $650 \text{ mOsm} \pm 98 \text{ mOsm}$, about 650 mOsm, or 650 mOsm.

[0018] In one embodiment, at least 95% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 24 months of storage at -80°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 57% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 24 months of storage at -80°C is of the non-basic and non-acidic form (*i.e.*, main peak or main charge form or "region 2 peak"), as determined by ion exchange chromatography. In one embodiment, anti-DII4 on average maintains from about 65% to about 135% of its binding activity after 24 months of storage at -80°C , relative to the binding activity of the antibody prior to storage.

[0019] In one embodiment, at least 95% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 18 months of storage at -30°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 57% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 18 months of storage at -30°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after 18 months of storage at -30°C , relative to the binding activity of the antibody prior to storage.

[0020] In one embodiment, at least 95% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after six months of storage at -20°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 60% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after six months of storage at -20°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after six months of storage at -20°C , relative to the binding activity of the antibody prior to storage.

[0021] In one embodiment, at least 95% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after six months of storage at 5°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 61% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after six months of storage at 5°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after six months of storage at 5°C, relative to the binding activity of the antibody prior to storage.

[0022] In one embodiment, at least 97% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after six months of thermal stress at 25°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 53% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after six months of storage at 25°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after six months of storage at 25°C, relative to the binding activity of the antibody prior to storage.

[0023] In one embodiment, at least 94% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 28 days of thermal stress at 45°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 45% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 28 days of thermal stress at 45°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after 28 days of thermal stress at 45°C, relative to the binding activity of the antibody prior to application of the thermal stress condition.

[0024] In one embodiment, at least 96% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 28 days of thermal stress at 37°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 51% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 28 days of thermal stress at 37°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after 28 days of thermal stress at 37°C, relative to the binding activity of the antibody prior to application of the thermal stress condition.

[0025] In one embodiment, at least 97% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 28 days of thermal stress at 25°C is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 57% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 28 days of thermal stress at 25°C is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 antibody maintains on average about 65% to about 135% of its binding activity after 28 days of thermal stress at 25°C, relative to the binding activity of the antibody prior to application of the thermal stress condition.

[0026] In one embodiment, at least 98% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 120 minutes of agitation stress is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 59% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after 120 minutes of agitation stress is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 maintains on average about 65% to about 135% of its binding activity after 120 minutes of agitation stress, relative to the binding activity of the antibody prior to application of the agitation stress condition.

[0027] In one embodiment, at least 98% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after eight freeze/thaw cycles is non-aggregated and un-degraded, as determined by size exclusion chromatography. In one embodiment, at least 58% of the anti-DII4 antibody recovered from the liquid pharmaceutical formulation after eight freeze/thaw cycles is of the main charge form, as determined by ion exchange chromatography. In one embodiment, the anti-DII4 maintains on average about 65% to about 135% of its binding activity after eight freeze/thaw cycles, relative to the binding activity of the antibody prior to application of the freeze/thaw stress condition.

[0028] In one aspect, a liquid pharmaceutical formulation is provided, comprising: (i) from 20 ± 3 mg/ml to 75 ± 11.25 mg/ml of a human antibody that specifically binds to human DII4; (ii) from $5 \text{ mM} \pm 0.75 \text{ mM}$ to $50 \text{ mM} \pm 7.5 \text{ mM}$ phosphate; (iii) from $0.005\% \pm .000075\%$ to $1\% \pm 0.15\%$ (w/v) polysorbate 20; (iv) from $5\% \pm 0.75\%$ to $40\% \pm 6\%$ (w/v) sucrose; and (v) from $50 \text{ mM} \pm 7.5 \text{ mM}$ to $250 \text{ mM} \pm 37.5 \text{ mM}$ sodium chloride, at a pH of from about 5.5 to about 6.5. The anti-DII4 antibody of this aspect comprises a heavy chain variable region (HCVR) and a light chain variable region (LCVR) such that the HCVR/LCVR combination comprises heavy and light chain complementarity determining regions (HCDR1-HCDR2-HCDR3 / LCDR1-LCDR2-LCDR3), which comprise the amino acid sequences of SEQ ID NOs:2 - 3 - 4 / SEQ ID NOs:6 - 7 - 8, respectively. In a particular embodiment, the anti-DII4 antibody comprises a heavy chain variable region (HCVR) and light chain variable region (LCVR) comprising an amino acid sequence of SEQ ID NO: 1 and SEQ ID NO: 5, respectively (antibody REGN421 of U.S. Pat. Nos. 7,488,806, 7,534,868, and 7,919,593, and U.S. Pat. App. Pub. No. 2011-0150905).

[0029] In one embodiment of the invention the liquid formulation comprises (i) $25 \text{ mg/ml} \pm 3.75 \text{ mg/ml}$ of antibody REGN421; (ii) $10 \pm 1.5 \text{ mM}$ phosphate; (iii) $0.2\% \pm 0.03\%$ (w/v) polysorbate 20; (iv) $10\% \pm 1.5\%$ (w/v) sucrose; and

(v) 150 mM $\pm$ 22.5 mM sodium chloride at a pH of 6.0 $\pm$ 0.5. In one embodiment, after thermal stress of the formulation at 45° for 28 days, $\geq$ 94% of the antibody retains its native conformation, $\geq$ 45% of the antibody is of the main charge form, and $\geq$ 98% of the antibody retains its binding activity. In another embodiment, after thermal stress of the formulation at 37° for 28 days, $\geq$ 97% of the antibody is native and $\geq$ 59% of the antibody is of the main charge form, and about 100% of the antibody retains its binding activity. In one embodiment, after thermal stress of the formulation at 25° for 28 days, $\geq$ 97% of the antibody is native and $\geq$ 57% of the antibody is of the main charge form, and about 100% of the antibody retains its binding activity. In one embodiment of this particular formulation, after eight cycles of freezing followed by thawing, $\geq$ 98% of the antibody is native and $\geq$ 59% of the antibody is of the main charge form, and about 100% of the antibody retains its binding activity. In one embodiment of this particular formulation, after 120 minutes of agitation stress, $\geq$ 98% of the antibody is native and $\geq$ 58% of the antibody is of the main charge form, and at least about 98% of the antibody retains its potency.

[0030] In one aspect, a liquid pharmaceutical formulation of any of the preceding aspects is provided in a container. In one embodiment, the container is a polycarbonate vial. In another embodiment, the container is a glass vial, which in some embodiments is a type 1 borosilicate glass vial with a fluorocarbon-coated butyl rubber stopper. In yet another embodiment, the container is a bag, such as an intravenous drip (IV) bag, which in some embodiments is made of polyvinyl chloride or polyolefin. In another embodiment, the container is an injection device, such as a microinfuser or a syringe. Described is a pharmaceutical formulation comprising (a) 25 mg/mL $\pm$ 3.75 mg/mL of an anti-Ang-2 antibody, (b) 10 mM $\pm$ 1.5 mM phosphate, pH 6 $\pm$ 0.5, (c) 0.2% $\pm$ 0.03% polysorbate 20, (d) 10% w/v $\pm$ 1.5% sucrose, and (e) 150 mM $\pm$ 22.5 mM sodium chloride is provided, wherein (a) the antibody comprises an HCVD of SEQ ID NO: 1 and an LCVD of SEQ ID NO: 5, (b) the antibody has a molecular mass of about 146 kDa, about 147 kDa, or about 150 kDa, (c) from about 82% to about 87% of the antibodies contain fucosylated oligosaccharide chains, and (d) the heavy chains of the antibody lack a C-terminal lysine. The pharmaceutical formulation consists of (a) 25 mg/mL $\pm$ 3.75 mg/mL of an anti-Ang-2 antibody, (b) 10 mM $\pm$ 1.5 mM phosphate, pH 6 $\pm$ 0.5, (c) 0.2% $\pm$ 0.03% polysorbate 20, (d) 10% $\pm$ 1.5% sucrose, and (e) 150 mM $\pm$ 22.5 mM sodium chloride is provided, wherein (a) the antibody comprises an HCVD of SEQ ID NO: 1 and an LCVD of SEQ ID NO: 5, (b) the antibody has a molecular mass of about 146 kDa, about 147 kDa, or about 150 kDa, (c) from about 82% to about 87% of the antibodies contain fucosylated oligosaccharide chains, and (d) the heavy chains of the antibody lack a C-terminal lysine.

[0031] In one aspect, a kit comprising a pharmaceutical composition of any one of the preceding aspects, a container, and instructions is provided. In one embodiment, the container is a prefilled syringe. In one embodiment, the container is a borosilicate vial fitted with a FLUROTEC-coated 4023/50-rubber stopper.

[0032] Other embodiments of the present invention will become apparent from a review of the ensuing detailed description.

DESCRIPTION

[0033] Before the present invention is described, it is to be understood that this invention is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0034] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein, the term "about", when used in reference to a particular recited numerical value or range of values, means that the value may vary from the recited value by no more than 5%. For example, as used herein, the expression "about 100" includes 95 and 105 and all values in between (e.g., 95.00, 95.01, 95.02, 95.03, 95.04, ..., 104.96, 104.97, 104.98, 104.99, 105.00).

[0035] Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described.

PHARMACEUTICAL FORMULATIONS

[0036] As used herein, the expression "pharmaceutical formulation" means a combination of at least one active ingredient (e.g., a small molecule, macromolecule, compound, etc. which is capable of exerting a biological effect in a human or non-human animal), and at least one inactive ingredient which, when combined with the active ingredient or one or more additional inactive ingredients, is suitable for therapeutic administration to a human or non-human animal. The term "formulation", as used herein, means "pharmaceutical formulation" unless specifically indicated otherwise. The present invention provides pharmaceutical formulations comprising at least one therapeutic polypeptide. According to certain embodiments of the present invention, the therapeutic polypeptide is an antibody, or an antigen-binding fragment thereof, which binds specifically to human delta-like ligand-4 (DIL4) protein. More specifically, the present invention includes pharmaceutical formulations that comprise: (i) a human antibody that specifically binds to human DIL4 (ii) a

phosphate buffer; (iii) an organic cosolvent that is a non-ionic surfactant; and (iv) a stabilizer that is a carbohydrate or an inorganic salt, or a combination of carbohydrate and inorganic salt. Specific exemplary components and formulations included within the present invention are described in detail below.

ANTIBODIES THAT BIND SPECIFICALLY TO DLL4

[0037] The pharmaceutical formulations of the present invention may comprise a human antibody, that binds specifically to human Dll4. As used herein, the term "Dll4" means a human delta-like ligand 4, which is a vascular-specific ligand of Notch. The Dll4-Notch signaling pathway is generally understood to modulate angiogenesis by blocking blood vessel branching and promoting blood vessel maturation (Dll4 as a negative regulator of branching; reviewed in Sainsin and Harris, Trends in Molecular Medicine, Vol. 13(9):389-395, Sep. 2007.) Agents that antagonize the Dll4-Notch pathway have been shown to inhibit the rate of tumor growth by "triggering excessive but nonfunctional angiogenesis." (See Sainsin at 389.) Dll4 is upregulated by VEGF and is thought to promote the organized development of functional neo-vessels. An exemplary human Dll4 amino acid sequence is described in SEQ ID NO: 9. Antibodies to human Dll4 are described in patents no. 7,488,806, 7,534,868, and 7,919,593.

[0038] The term "antibody", as used herein, is generally intended to refer to immunoglobulin molecules comprising four polypeptide chains: two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, as well as multimers thereof (e.g., IgM); however, immunoglobulin molecules consisting of only heavy chains (*i.e.*, lacking light chains) are also encompassed within the definition of the term "antibody". Each heavy chain comprises a heavy chain variable region (abbreviated herein as HCVR or V_H) and a heavy chain constant region. The heavy chain constant region comprises three domains, CH1, CH2 and CH3. Each light chain comprises a light chain variable region (abbreviated herein as LCVR or V_L) and a light chain constant region. The light chain constant region comprises one domain (CL1). The V_H and V_L regions can be further subdivided into regions of hypervariability, termed complementary determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

[0039] Unless specifically indicated otherwise, the term "antibody", as used herein, shall be understood to encompass complete antibody molecules as well as antigen-binding fragments thereof. The term "antigen-binding portion" or "antigen-binding fragment" of an antibody (or simply "antibody portion" or "antibody fragment"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to human Dll4 or an epitope thereof.

[0040] An "isolated antibody", as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds human Dll4 is substantially free of antibodies that specifically bind antigens other than human Dll4), with the notable exception of bi-specific (or multi-specific) antibodies that specifically bind Dll4 on the one hand, and another epitope on the other. Moreover, an isolated antibody may be substantially free of other cellular material or chemicals.

[0041] The term "specifically binds", or the like, means that an antibody or antigen-binding fragment thereof forms a complex with an antigen that is relatively stable under physiologic conditions. Specific binding can be characterized by a dissociation constant of at least about 1×10^{-6} M or greater. Methods for determining whether two molecules specifically bind are well known in the art and include, for example, equilibrium dialysis, surface plasmon resonance, and the like. An isolated antibody that specifically binds human Dll4 may, however, have cross-reactivity to other antigens, such as Dll4 molecules from other species (orthologs). In the context of the present invention, multispecific (e.g., bispecific) antibodies that bind to human Dll4 as well as one or more additional antigens are deemed to "specifically bind" human Dll4.

[0042] Exemplary anti-human Dll4 antibodies that may be included in the pharmaceutical formulations of the present invention are set forth in U.S. Patents No. 7,488,806, 7,534,868, 7,919,593, and U.S. Patent Application No. 2011-0150905.

[0043] According to certain embodiments of the present invention, the anti-human Dll4 antibody is a human IgG1 comprising a heavy chain variable region of the IGHV3-33.01 subtype and a light chain variable region of the IGKV3-11.01 subtype (see Barbie and Lefranc, The Human Immunoglobulin Kappa Variable (IGKV) Genes and Joining (IGKJ) Segments, Exp. Clin. Immunogenet. 1998; 15:171-183; and Scaviner, D. et al., Protein Displays of the Human Immunoglobulin Heavy, Kappa and Lambda Variable and Joining Regions, Exp. Clin. Immunogenet., 1999; 16:234-240). The germline IGHV3-33 and IGKV3-11 sequences, and the amino acid position assignment numbers presented herein comport with the international Immunogenetics (IMGT) information system, as described in Lefranc, M.-P., et al., IMGT®, the international ImMunoGeneTics information system®, Nucl. Acids Res, 37, D1006-D1012 (2009).

[0044] In some embodiments, the anti-human Dll4 antibody comprises one or more amino acid substitutions in one or more framework regions relative to the canonical heavy chain variable region, which is reasonably expected to result in an altered charge distribution across the exposed surface of the antibody, and therefore affect its interaction with the surrounding solvent and excipients. In some embodiments, the amino acid substitution comprises a substitution of serine for alanine, methionine for threonine, and/or glutamic acid for glutamine at IMGT positions 54, 86, and 90, respectively,

of IGHV3-33.

[0045] In some embodiments, the anti-human DII4 antibody comprises one or more amino acid substitutions in one or more CDRs relative to the canonical light chain variable region, which is reasonably expected to result in an altered charge distribution across the exposed surface of the antibody, and therefore affect its interaction with the solvent environment. In some embodiments, the amino acid substitution comprises the substitution of histidine for glutamine at IMGT position 106 of IGKV3-11.

[0046] According to certain embodiments of the present invention, the anti-human DII4 antibody, or antigen-binding fragment thereof, comprises a heavy chain complementary determining region (HCDR) 1 of SEQ ID NO: 2, an HCDR2 of SEQ ID NO: 3, and an HCDR3 of SEQ ID NO: 4. In certain embodiments, the anti-human DII4 antibody, or antigen-binding fragment thereof, comprises an HCVD of SEQ ID NO: 1.

[0047] According to certain embodiments of the present invention, the anti-human DII4, or antigen-binding fragment thereof, comprises a light (kappa) chain complementary determining region (LCDR) 1 of SEQ ID NO: 6, an LCDR2 of SEQ ID NO: 7, and an LCDR3 of SEQ ID NO: 8. In certain embodiments, the anti-human DII4 antibody, or antigen-binding fragment thereof, comprises an LCVD of SEQ ID NO: 5.

[0048] The non-limiting, exemplary antibody used in the Examples herein is referred to as REGN421, as in US 7,488,806, US 7,534,868, US 7,919,593, and US 2011-0150905. This antibody comprises an HCVR/LCVR amino acid sequence pair having SEQ ID NOs: 1/5, and HCDR1-HCDR2-HCDR3 / LCDR1-LCDR2-LCDR3 domains represented by SEQ ID NOs: 2 - 3 - 4 / SEQ ID NOs: 6 - 7 - 8.

[0049] The amount of antibody, or antigen-binding fragment thereof, contained within the pharmaceutical formulations may vary depending on the specific properties desired of the formulations, as well as the particular circumstances and purposes for which the formulations are intended to be used. In certain embodiments, the pharmaceutical formulations are liquid formulations that may contain 20 mg/mL $\pm$ 3 mg/mL to 75 mg/mL $\pm$ 11.25 mg/mL of antibody; 25 $\pm$ 3.75 mg/mL to 70 $\pm$ 10.5 mg/mL of antibody; 30 $\pm$ 4.5 mg/mL to 65 $\pm$ 9.75 mg/mL of antibody; 35 $\pm$ 5.25 mg/mL to 60 $\pm$ 9 mg/mL of antibody; 40 $\pm$ 6 mg/mL to 55 $\pm$ 8.25 mg/mL of antibody; 45 $\pm$ 6.75 mg/mL to 50 $\pm$ 7.5 mg/mL of antibody; 25 mg/mL $\pm$ 3.75 mg/mL; about 25 mg/mL; 25 mg/mL; 50 mg/mL $\pm$ 7.5 mg/mL; about 50 mg/mL; or 50 mg/mL of an antibody or an antigen-binding fragment thereof, that binds specifically to human DII4.

EXCIPIENTS AND PH

[0050] The pharmaceutical formulations comprise one or more excipients. The term "excipient", as used herein, means any non-therapeutic agent added to the formulation to provide a desired consistency, viscosity or stabilizing effect.

[0051] In certain embodiments, the pharmaceutical formulation comprises at least one organic cosolvent in a type and in an amount that stabilizes the human DII4 antibody under conditions of rough handling or agitation, such as, *e.g.*, vortexing. In some embodiments, what is meant by "stabilizes" is the maintenance of at least 95% of the DII4 antibody in its native state, *i.e.*, not fragmented or aggregated, of antibody (on a molar basis) over the course of rough handling, such as by vortexing the antibody-organic cosolvent solution for about 60 minutes or about 120 minutes.

[0052] In certain embodiments, the organic cosolvent is a non-ionic surfactant, such as an alkyl poly(ethylene oxide). Specific non-ionic surfactants that can be included in the formulations of the present invention include, *e.g.*, polysorbates such as polysorbate 20, polysorbate 28, polysorbate 40, polysorbate 60, polysorbate 65, polysorbate 80, polysorbate 81, and polysorbate 85; poloxamers such as poloxamer 181, poloxamer 188, poloxamer 407; or polyethylene glycol (PEG). Polysorbate 20 is also known as TWEEN 20, sorbitan monolaurate and polyoxyethylenesorbitan monolaurate. Poloxamer 188 is also known as PLURONIC F68.

[0053] The amount of non-ionic surfactant contained within the pharmaceutical formulations may vary depending on the specific properties desired of the formulations, as well as the particular circumstances and purposes for which the formulations are intended. In certain embodiments, the formulations may contain 0.01% $\pm$ 0.0015% to 1% $\pm$ 0.15% surfactant. For example, the formulations of the present invention may comprise about 0.0085%; about 0.01%; about 0.02%; about 0.03%; about 0.04%; about 0.05%; about 0.06%; about 0.07%; about 0.08%; about 0.09%; about 0.1%; about 0.11%; about 0.12%; about 0.13%; about 0.14%; about 0.15%; about 0.16%; about 0.17%; about 0.18%; about 0.19%; about 0.20%; about 0.21%; about 0.22%; about 0.23%; about 0.24%; about 0.25%; about 0.3%; about 0.4%; about 0.5%; about 0.6%; about 0.7%; about 0.8%; about 0.9%; about 1%; about 1.1%; about 1.15%; or about 1.2% polysorbate 20 or poloxamer 188.

[0054] The pharmaceutical also comprise one or more stabilizers in a type and in an amount that stabilizes the human DII4 antibody under conditions of thermal stress. In some embodiments, what is meant by "stabilizes" is maintaining greater than about 94% of the antibody in a native conformation when the solution containing the antibody and the thermal stabilizer is kept at about 45°C for up to about 28 days. In some embodiments, what is meant by "stabilizes" is maintaining greater than about 96% of the antibody in a native conformation when the solution containing the antibody and the thermal stabilizer is kept at about 37°C for up to about 28 days. As used herein, "native" means the major form of the antibody by size exclusion, which is generally an intact monomer of the antibody.

[0055] In certain embodiments, the thermal stabilizer is a sugar or sugar alcohol selected from sucrose, sorbitol, glycerol, trehalose and mannitol, or any combination thereof, the amount of which contained within the formulation can vary depending on the specific circumstances and intended purposes for which the formulation is used. In certain embodiments, the formulations may contain about 5% to about 40% sugar or sugar alcohol; about 1% to about 20% sugar or sugar alcohol; about 5 % to about 15% sugar or sugar alcohol; about 7.5% to about 12.5% sugar or sugar alcohol; about 10% sugar or sugar alcohol; $10\% \pm 1.5\%$ sugar or sugar alcohol; or 10% sugar or sugar alcohol. For example, the pharmaceutical formulations of the present invention may comprise $4\% \pm 0.6\%$; $5\% \pm 0.75\%$; $6\% \pm 0.9\%$; $7\% \pm 1.05\%$; $8\% \pm 1.2\%$; $9\% \pm 1.35\%$; $10\% \pm 1.5\%$; $11\% \pm 1.65\%$; $12\% \pm 1.8\%$; $13\% \pm 1.95\%$; or about $14\% \pm 2.1\%$ sugar or sugar alcohol (e.g., sucrose, trehalose or mannitol).

[0056] The pharmaceutical formulations may also comprise a buffer or buffer system, which serves to maintain a stable pH and to help stabilize the human DII4 antibody. In some embodiments, what is meant by "stabilizes" is wherein at least 94% of the antibody is in its native conformation as determined by size exclusion chromatography when the solution containing the antibody and the buffer is kept at about 45°C for up to about 28 days. By "native" or "native conformation", what is meant is the antibody fraction that is not aggregated or degraded. This is generally determined by an assay that measures the relative size of the antibody entity, such as a size exclusion chromatographic assay. The non-aggregated and non-fragmented antibody elutes at a fraction that equates to the native antibody, and is generally the main elution fraction. Aggregated antibody elutes at a fraction that indicates a size greater than the native antibody. Fragmented antibody elutes at a fraction that indicates a size less than the native antibody.

[0057] In some embodiments, what is meant by "stabilizes" is wherein at least 45% of the antibody is in its main charge form as determined by cation exchange chromatography when the solution containing the antibody and the buffer is kept at about 45°C for up to about 28 days. By "main charge" or "main charge form", what is meant is the fraction of antibody that elutes from an ion exchange resin in the main peak, which is generally flanked by more "basic" peaks on one side and more "acidic" peaks on the other side.

[0058] The pharmaceutical formulations may have a pH of from about 5.5 to about 6.5. For example, the formulations of the present invention may have a pH of about 5.5; about 5.6; about 5.7; about 5.8; about 5.9; about 6.0; about 6.1; about 6.2; about 6.3; about 6.4; or about 6.5. In some embodiments, the pH is 6.0 ± 0.5 ; 6.0 ± 0.4 ; 6.0 ± 0.3 ; 6.0 ± 0.2 ; 6.0 ± 0.1 ; about 6.0; or 6.0.

[0059] In some embodiments, the buffer or buffer system comprises at least one buffer that has a buffering range that overlaps fully or in part the range of pH 5.5 - 7.4, such as, e.g., a buffer having a useful buffering range of pH 4.8 to pH 8.8. In one embodiment, the buffer has a pKa of about 7.21. In certain embodiments, the buffer comprises a phosphate buffer. In certain embodiments, the phosphate is present at a concentration of $5 \text{ mM} \pm 0.75 \text{ mM}$ to $15 \text{ mM} \pm 2.25 \text{ mM}$; $6 \text{ mM} \pm 0.9 \text{ mM}$ to $14 \text{ mM} \pm 2.1 \text{ mM}$; $7 \text{ mM} \pm 1.05 \text{ mM}$ to $13 \text{ mM} \pm 1.95 \text{ mM}$; $8 \text{ mM} \pm 1.2 \text{ mM}$ to $12 \text{ mM} \pm 1.8 \text{ mM}$; $9 \text{ mM} \pm 1.35 \text{ mM}$ to $11 \text{ mM} \pm 1.65 \text{ mM}$; $10 \text{ mM} \pm 1.5 \text{ mM}$; or about 10 mM. In certain embodiments, the buffer system comprises phosphate at $10 \text{ mM} \pm 1.5 \text{ mM}$, at a pH of 6.0 ± 0.5 .

EXEMPLARY FORMULATIONS

[0060] According to one aspect the liquid pharmaceutical formulation has low viscosity (i.e, less than 10 cPoise, or about 1.5 cPoise) and has an osmolality between 600 and 700 mOsm, or about 650 mOsm; and comprises: (i) $25 \text{ mg/mL} \pm 3.75 \text{ mg/mL}$, or $50 \text{ mg/mL} \pm 7.5 \text{ mg/mL}$ of a human antibody that specifically binds to human DII4 (e.g., REGN421); (ii) a buffer system that buffers at about $\text{pH } 6.0 \pm 0.5$; (iii) a thermal stabilizer comprising a sugar and a salt, which serves as a; and (iv) an organic cosolvent.

[0061] According to one embodiment, the pharmaceutical formulation comprises: (i) $20 \pm 3 \text{ mg/mL}$ to $60 \pm 9 \text{ mg/mL}$ human IgG1 antibody that specifically binds to human DII4 and which comprises a substituted IGHV3-33 type heavy chain variable region and a substituted IGKV3-11 type light chain variable region; (ii) a phosphate buffer, which buffers at $\text{pH } 6.0 \pm 0.5$; (iii) sucrose and sodium chloride; and (iv) a non-ionic detergent, such as a polysorbate.

[0062] According to one embodiment, the pharmaceutical formulation comprises: (i) $25 \text{ mg/mL} \pm 3.75 \text{ mg/mL}$ human IgG1 antibody that specifically binds to human DII4, and which comprises an HCDR1 of SEQ ID NO: 2, an HCDR2 of SEQ ID NO: 3, an HCDR3 of SEQ ID NO: 4, an LCDR1 of SEQ ID NO: 6, an LCDR2 of SEQ ID NO: 7, and an LCDR3 of SEQ ID NO: 8; (ii) $10 \text{ mM} \pm 1.5 \text{ mM}$ phosphate, $\text{pH } 6.0 \pm 0.5$; (iii) $10\% \pm 1.5\%$ sucrose; (iv) $150 \text{ mM} \pm 22.5 \text{ mM}$ sodium chloride; and (v) $0.2\% \pm 0.03\%$ polysorbate 20.

[0063] According to one embodiment, the pharmaceutical formulation comprises: (i) $50 \text{ mg/mL} \pm 7.5 \text{ mg/mL}$ human IgG1 antibody that specifically binds to human DII4, and which comprises an HCDR1 of SEQ ID NO: 2, an HCDR2 of SEQ ID NO: 3, an HCDR3 of SEQ ID NO: 4, an LCDR1 of SEQ ID NO: 6, an LCDR2 of SEQ ID NO: 7, and an LCDR3 of SEQ ID NO: 8; (ii) $10 \text{ mM} \pm 1.5 \text{ mM}$ phosphate, $\text{pH } 6.0 \pm 0.5$; (iii) $10\% \pm 1.5\%$ sucrose; (iv) $150 \text{ mM} \pm 22.5 \text{ mM}$ sodium chloride; and (v) $0.2\% \pm 0.03\%$ polysorbate 20.

[0064] According to one embodiment, the pharmaceutical formulation comprises: (i) $25 \text{ mg/mL} \pm 3.75 \text{ mg/mL}$ human IgG1 antibody that specifically binds to human DII4, and which comprises a heavy chain variable domain of SEQ ID NO:

1, and a light chain variable domain of SEQ ID NO: 5, at a concentration of; (ii) 10 mM $\pm$ 1.5 mM phosphate pH 6.0 $\pm$ 0.5; (iii) 10% $\pm$ 1.5% sucrose; (iv) 150 mM $\pm$ 22.5 mM sodium chloride; and (v) 0.2% $\pm$ 0.03% polysorbate 20.

[0065] According to one embodiment, the pharmaceutical formulation comprises: (i) 50 mg/ml $\pm$ 7.5 mg/mL human IgG1 antibody that specifically binds to human DII4, and which comprises a heavy chain variable domain of SEQ ID NO: 1, and a light chain variable domain of SEQ ID NO: 5; (ii) 10 mM $\pm$ 1.5 mM phosphate, pH 6.0 $\pm$ 0.5; (iii) 10% $\pm$ 1.5% sucrose; (iv) 150 mM $\pm$ 22.5 mM sodium chloride; and (v) 0.2% $\pm$ 0.03% polysorbate 20.

[0066] Additional non-limiting examples of pharmaceutical formulations encompassed by the present invention are set forth elsewhere herein, including the working Examples presented below.

STABILITY AND VISCOSITY OF THE PHARMACEUTICAL FORMULATIONS

[0067] The pharmaceutical formulations of the present invention typically exhibit high levels of stability. The term "stable", as used herein in reference to the pharmaceutical formulations, means that the antibodies within the pharmaceutical formulations retain an acceptable degree of chemical structure or biological function after storage under defined conditions. A formulation may be stable even though the antibody contained therein does not maintain 100% of its chemical structure or biological function after storage for a defined amount of time. Under certain circumstances, maintenance of at least about 90%, 95%, 96%, 97%, 98% or 99% of an antibody's structure or function after storage for a defined amount of time may be regarded as "stable".

[0068] Stability can be measured, *inter alia*, by determining the percentage of native antibody that remains in the formulation after storage for a defined amount of time at a defined temperature. The percentage of native antibody can be determined by, *inter alia*, size exclusion chromatography (e.g., size exclusion high performance liquid chromatography [SE-HPLC]), such that native means non-aggregated and non-fragmented. An "acceptable degree of stability", as that phrase is used herein, means that at least 90% of the native form of the antibody can be detected in the formulation after storage for a defined amount of time at a given temperature. In certain embodiments, at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the native form of the antibody can be detected in the formulation after storage for a defined amount of time at a defined temperature. The defined amount of time after which stability is measured can be at least 14 days, at least 28 days, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, at least 7 months, at least 8 months, at least 9 months, at least 10 months, at least 11 months, at least 12 months, at least 18 months, at least 24 months, or more. The defined temperature at which the pharmaceutical formulation may be stored when assessing stability can be any temperature from about -80°C to about 45°C, e.g., storage at about -80°C, about -30°C, about -20°C, about 0°C, about 4°-8°C, about 5°C. Thermal stress may be applied to the pharmaceutical formulation to assess stability. Thermal stress includes for example holding the formulation for about 14 days or about 28 days at about 25°C, about 35°C, about 37°C, or about 45°C.

[0069] For example, a pharmaceutical formulation may be deemed stable if after six months of storage at 5°C, at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 98.5%, 99% or 99.5% of antibody detected by SE-HPLC is native (*i.e.*, in the native peak fraction). A pharmaceutical formulation may also be deemed stable if after six months of storage at 25°C at least 97%, 97.5%, 98%, 98.5%, 99% or 99.5% of antibody detected by SE-HPLC is native. A pharmaceutical formulation may also be deemed stable if after 28 days of storage at 37°C, at least 96%, 96.5%, 97%, 97.5%, 98%, 98.5%, 99% or 99.5% of antibody detected by SE-HPLC is native. A pharmaceutical formulation may also be deemed stable if after 28 days of storage at 45°C at least 94%, 95%, 96%, 97%, 98% or 99% of antibody detected by SE-HPLC is native. A pharmaceutical formulation may also be deemed stable if after six months of storage at -20°C at least 98.5%, 99% or 99.5% of antibody detected by SE-HPLC is native. A pharmaceutical formulation may also be deemed stable if after 18 months of storage at -30°C at least 99% or 99.5% of antibody detected by SE-HPLC is native. A pharmaceutical formulation may also be deemed stable if after six months of storage at -80°C at least 99% or 99.5% of antibody detected by SE-HPLC is native.

[0070] Stability can be measured, *inter alia*, by determining the percentage of antibody that migrates in the main fraction of antibody ("main charge form") during ion exchange, relative to the total combined peak area, wherein stability is proportional to the fraction of antibody in the main charge form. While not wishing to be bound by theory, deamidation of the antibody may cause the antibody to become more negatively charged and thus more acidic relative to the non-deamidated antibody (see, e.g., Robinson, N., Protein Deamidation, PNAS, April 16, 2002, 99(8):5283-5288). An "acceptable degree of stability", as that phrase is used herein, means that at least 53% of the antibody is in a main charge form detected in the formulation after storage for a defined amount of time at a defined temperature. In certain embodiments an acceptable degree of stability means that at least about 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% of the antibody can be detected in a main charge form after storage for a defined amount of time at a given temperature, or under thermal, freeze/thaw, or agitation stress. The defined amount of time after which stability is measured can be at least 2 weeks, at least 28 days, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, at least 7 months, at least 8 months, at least 9 months, at least 10 months, at least 11 months, at least 12 months, at least 18 months, at least 24 months,

or more. The temperature at which the pharmaceutical formulation may be stored when assessing stability can be any temperature from about -80°C to about 45°C, e.g., storage at about -80°C, about -30°C, about -20°C, about 0°C, about 4°C, about 5°C, about 25°C, about 37°C, or about 45°C. For example, a pharmaceutical formulation may be deemed stable if after three months of storage at -80°C, -30°C, or -20°C no less than about 57%, 58%, 59%, 60%, or 61% of the antibody is in a main charge form. A pharmaceutical formulation may also be deemed stable if after six months of storage at 5°C, no less than about 59%, 59.1%, 59.2%, 59.3%, 59.4%, 59.5%, 59.6%, 59.7%, 59.8%, 59.9%, 60%, 60.1%, 60.2%, 60.3%, 60.4%, 60.5%, 60.6%, 60.7%, 60.8%, 60.9%, 61%, or 61.1% of the antibody is in a main charge form. A pharmaceutical formulation may also be deemed stable if after six months of storage at 25°C, no less than about 53% of the antibody is in a main charge form. A pharmaceutical formulation may also be deemed stable if after 28 days of storage at 37°C, no less than about 50%, 50.5%, 51%, or 51.5% of the antibody is in a main charge form. A pharmaceutical formulation may also be deemed stable if after 28 days of storage at 45°C, no less than about 45%, 45.1%, 45.2%, 45.3%, 45.4%, 45.5%, 45.6%, 45.7%, 45.8%, 45.9%, or 50% of the antibody can be detected in a main charge form.

[0071] Measuring the binding affinity of the antibody to its target may also be used to assess stability. For example, a formulation of the present invention may be regarded as stable if, after storage at e.g., -80°C, -30°C, -20°C, 5°C, 25°C, 37°C, 45°C, etc. for a defined amount of time (e.g., 14 days to 24 months), the anti-Dll4 antibody contained within the formulation maintains at least 50% and up to about 150% of the potency, as measured as binding affinity, of the antibody prior to said storage. Binding affinity may be determined by e.g., ELISA or plasmon resonance. Biological activity may be determined by a Dll4 activity assay, such as e.g., by contacting a cell that expresses Dll4 with the formulation comprising the anti Dll4 antibody. The binding of the antibody to such a cell may be measured directly, such as *via* FACS analysis. Alternatively, the downstream activity of the Dll4/Notch signalling pathway may be measured in the presence of the antibody, and compared to the activity of the Dll4/Notch signaling pathway in the absence of antibody. In some embodiments, the Dll4 may be endogenous to the cell. In other embodiments, the Dll4 may be ectopically (heterologously) expressed in the cell.

[0072] Additional methods for assessing the stability of an antibody in formulation are demonstrated in the Examples presented below.

CONTAINERS AND METHODS OF ADMINISTRATION

[0073] The pharmaceutical formulations of the present invention may be contained within any container suitable for storage or administration of medicines and other therapeutic compositions. For example, the pharmaceutical formulations may be contained within a sealed and sterilized plastic or glass container having a defined volume such as a vial, ampule, syringe, cartridge, bottle, or IV bag. Different types of vials can be used to contain the formulations of the present invention including, e.g., clear and opaque (e.g., amber) glass or plastic vials. Likewise, any type of syringe can be used to contain or administer the pharmaceutical formulations of the present invention.

[0074] The pharmaceutical formulations of the present invention may be contained within "normal tungsten" syringes or "low tungsten" syringes. As will be appreciated by persons of ordinary skill in the art, the process of making glass syringes generally involves the use of a hot tungsten rod which functions to pierce the glass thereby creating a hole from which liquids can be drawn and expelled from the syringe. This process results in the deposition of trace amounts of tungsten on the interior surface of the syringe. Subsequent washing and other processing steps can be used to reduce the amount of tungsten in the syringe. As used herein, the term "normal tungsten" means that the syringe contains greater than or equal to 500 parts per billion (ppb) of tungsten. The term "low tungsten" means that the syringe contains less than 500 ppb of tungsten. For example, a low tungsten syringe, according to the present invention, can contain less than about 490, 480, 470, 460, 450, 440, 430, 420, 410, 390, 350, 300, 250, 200, 150, 100, 90, 80, 70, 60, 50, 40, 30, 20, 10 or fewer ppb of tungsten.

[0075] The rubber plungers used in syringes, and the rubber stoppers used to close the openings of vials, may be coated to prevent contamination of the medicinal contents of the syringe or vial, or to preserve their stability. Thus, pharmaceutical formulations of the present invention, according to certain embodiments, may be contained within a syringe that comprises a coated plunger, or within a vial that is sealed with a coated rubber stopper. For example, the plunger or stopper may be coated with a fluorocarbon film. Examples of coated stoppers or plungers suitable for use with vials and syringes containing the pharmaceutical formulations of the present invention are mentioned in, e.g., U.S. Patent Nos. 4,997,423; 5,908,686; 6,286,699; 6,645,635; and 7,226,554. Particular exemplary coated rubber stoppers and plungers that can be used in the context of the present invention are commercially available under the tradename "FluroTec®", available from West Pharmaceutical Services, Inc. (Lionville, PA). FluroTec® is an example of a fluorocarbon coating used to minimize or prevent drug product from adhering to the rubber surfaces.

[0076] According to certain embodiments of the present invention, the pharmaceutical formulations may be contained within a low tungsten syringe that comprises a fluorocarbon-coated plunger.

[0077] The pharmaceutical formulations can be administered to a patient by parenteral routes such as injection (e.g., subcutaneous, intravenous, intramuscular, intraperitoneal, etc.) or percutaneous, mucosal, nasal, pulmonary or oral

administration. Numerous reusable pen or autoinjector delivery devices can be used to subcutaneously deliver the pharmaceutical formulations of the present invention. Examples include, but are not limited to AUTOPEN™ (Owen Mumford, Inc., Woodstock, UK), DISETRONIC™ pen (Disetronic Medical Systems, Bergdorf, Switzerland), HUMALOG MIX 75/25™ pen, HUMALOG™ pen, HUMALIN 70/30™ pen (Eli Lilly and Co., Indianapolis, IN), NOVOPEN™ I, II and III (Novo Nordisk, Copenhagen, Denmark), NOVOPEN JUNIOR™ (Novo Nordisk, Copenhagen, Denmark), BD™ pen (Becton Dickinson, Franklin Lakes, NJ), OPTIPEN™, OPTIPEN PRO™, OPTIPEN STARLET™, and OPTICLIK™ (sanofi-aventis, Frankfurt, Germany). Examples of disposable pen or autoinjector delivery devices having applications in subcutaneous delivery of a pharmaceutical composition of the present invention include, but are not limited to the SOLOSTAR™ pen (sanofi-aventis), the FLEXPEN™ (Novo Nordisk), and the KWIKPEN™ (Eli Lilly), the SURECLICK™ Autoinjector (Amgen, Thousand Oaks, CA), the PENLET™ (Haselmeier, Stuttgart, Germany), the EPIPEN (Dey, L.P.), and the HUMIRA™ Pen (Abbott Labs, Abbott Park, IL).

[0078] The use of a microinfusor to deliver the pharmaceutical formulations of the present invention is also contemplated herein. As used herein, the term "microinfusor" means a subcutaneous delivery device designed to slowly administer large volumes (e.g., up to about 2.5 mL or more) of a therapeutic formulation over a prolonged period of time (e.g., about 10, 15, 20, 25, 30 or more minutes). See, e.g., U.S. 6,629,949; US 6,659,982; and Meehan et al., J. Controlled Release 46:107-116 (1996). Microinfusors are particularly useful for the delivery of large doses of therapeutic proteins contained within high concentration (e.g., about 100, 125, 150, 175, 200 or more mg/mL) or viscous solutions.

[0079] In one embodiment, the pharmaceutical formulation is administered via an IV drip, such that the formulation is diluted in an IV bag containing a physiologically acceptable solution. In one embodiment, the pharmaceutical composition is a compounded sterile preparation in an intravenous infusion bag, such that a single dose of drug product is diluted into 100 mL, 250 mL (or other like amount suitable for intravenous drip delivery) of a physiological buffer (e.g., 0.9% saline). In some embodiments, the infusion bag is made of a polyvinyl chloride (e.g., VIAFLEX, Baxter, Deerfield, Illinois). In some embodiments, the infusion bag is made of a polyolefin (EXCEL IV Bags, Braun Medical Inc., Bethlehem, Pennsylvania).

THERAPEUTIC USES OF THE PHARMACEUTICAL FORMULATIONS

[0080] The pharmaceutical formulations of the present invention are useful, *inter alia*, for the treatment, prevention or amelioration of any disease or disorder associated with Dll4 activity, including diseases or disorders mediated by Dll4 or the Notch signaling pathway. Exemplary, non-limiting diseases and disorders that can be treated or prevented by the administration of the pharmaceutical formulations of the present invention include various diseases involving angiogenesis, which is the biological process whereby new blood vessels are formed. Aberrant angiogenesis is associated with several disease conditions including, e.g., proliferative retinopathies, rheumatoid arthritis and psoriasis. In addition, it is well established that angiogenesis is critical for tumor growth and maintenance.

EXAMPLES

[0081] The following examples are presented so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by mole, molecular weight is average molecular weight, percent concentration (%) means the mass of the solute in grams divided by the volume of the solution in milliliters times 100% (e.g., 10% sucrose means 0.1 gram of sucrose per milliliter of solution), temperature is in degrees Centigrade, and pressure is at or near atmospheric pressure.

[0082] Initial formulation development activities involved screening organic cosolvents, thermal stabilizers, and buffers in liquid and lyophilized formulations of anti-Dll4 antibodies to identify excipients that are compatible with the protein and enhance its stability, while maintaining near physiologic osmolality and low viscosity for intravenous and subcutaneous injection. Buffer conditions were also examined to determine the optimal pH for maximum protein stability.

EXAMPLE 1: EXEMPLARY ANTI-DLL4 FORMULATION

[0083] Formulation development activities included the screening of buffers, organic cosolvents, and thermal stabilizers in liquid formulations of the anti-Dll4 antibody to identify excipients that enhance the stability of the protein. Buffer conditions were also examined to determine the optimal pH for maximum protein stability. Results generated from these studies were used to develop a stable, liquid formulation suitable for clinical use. Anti-Dll4 (e.g., REGN421) was formulated at 25 ± 3.75 mg/ml or 50 ± 7.5 mg/ml. In one embodiment, the anti-Dll4 antibody is formulated in 10 ± 1.5 mM phosphate (pH 6.0 ± 0.5), $0.2\% \pm 0.03\%$ polysorbate 20, $10\% \pm 1.5\%$ sucrose, and $150 \text{ mM} \pm 22.5 \text{ mM}$ sodium chloride. The

phosphate buffer, which is a combination of monobasic phosphate and dibasic phosphate, was selected for its buffering capacity at the optimal pH for anti-DII4 stability and low rate of aggregation. Anti-DII4 antibody REGN421 was observed to have optimum stability and the lowest rate of aggregation and formation of charge variants at pH 6.0. Polysorbate 20 was selected as the agitation stabilizing agent, which reduced the rate of aggregation and precipitation when the antibody was agitated and otherwise handled. Sucrose, which was observed to reduce the rate of aggregation of the antibody in solution, was selected as a component of the thermal stabilizing agent. Sodium chloride, which was observed to reduce the formation of charged species of the antibody in solution, was selected as another component of the thermal stabilizing agent.

EXAMPLE 2: STRESS STABILITY STUDY METHODOLOGY

[0084] Optical density (405 nm) and reverse-phase high-performance liquid chromatography (RP-HPLC) were used in development studies to assess the physical stability of the formulated anti-DII4 antibody. Physical stability is defined as the recovery of soluble forms of the antibody in solution. Loss of antibody could be due to either precipitation or adsorption. The presence of particulates can be detected by visual inspection or by optical density (OD) measurements at 405 nm (turbidity measurements), where an increase in OD indicates an increase in turbidity due to the formation of particulates. The presence of particulates as determined visually or by OD measurements indicates that the sample has failed to maintain stability. Recovery of anti-DII4 antibody was measured either by RP-HPLC and by UV spectrophotometry at 280 nm. The concentration of each test sample is determined from the area of the eluted antibody peak compared to a standard curve generated using antibody of defined concentration.

[0085] Size exclusion-HPLC (SE-HPLC) was used to assess the purity of the anti-DII4 antibody and to provide a measure of the relative amounts of antibody aggregates, fragments and impurities that differ in molecular size from the subject anti-DII4 antibody. Purified anti-DII4 antibody generally eluted as a small aggregate peak and a large native dimer peak followed by a small low molecular weight peak. This method employed a high-resolution silica-based size exclusion column and a phosphate - sodium chloride buffer. Absorbance at 215 nm was recorded and integrated for purity determination. The purity specification for the anti-DII4 antibody by SE-HPLC required that the area of the main peak was $\geq 90.0\%$ of the total peak area. The percentages of the total peak area for the high and low molecular weight peaks areas are reported in the Tables.

[0086] Antibody charge variants were analyzed by cation exchange chromatography using a weak cation exchange column from Dionex (PROPAC® WCX-10, 4 x 250 mm) over a shallow salt gradient in phosphate buffer at pH 6.0 (40 mM to 120 mM NaCl over 44 minutes at 1mL/ min). Absorbance at 215 nm was recorded and integrated to assess charge heterogeneity. Chromatograms of anti-DII4 antibody were integrated by dividing the peaks into three major charge groups: acidic, main and basic. The group of peaks that eluted first have been designated as the acidic species (this was determined by analyzing material exposed to high pH and temperature). The major peak in the chromatogram has been designated as the main peak, main charge form, or major charge variant. The final group of peaks was reported as the basic species. The designation was made based on the extended column retention most likely due to the high level of positive charge. The peak pattern of the antibody test sample was compared to the peak pattern of the reference standard and percent peak areas of the acidic, main and basic species were reported.

[0087] The activity or potency of the anti-DII4 antibody was assessed using a binding assay. The ability of that antibody to block the binding of biotinylated-hDII4-hFc to Notch1 was measured using a specific and sensitive ELISA. The ELISA measured unbound biotinylated hDII4-hFc in a solution composed of a mixture of biotin-hDII4-hFc and anti-DII4 antibody. The mixtures were added to microtiter plates coated with human Notch1. A fixed amount of biotin-hDII4-hFc was titrated with various amounts of antibody; and the antibody:biotin hDII4 hFc complex was washed off. The amount of biotin-hDII4-hFc remaining bound to the plate was detected using Streptavidin conjugated to horseradish peroxidase (HRP). The analyzed samples were compared to a reference standard to determine relative potency. HRP colorimetric substrates were used for detection and OD450 nm was measured. The OD450 nm values were plotted against the concentrations of antibody in the solutions and an IC_{50} was determined using non-linear regression analysis. The IC_{50} was defined as the antibody concentration required to block 50% of the maximal binding of biotin-hDII4-hFc to hNotch1-hFc coated plates, and these values reflect the blocking potency of the antibody against hDII4. The potency was determined by comparison to a reference standard sample run on the same plate and expressed as a percent of the reference standard IC_{50} value.

EXAMPLE 3: BUFFER AND PH

[0088] The effect of pH and buffer on the stability of anti-DII4 was considered. REGN421 antibody ("REGN421") was the most stable when formulated with either a 10 mM phosphate buffer or a 10 mM histidine buffer. Phosphate was chosen as the buffer for formulation due to concerns over the potential risk of histidine oxidation during storage in a liquid state. The mixture of monobasic phosphate and dibasic phosphate results in a pH of 6.0 when all excipients are

added together with the REGN421 antibody.

[0089] Analysis of buffered REGN421 incubated at elevated temperatures revealed that the main protein degradation pathways were the formation of aggregates, cleavage products, and charge variants. Therefore, to assess the effect of pH and buffer on REGN421 stability, REGN421 test samples were incubated at 45°C in various buffers at pH values ranging from 5.0 to 8.0 to identify conditions that increase the thermal stability of the protein (Table 1). REGN421 had maximum stability, as determined by both SE-HPLC and Cation Exchange- High Performance Liquid Chromatography (CEX-HPLC) analysis, when the protein was formulated at pH 6.0. Formulating REGN421 in phosphate or histidine buffer at pH 6.0 resulted in greater protein stability compared to formulating at pH 6.0 in succinate or citrate buffer. Phosphate was chosen for the REGN421 formulation because of concerns over the potential risk of histidine oxidation during long term storage in a liquid state.

[0090] For the results shown in Table 1, 0.3 mL of 25 mg/mL REGN421 in 10 mM test buffer containing 0.2% polysorbate 20 in a 2 mL Type 1 borosilicate glass vial with a FluroTec® coated 4432/50 butyl rubber stopper were tested for 28 days at 45. Turbidity was reported as the change in OD₄₀₅ relative compared to starting material. Starting material results represent the average results of the starting material for all 14 formulations. OD = Optical density; RP-HPLC = Reverse phase high performance liquid chromatography; SE-HPLC = Size exclusion high performance liquid chromatography; CEX-HPLC = cation exchange high performance liquid chromatography.

Table 1: Effect of Buffer and pH on the Stability of REGN421 Incubated at 45°C for 28 Days

Buffer and pH	Visual Appearance	Turbidity (OD 405 nm)	% Total REGN421 Recovered (RP-HPLC)	% Native REGN421 Recovered (SE-HPLC)	% Native REGN421 Recovered (CEX-HPLC)
Starting Material* (no 45°C incubation)	Pass	0.00	100	97.6	58.5
pH 8.0, Tris	Fail	0.30	107	85.6	21.0
pH 8.0, Phosphate	Fail	0.27	100	65.7	5.6
pH 7.5, Phosphate	Fail	0.12	104	80.1	14.6
pH 7.0, Phosphate	Fail	0.03	104	83.4	24.9
pH 6.5, Phosphate	Pass	0.00	100	93.1	35.3
pH 6.0, Phosphate	Pass	0.00	104	93.9	38.3
pH 6.0, Histidine	Pass	0.00	103	94.4	41.6
pH 6.0, Succinate	Pass	0.01	102	92.3	40.7
pH 6.0, Citrate	Pass	0.01	99	92.4	36.7
pH 5.5, Citrate	Fail	0.01	105	91.1	34.6
pH 5.0, Citrate	Fail	0.02	103	86.7	26.9
pH 5.0, Acetate	Pass	0.01	107	92.3	34.6

EXAMPLE 4: SELECTION OF PROTECTANTS AGAINST AGITATION STRESS

[0091] 0.2% polysorbate 20 was chosen as the organic cosolvent because it stabilized REGN421 when agitated in the liquid state. Studies that were initially performed to monitor the stability of REGN421 following agitation in the absence of organic cosolvents yielded varying results. Analysis of agitated REGN421 via SE-HPLC demonstrated an increase in aggregation of the antibody in one particular study, but no significant increase in aggregation was observed in the majority of other agitation studies performed on REGN421. Results from two representative stability studies are shown in Table 2. The conservative approach of adding polysorbate 20 to the REGN421 formulation was nonetheless taken

to prevent potential agitation dependent protein instability that may occur during the manufacture, shipping, and handling of REGN421. Polysorbate 20 was chosen over other organic cosolvents because it did not change the thermal stability of REGN421.

[0092] For the antibody stability results shown in Table 2, 0.3 mL of 25 mg/mL REGN421, in 10 mM phosphate, pH 6.0 in a 2 mL Type 1 borosilicate glass vial with a FluroTec® coated 4432/50 butyl rubber stopper was combined with the organic cosolvents and subjected to 120 minutes of vortexing. Turbidity was reported as the change in OD₄₀₅ relative compared to starting material. Starting material results represent the average results of the starting material for all nine formulations. OD = Optical density; RP-HPLC = Reverse phase high performance liquid chromatography; SE-HPLC = Size exclusion high performance liquid chromatography.

Table 2: Effect of Organic Cosolvents on Antibody Stability - Agitation

Organic Cosolvent	Stress	Visual Appearance	% Total REGN421 Recovered (RP-HPLC)	% Native REGN421 Recovered (SE-HPLC)	% REGN421 Aggregate Recovered (SE-HPLC)
Starting Material*	None	Pass	100	96.6	1.5
No Cosolvent	Vortex 120min	Pass	101	94.9	3.0
Starting Material *	None	Pass	100	97.8	0.7
No Cosolvent	Vortex 120min	Pass	101	97.4	0.9
0.2% Polysorbate 20	Vortex 120min	Pass	99	98.0	0.7
0.2% Polysorbate 80	Vortex 120min	Pass	100	97.9	0.8
0.2% PEG 3350	Vortex 120min	Pass	101	98.0	0.7

EXAMPLE 5: SELECTION OF PROTECTANTS AGAINST THERMAL STRESS

[0093] Stabilizers such as sugars, amino acids, and inorganic salts were examined for their ability to increase the thermal stability of REGN421. A summary of the thermal stabilizers that were examined is presented in Table 3. Formulations containing sucrose and trehalose showed the least amount of REGN421 aggregate formation as determined by SE-HPLC analysis. The addition of sodium chloride to the formulation was shown to decrease the formation of charge variants (CEX-HPLC) but resulted in an increased amount of aggregate formation (SE-HPLC). A subsequent study was conducted to examine the thermal stability of REGN421 following formulation with a combination of 10% sucrose and 150 mM sodium chloride (NaCl). The goal of this study was to reduce NaCl-induced aggregation by including sucrose in the formulation, thereby maintaining the protective effects of NaCl on charge variant formation within the formulation. The sucrose concentration was modulated from 20% (no NaCl) to 10% (with NaCl) to control the osmolality of the formulation. The thermal stability of REGN421 in this formulation as determined by CEX-HPLC analysis (formation of charge variants) was similar to results using NaCl as a single stabilizing agent (Table 3), demonstrating the protective effect of NaCl on charge variant formation. Furthermore, the rate of aggregate formation was significantly decreased compared to the formulation using NaCl as a single stabilizing agent, demonstrating the stabilizing effects of sucrose on the formulation. Based on these results, REGN421 was formulated with 10% sucrose and 150 mM NaCl.

[0094] For the antibody stability results shown in Table 3, 0.3 mL of 10 mM phosphate, pH 6.0, 0.2% Polysorbate 20, and 25 mg/mL REGN421, plus the indicated thermal stabilizer, in a 2 mL Type 1 borosilicate glass vial with a FluroTec® coated 4432/50 butyl rubber stopper were subjected to 45°C for 28 days. Turbidity was reported as the change in OD₄₀₅ relative compared to starting material. Starting material results represent the average results of the starting material for all ten formulations not incubated at 45°C. OD = Optical density; RP = Reverse phase high performance liquid chromatography; SE = Size exclusion high performance liquid chromatography; CEX = cation exchange high performance liquid chromatography.

Table 3: Effect of Thermal Stabilizers on the Stability

Thermal Stabilizer	Visual Appearance	Turbidity (OD 405 nm)	% Total REGN421 Recovered (RP-HPLC)	% Native REGN421 Recovered (SE-HPLC)	% REGN421 Aggregate Recovered (SE-HPLC)	% Native REGN421 Recovered (CEX-HPLC)
Starting Material * (no 45°C incubation)	Pass	0.00	100	97.7	1.0	58.4
No Thermal Stabilizer	Fail	0.00	100	94.0	2.0	41.0
150 mM NaCl	Pass	0.01	100	91.4	4.4	44.7
150 mM NaCl +10% Sucrose	Pass	0.00	100	94.0	2.1	45.4
20% Sucrose	Pass	0.00	98	94.5	1.2	40.8
20% Sorbitol	Pass	0.04	99	92.9	2.6	26.9
10% Mannitol	Pass	0.00	99	93.8	2.0	39.5
20% Trehalose	Pass	0.00	99	94.7	1.2	40.2
5% Glycerol	Pass	0.02	100	85.4	10.9	NP
3% Arginine	Fail	0.02	100	90.1	5.7	42.7
3% Glycine	Pass	0.01	106	91.1	5.0	31.9

EXAMPLE 6: STRESS STABILITY STUDIES ON REGN421 FORMULATION

[0095] Accelerated stress studies were performed by holding REGN421 at 25°C for 6 months. At this accelerated condition, a shift in the charge variant profile toward more acidic species was observed by cation exchange chromatography (CEX-HPLC). A 2% to 5% decrease in purity was observed by SDS PAGE after six months of storage, along with a 1.6% reduction in purity and a 1.2% increase in high molecular weight aggregate formation detected by SE-HPLC. There was no loss of potency of REGN421 as determined by binding assay after six months of storage at 25°C.

[0096] Additional accelerated and stress stability studies performed on REGN421 antibody indicate that the protein is physically stable when vortexed for 60 minutes or 120 minutes, frozen and thawed from -80°C to ambient temperature for eight cycles, incubated at 45°C for 28 days, incubated at 37°C for 28 days, or incubated at 25°C for 28 days (Table 4). During these studies, the solution remained visibly clear, no loss of protein was observed, and no change in pH occurred after these stresses. However, a decrease in the purity of REGN421 was detected by size exclusion HPLC and cation exchange HPLC when the protein was incubated at $\geq 25^\circ\text{C}$ for 28 days, indicating that changes in molecular weight and charge variants occur when REGN421 is exposed to stress conditions. A smaller level of chemical degradation was detected when the protein was incubated at 37°C compared to incubation at 45°C. REGN421 was physically and chemically stable over the 14 day assessment period at 25°C and after agitation and freeze/thaw cycles. No significant loss of potency, as determined using the DII4/Notch binding assay, was observed for any of the stressed samples.

Table 4: Stress Stability of 25 mg/mL Anti-DII4 Antibody

Stress Test	No Stress	Agitation		45°C Incubation		37°C Incubation		25°C Incubation		Freeze/Thaw
Length of Stress	0 min	60 min	120 min	14 days	28 days	14 days	28 days	14 days	28 days	8 cycles
Visual Appearance	Pass	Pass	Pass	Pass	Pass	Pass	Pass	Pass	Pass	Pass
Stress Test	No Stress	Agitation		45°C Incubation		37°C Incubation		25°C Incubation		Freeze/Thaw
Turbidity (OD 405 nm)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00
pH	5.9	5.9	5.9	6.0	6.0	6.0	6.0	6.0	6.0	6.0
% Total REGN421 Recovered (RP-HPLC)	100	99	96	96	100	98	100	97	101	99
% Native REGN421 Recovered (SE-HPLC)	98.0	97.9	98.0	95.8	94.0	96.9	96.0	97.9	97.7	98.0
% Native REGN421 Recovered (CEX-HPLC)	58.4	58.7	59.4	52.2	45.4	55.5	51.5	59.3	57.7	58.8
Binding Assay (% Relative Potency)	89	NP	94	NP	88	NP	101	NP	94	88

[0097] For the antibody stability results shown in Table 4, 0.3 mL of 10 mM Histidine, pH 6.0, 0.2% Polysorbate 20, 10% Sucrose, 150 mM NaCl, and 25 mg/mL HIH685P, in a 2 mL Type 1 borosilicate glass vial with a FluroTec® coated 4432/50 butyl rubber stopper was combined with the organic cosolvents and subjected to the various designated stresses. Turbidity was reported as the relative change in OD at 405 nm as compared to the starting material. The acceptance criteria for the binding assay was 50-150% of reference standard. OD = Optical density; RP = Reverse phase high performance liquid chromatography; SE = Size exclusion high performance liquid chromatography; CEX = cation exchange high performance liquid chromatography.

EXAMPLE 7: STORAGE STABILITY OF FORMULATED ANTI-DLL4 ANTIBODY

[0098] No significant loss of REGN421 was observed by UV spectrophotometry, or as determined by SDS-PAGE, after 24 months of storage at -80°C, -30°C, or -20°C. A 0.8% decrease in purity and a 0.4% increase in aggregate formation were detected by SE-HPLC after storage at 5°C for 24 months. There was no significant change in charge variants as detected by CEX HPLC after 24 months of storage at the tested conditions. There was no change in potency as determined by binding assay at the end of the storage period. REGN421 continued to meet the acceptance criteria for all the analyses after 24 months of storage at 5°C. See Table 5.

Table 5: Stability of 25 mg/mL REGN421 (10 mM phosphate, pH 6.0, 0.2% polysorbate 20, 10% sucrose, and 150 mM NaCl)

Storage Temperature/duration	-80°C/24 mo.	-30°C/24 mo.	-20°C/12 mo.	5°C/24 mo.
Appearance	Pass	Pass	Pass	Pass
pH	6.1	6.1	5.9	5.9
Turbidity (OD 405 nm) ¹	0.00	0.00	0.00	0.00
% Total REGN421 Recovered (RP-HPLC)	105	105	99	105
Purity by Size-Exclusion-HPLC	99.4	98.8	98.3	97.7
% main peak purity				
% HMW species	0.4	0.5	NA	NA

(continued)

Storage Temperature/duration	-80°C/24 mo.	-30°C/24 mo.	-20°C/12 mo.	5°C/24 mo.
Charged Variant Analysis by CE-HPLC	18	18	NA	NA
% region 1 (acidic)	57	57	59	58.3
2 % region (main)	25	25	NA	NA
% region 3 (basic)				
Binding Assay (% Ref. Std.)	100	99	95.3	NP

EXAMPLE 8: COMPATIBILITY WITH IV DELIVERY DEVICE

[0099] The 25 mg/mL REGN421 formulation was diluted in an intravenous (IV) bag composed of polyvinyl chloride (PVC) with Di-(2-ethylhexyl)-phthalate (DEHP) containing normal saline. Three different doses of REGN421 were examined in this study including a low dose (0.1 mg/kg and 40 kg patient), a middle dose (4 mg/kg and 40 kg patient), and a high dose (16 mg/kg and 120 kg patient). For the low dose, additional "placebo" was added to the bag to help stabilize the REGN421 against agitation.

[0100] The IV bags containing diluted REGN421 were first held for 24 hours at 5°C and then held for an additional 24 hours at 25°C. After these incubations were complete, the IV bags were connected to infusion sets, primed with the diluted antibody, and held for one hour at ambient temperature. Diluted antibody was pumped through the infusion sets at rates ranging from 25 mL/hr to 500 mL/hr. Three infusion pumps and three infusion sets representing the major suppliers (Alaris, Baxter, and Hospira) were utilized in this investigation. These infusion sets represent all of the basic materials (PVC with DEHP, PVC with TriOctyl-TriMellitate (TOTM), and polyolefins) that comprise infusion sets with inline filters. All infusion sets contained a 0.2 µm polyethersulfone filter.

[0101] REGN421 was physically stable when held for 24 hours at 5°C, held for 24 hours at 25°C (see Table 6), held for 1 hour at ambient temperature in the various infusion sets, and pumped through the infusion sets utilizing the various infusion pumps at a rate of either 25 mL/hr or 500 mL/hr. Precipitates were not detected by visual inspection or turbidity, the pH of the solution was stable, and there was no decrease in protein concentration as determined by RP-HPLC detected in any sample tested. The protein was also chemically stable. No increase in molecular weight species or charge variants were observed as determined by SE-HPLC and CEX-HPLC, respectively, in this compatibility study.

Table 6

Saline IV Bag (mL)	REGN421 (mg)	IV Infusion Set	Total REGN421 recovered (mg/mL)	% Native (SE-HPLC)	% Main Charge Form (CEX-HPLC)
100	8	Alaris PVC w/ DEHP	0.059	97.6	60.3
100	8	Baxter PVC w/ TOTM	0.060	97.3	61.8
100	8	Hospira Polyethylene lined PVC	0.061	97.4	61.3
100	160	Alaris Low Sorbing Polyethylene lined PVC	1.40	98.1	59.8
100	160	Baxter PVC w/ DEHP	1.51	97.5	60.3
100	160	Hospira PVC w/ TOTM	1.41	97.7	59.9
250	1920	Alaris Polypropylene	6.87	97.7	60.6
250	1920	Baxter PVC w/ TOTM	6.85	97.9	60.4
250	1920	Hospira PVC w/ DEHP	6.77	98.0	60.2

[0102] This data support the following conclusions: (a) REGN421 was stable following dilution of 4 mg of REGN421 in a 50 mL IV bag containing saline or dilution of 1920 mg of REGN421 in a 250 mL IV bag of saline. The saline containing IV bag of saline was composed of PVC containing DEHP. (b) Diluted REGN421 was stable following incubation in an IV bag for periods of up to 24 hours at 5°C or 24 hours at 25°C. (c) Diluted REGN421 can be administered using an infusion pump. (d) Diluted REGN421 can be administered with an infusion set composed of either PVC containing DEHP, PVC containing TOTM, or polyolefin. (e) The use of an inline 0.2 µm polyethersulfone filter is compatible with REGN421.

(f) Diluted REGN421 can be administered at a rate ranging from 25 mL/hr to 500 mL/hr.

EXAMPLE 9: MOLECULAR MASS DETERMINATION

[0103] Formulated REGN421 was characterized with respect to protein structure and activity. The predicted protein molecular weight of native REGN421 without glycosylation is approximately 146 kDa. Non-reducing SDS-PAGE was performed to confirm the molecular weight of the heterotetrameric molecule and to examine levels of high and low molecular weight protein forms that co-purified with the intact antibody. An alkylating reagent was added during sample preparation to prevent disulfide shuffling and to minimize antibody degradation due to heating. The observed molecular weight of the glycosylated antibody determined following electrophoresis and Coomassie staining was observed to be approximately 150 kDa. The mass of REGN421 was also examined using SDS-PAGE under reducing conditions. Samples were reduced and heated prior to electrophoresis, and protein bands were detected by staining with Coomassie blue. REGN421 was detected as two main bands corresponding to the antibody heavy chain (approximately 50 kDa) and light chain (approximately 25 kDa).

[0104] Capillary electrophoresis-SDS (CE-SDS) was also employed to assess the mass of REGN421. CE-SDS is able to resolve reduced IgG polypeptide chains by size and allows for quantification of the heterogeneity and variants that may exist in REGN421 antibody. The percentage of non-glycosylated heavy chain is calculated from the ratio of corrected peak area of non-glycosylated heavy chain to the sum of the corrected peak areas corresponding to non-glycosylated and glycosylated heavy chains. Two principal peaks were observed in the CE-SDS reducing electropherograms for REGN421 antibody. The two principal peaks for drug substance lots align with two principal peaks identified as the reduced light (LC) and glycosylated heavy (HC) chains in the IgG control standard. The molecular weights of peaks labeled HC and LC for REGN421 and the IgG control standard are approximately 60 kDa and 25 kDa, respectively. The molecular weights of the heavy and light chains determined by this method are slightly greater than expected for human IgG1, suggesting that this technique may slightly overestimate the molecular weights of the heavy and light chains. The relative abundance of the non-glycosylated heavy chain (NGHC) form of the tested lots of REGN421 represent approximately 1.5% of the total peak area detected.

EXAMPLE 10: FUCOSYLATION ANALYSIS

[0105] REGN421 drug substance lots were treated with PNGase F to release N-linked oligosaccharides from the antibody heavy chain. Free oligosaccharides were subsequently derivatized at the reducing end with the fluorescent reagent, anthranilic acid. Modified oligosaccharides were analyzed by reverse phase HPLC and MALDI-TOF mass spectrometry. Examination of the HPLC chromatogram revealed that the modified oligosaccharides were separated into two main groups, non-fucosylated bi-antennary species and fucosylated bi-antennary species. Within each group (fucosylated vs non-fucosylated), the oligosaccharides were further separated into digalactosyl (G2), monogalactosyl (G1) or agalactosyl (G0) forms. Oligosaccharide structure assignments were achieved through the mass analysis of each fractionated oligosaccharide using MALDI-TOF mass spectrometry. REGN421 lots that were analyzed contained between 82.4 and 87.0% ($84.6\% \pm 1.47\%$; $\text{av} \pm \text{SD}$) of fucosylated oligosaccharide chains (Table 7), based on the integration of corresponding peaks in the chromatograms. Mass analysis of pooled oligosaccharides demonstrated that fucosylated bi-antennary oligosaccharides are the dominant species, which agrees with the HPLC analysis. The observed difference in the amount of galactosylation within the N-linked sugar chains between various tested lots of REGN421 reflects the heterogeneity that can occur during antibody production.

Table 7: REGN421 Glycosylation Patterns

	Glycan Form	REGN421 Lots					Glycan Formulae
		Lot 1	Lot 2	Lot 3	Lot 4	Lot 5	
Nonfucosylated	G0 (%)	6.9	7.1	7.9	5.2	3.8	(GlcNAc) ₂ (Man) ₃ (GlcNAc) ₂
	G1 (%)	3.3	3.9	4.3	3.9	3.7	(GlcNAc) ₂ (Man) ₃ (GlcNAc) ₂ (Gal) ₁
	G2	0.9	0.8	1.0	1.8	2.1	(GlcNAc) ₂ (Man) ₃ (GlcNAc) ₂ (Gal) ₂

	Glycan Form	REGN421 Lots					Glycan Formulae
		Lot 1	Lot 2	Lot 3	Lot 4	Lot 5	
	(%)						
	Man-5 (%)	4.5	3.5	4.0	4.6	3.6	(GlcNAc) ₂ (Man) ₅
Fucosylated	G0F (%)	36.8	37.2	35.9	29.4	25.2	Fuc(GlcNAc) ₂ (Man) ₃ (GlcNAc) ₂
	G1F (%)	37.1	37.3	36.0	40.7	45.0	Fuc(GlcNAc) ₂ (Man) ₃ (GlcNAc) ₂ (Gal) ₁
	G2F (%)	10.4	10.3	10.9	14.3	16.7	Fuc(GlcNAc) ₂ (Man) ₃ (GlcNAc) ₂ (Gal) ₂
	% Fucosylation	84.3	84.8	82.8	84.4	86.9	

EXAMPLE 11: HEAVY CHAIN C-TERMINAL LYSINE ANALYSIS

[0106] Peptide mapping was used to confirm the C-terminal sequence of REGN421. REGN421 was denatured in 6.0 M urea, 100 mM Tris, pH 7.5, reduced with 5 mM DTT and alkylated with 12.5 mM iodoacetamide. The protein was then diluted 6-fold to bring the urea concentration down to 1.0 M, and subsequently digested with trypsin (1:20 enzyme to substrate ratio) at 37°C for three hours. Peptides were separated by HPLC and analyzed by mass spectroscopy. As standards for identification of the native C-terminal peptide from REGN421, three synthetic peptides corresponding to the three most likely putative C-terminal peptide sequences (SLSLSP, SLSLSPG, and SLSLSPGK; respectively SEQ ID NO: 10, SEQ ID NO: 11, and SEQ ID NO: 12) were generated and analyzed by HPLC using an identical solvent elution gradient compared to the gradient used to elute the REGN421 trypsin digested peptides to facilitate the identification of the actual C-terminal peptide in the tryptic map of REGN421. The expected trypsin derived C-terminal peptide of SLSLSPGK (SEQ ID NO:12) was not found in the tryptic maps of any of the tested REGN421 lots, indicating that the C-terminal Lys was removed. In the tryptic map, there is only one peptide (elution at 46.60 min) that co-eluted with one of the synthetic peptides (SLSLSPG; SEQ ID NO:11). This peptide corresponds to the C-terminal peptide of REGN421

heavy chain through mass analysis (expected mass 659.69, observed mass 660.2). The results suggest that post-translational processing leads to complete removal of the expected Lys from the heavy chain C-terminus.

SEQUENCE LISTING

[0107]

<110> WALSH, Scott
DIX, Daniel

<120> Stabilized Formulations Containing Anti-Dll4 Antibodies

<130> 7400P

<160> 12

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 123

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 1

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr
			20					25					30		
Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35				40						45			
Ser	Phe	Leu	Trp	Tyr	Asp	Gly	Thr	Asn	Lys	Asn	Tyr	Val	Glu	Ser	Val
	50				55						60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Met	Leu	Tyr
65					70				75					80	
Leu	Glu	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Arg	Asp	His	Asp	Phe	Arg	Ser	Gly	Tyr	Glu	Gly	Trp	Phe	Asp	Pro
			100					105					110		
Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					
		115					120								

<210> 2

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 2

Gly	Phe	Thr	Phe	Ser	Ser	Tyr	Gly
1				5			

<210> 3

EP 2 854 849 B1

<211> 8
<212> PRT
<213> Artificial Sequence

5 <220>
<223> Synthetic

<400> 3

10 Leu Trp Tyr Asp Gly Thr Asn Lys
1 5

<210> 4
<211> 16
15 <212> PRT
<213> Artificial Sequence

20 <220>
<223> Synthetic

<400> 4

25 Ala Arg Asp His Asp Phe Arg Ser Gly Tyr Glu Gly Trp Phe Asp Pro
1 5 10 15

<210> 5
<211> 107
<212> PRT
<213> Artificial Sequence

30 <220>
<223> Synthetic

<400> 5

35 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15
Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Tyr
20 25 30
40 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45
Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60
45 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80
Glu Asp Phe Ala Val Tyr Tyr Cys Gln His Arg Ser Asn Trp Pro Pro
85 90 95
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

50 <210> 6
<211> 6
<212> PRT
<213> Artificial Sequence

55 <220>
<223> Synthetic

EP 2 854 849 B1

<400> 6

Gln Ser Val Ser Ser Tyr
1 5

5

<210> 7
<211> 3
<212> PRT
<213> Artificial Sequence

10

<220>
<223> Synthetic

<400> 7

15

Asp Ala Ser
1

<210> 8
<211> 9
<212> PRT
<213> Artificial Sequence

20

<220>
<223> Synthetic

25

<400> 8

Gln His Arg Ser Asn Trp Pro Pro Thr
1 5

30

<210> 9
<211> 685
<212> PRT
<213> Homo sapiens

35

<400> 9

40

45

50

55

EP 2 854 849 B1

	Met	Ala	Ala	Ala	Ser	Arg	Ser	Ala	Ser	Gly	Trp	Ala	Leu	Leu	Leu	Leu
	1				5					10					15	
	Val	Ala	Leu	Trp	Gln	Gln	Arg	Ala	Ala	Gly	Ser	Gly	Val	Phe	Gln	Leu
5				20					25					30		
	Gln	Leu	Gln	Glu	Phe	Ile	Asn	Glu	Arg	Gly	Val	Leu	Ala	Ser	Gly	Arg
			35					40					45			
	Pro	Cys	Glu	Pro	Gly	Cys	Arg	Thr	Phe	Phe	Arg	Val	Cys	Leu	Lys	His
		50					55					60				
10	Phe	Gln	Ala	Val	Val	Ser	Pro	Gly	Pro	Cys	Thr	Phe	Gly	Thr	Val	Ser
	65					70					75					80
	Thr	Pro	Val	Leu	Gly	Thr	Asn	Ser	Phe	Ala	Val	Arg	Asp	Asp	Ser	Ser
					85					90					95	
	Gly	Gly	Gly	Arg	Asn	Pro	Leu	Gln	Leu	Pro	Phe	Asn	Phe	Thr	Trp	Pro
				100					105					110		
15	Gly	Thr	Phe	Ser	Leu	Ile	Ile	Glu	Ala	Trp	His	Ala	Pro	Gly	Asp	Asp
			115					120					125			
	Leu	Arg	Pro	Glu	Ala	Leu	Pro	Pro	Asp	Ala	Leu	Ile	Ser	Lys	Ile	Ala
		130					135					140				
20	Ile	Gln	Gly	Ser	Leu	Ala	Val	Gly	Gln	Asn	Trp	Leu	Leu	Asp	Glu	Gln
	145					150				155					160	
	Thr	Ser	Thr	Leu	Thr	Arg	Leu	Arg	Tyr	Ser	Tyr	Arg	Val	Ile	Cys	Ser
					165					170					175	
	Asp	Asn	Tyr	Tyr	Gly	Asp	Asn	Cys	Ser	Arg	Leu	Cys	Lys	Lys	Arg	Asn
			180					185					190			
25	Asp	His	Phe	Gly	His	Tyr	Val	Cys	Gln	Pro	Asp	Gly	Asn	Leu	Ser	Cys
			195					200					205			
	Leu	Pro	Gly	Trp	Thr	Gly	Glu	Tyr	Cys	Gln	Gln	Pro	Ile	Cys	Leu	Ser
		210					215					220				
	Gly	Cys	His	Glu	Gln	Asn	Gly	Tyr	Cys	Ser	Lys	Pro	Ala	Glu	Cys	Leu
	225					230					235				240	
30	Cys	Arg	Pro	Gly	Trp	Gln	Gly	Arg	Leu	Cys	Asn	Glu	Cys	Ile	Pro	His
					245					250					255	
	Asn	Gly	Cys	Arg	His	Gly	Thr	Cys	Ser	Thr	Pro	Trp	Gln	Cys	Thr	Cys
				260					265					270		
	Asp	Glu	Gly	Trp	Gly	Gly	Leu	Phe	Cys	Asp	Gln	Asp	Leu	Asn	Tyr	Cys
			275					280					285			
35	Thr	His	His	Ser	Pro	Cys	Lys	Asn	Gly	Ala	Thr	Cys	Ser	Asn	Ser	Gly
		290					295					300				
	Gln	Arg	Ser	Tyr	Thr	Cys	Thr	Cys	Arg	Pro	Gly	Tyr	Thr	Gly	Val	Asp
	305					310					315				320	
40	Cys	Glu	Leu	Glu	Leu	Ser	Glu	Cys	Asp	Ser	Asn	Pro	Cys	Arg	Asn	Gly
					325					330					335	
	Gly	Ser	Cys	Lys	Asp	Gln	Glu	Asp	Gly	Tyr	His	Cys	Leu	Cys	Pro	Pro
				340					345					350		
	Gly	Tyr	Tyr	Gly	Leu	His	Cys	Glu	His	Ser	Thr	Leu	Ser	Cys	Ala	Asp
			355					360					365			
45	Ser	Pro	Cys	Phe	Asn	Gly	Gly	Ser	Cys	Arg	Glu	Arg	Asn	Gln	Gly	Ala

EP 2 854 849 B1

		370				375				380						
	Asn	Tyr	Ala	Cys	Glu	Cys	Pro	Pro	Asn	Phe	Thr	Gly	Ser	Asn	Cys	Glu
	385					390					395					400
5	Lys	Lys	Val	Asp	Arg	Cys	Thr	Ser	Asn	Pro	Cys	Ala	Asn	Gly	Gly	Gln
					405					410					415	
	Cys	Leu	Asn	Arg	Gly	Pro	Ser	Arg	Met	Cys	Arg	Cys	Arg	Pro	Gly	Phe
				420					425					430		
	Thr	Gly	Thr	Tyr	Cys	Glu	Leu	His	Val	Ser	Asp	Cys	Ala	Arg	Asn	Pro
			435					440					445			
10	Cys	Ala	His	Gly	Gly	Thr	Cys	His	Asp	Leu	Glu	Asn	Gly	Leu	Met	Cys
		450					455					460				
	Thr	Cys	Pro	Ala	Gly	Phe	Ser	Gly	Arg	Arg	Cys	Glu	Val	Arg	Thr	Ser
	465					470					475					480
	Ile	Asp	Ala	Cys	Ala	Ser	Ser	Pro	Cys	Phe	Asn	Arg	Ala	Thr	Cys	Tyr
				485						490					495	
15	Thr	Asp	Leu	Ser	Thr	Asp	Thr	Phe	Val	Cys	Asn	Cys	Pro	Tyr	Gly	Phe
			500					505						510		
	Val	Gly	Ser	Arg	Cys	Glu	Phe	Pro	Val	Gly	Leu	Pro	Pro	Ser	Phe	Pro
			515					520					525			
20	Trp	Val	Ala	Val	Ser	Leu	Gly	Val	Gly	Leu	Ala	Val	Leu	Leu	Val	Leu
		530					535					540				
	Leu	Gly	Met	Val	Ala	Val	Ala	Val	Arg	Gln	Leu	Arg	Leu	Arg	Arg	Pro
	545					550				555						560
	Asp	Asp	Gly	Ser	Arg	Glu	Ala	Met	Asn	Asn	Leu	Ser	Asp	Phe	Gln	Lys
				565					570						575	
25	Asp	Asn	Leu	Ile	Pro	Ala	Ala	Gln	Leu	Lys	Asn	Thr	Asn	Gln	Lys	Lys
			580					585						590		
	Glu	Leu	Glu	Val	Asp	Cys	Gly	Leu	Asp	Lys	Ser	Asn	Cys	Gly	Lys	Gln
			595				600						605			
	Gln	Asn	His	Thr	Leu	Asp	Tyr	Asn	Leu	Ala	Pro	Gly	Pro	Leu	Gly	Arg
		610					615					620				
30	Gly	Thr	Met	Pro	Gly	Lys	Phe	Pro	His	Ser	Asp	Lys	Ser	Leu	Gly	Glu
	625					630					635					640
	Lys	Ala	Pro	Leu	Arg	Leu	His	Ser	Glu	Lys	Pro	Glu	Cys	Arg	Ile	Ser
				645					650						655	
	Ala	Ile	Cys	Ser	Pro	Arg	Asp	Ser	Met	Tyr	Gln	Ser	Val	Cys	Leu	Ile
35			660					665					670			
	Ser	Glu	Glu	Arg	Asn	Glu	Cys	Val	Ile	Ala	Thr	Glu	Val			
			675				680					685				

<210> 10
 <211> 6
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 10

Ser Leu Ser Leu Ser Pro
 1 5

<210> 11
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic

<400> 11

Ser Leu Ser Leu Ser Pro Gly
1 5

<210> 12

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic

<400> 12

Ser Leu Ser Leu Ser Pro Gly Lys
1 5

Claims

1. A pharmaceutical formulation comprising:

- (a) 25 mg/mL $\pm$ 3.75 mg/mL of a human IgG1 antibody that binds specifically to human delta-like ligand 4 (DIL4);
- (b) 10 mM $\pm$ 1.5 mM phosphate, pH 6 $\pm$ 0.5;
- (c) 0.2% $\pm$ 0.03% polysorbate 20;
- (d) 150 mM $\pm$ 22.5 mM sodium chloride; and
- (e) 10% $\pm$ 1.5% sucrose, wherein:

- (a) the antibody comprises a heavy chain variable domain (HCVD) of SEQ ID NO: 1 and a light chain variable domain (LCVD) of SEQ ID NO: 5;
- (b) the antibody has a molecular weight of about 150 kDa; and
- (c) about 82% to about 87% of the antibodies are fucosylated.

2. The pharmaceutical formulation of claim 1 consisting of (a) 25 mg/mL $\pm$ 3.75 mg/mL of the antibody, (b) 10 mM $\pm$ 3 mM phosphate, pH 6 $\pm$ 0.5, (c) 0.2% $\pm$ 0.03% polysorbate 20, (d) 150 mM $\pm$ 22.5 mM sodium chloride, and (e) 10% $\pm$ 1.5% sucrose, in water.

3. The pharmaceutical formulation of claim 1 or 2, wherein at least 94% of the antibody has native conformation after 28 days of storage at 45°C; and/or wherein at least 45% of the antibody is the main charge form of the antibody after 28 days of storage at 45°C; and/or wherein at least 97% of the antibody has native conformation after 28 days of storage at 25°C; and/or wherein at least 57% of the antibody is the main charge form of the antibody after 28 days of storage at 25°C; and/or wherein at least 97% of the antibody has native conformation after 28 days of storage at 37°C; and/or wherein at least 51% of the antibody is the main charge form of the antibody after 28 days of storage at 37°C; and/or wherein at least 98% of the antibody has native conformation after six months of storage at 5°C; and/or wherein at least 61% of the antibody is the main charge form of the antibody after six months of storage at 5°C; and/or wherein the percent relative potency of the antibody after six months of storage at 5°C is at least 100% of the potency of the antibody prior to storage; and/or wherein at least 99% of the antibody has native conformation after six months of storage at -80°C; and/or wherein at least 57% of the antibody is the main charge form of the antibody after six months of storage at -80°C; and/or wherein the percent relative potency of the antibody after six months of storage at -80°C is at least 100% of the potency of the antibody prior to storage; and/or wherein at least 99% of the antibody has native conformation after six months of storage at -30°C; and/or wherein at least 57% of the antibody is the main charge variant of the antibody after six months of storage at -30°C; and/or wherein the percent relative potency of the antibody after six months of storage at -30°C is at least 65% of the

potency of the antibody prior to storage; and/or wherein at least 98% of the antibody has native conformation after six months of storage at -20°C; and/or wherein at least 60% of the antibody is the main charge variant of the antibody after six months of storage at -20°C; and/or wherein the percent relative potency of the antibody after six months of storage at -20°C is at least 100% of the potency of the antibody prior to storage.

4. A pharmaceutical composition of any one of claims 1-3, wherein said composition is contained in a container, preferably
 - (a) wherein the container is a vial, more preferably wherein the vial is glass; or
 - (b) wherein the container is an IV drip bag, preferably wherein the bag is made of polyvinyl chloride or polyolefin.
5. A kit comprising a pharmaceutical composition of any one of claims 1-3, a container, and instructions, preferably wherein the container is a glass vial fitted with a fluorocarbon-coated rubber stopper.

Patentansprüche

1. Pharmazeutische Formulierung, umfassend:

- (a) 25 mg/ml $\pm$ 3,75 mg/ml eines menschlichen IgG1-Antikörpers, der an menschlichen Delta-ähnlichen Liganden 4 (DIL4) spezifisch bindet;
- (b) 10 mM $\pm$ 1,5 mM Phosphat, pH 6 $\pm$ 0,5,
- (c) 0,2 % $\pm$ 0,03 % Polysorbat 20;
- (d) 150 mM $\pm$ 22,5 mM Natriumchlorid; und
- (e) 10 % $\pm$ 1,5 % Sucrose, wobei:

- (a) der Antikörper eine variable Region der schweren Kette (heavy chain variable domain, HCVD) von SEQ ID NO: 1 und eine variable Region der leichten Kette (light chain variable domain, LCVD) von SEQ ID NO: 5 umfasst;
- (b) der Antikörper ein Molekulargewicht von ungefähr 150 kDa hat; und
- (c) ungefähr 82 % bis ungefähr 87 % der Antikörper fucosyliert sind.

2. Pharmazeutische Formulierung nach Anspruch 1, bestehend aus (a) 25 mg/ml $\pm$ 3,75 mg/ml des Antikörpers, (b) 10 mM $\pm$ 3 mM Phosphat, pH 6 $\pm$ 0,5, (c) 0,2 % $\pm$ 0,03 % Polysorbat 20, (d) 150 mM $\pm$ 22,5 mM Natriumchlorid und (e) 10 % $\pm$ 1,5 % Sucrose, in Wasser.
3. Pharmazeutische Formulierung nach Anspruch 1 oder 2, wobei mindestens 94 % des Antikörpers nach 28 Tagen Lagerung bei 45 °C native Konformation haben; und/oder wobei mindestens 45 % des Antikörpers nach 28 Tagen Lagerung bei 45 °C die Hauptladungsform des Antikörpers sind; und/oder wobei mindestens 97 % des Antikörpers nach 28 Tagen Lagerung bei 25 °C native Konformation haben; und/oder wobei mindestens 57 % des Antikörpers nach 28 Tagen Lagerung bei 25 °C die Hauptladungsform des Antikörpers sind; und/oder wobei mindestens 97 % des Antikörpers nach 28 Tagen Lagerung bei 37 °C native Konformation haben; und/oder wobei mindestens 51 % des Antikörpers nach 28 Tagen Lagerung bei 37 °C die Hauptladungsform des Antikörpers sind; und/oder wobei mindestens 98 % des Antikörpers nach sechs Monaten Lagerung bei 5 °C native Konformation haben; und/oder wobei mindestens 61 % des Antikörpers nach sechs Monaten Lagerung bei 5 °C die Hauptladungsform des Antikörpers sind; und/oder wobei die prozentrelative Wirksamkeit des Antikörpers nach sechs Monaten Lagerung bei 5 °C mindestens 100 % der Wirksamkeit des Antikörpers vor der Lagerung ist; und/oder wobei mindestens 99 % des Antikörpers nach sechs Monaten Lagerung bei -80 °C native Konformation haben; und/oder wobei mindestens 57 % des Antikörpers nach sechs Monaten Lagerung bei -80 °C die Hauptladungsform des Antikörpers sind; und/oder wobei die prozentrelative Wirksamkeit des Antikörpers nach sechs Monaten Lagerung bei -80 °C mindestens 100

% der Wirksamkeit des Antikörpers vor der Lagerung ist; und/oder
wobei mindestens 99 % des Antikörpers nach sechs Monaten Lagerung bei -30 °C native Konformation haben;
und/oder
wobei mindestens 57 % des Antikörpers nach sechs Monaten Lagerung bei -30 °C die Hauptladungsform des
Antikörpers sind; und/oder
wobei die prozentrelative Wirksamkeit des Antikörpers nach sechs Monaten Lagerung bei -30 °C mindestens 65 %
der Wirksamkeit des Antikörpers vor der Lagerung ist; und/oder
wobei mindestens 98 % des Antikörpers nach sechs Monaten Lagerung bei -20 °C native Konformation haben;
und/oder
wobei mindestens 60 % des Antikörpers nach sechs Monaten Lagerung bei -20 °C die Hauptladungsform des
Antikörpers sind; und/oder
wobei die prozentrelative Wirksamkeit des Antikörpers nach sechs Monaten Lagerung bei -20 °C mindestens 100
% der Wirksamkeit des Antikörpers vor der Lagerung ist.

4. Pharmazeutische Zusammensetzung nach einem der Ansprüche 1 bis 3, wobei die Zusammensetzung in einem
Behälter enthalten ist, bevorzugt

- (a) wobei der Behälter eine Ampulle ist, noch bevorzugt wobei die Ampulle Glas ist; oder
- (b) wobei der Behälter ein IV-Infusionsbeutel ist, bevorzugt wobei der Beutel aus Polyvinylchlorid oder Polyolefin
hergestellt ist.

5. Kit, umfassend eine pharmazeutische Zusammensetzung nach einem der Ansprüche 1 bis 3, einen Behälter und
Anweisungen, bevorzugt wobei der Behälter eine Glasampulle ist, die mit einem mit Fluorkohlenstoff beschichteten
Gummistopfen ausgestattet ist.

Revendications

1. Formulation pharmaceutique comprenant :

- (a) 25 mg/mL $\pm$ 3,75 mg/mL d'un anticorps IgG1 humain qui se lie spécifiquement au ligand de type delta 4
(DII4) humain ;
- (b) du phosphate 10 mM $\pm$ 1,5 mM, pH 6 $\pm$ 0,5 ;
- (c) du polysorbate 20 à 0,2 % $\pm$ 0,03 % ;
- (d) du chlorure de sodium 150 mM $\pm$ 22,5 mM ; et
- (e) du saccharose à 10 % $\pm$ 1,5 %,

- (a) l'anticorps comprenant un domaine variable de chaîne lourde (HCVD) de SEQ ID NO : 1 et un domaine
variable de chaîne légère (LCVD) de SEQ ID NO : 5 ;
- (b) l'anticorps ayant un poids moléculaire d'environ 150 kDa ; et
- (c) environ 82 % à environ 87 % des anticorps étant fucosylés.

2. Formulation pharmaceutique de la revendication 1, constituée de (a) 25 mg/mL $\pm$ 3,75 mg/mL de l'anticorps, (b)
du phosphate 10 mM $\pm$ 3 mM, pH 6 $\pm$ 0,5, (c) du polysorbate 20 à 0,2 % $\pm$ 0,03 %, (d) du chlorure de sodium 150
mM $\pm$ 22,5 mM, et (e) du saccharose à 10 % $\pm$ 1,5 %, dans de l'eau.

3. Formulation pharmaceutique de la revendication 1 ou 2, dans laquelle au moins 94 % de l'anticorps présente une
conformation native après 28 jours de stockage à 45 °C ; et/ou
dans laquelle au moins 45 % de l'anticorps constitue la forme de charge principale de l'anticorps après 28 jours de
stockage à 45 °C ; et/ou
dans laquelle au moins 97 % de l'anticorps présente une conformation native après 28 jours de stockage à 25 °C ;
et/ou
dans laquelle au moins 57 % de l'anticorps constitue la forme de charge principale de l'anticorps après 28 jours de
stockage à 25 °C ; et/ou
dans laquelle au moins 97 % de l'anticorps présente une conformation native après 28 jours de stockage à 37 °C ;
et/ou
dans laquelle au moins 51 % de l'anticorps constitue la forme de charge principale de l'anticorps après 28 jours de
stockage à 37 °C ; et/ou

dans laquelle au moins 98 % de l'anticorps présente une conformation native après six mois de stockage à 5 °C ; et/ou dans laquelle au moins 61 % de l'anticorps constitue la forme de charge principale de l'anticorps après six mois de stockage à 5 °C ; et/ou

dans laquelle le pourcentage de pouvoir relatif de l'anticorps après six mois de stockage à 5 °C est d'au moins 100 % du pouvoir de l'anticorps avant le stockage ; et/ou

dans laquelle au moins 99 % de l'anticorps présente une conformation native après six mois de stockage à -80 °C ; et/ou

dans laquelle au moins 57 % de l'anticorps constitue la forme de charge principale de l'anticorps après six mois de stockage à -80 °C ; et/ou

dans laquelle le pourcentage de pouvoir relatif de l'anticorps après six mois de stockage à -80 °C est d'au moins 100 % du pouvoir de l'anticorps avant le stockage ; et/ou

dans laquelle au moins 99 % de l'anticorps présente une conformation native après six mois de stockage à -30 °C ; et/ou

dans laquelle au moins 57 % de l'anticorps constitue la variante de charge principale de l'anticorps après six mois de stockage à -30 °C ; et/ou

dans laquelle le pourcentage de pouvoir relatif de l'anticorps après six mois de stockage à -30 °C est d'au moins 65 % du pouvoir de l'anticorps avant le stockage ; et/ou

dans laquelle au moins 98 % de l'anticorps présente une conformation native après six mois de stockage à -20 °C ; et/ou

dans laquelle au moins 60 % de l'anticorps constitue la variante de charge principale de l'anticorps après six mois de stockage à -20 °C ; et/ou

dans laquelle le pourcentage de pouvoir relatif de l'anticorps après six mois de stockage à -20 °C est d'au moins 100 % du pouvoir de l'anticorps avant le stockage.

4. Composition pharmaceutique de l'une quelconque des revendications 1 à 3, ladite composition étant contenue dans un contenant, de préférence

(a) le contenant étant une fiole, de manière davantage préférée la fiole étant en verre ; ou

(b) le contenant étant une poche de perfusion IV, de préférence la poche étant constituée de polychlorure de vinyle ou de polyoléfine.

5. Kit comprenant une composition pharmaceutique de l'une quelconque des revendications 1 à 3, un contenant et des instructions, le contenant étant de préférence une fiole en verre munie d'un bouchon en caoutchouc revêtu de fluorocarbène.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61653478 PCT [0001]
- US 6949245 B [0005]
- WO 9402602 A, Abgenix [0005]
- US 6596541 B [0005]
- US 7488806 B [0005] [0028] [0042] [0048]
- US 7534868 B [0005] [0028] [0042] [0048]
- US 7919593 B [0005] [0028] [0042] [0048]
- US 20110150905 A1 [0005]
- US 20110150905 A [0028] [0042] [0048]
- WO 7488806 A [0037]
- WO 7534868 A [0037]
- WO 7919593 A [0037]
- US 4997423 A [0075]
- US 5908686 A [0075]
- US 6286699 B [0075]
- US 6645635 B [0075]
- US 7226554 B [0075]
- US 6629949 B [0078]
- US 6659982 B [0078]

Non-patent literature cited in the description

- **SHUTTER et al.** *Genes Develop.*, 2000, vol. 14, 1313-1318 [0004]
- **SAINSIN ; HARRIS.** *Trends in Molecular Medicine*, September 2007, vol. 13 (9), 389-395 [0037]
- **BARBIE ; LEFRANC.** The Human Immunoglobulin Kappa Variable (IGKV) Genes and Joining (IGKJ) Segments. *Exp. Clin. Immunogenet.*, 1998, vol. 15, 171-183 [0043]
- **SCAVINER, D. et al.** Protein Displays of the Human Immunoglobulin Heavy, Kappa and Lambda Variable and Joining Regions. *Exp. Clin. Immunogenet.*, 1999, vol. 16, 234-240 [0043]
- **LEFRANC, M.-P. et al.** IMGT®, the international Immunogenetics information system®. *Nucl. Acids Res*, 2009, vol. 37, D1006-D1012 [0043]
- **ROBINSON, N.** Protein Deamidation. *PNAS*, 16 April 2002, vol. 99 (8), 5283-5288 [0070]
- **MEEHAN et al.** *J. Controlled Release*, 1996, vol. 46, 107-116 [0078]