

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3804716 T3**

(12) **Oversættelse af
europæisk patentskrift**

- (51) Int.Cl.: **A 61 K 9/14 (2006.01)** **A 61 K 9/20 (2006.01)** **A 61 K 9/48 (2006.01)**
A 61 K 31/17 (2006.01) **A 61 K 31/381 (2006.01)** **A 61 K 31/415 (2006.01)**
A 61 K 31/426 (2006.01) **A 61 K 31/427 (2006.01)** **A 61 K 31/437 (2006.01)**
A 61 K 31/444 (2006.01) **A 61 K 31/497 (2006.01)** **A 61 K 31/506 (2006.01)**
A 61 K 31/70 (2006.01) **A 61 K 45/06 (2006.01)** **A 61 P 1/16 (2006.01)**
A 61 P 3/00 (2006.01) **A 61 P 3/04 (2006.01)** **A 61 P 5/50 (2006.01)**
A 61 P 9/12 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2025-03-03**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-12-25**
- (86) Europæisk ansøgning nr.: **19811691.5**
- (86) Europæisk indleveringsdag: **2019-05-28**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-04-14**
- (86) International ansøgning nr.: **CN2019088863**
- (87) Internationalt publikationsnr.: **WO2019228364**
- (30) Prioritet: **2018-05-31 CN 201810556685**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Hua Medicine (Shanghai) Ltd., 275 Ai Di Sheng Road, Pudong, Shanghai 201203, Kina**
- (72) Opfinder: **CHEN, Li, Building C-40, No. 2 Xinchu Road, Xinqiao Town, Songjiang, Shanghai 201612, Kina**
LI, Yongguo, Rm.203, No.3, Lane 300, Nan Dan Dong Road, Xuhui, Shanghai 200030, Kina
WANG, Gaosen, Rm. 603, No.10, Lane 240, An Shun Road, Changning, Shanghai 200051, Kina
GAO, Huisheng, Rm.301, No.143, Lane 229, Guanyue Road, Zhoupu, Town, Pudong, Shanghai 201318, Kina
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **FARMACEUTISK KOMBINATION, SAMMENSÆTNING OG KOMBINATIONSPRÆPARAT OMFATTENDE GLUCOKINASE-AKTIVATOR OG SGLT-2-INHIBITOR OG FREMGANGSMÅDER TIL FREMSTILLING OG ANVENDELSER DERAFF**
- (56) Fremdragne publikationer:
EP-A1- 1 734 040
WO-A1-2015/176640
CN-A- 106 474 480
CN-A- 107 854 435
US-A1- 2012 142 636
US-A1- 2012 294 936
ZHU XIAO-XUE ET AL: "Dorzagliatin (HMS5552), a novel dual-acting glucokinase activator, improves glycaemic control and pancreatic [beta]-cell function in patients with type 2 diabetes: A 28-day treatment study using

Fortsættes ...

biomarker-guided patient selection", vol. 20, no. 9, 30 May 2018 (2018-05-30), GB, pages 2113 - 2120, XP055880617, ISSN: 1462-8902, Retrieved from the Internet

<URL:https://api.wiley.com/onlinelibrary/tdm/v1/articles/10.1111%2Fdom.13338> DOI: 10.1111/dom.13338

MINI ET AL: "Recent Updates on Glucokinase Activators for the Treatment of Type 2 Diabetes Mellitus", REVIEWS IN MEDICINAL CHEMISTRY, 1 January 2014 (2014-01-01), pages 585 - 602, XP055880851, Retrieved from the Internet <URL:http://www.eurekaselect.com/article/61496> [retrieved on 20220119]

SCHEEN ANDRÉ J.: "Investigational insulin secretagogues for type 2 diabetes", vol. 25, no. 4, 27 February 2016 (2016-02-27), UK, pages 405 - 422, XP055883340, ISSN: 1354-3784, Retrieved from the Internet

<URL:https://www.tandfonline.com/doi/pdf/10.1517/13543784.2016.1152260?needAccess=true> DOI: 10.1517/13543784.2016.1152260

BLONDE LAWRENCE ET AL: "Fixed-Dose Combination Therapy in Type 2 Diabetes Mellitus", ENDOCRINE PRACTICE, vol. 20, no. 12, 1 December 2014 (2014-12-01), US, pages 1322 - 1332, XP055912325, ISSN: 1530-891X, Retrieved from the Internet <URL:http://dx.doi.org/10.4158/EP14259.RA> DOI: 10.4158/EP14259.RA

CHIKARA GAURAV ET AL: "A Narrative Review of Potential Future Antidiabetic Drugs: Should We Expect More?", INDIAN JOURNAL OF CLINICAL BIOCHEMISTRY, SPRINGER PRIVATE LTD, IN, vol. 33, no. 2, 8 June 2017 (2017-06-08), pages 121 - 131, XP036476265, ISSN: 0970-1915, [retrieved on 20170608], DOI:

10.1007/S12291-017-0668-Z

WANG YE; WENDONG LIU; FANG WANG: "Research Progress in application of therapeutic drugs for Type2 diabetes mellitus", PROGRESS IN PHARMACEUTICAL SCIENCES, vol. 41, no. 6, 25 June 2017 (2017-06-25), pages 434 - 443, XP055755746, ISSN: 1001-5094

DESCRIPTION

Description

PRIORITY APPLICATION

[0001] This application claims the priority of the Chinese patent application No. 201810556685.6 filed on May 31, 2018.

TECHNICAL FIELD

[0002] The present disclosure relates to a pharmaceutical combination, a composition and a fixed dose combination (FDC) formulation comprising a glucokinase activator (GKA) drug and a partner drug, to a preparation method thereof and to a use thereof in the treatment of some diseases.

[0003] More specifically, the present disclosure relates to a pharmaceutical combination, a pharmaceutical composition, or an oral solid formulation of a fixed dose combination comprising a glucokinase activator drug and a partner drug, and a preparation method thereof. The present disclosure also relates to the use of a pharmaceutical combination, a pharmaceutical composition or a fixed dose combination formulation comprising a glucokinase activator for the treatment and/or prevention of one or more diseases and medical disorders, including but not limited to type I diabetes, type II diabetes, diabetic nephropathy, decreased renal function, impaired glucose tolerance, impaired fasting blood glucose, obesity and hypertension. In addition, the present disclosure also relates to a method for treating and/or preventing one or more diseases and medical disorders, comprising administering a therapeutically effective amount of the pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation disclosed herein to a subject in need thereof.

BACKGROUND

[0004] Diabetes mellitus has become a prevalent disease worldwide, with 425 million patients over the world, and 120 million patients in China (International Diabetes Federation, Diabetes Atlas, 2015). Type II diabetes, i.e., non-insulin dependent diabetes mellitus (NIDDM), accounts for more than 90% of the patients with diabetes. Type II diabetes is a hyperglycemic, chronic, metabolic dysfunction resulting from an imbalance of blood glucose homeostasis in human body caused by insulin secretion disorder and insulin resistance. The blood glucose balance of the human body is mainly coordinated by two hormones that control blood glucose, including insulin and glucagon.

[0005] Glucose sensor glucokinase (GK) senses blood glucose changes, regulates the secretion of

the messenger glucose-controlling hormones, insulin and glucagon, and GLP-1 (glucagon-like peptide-1), and constitutes a sensing system for steady-state regulation of human blood glucose. Glucose-controlling hormones control glucose storage during glucose uptake and glucose supply during fasting, constituting the steady-state regulation of human blood glucose. Organs involved in glucose storage are mainly liver, muscle and fat. Under the action of blood glucose and insulin, glucose is taken up and converted into liver glycogen, muscle glycogen and triglycerides. The main organ involved in glucose supply is liver. Under the action of blood glucose and glucagon, liver supplies glucose to a human body through liver glucose synthesis and liver glucose output. Insulin can also effectively regulate the activity of sodium-glucose cotransporter SGLT-2. When blood glucose rises, the glucose excreted by kidneys is reabsorbed for body's glucose storage. Glucose uptake and liver glucose output, as well as the use of glucose by various organs, constitute an operating system for a steady-state balance of the human blood glucose. The coordinated operation of the sensing system and the operating system of glucose constitute a random regulation of a steady-state of the human blood glucose.

[0006] In diabetic patients, the impaired function and expression of glucokinase, and the dysfunction of the sensor, result in the dysfunction of the early phase secretion of glucose-controlling hormones, affecting glucose uptake and output, and resulting in post-prandial hyperglycemia and pre-prandial hypoglycemia. Abnormal signaling of glucose-controlling hormones cause abnormal functions and expressions of key proteins in the execution system of glucose uptake and output, forming abnormal operating state, leading to type II diabetes.

[0007] Existing oral hypoglycemic drugs for diabetes usually act on a single glucose-controlling organ and cannot effectively treat the problem of imbalanced blood glucose homeostasis. Glucokinase activators represent a class of new drugs developed to treat or improve the disease state of patients with type II diabetes. For example, ((S)-2-[4-((2-chloro-phenoxy)-2-oxo-2,5-dihydro-pyrrol-1-yl)-4-methyl-pentanoic acid [1-((R)-2,3-dihydroxy-propyl)-1H-pyrazol-3-yl]-amide (referred to as HMS5552 hereinafter) can effectively improve the function of the glucose sensor in diabetic patients, and is currently the most promising drug for the treatment of diabetes to solve the above clinical needs. US 2012/0142636 A1 discloses the use of a phosphonate and phosphinate activator in combination with a further anti-diabetic agent. Zhu Xiao-Xue et al., Diabetes, Obesity and Metabolism, 2018, 20, 2113-2120, investigate the pharmacokinetics and pharmacodynamics of a dual-acting glucokinase activator, dorzagliatin (HMS5552), and its safety, tolerability and effect on pancreatic β -cell function in Chinese patients with type 2 diabetes. EP 1 734 040 A1 teaches a class of anti-diabetic quinazoline compounds which may be used in combination with other drugs that are used in the treatment or prevention of type 2 diabetes mellitus.

SUMMARY


[0008] Diabetic patients often encounter such a situation during treatment that SGLT-2 inhibitors alone are not effective, and blood glucose cannot be controlled to an ideal level, especially after a period of use. In this regard, the inventors found that the combination of a SGLT-2 inhibitor and a glucokinase activator can significantly improve the hypoglycemic effect of the SGLT-2 inhibitor and reduce the safety risk, thus the pharmaceutical combination, composition and combination formulation comprising a glucokinase activator and a SGLT-2 inhibitor disclosed herein were

obtained.

[0009] More specifically, the combination of a SGLT-2 inhibitor and a glucokinase activator can improve the function of multiple organs of patients in middle and late stages, and diabetes and accompanying diseases and complications were well treated. The number of pills taken by patients was reduced and the compliance of patients was improved, the total dose of drugs that achieve the same therapeutic effect was reduced, and the maximum efficacy was achieved with the lowest dose. It has a good effect and practical significance for the treatment or prevention of one or more of type I diabetes, type II diabetes, hyperglycemia, impaired glucose tolerance, obesity and other symptoms.

[0010] On the other hand, the fixed dose combination formulation comprising a glucokinase activator and a partner drug (the second or more active pharmaceutical ingredients) disclosed herein not only has better therapeutic effects than the single use of each of these two or more drugs, but also solves the technical challenges that usually exist in combination formulations. The fixed dose combination formulation disclosed herein can provide the simultaneous release of two or more active ingredients with uniform content, and can optimize the dissolution rate of active ingredients contained in the formulation, especially making the active ingredients contained in the formulation quickly released in the pH environment of a small intestine. This is beneficial to the timely or simultaneous arrival of drugs in gut, pancreatic islet and liver target organs, achieving clinical advantages of multi-organs targeting, and synergistic hypoglycemic effect, and exhibiting a better therapeutic effect and reduced toxic or side effects. In addition, the fixed dose combination formulation comprising a glucokinase activator and a partner drug (the second or more active pharmaceutical ingredients) disclosed herein also has a short disintegration time, and good dissolution characteristics, and/or gives high bioavailability of the glucokinase activator in patients.

[0011] The present disclosure provides a pharmaceutical combination, a pharmaceutical composition, and a fixed dose combination formulation comprising a glucokinase activator, such as HMS5552 with the following structure, or an isotope labeled compound thereof, or a pharmaceutically acceptable salt thereof, and other oral hypoglycemic drug which is a SGLT-2 inhibitor, especially a solid formulation, such as an oral solid formulation such as a tablet,



[0012] Specifically, the present disclosure also provides a pharmaceutical combination, a pharmaceutical composition or a fixed dose combination formulation comprising a glucokinase activator drug such as HMS5552 or a pharmaceutically acceptable salt thereof and a SGLT-2 inhibitor. Further, examples of the SGLT-2 inhibitor include, but are not limited to: canagliflozin, dapagliflozin, empagliflozin, ipragliflozin, luseogliflozin and tofogliflozin, and a pharmaceutically acceptable salt thereof. Preferably, the SGLT-2 inhibitor is selected from the group consisting of empagliflozin, dapagliflozin and canagliflozin.

[0013] More specifically, the present disclosure also provides a fixed dose combination solid formulation comprising a glucokinase activator drug such as HMS5552 or a pharmaceutically acceptable salt thereof and a partner drug which is a SGLT-2 inhibitor such as empagliflozin. The solid formulation is preferably a tablet, or more preferably a coated tablet. In one embodiment, the glucokinase activator, such as HMS5552, is present in the form of a solid dispersion.

[0014] More specifically, the present disclosure also provides a fixed dose combination solid formulation comprising a glucokinase activator drug such as HMS5552 or a pharmaceutically acceptable salt thereof and a partner drug which is a SGLT-2 inhibitor such as dapagliflozin. The solid formulation is preferably a tablet, or more preferably a coated tablet. In one embodiment, the glucokinase activator, such as HMS5552, is present in the form of a solid dispersion.

[0015] More specifically, the present disclosure also provides a fixed dose combination solid formulation comprising a glucokinase activator drug such as HMS5552 or a pharmaceutically acceptable salt thereof and a partner drug which is a SGLT-2 inhibitor such as canagliflozin. The solid formulation is preferably a tablet, or more preferably a coated tablet. In one embodiment, the glucokinase activator, such as HMS5552, is present in the form of a solid dispersion.

[0016] The present disclosure also provides a pharmaceutical combination, a pharmaceutical composition or a fixed dose combination formulation of a glucokinase activator drug and a partner drug which is a SGLT-2 inhibitor (the second or more active pharmaceutical ingredients) prepared by a dry or wet processing method. The release mode of the pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation disclosed herein is rapid release of the two or more active pharmaceutical ingredients.

[0017] The present disclosure also provides a pharmaceutical formulation comprising a glucokinase activator drug and a partner drug which is a SGLT-2 inhibitor (the second or more active pharmaceutical ingredients), which has a short disintegration time and good dissolution characteristics and/or gives high bioavailability of the glucokinase activator in patients. The present disclosure also provides a method of preparing a pharmaceutical composition or a pharmaceutical formulation of a fixed dose combination of a glucokinase activator drug and a partner drug (the second or more active pharmaceutical ingredients, such as empagliflozin, dapagliflozin, and canagliflozin) by a dry or wet processing method. The dry processing method includes dry compression (tableting) and dry granulation; and the wet processing method includes wet granulation.

[0018] The present disclosure also provides a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation comprising a glucokinase activator drug and a partner drug which is a SGLT-2 inhibitor (the second or more active pharmaceutical ingredients), and a method for preventing, slowing down, delaying or treating a metabolic disorder (especially type II diabetes).

[0019] The present disclosure also provides a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation comprising a glucokinase activator drug and a partner drug which is a SGLT-2 inhibitor (the second or more active pharmaceutical ingredients), and a method for improving blood glucose control in patients in need thereof, especially patients with type

II diabetes.

[0020] The present disclosure also provides a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation comprising a glucokinase activator drug and a partner drug, which is a SGLT-2 inhibitor, and a method for improving blood glucose control in patients with insufficient blood glucose control.

[0021] The present disclosure also provides a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation comprising a glucokinase activator drug and a partner drug, which is a SGLT-2 inhibitor, and a method for preventing, slowing down or delaying diabetic nephropathy, decreased renal function, impaired glucose tolerance (IGT), impaired fasting blood glucose (IFG), hypertension, insulin resistance and/or progression of metabolic syndrome to type II diabetes.


[0022] The present disclosure also provides a pharmaceutical combination, a pharmaceutical composition and a pharmaceutical formulation comprising a glucokinase activator drug and a partner drug, which is a SGLT-2 inhibitor, and a method for preventing, slowing down, delaying or treating a disease or a disorder including diabetes complications.

[0023] Other objects of the present disclosure will be apparent to those skilled in the art from the description and examples.

BRIEF DESCRIPTION OF THE EMBODIMENTS

[0024] The first aspect of the present disclosure provides a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation comprising the following components, a preparation method thereof, and a use thereof for treating diabetes and related diseases:

1. (a) a glucokinase activator, which is a compound selected from the following formula, or a pharmaceutically acceptable salt, an isotope labeled compound, a crystalline form, a hydrate, or a solvate thereof; preferably, the glucokinase activator is HMS5552; more preferably, HMS5552 is present in the form of a solid dispersion,



2. (b) a SGLT-2 inhibitor; and
3. (c) one or more excipients.

[0025] Another aspect of the present disclosure provides a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation comprising the following components, a preparation method thereof, and a use thereof for the treatment of diabetes and related diseases:

1. (a) a glucokinase activator, which is HMS5552 compound, or a pharmaceutically acceptable salt, an isotope labeled compound, a crystalline form, a hydrate, or a solvate thereof; preferably, the glucokinase activator is HMS5552; more preferably, HMS5552 is present in the form of a solid dispersion;
2. (b) empagliflozin, dapagliflozin, canagliflozin, or a pharmaceutically acceptable salt, an isotope labeled compound, a crystalline form, a hydrate, a solvate, a diastereomeric or enantiomeric form thereof; and
3. (c) one or more excipients.

[0026] In particular, one aspect of the present disclosure also relates to a pharmaceutical combination, a pharmaceutical composition, and a pharmaceutical formulation comprising a fixed dose combination of a HMS5552 solid dispersion and a partner drug which is a SGLT-2 inhibitor (e.g., empagliflozin, dapagliflozin, canagliflozin), and a preparation method thereof and a use thereof.

DEFINITION

[0027] Unless otherwise specified, all technical and scientific terms used herein have the same meanings as commonly understood by those skilled in the art to which the present disclosure belongs, but in case of conflict, the definitions in this specification shall prevail.

[0028] As used in the specification and claims, the singular forms "a", "an" and "the (said)" include plural forms, unless the context clearly dictates otherwise.

[0029] Unless otherwise specified, the percentages (%) in the specification are all weight percentages (% by weight).

[0030] All numerical values or expressions related to component amounts, process conditions, etc. used in the specification and claims should be understood to be modified by "about" in all cases. The term "about" when referring to a quantity or a range of numerical values means that the quantity or the range of numerical values referred to is an approximate value within experimental variability (or within statistical experimental error). Therefore, the quantity or the range of numerical values can be varied between, for example, ± 5 percent of the quantity or the range of numerical values.

[0031] All ranges involving the same components or properties include endpoints, which can be independently combined. Since these ranges are continuous, they include every value between the minimal and maximal values. It should also be understood that any numerical range cited in this application is intended to include all sub-ranges within that range.

[0032] When the present disclosure uses a range to define physical properties such as molecular weight or chemical properties, it shall include all combinations and sub-combinations of the range and specific embodiments therein. The term "comprising" (and related terms such as "containing"

or "including" or "having") includes embodiments which are, for example, any combinations of substances, compositions, methods, or processes, etc., and are "consisted of the described features" or "essentially consisted of the described features".

[0033] As used in the specification and claims, "and/or" should be understood as "one or both" of the associated components, that is, the components coexist in some cases and exist separately in other cases. Multiple components listed with "and/or" should be understood in the same way, that is, "one or more" of the associated components. In addition to the components specifically identified in the "and/or" clause, other components may optionally be present, whether associated or not associated with those specifically identified components. Therefore, as a non-limiting example, when "A and/or B" is used to connect an open ending word such as "comprising", in one embodiment, it may only refer to A (optionally comprising components other than B); in another embodiment, it can only refer to B (optionally comprising components other than A); in yet another embodiment, it refers to A and B (optionally comprising other components), etc.

[0034] It should be understood that, unless explicitly indicated to the contrary, in any method that includes more than one step or one act claimed herein, the order of the steps and acts of the method is not necessarily limited to the order of the steps and acts of the method described.

[0035] The abbreviations used in the present disclosure have the common meanings in the fields of chemistry, biology and formulation.

[0036] Unless otherwise specified, the term "SGLT-2 inhibitor" or any substance thereof (e.g., "empagliflozin", "dapagliflozin", "canagliflozin") in the context of the present disclosure is also intended to include any pharmaceutically acceptable salt, crystalline form, hydrate, solvate, diastereomer or enantiomer thereof.

[0037] HMS552 (its former name is RO5305552, and English name is Dorzagliatin) has a chemical name of (S)-2-[4-(2-chloro-phenoxy)-2-oxo-2,5-dihydro-pyrrol-1-yl]-4-methyl-pentanoic acid [1-((R)-2,3-dihydroxy-propyl)-1H-pyrazol-3-yl]-amide.

[0038] Unless otherwise specified, % by weight (wt %) represents a percentage of the total weight of a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation.

[0039] A solid dispersion (SD) refers to a solid dispersion system formed by highly dispersing one or more active pharmaceutical ingredients into inactive adjuvants or carriers.

[0040] EUDRAGIT is the trade name of a synthetic pharmaceutical adjuvant. It includes copolymers of methacrylic acid and copolymers of methacrylate ester, collectively referred to as polyacrylic resins. Adjuvants based on polyacrylic resin are divided into different models according to their composition, proportion and degree of polymerization. Among them, Eudragit E is a polymer of dimethylaminoethyl methacrylate and methacrylate; Eudragit L is a polymer of methacrylic acid and methyl methacrylate with free carboxyl:ester of 1:1; and Eudragit S is a polymer of methacrylic acid and methyl methacrylate with free carboxyl:ester of 1:2.

[0041] The term "tablet" as used herein is intended to include compressed pharmaceutical

formulations of all shapes and sizes, whether coated or not.

[0042] The terms "effective amount" or "therapeutically effective amount" refer to an amount of the agent sufficient to provide the desired biological result. The biological result may be a reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an "effective amount" for therapeutic use is the necessary amount of the composition comprising a compound as the active ingredient as disclosed herein for providing a clinically significant decrease in a disease. An appropriate "effective" amount in any individual embodiment may be determined by one of ordinary skill in the art using routine experimentation. Thus, the expression "effective amount" generally refers to the quantity for which the active substance has therapeutic effects.

[0043] As used herein, the terms "treat" or "treatment" are synonymous with the term "prevent" and are meant to indicate a postponement of development of diseases, preventing the development of diseases, and/or reducing severity of such symptoms that will or are expected to develop. Thus, these terms include ameliorating existing disease symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disorder or disease, e.g., preventing the development of the disorder or disease, relieving the disorder or disease, causing a regression of the disorder or disease, relieving a condition caused by the disease or disorder, or stopping the symptoms of the disease or disorder.

[0044] By "pharmaceutically acceptable" or "pharmacologically acceptable", it is meant a material which is not biologically or otherwise undesirable, i.e., the material may be administered to an individual without causing a minimum of undesirable biological effects or interacting in a deleterious manner with any other components of the composition in which it is contained.

[0045] As used herein, the term "subject" encompasses mammals and non-mammals. Examples of mammals include, but are not limited to, any member of the mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fish and the like. In one embodiment of the present disclosure, the mammal is a human.

[0046] A compound as an active ingredient in a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation (e.g., a fixed dose combination formulation) comprising a glucokinase activator disclosed herein may form a salt. Reference to a compound disclosed herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as used herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compound may be formed, for example, by reacting the compound with an amount, such as an equivalent amount, of acid or base in a medium such as a medium from which the salt precipitates

or in an aqueous medium (lyophilization after reaction).

[0047] Various compounds and salts, solvates, esters and prodrugs thereof, and polymorphs thereof are intended to be included in the disclosure.

[0048] It is to be understood that the terminology employed herein is for the purpose of describing particular embodiments, and is not intended to be limiting. Further, although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods, devices and materials are described below.

BRIEF DESCRIPTION OF THE DRAWINGS

[0049]

Figure 1 shows the effect on blood glucose after glucose challenge in normal mice ($X \pm s$, $n = 10$) with the administration of empagliflozin alone and the combination of HMS5552 and empagliflozin; and

Figure 2 shows the effect on $AUC_{0-120 \text{ min}}$ after glucose challenge in normal mice ($X \pm s$, $n = 10$; ***, $P < 0.001$) with the administration of empagliflozin alone and the single administration of the combination of HMS5552 and empagliflozin.

DETAILED DESCRIPTION OF THE EMBODIMENTS

[0050] One aspect of the present disclosure relates to a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation such as a fixed dose combination formulation of a glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and a partner drug which is a SGLT-2 inhibitor (e.g., empagliflozin, dapagliflozin, canagliflozin). The formulation can be powder, granule, tablet, capsule, sachet or other solid forms. Specifically, one aspect of the present disclosure relates to a tablet comprising a fixed dose combination of a glucokinase activator and a partner drug which is a SGLT-2 inhibitor (e.g., empagliflozin, dapagliflozin, canagliflozin).

[0051] In a specific aspect of the present disclosure, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation comprises:

1. (1) a glucokinase activator or a pharmaceutically acceptable salt thereof, or an isotope labeled analogue, a crystalline form, a hydrate, or a solvate thereof; preferably, the glucokinase activator is HMS5552; more preferably, HMS5552 is present in the form of a solid dispersion, such as a solid dispersion (e.g., a spray-dried powder) comprising a polymer carrier;
2. (2) a SGLT-2 inhibitor; preferably selected from: empagliflozin, dapagliflozin, canagliflozin, or a pharmaceutically acceptable salt, an isotope labeled compound, a crystalline form, a

- hydrate, or a solvate thereof; and/or
3. (3) filler(s); and/or
 4. (4) binder(s); and/or
 5. (5) disintegrant(s); and/or
 6. (6) lubricant(s) or glidant(s); and/or
 7. (7) coating agent(s).


[0052] In one embodiment disclosed herein, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation may also contain one or more excipients selected from the group consisting of one or more binders; one or more diluents (fillers); one or more disintegrants; one or more lubricants; one or more glidants; one or more surfactants or wetting agents; one or more antioxidants; and one or more coating agents.


Pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation

Glucokinase activator + SGLT-2 inhibitor

[0053] In one embodiment, the present disclosure relates to a pharmaceutical combination, a pharmaceutical composition or a pharmaceutical formulation (preferably, a fixed dose combination formulation), which comprises:

1. (a) a glucokinase activator, which is a compound represented by the following formulae, or a pharmaceutically acceptable salt, an isotope labeled compound, a crystalline form, a hydrate, or a solvate thereof





2. (b) a SGLT-2 inhibitor;

preferably, examples of the SGLT-2 inhibitor include but are not limited to: canagliflozin, dapagliflozin, empagliflozin, ipragliflozin, luseogliflozin and tofogliflozin, or a pharmaceutically acceptable salt thereof;

more preferably, the SGLT-2 inhibitor is selected from the group consisting of empagliflozin, dapagliflozin and canagliflozin; and

3. (c) one or more excipients;

wherein the above-mentioned drugs (a) and (b) are used simultaneously, separately or sequentially.


[0054] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the weight ratio of the glucokinase activator to the SGLT-2 inhibitor is about 30:1 to 1:30, preferably about 20:1 to 1:12, or more preferably about 0.75:1, about 1:2, about 1:1, about 1:4, about 1:6, about 1:12, about 2:1, about 2.5:1, about 3:1, about 5:1, about 6:1, about 7.5:1, about 10:1, about 15:1, or about 20:1.

[0055] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the glucokinase activator is present in a dose (preferably, a unit dose) ranging from 1 mg to about 200 mg, or preferably from about 25 mg to about 100 mg, preferably, wherein the dose (preferably, a unit dose) of the glucokinase activator is about 25 mg, about 50 mg, about 75 mg or about 100 mg.

[0056] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation, the SGLT-2 inhibitor is present in a dose (preferably, a unit dose) ranging from about 1 mg to 500 mg, or preferably from about 5 mg to about 300 mg, preferably, wherein the dose (preferably, a unit dose) of the SGLT-2 inhibitor is about 2.5 mg, about 5 mg, about 10 mg, about 12.5 mg, about 20 mg, about 25 mg, about 100 mg, about 200 mg or about 300 mg, or still preferably about 5 mg, about 10 mg, about 12.5 mg, about 25 mg, about 100 mg or about 300 mg; preferably, the SGLT-2 inhibitor is empagliflozin, with a dose (preferably, a unit dose) of about 0.5 mg to about 50 mg, preferably about 1 mg to about 25 mg; or preferably about 5 mg, about 10 mg, about 12.5 mg or about 25 mg; preferably, the SGLT-2 inhibitor is dapagliflozin, with a dose (preferably, a unit dose) of about 1 mg to about 50 mg, preferably about 2.5 mg, about 5 mg, about 10 mg or about 25 mg; or preferably about 2.5 mg, about 5 mg or about 10 mg; preferably, the SGLT-2 inhibitor is canagliflozin, with a dose (preferably, a unit dose) of about 50 mg to about 500 mg, and the preferable dose of canagliflozin is about 100 mg or about 300 mg.

[0057] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the above-mentioned glucokinase activator is the compound HMS5552, or an isotope labeled

compound, or a pharmaceutically acceptable salt thereof



[0058] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the glucokinase activator is present in the form of a solid dispersion.

[0059] In one embodiment, the solid dispersion is obtained by spray drying, hot melting or freeze drying of a glucokinase activator, or an isotope labeled compound, or a pharmaceutically acceptable salt thereof, together with a polymer carrier.

[0060] In one embodiment, the amount of the glucokinase activator in the solid dispersion, based on the total weight of the solid dispersion, may vary from about 1% to about 99% by weight, or preferably from 10% to 90% by weight. In one embodiment, the amount of the glucokinase activator is about 1% by weight, about 2% by weight, about 3% by weight, about 4% by weight, about 5% by weight, about 6% by weight, about 7% by weight, about 8% by weight, about 9% by weight, about 10% by weight, about 11% by weight, about 12% by weight, about 13% by weight, about 14% by weight, about 15% by weight, about 16% by weight, about 17% by weight, about 18% by weight, about 19% by weight, about 20% by weight, about 21% by weight, about 22% by weight, about 23% by weight, about 24% by weight, about 25% by weight, about 26% by weight, about 27% by weight, about 28% by weight, about 29% by weight, about 30% by weight, about 31% by weight, about 32% by weight, about 33% by weight, about 34% by weight, about 35% by weight, about 36% by weight, about 37% by weight, about 38% by weight, about 39% by weight, about 40% by weight, about 41% by weight, about 42% by weight, about 43% by weight, about 44% by weight, about 45% by weight, about 46% by weight, about 47% by weight, about 48% by weight, about 49% by weight, about 50% by weight, about 51% by weight, about 52% by weight, about 53% by weight, about 54% by weight, about 55% by weight, about 56% by weight, about 57% by weight, about 58% by weight, about 59% by weight, about 60% by weight, about 61% by weight, about 62% by weight, about 63% by weight, about 64% by weight, about 65% by weight, about 66% by weight, about 67% by weight, about 68% by weight, about 69% by weight, about 70% by weight, about 71% by weight, about 72% by weight, about 73% by weight, about 74% by weight, about 75% by weight, about 76% by weight, about 77% by weight, about 78% by weight, about 79% by weight, about 80% by weight, about 81% by weight, about 82% by weight, about 83% by weight, about 84% by weight, about 85% by weight, about 86% by weight, about 87% by weight, about 88% by weight, about 89% by weight, about 90% by weight, about 91% by weight, about 92% by weight, about 93% by weight, about 94% by weight, about 95% by weight, about 96% by weight, about 97% by weight, about 98% by weight, or about 99% by weight, or any range therebetween.

[0061] In one embodiment, the amount of the glucokinase activator in the solid dispersion, based on the total weight of the solid dispersion, is about 1% to about 20% by weight, about 2% to about 40% by weight, about 30% to about 60% by weight, about 60% to about 80% by weight, about 70%

to about 90% by weight, or about 80% to about 100% by weight.

[0062] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the glucokinase activator is present in the form of a solid dispersion, in which the weight ratio of the solid dispersion of the glucokinase activator to the SGLT-2 inhibitor is about 60:1 to 1:15, preferably about 40:1 to 1:6, or more preferably about 1.5:1, about 1:1, about 2:1, about 1:2, about 1:3, about 1:6, about 4:1, about 5:1, about 6:1, about 10:1, about 12:1, about 15:1, about 20:1, about 30:1, or about 40:1 or any range therebetween.

[0063] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the glucokinase activator is the compound HMS5552, an isotope labeled compound thereof or a pharmaceutically acceptable salt thereof, which is combined with a polymer carrier to obtain a solid dispersion by spray drying, hot melting or freeze drying, etc.

[0064] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the polymer carrier in the solid dispersion is selected from a polypropylene resin-based polymer, which is a polymeric compound derived from the polymerization of acrylic acid (or methacrylic acid and esters thereof such as methyl ester, ethyl esters and the like) (as monomer), or derived from the polymerization of two monomers (binary polymerization) or three monomers (ternary polymerization) in a certain ratio using acrylic acid and methacrylic acid (or its ester such as methyl ester, ethyl ester, dimethylaminoethyl ester, etc.).

[0065] In one embodiment, the polymer carrier used in the solid dispersion in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) is selected from the group consisting of copolymer of butyl methacrylate, dimethylaminoethyl methacrylate and methyl methacrylate; copolymer of methacrylic acid and ethyl acrylate; copolymer of methacrylic acid and methyl methacrylate; copolymer of ethyl acrylate, methyl methacrylate and chlorotrimethylamino ethyl methacrylate; copolymer of ethyl acrylate and methyl methacrylate; copolymer of methacrylic acid, methyl acrylate and methyl methacrylate, and copolymer of methacrylic acid and butyl acrylate.

[0066] In one embodiment, the above-mentioned polymer carrier is selected from the group consisting of copolymer of butyl methacrylate, dimethylaminoethyl methacrylate and methyl methacrylate (1:2:1), copolymer of methacrylic acid and ethyl acrylate (1:1), copolymer of methacrylic acid and methyl methacrylate (1:2), copolymer of ethyl acrylate, methyl methacrylate and chlorotrimethylamino ethyl methacrylate (1:2:0.2), copolymer of ethyl acrylate, methyl methacrylate and chlorotrimethylamino ethyl methacrylate (1:2:0.1), copolymer of ethyl acrylate and methyl methacrylate (2:1), copolymer of methacrylic acid and butyl acrylate (35:65), copolymer of methacrylic acid and methyl methacrylate (1:1), copolymer of methacrylic acid and methyl methacrylate (35:65).

[0067] In one embodiment, the above-mentioned polymer carrier is Eudragit, including Eudragit E, Eudragit L, Eudragit S, Eudragit RL and Eudragit RS, wherein Eudragit E is produced by the

polymerization of dimethylamino methacrylate and other neutral methacrylates, including copolymers of dimethylaminoethyl methacrylate and methacrylate; Eudragit L and Eudragit S is produced by the polymerization of methacrylic acid and methacrylates in various ratios, including a copolymer of methacrylic acid and methyl methacrylate with 1:1 of free carboxyl:ester or a copolymer of methacrylic acid and methyl methacrylate with 1:2 of free carboxyl:ester; Eudragit RL and Eudragit RS type is a copolymer of acrylic acid containing some quaternary amine groups and methacrylate, including the copolymer of acrylic acid containing 10% quaternary amine group and methacrylate and the copolymer of acrylic acid containing 5% quaternary amine group and methacrylate.

[0068] In one embodiment, the above-mentioned polymer carrier is selected from the group consisting of:

Eudragit E100, which is copolymer of butyl methacrylate, dimethylaminoethyl methacrylate and methyl methacrylate (1:2:1), including Eudragit E PO;

Eudragit L100, methacrylic acid copolymer of type A, which is an anionic copolymer of methacrylic acid and methyl methacrylate (1:1); and

Eudragit S 100, which is copolymer of methacrylic acid and methyl methacrylate (1:2).

[0069] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the polymer carrier in the solid dispersion of HMS5552 is methacrylic acid copolymer of type A (an anionic copolymer of methacrylic acid and methyl methacrylate (1:1)), preferably Eudragit, or more preferably Eudragit L100.

[0070] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the weight ratio of HMS5552 to Eudragit L100 in the solid dispersion of HMS5552 is about 1:10 to 10:1, about 1:9 to 9:1, about 2:3 to 9:1, about 3:4 to 9:1, about 4:5 to 9:1, about 5:6 to 9:1, or about 1:1 to 9:1; about 2:3 to 4:1, about 3:4 to 4:1, about 4:5 to 4:1, about 5:6 to 4:1, or about 1:1 to 4:1; about 2:3 to 7:3, about 3:4 to 7:3, about 4:5 to 7:3, about 5:6 to 7:3, or about 1:1 to 7:3; about 2:3 to 3:2, about 3:4 to 4:3, about 4:5 to 5:4, or about 5:6 to 6:5; about 1:4 to 4:1, about 3:7 to 7:3, about 2:3 to 3:2, about 3:4 to 4:3, about 4:5 to 5:4, or about 5:6 to 6:5, or any range therebetween.

[0071] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the weight ratio of HMS5552 to Eudragit L100 in the solid dispersion of HMS5552 is about 1:1, about 2:3, about 3:2, about 1:4, about 4:1, about 3:4, about 4:3, about 4:5, about 5:4, about 5:6, about 6:5, about 7:3, about 3:7, about 1:9, about 9:1, or any range therebetween.

[0072] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the second active ingredient is empagliflozin. The above-mentioned pharmaceutical composition or fixed dose combination formulation of the glucokinase activator (HMS5552 or an isotope labeled

compound or a pharmaceutically acceptable salt thereof) and empagliflozin contains (by weight): about 1-48% of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 1-35% of empagliflozin; about 0-90% of filler(s); about 1-10% of binder(s); about 1-10% of disintegrant(s); about 0.1-5% of lubricant(s); about 0-3% of glidant(s); and about 0-5% of coating agent(s). The pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) is prepared by a wet granulation method or a dry granulation method, preferably by a wet granulation method.

[0073] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and empagliflozin, the dose (preferably, a unit dosage) of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) is about 1 mg to 200 mg. Preferably, the dose (preferably, a unit dose) of the glucokinase activator is about 5 mg to 100 mg. Preferably, the dose (preferably, a unit dose) of the glucokinase activator is about 5 mg, about 10 mg, about 20 mg, about 25 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 75 mg, about 80 mg, about 90 mg, about 100 mg, or any range therebetween. More preferably, the dose (preferably, a unit dosage) of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound analogue or a pharmaceutically acceptable salt thereof) is about 25 mg, about 50 mg, about 75 mg, or about 100 mg. Preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0074] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and empagliflozin, the dose of empagliflozin (preferably, a unit dose) is about 0.5 mg to about 50 mg, or preferably about 1 mg to about 25 mg. Preferably, the dose (preferably, a unit dose) of empagliflozin is about 5 mg, about 10 mg, about 12.5 mg, or about 25 mg. Preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0075] In the pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation of the present disclosure, the specific embodiments of the doses (preferably, unit doses) of HMS5552 (or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and empagliflozin are as follows:

1. (1) about 75 mg of HMS5552 and about 5 mg of empagliflozin;
2. (2) about 75 mg of HMS5552 and about 10 mg of empagliflozin;
3. (3) about 75 mg of HMS5552 and about 12.5 mg of empagliflozin;
4. (4) about 75 mg of HMS5552 and about 25 mg of empagliflozin;
5. (5) about 50 mg of HMS5552 and about 10 mg of empagliflozin;
6. (6) about 50 mg of HMS5552 and about 25 mg of empagliflozin;
7. (7) about 25 mg of HMS5552 and about 10 mg of empagliflozin;

8. (8) about 25 mg of HMS5552 and about 25 mg of empagliflozin;
9. (9) about 100 mg of HMS5552 and about 5 mg of empagliflozin;
10. (10) about 100 mg of HMS5552 and about 10 mg of empagliflozin;
11. (11) about 100 mg of HMS5552 and about 12.5 mg of empagliflozin; and
12. (12) about 100 mg of HMS5552 and about 25 mg of empagliflozin;

preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0076] In one embodiment, the preferable dosage form of the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the present disclosure is a tablet.

[0077] In one embodiment, the above-mentioned tablet is a fixed dose combination tablet of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and empagliflozin.

[0078] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg empagliflozin) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the glucokinase activator is present in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0079] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg empagliflozin) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0080] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/12.5 mg empagliflozin) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled

compound or a pharmaceutically acceptable salt thereof); about 12.5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0081] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/25 mg empagliflozin) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 25 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0082] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg empagliflozin) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0083] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg empagliflozin) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0084] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/12.5 mg empagliflozin) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 12.5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0085] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/25 mg empagliflozin) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 25 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0086] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg empagliflozin) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0087] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg empagliflozin) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5%

of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0088] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/12.5 mg empagliflozin) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 12.5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0089] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/25 mg empagliflozin) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 25 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0090] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg empagliflozin) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0091] In one embodiment, the pharmaceutical combination, pharmaceutical composition or

pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg empagliflozin) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0092] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/12.5 mg empagliflozin) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 12.5 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0093] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/25 mg empagliflozin) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 25 mg of empagliflozin; about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0094] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the second active ingredient is dapagliflozin (or dapagliflozin propylene glycol monohydrate). The above-mentioned pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and dapagliflozin (or dapagliflozin propylene glycol monohydrate) contains (by weight): about 1-98% of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 0.1-30% dapagliflozin or dapagliflozin propylene glycol monohydrate; about 0-85% of filler(s); about 1-25% of

binder(s); about 1-15% of disintegrant(s); about 0.1-10% of lubricant(s); about 0-3% of glidant(s); and about 0-5% of coating agent(s). The pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) is prepared by a wet granulation method or a dry granulation method, preferably by a wet granulation method.

[0095] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and dapagliflozin or dapagliflozin propylene glycol monohydrate, the dose (preferably, a unit dose) of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) is about 1 mg to 200 mg. The preferable dose (preferably, a unit dose) of the glucokinase activator is about 5 mg to 100 mg. Preferably, the dose (preferably, a unit dose) of the glucokinase activator is about 5 mg, about 10 mg, about 20 mg, about 25 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 75 mg, about 80 mg, about 90 mg, about 100 mg, or any range therebetween. More preferably, the dose (preferably, a unit dose) of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) is about 25 mg, about 50 mg, about 75 mg, or about 100 mg. Preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0096] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and dapagliflozin or dapagliflozin propylene glycol monohydrate, based on the amount of dapagliflozin, the dose (preferably, a unit dose) of dapagliflozin is about 1 mg to about 50 mg, preferably about 2.5 mg, about 5 mg, about 10 mg or about 25 mg, or preferably, the dose (preferably, a unit dosage) is about 2.5 mg, about 5 mg or about 10 mg. Preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0097] In the pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation of the present disclosure, the specific embodiments of the doses (preferably, unit doses) of HMS5552 (or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and dapagliflozin are as follows:

1. (1) about 25 mg of HMS5552 and about 5 mg of dapagliflozin (or about 6.15 mg of dapagliflozin propylene glycol monohydrate);
2. (2) about 50 mg of HMS5552 and about 5 mg of dapagliflozin (or about 6.15 mg of dapagliflozin propylene glycol monohydrate);
3. (3) about 75 mg of HMS5552 and about 5 mg of dapagliflozin (or about 6.15 mg of dapagliflozin propylene glycol monohydrate);
4. (4) about 100 mg of HMS5552 and about 5 mg of dapagliflozin (or about 6.15 mg of dapagliflozin propylene glycol monohydrate);
5. (5) about 25 mg of HMS5552 and about 10 mg of dapagliflozin (or about 12.3 mg of dapagliflozin propylene glycol monohydrate);

6. (6) about 50 mg of HMS5552 and about 10 mg of dapagliflozin (or about 12.3 mg of dapagliflozin propylene glycol monohydrate);
7. (7) about 75 mg of HMS5552 and about 10 mg of dapagliflozin (or about 12.3 mg of dapagliflozin propylene glycol monohydrate);
8. (8) about 100 mg of HMS5552 and about 10 mg of dapagliflozin (or about 12.3 mg of dapagliflozin propylene glycol monohydrate);

preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0098] In one embodiment, the preferable dosage form of the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the present disclosure is a tablet.

[0099] In one embodiment, the above-mentioned tablet is a fixed dose combination tablet of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and dapagliflozin (or dapagliflozin propylene glycol monohydrate).

[0100] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0101] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0102] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L 100.

[0103] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/5 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 5 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0104] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L 100.

[0105] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg dapagliflozin (or an amount of dapagliflozin propylene glycol

monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0106] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L 100.

[0107] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/10 mg dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin)) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10 mg of dapagliflozin (or an amount of dapagliflozin propylene glycol monohydrate that can obtain said amount of dapagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0108] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), the second active ingredient is canagliflozin (or canagliflozin hemihydrate). The above-mentioned pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and canagliflozin (or canagliflozin hemihydrate) contains (by weight): about 1-80% of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 10-80% of dapagliflozin or canagliflozin

hemihydrate; about 0-85% of filler(s); about 1-25% of binder(s); about 1-15% of disintegrant(s); about 0.1-10% of lubricant(s); about 0-3% of glidant(s); and about 0-5% of coating agent(s). The pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) is prepared by a wet granulation method or a dry granulation method, preferably by a wet granulation method.

[0109] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and canagliflozin (or canagliflozin hemihydrate), the dose (preferably, a unit dose) of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) is about 1 mg to 200 mg. The preferable dose (preferably, a unit dose) of the glucokinase activator is about 5 mg to 100 mg. Preferably, the dose (preferably, a unit dose) of the glucokinase activator is about 5 mg, about 10 mg, about 20 mg, about 25 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 75 mg, about 80 mg, about 90 mg, about 100 mg, or any range therebetween. More preferably, the dose (preferably, a unit dosage) of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) is about 25 mg, about 50 mg, about 75 mg, or about 100 mg. Preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0110] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and canagliflozin (or canagliflozin hemihydrate), based on the amount of canagliflozin, the dose (preferably, a unit dose) of canagliflozin is about 50 mg to about 500 mg, or preferably, the dose (preferably, a unit dose) is about 100 mg or about 300 mg. Preferably, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), HMS5552 is present in the form of a solid dispersion.

[0111] In the pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation of the present disclosure, the specific embodiments of the doses (preferably, unit doses) of HMS5552 (or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and canagliflozin (or canagliflozin hemihydrate) are as follows:

1. (1) about 25 mg of HMS5552 and about 100 mg of canagliflozin (or about 101.93 mg of canagliflozin hemihydrate);
2. (2) about 50 mg of HMS5552 and about 100 mg of canagliflozin (or about 101.93 mg of canagliflozin hemihydrate);
3. (3) about 75 mg of HMS5552 and about 100 mg of canagliflozin (or about 101.93 mg of canagliflozin hemihydrate);
4. (4) about 100 mg of HMS5552 and about 100 mg of canagliflozin (or about 101.93 mg of canagliflozin hemihydrate);
5. (5) about 25 mg of HMS5552 and about 300 mg of canagliflozin (or about 305.78 mg of canagliflozin hemihydrate);

6. (6) about 50 mg of HMS5552 and about 300 mg of canagliflozin (or about 305.78 mg of canagliflozin hemihydrate);
7. (7) about 75 mg of HMS5552 and about 300 mg of canagliflozin (or about 305.78 mg of canagliflozin hemihydrate);
8. (8) about 100 mg of HMS5552 and about 300 mg of canagliflozin (or about 305.78 mg canagliflozin hemihydrate).

[0112] In one embodiment, the preferable dosage form of the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) of the present disclosure is a tablet.

[0113] In one embodiment, the above-mentioned tablet is a fixed dose combination tablet of the glucokinase activator (HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof) and canagliflozin (or canagliflozin hemihydrate).

[0114] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/100 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled analogue or a pharmaceutically acceptable salt thereof); about 100 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0 to 70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0115] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/100 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 100 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0116] In one embodiment, the pharmaceutical combination, pharmaceutical composition or

pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/100 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 100 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0 to 70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0117] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/100 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 100 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0 to 70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0118] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 25 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/300 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 25 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 300 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0119] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 50 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/300 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by

weight): about 50 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 300 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0120] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 75 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/300 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 75 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 300 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0121] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination tablet, which is a tablet of 100 mg glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof)/300 mg canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin)) contains the components with the following amounts (by weight): about 100 mg of the glucokinase activator (preferably, HMS5552 or an isotope labeled compound or a pharmaceutically acceptable salt thereof); about 300 mg of canagliflozin (or an amount of canagliflozin hemihydrate that can obtain said amount of canagliflozin); about 0-70% of optional filler(s); about 2-8% of binder(s); about 1-5% of disintegrant(s); about 0.5-3% of lubricant(s); about 0-0.5% of glidant(s) and about 0-5% of coating agent(s); preferably, the above-mentioned glucokinase activator is in the form of a solid dispersion as described above, or preferably, the solid dispersion contains a glucokinase activator and a polymer carrier, or preferably contains about 1:1 of the glucokinase activator and Eudragit L100.

[0122] In one embodiment, the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) further comprises other excipients, wherein the excipients include but are not limited to one or a mixture of diluents, flavoring agents (flavors), sweetening agents, and coloring agents.

[0123] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) disclosed herein contains optionally one or more fillers (diluents). Examples of fillers include, but are not limited to, cellulose derivatives such as microcrystalline cellulose or lignocellulose (including microcrystalline

cellulose and silicified microcrystalline cellulose), lactose, anhydrous lactose or lactose monohydrate, sucrose, starch, pregelatinized starch, dextrose, mannitol (including mannitol Pearlitol SD 200), fructose, xylitol, sorbitol, corn starch, modified corn starch, inorganic salts such as calcium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, dextrin/glucose binder, maltodextrin, compressible sugar and other known compatibilizers or fillers/or a mixture of two or more of them.

[0124] Examples of preferable fillers (diluent) include microcrystalline cellulose (MCC), silicified microcrystalline cellulose (SMCC), lactose, mannitol, sorbitol, calcium dihydrogen phosphate (dihydrate), corn starch, pregelatinized starch and powdered cellulose. Other preferable fillers (diluent) are microcrystalline cellulose and silicified microcrystalline cellulose. Microcrystalline cellulose can be obtained from several suppliers, including Avicel PH 101, Avicel PH 102, Avicel PH 103, Avicel PH 105, and Avicel PH 200 manufactured by FMC Corporation.

[0125] In one embodiment, the pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) disclosed herein contains optionally one or more binders. Examples include, but are not limited to, carboxymethylcellulose (including sodium carboxymethylcellulose), hydroxypropyl cellulose (including hydroxypropyl cellulose EXF), corn starch, pregelatinized starch, modified corn starch, polyvinyl pyrrolidone (PVP), hydroxypropylmethyl cellulose (HPMC) (including hydroxypropylmethyl cellulose 2208), lactose, sucrose, gum arabic, ethylcellulose, cellulose acetate and wax binders such as carnauba wax, paraffin wax, cetyl wax, polyethylene or microcrystalline wax and other conventional binder and/or a mixture of two or more of them. Further, in addition to the above binders, binders suitable for use in the present disclosure include, but are not limited to, alginate, alginate acid, microcrystalline cellulose, dextrin, gelatin, amylopectin, liquid glucose, guar gum, methylcellulose, polyethylene oxide, povidone and syrup, and the combination of them.

[0126] Preferable embodiments of the binder include hydroxypropyl cellulose (HPC), hydroxypropyl methyl cellulose (HMPC), polyvinylpyrrolidone (Povidone), hydroxyethyl cellulose, starch 1500 and Polyvidon. Other preferable binders are hydroxypropyl cellulose, hydroxypropyl methyl cellulose and polyvinylpyrrolidone.

[0127] In one embodiment, the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) contains optionally one or more disintegrants. Examples of disintegrants suitable for use in the present disclosure include, but are not limited to, croscarmellose sodium, crospovidone, lactose, sucrose, starch, potato starch, pregelatinized starch, corn starch, sodium carboxymethyl starch, sodium starch glycolate, microcrystalline cellulose, light silicic acid anhydride, low-substituted hydroxypropyl cellulose and other known disintegrants.

[0128] In one embodiment, the disintegrant is selected from one or more of modified starch, modified cellulose polymer or polycarboxylic acid, specifically selected from the group consisting of croscarmellose sodium, crospovidone, sodium starch glycolate, polyacrylamide and CMC Calcium. In one embodiment, the disintegrant is crospovidone. In another embodiment, the disintegrant is sodium starch glycolate. In another embodiment, the disintegrant is croscarmellose sodium. Croscarmellose sodium NF type A is available in the market under the trade name "Ac-di-

sol".

[0129] In one embodiment, the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) contains one or more lubricants. Examples of lubricants suitable for use in the present disclosure include, but are not limited to, magnesium stearate, zinc stearate, calcium stearate, talc, carnauba wax, stearic acid, palmitic acid, sodium stearyl fumarate, sodium lauryl sulphate, glyceryl palmitate stearate, palmitic acid, myristic acid and hydrogenated vegetable oils (including hydrogenated castor oil) and fats and other known lubricants and/or a mixture of two or more of them.

[0130] In one embodiment, embodiments of the lubricant include magnesium stearate, calcium stearate, stearic acid, sodium stearyl fumarate, hydrogenated castor oil, and a mixture thereof. Another preferable lubricant is magnesium stearate, or sodium stearyl fumarate, or a mixture thereof.

[0131] In one embodiment, the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) contains one or more glidants and/or anti-adherents. Examples of glidants and/or anti-adherents suitable for use in the present disclosure include, but are not limited to, silicon dioxide, colloidal silicon dioxide, magnesium silicate, calcium phosphate, magnesium trisilicate, talc and other forms of silicon dioxide such as aggregated silicate and hydrated silica gel.

[0132] In one embodiment, embodiments of the glidant include colloidal silicon dioxide, calcium phosphate, magnesium silicate, and talc, or a mixture thereof. The preferable glidant is colloidal silicon dioxide.

[0133] In one embodiment, the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) may also optionally contain one or more surfactants or wetting agents. The surfactant can be an anionic, cationic or neutral surfactant. The anionic surfactant includes sodium lauryl sulfate, sodium lauryl sulfonate, sodium oleyl sulfate, and sodium laurate mixed with stearate and talc. The cationic surfactant includes benzalkonium chloride and alkyl trimethyl ammonium bromide. The neutral surfactant includes glycerol monooleate, polyoxyethylene sorbitan fatty acid ester, polyvinyl alcohol and sorbitan ester. Embodiments of the wetting agent include poloxamer, polyoxyethylene alkyl ether, polyoxyethylene castor oil derivative, and polyoxyethylene stearate.

[0134] In one embodiment, the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation) may also optionally contain an antioxidant to render chemical stability. Examples of the antioxidant suitable for use in the present disclosure include, but are not limited to, tocopherol, ascorbic acid, gallic acid ester, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, thioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite and sodium sulfite and a combination thereof.

[0135] In one embodiment, the antioxidant is selected from the group consisting of α -tocopherol, γ -tocopherol, δ -tocopherol, extracts from natural sources enriched in tocopherol, L-ascorbic acid and

sodium or calcium salt thereof, ascorbyl palmitate, propyl gallate, octyl gallate, lauryl gallate, butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA). In one embodiment, the antioxidant is BHT or BHA.

[0136] In one embodiment, the preferable formulation of the above-mentioned fixed dose combination formulation is a tablet prepared by a compression method. The tablet may be coated, and preferable examples of coating substrates include sugar coating substrates, water-soluble film coating substrates, enteric film coating substrates, and the like.

[0137] Sucrose is used as the sugar coating substrate. In addition, one or more selected from talc powder, precipitated calcium carbonate, gelatin, gum arabic, amylopectin, carnauba wax, and the like can also be used in combination.

[0138] Examples of the water-soluble film coating substrate include cellulosic polymers such as hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hydroxyethyl cellulose, methyl hydroxyethyl cellulose, and the like; synthetic polymers such as polyvinyl acetal diethylaminoacetate, aminoalkyl methacrylate copolymer E [Eudragit E (trade name)], polyvinylpyrrolidone, and the like.

[0139] Examples of the enteric film coating substrate include cellulosic polymers such as hydroxypropyl methyl cellulose phthalate, hydroxypropyl methyl cellulose acetate succinate, carboxymethyl ethyl cellulose, cellulose acetate phthalate, and the like; acrylic polymer, such as methacrylic acid copolymer L [Eudragit L (trade name)], methacrylic acid copolymer LD [Eudragit L-30D55 (trade name)], methacrylic acid copolymer S [Eudragit S (trade name)], and the like.

[0140] Preferable examples of coating additives include: plasticizers such as polyvinyl alcohol (PVA), polyethylene glycol (PEG), propylene glycol, triethyl citrate, castor oil, polysorbate, and the like, or a mixture of two or more of them; opacifiers such as titanium dioxide and the like; coloring agents, dyes and lakes such as iron oxide red (ferric oxide), iron oxide yellow, and the like; and glidants such as talc and the like.

[0141] In one embodiment, the tablet may be coated with, for example, a mixture of hydroxypropylcellulose and hydroxypropylmethylcellulose, wherein the mixture contains titanium dioxide and/or other coloring agents, such as iron oxide, dyes and lakes; a mixture of polyvinyl alcohol (PVA) and polyethylene glycol (PEG); or any other suitable immediate-release coating agent. The coating provides taste masking and additional stability to the final tablet. The commercially available coating material is Opadry[®] such as Opadry 03K12429 which is a pre-formulated powder mixture provided by Colorcon.

[0142] In one embodiment, in the above-mentioned pharmaceutical combination, pharmaceutical composition or pharmaceutical formulation (preferably, a fixed dose combination formulation), sweetening agents and/or flavoring agents may also be added as needed.

[0143] In one embodiment, the above-mentioned binder is polyvinylpyrrolidone, hydroxypropyl cellulose or hydroxypropyl methyl cellulose, the above-mentioned filler is microcrystalline cellulose, silicified microcrystalline cellulose, lactose, calcium dihydrogen phosphate, mannitol, corn starch or pregelatinized starch, the above-mentioned disintegrant is croscarmellose sodium, crospovidone or

sodium starch glycolate, the above-mentioned lubricant is magnesium stearate or sodium stearyl fumarate, and the above-mentioned glidant is colloidal silicon dioxide.

[0144] In one embodiment, the above-mentioned binder is hydroxypropyl cellulose, the above-mentioned filler is microcrystalline cellulose, silicified microcrystalline cellulose or lactose, the above-mentioned disintegrant is croscarmellose sodium, crospovidone or sodium starch glycolate, the above-mentioned lubricant is magnesium stearate or sodium stearyl fumarate, and the above-mentioned glidant is colloidal silicon dioxide.

[0145] In one embodiment, the above-mentioned binder is polyvinylpyrrolidone, the above-mentioned filler is microcrystalline cellulose or silicified microcrystalline cellulose, the above-mentioned disintegrant is croscarmellose sodium or crospovidone, the above-mentioned lubricant is magnesium stearate or sodium stearyl fumarate, and the above-mentioned glidant is colloidal silicon dioxide.

[0146] In one embodiment, the above-mentioned binder is hydroxypropyl methyl cellulose, the above-mentioned filler is microcrystalline cellulose, silicified microcrystalline cellulose or lactose, the above-mentioned disintegrant is croscarmellose sodium, crospovidone or sodium starch glycolate, the above-mentioned lubricant is magnesium stearate or sodium stearyl fumarate, and the above-mentioned glidant is colloidal silicon dioxide.

[0147] In one embodiment, the above-mentioned binder is hydroxypropyl cellulose, the above-mentioned filler is microcrystalline cellulose, silicified microcrystalline cellulose or lactose, the above-mentioned disintegrant is croscarmellose sodium, and the above-mentioned lubricant is magnesium stearate or sodium stearyl fumarate.

[0148] In one embodiment, the above-mentioned binder is polyvinylpyrrolidone, the above-mentioned lubricant is magnesium stearate, and the above-mentioned glidant is colloidal silicon dioxide.

Preparation method

[0149] In one embodiment, the pharmaceutical composition or fixed dose combination formulation disclosed herein is prepared by wet granulation (high shear and/or fluidized bed). The granulation is a method in which binder(s) is added into a solvent to prepare a binder solution, and then added or directly added into a granulator to prepare wet granules. The wet granulation method includes the steps of:

1. (1) adding an active pharmaceutical ingredient glucokinase activator (preferably, HMS5552) and a partner drug which is a SGLT-2 inhibitor (preferably, empagliflozin, dapagliflozin, canagliflozin) into a granulator;
2. (2) adding optional filler(s) (e.g., microcrystalline cellulose, silicified microcrystalline cellulose, or lactose) into the mixture obtained in step (1);
3. (3) adding optional disintegrant(s) (e.g., croscarmellose sodium, crospovidone, or sodium starch glycolate) into the mixture obtained in step (1) or (2);

4. (4) for high-shear granulation, adding binder(s) (e.g., hydroxypropyl cellulose, polyvinylpyrrolidone or hydroxypropyl methyl cellulose) into pure water to prepare a binder solution, and then adding it into a granulator for granulation with stirring. For fluidized bed granulation, two active pharmaceutical ingredients are added into a fluidized bed, and a binder solution that is an aqueous solution prepared from binder(s) and pure water is sprayed into the fluidized bed by compressed air;
5. (5) sizing the obtained wet granules in a suitable mill to obtain wet granules of suitable size;
6. (6) for granules prepared by high-shear granulation, drying with a tray in an oven or drying in a fluidized bed dryer; and for granules obtained by granulating in a fluidized bed, drying in a fluidized bed;
7. (7) sizing the granules on a suitable grinder to obtain dry granules of suitable size;
8. (8) in a suitable mixer, adding optional filler(s) (diluent(s), such as microcrystalline cellulose) and optional disintegrant(s) (e.g., croscarmellose sodium), and mixing with the dry granules;
9. (9) adding lubricant(s) (e.g., magnesium stearate and sodium stearyl fumarate) into the mixture in step (8);
10. (10) adding optional glidant(s) (e.g., colloidal silicon dioxide) into the mixture in step (9);
11. (11) filling the mixture of lubricated granules in step (9) or (10) into vials, pouches or capsules, or compressing it into a tablet with a desired shape; and
12. (12) optionally, film-coating the obtained tablets.

[0150] In another embodiment, the pharmaceutical composition disclosed herein is prepared by wet granulation (high shear and/or fluidized bed). The granulation is a method in which binder(s) and a second active ingredient are added into a solvent to prepare a binder solution (or suspension), and then added into a granulator to prepare wet granules. The wet granulation method includes the steps of:

1. (1) adding the active pharmaceutical ingredient glucokinase activator (preferably, HMS5552) into a granulator;
2. (2) adding optional filler(s) (e.g., microcrystalline cellulose, silicified microcrystalline cellulose, or lactose) into the mixture in step (1);
3. (3) adding optional disintegrant(s) (e.g., croscarmellose sodium, crospovidone, or sodium starch glycolate) into the mixture obtained in step (1) or (2);
4. (4) for high-shear granulation, adding binder(s) (e.g., hydroxypropyl cellulose, polyvinylpyrrolidone or hydroxypropyl methyl cellulose) into a solvent for uniform dispersion or dissolution, and then adding the second active ingredient (preferably, empagliflozin, dapagliflozin, canagliflozin) in a formulated amount for dispersion or dissolution, so as to form a uniform binder system. The system is added into a granulator for granulation with stirring. For fluidized bed granulation, an active pharmaceutical ingredient such as HMS5552 is added into a fluidized bed, and a binder system that is a solution or suspension prepared by binder(s) and pure water or an organic solvent (e.g., ethanol) is sprayed into the fluidized bed by compressed air;
5. (5) sizing the obtained wet granules in a suitable mill to obtain wet granules of suitable size;
6. (6) for granules prepared by high-shear granulation, drying with a tray in an oven or drying in a fluidized bed dryer; and for granules obtained by granulating in a fluidized bed, drying in a fluidized bed;

7. (7) sizing the granules on a suitable grinder to obtain dry granules of suitable size;
8. (8) in a suitable mixer, adding optional filler(s) (diluent, such as microcrystalline cellulose) and optional disintegrant(s) (e.g., croscarmellose sodium), and mixing with the dry granules;
9. (9) adding lubricant(s) (e.g., magnesium stearate and sodium stearyl fumarate) into the mixture in step (8);
10. (10) optionally, adding optional glidant(s) (e.g., colloidal silicon dioxide) into the mixture in step (9);
11. (11) filling the mixture of lubricated granules in (9) or (10) into vials, pouches or capsules, or compressing it into a tablet with a desired shape; and
12. (12) optionally, film-coating the obtained tablets.

[0151] The dry processing (direct compression or dry granulation) method includes the steps of:

1. (1) adding the active pharmaceutical ingredient glucokinase activator (preferably, HMS5552) and the partner drug which is a SGLT-2 inhibitor (preferably, empagliflozin, dapagliflozin, canagliflozin) into a mixing tank;
2. (2) adding optional filler(s) (e.g., microcrystalline cellulose, silicified microcrystalline cellulose, or lactose) into the mixture in step (1);
3. (3) adding optional binder(s) (e.g., hydroxypropyl cellulose, polyvinylpyrrolidone or hydroxypropyl methyl cellulose) into the mixture obtained in step (1) or (2);
4. (4) adding lubricant(s) or glidant(s) into the mixture in step (3), and mixing;
5. (5) filling the mixture in step (4) into vials, pouches or capsules, compressing the mixture in step (4) into a tablet with a desired shape, or processing the mixture in step (4) by a roller compressor;
6. (6) mixing the mixture in step (3) in advance, and then rolled with a roller if processing the mixture in step (4) by a roller compressor; if necessary, sizing the granules on a suitable grinder to obtain granules of the required size;
7. (7) in a suitable mixer, adding optional diluent(s) to the granules obtained in step (6) to improve the compression performance;
8. (8) adding optional disintegrant(s) (e.g., croscarmellose sodium, crospovidone, or sodium starch glycolate) into the mixture in step (7);
9. (9) adding optional lubricant(s) or glidant(s) into the mixture in step (8);
10. (10) filling the mixture of lubricated granules in step (9) or (10) into vials, pouches or capsules, or compressing it into a tablet with a desired shape; and
11. (11) optionally, film-coating the tablets obtained in step (5) or step (10).

[0152] In one embodiment disclosed herein, the glucokinase activator in the pharmaceutical combination, pharmaceutical composition or fixed dose combination formulation disclosed herein is in the form of a solid dispersion, which can be prepared by a method selected from the group consisting of spray drying method, fluidized bed drying method, solvent method, melt extrusion method and the like.

[0153] One embodiment disclosed herein is a method of preparing a solid dispersion of a

glucokinase activator by a spray drying method, which includes the steps of:

1. (1) formulating a spray drying solution, including dissolving a polymer carrier and a glucokinase activator (preferably, HMS5552) in a solvent;
2. (2) spray drying, including controlling the temperature of the inlet air, the amount of the inlet air, the flow rate and pressure of the atomized airflow, the spraying speed of the solution, and the like.

[0154] In the embodiment disclosed herein, the solvent used in the preparation of the solid dispersion of the glucokinase activator includes but is not limited to alkanols, esters, nitriles, cycloalkanes, aromatic hydrocarbons, ketones and the like. Specifically, the solvent is selected from the group consisting of anhydrous ethanol, methanol, isopropanol, ethyl acetate, acetone, acetonitrile, isobutanol, n-hexane, benzene and toluene. The solvent can be a single solvent, a mixed solvent, or a mixture of an organic solvent and water.

[0155] Methods and uses for treating and/or preventing diseases. Any references to methods of treatment by therapy or surgery or in vivo diagnosis methods of this description is to be interpreted as a reference to compounds, pharmaceutical compositions and medicaments of the present invention for use in those methods.

[0156] Another embodiment disclosed herein relates to a method or use of a composition or formulation (preferably, a fixed dose combination pharmaceutical composition or a fixed dose combination formulation) comprising a glucokinase activator disclosed herein for the treatment and/or prevention of the following diseases and medical disorders, especially one or more diseases selected from the group consisting of type I diabetes, type II diabetes, impaired glucose tolerance, impaired fasting blood glucose, and hyperglycemia, including administering to a subject a therapeutically effective amount of the composition or preparation (preferably, a fixed dose combination pharmaceutical composition or a fixed dose combination formulation) disclosed herein:

- preventing, slowing the progression of, delaying or treating a metabolic disorder selected from the group consisting of type 1 diabetes mellitus, type 2 diabetes mellitus, impaired glucose tolerance, impaired fasting blood glucose, hyperglycemia, postprandial hyperglycemia, overweight, obesity and metabolic syndrome; or
- improving glycemic control and/or for reducing of fasting plasma glucose, of postprandial plasma glucose and/or of glycosylated hemoglobin HbA1c; or
- preventing, slowing, delaying or reversing progression from impaired glucose tolerance, insulin resistance and/or from metabolic syndrome to type 2 diabetes mellitus; or
- preventing, slowing the progression of, delaying or treating of a condition or disorder selected from the group consisting of complications of diabetes mellitus such as cataracts and micro- and macrovascular diseases, such as nephropathy, retinopathy, neuropathy, learning and memory dysfunction, neurodegenerative or cognitive disorders, cardio- or cerebrovascular diseases, tissue ischaemia, diabetic foot or ulcer, arteriosclerosis, hypertension, endothelial dysfunction, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable angina pectoris, stroke, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders and vascular restenosis; or

- reducing body weight and/or body fat or preventing an increase in body weight and/or body fat or facilitating a reduction in body weight and/or body fat; or
- preventing, slowing, delaying or treating the degeneration of pancreatic beta cells and/or the decline of the functionality of pancreatic beta cells and/or for improving and/or restoring or protecting the functionality of pancreatic beta cells and/or restoring the functionality of pancreatic insulin secretion; or
- for preventing, slowing, delaying or treating diseases or conditions attributed to an abnormal accumulation of liver or ectopic fat; or
- maintaining and/or improving the insulin sensitivity and/or for treating or preventing hyperinsulinemia and/or insulin resistance; or
- for preventing, slowing progression of, delaying, or treating new onset diabetes after transplantation (NODAT) and/or post-transplant metabolic syndrome (PTMS); or
- for preventing, delaying, or reducing NODAT and/or PTMS associated complications including micro- and macrovascular diseases and events, graft rejection, infection, and death; or
- for treating hyperuricemia and hyperuricemia associated conditions; or
- diabetic nephropathy, decreased renal function.

[0157] The present disclosure also provides a method for the treatment of type II diabetes by orally administering a therapeutically effective amount of a pharmaceutical composition or formulation (preferably, a fixed dose combination pharmaceutical composition or a fixed dose combination formulation) comprising a glucokinase activator and a partner drug disclosed herein to a subject in need of the treatment. In one embodiment, the subject in need of the treatment is a human. In another embodiment, the pharmaceutical composition is in the form of a tablet. The composition or formulation (preferably, a fixed dose combination pharmaceutical composition or a fixed dose combination formulation) comprising a glucokinase activator disclosed herein can be administered once a day (QD), twice a day or three times a day (TID).

[0158] The following examples further describe and illustrate embodiments within the scope of the present disclosure.

Examples

Preparation of the combination tablet of the glucokinase activator

[0159] The chemicals used in the present disclosure can be purchased from companies such as Shin-Etsu Japan, Evonik Germany, J.T. Baker US, SCR China, Ashland US, FMC US, JRS Germany, Colorcon US, Capsugel, BASF, Zhenxing China, and the like. A producing equipment, analytical testing instrument and the like can be purchased from companies such as Sartorius, Nikon, Sympatec, Bruker, Gea Niro, Korsch, Erweka, Agilent, Quadro Engineering, Canada; Waters, US; TA, US; SOTAX, Switzerland; Mettler Toledo Instrument Newark, DE.

I. Preparation of the solid dispersion of the glucokinase activator

1.1 Preparation of the solution of the solid dispersion used for spray drying**Example 1A (weight ratio of active ingredients to polymer carriers is 1:9)**

[0160] 6.75g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 0.75g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 2A (weight ratio of active ingredients to polymer carriers is 3:7)

[0161] 5.25g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 2.25g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 3A (weight ratio of active ingredients to polymer carriers is 5:5)

[0162] 3.75g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 3.75g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 4A (weight ratio of active ingredients to polymer carriers is 7:3)

[0163] 2.25g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 5.25g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 5A (weight ratio of active ingredients to polymer carriers is 8:2)

[0164] 1.5g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 6g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 6A (weight ratio of active ingredients to polymer carriers is 9:1)

[0165] 0.75g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 6.75 g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 7A (weight ratio of active ingredients to polymer carriers is 6:4)

[0166] 3.0 g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 4.5 g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 8A (weight ratio of active ingredients to polymer carriers is 4:6)

[0167] 4.5g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (J.T.Baker) under stirring. After it was completely dissolved, 3.0 g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 50 ml solution.

Example 9A (weight ratio of active ingredients to polymer carriers is 5:5)

[0168] 187.5 g Eudragit L100 (Evonik Germany) was weighed, and added to anhydrous ethanol (Zhenxing China). After it was completely dissolved, 187.5 g of the compound HMS5552 was added. Stirring was continued after adding sufficient amount of anhydrous ethanol to obtain 2500ml solution.

1.2 Preparation of the solid dispersion of the glucokinase activator

[0169] The solid dispersion of the glucokinase activator was prepared by spray drying the solution prepared above. The numbering of the obtained solid dispersion corresponds to the numbering of the above examples. The spray drying devices that are suitable for the present disclosure include, but are not limited to, the spray drying devices produced by Niro GEA Process Engineering Inc., Buchi Labortechnik AG, ProCept and SPX ANHYDROUS companies. The spray drying can be performed by selecting an appropriate inlet air temperature of dry gas, inlet amount, feed rate, and atomization pressure, so that the droplets are sufficiently dried as they reach the device wall. This can make sure that the dried droplets are essentially solid and in a form of a fine powder, which will not stick to the wall, and is not difficult to collect in the cyclone. The resulting powder is subjected to

a secondary drying to make sure the product meets quality requirement.

Description of the production process for the preparation of the solid dispersion of the glucokinase activator by spray drying

[0170] The solid dispersions were prepared by the spray drying the solution prepared in the above Examples **1A-8A**, wherein the inlet air temperature of the spray dryer was 90-150 °C, the flow rate of the inlet air was 0.3-0.5 m³/min, the flow rate of the air flow was 15-30 L/min, and the spray rate of above solutions were 5-7 mL/min. Solid dispersions **1-8** were obtained by spray drying.

[0171] The solid dispersion was prepared by spray drying the solution prepared in the above Example **9A**, wherein the inlet air temperature of the spray dryer was 90-150 °C, the flow rate of the inlet air was 20-30 kg/h, the flow rate of the air flow was 3-30 kg/h, and the spray rate of above solutions were 5-200 mL/min. Solid dispersion **9** was obtained by spray drying.

[0172] Solid dispersions **1-9** were prepared according to the process described above, wherein: mass percent of the compound HMS5552 in solid dispersion **1** was 10%; mass percent of the compound HMS5552 in solid dispersion **2** was 30%; mass percent of the compound HMS5552 in solid dispersion **3** was 50%; mass percent of the compound HMS5552 in solid dispersion **4** was 70%; mass percent of the compound HMS5552 in solid dispersion **5** was 80%; mass percent of the compound HMS5552 in solid dispersion **6** was 90%; mass percent of the compound HMS5552 in solid dispersion **7** was 60%; mass percent of the compound HMS5552 in solid dispersion **8** was 40%; and mass percent of the compound HMS5552 in solid dispersion **9** was 50%.

II. Preparation of the combination tablet

2.1 Preparation of the combination tablet by high-shear wet granulation

[0173] The HMS5552 solid dispersion prepared according to the above preparation examples of the solid dispersion of the glucokinase activator and the partner drug which is a SGLT-2 inhibitor were added into a high-shear wet granulator. The filler (e.g., microcrystalline cellulose, silicified microcrystalline cellulose, or lactose) and disintegrant (e.g., croscarmellose sodium, crospovidone, or sodium starch glycolate) were added. A part of the binder powder was added and mixed for 5 minutes with high-shear stirring. A prepared solution of the binder (e.g., hydroxypropyl cellulose, polyvinylpyrrolidone or hydroxypropyl methyl cellulose) was added to the above dry mixture under high-shear stirring for 1 to 6 minutes for granulation. Wet granules are sized on a Comil mill to obtain wet granules of suitable size. The wet granules were dried with a tray in an oven at about 60°C or in a fluidized bed dryer (with the inlet air temperature of 40-60°C) for 20-40 minutes. Then, the dried material was ground using a grinder to obtain granules of suitable size. After grinding, the microcrystalline cellulose or silicified microcrystalline cellulose (for the filler comprising an extragranular part) and the disintegrant (e.g., croscarmellose sodium, crospovidone, or sodium starch glycolate) were added to the granules and the mixture was mixed in a barrel mixer. Then,

the lubricant (magnesium stearate or sodium stearyl fumarate) and/or optional glidant (micronized silica gel) were added to the mixture and mixed well additionally. The lubricated mixture was compressed with a rotary tablet press to obtain tablets (plain tablets, uncoated tablet cores) of different tablet weights and tablet shapes corresponding to different strengths. Optionally, the obtained tablets were film-coated with Opadry® II, and the weight increased by about 3%, thereby obtaining film-coated tablets.

Example 1B A combination tablet of HMS5552 + empagliflozin (dose strength: 75 mg/5 mg)

[0174]

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	5.00	1.92
HMS5552 solid dispersion*	150.00	57.69
Microcrystalline cellulose	88.10	33.88
Hydroxypropyl cellulose	7.80	3.00
Croscarmellose sodium	6.50	2.50
Magnesium stearate	2.60	1.00
Total weight of a tablet core	260.0	100.0
Opadry	7.80	3.00
Total weight of a coated tablet	267.80	-
*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552.		

Example 2B A combination tablet of HMS5552+empagliflozin (dose strength: 75 mg/12.5 mg)

[0175]

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	12.50	4.46
HMS5552 solid dispersion*	150.00	53.57
Microcrystalline cellulose	99.30	35.47
Hydroxypropyl cellulose	8.40	3.00
Croscarmellose sodium	7.00	2.50
Magnesium stearate	2.80	1.00
Total weight of a tablet core	280.00	100.0
Opadry	8.40	3.00
Total weight of a coated tablet	288.40	-
*150.0 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552.		

Example 3B A combination tablet of HMS5552 + empagliflozin (dose strength: 75 mg/10 mg)**[0176]**

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	10.00	3.85
HMS5552 solid dispersion*	150.00	57.69
Microcrystalline cellulose	84.40	32.46
Hydroxypropyl cellulose	7.80	3.00
Croscarmellose sodium	5.20	2.00
Magnesium stearate	2.60	1.00
Total weight of a tablet core	260.00	100.00
Opadry	7.80	3.00
Total weight of a coated tablet	267.80	-
*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552.		

Example 4B A combination tablet of HMS5552 + empagliflozin (dose strength: 75 mg/25 mg)**[0177]**

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	25.00	8.92
HMS5552 solid dispersion*	150.00	53.58
Microcrystalline cellulose	88.20	31.50
Hydroxypropyl cellulose	8.40	3.00
Croscarmellose sodium	5.60	2.00
Magnesium stearate	2.80	1.00
Total weight of a tablet core	280.00	100.00
Opadry	8.40	3.00
Total weight of a coated tablet	288.40	-
*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552.		

Example 5B A combination tablet of HMS5552 + empagliflozin (dose strength: 100 mg/10 mg)**[0178]**

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	10.00	3.33
HMS5552 solid dispersion*	200.00	66.67
Microcrystalline cellulose	72.00	24.00
Hydroxypropyl cellulose	9.00	3.00
Croscarmellose sodium	6.00	2.00
Magnesium stearate	3.00	1.00
Total weight of a tablet core	300.00	100.00
Opadry	9.00	3.00
Total weight of a coated tablet	309.00	-

*200.00 mg of the HMS5552 solid dispersion contained 100 mg of HMS5552.

Example 6B A combination tablet of HMS5552 + dapagliflozin (dose strength: 75 mg/5 mg)

[0179]

Formula composition	Unit formula amount/mg	% (w/w)
Dapagliflozin propylene glycol monohydrate**	6.15	2.36
HMS5552 solid dispersion*	150.00	57.70
Microcrystalline cellulose	88.25	33.94
Hydroxypropyl cellulose	7.80	3.00
Croscarmellose sodium	5.20	2.00
Magnesium stearate	2.60	1.00
Total weight of a tablet core	260.00	100.00
Opadry	7.80	3.00
Total weight of a coated tablet	267.80	-

*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552;
**6.15 mg of dapagliflozin propylene glycol monohydrate was equivalent to 5.0 mg of dapagliflozin free base anhydrate.

Example 7B A combination tablet of HMS5552 + dapagliflozin (dose strength: 50 mg/5 mg)

[0180]

Formula composition	Unit formula amount/mg	% (w/w)
Dapagliflozin propylene glycol monohydrate**	6.15	2.80

Formula composition	Unit formula amount/mg	% (w/w)
HMS5552 solid dispersion*	100.00	45.45
Microcrystalline cellulose	100.65	45.75
Hydroxypropyl cellulose	6.60	3.00
Croscarmellose sodium	4.40	2.00
Magnesium stearate	2.20	1.00
Total weight of a tablet core	220.00	100.00
Opadry	6.60	3.00
Total weight of a coated tablet	266.60	-
*100.00 mg of the HMS5552 solid dispersion contained 50 mg of HMS5552; **6.15 mg of dapagliflozin propylene glycol monohydrate was equivalent to 5.0 mg of dapagliflozin free base anhydrate.		

Example 8B A combination tablet of HMS5552 + dapagliflozin (dose strength: 75 mg/10 mg)

[0181]

Formula composition	Unit formula amount/mg	% (w/w)
Dapagliflozin propylene glycol monohydrate**	12.30	4.56
HMS5552 solid dispersion*	150.00	55.56
Microcrystalline cellulose	91.50	33.89
Hydroxypropyl cellulose	8.10	3.00
Croscarmellose sodium	5.40	2.00
Magnesium stearate	2.70	1.00
Total weight of a tablet core	270.00	100.00
Opadry	8.10	3.00
Total weight of a coated tablet	278.10	-
*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552; ** 12.30 mg of dapagliflozin propylene glycol monohydrate was equivalent to 10 mg of dapagliflozin free base anhydrate.		

Example 9B A combination tablet of HMS5552 + dapagliflozin (dose strength: 100 mg/5 mg)

[0182]

Formula composition	Unit formula amount/mg	% (w/w)
Dapagliflozin propylene glycol monohydrate**	6.15	2.05
HMS5552 solid dispersion*	200.00	66.67
Microcrystalline cellulose	75.55	25.18
Hydroxypropyl cellulose	9.00	3.00
Croscarmellose sodium	6.30	2.00
Magnesium stearate	3.00	1.00
Total weight of a tablet core	300.00	100.00
Opadry	9.00	3.00
Total weight of a coated tablet	309.00	-
*200.00 mg of the HMS5552 solid dispersion contained 100 mg of HMS5552; **6.15 mg of dapagliflozin propylene glycol monohydrate was equivalent to 5.0 mg of dapagliflozin free base anhydrate.		

Example 10B A combination tablet of HMS5552+ canagliflozin (dose strength: 75 mg/100 mg)

[0183]

Formula composition	Unit formula amount/mg	% (w/w)
Canagliflozin hemihydrate**	101.93	16.99
HMS5552 solid dispersion*	150.00	25.00
Microcrystalline cellulose	294.07	49.01
Hydroxypropyl cellulose	18.00	3.00
Croscarmellose sodium	30.00	5.00
Magnesium stearate	6.00	1.00
Total weight of a tablet core	600.00	100.00
Opadry	18.00	3.00
Total weight of a coated tablet	618.00	-
*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552; **101.93 mg of canagliflozin hemihydrate was equivalent to 100 mg of canagliflozin free base anhydrate.		

Example 11B A combination tablet of HMS5552+ canagliflozin (dose strength: 50 mg/100 mg)

[0184]

Formula composition	Unit formula amount/mg	% (w/w)
Canagliflozin hemihydrate**	101.93	16.99
HMS5552 solid dispersion*	100.00	16.67
Microcrystalline cellulose	344.07	57.35
Hydroxypropyl cellulose	18.00	3.00
Croscarmellose sodium	30.00	5.00
Magnesium stearate	6.00	1.00
Total weight of a tablet core	600.0	100.00
Opadry	18.00	3.00
Total weight of a coated tablet	618.00	-
* 100.00 mg of the HMS5552 solid dispersion contained 50 mg of HMS5552; **101.93 mg of canagliflozin hemihydrate was equivalent to 100 mg of canagliflozin free base anhydrate.		

Example 12B A combination tablet of HMS5552+ canagliflozin (dose strength: 100 mg/100 mg)

[0185]

Formula composition	Unit formula amount/mg	% (w/w)
Canagliflozin hemihydrate**	101.93	16.99
HMS5552 solid dispersion*	200.00	33.33
Microcrystalline cellulose	244.07	40.68
Hydroxypropyl cellulose	18.00	3.00
Croscarmellose sodium	30.0	5.00
Magnesium stearate	6.00	1.00
Total weight of a tablet core	600.00	100.00
Opadry	18.00	3.00
Total weight of a coated tablet	618.00	-
*200.00 mg of the HMS5552 solid dispersion contained 100 mg of HMS5552; **101.93 mg of canagliflozin hemihydrate was equivalent to 100 mg of canagliflozin free base anhydrate.		

Example 13B A combination tablet of HMS5552+ canagliflozin (dose strength: 75 mg/300 mg)

[0186]

Formula composition	Unit formula amount/mg	% (w/w)
Canagliflozin hemihydrate**	305.78	40.77
HMS5552 solid dispersion*	150.00	20.00
Microcrystalline cellulose	226.72	30.23
Hydroxypropyl cellulose	22.50	3.00
Croscarmellose sodium	37.50	5.00
Magnesium stearate	7.50	1.00
Total weight of a tablet core	750.00	100.00
Opadry	22.50	3.00
Total weight of a coated tablet	772.50	-
*200.00 mg of the HMS5552 solid dispersion contained 100 mg of HMS5552; **305.78 mg of canagliflozin hemihydrate was equivalent to 300 mg of canagliflozin free base anhydrate.		

Example 14B A combination tablet of HMS5552+ canagliflozin (dose strength: 50 mg/300 mg)

[0187]

Formula composition	Unit formula amount/mg	% (w/w)
Canagliflozin hemihydrate**	305.78	40.77
HMS5552 solid dispersion*	100.00	13.33
Microcrystalline cellulose	276.72	36.90
Hydroxypropyl cellulose	22.50	3.00
Croscarmellose sodium	37.50	5.00
Magnesium stearate	7.50	1.00
Total weight of a tablet core	750.00	100.00
Opadry	22.50	3.00
Total weight of a coated tablet	772.50	-
* 100.00 mg of the HMS5552 solid dispersion contained 50 mg of HMS5552; **305.78 mg of canagliflozin hemihydrate was equivalent to 300 mg of canagliflozin free base anhydrate.		

Example 15B A combination tablet of HMS5552+ canagliflozin (dose strength: 25 mg/300 mg)

[0188]

Formula composition	Unit formula amount/mg	% (w/w)
Canagliflozin hemihydrate**	305.78	40.77
HMS5552 solid dispersion*	50.00	6.67
Microcrystalline cellulose	326.72	43.56
Hydroxypropyl cellulose	22.50	3.00
Croscarmellose sodium	37.50	5.00
Magnesium stearate	7.50	1.00
Total weight of a tablet core	750.00	100.00
Opadry	22.50	3.00
Total weight of a coated tablet	772.50	-
*50.00 mg of the HMS5552 solid dispersion contained 25 mg of HMS5552; **305.78 mg of canagliflozin hemihydrate was equivalent to 300 mg of canagliflozin free base anhydrate.		

2.2 Preparation of the combination tablet by fluidized bed wet granulation

[0189] The HMS5552 solid dispersion prepared according to the above preparation examples of the solid dispersion of the glucokinase activator and the partner drug which is a SGLT-2 inhibitor were added into a fluidized bed granulator. The optional filler (e.g., microcrystalline cellulose) was added. The prepared solution of the binder (e.g., polyvinylpyrrolidone) was sprayed into the mixture in the fluidized bed during 20 to 60 minutes for granulation, and then drying was continued in a fluidized bed dryer (with the inlet air temperature of 40-60°C). Then, the dried material was ground using a grinder to obtain granules of suitable size. After grinding, the microcrystalline cellulose or silicified microcrystalline cellulose (for the formulation comprising an extragranular filler) was added to the granules, and the mixture was mixed in a barrel mixer. Then, the lubricant (magnesium stearate) and/or optional glidant (micronized silica gel) were added to the mixture and mixed well additionally. The lubricated mixture was compressed with a rotary tablet press to obtain tablets (plain tablets, uncoated tablet cores) of different tablet weights and tablet shapes corresponding to different strengths. Optionally, the obtained tablets were film-coated according to the above preparation examples of the solid dispersion of the glucokinase activator, and the weight increased by about 3%, thereby obtaining film-coated tablets.

Example 16B A combination tablet of HMS5552+empagliflozin (dose strength: 50 mg/10 mg)

[0190]

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	10.00	4.50
HMS5552 solid dispersion*	100.00	45.50
Microcrystalline cellulose	96.80	44.00

Formula composition	Unit formula amount/mg	% (w/w)
Povidone	6.60	3.00
Croscarmellose sodium	4.40	2.00
Magnesium stearate	2.20	1.00
Total weight of a tablet core	220.00	100.0
Opadry	6.60	3.00
Total weight of a coated tablet	226.60	-
* 100.00 mg of the HMS5552 solid dispersion contained 50 mg of HMS5552;		

2.3 Preparation of the combination tablet by dry rolling granulation

[0191] The HMS5552 solid dispersion prepared according to the above preparation examples of the solid dispersion of the glucokinase activator and the partner drug which is a SGLT-2 inhibitor were added into a mixing tank. The filler (e.g., microcrystalline cellulose) and the binder (e.g., hydroxypropyl cellulose) were added and mixed well. Then, the mixture was rolled by a roller compaction granulator, and the obtained bar was crushed and sized by a crusher to obtain granules of suitable size. After grinding, the optional microcrystalline cellulose or silicified microcrystalline cellulose (for the filler comprising an extragranular part) and the disintegrant (e.g., croscarmellose sodium) were added to the granules, and the mixture was mixed in a barrel mixer. Then, the lubricant (magnesium stearate or sodium stearyl fumarate) and/or optional glidant (micronized silica gel) were added to the mixture and mixed well additionally. The lubricated mixture was compressed with a rotary tablet press to obtain tablets (plain tablets, uncoated tablet cores) of different tablet weights and tablet shapes corresponding to different strengths. Optionally, the obtained tablets were film-coated with Opadry® II, and the weight increased by about 3%, thereby obtaining film-coated tablets.

Example 17B A combination tablet of HMS5552 + dapagliflozin (dose strength: 50 mg/10 mg)

[0192]

Formula composition	Unit formula amount/mg	% (w/w)
Dapagliflozin propylene glycol monohydrate**	12.30	5.13
HMS5552 solid dispersion*	100.00	41.67
Microcrystalline cellulose	113.06	47.11
Hydroxypropyl cellulose	7.20	3.00
Croscarmellose sodium	5.04	2.10
Magnesium stearate	2.40	1.00
Total weight of a tablet core	240.00	100.00

Formula composition	Unit formula amount/mg	% (w/w)
Opadry	7.20	3.00
Total weight of a coated tablet	247.20	-

*150.00 mg of the HMS5552 solid dispersion contained 75 mg of HMS5552;
** 12.30 mg of dapagliflozin propylene glycol monohydrate was equivalent to 10 mg of dapagliflozin free base anhydrate.

Example 18B A combination tablet of HMS5552+empagliflozin (dose strength: 50 mg/25 mg)

[0193]

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	25.00	10.40
HMS5552 solid dispersion*	100.00	41.70
Microcrystalline cellulose	100.30	41.80
Hydroxypropyl cellulose	7.20	3.00
Croscarmellose sodium	5.10	2.13
Magnesium stearate	2.40	1.00
Total weight of a tablet core	240.00	100.00
Opadry	7.20	3.00
Total weight of a coated tablet	247.20	-

* 100.00 mg of the HMS5552 solid dispersion contained 50 mg of HMS5552;

2.4 Preparation of the combination tablet by direct compression of the powder mixture

[0194] The HMS5552 solid dispersion prepared according to the above preparation examples of the solid dispersion of the glucokinase activator and the partner drug which is a SGLT-2 inhibitor were premixed uniformly according to the principle of geometric progression, and then added to a mixing tank. The filler (e.g., microcrystalline cellulose), the disintegrant (e.g., croscarmellose sodium) and optional glidant (micronized silica gel) were added to the granules and mixed in a barrel mixer. Then, the lubricant (magnesium stearate or sodium stearyl fumarate) was added to the mixture and mixed well additionally. The lubricated mixture was compressed with a rotary tablet press to obtain tablets (plain tablets, uncoated tablet cores) of different tablet weights and tablet shapes corresponding to different strengths. Optionally, the obtained tablets were film-coated with Opadry® II, and the weight increased by about 3%, thereby obtaining film-coated tablets.

[0195] The formula composition of the combination tablet described in the above preparation process was:

Example 19B A combination tablet of HMS5552 + empagliflozin (dose strength: 25 mg/25 mg)

[0196]

Formula composition	Unit formula amount/mg	% (w/w)
Empagliflozin	25.00	6.25
HMS5552 solid dispersion*	50.00	12.50
Microcrystalline cellulose	309.00	77.25
Croscarmellose sodium	8.00	2.00
Micronized silica gel	4.00	1.00
Magnesium stearate	4.00	1.00
Total weight of a tablet core	400.00	100.00
Opadry	12.00	3.00
Total weight of a coated tablet	412.00	-
*50.00 mg of the HMS5552 solid dispersion contained 25 mg of HMS5552.		

Example 20B A combination tablet of HMS5552 + dapagliflozin (dose strength: 25 mg/10 mg)

[0197]

Formula composition	Unit formula amount/mg	% (w/w)
Dapagliflozin propylene glycol monohydrate**	12.30	3.075
HMS5552 solid dispersion*	50.00	12.50
Microcrystalline cellulose	313.70	69.04
Hydroxypropyl cellulose	8.00	2.00
Croscarmellose sodium	8.00	2.00
micronized silica gel	4.00	1.00
Magnesium stearate	4.00	1.00
Total weight of a tablet core	400.00	100.00
Opadry	12.00	3.00
Total weight of a coated tablet	412.00	-
*50.00 mg of the HMS5552 solid dispersion contained 25 mg of HMS5552;		
** 12.30 mg of dapagliflozin propylene glycol monohydrate was equivalent to 10 mg of dapagliflozin free base anhydrate.		

III. Dissolution test in vitro of the combination formulation comprising the glucokinase

activator

[0198] The dissolution rate of the tablet was tested by the paddle method of the Chinese Pharmacopoeia (2015 edition). The dissolution of HMS5552 and another partner drug which is a SGLT-2 inhibitor in the medium of pH 6.8 was tested, respectively. At 5 minutes, 15 minutes, 30 minutes, 45 minutes, and 60 minutes, respectively, 5 ml of samples were taken for HPLC analysis.

[0199] According to the above test method, several tablets of above-mentioned fixed dose strengths and their corresponding single tablets were tested for their dissolution, and the results were shown below.

Table 1 The dissolution results of the fixed dose combination tablets prepared in Example 1B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 1B-HMS5552, 75 mg	77.1	89.5	95.6	97.3	98.3
Example 1B-empagliflozin, 5 mg	53.8	85.3	96.3	99.0	100

Table 2 The dissolution results of the fixed dose combination tablets prepared in Example 4B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 4B-HMS5552, 75 mg	58.8	80.0	88.8	92.9	95.1
Example 4B-empagliflozin, 25 mg	63.7	84.1	92.0	95.6	97.5

Table 3 The dissolution results of the fixed dose combination tablets prepared in Example 5B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 5B-HMS5552, 100 mg	82.2	95.0	97.4	97.9	98.0
Example 5B-empagliflozin, 10 mg	84.1	94.0	96.1	96.7	97.2

Table 4 The dissolution results of the fixed dose combination tablets prepared in Example 7B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 7B-HMS5552, 50 mg	54.8	87.8	93.7	95.5	97.1
Example 7B-dapagliflozin, 5 mg	38.7	77.6	94.4	99.8	103.1

Table 5 The dissolution results of the fixed dose combination tablets prepared in Example 8B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 8B-HMS5552, 75 mg	56.8	88.8	95.7	97.6	99.2

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 8B-dapagliflozin, 10 mg	32.5	68.7	97.0	93.0	96.6

Table 6 The dissolution results of the fixed dose combination tablets prepared in Example 9B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 9B-HMS5552, 100 mg	76.8	98.8	101.4	101.8	101.9
Example 9B-dapagliflozin, 5 mg	49.0	82.3	95.9	98.6	99.3

Table 7 The dissolution results of the fixed dose combination tablets prepared in Example 19B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 19B-HMS5552, 25 mg	71.0	88.9	93.9	94.9	96.2
Example 19B-empagliflozin, 25 mg	67.4	87.7	94.3	95.8	97.3

Table 8 The dissolution results of the fixed dose combination tablets prepared in Example 20B

Time point	Dissolution rate (%)				
	5 min	15 min	30 min	45 min	60 min
Example 20B-HMS5552, 25 mg	61.0	82.8	88.5	90.5	94.3
Example 20B-dapagliflozin, 10 mg	48.7	80.0	92.5	96.6	101.9

[0200] It can be seen from the above dissolution results of the fixed dose combination formulations that the dissolution of the fixed dose combination formulations disclosed herein meets the requirements of a fast-release formulation.

IV. Physical properties of the combination formulations containing the glucokinase activator

[0201] According to the relevant instruments and methods described in the Pharmacopoeia, the physical properties of several tablets of above-mentioned fixed-dose strengths were tested. The results were described below.

Table 9 Physical properties of the fixed dose combination tablet cores prepared in different examples

Ingredient	Dose strength (HMS5552/empagliflozin), mg			
	75/5	75/12.5	75/10	75/25
Corresponding Example	Example 1B	Example 2B	Example 3B	Example 4B
Tablet shape	Round biconvex	Round biconvex	Round biconvex	Round biconvex

Ingredient	Dose strength (HMS5552/empagliflozin), mg			
	75/5	75/12.5	75/10	75/25
Tablet core size/mm	8.73	8.73	8.73	8.73
Colour	white	white	white	white
Weight/mg	260	280	260	280
Hardness/kp (average value)	≥7	≥7	≥7	≥7
Disintegration time/minute	≤10	≤10	≤10	≤10
Fragility/%	≤0.5	≤0.5	≤0.5	≤0.5

Table 10 Physical properties of the fixed dose combination tablet cores prepared in different examples

Ingredient	Dose strength (HMS5552/empagliflozin), mg			
	100/10	50/10	50/25	25/25
Corresponding Example	Example 5B	Example 16B	Example 18B	Example 19B
Tablet shape	Round biconvex	Round biconvex	Round biconvex	Round biconvex
Tablet core size/mm	8.73	8.73	8.73	8.73
Colour	white	white	white	white
Weight/mg	300	220	240	400
Hardness/kp (average value)	≥7	≥7	≥7	≥7
Disintegration time/minute	≤10	≤10	≤10	≤10
Fragility/%	≤0.5	≤0.5	≤0.5	≤0.5

Table 11 Physical properties of the fixed dose combination tablet cores prepared in different examples

Ingredient	Dose strength (HMS5552/dapagliflozin), mg					
	75/5	50/5	75/10	100/5	50/10	25/10
Corresponding Example	Example 6B	Example 7B	Example 8B	Example 9B	Example 17B	Example 20B
Tablet shape	Round biconvex	Round biconvex	Round biconvex	Round biconvex	Round biconvex	Round biconvex
Tablet core size/mm	8.73	8.73	8.73	8.73	8.73	8.73
Colour	white	white	white	white	white	white
Weight/mg	260	220	270	300	240	400
Hardness/kp (average value)	≥7	≥7	≥7	≥7	≥7	≥10
Disintegration time/minute	≤10	≤10	≤10	≤10	≤10	≤10

Ingredient	Dose strength (HMS5552/dapagliflozin), mg					
	75/5	50/5	75/10	100/5	50/10	25/10
Fragility/%	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5

Table 12 Physical properties of the fixed dose combination tablet cores prepared in different examples

Ingredient	Dose strength (HMS5552/canagliflozin), mg					
	75/100	50/100	100/100	75/300	50/300	25/300
Corresponding Example	Example 10B	Example 11B	Example 12B	Example 13B	Example 14B	Example 15B
Tablet shape	Oval biconvex	Oval biconvex	Oval biconvex	Oval biconvex	Oval biconvex	Oval biconvex
Tablet core size/mm	16.60*8.45	16.60*8.45	16.60*8.45	19.05*9.73	19.05*9.73	19.05*9.73
Colour	white	white	white	white	white	white
Weight/mg	600	600	600	750	750	750
Hardness/kp (average value)	≥15	≥15	≥15	≥15	≥15	≥15
Disintegration time/minute	≤10	≤10	≤10	≤10	≤10	≤10
Fragility/%	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5	≤0.5

V. Study on pharmacodynamics of the combination formulations comprising the glucokinase activator

Example 1C

Study on the effect of the combination of the glucokinase activator and the partner drug on glucose/sucrose tolerance in normal mice

[0202] Normal male C57BL/6J mice, after being fasted for 6 hours, were orally administered solvent control, 10 mg/kg of empagliflozin, or a combination of 10 mg/kg of HMS5552 and 10 mg/kg of empagliflozin, respectively; and after 1 hour, were orally given 2 g/kg of glucose. Blood was taken from the tail vein before being administered (-60 minutes), before being given the glucose (0 minutes), and at 15, 30, 60, and 120 minutes after being given the glucose, and the glucose content in the whole blood was determined. The area under the curve for blood glucose between 0 and 120 minutes (AUC_{0-120 min}, mmol/L*min) was analyzed and compared with that of the solvent control group. The results showed that the hypoglycemic effect of the combination of 10 mg/kg of HMS5552 and 10 mg/kg of empagliflozin is significantly better than that of the monotherapy of 10

mg /kg of empagliflozin, with statistically significant difference of $P < 0.001$.

[0203] Empagliflozin, a sodium-glucose cotransporter 2 (SGLT-2) inhibitor, reduces blood glucose by reducing renal glucose reabsorption, lowering renal glucose threshold, and promoting excretion of glucose from urine, while reducing cardiovascular risk and cardiovascular death, and is used for type 2 diabetes accompanied with a cardiovascular disease. HMS5552, a novel glucokinase activator, can improve pancreatic islet function in patients with type 2 diabetes, promote the secretion of incretin, reduce insulin resistance, and has dual therapeutic effects of reducing fasting and postprandial blood glucose. For patients whose blood glucose control fails with a SGLT-2 inhibitor, obesity patients and diabetes patients accompanied with a cardiovascular disease, the combination of a SGLT-2 inhibitor and HMS5552 has better efficacy of blood glucose control and reduces the risk of diabetes complications.

[0204] The above study on the effectiveness of HMS5552 combined with the existing oral diabetes drug (a SGLT-2 inhibitor) shows that the combined use can improve the efficacy of HMS5552 or existing hypoglycemic drugs, reduce safety risks, and improve medical effects. Oral fixed dose combination formulations developed with HMS5552 and existing oral diabetes drugs are currently the most promising combination drugs for diabetes treatment to solve the above clinical needs.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [CN201810556685](#) [0001]
- [US20120142636A1](#) [0007]
- [EP1734040A1](#) [0007]

Non-patent literature cited in the description


- **INTERNATIONAL DIABETES FEDERATION** Diabetes Atlas, 2015, [0004]
- **ZHU XIAO-XUE et al.** Diabetes, Obesity and Metabolism, 2018, vol. 20, 2113-2120 [0007]

- Chinese Pharmacopoeia 20150000 [\[0198\]](#)

Patentkrav

1. Farmaceutisk sammensætning, omfattende:

- (a) en glucokinase-aktivator, hvor glucokinase-aktivatoren er en forbindelse repræsenteret med følgende formler, eller et farmaceutisk acceptabelt salt, en isotopmærket forbindelse, en krystallinsk form, et hydrat eller et solvat deraf,



og

(b) en SGLT-2-inhibitor.


10

2. Den farmaceutiske sammensætning ifølge krav 1, yderligere omfattende en eller flere excipienser.

3. Den farmaceutiske sammensætning ifølge krav 1 eller 2, hvor vægtforholdet mellem glucokinase-aktivatoren og SGLT-2-inhibitoren er ca. 30:1 til 1:30, fortrinsvis ca. 20:1 til 1:12, eller mere fortrinsvis ca. 0,75:1, ca. 1:2, ca. 1:1, ca. 1:4, ca. 1:6, ca. 1:12, ca. 2:1, ca. 2,5:1, ca. 3:1, ca. 5:1, ca. 6:1, ca. 7,5:1, ca. 10:1, ca. 15:1 eller ca. 20:1, eventuelt hvor glucokinase-aktivatoren er ca. 1-96

vægtprocent; og SGLT-2-inhibitoren er ca. 0,1-80 vægtprocent.

- 4.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 3, hvor glucokinase-aktivatoren er forbindelsen HMS5552 repræsenteret med følgende formel, eller en isotopmærket forbindelse eller et farmaceutisk acceptabelt salt deraf,



- 5.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 4, hvor glucokinase-aktivatoren er i form af en fast dispersion; eller fortrinsvis, glucokinase-aktivatoren er i form af en fast dispersion indeholdende en polymerbærer, hvor polymerbæreren er methacrylsyrecopolymer af type A, som er en anionisk copolymer af methacrylsyre og methylmethacrylat (1:1), fortrinsvis Eudragit, eller mere fortrinsvis Eudragit L100.

15

- 6.** Den farmaceutiske sammensætning ifølge krav 5, hvor vægtforholdet mellem glucokinase-aktivatoren og polymerbæreren er ca. 1:10 til 10:1, fortrinsvis ca. 1:9 til 9:1, ca. 1:4 til 4:1, ca. 3:7 til 7:3, ca. 2:3 til 3:2, ca. 3:4 til 4:3, ca. 4:5 til 5:4 eller ca. 5:6 til 6:5, eller mere fortrinsvis ca. 1:1, ca. 2:3, ca. 3:4, ca. 4:5 eller ca. 5:6.

20

- 7.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 6, hvor SGLT-2-inhibitoren er valgt fra gruppen bestående af canagliflozin, dapagliflozin (eller dapagliflozin-propylenglycolmonohydrat), empagliflozin, ipragliflozin, luseogliflozin og tofogliflozin, og et farmaceutisk acceptabelt salt deraf; eller fortrinsvis, SGLT-2-inhibitoren er valgt fra gruppen bestående af empagliflozin, dapagliflozin (eller dapagliflozin-propylenglycolmonohydrat) og canagliflozin (eller canagliflozin-hemihydrat).

25

8. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 7, hvor glucokinase-aktivatoren er til stede i en dosis i området fra ca. 1 mg til ca. 200 mg, eller fortrinsvis ca. 25 mg til ca. 100 mg, eller mere fortrinsvis, hvor dosen af glucokinase-aktivatoren er ca. 25 mg, ca. 50 mg, ca. 75 mg eller ca. 5 100 mg.

9. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 8, hvor SGLT-2-inhibitoren er til stede i en dosis i området fra ca. 1 mg til 500 mg, eller fortrinsvis ca. 5 mg til ca. 300 mg, eller mere fortrinsvis, hvor dosen 10 af SGLT-2-inhibitoren er ca. 2,5 mg, ca. 5 mg, ca. 10 mg, ca. 12,5 mg, ca. 20 mg, ca. 25 mg, ca. 100 mg, ca. 200 mg eller ca. 300 mg, eller mere fortrinsvis ca. 5 mg, ca. 10 mg, ca. 12,5 mg, ca. 25 mg, ca. 100 mg eller ca. 300 mg; fortrinsvis er SGLT-2-inhibitoren empagliflozin, med en dosis på ca. 0,5 mg til ca. 50 mg, fortrinsvis ca. 1 mg til ca. 25 mg; eller fortrinsvis ca. 5 mg, ca. 10 mg, ca. 15 12,5 mg eller ca. 25 mg; fortrinsvis er SGLT-2-inhibitoren dapagliflozin, med en dosis på ca. 1 mg til ca. 50 mg, fortrinsvis ca. 2,5 mg, ca. 5 mg, ca. 10 mg eller ca. 25 mg; eller fortrinsvis ca. 2,5 mg, ca. 5 mg eller ca. 10 mg; fortrinsvis er SGLT-2-inhibitoren canagliflozin, med en dosis på ca. 50 mg til ca. 500 mg, eller fortrinsvis ca. 100 mg eller ca. 300 mg.

20

10. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 2 til 9, hvor den ene eller flere excipienser er valgt fra gruppen bestående af bindemidler, fyldstoffer, desintegreringsmidler, smøremidler, glidemidler, overfladeaktive midler, befugtningsmidler, antioxidanter, smagsstoffer, 25 sødemidler, farvestoffer og overtræksmidler, eventuelt hvor bindemidlet er valgt fra gruppen bestående af polyvinylpyrrolidon, hydroxypropylcellulose og hydroxypropylmethylcellulose; fyldstoffet er valgt fra gruppen bestående af mikrokrySTALLINSK cellulose, silicificeret mikrokrySTALLINSK cellulose, lactose, calcium-dihydrogenphosphat, mannitol, majsstivelse og prægelatineret stivelse; 30 desintegreringsmidlet er valgt fra gruppen bestående af croscarmellosenatrium, crospovidon og natriumstivelsesglycolat; smøremidlet er valgt fra gruppen bestående af magnesiumstearat og natriumstearylfumarat; og glidemidlet er valgt fra gruppen bestående af kolloidt siliciumdioxid og talkum.

11. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 10, som er i form valgt fra gruppen bestående af en tablet, en kapsel, en pille og en pastil, fortrinsvis en tablet, eller mere fortrinsvis en overtrukket tablet.

5 **12.** Den farmaceutiske sammensætning ifølge krav 11, hvor den overtrukne tablet er en filmovertrukket tablet, hvor filmovertræksmidlet omfatter:
 filmovertrækssubstrat(er), som er eventuelt hydroxypropylcellulose,
 hydroxypropylmethylcellulose eller en blanding deraf;
 blødgøringsmiddel/-midler, som er eventuelt polyvinylalkohol, polyethylenglycol,
 10 propylenglycol, polysorbat eller en blanding deraf;
 farvestof(fer), som er eventuelt jernoxid rød, jernoxid gul eller en blanding deraf;
 opakiseringsmiddel/-midler, som er eventuelt titandioxid, og
 glidemiddel/-midler, eventuelt hvor filmovertræksmidlet er Opadry.

15 **13.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 12, omfattende efter vægt:

- ca. 1-96%, eller fortrinsvis ca. 1-48% af glucokinase-aktivatoren, fortrinsvis HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion
- 20 indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 0.1-80%, eller fortrinsvis ca. 1-50% af SGLT-2-inhibitoren;
- ca. 0-80% fyldstof(fer);
- ca. 1-25% bindemiddel/-midler;
- ca. 0-15% desintegreringsmiddel/-midler;
- 25 - ca. 0,1-10% smøremiddel/-midler;
- ca. 0-3% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler.

14. Den farmaceutiske sammensætning ifølge krav 13, hvor doserne af de aktive
 30 ingredienser er efter vægt:

- ca. 25 mg, ca. 50 mg, ca. 75 mg eller ca. 100 mg af glucokinase-aktivatoren, eller fortrinsvis HMS5552; og
- ca. 5 mg, ca. 10 mg, ca. 12,5 mg eller ca. 25 mg empagliflozin.

- 15.** Den farmaceutiske sammensætning ifølge krav 14, som er
- (i) en tablet af 75 mg HMS5552/5 mg empagliflozin, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
 - ca. 5 mg empagliflozin;
 - ca. 0-70% fyldstof(fer);
 - ca. 2-8% bindemiddel/-midler;
- 10 - ca. 1-5% desintegreringsmiddel/-midler;
- ca. 0,5-3% smøremiddel/-midler;
 - ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- (ii) en tablet af 75 mg HMS5552/12,5 mg empagliflozin, og omfatter
- 15 komponenterne med følgende mængder efter vægt:
- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
 - ca. 12,5 mg empagliflozin;
- 20 - ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
 - ca. 1-5% desintegreringsmiddel/-midler;
 - ca. 0,5-3% smøremiddel/-midler;
 - ca. 0-0,5% glidemiddel/-midler; og
- 25 - ca. 0-5% overtræksmiddel/-midler;
- (iii) en tablet af 75 mg HMS5552/10 mg empagliflozin, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- 30 - ca. 10 mg empagliflozin;
- ca. 0-70% fyldstof(fer);
 - ca. 2-8% bindemiddel/-midler;
 - ca. 1-5% desintegreringsmiddel/-midler;
- 35 - ca. 0,5-3% smøremiddel/-midler;

- ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- (iv) en tablet af 75 mg HMS5552/25 mg empagliflozin, og omfatter komponenterne med følgende mængder efter vægt:
- 5 - ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
 - ca. 25 mg empagliflozin;
 - ca. 0-70% fyldstof(fer);
- 10 - ca. 2-8% bindemiddel/-midler;
- ca. 1-5% desintegreringsmiddel/-midler;
 - ca. 0,5-3% smøremiddel/-midler;
 - ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- 15 (v) en tablet af 50 mg HMS5552/25 mg empagliflozin, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 50 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- 20 - ca. 25 mg empagliflozin;
- ca. 0-70% fyldstof(fer);
 - ca. 2-8% bindemiddel/-midler;
 - ca. 1-5% desintegreringsmiddel/-midler;
 - ca. 0,5-3% smøremiddel/-midler;
- 25 - ca. 0-0,5% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler;
- (vi) en tablet af 50 mg HMS5552/10 mg empagliflozin, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 50 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en
- 30 fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 10 mg empagliflozin;
 - ca. 0-70% fyldstof(fer);
 - ca. 2-8% bindemiddel/-midler;
- 35 - ca. 1-5% desintegreringsmiddel/-midler;

- ca. 0,5-3% smøremiddel/-midler;
 - ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- (vii) en tablet af 100 mg HMS5552/10 mg empagliflozin, og omfatter
- 5 komponenterne med følgende mængder efter vægt:
- ca. 100 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
 - ca. 10 mg empagliflozin;
- 10 - ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
 - ca. 1-5% desintegreringsmiddel/-midler;
 - ca. 0,5-3% smøremiddel/-midler;
 - ca. 0-0,5% glidemiddel/-midler; og
- 15 - ca. 0-5% overtræksmiddel/-midler; eller
- (viii) en tablet af 25 mg HMS5552/25 mg empagliflozin, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 25 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- 20 - ca. 25 mg empagliflozin;
- ca. 0-70% fyldstof(fer);
 - ca. 2-8% bindemiddel/-midler;
 - ca. 1-5% desintegreringsmiddel/-midler;
- 25 - ca. 0,5-3% smøremiddel/-midler;
- ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler.
- 16.** Den farmaceutiske sammensætning ifølge krav 14, omfattende:
- 30 (i) ca. 150 mg af den faste dispersion, ca. 5,00 mg empagliflozin, ca. 88,10 mg mikrokrySTALLINSK cellulose, ca. 7,80 mg hydroxypropylcellulose, ca. 6,50 mg croscarmellosenatrium, ca. 2,60 mg magnesiumstearat og ca. 7,80 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 75 mg HMS5552;

- (ii) ca. 150 mg af den faste dispersion, ca. 12,50 mg empagliflozin, ca. 99,30 mg mikrokrySTALLinsk cellulose, ca. 8,40 mg hydroxypropylcellulose, ca. 7,00 mg croscarmellosenatrium, ca. 2,80 mg magnesiumstearat og ca. 8,40 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og
5 indeholder ca. 75 mg HMS5552;
- (iii) ca. 150 mg af den faste dispersion, ca. 10,00 mg empagliflozin, ca. 84,40 mg mikrokrySTALLinsk cellulose, ca. 7,80 mg hydroxypropylcellulose, ca. 5,20 mg croscarmellosenatrium, ca. 2,60 mg magnesiumstearat og ca. 7,80 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og
10 indeholder ca. 75 mg HMS5552;
- (iv) ca. 150 mg af den faste dispersion, ca. 25,00 mg empagliflozin, ca. 88,20 mg mikrokrySTALLinsk cellulose, ca. 8,40 mg hydroxypropylcellulose, ca. 5,60 mg croscarmellosenatrium, ca. 2,80 mg magnesiumstearat og ca. 8,40 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og
15 indeholder ca. 75 mg HMS5552;
- (v) ca. 200 mg af den faste dispersion, ca. 10,00 mg empagliflozin, ca. 72,00 mg mikrokrySTALLinsk cellulose, ca. 9,00 mg hydroxypropylcellulose, ca. 6,00 mg croscarmellosenatrium, ca. 3,00 mg magnesiumstearat og ca. 9,00 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og
20 indeholder ca. 100 mg HMS5552;
- (vi) ca. 100 mg af den faste dispersion, ca. 10,00 mg empagliflozin, ca. 96,80 mg mikrokrySTALLinsk cellulose, ca. 6,60 mg povidon, ca. 4,40 mg croscarmellose-
natrium, ca. 2,20 mg magnesiumstearat og ca. 6,60 mg Opadry, hvor den faste
dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 50
25 mg HMS5552;
- (vii) ca. 100 mg af den faste dispersion, ca. 25,00 mg empagliflozin, ca.
100,30 mg mikrokrySTALLinsk cellulose, ca. 7,20 mg hydroxypropylcellulose, ca.
5,10 mg croscarmellosenatrium, ca. 2,40 mg magnesiumstearat, og ca. 7,20 mg
Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit
30 L100, og indeholder ca. 50 mg HMS5552; eller
- (viii) ca. 50 mg af den faste dispersion, ca. 25,00 mg empagliflozin, ca.
309,00 mg mikrokrySTALLinsk cellulose, ca. 8,00 mg croscarmellosenatrium, ca.
4,00 mg mikroniseret silicagel, ca. 4,00 mg magnesiumstearat og ca. 12,00 mg
Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit
35 L100, og indeholder ca. 25 mg HMS5552.

17. Den farmaceutiske sammensætning ifølge krav 13, hvor doserne af de aktive ingredienser er efter vægt:

- ca. 25 mg, ca. 50 mg, ca. 75 mg eller ca. 100 mg af glucokinase-aktivatoren, eller
 5 fortrinsvis HMS5552; og
 ca. 2.5 mg, ca. 5 mg, eller ca. 10 mg dapagliflozin eller en tilsvarende mængde af dapagliflozin-propylenglycolmonohydrat.

18. Den farmaceutiske sammensætning ifølge krav 17, som er

10 (i) en tablet af 75 mg HMS5552/5 mg dapagliflozin eller en tilsvarende mængde af dapagliflozin-propylenglycolmonohydrat, og omfatter komponenterne med følgende mængder efter vægt:

- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en
 15 fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 5 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylenglycolmonohydrat;
- ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
- 20 - ca. 1-8% desintegreringsmiddel/-midler;
- ca. 0,5-3% smøremiddel/-midler;
- ca. 0-0,5% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler;

(ii) en tablet af 75 mg HMS5552/10 mg dapagliflozin eller en tilsvarende mængde
 25 dapagliflozin-propylenglycolmonohydrat, og omfatter komponenterne med følgende mængder efter vægt:

- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- 30 - ca. 10 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylenglycolmonohydrat;
- ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
- ca. 1-8% desintegreringsmiddel/-midler;
- 35 - ca. 0,5-3% smøremiddel/-midler;

- ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- (iii) en tablet af 50 mg HMS5552/5 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylenglycolmonohydrat, og omfatter komponenterne med
- 5 følgende mængder efter vægt:
- ca. 50 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
 - ca. 5 mg dapagliflozin eller en tilsvarende mængde af dapagliflozin-propylen-
- 10 glycolmonohydrat;
- ca. 0-70% fyldstof(fer);
 - ca. 2-8% bindemiddel/-midler;
 - ca. 1-8% desintegreringsmiddel/-midler;
 - ca. 0,5-3% smøremiddel/-midler;
- 15 - ca. 0-0,5% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler;
- (iv) en tablet af 50 mg HMS5552/10 mg dapagliflozin eller en tilsvarende mængde af dapagliflozin-propylenglycolmonohydrat, og omfatter komponenterne med følgende mængder efter vægt:
- 20 - ca. 50 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 10 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylen-glycolmonohydrat;
- 25 - ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
 - ca. 1-8% desintegreringsmiddel/-midler;
 - ca. 0,5-3% smøremiddel/-midler;
 - ca. 0-0,5% glidemiddel/-midler; og
- 30 - ca. 0-5% overtræksmiddel/-midler;
- (v) en tablet af 100 mg HMS5552/5 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylenglycolmonohydrat, og omfatter komponenterne med følgende mængder efter vægt:

- ca. 100 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 HMS5552 og Eudragit L100;
- ca. 5 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylen-glycolmonohydrat;
- ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
- ca. 1-8% desintegreringsmiddel/-midler;
- ca. 0,5-3% smøremiddel/-midler;
- 10 - ca. 0-0,5% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler; eller
- (vi) en tablet af 25 mg HMS5552/10 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylenglycolmonohydrat, og omfatter komponenterne med følgende mængder efter vægt:
- 15 - ca. 25 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 10 mg dapagliflozin eller en tilsvarende mængde dapagliflozin-propylen-glycolmonohydrat;
- 20 - ca. 0-70% fyldstof(fer);
- ca. 2-8% bindemiddel/-midler;
- ca. 1-8% desintegreringsmiddel/-midler;
- ca. 0,5-3% smøremiddel/-midler;
- ca. 0-0,5% glidemiddel/-midler; og
- 25 - ca. 0-5% overtræksmiddel/-midler.

19. Den farmaceutiske sammensætning ifølge krav 17, omfattende

- (i) ca. 150 mg af den faste dispersion, ca. 6,15 mg dapagliflozin-propylenglycolmonohydrat, ca. 88,25 mg mikrokrySTALLinsk cellulose, ca. 7,80 mg
- 30 hydroxypropylcellulose, ca. 5,20 mg croscarmellosenatrium, ca. 2,60 mg magnesiumstearat og 7,80 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 75 mg HMS5552;
- (ii) ca. 100 mg af den faste dispersion, ca. 6,15 mg dapagliflozin-propylen-glycolmonohydrat, ca. 100,65 mg mikrokrySTALLinsk cellulose, ca. 6,60 mg
- 35 hydroxypropylcellulose, ca. 4,40 mg croscarmellosenatrium, ca. 2,20 mg

- magnesiumstearat og 6,60 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 50 mg HMS5552;
- (iii) ca. 150 mg af den faste dispersion, ca. 12,30 mg dapagliflozin-propylen-glycolmonohydrat, ca. 91,50 mg mikrokrystallinsk cellulose, ca. 8,10 mg
- 5 hydroxypropylcellulose, ca. 5,40 mg croscarmellosenatrium, ca. 2,70 mg magnesiumstearat og 8,10 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 75 mg HMS5552;
- (iv) ca. 200 mg af den faste dispersion, ca. 6,15 mg dapagliflozin-propylenglycolmonohydrat, ca. 75,55 mg mikrokrystallinsk cellulose, ca. 9,00 mg
- 10 hydroxypropylcellulose, ca. 6,30 mg croscarmellosenatrium, ca. 3,00 mg magnesiumstearat og 9,00 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 100 mg HMS5552;
- (v) ca. 100 mg af den faste dispersion, ca. 12,30 mg dapagliflozin-propylenglycolmonohydrat, ca. 113,06 mg mikrokrystallinsk cellulose, ca. 7,20 mg
- 15 hydroxypropylcellulose, ca. 5,04 mg croscarmellosenatrium, ca. 2,40 mg magnesiumstearat og 7,20 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 50 mg HMS5552; eller
- (vi) ca. 50 mg af den faste dispersion, ca. 12,30 mg dapagliflozin-propylenglycolmonohydrat, ca. 313,70 mg mikrokrystallinsk cellulose, ca. 8,00 mg
- 20 hydroxypropylcellulose, ca. 8,00 mg croscarmellosenatrium, ca. 4,00 mg mikroniseret silicagel, ca. 4,00 mg magnesiumstearat og 12,00 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 25 mg HMS5552.
- 25 **20.** Den farmaceutiske sammensætning ifølge krav 13, hvor doserne af de aktive ingredienser er efter vægt:
ca. 25 mg, ca. 50 mg, ca. 75 mg eller ca. 100 mg af glucokinase-aktivatoren, eller fortrinsvis HMS5552; og
ca. 100 mg eller ca. 300 mg canagliflozin eller en tilsvarende mængde
- 30 canagliflozin-hemihydrat.
- 21.** Den farmaceutiske sammensætning ifølge krav 20, som er
- (i) en tablet af 75 mg HMS5552/100 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat, og omfatter komponenterne med følgende mængder
- 35 efter vægt:

- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 100 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat;
- 5 - ca. 0-70% fyldstof(fer), såsom mikrokrySTALLINSK cellulose;
- ca. 2-8% bindemiddel/-midler, såsom hydroxypropylcellulose;
- ca. 1-5% desintegreringsmiddel/-midler, såsom croscarmellosenatrium;
- ca. 0,5-3% smøremiddel/-midler, såsom natriumstearylfumarat eller magnesiumstearat;
- 10 - ca. 0-0,5% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler;
- (ii) en tablet af 50 mg HMS5552/100 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat, og omfatter komponenterne med følgende mængder efter vægt:
- 15 - ca. 50 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 100 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat;
- ca. 0-70% fyldstof(fer), såsom mikrokrySTALLINSK cellulose;
- 20 - ca. 2-8% bindemiddel/-midler, såsom hydroxypropylcellulose;
- ca. 1-5% desintegreringsmiddel/-midler, såsom croscarmellosenatrium;
- ca. 0,5-3% smøremiddel/-midler, såsom natriumstearylfumarat eller magnesiumstearat;
- ca. 0-0,5% glidemiddel/-midler; og
- 25 - ca. 0-5% overtræksmiddel/-midler;
- (iii) en tablet af 100 mg HMS5552/100 mg canagliflozin-hemihydrat eller en tilsvarende mængde canagliflozin-hemihydrat, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 100 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 100 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat;
- ca. 0-70% fyldstof(fer), såsom mikrokrySTALLINSK cellulose;
- ca. 2-8% bindemiddel/-midler, såsom hydroxypropylcellulose;
- 35 - ca. 1-5% desintegreringsmiddel/-midler, såsom croscarmellosenatrium;

- ca. 0,5-3% smøremiddel/-midler, såsom natriumstearylfumarat eller magnesiumstearat;
 - ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- 5 (iv) en tablet af 75 mg HMS5552/300 mg canagliflozin-hemihydrat eller en tilsvarende mængde canagliflozin-hemihydrat, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 75 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en
- 10 fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 300 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat;
 - ca. 0-70% fyldstof(fer), såsom mikrokrySTALLINSK cellulose;
 - ca. 2-8% bindemiddel/-midler, såsom hydroxypropylcellulose;
 - ca. 1-5% desintegreringsmiddel/-midler, såsom croscarmellosenatrium;
- 15 - ca. 0,5-3% smøremiddel/-midler, såsom natriumstearylfumarat eller magnesiumstearat;
- ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler;
- (v) en tablet af 50 mg HMS5552/300 mg canagliflozin eller en tilsvarende
- 20 mængde canagliflozin-hemihydrat, og omfatter komponenterne med følgende mængder efter vægt:
- ca. 50 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- 25 - ca. 300 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat;
- ca. 0-70% fyldstof(fer), såsom mikrokrySTALLINSK cellulose;
 - ca. 2-8% bindemiddel/-midler, såsom hydroxypropylcellulose;
 - ca. 1-5% desintegreringsmiddel/-midler, såsom croscarmellosenatrium;
 - ca. 0,5-3% smøremiddel/-midler, såsom natriumstearylfumarat eller
- 30 magnesiumstearat;
- ca. 0-0,5% glidemiddel/-midler; og
 - ca. 0-5% overtræksmiddel/-midler; eller
- (vi) en tablet af 25 mg HMS5552/300 mg canagliflozin eller en tilsvarende
- 35 mængde canagliflozin-hemihydrat, og omfatter komponenterne med følgende mængder efter vægt:

- ca. 25 mg HMS5552, fortrinsvis en fast dispersion af HMS5552, fortrinsvis en fast dispersion indeholdende HMS5552 og en polymerbærer, eller fortrinsvis en fast dispersion indeholdende ca. 1:1 af HMS5552 og Eudragit L100;
- ca. 300 mg canagliflozin eller en tilsvarende mængde canagliflozin-hemihydrat;
- 5 - ca. 0-70% fyldstof(fer), såsom mikrokrySTALLinsk cellulose;
- ca. 2-8% bindemiddel/-midler, såsom hydroxypropylcellulose;
- ca. 1-5% desintegreringsmiddel/-midler, såsom croscarmellosenatrium;
- ca. 0,5-3% smøremiddel/-midler, såsom natriumstearylfumarat eller magnesiumstearat;
- 10 - ca. 0-0,5% glidemiddel/-midler; og
- ca. 0-5% overtræksmiddel/-midler.

22. Den farmaceutiske sammensætning ifølge krav 20, omfattende

- (i) ca. 150 mg af den faste dispersion, ca. 101,93 mg canagliflozin-hemihydrat,
- 15 ca. 294,07 mg mikrokrySTALLinsk cellulose, ca. 18,00 mg hydroxypropylcellulose, ca. 30,00 mg croscarmellosenatrium, ca. 6,00 mg magnesiumstearat og ca. 18,00 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 75 mg HMS5552;
- (ii) ca. 100 mg af den faste dispersion, ca. 101,93 mg canagliflozin-hemihydrat,
- 20 ca. 344,07 mg mikrokrySTALLinsk cellulose, ca. 18,00 mg hydroxypropylcellulose, ca. 30,00 mg croscarmellosenatrium, ca. 6,00 mg magnesiumstearat og ca. 18,00 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og EudragitL100, og indeholder ca. 50 mg HMS5552;
- (iii) ca. 200 mg af den faste dispersion, ca. 101,93 mg canagliflozin-hemihydrat,
- 25 ca. 244,07 mg mikrokrySTALLinsk cellulose, ca. 18,00 mg hydroxypropylcellulose, ca. 30,0 mg croscarmellosenatrium, ca. 6,00 mg magnesiumstearat og ca. 18,00 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 100 mg HMS5552;
- (iv) ca. 150 mg af den faste dispersion, ca. 305,78 mg canagliflozin-hemihydrat,
- 30 ca. 226,72 mg mikrokrySTALLinsk cellulose, ca. 22,50 mg hydroxypropylcellulose, ca. 37,50 mg croscarmellosenatrium, ca. 7,50 mg magnesiumstearat og ca. 22,50 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og EudragitL100, og indeholder ca. 75 mg HMS5552;
- (v) ca. 100 mg af den faste dispersion, ca. 305,78 mg canagliflozin-hemihydrat,
- 35 ca. 276,72 mg mikrokrySTALLinsk cellulose, ca. 22,50 mg hydroxypropylcellulose,

ca. 37,50 mg croscarmellosenatrium, ca. 7,50 mg magnesiumstearat og ca. 22,50 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og Eudragit L100, og indeholder ca. 50 mg HMS5552; eller
(vi) ca. 50 mg af den faste dispersion, ca. 305,78 mg canagliflozin-hemihydrat,
5 ca. 326,72 mg mikrokrySTALLINSK cellulose, ca. 22,50 mg hydroxypropylcellulose, ca. 37,50 mg croscarmellosenatrium, ca. 7,50 mg magnesiumstearat og ca. 22,50 mg Opadry, hvor den faste dispersion indeholder ca. 1:1 af HMS5552 og EudragitL100, og indeholder ca. 25 mg HMS5552.

10 **23.** Fremgangsmåde til fremstilling af den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1 til 22, omfattende inkorporering af de aktive ingredienser i en eller flere excipienser til granulering, fortrinsvis yderligere
fyldning af den opnåede granulatløsning i et hætteglas, en pose eller en kapsel,
15 yderligere overtræk af den opnåede tablet; eventuelt hvor fremstillingen udføres ved vådgranulering (høj forskydning og/eller fluidiseret leje), eller ved tørbehandling (direkte komprimering eller tørgranulering).

24. Farmaceutisk sammensætning som defineret i et hvilket som helst af kravene
20 1 til 22, til anvendelse i en fremgangsmåde til at forebygge, bremse udviklingen af, forsinke eller behandle en eller flere stofskiftesygdomme valgt fra gruppen bestående af: type I diabetes, type II diabetes, nedsat glucoSetolerance, nedsat fastende blodsukker, hyperglykæmi, postprandial hyperglykæmi, overvægt, fedme, hypertension, insulinresistens, diabetisk nefropati, nedsat nyrefunktion
25 og/eller stofskiftesyndrom; eller at forbedre blodsukkerkontrol og/eller reducere fastende plasmaglucoSe, postprandial plasmaglucoSe og/eller glycosyleret hæmoglobin HbA1c; eller at forebygge, bremse, forsinke eller vende komplikationer af diabetes mellitus.

DRAWINGS

Drawing




FIG. 1




FIG. 2