




(86) **Date de dépôt PCT/PCT Filing Date:** 2007/04/12
(87) **Date publication PCT/PCT Publication Date:** 2007/10/25
(45) **Date de délivrance/Issue Date:** 2016/08/23
(85) **Entrée phase nationale/National Entry:** 2008/10/09
(86) **N° demande PCT/PCT Application No.:** JP 2007/058049
(87) **N° publication PCT/PCT Publication No.:** 2007/119774
(30) **Priorité/Priority:** 2006/04/13 (JP2006-110467)

(51) **Cl.Int./Int.Cl. C12P 21/00** (2006.01),
C07K 14/47 (2006.01), **C12N 15/12** (2006.01),
C12N 5/10 (2006.01), **C12P 21/08** (2006.01)
(72) **Inventeurs/Inventors:**
TABUCHI, HISAHIRO, JP;
SUGIYAMA, TOMOYA, JP;
TANAKA, SAEKO, JP;
TAINAKA, SATOSHI, JP
(73) **Propriétaire/Owner:**
CHUGAI SEIYAKU KABUSHIKI KAISHA, JP
(74) **Agent:** NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) **Titre : GENE TRANSPORTEUR DE LA TAURINE**
(54) **Title: TAURINE TRANSPORTER GENE**



(57) **Abrégé/Abstract:**

The present invention provides a method capable of producing a natural or recombinant protein at low cost.
The present invention relates to a method of producing a polypeptide, comprising culturing a cell which strongly expresses a taurine transporter and has a transferred DNA encoding a desired polypeptide and thereby allowing the cell to produce the polypeptide. Hamster taurine transporter, a DNA encoding the same, a recombinant vector and a transformed cell are also provided.

ABSTRACT

The present invention provides a method capable of producing a natural or recombinant protein at low cost.

The present invention relates to a method of producing a polypeptide, comprising culturing a cell which strongly expresses a taurine transporter and has a transferred DNA encoding a desired polypeptide and thereby allowing the cell to produce the polypeptide. Hamster taurine transporter, a DNA encoding the same, a recombinant vector and a transformed cell are also provided.

DESCRIPTION

TAURINE TRANSPORTER GENE

TECHNICAL FIELD

5 The present invention relates to a hamster taurine transporter and a gene encoding the hamster taurine transporter, as well as a method of producing a polypeptide using a cell which strongly expresses a taurine transporter.

BACKGROUND ART

10 When proteins useful as pharmaceuticals are produced with the recombinant DNA technique, use of animal cells enables complicated post-translational modification and folding which prokaryotic cells can not perform. Therefore, animal cells are frequently used as host cells for producing recombinant proteins.

15 Recently, a large number of biopharmaceuticals, such as antibodies and physiologically active proteins, have been developed. Techniques that permit efficient production of recombinant proteins by animal cells lead to cost reduction of biopharmaceuticals and promise their stable supply to patients.

 Under these circumstances, a method of protein production with higher production efficiency is desired.

20 Taurine is a type of amino acid present in high concentrations in fishes, shell fishes and mollusks and is an important nutrient for the growth of mammals. Although taurine is not used in protein synthesis, it has many functions such as normalization of hypercholesterolemia, lowering of blood pressure, detoxication effect, maintenance of immune function, stabilization of biological membranes, regulation of neural excitability, antioxidation, etc. It is known that taurine contributes to osmoregulation and stabilization of cell membrane in cultured cells (Non-Patent Document 1). However, addition of taurine to the medium of astrocyte primary culture where taurine transporter was functioning did not increase taurine uptake into cells (Non-Patent Document 2). Thus, addition of taurine to the medium alone was insufficient.

25 On the other hand, it is totally unknown whether or not uptake of taurine and other amino acids into cultured cells via taurine transporter contributes to improvement of the production of a desired recombinant protein in the cultured cells.

 Several taurine transporters (human: Non-Patent Document 3; mouse: Non-Patent Document 4; and rat: Non-Patent Document 5) and their involvement in the uptake of taurine

and other amino acids (e.g., β -alanine) into cells are known (Non-Patent Document 6). However, with respect to hamster taurine transporter, even its existence has not been known yet.

[Non-Patent Document 1]

5 Ian Henry Lambert, Neurochemical Research (2004) 29(1), 27-63

[Non-Patent Document 2]

Journal of Neurochemistry (2000), 75(3), 919-924

[Non-Patent Document 3]

Uchida, S. et al., Proc. Natl. Acad. Sci. U.S.A. (1992) 89 (17), 8230-8234

10 [Non-Patent Document 4]

Liu, Q.R. et al., Proc. Natl. Acad. Sci. U.S.A. (1992) 89 (24), 12145-12149

[Non-Patent Document 5]

Smith, K.E. et al., Mol. Pharmacol. (1992) 42 (4), 563-569

[Non-Patent Document 6]

15 Ryo Shioda et al., Investigative Ophthalmology & Visual Science (2002) 43 (9), 2916

DISCLOSURE OF THE INVENTION

PROBLEM FOR SOLUTION BY THE INVENTION

20 It is an object of the present invention to provide a method which is capable of producing a natural or recombinant protein at low cost.

MEANS TO SOLVE THE PROBLEM

25 As a result of extensive and intensive researches toward the solution of the above problem, the present inventors have found that it is possible to increase the yield of a desired polypeptide by using a cell that strongly expresses a taurine transporter. Thus, the present invention has been achieved.

The present invention may be summarized as follows.

- (1) A method of producing a polypeptide, comprising culturing a cell which strongly expresses a taurine transporter and has a transferred DNA encoding a desired polypeptide and thereby allowing the cell to produce said polypeptide.
- 30 (2) The method of (1) above, wherein the cell which strongly expresses a taurine transporter is a cell into which a DNA encoding the taurine transporter has been transferred.
- (3) The method of (2) above, wherein the cell is Chinese hamster ovary cells.
- (4) The method of any one of (1) to (3) above, wherein the desired polypeptide is an antibody.
- 35 (5) The method of any one of (2) to (4) above, wherein the DNA encoding the taurine

transporter is any one of the following (a) to (e):

(a) a DNA encoding a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8;

5 (b) a DNA encoding a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;

(c) a DNA encoding a polypeptide having 70% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 and yet having
10 taurine transporter activity;

(d) a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7;

(e) a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.

15 (6) The method of any one of (1) to (5) above, comprising culturing the cells in a medium with a taurine concentration of 0 to 100 g/L.

(7) A method of preparing a pharmaceutical containing a polypeptide prepared by the method of any one of (1) to (6) above.

(8) A DNA encoding a taurine transporter, which is any one of the following (a) to (e),
20 provided that DNA having the nucleotide sequence as shown in SEQ ID NO: 3, DNA having the nucleotide sequence as shown in SEQ ID NO: 5, and DNA having the nucleotide sequence as shown in SEQ ID NO: 7 are excluded:

(a) a DNA encoding a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2;

25 (b) a DNA encoding a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;

(c) a DNA encoding a polypeptide having 97% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2 and yet having taurine
30 transporter activity;

(d) a DNA having the nucleotide sequence as shown in SEQ ID NO: 1;

(e) a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.

35 (9) A polypeptide which is any one of the following (A) to (E), provided that a polypeptide

having the amino acid sequence as shown in SEQ ID NO: 4, a polypeptide having the amino acid sequence as shown in SEQ ID NO: 6, and a polypeptide having the amino acid sequence as shown in SEQ ID NO: 8 are excluded:

- (A) a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2;
- 5 (B) a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;
- (C) a polypeptide having 97% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2 and yet having taurine transporter activity;
- 10 (D) a polypeptide encoded by a DNA having the nucleotide sequence as shown in SEQ ID NO: 1;
- (E) a polypeptide encoded by a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.
- 15 (10) A recombinant vector comprising the DNA of (8) above.
- (11) A cell into which the DNA of (8) above has been transferred.
- (12) A cell which strongly expresses a taurine transporter and yet has a transferred DNA encoding a desired polypeptide.
- (13) A cell into which a DNA encoding a taurine transporter has been transferred.

20

EFFECT OF THE INVENTION

According to the present invention, it has become possible to produce a desired polypeptide at low cost.

25 The present specification encompasses the contents disclosed in the specification and/or the drawings of Japanese Patent Application No. 2006-110467 based on which the present patent application claims priority.

BRIEF DESCRIPTION OF THE DRAWINGS

30 Fig. 1 shows the nucleotide sequence of a newly cloned, CHO cell-derived hamster taurine transporter gene and the amino acid sequence deduced therefrom.

Fig. 2 is a taurine transporter membrane topology which was created based on the transmembrane regions and orientations predicted by TMpred program from the amino acid sequence of a newly cloned, CHO cell-derived hamster TauT with reference to FIG. 5 of Shinichi Uchida et al., Proc. Natl. Acad. Sci. USA Vol. 89, pp. 8230-8234, September 1992.

35 Mark © indicates hamster TauT specific amino acid residues. A large number of amino

acid residues different from those in human TauT are present in the 2nd loop (EX: extra-cell membrane region), the 12th transmembrane region (TM) and the C-terminal (IC: intracellular region).

Fig. 3 shows a plasmid which was used for expressing hamster TauT (622 amino acids).

Fig. 4 shows viable cell density plots on day 7 of 50 ml shaker flask batch culture (n=7). The viable cell density in pHyg/TauT-transferred cell was superior to that in pHyg-transferred cell ($P<0.05$).

Fig. 5 shows lactate yield plots on day 7 of 50 ml shaker flask batch culture (n=7). pHyg/TauT-transferred cell produced less lactate, and was superior to pHyg-transferred cell ($P<0.05$).

Fig. 6 shows anti-glypican-3 antibody yield plots on day 7 of 50 ml shaker flask batch culture (n=7). Four out of the 7 strains of pHyg/TauT-transferred cell showed antibody yields higher than the highest yield in pHyg-transferred cell.

Fig. 7 shows anti-glypican-3 antibody yield plots on day 7 of 50 ml shaker flask fed-batch culture (n=7). The antibody yield in pHyg/TauT-transferred cell was superior to that in pHyg-transferred cell ($P<0.01$).

Fig. 8 is a graph showing the survival ratio of a pHyg/TauT-transferred cell T10 (which showed high growth ability during the expansion process in static culture) in 1 L jar fed-batch culture. The survival ratio of T10 was 80% or more even on day 32 of the culture. On the other hand, the survival ratio of the parent strain became less than 80% on day 19.

Fig. 9 is a graph showing the antibody yield of a pHyg/TauT-transferred cell T10 (which showed high growth ability during the expansion process in static culture) in 1 L jar fed-batch culture. The anti-glypican-3 antibody yield of T10 was 2.9 g/L on day 35 of the culture.

Fig. 10 shows the results of flow cytometric analysis indicating that TauT-transferred T10 cell is expressing TauT molecules on its cell membrane. As a primary antibody, rabbit anti-rat taurine transporter antibody (Alpha Diagnostics, US) was used (antibody $\pm$). As a secondary antibody, donkey anti-rabbit IgG antibody-Pycoerythrin-conjugate (Abcam, UK) was used for labelling.

Fig. 11 is a graph showing intracellular ammonia contents (concentration ratios) in 1 L jar fed-batch culture. The ammonia inhibition in pHyg/TauT-transferred strains was remarkable compared to the parent strain.

Fig. 12 is a graph showing that taurine is taken into cells depending on the taurine concentration in the medium. No difference was observed in taurine uptake between

pHyg/TauT-transferred strains and the parent strain.

Fig. 13 is a graph showing the consumption of glutamine in the medium. Compared to the parent strain, pHyg/TauT-transferred strains showed a remarkably high glutamine consumption/cell without depending on the taurine concentration in the medium.

5 Fig. 14 is a graph showing that the anti-glypican-3 antibody yields of pHyg/TauT-transferred strains are almost equal without depending on the initial taurine concentration in the medium.

BEST MODE FOR CARRYING OUT THE INVENTION

10 Hereinbelow, embodiments of the present invention will be described in more detail.

The present invention provides a method of producing a polypeptide, comprising culturing a cell which strongly expresses a taurine transporter and has a transferred DNA encoding a desired polypeptide and thereby allowing the cell to produce the polypeptide.

15 In the method of the present invention, the cell may be either a natural cell capable of producing the desired polypeptide or a transformed cell into which a DNA encoding the desired polypeptide has been transferred. Preferably, a transformed cell into which a DNA encoding the desired polypeptide has been transferred is used.

In the method of the present invention, the desired polypeptide is not particularly
20 limited. The polypeptide may be any polypeptide such as an antibody (e.g., anti-IL-6 receptor antibody, anti-glypican-3 antibody, anti-CD3 antibody, anti-CD20 antibody, anti-GPIIb/IIIa antibody, anti-TNF antibody, anti-CD25 antibody, anti-EGFR antibody, anti-Her2/neu antibody, anti-RSV antibody, anti-CD33 antibody, anti-CD52 antibody, anti-IgE antibody, anti-CD11a antibody, anti-VEGF antibody, anti-VLA4 antibody, and the
25 like) or a physiologically active protein (e.g., granulocyte-colony stimulating factor (G-CSF), granulocyte macrophage-colony stimulating factor (GM-CSF), erythropoietin, interferon, interleukin such as IL-1 or IL-6, t-PA, urokinase, serum albumin, blood coagulation factor, PTH, and the like). An antibody is particularly preferred, and may be any antibody such as
30 a natural antibody, a low molecular sized antibody (e.g., Fab, scFv, sc(Fv)₂), a chimeric antibody, a humanized antibody, etc.

The present inventors have found that, by using a cell which strongly expresses a taurine transporter, intracellular uptake of not only taurine and β -alanine but also glutamine is specifically promoted via the strongly expressed taurine transporter.

It is known that taurine transporter is a membrane protein having the function of
35 taking up amino acids (such as taurine and β -alanine) into cells. However, it is not known

that cells begin to take up glutamine specifically when they are allowed to express a taurine transporter strongly. Since it is known that glutamine is involved in antibody production in hybridomas (Yeon-Ho Jeong et al, Enzyme and Microbial Technology (1995) 17, 47-55), the effect of enhancement of protein (e.g., antibody) production by a cell which has been
 5 allowed to strongly express a taurine transporter may be caused by the taurine transporter-mediated specific uptake of glutamine into cells.

A cell which strongly expresses a taurine transporter is not particularly limited as long as the cell has an increased expression level of a taurine transporter compared to a corresponding natural cell. The natural cell is not particularly limited. A cell which is
 10 used as a host in the production of a recombinant protein (e.g., CHO cells) may be used.

As a cell which strongly expresses a taurine transporter, a cell into which a taurine transporter gene has been artificially transferred may be given. A cell into which a taurine transporter gene has been artificially transferred can be prepared by methods known to those skilled in the art. For example, such a cell may be prepared by incorporating a taurine
 15 transporter gene into a vector and transforming the vector into a cell.

As a taurine transporter to be strongly expressed in a cell, a taurine transporter derived from any organism may be used. Specifically, a taurine transporter derived from human or a rodent (such as mouse, rat or hamster) may be used. Preferably, a taurine transporter derived from human, a rodent or the same species as the host cell may be used.
 20 For example, when the cell which is allowed to strongly express a taurine transporter is Chinese hamster ovary cells (CHO cells), the taurine transporter is preferably derived from human or hamster.

Further, as a taurine transporter gene to be strongly expressed in a cell, any one of the following DNAs (a) to (e) encoding a taurine transporter may be used.

- 25 (a) a DNA encoding a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8;
- (b) a DNA encoding a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;
- 30 (c) a DNA encoding a polypeptide having 70% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 and yet having taurine transporter activity;
- (d) a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7;
- (e) a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide
 35 sequence as shown in SEQ ID NO: 1, 3, 5 or 7 under stringent conditions and yet encodes a

polypeptide having taurine transporter activity.

The cell which strongly expresses a taurine transporter may be any cell. Preferably, CHO cells are used; CHO-dhfr- cells are particularly preferred.

Production of a desired polypeptide may be performed by transferring a gene
5 encoding the desired polypeptide into a cell which strongly expresses a taurine transporter and culturing the resultant cell.

When a desired polypeptide is produced using a cell into which a taurine transporter gene has been artificially transferred, the order of the transfer of a taurine transporter gene and the transfer of a gene encoding a desired polypeptide is not particularly
10 limited. A gene encoding a desired polypeptide may be transferred after the transfer of a taurine transporter gene. Alternatively, a taurine transporter gene may be transferred after the transfer of a gene encoding a desired polypeptide. It is also possible to transfer a taurine transporter gene and a gene encoding a desired polypeptide simultaneously.

A taurine transporter gene and a gene encoding a desired polypeptide may be
15 transferred simultaneously in a single vector. Alternatively, they may be transferred separately using a plurality of vectors.

For culturing the cell which strongly expresses a taurine transporter, media used in conventional cell culture (preferably, animal cell culture) may be used. These media usually contain amino acids, vitamins, lipid factors, energy sources, osmotic regulators, iron
20 sources and pH regulators. The contents of these components are usually as follows: amino acids 0.05-1500 mg/L, vitamins 0.001-10 mg/L, lipid factors 0-200 mg/L, energy sources 1-20 g/L, osmotic regulators 0.1-10000 mg/L, iron sources 0.1-500 mg/L, pH regulators 1-10000 mg/L, trace metal elements 0.00001-200 mg/L, surfactants 0-5000 mg/L, growth cofactors 0.05-10000 µg/L and nucleosides 0.001-50 mg/L. However, the contents are not
25 limited to these ranges and may be appropriately selected depending on the type of the cell to be cultured, the type of the desired polypeptide, and so on.

In addition to these components, trace metal elements, surfactants, growth cofactors, nucleosides, and the like may be added.

Specific examples of such components include amino acids, such as L-alanine,
30 L-arginine, L-asparagine, L-aspartic acid, L-cysteine, L-cystine, L-glutamine, L-glutamic acid, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-ornithine, L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine, and L-valine, preferably, L-alanine, L-arginine, L-asparagine, L-aspartic acid, L-cystine, L-glutamine, L-glutamic acid, glycine, L-histidine, L-isoleucine, L-leucine, L-lysine, L-methionine,
35 L-phenylalanine, L-proline, L-serine, L-threonine, L-tryptophan, L-tyrosine and L-valine;

vitamins, such as i-inositol, biotin, folic acid, lipoic acid, nicotinamide, nicotinic acid, p-aminobenzoic acid, calcium pantothenate, pyridoxal hydrochloride, pyridoxine hydrochloride, riboflavin, thiamine hydrochloride, vitamin B₁₂ and ascorbic acid, preferably, biotin, folic acid, lipoic acid, nicotinamide, calcium pantothenate, pyridoxal hydrochloride, riboflavin, thiamine hydrochloride, vitamin B₁₂ and ascorbic acid; lipid factors, such as choline chloride, choline tartrate, linoleic acid, oleic acid and cholesterol, preferably, choline chloride; energy sources, such as glucose, galactose, mannose, and fructose, preferably, glucose; osmotic regulators, such as sodium chloride, potassium chloride, and potassium nitrate, preferably, sodium chloride; iron sources, such as iron EDTA, ferric citrate, ferrous chloride, ferric chloride, ferrous sulfate, ferric sulfate, and ferric nitrate, preferably, ferric chloride, iron EDTA, and ferric citrate; and pH regulators, such as sodium hydrogencarbonate, calcium chloride, sodium dihydrogen-phosphate, HEPES and MOPS, preferably, sodium hydrogencarbonate. Culture media containing any of these components may be given as examples.

Besides the above components, there may be added trace metal elements, such as copper sulfate, manganese sulfate, zinc sulfate, magnesium sulfate, nickel chloride, tin chloride, magnesium chloride and sodium subsilicate, preferably, copper sulfate, zinc sulfate and magnesium sulfate; surfactants, such as Tween 80 and Pluronic F68; growth cofactors, such as recombinant insulin, recombinant IGF-1, recombinant EGF, recombinant FGF, recombinant PDGF, recombinant TGF- α , ethanolamine hydrochloride, sodium selenite, retinoic acid and putrescine dihydrochloride, preferably, sodium selenite, ethanolamine hydrochloride, recombinant IGF-1 and putrescine dihydrochloride; and nucleosides, such as deoxyadenosine, deoxycytidine, deoxyguanosine, adenosine, cytidine, guanosine and uridine. In preferable examples of above media, antibiotics, such as streptomycin, penicillin-G potassium and gentamicin, and pH-indicators, such as Phenol Red, may be contained.

The pH of the medium varies depending on the cell to be cultured. Generally, pH 6.8-7.6 is appropriate. In many cases, pH 7.0-7.4 is appropriate.

It is also possible to use a commercial medium for animal cell culture, e.g., D-MEM (Dulbecco's Modified Eagle Medium), D-MEM/F-12 1:1 Mixture (Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12), RPMI1640, CHO-S-SFMII (Invitrogen), CHO-SF (Sigma-Aldrich), EX-CELL 301 (JRH Biosciences), CD-CHO (Invitrogen), IS CHO-V (Irvine Scientific), PF-ACF-CHO (Sigma-Aldrich) or the like.

Alternatively, the medium may be a serum-free medium.

When the cell which strongly expresses a taurine transporter is CHO cells, CHO cells may be cultured by methods known to those skilled in the art. For example, CHO

cells may be cultured usually in an atmosphere with a CO₂ concentration in the gas phase of 0 to 40%, preferably 2 to 10%, at 30 to 39°C, preferably about 37°C.

As is clear from the Examples described later, production of waste products (such as lactate) which turn to be cell growth inhibitory substances can be inhibited in a cell strongly expressing a taurine transporter. As a result, the cell shows the effect of maintaining a high survival ratio. The cell of the present invention is capable of culturing for three months or a still longer period.

Further, when a desired polypeptide, such as an antibody, is produced in cultured cells, the cells come into a highly concentrated state (about 1×10^7 cells/ml) at the late-stage of culture, and the influence of waste products such as lactate becomes extremely high. When a desired polypeptide is produced using the cell of the present invention, a high survival ratio is maintained even at the late-stage of culture, and an improvement can be expected in the yield of the desired polypeptide.

An appropriate culture period for producing a desired polypeptide using the cell of the present invention is usually 1 day to 3 months, preferably 1 day to 2 months, more preferably 1 day to 1 month.

With respect to various culture devices for animal cell culture, a fermentor type tank culture device, an air lift type culture device, a culture flask type culture device, a spinner flask type culture device, a microcarrier type culture device, a fluidized bed type culture device, a hollow fiber type culture device, a roller bottle type culture device, a packed bed type culture device, or the like may be used.

Culture may be performed by any culture method such as batch culture, fed-batch culture or continuous culture. Preferably, fed-batch culture or continuous culture is used. Fed-batch culture is more preferred.

When the cell of the present invention is cultured, taurine may be added to the medium in order to promote taurine uptake into cells. The concentration of taurine added to the medium is not particularly limited. The concentration is usually 0-100 g/L, preferably 0-20 g/L, more preferably 0-10 g/L.

When the polypeptide produced according to the method of the present invention has a biological activity useful as a pharmaceutical, it is possible to produce a pharmaceutical by mixing this polypeptide with pharmaceutically acceptable carriers or additives and formulating into a preparation.

Specific examples of pharmaceutically acceptable carriers and additives include water, organic solvents that are pharmaceutically acceptable, collagen, polyvinyl alcohol, polyvinylpyrrolidone, carboxyvinyl polymer, carboxymethylcellulose sodium, sodium

polyacrylate, sodium alginate, water-soluble dextran, carboxymethyl starch sodium, pectin, methylcellulose, ethyl cellulose, xanthan gum, gum Arabic, casein, agar-agar, polyethylene glycol, diglycerin, glycerin, propylene glycol, petrolatum, paraffin, stearyl alcohol, stearic acid, human serum albumin (HSA), mannitol, sorbitol, lactose, and surfactants that are
 5 acceptable as pharmaceutical additives.

Actual additives may be selected from the above-mentioned additives singly or in combination according to the dosage form of the therapeutic of the present invention, but are not limited to those listed above. For example, when a polypeptide is used in an injectable formulation, the purified polypeptide may be dissolved in a solvent such as physiological
 10 saline, buffer or a glucose solution, and then an adsorption inhibitor such as Tween 80, Tween 20, gelatin or human serum albumin may be added to the solution. Alternatively, a freeze-dried agent may be used to prepare a dosage form which is dissolved and reconstituted prior to use. Examples of the excipient useful for freeze-drying include sugar alcohols and saccharides such as mannitol and glucose.

15 Effective doses of the polypeptide may be appropriately selected depending on the type of the polypeptide, the type of the disease to be treated or prevented, the age of the patient, the severity of the disease, etc. For example, when the polypeptide is anti-glypican antibody, the effective dose of anti-glypican antibody is selected within a range of 0.001 mg to 1000 mg per kg of body weight per administration. Alternatively, a dose of 0.01-100000
 20 mg/body may be selected per patient. However, effective dose is not limited to these ranges.

The polypeptide may be administered either orally or parenterally, but parenteral administration is preferred. Specifically, injection (e.g., systemic or local administration by intravenous injection, intramuscular injection, intraperitoneal injection, subcutaneous
 25 injection, etc.), transnasal administration, transpulmonary administration, transdermal administration and the like may be enumerated.

The present invention provides a novel polypeptide which is any one of the following (A) to (E), provided that a polypeptide having the amino acid sequence as shown in SEQ ID NO: 4, a polypeptide having the amino acid sequence as shown in SEQ ID NO: 6,
 30 and a polypeptide having the amino acid sequence as shown in SEQ ID NO: 8 are excluded:

- (A) a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2;
- (B) a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;
- 35 (C) a polypeptide having 97% or more amino acid sequence homology with the amino acid

sequence as shown in SEQ ID NO: 2 and yet having taurine transporter activity;

(D) a polypeptide encoded by a DNA having the nucleotide sequence as shown in SEQ ID NO: 1; or

(E) a polypeptide encoded by a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.

The novel polypeptides of the present invention are hamster taurine transporter and those polypeptides which are functionally equivalent thereto.

In the present invention, the expression "functionally equivalent to hamster taurine transporter" means having activities similar to the activities of hamster taurine transporter, such as taurine-binding activity, activity to transport taurine into cells, etc. Such a polypeptide encompasses, for example, mutants of hamster taurine transporter.

As methods well-known to those skilled in the art for preparing polypeptides functionally equivalent to a specific polypeptide, methods of introducing mutations into polypeptides may be given. For example, those skilled in the art could prepare polypeptides functionally equivalent to hamster taurine transporter by appropriately introducing mutations into amino acids of hamster taurine transporter by site-directed mutagenesis (Hashimoto-Gotoh, T. et al. (1995) *Gene* 152, 271-275; Zoller, MJ, and Smith, M.(1983) *Methods Enzymol.* 100, 468-500; Kramer, W. et al. (1984) *Nucleic Acids Res.* 12, 9441-9456; Kramer W, and Fritz HJ (1987) *Methods. Enzymol.* 154, 350-367; Kunkel, TA (1985) *Proc Natl Acad Sci USA.* 82, 488-492; Kunkel (1988) *Methods Enzymol.* 85, 2763-2766). Mutations in amino acids may also occur in nature. Thus, a polypeptide which has an amino acid derived from the amino acid sequence of the hamster taurine transporter of the present invention by mutation of one or more amino acids and is functionally equivalent to hamster taurine transporter is also included in the polypeptide of the present invention.

Specific examples of polypeptides functionally equivalent to the hamster taurine transporter of the present invention include, but are not limited to, a polypeptide having an amino acid sequence derived from the amino acid sequence of the hamster taurine transporter by deletion of one or more amino acids, preferably 1-30 amino acids, more preferably 1-20 amino acids, even more preferably 1-10 amino acids, the most preferably 1-5 amino acids; a polypeptide having an amino acid sequence derived from the amino acid sequence of the hamster taurine transporter by addition of one or more amino acids, preferably 1-30 amino acids, more preferably 1-20 amino acids, even more preferably 1-10 amino acids, the most preferably 1-5 amino acids; and a polypeptide having an amino acid

sequence derived from the amino acid sequence of the hamster taurine transporter by substitution of one or more amino acids, preferably 1-30 amino acids, more preferably 1-20 amino acids, even more preferably 1-10 amino acids, the most preferably 1-5 amino acids, with other amino acids.

5 Amino acid residues to be mutated are not particularly limited. Preferably, amino acid residues are mutated to other amino acids in which the nature of the initial amino acid side chain is conserved. Specific examples of the nature of amino acid side chain include hydrophobic amino acids (A, I, L, M, F, P, W, Y and V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S and T), amino acids with an aliphatic side chain (G, A, V, L, I and P), amino
10 acids with a hydroxyl group-containing side chain (S, T and Y), amino acids with a sulfur atom-containing side chain (C and M), amino acids with a carboxylic acid and amide-containing side chain (D, N, E and Q), amino acids with a base-containing side chain (R, K and H) and amino acids with an aromatic-containing side chain (H, F, Y and W) (In parentheses are one-letter codes for amino acids).

15 It has been reported that a polypeptide having an amino acid sequence derived from an original amino acid sequence by modification (such as deletion, addition and/or substitution of one or more amino acids) maintains the biological activity of the original polypeptide (Mark, D. F. et al., Proc. Natl. Acad. Sci. USA (1984) 81, 5662-5666; Zoller, M. J. & Smith, M. Nucleic Acids Research (1982) 10, 6487-6500; Wang, A. et al., Science 224,
20 1431-1433; Dalbadie-McFarland, G et al., Proc. Natl. Acad. Sci. USA (1982) 79, 6409-6413).

As one example of the polypeptide in which one or more amino acid residues are added to the hamster taurine transporter of the present invention, a fusion polypeptide comprising the hamster taurine transporter may be given. Such a fusion polypeptide is
25 composed of the protein of the invention (hamster taurine transporter) and other polypeptide fused thereto. Such a fusion polypeptide is included in the present invention. Such a fusion polypeptide may be prepared by linking a gene encoding the hamster taurine transporter of the present invention in frame with a gene encoding the other polypeptide, transferring the resultant DNA into an expression vector and expressing the DNA in a host
30 cell. Techniques known to those skilled in the art may be used. There is no limitation on the polypeptide to be fused to the polypeptide of the present invention.

Examples of polypeptides to be fused to the polypeptide of the present invention include, but are not limited to, FLAG (Hopp, T. P. et al., BioTechnology (1988) 6, 1204-1210), 6xHis comprising six histidine (His) residues, 10xHis, influenza hemagglutinin
35 (HA), human c-myc fragment, VSV-GP fragment, p18HIV fragment, T7-tag, HSV-tag, E-tag,

SV40T antigen fragment, lck tag, α -tubulin fragment, B-tag, protein C fragment, glutathione-S-transferase (GST), influenza hemagglutinin (HA), immunoglobulin constant region, β -galactosidase and maltose-binding protein (MBP).

5 A commercially available gene encoding such polypeptide is fused to the gene encoding the polypeptide of the present invention. The fused gene thus prepared is expressed to prepare a fused polypeptide.

10 An alternative method known to those skilled in the art for preparing polypeptides functionally equivalent to a specific polypeptide is a method using the hybridization technique (Sambrook, J et al., Molecular Cloning 2nd ed., 9.47-9.58, Cold Spring Harbor Lab. Press, 1989). Those skilled in the art could routinely isolate a DNA highly homologous to the DNA sequence of the hamster taurine transporter of the present invention based on that DNA sequence or a part thereof, and isolate polypeptides functionally equivalent to the hamster taurine transporter from that DNA. Thus, a polypeptide which is encoded by a DNA hybridizing to the DNA, or a part thereof, encoding the hamster taurine
15 transporter of the present invention and is functionally equivalent to the hamster taurine transporter of the present invention is also included in the polypeptide of the present invention.

Hybridization conditions for isolating a DNA encoding a polypeptide functionally equivalent to the hamster taurine transporter of the present invention can be appropriately
20 selected by those skilled in the art. For example, low stringent hybridization conditions may be given. Low stringent hybridization conditions are, for example, 42°C, 2 x SSC and 0.1% SDS, preferably 50°C, 2 x SSC and 0.1% SDS. More preferably, high stringent conditions may be given. For example, high stringent conditions are 65°C, 2 x SSC and 0.1% SDS. Under these conditions, as the hybridization temperature is lowered, not only
25 DNAs with high homology but also DNAs with only low homology are obtained. Conversely, it is expected that only those DNAs with high homology are obtained as the hybridization temperature is elevated. However, not only the temperature but also a plurality of factors (such as salt concentrations) affect the stringency of hybridization. Those skilled in the art could appropriately select these factors to realize similar stringency.

30 The polypeptide encoded by a DNA isolated by these hybridization techniques usually has high homology with the hamster taurine transporter of the present invention in the amino acid sequence. The polypeptide of the present invention also include those polypeptides which are functionally equivalent to the hamster taurine transporter of the present invention and have high homology with the amino acid sequence of the hamster
35 taurine transporter of the present invention. The term "high homology" refers to usually

97% or more homology, preferably 98% or more homology, more preferably 99% or more homology. For determination of the homology of polypeptides, the algorithm described in Wilbur, W. J. and Lipman, D. J., Proc. Natl. Acad. Sci. USA (1983) 80, 726-730 may be followed.

5 The polypeptide of the present invention may vary in amino acid sequence, molecular weight, isoelectric point, presence or absence of sugar chains, morphology, etc. depending on the cell or host that produce the polypeptide or the purification method that will be described later. However, as long as the resultant polypeptide has functions equivalent to the functions of the hamster taurine transporter of the present invention, the
10 polypeptide is included in the present invention. For example, when the polypeptide of the present invention is expressed in a prokaryote (e.g., *Escherichia coli*), a methionine residue is added to the N-terminus of the initial amino acid sequence of the polypeptide. When the polypeptide of the present invention is expressed in a eukaryote (e.g., a mammalian cell), the N-terminal signal sequence is removed. The polypeptide of the present invention includes
15 such polypeptides.

 The polypeptide of the present invention may be prepared as a recombinant polypeptide or a natural polypeptide by methods known to those skilled in the art. A recombinant polypeptide may be prepared by incorporating a DNA encoding the polypeptide of the present invention in an appropriate expression vector, introducing the vector into an
20 appropriate host cell, collecting the resultant transformant, extracting a crude polypeptide, and then purifying the polypeptide by chromatography (such as ion exchange, reversed phase or gel filtration chromatography, or affinity chromatography in which an antibody to the polypeptide of the present invention is fixed in a column) or a combination of these chromatographic techniques.

25 When the polypeptide of the present invention is expressed in a host cell (e.g., animal cell or *E. coli*) as a fusion polypeptide with glutathione-S-transferase polypeptide or as a recombinant polypeptide with histidine residues added thereto, the expressed polypeptide may be purified with a glutathione column or a nickel column.

 After purification of a fusion polypeptide, regions other than the polypeptide of
30 interest may be cut off by thrombin or factor Xa and removed from the fusion polypeptide.

 When the polypeptide of the present invention is a natural polypeptide, the polypeptide may be isolated by purification methods known to those skilled in the art. For example, an extract from tissues or cells expressing the polypeptide of the present invention may be applied to an affinity column to which an antibody to the hamster taurine transporter
35 described later is bound. The antibody may be either a polyclonal antibody or a

monoclonal antibody.

Further, the present invention provides a DNA encoding a taurine transporter, which is any one of the following (a) to (e), provided that DNA having the nucleotide sequence as shown in SEQ ID NO: 3, DNA having the nucleotide sequence as shown in
 5 SEQ ID NO: 5, and DNA having the nucleotide sequence as shown in SEQ ID NO: 7 are excluded:

(a) a DNA encoding a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2;

(b) a DNA encoding a polypeptide which has an amino acid sequence derived from the
 10 amino acid sequence as shown in SEQ ID NO: 2 by substitution, deletion, addition and/or insertion of one or more (several) amino acid residues and yet has taurine transporter activity;

(c) a DNA encoding a polypeptide having 97% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2 and yet having taurine transporter
 15 activity;

(d) a DNA having the nucleotide sequence as shown in SEQ ID NO: 1; or

(e) a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.

20 The DNA of the present invention is used in the *in vivo* or *in vitro* production of the polypeptide of the present invention as described above. Further, the DNA of the present invention may be used in the creation of a cell which strongly expresses the hamster taurine transporter. The DNA of the present invention may take any form as long as it is capable of encoding the polypeptide of the present invention. That is, the DNA may be, for example, a
 25 cDNA synthesized from mRNA, a genomic DNA or a chemically synthesized DNA. It should be noted that, as long as the DNA is capable of encoding the polypeptide of the present invention, the DNA may have any nucleotide sequence based on the degeneracy of genetic codes.

The DNA of the present invention may be prepared by methods known to those
 30 skilled in the art. For example, the DNA may be prepared by preparing a cDNA library from a cell expressing the polypeptide of the present invention and performing hybridization using a part of the DNA sequence of the present invention (e.g., SEQ ID NO: 1) as a probe. The cDNA library may be prepared, for example, by the method described in Sambrook, J. et al., Molecular Cloning, Cold Spring Harbor Laboratory Press (1989). Alternatively, a
 35 commercial cDNA library may be used. It is also possible to prepare the DNA of the

present invention by preparing RNA from a cell expressing the polypeptide of the present invention, synthesizing oligo DNA molecules based on the DNA sequence of the present invention (e.g., SEQ ID NO: 1), and performing PCR using the oligo DNA molecules as primers to thereby amplify a cDNA encoding the taurine transporter.

5 Further, by determining the nucleotide sequence of the resultant cDNA, it is possible to determine the translation region encoding the polypeptide and to obtain the amino acid sequence of the polypeptide of the present invention. Further, by screening a genomic library using the resultant cDNA as a probe, it is possible to isolate a genomic DNA.

Specifically, the following procedures may be used. First, mRNA is isolated from
10 cells, tissues or the like expressing the polypeptide of the present invention. For the isolation of mRNA, the total RNA is prepared by known methods, for example, the guanidine ultracentrifugation method (Chirgwin, J. M. et al., Biochemistry (1979) 18, 5294-5299), the AGPC method (Chomczynski, P. and Sacchi, N., Anal. Biochem. (1987) 162, 156-159) or the like, and then mRNA is purified from the total RNA using mRNA
15 Purification Kit (Pharmacia), etc. Alternatively, mRNA may be prepared directly using QuickPrep mRNA Purification Kit (Pharmacia).

From the resultant mRNA, cDNA is synthesized using a reverse transcriptase. Alternatively, cDNA may be synthesized using a kit such as AMV Reverse Transcriptase First-Strand cDNA Synthesis Kit (SEIKAGAKU CORPORATION). It is also possible to
20 synthesize and amplify cDNA according to the 5'-RACE method (Frohman, M. A. et al., Proc. Natl. Acad. Sci. USA (1988) 85, 8998-9002; Belyavsky, A. et al., Nucleic Acids Res. (1989) 17, 2919-2932) using 5'-Ampli FINDER RACE Kit (Clontech) and polymerase chain reaction (PCR) with primers.

A DNA fragment of interest is prepared from the resultant PCR product and ligated
25 to a vector DNA to thereby prepare a recombinant vector. The vector is introduced into a host (e.g., E. coli), followed by selection of resultant colonies to thereby obtain a desired recombinant vector. The nucleotide sequence of the DNA of interest may be confirmed by a known method such as the dideoxynucleotide chain termination method.

Further, a nucleotide sequence of a higher expression efficiency can be designed for
30 the DNA of the present invention by considering the frequency of codon usage in the host to be used for expression (Grantham, R. et al., Nucleic Acids Research (1981) 9, p. 43-74). Further, the DNA of the present invention can be modified using commercially available kits or known methods. Examples of such modifications include, but are not limited to, digestion with restriction enzymes, insertion of synthetic oligonucleotides or appropriate
35 DNA fragments, addition of linkers, and insertion of an initiation codon (ATG) and/or a

termination codon (TAA, TGA or TAG).

The DNA of the present invention also includes a DNA which hybridizes to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1 under stringent conditions and encodes a polypeptide functionally equivalent to the hamster taurine transporter.

5 Stringent conditions can be appropriately selected by those skilled in the art, including, for example, low stringent conditions. Low stringent conditions refer to, for example, 42° C, 2 x SSC and 0.1% SDS, preferably 50° C, 2 x SSC and 0.1% SDS. More preferably, high stringent conditions may be selected. High stringent conditions refer to, for example, 65° C, 2 x SSC and 0.1% SDS. Under these conditions, as the hybridization
10 temperature is elevated, DNAs with a higher homology can be obtained. The above-described DNA which hybridizes is preferably a DNA derived from nature, e.g., cDNA or chromosomal DNA.

These DNAs isolated by hybridization techniques usually have a high nucleotide sequence identity with a DNA encoding the hamster taurine transporter of the present
15 invention. The DNA of the present invention also includes a DNA which encodes a polypeptide functionally equivalent to the hamster taurine transporter of the present invention and has high identity with a DNA encoding the hamster taurine transporter of the present invention. The term "high identity" refers to usually 96% or more homology, preferably 98% or more homology, more preferably 99% or more identity. The identity of
20 nucleotide sequences may be determined by algorithm BLAST (Karlin and Altschul, Proc. Natl. Acad. Sci. USA 90:5873-5877, 1993). Based on this algorithm, programs such as BLASTN and BLASTX have been developed (Altschul et al. J. Mol. Biol. 215:403-410, 1990). When nucleotide sequences are analyzed by BLASTN based on BLAST, parameters may be set as score = 100 and wordlength = 12, for example. Specific
25 procedures for these analysis methods are known (<http://www.ncbi.nlm.nih.gov>).

Further, the present invention provides a vector into which the DNA of the present invention has been inserted. The vector of the present invention is useful for retaining the DNA of the present invention within the host cell and for permitting expression of the polypeptide of the present invention (i.e., hamster taurine transporter or a polypeptide
30 functionally equivalent thereto). The vector of the present invention is also useful for permitting the host cell to strongly express the taurine transporter. By permitting the host cell to strongly express the taurine transporter, taurine uptake into the host cell is promoted, which leads to an increased production of a desired polypeptide in the host cell.

When the host cell to be used is *E. coli*, it is preferable that the vector has a
35 replication origin ("ori") so that the vector is largely amplified in *E. coli* (e.g., JM109, DH5α,

HB101 and XL1-Blue) and prepared in large quantity, and also genes for selecting transformed *E. coli* (e.g., drug resistance genes that enable discrimination of transformant with some drugs such as ampicillin, tetracycline, kanamycin or chloramphenicol). Examples of preferable vectors include, but are not limited to, M13 vectors, pUC vectors, pBR322, pBluescript and pCR-Script. In addition to these vectors, pGEM-T, pDIRECT, pT7, etc. may be enumerated when the vector is used for the purpose of subcloning a cDNA and cutting off the subcloned cDNA. When the vector is used for the purpose of producing the polypeptide of the present invention, an expression vector is especially useful. When expression in *E. coli* is intended, the expression vector preferably has the above-described features so that the vector is amplified in *E. coli*, and it also preferably has a promoter which allows efficient expression in *E. coli* such as JM109, DH5 α , HB101 or XL1-Blue, e.g., lacZ promoter (Ward et al, Nature (1989) 341, 544-546; FASEB J. (1992) 6, 2422-2427), araB promoter (Better et al, Science (1988) 240, 1041-1043) or T7 promoter. Specific examples of such vector include, in addition to those listed above, pGEX-5X-1 (Pharmacia), QIAexpress system (Qiagen), pEGFP, or pET (for its host, T7 RNA polymerase-expressing BL21 is preferred).

The vector may comprise signal sequences for polypeptide secretion. When the polypeptide is to be produced in the periplasm of *E. coli*, pelB signal sequence (Lei, S. P. et al., J. Bacteriol. (1987) 169, 4379) may be used as a signal sequence for polypeptide secretion. Introduction of the vector into a host cell may be performed, for example, by the calcium chloride method or electroporation.

In cases where a host cell other than *E. coli* is used, vectors useful for producing the polypeptide of the present invention include, but are not limited to, mammal-derived expression vectors [e.g., pcDNA3 from Invitrogen; pEGF-BOS (Nucleic Acids. Res. 1990, 18(17), p. 5322); pEF, pCDM8], insect cell-derived expression vectors (e.g., Bac-to-BAC baculovirus expression system from GIBCO BRL; pBacPAK8), plant-derived expression vectors (e.g., pMH1, pMH2), animal virus-derived expression vectors (e.g., pHSV, pMV, pAdexLcw), retrovirus-derived expression vectors (e.g., pZIpneo), yeast-derived expression vectors (e.g., Pichia Expression Kit from Invitrogen; pNV11; SP-Q01), and *Bacillus subtilis*-derived expression vectors (e.g., pPL608, pKTH50).

When expression of the polypeptide in animal cells (such as CHO cells, COS cells, NIH3T3 cells, etc.) is intended, the vector preferably has a promoter necessary for expressing the polypeptide in those cells. Examples of such promoter include, but are not limited to, SV40 promoter (Mulligan et al, Nature (1979) 277, 108), MMLV-LTR promoter, EF1 α promoter (Mizushima et al., Nucleic Acids Res. (1990) 18, 5322) and CMV promoter.

More preferably, the vector also has genes for selecting transformed cells (e.g., drug resistance genes that enable discrimination with drugs such as neomycin or G418). Examples of vectors having such properties include, but are not limited to, pMAM, pDR2, pBK-RSV, pBK-CMV, pOPRSV and pOP13.

5 Further, when stable expression of a gene of interest and intracellular amplification of the copy number of the gene are intended, the following method may be used. Briefly, into CHO cells lacking a nucleic acid synthesis pathway, a vector having DHFR gene that complements the lack (e.g., pCHOI) is introduced, followed by amplification with methotrexate (MTX). On the other hand, when tentative expression of a gene of interest is
10 intended, a method may be used in which COS cells carrying a gene expressing SV40T antigen on the chromosome is transformed with a vector having the replication origin of SV40 (e.g., pcD). As the replication origin, a replication origin derived from polyomavirus, adenovirus or bovine papillomavirus (BPV) may also be used. Further, the expression vector may contain selectable markers for amplifying the copy number of the gene in a host
15 cell system. Examples of such selectable markers include, but are not limited to, aminoglycoside phosphotransferase (APH) gene, thymidine kinase (TK) gene, *E. coli* xanthine-guanine phosphoribosyl transferase (Ecogpt) gene and dihydrofolate reductase (dhfr) gene.

The present invention also provides a host cell into which the vector of the present
20 invention has been transferred. The host cell into which the vector of the present invention is transferred is not particularly limited. For example, *E. coli* or various animal cells may be used. The host cell of the present invention may be used, for example, as a production system for the preparation or expression of the polypeptide of the present invention. Further, the host cell of the present invention is capable of expressing the taurine transporter
25 strongly, promoting taurine uptake, and increasing the yield of the desired polypeptide. For the production of the polypeptide, there are *in vivo* and *in vitro* production systems. Examples of *in vitro* production systems include systems using eukaryotes and systems using prokaryotes.

When eukaryotes are used, animal cells, plant cells, fungal cells, etc. may be used
30 as the host. Specific examples of animal cells include mammalian cells, such as CHO cells (J. Exp. Med. (1995) 108, 945), COS cells, 3T3 cells, myeloma cells, BHK (baby hamster kidney) cells, HeLa cells and Vero cells; amphibian cells, such as oocytes of *Xenopus laevis* (Valle, et al., Nature (1981) 291, 358-340); or insect cells, such as sf9, sf21 and Tn5 cells. Among CHO cells, dhfr-CHO lacking DHFR gene (Proc. Natl. Acad. Sci. USA (1980) 77,
35 4216-4420) and CHO K-1 (Proc. Natl. Acad. Sci. USA (1968) 60, 1275) are used with

particular advantage. When high expression is intended in an animal cell, CHO cells are especially preferred. Introduction of the expression vector into the host cell may be performed by such methods as the calcium phosphate method, the DEAE dextran method, a method using a cationic ribosome DOTAP (Boehringer-Mannheim), electroporation,
 5 lipofection, etc.

As plant cells for polypeptide production, a *Nicotiana tabacum*-derived cell is known as a polypeptide production system and this may be subjected to callus culture. As fungal cells for polypeptide production, specific examples include yeast belonging to the genus *Saccharomyces*, e.g., *Saccharomyces cerevisiae*, and filamentous fungi belonging to
 10 the genus *Aspergillus*, e.g., *Aspergillus niger*.

When prokaryotes are used, production systems using bacterial cells are known. Specific examples of such bacterial cells include *E. coli* (such as JM109, DH5 α , HB101) and *Bacillus subtilis*.

The polypeptide encoded by a gene of interest may be obtained by transforming
 15 these cells with the gene of interest and culturing the transformed cells *in vitro*. The culture may be performed by known methods. For example, as a culture broth for animal cells, a medium such as DMEM, MEM, RPMI1640 or IMDM may be used. A serum supplement such as fetal calf serum (FCS) may be used jointly. Alternatively, serum-free culture may be performed. The pH during culture is preferably about 6 to 8. The culture is usually
 20 performed at about 30-40°C for about 15-200 hours. If necessary, replacement of the medium, aeration and agitation are carried out.

On the other hand, *in vivo* production systems include those using animals or plants. A gene of interest is transferred into these animals or plants to produce the polypeptide in the animal bodies or plant bodies. Then, the polypeptide is collected. The term "host" as
 25 used herein includes such animals or plants.

When animals are used, available production systems include those using mammals or insects. Goat, pig, sheep, mouse and cattle may be used as mammals (Vicki Glaser, SPECTRUM Biotechnology Applications, 1993). When mammals are used, transgenic animals may be used.

30 First, a gene of interest is fused to a gene encoding a polypeptide produced inherently in milk (such as goat β -casein) to thereby prepare a fusion gene. A DNA fragment containing this fusion gene is injected into a goat embryo, which is then implanted in the uterus of a female goat. The polypeptide of interest can be obtained from the milk produced by transgenic goats born from the goat which accepted the embryo or the offspring
 35 of the transgenic goats. In order to increase the yield of milk containing the polypeptide

produced by the transgenic goats, hormones may be appropriately administered to the transgenic goats (Ebert, K. M. et al., Bio/Technology (1994) 12, 699-702).

Examples of insects which may be used include silkworm. In this case, silkworm is infected with baculovirus carrying a transferred gene encoding the polypeptide of interest.
5 The polypeptide of interest can be obtained from the body fluid of the silkworm (Susumu, M. et al., Nature (1985) 315, 592-594).

Furthermore, when plants are used, tobacco can typically be used. When tobacco is used, a gene encoding the polypeptide of interest is inserted into a plant expression vector (e.g., pMON 530), which is then transferred into a bacterium such as *Agrobacterium*
10 *tumefaciens*. A tobacco plant (e.g., *Nicotiana tabacum*) is infected with the resultant bacterium. The polypeptide of interest can be obtained from leaves of this plant (Julian, K.-C. Ma et al., Eur. J. Immunol. (1994) 24, 131-138).

The polypeptide thus obtained can be isolated from the inside of the host cell or from its outside (e.g., medium), and purified to a substantially pure and homogeneous
15 polypeptide. Isolation and purification of polypeptides can be performed using conventional isolation and purification methods for polypeptides, and are not limited in any way. For example, polypeptides can be isolated and purified by appropriate selection and combination of various tools and techniques, such as chromatography columns, filters, ultrafiltration, salting-out, precipitation with solvent, extraction with solvent, distillation,
20 immunoprecipitation, SDS-polyacrylamide gel electrophoresis, isoelectric focusing, dialysis, recrystallization, etc.

Examples of chromatography include affinity chromatography, ion exchange chromatography, hydrophobic chromatography, gel filtration, reverse-phase chromatography, adsorption chromatography, etc. (Strategies for Protein Purification and Characterization: A
25 Laboratory Course Manual. Ed. Daniel R. Marshak et al., Cold Spring Harbor Laboratory Press, 1996). These chromatographic techniques can be carried out using liquid phase chromatography, for example, HPLC, FPLC, etc. The present invention also includes those polypeptides highly purified using these purification methods.

Before or after the purification, it is also possible to give optional modifications to
30 the polypeptide or remove a partial peptide therefrom by reacting the polypeptide with an appropriate polypeptide modification enzyme. Examples of such enzyme include, but are not limited to, trypsin, chymotrypsin, lysyl endopeptidase, protein kinase and glucosidase.

The present invention also provides an antibody that binds to the polypeptide of the present invention. The form of the antibody of the present invention is not particularly
35 limited; the antibody of the present invention includes polyclonal antibodies and monoclonal

antibodies. In addition, antisera obtained by immunizing immune animals such as rabbit with the polypeptide of the present invention, and polyclonal antibodies and monoclonal antibodies of every class are also included in the antibody of the present invention.

5 The polypeptide used as a sensitizing antigen may be an intact polypeptide or a partial peptide thereof. Examples of such partial peptides include amino (N)-terminal fragments and carboxyl (C)-terminal fragments of the polypeptide. The term "antibody" as used herein means an antibody that reacts with the full length polypeptide or fragments thereof.

10 After inserting a gene encoding the polypeptide of the present invention or a fragment thereof into a known expression vector and transforming the host cell described earlier in this specification with this expression vector, the desired polypeptide or a fragment thereof is obtained from inside and outside of the host cells by a known method. This polypeptide or a fragment thereof may be used as a sensitizing antigen. Alternatively, cells expressing the polypeptide of the present invention or a lysate thereof, or a chemically synthesized polypeptide of the present invention may also be used as a sensitizing antigen.

15 Although there is no particular limitation on the species of mammals to be immunized with a sensitizing antigen, it is preferable to select mammals taking into consideration the compatibility with the parent cell to be used for cell fusion, and animals of the rodent, lagomorph and primate are generally used.

20 For example, mouse, rat, hamster and the like are used as rodent animals; rabbit is used as a lagomorph animal, and monkey is used as a primate animal. Among monkeys, catarrhine monkeys (old world monkeys) are used, as exemplified by cynomolgus monkey (*Macaca fascicularis*), rhesus monkey, baboon, chimpanzee and the like.

25 Immunization of animals with a sensitizing antigen is carried out according to any known method. Generally, immunization is performed by intraperitoneal or subcutaneous injection of the antigen to mammals. Specifically, the antigen is appropriately diluted and suspended in PBS (Phosphate-Buffered Saline), physiological saline or the like, optionally mixed with a suitable amount of usual adjuvant (e.g., Freund's complete adjuvant), emulsified and then administered to mammals, preferably followed by several booster injections of the antigen mixed with an appropriate amount of Freund's incomplete adjuvant

30 every 4 to 21 days. In addition, suitable carriers can be used at the time of immunization with the antigen. Subsequently, elevation of the level of a desired antibody in the sera of animals is confirmed by a conventional method.

35 In order to obtain polyclonal antibodies to the polypeptide of the present invention, the blood of mammals sensitized with the antigen is withdrawn after confirming elevation of

the level of a desired antibody in sera. Then, sera are isolated from the blood by a known method. A serum containing a polyclonal antibody may be used as the polyclonal antibody. If necessary, a fraction containing the polyclonal antibody may be further isolated from the serum and used as the polyclonal antibody. For example, a fraction recognizing only the polypeptide of the present invention is obtained by using an affinity column to which the polypeptide of the present invention has been coupled, and further purified by using protein A or protein G column to prepare immunoglobulin G or M.

Monoclonal antibodies can be obtained by removing immunocytes from the mammals sensitized with the above-described antigen, after confirming the elevation of the level of a desired antibody in their sera, and then subjecting the immunocytes to cell fusion. Immunocytes preferably used for cell fusion are spleen cells. Parent cells to be fused to the above-described immunocytes are preferably mammalian myeloma cells, more preferably myeloma cells which have acquired characteristics for the selection of fused cells with drugs.

Cell fusion between the above-described immunocytes and myeloma cells can be carried out according to a known method, for example, the method of Milstein et al. (Galfre, G. and Milstein, C., *Methods Enzymol.* (1981) 73, 3-46).

Hybridomas thus obtained by cell fusion are selected by culturing them in a conventional selection medium, for example, HAT culture medium (culture broth containing hypoxanthine, aminopterin and thymidine). Culturing in the HAT medium is continued for a sufficient time to kill other cells (non-fused cells) than desired hybridomas, usually for several days to several weeks. Then, conventional limiting culture-dilution method is performed to carry out screening and cloning of hybridomas producing the antibody of interest.

Subsequently, the resultant hybridomas are transplanted into the abdominal cavities of mice, and abdominal dropsies are collected from the mice to thereby obtain monoclonal antibodies. These monoclonal antibodies may be purified by ammonium sulfate precipitation, with protein A or protein G column, by DEAE ion exchange chromatography, with affinity column to which the polypeptide of the present invention has been coupled, etc. The antibody of the present invention can be used for purification and detection of the polypeptide of the present invention.

Further, the thus obtained monoclonal antibodies can also be prepared as recombinant antibodies using recombinant DNA techniques (see, for example, Borrebaeck, C. A. K. and Larrick, J. W., *THERAPEUTIC MONOCLONAL ANTIBODIES*, Published in the United Kingdom by MACMILLAN PUBLISHERS LTD, 1990). A recombinant antibody is produced by cloning a gene encoding the recombinant antibody from hybridomas

or the antibody-producing immunocytes (such as sensitized lymphocytes), incorporating the gene into an appropriate vector, and transferring the vector into host cells for antibody production. The produced recombinant antibody is also included in the present invention.

The antibody of the present invention may be antibody fragments or modified
 5 antibodies as long as they are capable of binding to the polypeptide of the present invention. Examples of such antibody fragments include Fab, F(ab')₂, Fv, or a single-chain Fv (scFv) prepared by linking the Fv of H-chain to the Fv of L-chain via a suitable linker (Huston, J. S. et al., Proc. Natl. Acad. Sci. USA (1988) 85, 5879-5883). Specifically, antibody fragments are produced by digesting the antibody with enzymes, for example, papain or pepsin; or by
 10 constructing a gene encoding such a fragment, inserting the gene into an expression vector, and expressing it in a suitable host cell (see, for example, Co, M. S. et al., J. Immunol. (1994) 152, 2968-2976; Better, M. and Horwitz, A. H., Methods Enzymol. (1989) 178, 476-496; Pluckthun, A. and Skerra, A., Methods Enzymol. (1989) 178, 497-515; Lamoyi, E., Methods Enzymol. (1986) 121, 652-663; Rousseaux, J. et al., Methods Enzymol. (1986) 121, 663-669;
 15 Bird, R. E. and Walker, B. W., Trends Biotechnol. (1991) 9, 132-137).

As a modified antibody, the antibody bound to various molecules such as polyethylene glycol (PEG) may be also used. The "antibody" of present invention also includes these modified antibodies. These modified antibodies can be prepared by chemically modifying the antibody obtained as described above. These modification
 20 methods have been already established in the art.

Antibodies obtained as described above can be purified to homogeneity. For the isolation and purification of the antibody used in present invention, any methods of isolation and purification used for conventional polypeptides may be used. For example, chromatography columns such as affinity chromatography columns, filters, ultrafiltration,
 25 salting-out, dialysis, SDS-polyacrylamide gel electrophoresis, isoelectric focusing, etc. may be used independently or in appropriate combinations (Antibodies: A Laboratory Manual. Ed Harlow and David Lane, Cold Spring Harbor Laboratory, 1988) but these are not the sole examples. Concentration of the antibody obtained as described above can be determined by measuring absorbance or by a method such as enzyme-linked immunosorbent assay
 30 (ELISA).

Examples of columns used in affinity chromatography include protein A column and protein G column. As columns using protein A, Hyper D, POROS, Sepharose F. F.TM (Pharmacia), etc. may be given.

Examples of chromatography other than affinity chromatography include ion
 35 exchange chromatography, hydrophobic chromatography, gel filtration, reversed-phase

chromatography, adsorption chromatography, etc. (Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Ed Daniel R. Marshak et al., Cold Spring Harbor Laboratory Press, 1996). These chromatographic techniques can be carried out using liquid phase chromatography such as HPLC, FPLC, etc.

5 Furthermore, absorbance measurement, ELISA, EIA (enzyme immunoassay), RIA (radioimmunoassay) or fluorescent antibody technique can be used as methods for measuring the antigen-binding activity of the antibody of present invention. When ELISA is used, the polypeptide of the present invention is added to plates on which the antibody of the present invention has been immobilized, and then a sample containing an antibody of interest
10 (e.g., culture supernatant of antibody-producing cells or purified antibody) is added to the plates. A secondary antibody labeled with an enzyme (e.g., alkaline phosphatase) that recognizes the antibody is added to the plates. After incubating and washing the plates, a substrate for the enzyme (e.g., p-nitrophenyl phosphate) is added to determine the absorbance to thereby evaluate the antigen-binding activity. Instead of the entire
15 polypeptide, a fragment thereof may be used. For example, fragments comprising its C-terminal or N-terminal may be used. For evaluating the activity of the antibody of the present invention, BIAcore (Pharmacia) may be used.

EXAMPLES

20 Hereinbelow, the present invention will be described in more detail with reference to the following Examples. It should be noted that these Examples are provided only for illustrating the present invention and not for limiting the scope of the present invention.

[EXAMPLE 1] Cloning of CHO Cell-Derived Hamster Taurine Transporter Gene

25 Total RNA was extracted from anti-IL-6 receptor antibody-producing cells (A CHO DXB11 cell line into which an anti-IL-6 receptor antibody gene had been transferred) (Japanese Unexamined Patent Publication No. Hei 8-99902), and then cDNA was synthesized therefrom in a poly(A) dependent manner. Hamster taurine transporter (TauT) gene was obtained by PCR using as a template the cDNA fragmented with three restriction
30 enzymes, Sall, XhoI and EcoRI. As PCR primers, those containing the 5'-end and the 3'-end sequence conserved between rat and mouse TauTs were designed. The nucleotide sequence of the cloned gene was determined. From its homology with other TauT genes of known species, the cloned gene was confirmed to encode hamster TauT (Fig. 1). The amino acid sequence of hamster TauT has high homology with mouse TauT (96% identity),
35 rat TauT (96% identity) and human TauT (93% identity); it was predicted that hamster TauT

is a transporter with 12 transmembrane regions (Fig. 2).

[EXAMPLE 2] Increase in Viable Cell Density, Inhibition of Lactate Production and Increase in Antibody Yield, as Caused by Transfer of Hamster Taurine Transporter

CMV promoter expression plasmid pHyg/TauT (Fig. 3) was constructed by adding Kozak sequence to the hamster TauT (hereinafter, TauT) gene obtained by cloning in Example 1. Control plasmid pHyg without pHyg/TauT or TauT gene was introduced by electroporation into the parent strain anti-glypican-3 antibody producing CHO cell (see WO 2006/006693). After selection of expression plasmid-transferred cells in the presence of hygromycin (400 µg/ml), all of the stably growing cell strains were expanded (pHyg/TauT: 8 strains; pHyg: 7 strains). TauT mRNA was prepared. Subsequently, 7 strains were confirmed to express TauT more strongly than the parent strain by the TaqMan method; they were selected as pHyg/TauT transferred cells. The mean mRNA expression level of these transferred cells (7 strains) was about 40 times larger than the control (7 strains). Cells of the total 14 strains were subjected to batch culture and fed-batch culture in 50 ml shaker flasks with an initial cell density of 2×10^5 cells/ml. On day 7 of culture (late-stage), viable cell densities, lactate yields and anti-glypican-3 antibody yields in those strains were compared. In batch culture, growth inhibitory substances such as lactate accumulate in culture broth as cells grow and their growth is inhibited. However, the viable cell densities (Fig. 4) and lactate yields (Fig. 5) in pHyg/TauT transferred cells were superior to those in pHyg transferred cells (t test; $p < 0.05$). With respect to anti-glypican-3 antibody yield, 4 out of the 7 strains of pHyg/TauT-transferred cell showed antibody yields higher than the highest yield in pHyg-transferred cell (Fig. 6). Further, since superiority of pHyg/TauT transferred cells in anti-glypican-3 antibody yield became more evident (t test; $P < 0.01$; Fig. 7) in fed-batch culture, pHyg/TauT transferred T10 strain (which showed the highest growth ability among the above 4 strains) and the parent strain were subjected to fed-batch culture in 1 L jar. As a result, the viable ratio of T10 was maintained at 80% or more even on day 32 of culture (Fig. 8), with inhibited lactate production. Consequently, its anti-glypican-3 antibody yield achieved 2.9 g/L on day 35 of culture (Fig. 9). It was confirmed by flow cytometric analysis that TauT-transferred T10 cell was expressing TauT molecules on the cell membrane (Fig. 10). These results suggest that by artificially expressing hamster Taut, it is possible to raise the potential of antibody-producing cells and create strains capable of enhanced antibody production.

[EXAMPLE 3] Inhibition of Ammonia Production, Taurine Uptake, Increase in

Glutamine Consumption and Taurine Non-Dependent Antibody Yield in Hamster TauT Transferred Strains

The parent strain and pHyg/TauT transferred strain were fed-batch cultured in 1 L jar with an initial cell density of 2×10^5 cells/ml. A part of the culture broth containing 450×10^5 cells was taken from the jar at appropriate time points. After the culture supernatant was separated by centrifugation, 1 ml of cooled sterile water containing a protease inhibitor (Complete Mini; Roche Diagnostics; Protease inhibitor cocktail tablets) was added to the cell pellet. Then, the cells were completely disrupted on ice in a sonicator (MISONIX ASTRASON MODEL XL2020) with a set of 5 seconds pulse-on and 5 seconds pulse-off being repeated 12 times. The total volume of the thus treated cells was applied to a centrifugal filter unit to thereby prepare a filtrate with a molecular weight of 5000 or less. This filtrate was used as a sample for determining intracellular amino acids. Each sample was subjected to detection and comparison of absorbance at 570 nm using a ninhydrin reagent L-8500 set (Wako PureChemical Industries) and an improved model of Hitachi fully automated amino acid analyzer (L-8500). Thus, various amino acid concentrations in samples were determined. Since the concentrations of amino acids and ammonia in culture broth were directly measured values, concentration comparisons in the order of μM were performed. On the other hand, since intracellular concentrations were obtained after addition of 1 ml of cooled sterile water to the cell pellet and sonication thereof, the measured concentrations of various amino acids and ammonia were converted into values per cell, followed by comparison of the converted values. To determine the ammonia concentration ratios shown in Fig. 11, the detected ammonia value per 450×10^5 cells in the parent strain at the start of 1 L jar fed-batch culture was taken as 1 and compared with detected values at the start of the culture and on days 6, 12 and 18 of the culture in the transferred strain. The taurine values in Fig. 12 and the glutamine values in Fig. 13 were also determined by the above-described amino acid analysis.

As a result, the intracellular ammonia in pHyg/TauT transferred strain was maintained at a low concentration at the late stage of culture; it is believed that this contributes to high antibody yield (Fig. 11).

Intracellular taurine concentration ratios were determined in the same manner as described above for ammonia concentrations (Fig. 12), except that the detected ammonia value per 200×10^5 cells in the parent strain on day 4 of 50 ml shaker batch culture was taken as 1.

As a result, it was found that pHyg/TauT transferred strain had taken up taurine in a manner dependent on the amount of taurine added and that its uptake was almost equal to

that by the parent strain. However, as shown in Fig. 13, glutamine consumption in pHyg/TauT transferred strain was remarkably high compared to the parent strain and was not dependent on the initial taurine concentration. It has been reported that glutamine improves cell growth, survival ratio and antibody production ability in hybridomas to thereby raise their antibody yields (Enzyme and Microbial Technology 17:47-55, 1995). Therefore, the antibody production enhancement effect of pHyg/TauT transferred strain may be caused by taurine transporter-mediated uptake of amino acids other than taurine (e.g., glutamine). The glutamine concentrations were obtained by converting the values determined by amino acid analysis of the culture broth on day 4 of culture in Fig. 12 into values per 1×10^5 cells.

Actually, anti-glypican-3 antibody yield was not dependent on the initial taurine concentration (0-500 mM (62.575 g/L)) at the start of 50 ml shaker fed-batch culture (Fig. 14). No significant difference was observed in the parent strains in the effect of initial taurine concentration on antibody yield.

The results described so far suggest that TauT strongly expressing strains have high antibody production ability even if the medium does not contain taurine at the start of culture and that there is a possibility that such strains also promote uptake of amino acids other than taurine.

The present invention is applicable to any antibody-producing cell.

20

INDUSTRIAL APPLICABILITY

The present invention is applicable to production of polypeptides.

25 SEQUENCE LISTING FREE TEXT

<SEQ ID NO: 1>

SEQ ID NO: 1 shows the nucleotide sequence of a gene encoding hamster taurine transporter.

<SEQ ID NO: 2>

30 SEQ ID NO: 2 shows the amino acid sequence of hamster taurine transporter.

<SEQ ID NO: 3>

SEQ ID NO: 3 shows the nucleotide sequence of a gene encoding rat taurine transporter.

<SEQ ID NO: 4>

35 SEQ ID NO: 4 shows the amino acid sequence of rat taurine transporter.

<SEQ ID NO: 5>

SEQ ID NO: 5 shows the nucleotide sequence of a gene encoding mouse taurine transporter.

<SEQ ID NO: 6>

5 SEQ ID NO: 6 shows the amino acid sequence of mouse taurine transporter.

<SEQ ID NO: 7>

SEQ ID NO: 7 shows the nucleotide sequence of a gene encoding human taurine transporter.

<SEQ ID NO: 8>

10 SEQ ID NO: 8 shows the amino acid sequence of human taurine transporter.

CLAIMS

1. A method of producing a polypeptide by a mammalian cell, comprising culturing the mammalian cell which has a transferred DNA encoding a taurine transporter and a transferred DNA encoding a desired polypeptide and thereby the polypeptide production by the mammalian cell is enhanced.

2. The method of claim 1, wherein the cell is Chinese hamster ovary cells.

3. The method of claim 1 or 2, wherein the desired polypeptide is an antibody.

4. The method of any one of claims 1 to 3, wherein the DNA encoding the taurine transporter is any one of the following (a) to (e):

(a) a DNA encoding a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8;

(b) a DNA encoding a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;

(c) a DNA encoding a polypeptide having 70% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 and yet having taurine transporter activity;

(d) a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7; or

(e) a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.

5. The method of any one of claims 1 to 4, comprising culturing the cells in a medium with a taurine concentration of 0 to 100 g/L.

6. Use of a mammalian cell, comprising a transferred DNA encoding a taurine transporter and a transferred DNA encoding desired polypeptide for enhancing the production of the polypeptide.

7. The use of claim 6, wherein the cell is a Chinese hamster ovary cells.

8. The use of claim 6 or 7, wherein the desired polypeptide is an antibody.

9. The use of any one of claims 1 to 3, wherein the DNA encoding the taurine transporter is any one of the following (a) to (e):

- (a) a DNA encoding a polypeptide having the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8;
- (b) a DNA encoding a polypeptide which has an amino acid sequence derived from the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 by substitution, deletion, addition and/or insertion of one or more amino acid residues and yet has taurine transporter activity;
- (c) a DNA encoding a polypeptide having 70% or more amino acid sequence homology with the amino acid sequence as shown in SEQ ID NO: 2, 4, 6 or 8 and yet having taurine transporter activity;
- (d) a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7; or
- (e) a DNA which hybridizes to a DNA complementary to a DNA having the nucleotide sequence as shown in SEQ ID NO: 1, 3, 5 or 7 under stringent conditions and yet encodes a polypeptide having taurine transporter activity.

[illegible]

Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14

