



(11) **EP 1 219 299 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
14.02.2007 Bulletin 2007/07

(51) Int Cl.:
A61K 39/35 ^(2006.01) **A61K 39/36** ^(2006.01)
C07K 14/415 ^(2006.01)

(21) Application number: **00128659.0**

(22) Date of filing: **28.12.2000**

(54) **Allergy vaccines and their preparation**

Allergievakzine sowie ihre Herstellung

Vaccins contre l'allergie et sa préparation

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
LT LV MK SI

(43) Date of publication of application:
03.07.2002 Bulletin 2002/27

(73) Proprietor: **BIOMAY Produktions- und Handels-
Aktiengesellschaft**
1090 Wien (AT)

(72) Inventors:
• **Focke, Margarete, Dr.**
1140 Wien (AT)
• **Mahler, Vera, Dr.**
91054 Erlangen (DE)
• **Sperr, Wolfgang R., Dr.**
1190 Wien (AT)
• **Valent, Peter, Dr.**
1170 Wien (AT)
• **Kraft, Dietrich, Dr.**
1170 Wien (AT)
• **Valenta, Rudolf, Dr.**
2604 Theresienfeld (AT)

(74) Representative: **Keller, Günter et al**
Lederer & Keller
Patentanwälte
Prinzregentenstrasse 16
80538 München (DE)

(56) References cited:
EP-A- 0 714 662 **WO-A-96/27005**
WO-A-97/05258 **WO-A1-97/24139**

- **23rd Symposium of the Collegium Internationale Allergologicum, May 18-23, 2000, Hakone Japan & M. FOCKE ET AL.: "Nonallergic peptides from surface-exposed areas or B-cell epitopes of allergens for specific immunotherapy" INT. ARCH. ALLERGY IMMUNOL., vol. 124, 2001, pages 298-399, XP001016272**
- **23rd Symposium of the Collegium Internationale Allergologicum, May 18-23, 2000, Hakone Japan & E. GANGLBERGER ET AL.: "IgE momotopes of birch pollen allergen Bet v 1 induce blocking IgG in mice" INT. ARCH. ALLERGY IMMUNOL., vol. 124, 2001, pages 395-397, XP001016271**
- **M. GAJHEDE ET AL.: "X-ray and NMR structure of Ber v 1 the origin of birch pollen allergy" NATURE STRUCTURAL BIOLOGY, vol. 3, no. 12, 1996, pages 1040-1045, XP001016261**
- **R. FURMONAVICIENE ET AL.: "The use of phage-peptide libraries to define th epitope specificity of a mouse monoclonal ant p1 antibody representative of a major component of the human immunoglobulin E anti-Der p 1 response" CLIN. EXP. ALLERGY, vol. 29, 1999, pages 1563-1571, XP001016249**
- **E. GANGLBERGER ET AL.: "Allergen mimotopes for 3-dimensional apitope search and induction of antibodies inhibiting human IgE" FASEB J., vol. 14, no. 14, November 2000 (2000-11), pages 2177-2184, XP002177915**

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 219 299 B1

Description

[0001] Almost 500 million individuals suffer from Type I allergy, a genetically determined hypersensitivity disease which is based on the formation of IgE antibodies against *per se* harmless antigens (i.e., allergens). In sensitized patients, allergen contact can induce a variety of symptoms ranging from allergic rhinoconjunctivitis, atopic dermatitis, diarrhoea, bronchial asthma, and sometimes life-threatening anaphylaxis. The immediate symptoms of allergic disease are caused by crosslinking of effector cell-bound IgE antibodies by allergens which triggers the release of biologically active mediators (e.g., histamine, leukotrienes). In addition, allergen presentation to T cells may be also mediated by IgE antibodies and induce chronic symptoms of allergic disease (e.g., atopic dermatitis, chronic asthma) via T cell activation and release of proinflammatory cytokines.

[0002] The only curative approach towards allergy treatment, allergen-specific immunotherapy, is based on the continuous administration of the disease-eliciting allergens to the patients in order to induce allergen-specific non-responsiveness. Multiple studies document the clinical efficacy of specific immunotherapy, but two major disadvantages must be overcome: First, systemic administration of allergens can induce life-threatening anaphylactic side effects. Second, immunotherapy is performed with allergen extracts consisting of difficult to standardize mixtures of allergenic and non-allergenic components which cannot be tailored to the individual patients' specific sensitization profile.

[0003] WO 96/27005 A2 discloses recombinant DNA molecules that code for polypeptides with antigenic properties of allergens Clah8 and Clah12.

[0004] EP-A-0714662 A2 describes peptides containing monovalent B cell epitope on an allergen molecule with which an IgE antibody specifically combines and having the effect of inhibiting the formation of a cross-linked structure by the combination of the allergen with the IgE antibody.

[0005] WO 97/05258 A2 discloses a recombinant DNA molecule which encodes a polypeptide antigen property of the co-factor-independent phosphoglycerate mutase of birch, mugwort or timothy grass pollen.

[0006] Ganglberger et al. (2000) FASEB J. 14, 2177-2184 describe allergen mimotopes for 3-dimensional epitope search and induction of antibodies inhibiting human IgE. The peptides described are derived from a phage library and show no similarity to the primary sequence of Bet v 1.

[0007] WO 97/24139 A1 discloses various peanut allergens including Ara h I and Ara h II.

[0008] It is an object of the invention to provide an advantageous pharmaceutical composition for the treatment of allergic disorders.

[0009] Surprisingly, it has been found that the administration of peptides comprising solvent-exposed amino acids of an allergen lead to the production of protective antibodies.

[0010] Therefore, the present invention relates to a method for the preparation of a pharmaceutical composition which comprises as a first step the determination which amino acids of a given allergenic protein are solvent-exposed on the surface of the allergenic protein. As a second step the method comprises the preparation of a peptide having a length of 8 to 50 amino acids wherein at least three consecutive amino acids of the peptide are identical to at least three amino acids which appear on the molecular surface of the allergenic protein within a surface patch of 500 square Angström, wherein the at least three amino acids are solvent-exposed amino acids of the allergenic protein. The peptide is then admixed with a pharmaceutically acceptable carrier (e.g., KLH) or diluent, e.g. with a buffer and/or salt solutions.

[0011] More preferably, the present invention relates to a method for the preparation of a pharmaceutical composition which comprises as a first step the determination which amino acids of a given allergenic protein are solvent-exposed on the surface of the allergenic protein. As a second step the method comprises the preparation of a peptide having a length of 8 to 50 amino acids wherein at least five consecutive amino acids of the peptide are identical to at least five consecutive amino acids of the amino acid sequence of the allergenic protein with the at least five consecutive amino acids being solvent-exposed amino acids of the allergenic protein. The peptide may is then admixed with a pharmaceutically acceptable carrier or diluent, e.g. with a buffer and/or salt solution.

[0012] In another preferred embodiment the method comprises as a further step the addition of an adjuvant. Adjuvants that are known in the art and may be used according to the invention are e.g. $Al(OH)_3$.

[0013] The peptide may be prepared by synthetic methods known in the art such as solid phase synthesis. The peptide may also be prepared by expression of recombinant DNA. A cDNA sequence encoding the peptide may be introduced into a host such as E. coli cells and expressed. Following expression, the peptides are recovered by known purification methods. The method of recombinant production of peptides may be preferred when longer peptides are used for the preparation of the pharmaceutical composition. The method of synthetically preparing the peptides is preferred when shorter peptides are employed.

[0014] In the first step of the method of the invention, it is determined which amino acids of a given allergenic protein are solvent-exposed on the surface of the allergenic protein. Any known allergenic protein may be used for the design of the peptides. An allergen source may be selected against which a high percentage of allergic patients is sensitized. Preferred allergenic proteins are derived from plants. A preferred allergen source is the birch pollen, because it is widely distributed in Europe, North America and Australia. Almost 25% of allergic patients suffer from allergy to the major birch

pollen allergen Bet v 1. Bet v 1 contains most of the IgE epitopes present in pollens of trees belonging to the *Fagales* order (birch, alder, hazel, oak, hornbeam) and in plant-derived fruit. The cDNA and amino acid sequence as well as the three-dimensional structure of Bet v 1 have been determined and recombinant Bet v 1 which equals the natural Bet v 1 wild type has been produced by recombinant DNA technology. Further allergenic proteins that can be envisaged are e.g. the major grass pollen (e.g. group 1, group 2, group 5 etc.) mite (e.g., Der p 2), bee venom (e.g., phospholipase) and animal hair dander allergens (e.g., Cat: Fel d 1).

[0015] In one embodiment the solvent-exposed amino acids are determined by examination of the three-dimensional structure of the allergenic protein. Amino acids which are on the surface of the protein and not buried in the protein are considered as being solvent-exposed. Within the meaning of this application an amino acid is solvent exposed if it has a relative solvent accessibility of at least 16%, preferably at least 25%. Solvent accessibility can be determined according to residue hydrophobicity as described in Rost and Sander (1994) *Proteins* 20, 216-226. The three-dimensional structure of the protein may be determined by X-ray crystallography or NMR.

[0016] If the amino acid sequence of the protein is known, it is also possible to perform a hydrophilicity analysis to determine which amino acids within the sequence have a high hydrophilicity index and thus a high probability of being exposed on the surface. Another possibility is to perform a "surface probability analysis" to find out which amino acids have the highest probability to be exposed to the solvent.

[0017] If neither the sequence nor the three-dimensional structure of the allergenic protein is known, it is usually necessary to clone the gene encoding the allergenic protein. This can be done by methods known in the art.

[0018] In some cases IgE epitopes have been determined for an allergen. Such epitopes may also be used for the design of peptides, since these epitopes are on the surface of the allergenic protein.

[0019] Once it is known which amino acids within the amino acid sequence of the allergenic protein are solvent-exposed, a peptide is prepared comprising at least five consecutive amino acids which are identical to at least five consecutive solvent-exposed amino acids of the allergenic protein. The peptide has a length of 8 to 50 amino acids. Preferably, the minimum length of the peptide is twelve, more preferably 15, still more preferably 18, most preferably 21 amino acids. The maximum length of a peptide preferably is 45, more preferably 40, most preferably 35 amino acids. The most preferred length of the peptide is within a range of 21 to 35 amino acids.

[0020] It is also preferred that at least eight consecutive amino acids of the peptide are identical to at least eight consecutive solvent-exposed amino acids of the allergenic protein. Most preferably at least ten consecutive amino acids of the peptide are identical to at least 10 consecutive solvent-exposed amino acids of the allergenic protein.

[0021] The peptides used for the preparation of the pharmaceutical composition may also comprise amino acids which are not derived from the allergenic protein. N-terminal or C-terminal to the at least five consecutive solvent-exposed amino acids, amino acids derived from another source or artificially designed amino acids may be added. It is preferred, however, that almost the complete sequence of the peptide is derived from the allergenic protein. In a particular embodiment, only one amino acid within the peptide sequence is not derived from the amino acid sequence of the allergenic protein. Preferably, this is the N-terminal or the C-terminal amino acid of the peptide. Most preferably, this amino acid is a cysteine residue which allows coupling of the peptide to various carrier molecules such as keyhole limpet hemocyanin (KLH).

[0022] In a further embodiment the peptide comprises at least the N-terminal or C-terminal five amino acids of the allergenic protein. The N- and C-termini often are solvent-exposed.

[0023] In another embodiment the complete peptide sequence is derived from the amino acid sequence of the allergenic protein.

[0024] Preferably, the peptides prepared according to the invention upon administration lead to the production of IgG antibodies which react with the protein from which the peptides are derived. More preferably, these IgG antibodies are "blocking antibodies" or "protective antibodies" which prevent IgE antibodies from binding to the respective allergenic protein from which the peptides are derived. A significant reduction of allergic symptoms may be achieved in this way.

[0025] Another preferred feature of the peptides is that they do not elicit an IgE response against themselves upon administration, i.e. they are hypoallergenic or non-allergenic.

[0026] The amount of the peptide administered can be 0.1 μ g to 500 μ g. The amount may vary depending on the mode of treatment. During immunotherapy treatment about 50 μ g can be administered in a volume of 100 μ l per injection. In a particular embodiment, more than one peptide is contained in the pharmaceutical composition. It is preferred that two, three, four, five or six different peptides are contained. These different peptides may be derived from the same allergen, it is also possible, however, that they are derived from different proteins. Thus, by administration of a single pharmaceutical composition an antiallergic effect with respect to different allergens can be achieved.

[0027] Another aspect of the present invention is a pharmaceutical composition as defined in the claims, prepared by a method described supra. The pharmaceutical composition contains a peptide as defined in the claims. The preferred properties and characteristics of the peptide contained in the pharmaceutical composition of the invention have been described for the method for its preparation.

[0028] The pharmaceutical composition preferably is a vaccine composition. It may be administered to allergic patients

via various routes. Preferably, the pharmaceutical composition is administered by subcutaneous injection. Other modes of administration are intramuscular or intravenous injection. Other preferred modes are sublingual, oral or nasal administration.

[0029] The systemic administration of allergens to allergic patients in the course of specific immunotherapy as practiced in the art harbors the risk of inducing life-threatening anaphylactic reactions. Furthermore, allergen extracts currently used for treatment contain a variety of allergenic and non-allergenic components which cannot be tailored according to the individual patient's sensitization profile. Improvement of safety and specificity of allergy vaccines has therefore been a long sought goal. The vaccination strategy of the present invention allows to generate safe, hypoallergenic allergy vaccines whenever the sequence, three-dimensional structure and/or IgE epitopes of an allergen molecule is available.

[0030] Birch pollen allergic patients IgE antibodies primarily recognize conformational Bet v 1 IgE epitopes. Surprisingly, it has been found by the inventors that hypoallergenic peptides can also be synthesized for allergens containing continuous (i.e. sequential, linear) IgE epitopes. The inventors succeeded in producing a set of hypoallergenic peptides for the major timothy grass pollen allergen, Phl p 1, which contains continuous epitopes.

[0031] Unexpectedly, peptide immunization of mice and rabbits with the peptides used in the present invention resulted in the production of IgG antibodies specific for the complete Bet v 1 wild type allergy although the peptides were rather short in length (25 to 32 amino acids) and despite the fact that certain of the peptides (P1-P3) lacked Bet v 1 specific T cell epitopes.

[0032] It was also found that peptide-induced IgG antibodies crossreacted with Bet v 1 related allergens from alder, hazel and hornbeam pollen, indicating that peptide immunotherapy may also protect against allergy to allergen sources containing a related allergen.

[0033] The peptide immunotherapy approach of the present invention is different from previously made attempts to use T cell epitope containing peptides for a modulation of allergen specific T cell responses. While administration of free, soluble T cell epitope containing peptides either aims at inducing tolerance or at switching towards Th1-responses, the inventors used adjuvant-bound, surface-exposed peptides for the focusing of blocking IgG antibodies to solvent-exposed allergen domains which can be principle targets for IgE antibodies. Since preferably adjuvant-bound peptides are used for the induction of blocking antibodies, it is unlikely that injection of peptides even when they contain T cell epitopes will cause T cell-mediated systemic reaction as has been reported for the major cat allergen, Fel d 1.

[0034] Hypoallergenic surface-exposed allergen peptides can be produced in a well controlled manner and represent therapeutic reagents which can be easily purified and standardized. Such peptides can be administered in high doses with low risk of inducing anaphylactic side effects, particularly when they are bound to adjuvants. Administration of high peptide doses is expected to induce a vigorous production of IgG antibodies which is accompanied by a Th1 immune response. The approach of selecting surface-exposed peptides according to the three-dimensional structure of important crossreactive allergens is of general applicability and allows a rational design of safe peptide-based vaccines for many other forms of type I allergy.

Description of the tables and figures:

[0035] Table 1 summarizes the characteristics of non-allergenic Bet v 1-derived synthetic peptides. Position, sequence, length, molecular weight, isoelectric point, residual fold, presence of T cell epitopes and AGADIR prediction of six Bet v 1-derived peptides are displayed.

[0036] Table 2 shows the serological characterization of 60 birch pollen allergic patients and a non-allergic control individual (NHS). Sex, age, total serum IgE levels (kU/L), birch pollen extract-specific IgE levels (kUA/L) are displayed. IgE antibodies specific for rBet v 1 and the Bet v 1-derived peptides were measured by ELISA and OD (optical densities) are shown. Dashes indicate reactivities below the cut-off level (OD = 0.050).

[0037] Table 3 summarizes immediate type skin reactions to complete rBet v 1 and Bet v 1-derived peptides. Five birch pollen allergic patients (#1-5) and an allergic individual without birch pollen allergy (#6) were tested. The mean wheal diameters (mm) are displayed for different concentrations of rBet v 1, for birch pollen extract, histamine, the individual peptides and a mixture of the six peptides.

[0038] Table 4. IgG reactivity of anti-peptide antisera with complete rBet v 1 wildtype. The upper panel shows the mean IgG reactivities (OD values) to rBet v 1 in 4 serum samples (bleeding 1: preimmune serum; bleedings 2-4: serum samples obtained in monthly intervals) of mice which were immunized with KLH-coupled peptides (P1-KLH to P6-KLH). In the lower panel, IgG reactivities to rBet v 1 are shown for 3 serum samples (bleeding 1: preimmune serum; bleedings 2-3: serum samples collected in monthly intervals) obtained from 6 rabbits of which each was immunized with one of the KLH-conjugated peptides.

[0039] Table 5. Crossreactivity of anti-Bet v 1 peptide antisera with natural allergens from birch, alder, hazel and hornbeam. IgG reactivities of peptide antisera (anti-P1 to anti-P6) to pollen extracts from birch, alder, hazel and hornbeam (mean OD values +/- SEM) are displayed for 6 groups of mice which were immunized with Bet v 1 peptides (P1-P6).

[0040] Table 6. Rabbit anti-Bet v 1 peptide antisera inhibit serum IgE binding of birch pollen allergic patients to rBet

v 1. The percentage inhibition of IgE binding to complete rBet v 1 obtained with the individual anti-peptide antisera (anti-P1 to anti-P6) and a mixture of the anti-peptide antisera (anti-P1 to anti-P6) are displayed for sera from 60 birch pollen allergic patients. The bottom line summarizes the mean percentages of inhibition obtained for all serum samples.

5 Figure 1 shows ¹H ¹D-spectra of the amide regions of the six peptides. The spectra were recorded at 20°C on a 600 MHz spectrometer.

10 Figure 2 shows induction of histamine release from basophils of a birch pollen allergic patient. Granulocytes of a birch pollen allergic patient were incubated with various concentrations (x-axis) of rBet v 1 wildtype, Bet v 1 derived peptides (P1, P2, P3, P4, P5, P6) and an equimolar mix of the 6 peptides (P1-P6). The percentage of histamine released into the cell-free culture supernatant is displayed on the y-axis.

15 Figure 3 shows the reactivity of peptide antisera with nitrocellulose-blotted natural Bet v 1. Nitrocellulose-blotted birch pollen extract was incubated with sera from six peptide-immunized mice (lanes i; panels 1-6: peptides 1-6). Lanes p show the corresponding preimmune sera. The position of nBet v 1 (17 kDa) is indicated.

[0041] The following examples further illustrate the invention.

Example 1: Characteristics of six synthetic peptides spanning the surface of the Bet v 1 allergen

20 [0042] Six Bet v 1-derived peptides of a length between 25 (P1: 2809.3 Dalton) to 32 (P5: 3556.1 Dalton) amino acids spanning solvent-exposed areas of the Bet v 1 molecule were selected according to the three-dimensional structure of Bet v 1 determined by x-ray crystallography and NMR (Gajhede, M. et al. (1996 X-ray and NMR structure of Bet v 1, the origin of birch pollen allergy. *Nat. Struct. Biol.* 3, 1040-1045).

25 [0043] The isoelectric points calculated for the 6 peptides ranges from 4.70 (P6) to 8.63 (P2) (Table 1). Three of the peptides (P4, P5, P6) comprise sequence motifs which were found to stimulate Bet v 1-specific human T cells whereas the other three peptides (P1, P2, P3) do not contain Bet v 1-specific T cell epitopes (Table 1).

30

35

40

45

50

55

Table 1 Characteristics of non-allergenic Bet v 1 - derived synthetic peptides

	Position aa	Sequence	Number of aa	Molecular Weight	Isoelectric Point (pI)	Fold (NMR)	T cell epitope	Agadir %
Peptide 1 SEQ ID NO.1	1-24	MGVFN YETETTSVI PAARLFKAFIC	25	2809.3	6.28	-	-	0.86
Peptide 2 SEQ ID NO.2	30-59	LFPKVAPQAISSE NIEGN GGP GTIKKI SFC	31	3202.7	8.63	-	-	0.56
Peptide 3 SEQ ID NO.3	50-79	CGPGTIKKISFPEG FPFKYVKDRVDEV DHTN	31	3525	7.00	-	-	0.5
Peptide 4 SEQ ID NO.4	110-139	DGGSILKISNKYHT KGDHEVKAEQVKA SKEC	31	3400.8	8.44	-	+	0.85
Peptide 5 SEQ ID NO.5	130-160	CKAEQVKASKEMG ETLLRAVESYLLAH SDAYN	32	3556.1	5.49	+/-	+	12.09
Peptide 6 SEQ ID NO.6	75-104	CVDHTNFKYNYSVIEGGPI GDTLEK I SNEIK	31	3484.9	4.70	-	+	1

a) Peptide synthesis

[0044] Peptides were synthesized using Fmoc (9-fluorenylmethoxycarbonyl)-strategy with HBTU (2-(1H-Benzotriazol-1-yl) 1,1,3,3-tetramethyluronium hexafluorophosphate)-activation (0.1 mmol small-scale cycles) on the Applied Biosystems peptide synthesizer Model 433A (Foster City, CA). Preloaded PEG-PS (polyethyleneglycol polystyrene) resins (0.15-0.2 mmol/g loading) (PerSeptive Biosystems, Warrington, UK) were used as solid phase to build up the peptides. Chemicals were purchased from Applied Biosystems. Coupling of amino acids was confirmed by conductivity monitoring in a feedback control system. One cysteine residue was added to each peptide to facilitate coupling of the peptides to carriers. Peptides were cleaved from the resins with a mixture of: 250 µl dest. water, 250 µl Triisopropylsilan (Fluka, Buchs, Switzerland), 9.5 ml TFA for 2h and precipitated in tert-Butylmethylether (Fluka, Buchs, Switzerland). The identity of the peptides was checked by mass-spectrometry and they were purified to >90% purity by preparative HPLC (PiChem, Graz, Austria).

b) Secondary structure analysis

[0045] The six Bet v 1-derived peptides resemble almost all primary structure elements of the Bet v 1 allergen but according to AGADIR prediction and NMR analysis lack secondary structure. The AGADIR program is based on the helix-to-coil transition theory and estimates the propensity of a sequence to adopt an alpha-helical conformation irrespective of other secondary structures (Munoz, V. & Serrano, L. (1994) Elucidating the folding problem of helical peptides using empirical parameters. *Nat. Struct. Biol.* 1, 399-409). It has been shown to correctly estimate the helical behavior of a database of peptides as compared to CD (circular dichroism) and NMR data. Because it calculates the helical tendency for each amino acid in a given sequence, AGADIR may predict both the average helical population of a peptide and the propensity to populate the helical conformation for each residue. Results from these calculations are reported in Table 1. Of the peptides, only P5 shows an appreciable tendency to form a helical conformation.

c) NMR analysis

[0046] NMR was then used to check for residual secondary or tertiary structure. Compared to other spectroscopical approaches, this technique has the advantage that it provides positional information along the sequence.

[0047] Peptides were examined for tertiary and/or secondary structure by one and two-dimensional nuclear magnetic resonance (NMR) on a UNITY VARIAN 600 MHz spectrometer equipped with z-shielded gradient coils using 0.8-1.0 mM samples in 90% H₂O/10% D₂O, 20 mM potassium phosphate buffer pH 7.0, at 20°C. Water suppression was achieved with WATERGATE. Mixing times used were 70 ms and 150 or 200 ms for TOCSY and NOESY experiments, respectively.

[0048] The appearance of the 1D experiments clearly shows absence of tertiary structure in all of the six peptides: all the resonances have poor spectral dispersion as expected for a mixture of amino acids or peptides without tertiary fold (Figure 1). The absence of ring current shifted peaks in the region of the spectra below 0.5 ppm typical of well folded proteins is also a feature common to the spectra of the Bet v 1 peptides (data not shown). However, the spectrum of P5 contains unusually broad resonances suggesting aggregation of this peptide. This peptide (P5), that in the Bet v 1 structure spans the whole C-terminal helix probably tends to form helical structures that aggregate in water solution.

Example 2: Surface exposed Bet v 1-derived peptides lack IgE binding capacity and allergenic activity

a) IgE binding capacity

[0049] The IgE binding capacity of the six Bet v 1-derived peptides was compared with that of the complete rBet v 1 wildtype allergen using sera from 60 birch pollen allergic patients (Table 2).

Characterization of allergic patients and patients' sera:

[0050] Birch pollen allergic patients suffering from allergic rhinoconjunctivitis and/or asthma to birch pollen and related allergens were selected according to case history indicative for seasonal birch pollinosis and characterized by skin prick testing with birch pollen extract and serological CAP RAST (Pharmacia Diagnostics, Uppsala, Sweden) testing. Total IgE levels in the sera were determined by CAP RIST measurements (Pharmacia). IgE antibodies specific for rBet v 1 were determined by ELISA (Niederberger et al. (1998) IgE antibodies to recombinant pollen allergens (Phl p 1, Phl p 2, Phl p 5, and Bet v 2) account for a high percentage of grass pollen-specific IgE. *J. Allergy Clin. Immunol.* 101, 258-264). Sera from 60 birch pollen allergic patients and a non-atopic individual were used for IgE competition studies. The birch pollen allergic patients group consisted of 25 males and 35 females with a mean age of 37 years (ranging from 19-60 years) (Table 2).

Result:

[0051] Total IgE levels and IgE levels specific for birch pollen extract ranged from 24-2000kU/L (mean: 300) and 3-100kUA/L (mean: 47), respectively. All patients contained rBet v1-specific IgE antibodies ranging between 0.081-2.530 OD units (mean: 1.416 OD units) (Table 2). When tested for IgE reactivity to ELISA plate bound peptides (P1-P6) no serum showed significant IgE reactivity above the cut off level determined with serum from a non-atopic person (NHS) (Table 2).

Table 2 Serological characterization of birch pollen allergic patients

	Patient	sex	age	total IgE (kU/L)	IgE birch (kUA/L)	IgE (OD)						
						rBet v 1	P1	P2	P3	P4	P5	P6
10	1	w	27	204	27.2	0.829	-	-	-	-	-	-
	2	w	54	617	69	1.380	-	-	-	-	-	-
15	3	m	55	592	100.0	2.144	-	-	-	-	-	-
	4	w	35	109	34.9	0.701	-	-	-	-	-	-
	5	w	35	41.5	21.1	0.344	-	-	-	-	-	-
	6	m	27	1642	77.0	2.300	-	-	-	-	-	-
20	7	m	24	2000	96.9	2.245	-	-	-	-	-	-
	8	m	60	235	49.6	0.950	-	-	-	-	-	-
	9	w	56	35.4	8.7	0.196	-	-	-	-	-	-
	10	w	32	46.6	5.8	0.097	-	-	-	-	-	-
25	11	w	23	121	23.4	0.428	-	-	-	-	-	-
	12	m	24	122	176	0.271	-	-	-	-	-	-
	13	w	21	238	55.4	0.281	-	-	-	-	-	-
	14	w	26	124	41.7	0.704	-	-	-	-	-	-
	15	m	38	95.3	41.6	0.606	-	-	-	-	-	-
	16	m	37	147	11.7	0.215	-	-	-	-	-	-
30	17	m	20	206	5.0	0.622	-	-	-	-	-	-
	18	w	43	135	50.9	0.683	-	-	-	-	-	-
	19	w	39	267	79.8	1.166	-	-	-	-	-	-
	20	w	50	28.1	13.2	0.249	-	-	-	-	-	-
35	21	m	60	36.9	26.1	0.461	-	-	-	-	-	-
	22	w	44	59.7	23.5	0.258	-	-	-	-	-	-
	23	m	34	153	30.8	1.555	-	-	-	-	-	-
	24	w	35	84.6	16.4	0.237	-	-	-	-	-	-
40	25	w	55	152	39.2	0.5	-	-	-	-	-	-
	26	m	53	31.5	8.5	0.191	-	-	-	-	-	-
	27	m	41	89.9	33.0	0.535	-	-	-	-	-	-
	28	w	37	61.6	39.8	0.492	-	-	-	-	-	-
	29	w	50	65.2	10.0	0.154	-	-	-	-	-	-
	30	m	41	260	87.5	0.081	-	-	-	-	-	-
45	31	m	52	450	73.5	1.168	-	-	-	-	-	-
	32	w	34	70.2	13.7	0.195	-	-	-	-	-	-
	33	w	30	293	9.9	0.106	-	-	-	-	-	-
	34	m	24	309	32.0	0.377	-	-	-	-	-	-
50	35	w	39	273	84.6	0.601	-	-	-	-	-	-
	36	m	39	257	31.1	0.306	-	-	-	-	-	-
	37	m	49	208	68.9	0.227	-	-	-	-	-	-
	38	w	46	208	79.7	1.571	-	-	-	-	-	-
55	39	m	53	125	51.3	0.620	-	-	-	-	-	-
	40	w	23	121	5.7	0.084	-	-	-	-	-	-
	41	w	29	490	100.0	1.193	-	-	-	-	-	-
	42	w	51	370	100.0	1.119	-	-	-	-	-	-

EP 1 219 299 B1

(continued)												
	Patient	sex	age	total IgE (kU/L)	IgE birch (kUA/L)	IgE (OD)						
						rBet v 1	P1	P2	P3	P4	P5	P6
5	43	m	19	1234	100.0	0.383	-	-	-	-	-	-
	44	w	23	589	100.0	2.386	-	-	-	-	-	-
	45	w	25	278	54.5	0.289	-	-	-	-	-	-
	46	w	35	184	47.5	0.892	-	-	-	-	-	-
	47	m	23	207	30.2	0.381	-	-	-	-	-	-
10	48	w	33	200	47.5	1.422	-	-	-	-	-	-
	49	w	24	297	23.4	0.379	-	-	-	-	-	-
	50	w	26	24	11.5	0.220	-	-	-	-	-	-
	51	m	29	543	100.0	0.307	-	-	-	-	-	-
	52	w	48	178	17.5	0.208	-	-	-	-	-	-
15	53	w	33	152	100.0	1.256	-	-	-	-	-	-
	54	w	36	187	64.7	1.310	-	-	-	-	-	-
	55	m	26	2000	100.0	2.489	-	-	-	-	-	-
	56	m	35	1733	100.0	2.530	-	-	-	-	-	-
	57	w	28	270	62.5	0.741	-	-	-	-	-	-
20	58	m	41	80	3.0	0.072	-	-	-	-	-	-
	59	m	49	41.9	12.9	0.168	-	-	-	-	-	-
	60	w	35	84.6	16.4	0.237	-	-	-	-	-	-
	NHS	W	40	5	<0.35	0.048	-	-	-	-	-	-
25												

b) Histamine release

[0052] Next the Bet v 1-derived peptides were compared with complete rBet v 1 for their capacity to induce histamine release from basophilic granulocytes of 3 birch pollen allergic individuals.

Basophil histamine-release assay:

[0053] Granulocytes were isolated from heparinized blood samples of 3 birch pollen allergic individuals by dextran sedimentation (Valent, P. et al. (1989) Interleukin 3 activates human blood basophils via high-affinity binding sites. *Proc. Natl. Sci. USA* 86, 5542-5547). Cells were incubated with increasing concentrations (10^{-3} - $10\mu\text{g/ml}$) of each peptide, an equimolar mix of the 6 peptides, and, for control purposes, with complete rBet v 1. Histamine released in the cell-free culture supernatant was measured by radioimmunoassay (Immunotech, Marseille, France). Total histamine was determined after freezing and thawing of cells. Results are displayed as mean values of triplicate determinations.

Result:

[0054] As exemplified in Figure 2 it was found that none of the peptides induced release of histamine up to concentrations of $10\mu\text{g/ml}$ whereas complete rBet v 1 wildtype induced a dose-dependent release of histamine with a maximum release already at a concentration of 1 ng/ml .

c) Allergenic activity in vivo

[0055] *In vivo* testing in birch pollen allergic patients confirmed the lack of allergenic activity of the Bet v 1-derived peptides.

Skin Prick testing of allergic patients:

[0056] The *in vivo* allergenic activity of the peptides was studied by skin prick testing (SPT) in 5 birch pollen allergic patients and a non-atopic grass pollen allergic patient without birch pollen allergy. SPTs were performed on the individuals forearms as described (Vrtala, S. et al. (1997) *J. Clin Invest.* 99, 1673-168; Dreborg S: (1989) *J. Am. Acad. Dermatol.* 21, 820-821). Twenty microliter aliquots containing 6 concentrations of complete rBet v 1 ($0.78\mu\text{g/ml}$, $1.56\mu\text{g/ml}$, $3.12\mu\text{g/ml}$, $6.25\mu\text{g/ml}$, $12.5\mu\text{g/ml}$, $25\mu\text{g/ml}$), $100\mu\text{g/ml}$ of each peptide and an equimolar peptide mix were applied. In

addition standardized skin prick solutions (birch pollen extract and histamine) (Allergopharma, Reinbek, Germany) were tested. Reactions were recorded 20 minutes after SPT by photography and by transferring the ballpoint pen-surrounded wheal area with a scotch tape to paper. The mean wheal diameter (Dm) was calculated by measuring the maximal longitudinal and transversal diameter and dividing their sum by 2.

Result:

[0057] Skin prick tests were performed in 5 birch pollen allergic patients and an allergic individual without birch pollen allergy with rBet v 1 and Bet v 1-derived peptides (Table 3). None of the Bet v 1-derived peptides induced any immediate skin reaction when applied at a concentration of 100 µg/ml or as peptide mixture containing 100 µg/ml of each peptide (Table 3). Complete rBet v 1 wildtype already induced immediate type skin reactions in three patients (Table 3: #1-3) at a concentration of <0.78 µg/ml. A concentration of 3.12 µg/ml was sufficient to induce wheal reactions in all five birch pollen allergic patients (Table 3). All birch pollen allergic patients displayed immediate skin reactions to birch pollen extract. The grass pollen allergic patient without birch pollen allergy (Table 3: #6) showed no reactions to birch pollen extract, rBet v 1 and Bet v 1-derived peptides (Table 3) but exhibited a wheal reaction after prick testing with timothy grass pollen extract (5.5 mm mean diameter) (data not shown). All individuals reacted after testing with histamine, used as a positive control (Table 3). Since local and even systemic late phase reactions to allergens and in particular to allergen-derived non-allergenic peptides have been reported we checked the application sites 24 - 48 hours after skin prick testing but found no evidence for peptide-induced late phase reactions in any of the tested individuals (data not shown).

55
50
45
40
35
30
25
20
15
10
5

Table 3 *Induction of immediate skin reaction with rBet v 1 and Bet v 1-derived peptides*

mean wheal diameter

Individual	rBet v 1 µg/ml								birch	histamine	100µg/ml						
	0.19	0.39	0.78	1.56	3.12	6.25	12.5	25			P1	P2	P3	P4	P5	P6	P1-P6 mix
1	0	2.5	4	4.5	5	6	13	n.d.	5.5	5	0	0	0	0	0	0	0
2	0	2	3.5	5	5.5	7	7.5	8	7.5	6	0	0	0	0	0	0	0
3	n.d.	n.d.	2	3	4	6	6.5	8	8	7	0	0	0	0	0	0	0
4	n.d.	n.d.	0	0	2	3	4	4.5	8	5	0	0	0	0	0	0	0
5	n.d.	n.d.	0	2	2.5	4	11	5	9	5.5	0	0	0	0	0	0	0
6	n.d.	n.d.	0	0	0	0	0	0	0	4	0	0	0	0	0	0	0

Example 3: Immunization with Bet v 1-derived peptides induces IgG antibodies reactive with complete Bet v 1 and Bet v 1-related plant allergens

a) Induction of IgG antibodies

[0058] In order to test whether immunization with Bet v 1-derived peptides will induce IgG antibodies that react with the complete Bet v 1 molecule and Bet v 1-related allergens mice and rabbits were immunized with the KLH-conjugated peptides using different adjuvants (Aluminiumhydroxide, Freund's adjuvant).

Immunization of mice and rabbits:

[0059] HPLC-purified peptides were coupled to KLH (keyhole limpet hemocyanin, MW 4.5×10^3 - 1.3×10^7 , Pierce, Rockford, IL) according to the manufacturers advice and purified using a Conjugation Kit (Sigma, St. Louis). KLH-conjugated peptides were used to immunize mice and rabbits using $\text{Al}(\text{OH})_3$ and CFA as adjuvant, respectively. Six week old female BALB/c mice were purchased from Charles River (Kißlegg, Germany). Animals were maintained in the animal care unit of the Department of Pathophysiology, University of Vienna, according to the local guidelines for animal care. Groups of 5 animals were immunized with conjugated and unconjugated peptides adsorbed to AluGel (Serva, Heidelberg, Germany). Each mouse received four 100 μ l-injections (20 μ g peptide or peptide-KLH conjugate adsorbed to 15 μ g of $\text{Al}(\text{OH})_3$) subcutaneously in the neck in four week intervals. Shortly before each injection and four weeks after the last immunization, mice were bled from the tail veins. Sera were stored at -20°C until analysis.

[0060] In parallel six rabbits were immunized with one of the peptide-KLH conjugates (200 μ g/injection) using Freund's complete and incomplete adjuvants (Charles River, Kißlegg, Germany). Serum samples were obtained in four week intervals.

Crossreactivity of anti-peptide antisera:

[0061] Reactivity of peptide-induced IgG antibodies to rBet v 1, natural Bet v 1 and Bet v 1-related pollen and plant food allergens was studied by ELISA and immunoblotting. For ELISA detection, plates (Nunc Maxisorp, Roskilde, Denmark) were coated with pollen allergen extracts (100 μ g/ml: *Betula verrucosa*, *Alnus glutinosa*, *Corylus avellana* and *Carpinus betulus*) or purified allergens (5 μ g/ml: rBet v 1). ELISA plates were washed and blocked as described (Niederberger, V. et al. (1998). *J. Allergy Clin. Immunol.* 101, 258-264) and subsequently incubated with mouse anti-peptide antisera diluted 1:500. Bound mouse IgG₁ antibodies were detected with a monoclonal rat anti mouse IgG₁ antibody (Pharmingen, San Diego, CA) diluted 1:1000 followed by a 1:2000 diluted HRP-coupled sheep anti-rat Ig antiserum (Amersham Pharmacia Biotech, Uppsala, Sweden). Results represent means of duplicate determination with an error of <5% and are displayed as means $\pm$ SEM for the group of 5 mice. ELISAs with rabbit anti-peptide antisera were performed as described for the mouse sera with the exception that rabbit sera were diluted 1:1000 and bound rabbit antibodies were detected with a HRP-coupled goat anti-rabbit Ig antiserum (Jackson Immunresearch, Pennsylvania).

[0062] Immunoblotting experiments were performed with nitrocellulose-blotted allergens. Pollen extracts and plant food extracts were separated by 12.5% preparative SDS-PAGE (12 μ g/cm gel) and blotted onto nitrocellulose membranes (Schleicher & Schuell, Dassel, Germany). Nitrocellulose membranes were exposed to 1:1000 diluted rabbit sera (pre-immune, immune sera) and in certain experiments, with sera from mice which were immunized with unconjugated $\text{Al}(\text{OH})_3$ -adsorbed peptides. Bound rabbit antibodies were detected with a 1:2000 diluted ^{125}I -labeled donkey anti-rabbit IgG antiserum (Amersham Pharmacia Biotech). Bound mouse antibodies were detected with a 1:2500 diluted rabbit anti-mouse IgG antiserum (Jackson Immunresearch) followed by 1:2000 diluted ^{125}I -labeled donkey anti-rabbit IgG antiserum (Amersham).

Result:

[0063] Table 4A shows the development of IgG anti-rBet v 1 antibody responses in groups of mice which were immunized with $\text{Al}(\text{OH})_3$ -adsorbed, KLH-conjugated peptides. All 6 peptides induced IgG anti-rBet v 1 antibody responses which could be detected 4-8 weeks after the first immunization (Table 4A: bleeding 2-3). Peptide-induced IgG anti-rBet v 1 responses constantly increased during the immunization. Using unconjugated peptides, anti-rBet v 1 IgG antibody responses could be induced, albeit of a lower magnitude (data not shown).

[0064] Similar results were obtained after immunization of rabbits with KLH-conjugated peptides which were administered s. c. together with Freund's adjuvant (Table 4B). All 6 peptides induced IgG anti-rBet v 1 antibody responses which increased during the course of immunization. Peptide-induced mouse IgG antibodies reacted also with nitrocellulose-blotted pollen-derived natural Bet v 1 whereas no signals were obtained with the corresponding preimmune sera (Figure 3).

Table 4 IgG reactivity of mouse anti-peptide antisera with rBet v 1

bleeding	P1-KLH	P2-KLH	P3-KLH	P4-KLH	P5-KLH	P6-KLH
1	0.040	0.040	0.040	0.040	0.042	0.040
2	0.057	0.335	1.150	0.433	0.872	0.152
3	0.289	0.858	1.380	1.177	1.621	0.705
4	0.394	1.182	2.025	1.965	2.274	1.337

IgG reactivity of rabbit anti-peptide antisera with rBet v 1

bleeding	P1-KLH	P2-KLH	P3-KLH	P4-KLH	P5-KLH	P6-KLH
1	0.170	0.044	0.046	0.044	0.033	0.051
2	0.642	0.702	0.511	0.501	0.917	0.997
3	0.751	1.201	0.961	0.926	1.503	0.751

b) Reactivity of induced IgG antibodies with Bet v 1-related plant allergens

[0065] Next it was investigated whether anti-Bet v 1 peptide antisera crossreact with Bet v 1-related allergens present in pollens of alder, hazel and hornbeam.

Allergen extracts:

[0066] Pollen from trees of the order *Fagales* (birch: *Betula verrucosa*; alder: *Alnus glutinosa*; hazel: *Corylus avellana*; hornbeam: *Carpinus betulus*) were purchased from Allergon (Välinge, Sweden). Aqueous pollen extracts were prepared and checked for quantity and quality of proteins by SDS-PAGE and Coomassie blue staining (Vrtala, S. et al. (1993) *Int. Arch. Allergy Immunol.* 102, 160-169). Purified recombinant Bet v 1 was purchased from BIOMAY (Linz, Austria).

Result:

[0067] Table 5 shows that almost all anti-Bet v 1 peptide antisera contained IgG antibodies to natural Bet v 1 and Bet v 1-related *Fagales* pollen allergens. Only anti-peptide 4 antibodies showed weak reactivity with the major hazel pollen allergen, Cor a 1, and anti-peptide 6 antibodies reacted weakly with the major alder and hazel pollen allergen, Aln g 1 and Cor a 1, respectively.

Table 5 Crossreactivity of anti-Bet v 1 peptide antisera with natural allergens from birch, alder, hazel and hornbeam

	anti-P1	anti-P2	anti-P3	anti-P4	anti-P5	anti-P6
birch	0.301±0.198	0.841±0.465	1.330±0.172	1.135±0.312	1.945±0.329	0.898±0.434
alder	0.307±0.232	0.875±0.453	0.619±0.314	0.330±0.265	1.613±0.477	0.084±0.032
hazel	0.217±0.200	0.350±0.297	0.071±0.018	0.073±0.022	1.239±0.397	0.060±0.007
hornbeam	0.253±0.209	0.559±0.373	0.183±0.117	0.233±0.145	1.369±0.234	0.449±0.490

Example 4: Anti-peptide antisera inhibit the binding of birch pollen allergic patients IgE to complete Bet v 1

[0068] The capacity of anti-Bet v 1 peptide antibodies to inhibit the binding of allergic patients IgE to complete rBet v 1 was studied by ELISA competition using sera from 60 birch pollen allergic patients' (Tables 2 and 6).

ELISA detection of peptide and rBet v 1-specific IgE antibodies:

[0069] inhibition of allergic patients IgE binding to rBet v 1 by peptide-induced IgG. ELISA plates (Nunc Maxisorp, Roskilde, Denmark) were coated with Bet v 1-derived peptides (5µg/ml) or rBet v 1 as control (5µg/ml), washed and blocked as described in Example 3. Subsequently, plates were incubated with 1:3 diluted sera from 60 birch pollen allergic patients and from a non-atopic individual overnight at 4°C. Bound IgE antibodies were detected with a 1:1000 diluted alkaline-phosphatase-coupled mouse monoclonal anti-human IgE antibody (Pharmingen, San Diego, CA).

[0070] The ability of peptide-induced rabbit Ig to inhibit the binding of allergic patients' IgE to complete rBet v 1 was investigated by ELISA competition assay as described (Denepoux et al. (2000) *FEBS Lett.* 465, 39-46). ELISA plates were coated with rBet v 1 (1 µg/ml) and preincubated either with a 1:250 dilution of each of the anti-peptide antisera (anti-P1-anti-P6), a mixture containing equal volumes of the anti-peptide antisera and, for control purposes, with the corresponding preimmune sera and a mixture of the preimmune sera as described. After washing plates were incubated with 1:3 diluted sera from 60 birch pollen allergic patients and bound IgE antibodies were detected with the alkaline phosphatase-coupled mouse monoclonal anti-human IgE antibody (Pharmingen). The percentage inhibition of IgE binding achieved by preincubation with the anti-peptide antisera was calculated as follows: %inhibition of IgE binding = $100 - OD_i/OD_p \times 100$. OD_i and OD_p represent the extinctions after preincubation with the rabbits immune and preimmune serum, respectively.

Result:

[0071] All birch pollen allergic patients tested exhibited immediate type skin reactions to birch pollen extract. For all but three sera (serum #9, 10, 38), preincubation with peptide-induced rabbit IgG inhibited considerably the binding of human IgE to Bet v 1. The strongest inhibition of IgE binding was observed after preincubation with anti-P2 (68% average inhibition), anti-P6 (60% average inhibition), anti-P3 (53.6% average inhibition) and anti-P5 (41.6% average inhibition). Anti-P4 antibodies exhibited a lower capacity to inhibit serum IgE binding to Bet v 1 (average inhibition 28.5%) and anti-P1 antibodies raised against the N-terminus of Bet v 1 weakly inhibited IgE binding to Bet v 1 (average inhibition 11.4%). Interestingly, a mixture of anti-P1 to anti-P6 antibodies did not exhibit a stronger inhibition of IgE binding (average inhibition 56.9%) than antibodies induced against single peptides (e. g., anti-P2-, anti-P6 antibodies) (Table 6). No association between the levels of rBet v 1-specific IgE in the sera and the degree of inhibition of IgE binding by the anti-peptide antisera was observed.

Table 6 Rabbit anti-Bet v 1 peptide antisera inhibit serum IgE binding of birch pollen allergic patients to rBet v 1

		% inhibition						
	Patient	anti-P1	anti-P2	anti-P3	anti-P4	anti-P5	anti-P6	anti-P1-6
	1	16	82	76	59	72	80	86
	2	0	88	78	49	70	76	85
	3	16	92	86	50	76	85	92
	4	39	66	47	28	38	60	67
	5	0	75	44	60	31	62	48
	6	0	88	75	38	59	76	90
	7	20	92	80	40	68	86	93
	8	17	68	69	44	66	65	67
	9	13	0	0	0	0	0	0
	10	0	0	0	0	0	0	0
	11	34	70	55	0	13	56	62
	12	10	44	41	24	27	28	0
	13	18	54	62	31	33	53	0
	14	12	72	51	0	52	61	48
	15	0	51	6	18	14	51	6
	16	13	81	56	27	52	72	69
	17	13	87	75	48	46	80	86
	18	15	89	73	44	65	85	90
	19	46	82	35	0	51	61	92
	20	0	85	61	34	59	77	84
	21	14	69	54	13	44	56	53
	22	0	72	62	35	52	65	71
	23	17	66	74	51	69	81	87
	24	11	68	39	19	43	51	37
	25	15	83	73	48	67	77	78
	26	8	51	37	31	29	51	0

EP 1 219 299 B1

(continued)

% inhibition

	Patient	anti-P1	anti-P2	anti-P3	anti-P4	anti-P5	anti-P6	anti-P1-6
5	27	0	79	65	33	60	78	84
	28	5	72	59	17	45	70	75
	29	0	66	44	21	38	41	33
	30	0	40	18	0	0	31	13
	31	6	79	80	25	66	46	82
10	32	7	79	67	38	51	68	78
	33	0	22	21	4	20	21	0
	34	0	62	44	7	47	51	65
	35	0	72	57	16	45	64	61
15	36	13	69	52	12	46	57	54
	37	2	22	23	7	12	14	0
	38	0	0	0	0	0	0	0
	39	0	89	68	0	50	80	87
	40	9	78	63	32	61	79	66
20	41	16	90	80	64	49	87	91
	42	11	89	76	49	45	79	86
	43	25	80	69	28	3	81	74
	44	0	67	28	17	8	50	62
25	45	23	82	73	37	76	76	80
	46	23	84	85	47	76	82	86
	47	5	73	64	18	32	60	42
	48	7	82	65	38	20	71	83
	49	2	74	68	38	56	63	48
30	50	13	64	42	35	29	62	58
	51	20	47	42	15	0	28	27
	52	14	55	38	35	15	48	13
	53	7	87	69	52	34	80	85
35	54	37	82	81	71	79	78	74
	55	16	82	52	29	39	72	85
	56	21	87	74	51	61	81	88
	57	30	75	60	45	55	66	65
	58	4	23	0	0	0	28	9
40	59	7	62	43	17	41	59	13
	60	11	68	39	19	43	51	37
	mean	11.4	68.0	53.6	28.5	41.6	60.0	56.9

45 SEQUENCE LISTING

[0072]

50 <110> SHAN-Beteiligungsgesellschaft m.b.H.

<120> Allergy vaccines and their preparation

<130> solvent-exposed peptides

55 <140>

<141>

<160> 6

EP 1 219 299 B1

<170> PatentIn Ver. 2.1

<210> 1

<211> 25

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: solvent-exposed peptide

<400> 1

Met Gly Val Phe Asn Tyr Glu Thr Glu Thr Thr Ser Val Ile Pro Ala
1 5 10 15

Ala Arg Leu Phe Lys Ala Phe Ile Cys
20 25

<210> 2

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: solvent-exposed peptide

<400> 2

Leu Phe Pro Lys Val Ala Pro Gln Ala Ile Ser Ser Val Glu Asn Ile
1 5 10 15

Glu Gly Asn Gly Gly Pro Gly Thr Ile Lys Lys Ile Ser Phe Cys
20 25 30

<210> 3 '

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: solvent-exposed peptide

<400> 3

Cys Gly Pro Gly Thr Ile Lys Lys Ile Ser Phe Pro Glu Gly Phe Pro
1 5 10 15

Phe Lys Tyr Val Lys Asp Arg Val Asp Glu Val Asp His Thr Asn
20 25 30

<210> 4

<211> 31

EP 1 219 299 B1

<212> PRT

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence: solvent-exposed peptide

<400> 4

10 Asp Gly Gly Ser Ile Leu Lys Ile Ser Asn Lys Tyr His Thr Lys Gly
1 5 10 15

15 Asp His Glu Val Lys Ala Glu Gln Val Lys Ala Ser Lys Glu Cys
20 25 30

<210> 5

<211> 32

20 <212> PRT

<213> Artificial Sequence

<220>

25 <223> Description of Artificial Sequence: solvent-exposed peptide

<400> 5

30 Cys Lys Ala Glu Gln Val Lys Ala Ser Lys Glu Met Gly Glu Thr Leu
1 5 10 15

35 Leu Arg Ala Val Glu Ser Tyr Leu Leu Ala His Ser Asp Ala Tyr Asn
20 25 30

<210> 6

<211> 31

40 <212> PRT

<213> Artificial Sequence

<220>

45 <223> Description of Artificial Sequence: solvent-exposed peptide

<400> 6

50 Cys Val Asp His Thr Asn Phe Lys Tyr Asn Tyr Ser Val Ile Glu Gly
1 5 10 15

Gly Pro Ile Gly Asp Thr Leu Glu Lys Ile Ser Asn Glu Ile Lys
20 25 30

55

Claims

1. A peptide wherein the amino acid sequence of the peptide is identical to an amino acid sequence which is part of the birch pollen allergen Bet v 1, or all amino acids of the peptide except the N-terminal or C-terminal amino acid of the peptide are identical to an amino acid sequence which is part of the birch pollen allergen Bet v 1, **characterized in that** said amino acid sequence which is part of the birch pollen allergen Bet v 1 is selected from the group consisting of amino acid positions 1 to 24, 30 to 59, 50 to 79, 110 to 139, 130 to 160 and 75 to 104 of the birch pollen allergen Bet v 1.
2. A peptide according to claim 1, consisting of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6.
3. A peptide according to claim 1 or 2, **characterized in that** the peptide upon administration is capable of inducing IgG antibodies which react with the birch pollen allergen Bet v 1.
4. A peptide according to claim 3 **characterized in that** the induced IgG antibodies can reduce or prevent binding of IgE antibodies to the birch pollen allergen Bet v 1.
5. A peptide according to any of claims 1 to 4 **characterized in that** the peptide upon administration does not induce an IgE response.
6. A pharmaceutical composition containing a peptide according to any one of claims 1 to 5 and a pharmaceutically acceptable carrier or diluent.
7. A pharmaceutical composition according to claim 6, further containing an adjuvant.
8. A pharmaceutical composition according to claim 6 or 7, **characterized in that** it is a vaccine composition.
9. A method for the preparation of a pharmaceutical composition comprising the following steps:
 - a) determining which amino acids of a given allergenic protein are solvent-exposed on the surface of the allergenic protein;
 - b) preparing a peptide having a length of 8 to 50 amino acids wherein at least three consecutive amino acids of the peptide are identical to at least three amino acids which appear on the molecular surface of an allergenic protein within a surface patch of 500 square Angström wherein the at least three amino acids are solvent-exposed amino acids in the allergenic protein; and
 - c) admixing the peptide with a pharmaceutically acceptable carrier or diluent;
 wherein the peptide lacks IgE binding capacity.
10. A method according to claim 9 for the preparation of a pharmaceutical composition, comprising the following steps:
 - a) determining which amino acids of a given allergenic protein are solvent-exposed on the surface of the allergenic protein;
 - b) preparing a peptide having a length of 8 to 50 amino acids wherein at least five consecutive amino acids of the peptide are identical to at least five consecutive amino acids of the amino acid sequence of the allergenic protein wherein the at least five consecutive amino acids are solvent-exposed amino acids in the allergenic protein; and
 - c) admixing the peptide with a pharmaceutically acceptable carrier or diluent.
11. A method according to claim 9 or 10, further comprising the addition of an adjuvant.
12. A method according to any one of claims 9 to 11, **characterized in that** all amino acids of the peptide except one are identical to an amino acid sequence which is part of the allergenic protein.
13. A method according to claim 12 **characterized in that** the one amino acid which deviates from the amino acid sequence of the allergenic protein is the N-terminal or C-terminal amino acid of the peptide.

14. A method according to any of claims 9 to 13 **characterized in that** the amino acid sequence of the peptide is identical to an amino acid sequence which is part of the allergenic protein.
15. A method according to any of claims 9 to 14 **characterized in that** the allergenic protein is the birch pollen allergen Bet v 1.
16. A method according to any of claims 9 to 15 **characterized in that** the solvent-exposed amino acids of the allergenic protein are determined by determining the hydrophilicity profile of the amino acid sequence of the allergenic protein.
17. A method according to any of claims 9 to 16 **characterized in that** the solvent-exposed amino acids are determined from the three-dimensional structure of the allergenic protein.
18. A method according to any of claims 9 to 17 wherein the length of the peptide is within a range of 21 to 35 amino acids.
19. A method according to claim 18 wherein the peptide comprises amino acid positions 1 to 24, 30 to 59, 50 to 79, 110 to 139, 130 to 160 or 75 to 104 of the birch pollen allergen Bet v 1.
20. A method according to any one of claims 9 to 19 wherein the peptide consists of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:6.
21. The use of a peptide according to any one of claims 1 to 5 for the preparation of a medicament for the treatment of an allergic disease.

Patentansprüche

1. Peptid, bei dem die Aminosäuresequenz des Peptids identisch ist mit einer Aminosäuresequenz, die Teil des Birkenpollenallergens Bet v1 ist oder bei dem alle Aminosäuren des Peptids außer der N-terminalen oder C-terminalen Aminosäure des Peptids identisch sind mit einer Aminosäuresequenz, die Teil des Birkenpollenallergens Bet v1 ist, **dadurch gekennzeichnet, dass** diese Aminosäuresequenz, die Teil des Birkenpollenallergens Bet v1 ist, ausgewählt ist aus der Gruppe bestehend aus den Aminosäurepositionen 1 bis 24, 30 bis 59, 50 bis 79, 110 bis 139, 130 bis 160 und 75 bis 104 des Birkenpollenallergens Bet v1.
2. Peptid nach Anspruch 1 bestehend aus einer Aminosäuresequenz ausgewählt aus der Gruppe bestehend aus SEQ ID Nr. 1, SEQ ID Nr. 2, SEQ ID Nr. 3, SEQ ID Nr. 4, SEQ ID Nr. 5 und SEQ ID Nr. 6.
3. Peptid nach Anspruch 1 oder Anspruch 2, **dadurch gekennzeichnet, dass** das Peptid bei Verabreichung IgG-Antikörper induzieren kann, die mit dem Birkenpollenallergen Bet v1 reagieren.
4. Peptid nach Anspruch 3, **dadurch gekennzeichnet, dass** die induzierten IgG-Antikörper das Binden von IgE-Antikörpern an das Birkenpollenallergen Bet v1 vermindern oder verhindern können.
5. Peptid nach einem der Ansprüche 1 bis 4, **dadurch gekennzeichnet, dass** das Peptid bei Verabreichung keine IgE-Antwort induziert.
6. Pharmazeutische Zusammensetzung enthaltend ein Peptid gemäß einem der Ansprüche 1 bis 5 und einen pharmazeutisch annehmbaren Träger oder ein pharmazeutisch annehmbares Verdünnungsmittel.
7. Pharmazeutische Zusammensetzung nach Anspruch 6, die weiterhin ein Adjuvans enthält.
8. Pharmazeutische Zusammensetzung nach Anspruch 6 oder Anspruch 7, **dadurch gekennzeichnet, dass** sie eine Impfzusammensetzung ist.
9. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, das die folgenden Stufen umfasst:
 - a) dass bestimmt wird, welche Aminosäuren eines gegebenen allergenen Proteins an der Oberfläche des allergenen Proteins einem Lösungsmittel ausgesetzt sind;

- b) ein Peptid hergestellt wird mit einer Länge von 8 bis 50 Aminosäuren, wobei mindestens drei aufeinander folgende Aminosäuren des Peptids identisch sind mit mindestens drei Aminosäuren, die auf der Moleküloberfläche eines allergenen Proteins erscheinen innerhalb eines Oberflächenstücks von 500 Å², wobei mindestens drei Aminosäuren in dem allergenen Protein Aminosäuren sind, die dem Lösungsmittel ausgesetzt sind und
- c) das Peptid mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel vermischt wird;

wobei dem Peptid die IgE-Bindungskapazität fehlt.

10. Verfahren nach Anspruch 9 zur Herstellung einer pharmazeutischen Zusammensetzung, das die folgenden Stufen umfasst:

- a) dass bestimmt wird, welche Aminosäuren eines gegebenen allergenen Proteins auf der Oberfläche des allergenen Proteins einem Lösungsmittel ausgesetzt sind;
- b) ein Peptid mit einer Länge von 8 bis 50 Aminosäuren hergestellt wird, wobei mindestens fünf aufeinander folgende Aminosäuren des Peptids identisch sind mit mindestens fünf aufeinander folgenden Aminosäuren der Aminosäuresequenz des allergenen Proteins, wobei die mindestens fünf aufeinander folgenden Aminosäuren in dem allergenen Protein dem Lösungsmittel ausgesetzte Aminosäuren sind und
- c) das Peptid mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel vermischt wird.

11. Verfahren nach Anspruch 9 oder Anspruch 10, das weiterhin die Zugabe eines Adjuvans beinhaltet.

12. Verfahren nach einem der Ansprüche 9 bis 11, **dadurch gekennzeichnet, dass** alle Aminosäuren des Peptids außer einer identisch sind mit einer Aminosäuresequenz, die Teil des allergenen Proteins ist.

13. Verfahren nach Anspruch 12, **dadurch gekennzeichnet, dass** die eine Aminosäure, die von der Aminosäuresequenz des allergenen Proteins abweicht, die N-terminale oder C-terminale Aminosäure des Peptids ist.

14. Verfahren nach einem der Ansprüche 9 bis 13, **dadurch gekennzeichnet, dass** die Aminosäuresequenz des Peptids identisch ist mit einer Aminosäuresequenz, die Teil des allergenen Proteins ist.

15. Verfahren nach einem der Ansprüche 9 bis 14, **dadurch gekennzeichnet, dass** das allergene Protein das Birkenpollenallergen Bet v1 ist.

16. Verfahren nach einem der Ansprüche 9 bis 15, **dadurch gekennzeichnet, dass** die dem Lösungsmittel ausgesetzten Aminosäuren des allergenen Proteins bestimmt werden, indem das Hydrophilieprofil der Aminosäuresequenz des allergenen Proteins bestimmt wird.

17. Verfahren nach einem der Ansprüche 9 bis 16, **dadurch gekennzeichnet, dass** die dem Lösungsmittel ausgesetzten Aminosäuren aus der dreidimensionalen Struktur des allergenen Proteins bestimmt werden.

18. Verfahren nach einem der Ansprüche 9 bis 17, wobei die Länge des Peptids in einem Bereich von 21 bis 35 Aminosäuren liegt.

19. Verfahren nach Anspruch 18, wobei das Peptid die Aminosäurepositionen 1 bis 24, 30 bis 59, 50 bis 79, 110 bis 139, 130 bis 160 oder 75 bis 104 des Birkenpollenallergens Bet v1 enthält.

20. Verfahren nach einem der Ansprüche 9 bis 19, wobei das Peptid aus einer Aminosäuresequenz besteht, die ausgewählt ist aus der Gruppe bestehend aus SEQ ID Nr. 1, SEQ ID Nr. 2, SEQ ID Nr. 3, SEQ ID Nr. 4, SEQ ID Nr. 5 und SEQ ID Nr. 6.

21. Verwendung eines Peptids nach einem der Ansprüche 1 bis 5 zur Herstellung eines Arzneimittels zur Behandlung einer allergischen Krankheit.

Revendications

1. Peptide dans lequel la séquence d'acides aminés du peptide est identique à une séquence d'acides aminés qui constitue une partie de l'allergène du pollen de bouleau Bet v 1, ou dans lequel tous les acides aminés du peptide

sauf l'acide aminé N-terminal ou C-terminal du peptide sont identiques à une séquence d'acides aminés qui fait partie de l'allergène du pollen de bouleau Bet v 1, **caractérisé en ce que** ladite séquence d'acides aminés qui constitue une partie de l'allergène du pollen de bouleau Bet v 1 est sélectionnée dans le groupe consistant en les positions d'acides aminés 1 à 24, 30 à 59, 50 à 79, 110 à 139, 130 à 160 et 75 à 104 de l'allergène du pollen de bouleau Bet v 1.

2. Peptide selon la revendication 1 consistant en une séquence d'acides aminés sélectionnée dans le groupe consistant en SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5 et SEQ ID N° 6.

3. Peptide selon la revendication 1 ou 2, **caractérisé en ce que** le peptide, lorsqu'il est administré, est capable d'induire des anticorps IgG qui réagissent avec l'allergène du pollen de bouleau Bet v 1.

4. Peptide selon la revendication 3, **caractérisé en ce que** les anticorps IgG induits peuvent réduire ou empêcher la liaison d'anticorps IgE à l'allergène du pollen de bouleau Bet v 1.

5. Peptide selon l'une quelconque des revendications 1 à 4, **caractérisé en ce que** le peptide, lorsqu'il est administré, n'induit pas de réponse par IgE.

6. Composition pharmaceutique contenant un peptide selon l'une quelconque des revendications 1 à 5 et un véhicule ou un diluant pharmaceutiquement acceptables.

7. Composition pharmaceutique selon la revendication 6, contenant en outre un adjuvant.

8. Composition pharmaceutique selon la revendication 6 ou 7, **caractérisée en ce qu'il** s'agit d'une composition de vaccin.

9. Procédé pour la préparation d'une composition pharmaceutique, comprenant les étapes suivantes, consistant à :

a) déterminer quels acides aminés d'une protéine allergénique donnée sont exposés au solvant sur la surface de la protéine allergénique ;

b) préparer un peptide ayant une longueur de 8 à 50 acides aminés dans lequel au moins trois acides aminés consécutifs du peptide sont identiques à au moins trois acides aminés qui apparaissent sur la surface moléculaire d'une protéine allergénique dans une surface de 500 Angström carrés dans lequel les au moins trois acides aminés sont, dans la protéine allergénique, des acides aminés exposés au solvant ; et

c) mélanger le peptide avec un véhicule ou un diluant pharmaceutiquement acceptables ;

dans lequel le peptide ne présente pas de capacité de liaison aux IgE.

10. Procédé selon la revendication 9 pour la préparation d'une composition pharmaceutique, comprenant les étapes suivantes, consistant à :

a) déterminer quels acides aminés d'une protéine allergénique donnée sont exposés au solvant sur la surface de la protéine allergénique ;

b) préparer un peptide ayant une longueur de 8 à 50 acides aminés dans lequel au moins cinq acides aminés consécutifs du peptide sont identiques à au moins cinq acides aminés consécutifs de la séquence d'acides aminés de la protéine allergénique, dans lequel les au moins cinq acides aminés consécutifs sont, dans la protéine allergénique, des acides aminés exposés au solvant ; et

c) mélanger le peptide avec un véhicule ou un diluant pharmaceutiquement acceptables.

11. Procédé selon la revendication 9 ou 10, comprenant en outre l'ajout d'un adjuvant.

12. Procédé selon l'une quelconque des revendications 9 à 11, **caractérisé en ce que** tous les acides aminés du peptide, sauf un, sont identiques à une séquence d'acides aminés qui constitue une partie de la protéine allergénique.

13. Procédé selon la revendication 12 **caractérisé en ce que** l'acide aminé qui dévie de la séquence d'acides aminés de la protéine allergénique est l'acide aminé N-terminal ou C-terminal du peptide.

14. Procédé selon l'une quelconque des revendications 9 à 13, **caractérisé en ce que** la séquence d'acides aminés

EP 1 219 299 B1

du peptide est identique à une séquence d'acides aminés qui constitue une partie de la protéine allergénique.

5 15. Procédé selon l'une quelconque des revendications 9 à 14, **caractérisé en ce que** la protéine allergénique est l'allergène du pollen de bouleau Bet V 1.

16. Procédé selon l'une quelconque des revendications 9 à 15, **caractérisé en ce que** les acides aminés exposés au solvant de la protéine allergénique sont déterminés en déterminant le profil d'hydrophilie de la séquence d'acides aminés de la protéine allergénique.

10 17. Procédé selon l'une quelconque des revendications 9 à 16, **caractérisé en ce que** les acides aminés exposés au solvant sont déterminés à partir de la structure en trois dimensions de la protéine allergénique.

18. Procédé selon l'une quelconque des revendications 9 à 17 dans lequel la longueur du peptide est dans une plage de 21 à 35 acides aminés.

15 19. Procédé selon la revendication 18, dans lequel le peptide comprend les positions d'acides aminés 1 à 24, 30 à 59, 50 à 79, 110 à 139, 130 à 160 ou 75 à 104 de l'allergène du pollen de bouleau Bet v 1.

20 20. Procédé selon l'une quelconque des revendications 9 à 19, dans lequel le peptide consiste en une séquence d'acides aminés choisie dans le groupe consistant en SEQ ID N° 1, SEQ ID N° 2, SEQ ID N° 3, SEQ ID N° 4, SEQ ID N° 5 et SEQ ID N° 6.

25 21. Utilisation d'un peptide selon l'une quelconque des revendications 1 à 5 pour la préparation d'un médicament pour le traitement d'une maladie allergique.

Figure 1




Figure 2




Figure 3

