



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 405/04, 257/04	A1	(11) International Publication Number: WO 94/12492 (43) International Publication Date: 9 June 1994 (09.06.94)
(21) International Application Number: PCT/EP93/03257 (22) International Filing Date: 19 November 1993 (19.11.93) (30) Priority Data: 9224922.6 27 November 1992 (27.11.92) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brendford, Middlesex 9EP TW8 (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): JOHNSON, Graham [GB/GB]; SmithKline Beecham Pharmaceuticals, Clarendon Road, Worthing, West Sussex BN14 8QH (GB). SMITH, Neil [GB/GB]; SmithKline Beecham Pharmaceuticals, Clarendon Road, Worthing, West Sussex BN14 8QH (GB). GEEN, Richard, Graham [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). MANN, Inderjit, Singh [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). NOVACK, Vance [US/US]; SmithKline Beecham Corporation, P.O. Box 1539, King of Prussia, Philadelphia, PA 19406-0939 (US).	(74) Agent: GIDDINGS, Peter, J.; Corporate Intellectual Property, SmithKline Beecham, Mundells, Welwyn Garden City, Hertfordshire AL7 1EY (GB). (81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: PROCESS FOR PREPARING BENZOPYRAN COMPOUNDS (57) Abstract A process for preparing substituted benzopyran derivatives which comprises a final step cyclisation of the corresponding open-chain intermediates.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				


Process for preparing benzopyran compounds

The present invention relates to a new process for preparing certain substituted benzopyran compounds, intermediates useful in the process, and to a process for the preparation of the intermediates.

Substituted benzopyran compounds are known in the art. For example, EP 0 173 516-A discloses a class of substituted benzopyran compounds which are described as compounds having activity as leukotriene antagonists and useful in therapy in the treatment of, for example, diseases induced by leukotrienes and 5- α -reductase.

The present invention relates to a new process for preparing certain of the benzopyran compounds described in EP 0 173 516-A, and in particular provides an efficient route to the compounds in which far fewer reaction steps are involved than has hitherto been described. A reduction in the number of reaction steps involved in preparing end products generally results in a much more efficient and cost-effective route than one involving large numbers of steps.


The present invention therefore provides in a first aspect a process for preparing a compound of structure (I):



(I)

in which,

R¹ is C₁₋₂₀alkyl, C₂₋₂₀alkenyl, C₂₋₂₀alkynyl, or a group of structure:



each of which may be substituted by one or two substituents selected independently from C₁₋₂₀alkyl, C₂₋₂₀alkenyl or C₂₋₂₀alkynyl, up to 5 carbon atom(s) of which may optionally be replaced by oxygen atom(s), sulphur atom(s), halogen atom(s), nitrogen atom(s), benzene ring(s), thiophene ring(s), naphthalene ring(s), carbocyclic ring(s) of from 4 to 7 carbon atom(s), carbonyl group(s), carbonyloxy group(s), hydroxy group(s), carboxy group(s), azido group(s) and/or nitro group(s);


R² is hydrogen or C₁₋₆alkyl;

R³ is hydrogen, halogen, hydroxy, nitro, a group of general formula -COOR⁴ (wherein R⁴ represents hydrogen or C₁₋₆alkyl), C₁₋₆alkyl, C₁₋₆alkoxy or C₁₋₆alkylthio;

A is a single bond or a methylene, ethylene, trimethylene, tetramethylene, vinylene, propenylene, butenylene, butadienylene or ethynylene group optionally being substituted by one, two or three C₁₋₁₀alkyl and/or phenyl group(s); and

X is oxygen or sulphur;


or a salt, solvate or hydrate thereof, which comprises cyclisation of a compound of structure (II)



or a salt, hydrate or solvate thereof, in which R¹, R², R³, A and X are as

described for structure (I), and optionally thereafter forming a salt, solvate or hydrate thereof.


Suitably, R^1 is C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, or a group of structure:



each of which may be substituted by one or two substituents selected independently from C_{1-20} alkyl, C_{2-20} alkenyl or C_{2-20} alkynyl, up to 5 carbon atom(s) of which may optionally be replaced by oxygen atom(s), sulphur atom(s), halogen atom(s), nitrogen atom(s), benzene ring(s), thiophene ring(s), naphthalene ring(s), carbocyclic ring(s) of from 4 to 7 carbon atom(s), carbonyl group(s), carbonyloxy group(s), hydroxy group(s), carboxy group(s), azido group(s) and/or nitro group(s).

Preferably R^1 is a group of structure (i) substituted or unsubstituted by one or two substituents selected independently from C_{1-20} alkyl, C_{2-20} alkenyl or C_{2-20} alkynyl, up to 5 carbon atom(s) of which may optionally be replaced by oxygen atom(s), sulphur atom(s), halogen atom(s), nitrogen atom(s), benzene ring(s), thiophene ring(s), naphthalene ring(s), carbocyclic ring(s) of from 4 to 7 carbon atom(s), carbonyl group(s), carbonyloxy group(s), hydroxy group(s), carboxy group(s), azido group(s) and/or nitro group(s).

More preferably R^1 is a group of structure (i) substituted in the para position of the ring by a single substituent selected from the above, in particular R^1 is a group of structure



Suitably, R^2 is hydrogen or C_{1-6} alkyl; preferably R^2 is hydrogen.

Suitably, R^3 is hydrogen, halogen, hydroxy, nitro, a group of general formula $-COOR^4$ (wherein R^4 represents hydrogen or C_{1-6} alkyl) or C_{1-6} alkyl, C_{1-6} alkoxy or C_{1-6} alkylthio. Preferably R^3 is hydrogen.

Suitably A is a single bond or a methylene, ethylene, trimethylene, tetramethylene, vinylene, propenylene, butenylene, butadienylene or ethynylene group optionally being substituted by one, two or three C₁₋₁₀alkyl and/or phenyl group(s). Preferably A is a single bond.


5

Suitably, the cyclisation of the compound of structure (II) is carried out in the presence of an acid. For example, the cyclisation can be carried out in the presence of sulphuric acid, in methanol or in acetic acid as the solvent medium. Preferably the reaction is carried out in a methanol/tetrahydrofuran solvent mixture in the presence of hydrochloric acid. Alternative acid/solvent conditions will be apparent to those skilled in the art and include, for example, acids such as hydrobromic or hydroiodic acid, perchloric acid or p-toluene sulphonic acid, and Lewis Acids for example aluminium trichloride, in suitable solvents such as water, C₁₋₄alkanols such as ethanol or methanol, and unsaturated carbocyclic hydrocarbons such as benzene or toluene.

10

It is to be noted that, although for the sake of convenience structure (II) is represented as the 'di-keto' form, the compounds of structure (II) can exist also in the 'keto-enol' form and in the 'cyclic hydroxy chromanone' form (IIB)

20




(IIB)

25

It is intended that structure (II) encompasses all of the tautomeric forms of the compounds of structure (II).


In a preferred aspect there is therefore provided a process for the preparation of a compound of structure (IA) or a salt, solvate or hydrate thereof:

30



(IA)

- 5 which comprises cyclisation of a compound of structure (IIA) or a salt, solvate or hydrate thereof:



(IIA)

10


and optionally thereafter forming a salt, solvate or hydrate thereof. Most preferably, the cyclisation is carried out in the presence of hydrochloric acid in methanol/tetrahydrofuran as solvent,

15

As for compounds of structure (II), the compounds of structure (IIA) can, of course, exist in the corresponding keto-enol and cyclic chromanone forms, each of which are intended to be encompassed by the structure (IIA).


20

The compounds of structure (II) (and in particular structure (IIA)) are novel and form a further aspect of the invention. The compounds of structure (II) can be prepared by the reaction of a compound of structure (III)



(III)

- 5 in which R¹, R², R³, A and X are as described for structure (I) in claim 1 with a compound of structure (IV) or a salt thereof



(IV)

10

in which Z is an activated leaving group.

- Suitably, activated leaving groups Z include, for example, activated
 15 amides of structure N(R⁵)(OR⁵) in which R⁵ is C₁₋₆alkyl, halogen groups, groups of structure R⁶O, R⁶S or R⁶SO₂O in which R⁶ is C₁₋₆alkyl, optionally substituted phenyl or optionally substituted phenylC₁₋₆alkyl, or groups of

structure $R^7\overset{\overset{X}{\parallel}}{C}-X-$ in which R⁷ is C₁₋₆alkyl, optionally substituted phenyl


or optionally substituted phenylC₁₋₆alkyl, and each group X is, independently, oxygen or sulphur. Preferably Z is R⁶O.

20


Suitably, R⁶ is C₁₋₆alkyl, optionally substituted phenyl or optionally substituted phenylC₁₋₆alkyl. Preferably, R⁶ is C₁₋₆alkyl, for example, methyl, ethyl, i-butyl or t-butyl; most preferably R⁶ is ethyl.

Suitably, the reaction is carried out in an organic solvent such as for example, dimethylformamide, ethereal solvents such as tetrahydrofuran, toluene or benzene, hexanes or C₁₋₆alkanols such as methanol or ethanol, in the presence of a base, for example an alkali metal alkoxide such as potassium t-butoxide, sodium methoxide or potassium methoxide, hydrides such as sodium hydride, or an amide base such as potassium amide or sodium amide. Preferably, the reaction is carried out in tetrahydrofuran as a solvent in the presence of sodium methoxide as a base.

The process for preparing the compounds of structure (II) is novel and forms a further aspect of the invention. In particular, the process is preferred to be used in the preparation of the compounds of structure (IIA) by reaction of the following compounds of structure (IIIA) and a compound of structure (IVA) or a salt thereof, in the presence of sodium methoxide in tetrahydrofuran as a solvent:



(IIIA)



(IVA)

The compounds of structure (III) and (IV) are prepared from commercially available starting materials by standard techniques as hereinafter described. For example, the preparation of compounds of structure (III) is described in EP 0 173 516-A. Compounds of structure (IV), for example in which Z is R⁶O, can be prepared from sodium azide and an appropriate alkyl cyanoformate such as ethyl cyanoformate by known methods or from tetrazole-5-carboxylic acid disodium salt (commercially available) by reaction with the appropriate alkyl, aryl or arylalkyl haloformate, for example, ethylchloroformate or isobutylchloroformate. The preparation of compounds of structure (IV) from the corresponding tetrazole-5-carboxylic acid disodium salt is novel and forms a still further aspect of the invention. It is to be noted that the compounds of structure (IV) can be prepared and then isolated before

reaction with appropriate compounds of structure (III), or can be prepared 'in situ' and further reacted with the compounds of structure (III) without prior isolation.

5 The present invention is in particular useful in the preparation of the compounds of structure (IA) beginning from compounds (IIIA) and (IVA) to form the intermediates of structure (IIA) which undergo cyclisation under the conditions described herein to form the desired product. The compounds of structure (II) can be isolated from the reaction mixture before cyclisation to
10 the compounds of structure (I) or alternatively, as described in the examples, the reaction between compounds of structures (III) and (IV) followed by the cyclisation of the compounds of structure (II) so formed can be carried to completion in 'one pot', that is to say without the isolation of the intermediates so formed.

15

 The following examples serve to illustrate the invention.
Temperatures are recorded in degrees Celsius (°C).

EXAMPLES

1. Preparation of Ethyl-1H-Tetrazole-5-Carboxylate

- 5 Trifluoroacetic acid (24.47 g, 0.21 M) was added dropwise over 0.5 hr under nitrogen to a stirred suspension of sodium azide (12.59 g, 0.19 M) in 2,6-lutidine (100 ml) at 8 to 12°. After stirring for 7 minutes ethyl cyanofornate (20.4 g, 0.20 M) was added in one portion. The mixture was heated and stirred at 75° for 6 hours and then, after cooling, stirred at 20° for 16 hr.
- 10 After cooling to 10° the mixture was added to ice (250 g) and 11 molar hydrochloric acid (100 ml) keeping the temperature below 20°. The product was extracted into ethyl acetate (1 x 250, 1 x 200, 2 x 100 ml) and the combined extracts dried over magnesium sulphate. After evaporation of the solvent under reduced pressure the oily product (38.22 g) was taken up in
- 15 ether (50 ml) and hexane (25 ml) added. Storage at 4° for 2-3 days produced crystalline ethyl-1H-tetrazole-5-carboxylate that was filtered off, washed with chilled ether, and air dried, 14.12 g (52.6% yield), m.p. 88 - 93°.

NMR: (270 MHz, solution in CDCl₃)

- 20 δ 13.6 - 13.8 (s, 1H); 4.6 - 4.5 (q, 2H); 1.5 - 1.4 (t, 3H).

Workup variation

- On a larger scale (83.4 g sodium azide) the reaction was worked up
- 25 differently in order to prevent the liberation of any hydrazoic acid.

- After stirring at 75° and 20° a solution of sodium nitrite (63 g) in water (300 ml) was added over 10 minutes at 20 to 30°. The mixture was stirred at 20 - 25° for 20 minutes and then a chilled mixture of water (1.5 L) and 11
- 30 molar hydrochloric acid (690 ml) added keeping the temperature between 25 and 30°. The product was then extracted into ethyl acetate and crystallised as described above.

2. Preparation of i-Butyl-1H-Tetrazole-5-Carboxylate

35

To a stirred suspension of tetrazole-5-carboxylic acid disodium salt (15.8 g, 0.1 mol) in dimethylformamide (100 ml) under a nitrogen atmosphere

at 5° was added isobutyl chloroformate (13.6 g, 13 ml, 0.1 mol) dropwise over 15 minutes. The mixture was stirred at 5-10° for 2 hours, then at 20° for 2 hours. The mixture was added to water (500 ml) and extracted with ethyl acetate (2 x 200 ml). The aqueous phase was then acidified to pH1 with conc. HCl and further extracted with ethyl acetate (2 x 200 ml). The latter extracts were washed with water (2 x 200 ml), dried (MgSO₄) and evaporated to give the title compound as a gum (8.6 g, 50.5%).

¹H NMR (CDCl₃): δ 0.95 (d, 6H, CH₃), 2.08 (tq, 1H, CH), 4.25 (d, 2H, CH₂).

3. Preparation of Methyl 4-(4-Phenylbutoxy)benzoate

A solution of methyl 4-hydroxybenzoate (13.4 kg, 88 mol) in DMF (52 L) was added dropwise to a mixture of NaOMe (4.8 kg, 89 mol) and DMF (50 L) at room temperature under a gentle stream of nitrogen. The reaction mixture was heated at 60-70° for 1 hr with stirring and then cooled to room temperature. To this mixture was added dropwise a solution of 4-phenylbutyl bromide (16.92 kg, 79.4 mol) in DMF (5 L). The resulting mixture was heated at 60-70° for 1 hr with constant stirring and cooled to room temperature. After an addition of 1N-NaOH (110L) was added, and the product was extracted twice with ethyl acetate (50 L and 80 L). The extracts were washed with 1N-NaOH (110 L) and saturated brine (20 L) successively, and then concentrated to dryness *in vacuo* to give the title compound in quantitative yield.

4. Preparation of 4-(4-Phenylbutoxy)benzoic Acid

To a solution of the compound from Example 3 in MeOH (50 L) was added 3N-NaOH (46 L). The mixture was heated under reflux for 1.5 hrs. Upon termination of the reaction, the MeOH was removed by distillation *in vacuo*. Ice-water (120 L) was added to the residue, and the neutral materials were extracted with ether (30 L x 3). The combined ethereal extracts were washed with 2N-NaOH (25 L). The aqueous layers were combined and adjusted to pH 2-3 with concentrated HCl (16 L). Precipitated solids were collected by centrifugal filtration, washed with water and dried by heating at 70-80° under a stream of air to obtain the title compound, (17.67 kg, 65.4 mol, 82% yield from 4-hydroxybenzoate).

5. **Preparation of 3-[4-(4-Phenylbutoxy)benzoylamino]-2-hydroxy-acetophenone**

5 To a solution of the compound from Example 4 (18.1 g, 67 mmol) in CH₂Cl₂ (45 ml) was added a catalytic amount of DMF (0.45 ml) followed by thionyl chloride (6.26 ml, 85.8 mmol) at room temperature under a stream of nitrogen. After reflux for 2 hr, the mixture was cooled to room temperature and was added to a solution of 3'-amino-2'-hydroxyacetophenone hydrochloride
10 (12 g, 64 mmol) and pyridine (15.5 ml, 192 mmol) in CH₂Cl₂ (90 ml) while maintaining the temperature between 0-3°. The mixture was stirred at 0-3° for 2 hr, and poured into 2N-HCl (200 ml). The aqueous layer was separated. The product in aqueous layer was extracted twice with CH₂Cl₂ (150 ml and 100 ml). The CH₂Cl₂ layers were combined, washed successively with water,
15 saturated NaHCO₃ (150 ml), and saturated brine (150 ml), dried over MgSO₄. The resulting solution was concentrated *in vacuo* until some of the crystals were precipitated. Ethyl acetate (150 ml) was added to the residue, and the solution was concentrated *in vacuo* until about a half of the ethyl acetate was distilled out. The mixture was cooled to approximately 0°. Precipitated
20 crystals were collected by filtration and dried *in vacuo* to afford the title compound, (21.6 g, 53.6 mmol, 90% yield).

6. **Preparation of 2-[4-(4-Phenylbutoxy)benzoylamino]-6-[1,3-dioxo-3-(tetrazol-5-yl)propyl] phenol**

25 Under a nitrogen atmosphere, potassium *tert*-butoxide (31.36 g, 0.28 mol) was dissolved in dry DMF (160 ml) by stirring. To the resulting solution were added the hydroxy acetophenone compound from Example 5 (16.12 g, 0.04 mol) followed by 5-ethoxycarbonyl tetrazole from Example 1 (7.39 g, 0.052
30 mol, 1.3 equiv.) at room temperature. The reaction temperature rises to approximately 45°. The mixture was stirred for 3 hours at 40° (oil bath), then cooled to 30° and poured into cold 1N HCl (800 ml). The resulting precipitate was filtered, washed with water (500 ml), and then dried at 70° in a fan oven to obtain the title compound (19.4 g, 97%). Purification was carried out using
35 either of the following procedures.

Procedure 1: A stirred slurry of crude product (10 g) in ethyl acetate (150 ml) was heated at 60° for 2 hours. After cooling to room temperature, the mixture was transferred to a refrigerator and left for 2 hours. The product was then filtered, washed with cold ethyl acetate (15 ml), and dried at 70° in a fan oven to afford purified product (8.5 g, 85%).

Procedure 2: A stirred slurry of crude product (5 g) in acetone (50 ml) was heated under reflux for 2 hours. After cooling to room temperature, the mixture was transferred to a refrigerator and left for 2 hours. The product was then filtered, washed with cold acetone (5-10 ml), and dried at 70° in a fan oven to afford purified product (4.1 g, 82%).

7. **Preparation of 4-Oxo-8-[4-(4-phenylbutoxy)benzoylamino]-2-tetrazol-5-yl-4H-1-benzopyran hemihydrate**

To a stirred slurry of purified product from Example 6 (7.984 g, 0.016 mol) in methanol (72 ml) was added concentrated sulphuric acid (0.6 ml), and the reaction heated to reflux and stirred for 3 hours. The mixture was allowed to cool to room temperature and then transferred to a refrigerator for 2 hours. The thick mixture was then filtered, washed with cold methanol (40 ml) and water (90 ml) followed again by cold methanol (30 ml). The product was dried at 70° in a fan oven and then left to stand for 24 hours at room temperature to afford the title compound (7.36 g, 96%):

25 **Examples 8 and 9**

These two examples illustrate the 'one-pot' procedure for the preparation of compounds (I) from the intermediate compounds of structures (III) and (IV).

30 8. **Preparation of 4-Oxo-8-[4-(4-phenylbutoxy)benzoylamino]-2-tetrazol-5-yl-4H-1-benzopyran hemihydrate.**

To a stirred suspension of sodium methoxide (15 g, 0.28 mole) in dry THF under a nitrogen atmosphere was added, in portions, the hydroxy-acetophenone compound from example 5 (16 g, 0.04 mole) at about 25°C. A solution of ethyl tetrazole-5-carboxylate from example 1 (7.3 g, 0.05 mole) in THF was then added while maintaining the reaction temperature at about

25 °C. The reaction mixture was stirred at reflux for about 100 minutes to ensure complete formation of the diketone compound from example 6. Methanol was added to the reaction mixture followed by concentrated hydrochloric acid (28 ml, 0.34 mole), and subsequent heating of the reaction mixture at reflux for about 2 hours resulted in the formation of the title compound which crystallised out of solution. After cooling to about 20 °C, the product was isolated by filtration and washed with methanol. The isolated solid was purified by conversion to the sodium salt in methanol and reprecipitating the title compound with hydrochloric acid. The reprecipitated product was isolated by filtration, washed with aqueous methanol, dried and then rehydrated at room temperature to give the title compound (18.56 g, 94%).


9. Preparation of 4-Oxo-8-[4-(4-phenylbutoxy)benzoylamino]-2-tetrazol-5-yl-4H-1-benzopyran hemihydrate.

To a stirred suspension of sodium methoxide (14.1 kg, 261 mole) in dry THF under a nitrogen atmosphere was added, in portions, the hydroxy-acetophenone compound from example 5 (15.0 kg, 37.2 mole) at about 25 °C. A solution of ethyl tetrazole-5-carboxylate from example 1 (6.8 kg, 47.9 mole) in THF was then added while maintaining the reaction temperature at about 25 °C. The reaction mixture was stirred at reflux for about 100 minutes to ensure complete formation of the diketone compound from example 6. Methanol was added to the reaction mixture followed by concentrated hydrochloric acid (31.4 kg, 314 mole), and subsequent heating of the reaction mixture at reflux for about 2 hours resulted in the formation of the title compound which crystallised out of solution. After cooling to about 20 °C, the product was isolated by filtration and washed with methanol. The isolated solid was purified by conversion to the sodium salt in methanol and reprecipitating the title compound with hydrochloric acid. The reprecipitated product was isolated by filtration, washed with aqueous methanol, dried and then rehydrated at room temperature to give the title compound (15.5 kg, 85%).

35

CLAIMS:


1. A process for preparing a compound of structure (I):



(I)

in which,

- 10 R^1 is C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, or a group of structure:



- 15 each of which may be substituted by one or two substituents selected independently from C_{1-20} alkyl, C_{2-20} alkenyl or C_{2-20} alkynyl, up to 5 carbon atom(s) of which may optionally be replaced by oxygen atom(s), sulphur atom(s), halogen atom(s), nitrogen atom(s), benzene ring(s), thiophene ring(s), naphthalene ring(s), carbocyclic ring(s) of from 4 to 7 carbon atom(s), carbonyl group(s), carbonyloxy group(s), hydroxy group(s), carboxy group(s), azido group(s) and/or nitro group(s);

R^2 is hydrogen or C_{1-6} alkyl;

- 25 R^3 is hydrogen, halogen, hydroxy, nitro, a group of general formula $-COOR^4$ (wherein R^4 represents hydrogen or C_{1-6} alkyl), or C_{1-6} alkyl, C_{1-6} alkoxy or C_{1-6} alkylthio;


A is a single bond or a methylene, ethylene, trimethylene, tetramethylene, vinylene, propenylene, butenylene, butadienylene or ethynylene group

optionally being substituted by one, two or three C₁₋₁₀alkyl and/or phenyl group(s); and

X is oxygen or sulphur;

5


or a salt, hydrate or solvate thereof, which comprises cyclisation of a compound of structure (II)



10


(II)

or a salt, hydrate or solvate thereof, in which R¹, R², R³, A and X are as described for structure (I), and optionally thereafter forming a salt, hydrate or
15 solvate thereof.

2. A process according to claim 1 in which R¹ is
, A is a single bond, X is oxygen, R² is hydrogen and
 R³ is hydrogen.

3. A process according to claim 2 in which the cyclisation reaction
 20 is carried out in the presence of hydrochloric acid in methanol/tetrahydrofuran as solvent.


4. A compound of structure (II):




in which R^1 , R^2 , R^3 , A and X are as described for structure (I), or a salt,
 5 hydrate or solvate thereof.

5. 2-[4-(4-Phenylbutoxy)benzoylamino]-6-[1,3-dioxo-3-(tetrazol-5-yl)propyl] phenol.


10 6. A process for preparing a compound of structure (II) which comprises reaction of a compound of structure (III):




15 in which R^1 , R^2 , R^3 , A and X are as described for structure (I) in claim 1 with a compound of structure (IV) or a salt thereof:



20 in which Z is an activated leaving group.


7. A process according to claim 6 in which, in structure (III), R^1 is , A is a single bond, X is oxygen, R^2 is hydrogen and R^3 is hydrogen.

8. A process for preparing a compound of structure (IA):




(IA)

or a salt, hydrate or solvate thereof, which comprises reaction of a compound of structure (IIIA) with a compound of structure (IVA) or a salt thereof:




(IIIA)



(IVA)

followed by cyclisation of the intermediate compound of structure (IIA)




(IIA)

or a salt, hydrate or solvate thereof, so formed.

9. A process for preparing a compound of structure (IV):

5



10 in which Z is a group R^6O in which R^6 is C_{1-6} alkyl, optionally substituted phenyl or optionally substituted phenyl C_{1-6} alkyl, which comprises reaction of tetrazole-5-carboxylic acid disodium salt with the appropriate alkyl, aryl or arylalkyl haloformate.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 93/03257

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 5 C07D405/04 C07D257/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 115, no. 17, 1991, Columbus, Ohio, US; abstract no. 182817v, page 874 ; see abstract	1-8
A	& JP,A,0 395 144 (ONO PHARMACEUTICAL) 19 April 1991 -----	1-8

☐ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

4 February 1994

Date of mailing of the international search report

- 9. 02. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+ 31-70) 340-3016

Authorized officer

Francois, J

Information on patent family members

PC1/EP 93/03257

Form PCT/ISA/210 (patent family annex) (July 1992)