

**(19) World Intellectual Property Organization
International Bureau**



A standard linear barcode is located at the bottom of the page, spanning most of the width. It is used for tracking and identification of the document.

**(43) International Publication Date
30 October 2008 (30.10.2008)**

PCT

(10) International Publication Number
WO 2008/128984 A1

(51) International Patent Classification:

C07D 403/12 (2006.01) **A61K 31/506** (2006.01)
C07D 401/14 (2006.01) **A61P 25/00** (2006.01)

(21) International Application Number:

PCT/EP2008/054714

(22) International Filing Date: 18 April 2008 (18.04.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/912,942 20 April 2007 (20.04.2007) US

(71) Applicant (for all designated States except US): PROBIO-

DRUG AG [—/DE]; Weinbergweg 22, 06120 Halle/Saale

(72) Inventors; and

(75) **Inventors/Applicants (for US only): BUCHHOLZ, Mirko [DE/DE]; Brandenburger Strasse 6, 06114 Halle/Saale (DE). HEISER, Ulrich [DE/DE]; Franz-Schubert-Strasse 5, 06108 Halle/Saale (DE). SOMMER, Robert [DE/DE]; Schleiermacherstrasse 20, 06114 Halle/Saale (DE).**

(74) **Agent: HOFFMANN, Matthias**; Probiot drug AG, Weinbergweg 22, 06120 Halle / Saale (DE).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.



(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: AMINOPYRIMIDINE DERIVATIVES AS GLUTAMINYL CYCLASE INHIBITORS



WO 2008/128984 A1

(57) Abstract: The present invention relates to compounds of formula (I), combinations and uses thereof for disease therapy, or a pharmaceutically acceptable salt, solvate or polymorph thereof, including all tautomers and stereoisomers thereof wherein: A represents formula (A) or formula (B) and B, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, Y and Z are as defined throughout the description and the claims.



- *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

AMINOPYRIMIDINE DERIVATIVES AS GLUTAMINYL CYCLASE INHIBITORS

Field of the invention

The invention relates to novel pyrimidine derivatives as inhibitors of glutaminyl cyclase (QC, EC

5 2.3.2.5). QC catalyzes the intramolecular cyclization of N-terminal glutamine residues into pyroglutamic acid (5-oxo-prolyl, pGlu*) under liberation of ammonia and the intramolecular cyclization of N-terminal glutamate residues into pyroglutamic acid under liberation of water.

Background of the invention

10 Glutaminyl cyclase (QC, EC 2.3.2.5) catalyzes the intramolecular cyclization of N-terminal glutamine residues into pyroglutamic acid (pGlu*) liberating ammonia. A QC was first isolated by Messer from the latex of the tropical plant *Carica papaya* in 1963 (Messer, M. 1963 Nature 4874, 1299). 24 years later, a corresponding enzymatic activity was discovered in animal pituitary (Busby, W. H. J. et al. 1987 J Biol Chem 262, 8532-8536; Fischer, W. H. and Spiess, J. 15 1987 Proc Natl Acad Sci U S A 84, 3628-3632). For the mammalian QC, the conversion of Gln into pGlu by QC could be shown for the precursors of TRH and GnRH (Busby, W. H. J. et al. 1987 J Biol Chem 262, 8532-8536; Fischer, W. H. and Spiess, J. 1987 Proc Natl Acad Sci U S A 84, 3628-3632). In addition, initial localization experiments of QC revealed a co-localization 20 with its putative products of catalysis in bovine pituitary, further improving the suggested function in peptide hormone synthesis (Bockers, T. M. et al. 1995 J Neuroendocrinol 7, 445-453). In contrast, the physiological function of the plant QC is less clear. In the case of the enzyme from *C. papaya*, a role in the plant defense against pathogenic microorganisms was suggested (El Moussaoui, A. et al. 2001 Cell Mol Life Sci 58, 556-570). Putative QCs from other plants were identified by sequence comparisons recently (Dahl, S. W. et al. 2000 Protein Expr 25 Purif 20, 27-36). The physiological function of these enzymes, however, is still ambiguous.

The QCs known from plants and animals show a strict specificity for L-Glutamine in the N-terminal position of the substrates and their kinetic behavior was found to obey the Michaelis-Menten equation (Pohl, T. et al. 1991 Proc Natl Acad Sci U S A 88, 10059-10063; Consalvo, A. P. et al. 1988 Anal Biochem 175, 131-138; Gololobov, M. Y. et al. 1996 Biol Chem Hoppe

30 Seyler 377, 395-398). A comparison of the primary structures of the QCs from *C. papaya* and that of the highly conserved QC from mammals, however, did not reveal any sequence homology (Dahl, S. W. et al. 2000 Protein Expr Purif 20, 27-36). Whereas the plant QCs appear to belong to a new enzyme family (Dahl, S. W. et al. 2000 Protein Expr Purif 20, 27-36), the mammalian QCs were found to have a pronounced sequence homology to bacterial aminopeptidases (Bateman, R. C. et al. 2001 Biochemistry 40, 11246-11250), leading to the 35 conclusion that the QCs from plants and animals have different evolutionary origins.

Recently, it was shown that recombinant human QC as well as QC-activity from brain extracts catalyze both, the N-terminal glutamyl as well as glutamate cyclization. Most striking is the finding, that cyclase-catalyzed Glu₁-conversion is favored around pH 6.0 while Gln₁-conversion to pGlu-derivatives occurs with a pH-optimum of around 8.0. Since the formation of pGlu-A β -related peptides can be suppressed by inhibition of recombinant human QC and QC-activity from pig pituitary extracts, the enzyme QC is a target in drug development for treatment of Alzheimer's disease.

First inhibitors of QC are described in WO 2004/098625, WO 2004/098591, WO 2005/039548 and WO 2005/075436.

EP 02 011 349.4 discloses polynucleotides encoding insect glutamyl cyclase, as well as polypeptides encoded thereby and their use in methods of screening for agents that reduce glutamyl cyclase activity. Such agents are useful as pesticides.

The terms "k_i" or "K_i" and "K_D" are binding constants, which describe the binding of an inhibitor to and the subsequent release from an enzyme. Another measure is the "IC₅₀" value, which reflects the inhibitor concentration, which at a given substrate concentration results in 50 % enzyme activity.

The term "DP IV-inhibitor" or "dipeptidyl peptidase IV inhibitor" is generally known to a person skilled in the art and means enzyme inhibitors, which inhibit the catalytic activity of DP IV or DP IV-like enzymes.

"DP IV-activity" is defined as the catalytic activity of dipeptidyl peptidase IV (DP IV) and DP IV-like enzymes. These enzymes are post-proline (to a lesser extent post-alanine, post-serine or post-glycine) cleaving serine proteases found in various tissues of the body of a mammal including kidney, liver, and intestine, where they remove dipeptides from the N-terminus of biologically active peptides with a high specificity when proline or alanine form the residues that are adjacent to the N-terminal amino acid in their sequence.

The term "PEP-inhibitor" or "prolyl endopeptidase inhibitor" is generally known to a person skilled in the art and means enzyme inhibitors, which inhibit the catalytic activity of prolyl endopeptidase (PEP, prolyl oligopeptidase, POP).


"PEP-activity" is defined as the catalytic activity of an endoprotease that is capable to hydrolyze post proline bonds in peptides or proteins were the proline is in amino acid position 3 or higher counted from the N-terminus of a peptide or protein substrate.

5 The term "QC" as used herein comprises glutaminyl cyclase (QC) and QC-like enzymes. QC and QC-like enzymes have identical or similar enzymatic activity, further defined as QC activity. In this regard, QC-like enzymes can fundamentally differ in their molecular structure from QC. Examples of QC-like enzymes are the glutaminyl-peptide cyclotransferase-like proteins (QPCTLs) from human (GenBank NM_017659), mouse (GenBank BC058181), Macaca fascicularis (GenBank AB168255), Macaca mulatta (GenBank XM_001110995), Canis familiaris (GenBank XM_541552), Rattus norvegicus (GenBank XM_001066591), Mus musculus (GenBank BC058181) and Bos taurus (GenBank BT026254).

10


15 The term "QC activity" as used herein is defined as intramolecular cyclization of N-terminal glutamine residues into pyroglutamic acid (pGlu*) or of N-terminal L-homoglutamine or L- β -homoglutamine to a cyclic pyro-homoglutamine derivative under liberation of ammonia. See therefore schemes 1 and 2.

Scheme 1: Cyclization of glutamine by QC



20


Scheme 2: Cyclization of L-homoglutamine by QC



The term "EC" as used herein comprises the activity of QC and QC-like enzymes as glutamate cyclase (EC), further defined as EC activity.

The term "EC activity" as used herein is defined as intramolecular cyclization of N-terminal glutamate residues into pyroglutamic acid (pGlu*) by QC. See therefore scheme 3.

Scheme 3: N-terminal cyclization of uncharged glutamyl peptides by QC (EC)



10 The term "QC-inhibitor" "glutaminyl cyclase inhibitor" is generally known to a person skilled in the art and means enzyme inhibitors, which inhibit the catalytic activity of glutaminyl cyclase (QC) or its glutamyl cyclase (EC) activity.

Potency of QC inhibition

15 In light of the correlation with QC inhibition, in preferred embodiments, the subject method and medical use utilize an agent with an IC_{50} for QC inhibition of 10 μ M or less, more preferably of 1 μ M or less, even more preferably of 0.1 μ M or less or 0.01 μ M or less, or most preferably 0.001 μ M or less. Indeed, inhibitors with K_i values in the lower micromolar, preferably the nanomolar and even more preferably the picomolar range are contemplated. Thus, while the active agents 20 are described herein, for convenience, as "QC inhibitors", it will be understood that such nomenclature is not intending to limit the subject of the invention to a particular mechanism of action.

Molecular weight of QC inhibitors

25 In general, the QC inhibitors of the subject method or medical use will be small molecules, e.g., with molecular weights of 500 g/mole or less, 400 g/mole or less, preferably of 350 g/mole or less, and even more preferably of 300 g/mole or less and even of 250 g/mole or less.

The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term "pharmaceutically acceptable" embraces both human and veterinary use: For example the term "pharmaceutically acceptable" embraces a veterinarian acceptable compound or a compound acceptable in human medicine and health care.

Throughout the description and the claims the expression "alkyl", unless specifically limited, denotes a C₁₋₁₂ alkyl group, suitably a C₁₋₆ alkyl group, e.g. C₁₋₄ alkyl group. Alkyl groups may be straight chain or branched. Suitable alkyl groups include, for example, methyl, ethyl, propyl (e.g. n-propyl and isopropyl), butyl (e.g. n-butyl, iso-butyl, sec-butyl and tert-butyl), pentyl (e.g. n-pentyl), hexyl (e.g. n-hexyl), heptyl (e.g. n-heptyl) and octyl (e.g. n-octyl). The expression "alk", for example in the expressions "alkoxy", "haloalkyl" and "thioalkyl" should be interpreted in accordance with the definition of "alkyl". Exemplary alkoxy groups include methoxy, ethoxy, propoxy (e.g. n-propoxy), butoxy (e.g. n-butoxy), pentoxy (e.g. n-pentoxy), hexoxy (e.g. n-hexaoxy), heptoxy (e.g. n-heptoxy) and octoxy (e.g. n-octoxy). Exemplary thioalkyl groups include methylthio-. Exemplary haloalkyl groups include fluoroalkyl e.g. CF₃.

The expression "alkenyl", unless specifically limited, denotes a C₂₋₁₂ alkenyl group, suitably a C₂₋₆ alkenyl group, e.g. a C₂₋₄ alkenyl group, which contains at least one double bond at any desired location and which does not contain any triple bonds. Alkenyl groups may be straight chain or branched. Exemplary alkenyl groups including one double bond include propenyl and butenyl. Exemplary alkenyl groups including two double bonds include pentadienyl, e.g. (1E, 3E)-pentadienyl.

The expression "alkynyl", unless specifically limited, denotes a C₂₋₁₂ alkynyl group, suitably a C₂₋₆ alkynyl group, e.g. a C₂₋₄ alkynyl group, which contains at least one triple bond at any desired location and may or may not also contain one or more double bonds. Alkynyl groups may be straight chain or branched. Exemplary alkynyl groups include propynyl and butynyl.

The expression "alkylene" denotes a chain of formula -(CH₂)_n- wherein n is an integer e.g. 2-5, unless specifically limited.

The expression "cycloalkyl", unless specifically limited, denotes a C₃₋₁₀ cycloalkyl group (i.e. 3 to 10 ring carbon atoms), more suitably a C₃₋₈ cycloalkyl group, e.g. a C₃₋₆ cycloalkyl group. Exemplary cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl

5 and cyclooctyl. A most suitable number of ring carbon atoms is three to six.

The expression "cycloalkenyl", unless specifically limited, denotes a C₅₋₁₀ cycloalkenyl group (i.e. 5 to 10 ring carbon atoms), more suitably a C₅₋₈ cycloalkenyl group e.g. a C₅₋₆ cycloalkenyl group. Exemplary cycloalkenyl groups include cyclopropenyl, cyclohexenyl, cycloheptenyl and

10 cyclooctenyl. A most suitable number of ring carbon atoms is five to six.

The expression "carbocyclyl", unless specifically limited, denotes any ring system in which all the ring atoms are carbon and which contains between three and twelve ring carbon atoms, suitably between three and ten carbon atoms and more suitably between three and eight carbon

15 atoms. Carbocyclyl groups may be saturated or partially unsaturated, but do not include aromatic rings. Examples of carbocyclyl groups include monocyclic, bicyclic, and tricyclic ring systems, in particular monocyclic and bicyclic ring systems. Other carbocyclyl groups include bridged ring systems (e.g. bicyclo[2,2,1]heptenyl). A specific example of a carbocyclyl group is a cycloalkyl group. A further example of a carbocyclyl group is a cycloalkenyl group.

20 The expression "heterocyclyl", unless specifically limited, refers to a carbocyclyl group wherein one or more (e.g. 1, 2 or 3) ring atoms are replaced by heteroatoms selected from N, S and O. A specific example of a heterocyclyl group is a cycloalkyl group (e.g. cyclopentyl or more particularly cyclohexyl) wherein one or more (e.g. 1, 2 or 3, particularly 1 or 2, especially 1) ring

25 atoms are replaced by heteroatoms selected from N, S or O. Exemplary heterocyclyl groups containing one hetero atom include pyrrolidine, tetrahydrofuran and piperidine, and exemplary heterocyclyl groups containing two hetero atoms include morpholine and piperazine. A further specific example of a heterocyclyl group is a cycloalkenyl group (e.g. a cyclohexenyl group) wherein one or more (e.g. 1, 2 or 3, particularly 1 or 2, especially 1) ring atoms are replaced by

30 heteroatoms selected from N, S and O. An example of such a group is dihydropyranyl (e.g. 3,4-dihydro-2H-pyran-2-yl-).

The expression "aryl", unless specifically limited, denotes a C₆₋₁₂ aryl group, suitably a C₆₋₁₀ aryl group, more suitably a C₆₋₈ aryl group. Aryl groups will contain at least one aromatic ring (e.g.

35 one, two or three rings). An example of a typical aryl group with one aromatic ring is phenyl. An example of a typical aryl group with two aromatic rings is naphthyl.

The expression "heteroaryl", unless specifically limited, denotes an aryl residue, wherein one or more (e.g. 1, 2, 3, or 4, suitably 1, 2 or 3) ring atoms are replaced by heteroatoms selected from N, S and O, or else a 5-membered aromatic ring containing one or more (e.g. 1, 2, 3, or 4, suitably 1, 2 or 3) ring atoms selected from N, S and O. Exemplary monocyclic heteroaryl

5 groups having one heteroatom include: five membered rings (e.g. pyrrole, furan, thiophene); and six membered rings (e.g. pyridine, such as pyridin-2-yl, pyridin-3-yl and pyridin-4-yl). Exemplary monocyclic heteroaryl groups having two heteroatoms include: five membered rings (e.g. pyrazole, oxazole, isoxazole, thiazole, isothiazole, imidazole, such as imidazol-1-yl, imidazol-2-yl imidazol-4-yl); six membered rings (e.g. pyridazine, pyrimidine, pyrazine).

10 Exemplary monocyclic heteroaryl groups having three heteroatoms include: 1,2,3-triazole and 1,2,4-triazole. Exemplary monocyclic heteroaryl groups having four heteroatoms include tetrazole. Exemplary bicyclic heteroaryl groups include: indole (e.g. indol-6-yl), benzofuran, benzthiophene, quinoline, isoquinoline, indazole, benzimidazole, benzthiazole, quinazoline and purine.

15 The expression "-alkylaryl", unless specifically limited, denotes an aryl residue which is connected via an alkylene moiety e.g. a C₁₋₄alkylene moiety.

20 The expression "-alkylheteroaryl", unless specifically limited, denotes a heteroaryl residue which is connected via an alkylene moiety e.g. a C₁₋₄alkylene moiety.

The term "halogen" or "halo" comprises fluorine (F), chlorine (Cl) and bromine (Br).

The term "amino" refers to the group -NH₂.

25 Stereoisomers:
All possible stereoisomers of the claimed compounds are included in the present invention.
Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention.

Preparation and isolation of stereoisomers:
35 Where the processes for the preparation of the compounds according to the invention give rise to a mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or

individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their components enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed

5 by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

10 **Pharmaceutically acceptable salts:**

In view of the close relationship between the free compounds and the compounds in the form of their salts or solvates, whenever a compound is referred to in this context, a corresponding salt, solvate or polymorph is also intended, provided such is possible or appropriate under the circumstances.

15 Salts and solvates of the compounds of formula (I) and physiologically functional derivatives thereof which are suitable for use in medicine are those wherein the counter-ion or associated solvent is pharmaceutically acceptable. However, salts and solvates having non-pharmaceutically acceptable counter-ions or associated solvents are within the scope of the
20 present invention, for example, for use as intermediates in the preparation of other compounds and their pharmaceutically acceptable salts and solvates.

Suitable salts according to the invention include those formed with both organic and inorganic acids or bases. Pharmaceutically acceptable acid addition salts include those formed from

25 hydrochloric, hydrobromic, sulfuric, nitric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, triphenylacetic, sulfamic, sulfanilic, succinic, oxalic, fumaric, maleic, malic, mandelic, glutamic, aspartic, oxaloacetic, methanesulfonic, ethanesulfonic, arylsulfonic (for example p-toluenesulfonic, benzenesulfonic, naphthalenesulfonic or naphthalenedisulfonic), salicylic, glutaric, gluconic, tricarballylic, cinnamic, substituted cinnamic (for example, phenyl,
30 methyl, methoxy or halo substituted cinnamic, including 4-methyl and 4-methoxycinnamic acid), ascorbic, oleic, naphthoic, hydroxynaphthoic (for example 1- or 3-hydroxy-2-naphthoic), naphthaleneacrylic (for example naphthalene-2-acrylic), benzoic, 4-methoxybenzoic, 2- or 4-hydroxybenzoic, 4-chlorobenzoic, 4-phenylbenzoic, benzeneacrylic (for example 1,4-benzenediacylic), isethionic acids, perchloric, propionic, glycolic, hydroxyethanesulfonic,
35 pamoic, cyclohexanesulfamic, salicylic, saccharinic and trifluoroacetic acid. Pharmaceutically acceptable base salts include ammonium salts, alkali metal salts such as those of sodium and

potassium, alkaline earth metal salts such as those of calcium and magnesium and salts with organic bases such as dicyclohexylamine and *N*-methyl-D-glucamine.

All pharmaceutically acceptable acid addition salt forms of the compounds of the present invention are intended to be embraced by the scope of this invention.

5

Polymorph crystal forms:

Furthermore, some of the crystalline forms of the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e. hydrates) or common organic solvents, and such solvates are

10 also intended to be encompassed within the scope of this invention. The compounds, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization.

Prodrugs:

15 The present invention further includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds which are readily convertible *in vivo* into the desired therapeutically active compound. Thus, in these cases, the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with prodrug versions of one or 20 more of the claimed compounds, but which converts to the above specified compound *in vivo* after administration to the subject. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

25 Protective Groups:

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973;

30 and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991, fully incorporated herein by reference. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

As used herein, the term "composition" is intended to encompass a product comprising the 35 claimed compounds in the therapeutically effective amounts, as well as any product which results, directly or indirectly, from combinations of the claimed compounds.

Carriers and Additives for galenic formulations:

Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives may advantageously include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as,

5 for example, powders, capsules, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like.

Carriers, which can be added to the mixture, include necessary and inert pharmaceutical

10 excipients, including, but not limited to, suitable binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, coatings, disintegrating agents, dyes and coloring agents.

Soluble polymers as targetable drug carriers can include polyvinylpyrrolidone, pyran copolymer,

15 polyhydroxypropylmethacrylamidephenol, polyhydroxyethylaspartamide-phenol, or polyethyleneoxidepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polyactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-20 linked or amphipathic block copolymers of hydrogels.

Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or

betalactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium


oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium

25 chloride and the like.

Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

30 **Summary of the invention**

According to the invention there are provided compounds of formula (I),



or a pharmaceutically acceptable salt, solvate or polymorph thereof, including all tautomers and stereoisomers thereof wherein:

R¹ represents alkyl; haloalkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl; or

5 optionally substituted heteroaryl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

and in which any of the aforesaid aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl), -C(O)OC₁₋₆alkyl, SOC₁₋₄alkyl and -SOC₃₋₆cycloalkyl;

10 or R¹ represents -carbocyclyl-phenyl; -carbocyclyl-(monocyclic heteroaryl); -heterocyclyl-phenyl;

15 -carbocyclyl fused to phenyl; carbocyclyl fused to monocyclic heteroaryl or heterocyclyl fused to phenyl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

and in which any of the aforesaid phenyl and heteroaryl groups may optionally be substituted by

20 one or more substituents selected from C₁₋₆alkyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl) and -SOC₁₋₄alkyl;

and

25 R² represents alkyl; haloalkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

and in which any of the aforesaid aryl and heteroaryl groups may optionally be substituted by

30 one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl) and -SOC₁₋₄alkyl;

$\text{C}_1\text{-alkenyl}$, $-\text{SO}_2\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-6}\text{alkoxy-}$, $-\text{O}-\text{C}_{3-8}\text{cycloalkyl}$, $\text{C}_{3-8}\text{cycloalkyl}$, $-\text{SO}_2\text{C}_{3-8}\text{cycloalkyl}$, $\text{C}_{3-6}\text{alkenyloxy-}$, $\text{C}_{3-6}\text{alkynyl}$, $-\text{C}(\text{O})\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-6}\text{alkoxy-C}_{1-6}\text{alkyl-}$, nitro, halogen, cyano, hydroxyl, $-\text{C}(\text{O})\text{OH}$, $-\text{NH}_2$, $-\text{NHC}_{1-4}\text{alkyl}$, $-\text{N}(\text{C}_{1-4}\text{alkyl})(\text{C}_{1-4}\text{alkyl})$, $-\text{C}(\text{O})\text{N}(\text{C}_{1-4}\text{alkyl})(\text{C}_{1-4}\text{alkyl})$, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_{1-4}\text{alkyl})$, $-\text{C}(\text{O})\text{OC}_{1-6}\text{alkyl}$, $\text{SOC}_{1-4}\text{alkyl}$ and $-\text{SOC}_{3-6}\text{cycloalkyl}$;


5 or R^2 represents $-\text{carbocyclyl-phenyl}$; $-\text{carbocyclyl-(monocyclic heteroaryl)}$; $-\text{heterocyclyl-phenyl}$; $-\text{carbocyclyl fused to phenyl}$; $-\text{carbocyclyl fused to monocyclic heteroaryl}$ or $-\text{heterocyclyl fused to phenyl}$;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

10 and in which any of the aforesaid phenyl and heteroaryl groups may optionally be substituted by one or more substituents selected from $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-6}\text{haloalkyl}$, $-\text{C}_{1-6}\text{thioalkyl}$, $-\text{SO}_2\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-6}\text{alkoxy-}$, $-\text{C}(\text{O})\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-6}\text{alkoxy-C}_{1-6}\text{alkyl-}$, nitro, halogen, cyano, hydroxyl, $-\text{C}(\text{O})\text{OH}$, $-\text{NH}_2$, $-\text{NHC}_{1-4}\text{alkyl}$, $-\text{N}(\text{C}_{1-4}\text{alkyl})(\text{C}_{1-4}\text{alkyl})$, $-\text{C}(\text{O})\text{N}(\text{C}_{1-4}\text{alkyl})(\text{C}_{1-4}\text{alkyl})$, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_{1-4}\text{alkyl})$ and $-\text{SOC}_{1-4}\text{alkyl}$;

15 and

A represents




wherein Y represents a C_{2-5} alkylene chain, which may optionally be substituted by one or two methyl groups or may optionally be substituted by two alkylene substituents at the same

20 position wherein the two alkylene substituents are joined to each other to form a C_{3-5} spirocycloalkyl group and

R^3 , R^4 and R^5 independently represent H or $\text{C}_{1-2}\text{alkyl}$; or

A represents



25 wherein Z represents a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{-CH}_2-$, $-\text{CH}(\text{Me})-$, $-\text{CH}(\text{Me})\text{-CH}_2-$ or $-\text{CH}_2\text{-CH}(\text{Me})-$ and R^6 and R^7 independently represent H or $\text{C}_{1-2}\text{alkyl}$,

B represents H or methyl.

Typically R¹ represents alkyl; haloalkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;
which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

5 and in which any of the aforesaid phenyl, aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂ and -C(O)NH(C₁₋₄alkyl) and

10 typically R² represents alkyl; haloalkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;
which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

15 and in which any of the aforesaid phenyl, aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂ and -C(O)NH(C₁₋₄alkyl).

20

More typically R¹ represents alkyl; haloalkyl; alkenyl; carbocyclyl; optionally substituted aryl or optionally substituted heteroaryl;
which any of the aforesaid carbocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

25 and in which any of the aforesaid phenyl, aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂ and -C(O)NH(C₁₋₄alkyl) and

30 typically R² represents alkyl; haloalkyl; alkenyl; carbocyclyl; optionally substituted aryl or optionally substituted heteroaryl;
which any of the aforesaid carbocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

35 and in which any of the aforesaid phenyl, aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂ and -C(O)NH(C₁₋₄alkyl).

$\text{C}_1\text{-}_6$ haloalkyl, $-\text{C}_1\text{-}_6$ thioalkyl, $-\text{SO}_2\text{C}_1\text{-}_4$ alkyl, $\text{C}_1\text{-}_6$ alkoxy-, $-\text{O}-\text{C}_3\text{-}_8$ cycloalkyl, $\text{C}_3\text{-}_8$ cycloalkyl, $-\text{SO}_2\text{C}_3\text{-}_8$ cycloalkyl, $\text{C}_3\text{-}_6$ alkenyloxy-, $\text{C}_3\text{-}_6$ alkynyoxy-, $-\text{C}(\text{O})\text{C}_1\text{-}_6$ alkyl, $\text{C}_1\text{-}_6$ alkoxy- $\text{C}_1\text{-}_6$ alkyl-, nitro, halogen, cyano, hydroxyl, $-\text{C}(\text{O})\text{OH}$, $-\text{NH}_2$, $-\text{NHC}_1\text{-}_4$ alkyl, $-\text{N}(\text{C}_1\text{-}_4\text{alkyl})(\text{C}_1\text{-}_4\text{alkyl})$, $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-}_4\text{alkyl})(\text{C}_1\text{-}_4\text{alkyl})$, $-\text{C}(\text{O})\text{NH}_2$ and $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-}_4\text{alkyl})$.

5

Detailed description of the invention

In one embodiment, the following compounds are disclaimed from the definition of formula (I):

N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(3-pyridinyl)-4-pyrimidinamine;

6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;

10 N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(4-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine;

2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine;

2-(3-chloro-4-fluorophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;

2-(4-bromophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-methyl-4-pyrimidinamine;

15 2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine;

2-(4-bromophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;

6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-4-pyridinyl-4-pyrimidinamine,

N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-6-(4-pyridinyl)-4-pyrimidinamine;

20 N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-(4-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(4-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-(3-pyridinyl)-2-(4-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-3-pyridinyl-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-(3-pyridinyl)-4-pyrimidinamine;

25 N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(2-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(3-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(2-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(3-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(4-pyridinyl)-4-pyrimidinamine;

30 N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-(2-pyridinyl)-2-(4-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-(2-pyridinyl)-2-(3-pyridinyl)-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(2-pyridinyl)-4-pyrimidinamine

35 N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-2-pyridinyl-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-4-pyrimidinamine;

5 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-4-pyrimidinamine;
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;
10 6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine; and
N-[3-(1H-imidazol-1-yl)propyl]-6-(methoxymethyl)-2-(2-thienyl)-4-pyrimidinamine.

In another embodiment, the following compounds are disclaimed from the definition of formula

15 (I):
N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(3-pyridinyl)-4-pyrimidinamine;
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine;
20 2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine;
2-(3-chloro-4-fluorophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;
2-(4-bromophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-methyl-4-pyrimidinamine;
2-(4-bromophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine; and
2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine

25
In a further embodiment, the following compounds are disclaimed from the definition of formula (I):
N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(3-pyridinyl)-4-pyrimidinamine;
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
30 N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine;
2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine;
2-(3-chloro-4-fluorophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;
2-(4-bromophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-methyl-4-pyrimidinamine;
35 2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine;
2-(4-bromophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-4-pyridinyl-4-pyrimidinamine,
N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(4-pyridinyl)-4-pyrimidinamine;

5 N-[3-(1H-imidazol-1-yl)propyl]-6-(3-pyridinyl)-2-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-3-pyridinyl-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-(3-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(2-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(3-pyridinyl)-4-pyrimidinamine;

10 N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(2-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(3-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(2-pyridinyl)-2-(4-pyridinyl)-4-pyrimidinamine;

15 N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(2-pyridinyl)-2-(3-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(2-pyridinyl)-4-pyrimidinamine
N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-2-pyridinyl-4-pyrimidinamine;

20 N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-4-pyrimidinamine;

25 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-4-pyrimidinamine;
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;

30 6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-4-pyrimidinamine;
6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(methoxymethyl)-2-(2-thienyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(3-pyridinyl)-4-pyrimidinamine;
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;

35 N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine;
2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine;

2-(3-chloro-4-fluorophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;
2-(4-bromophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-methyl-4-pyrimidinamine;
2-(4-bromophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine; and
2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine

5

Certain known compounds

N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(3-pyridinyl)-4-pyrimidinamine,
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine and
N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(4-pyridinyl)-4-pyrimidinamine

10 are disclosed in Desai, P. V. et al., Journal of Medicinal Chemistry (2006), 49(5), p1576-1584 as potential parasitic cysteine protease inhibitors.

N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine and
2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine

15 are disclosed in Crespo Crespo, M. I. et al., WO 2005/058883 as potential adenosine receptor antagonists.

2-(3-chloro-4-fluorophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine,
2-(4-bromophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-methyl-4-pyrimidinamine and
20 2-(4-bromophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine
are disclosed in Yonetoku, Y. et al., WO 2003/026661 as potential treatments for insulin-dependent diabetes, non insulin-dependent diabetes, insulin-resistant diabetes and obesity.

25 2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine, which is disclosed in Schindler, U et al, DE 19836697 as a potential guanylate cyclase inhibitor.

When carbocyclyl and heterocyclyl are substituted, they are typically substituted by 1 or 2 substituents (e.g. 1 substituent). Typically the substituent is methyl. More typically carbocyclyl and heterocyclyl groups are unsubstituted.

30 When aryl and heteroaryl are substituted, they are typically substituted by 1, 2 or 3 (e.g. 1 or 2) substituents. Substituents for aryl and heteroaryl are selected from C₁₋₆alkyl (e.g. methyl), C₂₋₆alkenyl (e.g. buten-3-yl), C₂₋₆alkynyl (e.g. butyn-3-yl), C₁₋₆haloalkyl (e.g. fluoromethyl, trifluoromethyl), -C₁₋₆thioalkyl (e.g. -S-methyl), -SO₂C₁₋₄alkyl (e.g. -SO₂methyl), C₁₋₆alkoxy- (e.g. 35 methoxy, ethoxy), -O-C₃₋₈cycloalkyl (e.g. -O-cyclopentyl), C₃₋₈cycloalkyl (e.g. cyclopropyl, cyclohexyl), -SO₂C₃₋₈cycloalkyl (e.g. -SO₂cyclohexyl), C₃₋₆alkenyloxy- (e.g. -O-buten-2-yl), C₃₋₆alkynyloxy- (e.g. -O-buten-2-yl), -C(O)C₁₋₆alkyl (e.g. -C(O)ethyl), -C(O)OC₁₋₆alkyl (e.g.

-C(O)O-methyl), C₁₋₆alkoxy-C₁₋₆alkyl- (e.g. methoxy-ethyl-), nitro, halogen (e.g. fluoro, chloro, bromo), cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl (e.g. -NHmethyl), -N(C₁₋₄alkyl)(C₁₋₄alkyl) (e.g. -N(methyl)₂), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl) (e.g. -C(O)N(methyl)₂), -C(O)NH₂ and

-C(O)NH(C₁₋₄alkyl) (e.g. -C(O)NHmethyl). Further suitable examples are -C(O)OC₁₋₆alkyl (e.g. -

5 C(O)OMe, -SOC₁₋₄alkyl (e.g. SOMe) and -SOC₃₋₆cycloalkyl (e.g. -SO-cyclopropyl). More typically, substituents will be selected from C₁₋₆alkyl (e.g. methyl), C₁₋₆haloalkyl (e.g. C₁₋₆fluoroalkyl, e.g. CF₃), C₁₋₆alkoxy (e.g. OMe), halogen and hydroxy.

When R¹ represents alkyl, examples include methyl, ethyl, propyl, butyl, pentyl and hexyl,

10 especially methyl.

When R¹ represents haloalkyl, examples include fluoroalkyl e.g. trifluoromethyl.

When R¹ represents alkenyl, examples include propen-2-yl.

15 When R¹ represents optionally substituted carbocyclyl, examples include cyclopropyl, cyclopentyl, cyclohexyl and adamant-1-yl.

When R¹ represents optionally substituted heterocyclyl, exemplary heterocyclyl groups

20 containing two hetero atoms include pyrrolidine, tetrahydrofuran and piperidine, and exemplary heterocyclyl groups containing two hetero atoms include morpholine and piperazine. Further exemplary heterocyclyl groups include pyrrolidone, 2-methyl-pyrrolidine, 3-methyl-pyrrolidine, 3-methyl-tetrahydrofuran-, 3-methyl-piperidine-, 3-methyl-morpholine and 3-methyl-piperazine.

25 When R¹ represents optionally substituted aryl, examples include optionally substituted phenyl e.g. 2-methoxyphenyl; 3,4,5-trimethoxyphenyl; 3,4,5-bis(trifluoromethyl)phenyl; 3-methoxyphenyl; 4-methoxyphenyl; 4-methylphenyl; 4-nitrophenyl and unsubstituted phenyl.

When R¹ represents optionally substituted heteroaryl, examples include optionally substituted

30 pyridinyl e.g. pyridin-4-yl, optionally substituted furanyl e.g. furan-3-yl and quinolinyl e.g. quinolin-6-yl

When R¹ represents -carbocyclyl-phenyl, examples include 2-phenylcyclopropyl- and 2-tolylcyclopropyl-.

35 When R¹ represents -carbocyclyl-(monocyclic heteroaryl), examples include 2-pyridinyl-cyclopropyl-, 2-furanyl-cyclopropyl- 2-thiophenyl-cyclopropyl- and 2-pyrrolyl-cyclopropyl-.

When R¹ represents -heterocycl-phenyl, examples include 4-phenyl-tetrahydrofuran-2-yl-, 4-phenyl-tetrahydrothien-2-yl-, 4-phenyl-pyrrolidin-2-yl-, 4-phenyl-piperidin-2-yl, 5-phenyl-piperazin-2-yl, 5-phenyl-1,4-dioxan-2-yl and 5-phenyl-morpholin-2-yl.

5 When R¹ represents -carbocycl fused to phenyl, examples include 2,3-dihydro-1H-inden-2-yl. Further examples include tetralin-2-yl.

When R¹ represents carbocycl fused to monocyclic heteroaryl, examples include 5-membered 10 carbocycl fused to 5- or 6-membered monocyclic heteroaryl and 6-membered carbocycl fused to 5- or 6-membered monocyclic heteroaryl. The carbocycl ring may be fully or partially unsaturated.

Examples include 4,5-dihydrobenzofuran-5-yl, 4,5,6,7-tetrahydrobenzofuran-4-yl, 4,5,6,7-tetrahydrobenzofuran-5-yl, 4,5,6,7-tetrahydrobenzofuran-6-yl, 4,5,6,7-tetrahydrobenzofuran-7-15 yl, 4,5,6,7-tetrahydroindol-5-yl, 4,5,6,7-tetrahydroisoindol-5-yl, 4,5,6,7-tetrahydrobenzothiophen-5-yl, 4,5,6,7-tetrahydrobenzoimidazol-5-yl, 5,6,7,8-tetrahydroquinolin-6-yl, and 5,6,7,8-tetrahydroisoquinolin-6-yl.

When R¹ represents heterocycl fused to phenyl, examples include chromene, chromane, 20 isochromane, indoline, 3H-indole and isoindoline.

Suitably R¹ represents alkyl, optionally substituted aryl; carbocycl; haloalkyl or optionally substituted heteroaryl.

25 More suitably R¹ represents alkyl, optionally substituted aryl; carbocycl or haloalkyl.

Most suitably R¹ represents optionally substituted aryl, particularly substituted phenyl.

Particularly suitably, when R¹ is substituted phenyl, phenyl is substituted by one or more e.g. 1, 30 2 or 3 (e.g. at position(s) 4- or 2,4-, or 3,4-, or 3,4,5-) -C₁₋₄alkoxy groups e.g. methoxy or ethoxy groups, especially methoxy groups. For example, 3,4,5-trimethoxyphenyl is particularly suitable.

When R² represents alkyl, examples include methyl, ethyl, propyl, butyl, pentyl and hexyl, 35 especially methyl.

When R² represents haloalkyl, examples include fluoroalkyl e.g. trifluoromethyl.

When R^2 represents alkenyl, examples include propen-2-yl.

When R^2 represents optionally substituted carbocyclyl, examples include cyclopropyl,

5 cyclopentyl and cyclohexyl.

When R^2 represents optionally substituted heterocyclyl, exemplary heterocyclyl groups containing two hetero atoms include pyrrolidine, tetrahydrofuran and piperidine, and exemplary heterocyclyl groups containing two hetero atoms include morpholine and piperazine. Further

10 exemplary heterocyclyl groups include pyrrolidone, 2-methyl-pyrrolidine, 3-methyl-pyrrolidine, 3-methyl-tetrahydrofuran-, 3-methyl-piperidine-, 3-methyl-morpholine and 3-methyl-piperazine.

When R^2 represents optionally substituted aryl, examples include optionally substituted phenyl e.g. unsubstituted phenyl.

15

When R^2 represents optionally substituted heteroaryl, examples include optionally substituted pyridin-4-yl e.g. unsubstituted pyridin-4-yl.

When R^2 represents -carbocyclyl-phenyl, examples include 2-phenylcyclopropyl- and 2-

20 tolylcyclopropyl-.

When R^2 represents -carbocyclyl-(monocyclic heteroaryl), examples include 2-pyridinyl-cyclopropyl-, 2-furanyl-cyclopropyl- 2-thiophenyl-cyclopropyl- and 2-pyrrolyl-cyclopropyl-.

25 When R^2 represents -heterocyclyl-phenyl, examples include 4-phenyl-tetrahydrofuran-2-yl-, 4-phenyl-tetrahydrothien-2-yl-, 4-phenyl-pyrrolidin-2-yl-, 4-phenyl-piperidin-2-yl, 5-phenyl-piperazin-2-yl, 5-phenyl-1,4-dioxan-2-yl and 5-phenyl-morpholin-2-yl.

When R^2 represents -carbocyclyl fused to phenyl, examples include 2,3-dihydro-1H-inden-2-yl.

30 Further examples include tetralin-2-yl.

When R^2 represents carbocyclyl fused to monocyclic heteroaryl, examples include 5-membered carbocyclyl fused to 5- or 6-membered monocyclic heteroaryl and 6-membered carbocyclyl fused to 5- or 6-membered monocyclic heteroaryl. The carbocyclyl ring may be fully or partially

35 unsaturated.

Examples include 4,5-dihydrobenzofuran-5-yl, 4,5,6,7-tetrahydrobenzofuran-4-yl, 4,5,6,7-tetrahydrobenzofuran-5-yl, 4,5,6,7-tetrahydrobenzofuran-6-yl, 4,5,6,7-tetrahydrobenzofuran-7-

yl, 4,5,6,7-tetrahydroindol-5-yl, 4,5,6,7-tetrahydroisoindol-5-yl, 4,5,6,7-tetrahydrobenzothiophen-5-yl, 4,5,6,7-tetrahydrobenzoimidazol-5-yl, 5,6,7,8-tetrahydroquinolin-6-yl, and 5,6,7,8-tetrahydroisoquinolin-6-yl,


5 Suitably R^2 represents alkyl; optionally substituted aryl, or optionally substituted heteroaryl.

More suitably R^2 represents alkyl or optionally substituted aryl.

Most suitably R^2 represents alkyl or optionally substituted phenyl (e.g. unsubstituted phenyl).

10

Suitably A represents



R^3 suitably represents H.

R^4 suitably represents H or methyl.

15 R^5 suitably represents H or methyl.

In one embodiment of the invention, R^4 represents H and R^5 represents methyl. In another embodiment, R^4 represents methyl and R^5 represents H. In a third embodiment, R^4 represents H and R^5 represents H.

20 Suitably R^3 , R^4 and R^5 do not all represent H.

Suitably Y represents an unsubstituted C_{2-5} alkylene chain. In one embodiment, Y represents $-(CH_2)_2-$. In another embodiment, Y represents $-(CH_2)_3-$. In a third embodiment, Y represents $-(CH_2)_4-$. In further embodiment, Y represents $-(CH_2)_5-$. More suitably Y represents $-(CH_2)_2-$,

25 $-(CH_2)_4-$ or $-(CH_2)_5-$. In one embodiment, Y represents $-(CH_2)_3-$. In another embodiment, Y represents $-(CH_2)_4-$.


Alternatively Y represents a substituted C_{2-5} alkylene chain.

In one embodiment Y does not represent $-(CH_2)_3-$.

30

When Y represents a C₂₋₅ alkylene chain, which is substituted by two alkylene substituents at the same position wherein the two alkylene substituents are joined to each other to form a C₃-spiro-cycloalkyl group, the spiro-cycloalkyl group is suitably C₃spiro-cycloalkyl.

5 Alternatively A represents




In one embodiment R⁶ represents H and R⁷ represents H. In another embodiment R⁶ represents H and R⁷ represents C₁₋₂alkyl. In a third embodiment R⁶ represents C₁₋₂alkyl and R⁷ represents H.

Suitably Z represents a bond, -CH₂- or -CH₂CH₂-.


10 In one embodiment Z represents a bond. In another embodiment, Z represents -CH₂-.

In a third embodiment, Z represents -CH₂CH₂-.


15 More suitably A represents




or




or



20 Most suitably A represents




or




Most preferably A represents

5




particularly




B suitably represents H.

10

When benzimidazolyl is shown as benzimidazol-5-yl, which is represented as:




the person skilled in the art will appreciate that benzimidazol-6-yl, which is represented as:



is an equivalent structure. As employed herein, the two forms of benzimidazolyl are covered by the term "benzimidazol-5-yl".


5 Processes

A process for preparation of a compound of formula (I)



wherein A, B, R¹ and R² are as defined above

comprises the reaction of a compound of formula (II)




10

(II)


wherein X represents a leaving group (e.g. -O-tosylate, -O-mesylate or Cl).

with a compound of formula (III).




(III)

15 When A represents



and X represents -O-tosylate or -O-mesylate, the reaction may typically be carried out in an aprotic organic solvent (e.g. toluene) in the presence of a base (e.g. triethylamine).

Alternatively when A represents




5

and X represents Cl, the reaction may typically be carried out under Ullman-type conditions e.g. in the presence of copper iodide, N,N-diethylsalicylamide and potassium triphosphate at elevated temperature in a polar organic solvent (e.g. dimethylformamide or N-methylpyrrolidone).

10


When A represents



X may typically represent Cl and the reaction may typically be carried out in the presence of a base (e.g. sodium carbonate) at elevated temperature in a protic organic solvent (e.g. ethanol).

15

A compound of formula (II) may be prepared from a compound of formula (IV).




When X represents -O-tosylate, the reaction may typically be carried out in the presence of paratoluenesulfonyl chloride and a base (e.g. N-methyl morpholine) in an aprotic organic solvent (e.g. chloroform).

5 When X represents -O-mesylate, the reaction may typically be carried out in the presence of methanesulfonyl chloride and a base (e.g. triethylamine) in an aprotic organic solvent (e.g. chloroform).

When X represents Cl, the reaction may typically be carried out in the presence of phosphorous oxychloride at elevated temperature.


10

A compound of formula (IV) may be prepared from a compound of formula (V)



(V)

and a compound of formula (VI)



(VI)

15 wherein L represents a leaving group such as -O-ethyl.

The reaction may typically be carried out in the presence potassium tert-butoxide in an organic solvent (e.g. a polar, aprotic organic solvent such as ethanol).

Compounds of formulae (III), (V) and (VI) are either known or may be prepared by conventional methods known *per se*.

Therapeutic uses

5 In one embodiment, the compounds of formula (I) are suitable for use as a medicament.

In a specific embodiment, the compounds of formula (I) are suitable for use as a medicament, wherein the following compounds are disclaimed from the first pharmaceutical use and pharmaceutical composition claims:

10 N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(3-pyridinyl)-4-pyrimidinamine;
6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine;
2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine;
15 2-(3-chloro-4-fluorophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine;
2-(4-bromophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-methyl-4-pyrimidinamine;
2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine; and
2-(4-bromophenyl)-6-ethyl-N-[3-(1H-imidazol-1-yl)propyl]-4-pyrimidinamine.

20 In another embodiment the following compounds are also disclaimed from the first pharmaceutical use and pharmaceutical composition claims:

6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-4-pyridinyl-4-pyrimidinamine,
N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-6-(4-pyridinyl)-4-pyrimidinamine;
25 N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(3-pyridinyl)-2-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-3-pyridinyl-4-pyrimidinamine;

30 N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-(3-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-(2-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(2-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-(3-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(4-pyridinyl)-4-pyrimidinamine;
35 N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-6-(2-pyridinyl)-2-(4-pyridinyl)-4-pyrimidinamine;
N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;

N-[3-(1H-imidazol-1-yl)propyl]-6-(2-pyridinyl)-2-(3-pyridinyl)-4-pyrimidinamine;
 N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-2-(2-pyridinyl)-4-pyrimidinamine
 N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
 N-[3-(1H-imidazol-1-yl)propyl]-2,6-di-2-pyridinyl-4-pyrimidinamine;

5 N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
 N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-6-[3-(trifluoromethyl)phenyl]-4-pyrimidinamine;
 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-4-pyrimidinamine;
 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-4-pyrimidinamine;
 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-4-pyrimidinamine;

10 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine;
 6-(2-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;
 6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(4-pyridinyl)-4-pyrimidinamine;
 6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(3-pyridinyl)-4-pyrimidinamine;
 6-(5-bromo-3-pyridinyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-methyl-4-pyrimidinamine;

15 6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-(2-pyridinyl)-4-pyrimidinamine;
 6-(4-fluorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-2-phenyl-4-pyrimidinamine; and
 N-[3-(1H-imidazol-1-yl)propyl]-6-(methoxymethyl)-2-(2-thienyl)-4-pyrimidinamine.

Physiological substrates of QC (EC) in mammals are, e.g. amyloid beta-peptides (3-40), (3-42), (11-40) and (11-42), ABri, ADan, Gastrin, Neurotensin, FPP, CCL 2, CCL 7, CCL 8, CCL 16, CCL 18, Fractalkine, Orexin A, [Gln³]-glucagon(3-29), [Gln⁵]-substance P(5-11) and the peptide QYNAD. For further details see table 1. The compounds and/or combinations according to the present invention and pharmaceutical compositions comprising at least one inhibitor of QC (EC) are useful for the treatment of conditions that can be treated by modulation of QC activity.

25

Table 1: Amino acid sequences of physiological active peptides with an N-terminal glutamine residue, which are prone to be cyclized to final pGlu

Peptide	Amino acid sequence	Function
Abeta(1-42)	Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala	Plays a role in neurodegeneration, e.g. in Alzheimer's Disease, Familial British Dementia, Familial Danish Dementia, Down Syndrome

Peptide	Amino acid sequence	Function
Abeta(1-40)	Asp-Ala-Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Val	Plays a role in neurodegeneration, e.g. in Alzheimer's Disease, Familial British Dementia, Familial Danish Dementia, Down Syndrome
Abeta(3-42)	Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala	Plays a role in neurodegeneration, e.g. in Alzheimer's Disease, Familial British Dementia, Familial Danish Dementia, Down Syndrome
Abeta(3-40)	Glu-Phe-Arg-His-Asp-Ser-Gly-Tyr-Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val	Plays a role in neurodegeneration, e.g. in Alzheimer's Disease, Familial British Dementia, Familial Danish Dementia, Down Syndrome
Abeta(11-42)	Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val-Ile-Ala	Plays a role in neurodegeneration, e.g. in Alzheimer's Disease, Familial British Dementia, Familial Danish Dementia, Down Syndrome
Abeta(11-40)	Glu-Val-His-His-Gln-Lys-Leu-Val-Phe-Phe-Ala-Glu-Asp-Val-Gly-Ser-Asn-Lys-Gly-Ala-Ile-Ile-Gly-Leu-Met-Val-Gly-Gly-Val-Val	Plays a role in neurodegeneration, e.g. in Alzheimer's Disease, Familial British Dementia, Familial Danish Dementia, Down Syndrome
ABri	EASNCFA IRHFENKFAV ETLIC SRTVKKNIIIEEN	Pyroglutamated form plays a role in Familial British Dementia
ADan	EASNCFA IRHFENKFAV ETLIC FNLFLNSQEKHY	Pyroglutamated form plays a role in Familial Danish Dementia

Peptide	Amino acid sequence	Function
Gastrin 17 Swiss-Prot: P01350	QGPWL EEEEEAYGWM DF (amide)	Gastrin stimulates the stomach mucosa to produce and secrete hydrochloric acid and the pancreas to secrete its digestive enzymes. It also stimulates smooth muscle contraction and increases blood circulation and water secretion in the stomach and intestine.
Neurotensin Swiss-Prot: P30990	QLYENKPRRP YIL	Neurotensin plays an endocrine or paracrine role in the regulation of fat metabolism. It causes contraction of smooth muscle.
FPP	QEP amide	A tripeptide related to thyrotrophin releasing hormone (TRH), is found in seminal plasma. Recent evidence obtained <i>in vitro</i> and <i>in vivo</i> showed that FPP plays an important role in regulating sperm fertility.
TRH Swiss-Prot: P20396	QHP amide	TRH functions as a regulator of the biosynthesis of TSH in the anterior pituitary gland and as a neurotransmitter/neuromodulator in the central and peripheral nervous systems.
GnRH Swiss-Prot: P01148	QHWSYGL RP(G) amide	Stimulates the secretion of gonadotropins; it stimulates the secretion of both luteinizing and follicle-stimulating hormones.

Peptide	Amino acid sequence	Function
CCL16 (small inducible cytokine A16) Swiss-Prot: O15467	QPKVPEW VNTPSTCCLK YYEKVLPRRL VVGYRKALNC HLPAIIFVTK RNREVCTNPN DDWVQEYIKD PNLPLL PTRN LSTVKIITAK NGQPQLLNSQ	Shows chemotactic activity for lymphocytes and monocytes but not neutrophils. Also shows potent myelosuppressive activity, suppresses proliferation of myeloid progenitor cells. Recombinant SCYA16 shows chemotactic activity for monocytes and THP-1 monocytes, but not for resting lymphocytes and neutrophils. Induces a calcium flux in THP-1 cells that were desensitized by prior expression to RANTES.
CCL8 (small inducible cytokine A8) Swiss-Prot: P80075	QPDSVSI PITCCFNVIN RKIPIQRLES YTRITNIQCP KEAVIFKTKR GKEVCADPKE RWVRDSMKHL DQIFQNLKP	Chemotactic factor that attracts monocytes, lymphocytes, basophils and eosinophils. May play a role in neoplasia and inflammatory host responses. This protein can bind heparin.

Peptide	Amino acid sequence	Function
CCL2 (MCP-1, small inducible cytokine A2) Swiss-Prot: P13500	QPDAINA PVTCCYNFTN RKISVQRLAS YRRITSSKCP KEAVIFKTIV AKEICADPKQ KWVQDSMDHL DKQTQTPKT	Chemotactic factor that attracts monocytes and basophils but not neutrophils or eosinophils. Augments monocyte anti-tumor activity. Has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis or atherosclerosis. May be involved in the recruitment of monocytes into the arterial wall during the disease process of atherosclerosis. Binds to CCR2 and CCR4.
CCL18 (small inducible cytokine A18) Swiss-Prot: P55774	QVGTNKELC CLVYTSWQIP QKFIVDYSET SPQCPKPGVI LLTKRGRQIC ADPNKKWVQK YISDLKLNA	Chemotactic factor that attracts lymphocytes but not monocytes or granulocytes. May be involved in B cell migration into B cell follicles in lymph nodes. Attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes, has chemotactic activity for naive T cells, CD4+ and CD8+ T cells and thus may play a role in both humoral and cell-mediated immunity responses.

Peptide	Amino acid sequence	Function
Fractalkine (neurotactin) Swiss-Prot: P78423	QHHGVT KCNITCSKMT SKIPVALLIH YQQQNQASCAGK RAIILETRQH RLFCADPKEQ WVKDAMQHLD RQAAALTRNG GTFEKQIGEV KPRTPAAGG MDESVVLEPE ATGESSSLEP TPSSQEAQRA LGTSPELPTG VTGSSGTRLP PTPKAQDGGP VGTELFRVPP VSTAATWQSS APHQPGPSLW AEAKTSEAPS TQDPSTQAST ASSPAPEENA PSEGQRVWGQ GQSPRPENSL EREEMGPVPA HTDAFQDWGP GSMAHVSVP VSSEGTPSRE PVASGSWTPK AEEPIHATMD PQRLGVLITP VPDAQAAATTRR QAVGLLAFLG LLFCLGVAMF TYQSLQGCPK KMAGEMAEGL RYIPRSCGSN SYVLVPV	The soluble form is chemotactic for T cells and monocytes, but not for neutrophils. The membrane-bound form promotes adhesion of those leukocytes to endothelial cells. May play a role in regulating leukocyte adhesion and migration processes at the endothelium binds to CX3CR1.
CCL7 inducible A7) Swiss-Prot: P80098	(small cytokine QPVGINT STTCCYRFIN KKIPKQRLES YRRRTSSHCP REAVIFKTKL DKEICADPTQ KWVQDFMKHL DKKTQTPKL	Chemotactic factor that attracts monocytes and eosinophils, but not neutrophils. Augments monocyte anti-tumor activity. Also induces the release of gelatinase B. This protein can bind heparin. Binds to CCR1, CCR2 and CCR3.

Peptide	Amino acid sequence	Function
Orexin A (Hypocretin-1) Swiss-Prot O43612	QPLPDCCRQK TCSCRLYELL HGAGNHAAGI LTL	Neuropeptide that plays a significant role in the regulation of food intake and sleep-wakefulness, possibly by coordinating the complex behavioral and physiologic responses of these complementary homeostatic functions. It plays also a broader role in the homeostatic regulation of energy metabolism, autonomic function, hormonal balance and the regulation of body fluids. Orexin-A binds to both OX1R and OX2R with a high affinity.
Substance P	RPK PQQFFGLM	Belongs to the tachykinins. Tachykinins are active peptides which excite neurons, evoke behavioral responses, are potent vasodilators and secretagogues, and contract (directly or indirectly) many smooth muscles.
QYNAD	Gln-Tyr-Asn-Ala-Asp	Acts on voltage-gated sodium channels.

Glutamate is found in positions 3, 11 and 22 of the amyloid β -peptide. Among them the mutation from glutamic acid (E) to glutamine (Q) in position 22 (corresponding to amyloid precursor protein APP 693, Swissprot P05067) has been described as the so called Dutch type 5 cerebroarterial amyloidosis mutation.

The beta-amyloid peptides with a pyroglutamic acid residue in position 3, 11 and/or 22 have been described to be more cytotoxic and hydrophobic than the amyloid beta-peptides 1-40(42/43) (Saido T.C. 2000 Medical Hypotheses 54(3): 427-429).

10 The multiple N-terminal variations, e.g. Abeta(3-40), Abeta(3-42), Abeta(11-40) and Abeta (11-42) can be generated by the beta-secretase enzyme beta-site amyloid precursor protein-

cleaving enzyme (BACE) at different sites (Huse J.T. et al. 2002 *J. Biol. Chem.* 277 (18): 16278-16284), and/or by aminopeptidase or dipeptidylaminopeptidase processing from the full length peptides Abeta(1-40) and Abeta(1-42). In all cases, cyclization of the then N-terminal occurring glutamic acid residue is catalyzed by QC.

5 Transepithelial transducing cells, particularly the gastrin (G) cell, co-ordinate gastric acid secretion with the arrival of food in the stomach. Recent work showed that multiple active products are generated from the gastrin precursor, and that there are multiple control points in gastrin biosynthesis. Biosynthetic precursors and intermediates (progastrin and Gly-gastrins) 10 are putative growth factors; their products, the amidated gastrins, regulate epithelial cell proliferation, the differentiation of acid-producing parietal cells and histamine-secreting enterochromaffin-like (ECL) cells, and the expression of genes associated with histamine synthesis and storage in ECL cells, as well as acutely stimulating acid secretion. Gastrin also stimulates the production of members of the epidermal growth factor (EGF) family, which in turn 15 inhibit parietal cell function but stimulate the growth of surface epithelial cells. Plasma gastrin concentrations are elevated in subjects with *Helicobacter pylori*, who are known to have increased risk of duodenal ulcer disease and gastric cancer (Dockray, G.J. 1999 *J Physiol* 15 315-324).

20 The peptide hormone gastrin, released from antral G cells, is known to stimulate the synthesis and release of histamine from ECL cells in the oxyntic mucosa via CCK-2 receptors. The mobilized histamine induces acid secretion by binding to the H(2) receptors located on parietal cells. Recent studies suggest that gastrin, in both its fully amidated and less processed forms (progastrin and glycine-extended gastrin), is also a growth factor for the gastrointestinal tract. It 25 has been established that the major trophic effect of amidated gastrin is for the oxyntic mucosa of stomach, where it causes increased proliferation of gastric stem cells and ECL cells, resulting in increased parietal and ECL cell mass. On the other hand, the major trophic target of the less processed gastrin (e.g. glycine-extended gastrin) appears to be the colonic mucosa (Koh, T.J. and Chen, D. 2000 *Regul Pept* 93:37-44).

30 Neurotensin (NT) is a neuropeptide implicated in the pathophysiology of schizophrenia that specifically modulates neurotransmitter systems previously demonstrated to be misregulated in this disorder. Clinical studies in which cerebrospinal fluid (CSF) NT concentrations have been measured revealed a subset of schizophrenic patients with decreased CSF NT concentrations 35 that are restored by effective antipsychotic drug treatment. Considerable evidence also exists concordant with the involvement of NT systems in the mechanism of action of antipsychotic drugs. The behavioral and biochemical effects of centrally administered NT remarkably

resemble those of systemically administered antipsychotic drugs, and antipsychotic drugs increase NT neurotransmission. This concatenation of findings led to the hypothesis that NT functions as an endogenous antipsychotic. Moreover, typical and atypical antipsychotic drugs differentially alter NT neurotransmission in nigrostriatal and mesolimbic dopamine terminal 5 regions, and these effects are predictive of side effect liability and efficacy, respectively (Binder, E. B. et al. 2001 *Biol Psychiatry* 50 856-872).

Fertilization promoting peptide (FPP), a tripeptide related to thyrotrophin releasing hormone (TRH), is found in seminal plasma. Recent evidence obtained *in vitro* and *in vivo* showed that 10 FPP plays an important role in regulating sperm fertility. Specifically, FPP initially stimulates nonfertilizing (uncapacitated) spermatozoa to "switch on" and become fertile more quickly, but then arrests capacitation so that spermatozoa do not undergo spontaneous acrosome loss and therefore do not lose fertilizing potential. These responses are mimicked, and indeed augmented, by adenosine, known to regulate the adenylyl cyclase (AC)/cAMP signal 15 transduction pathway. Both FPP and adenosine have been shown to stimulate cAMP production in uncapacitated cells but inhibit it in capacitated cells, with FPP receptors somehow interacting with adenosine receptors and G proteins to achieve regulation of AC. These events affect the tyrosine phosphorylation state of various proteins, some being important in the initial "switching on," others possibly being involved in the acrosome reaction itself. Calcitonin and 20 angiotensin II, also found in seminal plasma, have similar effects *in vitro* on uncapacitated spermatozoa and can augment responses to FPP. These molecules have similar effects *in vivo*, affecting fertility by stimulating and then maintaining fertilizing potential. Either reductions in the availability of FPP, adenosine, calcitonin, and angiotensin II or defects in their receptors 25 contribute to male infertility (Fraser, L.R. and Adeoya-Osiguwa, S. A. 2001 *Vitam Horm* 63, 1-28).

CCL2 (MCP-1), CCL7, CCL8, CCL16, CCL18 and fractalkine play an important role in pathophysiological conditions, such as suppression of proliferation of myeloid progenitor cells, neoplasia, inflammatory host responses, cancer, psoriasis, rheumatoid arthritis, atherosclerosis, 30 vasculitis, humoral and cell-mediated immunity responses, leukocyte adhesion and migration processes at the endothelium, inflammatory bowel disease, restenosis, pulmonary fibrosis, pulmonary hypertension, liver fibrosis, liver cirrhosis, nephrosclerosis, ventricular remodeling, heart failure, arteriopathy after organ transplantations and failure of vein grafts.

35 A number of studies have underlined in particular the crucial role of MCP-1 for the development of atherosclerosis (Gu, L., et al., (1998) *Mol.Cell* 2, 275-281; Gosling, J., et al., (1999) *J Clin.Invest* 103, 773-778); rheumatoid arthritis (Gong, J. H., et al., (1997) *J Exp.Med* 186, 131-

137; Ogata, H., et al., (1997) *J Pathol.* 182, 106-114); pancreatitis (Bhatia, M., et al., (2005) *Am.J Physiol Gastrointest.Liver Physiol* 288, G1259-G1265); Alzheimer's disease (Yamamoto, M., et al., (2005) *Am.J Pathol.* 166, 1475-1485); lung fibrosis (Inoshima, I., et al., (2004) *Am.J Physiol Lung Cell Mol.Physiol* 286, L1038-L1044); renal fibrosis (Wada, T., et al., (2004) *J Am.Soc.Nephrol.* 15, 940-948), and graft rejection (Saiura, A., et al., (2004) *Arterioscler. Thromb. Vasc. Biol.* 24, 1886-1890). Furthermore, MCP-1 might also play a role in gestosis (Katabuchi, H., et al., (2003) *Med Electron Microsc.* 36, 253-262), as a paracrine factor in tumor development (Ohta, M., et al., (2003) *Int.J Oncol.* 22, 773-778; Li, S., et al., (2005) *J Exp.Med* 202, 617-624), neuropathic pain (White, F. A., et al., (2005) *Proc. Natl. Acad.Sci.U.S.A*) and AIDS (Park, I. W., Wang, J. F., and Groopman, J. E. (2001) *Blood* 97, 352-358; Coll, B., et al., (2006) *Cytokine* 34, 51-55).

MCP-1 levels are increased in CSF of AD patients and patients showing mild cognitive impairment (MCI) (Galimberti, D., et al., (2006) *Arch.Neurol.* 63, 538-543). Furthermore, MCP-1 shows an increased level in serum of patients with MCI and early AD (Clerici, F., et al., (2006) *Neurobiol.Aging* 27, 1763-1768).

Several cytotoxic T lymphocyte peptide-based vaccines against hepatitis B, human immunodeficiency virus and melanoma were recently studied in clinical trials. One interesting melanoma vaccine candidate alone or in combination with other tumor antigens, is the decapeptide ELA. This peptide is a Melan-A/MART-1 antigen immunodominant peptide analog, with an N-terminal glutamic acid. It has been reported that the amino group and gamma-carboxylic group of glutamic acids, as well as the amino group and gamma-carboxamide group of glutamines, condense easily to form pyroglutamic derivatives. To overcome this stability problem, several peptides of pharmaceutical interest have been developed with a pyroglutamic acid instead of N-terminal glutamine or glutamic acid, without loss of pharmacological properties. Unfortunately compared with ELA, the pyroglutamic acid derivative (PyrELA) and also the N-terminal acetyl-capped derivative (AcELA) failed to elicit cytotoxic T lymphocyte (CTL) activity. Despite the apparent minor modifications introduced in PyrELA and AcELA, these two derivatives probably have lower affinity than ELA for the specific class I major histocompatibility complex. Consequently, in order to conserve full activity of ELA, the formation of PyrELA must be avoided (Beck A. et al. 2001, *J Pept Res* 57(6):528-38.).

Orexin A is a neuropeptide that plays a significant role in the regulation of food intake and sleep-wakefulness, possibly by coordinating the complex behavioral and physiologic responses of these complementary homeostatic functions. It plays also a role in the homeostatic regulation of energy metabolism, autonomic function, hormonal balance and the regulation of body fluids.

Recently, increased levels of the pentapeptide QYNAD were identified in the cerebrospinal fluid (CSF) of patients suffering from multiple sclerosis or Guillain-Barré syndrome compared to healthy individuals (Brinkmeier H. et al. 2000, *Nature Medicine* 6, 808-811). There is a big

5 controversy in the literature about the mechanism of action of the pentapeptide Gln-Tyr-Asn-Ala-Asp (QYNAD), especially its efficacy to interact with and block sodium channels resulting in the promotion of axonal dysfunction, which are involved in inflammatory autoimmune diseases of the central nervous system. But recently, it could be demonstrated that not QYNAD, but its cyclized, pyroglutamated form, pEYNAD, is the active form, which blocks sodium channels
10 resulting in the promotion of axonal dysfunction. Sodium channels are expressed at high density in myelinated axons and play an obligatory role in conducting action potentials along axons within the mammalian brain and spinal cord. Therefore, it is speculated that they are involved in several aspects of the pathophysiology of inflammatory autoimmune diseases, especially multiple sclerosis, the Guillain-Barré syndrome and chronic inflammatory demyelinizing
15 polyradiculoneuropathy.

Furthermore, QYNAD is a substrate of the enzyme glutaminyl cyclase (QC, EC 2.3.2.5), which is also present in the brain of mammals, especially in human brain. Glutaminyl cyclase catalyzes effectively the formation of pEYNAD from its precursor QYNAD.

20 Accordingly, the present invention provides the use of the compounds of formula (I) for the preparation of a medicament for the prevention or alleviation or treatment of a disease selected from the group consisting of mild cognitive impairment, Alzheimer's disease, Familial British Dementia, Familial Danish Dementia, neurodegeneration in Down Syndrome, Huntington's disease, Kennedy's disease, ulcer disease, duodenal cancer with or w/o *Helicobacter pylori*
25 infections, colorectal cancer, Zollinger-Ellison syndrome, gastric cancer with or without *Helicobacter pylori* infections, pathogenic psychotic conditions, schizophrenia, infertility, neoplasia, inflammatory host responses, cancer, malign metastasis, melanoma, psoriasis, rheumatoid arthritis, atherosclerosis, pancreatitis, restenosis, impaired humoral and cell-mediated immune responses, leukocyte adhesion and migration processes in the endothelium, impaired food intake, impaired sleep-wakefulness, impaired homeostatic regulation of energy metabolism, impaired autonomic function, impaired hormonal balance or impaired regulation of body fluids, multiple sclerosis, the Guillain-Barré syndrome and chronic inflammatory demyelinizing polyradiculoneuropathy.

30 35 Furthermore, by administration of a compound according to the present invention to a mammal it can be possible to stimulate the proliferation of myeloid progenitor cells.

In addition, the administration of a QC inhibitor according to the present invention can lead to suppression of male fertility.

In a preferred embodiment, the present invention provides the use of inhibitors of QC (EC) 5 activity in combination with other agents, especially for the treatment of neuronal diseases, atherosclerosis and multiple sclerosis.

The present invention also provides a method of treatment of the aforementioned diseases comprising the administration of a therapeutically active amount of at least one compound of 10 formula (I) to a mammal, preferably a human.

Most preferably, said method and corresponding uses are for the treatment of a disease selected from the group consisting of mild cognitive impairment, Alzheimer's disease, Familial British Dementia, Familial Danish Dementia, neurodegeneration in Down Syndrome, Parkinson 15 disease and Chorea Huntington, comprising the administration of a therapeutically active amount of at least one compound of formula (I) to a mammal, preferably a human.

Even preferably, the present invention provides a method of treatment and corresponding uses for the treatment of rheumatoid arthritis, atherosclerosis, pancreatitis and restenosis.

20 **Pharmaceutical combinations**
In a preferred embodiment, the present invention provides a composition, preferably a pharmaceutical composition, comprising at least one QC inhibitor optionally in combination with at least one other agent selected from the group consisting of nootropic agents, 25 neuroprotectants, antiparkinsonian drugs, amyloid protein deposition inhibitors, beta amyloid synthesis inhibitors, antidepressants, anxiolytic drugs, antipsychotic drugs and anti-multiple sclerosis drugs.

Most preferably, said QC inhibitor is a compound of formula (I) of the present invention.

30 More specifically, the aforementioned other agent is selected from the group consisting of beta-amyloid antibodies, cysteine protease inhibitors, PEP-inhibitors, LiCl, acetylcholinesterase (AChE) inhibitors, PIMT enhancers, inhibitors of beta secretases, inhibitors of gamma secretases, inhibitors of aminopeptidases, preferably inhibitors of dipeptidyl peptidases, most 35 preferably DP IV inhibitors; inhibitors of neutral endopeptidase, inhibitors of Phosphodiesterase-4 (PDE-4), TNFalpha inhibitors, muscarinic M1 receptor antagonists, NMDA receptor antagonists, sigma-1 receptor inhibitors, histamine H3 antagonists, immunomodulatory

agents, immunosuppressive agents, MCP-1 antagonists or an agent selected from the group consisting of antegren (natalizumab), Neurelan (fampridine-SR), campath (alemtuzumab), IR 208, NBI 5788/MSP 771 (tadalafil), paclitaxel, Anergix.MS (AG 284), SH636, Differin (CD 271, adapalene), BAY 361677 (interleukin-4), matrix-metalloproteinase-inhibitors (e.g. BB 5 76163), interferon-tau (trophoblastin) and SAIK-MS.

Furthermore, the other agent may be, for example, an anti-anxiety drug or antidepressant selected from the group consisting of

- (a) Benzodiazepines, e.g. alprazolam, chlordiazepoxide, clobazam, clonazepam, 10 clorazepate, diazepam, fludiazepam, loflazepate, lorazepam, methaqualone, oxazepam, prazepam, traxene,
- (b) Selective serotonin re-uptake inhibitors (SSRI's), e.g. citalopram, fluoxetine, fluvoxamine, escitalopram, sertraline, paroxetine,
- (c) Tricyclic antidepressants, e.g. amitriptyline, clomipramine, desipramine, doxepin, 15 imipramine
- (d) Monoamine oxidase (MAO) inhibitors,
- (e) Azapirones, e.g. buspirone, tandospirone,
- (f) Serotonin-norepinephrine reuptake inhibitors (SNRI's), e.g. venlafaxine, duloxetine,
- (g) Mirtazapine,
- (h) Norepinephrine reuptake inhibitors (NRI's), e.g. reboxetine,
- (i) Bupropione,
- (j) Nefazodone,
- (k) beta-blockers,
- (l) NPY-receptor ligands: NPY agonists or antagonists.

25 In a further embodiment, the other agent may be, for example, an anti-multiple sclerosis drug selected from the group consisting of

- a) dihydroorotate dehydrogenase inhibitors, e.g. SC-12267, teriflunomide, MNA-715, HMR- 30 1279 (syn. to HMR-1715, MNA-279),
- b) autoimmune suppressant, e.g. laquinimod,
- c) paclitaxel,
- d) antibodies, e.g. AGT-1, anti-granulocyte-macrophage colony-stimulating factor (GM-CSF) monoclonal antibody, Nogo receptor modulators, ABT-874, alemtuzumab (CAMPATH), anti-OX40 antibody, CNTO-1275, DN-1921, natalizumab (syn. to AN- 35 100226, Antegren, VLA-4 Mab), daclizumab (syn. to Zenepax, Ro-34-7375, SMART

anti-Tac), J-695, priliximab (syn. to Centara, CEN-000029, cM-T412), MRA, Dantes, anti-IL-12-antibody,

5 e) peptide nucleic acid (PNA) preparations, e.g. reticulose,

f) interferon alpha, e.g. Alfaferone, human alpha interferon (syn. to Omnipferon, Alpha Leukoferon),

g) interferon beta, e.g. Frone, interferon beta-1a like Avonex, Betron (Rebif), interferon beta analogs, interferon beta-transferrin fusion protein, recombinant interferon beta-1b like Betaseron,

h) interferon tau,

10 i) peptides, e.g. AT-008, AnergiX.MS, Immunokine (alpha-Immunokine-NNSO3), cyclic peptides like ZD-7349,

j) therapeutic enzymes, e.g. soluble CD8 (sCD8),

k) multiple sclerosis-specific autoantigen-encoding plasmid and cytokine-encoding plasmid, e.g. BHT-3009;

15 l) inhibitor of TNF-alpha, e.g. BLX-1002, thalidomide, SH-636,

m) TNF antagonists, e.g. solimastat, lenercept (syn. to RO-45-2081, Tenefuse), onercept (sTNFR1), CC-1069,

n) TNF alpha, e.g. etanercept (syn. to Enbrel, TNR-001)

o) CD28 antagonists, e.g. abatacept,

20 p) Lck tyrosine kinase inhibitors,

q) cathepsin K inhibitors,

r) analogs of the neuron-targeting membrane transporter protein taurine and the plant-derived calpain inhibitor leupeptin, e.g. Neurodur,

s) chemokine receptor-1 (CCR1) antagonist, e.g. BX-471,

25 t) CCR2 antagonists,

u) AMPA receptor antagonists, e.g. ER-167288-01 and ER-099487, E-2007, talampanel,

v) potassium channel blockers, e.g. fampridine,

w) tosyl-proline-phenylalanine small-molecule antagonists of the VLA-4/VCAM interaction, e.g. TBC-3342,

30 x) cell adhesion molecule inhibitors, e.g. TBC-772,

y) antisense oligonucleotides, e.g. EN-101,

z) antagonists of free immunoglobulin light chain (IgLC) binding to mast cell receptors, e.g. F-991,

aa) apoptosis inducing antigens, e.g. Apogen MS,

35 bb) alpha-2 adrenoceptor agonist, e.g. tizanidine (syn. to Zanaflex, Ternelin, Sirdalvo, Sirdalud, Mionidine),

- cc) copolymer of L-tyrosine, L-lysine, L-glutamic acid and L-alanine, e.g. glatiramer acetate (syn. to Copaxone, COP-1, copolymer-1),
- dd) topoisomerase II modulators, e.g. mitoxantrone hydrochloride,
- ee) adenosine deaminase inhibitor, e.g. cladribine (syn. to Leustatin, Mylinax, RWJ-26251),
- 5 ff) interleukin-10, e.g. ilodecakin (syn. to Tenovil, Sch-52000, CSIF),
- gg) interleukin-12 antagonists, e.g. lisofylline (syn. to CT-1501R, LSF, lysofylline),
- hh) Ethanaminum, e.g. SRI-62-834 (syn. to CRC-8605, NSC-614383),
- ii) immunomodulators, e.g. SAIK-MS, PNU-156804, alpha-fetoprotein peptide (AFP), IPDS,
- jj) retinoid receptor agonists, e.g. adapalene (syn. to Differin, CD-271),
- 10 kk) TGF-beta, e.g. GDF-1 (growth and differentiation factor 1),
- ll) TGF-beta-2, e.g. BetaKine,
- mm) MMP inhibitors, e.g. glycomed,
- nn) phosphodiesterase 4 (PDE4) inhibitors, e.g. RPR-122818,
- oo) purine nucleoside phosphorylase inhibitors, e.g. 9-(3-pyridylmethyl)-9-deazaguanine,
- 15 peldesine (syn. to BCX-34, TO-200),
- pp) alpha-4/beta-1 integrin antagonists, e.g. ISIS-104278,
- qq) antisense alpha-4 integrin (CD49d), e.g. ISIS-17044, ISIS-27104,
- rr) cytokine-inducing agents, e.g. nucleosides, ICN-17261,
- ss) cytokine inhibitors,
- 20 tt) heat shock protein vaccines, e.g. HSPPC-96,
- uu) neuregulin growth factors, e.g. GGF-2 (syn. to neuregulin, glial growth factor 2),
- vv) cathepsin S – inhibitors,
- ww) broprimine analogs, e.g. PNU-56169, PNU-63693,
- xx) Monocyte chemoattractant protein-1 inhibitors, e.g. benzimidazoles like MCP-1
- 25 inhibitors, LKS-1456, PD-064036, PD-064126, PD-084486, PD-172084, PD-172386.

Further, the present invention provides pharmaceutical compositions e.g. for parenteral, enteral or oral administration, comprising at least one QC inhibitor, optionally in combination with at least one of the other aforementioned agents.

30 These combinations provide a particularly beneficial effect. Such combinations are therefore shown to be effective and useful for the treatment of the aforementioned diseases. Accordingly, the invention provides a method for the treatment of these conditions.

35 The method comprises either co-administration of at least one QC inhibitor and at least one of the other agents or the sequential administration thereof.

Co-administration includes administration of a formulation, which comprises at least one QC inhibitor and at least one of the other agents or the essentially simultaneous administration of separate formulations of each agent.

5 Beta-amyloid antibodies and compositions containing the same are described, e.g. in WO 2006/137354, WO 2006/118959, WO 2006/103116, WO 2006/095041, WO 2006/081171, WO 2006/066233, WO 2006/066171, WO 2006/066089, WO 2006/066049, WO 2006/055178, WO 2006/046644, WO 2006/039470, WO 2006/036291, WO 2006/026408, WO 2006/016644, WO 10 2006/014638, WO 2006/014478, WO 2006/008661, WO 2005/123775, WO 2005/120571, WO 2005/105998, WO 2005/081872, WO 2005/080435, WO 2005/028511, WO 2005/025616, WO 2005/025516, WO 2005/023858, WO 2005/018424, WO 2005/011599, WO 2005/000193, WO 2004/108895, WO 2004/098631, WO 2004/080419, WO 2004/071408, WO 2004/069182, WO 2004/067561, WO 2004/044204, WO 2004/032868, WO 2004/031400, WO 2004/029630, WO 15 2004/029629, WO 2004/024770, WO 2004/024090, WO 2003/104437, WO 2003/089460, WO 2003/086310, WO 2003/077858, WO 2003/074081, WO 2003/070760, WO 2003/063760, WO 2003/055514, WO 2003/051374, WO 2003/048204, WO 2003/045128, WO 2003/040183, WO 2003/039467, WO 2003/016466, WO 2003/015691, WO 2003/014162, WO 2003/012141, WO 2002/088307, WO 2002/088306, WO 2002/074240, WO 2002/046237, WO 2002/046222, WO 20 2002/041842, WO 2001/062801, WO 2001/012598, WO 2000/077178, WO 2000/072880, WO 2000/063250, WO 1999/060024, WO 1999/027944, WO 1998/044955, WO 1996/025435, WO 1994/017197, WO 1990/014840, WO 1990/012871, WO 1990/012870, WO 1989/006242.

25 The beta-amyloid antibodies may be selected from, for example, polyclonal, monoclonal, chimeric or humanized antibodies. Furthermore, said antibodies may be useful to develop active and passive immune therapies, i.e. vaccines and monoclonal antibodies.

Suitable examples of beta-amyloid antibodies are ACU-5A5, huC091 (Acumen/Merck); PF-4360365, RI-1014, RI-1219, RI-409, RN-1219 (Rinat Neuroscience Corp (Pfizer Inc)); the nanobody therapeutics of Ablynx/Boehringer Ingelheim; beta-amyloid-specific humanized 30 monoclonal antibodies of Intellect Neurosciences/IBL; m266, m266.2 (Eli Lilly & Co.); AAB-02 (Elan); bapineuzumab (Elan); BAN-2401 (Bioarctic Neuroscience AB); ABP-102 (Abiogen Pharma SpA); BA-27, BC-05 (Takeda); R-1450 (Roche); ESB-212 (ESBATech AG); AZD-3102 (AstraZeneca) and beta-amyloid antibodies of Mindset BioPharmaceuticals Inc.

35 Especially preferred are antibodies, which recognize the N-terminus of the A β peptide. A suitable antibody, which recognizes the A β -N-Terminus is, for example Acl-24 (AC Immune SA).

A monoclonal antibody against beta-amyloid peptide is disclosed in WO 2007/068412. Respective chimeric and humanized antibodies are disclosed in WO 2008/011348. A method for producing a vaccine composition for treating an amyloid-associated disease is disclosed in WO 2007/068411.

5 Suitable cysteine protease inhibitors are inhibitors of cathepsin B. Inhibitors of cathepsin B and compositions containing such inhibitors are described, e.g. in WO 2006/060473, WO 2006/042103, WO 2006/039807, WO 2006/021413, WO 2006/021409, WO 2005/097103, WO 2005/007199, WO2004/084830, WO 2004/078908, WO 2004/026851, WO 2002/094881, WO 10 2002/027418, WO 2002/021509, WO 1998/046559, WO 1996/021655.

Examples of suitable PIMT enhancers are 10-aminoaliphatyl-dibenz[b, f] oxepines described in WO 98/15647 and WO 03/057204, respectively. Further useful according to the present invention are modulators of PIMT activity described in WO 2004/039773.

15 Inhibitors of beta secretase and compositions containing such inhibitors are described, e.g. in WO03/059346, WO2006/099352, WO2006/078576, WO2006/060109, WO2006/057983, WO2006/057945, WO2006/055434, WO2006/044497, WO2006/034296, WO2006/034277, WO2006/029850, WO2006/026204, WO2006/014944, WO2006/014762, WO2006/002004, US 20 7,109,217, WO2005/113484, WO2005/103043, WO2005/103020, WO2005/065195, WO2005/051914, WO2005/044830, WO2005/032471, WO2005/018545, WO2005/004803, WO2005/004802, WO2004/062625, WO2004/043916, WO2004/013098, WO03/099202, WO03/043987, WO03/039454, US 6,562,783, WO02/098849 and WO02/096897.

25 Suitable examples of beta secretase inhibitors for the purpose of the present invention are WY-25105 (Wyeth); Posiphen, (+)-phenserine (TorreyPines / NIH); LSN-2434074, LY-2070275, LY-2070273, LY-2070102 (Eli Lilly & Co.); PNU-159775A, PNU-178025A, PNU-17820A, PNU-33312, PNU-38773, PNU-90530 (Elan / Pfizer); KMI-370, KMI-358, kmi-008 (Kyoto University); OM-99-2, OM-003 (Athenagen Inc.); AZ-12304146 (AstraZeneca / Astex); GW-840736X 30 (GlaxoSmithKline plc.), DNP-004089 (De Novo Pharmaceuticals Ltd.) and CT-21166 (CoMentis Inc.).

Inhibitors of gamma secretase and compositions containing such inhibitors are described, e.g. in WO2005/008250, WO2006/004880, US 7,122,675, US 7,030,239, US 6,992,081, US 35 6,982,264, WO2005/097768, WO2005/028440, WO2004/101562, US 6,756,511, US 6,683,091, WO03/066592, WO03/014075, WO03/013527, WO02/36555, WO01/53255, US 7,109,217, US 7,101,895, US 7,049,296, US 7,034,182, US 6,984,626, WO2005/040126, WO2005/030731,

WO2005/014553, US 6,890,956, EP 1334085, EP 1263774, WO2004/101538, WO2004/00958, WO2004/089911, WO2004/073630, WO2004/069826, WO2004/039370, WO2004/031139, WO2004/031137, US 6,713,276, US 6,686,449, WO03/091278, US 6,649,196, US 6,448,229, WO01/77144 and WO01/66564.

5

Suitable gamma secretase inhibitors for the purpose of the present invention are GSI-953, WAY-GSI-A, WAY-GSI-B (Wyeth); MK-0752, MRK-560, L-852505, L-685-458, L-852631, L-852646 (Merck & Co. Inc.); LY-450139, LY-411575, AN-37124 (Eli Lilly & Co.); BMS-299897, BMS-433796 (Bristol-Myers Squibb Co.); E-2012 (Eisai Co. Ltd.); EHT-0206, EHT-206

10 (ExonHit Therapeutics SA); and NGX-555 (TorreyPines Therapeutics Inc.).

DP IV-inhibitors and compositions containing such inhibitors are described, e.g. in US6,011,155; US6,107,317; US6,110,949; US6,124,305; US6,172,081; WO99/61431, WO99/67278, WO99/67279, DE19834591, WO97/40832, WO95/15309, WO98/19998,

15 WO00/07617, WO99/38501, WO99/46272, WO99/38501, WO01/68603, WO01/40180, WO01/81337, WO01/81304, WO01/55105, WO02/02560, WO01/34594, WO02/38541, WO02/083128, WO03/072556, WO03/002593, WO03/000250, WO03/000180, WO03/000181, EP1258476, WO03/002553, WO03/002531, WO03/002530, WO03/004496, WO03/004498, WO03/024942, WO03/024965, WO03/033524, WO03/035057, WO03/035067, WO03/037327,

20 WO03/040174, WO03/045977, WO03/055881, WO03/057144, WO03/057666, WO03/068748, WO03/068757, WO03/082817, WO03/101449, WO03/101958, WO03/104229, WO03/74500, WO2004/007446, WO2004/007468, WO2004/018467, WO2004/018468, WO2004/018469, WO2004/026822, WO2004/032836, WO2004/033455, WO2004/037169, WO2004/041795, WO2004/043940, WO2004/048352, WO2004/050022, WO2004/052850, WO2004/058266,

25 WO2004/064778, WO2004/069162, WO2004/071454, WO2004/076433, WO2004/076434, WO2004/087053, WO2004/089362, WO2004/099185, WO2004/103276, WO2004/103993, WO2004/108730, WO2004/110436, WO2004/111041, WO2004/112701, WO2005/000846, WO2005/000848, WO2005/011581, WO2005/016911, WO2005/023762, WO2005/025554, WO2005/026148, WO2005/030751, WO2005/033106, WO2005/037828, WO2005/040095,

30 WO2005/044195, WO2005/047297, WO2005/051950, WO2005/056003, WO2005/056013, WO2005/058849, WO2005/075426, WO2005/082348, WO2005/085246, WO2005/087235, WO2005/095339, WO2005/095343, WO2005/095381, WO2005/108382, WO2005/113510, WO2005/116014, WO2005/116029, WO2005/118555, WO2005/120494, WO2005/121089, WO2005/121131, WO2005/123685, WO2006/995613; WO2006/009886; WO2006/013104;

35 WO2006/017292; WO2006/019965; WO2006/020017; WO2006/023750; WO2006/039325; WO2006/041976; WO2006/047248; WO2006/058064; WO2006/058628; WO2006/066747; WO2006/066770 and WO2006/068978.

Suitable DP IV-inhibitors for the purpose of the present invention are for example Sitagliptin, des-fluoro-sitagliptin (Merck & Co. Inc.); vildagliptin, DPP-728, SDZ-272-070 (Novartis) ; ABT-279, ABT-341 (Abbott Laboratories); denagliptin, TA-6666 (GlaxoSmithKline plc.); SYR-322

5 (Takeda San Diego Inc.); talabostat (Point Therapeutics Inc.); Ro-0730699, R-1499, R-1438 (Roche Holding AG); FE-999011 (Ferring Pharmaceuticals); TS-021 (Taisho Pharmaceutical Co. Ltd.); GRC-8200 (Glenmark Pharmaceuticals Ltd.); ALS-2-0426 (Alantos Pharmaceuticals Holding Inc.); ARI-2243 (Arisaph Pharmaceuticals Inc.); SSR-162369 (Sanofi-Synthelabo); MP-513 (Mitsubishi Pharma Corp.); DP-893, CP-867534-01 (Pfizer Inc.); TSL-225, TMC-2A
10 (Tanabe Seiyaku Co. Ltd.); PHX-1149 (Phenomenix Corp.); saxagliptin (Bristol-Myers Squibb Co.); PSN-9301 ((OSI) Prosidion), S-40755 (Servier); KRP-104 (ActivX Biosciences Inc.); sulphostin (Zaidan Hojin); KR-62436 (Korea Research Institute of Chemical Technology); P32/98 (Probiodrug AG); BI-A, BI-B (Boehringer Ingelheim Corp.); SK-0403 (Sanwa Kagaku Kenkyusho Co. Ltd.); and NNC-72-2138 (Novo Nordisk A/S).

15 Other preferred DP IV-inhibitors are
(i) dipeptide-like compounds, disclosed in WO 99/61431, e.g. N-valyl prolyl, O-benzoyl hydroxylamine, alanyl pyrrolidine, isoleucyl thiazolidine like L-allo-isoleucyl thiazolidine, L-threo-isoleucyl pyrrolidine and salts thereof, especially the fumaric salts, and L-allo-isoleucyl
20 pyrrolidine and salts thereof;
(ii) peptide structures, disclosed in WO 03/002593, e.g. tripeptides;
(iii) peptidylketones, disclosed in WO 03/033524;
(vi) substituted aminoketones, disclosed in WO 03/040174;
(v) topically active DP IV-inhibitors, disclosed in WO 01/14318;
25 (vi) prodrugs of DP IV-inhibitors, disclosed in WO 99/67278 and WO 99/67279; and
(v) glutamyl based DP IV-inhibitors, disclosed in WO 03/072556 and WO 2004/099134.

Suitable beta amyloid synthesis inhibitors for the purpose of the present invention are for example Bisnorcymserine (Axonyx Inc.); (R)-flurbiprofen (MCP-7869; Flurizan) (Myriad Genetics); nitroflurbiprofen (NicOx); BGC-20-0406 (Sankyo Co. Ltd.) and BGC-20-0466 (BTG plc.).

Suitable amyloid protein deposition inhibitors for the purpose of the present invention are for example SP-233 (Samaritan Pharmaceuticals); AZD-103 (Ellipsis Neurotherapeutics Inc.); AAB-35 001 (Bapineuzumab), AAB-002, ACC-001 (Elan Corp plc.); Colostrinin (ReGen Therapeutics plc.); Tramiprosate (Neurochem); AdPED1-(amyloid-beta1-6)11 (Vaxin Inc.); MPI-127585, MPI-423948 (Mayo Foundation); SP-08 (Georgetown University); ACU-5A5 (Acumen / Merck);

Transthyretin (State University of New York); PTI-777, DP-74, DP 68, Exebryl (ProteoTech Inc.); m266 (Eli Lilly & Co.); EGb-761 (Dr. Willmar Schwabe GmbH); SPI-014 (Satori Pharmaceuticals Inc.); ALS-633, ALS-499 (Advanced Life Sciences Inc.); AGT-160 (ArmaGen Technologies Inc.); TAK-070 (Takeda Pharmaceutical Co. Ltd.); CHF-5022, CHF-5074, CHF-5096 and CHF-5 5105 (Chiesi Farmaceutici SpA.).

Suitable PDE-4 inhibitors for the purpose of the present invention are for example Doxofylline (Instituto Biologico Chemioterapica ABC SpA.); idudilast eye drops, tipelukast, ibudilast (Kyorin Pharmaceutical Co. Ltd.); theophylline (Elan Corp.); cilomilast (GlaxoSmithKline plc.); Atopik 10 (Barrier Therapeutics Inc.); tofimilast, CI-1044, PD-189659, CP-220629, PDE 4d inhibitor BHN (Pfizer Inc.); arofylline, LAS-37779 (Almirall Prodesfarma SA.); roflumilast, hydroxypumafentrine (Altana AG), tetomilast (Otska Pharmaceutical Co. Ltd.); tipelukast, ibudilast (Kyorin Pharmaceutical), CC-10004 (Celgene Corp.); HT-0712, IPL-4088 (Inflazyme Pharmaceuticals Ltd.); MEM-1414, MEM-1917 (Memory Pharmaceuticals Corp.); oglemilast, GRC-4039 15 (Glenmark Pharmaceuticals Ltd.); AWD-12-281, ELB-353, ELB-526 (Elbion AG); EHT-0202 (ExonHit Therapeutics SA.); ND-1251 (Neuro3d SA.); 4AZA-PDE4 (4 AZA Bioscience NV.); AVE-8112 (Sanofi-Aventis); CR-3465 (Rottapharm SpA.); GP-0203, NCS-613 (Centre National de la Recherche Scientifique); KF-19514 (Kyowa Hakko Kogyo Co. Ltd.); ONO-6126 (Ono Pharmaceutical Co. Ltd.); OS-0217 (Dainippon Pharmaceutical Co. Ltd.); IBFB-130011, IBFB- 20 150007, IBFB-130020, IBFB-140301 (IBFB Pharma GmbH); IC-485 (ICOS Corp.); RBx-14016 and RBx-11082 (Ranbaxy Laboratories Ltd.). A preferred PDE-4-inhibitor is Rolipram.

MAO inhibitors and compositions containing such inhibitors are described, e.g. in WO2006/091988, WO2005/007614, WO2004/089351, WO01/26656, WO01/12176, 25 WO99/57120, WO99/57119, WO99/13878, WO98/40102, WO98/01157, WO96/20946, WO94/07890 and WO92/21333.

Suitable MAO-inhibitors for the purpose of the present invention are for example Linezolid (Pharmacia Corp.); RWJ-416457 (RW Johnson Pharmaceutical Research Institute); bupropine 30 (Altana AG); GPX-325 (BioResearch Ireland); isocarboxazid; phenelzine; tranylcypromine; indantadol (Chiesi Farmaceutici SpA.); moclobemide (Roche Holding AG); SL-25.1131 (Sanofi-Synthelabo); CX-1370 (Burroughs Wellcome Co.); CX-157 (Krenitsky Pharmaceuticals Inc.); desoxypheganine (HF Arzneimittelforschung GmbH & Co. KG); bifemelane (Mitsubishi-Tokyo Pharmaceuticals Inc.); RS-1636 (Sankyo Co. Ltd.); esuprone (BASF AG); rasagiline (Teva Pharmaceutical Industries Ltd.); ladostigil (Hebrew University of Jerusalem); safinamide (Pfizer) 35 and NW-1048 (Newron Pharmaceuticals SpA.).

Suitable histamine H3 antagonists for the purpose of the present invention are, e.g. ABT-239, ABT-834 (Abbott Laboratories); 3874-H1 (Aventis Pharma); UCL-2173 (Berlin Free University), UCL-1470 (BioProjet, Societe Civile de Recherche); DWP-302 (Daewoong Pharmaceutical Co Ltd); GSK-189254A, GSK-207040A (GlaxoSmithKline Inc.); cipralisant, GT-2203 (Gliatech Inc.);

5 Ciproxifan (INSERM), 1S,2S-2-(2-Aminoethyl)-1-(1H-imidazol-4-yl)cyclopropane (Hokkaido University); JNJ-17216498, JNJ-5207852 (Johnson & Johnson); NNC-0038-0000-1049 (Novo Nordisk A/S); and Sch-79687 (Schering-Plough).

PEP inhibitors and compositions containing such inhibitors are described, e.g. in JP 01042465,

10 JP 03031298, JP 04208299, WO 00/71144, US 5,847,155; JP 09040693, JP 10077300, JP 05331072, JP 05015314, WO 95/15310, WO 93/00361, EP 0556482, JP 06234693, JP 01068396, EP 0709373, US 5,965,556, US 5,756,763, US 6,121,311, JP 63264454, JP 64000069, JP 63162672, EP 0268190, EP 0277588, EP 0275482, US 4,977,180, US 5,091,406, US 4,983,624, US 5,112,847, US 5,100,904, US 5,254,550, US 5,262,431, US 15 5,340,832, US 4,956,380, EP 0303434, JP 03056486, JP 01143897, JP 1226880, EP 0280956, US 4,857,537, EP 0461677, EP 0345428, JP 02275858, US 5,506,256, JP 06192298, EP 0618193, JP 03255080, EP 0468469, US 5,118,811, JP 05025125, WO 9313065, JP 05201970, WO 9412474, EP 0670309, EP 0451547, JP 06339390, US 5,073,549, US 4,999,349, EP 0268281, US 4,743,616, EP 0232849, EP 0224272, JP 62114978, JP 62114957, 20 US 4,757,083, US 4,810,721, US 5,198,458, US 4,826,870, EP 0201742, EP 0201741, US 4,873,342, EP 0172458, JP 61037764, EP 0201743, US 4,772,587, EP 0372484, US 5,028,604, WO 91/18877, JP 04009367, JP 04235162, US 5,407,950, WO 95/01352, JP 01250370, JP 02207070, US 5,221,752, EP 0468339, JP 04211648, WO 99/46272, WO 2006/058720 and PCT/EP2006/061428.

25 Suitable prolyl endopeptidase inhibitors for the purpose of the present invention are, e.g. Fmoc-Ala-Pyrr-CN, Z-Phe-Pro-Benzothiazole (Probiodrug), Z-321 (Zeria Pharmaceutical Co Ltd.); ONO-1603 (Ono Pharmaceutical Co Ltd); JTP-4819 (Japan Tobacco Inc.) and S-17092 (Servier).

30 Other suitable compounds that can be used according to the present invention in combination with QC-inhibitors are NPY, an NPY mimetic or an NPY agonist or antagonist or a ligand of the NPY receptors.

35 Preferred according to the present invention are antagonists of the NPY receptors.

Suitable ligands or antagonists of the NPY receptors are 3a, 4,5,9b-tetrahydro-1h-benz[e]indol-2-yl amine-derived compounds as disclosed in WO 00/68197.

NPY receptor antagonists which may be mentioned include those disclosed in European patent applications EP 0 614 911, EP 0 747 357, EP 0 747 356 and EP 0 747 378; international patent applications WO 94/17035, WO 97/19911, WO 97/19913, WO 96/12489, WO 97/19914, WO 96/22305, WO 96/40660, WO 96/12490, WO 97/09308, WO 97/20820, WO 97/20821, WO 97/20822, WO 97/20823, WO 97/19682, WO 97/25041, WO 97/34843, WO 97/46250, WO 98/03492, WO 98/03493, WO 98/03494 and WO 98/07420; WO 00/30674, US patents Nos. 5,552,411, 5,663,192 and 5,567,714; 6,114,336, Japanese patent application JP 09157253; international patent applications WO 94/00486, WO 93/12139, WO 95/00161 and WO 99/15498; US Patent No. 5,328,899; German patent application DE 393 97 97; European patent applications EP 355 794 and EP 355 793; and Japanese patent applications JP 06116284 and JP 07267988. Preferred NPY antagonists include those compounds that are specifically disclosed in these patent documents. More preferred compounds include amino acid and non-peptide-based NPY antagonists. Amino acid and non-peptide-based NPY antagonists which may be mentioned include those disclosed in European patent applications EP 0 614 911, EP 0 747 357, EP 0 747 356 and EP 0 747 378; international patent applications WO 94/17035, WO 97/19911, WO 97/19913, WO 96/12489, WO 97/19914, WO 96/22305, WO 96/40660, WO 96/12490, WO 97/09308, WO 97/20820, WO 97/20821, WO 97/20822, WO 97/20823, WO 97/19682, WO 97/25041, WO 97/34843, WO 97/46250, WO 98/03492, WO 98/03493, WO 98/03494, WO 98/07420 and WO 99/15498 ; US patents Nos. 5,552,411, 5,663,192 and 5,567,714; and Japanese patent application JP 09157253. Preferred amino acid and non-peptide-based NPY antagonists include those compounds that are specifically disclosed in these patent documents.

Particularly preferred compounds include amino acid-based NPY antagonists. Amino acid-based compounds, which may be mentioned include those disclosed in international patent applications WO 94/17035, WO 97/19911, WO 97/19913, WO 97/19914 or, preferably, WO 99/15498. Preferred amino acid-based NPY antagonists include those that are specifically disclosed in these patent documents, for example BIBP3226 and, especially, (R)-N2-(diphenylacetyl)-(R)-N-[1-(4-hydroxy- phenyl) ethyl] arginine amide (Example 4 of international patent application WO 99/15498).

35 M1 receptor agonists and compositions containing such inhibitors are described, e.g. in WO2004/087158, WO91/10664.

Suitable M1 receptor antagonists for the purpose of the present invention are for example CDD-0102 (Cognitive Pharmaceuticals); Cevimeline (Evoxac) (Snow Brand Milk Products Co. Ltd.); NGX-267 (TorreyPines Therapeutics); sabcomeline (GlaxoSmithKline); alvameline (H Lundbeck A/S); LY-593093 (Eli Lilly & Co.); VRTX-3 (Vertex Pharmaceuticals Inc.); WAY-132983 (Wyeth) 5 and CI-101 7/ (PD-151832) (Pfizer Inc.).

Acetylcholinesterase inhibitors and compositions containing such inhibitors are described, e.g. in WO2006/071274, WO2006/070394, WO2006/040688, WO2005/092009, WO2005/079789, WO2005/039580, WO2005/027975, WO2004/084884, WO2004/037234, WO2004/032929,

10 WO03/101458, WO03/091220, WO03/082820, WO03/020289, WO02/32412, WO01/85145, WO01/78728, WO01/66096, WO00/02549, WO01/00215, WO00/15205, WO00/23057, WO00/33840, WO00/30446, WO00/23057, WO00/15205, WO00/09483, WO00/07600, WO00/02549, WO99/47131, WO99/07359, WO98/30243, WO97/38993, WO97/13754, WO94/29255, WO94/20476, WO94/19356, WO93/03034 and WO92/19238.

15 Suitable acetylcholinesterase inhibitors for the purpose of the present invention are for example Donepezil (Eisai Co. Ltd.); rivastigmine (Novartis AG); (-)-phenserine (TorreyPines Therapeutics); Iadostigil (Hebrew University of Jerusalem); huperzine A (Mayo Foundation); galantamine (Johnson & Johnson); Memoquin (Universita di Bologna); SP-004 (Samaritan 20 Pharmaceuticals Inc.); BGC-20-1259 (Sankyo Co. Ltd.); physostigmine (Forest Laboratories Inc.); NP-0361 (Neuropharma SA); ZT-1 (Debiopharm); tacrine (Warner-Lambert Co.); metrifonate (Bayer Corp.) and INM-176 (WhanInn).

NMDA receptor antagonists and compositions containing such inhibitors are described, e.g. in

25 WO2006/094674, WO2006/058236, WO2006/058059, WO2006/010965, WO2005/000216, WO2005/102390, WO2005/079779, WO2005/079756, WO2005/072705, WO2005/070429, WO2005/055996, WO2005/035522, WO2005/009421, WO2005/000216, WO2004/092189, WO2004/039371, WO2004/028522, WO2004/009062, WO03/010159, WO02/072542, WO02/34718, WO01/98262, WO01/94321, WO01/92204, WO01/81295, WO01/32640, 30 WO01/10833, WO01/10831, WO00/56711, WO00/29023, WO00/00197, WO99/53922, WO99/48891, WO99/45963, WO99/01416, WO99/07413, WO99/01416, WO98/50075, WO98/50044, WO98/10757, WO98/05337, WO97/32873, WO97/23216, WO97/23215, WO97/23214, WO96/14318, WO96/08485, WO95/31986, WO95/26352, WO95/26350, WO95/26349, WO95/26342, WO95/12594, WO95/02602, WO95/02601, WO94/20109, 35 WO94/13641, WO94/09016 and WO93/25534.

Suitable NMDA receptor antagonists for the purpose of the present invention are for example Memantine (Merz & Co. GmbH); topiramate (Johnson & Johnson); AVP-923 (Neurodex) (Center for Neurologic Study); EN-3231 (Endo Pharmaceuticals Holdings Inc.); neramexane (MRZ-2/579) (Merz and Forest); CNS-5161 (CeNeS Pharmaceuticals Inc.); dexanabinol (HU-211;

5 Sinnabidol; PA-50211) (Pharmos); EpiCept NP-1 (Dalhousie University); indantadol (V-3381; CNP-3381) (Vernalis); perzinfotel (EAA-090, WAY-126090, EAA-129) (Wyeth); RGH-896 (Gedeon Richter Ltd.); traxoprodil (CP-101606), besonprodil (PD-196860, CI-1041) (Pfizer Inc.); CGX-1007 (Cognetix Inc.); delucemine (NPS-1506) (NPS Pharmaceuticals Inc.); EVT-101 (Roche Holding AG); acamprosate (Synchroneuron LLC.); CR-3991, CR-2249, CR-3394 (Rottapharm SpA.); AV-101 (4-Cl-kynurenine (4-Cl-KYN)), 7-chloro-kynurenic acid (7-Cl-KYNA) (VistaGen); NPS-1407 (NPS Pharmaceuticals Inc.); YT-1006 (Yaupon Therapeutics Inc.); ED-1812 (Sosei R&D Ltd.); himantane (hydrochloride N-2-(adamantly)-hexamethylen-imine) (RAMS); Lancicemine (AR-R-15896) (AstraZeneca); EVT-102, Ro-25-6981 and Ro-63-1908 (Hoffmann-La Roche AG / Evotec).

10 Furthermore, the present invention relates to combination therapies useful for the treatment of atherosclerosis, restenosis or arthritis, administering a QC inhibitor in combination with another therapeutic agent selected from the group consisting of inhibitors of the angiotensin converting enzyme (ACE); angiotensin II receptor blockers; diuretics; calcium channel blockers (CCB);

15 beta-blockers; platelet aggregation inhibitors; cholesterol absorption modulators; HMG-Co-A reductase inhibitors; high density lipoprotein (HDL) increasing compounds; renin inhibitors; IL-6 inhibitors; antiinflammatory corticosteroids; antiproliferative agents; nitric oxide donors; inhibitors of extracellular matrix synthesis; growth factor or cytokine signal transduction inhibitors; MCP-1 antagonists and tyrosine kinase inhibitors providing beneficial or synergistic therapeutic effects


20 over each monotherapy component alone.

25 Angiotensin II receptor blockers are understood to be those active agents that bind to the AT1 - receptor subtype of angiotensin II receptor but do not result in activation of the receptor. As a consequence of the blockade of the AT1 receptor, these antagonists can, e.g. be employed as


30 antihypertensive agents.

35 Suitable angiotensin II receptor blockers which may be employed in the combination of the present invention include AT₁ receptor antagonists having differing structural features, preferred are those with non-peptidic structures. For example, mention may be made of the compounds that are selected from the group consisting of valsartan (EP 443983), losartan (EP 253310), candesartan (EP 459136), eprosartan (EP 403159), irbesartan (EP 454511), olmesartan (EP


503785), tasosartan (EP 539086), telmisartan (EP 522314), the compound with the designation E-41 77 of the formula



5 the compound with the designation SC-52458 of the following formula



and the compound with the designation the compound ZD-8731 of the formula



10

or, in each case, a pharmaceutically acceptable salt thereof.

Preferred AT1-receptor antagonists are those agents that have been approved and reached the market, most preferred is valsartan, or a pharmaceutically acceptable salt thereof.

15

The interruption of the enzymatic degradation of angiotensin to angiotensin II with ACE inhibitors is a successful variant for the regulation of blood pressure and thus also makes available a therapeutic method for the treatment of hypertension.

20 A suitable ACE inhibitor to be employed in the combination of the present invention is, e.g. a compound selected from the group consisting alacepril, benazepril, benazeprilat; captopril,

ceronapril, cilazapril, delapril, enalapril, enaprilat, fosinopril, imidapril, lisinopril, moveltopril, perindopril, quinapril, ramipril, spirapril, temocapril and trandolapril, or in each case, a pharmaceutically acceptable salt thereof.

5 Preferred ACE inhibitors are those agents that have been marketed, most preferred are benazepril and enalapril.

A diuretic is, for example, a thiazide derivative selected from the group consisting of chlorothiazide, hydrochlorothiazide, methylclothiazide, and chlorothalidon. The most preferred

10 diuretic is hydrochlorothiazide. A diuretic furthermore comprises a potassium sparing diuretic such as amiloride or triameterine, or a pharmaceutically acceptable salt thereof.

The class of CCBs essentially comprises dihydropyridines (DHPs) and non-DHPs, such as diltiazem-type and verapamil-type CCBs.

15 A CCB useful in said combination is preferably a DHP representative selected from the group consisting of amlodipine, felodipine, ryosidine, isradipine, lacidipine, nicardipine, nifedipine, niguldipine, niludipine, nimodipine, nisoldipine, nitrendipine and nivaldipine, and is preferably a non-DHP representative selected from the group consisting of flunarizine, prenylamine,

20 diltiazem, fendiline, gallopamil, mibefradil, anipamil, tiapamil and verapamil, and in each case, a pharmaceutically acceptable salt thereof. All these CCBs are therapeutically used, e.g. as anti-hypertensive, anti-angina pectoris or anti-arrhythmic drugs.

Preferred CCBs comprise amlodipine, diltiazem, isradipine, nicardipine, nifedipine, nimodipine, 25 nisoldipine, nitrendipine and verapamil or, e.g. dependent on the specific CCB, a pharmaceutically acceptable salt thereof. Especially preferred as DHP is amlodipine or a pharmaceutically acceptable salt thereof, especially the besylate. An especially preferred representative of non-DHPs is verapamil or a pharmaceutically acceptable salt, especially the hydrochloride, thereof.

30 Beta-blockers suitable for use in the present invention include beta-adrenergic blocking agents (beta-blockers), which compete with epinephrine for beta-adrenergic receptors and interfere with the action of epinephrine. Preferably, the beta-blockers are selective for the beta-adrenergic receptor as compared to the alpha-adrenergic receptors, and so do not have a 35 significant alpha-blocking effect. Suitable beta-blockers include compounds selected from acebutolol, atenolol, betaxolol, bisoprolol, carteolol, carvedilol, esmolol, labetalol, metoprolol, nadolol, oxprenolol, penbutolol, pindolol, propranolol, sotalol and timolol. Where the beta-

blocker is an acid or base or otherwise capable of forming pharmaceutically acceptable salts or prodrugs, these forms are considered to be encompassed herein, and it is understood that the compounds may be administered in free form or in the form of a pharmaceutically acceptable salt or a prodrug, such as a physiologically hydrolyzable and acceptable ester. For example,

5 metoprolol is suitably administered as its tartrate salt, propranolol is suitably administered as the hydrochloride salt, and so forth.

Platelet aggregation inhibitors include PLAVIX® (clopidogrel bisulfate), PLETAL® (cilostazol) and aspirin.

10 Cholesterol absorption modulators include ZETIA® (ezetimibe) and KT6-971 (Kotobuki Pharmaceutical Co. Japan).

15 HMG-Co-A reductase inhibitors (also called beta-hydroxy-beta-methylglutaryl-co-enzyme-A reductase inhibitors or statins) are understood to be those active agents which may be used to lower lipid levels including cholesterol in blood.

20 The class of HMG-Co-A reductase inhibitors comprises compounds having differing structural features. For example, mention may be made of the compounds, which are selected from the group consisting of atorvastatin, cerivastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin and simvastatin, or in each case, a pharmaceutically acceptable salt thereof.

25 Preferred HMG-Co-A reductase inhibitors are those agents, which have been marketed, most preferred is atorvastatin, pitavastatin or simvastatin, or a pharmaceutically acceptable salt thereof.

HDL-increasing compounds include, but are not limited to, cholesterol ester transfer protein (CETP) inhibitors. Examples of CETP inhibitors include JTT705 disclosed in Example 26 of U.S. Patent No. 6,426,365 issued July 30, 2002, and pharmaceutically acceptable salts thereof.

30 Inhibition of interleukin 6 mediated inflammation may be achieved indirectly through regulation of endogenous cholesterol synthesis and isoprenoid depletion or by direct inhibition of the signal transduction pathway utilizing interleukin-6 inhibitor/antibody, interleukin-6 receptor inhibitor/antibody, interleukin-6 antisense oligonucleotide (ASON), gp130 protein inhibitor/antibody, tyrosine kinase inhibitors/antibodies, serine/threonine kinase inhibitors/antibodies, mitogen-activated protein (MAP) kinase inhibitors/antibodies, phosphatidylinositol 3-kinase (PI3K) inhibitors/antibodies, Nuclear factor kappaB (NF- κ B)

inhibitors/antibodies, I_KB kinase (IKK) inhibitors/antibodies, activator protein-1 (AP-1) inhibitors/antibodies, STAT transcription factors inhibitors/antibodies, altered IL-6, partial peptides of IL-6 or IL-6 receptor, or SOCS (suppressors of cytokine signaling) protein, PPAR gamma and/or PPAR beta/delta activators/ligands or a functional fragment thereof.

5

A suitable antiinflammatory corticosteroid is dexamethasone.

Suitable antiproliferative agents are cladribine, rapamycin, vincristine and taxol.

10 A suitable inhibitor of extracellular matrix synthesis is halofuginone.

A suitable growth factor or cytokine signal transduction inhibitor is, e.g. the ras inhibitor R115777.

15 A suitable tyrosine kinase inhibitor is tyrphostin.

Suitable renin inhibitors are described, e.g. in WO 2006/116435. A preferred renin inhibitor is aliskiren, preferably in the form of the hemi-fumarate salt thereof.

20 MCP-1 antagonists may, e.g. be selected from anti-MCP-1 antibodies, preferably monoclonal or humanized monoclonal antibodies, MCP-1 expression inhibitors, CCR2-antagonists, TNF-alpha inhibitors, VCAM-1 gene expression inhibitors and anti-C5a monoclonal antibodies.

MCP-1 antagonists and compositions containing such inhibitors are described, e.g. in
25 WO02/070509, WO02/081463, WO02/060900, US2006/670364, US2006/677365, WO2006/097624, US2006/316449, WO2004/056727, WO03/053368, WO00/198289, WO00/157226, WO00/046195, WO00/046196, WO00/046199, WO00/046198, WO00/046197, WO99/046991, WO99/007351, WO98/006703, WO97/012615, WO2005/105133, WO03/037376, WO2006/125202, WO2006/085961, WO2004/024921, WO2006/074265.

30

Suitable MCP-1 antagonists are, for instance, C-243 (Telik Inc.); NOX-E36 (Noxxon Pharma AG); AP-761 (Actimis Pharmaceuticals Inc.); ABN-912, NIBR-177 (Novartis AG); CC-11006 (Celgene Corp.); SSR-150106 (Sanofi-Aventis); MLN-1202 (Millenium Pharmaceuticals Inc.); AGI-1067, AGIX-4207, AGI-1096 (AtherioGenics Inc.); PRS-211095, PRS-211092 (Pharmos

35 Corp.); anti-C5a monoclonal antibodies, e.g. neutrazumab (G2 Therapies Ltd.); AZD-6942 (AstraZeneca plc.); 2-mercaptoimidazoles (Johnson & Johnson); TEI-E00526, TEI-6122

(Deltagen); RS-504393 (Roche Holding AG); SB-282241, SB-380732, ADR-7 (GlaxoSmithKline); anti-MCP-1 monoclonal antibodies(Johnson & Johnson).

5 Combinations of QC-inhibitors with MCP-1 antagonists may be useful for the treatment of inflammatory diseases in general, including neurodegenerative diseases.

Combinations of QC-inhibitors with MCP-1 antagonists are preferred for the treatment of Alzheimer's disease.

10 Most preferably the QC inhibitor is combined with one or more compounds selected from the following group:

PF-4360365, m266, bapineuzumab, R-1450, Posiphen, (+)-phenserine, MK-0752, LY-450139, E-2012, (R)-flurbiprofen, AZD-103, AAB-001 (Bapineuzumab), Tramiprosate, EGb-761, TAK-070, Doxofylline, theophylline, cilomilast, tofimilast, roflumilast, tetomilast, tipelukast, ibudilast,

15 HT-0712, MEM-1414, oglemilast, Linezolid, budipine, isocarboxazid, phenelzine, tranylcypromine, indantadol, moclobemide, rasagiline, ladostigil, safinamide, ABT-239, ABT-834, GSK-189254A, Ciproxifan, JNJ-17216498, Fmoc-Ala-Pyrr-CN, Z-Phe-Pro-Benzothiazole, Z-321, ONO-1603, JTP-4819, S-17092, BIBP3226; (R)-N2-(diphenylacetyl)-(R)-N-[1-(4-hydroxyphenyl) ethyl] arginine amide, Cevimeline, sabcomeline, (PD-151832), Donepezil,

20 rivastigmine, (-)-phenserine, ladostigil, galantamine, tacrine, metrifonate, Memantine, topiramate, AVP-923, EN-3231, neramexane, valsartan, benazepril, enalapril, hydrochlorothiazide, amlodipine, diltiazem, isradipine, nicardipine, nifedipine, nimodipine, nisoldipine, nitrendipine, verapamil, amlodipine, acebutolol, atenolol, betaxolol, bisoprolol, carteolol, carvedilol, esmolol, labetalol, metoprolol, nadolol, oxprenolol, penbutolol, pindolol,

25 propranolol, sotalol, timolol, PLAVIX® (clopidogrel bisulfate), PLETAL® (cilostazol), aspirin, ZETIA® (ezetimibe) and KT6-971, statins, atorvastatin, pitavastatin or simvastatin; dexamethasone, cladribine, rapamycin, vincristine, taxol, aliskiren, C-243, ABN-912, SSR-150106, MLN-1202 and betaferon.

30 In particular, the following combinations are considered:

35 - a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with Atorvastatin for the treatment and/or prevention of atherosclerosis,

- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with immunosuppressive agents, preferably rapamycin for the prevention and/or treatment of restenosis,

- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with immunosuppressive agents, preferably paclitaxel for the prevention and/or treatment of restenosis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with AChE inhibitors, preferably Donepezil, for the prevention and/or treatment of Alzheimer's disease,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with interferones, preferably Aronex, for the prevention and/or treatment of multiple sclerosis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with interferones, preferably betaferon, for the prevention and/or treatment of multiple sclerosis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with interferones, preferably Rebif, for the prevention and/or treatment of multiple sclerosis
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with Copaxone, for the prevention and/or treatment of multiple sclerosis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with dexamethasone, for the prevention and/or treatment of restenosis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with dexamethasone, for the prevention and/or treatment of atherosclerosis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with dexamethasone, for the prevention and/or treatment of rheumatid arthritis,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with HMG-Co-A-reductase inhibitors, for the prevention and/or treatment of restenosis, wherein the HMG-Co-A-reductase inhibitor is selected from atorvastatin, cerivastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin and simvastatin,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with HMG-Co-A reductase inhibitors, for the prevention and/or treatment of atherosclerosis wherein the HMG-Co-A-reductase inhibitor is selected from atorvastatin, cerivastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin and simvastatin,

- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with HMG-Co-A reductase inhibitors, for the prevention and/or treatment of rheumatoid arthritis wherein the HMG-Co-A-reductase inhibitor is selected from atorvastatin, cerivastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin and simvastatin,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with amyloid-beta antibodies for the prevention and/or treatment of mild cognitive impairment, wherein the amyloid-beta antibody is Acl-24,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with amyloid-beta antibodies for the prevention and/or treatment of Alzheimer's disease, wherein the amyloid-beta antibody is Acl-24,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with amyloid-beta antibodies for the prevention and/or treatment of neurodegeneration in Down Syndrome, wherein the amyloid-beta antibody is Acl-24,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with beta-secretase inhibitors for the prevention and/or treatment of mild cognitive impairment, wherein the beta-secretase inhibitor is selected from WY-25105, GW-840736X and CTS-21166,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with beta-secretase inhibitors for the prevention and/or treatment of Alzheimer's disease, wherein the beta-secretase inhibitor is selected from WY-25105, GW-840736X and CTS-21166,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with beta-secretase inhibitors for the prevention and/or treatment of neurodegeneration in Down Syndrome, wherein the beta-secretase inhibitor is selected from WY-25105, GW-840736X and CTS-21166,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with gamma-secretase inhibitors for the prevention and/or treatment of mild cognitive impairment, wherein the gamma-secretase inhibitor is selected from LY-450139, LY-411575 and AN-37124,
- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with gamma-secretase

inhibitors for the prevention and/or treatment of Alzheimer's disease, wherein the gamma-secretase inhibitor is selected from LY-450139, LY-411575 and AN-37124,

- a QC inhibitor, preferably a QC inhibitor of formula (I), more preferably a QC inhibitor selected from any one of examples 1-36, in combination with gamma-secretase inhibitors for the prevention and/or treatment of neurodegeneration in Down Syndrome, wherein the gamma-secretase inhibitor is selected from LY-450139, LY-411575 and AN-37124.

Such a combination therapy is in particular useful for AD, FAD, FDD and neurodegeneration in

10 Down syndrome as well as atherosclerosis, rheumatoid arthritis, restenosis and pancreatitis.

Such combination therapies might result in a better therapeutic effect (less proliferation as well as less inflammation, a stimulus for proliferation) than would occur with either agent alone.

15 With regard to the specific combination of inhibitors of QC and further compounds it is referred in particular to WO 2004/098625 in this regard, which is incorporated herein by reference.

Pharmaceutical compositions

To prepare the pharmaceutical compositions of this invention, at least one compound of formula

20 (I) optionally in combination with at least one of the other aforementioned agents can be used as the active ingredient(s). The active ingredient(s) is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage

25 form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents,

30 lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as aiding solubility or for

35 preservation, may be included.

Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient(s) necessary to deliver an effective dose as described above. The 5 pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, from about 0.03 mg to 100 mg/kg (preferred 0.1 – 30 mg/kg) and may be given at a dosage of from about 0.1 – 300 mg/kg per day (preferred 1 – 50 mg/kg per day) of each active ingredient or combination thereof. The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition 10 being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, 15 drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing 20 solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. 25 When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of each active ingredient or combinations 30 thereof of the present invention.

The tablets or pills of the compositions of the present invention can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the 35 latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material

can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

This liquid forms in which the compositions of the present invention may be incorporated for 5 administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, 10 methylcellulose, polyvinylpyrrolidone or gelatin.

The pharmaceutical composition may contain between about 0.01 mg and 100 mg, preferably about 5 to 50 mg, of each compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical 15 excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixirs, emulsions, and suspensions. Forms useful for parenteral 20 administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form 25 via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

30 For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as 35 glucose or betalactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium

acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitable flavored suspending or dispersing agents such as the synthetic and

5 natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

10 The compounds or combinations of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

15 Compounds or combinations of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxyethylaspartamid-ephenol, or polyethyl
20 eneoxidepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polyactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

25 Compounds or combinations of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of the addressed disorders is required.

30 The daily dosage of the products may be varied over a wide range from 0.01 to 1.000 mg per mammal per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of each active ingredient or combinations thereof for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is
35 ordinarily supplied at a dosage level of from about 0.1 mg/kg to about 300 mg/kg of body weight per day. Preferably, the range is from about 1 to about 50 mg/kg of body weight per day. The compounds or combinations may be administered on a regimen of 1 to 4 times per day.

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of disease condition. In addition,

5 factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

In a further aspect, the invention also provides a process for preparing a pharmaceutical composition comprising at least one compound of formula (I), optionally in combination with at

10 least one of the other aforementioned agents and a pharmaceutically acceptable carrier.

The compositions are preferably in a unit dosage form in an amount appropriate for the relevant daily dosage.

15 Suitable dosages, including especially unit dosages, of the the compounds of the present invention include the known dosages including unit doses for these compounds as described or referred to in reference text such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition page 341 and pages 20 cited therein) or the above mentioned publications.

Examples

Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
1	N-(3-(1H-imidazol-1-yl)propyl)-6-methyl-2-phenylpyrimidin-4-amine		293.37	294.5	18	47.8
2	N-(3-(1H-imidazol-1-yl)propyl)-2,6-diphenylpyrimidin-4-amine		355.43	356.2	3.9	49.8

Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
3	N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-phenylpyrimidin-4-amine		293.36	294.3	4.6	28.06
4	N-(3-(1H-imidazol-1-yl)propyl)-2,6-dimethylpyrimidin-4-amine		231.29	232.3	9.31	30.3
5	N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-p-tolylpyrimidin-4-amine		307.39	308.2	3.52	13.3
6	N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-methylpyrimidin-4-amine		429.36	430.3	4.69	14.7
7	N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-phenylpyrimidin-4-amine		491.43	492.5		91.1
8	N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxyphenyl)-2-phenylpyrimidin-4-amine		445.51	446.4	1.16	7.08
9	N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-6-p-tolylpyrimidin-4-amine		369.46	370.4		14.4

Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
10	N-(3-(1H-imidazol-1-yl)propyl)-6-(4-nitrophenyl)-2-phenylpyrimidin-4-amine		400.43	401.4		12.4
11	N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-phenylpyrimidin-4-amine		385.46	386.2		22.6
12	N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-phenylpyrimidin-4-amine		385.46	386.2	3.75	9.37
13	N-(3-(1H-imidazol-1-yl)propyl)-6-(furan-3-yl)-2-phenylpyrimidin-4-amine		345.39	346.1	14.92	40.34
14	N-(3-(1H-imidazol-1-yl)propyl)-6-(adamant-1-yl)-2-phenylpyrimidin-4-amine		413.55	414.5		115.7
15	N-(3-(1H-imidazol-1-yl)propyl)-6-(trifluoromethyl)-2-phenylpyrimidin-4-amine		347.33	348.4	n.a.	n.a.
16	N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxyphenyl)-2-(pyridin-4-		446.50	447.4		


Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
	yl)pyrimidin-4-amine					
17	N-(3-(1H-imidazol-1-yl)propyl)-6-(furan-3-yl)-2-(pyridin-4-yl)pyrimidin-4-amine		346.38	347.2	3.70	27.29
18	N-(3-(1H-imidazol-1-yl)propyl)-6-(trifluoromethyl)-2-(pyridin-4-yl)pyrimidin-4-amine		348.32	349.2	4.19	36.75
19	N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-methylpyrimidin-4-amine		323.39	324.4	2.7	20.87
20	N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxy-phenyl)-2-methylpyrimidin-4-amine		383.44	384.2	0.72	6.43
21	N-(3-(1H-imidazol-1-yl)propyl)-6-(4-methoxyphenyl)-2-methylpyrimidin-4-amine		323.39	324.4	2.26	18.5
22	N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-methylpyrimidin-4-amine		323.39	324.4	2.16	19.3
23	N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-(pyridin-4-		492.41	493.3	n.a.	n.a.

Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
	yl)pyrimidin-4-amine					
24	N-(3-(1H-imidazol-1-yl)propyl)-6-(4-nitrophenyl)-2-(pyridin-4-yl)pyrimidin-4-amine		401.42	402.3		16.9
25	N-(3-(1H-imidazol-1-yl)propyl)-2-(pyridin-4-yl)-6-p-tolylpyrimidin-4-amine		370.45	371.2		30.6
26	N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine		386.44	387.2	0.83	8.18
27	N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine		386.44	387.3		19.0
28	6-(2-methoxyphenyl)-N-(3-(5-methyl-1H-imidazol-1-yl)propyl)-2-(pyridin-4-yl)pyrimidin-4-amine		400.47	401.2	0.281	1.09
29	6-(3,4,5-trimethoxyphenyl)-2-methyl-N-(3-(5-methyl-1H-imidazol-1-yl)propyl)pyrimidin-		397.47	398.5	0.116	0.886

Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
	4-amine					
30	N-(2-phenyl-6-(quinolin-6-yl)pyrimidin-4-yl)-3H-benzo[d]imidazol-5-amine		414.46	415.2		100
31	N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-(quinolin-6-yl)pyrimidin-4-amine		344.41	245.1	3.56	31.4
32	N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-6-(2-phenylcyclopropyl)pyrimidin-4-amine		395.49	396.3	n.a.	n.a.
33	N-(3-(1H-imidazol-1-yl)propyl)-6-(2-phenylcyclopropyl)-2-(pyridin-4-yl)pyrimidin-4-amine		396.48	397.3	3.06	26.5
34	N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-methylpyrimidin-4-amine		333.43	334.1	6.82	48.3
35	N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-phenylpyrimidin-4-amine		395.49	396.1	n.a.	n.a.

Ex.	Name	Structure	MW	Det [M+H] ⁺	K _i [μM]	IC ₅₀ [μM]
36	N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-(pyridin-4-yl)pyrimidin-4-amine		396.48	397.3	3.52	26.8

General synthesis description



5 Method A

The gamma-ketoester I (1.0 eq), the corresponding amidine hydrochloride II (1.3 eq) and potassium-*tert*-butylate (1.3 eq) were dissolved in 10 mL of dry ethanol. The mixture was stirred under reflux for 24 h. After that the mixture was filtered and the filter cake was washed four times by means of ethanol. The remaining solid 3*H*-pyridine-4-one (III) was dried and used without further purification.

Method B

The 3*H*-pyridin-4-one (III) (1eq) was dissolved in 10 mL of chloroform. After that tosyl chloride (1 eq) and N-methyl morpholine (1.4 eq) were added. The mixture was stirred for 30 min at 0 °C and 3 h at room temperature. The solvent was removed and the remaining solid, raw pyrimidine (IV) was dried and used without further purification.

Method C

Same procedure as describe for Method B, mesyl chloride (1eq) was used instead of tosyl chloride and triethylamine (1.4 eq) was used instead of N-methyl morpholine.

Method D

The 3*H*-pyridin-4-one (III) (1eq) was dissolved with POCl₃ (18 eq) and heated up to 90 °C for 4h. The excess of POCl₃ was removed by means of distillation under reduced pressure. The remaining solid was dissolved in 5 mL of water and the pH-value of the solution was adjusted to a neutral pH-value, by adding a saturated aqueous solution of NaHCO₃. After that the aqueous layer was extracted 4 times by means of 5 mL of CHCl₃. The combined organic layers was dried over Na₂SO₄, filtered and the solvent was removed. The remaining solid was dried and used without further purification.

Method E

The pyrimidines (IV, resulting from method B of C) (1eq) were dissolved in 5 mL of toluol. After that triethylamine (5 eq) and 3-(1H-imidazol-1-yl)propan-1-amine (1.3 eq) or 3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5) (1.3 eq) was added. The reaction mixture was stirred under

5 reflux for 24 h, then the solvent was removed. The remains was dissolved in 10 mL of chloroform and washed 3 times by means of 5mL of a saturated aqueous solution of NaHCO_3 . The organic layer was dried, filtered and the solvent was removed. The remaining oil was subjected to centrifugal force chromatography on a Chromatotron® Device (Harrison Research Ltd), utilizing silica gel plates of 2 mm layer thickness and a $\text{CHCl}_3/\text{MeOH}$ -gradient.

10 Method F

The pyrimidines (IV, resulting from method D) (1eq), dry CuI (0.05 eq), dry N,N-diethylsalicylamide (0.2 eq) and dry K_3PO_4 (2.0 eq) were dissolved in 10 mL of dry dimethyl formamide and kept under an argon atmosphere. 3-(1H-imidazol-1-yl)propan-1-amine (1.5 eq) or 3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5) (1.5 eq) were added dropwise. The mixture 15 was stirred for 20 h at 90 C. After that 20 mL of a saturated, aqueous solution of NaHCO_3 was added and the solution was extracted four times by means of 10 mL of ethyl-acetate. The organic layers were combined and the solvent was removed. The remaining oil was subjected to centrifugal force chromatography on a Chromatotron® Device (Harrison Research Ltd), utilizing silica gel plates of 2 mm layer thickness and a $\text{CHCl}_3/\text{MeOH}$ -gradient.

20 Method G

The pyrimidines (IV, resulting from method D) (1eq), dry CuI (0.05 eq), dry N,N-diethylsalicylamide (0.2 eq) and dry K_3PO_4 (2.0 eq) were dissolved in 10 mL of dry N-methyl-pyrrolidone and kept under an argon atmosphere. 3-(1H-imidazol-1-yl)propan-1-amine (1.5 eq) or 3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5) (1.5 eq) were added dropwise. The mixture 25 was stirred for 20 h at 90 C. After that 20 mL of a saturated, aqueous solution of NaHCO_3 was added and the solution was extracted four times by means of 10 mL of ethyl-acetate. The organic layers were combined and the solvent was removed. The remaining oil was subjected to centrifugal force chromatography on a Chromatotron® Device (Harrison Research Ltd), utilizing silica gel plates of 2 mm layer thickness and a $\text{CHCl}_3/\text{MeOH}$ -gradient.

30 Method H

5-Aminobenzimidazole (e.g. 1.0 mmol, 1 eq) and the corresponding pyrimidine (IV, resulting from method D) (1 eq) were dissolved in 10 mL of ethanol. Na_2CO_3 (2 eq) was added and the mixture was stirred for 48 h under reflux. The solvent was removed and the remains was redissolved in 10 mL of MeCN and 0.5 mL of triflouracetic acid and subjected to preparation by 35 means of semi-preparative HPLC.

Method I


The pyrimidines (IV, resulting from method D) (1eq), dry CuI (0.05 eq), dry N,N-diethylsalicylalamide (0.2 eq) and dry K₃PO₄ (2.0 eq) were dissolved in 10 mL of dry N-methylpyrrolidone and kept under an argon atmosphere. 3-(1H-imidazol-1-yl)propan-1-amine (1.5 eq)

5 or 3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5) (1.5 eq) were added dropwise. The mixture was stirred for 20 h at 90 C. After that 20 mL of a saturated, aqueous solution of NaHCO₃ was added and the solution was extracted four times by means of 10 mL of ethyl-acetate. The organic layers were combined and the solvent was removed. The remaining residue was purified by means of semi-preparative HPLC.

10 Semi-preparative HPLC-method

The system consisted of Alpha Crom Varian PrepStar device (model 218) equipped with a Luna® 10μ C18(2) 100A semi-preparative column (Phenomenex, length: 250 mm, diameter: 21,2 mm). The compounds were purified using a gradient at a flow rate of 21 mL/min; whereby eluent (A) was acetonitrile, eluent (B) was water, both containing 0.04% (v/v) trifluoro acetic acid

15 applying the following gradient: 0 min – 32 min. 20 -95% (A)

Detailed synthesis description**Synthesis of 3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5)**

20 **Figure 1:** Synthesis of 3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5)

3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (5)**4-methyl-1-trityl-1H-imidazole (2)**

1 equivalent (i.e. 36.53 mmol, 1eq.) of 4-methyl-1H-imidazole (1) was dissolved in 120 mL of
25 dimethylformamide, triethylamine (73.06 mmol, 2eq.) and chlorotriphenylmethane (40.1mmol,

1.1eq) where added. The mixture was stirred for 3.5 h. The precipitate filtered off and was washed by means of ice-cooled dimethylformamide (2x50 mL) and water (2x50 mL). After removal of the solvent the remaining product was dried over P_4O_{10} .

Yield: 10.65 g (98.2%). The product was used without further purification.

5 1-Trityl-3-[3-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)propyl]-1,4-dimethyl-1H-imidazol-3-ium bromide (**3**)
4-Methyl-1-trityl-1H-imidazole (i.e 32.85 mmol, 1eq.) was suspended in acetonitrile (10 mL) and 2-(3-bromopropyl)isoindoline-1,3-dione (32.85 mmol, 1eq.) was added. The mixture was kept under reflux over night. The organic sovent was removed by meaning of reduced pressure
10 resulting in 20.1 g of. Purification was done by flash-chromatography using silica gel and a $CHCl_3/MeOH$ -gradient.
Yield: 10.65g (63.44%).

*2-(3-(5-methyl-1H-imidazol-1-yl)propyl)isoindoline-1,3-dione (**4**)*

15 1-Trityl-3-[3-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)propyl]-1,4-dimethyl-1H-imidazol-3-ium bromide (i.e. 7.86 mmol) was dissolved in a stirred solution containing methanol (20 mL) and trifluoracetic acid (4 mL). The mixture was kept under reflux over night. After that the solvent was removed by means of reduced pressure and the remaining oil was purified by flash-chromatography using silica gel and a $CHCl_3/MeOH$ -gradient.
20 Yield: 2.05g (97.0%).

*3-(5-methyl-1H-imidazol-1-yl)propan-1-amine (**I**)*

2-(3-(5-methyl-1H-imidazol-1-yl)propyl)isoindoline-1,3-dione (i.e. 8.92 mmol, 1eq.) and hydazine monohydrate (17.84 mmol, 2eq.) were dissolved in dry $EtOH$ (50 mL). The mixture was kept under reflux over night, then mixture was concentrated down to a volume of 25 mL. After that hydrochloric acid (conc., 55 mL) was added and the mixture was heated up to 50 °C for and kept at this temperature for 30 min. The formed precipitate was filtered off. The filtrate was cooled down to 0 °C and solid NaOH was added until a final pH-value of 10-12 is reached. The aqueous solution was extracted by means of $CHCl_3$ (3x50 mL). The combined organic layers
25 were dried over Na_2SO_4 , filtered and the solvent was removed. The product was purified by meaning of flash-chromatography using silica gel and a $CHCl_3/MeOH$ -gradient.
30 Yield: 0.74 g (60%). viscous oil

Yield over all steps: 36.3 %

35 1H -NMR ($CDCl_3$, 499.78 MHz): δ 1.79-1.847 (m, 2H); 2.179 (s, 3H); 2.694-2.721 (m, 2H); 3.891-3.920 (m, 2H); 6.731 (s, H); 7.240 (s, solv.); 7.380 (s; H); ESI-MS m/z: 140.3 ($M+H$)⁺, 279.4 (2 $M+H$)⁺; HPLC (λ = 214 nm) rt: dead time (100%)

Semi-preparative HPLC-method

The system consisted of Merck-Hitachi device (model LaChrom) equipped with a SP250/21 Luna[®] 100-7 C18 semi-preparative column (Phenomenex, length: 250 mm, diameter: 21 mm).

The compounds were purified using a gradient at a flow rate of 6 mL/min; whereby eluent (A) was acetonitrile, eluent (B) was water, both containing 0.1 % (v/v) trifluoro acetic acid applying the following gradient: 0 min – 40 min. 40 -95 % (A)

Synthesis of examples**Example 1: N-(3-(1H-imidazol-1-yl)propyl)-6-methyl-2-phenylpyrimidin-4-amine**

10 Example 1 was synthesized according to the methods A, B and E.

Yield (over all steps): 0.03 g (0.02 %). ¹H NMR: (CDCl₃) δ 2.16-2.20 (m, 2H); 2.40 (s, 3H); 3.45-3.46 (d, 2H); 4.10-4.12 (t, 2H); 5.31 (br s, 1H); 6.12 (s, 1H); 6.95 (s, 1H); 7.09 (s, 1H); 7.41-7.45 (m, 3H); 7.79 (s, 1H); 8.31-8.35 (m, 2H). MS m/z 294.5(M+H)⁺; HPLC (λ = 214 nm, [D]) rt: 15.01 min (99.5%).

15

Example 2: N-(3-(1H-imidazol-1-yl)propyl)-2,6-diphenylpyrimidin-4-amine

The compound was synthesized according to the methods A, C and E.

Yield (over all steps): 0.02 g (0.56 %). ¹H NMR: (CDCl₃) δ 2.19-2.23 (m, 2H); 3.53-3.57 (m, 2H); 4.10-4.13 (t, 2H); 5.06 (br s, 1H); 6.61 (s, 1H); 6.95 (s, 1H); 7.09 (s, 1H); 7.43-7.51 (m, 6H); 20 7.61 (s, 1H); 8.09-8.11 (m, 2H); 8.48-8.50 (m, 2H). MS m/z 356.2 (M+H)⁺; HPLC (λ = 214 nm, [D]) rt: 22.29 min (90.2%).

Example 3: N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and E.

25 Yield (over all steps): 0.02 g (2.27 %). ¹H NMR: (CDCl₃) δ 2.11-2.15 (m, 2H); 2.57 (s, 3H); 3.41-3.47 (m, 2H); 4.06-4.09 (t, 2H); 4.94 (br s, 1H); 6.45 (s, 1H); 6.95 (s, 1H); 7.09 (s, 1H); 7.43-7.44 (m, 3H); 7.53 (s, 1H); 7.91-7.93 (m, 2H). MS m/z 294.3 (M+H)⁺; ESI-FTICR-MS: m/z 316.15311 ([M+Na]⁺, calc. for C₂₃H₂₄N₅Na⁺ 316,15326) HPLC (λ = 214 nm, [D]) rt: 6.99 min (95.9%).

30 **Example 4: N-(3-(1H-imidazol-1-yl)propyl)-2,6-dimethylpyrimidin-4-amine**

The compound was synthesized according to the methods A, D and E.

Yield (over all steps): 0.1 g (3.09 %). ¹H NMR: (CDCl₃) δ 2.05-2.10 (m, 2H); 2.29 (s, 3H); 2.46 (s, 3H); 3.28-3.32 (m, 2H); 4.02-4.05 (t, 2H); 4.83 (br s, 1H); 5.93 (s, 1H); 6.92 (s, 1H); 7.07 (s, 1H); 7.50 (s, 1H). MS m/z 232.3 (M+H)⁺; ESI-FTICR-MS: m/z 232.15559 ([M+H]⁺, calc. for 35 C₁₂H₁₇N₅⁺ 232.15567) HPLC (λ = 214 nm, [C]) rt: 2.53 min (95.6%).

Example 5: N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-p-tolylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and E

Yield (over all steps): 0.005 g (1.72 %). ^1H NMR: (CDCl_3) δ 2.10-2.13 (m, 2H); 2.38 (s, 3H); 2.55 (s, 3H); 3.37-3.41 (m, 2H); 4.05-4.08 (m, 2H); 4.86 (br s, 1H); 6.42 (s, 1H); 6.94 (s, 1H); 7.08 (s, 1H); 7.23 (s, 2H); 7.51 (s, 1H); 7.81-7.82 (m, 2H). MS m/z 308.2 ($\text{M}+\text{H}$) $^+$; HPLC (λ = 214 nm, [C]) rt: 8.08 min (91.1%).

Example 6: N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-methyl-pyrimidin-4-amine

The compound was synthesized according to the methods A, D and E.

Yield (over all steps): 0.015 g (0.2 %). ^1H NMR: (CDCl_3) δ 2.16 (br s, 2H); 2.58 (s, 3H); 3.47 (s, 2H); 4.09-4.10 (m, 2H); 5.12 (br s, 1H); 6.51 (s, 1H); 6.98 (br s, 1H); 7.11 (br s, 1H); 7.64 (br s, 1H); 7.91 (s, 1H); 8.39 (s, 2H). MS m/z 430.3 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 430.14645 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{19}\text{H}_{18}\text{F}_6\text{N}_5^+$ 430.14609) HPLC (λ = 214 nm, [C]) rt: 12.51 min (75.1 %).

Example 7: N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-phenyl-pyrimidin-4-amine

The compound was synthesized according to the methods A, D and E.

Yield (over all steps): 0.006 g (0.9 %). ^1H NMR: (CDCl_3) δ 2.23-2.28 (m, 2H); 3.06-3.11 (m, 2H); 4.18-4.21 (m, 2H); 5.84 (br s, 1H); 6.77 (s, 1H); 6.99 (s, 1H); 7.13 (s, 1H); 7.47-7.49 (m, 3H); 7.91 (s, 1H); 8.10 (s, 1H); 8.45-8.51 (m, 4H). MS m/z 492.5 ($\text{M}+\text{H}$) $^+$; HPLC (λ = 214 nm, [C]) rt: 21.60 min (82.1 %).

Example 8: N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxyphenyl)-2-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and F.

Yield (over all steps): 0.069 g (8.6 %). ^1H NMR: (CDCl_3) δ 2.19-2.23 (m, 2H); 3.53-3.59 (m, 2H); 3.90 (s, 3H); 3.97 (s, 6H); 4.09-4.13 (m, 2H); 4.93 (br s, 1H); 6.52 (s, 1H); 6.95 (s, 1H); 7.09 (s, 1H); 7.34 (s, 2H); 7.46-7.48 (m, 3H); 7.53 (s, 1H); 8.45-8.48 (m, 2H). MS m/z 446.4 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 468.20050 ($[\text{M}+\text{Na}]^+$, calc. for $\text{C}_{25}\text{H}_{27}\text{N}_5\text{Na}^+$ 468.20061) HPLC (λ = 214 nm, [C]) rt: 11.25 min (95.1 %).

Example 9: N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-6-p-tolylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and F.

Yield (over all steps): 0.06 g (6.8 %). ^1H NMR: (CDCl_3) δ 2.18-2.21 (m, 2H); 2.41 (s, 3H); 3.51-3.55 (m, 2H); 4.08-4.11 (m, 2H); 4.96 (br s, 1H); 6.56 (s, 1H); 6.66 (br s, 1H); 7.10 (br s, 1H); 7.24-7.28 (m, 2H); 7.45-7.49 (m, 3H); 7.55 (br s, 1H); 7.99-8.01 (m, 2H); 8.47-8.50 (m, 2H). MS m/z 370.4 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 370.20202 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{23}\text{H}_{24}\text{N}_5^+$ 370.20262)

5 HPLC ($\lambda = 214$ nm, [C]) rt: 11.49 min (98.1 %).

Example 10: N-(3-(1H-imidazol-1-yl)propyl)-6-(4-nitrophenyl)-2-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.23 g (24.0 %). ^1H NMR: (CDCl_3) δ 2.22-2.24 (m, 2H); 3.57-3.58 (d, 2H); 4.11-4.14 (m, 2H); 5.15 (br s, 1H); 6.63 (s, 1H); 7.24 (s, 1H); 7.47-7.49 (m, 3H); 8.24-8.33 (m, 4H); 8.46-8.48 (m, 2H). MS m/z 401.4 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 401.17124 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{22}\text{H}_{21}\text{N}_6\text{O}_2^+$ 401.17205) HPLC ($\lambda = 214$ nm, [C]) rt: 14.13 min (99.4 %).

Example 11: N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.021 g (2.4 %). ^1H NMR: (CDCl_3) δ 22.18-2.22 (m, 2H); 3.53-3.54 (d, 2H); 3.88 (s, 3H); 4.09-4.12 (m, 2H); 4.94 (br s, 1H); 6.94-7.00 (m, 3H); 7.08-7.12 (m, 2H); 7.37-7.45 (m, 4H); 8.16-8.18 (m, 1H); 8.44-8.46 (m, 2H). MS m/z 386.2 ($\text{M}+\text{H}$) $^+$; HPLC ($\lambda = 214$ nm, [C]) rt: 11.15 min (89.7 %).

Example 12: N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

25 Yield (over all steps): 0.089 g (10.3 %). ^1H NMR: (CDCl_3) δ 2.17-2.23 (m, 2H); 3.53-3.54 (d, 2H); 3.90 (s, 3H); 4.08-4.12 (m, 2H); 5.00 (br s, 1H); 6.57 (s, 1H); 6.99-7.01 (m, 1H); 7.36-7.40 (t, 1H); 7.45-7.49 (m, 3H); 7.61-7.63 (d, 2H); 7.71 (s, 1H); 8.47-8.50 (m, 2H). MS m/z 386.2 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 386.19681 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{23}\text{H}_{24}\text{N}_5\text{O}^+$ 386.19754) HPLC ($\lambda = 214$ nm, [C]) rt: 11.28 min (98.8 %).

30

Example 13: N-(3-(1H-imidazol-1-yl)propyl)-6-(furan-3-yl)-2-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.048 g (5.1 %). ^1H NMR: (CDCl_3) δ 2.17-2.20 (m, 2H); 2.49-2.51 (d, 2H); 4.08-4.10 (m, 2H); 5.04 (br s, 1H); 6.30 (s, 1H); 6.86 (s, 1H); 7.01 (br s, 1H); 7.13 (br s, 1H); 7.43-7.48 (m, 4H); 7.65 (br s, 1H); 8.17 (s, 1H); 8.41-8.43 (m, 2H). MS m/z 386.2 ($\text{M}+\text{H}$) $^+$; ESI-

35

FTICR-MS: m/z 346.16570 ([M+H]⁺, calc. for C₂₀H₂₀N₅O⁺ 346.16623) HPLC (λ = 214 nm, [B]) rt: 8.55 min (96.6 %).

Example 14: N-(3-(1H-imidazol-1-yl)propyl)-6-(adamant-1 yl)-2-phenylpyrimidin-4-amine

5 The compound was synthesized according to the methods A, D and G.
Yield (over all steps): 0.076 g (9.2 %). ¹H NMR: (CDCl₃) δ 1.74-1.81 (m, 6H); 1.96-2.04 (m, 6H); 2.14-2.18 (m, 2H); 2.82 (s, 3H); 3.45-3.50 (m, 2H); 4.06-4.09 (t, 2H); 4.76 (br s, 1H); 6.08 (s, 1H); 6.94 (br s, 1H); 7.08 (br s, 1H); 7.39-7.42 (m, 3H); 7.51 (br s, 1H); 8.40-8.42 (m, 2H). MS m/z 414.5 (M+H)⁺; ESI-FTICR-MS: m/z 414.26441 ([M+H]⁺, calc. for C₂₆H₃₂N₅⁺ 414.26522)

10 HPLC (λ = 214 nm, [B]) rt: 8.55 min (91.7 %).

Example 15: N-(3-(1H-imidazol-1-yl)propyl)-6-(trifluoromethyl)-2-phenylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.076 g (8.0 %). ¹H NMR: (CDCl₃) δ 2.17-2.23 (m, 2H); 3.47-3.62 (m, 2H); 4.09-4.12 (m, 2H); 5.67 (br s, 1H); 6.55 (s, 1H); 6.94 (s, 1H); 7.10 (s, 1H); 7.42-7.49 (m, 3H); 7.68 (br s, 1H); 8.36-8.39 (m, 2H). MS m/z 348.4 (M+H)⁺; ESI-FTICR-MS: m/z 348.14242 ([M+H]⁺, calc. for C₁₇H₁₇F₃N₅⁺ 348,14305) HPLC (λ = 214 nm, [B]) rt: 13.87 min (100 %).

Example 16: N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine

20 The compound was synthesized according to the methods A, D and G.
Yield (over all steps): 0.033 g (4.1 %). ¹H NMR: (DMSO-d₆) δ 2.02-2.08 (m, 2H); 3.44 (br s, 2H); 3.73 (s, 3H); 3.90 (s, 6H); 4.08-4.11 (t, 2H); 6.93-6.95 (m, 2H); 7.23 (s, 1H); 7.42 (s, 2H); 7.66-7.69 (m, 2H); 8.25-8.26 (m, 2H); 8.72-8.73 (m, 2H). MS m/z 474.4(M+H)⁺; ESI-FTICR-MS: m/z 447.21458 ([M+H]⁺, calc. for C₂₄H₂₇N₆O₃⁺ 447.21391) HPLC (λ = 214 nm, [D]) rt: 10.75 min (98.9 %).

Example 17: N-(3-(1H-imidazol-1-yl)propyl)-6-(furan-3-yl)-2-(pyridin-4-yl)pyrimidin-4-amine

30 The compound was synthesized according to the methods A, D and G.
Yield (over all steps): 0.021 g (3.4 %). ¹H NMR: (CDCl₃) δ 2.17-2.22 (m, 2H); 3.51-3.52 (d, 2H); 4.09-4.12 (t, 2H); 5.06 (br s, 1H); 6.35 (s, 1H); 6.84-6.85 (m, 1H); 6.95 (s, 1H); 7.11 (s, 1H); 7.49-7.51 (m, 1H); 7.54 (s, 1H); 8.17 (m, 1H); 8.23-8.24 (m, 2H); 8.72-8.73 (m, 2H). MS m/z 347.2 (M+H)⁺; ESI-FTICR-MS: m/z 347.16168 ([M+H]⁺, calc. for C₁₉H₁₉N₆O⁺ 347.16149) HPLC (λ = 214 nm, [B]) rt: 8.99 min (99.5 %).

Example 18: N-(3-(1H-imidazol-1-yl)propyl)-6-(trifluoromethyl)-2-(pyridin-4-yl)pyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.21 g (22.4 %). ^1H NMR: (DMSO-d₆) δ 2.02-2.07 (m, 2H); 3.45-3.49 (m, 2H); 4.06-4.09 (t, 2H); 6.88 (s, 1H); 6.92 (s, 1H); 7.21 (s, 1H); 7.65 (s, 1H); 8.10-8.11 (m, 2H); 8.27-8.29 (m, 1H); 8.73-8.74 (m, 2H). MS m/z 349.2 (M+H)⁺; ESI-FTICR-MS: m/z 349.13869 ([M+H]⁺, calc. for C₁₆H₁₆F₃N₆⁺ 349,13830) HPLC (λ = 214 nm, [D]) rt: 8.64 min (99.2 %).

Example 19: N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-methylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.07 g (9.3 %). ^1H NMR: (CDCl₃) δ 2.09-2.15 (m, 2H); 2.56 (s, 3H); 3.38-3.41 (m, 2H); 3.86 (s, 3H); 4.05-4.08 (t, 2H); 4.89 (br s, 1H); 6.43 (s, 1H); 6.94-6.98 (m, 2H); 7.08 (s, 1H); 7.32-7.35 (t, 1H); 7.45-7.47 (m, 1H); 7.50-7.51 (m, 2H). MS m/z 324.4 (M+H)⁺; ESI-FTICR-MS: m/z 324.18209 ([M+H]⁺, calc. for C₁₈H₂₂N₅O⁺ 324,18189) HPLC (λ = 214 nm, [B]) rt: 7.52 min (97.0 %).

Example 20: N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxy-phenyl)-2-methylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.10 g (14.5 %). ^1H NMR: (CDCl₃) δ 2.09-2.15 (m, 2H); 2.56 (s, 3H); 3.39-3.43 (m, 2H); 3.87 (s, 3H); 3.93 (s, 6H); 4.06-4.08 (t, 2H); 4.84 (br s, 1H); 6.37 (s, 1H); 6.94 (s, 1H); 7.07 (s, 1H); 7.15 (s, 2H); 7.51 (s, 1H). MS m/z 384 (M+H)⁺; ESI-FTICR-MS: m/z 384.20342 ([M+H]⁺, calc. for C₂₀H₂₆N₅O₃ 384.20342) HPLC (λ = 214 nm, [B]) rt: 8.24 min (100 %).

Example 21: N-(3-(1H-imidazol-1-yl)propyl)-6-(4-methoxyphenyl)-2-methylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.3 g (10.5 %). ^1H NMR: (CDCl₃) δ 2.09-2.14 (m, 2H); 2.55 (s, 3H); 3.36-3.40 (m, 2H); 3.84 (s, 3H); 4.05-4.08 (t, 2H); 4.82 (br s, 1H); 6.37 (s, 1H); 6.93-6.96 (m, 3H); 7.09 (br s, 1H); 7.52 (br s, 1H); 7.88-7.91 (m, 2H). MS m/z 324.4 (M+H)⁺; ESI-FTICR-MS: m/z 324.18203 ([M+H]⁺, calc. for C₁₈H₂₂N₅O⁺ 324,18189) HPLC (λ = 214 nm, [B]) rt: 6.19 min (99.2 %).

Example 22: N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-methylpyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

5 Yield (over all steps): 0.29 g (39.3 %). ^1H NMR: (CDCl_3) δ 2.09-2.13 (m, 2H); 2.55 (s, 3H); 3.35-3.39 (m, 2H); 3.83 (s, 3H); 4.04-4.07 (t, 2H); 4.84 (br s, 1H); 6.68 (s, 1H); 6.94-6.96 (m, 2H); 7.02-7.07 (m, 2H); 7.33-7.37 (m, 1H); 7.50 (br s, 1H); 7.81-7.83 (m, 1H). MS m/z 324.4 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 324.18195 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{18}\text{H}_{22}\text{N}_5\text{O}^+$ 324,18189) HPLC (λ = 214 nm, [C]) rt: 8.90 min (99.4 %).

10

Example 23: N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-(pyridin-4-yl)pyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

15 Yield (over all steps): 0.058 g (7.9 %). ^1H NMR: (CDCl_3) δ 2.21-2.27 (m, 2H); 3.60-3.62 (m, 2H); 4.12-4.15 (t, 2H); 5.35 (br s, 1H); 6.73 (s, 1H); 6.97 (s, 1H); 7.12 (s, 1H); 7.56 (s, 1H); 7.97 (s, 1H); 8.28-2.29 (m, 2H); 8.51 (s, 2H); 8.77-8.78 (m, 2H). MS m/z 493.3 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 493,15749 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{23}\text{H}_{19}\text{F}_6\text{N}_6^+$ 493,15699) HPLC (λ = 214 nm, slowlong) rt: 16.40 min (100 %).

20 **Example 24: N-(3-(1H-imidazol-1-yl)propyl)-6-(4-nitrophenyl)-2-(pyridin-4-yl)pyrimidin-4-amine**

The compound was synthesized according to the methods A, D and G.

25 Yield (over all steps): 0.044 g (1.3 %). ^1H NMR: (DMSO-d_6) δ 2.04-2.08 (m, 2H); 3.48 (br s, 2H); 4.08-4.11 (t, 2H); 6.93 (s, 1H); 7.08 (s, 1H); 7.23 (s, 1H); 7.67 (s, 1H); 7.90-7.92 (m, 1H); 8.25-8.26 (d, 2H); 8.37 (s, 3H); 8.74-8.75 (m, 2H). MS m/z 402.3 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 424,14977 ($[\text{M}+\text{Na}]^+$, calc. for $\text{C}_{21}\text{H}_{19}\text{N}_7\text{NaO}_2^+$ 424,14924) HPLC (λ = 214 nm, [C]) rt: 11.55 min (100 %).

Example 25: N-(3-(1H-imidazol-1-yl)propyl)-2-(pyridin-4-yl)-6-p-tolylpyrimidin-4-amine

30 The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.15 g (16.9 %). ^1H NMR: (CDCl_3) δ 2.18-2.23 (m, 2H); 2.41 (s, 3H); 3.54-3.55 (d, 2H); 4.09-4.12 (t, 2H); 5.07 (br s, 1H); 6.63 (s, 1H); 6.95 (s, 1H); 7.10 (s, 1H); 7.28-7.29 (d, 2H); 7.52 (s, 1H); 7.97-7.99 (d, 2H); 8.30-8.31 (m, 2H); 8.73-8.74 (m, 2H). MS m/z 371.2 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 371,19822 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{22}\text{H}_{23}\text{N}_6^+$ 371.19787) HPLC (λ = 214 nm, [C]) rt: 11.95 min (100 %)

Example 26: N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

5 Yield (over all steps): 0.11 g (13.1 %). ^1H NMR: (CDCl_3) δ 2.18-2.23 (m, 2H); 3.53-3.54 (m, 2H); 3.88 (s, 3H); 4.09-4.12 (t, 2H); 5.05 (br s, 1H); 6.95 (s, 1H); 6.99-7.03 (m, 2H); 7.09-7.12 (m, 2H); 7.39-7.43 (m, 1H); 7.52 (s, 1H); 8.15-8.17 (m, 1H); 8.27-8.28 (m, 2H); 8.71-8.72 (m, 2H). MS m/z 387.2 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 387.19318 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{22}\text{H}_{23}\text{N}_6\text{O}^+$ 387.19279) HPLC (λ = 214 nm, [C]) rt: 9.97 min (96.9 %)

10

Example 27: N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine

The compound was synthesized according to the methods A, D and G.

15 Yield (over all steps): 0.10 g (12.8 %). ^1H NMR: (CDCl_3) δ 2.03-2.08 (m, 2H); 3.45 (br s, 2H); 3.85 (s, 3H); 4.08-4.11 (t, 2H); 6.93-6.95 (d, 2H); 7.08-7.10 (m, 1H); 7.22 (s, 1H); 7.43-7.47 (t, 1H); 7.67-7.73 (m, 3H); 8.24-8.25 (m, 2H); 8.72-8.73 (m, 2H). MS m/z 387.3 ($\text{M}+\text{H}$) $^+$; ESI-FTICR-MS: m/z 387.19311 ($[\text{M}+\text{H}]^+$, calc. for $\text{C}_{22}\text{H}_{23}\text{N}_6\text{O}^+$ 387.19279) HPLC (λ = 214 nm, [C]) rt: 11.39 min (100 %).

20 **Example 28: 6-(2-Methoxyphenyl)-N-(3-(5-methyl-1H-imidazol-1-yl)propyl)-2-(pyridin-4-yl)pyrimidin-4-amine**

The compound was synthesized according to the methods A, D and G

25 Yield (over all steps): 0.031 g (7.0 %). ^1H NMR: (CDCl_3) δ 2.17 (s, 2H); 2.22 (s, 3H); 3.54 (br s, 2H); 3.88 (s, 3H); 4.02 (br s, 2H); 5.12 (br s, 1H); 6.98-7.12 (m, 3H); 7.38-7.43 (m, 1H); 8.14-8.17 (m, 1H); 8.28 (br s, 2H); 8.71 (br s, 1H). MS m/z 401.2 ($\text{M}+\text{H}$) $^+$; HPLC (λ = 214 nm, [C]) rt: 10.75 min (97.0 %).

Example 29: 6-(3,4,5-Trimethoxyphenyl)-2-methyl-N-(3-(5-methyl-1H-imidazol-1-yl)-propyl)pyrimidin-4-amine

30 The compound was synthesized according to the methods A, D and G.

Yield (over all steps): 0.21 g (15.1 %). ^1H NMR: (DMSO-d_6) δ 1.99-2.02 (m, 2H); 2.17 (s, 3H); 3.27-3.31 (m, 2H); 3.44 (br s, 1H); 3.88 (s, 3H); 3.99-4.02 (t, 2H); 7.08-7.18 (m, 3H); 7.43-7.47 (m, 1H); 7.71 (br s, 1H); 8.07-8.09 (d, 1H); 8.20-8.21 (d, 2H); 8.71 (br s, 2H). MS m/z 395.5 ($\text{M}+\text{H}$) $^+$; HPLC (λ = 214 nm, [C]) rt: 8.83 min (99.5 %).

35

Example 30: N-(2-phenyl-6-(quinolin-6-yl)pyrimidin-4-yl)-3H-benzo[d]imidazol-5-amine

The compound was synthesized according to the methods A, D and H.

Yield (over all steps): 0.097 g (11.4%). ^1H NMR: (DMSO-d₆) δ 7.43 (s, 1H); 7.57-7.58 (m, 3H); 7.63-7.66 (br m, 1H); 7.69-7.70 (m, 1H); 7.84-7.86 (m, 1H); 8.19-8.21 (m, 1H); 8.50-8.54 (m, 1H); 8.57-8.60 (m, 2H); 8.62-8.64 (m, 1H); 8.76 (s, 1H); 8.85 (s, 1H); 8.98-9.00 (m, 1H); 9.44 (s, 1H); 10.25 (s, 1H). MS m/z 415.2 (M+H)⁺; HPLC (λ = 214 nm, gradientH1) rt: 12.72 min (100%).

Example 31: N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-(quinolin-6-yl)pyrimidin-4-amine

The compound was synthesized according to the methods A, D and I.

Yield (over all steps): 0.089 g (8.4%). ^1H NMR (DMSO-d₆): δ 2.15-2.20 (m, 2H); 2.60 (s, 3H); 3.52 (br s, 2H); 4.28-4.31 (t, 2H, J=6.6 Hz); 7.01 (br s, H); 7.69-7.72 (m, 2H); 7.82 (s, H); 8.24-8.26 (d, H, J=8.7 Hz); 8.55-8.56 (d, H, J=8.7 Hz); 9.05-9.07 (dd, H, J=1.7 Hz); 9.13 (s, H). MS m/z 245.1 (M+H)⁺. HPLC (λ = 214 nm, [C]) rt: 4.99 min (91.0%).

15 **Example 32: N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-6-(2-phenylcyclopropyl)pyrimidin-4-amine**

The compound was synthesized according to the methods A, D and I.

Yield (over all steps): 0.376 g (29.4%). ^1H NMR (DMSO-d₆): δ 1.51 (br s, H); 1.78 (br s, H); 2.10-2.17 (quintuplett, 2H, J=7.0 Hz); 2.28 (br s, H); 2.51-2.54 (m, H); 3.44 (m, 2H); 4.27-4.30 (t, 2H, J=7.0 Hz); 6.38 (br s, H); 7.16-7.21 (m, 3H); 7.27-7.31 (m, 2H); 7.48-7.50 (m, 3H); 7.69-7.72 (t, H, J=1.7 Hz); 7.80-7.83 (t, H, J=1.7 Hz); 8.21-8.23 (m, 2H); 9.13 (s, H). MS m/z 396.3 (M+H)⁺. HPLC (λ = 214 nm, [C]) rt: 13.31 min (100%).

25 **Example 33: N-(3-(1H-imidazol-1-yl)propyl)-6-(2-phenylcyclopropyl)-2-(pyridin-4-yl)-pyrimidin-4-amine**

The compound was synthesized according to the methods A, D and I.

Yield (over all steps): 0.044 g (3.4%). ^1H NMR (DMSO-d₆): δ 1.50 (br s, H); 1.79 (br s, H); 2.10-2.16 (quintuplett, 2H, J=7.1 Hz); 2.27-2.32 (m, H); 2.51-2.55 (m, H); 3.46 (br s, H); 4.26-4.30 (t, 2H, J=7.1 Hz); 6.51 (br s, H); 7.18-7.20 (m, 3H); 7.26-7.30 (m, 2H); 7.71 (s, H); 7.81 (s, H); 8.30-8.32 (d, H, J=5.4 Hz); 8.80-8.82 (d, H, J=5.8 Hz); 9.12 (s, H). MS m/z 397.3 (M+H)⁺. HPLC (λ = 214 nm, [C]) rt: 13.27 min (100%).

35 **Example 34: N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-methyl-pyrimidin-4-amine**

The compound was synthesized according to the methods A, D and I.

Yield (over all steps): 0.248 g (23.0%). ^1H NMR (DMSO-d₆): δ 2.06-2.13 (m, 2H); 3.05-3.10 (m, 2H); 3.31-3.44 (m, 4H); 3.65-3.73 (m, H); 4.21-4.29 (t, 2H, J=7.1 Hz); 6.43 (s, H); 7.18-7.22 (m, 2H); 7.26-7.31 (m, 2H); 7.68 (s, H); 7.77 (s, H); 9.08 (s, H). MS m/z 334.1 (M+H)⁺. HPLC (λ = 214 nm, [C]) rt: 8.85 min (100%).

5

Example 35: N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-phenyl-pyrimidin-4-amine

The compound was synthesized according to the methods A, D and I.

Yield (over all steps): 0.589 g (46.1%). ^1H NMR (DMSO-d₆): δ 2.12-2.15 (quintuplet, 2H, J=7.1 Hz); 3.16-3.22 (m, 2H); 3.28-3.34 (m, 2H); 3.48 (br s, 2H); 3.70-3.74 (m, H); 4.27-3.30 (t, 2H, J=7.1 Hz); 6.39 (br s, H); 7.16-7.18 (m, 2H); 7.24-7.28 (m, 2H); 7.50-7.53 (m, 3H); 7.70 (s, H); 7.81 (s, H); 8.19-8.21 (d, 2H, J=6.6 Hz); 9.13 (s, H). MS m/z 396.1 (M+H)⁺. HPLC (λ = 214 nm, [C]) rt: 12.19 min (100%).

15 **Example 36: N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-(pyridin-4-yl)-pyrimidin-4-amine**

The compound was synthesized according to the methods A, D and I.

Yield (over all steps): 0.275 g (21.5%). ^1H NMR (DMSO-d₆): δ 2.10-2.17 (quintuplet, 2H, J=7.1); 3.17-3.30 (m, 4H); 3.46 (br s, 2H); 3.63-3.72 (m, H); 4.27-3.31 (t, 2H, J=7.1); 6.44 (br s, H); 7.15-7.18 (m, 2H); 7.23-7.27 (m, 2H); 7.71 (t, H, J=1.7); 7.81 (t, H, J=1.7); 8.24-8.26 (d, 2H, J=6.2); 8.76-8.77 (d, 2H, J=6.2); 9.12 (s, H). MS m/z 397.3 (M+H)⁺. HPLC (λ = 214 nm, [C]) rt: 12.08 min (100%).

Activity screening

25 **Fluorometric assays**

All measurements were performed with a BioAssay Reader HTS-7000Plus for microplates (Perkin Elmer) at 30 °C. QC activity was evaluated fluorometrically using H-Gln- β NA. The samples consisted of 0.2 mM fluorogenic substrate, 0.25 U pyroglutamyl aminopeptidase (Unizyme, Hørsholm, Denmark) in 0.2 M Tris/HCl, pH 8.0 containing 20 mM EDTA and an 30 appropriately diluted aliquot of QC in a final volume of 250 μ l. Excitation/emission wavelengths were 320/410 nm. The assay reactions were initiated by addition of glutaminyl cyclase. QC activity was determined from a standard curve of β -naphthylamine under assay conditions. One unit is defined as the amount of QC catalyzing the formation of 1 μ mol pGlu- β NA from H-Gln- β NA per minute under the described conditions.

35

In a second fluorometric assay, QC was activity determined using H-Gln-AMC as substrate. Reactions were carried out at 30 °C utilizing the NOVOStar reader for microplates (BMG labtechnologies). The samples consisted of varying concentrations of the fluorogenic substrate, 0.1 U pyroglutamyl aminopeptidase (Qiagen) in 0.05 M Tris/HCl, pH 8.0 containing 5 mM EDTA and an appropriately diluted aliquot of QC in a final volume of 250 µl. Excitation/emission wavelengths were 380/460 nm. The assay reactions were initiated by addition of glutaminyl cyclase. QC activity was determined from a standard curve of 7-amino-4-methylcoumarin under assay conditions. The kinetic data were evaluated using GraFit sofware.

10 **Spectrophotometric assay of QC**

This novel assay was used to determine the kinetic parameters for most of the QC substrates. QC activity was analyzed spectrophotometrically using a continuous method, that was derived by adapting a previous discontinuous assay (Bateman, R. C. J. 1989 *J Neurosci Methods* 30, 23-28) utilizing glutamate dehydrogenase as auxiliary enzyme. Samples consisted of the 15 respective QC substrate, 0.3 mM NADH, 14 mM α-Ketoglutaric acid and 30 U/ml glutamate dehydrogenase in a final volume of 250 µl. Reactions were started by addition of QC and pursued by monitoring of the decrease in absorbance at 340 nm for 8-15 min.

20 The initial velocities were evaluated and the enzymatic activity was determined from a standard curve of ammonia under assay conditions. All samples were measured at 30 °C, using either the SPECTRAFluor Plus or the Sunrise (both from TECAN) reader for microplates. Kinetic data was evaluated using GraFit software.

Inhibitor assay

25 For inhibitor testing, the sample composition was the same as described above, except of the putative inhibitory compound added. For a rapid test of QC-inhibition, samples contained 4 mM of the respective inhibitor and a substrate concentration at 1 K_m . For detailed investigations of the inhibition and determination of K_i -values, influence of the inhibitor on the auxiliary enzymes was investigated first. In every case, there was no influence on either enzyme detected, thus 30 enabling the reliable determination of the QC inhibition. The inhibitory constant was evaluated by fitting the set of progress curves to the general equation for competitive inhibition using GraFit software.

Analytical methods

35 The analytical HPLC-system consisted of a Merck-Hitachi device (model LaChrom®) utilizing a Li-Chrospher® 100 RP 18 (5 µm), analytical column (length: 125 mm, diameter: 4 mm), and a

diode array detector (DAD) with $\lambda = 214$ nm as the reporting wavelength. The compounds were analyzed using a gradient at a flow rate of 1 mL/min; whereby eluent (A) was acetonitrile, eluent (B) was water, both containing 0.1 % (v/v) trifluoro acetic acid applying the following gradient: Method [A]: 0 min - 5 min \rightarrow 5% (A), 5 min - 17 min \rightarrow 5 - 15% (A), 15 min - 27 min \rightarrow 15 - 95% 5 (A) 27 min - 30 min \rightarrow 95% (A). Method [B]: 0 min - 15 min \rightarrow 5 - 50 % (A), 15 min - 20 min \rightarrow 50 - 95 % (A), 20 min - 23 min \rightarrow 95 % (A). Method [C]: 0 min - 20 min \rightarrow 5 - 60 % (A), 20 min - 25 min \rightarrow 60 - 95 % (A), 25 min - 30 min \rightarrow 95 % (A). The purities of all reported compounds were determined by the percentage of the peak area at 214 nm.

10 ESI-Mass spectra were obtained with a SCIEX API 365 spectrometer (Perkin Elmer) utilizing the positive ionization mode.

The high resolution positive ion ESI mass spectra were obtained from a Bruker Apex III 70e Fourier transform ion cyclotron resonance mass spectrometer (Bruker Daltonics, Billerica, USA) equipped with an Infinity™ cell, a 7.0 Tesla superconducting magnet (Bruker, Karlsruhe, Germany), an RF-only hexapole ion guide and an external electrospray ion source (API Apollo, 15 voltages: endplate, -3.700V; capillary, -4.400V; capillary exit, 100V; skimmer 1.15 V; skimmer 2.6 V). Nitrogen was used as drying gas at 150°C. The sample solutions were introduced continuously via a syringe pump with a flow rate of 120 μ l h⁻¹. All data were acquired with 256 k data points and zero filled to 1024 k by averaging 32 scans.

10 The melting points were detected utilizing a Kofler melting point device. They are not corrected 20 The ¹H NMR-Spectra (500 MHz) were recorded at a BRUKER AC 500. The solvent was DMSO-D₆, unless otherwise specified. Chemical shifts are expressed as parts per million (ppm) downfiled from tetramethylsilan. Splitting patterns have been designated as follows: s (singlet), d (doublet), dd (doublet of doublet), t (triplet), m (multiplet) and br (broad signal).

25 **MALDI-TOF mass spectrometry**

Matrix-assisted laser desorption/ionization mass spectrometry was carried out using the Hewlett-Packard G2025 LD-TOF System with a linear time of flight analyzer. The instrument was equipped with a 337 nm nitrogen laser, a potential acceleration source (5 kV) and a 1.0 m flight tube. Detector operation was in the positive-ion mode and signals are recorded and 30 filtered using LeCroy 9350M digital storage oscilloscope linked to a personal computer. Samples (5 μ l) were mixed with equal volumes of the matrix solution. For matrix solution DHAP/DAHC was used, prepared by solving 30 mg 2',6'-dihydroxyacetophenone (Aldrich) and 44 mg diammonium hydrogen citrate (Fluka) in 1 ml acetonitrile/0.1% TFA in water (1/1, v/v). A small volume (\approx 1 μ l) of the matrix-analyte-mixture was transferred to a probe tip and 35 immediately evaporated in a vacuum chamber (Hewlett-Packard G2024A sample prep accessory) to ensure rapid and homogeneous sample crystallization.

For long-term testing of Glu¹-cyclization, A β -derived peptides were incubated in 100 μ l 0.1 M sodium acetate buffer, pH 5.2 or 0.1 M Bis-Tris buffer, pH 6.5 at 30 °C. Peptides were applied in 0.5 mM [A β (3-11)a] or 0.15 mM [A β (3-21)a] concentrations, and 0.2 U QC is added all 24 hours.

In case of A β (3-21)a, the assays contained 1 % DMSO. At different times, samples are removed

5 from the assay tube, peptides extracted using ZipTips (Millipore) according to the manufacturer's recommendations, mixed with matrix solution (1:1 v/v) and subsequently the mass spectra recorded. Negative controls either contain no QC or heat deactivated enzyme. For the inhibitor studies the sample composition was the same as described above, with exception of the inhibitory compound added (5 mM or 2 mM of a test compound of the invention).

10

The first QC inhibitors were disclosed in WO 2004/098591 and WO 2005/075436. There are no other potent QC inhibitors known in the art. The same holds true for combinations and compositions for the treatment of neuronal diseases comprising QC inhibitors. Compounds and

15 combinations of the invention may have the advantage that they are, for example, more potent, more selective, have fewer side-effects, have better formulation and stability properties, have better pharmacokinetic properties, be more bioavailable, be able to cross blood brain barrier and are more effective in the brain of mammals, are more compatible or effective in combination with other drugs or be more readily synthesized than other compounds of the prior art.

20 Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer, step, group of integers or group of steps but not to the exclusion of any other integer, step, group of integers or group of steps.


25 All patents and patent applications mentioned throughout the specification of the present invention are herein incorporated in their entirety by reference.

The invention embraces all combinations of preferred and more preferred groups and embodiments of groups recited above.

30

Claims

1. A compound of formula (I),



or a pharmaceutically acceptable salt, solvate or polymorph thereof, including all tautomers and stereoisomers thereof wherein:

R¹ represents alkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo; and in which any of the aforesaid aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl), -C(O)OC₁₋₆alkyl, SOC₁₋₄alkyl and -SOC₃₋₆cycloalkyl;

or R¹ represents -carbocyclyl-phenyl; -carbocyclyl-(monocyclic heteroaryl); -heterocyclyl-phenyl; -carbocyclyl fused to phenyl; carbocyclyl fused to monocyclic heteroaryl or heterocyclyl fused to phenyl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;


and in which any of the aforesaid phenyl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl) and -SOC₁₋₄alkyl;

and

R² represents alkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo; and in which any of the aforesaid aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl), -C(O)OC₁₋₆alkyl, SOC₁₋₄alkyl and -SOC₃₋₆cycloalkyl; or R¹ represents -carbocyclyl-phenyl; -carbocyclyl-(monocyclic heteroaryl); -heterocyclyl-phenyl; -carbocyclyl fused to phenyl; carbocyclyl fused to monocyclic heteroaryl or heterocyclyl fused to phenyl; which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo; and in which any of the aforesaid phenyl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂, -C(O)NH(C₁₋₄alkyl) and -SOC₁₋₄alkyl; and


A represents



wherein Y represents a C₂₋₅ alkylene chain, which may optionally be substituted by one or two methyl groups or may optionally be substituted by two alkylene substituents at the same position wherein the two alkylene substituents are joined to each other to form a C₃₋₅ spiro-cycloalkyl group and

R³, R⁴ and R⁵ independently represent H or C₁₋₂alkyl; or

A represents



wherein Z represents a bond, -CH₂-, -CH₂-CH₂-, -CH(Me)-, -CH(Me)-CH₂- or -CH₂-CH(Me)- and

R⁶ and R⁷ independently represent H or C₁₋₂alkyl,
and

B represents H or methyl.

2. A compound according to claim 1 wherein:

R¹ represents alkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;


and in which any of the aforesaid aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂ and -C(O)NH(C₁₋₄alkyl); and

R² represents alkyl; alkenyl; carbocyclyl; heterocyclyl; optionally substituted aryl or optionally substituted heteroaryl;

which any of the aforesaid carbocyclyl and heterocyclyl groups may optionally be substituted by one or more groups selected from methyl and oxo;

and in which any of the aforesaid aryl and heteroaryl groups may optionally be substituted by one or more substituents selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆haloalkyl, -C₁₋₆thioalkyl, -SO₂C₁₋₄alkyl, C₁₋₆alkoxy-, -O-C₃₋₈cycloalkyl, C₃₋₈cycloalkyl, -SO₂C₃₋₈cycloalkyl, C₃₋₆alkenyloxy-, C₃₋₆alkynyoxy-, -C(O)C₁₋₆alkyl, C₁₋₆alkoxy-C₁₋₆alkyl-, nitro, halogen, cyano, hydroxyl, -C(O)OH, -NH₂, -NHC₁₋₄alkyl, -N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)N(C₁₋₄alkyl)(C₁₋₄alkyl), -C(O)NH₂ and -C(O)NH(C₁₋₄alkyl); and


A represents



wherein Y represents a C_{2-5} alkylene chain, which may optionally be substituted by one or two methyl groups or may optionally be substituted by two alkylene substituents at the same position wherein the two alkylene substituents are joined to each other to form a C_{3-5} spiro-cycloalkyl group and

R^3 , R^4 and R^5 independently represent H or C_{1-2} alkyl; or

A represents




wherein Z represents a bond, $-CH_2-$, $-CH_2-CH_2-$, $-CH(Me)-$, $-CH(Me)-CH_2-$ or $-CH_2-CH(Me)-$ and

R^6 and R^7 independently represent H or C_{1-2} alkyl,


and

B represents H or methyl.


3. A compound according to claim 1 or claim 2 wherein R^2 represents alkyl.
4. A compound according to any one of claims 1 to 3 wherein R^2 represents optionally substituted aryl.
5. A compound according to any one of claims 1 to 4 wherein R^2 represents optionally substituted heteroaryl.
6. A compound according to any one of claims 1 to 5 wherein R^1 represents alkyl, optionally substituted aryl; carbocyclyl; haloalkyl or optionally substituted heteroaryl.
7. A compound according to claim 6 wherein R^1 represents optionally substituted aryl.
8. A compound according to any one of claims 1 to 7 wherein A represents




9. A compound according to claim 8 wherein R^3 , R^4 and R^5 do not all represent H.
10. A compound according to claim 8 wherein R^3 represents H, R^4 represents H and R^5 represents methyl.
11. A compound according to any one of claims 8 to 10 wherein Y represents an unsubstituted C_{2-5} alkylene chain.
12. A compound according to any one of claims 8 to 11 wherein Y does not represent $-(CH_2)_3-$.
13. A compound according to claim 11 wherein Y represents $-(CH_2)_4-$.
14. A compound according to claim 8 or claim 13 wherein A represents



15. A compound according to claim 8 or claim 13 wherein A represents



16. A compound according to any one of claims 1 to 7 wherein A represents



17. A compound according to any one of claims 16 wherein Z represents a bond, $-CH_2-$ or $-CH_2CH_2-$.
18. A compound according to any one of claims 1 to 17 wherein B represents H.
19. A compound according to claim 1 wherein the compound of formula (I) is represented

by any one of Examples 1 to 30:

- 1) N-(3-(1H-imidazol-1-yl)propyl)-6-methyl-2-phenylpyrimidin-4-amine;
- 2) N-(3-(1H-imidazol-1-yl)propyl)-2,6-diphenylpyrimidin-4-amine;
- 3) N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-phenylpyrimidin-4-amine;
- 4) N-(3-(1H-imidazol-1-yl)propyl)-2,6-dimethylpyrimidin-4-amine;
- 5) N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-p-tolylpyrimidin-4-amine;
- 6) N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-methylpyrimidin-4-amine;
- 7) N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-phenylpyrimidin-4-amine;
- 8) N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxyphenyl)-2-phenylpyrimidin-4-amine;
- 9) N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-6-p-tolylpyrimidin-4-amine;
- 10) N-(3-(1H-imidazol-1-yl)propyl)-6-(4-nitrophenyl)-2-phenylpyrimidin-4-amine;
- 11) N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-phenylpyrimidin-4-amine;
- 12) N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-phenylpyrimidin-4-amine;
- 13) N-(3-(1H-imidazol-1-yl)propyl)-6-(furan-3-yl)-2-phenylpyrimidin-4-amine;
- 14) N-(3-(1H-imidazol-1-yl)propyl)-6-(adamant-1-yl)-2-phenylpyrimidin-4-amine;
- 15) N-(3-(1H-imidazol-1-yl)propyl)-6-(trifluoromethyl)-2-phenylpyrimidin-4-amine;
- 16) N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine;
- 17) N-(3-(1H-imidazol-1-yl)propyl)-6-(furan-3-yl)-2-(pyridin-4-yl)pyrimidin-4-amine;
- 18) N-(3-(1H-imidazol-1-yl)propyl)-6-(trifluoromethyl)-2-(pyridin-4-yl)pyrimidin-4-amine;
- 19) N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-methylpyrimidin-4-amine;
- 20) N-(3-(1H-imidazol-1-yl)propyl)-6-(3,4,5-trimethoxy-phenyl)-2-methylpyrimidin-4-amine;
- 21) N-(3-(1H-imidazol-1-yl)propyl)-6-(4-methoxyphenyl)-2-methylpyrimidin-4-amine;
- 22) N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-methylpyrimidin-4-amine;
- 23) N-(3-(1H-imidazol-1-yl)propyl)-6-(3,5-bis(trifluoromethyl)phenyl)-2-(pyridin-4-yl)pyrimidin-4-amine;

24) N-(3-(1H-imidazol-1-yl)propyl)-6-(4-nitrophenyl)-2-(pyridin-4-yl)pyrimidin-4-amine;

25) N-(3-(1H-imidazol-1-yl)propyl)-2-(pyridin-4-yl)-6-p-tolylpyrimidin-4-amine

26) N-(3-(1H-imidazol-1-yl)propyl)-6-(2-methoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine;

27) N-(3-(1H-imidazol-1-yl)propyl)-6-(3-methoxyphenyl)-2-(pyridin-4-yl)pyrimidin-4-amine;

28) 6-(2-methoxyphenyl)-N-(3-(5-methyl-1H-imidazol-1-yl)propyl)-2-(pyridin-4-yl)pyrimidin-4-amine;

29) 6-(3,4,5-trimethoxyphenyl)-2-methyl-N-(3-(5-methyl-1H-imidazol-1-yl)propyl)pyrimidin-4-amine;

30) N-(2-phenyl-6-(quinolin-6-yl)pyrimidin-4-yl)-3H-benzo[d]imidazol-5-amine;
or a pharmaceutically acceptable salt, solvate or polymorph thereof, including all tautomers and stereoisomers thereof;
or any one of Examples 31 to 36

31) N-(3-(1H-imidazol-1-yl)propyl)-2-methyl-6-(quinolin-6-yl)pyrimidin-4-amine;

32) N-(3-(1H-imidazol-1-yl)propyl)-2-phenyl-6-(2-phenylcyclopropyl)pyrimidin-4-amine;

33) N-(3-(1H-imidazol-1-yl)propyl)-6-(2-phenylcyclopropyl)-2-(pyridin-4-yl)pyrimidin-4-amine;

34) N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-methylpyrimidin-4-amine;

35) N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-phenylpyrimidin-4-amine;

36) N-(3-(1H-imidazol-1-yl)propyl)-6-(2,3-dihydro-1H-inden-2-yl)-2-(pyridin-4-yl)pyrimidin-4-amine;
or a pharmaceutically acceptable salt, solvate or polymorph thereof, including all tautomers and stereoisomers thereof.

20. A compound according to claims 1 to 19 for use as a medicament.

21. A pharmaceutical composition comprising a compound according to any one of claims 1 to 19 optionally in combination with one or more therapeutically acceptable diluents or carriers.

22. The pharmaceutical composition of claim 21 which comprises additionally at least one compound, selected from the group consisting of neuroprotectants, antiparkinsonian drugs, amyloid protein deposition inhibitors, beta amyloid synthesis inhibitors, antidepressants, anxiolytic drugs, antipsychotic drugs and anti-multiple

sclerosis drugs.

23. The pharmaceutical composition according to any one of claims 21 or 22, which comprises additionally at least one compound, selected from the group consisting of PEP-inhibitors, LiCl, inhibitors of inhibitors of DP IV or DP IV-like enzymes, acetylcholinesterase (ACE) inhibitors, PIMT enhancers, inhibitors of beta secretases, inhibitors of gamma secretases, inhibitors of neutral endopeptidase, inhibitors of Phosphodiesterase-4 (PDE-4), TNFalpha inhibitors, muscarinic M1 receptor antagonists, NMDA receptor antagonists, sigma-1 receptor inhibitors, histamine H3 antagonists, immunomodulatory agents, immunosuppressive agents or an agent selected from the group consisting of antegren (natalizumab), Neurelan (fampridine-SR), campath (alemtuzumab), IR 208, NBI 5788/MSP 771 (tiplimotide), paclitaxel, Anergix.MS (AG 284), SH636, Differin (CD 271, adapalene), BAY 361677 (interleukin-4), matrix-metalloproteinase-inhibitors, interferon-tau (trophoblastin) and SAIK-MS.

24. The compound of any one of claims 1 to 19 for use in the treatment of a disease selected from the group consisting of Kennedy's disease, duodenal cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zolliger-Ellison syndrome, gastric cancer with or without *Helicobacter pylori* infections, pathogenic psychotic conditions, schizophrenia, infertility, neoplasia, inflammatory host responses, cancer, malign metastasis, melanoma, psoriasis, impaired humoral and cell-mediated immune responses, leukocyte adhesion and migration processes in the endothelium, impaired food intake, impaired sleep-wakefulness, impaired homeostatic regulation of energy metabolism, impaired autonomic function, impaired hormonal balance or impaired regulation of body fluids, multiple sclerosis, the Guillain-Barré syndrome and chronic inflammatory demyelinizing polyradiculoneuropathy.

25. The compound of any one of claims 1 to 19 for use in the treatment of a disease selected from the group consisting of mild cognitive impairment, Alzheimer's disease, Familial British Dementia, Familial Danish Dementia, neurodegeneration in Down Syndrome, Parkinson's disease and Huntington's disease.

26. The compound of any one of claims 1 to 19 for use in the treatment of a disease selected from the group consisting of rheumatoid arthritis, atherosclerosis, pancreatitis and restenosis.


27. A method of treatment or prevention of a disease selected from the group consisting of Kennedy's disease, duodenal cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zolliger-Ellison syndrome, gastric cancer with or without *Helicobacter pylori* infections, pathogenic psychotic conditions, schizophrenia,

infertility, neoplasia, inflammatory host responses, cancer, malign metastasis, melanoma, psoriasis, impaired humoral and cell-mediated immune responses, leukocyte adhesion and migration processes in the endothelium, impaired food intake, impaired sleep-wakefulness, impaired homeostatic regulation of energy metabolism, impaired autonomic function, impaired hormonal balance or impaired regulation of body fluids, multiple sclerosis, the Guillain-Barré syndrome and chronic inflammatory demyelinizing polyradiculoneuropathy, which comprises administering to a subject an effective amount of a compound of formula (I) according to any one of claims 1 to 19 or a pharmaceutical composition according to any one of claims 21 to 23.

28. A method of treatment or prevention of a disease selected from the group consisting of mild cognitive impairment, Alzheimer's disease, Familial British Dementia, Familial Danish Dementia, neurodegeneration in Down Syndrome, Parkinson's disease and Huntington's disease, which comprises administering to a subject an effective amount of a compound of formula (I) according to any one of claims 1 to 19 or a pharmaceutical composition according to any one of claims 21 to 23.
29. A method of treatment or prevention of a disease selected from the group consisting of rheumatoid arthritis, atherosclerosis, pancreatitis and restenosis, which comprises administering to a subject an effective amount of a compound of formula (I) according to any one of claims 1 to 19 or a pharmaceutical composition according to any one of claims 21 to 23.
30. Use of a compound of formula (I) according to any one of claims 1 to 19 in the manufacture of a medicament for the treatment of a disease selected from the group consisting of Kennedy's disease, duodenal cancer with or w/o *Helicobacter pylori* infections, colorectal cancer, Zollinger-Ellison syndrome, gastric cancer with or without *Helicobacter pylori* infections, pathogenic psychotic conditions, schizophrenia, infertility, neoplasia, inflammatory host responses, cancer, malign metastasis, melanoma, psoriasis, impaired humoral and cell-mediated immune responses, leukocyte adhesion and migration processes in the endothelium, impaired food intake, impaired sleep-wakefulness, impaired homeostatic regulation of energy metabolism, impaired autonomic function, impaired hormonal balance or impaired regulation of body fluids, multiple sclerosis, the Guillain-Barré syndrome and chronic inflammatory demyelinizing polyradiculoneuropathy;
wherein the following compounds are disclaimed from the definition of formula (I):
N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-2-(2-thienyl)-4-pyrimidinamine,
2-(2-furanyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-(1H-pyrazol-1-yl)-4-pyrimidinamine and


2-(4-chlorophenyl)-N-[3-(1H-imidazol-1-yl)propyl]-6-phenyl-4-pyrimidinamine.

31. The use of a compound of formula (I) according to any one of claims 1 to 19 for the treatment of a disease selected from the group consisting of mild cognitive impairment, Alzheimer's disease, Familial British Dementia, Familial Danish Dementia, neurodegeneration in Down Syndrome, Parkinson's disease and Huntington's disease.
32. The use of a compound of formula (I) according to any one of claims 1 to 19 for the treatment of a disease selected from the group consisting of rheumatoid arthritis, atherosclerosis, pancreatitis and restenosis.
33. A process for preparation of a compound of formula (I) according to any one of claims 1 to 19 which comprises reaction of a compound of formula (II)



(II)

wherein X represents a leaving group
with a compound of formula (III)



(III)

wherein R¹, X, A and B are defined as in any one of claims 1 to 19.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2008/054714

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D403/12 C07D401/14 A61K31/506 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, BEILSTEIN Data, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DESAI PRASHANT V ET AL: "Identification of novel parasitic cysteine protease inhibitors by use of virtual screening. 2. The available chemical directory" JOURNAL OF MEDICINAL CHEMISTRY, vol. 49, no. 5, March 2006 (2006-03), pages 1576-1584, XP002492846 ISSN: 0022-2623 page 1582; compounds 23-26</p> <p>-----</p> <p style="text-align: center;">-/--</p>	1-4, 6, 8, 11, 18, 20, 21, 33

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
20 August 2008	08/09/2008

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016	Authorized officer Steedijk, Martin
---	--

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2008/054714

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2005/058883 A (ALMIRALL PRODESFARMA SA [ES]; CRESPO CRESPO MARIA ISABEL [ES]; PRAT QU) 30 June 2005 (2005-06-30) claim 1 examples 128,138 pages 8,44 -----	1,2,5,6, 8,11,12, 20-22, 25,26, 28,29, 31-33
X	KATIYAR S B ET AL: "Syntheses of 2,4,6-trisubstituted pyrimidine derivatives as a new class of antifilarial topoisomerase II inhibitors" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 15, no. 1, 3 January 2005 (2005-01-03), pages 47-50, XP004694208 ISSN: 0960-894X page 49; compound 24 -----	1,2,4,6, 8,11,18, 20,21,33
X	WO 2004/048365 A (CHIRON CORP [US]; NUSS JOHN M [US]; PECCHI SABINA [US]; RENHOWE PAUL A) 10 June 2004 (2004-06-10) claim 1 examples 68,94 -----	1,2,6-8, 11,12, 16-18, 20,21, 24,27, 30,33
X	WO 03/026661 A (YAMANOUCHI PHARMA CO LTD [JP]; YONETOKU YASUHIRO [JP]; MARUYAMA TATSUY) 3 April 2003 (2003-04-03) abstract pages 55-56; examples 214,215,259 -----	1,2,4,6, 8,11,12, 18,20, 21,24, 27,30,33
X	WO 00/09496 A (AVENTIS PHARMA GMBH [DE]; SCHINDLER URSULA [DE]; SCHOENAFINGER KARL [D]) 24 February 2000 (2000-02-24) claim 1 page 28; example 58 page 19 -----	1,2,4, 6-8,11, 12,20, 21,24, 26,27, 29,30, 32,33

-/-

INTERNATIONAL SEARCH REPORT

International application No PCT/EP2008/054714

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CHEMCATS [Online] 13 September 2006 (2006-09-13), XP002492847 retrieved from STN Database accession no. 2019237087 abstract see RN 887465-12-5, see also RN: 887464-12-2 to 887423-54-3 -----	1-8,11, 18
A	WO 2004/098591 A (PROBIODRUG AG [DE]; DEMUTH HANS-ULRICH [DE]; HEISER ULRICH [DE]; BUCHH) 18 November 2004 (2004-11-18) claims 1-4 -----	1-33
P,X	WO 2007/084786 A (NOVARTIS AG [CH]; BURGER MATTHEW [US]; NI ZHI-JIE [US]; PECCHI SABINA) 26 July 2007 (2007-07-26) claim 1 page 222; example 521 pages 37-40 pages 44-49 -----	1,2,6, 16-18, 20,21, 23,24, 26,27, 29,30, 32,33

INTERNATIONAL SEARCH REPORT

Information on patent family members

 International application No
 PCT/EP2008/054714

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO 2005058883	A 30-06-2005	AU 2004299461 A1		30-06-2005
		CA 2551944 A1		30-06-2005
		JP 2007514003 T		31-05-2007
WO 2004048365	A 10-06-2004	AU 2003295776 A1		18-06-2004
		CA 2507100 A1		10-06-2004
		EP 1575940 A1		21-09-2005
		JP 2006514118 T		27-04-2006
		KR 20050085114 A		29-08-2005
		MX PA05005477 A		25-07-2005
WO 03026661	A 03-04-2003	NONE		
WO 0009496	A 24-02-2000	AT 240315 T		15-05-2003
		AU 760988 B2		29-05-2003
		AU 5730799 A		06-03-2000
		BR 9913003 A		08-05-2001
		CA 2340405 A1		24-02-2000
		DE 19836697 A1		17-02-2000
		DK 1112266 T3		15-09-2003
		EP 1112266 A1		04-07-2001
		ES 2196849 T3		16-12-2003
		JP 2002522536 T		23-07-2002
		PT 1112266 T		30-09-2003
		US 6844347 B1		18-01-2005
WO 2004093591	A 18-11-2004	EP 1620091 A2		01-02-2006
		JP 2006525276 T		09-11-2006
WO 2007084786	A 26-07-2007	AR 059087 A1		12-03-2008
		AU 2007206048 A1		26-07-2007