

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
13 February 2003 (13.02.2003)

PCT

(10) International Publication Number  
**WO 03/012031 A2**

(51) International Patent Classification<sup>7</sup>: **C12N**  
(21) International Application Number: PCT/US02/22676  
(22) International Filing Date: 16 July 2002 (16.07.2002)  
(25) Filing Language: English  
(26) Publication Language: English  
(30) Priority Data:  
09/918,187 30 July 2001 (30.07.2001) US  
(71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and  
(75) Inventors/Applicants (for US only): **CROOKE, Rosanne, M.** [US/US]; 3211 Piraqua Street, Carlsbad, CA 92009 (US). **GRAHAM, Mark, J.** [US/US]; 2305 S. Ola Vista, San Clemente, CA 92672 (US).  
(74) Agents: **LICATA, Jane, Massey** et al.; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

**Published:**  
— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



**WO 03/012031 A2**

(54) Title: ANTISENSE MODULATION OF STEAROYL-COA DESATURASE EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of stearoyl-CoA desaturase. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding stearoyl-CoA desaturase. Methods of using these compounds for modulation of stearoyl-CoA desaturase expression and for treatment of diseases associated with expression of stearoyl-CoA desaturase are provided.

**ANTISENSE MODULATION OF STEAROYL-COA DESATURASE EXPRESSION****FIELD OF THE INVENTION**

The present invention provides compositions and  
5 methods for modulating the expression of stearoyl-CoA  
desaturase. In particular, this invention relates to  
compounds, particularly oligonucleotides, specifically  
hybridizable with nucleic acids encoding stearoyl-CoA  
desaturase. Such compounds have been shown to modulate  
10 the expression of stearoyl-CoA desaturase.

**BACKGROUND OF THE INVENTION**

Saturated fatty acids are known to be the precursors  
of unsaturated fatty acids in higher organisms. However,  
the control mechanisms that govern the conversion of  
15 saturated fatty acids to unsaturated fatty acids are not  
well understood. The relative amounts of different fatty  
acids has an effect on the physical properties of  
membranes. Furthermore, regulation of unsaturated fatty  
acids is important because they play a role in cellular  
20 activity, metabolism and nuclear events that govern gene  
transcription.

A critical committed step in the biosynthesis of  
mono-unsaturated fatty acids is the introduction of the  
first *cis*-double bond in the delta-9 position (between  
25 carbons 9 and 10). This oxidative reaction is catalyzed by  
stearoyl-CoA desaturase (SCD, also known as delta-9-  
desaturase) and involves cytochrome  $b_5$ , NADH (P)-cytochrome  
 $b_5$  reductase and molecular oxygen (Ntambi, *J. Lipid Res.*,  
**1999**, *40*, 1549-1558). Although the insertion of the double  
30 bond occurs in several different methylene-interrupted  
fatty acyl-CoA substrates, the preferred substrates are  
palmitoyl- and stearoyl-CoA which are converted to

palmitoleoyl- and oleoyl-CoA respectively (Ntambi, *J. Lipid Res.*, **1999**, *40*, 1549-1558).

It has been recognized that, irrespective of diet, the major storage fatty acids in human adipose tissue are  
5 oleic acid, an 18 carbon unsaturated fatty acid, and palmitoleic acid, a 16 carbon unsaturated fatty acid (Ntambi, *J. Lipid Res.*, **1999**, *40*, 1549-1558). During the *de novo* synthesis of fatty acids, the fatty acid synthase enzyme stops at palmitoleic acid but the end product of  
10 the pathway is usually oleic acid (Ntambi, *J. Lipid Res.*, **1999**, *40*, 1549-1558).

The stearoyl-CoA desaturase gene was partially characterized in 1994 via isolation of a 0.76 kb partial  
cDNA from human adipose tissue (Li et al., *Int. J. Cancer*,  
15 **1994**, *57*, 348-352). Increased levels of stearoyl-CoA desaturase mRNA were found in colonic and esophageal carcinomas and in hepatocellular carcinoma (Li et al., *Int. J. Cancer*, **1994**, *57*, 348-352). The gene was fully characterized in 1999 and it was found that alternative  
20 usage of polyadenylation sites generates two transcripts of 3.9 and 5.2 kb (Zhang et al., *Biochem. J.*, **1999**, *340*, 255-264). Two loci for the stearoyl-CoA desaturase gene were mapped to chromosomes 10 and 17 and it was determined that the chromosome 17 loci encodes a transcriptionally  
25 inactive pseudogene (Ntambi, *J. Lipid Res.*, **1999**, *40*, 1549-1558).

Disclosed and claimed in PCT publication WO 00/09754 is a nucleic acid molecule encoding the human stearoyl-CoA desaturase and a nucleic acid molecule which, under  
30 suitable conditions, specifically hybridizes to said nucleic acid molecule encoding the human stearoyl-CoA desaturase (Stenn et al., **2000**).

Stearoyl-CoA desaturase affects the ratio of stearate to oleate which, in turn, affects cell membrane fluidity. Alterations of this ratio have been implicated in various disease states including cardiovascular disease, obesity, 5 non-insulin-dependent diabetes mellitus, skin disease, hypertension, neurological diseases, immune disorders and cancer (Ntambi, *J. Lipid Res.*, 1999, 40, 1549-1558). Stearoyl-CoA desaturase has been viewed as a lipogenic enzyme not only for its key role in the biosynthesis of 10 monounsaturated fatty acids, but also for its pattern of regulation by diet and insulin (Ntambi, *J. Lipid Res.*, 1999, 40, 1549-1558).

The regulation of stearoyl-CoA desaturase is therefore of considerable physiologic importance and its 15 activity is sensitive to dietary changes, hormonal imbalance, developmental processes, temperature changes, metals, alcohol, peroxisomal proliferators and phenolic compounds (Ntambi, *J. Lipid Res.*, 1999, 40, 1549-1558).

Animal models have been very useful in investigations 20 of the regulation of stearoyl-CoA by polyunsaturated fatty acids (PUFAs). For example, in adipose tissue of lean and obese Zucker rats, Jones et al. observed a 75% decrease in stearoyl-CoA desaturase mRNA when both groups were fed a diet high in PUFAs relative to a control diet (Jones et 25 al., *Am. J. Physiol.*, 1996, 271, E44-49).

Similar results have been obtained with tissue culture systems. In the murine 3T3-L1 adipocyte cell line, arachidonic linoleic, linolenic, and eicosapentanoic acids decreased stearoyl-CoA desaturase expression in a 30 dose-dependent manner (Sessler et al., *J. Biol. Chem.*, 1996, 271, 29854-29858).

The molecular mechanisms by which PUFAs regulate stearoyl-CoA desaturase gene expression in different

tissues are still poorly understood. The current understanding of the regulatory mechanism involves binding of PUFAs to a putative PUFA-binding protein, after which repression transcription occurs via binding of the PUFA-  
5 binding protein to a *cis*-acting PUFA response element of the stearoyl-CoA desaturase gene (SREBP) (Ntambi, *J. Lipid Res.*, **1999**, *40*, 1549-1558; Zhang et al., *Biochem. J.*, **2001**, *357*, 183-193).

Cholesterol has also been identified as a regulator  
10 of stearoyl-CoA desaturase gene expression by a mechanism involving repression of the maturation of the sterol regulatory element binding protein (Bene et al., *Biochem. Biophys. Res. Commun.*, **2001**, *284*, 1194-1198; Ntambi, *J. Lipid Res.*, **1999**, *40*, 1549-1558).

15 Thiazolidinediones have been employed as regulators of stearoyl-CoA desaturase activity in murine 3T3-L1 adipocytes (Kim et al., *J. Lipid Res.*, **2000**, *41*, 1310-1316), and in diabetic rodents (Singh Ahuja et al., *Mol. Pharmacol.*, **2001**, *59*, 765-773).

20 Disclosed and claimed in PCT publication WO 99/63979 are compositions comprising a saponin in an amount effective to inhibit stearoyl-CoA desaturase enzyme activity wherein said saponin is derived from a source selected from the group consisting of *Quillaja saponaria*,  
25 *Panax trifolium*, *Panax quinquefolium* and *Glycyrrhiza glabra* (Chavali and Forse, **1999**).

Disclosed and claimed in European patent EP 238198 is an inhibitor of stearoyl-CoA desaturase prepared in a form suitable for oral, parenteral, rectal or dermal  
30 administration for use in modifying the lipid structure of cell membranes wherein said inhibitor consists of a saturated fatty acid having from 12 to 28 carbon atoms in the alkyl chain, e.g. stearic acid, or a pharmaceutically

acceptable derivative thereof prepared in a form suitable for parenteral, rectal or dermal administration (Wood et al., 1987). A stearoyl-CoA desaturase antisense vector has been used to reduce expression levels of stearoyl-CoA desaturase in chicken LMH hepatoma cells (Diot et al.,  
5 *Arch. Biochem. Biophys.*, 2000, 380, 243-250).

To date, investigative strategies aimed at inhibiting stearoyl-CoA desaturase function include the previously cited use of polyunsaturated fatty acids, saturated fatty  
10 acids, thiazolidinediones, cholesterol, and an antisense vector. However, these strategies are untested as therapeutic protocols. Consequently, there remains a long felt need for additional agents capable of effectively inhibiting stearoyl-CoA desaturase.

15 Antisense technology is emerging as an effective means of reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic and research applications involving modulation of stearoyl-CoA desaturase  
20 expression.

The present invention provides compositions and methods for modulating stearoyl-CoA desaturase expression, including inhibition of both isoforms of stearoyl-CoA desaturase.

## 25 **SUMMARY OF THE INVENTION**

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding stearoyl-CoA desaturase, and which modulate the expression of stearoyl-  
30 CoA desaturase. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of stearoyl-CoA desaturase in cells or tissues

comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being  
5 prone to a disease or condition associated with expression of stearoyl-CoA desaturase, by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

#### 10 DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding stearoyl-CoA desaturase, ultimately modulating the amount  
15 of stearoyl-CoA desaturase produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding stearoyl-CoA desaturase. As used herein, the terms "target nucleic acid" and "nucleic acid encoding stearoyl-  
20 CoA desaturase" encompass DNA encoding stearoyl-CoA desaturase, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of  
25 the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with  
30 include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and

catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of stearoyl-CoA desaturase. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

10 It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding stearoyl-CoA desaturase. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have

a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding stearyl-CoA desaturase, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between

the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective,

and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently  
5 complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick,  
10 Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds.

"Complementary," as used herein, refers to the capacity  
15 for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered  
20 to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus,  
25 "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that  
30 the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to

the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of *in vivo* assays or therapeutic treatment, and in the case of *in vitro* assays, under conditions in which the assays are performed.

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with other antisense compounds or therapeutics, can be used as tools in differential and/or

combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

Expression patterns within cells or tissues treated  
5 with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway,  
10 cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds which affect expression patterns.

15 Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, 2000, 480, 17-24; Celis, et al., *FEBS Lett.*, 2000, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., *Drug Discov. Today*, 2000, 5,  
20 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, 1999, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U. S. A.*, 2000, 97, 1976-81), protein arrays and proteomics (Celis, et  
25 al., *FEBS Lett.*, 2000, 480, 2-16; Jungblut, et al., *Electrophoresis*, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, 2000, 480, 2-16; Larsson, et al., *J. Biotechnol.*, 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al.,  
30 *Anal. Biochem.*, 2000, 286, 91-98; Larson, et al., *Cytometry*, 2000, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, 2000, 3, 316-21), comparative genomic

hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*,  
1998, 31, 286-96), FISH (fluorescent in situ  
hybridization) techniques (Going and Gusterson, *Eur. J.*  
*Cancer*, 1999, 35, 1895-904) and mass spectrometry methods  
5 (reviewed in (To, *Comb. Chem. High Throughput Screen*,  
2000, 3, 235-41).

The specificity and sensitivity of antisense is also  
harnessed by those of skill in the art for therapeutic  
uses. Antisense oligonucleotides have been employed as  
10 therapeutic moieties in the treatment of disease states in  
animals and man. Antisense oligonucleotide drugs,  
including ribozymes, have been safely and effectively  
administered to humans and numerous clinical trials are  
presently underway. It is thus established that  
15 oligonucleotides can be useful therapeutic modalities that  
can be configured to be useful in treatment regimes for  
treatment of cells, tissues and animals, especially  
humans.

In the context of this invention, the term  
20 "oligonucleotide" refers to an oligomer or polymer of  
ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or  
mimetics thereof. This term includes oligonucleotides  
composed of naturally-occurring nucleobases, sugars and  
covalent internucleoside (backbone) linkages as well as  
25 oligonucleotides having non-naturally-occurring portions  
which function similarly. Such modified or substituted  
oligonucleotides are often preferred over native forms  
because of desirable properties such as, for example,  
enhanced cellular uptake, enhanced affinity for nucleic  
30 acid target and increased stability in the presence of  
nucleases.

While antisense oligonucleotides are a preferred form  
of antisense compound, the present invention comprehends

other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50  
5 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide  
10 sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

As is known in the art, a nucleoside is a base-sugar  
15 combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar  
20 portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to  
25 form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are  
30 commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages

include, but are not limited to, U.S.: 3,687,808;  
4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897;  
5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131;  
5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677;  
5 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111;  
5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555;  
5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of  
which are commonly owned with this application, and each  
of which is herein incorporated by reference.

10 Preferred modified oligonucleotide backbones that do  
not include a phosphorus atom therein have backbones that  
are formed by short chain alkyl or cycloalkyl  
internucleoside linkages, mixed heteroatom and alkyl or  
cycloalkyl internucleoside linkages, or one or more short  
15 chain heteroatomic or heterocyclic internucleoside  
linkages. These include those having morpholino linkages  
(formed in part from the sugar portion of a nucleoside);  
siloxane backbones; sulfide, sulfoxide and sulfone  
backbones; formacetyl and thioformacetyl backbones;  
20 methylene formacetyl and thioformacetyl backbones;  
riboacetyl backbones; alkene containing backbones;  
sulfamate backbones; methyleneimino and methylenehydrazino  
backbones; sulfonate and sulfonamide backbones; amide  
backbones; and others having mixed N, O, S and CH<sub>2</sub>  
25 component parts.

Representative United States patents that teach the  
preparation of the above oligonucleosides include, but are  
not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444;  
5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564;  
30 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677;  
5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289;  
5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070;  
5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and

5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the  
5 sugar and the internucleoside linkage, i.e., the backbone,  
of the nucleotide units are replaced with novel groups.  
The base units are maintained for hybridization with an  
appropriate nucleic acid target compound. One such  
oligomeric compound, an oligonucleotide mimetic that has  
10 been shown to have excellent hybridization properties, is  
referred to as a peptide nucleic acid (PNA). In PNA  
compounds, the sugar-backbone of an oligonucleotide is  
replaced with an amide containing backbone, in particular  
an aminoethylglycine backbone. The nucleobases are  
15 retained and are bound directly or indirectly to aza  
nitrogen atoms of the amide portion of the backbone.  
Representative United States patents that teach the  
preparation of PNA compounds include, but are not limited  
to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of  
20 which is herein incorporated by reference. Further  
teaching of PNA compounds can be found in Nielsen et al.,  
*Science*, **1991**, 254, 1497-1500.

Most preferred embodiments of the invention are  
oligonucleotides with phosphorothioate backbones and  
25 oligonucleosides with heteroatom backbones, and in  
particular  $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$  [known as a  
methylene (methyylimino) or MMI backbone],  $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-$   
 $\text{CH}_2-$ ,  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$  and  $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$  [wherein  
the native phosphodiester backbone is represented as  $-\text{O}-\text{P}-$   
30  $\text{O}-\text{CH}_2-$ ] of the above referenced U.S. patent 5,489,677, and  
the amide backbones of the above referenced U.S. patent  
5,602,240. Also preferred are oligonucleotides having  
morpholino backbone structures of the above-referenced

U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, **1995**, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, also described in examples hereinbelow.

A further preferred modification includes Locked

Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene  $(-\text{CH}_2-)_n$  group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNA  
5 LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 2'-allyl (2'-CH<sub>2</sub>-CH=CH<sub>2</sub>), 2'-O-allyl (2'-O-CH<sub>2</sub>-CH=CH<sub>2</sub>) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T),

cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other  
5 alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ( $-C\equiv C-CH_3$ ) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo  
10 uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-  
15 methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-  
20 2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole  
25 cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further  
30 nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990,

those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B. ,  
5 ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyl-  
10 adenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278)  
15 and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified  
20 nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121,  
25 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent  
30 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which

enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers.

Typical conjugates groups include cholesterol, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion.

Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, *86*, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Let.*, **1994**, *4*, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, **1992**, *660*, 306-309; Manoharan et al., *Bioorg. Med. Chem. Let.*, **1993**, *3*, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, **1992**, *20*, 533-538), an aliphatic chain, e.g., dodecandiol

or undecyl residues (Saison-Behmoaras et al., *EMBO J.*,  
1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990,  
259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-  
54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or  
5 triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-  
phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995,  
36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18,  
3777-3783), a polyamine or a polyethylene glycol chain  
(Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14,  
10 969-973), or adamantane acetic acid (Manoharan et al.,  
*Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmityl moiety  
(Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-  
237), or an octadecylamine or hexylamino-carbonyl-  
oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp.*  
15 *Ther.*, 1996, 277, 923-937. Oligonucleotides of the  
invention may also be conjugated to active drug  
substances, for example, aspirin, warfarin, phenylbuta-  
zone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-  
pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodo-  
20 benzoic acid, flufenamic acid, folinic acid, a  
benzothiadiazide, chlorothiazide, a diazepine, indo-  
methicin, a barbiturate, a cephalosporin, a sulfa drug, an  
antidiabetic, an antibacterial or an antibiotic.  
Oligonucleotide-drug conjugates and their preparation are  
25 described in United States Patent Application 09/334,130  
(filed June 15, 1999) which is incorporated herein by  
reference in its entirety.

Representative United States patents that teach the  
preparation of such oligonucleotide conjugates include,  
30 but are not limited to, U.S.: 4,828,979; 4,948,882;  
5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538;  
5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124;  
5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439;

5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025;  
4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335;  
4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136;  
5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469;  
5 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098;  
5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475;  
5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142;  
5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923;  
5,599,928 and 5,688,941, certain of which are commonly  
10 owned with the instant application, and each of which is  
herein incorporated by reference.

It is not necessary for all positions in a given  
compound to be uniformly modified, and in fact more than  
one of the aforementioned modifications may be  
15 incorporated in a single compound or even at a single  
nucleoside within an oligonucleotide. The present  
invention also includes antisense compounds which are  
chimeric compounds. "Chimeric" antisense compounds or  
"chimeras," in the context of this invention, are  
20 antisense compounds, particularly oligonucleotides, which  
contain two or more chemically distinct regions, each made  
up of at least one monomer unit, i.e., a nucleotide in the  
case of an oligonucleotide compound. These  
oligonucleotides typically contain at least one region  
25 wherein the oligonucleotide is modified so as to confer  
upon the oligonucleotide increased resistance to nuclease  
degradation, increased cellular uptake, and/or increased  
binding affinity for the target nucleic acid. An  
additional region of the oligonucleotide may serve as a  
30 substrate for enzymes capable of cleaving RNA:DNA or  
RNA:RNA hybrids. By way of example, RNase H is a cellular  
endonuclease which cleaves the RNA strand of an RNA:DNA  
duplex. Activation of RNase H, therefore, results in

cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric  
5 oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in  
10 the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as  
15 described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.:  
5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878;  
20 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355;  
5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this  
25 invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art  
30 may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized *in vitro* and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the *in vivo* synthesis of  
5 antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted  
10 molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to,  
15 U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528;  
20 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon  
25 administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the  
30 invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an

active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE  
5 [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

The term "pharmaceutically acceptable salts" refers  
10 to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are  
15 formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are  
20 N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic  
25 compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional  
30 manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical

addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable

cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

5 For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts  
10 formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid,  
15 gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the  
20 like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an  
25 animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of stearoyl-CoA desaturase is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized  
30 in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may

also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds  
5 hybridize to nucleic acids encoding stearoyl-CoA desaturase, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding stearoyl-CoA desaturase can be detected by  
10 means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of stearoyl-CoA desaturase  
15 in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered  
20 in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or  
25 insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection  
30 or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1-10</sub> alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States

patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate,. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-

20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include

5 poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized

10 polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylamino-methylethylene

15 P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcynaoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-

20 dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in detail in United States applications 08/886,829 (filed

25 July 1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed February 23, 1999), 09/082,624 (filed May 21, 1998) and 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

Compositions and formulations for parenteral,

30 intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and

other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may  
5 be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit  
10 dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations  
15 are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be  
20 formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed  
25 media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

30 In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams,

jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to

the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199).

Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations.

Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgate, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl

gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

5           The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral  
10 delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245;  
15 Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are  
20 among the materials that have commonly been administered orally as o/w emulsions.

          In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be  
25 defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically  
30 microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a

transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active

5 molecules (Leung and Shah, *in: Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215).

Microemulsions commonly are prepared via a combination of three to five components that include oil, water,

10 surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the

15 surfactant molecules (Schott, *in Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a

20 comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, *in Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, *in Pharmaceutical Dosage Forms*, Lieberman, Rieger

25 and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed

30 spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers,

polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate  
5 (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into  
10 the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art.

15 The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355,  
20 Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

25 Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides  
30 (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from

enzymatic hydrolysis, possible enhancement of drug  
absorption due to surfactant-induced alterations in  
membrane fluidity and permeability, ease of preparation,  
ease of oral administration over solid dosage forms,  
5 improved clinical potency, and decreased toxicity  
(Constantinides et al., *Pharmaceutical Research*, 1994, 11,  
1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143).  
Often microemulsions may form spontaneously when their  
components are brought together at ambient temperature.  
10 This may be particularly advantageous when formulating  
thermolabile drugs, peptides or oligonucleotides.  
Microemulsions have also been effective in the transdermal  
delivery of active components in both cosmetic and  
pharmaceutical applications. It is expected that the  
15 microemulsion compositions and formulations of the present  
invention will facilitate the increased systemic  
absorption of oligonucleotides and nucleic acids from the  
gastrointestinal tract, as well as improve the local  
cellular uptake of oligonucleotides and nucleic acids  
20 within the gastrointestinal tract, vagina, buccal cavity  
and other areas of administration.

Microemulsions of the present invention may also  
contain additional components and additives such as  
sorbitan monostearate (Grill 3), Labrasol, and penetration  
25 enhancers to improve the properties of the formulation and  
to enhance the absorption of the oligonucleotides and  
nucleic acids of the present invention. Penetration  
enhancers used in the microemulsions of the present  
invention may be classified as belonging to one of five  
30 broad categories - surfactants, fatty acids, bile salts,  
chelating agents, and non-chelating non-surfactants (Lee  
et al., *Critical Reviews in Therapeutic Drug Carrier*

*Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

There are many organized surfactant structures  
5 besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the  
10 standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles  
15 which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as  
20 efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a  
25 suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and  
30 biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical*

*Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size  
5 and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the  
10 liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of  
15 extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the  
20 administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of  
25 liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper  
30 epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable

complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, **1987**, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, **1992**, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin

resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an  
5 additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral*  
10 *Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal  
15 formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin.  
20 Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized"  
25 liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized  
30 liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G<sub>M1</sub>, or (B) is derivatized with one or more hydrophilic polymers, such as

a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, **1987**, 223, 42; Wu et al., *Cancer Research*, **1993**, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, **1987**, 507, 64) reported the ability of monosialoganglioside  $G_{M1}$ , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, **1988**, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside  $G_{M1}$  or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-*sn*-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, **1980**, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, **1984**, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols

(e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov *et al.* (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume *et al.* (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle *et al.* (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin *et al.* (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin *et al.*) and in WO 94/20073 (Zalipsky *et al.*) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi *et al.*). U.S. Patent Nos. 5,540,935 (Miyazaki *et al.*) and 5,556,948 (Tagawa *et al.*) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry *et al.* discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa *et al.* discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman *et al.*

describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love *et al.* discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

5           Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to  
10 penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, *e.g.* they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without  
15 fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated  
20 delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

          Surfactants find wide application in formulations such as emulsions (including microemulsions) and  
25 liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the  
30 most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values.

5 In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated  
10 esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

15 If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl  
20 sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

25 If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of  
30 this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include

acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in  
5 *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient  
10 delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even  
15 non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic  
20 drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in*  
25 *Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are  
30 chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of

oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and diglycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic

derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucholic acid (sodium glucholate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate

(EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacycloalkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol,

pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

#### Carriers

Certain compositions of the present invention also  
5 incorporate carrier compounds in the formulation. As used  
herein, "carrier compound" or "carrier" can refer to a  
nucleic acid, or analog thereof, which is inert (*i.e.*,  
does not possess biological activity *per se*) but is  
recognized as a nucleic acid by *in vivo* processes that  
10 reduce the bioavailability of a nucleic acid having  
biological activity by, for example, degrading the  
biologically active nucleic acid or promoting its removal  
from circulation. The coadministration of a nucleic acid  
and a carrier compound, typically with an excess of the  
15 latter substance, can result in a substantial reduction of  
the amount of nucleic acid recovered in the liver, kidney  
or other extracirculatory reservoirs, presumably due to  
competition between the carrier compound and the nucleic  
acid for a common receptor. For example, the recovery of  
20 a partially phosphorothioate oligonucleotide in hepatic  
tissue can be reduced when it is coadministered with  
polyinosinic acid, dextran sulfate, polycytidic acid or 4-  
acetamido-4'isothiocyano-stilbene-2,2'-disulfonic acid  
(Miyao *et al.*, *Antisense Res. Dev.*, 1995, 5, 115-121;  
25 Takakura *et al.*, *Antisense & Nucl. Acid Drug Dev.*, 1996,  
6, 177-183).

#### Excipients

In contrast to a carrier compound, a "pharmaceutical  
carrier" or "excipient" is a pharmaceutically acceptable  
30 solvent, suspending agent or any other pharmacologically  
inert vehicle for delivering one or more nucleic acids to  
an animal. The excipient may be liquid or solid and is  
selected, with the planned manner of administration in

mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, 5 binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium 10 hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch 15 glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be 20 used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, 25 hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid 30 or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration

which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, 5 magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Other Components

The compositions of the present invention may 10 additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, 15 antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, 20 opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary 25 agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

30 Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not

limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on  $EC_{50}$ s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even

once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following  
5 successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once  
10 or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the  
15 same.

#### **EXAMPLES**

##### **Example 1**

##### **Nucleoside Phosphoramidites for Oligonucleotide Synthesis Deoxy and 2'-alkoxy amidites**

20 2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent  
25 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

30 Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially

available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

### **2'-Fluoro amidites**

#### **2'-Fluorodeoxyadenosine amidites**

5        2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was  
10 synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-  
15 arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-  
20 dimethoxytrityl- (DMT) and 5'-DMT-3'-phosphoramidite intermediates.

#### **2'-Fluorodeoxyguanosine**

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS)  
25 protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected  
30 arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and

5'-DMT-3'-phosphoramidites.

**2'-Fluorouridine**

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

**2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

**2'-O-(2-Methoxyethyl) modified amidites**

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

**2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-**

**methyluridine]**

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL).

The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was

decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

10            **2'-O-Methoxyethyl-5-methyluridine**

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

30            2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of

dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional  
5 one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL). The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500  
10 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were  
15 evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

20 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours.  
25 The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and extracted with 2x200 mL of saturated  
30 sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product).

The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

5           **3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set  
10 aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting  
15 mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The  
20 residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title  
25 compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141  
30 M) in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and

transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to  
5 dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

10 **N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the  
15 reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a  
20 residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

25 **N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxytetra(isopropyl)phosphite (40.5 mL, 0.123 M) were added  
30 with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and

saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column  
5 using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

**2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites**

10 **2'-(Dimethylaminooxyethoxy) nucleoside amidites**

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine  
15 nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

20 **5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine**

O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry  
25 pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22,  
30 ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine

(1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to  
5 -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

10 **5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first  
15 until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160  
20 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional  
25 side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution  
30 can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate

fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine**

5'-O-tert-Butylidiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

**5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine**

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h

the mixture was filtered, the filtrate was washed with ice cold  $\text{CH}_2\text{Cl}_2$  and the combined organic phase was washed with water, brine and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinooxy) ethyl]-5-methyluridine as white foam (1.95 g, 78%).

**5'-O-tert-Butylidiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine**

5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in  $\text{CH}_2\text{Cl}_2$ ). Aqueous  $\text{NaHCO}_3$  solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes,

the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was  
5 dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness . The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam  
10 (14.6g, 80%).

**2'-O-(dimethylaminoxyethyl)-5-methyluridine**

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF  
15 was then added to 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and  
20 eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

**5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine**

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight  
25 at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction  
30 mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to get

5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine  
(1.13g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-  
methyluridine-3'-[(2-cyanoethyl)-N,N-  
5 diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine  
(1.08g, 1.67mmol) was co-evaporated with toluene (20mL).  
To the residue N,N-diisopropylamine tetrazonide (0.29g,  
1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum  
10 overnight at 40°C. Then the reaction mixture was dissolved  
in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-  
N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol)  
was added. The reaction mixture was stirred at ambient  
temperature for 4 hrs under inert atmosphere. The  
15 progress of the reaction was monitored by TLC  
(hexane:ethyl acetate 1:1). The solvent was evaporated,  
then the residue was dissolved in ethyl acetate (70mL) and  
washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl acetate layer  
was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. Residue  
20 obtained was chromatographed (ethyl acetate as eluent) to  
get 5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-  
methyluridine-3'-[(2-cyanoethyl)-N,N-  
diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

**2'-(Aminoxyethoxy) nucleoside amidites**

25 2'-(Aminoxyethoxy) nucleoside amidites [also known  
in the art as 2'-O-(aminoxyethyl) nucleoside amidites]  
are prepared as described in the following paragraphs.  
Adenosine, cytidine and thymidine nucleoside amidites are  
prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

The 2'-O-aminoxyethyl guanosine analog may be  
5 obtained by selective 2'-O-alkylation of diaminopurine  
riboside. Multigram quantities of diaminopurine riboside  
may be purchased from Schering AG (Berlin) to provide 2'-  
O-(2-ethylacetyl) diaminopurine riboside along with a  
minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl)  
10 diaminopurine riboside may be resolved and converted to  
2'-O-(2-ethylacetyl)guanosine by treatment with adenosine  
deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J.,  
WO 94/02501 A1 940203.) Standard protection procedures  
should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-  
15 dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-  
diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-  
dimethoxytrityl)guanosine which may be reduced to provide  
2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-  
hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As  
20 before the hydroxyl group may be displaced by N-  
hydroxyphthalimide via a Mitsunobu reaction, and the  
protected nucleoside may phosphitylated as usual to yield  
2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-([2-  
phthalmidoxy]ethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-  
25 3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside  
amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites  
(also known in the art as 2'-O-dimethylaminoethoxyethyl,  
30 i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside  
amidites) are prepared as follows. Other nucleoside  
amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2 [2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-,2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH<sub>2</sub>Cl<sub>2</sub>:Et<sub>3</sub>N (20:1, v/v, with 1% triethylamine) gives the title compound.

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

Diisopropylaminotetrazolide (0.6 g) and 2-  
5 cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and  
10 the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

**Example 2**

**Oligonucleotide synthesis**

15 Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the  
20 phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the  
25 capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphate oligonucleotides are prepared  
30 as described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by

reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

5 Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902  
10 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

15 Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both  
20 herein incorporated by reference.

### **Example 3**

#### **Oligonucleoside Synthesis**

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedi-  
25 methylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligo-  
30 nucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of

which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

5 Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

#### **Example 4**

##### **PNA Synthesis**

10 Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance  
15 with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

#### **Example 5**

##### **Synthesis of Chimeric Oligonucleotides**

Chimeric oligonucleotides, oligonucleosides or mixed  
20 oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap"  
25 segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

30 **[2'-O-Me] -- [2'-deoxy] -- [2'-O-Me]**

##### **Chimeric Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligo-

nucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphor-  
5 amidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl.  
10 The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect  
15 all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being  
20 desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

**[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-**  
25 **(Methoxyethyl)] Chimeric Phosphorothioate**  
**Oligonucleotides**

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxy-ethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl  
30 chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

**[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy  
Phosphorothioate]--[2'-O-(2-Methoxyethyl)  
Phosphodiester] Chimeric Oligonucleotides**

[2'-O-(2-methoxyethyl phosphodiester)--[2'-deoxy  
5 phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester]  
chimeric oligonucleotides are prepared as per the above  
procedure for the 2'-O-methyl chimeric oligonucleotide  
with the substitution of 2'-O-(methoxyethyl) amidites for  
the 2'-O-methyl amidites, oxidization with iodine to  
10 generate the phosphodiester internucleotide linkages  
within the wing portions of the chimeric structures and  
sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1  
dioxide (Beaucage Reagent) to generate the  
phosphorothioate internucleotide linkages for the center  
15 gap.

Other chimeric oligonucleotides, chimeric oligo-  
nucleosides and mixed chimeric oligonucleotides/oligo-  
nucleosides are synthesized according to United States  
patent 5,623,065, herein incorporated by reference.

20 **Example 6**

**Oligonucleotide Isolation**

After cleavage from the controlled pore glass column  
(Applied Biosystems) and deblocking in concentrated  
ammonium hydroxide at 55°C for 18 hours, the  
25 oligonucleotides or oligonucleosides are purified by  
precipitation twice out of 0.5 M NaCl with 2.5 volumes  
ethanol. Synthesized oligonucleotides were analyzed by  
polyacrylamide gel electrophoresis on denaturing gels and  
judged to be at least 85% full length material. The  
30 relative amounts of phosphorothioate and phosphodiester  
linkages obtained in synthesis were periodically checked  
by <sup>31</sup>P nuclear magnetic resonance spectroscopy, and for  
some studies oligonucleotides were purified by HPLC, as

described by Chiang et al., *J. Biol. Chem.* **1991**, *266*, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

5 **Example 7**

**Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences  
10 simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3, H-1,2 benzodithiole-3-one 1,1 dioxide  
15 (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per  
20 known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature  
25 (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

30 **Example 8**

**Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption

spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

#### **Example 9**

##### **Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 7 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

##### **25 T-24 cells:**

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life

Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well  
5 for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

10 A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg,  
15 MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached  
20 90% confluence.

NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium  
25 (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

HEK cells:

Human embryonic keratinocytes (HEK) were obtained  
30 from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely

maintained for up to 10 passages as recommended by the supplier.

HepG2 cells:

The human hepatoblastoma cell line HepG2 was obtained  
5 from the American Type Culture Collection (Manassas, VA).  
HepG2 cells were routinely cultured in Eagle's MEM  
supplemented with 10% fetal calf serum, non-essential  
amino acids, and 1 mM sodium pyruvate (Gibco/Life  
Technologies, Gaithersburg, MD). Cells were routinely  
10 passaged by trypsinization and dilution when they reached  
90% confluence. Cells were seeded into 96-well plates  
(Falcon-Primaria #3872) at a density of 7000 cells/well  
for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be  
15 seeded onto 100 mm or other standard tissue culture plates  
and treated similarly, using appropriate volumes of medium  
and oligonucleotide.

AML12 cells:

AML12 (alpha mouse liver 12) cell line was  
20 established from hepatocytes from a mouse (CD1 strain,  
line MT42) transgenic for human TGF alpha. Cells are  
cultured in a 1:1 mixture of Dulbecco's modified Eagle's  
medium and Ham's F12 medium with 0.005 mg/ml insulin,  
0.005 mg/ml transferrin, 5 ng/ml selenium, and 40 ng/ml  
25 dexamethasone, and 90%; 10% fetal bovine serum. For  
subculturing, spent medium is removed and fresh media of  
0.25% trypsin, 0.03% EDTA solution is added. Fresh trypsin  
solution (1 to 2 ml) is added and the culture is left to  
sit at room temperature until the cells detach.

30 Cells were routinely passaged by trypsinization and  
dilution when they reached 90% confluence. Cells were  
seeded into 96-well plates (Falcon-Primaria #3872) at a  
density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

5 Primary mouse hepatocytes:

Primary mouse hepatocytes were prepared from CD-1 mice purchased from Charles River Labs (Wilmington, MA) and were routinely cultured in Hepatocyte Attachment Media (Gibco) supplemented with 10% Fetal Bovine Serum (Gibco/Life Technologies, Gaithersburg, MD), 250nM dexamethasone (Sigma), and 10nM bovine insulin (Sigma). Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 10000 cells/well for use in RT-PCR analysis.

15 For Northern blotting or other analyses, cells are plated onto 100 mm or other standard tissue culture plates coated with rat tail collagen (200ug/mL) (Becton Dickinson) and treated similarly using appropriate volumes of medium and oligonucleotide.

20 Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM<sup>TM</sup>-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM<sup>TM</sup>-1 containing 3.75  $\mu$ g/mL LIPOFECTIN<sup>TM</sup> (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

30 The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control

oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCCAAGGA**, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

#### **Example 10**

#### **Analysis of oligonucleotide inhibition of stearoyl-CoA desaturase expression**

Antisense modulation of stearoyl-CoA desaturase expression can be assayed in a variety of ways known in the art. For example, stearoyl-CoA desaturase mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation

are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions.

Protein levels of stearyl-CoA desaturase can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to stearyl-CoA desaturase can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current*

*Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al.,

5 *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

#### **Example 11**

##### **Poly(A)+ mRNA isolation**

Poly(A)+ mRNA was isolated according to Miura et al.,

10 *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates,

15 growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room

20 temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl).

25 After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate

30 was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all

solutions.

### Example 12

#### Total RNA Isolation

Total RNA was isolated using an RNEASY 96™ kit and  
5 buffers purchased from Qiagen Inc. (Valencia CA) following  
the manufacturer's recommended procedures. Briefly, for  
cells grown on 96-well plates, growth medium was removed  
from the cells and each well was washed with 200 µL cold  
PBS. 100 µL Buffer RLT was added to each well and the  
10 plate vigorously agitated for 20 seconds. 100 µL of 70%  
ethanol was then added to each well and the contents mixed  
by pipetting three times up and down. The samples were  
then transferred to the RNEASY 96™ well plate attached to  
a QIAVAC™ manifold fitted with a waste collection tray and  
15 attached to a vacuum source. Vacuum was applied for 15  
seconds. 1 mL of Buffer RW1 was added to each well of the  
RNEASY 96™ plate and the vacuum again applied for 15  
seconds. 1 mL of Buffer RPE was then added to each well  
of the RNEASY 96™ plate and the vacuum applied for a  
20 period of 15 seconds. The Buffer RPE wash was then  
repeated and the vacuum was applied for an additional 10  
minutes. The plate was then removed from the QIAVAC™  
manifold and blotted dry on paper towels. The plate was  
then re-attached to the QIAVAC™ manifold fitted with a  
25 collection tube rack containing 1.2 mL collection tubes.  
RNA was then eluted by pipetting 60 µL water into each  
well, incubating 1 minute, and then applying the vacuum  
for 30 seconds. The elution step was repeated with an  
additional 60 µL water.

30 The repetitive pipetting and elution steps may be  
automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc.,

Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

5 **Example 13**

**Real-time Quantitative PCR Analysis of stearyl-CoA desaturase mRNA Levels**

Quantitation of stearyl-CoA desaturase mRNA levels was determined by real-time quantitative PCR using the ABI  
10 PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products  
15 in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe  
20 that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of  
25 the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3'  
30 quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR

amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With  
5 each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions  
10 containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

Prior to quantitative PCR analysis, primer-probe sets  
15 specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA  
20 isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA  
25 signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the  
30 single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

PCR reagents were obtained from PE-Applied

Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN<sup>TM</sup> buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse  
5 primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD<sup>TM</sup>, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25  $\mu$ L total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C  
10 to activate the AMPLITAQ GOLD<sup>TM</sup>, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR  
15 are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreen<sup>TM</sup> (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing,  
20 or separately. Total RNA is quantified using RiboGreen<sup>TM</sup> RNA quantification reagent from Molecular Probes. Methods of RNA quantification by RiboGreen<sup>TM</sup> are taught in Jones, L.J., et al, *Analytical Biochemistry*, 1998, 265, 368-374.

In this assay, 175  $\mu$ L of RiboGreen<sup>TM</sup> working reagent  
25 (RiboGreen<sup>TM</sup> reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25uL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

30 Probes and primers to human stearoyl-CoA desaturase were designed to hybridize to a human stearoyl-CoA desaturase sequence, using published sequence information

(GenBank accession number AF)97514, incorporated herein as SEQ ID NO: 3). For human stearyl-CoA desaturase the PCR primers were:

forward primer: GATCCCGGCATCCGAGA (SEQ ID NO: 4)

5 reverse primer: GGTATAGGAGCTAGAGATATCGTCCTG (SEQ ID NO: 5)

and the PCR probe was: FAM-CCAAGATGCCGGCCCACTTGC-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

10 For human GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 7)

reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 8) and

the PCR probe was: 5' JOE-CAAGCTTCCCGTTCTCAGCC-TAMRA 3' (SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster

15 City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

#### **Example 14**

#### **Northern blot analysis of stearyl-CoA desaturase mRNA levels**

20 Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was

25 fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer

30 using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400

(Stratagene, Inc, La Jolla, CA) and then robed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions.

5 To detect human stearoyl-CoA desaturase, a human stearoyl-CoA desaturase specific probe was prepared by PCR using the forward primer GATCCCGGCATCCGAGA (SEQ ID NO: 4) and the reverse primer GGTATAGGAGCTAGAGATATCGTCCTG (SEQ ID NO: 5). To normalize for variations in loading and  
10 transfer efficiency membranes were stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3  
15 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

#### **Example 15**

**Antisense inhibition of human stearoyl-CoA desaturase  
expression by chimeric phosphorothioate oligonucleotides  
20 having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human stearoyl-CoA desaturase RNA, using published sequence (GenBank accession number AF097514, incorporated  
25 herein as SEQ ID NO: 3). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides  
30 in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides.

The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human

5 stearyl-CoA desaturase mRNA levels in HepG2 cells by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1

10 Inhibition of human stearyl-CoA desaturase mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
147899	5'UTR	3	9	GTCCGGTATTTCTCAGCCC	N.D.	10
147900	5'UTR	3	72	CCGCGGTGCGTGGAGGTCCC	N.D.	11
147901	5'UTR	3	121	TACGCGCTGAGCCGCGCGC	N.D.	12
147902	5'UTR	3	141	GCGGTTTCGAAGCCCGCCGG	N.D.	13
147903	Coding	3	311	CCTCCATTCTGCAGGACCCT	N.D.	14
147904	Coding	3	471	TCCCAAGTGTAGCAGAGACA	N.D.	15
147905	Coding	3	571	CTCCTGCTGTTATGCCCAGG	N.D.	16
147906	Coding	3	691	CACGGTGGTCACGAGCCCAT	N.D.	17
147907	Coding	3	771	CAGCCAACCCACGTGAGAGA	22	18
147908	Coding	3	824	GACAAGTCTAGCGTACTCCC	49	19
147909	Coding	3	1011	GTTCACCAGCCAGGTGGCAT	10	20
147910	Coding	3	1111	TGTGGAAGCCCTCACCCACA	0	21
147911	Coding	3	1171	AGTTGATGTGCCAGCGGTAC	6	22
147912	Stop Codon	3	1307	GGACCCCAAACCTCAGCCACT	25	23
147913	3'UTR	3	1581	TGCCTGGGAGGCAATAAGGG	8	24
147914	3'UTR	3	1861	ATACATGCTAACTCTCTCCC	10	25
147915	3'UTR	3	1941	AAGTCCTCATTAGGTAGGCA	37	26
147916	3'UTR	3	2241	TGTAATGAGCAGCTCATGGA	0	27
147917	3'UTR	3	2616	TCAGTAACCTTCTCAAGCCC	0	28
147918	3'UTR	3	2980	GGAGCTGCCTGGACAGCAAG	16	29
147919	3'UTR	3	3011	TCAGTGACCCTGAGCATTCT	30	30
147920	3'UTR	3	3231	TGGCTGGCCCACTGGCTCAA	11	31
147921	3'UTR	3	3291	GCATGCCCTCTGGTTCTGAC	7	32
147922	3'UTR	3	3471	GCTTTGCAGTTCACCCTGAC	23	33
147923	3'UTR	3	3502	GTGGTATCTCAAATCCCAGG	0	34
147924	3'UTR	3	3791	TAGTCCAGGCTAACCCTGT	0	35
147925	3'UTR	3	3851	GTGATCTTCCCTTAGATCCT	0	36
147926	3'UTR	3	4101	CTCAGCAGACACACTCCCAC	3	37
147927	3'UTR	3	4226	GCTAAGTTGTCAGCACACCC	0	38
147928	3'UTR	3	4406	AAGTTTCAGAATGAAGCCC	25	39
147929	3'UTR	3	4571	AGAGAATACACCCAAGATAC	0	40

147930	3'UTR	3	4708	TAGTTAAGTGACTTGCCAG	0	41
147931	3'UTR	3	4771	GCCCTTTGAGGTAGGTCAGT	4	42
147932	3'UTR	3	4921	CCATATAGACTAATGACAGC	10	43
147933	3'UTR	3	5021	CTGTATGTTTCCGTGGCAAT	11	44
168231	5'UTR	3	101	CTTGCACGCTAGCTGGTTGT	N.D.	45
168232	Coding	3	331	GCATCGTCTCCAACCTTATCT	N.D.	46
168233	Coding	3	451	TAAGGATGATGTTTCTCCAG	N.D.	47
168234	Coding	3	526	CCCAAAGCCAGGTGTAGAAC	N.D.	48
168235	Coding	3	601	TGTAAGAGCGGTGGCTCCAC	N.D.	49
168236	Coding	3	661	CATTCTGGAATGCCATTGTG	N.D.	50
168237	Coding	3	731	TTATGAGGATCAGCATGTGT	N.D.	51
168238	Coding	3	861	CCTCTGGAACATCACCAGTT	N.D.	52
168239	Coding	3	901	GGATGAAGCACATCAGCAGC	N.D.	53
168240	Coding	3	936	TTCACCCAGAAATACCAGG	N.D.	54
168241	Coding	3	1082	GAAACCAGGATATTCTCCCG	N.D.	55
168242	Coding	3	1151	TCACTGGCAGAGTAGTCATA	N.D.	56
168243	Coding	3	1261	TAATCCTGGCCAAGATGGCG	N.D.	57
168244	3'UTR	3	1401	TCATCATCTTTAGCATCCTG	N.D.	58
168245	3'UTR	3	1601	GCAAAGACTGACCAGCTGCT	N.D.	59
168246	3'UTR	3	1748	GACTACCCAGAAGATTCTGT	N.D.	60
168247	3'UTR	3	1881	CTTCCCTCATCCTTACATTC	N.D.	61
168248	3'UTR	3	1985	CCCAGCCAGGAGAGAAAGG	N.D.	62
168249	3'UTR	3	2102	CTTCCCAGCAGAGACCACT	N.D.	63
168250	3'UTR	3	2281	CCAATATCCTGAAGATGGCA	N.D.	64
168251	3'UTR	3	2481	CCCAACTAATTCCTCCTCTC	N.D.	65
168252	3'UTR	3	2541	TATAGATCCTGTCCCTCAGC	N.D.	66
168253	3'UTR	3	2631	CTCCCAATAACTCACTCAGT	N.D.	67
168254	3'UTR	3	2826	AAGAGATTCCTAACCCTGCC	N.D.	68
168255	3'UTR	3	2941	CACACAAAGGAGGCTGCCTG	N.D.	69
168256	3'UTR	3	3051	AAGTGGCAGCTAGCTCTACT	N.D.	70
168257	3'UTR	3	3321	CACCCTACCAAGTAAGCAG	N.D.	71
168258	3'UTR	3	3401	TGCTTCTTCCCAGTGAGAAC	N.D.	72
168259	3'UTR	3	3941	ATCAAGCAGGCATCTGATGA	N.D.	73
168260	3'UTR	3	4052	CCCTCAGCCTGAGGTGCCAT	N.D.	74
168261	3'UTR	3	4357	ATAATCCTCCACTCAGGCC	N.D.	75
168262	3'UTR	3	4431	CACTTAAGAAAAGCAGCCCT	N.D.	76
168263	3'UTR	3	4681	CAGCAAGTCAGTGGCACAGT	N.D.	77
168264	3'UTR	3	4971	GGCTAGTTATCCACCGCTTC	N.D.	78
168265	3'UTR	3	5044	CCCAATCACAGAAAAGGCAT	N.D.	79
168266	3'UTR	3	5081	AACTACTATATCCACATAA	N.D.	80

As shown in Table 1, SEQ ID NOs 18, 19, 20, 23, 25, 26, 29, 30, 31, 33, 39, 43 and 44 demonstrated at least 10% inhibition of human stearoyl-CoA desaturase expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

**Example 16****Western blot analysis of stearyl-CoA desaturase protein levels**

Western blot analysis (immunoblot analysis) is  
5 carried out using standard methods. Cells are harvested  
16-20 h after oligonucleotide treatment, washed once with  
PBS, suspended in Laemmli buffer (100 ul/well), boiled for  
5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run  
for 1.5 hours at 150 V, and transferred to membrane for  
10 western blotting. Appropriate primary antibody directed  
to stearyl-CoA desaturase is used, with a radiolabelled  
or fluorescently labeled secondary antibody directed  
against the primary antibody species. Bands are  
visualized using a PHOSPHORIMAGER™ (Molecular Dynamics,  
15 Sunnyvale CA).

**What is claimed is:**

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding human  
5 stearoyl-CoA desaturase, wherein said compound specifically hybridizes with a nucleic acid molecule encoding human stearoyl-CoA desaturase and inhibits the expression of human stearoyl-CoA desaturase.
2. The compound of claim 1 which is an antisense  
10 oligonucleotide.
3. The compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 10,  
11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 25, 26, 29,  
30, 31, 33, 39, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52,  
15 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66,  
67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80.
4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.
- 20 5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.
6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.
- 25 7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.
8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.
- 30 9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.
10. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding human stearoyl-CoA desaturase.

5 12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

10 14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15 15. A method of inhibiting the expression of human stearoyl-CoA desaturase in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of human stearoyl-CoA desaturase is inhibited.

20 16. A method of treating a human having a disease or condition associated with human stearoyl-CoA desaturase comprising administering to said human a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of human stearoyl-CoA desaturase is inhibited.

17. The method of claim 16 wherein the condition involves abnormal lipid metabolism.

25 18. The method of claim 16 wherein the condition involves abnormal cholesterol metabolism.

19. The method of claim 16 wherein the condition is atherosclerosis.

20. The method of claim 16 wherein the disease is cardiovascular disease.

SEQUENCE LISTING

<110> Isis Pharmaceuticals, Inc.  
 Rosanne M. Crooke  
 Mark J. Graham

<120> ANTISENSE MODULATION OF STEAROYL-COA DESATURASE  
 EXPRESSION

<130> ISPH-0695

<150> 09/918,187

<151> 2001-07-30

<160> 80

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctctcaggg 20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2

atgcattctg cccccaagga 20

<210> 3

<211> 5221

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (236)...(1315)

<400> 3

ataaaagggg gctgaggaaa taccggacac ggtcaccogt tgccagctct agcctttaa 60

ttcccggctc ggggacctcc acgcaccgcg gctagcgccg acaaccagct agcgtgcaag 120

gcgccgcggc tcagcgcgta ccggcgggct tcgaaaccgc agtctctcgg cgaccccgaa 180

ctccgctccg gacctcagc cccctggaaa gtgatcccgg catccgagag ccaag atg 238

Met

1



ttt caa aac agt gtg ttc gtt gcc act ttc ttg cga tat gct gtg gtg	1006
Phe Gln Asn Ser Val Phe Val Ala Thr Phe Leu Arg Tyr Ala Val Val	
245 250 255	
ctt aat gcc acc tgg ctg gtg aac agt gct gcc cac ctc ttc gga tat	1054
Leu Asn Ala Thr Trp Leu Val Asn Ser Ala Ala His Leu Phe Gly Tyr	
260 265 270	
cgf cct tat gac aag aac att agc ccc cgg gag aat atc ctg gtt tca	1102
Arg Pro Tyr Asp Lys Asn Ile Ser Pro Arg Glu Asn Ile Leu Val Ser	
275 280 285	
ctt gga gct gtg ggt gag ggc ttc cac aac tac cac cac tcc ttt ccc	1150
Leu Gly Ala Val Gly Glu Gly Phe His Asn Tyr His His Ser Phe Pro	
290 295 300 305	
tat gac tac tct gcc agt gag tac cgc tgg cac atc aac ttc acc aca	1198
Tyr Asp Tyr Ser Ala Ser Glu Tyr Arg Trp His Ile Asn Phe Thr Thr	
310 315 320	
ttc ttc att gat tgc atg gcc gcc ctc ggt ctg gcc tat gac cgg aag	1246
Phe Phe Ile Asp Cys Met Ala Ala Leu Gly Leu Ala Tyr Asp Arg Lys	
325 330 335	
aaa gtc tcc aag gcc gcc atc ttg gcc agg att aaa aga acc gga gat	1294
Lys Val Ser Lys Ala Ala Ile Leu Ala Arg Ile Lys Arg Thr Gly Asp	
340 345 350	
gga aac tac aag agt ggc tga gtttgggggtc cctcagggttt cctttttcaa	1345
Gly Asn Tyr Lys Ser Gly	
355	
aaaccagcca ggcagaggtt ttaatgtctg tttattaact actgaataat gctaccagga	1405
tgctaaagat gatgatgta acccattcca gtacagtatt cttttaaaat tcaaaagtat	1465
tgaagccaa caactctgcc tttatgatgc taagctgata ttatttcttc tcttatcctc	1525
tctctcttct aggccattg tcctccttt cactttattg ctatcgccct cctttccctt	1585
attgcctccc aggcaagcag ctggtcagtc tttgctcagt gtccagcttc caaagcctag	1645
acaacctttc tgtagcctaa aacgaatggc ctttgcctca gataactctc tttccttgag	1705
ctgtttgtgag ctttgaagta ggtggcttga gctagagata aaacagaatc ttctgggtag	1765
tcccctggtg attatcttca gcccaggctt ttgctagatg gaatggaaaa gcaacttcat	1825
ttgacacaaa gcttctaaag caggtaaatt gtcgggggag agagttagca tgtatgaatg	1885
taaggatgag ggaagcgaag caagaggaac ctctcgccat gatcagacat acagctgctt	1945
acctaatgag gacttcaagc cccaccacat agcatgcttc ctttctctcc tggctcgggg	2005
taaaaagtggt ctgctgggtgt tggcaatgct aattcaatgc cgcaacatat agttgaggcc	2065
gaggataaag aaaagacatt ttaagtttgt agtaaaagtg gtctctgctg ggggaagggtt	2125
ttcttttctt tttttcttta ataacaagga gatttcttag ttcatatatc aagaagtctt	2185
gaagttgggt gtttccagaa ttggtaaaaa cagcagctca tggaaatttg agtattccat	2245
gagctgctca ttacagttct ttctctttc tgctctgcca tcttcaggat attggttctt	2305
cccctcatag taataagatg gctgtggcat ttccaaacat ccaaaaaaag ggaaggattt	2365
aaggaggtga agtcgggtca aaaataaaat atatatacat atatacattg cttagaacgt	2425
taaaactatta gagtatttcc cttccaaaaga gggatgtttg gaaaaaactc tgaaggagag	2485
gagggaattag ttgggatgcc aatttctctt ccactgctgg acatgagatg gagaggctga	2545
gggacaggat ctataggcag cttctaagag cgaacttcac ataggaaggg atctgagaac	2605
acgttgccag gggcttgaga aggttactga gtgagttatt gggagtctta ataaaaataa	2665
ctagatatta ggtccattca ttaattagtt ccagtttctc cttgaaatga gtaaaaaacta	2725
gaaggcttct ctccacagtg ttgtgcccct tcaactcattt ttttttgagg agaaggggggt	2785
ctctgttaac atctagccta aagtatacaa ctgcctgggg ggcagggtta ggaatctctt	2845

```

cactaccctg attccttgatt cctggctcta ccctgtctgt cccttttctt tgaccagatc 2905
tttctcttcc ctgaaacgttt tcttctttcc ctggacagggc agcctccttt gtgtgtattc 2965
agaggcagtg atgacttgct gtccagggcag ctccctcctg cacacagaat gctcagggtc 3025
actgaaccac tgcttctctt ttgaaagtag agctagctgc cactttcacg tggcctccgc 3085
agtgtctcca cctacacccc tgtgctcccc tgccacactg atggctcaag acaaggctgg 3145
caaaccctcc cagaaacatc tctggcccag aaagcctctc tctccctccc tctctcatga 3205
ggcacagcca agccaagcgc tcatgttgag ccagtggggc agccacagag caaaagaggg 3265
tttattttca gtcccctctc tctgggtcag aaccagaggg catgctgaat gccccctgct 3325
tacttgggtga gggcgccccg cctgagtcag tgetctcagc tggcagtgca atgctttag 3385
aagtaggagg aaacagttct cactgggaag aagcaagggc aagaacccaa gtgcctcacc 3445
tcgaaaggag gccctgttcc ctggagtcag ggtgaactgc aaagctttgg ctgagacctg 3505
ggatttgaga taccacaaac cctgctgaac acagtgtctg ttcagcaaac taaccagcat 3565
tccctacagc ctagggcaga caatagtata gaagtctgga aaaaaacaaa aacagaattt 3625
gagaaccttg gacctcctc gtccctgtag ctcaactcagc aaagcagaag tctggctttg 3685
ctctattaag attggaaatg tacactacca aacactcagt ccactgttga gccccagtgc 3745
tggaaggagg gaagcgcttt cttctgtggt aattgcgtag aggctacagg ggttagctg 3805
gactaaaggc atcctgtctt tttgagctat tcacctcagt agaaaaggat ctaaggggag 3865
atcactgtag tttagttctg ttgacctgtg cacctacccc ttggaaatgt ctgctgggat 3925
ttctaattcc acaggtcatc agatgcctgc ttgataatat ataaacaata aaaacaactt 3985
tcacttcttc ctattgtaat cgtgtgccat ggatctgatc tgtacatga ccctacataa 4045
ggctggatgg cacctcaggc tgagggcccc aatgtatgtg tggctgtggg tgtgggtggg 4105
agtgtgtctg ctgagtaagg aacacgattt tcaagattct aaagctcaat tcaagtgaca 4165
cattaatgat aaactcagat ctgatcaaga gtcocggattt ctaacagtcc ctgctttggg 4225
gggtgtgctg acaacttagc tcaggtgcct tacatctttt ctaatcacag tgttgcatat 4285
gagcctgccc tcactccctc tgcagaatcc ctttgcacct gagaccctac tgaagtggct 4345
ggtagaaaaa ggggcctgag tggaggatta tcagtatcac gatttgcagg attcccttct 4405
gggcttcatt ctggaaactt ttgttagggc tgcttttctt aagtgccac atttgatgga 4465
gggtggaat aatttgaatg tatttgattt ataagttttt tttttttttt gggtaaaaag 4525
atggtttagt catttaaaat ggaaaatfff ctccctgggt tgctagtatc ttgggtgtat 4585
tctctgtaag tgtagtcaa ataggtcatc atgaaagggt aaaaaagcga ggtggccatg 4645
ttatgctggg ggttaaggcc agggcctctc caaccactgt gccactgact tgctgtgtga 4705
ccctgggcaa gtcaactaac tataagggtc ctcagttttc cttctgttaa aatggggata 4765
ataaactga cctacctcaa agggcagttt tgaggcatga ctaatgcttt ttagaaagca 4825
ttttgggatc cttcagcaca ggaattctca agacctgagt attttttata ataggaatgt 4885
ccacatgaa cttgatacgt ccgtgtgtcc cagatgctgt cattagteta tatggttctc 4945
caagaaactg aatgaatcca ttggagaagc ggtggataac tagccagaca aaatttgaga 5005
atacataaac aacgcattgc cacggaaaca tacagaggat gccttttctg tgattgggtg 5065
ggattttttc ctttttatg tgggatatag tagttacttg tgacaaaaat aattttggaa 5125
taatttctat taatatcaag tctgaagcta attgtactaa tctgagattg tgtttgttca 5185
taataaaagt gaagtgaatc taaaaaaaaa aaaaaa 5221

```

<210> 4  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

<400> 4  
 gatccccgca tccgaga 17

<210> 5  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

<400> 5  
ggtataggag ctagagatat cgtcctg 27

<210> 6  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 6  
ccaagatgcc ggcccacttg c 21

<210> 7  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 7  
gaaggtgaag gtcggagtc 19

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 8  
gaagatggtg atgggatttc 20

<210> 9  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 9  
caagcttccc gttctcagcc 20

<210> 10  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 10  
gtccggtatt tcctcagccc 20

<210> 11

<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 11  
ccgcggtgcg tggaggtccc 20

<210> 12  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 12  
tacgcgctga gccgcggcgc 20

<210> 13  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 13  
gcggtttcga agcccgccgg 20

<210> 14  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 14  
cctccattct gcaggaccct 20

<210> 15  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 15  
tccaagtgt agcagagaca 20

<210> 16  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 16  
ctcctgctgt tatgcccagg 20

<210> 17  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 17  
cacggtggtc acgagcccat 20

<210> 18  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 18  
cagccaacc acgtgagaga 20

<210> 19  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 19  
gacaagtcta gcgtactccc 20

<210> 20  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 20  
gttcaccagc caggtggcat 20

<210> 21  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 21

tgtggaagcc ctcacccaca 20

<210> 22  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 22  
agttgatgtg ccagcggtag 20

<210> 23  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 23  
ggacccccaaa ctcagccact 20

<210> 24  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 24  
tgcctggggag gcaataaggg 20

<210> 25  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 25  
atacatgcta actctctccc 20

<210> 26  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 26  
aagtcctcat taggtaggca 20

<210> 27  
<211> 20

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 27  
tgtaatgagc agctcatgga 20

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 28  
tcagtaacct tctcaagccc 20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 29  
ggagctgcct ggacagcaag 20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 30  
tcagtgacct tgagcattct 20

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 31  
tggctggccc actggctcaa 20

<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 32  
gcatgccctc tggttctgac 20

<210> 33  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 33  
gctttgcagt tcaccctgac 20

<210> 34  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 34  
gtggtatctc aaatcccagg 20

<210> 35  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 35  
tagtccaggc taaccctgt 20

<210> 36  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 36  
gtgatcttcc cttagatcct 20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 37  
ctcagcagac aactcccac 20

<210> 38  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 38  
gctaagttgt cagcacaccc 20

<210> 39  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 39  
aagtttccag aatgaagccc 20

<210> 40  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 40  
agagaataca cccaagatac 20

<210> 41  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 41  
tagttaagtg acttgcccag 20

<210> 42  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 42  
gccctttgag gtaggtcagt 20

<210> 43  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 43  
ccatatagac taatgacagc 20

<210> 44  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 44  
ctgtatgttt ccgtggcaat 20

<210> 45  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 45  
cttgcacgct agctggttgt 20

<210> 46  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 46  
gcatcgtctc caacttatct 20

<210> 47  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 47  
taaggatgat gtttctccag 20

<210> 48  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 48

cccaaagcca ggtgtagaac 20

<210> 49  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 49  
tgtaagagcg gtggctccac 20

<210> 50  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 50  
cattctggaa tgccattgtg 20

<210> 51  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 51  
ttatgaggat cagcatgtgt 20

<210> 52  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 52  
cctctggaac atcaccagtt 20

<210> 53  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 53  
ggatgaagca catcagcagc 20

<210> 54  
<211> 20

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 54  
ttcacccag aaataccagg 20

<210> 55  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 55  
gaaaccagga tattctcccg 20

<210> 56  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 56  
tcactggcag agtagtcata 20

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 57  
taatcctggc caagatggcg 20

<210> 58  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 58  
tcatcatctt tagcatcctg 20

<210> 59  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 59  
gcaaagactg accagctgct 20

<210> 60  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 60  
gactaccag aagattctgt 20

<210> 61  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 61  
cttcctcat cttacattc 20

<210> 62  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 62  
cccgagccag gagagaaagg 20

<210> 63  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 63  
cttccccagc agagaccact 20

<210> 64  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 64  
ccaatattcct gaagatggca 20

<210> 65  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 65  
cccaactaat tcctcctctc 20

<210> 66  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 66  
tatagatcct gtcctcagc 20

<210> 67  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 67  
ctcccaataa ctcaactcagt 20

<210> 68  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 68  
aagagattcc taaccctgcc 20

<210> 69  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 69  
cacacaaagg aggctgcctg 20

<210> 70  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 70  
aagtggcagc tagctctact 20

<210> 71  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 71  
cacctcacc aagtaagcag 20

<210> 72  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 72  
tgcttcttcc cagtgagaac 20

<210> 73  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 73  
atcaagcagg catctgatga 20

<210> 74  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 74  
ccctcagcct gagtgccat 20

<210> 75  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 75

ataatcctcc actcaggccc 20

<210> 76  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 76  
cacttaagaa aagcagccct 20

<210> 77  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 77  
cagcaagtca gtggcacagt 20

<210> 78  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 78  
ggctagttat ccaccgcttc 20

<210> 79  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 79  
cccaatcaca gaaaaggcat 20

<210> 80  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 80  
aactactata tcccacataa 20