



(51) International Patent Classification:

C07D 417/12 (2006.01) C07J 19/00 (2006.01)
C07D 487/04 (2006.01) C07D 493/10 (2006.01)
C07D 417/04 (2006.01) C12Q 1/66 (2006.01)
C07D 417/14 (2006.01) G01N 33/58 (2006.01)
C07J 9/00 (2006.01)

(21) International Application Number:

PCT/US2015/013617

(22) International Filing Date:

29 January 2015 (29.01.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/933,122 29 January 2014 (29.01.2014) US

(71) Applicant: **PROMEGA CORPORATION** [US/US];
2800 Woods Hollow Road, Madison, WI 53711-5399
(US).

(72) Inventor; and

(71) Applicant : **MUSTAFA, Dana** [JO/US]; 1422 Woodside
Drive, Apt. F, San Luis Obispo, CA 93401 (US).

(72) Inventors: **ZHOU, Wenhui**; 3936 Hollyhock Way, San
Luis Obispo, CA 93401 (US). **MEISENHERMIER,
Poncho**; 2868 Victoria Avenue, San Luis Obispo, CA
93401 (US). **DUELLMAN, Sarah**; 2814 Sunflower Drive,
Fitchburg, WI 53711 (US). **MA, Dongping**; 1226 Day-
flower Drive, Madison, WI 53719 (US). **CALL, James, J.**;
546 Harvest Lane, Verona, WI 53593 (US).

(74) Agent: **YEH, Sansun**; Michael Best & Friedrich LLP, 100
East Wisconsin Avenue, Suite 3300, Milwaukee, WI
53202-4108 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,

[Continued on next page]

(54) Title: QUINONE-MASKED PROBES AS LABELING REAGENTS FOR CELL UPTAKE MEASUREMENTS





FIG. 1



(57) Abstract: Provided are labeling reagents and methods of using the reagents for cell uptake measurements. The labeling reagents can be quinone-masked probes including fluorophores and/or luminophores.

MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,

Published:

— *with international search report (Art. 21(3))*

QUINONE-MASKED PROBES AS LABELING REAGENTS FOR CELL UPTAKE MEASUREMENTS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application 61/933,122, filed January 29, 2014, and is herein incorporated by reference in its entirety.

TECHNICAL FIELD


[0002] The present disclosure relates generally to compounds, compositions, and methods of using the compounds and compositions as labeling reagents. More particularly, the present disclosure relates to quinone-masked probes as labeling reagents for cell uptake measurements.

BACKGROUND

[0003] Quantification of the uptake of biomolecules and small molecules of interests in biological samples (i.e., cells, tissues and organs) is becoming a vital part for drug delivery, effectiveness of therapeutic treatments, profiling pharmacokinetics, drug-drug interactions and drug toxicity, but also represents an important indicator for cellular metabolic status, such as fatty acids and glucose, and homeostasis of cholesterol to maintain normal physiological functions. Various techniques such as fluorescent-labeled or isotope-labeled molecules of interests for uptake quantification of cells and tissues already exist, but still needs for new uptake techniques in order to make the uptake measurements simple, reliable and reproducible in high throughput formats.

SUMMARY

[0004] In some aspects, disclosed is a compound of formula (I), or a salt thereof,



(I)

wherein

A is a reporter moiety;

R^{14} is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R^9 and R^{10} are independently selected from alkyl;

R^{11} , R^{12} and R^{13} are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R^{11} and R^{12} can form a fused phenyl ring;

X is O;

L is $-(CH_2)_mC(R^{17})_2(CH_2)_n-Y-C(O)-$;

R^{17} is independently H, alkyl or both R^{17} together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;

n is an integer from 0-2;

Y is O or NR^{15} ;

R^{15} is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl- $N(R^{23})C(O)R^{24}$, -alkyl- SO_3R^{25} , -alkyl- $SO_2N(R^{26})(R^{27})$, -alkyl- COR^{28} , -alkyl- CO_2R^{29} , -alkyl- $OC(O)R^{29}$, -alkyl- $OC(O)N(R^{30})(R^{31})$, -alkyl- $CON(R^{30})(R^{31})$, or polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with one or more suitable substituents; and


R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[0005] In certain embodiments, R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-haloalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-hydroxyalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-aminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkylaminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{di(alkyl)aminoalkyl}$, $-(C_2-C_6\text{-$

alkoxy)_x-azidoalkyl, -(C₂-C₆-alkoxy)_x-cyanoalkyl, -(C₂-C₆-alkoxy)_x-alkenyl, -(C₂-C₆-alkoxy)_x-alkynyl, -(C₂-C₆-alkoxy)_x-N(R²³)C(O)R²⁴, -(C₂-C₆-alkoxy)_x-SO₃R²⁵, -(C₂-C₆-alkoxy)_x-alkyl-SO₂N(R²⁶)(R²⁷), -(C₂-C₆-alkoxy)_x-alkyl-COR²⁸, -(C₂-C₆-alkoxy)_x-alkyl-OC(O)R²⁹, -(C₂-C₆-alkoxy)_x-alkyl-OC(O)N(R³⁰)(R³¹), -(C₂-C₆-alkoxy)_x-alkyl-CO₂R²⁹, and -(C₂-C₆-alkoxy)_x-alkyl-CON(R³⁰)(R³¹), wherein R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl; and x is an integer selected from 1 to 20.

[0006] In certain embodiments, R¹⁵ is polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of alkyl, alkoxyalkyl, haloalkyl, haloalkoxyalkyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, azidoalkyl, cyanoalkyl, alkoxy, haloalkoxy, alkoxyalkoxy, alkenyl, alkynyl, -N(R²³)C(O)R²⁴, -SO₃R²⁵, -SO₂N(R²⁶)(R²⁷), -COR²⁸, -CO₂R²⁹, and -CON(R³⁰)(R³¹), wherein R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[0007] In certain embodiments, R¹⁵ is



wherein


R⁴⁰ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein said aryl, heteroaryl, heterocyclyl, and cycloalkyl are unsubstituted or substituted with one or more suitable substituents; and

x is an integer selected from 1 to 20.

[0008] In certain embodiments, R⁴⁰ is hydrogen.


[0009] In certain embodiments, R⁴⁰ is a 5- or 6-membered heterocyclyl, having 1, 2, or 3 heteroatoms independently selected from the group consisting of N, O, and S.

[0010] In certain embodiments, R⁴⁰ is



[0011] In certain embodiments, x is 2, 3, or 4.

[0012] In certain embodiments, the compound of formula (I) has formula (I-vi), or a salt thereof,




(I-vi)

wherein

R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{40} , x, and A are as defined above.

[0013] In some aspects, disclosed is a compound of formula (I), or a salt thereof,

A compound of formula (I), or a salt thereof,



(I)

wherein

A is a reporter moiety;

R^{14} is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R^9 and R^{10} are independently selected from alkyl;

R^{11} , R^{12} and R^{13} are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R^{11} and R^{12} can form a fused phenyl ring;

X is O;

L is $-(CH_2)_mC(R^{17})_2(CH_2)_n-Y-C(O)-$;

R^{17} is independently H, alkyl or both R^{17} together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;


n is an integer from 0-2;

Y is O or NR^{15} ;

R^{15} is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl- $N(R^{23})C(O)R^{24}$, -alkyl- SO_3R^{25} , -alkyl- $SO_2N(R^{26})(R^{27})$, -alkyl- COR^{28} , -alkyl- CO_2R^{29} , -alkyl- $OC(O)R^{29}$, -alkyl- $OC(O)N(R^{30})(R^{31})$, -alkyl-amide, or polyalkoxyalkyl, wherein the polyalkoxyalkyl and -alkyl-amide are unsubstituted or substituted with one or more suitable substituents; and


R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[0014] In certain embodiments, a compound of the invention is selected from the group consisting of:




PBI 5463

;




PBI 5470

;




PBI 5471


;




PBI 5508




PBI 5915





PBI 5916







PBI 5917



PBI 5918









PBI 5666

;





PBI 5684


;



PBI 5825


;






PBI 5682

;





PBI 5647







PBI 5668



;











;











;















; and

or a salt thereof.

[0015] In certain embodiments, the reporter moiety is a bioluminescent or fluorescent reporter moiety. In certain embodiments, the bioluminescent reporter moiety is a luciferin, a luciferin derivative or analog, or a luciferin precursor analog, coelenterazine or a coelenterazine derivative or analog.

[0016] In other aspects, disclosed is a method for evaluating cellular uptake of an agent comprising: a) contacting a sample comprising cells with a labeled agent, wherein the labeled

agent is derived from an agent and a compound of formula (I); and b) detecting light emission, whereby the detection of light emission indicates cellular uptake of the agent.

[0017] In certain embodiments, the compound is a prosubstrate for luciferase or an alternative pre-prosubstrate that does not react with luciferase.

[0018] In certain embodiments, the light emission is fluorescence or luminescence.

[0019] In certain embodiments, the light emission is luminescence, and the cell comprises a luciferase.

[0020] In certain embodiments, the light emission is detected in the cell, outside the cell, e.g., in the cell medium or in a biological fluid, e.g., bodily fluid, e.g., blood, urine, etc.

[0021] In certain embodiments, the compound that does not react with luciferase fluoresces in the cell.

[0022] In certain embodiments, the agent is a biological agent or a non-biological agent.

[0023] In certain embodiments, the agent is selected from the group consisting of a protein, a nucleic acid, a lipid, a sugar, a therapeutic drug, a small molecule, a nanoparticle, antibody, and any combination thereof.

[0024] In certain embodiments, the protein is an antibody or a lipoprotein.

[0025] In certain embodiments, the cell is a eukaryotic cell.

[0026] In certain embodiments, the cell is a prokaryotic cell.

[0027] In certain embodiments, the cell is in an animal.

[0028] In certain embodiments, the cell is growing in culture medium.

[0029] In certain embodiments, the cell is a live cell.

[0030] In certain embodiments, the cell expresses a luciferase enzyme.

[0031] In another aspect, disclosed is a labeled agent derived from an agent and a compound according to formula (I).

[0032] In certain embodiments, the compound in a cell is a substrate for luciferase or an alternative substrate that does not react with luciferase.

[0033] In other aspects, disclosed is a kit comprising a compound of formula (I), an agent, a labeled agent derived from an agent and a compound according to formula (I), or any combination thereof.

[0034] In certain embodiments, the compound is a prosubstrate for luciferase.

[0035] In certain embodiments, the kit further comprises a detection reagent. The detection reagent may comprise a luciferase enzyme.

[0036] The compounds, compositions, methods, and processes are further described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

- [0037] FIG. 1 shows the transporter assay with PBI-5651.
- [0038] FIG. 2 shows the transporter assay with PBI-5648.
- [0039] FIG. 3 shows the transporter assay with PBI-5657.
- [0040] FIG. 4 shows the transporter assay with PBI-5658.
- [0041] FIG. 5 shows the transporter assay with PBI-5665.
- [0042] FIG. 6 shows the transporter assay with PBI-5666.
- [0043] FIG. 7 shows the transporter assay with PBI-5824.
- [0044] FIG. 8 shows the transporter assay with PBI-5684.
- [0045] FIG. 9 shows the transporter assay with PBI-5825.
- [0046] FIG. 10 shows the transporter assay with PBI-5683.
- [0047] FIG. 11 shows the transporter assay with PBI-5826.
- [0048] FIG. 12 shows the transporter assay with PBI-5682.
- [0049] FIG. 13 shows the transporter assay with various amounts of PBI-5651.
- [0050] FIG. 14 shows examples of monoclonal antibody (mAb) quinone-luciferin conjugates.
- [0051] FIG. 15 shows examples of labeled solid-liquid nanoparticles (SLNs).
- [0052] FIG. 16 shows an example of labeled Poly(amido amine) (PAMAM) dendrimer.
- [0053] FIG. 17 shows an example of labeled oligonucleotide and peptide-oligonucleotide conjugates for antisense inhibition of P-glycoprotein expression.

DETAILED DESCRIPTION

[0054] Disclosed herein are quinone-masked probes including fluorophores and/or luminophores. The compounds can be used as labeling reagents for cell uptake measurements. These labeling reagents can be broadly used to label a variety of targets of interest such as peptides, proteins including but not limited to antibodies or low density lipoproteins (LDL), nucleotides, sugars, lipids, specific substrates for transporters, or nanoparticles and microsomes

including but not limited to drug carriers. The labeling reagents may also be used to label detergents. Upon uptake of labeled targets of interest by cells, quinone can be reduced by intracellular reduction potential to release the fluorophore or luminophore. The intensity of fluorescence or luminescence can be correlated to the amount of the target(s) of interest taken up by cells. Also disclosed are labeled agents.

[0055] The disclosed labeling reagents and labeled agents have multiple advantages over current techniques: they allow for either imaging or plate-based detection; they are less sensitive to extracellular reductants (e.g. free thiols, DTT, GSH); they avoid the need for fixed cells; and no washing step is required opposed to traditional isotope and fluorescence methods.

1. Definition of Terms

[0056] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. In case of conflict, the present document, including definitions, will control. Preferred methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in practice or testing of the present invention. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples disclosed herein are illustrative only and not intended to be limiting.

[0057] The terms “comprise(s),” “include(s),” “having,” “has,” “can,” “contain(s),” and variants thereof, as used herein, are intended to be open-ended transitional phrases, terms, or words that do not preclude the possibility of additional acts or structures. The present disclosure also contemplates other embodiments “comprising,” “consisting of” and “consisting essentially of,” the embodiments or elements presented herein, whether explicitly set forth or not.

[0058] As used herein, the term “suitable substituent” is intended to mean a chemically acceptable functional group e.g., a moiety that does not negate the activity of the inventive compounds. Illustrative examples of suitable substituents include, but are not limited to halo groups, perfluoroalkyl groups, perfluoroalkoxy groups, alkyl groups, alkenyl groups, alkynyl groups, hydroxy groups, halo groups, oxo groups, mercapto groups, alkylthio groups, alkoxy groups, nitro groups, azidealkyl groups, aryl or heteroaryl groups, aryloxy or heteroaryloxy groups, aralkyl or heteroaralkyl groups, aralkoxy or heteroaralkoxy groups, HO—(C=O)—

groups, heterocyclic groups, cycloalkyl groups, amino groups, alkyl - and dialkylamino groups, carbamoyl groups, alkylcarbonyl groups, alkylcarbonyloxy groups, alkoxy carbonyl groups, alkylaminocarbonyl groups, dialkylamino carbonyl groups, arylcarbonyl groups, aryloxy carbonyl groups, alkylsulfonyl groups, arylsulfonyl groups and the like. The substituents can be substituted by additional substituents.

[0059] As used herein, the term “alkenyl” refers a straight or branched hydrocarbon chain containing from 2 to 10 carbons and containing at least one carbon-carbon double bond formed by the removal of two hydrogens. Representative examples of alkenyl include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl. Alkenyl groups of the present invention may be unsubstituted or substituted by one or more suitable substituents, preferably 1 to 3 suitable substituents, as defined above.

[0060] As used herein, the term “alkoxy” refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy, tert-butoxy, pentyloxy, and hexyloxy.

[0061] As used herein, the term “alkoxyalkoxy” refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through another alkoxy group, as defined herein. Representative examples of alkoxyalkoxy include, but are not limited to, tert-butoxymethoxy, 2-ethoxyethoxy, 2-methoxyethoxy, and methoxymethoxy.

[0062] As used herein, the term “alkoxyalkyl” refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkoxyalkyl include, but are not limited to, tert-butoxymethyl, 2-ethoxyethyl, 2-methoxyethyl, and methoxymethyl.

[0063] As used herein, the term “alkoxycarbonyl” refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkoxy carbonyl include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, and tert-butoxycarbonyl.

[0064] As used herein, the term “alkyl” refers to a linear or branched hydrocarbon radical, preferably having 1 to 30 carbon atoms, 1 to 10 carbon atoms, 1 to 6 carbon atoms, or 1 to 4 carbon atoms. The term “C₁-C₆-alkyl” is defined to include alkyl groups having 1, 2, 3, 4, 5, or 6

carbons in a linear or branched arrangement. For example, “C₁-C₆-alkyl” specifically includes methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl, pentyl, and hexyl. Alkyl groups of the present invention may be unsubstituted or substituted by one or more suitable substituents, preferably 1 to 3 suitable substituents, as defined above.

[0065] As used herein, the term “alkylamino” refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an amino group, as defined herein.

Representative examples of alkylamino include, but are not limited to, methylamino, ethylamino, and sec-butylamino.

[0066] As used herein, the term “alkylaminoalkyl” refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an aminoalkyl group, as defined herein.

Representative examples of alkylaminoalkyl groups include, but are not limited to, methylaminoethyl and methylamino-2-propyl.

[0067] As used herein, the term “alkylcarbonyl” refers to an alkyl group appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkylcarbonyl include, but are not limited to, acetyl, 1-oxopropyl, 2,2-dimethyl-1-oxopropyl, 1-oxobutyl, and 1-oxopentyl.

[0068] As used herein, the term “alkylcarbonylalkoxy” refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of alkylcarbonylalkoxy include, but are not limited to, 3-oxopentyloxy, 3-oxobutoxy and 2-oxopropoxy.

[0069] As used herein, the term “alkylcarbonylalkoxyalkyl” refers to an alkylcarbonylalkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group.

[0070] As used herein, the term “alkylcarbonyloxy” refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom.

Representative examples of alkylcarbonyloxy include, but are not limited to, acetyloxy, ethylcarbonyloxy, and tert-butylcarbonyloxy.

[0071] As used herein, the term “alkynyl” refers to a straight or branched hydrocarbon radical having 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons, and having one or more carbon-carbon triple bonds. Alkynyl groups of the present invention include, but are not limited to, ethynyl, propynyl, and butynyl. Alkynyl groups of the present invention may be unsubstituted or substituted by one or more suitable substituents, preferably 1 to 3 suitable substituents, as defined above.

[0072] As used herein, the term “amidoalkyl” refers to an amide group, appended to the parent molecular moiety through an alkyl group as defined herein. The term amidoalkyl includes $-\text{CH}_2\text{CONH}_2$; $-\text{CH}_2\text{CH}_2\text{CONH}_2$; $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CONH}_2$ and the like.

[0073] As used herein, the term “amino” refers to an $-\text{NH}_2$ group.

[0074] As used herein, the term “amino acid” refers to both natural and unnatural amino acids. It also includes protected natural and unnatural amino acids.

[0075] As used herein, the term “aminoalkyl” refers to at least one amino group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of aminoalkyl include, but are not limited to, aminomethyl, 2-aminoethyl, and 2-aminopropyl.

[0076] As used herein, the term “aryl” means monocyclic, bicyclic, or tricyclic aromatic radicals. Representative examples of the aryl groups include, but are not limited to, phenyl, dihydroindenyl, indenyl, naphthyl, dihydronaphthalenyl, and tetrahydronaphthalenyl. Aryl groups of the present invention may be optionally substituted by one or more suitable substituents, preferably 1 to 5 suitable substituents, as defined above.

[0077] As used herein, the term “arylalkyl” refers to an aryl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of arylalkyl include, but are not limited to, phenylmethyl and phenylethyl.

[0078] As used herein, the term “arylcarbonyl” refers to an aryl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein.

[0079] As use herein, the term “azide” refers to an $-\text{N}=\text{N}^+=\text{N}^-$ ($-\text{N}_3$) group.

[0080] As used herein, the term “azidealkyl” or “azidoalkyl” refers to an azide group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of azidealkyl include, but are not limited to, azidemethyl and azideethyl.

[0081] As used herein, the term “bioluminescence” or “luminescence” may refer to light produced as a result of a reaction between an enzyme and a substrate that generates light. Examples of such enzymes (bioluminescent enzymes) include firefly luciferase, e.g. *Photinus pyralis* or *Photuris pennsylvanica*, click beetle luciferase, *Renilla* luciferase, cypridina luciferase,

Oplophorus luciferase, e.g., *Oplophorous gracilirostris*, Aequorin photoprotein, obelin photoprotein and the like.

[0082] As used herein, the term “derivative” may refer to a compound that is derived from a similar compound by some chemical or physical process. The derivative is a compound of similar chemical structure. The derivative may be a structural analogue.

[0083] As used herein, the term “reporter moiety” may refer to a moiety that, under appropriate conditions directly or indirectly generates a detectable signal. Exemplary reporter moieties include, but are not limited to, fluorophores, luminescent molecules, dyes, radiolabels and substrates for enzymes such as luciferase. In some embodiments, a reporter moiety may indirectly generate a detectable signal, for example, when the reporter moiety is a substrate for an enzyme. The reaction of the enzyme with the substrate then produces a detectable signal such as fluorescence or luminescence. As used herein, the term “bioluminescent reporter moiety” may refer to a moiety that is a substrate for a luciferase. For example, the bioluminescent reporter moiety can be a luciferin, a luciferin derivative, e.g., pre-luciferin, aminoluciferin, quionolyl-luciferin, naphthyl luciferin, fluoroluciferin, chloroluciferin, precursors of luciferin derivatives, a coelenterazine or a coelenterazine derivative or analog, e.g., furimazine. The luminescent signal generated may be detected using a luminometer. As used herein, the term “fluorescent reporter moiety” may refer to a moiety that fluoresces. For example, the fluorescent reporter moiety may be a fluorophore, such as coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, sily xanthene, or carbopyronine. Fluorescence may be detected using a fluorometer.

[0084] As used herein, the term “carbonyl” or “(C=O)” (as used in phrases such as alkylcarbonyl, alkyl -(C=O)— or alkoxy carbonyl) refers to the joiner of the >C=O moiety to a second moiety such as an alkyl or amino group (i.e. an amido group). Alkoxy carbonylamino (i.e. alkoxy(C=O)—NH—) refers to an alkyl carbamate group. The carbonyl group is also equivalently defined herein as (C=O). Alkylcarbonylamino refers to groups such as acetamide.

[0085] As used herein, the term “carboxy” refers to a —C(O)—OH group.

[0086] As used herein, the term “carboxyalkoxyalkyl” refers to —alkyl—O—alkyl—CO₂H.

[0087] As used herein, the term “carboxyalkyl” refers to a carboxy group as defined herein, appended to the parent molecular moiety through an alkyl group as defined herein.

[0088] As used herein, the term “cycloalkyl” refers to a mono, bicyclic or tricyclic carbocyclic radical (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclopentenyl, cyclohexenyl, bicyclo[2.2.1]heptanyl, bicyclo[3.2.1]octanyl and bicyclo[5.2.0]nonanyl, etc.); optionally containing 1 or 2 double bonds. Cycloalkyl groups of the present invention may be unsubstituted or substituted by one or more suitable substituents, preferably 1 to 5 suitable substituents, as defined above.

[0089] As used herein, the term “di(alkyl)amino” refers to two independently selected alkyl groups, as defined herein, appended to the parent molecular moiety through an amino group, as defined herein. Representative examples of di(alkyl)amino include, but are not limited to, N,N-dimethylamino, N-ethyl-N-methylamino, and N-isopropyl-N-methylamino.

[0090] As used herein, the term “di(alkyl)aminoalkyl” refers to a di(alkyl)amino group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of di(alkyl)aminoalkyl include, but are not limited to, N,N-dimethylaminoethyl and N,N-methyl(2-propyl)aminoethyl.

[0091] As used herein, the term “halogen” or “halo” refers to a fluoro, chloro, bromo or iodo radical.

[0092] As used herein, the term “haloalkoxy” refers to an alkoxy group, as defined herein, substituted by one, two, three, or four halogen atoms. Representative examples of haloalkoxy include, but are not limited to, chloromethoxy, 2-fluoroethoxy, trifluoromethoxy, and pentafluoroethoxy.

[0093] As used herein, the term “haloalkyl” refers to an alkyl group, as defined herein, substituted by one, two, three, or four halogen atoms. Representative examples of haloalkyl include, but are not limited to, chloromethyl, 2-fluoroethyl, trifluoromethyl, pentafluoroethyl, 2-chloro-3-fluoropentyl, and 4,4,4-trifluorobutyl.

[0094] As used herein, the term “heteroaryl” refers to a monocyclic heteroaryl or a bicyclic heteroaryl. The monocyclic heteroaryl is a five- or six-membered ring. The five-membered ring contains two double bonds. The five-membered ring may contain one heteroatom selected from O or S; or one, two, three, or four nitrogen atoms and optionally one oxygen or sulfur atom. The six-membered ring contains three double bonds and one, two, three or four nitrogen atoms. Representative examples of monocyclic heteroaryl include, but are not limited to, furanyl, imidazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, 1,3-oxazolyl, pyridinyl, pyridazinyl,

pyrimidinyl, pyrazinyl, pyrazolyl, pyrrolyl, tetrazolyl, thiadiazolyl, 1,3-thiazolyl, thienyl, triazolyl, and triazinyl. The bicyclic heteroaryl includes a monocyclic heteroaryl fused to a phenyl, or a monocyclic heteroaryl fused to a monocyclic cycloalkyl, or a monocyclic heteroaryl fused to a monocyclic cycloalkenyl, or a monocyclic heteroaryl fused to a monocyclic heteroaryl, or a monocyclic heteroaryl fused to a monocyclic heterocycle. Representative examples of bicyclic heteroaryl groups include, but are not limited to, benzofuranyl, benzothienyl, benzoxazolyl, benzimidazolyl, benzoxadiazolyl, 6,7-dihydro-1,3-benzothiazolyl, imidazo[1,2-a]pyridinyl, indazolyl, indolyl, isoindolyl, isoquinolyl, naphthyridinyl, pyridoimidazolyl, quinazolyl, quinolyl, thiazolo[5,4-b]pyridin-2-yl, thiazolo[5,4-d]pyrimidin-2-yl, and 5,6,7,8-tetrahydroquinolin-5-yl. Heteroaryl groups of the present invention may be unsubstituted or substituted by one or more suitable substituents, preferably 1 to 5 suitable substituents, as defined above.

[0095] As used herein, the term “heterocycle” or “heterocyclyl” refers to a monocyclic heterocycle, a bicyclic heterocycle, or a tricyclic heterocycle. The monocyclic heterocycle is a three-, four-, five-, six-, seven-, or eight-membered ring containing at least one heteroatom independently selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur. The three- or four-membered ring contains zero or one double bond, and one heteroatom selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur. The five-membered ring contains zero or one double bond and one, two or three heteroatoms selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur. The six-membered ring contains zero, one or two double bonds and one, two, or three heteroatoms selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur. The seven- and eight-membered rings contains zero, one, two, or three double bonds and one, two, or three heteroatoms selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur. Representative examples of monocyclic heterocycles include, but are not limited to, azetidiny, azepanyl, aziridiny, diazepanyl, 1,3-dioxanyl, 1,3-dioxolanyl, 1,3-dithiolanyl, 1,3-dithianyl, imidazoliny, imidazolidiny, isothiazoliny, isothiazolidiny, isoxazoliny, isoxazolidiny, morpholiny, oxadiazoliny, oxadiazolidiny, oxazoliny, oxazolidiny, phosphinane, piperaziny, piperidiny, pyranyl, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidiny, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyridiny, tetrahydropyrimidiny, tetrahydrothienyl, thiadiazoliny, thiadiazolidiny, thiazoliny, thiazolidiny, thiomorpholiny, 1,1-dioxidothiomorpholiny (thiomorpholine sulfone),

thiopyranyl, trithianyl, and 2,5-dioxo-pyrrolidinyl. The bicyclic heterocycle is a monocyclic heterocycle fused to a phenyl group, or a monocyclic heterocycle fused to a monocyclic cycloalkyl, or a monocyclic heterocycle fused to a monocyclic cycloalkenyl, or a monocyclic heterocycle fused to a monocyclic heterocycle, or a bridged monocyclic heterocycle ring system in which two non-adjacent atoms of the ring are linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, or an alkenylene bridge of two, three, or four carbon atoms. Representative examples of bicyclic heterocycles include, but are not limited to, benzopyranyl, benzothiopyranyl, chromanyl, 2,3-dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, azabicyclo[2.2.1]heptyl (including 2-azabicyclo[2.2.1]hept-2-yl), 2,3-dihydro-1H-indolyl, isoindolyl, octahydrocyclopenta[c]pyrrolyl, octahydropyrrolopyridinyl, 9-phosphabicyclo[3.3.1]nonane, 8-phosphabicyclo[3.2.1]octane, and tetrahydroisoquinolyl. Tricyclic heterocycles are exemplified by a bicyclic heterocycle fused to a phenyl group, or a bicyclic heterocycle fused to a monocyclic cycloalkyl, or a bicyclic heterocycle fused to a monocyclic cycloalkenyl, or a bicyclic heterocycle fused to a monocyclic heterocycle, or a bicyclic heterocycle in which two non-adjacent atoms of the bicyclic ring are linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, or an alkenylene bridge of two, three, or four carbon atoms. Examples of tricyclic heterocycles include, but are not limited to, octahydro-2,5-epoxypentalene, hexahydro-2H-2,5-methanocyclopenta[b]furan, hexahydro-1H-1,4-methanocyclopenta[c]furan, aza-admantane (1-azatricyclo[3.3.1.1^{3,7}]decane), oxa-adamantane (2-oxatricyclo[3.3.1.1^{3,7}]decane), and 2,4,6-trioxa-8-phosphatricyclo[3.3.1.1^{3,7}]decane. Heterocyclic groups of the present invention may be unsubstituted or substituted by one or more suitable substituents, preferably 1 to 3 suitable substituents, as defined above. Heterocyclic groups of the present invention may be can contain one or more oxo groups (=O) or thioxo (=S) groups attached to the ring.

[0096] As used herein, the term “hydroxy” refers to an -OH group.

[0097] As used herein, the term “hydroxyalkoxy” refers to an alkoxy group, as defined herein, substituted by at least one hydroxy group. Representative examples of hydroxyalkoxy include, but are not limited to, hydroxyethoxy, and 2-hydroxypropoxy.

[0098] As used herein, the term “hydroxyalkyl” refers to an alkyl group, as defined herein, substituted by at least one hydroxy group. Representative examples of hydroxyalkyl include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2,3-dihydroxypropyl, 2,3-

dihydroxypentyl, 4-hydroxybutyl, 2-ethyl-4-hydroxyheptyl, 3,4-dihydroxybutyl, and 5-hydroxypentyl.

[0099] As used herein, the term “hydroxycarbonyl” refers to a hydroxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein.

[00100] The term “linker” may refer to a chain of 2 to 50 atoms that link a substrate moiety to the parent molecular moiety. Linkers may include one or more heteroatoms. Linkers may also be substituted by oxo groups, amino groups, alkyl groups, halogens and nitro groups. Linkers may also contain aryl groups. The linkers may be “traceless” or “self-immolative” linkers. The term “traceless linker” or “self-immolative linker” refers to a linker wherein cleavage of the substrate moiety from the linker results in spontaneous cleavage of the linker from the parent molecular moiety.

[00101] The term “lower cycloalkyl” refers to a monovalent moiety obtained by removing a hydrogen atom from a hydrocarbon compound having from 3 to 6 carbon atoms. Examples of saturated lower cycloalkyl groups include, but are not limited to, groups such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. Examples of unsaturated lower cycloalkyl groups which have one or more carbon-carbon double bonds include, but are not limited to, groups such as cyclopropenyl, cyclobutenyl, cyclopentenyl and cyclohexenyl.

[00102] The term “luminescent enzyme”, “bioluminescent enzyme”, or “luciferase” as used interchangeably herein refers to a class of oxidative enzymes used in bioluminescence wherein the enzyme produces and emits light when given a substrate. The luciferase may be a naturally occurring, recombinant or mutant luciferase that uses a luciferase substrate. The luciferase substrate may be luciferin, a luciferin derivative or analog, a preluciferin derivative or analog, a coelenterazine, or a coelenterazine derivative or analog. The luminescent enzyme, if naturally occurring, may be obtained easily by the skilled person from an organism. If the luminescent enzyme is one that occurs naturally or is a recombinant or mutant luminescent enzyme, e.g. one which retains activity in a luciferase-coelenterazine or luciferase-luciferin reaction of a naturally occurring luminescent enzyme, it can be obtained readily from a culture of bacteria, yeast, mammalian cells, insect cells, plant cells, or the like, transformed to express a nucleic acid encoding the luminescent enzyme. Further, the recombinant or mutant luminescent enzyme can be derived from an in vitro cell-free system using a nucleic acid encoding the luciferase. Suitable luminescent enzymes include luciferases derived from bioluminescent decapods, such

as from the Oplophoroidea (e.g. *Oplophorus*-derived luciferases), beetle luciferases (e.g., *Photinus pyralis*, *Photuris pennsylvanica*, etc.), marine organisms such as cnidarians (e.g., *Renilla* luciferase), Aristeidae, Solenoceridae, Luciferidae, Sergestidae, Pasipheidae and Thalassocarididae decapoda families, copepod luciferases, such as *Gaussia* luciferase, such as *Gaussia princeps* luciferase, *Metridia* luciferases, such as *Metridia longa* and *Metridia pacifica* luciferases, *Vargula* luciferases, such as *Vargula hilgendorffii* luciferase, *Pleuromamma xiphias* luciferase, and photoproteins, such as *Aequorin*, and variants, recombinants, and mutants thereof.

[00103] A “luminescent reaction mixture” contains materials that will allow the luminescent enzyme to generate a light signal, i.e., luminescence. The mixture may also contain the enzyme, e.g., the luciferase enzyme or luciferase. The materials, and the particular concentrations and/or amounts, needed to generate a luminescent signal will vary depending on the luminescent enzyme used as well as the type of assay being performed. Often other materials will be added to the solution including: a buffer to maintain the reaction at the proper pH, an additive such as PRIONEX or Bovine serum albumin (BSA) to help maintain enzyme activity, reducing agents, detergents, etc.

[00104] As used herein, the term “methylenedioxy” refers to a —OCH₂O— group wherein the oxygen atoms of the methylenedioxy are attached to the parent molecular moiety through two adjacent carbon atoms.

[00105] As used herein, the term “natural amino acid” refers to any one of the common, naturally occurring L-amino acids found in proteins, including glycine (Gly), alanine (Ala), valine (Val), leucine (Leu), isoleucine (Ile), lysine (Lys), arginine (Arg), histidine (His), proline (Pro), serine (Ser), threonine (Thr), phenylalanine (Phe), tyrosine (Tyr), tryptophan (Trp), aspartic acid (Asp), glutamic acid (Glu), asparagine (Asn), glutamine (Gln), cysteine (Cys) and methionine (Met).

[00106] As used herein, the term “unnatural amino acid” refers to all amino acids which are not natural amino acids as described above. Such amino acids include the D-isomers of any of the 20 naturally occurring amino acids described above. Unnatural amino acids also include homoserine, ornithine, norleucine, and thyroxine. Additional unnatural amino acids are well known to one of ordinary skill in the art. An unnatural amino acid may be a D- or L-isomer. An unnatural amino acid may also be an alpha amino acid or a beta amino acid. An unnatural amino acid may also be a post-translationally modified amino acid, such as a phosphorylated serine,

threonine or tyrosine, an acylated lysine, or an alkylated lysine or arginine. Many forms of post-translationally modified amino acids are known.

[00107] As used herein, the term “protected amino acid,” refers to an amino acid side chain as described above which additionally contains a protected functional group. Protecting groups are well known in the art and are intended to protect such functional groups as amino, hydroxy, thio or carboxy against undesirable reactions during synthetic procedures. The protecting groups may be removed by a chemical reaction following the synthesis. Examples of protected amino acid side chains include benzyloxymethyl derived from serine, (4-methoxyphenyl)methyl derived from tyrosine, and tert-butylpropanoate derived from glutamate.

[00108] As used herein, the term “amino acid side chain,” refers to the group attached to the α -carbon of an amino acid. It is the characterizing portion of an amino acid and is derived from a corresponding amino acid by elimination of the $\text{NH}_2\text{CH}(\text{O})\text{OH}$ moiety. For example, the amino acid side chain of alanine is methyl, and the amino acid side chain of phenylalanine is phenylmethyl. An amino acid side chain may be a natural amino acid side chain or an unnatural amino acid side chain. In some embodiments, an amino acid side chain may be a protected amino acid side chain.

[00109] As used herein, the term “amino protecting group,” refers to a moiety that prevents chemical reactions from occurring on the nitrogen atom to which that protecting group is attached. An amino protecting group must also be removable by a chemical reaction. Such groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, the entirety of which is incorporated herein by reference. Suitable amino protecting groups include, but are not limited to, carbobenzyloxy ($-\text{NHCO}-\text{OCH}_2\text{C}_6\text{H}_5$ or $-\text{NH-Cbz}$); t-butyloxycarbonyl ($-\text{NHCO}-\text{OC}(\text{CH}_3)_3$ or $-\text{NH-Boc}$); 9-fluorenylmethyloxycarbonyl ($-\text{NH-Fmoc}$), 2,2,2-trichloroethyloxycarbonyl ($-\text{NH-Troc}$), and allyloxycarbonyl ($-\text{NH-Alloc}$). (In each of the above, the $-\text{NH}-$ represents the nitrogen from the amino group that is being protected.)

[00110] As used herein, the term “amino blocking group,” refers to a moiety that prevents chemical reactions from occurring on the nitrogen atom to which that blocking group is attached. In contrast to an amino protecting group, an amino blocking group is not intended to be removed by a chemical reaction. Such groups include, for example, acyl groups such as acetyl ($-\text{NHCOCH}_3$)

NHCO—CH₃), and succinyl (—NH—CO—CH₂—CH₂—COO⁻). (In each of the above, the —NH— represents the nitrogen from the amino group that is being blocked.)

[00111] As used herein, the term “aminoluciferin” refers to (4S)-2-(6-amino-1,3-benzothiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid, or a substituted analog of this molecule.

[00112] As used herein, the term “nitrogen protecting group” refers to groups intended to protect an amino group against undesirable reactions during synthetic procedures.

Representative nitrogen protecting groups include acetyl, benzoyl, benzyl, benzyloxycarbonyl (Cbz), formyl, phenylsulfonyl, tert-butoxycarbonyl (Boc), tert-butylacetyl, trifluoroacetyl, and triphenylmethyl (trityl).

[00113] As used herein, the term “oxo” refers to a double bonded oxygen (=O) radical wherein the bond partner is a carbon atom. Such a radical can also be thought as a carbonyl group.

[00114] The term “peptide” or “polypeptide” refers to a sequence of at least two amino acids. In some embodiments, a peptide may contain no more than 80 amino acids, or no more than 35 amino acids, or no more than 10 amino acids.

[00115] The term “saccharide” refers to a sugar or other carbohydrate, especially a simple sugar. It includes both the alpha- and the beta-anomers. The saccharide can be a C₆-polyhydroxy compound, typically a C₆-pentahydroxy, and often a cyclic glycol. It includes the known simple sugars and their derivatives, as well as polysaccharides with two or more monosaccharide residues. The saccharide can include protecting groups on the hydroxyl groups. The hydroxyl groups of the saccharide can be replaced with one or more acetamido, halo or amino groups. Additionally, one or more of the carbon atoms can be oxidized, for example to keto or carbonyl groups. Suitable saccharides include galactose, glucose, glucuronic acid and neuraminic acid.

[00116] As used herein, the term “sulfonyl” refers to an >S(O)₂ group.

[00117] A prefix attached to a multi-component substituent only applies to the first component it precedes. To illustrate, the term “cycloalkylalkyl” contains two components: alkyl and cycloalkyl. Thus, the C₁-C₆-prefix on C₁-C₆-cycloalkylalkyl means that the alkyl component of the cycloalkylalkyl contains from 1 to 6 carbon atoms; the C₁-C₆-prefix does not describe the cycloalkyl component. To illustrate further, the term “halo-C₁-C₆-alkyl” refers to halomethyl, haloethyl, halopropyl, halobutyl, halopentyl, and haloethyl. To illustrate further, the prefix “halo” on haloalkoxyalkyl indicates that only the alkoxy component of the alkoxyalkyl

substituent is substituted with one or more halogen radicals. If the halogen substitution may only occur on the alkyl component, the substituent would instead be described as “alkoxyhaloalkyl.”

[00118] A substituent is “substitutable” if it comprises at least one carbon or nitrogen atom that is bonded to one or more hydrogen atoms. Thus, for example, hydrogen, halogen, and cyano do not fall within this definition. In addition, a sulfur atom in a heterocyclyl containing such atom is substitutable with one or two oxo substituents.

[00119] If a substituent is described as being “substituted”, a non-hydrogen radical is in the place of hydrogen radical on a carbon or nitrogen of the substituent. Thus, for example, a substituted alkyl substituent is an alkyl substituent in which at least one non-hydrogen radical is in the place of a hydrogen radical on the alkyl substituent. To illustrate, monofluoroalkyl is alkyl substituted with a fluoro radical, and difluoroalkyl is alkyl substituted with two fluoro radicals. It should be recognized that if there is more than one substitution on a substituent, each non-hydrogen radical may be identical or different (unless otherwise stated).


[00120] When a substituent is referred to as “unsubstituted” or not referred to as “substituted” or “optionally substituted”, it means that the substituent does not have any substituents. If a substituent is described as being “optionally substituted”, the substituent may be either (1) not substituted or (2) substituted. If a substituent is described as being optionally substituted with up to a particular number of non-hydrogen radicals, that substituent may be either (1) not substituted; or (2) substituted by up to that particular number of non-hydrogen radicals or by up to the maximum number of substitutable positions on the substituent, whichever is less. Thus, for example, if a substituent is described as a heteroaryl optionally substituted with up to 3 non-hydrogen radicals, then any heteroaryl with less than 3 substitutable positions would be optionally substituted by up to only as many non-hydrogen radicals as the heteroaryl has substitutable positions. To illustrate, tetrazolyl (which has only one substitutable position) would be optionally substituted with up to one non-hydrogen radical. To illustrate further, if an amino nitrogen is described as being optionally substituted with up to 2 non-hydrogen radicals, then a primary amino nitrogen will be optionally substituted with up to 2 non-hydrogen radicals, whereas a secondary amino nitrogen will be optionally substituted with up to only 1 non-hydrogen radical.

[00121] If substituents are described as being “independently selected” from a group, each substituent is selected independent of the other. Each substituent, therefore, may be identical to or different from the other substituent(s).

2. Labeling Reagents

[00122] Disclosed are compounds that may be labeling reagents. In certain embodiments, the compounds of formula (I) are labeling reagents. The labeling reagents can be used for cell uptake measurements.

[00123] The labeling reagents of the disclosure are compounds of formula (I), or a salt thereof,



(I),

wherein

A is a reporter moiety;

R¹⁴ is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R⁹ and R¹⁰ are independently selected from alkyl;

R¹¹, R¹² and R¹³ are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R¹¹ and R¹² can form a fused phenyl ring;

X is O;

L is $-(\text{CH}_2)_m\text{C}(\text{R}^{17})_2(\text{CH}_2)_n\text{-Y-C(O)-}$;

R¹⁷ is independently H, alkyl or both R¹⁷ together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;

n is an integer from 0-2;

Y is O or NR¹⁵;

R¹⁵ is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl-N(R²³)C(O)R²⁴, -alkyl-SO₃R²⁵, -alkyl-SO₂N(R²⁶)(R²⁷), -alkyl-COR²⁸, -alkyl-

CO₂R²⁹, -alkyl-OC(O)R²⁹, -alkyl-OC(O)N(R³⁰)(R³¹), -alkyl-CON(R³⁰)(R³¹), or polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with one or more suitable substituents; and

R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00124] In certain embodiments, R¹⁴ is H, C₁-C₃₀-alkyl, hydroxy-C₁-C₃₀-alkyl, C₁-C₃₀-alkoxy, carboxy-C₁-C₃₀-alkyl, or amido-C₁-C₃₀-alkyl; R⁹ and R¹⁰ are independently selected from C₁-C₄-alkyl; R¹¹, R¹² and R¹³ are independently selected from H, C₁-C₄-alkyl, C₁-C₄-alkoxy, bromo, chloro or amino, or R¹¹ and R¹² can form a fused phenyl ring; X is O; L is -(CH₂)_mC(R¹⁷)₂(CH₂)_n-Y-C(O)-; R¹⁷ is independently H, C₁-C₄-alkyl or both R¹⁷ together can form an alkyl ring having from 3-7 carbons; m is an integer from 0-2; n is an integer from 0-2; Y is O or NR¹⁵; R¹⁵ is H, C₁-C₃₀-alkyl, hydroxy-C₁-C₃₀-alkyl, azido-C₁-C₃₀-alkyl, cyano-C₁-C₃₀-alkyl, halo-C₁-C₃₀-alkyl, C₂-C₃₀-alkenyl, C₂-C₃₀-alkynyl, -C₁-C₃₀-alkyl-N(R²³)C(O)R²⁴, -C₁-C₃₀-alkyl-SO₃R²⁵, -C₁-C₃₀-alkyl-SO₂N(R²⁶)(R²⁷), -C₁-C₃₀-alkyl-COR²⁸, -C₁-C₃₀-alkyl-CO₂R²⁹, -C₁-C₃₀-alkyl-OC(O)R²⁹, -C₁-C₃₀-alkyl-OC(O)N(R³⁰)(R³¹), -C₁-C₃₀-alkyl-CON(R³⁰)(R³¹), or polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with one or more suitable substituents; and R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, C₁-C₄-alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00125] In certain embodiments, R¹⁴ is H, C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, -C₁-C₆-alkyl-CO₂H, or -C₁-C₆-alkyl-amide.

[00126] In certain embodiments, R¹⁴ is methyl.

[00127] In certain embodiments, R¹⁴ is carboxyalkyl. In certain embodiments, R¹⁴ is -C₂-C₃₀-alkyl-CO₂H. In certain embodiments, R¹⁴ is -CH₂CO₂H; -(CH₂)₂CO₂H; -(CH₂)₃CO₂H; -(CH₂)₄CO₂H; -(CH₂)₅CO₂H; -(CH₂)₆CO₂H; -(CH₂)₇CO₂H; -(CH₂)₈CO₂H; -(CH₂)₉CO₂H; -(CH₂)₁₀CO₂H; -(CH₂)₁₁CO₂H; -(CH₂)₁₂CO₂H; -(CH₂)₁₃CO₂H; -(CH₂)₁₄CO₂H; -(CH₂)₁₅CO₂H; -(CH₂)₁₆CO₂H; -(CH₂)₁₇CO₂H; -(CH₂)₁₈CO₂H; -(CH₂)₁₉CO₂H; -(CH₂)₂₀CO₂H; -(CH₂)₂₁CO₂H; -(CH₂)₂₂CO₂H; -(CH₂)₂₃CO₂H; -(CH₂)₂₄CO₂H; -(CH₂)₂₅CO₂H; -(CH₂)₂₆CO₂H; -(CH₂)₂₇CO₂H; -(CH₂)₂₈CO₂H; -(CH₂)₂₉CO₂H; or -(CH₂)₃₀CO₂H. In certain embodiments, R¹⁴ is -(CH₂)₁₅CO₂H.

[00128] In certain embodiments, R⁹, R¹⁰, R¹¹, R¹², R¹³, and R¹⁴ are each methyl.

[00129] In certain embodiments, in the variable L, m is 1, R¹⁷ is hydrogen at each occurrence, and n is 0. In certain embodiments, in the variable L, m is 1, R¹⁷ is hydrogen at each occurrence, n is 0, and Y is NR¹⁵. In certain embodiments, L is -CH₂CH₂-NR¹⁵-C(O)-.


[00130] In certain embodiments, A is a bioluminescent reporter moiety. In certain embodiments, A is a luciferin, a luciferin derivative or analog, a preluciferin or analog, coelenterazine or a coelenterazine derivative or analog. In some embodiments, A is luciferin, pro-luciferin, aminoluciferin, quionolyl-luciferin, naphthyl luciferin, chloroluciferin, coelenterazine, furimazine, coelenterazine-n, coelenterazine-f, coelenterazine-h, coelenterazine-hcp, coelenterazine-cp, coelenterazine-c, coelenterazine-e, coelenterazine-fcp, bis-deoxycoelenterazine (“coelenterazine-hh”), coelenterazine-i, coelenterazine-icp, coelenterazine-v, and 2-methyl coelenterazine, in addition to those disclosed in WO 2003/040100, U.S. Patent Publication No. 20080248511, and U.S. Patent Publication No. US 20120117667, the disclosures of which are incorporated by reference herein.

[00131] In certain embodiments, A is a fluorescent reporter moiety. In certain embodiments, A is a coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthene, or carbopyronine. In some embodiments, A is rhodamine 123, rhodamine X, Alexa dyes (e.g., Alexa Fluor-350, -430, -488, -and -660), DyLight 594, , sulforhodamine B, sulforhodamine 101, sulfonyl chloride derivative of sulforhodamine 101 (Texas Red), fluorescein, 6-carboxyfluorescein (6-FAM), 5-carboxyfluorescein (5-Fam), 5- or 6-carboxy-4,7,2',7'-tetrachlorofluorescein (TET), 5- or 6-carboxy-4'5'2'4'5'7' hexachlorofluorescein (HEX), 5' or 6'-carboxy-4', 5'-dichloro-2',7'-dimethoxyfluorescein (JOE), 6-JOE, 5-carboxy-2',4',5',7'-tetrachlorofluorescein (ZOE) rhodol, fluorescein isothiocyanate, coumarin, 7-amino-4-methylcoumarin, aminocoumarin, hydroxycoumarin,, silyl xanthene, or carbopyronine. In certain embodiments, A is a bicyclic aryl or heteroaryl, each of which are independently unsubstituted or substituted with one or more suitable substituents. In certain embodiments, A is a bicyclic aryl or heteroaryl, each of which are independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, cyanoalkyl, alkoxy, haloalkoxy, cyano, hydroxy, amino, alkylamino, di(alkyl)amino, aryl, heteroaryl, heterocyclyl, cycloalkyl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and cycloalkylalkyl, wherein the aryl, heteroaryl, heterocyclyl, cycloalkyl, aryl of the arylalkyl, heteroaryl of the

heteroarylalkyl, heterocyclyl of the heterocyclylalkyl, and cycloalkyl of the cycloalkylalkyl are each independently unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, cyanoalkyl, alkoxy, haloalkoxy, cyano, hydroxy, amino, alkylamino, and di(alkyl)amino.


[00132] In certain embodiments, A is a luciferin, luciferin derivative or analog (e.g., luciferin ester), preluciferin, preluciferin derivative or analog, cyanobenzothiazole, coelenterazine, a coelenterazine derivative (e.g., furimazine) or analog, or a fluorophore.

[00133] In certain embodiments, A is selected from the group consisting of:




wherein R^1 is H, C_1 - C_4 -alkyl, hydroxy- C_1 - C_4 -alkyl, C_3 - C_7 -cycloalkyl, aryl, benzyl or substituted benzyl, heterocyclyl, heteroaryl, or $-(CH_2)_q-P(Ph)_3$, wherein q is an integer selected from 1, 2, 3, 4, 5, and 6; R^2 is $-CH_2$ -aryl or $-CH_2$ -heteroaryl; R^4 is hydrogen, halogen, methyl, or trifluoromethyl; R^5 is hydrogen, halogen, methyl, or trifluoromethyl; R^7 is aryl (e.g., phenyl), substituted aryl (e.g., 4-hydroxyphenyl), $-CH_2$ -aryl, or $-CH_2$ -heteroaryl; and R^8 is $-CH_2$ -aryl or $-CH_2$ -heteroaryl. In certain embodiments, the $-CH_2$ -aryl is benzyl. In certain embodiments, the $-CH_2$ -heteroaryl is furylmethyl.


[00134] In certain embodiments, A is selected from the group consisting of:




coelenterazine HH ;




coelenterazine H ;




coelenterazine ;




Furimazine ;




Furimazine H ;



preluciferin ;



luciferin ; and



luciferin ester ;

wherein R¹ is C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, aryl, benzyl or substituted benzyl, heterocyclyl, heteroaryl, or -(CH₂)_q-P(Ph)₃, wherein q is an integer selected from 1, 2, 3, 4, 5, and 6.


[00135] In certain embodiments, R¹⁵ is H, C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, C₁-C₆-alkoxy, carboxy-C₁-C₆-alkyl, amido-C₁-C₆-alkyl, or polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with one or more suitable substituents.

[00136] In certain embodiments, R¹⁵ is polyalkoxyalkyl having 1, 2, 3, 4, 5, or 6 repeating alkoxy units.

[00137] In certain embodiments, R¹⁵ is polyalkoxyalkyl, wherein the polyalkoxyalkyl is -(C₂-C₆-alkoxy)_x-C₁-C₆-alkyl, -(C₂-C₆-alkoxy)_x-C₂-C₆-haloalkyl, -(C₂-C₆-alkoxy)_x-hydroxy-C₂-C₆-alkyl, -(C₂-C₆-alkoxy)_x-amino-C₂-C₆-alkyl, -(C₂-C₆-alkoxy)_x-C₁-C₆-alkylamino-C₂-C₆-alkyl, -(C₂-C₆-alkoxy)_xdi(C₁-C₆-alkyl)amino-C₂-C₆-alkyl, -(C₂-C₆-alkoxy)_x-azido-C₂-C₆-alkyl, -(C₂-C₆-alkoxy)_x-cyano-C₁-C₆-alkyl, -(C₂-C₆-alkoxy)_x-C₂-C₆-alkenyl, -(C₂-C₆-alkoxy)_x-C₂-C₄-alkynyl, -(C₂-C₆-alkoxy)_x-N(R²³)C(O)R²⁴, -(C₂-C₆-alkoxy)_x-SO₃R²⁵, -(C₂-C₆-alkoxy)_x-C₂-C₆-alkyl-SO₂N(R²⁶)(R²⁷), -(C₂-C₆-alkoxy)_x-C₂-C₆-alkyl-COR²⁸, -(C₂-C₆-alkoxy)_x-C₂-C₆-alkyl-OC(O)R²⁹, -(C₂-C₆-alkoxy)_x-C₂-C₆-alkyl-OC(O)N(R³⁰)(R³¹), -(C₂-C₆-alkoxy)_x-C₂-C₆-alkyl-CO₂R²⁹, or -(C₂-C₆-alkoxy)_x-C₂-C₆-alkyl-CON(R³⁰)(R³¹), wherein R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, C₁-C₄-alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl; and x is an integer selected from 1 to 20.

[00138] In certain embodiments, R¹⁵ is polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, halo-C₁-C₆-alkyl, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, amino-C₁-C₆-alkyl, C₁-C₆-alkylamino-C₁-C₆-alkyl, di(C₁-C₆-alkyl)amino-C₁-C₆-alkyl, azido-C₁-C₆-alkyl, cyano-C₁-C₆-alkyl, C₁-C₆-alkoxy, halo-C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₆-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkynyl, -N(R²³)C(O)R²⁴, -SO₃R²⁵, -SO₂N(R²⁶)(R²⁷), -COR²⁸, -CO₂R²⁹, and -CON(R³⁰)(R³¹), wherein R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, C₁-C₄-alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00139] In certain embodiments, R¹⁵ is a substituted polyalkoxyalkyl of formula:




wherein

R^{40} is selected from the group consisting of hydrogen, C_1 - C_4 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein the aryl, heteroaryl, heterocyclyl, and cycloalkyl are each independently unsubstituted or substituted with one or more suitable substituents; and

x is an integer selected from 1 to 20.

[00140] In certain embodiments, R^{40} is hydrogen.

[00141] In certain embodiments, R^{40} is a 5- or 6-membered heterocyclyl, having 1, 2, or 3 heteroatoms independently selected from the group consisting of N, O, and S, wherein said heterocyclyl is unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, cyanoalkyl, alkoxy, haloalkoxy, cyano, hydroxy, amino, alkylamino, di(alkyl)amino. In certain embodiments, R^{40} is




[00142] In certain embodiments, $-OR^{40}$ is defined as a leaving group.

[00143] In certain embodiments, x is 2, 3, or 4.


[00144] In certain embodiments, R^{15} is a carboxyalkyl selected from the group consisting of: $-CH_2CO_2H$; $-(CH_2)_2CO_2H$; $-(CH_2)_3CO_2H$; $-(CH_2)_4CO_2H$; $-(CH_2)_5CO_2H$; $-(CH_2)_6CO_2H$; $-(CH_2)_7CO_2H$; $-(CH_2)_8CO_2H$; $-(CH_2)_9CO_2H$; $-(CH_2)_{10}CO_2H$; $-(CH_2)_{11}CO_2H$; $-(CH_2)_{12}CO_2H$; $-(CH_2)_{13}CO_2H$; $-(CH_2)_{14}CO_2H$; $-(CH_2)_{15}CO_2H$; $-(CH_2)_{16}CO_2H$; $-(CH_2)_{17}CO_2H$; $-(CH_2)_{18}CO_2H$; $-(CH_2)_{19}CO_2H$; $-(CH_2)_{20}CO_2H$; $-(CH_2)_{21}CO_2H$; $-(CH_2)_{22}CO_2H$; $-(CH_2)_{23}CO_2H$; $-(CH_2)_{24}CO_2H$; $-(CH_2)_{25}CO_2H$; $-(CH_2)_{26}CO_2H$; $-(CH_2)_{27}CO_2H$; $-(CH_2)_{28}CO_2H$; $-(CH_2)_{29}CO_2H$; and $-(CH_2)_{30}CO_2H$. In certain embodiments, R^{15} is $-(CH_2)_{16}CO_2H$.

[00145] In certain embodiments, R^{15} is




each x is independently 0 to 20. In certain embodiments, each x is independently 0, 1, 2, 3, or 4.

[00146] In certain embodiments, compounds of formula (I) have formula (I-i), or a salt thereof,




(I-i),

wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{17} , m, n, and A are as defined above. In certain embodiments, A is selected from the group consisting of:



wherein R^{200} is phenyl, 4-hydroxyphenyl, or furyl; and R^{215} is hydrogen or hydroxy.


[00147] In certain embodiments, compounds of formula (I) have formula (I-ii), or a salt thereof,



(I-ii),

wherein R^{200} is phenyl, 4-hydroxyphenyl, or furyl; R^{215} is hydrogen or hydroxy; and R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , and R^{15} are as defined above.


[00148] In certain embodiments, compounds of formula (I) have formula (I-iv), or a salt thereof,



(I-iv),

wherein R²⁰⁰ is phenyl, 4-hydroxyphenyl, or furyl; R²¹⁵ is hydrogen or hydroxy; and R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R⁴⁰, and x are as defined above.


[00149] In certain embodiments, compounds of formula (I) have formula (I-iii), or a salt thereof,



(I-iii),

wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , and R^{15} are as defined above.


[00150] In certain embodiments, compounds of the invention have formula (I-v), or a salt thereof,



(I-v),


wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{40} , and x are as defined above.

[00151] In certain embodiments, compounds of formula (I) have formula (I-vi), or a salt thereof,




(I-vi),

wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{40} , x , and A are as defined above. In certain embodiments, A is selected from the group consisting of:



wherein R^{200} is phenyl, 4-hydroxyphenyl, or furyl; and R^{215} is hydrogen or hydroxy.


[00152] In certain embodiments, compounds of formula (I) have formula (I-vii), or a salt thereof,



(I-vii),

wherein z is an integer selected from 1 to 30; and R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, and A are as defined above. In certain embodiments, z is 16.


[00153] In certain embodiments, compounds of formula (I) have formula (I-viii), or a salt thereof,




(I-viii),

wherein y and z are each an integer independently selected from 1 to 30; and R⁹, R¹⁰, R¹¹, R¹², R¹³, and A are as defined above. In certain embodiments, y is 15 and z is 16.


[00154] In certain embodiments, the compound of formula (I) is selected from the group consisting of:




5463: 16-(((8-benzyl-2-(furan-2-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3-yl)oxy)carbonyl)-19,22-dimethyl-20-oxo-22-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-4,7,10,13-tetraoxa-16,19-diazatricosan-1-oic acid;




5470: 13-(((2,8-dibenzyl-6-phenylimidazo[1,2-a]pyrazin-3-yl)oxy)carbonyl)-16,19-dimethyl-17-oxo-19-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-4,7,10-trioxa-13,16-diazaicosan-1-oic acid;




5471: 10-(((2,8-dibenzyl-6-phenylimidazo[1,2-a]pyrazin-3-yl)oxy)carbonyl)-13,16-dimethyl-14-oxo-16-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-4,7-dioxa-10,13-diazaheptadecan-1-oic acid;




5508: 2,5-dioxopyrrolidin-1-yl 10-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)-13,16-dimethyl-14-oxo-16-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-4,7-dioxa-10,13-diazaheptadecan-1-oate;




(S)-2-(6-(((2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)(2-(2-(3-((2,5-dioxopyrrolidin-1-yl)oxy)-3-oxopropoxy)ethoxy)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




17-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)heptadecanoic acid;




5915: 2-cyanobenzo[d]thiazol-6-yl (2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)(2-(2-(3-((2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl)amino)-3-oxopropoxy)ethoxy)ethyl)carbamate;



5916: 2-cyanobenzo[d]thiazol-6-yl (2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)(15-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)-12-oxo-3,6,9-trioxa-13-azapentadecyl)carbamate;



5917: 2-cyanobenzo[d]thiazol-6-yl (2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)(4-((2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl)amino)-4-oxobutyl)carbamate; and



5918: 2-cyanobenzo[d]thiazol-6-yl (2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)(6-((2-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)ethyl)amino)-6-oxohexyl)carbamate; or a salt thereof.

3. Agents

[00155] The disclosed compounds may be used as labels that allow for the detection of an agent taken up by cells, tissues or organs. An agent may include a variety of compositions of matter. The disclosed compounds can be used to label a broad range of molecules, including but not limited to, biological agents or biomolecules such as polypeptides, polypeptide-based toxins, amino acids, nucleotides, polynucleotides including nucleic acids such as DNA and RNA, lipids, sugars, carbohydrates, detergents, and enzyme substrates, and any combination thereof. The disclosed compounds can also be used to label biological molecules, including but not limited to, antibodies, nanobodies, haptens, small molecules, drugs, drug compounds, ion-complexing agents, such as metal chelators, microparticles, synthetic or natural polymers, cells, viruses, other fluorescent molecules, such as FAM diamine, or surfaces, and any combination thereof. The drug, or derivative of a drug, may be a bile acid, such as ursodeoxycholic acid (UDCA), chenodeoxycholic acid (CDCA), lithocholic acid (LCA), and deoxycholic acid (DCA), a statin,

such as rosuvastatin, or an estradiol, such as estradiol glucuronide. The agent may be an antibody. The resulting labeled agents may be referred to as conjugates or tracers.

[00156] The polypeptide may be an antibody, an antibody-like moiety, such as a centyrin (Centyrex), or lipoprotein. The antibody may be a multispecific antibody, a human antibody, a humanized antibody (fully or partially humanized), an animal antibody such as, but not limited to, a bird (for example, a duck or a goose), a shark, a whale, and a mammal, including a non-primate (for example, a cow, a pig, a camel, a llama, a horse, a goat, a rabbit, a sheep, a hamster, a guinea pig, a cat, a dog, a rat, a mouse, etc.) or a non-human primate (for example, a monkey, a chimpanzee, etc.), a recombinant antibody, a chimeric antibody, a single-chain Fv (“scFv”), a single chain antibody, a single domain antibody, a Fab fragment, a F(ab') fragment, a F(ab')₂ fragment, a disulfide-linked Fv (“sdFv”), and an anti-idiotypic (“anti-Id”) antibody, a dual-domain antibody, a dual variable domain (DVD) or a triple variable domain (TVD) antibody (dual-variable domain immunoglobulin, and functionally active epitope-binding fragment of any of the above. In particular, an antibody includes an immunoglobulin molecule and an immunologically active fragment of an immunoglobulin molecule, namely, a molecule that contain an analyte-binding site. Immunoglobulin molecules can be of any type (for example, IgG, IgE, IgM, IgD, IgA, and IgY), class (for example, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2), or subclass. The antibody may be a therapeutic antibody such as Trastuzumab (Herceptin), Abciximab (ReoPro), Adalimumab (Humira), Alemtuzumab (Campath), Basiliximab (Simulect), Belimumab (Benlysta), Bevacizumab (Avastin), Brentuximab vedotin (Adcetris), Canakinumab (Ilaris), Cetuximab (Erbix), Certolizumab pegol (Cimzia), Daclizumab (Zenapax), Denosumab (Prolia, Xgeva), Eculizumab (Soliris), Efalizumab (Raptiva), Gemtuzumab (Mylotarg), Golimumab (Simponi), Ibritumomab tiuxetan (Zevalin), Infliximab (Remicade), Ipilimumab (MDX-101; Yervoy), Muromonab-CD3 (Orthoclone OKT3), Natalizumab (Tysabri), Ofatumumab (Arzerra), Omalizumab (Xolair), Palivizumab (Synagis), Panitumumab (Vectibix), Ranibizumab (Lucentis), Rituximab (Rituxan, Mabthera), Tocilizumab (or Atlizumab) (Actemra and RoActemra), and Tositumomab (Bexxar). Further example antibodies are also found in Table 1.

Table 1

Antigens	Antibodies
CD30	Ch IgG1
CD22	Hu IgG4
CD33	Hu IgG4
CD19	Hu IgG1
CD138	Ch IgG4
CD22	Hu IgG1
CD79b	Hu IgG1
CD74	Hu IgG1
HER2	Hu IgG1
GPNMB	Hu IgG2
PSMA	Hu IgG1
CD56	Hu IgG1
SLC44A4	Hu IgG2
CA6	Hu IgG1
CA-IX	Hu IgG1
Mesothelin	Hu IgG1
CD70	Hu IgG1
CD66e/CEACAM5	Hu IgG1
Nectin-4	Hu IgG1

Ch: chimeric; Hz: humanized; Hu: fully human; GPNMB: Glycoprotein NMB; PSMA: prostate specific membrane antigen.

[00157] The polypeptide may be Cholecystockinin (CCK), such as herring CCK-8, i.e., DYMGWMDF (SEQ ID NO: 1).

[00158] The lipoprotein may be high-density lipoprotein (HDL), low-density lipoprotein (LDL), intermediate-density lipoprotein, oxidized LDL, very low-density lipoprotein (VLDL), chylomicrons, chylomicron remnants, and non-natural lipoproteins.

[00159] Additionally, the compounds may be used to label non-biological agents, including but not limited to, solid supports/surfaces, nanoparticles, such as [60] fullerene, core-shell nanoparticles, liposome, dendrimer, and gold nanoparticles, and detergents. The detergent may be comprised within a liposome and/or solid-lipid nanoparticle (SLN) that may optionally be used to test drug delivering efficiency. The detergent may be a transfecting reagent, such as Lipofectamine® 2000 or Fugene 6.


a. Methods of Labeling an Agent

[00160] The disclosed compounds may be used to label an agent. In some aspects, the compounds can be conjugated with a nucleoside, nucleotide, or a polynucleotide. The


compounds of the invention may be conjugated with a nucleoside, nucleotide, or polynucleotide in any way known to one of ordinary skill in the art such as through a phosphoramidite, an activated ester or a reactive platinum complex. In certain embodiments, the labeling compounds may be conjugated to an agent using an activated carboxylic acid such as an NHS ester, pentafluorobenzene ester, an anhydride, an acetyl chloride; or using direct carboxylic acid and amine coupling reactions, or click chemistry, or maleimide, or activated carbonate, or phosphoramidite.

[00161] For example, as shown in Scheme 1, a labeling reagent can be attached to dU allylamine phosphoramidite and further incorporated in oligomers of interest by a traditional phosphoramidite chemistry. Alternatively, dU allylamine modified oligomers of interest can be labeled with the labeling reagent through an activated ester by post labeling. If those labeled oligos are primers of interest, they can also be used to amplify a sequence of interest through PCR.

Scheme 1




[00162] In certain embodiments, a labeled agent may have formula (II), or a salt thereof,



(II),

wherein the linker is a bond or a carbon chain that covalently attaches the parent molecular moiety to the agent, wherein the carbon chain is optionally interrupted by one or more nitrogen atoms, oxygen atoms, carbonyl groups, (substituted) aromatic rings, or peptide bonds; the agent is a target of interest; and R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{17} , and A are as defined above.


[00163] In certain embodiments, the labeled agent of formula (II) may have formula (II-i), or a salt thereof,



(II-i),

wherein the linker, the agent, R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{17} , and A are as defined above.


[00164] In certain embodiments, the labeled agent of formula (II) may have formula (II-ii), or a salt thereof,



(II-ii),

wherein the agent, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, x, and A are as defined above.


[00165] In certain embodiments, the labeled agent of formula (II) may have formula (II-iii),



(II-iii),


wherein R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, and A are as defined above.

[00166] In certain embodiments, the labeled agent may have formula (III),



(II-iv),

wherein each R is independently selected from hydrogen, alkyl, alkenyl, and a group of formula (a):



(a),


wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , z , and A are as defined above; wherein at least one R group is formula (a).

[00167] In certain embodiments, the agent in the above formulas is selected from the group consisting of: a polypeptide, a polypeptide-based toxin, an amino acid, a nucleotide, a polynucleotide, including nucleic acids such as DNA and RNA, a lipid, a carbohydrate, and an enzyme substrate, or any combination thereof. In certain embodiments, the agent is a peptide, an antibody, a lipoprotein, a protein, a nucleotide, a sugar, a fatty acid, a detergent, or a lipid. In certain embodiments, the agent is an antibody. In certain embodiments, the agent in the above formulas is selected from the group consisting of: a therapeutic drug, a small molecule, and a nanoparticle, or any combination thereof.

[00168] In other aspects, the disclosed labeling reagents can be conjugated with an amino acid, an amino acid analog, or a polypeptide. In other aspects, the disclosed labeling reagents can be conjugated with a small molecule, e.g., a drug or drug compound. In some aspects, the conjugated small molecule can be used as a fluorescent tracer.

4. Labeled Agents

[00169] In certain embodiments, the compounds of the present disclosure are labeled agents. The labeled agents are compounds of formula (I), or a salt thereof,



(I)

wherein

A is a reporter moiety;

R¹⁴ is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R⁹ and R¹⁰ are independently selected from alkyl;

R¹¹, R¹² and R¹³ are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R¹¹ and R¹² can form a fused phenyl ring;

X is O;

L is $-(\text{CH}_2)_m\text{C}(\text{R}^{17})_2(\text{CH}_2)_n\text{-Y-C(O)-}$;

R¹⁷ is independently H, alkyl or both R¹⁷ together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;

n is an integer from 0-2;

Y is O or NR¹⁵;

R¹⁵ is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl-N(R²³)C(O)R²⁴, -alkyl-SO₃R²⁵, -alkyl-SO₂N(R²⁶)(R²⁷), -alkyl-COR²⁸, -alkyl-CO₂R²⁹, -alkyl-OC(O)R²⁹, -alkyl-OC(O)N(R³⁰)(R³¹), -alkyl-amide, or polyalkoxyalkyl, wherein

the polyalkoxyalkyl and -alkyl-amide are unsubstituted or substituted with one or more suitable substituents; and

R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00170] In certain embodiments, the agent being labeled is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

[00171] In certain embodiments, Y is NR^{15} .

[00172] In certain embodiments, R^{15} comprises the agent.

[00173] In certain embodiments, R^{15} is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl- $N(R^{23})C(O)R^{24}$, -alkyl- SO_3R^{25} , -alkyl- $SO_2N(R^{26})(R^{27})$, -alkyl- COR^{28} , -alkyl- CO_2R^{29} , -alkyl- $OC(O)R^{29}$, -alkyl- $OC(O)N(R^{30})(R^{31})$, -alkyl-amide, or polyalkoxyalkyl, wherein the polyalkoxyalkyl and -alkyl-amide are unsubstituted or substituted with one or more suitable substituents; and R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00174] In certain embodiments, R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is selected from the group consisting of $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-haloalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-hydroxyalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-aminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkylaminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{di(alkyl)aminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-azidoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-cyanoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkenyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkynyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-N}(R^{23})C(O)R^{24}$, $-(C_2-C_6\text{-alkoxy})_x\text{-SO}_3R^{25}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-SO}_2N(R^{26})(R^{27})$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-COR}^{28}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-OC}(O)R^{29}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-OC}(O)N(R^{30})(R^{31})$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CO}_2R^{29}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CON}(R^{32})(R^{33})$ and $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-(CO)-NR}^{34}\text{-}(CR^aR^b)_p\text{-NR}^{35}\text{(CO)-T}$, wherein

R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl;

x is an integer selected from 1 to 20;

R^{32} and R^{33} are each independently selected from hydrogen, alkyl, carboxy, a peptide, a drug, a biologically active moiety, and a dye;

R^{34} and R^{35} are each independently selected from hydrogen, and alkyl;
 R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;
 p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.


[00175] In certain embodiments, R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CON}(R^{32})(R^{33})$ or $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-(CO)-NR}^{34}\text{-(CR}^a\text{R}^b)_p\text{-NR}^{35}\text{(CO)-T}$;
 x is an integer selected from 1 to 20;


R^{32} and R^{33} are each independently selected from hydrogen, alkyl, carboxy, aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof;

R^{34} and R^{35} are each independently selected from hydrogen, and alkyl;
 R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;
 p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

[00176] In certain embodiments, R^{15} is





wherein

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof; and

x is an integer selected from 1 to 20.

[00177] In certain embodiments, x is 1 to 10, 1 to 8, 1 to 6, 1 to 5, 1 to 4, 1 to 3, or 1 to 2. In certain embodiments, x is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20.

[00178] In certain embodiments, R¹⁵ is -alkyl-amide, wherein the -alkyl-amide is -alkyl-CON(R³²)(R³³) or -alkyl-(CO)-NR³⁴-(CR^aR^b)_p-NR³⁵(CO)-T, wherein

R³² and R³³ are each independently selected from hydrogen, alkyl, carboxy, aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heterocycle, alkene, polyol, alkenylpolyol, a peptide, a drug, a derivative of a drug, a biologically active moiety, and a dye;


R³⁴ and R³⁵ are each independently selected from hydrogen, and alkyl;

R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;

p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

[00179] In certain embodiments, R¹⁵ is




wherein


T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof; and

x is 0 to 20.

[00180] In certain embodiments, x is 0 to 10, 0 to 8, 0 to 6, 0 to 5, 0 to 4, 0 to 3, or 0 to 2. In certain embodiments, x is 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20.

[00181] In certain embodiments, R¹⁵ is







wherein x is 1 to 20 and T is an antibody.

[00182] In certain embodiments, when T is a monoclonal antibody, examples of antibodies include those in Table 1.

[00183] In certain embodiments, R¹⁵ is




wherein T comprises a C₆₀ fullurene. For example, T may be



[00184] In certain embodiments, R^{15} is $-\text{alkyl-T}$, wherein T is a solid-liquid nanoparticle (SLN). SLNs mainly comprise lipids that are in solid phase at room temperature and surfactant for emulsification, the mean diameters of which range from 50nm to 1000 nm for colloid drug delivery application. Solid lipids that utilized in SLN include fatty acids, triglycerides, steroids and waxes. Several types of surfactants are commonly used as emulsifiers to stabilize lipid dispersion.

[00185] In certain embodiments, R^{15} is



wherein T is $-\text{NH-PEG}_{3000}$ -gold nanoparticle, a PAMAM dendrimer, an oligonucleotide, or a peptide-oligonucleotide conjugate. The oligonucleotide or peptide-oligonucleotide conjugate may be one from Table 2.

Table 2

Name	Sequence
Ant-20	NH2RQIKIWFQNRRMKWKKGGC(COOH)-S-S-5'CCA-TCC-CGA-CCT-CGC-GCT-CC-3'-NH-Quinone-label
Ant mismatch	NH2RQIKIWFQNRRMKWKKGGCCOOHSS5'CCA-TAC-CAA-CAT-CAC-GCT-CC-3'-NH-Quinone-label
Tat-20	NH2RKKRRQRRRPPQC(COOH)-S-S-5'CCA-TCCCGA-CCT-CGC-GCT-CC-3'-NH-Quinone-label
Tat 20mismatch	NH2RKKRRQRRRPPQC(COOH)-S-S-5'CCA-TAC-CAA-CAT-CAC-GCT-CC-3'-NH-Quinone-label
20	5'CAA-TCC-CGA-CCT-CGC-GCT-CC-3'-NH-Quinone-label

[00186] In certain embodiments, R^{14} is H, C_1 - C_{30} -alkyl, hydroxy- C_1 - C_{30} -alkyl, C_1 - C_{30} -alkoxy, carboxy- C_1 - C_{30} -alkyl, or amido- C_1 - C_{30} -alkyl; R^9 and R^{10} are independently selected from C_1 - C_4 -alkyl; R^{11} , R^{12} and R^{13} are independently selected from H, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, bromo, chloro or amino, or R^{11} and R^{12} can form a fused phenyl ring; X is O; L

is $-(\text{CH}_2)_m\text{C}(\text{R}^{17})_2(\text{CH}_2)_n\text{-Y-C(O)-}$; R^{17} is independently H, $\text{C}_1\text{-C}_4\text{-alkyl}$ or both R^{17} together can form an alkyl ring having from 3-7 carbons; m is an integer from 0-2; n is an integer from 0-2; Y is O or NR^{15} ; R^{15} is H, $\text{C}_1\text{-C}_{30}\text{-alkyl}$, hydroxy- $\text{C}_1\text{-C}_{30}\text{-alkyl}$, azido- $\text{C}_1\text{-C}_{30}\text{-alkyl}$, cyano- $\text{C}_1\text{-C}_{30}\text{-alkyl}$, halo- $\text{C}_1\text{-C}_{30}\text{-alkyl}$, $\text{C}_2\text{-C}_{30}\text{-alkenyl}$, $\text{C}_2\text{-C}_{30}\text{-alkynyl}$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-N}(\text{R}^{23})\text{C(O)}\text{R}^{24}$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-SO}_3\text{R}^{25}$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-SO}_2\text{N}(\text{R}^{26})(\text{R}^{27})$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-COR}^{28}$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-CO}_2\text{R}^{29}$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-OC(O)}\text{R}^{29}$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-OC(O)}\text{N}(\text{R}^{30})(\text{R}^{31})$, $-\text{C}_1\text{-C}_{30}\text{-alkyl-CON}(\text{R}^{30})(\text{R}^{31})$, or polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with one or more suitable substituents; and R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, $\text{C}_1\text{-C}_4\text{-alkyl}$, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00187] In certain embodiments, R^{14} is H, $\text{C}_1\text{-C}_6\text{-alkyl}$, hydroxy- $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-alkoxy}$, $-\text{C}_1\text{-C}_6\text{-alkyl-CO}_2\text{H}$, or $-\text{C}_1\text{-C}_6\text{-alkyl-amide}$.

[00188] In certain embodiments, R^{14} is methyl.

[00189] In certain embodiments, R^{14} is carboxyalkyl. In certain embodiments, R^{14} is $-\text{C}_2\text{-C}_{30}\text{-alkyl-CO}_2\text{H}$. In certain embodiments, R^{14}

is $-\text{CH}_2\text{CO}_2\text{H}$; $-(\text{CH}_2)_2\text{CO}_2\text{H}$; $-(\text{CH}_2)_3\text{CO}_2\text{H}$; $-(\text{CH}_2)_4\text{CO}_2\text{H}$; $-(\text{CH}_2)_5\text{CO}_2\text{H}$; $-(\text{CH}_2)_6\text{CO}_2\text{H}$; $-(\text{CH}_2)_7\text{CO}_2\text{H}$; $-(\text{CH}_2)_8\text{CO}_2\text{H}$; $-(\text{CH}_2)_9\text{CO}_2\text{H}$; $-(\text{CH}_2)_{10}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{11}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{12}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{13}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{14}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{15}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{16}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{17}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{18}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{19}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{20}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{21}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{22}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{23}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{24}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{25}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{26}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{27}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{28}\text{CO}_2\text{H}$; $-(\text{CH}_2)_{29}\text{CO}_2\text{H}$; or $-(\text{CH}_2)_{30}\text{CO}_2\text{H}$. In certain embodiments, R^{14} is $-(\text{CH}_2)_{15}\text{CO}_2\text{H}$.

[00190] In certain embodiments, R^9 , R^{10} , R^{11} , R^{12} , R^{13} , and R^{14} are each methyl.

[00191] In certain embodiments, in the variable L, m is 1, R^{17} is hydrogen at each occurrence, and n is 0. In certain embodiments, in the variable L, m is 1, R^{17} is hydrogen at each occurrence, n is 0, and Y is NR^{15} . In certain embodiments, L is $-\text{CH}_2\text{CH}_2\text{-NR}^{15}\text{-C(O)-}$.


[00192] In certain embodiments, A is a bioluminescent reporter moiety. In certain embodiments, A is a luciferin, a luciferin derivative or analog, a pre-luciferin or analog, coelenterazine or a coelenterazine derivative or analog. In some embodiments, A is luciferin, pro-luciferin, aminoluciferin, quionolyl-luciferin, naphthyl luciferin, chloroluciferin, coelenterazine, furimazine, coelenterazine-n, coelenterazine-f, coelenterazine-h, coelenterazine-hcp, coelenterazine-cp, coelenterazine-c, coelenterazine-e, coelenterazine-fcp, bis-deoxycoelenterazine (“coelenterazine-hh”), coelenterazine-i, coelenterazine-icp, coelenterazine-

v, and 2-methyl coelenterazine, in addition to those disclosed in WO 2003/040100, U.S. Patent Publication No. 20080248511, and U.S. Patent Publication No. US 20120117667, the disclosures of which are incorporated by reference herein.


[00193] In certain embodiments, A is a fluorescent reporter moiety. In certain embodiments, A is a coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthene, or carbopyronine. In some embodiments, A is rhodamine 123, rhodamine X, Alexa dyes (e.g., Alexa Fluor-350, -430, -488, -and -660), DyLight 594, , sulforhodamine B, sulforhodamine 101, sulfonyl chloride derivative of sulforhodamine 101 (Texas Red), fluorescein, 6-carboxyfluorescein (6-FAM), 5-carboxyfluorescein (5-Fam), 5- or 6-carboxy-4,7,2',7'-tetrachlorofluorescein (TET), 5- or 6-carboxy-4'5'2'4'5'7' hexachlorofluorescein (HEX), 5' or 6'-carboxy-4', 5'-dichloro-2,'7'-dimethoxyfluorescein (JOE), 6-JOE, 5-carboxy-2',4',5',7'-tetrachlorofluorescein (ZOE) rhodol, fluorescein isothiocyanate, coumarin, 7-amino-4-methylcoumarin, aminocoumarin, hydroxycoumarin,, silyl xanthene, or carbopyronine. In certain embodiments, A is a bicyclic aryl or heteroaryl, each of which are independently unsubstituted or substituted with one or more suitable substituents. In certain embodiments, A is a bicyclic aryl or heteroaryl, each of which are independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, cyanoalkyl, alkoxy, haloalkoxy, cyano, hydroxy, amino, alkylamino, di(alkyl)amino, aryl, heteroaryl, heterocyclyl, cycloalkyl, arylalkyl, heteroarylalkyl, heterocyclylalkyl, and cycloalkylalkyl, wherein the aryl, heteroaryl, heterocyclyl, cycloalkyl, aryl of the arylalkyl, heteroaryl of the heteroarylalkyl, heterocyclyl of the heterocyclylalkyl, and cycloalkyl of the cycloalkylalkyl are each independently unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, cyanoalkyl, alkoxy, haloalkoxy, cyano, hydroxy, amino, alkylamino, and di(alkyl)amino.

[00194] In certain embodiments, A is a luciferin, luciferin derivative or analog (e.g., luciferin ester), preluciferin, preluciferin derivative or analog, cyanobenzothiazole, coelenterazine, a coelenterazine derivative or analog, or a fluorophore.


[00195] In certain embodiments, A is selected from the group consisting of:



a coelenterazine ;




a preluciferin ; and




a luciferin ;

wherein R¹ is H, C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, aryl, benzyl or substituted benzyl, heterocyclyl, heteroaryl, or -(CH₂)_q-P(Ph)₃, wherein q is an integer selected from 1, 2, 3, 4, 5, and 6; R² is -CH₂-aryl or -CH₂-heteroaryl; R⁴ is hydrogen, halogen, methyl, or trifluoromethyl; R⁵ is hydrogen, halogen, methyl, or trifluoromethyl; R⁷ is aryl (e.g., phenyl), substituted aryl (e.g., 4-hydroxyphenyl), -CH₂-aryl, or -CH₂-heteroaryl; and R⁸ is -CH₂-aryl or -CH₂-heteroaryl. In certain embodiments, the -CH₂-aryl is benzyl. In certain embodiments, the -CH₂-heteroaryl is furylmethyl.



[00196] In certain embodiments, A is selected from the group consisting of:



coelenterazine HH ;




coelenterazine H ;




wherein R¹ is C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, aryl, benzyl or substituted benzyl, heterocyclyl, heteroaryl, or -(CH₂)_q-P(Ph)₃, wherein q is an integer selected from 1, 2, 3, 4, 5, and 6.

[00197] In certain embodiments, compounds of formula (I) have formula (I-i), or a salt thereof,




(I-i),

wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{17} , m , n , and A are as defined above. In certain embodiments, A is selected from the group consisting of:



wherein R^{200} is phenyl, 4-hydroxyphenyl, or furyl; and R^{215} is hydrogen or hydroxy.


[00198] In certain embodiments, compounds of formula (I) have formula (I-ii), or a salt thereof,



(I-ii),

wherein R^{200} is phenyl, 4-hydroxyphenyl, or furyl; R^{215} is hydrogen or hydroxy; and R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , and R^{15} are as defined above.


[00199] In certain embodiments, compounds of formula (I) have formula (I-iv), or a salt thereof,




(I-iii),

wherein R^9 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , and R^{15} are as defined above.


[00200] In certain embodiments, the compound of formula (I) is selected from the group consisting of:




5651: 2-(6-(((20R)-15-carboxy-20-((3R,5S,7S,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-9,17-dioxo-3,6-dioxa-10,16-diazahenicosyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5657: 2-(6-(((4-(((5-carboxy-5-((R)-4-((3R,5S,7R,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5658: 2-(6-(((4-(((5-carboxy-5-((R)-4-((3R,5S,7S,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5665: 2-(6-(((4-((5-carboxy-5-((R)-4-((3R,5S,7R,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5666: 2-(6-(((4-((5-carboxy-5-((R)-4-((3R,5S,7S,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5683: 2-cyanobenzo[d]thiazol-6-yl (6-((2-(3',6'-dihydroxy-3-oxo-3H-spiro[isobenzofuran-1,9'-xanthen]-5-ylcarboxamido)ethyl)amino)-6-oxohexyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamate;




5826: 2-(6-(((1-(3',6'-dihydroxy-3-oxo-3H-spiro[isobenzofuran-1,9'-xanthen]-5-yl)-1,6-dioxo-9,12-dioxa-2,5-diazatetradecan-14-yl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5682: 2-cyanobenzo[d]thiazol-6-yl (1-(3',6'-dihydroxy-3-oxo-3H-spiro[isobenzofuran-1,9'-xanthen]-5-yl)-1,6-dioxo-9,12-dioxa-2,5-diazatetradecan-14-yl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamate;




5647: 8-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)-23-((R)-4-((3R,5R,8R,9S,10S,13R,14S,17R)-3-hydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)-2,5-dimethyl-4,17-dioxo-2-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-11,14-dioxa-5,8,18-triazatetracosan-24-oic acid;




5668: 2-(6-((((20R)-15-carboxy-20-((3R,5R,8R,9S,10S,13R,14S,17R)-3-hydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-9,17-dioxo-3,6-dioxo-10,16-diazahenicosyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5625: 8-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)-23-((R)-4-((3R,5R,8R,9S,10S,12S,13R,14S,17R)-3,12-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)-2,5-dimethyl-4,17-dioxo-2-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-11,14-dioxo-5,8,18-triazatetracosan-24-oic acid;




5649: 2-(6-((((20R)-15-carboxy-20-((3R,5R,8R,9S,10S,12S,13R,14S,17R)-3,12-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-9,17-dioxo-3,6-dioxa-10,16-diazahenicosyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbonyloxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5659: 6-(6-((((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)hexanamido)-2-((R)-4-((3R,5R,8R,9S,10S,13R,14S,17R)-3-hydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5663: 2-(6-(((6-((5-carboxy-5-((R)-4-((3R,5R,8R,9S,10S,13R,14S,17R)-3-hydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-6-oxohexyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5652: 6-(4-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)butanamido)-2-((R)-4-((3R,5R,8R,9S,10S,13R,14S,17R)-3-hydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5669: 2-(6-(((4-((5-carboxy-5-((R)-4-((3R,5R,8R,9S,10S,13R,14S,17R)-3-hydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5653: 6-(4-((((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)butanamido)-2-((R)-4-((3R,5R,8R,9S,10S,12S,13R,14S,17R)-3,12-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5656: 2-(6-(((4-((5-carboxy-5-((R)-4-((3R,5R,8R,9S,10S,12S,13R,14S,17R)-3,12-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5660: 6-(6-((((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)hexanamido)-2-((R)-4-((3R,5R,8R,9S,10S,12S,13R,14S,17R)-3,12-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5664: 2-(6-(((6-((5-carboxy-5-((R)-4-((3R,5R,8R,9S,10S,12S,13R,14S,17R)-3,12-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)pentyl)amino)-6-oxohexyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5667: 8-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)-23-((R)-4-((3R,5S,7R,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)-2,5-dimethyl-4,17-dioxo-2-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-11,14-dioxa-5,8,18-triazatetracosan-24-oic acid;




5650: 2-(6-((((20R)-15-carboxy-20-((3R,5S,7R,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-9,17-dioxo-3,6-dioxa-10,16-diazahenicosyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5648: 8-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)-23-((R)-4-((3R,5S,7S,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)-2,5-dimethyl-4,17-dioxo-2-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-11,14-dioxa-5,8,18-triazatetracosan-24-oic acid;




5654: 6-(4-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)butanamido)-2-((R)-4-((3R,5S,7R,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5655: 6-(4-((((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)butanamido)-2-((R)-4-((3R,5S,7S,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5661: 6-(6-(((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)hexanamido)-2-((R)-4-((3R,5S,7R,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5662: 6-(6-((((2-cyanobenzo[d]thiazol-6-yl)oxy)carbonyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)amino)hexanamido)-2-((R)-4-((3R,5S,7S,8R,9S,10S,13R,14S,17R)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)pentanamido)hexanoic acid;




5684: 2-cyanobenzo[d]thiazol-6-yl (4-((2-(3',6'-dihydroxy-3-oxo-3H-spiro[isobenzofuran-1,9'-xanthene]-5-carboxamido)ethyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamate;




5824: 2-(6-(((4-((2-(3-carboxy-4-(3,6-dihydroxy-9H-xanthen-9-yl)benzamido)ethyl)amino)-4-oxobutyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamide)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)thiazole-4-carboxylic acid;




5825: 2-(6-(((6-((2-(3',6'-dihydroxy-3-oxo-3H-spiro[isobenzofuran-1,9'-xanthen]-5-ylcarboxamido)ethyl)amino)-6-oxohexyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5830: 2-(6-(((2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)((16R,18S,E)-20-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methylmethylsulfonamido)pyrimidin-5-yl)-16,18-dihydroxy-9,14-dioxo-3,6-dioxa-10,13-diazaicos-19-en-1-yl)carbonyloxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5831: 2-(6-(((2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)((19R,21S,E)-23-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methylmethylsulfonamido)pyrimidin-5-yl)-19,21-dihydroxy-12,17-dioxo-3,6,9-trioxa-13,16-diazatricos-22-en-1-yl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5827: 2-(6-((((S)-18-((3S,5R,7S,8S,9R,10R,13S,14R,17S)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-9,15-dioxo-3,6-dioxa-10,14-diazanonadecyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;



5828: 2-(6-((((S)-21-((3S,5R,7S,8S,9R,10R,13S,14R,17S)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-12,18-dioxo-3,6,9-trioxa-13,17-diazadocosyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5829: 2-(6-(((S)-24-((3S,5R,7S,8S,9R,10R,13S,14R,17S)-3,7-dihydroxy-10,13-dimethylhexadecahydro-1H-cyclopenta[a]phenanthren-17-yl)-15,21-dioxo-3,6,9,12-tetraoxa-16,20-diazapentacosyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;




5832: (3R,6R,9R,15R,18R,21R)-9-((1H-indol-3-yl)methyl)-3-(((R)-1-amino-1-oxo-3-phenylpropan-2-yl)carbamoyl)-21-(10-(((2-(4-carboxy-4,5-dihydrothiazol-2-yl)benzo[d]thiazol-

6-yl)oxy)carbonyl)-13,16-dimethyl-14-oxo-16-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)-4,7-dioxa-10,13-diazaheptadecanamido)-18-(4-hydroxybenzyl)-6,15-bis(2-(methylthio)ethyl)-5,8,11,14,17,20-hexaoxo-4,7,10,13,16,19-hexaazatricosane-1,23-dioic acid;




2-(6-(((2-(2-(3-((3-(((8R,9S,13S,14S,17S)-17-(((3S,4R,6R)-6-carboxy-3,4,5-trihydroxytetrahydro-2H-pyran-2-yl)oxy)-13-methyl-7,8,9,11,12,13,14,15,16,17-decahydro-6H-cyclopenta[a]phenanthren-3-yl)oxy)propyl)amino)-3-oxopropoxy)ethoxy)ethyl)(2-(N,3-dimethyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanamido)ethyl)carbamoyl)oxy)benzo[d]thiazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid;





5907


;



5908

;






;

or a salt thereof.

[00201] In certain embodiments, the labeled agent may be a labeled antibody. The labeled antibody may be a labeled monoclonal antibody (mAb) (see FIG. 14). The labeled antibody may be a Herceptin labeled with PBI-5508.

[00202] In certain embodiments, the labeled agent may be a labeled therapeutic drug. The labeled therapeutic drug may be a labeled bile acid. The labeled bile acid may be a labeled UDCA, such as PBI-5651, PBI-5658, PBI-5666, PBI-5648, PBI-5655, or PBI-5662, a labeled CDCA such as PBI-5657, PBI-5665, PBI-5667, PBI-5650, PBI-5654, PBI-5661, PBI-5827, PBI-5828, or PBI-5829, a labeled LCA, such as PBI-5647, PBI-5668, PBI-5659, PBI-5663, PBI-5652, or PBI-5669, or a labeled DCA such as PBI-5625, PBI-5649, PBI-5653, PBI-5656, PBI-5660, or PBI-5664. The labeled drug may be a labeled statin. The labeled statin may be a labeled rosuvastatin, such as PBI-5830 or PBI-5831. The labeled drug may be a labeled estradiol. The labeled estradiol may be a labeled estradiol glucuronide, such as



[00203] In certain embodiments, the labeled agent may be a solid-liquid nanoparticle. For example, the labeled agent may be a labeled gold nanoparticle (see e.g., FIG. 15). The labeled gold nanoparticle may be PBI-5907, PBI-5908, PBI-5909, PBI-5910, PBI-5911, or PBI-5912.

[00204] In certain embodiments, the labeled agent may be a labeled peptide. The peptide may be cholecystokinin (CCK), such as CCK-8. The labeled peptide may be PBI-5832.

[00205] In certain embodiments, the labeled agent may be a labeled fluorescent molecule. The labeled fluorescent molecule may be a labeled FAM diamine, such as PBI-5683, PBI-5826, PBI-5682, PBI-5684, PBI-5824, or PBI-5825.

[00206] In certain embodiments, the labeled agent may be a labeled PAMAM dendrimer (see e.g., FIG. 16). In certain embodiments, the labeled agent may be a labeled oligonucleotide or a labeled peptide-oligonucleotide conjugate. For example, the labeled agent may be a labeled oligonucleotide or a peptide-oligonucleotide conjugate for antisense inhibition of P-glycoprotein expression (see e.g., FIG. 17).

5. Methods

[00207] The compounds of the present invention provide an effective tool for covalently labeling agents for a wide variety of cell uptake applications. The labeled agents allow quantification of the uptake of a biomolecule(s) of interest in biological samples (i.e., cells, tissues, and organs), monitoring effectiveness for drug delivery, effectiveness of therapeutic treatments, profiling pharmacokinetics, drug-drug interactions and drug toxicity, monitoring

cellular metabolic status, such as fatty acids and glucose, and detecting homeostasis of cholesterol.

[00208] The cell uptake measurement is often a significant part of scientific research and development, drug design, screening and optimization, phylogenetic classification, genotyping individuals, parental and forensic identification, environmental studies, diagnosis, prognosis, and/or treatment of disease conditions.

a. Methods of Evaluating Cellular Uptake

[00209] In some aspects, provided are methods for evaluating cellular uptake of an agent. The methods may comprise contacting a sample comprising a cell with a labeled agent as detailed above and detecting light emission, whereby the detection of light emission indicates cellular uptake of the agent. The cellular uptake of the agent results in the reduction of the compound and the generation of a released reporter moiety.

(1) Fluorescent reporter moiety

[00210] In some embodiments, the reporter moiety may include a fluorescent reporter moiety. The fluorescent reporter moiety may include a fluorophore. Light emission is detected by exposing the sample to a wavelength of light and detecting the fluorescence generated by the released reported moiety. An increase in fluorescence or a change in fluorescence wavelength as compared to the fluorescence or fluorescence wavelength of a control sample indicates cellular uptake of the agent. The control sample may be sample medium only, without cells, a sample with cells but without experimental treatment, or a sample not contacted with labeled agent. Fluorescence may be detected inside or outside the cell. Non-reducible analogues of the disclosed quinone labeling moieties may be used as control labels.

(2) Bioluminescent reporter moiety

[00211] In some aspects, the reporter moiety may include a bioluminescent reporter moiety. The bioluminescent reporter moiety may include a prosubstrate for a luciferase. In some embodiments, the cell includes a luciferase. The luciferase may be expressed in the cell. Light emission is detected by detecting luminescence produced by the luciferase utilizing the released reporter moiety. The detection of any light emission may indicate the cellular uptake of the agent. Alternatively, luminescence of the sample may be compared to the luminescence of a control sample, wherein cellular uptake of the agent is indicated if the luminescence of the sample is higher than the luminescence of the control sample. The control sample may be a

sample that is not contacted with a labeled agent. Luminescence may be detected inside or outside the cell.

[00212] In some embodiments, the cell does not include or express a luciferase, and luciferase is added to the sample. Light emission is detected by detecting luminescence produced by the luciferase utilizing the released reporter moiety that may exit the cell or be present in a cell lysate. The detection of any light emission may indicate the cellular uptake of the agent. Alternatively, luminescence of the sample may be compared to the luminescence of a control sample, wherein cellular uptake of the agent is indicated if the luminescence of the sample is higher than the luminescence of the control sample. The control sample may be sample medium only, without cells, a sample with cells but without experimental treatment, or a sample not contacted with labeled agent. Luminescence may be detected inside or outside the cell. Non-reducible analogues of the disclosed quinone labeling moieties may be used as control labels.

(3) Sample

[00213] The labeled agents may be used with samples containing biological components. The sample may comprise cells, tissues, or organs *in vitro* or *in vivo*. The compounds are generally non-toxic to living cells and other biological components within the concentrations of use.

[00214] Cells may include eukaryotic cells, e.g., yeast, avian, plant, insect or mammalian cells, including but not limited to human, simian, murine, canine, bovine, equine, feline, ovine, caprine or swine cells, or prokaryotic cells, or cells from two or more different organisms, or cell lysates or supernatants thereof. The cells may not have been genetically modified via recombinant techniques (non-recombinant cells), or may be recombinant cells which are transiently transfected with recombinant DNA and/or the genome of which is stably augmented with a recombinant DNA, or which genome has been modified to disrupt a gene, e.g., disrupt a promoter, intron or open reading frame, or replace one DNA fragment with another. The recombinant DNA or replacement DNA fragment may encode a molecule to be detected by the methods of the invention, a moiety which alters the level or activity of the molecule to be detected, and/or a gene product unrelated to the molecule or moiety that alters the level or activity of the molecule. The cell may or may not express a luciferase.

(4) Contact

[00215] The labeled agents may be combined with the sample in a way that facilitates contact between the compound and the sample components of interest. Typically, the labeled agent or a solution containing the labeled agent is simply added to the sample.

[00216] The cell uptake levels for selected labeled agents can be monitored with/without treatments that permeabilize the plasma membrane, such as electroporation, shock treatments or high extracellular ATP. Alternatively, selected labeled agents can be monitored physically inserted into cells, e.g., by pressure microinjection, scrape loading, patch clamp methods, or phagocytosis.

[00217] An additional detection reagent typically produces a detectable response due to the presence of a specific cell component, intracellular substance, or cellular condition, according to methods generally known in the art. When the additional detection reagent has, or yields a product with, spectral properties that differ from those of the subject labeled agents, multi-color applications are possible. This is particularly useful where the additional detection reagent is a dye or dye conjugate having spectral properties that are detectably distinct from those of the labeled agents.

[00218] In certain embodiments, washing steps are unnecessary when using the disclosed labeling reagents and labeled agents.

(5) Light Detection

[00219] The labeled agents are generally utilized by combining a labeled agent as described above with a sample of interest comprising a cell under conditions selected to yield a detectable optical response or light output. Typically, a specified characteristic of the sample is determined by comparing the optical response with a standard or expected response. The sample may be illuminated at a wavelength selected to elicit the optical response. Alternatively, the light emission from the sample may be measured in a reading device that can measure the light output (luminescence) generated by the luciferase and bioluminescent reporter moiety, e.g., using a luminometer or photomultiplier. The optical response or light output may also be measured over time, for example in the same reaction chamber for a period of seconds, minutes, hours, etc.

[00220] A detectable optical response means a change in, or occurrence of, an optical signal that is detectable either by observation or instrumentally. Typically, the detectable response is a change in fluorescence or luminescence, such as a change in the intensity, excitation or emission

wavelength distribution of fluorescence or luminescence, fluorescence or luminescence lifetime, fluorescence or luminescence polarization, or a combination thereof. The degree and/or location of the signal, compared with a standard or expected response, indicates whether, and to what degree, the sample possesses a given characteristic.

[00221] At any time after or during contact with the labeled agent, the sample is illuminated with a wavelength of light selected to give a detectable optical response and observed with a means for detecting the optical response. Equipment that is useful for illuminating the compounds of the invention includes, but is not limited to, hand-held ultraviolet lamps, mercury arc lamps, xenon lamps, lasers and laser diodes. These illumination sources are optionally integrated into laser scanners, fluorescence microplate readers, standard or minifluorometers, or chromatographic detectors.

[00222] The optical response or light output may be optionally detected by visual inspection or by use of any of the following devices: CCD cameras, video cameras, photographic film, laser-scanning devices, fluorometers, photodiodes, quantum counters, epifluorescence microscopes, scanning microscopes, flow cytometers, fluorescence microplate readers, or by means for amplifying the signal such as photomultiplier tubes. Where the sample is examined using a flow cytometer, examination of the sample optionally includes sorting portions of the sample according to their fluorescence response.

6. Kits


[00223] One aspect of the invention is the formulation of kits that facilitate the practice of various assays using any of the compounds of the invention, as described above. The kits of the invention may comprise a labeling reagent, an agent, a labeled agent, or any combination thereof. The kit optionally further comprises one or more buffering agents, typically present as an aqueous solution. The kits of the invention optionally further comprise at least one component for detecting the released labeled reagent, such as a detection reagent. The detection reagent may contain a luciferase, as described above, and/or other reagents required to conduct the detection assay such as buffers, salts, enzymes, enzyme co-factors, such as D-cysteine, and the like buffers. For example, the detection reagent may be Luciferin Detection Reagent (Promega Corporation). The kits of the invention optionally further comprise a purification medium for purifying the resulting labeled agent, luminescence and/or fluorescence standards, enzymes,

enzyme inhibitors, organic solvent, constructs for expression of fusion proteins, e.g., fusion proteins comprising a luciferase protein fused to a protein or target of interest, fusion proteins, or instructions for carrying out an assay of the invention. In other embodiments, the kit also includes a genetically-modified cell or a vector for gene fusion, e.g., fusion comprising a luciferase protein fused to a protein or target of interest. Instructions for use optionally may be included.

7. Examples

[00224] The foregoing may be better understood by reference to the following examples, which are presented for purposes of illustration and are not intended to limit the scope of the invention.

EXAMPLE 1



[00225] **Synthesis of N-[(PEG)₄.COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine.** To the solution of N-*t*-BuOOC-(PEG)₄-amine (0.557 g, 1.73 mmol) in 20 ml of methanol, (N-methyl)-N-Boc acetaldehyde (0.3 g, 1.73 mmol) was added. The mixture was stirred at room temperature for 3 hours. NaBH₄ (0.196g, 5.2 mmol) was added to the mixture at 0°C, and the resultant mixture was stirred at 0°C for 1 hour and 1 hour at room temperature. The reaction was quenched by adding 5 ml of water. After removal of solvent, 5 ml of water was added, and the mixture was extracted three times with methylene chloride. The combined organic layer was dried over Na₂SO₄, and the product was purified by flash silica chromatography using heptane/ethyl acetate

to methylene chloride/methanol to give a yield of 64.5% (0.535 g). ¹H NMR (300 MHz, CD₂Cl₂) δ ppm: 3.5-3.8 (m, 18H, CH₂), 3.45 (br, 2H, CH₂), 2.98 (br, 2H, CH₂), 2.83 (s, 3H, NCH₃), 2.46 (t, 2H, COCH₂), 1.42 (s, 18H, CH₃); MS-ESI (m/e): 479.6 [M+H].

[00226] Synthesis of N-[(PEG)₄COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine


furimazine carbamate. To the mixture of furimazine (0.20 g, 0.526 mmol) and bis(pentafluorophenyl) dicarbonate (0.276 g, 0.630 mmol) in 10 ml of dry THF, TEA (0.106 mg, 1.05 mmol) was added at room temperature under argon. The mixture was stirred for 2-3 minutes, and N-[(PEG)₄COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene diamine (0.553 mg, 1.16 mmol) was added. The resulted mixture was stirred at room temperature for 30 minutes. The compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 25.8% (0.12 g). MS-ESI (m/e): 886.6 [M+H].

[00227] Synthesis of N-[(PEG)₄COOH]-(N'-methyl)-ethylenediamine furimazine

carbamate. N-[(PEG)₄COO-*t*-Bu]-N'-Boc-N-methyl-ethylenediamine furimazine carbamate (0.12 g, 0.136 mmol) and triisopropylsilane (50ul) were dissolved in 10 ml of methylene chloride and TFA (1:1 in volume), and the mixture was stirred at room temperature for 2 hours. After removal of the solvent, the residue was dried under high vacuum overnight, and the product was used directly in next step.

[00228] Synthesis of #5463. To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (107 mg, 0.428 mmol) and isobutyl chloroformate (58.4 mg, 0.427 mmol) in 10 ml dry THF, N-methyl morpholine (86.5 mg, 0.855 mmol) was added at 0°C. The resultant mixture was stirred 30 minutes at 0°C, and N'-methyl-N-[(PEG)₄COOH]ethylene-diamine] furimazine carbamate in 5 ml of CH₂Cl₂ was added, and the resultant mixture was stirred for 1 hour. The compound was directly purified with flash silica column chromatography using heptane and ethyl acetate as eluent to give the product in a yield of 76.6% (105 mg). MS-ESI (m/e): 962.5 [M+H].

EXAMPLE 2



[00229] Synthesis of N-[(PEG)₃-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine. To the solution of *t*-Bu-(PEG)₃-amine (1.01 g, 3.64 mmol) in 20 ml of methanol, (N-methyl)-N-Boc acetaldehyde (0.63 g, 3.64 mmol) was added. The mixture was stirred at room temperature for 3 hours. NaBH₄ (0.412g, 10.9 mmol) was added to the mixture at 0°C, and the resultant mixture was stirred at 0 °C for 1 hour and 1 hour at room temperature. The reaction was quenched by adding 5 ml of water. After removal of solvent, 5 ml of water was added, and the mixture was extracted three times with methylene chloride. The combined organic layer was dried over Na₂SO₄, and the product was purified by flash silica chromatography using heptane/ethyl acetate to methylene chloride/methanol to give a yield of 72% (1.14 g). ¹H NMR (300 MHz, CD₂Cl₂) δ ppm: 3.68 (t, 2H, OCH₂), 3.56-3.60 (br, 10H, CH₂), 3.53 (t, 2H, OCH₂), 3.29 (t, 2H, CH₂), 2.85 (s, 3H, NCH₃), 2.7-2.8 (m, 4H, NCH₂), 2.47 (t, 2H, COCH₂), 1.44 (s, 18H, CH₃); MS-ESI (m/e): 435.5 [M+H].


[00230] Synthesis of N-[(PEG)₃ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine coelenterazine-H, H- carbamate. To the mixture of coelenterazine H, H (0.60 g, 1.54 mmol) and bis(penta-fluorophenyl) dicarbonate (0.74 g, 1.69 mmol) in 30 ml of dry THF, TEA (0.311 g, 3.07 mmol) was added at room temperature under argon. The mixture was stirred for 2-3 minutes, and N-[(PEG)₃COO-*t*-Bu]-N'-Boc-N'-methyl ethylenediamine (1.34g, 3.07 mmol) was

added. The resulted mixture was stirred at room temperature for 30 minutes. The compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 70.3% (0.92 g). MS-ESI (m/e): 852.7 [M+H].

[00231] Synthesis of N-(PEG)₃COOH-N'-methyl-ethylenediamine coelenterazine H, H-carbamate. N-[(PEG)₃COO-*t*-Bu]-N'-Boc-N-methyl-ethylenediamine coelenterazine H, H-carbamate (0.92 g, 0.136 mmol) and triisopropylsilane (100ul) were dissolved in 30 ml of methylene chloride and TFA (1:1 in volume), and the mixture was stirred at room temperature for 2 hours. After removal of the solvent, the residue was dried under high vacuum overnight, and the product was used directly in next step.

[00232] Synthesis of #5470. To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (810 mg, 3.24 mmol) and isobutyl chloroformate (442 mg, 3.24 mmol) in 30 ml dry THF, N-methyl morpholine (655 mg, 6.48 mmol) was added at 0°C. The resultant mixture was stirred 30 minutes at 0°C, and N'-methyl-N-[(PEG)₃COOH]ethylenediamine coelenterazine H, H-carbamate (0.92 g) in 10 ml of CH₂Cl₂ was added, and the resultant mixture was stirred for 1 hour. The compound was directly purified with flash silica column chromatography using heptane and ethyl acetate as eluent to give the product in a yield of 61% (611 mg). MS-ESI (m/e): 928.7 [M+H]; HPLC 98.2% at 254 nm.

EXAMPLE 3



[00233] Synthesis of N-[(PEG)₂-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine. To the solution of N-(PEG)₂COO-*t*-Bu-amine (1.08 g, 4.62 mmol) in 50 ml of methanol, (N-methyl)-N-Boc acetaldehyde (0.8 g, 4.62 mmol) was added. The mixture was stirred at room temperature for 3 hours. NaBH₄ (0.524, 13.9 mmol) was added to the mixture at 0°C, and the resultant mixture was stirred at 0°C for 1 hour and 1 hour at room temperature. The reaction was quenched by adding 5 ml of water. After removal of solvent, 5 ml of water was added, and the mixture was extracted three times with methylene chloride. The combined organic layer was dried over Na₂SO₄, and the product was purified by flash silica chromatography using heptane/ethyl acetate to methylene chloride/methanol to give a yield of 77.6% (1.40 g). ¹H NMR (300 MHz, CD₂Cl₂) δ ppm: 3.67 (t, 2H, OCH₂), 3.56-3.60 (br, 6H, OCH₂), 3.29 (t, 2H, CH₂), 2.85 (s, 3H, NCH₃), 2.7-2.8 (m, 4H, NCH₂), 2.46 (t, 2H, COCH₂), 1.45 (s, 18H, CH₃); MS-ESI (m/e): 391.4 [M+H].


[00234] Synthesis of N-[(PEG)₂ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine coelenterazine-H, H- carbamate. To the mixture of coelenterazine H, H (0.60 g, 1.54 mmol) and bis(penta-fluorophenyl) dicarbonate (0.74 g, 1.69 mmol) in 30 ml of dry THF, TEA (0.311 g, 3.07 mmol) was added at room temperature under argon. The mixture was stirred for 2-3 minutes, and N-[(PEG)₂COO*t*Bu]-N'-BOC-N'-methyl ethylenediamine (1.20g, 3.07 mmol) was added. The resulted mixture was stirred at room temperature for 30 minutes. The compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 72.8% (0.92 g). MS-ESI (m/e): 808.7 [M+H].

[00235] Synthesis of N-[(PEG)₂COOH]-N'-methyl-ethylenediamine coelenterazine H, H-carbamate. N-[(PEG)₂COO-*t*-Bu]-N'-Boc-N-methyl-ethylenediamine coelenterazine H, H-carbamate (0.92 g, 0.136 mmol) and triisopropylsilane (100ul) were dissolved in 30 ml of methylene chloride and TFA (1:1 in volume), and the mixture was stirred at room temperature for 2 hours. After removal of the solvent, the residue was dried under high vacuum overnight, and the product was used directly in next step.

[00236] Synthesis of #5471. To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (855 mg, 3.42 mmol) and isobutyl chloroformate (466 mg, 3.42 mmol) in 40 ml dry THF, N-methyl morpholine (691 mg, 6.83 mmol) was added at 0°C. The resultant mixture was stirred 30 minutes at 0°C, and N'-methyl-N[(PEG)₂COOH]ethylene-diamine] coelenterazine H, H-carbamate (0.92 g) in 10 ml of CH₂Cl₂

was added and the resultant mixture was stirred for 1 hour. The compound was directly purified with flash silica column chromatography using heptane and ethyl acetate as eluent to give the product in a yield of 68.5% (690 mg). MS-ESI (m/e): 884.7 [M+H]; HPLC 92.2% at 254 nm.

EXAMPLE 4



[00237] Synthesis of N-[(PEG)₂ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine-2-cyanobenzothiazole-6-carbamate. To the mixture of 6-hydroxyl-2-cyanobenzothiazole (1.0 g, 5.68 mmol) and bis(penta-fluorophenyl) dicarbonate (2.68 g, 1.69 mmol) in 100 ml of dry THF, TEA (2.22 g, 5.68 mmol) was added at room temperature. The mixture was stirred for 10 minutes, and N-[(PEG)₂COOtBu]-N'-BOC-N'-methyl ethylenediamine (2.22g, 5.68 mmol) was added. The resultant mixture was stirred at room temperature for 30 minutes. The compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 100% (3.4 g). MS-ESI (m/e): 593.3 [M+H].


[00238] Synthesis of N-[(PEG)₂COOH]-N'-methyl-ethylenediamine-2-cyanobenzothiazole-6- carbamate. N-*t*-BuOOC-(PEG)₂-N'-Boc-N-methyl-ethylenediamine-2-cyanobenzothiazole-6- carbamate (2.0 g) and triisopropylsilane (100ul) were dissolved in 60 ml of methylene chloride and TFA (1:1 in volume), and the mixture was stirred at room temperature


for 4 hours. After removal of the solvent, the residue was dried under high vacuum overnight, and the product was used directly in next step.


[00239] Synthesis of N-[(PEG)₂COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate. To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (1.27 g, 5.07 mmol) and isobutyl chloroformate (691 mg, 5.07 mmol) in 50 ml dry THF was added N-methyl morpholine (932 mg, 9.21 mmol) at 0 °C. The resultant mixture was stirred 30 minutes at 0°C, and N'-methyl-N[(PEG)₂COOH]ethylenediamine]-2-cyanobenzothiazole-6-carbamate (1.07 g, 4.61 mmol) in 15 ml of CH₂Cl₂ was added and the resultant mixture was stirred for 1 hour. The compound was directly purified with flash silica column chromatography using heptane and ethyl acetate as eluent and further purified by HPLC using 0.1% of formic acid/acetonitrile as eluent. MS-ESI (m/e): 668.7 [M+H]; HPLC 92.6% at 330nm.

[00240] **Synthesis of #5508.** To the solution of N-[(PEG)₂COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate (210 mg, 0.314 mmol) and TSTU (473 mg, 1.57 mmol) in 20 ml acetonitrile and methylene chloride (1:1), DIPEA (325 mg, 2.51 mmol) was added at room temperature. The mixture was stirred for 30 minutes. 120 ml of methylene chloride was added, and the resultant mixture was washed three times with citric acid (30%) solution and twice with water. The organic layer was dried over Na₂SO₄. The compound was purified with heptane and ethyl acetate to give a yield of 87.3% (210 mg). MS-ESI (m/e): 593.3 [M+H]; HPLC 92.6% at 330nm. MS-ESI (m/e): 766.3 [M+H]; HPLC 98.2% at 254 nm.

EXAMPLE 5







[00241] Method 3: Synthesis of UDCA-lysCOOH. To a mixture of lysine (2.44 g, 7.20 mmol) and UDCA (1.13 g, 2.88 mmol), 120 mL methylene chloride and 4-dimethylaminopyridine (1.06 g, 8.64 mmol) was added. Lastly, *N,N'*-dicyclohexylcarbodiimide (1.78 g, 8.64 mmol) was added and stirred at room temperature for several days. The reaction mixture was then filtered, and the solvent was removed. The product was then purified by flash silica column chromatography using a heptane/ethyl acetate gradient to give a yield of 90% (1.76 g) of the intermediate with protected lysine. This compound was then dissolved in a mixture of 40 mL methylene chloride and 30 mL TFA together with 10% triisopropylsilane (TIPS), and the mixture was stirred at room temperature for 5 hours. The solvents were co-evaporated with toluene, and the product was purified by flash silica column chromatography using a gradient of methylene chloride/methanol to give UDCA-lysCOOH in a 67 % yield (901 mg). MS-ESI (m/e): 521.5 [M+H].

[00242] Synthesis of N-[(PEG)₂-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine. To the solution of N-(PEG)₂COO-*t*-Bu-amine (3.16 g, 13.54 mmol) in 100 ml of methanol, (N-methyl)-N-Boc acetaldehyde (2.35 g, 13.54 mmol) was added. The mixture was stirred at room temperature for 3 hours. NaBH₄ (1.54 g, 40.63 mmol) was added to the mixture at 0°C, and the resultant mixture was stirred at 0°C for 1 hour and 1 hour at room temperature. The reaction was quenched by adding 15 ml of water. After removal of solvent, 15 ml of water was added, and the mixture was extracted three times with methylene chloride. The combined organic layer was dried over Na₂SO₄, and the product was purified by flash silica chromatography using heptane/ethyl acetate to methylene chloride/methanol to give a yield of 67 % (3.56 g).

[00243] Synthesis of N-[(PEG)₂ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[*d*]thiazole-2-carbonitrile- carbamate. To the mixture of 2-cyano-6-hydroxybenzothiazole (0.803 g, 4.56 mmol) and bis(pentafluorophenyl) carbonate (1.98 g, 5.02

mmol) in 40 ml of dry THF, TEA (0.293 g, 2.89 mmol) was added at room temperature under nitrogen. The mixture was stirred for 1 hour, and N-*t*-butylCOO (PEG)₃-N'-BOC-N'-methyl ethylenediamine (3.56g, 9.12 mmol) in 18 mL dry THF was added. The resulting mixture was stirred at room temperature for over 30 minutes after which the solvent was removed. The product mixture was dissolved in methylene chloride and washed with sat K₂CO₃ solution three times and once with water, then the organic layer was dried over Na₂SO₄. The compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 70 % (1.90 g).

[00244] Synthesis of N-[(PEG)₂ COO-*t*-Bu]-N'-methyl-ethylene-diamine-2-cyanobenzothiazole-6-carbamate. N-[(PEG)₂ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[*d*]thiazole-2-carbonitrile- carbamate (1.90 g, 3.21 mmol) in 30 mL CH₂Cl₂ was dissolved over ice and then 26 mL (30 %) thioanisole was added. TFA (30mL) was slowly added over 10-15 min while on ice. The mixture was then stirred on ice for 20 min and then at room temperature for 5 hours. After the reaction, the solvent was removed, and the product was purified by flash column chromatography using a gradient of heptane/ethyl acetate followed by methylene chloride/methanol to give a yield of 57 % (0.797 g).

[00245] Synthesis of N-[(PEG)₂COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate. To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (1.37 g, 5.48 mmol) and isobutyl chloroformate (719 uL, 5.48 mmol) in 40 ml dry THF, N-methyl morpholine (1.53 mL, 10.96 mmol) was added at 0°C. The resultant mixture was stirred for 1 hour at 0°C, N-[(PEG)₂ COO-*t*-Bu]-N'-methyl-ethylene-diamine-2-cyanobenzothiazole-6-carbamate (0.797 g, 1.83 mmol) in 20 ml of dry THF added, and the resultant mixture was stirred for 2 hours. The reaction mixture was then acidified with acetic acid, dried down, then dissolved in methylene chloride and washed twice with water. The organic layer was dried over Na₂SO₄, and then the solvent was removed. The compound was directly purified with flash silica column using heptane and ethyl acetate as eluent, followed by 100% THF, to give the product in a yield of 48.8 % (0.596 g).


[00246] Synthesis of N-[(PEG)₂COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate NHS ester. To a solution of N-[(PEG)₂COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-

trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate (0.315 g, 0.471 mmol) in 70 mL acetonitrile, 709 mg (2.36 mmol) of TSTU and 660 μ L of DIPEA was added and then stirred for 1 hour at room temperature. The product mixture was diluted in a large volume of methylene chloride, and the product mixture washed twice with 30% citric acid solution and then once with water. The organic layer was dried with Na_2SO_4 , filtered, and the solvent was removed. The product was then purified by flash column chromatography using a heptane/ ethyl acetate gradient to give the product in 109% yield (394 mg).

[00247] Method 1: Synthesis of #5648. Dissolved NHS ester (100 mg, 0.130 mmol) in 2 mL DMF and added UDCA-lysCOOH synthesized by **Method 3** (~136 mg, 0.261 mmol), followed by 45 μ L of DIPEA and 3 more mL of DMF. The reaction was stirred at room temperature for 1 hour and then acidified with acetic acid and dried down. The product was purified by flash column chromatography using a gradient of heptane/ ethyl acetate, switched to methylene chloride/ methanol to give product in a yield of 40% (61 mg). MS-ESI (m/e): 1172.09 [M+H].

[00248] Method 2: Synthesis of #5651. Dissolved 12 mg of d-cysteine (0.077 mmol) in 1 mL of water and adjusted pH to 8-8.5 with TEA and then added 30 mg (0.025 mmol) of #5648 in 1 mL methanol and stirred for an hour at room temperature. The product was acidified with acetic acid and then extracted into methylene chloride. The solvent was removed, and the product was purified by flash column chromatography using a gradient of heptane/THF followed by methylene chloride/methanol to give the product in a yield of 70 % (22.8 mg). MS-ESI (m/e): 1276.93 [M+H].


EXAMPLE 6



[00249] **Synthesis of #5667.** Coupled the NHS ester in Example 9 and CDCA-lysCOOH (synthesized by **Method 3**) utilizing **Method 1** to give product in a yield of 25% (19 mg). MS-ESI (m/e): 1172.10 [M+H].

[00250] **Synthesis of #5650.** Synthesized the luciferin version of #5667 utilizing **Method 2** to give the product in a yield of 55 % (8.40 mg). MS-ESI (m/e): 1275.77 [M+H].


EXAMPLE 7



[00251] **Synthesis of #5647.** Method 1 was utilized with slight adjustments. NHS ester (100 mg, 0.130 mmol) was dissolved in 1 mL DMF. LCA-lysCOOH (synthesized by Method 3) (~131 mg, 0.261 mmol) was dissolved in 15 μ L of 50% TFA in DMF and then diluted with 4 mL DMF. The pH was increased to 5-6 with 23 μ L of DIPEA, and then the solution was added to the dissolved NHS ester, and 55 μ L of DIPEA was added. The reaction was stirred at room temperature overnight and then acidified with acetic acid and dried down. The product was extracted into ethyl acetate and then purified by flash column chromatography using a gradient of heptane/ethyl acetate, followed by methylene chloride/methanol to give product in a yield of 45% (68 mg). MS-ESI (m/e): 1155.94 [M+H].

[00252] **Synthesis of #5668.** The luciferin version of #5647 was synthesized using Method 2 to give the product in a yield of 50 % (16.20 mg). MS-ESI (m/e): 1260.27 [M+H].


EXAMPLE 8





[00253] **Synthesis of #5625.** Utilized **Method 1** to couple DCA-lysCOOH (synthesized by **Method 3**) to the NHS ester to give product in a yield of 15% (26.8 mg). MS-ESI (m/e): 1169.9 [M-H].

[00254] **Synthesis of #5649.** Synthesized the luciferin version of #5625 using **Method 2** to give the product in a yield of 69 % (12 mg). MS-ESI (m/e): 1176.79 [M+H].

EXAMPLE 9







[00255] Synthesis of 4-(((benzyloxy)carbonyl)amino)butanoic acid. γ -Amino butyric acid (2.90 g, 28.14 mmol) was dissolved in 15 % aqueous NaOH (100 mL) and dioxane (100 mL). After the mixture was cooled to 0°C, benzyl chloroformate (8 mL, 56.27 mmol) was added with vigorous stirring. The mixture was stirred for 3 days at room temperature. The solvent was removed, and the product mixture was dissolved in a small amount of water and acidified with acetic acid. The product was then extracted into methylene chloride (3 x 100 mL), and the organic layer was dried with sodium sulfate, filtered, and then dried down. The product was purified by flash column chromatography using a gradient of heptane/ethyl acetate to give the product in a 51 % yield (3.44 g).

[00256] Synthesis of *tert*-butyl 4-(((benzyloxy)carbonyl)amino)butanoate. To 4-(((benzyloxy)carbonyl)amino)butanoic acid (3.44 g, 14.50 mmol) in 100 mL methylene chloride, anhydrous *t*-butanol (4.2 mL, 43.50 mmol) was added, followed by DMAP (1.77 g, 14.50 mmol) and DCC (8.97 g, 43.50 mmol). The reaction mixture was stirred at room temperature for 2 days, after which the precipitate of DCU was filtered, and the solvent was removed. The product was purified by flash column chromatography using heptane/ethyl acetate to give a yield of 49 % (2.09 g).

[00257] Synthesis of *tert*-butyl 4-aminobutanoate. A mixture of *tert*-butyl 4-(((benzyloxy)carbonyl)amino)butanoate (2.09 g, 7.12 mmol) and 10% Pd/C (75.8 mg, 0.712 mmol) in 100 mL methanol was stirred at room temperature under hydrogen for 5 hours. The mixture was then filtered, and the solvent was removed to give a 93 % yield (1.06 g) of product.

[00258] Synthesis of N-[(CH₂)₃-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine. To the solution of *tert*-butyl 4-aminobutanoate (1.06g, 6.66 mmol) in 50 ml of methanol, (N-methyl)-N-BOC acetaldehyde (1.15 g, 6.66 mmol) was added. The mixture was stirred at room temperature for 3 hours. NaBH₄ (0.755 g, 19.97 mmol) was added to the mixture at 0°C, and the resultant mixture was stirred at 0°C for 1 hour and at room temperature for 1 hour. The reaction was quenched by adding 5 ml of water. After removal of solvent, 5 ml of water was added, and the mixture was extracted three times with methylene chloride. The combined organic layer was dried over Na₂SO₄, and the solvent was removed. The product was purified by flash column chromatography using heptane/ethyl acetate followed by methylene chloride/methanol to give a yield of 105 % (2.23 g).

[00259] Synthesis of N-[(CH₂)₃ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[d]thiazole-2-carbonitrile-carbamate. To the mixture of 2-cyano-6-hydroxybenzothiazole (0.587 g, 3.33 mmol) and bis(pentafluorophenyl) carbonate (1.44 g, 3.66 mmol) in 30 ml of dry THF, TEA (930 μL, 6.66 mmol) was added at room temperature under nitrogen. The mixture was stirred for 30 min minutes, and N-[(CH₂)₃-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine (2.11 g, 6.66 mmol) in 20 mL dry THF was added. The resulting mixture was stirred at room temperature for 2 hours. The reaction mixture was then dried down, and the product mixture was dissolved in methylene chloride and washed with saturated K₂CO₃ solution 3 times and once with water, then dried over Na₂SO₄. The solvent was removed, and the compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 86 % (1.5 g)

[00260] Synthesis of N-[(CH₂)₃ COO-*t*-Bu]-N'-methyl-ethylene-diamine-2-cyanobenzothiazole-6-carbamate. N-[(CH₂)₃ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[d]thiazole-2-carbonitrile-carbamate was dissolved in 15 mL methylene chloride over ice. Thioanisole (8 mL, 21%) was added, followed by the slow addition of 15 mL TFA over 10-15 min on ice. The mixture was stirred on ice for 30 min, then stirred at room temperature for 4 hours. The TFA was then co-evaporated with toluene, and the product was purified by flash column chromatography using heptane/ethyl acetate, followed by methylene chloride/ methanol to elute product the product in a 73 % yield (0.765 g).

[00261] Synthesis of N-[(CH₂)₃COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-

carbamate. To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (1.59 g, 6.33 mmol) and isobutyl chloroformate (831 μ L, 6.33 mmol) in 40 ml dry THF, N-methyl morpholine (1.76 mL, 12.67 mmol) was added at 0°C. The resulting mixture was stirred for 1 hour at 0°C, and then N-[(CH₂)₃ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[*d*]thiazole-2-carbonitrile-carbamate in 10 ml dry THF was added. The resulting mixture was stirred for 2 hours after which it was acidified with acetic acid then dried down. The product was then purified by flash column chromatography using heptane/ethyl acetate gradient followed by methylene chloride/methanol to give the product in a 142 % yield (1.79 g).

[00262] Synthesis of N-[(CH₂)₃COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate NHS ester (C4 NHS ester). TSTU (2.3 g, 7.64 mmol) was added to N-[(CH₂)₃COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6- carbamate (1.26 g, 2.12 mmol), followed by 70 mL of acetonitrile. DIPEA (2.95 mL, 16.95 mmol) was added, and the mixture was stirred for 1 hour at room temperature. The product mixture was then diluted in a large volume of methylene chloride and washed with 30% citric acid solution twice, then once with water. The organic layer was dried over Na₂SO₄, filtered, and the solvent was removed. The product was purified by flash column chromatography using heptane/ethyl acetate to give a 70 % yield (1.03 g).

[00263] Synthesis of #5654. C4 NHS ester was coupled to CDCA-lysineCOOH using **Method 1** to give the product in a 41% yield (65 mg). MS-ESI (m/e): 1098.49 [M+H].

[00264] Synthesis of #5657. The luciferin version of #5654 was prepared using **Method 2** to give a yield of 40 % (17.3 mg). MS-ESI (m/e): 1203.37 [M+H].

[00265] Synthesis of #5655. C4 NHS ester was coupled to UDCA-lysineCOOH using **Method 1** to give the product in a 49% yield (78 mg). MS-ESI (m/e): 1098.40 [M+H].

[00266] Synthesis of #5658. The luciferin version of #5655 was prepared using **Method 2** to give a yield of 69 % (28 mg). MS-ESI (m/e): 1203.14 [M+H].


[00267] Synthesis of #5652. C4 NHS ester was coupled to LCA-lysineCOOH using **Method 1** to give the product in a 78% yield (172 mg). MS-ESI (m/e): 1079.71 [M-H].


[00268] **Synthesis of #5669.** The luciferin version of #5652 was prepared using **Method 2** to give a yield of 24 % (6.5 mg). MS-ESI (m/e): 1086.30 [M+H].

[00269] **Synthesis of #5653.** C4 NHS ester was coupled to DCA-lysineCOOH using **Method 1** to give the product in a 71 % yield (112 mg). MS-ESI (m/e): 1095.59 [M-H].

[00270] **Synthesis of #5656.** The luciferin version of #5653 was prepared using **Method 2** to give a yield > 99 % yield (84 mg). MS-ESI (m/e): 1202.34 [M+H].

EXAMPLE 10





[00271] Synthesis of 6-(((benzyloxy)carbonyl)amino)hexanoic acid. 6-Aminohexanoic acid (4.61 g, 35.17 mmol) was dissolved in a mixture of 15% aqueous NaOH (100 mL) and dioxane (100 mL). After the mixture was cooled to 0°C, benzyl chloroformate (10 mL, 70.34 mmol) was added with vigorous stirring. The mixture was then stirred at room temperature for 2 days. The solvent was removed, the product mixture dissolved in water and acidified with acetic acid, and the product extracted into methylene chloride. The organic layer was dried with Na₂SO₄, filtered, and the solvent was removed. The product was purified by flash column chromatography using a gradient of heptane/ethyl acetate to give 81 % (7.62 g).

[00272] Synthesis of *tert*-butyl 6-(((benzyloxy)carbonyl)amino)hexanoate. 6-(((Benzyloxy)carbonyl)amino)hexanoic acid (7.62 g, 28.72 mmol) was dissolved in 200 mL methylene chloride, and anhydrous *t*-butanol (8.2 mL, 86.16 mmol) was added, followed by DMAP (3.51 g, 28.72 mmol). DCC (17.78 g, 86.16 mmol) was added, and the mixture was stirred at room temperature for 2 days. The mixture was filtered and then dried down, and the product was purified by flash column chromatography using heptane/ethyl acetate to give a yield of 63 % (5.86 g).

[00273] Synthesis of *tert*-butyl 6-aminohexanoate. To a mixture of *tert*-butyl 6-(((benzyloxy)carbonyl)amino)hexanoate (5.86 g, 18.23 mmol) and 10 % Pd/C (194 mg, 1.82 mmol), 250 mL of methanol was added. The mixture was stirred at room temperature for 5 hours

under a hydrogen atmosphere, after which it was filtered and the solvent was removed to give the product in 131 % yield (4.48 g).

[00274] Synthesis of N-[(CH₂)₅-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine. To the solution of *tert*-butyl 6-aminohexanoate (3.41 g, 18.21 mmol) in 80 mL of methanol, (N-methyl)-N-BOC acetaldehyde (3.15 g, 18.21 mmol) was added. The mixture was stirred at room temperature for 3 hours. NaBH₄ (2.07 g, 54.62 mmol) was added to the mixture at 0°C, and the resultant mixture was stirred at 0°C for 1 hour and at room temperature for 1 hour. The reaction was quenched by adding 10 mL of water. After removal of solvent, 30 mL of water was added, and the mixture was extracted three times with methylene chloride. The combined organic layer was dried over Na₂SO₄ and dried down. The product was purified by flash column chromatography using heptane/ethyl acetate followed by methylene chloride/methanol.

[00275] Synthesis of N-[(CH₂)₅ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[*d*]thiazole-2-carbonitrile-carbamate. To the mixture of 2-cyano-6-hydroxybenzothiazole (0.366 g, 2.08 mmol) and bis(pentafluorophenyl) carbonate (0.902 g, 2.29 mmol) in 20 ml of dry THF, TEA (580 uL, 4.16 mmol) was added at room temperature under nitrogen. The mixture was stirred for 30 min minutes, and N-[(CH₂)₅-COO-*t*-Bu]-N'-Boc-N'-methyl-ethylenediamine (1.43 g, 4.16 mmol) in 10 mL dry THF was added. The resulting mixture was stirred at room temperature for over 30 minutes. The solvent was removed, and then the product mixture was dissolved in methylene chloride and washed with saturated K₂CO₃ solution 3 times and once with water. The organic layer was dried over Na₂SO₄, and the solvent was removed. The compound was purified by flash column chromatography using heptane/ethyl acetate as eluent to give the product in a yield of 77 % (0.875 g).

[00276] Synthesis of N-[(CH₂)₅ COO-*t*-Bu]-N'-methyl-ethylene-diamine-2-cyanobenzothiazole-6-carbamate. N-[(CH₂)₅ COO-*t*-Bu]-N'-Boc-N'-methyl-ethylene-diamine 6-hydroxybenzo[*d*]thiazole-2-carbonitrile-carbamate was dissolved in 10 mL methylene chloride over ice, and 5 mL (20 %) thioanisole was added. 10 mL of TFA were slowly added over ice, after which the reaction mixture was stirred on ice for 30 min, then stirred at room temperature for 5 hours. The solvents were then co-evaporated with toluene, and the product was purified by flash column chromatography using heptane/ethyl acetate followed by methylene chloride/methanol to give the product in a 94 % yield (0.589 mg).

[00277] **Synthesis of N-[(CH₂)₅COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate.** To the solution of 3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic acid (1.13 g, 4.53 mmol) and isobutyl chloroformate (594 μ L, 4.53 mmol) in 35 ml dry THF, N-methyl morpholine (1.26 mL, 9.05 mmol) was added at 0°C. The resultant mixture was stirred for 1 hr at 0°C, after which N-[(CH₂)₅ COO-*t*-Bu]-N'-methyl-ethylene-diamine-2-cyanobenzothiazole-6-carbamate in 5 ml of dry THF was added, and the resultant mixture was stirred for 2 hours at room temperature. The product mixture was acidified with acetic acid then dried down. The product was purified by flash column chromatography using heptane/ethyl acetate, then methylene chloride/methanol to give a yield of 87% (0.817 g).

[00278] **Synthesis of N-[(CH₂)₅COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate NHS ester (C6 NHS ester).** To the mixture of TSTU (2.76 g, 9.16 mmol) and N-[(CH₂)₅COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6-carbamate (0.815 g, 1.31 mmol), 60 mL of acetonitrile was added. DIPEA (1.82 mL, 10.47 mmol) was added, and the reaction mixture was stirred for 1 hour at room temperature. The product mixture was diluted with a large volume of methylene chloride, then washed with 30% citric acid solution twice, then once with water, dried with Na₂SO₄, filtered, and the solvent was removed. The product was purified by flash column chromatography using heptane/ethyl acetate to give a 76 % yield (0.717 g).

[00279] **Synthesis of #5661.** C6 NHS ester and CDCA-lysCOOH were coupled using **Method 1** to give product #5661 in a 71 % yield (111 mg). MS-ESI (m/e): 1126.48 [M+H].

[00280] **Synthesis of #5665.** The luciferin version of #5661 was prepared using **Method 2** to give a 38 % yield (21 mg). MS-ESI (m/e): 1230.57 [M+H].

[00281] **Synthesis of #5662.** C6 NHS ester and UDCA-lysCOOH were coupled using **Method 1** to give product #5661 in a 78 % yield (122 mg). MS-ESI (m/e): 1126.48 [M+H].

[00282] **Synthesis of #5666.** The luciferin version of #5662 was prepared using **Method 2** to give a 20 % yield (16 mg). MS-ESI (m/e): 1230.57 [M+H].


[00283] **Synthesis of #5660.** C6 NHS ester and DCA-lysCOOH were coupled using **Method 1** to give product #5661 in a 38 % yield (59 mg). MS-ESI (m/e): 1126.67 [M+H].

[00284] **Synthesis of #5664.** The luciferin version of #5660 was prepared using **Method 2** to give a 98 % yield (27 mg). MS-ESI (m/e): 1230.62 [M+H].

[00285] **Synthesis of #5659.** C6 NHS ester and LCA-lysCOOH were coupled using **Method 1** to give product #5661 in a 33 % yield (51 mg). MS-ESI (m/e): 1109.55 [M+H].

[00286] **Synthesis of #5663.** The luciferin version of #5659 was prepared using **Method 2** to give a 36 % yield (12 mg). MS-ESI (m/e): 1212.30 [M-H].

EXAMPLE 11



[00287] **Synthesis of FAM diamine.** Diethyl amine (317 mg, 5.28 mmol) was dissolved in 7 mL DMF, and then 500 mg of FAM NHS ester dissolved in 7 mL DMF was added drop-wise and stirred at room temperature for 30 min. The solvent was then removed, and the product mixture was re-dissolved in methanol and water, and then purified by reverse phase HPLC on a

C18 column using 0.1% formic acid and methanol to give FAM diamine in a yield of 77 % (342 mg). MS-ESI (m/e): 442.9 [M+H].

[00288] Synthesis of # 5682. PEG2 NHS ester and FAM diamine were coupled using **Method 1** described above. The product was purified by reverse phase HPLC on a C18 column using 0.1% formic acid and acetonitrile to give #5682 in a 47 % yield (15 mg). MS-ESI (m/e): 1069.81 [M+H].

[00289] Synthesis of # 5826. The luciferin version of # 5682 was prepared using **Method 2** described above. The product was then purified by reverse phase HPLC on a C18 column using 0.1% formic acid and acetonitrile to give # 5826 in a 89 % yield (42 mg). MS-ESI (m/e): 1175.20 [M+H].



[00290] Synthesis of # 5684. C4 NHS ester and FAM diamine were coupled using **Method 1** described above. The product was purified by reverse phase HPLC on a C18 column using 0.1% formic acid and acetonitrile to give #5682 in a 46 % yield (41 mg). MS-ESI (m/e): 995.98 [M+H].


[00291] Synthesis of # 5824. The luciferin version of # 5684 was prepared using **Method 2** described above. The product was then purified by reverse phase HPLC on a C18 column using 0.1% formic acid and acetonitrile to give # 5824 in > 99% yield (67 mg). MS-ESI (m/e): 1099.37 [M+H].

[00292] Synthesis of # 5683. C6 NHS ester and FAM diamine were coupled using **Method 1** described above. The product was purified by reverse phase HPLC on a C18 column using 0.1% formic acid and acetonitrile to give #5682 in a 78 % yield (19 mg). MS-ESI (m/e): 1024.38 [M+H].

[00293] Synthesis of # 5825. The luciferin version of # 5683 was prepared using **Method 2** described above. The product was then purified by reverse phase HPLC on a C18 column using 0.1% formic acid and acetonitrile to give # 5825 in a 67 % yield (36 mg). MS-ESI (m/e): 1128.98 [M+H].

EXAMPLE 12






[00294] Synthesis of #5907, #5909, and #5910: 1 mL of 20nm gold nanoparticles (GNPs) (OD = 50) coated with Peg3000 and functionalized with free amine groups were purchased from Cytodyagnostics Inc. 10 μ L of GNP solution was added to a small eppendorf tube, and 1 mg of N-[(PEG)₂COOH]-N'-methyl-N'-[3-methyl-3-(2,4,5-trimethyl-3,6-dioxocyclohexa-1,4-dien-1-yl)butanoic amido]-ethylenediamine-2-cyanobenzothiazole-6- carbamate NHS ester dissolved in 10 μ L DMF was added. 500 μ L of PBS buffer and 500 μ L of DMF was added, then sonicated briefly and placed on shaker at room temperature for 4 hours. After the reaction, 1 mL PBST (0.05% Tween80 in PBS buffer) was added and centrifuged at 14,000 rpm for 30 min. The supernatant was removed, and 350 μ L methanol added. The pellet was resuspended and then spun down again. The methanol wash was repeated twice, and the pellet washed similarly with DMF three times.

[00295] Synthesis of #5908, #5911, and #5912: 1 mg of d-cysteine was dissolved in 350 μ L PBS buffer and added to eppendorf tube containing modified GNPs from #5907, #5909, or #5910, respectively. 350 μ L methanol was added, and then the solution placed on shaker at room temperature for 3 hours. After the reaction, the contents were centrifuged at 14,000 rpm for 30 min, and then most of the supernatant removed. The pellet was washed three times with a 1:1 mixture of water and methanol, and then three times with DMF.



EXAMPLE 13




[00296] Synthesis of #5915. Approximately 100mg (392 μ mol, 2 equiv.) of maleimide TFA salt was added to a vial together with 2mL DMF and 150 mg (196 μ mol, 1 equiv.) of the

respective NHS ester dissolved in 1mL DMF. 70 μ L DIPEA (392 μ mol, 2 equiv.) was added and then stirred at room temperature for over an hour. The reaction mixture was dried down, and then purified by flash column chromatography using a gradient of methylene chloride/ methanol. MS-ESI (m/e): 791.9 [M+H].

[00297] Compounds #5916, #5917, and #5918 were synthesized in a similar manner using the corresponding NHS esters.





EXAMPLE 14

[00298] SLNs may be attached to the quinone probe through the scheme shown in FIG. 15.

EXAMPLE 15

Antibody Labeling

[00299] To demonstrate labeling of an antibody with the compounds of the present invention, the monoclonal antibody, Herceptin, was labeled with PBI-5508 (Example 4). Lyophilized Herceptin was dissolved in 0.1 M Sodium Bicarbonate, pH 8.6 at 10 mg/ml, and then diluted to 1 mg/ml in 100 μ l of the same buffer. A 50 mM stock of the pro-luciferin, PBI-5508, was prepared in 100% DMSO. For labeling of the antibody, 3.3 μ l of PBI-5508 (3.3 nmol) was added to 100 μ l of diluted Herceptin (0.66 nmol) for 5-fold molar excess of PBI-5508. The sample was then covered in foil and incubated for 60 min at room temperature on a tube rotator. To remove free PBI-5508, 100 μ l of the labeling reaction was placed onto an equilibrated G-25 Sequencing MicroSpin Column (Amersham, 200 μ l resin) and eluted by spinning for 30 sec at 3000 rpm. The labeled Herceptin was contained in the flow-through. Labeling efficiency was determined by spectrophotometry by taking the absorbance of the Herceptin-PBI-5508 molecule at 280 nm and 320 nm. The degree of labeling was 2.3 PBI-5508 molecules per one Herceptin antibody molecule.

EXAMPLE 16

[00300] In a white, 384-well, low volume assay plate, in triplicate, 5 μ l of Herceptin:PBI-5508 (Example 15) was plated at varying concentrations, or 5 μ l of 0.1 M sodium bicarbonate pH 8.6 buffer was plated as a control. 5 μ l of a solution containing Luciferin Detection Reagent (Promega, V859/865), reductase (Promega, NAD(P)H-Glo kit at the recommended concentration), 50 μ M NADH and 10 mM D-cysteine was added to the assay wells. The reaction was incubated at room temperature for 45 min, and luminescence (relative light unit, RLU) was measured on a Tecan M1000Pro.

[00301] The results in Table 3 show that the Herceptin:PBI-5508 molecule is a substrate for diaphorase.

Table 3

	Avg RLU	SD
Herceptin: 5508, 1x	15513	166
Herceptin: 5508, 0.5x	6526	570
Buffer Control	11	5

EXAMPLE 17

[00302] In a 384-well assay plate, SKBR3 cells were plated at 10,000 cells/well in 20 μ l McCoy media. Other wells received media only as a control. Media and cells were incubated at 37°C /5% CO₂ for 2 hours. Herceptin:PBI-5508 was added to a final concentration of 15 ng/ μ l plus or minus 500 mM sucrose (final concentration) in 20 μ l final volume. The reaction was incubated at 37°C /5%CO₂ for 6 hours. Then, 40 μ l of Luciferin Detection Reagent (Promega, V865/859) plus 10 mM D-cysteine was added to all wells. The reaction was incubated at room temperature for 20 min, and luminescence (RLU) was measured on a Tecan M1000Pro.

[00303] The results in Table 4 show that Herceptin:PBI-5508 was reduced in cells after internalization. Sucrose was used to increase the release of molecules from the endosome.

Table 4

Cells	Herceptin: 5508	Sucrose	Avg RLU at 6 h
Y	Y	N	58
Y	Y	Y	143
N	Y	N	88
N	Y	Y	8

EXAMPLE 18

[00304] Experiments were conducted during development of embodiments of the present disclosure to demonstrate an uptake transporter assay using a variety of compounds described herein.

[00305] Stock solutions of each compound were prepared as 10mM in methanol. Corning TransportoCells OATP1B1*1a, OATP1B3, and Control Cells were used as cell model for cell-based uptake transporter assay. Cells were cultured according to manufacturer's recommendations. Briefly, cells were thawed from liquid nitrogen storage and cultured in Dulbecco's Modified Eagle's medium supplemented with fetal bovine serum (FBS), MEM non-essential amino acid, and sodium butyrate. Cells were then plated in Corning 96-well, poly-D-Lysine white/clear plate at 100,000 cells/well in 100µl medium and incubated for 24 hours at 37°C with 5% CO₂.

[00306] The uptake transporter assay was carried out by adding 100µl of Hank's balanced salt solution containing 10µM of one of the compounds to all the cells and incubated at 37°C for 30 minutes. For cells treated with PBI-5651, PBI-5657, PBI-5658, PBI-5665, PBI-5666, PBI-5824, PBI-5825, and PBI -5826, an equal volume of Luciferin Detection Reagent (Promega) was added. For cells treated with PBI-5648, PBI-5682, PBI-5683, and PBI-5684, an equal volume of Luciferin Detection Reagent containing D-cysteine (Promega) was added. After a 20 minute incubation at room temperature (20°-25°C), luminescence was detected on a GloMax® luminometer.

[00307] FIGS. 1-12 demonstrates increased luminescence (RLU) over TransportoCells Control Cells and no-cell control from TransportoCells OATP1B1*1a and OATP1B3 using the compounds of the present invention.

EXAMPLE 19

[00308] Experiments were conducted during development of embodiments of the present invention to demonstrate an uptake transporter assay using different concentrations of PBI-5651.

[00309] PBI-5651 stock solution was prepared as 10mM in methanol. Corning TransportoCells OATP1B1*1a, OATP1B3, and Control Cells were used as cell model for cell-based uptake transporter assay. Cells were cultured according to manufacturer's recommendations. Briefly, cells were thawed from liquid nitrogen storage and cultured in Dulbecco's Modified Eagle's medium supplemented with fetal bovine serum (FBS), MEM non-essential amino acid,

and sodium butyrate. Cells were then plated in Corning 96-well, poly-D-Lysine white/clear plate at 100,000 cells/well in 100µl medium and incubated for 24 hours at 37°C with 5% CO₂.

[00310] The uptake transporter assay was carried out by adding 100µl of Hank's balanced salt solution containing various concentrations (See FIG. 13) of PBI-5651 to all the cells and incubated at 37°C for 30 minutes. An equal volume of Luciferin Detection Reagent (Promega) was added to the cells. After a 20 minute incubation at room temperature (20°-25°C), luminescence was detected on a GloMax® luminometer.


[00311] FIG. 13 demonstrates dynamic response of luminescence (RLU) from TransportoCells OATP1B1*1a and OATP1B3.

[00312] Any ranges given either in absolute terms or in approximate terms are intended to encompass both, and any definitions used herein are intended to be clarifying and not limiting. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Moreover, all ranges disclosed herein are to be understood to encompass any and all subranges (including all fractional and whole values) subsumed therein.

[00313] Furthermore, the invention encompasses any and all possible combinations of some or all of the various embodiments described herein. Any and all patents, patent applications, scientific papers, and other references cited in this application, as well as any references cited therein, are hereby incorporated by reference in their entirety.

[00314] For reasons of completeness, various aspects of the present disclosure are set out in the following numbered clauses:

[00315] Clause 1. A compound of formula (I), or a salt thereof,



(I)

wherein

A is a reporter moiety;

R¹⁴ is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R⁹ and R¹⁰ are independently selected from alkyl;

R¹¹, R¹² and R¹³ are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R¹¹ and R¹² can form a fused phenyl ring;

X is O;

L is $-(\text{CH}_2)_m\text{C}(\text{R}^{17})_2(\text{CH}_2)_n\text{-Y-C(O)-}$;

R¹⁷ is independently H, alkyl or both R¹⁷ together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;

n is an integer from 0-2;


Y is O or NR¹⁵;

R¹⁵ is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl-N(R²³)C(O)R²⁴, -alkyl-SO₃R²⁵, -alkyl-SO₂N(R²⁶)(R²⁷), -alkyl-COR²⁸, -alkyl-CO₂R²⁹, -alkyl-OC(O)R²⁹, -alkyl-OC(O)N(R³⁰)(R³¹), -alkyl-CON(R³⁰)(R³¹), or polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with one or more suitable substituents; and R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl

[00316] Clause 2. The compound of claim 1, wherein R¹⁵ is polyalkoxyalkyl, wherein the polyalkoxyalkyl is $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-haloalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-hydroxyalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-aminoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkylaminoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{di(alkyl)aminoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-azidoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-cyanoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkenyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkynyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-N}(\text{R}^{23})\text{C}(\text{O})\text{R}^{24}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-SO}_3\text{R}^{25}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-SO}_2\text{N}(\text{R}^{26})\text{(R}^{27})$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-COR}^{28}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-OC}(\text{O})\text{R}^{29}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-OC}(\text{O})\text{N}(\text{R}^{30})\text{(R}^{31})$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-CO}_2\text{R}^{29}$, and $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-CON}(\text{R}^{30})\text{(R}^{31})$, wherein R²³, R²⁴, R²⁵, R²⁶, R²⁷, R²⁸, R²⁹, R³⁰, and R³¹ are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl; and x is an integer selected from 1 to 20.

[00317] Clause 3. The compound of clause 1, wherein R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of alkyl, alkoxyalkyl, haloalkyl, haloalkoxyalkyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, azidoalkyl, cyanoalkyl, alkoxy, haloalkoxy, alkoxyalkoxy, alkenyl, alkynyl, $-N(R^{23})C(O)R^{24}$, $-SO_3R^{25}$, $-SO_2N(R^{26})(R^{27})$, $-COR^{28}$, $-CO_2R^{29}$, and $-CON(R^{30})(R^{31})$, wherein R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00318] Clause 4. The compound of clause 1, wherein R^{15} is



wherein


R^{40} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein said aryl, heteroaryl, heterocyclyl, and cycloalkyl are unsubstituted or substituted with one or more suitable substituents; and

x is an integer selected from 1 to 20.

[00319] Clause 5. The compound of clause 4, wherein R^{40} is hydrogen.


[00320] Clause 6. The compound of clause 4, wherein R^{40} is a 5- or 6-membered heterocyclyl, having 1, 2, or 3 heteroatoms independently selected from the group consisting of N, O, and S.

[00321] Clause 7. The compound of clause 4, wherein R^{40} is



[00322] Clause 8. The compound of clause 4, wherein x is 2, 3, or 4.

[00323] Clause 9. The compound of clause 4, having formula (I-vi), or a salt thereof,




(I-vi)

wherein

R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R⁴⁰, x, and A are as defined above.

[00324] Clause 10. The compound of clause 1, selected from the group consisting of:



[00325] Clause 11. The compound of clause 1, wherein the reporter moiety comprises a bioluminescent reporter moiety or fluorescent reporter moiety.

[00326] Clause 12. The compound of clause 11, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00327] Clause 13. The compound of clause 12, wherein the substrate comprises luciferin, a luciferin derivative or analog, a preluciferin or a preluciferin analog, a coelenterazine or a coelenterazine derivative or analog.

[00328] Clause 14. The compound of clause 12, wherein the substrate is furimazine, coelenterazine H,H, or luciferin.

[00329] Clause 15. The compound of clause 12, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase or a variant thereof.

[00330] Clause 16. The compound of clause 11, wherein the reporter moiety comprises a fluorescent reporter moiety.

[00331] Clause 17. The compound of clause 16, wherein the fluorescent reporter moiety comprises a fluorophore.

[00332] Clause 18. The compound of clause 17, wherein the fluorophore is coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthene or carbopyronine.

[00333] Clause 19. A method for evaluating cellular uptake of an agent, the method comprising:

a) contacting a sample with a labeled agent, wherein the labeled agent comprises an agent and a compound of any one of clauses 1 to 18, wherein the sample comprises a cell; and

b) detecting light emission,

whereby the detection of light emission indicates cellular uptake of the agent.

[00334] Clause 20. The method of clause 19, wherein the cellular uptake of the agent results in the reduction of the compound and the release of the reporter moiety, thereby generating a released reporter moiety.

[00335] Clause 21. The method of clause 20, wherein the reporter moiety comprises a fluorescent reporter moiety.

[00336] Clause 22. The method of clause 21, wherein detecting light emission comprises exposing the sample to a wavelength of light to generate fluorescence and detecting light emission comprises detecting the fluorescence level of the sample, wherein an increase in fluorescence level or a change in fluorescence wavelength as compared to the fluorescence level of a control sample indicates cellular uptake of the agent.

[00337] Clause 23. The method of clause 22, wherein the control sample is a sample that is not contacted with a labeled agent.

[00338] Clause 24. The method of clause 22, wherein the fluorescence level is detected inside the cell.

[00339] Clause 25. The method of clause 22, wherein the fluorescence level is detected outside the cell.

[00340] Clause 26. The method of clause 20, wherein the reporter moiety comprises a bioluminescent reporter moiety.

[00341] Clause 27. The method of clause 26, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00342] Clause 28. The method of clause 27, wherein the cell comprises a luciferase.

[00343] Clause 29. The method of clause 28, wherein the cell expresses the luciferase.

[00344] Clause 30. The method of clause 28, wherein detecting light emission comprises detecting the luminescence level produced by the luciferase and the released reporter moiety.

[00345] Clause 31. The method of clause 27, further comprising contacting the sample with a luciferase and wherein detecting light emission comprises detecting luminescence level produced by the luciferase and the released reporter moiety.

[00346] Clause 32. The method of clause 30 or 31, wherein the light emission is detected inside the cell or detected outside the cell.

[00347] Clause 33. A method for evaluating cellular uptake of an agent, the method comprising:

(a) contacting a sample with a labeled agent, wherein the labeled agent comprises an agent and a compound of any one of clause 1 to 11, wherein the sample comprises a cell, wherein the cellular uptake of the agent results in the reduction of the compound and the release of the reporter moiety, thereby generating a released reporter moiety;

(b) exposing the sample to a wavelength of light to generate fluorescence,

(c) detecting the fluorescence level of the sample, and

(d) comparing the fluorescence level of the sample to a fluorescence level of a control sample, wherein an increase in fluorescence level or a change in fluorescence wavelength as compared to the fluorescence level of the control sample indicates cellular uptake of the agent.

[00348] Clause 34. The method of clause 33, wherein the reporter moiety comprises a fluorescent reporter moiety.

[00349] Clause 35. The method of clause 34, wherein the fluorescent reporter moiety comprises a fluorophore.

[00350] Clause 36. The method of clause 35, wherein the fluorophore is coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthen or carbopyronine.

[00351] Clause 37. The method of clause 33, wherein the fluorescence level is detected inside the cell.

[00352] Clause 38. The method of clause 33, wherein the fluorescence level is detected in the sample.

[00353] Clause 39. The method of clause 33, wherein the control sample is sample medium with labeled agent but without cells, a sample with labeled agent and cells but without experimental treatment, or a sample not contacted with labeled agent.

[00354] Clause 40. A method for evaluating cellular uptake of an agent, the method comprising:

a) contacting a sample with a labeled agent, wherein the labeled agent comprises an agent and a compound of any one of clause 1 to 11, wherein the sample comprises a cell, wherein the cell comprises luciferase, wherein the cellular uptake of the agent results in the reduction of the compound and the release of the reporter moiety, thereby generating a released reporter moiety;

b) detecting a luminescence level of the sample; and

c) comparing the luminescence level of the sample to a luminescence level of a control sample, wherein cellular uptake of the agent is indicated if the luminescence level of the sample is higher or lower than the luminescence level of the control sample.

[00355] Clause 41. The method of clause 40, wherein the luminescence level is produced by the luciferase and the released reporter moiety.

[00356] Clause 42. The method of clause 40, wherein the luminescence is detected inside the cell or detected outside in the cell medium.

[00357] Clause 43. The method of clause 40, wherein the control sample is a sample that is not contacted with a labeled agent.

[00358] Clause 44. A method for evaluating cellular uptake of an agent, the method comprising:

a) contacting a sample with a labeled agent, wherein the labeled agent comprises an agent and a compound of any one of clause 1 to 11, wherein the sample comprises a cell and a

cell medium, wherein the cellular uptake of the agent results in the reduction of the compound and the release of the reporter moiety, thereby generating a released reporter moiety;

b) contacting the sample with a luciferase;

c) detecting a luminescence level of the sample; and

d) comparing the luminescence level of the sample to a luminescence level of a control sample, wherein cellular uptake of the agent is indicated if the luminescence level of the sample is higher or lower than the luminescence level of the control sample.

[00359] Clause 45. The method of clause 44, wherein the luminescence level is produced by the luciferase and the release reporter moiety.

[00360] Clause 46. The method of clause 44, wherein the luminescence is detected in the cell or detected in the cell medium.

[00361] Clause 47. The method of clause 44, wherein the control sample is sample medium with labeled agent but without cells, a sample with labeled agent and cells but without experimental treatment, or a sample not contacted with labeled agent.

[00362] Clause 48. The method of any one of clauses 19-47, wherein the agent is a biological agent.

[00363] Clause 49. The method of clause 48, wherein the biological agent is selected from the group consisting of a protein, a nucleic acid, a lipid, and a sugar.

[00364] Clause 50. The method of clause 49, wherein the protein is an antibody or a lipoprotein.

[00365] Clause 51. The method of clause 50, wherein the antibody is Herceptin.

[00366] Clause 52. The method of any one of clauses 19-47, wherein the agent is a non-biological agent.

[00367] Clause 53. The method of clause 52, wherein the non-biological agent is selected from the group consisting of a therapeutic drug, a small molecule, and a nanoparticle, or any combination thereof.

[00368] Clause 54. The method of any one of clauses 19-53, wherein the cell is a eukaryotic cell or a prokaryotic cell.

[00369] Clause 55. The method of clause 54, wherein the cell is in an animal.

[00370] Clause 56. The method of clause 54, wherein the cell is growing in culture medium.

[00371] Clause 57. A labeled agent derived from an agent and a compound according to any one of clauses 1 to 10.

[00372] Clause 58. The labeled agent of clause 57, wherein the agent is a biological agent.

[00373] Clause 59. The labeled agent of clause 58, wherein the biological agent is selected from the group consisting of a protein, a nucleic acid, a lipid, and a sugar.

[00374] Clause 60. The labeled agent of clause 59, wherein the protein is an antibody or a lipoprotein.

[00375] Clause 61. The labeled agent of clause 60, wherein the antibody is Herceptin.

[00376] Clause 62. The labeled agent of clause 57, wherein the reporter moiety comprises a bioluminescent reporter moiety or fluorescent reporter moiety.

[00377] Clause 63. The labeled agent of any one of clauses 57, wherein the agent is a non-biological agent.

[00378] Clause 64. The labeled agent of clause 63, wherein the non-biological agent is selected from the group consisting of a therapeutic drug, a small molecule, and a nanoparticle, or any combination thereof.

[00379] Clause 65. The labeled agent of clause 62, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00380] Clause 66. The labeled agent of clause 65, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.

[00381] Clause 67. The labeled agent of clause 66, wherein the substrate is furimazine, coelenterazine H,H, luciferin, or pre-luciferin.

[00382] Clause 68. The labeled agent of clause 66, wherein the luciferase is a beetle luciferase, a Renilla luciferase, or an Oplophorus luciferase.

[00383] Clause 69. The labeled agent of clause 62, wherein the reporter moiety comprises a fluorescent reporter moiety.

[00384] Clause 70. The labeled agent of clause 69, wherein the fluorescent reporter moiety comprises a fluorophore.

[00385] Clause 71. The labeled agent of clause 70, wherein the fluorophore is coumarin, a rhodamine, R110, fluorescein, DDAO, resorufin or cresyl violet.

[00386] Clause 72. A kit comprising a compound according to any one of clauses 1 to 18, an agent, or any combination thereof.

[00387] Clause 73. A kit comprising a labeled agent comprising an agent and a compound according to any one of clauses 1 to 18.

[00388] Clause 74. The kit of clause 72 or 73, wherein the agent is a biological agent.

[00389] Clause 75. The kit of clause 74, wherein the biological agent is selected from the group consisting of a protein, a nucleic acid, a lipid, and a sugar.

[00390] Clause 76. The kit of clause 75, wherein the protein is an antibody or a lipoprotein.

[00391] Clause 77. The kit of clause 72 or 73, wherein the agent is a non-biological agent.


[00392] Clause 78. The kit of clause 77, wherein the non-biological agent is selected from the group consisting of a therapeutic drug, a small molecule, and a nanoparticle, or any combination thereof.

[00393] Clause 79. The kit of clause 72 or 73, wherein the compound is a substrate for luciferase.

[00394] Clause 80. The kit of clause 79, further comprising a detection reagent.

[00395] Clause 81. The kit of clause 80, wherein the detection reagent comprises a luciferase enzyme.

[00396] Clause 82. A compound of formula (I), or a salt thereof,



(I)

wherein

A is a reporter moiety;

R¹⁴ is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R⁹ and R¹⁰ are independently selected from alkyl;

R¹¹, R¹² and R¹³ are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R¹¹ and R¹² can form a fused phenyl ring;

X is O;

L is $-(\text{CH}_2)_m\text{C}(\text{R}^{17})_2(\text{CH}_2)_n\text{-Y-C(O)-}$;

R^{17} is independently H, alkyl or both R^{17} together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;

n is an integer from 0-2;

Y is O or NR^{15} ;

R^{15} is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl- $\text{N}(\text{R}^{23})\text{C(O)}\text{R}^{24}$, -alkyl- SO_3R^{25} , -alkyl- $\text{SO}_2\text{N}(\text{R}^{26})(\text{R}^{27})$, -alkyl- COR^{28} , -alkyl- CO_2R^{29} , -alkyl- $\text{OC(O)}\text{R}^{29}$, -alkyl- $\text{OC(O)}\text{N}(\text{R}^{30})(\text{R}^{31})$, -alkyl-amide, or polyalkoxyalkyl, wherein the polyalkoxyalkyl and -alkyl-amide are unsubstituted or substituted with one or more suitable substituents; and

R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00397] Clause 83. The compound of clause 82, wherein R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is selected from the group consisting of $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-haloalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-hydroxyalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-aminoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkylaminoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{di(alkyl)aminoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-azidoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-cyanoalkyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkenyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkynyl}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-N}(\text{R}^{23})\text{C(O)}\text{R}^{24}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-SO}_3\text{R}^{25}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-SO}_2\text{N}(\text{R}^{26})(\text{R}^{27})$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-COR}^{28}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-OC(O)}\text{R}^{29}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-OC(O)}\text{N}(\text{R}^{30})(\text{R}^{31})$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-CO}_2\text{R}^{29}$, $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-CON}(\text{R}^{32})(\text{R}^{33})$ and $-(\text{C}_2\text{-C}_6\text{-alkoxy})_x\text{-alkyl-(CO)-NR}^{34}\text{-(CR}^a\text{R}^b)_p\text{-NR}^{35}\text{(CO)-T}$, wherein

R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl;

x is an integer selected from 1 to 20;

R^{32} and R^{33} are each independently selected from hydrogen, alkyl, carboxy, a peptide, a drug, a biologically active moiety, and a dye;

R^{34} and R^{35} are each independently selected from hydrogen, and alkyl;

R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;

p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

[00398] Clause 84. The compound of clause 82, wherein R¹⁵ is polyalkoxyalkyl, wherein the polyalkoxyalkyl is $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CON}(R^{32})(R^{33})$ or $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-(CO)-NR}^{34}\text{-(CR}^a\text{R}^b)_p\text{-NR}^{35}\text{(CO)-T}$;

x is an integer selected from 1 to 20;

R³² and R³³ are each independently selected from hydrogen, alkyl, carboxy, aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof;


R³⁴ and R³⁵ are each independently selected from hydrogen, and alkyl;


R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;

p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

[00399] Clause 85. The compound of clause 82, wherein R¹⁵ is





wherein

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof; and

x is an integer selected from 1 to 20.

[00400] Clause 86. The compound of clause 82, wherein R^{15} is -alkyl-amide, wherein the -alkyl-amide is -alkyl-CON(R^{32})(R^{33}) or -alkyl-(CO)-NR³⁴-(CR^aR^b)_p-NR³⁵(CO)-T, wherein

R^{32} and R^{33} are each independently selected from hydrogen, alkyl, carboxy, aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heterocycle, alkene, polyol, alkenylpolyol, a peptide, a drug, a derivative of a drug, a biologically active moiety, and a dye;


R^{34} and R^{35} are each independently selected from hydrogen, and alkyl;

R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;

p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

[00401] Clause 87. The compound of clause 82, wherein R^{15} is



wherein


T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof; and

x is 0 to 20.

[00402] Clause 88. The compound of clause 82, wherein R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of alkyl, alkoxyalkyl, haloalkyl, haloalkoxyalkyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, azidoalkyl, cyanoalkyl, alkoxy, haloalkoxy, alkoxyalkoxy, alkenyl, alkynyl, $-N(R^{23})C(O)R^{24}$, $-SO_3R^{25}$, $-SO_2N(R^{26})(R^{27})$, $-COR^{28}$, $-CO_2R^{29}$,

and $-\text{CON}(\text{R}^{30})(\text{R}^{31})$, wherein R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

[00403] Clause 89. The compound of clause 82, wherein R^{15} is



wherein


R^{40} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein said aryl, heteroaryl, heterocyclyl, and cycloalkyl are unsubstituted or substituted with one or more suitable substituents; and

x is an integer selected from 1 to 20.

[00404] Clause 90. The compound of clause 89, wherein R^{40} is hydrogen.


[00405] Clause 91. The compound of clause 89, wherein R^{40} is a 5- or 6-membered heterocyclyl, having 1, 2, or 3 heteroatoms independently selected from the group consisting of N, O, and S.

[00406] Clause 92. The compound of clause 89, wherein R^{40} is




[00407] Clause 93. The compound of clause 89, wherein x is 2, 3, or 4.

[00408] Clause 94. The compound of clause 89, having formula (I-vi), or a salt thereof,




(I-vi).

[00409] Clause 95. The compound of clause 82, wherein R¹⁵ is




each x is independently 0 to 20.

[00410] Clause 96. The compound of clause 82, selected from the group consisting of:




PBI 5463

;




PBI 5470

;




PBI 5471

;




PBI 5508

;








;




PBI 5915

;









PBI 5666

;




PBI 5684

;




PBI 5825


;




;




;




;




;









;













;





















;






;




;



;

;



and

or a salt thereof.

[00411] Clause 97. The compound of any one of clauses 83-87, wherein T is a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, a small molecule, a fluorescent molecule, a polymer-linked nanoparticle, a detergent, or a sugar.

[00412] Clause 98. The compound of clause 97, wherein T is a protein.

[00413] Clause 99. The compound of clause 98, wherein the protein is an antibody.


[00414] Clause 100. The compound of clause 97, wherein T is a polypeptide.

[00415] Clause 101. The compound of clause 100, wherein the polypeptide is a polypeptide toxin.

[00416] Clause 102. The compound of clause 97, wherein T is a therapeutic drug.

[00417] Clause 103. The compound of clause 102, wherein the compound is PBI-5651, PBI-5658, PBI-5666, PBI-5648, PBI-5655, PBI-5662, PBI-5657, PBI-5665, PBI-5667, PBI-5650,

PBI-5654, PBI-5661, PBI-5827, PBI-5828, PBI-5829, PBI-5647, PBI-5668, PBI-5659, PBI-5663, PBI-5652, PBI-5669, PBI-5625, PBI-5649, PBI-5653, PBI-5656, PBI-5660, PBI-5664, PBI-5830, PBI-5831 or



[00418] Clause 104. The compound of clause 97, wherein T is a fluorescent molecule.

[00419] Clause 105. The compound of clause 104, wherein the fluorescent molecule is FAM diamine.

[00420] Clause 106. The compound of clause 105, wherein the compound is PBI-5683, PBI-5826, PBI-5682, PBI-5684, PBI-5824, or PBI-5825.

[00421] Clause 107. The compound of clause 97, wherein T is a polymer-linked nanoparticle or a detergent.

[00422] Clause 108. The compound of clause 107, wherein the nanoparticle is a gold nanoparticle.

[00423] Clause 109. The compound of clause 108, wherein the compound is PBI-5907, PBI-5908, PBI-5909, PBI-5910, PBI-5911, or PBI-5912.

[00424] Clause 110. The compound of clause 82, wherein the reporter moiety comprises a bioluminescent reporter moiety or fluorescent reporter moiety.

[00425] Clause 111. The compound of clause 110, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00426] Clause 112. The compound of clause 111, wherein the substrate comprises luciferin, a luciferin derivative or analog, a preluciferin or a preluciferin analog, a coelenterazine or a coelenterazine derivative or analog.

[00427] Clause 113. The compound of clause 111, wherein the substrate is furimazine, coelenterazine H,H, or luciferin.

[00428] Clause 114. The compound of clause 111, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.

[00429] Clause 115. The compound of clause 110, wherein the reporter moiety comprises a fluorescent reporter moiety.

[00430] Clause 116. The compound of clause 115, wherein the fluorescent reporter moiety comprises a fluorophore.

[00431] Clause 117. The compound of clause 116, wherein the fluorophore is coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthene or carbopyronine.

[00432] Clause 118. A method for evaluating cellular uptake of an agent, the method comprising:

a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of clauses 82-117 or ii) a compound of any one of clauses 82-117, wherein the sample comprises a cell; and

b) detecting light emission,

whereby the detection of light emission indicates cellular uptake of the agent.

[00433] Clause 119. The method of clause 118, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released reporter moiety.

[00434] Clause 120. The method of clause 119, wherein the reporter moiety comprises a fluorescent reporter moiety.

[00435] Clause 121. The method of clause 120, wherein detecting light emission comprises exposing the sample to a wavelength of light and detecting fluorescence level in the sample, wherein an increase in fluorescence or a change in fluorescence wavelength as compared to the fluorescence or fluorescence wavelength of a control sample indicates cellular uptake of the agent.

[00436] Clause 122. The method of clause 121, wherein the control sample is a sample that is not contacted with a labeled agent.

[00437] Clause 123. The method of clause 121, wherein fluorescence is detected inside the cell.

[00438] Clause 124. The method of clause 121, wherein fluorescence is detected outside the cell.

[00439] Clause 125. The method of clause 119, wherein the reporter moiety comprises a bioluminescent reporter moiety.

[00440] Clause 126. The method of clause 125, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00441] Clause 127. The method of clause 126, wherein the cell comprises a luciferase.

[00442] Clause 128. The method of clause 127, wherein the cell expresses the luciferase.

[00443] Clause 129. The method of clause 127, wherein detecting light emission comprises detecting the luminescence produced by the luciferase utilizing the released reporter moiety.

[00444] Clause 130. The method of clause 126, further comprising contacting the sample with a luciferase and wherein detecting light emission comprises detecting the luminescence produced by the luciferase utilizing the released reporter moiety.

[00445] Clause 131. The method of clause 129 or 130, wherein the light emission is detected inside or outside the cell.

[00446] Clause 132. A method for evaluating cellular uptake of an agent, the method comprising:

(a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of clauses 82-96, 110, or 115-117 or ii) a compound of any one of clauses 82-96, 110, or 115-117, wherein the sample comprises a cell, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released fluorescent reporter moiety;

(b) exposing the sample to a wavelength of light to generate fluorescence,

(c) detecting fluorescence in the sample, and

(d) comparing the fluorescence or fluorescent wavelength of the sample to the fluorescence of a control sample, wherein an increase in fluorescence or a change in the fluorescent wavelength as compared to fluorescence or fluorescent wavelength of the control sample indicates cellular uptake of the agent.

[00447] Clause 133. The method of clause 132, wherein the fluorescent reporter moiety comprises a fluorophore.

[00448] Clause 134. The method of clause 133, wherein the fluorophore is coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthen or carbopyronine.

[00449] Clause 135. The method of clause 132, wherein the fluorescence level is detected inside the cell.

[00450] Clause 136. The method of clause 132, wherein the fluorescence level is detected in the sample.

[00451] Clause 137. The method of clause 132, wherein the control sample is sample medium only with labeled agent, a sample with labeled agent and cells but without experimental treatment, or a sample not contacted with labeled agent.

[00452] Clause 137. A method for evaluating cellular uptake of an agent, the method comprising:

a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of clause 82-96, or ii) a compound of any one of clauses 82-96, wherein the sample comprises a cell, wherein the cell comprises luciferase, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released bioluminescent reporter moiety;

b) detecting luminescence in the sample; and

c) comparing the luminescence of the sample to luminescence of a control sample, wherein cellular uptake of the agent is indicated if the luminescence of the sample is changed compared to the luminescence of the control sample.

[00453] Clause 149. The method of clause 138, wherein the luminescence level is produced by the luciferase utilizing the released bioluminescent reporter moiety.

[00454] Clause 140. The method of clause 139, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00455] Clause 141. The method of clause 140, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.

[00456] Clause 142. The method of clause 141, wherein the substrate is furimazine, coelenterazine H,H, lucifeirn, or pre-luciferin.

[00457] Clause 143. The method of clause 141, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.

[00458] Clause 14482. The method of clause 138, wherein the luminescence is detected inside the cell or detected outside in the cell medium.

[00459] Clause 145. The method of clause 138, wherein the control sample is a sample that is not contacted with a labeled agent.

[00460] Clause 146. A method for evaluating cellular uptake of an agent, the method comprising:

[00461] a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of clauses 82-96 or ii) a compound of any one of clauses 82-96, wherein the sample comprises a cell and cell medium, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released bioluminescent reporter moiety;

b) contacting the sample with a luciferase;

c) detecting luminescence in the sample; and

d) comparing the luminescence of the sample to luminescence in a control sample, wherein cellular uptake of the agent is indicated if the luminescence in the sample is changed compared to the luminescence of the control sample.

[00462] Clause 147. The method of clause 146, wherein the luminescence level is produced by the luciferase utilizing the released bioluminescent reporter moiety.

[00463] Clause 148. The method of clause 147, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

[00464] Clause 149. The method of clause 148, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.

[00465] Clause 150. The method of clause 149, wherein the substrate is furimazine, coelenterazine H,H, luciferin, or pre-luciferin.

[00466] Clause 151. The method of clause 149, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.

[00467] Clause 152. The method of clause 146, wherein the luminescence is detected in the cell or detected in the cell medium.

[00468] Clause 153. The method of clause 146, wherein the control sample is sample medium with labeled agent but without cells, a sample with labeled agent and cells but without experimental treatment, or a sample not contacted with labeled agent.

- [00469] Clause 154. The method of any one of clauses 118-153, wherein the agent is a biological agent.
- [00470] Clause 155. The method of clause 154, wherein the biological agent is selected from the group consisting of a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, small molecule, a fluorescent molecule, and a sugar.
- [00471] Clause 156. The method of clause 155, wherein the protein is an antibody or a lipoprotein.
- [00472] Clause 157. The method of any one of clauses 118-153, wherein the agent is a non-biological agent.
- [00473] Clause 158. The method of clause 157, wherein the non-biological agent is a polymer-linked nanoparticle or a detergent.
- [00474] Clause 159. The method of any one of clauses 118-158, wherein the cell is a eukaryotic cell or a prokaryotic cell.
- [00475] Clause 160. The method of clause 159, wherein the cell is in an animal.
- [00476] Clause 161. The method of clause 159, wherein the cell is growing in culture medium.
- [00477] Clause 162. A labeled agent derived from i) an agent and a compound according to any one of clauses 82-96 or ii) a compound according to any one of clauses 82-117.
- [00478] Clause 163. The labeled agent of clause 162, wherein the agent is a biological agent.
- [00479] Clause 164. The labeled agent of clause 163, wherein the biological agent is selected from the group consisting of a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, small molecule, a fluorescent molecule, and a sugar.
- [00480] Clause 165. The labeled agent of clause 164, wherein the protein is an antibody or a lipoprotein.
- [00481] Clause 166. The labeled agent of clause 164, wherein the polypeptide is a polypeptide toxin.
- [00482] Clause 167. The labeled agent of clause 164, wherein the fluorescent molecule is FAM diamine.
- [00483] Clause 168. The labeled agent of clause 162, wherein the reporter moiety comprises a bioluminescent reporter moiety or fluorescent reporter moiety.

- [00484] Clause 169. The labeled agent of any one of clauses 162, wherein the agent is a non-biological agent.
- [00485] Clause 170. The labeled agent of clause 169, wherein the non-biological agent is a polymer-linked nanoparticle or detergent.
- [00486] Clause 171. The labeled agent of clause 170, wherein the nanoparticle is a gold nanoparticle.
- [00487] Clause 172. The labeled agent of clause 168, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.
- [00488] Clause 173. The labeled agent of clause 172, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.
- [00489] Clause 174. The labeled agent of clause 173, wherein the substrate is furimazine, coelenterazine H,H, luciferin, or pre-luciferin.
- [00490] Clause 175. The labeled agent of clause 174, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.
- [00491] Clause 176. The labeled agent of clause 168, wherein the reporter moiety comprises a fluorescent reporter moiety.
- [00492] Clause 177. The labeled agent of clause 176, wherein the fluorescent reporter moiety comprises a fluorophore.
- [00493] Clause 178. The labeled agent of clause 177, wherein the fluorophore is coumarin, a rhodamine, R110, fluorescein, DDAO, resorufin or cresyl violet.
- [00494] Clause 179. A kit comprising a compound according to any one of clauses 82-117, an agent, or any combination thereof.
- [00495] Clause 180. A kit comprising a labeled agent comprising an agent and a compound according to any one of clauses 82-117.
- [00496] Clause 181. The kit of clause 179 or 180, wherein the agent is a biological agent.
- [00497] Clause 182. The kit of clause 181, wherein the biological agent is selected from the group consisting of a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, small molecule, a fluorescent molecule, and a sugar.
- [00498] Clause 183. The kit of clause 182, wherein the protein is an antibody or a lipoprotein.
- [00499] Clause 184. The kit of clause 179 or 180, wherein the agent is a non-biological agent.

[00500] Clause 185. The kit of clause 184, wherein the non-biological agent is a polymer-linked nanoparticle or detergent.

[00501] Clause 186. The kit of clause 185, wherein the nanoparticle is a gold nanoparticle.

[00502] Clause 187. The kit of clause 179 or 180, wherein the compound is a substrate for luciferase.


[00503] Clause 188. The kit of clause 187, further comprising a detection reagent.

[00504] Clause 189. The kit of clause 188, wherein the detection reagent comprises a luciferase enzyme.

CLAIMS

What is claimed is:

1. A compound of formula (I), or a salt thereof,



(I)

wherein

A is a reporter moiety;

R¹⁴ is H, alkyl, hydroxyalkyl, alkoxy, carboxyalkyl, or amidoalkyl;

R⁹ and R¹⁰ are independently selected from alkyl;

R¹¹, R¹² and R¹³ are independently selected from H, alkyl, alkoxy, bromo, chloro or amino, or R¹¹ and R¹² can form a fused phenyl ring;

X is O;

L is $-(\text{CH}_2)_m\text{C}(\text{R}^{17})_2(\text{CH}_2)_n\text{-Y-C(O)-}$;

R¹⁷ is independently H, alkyl or both R¹⁷ together can form an alkyl ring having from 3-7 carbons;

m is an integer from 0-2;

n is an integer from 0-2;

Y is O or NR¹⁵;

R¹⁵ is H, alkyl, hydroxyalkyl, azidoalkyl, cyanoalkyl, haloalkyl, alkenyl, alkynyl, -alkyl-N(R²³)C(O)R²⁴, -alkyl-SO₃R²⁵, -alkyl-SO₂N(R²⁶)(R²⁷), -alkyl-COR²⁸, -alkyl-CO₂R²⁹, -alkyl-OC(O)R²⁹, -alkyl-OC(O)N(R³⁰)(R³¹), -alkyl-amide, or polyalkoxyalkyl, wherein the polyalkoxyalkyl and -alkyl-amide are unsubstituted or substituted with one or more suitable substituents; and

R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

2. The compound of claim 1, wherein R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is selected from the group consisting of $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-haloalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-hydroxyalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-aminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkylaminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-di(alkyl)aminoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-azidoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-cyanoalkyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkenyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkynyl}$, $-(C_2-C_6\text{-alkoxy})_x\text{-N}(R^{23})C(O)R^{24}$, $-(C_2-C_6\text{-alkoxy})_x\text{-SO}_3R^{25}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-SO}_2N(R^{26})(R^{27})$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-COR}^{28}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-OC}(O)R^{29}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-OC}(O)N(R^{30})(R^{31})$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CO}_2R^{29}$, $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CON}(R^{32})(R^{33})$ and $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-(CO)-NR}^{34}\text{-}(CR^aR^b)_p\text{-NR}^{35}\text{(CO)-T}$, wherein

R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl;

x is an integer selected from 1 to 20;

R^{32} and R^{33} are each independently selected from hydrogen, alkyl, carboxy, a peptide, a drug, a biologically active moiety, and a dye;

R^{34} and R^{35} are each independently selected from hydrogen, and alkyl;

R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;

p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

3. The compound of claim 1, wherein R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-CON}(R^{32})(R^{33})$ or $-(C_2-C_6\text{-alkoxy})_x\text{-alkyl-(CO)-NR}^{34}\text{-}(CR^aR^b)_p\text{-NR}^{35}\text{(CO)-T}$;

x is an integer selected from 1 to 20;

R^{32} and R^{33} are each independently selected from hydrogen, alkyl, carboxy, aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar,

carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof;





R^{34} and R^{35} are each independently selected from hydrogen, and alkyl;

R^a and R^b are each independently selected from hydrogen, alkyl, and carboxy;


p is 0 to 6; and

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof.

4. The compound of claim 1, wherein R^{15} is



wherein




wherein

T is selected from aryl, arylalkyl, cycloalkylalkyl, heteroaryl, heterocycle, hydroxyalkyl, protein, polypeptide, polypeptide-based toxin, amino acid, nucleotide, polynucleotide, lipid, sugar, carbohydrate, enzyme substrate, drug, derivative of a drug, polymer-linked nanoparticle, antibody, detergent, and a dye, or a combination thereof; and

x is 0 to 20.

7. The compound of claim 1, wherein R^{15} is polyalkoxyalkyl, wherein the polyalkoxyalkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of alkyl, alkoxyalkyl, haloalkyl, haloalkoxyalkyl, hydroxyalkyl, aminoalkyl, alkylaminoalkyl, di(alkyl)aminoalkyl, azidoalkyl, cyanoalkyl, alkoxy, haloalkoxy, alkoxyalkoxy, alkenyl, alkynyl, $-N(R^{23})C(O)R^{24}$, $-SO_3R^{25}$, $-SO_2N(R^{26})(R^{27})$, $-COR^{28}$, $-CO_2R^{29}$, and $-CON(R^{30})(R^{31})$, wherein R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , and R^{31} are each independently selected from hydrogen, alkyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl.

8. The compound of claim 1, wherein R^{15} is




wherein


R^{40} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein said aryl, heteroaryl, heterocyclyl, and cycloalkyl are unsubstituted or substituted with one or more suitable substituents; and

x is an integer selected from 1 to 20.

9. The compound of claim 8, wherein R^{40} is hydrogen.
10. The compound of claim 8, wherein R^{40} is a 5- or 6-membered heterocyclyl, having 1, 2, or 3 heteroatoms independently selected from the group consisting of N, O, and S.
11. The compound of claim 8, wherein R^{40} is




12. The compound of claim 8, wherein x is 2, 3, or 4.
13. The compound of claim 8, having formula (I-vi), or a salt thereof,




(I-vi).





14. The compound of claim 1, wherein R^{15} is




each x is independently 0 to 20.


15. The compound of claim 1, selected from the group consisting of:








PBI 5916





PBI 5917




;








PBI 5918


;




























;



























;

;




and

or a salt thereof.

16. The compound of any one of claims 2-6, wherein T is a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, a small molecule, a fluorescent molecule, a polymer-linked nanoparticle, a detergent, or a sugar.
17. The compound of claim 16, wherein T is a protein.
18. The compound of claim 17, wherein the protein is an antibody.
19. The compound of claim 16, wherein T is a polypeptide.
20. The compound of claim 19, wherein the polypeptide is a polypeptide toxin.
21. The compound of claim 16, wherein T is a therapeutic drug.

22. The compound of claim 21, wherein the compound is PBI-5651, PBI-5658, PBI-5666, PBI-5648, PBI-5655, PBI-5662, PBI-5657, PBI-5665, PBI-5667, PBI-5650, PBI-5654, PBI-5661, PBI-5827, PBI-5828, PBI-5829, PBI-5647, PBI-5668, PBI-5659, PBI-5663, PBI-5652, PBI-5669, PBI-5625, PBI-5649, PBI-5653, PBI-5656, PBI-5660, PBI-5664, PBI-5830, PBI-5831 or



23. The compound of claim 16, wherein T is a fluorescent molecule.
24. The compound of claim 23, wherein the fluorescent molecule is FAM diamine.
25. The compound of claim 24, wherein the compound is PBI-5683, PBI-5826, PBI-5682, PBI-5684, PBI-5824, or PBI-5825.
26. The compound of claim 16, wherein T is a polymer-linked nanoparticle or a detergent.
27. The compound of claim 26, wherein the nanoparticle is a gold nanoparticle.
28. The compound of claim 27, wherein the compound is PBI-5907, PBI-5908, PBI-5909, PBI-5910, PBI-5911, or PBI-5912.
29. The compound of claim 1, wherein the reporter moiety comprises a bioluminescent reporter moiety or fluorescent reporter moiety.

30. The compound of claim 29, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.
31. The compound of claim 30, wherein the substrate comprises luciferin, a luciferin derivative or analog, a preluciferin or a preluciferin analog, a coelenterazine or a coelenterazine derivative or analog.
32. The compound of claim 30, wherein the substrate is furimazine, coelenterazine H,H, or luciferin.
33. The compound of claim 30, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.
34. The compound of claim 29, wherein the reporter moiety comprises a fluorescent reporter moiety.
35. The compound of claim 34, wherein the fluorescent reporter moiety comprises a fluorophore.
36. The compound of claim 35, wherein the fluorophore is coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthene or carbopyronine.
37. A method for evaluating cellular uptake of an agent, the method comprising:
a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of claims 1 to 36 or ii) a compound of any one of claims 1-36, wherein the sample comprises a cell; and
b) detecting light emission,
whereby the detection of light emission indicates cellular uptake of the agent.
38. The method of claim 37, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released reporter moiety.
39. The method of claim 38, wherein the reporter moiety comprises a fluorescent reporter moiety.
40. The method of claim 39, wherein detecting light emission comprises exposing the sample to a wavelength of light and detecting fluorescence level in the sample, wherein an increase in

fluorescence or a change in fluorescence wavelength as compared to the fluorescence or fluorescence wavelength of a control sample indicates cellular uptake of the agent.

41. The method of claim 40, wherein the control sample is a sample that is not contacted with a labeled agent.

42. The method of claim 40, wherein fluorescence is detected inside the cell.

43. The method of claim 40, wherein fluorescence is detected outside the cell.

44. The method of claim 38, wherein the reporter moiety comprises a bioluminescent reporter moiety.

45. The method of claim 44, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.

46. The method of claim 45, wherein the cell comprises a luciferase.

47. The method of claim 46, wherein the cell expresses the luciferase.

48. The method of claim 46, wherein detecting light emission comprises detecting the luminescence produced by the luciferase utilizing the released reporter moiety.

49. The method of claim 45, further comprising contacting the sample with a luciferase and wherein detecting light emission comprises detecting the luminescence produced by the luciferase utilizing the released reporter moiety.

50. The method of claim 48 or 49, wherein the light emission is detected inside or outside the cell.

51. A method for evaluating cellular uptake of an agent, the method comprising:

(a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of claims 1-15, 29, or 34-36 or ii) a compound of any one of claims 1-15, 29, or 34-36, wherein the sample comprises a cell, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released fluorescent reporter moiety;

(b) exposing the sample to a wavelength of light to generate fluorescence,

(c) detecting fluorescence in the sample, and

(d) comparing the fluorescence or fluorescent wavelength of the sample to the fluorescence of a control sample, wherein an increase in fluorescence or a change in the fluorescent wavelength as compared to fluorescence or fluorescent wavelength of the control sample indicates cellular uptake of the agent.

52. The method of claim 51, wherein the fluorescent reporter moiety comprises a fluorophore.
53. The method of claim 52, wherein the fluorophore is coumarin, R110, fluorescein, DDAO, resorufin, cresyl violet, silyl xanthen or carbopyronine.
54. The method of claim 51, wherein the fluorescence level is detected inside the cell.
55. The method of claim 51, wherein the fluorescence level is detected in the sample.
56. The method of claim 51, wherein the control sample is sample medium only with labeled agent, a sample with labeled agent and cells but without experimental treatment, or a sample not contacted with labeled agent.
57. A method for evaluating cellular uptake of an agent, the method comprising:
- a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of claims 1-15, or ii) a compound of any one of claims 1-15, wherein the sample comprises a cell, wherein the cell comprises luciferase, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released bioluminescent reporter moiety;
 - b) detecting luminescence in the sample; and
 - c) comparing the luminescence of the sample to luminescence of a control sample, wherein cellular uptake of the agent is indicated if the luminescence of the sample is changed compared to the luminescence of the control sample.
58. The method of claim 57, wherein the luminescence level is produced by the luciferase utilizing the released bioluminescent reporter moiety.
59. The method of claim 58, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.
60. The method of claim 59, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.

61. The method of claim 60, wherein the substrate is furimazine, coelenterazine H,H, lucifeirn, or pre-luciferin.
62. The method of claim 60, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.
63. The method of claim 57, wherein the luminescence is detected inside the cell or detected outside in the cell medium.
64. The method of claim 57, wherein the control sample is a sample that is not contacted with a labeled agent.
65. A method for evaluating cellular uptake of an agent, the method comprising:
- a) contacting a sample with a labeled agent, wherein the labeled agent comprises i) an agent and a compound of any one of claims 1-15 or ii) a compound of any one of claims 1-15, wherein the sample comprises a cell and cell medium, wherein the cellular uptake of the agent results in the reduction of the compound and the generation of a released bioluminescent reporter moiety;
 - b) contacting the sample with a luciferase;
 - c) detecting luminescence in the sample; and
 - d) comparing the luminescence of the sample to luminescence in a control sample, wherein cellular uptake of the agent is indicated if the luminescence in the sample is changed compared to the luminescence of the control sample.
66. The method of claim 65, wherein the luminescence level is produced by the luciferase utilizing the released bioluminescent reporter moiety.
67. The method of claim 66, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.
68. The method of claim 67, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.
69. The method of claim 68, wherein the substrate is furimazine, coelenterazine H,H, luciferin, or pre-luciferin.
70. The method of claim 68, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.

71. The method of claim 65, wherein the luminescence is detected in the cell or detected in the cell medium.
72. The method of claim 65, wherein the control sample is sample medium with labeled agent but without cells, a sample with labeled agent and cells but without experimental treatment, or a sample not contacted with labeled agent.
73. The method of any one of claims 37-72, wherein the agent is a biological agent.
74. The method of claim 73, wherein the biological agent is selected from the group consisting of a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, small molecule, a fluorescent molecule, a nanoparticle, and a sugar.
75. The method of claim 74, wherein the protein is an antibody or a lipoprotein.
76. The method of any one of claims 37-72, wherein the agent is a non-biological agent.
77. The method of claim 76, wherein the non-biological agent is a polymer-linked nanoparticle or a detergent.
78. The method of any one of claims 37-77, wherein the cell is a eukaryotic cell or a prokaryotic cell.
79. The method of claim 78, wherein the cell is in an animal.
80. The method of claim 78, wherein the cell is growing in culture medium.
81. A labeled agent derived from i) an agent and a compound according to any one of claims 1-15 or ii) a compound according to any one of claims 1-36.
82. The labeled agent of claim 81, wherein the agent is a biological agent.
83. The labeled agent of claim 82, wherein the biological agent is selected from the group consisting of a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, small molecule, a fluorescent molecule, a nanoparticle, and a sugar.
84. The labeled agent of claim 83, wherein the protein is an antibody or a lipoprotein.
85. The labeled agent of claim 83, wherein the polypeptide is a polypeptide toxin.
86. The labeled agent of claim 83, wherein the fluorescent molecule is FAM diamine.

87. The labeled agent of claim 81, wherein the reporter moiety comprises a bioluminescent reporter moiety or fluorescent reporter moiety.
88. The labeled agent of any one of claims 81, wherein the agent is a non-biological agent.
89. The labeled agent of claim 88, wherein the non-biological agent is a polymer-linked nanoparticle or a detergent.
90. The labeled agent of claim 89, wherein the nanoparticle is a gold nanoparticle.
91. The labeled agent of claim 87, wherein the bioluminescent reporter moiety comprises a substrate for a luciferase.
92. The labeled agent of claim 91, wherein the substrate comprises luciferin, a luciferin derivative, a coelenterazine or a coelenterazine derivative.
93. The labeled agent of claim 92, wherein the substrate is furimazine, coelenterazine H,H, luciferin, or pre-luciferin.
94. The labeled agent of claim 93, wherein the luciferase is a beetle luciferase, a Renilla luciferase, an Oplophorus luciferase, or a variant thereof.
95. The labeled agent of claim 87, wherein the reporter moiety comprises a fluorescent reporter moiety.
96. The labeled agent of claim 95, wherein the fluorescent reporter moiety comprises a fluorophore.
97. The labeled agent of claim 96, wherein the fluorophore is coumarin, a rhodamine, R110, fluorescein, DDAO, resorufin or cresyl violet.
98. A kit comprising a compound according to any one of claims 1 to 36, an agent, or any combination thereof.
99. A kit comprising a labeled agent comprising an agent and a compound according to any one of claims 1 to 36.
100. The kit of claim 98 or 99, wherein the agent is a biological agent.

101. The kit of claim 100, wherein the biological agent is selected from the group consisting of a protein, a polypeptide, a oligonucleotide, a protein-oligonucleotide conjugate, a lipid, a therapeutic drug, small molecule, a fluorescent molecule, a nanoparticle, and a sugar.
102. The kit of claim 101, wherein the protein is an antibody or a lipoprotein.
103. The kit of claim 98 or 99, wherein the agent is a non-biological agent.
104. The kit of claim 103, wherein the non-biological agent is a polymer-linked nanoparticle or a detergent.
105. The kit of claim 104, wherein the nanoparticle is a gold nanoparticle.
106. The kit of claim 98 or 99, wherein the compound is a substrate for luciferase.
107. The kit of claim 106, further comprising a detection reagent.
108. The kit of claim 107, wherein the detection reagent comprises a luciferase enzyme.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13





FIG. 14



1. Mixed with non-labeled fatty acids, steroids or triglycerides
2. surfactants





FIG. 15



Name	Sequence
Ant-20	NH2RQIKIWFQRRMKWKGGC(COOH)S-S-5'CCA-TCC-CGA-CCT-CGC-GCT-CC-3'-NH-Quinone-label
Ant mismatch	NH2RQIKIWFQRRMKWKGGCCOOHSS'CCA-TAC-CAA-CAT-CAC-GCT-CC-3'-NH-Quinone-label
Tat-20	NH2RKKRRQRRRPPQC(COOH)-S-S-5'CCA-TCCCGA-CCT-CGC-GCT-CC-3'NH-Quinone-label
Tat 20mismatch	NH2RKKRRQRRRPPQC(COOH)-S-S-5'CCA-TAC-CAA-CAT-CAC-GCT-CC-3'NH-Quinone-label
20	5'CAA-TCC-CGA-CCT-CGC-GCT-CC-3'-NH-Quinone-label

FIG. 17

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2015/013617

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D417/12 C07D487/04 C07D417/04 C07D417/14 C07J9/00 C07J19/00 C07D493/10 C12Q1/66 G01N33/58 ADD. According to International Patent Classification (IPC) or to both national classification and IPC					
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D C07J C12Q G01N Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data					
C. DOCUMENTS CONSIDERED TO BE RELEVANT					
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.			
X,P	ROUBINET BENOÎT ET AL: "New insights into the water-solubilization of thiol-sensitive fluorogenic probes based on long-wavelength 7-hydroxycoumarin scaffolds", DYES AND PIGMENTS, vol. 110, 12 February 2014 (2014-02-12), - 12 February 2014 (2014-02-12), pages 270-284, XP029039923, ISSN: 0143-7208, DOI: 10.1016/J.DYEPIG.2014.02.004 abstract page 270 page 280; example 23 ----- -/--	1-108			
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.		<input checked="" type="checkbox"/> See patent family annex.			
* Special categories of cited documents : <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;"> "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td style="width: 50%; border: none; vertical-align: top;"> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family </td> </tr> </table>				"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family				
Date of the actual completion of the international search		Date of mailing of the international search report			
11 March 2015		24/03/2015			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Stix-Malaun, Elke			

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2015/013617

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/033515 A1 (PROMEGA CORP [US]; BENINK HELENE [US]; CALI JAMES J [US]; DUELLMAN SAR) 7 March 2013 (2013-03-07) page 1 examples claims figures	1-108
X	----- US 2012/107849 A1 (KLAUBERT DIETER H [US] ET AL) 3 May 2012 (2012-05-03) page 11 - page 12; examples 4-6 claims 12,41-43	1-108
X	----- US 2010/047839 A1 (HUANG SHENG-TUNG [TW] ET AL) 25 February 2010 (2010-02-25) page 2; example claims figure 3	1-108

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2015/013617

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013033515 A1	07-03-2013	EP 2751089 A1	09-07-2014
		JP 2014526443 A	06-10-2014
		US 2013130289 A1	23-05-2013
		US 2014093894 A1	03-04-2014
		WO 2013033515 A1	07-03-2013

US 2012107849 A1	03-05-2012	AU 2011323455 A1	30-05-2013
		CA 2815381 A1	10-05-2012
		CN 103180324 A	26-06-2013
		EP 2635583 A1	11-09-2013
		JP 2013544242 A	12-12-2013
		KR 20140003430 A	09-01-2014
		SG 189494 A1	31-05-2013
		US 2012107849 A1	03-05-2012
WO 2012061477 A1	10-05-2012		

US 2010047839 A1	25-02-2010	TW 201009336 A	01-03-2010
		US 2010047839 A1	25-02-2010
