



(51) International Patent Classification:

A61K 45/06 (2006.01) *A61K 31/4985* (2006.01)
A61K 31/475 (2006.01) *A61P 35/00* (2006.01)

(21) International Application Number:

PCT/US2015/040029

(22) International Filing Date:

10 July 2015 (10.07.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/024,424 14 July 2014 (14.07.2014) US

(71) Applicant: GILEAD SCIENCES, INC. [US/US]; 333 Lakeside Drive, Foster City, CA 94404 (US).

(72) Inventors: DI PAOLO, Julie; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, CA 94404 (US).
 CLARKE, Astrid; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, CA 94404 (US).

(74) Agents: ECK, Steven, R. et al.; Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, CA 94404 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: COMBINATIONS FOR TREATING CANCERS



(I)

(57) Abstract: Disclosed are combinations for treating a cancer in a subject (e.g., a human) in need thereof, comprising a therapeutically effective amount of a compound of formula (I); or a pharmaceutically acceptable salt thereof, and a vinca-alkaloid, or a pharmaceutically acceptable salt thereof. The subject may be very high risk or high risk for the cancer and may not respond to either agent administered as a sole therapy. The subject who has the cancer may also be refractory to at least one chemotherapy treatment, or is in relapse after treatment with chemotherapy, or both. The cancer may be a hematologic malignancy, such as leukemia or lymphoma, or a solid tumor cancer, such as pancreatic, lung and colon cancer.

COMBINATION METHODS FOR TREATING CANCERS

FIELD


[0001] The present disclosure relates generally to the field of therapeutics and medicinal chemistry, and more specifically to the use of certain Spleen Tyrosine Kinase (Syk) inhibitors in combination with the use of certain vinca-alkaloids in the treatment of cancer including, for example, leukemia, lymphoma and solid-cell tumors.

BACKGROUND

[0002] A number of imidazopyrazine compounds are under investigation for inhibiting Spleen Tyrosine Kinase (Syk) activity. Syk is a non-receptor tyrosine kinase that plays critical roles in immunoreceptor- and integrin-mediated signaling in a variety of cell types, including B-cells, macrophages, monocytes, mast cells, eosinophils, basophils, neutrophils, dendritic cells, T-cells, natural killer cells, platelets, and osteoclasts.

[0003] Syk has been reported to play an important role in signaling through the B-cell receptor, known to be an important survival signal in B-cells. As such, inhibition of Syk activity may be useful for treating certain types of hematologic malignancies. Examples of such hematologic malignancies include cancer, such as B-cell lymphoma and leukemia. Furthermore, there are reports of Syk expression in certain solid cancer (tumor) cell lines. Examples of such solid cancer tumors include pancreatic cancer, lung cancer, colon and colorectal cancer, ovarian cancer and hepatocellular cancer. Additionally, the inhibition of Syk activity is believed to be useful for treating of other diseases and conditions, including inflammatory diseases (*e.g.*, rheumatoid arthritis), allergic disorders and autoimmune diseases.

[0004] One such compound that has been found to inhibit Syk activity is represented by formula I:




or a pharmaceutically acceptable salt thereof. The compound of formula has the chemical name 6-(1H-indazol-6-yl)-N-(4-morpholinophenyl)imidazo[1,2-a]pyrazin-8-amine. This compound and its synthesis have been described in U.S. Patent Nos. 8,450,321 and 8,455,493, which are hereby incorporated by reference in their entirety and specifically with respect to the method of making this compound. *See e.g.*, U.S. Patent No. 8,450,321, Examples 1 and 2.

[0005] Vinca-alkaloids are a subset of drugs derived from the Madagascar periwinkle plant that were discovered in the 1950's, and have a variety of uses from treating diabetes, high blood pressure, and cancer. There are approximately 10 (ten) vinca-alkaloids either currently in use or in development for these indications, including the four major vinca-alkaloids in clinical oncology use: vinblastine, vinorelbine, vincristine, and vindesine. Each of these major vinca-alkaloids has been reported to cause serious side effects, most notably neuropathy. One of the more commonly known vinca-alkaloids is vincristine (VCR), also known as leurocristine and marketed as Oncovin. As with other vinca-alkaloids, vincristine is useful in cancer chemotherapy as a mitotic inhibitor and is commonly used in the standard of care regimen CHOP (Cyclophosphamide, Hydroxydaunorubicin (also known as doxyrubicin), Oncovin (vincristine) and Prednisone) for non-Hodgkin's disease or R-CHOP (CHOP in combination with Rituxan® (also known as rituximab) for B-cell lymphomas. However, vincristine also shares several serious side effects with the other vinca-alkaloids, the most serious of which is chemotherapy-induced peripheral neuropathy, a progressive, enduring, often irreversible neuropathy. This neuropathy can be so severe as to result in the reduction or even cessation of use of vincristine.

[0006] What is desired are methods for treating diseases responsive to the inhibition of Syk in subjects in need of such treatment, including in subjects who may be considered at risk for the disease, are refractory to standard treatments, and/or are in relapse after standard treatments, wherein a treatment regimen of a Syk inhibitor alone does not result in inhibition of cell activity, especially for subjects who may be sensitive to neuropathy secondary to standard dosing levels of vincristine or vinca-alkaloid containing treatment regimens.

SUMMARY

[0007] Provided herein are methods for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I:



or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof.

[0008] In some embodiments, the subject is a human who has a cancer responsive to Syk activity. In another embodiment, the subject is a human who has a solid cancer tumor which expresses Syk. In some embodiments, the subject is a human who has a 17p deletion, a TP53 mutation, NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof. In one embodiment, the subject is a human who has a 17p deletion, a TP53 mutation, or a combination thereof. In another embodiment, the subject is a human who has NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof.

[0009] In some embodiments, the vinca-alkaloid is selected from the group consisting of vincristine, vinblastine, vindesine, vinorelbine, desoxyvincaminol, vincaminol, vinburnine, vincamajine, and vineridine, and pharmaceutically acceptable salts thereof. In certain embodiments, at least one vinca-alkaloid is selected from the group consisting of vincristine, vinblastine, vindesine, vinorelbine, desoxyvincaminol, vincaminol, vinburnine, vincamajine, and vineridine and pharmaceutically acceptable salts thereof. In some embodiments, the vinca-alkaloid is selected from the group consisting of vincristine, vinblastine, vindesine, and vinorelbine, and pharmaceutically acceptable salts thereof. In other embodiments, the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine, and pharmaceutically acceptable salts thereof. In one embodiment, the vinca-alkaloid is vincristine and pharmaceutically acceptable salts thereof. In another embodiment, the vinca-alkaloid is vinblastine and pharmaceutically acceptable salts thereof.


[0010] Provided herein are also methods for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt thereof.

[0011] Provided herein are also methods for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, wherein: the subject is a human who is not undergoing any other anti-cancer treatments; and the subject is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof.


[0012] Provided herein are also figures and examples illustrating that the combination of the compound of formula I, or a pharmaceutically acceptable salt thereof, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt thereto, has unexpected improvements over the effects of the compound of formula I, or the vinca-alkaloid, alone in monotherapy or administered as a sole agent in the treatment of certain cancers and their respective cell lines.

[0013] In some embodiments, the subject is not undergoing any other anti-cancer treatments using one or more PI3K inhibitors. Such PI3K inhibitors may include, in certain embodiments, Compounds A, B and C, whose structures are provided below.


Compound A



Compound B



Compound C



[0014] Compound A has the chemical name (S)-2-(1-((9H-purin-6-yl)amino)propyl)-5-fluoro-3-phenylquinazolin-4(3H)-one, Compound B is named (S)-2-(1-((9H-purin-6-

yl)amino)ethyl)-5-fluoro-3-phenylquinazolin-4(3H)-one, and Compound C is (S)-2-(1-((9H-purin-6-yl)amino)ethyl)-3-(2,6-difluorophenyl)quinazolin-4(3H)-one.

[0015] In some embodiments, the subject is refractory to at least one anti-cancer treatment. In other embodiments, the subject is in relapse after treatment with at least one anti-cancer treatment.

[0016] In some embodiments, about 100 mg to 800 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to subject twice daily. In other embodiments, about 200 mg to 400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the subject twice daily. In one embodiment, about 400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to subject twice daily.

[0017] In one variation, the subject is a human who has a 17p deletion, a TP53 mutation, or a combination thereof; and about 100 mg to 800 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the subject twice daily. In another variation, the subject is a human who has a 17p deletion, a TP53 mutation, or a combination thereof; and about 200 mg to 400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the subject twice daily. In yet another variation, the subject is a human who has a 17p deletion, a TP53 mutation, or a combination thereof; and about 400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the subject twice daily.

[0018] In other embodiments, the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered to the subject once a week at clinically approved or sub-clinically approved amounts. In some embodiments, the amount of the vinca-alkaloid is administered to the subject once a week at a sub-clinically approved amount. In other embodiments, the vinca-alkaloid is vincristine and the amount of vincristine, or a pharmaceutically acceptable salt thereof, is administered at a dose between 0.1 mg-M² and 1.5 mg-M². In other embodiments, the vinca-alkaloid is administered to the subject once a week at a dose of between 0.25 mg-M² and 1.0 mg-M² and the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine. In other embodiments, the vinca-alkaloid is administered to the subject once daily at a dose of between 0.1 mg-M² and 0.2 mg-M² and the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine.

[0019] In certain embodiments, the compound of formula (I) or a pharmaceutically acceptable salt thereof is administered before the vinca-alkaloid, or a pharmaceutically acceptable salt thereof. In other embodiments, the vinca alkaloid, or a pharmaceutically acceptable salt thereof, is administered before the compound of formula (I) or a pharmaceutically acceptable salt thereof.

[0020] In one embodiment, the compound of formula (I) or a pharmaceutically acceptable salt thereof, and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, are administered simultaneously, wherein the vinca-alkaloid is administered via IV and the compound of formula I is administered via tablet. In certain embodiments, the compound of formula I and the at least one vinca-alkaloid, or pharmaceutically acceptable salts thereof, is independently administered twice a day. In other embodiments, the compound of formula (I) or a pharmaceutically acceptable salt thereof, and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, are administered once a day. In other embodiments, the compound of formula (I) or a pharmaceutically acceptable salt thereof, and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, are administered once a week. In one embodiment, the compound of formula (I) or a pharmaceutically acceptable salt thereof, is administered twice a day, and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered once a week. In one embodiment, the compound of formula (I) or a pharmaceutically acceptable salt thereof, is administered once a day, and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered once a week

[0021] In some embodiments, the cancer is a hematologic malignancy. In certain embodiments, the cancer is a leukemia. In one embodiment, the leukemia is chronic lymphocytic leukemia (CLL). In certain embodiments, the cancer is a lymphoma. In one embodiment, the lymphoma is non-Hodgkin's lymphoma (NHL). In one variation, the NHL is diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), follicular lymphoma (FL), small lymphocytic lymphoma (SLL), lymphoplasmacytic lymphoma (LPL), and/or marginal zone lymphoma (MZL). Thus, it is understood that in one aspect the subject is a human who has a hematologic malignancy, such as leukemia or lymphoma.

[0022] In certain embodiments, the cancer is selected from the group consisting of acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), myelodysplastic syndrome (MDS), myeloproliferative disease (MPD), chronic myeloid leukemia (CML), multiple myeloma

(MM), non-Hodgkin's lymphoma (NHL), indolent non-Hodgkin's lymphoma (iNHL), refractory iNHL, mantle cell lymphoma (MCL), follicular lymphoma (FL), Waldstrom's macroglobulinemia (WM), T-cell lymphoma, B-cell lymphoma, diffuse large B-cell lymphoma (DLBCL), lymphoplasmacytic lymphoma (LPL), and marginal zone lymphoma (MZL).

[0023] In some embodiments, the cancer is a solid tumor cancer (or solid cancer tumor). In certain embodiments the cancer is a solid tumor and expresses spleen tyrosine kinase (Syk) activity. In other embodiments, the solid tumor cancer is selected from the group consisting of pancreatic, lung, colorectal cancer, ovarian, breast, esophageal, adenocarcinoma and hepatocellular.

DESCRIPTION OF THE FIGURES

[0024] FIG. 1 depicts the chemical structure of the compound of Formula I.

[0025] FIG. 2 details the inhibition effects of the combination of the compound of formula I and vincristine as compared to vincristine alone in seventeen (17) malignant B-cell lines, representing 4 hematological cancer types: DLBCL, MM, FL, and MCL. (Figures 2(a) – 2(q))

[0026] FIG. 3 summarizes the effect of the combination of the compound of Formula I and vincristine versus the combination of A) the compound of Formula I and combretastatin A4, B) the compound of Formula I and colchicine, C) the compound of Formula I and doxorubicin and D) the compound of formula I and taxol.

[0027] FIG. 4 depicts and summarizes the inhibitory effects of the combination of a compound of Figure 1 and one of two of the vinca-alkaloids, vincristine and vinblastine respectively, in the DLBCL cell line, DHL-10 when compounds were co-administered.

[0028] FIG. 5 summarizes the inhibition of cell viability by the combination of the compound of Formula I and vincristine in the Syk-expressing malignant colon cell line, MiaPaca (FIG 5a), and in the non-Syk expressing malignant colon cell line, HepG2 (FIG 5b).


[0029] FIG. 6 depicts the level of Syk expression in the MiaPaca and HepG2 malignant colon cell lines.

[0030] FIG. 7 summarizes the levels of Syk expression in malignant cell lines from lung, pancreas, and colon.

DETAILED DESCRIPTION

[0031] The following description sets forth exemplary compositions and methods. It should be recognized, however, that such description is not intended as a limitation on the scope of the present disclosure but is instead provided as a description of exemplary embodiments.

[0032] Provided herein are methods for treating cancer in a certain population of subjects (*e.g.*, humans) in need thereof, comprising administering to such subjects a therapeutically effective amount of a compound of formula I:



or a pharmaceutically acceptable salt, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof.

Provided herein are also figures and examples illustrating that the combination of the compound of formula I, or a pharmaceutically acceptable salt thereof, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, has unexpected improvements over the effects of the compound of formula I, or the vinca-alkaloid, alone in monotherapy or administered as a sole agent in the treatment of certain cancers and their respective cell lines.

[0033] The compound of formula I may also be referred to by its compound name: 6-(1H-indazol-6-yl)-N-(4-morpholinophenyl)imidazo[1,2-a]pyrazin-8-amine. The compound name provided is named using ChemBioDraw Ultra 12.0, and one skilled in the art understands that the compound structure may be named or identified using other commonly recognized nomenclature systems and symbols including CAS and IUPAC. One method for

synthesizing the compound of formula I has been previously described in, for example, U.S. Patent No. 8,450,321.

[0034] Any formula or structure given herein, including the compound of formula I and pharmaceutically acceptable salts thereof, is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds, or salts thereof. Isotopically labeled compounds or salts thereof have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the disclosure include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as, but not limited to ^2H (deuterium, D), ^3H (tritium), ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}F , ^{31}P , ^{32}P , ^{35}S , ^{36}Cl and ^{125}I . Various isotopically labeled compounds or salts thereof of the present disclosure, for example those into which radioactive isotopes such as ^3H , ^{13}C and ^{14}C are incorporated. Such isotopically labeled compounds or salts thereof may be useful in metabolic studies, reaction kinetic studies, detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays or in radioactive treatment of subjects (*e.g.* humans).

[0035] The disclosure also includes the compound of formula I and pharmaceutically acceptable salts thereof, in which from 1 to n hydrogens attached to a carbon atom is/are replaced by deuterium, in which n is the number of hydrogens in the molecule. Such compounds may exhibit increased resistance to metabolism and are thus useful for increasing the half-life of the compound of formula I, or pharmaceutically acceptable salts thereof when administered to a mammal. See, for example, Foster, "Deuterium Isotope Effects in Studies of Drug Metabolism", Trends Pharmacol. Sci. 5(12):524-527 (1984). Such compounds are synthesized by means well known in the art, for example by employing starting materials in which one or more hydrogens have been replaced by deuterium.

[0036] Deuterium labeled or substituted therapeutic compounds of the disclosure may have improved DMPK (drug metabolism and pharmacokinetics) properties, relating to distribution, metabolism and excretion (ADME). Substitution with heavier isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life, reduced dosage requirements and/or an improvement in therapeutic index. An ^{18}F labeled compound may be useful for PET or

SPECT studies. Isotopically labeled compounds of this disclosure and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent. It is understood that deuterium in this context is regarded as a substituent in the compound of formula I and pharmaceutically acceptable salts thereof.

[0037] The concentration of such a heavier isotope, specifically deuterium, may be defined by an isotopic enrichment factor. In the compounds of this disclosure any atom not specifically designated as a particular isotope is meant to represent any stable isotope of that atom. Unless otherwise stated, when a position is designated specifically as “H” or “hydrogen”, the position is understood to have hydrogen at its natural abundance isotopic composition. Accordingly, in the compounds or salts thereof of this disclosure any atom specifically designated as a deuterium (D) is meant to represent deuterium.

[0038] Vinca alkaloids, and pharmaceutically acceptable salts thereof, are derived from the Madagascar periwinkle plant, and have been used to treat diabetes, high blood pressure, and various cancers. Examples of vinca-alkaloids include vincristine, vinblastine, vindesine, vinorelbine, desoxyvincaminol, vincaminol, vinburnine, vincamajine, and vineridine. Typically, there have been four major vinca alkaloids in clinical use: vinblastine, vinorelbine, vincristine, and vindesine. All vinca alkaloids are administered intravenously (IV).

[0039] The vinca-alkaloids, and pharmaceutically acceptable salts thereof, of the present disclosure are cytotoxics – they halt the division of cells and cause cell death. During cell division, vinca alkaloid molecules bind to the building blocks of a protein called tubulin, inhibiting its formation. Tubulin protein normally works in cells to create microtubules. These microtubules provide cells with both the structure and flexibility they need to divide and replicate. Without microtubules, cells cannot divide. As opposed to a Syk inhibitor, which inhibits spleen tyrosine kinase, vinca-alkaloids mechanism occupying tubulin’s building block structure, thus preventing, in theory, the formation of microtubules and thus interfering with cancer cells’ ability to divide.

[0040] One of the vinca-alkaloids of this disclosure, vinblastine, inhibits angiogenesis, or the process by which new blood vessels grow from pre-existing ones. Angiogenesis is an essential step in a tumor’s transition to malignancy. Vinblastine is generally applied to treat

Hodgkin's disease, non-Hodgkin's lymphoma, breast cancer, and germ cell tumors. Side effects of vinblastine include: toxicity to white blood cells, nausea, vomiting, constipation, dyspnea, chest or tumor pain, wheezing, and fever. Vinblastine is also occasionally associated with antidiuretic hormone secretion and angina.

[0041] Another vinca alkaloid of this disclosure is vinorelbine, which is similar in its effects to vinblastine. Vinorelbine has exhibited significant antitumor activity in patients with breast cancer and antiproliferation effects on osteosarcoma (bone tumor cells). Vinorelbine treatment can result in side effects including decreased resistance to infection, bruising or bleeding, anemia, constipation, diarrhea, nausea, numbness or tingling in the hands and feet, fatigue (also called peripheral neuropathy), and inflammation at the injection site. Less common side effects include hair loss and allergic reaction.

[0042] Another example or embodiment of the vinca alkaloids of this disclosure is vincristine, or pharmaceutically acceptable salts thereof. Vincristine has a high affinity for tubulin dimers (dimers are building blocks of a protein only two blocks long) and can attach and reattach at different sites quickly, thus in theory preventing the dimers' ability to reassemble (build) the tubules, thus destabilizing the tubulin and inhibiting microtubule formation. Vincristine is FDA approved to treat acute leukemia, rhabdomyosarcoma, neuroblastoma, Wilm's tumor, Hodgkin's disease, and other lymphomas. Vincristine's most common side effects are: peripheral neuropathy, suppression of bone marrow activity, constipation, nervous system toxicity, nausea, and vomiting, with neuropathy being the most common and serious side effect. As a result, there are reports of some subjects being treated with vincristine for oncology having had to stop vincristine treatment.

[0043] The fourth common vinca-alkaloid is vindesine, or pharmaceutically acceptable salts thereof. Vindesine has a serum half-life of only 24 hours, but similar effects (intended and side) to that of vinblastine. Vindesine is commonly administered at a dose of 3 milligrams per square meter of body surface during treatment for melanoma, lung cancers, and (combined with other drugs) uterine cancers. Additional side effects from vindesine include: anemia, blood cell toxicity, fatigue, tingling or pricking sensations in the skin, and skin toxicity

Pharmaceutically Acceptable Salts

[0044] In some embodiments of the methods described herein, a pharmaceutically acceptable salt of the compound of formula I is administered to the subject (*e.g.*, a human).

[0045] As used herein, by “pharmaceutically acceptable” refers to a material that is not biologically or otherwise undesirable, *e.g.*, the material may be incorporated into a pharmaceutical composition administered to a patient without causing any significant undesirable biological effects or interacting in a deleterious manner with any of the other components of the composition in which it is contained. Pharmaceutically acceptable vehicles (*e.g.*, carriers, adjuvants, and/or other excipients) have preferably met the required standards of toxicological and manufacturing testing and/or are included on the Inactive Ingredient Guide prepared by the U.S. Food and Drug administration.

[0046] “Pharmaceutically acceptable salts” include, for example, salts with inorganic acids and salts with an organic acid. Examples of salts may include hydrochlorate, phosphate, diphosphate, hydrobromate, sulfate, sulfinate, nitrate, malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, mesylate, p-toluenesulfonate, 2-hydroxyethylsulfonate, benzoate, salicylate, stearate, and alkanoate (such as acetate, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4). In addition, if the compounds described herein are obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt. Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds. Those skilled in the art will recognize various synthetic methodologies that may be used to prepare nontoxic pharmaceutically acceptable addition salts.

[0047] The terms “effective amount”, “pharmaceutically effective amount”, and “therapeutically effective amount” refer to an amount that may be effective to elicit the desired biological or medical response, including the amount of a compound that, when administered to a subject for treating a disease, is sufficient to effect such treatment for the disease. The effective amount will vary depending on the compound, the disease and its severity and the age, weight, etc., of the subject to be treated. The effective amount can include a range of amounts. A pharmaceutically effective amount includes amounts of an agent which are effective when combined with other agents.

[0048] In some embodiments of the disclosure, the pharmaceutically acceptable salt of the compound of formula I is a bis-mesylate salt. In some embodiments of the disclosure, the pharmaceutically acceptable salt of the vinca-alkaloids is a sulfate salt. In some embodiments of the disclosure, the pharmaceutically acceptable salt of the compound of formula I is a bis-mesylate salt and the pharmaceutically acceptable salt of the vinca-alkaloids is a sulfate salt. In one embodiment of the disclosure, the pharmaceutically acceptable salt of the compound of formula I is a bis-mesylate salt and the vinca-alkaloid is vincristine, wherein the pharmaceutically acceptable salt is vincristine sulfate.

Pharmaceutical Compositions

[0049] In some embodiments of the methods described herein, the compound of formula I, or a pharmaceutically acceptable salt, is present in a pharmaceutical composition comprising the compound of formula I, or a pharmaceutically acceptable salt, and at least one pharmaceutically acceptable vehicle. Pharmaceutically acceptable vehicles may include pharmaceutically acceptable carriers, adjuvants and/or other excipients, and other ingredients can be deemed pharmaceutically acceptable insofar as they are compatible with other ingredients of the formulation and not deleterious to the recipient thereof.

[0050] The pharmaceutical compositions of the compound of formula I described herein can be manufactured using any conventional method, *e.g.*, mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, melt-spinning, spray-drying, or lyophilizing processes. An optimal pharmaceutical formulation can be determined by one of skill in the art depending on the route of administration and the desired dosage. Such formulations can influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the administered agent. Depending on the condition being treated, these pharmaceutical compositions can be formulated and administered systemically or locally.

[0051] The term “carrier” refers to diluents, disintegrants, precipitation inhibitors, surfactants, glidants, binders, lubricants, and other excipients and vehicles with which the compound is administered. Carriers are generally described herein and also in “Remington's Pharmaceutical Sciences” by E.W. Martin. Examples of carriers include, but are not limited to, aluminum monostearate, aluminum stearate, carboxymethylcellulose, carboxymethylcellulose sodium, crospovidone, glyceryl isostearate, glyceryl monostearate,

hydroxyethyl cellulose, hydroxyethyl cellulose, hydroxymethyl cellulose, hydroxyoctacosanyl hydroxystearate, hydroxypropyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, lactose monohydrate, magnesium stearate, mannitol, microcrystalline cellulose, poloxamer 124, poloxamer 181, poloxamer 182, poloxamer 188, poloxamer 237, poloxamer 407, povidone, silicon dioxide, colloidal silicon dioxide, silicone, silicone adhesive 4102, and silicone emulsion. It should be understood, however, that the carriers selected for the pharmaceutical compositions, and the amounts of such carriers in the composition, may vary depending on the method of formulation (*e.g.*, dry granulation formulation, solid dispersion formulation).

[0052] The term “diluent” generally refers to a substance that are used to dilute the compound of interest prior to delivery. Diluents can also serve to stabilize compounds. Examples of diluents may include starch, saccharides, disaccharides, sucrose, lactose, polysaccharides, cellulose, cellulose ethers, hydroxypropyl cellulose, sugar alcohols, xylitol, sorbitol, maltitol, microcrystalline cellulose, calcium or sodium carbonate, lactose, lactose monohydrate, dicalcium phosphate, cellulose, compressible sugars, dibasic calcium phosphate dehydrate, mannitol, microcrystalline cellulose, and tribasic calcium phosphate.

[0053] The term “disintegrant” generally refers to a substance which, upon addition to a solid preparation, facilitates its break-up or disintegration after administration and permits the release of an active ingredient as efficiently as possible to allow for its rapid dissolution. Examples of disintegrants may include maize starch, sodium starch glycolate, croscarmellose sodium, crospovidone, microcrystalline cellulose, modified corn starch, sodium carboxymethyl starch, povidone, pregelatinized starch, and alginic acid.

[0054] The term “precipitation inhibitors” generally refers to a substance that prevents or inhibits precipitation of the active agent from a supersaturated solution. One example of a precipitation inhibitor includes hydroxypropylmethylcellulose (HPMC).

[0055] The term “surfactants” generally refers to a substance that lowers the surface tension between a liquid and a solid that could improve the wetting of the active agent or improve the solubility of the active agent. Examples of surfactants include poloxamer and sodium lauryl sulfate.

[0056] The term “glidant” generally refers to substances used in tablet and capsule formulations to improve flow-properties during tablet compression and to produce an anti-

caking effect. Examples of glidants may include colloidal silicon dioxide, talc, fumed silica, starch, starch derivatives, and bentonite.

[0057] The term “binder” generally refers to any pharmaceutically acceptable film which can be used to bind together the active and inert components of the carrier together to maintain cohesive and discrete portions. Examples of binders may include hydroxypropylcellulose, hydroxypropylmethylcellulose, povidone, copovidone, and ethyl cellulose.

[0058] The term “lubricant” generally refers to a substance that is added to a powder blend to prevent the compacted powder mass from sticking to the equipment during the tableting or encapsulation process. A lubricant can aid the ejection of the tablet from the dies, and can improve powder flow. Examples of lubricants may include magnesium stearate, stearic acid, silica, fats, calcium stearate, polyethylene glycol, sodium stearyl fumarate, or talc; and solubilizers such as fatty acids including lauric acid, oleic acid, and C₈/C₁₀ fatty acid.

Methods of Treatment

[0059] Provided herein are methods for using a compound of formula I, or a pharmaceutically acceptable salt thereof, to selectively or specifically inhibit Syk activity therapeutically or prophylactically, in combination with a vinca-alkaloid, or pharmaceutically acceptable salt thereof, to selectively or specifically inhibit tubulin or microtubule formation therapeutically or prophylactically. The method comprises administering a compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca alkaloid, or a pharmaceutically acceptable salt thereof, to a subject (*e.g.*, a human) in need thereof in an amount sufficient to inhibit Syk activity and/or inhibit tubulin or microtubule formation. The method can be employed to treat subjects (*e.g.*, humans) suffering from, or subject to, a condition whose symptoms or pathology is mediated by Syk expression or activity.

[0060] “Treatment” or “treating” is an approach for obtaining beneficial or desired results including clinical results. Beneficial or desired clinical results may include one or more of the following:

- (i) decreasing one or more symptoms resulting from the disease;

- (ii) diminishing the extent of the disease and/or stabilizing the disease (*e.g.*, delaying the worsening of the disease);
- (iii) delaying the spread (*e.g.*, metastasis) of the disease;
- (iv) delaying or slowing the recurrence of the disease and/or the progression of the disease;
- (v) ameliorating the disease state and/or providing a remission (whether partial or total) of the disease and/or decreasing the dose of one or more other medications required to treat the disease;
- (vi) increasing the quality of life, and/or
- (vii) prolonging survival.

[0061] “Delaying” the development of a disease or condition means to defer, hinder, slow, retard, stabilize, and/or postpone development of the disease or condition. This delay can be of varying lengths of time, depending on the history of the disease or condition, and/or subject being treated. A method that “delays” development of a disease or condition is a method that reduces probability of disease or condition development in a given time frame and/or reduces the extent of the disease or condition in a given time frame, when compared to not using the method. Such comparisons are typically based on clinical studies, using a statistically significant number of subjects. Disease or condition development can be detectable using standard methods, such as routine physical exams, mammography, imaging, or biopsy. Development may also refer to disease or condition progression that may be initially undetectable and includes occurrence, recurrence, and onset.

[0062] The compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca alkaloid, or a pharmaceutically acceptable salt thereof, may, in some embodiments, be administered to a subject (*e.g.*, a human) who is at risk or has a family history of the disease or condition.

[0063] The term “inhibition” indicates a decrease in the baseline activity of a biological activity or process. “Inhibition of activity of Syk activity” refers to a decrease in activity of Syk as a direct or indirect response to the presence of the compound of formula I, or a pharmaceutically acceptable salt thereof, relative to the activity of Syk in the absence of such

compound or a pharmaceutically acceptable salt thereof. In some embodiments, the inhibition of Syk activity may be compared in the same subject prior to treatment, or other subjects not receiving the treatment. "Inhibition of activity of tubulin formation" refers to a decrease in tubulin formation as a direct or indirect response to the presence of a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, relative to the activity of tubulin formation in the absence of such vinca-alkaloid or a pharmaceutically acceptable salt thereof. In some embodiments, the inhibition of tubulin formation may be compared in the same subject prior to treatment, or other subjects not receiving the treatment.

Diseases

[0064] In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is used in the treatment of cancer. In certain embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is used in the treatment of a hematologic malignancy. In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, inhibits the growth or proliferation of cancer cells of hematopoietic origin. In some embodiments, the cancer cells are of lymphoid origin, and in certain embodiments, the cancer cells are related to or derived from B lymphocytes or B lymphocyte progenitors.

[0065] Hematologic malignancies amenable to treatment using the method disclosed in the present disclosure include, without limitation, lymphomas (*e.g.*, malignant neoplasms of lymphoid and reticuloendothelial tissues, such as Burkitt's lymphoma, Hodgkins' lymphoma, non-Hodgkins' lymphomas, lymphocytic lymphomas); multiple myelomas; leukemias (*e.g.*, lymphocytic leukemias, chronic myeloid (myelogenous) leukemias). Other cancer cells, of hematopoietic origin or otherwise, that express Syk also can be treated by administration of the polymorphs and compositions thereof described herein.

[0066] In particular embodiments, the hematologic malignancy is leukemia or lymphoma. In certain embodiments, the hematologic malignancy is acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), myelodysplastic syndrome (MDS), myeloproliferative disease (MPD),

chronic myeloid leukemia (CML), multiple myeloma (MM), non-Hodgkin's lymphoma (NHL), indolent non-Hodgkin's lymphoma (iNHL), refractory iNHL, mantle cell lymphoma (MCL), follicular lymphoma (FL), Waldstrom's macroglobulinemia (WM), T-cell lymphoma, B-cell lymphoma, diffuse large B-cell lymphoma (DLBCL), lymphoplasmacytic lymphoma (LPL), and marginal zone lymphoma (MZL).

[0067] In one embodiment, the cancer is T-cell acute lymphoblastic leukemia (T-ALL), or B-cell acute lymphoblastic leukemia (B-ALL). In another embodiment, the cancer is chronic lymphocytic leukemia (CLL). In yet another embodiment, the cancer is non-Hodgkin's lymphoma (NHL). In one embodiment, the NHL is diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), follicular lymphoma (FL), small lymphocytic lymphoma (SLL), lymphoplasmacytic lymphoma (LPL), and marginal zone lymphoma (MZL). In one embodiment, the cancer is indolent non-Hodgkin's lymphoma (iNHL).

[0068] In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is used in the treatment of a solid tumor cancer. In certain embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is used in the treatment of certain solid tumor cancers, such as pancreatic cancer, lung cancer, colon cancer, colo-rectal cancer, breast cancer, hepatocellular cancer. In certain embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is used in the treatment of certain solid tumor cancers which have an expression of Syk activity or in which Syk is expressed. Other solid tumor cancer cells that express Syk also can be treated by administration of the polymorphs and compositions thereof described herein.

[0069] In yet another aspect, provided are methods of treating a subject (*e.g.*, a human) having a Syk-mediated disorder by administering a compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, to the subject. Provided are also methods of modulating Syk in a subject (*e.g.*, a human) by administering a compound of formula I, or a pharmaceutically acceptable salt thereof, or a

pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, to the subject.

[0070] In any of the methods described herein, the compound of formula I, or a pharmaceutically acceptable salt thereof, may be administered to the individual as a unit dosage, for example in the form of a tablet. In any of the methods described herein, the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, may be administered to the individual via IV (intravenous) delivery.

[0071] Any of the methods of treatment provided herein may be used to treat cancer at an advanced stage. Any of the methods of treatment provided herein may be used to treat cancer at locally advanced stage. Any of the methods of treatment provided herein may be used to treat early stage cancer. Any of the methods of treatment provided herein may be used to treat cancer in remission. In some of the embodiments of any of the methods of treatment provided herein, the cancer has reoccurred after remission. In some embodiments of any of the methods of treatment provided herein, the cancer is progressive cancer.

Subjects

[0072] Any of the methods of treatment provided may be used to treat a subject who has been diagnosed with or is suspected of having cancer. "Subject" refers to an animal, such as a mammal (including a human), that has been or will be the object of treatment, observation or experiment. The methods described herein may be useful in human therapy and/or veterinary applications. In some embodiments, the subject is a mammal. In one embodiment, the subject is a human.

[0073] In some of the embodiments of any of the methods provided herein, the subject is a human who is at risk of developing a cancer (e.g., a human who is genetically or otherwise predisposed to developing a cancer) and who has or has not been diagnosed with the cancer. As used herein, an "at risk" subject is a subject who is at risk of developing cancer (e.g., a hematologic malignancy, or a solid tumor cancer). The subject may or may not have detectable disease, and may or may not have displayed detectable disease prior to the treatment methods described herein. An at risk subject may have one or more so-called risk factors, which are measurable parameters that correlate with development of cancer, such as described herein. A subject having one or more of these risk factors has a higher probability of developing cancer than an individual without these risk factor(s).

[0074] These risk factors may include, for example, age, sex, race, diet, history of previous disease, presence of precursor disease, genetic (*e.g.*, hereditary) considerations, and environmental exposure. In some embodiments, a subject at risk for cancer includes, for example, a subject whose relatives have experienced this disease, and those whose risk is determined by analysis of genetic or biochemical markers. Prior history of having cancer may also be a risk factor for instances of cancer recurrence.

[0075] Provided herein are methods for treating a subject (*e.g.*, a human) who is at “very high risk” or “high risk” for cancer (*e.g.*, a hematologic malignancy). Such subjects may be identified by the presence of certain genetic deletions and/or mutations. In one aspect, a very high risk subject is a human who has a 17p deletion, a TP53 mutation, or a combination thereof. In one aspect, a high risk subject is a human who has NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof. Thus, it is understood that methods of treatment as detailed herein may, in some instances, employ selecting a subject who is at very high risk or at high risk for cancer by detecting the presence or absence of one or more 17p deletion, a TP53 mutation, NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof.

[0076] Provided herein are also methods for treating a subject (*e.g.*, a human) who exhibits one or more symptoms associated with cancer (*e.g.*, a hematologic malignancy or a solid tumor cancer). In some embodiments, the subject is at an early stage of cancer. In other embodiments, the subject is at an advanced stage of cancer.

[0077] Provided herein are also methods for treating a subject (*e.g.*, a human) who is undergoing one or more standard therapies for treating cancer (*e.g.*, a hematologic malignancy or a solid tumor cancer), such as chemotherapy, radiotherapy, immunotherapy, and/or surgery. Thus, in some foregoing embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, administered before, during, or after administration of chemotherapy, radiotherapy, immunotherapy, and/or surgery.

[0078] In certain embodiments, the subject may be a human who is (i) refractory to at least one anti-cancer therapy, or (ii) in relapse after treatment with at least one anti-cancer therapy, or both (i) and (ii). In some of embodiments, the subject is refractory to at least two, at least three, or at least four anti-cancer therapy (including, for example, standard or experimental chemotherapies).

[0079] In certain embodiments, the subject is refractory to at least one, at least two, at least three, or at least four anti-cancer therapy (including, for example, standard or experimental chemotherapy) selected from fludarabine, rituximab, obinutuzumab, alkylating agents, alemtuzumab and other chemotherapy treatments such as CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone); R-CHOP (rituximab-CHOP); hyperCVAD (hyperfractionated cyclophosphamide, vincristine, doxorubicin, dexamethasone, methotrexate, cytarabine); R-hyperCVAD (rituximab-hyperCVAD); FCM (fludarabine, cyclophosphamide, mitoxantrone); R-FCM (rituximab, fludarabine, cyclophosphamide, mitoxantrone); bortezomib and rituximab; temsirolimus and rituximab; temsirolimus and Velcade[®]; Iodine-131 tositumomab (Bexxar[®]) and CHOP; CVP (cyclophosphamide, vincristine, prednisone); R-CVP (rituximab-CVP); ICE (cyclophosphamide, carboplatin, etoposide); R-ICE (rituximab-ICE); FCR (fludarabine, cyclophosphamide, rituximab); FR (fludarabine, rituximab); and D.T. PACE (dexamethasone, thalidomide, cisplatin, Adriamycin[®], cyclophosphamide, etoposide).

[0080] Other examples of chemotherapy treatments (including standard or experimental chemotherapies) are described below. In addition, treatment of certain lymphomas is reviewed in Cheson, B.D., Leonard, J.P., “Monoclonal Antibody Therapy for B-Cell Non-Hodgkin’s Lymphoma” *The New England Journal of Medicine* 2008, 359(6), p. 613-626; and Wierda, W.G., “Current and Investigational Therapies for Patients with CLL” *Hematology* 2006, p. 285-294. Lymphoma incidence patterns in the United States is profiled in Morton, L.M., *et al.* “Lymphoma Incidence Patterns by WHO Subtype in the United States, 1992-2001” *Blood* 2006, 107(1), p. 265-276.

[0081] For example, treatment of non-Hodgkin’s lymphomas (NHL), especially of B-cell origin, include the use of monoclonal antibodies, standard chemotherapy approaches (*e.g.*, CHOP, CVP, FCM, MCP, and the like), radioimmunotherapy, and combinations thereof, especially integration of an antibody therapy with chemotherapy. Examples of unconjugated monoclonal antibodies for Non-Hodgkin’s lymphoma/B-cell cancers include rituximab, alemtuzumab, human or humanized anti-CD20 antibodies, lumiliximab, anti-TRAIL, bevacizumab, galiximab, epratuzumab, SGN-40, and anti-CD74. Examples of experimental antibody agents used in treatment of Non-Hodgkin’s lymphoma/B-cell cancers include ofatumumab, ha20, PRO131921, alemtuzumab, galiximab, SGN-40, CHIR-12.12, epratuzumab, lumiliximab, apolizumab, milatuzumab, and bevacizumab. Examples of

standard regimens of chemotherapy for Non-Hodgkin's lymphoma/B-cell cancers include CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone), FCM (fludarabine, cyclophosphamide, mitoxantrone), CVP (cyclophosphamide, vincristine and prednisone), MCP (mitoxantrone, chlorambucil, and prednisolone), R-CHOP (rituximab plus CHOP), R-FCM (rituximab plus FCM), R-CVP (rituximab plus CVP), and R-MCP (R-MCP). Examples of radioimmunotherapy for Non-Hodgkin's lymphoma/B-cell cancers include yttrium-90-labeled ibritumomab tiuxetan, and iodine-131-labeled tositumomab.

[0082] In another example, therapeutic treatments for mantle cell lymphoma (MCL) include combination chemotherapies such as CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone), hyperCVAD (hyperfractionated cyclophosphamide, vincristine, doxorubicin, dexamethasone, methotrexate, cytarabine) and FCM (fludarabine, cyclophosphamide, mitoxantrone). In addition, these regimens can be supplemented with the monoclonal antibody rituximab (Rituxan) to form combination therapies R-CHOP, hyperCVAD-R, and R-FCM. Other approaches include combining any of the abovementioned therapies with stem cell transplantation or treatment with ICE (ifosfamide, carboplatin and etoposide). Other approaches to treating mantle cell lymphoma includes immunotherapy such as using monoclonal antibodies like Rituximab (Rituxan). Rituximab can be used for treating indolent B-cell cancers, including marginal-zone lymphoma, WM, CLL and small lymphocytic lymphoma. A combination of Rituximab and chemotherapy agents is especially effective. A modified approach is radioimmunotherapy, wherein a monoclonal antibody is combined with a radioisotope particle, such as Iodine-131 tositumomab (Bexxar[®]) and Yttrium-90 ibritumomab tiuxetan (Zevalin[®]). In another example, Bexxar[®] is used in sequential treatment with CHOP. Another immunotherapy example includes using cancer vaccines, which is based upon the genetic makeup of an individual patient's tumor. A lymphoma vaccine example is GTOP-99 (MyVax[®]). Yet other approaches to treating mantle cell lymphoma includes autologous stem cell transplantation coupled with high-dose chemotherapy, or treating mantle cell lymphoma includes administering proteasome inhibitors, such as Velcade[®] (bortezomib or PS-341), or antiangiogenesis agents, such as thalidomide, especially in combination with Rituxan. Another treatment approach is administering drugs that lead to the degradation of Bcl-2 protein and increase cancer cell sensitivity to chemotherapy, such as oblimersen (Genasense) in combination with other chemotherapeutic agents. Another treatment approach includes administering mTOR inhibitors, which can lead to inhibition of cell growth and even cell

death; a non-limiting example is Temsirolimus (CCI-779), and Temsirolimus in combination with Rituxan[®], Velcade[®] or other chemotherapeutic agents.

[0083] Other recent therapies for MCL have been disclosed (*Nature Reviews*; Jares, P. 2007). Such examples include Flavopiridol, PD0332991, R-roscovitine (Selicilib, CYC202), Styryl sulphones, Obatoclax (GX15-070), TRAIL, Anti-TRAIL DR4 and DR5 antibodies, Temsirolimus (CCI-779), Everolimus (RAD001), BMS-345541, Curcumin, Vorinostat (SAHA), Thalidomide, lenalidomide (Revlimid[®], CC-5013), and Geldanamycin (17-AAG).

[0084] Examples of other therapeutic agents used to treat Waldenstrom's Macroglobulinemia (WM) include perifosine, bortezomib (Velcade[®]), rituximab, sildenafil citrate (Viagra[®]), CC-5103, thalidomide, epratuzumab (hLL2- anti-CD22 humanized antibody), simvastatin, enzastaurin, campath-1H, dexamethasone, DT PACE, oblimersen, antineoplaston A10, antineoplaston AS2-1, alemtuzumab, beta alethine, cyclophosphamide, doxorubicin hydrochloride, prednisone, vincristine sulfate, fludarabine, filgrastim, melphalan, recombinant interferon alfa, carmustine, cisplatin, cyclophosphamide, cytarabine, etoposide, melphalan, dolastatin 10, indium In 111 monoclonal antibody MN-14, yttrium Y 90 humanized epratuzumab, anti-thymocyte globulin, busulfan, cyclosporine, methotrexate, mycophenolate mofetil, therapeutic allogeneic lymphocytes, Yttrium Y 90 ibritumomab tiuxetan, sirolimus, tacrolimus, carboplatin, thiotepa, paclitaxel, aldesleukin, recombinant interferon alfa, docetaxel, ifosfamide, mesna, recombinant interleukin-12, recombinant interleukin-11, Bcl-2 family protein inhibitor ABT-263, denileukin diftitox, tanespimycin, everolimus, pegfilgrastim, vorinostat, alvocidib, recombinant flt3 ligand, recombinant human thrombopoietin, lymphokine-activated killer cells, amifostine trihydrate, aminocamptothecin, irinotecan hydrochloride, caspofungin acetate, clofarabine, epoetin alfa, nelarabine, pentostatin, sargramostim, vinorelbine ditartrate, WT-1 analog peptide vaccine, WT1 126-134 peptide vaccine, fenretinide, ixabepilone, oxaliplatin, monoclonal antibody CD19, monoclonal antibody CD20, omega-3 fatty acids, mitoxantrone hydrochloride, octreotide acetate, tositumomab and iodine I-131 tositumomab, motexafin gadolinium, arsenic trioxide, tipifarnib, autologous human tumor-derived HSPPC-96, veltuzumab, bryostatin 1, and PEGylated liposomal doxorubicin hydrochloride, and any combination thereof.

[0085] Examples of therapeutic procedures used to treat WM include peripheral blood stem cell transplantation, autologous hematopoietic stem cell transplantation, autologous bone marrow transplantation, antibody therapy, biological therapy, enzyme inhibitor therapy,

total body irradiation, infusion of stem cells, bone marrow ablation with stem cell support, *in vitro*-treated peripheral blood stem cell transplantation, umbilical cord blood transplantation, immunoenzyme technique, pharmacological study, low-LET cobalt-60 gamma ray therapy, bleomycin, conventional surgery, radiation therapy, and nonmyeloablative allogeneic hematopoietic stem cell transplantation.

[0086] Examples of other therapeutic agents used to treat diffuse large B-cell lymphoma (DLBCL) drug therapies (*Blood* 2005 Abramson, J.) include cyclophosphamide, doxorubicin, vincristine, prednisone, anti-CD20 monoclonal antibodies, etoposide, bleomycin, many of the agents listed for Waldenstrom's, and any combination thereof, such as ICE and R-ICE.

[0087] Examples of other therapeutic agents used to treat chronic lymphocytic leukemia (CLL) (Spectrum, 2006, Fernandes, D.) include Chlorambucil (Leukeran), Cyclophosphamide (Cyloxan, Endoxan, Endoxana, Cyclostin), Fludarabine (Fludara), Pentostatin (Nipent), Cladribine (Leustarin), Doxorubicin (Adriamycin[®], Adriblastine), Vincristine (Oncovin), Prednisone, Prednisolone, Alemtuzumab (Campath, MabCampath), many of the agents listed for Waldenstrom's, and combination chemotherapy and chemoimmunotherapy, including the common combination regimen: CVP (cyclophosphamide, vincristine, prednisone); R-CVP (rituximab-CVP); ICE (ifosfamide, carboplatin, etoposide); R-ICE (rituximab-ICE); FCR (fludarabine, cyclophosphamide, rituximab); and FR (fludarabine, rituximab).


[0088] In yet another aspect, provided is a method of sensitizing a subject (*e.g.*, a human) who is (i) refractory to at least one chemotherapy treatment, or (ii) in relapse after treatment with chemotherapy, or both (i) and (ii), wherein the method comprises administering a compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, to the subject. A subject who is sensitized is a subject who is responsive to the treatment involving administration of the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, to the subject or who has not developed resistance to such treatment. In one embodiment of the disclosure, a subject who is sensitized is a subject who is responsive to the treatment involving administration of the compound of formula I, or a pharmaceutically acceptable salt thereof to the subject or who has not developed resistance to such treatment.

Monotherapy and Combination Therapies


Monotherapy

[0089] In one aspect, the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered as a monotherapy (i.e. the only treatment regimen) to the subject (*e.g.*, a human). Provided herein are methods of treatment in which the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, administered to a subject (*e.g.*, a human) is the only anti-cancer therapy regimen administered to the subject. Provided herein are methods of treatment in which the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, administered to a subject (*e.g.*, a human), wherein the subject is not undergoing any other anti-cancer treatments. In one variation, the subject is not undergoing any other anti-cancer treatments using one or more PI3K inhibitors. Such PI3K inhibitors may include, in certain embodiments, Compounds A, B and C, whose structures are provided below.


Compound A



Compound B



Compound C



In one variation, the subject is not undergoing any other anti-cancer treatments using Compound A, or a pharmaceutically acceptable salt thereof. In another variation, the subject is not undergoing any other anti-cancer treatments using Compound B, or a pharmaceutically acceptable salt thereof. In yet another variation, the subject is not undergoing any other anti-cancer treatments using Compound C, or a pharmaceutically acceptable salt thereof.

[0090] In some embodiments where the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered as a monotherapy treatment regimen to the subject,

the subject may be a human who is (i) refractory to at least one anti-cancer therapy, or (ii) in relapse after treatment with at least one anti-cancer therapy, or both (i) and (ii). In some of embodiments, the subject is refractory to at least two, at least three, or at least four anti-cancer therapy (including, for example, standard or experimental chemotherapies). For example, in certain embodiments, the subject may be a human who is (i) refractory to a therapy using an anti-CD20 antibody, an alkylating agent (*e.g.*, bendamustine), a purine analog (*e.g.*, fludarabine), an anthracycline, or any combination thereof; (ii) in relapse after treatment with an anti-CD20 antibody, an alkylating agent (*e.g.*, bendamustine), a purine analog (*e.g.*, fludarabine), an anthracycline, or any combination thereof, or both (i) and (ii).

[0091] A human subject who is refractory to at least one anti-cancer therapy and/or is in relapse after treatment with at least one anti-cancer therapy, as described above, may have undergone one or more prior therapies. In some embodiments, such subjects have undergone one, two, three, or four, or at least one, at least two, at least three, at least four, or at least five, or between one and ten, between one and nine, between one and eight, between one and seven, between one and six, between one and five, or between one and four, anti-cancer therapies prior to treatment using the methods described herein (*e.g.*, prior to the administration of the compound of formula I, or a pharmaceutically acceptable salt thereof, as a monotherapy).

[0092] It should be understood that when a subject (*e.g.* a human) is treated with the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, as a monotherapy treatment regimen as described by this disclosure, the subject may also undergo one or more other therapies that are not anti-cancer therapies.

[0093] In some embodiments, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein: the vinca-alkaloid is selected from the group consisting of vincristine, vindesine, vinorelbine and vinblastine, and the subject is a human who is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof. In certain other embodiments, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically

effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein the vinca-alkaloid is selected from the group consisting of vincristine, vindesine, vinorelbine and vinblastine, and wherein further the subject is a human who is not undergoing any other anti-cancer treatments; and the subject is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof.

[0094] In some embodiments, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein: the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine, and the subject is a human who is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof. In certain other embodiments, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine, and wherein further the subject is a human who is not undergoing any other anti-cancer treatments; and the subject is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof.

[0095] In one embodiment, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein: the vinca-alkaloid is vincristine, and the subject is a human who is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof. In one other embodiment, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt,

wherein the vinca-alkaloid is vincristine, and wherein further the subject is a human who is not undergoing any other anti-cancer treatments; and the subject is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof.

[0096] In one embodiment, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein: the vinca-alkaloid is vinblastine, and the subject is a human who is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof. In one other embodiment, there is provided a method for treating cancer in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt, wherein the vinca-alkaloid is vinblastine, and wherein further the subject is a human who is not undergoing any other anti-cancer treatments; and the subject is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof.

[0097] In yet other embodiments where the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered as a monotherapy treatment regimen to the subject, the subject may have a 17p deletion, a TP53 mutation, NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof. In one embodiment where the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered as a monotherapy treatment regimen to the subject, the subject has a 17p deletion, a TP53 mutation, or a combination thereof. In another embodiment where the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered as a monotherapy treatment regimen to the subject, the subject has NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof.

Additional Combination therapies

[0098] Provided herein are also methods of treatment in which the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, administered to a subject (*e.g.*, a human) is given to a subject (*e.g.*, a human) in additional combination with one or more additional therapies, including one or more of the anti-cancer therapies described above. Thus, in some embodiments, the method for treating cancer in a subject (*e.g.*, a human) in need thereof, comprises administering to the subject a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, together with one or more additional therapies, which can be useful for treating the cancer. The one or more additional therapies may involve the administration of one or more therapeutic agents.

[0099] In some embodiments, the one or more additional therapies involve the use of a phosphatidylinositol 3-kinase (PI3K) inhibitor, including for example, Compounds A, B and C, or a pharmaceutically acceptable salt of such compounds.

[00100] In other embodiments of the methods described above involving the use of the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, in additional combination with one or more additional therapies, the one or more additional therapies is other than a therapy using Compound A, Compound B, or Compound C, or a pharmaceutically acceptable salt of such compounds. In one embodiment of the methods described above involving the use of the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, in additional combination with one or more additional therapies, the one or more additional therapies is other than a therapy using Compound A, or a pharmaceutically acceptable salt thereof. In another embodiment of the methods described above involving the use of the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, in additional combination with one or more additional therapies, the one or more additional therapies is other than a therapy using Compound B, or a pharmaceutically acceptable salt thereof. In yet another embodiment of the methods described above involving the use of the compound of formula I, or a

pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, in additional combination with one or more additional therapies, the one or more additional therapies is other than a therapy using Compound C, or a pharmaceutically acceptable salt thereof.

[00101] In other embodiments, the one or more additional therapeutic agent may be an inhibitors of lysyl oxidase-like 2 (LOXL2) and a substance that bind to LOXL2, including for example, a humanized monoclonal antibody (mAb) with an immunoglobulin IgG4 isotype directed against human LOXL2.

[00102] In other embodiments, the one or more additional therapeutic agent may be an anti-inflammatory agent. Treatment with the one or more additional therapeutic agent may be prior to, concomitant with, or following treatment with the pharmaceutical composition described herein. In some embodiments, the pharmaceutical composition described herein, is combined with another therapeutic agent in a single dosage form, which is then administered prior to, concomitant with or subsequent to administration with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, of this disclosure. Suitable antitumor therapeutics that may be used in combination with at least one chemical entity described herein include, but are not limited to, chemotherapeutic agents, for example mitomycin C, carboplatin, taxol, cisplatin, paclitaxel, etoposide, doxorubicin, or a combination comprising at least one of the foregoing chemotherapeutic agents. Radiotherapeutic antitumor agents may also be used, alone or in combination with chemotherapeutic agents.

[00103] The compound of formula I, or a pharmaceutically acceptable salt thereof, can be useful as chemosensitizing agents, and, thus, when used in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, can be useful in combination with other chemotherapeutic drugs, in particular, drugs that induce apoptosis.

[00104] A method for increasing sensitivity of cancer cells to chemotherapy, comprising administering to a subject (*e.g.*, human) undergoing chemotherapy a chemotherapeutic agent together with a compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, in an amount sufficient to increase the sensitivity of cancer cells to the chemotherapeutic agent is also provided herein. Examples of other chemotherapeutic drugs that can be used in combination with chemical entities described

herein include topoisomerase I inhibitors (camptothecin or topotecan), topoisomerase II inhibitors (e.g. daunomycin and etoposide), alkylating agents (e.g. cyclophosphamide, melphalan and BCNU), tubulin directed agents (e.g. taxol and vinblastine, or other vinca-alkaloids), and biological agents (e.g. antibodies such as anti CD20 antibody, IDEC 8, immunotoxins, and cytokines). In one embodiment of the method for increasing sensitivity of cancer cells to chemotherapy, the chemotherapeutic agent is other than Compound A, or a pharmaceutically acceptable salt thereof. In another embodiment of the method for increasing sensitivity of cancer cells to chemotherapy, the chemotherapeutic agent is other than Compound B, or a pharmaceutically acceptable salt thereof. In yet another embodiment of the method for increasing sensitivity of cancer cells to chemotherapy, the chemotherapeutic agent is other than Compound C, or a pharmaceutically acceptable salt thereof.

[00105] In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is used in combination with Rituxan® (Rituximab) or other agents that work by selectively depleting CD20+ B-cells.

[00106] Included herein are methods of treatment in which the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered in combination with an anti-inflammatory agent. Anti-inflammatory agents include but are not limited to NSAIDs, non-specific and COX-2 specific cyclooxygenase enzyme inhibitors, gold compounds, corticosteroids, methotrexate, tumor necrosis factor receptor (TNF) receptors antagonists, immunosuppressants and methotrexate. Examples of NSAIDs include, but are not limited to ibuprofen, flurbiprofen, naproxen and naproxen sodium, diclofenac, combinations of diclofenac sodium and misoprostol, sulindac, oxaprozin, diflunisal, piroxicam, indomethacin, etodolac, fenoprofen calcium, ketoprofen, sodium nabumetone, sulfasalazine, tolmetin sodium, and hydroxychloroquine. Examples of NSAIDs also include COX-2 specific inhibitors (i.e., a compound that inhibits COX-2 with an IC₅₀ that is at least 50-fold lower than the IC₅₀ for COX-1) such as celecoxib, valdecoxib, lumiracoxib, etoricoxib and/or rofecoxib.

[00107] In a further embodiment, the anti-inflammatory agent is a salicylate. Salicylates include but are not limited to acetylsalicylic acid or aspirin, sodium salicylate, and choline and magnesium salicylates. The anti-inflammatory agent may also be a corticosteroid. For

example, the corticosteroid may be chosen from cortisone, dexamethasone, methylprednisolone, prednisolone, prednisolone sodium phosphate, and prednisone. In some embodiments, the anti-inflammatory therapeutic agent is a gold compound such as gold sodium thiomalate or auranofin. In some embodiments, the anti-inflammatory agent is a metabolic inhibitor such as a dihydrofolate reductase inhibitor, such as methotrexate or a dihydroorotate dehydrogenase inhibitor, such as leflunomide.

[00108] In some embodiments, combinations in which at least one anti-inflammatory compound is an anti-C5 monoclonal antibody (such as eculizumab or pexelizumab), a TNF antagonist, such as entanercept, or infliximab, which is an anti-TNF alpha monoclonal antibody are used.

[00109] In some embodiments, combinations in which at least one therapeutic agent is an immunosuppressant compound such as methotrexate, leflunomide, cyclosporine, tacrolimus, azathioprine, or mycophenolate mofetil are used.

[00110] It should be understood that any combinations of the additional therapeutic agents described above may be used, as if each and every combination was individually listed. For example, in certain embodiments, the additional therapeutic agents include a PI3K inhibitor and a LOXL2 inhibitor, such as simtuzumab.

Dosing Regimen and Modes of Administration

[00111] In the methods provided herein, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered in a therapeutically effective amount to achieve its intended purpose. As used herein, a “therapeutically effective amount” when referring to the compound of Formula I is an amount sufficient to modulate Syk expression or activity, and thereby treat a subject (*e.g.*, a human) suffering an indication, or to ameliorate or alleviate the existing symptoms of the indication. For example, a therapeutically effective amount of the compound of Formula I or a pharmaceutically acceptable salt thereof may be an amount sufficient to decrease a symptom of a disease or condition responsive to inhibition of Syk activity.

[00112] In the methods provided herein, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered in a therapeutically effective amount to achieve its intended purpose. As used herein, a

“therapeutically effective amount” when referring to a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is an amount sufficient to inhibit tubulin growth or formation, or to inhibit or reduce microtubule formation, or to interfere with spindle formation, and thereby treat a subject (e.g. a human) suffering an indication, or to ameliorate or alleviate the existing symptoms of the indication. For example, a therapeutically effective amount of a vinca-alkaloid or a pharmaceutically acceptable salt thereof may be an amount sufficient to decrease a symptom of a disease or condition responsive to inhibit of tubulin activity and/or formation.

[00113] Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. In some embodiments, a therapeutically effective amount of the compound of formula I, or a pharmaceutically acceptable salt thereof, may (i) reduce the number of cancer cells; (ii) reduce tumor size; (iii) inhibit, retard, slow to some extent, and preferably stop cancer cell infiltration into peripheral organs; (iv) inhibit (*e.g.*, slow to some extent and preferably stop) tumor metastasis; (v) inhibit tumor growth; (vi) delay occurrence and/or recurrence of a tumor; and/or (vii) relieve to some extent one or more of the symptoms associated with the cancer. In various embodiments, the amount is sufficient to ameliorate, palliate, lessen, and/or delay one or more of symptoms of cancer.

[00114] The therapeutically effective amount may vary depending on the subject, and disease or condition being treated, the weight and age of the subject, the severity of the disease or condition, and the manner of administering, which can readily be determined by one or ordinary skill in the art.

[00115] The dosing regimen of the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with a vinca-alkaloid, or a pharmaceutically acceptable salt thereof, in the methods provided herein may vary depending upon the indication, route of administration, and severity of the condition, for example. Depending on the route of administration, a suitable dose can be calculated according to body weight, body surface area, or organ size. The final dosing regimen is determined by the attending physician in view of good medical practice, considering various factors that modify the action of drugs, *e.g.*, the specific activity of the compound, the identity and severity of the disease state, the responsiveness of the patient, the age, condition, body weight, sex, and diet of the patient, and the severity of any infection. Additional factors that can be taken into account include

time and frequency of administration, drug combinations, reaction sensitivities, and tolerance/response to therapy. Further refinement of the doses appropriate for treatment involving any of the formulations mentioned herein is done routinely by the skilled practitioner without undue experimentation, especially in light of the dosing information and assays disclosed, as well as the pharmacokinetic data observed in human clinical trials. Appropriate doses can be ascertained through use of established assays for determining concentration of the agent in a body fluid or other sample together with dose response data.

[00116] The formulation and route of administration chosen may be tailored to the individual subject, the nature of the condition to be treated in the subject, and generally, the judgment of the attending practitioner. For example, the therapeutic index of the compound of formula I, or a pharmaceutically acceptable salt thereof, may be enhanced by modifying or derivatizing the compound for targeted delivery to cancer cells expressing a marker that identifies the cells as such. For example, the compounds can be linked to an antibody that recognizes a marker that is selective or specific for cancer cells, so that the compounds are brought into the vicinity of the cells to exert their effects locally, as previously described. *See e.g.*, Pietersz et al., *Immunol. Rev.*, 129:57 (1992); Trail et al., *Science*, 261:212 (1993); and Rowlinson-Busza et al., *Curr. Opin. Oncol.*, 4:1142 (1992). A similar analysis may be applied to treatment with the vinca-alkaloid, or its pharmaceutically acceptable salt, of this disclosure and methods herein.

Dosing Regimen

[00117] The therapeutically effective amount of the compound of formula I, or a pharmaceutically acceptable salt thereof, may be provided in a single dose or multiple doses to achieve the desired treatment endpoint. As used herein, “dose” refers to the total amount of an active ingredient (*e.g.*, the compound of formula I, or a pharmaceutically acceptable salt thereof,) to be taken each time by a subject (*e.g.*, a human).

[00118] Exemplary doses of the compound of formula I, or a pharmaceutically acceptable salt thereof, for a human subject may be between about 0.01 mg to about 1800 mg, or between about 0.01 mg to about 1500 mg, or between about 10 mg to about 1500 mg, or between about 10 mg to about 1300 mg, or between about 10 mg to about 1000 mg, or between about 10 mg to about 800 mg, or between about 10 mg to about 600 mg, or between about 10 mg to about 300 mg, or between about 10 mg to about 200 mg, or between about 10

mg to about 100 mg, or between about 100 mg to about 800 mg, or between about 100 mg to about 600 mg, or between about 100 mg to about 300 mg, or between about 100 mg to about 200 mg, or between about 200 mg to about 350 mg, or between about 250 mg to about 300 mg, or between about 200 mg to about 400 mg, or between about 400 mg to about 600 mg, or between about 400 mg to about 800 mg, or between about 600 mg or about 800 mg, or between about 800 mg to about 1200 mg, or between about 1200 mg to about 1600, or between about 50 mg to about 200 mg, or about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 125 mg, or about 150 mg, or about 175 mg, about 200 mg, about 250 mg, about 300 mg, about 350 mg, about 400 mg, about 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, or about 750 mg, or about 800 mg, or about 850 mg, or about 900 mg, or about 950 mg, or about 1000 mg, or about 1100 mg, or about 1200 mg, or about 1300 mg, or about 1400 mg, or about 1500 mg, or about 1600 mg, or about 1800 mg. In one embodiment, the dose of the compound of formula I, or a pharmaceutically acceptable salt thereof, administered to the subject in the methods provided herein is about 400 mg. In one embodiment, the dose of the compound of formula I, or a pharmaceutically acceptable salt thereof, administered to the subject in the methods provided herein is about 800 mg.

[00119] In other embodiments, the methods provided comprise continuing to treat the subject (*e.g.*, a human) by administering the doses of the compound of formula I, or a pharmaceutically acceptable salt thereof, at which clinical efficacy is achieved or reducing the doses by increments to a level at which efficacy can be maintained. In a particular embodiment, the methods provided comprise administering to the subject (*e.g.*, a human) an initial daily dose of 100 mg to 1000 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, and administering subsequent daily doses of the compound of formula I, or a pharmaceutically acceptable salt thereof, over at least 6 days, wherein each subsequent daily dose is increased by 50 mg to 400 mg. Thus, it should also be understood that the dose of the compound of formula I, or a pharmaceutically acceptable salt thereof, may be increased by increments until clinical efficacy is achieved. Increments of about 25 mg, about 50 mg, about 100 mg, or about 125mg, or about 150 mg, or about 200 mg, or about 250 mg, or about 300 mg can be used to increase the dose. The dose can be increased daily, every other day, two, three, four, five or six times per week, or once per week.

[0100] The frequency of dosing will depend on the pharmacokinetic parameters of the compound administered, the route of administration, and the particular disease treated. The

dose and frequency of dosing may also depend on pharmacokinetic and pharmacodynamic, as well as toxicity and therapeutic efficiency data. For example, pharmacokinetic and pharmacodynamic information about the compound of formula I, or a pharmaceutically acceptable salt thereof, can be collected through preclinical *in vitro* and *in vivo* studies, later confirmed in humans during the course of clinical trials. Thus, for the compound of formula I, or a pharmaceutically acceptable salt thereof, used in the methods provided herein, a therapeutically effective dose can be estimated initially from biochemical and/or cell-based assays. Then, dosage can be formulated in animal models to achieve a desirable circulating concentration range that modulates Syk expression or activity. As human studies are conducted further information will emerge regarding the appropriate dosage levels and duration of treatment for various diseases and conditions.

[0101] Toxicity and therapeutic efficacy of the compound of formula I, or a pharmaceutically acceptable salt thereof, can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the “therapeutic index”, which typically is expressed as the ratio LD₅₀/ED₅₀. Compounds that exhibit large therapeutic indices, *i.e.*, the toxic dose is substantially higher than the effective dose, are preferred. The data obtained from such cell culture assays and additional animal studies can be used in formulating a range of dosage for human use. The doses of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity.

[0102] The administration of the compound of formula I, or a pharmaceutically acceptable salt thereof, may be administered under fed conditions. The term fed conditions or variations thereof refers to the consumption or uptake of food, in either solid or liquid forms, or calories, in any suitable form, before or at the same time when the compounds or pharmaceutical compositions thereof are administered. For example, the compound of formula I, or a pharmaceutically acceptable salt thereof, may be administered to the subject (*e.g.*, a human) within minutes or hours of consuming calories (*e.g.*, a meal). In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, may be administered to the subject (*e.g.*, a human) within 5-10 minutes, about 30 minutes, or about 60 minutes consuming calories.

[0103] The therapeutically effective amount of the vinca-alkaloid of this disclosure, or its pharmaceutically acceptable salt thereof, may be provided in a single dose or multiple doses to achieve the desired treatment endpoint. As used herein, “dose” refers to the total amount of an active ingredient (*e.g.*, vincristine or vinblastine, for example), or a pharmaceutically acceptable salt thereof,) to be taken each time by a subject (*e.g.*, a human).

[0104] Exemplary doses of the vinca-alkaloid of this disclosure, or its pharmaceutically acceptable salt thereof, for a human subject may be between about 0.01 mg-M² to about 3.0 mg-M², depending on the identity of the vinca-alkaloid, or between about 0.01 mg-M² to about 2.5 mg-M², or between about 0.01 mg-M² to about 2.0 mg-M², or between about 0.01 mg-M² to about 1.9 mg-M², or between about 0.01 mg-M² to about 1.8 mg-M², or between about 0.01 mg-M² to about 1.7 mg-M², or between about 0.01 mg-M² to about 1.6 mg-M², or between about 0.01 mg-M² to about 1.5 mg-M², or between about 0.01 mg-M² to about 1.4 mg-M², or between about 0.01 mg-M² to about 1.3 mg-M², or between about 0.01 mg-M² to about 1.2 mg-M², or between about 0.01 mg-M² to about 1.1 mg-M², or between about 0.01 mg-M² to about 1.0 mg-M², or between about 0.01 mg-M² to about 0.9 mg-M², or between about 0.01 mg-M² to about 0.8 mg-M², or between about 0.01 mg-M² to about 0.7 mg-M², or between about 0.01 mg-M² to about 0.6 mg-M², or between about 0.01 mg-M² to about 0.5 mg-M², or between about 0.01 mg-M² to about 0.45 mg-M², or between about 0.01 mg-M² to about 0.4 mg-M², or between about 0.01 mg-M² to about 0.35 mg-M², or between about 0.01 mg-M² to about 0.33 mg-M², or between about 0.01 mg-M² to about 0.3 mg-M², or between about 0.01 mg-M² to about 0.25 mg-M², or between about 0.01 mg-M² to about 0.2 mg-M², or between about 0.01 mg-M² to about 0.15 mg-M², or between about 0.01 mg-M² to about 0.01 mg-M², or between about 0.1 mg-M² to about 1.8 mg-M², or between about 0.15 mg-M² to about 1.7 mg-M², or between about 0.2 mg-M² to about 1.6 mg-M², or between about 0.25 mg-M² to about 1.5 mg-M², or between about 0.3 mg-M² to about 1.4 mg-M², or between about 0.33 mg-M² to about 1.3 mg-M², or between about 0.35 mg-M² to about 1.2 mg-M², or between about 0.4 mg-M² to about 1.1 mg-M², or between about 0.45 mg-M² to about 1.0 mg-M², or between about 0.5 mg-M² to about 0.9 mg-M², or between about 0.6 mg-M² to about 0.8 mg-M². In one embodiment, the dose of the vinca-alkaloid of this disclosure, or its pharmaceutically acceptable salt thereof, administered to the subject in the methods provided herein is about 1.5 mg-M². In one embodiment, the dose of the vinca-alkaloid of this disclosure, or its pharmaceutically acceptable salt thereof, administered to the subject in the methods provided herein is about 1.0 mg-M². In one embodiment, the dose of the vinca-

alkaloid of this disclosure, or its pharmaceutically acceptable salt thereof, administered to the subject in the methods provided herein is about 0.5 mg-M².

[0105] In other embodiments, the methods provided comprise continuing to treat the subject (*e.g.*, a human) by administering the doses of the vinca-alkaloid of this disclosure, or a pharmaceutically acceptable salt thereof, at which clinical efficacy is achieved or reducing the doses by increments to a level at which efficacy can be maintained. The frequency of dosing will depend on the pharmacokinetic parameters of the compound administered, the route of administration, and the particular disease treated. The dose and frequency of dosing may also depend on pharmacokinetic and pharmacodynamic, as well as toxicity and therapeutic efficiency data.

Modes of Administration

[0106] The pharmaceutical compositions of the compound of formula I may be administered in either single or multiple doses by any of the accepted modes of administration of agents having similar utilities, for example as described in those patents and patent applications incorporated by reference, including rectal, buccal, intranasal and transdermal routes, by intra-arterial injection, intravenously, intraperitoneally, parenterally, intramuscularly, subcutaneously, orally, topically, as an inhalant, or via an impregnated or coated device such as a stent, for example, or an artery-inserted cylindrical polymer.

[0107] One mode for administration is parenteral, particularly by injection. The forms in which the compound of formula I, or a pharmaceutically acceptable salt thereof, may be incorporated for administration by injection include aqueous or oil suspensions, or emulsions, with sesame oil, corn oil, cottonseed oil, or peanut oil, as well as elixirs, mannitol, dextrose, or a sterile aqueous solution, and similar pharmaceutical vehicles. Aqueous solutions in saline may also conventionally be used for injection. Ethanol, glycerol, propylene glycol, liquid polyethylene glycol, and the like (and suitable mixtures thereof), cyclodextrin derivatives, and vegetable oils may also be employed. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like.

[0108] Sterile injectable solutions are prepared by incorporating a compound according to the present disclosure in the required amount in the appropriate solvent with various other ingredients as enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. In certain embodiments, for parenteral administration, sterile injectable solutions are prepared containing a therapeutically effective amount, e.g., 0.1 to 1000 mg, of the compound of formula I, or a pharmaceutically acceptable salt thereof. It will be understood, however, that the amount of the compound actually administered usually will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

[0109] Oral administration is another route for administration of the compound of formula I, or a pharmaceutically acceptable salt thereof. Administration may be via capsule or enteric coated tablets, or the like. In making the pharmaceutical compositions that include the compound of formula I, or a pharmaceutically acceptable salt thereof, the active ingredient is usually diluted by an excipient and/or enclosed within such a carrier that can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be in the form of a solid, semi-solid, or liquid material (as above), which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, sterile injectable solutions, and sterile packaged powders.

[0110] Some examples of suitable excipients in an oral formulation include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally

include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl and propylhydroxybenzoates; sweetening agents; and flavoring agents.

[0111] The pharmaceutical compositions of the compound of formula I described herein can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art. Controlled release drug delivery systems for oral administration include osmotic pump systems and dissolutional systems containing polymer-coated reservoirs or drug-polymer matrix formulations. Examples of controlled release systems are given in U.S. Patent Nos. 3,845,770; 4,326,525; 4,902,514; and 5,616,345. Another formulation for use in the methods of the present disclosure employs transdermal delivery devices (patches). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present disclosure in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent Nos. 5,023,252, 4,992,445 and 5,001,139. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

[0112] In some embodiments, the compositions of the compound of formula I described herein are formulated in a unit dosage form. The term “unit dosage forms” refers to physically discrete units suitable as unitary dosages for human subjects, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient (e.g., a tablet, capsule, ampoule).

[0113] In other embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered orally at a unit dosage of about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 250 mg, about 300 mg, about 350 mg, about 400 mg, about 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 800 mg, about 900 mg, about 1100 mg, about 1200 mg, about 1300 mg, about 1400 mg, about 1500 mg, or about 1600 mg, about 1700 mg, or about 1800 mg. In other embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered orally at a unit dosage of about 200 mg, about 600 mg, or about 800 mg, or about 900 mg, or about 1200 mg. In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a

pharmaceutical composition thereof, is administered orally at a unit dosage of about 200 mg, or about 800 mg.

[0114] The dosages for oral administration described above for the compound of formula I may be administered once daily or twice daily (BID). For example, in certain embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered orally at a unit dosage of about 50 mg BID, about 100 mg BID, about 150 mg BID, about 200 mg BID, about 250 mg BID, about 300 mg BID, about 350 mg BID, about 400 mg BID, about 450 mg BID, about 500 mg BID, about 550 mg BID, about 600 mg BID, about 650 mg BID, about 700 mg BID, about 800 mg BID, about 900 mg BID, about 1100 mg BID, about 1200 mg BID, about 1300 mg BID, about 1400 mg BID, about 1500 mg BID, or about 1600 mg BID, about 1700 mg BID, or about 1800 mg BID. In other embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered orally at a unit dosage of about 200 mg BID, about 400 mg BID, or about 600 mg BID, or about 800 mg BID, or about 1000 mg BID. In some embodiments, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered orally at a unit dosage of about 200 mg BID, or about 800 mg BID. In one embodiment, the compound of formula I, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof, is administered orally at a unit dosage of about 800 mg BID.

[0115] The vinca-alkaloid of the disclosure, or the pharmaceutically acceptable salts thereof, are administered via IV. In one embodiment, the vinca-alkaloid is vincristine sulfate and the amount of the vial is 1mg/1ml. In some embodiments, the vinca-alkaloid is vincristine sulfate and the vial is 2ml containing either 1mg or 2mg of vincristine sulfate. In another embodiment, "Vincristine Sulfate", USP is a white to off-white powder. It is soluble in methanol, freely soluble in water, but only slightly soluble in 95% ethanol. In 98% ethanol, Vincristine Sulfate, USP has an ultraviolet spectrum with maxima at 221 nm (ϵ +47,100).

[0116] "Vincristine Sulfate Injection", USP is a sterile, preservative-free, single use only solution available for intravenous use in 2 mL (1 mg and 2 mg) vials. Each mL contains 1 mg Vincristine Sulfate, USP, 100 mg mannitol and Water for Injection, USP. Q.S. Sulfuric acid or sodium hydroxide have been added for pH control. The pH of Vincristine Sulfate Injection, USP ranges from 4.0 to 5.0.

Articles of Manufacture and Kits

[0117] Compositions (including, for example, formulations and unit dosages) comprising the compound of formula I, or a pharmaceutically acceptable salt thereof, can be prepared and placed in an appropriate container, and labeled for treatment of an indicated condition. Accordingly, provided is also an article of manufacture, such as a container comprising a unit dosage form of the compound of formula I, or a pharmaceutically acceptable salt thereof, and a label containing instructions for use of the compounds. In some embodiments, the article of manufacture is a container comprising a unit dosage form of the compound of formula I, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable vehicle. In another embodiment, the article of manufacture is a container comprising a unit dosage form of the compound of formula I, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable vehicle, and a vial containing a vinca-alkaloid, or a pharmaceutically acceptable salt.

[0118] Kits also are contemplated. For example, a kit can comprise unit dosage forms of the compound of formula I, or a pharmaceutically acceptable salt thereof, and a package insert containing instructions for use of the composition in treatment of a medical condition. In some embodiments, the kits comprises a unit dosage form of the compound of formula I, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable vehicle, and a vial containing a solution of vinca-alkaloid, or a pharmaceutically acceptable salt thereof.

[0119] The instructions for use in the kit may be for treating a cancer, including, for example, a hematologic malignancy or solid tumor cancer malignancy. In some embodiments, the instructions for use in the kit may be for treating cancer, such as leukemia or lymphoma, including relapsed and refractory leukemia or lymphoma. In certain embodiments, the instructions for use in the kit may be for treating acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), myelodysplastic syndrome (MDS), myeloproliferative disease (MPD), chronic myeloid leukemia (CML), multiple myeloma (MM), non-Hodgkin's lymphoma (NHL), indolent non-Hodgkin's lymphoma (iNHL), refractory iNHL, mantle cell lymphoma (MCL), follicular lymphoma (FL), Waldstrom's macroglobulinemia (WM), T-cell lymphoma, B-cell lymphoma, diffuse large B-cell lymphoma (DLBCL), lymphoplasmacytic lymphoma (LPL), or marginal zone lymphoma (MZL). In one

embodiment, the instructions for use in the kit may be for treating chronic lymphocytic leukemia (CLL) or non-Hodgkin's lymphoma (NHL). In one embodiment, the NHL is diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), follicular lymphoma (FL), small lymphocytic lymphoma (SLL), lymphoplasmacytic lymphoma (LPL), and marginal zone lymphoma (MZL). In one embodiment, the hematologic malignancy is indolent non-Hodgkin's lymphoma (iNHL). In certain embodiments, diseases or conditions indicated on the label can include, for example, treatment of cancer.

[0120] In certain embodiments of the article of manufacture or the kit, the unit dosage for the compound of formula I has about 50 mg, about 100 mg, about 150 mg, about 200 mg, about 250 mg, about 300 mg, about 350 mg, about 400 mg, about 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 800 mg, about 900 mg, about 1100 mg, about 1200 mg, about 1300 mg, about 1400 mg, about 1500 mg, or about 1600 mg, about 1700 mg, or about 1800 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof. In one embodiment, the unit dosage has about 400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof. In one embodiment of the article of manufacture or the kit, the unit dosage of the compound of formula I, or a pharmaceutically acceptable salt thereof, is a tablet.

EXAMPLES

[0121] The following examples are included to illustrate embodiments of the disclosure, and are not intended to limit the scope of the disclosure. It should be appreciated by those of skill in the art that the techniques disclosed herein represent techniques that apply in the practice of the disclosure. Those of skill in the art would appreciate that, in light of the present disclosure, changes can be made in the examples herein without departing from the spirit and scope of the disclosure.

Example 1

Effect of the Compound of Formula I Compared to Doxorubicin, Vincristine and Cyclophosphamide in 10 Diffuse large B-cell lymphoma (DLBCL), 4 multiple myeloma (MM), 2 follicular lymphoma (FL), and 1 mantle cell lymphoma (MCL) cell lines

[0122] This Example evaluates the efficacy of the compound of Formula I (Figure 1), or a pharmaceutically acceptable salt thereof, to inhibit malignant B-cell viability in 10 diffuse large B-cell lymphoma (DLBCL), 4 multiple myeloma (MM), 2 follicular lymphoma (FL), and 1 mantle cell lymphoma (MCL) cell lines. This Example also evaluates and compares the efficacy of doxorubicin, vincristine and cyclophosphamide to inhibit malignant B-cell viability in the same 10 diffuse large B-cell lymphoma (DLBCL), 4 multiple myeloma (MM), 2 follicular lymphoma (FL), and 1 mantle cell lymphoma (MCL) cell lines, to the efficacy of the compound of formula I, or a pharmaceutically acceptable salt thereof.

[0123] Cell Titer Glo Viability Assay: All procedures were performed at Gilead Sciences, Inc. in Branford, The compound of formula (I), cyclophosphamide, doxorubicin and vincristine sulfate were dissolved in DMSO to prepare stock solutions that were serially diluted three-fold in DMSO in a 96 well plate format at 1000X so the final starting assay concentrations tested would be 10 μ M, 200 μ M, 10 μ M, and 1 μ M respectively, for each compound. Compound plates were diluted 1:100 in RPMI without serum or additives creating a 10X stock. Cells were seeded into a 96-well plate at 10,000 – 20,000 cells per well in 100 μ l of growth media appropriate to the cell line supplemented with 100 U/L penicillin-streptomycin. OCI-Ly3, OCI-Ly4 and OCI-Ly7 were grown in Iscove's + 20% FBS, OCI-Ly19 was grown in alphaMEM + 20% FBS, and all other cell lines were grown in RPMI + 10% FBS. To each well, 11 μ l of 10X compound was added in a final concentration of 0.1% DMSO and cells were incubated at 37°C in 5% CO₂ for 72 hours. Cell viability was assessed with Cell Titer Glo (Promega, Madison, WI) following the manufacturer's protocol on an Envision plate reader (Perkin Elmer, Waltham, MA). EC₅₀ values were determined using a four parameter variable slope model with GraphPad Prism 6.0 software. All EC₅₀ values represent mean values of from duplicate dose response curves. The results are reported in Table 1 below.

[0124] Table 1 summarizes the efficacy of the four compounds tested separately in the Cell Titer Glow assay for each respective cell line. The test results for the compound of formula I and the test results for cyclophosphamide did not show inhibition cell viability in any of the cell lines tested at concentrations <3000 nM and <10000 nM, respectively. Doxorubicin and vincristine EC₅₀ values to inhibit cell viability were calculated and ranged from 3.1 - 557 nM and < 1 nM -103 nM respectively, as shown in Table 1.

Table 1

Cell line	Disease	Inhibition of Viability, EC ₅₀ (nM)			
		Compound of Formula I	Vincristine	Doxorubicin	Cyclophosphamide
SU-DHL-2	DLBCL	>3000	65	216	>10000
SU-DHL-4	DLBCL	>3000	75	215	>10000
SU-DHL-6	DLBCL	>3000	12	125	>10000
SU-DHL-10	DLBCL	>3000	45	195	>10000
Karpas 422	DLBCL	>3000	62	305	>10000
OCI-LY3	DLBCL	>3000	12	26	>10000
OCI-LY4	DLBCL	>3000	< 1	37	>10000
OCI-LY7	DLBCL	>3000	4.6	199	>10000
OCI-LY19	DLBCL	>3000	1.2	3.1	>10000
Pfeiffer	DLBCL	>3000	4.0	198	>10000
Mino	MCL	>3000	42	114	>10000
KMS-11	MM	>3000	103	469	>10000
MM1.S	MM	>3000	59	66	>10000
OPM-2	MM	>3000	97	557	>10000
RPMI-8266	MM	>3000	67	176	>10000
WSU-FSCCL	FL	>3000	58	52	>10000
WSU-NHL	FL	>3000	39	128	>10000

Example 2

Comparison Effect of the Compound of Formula I in Combination with Vincristine versus Effect of Vincristine alone in 10 Diffuse large B-cell lymphoma (DLBCL), 4 multiple myeloma (MM), 2 follicular lymphoma (FL), and 1 mantle cell lymphoma (MCL) cell lines

[0125] This Example evaluates the efficacy of the compound of Formula I, or a pharmaceutically acceptable salt thereof, in combination with vincristine to inhibit malignant B-cell viability in 10 diffuse large B-cell lymphoma (DLBCL), 4 multiple myeloma (MM), 2 follicular lymphoma (FL), and 1 mantle cell lymphoma (MCL) cell lines. This Example also evaluates and compares the efficacy of vincristine as a single agent to inhibit malignant B-cell viability in the same 10 diffuse large B-cell lymphoma (DLBCL), 4 multiple myeloma (MM), 2 follicular lymphoma (FL), and 1 mantle cell lymphoma (MCL) cell lines, to the

efficacy of the compound of formula I, or a pharmaceutically acceptable salt thereof, in combination with vincristine.

[0126] Cell Titer Glo Viability Assay: All procedures were performed at Gilead Sciences, Inc. in Branford, CT. Vincristine was tested alone as a single agent, or in combination with 300nM of the compound of Formula I, at 4 concentrations within the range of 0.03- 10 nM shown to be $< EC_{50}$ for cell viability as a single agent. The compound of Formula I and vincristine sulfate were dissolved in DMSO to prepare stock solutions in separate 96 well plates at 1000X for each compound so the final starting assay concentrations tested would be 300 nM for the compound of Formula I and 10 nM, 3 nM, 1 nM and 0.3 nM for vincristine. Compound plates were diluted and mixed together sequentially by 1:100 in RPMI without serum or additives creating a 10X stock in 2% DMSO of combinations or single agents. Cells were seeded into a 96-well plate at 10,000 – 20,000 cells per well in 100 μ l of growth media appropriate to the cell line supplemented with 100 U/L penicillin-streptomycin. OCI-Ly3, OCI-Ly4 and OCI-Ly7 were grown in Iscove's + 20% FBS, OCI-Ly19 was grown in alphaMEM + 20% FBS, and all other cell lines were grown in RPMI + 10% FBS. To each well containing cells, 11 μ l of 10X compound mixture in serum-free media was added in a final concentration of 0.2% DMSO and cells were incubated at 37°C in 5% CO₂ for 72 hours. Cell viability was assessed with Cell Titer Glo (CTG, Promega, Madison, WI) following the manufacturer's protocol on an Envision plate reader (Perkin Elmer, Waltham, MA). CTG signals were recorded for individual compound treatments and combinations. The results are shown in Figure 2 below.

[0127] FIG. 2 details the inhibition effects of the combination of the compound of formula I (FIG. 1) and vincristine as compared to vincristine alone in the 17 malignant B-cell lines, representing 4 hematological cancer types: DLBCL, MM, FL, and MCL (Figure 2).

Example 3

Comparison Effect of the Combination of the Compound of Formula I with Vincristine versus Effect of the Combination of the Compound of Formula I with A) obretastatin A4, B) colchicine, C) doxorubicin, and D) taxol in the malignant DLBCL B-cell line, DHL-10

[0128] This Example evaluates the efficacy of the compound of Formula I (Figure 1), or a pharmaceutically acceptable salt thereof, in combination with vincristine to inhibit malignant B-cell viability in the malignant diffuse large B-cell lymphoma (DLBCL) cell line DHL-10. This Example also evaluates and compares the efficacy of the compound of Formula I, or a pharmaceutically acceptable salt thereof, in combination with A) cobretastatin A4, B) colchicine, C) doxorubicin and D) taxol to inhibit malignant B-cell viability in the malignant diffuse large B-cell lymphoma (DLBCL) cell line DHL-10, to the efficacy of the combination of the compound of formula I, or a pharmaceutically acceptable salt thereof, with vincristine.

[0129] Cell Titer Glo Viability Assay: All procedures were performed at Gilead Sciences, Inc. in Branford, CT. Vincristine, combretastatin A4, colchicine, doxorubicin, and taxol were dissolved in DMSO to prepare stock solutions that were serially diluted three-fold in DMSO in a 96 well plate format at 1000X, so the final starting assay concentrations tested would be 1 μ M for all compounds except Doxorubicin which started at 10 μ M. Compound plates were diluted 1:100 in RPMI without serum or additives but containing either 1% DMSO or 3 μ M of the compound of Formula I creating a 10X stock. The DLBCL cell line, DHL-10 was plated at 10,000 cells per well in RPMI supplemented with 10 % FBS and 100 U/L penicillin-streptomycin. To each well containing cells, 11 μ l of 10X compound mixture in serum-free media was added in a final concentration of 0.2% DMSO and cells were incubated at 37°C in 5% CO₂ for 72 hours. Cell viability was assessed using Cell Titer Glo (CTG, Promega, Madison, WI) following the manufacturer's protocol on an Envision plate reader (Perkin Elmer, Waltham, MA). CTG signals were recorded for individual compound treatments and combinations. . EC₅₀ values were determined using a four parameter variable slope model with GraphPad Prism 6.0 software. Results are shown in Figure 2. The shifts in EC₅₀ values for inhibition of viable DHL-10 cells for the compounds alone or in combination with 300 nM of the compound of Formula I are shown in Table 2.

[0130] FIG. 3 summarizes the effect of the combination of the compound of Formula I and vincristine versus the combination of the compound of Formula I and combretastatin A4, the compound of Formula I and colchicine, the compound of Formula I and doxorubicin and the compound of formula I and taxol.

Table 2

Compound	Inhibition of cell viability EC ₅₀ (nM)		Fold shift
	No Compound of Formula I	+ 300 nM Compound of Formula I	
Vincristine	14.1	3.5	4.0
Colchicine	13.5	14.2	1.0
Combretastatin A4	5.9	4.8	1.2
Doxorubicin	152	161	0.9
Taxol	7.1	5.6	1.3

Example 4

Comparison Effect of the Combination of the Compound of Formula I with Vincristine versus Effect of the Combination of the Compound of Formula I with Vinblastine in the malignant DLBCL B-cell line, DHL-10

[0131] This Example evaluates the efficacy of the compound of Formula I, or a pharmaceutically acceptable salt thereof, in combination with vincristine to inhibit malignant B-cell viability in the malignant diffuse large B-cell lymphoma (DLBCL) cell line DHL-10. This Example also evaluates and compares the efficacy of the compound of Formula I, or a pharmaceutically acceptable salt thereof, in combination with vinblastine to inhibit malignant B-cell viability in the malignant diffuse large B-cell lymphoma (DLBCL) cell line DHL-10.

[0132] Cell Titer Glo Viability Assay: Vincristine and vinblastine were tested alone at 3 nM, or in combination with 100nM of the compound of Formula I. The compound of Formula I, vinblastine sulfate and vincristine sulfate were dissolved in DMSO to prepare stock solutions in separate 96 well plates at 1000X for each compound so the final starting assay concentrations tested would be 100 nM for the compound of Formula I and 3 nM for vinblastine and vincristine. Compound plates were diluted and mixed together sequentially by 1:100 in RPMI without serum or additives creating a 10X stock in 2% DMSO of combinations or single agents. DHL-10 cells were seeded into a 96-well plate at 10,000 cells per well in 100 μ l of in RPMI + 10% FBS supplemented with 100 U/L penicillin-streptomycin. To each well containing cells, 11 μ l of 10X compound mixture in serum-free

media was added in a final concentration of 0.2% DMSO and cells were incubated at 37°C in 5% CO₂ for 72 hours. Cell viability was assessed using Cell Titer Glo (CTG, Promega, Madison, WI) following the manufacturer's protocol and read on an Envision plate reader (Perkin Elmer, Waltham, MA). CTG signals were recorded for individual compound treatment and combinations. Results are shown in Figure 4.

[0133] FIG. 4 depicts and summarizes the inhibitory effects of the combination of the compound of Figure 1 (the compound of Formula I) and one of two of the vinca alkaloids, vincristine and vinblastine respectively, in the DLBCL cell line, DHL-10 when compounds were co-administered (Figure 4).

Example 5


Comparison Effect of the Compound of Formula I in Combination with Vincristine in a Syk-expressing Solid Tumor Cell Line versus Effect of the Compound of Formula I in Combination with Vincristine in a non Syk-expressing Solid Tumor Cell line

[0134] This Example evaluates the efficacy of the compound of Formula I, or a pharmaceutically acceptable salt thereof, in combination with vincristine to inhibit cell viability in the Syk-expressing pancreatic cell line, MIA PaCa-2. This Example also evaluates the efficacy of the combination of the compound of Formula I and vincristine to inhibit cell viability in the non-Syk expressing cell line, HepG2.

[0135] Cell Titer Glo Viability Assay: Vincristine was tested alone, or in combination with 4 concentrations of the compound of Formula I (300, 100, 33, and 11 nM), in the malignant pancreatic cell line, MIA PaCa-2 and in the hepatocellular carcinoma HepG2 cell line. The compound of Formula I and vincristine sulfate were dissolved in DMSO to prepare stock solutions in separate 96 well plates at 1000X for each compound so the final starting assay concentrations tested would be 300 - 11 nM for the compound of Formula I and 3 nM for vincristine. Compound plates were diluted and mixed together sequentially by 1:100 in RPMI without serum or additives creating a 10X stock in 2% DMSO of combinations or single agents. MIA PaCa-2 and HepG2 cells were seeded into 96-well plates at 5,000 cells per well in 100 µl of in RPMI + 10% FBS supplemented with 100 U/L penicillin-

streptomycin. To each well containing cells, 11 μ l of 10X compound mixture in serum-free media was added in a final concentration of 0.2% DMSO and cells were incubated at 37°C in 5% CO₂ for 72 hours. Cell viability was assessed using Cell Titer Glo (CTG, Promega, Madison, WI) following the manufacturer's protocol on an Envision plate reader (Perkin Elmer, Waltham, MA). CTG signals were recorded for individual compound treatment and combinations. Results are shown in Figure 5.

[0136] FIG. 5 summarizes the inhibition of cell viability by the combination of the compound of Formula I and vincristine in the Syk-expressing malignant pancreatic cell line, MiaPaca (Figure 5a) and in the non-Syk expressing malignant colon cell line, HepG2 (Figure 5b).



4.

[0137] **Syk Protein Assay:** Cell lines were grown logarithmically overnight in RPMI supplemented with 10% FBS and 100U/L penicillin-streptomycin. 1×10^7 were collected by centrifugation at 300 x g at room temperature for 8 minutes in 50 mL tubes. Cell pellets were lysed on ice for 15 minutes in 200 μ l of 1X RIPA buffer (Cell Signaling Technology, Danvers MA) containing protease (Roche, Palo Alto CA) and phosphatase inhibitors (Sigma, Saint Louis MO; Santa Cruz Technologies, Dallas TX). Cells lysates were transferred to 96-well V-bottom plates and used directly or frozen at -80°C for use the next day. Proteins were separated with 4-12% SDS-Bis/Tris gels and blotted onto nitrocellulose. Blots were blocked in Rockland Odyssey blocking buffer and incubated with a total Syk antibody, 4D10 (Santa Cruz) and pSyk-Y_{525/6} (Cell Signaling Technologies). The primary antibodies were diluted 1:1000 and incubated for 1 hour at room temperature. Blots were washed 3 times 5 minutes in Tris-buffered saline containing 1.0% Tween (TBS-T). Blots were then incubated

goat α mouse IgG (H + L), AlexaFluor 680 (Life Sciences, Inc) and Goat α Rabbit IgG (H + L), DyLight 800 (Thermo Scientific), each diluted 1:20,000 in blocking buffer, for 1 hour at room temperature. Blots were washed 3 times 5 minutes in TBS-T and analyzed on an Odyssey gel imager (LI-COR).

[0138] FIG 6 depicts the level of Syk expression in the MiaPaca and HepG2 malignant colon cell lines (Figure 6).

Example 6

Determination of Syk Expression in Certain Malignant Solid Tumor Cell Lines

(Figure 7)

[0139] Syk Protein Assay: Cell lines were grown logarithmically in overnight and 1×10^7 cells were collected by centrifugation at $300 \times g$ at room temperature for 8 minutes in 50 mL tubes. Cell pellets were lysed on ice for 15 minutes in 200 μ L of 1X RIPA buffer (Cell Signaling Technology, Danvers MA) containing protease (Roche, Palo Alto CA) and phosphatase inhibitors (Sigma, Saint Louis MO; Santa Cruz Technologies, Dallas TX). Cells lysates were transferred to 96-well V-bottom plates and used directly or frozen at -80°C for use the next day. Proteins were separated with 4-12% SDS-Bis/Tris gels and blotted onto nitrocellulose. Blots were blocked in Rockland Odyssey blocking buffer and incubated with a total Syk antibody, 4D10 (Santa Cruz). The primary antibody was diluted 1:1000 and incubated for 1 hour at room temperature. Blots were washed 3 times 5 minutes in Tris-buffered saline containing 1.0% Tween (TBS-T). Blots were then incubated goat α mouse IgG (H + L), AlexaFluor 680 (Life Sciences, Inc), diluted 1:20,000 in blocking buffer, for 1 hour at room temperature. Blots were washed 3 times 5 minutes in TBS-T and analyzed on an Odyssey gel imager (LI-COR). See Figure 7.

Example 7

Efficacy evaluation of entosplentinib and vincristine individually and in combination in the SU-DHL-10 Mouse Xenograft Model of Diffuse Large B Cell Lymphoma

Entosplentinib was evaluated for efficacy in vivo as a single agent and in combination with vincristine in a subcutaneous cell line tumor xenograft model in male SCID beige mice using the human diffuse large B-cell lymphoma cell line, SU-DHL-10 in a 2 x 2 dose level matrix of entosplentinib and vincristine.

FORMULATION - ENTOSPLENTINIB WAS FORMULATED IN A COMPLETE VEHICLE OF 0.5% HPMC-0.2% TWEEN 80 (PH 3.5 50 MM ACETATE BUFFER, AND THE LOWER DOSE WAS PREPARED BY DIRECT DILUTION). THE HIGH DOSE FORMULATION WAS MIXED USING A STIR BAR (AND POLYTRONED BRIEFLY AS NEEDED TO BREAK UP CLUMPS OF COMPOUND) TO FORM A YELLOW SOLUTION (PH=1.8), WHICH BECAME AN OPAQUE SUSPENSION WITHIN 30 MINUTES. THE DOSING SUSPENSION WAS PREPARED FRESH DAILY AND STORED AT 4°C PROTECTED FROM LIGHT BETWEEN TREATMENTS.

The formulation for the highest dose group of vincristine was prepared by diluting the stock solution with 0.9% sterile saline, and the lower dose was prepared by direct dilution of that stock with saline. The high dose formulation had a pH value of 7.08. The dosing formulations were prepared just prior to each treatment.

Animals - Male Harlan SCID beige mice (C.B-17/IcrHsd-*Prkdc*^{scid}*Lyst*^{bg-J}) were used in this study. They were 6-7 weeks old on Day 1 of the experiment and were fed irradiated Harlan 2918.15 Rodent Diet and water *ad libitum*. Animals were housed in static cages with Bed-O'Cobs™ bedding (The Andersons Lab Bedding Products) inside Biobubble® Clean Rooms that provide H.E.P.A filtered air into the bubble environment at 100 complete air changes per hour. All treatments, body weight determinations, and tumor measurements were carried out in the bubble environment. The environment was controlled to a temperature range of 70°±2°F and a humidity range of 30-70%.

Cells - SU-DHL-10 cells were obtained from Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures (DSMZ). They were grown in RPMI 1640 medium which was modified with 1% 100mM Na pyruvate, 1% 1M HEPES buffer, 1% of a 45% glucose solution and supplemented with 10% non-heat-inactivated Fetal Bovine Serum (FBS) and 1% 100X Penicillin/Streptomycin/L-Glutamine (PSG). The growth environment was maintained in an incubator with a 5% CO₂ atmosphere at 37°C. The cells were centrifuged at 200rcf for 8 minutes at 8°C, the supernatant aspirated, and the pellet re-suspended in cold Dulbecco's

Phosphate Buffered Saline (DPBS) by pipetting. An aliquot of the homogeneous cell suspension was diluted with a trypan blue solution and counted using a Luna automated cell counter. The pre-implantation cell viability was 91%. The cell suspension was centrifuged at 200g for 8 minutes at 8°C. The supernatant was aspirated and the cell pellet was re-suspended in cold 50% serum-free medium:50% Matrigel® Matrix (Corning) to generate a final concentration of 2.5×10^7 cells/ml. The cell suspension was maintained on wet ice during implantation. Following implantation, an aliquot of the remaining cells was diluted with a trypan blue solution and counted to determine the post-implantation cell viability (91%). Test animals were implanted subcutaneously, high in the axilla (just under the fore limb) on Day 0 with 5.0×10^6 cells in 0.2ml of 50% serum-free medium : 50% Matrigel® Matrix using a 27-gauge needle and syringe.

All mice were sorted into study groups based on tumor burden (estimated from caliper measurements). Efficacy was evaluated by tumor volume/weight monitored during and after treatment. Tumor burden (mg) was measured three times weekly by using caliper measurements and converting to tumor mass by the formula for the volume of a prolate ellipsoid assuming unit density as:

$$\text{Tumor burden (mg)} = (L \times W^2)/2$$

where L and W are the respective orthogonal tumor length and width measurements (mm).

The criteria used for a successful distribution in this study was a mean tumor burden for all groups within 10% of the overall mean tumor burden for the study population. Treatment began on Day 16 at an overall mean tumor burden of 197 mm^3 .

Group 1: Vehicle control (entospletinib) 0.5%HPMC in 0.2%Tween 80 in 50mM acetate buffer, pH 3.5 was dosed orally every twelve hours for 19 days (Days 16-32), and vehicle control (vincristine) 0.9% saline was dosed intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 2: Entospletinib was dosed orally at 75mg/kg every twelve hours for 6.5 days (Days 16-22); then lowered to 50mg/kg every twelve hours for 10.5 days (Days 22-32), in combination with saline intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 3: Entosplentinib was dosed orally at 25mg/kg every twelve hours for 19 days (Days 16-32), in combination with saline intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 4: Vehicle for entosplentinib was dosed orally every twelve hours for 19 days (Days 16-32), in combination with vincristine at 0.5mg/kg intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 5: Vehicle for entosplentinib was dosed orally every twelve hours for 19 days (Days 16-32), in combination with vincristine at 0.15mg/kg intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 6: Entosplentinib was dosed orally at 75mg/kg every twelve hours for 6.5 days (Days 16-22); then lowered due to body weight loss to 50mg/kg every twelve hours for 10.5 days (Days 22-32), in combination with vincristine at 0.5mg/kg intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 7: Entosplentinib was dosed orally at 25mg/kg every twelve hours for 19 days (Days 16-32), in combination with vincristine at 0.5mg/kg intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 8: Entosplentinib was dosed orally at 75mg/kg every twelve hours for 6.5 days (Days 16-22); then lowered to 50mg/kg every twelve hours for 10.5 days (Days 22-32), in combination with vincristine at 0.15mg/kg intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

Group 9: Entosplentinib was dosed orally at 25mg/kg every twelve hours for 19 days (Days 16-32), in combination with vincristine at 0.15mg/kg intravenously every 7 days, beginning on the third day of oral dosing (Days 18, 25, and 32).

All mice were dosed according to individual body weight on the day of treatment (0.1ml/20g).

On the 16th day post-tumor implantation, animals with established tumors (n=9 animals per group) were treated for 3 days by oral gavage with either vehicle or entosplentinib at 75mg/kg or 25 mg/kg twice daily (BID). On the third day, the animals received an intravenous dose of saline or vincristine and plasma and tumor samples were

collected 0.5, 2 or 12 hours post dose. An additional 12 animals per group were treated for 19 days (starting on Day 16 of study) by oral gavage with either vehicle or entosplentinib at the planned dose levels of 25 mg/kg and 75 mg/kg BID and intravenous administration of either saline or vincristine (0.5 mg/kg and 0.15 mg/kg every 7 days (Q7D) for 3 doses starting on the third day after the start of entosplentinib (Day 19 of study). The 75 mg/kg BID dose level of entosplentinib was administered for 6.5 days and then reduced to 50 mg/kg BID due to the higher than planned plasma concentrations measured in plasma samples collected on the third day of dosing, and weight loss in group being co-treated with 0.5 mg/kg vincristine. This dose level is designated 75/50 mg/kg.

Efficacy was evaluated by tumor volume measurements. Plasma levels of entosplentinib and vincristine were evaluated on the third day of dosing entosplentinib (Day 19) and the last day of dosing (Day 32). Tolerability was evaluated by daily observations, body weight (3 times per week) and circulating blood cell counts at the end of the study.

All dose levels and combinations evaluated were tolerated, following the 75 mg/kg entosplentinib dose reduction to 50 mg/kg. The doses of 25 and 50 mg/kg BID entosplentinib resulted in plasma concentrations at the end of study that exceed the in vitro pervanadate-stimulated mouse whole blood EC_{50} and EC_{80} values for inhibition of SYK, respectively, at C_{min} (trough). Vincristine administered at both 0.15 and 0.5 mg/kg resulted in significant inhibition of tumor growth. Entosplentinib dosed alone at 25 or 75/50 mg/kg inhibited tumor growth but to a lesser degree than that seen with vincristine.

The addition of entosplentinib at the 75/50 mg/kg level resulted in significant improvement in tumor growth inhibition over vincristine alone with an increase in % tumor growth inhibition (TGI) from 85% to 96% for 0.5 mg/kg vincristine and an increase in %TGI from 42% to 71% for 0.15 mg/kg vincristine. The addition of 25 mg/kg entosplentinib to either dose of vincristine did not significantly increase the tumor growth inhibition. While the groups receiving either entosplentinib or vincristine as single agents had no complete or partial regressions, 50% of the mice receiving the combination of 75/50 mg/kg entosplentinib with 0.5 mg/kg vincristine had partial responses and 8% had complete regression and 8% were tumor free at the end of study. The tumor growth inhibition and tolerability of entosplentinib and vincristine alone and in combination are summarized in the table below.

Summary of Efficacy with Entosplentinib and Vincristine at Tolerated Doses

Treatment (Group number)	Day 31 % Tumor Growth Inhibition^a	Significance (P value)
Vehicle for ENTO BID + Vehicle for VCR Q7Dx3 (1)	NA ^b	
ENTO 75/50 mg/kg BID + Vehicle for VCR Q7Dx3 (2)	39	<0.00001 vs. vehicle
ENTO 25 mg/kg BID + Vehicle for VCR Q7Dx3 (3)	20	0.0455 vs. vehicle
Vehicle for ENTO + VCR 0.5 mg/kg Q7Dx3 (4)	85	<0.00001 vs. vehicle
Vehicle for ENTO + VCR 0.15 mg/kg Q7Dx3 (5)	42	<0.00001 vs. vehicle
ENTO 75/50 mg/kg BID + VCR 0.5 mg/kg Q7Dx3 (6)	96	0.001 vs. VCR 0.5 mg/kg
ENTO 25 mg/kg BID + VCR 0.5 mg/kg Q7Dx3 (7)	90	0.3866 vs. VCR 0.5 mg/kg
ENTO 75/50 mg/kg BID + VCR 0.15 mg/kg Q7Dx3 (8)	71	<0.00001 vs. VCR 0.15 mg/kg
ENTO 25 mg/kg BID + VCR 0.15 mg/kg Q7Dx3 (9)	58	0.0754 vs. VCR 0.15 mg/kg

a The percentage of tumor growth inhibition (%TGI) was calculated using the delta of the mean treated tumor values divided by the delta of the mean control tumor values expressed as a percentage and then subtracted from 100%.

b NA does not apply since this is the control


The administration of combinations of entosplentinib with vincristine to animals bearing the SU-DHL-10 subcutaneous xenograft demonstrated that a dose of entosplentinib which results in a calculated EC₈₀ trough coverage of SYK resulted in significant increase in tumor growth

inhibition as well as an increase to 50% partial tumor regressions compared to 0% with entospletinib or vincristine treatment alone, with 8% complete regression.

CLAIMS

What is claimed is:

1. A method for treating cancer in a human in need thereof, comprising administering to the human a therapeutically effective amount of a compound of formula I:



or a pharmaceutically acceptable salt thereof, and a therapeutically effective amount of a vinca-alkaloid, or a pharmaceutically acceptable salt.

2. The method of claim 1, wherein the vinca-alkaloid is selected from the group consisting of vincristine, vinblastine, vindesine, vinorelbine, desoxyvincaminol, vincaminol, vinburnine, vincamajine, and vineridine.

3. The method of claim 2, wherein the vinca-alkaloid is selected from the group consisting of vincristine, vinblastine, vindesine, and vinorelbine.

4. The method of claim 1 wherein the vinca-alkalod is vincristine.

5. The method of claim 1 wherein the vinca-alkalod is vinblastine.

6. The method of claim 1, wherein the compound of formula (I) is present in a pharmaceutical composition comprising the compound of formula (I), and at least one pharmaceutically acceptable excipient

7. The method of claim 1, wherein the compound of formula (I) or a pharmaceutically acceptable salt thereof is administered before the vinca-alkaloid, or a pharmaceutically acceptable salt thereof.

8. The method of claim 1, wherein the vinca alkaloid, or a pharmaceutically acceptable salt thereof, is administered before the compound of formula (I) or a pharmaceutically acceptable salt thereof.

9. The method of claim 1, wherein the compound of formula (I) or a pharmaceutically acceptable salt thereof, and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, are administered simultaneously.
10. The method of claim 1, wherein each of the compound of formula I and the vinca-alkaloid, or pharmaceutically acceptable salts thereof, is independently administered, wherein the compound of formula I is administered twice a day and wherein further the vinca-alkaloid is administered once a week.
11. The method of claim 1, wherein the compound of formula (I) or a pharmaceutically acceptable salt thereof is administered at a dose between 100 mg and 800 mg and the vinca-alkaloid, or a pharmaceutically acceptable salt thereof, is administered at a dose between about 0.1 mg-M^2 and 1.5 mg-M^2 .
12. The method of claim 1, wherein the compound of formula (I) or a pharmaceutically acceptable salt thereof is administered at a dose between 200 mg and 400 mg and the vinca-alkaloid or a pharmaceutically acceptable salt thereof is administered at a dose between 0.25 mg-M^2 and 1.0 mg-M^2 , wherein the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine.
13. The method of claim 1, wherein the human who has cancer is (i) refractory to at least one chemotherapy treatment, or (ii) is in relapse after treatment with chemotherapy, or a combination thereof.
14. The method of claim 1, wherein the human has not previously been treated for the cancer.
15. The method of claim 1, wherein the human has a 17p deletion, a TP53 mutation, or a combination thereof.
16. The method of claim 15, wherein the human further has NOTCH1, a SF3B1 mutation, a 11q deletion, or any combination thereof.
17. The method of claim 1 wherein:

the vinca-alkaloid is selected from the group consisting of vincristine and vinblastine,

and

the human is (i) refractory to at least one anti-cancer treatment, or (ii) in relapse after treatment with at least one anti-cancer therapy, or a combination thereof.

18. The method of Claim 17 wherein the human is not undergoing any other anti-cancer treatments.

19. The method of claim 17, wherein the human is not undergoing any other anti-cancer treatments using one or more PI3K inhibitors.

20. The method of claim 18, wherein the human is refractory to at least one anti-cancer treatment.

21. The method of claim 18, wherein the human is in relapse after treatment with at least one anti-cancer treatment.

22. The method of claim 18, wherein about 200 mg to about 800 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the human twice daily.

23. The method of claim 22, wherein 100-400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the human twice daily.

24. The method of claim 18, wherein the human has a 17p deletion, a TP53 mutation, or a combination thereof and about 400 mg of the compound of formula I, or a pharmaceutically acceptable salt thereof, is administered to the human twice daily.

25. The method of claim 1, wherein the cancer is a hematologic malignancy.

26. The method of claim 1, wherein the cancer is a leukemia.

27. The method of claim 1, wherein the leukemia is chronic lymphocytic leukemia (CLL).

28. The method of claim 1, wherein the cancer is a lymphoma.

29. The method of claim 1, wherein the lymphoma is non-Hodgkin's lymphoma (NHL).

30. The method of claim 29, wherein the NHL is diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), follicular lymphoma (FL), small lymphocytic

lymphoma (SLL), lymphoplasmacytic lymphoma (LPL), and marginal zone lymphoma (MZL).

31. The method of claim 1, wherein the cancer is selected from the group consisting of acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), Burkitt's lymphoma (BL), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), myelodysplastic syndrome (MDS), myeloproliferative disease (MPD), chronic myeloid leukemia (CML), multiple myeloma (MM), non-Hodgkin's lymphoma (NHL), indolent non-Hodgkin's lymphoma (iNHL), refractory iNHL, mantle cell lymphoma (MCL), follicular lymphoma (FL), Waldstrom's macroglobulinemia (WM), T-cell lymphoma, B-cell lymphoma, Hodgkin's lymphoma, diffuse large B-cell lymphoma (DLBCL), lymphoplasmacytic lymphoma (LPL), and marginal zone lymphoma (MZL).

32. The method of claim 1, wherein the cancer is a solid tumor.

33. The method of claim 1, wherein the cancer is a solid tumor and expresses spleen tyrosine kinase (Syk) activity.

34. The method of claim 32, wherein the solid tumor cancer is selected from the group consisting of pancreatic, lung, colorectal cancer, ovarian, breast, esophageal, adenocarcinoma and hepatocellular.

35. The method of claim 1 wherein each of the compound of formula I and the vinca-alkaloid, or pharmaceutically acceptable salts thereof, is independently administered, wherein the compound of formula I is administered once.

36. The method of claim 1 wherein each of the compound of formula I and the vinca-alkaloid, or pharmaceutically acceptable salts thereof, is independently administered, wherein the compound of formula I is administered once a day and wherein further the vinca-alkaloid is administered once a week.




Figure 2 (a) – 2(f)




Figure 2(g)- 2(n)




FIGURE 2(o) – 2(q)




FIGURE 3




FIGURE 4




FIGURE 5(a) – 5(b)




FIGURE 6





FIGURE 7



INTERNATIONAL SEARCH REPORT

International application No

PCT/US2015/040029

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K45/06 A61K31/475 A61K31/4985 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, CHEM ABS Data, EMBASE, SCISEARCH, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 8 450 321 B2 (MITCHELL SCOTT A [US] ET AL) 28 May 2013 (2013-05-28) cited in the application column 1, lines 20-24 column 51, line 34 - column 53, line 15 column 53, line 60 - column 54, line 14 claims 1, 2	1-3, 5-36
Y	----- KEVIN S. CURRIE ET AL: "Discovery of GS-9973, a Selective and Orally Efficacious Inhibitor of Spleen Tyrosine Kinase", JOURNAL OF MEDICINAL CHEMISTRY, vol. 57, no. 9, 8 May 2014 (2014-05-08), pages 3856-3873, XP055143746, ISSN: 0022-2623, DOI: 10.1021/jm500228a abstract page 3865, left-hand column, paragraph 2 ----- -/-	1-3, 6-16, 25-36



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

1 October 2015

Date of mailing of the international search report

09/10/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Herdemann, Matthias

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2015/040029

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Anonymous: "Vinca Alkaloids for Cancer Treatment", 24 May 2014 (2014-05-24), XP055213453, Mountain Lakes, New Jersey 07046, US Retrieved from the Internet: URL:https://web.archive.org/web/20140524022911/http://chemoth.com/types/vinca-alkaloids [retrieved on 2015-09-15] the whole document	1-3, 6-16, 25-36
Y	SORENSEN J B ET AL: "Vinca alkaloids in the treatment of non-small cell lung cancer", CANCER TREATMENT REVIEWS, SAUNDERS, US, vol. 14, no. 1, 1 March 1987 (1987-03-01), pages 29-51, XP023271021, ISSN: 0305-7372, DOI: 10.1016/0305-7372(87)90049-1 [retrieved on 1987-03-01] the whole document	1-3, 6-16, 25-36
X,P	Anonymous: "NCT02404220 on 2015_06_19: ClinicalTrials.gov Archive", ClinicalTrials.gov, 19 June 2015 (2015-06-19), pages 1-4, XP055213378, Retrieved from the Internet: URL:https://clinicaltrials.gov/archive/NCT02404220/2015_06_19 [retrieved on 2015-09-15] the whole document	1-4, 6-17, 19-36
X,P	WO 2015/084992 A1 (GILEAD SCIENCES INC [US]) 11 June 2015 (2015-06-11) claims 1-18 page 23, lines 14-30	1-3,5-36
X,P	WO 2015/017460 A1 (GILEAD CONNECTICUT INC [US]) 5 February 2015 (2015-02-05) paragraphs [0187], [0191], [0192] claims 1, 28-33	1-36
X,P	WO 2015/017466 A1 (GILEAD CONNECTICUT INC [US]) 5 February 2015 (2015-02-05) paragraphs [0335], [0339], [0340] claims 1, 32-37	1-36

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2015/040029

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
US 8450321	B2	28-05-2013	US 2012220582 A1	30-08-2012
			US 2013267496 A1	10-10-2013

WO 2015084992	A1	11-06-2015	US 2015150881 A1	04-06-2015
			WO 2015084992 A1	11-06-2015

WO 2015017460	A1	05-02-2015	US 2015038505 A1	05-02-2015
			UY 35683 A	27-02-2015
			WO 2015017460 A1	05-02-2015

WO 2015017466	A1	05-02-2015	US 2015038504 A1	05-02-2015
			UY 35684 A	27-02-2015
			WO 2015017466 A1	05-02-2015
