



(86) Date de dépôt PCT/PCT Filing Date: 2001/11/26
(87) Date publication PCT/PCT Publication Date: 2003/01/16
(85) Entrée phase nationale/National Entry: 2003/05/21
(86) N° demande PCT/PCT Application No.: US 2001/043151
(87) N° publication PCT/PCT Publication No.: 2003/004055
(30) Priorité/Priority: 2000/11/27 (09/724,315) US

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 39/39, A61K 48/00
(71) Demandeur/Applicant:
POWERJECT VACCINES, INC., US
(72) Inventeurs/Inventors:
HAYNES, JOEL R., US;
ARRINGTON, JOSHUA E., US
(74) Agent: OGILVY RENAULT

(54) Titre : ADJUVANTS COMPORTANT DES ACIDES NUCLEIQUES
(54) Title: NUCLEIC ACID ADJUVANTS

(57) **Abrégé/Abstract:**

Recombinant nucleic acid molecules are described. The molecules have two nucleic acid sequences, wherein the first nucleic acid sequence is a truncated A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence is a truncated B subunit coding region. Vectors and compositions containing these molecules are also described. Methods for enhancing an immune response against an antigen of interest using these recombinant nucleic acid molecules and compositions are also described.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004055 A2

(51) International Patent Classification⁷: **A61K 39/39**,
48/00

(21) International Application Number: PCT/US01/43151

(22) International Filing Date:
26 November 2001 (26.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/724,315 27 November 2000 (27.11.2000) US

(71) Applicant: **POWDERJECT VACCINES, INC.**
[US/US]; 585 Science Drive, Madison, WI 53711 (US).

(72) Inventors: **HAYNES, Joel, R.**; 585 Science Drive, Madison, WI 53711 (US). **ARRINGTON, Joshua, E.**; 585 Science Drive, Madison, WI 53711 (US).

(74) Agent: **MCCRACKEN, Thomas, P.**; PowderJect Technologies Inc., 6511 Dumbarton Circle, Fremont, CA 94555 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID ADJUVANTS

(57) Abstract: Recombinant nucleic acid molecules are described. The molecules have two nucleic acid sequences, wherein the first nucleic acid sequence is a truncated A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence is a truncated B subunit coding region. Vectors and compositions containing these molecules are also described. Methods for enhancing an immune response against an antigen of interest using these recombinant nucleic acid molecules and compositions are also described.



WO 03/004055 A2

NUCLEIC ACID ADJUVANTS

Technical Field

5 The invention relates to the fields of molecular biology and immunology, and generally relates to nucleic acid immunization techniques. More specifically, the invention relates to polynucleotides encoding an adjuvant, and to immunization strategies employing such polynucleotides.

Background

10 Techniques for the injection of DNA and mRNA into mammalian tissue for the purposes of immunization against an expression product have been described in the art. The techniques, termed “nucleic acid immunization” herein, have been shown to elicit both humoral and cell-mediated immune responses. For example, sera from mice immunized with a DNA construct encoding the
15 envelope glycoprotein, gp160, were shown to react with recombinant gp160 in immunoassays, and lymphocytes from the injected mice were shown to proliferate in response to recombinant gp120. Wang et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:4156-4160. Similarly, mice immunized with a human growth hormone (hGH) gene demonstrated an antibody-based immune response. Tang et
20 al. (1992) *Nature* 356:152-154. Intramuscular injection of DNA encoding influenza nucleoprotein driven by a mammalian promoter has been shown to elicit a CD8+ CTL response that can protect mice against subsequent lethal challenge with virus. Ulmer et al. (1993) *Science* 259:1745-1749. Immunohistochemical studies of the injection site revealed that the DNA was
25 taken up by myeloblasts, and cytoplasmic production of viral protein could be demonstrated for at least 6 months.

Summary of the Invention

30 It is a primary object of the invention to provide a polynucleotide adjuvant composition containing first and second nucleic acid sequences, wherein the first nucleic acid sequence is a truncated A subunit coding region obtained or derived

from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence is a truncated B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin. Each of the truncated subunit coding regions has a 5' deletion and encodes a subunit peptide not having an amino terminal bacterial signal peptide.

The first and second nucleic acid sequences may be present in the same or in different nucleic acid constructs. The truncated subunit coding regions may be obtained or derived from the same bacterial ADP-ribosylating exotoxin and, in certain preferred embodiments, the bacterial ADP-ribosylating exotoxin is a cholera toxin (CT) or an *E. coli* heat labile enterotoxin (LT). In addition, at least one of the truncated subunit coding regions may be genetically modified to detoxify the subunit peptide encoded thereby, for example wherein the truncated A subunit coding region has been genetically modified to disrupt or inactivate ADP-ribosyl transferase activity in the subunit peptide expression product.

It is also a primary object of the invention to provide a polynucleotide adjuvant composition containing first and second nucleic acid sequences, wherein the first nucleic acid sequence is a modified A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence is a B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin. The modified A subunit coding region and said B subunit coding region each encode a mature subunit peptide, and the modified A subunit coding region has been genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby.

As above, the first and second nucleic acid sequences may be present in the same or in different nucleic acid constructs. The truncated subunit coding regions may be obtained or derived from the same bacterial ADP-ribosylating exotoxin and, in certain preferred embodiments, the bacterial ADP-ribosylating exotoxin is a cholera toxin (CT) or an *E. coli* heat labile enterotoxin (LT). In addition, at least one of the truncated subunit coding regions may be genetically modified to detoxify the subunit peptide encoded thereby, for example wherein the truncated A subunit coding region has been genetically modified to disrupt or

inactivate ADP-ribosyl transferase activity in the subunit peptide expression product.

In certain aspects of the invention, the above compositions can be provided in particulate form, for example wherein the compositions are particulates suitable for delivery from a particle delivery device. In this regard, the present compositions may be coated onto the same or a different core carrier particle and thus suitable for delivery using a particle-mediated transfection technique. Preferred core carrier particles will have an average diameter of about 0.1 to 10 μm , and may comprise a metal such as gold. Accordingly, it is a still further object of the invention to provide a particle delivery device loaded with (e.g., containing) a particulate composition as defined herein.

It is also a primary object of the invention to provide for the use of a composition containing a first and second nucleic acid sequence, where each sequence includes a coding region for a subunit from a bacterial ADP-ribosylating exotoxin in the manufacture of a medicament for enhancing an immune response in a vertebrate subject against an antigen of interest in the said subject. The antigen of interest and the composition are administered to the subject such that the toxin subunits encoded by the first and second nucleic acid sequences are expressed in an amount sufficient to elicit an enhanced immune response against the antigen. The first nucleic acid sequence contains a truncated A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence contains a truncated B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, however with the proviso that each of the truncated subunit coding regions has a 5' deletion and encodes a subunit peptide not having an amino terminal bacterial signal peptide.

It is a related primary object of the invention to provide a method for enhancing an immune response against an antigen of interest in a subject. The method generally entails: (a) administering the antigen of interest to the subject; (b) providing an adjuvant composition comprising first and second nucleic acid sequences, wherein the first nucleic acid sequence is a truncated A subunit coding

region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence is a truncated B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin; and (c) administering the adjuvant composition to the subject, whereby upon introduction to the subject, the first and second nucleic acid sequences are expressed to provide subunit peptides in an amount sufficient to elicit an enhanced immune response against the antigen of interest. The subunit coding regions are truncated in that each coding region has a 5' deletion and encodes a subunit peptide not having an amino terminal bacterial signal peptide.

It is yet a further primary object of the invention to provide for the use of a composition comprising a first and second nucleic acid sequence, where each sequence includes a coding region for a subunit from a bacterial ADP-ribosylating exotoxin in the manufacture of a medicament for enhancing an immune response in a vertebrate subject against an antigen of interest in the said subject. The antigen of interest and the composition are administered to the subject such that the toxin subunits encoded by the first and second nucleic acid sequences are expressed in an amount sufficient to elicit an enhanced immune response against the antigen. The first nucleic acid sequence contains a modified A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the second nucleic acid sequence contains a B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, however with the proviso that the modified A subunit coding region and B subunit coding region each encode a mature subunit peptide, and with the further proviso that the modified A subunit coding region has been genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby.

It is a related primary object of the invention to provide a method for enhancing an immune response against an antigen of interest in a subject, wherein the method entails: (a) administering the antigen of interest to the subject; (b) providing an adjuvant composition comprising first and second nucleic acid sequences, wherein the first nucleic acid sequence is a modified A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and the

second nucleic acid sequence is a B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin; and (c) administering the adjuvant composition to the subject, whereby upon introduction to the subject, the first and second nucleic acid sequences are expressed to provide subunit peptides in an amount sufficient to elicit an enhanced immune response against the antigen of interest. The subunit coding regions are modified in that each encodes a mature subunit peptide, but the A subunit coding region has been genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby.

In the uses and methods of the invention, administering the adjuvant compositions entails transfecting cells of the subject with a polynucleotide adjuvant composition according to the present invention. Expression cassettes and/or vectors containing the nucleic acid molecules of the present invention can be used to transfect the cells, and transfection is carried out under conditions that permit expression of the subunit peptides within the subject. The method may further entail one or more steps of administering at least one secondary composition to the subject.

The transfection procedure carried out during the immunization can be conducted either *in vivo*, or *ex vivo* (e.g., to obtain transfected cells which are subsequently introduced into the subject prior to carrying out the secondary immunization step). When *in vivo* transfection is used, the nucleic acid molecules can be administered to the subject by way of intramuscular or intradermal injection of plasmid DNA or, preferably, administered to the subject using a particle-mediated delivery technique. Vaccine compositions (containing the antigen of interest) can be provided in the form of any suitable vaccine composition, for example, in the form of a peptide subunit composition, in the form of a nucleic acid vaccine composition, or in the form of a whole or split virus influenza vaccine composition.

In certain methods, the antigen of interest and the adjuvant composition are administered to the same site in the subject. For example, the adjuvant composition and the antigen of interest can be administered concurrently (e.g.,

provided in a single vaccine composition). In certain preferred embodiments, the adjuvant and, optionally the antigen of interest, is administered in particulate form, for example wherein the adjuvant composition has been coated onto a core carrier particle and delivered to the subject using a particle-mediated delivery technique.

In these methods, administration of the polynucleotide adjuvant compositions of the present invention preferably results in an augmented cellular immune response against the co-administered antigen of interest. Such an enhanced immune response may be generally characterized by increased titers of interferon-producing CD4⁺ and/or CD8⁺ T lymphocytes, increased antigen-specific cytotoxic T lymphocyte (CTL) activity, and a T helper 1-like immune response (Th1) against the antigen of interest (characterized by increased antigen-specific antibody titers of the subclasses typically associated with cellular immunity (e.g., IgG2a), usually with a concomitant reduction of antibody titers of the subclasses typically associated with humoral immunity (e.g., IgG1)) instead of a T helper 2-like immune response (Th2) such as that normally produced when immunizing a subject using a bacterial ADP-ribosylating exotoxin adjuvant such as CT or LT.

Advantages of the present invention include, but are not limited to the ability of the present adjuvant compositions to provide significant adjuvant effect and thereby enhance the immunogenicity of a co-administered antigen in an immunized subject, as well as the ability to favor a Th1-like immune response against the co-administered antigen which is beneficial in a vaccine product.

These and other objects, aspects, embodiments and advantages of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

Brief Description of the Drawings

Figure 1 is a restriction map and functional map of plasmid pPJV2002 that contains a truncated coding sequence for a Cholera Toxin (CT) subunit A (CTA) peptide, wherein the plasmid further contains the human cytomegalovirus

(hCMV) immediate early promoter and associated intron A sequence, and the coding sequence for the signal peptide of human tissue plasminogen activator to allow for secretion from mammalian cells of the truncated CTA expression product. The figure further contains the complete nucleic acid sequence (SEQ ID NO: 1) for the pPJV2002 plasmid.

Figure 2 is a restriction map and functional map of plasmid pPJV2003 that contains a truncated coding sequence for a Cholera Toxin (CT) subunit B (CTB) peptide, wherein the plasmid further contains the hCMV immediate early promoter and associated intron A sequence, and the coding sequence for the signal peptide of human tissue plasminogen activator to allow for secretion from mammalian cells of the truncated CTB expression product. The figure further contains the complete nucleic acid sequence (SEQ ID NO: 2) for the pPJV2003 plasmid.

Figure 3 is a restriction map and functional map of plasmid pPJV2006 that contains a truncated coding sequence for a CTA peptide, wherein the truncated CTA coding sequence has been further modified to delete a C-terminal KDEL motif in the subunit peptide encoded thereby. The plasmid further contains the hCMV immediate early promoter and associated intron A sequence, and the coding sequence for the signal peptide of human tissue plasminogen activator to allow for secretion from mammalian cells of the truncated CTA expression product. The figure further contains the complete nucleic acid sequence (SEQ ID NO: 3) for the pPJV2006 plasmid.

Figure 4 is a restriction map and functional map of plasmid pPJV2004 that contains a truncated coding sequence for an *E. coli* heat labile enterotoxin (LT) subunit A (LTA) peptide, wherein the plasmid further contains the hCMV immediate early promoter and associated intron A sequence, and the coding sequence for the signal peptide of human tissue plasminogen activator to allow for secretion from mammalian cells of the truncated LTA expression product. The figure further contains the complete nucleic acid sequence (SEQ ID NO: 4) for the pPJV2004 plasmid.

Figure 5 is a restriction map and functional map of plasmid pPJV2005 that contains a truncated coding sequence for an LT subunit B (LTB) peptide, wherein the plasmid further contains the hCMV immediate early promoter and associated intron A sequence, and the coding sequence for the signal peptide of human tissue plasminogen activator to allow for secretion from mammalian cells of the truncated LTB expression product. The figure further contains the complete nucleic acid sequence (SEQ ID NO: 5) for the pPJV2005 plasmid.

Figure 6 is a restriction map and functional map of plasmid pPJV2007 that contains a truncated coding sequence for an LTA peptide, wherein the truncated LTA coding sequence has been further modified to delete a C-terminal RDEL motif in the subunit peptide encoded thereby. The plasmid further contains the hCMV immediate early promoter and associated intron A sequence, and the coding sequence for the signal peptide of human tissue plasminogen activator to allow for secretion from mammalian cells of the truncated LTA expression product. The figure further contains the complete nucleic acid sequence (SEQ ID NO: 6) for the pPJV2007 plasmid.

Figure 7 depicts the results from the ELISA carried out in Example 5. The histogram represents the log reciprocal titer of anti-gp120 antibody present in the animals receiving, from left to right in the figure, Formulation #1 containing the empty pWRG7054 vector ("EmpVec"); Formulation #2 containing the EmpVec (pWRG7054) and the pCIA-EnvT plasmid ("gp120"); Formulation #3 containing the EmpVec (pWRG7054) combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B"); Formulation #4 containing the pCIA-EnvT plasmid ("gp120") combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B"); Formulation #5 containing the pCIA-EnvT plasmid ("gp120") combined with the pPJV2006 and pPJV2003 adjuvant vectors ("CTA-KDEL/B"); or no vaccine and/or adjuvant composition ("naive").

Figure 8 depicts the results from the *in situ* ELISA carried out in Example 5. The histogram represents the relative level of gp120-specific IFN- γ production in splenocytes obtained from animals receiving, from left to right in the figure, Formulation #1 containing the empty pWRG7054 vector ("EmpVec");

Formulation #2 containing the EmpVec (pWRG7054) and the pCIA-EnvT plasmid ("gp120"); Formulation #3 containing the EmpVec (pWRG7054) combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B"); Formulation #4 containing the pCIA-EnvT plasmid ("gp120") combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B"); or Formulation #5 containing the pCIA-EnvT plasmid ("gp120") combined with the pPJV2006 and pPJV2003 adjuvant vectors ("CTA-KDEL/B").

Figure 9 depicts the results from the ELISPOT assay carried out in Example 5. The histogram represents the relative levels of IFN- γ -producing splenocytes obtained from animals receiving, from left to right in the figure, Formulation #1 containing the empty pWRG7054 vector ("EmpVec"); Formulation #2 containing the EmpVec (pWRG7054) and the pCIA-EnvT plasmid ("gp120"); Formulation #3 containing the EmpVec (pWRG7054) combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B"); Formulation #4 containing the pCIA-EnvT plasmid ("gp120") combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B"); or Formulation #5 containing the pCIA-EnvT plasmid ("gp120") combined with the pPJV2006 and pPJV2003 adjuvant vectors ("CTA-KDEL/B").

Figure 10 depicts the results from the ELISA carried out in Example 6. In this figure, the geometric mean absorbance values represent the titer of anti-HBcAg antibody present in serum samples (at four different dilutions) taken at the booster immunization (week 6 of the study) from animals receiving either Formulation #1 containing the HBcAg/HBsAg vector plasmid (pWRG7193); or Formulation #2 containing the HBcAg/HBsAg vector plasmid (pWRG7193) combined with the pPJV2002 and pPJV2003 adjuvant vectors ("CTA/B").

Figure 11 depicts the results from the ELISA carried out in Example 6. In this figure, the geometric mean absorbance values represent the titer of anti-HBcAg antibody present in serum samples (at four different dilutions) taken 2 weeks following the booster immunization (week 8 of the study) from animals receiving either Formulation #1 containing the HBcAg/HBsAg vector plasmid (pWRG7193); or Formulation #2 containing the HBcAg/HBsAg vector plasmid

(pWRG7193) combined with the pPJV2002 and pPJV2003 adjuvant vectors (“CTA/B”).

Figure 12 depicts the results from the ELISA carried out in Example 7. The histogram represents the log IgG1::IgG2a ratios from each immunization group receiving, from left to right in the figure, Formulation #1 containing the pM2-FL plasmid (“M2”) combined with the empty vector plasmid control (pWRG7054); Formulation #2 containing the pM2-FL plasmid combined with the pPJV2002 and pPJV2003 CTA/B adjuvant vectors (“M2 + CT”); or Formulation #7 containing the pM2-FL plasmid combined with the pPJV2004 and pPJV2005 LTA/B adjuvant vectors (“M2 + LT”).

Figures 13A-13D depict the results from the IFN- γ and the IL4 ELISPOT assays used to assess the immune response to the Hepatitis B virus surface and core antigens in the first study of Example 8. The histograms represent the number of spots per 1×10^5 spleen cells from the various experimental groups.

Figure 14 depicts the survival results from the HSV-2 virus challenge study carried out in Example 9.

Detailed Description of the Invention

Before describing the present invention in detail, it is to be understood that this invention is not limited to particularly exemplified molecules or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting. In addition, the practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology, recombinant DNA techniques and immunology all of which are within the ordinary skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, et al., *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *A Practical Guide to Molecular Cloning* (1984); and

Fundamental Virology, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.).

All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

5 It must be noted that, as used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the content clearly dictates otherwise.

Definitions

10 Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although a number of methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred materials and methods are described herein.

15 In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

The term “adjuvant” intends any material or composition capable of specifically or non-specifically altering, enhancing, directing, redirecting, potentiating or initiating an antigen-specific immune response. Thus,
20 coadministration of an adjuvant with an antigen may result in a lower dose or fewer doses of antigen being necessary to achieve a desired immune response in the subject to which the antigen is administered, or coadministration may result in a qualitatively and/or quantitatively different immune response in the subject. The effectiveness of an adjuvant can be determined by administering the adjuvant
25 with a vaccine composition in parallel with a vaccine composition alone to animals and comparing antibody and/or cellular-mediated immunity in the two groups using standard assays such as radioimmunoassay, ELISAs, CTL assays, and the like, all well known in the art. Typically, in a vaccine composition, the adjuvant is a separate moiety from the antigen, although a single molecule can
30 have both adjuvant and antigen properties.

An “adjuvant composition” intends any pharmaceutical composition containing an adjuvant. Adjuvant compositions can be delivered in the methods of the invention while in any suitable pharmaceutical form, for example, as a liquid, powder, cream, lotion, emulsion, gel or the like. However, preferred
5 adjuvant compositions will be in particulate form. It is intended, although not always explicitly stated, that molecules having similar biological activity as wild-type or purified peptide or chemical adjuvants.

The term “peptide” is used in its broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics.
10 The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used herein, the term “amino acid” refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an “oligopeptide” if the peptide chain is short. If the peptide chain is long, the peptide is typically referred to as a
15 “polypeptide” or a “protein”.

An “antigen” refers to any agent, generally a macromolecule, which can elicit an immunological response in an individual. The term may be used to refer to an individual macromolecule or to a homogeneous or heterogeneous
20 population of antigenic macromolecules. As used herein, “antigen” is generally used to refer to a peptide or carbohydrate molecule that contains one or more epitopes. For purposes of the present invention, antigens can be obtained or derived from any appropriate source. Furthermore, for purposes of the present invention, an “antigen” includes a peptide having modifications, such as
25 deletions, additions and substitutions (generally conservative in nature) to the native sequence, so long as the peptide maintains sufficient immunogenicity. These modifications may be deliberate, for example through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the antigens.

30 An “immune response” against an antigen of interest is the development in an individual of a humoral and/or a cellular immune response to that antigen.

For purposes of the present invention, a “humoral immune response” refers to an immune response mediated by antibody molecules, while a “cellular immune response” is one mediated by T-lymphocytes and/or other white blood cells.

5 The term “nucleic acid immunization” is used herein to refer to the introduction of a nucleic acid molecule encoding one or more selected antigens into a host cell for the *in vivo* expression of the antigen or antigens. The term also encompasses introduction of a nucleic acid molecule encoding one or more selected adjuvants into a host cell for the *in vivo* expression of the adjuvant or adjuvants. The nucleic acid molecule can be introduced directly into the
10 recipient subject, such as by standard intramuscular or intradermal injection; transdermal particle delivery; inhalation; topically, or by oral, intranasal or mucosal modes of administration. The molecule alternatively can be introduced *ex vivo* into cells which have been removed from a subject. In this latter case, cells containing the nucleic acid molecule of interest are re-introduced into the
15 subject such that an immune response can be mounted against the antigen encoded by the nucleic acid molecule, or such that the adjuvant encoded by the nucleic acid molecule can exert its adjuvant effect.

The terms “nucleic acid molecule” and “polynucleotide” are used interchangeably herein and refer to a polymeric form of nucleotides of any length,
20 either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown. Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched
25 polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine (T) when the polynucleotide is RNA). Thus, the term nucleic
30 acid sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical representation can be input into databases in a computer having

a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

A “vector” is capable of transferring nucleic acid sequences to target cells (e.g., viral vectors, non-viral vectors, particulate carriers, and liposomes).

5 Typically, “vector construct,” “expression vector,” and “gene transfer vector,” mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors. A “plasmid” is a vector in the form of an extrachromosomal genetic element.

10 A nucleic acid sequence which “encodes” a selected adjuvant and/or antigen is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a
15 translation stop codon at the 3' (carboxy) terminus. For the purposes of the invention, such nucleic acid sequences can include, but are not limited to, cDNA from viral, procaryotic or eucaryotic mRNA, genomic sequences from viral or procaryotic DNA or RNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence.

20 A “promoter” is a nucleotide sequence which initiates and regulates transcription of a polypeptide-encoding polynucleotide. Promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably
25 linked to the promoter is repressed by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters. It is intended that the term “promoter” or “control element” includes full-length promoter regions and functional (e.g., controls transcription or translation) segments of these regions.

“Operably linked” refers to an arrangement of elements wherein the
30 components so described are configured so as to perform their usual function. Thus, a given promoter operably linked to a nucleic acid sequence is capable of

effecting the expression of that sequence when the proper enzymes are present. The promoter need not be contiguous with the sequence, so long as it functions to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the nucleic acid sequence and the promoter sequence can still be considered “operably linked” to the coding sequence.

“Recombinant” is used herein to describe a nucleic acid molecule (polynucleotide) of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation is not associated with all or a portion of the polynucleotide with which it is associated in nature and/or is linked to a polynucleotide other than that to which it is linked in nature. Two nucleic acid sequences which are contained within a single recombinant nucleic acid molecule are “heterologous” relative to each other when they are not normally associated with each other in nature.

An “isolated polynucleotide” is a nucleic acid molecule separate and discrete from the whole organism with which the molecule is found in nature; or a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences (as defined below) in association therewith. A sequence is “derived or obtained from” a molecule if it has the same or substantially the same base pair sequence as a region of the source molecule, its cDNA, complements thereof, or if it displays sequence identity as described below.

Techniques for determining nucleic acid and amino acid “sequence identity” or “sequence homology” also are known in the art. Typically, such techniques include determining the nucleotide sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. In general, “identity” refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more sequences (polynucleotide or amino acid) can be compared by determining their “percent identity.” The percent identity of two sequences,

whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences divided by the length of the shorter sequences and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, Atlas of Protein Sequences and Structure, M.O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA, and normalized by Gribskov, Nucl. Acids Res. 14(6):6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, WI) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, WI). A preferred method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details of these programs can be found at the following internet address:
<http://www.ncbi.nlm.gov/cgi-bin/BLAST>.

Alternatively, homology can be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are “substantially homologous” to each other when the sequences exhibit at least about 80%-85%, preferably at least about 90%, and most preferably at least about 95%-98% sequence identity over a defined length of the molecules, as determined using the methods above. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. For example, stringent hybridization conditions can include 50% formamide, 5x Denhardt’s Solution, 5x SSC, 0.1% SDS and 100 µg/ml denatured salmon sperm DNA and the washing conditions can include 2x SSC, 0.1% SDS at 37°C followed by 1x SSC, 0.1% SDS at 68°C. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning, supra*; *Nucleic Acid Hybridization, supra*.

The term “transdermal” delivery intends intradermal (e.g., into the dermis or epidermis), transdermal (e.g., “percutaneous”) and transmucosal administration, i.e., delivery by passage of an agent into or through skin or mucosal tissue. See, e.g., *Transdermal Drug Delivery: Developmental Issues and Research Initiatives*, Hadgraft and Guy (eds.), Marcel Dekker, Inc., (1989); *Controlled Drug Delivery: Fundamentals and Applications*, Robinson and Lee (eds.), Marcel Dekker Inc., (1987); and *Transdermal Delivery of Drugs*, Vols. 1-3, Kydonieus and Berner (eds.), CRC Press, (1987). Thus, the term encompasses delivery of an agent using a particle delivery device (e.g., a needleless syringe) such as those described in U.S. Patent No. 5,630,796, as well as delivery using particle-mediated delivery devices such as those described in U.S. Patent No. 5,865,796.

By “core carrier” is meant a carrier on which a guest nucleic acid (e.g., DNA, RNA) is coated in order to impart a defined particle size as well as a sufficiently high density to achieve the momentum required for cell membrane penetration, such that the guest molecule can be delivered using particle-mediated techniques (see, e.g., U.S. Patent No. 5,100,792). Core carriers typically include materials such as tungsten, gold, platinum, ferrite, polystyrene and latex. See e.g., *Particle Bombardment Technology for Gene Transfer*, (1994) Yang, N. ed., Oxford University Press, New York, NY pages 10-11.

By “particle delivery device” is meant an instrument which delivers a particulate composition transdermally without the aid of a conventional needle to pierce the skin. Particle delivery devices for use with the present invention are discussed throughout this document.

As used herein, the term “treatment” includes any of following: the prevention of infection or reinfection; the reduction or elimination of symptoms; and the reduction or complete elimination of a pathogen. Treatment may be effected prophylactically (prior to infection) or therapeutically (following infection).

The terms “individual” and “subject” are used interchangeably herein to refer to any member of the subphylum cordata, including, without limitation, humans and other primates, including non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The terms do not denote a particular age. Thus, both adult and newborn individuals are intended to be covered. The methods described herein are intended for use in any of the above vertebrate species, since the immune systems of all of these vertebrates operate similarly.

General Overview

Before describing the present invention in detail, it is to be understood

that this invention is not limited to particular formulations or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

5 The present invention provides novel compositions containing nucleic acid sequences, wherein a first sequence in the composition is a coding sequence for an A subunit obtained or derived from an ADP-ribosylating bacterial toxin, and a second sequence in the composition is a coding sequence for a B subunit obtained or derived from an ADP-ribosylating bacterial toxin. The first and
10 second sequences are useful in immunization methods wherein they are delivered to a subject in order to provide for an adjuvant effect (against a co-administered antigen of interest) in the immunized subject. ADP-ribosylating bacterial toxins are a family of related bacterial exotoxins and include diphtheria toxin (DT), pertussis toxin (PT), cholera toxin (CT), the *E. coli* heat-labile toxins (LT1 and
15 LT2), *Pseudomonas* endotoxin A, *Pseudomonas* exotoxin S, *B. cereus* exoenzyme, *B. sphaericus* toxin, *C. botulinum* C2 and C3 toxins, *C. limosum* exoenzyme, as well as toxins from *C. perfringens*, *C. spiriforma* and *C. difficile*, *Staphylococcus aureus* EDIN, and ADP-ribosylating bacterial toxin mutants such as CRM₁₉₇, a non-toxic diphtheria toxin mutant (see, e.g., Bixler et al. (1989)
20 *Adv. Exp. Med. Biol.* 251:175; and Constantino et al. (1992) *Vaccine*). Most ADP-ribosylating bacterial toxins are organized as an A:B multimer, wherein the A subunit contains the ADP-ribosyltransferase activity, and the B subunit acts as the binding moiety. Preferred ADP-ribosylating bacterial toxins for use in the compositions of the present invention include cholera toxin and the *E. coli* heat-labile toxins.
25 labile toxins.

 Cholera toxin (CT) and the related *E. coli* heat labile enterotoxins (LT) are secretion products of their respective enterotoxic bacterial strains that are potent immunogens and exhibit strong toxicity when administered systemically, orally, or mucosally. Both CT and LT are known to provide adjuvant effects for antigen
30 when administered via the intramuscular or oral routes. These adjuvant effects have been observed at doses below that required for toxicity. The two toxins are

extremely similar molecules, and are at least about 70-80% homologous at the amino acid level.

5 The CT and LT toxins are hexamers, composed of a single molecule of an A subunit surrounded by a doughnut-shaped ring composed of 5 molecules of the B subunit. The A subunit possesses an ADP-ribosylase activity resulting in G protein modifications that lead to cAMP and protein kinase A upregulation following internalization of the A subunit in a mammalian cell. The A subunit can be nicked by exogenous proteases yielding the A1 and A2 fragments linked via a single disulfide bridge. The A subunit of CT contains a C-terminal KDEL peptide motif (RDEL for the A subunit of LT) that is associated with retrieval of proteins from the trans-Golgi network into the ER. This is likely important for delivery of the A fragment to the correct cellular compartment for toxicity following internalization, and retention of the toxin within that cellular compartment. The A1 subunit enters the cell cytoplasm and triggers Cl-efflux by catalysing ADP-ribosylation of a G protein, which activates adenylate cyclase leading to elevated levels of cAMP. Elevated cAMP causes protein kinase A to phosphorylate and open the cystic fibrosis transmembrane conductance regulator chloride channel.

20 The A2 fragment's main role is in interacting with the B subunit. Toxin internalization is mediated by the five copies of the B subunit which possess binding activity for the GM1 ganglioside, a glycosphingolipid found ubiquitously on the surface of mammalian cells.

25 The relative importance of the A and B subunits for adjuvanticity is controversial. There is speculation in the field that the toxic activity of the A subunit can modulate the adjuvant effects associated with the B subunit. Some studies demonstrate that mutations in the A subunit abrogate adjuvant activity while others show that A subunit mutations that block ADP-ribosylase activity have no effect on adjuvanticity. Other reports show varying levels of adjuvant effects for purified B subunits or recombinant B subunit preparations. It is likely that the A and B subunits have separate functions that independently contribute to adjuvanticity. These functions are ADP-ribosylase activity and receptor

30

triggering activity, respectively. Regarding the B subunit of LT in particular, binding to the GM1 receptor on B cells results in polyclonal activation, occurs in the absence of significant proliferation, and involves the upregulation of a number of important molecules such as MHC class II, B7, CD40, ICAM-1 and IL2-R α . For T cells, addition of CT or LT holotoxins or recombinant B subunits to concanavilin A-stimulated T cells results in a reduction in thymidine incorporation. Affects of these molecules on macrophages and dendritic cells have not been reported. In PBMC cultures, EtxB stimulates high level TNF- α and IL-10 production but not IL-12. This is consistent with the observations of predominately Th2-like responses associated with the use of CT and LT forms as adjuvants.

It is therefore a surprising feature of the present invention that administration of the present polynucleotide adjuvant compositions preferentially produces an augmented Th1-like immune response against the co-administered antigen of interest, rather than the Th2-like response that would be expected from the use of an ADP-ribosylating exotoxin adjuvant composition.

Coding Sequences for ADP-ribosylating Exotoxin Subunits

In one embodiment, a composition is provided which includes one or more recombinant nucleic acid molecules, said one or more molecules containing first and second nucleic acid sequences wherein (a) the first nucleic acid sequence is a coding region for a truncated A subunit peptide obtained or derived from an ADP-ribosylating exotoxin, and (b) the second nucleic acid sequence is a coding region for a truncated B subunit peptide obtained or derived from an ADP-ribosylating exotoxin. The coding regions provide for "truncated" subunit peptides in that each said subunit coding region has been genetically altered so as to create a 5' deletion, whereby the encoded subunit peptide does not have an amino terminal bacterial signal peptide.

In another embodiment, a composition is provided which includes one or more recombinant nucleic acid molecules, said one or more molecules containing first and second nucleic acid sequences wherein (a) the first nucleic acid sequence

is a coding region for a modified, mature A subunit peptide obtained or derived from an ADP-ribosylating exotoxin, and (b) the second nucleic acid sequence is a coding region for a mature B subunit peptide obtained or derived from an ADP-ribosylating exotoxin. The first nucleic acid sequence contains a coding region that provides for a “modified” A subunit peptide in that the said coding region has been genetically altered so as to delete a four amino acid residue C-terminal KDEL or RDEL motif normally found in the encoded subunit peptide.

In still a further embodiment, a composition is provided which includes one or more recombinant nucleic acid molecules, said one or more molecules containing first and second nucleic acid sequences wherein (a) the first nucleic acid sequence is a coding region for a truncated and modified A subunit peptide obtained or derived from an ADP-ribosylating exotoxin, and (b) the second nucleic acid sequence is a coding region for a truncated B subunit peptide obtained or derived from an ADP-ribosylating exotoxin.

In particularly preferred embodiments, the ADP-ribosylating exotoxin peptide subunit coding sequences are obtained or derived from a cholera toxin (CT). In other particularly preferred embodiments, the ADP-ribosylating exotoxin subunit peptide coding sequences are obtained or derived from an *E. coli* heat labile enterotoxin (LT), for example LT1 or LT2.

Portions of the genomes of various enterotoxic bacterial species, particularly those portions containing the coding sequences for ADP-ribosylating exotoxins, are generally known and the sequences therefor are publically available, for example on the World Wide Web, and/or are deposited with repositories such as the GenBank database. For example, a GenBank entry for the complete sequences of the CT subunit A and B genes can be found at Locus VIBCTXABB (Accession No. D30053), while a GenBank entry for the complete sequences of the LT subunit A and B genes can be found at Locus AB0116677 (Accession No. AB011677). Active variants or fragments of these toxin sequences may also be used in the compositions and methods of the present invention. For a general discussion of ADP-ribosylating exotoxins, see e.g., Krueger et al. (1995) *Clin. Microbiol. Rev.* 8:34-47. Furthermore, in certain

aspects of the invention, one or both of the toxin subunit peptide coding regions can be further genetically modified to detoxify the subunit peptide(s) encoded thereby. For example, genetically altered toxin mutants which have disrupted ADP-ribosylating activity, trypsin cleavage site mutations, or disrupted binding
5 activity are known in the art. See, e.g., Burnette et al. (1994) "Recombinant microbial ADP-ribosylating toxins of *Bordetella pertussis*, *Vibrio cholerae*, and enterotoxigenic *Escherichia coli*: structure, function and toxoid vaccine development," in *Bioprocess Technology*, Burnette et al. eds., pp. 185-203; Rappuoli et al. (1995) *Int. Archiv. Allergy Immunol.* 108:327-333; and Rappuoli
10 et al. (1996) *Ad. Exp. Med. Biol.* 397:55-60. Sequences encoding the selected toxin subunits are typically inserted into an appropriate vector (e.g., a plasmid backbone) using known techniques and as described below in the Examples.

The sequence or sequences encoding the ADP-ribosylating exotoxin subunit peptides of interest can be obtained and/or prepared using known
15 methods. For example, substantially pure preparations can be obtained using standard molecular biological tools. That is, polynucleotide sequences coding for the above-described toxin subunits can be obtained using recombinant methods, such as by screening cDNA libraries from cells expressing the toxin subunits, or by deriving the coding sequence for the toxin subunits from a vector known to
20 include the same. The toxin subunit coding sequences from various bacterial strains are on deposit with the American Type Culture Collection ATCC, and yet others are available from national and international health organizations such as the Centers of Disease Control (Atlanta, GA). See, e.g., Sambrook et al., *supra*, for a description of techniques used to obtain and isolate nucleic acid molecules.
25 Polynucleotide sequences can also be produced synthetically, rather than cloned.

Yet another convenient method for isolating specific nucleic acid molecules is by the polymerase chain reaction (PCR). Mullis et al. (1987) *Methods Enzymol.* 155:335-350. This technique uses DNA polymerase, usually a
thermostable DNA polymerase, to replicate a desired region of DNA. The region
30 of DNA to be replicated is identified by oligonucleotides of specified sequence complementary to opposite ends and opposite strands of the desired DNA to

prime the replication reaction. The product of the first round of replication is itself a template for subsequent replication, thus repeated successive cycles of replication result in geometric amplification of the DNA fragment delimited by the primer pair used.

5 Once the relevant sequences for the ADP-ribosylating exotoxin subunits of interest have been obtained, they can be linked together to provide one or more contiguous nucleic acid molecules using standard cloning or molecular biology techniques. More particularly, after sequence information for the selected toxin subunit combination has been obtained, the coding sequences can be combined
10 with each other or with other sequences to form a hybrid sequence, or handled separately. In hybrid sequences, the subunit peptide coding sequences can be positioned in any manner relative to each other, and be included in a single molecule in any number ways, for example, as a single copy, randomly repeated in the molecule as multiple copies, or included in the molecule as multiple
15 tandem repeats or otherwise ordered repeat motifs.

 Although any number of routine molecular biology techniques can be used to construct such recombinant nucleic acid molecules, one convenient method entails using one or more unique restriction sites in a shuttle or cloning
20 vector (or inserting one or more unique restriction sites into a suitable vector sequence) and standard cloning techniques to direct the subunit peptide coding sequence or sequences into particular target locations within a vector.

 Alternatively, hybrid molecules can be produced synthetically rather than cloned. The nucleotide sequence can be designed with the appropriate codons for the particular amino acid sequence desired. In general, one will select preferred
25 codons for the intended host in which the sequence will be expressed. The complete sequence can then be assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence. See, e.g., Edge (1981) *Nature* 292:756; Nambair et al. (1984) *Science* (1984) 223:1299; Jay et al. (1984) *J. Biol. Chem.* 259:6311.

30 Once the relevant ADP-ribosylating exotoxin peptide coding sequences have been obtained or constructed, they can be inserted into one or more suitable

vectors which include control sequences operably linked to the inserted sequence or sequences, thus providing expression cassettes that allow for expression of the toxin subunit peptides *in vivo* in a targeted subject species.

5 Typical promoters for mammalian cell expression include the SV40 early promoter, a CMV promoter such as the CMV immediate early promoter, the mouse mammary tumor virus LTR promoter, the adenovirus major late promoter (Ad MLP), and other suitably efficient promoter systems. Nonviral promoters, such as a promoter derived from the murine metallothionein gene, may also be used for mammalian expression. Inducible, repressible or otherwise controllable
10 promoters may also be used. Typically, transcription termination and polyadenylation sequences will also be present, located 3' to each translation stop codon. Preferably, a sequence for optimization of initiation of translation, located 5' to each coding sequence, is also present. Examples of transcription terminator/polyadenylation signals include those derived from SV40, as described
15 in Sambrook et al., *supra*, as well as a bovine growth hormone terminator sequence. Introns, containing splice donor and acceptor sites, may also be designed into the expression cassette.

In addition, enhancer elements may be included within the expression cassettes in order to increase expression levels. Examples of suitable enhancers
20 include the SV40 early gene enhancer (Dijkema et al. (1985) *EMBO J.* 4:761), the enhancer/promoter derived from the long terminal repeat (LTR) of the Rous Sarcoma Virus (Gorman et al. (1982) *Proc. Natl. Acad. Sci. USA* 79:6777), and elements derived from human or murine CMV (Boshart et al. (1985) *Cell* 41:521), for example, elements included in the CMV intron A sequence.

25 In some particular embodiments, a further ancillary sequence can be included which provides for secretion of the attached ADP-ribosylating exotoxin subunit peptide from a mammalian cell. Such secretion leader sequences are known to those skilled in the art, and include, for example, the tissue plasminogen activator (tpa) leader signal sequence.

30 After one or more suitable expression cassettes (or nucleic acid constructs such as plasmid vectors) have been produced that contain the ADP-ribosylating

exotoxin subunit peptide coding sequences of interest, the expression cassettes can be provided in a suitable transfection vector such as a DNA plasmid vector or a viral vector for subsequent administration to a subject. In this regard, the polynucleotide compositions of the invention can be used as standalone adjuvant compositions, or co-administered with an antigen of interest, e.g., as part of a multi-component vaccine composition. For example, in a multi-component vaccine composition, the present nucleic acid molecules (containing the coding sequences for the ADP-ribosylating exotoxin subunit peptides of interest) can be combined with additional nucleic acid molecules encoding one or more antigens known to be important for providing protection against a pathogen, for example, molecules containing sequences that encode influenza HA or NA antigens, or molecules containing sequences that encode HIV antigens. Alternatively, the multi-component vaccine composition may contain, in addition to the polynucleotides encoding the toxin subunits, antigen in the form of conventional whole virus, split virus, polysaccharide, or purified subunit vaccine preparations as described herein below.

Antigens

The compositions and methods described herein are useful in adjuvanting an immune response against a wide variety of co-administered antigens, for example antigens obtained or derived from diseased cells or tissues, of from human or animal pathogens. For the purposes of the present invention, an adjuvant is used to either enhance the immune response to a specific antigen, e.g., when an adjuvant is co-administered with a vaccine composition, the immune response is greater than the immune response elicited by an equivalent amount of the vaccine composition administered without the adjuvant, or the adjuvant is used to direct a particular type or class of immune response against a co-administered antigen. Co-administration of an "effective amount" of the adjuvant compositions of the present invention will be that amount which enhances an immunological response to the co-administered antigen such that, for example,

lower or fewer doses of the antigen are required to generate an efficient immune response.

As used herein, the term “co-administered,” such as when an ADP-ribosylating exotoxin subunit encoding adjuvant composition according to the present invention is “co-administered” with an antigen of interest (e.g., a vaccine composition), intends either the simultaneous or concurrent administration of the adjuvant composition and the antigen, e.g., when the two are present in the same composition or administered in separate compositions at nearly the same time but at different sites, as well as the delivery of the adjuvant composition and antigen in separate compositions at different times, including delivery to different sites. For example, the adjuvant composition may be delivered prior or subsequent to delivery of the antigen at the same or a different site. The timing between adjuvant and antigen deliveries can range from about several minutes apart, to several hours apart, to several days apart.

For the purposes of the instant invention, the term “pathogen” is used in a broad sense to refer to a specific causative agent of a disease or condition, and includes any agent that provides a source of a molecule that elicits an immune response. Thus, pathogens include, but are not limited to, viruses, bacteria, fungi, protozoa, parasites, cancer cells and the like. Typically, the immune response is elicited by one or more peptide or carbohydrate antigens produced by the pathogen. Methods for identifying suitable antigens, obtaining and preparing such molecules, and then determining suitable dosages, assaying for suitable immunogenicity and treating with such antigens are well known in the art. See e.g., Plotkin et al. (1994) *Vaccines*, 2nd Edition, W.B. Saunders, Philadelphia, PA. Non-limiting examples of sources for antigens that can be used to vaccinate vertebrate subjects, particularly, humans and non-human mammals, thus include viruses, bacteria, fungi, and other pathogenic organisms.

Suitable viral antigens include, but are not limited to, those obtained or derived from the hepatitis family of viruses, including hepatitis A virus (HAV), hepatitis B virus (HBV), hepatitis C virus (HCV), the delta hepatitis virus (HDV), hepatitis E virus (HEV) and hepatitis G virus (HGV). See, e.g., International

Publication Nos. WO 89/04669; WO 90/11089; and WO 90/14436. The HCV genome encodes several viral proteins, including E1 and E2. See, e.g., Houghton et al. (1991) *Hepatology* 14:381-388. Genomic fragments containing sequences encoding these proteins, as well as antigenic fragments thereof, will find use in
5 the present methods. Similarly, the coding sequence for the δ -antigen from HDV is known (see, e.g., U.S. Patent No. 5,378,814).

In like manner, a wide variety of proteins from the herpesvirus family can be used as antigens in the present invention, including proteins derived from herpes simplex virus (HSV) types 1 and 2, such as HSV-1 and HSV-2
10 glycoproteins gB, gD and gH; antigens from varicella zoster virus (VZV), Epstein-Barr virus (EBV) and cytomegalovirus (CMV) including CMV gB and gH; and antigens from other human herpesviruses such as HHV6 and HHV7. (See, e.g. Chee et al. (1990) *Cytomegaloviruses* (J.K. McDougall, ed., Springer-Verlag, pp. 125-169; McGeoch et al. (1988) *J. Gen. Virol.* 69:1531-1574; U.S.
15 Patent No. 5,171,568; Baer et al. (1984) *Nature* 310:207-211; and Davison et al. (1986) *J. Gen. Virol.* 67:1759-1816.)

Human immunodeficiency virus (HIV) antigens, such as gp120 molecules for a multitude of HIV-1 and HIV-2 isolates, including members of the various genetic subtypes of HIV, are known and reported (see, e.g., Myers et al., Los
20 Alamos Database, Los Alamos National Laboratory, Los Alamos, New Mexico (1992); and Modrow et al. (1987) *J. Virol.* 61:570-578) and antigen-containing genomic fragments derived or obtained from any of these isolates will find use in the present invention. Furthermore, other immunogenic proteins derived or
25 obtained from any of the various HIV isolates will find use herein, including fragments containing one or more of the various envelope proteins such as gp160 and gp41, gag antigens such as p24gag and p55gag, as well as proteins derived from the pol, env, tat, vif, rev, nef, vpr, vpu and LTR regions of HIV.

Antigens derived or obtained from other viruses will also find use herein, such as without limitation, antigens from members of the families Picornaviridae
30 (e.g., polioviruses, rhinoviruses, etc.); Caliciviridae; Togaviridae (e.g., rubella virus, dengue virus, etc.); Flaviviridae; Coronaviridae; Reoviridae (e.g., rotavirus,

etc.); Birnaviridae; Rhabdoviridae (e.g., rabies virus, etc.); Orthomyxoviridae (e.g., influenza virus types A, B and C, etc.); Filoviridae; Paramyxoviridae (e.g., mumps virus, measles virus, respiratory syncytial virus, parainfluenza virus, etc.); Bunyaviridae; Arenaviridae; Retroviridae (e.g., HTLV-I; HTLV-II; HIV-1 (also known as HTLV-III, LAV, ARV, hTLR, etc.)), including but not limited to antigens from the isolates HIV_{IIIb}, HIV_{SF2}, HIV_{LAV}, HIV_{LAI}, HIV_{MN}); HIV-1_{CM235}, HIV-1_{US4}; HIV-2, among others; simian immunodeficiency virus (SIV); Papillomavirus, the tick-bourne encephalitis viruses; and the like. See, e.g. Virology, 3rd Edition (W.K. Joklik ed. 1988); *Fundamental Virology*, 2nd Edition (B.N. Fields and D.M. Knipe, eds. 1991), for a description of these and other viruses.

In some contexts, it may be preferable that the selected viral antigens are obtained or derived from a viral pathogen that typically enters the body via a mucosal surface and is known to cause or is associated with human disease, such as, but not limited to, HIV (AIDS), influenza viruses (Flu), herpes simplex viruses (genital infection, cold sores, STDs), rotaviruses (diarrhea), parainfluenza viruses (respiratory infections), poliovirus (poliomyelitis), respiratory syncytial virus (respiratory infections), measles and mumps viruses (measles, mumps), rubella virus (rubella), and rhinoviruses (common cold).

Genomic fragments containing bacterial and parasitic antigens can be obtained or derived from known causative agents responsible for diseases including, but not limited to, Diphtheria, Pertussis, Tetanus, Tuberculosis, Bacterial or Fungal Pneumonia, Otitis Media, Gonorrhea, Cholera, Typhoid, Meningitis, Mononucleosis, Plague, Shigellosis or Salmonellosis, Legionaire's Disease, Lyme Disease, Leprosy, Malaria, Hookworm, Onchocerciasis, Schistosomiasis, Trypanosomiasis, Leishmaniasis, Giardia, Amoebiasis, Filariasis, Borelia, and Trichinosis. Still further antigens can be obtained or derived from unconventional viruses such as the causative agents of kuru, Creutzfeldt-Jakob disease (CJD), scrapie, transmissible mink encephalopathy, and chronic wasting diseases, or from proteinaceous infectious particles such as prions that are associated with mad cow disease.

Specific pathogens can include *M. tuberculosis*, *Chlamydia*, *N. gonorrhoeae*, *Shigella*, *Salmonella*, *Vibrio Cholera*, *Treponema pallidum*, *Pseudomonas*, *Bordetella pertussis*, *Brucella*, *Francisella tularensis*, *Helicobacter pylori*, *Leptospira interrogans*, *Legionella pneumophila*, *Yersinia pestis*, Streptococcus (types A and B), *Pneumococcus*, *Meningococcus*, *Hemophilus influenza* (type b), *Toxoplasma gondii*, *Compylobacteriosis*, *Moraxella catarrhalis*, *Donovanosis*, and *Actinomyces*; fungal pathogens including Candidiasis and Aspergillosis; parasitic pathogens including Taenia, Flukes, Roundworms, Amebiasis, Giardiasis, Cryptosporidium, Schistosoma, *Pneumocystis carinii*, Trichomoniasis and Trichinosis. Thus, the present invention can also be used to provide a suitable immune response against numerous veterinary diseases, such as Foot and Mouth diseases, Coronavirus, *Pasteurella multocida*, *Helicobacter*, *Strongylus vulgaris*, *Actinobacillus pleuropneumonia*, Bovine viral diarrhoea virus (BVDV), *Klebsiella pneumoniae*, *E. coli*, *Bordetella pertussis*, *Bordetella parapertussis* and *bronchiseptica*.

In some embodiments, the antigen of interest can be an allergen. An “allergen” is an antigen which can initiate a state of hypersensitivity, or which can provoke an immediate hypersensitivity reaction in an individual already sensitized with the allergen. Allergens are commonly proteins or chemicals bound to proteins which have the property of being allergenic; however, allergens can also include organic or inorganic materials derived from a variety of man-made or natural sources such as plant materials, metals, ingredients in cosmetics or detergents, latexes, or the like. Classes of suitable allergens for use in the methods of the invention can include, but are not limited to, pollens, animal dander, grasses, molds, dusts, antibiotics, stinging insect venoms, and a variety of environmental (including chemicals and metals), drug and food allergens. Common tree allergens include pollens from cottonwood, poplar, ash, birch, maple, oak, elm, hickory, and pecan trees; common plant allergens include those from rye, ragweed, English plantain, sorrel-dock and pigweed; plant contact allergens include those from poison oak, poison ivy and nettles; common grass allergens include Timothy, Johnson, Bermuda, fescue and bluegrass allergens;

common allergens can also be obtained from molds or fungi such as *Alternaria*, *Fusarium*, *Hormodendrum*, *Aspergillus*, *Micropolyspora*, *Mucor* and thermophilic actinomycetes; penicillin and tetracycline are common antibiotic allergens; epidermal allergens can be obtained from house or organic dusts (typically fungal in origin), from insects such as house mites (*dermatophagoides pterosinyssis*), or from animal sources such as feathers, and cat and dog dander; common food allergens include milk and cheese (diary), egg, wheat, nut (e.g., peanut), seafood (e.g., shellfish), pea, bean and gluten allergens; common environmental allergens include metals (nickel and gold), chemicals (formaldehyde, trinitrophenol and turpentine), Latex, rubber, fiber (cotton or wool), burlap, hair dye, cosmetic, detergent and perfume allergens; common drug allergens include local anesthetic and salicylate allergens; antibiotic allergens include penicillin and sulfonamide allergens; and common insect allergens include bee, wasp and ant venom, and cockroach calyx allergens. Particularly well characterized allergens include, but are not limited to, the major and cryptic epitopes of the Der pI allergen (Hoyne et al. (1994) *Immunology* 83:190-195), bee venom phospholipase A2 (PLA) (Akdis et al. (1996) *J. Clin. Invest.* 98:1676-1683), birch pollen allergen Bet v 1 (Bauer et al. (1997) *Clin. Exp. Immunol.* 107:536-541), and the multi-epitopic recombinant grass allergen rKBG8.3 (Cao et al. (1997) *Immunology* 90:46-51). These and other suitable allergens are commercially available and/or can be readily prepared as extracts following known techniques.

In certain other embodiments, the antigen of interest can be a tumor-specific antigen. For the purposes of the present invention, tumor-specific antigens include, but are not limited to, any of the various MAGEs (melanoma associated antigen E), including MAGE 1, MAGE 2, MAGE 3 (HLA-A1 peptide), MAGE 4, etc.; any of the various tyrosinases (HLA-A2 peptide); mutant ras; mutant p53; and p97 melanoma antigen. Other tumor-specific antigens include the Ras peptide and p53 peptide associated with advanced cancers, the HPV 16/18 and E6/E7 antigens associated with cervical cancers, MUC1-KLH antigen associated with breast carcinoma, CEA (carcinoembryonic antigen)

associated with colorectal cancer, gp100 or MART1 antigens associated with melanoma, and the PSA antigen associated with prostate cancer. The *p53* gene sequence is known (see e.g., Harris et al. (1986) *Mol. Cell. Biol.* 6:4650-4656) and is deposited with GenBank under Accession No. M14694. Thus, the
5 adjuvant compositions of the present invention can be used to carry out immunotherapeutic methods for treating cervical, breast, colorectal, prostate, lung cancers, and melanomas.

Antigens for use with the present invention can be obtained or produced using a variety of methods known to those of skill in the art. In particular, the
10 antigens can be isolated directly from native sources, using standard purification techniques. Alternatively, the antigens can be produced recombinantly using known techniques. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Vols. I, II and III, Second Edition (1989); DNA Cloning, Vols. I and II (D.N. Glover ed. 1985). Antigens for use herein may also be
15 synthesized, based on described amino acid sequences, via chemical polymer syntheses such as solid phase peptide synthesis. Such methods are known to those of skill in the art. See, e.g., J. M. Stewart and J. D. Young, *Solid Phase Peptide Synthesis*, 2nd Ed., Pierce Chemical Co., Rockford, IL (1984) and G. Barany and R. B. Merrifield, *The Peptides: Analysis, Synthesis, Biology*, editors
20 E. Gross and J. Meienhofer, Vol. 2, Academic Press, New York, (1980), pp. 3-254, for solid phase peptide synthesis techniques; and M. Bodansky, *Principles of Peptide Synthesis*, Springer-Verlag, Berlin (1984) and E. Gross and J. Meienhofer, Eds., *The Peptides: Analysis, Synthesis, Biology, supra*, Vol. 1, for classical solution synthesis.

25 If desired, polynucleotide sequences coding for the above-described antigens, can be obtained using recombinant methods, such as by screening cDNA and genomic libraries from cells expressing the gene, or by deriving the gene from a vector known to include the same. Furthermore, the desired gene can be isolated directly from cells and tissues containing the same, using standard
30 techniques, such as phenol extraction and PCR of cDNA or genomic DNA. See, e.g., Sambrook et al., *supra*, for a description of techniques used to obtain and

isolate DNA. Polynucleotide sequences can also be produced synthetically, rather than cloned.

In those compositions wherein the antigen component will be provided by way of a nucleic acid sequence encoding an antigen of interest, the coding
5 sequence for the selected antigen can be combined with one or both of the coding sequences for the ADP-ribosylating exotoxin subunit peptides to provide a single construct carrying all three coding sequences, combined with just one of the coding sequences for a toxin subunit to provide, e.g., a two plasmid composition, or provided in a separate construct from either a single construct, or multiple
10 constructs carrying the toxin subunit coding sequences. In each of the above cases, the antigen can be operably linked to and under the transcriptional control of the same or different control elements, e.g., promoters and enhancers. In those constructs where a single control element is used to direct transcription of two or more coding sequences, an internal ribosome entry sequence (IRES) can be used
15 to facilitate transcription of the multiple sequences.

Administration of Polynucleotides

The polynucleotides (nucleic acid molecules containing coding sequences for the selected ADP-ribosylating exotoxin subunit peptides) described herein
20 may be administered by any suitable method. In a preferred embodiment, described below, the polynucleotides are administered by coating one or more suitable constructs (e.g., one or more DNA plasmid constructs) containing the coding sequences for the ADP-ribosylating exotoxin subunit peptides of interest (and, in certain embodiments, a coding sequence for an antigen of interest on the
25 same or a different construct) onto core carrier particles and then administering the coated particles to the subject or cells. However, the polynucleotides of the present invention may also be delivered using a viral vector or using non-viral systems, e.g., naked nucleic acid delivery.

Viral Vectors

30 A number of viral based systems have been used for gene delivery. For example, retroviral systems are known and generally employ packaging lines

which have an integrated defective provirus (the “helper”) that expresses all of the genes of the virus but cannot package its own genome due to a deletion of the packaging signal, known as the *psi* sequence. Thus, the cell line produces empty viral shells. Producer lines can be derived from the packaging lines which, in addition to the helper, contain a viral vector which includes sequences required in *cis* for replication and packaging of the virus, known as the long terminal repeats (LTRs). The polynucleotide molecule(s) of interest can be inserted into the vector and packaged in the viral shells synthesized by the retroviral helper. The recombinant virus can then be isolated and delivered to a subject. (See, e.g., U.S. Patent No. 5,219,740.) Representative retroviral vectors include but are not limited to vectors such as the LHL, N2, LNSAL, LSHL and LHL2 vectors described in e.g., U.S. Patent No. 5,219,740, incorporated herein by reference in its entirety, as well as derivatives of these vectors, such as the modified N2 vector described herein. Retroviral vectors can be constructed using techniques well known in the art. See, e.g., U.S. Patent No 5,219,740; Mann et al. (1983) *Cell* 33:153-159.

Adenovirus based systems have been developed for gene delivery and are suitable for delivering the nucleic acid molecules described herein. Human adenoviruses are double-stranded DNA viruses which enter cells by receptor-mediated endocytosis. These viruses are particularly well suited for genetic transfer because they are easy to grow and manipulate and they exhibit a broad host range *in vivo* and *in vitro*. For example, adenoviruses can infect human cells of hematopoietic, lymphoid and myeloid origin. Furthermore, adenoviruses infect quiescent as well as replicating target cells. Unlike retroviruses which integrate into the host genome, adenoviruses persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis. The virus is easily produced at high titers and is stable so that it can be purified and stored. Even in the replication-competent form, adenoviruses cause only low level morbidity and are not associated with human malignancies. Accordingly, adenovirus vectors have been developed which make use of these advantages. For a description of adenovirus vectors and their uses *see, e.g.,* Haj-Ahmad and Graham (1986) *J.*

Virol. 57:267-274; Bett et al. (1993) *J. Virol.* 67:5911-5921; Mittereder et al. (1994) *Human Gene Therapy* 5:717-729; Seth et al. (1994) *J. Virol.* 68:933-940; Barr et al. (1994) *Gene Therapy* 1:51-58; Berkner, K.L. (1988) *BioTechniques* 6:616-629; Rich et al. (1993) *Human Gene Therapy* 4:461-476.

5 Adeno-associated viral vectors (AAV) can also be used to administer the polynucleotide molecules described herein. In this regard, AAV vectors can be derived from any AAV serotype, including without limitation, AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAVX7, etc. AAV vectors can have one or more of the AAV wild-type genes deleted in whole or in part, preferably the *rep* and/or
10 *cap* genes, but retain one or more functional flanking inverted terminal repeat (ITR) sequences. A functional ITR sequence is generally deemed necessary for the rescue, replication and packaging of the AAV virion. Thus, an AAV vector includes at least those sequences required in *cis* for replication and packaging (e.g., a functional ITR) of the virus. The ITR need not be the wild-type
15 nucleotide sequence, and may be altered, e.g., by the insertion, deletion or substitution of nucleotides, so long as the sequence provides for functional rescue, replication and packaging.

AAV expression vectors are constructed using known techniques to at least provide as operatively linked components in the direction of transcription,
20 control elements including a transcriptional initiation region, the DNA of interest and a transcriptional termination region. The control elements are selected to be functional in a mammalian cell. The resulting construct which contains the operatively linked components is bounded (5' and 3') with functional AAV ITR sequences. Suitable AAV constructs can be designed using techniques well
25 known in the art. *See, e.g.*, U.S. Patent Nos. 5,173,414 and 5,139,941; International Publication Nos. WO 92/01070 (published 23 January 1992) and WO 93/03769 (published 4 March 1993); Lebkowski et al. (1988) *Molec. Cell. Biol.* 8:3988-3996; Vincent et al. (1990) *Vaccines 90* (Cold Spring Harbor Laboratory Press); Carter, B.J. (1992) *Current Opinion in Biotechnology* 3:533-
30 539; Muzyczka, N. (1992) *Current Topics in Microbiol. and Immunol.* 158:97-129; Kotin, R.M. (1994) *Human Gene Therapy* 5:793-801; Shelling and Smith

(1994) *Gene Therapy* 1:165-169; and Zhou et al. (1994) *J. Exp. Med.* 179:1867-1875.

Conventional Pharmaceutical Preparations

5 Formulation of a preparation comprising the polynucleotide molecules of the present invention, with or without addition of an antigen composition, can be carried out using standard pharmaceutical formulation chemistries and methodologies all of which are readily available to the ordinarily skilled artisan. For example, compositions containing one or more suitable vectors can be combined with one or more pharmaceutically acceptable excipients or vehicles to
10 provide a liquid preparation.

Auxiliary substances, such as wetting or emulsifying agents, pH buffering substances and the like, may be present in the excipient or vehicle. These excipients, vehicles and auxiliary substances are generally pharmaceutical agents that do not induce an immune response in the individual receiving the
15 composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, polyethyleneglycol, hyaluronic acid, glycerol and ethanol. Pharmaceutically acceptable salts can also be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates,
20 and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. It is also preferred, although not required, that the preparation will contain a pharmaceutically acceptable excipient that serves as a stabilizer, particularly for peptide, protein or other like antigen molecules if they are to be included in the vaccine composition. Examples of suitable carriers that
25 also act as stabilizers for peptides include, without limitation, pharmaceutical grades of dextrose, sucrose, lactose, trehalose, mannitol, sorbitol, inositol, dextran, and the like. Other suitable carriers include, again without limitation, starch, cellulose, sodium or calcium phosphates, citric acid, tartaric acid, glycine, high molecular weight polyethylene glycols (PEGs), and combination thereof. A
30 thorough discussion of pharmaceutically acceptable excipients, vehicles and

auxiliary substances is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991), incorporated herein by reference.

Certain facilitators of nucleic acid uptake and/or expression ("transfection facilitating agents") can also be included in the compositions, for example, 5 facilitators such as bupivacaine, cardiotoxin and sucrose, and transfection facilitating vehicles such as liposomal or lipid preparations that are routinely used to deliver nucleic acid molecules. Anionic and neutral liposomes are widely available and well known for delivering nucleic acid molecules (see, e.g., *Liposomes: A Practical Approach*, (1990) RPC New Ed., IRL Press). Cationic 10 lipid preparations are also well known vehicles for use in delivery of nucleic acid molecules. Suitable lipid preparations include DOTMA (N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride), available under the tradename Lipofectin™, and DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane), see, e.g., Felgner et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7416; Malone et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:6077-6081; US Patent Nos 5,283,185 and 5,527,928, and International Publication Nos WO 90/11092, WO 91/15501 and WO 95/26356. These cationic lipids may preferably be used in association with a neutral lipid, for example DOPE (dioleyl phosphatidylethanolamine). Still further transfection-facilitating 20 compositions that can be added to the above lipid or liposome preparations include spermine derivatives (see, e.g., International Publication No. WO 93/18759) and membrane-permeabilizing compounds such as GALA, Gramicidine S and cationic bile salts (see, e.g., International Publication No. WO 93/19768).

25 Alternatively, the nucleic acid molecules of the present invention may be encapsulated, adsorbed to, or associated with, particulate carriers. Suitable particulate carriers include those derived from polymethyl methacrylate polymers, as well as PLG microparticles derived from poly(lactides) and poly(lactide-co-glycolides). See, e.g., Jeffery et al. (1993) *Pharm. Res.* 10:362-368. Other 30 particulate systems and polymers can also be used, for example, polymers such as

polylysine, polyarginine, polyornithine, spermine, spermidine, as well as conjugates of these molecules.

The formulated compositions will thus typically include one or more polynucleotide molecule (e.g., plasmid vector) containing the coding sequences for the selected ADP-ribosylating exotoxin subunit peptides in an amount sufficient to adjuvant an immunological response against a co-administered antigen. An appropriate effective amount can be readily determined by one of skill in the art. Such an amount will fall in a relatively broad range that can be determined through routine trials. For example, suitable adjuvant effect may be obtained using as little as 0.1 μ g of DNA, while in other administrations, up to 2mg of DNA may be used. It is generally expected that an effective dose of polynucleotides containing the ADP-ribosylating exotoxin subunit peptide coding sequences of interest will fall within a range of about 1 μ g to 1000 μ g, however, doses above and below this range may also be found effective. The compositions may thus contain from about 0.1% to about 99.9% of the polynucleotide molecules.

Administration of Conventional Pharmaceutical Preparations

Administration of the above-described pharmaceutical preparations can be effected in one dose, continuously or intermittently throughout the course of treatment. That is, once suitably formulated, the compositions of the present invention can be administered to a subject *in vivo* using a variety of known routes and techniques. For example, the liquid preparations can be provided as an injectable solution, suspension or emulsion and administered via parenteral, subcutaneous, intradermal, intramuscular, intravenous injection using a conventional needle and syringe, or using a liquid jet injection system. Liquid preparations can also be administered topically to skin or mucosal tissue, or provided as a finely divided spray suitable for respiratory or pulmonary administration. Other modes of administration include oral administration, suppositories, and active or passive transdermal delivery techniques.

Alternatively, the compositions can be administered *ex vivo*, for example delivery and reimplantation of transformed cells into a subject are known (e.g.,

dextran-mediated transfection, calcium phosphate precipitation, electroporation, and direct microinjection of into nuclei). However, delivery will most typically be via conventional needle and syringe for the liquid compositions and for liquid suspensions containing particulate compositions. Methods of determining the most effective means and dosages of administration are well known to those of skill in the art and will vary with the delivery vehicle, the composition of the therapy, the target cells, and the subject being treated. Single and multiple administrations can be carried out with the dose level and pattern being selected by the attending physician. It should be understood that more than one subunit coding region and/or antigen coding region can be carried by a single polynucleotide vector construct. Alternatively, separate vectors (*e.g.*, viral vectors, plasmids, or any combination thereof), each expressing one or more toxin subunit peptide and/or antigen derived from any pathogen can also be delivered to a subject as described herein.

Furthermore, it is also intended that the polynucleotides delivered by the methods of the present invention be combined with other suitable compositions and therapies. For instance, in order to further augment an immune response in a subject, the compositions and methods described herein can be combined with delivery of ancillary substances (*e.g.*, other adjuvants), such as pharmacological agents, cytokines, or the like. Ancillary substances may be administered, for example, as proteins or other macromolecules at the same time, prior to, or subsequent to, administration of the polynucleotide molecules described herein. The nucleic acid molecule compositions may also be administered directly to the subject or, alternatively, delivered *ex vivo*, to cells derived from the subject, using methods known to those skilled in the art.

Coated Particles

In one embodiment, polynucleotide constructs containing the coding sequences for the selected ADP-ribosylating exotoxin subunit peptides, and other ancillary components (such as coding sequences for one or more antigens) are delivered using carrier particles. Particle-mediated delivery methods for administering such nucleic acid preparations are known in the art. Thus, once

prepared and suitably purified, the above-described plasmid vector constructs can be coated onto carrier particles (*e.g.*, core carriers) using a variety of techniques known in the art. Carrier particles are selected from materials which have a suitable density in the range of particle sizes typically used for intracellular
5 delivery from an appropriate particle delivery device. The optimum carrier particle size will, of course, depend upon the diameter of the target cells.

For the purposes of the present invention, tungsten, gold, platinum and iridium core carrier particles can be used. Tungsten and gold particles are preferred. Tungsten particles are readily available in average sizes of 0.5 to 2.0
10 μm in diameter. Although such particles have optimal density for use in particle delivery methods, and allow highly efficient coating with DNA, tungsten may potentially be toxic to certain cell types. Accordingly, gold particles or microcrystalline gold (*e.g.*, gold powder A1570, available from Engelhard Corp., East Newark, NJ) will also find use with the present methods. Gold particles
15 provide uniformity in size (available from Alpha Chemicals in particle sizes of 1-3 μm , or available from Degussa, South Plainfield, NJ in a range of particle sizes including 0.95 μm) and reduced toxicity.

A number of methods are known and have been described for coating or precipitating DNA or RNA onto gold or tungsten particles. Most such methods
20 generally combine a predetermined amount of gold or tungsten with plasmid DNA, CaCl_2 and spermidine. The resulting solution is vortexed continually during the coating procedure to ensure uniformity of the reaction mixture. After precipitation of the nucleic acid, the coated particles can be transferred to suitable membranes and allowed to dry prior to use, coated onto surfaces of a sample
25 module or cassette, or loaded into a delivery cassette for use in a suitable particle delivery device.

Peptide antigens can also be coated onto the same or similar core carrier particles. For example, peptides can be attached to a carrier particle by simply mixing the two components in an empirically determined ratio, by ammonium
30 sulfate precipitation or other solvent precipitation methods familiar to those skilled in the art, or by chemical coupling of the peptide to the carrier particle.

The coupling of L-cysteine residues to gold has been previously described (Brown et al., *Chemical Society Reviews* 9:271-311 (1980)). Other methods would include, for example, dissolving the peptide in absolute ethanol, water, or an alcohol/water mixture, adding the solution to a quantity of carrier particles, and then drying the mixture under a stream of air or nitrogen gas while vortexing. Alternatively, the antigen can be dried onto carrier particles by centrifugation under vacuum. Once dried, the coated particles can be resuspended in a suitable solvent (e.g., ethyl acetate or acetone), and triturated (e.g., by sonication) to provide a substantially uniform suspension. The core carrier particles coated with the antigen can then be combined with core carrier particles carrying the ADP-ribosylating exotoxin subunit peptide construct(s) and administered in a single particle injection step, or administered separately from the toxin subunit compositions.

Administration of Coated Particles

Following their formation, core carrier particles coated with the nucleic acid preparations of the present invention, alone or in combination with e.g., antigen preparations, are delivered to a subject using particle-mediated delivery techniques.

Various particle delivery devices suitable for particle-mediated delivery techniques are known in the art, and are all suited for use in the practice of the invention. Current device designs employ an explosive, electric or gaseous discharge to propel the coated core carrier particles toward target cells. The coated particles can themselves be releasably attached to a movable carrier sheet, or removably attached to a surface along which a gas stream passes, lifting the particles from the surface and accelerating them toward the target. An example of a gaseous discharge device is described in U.S. Patent No. 5,204,253. An explosive-type device is described in U.S. Patent No. 4,945,050. One example of an electric discharge-type particle acceleration apparatus is described in U.S. Patent No. 5,120,657. Another electric discharge apparatus suitable for use herein is described in U.S. Patent No. 5,149,655. The disclosure of all of these patents is incorporated herein by reference in their entireties.

The coated particles are administered to the subject to be treated in a manner compatible with the dosage formulation, and in an amount that will be effective to bring about the desired adjuvant effect/immune response. The amount of the composition to be delivered which, in the case of nucleic acid molecules is generally in the range of from 0.001 to 1000 μg , more typically 0.01 to 10.0 μg of nucleic acid molecule per dose, and in the case of peptide or protein molecules is 1 μg to 5 mg, more typically 1 to 50 μg of peptide, depends on the subject to be treated. The exact amount necessary will vary depending on the age and general condition of the individual being immunized and the particular nucleotide sequence or peptide selected, as well as other factors. An appropriate effective amount can be readily determined by one of skill in the art upon reading the instant specification.

Particulate Compositions

Alternatively, polynucleotides carrying the selected ADP-ribosylating exotoxin subunit peptide coding sequences, as well as one or more selected antigen moiety, can be formulated as a particulate composition. More particularly, formulation of particles comprising one or more polynucleotide molecules can be carried out using standard pharmaceutical formulation chemistries and methodologies all of which are readily available to the reasonably skilled artisan. For example, one or more vector construct and/or antigen component can be combined with one or more pharmaceutically acceptable excipients or vehicles to provide a vaccine composition. Auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in the excipient or vehicle. These excipients, vehicles and auxiliary substances are generally pharmaceutical agents that do not themselves induce an immune response in the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, polyethyleneglycol, hyaluronic acid, glycerol and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids

such as acetates, propionates, malonates, benzoates, and the like. It is also preferred, although not required, that the nucleic acid composition will contain a pharmaceutically acceptable carrier that serves as a stabilizer, particularly for peptide, protein or other like antigens or ancillary materials. Examples of
5 suitable carriers that also act as stabilizers for peptides include, without limitation, pharmaceutical grades of dextrose, sucrose, lactose, trehalose, mannitol, sorbitol, inositol, dextran, and the like. Other suitable carriers include, again without limitation, starch, cellulose, sodium or calcium phosphates, citric acid, tartaric acid, glycine, high molecular weight polyethylene glycols (PEGs),
10 and combination thereof. A thorough discussion of pharmaceutically acceptable excipients, carriers, stabilizers and other auxiliary substances is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991), incorporated herein by reference.

The formulated compositions will include an amount of the toxin subunit
15 peptide-encoding polynucleotides which is sufficient to provide the desired adjuvant effect against a co-administered antigen, as defined above. An appropriate effective amount can be readily determined by one of skill in the art. Such an amount will fall in a relatively broad range, generally within the range of about 0.001 μ g to 25 mg or more of the nucleic acid construct of interest, and
20 specific suitable amounts can be determined through routine trials. The compositions may contain from about 0.1% to about 99.9% of the nucleic acid molecule(s). If an antigen component is included in the composition, or the methods are used to provide a particulate antigen composition, the antigen will be present in a suitable amount as described above. The compositions are then
25 prepared as particles using standard techniques, such as by simple evaporation (air drying), vacuum drying, spray drying, freeze drying (lyophilization), spray-freeze drying, spray coating, precipitation, supercritical fluid particle formation, and the like. If desired, the resultant particles can be densified using the techniques described in commonly owned International Publication No. WO
30 97/48485, incorporated herein by reference.

Single unit dosages or multidose containers, in which the particles may be

packaged prior to use, can comprise a hermetically sealed container enclosing a suitable amount of the particles comprising a suitable nucleic acid construct and/or a selected antigen (e.g., to provide a multicomponent vaccine composition). The particulate compositions can be packaged as a sterile
5 formulation, and the hermetically sealed container can thus be designed to preserve sterility of the formulation until use in the methods of the invention. If desired, the containers can be adapted for direct use in a particle delivery device. Such containers can take the form of capsules, foil pouches, sachets, cassettes, and the like. Appropriate particle delivery devices (e.g., needleless syringes) are
10 described herein.

The container in which the particles are packaged can further be labelled to identify the composition and provide relevant dosage information. In addition, the container can be labelled with a notice in the form prescribed by a governmental agency, for example the Food and Drug Administration, wherein
15 the notice indicates approval by the agency under Federal law of the manufacture, use or sale of the adjuvant, antigen (or vaccine composition) contained therein for human administration.

The particulate compositions can then be administered using a transdermal delivery technique. Preferably, the particulate compositions will be
20 delivered via a powder injection method, e.g., delivered from a needleless syringe system such as those described in commonly owned International Publication Nos. WO 94/24263, WO 96/04947, WO 96/12513, and WO 96/20022, all of which are incorporated herein by reference. Delivery of particles from such needleless syringe systems is typically practised with particles having an
25 approximate size generally ranging from 0.1 to 250 μm , preferably ranging from about 10-70 μm . Particles larger than about 250 μm can also be delivered from the devices, with the upper limitation being the point at which the size of the particles would cause untoward damage to the skin cells. The actual distance which the delivered particles will penetrate a target surface depends upon particle
30 size (e.g., the nominal particle diameter assuming a roughly spherical particle geometry), particle density, the initial velocity at which the particle impacts the

surface, and the density and kinematic viscosity of the targeted skin tissue. In this regard, optimal particle densities for use in needleless injection generally range between about 0.1 and 25 g/cm³, preferably between about 0.9 and 1.5 g/cm³, and injection velocities generally range between about 100 and 3,000 m/sec, or
5 greater. With appropriate gas pressure, particles having an average diameter of 10-70 μm can be accelerated through the nozzle at velocities approaching the supersonic speeds of a driving gas flow.

Compositions containing a therapeutically effective amount of the powdered molecules described herein can be delivered to any suitable target
10 tissue via the above-described particle delivery devices. For example, the compositions can be delivered to muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland and connective tissues. For nucleic acid molecules, delivery is preferably
15 to, and the molecules expressed in, terminally differentiated cells; however, the molecules can also be delivered to non-differentiated, or partially differentiated cells such as stem cells of blood and skin fibroblasts.

If desired, these needleless syringe systems can be provided in a preloaded condition containing a suitable dosage of the particles comprising the ADP-
20 ribosylating exotoxin subunit peptide coding sequences of interest. The loaded syringe can be packaged in a hermetically sealed container, which may further be labelled as described above.

Thus, the method can be used to obtain nucleic acid particles having a size ranging from about 10 to about 250 μm, preferably about 10 to about 150 μm,
25 and most preferably about 20 to about 60 μm; and a particle density ranging from about 0.1 to about 25 g/cm³, and a bulk density of about 0.5 to about 3.0 g/cm³, or greater.

Similarly, particles of selected antigens having a size ranging from about 0.1 to about 250 μm, preferably about 0.1 to about 150 μm, and most preferably
30 about 20 to about 60 μm; a particle density ranging from about 0.1 to about 25

g/cm³, and a bulk density of preferably about 0.5 to about 3.0 g/cm³, and most preferably about 0.8 to about 1.5 g/cm³ can be obtained.

Enhancing Immune Responses

5 In another embodiment of the invention, a method for enhancing an immune response against a co-administered antigen of interest is provided. In essence, the method entails (a) administering an antigen of interest to a subject, (b) providing an adjuvant composition according to the present invention, wherein the said composition contains one or more nucleic acid molecules
10 containing selected coding sequences for ADP-ribosylating exotoxin subunit peptides, and (c) co-administering the adjuvant composition to the subject, whereby the toxin subunit peptides are expressed from their respective coding sequences in an amount sufficient to elicit an enhanced immune response against the co-administered antigen. As described in detail herein above, the coding
15 sequences are operably linked to the same or different regulatory sequences to provide one or more expression cassettes. These expression cassettes are then provided in a suitable vector, for example a plasmid vector construct or a viral vector.

In one aspect, the method entails administering the polynucleotide
20 composition to a subject using standard gene delivery techniques that are known in the art. See, e.g., U.S. Patent Nos. 5,399,346, 5,580,859, 5,589,466. Typically, the polynucleotide vaccine composition is combined with a pharmaceutically acceptable excipient or vehicle to provide a liquid preparation (as described herein above) and then used as an injectable solution,
25 suspension or emulsion for administration via parenteral, subcutaneous, intradermal, intramuscular, intravenous injection using a conventional needle and syringe, or using a liquid jet injection system. It is preferred that the composition be administered to skin or mucosal tissue of the subject. Liquid preparations can also be administered topically to skin or mucosal tissue, or provided as a finely
30 divided spray suitable for respiratory or pulmonary administration. Other modes of administration include oral administration, suppositories, and active or passive

transdermal delivery techniques. The polynucleotide compositions can alternatively be delivered *ex vivo* to cells derived from the subject, whereafter the cells are reimplanted in the subject. Upon introduction into the subject, the nucleic acid sequence is expressed to provide the ADP-ribosylating exotoxin subunit peptides *in situ* in an amount sufficient to enhance an immune response against the co-administered antigen in the vaccinated subject. This enhanced immune response can be characterized as an enhanced humoral (antibody) response, an enhanced cellular (CTL) response, or be characterized as enhancing both humoral and cellular responses against the co-administered antigen.

In certain preferred embodiments, the enhanced immune response is characterized as an augmented Th1-like (cellular) immune response against the co-administered antigen, rather than a Th2-like response. In this regard, the augmented Th1-like immune response can be qualified by one or more of the following: increased titers of interferon- γ producing CD4⁺ helper T lymphocytes and/or CD8⁺ CTLs; increased antigen-specific CTL activity; of increased titers of antigen-specific antibodies of the subclass typically associated with cellular immunity (e.g., IgG2a).

It is preferred that the polynucleotide adjuvant compositions of the present invention be delivered in particulate form. For example, the compositions can be administered using a particle delivery device as described in detail herein above. In certain embodiments, the polynucleotide compositions can be coated onto core carrier particles using a variety of techniques known in the art and delivered using a particle-mediated delivery method. Carrier particles are selected from materials which have a suitable density in the range of particle sizes typically used for intracellular delivery from a particle delivery device. The optimum carrier particle size will, of course, depend on the diameter of the target cells.

These methods can alternatively be modified by co-administration of additional or ancillary components to the subject. For example, a secondary vaccine composition can be administered, wherein the secondary composition can comprise a nucleic acid vaccine, or the secondary vaccine composition can

comprise a conventional vaccine such as a whole virus, split virus, or subunit vaccine. The secondary vaccine composition can be combined with the polynucleotide ADP-ribosylating exotoxin subunit peptide compositions to form a single composition, or the secondary vaccine composition can be administered
5 separately to the same or to a different site, either concurrently, sequentially, or separated by a significant passage of time such as in a boosting step some days after the initial composition has been administered.

As above, the secondary vaccine composition and/or other ancillary component can be administered by injection using either a conventional syringe,
10 or using a particle-mediated delivery system as also described above. Administration will typically be either subcutaneously, epidermally, intradermally, intramucosally (e.g., nasally, rectally and/or vaginally), intraperitoneally, intravenously, orally or intramuscularly. Other modes of administration include topical, oral and pulmonary administration, suppositories,
15 and transdermal applications. Dosage treatment may be a single dose schedule or a multiple dose schedule.

Experimental

Below are examples of specific embodiments for carrying out the present
20 invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation
25 should, of course, be allowed for.

Example 1
Development of Expression Vectors Encoding
the A and B Subunits of CT.

5 Genomic DNA from *Vibrio cholerae* was used as a template for
polymerase chain reactions (PCR) to generate DNA fragments containing the
coding sequences for both the A and B subunits of cholera toxin (CT). For PCR
generation of the A subunit-encoding fragment (CTA), the following two
oligodeoxyribonucleotide primers were used:

10

Primer 1: 5'— GGA GCT AGC AAT GAT GAT AAG TTA TAT CGG —3'
(SEQ ID NO: 7); and

15

Primer 2: 5'— CCT GGA TCC TCA TAA TTC ATC CTT AAT TCT —3' (SEQ
ID NO: 8).

20

In order to facilitate insertion into an expression vector, Primers 1 and 2
contain extra sequences at their 5' ends (outside the region of homology to the
CTA coding sequence) which include recognition sites for *Nhe*I and *Bam*HI,
respectively. Primers 1 and 2 were designed to lead to PCR generation of a
fragment of the A subunit coding sequence starting at nucleotide position 164 and
ending at nucleotide position 886 (GenBank Accession #D30053). This region
encompasses the entire coding sequence for the mature subunit A peptide but
does not include the sequence encoding the bacterial signal peptide found at the
amino terminus of the pre-subunit A peptide.

25

For PCR generation of the B subunit-encoding fragment (CTB), the
following two oligodeoxyribonucleotide primers were used:

30

Primer 3: 5'— GGA GCT AGC ACA CCT CAA AAT ATT ACT GAT —3'
(SEQ ID NO: 9); and

Primer 4: 5'— CCT GGA TCC TTA ATT TGC CAT ACT AAT TGC —3' (SEQ ID NO: 10).

5 In order to facilitate insertion into an expression vector, Primers 3 and 4 contain extra sequences at their 5' ends (outside the region of homology to the CTB coding sequence) which include recognition sites for *NheI* and *BamHI*, respectively. Primers 3 and 4 were designed to lead to PCR generation of a fragment of the B subunit coding sequence starting at nucleotide position 946 and ending at nucleotide position 1257 (GenBank Accession #D30053). This region
10 encompasses the entire coding sequence for the mature subunit B peptide but does not include the sequence encoding the bacterial signal peptide found at the amino terminus of the pre-subunit B peptide.

In addition to the two PCR reactions described above, a third PCR reaction was performed to generate a coding sequence for a modified form of subunit A of CT in which the C-terminal four amino acid KDEL motif was
15 eliminated. This reaction involved the use of Primer 1 (SEQ ID NO: 7) and the following primer:

Primer 5: 5'— CCT GGA TCC TCA AAT TCT ATT ATG TGT ATC —3' (SEQ ID NO: 11).
20

All PCR reactions were performed using Pfu Turbo DNA polymerase (obtained from Strategene, La Jolla, CA) along with the PCR reaction buffer supplied by the manufacturer. PCR conditions were as follows: 95°C for 2 min., 30 cycles of
25 (95°C for 1 min., 55°C for 2 min. 15 sec., 72°C for 1 min.), 72°C for 5 min., and 4°C hold.

Following completion of the PCR reactions, the newly synthesized fragments were digested with *NheI* and *BamHI* enzymes to generate cohesive ends, and the individual fragments were inserted into the *NheI*- and *BamHI*-
30 cleaved pWRG7054 expression vector resulting in the following clones: pPJV2002, pPJV2003, and pPJV2006 encoding CTA, CTB, and modified CTA

(minus KDEL), respectively. The parental cloning vector pWRG7054 contains the human cytomegalovirus immediate early promoter with the associated intron A sequence. In addition, the coding sequence for the signal peptide of human tissue plasminogen activator is included in pWRG7054 to allow for the secretion from mammalian cells of any protein whose coding sequence is inserted at the *NheI* site in the appropriate reading frame. (See, e.g., Chapman et al. (1991) *Nuc. Acids Res.* 19:3979-3986, and Burke et al. (1986) *J. Biol. Chem.* 261:12574-12578.)

The restriction maps and full sequences of plasmids pPJV2002, pPJV2003, and pPJV2006 are depicted in Figures 1, 2, and 3, respectively.

Example 2

Development of Expression Vectors Encoding the A and B Subunits of *E. coli* Heat Labile Enterotoxin.

Genomic DNA from *E. coli* strain E078:H11 (American Type Culture Collection #35401) was used as a template for polymerase chain reactions to generate DNA fragments containing the coding sequences for both the A and B subunits of the heat labile enterotoxin (LT). For PCR generation of the A subunit-encoding fragment, the following two oligodeoxyribonucleotide primers were used:

Primer 6: 5'— GGA GCT AGC AAT GGC GAC AAA TTA TAC CGT —3' (SEQ ID NO: 12); and

Primer 7: 5'— CCT GGA TCC TCA TAA TTC ATC CCG AAT TCT —3' (SEQ ID NO: 13).

In order to facilitate insertion into an expression vector, Primers 6 and 7 contain extra sequences at their 5' ends (outside the region of homology to the LTA coding sequence) which include recognition sites for *NheI* and *BamHI*,

respectively. Primers 6 and 7 were designed to lead to PCR generation of a fragment of the A subunit coding sequence starting at nucleotide position 145 and ending at nucleotide position 867 of the sequence found in the GenBank database (Accession #AB011677). This region encompasses the entire coding sequence for the mature subunit A peptide but does not include the sequence encoding the bacterial signal peptide found at the amino terminus of the pre-subunit A peptide.

For PCR generation of the B subunit-encoding fragment, the following two oligodeoxyribonucleotide primers were used:

Primer 8: 5'— GGA GCT AGC GCT CCC CAG TCT ATT ACA GAA —3' (SEQ ID NO: 14); and

Primer 9: 5'— CCT GGA TCC CTA GTT TTC CAT ACT GAT TGC —3' (SEQ ID NO: 15).

In order to facilitate insertion into an expression vector, Primers 8 and 9 contain extra sequences at their 5' ends (outside the region of homology to the LTB coding sequence) which include recognition sites for *NheI* and *BamHI*, respectively. Primers 8 and 9 were designed to lead to PCR generation of a fragment of the B subunit coding sequence starting at nucleotide position 927 and ending at nucleotide position 1238 of the sequence found in the GenBank database (Accession #AB011677). This region encompasses the entire coding sequence for the mature subunit B peptide but does not include the sequence encoding the bacterial signal peptide found at the amino terminus of the pre-subunit B peptide.

In addition to the two PCR reactions described above, a third PCR reaction was performed to generate a coding sequence for a modified form of subunit A of LT in which the C-terminal four amino acid RDEL motif was eliminated. This reaction involved the use of Primer 6 (SEQ ID NO: 12) and the following primer:

Primer 10: 5'— CCT GGA TCC TCA AAT TCT GTT ATA TAT GTC —3'
(SEQ ID NO: 16).

5 All PCR reactions were performed using Pfu Turbo DNA polymerase (obtained from Strategene, La Jolla, CA) along with the PCR reaction buffer supplied by the manufacturer. PCR conditions were as follows: 95°C for 2 min., 30 cycles of (95°C for 1 min., 55°C for 2 min. 15 sec., 72°C for 1 min.), 72°C for 5 min., 4°C hold.

10 Following completion of the PCR reactions, the newly synthesized fragments were digested with *NheI* and *BamHI* to generate cohesive ends and the individual fragments were inserted into the *NheI*- and *BamHI*-cleaved pWRG7054 expression vector resulting in clones pPJV2004, pPJV2005, and pPJV2007 encoding LTA, LTB, and modified LTA (minus RDEL), respectively.

15 The restriction maps and complete sequences of plasmids pPJV2004, pPJV2005, and pPJV2007 are shown in Figures 4, 5 and 6, respectively.

Example 3

Enhancement of Antigen-Specific Antibody Response to a DNA Vaccine Using Plasmid Vectors Encoding CTA and/or CTB.

20

A DNA vaccine vector encoding the M2 protein of influenza A virus was employed to test the adjuvant effects of the pPJV2002, pPJV2003, and pPJV2006 adjuvant vectors in the context of particle-mediated DNA vaccination. Particle-mediated DNA vaccination was performed by precipitating the M2 DNA vaccine vector, with or without various combinations of the pPJV2002, pPJV2003 and pPJV2006 adjuvant vectors, onto microscopic gold particles and accelerating the coated gold particles into the epidermis of mice using a PowderJect® XR-1 particle delivery device (PowderJect Vaccines, Inc. Madison, WI).

25 More particularly, the sequence from the RNA segment #7 (that encodes the M2 protein) of influenza virus strain A/Kagoshima/10/95 (H3N2) was used as a model to design PCR primers to facilitate cloning of the mature M2 coding

sequence from A/Sydney/5/97 (H3N2). The A/Kagoshima sequence was used for primer design since the sequence of RNA segment 7 of A/Sydney has not yet been determined. The high degree of conservation among M2 sequences was expected to facilitate the use of primers designed from a different viral strain.

5 Since M2 is translated from a spliced RNA, it was deemed necessary that nucleotide positions 27 to 714 in the coding region of segment 7 RNA were spliced out. Accordingly, a set of PCR primers was generated and designed to generate the complete M2 coding sequence but ensure that the intron was cleanly eliminated from resulting M2 coding sequence clone. The PCR primers used to
10 generate the full-length M2 coding sequence clone were as follows:

Primer 11: 5'—CCC AAG CTT CCA CCA TGA GCC TTC TAA CCG AGG
TCG AAA CAC CTA TCA GAA ACG AAT GGG AGT GC—3' (SEQ ID NO:
17); and

15

Primer 12: 5'—CCC GGA TCC TTA CTC CAG CTC TAT GCT G—3' (SEQ ID
NO: 18).

Primer 11 (SEQ ID NO: 17) contains additional sequences at its 5' end
20 that include a recognition site for *Hind*III and a Kozak consensus sequence to facilitate mRNA translation initiation. Also, Primer 12 (SEQ ID NO:) contains additional sequences at its 5' end that includes a recognition sequence for *Bam*HI.

Viral RNA was isolated from a sample of A/Sydney/5/97 (H3N2) that was grown in embryonated chicken eggs. The viral RNA isolation process used
25 standard techniques known to those skilled in the art. RNA from this virus was used in a reverse transcriptase / polymerase chain reaction (RT-PCR) using an RT-PCR kit obtained from Stratagene (La Jolla, CA). The RT reaction step was completed by adding 5.9μl of RNase-free water to a reaction tube. To this tube was added 1.0μl 10X MMLV-RT buffer and 1.0μl dNTP mix from the kit. Also,
30 1μl of A/Sydney/5/97 RNA and 0.6μl (0.6μg) of Primer 11 (SEQ ID NO: 17) was added. The reaction was heated to 65°C for 5 minutes to denature the RNA,

after which 0.5 μ l of reverse transcriptase from the kit was added. The reaction was incubated at 37°C for 15 minutes to complete the reverse transcription step.

The PCR reaction step was completed by addition of the following components to a new reaction tube: 40 μ l water; 5 μ l 10X ultra HF buffer from the kit; 1.0 μ l dNTP mix from the kit; 1.0 μ l Primer 11 (SEQ ID NO: 17) (1.0 μ g); 1.0 μ l Primer 12 (SEQ ID NO: 18) (1.0 μ g); 1 μ l of the reverse transcriptase reaction mix from above; and 1 μ l Turbo PFU polymerase from the kit. The PCR reaction was carried out using the following incubation scheme: 1 minute @ 95°C; followed by 30 cycles of (30 sec @ 95°C, 30 sec @ 46°C, 3 min @ 68°C), followed by 10 minutes @ 68°C. PCR products were electrophoresed on a 2% agarose gel revealing a single DNA band of the expected size of approximately 300 bp.

The approximately 300 bp band was isolated from the gel and digested with *Hind*III and *Bam*HI in order to generate the necessary sticky ends for insertion into the pWRG7077 DNA vaccine expression vector (Schmaljohn et al. (1997) *J. Virol.* 71:9563-9569). The pWRG7077 DNA was digested partially with *Hind*III and completely with *Bam*HI to facilitate insertion of the M2 coding insert. The requirement for a partial *Hind*III digestion of the vector was due to the presence of a second *Hind*III site in the Kanamycin resistance marker of this plasmid. The resulting M2 DNA vaccine vector was termed pM2-FL. The pM2-FL vector contains the immediate early promoter from human cytomegalovirus (hCMV) and its associated intron A sequence to drive transcription from the M2 coding sequence. This vector also includes a polyadenylation sequence from the bovine growth hormone gene.

The pM2-FL plasmid was then precipitated onto 2 micron gold particles as single vector, or mixed vector plus adjuvant vector samples (i.e., the M2 plasmid vector was combined with pPJV2002, pPJV2003, and/or pPJV2006 adjuvant vectors). Specifically, plasmid DNA (single M2 vector or M2 vector plus one or more adjuvant vectors) was mixed with 2 micron gold particles (Degussa, Lot 65-0) in a small centrifuge tube containing 400 μ l of 50 mM spermidine. The DNA-to-gold ratio varied from 2.5 μ g DNA per mg of gold to

4.0 µg of DNA per mg of gold, and a single batch contained 26 mg of gold.

DNA was precipitated onto the gold particles by addition of a 1/10 volume of 10% CaCl₂ during continuous agitation of the tube on a rotary mixer. DNA-gold complexes were washed three times with absolute ethanol, and then injected into a TEFZEL® tube (McMaster-Carr) housed in a tube turner coating machine (PowderJect Vaccines, Inc., Madison WI) which coats the inside of the tube with the gold/DNA complex. This tube turner machine is described in US Patent No. 5,733,600. See also PCT patent application PCT/US95/00780 and US Patent Nos. 5,780,100; 5,865,796 and 5,584,807. After the coating procedure was completed, the tubes were cut into 0.5 inch “cartridges” suitable for loading into a particle delivery device.

The following DNA-gold formulations were generated for a mouse DNA vaccine adjuvant trial.

Formulation #1: pM2-FL DNA vector alone, 2.5 µg DNA per mg gold, 0.5 mg gold per cartridge;

Formulation #2: pM2-FL DNA vector precipitated onto one batch of gold (2 µg pM2-FL DNA per mg gold), pPJV2002 and pPJV2003 DNA vectors coprecipitated onto a second batch of gold (1.75 µg of each DNA adjuvant vector per mg gold), the gold batches were mixed equally, 0.5 mg gold per cartridge;

Formulation #3: pM2-FL DNA vector, pPJV2002 and pPJV2003 DNA vectors all coprecipitated onto single batch of gold (2 µg pM2-FL DNA per mg gold, and 1 µg each of pPJV2002 and pPJV2003 per mg gold), 0.5 mg gold per cartridge;

Formulation #4: pM2-FL DNA vector, pPJV2006 and pPJV2003 DNA vectors all coprecipitated onto single batch of gold (2 µg pM2-FL DNA per mg gold, and 1 µg each of pPJV2006 and pPJV2003 DNA per mg gold), 0.5 mg gold per cartridge;

Formulation #5: pM2-FL DNA vector and pPJV2002 coprecipitated onto single batch of gold (2 μ g pM2-FL DNA per mg gold, and 1 μ g of pPJV2002 DNA per mg gold), 0.5 mg gold per cartridge; and

- 5 Formulation #6: pM2-FL DNA vector and pPJV2003 coprecipitated onto single batch of gold (2 μ g pM2-FL DNA per mg gold, and 1 μ g of pPJV2003 DNA per mg gold), 0.5 mg gold per cartridge.

10 These DNA vaccine formulations were then administered to six groups of mice as follows. Each experimental group contained 7 animals and each animal received two immunizations with the respective formulation with a 4 week resting period between immunizations. Each immunization consisted of two tandem deliveries to the abdominal epidermis (one cartridge per delivery) using a PowderJect® XR-1 particle delivery device (PowderJect Vaccines Inc., Madison,
15 WI) at a helium pressure of 400 p.s.i.. Serum samples were collected 4 weeks following the primary immunization (just before the boost) and two weeks following the second or booster immunization.

Individual serum samples were assayed for M2-specific antibody responses using an ELISA assay in which 96-well plates were pre-coated with an
20 M2 synthetic peptide consisting of the following sequence:
SLLTEVETPIRNEWECR (SEQ ID NO: 19). ELISA plates were coated with the M2 peptide overnight at 4°C using the peptide in phosphate buffered saline (PBS) at a concentration of 1 μ g/ml. On the next day, the plates were blocked with 5% nonfat dry milk in PBS for 1 hour at room temperature. Plates were then washed
25 three times with wash buffer (10 mM Tris Buffered Saline, 0.1% Brij-35). Diluted serum samples were then added to the wells and the plates were incubated for 2 hours at room temperature. The plates were washed three times with wash buffer and 100 μ l of a secondary antibody was added and plates were incubated for 1 hour at room temperature. The secondary antibody consisted of a
30 goat anti-mouse IgG (H+L) biotin-labeled antibody (Southern Biotechnology) that was diluted 1:8000 in 1% BSA/PBS/0.1% Tween-20. Plates were then

washed three times, and a streptavidin-horse radish peroxidase conjugate (Southern Biotechnology) diluted to 1:8000 in PBS/0.1% Tween-20 was added and the plate incubated for 1 hour at room temperature. Following three additional washes, 100 μ l of TMB substrate (Bio Rad, Hercules, CA) was added and color development was allowed to proceed for 30 minutes at room temperature. Color development was stopped by the addition of 1N H₂SO₄ and the plates were read of 450 nm. Endpoint dilution titers were determined by identifying the highest dilution of serum that still yielded an absorbance value that was two times the background absorbance value obtained using a non immune control sample.

Endpoint antibody titers for individual animals in each group and geometric mean titers for each experimental group are shown in Table 1 below.

TABLE 1

Formulation #	Individual Titers	Geometric Mean Titer
1	24,300 24,300 72,900 218,700 218,700 218,700 218,700	99,781
2	72,900 72,900 218,700 218,700 218,700 218,700 656,100	186,934

Formulation #	Individual Titers	Geometric Mean Titer
3	24,300 72,900 72,900 218,700 656,100 656,100 656,100	186,934
4	24,300 218,700 656,100 656,100 656,100 656,100 656,100 656,100	350,211
5	----* 72,900 218,700 218,700 218,700 656,100 656,100	262,645
6	24,300 72,900 218,700 218,700 1,968,300 1,968,300 5,904,900	409,722

5

* (deceased)

As can be seen, all experimental groups immunized with a formulation containing one or more of the CT-encoding adjuvant vectors (Formulations #2-6) exhibited an increased geometric mean titer following the booster immunization relative to control animals immunized with the M2 vector (Formulation #1) alone.

10

Example 4Enhancement of Antigen-Specific Cellular Responses to a DNA Vaccine
Encoding the Hepatitis B Surface Antigen (HBsAg) Using
Adjuvant Plasmid Vectors Encoding the CT-A and CT-B Subunit Peptides.

5

A Hepatitis B surface antigen (HBsAg) vector plasmid was constructed as follows. To generate the HbsAg coding region, the pAM6 construct (obtained from the American Type Culture Collection "ATCC") was cut with *NcoI* and treated with mung bean nuclease to remove the start codon of the X-antigen. The resultant DNA was then cut with *BamHI* and treated with T4 DNA polymerase to blunt-end the DNA and create an HBsAg expression cassette. The HBsAg expression cassette is present in the 1.2 kB fragment. The plasmid construct pPJV7077 (Schmaljohn et al. (1997) *J. Virol.* 71:9563-9569) which contains the full-length human CMV (Towne strain) immediate early promoter (with enhancer) was cut with *HindIII* and *BglII*, and then treated with T4 DNA polymerase and calf-alkaline phosphatase to create blunt-ended DNA, and the HBsAg expression cassette was ligated into the plasmid to yield the pWRG7128 construct.

The pWRG7128 plasmid was precipitated onto gold particles following the procedure described in Example 3 above, again using 2 μ g DNA per mg gold. Adjuvant plasmid vectors expressing the CTA and CTB subunit genes (pPJV2002 and pPJV2003, respectively) were mixed together and precipitated onto gold particles with each plasmid being present at 1 μ g DNA per mg gold such that the total amount was also 2 μ g DNA per mg gold. The gold particles coated with pWRG7128 and those coated with the pPJV2002 and pPJV2003 were mixed 1:1 and then loaded into TEFZEL® tubing as above. For immunization, 0.5 inch lengths of tubing representing 1 μ g DNA (0.5 μ g pWRG7128 plasmid and 0.25 μ g of each of pPJV2002 and pPJV2003 plasmid) were delivered into the epidermis of Balb/c mice using the PowderJect® XR-1 particle delivery device using the same delivery conditions as described above in Example 3. For controls, mice were immunized with gold particles coated only

with pWRG7128 (1 μ g DNA/0.5 mg gold per delivery). Mice (4 per experimental group) were immunized and boosted at 4 weeks, then sacrificed at 2 weeks post-boost. Immune responses were evaluated for serum antibody levels by ELISA. In addition, cellular immune responses were measured using an
5 ELISPOT assay to quantify CD8-specific IFN- γ secretion.

Serum samples of individual mice were tested for antibodies specific for HBsAg using an ELISA assay. For the ELISA, Falcon Pro Bind microtiter plates were coated overnight at 4°C with purified HBsAg (BioDesign) at 0.1 μ g per well in PBS (phosphate buffered saline, BioWhittaker). The plates were blocked
10 for 1 hour at room temperature (RT) with 5% dry milk/PBS then washed 3 times with wash buffer (10 mM Tris Buffered saline, 0.1% Brij-35), and serum samples diluted in dilution buffer (2% dry milk/PBS/0.05 % Tween 20) were added to the plate and incubated for 2 hours at RT. The plates were washed 3 times and a biotinylated goat anti-mouse antibody (Southern Biotechnology) diluted 1:8000
15 in dilution buffer was added to the plate and incubated for 1 hr at RT. Following the incubation, plates were washed 3 times, after which a Streptavidin-Horseradish peroxidase conjugate (Southern Biotechnology) diluted 1:8000 in PBS was added and the plate incubated a further 1 hr at RT. After an additional three washes, Plates were washed 3 times, then a TMB substrate solution
20 (BioRad) was added and the reaction was stopped with 1N H₂SO₄ after 30 minutes. Optical density was read at 450 nm. Endpoint titers were calculated by comparison of the samples with a standard of known titer.

For the cellular immune assays, single cell suspensions of splenocytes from the spleens of the immunized animals were cultured *in vitro* in the presence
25 of a peptide corresponding to a known CD8 epitope in Balb/c mice. The peptide was dissolved in DMSO (10 mg/ml) and diluted to 10 μ g/ml in culture. The sequence of the peptide was IPQSLDSWWTSL (SEQ ID NO: 20).

For IFN- γ ELISPOT assays, Millipore Multiscreen membrane filtration plates were coated with 50 μ l of 15 μ g/ml anti-IFN- γ antiserum (Pharmingen) in
30 sterile 0.1 M carbonate buffer (pH 9.6) overnight at 4°C. Plates were washed 6 times with sterile PBS and then blocked with tissue culture medium containing

10% fetal bovine serum (FBS) for 1-2 hr at RT. The medium was removed and spleen cells dispensed into the wells with a total of 1×10^6 cells per well. For wells in which less than 1×10^6 cells from immunized animals was added, cells from naïve animals were used to bring the total to 1×10^6 . Cells were incubated
5 overnight in a tissue culture incubator in the presence of the peptide as described above. The plates were then washed 2 time with PBS and 1 time with distilled water. This was followed by 3 washes with PBS. A biotinylated anti IFN- γ monoclonal antibody (Pharmingen) was added to the plate (50 μ l of a 1 μ g/ml solution in PBS) and incubated for 2 hr at RT. The plates were washed 6 times
10 with PBS after which 50 μ l of a Streptavidin Alkaline phosphatase conjugate (1:1000 in PBS, Pharmingen) was added and incubated for 2 hr at RT. The plates were washed 6 times with PBS and an alkaline phosphatase color substrate (BioRad) was added and the reaction was allowed to proceed until dark spots appeared. The reaction was stopped by washing with water 3 times. Plates were
15 air dried and spots counted under a microscope.

For the IFN- γ ELISA assays, the cells were cultured overnight in round bottom 96 well tissue culture plates in the presence of the peptide. Samples of the supernatant were taken and used for the determination of IFN- γ levels. High binding plates (Costar) were coated with 100 μ l of 0.5 ug/ml of anti-mouse IFN- γ antibody (Pharmingen) in bicarbonate buffer pH 9.6. Plates were blocked for 1
20 hr at RT with tissue culture medium containing 10% FBS then washed 3 times with a TBS wash buffer. Supernatant samples obtained from cultured cells were diluted in tissue culture medium and loaded onto the plate and incubated for 2 hr at RT. Plates were washed 3 times with wash buffer and a secondary antibody (0.5 μ g/ml of biotinylated rat anti-mouse INF- γ in PBS, Pharmingen) was added
25 to the plates and incubated for 1 hr at RT. Plates were washed 3 times, and a Streptavidin-horseradish peroxidase conjugate (1:2000 in PBS, Southern Biotechnology) was added for 1 hr at RT. Plates were washed 3 times, then a TMB substrate solution was added (BioRad) and the reaction was stopped with
30 1N H₂SO₄. Optical density was read at 450 nm.

The results of the ELISA are reported below in Table 2.

TABLE 2

Formulation	Individual Titers	Average Titers
pWRG7128	40,000 84,000 41,000 54,000	55,000
pWRG7128, pPJV2002 and pPJV2003	28,000 14,000 27,000 23,000	23,000

As can be seen, antibody levels in control mice immunized with pWRG7128 alone were found to be higher than in those immunized with pWRG7128 in combination with the CT adjuvant vectors (pPJV2002 and pPJV2003). The average endpoint titer for the control animals was 55,000, while the average titer in the animals immunized with the adjuvanted formulation was 23,000. However, the group receiving the adjuvanted formulation actually received ½ the amount of pWRG7128 as the controls, which may account for the reduction in antibody titers.

The cellular immune responses measured in the two groups of mice indicate a significant enhancement of cellular responses by the CT adjuvant vectors. More particularly, the results of the CD8-specific IFN- γ ELISA assay are depicted below in Table 3.

TABLE 3

IFN- γ ELISA			
Formulation	Number of Cells/Well		
5	1.0 x 10 ⁶	0.5 x 10 ⁶	0.1 x 10 ⁶
	0.77	0.213	0.001
	0.828	0.121	0.027
	1.35	0.312	0.006
	1.25	0.323	0.007
	(Average)	1.05	0.242
10	pWRG7128,	1.96	1.19
	pPJV2002 and	2.30	1.70
	pPJV2003	2.20	1.83
		2.44	2.33
			0.898
	(Average)	2.23	1.76
			0.404

In this IFN- γ ELISA assay, the number of cells per well represents the number of cells recovered from immunized animals plated per well. Total number of cells per well is constant (e.g., 1 x 10⁶) and is supplemented with cells from naïve animals. Values are the OD readings measured at 450 nm. The higher OD values in the ELISA found for the cells from mice treated with the CT adjuvanted formulation is indicative of a greater amount of IFN- γ secreted by these cells in response to antigen. Cells from naïve mice did not yield a measurable OD value in the IFN- γ ELISA.

The results of the CD8-specific IFN- γ ELISPOT assay are depicted below in Table 4.

TABLE 4

IFN- γ ELISPOTs	
Formulation	Number of Positive Cells/ 1×10^6 Cells
5	pWRG7128
	510
	410
	530
	590
	(Average) 510
10	pWRG7128, pPJV2002 and pPJV2003
	1,480
	2,300
	2,500
	3,500
	(Average) 2,445

In this IFN- γ ELISPOT assay, the numbers are the average of duplicate wells and correspond to number of positive cells per 1×10^6 cells. Cells from naïve mice did not yield any spots in this ELISPOT assay. In addition, the greater number of ELISPOTs found in the animals treated with the CT adjuvanted formulation indicates a superior response as compared with the animals receiving the antigen (pWRG7128) alone.

The above data demonstrate that the novel adjuvant compositions of the present invention have a potent ability to enhance the cellular immune responses to a coadministered antigen, in this case a HBsAg expressed from a DNA vaccine.

Example 5Enhancement of Humoral and Cellular Immune Responses to HIV-1 gp120
Antigen Using Simultaneous Delivery of a gp120 Antigen Vector
and CTA/CTB Adjuvant Vectors.

5

A plasmid vector encoding HIV-1 gp120 was constructed as follows. The vector was constructed starting with a Bluescript (Stratagene, La Jolla, CA) plasmid backbone, the human cytomegalovirus (hCMV) immediate early promoter (Fuller et al. (1994) *Aids Res. Hum Retroviruses* 10:1433) and the SV40 virus late polyadenylation site. The hCMV promoter is contained within a 619 base pair (bp) *AccII* fragment extending 522 bp upstream and 96 bp downstream from the immediate early transcription initiation site. The SV40 virus late polyadenylation sequence is contained within an approximately 800 bp *BamHI-BglII* fragment derived from pSV2dhfr (formerly available from Bethesda Research Laboratories, catalogue #5369 SS). Initially, a plasmid encoding HIV-1 gp160, termed "pC-Env" was constructed. This plasmid contains a 2565 bp *KpnI-XhoI* fragment from LAV-1_{BRU} (ATCC Accession No. 53069, GenBank Accession No. K02013), which begins at the sequence encoding amino acid position #4 of the mature gp160 amino terminus. The *env* coding sequence fragment was placed immediately downstream of, and fused in frame with a 160 bp synthetic fragment encoding the herpes simplex virus glycoprotein D (gD) signal peptide and none amino acids of the mature gD amino terminus as previously described (Fuller et al. (1994) *Aids Res. Hum Retroviruses* 10:1433).

The plasmid encoding HIV-1 gp120, termed "pCIA-Env/T" herein, was then constructed as follows. The pCIA-Env/T plasmid encodes a truncated form of HIV-1 gp160, and is identical to the pC-Env construct except that the *env* coding sequences are truncated at the *HindIII* site at nucleotide position 8188. This results in a truncated gp160 translation

product with the truncation point lying 128 amino acid residues downstream of the gp120/gp41 processing site.

A second plasmid vector encoding HIV-1 rev, termed "pC-rev" herein, was constructed as follows. This vector contains three discontinuous regions of the LAV-1_{BRU} provirus (nucleotide positions 678-1085, 5821-6379, and 8188-8944) placed directly between the hCMV promoter and the SV40 virus late polyadenylation sequence as described above. The three discontinuous regions contain the major 5' splicing donor, the first exon of the *rev* gene, and the second exon of the *rev* gene, respectively.

A third plasmid vector construct termed "pWRG7054" was used as an empty vector control in the study. The pWRG7054 construct contains an SIV *nef* coding region, the hCMV promoter immediate early with the Intron A region, a TPA leader sequence and the bovine growth hormone polyadenylation sequence. Construction of the pWRG7054 plasmid is described herein below in Example 6.

The following DNA-gold formulations were generated for a mouse DNA vaccine adjuvant trial.

Formulation #1: Empty vector control (pWRG7054 without the gp120 insert), 2.5 µg DNA per mg gold, 0.5 mg gold per cartridge;

Formulation #2: Empty vector control (pWRG7054), the gp120 (pCIA-EnvT) DNA vector, and the rev (pC-rev) DNA vector (to allow for expression of the HIV-1 gp120 molecule), all coprecipitated onto a single batch of gold (1.25 µg of each of pWRG7054 DNA and pCIA-EnvT DNA per mg gold, and 0.125 µg of pC-rev per mg gold), 0.5 mg gold per cartridge;

Formulation #3: Empty vector control (pWRG7054), CTA (pPJV2002) and CTB (pPJV2003) DNA vectors all coprecipitated onto a single batch of gold (1.25 µg

pWRG7054 DNA per mg gold, and 1 μ g each of pPJV2002 and pPJV2003 DNA per mg gold), 0.5 mg gold per cartridge;

5 Formulation #4: gp120 (pCIA-EnvT) DNA vector, HIV-1 rev (pC-rev) DNA vector, CTA (pPJV2002) and CTB (pPJV2003) DNA vectors all coprecipitated onto a single batch of gold (1.25 μ g pCIA-EnvT DNA per mg gold, 0.125 μ g pC-rev DNA per mg gold, and 1 μ g each of pPJV2002 and pPJV2003 DNA per mg gold), 0.5 mg gold per cartridge; and

10 Formulation #5: gp120 (pCIA-EnvT) DNA vector, HIV-1 rev (pC-rev) DNA vector, CTA -KDEL (pPJV2006) and CTB (pPJV2003) DNA vectors all coprecipitated onto a single batch of gold (1.25 μ g pCIA-EnvT DNA per mg gold, 0.125 μ g pC-rev DNA per mg gold, and 1 μ g each of pPJV2006 and pPJV2003 DNA per mg gold), 0.5 mg gold per cartridge.

15

The pWRG7054, pCIA-EnvT, pC-rev, pPJV2002, pPJV2003 and pPJV2006 plasmid vectors were precipitated onto gold particles following the procedures described in Example 3 above, again using 2 μ g DNA per mg gold. The coated gold particles were loaded into TEFZEL® tubing, again using the
20 procedures described in Example 3 above. For immunization, two 0.5 inch lengths of tubing were used to deliver a total payload of 1.0 mg gold into the epidermis of 5-6 week old female Balb/c mice using a particle delivery device operated under the same delivery conditions as described above in Example 3 (400 p.s.i. helium).

25 Each of the five experimental groups (one per each DNA vaccine formulation) consisted of 4 animals, and each animal received primary and booster immunizations with their respective formulations, timed at 0 and 5 weeks. Each immunization consisted of two tandem deliveries to the abdominal epidermis (one cartridge per delivery) using a PowderJect® XR-1 particle
30 delivery device (PowderJect Vaccines Inc., Madison, WI).

Serum antibody responses to the HIV gp120 antigen were tested using an ELISA assay on specimens collected at week 5 and week 6.5 (post-prime and post-boost, respectively). For the ELISA, Costar high binding EIA plates were coated with 0.3 μ g/well of recombinant HIV gp120 (Intracel) in 50 μ l PBS by incubation overnight at 4°C. Plates were washed three times and blocked with 2% BSA in PBS for 2 hours at room temperature. Serial dilutions of serum were added to the coated plates, and incubated at 37°C for one hour. After washing, the plates were incubated with a 1:1500 dilution of alkaline phosphatase conjugated goat anti-mouse IgG (H+L) (BioRad), followed by color development with p-nitrophenylphosphate (PNPP) (BioRad) and OD reading @ 405nm.

The results of the ELISA are depicted in Figure 7. As can be seen, there was an approximate 20-fold augmentation of gp120-specific immune responses in the groups that received adjuvant vectors (Formulation #4 containing the pPJV2002 and pPJV2003 vectors, or Formulation #5 containing the pPJV2006 and pPJV2003 vectors) in combination with the gp120 vector as compared with the group that received the gp120 vector without adjuvant (Formulation #2, pCIA-EnvT).

Following collection of the post-boost serum samples, animals were sacrificed and spleens were collected from each mouse. Splenocytes were isolated by crushing the spleens, passing the cells through a 70 μ m cell strainer, and lysing the red blood cells with ACK lysis buffer (BioWhittaker). Splenocytes were washed 3 times with RPMI-5% FCS and resuspended to a concentration of 1×10^7 cells/ml in RPMI-10% FCS supplemented with antibiotics, sodium pyruvate, and non-essential amino acids.

The amount of antigen-specific IFN- γ secreted by the splenocytes was determined using an *in situ* ELISA. Costar high binding plates were coated with 10 μ g/ml of anti-mouse IFN-gamma capture monoclonal antibody (mAb) (Pharmingen, San Diego, CA) in 50 μ l 0.1M bicarbonate buffer (pH 9.6). After overnight incubation at 4°C, the wells were washed 5 times with PBS-0.05% Tween-20 and blocked with 200 μ l of complete R10 medium at room temperature for 2 hours. 1×10^6 splenocytes were added to each well and were

stimulated in medium alone (negative control), or in medium with 1 μ g/ml of a HIV gp120 peptide having the following sequence: RIQRGPGRAFVITGK (SEQ ID NO: 21). Following a 24 hour incubation at 37°C in 5% CO₂, the plates were washed 2 times with deionized (DI) water to lyse the cells, then washed 3 times
5 with PBS-0.05% Tween 20, and incubated for 1 hour at room temperature with 50 μ l per well of 1 μ g/ml biotinylated anti-mouse IFN- γ detecting mAb (Pharmingen). The plates were then washed 5 times and incubated for 1 hour with 50 μ l per well of a 1:8000 dilution of strepavidin-horseradish peroxidase (HRP) solution (Southern Biotechnology). The plates were again washed 5 times and
10 colorimetric development was accomplished by the addition of TMB substrate (BioRad, Hercules, CA) for 30 minutes at room temperature. The reaction was stopped by the addition of 1N sulfuric acid. Absorbance at 450nm was read with an optical plate reader.

The results of this study are depicted in Figure 8. This figure shows the
15 relative level of antigen-specific IFN- γ production in the 5 different vaccine test groups receiving the above-described formulations (Formulations #1-5). Importantly, the two immunization groups that received the gp120 vector (pCIA-EnvT) in combination with the CT adjuvant vectors (i.e., Formulation #4 containing the pPJV2002 and pPJV2003 vectors, and Formulation #5 containing
20 the pPJV2006 and pPJV2003 vectors) displayed significantly higher IFN- γ production levels than did the gp120 without adjuvant group (Formulation #2 containing the empty vector control and pCIA-EnvT), (P < 0.000001 and P = 0.0068, respectively). These data demonstrate the ability of the present CT vector adjuvant combinations to markedly augment antigen-specific cellular immunity to
25 an HIV antigen in an animal model.

In addition to an increase in the level of IFN- γ production, the number of IFN- γ secreting HIV-gp120 peptide-specific splenocytes was also increased markedly, as determined by an ELISPOT method. In the ELISPOT assay, nitrocellulose plates (Millipore) were coated with an IFN- γ capture antibody,
30 washed, and blocked as described above for the T helper cell IFN- γ *in situ* ELISA. Splenocytes were added to pre-coated wells at an input cell number of 1 x

10⁶ cells/well, and stimulated in medium alone (negative control) or in medium containing 1 µg/ml of a peptide containing the immunodominant HIV-gp120 CTL epitope and having the following sequence: RGPGRAFVTI (SEQ ID NO: 22). The plates were incubated for 24 hours at 37°C in 5% CO₂, washed 2 times with DI water, 3 times with PBS, and incubated for 1 hour at room temperature with 50 µl per well of 1 µg/ml biotinylated anti-mouse IFN-γ detecting mAb (Pharmingen). The plates were then washed 5 times and incubated for 1 hour with 50 µl per well of a 1:1000 dilution of strepavidin-alkaline phosphatase (ALP) solution (Mabtech). The plates were again washed 5 times and colorimetric development was accomplished by the addition of ALP membrane substrate (BioRad, Hercules, CA) until spots formed (2-30 minutes, room temperature). The reaction was stopped by washing with DI water, and the plates were air-dried overnight. Spots were enumerated under a 40X microscope. Only large spots with fuzzy borders were scored as spot forming cells (SFC).

The results from this ELISPOT assay are depicted in Figure 9 which shows the relative levels of IFN-γ-producing splenocytes in the 5 different vaccine test groups (receiving Formulations #1-5, respectively). Importantly, the two immunization groups that received the gp120 vector in combination with the CT adjuvant vectors (i.e., Formulation #4 containing the pPJV2002 and pPJV2003 vectors, and Formulation #5 containing the pPJV2006 and pPJV2003 vectors) displayed significantly higher numbers of IFN-γ-producing cells than did the gp120 without adjuvant group (Formulation #2 containing the empty vector control and pCIA-EnvT), (P < 0.000001 and P = 0.0032, respectively). These data demonstrate the ability of the present CT adjuvant combinations to markedly augment antigen-specific cellular immunity against a coadministered HIV gp120 antigen in an animal model. In addition, the enhanced IFN-γ production seen in these studies indicates that use of the CT adjuvant vector combinations of the present invention provides a robust Th1-like immune response in the immunized animals.

Example 6Enhancement of Antibody Responses to Hepatitis B Core and Surface Antigens Using Simultaneous Delivery of a Vector Encoding HBcAg and HBsAg with CTA/CTB Adjuvant Vectors.

5

A vector plasmid containing coding sequences for both the Hepatitis B core antigen (HBcAg) and Hepatitis B surface antigen (HBsAg) was constructed as follows. The HBcAg and HBsAg coding sequences were both obtained from the HBV clone pAM6 (ATCC Accession No. 45020). To
10 generate the HBsAg coding region, the pAM6 construct was cut with *NcoI* and treated with mung bean nuclease to remove the start codon of the X-antigen. The resultant DNA was then cut with *BamHI* and treated with T4 DNA polymerase to blunt-end the DNA and create an HBsAg expression cassette. The HBsAg expression cassette is present in the 1.2 kB fragment.
15 The plasmid construct pPJV7077 (Schmaljohn et al. (1997) *J. Virol.* 71:9563-9569) which contains the full-length human CMV (Towne strain) immediate early promoter (with enhancer) was cut with *HindIII* and *BglII*, and then treated with T4 DNA polymerase and calf-alkaline phosphatase to create blunt-ended DNA, and the HBsAg expression cassette was ligated into the
20 plasmid to yield the pWRG7128 construct.

To generate the HBcAg coding region, the pAM6 construct was cut to create an HBcAg expression cassette, after which the HBcAg sequence was truncated by site directed mutagenesis to remove the C-terminal arginine-rich region from the core antigen particle (which deletion does not interfere with
25 particle formation). The truncated HBcAg sequence was then cloned into a plasmid construct containing the human elongation factor promoter ("hELF", Mizushima et al. (1990) *Nucl. Acids Res.* 18:5322) to provide a HBcAg vector construct.

Expression cassettes containing: (a) the CMV promoter/enhancer, the
30 Intron A- 5' untranslated region, and the human tissue plasminogen activator (hTPA) signal peptide ("CMV-IA-TPA"); or (b) the bovine growth hormone

polyA sequence (bGHpA) were each obtained from the JW4303 vector construct (gift of Dr. Harriet Robinson, University of Massachusetts) and inserted into a plasmid backbone. The resultant construct was cut with *NheI*, filled with polymerase and then cut with *BamHI* to generate a vector fragment containing the pUC19 origin of replication, the ampicillin resistance gene and the bGHpA sequence. The plasmid backbone was cut a second time with *Sall*, filled with polymerase, and cut with *BamHI* to liberate a vector fragment containing the CMV-IA-TPA vector fragment. The two vector fragments were ligated together to yield a construct termed pWRG7054.

The pWRG7054 construct was cut with *NheI*, filled with polymerase, and cut with *BamHI* to produce a vector fragment. The HBcAg vector construct was cut with *NcoI*, filled with polymerase, and cut with *BamHI* to produce an insert fragment. The two fragments were then ligated together to yield a construct termed pWRG7063.

PEL-Bos was cut with *EcoRI* and dephosphorylated with calf intestinal phosphatase to produce a vector fragment. The pWRG7063 plasmid was cut with *HindIII*, filled with polymerase, and cut with *EcoRI* to produce an insert fragment containing the hTPA signal peptide, the HBcAg antigen sequence and the bGHpA region. These two fragments were ligated together to provide a construct termed pWRG7145.

The pWRG7128 construct was cut with *EcoRI* and dephosphorylated with calf intestinal phosphatase to produce a vector fragment containing the HBsAg coding region under transcriptional control of the hCMV promoter. The pWRG7145 construct was cut with *MfeI* and *EcoRI* to produce an insert fragment comprised of the hELF promoter/intron, the hTPA signal peptide sequence, the HBcAG antigen sequence and the bGHpA region. These fragments were then ligated together to provide the pPJV7193 plasmid construct containing the HBcAg and HBsAg coding sequences.

The following DNA-gold formulations were then generated for a pig DNA vaccine adjuvant trial.

Formulation #1: Control, the HBcAg/HBsAg vector (pWRG7193) alone, 2 μ g DNA per mg gold, 0.5 mg gold per cartridge; and

Formulation #2: The HBcAg/HBsAg vector (pWRG7193), CTA -KDEL
5 (pPJV2006) and CTB (pPJV2003) DNA vectors all coprecipitated onto single batch of gold (1.0 μ g pWRG7193 DNA per mg gold, and 0.5 μ g each of pPJV2006 and pPJV2003 DNA per mg gold), 0.5 mg gold per cartridge.

The pWRG7193 plasmid either alone or with the pPJV2006 and
10 pPJV2003 adjuvant vectors was precipitated onto gold particles following the procedures described in Example 3 above, again at a final concentration of 2 μ g DNA per mg gold. The coated gold particles were loaded into TEFZEL® tubing using the procedures described in Example 3 above. Two experimental groups (of 5 domestic pigs each) received two immunizations with Formulation #1 and
15 Formulation #2, respectively, wherein the two immunizations were spaced 6 weeks apart. Each immunization consisted of two tandem 500 p.s.i. shots to the groin area using a PowderJect® XR-1 particle delivery device in which each shot utilized a single cartridge. Thus, in the control group (Formulation #1), each immunization consisted of the delivery of a total of 1 mg gold and 2 μ g of the
20 DNA vaccine vector pWRG7193. Similarly, in the adjuvant test group (Formulation #2), each immunization consisted of a total delivery of 1 mg of gold, 2 μ g of the DNA vaccine vector pWRG7193, and 1 μ g each of the adjuvant vectors pPJV2006 and pPJV2003. Blood samples were collected from each animal at the time of the booster immunization (week 6) and 2 weeks following
25 the booster immunization (week 8).

Detection of antibody responses specific for the core antigen was performed as follows: ELISA plates were coated with the hepatitis B core antigen (Biodesign) at 100 ng/ml in PBS. After coating overnight at 4°C, plates were blocked with 5% nonfat dry milk in PBS for 1 hour at room temperature.
30 Plates were then washed 3 times with PBS containing 0.05% Tween-20. Swine serum samples were serially diluted in 2% nonfat dry milk/PBS/0.01% Tween-20

and added to the ELISA plates. After incubation at room temperature for 2 hours, plates were washed 3 times with PBS/0.05% Tween-20. The secondary antibody consisted of a goat anti-swine IgG conjugated to horse radish peroxidase (Kirkegaard and Perry) that was diluted 1:2000 in 2% nonfat dry milk/PBS/0.01% Tween-20, and added to the plates for a 1 hour incubation at room temperature. Plates were then washed 5 times with PBS/0.05% Tween-20 and 100 μ l of TMB substrate as added. Color development proceeded for 15 minutes and was stopped by the addition of 100 μ l of 1N H₂SO₄. Plates were read at 450 nm. Figures 10 and 11 show the geometric mean absorbance values at 6 and 8 weeks, respectively, for each of the two groups of pigs at 4 different serum dilutions. These data demonstrate marked enhancement of antibody titers to the hepatitis B core antigen following use of CT adjuvant vectors pPJV2006 and pPJV2003.

In addition to elevated humoral responses to the hepatitis B core antigen, a measurable elevation of antibody responses specific for the hepatitis B surface antigen was observed in the adjuvant test group as well. Surface antigen-specific antibodies were quantified in serum samples from individual animals using a commercial assay kit (AUSAB, Abbott Laboratories). This kit allows for the quantification of antibody responses in terms of milli-international unit (mIU/ml) using a standard panel. The geometric mean surface antigen antibody titers in the control group (Formulation #1) and adjuvant test group (Formulation #2) were 285 and 662 mIU/ml, respectively, demonstrating the ability of the present adjuvant plasmids (pPJV2006 and pPJV2003) to augment immune responses to an antigen encoded by a separate vector.

Example 7Enhancement of Cellular Th1-Like Immune Response to a DNAVaccine Using Plasmid Vectors Encoding CT or LT Subunits.

5 The pM2-FL DNA vaccine vector encoding the M2 protein of influenza A virus was employed to test the adjuvant effects of the pPJV2002, pPJV2003, pPJV2004, pPJV2005, pPJV2006 and pPJV2007 adjuvant vectors in the context of particle-mediated DNA vaccination. Particle-mediated DNA vaccination was performed by precipitating the M2 DNA vaccine vector, either with or without
10 various combinations of the adjuvant vectors, onto microscopic gold particles and accelerating the coated gold particles into the epidermis of mice using a PowderJect® XR-1 particle delivery device (PowderJect Vaccines, Inc. Madison, WI). Construction of the pM2-FL DNA plasmid vector is described herein above in Example 3.

15 As above, the pM2-FL plasmid was precipitated onto 2 micron gold particles as single vector, or mixed vector plus adjuvant vector samples. Specifically, plasmid DNA (single pM2-FL vector or pM2-FL vector plus one or more adjuvant vectors) was mixed with 2 micron gold particles (Degussa) in a small centrifuge tube containing spermidine. Precipitation was carried out
20 following the methodologies of Example 3, and the coated gold particles were then coated onto the inside surface of a TEFZEL® tube as also described in Example 3. The tubing was then cut into 0.5 inch cartridges suitable for loading into the particle delivery device.

25 The following DNA-gold formulations were generated for a mouse DNA vaccine adjuvant trial.

Formulation #1: pM2-FL DNA vector combined with the pWRG7054 empty plasmid vector prior to precipitation onto the same batch of gold, 2.1 µg total DNA per mg gold (0.1 µg pM2-FL and 2.0 µg pWRG7054), 0.5 mg gold per
30 cartridge;

Formulation #2: pM2-FL DNA vector combined with the CTA and CTB (pPJV2002 and pPJV2003) DNA vectors prior to precipitation onto a single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g of each DNA adjuvant vector per mg gold, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

5

Formulation #3: pM2-FL DNA vector combined with the CTA -KDEL and CTB (pPJV2006 and pPJV2003) DNA vectors all coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g of each DNA adjuvant vector per mg gold, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

10

Formulation #4: pM2-FL DNA vector combined with the CTA -KDEL (pPJV2006) DNA vector and supplemented with the pWRG7054 empty plasmid vector all combined and coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g pPJV2006, 1.0 μ g pWRG7054, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

15

Formulation #5: pM2-FL DNA vector combined with the CTA (pPJV2002) DNA vector and supplemented with the pWRG7054 empty plasmid vector, all DNAs combined and coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g pPJV2002, 1.0 μ g pWRG7054, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

20

Formulation #6: pM2-FL DNA vector combined with the CTB (pPJV2003) DNA vector and supplemented with the pWRG7054 empty plasmid vector, all DNAs combined and coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g pPJV2003, 1.0 μ g pWRG7054, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

25

Formulation #7: pM2-FL DNA vector combined with the LTA and LTB (pPJV2004 and pPJV2005) DNA vectors prior to precipitation onto a single batch

30

of gold, 2.1 μ g total DNA per mg gold (1.0 μ g of each DNA adjuvant vector per mg gold, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

5 Formulation #8: pM2-FL DNA vector combined with the LTA -RDEL and LTB (pPJV2007 and pPJV2005) DNA vectors all coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g of each DNA adjuvant vector per mg gold, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

10 Formulation #9: pM2-FL DNA vector combined with the LTA -RDEL (pPJV2007) DNA vector and supplemented with the pWRG7054 empty plasmid vector all combined and coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g pPJV2007, 1.0 μ g pWRG7054, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge;

15 Formulation #10: pM2-FL DNA vector combined with the LTA (pPJV2004) DNA vector and supplemented with the pWRG7054 empty plasmid vector, all DNAs combined and coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g pPJV2004, 1.0 μ g pWRG7054, 0.1 μ g pM2-FL), 0.5 mg gold per cartridge; and

20 Formulation #11: pM2-FL DNA vector combined with the LTB (pPJV2005) DNA vector and supplemented with the pWRG7054 empty plasmid vector, all DNAs combined and coprecipitated onto single batch of gold, 2.1 μ g total DNA per mg gold (1.0 μ g pPJV2005, 1.0 μ g pWRG7054, 0.1 μ g pM2-FL), 0.5 mg
25 gold per cartridge.

These DNA vaccine formulations were then administered to eleven groups of mice as follows. Each experimental group contained 8 animals and each animal received two immunizations with their respective formulation with a 4 week
30 resting period between immunizations. Each immunization consisted of two tandem deliveries to the abdominal epidermis (one cartridge per delivery) using a

PowderJect® XR-1 particle delivery device (PowderJect Vaccines Inc., Madison, WI) at a helium pressure of 400 p.s.i.. Serum samples were collected two weeks following the second or booster immunization.

Individual serum samples were assayed for M2-specific antibody responses for both IgG1 and IgG2a subclasses using the ELISA assay of Example 3 above for the determination of total IgG titer, except that a secondary antibody conjugate specific for either IgG1 or IgG2a was employed. The goat anti-mouse IgG1-biotin conjugated antibody was obtained from Southern Biotechnology Associates, Inc. (catalogue #1070-08, 0.5 mg/ml concentration) and used at a 1/8000 dilution. The goat anti-mouse IgG2a-biotin conjugated antibody was obtained from the same source (catalogue #1080-08, 0.5 mg/ml concentration) and was also used at a 1/8000 dilution.

Geometric mean antibody titers for M2 antigen-specific IgG1 and IgG2a were determined for each experimental group, and the IgG1-to-IgG2a ratios were calculated. These data are reported below in Table 5.

TABLE 5

Formulation #	IgG1-to-IgG2a Ratio
1	160.99
2	3.44
3	13.59
4	63.20
5	2.63
6	30.97
7	0.02
8	0.50
9	9.00
10	4.53
11	27.00

As can be seen with reference to Table 5, addition of either of the A or B subunits, or the various combinations of A and B subunits to the M2 DNA vaccine formulation resulted in significant reductions in the IgG1-to-IgG2a ratio otherwise elicited with the M2 DNA vaccine (pM2-FL) in the absence of adjuvant (Formulation #1). The greatest ratio reduction resulted from use of the LTA plus LTB, and LTA -RDEL plus LTB vector combinations (Formulations #7 and #8, respectively). In both of these formulations, use of the polynucleotide adjuvants of the present invention resulted in an abundance of M2 antigen-specific IgG2a titers over IgG1, which characteristic is a hallmark of a Th1-like immune response in mice.

The results from experimental groups receiving Formulations #1, #2 and #7 were plotted as the log IgG1-to-IgG2a ratio in Figure 12. As can be seen in that figure, adjuvanting the pM2-FL DNA vaccine composition with the CTA plus CTB adjuvant vectors (Formulation #2) caused a 2 order of magnitude drop in the IgG1-to-IgG2a ratio relative to the pM2-FL vaccine composition without adjuvant (Formulation #1). In addition, a further 2 order of magnitude drop in this ratio was observed when the pM2-FL DNA vaccine was adjuvanted with the LTA plus LTB adjuvant vectors (Formulation #7).

Example 8

Addition of Signal Peptide Coding Sequences to Adjuvant Vectors Encoding CT or LT.

The vector constructs containing the CTA, CTB, LTA and LTB toxin subunits (pPJV2002, pPJV2003, pPJV2004, and pPJV2005, respectively) were modified to remove the tpa signal peptide coding sequences, and a dual DNA vaccine vector encoding both the Hepatitis B surface and core antigens was constructed for use in the following experiments.

The standard PCR conditions that were used for the construction/modification of the vectors were as follows: 1x PCR core buffer

with 1.5 mM MgCl₂ (Promega Corp., Madison, WI); 0.400 μM of each primer; 200 μM of each dNTP (USB Inc., Cleveland, OH); 2.5 μg Taq polymerase (Promega Corp., Madison, WI); 1.0 ng template DNA; water to 100 μl; and a mineral oil (Aldrich Chemical Inc., Milwaukee WI) overlay. A PTC-200 thermocycler (MJ Research Inc., Waltham, MA) was programmed to run the following routine: 4 minutes @ 95°C; 30 cycles of (1 minute @ 95°C/ 1 minute 15 seconds @ 55°C/ 1 minute @ 72°C); 10 minutes @ 72°C; 4°C hold. The amplification products were removed from the PCR reaction using the QLAquick® PCR Purification Kit (Qiagen Inc., Valencia CA) prior to cutting with restriction enzymes (New England Biolabs, Beverly, MA). All PCR products were sequenced after cloning to ensure fidelity of the amplification.

More specifically, a dual DNA vaccine vector encoding both the Hepatitis B surface and core antigens was constructed as follows. The pWRG7128 construct (Example 4) which contains the surface antigen coding sequence was modified using a series of standard molecular biology techniques to provide the dual (surface/core antigen) construct. Initially, the pWRG7128 construct was modified to remove the bovine growth hormone polyadenylation region and replace the same with the rabbit beta-globin polyadenylation region. A first insertion fragment containing the CMV promoter with exon 1 and exon 2 sequences, and a second insertion fragment containing a second rabbit beta-globin polyadenylation region were ligated into the modified pWRG7128 construct, and an adaptor constructed by annealing syntenic oligonucleotides was inserted between *Sph*I and *Pst*I sites located immediately upstream of the inserted CMV promoter.

Plasmid mpSmpCC (GlaxoSmithKline, UK) was PCR'd with the following primers: 5'—GCC GCT AGC ATG GAC ATT GAC CCT TAT AAA GA—3' (SEQ ID NO:23) and 5'—CCA GGA TCC TTA ACA TTG AGA TTC C—3' (SEQ ID NO:24) to generate a Hepatitis B-adw2 core antigen coding sequence. This PCR product was cut with *Nhe*I and *Bam*HI to generate an insertion fragment, which fragment was then modified to include a downstream *Bgl*II site and inserted into a cloning vector. The cloning vector was cut with *Pst*I

and *EcoR1* to generate a core antigen insertion fragment; the modified pWRG7128 plasmid was cut with *Pst1* and *Mfe1* to generate a vector fragment, and the core antigen insertion fragment was ligated into the vector fragment, resulting in the dual surface/core antigen plasmid construct.

5 The vector constructs containing the CTA, CTB, LTA and LTB toxin subunits (pPJV2002, pPJV2003, pPJV2004, and pPJV2005, respectively) were modified to remove the tpa signal peptide coding sequences by merely excising the tpa coding sequences using restriction enzymes to produce the following constructs: CTA w/o TPA, CTB w/o TPA, LTA w/o TPA, and LTB w/o TPA,
10 respectively.

 The dual surface/core antigen plasmid was combined with irrelevant plasmid DNA (for non-adjuvanted control) or with the toxin subunit vector constructs and precipitated onto 2 micron gold particles. Specifically, plasmid DNA (dual surface/core antigen plasmid vector) plus two adjuvant vectors (to
15 provide a CTA/CTB or LTA/LTB combination) was mixed with 2 micron gold particles (Degussa) in a small centrifuge tube containing spermidine. Precipitation was carried out following the methodologies of Example 3, and the coated gold particles were then coated onto the inside surface of a TEFZEL® tube as also described in Example 3. The tubing was then cut into 0.5 inch
20 cartridges suitable for loading into the particle delivery device.

 The following DNA-gold formulations were thus generated for a mouse DNA vaccine adjuvant trial:

Formulation #1: ("no adjuvant") 1 µg dual surface/core antigen DNA plasmid
25 vector, 1 µg irrelevant DNA plasmid vector;

Formulation #2: ("CT") 1 µg dual surface/core antigen DNA plasmid vector, 0.5 µg pPJV2002 (CTA) DNA plasmid vector, 0.5 µg pPJV2003 (CTB) DNA
30 plasmid vector;

Formulation #3: ("CT w/o TPA") 1 μ g dual surface/core antigen DNA plasmid vector, 0.5 μ g CTA w/o TPA DNA plasmid vector, 0.5 μ g CTB w/o TPA DNA plasmid vector;

5 Formulation #4: ("CTA") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g pPJV2002 (CTA) DNA plasmid vector;

Formulation #5: ("CTA w/o TPA") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g CTA w/o TPA DNA plasmid vector;

10

Formulation #6: ("CTB") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g pPJV2003 (CTB) DNA plasmid vector;

15

Formulation #7: ("CTB w/o TPA") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g CTB w/o TPA DNA plasmid vector;

Formulation #8: ("LT") 1 μ g dual surface/core antigen DNA plasmid vector, 0.5 μ g pPJV2004 (LTA) DNA plasmid vector, 0.5 μ g pPJV2005 (LTB) DNA plasmid vector;

20

Formulation #9: ("LT w/o TPA") 1 μ g dual surface/core antigen DNA plasmid vector, 0.5 μ g LTA w/o TPA DNA plasmid vector, 0.5 μ g LTB w/o TPA DNA plasmid vector;

25

Formulation #10: ("LTA") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g pPJV2004 (LTA) DNA plasmid vector;

Formulation #11: ("LTA w/o TPA") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g LTA w/o TPA DNA plasmid vector;

30

Formulation #12: ("LTB") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g pPJV2005 (LTB) DNA plasmid vector; and

5 Formulation #13: ("LTB w/o TPA") 1 μ g dual surface/core antigen DNA plasmid vector, 1 μ g LTB w/o TPA DNA plasmid vector.

In a first study, various of the above-described DNA vaccine formulations were administered to five groups of mice using the PowderJect® XR-1 particle delivery device (PowderJect Vaccines Inc., Madison, WI). Each experimental group contained 5 animals, and each animal received two immunizations with the respective formulation with a four week resting period between immunizations. The Formulations that were tested were as follows: Formulation #1 (no adjuvant); Formulation #2 (CT); Formulation #3 (CT w/o TPA); Formulation #8 (LT); and Formulation #9 (LT w/o TPA). All animals were sacrificed two weeks after the second immunization, and the spleens harvested for use in IFN- γ and IL-4 ELISPOT assays. For the cellular immune assays, single cell suspensions of splenocytes from the spleens of the immunized animals were cultured *in vitro* in the presence of a peptide corresponding to a known T cell epitope (from either the surface or the core antigen) in the mice. The peptide was dissolved in DMSO (10 mg/ml) and diluted to 10 μ g/ml in culture.

For the ELISPOT assays, Millipore Multiscreen membrane filtration plates were coated with 50 μ l of the appropriate antiserum (15 μ g/ml anti-IFN- γ or -IL-4 antiserum, Pharmingen) in sterile 0.1 M carbonate buffer (pH 9.6) overnight at 4°C. Plates were washed 6 times with sterile PBS and then blocked with tissue culture medium containing 10% fetal bovine serum (FBS) for 1-2 hr at RT. The medium was removed and spleen cells dispensed into the wells with a total of 1×10^6 cells per well. For wells in which less than 1×10^6 cells from immunized animals was added, cells from naïve animals were used to bring the total to 1×10^6 . Cells were incubated overnight in a tissue culture incubator in the presence of the peptide as described above. The plates were then washed 2 times with PBS and 1 time with distilled water. This was followed by 3 washes with

PBS. Biotinylated anti-IFN- γ or anti-IL-4 monoclonal antibody (Pharmingen) was added to the plate (50 μ l of a 1 μ g/ml solution in PBS) and incubated for 2 hr at RT. The plates were washed 6 times with PBS after which 50 μ l of a Streptavidin Alkaline phosphatase conjugate (1:1000 in PBS, Pharmingen) was added and incubated for 2 hr at RT. The plates were washed 6 times with PBS and an alkaline phosphatase color substrate (BioRad) was added and the reaction was allowed to proceed until dark spots appeared. The reaction was stopped by washing with water 3 times. Plates were air dried and spots counted under a microscope.

The adjuvant effect of the secreted or non-secreted toxin subunit coding sequences was assessed by determining IFN- γ and IL-4 ELISPOT responses to both the Hepatitis B surface and core antigens ("sAg" and "cAg") encoded by the dual surface/core antigen construct, and these results were compared as reported in Figures 13A-13D. As can be seen, the secreted (signal sequence containing) CT and LT adjuvant vectors (Formulations #2 and #8, respectively) produced significant increases ($P \leq 0.05$) in IFN- γ responses to both the surface and core antigens, and in the IL-4 response to surface antigen (see Figures 13A, 13B and 13C). With respect to the IL-4 responses to the core antigen, the lack of adjuvant effect by the LT vectors (Formulations #8 and #9) is consistent with observations that the LT toxin is more of a Th1 adjuvant than the CT toxin. Most importantly, the CT and LT vectors lacking the signal sequences (Formulations #3 and #9, respectively) exhibited weaker adjuvant effect, particularly as seen in the IFN- γ ELISPOT data for both the surface and core antigens (see Figures 13A and 13C), where there was a statistically significant drop in adjuvant activity by virtue of deletion of the signal sequences.

Finally, the clearly observable difference in adjuvant effect between the secreted (Formulations #2 and #8) and the non-secreted (Formulations #3 and #9) helps establish that the observed adjuvant effects are not due to CpG motifs within the adjuvant vectors since the signal-containing and non signal-containing vectors do not have any difference in bacterial DNA (CpG) content yet exhibit

significant differences in their ability to augment surface antigen-specific IFN- γ responses.

In a second study, the above-described DNA vaccine formulations were administered to eight groups of mice using the PowderJect® XR-1 particle
5 delivery device (PowderJect Vaccines Inc., Madison, WI). Each experimental group contained 5 animals, and each animal received two immunizations with the respective formulation with a four week resting period between immunizations. The Formulations that were tested were as follows: Formulation #1 (no adjuvant); Formulation #2 (CT); Formulation #4 (CTA); Formulation #5 (CTA w/o TPA);
10 Formulation #6 (CTB); Formulation #7 (CTB w/o TPA); Formulation #8 (LT); Formulation #10 (LTA); Formulation #11 (LTA w/o TPA); Formulation #12 (LTB); and Formulation #13 (LTB w/o TPA). All animals were sacrificed two weeks after the second immunization, and the spleens harvested for use in IFN- γ and IL-4 ELISPOT assays described herein above.

15 As a result of this second study (data not shown), it was again observed that there was no discernable adjuvant effect that could be attributed to CpG content in the various adjuvant plasmids. Although for the most part no statistically relevant adjuvant effect was observed with the various toxin subunit adjuvant vectors, the LT subunit vectors (Formulations #10-13) did show
20 adjuvant effect in the IFN- γ and IL-4 response to surface antigen (sAg) that was influenced by the presence/absence of the secretion signal sequence.

Example 9

25 Adjuvant Plasmid Vectors Encoding CTA/CTB or LTA/LTB Subunit Peptides in a Viral Challenge Study

In order to assess the ability of the adjuvant plasmid vectors of the present invention to provide for protective effect in a Herpes Simplex Virus type 2 (HSV-
30 2) viral challenge model, the following study was carried out. A DNA vaccine encoding an HSV-2 antigen was constructed and then combined with various

combinations of the present adjuvant plasmid vectors to provide vaccine compositions. After immunization, the immunized animals were challenged with HSV-2 virus, and the protective effect of the various vaccine compositions was determined.

5 With respect to the construction of the DNA antigen plasmid, standard PCR techniques were used to construct the plasmid. The standard PCR conditions that were used for the construction of the vector were as follows: 1x PCR core buffer with 1.5 mM MgCl₂ (Promega Corp., Madison, WI); 0.400 μM of each primer; 200 μM of each dNTP (USB Inc., Cleveland, OH); 2.5 μg Taq
10 polymerase (Promega Corp., Madison, WI); 1.0 ng template DNA; water to 100 μl; and a mineral oil (Aldrich Chemical Inc., Milwaukee WI) overlay. A PTC-200 thermocycler (MJ Research Inc., Waltham, MA) was programmed to run the following routine: 4 minutes @ 95°C; 30 cycles of (1 minute @ 95°C/ 1 minute 15 seconds @ 55°C/ 1 minute @ 72°C); 10 minutes @ 72°C; 4°C hold. The
15 amplification products were removed from the PCR reaction using the QIAquick® PCR Purification Kit (Qiagen Inc., Valencia CA) prior to cutting with restriction enzymes (New England Biolabs, Beverly, MA). All PCR products were sequenced after cloning to ensure fidelity of the amplification.

 More specifically, a DNA vaccine plasmid vector encoding the HSV-2
20 early ICP27 antigen was constructed as follows. HSV is a double-stranded DNA virus having a genome of about 150-160 kbp. The viral genome is packaged within an icosahedral nucleocapsid which is enveloped in a membrane. The membrane (or envelope) includes at least 10 virus-encoded glycoproteins, the most abundant of which are gB, gC, gD, and gE. The viral genome also encodes
25 over 70 other proteins, including a group of approximately five ICP antigens. These early proteins are synthesized early in the viral replication cycle, in contrast to the envelope glycoproteins which are only made late in the life cycle of the virus. For a review of the molecular structure and organization of HSV, see, for example, Roizman and Sears (1996) "Herpes simplex viruses and their
30 replication" in Fields Virology, 3rd ed., Fields et al. eds., Lippincott-Raven Publishers, Philadelphia, PA. The HSV-2 ICP27 antigen can be readily obtained

from the HSV-2 genome, for example the genomic region spanning from approximately nucleotide 114589 to 134980 of the HSV-2 genome, or an *EcoRI* fragment that spans nucleotides 110931 to 139697 of the HSV-2 genome. The sequence of the HSV-2 genome is available from published sources, for example the sequence deposited with GenBank under Accession Number NC_001798.

In order to construct the ICP27 vector used in the present study, the ICP27 coding region was PCR'd from the HSV-2 genome using the following primers: 5'—GCC ACT CTC TTC CGA CAC—3' (SEQ ID NO:25) and 5'—CAA GAA CAT CAC ACG GAA C—3' (SEQ ID NO:26) to obtain a nucleotide fragment containing nucleotide sequences 114523-116179 (GenBank) of HSV-2 which correspond to the ICP27 coding region. The ICP27 fragment was then cloned into the multiple cloning region of the pTarget vector (Promega Corp., Madison, WI).

The adjuvant plasmid vector constructs containing the CTA, CTB, LTA and LTB toxin subunits (pPJV2002, pPJV2003, pPJV2004, and pPJV2005, respectively) were combined to provide CTA/CTB (pPJV2002 + pPJV2003) and LTA/LTB (pPJV2004 + pPJV2005) adjuvant. The ICP27 antigen plasmid was combined with the toxin subunit vector construct pairs and precipitated onto 2 micron gold particles. Specifically, plasmid DNA (ICP27 antigen plasmid vector) plus two adjuvant vectors (to provide a CTA/CTB or LTA/LTB combination) was mixed with 2 micron gold particles (Degussa) in a small centrifuge tube containing spermidine. Precipitation was carried out following the methodologies of Example 3, and the coated gold particles were then coated onto the inside surface of a TEFZEL® tube as also described in Example 3. The tubing was then cut into 0.5 inch cartridges suitable for loading into the particle delivery device.

The following DNA-gold formulations were thus generated for the HSV-2 challenge study:

Formulation #1: (no adjuvant) 2 µg ICP27 antigen DNA plasmid vector;

Formulation #2: ("High CT") 900 ng ICP27 antigen DNA plasmid vector, 50 ng of pPJV2002 (CTA), 50 ng of pPJV2003 (CTB);

5 Formulation #3: ("Low CT") 500 ng ICP27 antigen DNA plasmid vector, 250 ng of pPJV2002 (CTA), 250 ng of pPJV2003 (CTB);

Formulation #4: ("High LT") 900 ng ICP27 antigen DNA plasmid vector, 50 ng of pPJV2004 (LTA), 50 ng of pPJV2005 (LTB); and

10 Formulation #5: ("Low LT") 500 ng ICP27 antigen DNA plasmid vector, 250 ng of pPJV2004 (LTA), 250 ng of pPJV2005 (LTB).

In the study, the above-described DNA vaccine formulations were administered to five different groups of mice using the PowderJect® XR-1 particle delivery device (PowderJect Vaccines Inc., Madison, WI). Each experimental group contained 12 animals, and each animal received two immunizations (single shot applied to the abdomen) with the respective formulation with a four week resting period between immunizations. A sixth group of mice was established as a negative (naive) control, and did not receive any vaccinations. 4 mice from each group were sacrificed 2 weeks after the second immunization and used for IFN- γ ELISPOT assays (data not shown).

Two weeks post second immunization, all remaining mice (8/group) were challenged with 1×10^6 PFU of HSV-2 virus, strain MS, via intra-nasal instillation. The survival graph depicting the results of the challenge study is depicted in Figure 14. As can be seen, 100% of the naive animals succumbed within 4 days post challenge. The naive animals are depicted on the graph by the (●) curve. In addition, 100% of the animals receiving the ICP27 antigen plasmid vector alone (Formulation #1) died within 7 days post challenge. The animals receiving Formulation #1 are depicted on the graph by the (▼) curve. In marked contrast, the 25% (2/8) of the animals receiving the ICP27 plasmid adjuvanted with the low dose CT (Formulation #3) were protected from the viral challenge,

and 38% (3/8) of the animals receiving the ICP27 adjuvanted with the high dose CT (Formulation #2) were protected from the viral challenge. The animals receiving Formulation #3 are depicted on the graph by the (■) curve. The animals receiving Formulation #2 are depicted on the graph by the (◆) curve.

5 Finally, both the low dose LT-adjuvanted (Formulation #5) and the high dose LT-adjuvanted (Formulation #4) ICP27 vaccine provided complete (100%) protection in the immunized animals. The animals receiving Formulation #5 are depicted on the graph by the (▲) curve. The animals receiving Formulation #4 are depicted on the graph by the (○) curve.

10

Accordingly, novel polynucleotide adjuvant molecules, compositions comprising those adjuvant molecules, and conventional and nucleic acid immunization techniques have been described. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit and the scope of the invention as defined by the appended claims.

15

SEQUENCE LISTING

<110> HAYNES, Joel R.
ARRINGTON, Joshua

<120> NUCLEIC ACID ADJUVANTS

<130> APF41.40

<140>

<141> 2001-11-26

<160> 22

<170> PatentIn Ver. 2.1

<210> 1

<211> 5500

<212> DNA

<213> pPJV2002 plasmid

<400> 1

```

gacgaaaggg cctcgtgata cgcctatttt tatagggttaa tgtcatgata ataatgggttt 60
cttagacgtc aggtggcact tttcggggaa atgtgcgcgg aacccctatt tgtttatttt 120
tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat 180
aatattgaaa aaggaagagt atgagtattc aacattttccg tgtcgcctt attccctttt 240
ttgcggcatt ttgccttcct gtttttgctc acccagaaac gctgggtgaaa gtaaaagatg 300
ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac agcggtaaga 360
tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactttt aaagttctgc 420
tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggg cgcgcatac 480
actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg 540
gcatgacagt aagagaatta tgcagtgctg ccataaccat gagtgataac actgcggcca 600
acttacttct gacaacgatc ggaggaccga aggagctaac cgcttttttg cacaacatgg 660
gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacg 720
acgagcgtga caccacgatg cctgtagcaa tggcaacaac gttgcgcaaa ctattaactg 780
gcgaactact tactctagct tcccggcaac aattaataga ctggatggag gcggataaag 840
ttgcaggacc acttctgcgc tcggcccttc cggctggctg gtttatgtg gataaatctg 900
gagccggtga gcgtgggtct cgcggtatca ttgcagcact ggggccagat ggtaagccct 960
cccgtatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa cgaaatagac 1020
agatcgctga gatagggtgcc tcaactgatta agcattggta actgtcagac caagtttact 1080
catatatact ttagattgat ttaaaacttc attttttaatt taaaaggatc taggtgaaga 1140
tccttttttg taatctcatg accaaaatcc cttaacgtga gttttcgttc cactgagcgt 1200
cagaccccggt agaaaagatc aaaggatctt cttgagatcc tttttttctg cgcgtaatct 1260
gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgg ttgtttgccg gatcaagagc 1320
taccaactct ttttccgaag gtaactggct tcagcagagc gcagatacca aatactgtcc 1380
ttctagtgtg gccgtagtta ggccaccact tcaagaactc tgtagcaccg cctacatacc 1440
tcgctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg 1500
ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga acgggggggt 1560
cgtgcacaca gcccagcttg gagcgaacga cctacaccga actgagatac ctacagcgtg 1620
agcattgaga aagcgccacg cttcccgaag ggagaaaggc ggacaggtat ccggtaaagc 1680
gcagggtcgg aacaggagag cgcacgaggg agcttccagg gggaaacgcc tggatatctt 1740
atagtcctgt cgggtttcgc cacctctgac ttgagcgtcg atttttgtga tgctcgtcag 1800
gggggcggag cctatggaaa aacgccagca acgcggcctt ttacagggtt ctggcctttt 1860

```

gctggccttt	tgctcacatg	ttcttttctg	cgttatcccc	tgattctgtg	gataaccgta	1920
ttaccgcctt	tgagtgagct	gataccgctc	gccgcagccg	aacgaccgag	cgcagcgagt	1980
cagtgagcga	ggaagcggaa	gagcgcccaa	tacgcaaacc	gcctctcccc	gcgcgttggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccgaactg	gaaagcgggc	agtgagcgca	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcacccca	ggctttacac	tttatgcttc	2160
cggctcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220
accatgatta	cgccaagcta	gtcgacataa	atcaatatgt	gctattggcc	attgcatacg	2280
ttgtatctat	atcataatat	gtacattttat	attggctcat	gtccaatatg	accgccatgt	2340
tgacattgat	tattgactag	ttattaatag	taatcaatta	cgggggtcatt	agttcatagc	2400
ccatatatgg	agttccgcgt	tacataactt	acggtaaattg	gcccgcctcg	tgaccgcca	2460
acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	2520
ctttccattg	acgtcaatgg	gtggagttat	tacggtaaac	tgcccacttg	gcagtacatc	2580
aagtgtatca	tatgccaagt	ccggccccct	attgacgtca	atgacggtaa	atggccccgc	2640
tggcattatg	cccagtacat	gaccttacgg	gactttccta	cttggcagta	catctacgta	2700
ttagtcatcg	ctattaccat	ggatgatgcg	ttttggcagt	acaccaatgg	gcgtggatag	2760
cggtttgact	cacggggatt	tccaagtctc	caccccattg	acgtcaatgg	gagtttgttt	2820
tggcaccaaa	atcaacggga	ctttccaaaa	tgctcgtaata	accccgcctc	gttgacgcaa	2880
atggggcggta	ggcgtgtacg	gtgggaggtc	tatataagca	gagctcgttt	agtgaaccgt	2940
cagatcgctt	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	3000
tccagcctcc	gcggccggga	acgggtgcatt	ggaacgcgga	ttccccgtgc	caagagtgc	3060
gtaagtaccg	cctatagact	ctataggcac	acccctttgg	ctcttatgca	tgctatactg	3120
tttttggctt	ggggcctata	cacccccgct	ccttatgcta	taggtgatgg	tatagcttag	3180
cctatagggtg	tgggttattg	accattattg	accactcccc	tattgggtgac	gatactttcc	3240
attactaatc	cataacatgg	ctcttttgcca	caactatctc	tattggctat	atgccaatac	3300
tctgtccttc	agagactgac	acggactctg	tattttttaca	ggatggggtc	ccattttatta	3360
tttacaattt	cacatatata	acaacgccgt	cccccggtgc	cgcagttttt	attaaacata	3420
gcgtgggatc	tccacgcgaa	tctcgggtac	gtgttccgga	catgggctct	tctccggtag	3480
cggcggagct	tccacatccg	agccctgggtc	ccatgcctcc	agcggctcat	ggtcgctcgg	3540
cagctccttg	ctcctaacag	tggaggccag	acttaggcac	agcacaatgc	ccaccaccac	3600
cagtgtgccg	cacaaggccg	tggcggtagg	gtatgtgtct	gaaaatgagc	tcggagattg	3660
ggctcgcacc	gtgacgcaga	tggaagactt	aaggcagcgg	cagaagaaga	tgcaggcagc	3720
tgagttgttg	tattctgata	agagtcagag	gtaactcccc	ttgcgggtgct	gttaacgggtg	3780
gagggcagtg	tagtctgagc	agtactcgtt	gctgccgcgc	gcgccaccag	acataatagc	3840
tgacagacta	acagactgtt	cctttccatg	ggtcttttct	gcagtcaccg	tccaagcttg	3900
caatcatgga	tgcaatgaag	agagggctct	gctgtgtgct	gctgctgtgt	ggagcagttc	3960
tcgttttcggc	tagcaatgat	gataagttat	atcgggcaga	ttctagacct	cctgatgaaa	4020
taaagcagtc	aggtgggtctt	atgccaagag	gacagagtga	gtactttgac	cgagggtactc	4080
aaatgaatat	caacctttat	gatcatgcaa	gaggaactca	gacgggattt	gttaggcacg	4140
atgatggata	tgtttccacc	tcaattagtt	tgagaagtgc	ccacttagtg	ggtcaaacta	4200
tattgtcttg	tcattctact	tattatatat	atgttatagc	cactgcaccc	aacatgttta	4260
acgttaatat	tgtattaggg	gcatacagtc	ctcatccaga	tgaacaagaa	gtttctgctt	4320
taggtgggat	tccatactcc	caaataatat	gatggtatcg	agttcathtt	gggggtgcttg	4380
atgaacaatt	acatcgtaat	aggggctaca	gagatagata	ttacagtaac	ttagatattg	4440
ctccagcagc	agatggttat	ggattggcag	gtttccctcc	ggagcataga	gcttggaggg	4500
aagagccgtg	gattcatcat	gcacegccgg	gttgtgggaa	tgctccaaga	tcacgatga	4560
gtaataactt	cgatgaaaaa	acccaaagtc	taggtgtaaa	attccttgac	gaataaccaat	4620
ctaaagttaa	aagacaaata	ttttcaggct	atcaatctga	tattgatata	cataatagaa	4680
ttaaggatga	attatgagga	tcctcgcaat	ccctaggagg	attaggcaag	ggcttgagct	4740
cacgctcttg	tgagggacag	aaatacaatc	aggggcagta	tatgaatact	ccatggagaa	4800
accagatctt	acgtatgatc	agcctcgact	gtgccttcta	gttgccagcc	atctgttgtt	4860
tgccccctcc	ccgtgccttc	cttgaccctg	gaagggtgcca	ctcccactgt	cctttcctaa	4920
taaaatgagg	aaattgcac	gcattgtctg	agtaggtgtc	attctattct	gggggggtggg	4980

WO 03/004055

PCT/US01/43151

gtggggcagg	acagcaaggg	ggaggattgg	gaagacaata	gcaggcatgc	tgggggatgcg	5040
gtgggctcta	tggcttctga	ggcggaaaga	accagctggg	gctcgacagc	tcgactctag	5100
aattcactgg	ccgtcgtttt	acaacgtcgt	gactgggaaa	accctggcgt	tacccaactt	5160
aatcgccctg	cagcacatcc	ccctttcgcc	agctggcgta	atagcgaaga	ggcccgccacc	5220
gatcgccctt	cccaacagtt	gcgcagcctg	aatggcgaat	ggcgcctgat	gcggtatttt	5280
ctccttacgc	atctgtgcgg	tatttcacac	cgcataatgg	gcactctcag	tacaatctgc	5340
tctgatgccg	catagttaag	ccagccccga	caccgcgcaa	caccgcgtga	cgcgccctga	5400
cgggcttgtc	tgctcccggc	atccgccttac	agacaagctg	tgaccgtctc	cgggagctgc	5460
atgtgtcaga	ggttttcacc	gtcatcaccg	aaacgcgcga			5500

<210> 2

<211> 5089

<212> DNA

<213> pPJV2003 plasmid

<400> 2

gacgaaaggg	cctcgtgata	cgccatatttt	tatagggttaa	tgtcatgata	ataatgggttt	60
cttagacgtc	aggtggcact	tttcgggggaa	atgtgcgcgg	aacccttatt	tgttttatttt	120
tctaaataca	ttcaaatatg	tatccgctca	tgagacaata	accctgataa	atgcttcaat	180
aatattgaaa	aaggaagagt	atgagtattc	aacattttccg	tgtcgccctt	attccctttt	240
ttgcggcatt	ttgccttcct	gttttttgctc	acccagaaac	gctggtgaaa	gtaaaagatg	300
ctgaagatca	gttgggtgca	cgagtgggtt	acatcgaact	ggatctcaac	agcggtaaga	360
tccttgagag	ttttcgcccc	gaagaacgtt	ttccaatgat	gagcactttt	aaagtctctgc	420
tatgtggcgc	ggtattatcc	cgtattgacg	ccgggcaaga	gcaactcggg	cgccgcatac	480
actattctca	gaatgacttg	gttgagtact	caccagtcac	agaaaagcat	cttacggatg	540
gcatgacagt	aagagaatta	tgcagtgtctg	ccataaccat	gagtataaac	actgcggcca	600
acttacttct	gacaacgac	ggaggaccga	aggagctaac	cgcttttttg	cacaacatgg	660
gggatcatgt	aactcgcctt	gatcgttggg	aaccggagct	gaatgaagcc	ataccaaacg	720
acgagcgtga	caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaaa	ctattaactg	780
gcgaactact	tactctagct	tcccggcaac	aattaataga	ctggatggag	gcggataaag	840
ttgcaggacc	acttctgcgc	tcggcccttc	cggctggctg	gtttatttgc	gataaatctg	900
gagccggtga	gcgtgggtct	cgcggtatca	ttgcagcact	ggggccagat	ggtaagccct	960
cccgtatcgt	agttatctac	acgacgggga	gtcaggcaac	tatggatgaa	cgaaatagac	1020
agatcgctga	gatagggtgcc	tcactgatta	agcattggta	actgtcagac	caagtttact	1080
catatatact	ttagattgat	ttaaaacttc	attttttaatt	taaaaggatc	taggtgaaga	1140
tccttttttga	taatctcatg	accaaaaatcc	cttaacgtga	gttttcgttc	cactgagcgt	1200
cagaccccg	agaaaagatc	aaaggatctt	cttgagatcc	tttttttctg	cgcgtaatct	1260
gctgcttgca	aacaaaaaaa	ccaccgctac	cagcgggtgg	ttgttttgccg	gatcaagagc	1320
taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	aatactgtcc	1380
ttctagtgtg	gccgtagtta	ggccaccact	tcaagaactc	tgtagcaccg	cctacatacc	1440
tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	tgtcttaccg	1500
ggttggactc	aagacgatag	ttaccggata	aggcgcagcg	gtcgggctga	acgggggggt	1560
cgtgcacaca	gcccagcttg	gagcgaacga	cctacaccga	actgagatac	ctacagcgtg	1620
agcattgaga	aagcgccacg	cttcccgaag	ggagaaaggc	ggacagggtat	ccggtaagcg	1680
gcagggtcgg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	tggtatcttt	1740
atagtcctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	attttttgta	tgctcgtcag	1800
gggggcggag	cctatggaaa	aacgccagca	acgcggcctt	tttacgggttc	ctggcctttt	1860
gctggccttt	tgctcacatg	ttcttttctg	cgttatcccc	tgattcttgtg	gataaccgta	1920
ttaccgcctt	tgagttagct	gataccgctc	gccgcagccg	aacgaccgag	cgcgagcagt	1980
cagttagcga	ggaagcggaa	gagcgcccaa	tacgcaaacc	gcctctcccc	gcgcgttggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccgaactg	gaaagcgggc	agttagcgcga	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcacccca	ggcttttacac	tttatgcttc	2160
cggctcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220

accatgatta	cgccaagcta	gtcgacataa	atcaatattg	gctattggcc	attgcatacg	2280
ttgtatctat	atcataatat	gtacatttat	attggctcat	gtccaatatg	accgccatgt	2340
tgacattgat	tattgactag	ttattaatag	taatcaatta	cggggtcatt	agttcatagc	2400
ccatatatgg	agttccgcgt	tacataactt	acggtaaagt	gcccgcctcg	tgaccgcca	2460
acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	2520
ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaag	tgcccacttg	gcagtacatc	2580
aagtgtatca	tatgccaagt	ccggccccct	attgacgtca	atgacggtaa	atggccccgc	2640
tggcattatg	cccagtacat	gaccttacgg	gactttccta	cttggcagta	catctacgta	2700
ttagtcatcg	ctattacat	ggatgatgcg	ttttggcagt	acaccaatgg	gcgtggatag	2760
cggtttgact	cacggggatt	tccaagtctc	caccccatgt	acgtcaatgg	gagtttggtt	2820
tggcaccaaa	atcaacggga	ctttccaaaa	tgtcgtaata	accccgcccc	gttgacgcaa	2880
atgggcggta	ggcgtgtacg	gtgggagggt	tatataagca	gagctcgttt	agtgaaccgt	2940
cagatcgctt	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	3000
tccagcctcc	gcggccggga	acgggtgcatt	ggaacgcgga	ttccccgtgc	caagagtgc	3060
gtaagtaccg	cctatagact	ctataggcac	acccctttgg	ctcttatgca	tgctatactg	3120
ttttttggctt	ggggcctata	cacccccgct	ccttatgcta	taggtgatgg	tatagcttag	3180
cctatagggtg	tgggttattg	accattattg	accactcccc	tattgggtgac	gatactttcc	3240
attactaatc	cataacatgg	ctcttttgcca	caactatctc	tattggctat	atgccaatat	3300
tctgtccttc	agagactgac	acggactctg	tattttttaca	ggatgggggtc	ccattttatta	3360
tttacaattt	cacatatata	acaacgccgt	cccccggtgc	cgcagttttt	attaaacata	3420
gcgtgggatc	tccacgcgaa	tctcgggtac	gtgttccgga	catgggctct	tctccggtag	3480
cggcggagct	tccacatccg	agccctgggtc	ccatgcctcc	agcggctcat	ggtcgctcgg	3540
cagctccttg	ctcctaacag	tggaggccag	acttaggcac	agcacaatgc	ccaccaccac	3600
cagtgtgccg	cacaaggccg	tggcggtagg	gtatgtgtct	gaaaatgagc	tcggagattg	3660
ggctcgcacc	gtgacgcaga	tggaagactt	aaggcagcgg	cagaagaaga	tgcaggcagc	3720
tgagtgtgtg	tattctgata	agagtccagag	gtaactcccc	ttgcgggtgct	gttaacgggtg	3780
gagggcagtg	tagtctgagc	agtactcgtt	gctgccgcgc	gcgccaccag	acataatagc	3840
tgacagacta	acagactggt	cctttccatg	ggtcctttct	gcagtcaccg	tccaagcttg	3900
caatcatgga	tgcaatgaag	agagggctct	gctgtgtgtg	gctgctgtgt	ggagcagctt	3960
tcgttttcggc	tagcacacct	caaaatatta	ctgattttgtg	tgcagaatac	cacaacacac	4020
aaatatatac	gctaaatgat	aagatatattt	cgtatacaga	atctctagct	ggaaaaagag	4080
agatggctat	cattactttt	aagaatgggtg	caatttttca	agtagaagta	ccaggtagtc	4140
aacatataga	ttcacaaaaa	aaagcgattg	aaaggatgaa	ggataccctg	aggattgcat	4200
atcttactga	agctaaagtc	gaaaagttat	gtgtatggaa	taataaaaacg	cctcatgcga	4260
ttgccgcaat	tagtatggca	aattaaggat	cctcgcgaat	cctaggagga	ttaggcaagg	4320
gcttgagctc	acgctcttgt	gagggacaga	aatacaatca	ggggcagtat	atgaatactc	4380
catggagaaa	cccagatcta	cgtatgatca	gcctcgcactg	tgcccttctag	ttgccagcca	4440
tctgttggtt	gcccctcccc	cgtgccttcc	ttgaccttgg	aagggtgccac	tcccactgtc	4500
ctttccctaat	aaaatgagga	aattgcatcg	cattgtctga	gtaggtgtca	ttctattctg	4560
gggggtgggg	tggggcagga	cagcaagggg	gaggattggg	aagacaatag	caggcatgct	4620
ggggatgcgg	tgggctctat	ggcttctgag	gcggaaagaa	ccagctgggg	ctcgacagct	4680
cgactctaga	attcactggc	cgtcgtttta	caacgtcgtg	actgggaaaa	ccctggcggt	4740
acccaactta	atcgcccttc	agcacatccc	ccttttcgcca	gctggcgtaa	tagcgaagag	4800
gcccgcaccg	atcgcccttc	ccaacagttg	cgcagcctga	atggcgaatg	gcgcctgatg	4860
cggtatatttc	tccttacgca	tctgtgcggt	atttcacacc	gcataatgggtg	cactctcagt	4920
acaatctgct	ctgatgccgc	atagttaagc	cagccccgac	acccgccaac	acccgctgac	4980
gcgccctgac	gggcttgtct	gctcccggca	tccgcttaca	gacaagctgt	gaccgtctcc	5040
gggagctgca	tgtgtcagag	gttttccaccg	tcatacccca	aacgcgcga		5089

<210> 3
 <211> 5488
 <212> DNA
 <213> plasmid pPJV2006

<400> 3

gacgaaaggg	cctcgtgata	cgcctatttt	tatagggttaa	tgtcatgata	ataatggttt	60
cttagacgtc	aggtggcact	tttcgggggaa	atgtgcgcgg	aacccctatt	tgtttatttt	120
tctaaataca	ttcaaatatg	tatccgctca	tgagacaata	accctgataa	atgcttcaat	180
aatattgaaa	aaggaagagt	atgagtattc	aacattttccg	tgtcgccctt	attccctttt	240
ttgcggcatt	ttgccttcct	gttttttgctc	acccagaaac	gctgggtgaaa	gtaaaagatg	300
ctgaagatca	gttgggtgca	cgagtgggtt	acatcgaact	ggatctcaac	agcggtaaga	360
tccttgagag	ttttcgcccc	gaagaacgtt	ttccaatgat	gagcactttt	aaagttctgc	420
tatgtggcgc	ggtattatcc	cgtattgacg	ccgggcaaga	gcaactcggg	cgccgcatac	480
actattctca	gaatgacttg	gttgagtact	caccagtcac	agaaaagcat	cttacggatg	540
gcatgacagt	aagagaatta	tgcagtgctg	ccataaccat	gagtataaac	actgcggcca	600
acttacttct	gacaacgata	ggaggaccga	aggagctaac	cgcttttttg	cacaacatgg	660
gggatcatgt	aactcgcctt	gacgttggtg	aaccggagct	gaatgaagcc	ataccaaacg	720
acgagcgtga	caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaaa	ctattaactg	780
gcgaactact	tactctagct	tcccggcaac	aattaataga	ctggatggag	gcggataaag	840
ttgcaggacc	acttctgcgc	tccggccttc	cggctggctg	gtttattgct	gataaatctg	900
gagccggtga	gcgtgggtct	cgcggtatca	ttgcagcact	ggggccagat	ggtaagccct	960
cccgatatcgt	agttatctac	acgacgggga	gtcaggcaac	tatggatgaa	cgaaatagac	1020
agatcgctga	gatagggtgc	tcactgatta	agcattggta	actgtcagac	caagtttact	1080
catatatact	ttagattgat	ttaaaacttc	attttttaatt	taaaaggatc	taggtgaaga	1140
tccttttttga	taatctcatg	acccaaaatcc	cttaacgtga	gttttcgctc	cactgagcgt	1200
cagaccccggt	agaaaagatc	aaaggatcct	cttgagatcc	ttttttttctg	cgcgtaatct	1260
gctgcttgca	aacaaaaaaa	ccaccgctac	cagcgggtgg	ttgtttgccc	gatcaagagc	1320
taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	aatactgtcc	1380
ttctagtgtg	gccgtagtta	ggccaccact	tcaagaactc	tgtagcaccg	cctacatacc	1440
tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	tgtcttaccg	1500
ggttggactc	aagacgatag	ttaccggata	aggcgcagcg	gtcgggctga	acgggggggt	1560
cgtgcacaca	gcccagcttg	gagcgaacga	cctacaccga	actgagatac	ctacagcgtg	1620
agcattgaga	aagcgccacg	cttcccgaag	ggagaaaggc	ggacagggtat	ccggtaagcg	1680
gcagggctcg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	tggatatctt	1740
atagtcctgt	cgggttttcgc	cacctctgac	ttgagcgtcg	attttttgta	tgctcgtcag	1800
gggggpcggag	cctatggaaa	aacgcccagca	acgcggcctt	tttacggctc	ctggcctttt	1860
gctggccttt	tgctcacatg	ttcttttcctg	cgttatcccc	tgattctgtg	gataaccgta	1920
ttaccgcctt	tgagtgagct	gataccgctc	gccgcagccg	aacgaccgag	cgcagcgagt	1980
cagtgagcga	ggaagcggaa	gagcgcctca	tacgcaaacc	gcctctcccc	gcgcgttggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccgaactg	gaaagcgggc	agtgagcgca	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcacccca	ggcttttacac	tttatgcttc	2160
cggctcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220
accatgatta	cgccaagcta	gtcgacataa	atcaatattg	gctattggcc	attgcatacg	2280
ttgtatctat	atcataatat	gtacatttat	attggctcat	gtccaatatg	accgccatgt	2340
tgacattgat	tattgactag	ttathtaata	taatcaatta	cgggggtcatt	agttcatagc	2400
ccatatatgg	agttccgcgt	tacataactt	acggtaaattg	gcccgcctcg	tgaccgcca	2460
acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	2520
ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaacc	tgcccacttg	gcagtacatc	2580
aagtgtatca	tatgccaaag	ccggccccct	attgacgtca	atgacggtaa	atggccccgc	2640
tggcattatg	cccagtacat	gaccttacgg	gactttccta	cttggcagta	catctacgta	2700
ttagtcatcg	ctattacat	ggtgatgcgg	ttttggcagt	acaccaatgg	gcgtggatag	2760
cggtttgact	cacgggggatt	tccaagtctc	caccccatgg	acgtcaatgg	gagtttggtt	2820
tggcaccaaa	atcaacggga	ctttccaaaa	tgtcgtaata	accccgcccc	gttgacgcaa	2880
atgggpcggta	ggcgtgtacg	gtgggaggtc	tatataagca	gagctcgttt	agtgaaccgt	2940
cagatcgctt	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	3000
tccagcctcc	gcggccggga	acggtgcatt	ggaacgcgga	ttccccgtgc	caagagtgc	3060

```

gtaagtaccg cctatagact ctataggcac acccctttgg ctcttatgca tgctatactg 3120
tttttggcctt ggggcctata cccccccgct ccttatgcta taggtgatgg tatagcttag 3180
cctatagggtg tgggttattg accattattg accactcccc tattgggtgac gatactttcc 3240
attactaatc cataacatgg ctcttttgcca caactatctc tattggctat atgccaatac 3300
tctgtccttc agagactgac acggactctg tattttttaca ggatggggtc ccatttatta 3360
tttaciaaatt cacatataca acaacgccgt cccccgtgcc cgcagttttt attaaacata 3420
gcgtgggac tccacgcgaa tctcgggtac gtgttccgga catgggctct tctccggtag 3480
cggcggagct tccacatccg agccctggtc ccatgcctcc agcggctcat ggtcgcctcg 3540
cagctccttg ctcttaacag tggaggccag acttaggcac agcacaatgc ccaccaccac 3600
cagtgtgccg cacaaggccg tggcggtagg gtatgtgtct gaaaatgagc tcggagattg 3660
ggctcgcacc gtgacgcaga tgggaagactt aaggcagcgg cagaagaaga tgcaggcagc 3720
tgagttgttg tattctgata agagtcagag gtaactcccg ttgcgggtgct gttaacgggtg 3780
gagggcagtg tagtctgagc agtactcgtt gctgccgcgc gcgccaccag acataatagc 3840
tgacagacta acagactgtt cctttccatg ggtcttttct gcagtcaccg tccaagcttg 3900
caatcatgga tgcaatgaag agagggctct gctgtgtgct gctgctgtgt ggagcagtct 3960
tcgtttcggc tagcaatgat gataagttat atcgggcaga ttctagacct cctgatgaaa 4020
taaagcagtc aggtggtctt atgccaagag gacagagtga gtactttgac cgaggctactc 4080
aatgaatat caacctttat gatcatgcaa gaggaactca gacgggattt gttaggcacg 4140
atgatggata tgtttccacc tcaattagtt tgagaagtgc ccacttagtg ggtcaaacta 4200
tattgtctgg tcattctact tattatatat atgttatagc cactgcacc aacatgttta 4260
acgttaatga tgtattaggg gcatacagtc ctcatccaga tgaacaagaa gtttctgtct 4320
taggtgggat tccatactcc caaatatatg gatggtatcg agttcatttt ggggtgcttg 4380
atgaacaatt acatcgtaat aggggctaca gagatagata ttacagtaac ttagatattg 4440
ctccagcagc agatggttat ggattggcag gtttccctcc ggagcataga gcttggaggg 4500
aagagccgtg gattcatcat gcaccgccgg gttgtgggaa tgctccaaga tcatcgatga 4560
gtaatacttg cgatgaaaaa acccaaagtc taggtgtaaa attccttgac gaataccaat 4620
ctaaagttaa aagacaaata ttttcaggct atcaatctga tattgataca cataatagaa 4680
tttgaggatc ctcgcaatcc ctaggaggat taggcaaggg cttgagctca cgctcttggt 4740
agggacagaa atacaatcag gggcagtata tgaatactcc atggagaaac ccagatctac 4800
gtatgatcag cctcgactgt gccttctagt tgccagccat ctgttgtttg cccctcccc 4860
gtgccttctc tgaccctgga aggtgccact cccactgtcc tttcctaata aaatgaggaa 4920
attgcatcgc attgtctgag taggtgtcat tctattctgg ggggtggggg ggggcaggac 4980
agcaaggggg aggttggga agacaatagc aggcattgct gggatgcggg gggctctatg 5040
gcttctgagg cggaaagaac cagctggggc tcgacagctc gactctagaa ttcactggcc 5100
gtcgttttac aacgtcgtga ctgggaaaac cctggcgtaa cccaacttaa tcgccttgca 5160
gcacatcccc ctttcgccag ctggcgtaat agcgaagagg cccgcaccga tcgccttcc 5220
caacagttgc gcagcctgaa tggcgaatgg cgcctgatgc ggtattttct ccttacgcat 5280
ctgtgcggta tttcacaccg catatgggtg actctcagta caatctgctc tgatgccgca 5340
tagttaagcc agccccgaca cccgccaaac cccgctgacg cgccctgacg ggcttgtctg 5400
ctcccggcat ccgcttacag acaagctgtg accgtctccg ggagctgcat gtgtcagagg 5460
ttttcacctg catcaccgaa acgcgcga 5488

```

<210> 4

<211> 5500

<212> DNA

<213> plasmid pPJV2004

<400> 4

```

gacgaaaggg cctcgtgata cgcctatttt tatagggttaa tgtcatgata ataatggttt 60
cttagacgtc aggtggcact tttcggggaa atgtgcgcgg aaccctatt tgtttatttt 120
tctaaatata ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat 180
aatattgaaa aaggaagagt atgagtattc aacatttccg tgctgccctt attccctttt 240

```

ttgcgggcatt	ttgccttcct	gttttttgctc	acccagaaac	gctgggtgaaa	gtaaaagatg	300
ctgaagatca	gttgggtgca	cgagtgggtt	acatcgaact	ggatctcaac	agcggtaaga	360
tccttgagag	ttttcgcccc	gaagaacgtt	ttccaatgat	gagcactttt	aaagttctgc	420
tatgtggcgc	ggtattatcc	cgtattgacg	ccgggcaaga	gcaactcggg	cgccgcatac	480
actattctca	gaatgacttg	gttgagtact	caccagtcac	agaaaagcat	cttacggatg	540
gcatgacagt	aagagaatta	tgcagtgctg	ccataaccat	gagtataaac	actgcggcca	600
acttacttct	gacaacgata	ggaggaccga	aggagctaac	cgcttttttg	cacaacatgg	660
gggatcatgt	aactcgcctt	gatcgttggg	aaccggagct	gaatgaagcc	ataccaaacg	720
acgagcgtga	caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaaa	ctattaactg	780
gcgaactact	tactctagct	tcccggcaac	aattaataga	ctggatggag	gcggataaag	840
ttgcaggacc	acttctgcgc	tccggccctc	cggctggctg	gtttattgct	gataaatctg	900
gagccgggtga	gcgtgggtct	cgcggtatca	ttgcagcact	ggggccagat	ggtaagccct	960
cccgtatcgt	agttatctac	acgacgggga	gtcaggcaac	tatggatgaa	cgaaatagac	1020
agatcgctga	gatagggtgcc	tcaactgatta	agcattggta	actgtcagac	caagtttact	1080
catatatact	ttagattgat	ttaaaacttc	attttttaatt	taaaaggatc	taggtgaaga	1140
tccttttttga	taatctcatg	accaaaatcc	cttaacgtga	gttttcgttc	cactgagcgt	1200
cagaccccgt	agaaaagatc	aaaggatctt	cttgagatcc	tttttttctg	cgcgtaatct	1260
gctgcttgca	aacaaaaaaa	ccaccgctac	cagcgggtgg	ttgtttgccc	gatcaagagc	1320
taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	aatactgtcc	1380
ttctagtgtg	gccgtagtta	ggccaccact	tcaagaactc	tgtagcaccg	cctacatacc	1440
tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	tgtcttaccg	1500
ggttggactc	aagacgatag	ttaccggata	aggcgcagcg	gtcgggctga	acgggggggt	1560
cgtgcacaca	gcccagcttg	gagcgaacga	cctacaccga	actgagatac	ctacagcgtg	1620
agcattgaga	aagcggcacg	cttcccgaag	ggagaaaggc	ggacaggat	ccggtaagcg	1680
gcagggtcgg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	tggatatctt	1740
atagtcctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	attttttgtg	tgctcgtcag	1800
gggggaggag	cctatggaaa	aacgccagca	acgcggcctt	tttacggttc	ctggcctttt	1860
gctggccttt	tgctcacatg	ttctttcctg	cgttatcccc	tgattctgtg	gataaccgta	1920
ttaccgcctt	tgagttagct	gataccgctc	gccgcagccg	aacgaccgag	cgacagcagt	1980
cagttagcga	ggaagcggaa	gagcgcccaa	tacgcaaacc	gcctctcccc	gcgcgttggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccgaact	gaaagcgggc	agttagcgcg	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcacccca	ggctttacac	tttatgcttc	2160
cggctcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220
accatgatta	cgccaagcta	gtcgacataa	atcaatatgt	gctattggcc	attgcatacg	2280
ttgtatctat	atcataatat	gtacatttat	attggctcat	gtccaatatg	accgccatgt	2340
tgacattgat	tattgactag	ttattaatag	taatcaatta	cgggggtcatt	agttcatagc	2400
ccatataatg	agttccgcgt	tacataactt	acggtaaatg	gcccgcctcg	tgaccgcccc	2460
acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	2520
ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaac	tgcccacttg	gcagtacatc	2580
aagtgtatca	tatgccaaag	ccggcccccct	attgacgtca	atgacggtaa	atggccccgc	2640
tggcattatg	cccagtacat	gaccttacgg	gacttttccta	cttggcagta	catctacgta	2700
ttagtcatcg	ctattaccat	ggtgatgcgg	ttttggcagt	acaccaatgg	gcgtggatag	2760
cggtttgact	cacggggatt	tccaagtctc	caccccatgt	acgtcaatgg	gagtttgttt	2820
tggcaccaaaa	atcaacggga	ctttccaaaa	tgtcgtaata	accccgcccc	gttgacgcaa	2880
atggggcggt	ggcgtgtacg	gtgggaggtc	tatataagca	gagctcgttt	agtgaaccgt	2940
cagatcgccct	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	3000
tccagccctcc	gcggccggga	acggtgcatt	ggaacgcgga	ttccccgtgc	caagagtgcg	3060
gtaagtaccg	cctatagact	ctataggcac	accccttttg	ctcttatgca	tgctataactg	3120
tttttggctt	ggggcctata	cacccccgct	ccttatgcta	taggtgatgg	tatagcttag	3180
cctatagggtg	tgggttattg	accattattg	accactcccc	tattgggtgac	gatactttcc	3240
attactaatc	cataacatgg	ctcttttgcca	caactatctc	tattgggtat	atgccaatac	3300
tctgtccttc	agagactgac	acggactctg	tattttttaca	ggatgggggtc	ccatttatta	3360

```

tttacaaatt cacaatatata acaacgccgt cccccgtgcc cgcagttttt attaaacata 3420
gcgtgggata tccacgcgaa tctcgggtac gtgttccgga catgggctct tctccggtag 3480
cggcggagct tccacatccg agccctgggt ccatgcctcc agcggctcat ggtcgtcgg 3540
cagctccttg ctctaacag tggaggccag acttaggcac agcacaatgc ccaccaccac 3600
cagtgtgccg cacaaggccg tggcggtagg gtatgtgtct gaaaatgagc tcggagattg 3660
ggctcgcacc gtgacgcaga tgggaagact aaggcagcgg cagaagaaga tgcaggcagc 3720
tgagttgttg tattctgata agagtcagag gtaactcccg ttgcgggtgct gttaacggtg 3780
gagggcagtg tagtctgagc agtactcgtt gctgccgcgc gcgccaccag acataatagc 3840
tgacagacta acagactgtt cctttccatg ggtcttttct gcagtcaccg tccaagcttg 3900
caatcatgga tgcaatgaag agagggtctt gctgtgtgtt gctgctgtgt ggagcagttt 3960
tcgtttcggc tagcaatggc gacaaattat accgtgctga ctctagacct ccagatgaaa 4020
taaaacgttc cggaggtctt atgccagag ggcataatga gtacttcgat agaggaactc 4080
aaatgaatat taatctttat gatcacgcga gaggaacaca aaccggcttt gtcagatatg 4140
atgacggata tgtttccact tctcttagtt tgagaagtgc tcacttagca ggacagtcta 4200
tattatcagg atattccact tactatatat atgttatagc gacagcacca aatatgttta 4260
atgttaatga tgtattaggt gtatacagcc ctcaccata tgaacaggag gtttctgcgt 4320
taggtggaat accatattct cagatatatg gatggtatcg tgtaattttt ggtgtgattg 4380
atgaacgatt acatcgtaac agggaaatata gagaccggtt ttacagaaat ctgaatatag 4440
ctccggcaga ggatggttac agattagcag gtttcccacc ggatcaccaa gcttggagag 4500
aagaaccctg gattcatcat gcaccacaag gttgtggaaa ttcatacaaga acaattacag 4560
gtgatacttg taatgaggag acccagaatc tgagcacaat atatctcagg aaatatcaat 4620
caaaagttaa gaggcagata ttttcagact atcagtcaga ggttgacata tataacagaa 4680
ttcgggatga attatgagga tcctcgcgat ccctaggagg attaggcaag ggcttgagct 4740
cacgctcttg tgagggacag aaatacaatc aggggcagta tatgaatact ccatggagaa 4800
accagatct acgtatgatc agcctcgact gtgccttcta gttgccagcc atctgttgtt 4860
tgccccctcc cctgccttc cttgaccctg gaagggtgcca ctcccactgt cctttcctaa 4920
taaaatgagg aaattgcac gcattgtctg agtaggtgtc attctattct ggggggtggg 4980
gtggggcagg acagcaagg ggaggattgg gaagacaata gcaggcatgc tgggggatgcg 5040
gtgggctcta tggcttctga ggcggaaga accagctggg gctcgacagc tcgactctag 5100
aattcactgg ccgtcgtttt acaacgtcgt gactgggaaa accctggcgt tacccaactt 5160
aatcgccctg cagcacatcc ccctttcgcc agctggcgta atagcgaaga ggcccgcacc 5220
gategccctt cccaacagtt gcgcagcctg aatggcgaat ggcgcctgat gcggtathtt 5280
ctccttacgc atctgtgcgg tatttcacac cgcataatgg cactctcag tacaatctgc 5340
tctgatgccg catagttaag ccagccccga caccgcgcaa caccgcgtga cgcgccctga 5400
cgggcttgtc tgctcccggc atccgcttac agacaagctg tgaccgtctc cgggagctgc 5460
atgtgtcaga ggttttcacc gtcatcaccg aaacgcgcga 5500

```

<210> 5

<211> 5089

<212> DNA

<213> plasmid pPJV2005

<400> 5

```

gacgaaaggg cctcgtgata cgcctathtt tatagggttaa tgtcatgata ataattggtt 60
cttagacgtc aggtggcact tttcggggaa atgtgcgcgg aaccctatt tgtttathtt 120
tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat 180
aatattgaaa aaggaagagt atgagtattc aacatttccg tgcgcctt attccctttt 240
ttgcggcatt ttgccttcct gtttttgctc acccagaaac gctggtgaaa gtaaaagatg 300
ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac agcggtaga 360
tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactttt aaagtctctg 420
tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggg cgccgcatac 480
actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg 540
gcatgacagt aagagaatta tgcagtgttg ccataaccat gaggatgata actgcggcca 600

```

acttactttct	gacaacgatc	ggaggaccga	aggagctaac	cgctttttttg	cacaacatgg	660
gggatcatgt	aactcgcctt	gatcgttggg	aaccggagct	gaatgaagcc	ataccaaacg	720
acgagcgtga	caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaaa	ctattaactg	780
gcgaactact	tactctagct	tcccggaac	aattaataga	ctggatggag	gcggataaag	840
ttgcaggacc	acttctgcgc	tccggcccttc	cggctggctg	gtttattgct	gataaatctg	900
gagccggtga	gcgtgggtct	cgcggtatca	ttgcagcact	ggggccagat	ggtaagccct	960
cccgtatcgt	agttatctac	acgacgggga	gtcaggcaac	tatggatgaa	cgaaatagac	1020
agatcgctga	gatagggtgcc	tcactgatta	agcattggta	actgtcagac	caagtttact	1080
catatatact	ttagattgat	ttaaaacttc	atttttaatt	taaaaggatc	taggtgaaga	1140
tccttttttga	taatctcatg	accaaatacc	cttaacgtga	gttttcgctc	cactgagcgt	1200
cagaccccggt	agaaaagatc	aaaggatctt	cttgagatcc	tttttttctg	cgcgtaatct	1260
gctgcttgca	aacaaaaaaa	ccaccgctac	cagcgggtgg	ttgtttgccg	gatcaagagc	1320
taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	aatactgtcc	1380
ttctagtgtg	gccgtagtta	ggccaccact	tcaagaactc	tgtagcaccg	cctacatacc	1440
tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	tgtcttaccg	1500
ggttggactc	aagacgatag	ttaccggata	aggcgcagcg	gtcgggctga	acgggggggt	1560
cgtgcacaca	gcccagcttg	gagcgaacga	cctacaccga	actgagatac	ctacagcgtg	1620
agcattgaga	aagcgcacg	cttcccgaag	ggagaaaggc	ggacagggtat	ccggtaagcg	1680
gcagggtcgg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	tggatatctt	1740
atagtccctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	atttttgtga	tgctcgtcag	1800
gggggcggag	cctatggaaa	aacgccagca	acgcggcctt	tttacggctc	ctggcctttt	1860
gctggccttt	tgctcacatg	ttcttttctg	cgttatcccc	tgattctgtg	gataaccgta	1920
ttaccgcctt	tgagtgagct	gataccgctc	gccgcagccg	aacgaccgag	cgcagcagag	1980
cagtgagcga	ggaagcggaa	gagcgcacca	tacgcaaacc	gcctctcccc	gcgcgttggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccgaactg	gaaagcgggc	agtgagcgca	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcacccca	ggctttacac	tttatgcttc	2160
cggctcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220
accatgatta	cgccaagcta	gtcgacataa	atcaatatgt	gctattggcc	attgcatacg	2280
ttgtatctat	atcataatat	gtacatttat	attggctcat	gtccaatatg	accgccatgt	2340
tgacattgat	tattgactag	ttattaatag	taatcaatta	cgggggtcatt	agttcatagc	2400
ccatataatgg	agttccgcgt	tacataactt	acggtaaattg	gcccgcctcg	tgaccgcca	2460
acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	2520
ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaac	tgcccacttg	gcagtacatc	2580
aagtgtatca	tatgccaaagt	ccggcccccct	attgacgtca	atgacggtaa	atggcccgcc	2640
tggcattatg	cccagttacat	gaccttacgg	gactttccta	cttggcagta	catctacgta	2700
ttagtcatcg	ctattaccat	ggtgatgcgg	ttttggcagt	acaccaatgg	gcgtggatag	2760
cggttttgact	cacgggggatt	tccaagtctc	cacccatttg	acgtcaatgg	gagttttgtt	2820
tggcaccaaa	atcaacggga	ctttccaaaa	tgctcgtaata	accccgcccc	gttgacgcaa	2880
atgggcggta	ggcgtgtacg	gtgggaggtc	tatataagca	gagctcgttt	agtgaaccgt	2940
cagatcgctt	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	3000
tccagcctcc	gcggccggga	acggtgcatt	ggaacgcgga	ttccccgtgc	caagagtgc	3060
gtaagtaccg	cctatagact	ctataggcac	accccttttg	ctcttatgca	tgctatactg	3120
tttttggctt	ggggcctata	cacccccgct	ccttatgcta	taggtgatgg	tatagcttag	3180
cctataggtg	tgggttattg	accattattg	accactcccc	tattgggtgac	gatactttcc	3240
attactaatc	cataacatgg	ctcttttgcca	caactatctc	tattggctat	atgccaatac	3300
tctgtccttc	agagactgac	acggactctg	tattttttaca	ggatgggggtc	ccatttatta	3360
tttacaaatt	cacatatata	acaacgccgt	cccccggtgc	cgcagttttt	attaaacata	3420
gcgtgggatc	tccacgcgaa	tctcgggtac	gtgttccgga	catgggctct	tctccggtag	3480
cggcggagct	tccacatccg	agccctgggtc	ccatgcctcc	agcggctcat	ggtcgctcgg	3540
cagctccttg	ctcctaacag	tggaggccag	acttaggcac	agcacaatgc	ccaccaccac	3600
cagtgtgccg	cacaaggccg	tggcggtagg	gtatgtgtct	gaaaatgagc	tcggagattg	3660
ggctcgcacc	gtgacgcaga	tggaagactt	aaggcagcgg	cagaagaaga	tgagggcagc	3720

tgagttgttg	tattctgata	agagtcagag	gtaactcccg	ttgcggtgct	gttaacggtg	3780
gagggcagtg	tagtctgagc	agtactcggt	gctgccgcgc	gcgccaccag	acataatagc	3840
tgacagacta	acagactggt	cctttccatg	ggctctttct	gcagtcaccg	tccaagcttg	3900
caatcatgga	tgcaatgaag	agagggctct	gctgtgtgct	gctgctgtgt	ggagcagtc	3960
tcgtttcggc	tagcgctccc	cagtctatta	cagaactatg	ttcgggaatat	cgcaacacac	4020
aaatatatac	gataaatgac	aagatactat	catatacgga	atcgatggca	ggcaaaagag	4080
aatgggttat	cattacatth	aagagcggcg	caacatttca	ggtcgaagtc	ccgggcagtc	4140
aacatataga	ctccccaaaa	aaagccattg	aaaggatgaa	ggacacatta	agaatcacat	4200
atctgaccga	gaccaaaatt	gataaattat	gtgtatggaa	taataaaaacc	cccaattcaa	4260
ttgcggcaat	cagtatggaa	aactagggat	cctcgcaatc	cctaggagga	ttaggcaagg	4320
gcttgagctc	acgctcttgt	gagggacaga	aatacaatca	ggggcagtat	atgaatactc	4380
catggagaaa	cccagatcta	cgtatgatca	gcctcgactg	tgccttctag	ttgccagcca	4440
tctgttgttt	gccccctccc	cgtgccttcc	ttgaccctgg	aagggtgccac	tcccactgtc	4500
ctttccta	aaaatgagga	aattgcatcg	cattgtctga	gtaggtgtca	ttctattctg	4560
gggggtgggg	tggggcagga	cagcaagggg	gaggattggg	aagacaatag	caggcatgct	4620
ggggatgcgg	tgggctctat	ggcttctgag	gcggaaagaa	ccagctgggg	ctcgacagct	4680
cgactctaga	attcactggc	cgtcgtttta	caacgtcgtg	actgggaaaa	ccctggcggt	4740
acccaactta	atcgccttgc	agcacatccc	cctttcgcga	gctggcgtaa	tagcgaagag	4800
gcccgcaccg	atcgccttcc	ccaacagttg	cgcagcctga	atggcgaaatg	gcgcctgatg	4860
cggtattttc	tccttacgca	tctgtgcggg	atctcacacc	gcatatgggtg	cactctcagt	4920
acaatctgct	ctgatgccgc	atagttaagc	cagccccgac	acccgccaac	acccgctgac	4980
gcgccctgac	gggcttgtct	gctcccggca	tccgcttaca	gacaagctgt	gaccgtctcc	5040
gggagctgca	tgtgtcagag	gttttcaccg	tcatacccca	aacgcgcga		5089

<210> 6

<211> 5488

<212> DNA

<213> plasmid pPJV2007

<400> 6

gacgaaaggg	cctcgtgata	cgcctattht	tatagggttaa	tgtcatgata	ataatggtht	60
cttagacgct	aggtggcact	tttcggggaa	atgtgcgcgg	aaccctatt	tgtttattht	120
tctaaataca	ttcaaataatg	tatccgctca	tgagacaata	accctgataa	atgcttcaat	180
aatattgaaa	aaggaagagt	atgagtatth	aacattthccg	tgtcgcctt	attcccttht	240
ttgcggcatt	ttgccttctt	gtttttgtct	accagaaaac	gctgggtgaaa	gtaaaagatg	300
ctgaagatca	gttgggtgca	cgagtgggtt	acatcgaact	ggatctcaac	agcggtaaga	360
tccttgagag	ttttcgcctt	gaagaacggt	ttccaatgat	gagcacttht	aaagttctgc	420
tatgtggcgc	ggtattatcc	cgtattgacg	ccgggcaaga	gcaactcggg	cgccgcatac	480
actattctca	gaatgacttg	gttgagtact	caccagtcac	agaaaagcat	cttacggatg	540
gcatgacagt	aagagaatta	tgcagtgtctg	ccataaccat	gagtgataac	actgcggcca	600
acttacttct	gacaacgatc	ggaggaccga	aggagctaac	cgcttttttg	cacaacatgg	660
gggatcatgt	aactcgcctt	gatcgttggg	aaccggagct	gaatgaagcc	ataccaaacg	720
acgagcgtga	caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaaa	ctattaactg	780
gcgaactact	tactctagct	tcccggcaac	aattaataga	ctggatggag	gcggataaag	840
ttgcaggacc	acttctgcgc	tcggcccttc	cggctggctg	gtttattgct	gataaatctg	900
gagccgggtga	gcgtgggtct	cgcggtatca	ttgcagcact	ggggccagat	ggtaagccct	960
cccgtatcgt	agttatctac	acgacgggga	gtcaggcaac	tatggatgaa	cgaaatagac	1020
agatcgctga	gatagggtgc	tcactgatta	agcattggta	actgtcagac	caagtttact	1080
catatatact	ttagattgat	ttaaaacttc	atthtttaatt	taaaaggatc	taggtgaaga	1140
tcctttttga	taatctcatg	accaaaatcc	cttaacgtga	gttttcgtht	cactgagcgt	1200
cagaccccg	agaaaagatc	aaaggatctt	cttgagatcc	ttttttctg	cgcgtaatct	1260
gctgcttgca	aacaaaaaaa	ccaccgctac	cagcgggtgg	ttgtttgccg	gatcaagagc	1320
taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	aatactgtcc	1380

ttctagtgtgta	gccgtagttaa	ggccaccact	tcaagaactc	tgtagcaccg	cctacatacc	1440
tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	tgtcttaccg	1500
ggttggaactc	aagacgatag	ttaccggata	aggcgcagcg	gtcgggctga	acgggggggtt	1560
cgtgcacaca	gcccagcttg	gagcgaacga	cctacaccga	actgagatac	ctacagcgtg	1620
agcattgaga	aagcgccacg	cttcccgaag	ggagaaaggc	ggacaggat	ccggtaagcg	1680
gcagggctcgg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	tggtatcttt	1740
atagtcctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	atTTTTgtga	tgctcgtcag	1800
gggggaggag	cctatggaaa	aacgccagca	acgcggcctt	tttacgggtt	ctggcctttt	1860
gctggccttt	tgctcacatg	ttctttcctg	cgttatcccc	tgattctgtg	gataaccgta	1920
ttaccgcctt	tgagtgagct	gataccgctc	gccgcagccg	aacgaccgag	cgcagcagag	1980
cagtgagcga	ggaagcggaa	gagcgcccaa	tacgcaaacc	gcctctcccc	gcgcgttggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccgactg	gaaagcgggc	agtgagcgca	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcacccca	ggctttacac	tttatgcttc	2160
cggctcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220
accatgatta	cgccaagcta	gtcgacataa	atcaatatgt	gctattggcc	attgcatacg	2280
ttgtatctat	atcataatat	gtacatttat	attggctcat	gtccaatatg	accgccatgt	2340
tgacattgat	tattgactag	ttattaatag	taatcaatta	cggggtcatt	agttcatagc	2400
ccatatatgg	agttccgcgt	tacataactt	acggtaaagt	gcccgcctcg	tgaccgcca	2460
acgacccccg	cccattgacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	2520
ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaac	tgcccacttg	gcagtacatc	2580
aagtgtatca	tatgccaaagt	ccggcccccct	attgacgtca	atgacggtaa	atggcccgcc	2640
tggcattatg	cccagtacat	gaccttacgg	gactttccta	cttggcagta	catctacgta	2700
ttagtcatcg	ctattaccat	ggtgatgcgg	ttttggcagt	acaccaatgg	gcgtggatag	2760
cggtttgact	cacggggatt	tccaagtctc	caccccatgt	acgtcaatgg	gagtttgttt	2820
tggcaccaaa	atcaacggga	ctttccaaaa	tgctcgtaata	accccgcccc	gttgacgcaa	2880
atgggaggga	ggcgtgtacg	gtgggagggtc	tatataagca	gagctcgttt	agtgaaccgt	2940
cagatcgctt	ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	3000
tccagcctcc	gcggccggga	acgggtgcatt	ggaacgcgga	ttccccgtgc	caagagtgc	3060
gtaagtaccg	cctatagact	ctataggcac	acccctttgg	ctcttatgca	tgctatactg	3120
tttttggtct	ggggcctata	cacccccgct	ccttatgcta	taggtgatgg	tatagcttag	3180
cctataggtg	tgggttattg	accattattg	accactcccc	tattgggtgac	gatactttcc	3240
attactaatc	cataacatgg	ctcttttgcca	caactatctc	tattggctat	atgccaatac	3300
tctgtccttc	agagactgac	acggactctg	tatTTTTtaca	ggatgggggtc	ccattttatta	3360
tttacaaatt	cacatatata	acaacgccgt	cccccggtgc	cgcagttttt	attaaacata	3420
gcgtgggata	tccacgcgaa	tctcgggtac	gtgttccgga	catgggctct	tctccggtag	3480
cggcggagct	tccacatccg	agccctgggtc	ccatgcctcc	agcggctcat	ggtcgctcgg	3540
cagctccttg	ctcctaacag	tggaggccag	acttaggcac	agcacaatgc	ccaccaccac	3600
cagtgtgccg	cacaaggccg	tggcggtagg	gtatgtgtct	gaaaatgagc	tcggagattg	3660
ggctcgcacc	gtgacgcaga	tggaagactt	aaggcagcgg	cagaagaaga	tgcaaggcagc	3720
tgagttgttg	tattctgata	agagtcagag	gtaactcccc	ttgcgggtgct	gttaacgggtg	3780
gagggcagtg	tagtctgagc	agtactcgtt	gctgccgcgc	gcgccaccag	acataatagc	3840
tgacagacta	acagactggt	cctttccatg	ggtcttttct	gcagtcaccg	tccaagcttg	3900
caatcatgga	tgcaatgaag	agagggtctt	gctgtgtgct	gctgctgtgt	ggagcagtct	3960
tcgtttcggc	tagcaatggc	gacaaattat	accgtgctga	ctctagaccc	ccagatgaaa	4020
taaaacgttc	cggaggtctt	atgccagat	ggcataatga	gtacttcgat	agaggaaactc	4080
aatgaatat	taatctttat	gatcacgcga	gaggaaacaca	aaccggcttt	gtcagatatg	4140
atgacggata	tgtttccact	tctcttagtt	tgagaagtgc	tcacttagca	ggacagtcta	4200
tattatcagg	atattccact	tactatatat	atgttatagc	gacagcacca	aatatgttta	4260
atgttaatga	tgtattaggg	gtatacagcc	ctcaccata	tgaacaggag	gtttctgcgt	4320
taggtggaat	accatattct	cagatatatg	gatggtatcg	tgtaattttt	ggtgtgattg	4380
atgaacgatt	acatcgtaac	agggaatata	gagaccggta	ttacagaaat	ctgaatatag	4440
ctccggcaga	ggatgggttac	agattagcag	gtttccacc	ggatcaccaa	gcttggagag	4500

```

aagaaccctg gattcatcat gcaccacaag gttgtggaaa ttcattcaaga acaattacag 4560
gtgatacttg taatgaggag acccagaatc tgagcacaat atatctcagg aaatatcaat 4620
caaaagttaa gaggcagata ttttcagact atcagtcaga gggtgacata tataacagaa 4680
tttgaggatc ctcgcaatcc ctaggaggat taggcaaggg cttgagctca cgctcttgtg 4740
agggacagaa atacaatcag gggcagtata tgaataactcc atggagaaac ccagatctac 4800
gtatgatcag cctcgactgt gccttctagt tgccagccat ctggtgtttg cccctcccc 4860
gtgccttcct tgaccctgga aggtgccact cccactgtcc tttcctaata aaatgaggaa 4920
attgcacgcg attgtctgag taggtgtcat tctattctgg ggggtggggg ggggcaggac 4980
agcaaggggg aggattggga agacaatagc aggcattgctg gggatgcggg gggctctatg 5040
gcttctgagg cggaaagaac cagctggggc tcgacagctc gactctagaa ttcactggcc 5100
gtcgttttac aacgtcgtga ctgggaaaac cctggcggtt cccaacttaa tcgccttgca 5160
gcacatcccc ctttcgccag ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc 5220
caacagttgc gcagcctgaa tggcgaatgg cgcctgatgc ggtattttct ccttacgcat 5280
ctgtgcggta tttcacaccg catatgggtg actctcagta caatctgctc tgatgccgca 5340
tagttaagcc agccccgaca cccgccaaac cccgctgacg cgccttgacg ggcttgtctg 5400
ctcccggcat ccgcttacag acaagctgtg accgtctccg ggagctgcat gtgtcagagg 5460
ttttcacctg catcaccgaa acgcgcga

```

<210> 7

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 7

ggagctagca atgatgataa gttatatcgg

30

<210> 8

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 8

cctggatcct cataattcat ccttaattct

30

<210> 9

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 9

ggagctagca cacctcaaaa tattactgat

30

<210> 10

<211> 30

<212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic
 construct

<400> 10
 cctggatcct taatttgcca tactaattgc 30
 <210> 11
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic
 construct

<400> 11
 cctggatcct caaattctat tatgtgtatc 30
 <210> 12
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic
 construct

<400> 12
 ggagctagca atggcgacaa attataccgt 30
 <210> 13
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic
 construct

<400> 13
 cctggatcct cataattcat cccgaattct 30
 <210> 14
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:synthetic
 construct

<400> 14

ggagctagcg ctccccagtc tattacagaa 30
 <210> 15
 <211> 30
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:synthetic
 construct

 <400> 15
 cctggatccc tagttttcca tactgattgc 30
 <210> 16
 <211> 30
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:synthetic
 construct

 <400> 16
 cctggatcct caaattctgt tatatatgtc 30

 <210> 17
 <211> 65
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:synthetic
 construct

 <400> 17
 cccaagcttc caccatgagc cttctaaccg aggtcgaaac acctatcaga aacgaatggg 60
 agtgc 65
 <210> 18
 <211> 28
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:synthetic
 construct

 <400> 18
 cccggatcct tactccagct ctatgctg 28
 <210> 19
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 19

Ser	Leu	Leu	Thr	Glu	Val	Glu	Thr	Pro	Ile	Arg	Asn	Glu	Trp	Glu	Cys
1				5					10					15	

Arg

<210> 20

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 20

Ile	Pro	Gln	Ser	Leu	Asp	Ser	Trp	Trp	Thr	Ser	Leu
1				5					10		

<210> 21

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 21

Arg	Ile	Gln	Arg	Gly	Pro	Gly	Arg	Ala	Phe	Val	Ile	Thr	Gly	Lys
1				5					10					15

<210> 22

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic
construct

<400> 22

Arg	Gly	Pro	Gly	Arg	Ala	Phe	Val	Thr	Ile
1				5					10

Claims

What is claimed is:

- 5 1. A composition comprising first and second nucleic acid sequences, wherein said first nucleic acid sequence is a truncated A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and said second nucleic acid sequence is a truncated B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, with the proviso
10 that each of said truncated subunit coding regions has a 5' deletion and encodes a subunit peptide not having an amino terminal bacterial signal peptide.
2. The composition of claim 1, wherein said first and second nucleic acid sequences are present in a single nucleic acid construct.
- 15 3. The composition of claim 2, wherein said nucleic acid construct is a plasmid vector.
4. The composition of claim 2, wherein the first and second nucleic acid sequences are operably linked to a transcriptional control element.
- 20 5. The composition of claim 4, wherein said transcriptional control element is a heterologous promoter.
6. The composition of claim 1 wherein said first and second nucleic acid sequences are present in separate nucleic acid constructs.
- 25 7. The composition of claim 6, wherein said separate nucleic acid constructs are plasmid vectors.

30

8. The composition of claim 1, wherein the truncated subunit coding regions are obtained or derived from the same bacterial ADP-ribosylating exotoxin.

5 9. The composition of claim 8, wherein said bacterial ADP-ribosylating exotoxin is a cholera toxin (CT).

10. The composition of claim 8, wherein said bacterial ADP-ribosylating exotoxin is an *E. coli* heat labile enterotoxin (LT).

10 11. The composition of claim 1, wherein at least one of the truncated subunit coding regions has been genetically modified to detoxify the subunit peptide encoded thereby.

15 12. The composition of claim 11, wherein the truncated A subunit coding region has been genetically modified to disrupt or inactivate ADP-ribosyl transferase activity in the subunit peptide encoded thereby.

20 13. The composition of claim 1, wherein the truncated A subunit coding region has been further genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby.

25 14. The composition of claim 1 further comprising an antigen of interest.

15. The composition of claim 14, wherein said antigen is from a bacterial, viral or parasitic pathogen.

30 16. The composition of claim 1, further comprising a third nucleic acid sequence that encodes an antigen of interest.

17. The composition of claim 16, wherein said antigen is from a bacterial, viral or parasitic pathogen.

18. The composition of claim 16, wherein said third nucleic acid sequence is present in a nucleic acid construct that does not contain said first or said second nucleic acid sequence.

19. The composition of claim 18, wherein the nucleic acid construct containing the third nucleic acid sequence is a plasmid vector.

20. The composition of claim 16, wherein said third nucleic acid sequence is present in a nucleic acid construct that also contains at least one of said first or said second nucleic acid sequence.

21. The composition of claim 20, wherein the nucleic acid construct containing the third nucleic acid sequence is a plasmid vector.

22. The composition of claim 1, wherein said composition is in a particulate form.

23. The composition of claim 22, wherein said particulate composition is suitable for transdermal delivery via a particle delivery device.

24. The composition of claim 1, further comprising a pharmaceutically acceptable vehicle or excipient.

25. The composition of claim 1, wherein the first and second nucleic acid sequences are coated onto a core carrier particle.

26. The composition of claim 25, wherein the core carrier particle has an average diameter of about 0.1 to about 10 μ m.

27. The composition of claim 25, wherein the core carrier particle comprises a metal.

28. The composition of claim 27 wherein the metal is gold.

5

29. A particle delivery device, wherein said device is loaded with a particulate vaccine composition as defined in claim 23.

30. The composition of claim 1 further comprising a transfection
10 facilitating agent.

31. The composition of claim 30, wherein the transfection facilitating agent is a liposome.

15 32. A composition comprising first and second nucleic acid sequences, wherein said first nucleic acid sequence is a modified A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and said second nucleic acid sequence is a B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, with the proviso that said
20 modified A subunit coding region and said B subunit coding region each encode a mature subunit peptide, and with the further proviso that the modified A subunit coding region has been genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby.

25 33. The composition of claim 32, wherein said first and second nucleic acid sequences are present in a single nucleic acid construct.

34. The composition of claim 33, wherein said nucleic acid construct is a plasmid vector.

30

35. The composition of claim 33, wherein the first and second nucleic acid sequences are operably linked to a transcriptional control element.

36. The composition of claim 35; wherein said transcriptional control element is a heterologous promoter.

37. The composition of claim 32, wherein said first and second nucleic acid sequences are present in separate nucleic acid constructs.

38. The composition of claim 37, wherein said separate nucleic acid constructs are plasmid vectors.

39. The composition of claim 32, wherein the B and modified A subunit coding regions are obtained or derived from the same bacterial ADP-ribosylating exotoxin.

40. The composition of claim 39, wherein said bacterial ADP-ribosylating exotoxin is a cholera toxin (CT).

41. The composition of claim 39, wherein said bacterial ADP-ribosylating exotoxin is an *E. coli* heat labile enterotoxin (LT).

42. The composition of claim 32, wherein at least one of the B or modified A subunit coding regions has been genetically modified to detoxify the subunit peptide encoded thereby.

43. The composition of claim 42, wherein the modified A subunit coding region has been genetically modified to disrupt or inactivate ADP-ribosyl transferase activity in the subunit peptide encoded thereby.

44. The composition of claim 32, wherein the modified A subunit coding region and the B subunit coding region have each been truncated by a 5' deletion whereby each of said truncated subunit coding regions encodes a subunit peptide not having an amino terminal bacterial signal peptide.

5

45. The composition of claim 32 further comprising an antigen of interest.

46. The composition of claim 45, wherein said antigen is from a bacterial, viral or parasitic pathogen.

10

47. The composition of claim 32 further comprising a third nucleic acid sequence that encodes an antigen of interest.

48. The composition of claim 47, wherein said antigen is from a bacterial, viral or parasitic pathogen.

15

49. The composition of claim 47, wherein said third nucleic acid sequence is present in a nucleic acid construct that does not contain said first or said second nucleic acid sequence.

20

50. The composition of claim 49, wherein the nucleic acid construct containing the third nucleic acid sequence is a plasmid vector.

51. The composition of claim 47, wherein said third nucleic acid sequence is present in a nucleic acid construct that also contains at least one of said first or said second nucleic acid sequence.

25

52. The composition of claim 51, wherein the nucleic acid construct containing the third nucleic acid sequence is a plasmid vector.

30

53. The composition of claim 32, wherein said composition is in a particulate form.

54. The composition of claim 53, wherein said particulate composition is suitable for transdermal delivery via a particle delivery device.

55. The composition of claim 32 further comprising a pharmaceutically acceptable vehicle or excipient.

56. A composition according to claim 55, wherein the first and second nucleic acid sequences are coated onto a core carrier particle.

57. The composition of claim 56, wherein the core carrier particle has an average diameter of about 0.1 to about 10 μ m.

58. The composition of claim 56, wherein the core carrier particle comprises a metal.

59. The composition of claim 58, wherein the metal is gold.

60. A particle delivery device, wherein said device is loaded with a particulate vaccine composition as defined in claim 54.

61. The composition of claim 32 further comprising a transfection facilitating agent.

62. The composition of claim 61, wherein the transfection facilitating agent is a liposome.

63. Use of a composition comprising a first and second nucleic acid sequence, each said sequence including a coding region for a subunit from a

bacterial ADP-ribosylating exotoxin in the manufacture of a medicament for enhancing an immune response in a vertebrate subject against an antigen of interest in the said subject by administering the antigen of interest and the said composition to the subject whereby the toxin subunits encoded by the first and second nucleic acid sequences are expressed in an amount sufficient to elicit an enhanced immune response against the antigen, wherein said first nucleic acid sequence contains a truncated A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and said second nucleic acid sequence contains a truncated B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, with the proviso that each of said truncated subunit coding regions has a 5' deletion and encodes a subunit peptide not having an amino terminal bacterial signal peptide.

64. Use according to claim 63, wherein the antigen of interest and the composition are administered to the same site in the subject.

65. Use according to claim 63, wherein the antigen of interest and the composition are administered concurrently.

66. Use according to claim 65, wherein the antigen of interest and the composition are combined to provide a single vaccine composition.

67. Use according to claim 63, wherein the antigen of interest is from a bacterial, viral or parasitic pathogen.

68. Use according to claim 67, wherein a third nucleic acid sequence is administered to the subject and the third sequence encodes said antigen of interest.

69. Use according to claim 63, wherein the first and second nucleic acid sequences are administered to the subject in particulate form.

70. Use according to claim 69, wherein the composition containing the first and second nucleic acid sequences are coated onto a core carrier particle and administered to the subject using a particle-mediated delivery technique.

5 71. Use according to claim 63, wherein the subject is human.

72. Use of a composition comprising a first and second nucleic acid sequence, each said sequence including a coding region for a subunit from a bacterial ADP-ribosylating exotoxin in the manufacture of a medicament for enhancing an immune response in a vertebrate subject against an antigen of interest in the said subject by administering the antigen of interest and the said composition to the subject whereby the toxin subunits encoded by the first and second nucleic acid sequences are expressed in an amount sufficient to elicit an enhanced immune response against the antigen, wherein said first nucleic acid sequence contains a modified A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and said second nucleic acid sequence contains a B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, with the proviso that said modified A subunit coding region and said B subunit coding region each encode a mature subunit peptide, and with the further proviso that the modified A subunit coding region has been genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby.

73. Use according to claim 72, wherein the antigen of interest and the composition are administered to the same site in the subject.

74. Use according to claim 72, wherein the antigen of interest and the composition are administered concurrently.

75. Use according to claim 74, wherein the antigen of interest and the composition are combined to provide a single vaccine composition.

76. Use according to claim 72, wherein the antigen of interest is from a bacterial, viral or parasitic pathogen.

77. Use according to claim 76, wherein a third nucleic acid sequence is administered to the subject and the third sequence encodes said antigen of interest.

78. Use according to claim 72, wherein the first and second nucleic acid sequences are administered to the subject in particulate form.

79. Use according to claim 78, wherein the composition comprising the first and second nucleic acid sequences are coated onto a core carrier particle and administered to the subject using a particle-mediated delivery technique.

80. Use according to claim 72, wherein the subject is human.

81. A method for enhancing an immune response against an antigen of interest in a subject, the method comprising:

(a) administering the antigen of interest to the subject;

(b) providing an adjuvant composition comprising first and second nucleic acid sequences, wherein said first nucleic acid sequence is a truncated A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and said second nucleic acid sequence is a truncated B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, with the proviso that each of said truncated subunit coding regions has a 5' deletion and encodes a subunit peptide not having an amino terminal bacterial signal peptide; and


(c) administering said adjuvant composition to the subject, whereby upon introduction to the subject, the first and second nucleic acid sequences are expressed to provide subunit peptides in an amount sufficient to elicit said enhanced immune response against the antigen of interest.

82. A method for enhancing an immune response against an antigen of interest in a subject, the method comprising:

(a) administering the antigen of interest to the subject;

(b) providing an adjuvant composition comprising first and second
5 nucleic acid sequences, wherein said first nucleic acid sequence is a modified A subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, and said second nucleic acid sequence is a B subunit coding region obtained or derived from a bacterial ADP-ribosylating exotoxin, with the proviso that said modified A subunit coding region and said B subunit coding region each
10 encode a mature subunit peptide, and with the further proviso that the modified A subunit coding region has been genetically modified so as to delete a C-terminal KDEL or RDEL motif in the subunit peptide encoded thereby; and

(c) administering said adjuvant composition to the subject, whereby
upon introduction to the subject, the first and second nucleic acid sequences are
15 expressed to provide subunit peptides in an amount sufficient to elicit said enhanced immune response against the antigen of interest.



Molecule: pPJV2002, 5500 bps DNA Circular
 File Name: pPJV2002.cm5,

Description: Ligation of CTA PCR frag Nhe Bam cut into 7054 Nhe Bam Vector

Notes:

Molecule Features:

Type	Start	End	Name	Description
REGION	2242	3060	CMVpro	
REGION	3061	3884	intronA	
GENE	3906	3969	TPAsigCDS'	
GENE	3975	4697	CTA CDS	
REGION	4805	5101	bGHpA	

Enzymes (15 sites)

Sall	2241,	MscI	2266,	SpeI	2356,	SacII	3009
NsiI	3106,	PstI	3879,	HindIII	3894,	NheI	3969
ClaI	4553,	BamHI	4698,	BglII	4805,	EcoRI	5100

FIGURE 1-1

```

1   GACGAAAGGG CCTCGTGATA CGCCTATTTT TATAGGTTAA TGTCATGATA ATAATGGTTT
61  CTTAGACGTC AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTTTATTTT
121 TCTAAATACA TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT
181 AATATTGAAA AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTCGCCCTT ATTCCCTTTT
241 TTGCGGCATT TTGCCCTCCT GTTTTTTGCTC ACCCAGAAAC GCTGGTGAAA GTAAAAGATG
301 CTGAAGATCA GTTGGGTGCA CGAGTGGGTT ACATCGAACT GGATCTCAAC AGCGGTAAGA
361 TCCTTGAGAG TTTTCGCCCC GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCTGC
421 TATGTGGCGC GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC
481 ACTATTCTCA GAATGACTTG GTTGAGTACT CACCAGTCAC AGAAAAGCAT CTTACGGATG
541 GCATGACAGT AAGAGAATTA TGCAGTGCTG CCATAACCAT GAGTGATAAC ACTGCGGCCA
601 ACTTACTTCT GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTTTG CACAACATGG
661 GGGATCATGT AACTCGCCTT GATCGTTGGG AACCGGAGCT GAATGAAGCC ATACCAAACG
721 ACGAGCGTGA CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAA CTATTAAC TG
781 GCGAACTACT TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG
841 TTGCAGGACC ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG
901 GAGCCGGTGA GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT
961 CCCGTATCGT AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC
1021 AGATCGCTGA GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT
1081 CATATATACT TTAGATTGAT TTAAAACTTC ATTTTAAATT TAAAAGGATC TAGGTGAAGA
1141 TCCTTTTTTG TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTT CACTGAGCGT
1201 CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT
1261 GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC
1321 TACCAACTCT TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC
1381 TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACCG CCTACATACC
1441 TCGCTCTGCT AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG
1501 GGTTGGACTC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGTT
1561 CGTGACACCA GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG
1621 AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG
1681 GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT
1741 ATAGTCCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG
1801 GGGGGCGGAG CCTATGGAAG AACGCCAGCA ACGCGCCTT TTTACGGTTC CTGGCCTTTT
1861 GCTGGCCTTT TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA
1921 TTACCGCCTT TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT
1981 CAGTGAGCGA GGAAGCGGAA GAGCGCCCAA TACGCAAACC GCCTCTCCCC GCGCGTTGGC
2041 CGATTCAATTA ATGCAGCTGG CACGACAGGT TTCCCGACTG GAAAGCGGGC AGTGAGCGCA
2101 ACGCAATTAA TGTGAGTTAG CTCACTCATT AGGCACCCCA GGCTTTACAC TTTATGCTTC
2161 CGGCTCGTAT GTTGTGTGGA ATTGTGAGCG GATAACAATT TCACACAGGA AACAGCTATG
2221 ACCATGATTA CGCCAAGCTA GTCGACATAA ATCAATATTG GCTATTGGCC ATTGCATACG
2281 TTGTATCTAT ATCATAATAT GTACATTTAT ATTGGCTCAT GTCCAATATG ACCGCCATGT
2341 TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCATT AGTTCATAGC
2401 CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCCCGCCTCG TGACCGCCCA
2461 ACGACCCCCG CCCATTGACG TCAATAATGA CGTATGTTCC CATAGTAACG CCAATAGGGA
2521 CTTTCCATTG ACGTCAATGG GTGGAGTATT TACGGTAAAC TGCCCACTTG GCAGTACATC
2581 AAGTGATATCA TATGCCAAGT CCGGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCGCC
2641 TGGCATTATG CCCAGTACAT GACCTTACGG GACTTTCCTA CTTGGCAGTA CATCTACGTA
2701 TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACACCAATGG GCGTGGATAG
2761 CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCATTG ACGTCAATGG GAGTTTGTTT
2821 TGGCACCAA ATCAACGGGA CTTTCCAAA TGTCGTAATA ACCCGCCCC GTTGACGCAA
2881 ATGGGCGGTA GCGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT
2941 CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA CCGGGACCGA
3001 TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCGGTGC CAAGAGTGAC
3061 GTAAGTACCG CCTATAGACT CTATAGGCAC ACCCCTTTGG CTCTTATGCA TGCTATACTG
3121 TTTTGGCTT GGGGCCTATA CACCCCGCT CTTATGCTA TAGGTGATGG TATAGCTTAG
3181 CCTATAGGTG TGGGTTATTG ACCATTATTG ACCACTCCCC TATTGGTGAC GATACTTTCC
3241 ATTACTAATC CATAACATGG CTCTTTGCCA CAACTATCTC TATTGGCTAT ATGCCAATAC
3301 TCTGTCCTTC AGAGACTGAC ACGGACTCTG TATTTTTACA GGATGGGGTC CCATTTATTA
3361 TTTACAAATT CACATATACA ACAACGCCGT CCCCCTGCC CGCAGTTTTT ATTAAACATA
3421 GCGTGGGATC TCCACGCGAA TCTCGGGTAC GTGTTCCGGA CATGGGCTCT TCTCCGGTAG
3481 CGGCGGAGCT TCCACATCCG AGCCCTGGTC CCATGCCTCC AGCGGCTCAT GGTGCTCGG
3541 CAGCTCCTTG CTCCTAACAG TGGAGGCCAG ACTTAGGCAC AGCACAATGC CCACCACCAC
3601 CAGTGTGCCG CACAAGGCCG TGGCGGTAGG GTATGTGTCT GAAAATGAGC TCGGAGATTG

```


FIGURE 1-2

```

3661  GGCTCGCACC GTGACGCAGA TGGAAGACTT AAGGCAGCGG CAGAAGAAGA TGCAGGCAGC
3721  TGAGTTGTTG TATTCTGATA AGAGTCAGAG GTAACCTCCG TTGCGGTGCT GTTAACGGTG
3781  GAGGGCAGTG TAGTCTGAGC AGTACTCGTT GCTGCCGCGC GCGCCACCAG ACATAATAGC
3841  TGACAGACTA ACAGACTGTT CCTTTCCATG GGTCTTTTCT GCAGTCACCG TCCAAGCTTG
3901  CAATCATGGA TGCAATGAAG AGAGGGCTCT GCTGTGTGCT GCTGCTGTGT GGAGCAGTCT
3961  TCGTTTCGGC TAGCAATGAT GATAAGTTAT ATCGGGCAGA TTCTAGACCT CCTGATGAAA
4021  TAAAGCAGTC AGGTGGTCTT ATGCCAAGAG GACAGAGTGA GTACTTTGAC CGAGGTACTC
4081  AAATGAATAT CAACCTTTAT GATCATGCAA GAGGAACCTCA GACGGGATTT GTTAGGCACG
4141  ATGATGGATA TGTTCACACC TCAATTAGTT TGAGAAGTGC CCACTTAGTG GGTCAAACCTA
4201  TATTGTCTGG TCATTCTACT TATTATATAT ATGTTATAGC CACTGCACCC AACATGTTTA
4261  ACGTTAATGA TGTATTAGGG GCATACAGTC CTCATCCAGA TGAACAAGAA GTTTCTGCTT
4321  TAGGTGGGAT TCCATACTCC CAAATATATG GATGGTATCG AGTTCATTTT GGGGTGCTTG
4381  ATGAACAATT ACATCGTAAT AGGGGCTACA GAGATAGATA TTACAGTAAC TTAGATATTG
4441  CTCCAGCAGC AGATGGTTAT GGATTGGCAG GTTTCCTCC GGAGCATAGA GCTTGGAGGG
4501  AAGAGCCGTG GATTCATCAT GCACCGCCGG GTTGTGGGAA TGCTCCAAGA TCATCGATGA
4561  GTAATACTTG CGATGAAAAA ACCCAAAGTC TAGGTGTAAA ATTCCTTGAC GAATACCAAT
4621  CTAAAGTTAA AAGACAAATA TTTTCAGGCT ATCAATCTGA TATTGATACA CATAATAGAA
4681  TTAAGGATGA ATTATGAGGA TCCTCGCAAT CCCTAGGAGG ATTAGGCAAG GGCTTGAGCT
4741  CACGCTCTTG TGAGGGACAG AAATACAATC AGGGGCAGTA TATGAATACT CCATGGAGAA
4801  ACCCAGATCT ACGTATGATC AGCCTCGACT GTGCCTTCTA GTTGCCAGCC ATCTGTTGTT
4861  TGCCCCTCCC CCGTGCCTTC CTTGACCCTG GAAGGTGCCA CTCCCCTGT CTTTCCTAA
4921  TAAAATGAGG AAATTGCATC GCATTGTCTG AGTAGGTGTC ATTCTATTCT GGGGGGTGGG
4981  GTGGGGCAGG ACAGCAAGGG GGAGGATTGG GAAGACAATA GCAGGCATGC TGGGGATGCG
5041  GTGGGGCTCTA TGGCTTCTGA GCGGAAAGA ACCAGCTGGG GCTCGACAGC TCGACTCTAG
5101  AATTCACCTG CCGTCGTTTT ACAACGTCGT GACTGGGAAA ACCCTGGCGT TACCCAACTT
5161  AATCGCCTTG CAGCACATCC CCCTTTCGCC AGCTGGCGTA ATAGCGAAGA GGCCCGCACC
5221  GATCGCCCTT CCAACAGTT GCGCAGCCTG AATGGCGAAT GGCGCCTGAT GCGGTATTTT
5281  CTCCTTACGC ATCTGTGCGG TATTTACAC ACATATGGT GCACTCTCAG TACAATCTGC
5341  TCTGATGCCG CATAGTTAAG CCAGCCCCGA CACCCGCCAA CACCCGCTGA CGCGCCCTGA
5401  CGGGCTTGTC TGCTCCCGGC ATCCGCTTAC AGACAAGCTG TGACCGTCTC CGGGAGCTGC
5461  ATGTGTCAGA GGTTTTCACC GTCATCACCG AAACGCGCGA

```

FIGURE 1-3



Molecule: pPJV2003, 5089 bps DNA Circular
 File Name: pPJV2003.cm5,

Description: Ligation of CTB nhe bam cut frag into 7054 Nhe Bam Vector

Notes:

Molecule Features:

Type	Start	End	Name	Description
REGION	2242	3060	CMVpro	
REGION	3061	3884	intronA	
GENE	3906	3969	TPAsigCDS'	
GENE	3975	4286	CTB CDS	
REGION	4394	4690	bGHpA	

Enzymes (16 sites)

PciI	1876,	SalI	2241,	MscI	2266,	SpeI	2356
SacII	3009,	NsiI	3106,	PstI	3879,	HindIII	3894
NheI	3969,	BamHI	4287,	BglII	4394,	XbaI	4685

FIGURE 2-1

```

1   GACGAAAGGG CCTCGTGATA CGCCTATTTT TATAGGTTAA TGTCATGATA ATAATGGTTT
61  CTTAGACGTC AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTTTATTTT
121 TCTAAATACA TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT
181 AATATTGAAA AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTGCGCCCTT ATTCCCTTTT
241 TTGCGGCATT TTGCCTTCCT GTTTTTGCTC ACCCAGAAAC GCTGGTGAAA GTAAAAGATG
301 CTGAAGATCA GTTGGGTGCA CGAGTGGGTT ACATCGAACT GGATCTCAAC AGCGGTAAGA
361 TCCTTGAGAG TTTTCGCCCC GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCTGC
421 TATGTGGCGC GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC
481 ACTATTCTCA GAATGACTTG GTTGAGTACT CACCAGTCAC AGAAAAGCAT CTTACGGATG
541 GCATGACAGT AAGAGAATTA TGCAGTGCTG CCATAACCAT GAGTGATAAC ACTGCGGCCA
601 ACTTACTTCT GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTTTTG CACAACATGG
661 GGGATCATGT AACTCGCCTT GATCGTTGGG AACCGGAGCT GAATGAAGCC ATACCAAACG
721 ACGAGCGTGA CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAAA CTATTAAC TG
781 GCGAACTACT TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG
841 TTGCAGGACC ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG
901 GAGCCGGTGA GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT
961 CCCGTATCGT AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC
1021 AGATCGCTGA GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT
1081 CATATATACT TTAGATTGAT TTAAAACTTC ATTTTAAATT TAAAAGGATC TAGGTGAAGA
1141 TCCTTTTTTG TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTT CACTGAGCGT
1201 CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTTCTG CGCGTAATCT
1261 GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC
1321 TACCAACTCT TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC
1381 TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACCG CCTACATACC
1441 TCGCTCTGCT AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG
1501 GGTGACTC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGTT
1561 CGTGACACA GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG
1621 AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG
1681 GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT
1741 ATAGTCCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG
1801 GGGGGCGGAG CCTATGGAAA AACGCCAGCA ACGCGGCCTT TTTACGGTTC CTGGCCTTTT
1861 GCTGGCCTTT TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA
1921 TTACCGCCTT TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT
1981 CAGTGAGCGA GGAAGCGGAA GAGCGCCCAA TACGCAAACC GCCTCTCCCC GCGCGTTGGC
2041 CGATTCAATTA ATGCAGCTGG CACGACAGGT TTCCCGACTG GAAAGCGGGC AGTGAGCGCA
2101 ACGCAATTAA TGTGAGTTAG CTCACTCATT AGGCACCCCA GGCTTTACAC TTTATGCTTC
2161 CGGCTCGTAT GTTGTGTGGA ATTGTGAGCG GATAACAATT TCACACAGGA AACAGCTATG
2221 ACCATGATTA CGCCAAGCTA GTCGACATAA ATCAATATTG GCTATTGGCC ATTGCATACG
2281 TTGTATCTAT ATCATAATAT GTACATTTAT ATTGGCTCAT GTCCAATATG ACCGCCATGT
2341 TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCATT AGTTCATAGC
2401 CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCGCGCCTCG TGACCGCCCA
2461 ACGACCCCGG CCCATTGACG TCAATAATGA CGTATGTTCC CATAGTAACG CCAATAGGGA
2521 CTTTCCATTG ACGTCAATGG GTGGAGTATT TACGGTAAAC TGCCCACTTG GCAGTACATC
2581 AAGTGTATCA TATGCCAAGT CCGGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCGCC
2641 TGGCATTATG CCCAGTACAT GACCTTACGG GACTTTCCTA CTTGGCAGTA CATCTACGTA
2701 TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACACCAATGG GCGTGGATAG
2761 CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCCATTG ACGTCAATGG GAGTTTGT TT
2821 TGGCACCAAA ATCAACGGGA CTTTCCAAAA TGTCGTAATA ACCCGCCCC GTTGACGCAA
2881 ATGGGCGGTA GCGGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT
2941 CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA CCGGGACCGA
3001 TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCCGTGC CAAGAGTGAC
3061 GTAAGTACCG CCTATAGACT CTATAGGCAC ACCCCTTTTG CTCTTATGCA TGCTATACTG
3121 TTTTGGCTT GGGGCCTATA CACCCCGCT CTTTATGCTA TAGGTGATGG TATAGCTTAG
3181 CCTATAGGTG TGGGTTATTG ACCATTATTG ACCACTCCCC TATTGGTGAC GATACTTTCC
3241 ATTACTAATC CATAACATGG CTCTTTGCCA CAACTATCTC TATTGGCTAT ATGCCAATAC
3301 TCTGTCCTTC AGAGACTGAC ACGGACTCTG TATTTTTTACA GGATGGGGTC CCATTTATTA
3361 TTTACAAATT CACATATACA ACAACGCCGT CCCCCTGACC CGCAGTTTTT ATTAAACATA
3421 GCGTGGGATC TCCACGCGAA TCTCGGGTAC GTGTTCCGGA CATGGGCTCT TCTCCGGTAG
3481 CGGCGGAGCT TCCACATCCG AGCCCTGGTC CCATGCCTCC AGCGGCTCAT GGTCGCTCGG
3541 CAGCTCCTTG CTCCTAACAG TGGAGGCCAG ACTTAGGCAC AGCACAATGC CCACCACCAC
3601 CAGTGTGCCG CACAAGGCCG TGGCGGTAGG GTATGTGTCT GAAAATGAGC TCGGAGATTG

```


FIGURE 2-2

```

3661  GGCTCGCACC  GTGCGCAGA  TGGAAGACTT  AAGGCAGCGG  CAGAAGAAG  TGCAGGCAGC
3721  TGAGTTGTTG  TATTCTGATA  AGAGTCAGAG  GTAACCTCCG  TTGCGGTGCT  GTTAACGGTG
3781  GAGGGCAGTG  TAGTCTGAGC  AGTACTCGTT  GCTGCCGCGC  GCGCCACCAG  ACATAATAGC
3841  TGACAGACTA  ACAGACTGTT  CCTTTCCATG  GGTCTTTTCT  GCAGTCACCG  TCCAAGCTTG
3901  CAATCATGGA  TGCAATGAAG  AGAGGGCTCT  GCTGTGTGCT  GCTGCTGTGT  GGAGCAGTCT
3961  TCGTTTCGGC  TAGCACACCT  CAAAATATTA  CTGATTTGTG  TGCAGAATAC  CACAACACAC
4021  AAATATATAC  GCTAAATGAT  AAGATATTTT  CGTATACAGA  ATCTCTAGCT  GGAAAAGAG
4081  AGATGGCTAT  CATTACTTTT  AAGAATGGTG  CAATTTTTC  AGTAGAAGTA  CCAGGTAGTC
4141  AACATATAGA  TTCACAAAAA  AAAGCGATTG  AAAGGATGAA  GGATACCCTG  AGGATTGCAT
4201  ATCTTACTGA  AGCTAAAGTC  GAAAAGTTAT  GTGTATGGAA  TAATAAAACG  CCTCATGCGA
4261  TTGCCGCAAT  TAGTATGGCA  AATTAAGGAT  CCTCGCAATC  CCTAGGAGGA  TTAGGCAAGG
4321  GCTTGAGCTC  ACGCTCTTGT  GAGGGACAGA  AATACAATCA  GGGGCAGTAT  ATGAATACTC
4381  CATGGAGAAA  CCCAGATCTA  CGTATGATCA  GCCTCGACTG  TGCCTTCTAG  TTGCCAGCCA
4441  TCTGTTGTTT  GCCCCTCCCC  CGTGCCTTCC  TTGACCCTGG  AAGGTGCCAC  TCCCAGTCTC
4501  CTTTCCTAAT  AAAATGAGGA  AATTGCATCG  CATTGTCTGA  GTAGGTGTCA  TTCTATTCTG
4561  GGGGGTGGGG  TGGGGCAGGA  CAGCAAGGGG  GAGGATTGGG  AAGACAATAG  CAGGCATGCT
4621  GGGGATGCGG  TGGGCTCTAT  GGCTTCTGAG  GCGGAAAGAA  CCAGCTGGGG  CTCGACAGCT
4681  CGACTCTAGA  ATTCACTGGC  CGTCGTTTTA  CAACGTCGTG  ACTGGGAAAA  CCCTGGCGTT
4741  ACCCAACTTA  ATCGCCTTGC  AGCACATCCC  CCTTTCGCCA  GCTGGCGTAA  TAGCGAAGAG
4801  GCGCGCACCG  ATCGCCCTTC  CCAACAGTTG  CGCAGCCTGA  ATGGCGAATG  GCGCCTGATG
4861  CGGTATTTTC  TCCTTACGCA  TCTGTGCGGT  ATTTACACAC  GCATATGGTG  CACTCTCAGT
4921  ACAATCTGCT  CTGATGCCGC  ATAGTTAAGC  CAGCCCCGAC  ACCCGCCAAC  ACCCGCTGAC
4981  GCGCCCTGAC  GGGCTTGTCT  GCTCCCGGCA  TCCGCTTACA  GACAAGCTGT  GACCGTCTCC
5041  GGGAGCTGCA  TGTGTCAGAG  GTTTTCACCG  TCATCACCGA  AACGCGCGA

```

FIGURE 2-3



Molecule: pPJV2006, 5488 bps DNA Circular
 File Name: pPJV2006.cm5,

Description: Ligation of CTA-KDEL PCR Frag cut w/ Nhe Bam into 7054 Nhe Bam Vector

Notes:

Molecule Features:

Type	Start	End	Name	Description
REGION	2242	3060	CMVpro	
REGION	3061	3884	intronA	
GENE	3906	3969	TPAsigCDS'	
GENE	3975	4685	CTA-KDEL CDS	
REGION	4793	5089	bGHpA	

Enzymes (15 sites)

SalI	2241,	MscI	2266,	SpeI	2356,	SacII	3009
NsiI	3106,	PstI	3879,	HindIII	3894,	NheI	3969
ClaI	4553,	BamHI	4686,	BglII	4793,	EcoRI	5088

FIGURE 3-1

```

1   GACGAAAGGG CCTCGTGATA CGCCTATTTT TATAGGTTAA TGTCATGATA ATAATGGTTT
61  CTTAGACGTC AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTTTATTTT
121 TCTAAATACA TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT
181 AATATTGAAA AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTCGCCCTT ATTCCCTTTT
241 TTGCGGCATT TTGCCTTCCT GTTTTTGCTC ACCCAGAAAC GCTGGTGAAA GTAAAAGATG
301 CTGAAGATCA GTTGGGTGCA CGAGTGGGTT ACATCGAACT GGATCTCAAC AGCGGTAAGA
361 TCCTTGAGAG TTTTCGCCCC GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCTGC
421 TATGTGGCGC GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC
481 ACTATTCTCA GAATGACTTG GTTGAGTACT CACCACTCAC AGAAAAGCAT CTTACGGATG
541 GCATGACAGT AAGAGAATTA TGCAGTGCTG CCATAACCAT GAGTGATAAC ACTGCGGCCA
601 ACTTACTTCT GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTTTG CACAACATGG
661 GGGATCATGT AACTCGCCTT GATCGTTGGG AACC GGAGCT GAATGAAGCC ATACCAAACG
721 ACGAGCGTGA CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAA CTATTAACTG
781 GCGAACTACT TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG
841 TTGCAGGACC ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG
901 GAGCCGGTGA GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT
961 CCCGTATCGT AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC
1021 AGATCGCTGA GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT
1081 CATATATACT TTAGATTGAT TTA AAACTTC ATTTTAAATT TAAAAGGATC TAGGTGAAGA
1141 TCCTTTTTGA TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTC CACTGAGCGT
1201 CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT
1261 GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC
1321 TACCAACTCT TTTTCCGAAG GTA ACTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC
1381 TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACC GCTACATACC
1441 TCGCTCTGCT AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG
1501 GGTTGGACTC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGTT
1561 CGTGACACACA GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG
1621 AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG
1681 GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT
1741 ATAGTCCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG
1801 GGGGGCGGAG CCTATGGA AAACGCCAGCA ACGCGGCCTT TTTACGGTTC CTGGCCTTTT
1861 GCTGGCCTTT TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA
1921 TTACCGCCTT TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT
1981 CAGTGAGCGA GGAAGCGGAA GAGCGCCCAA TACGCAAACC GCCTCTCCCC GCGCGTTGGC
2041 CGATTCAATTA ATGCAGCTGG CACGACAGGT TTCCCGACTG GAAAGCGGGC AGTGAGCGCA
2101 ACGCAATTAA TGTGAGTTAG CTCACTCATT AGGCACCCCA GGCTTTACAC TTTATGCTTC
2161 CGGCTCGTAT GTTGTGTGGA ATTGTGAGCG GATAACAATT TCACACAGGA AACAGCTATG
2221 ACCATGATTA CGCCAAGCTA GTCGACATAA ATCAATATTG GCTATTGGCC ATTGCATACG
2281 TTGTATCTAT ATCATAATAT GTACATTTAT ATTGGCTCAT GTCCAATATG ACCGCCATGT
2341 TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCAAT AGTTCATAGC
2401 CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCCCGCCTCG TGACCGCCCA
2461 ACGACCCCGG CCCATTGACG TCAATAATGA CGTATGTTCC CATAGTAACG CCAATAGGGA
2521 CTTTCCATTG ACGTCAATGG GTGGAGTATT TACGGTAAAC TGCCCACTTG GCAGTACATC
2581 AAGTGTATCA TATGCCAAGT CCGGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCGCC
2641 TGGCATTATG CCCAGTACAT GACCTTACGG GACTTTCCTA CTTGGCAGTA CATCTACGTA
2701 TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACACCAATGG GCGTGGATAG
2761 CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCCATTG ACGTCAATGG GAGTTTGTTC
2821 TGGCACCAA ATCAACGGGA CTTTCCAAAA TGTCGTAATA ACCCCGCCCC GTTGACGCAA
2881 ATGGGCGGTA GGCGTGACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT
2941 CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA CCGGGACCGA
3001 TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCGGTGC CAAGAGTGAC
3061 GTAAGTACCG CCTATAGACT CTATAGGCAC ACCCCTTTGG CTCTTATGCA TGCTATACTG
3121 TTTTGGCTT GGGGCCTATA CACCCCGCT CTTATGCTA TAGGTGATGG TATAGCTTAG
3181 CCTATAGGTG TGGGTTATTG ACCATTATTG ACCACTCCCC TATTGGTGAC GATACTTTCC
3241 ATTACTAATC CATAACATGG CTCTTTGCCA CAACTATCTC TATTGGCTAT ATGCCAATAC
3301 TCTGTCCTTC AGAGACTGAC ACGGACTCTG TATTTTACA GGATGGGGTC CCATTTATTA
3361 TTTACAAATT CACATATACA ACAACGCCGT CCCCCGTGCC CGCAGTTTTT ATTAAACATA
3421 GCGTGGGATC TCCACGCGAA TCTCGGGTAC GTGTTCCGGA CATGGGCTCT TCTCCGGTAG
3481 CGGCGGAGCT TCCACATCCG AGCCCTGGTC CCATGCCTCC AGCGGCTCAT GGTCGCTCGG
3541 CAGCTCCTTG CTCCTAACAG TGGAGGCCAG ACTTAGGCAC AGCACAATGC CCACCACCAC
3601 CAGTGTGCCG CACAAGGCCG TGGCGGTAGG GTATGTGTCT GAAAATGAGC TCGGAGATTG

```


FIGURE 3-2

```

3661  GGCTCGCACC GTG|   AGA TGAAGACTT AAGGCAGCGG CAGAA  G|  GCAGGCAGC
3721  TGAGTTGTTG TATTCTGATA AGAGTCAGAG GTAAC TCCCG TTGCGGTGC|  TTAACGGTG
3781  GAGGGCAGTG TAGTCTGAGC AGTACTCGTT GCTGCCGCGC GCGCCACCAG ACATAATAGC
3841  TGACAGACTA ACAGACTGTT CCTTTCCATG GGTCTTTTCT GCAGTCACCG TCCAAGCTTG
3901  CAATCATGGA TGCAATGAAG AGAGGGCTCT GCTGTGTGCT GCTGCTGTGT GGAGCAGTCT
3961  TCGTTTCGGC TAGCAATGAT GATAAGTTAT ATCGGGCAGA TTCTAGACCT CCTGATGAAA
4021  TAAAGCAGTC AGGTGGTCTT ATGCCAAGAG GACAGAGTGA GTACTTTGAC CGAGGTACTC
4081  AAATGAATAT CAACCTTTAT GATCATGCAA GAGGAACTCA GACGGGATTT GTTAGGCACG
4141  ATGATGGATA TGTTTCCACC TCAATTAGTT TGAGAAGTGC CCACTTAGTG GGTCAAAC TA
4201  TATTGTCTGG TCATTCTACT TATTATATAT ATGTTATAGC CACTGCACCC AACATGTTTA
4261  ACGTTAATGA TGTATTAGGG GCATACAGTC CTCATCCAGA TGAACAAGAA GTTCTGCTT
4321  TAGGTGGGAT TCCATACTCC CAAATATATG GATGGTATCG AGTTCATTTT GGGGTGCTTG
4381  ATGAACAATT ACATCGTAAT AGGGGCTACA GAGATAGATA TTACAGTAAC TTAGATATTG
4441  CTCCAGCAGC AGATGGTTAT GGATTGGCAG GTTTCCTCC GGAGCATAGA GCTTGGAGGG
4501  AAGAGCCGTG GATTCATCAT GCACCGCCGG GTTGTGGGAA TGCTCCAAGA TCATCGATGA
4561  GTAATACTTG CGATGAAAAA ACCCAAAGTC TAGGTGTAAA ATTCCTTGAC GAATACCAAT
4621  CTAAAGTTAA AAGACAAATA TTTTCAGGCT ATCAATCTGA TATTGATACA CATAATAGAA
4681  TTTGAGGATC CTCGCAATCC CTAGGAGGAT TAGGCAAGGG CTTGAGCTCA CGCTCTTG TG
4741  AGGGACAGAA ATACAATCAG GGGCAGTATA TGAATACTCC ATGGAGAAAC CCAGATCTAC
4801  GTATGATCAG CCTCGACTGT GCCTTCTAGT TGCCAGCCAT CTGTTGTTTG CCCCTCCCCC
4861  GTGCCTTCCT TGACCCTGGA AGGTGCCACT CCCACTGTCC TTTCCTAATA AAATGAGGAA
4921  ATTGCATCGC ATTGTCTGAG TAGGTGT CAT TCTATTCTGG GGGGTGGGGT GGGGCAGGAC
4981  AGCAAGGGGG AGGATTGGGA AGACAATAGC AGGCATGCTG GGGATGCGGT GGGCTCTATG
5041  GCTTCTGAGG CGGAAAGAAC CAGCTGGGGC TCGACAGCTC GACTCTAGAA TTCCTGGCC
5101  GTCGTTTTAC AACGTCGTGA CTGGGAAAAC CCTGGCGTTA CCCAACTTAA TCGCCTTGCA
5161  GCACATCCCC CTTTCGCCAG CTGGCGTAAT AGCGAAGAGG CCCGCACCGA TCGCCCTTCC
5221  CAACAGTTGC GCAGCCTGAA TGGCGAATGG CGCCTGATGC GGTATTTTCT CCTTACGCAT
5281  CTGTGCGGTA TTTACACCG CATATGGTGC ACTCTCAGTA CAATCTGCTC TGATGCCGCA
5341  TAGTTAAGCC AGCCCCGACA CCCGCCAACA CCCGCTGACG CGCCCTGACG GGCTTGTCTG
5401  CTCCCGGCAT CCGCTTACAG ACAAGCTGTG ACCGTCTCCG GGAGCTGCAT GTGTCAGAGG
5461  TTTTCACCGT CATCACCGAA ACGCGCGA

```

FIGURE 3-3



Molecule: pPJV2004, 5500 bps DNA Circular
 File Name: pPJV2004.cm5,

Description: Ligation of LTA Nhe-Bam Insert into 7054 Nhe Bam Vector

Notes:

Molecule Features:

Type	Start	End	Name	Description
REGION	2242	3060	CMVpro	
REGION	3061	3884	intronA	
GENE	3906	3969	TPAsigCDS'	
GENE	3975	4697	LTA-CDS	
REGION	4805	5101	bGHpA	

Enzymes (13 sites)

PciI	1876,	SalI	2241,	MscI	2266,	SpeI	2356
SacII	3009,	NsiI	3106,	PstI	3879,	NheI	3969
BamHI	4698,	BglII	4805,	EheI	5261,	KasI	5261

FIGURE 4-1

```

1   GACGAAAGGG CCTCGTGATA CGCCTATTTT TATAGGTTAA TGTCATGATA ATAATGGTTT
61  CTTAGACGTC AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTTTATTTT
121 TCTAAATACA TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT
181 AATATTGAAA AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTCGCCCTT ATTCCCTTTT
241 TTGCGGCATT TTGCCTTCCT GTTTTGTGCT ACCCAGAAAC GCTGGTGAAA GTAAAAGATG
301 CTGAAGATCA GTTGGGTGCA CGAGTGGGTT ACATCGAACT GGATCTCAAC AGCGGTAAGA
361 TCCTTGAGAG TTTTCGCCCC GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCCTG
421 TATGTGGCGC GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC
481 ACTATTCTCA GAATGACTTG GTTGAGTACT CACCAGTCAC AGAAAAGCAT CTTACGGATG
541 GCATGACAGT AAGAGAATTA TGCAGTGCTG CCATAACCAT GAGTGATAAC ACTGCGGCCA
601 ACTTACTTCT GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTTTG CACAACATGG
661 GGGATCATGT AACTCGCCTT GATCGTTGGG AACC GGAGCT GAATGAAGCC ATACCAAACG
721 ACGAGCGTGA CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAAA CTATTAAC TG
781 GCGAACTACT TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG
841 TTGCAGGACC ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG
901 GAGCCGGTGA GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT
961 CCCGTATCGT AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC
1021 AGATCGCTGA GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT
1081 CATATATACT TTAGATTGAT TTAAAACTTC ATTTTAAATT TAAAAGGATC TAGGTGAAGA
1141 TCCTTTTTGA TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTT CACTGAGCGT
1201 CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT
1261 GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC
1321 TACCAACTCT TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC
1381 TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACCG CCTACATACC
1441 TCGCTCTGCT AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG
1501 GGTTGGACTC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGTT
1561 CGTGACACAC GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG
1621 AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG
1681 GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT
1741 ATAGTCCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG
1801 GGGGGCGGAG CCTATGGAAA AACGCCAGCA ACGCGGCCTT TTTACGGTTC CTGGCCTTTT
1861 GCTGGCCTTT TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA
1921 TTACCGCCTT TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT
1981 CAGTGAGCGA GGAAGCGGAA GAGCGCCCAA TACGCAAACC GCCTCTCCCC GCGCGTTGGC
2041 CGATTCATTA ATGCAGCTGG CACGACAGGT TTCCCGACTG GAAAGCGGGC AGTGAGCGCA
2101 ACGCAATTAA TGTGAGTTAG CTCACTCATT AGGCACCCCA GGCTTTACAC TTTATGCTTC
2161 CGGCTCGTAT GTTGTGTGGA ATTGTGAGCG GATAACAATT TCACACAGGA AACAGCTATG
2221 ACCATGATTA CGCCAAGCTA GTCGACATAA ATCAATATTG GCTATTGGCC ATTGCATACG
2281 TTGTATCTAT ATCATAATAT GTACATTTAT ATTGGCTCAT GTCCAATATG ACCGCCATGT
2341 TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCAAT AGTTCATAGC
2401 CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCCCGCCTCG TGACCGCCCA
2461 ACGACCCCCG CCCATTGACG TCAATAATGA CGTATGTTCC CATAGTAACG CCAATAGGGA
2521 CTTTCCATTG ACGTCAATGG GTGGAGTATT TACGGTAAAC TGCCCACTTG GCAGTACATC
2581 AAGTGTATCA TATGCCAAGT CCGGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCGCC
2641 TGGCATTATG CCCAGTACAT GACCTTACGG GACTTTCTTA CTTGGCAGTA CATCTACGTA
2701 TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACACCAATGG GCGTGGATAG
2761 CGGTTTGACT CACGGGGGATT TCCAAGTCTC CACCCCATTG ACGTCAATGG GAGTTTGT TT
2821 TGGCACCAAA ATCAACGGGA CTTTCCAAAA TGTCGTAATA ACCCGCCCCC GTTGACGCAA
2881 ATGGGCGGTA GGCGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT
2941 CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA CCGGGACCGA
3001 TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCCGTGC CAAGAGTGAC
3061 GTAAGTACCG CCTATAGACT CTATAGGCAC ACCCCTTTGG CTCTTATGCA TGCTATACTG
3121 TTTTGGGCTT GGGGCCTATA CACCCCGCTT CCTTATGCTA TAGGTGATGG TATAGCTTAG
3181 CCTATAGGTG TGGGTTATTG ACCATTATTG ACCACTCCCC TATTGGTGAC GATACTTTCC
3241 ATTACTAATC CATAACATGG CTCTTTGCCA CAACTATCTC TATTGGCTAT ATGCCAATAC
3301 TCTGTCCTTC AGAGACTGAC ACGGACTCTG TATTTTTACA GGATGGGGTC CCATTTATTA
3361 TTTACAAATT CACATATACA ACAACGCCGT CCCCCGTGCC CGCAGTTTTT ATTAAACATA
3421 GCGTGGGATC TCCACGCGAA TCTCGGGTAC GTGTTCCGGA CATGGGCTCT TCTCCGGTAG
3481 CGGCGGAGCT TCCACATCCG AGCCCTGGTC CCATGCCTCC AGCGGCTCAT GGTGCTCGG
3541 CAGCTCCTTG CTCCTAACAG TGGAGGCCAG ACTTAGGCAC AGCACAATGC CCACCACCAC
3601 CAGTGTGCCG CACAAGGCCG TGGCGGTAGG GTATGTGTCT GAAAATGAGC TCGGAGATTG

```


FIGURE 4-2

```

3661  GGCTCGGACC GTGACGCAGA TGGAAGACTT AAGGCAGCGG CAGAAGAAGA TGCAGGCAGC
3721  TGAGTTGTTG TATTCTGATA AGAGTCAGAG GTAAC TCCCG TTGCGGTGCT GTTAACGGTG
3781  GAGGGCAGTG TAGTCTGAGC AGTACTCGTT GCTGCCGCGC GCGCCACCAG ACATAATAGC
3841  TGACAGACTA ACAGACTGTT CCTTTCCATG GGTCTTTTCT GCAGTCACCG TCCAAGCTTG
3901  CAATCATGGA TGCAATGAAG AGAGGGCTCT GCTGTGTGCT GCTGCTGTGT GGAGCAGTCT
3961  TCGTTTCGGC TAGCAATGGC GACAAATTAT ACCGTGCTGA CTCTAGACCC CCAGATGAAA
4021  TAAAACGTTT CGGAGGTCTT ATGCCCAGAG GGCATAATGA GTACTTCGAT AGAGGAACTC
4081  AAATGAATAT TAATCTTTAT GATCACGCGA GAGGAACACA AACC GGCTTT GTGAGATATG
4141  ATGACGGATA TGTTTCCACT TCTCTTAGTT TGAGAAGTGC TCACTTAGCA GGACAGTCTA
4201  TATTATCAGG ATATTCCACT TACTATATAT ATGTTATAGC GACAGCACCA AATATGTTTA
4261  ATGTTAATGA TGTATTAGGC GTATACAGCC CTCACCCATA TGAACAGGAG GTTTCTGCGT
4321  TAGGTGGAAT ACCATATTCT CAGATATATG GATGGTATCG TGTTAATTTT GGTGTGATTG
4381  ATGAACGATT ACATCGTAAC AGGGAATATA GAGACCGGTA TTACAGAAAT CTGAATATAG
4441  CTCCGGCAGA GGATGGTTAC AGATTAGCAG GTTTCCCACC GGATCACCAA GCTTGAGAG
4501  AAGAACCCTG GATTCATCAT GCACCACAAG GTTGTGGAAT TTCATCAAGA ACAATTACAG
4561  GTGATACTTG TAATGAGGAG ACCCAGAATC TGAGCACAAT ATATCTCAGG AAATATCAAT
4621  CAAAAGTTAA GAGGCAGATA TTTTCAGACT ATCAGTCAGA GGTGACATA TATAACAGAA
4681  TTCGGGATGA ATTATGAGGA TCCTCGCAAT CCCTAGGAGG ATTAGGCAAG GGCTTGAGCT
4741  CACGCTCTTG TGAGGGACAG AAATACAATC AGGGGCAGTA TATGAATACT CCATGGAGAA
4801  ACCCAGATCT ACGTATGATC AGCCTCGACT GTGCCTTCTA GTTGCCAGCC ATCTGTTGTT
4861  TGCCCCTCCC CCGTGCCTTC CTTGACCCTG GAAGGTGCCA CTCCCCTGT CTTTCCTAA
4921  TAAAATGAGG AAATTGCATC GCATTGTCTG AGTAGGTGTC ATTCTATTCT GGGGGGTGGG
4981  GTGGGGCAGG ACAGCAAGGG GGAGGATTGG GAAGACAATA GCAGGCATGC TGGGGATGCG
5041  GTGGGCTCTA TGGCTTCTGA GCGGAAAGA ACCAGCTGGG GCTCGACAGC TCGACTCTAG
5101  AATTCACCTG CCGTCGTTTT ACAACGTCGT GACTGGGAAA ACCCTGGCGT TACCCAACTT
5161  AATCGCCTTG CAGCACATCC CCCTTTCGCC AGCTGGCGTA ATAGCGAAGA GGCCCGCACC
5221  GATCGCCCTT CCCAACAGTT GCGCAGCCTG AATGGCGAAT GGCGCCTGAT GCGGTATTTT
5281  CTCCTTACGC ATCTGTGCGG TATTTACAC CGCATATGGT GCACTCTCAG TACAATCTGC
5341  TCTGATGCCG CATAGTTAAG CCAGCCCCGA CACCCGCCAA CACCCGCTGA CGCGCCCTGA
5401  CGGGCTTGTC TGCTCCCGGC ATCCGCTTAC AGACAAGCTG TGACCGTCTC CGGGAGCTGC
5461  ATGTGTCAGA GGTTTTACAC GTCATCACCG AAACGCGCGA

```

FIGURE 4-3



Molecule: pPJV2005, 5089 bps DNA Circular
 File Name: pPJV2005.cm5,

Description: Ligation of LTB NheBam Frag into 7054 Nhe Bam Vector

Notes:

Molecule Features:

Type	Start	End	Name	Description
REGION	2242	3060	CMVpro	
REGION	3061	3884	intronA	
GENE	3906	3969	TPAsigCDS'	
GENE	3975	4286	LTB CDS	
REGION	4394	4690	bGHpA	

Enzymes (19 sites)

PciI	1876,	Sall	2241,	MscI	2266,	SpeI	2356
SacII	3009,	NsiI	3106,	PstI	3879,	HindIII	3894
NheI	3969,	ClaI	4061,	SmaI	4130,	MunI	4258

FIGURE 5-1

```

1   GACGAAAGGG CCTCGTGATA CGCCTATTTT TATAGGTTAA TGTCATGATA ATAATGGTTT
61  CTTAGACGTC AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTTTATTTT
121 TCTAAATACA TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT
181 AATATTGAAA AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTCGCCCTT ATTCCCTTTT
241 TTGCGGCATT TTGCCTTCCT GTTTTGTCTC ACCCAGAAAC GCTGGTGAAA GTAAAAGATG
301 CTGAAGATCA GTTGGGTGCA CGAGTGGGTT ACATCGAACT GGATCTCAAC AGCGGTAAGA
361 TCCTTGAGAG TTTCGCCCCG GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCTGC
421 TATGTGGCGC GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC
481 ACTATTCTCA GAATGACTTG GTTGAGTACT CACCAGTCAC AGAAAAGCAT CTTACGGATG
541 GCATGACAGT AAGAGAATTA TGCAGTGCTG CCATAACCAT GAGTGATAAC ACTGCGGCCA
601 ACTTACTTCT GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTGTG CACAACATGG
661 GGGATCATGT AACTCGCCTT GATCGTTGGG AACCGGAGCT GAATGAAGCC ATACCAAACG
721 ACGAGCGTGA CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAAA CTATTAAGTG
781 GCGAACTACT TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG
841 TTGCAGGACC ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG
901 GAGCCGGTGA GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT
961 CCCGTATCGT AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC
1021 AGATCGCTGA GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT
1081 CATATATACT TTAGATTGAT TTAAAACTTC ATTTTAAATT TAAAAGGATC TAGGTGAAGA
1141 TCCTTTTTGA TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTTT CACTGAGCGT
1201 CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT
1261 GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC
1321 TACCAACTCT TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC
1381 TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACC GCTACATACC
1441 TCGCTCTGCT AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG
1501 GGTGACTC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGTT
1561 CGTGACACA GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG
1621 AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG
1681 GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT
1741 ATAGTCCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG
1801 GGGGGCGGAG CCTATGGAAA AACGCCAGCA ACGCGGCCTT TTTACGGTTC CTGGCCTTTT
1861 GCTGGCCTTT TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA
1921 TTACCGCCTT TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT
1981 CAGTGAGCGA GGAAGCGGAA GAGCGCCCAA TACGCAAACC GCCTCTCCCC GCGCGTTGGC
2041 CGATTCAATTA ATGCAGCTGG CACGACAGGT TTCCCGACTG GAAAGCGGGC AGTGAGCGCA
2101 ACGCAATTAA TGTGAGTTAG CTCACTCATT AGGCACCCCA GGCTTTACAC TTTATGCTTC
2161 CGGCTCGTAT GTTGTGTGGA ATTGTGAGCG GATAACAATT TCACACAGGA AACAGCTATG
2221 ACCATGATTA CGCCAAGCTA GTCGACATAA ATCAATATTG GCTATTGGCC ATTGCATACG
2281 TTGTATCTAT ATCATAATAT GTACATTTAT ATTGGCTCAT GTCCAATATG ACCGCCATGT
2341 TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCATT AGTTCATAGC
2401 CCATATATGG AGTTCCGCGT TACATAACTT ACGGTAAATG GCCCGCCTCG TGACCGCCCA
2461 ACGACCCCGG CCCATTGACG TCAATAATGA CGTATGTTCC CATAGTAACG CCAATAGGGA
2521 CTTTCCATTG ACGTCAATGG GTGGAGTATT TACGGTAAAC TGCCCACTTG GCAGTACATC
2581 AAGTGTATCA TATGCCAAGT CCGGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCGCC
2641 TGGCATTATG CCCAGTACAT GACCTTACGG GACTTTCCTA CTTGGCAGTA CATCTACGTA
2701 TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACACCAATGG GCGTGGATAG
2761 CGGTTTGA CTACGGGGATT TCCAAGTCTC CACCCCATTTG ACGTCAATGG GAGTTTGTTC
2821 TGGCACCAA ATCAACGGGA CTTTCCAAAA TGTCGTAATA ACCCCGCCCC GTTGACGCAA
2881 ATGGGCGGTA GCGGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT
2941 CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA CCGGGACCGA
3001 TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCGGTGC CAAGAGTGAC
3061 GTAAGTACCG CCTATAGACT CTATAGGCAC ACCCCTTTGG CTCTTATGCA TGCTATACTG
3121 TTTTGGCTT GGGGCCTATA CACCCCGCT CTTTATGCTA TAGGTGATGG TATAGCTTAG
3181 CCTATAGGTG TGGGTTATTG ACCATTATTG ACCACTCCCC TATTGGTGAC GATACTTTCC
3241 ATTACTAATC CATAACATGG CTCTTTGCCA CAACTATCTC TATTGGCTAT ATGCCAATAC
3301 TCTGTCTTTC AGAGACTGAC ACGGACTCTG TATTTTACCA GGATGGGGTC CCATTTATTA
3361 TTTACAAATT CACATATACA ACAACGCCGT CCCCCTGCC CGCAGTTTTT ATTAACATA
3421 GCGTGGGATC TCCACGCGAA TCTCGGGTAC GTGTTCCGGA CATGGGCTCT TCTCCGCTAG
3481 CGGCGGAGCT TCCACATCCG AGCCCTGGTC CCATGCCTCC AGCGGCTCAT GGTCCGCTCG
3541 CAGCTCCTTG CTCCTAACAG TGGAGGCCAG ACTTAGGCAC AGCACAATGC CCACCACCAC
3601 CAGTGTGCCG CACAAGGCCG TGGCGGTAGG GTATGTGTCT GAAAATGAGC TCGGAGATTG

```


FIGURE 5-2

```

3661  GGCTCGCACC GTGACGCAGA TGGAAGACTT AAGGCAGCGG CAGAAGAAAG GCACGCAGC
3721  TGAGTTGTTG TATTCTGATA AGAGTCAGAG GTAACCTCCG TTGCGGTGCT GTTAACGGTG
3781  GAGGGCAGTG TAGTCTGAGC AGTACTCGTT GCTGCCGCGC GCGCCACCAG ACATAATAGC
3841  TGACAGACTA ACAGACTGTT CCTTTCCATG GGTCTTTTCT GCAGTCACCG TCCAAGCTTG
3901  CAATCATGGA TGCAATGAAG AGAGGGCTCT GCTGTGTGCT GCTGCTGTGT GGAGCAGTCT
3961  TCGTTTCGGC TAGCGCTCCC CAGTCTATTA CAGAACTATG TTCGGAATAT CGCAACACAC
4021  AAATATATAC GATAAATGAC AAGATACTAT CATATACGGA ATCGATGGCA GGCAAAAGAG
4081  AAATGGTTAT CATTACATTT AAGAGCGGCG CAACATTTCA GGTCGAAGTC CCGGGCAGTC
4141  AACATATAGA CTCCCAAAAA AAAGCCATTG AAAGGATGAA GGACACATTA AGAATCACAT
4201  ATCTGACCGA GACCAAAATT GATAAATTAT GTGTATGGAA TAATAAAACC CCCAATTCAA
4261  TTGCGGCAAT CAGTATGGAA AACTAGGGAT CCTCGCAATC CCTAGGAGGA TTAGGCAAGG
4321  GCTTGAGCTC ACGCTCTTGT GAGGGACAGA AATACAATCA GGGGCAGTAT ATGAATACTC
4381  CATGGAGAAA CCCAGATCTA CGTATGATCA GCCTCGACTG TGCCTTCTAG TTGCCAGCCA
4441  TCTGTTGTTT GCCCCTCCCC CGTGCCTTCC TTGACCCTGG AAGGTGCCAC TCCCACTGTC
4501  CTTTCCTAAT AAAATGAGGA AATTGCATCG CATTGTCTGA GTAGGTGTCA TTCTATTCTG
4561  GGGGGTGGGG TGGGGCAGGA CAGCAAGGGG GAGGATTGGG AAGACAATAG CAGGCATGCT
4621  GGGGATGCGG TGGGCTCTAT GGCTTCTGAG GCGGAAAGAA CCAGCTGGGG CTCGACAGCT
4681  CGACTCTAGA ATTCACTGGC CGTCGTTTTA CAACGTCGTG ACTGGGAAAA CCCTGGCGTT
4741  ACCCAACTTA ATCGCCTTGC AGCACATCCC CCTTTCGCCA GCTGGCGTAA TAGCGAAGAG
4801  GCCCGCACCG ATCGCCCTTC CCAACAGTTG CGCAGCCTGA ATGGCGAATG GCGCCTGATG
4861  CGGTATTTTC TCCTTACGCA TCTGTGCGGT ATTTACACACC GCATATGGTG CACTCTCAGT
4921  ACAATCTGCT CTGATGCCGC ATAGTTAAGC CAGCCCCGAC ACCCGCCAAC ACCCGCTGAC
4981  GCGCCCTGAC GGGCTTGTCT GCTCCCGGCA TCCGCTTACA GACAAGCTGT GACCGTCTCC
5041  GGGAGCTGCA TGTGTCAGAG GTTTTCACCG TCATCACCGA AACGCGCGA

```

FIGURE 5-3



Molecule: pPJV2007, 5488 bps DNA Circular
File Name: pPJV2007.cm5,

Description: Ligation of LTA-RDEL Nhe Bam insert into 7054 Nhe Bam Vector

Notes :

Molecule Features:

Type	Start	End	Name	Description
REGION	2242	3060	CMVpro	
REGION	3061	3884	intronA	
GENE	3906	3969	TPAsigCDS'	
GENE	3975	4685	LTA-RDEL CDS	
REGION	4793	5089	bGHpA	

Enzymes (14 sites)

PciI	1876,	SalI	2241,	MscI	2266,	SpeI	2356
SacII	3009,	NsiI	3106,	PstI	3879,	NheI	3969
BamHI	4686,	BglII	4793,	EcoRI	5088,	EheI	5249

FIGURE 6-1

```

1   GACGAAAGGG CCTCGTGATA CGCCTATTTT TATAGGTAA TGTCATGATA ATAATGGTTT
61  CTTAGACGTC AGGTGGCACT TTTCGGGGAA ATGTGCGCGG AACCCCTATT TGTTTATTTT
121 TCTAAATACA TTCAAATATG TATCCGCTCA TGAGACAATA ACCCTGATAA ATGCTTCAAT
181 AATATTGAAA AAGGAAGAGT ATGAGTATTC AACATTTCCG TGTCGCCCTT ATTCCTTTT
241 TTGCGGCATT TTGCCTTCCT GTTTTGTCTC ACCCAGAAAC GCTGGTGAAA GTAAAAGATG
301 CTGAAGATCA GTTGGGTGCA CGAGTGGGT ACATCGAACT GGATCTCAAC AGCGGTAAGA
361 TCCTTGAGAG TTTTCGCCCC GAAGAACGTT TTCCAATGAT GAGCACTTTT AAAGTTCTGC
421 TATGTGGCGC GGTATTATCC CGTATTGACG CCGGGCAAGA GCAACTCGGT CGCCGCATAC
481 ACTATTCTCA GAATGACTTG GTTGAGTACT CACCAGTCAC AGAAAAGCAT CTTACGGATG
541 GCATGACAGT AAGAGAATTA TGCAGTGCTG CCATAACCAT GAGTGATAAC ACTGCGGCCA
601 ACTTACTTCT GACAACGATC GGAGGACCGA AGGAGCTAAC CGCTTTTTTG CACAACATGG
661 GGGATCATGT AACTCGCCTT GATCGTTGGG AACCGGAGCT GAATGAAGCC ATACCAAACG
721 ACGAGCGTGA CACCACGATG CCTGTAGCAA TGGCAACAAC GTTGCGCAA CTATTAAGTG
781 GCGAACTACT TACTCTAGCT TCCCGGCAAC AATTAATAGA CTGGATGGAG GCGGATAAAG
841 TTGCAGGACC ACTTCTGCGC TCGGCCCTTC CGGCTGGCTG GTTTATTGCT GATAAATCTG
901 GAGCCGGTGA GCGTGGGTCT CGCGGTATCA TTGCAGCACT GGGGCCAGAT GGTAAGCCCT
961 CCCGTATCGT AGTTATCTAC ACGACGGGGA GTCAGGCAAC TATGGATGAA CGAAATAGAC
1021 AGATCGCTGA GATAGGTGCC TCACTGATTA AGCATTGGTA ACTGTCAGAC CAAGTTTACT
1081 CATATATACT TTAGATTGAT TTAAAACCTC ATTTTAAATT TAAAAGGATC TAGGTGAAGA
1141 TCCTTTTTGA TAATCTCATG ACCAAAATCC CTTAACGTGA GTTTTCGTC CACTGAGCGT
1201 CAGACCCCGT AGAAAAGATC AAAGGATCTT CTTGAGATCC TTTTTTCTG CGCGTAATCT
1261 GCTGCTTGCA AACAAAAAAA CCACCGCTAC CAGCGGTGGT TTGTTTGCCG GATCAAGAGC
1321 TACCAACTCT TTTTCCGAAG GTAAGTGGCT TCAGCAGAGC GCAGATACCA AATACTGTCC
1381 TTCTAGTGTA GCCGTAGTTA GGCCACCACT TCAAGAACTC TGTAGCACC CGTACATACC
1441 TCGCTCTGCT AATCCTGTTA CCAGTGGCTG CTGCCAGTGG CGATAAGTCG TGTCTTACCG
1501 GGTTGGACTC AAGACGATAG TTACCGGATA AGGCGCAGCG GTCGGGCTGA ACGGGGGGT
1561 CGTGACACACA GCCCAGCTTG GAGCGAACGA CCTACACCGA ACTGAGATAC CTACAGCGTG
1621 AGCATTGAGA AAGCGCCACG CTTCCCGAAG GGAGAAAGGC GGACAGGTAT CCGGTAAGCG
1681 GCAGGGTCGG AACAGGAGAG CGCACGAGGG AGCTTCCAGG GGGAAACGCC TGGTATCTTT
1741 ATAGTCTGT CGGGTTTCGC CACCTCTGAC TTGAGCGTCG ATTTTGTGA TGCTCGTCAG
1801 GGGGCGGAG CCTATGGAAA AACGCCAGCA ACGCGCCTT TTTACGGTC CTGGCCTTTT
1861 GCTGGCCTTT TGCTCACATG TTCTTTCCTG CGTTATCCCC TGATTCTGTG GATAACCGTA
1921 TTACCGCCTT TGAGTGAGCT GATACCGCTC GCCGCAGCCG AACGACCGAG CGCAGCGAGT
1981 CAGTGAGCGA GGAAGCGGAA GAGCGCCCAA TACGCAAACC GCCTCTCCCC GCGCGTTGGC
2041 CGATTCAATTA ATGCAGCTGG CACGACAGGT TTCCCGACTG GAAAGCGGGC AGTGAGCGCA
2101 ACGCAATTAA TGTGAGTTAG CTCACTCATT AGGCACCCCA GGCTTTACAC TTTATGCTTC
2161 CGGCTCGTAT GTTGTGTGGA ATTGTGAGCG GATAACAATT TCACACAGGA AACAGCTATG
2221 ACCATGATTA CGCCAAGCTA GTCGACATAA ATCAATATTG GCTATTGGCC ATTGCATACG
2281 TTGTATCTAT ATCATAATAT GTACATTTAT ATTGGCTCAT GTCCAATATG ACCGCCATGT
2341 TGACATTGAT TATTGACTAG TTATTAATAG TAATCAATTA CGGGGTCATT AGTTCATAGC
2401 CCATATATGG AGTTCCCGCT TACATAACTT ACGGTAAATG GCCCGCTCG TGACCGCCCA
2461 ACGACCCCGG CCCATTGACG TCAATAATGA CGTATGTTCC CATAGTAACG CCAATAGGGA
2521 CTTTCCATTG ACGTCAATGG GTGGAGTATT TACGGTAAAC TGCCCACTTG GCAGTACATC
2581 AAGTGTATCA TATGCCAAGT CCGGCCCCCT ATTGACGTCA ATGACGGTAA ATGGCCCCG
2641 TGGCATTATG CCCAGTACAT GACCTTACGG GACTTTCCTA CTTGGCAGTA CATCTACGTA
2701 TTAGTCATCG CTATTACCAT GGTGATGCGG TTTTGGCAGT ACACCAATGG GCGTGGATAG
2761 CGGTTTGACT CACGGGGATT TCCAAGTCTC CACCCCATTG ACGTCAATGG GAGTTTGT
2821 TGGCACCAAA ATCAACGGGA CTTTCCAAAA TGTCGTAATA ACCCGCCCC GTTGACGCAA
2881 ATGGGCGGTA GCGGTGTACG GTGGGAGGTC TATATAAGCA GAGCTCGTTT AGTGAACCGT
2941 CAGATCGCCT GGAGACGCCA TCCACGCTGT TTTGACCTCC ATAGAAGACA CCGGGACCGA
3001 TCCAGCCTCC GCGGCCGGGA ACGGTGCATT GGAACGCGGA TTCCCCGTGC CAAGAGTGAC
3061 GTAAGTACCG CCTATAGACT CTATAGGCAC ACCCCTTTGG CTCTTATGCA TGCTATACTG
3121 TTTTGGCTT GGGGCTTATA CACCCCGCT CTTTATGCTA TAGGTGATGG TATAGCTTAG
3181 CCTATAGGTG TGGGTTATTG ACCATTATTG ACCACTCCCC TATTGGTGAC GATACTTTCC
3241 ATTACTAATC CATAACATGG CTCTTTGCCA CAACTATCTC TATTGGCTAT ATGCCAATAC
3301 TCTGTCCTTC AGAGACTGAC ACGGACTCTG TATTTTTTACA GGATGGGGTC CCATTTATTA
3361 TTTACAAATT CACATATACA ACAACGCCGT CCCCCGTGCC CGCAGTTTTT ATTAAACATA
3421 GCGTGGGATC TCCACGCGAA TCTCGGGTAC GTGTTCCGGA CATGGGCTCT TCTCCGGTAG
3481 CGGCGGAGCT TCCACATCCG AGCCCTGGTC CCATGCCTCC AGCGGCTCAT GGTCGCTCGG
3541 CAGCTCCTTG CTCCTAACAG TGGAGGCCAG ACTTAGGCAC AGCACAATGC CCACCACCAC
3601 CAGTGTGCCG CACAAGGCCG TGGCGGTAGG GTATGTGTCT GAAAATGAGC TCGGAGATTG

```


FIGURE 6-2


```


3661  GGCTCGCACC GTGACGCAGA TGGAAGACTT AAGGCAGCGG CAGAAGAAGA GCAGGCAGC
3721  TGAGTTGTTG TATTCTGATA AGAGTCAGAG GTAAC TCCCG TTGCGGTGCT GTTAACGGTG
3781  GAGGGCAGTG TAGTCTGAGC AGTACTCGTT GCTGCCGCGC GCGCCACCAG ACATAATAGC
3841  TGACAGACTA ACAGACTGTT CCTTTCCATG GGTCTTTTCT GCAGTCACCG TCCAAGCTTG
3901  CAATCATGGA TGCAATGAAG AGAGGGCTCT GCTGTGTGCT GCTGCTGTGT GGAGCAGTCT
3961  TCGTTTCGGC TAGCAATGGC GACAAATTAT ACCGTGCTGA CTCTAGACCC CCAGATGAAA
4021  TAAAACGTTT CGGAGGTCTT ATGCCCAGAT GGCATAATGA G TACTTCGAT AGAGGAACTC
4081  AAATGAATAT TAATCTTTAT GATCACGCGA GAGGAACACA AACC GGCTTT GTCAGATATG
4141  ATGACGGATA TGTTTCCACT TCTCTTAGTT TGAGAAGTGC TCACTTAGCA GGACAGTCTA
4201  TATTATCAGG ATATTCCACT TACTATATAT ATGTTATAGC GACAGCACCA AATATGTTTA
4261  ATGTTAATGA TGTATTAGGC GTATACAGCC CTCACCCATA TGAACAGGAG GTTTCTGCGT
4321  TAGGTGGAAT ACCATATTCT CAGATATATG GATGGTATCG TGTTAATTTT GGTGTGATTG
4381  ATGAACGATT ACATCGTAAC AGGGAATATA GAGACCGGTA TTACAGAAAT CTGAATATAG
4441  CTCCGCGAGA GGATGGTTAC AGATTAGCAG GTTTCCACCC GGATCACCAA GCTTGAGAG
4501  AAGAACCCTG GATTCATCAT GCACCACAAG GTTGTGGAAA TTCATCAAGA ACAATTACAG
4561  GTGATACTTG TAATGAGGAG ACCCAGAATC TGAGCACAAT ATATCTCAGG AAATATCAAT
4621  CAAAAGTTAA GAGGCAGATA TTTTCAGACT ATCAGTCAGA GGTTGACATA TATAACAGAA
4681  TTTGAGGATC CTCGCAATCC CTAGGAGGAT TAGGCAAGGG CTTGAGCTCA CGCTCTTGTG
4741  AGGGACAGAA ATACAATCAG GGGCAGTATA TGAATACTCC ATGGAGAAAC CCAGATCTAC
4801  GTATGATCAG CCTCGACTGT GCCTTCTAGT TGCCAGCCAT CTGTTGTTTG CCCCTCCCCC
4861  GTGCCTTCCT TGACCCTGGA AGGTGCCACT CCCACTGTCC TTTCTAATA AAATGAGGAA
4921  ATTGCATCGC ATTGTCTGAG TAGGTGTCAT TCTATTCTGG GGGGTGGGGT GGGGCAGGAC
4981  AGCAAGGGGG AGGATTGGGA AGACAATAGC AGGCATGCTG GGGATGCGGT GGGCTCTATG
5041  GCTTCTGAGG CGGAAAGAAC CAGCTGGGGC TCGACAGCTC GACTCTAGAA TTCACTGGCC
5101  GTCGTTTTAC AACGTCGTGA CTGGGAAAAC CCTGGCGTTA CCCAACTTAA TCGCCTTGCA
5161  GCACATCCCC CTTTCGCCAG CTGGCGTAAT AGCGAAGAGG CCCGCACCGA TCGCCCTTCC
5221  CAACAGTTGC GCAGCCTGAA TGGCGAATGG CGCCTGATGC GGTATTTTCT CCTTACGCAT
5281  CTGTGCGGTA TTTCACACCG CATATGGTGC ACTCTCAGTA CAATCTGCTC TGATGCCGCA
5341  TAGTTAAGCC AGCCCCGACA CCCGCCAACA CCCGCTGACG CGCCCTGACG GGCTTGTCTG
5401  CTCCCGGCAT CCGCTTACAG ACAAGCTGTG ACCGTCTCCG GGAGCTGCAT GTGTCAGAGG
5461  TTTTCACCGT CATCACCGAA ACGCGCGA


```


FIGURE 6-3

**Figure 7**

**Figure 8**

**Figure 9**

**Figure 10**

**Figure 11**








Figure 12



FIGURES 13A-13D

Protection Against HSV-2 Challenge in Mice




FIGURE 14