



US00PP27480P3

(12) **United States Plant Patent**
Schoone

(10) **Patent No.:** **US PP27,480 P3**

(45) **Date of Patent:** **Dec. 20, 2016**

(54) **PHALAEOPSIS ORCHID PLANT NAMED**
‘SILIDREAM’

(50) Latin Name: *Phalaenopsis hybrida*
Varietal Denomination: **Silidream**

(71) Applicant: **Floricultura**, Heemskerk (NL)

(72) Inventor: **René Schoone**, Assendelft (NL)

(73) Assignee: **Floricultura**, Heemskerk (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/756,559**

(22) Filed: **Sep. 17, 2015**

(65) **Prior Publication Data**

US 2016/0095271 P1 Mar. 31, 2016

Related U.S. Application Data

(60) Provisional application No. 62/055,120, filed on Sep. 25, 2014.

(30) **Foreign Application Priority Data**

Sep. 30, 2014 (NL) PBR OPS1176

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./311**

(58) **Field of Classification Search**
USPC Plt./311
See application file for complete search history.

Primary Examiner — Annette Para

(74) *Attorney, Agent, or Firm* — Foley & Lardner LLP;
Sunit Talapatra

(57) **ABSTRACT**

A new and distinct *Phalaenopsis* plant named ‘Silidream’ particularly characterized by flowers which are white with some yellow/green in the labellum; plants which may be propagated economically and uniformly using tissue culture; plants which produce more than one inflorescence; long and sturdy inflorescences; and relatively short, dark-green foliage.

3 Drawing Sheets

1

Latin name of the genus and species of the plant claimed:
Phalaenopsis hybrida.

Variety denomination: ‘Silidream’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Phalaenopsis* plant, botanically known as *Phalaenopsis* of the Orchidaceae family, and hereinafter referred to by the cultivar name ‘Silidream’.

Phalaenopsis comprises a genus of about 55 species of herbaceous perennials many of which, or the hybrids thereof, are suitable for cultivation in the home or greenhouse. *Phalaenopsis* is predominantly epiphytic or rock-dwelling, and is native to tropical Asia, the Malay Archipelago, and Oceania. The species typically has 2-ranked, fleshy, oblong or elliptic leaves affixed to a short central stem (monopodial growth), which vary in size from 5 to 8 inches to over 2 feet. The leaves may be entirely green or mottled with silver grey.

Phalaenopsis orchids, often referred to as ‘Moth Orchids’ in the horticultural trade, are frequently used to furnish cut flowers for the florist trade or sold as flowering potted-plants for home or interiorscape.

Phalaenopsis produces upright or pendent lateral racemes, often with many showy flowers which open in succession beginning with the lowermost. The flowers possess three sepals and three petals; the lateral ones being alike. The lowermost petal, called the labellum, is three-lobed and is often more brightly-colored than the other flower segments. Flower colors include various shades of pink, white, yellow and red-brown.

2

Phalaenopsis orchids are typically propagated from seeds. Asexual propagation of *Phalaenopsis* is often done from off-shoots which frequently arise from the lower bracts of the inflorescence. The resulting plants are detached from the mother plant and may be planted in a suitable substrate.

The new *Phalaenopsis* ‘Silidream’ is a product of a controlled breeding program conducted by the inventor, René Schoone, in Strengweg, Heemskerk, The Netherlands. The objective of the breeding program was to develop a new *Phalaenopsis* cultivar particularly characterized by its attractive and unique colored flowers, economical propagation via tissue culture, rapid growth, and a plant dimension suitable for packaging and shipping to the market.

The new *Phalaenopsis* ‘Silidream’ originated from a cross made by the inventor in 2002 in Strengweg, Heemskerk, The Netherlands. The female or seed parent is the *Phalaenopsis* cultivar designated ‘Silibama’, unpatented. The male or pollen parent is the *Phalaenopsis* cultivar designated ‘Hawaiian Dream’, unpatented. The new *Phalaenopsis* ‘Silidream’ was discovered and selected by the inventor as a single flowering plant within the progeny of the stated cross in a controlled environment in 2010 in Strengweg, Heemskerk, The Netherlands.

Asexual reproduction of the new *Phalaenopsis* cultivar by tissue culture (mericlone) was first performed in July, 2010 in Cieweg 13, Heemskerk, The Netherlands, and has demonstrated that the combination of characteristics as herein disclosed for the new cultivar are firmly fixed and retained through successive generations of asexual reproduction. The new cultivar asexually reproduces true-to-type.

BRIEF DESCRIPTION OF THE INVENTION

The following traits have been repeatedly observed and are determined to be unique characteristics of ‘Silidream’, which in combination distinguish this *Phalaenopsis* as a new and distinct cultivar:

1. flowers which are white with some yellow/green in the labellum;
2. plant produces more than one inflorescence;
3. plants may be propagated economically and uniformly using tissue culture;
4. inflorescences are long and sturdy; and
5. relatively short, dark-green foliage.

In comparison with the parental cultivars of 'Silidream', the flowers of the female parent 'Silibama' are about 8 cm in size, the flowers of the male parent 'Hawaiian Dream' are about 10.5 cm in size, whereas the flowers of 'Silidream' are about 10 cm in size.

Presently, the commercial cultivar to which 'Silidream' can be meaningfully compared is 'Izzy'. The flowers of 'Silidream' are bigger in size than the flowers of 'Izzy'. Also has 'Silidream' more yellow in the labellum than 'Izzy'. The apex of the lateral sepals of 'Izzy' are more pointed than the apex of the lateral sepals of 'Silidream'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Phalaenopsis* 'Silidream' showing the colors as true as is reasonably possible with colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description, which accurately describe the color of 'Silidream'.

FIG. 1 shows a side view perspective of a typical flowering plant of 'Silidream' in a 12 cm pot, at 16 months of age.

FIG. 2 shows a close-up view of the typical flower of 'Silidream'.

FIG. 3 shows a close-up view of the typical leaves of 'Silidream'.

DETAILED BOTANICAL DESCRIPTION

The new *Phalaenopsis* cultivar 'Silidream' has not been observed under all possible environmental conditions. The phenotype of the new cultivar may vary with variations in environment such as temperature, light intensity, and day length without any change in the genotype of the plant.

The aforementioned photographs, together with the following observations, measurements and values describe plants of 'Silidream' as grown in a greenhouse in Strengweg, Heemskerk, The Netherlands, under conditions which closely approximate those generally used in commercial practice. Initially, the ideal temperature to grow plants of 'Silidream' is 27° C. during the day and at night. Then, during the flowering phase of 'Silidream', the ideal growing temperature is 20-22° C. during the day and 18° C. at night. Light levels for growing 'Silidream' are a minimum of 5,000 lux and a maximum of 10,000 lux. A balanced fertilizer with level of 200 ppm N, 87 ppm P, 168 ppm K is applied. Duration of growth of 'Silidream' from potting size is between 10 and 14 months.

Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.), 2007 edition, except where general colors of ordinary significance are used. Color values were taken under daylight conditions at approximately noon in Zaandammerweg, Assendelft, The Netherlands. The age of the 'Silidream' plants described is 12 months after potting.

Classification:

Botanical.—*Phalaenopsis hybrida*.

Parentage:

Female or seed parent.—*Phalaenopsis* cultivar designated 'Silibama', unpatented.

Male or pollen parent.—*Phalaenopsis* cultivar designated 'Hawaiian Dream', unpatented.

Propagation:

Type.—Tissue culture.

Rooting habit and description.—Fleshy; approximately 3 mm-6 mm wide and greyed/green in color (RHS 190B); freely branching. It takes 12 weeks for plants growing in tissue culture to initiate roots.

Plant:

Size at maturity.—Height (from bottom of pot to highest flower): about 60 cm. Spread: about 55 cm.

Growth habit.—Standard; green leaves (RHS 137A) and a relatively normal raceme.

Vigor.—Moderate.

Crop time.—Following asexual propagation, at about 26 weeks 2 leaves appear; at about 30 weeks 3-4 leaves appear; after a cold treatment of about 4-8 weeks at a temperature of about 19° C. about 1-3 racemes with flowers appear.

Foliage:

Quantity per plant.—About 6-8 leaves are produced before flowering.

Arrangement and attachment.—Half up/horizontal and on two sides.

Overall shape of leaf.—Oval; the tip is blunt and asymmetric.

Texture (upper & underside).—Smooth and leathery.

Pubescence.—None.

Mature leaf length.—About 18 cm.

Mature leaf width.—About 9 cm.

Mature leaf thickness.—About 2 mm.

Mature leaf color.—Upper side: green (RHS 137A). Under side: yellow/green (RHS 146A).

Leaf base.—Acute.

Margin.—Entire.

Venation.—Pattern: parallel. Color of midvein: upper side: green (RHS 137A). under side: green (RHS 146A).

Inflorescence description:

Appearance: Upright to slightly pendant, racemose inflorescence with bilaterally symmetrical flowers that open in succession beginning with the lowermost flower.

Raceme:

Quantity per plant.—About 1 to 3.

Number of flowers per raceme.—About 8 to 15.

Length.—About 65 cm.

Diameter.—About 7 mm.

Peduncle:

Length.—About 20 cm.

Diameter.—About 4 mm.

Strength.—Strong.

Aspect.—Upright.

Texture.—Glabrous and smooth.

Color.—Green (RHS 183A) with brown (RHS 200C).

Buds:

Height (from base to tip).—About 22 mm.

Diameter (at midpoint).—About 18 mm.

Shape.—Oval/egg-shaped with a bump on one side.

Color.—Yellow/green (RHS 145A) with a red/purple haze (RHS 70A).

Orientation.—Same as flowers (forward facing).

Flowering time: For an untreated plant (flowering plant that has not undergone cold-treatment where the plant grows at a temperature of 18° C. to 19° C. for about 4 to 8 weeks after a period of about 30 weeks at a temperature of 25° C.), 1-3 racemes appear with flower buds and flowers. 5
First flowers can be expected approximately 4 to 6 months after planting a plant with a leaf diameter of 3 to 5 cm. Flowers persistent.

Flowering longevity: On the plant: about 4 to 6 months; lastingness of cut flowers: has not been observed.

Fragrance: No fragrance.

Flower:

Rate of opening.—Flowers fully opened about 2 to 3 days after petal and sepal separation.

Orientation at opening.—Slanted upward and outward. 15

Shape.—Typical shape of *Phalaenopsis*; see FIG. 2.

Size (of single bloom).—Height: about 80 mm. Diameter: about 90 mm.

Quantity and arrangement.—Three petals and three sepals that are trimerous, overlapping and arranged in 2 whorls. Petals are more pronounced than sepals. 20

Petals:

Arrangement.—Inner whorl comprises 3 petals: 2 lateral petals and a labellum.

2 *lateral petals.*—Overall shape: broadly ovate and weakly cupped. Apex: oval. Margin: entire and weakly undulate. Base: broadly ovate. Length: about 55 mm. Width: about 43 mm. Texture: Upper surface: smooth and satiny. Under surface: smooth and satiny. Color (when fully opened): upper side: white (RHS NN155D). Under side: white (RHS NN155D). 30

Labellum.—Overall shape: 3-lobed with 2 prominent callosities at central junction of the lateral lobes and base of the midlobe. Lateral lobes of labellum fold upward about the column; the midlobe extends forward and is terminated by 2 filiform appendages at the apex. Lateral lobes of the labellum are ovate in shape while the midlobe is triangular with a bump and a rib on it. Margin: entire and weakly undulate. Apex: oval. Length: about 20 mm. Width (not flattened): about 25 mm. Depth of tube created by lateral lobes of labellum: about 17 mm. Texture: Upper & under surface: smooth and satiny. Color (when fully opened): Mid lobe, upper side: main color is white (RHS NN155D). At the base and on the midvein some green/yellow (RHS 1A) and also green/yellow on the upper edges until the corners. At the base and at the upper edge a little red/purple haze (RHS 71A). Under side: main color is white (RHS NN155D). At the upper edges yellow (RHS 9A) which runs into green/yellow (RHS 1C). Lateral lobes, upper side: main color is white (RHS NN155D). At the lower edges yellow (RHS 9B) with green/yellow (RHS 1C). From the base some red/purple stripes (RHS 50

72A). Under side: main color is white (RHS NN155D) with at the lower edges some yellow (RHS 9C) with a little green/yellow (RHS 1C). Cirrhi: about 20 mm. color: white (RHS NN155D) with yellow tips (RHS 2D). Pestle (Callosities): Length: about 5 mm. Width (not flattened): about 7 mm. Color: main color is yellow (RHS 6A) with red/purple spots (RHS 70A). In the center and on the sides some white (RHS NN155D).

10 Sepals:

Arrangement.—Outer whorl comprises 3 sepals, one dorsal and two lateral sepals.

Overall shape.—Elliptical and weakly cupped.

Margin.—Entire and weakly undulate.

Length.—About 45 mm.

Width.—About 27 mm.

Apex.—Oval.

Texture.—Upper and under surface: smooth and satiny.

Color (when fully opened).—Upper side, dorsal: white (RHS NN155D). Lateral: white (RHS NN155D).

Under side, dorsal: white (RHS NN155D) with a red/purple haze (RHS 70A). Lateral: white (RHS NN155D) with a yellow/green haze (RHS 145C).

Pedicle:

Length.—About 40 mm.

Diameter.—About 4 mm.

Texture.—Glabrous and smooth.

Color.—Starting at the flower yellow/green (RHS 145D), which runs into white (RHS NN155B), then yellow/green (RHS 145C) and close to the raceme yellow/green (RHS 144A).

Reproductive organs:

Arrangement.—The stamens, style and stigmas are fused into a single, short structure called the column, possessing one terminal anther with pollen grains united into a pollinia, which are covered by an anther cap. The stigma is located under the column behind the pollinia. The ovary is inferior with three carpels present. The plant has not produced seed.

Column.—Length: about 8 mm. Diameter: about 6 mm. Color: white (RHS NN155D).

Pollinia.—Quantity: two. Diameter: about 3 mm. Color: yellow/orange (RHS 23A).

Ovary.—Length: about 4 mm. Diameter: about 5 mm. Color: white (RHS NN155D).

Disease/pest resistance/susceptibility: No specific resistance or susceptibility observed.

Temperature tolerance: Tolerant to a low temperature of about 15° C. and to a high temperature about 30° C.

What is claimed is:

1. A new and distinct *Phalaenopsis* plant named 'Sili-dream', as illustrated and described herein.

* * * * *

FIG. 1




FIG. 2




FIG. 3

