

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/653, B27K 3/50 // (A01N 43/653, 59:20, 43:653)		A1	(11) International Publication Number: WO 96/01054
			(43) International Publication Date: 18 January 1996 (18.01.96)
(21) International Application Number: PCT/EP95/02501		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 27 June 1995 (27.06.95)			
(30) Priority Data: 94201898.7 1 July 1994 (01.07.94) EP (34) Countries for which the regional or international application was filed: AT et al. 08/455,418 31 May 1995 (31.05.95) US			
(60) Parent Application or Grant (63) Related by Continuation US 08/455,418 (CIP) Filed on 31 May 1995 (31.05.95)		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): VALCKE, Alex, Raymond, Albert [BE/BE]; Beulk 80, B-2152 Wechelderzande (BE). VAN DER FLAAS, Mark, Arthur, Josepha [BE/BE]; Goorbaan 14B, B-2230 Herselt (BE).			
(54) Title: SYNERGISTIC COMPOSITIONS CONTAINING METCONAZOLE AND ANOTHER TRIAZOLE			
(57) Abstract Synergistic fungicidal compositions containing a fungicidal triazole and metconazole for treating plants or the loci thereof, or for use in wood-preservation or protection of biodegradable materials. Method of treating plants comprising the administration of a fungicidal triazole and metconazole. Method of protecting wood, wood-products and biodegradable materials from fungal attack and destruction.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-


SYNERGISTIC COMPOSITIONS CONTAINING METCONAZOLE AND ANOTHER TRIAZOLE

5 This invention relates to synergistic compositions of metconazole and another triazole for use as fungicidal preservatives in material protection (in particular in wood), as well as for use as agrochemicals in the protection of plants, fruits and seeds. Imidazole and triazole derivatives are of particular interest as fungicidal agents and several such compounds are now widely used for this purpose. Further, combinations comprising
10 two or more such fungicidally active compounds are known from e.g. EP-A-0,237,764, EP-A-0,393,746 and WO-95/00303.

It now has been found that compositions comprising particular ratios of metconazole and another fungicidal triazole exhibit synergistic fungicidal activity.

15 The present invention is concerned with compositions comprising (I) metconazole, a salt, a stereoisomer or a stereoisomeric mixture thereof, and (II) at least one other fungicidal triazole, salt, stereoisomer or stereoisomeric mixture thereof, in quantities producing a mutual synergistic effect, and a carrier.

Metconazole (I) as mentioned hereinabove is the generic name of the compound 5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol,
20 which compound can be represented by the formula



25 This compound, its syntheses as well as its fungicidal properties are described in EP-0,267,778, EP-0,294,222, EP-0,329,397, EP-0,357,404, EP-0,488,395, EP-0,529,976 and EP-0,576,834. Its use in biocidal compositions for the preservation of industrial materials is described in EP-0,341,954.

30 The other fungicidal triazole (II) mentioned hereinabove is preferably selected from :
 azaconazole 1-[[2-(2,4-dichlorophenyl)-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole,
 bromuconazole 1-[4-bromo-2-(2,4-dichlorophenyl)tetrahydrofurfuryl]-1H-1,2,4-triazole,

-2-

- | | | |
|----|----------------|--|
| | cyproconazole | α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1H-1,2,4-triazole-1-ethanol, |
| | difenoconazole | 1-[2-[4-(4-chlorophenoxy)-2-chlorophenyl]-4-methyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole, |
| 5 | epoxiconazole | 1-[3-(2-chlorophenyl)-2-(4-fluorophenyl)oxiran-2-ylmethyl]-1H-1,2,4-triazole |
| | fenbuconazole | 4-(4-chlorophenyl)-2-phenyl-2-(1H-1,2,4-triazol-1-ylmethyl)butyronitrile, |
| | hexaconazole | α -butyl- α -(2,4-dichlorophenyl)-1H-1,2,4-triazole-1-ethanol, |
| 10 | penconazole | 1-[2-(2,4-dichlorophenyl)pentyl]-1H-1,2,4-triazole, |
| | propiconazole | 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole, |
| | tebuconazole | α -[2-(4-chlorophenyl)ethyl]- α -(1,1-dimethylethyl)-1H-1,2,4-triazol-1-ethanol, and |
| 15 | triticonazole | (E)-5-(4-chlorophenyl)methylene)-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol. |

The active ingredients for use in the mixtures or compositions according to the present invention may be used as stereochemical mixtures or as pure stereoisomers. In particular, metconazole may occur as 1,5-cis or 1,5-trans isomers. The 1,5-cis isomer is preferably used in the compositions of the present invention. Or use may be made of stereochemical mixtures containing predominantly (over 50%) the cis isomer.

The active ingredients (I) and (II) may be present in base or in salt form, the latter being obtained by reaction of the base form with an appropriate acid. Appropriate acids comprise, for example, inorganic acids, such as the hydrohalic acids, i.e. hydrofluoric, hydrochloric, hydrobromic and hydroiodic, sulfuric acid, nitric acid, phosphoric acid and the like; or organic acids, such as, for example, acetic, propanoic, hydroxyacetic, 2-oxopropanoic, 2-hydroxypropanoic, ethanedioic, propanedioic, butanedioic, (Z)-2-butenedioic, (E)-2-butenedioic, 2-hydroxybutanedioic, 2,3-dihydroxybutanedioic, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. The aforementioned salts are generally most suitable for preparing compositions for use as agrochemicals.

The term salt form also comprises metal salt complexes which the active ingredients (I) or (II) may form. These metal salt complexes are generally most suitable for preparing compositions for use in material protection, in particular in wood protection. One of the

-3-

ingredients may occur as a complex and the other not; or both ingredients may occur as a complex. Metal salt complexes as mentioned above consist of a complex formed between one or more active ingredients and one or more organic or inorganic metal salts.

Examples of organic or inorganic metal salts comprise the halogenides, nitrates, sulfates,

- 5 carbonates, hydroxides, oxides, borates, phosphates, acetates, trifluoroacetates, trichloroacetates, propionates, tartrates, sulfonates, e.g. methanesulfonates, 4-methylbenzenesulfonates, salicylates, benzoates and the like, of the metals of the second main group of the periodical system, e.g. magnesium or calcium, metals of the third or fourth main group, e.g. aluminum, tin, lead and the like metals, as well as
- 10 metals of the first to the eighth transition groups of the periodical system such as, for example, chromium, manganese, iron, cobalt, nickel, copper, zinc, silver, cadmium, antimony, mercury, bismuth and the like. Preferred are the metals pertaining to the transition elements of the fourth period, in particular copper and zinc, either used alone, in combination with each other or with one or more of the metals listed above. The metal
- 15 ions may be present in any of their possible valences, the most preferred metal copper being most advantageously used in its divalent form Cu(II). Suitable copper compounds are copper sulfate, acetate, hydroxide, oxide, borate, fluoride and in particular copper hydroxide carbonate $\text{Cu}(\text{OH})_2\text{CuCO}_3$. The ratio (w/w) of the amount of copper to the total amount of triazoles (metconazole (I) + the other triazole (II)) in preferred
- 20 compositions for wood-protection ranges from about 1 : 1 to about 20 : 1, and in particular is about 2 : 1 to about 5 : 1. The complexes can be mono- or polynuclear and may contain one or more parts of the active ingredients as ligands.

The metal salt complexes of the active ingredients can conveniently be prepared by dissolving the metal salt in suitable solvent e.g. ethanol, and adding thereto the active

25 ingredients. The thus obtained complexes may be isolated following art-known techniques, e.g. by filtration or evaporation, and may be further purified, e.g. by recrystallization. The term salt as used hereinabove also comprises the solvates which metconazole and the other fungicidal triazoles are able to form, e.g. hydrates, alcoholates and the like.

30 The synergistic mixtures according to the present invention are most useful to combat fungi or prevent the growth thereof in plants or the loci thereof; particularly in plant products, including wood; in pulpwood for paper manufacture; and also in biodegradable materials such as, for example, textiles of natural fibers, e.g. cotton, flax,

35 hemp, wool, silk and the like; textiles of synthetic fibers, e.g. polyamide, polyacrylonitrile or polyester fibers, or of mixtures of such fibers; coatings, e.g. oil paints, dispersion paints, lacquers, lacquer films, whitewash, finishing stains and the like; glues

-4-

and other such materials which are biodegradable by fungi.

The synergistic mixtures of the present invention are active against a broad range of fungi. As examples of such fungi there may be named Ascomycetes (e.g. Venturia, Podosphaera, Erysiphe, Monilinia, Uncinula, Aureobasidium, Sclerophoma);

- 5 Basidiomycetes (e.g. Hemileia, Rhizoctonia, Puccinia, Coniophora, Serpula, Poria, Uromyces, Gloeophyllum, Lentinus, Coriolus, Irpex); Fungi imperfecti (e.g. Botrytis, Helminthosporium, Rhynchosporium, Fusarium, Septoria, Cercospora, Alternaria, Pyricularia, Penicillium, Geotrichum).
- 10 Firstly, the synergistic mixtures according to the present invention possess advantageous curative, preventive and systemic fungicidal activity to protect plants, in particular culture plants. The present mixtures can be used to protect plants or parts of plants, e.g. fruits, blossoms, flowers, foliage, stems, roots, tubers of plants or culture plants infected, harmed or destroyed by micro-organisms, whereby later-growing parts of plants are
- 15 protected against such micro-organisms. They can further be used in seed disinfection (fruits, tubers, cereal grains), to treat plant cuttings as well as to combat phytopathogenous fungi occurring in the soil. The mixtures of the present invention are particularly attractive due to their good plant tolerance and lack of environmental problems (low application rates).
- 20 As examples of the wide variety of culture plants in which the combinations of active ingredients according to the present invention can be used, there may be named for example cereals, e.g. wheat, barley, rye, oats, rice, sorghum and the like ; beets, e.g. sugar beet and fodder beet ; pome and stone fruits and berries, e.g. apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries ;
- 25 leguminous plants, e.g. beans, lentils, peas, soy beans ; oleaginous plants, e.g. rape, mustard, poppy, olive, sunflower, coconut, castor-oil plant, cocoa, ground-nuts ; cucurbitaceae, e.g. pumpkins, gherkins, melons, cucumbers, squashes ; fibrous plants, e.g. cotton, flax, hemp, jute ; citrus fruits, e.g. orange, lemon, grapefruit, mandarin ;
- 30 vegetables, e.g. spinach, lettuce, asparagus, brassicaceae such as cabbages and turnips, carrots, onions, tomatoes, potatoes, hot and sweet peppers ; laurel-like plants, e.g. avocado, cinnamon, camphor tree ; or plants such as maize, tobacco, nuts, coffee, sugar-cane, tea, vines, hops, bananas, rubber plants, as well as ornamental plants, e.g. flowers, shrubs, deciduous trees and evergreen trees such as conifers. This enumeration of culture plants is given with the purpose of illustrating the invention and not of
- 35 delimiting it thereto.

A particular mode of administering a synergistic composition containing the active

-5-

ingredients (I) and (II), is the administration to the aboveground parts of plants, in particular to the leaves thereof (leaf-application). The number of applications and the administered doses are chosen in accordance with the biological and climatic conditions of life of the causative agent. The active ingredients can also be applied to the soil and get
5 into the plants through the root system (systemic activity), in which case the locus of the plants can be sprayed with a liquid composition; or the compounds can be added to the soil in a solid formulation e.g. in the form of a granulate (soil application). The compounds (I) and (II) can be coated on seeds by drenching them either consecutively with a coating composition of each of the active ingredients individually, or alternatively
10 by drenching the seeds with a coating composition comprising both active ingredients.

The synergistic mixtures according to the present invention are also useful as wood-preserving agents both against wood-destroying and wood-discoloring fungi. As wood which can be preserved with the synergistic compositions according to the present
15 invention is considered, for example, wood products such as timber, lumber, railway sleepers, telephone poles, fences, wood coverings, wicker-work, windows and doors, plywood, particle board, waferboards, chipboard, joinery, bridges or wood products which are generally used in housebuilding, construction and carpentry. Synergistic compositions according to the present invention can also advantageously be applied in
20 the cellulose and paper industry, in particular to protect pulpwood for paper manufacture from fungal attack.

Wood which is preserved from staining, discoloring and decay is meant to be protected from for example, moulding, rotting, loss of its useful mechanical properties such as
25 breaking strength, resistance to shock and shearing strength, or decrease of its optical or other useful properties due to the occurrence of odour, staining and spot formation. These phenomena are caused by a number of micro-organisms of which the following are typical examples :

Wood-discoloring fungi :

- 30 1: Ascomycetes : Ceratocystis e.g. Ceratocystis minor.
Aureobasidium e.g. Aureobasidium pullulans
Sclerophoma e.g. Sclerophoma pithyophila
Cladosporium e.g. Cladosporium herbarum
2: Deuteromycetes: Fungi imperfecti
35 Aspergillus e.g. Aspergillus niger
Dactylium e.g. Dactylium fusarioides
Penicillium e.g. P. brevicaulis, P. variabile, P. funiculosum or P.

-6-

glaucumScopularia e.g. Scopularia phycomycesTrichoderma e.g. Trichoderma viride or Trichoderma lignorum.Alternaria e.g. Alternaria tenius, Alternaria alternata5 3: Zygomycetes: Mucor e.g. Mucor spinorus.

Wood-destroying fungi

1: Soft-rot Fungi: Chaetomium e.g. Ch. globosum or Ch. alba-arenulumHumicola e.g. Humicola grisea10 Petriella e.g. Petriella setiferaTrichurus e.g. Trichurus spiralis.

2: White and brown rot Fungi :

Coniophora e.g. Coniophora puteanaCoriolus e.g. Coriolus versicolor15 Donkioporia e.g. Donkioporia expansaGlenospora e.g. Glenospora graphiiGloeophyllum e.g. Gl. abietinum, Gl. adoratum, Gl. protactum,Gl. sepiarium or Gl. trabeumLentinus e.g. L. cyathiformes, L. edodes, L. lepideus, L. grinus20 or L. squarroloususPaxillus e.g. Paxillus panuoidesPleurotus e.g. Pleurotus ostreatusPoria e.g. P. monticola, P. placenta, P. vaillantii or P. vaporariaSerpula (Merulius) e.g. Serpula himantoides or Serpula lacrymans25 Stereum e.g. Stereum hirsutumTrychophyton e.g. Trychophyton mentagrophytesTyromyces e.g. Tyromyces palustris.

30 In order to protect wood from decay it is treated with synergistic compositions according to the present invention. Such treatment is applied by several different procedures such as, for example, by treating the wood in closed pressure or vacuum systems, in thermal or dip systems and the like, or by a wide variety of surface treatments, e.g. by brushing, dipping, spraying or soaking the wood with a formulation containing the wood-preserving agents metconazole and a fungicidal triazole.

35

The amount of each of the active ingredients metconazole (I) and the fungicidal triazole (II) in the compositions according to the present invention is such that a mutual

-7-

synergistic fungicidal effect is obtained upon application. In particular, it is contemplated that in the compositions to be used directly, the concentration of metconazole taken as base equivalent, may range from 10 to 15000 ppm, in particular from 50 to 12000 ppm or from 50 to 6000 ppm, more in particular from 100 to 3000 ppm; and the concentration of the other azole taken as base equivalent is contemplated to range from 10 to 15000 ppm, in particular from 50 to 10000 ppm or from 100 to 8000 ppm, more in particular from 200 to 6000 ppm. In many instances said compositions to be used directly can be obtained from concentrates upon dilution with aqueous or organic media, such concentrates also being intended to be covered by the term composition as used in the definitions of the present invention. The content of the active ingredients in the above-indicated compositions is from 0.01 to 95%, preferably from 0.1 to 50% more preferably from 0.1 to 20% and in particular from 0.2 to 15% by weight. The compositions according to the invention are preferably used in the form of solutions or emulsions.

The ratio between the active ingredients of formula (I) and (II) in said synergistic compositions may vary within relatively broad ranges and will be dependent on the nature of the active ingredient (II). However, said ratio will be such that a mutual synergistic fungicidal effect is obtained. Particularly, it is contemplated that the weight ratio between the active ingredients (I) and (II) (metconazole : the other triazole) may be situated between 50:1 and 1:50, more particularly between 20:1 and 1:20. Preferably said ratio is between 10:1 and 1:10, more preferably between 5:1 and 1:5.

The active ingredients of formula (I) and (II) are used in unmodified form or together with adjuvants conventionally employed in the art of formulation. The formulations, i.e. the compositions, preparations or mixtures containing the active ingredients and, where appropriate, a solid or liquid adjuvant, are prepared following art-known procedures, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, e.g. solvents, solid carriers and, where appropriate, surface-active compounds (surfactants), to emulsifiable concentrates, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations in e.g. polymer substances. Both the nature of the compositions and the methods of application such as spraying, atomizing, dusting, scattering or pouring, brushing, dipping, soaking or impregnating, should be chosen in accordance with the intended objectives and the prevailing circumstances. It is evident that in general different compositions with different characteristics will be required for use in plant protection on the one hand, and for use in material protection on the other. Carriers and

adjuvants equally useful in both types of compositions are described first.

Appropriate carriers and adjuvants for use in the compositions of the present invention may be solid or liquid and correspond to suitable substances known in the art for preparing formulations for treating plants or the loci thereof, or for treating plant products, in particular for treating wood, such as, for example, natural or regenerated mineral substances, solvents, dispersants, surfactants, wetting agents, adhesives, thickeners, binders, fertilizers, anti-freeze agents, repellents, colour additives, corrosion inhibitors, water-repelling agents, siccatives, UV-stabilizers and other active ingredients.

10

The solid carriers used e.g. for dusts and dispersible powders are normally natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgite. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated absorbent carriers are of the porous type, for example pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are materials such as calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverised plant residues.

15

20

Suitable solvents are aromatic hydrocarbons, preferably the fractions containing 8 to 12 carbon atoms, e.g. dimethylbenzene mixtures or substituted naphthalenes, phthalates such as dibutyl phthalate or dioctyl phthalate, aliphatic or alicyclic hydrocarbons such as cyclohexane or paraffins, alcohols and glycols and their ethers and esters, such as ethanol, ethylene glycol, ethylene glycol monomethyl or monoethyl ether, ketones such as cyclohexanone, strongly polar solvents such as N-methyl-2-pyrrolidone, dimethylsulfoxide or dimethylformamide, as well as vegetable oils or epoxidised vegetable oils such as epoxidised coconut oil or soybean oil; or water, or a mixture of said solvents.

25

30

Suitable surface-active compounds to be used in the compositions of the present invention are non-ionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The term "surfactants" will also be understood as comprising mixtures of surfactants.

35

Suitable anionic surfactants can be both water-soluble soaps and water-soluble synthetic surface-active compounds.

Suitable soaps are the alkali metal salts, earth alkaline metal salts or unsubstituted or

substituted ammonium salts of higher fatty acids (C₁₀-C₂₂), e.g. the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures which can be obtained e.g. from coconut oil or tallow oil. In addition, there may also be mentioned fatty acid methyltaurin salts.

- 5 More frequently, however, so-called synthetic surfactants are used, especially fatty sulfonates, fatty sulfates, sulfonated benzimidazole derivatives or alkylaryl-sulfonates. The fatty sulfonates or sulfates are usually in the form of alkali metal salts, earth alkaline metal salts or unsubstituted or substituted ammonium salts and contain an alkyl radical having from 8 to 22 carbon atoms said alkyl also comprising radicals derived from acyl radicals, e.g. the sodium or calcium salt of lignosulfonic acid, of dodecylsulfate or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfuric acid esters and sulfonic acids of fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain 2 sulfonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulfonic acid, dibutyl-naphthalenesulfonic acid, or of a naphthalenesulfonic acid/formaldehyde condensation product. Also suitable are corresponding phosphates, e.g. salts of the phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 moles of ethylene oxide, or phospholipids.

- 20 Non-ionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, or saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 10 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols.

- 25 Further suitable non-ionic surfactants are the water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit. Representative examples of non-ionic surfactants are nonylphenol polyethoxy-ethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxy-polyethoxyethanol, polyethylene glycol and octylphenoxypolyethoxyethanol. Fatty acid esters of polyethylene sorbitan, such as polyoxyethylene sorbitan trioleate, are also suitable non-ionic surfactants. They are of particular utility in compositions for material, in particular wood, protection.
- 35

-10-

Cationic surfactants are preferably quaternary ammonium salts wherein at least one N-substituent is a C₈-C₂₂alkyl radical and the further substituents are unsubstituted or halogenated lower alkyl, benzyl or hydroxylower alkyl radicals. The salts are preferably in the form of halides, methylsulfates or ethylsulfates, e.g. stearyltrimethylammonium chloride or benzyldi(2-chloroethyl)ethylammonium bromide. They are particularly useful in compositions for agrochemical purposes.

The surfactants customarily employed in the art of formulation are described e.g. in the following publications : "McCutcheon's Detergents and Emulsifiers Annual", MC Publishing Corp., Ridgewood, New Jersey, 1981; H. Stache, "Tensid-Taschenbuch", 2nd Edition, C. Hanser Verlag, Munich & Vienna, 1981, M. and J. Ash, "Encyclopedia of Surfactants", Vol. I-III, Chemical Publishing Co., New York, 1980-81.

The following paragraphs in particular relate to compositions for use in plant protection. In general such compositions are designed so as to be innocuous to culture plants, to be easily and safely applicable, to have good bioavailability to the host plants and to remain (persist) only temporarily in the environment.

Particularly advantageous additives useful to improve the application and reduce the dose of the active ingredients, are the natural (animal or plant) or synthetic phospholipids of the cephalin or lecithin type such as, for example, phosphatidyl-ethanolamine, phosphatidyl serine, phosphatidylglycerine, lysolecithin, or cardiolipin. Such phospholipids may be obtained from animal or plant cells, in particular from brain-, heart- or liver tissue, egg yolks or soy beans. Appropriate such phospholipids are for instance, phosphatidylcholine mixtures. Synthetic phospholipids are for instance, dioctanylphosphatidylcholine and dipalmitoylphosphatidylcholine.

In case of liquid formulations, and particularly of aqueous or alcoholic formulations, it is recommendable to add an appropriate surfactant, either from the anionic, cationic, nonionic or amphoteric type. In particular said surfactants will be of the cationic type and more in particular said surfactant is a quaternary ammonium salt or a mixture of quaternary ammonium salts. Such quaternary ammonium surfactants comprise, for example, ammonium salts having four hydrocarbon radicals which may optionally be substituted with halo, phenyl, substituted phenyl or hydroxy; said hydrocarbon radicals in particular being alkyl or alkenyl radicals; they may also be derived from fatty acids or alcohols, e.g. cetyl, lauryl, palmityl, myristyl, oleyl and the like or from the hydrosylates from coconut oil, tallow oil, soy bean oil, or the hydrogenated forms

-11-

thereof, and the like.

Examples of such quaternary ammonium salts are of the trimethyl alkyl ammonium halide type, e.g. trimethyl decyl ammonium chloride, trimethyl dodecylammonium chloride, trimethyl tallow ammonium chloride, trimethyl oleyl ammonium chloride; or of
5 the dimethyl alkyl benzyl ammonium type, e.g. dimethyl decyl benzyl ammonium chloride, dimethyldodecyl benzyl ammonium chloride, dimethyl hexadecylbenzyl ammonium chloride (commonly designated as "cetalkonium chloride"), dimethyl octadecyl benzyl ammonium chloride, dimethyl coco benzyl ammonium chloride, dimethyl tallow benzyl ammonium chloride; and particularly the dimethyl C₈₋₁₈alkyl
10 benzyl ammonium chloride mixture which is commonly known as "benzalkonium chloride"; dimethyl dialkyl ammonium halides, e.g. dimethyl dioctyl ammonium chloride, dimethyl didecyl ammonium chloride, dimethyl didodecyl ammonium chloride, dimethyl dicoco ammonium chloride, dimethyl ditallow ammonium chloride, dimethyl octyl decyl ammonium chloride, dimethyl dodecyl octyl ammonium chloride, dimethyl
15 dihydrogenated tallow ammonium chloride.

As used in the foregoing enumeration of quaternary ammonium salts, the terms "coco", "tallow" and "hydrogenated tallow" designate those hydrocarbon radicals derived from the hydrosylates of coconut oil, tallow oil or hydrogenated tallow oil.

The weight ratio between said quaternary ammonium surfactants and the active
20 ingredient (I) is situated between 1:1 and 10:1. Excellent results are obtained when said ratio is about 5:1.

The following paragraphs in particular relate to compositions for use in wood protection. In general such compositions are designed so as to penetrate well in to wood, to persist
25 there for a long time, and to be industrially applicable.

A biocidally active quaternary ammonium compound or tertiary amine salt can advantageously be used in the formulation of emulsions of triazole compounds in aqueous solutions of metal salts. Micro-emulsions may thus be formed which have
30 particular utility in wood preservation. Additional advantages related to these adjuvants comprise their solubilizing effect on the triazole compounds, their contributory biocidal effect and their ability to promote penetration of the formulation into wood.

Binders are meant to comprise binding drying oils (e.g. linseed oil) and resins that are
35 water-dilutable or dilutable, dispersable or emulsifiable in organic solvents, e.g. acryl, vinyl, polyester, polyurethane, alkyd, phenolic, hydrocarbon and silicon resins. Mixtures of an alkyd resin with a drying oil are advantageously used as a binding

-12-

material. Part of the binding material may further be substituted with one or more fixing agents or one or more plasticizers. These adjuvants delay or prevent evaporation of the active ingredients as well precipitation or crystallisation thereof. About 0.01% to about 30 % of the binding material may thus be replaced. Suitable plasticizers are phthalic acid esters, e.g. the dibutyl, dioctyl and benzylbutyl phthalate esters; phosphoric acid esters, e.g. tributylphosphate; fatty acid esters, e.g. di(2-ethylhexyl)adipate, butylstearate, amylstearate, butyloleate; glycerolethers; glycolethers; glycerolesters; and p-toluenesulfonic acid esters. Suitable fixing agents are polyvinylalkyl ethers, e.g. polyvinylmethyl ethers, or ketones, e.g. benzophenone or ethylenebenzophenone.

10

In view of their solubility in organic solvents the active ingredients are well suited for application in non-aqueous media, which is of interest in wood-preservation. The wood or wood products to be protected can easily be impregnated with such solutions. As organic solvents there may be used aliphatic and aromatic hydrocarbons, their chlorinated derivatives, acid amides, mineral oils, alcohols, ethers, glycolethers, such as, for example, methylene chloride, propylene glycol, methoxyethanol, ethoxyethanol, N,N-dimethylform-amide and the like or mixtures of such solvents, to which there may be added dispersants or emulsifiers such as sulfated ricinus oil, fatty alcohol sulfates and other additives.

20

Particularly attractive formulations comprise water-dilutable wood-preservative liquids containing an appropriate amount of a suitable solvent, a suitable solubilizer and both the active ingredients. Preferably there is used 10-80% of a solvent, 20-80% of a solubilizer and from 0.01 to 10% of the active ingredients (I) and (II).

25 Preferred solubilizers to be used in the said water-dilutable wood-preservative liquids are selected from :

- i) addition products of 1 to 60 moles of ethylene oxide with 1 mole of a phenol which is further substituted with at least one C₁₋₁₅ alkyl group; and
- ii) addition products of 1 to 60 moles of ethylene oxide with 1 mole of ricinus oil.

30

The most preferred solubilizers are selected from :

- i) addition products of 1 to 60 moles of ethylene oxide with 1 mole of nonylphenol or octylphenol; and
- ii) addition products of 1 to 60 moles of ethylene oxide with 1 mole of ricinus oil.

35

Said suitable solvent should fulfil the requirements of sufficiently solubilizing the active ingredients and, when combined with the solubilizer, of being homogeneously miscible

-13-

with a predominantly aqueous medium. Preferred solvents are 2-butoxyethanol, butyl 2-hydroxyacetic acid ester and propyleneglycol monomethylether.

Preferred water-dilutable compositions for use in wood protection comprise (I)
5 metconazole, the other fungicidal triazole (II), a copper compound, suitable solvents
and/or solubility enhancers and optionally other adjuvants. Suitable solvents are e.g.
alcohols (ethanol, iso-propanol), glycols (ethylene and propylene glycol), glycolethers
(ethylene glycol monomethyl and monoethyl ether), dimethylformamide, n-
methylpyrrolidone, which yield homogenous concentrates. As solubilizers there may be
10 employed carboxylic acids, or the amine, alkali metal or copper salt forms corresponding
thereto, so that the amount of the organic solvent used in the homogenous concentrates
can be kept to a minimum. Examples of such acids are propionic, hexanoic, heptanoic,
2-ethylhexanoic, iso-octanoic, sebacic, cyclohexanoic, benzoic, 3-hydroxybenzoic and
4-hydroxybenzoic acid. To improve further the industrial applicability of such water-
15 dilutable, homogenous compositions comprising a carboxylic acid as solubilizer,
advantageous use may be made of polyethyleneimines (PEI, Polymin) derived from
ethyleneimine (aziridine) and having the formula $(C_2H_5N)_n$. The degree of
polymerization 'n' should be larger than 10 and preferably ranges from about 50 to about
1000, and in particular is about 150. The use of alkanolamines, in particular
20 monoethanolamine, but also di- and triethanolamine, as complexing agents for the
copper compound employed may be of particular benefit. Typically there may be used
about 4 molar equivalents of alkanolamine per mole of copper. Further useful additives
are, for example, boron derivatives, e.g. boric acid, its salts and esters, and fluorides,
e.g. potassium fluoride.

25

Water-dilutable homogenous concentrates in particular comprise by weight :

2.5 to 45 %, in particular 10 to 20 % copper compound,

5 to 50 %, in particular 20 to 40 % alkanolamine,

0.25 to 15 %, in particular 1 to 10 % triazoles (I + II),

30 0.5 to 30 %, in particular 5 to 15 % surfactant(s),

0 to 40 % other fungicidal compound(s),

0 to 40 % organic solvent(s),

0 to 40 % carboxylic acid(s), and

0 to 40 % polymin.

35

Said water-dilutable wood-preserving liquids have the advantage that almost
instantaneously homogeneous or quasi homogeneous solutions are formed by mixing

-14-

these liquids with predominantly aqueous media. These solutions have an extremely high physical stability, not only at ambient temperature, i.e. at temperatures comprised between 15°C and 35°C, but also at decreased temperatures. Thus, the physical stability of said solutions does not deteriorate after several freeze-thaw cycles. Said

5 homogeneous solutions further unite the advantages of moistening the wood-surface well and penetrating the wood to a high degree, resulting in a high uptake of the solution and its active ingredients by the wood, and, consequently, obtaining the desired preservation of the treated wood. Additionally, due to a more uniform uptake of the aqueous solution the wood-preserving liquids and the resulting aqueous solutions are

10 particularly useful in treatment techniques which require the possibility of a continuous process, such as, for example, impregnation or dip techniques.

In addition, the solutions formed with the wood-preserving liquids unite in themselves the hereinabove mentioned advantages with those which are characteristic of predominantly aqueous media, such as, for example, a relatively high flashpoint and

15 reduced toxicity, resulting in advantageous influence on the environment and the health and safety of the applicator, lack of irritation and the like benefits.

In the wood-preserving solutions which are used to be contacted with the wood, said solutions either being a composition as described hereinabove or prepared therefrom

20 upon dilution with a suitable solvent, the concentration of metconazole may vary between 100 and 10000 ppm, in particular between 200 and 5000 ppm and preferably between 500 and 1000 ppm; the concentration of the compound of formula (II) may vary between 100 and 15000 ppm, in particular between 300 and 7500 ppm and preferably between 750 and 1500 ppm.

25 In said wood-preserving solutions, the ratio between the active ingredients (I) and (II) (metconazole / a fungicidal triazole) will be such that a synergistic fungicidal effect is obtained with both active ingredients. Particularly, the weight ratio between (I) and (II) may range from 20:1 to 1:20, more particularly from 10:1 to 1:10 and preferably will range from about 5:1 to 1:5. When the fungicidal triazole (II) is propiconazole, the weight

30 ratio between (I) and (II) preferably ranges from 2:1 to 1:4, more particularly from 1:1 to 1:3 and preferably will be about 1:2.

Apart from both the aforementioned active ingredients of formula (I) and (II), the compositions according to the present invention may further contain other active

35 ingredients, e.g. other microbiocides, in particular insecticides, and also bactericides, acaricides, nematocides, herbicides, plant growth regulators, fertilizers or further fungicides. As antimicrobial agents, which may be used in combination with the active

- substances there may be considered products of the following classes : phenol derivatives such as 3,5-dichlorophenol, 2,5-dichlorophenol, 3,5-dibromophenol, 2,5-dibromophenol, 2,5-(resp. 3,5)-dichloro-4-bromophenol, 3,4,5-trichlorophenol, tribromophenol, tetrachlorophenol, 3-methyl-4-chlorophenol; chlorinated
- 5 hydroxydiphenylethers such as, for example, 2-hydroxy-3,2'4'-trichlorodiphenylether; phenylphenol (o-, m-, p-), 4-chloro-2-phenylphenol, 4-chloro-2-benzylphenol, dichlorophene, hexachlorophene; aldehydes such as formaldehyde, glutaraldehyde, salicylaldehyde; alcohols such as phenoxyethanol; antimicrobially active carboxylic acids and their derivatives; organometallic compounds such as tributyltin compounds; iodine
- 10 compounds such as iodophores, iodonium compounds; diiodomethyl-p-tolylsulfone, 3-iodo-2-propynyl-alkohol, 4-chlorophenyl-3-iodopropargyl-formal, 3-bromo-2,3-diiodo-2-propenylethylcarb-Rat, 2,3,3-triiodoallylalkohol, 3-bromo-2,3-diiodo-2-propenylalkohol, 3-iodo-2-propynyl-n-butylcarbamate, 3-iodo-2-propynyl-n-hexylcarbamate, 3-iodo-2-propynyl-cyclohexylcarbamate, 3-iodo-2-propynyl-phenyl-
- 15 carbamate; mono-, di- and polyamines such as dodecylamine or 1,10-di(n-heptyl)-1,10-diaminodecane; sulfonium- and phosphonium compounds; mercapto compounds as well as their alkali, earth alkaline and heavy metal salts such as 2-mercaptopyridine-N-oxide and its sodium, iron, manganese and zinc salt, 3-mercaptopyridazin-2-oxide, 2-mercaptoquinoxaline-1-oxide, 2-mercaptoquinoxaline-di-N-oxide, as well as the
- 20 symmetrical disulfides of said mercapto compounds; ureas such as tribromo- or trichlorocarbanilide, dichlorotrifluoromethyl-diphenylurea; tribromosalicylanilide; 2-bromo-2-nitro-1,3-dihydroxypropane (bronopol); dichlorobenzoxazolone; chlorohexidine; sulfenamides e.g. dichlofluanid, tolylfluanid, folpet, fluorfolpet; benzimidazoles e.g. carbendazim, benomyl, fuberidazole, thiabendazole; thiocyanates
- 25 e.g. thiocyanatomethylthiobenzothiazole, methylenebisthiocyanate; quaternary ammonium compounds e.g. benzyldimethyltetradecyl ammonium chloride, benzyl-dimethyldodecyl ammonium chloride, didecyldimethyl ammonium chloride; morpholine derivatives e.g. tridemorph, fenpropimorph, falimorph; azoles e.g. triadimefon, triadimenol, bitertanol, prochloraz; 2-(1-chlorocyclopropyl)-1-(2-chlorophenyl)-3-(1H-
- 30 1,2,4-triazol-1-yl)-propan-2-ol; isothiazolinones e.g. N-methylisothiazolin-3-one, 5-chloro-N-methyl-isothiazolin-3-one, 4,5-dichloro-N-octylisothiazolin-3-one, N-octyl-isothiazolin-3-one;
- benzisothiazolinone, cyclopenteneisothiazolinone; tetrachloro-4-methylsulfonylpyridine; metal salts e.g. tin, copper, zinc naphthenate, octoate, 2-ethylhexanoate, oleate,
- 35 phosphate, benzoate; oxides e.g. tributyltin oxide, Cu₂O, CuO, ZnO; dialkyldithiocarbamate e.g. Na- and Zn-salts of dialkyldithiocarbamates, tetramethylthiuramdisulfide; nitriles e.g. 2,4,5,6-tetrachloroisophthalodinitrile;

-16-

- benzothiazole e.g. 2-mercaptobenzothiazole; quinolines e.g. 8-hydroxyquinoline and its Cu-salts; boron compounds e.g. boric acid, boric acid esters, borax; formaldehyde and formaldehyde releasing compounds e.g. benzylalcohol mono(poly)hemiformal, oxazolidine, hexahydro-S-triazine, N-methylol-chloracetamide, paraformaldehyde;
- 5 tris-N-(cyclohexyldiazeniumdioxy)-aluminium,
N-cyclohexyldiazeniumdioxy)-tributyltin bisN-(cyclohexyldiazeniumdioxy)-copper.

- As insecticidal agents which may be used in the combination according to the present invention the following classes of products may be considered : insecticides having a
- 10 natural origin, e.g., nicotine, rotenone, pyrethrum and the like; chlorinated hydrocarbons, e.g., lindane, chlordane, endosulfan and the like; organic phosphor compounds, e.g. azinphos-ethyl, azinphos-methyl, 1-(4-chlorophenyl)-4(O-ethyl, S-propyl)phosphoryl-oxypyrazole, chlorpyrifos, coumaphos, demeton, demeton-S-methyl, diazinon, dichlorvos, dimethoate, ethoprophos, etrimfos,
- 15 fenitrothion, fenthion, heptenophos, parathion, parathionmethyl, phosalone, phoxim, pirimiphos-ethyl, pirimiphos-methyl, profenofos, prothiofos, sulfprofos, triazophos, trichlorphon; carbamates, e.g., aldicarb, bendiocarb, carbaryl, carbofuran, carbosulfan, cloethocarb, 2-(1-methylpropyl)phenylmethylcarbamate, butocarboxime, butoxycarboxime, fenoxycarb, isoprocarb, methomyl, methiocarb, oxamyl, pirimicarb,
- 20 promecarb, propoxur and thiodicarb; ; biological insecticides, e.g., products originating from *Bacillus thuringiensis*; synthetic pyrethroids, e.g., allethrin, alphamethrin, bioresmethrin, bifenthrin, cycloprothrin, cyfluthrin, cyhalothrin, cypermethrin, decamethrin, deltamethrin, fenpropathrin, fenfluthrin, fenvalerate, flucythrinate, flumethrin, fluvalinate, halothrin, permethrin, resmethrin and tralomethrin,
- 25 alpha-cyano-3-phenyl-2-methylbenzyl-2~2-dimethyl-3-(2-chloro-2-tri-fluoromethylvinyl)cyclopropanecarboxylate;
- organosilicon compounds such as dimethylphenylsilylmethyl-3-phenoxybenzylethers e.g. dimethyl(4-ethoxyphenyl)silylmethyl-3-phenoxybenzylether; or dimethylphenylsilylmethyl-2-phenoxy-6-pyridylmethylethers e.g.
- 30 dimethyl(9-ethoxy-phenyl)silylmethyl-2-phenoxy-6-pyridylmethylether or [(phenyl)-3-(3-phenoxyphenyl)propyl](dimethyl)silanes e.g. (4-ethoxyphenyl)[3-(4-fluoro-3-phenoxyphenyl)propyl]dimethylsilane, silafluofen;
- nitroimines and nitromethylenes e.g. 1~(6-chloro-3-pyridinylmethyl)-4,5-dihydro-N-nitro-1H-imidazol-2-amine (imidacloprid); benzoylureas e.g. lufenuron,
- 35 hexaflumuron, flufenoxuron.

Particularly interesting active ingredients for combination with the present metconazole /

-17-

- triazole synergistic mixtures are : dichlofluanid, tolylfluanid, benzyldimethyldodecyl ammonium chloride, didecyldimethyl ammonium chloride, 3-bromo-2,3-diiodo-2-propenylalcohol, 3-iodo-2-propinyl-n-butylcarbamate, o-phenylphenol, m-phenylphenol, p-phenylphenol, 3-methyl-4-chlorophenol,
- 5 thiocyanatomethylthiobenzothiazole, N-methyl-isothiazolin-3-one, 5-chloro-N-methylisothiazolin-3-one, 4,5-dichloro-N-octylisothiazolin-3-one, N-octyl-isothiazolin-3-one, benzylalkoholmono(poly)hemiformal, N-methylolchloracetamide, phoxim, cyfluthrin, permethrin, cypermethrin, deltamethrin, imidacloprid, silafluofen, lufenuron, bifenthrin, fenoxycarb, hexaflumuron,
- 10 flufenoxuron.

Susceptible material (in particular wood) destroying insects are, for example :

Beetles

Anobium punctatum
Apate monachus
Bostrychus capucinus
Chlorophores pilosus
Dendrobium pertinex
Dinoderus minutus
Ernobius mollis
Heterobostrychus brunneus
Hylotrupes bajulus
Lyctus africanus
Lyctus brunneus
Lyctus linearis
Lyctus planicollis
Lyctus pubescens
Minthea rugicollis
Priobium carpini
Ptilinus pecticornis
Sinoxylon spp.
Trogoxylon aequale
Tryptodendron spp.
Xestobium rufovillosum
Xyleborus spp.

Hymenoptera:

Sirex juvencus
Urocerus augur
Urocerus gigas
Urocerus gigas taignus.

Termites:

Coptotermes formosanus
Cryptotermes brevis
Heterotermes indicola
Kalotermes flavicollis
Mastotermes darwiniensis
Reticulitermes flavipes
Reticulitermes lucifugus
Reticulitermes santonensis
Zootermopsis nevadensis.

-18-

The synergistic mixtures or compositions to be used directly may also be obtained from separate compositions containing the active ingredients or from the technical active ingredients themselves, by mixing and/or diluting with aqueous or organic media and/or optionally further adding adjuvants such as those described hereinabove. Said separate
5 compositions generally are such as described hereinbefore for compositions containing both active ingredients. Of particular interest to some users may be preparation of custom-made formulations from both active ingredients in unmodified, technical form, thus allowing maximal flexibility in the application of the present synergistic mixtures of metconazole and a fungicidal triazole.

10

The present invention also concerns a method of combating fungi comprising treating plants or the loci thereof, or treating plant products such as wood; or pulpwood for paper manufacture, or treating biodegradable materials simultaneously, separately or sequentially with an effective amount of a synergistic, fungicidal composition as
15 described hereinabove.

The present invention also concerns a method of preserving wood, wood products and biodegradable materials from deterioration by fungi. This method comprises the application to or incorporation in said wood or wood products or in or to said
20 biodegradable materials, of a synergistic mixture or composition as defined hereinabove.

Metconazole (I) and the fungicidal triazole (II) can be applied to plants or to the loci thereof or to plant products, e.g. wood, or to biodegradable materials such as textiles, simultaneously, or can also be applied consecutively within a time period selected so that
25 both active ingredients are allowed to act synergistically as antifungals, e.g. within 24 hours. In such applications, the active ingredients are used optionally together with adjuvants conventionally employed in the art of formulation such as carriers, surfactants or other useful additives. Therefore, the present invention also concerns products containing (I) metconazole, a salt, a stereoisomer or a stereoisomeric mixture thereof,
30 and (II) a fungicidal triazole, a salt, a stereoisomer or a stereoisomeric mixture thereof, as a combination for simultaneous, separate or sequential use in fungicidal applications. Such products may consists of a package comprising containers with both active ingredients, preferably in formulated form. Such formulated forms in general have the same composition as described for the formulations containing both active ingredients.

35

The following examples are intended to illustrate the scope of the present invention in all its aspects. Unless otherwise stated all parts therein are by weight.

-19-

Examples Throughout all examples percentages are by weight.

A) COMPOSITION EXAMPLES (Plant protection)

5	<u>Example 1: Wettable powders</u>	a)	b)	c)
	metconazole	10%	25%	0.25%
	a fungicidal triazole (azaconazole)	10%	25%	0.25%
	sodium lignosulfonate	5%	5%	5%
	sodium laurylsulfate	3%	-	-
10	sodium diisobutyl-naphthalenesulfonate	-	6%	6%
	octylphenol polyethylene glycol ether (7-8 moles of ethylene oxide)	-	2%	2%
	highly dispersed silicic acid	5%	27%	27%
	kaolin	67%	10%	-
15	sodium chloride	-	-	59.5%

The active ingredients were thoroughly mixed with the adjuvants and the mixture was thoroughly ground in a suitable mill, affording wettable powders which could be diluted with water to give suspensions of the desired concentration.

20

	<u>Example 2: Emulsifiable concentrates</u>	a)	b)	c)	d)
	metconazole	5%	0.5%	7%	9%
	a fungicidal triazole (bromuconazole)	5%	0.5%	3%	1%
	octylphenol polyethylene glycol ether (4-5 moles of ethylene oxide)	3%	3%	3%	3%
25	calcium dodecylbenzenesulfonate	3%	3%	3%	3%
	castor oil polyglycol ether (36 moles of ethylene oxide)	4%	4%	4%	4%
	cyclohexanone	30%	10%	30%	30%
30	dimethylbenzene mixture	50%	79%	50%	50%
		e)	f)	g)	h)
	metconazole	5%	2.5%	4%	9%
	a fungicidal triazole (tebuconazole)	5%	2.5%	1%	1%
35	calcium dodecylbenzenesulfonate	5%	8%	8%	5%
	castor oil polyethylene glycol ether (36 moles of ethylene oxide)	5%	-	-	5%

-20-

tributylphenol polyethylene glycol ether

(30 moles of ethylene oxide)

- 12% 12% -

cyclohexanone

- 15% 15% -

dimethylbenzene mixture

80% 60% 60% 80%

5

Emulsions of any required concentration could be obtained from these concentrates by dilution with water.

Example 3: Dusts

	a)	b)	c)	d)
10 metconazole	0.05%	0.5%	0.075%	0.095%
a fungicidal triazole (cyproconazole)	0.05%	0.5%	0.025%	0.005%
talcum	99.9%	-	99.9%	99.9%
kaolin	-	99%	-	

- 15 Usable dusts were obtained by mixing the active ingredients with the carriers, and grinding the mixture in a suitable mill.

Example 4: Extruder granulates

	a)	b)	c)	d)
metconazole	5%	0.5%	9.5%	0.9%
20 a fungicidal triazole (difenoconazole)	5%	0.5%	0.5%	0.1%
sodium lignosulfate	2%	2%	2%	2%
carboxymethylcellulose	1%	1%	1%	1%
kaolin	87%	96%	87%	96%

- 25 The active ingredients were mixed and ground with the adjuvants, and the mixture was subsequently moistened with water. The mixture was extruded and dried in a stream of air.

	e)	f)	g)	h)
30 metconazole	2.5%	5%	4.5%	8%
a fungicidal triazole (epoxiconazole)	2.5%	5%	0.5%	2%
kaolin	94%	-	94%	-
highly dispersed silicic acid	1%	-	1%	-
attapulgate	-	90%	-	90%

35

The active ingredients were dissolved in dichloromethane, the solution was sprayed onto the carrier, and the solvent was subsequently evaporated off in vacuo.

-21-

<u>Example 5: Coated granulates</u>		a)	b)	c)
	metconazole	1.5%	4%	9%
	a fungicidal triazole (triticonazole)	1.5%	1%	1%
	polyethylene glycol (mol. wt. 200)	2%	2%	2%
5	kaolin	95%	93%	88%

The active ingredients were uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granulates were obtained in this manner.

10	<u>Example 6: Suspension concentrates</u>	a)	b)	c)	d)
	metconazole	20%	2.5%	40%	30%
	a fungicidal triazole (hexaconazole)	20%	2.5%	8%	1.5%
	ethylene glycol	10%	10%	10%	10%
	nonylphenol polyethylene glycol ether				
15	(15 moles of ethylene oxide)	6%	1%	5%	7.5%
	sodium lignosulfate	10%	5%	9%	11%
	carboxymethylcellulose	1%	1%	1%	1%
	37% aqueous formaldehyde solution	0.2%	0.2%	0.2%	0.2%
	silicone oil in the form of a 75%				
20	aqueous emulsion	0.8%	0.8%	0.8%	0.8%
	water	32%	77%	26%	38%

The active ingredients were intimately mixed with the adjuvants, giving a suspension concentrate from which suspension of any desired concentration could be obtained by dilution with water.

		e)	f)	g)
	metconazole	5%	2.5%	10%
	a fungicidal triazole (penconazole)	5%	2.5%	5%
30	polyethylene glycol (MG 400)	70%	-	-
	N-methyl-2-pyrrolidone	20%	-	-
	epoxidised coconut oil	-	1%	1%
	petroleum distillate (boiling range			
	160-190°C)	-	94%	84%
35				

These solutions were suitable for application in the form of microdrops.

-22-

B) COMPOSITION EXAMPLES (Wood protection)

<u>Example 7: Water-dilutable concentrates</u>		a)	b)	c)	d)
	metconazole	3	2	1	0.5
5	triazole (propiconazole)	1	2	3	3.5
	Cu(OH) ₂ CuCO ₃	14	14	14	14
	monoethanolamine	33.5	33.5	33.5	27
	water	19	10.5	10.5	10.5
	nonylphenol polyethylene glycol ether				
10	(10 moles of ethylene oxide)	10	10	10	10
	propylene glycol	6	6	6	6
	boric acid	7.5	-	-	-
	propionic acid	6	-	-	-
	benzoic acid	-	22	22	21
15	polymin (n = 150)	-	-	-	7.5

C. Biological examplesExample 8

The synergistic activity of the mixtures or compositions of (I) and (II) according to the present invention can be demonstrated by comparison with the activity of the active ingredients (I) and (II) alone. The efficacy of the active ingredients against mycelial growth and sporulation of Coriolus versicolor, Gloeophyllum trabeum, Coniophora puteana, Poria monticola, Coriolus versicolor and Lentinus edodus was determined in the poison plate assay. The required concentrations of the fungicide(s) were obtained by diluting the active ingredients (I), (II) or the combination of (I) and (II) dissolved in 50% aqueous ethanol with a calculated amount of sterile water and pouring said dilutions in Petri-dishes. Malt extract agar (3%) was added aseptically and uniform distribution was obtained by shaking. Each plate was inoculated with mycelium from the margin of an actively growing colony. After incubation at 22°C and 70% relative humidity for a period long enough to allow complete growth of controls, diameters of colonies were measured. Relative activities were calculated by taking the absence of fungal growth (diameter 0 mm) as 100%. From the activity of the active ingredients alone, the expected activities E were calculated by using the so-called formula of Colby: (Colby, S.R. Weeds 1967, 15: 20-22),

$$E = X + Y - \frac{X \cdot Y}{100}$$

-23-

wherein X and Y express the relative activities obtained for each of the active ingredients. A synergistic effect can be acknowledged if the found activity exceeds calculated activity.

- 5 The results are listed in the table 1 below and clearly demonstrate that the measured activity generally exceeds the calculated activity. Equal efficacy was observed whenever complete inhibition of fungal growth occurred by one of the active ingredients (I) or (II) alone.
- 10 Table 1 : Activity of propiconazole, metconazole and their mixture against various basidiomycetes

Propiconazole conc. (ppm)	Metconazole conc. (ppm)	Measured activity (%)	Calculated activity (%)
<i>Gloeophyllum trabeum</i>			
10		90	
5		86	
2.5		80	
	0.25	92	
10	0.25	100	99
5	0.25	100	99
2.5	0.25	100	98
<i>Coniophora puteana</i>			
1.25		46	
	0.5	18	
	0.25	14	
	0.125	0	
1.25	0.5	76	56
1.25	0.25	78	54
1.25	0.125	84	46

-24-

Table 1 (cont'd) : Activity of propiconazole, metconazole and their mixture against various basidiomycetes

Propiconazole conc. (ppm)	Metconazole conc. (ppm)	Measured activity (%)	Calculated activity (%)
<i>Poria monticola</i>			
1.25		72	
	0.25	28	
	0.125	8	
1.25	0.25	86	80
1.25	0.125	84	74
<i>Coriolus versicolor</i>			
2.5		80	
1.25		58	
	0.5	16	
	0.25	0	
2.5	0.5	100	83
2.5	0.25	100	80
1.25	0.5	100	65
1.25	0.25	100	58
<i>Lentinus elodus</i>			
5		79	
2.5		53	
1.25		19	
	0.125	66	
5	0.125	100	93
2.5	0.125	100	84
1.25	0.125	100	72

Example 9

A greenhouse experiment demonstrating synergistic action of metconazole and propiconazole against the powdery mildew *Erysiphe graminis* on wheat and barley.

5 MATERIAL AND METHODS

- Test compounds were micronised by milling them for 10 minutes in a "Pulverisette 5" ball mill (Fritsch GMBH) and subsequently suspended in distilled water containing 0.05% macrogol as a dispersant. Treatment was performed as follows : two week old wheat or barley plants grown in 8 cm diameter plastic pots in garden soil were sprayed
10 until runoff with suitable doses of the test compounds. Four hours later the treated plants were dusted with *Erysiphe graminis* conidia from infected wheat or barley plants respectively, and placed in the greenhouse. After one week the severity of the infection was estimated visually and expressed as % fungal attack as compared to untreated controls (100 % fungal attack).
- 15 Synergy was determined by comparing observed % fungal attack values with expected values, calculated according to Limpel's formula, often called Colby's formula (Richer, 1987; Colby, 1967; Limpel et al., 1962).

Limpel's formula:
$$E = \frac{A \cdot B}{100}$$

20 Where

E = the expected % fungal attack when product 1 at dose a is combined with product 2 at dose b, supposing that the combined effect of both products is purely additive,

A = the % fungal attack observed when product 1 is applied alone at dose a, and

B = the % fungal attack observed when product 2 is applied alone at dose b.

25

When the % fungal attack observed for a combination of compounds is appreciably lower than the expected value, synergy is assumed.

-26-

RESULTS

The results for *Erysiphe graminis* on wheat, expressed as % fungal attack, can be found in Table 2. Expected fungal attack values for each combination are added between brackets.

5

Table 2 : % *Erysiphe graminis* attack on wheat plants treated with different combinations of propiconazole and metconazole. Each value is the mean of six replicates.

		propiconazole (ppm)			
		1	0,5	0,25	0
metconazole (ppm)	1	6 (79)	8 (88)	12 (88)	88
	0,5	20 (90)	58 (100)	100 (100)	100
	0,25	27 (90)	80 (100)	100 (100)	100
	0	90	100	100	100

10

As can be seen in table 2, synergy occurred with the following propiconazole : metconazole ratios: 1:1, 1:2, 1:4, 2:1 and 4:1.

-27-

The results for *Erysiphe graminis* on barley can be found in Table 3. Again, expected fungal attack values for each combination are added between brackets.

5 Table 3 : *Erysiphe graminis* attack on barley plants treated with different combinations of propiconazole and metconazole. Each value is the mean of six replicates.

		propiconazole (ppm)			
		8	4	2	0
metconazole (ppm)	2	7 (30)	38 (100)	100 (100)	100
	1	100 (100)	100 (100)	100 (100)	100
	0,5	100 (100)	100 (100)	100 (100)	100
	0	30	100	100	100

As can be seen in Table 3, synergy occurred with 2:1 and 4:1 propiconazole:metconazole ratios.

10

REFERENCES

Colby, S.R. (1967) *Weeds* **15**: 20-22

Limpel, L.E. et al (1962) *Proc. N. E. Weed Control Conf.* **16**: 48-53

Richer, D.L. (1987) *Pestic. Sci.* **19**: 309-315

-28-

CLAIMS

1. A composition comprising (I) metconazole, a salt, a stereoisomer or a stereoisomeric mixture thereof, and (II) at least one other fungicidal triazole, salt,
5 stereoisomer or stereoisomeric mixture thereof, in quantities producing a mutual synergistic fungicidal effect, and a carrier.
2. A composition according to claim 1 wherein the fungicidal triazole (II) is selected from the group consisting of azaconazole, bromuconazole, cyproconazole,
10 difenoconazole, epoxiconazole, fenbuconazole, hexaconazole, penconazole, propiconazole, tebuconazole and triticonazole.
3. A composition according to claim 1 wherein the ratio by weight between both active ingredients I:II is from 50:1 to 1:50.
15
4. A composition according to claim 1 for use in plant protection further comprising a cationic surfactant.
5. A composition according to claim 1 for use in material protection further comprising
20 a copper compound.
6. A composition according to claim 5 further comprising a non-ionic surfactant.
7. A product containing (I) metconazole, a salt, a stereoisomer or stereoisomeric
25 mixture thereof and (II) at least one other fungicidal triazole, salt, stereoisomer or stereoisomeric mixture thereof as a combination for simultaneous, separate or sequential use in fungicidal applications.
8. A process for preparing a synergistic composition as claimed in any of claims 1 to
30 6, characterized in that the active ingredients are intimately mixed with the carrier.
9. A method of combating fungi comprising treating plants or the loci thereof, or treating wood; or pulpwood for paper manufacture; or treating biodegradable materials simultaneously, separately or sequentially with an effective amount of a
35 synergistic, fungicidal composition as claimed in any of claims 1 to 6.
10. A method of preserving wood, wood products and biodegradable materials,
40 characterized by treating the wood, wood products or biodegradable materials with an effective amount of a synergistic, fungicidal composition as claimed in any of claims 1 to 6.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 95/02501

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A01N43/653 B27K3/50
43:653)

//(A01N43/653,59:20,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N B27K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 393 746 (JANSSEN PHARMACEUTICA N.V.) 24 October 1990 cited in the application see the whole document ---	1-10
A	EP,A,0 341 954 (KUREHA KAGAKU KOGYO KABUSHIKI KAISHA) 15 November 1989 cited in the application see page 2, line 13 - page 3, line 21 see table 1, compounds no. 1 and 2 see claims ---	1-10
A	WO,A,93 02557 (HICKSON INTERNATIONAL PLC) 18 February 1993 see page 2, line 3 - page 4, line 29; examples 1,3 see page 20 - page 21 --- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

7 November 1995

Date of mailing of the international search report

29. 11. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Muellners, W

INTERNATIONAL SEARCH REPORT

International Application No

PC1/EP 95/02501

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>WO,A,95 00303 (BAYER AG) 5 January 1995 cited in the application see page 1 - page 2, line 14 see page 2, line 31 - page 3, line 13 see page 6, lines 5,6 and 20,21 see page 13, line 13 - line 21; claims 1-6 -----</p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC/EP 95/02501

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0393746	24-10-90	AT-T- 107128	15-07-94
		AU-B- 623514	14-05-92
		AU-B- 5373390	25-10-90
		DE-D- 69009839	21-07-94
		DE-T- 69009839	06-10-94
		ES-T- 2057356	16-10-94
		FI-B- 94206	28-04-95
		JP-A- 2292203	03-12-90
		NO-B- 175922	26-09-94
		PL-B- 163728	29-04-94
		RU-C- 2024228	15-12-94
		US-A- 5397795	14-03-95
		US-A- 5223524	29-06-93
EP-A-0341954	15-11-89	JP-A- 2042003	13-02-90
		JP-B- 6099251	07-12-94
		US-A- 5292764	08-03-94
WO-A-9302557	18-02-93	AU-B- 659203	11-05-95
		AU-A- 2379792	02-03-93
		CA-A- 2114644	18-02-93
		EP-A- 0641164	08-03-95
		JP-T- 7500092	05-01-95
		NO-A- 940324	31-01-94
		NZ-A- 243460	27-06-95
		PT-A- 100707	29-10-93
WO-A-9500303	05-01-95	DE-A- 4320495	22-12-94
		DE-A- 4406819	07-09-95
		AU-B- 7123194	17-01-95