

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 September 2002 (12.09.2002)

PCT

(10) International Publication Number
WO 02/070541 A2

- (51) International Patent Classification⁷: **C07K**
- (21) International Application Number: PCT/US02/06275
- (22) International Filing Date: 1 March 2002 (01.03.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/272,570 1 March 2001 (01.03.2001) US
- (71) Applicant (*for all designated States except US*): **SMITH KLINE BEECHAM CORPORATION** [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **XIANG, JIA-NING** [US/US]; 707 DeSoto Drive, Palo Alto, CA 94303 (US). **CHRISTENSEN, Siegfried, B.** [US/US]; 1250 South Collegeville Road, Collegeville, PA 19426 (US). **LEE, Jinhwa** [KR/US]; 1250 South Collegeville Road, Collegeville, PA 19426 (US). **MERCER, Daniel, J.** [US/US]; 1250 South Collegeville Road, Collegeville, PA 19526 (US).
- (74) Agents: **SIMON, Soma, G.** et al.; SMITHKLINE BEECHAM CORPORATION, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/070541 A2

(54) Title: PEPTIDE DEFORMYLASE INHIBITORS

(57) Abstract: Nobel PDF inhibitors and novel methods for their use are provided.


PEPTIDE DEFORMYLASE INHIBITORS

FIELD OF THE INVENTION

The present invention relates to the use of novel antibacterial compounds, and
 5 pharmaceutical compositions containing these compounds as peptide deformylase
 inhibitors.

BACKGROUND OF THE INVENTION

Bacterial initiator methionyl tRNA is modified by methionyl tRNA
 10 formyltransferase (FMT) to produce formyl-methionyl tRNA. The formyl methionine
 (f-met) is then incorporated at the N-termini of newly synthesized polypeptides.
 Polypeptide deformylase (PDF or Def) then deformylates primary translation products
 to produce N-methionyl polypeptides. Most intracellular proteins are further
 processed by methionine amino peptidase (MAP) to yield the mature peptide and free
 15 methionine, which is recycled. PDF and MAP are both essential for bacterial growth,
 and PDF is required for MAP activity. This series of reactions is referred to as the
 methionine cycle (Figure 1).



20

Figure 1. The methionine cycle.

To date, polypeptide deformylase homologous genes have been found in bacteria, in chloroplast-containing plants, in mice and in human. The plant proteins are nuclear encoded but appear to carry a chloroplast localisation signal. This is consistent with the observation that chloroplast RNA and protein synthesis processes are highly similar to those of eubacteria. While there is limited information on protein expression of mammalian PDF gene homologs (Bayer Aktiengesellschaft, Pat. WO2001/42431), no functional role for such proteins has been demonstrated to date (Meinzel, T., Parasitology Today 16(4), 165-168, 2000).

Polypeptide deformylase is found in all eubacteria for which high coverage genomic sequence information is available. Sequence diversity among PDF homologs is high, with as little as 20% identity between distantly related sequences. However, conservation around the active site is very high, with several completely conserved residues, including one cysteine and two histidines which are required to coordinate the active site metal (Meinzel, T. et al., J. Mol. Biol. 267, 749-761, 1997).


PDF is recognized to be an attractive antibacterial target, as this enzyme has been demonstrated to be essential for bacterial growth in vitro (Mazel, D. et al., EMBO J. 13 (4), 914-923, 1994), is not believed to be involved in eukaryotic protein synthesis (Rajagopalan et al., J. Am. Chem. Soc. 119, 12418-12419, 1997), and is universally conserved in prokaryotes (Kozak, M., Microbiol. Rev. 47, 1-45, 1983). Therefore PDF inhibitors can potentially serve as broad spectrum antibacterial agents.

SUMMARY OF THE INVENTION

The present invention involves novel anti-bacterial compounds represented by Formula (1) hereinbelow and their use as PDF inhibitors.

DETAILED DESCRIPTION OF THE INVENTION

In one aspect of the present invention, there is provided a compound of formula (1):



(1) X = O, NR₃ or a bond;

Y = O, CH₂ or a bond

5 wherein:

R represents:

C₂₋₆ alkyl (optionally substituted by alkoxy, halogen, or C₁₋₃ alkylsulfanyl),
 C₂₋₆ alkenyl (optionally substituted by alkoxy, halogen, or C₁₋₃
 alkylsulfanyl), C₂₋₆ alkynyl (optionally substituted by alkoxy, halogen, or C₁₋₃
 10 alkylsulfanyl), (CH₂)_n—C₃₋₆ carbocycle (optionally substituted by alkoxy,
 halogen, or C₁₋₃ alkylsulfanyl), (CH₂)_n—R₄ {where R₄ is phenyl, furan,
 benzofuran, thiophene, benzothiophene, tetrahydrofuran, tetrahydropyran,
 dioxane, 1,4-benzodioxane or benzo[1,3]dioxole; R₄ is optionally substituted
 by one or more Cl, Br, I, C₁₋₃ alkyl (optionally substituted by one to three F)
 15 or C₁₋₂ alkoxy (optionally substituted by one to three F)};

R₁ represents:

hydrogen, C₁₋₆ alkyl (optionally substituted by hydroxy, halogen, amino,
 guanidino, phenyl, pyridyl, pyrrolyl, indolyl, imidazolyl, furanyl,
 benzofuranyl, piperidinyl, morpholinyl, quinolinyl, piperazinyl or
 20 dimethylaminophenyl) or (CH₂)_n—C₃₋₇ carbocycle;

R₂ represents:

hydrogen (provided that X is not O), C₁₋₃ substituted alkyl, C₂₋₃ substituted
 alkenyl, C₂₋₃ substituted alkynyl, (CH₂)_n—C₃₋₆ substituted carbocycle, aryl,
 heteroaryl, heterocyclic, carboxy (provided that X is not NR₃ or O) or
 25 aminocarbonyl (provided that X is not NR₃ or O);

R₃ represents:

hydrogen, C₁₋₃ substituted alkyl, phenyl, or may be taken together with R₂
 and the nitrogen atom to which they are attached to form an optionally

substituted heterocyclic ring which is optionally fused to an aryl, a heteroaryl, or a second heterocyclic ring;


X represents O, NR₃ or a covalent bond;

Y represents O, CH₂ or a covalent bond;

5 n = 0-2;

or a salt, solvate, or physiologically functional derivative thereof.

In this invention the most preferred R₁ group is hydrogen. Furthermore, in this invention the most preferred absolute configuration of compounds of the formula (1) is indicated below:



10

X = O, NR₃ or a bond;

Y = O, CH₂ or a bond

15 In a second aspect of the present invention, there is provided a compound of Formula (1) wherein X = O, and R, R₁, R₂, R₃, R₄, Y and n are as defined above; or a salt, solvate, or physiologically functional derivative thereof.

In a third aspect of the present invention, there is provided a compound of Formula (1) wherein X = NR₃, and R, R₁, R₂, R₃, R₄, Y and n are as defined above; or a salt, solvate, or physiologically functional derivative thereof.

20 In a fourth aspect of the present invention, there is provided a compound of Formula (1) wherein X is a covalent bond, and R, R₁, R₂, R₃, R₄, Y and n are as defined above; or a salt, solvate, or physiologically functional derivative thereof.

25 As used herein, the term "alkyl" refers to a straight or branched chain saturated hydrocarbon radical. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, isopentyl, hexyl and the like.

As used herein, the term "substituted alkyl" refers to a straight or branched chain saturated hydrocarbon radical, optionally substituted with substituents selected from the group that includes C₁₋₃ alkyl (optionally substituted by one to three fluorines), C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₂ alkoxy (optionally substituted by one to three fluorines), sulfanyl, sulfinyl, sulfonyl, oxo, hydroxy, mercapto, amino, guanidino, carboxy, aminocarbonyl, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, aminosulfonyl, sulfonylamino, carboxamide, ureido, nitro, cyano and halogen, multiple degrees of substitution being allowed.

As used herein, the term "alkenyl" refers to a straight or branched chain hydrocarbon radical having at least one carbon-carbon double bond. Examples of "alkenyl" as used herein include, but are not limited to, ethenyl and propenyl.

As used herein, the term "substituted alkenyl" refers to a straight or branched chain hydrocarbon radical having at least one carbon-carbon double bond, optionally substituted with substituents selected from the group which includes C₁₋₃ alkyl (optionally substituted by one to three F), amino, aryl, cyano and halogen, multiple degrees of substitution being allowed.

As used herein, the term "alkynyl" refers to a straight or branched chain hydrocarbon radical having at least one carbon-carbon triple bond. Examples of "alkynyl" as used herein include, but are not limited to, acetylenyl and 1-propynyl.

As used herein, the term "substituted alkynyl" refers to a straight or branched chain hydrocarbon radical having at least one carbon-carbon triple bond, optionally substituted with substituents selected from the group which includes C₁₋₃ alkyl (optionally substituted by one to three F), amino, aryl and halogen, multiple degrees of substitution being allowed.

As used herein, the term "halogen" refers to fluorine (F), chlorine (Cl), bromine (Br), or iodine (I), and "halo" refers to the halogen radicals fluoro, chloro, bromo and iodo.

As used herein, the term "carbocycle" refers to a non-aromatic cyclic hydrocarbon radical having from three to seven carbon atoms. For carbocycles with five- to seven-membered rings, a ring double bond is allowed. Exemplary

“carbocycle” groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, and cycloheptyl.

As used herein, the term “substituted carbocycle” refers to a non-aromatic cyclic hydrocarbon radical having from three to seven carbon atoms, and which is optionally substituted with substituents selected from the group which includes C₁₋₃ alkyl (optionally substituted by one to three F), C₂₋₃ alkenyl, C₂₋₃ alkynyl, C₁₋₂ alkoxy (optionally substituted by one to three F), sulfanyl, sulfinyl, sulfonyl, oxo, hydroxy, mercapto, amino, guanidino, carboxy, aminocarbonyl, aryl, aryloxy, heteroaryl, heterocyclic, aminosulfonyl, sulfonylamino, carboxamide, nitro, ureido, cyano and halogen, multiple degrees of substitution being allowed. For carbocycles with five- to seven-membered rings, a ring double bond is allowed.

As used herein, the term “aryl” refers to an optionally substituted benzene ring or to an optionally substituted benzene ring fused to one or more optionally substituted benzene rings to form a ring system. Exemplary optional substituents include C₁₋₃ substituted alkyl, C₂₋₃ substituted alkenyl, C₂₋₃ substituted alkynyl, heteroaryl, heterocyclic, aryl, C₁₋₃ alkoxy (optionally substituted by one to three F), aryloxy, aralkoxy, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, sulfanyl, sulfinyl, sulfonyl, aminosulfonyl, sulfonylamino, carboxamide, aminocarbonyl, carboxy, oxo, hydroxy, mercapto, amino, nitro, cyano, halogen, or ureido, multiple degrees of substitution being allowed. Such a ring or ring system may be optionally fused to one or more optionally substituted aryl rings (including benzene rings), carbocycle rings or heterocyclic rings. Examples of “aryl” groups include, but are not limited to, phenyl, naphthyl, tetrahydronaphthyl, biphenyl, indanyl, anthracyl or phenanthryl, as well as substituted derivatives thereof.

As used herein, the term “heteroaryl” refers to an optionally substituted monocyclic five to six membered aromatic ring containing one or more heteroatomic substitutions selected from S, SO, SO₂, O, N, or N-oxide, or to such an aromatic ring fused to one or more optionally substituted rings, such as heteroaryl rings, aryl rings, heterocyclic rings, or carbocycle rings (e.g., a bicyclic or tricyclic ring system). Examples of optional substituents are selected from the group which includes C₁₋₃ substituted alkyl, C₂₋₃ substituted alkenyl, C₂₋₃ substituted alkynyl, heteroaryl,

heterocyclic, aryl, C₁₋₃ alkoxy (optionally substituted by one to three F), aryloxy, aralkoxy, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, sulfanyl, sulfinyl, sulfonyl, aminosulfonyl, sulfonylamino, carboxamide, aminocarbonyl, carboxy, oxo, hydroxy, mercapto, amino, nitro, cyano, halogen or ureido, multiple
5 degrees of substitution being allowed. Examples of "heteroaryl" groups used herein include, but are not limited to, benzoimidazolyl, benzothiazolyl, benzoisothiazolyl, benzothiophenyl, benzopyrazinyl, benzotriazolyl, benzo[1,4]dioxanyl, benzofuranyl, 9*H*-*a*-carbolinyl, cinnolinyl, furanyl, furo[2,3-*b*]pyridinyl, imidazolyl, imidazolidinyl, imidazopyridinyl, isoxazolyl, isothiazolyl, isoquinolinyl, indolyl, indazolyl,
10 indolizinyl, naphthyridinyl, oxazolyl, oxothiadiazolyl, oxadiazolyl, phthalazinyl, pyridyl, pyrrolyl, purinyl, pteridinyl, phenazinyl, pyrazolyl, pyridyl, pyrazolopyrimidinyl, pyrrolizinyl, pyridazyl, pyrazinyl, pyrimidyl, 4-oxo-1,2-dihydro-4*H*-pyrrolo[3,2,1-*ij*]-quinolin-4-yl, quinoxalanyl, quinazolanyl, quinolinyl, quinolizinyl, thiophenyl, triazolyl, triazinyl, tetrazolopyrimidinyl,
15 triazolopyrimidinyl, tetrazolyl, thiazolyl, thiazolidinyl, and substituted versions thereof.

As used herein, the term "heterocyclic" refers to a three to seven-membered ring containing one or more heteroatomic moieties selected from S, SO, SO₂, O, N, or N-oxide, optionally substituted with substituents selected from the group which
20 includes C₁₋₃ substituted alkyl, C₂₋₃ substituted alkenyl, C₂₋₃ substituted alkynyl, heteroaryl, heterocyclic, aryl, C₁₋₃ alkoxy (optionally substituted by one to three F), aryloxy, aralkoxy, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, sulfanyl, sulfinyl, sulfonyl, aminosulfonyl, sulfonylamino, carboxamide, aminocarbonyl, carboxy, oxo, hydroxy, mercapto, amino, nitro, cyano, halogen, or
25 ureido, multiple degrees of substitution being allowed. Such a ring can be saturated or have one or more degrees of unsaturation. Such a ring may be optionally fused to one or more other optionally substituted "heterocyclic" ring(s), aryl ring(s), heteroaryl ring(s), or carbocycle ring(s). Examples of "heterocyclic" moieties include, but are not limited to, 1,4-dioxanyl, 1,3-dioxanyl, pyrrolidinyl, pyrrolidin-2-onyl, piperidinyl,
30 imidazolidine-2,4-dione-piperidinyl, piperazinyl, piperazine-2,5-dionyl, morpholinyl, dihydropyranyl, dihydrocinnolinyl, 2,3-dihydrobenzo[1,4]dioxinyl, 3,4-dihydro-2*H*-

benzo[b][1,4]-dioxepinyl, tetrahydropyranyl, 2,3-dihydrofuranyl, 2,3-dihydrobenzofuranyl, dihydroisoxazolyl, tetrahydrobenzodiazepinyl, tetrahydroquinolinyl, tetrahydrofuranyl, tetrahydronaphthyridinyl, tetrahydropurinyl, tetrahydrothiopyranyl, tetrahydrothiophenyl, tetrahydroquinoxalinyl, 5 tetrahydropyridinyl, tetrahydrocarbolinyl, 4H-benzo[1,3]-dioxinyl, benzo[1,3]dioxonyl, 2,2-difluorobenzo-[1,3]-dioxonyl, 2,3-dihydro-phthalazine-1,4-dionyl, isoindole-1,3-dionyl, and the like.

As used herein, the term "alkoxy" refers to the group $-OR_a$, where R_a is alkyl as defined above. Exemplary alkoxy groups useful in the present invention include, 10 but are not limited to, methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, and t-butoxy.

As used herein the term "aralkoxy" refers to the group $-OR_aR_b$, where R_a is alkyl and R_b is aryl as defined above.

As used herein the term "aryloxy" refers to the group $-OR_a$, where R_a is aryl as 15 defined above.

As used herein, the term "mercapto" refers to the group $-SH$.

As used herein, the term "sulfanyl" refers to the group $-SR_a$, where R_a is substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "sulfinyl" refers to the group $-S(O)R_a$, where R_a is 20 substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "sulfonyl" refers to the group $-S(O)_2R_a$, where R_a is 25 substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "oxo" refers to the group $=O$.

As used herein, the term "hydroxy" refers to the group $-OH$.

As used herein, the term "amino" refers to the group $-NH_2$. The amino group is optionally substituted by substituted alkyl, substituted carbocycle, aryl, heteroaryl 30 or heterocyclic, as defined above.

As used herein, the term "cyano" refers to the group $-CN$.

As used herein, the term "aminosulfonyl" refers to the group $-S(O)_2NH_2$. The aminosulfonyl group is optionally substituted by substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "sulfonylamino" refers to the group $-NHS(O)_2R_a$ where R_a is substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "carboxamide" refers to the group $-NHC(O)R_a$ where R_a is substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "carboxy" refers to the group $-C(O)OH$. The carboxy group is optionally substituted by substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "aminocarbonyl" refers to the group $-C(O)NH_2$. The aminocarbonyl group is optionally substituted by substituted alkyl, substituted carbocycle, aryl, heteroaryl or heterocyclic, as defined above.

As used herein, the term "ureido" refers to the group $-NHC(O)NHR_a$ wherein R_a is hydrogen, alkyl, carbocycle or aryl as defined above.

As used herein, the term "guanidino" refers to the group $-NHC(=NH)NH_2$.

As used herein, the term "acyl" refers to the group $-C(O)R_a$, where R_a is alkyl, carbocycle, or heterocyclic as defined herein.

As used herein, the term "aroyl" refers to the group $-C(O)R_a$, where R_a is aryl as defined herein.

As used herein, the term "heteroaroyl" refers to the group $-C(O)R_a$, where R_a is heteroaryl as defined herein.

As used herein, the term "acyloxy" refers to the group $-OC(O)R_a$, where R_a is alkyl, carbocycle, or heterocyclic as defined herein.

As used herein, the term "aroyloxy" refers to the group $-OC(O)R_a$, where R_a is aryl as defined herein.


As used herein, the term "heteroaroyloxy" refers to the group $-OC(O)R_a$, where R_a is heteroaryl as defined herein.

Also included in the present invention are pharmaceutically acceptable salts and complexes, such as the hydrochloride, hydrobromide and trifluoroacetate salts and the sodium, potassium and magnesium salts. The compounds of the present invention may contain one or more asymmetric carbon atoms and may exist in
5 racemic and optically active forms. All of these compounds and diastereomers are contemplated to be within the scope of the present invention.

GENERAL SYNTHETIC SEQUENCE

The compounds and processes of the present invention will be better
10 understood in connection with the following synthetic schemes, which are merely illustrative of the methods by which the compounds of the invention may be prepared and are not intended to limit the scope of the invention as defined in the appended claims.


The present invention provides compounds of Formula (1) that can be
15 prepared from the common racemic intermediate (8), or common chiral intermediates (17) and (25).



(1) X = O, NR₃ or a bond;

Y = O, CH₂ or a bond

20




Scheme 1.


As shown in Scheme 1, intermediate (8) can be prepared by reacting the mono-substituted dialkyl malonate (2) with a base, such as potassium hydroxide, in an appropriate solvent, such as ethanol/water, to afford the mono-acid (3). Coupling of (3) with O-benzylhydroxylamine in the presence of a coupling reagent, such as 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDCI), and a base, such as 4-dimethylaminopyridine, (DMAP) in an appropriate solvent, such as dichloromethane, gives the amide (4). Reduction of the ester functionality of compound (4) with a reducing agent, such as lithium borohydride, in an appropriate solvent, such as tetrahydrofuran, at room temperature provides the alcohol (5). Treatment of the alcohol (5) under Mitsunobu conditions affords the lactam (6). The same transformation may be achieved by treating (5) with triphenylphosphine, carbon tetrachloride and a base, such as triethylamine, to obtain (6). Hydrolysis of the lactam (6) using, for example, lithium hydroxide in an appropriate solvent mixture, such as THF-H₂O-MeOH, gives acid (7). Formylation of the amine group of (7) is achieved using formic acid and acetic anhydride in a solvent, such as dichloromethane, to provide the formylated compound (8).

Any racemates can be resolved at the level of any intermediate during the synthesis or at the level of the final product using, for example, a chiral chromatography method, to provide compound (8) in each of two enantiomeric forms.

Alternatively, an enantiomer of intermediate (8), such as (17) in Scheme 2 or (25) in Scheme 3, can be prepared by reacting an appropriate acid chloride (9) with a chiral agent, such as Evans' chiral oxazolidinone, in the presence of a base, such as n-butyl lithium, to afford the chiral intermediate (10) in Scheme 2 or (18) in Scheme 3. Treatment of the compound (10) or (18) with a base, such as diisopropylethylamine, in the presence of a chelating agent, such as titanium tetrachloride, in a solvent, such as tetrahydrofuran, followed by addition of an electrophile, such as benzyloxymethylchloride, provides either of two chiral compounds (11) and (19), depending on the selection of chiral auxiliary.



Scheme 2.



Scheme 3.

Conversion of compound (11) or (19) to the corresponding hydroxyacid (13) or (21) can be achieved by a sequence comprising oxidative cleavage of the chiral oxazolidinone, using, for example, H_2O_2 and lithium hydroxide to the respective intermediates (12) and (20), followed by hydrogenolysis. Coupling of the acid (13) or (21) with benzyloxyamine in the presence of coupling agents, such as EDCI/DMAP, yields the amides (14) and (22). These can be cyclized to the azetidin-2-ones (15) or (23) using either Mitsunobu conditions or a combination of triphenylphosphine/carbon tetrachloride/triethylamine. Hydrolysis of the azetidin-2-one (15) or (23), using for example lithium hydroxide, in an appropriate solvent,




gives the corresponding acid (16) or (24). Conversion of compound (16) or (24) to the formate (17) or (25) can be achieved using an appropriate formylating agent, such as formic acid/acetic anhydride or methyl formate, in an appropriate solvent, such as dichloromethane.

5

SPECIFIC EMBODIMENTS

Second Embodiment

As the second embodiment of the present invention, the compounds of
 10 Formula (1) with X = O are disclosed, as in the racemic compound (34) and the chiral compounds (36) and (38). These compounds have preferentially R1 = H.



15

Preferred compounds useful in the present invention are selected from the group consisting of:

20 N-Butyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

N-Butyl-N-phenoxy carbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

N-Isobutyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

25 N-Isobutyl-N-phenoxy carbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

N-Phenethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-Cyclohexylmethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Benzyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-(3-pyridin-3-yl-propyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(2-Morpholin-4-yl-ethyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Hydroxy-butyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-
- 10 heptanoyl}-hydrazine.
- N-(4-Amino-butyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(Tetrahydro-pyran-4-yl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-Methyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3-Aminopropyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(t-Butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-
- 20 hydrazine.
- N-(3-Hydroxypropyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-(t-butoxycarbonyl)-N'-{(2S)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-Butyl-N-(phenoxy carbonyl)-N'-{(2S)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[2-(4-Dimethylaminophenyl)ethyl]-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxy amino)methyl]-heptanoyl}-hydrazine.
- N-(t-Butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30 N-Pentyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.


- N-[2-(1H-Indol-3-yl)-ethyl]-N-(t-butoxycarbonyl)-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Isopentyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-
 heptanoyl}-hydrazine.
- 5 N-Cyclohexyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-
 heptanoyl}-hydrazine.
- N-(1-Ethyl-propyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-
 heptanoyl}-hydrazine.
- N-Isopropyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-
 10 heptanoyl}-hydrazine.
- N-Propyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-
 hydrazine.
- N-Ethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-
 hydrazine.
- 15 N-Methoxycarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-{[1-(3,5-Dimethoxyphenyl)-1-methyl-ethoxy]carbonyl}-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

The following synthetic schemes are merely illustrative of the methods by
 20 which the compounds of the invention may be prepared and are not intended to limit
 the scope of the invention as defined in the appended claims.

As shown in Scheme 4, treatment of the alkyl halide R1X (26) with hydrazine
 in a solvent such as ethanol, at an elevated temperature, gives hydrazine derivative
 (27). Reacting (27) with the carbonate R2OC(O)OC(O)OR2 or the chloroformate
 25 R2OCOC(=O)Cl affords the intermediate (28). Alternatively, (28) can be prepared from the
 Boc-protected hydrazine (29) by reaction with the aldehyde or ketone R'COR",
 followed by reduction with hydrogen gas in the presence of palladium, to afford
 hydrazine derivative (30). Reacting hydrazine (30) with a carbonate
 R2OC(O)OC(O)OR2 or a chloroformate R2OCOC(=O)Cl, followed by removal of the Boc
 30 protecting group with an appropriate acid, such as trifluoroacetic acid, gives the
 hydrazine derivative (28) wherein R1 = CHR'R". Alternatively, the primary amino


group of (29) can be protected as the phthalimide (31). Reacting the compound (31) with an alcohol under Mitsunobu conditions gives compound (32), which, upon hydrazinolysis, is readily converted to a hydrazine of Formula (10) where R2 = t-butyl.

5



Scheme 4.

As shown in Scheme 5, coupling of the acid (8) with the hydrazine derivative (28) using conditions such as DMAP/EDCI or EDCI/HOAt/NMM provides the hydrazide (33). Hydrogenolysis to remove the benzyl group using a catalyst, such as 10% Pd/C, in an appropriate solvent, such as ethanol, gives desired compound (34). Similarly, coupling of the chiral acid (17) or (25) with the hydrazine derivative (28) provides the corresponding hydrazide (35) or (37). Hydrogenolysis of the benzyl group gives the final desired compound (36) or (38).



Scheme 5.

5 SYNTHETIC EXAMPLES

The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

As used herein the symbols and conventions used in these processes, schemes and examples are consistent with those used in the contemporary scientific literature, for example, the *Journal of the American Chemical Society* or the *Journal of Biological Chemistry*. Standard single-letter or three-letter abbreviations are generally used to designate amino acid residues, which are assumed to be in the L-configuration unless otherwise noted. Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification.

Hz (Hertz);
chromatography);

T_r (retention time);

MeOH (methanol);

20 EtOH (ethanol);

TFA (trifluoroacetic acid);

DMSO (dimethylsulfoxide);

acetate);

TLC (thin layer

RP (reverse phase);

i-PrOH (isopropanol);

TEA (triethylamine);

THF (tetrahydrofuran);

AcOEt or EtOAc (ethyl

- DCM (dichloromethane);
dimethylformamide);
- CDI (1,1-carbonyldiimidazole);
HOSu (*N*-hydroxysuccinimide);
5 HOBT (1-hydroxybenzotriazole);
- mCPBA (meta-chloroperbenzoic acid);
fluorenylmethoxycarbonyl);
- DCC (dicyclohexylcarbodiimide);
10 NMM (*N*-methyl morpholine);
azabenzotriazole);
- DMAP (4-dimethylaminopyridine);
TBAF (tetra-*n*-butylammonium fluoride);
HPLC (high pressure liquid chromatography);
- 15 BOP (bis(2-oxo-3-oxazolidinyl)phosphinic chloride);
EDCI (1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride);
HBTU (*O*-Benzotriazole-1-yl-*N,N,N',N'*- tetramethyluronium
hexafluorophosphate).

20 All references to ether are to diethyl ether; brine refers to a saturated aqueous solution of NaCl. Unless otherwise indicated, all temperatures are expressed in °C (degrees Centigrade). All reactions are conducted under an inert atmosphere at room temperature unless otherwise noted, and all solvents are highest available purity unless otherwise indicated.

25 ¹H NMR (hereinafter also "NMR") spectra were recorded on a Varian VXR-300, a Varian Unity-300, a Varian Unity-400 instrument, a Bruker AVANCE-400, a General Electric QE-300 or a Bruker AM 400 spectrometer. Chemical shifts are expressed in parts per million (ppm, δ units). Coupling constants are in units of hertz (Hz). Splitting patterns describe apparent multiplicities and are designated as s
30 (singlet), d (doublet), t (triplet), q (quartet), quint (quintet), m (multiplet), br (broad).

Mass spectra were run on an open access LC-MS system using electrospray ionization. LC conditions: 4.5% to 90% CH₃CN (0.02% TFA) in 3.2 min with a 0.4 min hold and 1.4 min re-equilibration; detection by MS, UV at 214 nm, and a light scattering detector (ELS). Column: 1 X 40 mm Aquasil (C18).

5 For preparative (prep) hplc; *ca* 50 mg of the final products were injected in 500 μ L of DMSO onto a 50 X 20 mm I. D. YMC CombiPrep ODS-A column at 20 mL/min with a 10 min gradient from 10% CH₃CN (0.1% TFA) to 90% CH₃CN (0.1% TFA) in H₂O (0.1% TFA) and a 2 min hold. Flash chromatography was run over Merck Silica gel 60 (230 - 400 mesh).

10 Infrared (IR) spectra were obtained on a Nicolet 510 FT-IR spectrometer using a 1-mm NaCl cell. Most of the reactions were monitored by thin-layer chromatography on 0.25 mm E. Merck silica gel plates (60F-254), visualized with UV light, 5% ethanolic phosphomolybdic acid or p-anisaldehyde solution.

The compounds disclosed in Examples 2 to 30 were prepared following the
15 general procedures described in Example 1.

Preparation 1

(4S)-Benzyl-3-heptanoyl-oxazolidin-2-one.

To a solution of (S)-(-)-4-benzyl-2-oxazolidinone (3.3 g, 18.6 mmol) in THF (50 mL)
20 at -78 °C was added dropwise n-BuLi (7.4 mL, 2.5M solution in hexane, 18.6 mmol). After stirring for 30 min at the same temperature, the reaction mixture was then treated with heptanoyl chloride (2.76 g, 18.6 mmol). The reaction mixture was stirred and allowed to warm to 10 °C over 5 h, and then quenched with saturated aqueous NH₄Cl solution (100 mL). The aqueous layer was extracted with EtOAc (100 mL x
25 2). The combined organic layers were washed with brine, and dried over MgSO₄. Removal of the solvent under reduced pressure yielded 4.63 g (86%) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.37-7.22 (m, 5H), 4.69 (m, 1H), 4.19 (m, 2H), 3.31 (dd, J = 13.4, 3.3 Hz, 1H), 2.95 (m, 2H), 2.79 (dd, J = 13.4, 9.7 Hz, 1H), 1.71 (m, 2H), 1.42-1.32 (m, 6H), 0.92 (t, J = 6.8 Hz, 3H). MH⁺ 290.

30

Preparation 2

(4S)-Benzyl-3-[(2R)-benzyloxymethylheptanoyl]oxazolidin-2-one.

To a solution of (S)-4-benzyl-3-heptanoyloxazolidin-2-one (4.63 g, 16.02 mmol) and titanium (IV) chloride (1.9 mL, 16.82 mmol) in dichloromethane (55 mL) at 0 °C was added dropwise diisopropylethylamine (3.1 mL, 17.62 mmol). After stirring at 0 °C
5 for 1 hour, the resulting titanium enolate was then reacted with benzylchloromethyl ether (TCI-America, 4.9 mL, 32.04 mmol) at 0 °C for 6 h. The reaction mixture was then quenched with water (100 mL). The aqueous layer was extracted with dichloromethane (100 mL x 2). The organic extracts were washed with brine, and dried over MgSO₄. After removing the solvent under reduced pressure, purification
10 by flash column chromatography using an eluting system of hexane/EtOAc (5:1) yielded 4.39 g (67%) of the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.38-7.21 (m, 10H), 4.74 (m, 1H), 4.57 (m, 2H), 4.28-4.13 (m, 3H), 3.82 (t, J = 8.7 Hz, 1H), 3.68 (dd, J = 9.0, 4.9 Hz, 1H), 3.25 (dd, J = 13.5, 3.1 Hz, 1H), 2.71 (dd, J = 13.5, 9.3 Hz, 1H), 1.74 (m, 1H), 1.54 (m, 1H), 1.31-1.28 (m, 6H), 0.89 (t, J = 6.7 Hz, 3H).
15 MH+ 410.

Preparation 3**(3R)-Benzyloxy-2-pentylpropionic acid.**

A 0.05 M solution of (S)-4-benzyl-3-[(R)-2-benzyloxymethylheptanoyl]oxazolidin-2-
20 one (2.0 g, 4.89 mmol) in a 3:1 mixture of THF and H₂O was treated with 30% H₂O₂ (4.5 mL, 39.12 mmol), followed by LiOH (0.48 g, 9.78 mmol) at 0 °C. The resulting mixture was stirred and allowed to warm to room temperature overnight. THF was then removed under vacuum. The residue was washed with dichloromethane (50 mL x 2) to remove (S)-4-benzyloxazolidin-2-one. The desired product was isolated by
25 EtOAc extraction of the acidified (pH 1~2) aqueous phase. No further purification was required. Standing under high vacuum yielded 1.16 g (95%) of the title compound. ¹H NMR (400 MHz, CHCl₃) δ 11.1 (br s, 1H), 7.36 (m, 5H), 4.57 (s, 2H), 3.69 (m, 1H), 3.58 (dd, J = 9.2, 5.2 Hz, 1H), 2.74 (m, 1H), 1.66 (m, 1H), 1.54 (m, 1H), 1.34-1.30 (m, 6H), 0.90 (t, J = 6.7 Hz, 3H). MH+ 251.

30

Preparation 4

3-Hydroxy-(2R)-pentylpropionic acid.

To a solution of (R)-3-benzyloxy-2-pentyl-propionic acid (1.54 g, 6.16 mmol) in EtOH (100 mL) was added 10% Pd/C (310 mg). The reaction mixture was subjected to hydrogenation overnight at room temperature. After the reaction was completed, 5 the reaction mixture was filtered through a pad of Celite, and washed with EtOH (50 mL x 3). Removal of the solvent provided the title compound (0.92 g, 93%). No further purification was required. ¹H NMR (400 MHz, CHCl₃) δ 6.30 (br s, 1H), 3.81 (d, J = 5.4 Hz, 2H), 2.64 (m, 1H), 1.69 (m, 1H), 1.56 (m, 1H), 1.41-1.27 (m, 6H), 0.91 (t, J = 7.7 Hz, 3H). MH+ 161.

10

Preparation 5**N-Benzyloxy-3-hydroxy-(2R)-pentylpropionamide.**

To a mixture of (R)-3-hydroxy-2-pentylpropionic acid (0.92 g, 5.75 mmol), O-benzyl hydroxylamine hydrochloride (0.92 g, 5.75 mmol) and 4-(dimethylamino)pyridine 15 (1.41 g, 11.50 mmol) in dichloromethane (25 mL) at 0 °C was added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1.11 g, 5.75 mmol). After stirring at room temperature overnight, the reaction was then quenched with 1N aqueous HCl solution (25 mL) and extracted using dichloromethane (25 mL x 2). The organic extracts were washed with water, brine, and dried over MgSO₄. Removal 20 of the solvent under reduced pressure yielded the title compound (1.43 g, 94%). ¹H NMR (400 MHz, CHCl₃) δ 9.22 (br s, 1H), 7.41-7.28 (m, 5H), 4.89 (q, J = 10.6 Hz, 2H), 3.70-3.37 (m, 3H), 2.17 (m, 1H), 1.54 (br s, 1H), 1.27 (m, 6H), 0.88 (t, J = 6.9 Hz, 3H). MH+ 266.

25

Preparation 6**1-benzyloxy-(3R)-pentylazetid-2-one.**

To a mixture of (R)-N-benzyloxy-3-hydroxy-2-pentylpropionamide (1.41 g, 5.32 mmol) and triphenylphosphine (1.68 g, 6.39 mmol) in THF (53 mL) was added dropwise diethyl azodicarboxylate (1.1 mL, 6.39 mmol) at 0 °C. The reaction mixture 30 was stirred and allowed to warm to room temperature overnight. The reaction was then quenched with water (50 mL). The aqueous layer was extracted with EtOAc (50

mL x 2). The combined organic layers were washed with brine, and dried over MgSO₄. After removing the solvent under vacuum, the residue was purified by flash column chromatography (hex:EtOAc 5/1) to provide the title compound (1.17 g, 89%). ¹H NMR (400 MHz, CHCl₃) δ 7.35-7.25 (m, 5H), 4.87 (s, 2H), 3.28 (t, J = 4.85 Hz, 1H), 2.84 (q, J 2.35 Hz, 1H), 2.77 (m, 1H), 1.62 (m, 1H), 1.36 (m, 1H), 1.25-1.16 (m, 6H), 0.88 (t, J = 6.9 Hz, 3H). MH+ 248.

Preparation 7

3-benzyloxyamino-(2R)-pentylpropionic acid.

10 To a mixture of (R)-1-benzyloxy-3-pentylazetidin-2-one (0.96 g, 3.89 mmol) in a mixture of THF-H₂O-MeOH (50 mL, 3:1:1 v/v) was added lithium hydroxide monohydrate (1.91 g, 38.9 mmol). After stirring at room temperature overnight, water (25 mL) was added to the mixture. The solution was acidified to pH 5-6 with 3N aqueous HCl solution. It was extracted with EtOAc (50 mL x 2). The combined
15 organic layers were dried over MgSO₄. Removal of the solvent under vacuum provided the title compound (0.98 g, 95%). ¹H NMR (400 MHz, CHCl₃) δ 9.80 (br s, 1H), 7.37 (m, 5H), 4.75 (m, 2H), 3.14 (m, 2H), 2.74 (m, 1H), 1.70 (m, 1H), 1.53 (m, 1H), 1.38-1.25 (m, 6H), 0.91 (t, J = 6.8 Hz, 3H). MH+ 266.

20

Preparation 8

(2R)-[(benzyloxyformylamino)methyl]heptanoic acid.

To a cold solution of (R)-3-Benzyloxyamino-2-pentylpropionic acid (1.03 g, 3.89 mmol) in HCO₂H (19 mL) and dichloromethane (19 mL) at 0 °C was added acetic anhydride (3.9 mL, 41.2 mmol). The mixture was stirred at 0 °C for 3 hours. The
25 volatiles were removed by evaporation under vacuum. Dichloromethane (50 mL) was added to it. It was washed with brine (50 mL x 2), and dried over MgSO₄. Filtration and evaporation under vacuum provided the title compound (1.08 g, 95%). ¹H NMR (400 MHz, CHCl₃) δ 8.07 (br s, 1H), 7.29 (m, 5H), 4.91-4.71 (m, 2H), 3.76 (m, 2H), 2.67 (m, 1H), 1.54 (m, 1H), 1.41(m, 1H), 1.20 (m, 6H), 0.80 (t, J = 7.0 Hz, 3H).
30 MH+ 294.

Preparation 9

Butylhydrazine.

1-Iodobutane (5.1 mL, 45.1 mmol) was added through a condenser for 30 min to a refluxing solution of hydrazine monohydrate (11.3 g, 225.5 mmol) in EtOH (100 mL). After stirring and refluxing for 18 hours, ethanol was removed by evaporation under vacuum. The residue was extracted with ether (50 mL x 2). The combined organic layers were dried (K_2CO_3), filtered and evaporated to yield 2.6 g (66%) of the title compound. 1H NMR (400 MHz, $CHCl_3$) δ 3.15 (br s, 2H), 2.73 (t, J = 7.2 Hz, 2H), 1.44 (m, 2H), 1.33 (m, 2H), 0.89 (t, J 7.2 Hz, 3H). MH+ 89.

10

Preparation 10

N-Butylhydrazinecarboxylic acid tert-butyl ester.

To a solution of butylhydrazine (510 mg, 5.80 mmol) and triethylamine (1.2 mL, 8.69 mmol) in dichloromethane (20 mL) at 0 °C was added di-t-butyl dicarbonate (1.26 g, 5.80 mmol). The reaction mixture was stirred and allowed to warm up to room temperature overnight. Water (20 mL) was then added to the reaction mixture. The aqueous layer was extracted with dichloromethane (20 mL x 2). The combined organic layers were dried ($MgSO_4$). Filtration and evaporation under vacuum provided the title compound (820 mg, 75%). 1H NMR (400 MHz, $CHCl_3$) δ 3.89 (br s, 2H), 3.29 (t, J = 7.1 Hz, 2H), 1.46 (m, 2H), 1.40 (s, 9H), 1.24 (m, 2H), 0.86 (t, J = 7.3 Hz, 3H). MH+ 189.

20

Preparation 11

N-Butyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(benzyloxyformylamino)methyl]heptanoyl}-hydrazine.

25

To a solution of (R)-2-[(benzyloxyformyl-amino)methyl]heptanoic acid (180 mg, 0.614 mmol), N-butyl-hydrazinecarboxylic acid tert-butyl ester (140 mg, 0.737 mmol) and 4-dimethylaminopyridine (90 mg, 0.737 mmol) in dichloromethane (6.5 mL) at 0 °C was added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (142 mg, 0.737 mmol). After stirring at room temperature overnight, the reaction was then quenched with aqueous 1N HCl and extracted with dichloromethane (15 mL x 2).

30

The organic extracts were washed with brine (30 mL), and dried over MgSO₄. Evaporation of the solvent under vacuum, followed by purification by flash column chromatography yielded 195 mg (69%) of the title compound. ¹H NMR (400 MHz, CHCl₃) δ 8.09 (br s, 1H), 7.25 (s, 5H), 4.80 (m, 2H), 4.10 (dd, J = 14.1, 4.0 Hz, 1H),
5 3.62 (m, 1H), 3.35 (m, 2H), 2.55 (m, 1H), 1.72 (m, 1H), 1.56 (m, 1H), 1.50 (s, 9H), 1.30 (m, 10H), 0.90 (m, 6H). MH+ 464.

Example 1

**N-Butyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-
10 heptanoyl}-hydrazine.**

To a solution of N'-{(R)-2-[(benzyloxy-formylamino)methyl]heptanoyl}-N-butylhydrazine carboxylic acid tert-butyl ester (195 mg, 0.421 mmol) in EtOH (15 mL) was added 10% Pd/C (60 mg). The reaction mixture was subjected to hydrogenation overnight at room temperature. After the reaction was completed, the
15 reaction mixture was filtered through a pad of Celite, and washed with EtOH (10 mL x 2). Removal of the solvent provided the crude product, which was further purified by HPLC to yield the title compound (52 mg, 33%). ¹H NMR (400 MHz, CHCl₃) δ 9.94 (s, 1H), 9.39 (s, 1H), 8.32 (s, 1H), 4.10 (dd, J = 14.1, 4.0 Hz, 1H), 3.62 (m, 1H), 3.35 (m, 2H), 2.55 (m, 1H), 1.72 (m, 1H), 1.56 (m, 1H), 1.50 (s, 9H), 1.30 (m, 10H),
20 0.90 (m, 6H). MH+ 374.

Preparation 12

N-Butylhydrazinecarboxylic acid phenyl ester.

To a solution of butylhydrazine (370 mg, 4.20 mmol) and triethylamine (0.88 mL,
25 6.30 mmol) in dichloromethane (15 mL) at 0 °C was added phenyl chloroformate (0.53 mL, 4.20 mmol). The reaction mixture was stirred and allowed to warm up to room temperature overnight. Water (20 mL) was then added to the reaction mixture. The aqueous layer was extracted with dichloromethane (20 mL x 2). The combined organic layers were dried (MgSO₄). Filtration and evaporation under vacuum,
30 followed by purification by flash column chromatography provided the title compound (250 mg, 29%). ¹H NMR (400 MHz, CHCl₃) δ 7.40-7.10 (m, 5H), 4.20 (t,

J = 7.1 Hz, 2H), 3.60 (br s, 2H), 1.71 (m, 2H), 1.41 (m, 2H), 0.98 (t, J = 7.3 Hz, 3H).
MH+ 209.

Example 2

5 **N-Butyl-N-phenoxy carbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

Purification by preparative HPLC yielded 49 mg (41%) of the title compound. ¹H NMR (400 MHz, CHCl₃) δ 9.78 (s, 1H), 9.39 (s, 1H), 8.27 (s, 1H), 7.40-7.10 (m, 5H), 4.20-3.30 (m, 4H), 2.70 (m, 1H), 1.80-1.20 (m, 10H), 0.90 (m, 6H). MH+ 394.

10

Example 3

N-Isobutyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15 Purification by preparative HPLC yielded 44 mg (22%, 2 steps) of the title compound. ¹H NMR (400 MHz, CHCl₃) δ 9.94 (s, 1H), 9.54 (s, 1H), 8.32 (s, 1H), 4.11 (dd, J = 3.9, 14.1 Hz, 1H), 3.46 (m, 1H), 3.35 (m, 1H), 3.12 (dd, J = 6.3, 14.1 Hz, 1H), 2.54 (m, 1H), 1.88 (m, 1H), 1.72 (m, 1H), 1.50 (s, 9H), 1.49-1.25 (m, 7H), 0.95-0.88 (m, 9H). MH+ 374.

20

Example 4

N-Isobutyl-N-phenoxy carbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

25 Purification by preparative HPLC yielded 44 mg (22%, 2 steps) of the title compound. ¹H NMR (400 MHz, CHCl₃) δ 9.97 (s, 1H), 9.38 (s, 1H), 8.27 (s, 1H), 7.41-7.13 (m, 5H), 4.08 (dd, J = 3.9, 14.1 Hz, 1H), 3.37 (m, 2H), 2.65 (m, 1H), 2.02 (m, 1H), 1.74 (m, 1H), 1.32-1.02 (m, 7H), 0.92-0.82 (m, 9H). MH+ 394.

Example 5

30 **N-Phenethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

^1H NMR (400 MHz, CDCl_3) δ 9.80 (s, 1H), 9.03 (s, 1H), 8.35 (s, 1H), 7.22 (s, 5H), 4.11-3.36 (m, 4H), 2.86 (t, 3H), 2.52 (m, 1H), 2.07 (m, 1H), 1.72 (m, 1H), 1.41-1.34 (m, 15H), 0.92 (t, 3H). MH+ 422.

5

Example 6

N-Cyclohexylmethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3) δ 8.44 (s, 1H), 7.70 (br s, 1H), 4.20-3.05 (m, 4H), 2.50 (m, 1H), 1.74-0.90 (m, 31H). MH+ 414.

10

Example 7

N-Benzyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3) δ 9.70 (br s, 1H), 8.77 (br s, 1H), 8.25 (s, 1H), 7.64-7.28 (m, 5H), 5.00 (d, $J = 14.7$ Hz, 1H), 2.40 (m, 1H), 1.66 (m, 1H), 1.56 (m, 1H), 1.52 (s, 9H), 1.48-1.10 (m, 6H), 0.87 (t, $J = 7.1$ Hz, 3H). MH+ 408.

15

Example 8

N-(3-pyridin-3-yl-propyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

20

^1H NMR (400 MHz, CDCl_3) δ 8.79-7.25 (m, 5H), 4.15-3.03 (m, 4H), 2.75 (m, 1H), 2.56 (m, 2H), 2.07-1.72 (m, 4H), 1.49 (s, 9H), 1.46-1.28 (m, 6H), 0.85 (t, $J = 6.9$ Hz, 3H). MH+ 437.

25

Example 9

N-(2-Morpholin-4-yl-ethyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3) δ 8.45 (br s, 1H), 7.88 (br s, 1H), 4.20-3.30 (m, 8H), 2.70 (m, 1H), 2.60-2.45 (m, 6H), 1.72 (m, 1H), 1.50 (s, 9H), 1.39 (m, 1H), 1.29 (m, 6H), 0.89 (t, $J = 7.1$ Hz, 3H). MH+ 431.

30

Example 10

N-(4-Hydroxy-butyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 5 ¹H NMR (400 MHz, CDCl₃) δ 9.04 (br s, 1H), 8.37 (br s, 1H), 4.07-3.46 (m, 6H), 2.54 (m, 1H), 1.64-1.30 (m, 12H), 1.47 (s, 9H), 0.89 (t, J = 6.9 Hz, 3H). MH+ 390.

Example 11

10 **N-(4-Amino-butyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD) δ 7.90 (s, 1H), 7.70 (s, 1H), 4.00 (m, 2H), 3.70-3.40 (m, 4H), 3.0 (m, 2H), 2.50 (m, 1H), 1.70 (m, 1H), 1.50 (m, 1H), 1.49 (s, 9H), 1.48-1.20 (m, 10H), 0.89 (t, J = 6.9 Hz, 3H). MH+ 389.

15

Example 12

N-(Tetrahydro-pyran-4-yl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- ¹H NMR (400 MHz, CDCl₃) δ 9.88 (br s, 1H), 8.31 (s, 1H), 4.16-4.00 (m, 4H), 3.44-3.39 (m, 3H), 2.60 (m, 1H), 1.97-1.26 (m, 12H), 0.90 (t, J = 6.9 Hz, 3H). MH+ 402.

20

Example 13

N-Methyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 25 ¹H NMR (400 MHz, CDCl₃) δ 9.74 (br s, 1H), 8.90 (br s, 1H), 8.29 (br s, 1H), 4.06 (m, 1H), 3.24 (m, 1H), 3.10 (s, 3H), 2.43 (m, 1H), 1.64 (m, 1H), 1.42 (s, 9H), 1.31 (m, 1H), 1.19 (m, 6H), 0.79 (t, J = 6.9 Hz, 3H). MH+ 332.

Example 14

**N-(3-Amino-propyl)-N-(t-butoxycarbonyl)-N'-{2-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 8.27 (s, 1H), 7.78 (s, 1H), 3.76-3.32 (m, 4H), 2.84 (m, 2H), 2.68 (m, 1H), 1.80 (m, 1H), 1.74 (m, 1H), 1.48 (s, 9H), 1.35 (m, 8H), 0.93 (t, J = 6.8 Hz, 3H). MH+ 375.

Example 15

N-(t-Butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.53 (s, 1H), 8.17 (s, 1H), 6.68 (s, 1H), 4.11 (m, 1H), 3.38 (m, 1H), 2.63 (m, 1H), 1.70 (m, 1H), 1.49 (s, 9H), 1.42 (m, 1H), 1.29 (m, 6H), 0.87 (t, J = 6.8 Hz, 3H). MH+ 318.

Example 16

**N-(3-Hydroxy-propyl)-N-(t-butoxycarbonyl)-N'-{2-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 9.34 (s, 1H), 8.33 (s, 1H), 7.83 (s, 1H), 3.83-3.43 (m, 6H), 2.81 (m, 1H), 1.78-1.65 (m, 2H), 1.52-1.29 (m, 8H), 1.45 (s, 9H), 0.87 (s, 9H). MH+ 376.

20

Example 17

N-Butyl-N-(t-butoxycarbonyl)-N'-{(2S)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 374.

25

Example 18

N-Butyl-N-(phenoxycarbonyl)-N'-{(2S)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.27 (s, 1H), 8.20 (s, 1H), 7.32-7.04 (m, 5H), 3.80-3.31 (m, 4H), 2.55 (s, 1H), 1.66 (s, 1H), 1.59-1.24 (m, 11H), 0.90-0.82 (m, 6H).
MH+ 394.

5

Example 19

N-[2-(4-Dimethylaminophenyl)ethyl]-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxy amino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.81 (s, 1H), 8.89 (s, 1H), 8.33 (s, 1H), 7.06 (d, J = 8.5 Hz, 2H), 6.71 (d, J = 8.5 Hz, 2H), 4.17-3.33 (m, 4H), 2.93 (s, 6H), 2.80 (m, 2H),
10 2.48 (m, 1H), 1.71 (m, 1H), 1.41 (s, 9H), 1.32 (m, 7H), 0.91 (t, 3H). MH+ 465.

Example 20

N-(t-Butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15 ¹H NMR (400 MHz, CDCl₃) δ 9.81 (s, 1H), 9.54 (s, 1H), 8.46 (s, 1H), 6.78 (s, 1H), 3.85-3.37 (m, 2H), 2.80-2.62 (m, 1H), 1.71 (m, 1H), 1.49 (s, 9H), 1.30-1.25 (m, 7H), 0.89 (t, 3H). MH+ 318.

Example 21

20 **N-Pentyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 9.10 (s, 1H), 8.43 (s, 1H), 4.14-3.05 (m, 4H), 2.86-2.48 (m, 1H), 1.61 (m, 1H), 1.41 (m, 9H), 1.25-1.16 (m, 13H), 0.84 (m, 6H). MH+ 388.

25

Example 22

N-[2-(1H-Indol-3-yl)-ethyl]-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.78 (s, 1H), 8.55 (s, 1H), 8.33 (s, 1H), 8.15 (s, 1H), 7.55-7.05 (m, 5H), 4.10-3.92 (m, 2H), 3.71-3.30 (m, 2H), 3.02 (m, 2H), 2.40 (m, 1H), 1.67 (m, 1H), 1.50-1.20 (m, 16H), 0.89 (m, 6H). MH+ 461.

5

Example 23

N-Isopentyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.92 (s, 1H), 8.39 (s, 1H), 4.05-3.63 (m, 2H), 3.31 (t, 2H), 2.54 (m, 1H), 1.70 (m, 1H), 1.55 (m, 1H), 1.50-1.18 (m, 18H), 0.88 (m, 9H).

10 MH+ 388.

Example 24

N-Cyclohexyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15 ¹H NMR (400 MHz, CDCl₃) δ 8.79 (s, 1H), 8.30 (s, 1H), 4.11 (m, 1H), 3.85 (m, 1H), 3.35 (m, 1H), 2.55 (m, 1H), 1.94 (m, 1H), 1.81-1.57 (m, 5H), 1.51-1.18 (m, 18H), 0.89 (m, 3H). MH+ 400.

Example 25

20 **N-(1-Ethyl-propyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 9.81 (s, 1H), 8.41 (s, 1H), 4.11-3.32 (m, 3H), 2.55 (m, 1H), 1.80 (m, 1H), 1.58-1.15 (m, 20H), 1.05 (m, 3H), 0.89 (m, 6H). MH+ 388.

25

Example 26

N-Isopropyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 8.92 (s, 1H), 8.31 (s, 1H), 4.34-4.08 (m, 2H), 3.37 (m, 1H), 2.76 (m, 1H), 1.73 (m, 1H), 1.51 (s, 9H), 1.31 (m, 7H), 1.11 (m, 6H), 0.88

30 (m, 3H). MH+ 360.

Example 27

N-Propyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 5 ^1H NMR (400 MHz, CDCl_3) δ 9.37 (s, 1H), 8.24 (s, 1H), 4.03-3.17 (m, 4H), 2.42 (m, 1H), 1.61 (m, 1H), 1.52-1.31 (m, 11H), 1.20 (m, 7H), 0.82 (m, 6H). MH+ 360.

Example 28

N-Ethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 10 ^1H NMR (400 MHz, CDCl_3) δ 9.79 (s, 1H), 8.05 (s, 1H), 4.21-3.25 (m, 4H), 2.48 (m, 1H), 1.51-1.02 (m, 20H), 0.88 (m, 3H). MH+ 346.

Example 29

- 15 **N-Methoxycarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

^1H NMR (400 MHz, CDCl_3) δ 8.35 (s, 1H), 8.05 (s, 1H), 7.64 (s, 1H), 7.50 (s, 1H), 3.68 (m, 3H), 3.37 (d, J = 5.5 Hz, 2H), 2.72 (m, 1H), 1.61 (m, 1H), 1.41-1.12 (m, 7H), 0.85 (m, 3H). MH+ 276.

20

Example 30


N-[[1-(3,5-Dimethoxyphenyl)-1-methyl-ethoxy]carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 25 ^1H NMR (400 MHz, CDCl_3) δ 8.71 (s, 1H), 8.21 (s, 1H), 7.76 (s, 1H), 6.62 (s, 1H), 6.36 (d, J = 4.7 Hz, 1H), 3.83-3.69 (m, 7H), 3.38 (m, 1H), 2.70 (m, 1H), 1.82-1.55 (m, 7H), 1.43-1.21 (m, 7H), 0.84 (m, 3H). MH+ 440.

Third Embodiment

As the third embodiment of the present invention, the compounds of Formula (1) with $X = NR_3$ are disclosed, as in the racemic compound (42) and the chiral compounds (44) and (46). These compounds have preferentially $R_1 = H$.

5



Preferred compounds useful in the present invention are selected from the
10 group consisting of:


- N-Butyl-N-[(4-methylpiperazin-1-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-diphenylaminocarbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-
15 heptanoyl}-hydrazine.
- N-Butyl-N-(t-butylamino)carbonyl-N'-{2-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-Butyl-N-phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-
hydrazine.
- 20 N-Butyl-N-[(3,5-dimethyl-4,5-dihydro-isoxazol-4-yl)aminocarbonyl]-N'-{2-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-[(1-morpholin-4-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-4-phenyl-butanoyl}-
25 hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-hexanoyl}-
hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-3-phenyl-
propanoyl}-hydrazine.

- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-3-(3,4-dichlorophenyl)-propanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-(3,4-Dichlorophenylaminocarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3,4-Dichlorophenylaminocarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-
- 10 heptanoyl}-hydrazine.
- N-[(1-Morpholin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Methoxyphenyl)aminocarbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(2,4-Dichlorophenyl)aminocarbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,6-Dichlorophenyl)aminocarbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(4-Methyl-piperazin-1-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
- 20 heptanoyl}-hydrazine.
- N-[(4-Chloro-3-trifluoromethylphenyl)aminocarbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Methyl-phenyl-amino)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
- 25 heptanoyl}-hydrazine.

The following synthetic schemes are merely illustrative of the methods by which the compounds of the invention may be prepared and are not intended to limit the scope of the invention as defined in the appended claims.

- As shown in Scheme 6, treatment of an alkyl halide R1X (26) with hydrazine
- 30 in a solvent such as ethanol, at an elevated temperature, gives the hydrazine derivative (27). Reacting compound (27) with the carbamyl chloride R2R3NC(O)Cl affords

intermediate (39). Alternatively, compound (39) can be prepared from the Boc-protected hydrazine (29) by reaction with the aldehyde or ketone R'COR'', followed by reduction with hydrogen gas in the presence of palladium, to afford hydrazine derivative (30). Reacting hydrazine (30) with the carbamyl chloride R₂R₃NC(O)Cl, followed by removing the Boc protecting group with an appropriate acid, such as trifluoroacetic acid, gives the hydrazine derivative (39) wherein R₁ = CHR'R''. Alternatively, reacting the isocyanate R₂NCO (40) with hydrazine affords compound (39) wherein R₁ = H and R₃ = H. Alternatively, reacting the isocyanate R₂NCO (40) with Boc-protected hydrazine, followed by acid treatment, affords the salt form of compound (39) wherein R₁ = H and R₃ = H. Alternatively, reacting the isocyanate R₂NCO (40) with Cbz-protected hydrazine, followed by hydrogenation, affords compound (39) wherein R₁ = H and R₃ = H.




15

Scheme 6.

As shown in Scheme 7, coupling of the carboxylic acid (8) with the hydrazine derivative (39) provides hydrazide (41). Hydrogenolysis to remove the benzyl group using a catalyst, such as 10% Pd/C, in an appropriate solvent, such as ethanol, gives compound (42). Similarly, coupling of the chiral acid (17) or (25) with hydrazine

20

derivative (39) provides the corresponding hydrazide (43) or (45). Hydrogenolysis of the benzyl group gives the final compounds (44) or (46).




5

Scheme 7.

Alternatively, as shown in Scheme 8, the carbamate (34) wherein R2 = *t*-butyl or benzyl can be converted to the hydrazide (47) by acid treatment or hydrogenolysis, respectively. Reaction of (47) with an isocyanate in an appropriate solvent, such as methylene chloride, and in the optional presence of an appropriate base, such as triethylamine, gives compound (42) wherein R3 = H. Similarly, appropriate deprotection of the chiral carbamates (36) and (38), followed by reaction with an isocyanate, gives the chiral hydrazide (44) and (46), wherein R3 = H.

15



Scheme 8.

5 SYNTHETIC EXAMPLES

The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. The same experimental general conditions and conventions described in the Experimental Section of the Second Embodiment are applicable here.

10 The compounds disclosed in Examples 31 to 51 were prepared following the general procedures described in Example 1.

Preparation 13.

N-[(4-Methylpiperazine)carbonyl]-N-butylhydrazine.

15 To a solution of butyl-hydrazine (200 mg, 2.27 mmol) and triethylamine (0.95 mL, 6.81 mmol) in dichloromethane (10 mL) at $-78\text{ }^\circ\text{C}$ was added 4-methyl-1-piperazine carbonyl chloride hydrochloride (0.45 g, 2.27 mmol). The reaction mixture was stirred and allowed to warm up to room temperature overnight. Saturated aqueous NaHCO_3 (20 mL) was then added to the reaction mixture. The aqueous layer was

20 extracted with dichloromethane (20 mL x 2). The combined organic layers were dried (MgSO_4). Filtration and evaporation under vacuum, followed by purification by flash column chromatography ($\text{CH}_2\text{Cl}_2:\text{MeOH}:\text{Et}_3\text{N} = 9:1:0.05$) provided the title

compound (350 mg, 72%). ^1H NMR (400 MHz, CHCl_3) δ 3.89 (br s, 2H), 3.41 (t, J = 4.9 Hz, 2H), 3.31 (t, J = 4.9 Hz, 2H), 3.14 (t, J = 7.6 Hz, 2H), 2.44-2.41 (m, 4H), 2.33 (s, 3H), 1.64 (m, 2H), 1.32 (m, 2H), 0.95 (t, J = 7.3 Hz, 3H). MH+ 215.

5

Example 31

N-Butyl-N-[(4-methylpiperazin-1-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

Purification by preparative HPLC yielded 44 mg (22%, 2 steps) of the title compound. MH+ 400.

10

Example 32

N-Butyl-N-diphenylaminocarbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

Purification by preparative HPLC yielded 85 mg (16%, 2 steps) of the title

15

compound. ^1H NMR (400 MHz, CHCl_3) δ 9.52 (s, 1H), 8.89 (s, 1H), 8.35 (s, 1H), 7.39-7.07 (m, 10H), 4.03 (m, 1H), 3.51-3.26 (m, 3H), 2.54 (m, 1H), 1.74 (m, 1H), 1.41 (m, 1H), 1.30-1.07 (m, 10H), 0.90 (t, J = 7.3 Hz, 3H), 0.76 (t, J = 7.3 Hz, 3H). MH+ 469.

20

Example 33

N-Butyl-N-(t-butylamino)carbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3) δ 8.25 (s, 1H), 7.81 (s, 1H), 4.10-3.20 (m, 6H), 3.01 (br s, 1H), 2.42 (m, 1H), 1.55-1.20 (m, 12H), 1.27 (s, 9H), 0.88-0.85 (m, 6H). MH+

25

373.

Example 34

N-Butyl-N-phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3) δ 8.26 (s, 1H), 7.81 (s, 1H), 7.25 (br s, 1H), 4.10-3.20 (m, 6H), 3.00 (br s, 1H), 2.42 (m, 1H), 1.60-1.15 (m, 12H), 0.88-0.83 (m, 6H).
MH+ 393.

5

Example 35

N-Butyl-N-[(3,5-dimethyl-4,5-dihydro-isoxazol-4-yl)aminocarbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3) δ 9.30 (s, 1H), 8.10 (s, 1H), 7.75 (s, 1H), 3.90-3.20 (m, 6H), 2.75 (m, 1H), 2.20-1.90 (m, 6H), 1.80-1.32 (m, 12H), 0.97-0.91 (m, 6H). MH+ 414.

10

Example 36

N-Butyl-N-[(1-morpholin-4-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15 ^1H NMR (400 MHz, CDCl_3) δ 8.80 (s, 1H), 7.80 (s, 1H), 3.80-3.20 (m, 12H), 2.75 (m, 1H), 1.80-1.30 (m, 14H), 0.96-0.90 (m, 6H). MH+ 387.

Example 37

20 **N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-4-phenylbutanoyl}-hydrazine.**

MH+ 371.

Example 38

25 **N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-hexanoyl}-hydrazine.**

^1H NMR (400 MHz, CDCl_3) δ 8.80 (br s, 1H), 7.84 (s, 1H), 7.29 (m, 5H), 3.90-3.40 (m, 2H), 2.86 (m, 1H), 1.70 (m, 1H), 1.50-1.15 (m, 5H), 0.89 (m, 3H). MH+ 323.

Example 39

N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-3-phenylpropanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.90 (br s, 1H), 8.40 (s, 1H), 7.70 (s, 1H), 7.30-7.00 (m, 10H), 4.15-3.55 (m, 4H), 3.10 (m, 1H), 2.85 (m, 1H), 2.70 (m, 1H). MH+ 357.

5

Example 40

N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-3-(3,4-dichlorophenyl)propanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 9.90 (br s, 1H), 8.50 (s, 1H), 7.70 (s, 1H), 7.28-7.05 (m, 8H), 4.10-3.45 (m, 2H), 3.10 (m, 1H), 2.95 (m, 1H), 2.70 (m, 1H). MH+ 425.

10

Example 41

N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 8.90 (s, 1H), 8.40 (s, 1H), 7.95 (s, 1H), 7.81 (s, 1H), 7.41-6.95 (m, 5H), 3.90-3.40 (m, 2H), 2.83 (m, 1H), 1.61 (m, 1H), 1.42-1.12 (m, 7H), 0.87 (m, 3H). MH+ 337.

15

Example 42

N-(3,4-Dichlorophenylaminocarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 8.29 (s, 1H), 7.75 (s, 1H), 7.52 (s, 1H), 7.22-7.15 (m, 4H), 3.98-3.29 (m, 2H), 2.85-2.43 (m, 1H), 1.59 (m, 1H), 1.41-1.15 (m, 7H), 0.80 (m, 3H). MH+ 406.

20

Example 43

N-Phenylaminocarbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 8.28 (s, 1H), 7.71 (s, 1H), 7.27-6.81 (m, 5H), 3.81-3.25 (m, 2H) 2.75-2.41 (m, 1H), 1.45 (m, 1H), 1.31-1.01 (m, 7H), 0.72 (m, 3H).
MH+ 337.

5

Example 44

N-(3,4-Dichlorophenylaminocarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃) δ 10.00 (s, 1H), 8.41 (s, 1H), 8.28 (s, 1H), 8.16 (d, J = 7 Hz, 1H), 7.74 (s, 1H), 7.38-6.81 (m, 3H), 3.81-3.38 (m, 2H), 2.81-2.52 (m, 1H),
10 1.68-1.07 (m, 8H), 0.78 (m, 3H). MH+ 405.

Example 45

N-[(1-Morpholin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15 ¹H NMR (400 MHz, CDCl₃) δ 9.29 (s, 1H), 8.25 (s, 1H), 8.15 (s, 1H), 4.08-3.49 (m, 6H), 3.35 (m, 4H), 2.78-2.55 (m, 1H), 1.52 (m, 1H), 1.41-1.08 (m, 7H), 0.79 (m, 3H). MH+ 331.

Example 46

20 **N-[(2-Methoxyphenyl)aminocarbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 8.33 (s, 1H), 7.99 (s, 1H), 7.05-6.70 (m, 4H) 4.14-3.32 (m, 5H), 2.85-2.59 (m, 1H), 1.60 (m, 1H), 1.42-1.18 (m, 7H), 0.87 (m, 3H).
MH+ 367.

25

Example 47

N-[(2,4-Dichlorophenyl)aminocarbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 405.

30

Example 48

N-[(2,6-Dichlorophenyl)aminocarbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 405.

5

Example 49

N-[(4-Methyl-piperazin-1-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 344.

10

Example 50

N-[(4-Chloro-3-trifluoromethylphenyl)aminocarbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 439.

15

Example 51

N-[(Methyl-phenyl-amino)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.




MH+ 351.

20

Fourth Embodiment

As the fourth embodiment of the present invention, the compounds of
Formula (1) where X is a covalent bond are disclosed, as in the racemic compound
(55) and the chiral compounds (57) and (59). These compounds have preferentially

25 R1 = H.



Preferred compounds useful in the present invention are selected from the group consisting of:

- 5 N-[(Phenylaminocarbonyl)-carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-Butyl-N-[[1-(t-butoxycarbonyl)-piperidin-4-yl]-carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-Butyl-N-[[1-(t-butoxycarbonyl)-pyrrolidin-(2S)-yl]carbonyl]-N'-{2-
 10 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-Butyl-N-[[1-(t-butylaminocarbonyl)piperidin-4-yl]carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-Butyl-N-[[1-(t-butylcarbonyl)piperidin-4-yl]carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 15 N-Butyl-N-[(1,2,3,4-tetrahydro-quinoxalin-2-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-(p-Methoxyphenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-Phenoxyacetyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 20 N-[(p-Methoxy-phenoxy)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-(2,6-Dichlorophenyl-acetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-(3,4-Dichlorophenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-
 25 hydrazine.
 N-(Ethoxycarbonyl)carbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 N-(2,4-Dichlorophenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
 30 N-[(Benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-(2,3-Dichlorophenoxyacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3,4-Dimethoxyphenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(1H-Indol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Methyl-pyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(5-Methoxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-
- 10 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,3-Dihydro-benzo[1,4]dioxin-(2S)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(1,2,3,4-Tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Tetrahydro-furan-(2S)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Tetrahydro-furan-(2R)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
- 20 heptanoyl}-hydrazine.
- N-[(3-Methyl-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Pyridin-2-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-{3-[3-(4-Methoxybenzyl)-1H-benzoimidazol-2-yl]-propanoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Pyrimidin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Methyl-5,6,7,8-tetrahydro-[1,8]naphthyridin-3-yl)carbonyl]-N'-{(2R)-
- 30 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-[(Isoquinolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(5,6,7,8-Tetrahydro-[1,8]naphthyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(3,4-Dihydro-2H-benzo[b][1,4]dioxepin-7-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1-Methyl-2,5-dioxo-imidazolidin-4-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-phenyl-propanoyl}-hydrazine.
- N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-(3,4-dichloro)phenyl-propanoyl}-hydrazine.
- N-[(4-Imidazol-1-yl)benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[[1-Methyl-5-oxo-2-S-(pyridin-3-yl)-pyrrolidin-(3S)-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1,2-Dihydro-cinnolin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-[4-(4-Acetylpiperazin-1-yl)phenoxyacetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenylacetyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[[1-Benzyl-5-oxo-pyrrolidin-(2S)-yl]-carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[[1-Benzyl-5-oxo-pyrrolidin-(2R)-yl]-carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(5S)-Benzyl-3,6-dioxo-piperazin-(2S)-yl]acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30

- N-[(Quinolin-8-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1,2,3,4-Tetrahydroquinolin-8-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-(N"-Acetyl-L-tyrosyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1-Acetyl-1,2,3,4-tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1H-Benzoimidazol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[[1-(2-Hydroxyacetyl)-1,2,3,4-tetrahydro-quinolin-6-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1H-Indol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-{4-[Methyl-(4,6-dimethylpyrimidin-2-yl)-amino]benzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1-Benzo[1,3]dioxol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-{4-(3,5-Dimethyl-pyrazol-1-yl)methyl]benzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-[4-(Morpholin-4-yl)-benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[4-Hydroxy-3-(morpholin-4-yl)methyl-benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-(3-Hydroxy-3-methyl-butanoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Methylamino-benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1-Isopropyl-1H-benzotriazol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30

- N-[(1,2,3,4-Tetrahydro-isoquinolin-(3S)-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(5-Chloro-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- 5 N-[[1-(Dimethylaminocarbonylmethyl)-3,4-dihydro-2H-quinolin-6-yl]carbonyl]-N'-
{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,2-Difluoro-benzo[1,3]dioxol-4-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(5-Amino-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
10 heptanoyl}-hydrazine.
- N-[(4-Oxo-1,2-dihydro-4H-pyrrolo[3,2,1-ij]quinolin-5-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(7-Hydroxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- 15 N-[(6-Methoxy-benzofuran-2-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-[(5-Acetamidobenzofuran-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,3-Dihydro-benzo[1,4]dioxin-6-yl)carbonyl]-N'-{(2R)-
20 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(3-Amino-4,6-dimethyl-furo[2,3-b]pyridin-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Methyl-5,6,7,8-tetrahydro-[1,6]naphthyridin-3-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(6-Fluoro-4H-benzo[1,3]dioxin-8-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(7-Amino-1H-indol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-[(1-Methyl-1,2,3,4-tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxy
30 amino)methyl]-heptanoyl}-hydrazine.

- N'-[(6,7,9,10,12,13,15,16-Octahydro-5,8,11,14,17-pentaoxa-benzocyclopentadecen-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(2-Benzo[1,3]dioxol-5-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Pentanoyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Benzoyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Trifluoroacetamido-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[(3-Hydroxy-naphthalen-2-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenylacetyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Furan-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Methoxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(1H-Indol-3-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Dimethylaminobenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-(2-Hydroxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Piperidin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(1,2,5,6-Tetrahydro-pyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(7-Methoxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(3-Chloro-4-methoxy-phenyl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30 N-[(1H-Pyrrol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-[(Quinolin-7-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Pyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-(4-Chloro-3-methoxy-benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3-Methoxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-
10 hydrazine.
- N-[(5-Methyl-2-phenyl-oxazol-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinoxalin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-4-phenylbutanoyl}-hydrazine.
- N-[(3-Methoxy-quinoxalin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,6-Dimethoxypyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
20 heptanoyl}-hydrazine.
- N-[(N"-Methylsulfonyl)-L-tyrosyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[[5-Oxo-pyrrolidin-(2S)-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(4-(Pyrrol-1-yl)benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Acetamidobenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(3-Cyclopentyloxy-4-methoxy)benzoyl]-N'-{(2R)-
30 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-cyclopentyl-propanoyl}-hydrazine.
- N-[(7-Methoxy-benzofuran-2-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-3-cyclopentyl-propanoyl}-hydrazine.
- 5 N-[3-(Morpholin-4-yl)propanoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,3-Dihydro-benzofuran-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(4,6-Dimethoxy-pyrimidin-2-yl)benzoyl]-N'-{(2R)-
- 10 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Trifluoromethyl-5,6,7,8-tetrahydro-naphthyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(9H-beta-Carbolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15


The following synthetic schemes are merely illustrative of the methods by which the compounds of the invention may be prepared and are not intended to limit the scope of the invention as defined in the appended claims.

As shown in Scheme 9, treatment of an alkyl halide R1X (26) with hydrazine

20 in a solvent such as ethanol, at an elevated temperature, gives the hydrazine derivative (27). Reacting compound (27) with the acid chloride R2C(O)Cl affords intermediate (50). Alternatively, compound (50) can be prepared from the Boc-protected hydrazine (29) by reaction with the aldehyde or ketone R'COR", followed by reduction with hydrogen gas in the presence of palladium, to afford hydrazine

25 derivative (30). Reacting hydrazine (30) with the acid chloride R2C(O)Cl, followed by removal of the Boc protecting group with an appropriate acid, such as trifluoroacetic acid, gives the hydrazine (50) wherein R1 = CHR'R". Alternatively, ester (51) can be submitted to hydrazinolysis to afford (50) wherein R1 = H. Alternatively, acid (52) can be reacted with ethyl chloroformate to form the


30 intermediate mixed anhydride (53), which upon hydrazinolysis yields (50) wherein R1 = H.



Scheme 9.


5 As shown in Scheme 10, coupling of the carboxylic acid (8) with the hydrazine derivative (50) provides acylated hydrazide (54). Hydrogenolysis to remove the benzyl group using a catalyst, such as 10% Pd/C, in an appropriate solvent, such as ethanol, gives compound (55). Similarly, coupling of the chiral acid (17) or (25) with hydrazine derivative (50) provides the corresponding hydrazide (56) or (58). Hydrogenolysis of the benzyl group gives the final compounds (57) or (59).

10



Scheme 10.

- 5 Alternatively, as shown in Scheme 11, the carbamate (34) wherein R2 = t-butyl or benzyl can be converted to the hydrazide (47) by acid treatment or hydrogenolysis, respectively. Reaction of (47) with the acyl chloride R2C(O)Cl or the mixed anhydride R2C(O)OC(O)OEt in an appropriate solvent, such as methylene chloride, and in the presence of an optional appropriate base, such as triethylamine,
- 10 gives compound (55). Similarly, appropriate deprotection of the chiral carbamates (36) and (38), followed by reaction with an acyl chloride or a mixed anhydride, gives the chiral acylated hydrazide (57) or (59).



Scheme 11.

SYNTHETIC EXAMPLES

The invention will now be described by reference to the following examples
5 which are merely illustrative and are not to be construed as a limitation of the scope
of the present invention. The same experimental general conditions and conventions
described in the Experimental Section of the Second Embodiment are applicable here.

The compounds disclosed in Examples 52 to 165 were prepared following
the general procedures described in Example 1.

10

Example 52

**N-[(Phenylaminocarbonyl)-carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 9.50 (s, 1H), 9.20 (s, 1H), 7.85 (s, 1H), 7.70-7.00 (m,
15 5H), 3.75 (m, 1H), 3.40 (m, 1H), 2.88 (m, 1H), 1.63 (m, 1H), 1.39-1.15 (m, 7H),
0.79 (t, J = 6.8 Hz, 3H). MH+ 365.

Example 53

**N-Butyl-N-[[1-(t-butoxycarbonyl)-piperidin-4-yl]-carbonyl]-N'-{2-
20 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃): δ 9.14 (br s, 1H), 8.33 (br s, 1H), 7.81 (br s, 1H),
4.30-3.20 (m, 8H), 2.80-2.50 (m, 2H), 1.92-1.22 (m, 14H), 1.47 (s, 9H), 0.97-0.90
(m, 6H). MH+ 485.

25

Example 54

**N-Butyl-N-[[1-(t-butoxycarbonyl)-pyrrolidin-(2S)-yl]carbonyl]-N'-{2-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃) δ 9.70 (br s, 1H), 8.21 (s, 1H), 7.80 (s, 1H), 4.51-3.20
(m, 6H), 2.86 (m, 1H), 2.18-1.41 (m, 6H), 1.38 (s, 9H), 1.38-1.17 (m, 10H), 0.88 (m,
30 6H). MH+ 471.

Example 55

N-Butyl-N-[(1-t-butylaminocarbonyl)piperidin-4-yl]carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.67 (br s, 1H), 8.38 (s, 1H), 4.15-3.47 (m, 8H),
5 2.89 (m, 1H), 2.67 (m, 1H), 1.75-1.26 (m, 16H), 1.37 (s, 9H), 0.99-0.90 (m, 6H).
MH+ 484.

Example 56

N-Butyl-N-[(1-t-butylcarbonyl)piperidin-4-yl]carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.30 (s, 1H), 8.25 (s, 1H), 7.80 (s, 1H), 4.39-3.34
10 (m, 8H), 2.95 (m, 1H), 2.79 (m, 1H), 1.89-1.25 (m, 16H), 1.29 (s, 9H), 0.99-0.90 (m,
6H). MH+ 469.

Example 57

N-Butyl-N-[(1,2,3,4-tetrahydro-quinoxalin-2-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.28 (s, 1H), 8.20 (s, 1H), 7.80 (s, 1H), 6.70-6.51
15 (m, 4H), 4.16-3.22 (m, 6H), 2.87 (m, 1H), 2.75 (m, 1H), 1.64-1.26 (m, 12H), 0.97-
20 0.87 (m, 6H). MH+ 434.

Example 58

N-(p-Methoxyphenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.20 (s, 1H), 8.30 (s, 1H), 7.50 (s, 1H), 7.24 (d, J =
25 16.3 Hz, 2H), 6.90 (d, J = 16.3 Hz, 2H), 3.95-3.38 (m, 4H), 3.77 (s, 3H), 2.83 (m,
1H), 1.64 (m, 1H), 1.34 (m, 1H), 1.29 (m, 6H), 0.86 (t, J = 6.3 Hz, 3H). MH+ 366.

Example 59

N-Phenoxyacetyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 352.

5

Example 60

N-[(p-Methoxy-phenoxy)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.39 (s, 1H), 8.53 (s, 1H), 7.48 (s, 1H), 6.79-6.73 (m, 4H), 4.49 (m, 2H), 3.73 (s, 3H), 3.44 (m, 2H), 2.82 (m, 1H), 1.64 (m, 1H), 1.25 (m, 1H), 1.24-1.19 (m, 6H), 0.80 (t, J = 6.7 Hz, 3H). MH+ 382.

10

Example 61

N-(2,6-Dichlorophenyl-acetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.35 (s, 1H), 7.80 (s, 1H), 7.30-7.10 (m, 3H), 4.09 (s, 2H), 3.85-3.50 (m, 2H), 2.90 (m, 1H), 1.80-1.28 (m, 6H), 0.89 (br s, 3H). MH+ 405.

15

Example 62

N-(3,4-Dichlorophenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.30 (s, 1H), 7.91 (s, 1H), 7.20-7.10 (m, 3H), 3.45-3.10 (m, 2H), 3.25 (br s, 2H), 1.85 (m, 1H), 1.62 (m, 1H), 1.45 (m, 1H), 1.32 (m, 6H), 0.91 (br s, 3H). MH+ 405.

20

Example 63

N-(Ethoxycarbonyl)carbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.96 (s, 1H), 9.27 (s, 1H), 7.77 (s, 1H), 4.37 (m, 2H), 3.90 (m, 1H), 3.51 (m, 1H), 2.96 (m, 1H), 1.71 (m, 1H), 1.41 (m, 1H), 1.40-1.32 (m, 9H), 0.91 (br s, 3H). MH+ 318.

5

Example 64

N-(2,4-Dichlorophenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.70 (s, 1H), 7.80 (s, 1H), 7.50-7.20 (m, 3H), 3.82 (m, 2H), 3.55 (m, 2H), 2.85 (m, 1H), 1.85 (m, 1H), 1.61 (m, 1H), 1.45-1.22 (m, 6H), 0.85 (br s, 3H). MH+ 405.

10

Example 65

N-[(Benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.87 (s, 1H), 8.68 (s, 1H), 7.78 (s, 1H), 7.54-7.06 (m, 5H), 4.15-3.30 (m, 2H), 2.95 (m, 1H), 1.85 (m, 1H), 1.62 (m, 1H), 1.27-1.18 (m, 6H), 0.84 (br s, 3H). MH+ 362.

15

Example 66

N-(2,3-Dichlorophenoxyacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.17 (s, 1H), 7.88 (br s, 1H), 7.28-7.14 (m, 3H), 4.73 (m, 2H), 3.76 (m, 1H), 3.49 (m, 1H), 2.90 (m, 1H), 1.74 (m, 1H), 1.41-1.28 (m, 7H), 0.90 (br s, 3H). MH+ 421.

20

Example 67

N-(3,4-Dimethoxyphenylacetyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3): δ 8.92 (s, 1H), 7.58 (s, 1H), 6.83 (m, 3H), 3.87 (s, 3H), 3.85 (s, 3H), 3.74-3.44 (m, 4H), 1.65 (m, 1H), 1.36 (m, 1H), 1.29-1.23 (m, 6H), 0.88 (t, $J = 6.5$ Hz, 3H). MH^+ 396.

5

Example 68

N-[(1H-Indol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3): δ 9.51 (s, 1H), 8.50 (s, 1H), 7.81 (s, 1H), 7.50-6.80 (m, 5H), 4.20-3.45 (m, 2H), 2.95 (m, 1H), 1.82 (m, 1H), 1.75-1.20 (m, 7H), 1.00 (br s, 3H). MH^+ 396.

10

Example 69

N-[(2-Methyl-pyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

15 ^1H NMR (400 MHz, CDCl_3): δ 8.50 (s, 1H), 7.80 (s, 1H), 7.70-7.00 (m, 3H), 3.84 (m, 1H), 3.50 (m, 1H), 2.58 (s, 1H), 1.73 (m, 1H), 1.44 (m, 1H), 1.43-1.27 (m, 6H), 0.89 (br s, 3H). MH^+ 337.

Example 70

20 **N-[(5-Methoxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

^1H NMR (400 MHz, CDCl_3): δ 9.50 (s, 1H), 8.50 (s, 1H), 7.85 (s, 1H), 7.50-6.80 (m, 4H), 3.90 (m, 1H), 3.80 (s, 3H), 3.55 (m, 1H), 3.00 (m, 1H), 1.85 (m, 1H), 1.60-1.25 (m, 7H), 0.95 (br s, 3H). MH^+ 392.

25

Example 71

N-[(2,3-Dihydro-benzo[1,4]dioxin-(2S)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CD_3OD) δ 8.50 (s, 1H), 7.80 (s, 1H), 7.30 (m, 4H), 4.55-3.20 (m, 5H), 2.85 (m, 1H), 1.85 (m, 1H), 1.65-1.24 (m, 7H), 0.93 (t, $J = 6.2$ Hz, 3 H). MH+ 380.

5

Example 72

N-[(Quinolin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CD_3OD): δ 8.51-7.72 (m, 6H), 3.88 (m, 1H), 3.61 (m, 1H), 2.99 (m, 1H), 1.69 (m, 1H), 1.58-1.26 (m, 7H), 0.97 (t, $J = 6.7$ Hz, 3H). MH+ 373.

10

Example 73

N-[(1,2,3,4-Tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CD_3OD): δ 8.36 (s, 1H), 7.48 (br s, 2H), 6.45 (d, $J = 9.04$ Hz, 1H), 3.96-3.32 (m, 4H), 2.91 (m, 1H), 2.77 (t, $J = 6.12$ Hz, 2H), 1.98 (m, 2H), 1.65 (m, 1H), 1.49 (m, 1H), 1.36 (m, 6H), 0.94 (br s, 3H). MH+ 377.

15

Example 74

N-[(Tetrahydro-furan-(2S)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CD_3OD): δ 8.32 (s, 1H), 4.44 (m, 1H), 4.03 (m, 1H), 3.88 (m, 3H), 3.50 (m, 1H), 2.85 (m, 1H), 2.35-1.93 (m, 4H), 1.49 (m, 1H), 1.40 (m, 1H), 1.35 (m, 6H), 0.92 (t, $J = 6.9$ Hz, 3H). MH+ 316.

20

25

Example 75

N-[(Tetrahydro-furan-(2R)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CD_3OD): δ 8.32 (s, 1H), 4.43 (m, 1H), 4.03 (m, 1H), 3.87 (m, 3H), 3.55 (m, 1H), 2.86 (m, 1H), 2.28-1.93 (m, 4H), 1.63 (m, 1H), 1.50 (m, 1H), 1.34 (m, 6H), 0.93 (br s, 3H). MH+ 316.

30

Example 76

N-[(3-Methyl-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 5 ^1H NMR (400 MHz, CDCl_3): δ 8.36 (s, 1H), 7.54-7.20 (m, 4H), 4.04 (m, 1H), 3.48 (m, 1H), 2.97 (s, 3H), 2.59 (m, 1H), 1.66 (m, 1H), 1.28 (m, 1H), 1.27 (m, 6H), 0.83 (t, J = 6.7 Hz, 3H). MH+ 376.

Example 77

- 10 **N-[(Pyridin-2-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

^1H NMR (400 MHz, CD_3OD): δ 8.48 (s, 1H), 7.90-7.33 (m, 4H), 4.00-3.50 (m, 4H), 2.87 (m, 1H), 1.62 (m, 1H), 1.46 (m, 1H), 1.33 (m, 6H), 0.90 (br s, 3H). MH+ 337.

15

Example 78

N-{3-[3-(4-Methoxybenzyl)-1H-benzoimidazol-2-yl]-propanoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 20 ^1H NMR (400 MHz, CD_3OD): δ 8.31 (s, 1H), 7.61 (m, 1H), 7.42 (m, 1H), 7.25 (m, 2H), 7.10 (d, J = 8.6 Hz, 2H), 6.88 (d, J = 8.6 Hz, 2H), 5.46 (s, 2H), 3.77 (s, 3H), 3.90-3.50 (m, 2H), 3.23 (m, 2H), 2.85 (m, 2H), 2.69 (m, 1H), 1.65 (m, 1H), 1.61 (m, 1H), 1.33 (m, 6H), 0.92 (m, 3H). MH+ 510.

Example 79

- 25 **N-[(Pyrimidin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

^1H NMR (400 MHz, CD_3OD): δ 9.26 (s, 1H), 8.84 (s, 1H), 8.72 (s, 1H), 7.95 (s, 1H), 3.85-3.40 (m, 2H), 2.86 (m, 1H), 1.67 (m, 1H), 1.47 (m, 1H), 1.37 (m, 6H), 0.94 (t, J = 6.8 Hz, 3H). MH+ 324.

30

Example 80

N-[(2-Methyl-5,6,7,8-tetrahydro-[1,8]naphthyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.92 (s, 1H), 7.44 (s, 1H), 3.90-3.60 (m, 2H), 3.43 (m, 2H), 2.94 (m, 1H), 2.75 (m, 2H), 2.47 (s, 3H), 1.98 (m, 2H), 1.66 (m, 1H), 1.45 (m, 1H), 1.37 (m, 6H), 0.94 (br s, 3H). MH+ 392.

Example 81

N-[(Isoquinolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 9.17 (s, 1H), 8.48 (s, 1H), 8.03-7.73 (m, 5H), 4.03-3.99 (m, 1H), 3.59 (m, 1H), 2.70 (m, 1H), 1.71 (m, 1H), 1.43 (m, 1H), 1.26 (m, 6H), 0.86 (t, J = 6.7 Hz, 3H). MH+ 373.

15

Example 82

N-[(5,6,7,8-Tetrahydro-[1,8]naphthyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.93 (br s, 1H), 7.31 (d, J = 7.2 Hz, 1H), 7.21 (d, J = 7.31 Hz, 1H), 3.82 (m, 1H), 3.55 (m, 1H), 3.43 (br s, 2H), 2.94 (m, 1H), 2.80 (br s, 2H), 1.92 (m, 2H), 1.65 (m, 1H), 1.60-1.09 (m, 7H), 0.94 (br s, 3H). MH+ 378.

20

Example 83

N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.86 (s, 1H), 7.60 (s, 1H), 7.33 (m, 5H), 3.90-3.42 (m, 4H), 2.83 (m, 1H), 1.67 (m, 1H), 1.32-1.20 (m, 5H), 0.89 (t, J = 6.7 Hz, 3H). MH+ 322.

25

Example 84

N-[(3,4-Dihydro-2H-benzo[b][1,4]dioxepin-7-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

30

¹H NMR (400 MHz, CDCl₃): δ 8.45 (d, J = 9.2 Hz, 1H), 7.82 (s, 1H), 7.45-7.28 (m, 2H), 4.30-3.40 (m, 6H), 2.96 (m, 1H), 2.22 (m, 2H), 1.70 (m, 1H), 1.55 (m, 1H), 1.35-1.23 (m, 6H), 0.88 (t, J = 6.7 Hz, 3H). MH+ 394.

5

Example 85

N-[(1-Methyl-2,5-dioxo-imidazolidin-4-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.90 (s, 1H), 4.39 (m, 1H), 3.84-3.50 (m, 2H), 2.97 (s, 3H), 2.85 (m, 1H), 2.67 (m, 1H), 1.62 (m, 1H), 1.49 (m, 1H), 1.34 (m, 6H), 0.93 (br s, 3H). MH+ 372.

10

Example 86

N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-phenyl-propanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.80 (br s, 1H), 8.10 (s, 1H), 7.22 (m, 10H), 3.96-3.43 (m, 4H), 3.15 (m, 1H), 3.02 (m, 1H), 2.70 (m, 1H). MH+ 356.

15

Example 87

N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-(3,4-dichloro)phenyl-propanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.36 (br s, 1H), 7.60 (s, 1H), 7.37-7.00 (m, 8H), 4.17-3.51 (m, 4H), 3.05 (m, 1H), 2.95 (m, 1H), 2.65 (m, 1H). MH+ 424.

20

Example 88

N-[(4-Imidazol-1-yl)benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.37-7.71 (m, 7H), 7.21 (s, 1H), 3.87 (m, 1H), 3.71-3.57 (m, 1H), 2.95 (m, 1H), 1.67 (m, 1H), 1.54 (m, 1H), 1.37 (m, 6H), 0.95 (t, J = 6.7 Hz, 3H). MH+ 388.

25
30

Example 89

N-[[1-Methyl-5-oxo-2-S-(pyridin-3-yl)-pyrrolidin-(3S)-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.57 (m, 1H), 7.90 (s, 1H), 7.84 (m, 1H), 7.55 (m, 1H), 3.79 (m, 1H), 3.53 (m, 1H), 3.08 (m, 1H), 2.92-2.69 (m, 3H), 2.68 (s, 3H), 1.62 (m, 1H), 1.47 (m, 1H), 1.35 (m, 6H), 0.93 (br s, 3H). MH+ 420.

Example 90

N-[(1,2-Dihydro-cinnolin-4-yl)carbonyl]-N'-{(2R)-

[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.91 (s, 1H), 7.22 (m, 2H), 6.98 (m, 1H), 6.81 (d, J = 7.8 Hz, 1H), 6.64 (m, 1H), 4.36 (br s, 1H), 3.79 (m, 1H), 3.53 (m, 1H), 2.87 (m, 1H), 1.63 (m, 1H), 1.48 (m, 1H), 1.34 (m, 6H), 0.92 (br s, 3H). MH+ 376.

Example 91

N-[4-(4-Acetylpiperazin-1-yl)phenoxyacetyl]-N'-{(2R)-

[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.92 (s, 1H), 4.60 (br s, 2H), 3.90-3.50 (m, 6H), 3.07 (m, 4H), 3.04 (m, 1H), 2.16 (s, 3H), 1.62 (m, 1H), 1.51 (m, 1H), 1.35 (m, 6H), 0.93 (br s, 3H). MH+ 478.

Example 92

N-Phenylacetyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.30 (s, 1H), 7.90 (s, 1H), 7.32 (br s, 5H), 3.84-3.47 (m, 4H), 2.84 (m, 1H), 1.62 (m, 1H), 1.44 (m, 1H), 1.32 (m, 6H), 0.90 (br s, 3H). MH+ 336.

Example 93

N-[[1-Benzyl-5-oxo-pyrrolidin-(2S)-yl]-carbonyl]-N'-{(2R)-

[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 9.20 (s, 1H), 7.63 (s, br s, 1H), 7.13 (m, 5H), 4.92 (m, 1H), 3.94-3.31 (m, 4H), 2.82 (m, 1H), 2.60-1.90 (m, 4H), 1.49 (m, 1H), 1.23 (m, 1H), 1.11 (m, 6H), 0.85 (t, J = 6.7 Hz, 3H). MH+ 419.

5

Example 94

N-{{1-Benzyl-5-oxo-pyrrolidin-(2R)-yl}-carbonyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 9.45 (br s, 1H), 7.82 (s, 1H), 7.30 (m, 5H), 5.06 (m, 1H), 4.10-3.50 (m, 4H), 2.95 (m, 1H), 2.85-2.10 (m, 4H), 1.67 (m, 1H), 1.41-1.26 (m, 7H), 0.87 (t, J = 6.7 Hz, 3H). MH+ 419.

10

Example 95

N-{{(5S)-Benzyl-3,6-dioxo-piperazin-(2S)-yl}acetyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.90 (s, 1H), 7.38-7.23 (m, 5H), 4.34-4.21 (m, 2H), 3.82 (m, 1H), 3.49 (m, 1H), 3.31 (m, 1H), 3.06 (m, 1H), 2.85 (m, 1H), 1.57 (m, 1H), 1.46 (m, 1H), 1.34 (m, 6H), 0.92 (t, J = 6.7 Hz, 3H). MH+ 462.

15

Example 96

N-{{(Quinolin-4-yl)carbonyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 9.11 (br s, 1H), 8.30-7.40 (m, 6H), 3.90 (m, 1H), 3.51 (m, 1H), 3.12 (m, 1H), 1.69 (m, 1H), 1.41 (m, 1H), 1.30 (m, 6H), 0.86 (t, J = 6.7 Hz, 3H). MH+ 373.

20

Example 97

N-{{(Quinolin-8-yl)carbonyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 9.03 (m, 1H), 8.66 (t, J = 6.3 Hz, 1H), 8.46 (m, 1H), 8.05 (d, J = 8.1 Hz, 1H), 7.84 (s, 1H), 7.67 (m, 2H), 3.84-3.30 (m, 2H), 1.66 (m, 1H), 1.43-1.18 (m, 7H), 0.82 (t, J = 6.7 Hz, 3H). MH+ 373.

5

Example 98

N-[(1,2,3,4-Tetrahydroquinolin-8-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.83 (s, 1H), 7.26 (d, J = 8.0 Hz, 1H), 6.91 (m, 1H), 6.34 (m, 1H), 3.71 (m, 1H), 3.50 (m, 1H), 3.26 (t, J = 5.6 Hz, 2H), 2.81 (m, 10 1H), 2.66 (t, J = 6.2 Hz, 2H), 1.77 (m, 2H), 1.55 (m, 1H), 1.40 (m, 1H), 1.38 (m, 1H), 1.25 (m, 6H), 0.83 (t, J = 6.7 Hz, 3H). MH+ 377.

Example 99

N-(N''-Acetyl-L-tyrosyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-15 hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.91 (s, 1H), 7.09 (d, J = 8.4 Hz, 2H), 6.71 (d, J = 8.4 Hz, 2H), 4.62 (m, 1H), 3.87 (m, 1H), 3.57 (m, 1H), 3.12 (dd, J = 14.0, 4.6 Hz, 1H), 2.85 (m, 2H), 2.72 (m, 1H), 1.91 (s, 3H), 1.65 (m, 1H), 1.48 (m, 1H), 1.34 (m, 6H), 0.92 (t, J = 6.7 Hz, 3H). MH+ 423.

20

Example 100

N-[(1-Acetyl-1,2,3,4-tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxy amino) methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.75 (m, 3H), 3.85 (m, 1H), 3.82 (t, 25 J = 6.4 Hz, 2H), 3.56 (m, 1H), 2.95 (m, 1H), 2.83 (t, J = 6.5 Hz, 2H), 2.05 (s, 3H), 2.00 (m, 2H), 1.65 (m, 1H), 1.52 (m, 1H), 1.37 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 419.

Example 101

N-[(1H-Benzoimidazol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.95 (br s, 1H), 7.70 (br s, 2H), 7.32 (br s, 2H), 3.98-3.50 (m, 2H), 2.98 (m, 1H), 1.69 (m, 1H), 1.55 (m, 1H), 1.45 (m, 6H), 0.88 (t, J = 6.7 Hz, 3H). MH+ 362.

Example 102

N-[[1-(2-Hydroxyacetyl)-1,2,3,4-tetrahydro-quinolin-6-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

10 ¹H NMR (400 MHz, CD₃OD): δ 7.81 (s, 1H), 7.70-7.50 (m, 3H), 4.27 (s, 2H), 3.75 (m, 1H), 3.60 (m, 2H), 3.55 (m, 1H), 2.81 (m, 1H), 2.72 (m, 2H), 2.85 (m, 2H), 3.55 (m, 1H), 2.81 (m, 1H), 2.72 (m, 2H), 2.85 (m, 2H), 1.55 (m, 1H), 1.45 (m, 1H), 1.25 (m, 6H), 0.82 (t, J = 6.7 Hz, 3H). MH+ 435.

15

Example 103

N-[(1H-Indol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.25 (br s, 1H), 7.95 (s, 1H), 7.68 (d, J = 8.5 Hz, 1H), 7.46 (m, 1H), 7.35 (br s, 1H), 6.59 (br s, 1H), 3.95 (m, 1H), 3.59 (m, 1H), 2.96 (m, 1H), 1.66 (m, 1H), 1.54 (m, 1H), 1.36 (m, 6H), 0.95 (t, J = 6.7 Hz, 3H). MH+ 361.

Example 104

N-{4-[Methyl-(4,6-dimethylpyrimidin-2-yl)-amino]benzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

25 ¹H NMR (400 MHz, CD₃OD): δ 7.96 (s, 1H), 7.88 (m, 2H), 7.46 (m, 2H), 6.57 (s, 1H), 3.91 (m, 1H), 3.62 (m, 1H), 3.57 (s, 3H), 2.96 (m, 1H), 2.29 (s, 6H), 1.68 (m, 1H), 1.52 (m, 1H), 1.37 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 457.

30

Example 105

N-[(1-Benzo[1,3]dioxol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.94 (s, 1H), 7.49 (m, 1H), 7.36 (s, 1H), 6.92 (dd, J = 8.2, 2.5 Hz, 1H), 6.06 (s, 2H), 3.92 (m, 1H), 3.58 (m, 1H), 2.93 (m, 1H), 1.63 (m, 1H), 1.49 (m, 1H), 1.38 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 366.

Example 106

N-[[4-(3,5-Dimethyl-pyrazol-1-yl)methyl]benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

10 ¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.85 (d, J = 8.2 Hz, 2H), 7.16 (d, J = 8.2 Hz, 2H), 5.97 (s, 1H), 5.33 (s, 2H), 3.88 (m, 1H), 3.57 (m, 1H), 2.95 (m, 1H), 2.22 (s, 3H), 2.20 (s, 3H), 1.59 (m, 1H), 1.44 (m, 1H), 1.26 (m, 6H), 0.93 (t, J = 6.7 Hz, 3H). MH+ 430.

15

Example 107

N-[4-(Morpholin-4-yl)-benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.81 (d, J = 6.6 Hz, 2H), 7.00 (d, J = 6.6 Hz, 2H), 3.96 (m, 1H), 3.84 (m, 4H), 3.60 (m, 1H), 3.28 (m, 4H), 2.94 (m, 1H), 1.67 (m, 1H), 1.53 (m, 1H), 1.37 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 407.

20

Example 108

N-[4-Hydroxy-3-(morpholin-4-yl)methyl-benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

25 ¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.85 (d, J = 8.4 Hz, 2H), 6.85 (d, J = 8.4 Hz, 2H), 3.85 (m, 2H), 3.80 (m, 1H), 3.74 (m, 4H), 3.55 (m, 1H), 2.93 (m, 1H), 2.68 (br s, 4H), 1.66 (m, 1H), 1.50 (m, 1H), 1.38 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 444.

30

Example 109

N-(3-Hydroxy-3-methyl-butanoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 3.82 (m, 1H), 3.55 (m, 1H), 2.86 (m, 1H), 2.42 (d, J = 5.7 Hz, 2H), 1.56 (m, 1H), 1.44 (m, 1H), 1.43-1.29 (m, 15 H), 0.93 (t, J = 6.7 Hz, 3H). MH+ 318.

Example 110

N-(4-Methylamino-benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

10 ¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.73 (d, J = 7.7 Hz, 2H), 6.60 (d, J = 7.7 Hz, 2H), 3.96 (m, 1H), 3.58 (m, 1H), 2.93 (m, 1H), 2.83 (s, 3H), 1.52 (m, 1H), 1.38 (m, 1H), 1.26 (m, 6H), 0.94 (t, J 6.7 Hz, 3H). MH+ 351.

Example 111

15 **N-[(1-Isopropyl-1H-benzotriazol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 8.75 (s, 1H), 8.09 (d, J = 8.7 Hz, 1H), 7.96 (s, 1H), 7.93 (s, 1H), 5.26 (m, 1H), 3.86 (m, 1H), 3.65 (m, 1H), 2.96 (m, 1H), 1.75 (d, J = 6.8 Hz, 6H), 1.70 (m, 1H), 1.56 (m, 1H), 1.28 (m, 6H), 0.95 (t, J = 6.7 Hz, 3H). MH+ 405.

Example 112

N-[(1,2,3,4-Tetrahydro-isoquinolin-(3S)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

25 ¹H NMR (400 MHz, CD₃OD): δ 7.90 (s, 1H), 7.20 (m, 4H), 4.30-3.40 (m, 5H), 3.10 (m, 1H), 2.90 (m, 1H), 2.72 (m, 1H), 1.65 (m, 1H), 1.50 (m, 1H), 1.36 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 377.

Example 113

**N-[(5-Chloro-benzofuran-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.79 (s, 1H), 7.62 (d, J = 8.8 Hz, 1H),
7.57 (d, J = 5.1 Hz, 1H), 7.49 (d, J = 8.8 Hz, 1H), 3.86 (m, 1H), 3.55 (m, 1H), 3.95
5 (m, 1H), 1.65 (m, 1H), 1.55 (m, 1H), 1.36 (m, 6H), 0.95 (t, J = 6.8 Hz, 3H). MH+
396.

Example 114

**N-{{[1-(Dimethylaminocarbonylmethyl)-3,4-dihydro-2H-quinolin-6-
10 yl]carbonyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 7.94 (br s, 1H), 7.55 (d, J = 8.3 Hz, 1H), 7.51 (s,
1H), 6.38 (d, J = 8.3 Hz, 1H), 4.28 (s, 2H), 3.95 (m, 1H), 3.63 (m, 1H), 3.41 (t, J =
5.4 Hz, 2H), 3.13 (s, 3H), 2.98 (s, 3H), 2.97 (m, 1H), 2.83 (t, J = 5.4 Hz, 2H), 1.99
15 (m, 4H), 1.65 (m, 1H), 1.50 (m, 1H), 1.36 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+
462.

Example 115

**N-[(2,2-Difluoro-benzo[1,3]dioxol-4-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.59 (m, 1H), 7.42 (m, 1H), 7.30 (m,
1H), 3.85 (m, 1H), 3.54 (m, 1H), 2.95 (m, 1H), 1.64 (m, 1H), 1.54 (m, 1H), 1.38 (m,
20 6H), 0.94 (t, J 6.7 Hz, 3H). MH+ 402.

Example 116

**25 N-[(5-Amino-benzofuran-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.35 (m, 2H), 6.96 (m, 2H), 3.84 (m,
1H), 3.54 (m, 1H), 2.82 (m, 1H), 1.67 (m, 1H), 1.38 (m, 1H), 1.37 (m, 6H), 0.95 (t, J
= 6.7 Hz, 3H). MH+ 377.

30

Example 117

N-[(4-Oxo-1,2-dihydro-4H-pyrrolo[3,2,1-ij]quinolin-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.90 (br s, 1H), 7.95 (s, 1H), 7.66 (m, 1H), 7.58 (m, 1H), 7.34 (m, 1H), 4.52 (br s, 2H), 3.85 (m, 1H), 3.54 (m, 1H), 3.52 (m, 2H), 2.92 (m, 1H), 1.64 (m, 1H), 1.36-1.24 (m, &H), 0.94 (t, J = 6.8 Hz, 3H). MH+ 415.

Example 118

N-[(7-Hydroxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-

10 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.55 (m, 1H), 7.21-7.13 (m, 2H), 6.92 (m, 1H), 3.89 (m, 1H), 3.68 (m, 1H), 2.92 (m, 1H), 1.66 (m, 1H), 1.54 (m, 1H), 1.37 (m, 6H), 0.94 (t, J = 6.8 Hz, 3H). MH+ 378.

15

Example 119

N-[(6-Methoxy-benzofuran-2-yl)acetyl]-N'-{(2R)-

[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.91 (s, 1H), 7.63 (s, 1H), 7.53 (d, J = 8.6 Hz, 1H), 7.05 (d, J = 1.9 Hz, 1H), 6.88 (dt, J = 8.6, 2.2 Hz, 1H), 3.84 (s, 3H), 3.76 (m, 1H), 3.64 (d, J = 7.2 Hz, 2H), 3.53 (m, 1H), 2.86 (m, 1H), 1.65 (m, 1H), 1.48 (m, 1H), 1.26 (m, 1H), 0.92 (t, J = 6.7 Hz, 3H). MH+ 406.

20

Example 120

N-[(5-Acetamidobenzofuran-2-yl)carbonyl]-N'-{(2R)-

25 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.36 (s, 1H), 8.04 (br s, 1H), 7.54 (m, 3H), 3.85 (m, 1H), 3.54 (m, 1H), 2.94 (m, 1H), 2.17 (s, 3H), 1.54 (m, 1H), 1.52 (m, 1H), 1.37 (m, 6H), 0.95 (t, J = 6.8 Hz, 3H). MH+ 419.

30

Example 121

N-[(2,3-Dihydro-benzo[1,4]dioxin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.94 (s, 1H), 7.42 (m, 2H), 6.92 (m, 1H), 4.31 (m, 4H), 3.85 (m, 1H), 3.55 (m, 1H), 2.93 (m, 1H), 1.65(m, 1H), 1.51 (m, 1H), 1.37 (m, 5
6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 380.

Example 122

N-[(3-Amino-4,6-dimethyl-furo[2,3-b]pyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

10 ¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.02 (s, 1H), 3.97 (m, 1H), 3.60 (m, 1H), 2.93 (m, 1H), 2.70 (s, 3H), 2.57 (s, 3H), 1.66 (m, 1H), 1.52 (m, 1H), 1.38 (m, 6H), 0.95 (t, J = 6.7 Hz, 3H). MH+ 406.

Example 123

15 **N-[(2-Methyl-5,6,7,8-tetrahydro-[1,6]naphthyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.69 (br s, 1H), 3.71 (m, 1H), 3.45 (m, 1H), 3.05 (m, 2H), 2.90 (m, 2H), 2.68 (m, 2H), 2.50 (s, 3H), 1.65 (m, 1H), 1.48 (m, 1H), 1.36 (m, 6H), 0.95 (t, J = 6.7 Hz, 3H). MH+ 392.

20

Example 124

N-[(6-Fluoro-4H-benzo[1,3]dioxin-8-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

25 ¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.55 (dd, J = 9.2, 2.8 Hz, 1H), 7.07 (dd, J = 7.8, 3.4 Hz, 1H), 5.40 (s, 2H), 4.97 (s, 2H), 3.86 (m, 1H), 3.65 (m, 1H), 2.95 (m, 1H), 1.67 (m, 1H), 1.51 (m, 1H), 1.26 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 398.

Example 125

N-[(7-Amino-1H-indol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.96 (s, 1H), 7.12 (d, J = 6.3 Hz, 1H), 7.03 (d, J = 8.1 Hz, 1H), 6.89 (m, 1H), 6.60 (m, 1H), 3.90 (m, 1H), 3.55 (m, 1H), 2.95 (m, 1H),
5 1.67 (m, 1H), 1.51 (m, 1H), 1.26 (m, 6H), 0.94 (t, J = 6.7 Hz, 3H). MH+ 376.

Example 126

N-[(1-Methyl-1,2,3,4-tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxy amino) methyl]-heptanoyl}-hydrazine.

10 ¹H NMR (400 MHz, CD₃OD): δ 7.94 (s, 1H), 7.61 (d, J = 8.7 Hz, 1H), 7.49 (s, 1H), 6.59 (d, J = 8.7 Hz, 1H), 3.85 (m, 1H), 3.55 (m, 1H), 3.36 (m, 2H), 2.97 (s, 3H), 2.80 (t, J = 6.3 Hz, 2H), 1.96 (m, 2H), 1.66 (m, 1H), 1.51 (m, 1H), 1.26 (m, 6H), 0.93 (t, J = 6.7 Hz, 3H). MH+ 391.

15

Example 127

N'-[(6,7,9,10,12,13,15,16-Octahydro-5,8,11,14,17-pentaoxa-benzocyclopentadecen-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 7.50 (m, 1H), 7.49 (s, 1H), 7.01 (dd, J 8.4, 3.1 Hz, 1H), 4.18 (m, 4H), 3.89 (m, 4H), 3.86 (m, 1H), 3.73 (br s, 8H), 3.56 (m, 1H), 2.96 (m, 1H), 1.67 (m, 1H), 1.51 (m, 1H), 1.26 (m, 6H), 0.93 (t, J = 6.7 Hz, 3H). MH+ 512.

Example 128

25 **N-[(2-Benzo[1,3]dioxol-5-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CD₃OD): δ 7.95 (s, 1H), 6.85 (s, 1H), 6.77 (m, 2H), 5.93 (s, 2H), 3.84 (m, 1H), 3.56 (m, 1H), 3.49 (d, J = 6.2 Hz, 2H), 2.86 (m, 1H), 1.60 (m, 1H), 1.48 (m, 1H), 1.34 (m, 6H), 0.92 (t, J = 6.9 Hz, 3H). MH+ 380.

30

Example 129**N-Pentanoyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃): δ 9.88 (m, 1H), 9.46 (s, 1H), 9.05 (d, J = 3.5 Hz, 2H), 7.75 (s, 1H), 3.98-3.43 (m, 2H), 2.88 (m, 1H), 2.28 (m, 2H), 1.67 (m, 2H), 1.51-1.22 (m, 10H), 0.89 (m, 3H). MH+ 302.

Example 130**N-Benzoyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃): δ 8.31 (s, 1H), 7.75 (m, 2H), 7.51-7.20 (m, 3H), 3.86-3.27 (m, 2H), 3.03 (s, 2H), 2.82-2.65 (m, 1H), 1.62 (m, 1H), 1.41-1.12 (m, 7H), 0.89 (m, 3H). MH+ 322.

Example 131**N-Trifluoroacetamido-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-****hydrazine.**

¹H NMR (400 MHz, CDCl₃): δ 9.98 (s, 1H), 9.08 (s, 1H), 4.25-3.45 (m, 2H), 2.91 (m, 1H), 1.69 (m, 1H), 1.61-1.21 (m, 7H), 0.88 (m, 3H). MH+ 314.

Example 132

20 N-[(3-Hydroxy-naphthalen-2-yl)carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.48 (s, 1H), 7.97 (s, 1H), 7.81 (m, 1H), 7.68 (m, 1H), 7.52 (m, 1H), 7.35-7.21 (m, 1H), 3.90-3.51 (m, 2H) 3.02 (m, 1H), 1.67 (m, 1H), 1.59-1.21 (m, 7H), 0.88 (m, 3H). MH+ 388.

25

Example 133**N-Phenylacetyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃): δ 9.67 (s, 1H), 9.33 (s, 1H), 8.93 (s, 1H), 8.21 (s, 1H), 7.15 (m, 5H), 3.72-3.25 (m, 6H) 2.75-2.50 (m, 1H), 1.52 (m, 1H), 1.32-1.09 (m, 7H), 0.78 (m, 3H). MH+ 336.

30

Example 134

N-[(Furan-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 5 ^1H NMR (400 MHz, CDCl_3): δ 8.40 (s, 1H), 7.78 (s, 1H), 7.31 (s, 1H), 7.21-7.11 (m, 3H), 6.35 (m, 1H), 5.31 (s, 1H), 4.15-3.43 (m, 2H) 3.00-2.65 (m, 1H), 1.75 (m, 1H), 1.62-1.20 (m, 7H), 0.90 (m, 3H). MH+ 312.

Example 135

- 10 **N-(4-Methoxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

^1H NMR (400 MHz, CDCl_3): δ 7.75 (m, 1H), 6.89 (d, 1H), 6.72 (d, 1H), 4.05-3.40 (m, 5H) 3.00-2.58 (m, 1H), 1.82-1.11 (m, 8H), 0.88 (m, 3H). MH+ 352.

- 15 **Example 136**

N-[(1H-Indol-3-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

^1H NMR (400 MHz, CDCl_3): δ 8.28 (s, 1H), 7.48 (m, 1H), 7.36-6.94 (m, 4H), 3.89-3.19 (m, 4H) 2.71-2.42 (m, 1H), 2.02-1.04 (m, 8H), 0.76 (m, 3H). MH+ 375.

20

Example 137

N-(4-Dimethylaminobenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- 25 ^1H NMR (400 MHz, CDCl_3): δ 8.40(s, 1H), 7.71 (m, 2H), 6.61 (m, 2H), 4.05-3.38 (m, 2H) 3.02 (d, J = 8 Hz, 6H), 2.93-2.55 (m, 1H), 1.68 (m, 1H), 1.52-1.21 (m, 7H), 0.89 (m, 3H). MH+ 365.

Example 138

- 30 **N-(2-Hydroxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**

¹H NMR (400 MHz, CDCl₃): δ 8.37 (s, 1H), 7.82-7.57 (m, 2H), 7.41-6.81 (m, 2H), 6.65 (m, 1H), 3.95-3.35 (m, 2H) 3.00-2.57 (m, 1H), 1.75-1.05 (m, 8H), 0.89 (m, 3H). MH+ 338.

5

Example 139

N-[(Piperidin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 8.38 (s, 1H), 7.81 (s, 1H), 4.04-3.32 (m, 5H), 3.12-2.72 (m, 2H), 2.62-2.28 (m, 2H), 2.05-1.85 (m, 5H), 1.60 (m, 1H) 1.50-1.18 (m, 7H), 0.85 (m, 3H). MH+ 329.

10

Example 140

N-[(1,2,5,6-Tetrahydro-pyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CDCl₃): δ 10.50 (s, 1H), 8.42 (s, 1H), 7.85 (m, 1H), 7.60-7.12 (m, 2H), 4.75 (m, 1H), 4.11 (m, 1H), 3.81-3.20 (m, 2H) 2.85-2.50 (m, 1H), 2.35-2.20 (m, 2H), 1.90 (m, 1H), 1.81-1.15 (m, 8H), 0.81 (m, 3H). MH+ 327.

15

Example 141

N-[(7-Methoxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 7.94 (s, 1H), 7.55 (m, 1H), 7.30 (m, 1H), 7.05 (m, 1H), 4.01 (s, 3H), 3.52-3.89 (m, 2H), 2.99 (M, 1H), 1.65 (m, 1H), 1.53 (m, 1H), 1.40 (m, 6H), 0.95 (m, 3H). MH+ 392.

25

Example 142

N-[(3-Chloro-4-methoxy-phenyl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
MH+ 400.

30

Example 143

N-[(1H-Pyrrol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 311.

5

Example 144

N-[(Quinolin-7-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 373.

10

Example 145

N-[(Pyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 323.

15

Example 146

N-(4-Chloro-3-methoxy-benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 386.

20

Example 147

N-(3-Methoxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 352.

25

Example 148

N-[(Quinolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 373.

30

Example 149

**N-[(5-Methyl-2-phenyl-oxazol-4-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**
MH+ 403.

5

Example 150

**N-[(Quinoxalin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.**
MH+ 374.

10

Example 151

**N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-4-phenylbutanoyl}-
hydrazine.**
MH+ 370.

15

Example 152

**N-[(3-Methoxy-quinoxalin-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**
MH+ 404.

20

Example 153

**N-[(2,6-Dimethoxypyridin-3-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**
MH+ 383.

25

Example 154

**N-[(N''-Methylsulfonyl)-L-tyrosyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.**
MH+ 459.

30

Example 155

**N-{{5-Oxo-pyrrolidin-(2S)-yl}carbonyl}-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**
MH+ 329.

5

Example 156

**N-{{4-(Pyrrol-1-yl)benzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.**
MH+ 387.

10

Example 157

**N-{{4-Acetamidobenzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-
hydrazine.**
MH+ 379.

15

Example 158

**N-{{3-Cyclopentyloxy-4-methoxy}benzoyl}-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.**
MH+ 436.

20

Example 159

**N-{{Phenylacetyl}-N'-{2-[(formylhydroxyamino)methyl]-3-cyclopentyl-
propanoyl}-hydrazine.**
MH+ 348.

25

Example 160

**N-{{7-Methoxy-benzofuran-2-yl}carbonyl}-N'-{2-
[(formylhydroxyamino)methyl]-3-cyclopentyl-propanoyl}-hydrazine.**
MH+ 404.

30

Example 161

N-[3-(Morpholin-4-yl)propanoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 359.

5

Example 162

N-[(2,3-Dihydro-benzofuran-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 364.

10

Example 163

N-[(4,6-Dimethoxy-pyrimidin-2-yl)benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 460.

15

Example 164

N-[(2-Trifluoromethyl-5,6,7,8-tetrahydro-naphthyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

MH+ 446.

20

Example 165

N-[(9H-beta-Carbolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

¹H NMR (400 MHz, CD₃OD): δ 8.20 (m, 1H), 8.59 (m, 1H), 8.05 (m, 1H), 7.80 (s, 1H), 7.46 (m, 2H), 7.16 (m, 1H), 7.75 (m, 1H), 3.43 (m, 1H), 2.89 (m, 1H), 1.56 (m, 1H), 1.43 (m, 1H), 1.30 (m, 6H), 0.83 (t, J = 6.7 Hz, 3H). MH+ 412.

25

COMPOSITIONS, ADMINISTRATION AND BIOLOGICAL ASSAYS

Compounds of Formula (1) and their pharmaceutically acceptable salts may be administered in a standard manner for antibiotics, for example orally, parenterally, sub-lingually, dermally, transdermally, rectally, via inhalation or via buccal administration.

30

Compositions of Formula (1) and their pharmaceutically acceptable salts which are active when given orally can be formulated as syrups, tablets, capsules, creams and lozenges. A syrup formulation will generally consist of a suspension or solution of the compound or salt in a liquid carrier for example, ethanol, peanut oil, olive oil, glycerine or water with a flavoring or coloring agent. Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include magnesium stearate, terra alba, talc, gelatin, acacia, stearic acid, starch, lactose and sucrose. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example, using the aforementioned carriers in a hard gelatin capsule shell. Where the composition is in the form of a soft gelatin shell capsule, any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example, aqueous gums, celluloses, silicates or oils, and incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of a compound or salt in a sterile aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil, for example, polyethylene glycol, polyvinylpyrrolidone, lecithin, arachis oil or sesame oil.

Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be administered as a dry powder or in the form of an aerosol using a conventional propellant such as dichlorodifluoromethane or trichlorofluoromethane.

A typical suppository formulation comprises a compound of Formula (1) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent, for example, polymeric glycols, gelatins, cocoa-butter or other low melting vegetable waxes or fats or their synthetic analogs.

Typical dermal and transdermal formulations comprise a conventional aqueous or non-aqueous vehicle, for example, a cream, ointment, lotion or paste or are in the form of a medicated plaster, patch or membrane.

Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer a single dose.

Each dosage unit for oral administration contains suitably from 0.1 mg to 500 mg/Kg, and preferably from 1 mg to 100 mg/Kg, and each dosage unit for parenteral administration contains suitably from 0.1 mg to 100 mg/Kg, of a compound of Formula (1) or a pharmaceutically acceptable salt thereof calculated as the free acid. Each dosage unit for intranasal administration contains suitably 1-400 mg and preferably 10 to 200 mg per person. A topical formulation contains suitably 0.01 to 5.0% of a compound of Formula (1).

The daily dosage regimen for oral administration is suitably about 0.01 mg/Kg to 40 mg/Kg of a compound of Formula (1) or a pharmaceutically acceptable salt thereof calculated as the free acid. The daily dosage regimen for parenteral administration is suitably about 0.001 mg/Kg to 40 mg/Kg of a compound of Formula (1) or a pharmaceutically acceptable salt thereof calculated as the free acid. The daily dosage regimen for intranasal administration and oral inhalation is suitably about 10 to about 500 mg/person. The active ingredient may be administered from 1 to 6 times a day, sufficient to exhibit the desired activity.

No unacceptable toxicological effects are expected when compounds of the present invention are administered in accordance with the present invention.

The biological activity of the compounds of Formula (1) are demonstrated by the following test:

Biological Assay


S. aureus or *E. coli* PDF activity is measured at 25°C, using a continuous enzyme-linked assay developed by Lazennec & Meinel ("Formate dehydrogenase-coupled spectrophotometric assay of peptide deformylase", *Anal. Biochem.* 1997, 244, pp.180-182), with minor modifications. The reaction mixture is contained in 50 μ L with 50 mM potassium phosphate buffer (pH 7.6), 15 mM NAD, 0.25 U formate dehydrogenase. The substrate peptide, f-Met-Ala-Ser, is included at the K_M concentration. The reaction is triggered with the addition of 10 nM Def1 enzyme, and absorbance is monitored for 20 min at 340 nm.

Antimicrobial Activity Assay

Whole-cell antimicrobial activity was determined by broth microdilution using the National Committee for Clinical Laboratory Standards (NCCLS) recommended procedure, Document M7-A4, "Methods for Dilution Susceptibility Tests for Bacteria that Grow Aerobically" (incorporated by reference herein). The compound was tested in serial two-fold dilutions ranging from 0.06 to 64 mcg/ml. A panel of 12 strains were evaluated in the assay. This panel consisted of the following laboratory strains: Staphylococcus aureus Oxford, Staphylococcus aureus WCUH29, Enterococcus faecalis I, Enterococcus faecalis 7, Haemophilus influenzae Q1, Haemophilus influenzae NEMC1, Moraxella catarrhalis 1502, Streptococcus pneumoniae 1629, Streptococcus pneumoniae N1387, Streptococcus pneumoniae N1387, E. coli 7623 (AcrABEFD+) and E. coli 120 (AcrAB-). The minimum inhibitory concentration (MIC) was determined as the lowest concentration of compound that inhibited visible growth. A mirror reader was used to assist in determining the MIC endpoint.

What is claimed is:

1. A compound according to Formula (1):



5

(1) X = O, NR₃ or a bond;

Y = O, CH₂ or a bond

wherein:

R represents:

10 C₂₋₆ alkyl (optionally substituted by alkoxy, halogen, or C₁₋₃ alkylsulfanyl),
 C₂₋₆ alkenyl (optionally substituted by alkoxy, halogen, or C₁₋₃
 alkylsulfanyl), C₂₋₆ alkynyl (optionally substituted by alkoxy, halogen, or C₁₋₃
 alkylsulfanyl), (CH₂)_n—C₃₋₆ carbocycle (optionally substituted by alkoxy,
 halogen, or C₁₋₃ alkylsulfanyl), (CH₂)_n—R₄ {where R₄ is phenyl, furan,
 benzofuran, thiophene, benzothiophene, tetrahydrofuran, tetrahydropyran,
 15 dioxane, 1,4-benzodioxane or benzo[1,3]dioxole; R₄ is optionally substituted
 by one or more Cl, Br, I, C₁₋₃ alkyl (optionally substituted by one to three F)
 or C₁₋₂ alkoxy (optionally substituted by one to three F)};

R₁ represents:

20 hydrogen, C₁₋₆ alkyl (optionally substituted by hydroxy, halogen, amino,
 guanidino, phenyl, pyridyl, pyrrolyl, indolyl, imidazolyl, furanyl,
 benzofuranyl, piperidinyl, morpholinyl, quinolinyl, piperazinyl or
 dimethylaminophenyl) or (CH₂)_n—C₃₋₇ carbocycle;

R₂ represents:

25 hydrogen (provided that X is not O), C₁₋₃ substituted alkyl, C₂₋₃ substituted
 alkenyl, C₂₋₃ substituted alkynyl, (CH₂)_n—C₃₋₆ substituted carbocycle, aryl,
 heteroaryl, heterocyclic, carboxy (provided that X is not NR₃ or O) or
 aminocarbonyl (provided that X is not NR₃ or O);

R₃ represents:

hydrogen, C₁₋₃ substituted alkyl, phenyl, or may be taken together with R₂ and the nitrogen atom to which they are attached to form an optionally substituted heterocyclic ring which is optionally fused to an aryl, a heteroaryl, or a second heterocyclic ring;


5 X represents O, NR₃ or a covalent bond;

Y represents O, CH₂ or a covalent bond;

n = 0-2;

or a salt, solvate, or physiologically functional derivative thereof.

10 2. A compound as claimed in claim 1, with the following absolute configuration:



X = O, NR₃ or a bond;

Y = O, CH₂ or a bond

15 or a salt, solvate or physiologically functional derivative thereof.

3. A compound as claimed in claim 2, wherein R₁ = H; or a salt, solvate or physiologically functional derivative thereof.

20 4. A compound as claimed in claim 1, wherein X = O; or a salt, solvate, or physiologically functional derivative thereof.

5. A compound according to claim 4 selected from the group consisting of:

25 N-Butyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

N-Butyl-N-phenoxy carbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-Isobutyl-N-(t-butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Isobutyl-N-phenoxy carbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-Phenethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Cyclohexylmethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Benzyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-(3-pyridin-3-yl-propyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(2-Morpholin-4-yl-ethyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-(4-Hydroxy-butyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Amino-butyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(Tetrahydro-pyran-4-yl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-Methyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3-Aminopropyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-(t-Butoxycarbonyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3-Hydroxypropyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-(t-butoxycarbonyl)-N'-{(2S)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30

- N-Butyl-N-(phenoxy carbonyl)-N'-{(2S)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[2-(4-Dimethylaminophenyl)ethyl]-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxy amino)methyl]-heptanoyl}-hydrazine.
- 5 N-(t-Butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Pentyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[2-(1H-Indol-3-yl)-ethyl]-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-Isopentyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Cyclohexyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-(1-Ethyl-propyl)-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Isopropyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Propyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-Ethyl-N-(t-butoxycarbonyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Methoxycarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[[1-(3,5-Dimethoxyphenyl)-1-methyl-ethoxy]carbonyl]-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
6. A compound as claimed in claim 1, wherein X = NR₃; or a salt, solvate, or physiologically functional derivative thereof.
- 30 7. A compound according to claim 6 selected from the group consisting of:

- N-Butyl-N-[(4-methylpiperazin-1-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-Butyl-N-diphenylaminocarbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-Butyl-N-(t-butylamino)carbonyl-N'-{2-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-Butyl-N-phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- 10 N-Butyl-N-[(3,5-dimethyl-4,5-dihydro-isoxazol-4-yl)aminocarbonyl]-N'-{2-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-[(1-morpholin-4-yl)carbonyl]-N'-{2-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-4-phenyl-
15 butanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-hexanoyl}-
hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-3-phenyl-
propanoyl}-hydrazine.
- 20 N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-3-(3,4-
dichlorophenyl)-propanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-
hydrazine.
- N-(3,4-Dichlorophenylaminocarbonyl)-N'-{2-
25 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenylaminocarbonyl-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-(3,4-Dichlorophenylaminocarbonyl)-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30 N-[(1-Morpholin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.

- N-[(2-Methoxyphenyl)aminocarbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,4-Dichlorophenyl)aminocarbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(2,6-Dichlorophenyl)aminocarbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(4-Methyl-piperazin-1-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(4-Chloro-3-trifluoromethylphenyl)aminocarbonyl]-N'-{(2R)-
 10 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Methyl-phenyl-amino)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
8. A compound as claimed in claim 1, wherein X is a covalent bond; or a salt,
 15 solvate, or physiologically functional derivative thereof.
9. A compound according to claim 8 selected from the group consisting of:
- N-[(Phenylaminocarbonyl)-carbonyl]-N'-{(2R)-
 20 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-{{[1-(t-butoxycarbonyl)-piperidin-4-yl]-carbonyl}}-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-{{[(1-t-butoxycarbonyl)-pyrrolidin-(2S)-yl]carbonyl}}-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-Butyl-N-{{[(1-t-butylaminocarbonyl)piperidin-4-yl]carbonyl}}-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-{{[(1-t-butylcarbonyl)piperidin-4-yl]carbonyl}}-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Butyl-N-[(1,2,3,4-tetrahydro-quinoxalin-2-yl)carbonyl]-N'-{2-
 30 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-(p-Methoxyphenylacetyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenoxyacetyl-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(p-Methoxy-phenoxy)acetyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(2,6-Dichlorophenyl-acetyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3,4-Dichlorophenylacetyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-(Ethoxycarbonyl)carbonyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(2,4-Dichlorophenylacetyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(Benzofuran-2-yl)carbonyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(2,3-Dichlorophenoxyacetyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3,4-Dimethoxyphenylacetyl)-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-[(1H-Indol-2-yl)carbonyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Methyl-pyridin-3-yl)carbonyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(5-Methoxy-benzofuran-2-yl)carbonyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,3-Dihydro-benzo[1,4]dioxin-(2S)-yl)carbonyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-2-yl)carbonyl]-N²-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30

- N-[(1,2,3,4-Tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Tetrahydro-furan-(2S)-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(Tetrahydro-furan-(2R)-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(3-Methyl-benzofuran-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[(Pyridin-2-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-[3-[3-(4-Methoxybenzyl)-1H-benzoimidazol-2-yl]-propanoyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Pyrimidin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- 15 N-[(2-Methyl-5,6,7,8-tetrahydro-[1,8]naphthyridin-3-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Isoquinolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- 20 N-[(5,6,7,8-Tetrahydro-[1,8]naphthyridin-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-
hydrazine.
- N-[(3,4-Dihydro-2H-benzo[b][1,4]dioxepin-7-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(1-Methyl-2,5-dioxo-imidazolidin-4-yl)acetyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-phenyl-
propanoyl}-hydrazine.
- 30 N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-(3,4-
dichloro)phenyl-propanoyl}-hydrazine.

- N-[(4-Imidazol-1-yl)benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[[1-Methyl-5-oxo-2-S-(pyridin-3-yl)-pyrrolidin-(3S)-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(1,2-Dihydro-cinnolin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[4-(4-Acetylpiperazin-1-yl)phenoxyacetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Phenylacetyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[[1-Benzyl-5-oxo-pyrrolidin-(2S)-yl]-carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[[1-Benzyl-5-oxo-pyrrolidin-(2R)-yl]-carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(5S)-Benzyl-3,6-dioxo-piperazin-(2S)-yl]acetyl-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-8-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-[(1,2,3,4-Tetrahydroquinolin-8-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(N^m-Acetyl-L-tyrosyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(1-Acetyl-1,2,3,4-tetrahydroquinolin-6-yl)carbonyl]-N'-{(2R)-[(formylhydroxy amino) methyl]-heptanoyl}-hydrazine.
- N-[(1H-Benzoimidazol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[[1-(2-Hydroxyacetyl)-1,2,3,4-tetrahydroquinolin-6-yl]carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30

- N-[(1H-Indol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-{4-[Methyl-(4,6-dimethylpyrimidin-2-yl)-amino]benzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine).
- 5 N-[(1-Benzo[1,3]dioxol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-{[4-(3,5-Dimethyl-pyrazol-1-yl)methyl]benzoyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[4-(Morpholin-4-yl)-benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[4-Hydroxy-3-(morpholin-4-yl)methyl-benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3-Hydroxy-3-methyl-butanoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-(4-Methylamino-benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1-Isopropyl-1H-benzotriazol-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1,2,3,4-Tetrahydro-isoquinolin-(3S)-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-[(5-Chloro-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-{[1-(Dimethylaminocarbonylmethyl)-3,4-dihydro-2H-quinolin-6-yl]carbonyl}-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-[(2,2-Difluoro-benzo[1,3]dioxol-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(5-Amino-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30 N-[(4-Oxo-1,2-dihydro-4H-pyrrolo[3,2,1-ij]quinolin-5-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-[(7-Hydroxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(6-Methoxy-benzofuran-2-yl)acetyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(5-Acetamidobenzofuran-2-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2,3-Dihydro-benzo[1,4]dioxin-6-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[(3-Amino-4,6-dimethyl-furo[2,3-b]pyridin-2-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Methyl-5,6,7,8-tetrahydro-[1,6]naphthyridin-3-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(6-Fluoro-4H-benzo[1,3]dioxin-8-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(7-Amino-1H-indol-2-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1-Methyl-1,2,3,4-tetrahydro-quinolin-6-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxy amino) methyl]-heptanoyl}-hydrazine.
- 20 N'-[(6,7,9,10,12,13,15,16-Octahydro-5,8,11,14,17-pentaoxa-
 benzocyclopentadecen-2-yl)carbonyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(2-Benzo[1,3]dioxol-5-yl)acetyl]-N'-{(2R)-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Pentanoyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-Benzoyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-Trifluoroacetamido-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-
 hydrazine.
- N-[(3-Hydroxy-naphthalen-2-yl)carbonyl]-N'-{2-
 [(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30 N-Phenylacetyl-N'-{2-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

- N-[(Furan-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Methoxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 5 N-[(1H-Indol-3-yl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(4-Dimethylaminobenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(2-Hydroxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[(Piperidin-4-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1,2,5,6-Tetrahydro-pyridin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(7-Methoxy-benzofuran-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(3-Chloro-4-methoxy-phenyl)acetyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(1H-Pyrrol-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-[(Quinolin-7-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Pyridin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 25 N-(4-Chloro-3-methoxy-benzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-(3-Methoxybenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinolin-3-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30

- N-[(5-Methyl-2-phenyl-oxazol-4-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(Quinoxalin-2-yl)carbonyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- 5 N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-4-phenylbutanoyl}-
hydrazine.
- N-[(3-Methoxy-quinoxalin-2-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 10 N-[(2,6-Dimethoxypyridin-3-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- N-[(N"-Methylsulfonyl)-L-tyrosyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-[5-Oxo-pyrrolidin-(2S)-yl]carbonyl}-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 15 N-[(4-(Pyrrol-1-yl)benzoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-(4-Acetamidobenzoyl)-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-[(3-Cyclopentyloxy-4-methoxy)benzoyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 20 N-(Phenylacetyl)-N'-{2-[(formylhydroxyamino)methyl]-3-cyclopentyl-
propanoyl}-hydrazine.
- N-[(7-Methoxy-benzofuran-2-yl)carbonyl]-N'-{2-
[(formylhydroxyamino)methyl]-3-cyclopentyl-propanoyl}-hydrazine.
- 25 N-[3-(Morpholin-4-yl)propanoyl]-N'-{(2R)-[(formylhydroxyamino)methyl]-
heptanoyl}-hydrazine.
- N-[(2,3-Dihydro-benzofuran-5-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.
- 30 N-[(4,6-Dimethoxy-pyrimidin-2-yl)benzoyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

N-[(2-Trifluoromethyl-5,6,7,8-tetrahydro-naphthyridin-2-yl)carbonyl]-N'-
{(2R)-[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

N-[(9H-beta-Carbolin-3-yl)carbonyl]-N'-{(2R)-
[(formylhydroxyamino)methyl]-heptanoyl}-hydrazine.

5

10. A method of treating a bacterial infection by administering to a subject in need of treatment a compound according to claim 1.