



US 20080171041A1

(19) **United States**(12) **Patent Application Publication**
Thompson et al.(10) **Pub. No.: US 2008/0171041 A1**(43) **Pub. Date: Jul. 17, 2008**(54) **METHOD FOR TREATING INFLAMMATION**(75) Inventors: **Penny Thompson**, Snohomish, WA (US); **Hal Blumberg**, Seattle, WA (US); **Yasmin A. Chandrasekher**, Mercer Island, WA (US); **Julia E. Novak**, Bainbridge Island, WA (US)Correspondence Address:
ZYMOGENETICS, INC.
INTELLECTUAL PROPERTY DEPARTMENT
1201 EASTLAKE AVENUE EAST
SEATTLE, WA 98102-3702(73) Assignee: **ZymoGenetics, Inc.**(21) Appl. No.: **12/043,046**(22) Filed: **Mar. 5, 2008****Related U.S. Application Data**


(60) Continuation of application No. 11/458,666, filed on Jul. 19, 2006, now Pat. No. 7,364,732, which is a continuation of application No. 10/424,658, filed on Apr. 28, 2003, now Pat. No. 7,189,394, which is a

division of application No. 09/746,359, filed on Dec. 22, 2000, now Pat. No. 6,610,286.

(60) Provisional application No. 60/171,969, filed on Dec. 23, 1999, provisional application No. 60/213,341, filed on Jun. 22, 2000.

Publication Classification(51) **Int. Cl.**
A61K 39/395 (2006.01)
A61P 29/00 (2006.01)
(52) **U.S. Cl.** **424/135.1; 424/158.1**(57) **ABSTRACT**

A method for treating IL-20 induced inflammation. An antagonist to IL-20 is administered to treat inflammation and associated diseases. The antagonist can be an antibody that binds to IL-20 or its receptor or a soluble receptor that binds to IL-20. Examples of such diseases are adult respiratory disease, psoriasis, eczema, contact dermatitis, atopic dermatitis, septic shock, multiple organ failure, inflammatory lung injury, bacterial pneumonia, inflammatory bowel disease, rheumatoid arthritis, asthma, ulcerative colitis and Crohn's disease.






FIG.1




FIG.2




FIG.3




FIG.4




FIG.5




FIG.6




FIG.7




FIG.8

METHOD FOR TREATING INFLAMMATION

REFERENCE TO RELATED APPLICATIONS

[0001] The present application is a continuation application of U.S. application Ser. No. 11/458,666, filed Jul. 19, 2006, which is a continuation application of U.S. application Ser. No. 10/424,658, filed Apr. 28, 2003, now U.S. Pat. No. 7,189,394, which is a divisional application of U.S. application Ser. No. 09/746,359, filed Dec. 22, 2000, now U.S. Pat. No. 6,610,286, which claims priority under 35 U.S.C. §119 (e) to U.S. Provisional Patent Application. 60/171,969, filed Dec. 23, 1999 and U.S. Provisional Application No. 60/213,341 filed Jun. 22, 2000.

BACKGROUND OF THE INVENTION

[0002] The teachings of all of the references cited herein are incorporated in their entirety herein by reference.

[0003] Inflammation normally is a localized, protective response to trauma or microbial invasion that destroys, dilutes, or walls-off the injurious agent and the injured tissue. It is characterized in the acute form by the classic signs of pain, heat, redness, swelling, and loss of function. Microscopically, it involves a complex series of events, including dilation of arterioles, capillaries, and venules, with increased permeability and blood flow, exudation of fluids, including plasma proteins, and leukocyte migration into the area of inflammation.

[0004] Diseases characterized by inflammation are significant causes of morbidity and mortality in humans. Commonly, inflammation occurs as a defensive response to invasion of the host by foreign, particularly microbial, material. Responses to mechanical trauma, toxins, and neoplasia also may result in inflammatory reactions. The accumulation and subsequent activation of leukocytes are central events in the pathogenesis of most forms of inflammation. Deficiencies of inflammation compromise the host. Excessive inflammation caused by abnormal recognition of host tissue as foreign or prolongation of the inflammatory process may lead to inflammatory diseases as diverse as diabetes, arteriosclerosis, cataracts, reperfusion injury, and cancer, to post-infectious syndromes such as in infectious meningitis, rheumatic fever, and to rheumatic diseases such as systemic lupus erythematosus and rheumatoid arthritis. The centrality of the inflammatory response in these varied disease processes makes its regulation a major element in the prevention control or cure of human disease.

[0005] An important cytokine in the inflammatory process is interleukin-8 (IL-8). IL-8 is a chemokine. It was identified as an agonist for neutrophils on the basis of two effects, chemotaxis and the release of granule enzymes. IL-8 binds to two receptors on neutrophils. IL-8 receptors are also found on monocytes, basophils, and eosinophils. In human fibroblasts, cytomegalovirus has been shown to induce the expression of IL-8 receptors and to replicate more rapidly when cells are exposed to IL-8. IL-8 is a potent chemoattractant for neutrophils; and the early stages of periodontal disease are characterized by the influx of neutrophils. IL-8 is a potent inducer of angiogenesis in several angiogenesis-dependent chronic inflammatory conditions, including rheumatoid arthritis, psoriasis, and idiopathic pulmonary fibrosis. Additionally, IL-8 is an important source of angiogenic activity in human lung cancer. Also, IL-8 expression correlates with experimental metastatic activity of some melanoma cell lines. Thus, an

effective method to treat inflammatory diseases would be to administer an agent that would inhibit IL-8.

BRIEF DESCRIPTION OF THE DRAWINGS

[0006] FIGS. 1-8 are schematic representations of a representative number of embodiments of the IL-20 soluble receptor.

DESCRIPTION OF THE INVENTION

[0007] The present invention fills this need by providing for a method for treating inflammation in which IL-20 plays a role, comprising administering to a mammal in need of treatment of inflammation an antagonist to IL-20 or an antagonist to the IL-20 receptor. The antagonist to IL-20 can be an antibody, antibody fragment or single-chain antibody that binds to IL-20, a soluble receptor that binds to IL-20. The antagonist to the IL-20 receptor can be an antibody, antibody fragment, single-chain antibody or small molecule that binds to the IL-20 receptor. Also an anti-sense nucleotide that binds to the mRNA that encodes IL-20 can be used as an antagonist. We have shown that IL-20 up-regulates IL-8. Inflammatory diseases in which IL-8 plays a significant role, and for which a decrease in IL-8 would be beneficial are, adult respiratory disease (ARD), septic shock, multiple organ failure, inflammatory lung injury such as asthma or bronchitis, bacterial pneumonia, psoriasis and inflammatory bowel disease such as ulcerative colitis and Crohn's disease, eczema, atopic dermatitis and contact dermatitis. Thus, antagonists to IL-20 can be used to treat these diseases.

[0008] The teachings of all the references cited herein are incorporated in their entirety by reference.

DEFINITIONS

[0009] Prior to setting forth the invention in detail, it may be helpful to the understanding thereof to define the following terms.

[0010] The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides. Where the context allows, these terms are used with reference to a particular sequence or portion of a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a polypeptide is located proximal to the carboxyl terminus of the reference sequence, but is not necessarily at the carboxyl terminus of the complete polypeptide.

[0011] The term "complement/anti-complement pair" denotes non-identical moieties that form a non-covalently associated, stable pair under appropriate conditions. For instance, biotin and avidin (or streptavidin) are prototypical members of a complement/anti-complement pair. Other exemplary complement/anti-complement pairs include receptor/ligand pairs, antibody/antigen (or hapten or epitope) pairs, sense/antisense polynucleotide pairs, and the like. Where subsequent dissociation of the complement/anti-complement pair is desirable, the complement/anti-complement pair preferably has a binding affinity of $<10^9 \text{ M}^{-1}$.

[0012] The term "complements of a polynucleotide molecule" is a polynucleotide molecule having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATG-CACGGG 3' is complementary to 5' CCCGTGCAT 3'.

[0013] The term "contig" denotes a polynucleotide that has a contiguous stretch of identical or complementary sequence

to another polynucleotide. Contiguous sequences are said to “overlap” a given stretch of polynucleotide sequence either in their entirety or along a partial stretch of the polynucleotide. For example, representative contigs to the polynucleotide sequence 5'-ATGGCTTAGCTT-3' are 5'-TAGCTTgagtct-3' and 3'-gtcgacTACCGA-5'.

[0014] The term “degenerate nucleotide sequence” denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

[0015] The term “expression vector” is used to denote a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

[0016] The term “isolated”, when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, *Nature* 316:774-78 (1985)).

[0017] An “isolated” polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. In a preferred form, the isolated polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, i.e. greater than 95% pure, more preferably greater than 99% pure. When used in this context, the term “isolated” does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

[0018] The term “operably linked”, when referring to DNA segments, indicates that the segments are arranged so that they function in concert for their intended purposes, e.g., transcription initiates in the promoter and proceeds through the coding segment to the terminator.

[0019] A “polynucleotide” is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized in vitro, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs (abbreviated “bp”), nucleotides (“nt”), or kilobases (“kb”). Where the context allows, the latter two terms may describe polynucleotides that are single-stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term “base pairs”. It will be recognized by those skilled in the art that the two strands of a

double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nucleotides in length.

[0020] A “polypeptide” is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as “peptides”.

[0021] The term “promoter” is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

[0022] A “protein” is a macromolecule comprising one or more polypeptide chains. A protein may also comprise non-peptidic components, such as carbohydrate groups. Carbohydrates and other non-peptidic substituents may be added to a protein by the cell in which the protein is produced, and will vary with the type of cell. Proteins are defined herein in terms of their amino acid backbone structures; substituents such as carbohydrate groups are generally not specified, but may be present nonetheless.

[0023] The term “receptor” denotes a cell-associated protein that binds to a bioactive molecule (i.e., a ligand) and mediates the effect of the ligand on the cell. Membrane-bound receptors are characterized by a multi-domain structure comprising an extracellular ligand-binding domain and an intracellular effector domain that is typically involved in signal transduction. Binding of ligand to receptor results in a conformational change in the receptor that causes an interaction between the effector domain and other molecule(s) in the cell. This interaction in turn leads to an alteration in the metabolism of the cell. Metabolic events that are linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids. In general, receptors can be membrane bound, cytosolic or nuclear, monomeric (e.g., thyroid stimulating hormone receptor, beta-adrenergic receptor) or multimeric (e.g., PDGF receptor, growth hormone receptor, IL-3 receptor, GM-CSF receptor, G-CSF receptor, erythropoietin receptor and IL-6 receptor).

[0024] The term “secretory signal sequence” denotes a DNA sequence that encodes a polypeptide (a “secretory peptide”) that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

[0025] Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as “about” X or “approximately” X, the stated value of X will be understood to be accurate to $\pm 10\%$.

INTRODUCTION

[0026] The present invention relates to the treatment of inflammation and inflammatory diseases in which IL-20 plays a role either in initiating or spreading or maintenance by

administering to the afflicted individual an antagonist to IL-20. That antagonist can be an antibody to IL-20, a soluble receptor that binds to IL-20, an anti-sense molecule or an antibody that binds to either the IL-20RA subunit or the IL-20RB subunit of the IL-20 receptor.

[0027] IL-20 is defined and methods for producing it and antibodies to IL-20 are contained in International Patent Application No. PCT/US98/25228, publication no. WO 99/27103, published Nov. 25, 1998 and U.S. patent application Ser. No. 09/313,458 filed May 17, 1999. The polynucleotide and polypeptide of human IL-20 are represented by SEQ ID NOs: 1-4, and mouse IL-20 by SEQ ID NOs: 5-9.

[0028] The receptor to IL-20 has been discovered and is a heterodimer comprised of the polypeptide termed 'IL-20RA' (formally called Zcytor7) and a polypeptide termed 'IL-20RB'. The IL-20RA polypeptide, nucleic acid that encodes it, antibodies to IL-20RA, and methods for producing it are disclosed in U.S. Pat. No. 5,945,511 issued Aug. 31, 1999. SEQ ID NOs: 10-12 are the IL-20RA polynucleotides and polypeptides. The extracellular domain of the human IL-20RA is comprised of a polypeptide selected from the group consisting of SEQ ID NOs: 12, 55, 63 and 65, the full-length receptor subunit being comprised of SEQ ID NO: 11. The extracellular domain of mouse IL-20RA is SEQ ID NO: 38, SEQ ID NO: 37 being the entire mouse IL-20RA.

[0029] The extracellular domain of IL-20RB (SEQ ID NOs: 13-14, and a variant SEQ ID NOs: 18 and 19) is comprised of a polypeptide selected from the group consisting of SEQ ID NOs: 15, 59, 61, 67, 68 and 69. Preferably, the extracellular domain of the IL-20RA polypeptide and the extracellular domain of the IL-20RB polypeptide are covalently linked together. In a preferred embodiment one extracellular subunit polypeptide has a constant region of a heavy chain of an immunoglobulin fused to its carboxy terminus and the other extracellular subunit has a constant light chain of an immunoglobulin (Ig) fused to its carboxy terminus such that the two polypeptides come together to form a soluble receptor and a disulfide bond is formed between the heavy and the light Ig chains. In another method, a peptide linker could be fused to the two carboxy-termini of the polypeptides to form a covalently bonded soluble receptor.

[0030] SEQ ID NOs: 22 and 23 are constructs of the extracellular domain of IL-20RA fused to a mutated human immunoglobulin gamma 1 constant region produced according to the procedure set forth in example 5. SEQ ID NO: 62 is the predicted mature sequence without the signal sequence. SEQ ID NOs: 20 and 21 are constructs of the extracellular domain of IL-20RB fused to wild type human immunoglobulin kappa light chain constant region produced according to the procedure of example 5. SEQ ID NO: 60 is the predicted mature sequence without the signal sequence. FIG. 1 depicts the heterotetramer produced by example 5.

[0031] SEQ ID NOs: 52 and 53 are constructs of the extracellular domain of IL-20RA fused to a mutated human immunoglobulin gamma 1 constant region produced according to the procedure set forth in example 12. SEQ ID NO: 54 is the predicted mature sequence without the signal sequence. SEQ ID NOs: 56 and 57 are constructs of the extracellular domain of IL-20RB fused to wild type human immunoglobulin kappa light chain constant region produced according to the procedure of example 12. SEQ ID NO: 58 is the predicted mature sequence without the signal sequence. The resultant heterotetramer is almost identical to that produced by example 5, the primary difference being the absence of a polypeptide

linker between the extracellular domains and the beginning of the Ig constant regions, 22 in FIG. 1. Hereinafter, the term "extracellular domain of a receptor" means the extracellular domain of the receptor or a portion of the extracellular domain that is necessary for binding to its ligand, in this case the ligand being IL-20.

[0032] One can link together the extracellular domains of IL-20RA and IL-20RB in a number of ways such that the resultant soluble receptor can bind to IL-20. FIGS. 1-8 illustrate a representative number of embodiments of the present invention. Common elements in each of the drawings are given the same number. FIG. 1 represents the embodiment of the present invention produced according to example 5 below. The soluble receptor construct, designated 10, is comprised of two IL-20 binding site polypeptide chains designated 12 and 14. Each binding site is comprised of the extracellular domain of IL-20RA, designated 16, and the extracellular domain of IL-20RB designated 18.

[0033] The extracellular domain, 16, of IL-20RA is linked to the constant heavy one (CH1) domain, 20, of the human immunoglobulin gamma 1 heavy chain constant region via linker 22, which is SEQ ID NO:72. The CH1 domain, 20, is then linked to the CH2 domain, 24, via hinge region 23. The CH2 domain, 24, is linked to the CH3 domain, 26, via hinge region 25.

[0034] Comparing the construct of FIG. 1 with SEQ ID NO:22, the extracellular domain, 16, of IL-20RA extends from amino acid residues 36, a valine, to and including amino acid residue 249, a glutamine of SEQ ID NO:22. Polypeptide linker, 22, extends from amino acid residue 250, a glycine to and including amino acid residue 264, a serine, of SEQ ID NO:22. The CH1 domain, 22 of FIG. 1, extends from amino acid residue 265, an alanine, to and including amino acid residue 362, a valine, of SEQ ID NO:22. Hinge region 23 of FIG. 1 extends from amino acid residue 363, a glutamic acid to and including amino acid residue 377, a proline, of SEQ ID NO: 22. Chains 12 and 14 are disulfide-bonded together by means of disulfide bonds 28 and 30. The disulfide bonds are formed between the heavy chains by the cysteine residues at positions 373 and 376 of SEQ ID NO: 22 of each of the two heavy chains.

[0035] Extracellular domain, 18, of IL-20RB is linked to the constant region of the human kappa light chain (CL), 34 of FIG. 1 via polypeptide linker 32, which is the polypeptide SEQ ID NO: 72. The extracellular domain, 18, of IL-20RB extends from amino acid residue 30, a valine, to and including amino acid residue 230, an alanine, of SEQ ID NO: 20. Polypeptide linker, 32, extends from amino acid residue 231, a glycine, to and including amino acid residue 245, a serine, of SEQ ID NO:20. The kappa constant light region, 34, extends from amino acid residue 246, an arginine, to and including the final amino acid residue 352, a cysteine, of SEQ ID NO:20. The cysteine at position 352 of SEQ ID NO: 20 forms a disulfide bond, 36 in FIG. 1, with the cysteine at position 367 of SEQ ID NO: 22. The constant light chain 34 is thus linked to the hinge region, 23, by disulfide bond, 36. In this way, the extracellular domain, 16, of IL-20RA is linked to the extracellular domain, 18, of IL-20RB to form a soluble receptor.

[0036] If the cysteine residues at positions 373 and 376 of SEQ ID NO:22 were changed to different amino acid residues, the two IL-20 binding polypeptides, 12 and 14, would not be disulfide bonded together and would form a construct shown in FIG. 2. having hinge region, 27.

[0037] FIG. 3 shows a very simple soluble receptor **38** of the present invention wherein extracellular domain, **16**, of IL-20RA is connected to the extracellular domain, **18**, of IL-20RB by means of a polypeptide linker, **40**. The polypeptide linker extends from the amino terminus of extracellular domain, **16**, of IL-20RA and is connected to the carboxyl terminus of the extracellular domain, **18**, of IL-20RB. The polypeptide linker should be between 100-240 amino acids in length, preferably about 170 amino acid residues in length. A suitable linker would be comprised of glycine and serine residues. A possible linker would be multiple units of SEQ ID NO: 72, preferably about 12.

[0038] FIG. 4 shows an embodiment that has the extracellular domain, **16**, of IL-20RA linked to the extracellular domain, **18**, of IL-20RB by means of linker **40**, as in FIG. 3. While the extracellular domain, **16**, of IL-20RA is linked to the CH1 domain, **20**, as in FIG. 1 by means of polypeptide linker **42**, which should be about 30 amino acid residues in length. An ideal linker would be comprised of glycine and serine as in SEQ ID NO: 72, and the hinge sequence, **23** of FIG. 1.

[0039] FIG. 5 shows another possible embodiment of the present invention. In this embodiment, a polypeptide linker **44** of about 15 amino acid residue, e.g. SEQ ID NO: 72, links the carboxyl terminus of the extracellular domain, **18**, of IL-20RB with the amino terminus of the extracellular domain, **16**, of IL-20RA. A polypeptide linker **46** of about 30 amino acid residues extends from the carboxy terminus of the extracellular domain, **16**, of IL-20RA to the CH2 domain. The carboxyl terminus of linker **46** would preferably be comprised of the hinge region extending from amino acid residue 363, a glutamic acid to and including amino acid residue 377, a proline, of SEQ ID NO: 22. Nonetheless, polypeptide linker **46** would ideally have at least one cysteine residue at its carboxyl terminus so a disulfide bond could be formed.

[0040] The soluble IL-20 receptor of FIG. 6 is identical to that of FIG. 1 except for the CH3 domain, **26** of FIG. 1, is not present on the embodiment of FIG. 6. The CH3 region begins at amino acid residue 488, a glycine, and extends to the last residue 594 of SEQ ID NO: 22.

[0041] FIG. 7 shows a soluble IL-20 receptor construct that is identical to the construct of FIG. 1 except both the CH2, and CH3 domains are absent. The CH2 and CH3 domains run from amino acid residue 378, an alanine, to the end of the polypeptide sequence of SEQ ID NO: 22.

[0042] FIG. 8 shows a construct wherein both IL-20RA, **16**, and IL-20RB have a polypeptide linker, **48**, fused to their respective carboxyl termini. Each polypeptide linker has two cysteine residues such that when they are expressed the cysteines form two disulfide bonds, **50** and **52**. In this case the polypeptide linker is comprised of the hinge region, **23** in FIG. 1. The hinge region is comprised of amino acid residues 363, a glutamine, to and including amino acid residue 377 of SEQ ID NO: 22.

[0043] In another aspect of the invention, a method is provided for producing a soluble receptor comprised of extracellular domains of IL-20RA and IL-20RB comprising (a) introducing into a host cell a first DNA sequence comprised of a transcriptional promoter operatively linked to a first secretory signal sequence followed downstream by and in proper reading frame the DNA that encodes the extracellular portion of IL-20RA and the DNA that encodes an immunoglobulin light chain constant region; (b) introducing into the host cell a second DNA construct comprised of a transcriptional pro-

motor operatively linked to a second secretory signal followed downstream by and in proper reading frame a DNA sequence that encodes the extracellular portion of IL-20RB and a DNA sequence that encodes an immunoglobulin heavy chain constant region domain selected from the group consisting of C_H1 , C_H2 , C_H3 and C_H4 ; (c) growing the host cell in an appropriate growth medium under physiological conditions to allow the secretion of a fusion protein comprised of the extracellular domain of IL-20RA and IL-20RB; and (d) isolating the polypeptide from the host cell. In one embodiment, the second DNA sequence further encodes an immunoglobulin heavy chain hinge region wherein the hinge region is joined to the heavy chain constant region domain. In another embodiment, the second DNA sequence further encodes an immunoglobulin variable region joined upstream of and in proper reading frame with the immunoglobulin heavy chain constant region.

[0044] In an alternative embodiment, a method is provided for producing a soluble receptor comprised of the extracellular domains of IL-20RA and IL-20RB comprising (a) introducing into a host cell a first DNA sequence comprised of a transcriptional promoter operatively linked to a first secretory signal sequence followed downstream by and in proper reading frame the DNA that encodes the extracellular portion of IL-20RB and the DNA that encodes an immunoglobulin light chain constant region; (b) introducing into the host cell a second DNA construct comprised of a transcriptional promoter operatively linked to a second secretory signal followed downstream by and in proper reading frame a DNA sequence that encodes the extracellular domain of IL-20RA and a DNA sequence that encodes an immunoglobulin heavy chain constant region domain selected from the group consisting of C_H1 , C_H2 , C_H3 and C_H4 ; (c) growing the host cell in an appropriate growth medium under physiological conditions to allow the secretion of a dimerized heterodimeric fusion protein comprised of the extracellular domain of IL-20RA and IL-20RB; and (d) isolating the dimerized polypeptide from the host cell. In one embodiment, the second DNA sequence further encodes an immunoglobulin heavy chain hinge region wherein the hinge region is joined to the heavy chain constant region domain. In another embodiment, the second DNA sequence further encodes an immunoglobulin variable region joined upstream of and in proper reading frame with the immunoglobulin heavy chain constant region. (See U.S. Pat. No. 5,843,725.)

[0045] As used herein, the term "antibodies" includes polyclonal antibodies, affinity-purified polyclonal antibodies, monoclonal antibodies, and antigen-binding fragments, such as $F(ab')_2$ and Fab proteolytic fragments. Genetically engineered intact antibodies or fragments, such as chimeric antibodies, Fv fragments, single chain antibodies and the like, as well as synthetic antigen-binding peptides and polypeptides, are also included. Non-human antibodies may be humanized by grafting non-human CDRs onto human framework and constant regions, or by incorporating the entire non-human variable domains (optionally "cloaking" them with a human-like surface by replacement of exposed residues, wherein the result is a "veneered" antibody). In some instances, humanized antibodies may retain non-human residues within the human variable region framework domains to enhance proper binding characteristics. Through humanizing antibodies, biological half-life may be increased, and the potential for adverse immune reactions upon administration to humans is reduced. The binding affinity of an antibody can be readily

determined by one of ordinary skill in the art, for example, by Scatchard analysis. A variety of assays known to those skilled in the art can be utilized to detect antibodies that bind to protein or peptide. Exemplary assays are described in detail in *Antibodies: A Laboratory Manual*, Harlow and Lane (Eds.) (Cold Spring Harbor Laboratory Press, 1988). Representative examples of such assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich assay.

[0046] We have shown that IL-20 up-regulates IL-8. Inflammatory diseases in which IL-8 plays a significant role, and for which a decrease in IL-8 would be beneficial are, adult respiratory disease (ARD), septic shock, multiple organ failure, inflammatory lung injury such as asthma or bronchitis, bacterial pneumonia, psoriasis and inflammatory bowel disease such as ulcerative colitis and Crohn's disease, eczema, atopic dermatitis and contact dermatitis. Thus, antagonists to IL-20 can be used to treat these diseases.

Biology of IL-20, its Receptor and its Role in Psoriasis

[0047] Two orphan class II cytokine receptors, both of which are expressed in skin, were identified as IL-20 receptor subunits. Both IL-20 receptor subunits are required for ligand binding, distinguishing their role from that of subunits in the four other known class II cytokine receptors. IL-20RA and IL-20RB are also coexpressed in a number of human tissues besides skin, including ovary, adrenal gland, testis, salivary gland, muscle, lung, kidney, heart and to a lesser degree the small intestine suggesting additional target tissues for IL-20 action. Additionally, we have detected IL-20RA mRNA but not IL-20RB mRNA in several human tissues including stomach, thyroid, pancreas, uterus, brain and prostate suggesting that IL-20RA may partner with other class II receptor subunits. We conclude that the IL-20 heterodimeric receptor is structurally similar to other class II cytokine receptors and is expressed in skin where we have demonstrated activity of the IL-20 ligand.

[0048] Two lines of evidence indicate that a role IL-20 and its receptor are involved in psoriasis. This multigenic skin disease is characterized by increased keratinocyte proliferation, altered keratinocyte differentiation, and infiltration of immune cells into the skin. The first line of evidence for a role of IL-20 in psoriasis is that the observed hyperkeratosis and thickened epidermis in the transgenic mice that resemble human psoriatic abnormalities. Decreased numbers of tonofilaments, thought to be related to defective keratinization, are a striking feature of human psoriasis. Intramitochondrial inclusions have been found in both chemically induced and naturally occurring hyperplastic skin conditions in mice. The cause of the inclusions and their effects on mitochondrial function, if any, are unknown. We conclude that IL-20 transgenic mice exhibit many of the characteristics observed in human psoriasis.

[0049] A second line of evidence that implicates the IL-20 receptor in psoriasis is that both IL-20RA and IL-20RB mRNA are markedly upregulated in human psoriatic skin compared to normal skin. Both IL-20 receptor subunits are expressed in keratinocytes throughout the epidermis and are also expressed in a subset of immune and endothelial cells. We propose that increased expression of an activated IL-20 receptor may alter the interactions between endothelial cells,

immune cells and keratinocytes, leading to dysregulation of keratinocyte proliferation and differentiation.

[0050] A crucial step in understanding the function of a novel cytokine is the identification and characterization of its cognate receptor. We have successfully used a structure-based approach to isolate a novel interleukin that ultimately led to the isolation of its receptor. IL-20 stimulates signal transduction in the human keratinocyte HaCaT cell line, supporting a direct action of this novel ligand in skin. In addition, IL-1 β , EGF and TNF- α , proteins known to be active in keratinocytes and to be involved with proliferative and pro-inflammatory signals in skin, enhance the response to IL-20. In both HaCaT and BHK cells expressing the IL-20 receptor, IL-20 signals through STAT3.

Use of Antagonist to IL-20 to Treat Psoriasis

[0051] As indicated in the discussion above and the examples below, IL-20 is involved in the pathology of psoriasis. The present invention is in particular a method for treating psoriasis by administering antagonists to IL-20. The antagonists to IL-20 can either be a soluble receptor that binds to IL-20 or antibodies, single chain antibodies or fragments of antibodies that bind to either IL-20 or the IL-20 receptor. The antagonists will thus prevent activation of the IL-20 receptor.

[0052] Psoriasis is one of the most common dermatologic diseases, affecting up to 1 to 2 percent of the world's population. It is a chronic inflammatory skin disorder characterized by erythematous, sharply demarcated papules and rounded plaques, covered by silvery micaceous scale. The skin lesions of psoriasis are variably pruritic. Traumatized areas often develop lesions of psoriasis. Additionally, other external factors may exacerbate psoriasis including infections, stress, and medications, e.g. lithium, beta blockers, and anti-malarials.

[0053] The most common variety of psoriasis is called plaque type. Patients with plaque-type psoriasis will have stable, slowly growing plaques, which remain basically unchanged for long periods of time. The most common areas for plaque psoriasis to occur are the elbows knees, gluteal cleft, and the scalp. Involvement tends to be symmetrical. Inverse psoriasis affects the intertriginous regions including the axilla, groin, submammary region, and navel, and it also tends to affect the scalp, palms, and soles. The individual lesions are sharply demarcated plaques but may be moist due to their location. Plaque-type psoriasis generally develops slowly and runs an indolent course. It rarely spontaneously remits.

[0054] Eruptive psoriasis (guttate psoriasis) is most common in children and young adults. It develops acutely in individuals without psoriasis or in those with chronic plaque psoriasis. Patients present with many small erythematous, scaling papules, frequently after upper respiratory tract infection with beta-hemolytic streptococci. Patients with psoriasis may also develop pustular lesions. These may be localized to the palms and soles or may be generalized and associated with fever, malaise, diarrhea, and arthralgias.

[0055] About half of all patients with psoriasis have fingernail involvement, appearing as punctate pitting, nail thickening or subungual hyperkeratosis. About 5 to 10 percent of patients with psoriasis have associated joint complaints, and these are most often found in patients with fingernail involvement. Although some have the coincident occurrence of classic psoriasis, although some have the coincident occurrence of classic rheumatoid arthritis, many have joint disease that falls into

one of five type associated with psoriasis: (1) disease limited to a single or a few small joints (70 percent of cases); (2) a seronegative rheumatoid arthritis-like disease; (3) involvement of the distal interphalangeal joints; (4) severe destructive arthritis with the development of "arthritis mutilans"; and (5) disease limited to the spine.

[0056] Psoriasis can be treated by administering antagonists to IL-20. The preferred antagonists are either a soluble receptor to IL-20 or antibodies, antibody fragments or single chain antibodies that bind to either the IL-20 receptor or to IL-20. The antagonists to IL-20 can be administered alone or in combination with other established therapies such as lubricants, keratolytics, topical corticosteroids, topical vitamin D derivatives, anthralin, systemic antimetabolites such as methotrexate, psoralen-ultraviolet-light therapy (PUVA), etretinate, isotretinoin, cyclosporine, and the topical vitamin D3 derivative calcipotriol. The antagonists, in particularly the soluble receptor or the antibodies that bind to IL-20 or the IL-20 receptor can be administered to individual subcutaneously, intravenously, or transdermally using a cream or transdermal patch that contains the antagonist of IL-20. If administered subcutaneously, the antagonist can be injected into one or more psoriatic plaques. If administered transdermally, the antagonists can be administered directly on the plaques using a cream containing the antagonist to IL-20.

Use of Antagonists to IL-20 to Treat Inflammatory Conditions of the Lung.

[0057] Antagonists to IL-20 can be administered to a person who has asthma, bronchitis or cystic fibrosis or other inflammatory lung disease to treat the disease. The antagonists can be administered by any suitable method including intravenous, subcutaneous, bronchial lavage, and the use of inhalant containing an antagonist to IL-20.

Administration of Antagonists to IL-20

[0058] The quantities of antagonists to IL-20 necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medications administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in vivo administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Methods for administration include oral, intravenous, peritoneal, intramuscular, transdermal or administration into the lung or trachea in spray form by means of a nebulizer or atomizer. Pharmaceutically acceptable carriers will include water, saline, buffers to name just a few. Dosage ranges would ordinarily be expected from 1 µg to 1000 µg per kilogram of body weight per day. A dosage for an average adult of the IL-20 soluble receptor would be about 25 mg given twice weekly as a subcutaneous injection. A dosage of an antibody that binds to IL-20 would be about 25 mg given twice weekly. A mixture of antibodies that bind to the IL-20RA and IL-20RB subunits would be about 25 mg given twice weekly for an adult. Injections could be given at the site of psoriatic lesions for the treatment of psoriasis. For subcutaneous or intravenous administration of the antagonist to IL-20, the antibody or soluble receptor can be in phosphate buffered saline. Also in skin diseases such as psoriasis, the antagonist to IL-20 can be administered via an ointment or

transdermal patch. The doses may be higher or lower as can be determined by a medical doctor with ordinary skill in the art. For a complete discussion of drug formulations and dosage ranges see *Remington's Pharmaceutical Sciences*, 18th Ed., (Mack Publishing Co., Easton, Pa., 1996), and *Goodman and Gilman's: The Pharmacological Bases of Therapeutics*, 9th Ed. (Pergamon Press 1996).

[0059] The invention is further illustrated by the following non-limiting examples.

EXAMPLE 1

Up-regulation of IL-8 by IL-20

Methods:

[0060] Normal Human Epidermal neonatal keratinocytes (NHEK) (from Clonetics) at passage 2 were plated and grown to confluency in 12 well tissue culture plates. KGM (Keratinocyte growth media) was purchased from Clonetics. When cells reached confluency, they were washed with KGM media minus growth factors=KBM (keratinocyte basal media). Cells were serum starved in KBM for 72 hours prior to the addition of test compounds. Thrombin at 1 I.U./mL and trypsin at 25 nM were used as positive controls. One mL of media/well was added. KBM only was used as the negative control.

[0061] IL-20 was made up in KBM media and added at varying concentrations, from 2.5 µg/mL down to 618 ng/mL in a first experiment and from 2.5 µg/mL down to 3 ng/mL in a second experiment.

[0062] Cells were incubated at 37° C., 5% CO₂ for 48 hours. Supernatants were removed and frozen at -80° C. for several days prior to assaying for IL-8 and GM-CSF levels. Human IL-8 Immunoassay kit # D8050 (Rand D Systems, Inc.) and human GM-CSF Immunoassay kit # HSGMO (Rand D Systems, Inc.) were used to determine cytokine production following manufacturer's instructions.

Results

[0063] The results indicated that the expression of IL-8 and GM-CSF were induced by IL-20.

EXAMPLE 2

Cloning of IL-20RB

Cloning of IL-20RB Coding Region

[0064] Two PCR primers were designed based on the sequence from International Patent Application No. PCT/US99/03735 (publication no. WO 99/46379) filed on Mar. 8, 1999. SEQ ID NO: 16 contains the ATG (Met) codon with an EcoRI restriction site, SEQ ID NO: 17 contains the stop codon (TAG) with an XhoI restriction site. The PCR amplification was carried out using a human keratinocyte (HaCaT) cDNA library DNA as a template and SEQ ID NO: 16 and SEQ ID NO: 17 as primers. The PCR reaction was performed as follows: incubation at 94° C. for 1 min followed by 30 cycles of 94° C. for 30 sec and 68° C. for 2 min, after additional 68° C. for 4 min, the reaction was stored at 4° C. The PCR products were run on 1% Agarose gel, and a 1 kb DNA band was observed. The PCR products were cut from the gel and the DNA was purified using a QIAquick Gel Extraction Kit (Qiagen). The purified DNA was digested with EcoRI and XhoI, and cloned into a pZP vector that was called pZP7N. A pZP plasmid is a mammalian expression vector containing an

expression cassette having the mouse metallothionein-1 promoter, human tPA leader peptide, multiple restriction sites for insertion of coding sequences, a Glu-Glu tag, and a human growth hormone terminator. The plasmid also has an *E. coli* origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, an enhancer and an origin of replication, as well as a DHFR gene, and the SV40 terminator. Several IL-20RB-pZP7N clones were sequenced. They all contain three non-conservative mutations compared with the sequence of IL-20RB in PCT/US99/03735: (sequence IL-20RB-pZP7N), 146Pro (CCC)-Thr (ACC), 148His (CAT)-Asp (GAT), and 171Thr (ACG)-Arg (AGG).

[0065] To verify the three substitutions in IL-20RB-pZP7N clone, PCR amplification was carried out using three difference cDNA sources—fetal skin marathon cDNA, HaCaT cDNA library DNA, and prostate smooth muscle cDNA library DNA—as templates. The PCR products were gel purified and sequenced. The sequence of each of the three PCR products was consistent with that of the IL-20RB-pZP7N clone. IL-20RB is SEQ ID NO: 13 and 14, and the mature extracellular domain is SEQ ID NO: 15.

EXAMPLE 3

Binding of IL-20 to IL-20RB/IL-20RA Heterodimer

[0066] A cell-based binding assay was used to verify IL-20 binds to IL-20RA-IL-20RB heterodimer.

[0067] Expression vectors containing known and orphan Class II cytokine receptors (including IL-20RA and IL-20RB) were transiently transfected into COS cells in various combinations, which were then assayed for their ability to bind biotin-labeled IL-20 protein. The results show IL-20RB-IL-20RA heterodimer is a receptor for IL-20. The procedure used is described below.

[0068] The COS cell transfection was performed in a 12-well tissue culture plate as follows: 0.5 μ g DNA was mixed with medium containing 5 μ l lipofectamine in 92 μ l serum free Dulbecco's modified Eagle's medium (DMEM) (55 mg sodium pyruvate, 146 mg L-glutamine, 5 mg transferrin, 2.5 mg insulin, 1 μ g selenium and 5 mg fetuin in 500 ml DMEM), incubated at room temperature for 30 minutes and then added to 400 μ l serum free DMEM media. This 500 μ l mixture was then added to 1.5×10^5 COS cells/well and incubated for 5 hours at 37° C. 500 μ l 20% fetal bovine serum (FBS) DMEM media was added and incubated overnight.

[0069] The assay, a modification of the "secretion trap" (Davis, S., et al., *Cell* 87: 1161-1169 (1996), was performed as follows: cells were rinsed with PBS/1% bovine serum albumin (BSA) and blocked for 1 hour with TNB (0.1 M Tris-HCl, 0.15 M NaCl and 0.5% Blocking Reagent (NEN Renaissance TSA-Direct Kit Cat# NEL701) in water). This was followed by a one-hour incubation with 3 μ g/ml biotinylated IL-20 protein in TNB. Cells were washed with PBS/1% BSA and incubated for another hour with 1:300 diluted streptavidin-HRP (NEN kit) in TNB. Following another wash, cells were fixed for 15 minutes with 1.8% Formaldehyde in phosphate-buffered saline (PBS). Cells were then washed with TNT (0.1 M Tris-HCl, 0.15 M NaCl, and 0.05% Tween-20 in water). Positive binding signals were detected following a five-minute incubation with fluorescein tyramide reagent diluted 1:50 in dilution buffer (NEN kit). Cells were washed with TNT, preserved with Vectashield Mounting

Media (Vector Labs) diluted 1:5 in TNT, and visualized using an FITC filter on an inverted fluorescent microscope.

EXAMPLE 4

Up-Regulation of Inflammatory Cytokines by IL-20

Cell Treatment

[0070] The human keratinocyte cell line, HaCaT was grown at 37° C. to several days post-confluence in T-75 tissue culture flasks. At this point, normal growth media (DMEM+10% FBS) was removed and replaced with serum-free media. Cells were then incubated for two days at 37° C. DMEM was then removed and four flasks of cells per treatment were treated with one of each of the following conditions for four hours at 37° C.: recombinant human (rh) IL-1 alpha at 5 ng/mL, rh IL-1 alpha at 20 ng/mL, rh IL-1 alpha at 5 ng/mL+IL-20 at 1 μ g/mL, IL-20 at 1 μ g/mL, or rh IL-10 at 10 ng/mL.

RNA Isolation

[0071] Following cytokine treatment, media was removed and cells were lysed using a guanidium thiocyanate solution. Total RNA was isolated from the cell lysate by an overnight spin on a cesium chloride gradient. The following day, the RNA pellet was resuspended in a TE/SDS solution and ethanol precipitated. RNA was then quantitated using a spectrophotometer, followed by a DNase treatment as per Section V.B. of Clontech's Atlas™ cDNA Expression Arrays User Manual (version PT3140-1/PR9X390, published Nov. 5, 1999). Quality of RNA samples was verified by purity calculations based on spec readings, and by visualization on agarose gel. Genomic contamination of the RNA samples was ruled out by PCR analysis of the beta-actin gene.

Probe Synthesis

[0072] Clontech's protocols for polyA+ enrichment, probe synthesis and hybridization to Atlas™ arrays were followed (see above, plus Atlas™ Pure Total RNA Labeling System User Manual, PT3231-1/PR96157, published Jun. 22, 1999). Briefly, polyA+ RNA was isolated from 50 mg of total RNA using streptavidin coated magnetic beads (by Clontech, Paolo Alto, Calif.) and a magnetic particle separator. PolyA+ RNA was then labeled with α^{32} P-dATP via RT-PCR. Clontech CDS primers specific to the 268 genes on the Atlas™ human cytokine/receptor array (Cat. #7744-1) were used in the reaction. Labeled probe was isolated using column chromatography and counted in scintillation fluid.

Array Membrane Hybridization

[0073] Atlas™ arrays were pre-hybridized with Clontech ExpressHyb plus 100 mg/mL heat denatured salmon sperm DNA for at least thirty minutes at 68° C. with continuous agitation. Membranes were then hybridized with 1.9×10^6 CPM/mL (a total of 1.14×10^7 CPM) overnight at 68° C. with continuous agitation. The following day, membranes were washed for thirty minutes $\times$ 4 in 2 $\times$ SSC, 1% SDS at 68° C., plus for thirty minutes $\times$ 1 in 0.1 $\times$ SSC, 0.5% SDS at 68° C., followed by one final room temperature wash for five minutes in 2 $\times$ SSC. Array membranes were then placed in Kodak plastic pouches sealed and exposed to a phosphor imager screen overnight at room temperature. The next day, phos-

phor screens were scanned on a phosphor imager and analyzed using Clontech's AtlasImage™ 1.0 software.

Results

Genes Up-Regulated by IL-20

- [0074] 1. Tumor necrosis factor (TNF) was up-regulated 1.9-2.4 fold by IL-20.
- [0075] 2. Placental growth factors 1 & 2 (PLGF) were up-regulated 1.9-2.0 fold by IL-20.
- [0076] 3. Coagulating factor II receptor was up-regulated 2.0-2.5 fold by IL-20.
- [0077] 4. Calcitonin receptor was up-regulated 2.2-2.3 fold by IL-20.
- [0078] 5. TNF-inducible hyaluronate-binding protein TSG-6 was up-regulated 2.1-2.2 fold by IL-20.
- [0079] 6. Vascular endothelial growth factor (VEGF) receptor-1 precursor, tyrosine-protein kinase receptor (FLT-1) (SFLT) was up-regulated 2.1-2.7 fold by IL-20.
- [0080] 7. MRP-8 (calcium binding protein in macrophages MIF-related) was up-regulated 2.9-4.1 fold by IL-20.
- [0081] 8. MRP-14 (calcium binding protein in macrophages MIF-related) was up-regulated 3.0-3.8 fold by IL-20.
- [0082] 9. Relaxin H2 was up-regulated 3.14 fold by IL-20.
- [0083] 10. Transforming growth factor beta (TGFβ) receptor III 300 kDa was up-regulated 2.4-3.6 fold by IL-20.

Genes Showing Synergy with IL-20+IL-1 Treatment

- [0084] 1. Bone morphogenic protein 2a was up-regulated 1.8 fold with IL-20 treatment alone, 2.5 fold with IL-1 treatment alone, and 8.2 fold with both IL-20 and IL-1 treatment together.
- [0085] 2. MRP-8 was up-regulated 2.9 fold with IL-20 treatment alone, 10.7 fold with IL-1 treatment alone and 18.0 fold with both IL-20 and IL-1 treatment together.
- [0086] 3. Erythroid differentiation protein (EDF) was up-regulated 1.9 fold with IL-20 treatment alone, 9.7 fold with IL-1 treatment alone and 19.0 fold with both IL-20 and IL-1 treatment together.
- [0087] 4. MRP-14 (calcium binding protein in macrophages, MIF related) was up-regulated 3.0 fold with IL-20 treatment alone, 12.2 fold with IL-1 treatment alone and 20.3 fold with both IL-20 and IL-1 treatment together.
- [0088] 5. Heparin-binding EGF-like growth factor was up-regulated 2.0 fold with IL-20 treatment alone, 14 fold with IL-1 treatment alone and 25.0 fold with both IL-20 and IL-1 treatment together.
- [0089] 6. Beta-thromboglobulin-like protein was up-regulated 1.5 fold with IL-20 treatment alone, 15 fold with IL-1 treatment alone and 27 fold with both IL-20 and IL-1 treatment together.
- [0090] 7. Brain-derived neurotrophic factor (BDNF) was up-regulated 1.7 fold with IL-20 treatment alone, 25 fold with IL-1 treatment alone and 48 fold with both IL-20 and IL-1 treatment together.
- [0091] 8. Monocyte chemotactic and activating factor MCAF was up-regulated 1.3 fold with IL-20 treatment

alone, 32 fold with IL-1 treatment alone and 56 fold with both IL-20 and IL-1 treatment together.

EXAMPLE 5

IL-20RA/RB Receptor-Ig Fusion Heterotetramer

[0092] The expression vector pEZE3 was used to express the recombinant IL-20 receptor-Ig fusion protein. The plasmid pEZE3 is derived from pDC312. pDC312 was obtained through license from Immunex Corporation. The plasmids pDC312 and pEZE3 contain an EASE segment as described in WO 97/25420. The presence of the EASE segment in an expression vector can improve expression of recombinant proteins two to eight fold in stable cell pools.

[0093] The plasmid pEZE3 is a tricistronic expression vector that may be used to express up to three different proteins in mammalian cells, preferably Chinese Hamster Ovary (CHO) cells. The pEZE3 expression unit contains the cytomegalovirus (CMV) enhancer/promoter, the adenovirus tripartite leader sequence, a multiple cloning site for insertion of the coding region for the first recombinant protein, the poliovirus type 2 internal ribosome entry site, a second multiple cloning site for insertion of the coding region for the second recombinant protein, an encephalomyocarditis virus internal ribosome entry site, a coding segment for mouse dihydrofolate reductase, and the SV40 transcription terminator. In addition, pEZE3 contains an *E. coli* origin of replication and the bacterial beta lactamase gene.

[0094] The IL-20 receptor-Ig fusion protein is a disulfide linked heterotetramer consisting of two chains of the extracellular domain of the human IL-20RB fused to the wild type human immunoglobulin kappa light chain constant region and two chains of the human IL-20RA protein extracellular domain fused to a mutated human immunoglobulin gamma 1 constant region. The human immunoglobulin gamma 1 constant region contains amino acid substitutions to reduce FcγRI binding and C1q complement fixation.

[0095] The human IL-20RB extracellular domain human immunoglobulin kappa light chain constant region fusion construct was generated by overlap PCR. The IL-20RB coding segment consists of amino acids 1 to 230. The template used for the PCR amplification of the IL-20RB segment was generated IL-20RB human kappa light chain constant region expression construct as described below in Example 12. Oligonucleotide primers SEQ ID NO: 24 and SEQ ID NO: 25 were used to amplify the IL-20RB segment. The entire wild type human immunoglobulin kappa light chain constant region was used. The template used for the PCR amplification of the wild type human immunoglobulin kappa light chain constant region segment was generated IL-20RB human kappa light chain constant region expression construct as described in Example 12. Oligonucleotide primers SEQ ID NO: 26 and SEQ ID NO: 27 were used to amplify the wild type human immunoglobulin kappa light chain constant region. The two protein coding domains were linked by overlap PCR using oligonucleotides SEQ ID NO: 24 and SEQ ID NO: 27. A (Gly₄Ser)₃ (SEQ ID NO: 72) peptide linker was inserted between the two protein domains. The (Gly₄Ser)₃ peptide linker was encoded on the PCR primers SEQ ID NO: 26 and SEQ ID NO: 25. The resultant IL-20RB extracellular domain/kappa light chain constant region fusion construct is shown by SEQ ID NOs: 20 and 21. The predicted mature polypeptide, minus the signal sequence, is SEQ ID NO: 60. The portion of the extracellular domain of IL-20RB that was

actually used was comprised of the amino acid sequence of SEQ ID NO: 61. N-terminal sequencing resulted in the predicted amino acid sequence.

[0096] The human IL-20RA extracellular domain human immunoglobulin gamma 1 heavy chain constant region fusion construct was generated by overlap PCR of four separate DNA fragments, each generated by separate PCR amplification reactions. The first fragment contained an optimized tPA (tissue plasminogen activator) signal sequence. The tPA signal sequence was amplified using oligonucleotide primers SEQ ID NO: 28 and SEQ ID NO: 29 using an in-house previously generated expression vector as the template. The second fragment contained the IL-20RA extracellular domain-coding region consisting of amino acids 30 to 243 of SEQ ID NO: 11. Oligonucleotide primers SEQ ID NO: 30 and SEQ ID NO: 31 were used to amplify this IL-20RA segment using a previously generated clone of IL-20RA as the template.

[0097] The human gamma 1 heavy chain constant region was generated from 2 segments. The first segment containing the CH1 domain was amplified using oligonucleotide primers SEQ ID NO: 32 and SEQ ID NO: 33 using a clone of the wild type human gamma 1 heavy chain constant region as the template. The second segment containing the remaining hinge, C_H2, and C_H3 domains of the human immunoglobulin gamma 1 heavy chain constant region was generated by PCR amplification using oligonucleotide primers SEQ ID NO: 34 and SEQ ID NO: 35. The template used for this PCR amplification was from a previously generated human gamma 1 Fc construct that contained codons for amino acid substitutions to reduce FcγRI binding and C1q complement fixation as described in Example 12.

[0098] The four protein coding domains were linked by overlap PCR using oligonucleotides SEQ ID NO: 28 and SEQ ID NO: 35. A (Gly₄Ser)₃ peptide linker was inserted between the IL-20RA and CH1 protein domains. The (Gly₄Ser)₃ peptide linker was encoded on the PCR primers SEQ ID NO: 32 and SEQ ID NO: 31. The IL-20RA extracellular domain/domain human immunoglobulin gamma 1 heavy constant region fusion protein and DNA sequence are shown in SEQ ID NOS: 22 and 23. The predicted mature polypeptide sequence, minus the signal sequence, is SEQ ID NO: 62. The portion of extracellular domain of IL-20RA that was actually used was comprised of SEQ ID NO: 63.

[0099] The IL-20RB extracellular domain human immunoglobulin kappa light chain constant region fusion coding segment was cloned into the second MCS while the human IL-20RA extracellular domain human immunoglobulin gamma 1 heavy chain constant region fusion coding segment was cloned into the first MCS of pEZE3. The plasmid was used to transfect CHO cells. The cells were selected in medium without hypoxanthine or thymidine and the transgene was amplified using methotrexate. The presence of protein was assayed by Western blotting using anti human gamma 1 heavy chain constant region and anti human kappa light chain antibodies. N-terminal sequencing revealed that the optimized tPA leader was not completely cleaved. The observed mass indicated that the first residue of the polypeptide sequence to be pyroglutamic acid, and the N-terminal

sequence appears to be pyroEEIHAELRRFRRVPCVSGG (SEQ ID NO: 64), the underlined portion being remnants of the tPA leader.

EXAMPLE 6

IL-20 Transgenic Phenotype

[0100] Both human and mouse IL-20 were overexpressed in transgenic mice using a variety of promoters. The liver-specific mouse albumin promoter, directing expression of human IL-20, was used initially in an attempt to achieve circulating levels of protein. Subsequent studies were conducted using the keratin 14 (K14) promoter, which primarily targets expression to the epidermis and other stratified squamous epithelia; the mouse metallothionein-1 promoter, which gives a broad expression pattern; and the EμLCK promoter, which drives expression in cells of the lymphoid lineage. Similar results were obtained in all four cases, possibly because these promoters all give rise to circulating levels of IL-20.

[0101] In all cases, transgenic pups expressing the IL-20 transgene were smaller than non-transgenic littermates, had a shiny appearance with tight, wrinkled skin and died within the first few days after birth. Pups had milk in their stomachs indicating that they were able to suckle. These mice had swollen extremities, tail, nostril and mouth regions and had difficulty moving. In addition, the mice were frail, lacked visible adipose tissue and had delayed ear and toe development. Low expression levels in liver (less than 100 mRNA molecules/cell) were sufficient for both the neonatal lethality and skin abnormalities. Transgenic mice without a visible phenotype either did not express the transgene, did not express it at detectable levels, or were mosaic.

[0102] Histologic analysis of the skin of the IL-20 transgenic mice showed a thickened epidermis, hyperkeratosis and a compact stratum corneum compared to non-transgenic littermates. Sero cellular crusts (scabs) were observed occasionally. Electron microscopic (EM) analysis of skin from transgenic mice showed intramitochondrial lipid inclusions, mottled keratohyaline granules, and relatively few tonofilaments similar to that observed in human psoriatic skin and in mouse skin disease models. In addition, many of the transgenic mice had apoptotic thymic lymphocytes. No other abnormalities were detected by histopathological analysis. These histological and EM results support and extend the observed gross skin alterations.

EXAMPLE 7

Specificity and Affinity of IL-20 for its Receptor

[0103] The specificity and affinity of IL-20 for its receptor was determined using BHK cells stably transfected with IL-20RA, IL-20RB or both receptor subunits. Binding assays using radiolabeled ligand demonstrated that IL-20 bound to BHK transfectants expressing both IL-20RA and IL-20RB but not to untransfected cells nor to transfectants expressing either receptor subunit alone. Binding of ¹²⁵I-labeled IL-20 was eliminated in the presence of 100-fold excess of unlabeled IL-20 but not with 100-fold excess of the unrelated

cytokine, IL-21. The binding affinity (K_D) of IL-20 to the IL-20RA/IL-20RB heterodimeric receptor was determined to be approximately 1.5 nM.

EXAMPLE 8

IL-20 receptor Activation

[0104] To determine if IL-20 binding leads to receptor activation, the factor-dependent pre-B cell line BaF3 was co-transfected with IL-20RA and IL-20RB and treated with IL-20 at various concentrations. IL-20 stimulated proliferation in a dose-dependent manner and gave a detectable signal at 1.1 pM, with a half maximal response at 3.4 pM. We note that the IL-20 concentration for the half maximal proliferative response in BaF3 cells is 1000× lower than that for half maximal binding affinity in BHK cells. Possible explanations for this large difference include the use of different cell lines, different receptor expression levels and different assay outputs. IL-20 also stimulated signal transduction in the biologically relevant human keratinocyte cell line HaCaT, which naturally expresses IL-20RA and IL-20RB. Therefore, IL-20 binds and activates the heterodimeric IL-20RA/IL-20RB receptor at concentrations expected for a cytokine. While the negative controls containing untransfected BaF3

EXAMPLE 9

Expression Analysis of IL-20RA and IL-20RB

[0105] RT-PCR analysis was performed on a variety of human tissues to determine the expression pattern of IL-20RA and IL-20RB. Both receptor subunits are most highly expressed in skin and testis. The significant result is that IL-20RA and IL-20RB are both expressed in skin, where they have been shown to mediate the IL-20-induced response. Both IL-20RA and IL-20RB are also both expressed in monocytes, lung, ovary, muscle, testis, adrenal gland, heart, salivary gland and placenta. IL-20RA is also in brain, kidney, liver, colon, small intestine, stomach, thyroid, pancreas, uterus and prostate while IL-20RB is not.

EXAMPLE 10

IL-20RA and IL-20RB mRNA are Up-Regulated in Psoriasis

[0106] In situ hybridization was used to determine whether IL-20 receptor expression is altered in psoriasis. Skin samples from four psoriasis patients and three unaffected patients were assayed with probes specific for the two-receptor subunit mRNAs. All four psoriatic skin samples had high levels of IL-20RA and IL-20RB mRNA in keratinocytes whereas normal skin samples did not have detectable levels of either receptor subunit mRNA. Positive signals in psoriatic skin were also observed in mononuclear immune cells and in endothelial cells in a subset of vessels. Therefore, both IL-20RA and IL-20RB are expressed in keratinocytes, immune cells and endothelial cells, the major cell types thought to interact in psoriasis.

EXAMPLE 11

Cloning of Mouse IL-20RA

[0107] A cross-species hybridization probe was generated which contained the full-length cDNA fragment encoding human IL-20RA. A Southern blot of mouse genomic DNA and Northern blots of mouse RNA were performed to dem-

onstrate that the human IL-20RA cDNA could specifically hybridize to mouse sequences. The Northern blot results indicated that mouse IL-20RA RNA was present in mouse embryo day 15 and 17 as well as heart, brain, lung, liver, kidney, testes, spleen, thymus, liver, stomach, and small intestine.

[0108] The human IL-20RA full length DNA hybridization probe was used to screen a mouse genomic library. The library, which was obtained from Clontech (Palo Alto, Calif.), was generated from an MboI partial digest of mouse genomic DNA and cloned into the BamHI site of Lambda bacteriophage EMBL3 SP6/T7. Positive bacteriophage was plaque purified and bacteriophage DNA was prepared using Promega's Wizard Lambda Preps DNA Purification System. Two genomic restriction enzyme fragments, a 5.7 kb EcoRI fragment and an 8.0 kb SacI fragment, were generated from the positive bacteriophage and subcloned into pBluescript. DNA sequence analysis revealed the presence of 3 exons from the mouse ortholog to human IL-20RA.

[0109] PCR primers from the 5' UTR, SEQ ID NO: 40, and 3' UTR, SEQ ID NO: 41, were designed to generate a full-length mouse IL-20RA sequence by PCR amplification. Mouse embryo 15 day plus 17 day cDNA was used as the template for the PCR amplification. PCR products were subcloned and sequenced for confirmation. The mouse sequences are SEQ ID NOs: 36 and 37. The mature extracellular domain is comprised of SEQ ID NO: 38.

EXAMPLE 12

Construction of an IL-20 Receptor Heterotetramer

[0110] A vector expressing a secreted hIL-20RA/hIL-20B heterodimer was constructed. In this construct, the extracellular domain of hIL-20RA was fused to the heavy chain of IgG gamma 1 (IgGγ1), while the extracellular portion of IL-20RB was fused to human kappa light chain (human κ light chain).

Construction of IgG Gamma 1 and Human κ Light Fusion Vectors

[0111] The heavy chain of IgGγ1 was cloned into the Zem229R mammalian expression vector (ATCC deposit No. 69447) such that any extracellular portion of a receptor having a 5' EcoRI and 3' NheI site can be cloned in, resulting in an N-terminal extracellular domain-C-terminal IgGγ1 fusion. The IgGγ1 fragment used in this construct was made by using PCR to isolate the IgGγ1 sequence from a Clontech human fetal liver cDNA library as template. A PCR reaction using oligos SEQ ID NO: 42 and SEQ ID NO: 43 was run as follows: 40 cycles of 94° for 60 sec., 53° C. for 60 sec., and 72° for 120 sec.; and 72° C. for 7 minutes. PCR products were separated by agarose gel electrophoresis and purified using a QiaQuick™ (Qiagen Inc., Valencia, Calif.) gel extraction kit. The isolated, 990 bp, DNA fragment was digested with MluI and EcoRI (Boehringer-Mannheim), extracted with QiaQuick™ gel extraction kit and ligated with oligos SEQ ID NO: 44 and SEQ ID NO: 45, which comprise an MluI/EcoRI linker, into Zem229R previously digested with MluI and EcoRI using standard molecular biology techniques disclosed herein. This generic cloning vector was called Vector#76 hIgGgamma1 w/ Ch1 #786 Zem229R (Vector #76). The polynucleotide sequence of the extracellular domain of hIL-20RA fused to the heavy chain of IgG gamma 1 is shown in SEQ ID NO: 52 and the corresponding polypeptide

sequence shown in SEQ ID NO: 53, the mature polypeptide, minus the signal sequence being comprised of SEQ ID NO: 54. The portion of the extracellular domain of IL-20RA used was comprised of SEQ ID NO: 55.

[0112] The human κ light chain was cloned in the Zem228R mammalian expression vector (ATCC deposit No. 69446) such that any extracellular portion of a receptor having a 5' EcoRI site and a 3' KpnI site can be cloned in, resulting in an N-terminal extracellular domain-C-terminal human κ light chain fusion. The human κ light chain fragment used in this construct was made by using PCR to isolate the human κ light chain sequence from the same Clontech hFetal Liver cDNA library used above. A PCR reaction was run using oligos SEQ ID NO: 46 and SEQ ID NO: 47. PCR products were separated by agarose gel electrophoresis and purified using a QiaQuick™ (Qiagen) gel extraction kit. The isolated, 315 bp, DNA fragment was digested with MluI and EcoRI (Boehringer-Mannheim), extracted with QiaQuick™ gel extraction kit and ligated with the MluI/EcoRI linker described above, into Zem228R previously digested with MluI and EcoRI using standard molecular biology techniques disclosed herein. This generic cloning vector was called Vector #77 hκlight #774 Zem228R (Vector #77). The polynucleotide sequence of the extracellular portion of IL-20RB fused to human kappa light chain is shown in SEQ ID NO: 56 and the corresponding polypeptide sequence shown in SEQ ID NO: 57, the mature polypeptide, minus the signal sequence, is comprised of SEQ ID NO: 58. The portion of the extracellular domain of IL-20RB actually used was comprised of SEQ ID NO: 59.

Insertion of hIL-20RA and IL-20RB Extracellular Domains into Fusion Vector Constructs

[0113] Using the construction vectors above, a construct having human IL-20RA fused to IgGγ1 was made. This construction was done by using PCR to obtain human IL-20RA receptor from hIL-20RA/IgG Vector #102 with oligos SEQ ID NO: 48 and SEQ ID NO: 49 under conditions described as follows: 30 cycles of 94° C. for 60 sec., 57° C. for 60 sec., and 72° C. for 120 sec.; and 72° C. for 7 min. The resulting PCR product was digested with EcoRI and NheI, gel purified, as described herein, and ligated into a previously EcoRI and NheI digested and band-purified Vector #76 (above). The resulting vector was sequenced to confirm that the human IL-20RA/IgG gamma 1 fusion (hIL-20RA/Ch1 IgG) was correct. The hIL-20RA/Ch1 IgG gamma 1 #1825 Zem229R vector was called vector #195. The IL-20RA/Ch1 IgGγ1 sequence thus obtained is depicted by SEQ ID NOs: 52 and 53. N-terminal sequencing indicated the presence of the predicted mature polypeptide sequence of SEQ ID NO: 54.

[0114] A separate construct having IL-20RB fused to κ light was also constructed. The IL-20RB/human κ light chain construction was performed as above by PCRing from DR1/7N-4 with oligos SEQ ID NO: 50 and SEQ ID NO: 51, digesting the resulting band with EcoRI and KpnI and then ligating this product into a previously EcoRI and KpnI digested and band-purified Vec#77 (above). The resulting vector was sequenced to confirm that the IL-20RB/human κ light chain fusion (IL-20RB/κlight) was correct. This IL-20RB/κlight construct is shown by SEQ ID NOs: 56 and 57. N-terminal sequencing of the resultant polypeptide indicated the presence of the predicted mature amino acid

sequence comprised of SEQ ID NO: 58. SEQ ID NO:59 is the mature portion of the extracellular domain of IL-20RB used.

Co-Expression of the Human IL-20RA and Human IL-20RB Receptors

[0115] Approximately 16 μg of each of vectors #194 and #195, above, were co-transfected into BHK-570 cells (ATCC No. CRL-10314) using Lipofectamine™ reagent (Gibco/BRL), as per manufacturer's instructions. The transfected cells were selected for 10 days in DMEM+5% FBS (Gibco/BRL) containing 1 μM of methotrexate (MTX) (Sigma, St. Louis, Mo.) and 0.5 mg/ml G418 (Gibco/BRL) for 10 days. The resulting pool of transfectants was selected again in 10 μM MTX and 0.5 mg/ml G418 for 10 days.

[0116] The resulting pool of doubly selected cells was used to generate protein. Three factories (Nunc, Denmark) of this pool were used to generate 8 L of serum free conditioned medium. This conditioned media was passed over a 1 ml protein-A column and eluted in (10) 750 microliter fractions. 4 of these fractions found to have the highest concentration were pooled and dialyzed (10 kD MW cutoff) against PBS. Finally, the dialyzed material was analyzed by BCA (Pierce) and found to have a concentration of 317 μg/ml. A total of 951 μg was obtained from this 8 L purification.

EXAMPLE 13

IL-20 Binding Activates STAT3 in the HaCaT Keratinocyte Cell Line

[0117] IL-20 binds cell lines transfected with both subunits of its receptor. However, these cell lines overexpress the IL-20 receptor relative to its normal level and their relevance to the physiological role of IL-20 is unclear. The human HaCaT keratinocyte cell line, which expresses endogenous IL-20RA and IL-20RB was used to examine IL-20 signal transduction in a biologically relevant cell type. HaCaT cells were infected with recombinant adenovirus containing a reporter construct to allow detection of intracellular signaling. The construct consists of the firefly luciferase gene driven by promoter/enhancer sequences comprised of the serum response element (SRE) and signal transducers and activators of transduction elements (STATs). This assay system detects productive ligand-receptor interactions and indicates possible downstream signal transduction components involved in receptor activation. Treatment with IL-20 alone resulted in a dose-dependent increase in luciferase activity with a half maximal response occurring at approximately 2.3 nM. Subsequent luciferase reporter assays using adenovirus vectors containing only the SRE element or only the STAT elements produced detectable reporter activation only through STATs.

[0118] To determine if other cytokines act in concert with IL-20, HaCaT cells were treated with IL-20 alone or in combination with a single submaximal dose of EGF, IL-1β, or TNFα. In the presence of each of these three proteins, IL-20 treatment resulted in a dose-dependent increase in luciferase activity. IL-20 in combination with IL-1β results in a half-maximal response at approximately 0.5 nM, about five-fold lower than with IL-20 alone. In addition, activation of the reporter gene is detectable at 0.1 nM IL-20, a dose that is at least tenfold lower than the IL-20 dose required alone.

[0119] BHK cells transfected with IL-20RA, IL-20RB or both receptor subunits were used to determine whether receptor pairing was required for IL-20 stimulation of STAT-luciferase. As was the case with binding assays, only cells

transfected with both receptor subunits responded to IL-20 and did so with a half-maximal response of 5.7 pM. We note that the IL-20 concentration for the half-maximal response in BHK cells is 400-fold lower than that for half-maximal response in HaCaT cells. It is likely that a lower concentration of IL-20 is needed for half-maximal response in BHK cells, as compared to HaCaT cells, due to higher receptor levels in the BHK IL-20 receptor transfectants.

[0120] A nuclear translocation assay was used to identify STAT proteins involved in IL-20 action. Both HaCaT cells, with endogenous IL-20 receptors, and BHK cells transfected with IL-20RA and IL-20RB, were treated with IL-20 protein and translocation of STAT3 and STAT1 transcription factors from the cytoplasm to the nucleus was assayed by immunofluorescence.

[0121] In unstimulated HaCaT cells, STAT3 staining was predominantly in the cytosol. Treatment of HaCaT cells with

IL-20 resulted in a distinct accumulation of STAT3 in the nucleus. Nuclear translocation of STAT3 in response to increasing concentrations of IL-20 occurred with a half-maximal IL-20 concentration of 7 nM. In contrast to STAT3 translocation, HaCaT cells treated with IL-20 did not show any detectable nuclear accumulation of STAT1.

[0122] BHK cells transfected with IL-20RA and IL-20RB were used to confirm that the IL-20 receptor was required for IL-20 stimulation of STAT3 nuclear translocation. In BHK cells lacking the IL-20 receptor, STAT3 remained cytosolic following treatment with IL-20. In contrast, in BHK cells transfected with the IL-20 receptor, STAT3 translocated to the nucleus in response to IL-20. Again, STAT1 remained cytosolic regardless of IL-20 treatment or IL-20 receptor expression. We conclude that the IL-20 receptor is required for IL-20-mediated STAT3 activation.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 72

<210> SEQ ID NO 1

<211> LENGTH: 176

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Lys Ala Ser Ser Leu Ala Phe Ser Leu Leu Ser Ala Ala Phe Tyr
 1             5             10             15

Leu Leu Trp Thr Pro Ser Thr Gly Leu Lys Thr Leu Asn Leu Gly Ser
 20             25             30

Cys Val Ile Ala Thr Asn Leu Gln Glu Ile Arg Asn Gly Phe Ser Asp
 35             40             45

Ile Arg Gly Ser Val Gln Ala Lys Asp Gly Asn Ile Asp Ile Arg Ile
 50             55             60

Leu Arg Arg Thr Glu Ser Leu Gln Asp Thr Lys Pro Ala Asn Arg Cys
 65             70             75             80

Cys Leu Leu Arg His Leu Leu Arg Leu Tyr Leu Asp Arg Val Phe Lys
 85             90             95

Asn Tyr Gln Thr Pro Asp His Tyr Thr Leu Arg Lys Ile Ser Ser Leu
100             105             110

Ala Asn Ser Phe Leu Thr Ile Lys Lys Asp Leu Arg Leu Cys His Ala
115             120             125

His Met Thr Cys His Cys Gly Glu Glu Ala Met Lys Lys Tyr Ser Gln
130             135             140

Ile Leu Ser His Phe Glu Lys Leu Glu Pro Gln Ala Ala Val Val Lys
145             150             155             160

Ala Leu Gly Glu Leu Asp Ile Leu Leu Gln Trp Met Glu Glu Thr Glu
165             170             175

```

<210> SEQ ID NO 2

<211> LENGTH: 152

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Leu Lys Thr Leu Asn Leu Gly Ser Cys Val Ile Ala Thr Asn Leu Gln

```

-continued

1	5	10	15
Glu Ile Arg	Asn Gly Phe Ser Asp	Ile Arg Gly Ser Val	Gln Ala Lys
	20	25	30
Asp Gly Asn	Ile Asp Ile Arg	Ile Leu Arg Arg Thr	Glu Ser Leu Gln
	35	40	45
Asp Thr Lys	Pro Ala Asn Arg Cys Cys	Leu Leu Arg His	Leu Leu Arg
	50	55	60
Leu Tyr Leu	Asp Arg Val Phe Lys Asn Tyr	Gln Thr Pro Asp	His Tyr
	65	70	75
Thr Leu Arg	Lys Ile Ser Ser Leu Ala	Asn Ser Phe Leu Thr	Ile Lys
	85	90	95
Lys Asp Leu	Arg Leu Cys His Ala His	Met Thr Cys His	Cys Gly Glu
	100	105	110
Glu Ala Met	Lys Lys Tyr Ser Gln Ile	Leu Ser His Phe	Glu Lys Leu
	115	120	125
Glu Pro Gln	Ala Ala Val Val Lys Ala	Leu Gly Glu Leu	Asp Ile Leu
	130	135	140
Leu Gln Trp	Met Glu Glu Thr Glu		
	145	150	

<210> SEQ ID NO 3
 <211> LENGTH: 151
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Met Lys Ala	Ser Ser Leu Ala Phe	Ser Leu Leu Ser Ala	Ala Phe Tyr
1	5	10	15
Leu Leu Trp	Thr Pro Ser Thr Gly	Leu Lys Thr Leu Asn	Leu Gly Ser
	20	25	30
Cys Val Ile	Ala Thr Asn Leu Gln	Glu Ile Arg Asn	Gly Phe Ser Asp
	35	40	45
Ile Arg Gly	Ser Val Gln Ala Lys	Asp Gly Asn Ile	Asp Ile Arg Ile
	50	55	60
Leu Arg Arg	Thr Glu Ser Leu Gln	Asp Thr Lys Pro	Ala Asn Arg Cys
	65	70	75
Cys Leu Leu	Arg His Leu Leu Arg	Leu Tyr Leu Asp	Arg Val Phe Lys
	85	90	95
Asn Tyr Gln	Thr Pro Asp His Tyr	Thr Leu Arg Lys	Ile Ser Ser Leu
	100	105	110
Ala Asn Ser	Phe Leu Thr Ile Lys	Lys Asp Leu Arg	Leu Cys Leu Glu
	115	120	125
Pro Gln Ala	Ala Val Val Lys Ala	Leu Gly Glu Leu	Asp Ile Leu Leu
	130	135	140
Gln Trp Met	Glu Glu Thr Glu		
	145	150	

<210> SEQ ID NO 4
 <211> LENGTH: 127
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Leu Lys Thr Leu Asn Leu Gly Ser Cys Val Ile Ala Thr Asn Leu Gln

-continued

1	5	10	15
Glu Ile Arg	Asn Gly Phe Ser Asp	Ile Arg Gly Ser Val	Gln Ala Lys
	20	25	30
Asp Gly Asn	Ile Asp Ile Arg	Ile Leu Arg Arg Thr	Glu Ser Leu Gln
	35	40	45
Asp Thr Lys	Pro Ala Asn Arg Cys Cys	Leu Leu Arg His	Leu Leu Arg
	50	55	60
Leu Tyr Leu	Asp Arg Val Phe Lys	Asn Tyr Gln Thr Pro	Asp His Tyr
	65	70	75
Thr Leu Arg	Lys Ile Ser Ser Leu	Ala Asn Ser Phe Leu	Thr Ile Lys
	85	90	95
Lys Asp Leu	Arg Leu Cys Leu Glu	Pro Gln Ala Ala Val	Val Lys Ala
	100	105	110
Leu Gly Glu	Leu Asp Ile Leu Leu	Gln Trp Met Glu Glu	Thr Glu
	115	120	125

<210> SEQ ID NO 5

<211> LENGTH: 176

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

Met Lys Gly	Phe Gly Leu Ala	Phe Gly Leu Phe	Ser Ala Val	Gly Phe
1	5	10	15	
Leu Leu Trp	Thr Pro Leu Thr	Gly Leu Lys Thr	Leu His Leu	Gly Ser
	20	25	30	
Cys Val Ile	Thr Ala Asn Leu	Gln Ala Ile Gln	Lys Glu Phe	Ser Glu
	35	40	45	
Ile Arg Asp	Ser Val Gln Ala	Glu Asp Thr Asn	Ile Asp Ile	Arg Ile
	50	55	60	
Leu Arg Thr	Thr Glu Ser Leu	Lys Asp Ile Lys	Ser Leu Asp	Arg Cys
	65	70	75	80
Cys Phe Leu	Arg His Leu Val	Arg Phe Tyr Leu	Asp Arg Val	Phe Lys
	85	90	95	
Val Tyr Gln	Thr Pro Asp His	His Thr Leu Arg	Lys Ile Ser	Ser Leu
	100	105	110	
Ala Asn Ser	Phe Leu Ile Ile	Lys Lys Asp Leu	Ser Val Cys	His Ser
	115	120	125	
His Met Ala	Cys His Cys Gly	Glu Glu Ala Met	Glu Lys Tyr	Asn Gln
	130	135	140	
Ile Leu Ser	His Phe Ile Glu	Leu Glu Leu Gln	Ala Ala Val	Val Lys
	145	150	155	160
Ala Leu Gly	Glu Leu Gly Ile	Leu Leu Arg Trp	Met Glu Glu	Met Leu
	165	170	175	

<210> SEQ ID NO 6

<211> LENGTH: 152

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Leu Lys Thr	Leu His Leu Gly	Ser Cys Val Ile	Thr Ala Asn	Leu Gln
1	5	10	15	
Ala Ile Gln	Lys Glu Phe Ser	Glu Ile Arg Asp	Ser Val Gln	Ala Glu

-continued

20					25					30					
Asp	Thr	Asn	Ile	Asp	Ile	Arg	Ile	Leu	Arg	Thr	Thr	Glu	Ser	Leu	Lys
		35					40					45			
Asp	Ile	Lys	Ser	Leu	Asp	Arg	Cys	Cys	Phe	Leu	Arg	His	Leu	Val	Arg
		50					55					60			
Phe	Tyr	Leu	Asp	Arg	Val	Phe	Lys	Val	Tyr	Gln	Thr	Pro	Asp	His	His
		65					70					75			80
Thr	Leu	Arg	Lys	Ile	Ser	Ser	Leu	Ala	Asn	Ser	Phe	Leu	Ile	Ile	Lys
			85						90					95	
Lys	Asp	Leu	Ser	Val	Cys	His	Ser	His	Met	Ala	Cys	His	Cys	Gly	Glu
			100					105					110		
Glu	Ala	Met	Glu	Lys	Tyr	Asn	Gln	Ile	Leu	Ser	His	Phe	Ile	Glu	Leu
		115					120					125			
Glu	Leu	Gln	Ala	Ala	Val	Val	Lys	Ala	Leu	Gly	Glu	Leu	Gly	Ile	Leu
		130					135					140			
Leu	Arg	Trp	Met	Glu	Glu	Met	Leu								
		145					150								

<210> SEQ ID NO 7
 <211> LENGTH: 144
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 7

Cys	Val	Ile	Thr	Ala	Asn	Leu	Gln	Ala	Ile	Gln	Lys	Glu	Phe	Ser	Glu
	1			5					10					15	
Ile	Arg	Asp	Ser	Val	Gln	Ala	Glu	Asp	Thr	Asn	Ile	Asp	Ile	Arg	Ile
		20					25					30			
Leu	Arg	Thr	Thr	Glu	Ser	Leu	Lys	Asp	Ile	Lys	Ser	Leu	Asp	Arg	Cys
		35					40					45			
Cys	Phe	Leu	Arg	His	Leu	Val	Arg	Phe	Tyr	Leu	Asp	Arg	Val	Phe	Lys
		50					55					60			
Val	Tyr	Gln	Thr	Pro	Asp	His	His	Thr	Leu	Arg	Lys	Ile	Ser	Ser	Leu
		65					70					75			80
Ala	Asn	Ser	Phe	Leu	Ile	Ile	Lys	Lys	Asp	Leu	Ser	Val	Cys	His	Ser
			85					90					95		
His	Met	Ala	Cys	His	Cys	Gly	Glu	Glu	Ala	Met	Glu	Lys	Tyr	Asn	Gln
		100					105					110			
Ile	Leu	Ser	His	Phe	Ile	Glu	Leu	Glu	Leu	Gln	Ala	Ala	Val	Val	Lys
		115					120					125			
Ala	Leu	Gly	Glu	Leu	Gly	Ile	Leu	Leu	Arg	Trp	Met	Glu	Glu	Met	Leu
		130					135					140			

<210> SEQ ID NO 8
 <211> LENGTH: 154
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 8

Met	Lys	Gly	Phe	Gly	Leu	Ala	Phe	Gly	Leu	Phe	Ser	Ala	Val	Gly	Phe
	1			5					10					15	
Leu	Leu	Trp	Thr	Pro	Leu	Thr	Gly	Leu	Lys	Thr	Leu	His	Leu	Gly	Ser
		20					25					30			
Cys	Val	Ile	Thr	Ala	Asn	Leu	Gln	Ala	Ile	Gln	Lys	Glu	Phe	Ser	Glu

-continued

35	40	45
Ile Arg Asp Ser Val Ser Leu Asp Arg Cys Cys Phe Leu Arg His Leu		
50	55	60
Val Arg Phe Tyr Leu Asp Arg Val Phe Lys Val Tyr Gln Thr Pro Asp		
65	70	75 80
His His Thr Leu Arg Lys Ile Ser Ser Leu Ala Asn Ser Phe Leu Ile		
	85	90 95
Ile Lys Lys Asp Leu Ser Val Cys His Ser His Met Ala Cys His Cys		
	100	105 110
Gly Glu Glu Ala Met Glu Lys Tyr Asn Gln Ile Leu Ser His Phe Ile		
	115	120 125
Glu Leu Glu Leu Gln Ala Ala Val Val Lys Ala Leu Gly Glu Leu Gly		
	130	135 140
Ile Leu Leu Arg Trp Met Glu Glu Met Leu		
145	150	

<210> SEQ ID NO 9
 <211> LENGTH: 130
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Leu Lys Thr Leu His Leu Gly Ser Cys Val Ile Thr Ala Asn Leu Gln	
1	5 10 15
Ala Ile Gln Lys Glu Phe Ser Glu Ile Arg Asp Ser Val Ser Leu Asp	
	20 25 30
Arg Cys Cys Phe Leu Arg His Leu Val Arg Phe Tyr Leu Asp Arg Val	
	35 40 45
Phe Lys Val Tyr Gln Thr Pro Asp His His Thr Leu Arg Lys Ile Ser	
	50 55 60
Ser Leu Ala Asn Ser Phe Leu Ile Ile Lys Lys Asp Leu Ser Val Cys	
65	70 75 80
His Ser His Met Ala Cys His Cys Gly Glu Glu Ala Met Glu Lys Tyr	
	85 90 95
Asn Gln Ile Leu Ser His Phe Ile Glu Leu Glu Leu Gln Ala Ala Val	
	100 105 110
Val Lys Ala Leu Gly Glu Leu Gly Ile Leu Leu Arg Trp Met Glu Glu	
	115 120 125
Met Leu	
130	

<210> SEQ ID NO 10
 <211> LENGTH: 3516
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (237) ... (1895)

<400> SEQUENCE: 10

tccagctggg tagccggggg agcgcgcgtg ggggctccgc gagtcgctcg cccttggttt	60
ctggggaagc ctgggggacg cggctgtggc ggaggcgccc tgggactcag gtcgcctgga	120
gcgtggcaag cagagcccca ggcgcggagc tgaggccgcg cgccgcgcgt tggccccagc	180
gggcgtggga ctgagcagtc tgctgcccc cgacatgtga ccagccccg ccgccc atg	239

															Met 1	
egg gct ccc ggc cgc ccg gcc ctg cgg ccg ctg ccg ctg ccg ccg ctg	287															
Arg Ala Pro Gly Arg Pro Ala Leu Arg Pro Leu Pro Leu Pro Pro Leu																
5 10 15																
ctg ctg ttg ctc ctg gcg gcg cct tgg gga cgg gca gtt ccc tgt gtc	335															
Leu Leu Leu Leu Leu Ala Ala Pro Trp Gly Arg Ala Val Pro Cys Val																
20 25 30																
tct ggt ggt ttg cct aaa cct gca aac atc acc ttc tta tcc atc aac	383															
Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn																
35 40 45																
atg aag aat gtc cta caa tgg act cca cca gag ggt ctt caa gga gtt	431															
Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly Val																
50 55 60 65																
aaa gtt act tac act gtg cag tat ttc ata tat ggg caa aag aaa tgg	479															
Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp																
70 75 80																
ctg aat aaa tca gaa tgc aga aat atc aat aga acc tac tgt gat ctt	527															
Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu																
85 90 95																
tct gct gaa act tct gac tac gaa cac cag tat tat gcc aaa gtt aag	575															
Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys																
100 105 110																
gcc att tgg gga aca aag tgt tcc aaa tgg gct gaa agt gga cgg ttc	623															
Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg Phe																
115 120 125																
tat cct ttt tta gaa aca caa att ggc cca cca gag gtg gca ctg act	671															
Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu Val Ala Leu Thr																
130 135 140 145																
aca gat gag aag tcc att tct gtt gtc ctg aca gct cca gag aag tgg	719															
Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala Pro Glu Lys Trp																
150 155 160																
aag aga aat cca gaa gac ctt cct gtt tcc atg caa caa ata tac tcc	767															
Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln Gln Ile Tyr Ser																
165 170 175																
aat ctg aag tat aac gtg tct vtg ttg aat act aaa tca aac aga acg	815															
Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys Ser Asn Arg Thr																
180 185 190																
tgg tcc cag tgt gtg acc aac cac acg ctg gtg ctc acc tgg ctg gag	863															
Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu Thr Trp Leu Glu																
195 200 205																
ccg aac act ctt tac tgc gta cac gtg gag tcc ttc gtc cca ggg ccc	911															
Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe Val Pro Gly Pro																
210 215 220 225																
cct cgc cgt gct cag cct tct gag aag cag tgt gcc agg act ttg aaa	959															
Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala Arg Thr Leu Lys																
230 235 240																
gat caa tca tca gag ttc aag gct aaa atc atc ttc tgg tat gtt ttg	1007															
Asp Gln Ser Ser Glu Phe Lys Ala Lys Ile Ile Phe Trp Tyr Val Leu																
245 250 255																
ccc ata tct att acc gtg ttt ctt ttt tct gtg atg ggc tat tcc atc	1055															
Pro Ile Ser Ile Thr Val Phe Leu Phe Ser Val Met Gly Tyr Ser Ile																
260 265 270																
tac cga tat atc cac gtt ggc aaa gag aaa cac cca gca aat ttg att	1103															
Tyr Arg Tyr Ile His Val Gly Lys Glu Lys His Pro Ala Asn Leu Ile																
275 280 285																
ttg att tat gga aat gaa ttt gac aaa aga ttc ttt gtg cct gct gaa	1151															

[illegible]

-continued

```

gcacacttgt acaattat tctgggtact tcccatatgc acatagcact gtaaaaaata 2285
tttcccaaag atcactcatt ttataaat acactttttca gaattggggt tattgcgagc 2345
aggaggagat acttaaaaca tgcacatata ccagggttgt ggtaagttgg tcacatgtga 2405
aaacctcaac tatttaata tcatgattca tttttgagt gaatacatca ggcacagacc 2465
ttcatgatat cacacactct tggctacttt aagaggccat ctttaatact ttatgagtag 2525
ttctggagtg taaacataaa cgagtattct tttgtagtca gaaaagtgct ctctcaataa 2585
tttagtaggg gcttattgtc tctcaaaact aacctaaaag aaaatgacac attttataat 2645
agaatattac atttatttct ggaagtgtgt tttcaaaaag atatttacat agtctgtaaa 2705
ctagaaagtg ttaggtaaag ctctagggtta ctgtgttact attataatat taaacattcg 2765
aataggcagt cgttcaaaga ctctttggaa tatctatgaa tgaatatact ctattcttat 2825
aatattaaaa cccataagta aatataggac atacaagaga atgaggttaa atgactatgt 2885
aaggagagtg ttattaaaat ttgatgaaat ttactgtagg aactaaacta tgccataaaa 2945
caatagcttt ctagttcatt tccagtaact gttcccatct cctttaccac ttgttaagaa 3005
aattaaattc ttcagtcacg ctgctttaa atgggacaaa atctattaag ttgaaccata 3065
tataattgtg gatatttggc tgtttttaat ctgacaagca gtaacttcat atggtttgcc 3125
ttaatatata tttgttttag tcatgaactc ataataccatt gatgctcttt catgagaaga 3185
gatatgacct atatttcctt attgatatta ttggtacagg cagacaaccc tggtaggaga 3245
gatggattct ggggtcatga cctttcgtga ttatccgcaa atgcaaacag ttcagatct 3305
aatggtttaa tttaggagtg aattatatta atcagagtgt tctgttattc tcaatcttta 3365
tagaaacgat tctgctggtt ttgaagaaca gatgtattac actaaactga aaagtagttc 3425
aagagtgaga aagaataaat tgttattaag agcaaaagaa aaataaagtg attgatgata 3485
aaaaaaaaa aaaaaaagcg gccgcctcga g 3516

```

<210> SEQ ID NO 11

<211> LENGTH: 553

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

```

Met Arg Ala Pro Gly Arg Pro Ala Leu Arg Pro Leu Pro Leu Pro Pro
 1           5           10          15
Leu Leu Leu Leu Leu Leu Ala Ala Pro Trp Gly Arg Ala Val Pro Cys
 20          25          30
Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile
 35          40          45
Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly
 50          55          60
Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys
 65          70          75          80
Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp
 85          90          95
Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val
100         105         110
Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg
115         120         125

```

-continued

Phe	Tyr	Pro	Phe	Leu	Glu	Thr	Gln	Ile	Gly	Pro	Pro	Glu	Val	Ala	Leu
130						135				140					
Thr	Thr	Asp	Glu	Lys	Ser	Ile	Ser	Val	Val	Leu	Thr	Ala	Pro	Glu	Lys
145				150						155					160
Trp	Lys	Arg	Asn	Pro	Glu	Asp	Leu	Pro	Val	Ser	Met	Gln	Gln	Ile	Tyr
			165						170					175	
Ser	Asn	Leu	Lys	Tyr	Asn	Val	Ser	Val	Leu	Asn	Thr	Lys	Ser	Asn	Arg
		180						185					190		
Thr	Trp	Ser	Gln	Cys	Val	Thr	Asn	His	Thr	Leu	Val	Leu	Thr	Trp	Leu
		195					200					205			
Glu	Pro	Asn	Thr	Leu	Tyr	Cys	Val	His	Val	Glu	Ser	Phe	Val	Pro	Gly
210						215					220				
Pro	Pro	Arg	Arg	Ala	Gln	Pro	Ser	Glu	Lys	Gln	Cys	Ala	Arg	Thr	Leu
225					230					235					240
Lys	Asp	Gln	Ser	Ser	Glu	Phe	Lys	Ala	Lys	Ile	Ile	Phe	Trp	Tyr	Val
			245						250					255	
Leu	Pro	Ile	Ser	Ile	Thr	Val	Phe	Leu	Phe	Ser	Val	Met	Gly	Tyr	Ser
		260					265						270		
Ile	Tyr	Arg	Tyr	Ile	His	Val	Gly	Lys	Glu	Lys	His	Pro	Ala	Asn	Leu
	275						280					285			
Ile	Leu	Ile	Tyr	Gly	Asn	Glu	Phe	Asp	Lys	Arg	Phe	Phe	Val	Pro	Ala
290					295						300				
Glu	Lys	Ile	Val	Ile	Asn	Phe	Ile	Thr	Leu	Asn	Ile	Ser	Asp	Asp	Ser
305					310					315					320
Lys	Ile	Ser	His	Gln	Asp	Met	Ser	Leu	Leu	Gly	Lys	Ser	Ser	Asp	Val
			325						330					335	
Ser	Ser	Leu	Asn	Asp	Pro	Gln	Pro	Ser	Gly	Asn	Leu	Arg	Pro	Pro	Gln
		340						345					350		
Glu	Glu	Glu	Glu	Val	Lys	His	Leu	Gly	Tyr	Ala	Ser	His	Leu	Met	Glu
	355						360					365			
Ile	Phe	Cys	Asp	Ser	Glu	Glu	Asn	Thr	Glu	Gly	Thr	Ser	Phe	Thr	Gln
370						375					380				
Gln	Glu	Ser	Leu	Ser	Arg	Thr	Ile	Pro	Pro	Asp	Lys	Thr	Val	Ile	Glu
385					390					395					400
Tyr	Glu	Tyr	Asp	Val	Arg	Thr	Thr	Asp	Ile	Cys	Ala	Gly	Pro	Glu	Glu
			405						410					415	
Gln	Glu	Leu	Ser	Leu	Gln	Glu	Glu	Val	Ser	Thr	Gln	Gly	Thr	Leu	Leu
		420						425					430		
Glu	Ser	Gln	Ala	Ala	Leu	Ala	Val	Leu	Gly	Pro	Gln	Thr	Leu	Gln	Tyr
	435						440					445			
Ser	Tyr	Thr	Pro	Gln	Leu	Gln	Asp	Leu	Asp	Pro	Leu	Ala	Gln	Glu	His
	450					455					460				
Thr	Asp	Ser	Glu	Glu	Gly	Pro	Glu	Glu	Glu	Pro	Ser	Thr	Thr	Leu	Val
465					470					475					480
Asp	Trp	Asp	Pro	Gln	Thr	Gly	Arg	Leu	Cys	Ile	Pro	Ser	Leu	Ser	Ser
			485						490					495	
Phe	Asp	Gln	Asp	Ser	Glu	Gly	Cys	Glu	Pro	Ser	Glu	Gly	Asp	Gly	Leu
		500						505					510		
Gly	Glu	Glu	Gly	Leu	Leu	Ser	Arg	Leu	Tyr	Glu	Glu	Pro	Ala	Pro	Asp
	515						520					525			

-continued

Arg Pro Pro Gly Glu Asn Glu Thr Tyr Leu Met Gln Phe Met Glu Glu
530 535 540

Trp Gly Leu Tyr Val Gln Met Glu Asn
545 550

<210> SEQ ID NO 12

<211> LENGTH: 221

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe
1 5 10 15

Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly
20 25 30

Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly
35 40 45

Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr
50 55 60

Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr
65 70 75 80

Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu
85 90 95

Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu
100 105 110

Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala
115 120 125

Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln
130 135 140

Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys
145 150 155 160

Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu
165 170 175

Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe
180 185 190

Val Pro Gly Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala
195 200 205

Arg Thr Leu Lys Asp Gln Ser Ser Glu Phe Lys Ala Lys
210 215 220

<210> SEQ ID NO 13

<211> LENGTH: 971

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (18)...(950)

<400> SEQUENCE: 13

gaattcgagt ctaccaa atg cag act ttc aca atg gtt cta gaa gaa atc 50
Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile
1 5 10

tgg aca agt ctt ttc atg tgg ttt ttc tac gca ttg att cca tgt ttg 98
Trp Thr Ser Leu Phe Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu
15 20 25

ctc aca gat gaa gtg gcc att ctg cct gcc cct cag aac ctc tct gta 146

-continued

Leu	Thr	Asp	Glu	Val	Ala	Ile	Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val		
		30					35					40					
ctc	tca	acc	aac	atg	aag	cat	ctc	ttg	atg	tgg	agc	cca	gtg	atc	gcg	194	
Leu	Ser	Thr	Asn	Met	Lys	His	Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala		
		45				50					55						
cct	gga	gaa	aca	gtg	tac	tat	tct	gtc	gaa	tac	cag	ggg	gag	tac	gag	242	
Pro	Gly	Glu	Thr	Val	Tyr	Tyr	Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu		
		60			65					70					75		
agc	ctg	tac	acg	agc	cac	atc	tgg	atc	ccc	agc	agc	tgg	tgc	tca	ctc	290	
Ser	Leu	Tyr	Thr	Ser	His	Ile	Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu		
				80					85					90			
act	gaa	ggt	cct	gag	tgt	gat	gtc	act	gat	gac	atc	acg	gcc	act	gtg	338	
Thr	Glu	Gly	Pro	Glu	Cys	Asp	Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val		
			95					100						105			
cca	tac	aac	ctt	cgt	gtc	agg	gcc	aca	ttg	ggc	tca	cag	acc	tca	gcc	386	
Pro	Tyr	Asn	Leu	Arg	Val	Arg	Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala		
		110					115					120					
tgg	agc	atc	ctg	aag	cat	ccc	ttt	aat	aga	aac	tca	acc	atc	ctt	acc	434	
Trp	Ser	Ile	Leu	Lys	His	Pro	Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr		
		125				130					135						
cga	cct	ggg	atg	gag	atc	acc	aaa	gat	ggc	ttc	cac	ctg	gtt	att	gag	482	
Arg	Pro	Gly	Met	Glu	Ile	Thr	Lys	Asp	Gly	Phe	His	Leu	Val	Ile	Glu		
		140			145				150					155			
ctg	gag	gac	ctg	ggg	ccc	cag	ttt	gag	ttc	ctt	gtg	gcc	tac	tgg	agg	530	
Leu	Glu	Asp	Leu	Gly	Pro	Gln	Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Arg		
			160					165						170			
agg	gag	cct	ggt	gcc	gag	gaa	cat	gtc	aaa	atg	gtg	agg	agt	ggg	ggt	578	
Arg	Glu	Pro	Gly	Ala	Glu	Glu	His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly		
		175						180						185			
att	cca	gtg	cac	cta	gaa	acc	atg	gag	cca	ggg	gct	gca	tac	tgt	gtg	626	
Ile	Pro	Val	His	Leu	Glu	Thr	Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val		
		190					195					200					
aag	gcc	cag	aca	ttc	gtg	aag	gcc	att	ggg	agg	tac	agc	gcc	ttc	agc	674	
Lys	Ala	Gln	Thr	Phe	Val	Lys	Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser		
		205					210				215						
cag	aca	gaa	tgt	gtg	gag	gtg	caa	gga	gag	gcc	att	ccc	ctg	gta	ctg	722	
Gln	Thr	Glu	Cys	Val	Glu	Val	Gln	Gly	Glu	Ala	Ile	Pro	Leu	Val	Leu		
		220				225				230				235			
gcc	ctg	ttt	gcc	ttt	gtt	ggc	ttc	atg	ctg	atc	ctt	gtg	gtc	gtg	cca	770	
Ala	Leu	Phe	Ala	Phe	Val	Gly	Phe	Met	Leu	Ile	Leu	Val	Val	Val	Pro		
			240					245						250			
ctg	ttc	gtc	tgg	aaa	atg	ggc	cgg	ctg	ctc	cag	tac	tcc	tgt	tgc	ccc	818	
Leu	Phe	Val	Trp	Lys	Met	Gly	Arg	Leu	Leu	Gln	Tyr	Ser	Cys	Cys	Pro		
			255					260						265			
gtg	gtg	gtc	ctc	cca	gac	acc	ttg	aaa	ata	acc	aat	tca	ccc	cag	aag	866	
Val	Val	Val	Leu	Pro	Asp	Thr	Leu	Lys	Ile	Thr	Asn	Ser	Pro	Gln	Lys		
		270					275					280					
tta	atc	agc	tgc	aga	agg	gag	gag	gtg	gat	gcc	tgt	gcc	acg	gct	gtg	914	
Leu	Ile	Ser	Cys	Arg	Arg	Glu	Glu	Val	Asp	Ala	Cys	Ala	Thr	Ala	Val		
		285				290					295						
atg	tct	cct	gag	gaa	ctc	ctc	agg	gcc	tgg	atc	tca	taggtttgcg				960	
Met	Ser	Pro	Glu	Glu	Leu	Leu	Arg	Ala	Trp	Ile	Ser						
		300				305				310							
gaaggctcga	g															971	

<210> SEQ ID NO 14

<211> LENGTH: 311

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser Leu Phe
 1             5             10             15

Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr Asp Glu Val
          20             25             30

Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met
 35             40             45

Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val
 50             55             60

Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser
 65             70             75             80

His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu
          85             90             95

Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg
          100            105            110

Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys
          115            120            125

His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu
          130            135            140

Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly
          145            150            155            160

Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu Pro Gly Ala
          165            170            175

Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu
          180            185            190

Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe
          195            200            205

Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val
          210            215            220

Glu Val Gln Gly Glu Ala Ile Pro Leu Val Leu Ala Leu Phe Ala Phe
          225            230            235            240

Val Gly Phe Met Leu Ile Leu Val Val Val Pro Leu Phe Val Trp Lys
          245            250            255

Met Gly Arg Leu Leu Gln Tyr Ser Cys Cys Pro Val Val Val Leu Pro
          260            265            270

Asp Thr Leu Lys Ile Thr Asn Ser Pro Gln Lys Leu Ile Ser Cys Arg
          275            280            285

Arg Glu Glu Val Asp Ala Cys Ala Thr Ala Val Met Ser Pro Glu Glu
          290            295            300

Leu Leu Arg Ala Trp Ile Ser
          305            310

<210> SEQ ID NO 15
<211> LENGTH: 203
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser
 1             5             10             15

```


-continued

Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly
 20 25 30
 Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu
 35 40 45
 Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu
 50 55 60
 Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr
 65 70 75 80
 Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser
 85 90 95
 Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro
 100 105 110
 Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu
 115 120 125
 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu
 130 135 140
 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro
 145 150 155 160
 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala
 165 170 175
 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr
 180 185 190
 Glu Cys Val Glu Val Gln Gly Glu Ala Ile Pro
 195 200

<210> SEQ ID NO 16
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

gcgaattcga gtctacacaa tgcagacttt cac 33

<210> SEQ ID NO 17
 <211> LENGTH: 32
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

cgctcgagcc ttccgcaaac ctatgagatc ca 32

<210> SEQ ID NO 18
 <211> LENGTH: 1382
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (132)...(1034)

<400> SEQUENCE: 18

tcgacccacg cgctccgcgt gcgactcaga cctcagctcc aacatatgca ttctgaagaa 60
 agatggctga gatggacaga atgctttatt ttggaaagaa acaatgttct aggtcaaaact 120
 gagtctacca a atg cag act ttc aca atg gtt cta gaa gaa atc tgg aca 170
 Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr
 1 5 10
 agt ctt ttc atg tgg ttt ttc tac gca ttg att cca tgt ttg ctc aca 218

Ser	Leu	Phe	Met	Trp	Phe	Phe	Tyr	Ala	Leu	Ile	Pro	Cys	Leu	Leu	Thr	
	15					20					25					
gat	gaa	gtg	gcc	att	ctg	cct	gcc	cct	cag	aac	ctc	tct	gta	ctc	tca	266
Asp	Glu	Val	Ala	Ile	Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val	Leu	Ser	
	30				35					40					45	
acc	aac	atg	aag	cat	ctc	ttg	atg	tgg	agc	cca	gtg	atc	gcg	cct	gga	314
Thr	Asn	Met	Lys	His	Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala	Pro	Gly	
				50					55					60		
gaa	aca	gtg	tac	tat	tct	gtc	gaa	tac	cag	ggg	gag	tac	gag	agc	ctg	362
Glu	Thr	Val	Tyr	Tyr	Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu	
			65					70					75			
tac	acg	agc	cac	atc	tgg	atc	ccc	agc	agc	tgg	tgc	tca	ctc	act	gaa	410
Tyr	Thr	Ser	His	Ile	Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu	
			80				85					90				
ggt	cct	gag	tgt	gat	gtc	act	gat	gac	atc	acg	gcc	act	gtg	cca	tac	458
Gly	Pro	Glu	Cys	Asp	Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr	
	95					100					105					
aac	ctt	cgt	gtc	agg	gcc	aca	ttg	ggc	tca	cag	acc	tca	gcc	tgg	agc	506
Asn	Leu	Arg	Val	Arg	Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala	Trp	Ser	
	110				115					120				125		
atc	ctg	aag	cat	ccc	ttt	aat	aga	aac	tca	acc	atc	ctt	acc	cga	cct	554
Ile	Leu	Lys	His	Pro	Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr	Arg	Pro	
				130				135						140		
ggg	atg	gag	atc	ccc	aaa	cat	ggc	ttc	cac	ctg	gtt	att	gag	ctg	gag	602
Gly	Met	Glu	Ile	Pro	Lys	His	Gly	Phe	His	Leu	Val	Ile	Glu	Leu	Glu	
			145					150					155			
gac	ctg	ggg	ccc	cag	ttt	gag	ttc	ctt	gtg	gcc	tac	tgg	acg	agg	gag	650
Asp	Leu	Gly	Pro	Gln	Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Thr	Arg	Glu	
			160				165					170				
cct	ggt	gcc	gag	gaa	cat	gtc	aaa	atg	gtg	agg	agt	ggg	ggt	att	cca	698
Pro	Gly	Ala	Glu	Glu	His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly	Ile	Pro	
			175			180					185					
gtg	cac	cta	gaa	acc	atg	gag	cca	ggg	gct	gca	tac	tgt	gtg	aag	gcc	746
Val	His	Leu	Glu	Thr	Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val	Lys	Ala	
			190			195				200				205		
cag	aca	ttc	gtg	aag	gcc	att	ggg	agg	tac	agc	gcc	ttc	agc	cag	aca	794
Gln	Thr	Phe	Val	Lys	Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser	Gln	Thr	
				210				215						220		
gaa	tgt	gtg	gag	gtg	caa	gga	gag	gcc	att	ccc	ctg	gta	ctg	gcc	ctg	842
Glu	Cys	Val	Glu	Val	Gln	Gly	Glu	Ala	Ile	Pro	Leu	Val	Leu	Ala	Leu	
			225					230					235			
ttt	gcc	ttt	gtt	ggc	ttc	atg	ctg	atc	ctt	gtg	gtc	gtg	cca	ctg	ttc	890
Phe	Ala	Phe	Val	Gly	Phe	Met	Leu	Ile	Leu	Val	Val	Val	Pro	Leu	Phe	
			240				245					250				
gtc	tgg	aaa	atg	ggc	cgg	ctg	ctc	cag	tac	tcc	tgt	tgc	ccc	gtg	gtg	938
Val	Trp	Lys	Met	Gly	Arg	Leu	Leu	Gln	Tyr	Ser	Cys	Cys	Pro	Val	Val	
			255			260					265					
gtc	ctc	cca	gac	acc	ttg	aaa	ata	acc	aat	tca	ccc	cag	gtt	aat	cag	986
Val	Leu	Pro	Asp	Thr	Leu	Lys	Ile	Thr	Asn	Ser	Pro	Gln	Val	Asn	Gln	
					275					280				285		
ctg	cag	aag	gga	gga	ggg	gga	tgc	ctg	tgc	cac	ggc	tgt	gat	gtc	tcc	1034
Leu	Gln	Lys	Gly	Gly	Gly	Gly	Cys	Leu	Cys	His	Gly	Cys	Asp	Val	Ser	
				290					295					300		
tgagggaactc ctcaggggcct ggatctcata tcagggtttgc ggaagggccc aggtgaagcc																1094
gagaacctgg tctgcatgac atggaaacca tgagggggaca agttgtgttt ctgttttccg																1154
ccacggacaa ggggatgagag aagtaggaag agcctgttgt ctacaagtct agaagcaacc																1212

-continued

```

atcagaggca ggggtgggttg tctaacagaa caactgactg aggctatggg ggttgtgacc 1274
tctagacttt gggcttcac ttgcttggt gagcaaccct gggaaaagtg acttcatccc 1334
ttcgggtccca agttttctca tctgtaatgg gggatcccta caaaactg 1382

```

```

<210> SEQ ID NO 19
<211> LENGTH: 301
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 19

```

```

Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser Leu Phe
  1             5             10             15
Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr Asp Glu Val
             20             25             30
Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met
             35             40             45
Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val
             50             55             60
Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser
             65             70             75             80
His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu
             85             90             95
Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg
             100            105            110
Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys
             115            120            125
His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu
             130            135            140
Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly
             145            150            155            160
Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu Pro Gly Ala
             165            170            175
Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu
             180            185            190
Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe
             195            200            205
Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val
             210            215            220
Glu Val Gln Gly Glu Ala Ile Pro Leu Val Leu Ala Leu Phe Ala Phe
             225            230            235            240
Val Gly Phe Met Leu Ile Leu Val Val Val Pro Leu Phe Val Trp Lys
             245            250            255
Met Gly Arg Leu Leu Gln Tyr Ser Cys Cys Pro Val Val Val Leu Pro
             260            265            270
Asp Thr Leu Lys Ile Thr Asn Ser Pro Gln Val Asn Gln Leu Gln Lys
             275            280            285
Gly Gly Gly Gly Cys Leu Cys His Gly Cys Asp Val Ser
             290            295            300

```

```

<210> SEQ ID NO 20
<211> LENGTH: 1081
<212> TYPE: DNA

```

-continued

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (9)...(1067)

<400> SEQUENCE: 20

```

ggccggcc atg cag act ttc aca atg gtt cta gaa gaa atc tgg aca agt      50
      Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser
        1              5              10

ctt ttc atg tgg ttt ttc tac gca ttg att cca tgt ttg ctc aca gat      98
Leu Phe Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr Asp
  15              20              25              30

gaa gtg gcc att ctg cct gcc cct cag aac ctc tct gta ctc tca acc     146
Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr
      35              40              45

aac atg aag cat ctc ttg atg tgg agc cca gtg atc gcg cct gga gaa     194
Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu
      50              55              60

aca gtg tac tat tct gtc gaa tac cag ggg gag tac gag agc ctg tac     242
Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr
      65              70              75

acg agc cac atc tgg atc ccc agc agc tgg tgc tca ctc act gaa ggt     290
Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly
      80              85              90

cct gag tgt gat gtc act gat gac atc acg gcc act gtg cca tac aac     338
Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn
      95              100              105              110

ctt cgt gtc agg gcc aca ttg ggc tca cag acc tca gcc tgg agc atc     386
Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile
      115              120              125

ctg aag cat ccc ttt aat aga aac tca acc atc ctt acc cga cct ggg     434
Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly
      130              135              140

atg gag atc ccc aaa cat ggc ttc cac ctg gtt att gag ctg gag gac     482
Met Glu Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu Asp
      145              150              155

ctg ggg ccc cag ttt gag ttc ctt gtg gcc tac tgg acg agg gag cct     530
Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu Pro
      160              165              170

ggg gcc gag gaa cat gtc aaa atg gtg agg agt ggg ggt att cca gtg     578
Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val
      175              180              185              190

cac cta gaa acc atg gag cca ggg gct gca tac tgt gtg aag gcc cag     626
His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln
      195              200              205

aca ttc gtg aag gcc att ggg agg tac agc gcc ttc agc cag aca gaa     674
Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu
      210              215              220

tgt gtg gag gtg caa gga gag gcc gga ggt ggt ggc agt gga ggc ggc     722
Cys Val Glu Val Gln Gly Glu Ala Gly Gly Gly Ser Gly Gly Gly
      225              230              235

ggt agc gga ggc ggt ggc agt cga act gtg gct gca cca tct gtc ttc     770
Gly Ser Gly Gly Gly Ser Arg Thr Val Ala Ala Pro Ser Val Phe
      240              245              250

atc ttc ccg cca tct gat gag cag ttg aaa tct gga act gcc tct gtt     818
Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val
      255              260              265              270

gtg tgc ctg ctg aat aac ttc tat ccc aga gag gcc aaa gta cag tgg     866

```

-continued

Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp		
				275					280					285			
aag	gtg	gat	aac	gcc	ctc	caa	tcg	ggt	aac	tcc	cag	gag	agt	gtc	aca	914	
Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr		
			290					295					300				
gag	cag	gac	agc	aag	gac	agc	acc	tac	agc	ctc	agc	agc	acc	ctg	acg	962	
Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr		
			305				310					315					
ctg	agc	aaa	gca	gac	tac	gag	aaa	cac	aaa	gtc	tac	gcc	tgc	gaa	gtc	1010	
Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val		
			320				325				330						
acc	cat	cag	ggc	ctg	agc	tcg	ccc	gtc	aca	aag	agc	ttc	aac	agg	gga	1058	
Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly		
				340						345				350			
gag	tgt	taa	tctagagcg	cgcc												1081	
Glu	Cys	*															

<210> SEQ ID NO 21

<211> LENGTH: 352

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Met	Gln	Thr	Phe	Thr	Met	Val	Leu	Glu	Glu	Ile	Trp	Thr	Ser	Leu	Phe		
1				5					10					15			
Met	Trp	Phe	Phe	Tyr	Ala	Leu	Ile	Pro	Cys	Leu	Leu	Thr	Asp	Glu	Val		
			20					25					30				
Ala	Ile	Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val	Leu	Ser	Thr	Asn	Met		
		35				40						45					
Lys	His	Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala	Pro	Gly	Glu	Thr	Val		
	50				55					60							
Tyr	Tyr	Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu	Tyr	Thr	Ser		
65				70				75						80			
His	Ile	Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu	Gly	Pro	Glu		
			85				90						95				
Cys	Asp	Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr	Asn	Leu	Arg		
		100					105						110				
Val	Arg	Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala	Trp	Ser	Ile	Leu	Lys		
		115				120					125						
His	Pro	Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr	Arg	Pro	Gly	Met	Glu		
	130				135						140						
Ile	Pro	Lys	His	Gly	Phe	His	Leu	Val	Ile	Glu	Leu	Glu	Asp	Leu	Gly		
145				150					155					160			
Pro	Gln	Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Thr	Arg	Glu	Pro	Gly	Ala		
			165					170						175			
Glu	Glu	His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly	Ile	Pro	Val	His	Leu		
		180					185					190					
Glu	Thr	Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val	Lys	Ala	Gln	Thr	Phe		
		195				200						205					
Val	Lys	Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser	Gln	Thr	Glu	Cys	Val		
	210				215						220						
Glu	Val	Gln	Gly	Glu	Ala	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser		
225				230				235						240			
Gly	Gly	Gly	Gly	Ser	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe		

245								250					255					
Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys			
			260					265					270					
Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val			
			275				280					285						
Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln			
			290			295					300							
Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser			
305					310					315					320			
Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His			
			325						330					335				
Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys			
			340					345					350					
<210> SEQ ID NO 22																		
<211> LENGTH: 1801																		
<212> TYPE: DNA																		
<213> ORGANISM: Homo sapiens																		
<220> FEATURE:																		
<221> NAME/KEY: CDS																		
<222> LOCATION: (8) ... (1789)																		
<400> SEQUENCE: 22																		
gtcgcacc	atg	gat	gca	atg	aag	aga	ggg	ctc	tgc	tgt	gtg	ctg	ctg	ctg		49		
	Met	Asp	Ala	Met	Lys	Arg	Gly	Leu	Cys	Cys	Val	Leu	Leu	Leu				
	1				5				10									
tgt	ggc	gcc	gtc	ttc	gtt	tcg	ctc	agc	cag	gaa	atc	cat	gcc	gag	ttg	97		
Cys	Gly	Ala	Val	Phe	Val	Ser	Leu	Ser	Gln	Glu	Ile	His	Ala	Glu	Leu			
15				20						25					30			
aga	cgc	ttc	cgt	aga	gtt	ccc	tgt	gtc	tct	ggg	ggg	ttg	cct	aaa	cct	145		
Arg	Arg	Phe	Arg	Arg	Val	Pro	Cys	Val	Ser	Gly	Gly	Leu	Pro	Lys	Pro			
				35				40						45				
gca	aac	atc	acc	ttc	tta	tcc	atc	aac	atg	aag	aat	gtc	cta	caa	tgg	193		
Ala	Asn	Ile	Thr	Phe	Leu	Ser	Ile	Asn	Met	Lys	Asn	Val	Leu	Gln	Trp			
			50					55					60					
act	cca	cca	gag	ggg	ctt	caa	gga	gtt	aaa	gtt	act	tac	act	gtg	cag	241		
Thr	Pro	Pro	Glu	Gly	Leu	Gln	Gly	Val	Lys	Val	Thr	Tyr	Thr	Val	Gln			
		65					70					75						
tat	ttc	ata	tat	ggg	caa	aag	aaa	tgg	ctg	aat	aaa	tca	gaa	tgc	aga	289		
Tyr	Phe	Ile	Tyr	Gly	Gln	Lys	Lys	Trp	Leu	Asn	Lys	Ser	Glu	Cys	Arg			
		80				85					90							
aat	atc	aat	aga	acc	tac	tgt	gat	ctt	tct	gct	gaa	act	tct	gac	tac	337		
Asn	Ile	Asn	Arg	Thr	Tyr	Cys	Asp	Leu	Ser	Ala	Glu	Thr	Ser	Asp	Tyr			
95					100					105				110				
gaa	cac	cag	tat	tat	gcc	aaa	gtt	aag	gcc	att	tgg	gga	aca	aag	tgt	385		
Glu	His	Gln	Tyr	Tyr	Ala	Lys	Val	Lys	Ala	Ile	Trp	Gly	Thr	Lys	Cys			
				115					120					125				
tcc	aaa	tgg	gct	gaa	agt	gga	cgg	ttc	tat	cct	ttt	tta	gaa	aca	caa	433		
Ser	Lys	Trp	Ala	Glu	Ser	Gly	Arg	Phe	Tyr	Pro	Phe	Leu	Glu	Thr	Gln			
			130				135						140					
att	ggc	cca	cca	gag	gtg	gca	ctg	act	aca	gat	gag	aag	tcc	att	tct	481		
Ile	Gly	Pro	Pro	Glu	Val	Ala	Leu	Thr	Thr	Asp	Glu	Lys	Ser	Ile	Ser			
		145				150						155						
gtt	gtc	ctg	aca	gct	cca	gag	aag	tgg	aag	aga	aat	cca	gaa	gac	ctt	529		
Val	Val	Leu	Thr	Ala	Pro	Glu	Lys	Trp	Lys	Arg	Asn	Pro	Glu	Asp	Leu			
		160				165					170							
cct	gtt	tcc	atg	caa	caa	ata	tac	tcc	aat	ctg	aag	tat	aac	gtg	tct	577		

-continued

Pro	Val	Ser	Met	Gln	Gln	Ile	Tyr	Ser	Asn	Leu	Lys	Tyr	Asn	Val	Ser	
175					180					185					190	
gtg	ttg	aat	act	aaa	tca	aac	aga	acg	tgg	tcc	cag	tgt	gtg	acc	aac	625
Val	Leu	Asn	Thr	Lys	Ser	Asn	Arg	Thr	Trp	Ser	Gln	Cys	Val	Thr	Asn	
				195					200					205		
cac	acg	ctg	gtg	ctc	acc	tgg	ctg	gag	ccg	aac	act	ctt	tac	tgc	gta	673
His	Thr	Leu	Val	Leu	Thr	Trp	Leu	Glu	Pro	Asn	Thr	Leu	Tyr	Cys	Val	
			210					215					220			
cac	gtg	gag	tcc	ttc	gtc	cca	ggg	ccc	cct	cgc	cgt	gct	cag	cct	tct	721
His	Val	Glu	Ser	Phe	Val	Pro	Gly	Pro	Pro	Arg	Arg	Ala	Gln	Pro	Ser	
		225					230					235				
gag	aag	cag	tgt	gcc	agg	act	ttg	aaa	gat	caa	ggg	gga	ggc	ggg	tca	769
Glu	Lys	Gln	Cys	Ala	Arg	Thr	Leu	Lys	Asp	Gln	Gly	Gly	Gly	Gly	Ser	
	240					245					250					
ggc	gga	ggg	ggc	tct	ggc	ggg	ggc	gga	tgc	gcc	tcc	acc	aag	ggc	cca	817
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Ser	Ala	Ser	Thr	Lys	Gly	Pro		
	255				260				265					270		
tgc	gtc	ttc	ccc	ctg	gca	ccc	tcc	tcc	aag	agc	acc	tct	ggg	ggc	aca	865
Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	
				275					280					285		
gcg	gcc	ctg	ggc	tgc	ctg	gtc	aag	gac	tac	ttc	ccc	gaa	ccg	gtg	acg	913
Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	
		290						295					300			
gtg	tgc	tgg	aac	tca	ggc	gcc	ctg	acc	agc	ggc	gtg	cac	acc	ttc	ccg	961
Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	
		305					310					315				
gct	gtc	cta	cag	tcc	tca	gga	ctc	tac	tcc	ctc	agc	agc	gtg	gtg	acc	1009
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	
		320				325					330					
gtg	ccc	tcc	agc	agc	ttg	ggc	acc	cag	acc	tac	atc	tgc	aac	gtg	aat	1057
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	
	335				340					345				350		
cac	aag	ccc	agc	aac	acc	aag	gtg	gac	aag	aaa	gtt	gag	ccc	aaa	tct	1105
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	
			355					360						365		
tgt	gac	aaa	act	cac	aca	tgc	cca	ccg	tgc	cca	gca	cct	gaa	gcc	gag	1153
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Glu	
		370					375						380			
ggg	gca	ccg	tca	gtc	ttc	ctc	ttc	ccc	cca	aaa	ccc	aag	gac	acc	ctc	1201
Gly	Ala	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	
		385				390						395				
atg	atc	tcc	cgg	acc	cct	gag	gtc	aca	tgc	gtg	gtg	gtg	gac	gtg	agc	1249
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	
		400				405					410					
cac	gaa	gac	cct	gag	gtc	aag	ttc	aac	tgg	tac	gtg	gac	ggc	gtg	gag	1297
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	
	415				420					425				430		
gtg	cat	aat	gcc	aag	aca	aag	ccg	cgg	gag	gag	cag	tac	aac	agc	acg	1345
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	
			435					440					445			
tac	cgt	gtg	gtc	agc	gtc	ctc	acc	gtc	ctg	cac	cag	gac	tgg	ctg	aat	1393
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	
		450					455						460			
ggc	aag	gag	tac	aag	tgc	aag	gtc	tcc	aac	aaa	gcc	ctc	cca	tcc	tcc	1441
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ser	Ser	
		465				470					475					
atc	gag	aaa	acc	atc	tcc	aaa	gcc	aaa	ggg	cag	ccc	cga	gaa	cca	cag	1489

```
<210> SEQ ID NO 23
<211> LENGTH: 594
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23
```

Met 1	Asp	Ala	Met	Lys 5	Arg	Gly	Leu	Cys 10	Cys	Val	Leu	Leu	Leu	Cys 15	Gly
Ala	Val	Phe	Val 20	Ser	Leu	Ser	Gln	Glu 25	Ile	His	Ala	Glu	Leu 30	Arg	Arg
Phe	Arg	Arg 35	Val	Pro	Cys	Val	Ser 40	Gly	Gly	Leu	Pro	Lys 45	Pro	Ala	Asn
Ile	Thr 50	Phe	Leu	Ser	Ile	Asn 55	Met	Lys	Asn	Val	Leu 60	Gln	Trp	Thr	Pro
Pro 65	Glu	Gly	Leu	Gln	Gly 70	Val	Lys	Val	Thr	Tyr 75	Thr	Val	Gln	Tyr	Phe 80
Ile	Tyr	Gly	Gln	Lys 85	Lys	Trp	Leu	Asn	Lys 90	Ser	Glu	Cys	Arg	Asn 95	Ile
Asn	Arg	Thr	Tyr 100	Cys	Asp	Leu	Ser	Ala 105	Glu	Thr	Ser	Asp	Tyr 110	Glu	His
Gln	Tyr	Tyr 115	Ala	Lys	Val	Lys	Ala 120	Ile	Trp	Gly	Thr	Lys 125	Cys	Ser	Lys
Trp	Ala 130	Glu	Ser	Gly	Arg	Phe 135	Tyr	Pro	Phe	Leu	Glu 140	Thr	Gln	Ile	Gly
Pro 145	Pro	Glu	Val	Ala	Leu 150	Thr	Thr	Asp	Glu	Lys 155	Ser	Ile	Ser	Val	Val 160
Leu	Thr	Ala	Pro	Glu 165	Lys	Trp	Lys	Arg	Asn	Pro 170	Glu	Asp	Leu	Pro 175	Val
Ser	Met	Gln	Gln 180	Ile	Tyr	Ser	Asn	Leu 185	Lys	Tyr	Asn	Val	Ser 190	Val	Leu
Asn	Thr	Lys 195	Ser	Asn	Arg	Thr	Trp 200	Ser	Gln	Cys	Val	Thr 205	Asn	His	Thr

-continued

Leu Val Leu Thr Trp	Leu Glu Pro Asn Thr	Leu Tyr Cys Val His Val
210	215	220
Glu Ser Phe Val Pro	Gly Pro Pro Arg Arg	Ala Gln Pro Ser Glu Lys
225	230	235 240
Gln Cys Ala Arg Thr	Leu Lys Asp Gln Gly Gly Gly Ser Gly Gly	
245	250	255
Gly Gly Ser Gly Gly Gly Gly Ser	Ala Ser Thr Lys Gly Pro Ser Val	
260	265	270
Phe Pro Leu Ala Pro Ser Ser	Lys Ser Thr Ser Gly Gly Thr Ala Ala	
275	280	285
Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro	Glu Pro Val Thr Val Ser	
290	295	300
Trp Asn Ser Gly Ala Leu Thr Ser Gly Val	His Thr Phe Pro Ala Val	
305	310	315 320
Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser	Val Val Thr Val Pro	
325	330	335
Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys	Asn Val Asn His Lys	
340	345	350
Pro Ser Asn Thr Lys Val Asp Lys Lys Val	Glu Pro Lys Ser Cys Asp	
355	360	365
Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro	Glu Ala Glu Gly Ala	
370	375	380
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro	Lys Asp Thr Leu Met Ile	
385	390	395 400
Ser Arg Thr Pro Glu Val Thr Cys Val Val Asp	Val Ser His Glu	
405	410	415
Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp	Gly Val Glu Val His	
420	425	430
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr	Asn Ser Thr Tyr Arg	
435	440	445
Val Val Ser Val Leu Thr Val Leu His Gln Asp	Trp Leu Asn Gly Lys	
450	455	460
Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro	Ser Ser Ile Glu	
465	470	475 480
Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro	Gln Val Tyr	
485	490	495
Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys	Asn Gln Val Ser Leu	
500	505	510
Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp	Ile Ala Val Glu Trp	
515	520	525
Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr	Pro Pro Val	
530	535	540
Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr	Val Asp	
545	550	555 560
Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser	Val Met His	
565	570	575
Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser	Pro	
580	585	590
Gly Lys		

-continued

<210> SEQ ID NO 24
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

ggcgcggccat gcagactttc acaatgggtt 29

<210> SEQ ID NO 25
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 25

tccgctaccg cgcctccac tgccaccacc tccggcctct ccttgcaact cc 52

<210> SEQ ID NO 26
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

gtggaggcgg cggtagcgga gccggtggca gtcgaactgt ggctgcacca tct 53

<210> SEQ ID NO 27
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

ggcgcgcctc tagattaaca ctctcccctg ttgaagct 38

<210> SEQ ID NO 28
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

gtcgaccatg gatgcaatga agagagggt 30

<210> SEQ ID NO 29
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

cacaggaac tctacggaag cgtctcaact 30

<210> SEQ ID NO 30
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

cttcgtaga gttccctgtg tctctggtgg ttt 33

<210> SEQ ID NO 31
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

-continued

gccagagcca cctccgctg aaccgcctcc accttgatct ttcaaagtc tgg 53

<210> SEQ ID NO 32
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

cagggcggagg tggctctggc ggtggcggat cggcctccac caagggccca t 51

<210> SEQ ID NO 33
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

ctgggcacgg tgggcatgtg 20

<210> SEQ ID NO 34
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

cacatgcca ccgtgcccag 20

<210> SEQ ID NO 35
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

agatctagat tatttaccg gagacaggga g 31

<210> SEQ ID NO 36
 <211> LENGTH: 1806
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (38)...(1675)

<400> SEQUENCE: 36

cgccgcgttc ccgagatgtg acccgaaactg acagccc atg cac act ccc ggg acc 55
Met His Thr Pro Gly Thr
1 5

ccg gcg ccg ggc cac ccg gac ccg ccg cca ctg ttg ctg ctc acg ctg 103
Pro Ala Pro Gly His Pro Asp Pro Pro Leu Leu Leu Leu Thr Leu
10 15 20

ctt ctg ctg ctg gcc gct tgg gga cgc gca gtt cct tgt gtc ttc tgt 151
Leu Leu Leu Leu Ala Ala Ser Gly Arg Ala Val Pro Cys Val Phe Cys
25 30 35

ggt ttg cct aaa cct aca aat atc acc ttc tta tcc atc aac atg aag 199
Gly Leu Pro Lys Pro Thr Asn Ile Thr Phe Leu Ser Ile Asn Met Lys
40 45 50

aat gtc ctg cat tgg aat cca cca gag agt cta cac gga gtt gaa gtc 247
Asn Val Leu His Trp Asn Pro Pro Glu Ser Leu His Gly Val Glu Val
55 60 65 70

aca tac act gtg caa tat ttc ata tat ggg cag aag aaa tgg ctg aat 295
Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn

75						80						85							
gcc	tct	aaa	tgc	ggg	agt	atc	aac	agg	acc	tac	tgt	gac	ctt	tct	gtt	343			
Ala	Ser	Lys	Cys	Gly	Ser	Ile	Asn	Arg	Thr	Tyr	Cys	Asp	Leu	Ser	Val				
90						95						100							
gag	acc	tca	gac	tat	gaa	cac	cag	ttc	tat	gcc	aaa	gtg	aag	gcc	att	391			
Glu	Thr	Ser	Asp	Tyr	Glu	His	Gln	Phe	Tyr	Ala	Lys	Val	Lys	Ala	Ile				
105						110						115							
tgg	gaa	gcc	agg	tgc	tcc	gaa	tgg	gcc	gag	acg	gaa	cgc	ttc	tat	cct	439			
Trp	Glu	Ala	Arg	Cys	Ser	Glu	Trp	Ala	Glu	Thr	Glu	Arg	Phe	Tyr	Pro				
120						125						130							
ttc	ttg	gaa	act	caa	gtc	agc	cca	cca	gag	att	gcc	ctg	aca	act	ggc	487			
Phe	Leu	Glu	Thr	Gln	Val	Ser	Pro	Pro	Glu	Ile	Ala	Leu	Thr	Thr	Gly				
135						140						145						150	
gag	aag	tcc	atc	tct	att	gcc	ctg	aca	gca	cca	gag	aag	tgg	aaa	aga	535			
Glu	Lys	Ser	Ile	Ser	Ile	Ala	Leu	Thr	Ala	Pro	Glu	Lys	Trp	Lys	Arg				
155						160						165							
aat	cca	caa	gac	cac	act	gtt	tct	atg	caa	cag	ata	tac	ccc	aat	ttg	583			
Asn	Pro	Gln	Asp	His	Thr	Val	Ser	Met	Gln	Gln	Ile	Tyr	Pro	Asn	Leu				
170						175						180							
aag	tac	aat	gtg	tct	gtg	tat	aac	act	aag	tcg	aga	aga	acg	tgg	tcc	631			
Lys	Tyr		Val	Ser	Val	Tyr	Asn	Thr	Lys	Ser	Arg	Arg	Thr	Trp	Ser				
185						190						195							
cag	tgt	gtc	acc	aac	agc	aca	ctg	gtc	ctc	agc	tgg	ctg	gag	ccc	aac	679			
Gln	Cys	Val	Thr	Asn	Ser	Thr	Leu	Val	Leu	Ser	Trp	Leu	Glu	Pro	Asn				
200						205						210							
act	ctg	tat	tgt	gtc	cac	gtg	gag	tcc	ctt	gtc	cca	ggg	ccc	cct	cgc	727			
Thr	Leu	Tyr	Cys	Val	His	Val	Glu	Ser	Leu	Val	Pro	Gly	Pro	Pro	Arg				
215						220						225						230	
ctc	ccg	atg	cct	tct	cag	aag	cag	tgc	atc	agt	act	ttg	gaa	gtt	caa	775			
Leu	Pro	Met	Pro		Ser	Gln	Lys	Gln	Cys	Ile	Ser	Thr	Leu	Glu	Val	Gln			
235						240						245							
aca	tca	gca	tgg	aag	gct	aaa	gtc	atc	ttc	tgg	tat	gtc	ttc	ctc	aca	823			
Thr	Ser	Ala	Trp	Lys	Ala	Lys	Val	Ile	Phe	Trp	Tyr	Val	Phe	Leu	Thr				
250						255						260							
tct	gtt	atc	gtg	ttt	ctt	ttc	tcc	gca	att	ggc	tac	ttg	gtt	tac	cgt	871			
Ser	Val	Ile	Val	Phe	Leu	Phe	Ser	Ala	Ile	Gly	Tyr	Leu	Val	Tyr	Arg				
265						270						275							
tac	atc	cat	gtt	ggc	aag	gaa	aaa	cac	cca	gca	aat	ttg	gta	ctg	att	919			
Tyr	Ile	His	Val	Gly	Lys	Glu	Lys	His	Pro	Ala	Asn	Leu	Val	Leu	Ile				
280						285						290							
tat	aga	aat	gaa	att	ggc	aca	aga	gtc	ttt	gaa	cct	act	gaa	aca	atc	967			
Tyr	Arg	Asn	Glu	Ile	Gly	Thr	Arg	Val	Phe	Glu	Pro	Thr	Glu	Thr	Ile				
295						300						305						310	
aca	ctt	aat	ttt	atc	acc	ttc	agt	atg	ttg	gat	gat	act	aaa	att	tct	1015			
Thr	Leu	Asn	Phe	Ile	Thr	Phe	Ser	Met	Leu	Asp	Asp	Thr	Lys	Ile	Ser				
315						320						325							
cca	aag	gat	atg	aat	tta	ctg	gac	aaa	agc	agt	gat	gac	atc	agt	gtt	1063			
Pro	Lys	Asp	Met	Asn	Leu	Leu	Asp	Lys	Ser	Ser	Asp	Asp	Ile	Ser	Val				
330						335						340							
aat	gac	cct	gag	cac	aat	gag	gcc	tgg	gag	ccg	cac	tgg	gag	gag	gtg	1111			
Asn	Asp	Pro	Glu	His	Asn	Glu	Ala	Trp	Glu	Pro	His	Trp	Glu	Glu	Val				
345						350						355							
gag	ggg	caa	cat	tta	gga	tgc	tct	tcg	cat	ttg	atg	gac	gct	gtc	tgt	1159			
Glu	Gly	Gln	His	Leu	Gly	Cys	Ser	Ser	His	Leu	Met	Asp	Ala	Val	Cys				
360						365						370							
ggt	gct	gag	caa	aga	gac	gga	gac	acc	tcc	cta	acc	cag	cat	ggg	tgg	1207			
Gly	Ala	Glu	Gln	Arg	Asp	Gly	Gly	Asp	Thr	Ser	Leu	Thr	Gln	His	Gly	Trp			

-continued

375	380	385	390	
ctt aac agc acc atc ccc aca gga gag aca gac act gag cct caa tac				1255
Leu Asn Ser Thr Ile Pro Thr Gly Glu Thr Asp Thr Glu Pro Gln Tyr				
	395	400	405	
aaa gtc cta agt gac ttc tac ggg gag ggt gaa atc caa ctg tcc tgt				1303
Lys Val Leu Ser Asp Phe Tyr Gly Glu Gly Glu Ile Gln Leu Ser Cys				
	410	415	420	
gag ccg gaa gag gcg gcc aga aca gag aaa ata tct gag cca ctg gtg				1351
Glu Pro Glu Glu Ala Ala Arg Thr Glu Lys Ile Ser Glu Pro Leu Val				
	425	430	435	
act tca gca aac ttg gac cca cag ctt gaa gac cta cat cac ctg ggt				1399
Thr Ser Ala Asn Leu Asp Pro Gln Leu Glu Asp Leu His His Leu Gly				
	440	445	450	
cag gag cat act gtc tcc gag gat ggg cca gag gaa gag aca tct ata				1447
Gln Glu His Thr Val Ser Glu Asp Gly Pro Glu Glu Glu Thr Ser Ile				
	455	460	465	470
aca gta gtg gat tgg gac cct caa act ggc agg ctg tgt atc cct tcc				1495
Thr Val Val Asp Trp Asp Pro Gln Thr Gly Arg Leu Cys Ile Pro Ser				
	475	480	485	
tta cct atc ttt ggc cgt gat cct gag aac tat ggt cat tat gag aga				1543
Leu Pro Ile Phe Gly Arg Asp Pro Glu Asn Tyr Gly His Tyr Glu Arg				
	490	495	500	
gac cag ctc tta gag ggt ggc ctt ttg tct aga ctc tat gag aac cag				1591
Asp Gln Leu Leu Glu Gly Gly Leu Leu Ser Arg Leu Tyr Glu Asn Gln				
	505	510	515	
gca cct gac aag cca gag aaa gaa aat gaa aac tgt ctc aca cgg ttt				1639
Ala Pro Asp Lys Pro Glu Lys Glu Asn Glu Asn Cys Leu Thr Arg Phe				
	520	525	530	
atg gag gaa tgg ggg tta cat gta caa atg gaa agc tagtgccagg				1685
Met Glu Glu Trp Gly Leu His Val Gln Met Glu Ser				
	535	540	545	
ctttctgttg actgccaaca aatgaaggaa ccattcccagg gggatgaacag tgttcaggtt				1745
atcagtggtca gcaatgagac tgttctctct gttcatgaac tttgtcagcc ctgcctcatc				1805
c				1806

<210> SEQ ID NO 37

<211> LENGTH: 546

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 37

Met His Thr Pro Gly Thr Pro Ala Pro Gly His Pro Asp Pro Pro Pro				
1	5	10	15	
Leu Leu Leu Leu Thr Leu Leu Leu Leu Leu Ala Ala Ser Gly Arg Ala				
	20	25	30	
Val Pro Cys Val Phe Cys Gly Leu Pro Lys Pro Thr Asn Ile Thr Phe				
	35	40	45	
Leu Ser Ile Asn Met Lys Asn Val Leu His Trp Asn Pro Pro Glu Ser				
	50	55	60	
Leu His Gly Val Glu Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly				
	65	70	75	80
Gln Lys Lys Trp Leu Asn Ala Ser Lys Cys Gly Ser Ile Asn Arg Thr				
	85	90	95	
Tyr Cys Asp Leu Ser Val Glu Thr Ser Asp Tyr Glu His Gln Phe Tyr				
	100	105	110	

-continued

Ala Lys Val Lys Ala Ile Trp Glu Ala Arg Cys Ser Glu Trp Ala Glu	115	120	125
Thr Glu Arg Phe Tyr Pro Phe Leu Glu Thr Gln Val Ser Pro Pro Glu	130	135	140
Ile Ala Leu Thr Thr Gly Glu Lys Ser Ile Ser Ile Ala Leu Thr Ala	145	150	155
Pro Glu Lys Trp Lys Arg Asn Pro Gln Asp His Thr Val Ser Met Gln	165	170	175
Gln Ile Tyr Pro Asn Leu Lys Tyr Asn Val Ser Val Tyr Asn Thr Lys	180	185	190
Ser Arg Arg Thr Trp Ser Gln Cys Val Thr Asn Ser Thr Leu Val Leu	195	200	205
Ser Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Leu	210	215	220
Val Pro Gly Pro Pro Arg Leu Pro Met Pro Ser Gln Lys Gln Cys Ile	225	230	235
Ser Thr Leu Glu Val Gln Thr Ser Ala Trp Lys Ala Lys Val Ile Phe	245	250	255
Trp Tyr Val Phe Leu Thr Ser Val Ile Val Phe Leu Phe Ser Ala Ile	260	265	270
Gly Tyr Leu Val Tyr Arg Tyr Ile His Val Gly Lys Glu Lys His Pro	275	280	285
Ala Asn Leu Val Leu Ile Tyr Arg Asn Glu Ile Gly Thr Arg Val Phe	290	295	300
Glu Pro Thr Glu Thr Ile Thr Leu Asn Phe Ile Thr Phe Ser Met Leu	305	310	315
Asp Asp Thr Lys Ile Ser Pro Lys Asp Met Asn Leu Leu Asp Lys Ser	325	330	335
Ser Asp Asp Ile Ser Val Asn Asp Pro Glu His Asn Glu Ala Trp Glu	340	345	350
Pro His Trp Glu Glu Val Glu Gly Gln His Leu Gly Cys Ser Ser His	355	360	365
Leu Met Asp Ala Val Cys Gly Ala Glu Gln Arg Asp Gly Asp Thr Ser	370	375	380
Leu Thr Gln His Gly Trp Leu Asn Ser Thr Ile Pro Thr Gly Glu Thr	385	390	395
Asp Thr Glu Pro Gln Tyr Lys Val Leu Ser Asp Phe Tyr Gly Glu Gly	405	410	415
Glu Ile Gln Leu Ser Cys Glu Pro Glu Glu Ala Ala Arg Thr Glu Lys	420	425	430
Ile Ser Glu Pro Leu Val Thr Ser Ala Asn Leu Asp Pro Gln Leu Glu	435	440	445
Asp Leu His His Leu Gly Gln Glu His Thr Val Ser Glu Asp Gly Pro	450	455	460
Glu Glu Glu Thr Ser Ile Thr Val Val Asp Trp Asp Pro Gln Thr Gly	465	470	475
Arg Leu Cys Ile Pro Ser Leu Pro Ile Phe Gly Arg Asp Pro Glu Asn	485	490	495
Tyr Gly His Tyr Glu Arg Asp Gln Leu Leu Glu Gly Gly Leu Leu Ser	500	505	510

<400> SEQUENCE: 38

<400> SEQUENCE: 39

Val	Pro	Cys	Val	Phe	Cys	Gly	Leu	Pro	Lys	Pro	Thr	Asn	Ile	Thr	Phe
1				5					10					15	
Leu	Ser	Ile	Asn	Met	Lys	Asn	Val	Leu	His	Trp	Asn	Pro	Pro	Glu	Ser
			20					25					30		
Leu	His	Gly	Val	Glu	Val	Thr	Tyr	Thr	Val	Gln	Tyr	Phe	Ile	Tyr	Gly
		35					40					45			

-continued

Gln	Lys	Lys	Trp	Leu	Asn	Ala	Ser	Lys	Cys	Gly	Ser	Ile	Asn	Arg	Thr
50						55				60					
Tyr	Cys	Asp	Leu	Ser	Val	Glu	Thr	Ser	Asp	Tyr	Glu	His	Gln	Phe	Tyr
65					70				75					80	
Ala	Lys	Val	Lys	Ala	Ile	Trp	Glu	Ala	Arg	Cys	Ser	Glu	Trp	Ala	Glu
			85						90					95	
Thr	Glu	Arg	Phe	Tyr	Pro	Phe	Leu	Glu	Thr	Gln	Val	Ser	Pro	Pro	Glu
		100					105						110		
Ile	Ala	Leu	Thr	Thr	Gly	Glu	Lys	Ser	Ile	Ser	Ile	Ala	Leu	Thr	Ala
		115				120						125			
Pro	Glu	Lys	Trp	Lys	Arg	Asn	Pro	Gln	Asp	His	Thr	Val	Ser	Met	Gln
	130					135					140				
Gln	Ile	Tyr	Pro	Asn	Leu	Lys	Tyr	Asn	Val	Ser	Val	Tyr	Asn	Thr	Lys
145					150					155					160
Ser	Arg	Arg	Thr	Trp	Ser	Gln	Cys	Val	Thr	Asn	Ser	Thr	Leu	Val	Leu
			165						170					175	
Ser	Trp	Leu	Glu	Pro	Asn	Thr	Leu	Tyr	Cys	Val	His	Val	Glu	Ser	Leu
		180					185						190		
Val	Pro	Gly	Pro	Pro	Arg	Leu	Pro	Met	Pro	Ser	Gln	Lys	Gln	Cys	Ile
		195					200					205			
Ser	Thr	Leu	Glu	Val	Gln	Thr	Ser	Ala	Trp	Lys	Ala	Lys	Val	Ile	Phe
	210					215					220				
Trp	Tyr	Val	Phe	Leu	Thr	Ser	Val	Ile	Val	Phe	Leu	Phe	Ser	Ala	Ile
225					230					235					240
Gly	Tyr	Leu	Val	Tyr	Arg	Tyr	Ile	His	Val	Gly	Lys	Glu	Lys	His	Pro
			245						250					255	
Ala	Asn	Leu	Val	Leu	Ile	Tyr	Arg	Asn	Glu	Ile	Gly	Thr	Arg	Val	Phe
		260						265					270		
Glu	Pro	Thr	Glu	Thr	Ile	Thr	Leu	Asn	Phe	Ile	Thr	Phe	Ser	Met	Leu
		275					280					285			
Asp	Asp	Thr	Lys	Ile	Ser	Pro	Lys	Asp	Met	Asn	Leu	Leu	Asp	Lys	Ser
	290					295					300				
Ser	Asp	Asp	Ile	Ser	Val	Asn	Asp	Pro	Glu	His	Asn	Glu	Ala	Trp	Glu
305					310					315					320
Pro	His	Trp	Glu	Glu	Val	Glu	Gly	Gln	His	Leu	Gly	Cys	Ser	Ser	His
			325					330						335	
Leu	Met	Asp	Ala	Val	Cys	Gly	Ala	Glu	Gln	Arg	Asp	Gly	Asp	Thr	Ser
		340					345						350		
Leu	Thr	Gln	His	Gly	Trp	Leu	Asn	Ser	Thr	Ile	Pro	Thr	Gly	Glu	Thr
		355					360					365			
Asp	Thr	Glu	Pro	Gln	Tyr	Lys	Val	Leu	Ser	Asp	Phe	Tyr	Gly	Glu	Gly
	370					375					380				
Glu	Ile	Gln	Leu	Ser	Cys	Glu	Pro	Glu	Glu	Ala	Ala	Arg	Thr	Glu	Lys
385					390					395					400
Ile	Ser	Glu	Pro	Leu	Val	Thr	Ser	Ala	Asn	Leu	Asp	Pro	Gln	Leu	Glu
			405						410					415	
Asp	Leu	His	His	Leu	Gly	Gln	Glu	His	Thr	Val	Ser	Glu	Asp	Gly	Pro
		420					425						430		
Glu	Glu	Glu	Thr	Ser	Ile	Thr	Val	Val	Asp	Trp	Asp	Pro	Gln	Thr	Gly
		435					440					445			
Arg	Leu	Cys	Ile	Pro	Ser	Leu	Pro	Ile	Phe	Gly	Arg	Asp	Pro	Glu	Asn

-continued

450	455	460	
Tyr Gly His Tyr Glu Arg Asp Gln Leu Leu Glu Gly Gly Leu Leu Ser			
465	470	475	480
Arg Leu Tyr Glu Asn Gln Ala Pro Asp Lys Pro Glu Lys Glu Asn Glu			
	485	490	495
Asn Cys Leu Thr Arg Phe Met Glu Glu Trp Gly Leu His Val Gln Met			
	500	505	510
Glu Ser			
<210> SEQ ID NO 40			
<211> LENGTH: 18			
<212> TYPE: DNA			
<213> ORGANISM: Mus musculus			
<400> SEQUENCE: 40			
cgccgcggttc ccgagatg			18
<210> SEQ ID NO 41			
<211> LENGTH: 24			
<212> TYPE: DNA			
<213> ORGANISM: Mus musculus			
<400> SEQUENCE: 41			
ggatgaggca gggctgacaa agtt			24
<210> SEQ ID NO 42			
<211> LENGTH: 36			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 42			
acttgtggaa ttcgctagca ccaagggccc atcggt			36
<210> SEQ ID NO 43			
<211> LENGTH: 32			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 43			
gcctagaacg cgttcattta cccggagaca gg			32
<210> SEQ ID NO 44			
<211> LENGTH: 8			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 44			
aattgaga			8
<210> SEQ ID NO 45			
<211> LENGTH: 8			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 45			
cgcggtctc			8
<210> SEQ ID NO 46			
<211> LENGTH: 37			

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46
gtcacttgaa ttcggtaccg cctctgttgt gtgcctg          37

<210> SEQ ID NO 47
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47
gacctgaacg cgtctaacac tctccctgt tg              32

<210> SEQ ID NO 48
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48
tcagtcggaa ttcgcagaag ccatgcgggc tcccggcc          38

<210> SEQ ID NO 49
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49
ctgtgacgct agcctctgat gattgatctt tcaaa          35

<210> SEQ ID NO 50
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50
gatgtctgaa ttcgcagaag ccatgcagac tttcacaatg gtt          43

<210> SEQ ID NO 51
<211> LENGTH: 86
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51
aagacggtac cagatttcaa ctgctcatca gatggcggga agatgaagac agatgggtgca          60
gccacagtgg cctctccttg cacctc                      86

<210> SEQ ID NO 52
<211> LENGTH: 1720
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)...(1713)

<400> SEQUENCE: 52
atg cgg gct ccc ggc cgc ccg gcc ctg cgg ccg ctg ctg ctg ttg ctc          48
Met Arg Ala Pro Gly Arg Pro Ala Leu Arg Pro Leu Leu Leu Leu Leu
  1             5             10             15

ctg gcg gcg cct tgg gga cgg gca gtt ccc tgt gtc tct ggt ggt ttg          96
Leu Ala Ala Pro Trp Gly Arg Ala Val Pro Cys Val Ser Gly Gly Leu

```

-continued

	20	25	30	
cct aaa cct gca aac atc acc ttc tta tcc atc aac atg aag aat gtc				144
Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn Val				
	35	40	45	
cta caa tgg act cca cca gag ggt ctt caa gga gtt aaa gtt act tac				192
Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly Val Lys Val Thr Tyr				
	50	55	60	
act gtg cag tat ttc ata tat ggg caa aag aaa tgg ctg aat aaa tca				240
Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn Lys Ser				
	65	70	75	80
gaa tgc aga aat atc aat aga acc tac tgt gat ctt tct gct gaa act				288
Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu Thr				
	85	90	95	
tct gac tac gaa cac cag tat tat gcc aaa gtt aag gcc att tgg gga				336
Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys Ala Ile Trp Gly				
	100	105	110	
aca aag tgt tcc aaa tgg gct gaa agt gga cgg ttc tat cct ttt tta				384
Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg Phe Tyr Pro Phe Leu				
	115	120	125	
gaa aca caa att ggc cca cca gag gtg gca ctg act aca gat gag aag				432
Glu Thr Gln Ile Gly Pro Pro Glu Val Ala Leu Thr Thr Asp Glu Lys				
	130	135	140	
tcc att tct gtt gtc ctg aca gct cca gag aag tgg aag aga aat cca				480
Ser Ile Ser Val Val Leu Thr Ala Pro Glu Lys Trp Lys Arg Asn Pro				
	145	150	155	160
gaa gac ctt cct gtt tcc atg caa caa ata tac tcc aat ctg aag tat				528
Glu Asp Leu Pro Val Ser Met Gln Gln Ile Tyr Ser Asn Leu Lys Tyr				
	165	170	175	
aac gtg tct gtg ttg aat act aaa tca aac aga acg tgg tcc cag tgt				576
Asn Val Ser Val Leu Asn Thr Lys Ser Asn Arg Thr Trp Ser Gln Cys				
	180	185	190	
gtg acc aac cac acg ctg gtg ctc acc tgg ctg gag ccg aac act ctt				624
Val Thr Asn His Thr Leu Val Leu Thr Trp Leu Glu Pro Asn Thr Leu				
	195	200	205	
tac tgc gta cac gtg gag tcc ttc gtc cca ggg ccc cct cgc cgt gct				672
Tyr Cys Val His Val Glu Ser Phe Val Pro Gly Pro Pro Arg Arg Ala				
	210	215	220	
cag cct tct gag aag cag tgt gcc agg act ttg aaa gat caa tca tca				720
Gln Pro Ser Glu Lys Gln Cys Ala Arg Thr Leu Lys Asp Gln Ser Ser				
	225	230	235	240
gag gct agc acc aag ggc cca tcg gtc ttc ccc ctg gca ccc tcc tcc				768
Glu Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser				
	245	250	255	
aag agc acc tct ggg ggc aca gcg gcc ctg ggc tgc ctg gtc aag gac				816
Lys Ser Thr Ser Gly Gly Thr Ala Leu Gly Cys Leu Val Lys Asp				
	260	265	270	
tac ttc ccc gaa ccg gtg acg gtg tcg tgg aac tca ggc gcc ctg acc				864
Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr				
	275	280	285	
agc ggc gtg cac acc ttc ccg gct gtc cta cag tcc tca gga ctc tac				912
Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr				
	290	295	300	
tcc ctc agc agc gtg gtg acc gtg ccc tcc agc agc ttg ggc acc cag				960
Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln				
	305	310	315	320
acc tac atc tgc aac gtg aat cac aag ccc agc aac acc aag gtg gac				1008
Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp				

-continued

325	330	335	
aag aaa gtt gag ccc aaa tct tgt gac aaa act cac aca tgc cca ccg Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro 340 345 350			1056
tgc cca gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro 355 360 365			1104
cca aaa ccc aag gac acc ctc atg atc tcc cgg acc cct gag gtc aca Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr 370 375 380			1152
tgc gtg gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc aac Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn 385 390 395 400			1200
tgg tac gtg gac ggc gtg gag gtg cat aat gcc aag aca aag ccg cgg Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg 405 410 415			1248
gag gag cag tac aac agc acg tac cgt gtg gtc agc gtc ctc acc gtc Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val 420 425 430			1296
ctg cac cag gac tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser 435 440 445			1344
aac aaa gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys 450 455 460			1392
ggg cag ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp 465 470 475 480			1440
gag ctg acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe 485 490 495			1488
tat ccc agc gac atc gcc gtg gag tgg gag agc aat ggg cag ccg gag Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu 500 505 510			1536
aac aac tac aag acc acg cct ccc gtg ctg gac tcc gac ggc tcc ttc Asn Asn Tyr Lys Thr Thr Pro Val Leu Asp Ser Asp Gly Ser Phe 515 520 525			1584
ttc ctc tac agc aag ctc acc gtg gac aag agc agg tgg cag cag ggg Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly 530 535 540			1632
aac gtc ttc tca tgc tcc gtg atg cat gag gct ctg cac aac cac tac Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr 545 550 555 560			1680
acg cag aag agc ctc tcc ctg tct ccg ggt aaa tgacgcg Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 565 570			1720

<210> SEQ ID NO 53

<211> LENGTH: 571

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

Met Arg Ala Pro Gly Arg Pro Ala Leu Arg Pro Leu Leu Leu Leu Leu
1 5 10 15

Leu Ala Ala Pro Trp Gly Arg Ala Val Pro Cys Val Ser Gly Gly Leu
20 25 30

Pro	Lys	Pro	Ala	Asn	Ile	Thr	Phe	Leu	Ser	Ile	Asn	Met	Lys	Asn	Val
	35							40				45			
Leu	Gln	Trp	Thr	Pro	Pro	Glu	Gly	Leu	Gln	Gly	Val	Lys	Val	Thr	Tyr
	50					55					60				
Thr	Val	Gln	Tyr	Phe	Ile	Tyr	Gly	Gln	Lys	Lys	Trp	Leu	Asn	Lys	Ser
65					70					75					80
Glu	Cys	Arg	Asn	Ile	Asn	Arg	Thr	Tyr	Cys	Asp	Leu	Ser	Ala	Glu	Thr
			85						90					95	
Ser	Asp	Tyr	Glu	His	Gln	Tyr	Tyr	Ala	Lys	Val	Lys	Ala	Ile	Trp	Gly
			100					105					110		
Thr	Lys	Cys	Ser	Lys	Trp	Ala	Glu	Ser	Gly	Arg	Phe	Tyr	Pro	Phe	Leu
		115					120					125			
Glu	Thr	Gln	Ile	Gly	Pro	Pro	Glu	Val	Ala	Leu	Thr	Thr	Asp	Glu	Lys
	130					135						140			
Ser	Ile	Ser	Val	Val	Leu	Thr	Ala	Pro	Glu	Lys	Trp	Lys	Arg	Asn	Pro
145					150					155					160
Glu	Asp	Leu	Pro	Val	Ser	Met	Gln	Gln	Ile	Tyr	Ser	Asn	Leu	Lys	Tyr
			165						170					175	
Asn	Val	Ser	Val	Leu	Asn	Thr	Lys	Ser	Asn	Arg	Thr	Trp	Ser	Gln	Cys
		180						185					190		
Val	Thr	Asn	His	Thr	Leu	Val	Leu	Thr	Trp	Leu	Glu	Pro	Asn	Thr	Leu
	195						200					205			
Tyr	Cys	Val	His	Val	Glu	Ser	Phe	Val	Pro	Gly	Pro	Pro	Arg	Arg	Ala
	210				215						220				
Gln	Pro	Ser	Glu	Lys	Gln	Cys	Ala	Arg	Thr	Leu	Lys	Asp	Gln	Ser	Ser
225					230						235				240
Glu	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser
			245						250					255	
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp
			260					265					270		
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr
	275						280					285			
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr
	290				295						300				
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln
305				310						315					320
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp
			325					330						335	
Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro
		340						345					350		
Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
	355						360					365			
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
	370				375							380			
Cys	Val	Val	Val	Asp	Val	Ser	His	G							

-continued

435					440					445					
Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys
450					455						460				
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp
465					470					475					480
Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe
				485					490					495	
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu
			500					505					510		
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe
		515					520					525			
Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly
	530					535					540				
Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr
545					550					555					560
Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys					
				565					570						

<210> SEQ ID NO 54

<211> LENGTH: 547

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 54

Val	Pro	Cys	Val	Ser	Gly	Gly	Leu	Pro	Lys	Pro	Ala	Asn	Ile	Thr	Phe
1				5					10					15	
Leu	Ser	Ile	Asn	Met	Lys	Asn	Val	Leu	Gln	Trp	Thr	Pro	Pro	Glu	Gly
			20					25					30		
Leu	Gln	Gly	Val	Lys	Val	Thr	Tyr	Thr	Val	Gln	Tyr	Phe	Ile	Tyr	Gly
		35					40					45			
Gln	Lys	Lys	Trp	Leu	Asn	Lys	Ser	Glu	Cys	Arg	Asn	Ile	Asn	Arg	Thr
		50				55					60				
Tyr	Cys	Asp	Leu	Ser	Ala	Glu	Thr	Ser	Asp	Tyr	Glu	His	Gln	Tyr	Tyr
65					70				75					80	
Ala	Lys	Val	Lys	Ala	Ile	Trp	Gly	Thr	Lys	Cys	Ser	Lys	Trp	Ala	Glu
			85					90						95	
Ser	Gly	Arg	Phe	Tyr	Pro	Phe	Leu	Glu	Thr	Gln	Ile	Gly	Pro	Pro	Glu
		100					105						110		
Val	Ala	Leu	Thr	Thr	Asp	Glu	Lys	Ser	Ile	Ser	Val	Val	Leu	Thr	Ala
		115				120						125			
Pro	Glu	Lys	Trp	Lys	Arg	Asn	Pro	Glu	Asp	Leu	Pro	Val	Ser	Met	Gln
	130					135					140				
Gln	Ile	Tyr	Ser	Asn	Leu	Lys	Tyr	Asn	Val	Ser	Val	Leu	Asn	Thr	Lys
145					150					155					160
Ser	Asn	Arg	Thr	Trp	Ser	Gln	Cys	Val	Thr	Asn	His	Thr	Leu	Val	Leu
			165						170					175	
Thr	Trp	Leu	Glu	Pro	Asn	Thr	Leu	Tyr	Cys	Val	His	Val	Glu	Ser	Phe
		180					185						190		
Val	Pro	Gly	Pro	Pro	Arg	Arg	Ala	Gln	Pro	Ser	Glu	Lys	Gln	Cys	Ala
		195					200					205			
Arg	Thr	Leu	Lys	Asp	Gln	Ser	Ser	Glu	Ala	Ser	Thr	Lys	Gly	Pro	Ser
	210					215						220			

-continued

Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	225	230	235	240
Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	245	250	255	
Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	260	265	270	
Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	275	280	285	
Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	290	295	300	
Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	305	310	315	320
Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	325	330	335	
Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	340	345	350	
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	355	360	365	
Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	370	375	380	
His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	385	390	395	400
Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	405	410	415	
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	420	425	430	
Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	435	440	445	
Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	450	455	460	
Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	465	470	475	480
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	485	490	495	
Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	500	505	510	
Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	515	520	525	
His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	530	535	540	
Pro	Gly	Lys														545			

<210> SEQ ID NO 55

<211> LENGTH: 217

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

Val	Pro	Cys	Val	Ser	Gly	Gly	Leu	Pro	Lys	Pro	Ala	Asn	Ile	Thr	Phe	1	5	10	15
Leu	Ser	Ile	Asn	Met	Lys	Asn	Val	Leu	Gln	Trp	Thr	Pro	Pro	Glu	Gly	20	25	30	

-continued

Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly
 35 40 45
 Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr
 50 55 60
 Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr
 65 70 75 80
 Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu
 85 90 95
 Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu
 100 105 110
 Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala
 115 120 125
 Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln
 130 135 140
 Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys
 145 150 155 160
 Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu
 165 170 175
 Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe
 180 185 190
 Val Pro Gly Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala
 195 200 205
 Arg Thr Leu Lys Asp Gln Ser Ser Glu
 210 215

<210> SEQ ID NO 56
 <211> LENGTH: 1011
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1) ... (1008)

<400> SEQUENCE: 56

atg cag act ttc aca atg gtt cta gaa gaa atc tgg aca agt ctt ttc	48
Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser Leu Phe	
1 5 10 15	
atg tgg ttt ttc tac gca ttg att cca tgt ttg ctc aca gat gaa gtg	96
Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr Asp Glu Val	
20 25 30	
gcc att ctg cct gcc cct cag aac ctc tct gta ctc tca acc aac atg	144
Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met	
35 40 45	
aag cat ctc ttg atg tgg agc cca gtg atc gcg cct gga gaa aca gtg	192
Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val	
50 55 60	
tac tat tct gtc gaa tac cag ggg gag tac gag agc ctg tac acg agc	240
Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser	
65 70 75 80	
cac atc tgg atc ccc agc agc tgg tgc tca ctc act gaa ggt cct gag	288
His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu	
85 90 95	
tgt gat gtc act gat gac atc acg gcc act gtg cca tac aac ctt cgt	336
Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg	
100 105 110	

-continued

gtc agg gcc aca ttg ggc tca cag acc tca gcc tgg agc atc ctg aag	384
Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys	
115 120 125	
cat ccc ttt aat aga aac tca acc atc ctt acc cga cct ggg atg gag	432
His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu	
130 135 140	
atc acc aaa gat ggc ttc cac ctg gtt att gag ctg gag gac ctg ggg	480
Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly	
145 150 155 160	
ccc cag ttt gag ttc ctt gtg gcc tac tgg agg agg gag cct ggt gcc	528
Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu Pro Gly Ala	
165 170 175	
gag gaa cat gtc aaa atg gtg agg agt ggg ggt att cca gtg cac cta	576
Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu	
180 185 190	
gaa acc atg gag cca ggg gct gca tac tgt gtg aag gcc cag aca ttc	624
Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe	
195 200 205	
gtg aag gcc att ggg agg tac agc gcc ttc agc cag aca gaa tgt gtg	672
Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val	
210 215 220	
gag gtg caa gga gag gcc act gtg gct gca cca tct gtc ttc atc ttc	720
Glu Val Gln Gly Glu Ala Thr Val Ala Ala Pro Ser Val Phe Ile Phe	
225 230 235 240	
ccg cca tct gat gag cag ttg aaa tct ggt acc gcc tct gtt gtg tgc	768
Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys	
245 250 255	
ctg ctg aat aac ttc tat ccc aga gag gcc aaa gta cag tgg aag gtg	816
Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val	
260 265 270	
gat aac gcc ctc caa tcg ggt aac tcc cag gag agt gtc aca gag cag	864
Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln	
275 280 285	
gac agc aag gac agc acc tac agc ctc agc agc acc ctg acg ctg agc	912
Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser	
290 295 300	
aaa gca gac tac gag aaa cac aaa gtc tac gcc tgc gaa gtc acc cat	960
Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His	
305 310 315 320	
cag ggc ctg agc tcg ccc gtc aca aag agc ttc aac agg gga gag tgt	1008
Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys	
325 330 335	
tag	1011

<210> SEQ ID NO 57

<211> LENGTH: 336

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 57

Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser Leu Phe
1 5 10 15

Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr Asp Glu Val
20 25 30

Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met
35 40 45

Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val

-continued

50					55					60					
Tyr	Tyr	Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu	Tyr	Thr	Ser
65					70					75					80
His	Ile	Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu	Gly	Pro	Glu
				85					90					95	
Cys	Asp	Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr	Asn	Leu	Arg
				100				105						110	
Val	Arg	Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala	Trp	Ser	Ile	Leu	Lys
				115				120						125	
His	Pro	Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr	Arg	Pro	Gly	Met	Glu
				130				135						140	
Ile	Thr	Lys	Asp	Gly	Phe	His	Leu	Val	Ile	Glu	Leu	Glu	Asp	Leu	Gly
145					150					155					160
Pro	Gln	Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Arg	Arg	Glu	Pro	Gly	Ala
				165					170					175	
Glu	Glu	His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly	Ile	Pro	Val	His	Leu
				180					185					190	
Glu	Thr	Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val	Lys	Ala	Gln	Thr	Phe
				195				200						205	
Val	Lys	Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser	Gln	Thr	Glu	Cys	Val
				210				215						220	
Glu	Val	Gln	Gly	Glu	Ala	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe
225					230					235					240
Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys
				245					250					255	
Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val
				260					265					270	
Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln
				275					280					285	
Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser
				290				295						300	
Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His
305					310					315					320
Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys
				325					330					335	

<210> SEQ ID NO 58

<211> LENGTH: 307

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

Asp	Glu	Val	Ala	Ile	Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val	Leu	Ser
1				5					10					15	
Thr	Asn	Met	Lys	His	Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala	Pro	Gly
				20				25					30		
Glu	Thr	Val	Tyr	Tyr	Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu
				35			40					45			
Tyr	Thr	Ser	His	Ile	Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu
				50			55				60				
Gly	Pro	Glu	Cys	Asp	Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr
65					70					75				80	

-continued

```

Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser
      85                      90                      95
Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro
      100                      105                      110
Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu
      115                      120                      125
Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu
      130                      135                      140
Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro
      145                      150                      155                      160
Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala
      165                      170                      175
Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr
      180                      185                      190
Glu Cys Val Glu Val Gln Gly Glu Ala Thr Val Ala Ala Pro Ser Val
      195                      200                      205
Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser
      210                      215                      220
Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln
      225                      230                      235                      240
Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val
      245                      250                      255
Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu
      260                      265                      270
Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu
      275                      280                      285
Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg
      290                      295                      300
Gly Glu Cys
305

```

<210> SEQ ID NO 59

<211> LENGTH: 201

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 59

```

Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser
  1                      5                      10                      15
Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly
      20                      25                      30
Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu
      35                      40                      45
Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu
      50                      55                      60
Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr
      65                      70                      75                      80
Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser
      85                      90                      95
Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro
      100                      105                      110
Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu
      115                      120                      125

```

-continued

Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu
 130 135 140
 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro
 145 150 155 160
 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala
 165 170 175
 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr
 180 185 190
 Glu Cys Val Glu Val Gln Gly Glu Ala
 195 200

<210> SEQ ID NO 60
 <211> LENGTH: 323
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser
 1 5 10 15
 Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly
 20 25 30
 Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu
 35 40 45
 Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu
 50 55 60
 Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr
 65 70 75 80
 Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser
 85 90 95
 Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro
 100 105 110
 Gly Met Glu Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu
 115 120 125
 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu
 130 135 140
 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro
 145 150 155 160
 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala
 165 170 175
 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr
 180 185 190
 Glu Cys Val Glu Val Gln Gly Glu Ala Gly Gly Gly Ser Gly Gly
 195 200 205
 Gly Gly Ser Gly Gly Gly Gly Ser Arg Thr Val Ala Ala Pro Ser Val
 210 215 220
 Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser
 225 230 235 240
 Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln
 245 250 255
 Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val
 260 265 270
 Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu

-continued

275	280	285
Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu		
290	295	300
Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg		
305	310	315 320
Gly Glu Cys		

<210> SEQ ID NO 61
 <211> LENGTH: 201
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser		
1	5	10 15
Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly		
20	25	30
Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu		
35	40	45
Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu		
50	55	60
Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr		
65	70	75 80
Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser		
85	90	95
Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro		
100	105	110
Gly Met Glu Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu		
115	120	125
Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu		
130	135	140
Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro		
145	150	155 160
Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala		
165	170	175
Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr		
180	185	190
Glu Cys Val Glu Val Gln Gly Glu Ala		
195	200	

<210> SEQ ID NO 62
 <211> LENGTH: 559
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe		
1	5	10 15
Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly		
20	25	30
Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly		
35	40	45
Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr		
50	55	60

-continued

Tyr	Cys	Asp	Leu	Ser	Ala	Glu	Thr	Ser	Asp	Tyr	Glu	His	Gln	Tyr	Tyr
65					70					75				80	
Ala	Lys	Val	Lys	Ala	Ile	Trp	Gly	Thr	Lys	Cys	Ser	Lys	Trp	Ala	Glu
			85						90					95	
Ser	Gly	Arg	Phe	Tyr	Pro	Phe	Leu	Glu	Thr	Gln	Ile	Gly	Pro	Pro	Glu
		100					105						110		
Val	Ala	Leu	Thr	Thr	Asp	Glu	Lys	Ser	Ile	Ser	Val	Val	Leu	Thr	Ala
		115				120						125			
Pro	Glu	Lys	Trp	Lys	Arg	Asn	Pro	Glu	Asp	Leu	Pro	Val	Ser	Met	Gln
	130					135					140				
Gln	Ile	Tyr	Ser	Asn	Leu	Lys	Tyr	Asn	Val	Ser	Val	Leu	Asn	Thr	Lys
145					150					155					160
Ser	Asn	Arg	Thr	Trp	Ser	Gln	Cys	Val	Thr	Asn	His	Thr	Leu	Val	Leu
			165						170					175	
Thr	Trp	Leu	Glu	Pro	Asn	Thr	Leu	Tyr	Cys	Val	His	Val	Glu	Ser	Phe
		180					185						190		
Val	Pro	Gly	Pro	Pro	Arg	Arg	Ala	Gln	Pro	Ser	Glu	Lys	Gln	Cys	Ala
		195					200					205			
Arg	Thr	Leu	Lys	Asp	Gln	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	
	210					215				220					
Gly	Gly	Gly	Gly	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
225					230					235					240
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
				245					250					255	
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
		260						265					270		
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
		275					280					285			
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
	290					295					300				
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
305					310					315					320
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
			325						330					335	
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Ala	Glu	Gly	Ala	Pro	Ser	Val
		340						345					350		
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
		355					360					365			
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
	370					375					380				
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
385					390				395						400
Thr	Lys	Pro	Arg	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	
			405					410					415		
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
		420						425					430		
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile
		435					440					445			
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
	450					455					460				

-continued

```

Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
465                470                475                480

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
                485                490                495

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
                500                505                510

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
                515                520                525

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
                530                535                540

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
545                550                555

```

```

<210> SEQ ID NO 63
<211> LENGTH: 214
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 63

```

```

Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe
  1                5                10                15

Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly
                20                25                30

Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly
                35                40                45

Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr
  50                55                60

Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr
  65                70                75                80

Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu
                85                90                95

Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu
                100                105                110

Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala
                115                120                125

Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln
                130                135                140

Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys
                145                150                155                160

Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu
                165                170                175

Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe
                180                185                190

Val Pro Gly Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala
                195                200                205

Arg Thr Leu Lys Asp Gln
                210

```

```

<210> SEQ ID NO 64
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 64

```

-continued

Glu Glu Ile His Ala Glu Leu Arg Arg Phe Arg Arg Val Pro Cys Val
1 5 10 15

Ser Gly Gly

<210> SEQ ID NO 65

<211> LENGTH: 207

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn
1 5 10 15

Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly Val Lys Val Thr
20 25 30

Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn Lys
35 40 45

Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu
50 55 60

Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys Ala Ile Trp
65 70 75 80

Gly Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg Phe Tyr Pro Phe
85 90 95

Leu Glu Thr Gln Ile Gly Pro Pro Glu Val Ala Leu Thr Thr Asp Glu
100 105 110

Lys Ser Ile Ser Val Val Leu Thr Ala Pro Glu Lys Trp Lys Arg Asn
115 120 125

Pro Glu Asp Leu Pro Val Ser Met Gln Gln Ile Tyr Ser Asn Leu Lys
130 135 140

Tyr Asn Val Ser Val Leu Asn Thr Lys Ser Asn Arg Thr Trp Ser Gln
145 150 155 160

Cys Val Thr Asn His Thr Leu Val Leu Thr Trp Leu Glu Pro Asn Thr
165 170 175

Leu Tyr Cys Val His Val Glu Ser Phe Val Pro Gly Pro Pro Arg Arg
180 185 190

Ala Gln Pro Ser Glu Lys Gln Cys Ala Arg Thr Leu Lys Asp Gln
195 200 205

<210> SEQ ID NO 66

<211> LENGTH: 150

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu Thr Ser
1 5 10 15

Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys Ala Ile Trp Gly Thr
20 25 30

Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg Phe Tyr Pro Phe Leu Glu
35 40 45

Thr Gln Ile Gly Pro Pro Glu Val Ala Leu Thr Thr Asp Glu Lys Ser
50 55 60

Ile Ser Val Val Leu Thr Ala Pro Glu Lys Trp Lys Arg Asn Pro Glu
65 70 75 80

Asp Leu Pro Val Ser Met Gln Gln Ile Tyr Ser Asn Leu Lys Tyr Asn

-continued

85					90					95					
Val	Ser	Val	Leu	Asn	Thr	Lys	Ser	Asn	Arg	Thr	Trp	Ser	Gln	Cys	Val
			100					105					110		
Thr	Asn	His	Thr	Leu	Val	Leu	Thr	Trp	Leu	Glu	Pro	Asn	Thr	Leu	Tyr
		115					120					125			
Cys	Val	His	Val	Glu	Ser	Phe	Val	Pro	Gly	Pro	Pro	Arg	Arg	Ala	Gln
	130					135						140			
Pro	Ser	Glu	Lys	Gln	Cys										
145				150											

<210> SEQ ID NO 67

<211> LENGTH: 196

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 67

Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val	Leu	Ser	Thr	Asn	Met	Lys	His
1				5					10					15	
Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala	Pro	Gly	Glu	Thr	Val	Tyr	Tyr
		20					25						30		
Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu	Tyr	Thr	Ser	His	Ile
	35					40						45			
Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu	Gly	Pro	Glu	Cys	Asp
50					55					60					
Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr	Asn	Leu	Arg	Val	Arg
65				70					75					80	
Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala	Trp	Ser	Ile	Leu	Lys	His	Pro
			85					90						95	
Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr	Arg	Pro	Gly	Met	Glu	Ile	Thr
	100						105						110		
Lys	Asp	Gly	Phe	His	Leu	Val	Ile	Glu	Leu	Glu	Asp	Leu	Gly	Pro	Gln
	115				120						125				
Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Arg	Arg	Glu	Pro	Gly	Ala	Glu	Glu
130					135					140					
His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly	Ile	Pro	Val	His	Leu	Glu	Thr
145				150					155					160	
Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val	Lys	Ala	Gln	Thr	Phe	Val	Lys
			165					170					175		
Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser	Gln	Thr	Glu	Cys	Val	Glu	Val
	180						185						190		
Gln	Gly	Glu	Ala												
	195														

<210> SEQ ID NO 68

<211> LENGTH: 203

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

Asp	Glu	Val	Ala	Ile	Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val	Leu	Ser
1				5					10					15	
Thr	Asn	Met	Lys	His	Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala	Pro	Gly
	20						25						30		
Glu	Thr	Val	Tyr	Tyr	Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu

-continued

35					40					45					
Tyr	Thr	Ser	His	Ile	Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu
50					55					60					
Gly	Pro	Glu	Cys	Asp	Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr
65					70					75					80
Asn	Leu	Arg	Val	Arg	Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala	Trp	Ser
			85						90					95	
Ile	Leu	Lys	His	Pro	Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr	Arg	Pro
			100					105					110		
Gly	Met	Glu	Ile	Pro	Lys	His	Gly	Phe	His	Leu	Val	Ile	Glu	Leu	Glu
			115				120					125			
Asp	Leu	Gly	Pro	Gln	Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Thr	Arg	Glu
			130				135					140			
Pro	Gly	Ala	Glu	Glu	His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly	Ile	Pro
					150					155					160
Val	His	Leu	Glu	Thr	Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val	Lys	Ala
				165					170					175	
Gln	Thr	Phe	Val	Lys	Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser	Gln	Thr
			180					185					190		
Glu	Cys	Val	Glu	Val	Gln	Gly	Glu	Ala	Ile	Pro					
			195				200								

<210> SEQ ID NO 69

<211> LENGTH: 196

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

Leu	Pro	Ala	Pro	Gln	Asn	Leu	Ser	Val	Leu	Ser	Thr	Asn	Met	Lys	His
1				5					10					15	
Leu	Leu	Met	Trp	Ser	Pro	Val	Ile	Ala	Pro	Gly	Glu	Thr	Val	Tyr	Tyr
			20					25					30		
Ser	Val	Glu	Tyr	Gln	Gly	Glu	Tyr	Glu	Ser	Leu	Tyr	Thr	Ser	His	Ile
			35					40					45		
Trp	Ile	Pro	Ser	Ser	Trp	Cys	Ser	Leu	Thr	Glu	Gly	Pro	Glu	Cys	Asp
			50				55					60			
Val	Thr	Asp	Asp	Ile	Thr	Ala	Thr	Val	Pro	Tyr	Asn	Leu	Arg	Val	Arg
				70					75					80	
Ala	Thr	Leu	Gly	Ser	Gln	Thr	Ser	Ala	Trp	Ser	Ile	Leu	Lys	His	Pro
				85					90					95	
Phe	Asn	Arg	Asn	Ser	Thr	Ile	Leu	Thr	Arg	Pro	Gly	Met	Glu	Ile	Pro
			100					105					110		
Lys	His	Gly	Phe	His	Leu	Val	Ile	Glu	Leu	Glu	Asp	Leu	Gly	Pro	Gln
			115				120					125			
Phe	Glu	Phe	Leu	Val	Ala	Tyr	Trp	Thr	Arg	Glu	Pro	Gly	Ala	Glu	Glu
			130				135					140			
His	Val	Lys	Met	Val	Arg	Ser	Gly	Gly	Ile	Pro	Val	His	Leu	Glu	Thr
				150					155					160	
Met	Glu	Pro	Gly	Ala	Ala	Tyr	Cys	Val	Lys	Ala	Gln	Thr	Phe	Val	Lys
				165					170					175	
Ala	Ile	Gly	Arg	Tyr	Ser	Ala	Phe	Ser	Gln	Thr	Glu	Cys	Val	Glu	Val
			180					185					190		

-continued

Gln Gly Glu Ala
195

<210> SEQ ID NO 70
<211> LENGTH: 135
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr
1 5 10 15
Ala Thr Val Pro Tyr Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln
20 25 30
Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr
35 40 45
Ile Leu Thr Arg Pro Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu
50 55 60
Val Ile Glu Leu Glu Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala
65 70 75 80
Tyr Trp Arg Arg Glu Pro Gly Ala Glu Glu His Val Lys Met Val Arg
85 90 95
Ser Gly Gly Ile Pro Val His Leu Glu Thr Met Glu Pro Gly Ala Ala
100 105 110
Tyr Cys Val Lys Ala Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser
115 120 125
Ala Phe Ser Gln Thr Glu Cys
130 135

<210> SEQ ID NO 71
<211> LENGTH: 135
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr
1 5 10 15
Ala Thr Val Pro Tyr Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln
20 25 30
Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr
35 40 45
Ile Leu Thr Arg Pro Gly Met Glu Ile Pro Lys His Gly Phe His Leu
50 55 60
Val Ile Glu Leu Glu Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala
65 70 75 80
Tyr Trp Thr Arg Glu Pro Gly Ala Glu Glu His Val Lys Met Val Arg
85 90 95
Ser Gly Gly Ile Pro Val His Leu Glu Thr Met Glu Pro Gly Ala Ala
100 105 110
Tyr Cys Val Lys Ala Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser
115 120 125
Ala Phe Ser Gln Thr Glu Cys
130 135

<210> SEQ ID NO 72
<211> LENGTH: 15

-continued

<212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 72

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 1 5 10 15

What is claimed is:

1. A method of suppressing or reducing IL-20 induced inflammation in a mammal comprising administering to the mammal an effective amount of a composition comprising an antibody, antibody fragment, or single chain antibody and a pharmaceutically acceptable carrier, wherein the antibody, antibody fragment, or single chain antibody antagonizes IL-20, and wherein the antibody, antibody fragment, or single chain antibody suppresses or reduces inflammation in inflammatory skin disease selected from the group consisting of psoriasis, eczema, atopic dermatitis, and contact dermatitis.

2. The method according to claim 1, wherein the inflammatory disorder is a defensive response to mechanical trauma, toxins, neoplasia, abnormal recognition of host tissue as foreign, or invasion of the host by foreign material.

3. The method according to claim 1, wherein the inflammatory disorder is an angiogenesis dependent chronic inflammatory condition.

4. The method according to claim 1, wherein the inflammatory skin disease is a type of psoriasis selected from the group consisting of plaque type psoriasis, inverse psoriasis, eruptive psoriasis, and psoriasis with fingernail involvement.

5. The method according to claim 1, wherein the composition is administered in combination with another therapeutic agent.

6. The method according to claim 12, wherein the therapeutic agent is selected from the group consisting of a lubricant, keratolytic, topical corticosteroid, topical vitamin D derivative, anthralin, systemic antimetabolite, methotrexate, psoralen-ultraviolet-light therapy (PUVA), etretinate, isotretinoin, cyclosporine, and a calcipotriol.

7. A method of suppressing or reducing IL-20 induced inflammation in a mammal comprising administering to the mammal an effective amount of a composition comprising an antibody, antibody fragment, or single chain antibody and a pharmaceutically acceptable carrier, wherein the antibody, antibody fragment, or single chain antibody antagonizes IL-20, and wherein the antibody, antibody fragment, or single chain antibody suppresses or reduces inflammation in an inflammatory lung disease selected from the group consisting of bronchitis, lung cancer, bacterial pneumonia, and idiopathic pulmonary fibrosis.

8. The method according to claim 7, wherein the inflammatory disorder is a defensive response to mechanical trauma, toxins, neoplasia, abnormal recognition of host tissue as foreign, or invasion of the host by foreign material.

9. The method according to claim 7, wherein the inflammatory disorder is an angiogenesis dependent chronic inflammatory condition.

10. The method according to claim 7, wherein the composition is administered in combination with another therapeutic agent.

11. The method according to claim 10, wherein the therapeutic agent is selected from the group consisting of a lubricant, keratolytic, topical corticosteroid, topical vitamin D derivative, anthralin, systemic antimetabolite, methotrexate, psoralen-ultraviolet-light therapy (PUVA), etretinate, isotretinoin, cyclosporine, and a calcipotriol.

12. A method of suppressing or reducing IL-20 induced inflammation in a mammal comprising administering to the mammal an effective amount of a composition comprising an antibody, antibody fragment, or single chain antibody and a pharmaceutically acceptable carrier, wherein the antibody, antibody fragment, or single chain antibody antagonizes IL-20, and wherein the antibody, antibody fragment, or single chain antibody suppresses or reduces inflammation in an inflammatory bowel disease selected from the group consisting of ulcerative colitis and Crohn's disease.

13. The method according to claim 12, wherein the inflammatory disorder is a defensive response to mechanical trauma, toxins, neoplasia, abnormal recognition of host tissue as foreign, or invasion of the host by foreign material.

14. The method according to claim 12, wherein the composition is administered in combination with another therapeutic agent.

15. The method according to claim 14, wherein the therapeutic agent is selected from the group consisting of a lubricant, keratolytic, topical corticosteroid, topical vitamin D derivative, anthralin, systemic antimetabolite, methotrexate, psoralen-ultraviolet-light therapy (PUVA), etretinate, isotretinoin, cyclosporine, and a calcipotriol.

* * * * *