



(22) Date de dépôt/Filing Date: 2010/05/11
(41) Mise à la disp. pub./Open to Public Insp.: 2010/11/18
(45) Date de délivrance/Issue Date: 2018/09/11
(62) Demande originale/Original Application: 2 761 627
(30) Priorité/Priority: 2009/05/14 (IT PD2009A 000135)

(51) Cl.Int./Int.Cl. *A61K 31/737* (2006.01),
A61P 17/00 (2006.01), *A61P 19/10* (2006.01),
A61P 31/12 (2006.01), *A61P 9/00* (2006.01)

(72) Inventeurs/Inventors:
GALESSO, DEVIS, IT;
ZANELATO, ANNA MARIA, IT

(73) Propriétaire/Owner:
FIDIA FARMACEUTICI S.P.A., IT

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : ACIDES HYALURONIQUES SULFATES COMME AGENTS DE REGULATION DE L'ACTIVITE CYTOKINIQUE
(54) Title: SULPHATED HYALURONIC ACIDS AS REGULATOR AGENTS OF THE CYTOKINE ACTIVITY

(57) **Abrégé/Abstract:**

The present invention relates to the new and surprising use of sulfated hyaluronic acid (HAS) as regulator agent of the cytokine activity (pro- and anti-inflammatory) and consequently the use of HAS for the preparation of a new medicine for the prevention and treatment of pathologies associated with the activation and/or deficiency of cytokines of a pro- and anti-inflammatory nature. The Applicant has in fact discovered its exclusive capacity in modulating the activity of these particular proteins, it has studied the action mechanism and demonstrated the substantial difference between the different sulfated types known in the state of the art, but above all it has demonstrated an unexpectedly high activity of HAS vs different types and strains of Herpes virus, HIV, Cytomegalovirus and the virus of vesicular stomatitis.

Abstract

The present invention relates to the new and surprising use of sulfated hyaluronic acid (HAS) as regulator agent of the cytokine activity (pro- and anti-inflammatory) and consequently the use of HAS for the preparation of a new medicine for the prevention and treatment of pathologies associated with the activation and/or deficiency of cytokines of a pro- and anti-inflammatory nature. The Applicant has in fact discovered its exclusive capacity in modulating the activity of these particular proteins, it has studied the action mechanism and demonstrated the substantial difference between the different sulfated types known in the state of the art, but above all it has demonstrated an unexpectedly high activity of HAS vs different types and strains of Herpes virus, HIV, Cytomegalovirus and the virus of vesicular stomatitis.

SULPHATED HYALURONIC ACIDS AS REGULATOR AGENTS OF THE CYTOKINE ACTIVITY

This application has been divided out of Canadian Patent
5 Application Serial No. 2,761,627 filed internationally
on May 11, 2010 as PCT/EP2010/003044 and published
internationally on November 18, 2010 as WO 2010/130466.

Field of the invention

10 For many years now, scientific/patent literature has
been studying and describing sulfated hyaluronic acid
which is obtained starting from hyaluronic acid (HA)
suitably sulfated according to what is described in the
state of the art (EP0940410B1 and EP0702699B1), to which
15 anticoagulant effects are attributed. HAS can also be
obtained by the de-acetylation and subsequent sulfation of
glucosamine of HA (defined as HA-NS) (EP0971961B1), for
the production of surgical articles and pharmaceutical
compositions. Patents EP0754460B1 and EP1385492B1 are also
20 known, in which the use of HAS is described in pathologies
such as, for example, ARDS, articular rheumatism,
rheumatoid arthritis and dermatitis.

An object of the present invention relates to the
new and surprising use of HAS as regulator agent of the
25 cytokine activity, as the Applicant has discovered the

exclusive capacity of HAS of modulating the activity of particular cytokines (both pro- and anti-inflammatory), it has studied its action mechanism and revealed the substantial difference between the two types of sulfated
5 product (HAS and HA-NS), but above all the Applicant has surprisingly found an unexpectedly high activity vs different types and strains of Herpes virus, HIV, Cytomega-

lovirus and the virus of vesicular stomatitis.

Since 1970, scientists have understood that selected populations of lymphoid cells can produce and release into the circulatory bed, molecules of a protein nature not
5 assimilable to antibodies, defined with the term "cytokines". They represent a new type of "hormone", capable of acting on different cell targets in numerous regions of the body.

The progression of scientific knowledge relating to
10 the synthesis and biological/biochemical functions of these proteins, has altered the "old" vision of the immune system (I.S.) of the same scientific world and has opened up new horizons in understanding its numerous functions, thus creating new prospects for the treatment
15 of different pathologies, topic and/or systemic, also comprising new therapeutic possibilities relating to the immunotherapy of cancer.

The central IS cell is the lymphocyte, it represents about 20% of all the white blood cells and, on the basis
20 of its various functions, forms 3 groups: lymphocyte B, lymphocyte T and killer lymphocyte. Many cytokines are soluble proteins produced by lymphocytes and/or monocytes, capable of acting against other cells/tissues also situated very far from their production site. They have
25 immunological functions, in fact, and also regulation

functions in the synthesis of other cytokines by different cells of the I.S. or target cells involved in the cascade of reactions initiated by the I.S..

Numerous different cytokines have been studied so far, also having numerous different acronyms, but those studied in particular by the Applicant are: Interleukin 1 and 2, Interleukin 6, 7 and 12, hereafter defined as IL-1, IL-2, IL-6, IL-7 and IL-12 which, with TNF, are defined as cytokines of a pro-inflammatory nature, whereas Interleukin 10 (IL-10) on the contrary, is a cytokine with strong anti-inflammatory properties.

The first cytokine to be studied was definitely IL-1 present in two forms α and β , a powerful inducer of inflammatory processes, both local and systemic. It is mainly produced by lymphocytes B, T and macrophages after bacterial stimulus or stimulation on the part of other agents including other cytokines; it is also secreted from alveolar, peritoneal macrophages, Kupffer cells, peripheral neutrophils, endothelial, epithelial and smooth muscle cells, fibroblasts, Langerhans cells of the skin, osteoclasts, synoviocytes and many other types of cell. It is also present in the cerebrospinal fluid where it is both produced and transported. Both forms bind the same receptor and have very similar, if not identical, biological activities. Many of their pro-inflammatory func-

tions are linked to the stimulation of other cytokines, such as IL-6 and IL-8, and their very synthesis can be induced by cytokines such as TNF, Interferon, bacterial endotoxins, viruses and different types of other anti-
5 gens. It is involved in septic shock but it should also be noted that recent studies have demonstrated that IL-1 is capable of activating the expression of some oncogenes and consequently of participating in the pathogenesis of neoplasias. Furthermore, an autocrin control system of
10 the growth of blast cells of the white series which are present in the circulatory bed of patients with the leukaemia pathology has been suggested for this cytokine: these blast cells in fact uncontrollably produce IL-1 which in turn stimulates the synthesis of those growth
15 factors which increase the proliferation of the same blast cells. Combined with other cytokines, IL-1 therefore represents one of the major mediators of inflammatory processes: it stimulates T-cells in fact to produce IL-2 and B-cells to produce immunoglobulins. It is consequently involved in numerous pathologies such as, for ex-
20 ample, astrogliosis and the demyelination of neuronal fibres, it is cytotoxic for Langerhans pancreatic cells which produce insulin, and is also involved in lithic processes of the bones, both activating osteoclasts and
25 suppressing the formation of new bone, both processes be-

ing involved in the pathology of osteoporosis. It is capable of functioning as an endogenous pyrogen as, by increasing the release of prostaglandins in the hypothalamic centre, it causes an increase in the body temperature.

5 It is also involved in the pathogenesis of rheumatoid arthritis and arthrosis: high quantities of IL-1 have in fact been found in the synovial fluid of patients affected by rheumatoid arthritis and/or osteoarthritis. Finally, it participates in the establishment of vascular damage such as venous thrombosis and is present in all vessels with pathologies of the arterio/arteriosclerotic type. Receptor antagonists are currently being experimented for this cytokine, as blockage of the receptor is proving to be an effective way of treating these pathologies in which IL-1 is among the protagonists.

10

15

TNF: The necrosis factor is part of the group of cytokines which promotes the reaction of acute systemic inflammation phase. TNF is therefore involved in an extremely wide number of processes such as cell proliferation, differentiation and apoptosis, carcinogenesis and viral replication.

20

It is mostly produced by macrophages and by a series of other cell types including mastocytes, lymphoid cells, muscular and endothelial cells, fibroblasts and nerve cells. Its synthesis can be stimulated by bacterial en-

25

dotoxins, other cytokines such as IL-2, Interferon and IL-1, and it can be inhibited by steroids.

By acting on numerous organs and systems, generally together with other cytokines, it participates in the establishment and regulation of many pathogenetic processes:

- it modulates the expression of many proteins and important cytokines, such as IL-1 and IL-6 thus resulting involved in many cutaneous pathologies such as, dermatitis, vitiligo and eczema;
- it stimulates the hypothalamo-hypophyseal-adrenal axis increasing the release of some types of hormones;
- it suppresses the appetite;
- it induces fever (it acts as endogenous pyrogen inducing the hypothalamus to produce prostaglandins);
- it stimulates the synthesis of collagenase in the synoviocytes and for this reason, large quantities of TNF have been found in the synovial fluids of patients suffering from arthrosis and rheumatoid arthritis;
- it activates the osteoclasts and therefore induces re-absorption of the bone, process char-

acterizing the osteoporosis pathology;

- it is also involved in pathologies of the Nervous System, such as astrocytosis and demyelination;
- 5 - it strongly attracts neutrophils and helps them attach themselves to the endothelial cells to extravasate;
- it stimulates the macrophagic production of molecules with an oxidizing action;
- 10 - it is involved in particular pathologies of the Cardio-circulatory System participating in the formation of venous thrombosis, in the pathogenesis of arteriosclerosis and vasculitis;
- it increases the resistance to insulin, it increases the protein catabolism in the muscles
15 whereas it suppresses the lipogenic metabolism in the adipose tissue.

High concentrations of TNF can induce shock-like symptoms, whereas prolonged exposure to low concentrations can lead to cachexia, a syndrome which causes the
20 depletion of the protein and lipid patrimony of the tissues (in particular muscular and adipose). The TNF is capable of binding itself to two receptors, TNF-R1 (receptor for type 1 TNF) and TNF-R2 (receptor for type 2 TNF),
25 which are expressed in all somatic cells excluding eryth-

rocytes. In short, the TNF promotes the inflammatory response which in turn triggers numerous pathologies also of an auto-immune nature, such as rheumatoid arthritis, Crohn disease, psoriasis and asthma. Scientific research
5 has so far tried to perfect "biological" drugs (such as, for example, monoclonal antibodies) which inhibit the synthesis of TNF and/or block its receptor.

IL-2: this is a highly pro-inflammatory, atherogenic cytokine, mainly produced by lymphocytes T, whose synthesis
10 is inhibited by steroids and cyclosporines. Leukemic cells synthesize the above cytokine and contemporaneously express its receptor, thus creating an autocrin system in the stimulation of their growth which causes a deterioration in the leukemia pathology. IL-2 has a central role
15 in regulating the immunological response: it stimulates in fact the synthesis of IFN in the peripheral leukocytes and induces the production of IL-1 and TNF. IL-2 can also damage the hematoencephalic barrier and integrity of the endothelium of the cerebral vessels, causing neuropsychi-
20 atric disorders such as disorientation and depression.

There are consequently numerous pathologies which have been associated with an aberrant production of IL-2, such as Hodgkin's lymphoma, the rejection of organ transplants, multiple sc(h)lerosis, rheumatoid arthritis, Lu-
25 pus erythematosus, diabetes and AIDS.

IL-6: produced by many cell types above all of I.S., with
TNF it is amongst the most important members of the group
of chemical mediators of the acute phase of the inflamma-
tory process, and is therefore involved in pathologies
5 with a strong inflammatory component, such as asthma
(where it participates in the emergence and maintenance
of the inflammatory process), chronic intestinal inflam-
mation (Crohn's disease), rheumatoid arthritis and
arthrosis. As previously affirmed, in fact, cytokines
10 such as TNF, IL-1 and IL-6 have proved to be greatly in-
volved in the degenerative articular osteoarthrosis pro-
cess as they have a primary role in regulating the ex-
pression of metalloproteases (responsible for cartilage
degradation), in the production of prostaglandins and in
15 osteoclastic activation and, for this reason, high cytokine
levels have been registered in the synovial fluids of pa-
tients suffering from arthrosis and rheumatoid arthritis
(R.A.). These discoveries have stimulated the use of inhibi-
tors in the above interleukins and/or receptor antagonists as
20 a new treatment strategy of the arthrosis pathology.

High concentrations of IL-6 have also been found in
the urine of patients subjected to transplants and its
presence represents one of the first signs of the rejec-
tion reaction of the organ. The serum level of this cyto-
25 kine is also drastically increased in many patients suf-

fering from tumors (such as, for example, myeloma, leukemia, cardiac myxomas, or in pathologies such as lymphadenopathies and hepatic cirrhosis) and can also be used as indicator in monitoring the size of the tumoral mass. Finally, recent studies have connected cancer with longevity and revealed how some tumors are influenced by the quali/quantitative situation of the cytokine proteins of the patient: in short, recent evidence has linked a low production profile of IL-10 and high secretion of IL-6 to a deterioration in the clinical survival of patients affected by tumoral pathologies, whereas a genotype capable of producing and maintaining high levels of IL-10 can facilitate survival. Consequently, persons with high anti-inflammatory cytokine levels and a low concentration of pro-inflammatory cytokines are genetically susceptible to having a greater longevity (Caruso C. et al., Ann N.Y. Acad. SCI., 2004, 1028:1-13).

IL-7: cytokine mainly produced by stromal cells of bone marrow, it is also secreted by the thymus and keratinocytes. IL-7 induces the synthesis of inflammatory cytokines such as IL-1, IL-6 and TNF, thus participating in the pathogenesis of some skin diseases (such as psoriasis and cutaneous lymphoma) and the osteoarticular system. High levels of IL-7 have in fact been found in patients suffering from R.A. as IL-1 and TNF (cytokines strongly

involved in the above pathology) can increase the stromal production of IL-7 which in turn stimulates the macrophagic synthesis of TNF. Finally, IL-7 can induce the maturation of osteoclasts and consequently increase bone resorption thus contributing to the degeneration of the joints.

IL-12: this protein also plays a central role in regulating the functions of the I.S. It acts in fact on the differentiation of the lymphocytes, it induces the synthesis of Interferon and TNF, and its production can be inhibited by IL-10. The over-production of this protein enters in the pathogenesis of diseases of (an) auto-immune nature such as colitis, arthritis, insulin-dependent diabetes, encephalomyelitis, psoriasis and multiple sclerosis (Brahmachari S. et al., Minerva Med., 2008, 99(2):105-118).

IL-10: mainly produced by lymphocytes, it is a cytokine of an anti-inflammatory nature, capable of inhibiting the synthesis of IL-2 and Interferon produced by lymphocytes T. The anti-inflammatory action of IL-10 is also revealed in the capacity of inhibiting the synthesis of IL-1, IL-6, IL-8, IL-12 and TNF in the macrophages stimulated with bacterial endotoxins. IL-10 deficiencies are associated with pathologies such as diabetes mellitus and chronic intestinal inflammations, such as Crohn's disease. Recent

evidence has led IL-10 to also be experimented as a new therapeutic approach for the treatment of Lupus erythematosus. Low IL-10 levels have been observed in cutaneous tissues of patients suffering of pathologies such as vitiligo, psoriasis, eczema and dermatitis in general. It should be noted that both corticosteroids and cyclosporine increase the production and/or release of this interleukin from the relative competent cells during conventional immunosuppression therapy for the treatment of inflammations and organ rejection (Zhou X. Et al., Current Drug Targets-Immune, Endocrine & Metabolic Disorders, 2005, 5(465475). Experimental data have also demonstrated its effectiveness in reducing the release of prostaglandins and cyclo-oxygenase induced *in vitro* by TNF on human synoviocytes, thus indicating the capacity of IL-10 of reducing inflammatory processes which involve articulations affected by osteoarthrosic degeneration (Alaaeddine N. et al., Arthritis & Rheumatism, 1999, 42:710-718). Recent studies have confirmed its therapeutic effectiveness towards the asthma pathology in experimental animal models of bronchial hyper-reactivity, showing how this cytokine has a high therapeutic potentiality in reducing the inflammation which characterizes the air passages of asthmatic patients, in which high concentrations of TNF, IL-1, IL-5, IL-6 and IL-8 have been found

in the bronchial-washing liquid and/or on a serum level
and/or tissue level (Stankiewicz W. et al., Mediators of
Inflammation, 2002, 11:307-312). For this interleukin,
the important role of regulator cytokine of the mainte-
5 nance of immunological homeostasis, has therefore been
assumed.

Asthma can be an extremely invalidating disease of
which approximately 200 million people in the world suf-
fer, with over 5,000 deaths per year. It is a pathology
10 which is based on a distorted response of the I.S. to en-
vironmental factors, consequently linked to an exacerbat-
ed production of pro-inflammatory cytokines for the
growth and differentiation of mast cells and eosinophils
with other types of cells of the I.S.. The causes of this
15 out-of-balance activity of the immune system are still
not completely known, there are however genetic, environ-
mental, viral and also nutritional factors which contrib-
ute in different ways to the development of this patholo-
gy. Consequently, finding an effective therapy for its
20 prevention and/or treatment which allows the suspension
or reduction of the use of steroids (conventional treat-
ment therapy), could represent a valid solution for both
the more serious forms (as it would in any case enable a
reduction in the use of steroids) and for less serious
25 cases, as the suspension of the steroid therapy could be

total.

DETAILED DESCRIPTION OF THE INVENTION

An object of the invention is the new and surprising use of HAS as regulator agent of the cytokine activity, as the Applicant has discovered its exclusive capacity in modulating the activity of particular cytokines, it has studied its action mechanism and revealed the substantial difference between the various types of sulfated products known in the state of the art, but above all the Applicant has discovered an unexpected activity vs different types and strains of Herpes virus, HIV, Cytomegalovirus and the virus of vesicular stomatitis. The sulfated hyaluronic acid suitable for the purposes of the present invention is prepared according to the process described in EP 702699 B1: the sulfation is effected by means of the complex SO₃-pyridine and involves the alcohol hydroxyls present in the polysaccharide chain starting from a HA deriving from any source, for example, by extraction from cockscombs, either fermentatively or technologically, and having a molecular weight ranging from 400 to 3x10⁶Da, in particular from 1x 10⁴Da to 1x 10⁶Da, even more in particular from 10,000 to 50,000 Da, 150,000 to 250,000 Da and 500,000 to 750,000 Da.

The derivative obtained maintains all the physical characteristics of the starting polymer unaltered, in par-

particular the molecular weight of the starting HA is not altered by the sulfation process thus allowing all the physico-chemical characteristics of the starting polysaccharide to be maintained. The sulfation involves the various hydroxyl groups of the disaccharide unit and it is therefore possible to obtain different sulfation degrees, from 0.5 to 3.5 (intended as the number of sulfate groups per disaccharide unit), by varying the quantity of SO₃-pyridine introduced as known in the state of the art.

10 The derivative used in all the experimentations effected generally has sulfation degree 1 or degree 3 and is defined hereafter as HAS1 and HAS3. All the free carboxyl groups of the HA can be salified with cations of an organic and/or inorganic origin.

15 Both degrees of HAS are soluble in water and they can also be sterilized with the normal techniques known to experts in the field, even if sterilization using an autoclave is preferable.

20 In the experimentations described hereunder, the Applicant demonstrated that:

- HAS is capable of both stimulating the production of new mRNA and the protein synthesis of anti-inflammatory cytokines (such as, for example, IL-10), thus increasing the immune defense capacity of the cells and therefore of the whole organism. The anti-inflammatory ac-

tion of the above cytokines is revealed in the capacity of inhibiting the synthesis of IL-1, IL-6, IL-8, IL-12 and TNF, all proteins of a pro-inflammatory nature.

- HAS is effective in both diminishing the synthesis of
5 new mRNA and in significantly reducing the protein syn-
thesis of IL-2, IL-7 and IL-12 on the part of the synoviocytes (for IL-12) and cell components of the I.S., both in situations in which an immune response is not demanded, and also in particular inflammatory stress events
10 in which the immune cell responds by producing a cytokine cascade and, above all in this case, the data presented reveal the greater inhibitory effect of HAS.
- HAS is effective in inhibiting the binding of TNF, IL-1 and IL-6 to their receptor. These results are of funda-
15 mental importance as they prove that the behaviour of the sulfate is completely similar to that of specific monoclonal antibodies for the receptors of the above pro-inflammatory proteins, consequently capable of blocking their function even if they do not have this
20 specificity. This receptor blockage represents the most effective way of antagonizing the pro-inflammatory and tumoral effects of the TNF, IL-1 and IL-6 factor, thus opening up new horizons in clinical experimentation, allowing the perfectioning of new therapeutic ap-
25 proaches in the treatment and/or prevention of an ex-

tremely high number of pathologies, considering the role that TNF, IL-1 and IL-6 have in the onset and progression of numerous diseases.

The Applicant therefore describes and claims the new
5 use of HAS for the preparation of a medicine:

- for the prevention and/or treatment of pathologies associated with immune deficiency and, in particular, deficiency of IL-10, stimulating the synthesis of anti-inflammatory cytokines as new local and systemic therapy
10 py of pathologies such as vitiligo, eczema, psoriasis and dermatitis;
- for the prevention and/or treatment of pathologies associated with the increase/activation of IL-1, IL-2, IL-6, IL-7, IL-8, IL-12 and TNF;
- 15 • for the prevention and/or treatment of asthma, rheumatoid arthritis, and arthrosis associated with the activation of IL-1, IL-6 and TNF;
- for the prevention and/or treatment of pathologies associated with endothelial damage in blood vessels;
- 20 • for the prevention and/or treatment of skin diseases such as, for example, dermatitis and cutaneous lymphoma;
- for the prevention and/or treatment of diseases of an autoimmune nature such as rheumatoid arthritis, Crohn's
25 disease (and all chronic intestinal inflammations),

- psoriasis and asthma, diabetes mellitus, encephalomyelitis, Multiple Sclerosis, Lupus erythematosus and for the treatment of rejection in organ transplants;
- for the prevention and/or treatment of neoplasias such
5 as, for example, leukaemia and Hodgkin's lymphoma;
 - for the prevention and/or treatment of astrogliosis, astrocytosis and demyelination of the neuronal fibres;
 - for the prevention and/or treatment of pathologies associated with the activation of osteoclasts, such as
10 the osteoporosis;
 - for the prevention and/or treatment of vascular pathologies of the arterio/atherosclerotic type, venous thrombosis and vasculitis, associated with the activation of TNF, IL-1 and IL-6;
 - 15 • for the prevention and/or treatment of febrile pathologies.

In a particular embodiment there is provided sulphated hyaluronic acid for use in the prevention and/or treatment of herpes genitalis, said sulphated hyaluronic acid (HA) being obtained by a sulfation involving the alcohol hydroxyls present in the HA polysaccharide chain, starting from a HA having a molecular weight ranging from 10,000 to 50,000 Da, 150,000 to 250,000 Da and 500,000 to 750,000 Da, said sulphated hyaluronic acid having a sulphation degree equal to 1 or 3.

10 The Applicant has also demonstrated, in the experimentations described hereunder, the powerful antiviral action of HAS vs different types of viruses:

- the results obtained show the effectiveness of HAS (both degree 1 and degree 3) in inhibiting the viral replication of the HIV 1 and 2 virus. The experimentation has also demonstrated that not all sulfated hyaluronic acids known in the state of the art act effectively vs viral replication, as only HAS in which the

sulfation has taken place with respect to the hydroxyl groups alone has proved to be active, where HA-NS is not effective in blocking viral cytopathogenicity.

- Experimental data confirm the antiviral action of HAS1 and HAS3 vs Herpes Simplex virus 1 and 2 and vs the virus of vesicular stomatitis (VSV). The first form, extremely widespread, is responsible for the appearance of characteristic febrile vesicles which normally affect the facial cutis (lips, nostrils); it is also called herpes simplex labialis. The infection caused by lip herpes can easily reappear as the virus survives inside the cells and is not even eliminated with the use of effective drugs. The second form is a genital infection, also known as herpes genitalis. Both are caught by physical and sexual contact. Due to the location of the virions in the nervous ganglia, where they can remain quiescent for a long period of time, the herpetic infection has recurring characteristics in correspondence with stressing events of the immune system and usually reappears in the primary site. The virus of vesicular stomatitis is an RNA-virus, it strikes mammals (mainly animals) and is used in the laboratory for studying the development of the life-cycle of the RNA-virus. A comparison between HA-NS1 and HAS1 shows once again that not all sulfated hyaluronic acids are

equivalent, as HA-NS1 has proved not to be active at all, whereas both HAS 1 and 3 show a very strong antiviral activity vs Herpes Simplex and also vs VSV. None of the samples tested prove to be cytotoxic towards the host cell, the minimum cytotoxic concentration obtained, in fact, is equal to that of reference drugs normally used in clinical practice for the treatment of Herpes, and on an average has proved to be 100 times higher than that revealed active in the inhibition of viral replication.

- Experimental data obtained for both HAS1 and HAS3 have revealed a clear and significant antiviral result vs Cytomegalovirus: this is a particular type of virus which enters some types of cells of our organism in which it reproduces itself parasitically causing their death. It belongs to the same family as *herpes labialis* and *herpes genitalis*, chickenpox and infective mononucleosis. Epithelial cells, mucous membranes, lymphonodes are the site of multiple primary infection. It remains in latent form for life in the peripheral blood, in the epithelium of the renal tubules and in the epithelium of the salivary glands. Serious forms are found in immunocompromised subjects (such as those affected by AIDS and transplant subjects in immunosuppressive therapy) involving various organs: pneumonia,

hepatitis, colitis, esophagitis, nephritis. The treatment therapy consists in the administration of drugs such as ganciclovir, valganciclovir and foscarnet (inhibitors of the synthesis of viral DNA). Also in this
5 case, HA-NS1 has proved to be non-active in inhibiting the proliferation of the virus confirming the absolute diversity, as antiviral capacity, between the two types of sulfated products.

The Applicant therefore describes and claims the new
10 use of HAS for the preparation of a medicine:

- for the prevention and/or treatment of HIV;
- for the prevention and/or treatment of herpes simplex labialis and herpes genitalis;
- for the prevention and/or treatment of the virus of ve-
15 sicular stomatitis;
- for the prevention and/or treatment of Cytomegalovirus.

Finally, the Applicant describes the preparation of various pharmaceutical formulations/compositions containing HAS as sole active principle, or in association with
20 other pharmacologically and/or biologically active agents such as, for example, steroids, hormones, proteins, trophic factors, vitamins, non-steroid anti-inflammatory drugs (FANS), chemotherapy drugs, calcium-antagonists, antibiotics, antiviral agents, anticoagulants and/or
25 fibrinolytic agents, local anaesthetics, enzymes such as,

for example, collagenase and/or hyaluronidase and/or other proteases; it can be formulated with polymers such as hyaluronic acid and its derivatives, carboxymethylcellulose and/or other polymers of a natural
5 (such as collagen) or synthetic nature.

The pharmaceutical composition in question can be administered systemically (endovenously or arterially, intramuscularly, intraperitoneally, hypodermically or orally), it can be used for a topic application by dermal
10 and/or transdermal absorption, it can be administered by inhalation/aerosol (especially for the treatment of asthmatic pathologies), intra-articularly or it can be administered directly in the site to be treated by direct injection.

15 The pharmaceutical composition in question can be formulated as an ointment, lipogel, hydrogel, lipstick, cream, vaginal ovules and bougies, foam, mucosal gel, ophthalmic preparations, douches, mouthwash, patches for dermal and/or transdermal absorption, especially of FANS
20 and hormones, solution for inhalant use.

Some examples of the preparation of HAS degree 1 and 3, pharmaceutical formulations containing it, are provided for purely descriptive and non-limiting purposes, together with the results obtained by experimentation in
25 *vitro*.

Example 1

Preparation of the tetrabutylammonium salt of hyaluronic acid (HA) having an average molecular weight equal to 200KD (ranging from 150,000 to 250,000 Da)

5 5.00 g of hyaluronic acid of a fermentative sodium salt origin (200 KD) were dissolved in 250 ml of water and the resulting solution was percolated through a glass column pre-filled with 100 cm³ of Dowex resin in the form of tetrabutylammonium (TBA). The eluted solution of HA-
10 TBA salt is collected and freeze-dried. 7.50 g of product are obtained.

Example 2

Synthesis of sulfated HA starting from HA having an average molecular weight of 200 KD and a sulfation degree
15 equal to 3 sulfate groups per repetitive unit

Method A

10.0 g of the TBA salt of hyaluronic acid having an average molecular weight of 200 KD prepared according to Example 1, are dissolved in 300 ml of dimethylsulfoxide
20 (DMSO); 26.0 g of the complex SO₃-pyridine (sulphur trioxide and pyridine, hereafter abbreviated as PySO₃) are dispersed in 150 ml of DMSO, and then added to the solution of HA. After 20 hours under mechanical stirring at a temperature of 21°C, the reaction is interrupted adding
25 0.1 volumes of water; the raw reaction product is iso-

lated by precipitation after the addition of 2 volumes of ethanol. The solid obtained is dispersed in 150 ml of water and the pH brought to neutrality with NaOH 1 M. The mixture is exhaustively dialysed against water through a
5 membrane with a cut-off of 12-14,000 Da. The dialysed product is subjected to freeze-drying. 9.7 g of product are obtained with a sulfation degree equal to 3 sulfate groups per repetitive unit (yield = 88%).

Method B

10 32.0 g of the TBA salt of hyaluronic acid having an average molecular weight of 200 KD prepared according to Example 1, are dissolved in 900 ml of N-Methyl-Pyrrolidone (NMP); 100 g of PySO₃ are dispersed in 600 ml of NMP, and then added to the solution of HA. After 20
15 hours under mechanical stirring at a temperature of 21 ± 1°C, the reaction is interrupted adding 0.5 volumes of water; The pH initially lower than 2.5, is brought to neutrality with NaOH (in solution). The reaction raw product is isolated by precipitation by the addition of
20 2.5 volumes of methanol and washed with 2 volumes of a methanol/water mixture 8/2. The solid is re-dissolved and exhaustively dialysed against water through a membrane with a cut-off of 12-14,000 Da. 30.4 g of product are obtained with a sulfation degree equal to 3 sulfate groups
25 per repetitive unit (yield = 86%).

Example 3

Synthesis of sulfated HA starting from HA having an average molecular weight of 200 KD and a sulfation degree equal to 1 sulfate group per repetitive unit

5 Using the procedure illustrated in Example 1, 10.0 g of TBA salt of HA are prepared, which are dissolved in 350 ml of DMSO. 10.0 g of the complex PySO_3 are dispersed in 100 ml of DMSO, and then added to the solution of HA. After 20 hours under mechanical stirring at a temperature
10 of 21°C, the reaction is interrupted adding 0.1 volumes of water; the raw reaction product is isolated by precipitation after the addition of 2.5 volumes of ethanol. The solid obtained is dispersed in 150 ml of water and the pH brought to neutrality with NaOH 1 moles/l. The mixture is
15 exhaustively dialysed against water through a membrane with a cut-off of 12-14,000 Da. The dialysed product is subjected to freeze-drying. 7.54 g of product are obtained with a sulfation degree equal to 1.0 sulfate group per repetitive unit (yield = 93%).

20 Example 4

Synthesis of sulfated HA starting from HA having a low molecular weight (average MW of 10 KD, ranging from 10,000 to 50,000 Da) and a sulfation degree equal to 3 sulfate groups per repetitive unit

25 Using the procedure illustrated in Example 1, 12.4 g

of TBA salt of low-molecular-weight hyaluronic acid are prepared, which are dissolved in 300 ml of NMP. 40 g of PySO_3 are dispersed in 100 ml of NMP, and then added to the solution of HA. After 20 hours under mechanical stirring at a temperature of 21°C , the reaction is interrupted adding 0.5 volumes of water. The pH initially lower than 2.5, is brought to neutrality with NaOH 4M. The reaction raw product is isolated by precipitation by the addition of 2.5 volumes of methanol and washed with 2 volumes of a methanol/water mixture 8/2. The solid is redissolved and exhaustively dialyzed against water through a membrane with a cut-off of 3,500 Da. 12.0 g of product are obtained with a sulfation degree equal to 3.0 sulfate groups per repetitive unit (yield = 85%).

15 **Example 5**

Synthesis of sulfated HA starting from HA having a low average molecular weight and a sulfation degree equal to 1 sulfate group per repetitive unit

Using the procedure illustrated in Example 1, 12.4 g of TBA salt of HA are dissolved in 300 ml of DMSO. 16.0 g of PySO_3 are dispersed in 100 ml of DMSO and then added to the solution of HA. After 20 hours under mechanical stirring at a temperature of 21°C , the reaction is interrupted adding 0.1 volumes of water; the reaction raw product is isolated by precipitation by the addition of

2.5 volumes of ethanol. The solid obtained is dispersed in 150 ml of water and the pH brought to neutrality with NaOH 1 moles/l. The mixture is exhaustively dialyzed against water through a membrane with a cut-off of 3,500
5 Da. The dialyzed product is subjected to freeze-drying. 9.04 g of product are obtained with a sulfation degree equal to 1.0 sulfate group per repetitive unit (yield = 90%).

Example 6

10 Synthesis of sulfated HA starting from HA having a molecular weight within the range of 500-730 KD Da and a sulfation degree equal to 3 sulfate groups per repetitive unit

21.0 g of hyaluronic acid sodium salt of an extractive origin (500-730 KD) are dissolved in 1.5 l of water and the resulting solution is percolated through a glass column pre-filled with 450 cm³ of Dowex resin in the form of TBA. The eluted solution of HA-TBA salt is collected and freeze-dried. 32.0 g of product are obtained, which
15 are dissolved in 1.35 l of NMP; 100 g of PySO₃ are dispersed in 650 ml of NMP, and then added to the solution of HA. After 20 hours under mechanical stirring at a temperature of $23 \pm 1^{\circ}\text{C}$, the reaction is interrupted adding 0.5 volumes of water. The pH initially lower than 2.5, is
20 brought to neutrality by the addition of NaOH (in solu-
25

tion at a concentration of 4 moles/l). The reaction raw product is isolated by precipitation by the addition of 2.5 volumes of methanol and washed with 3.5 volumes of a methanol/water mixture 8/2. The solid is re-dissolved and
5 exhaustively dialyzed against water through a membrane with a cut-off of 12-14,000 Da. 30.3 g of product are obtained with a sulfation degree equal to 3 sulfate groups per repetitive unit (yield = 83%).

Example 7

10 Synthesis of sulfated HA starting from HA having a molecular weight within the range of 500-730 KD and a sulfation degree equal to 1 sulfate group per repetitive unit

21.0 g of hyaluronic acid sodium salt of an extractive origin (500-730 KD) are dissolved in 1.5 l of water and the resulting solution is percolated through a glass column pre-filled with 450 cm³ of Dowex resin in the form of TBA. The eluted solution of HA-TBA salt is collected and freeze-dried. 32.0 g of product are obtained, which
15 are dissolved in 1.65 l of NMP; 40 g of PySO₃ are dispersed in 350 ml of NMP, and then added to the solution of HA. After 20 hours under mechanical stirring at a temperature of 25 ± 1°C, the reaction is interrupted adding 0.5 volumes of water. The pH initially lower than 2.5, is
20 brought to neutrality by the addition of NaOH (in solu-
25

tion at a concentration of 4 moles/l). The reaction raw product is isolated by precipitation by the addition of 3.5 volumes of methanol and washed with 3.5 volumes of a methanol/water mixture 8/2. The solid is re-dissolved and
5 exhaustively dialyzed against water through a membrane with a cut-off of 12-14,000 Da. 22.5 g of product are obtained with a sulfation degree equal to 1.0 sulfate group per repetitive unit (yield = 87%).

Example 8

10 Evaluation of the regulatory effect of HAS degree 1 and degree 3 on the gene expression of IL-10 and IL-12 in human synoviocytes

Human synoviocytes previously expanded in vitro and maintained in a culture at 37°C with a medium of DMEM
15 containing 10% of FCS, were seeded at a concentration of 20,000 cells per well. Sulfated HA degree 1 (HAS1) and degree 3 (HAS3) prepared as described in Examples 1-3, were then added to the culture medium at concentrations of 0.1 and 0.5 mg/ml (for both samples), whereas the con-
20 trol treatment is represented by non-sulfated HA having an average molecular weight (MW) of 200 KD. After 3 days of treatment, the *PCR Real Time* was effected to evaluate the gene expression of IL-10 and IL-12: the cell RNA was extracted using the "Trizol" method, following the indi-
25 cations of the supplier (TRIZOL reagent, LIFE Techonolo-

gies, GIBCO BRL). In short, the cells were lysed by the addition of 1.0 ml of Trizol and the total RNA was quantified by measuring its absorbance at 260 nm. The appropriate *primers* were selected for each gene to be amplified, using the software Primer3 (Roche Molecular Diagnostics, Pleasanton, CA, USA). The gene expression was evaluated by means of *PCR Real Time* effected with a Rotor-gene TM5500 (Corbett research, Sydney, Australia). The PCR reactions were effected using *primers* at 300 nm and SYBR Green (Invitroge, Carlsbad, CA, USA) at 40 cycles of 15s at 95°C, and 1 min. at 60°C. The value of *Fluorescence thresholds (Ct)*" was automatically determined by the software, evaluating an amplification coefficient for the genes studied between 92 and 110%. For each sample of cDNA, the gene expression value was expressed in terms of the ratio between the ct of the *house keeping* gene (i.e. the gene for the beta-Actin protein which represents the control gene as it is present in every cell and is not subjected to the influence of HAS) and the ct of the gene of interest (i.e. the gene for IL-10 and IL-12), consequently the house keeping ct/gene ct value is indicated in the axis of the ordinates, which therefore indicates the quantity of mRNA expressed by the gene which is being studied. The results obtained are expressed in Figure 1 and 2:

Fig. 1: the treatment of human synoviocytes with HAS1 and HAS3 caused a significant increase in the gene expression of the cytokine IL-10 vs the control treated with non-sulfated HA.

5 Fig. 2:

Also in this experiment, both sulfation degrees (degree 1 and degree 3) of HAS proved to be capable of significantly reducing the gene expression of IL-12, halving the synthesis of its mRNA vs the control treated with non-sulfated HA. The sulfated hyaluronic acid therefore proved to be:

- capable of stimulating the production of new mRNA for the synthesis of anti-inflammatory cytokines, thus increasing the defence capacity of the cell and consequently of the whole organism, vs those pathologies previously described in which IL-10 proved to be of fundamental importance for the resolution and/or improvement of diseases such as asthma, rheumatoid arthritis, arthrosis and all inflammations in which IL-10 is involved.
- effective in diminishing the synthesis of new mRNA of the highly pro-inflammatory cytokine IL-12, proving to be a valid anti-inflammatory agent capable of intervening on the expression of proteins involved in the pathogenesis of invalidating diseases such as psoriasis,

arthritis and all those previously described.

Example 9

Inhibition of the binding of TNF to its receptor expressed in monocyte lines: evaluation of the effectiveness of HAS degree 1 and degree 3 at different MW values

These experiments were effected to evaluate the effectiveness of the samples tested (prepared according to Examples 1-4) on the capacity of inhibiting the binding of TNF to its receptor expressed by cells of the I.S. normally used *in vitro* for this type of experiment, carried out with iodinated cytokine components for an evaluation in *Radioligand Binding assays*.

The experimental procedure was effected as described in Baglioni C. et al., J Biol Chem, 1985, 260:13395-13397.

In short, the line of human histiocytes of the lymphoma U937 was used, with characteristics of monocytes sensitive to the cytotoxic activity of TNF, expressing its relative receptor. The cells were initially incubated with ^{125}I -TNF 0.028nM (carried in water) contemporaneously with the samples to be analyzed (at a concentration of 1 mg/ml which proved to be the lowest concentration which causes the maximum inhibition), in an incubation buffer consisting of 50 mM Tris-HCL pH 7.4, 0.5 mM EDTA, at 4°C for 3 hours.

At the end of the incubation, the cells were centrifuged with dibutylphthalate/dinonylphthalate 2/1 and the pellet obtained was counted in a γ -counter.

The results obtained are expressed in Figure 3:

5 The results obtained show the effectiveness of HAS
in totally (100%) inhibiting the binding of TNF to its
receptor, for both degree 1 and degree 3, with a medium
and low MW. These results are of fundamental importance
as they prove that the behaviour of the sulfated product
10 is completely analogous to that of a monoclonal antibody
specific for the TNF receptor, capable therefore of
blocking its function. This receptor blockage consequently
represents the most effective way of antagonizing
the pro-inflammatory and tumoral effects of the TNF fac-
15 tor.

Example 10

Inhibition of the binding of the cytokine IL-1 to its re-
ceptor expressed in fibroblast lines: evaluation of the
effectiveness of HAS degree 3 at different MW values

20 These experiments were effected to evaluate the ef-
fectiveness of the samples tested (prepared according to
Examples 1-3 and 4) on the capacity of inhibiting the
binding of IL-1 to its receptor expressed by 3T3 cells of
mice, normally used *in vitro* for this type of experiment,
25 carried out with iodinated cytokine components for an

evaluation in *Radioligand Binding assays*.

The experimental procedure was effected as described in Chin J et al., J Exp Med, 1987, 165:70-86.

In short, the line of murine fibroblasts 3T3 was
5 used, sensitive to the cytotoxic activity of IL-1, expressing its relative receptor. The cells were initially incubated with ^{125}I -IL-1 10 pM (carried in water) contemporaneously with the samples to be analyzed (at a concentration of 1 mg/ml which proved to be the lowest concentration
10 tration which causes the maximum inhibition), in an incubation buffer consisting of RPMI 1640 containing 20 mM HEPES pH 7.2 and 1% BSA, at 37°C for 2 hours. At the end of the incubation, the cells were washed with phosphate buffer, then dissolved in 2.5 M of NaOH and counted in a
15 γ -counter.

The results obtained are expressed in Figure 4:

The results obtained show the effectiveness of HAS (with both medium and low MW) in inhibiting the binding of IL-1 to its receptor by 30%. These results are extremely significant as they prove that the behaviour of
20 the sulfated product is completely analogous to that of a monoclonal antibody specific for the receptor of the cytokine in question, capable therefore of blocking its function. This receptor blockage represents the most effective way of antagonizing the pro-inflammatory and tu-
25

moral effects of IL-1, as previously described.

Example 11

Inhibition of the binding of the cytokine IL-6 to its receptor expressed in human myeloma cells: evaluation of the effectiveness of HAS degree 3 at different MW values

These experiments were effected to evaluate the effectiveness of the samples tested (prepared according to Examples 1-3 and 4) on the capacity of inhibiting the binding of IL-6 to its receptor expressed in human myeloma U266, normally used *in vitro* for this type of experiment, carried out with iodinated cytokine components for an evaluation in *Radioligand Binding assays*.

The experimental procedure was effected as described in Taga T. et al., J Exp Med, 1987, 166:967-981.

In short, the line of human myeloma U266 was used, sensitive to the cytotoxic activity of IL-6, expressing its relative receptor. The cells were initially incubated with ^{125}I -IL-6 0.08nM (carried in water) contemporaneously with the samples to be analyzed (at a concentration of 1 mg/ml which proved to be the lowest concentration which causes the maximum inhibition), in an incubation buffer consisting of RPMI 1640 containing 25 mM HEPES pH 7.1 and 10% BSA, at 4°C for 16 hours. At the end of the incubation, the cells were washed with phosphate buffer, centrifuged at 9,000 rpm and the pellet counted in a γ -

counter.

The results obtained are expressed in Figure 5:

The results obtained show the effectiveness of HAS, with both medium and low MW, in inhibiting the binding of
 5 IL-6 to its receptor by up to 100%. These results consequently prove that the behaviour of the sulfated product, also in this case, is completely analogous to that of a monoclonal antibody specific for the receptor of the cytokine in question, capable therefore of blocking its
 10 function. This receptor blockage represents the most effective way of blocking the pro-inflammatory effects of IL-6.

Example 12

Evaluation of the inhibitory effect of HAS degree 1 and
 15 degree 3 on the protein synthesis of the cytokines IL-2, IL-7, IL-10 and IL-12 in human PBMC

For these experimentations, mononucleate cells of human peripheral blood (PBMC) were adopted, deriving from many different donors for evaluating the effect of HAS on
 20 the production of the cytokines listed above, using:

- non-sulfated HA (average MW: 200 KD),
- HAS1 and HAS3 (prepared as described in Examples 1-3 (average MW: 200KD)).

The separation of the PBMC (Bøyum A., *Scand J Clin*
 25 *Lab Invest* 21 Suppl, 1968, 97:77-89) was effected using

the product Ficoll-Paque PLUS (GE Healthcare) and following the protocol indicated by the supplier. At day zero 100,000 cells were seeded per well (using plates with 96 wells) in 200 µl of medium RPMI 1640, to which 10% of fetal bovine serum, HEPES 10 mM, Glutamine 2mM, 1% Penicillin-Streptomycin 100 U/ml, had been added. The effect of all the samples was evaluated on non-treated PBMC with agents which stimulate the synthesis of cytokines, or stimulated with Lipopolysaccharide LPS (10 µg/ml) (highly pro-inflammatory) or with Phytohemagglutinin PHA (10 µg/ml) (a substance capable of stimulating the lymphocytes to divide themselves), both agents capable of stimulating the synthesis of cytokines. The cells were treated separately with the three compounds at a concentration of 0.1 mg/ml or 1 mg/ml. After 24 hours of incubation at 37°C (5% CO₂), 100 µl of supernatant were taken from each well in order to analyze the production of IL-2, IL-7, IL-10 and IL-12.

The quantification of the inflammation mediators was effected by means of the *SearchLight*[®] technology, using a *Custom Human 9-Plex Array* plate following the protocol indicated by the supplier in the technical card.

The results obtained are expressed in Figures 6-9:

These graphs clearly show that HAS degree 1 and degree 3 are capable of significantly reducing the synthesis of

IL-2, IL-7 and IL-12 on the part of monocytes, both when the cells are not stimulated and also when, on the contrary, they are stimulated by specific and powerful inflammatory factors and/or mitogens. HAS therefore proves
 5 to be a molecule with precise pharmacological characteristics, capable of modulating/regulating the synthesis of cytokines with a marked anti-inflammatory activity, both in situations where an immune response is not stimulated, and in particular inflammatory stress events in which the
 10 immune cell responds by producing a cytokine cascade and, above all, in this case, the data presented reveal a greater HAS modulating effect.

Figure 9, on the other hand, confirms the evident stimulus for the production of IL-10 also for cells such
 15 as human monocytes (besides synoviocytes) belonging to the Immune System. It is consequently again confirmed that HAS is capable of modulating the synthesis of cytokines, stimulating those which are anti-inflammatory and inhibiting the synthesis of pro-inflammatory cytokines.

20 **Example 13**

Evaluation of the antiviral action of HAS degree 1 and degree 3 Vs HA-NS:

HIV-1 (III_B) and HIV-2 (ROD) virus

The activity of the samples tested was determined by
 25 evaluating the inhibition of the cytopathogenicity deter-

mined by the HIV-1 virus (obtained from T lymphocytes infected by the type III_B virus) and HIV-2 virus (obtained from T lymphocytes infected by the type ROD virus) in MT-4 cells. The experimental procedure was effected as described in Baba M. et al., ANTIMICROB. AGENTS CHEMOTHER., 1988, 32:1742-1745.

In short, human T lymphocytes called MT-4 infected by the HIV-1 virus strain III_B and HIV-2 strain ROD, were used for this assay. Different concentrations of the samples tested were adopted for this test, using dextran-sulfate as positive control, a polysaccharide known for its capacity of inhibiting different types of viruses. The samples tested were:

1. HS-NS1: sulfated hyaluronic acid prepared by sulfation of the glucosamine after deacetylation, degree 1 (EP0971961), to compare it with
2. HAS1: sulfated hyaluronic acid only in its hydroxyl groups, degree 1, and
3. HAS3: sulfated hyaluronic acid only in its hydroxyl groups, degree 3, prepared as described in Examples 1-3 (all the samples with average MW: 200KD).

The antiviral activity of the samples tested is expressed as the minimum concentration required for inhibiting the cytopathogenicity of the virus by 50%: IC50. The cells were treated with various concentrations of the

above samples added immediately after the infection with infective doses of HIV. After 5 days at 37°C, the number of vital cells is determined with the MTT test: tetrazole salt subjected to redox reaction only by the mitochondrial enzymes of vital cells. The infected and non-treated cells lose the progressive capacity of transforming the tetrazole salt to formazane, consequently, if these cells keep this capacity intact after infection and treatment, this means that this treatment has inhibited the viral action, thus blocking the pathogenicity of the virus (Dezinot F. et al., J Immunol. Methods, 1986, 22(89): 271-277).

The results obtained are expressed in Figure 10:

The results obtained demonstrate the effectiveness of HAS (both degree 1 and degree 3) in inhibiting the viral replication of the virus HIV 1 and 2, in a way which is completely comparable to the positive control (represented by the polysaccharide dextran-sulfate). The experimentation also demonstrated that not all sulfated hyaluronic acids known in the state of the art act effectively vs the replication of the virus studied, as the Applicant has proved that only the HAS whose sulfation has taken place with respect to its hydroxyl groups proves to be active in its powerful antiviral action as HA-NS is not effective in blocking the viral cytopatho-

genicity.

Herpes Simplex virus-1, Herpes Simplex virus-2, Vesicular
stomatitis virus

The activity of the samples tested was determined by
5 evaluating the inhibition of the cytopathogenicity caused
by the virus of Herpes Simplex Virus-1, (HSV-1: KOS, F
and McIntyre *strain*) and by the virus of Herpes Simplex
Virus-2, (HSV-2: G, 196 and Lyons *strain*) in E₆SM fibro-
blast cells which derive from muscular/embryonic skin
10 tissue. Furthermore, the anti-viral activity was tested
again vs E₆SM cells infected by the virus of vesicular
stomatitis (Vesicular stomatitis virus: VSV). HSV-1 is
the virus which preferentially infects the oral mucous,
whereas HSV-2 attacks the genital mucous. The experimen-
15 tal procedure was effected as described in Baba M. et
al., ANTIMICROB. AGENTS CHEMOTHER., 1988, 32:1742-1745.

In short, confluent cell cultures were exposed to
infective doses of the viruses listed above in the pres-
ence of the samples HS-NS1 (EP0971961), HAS1 and HAS3
20 prepared as described in Example 1 and 2 (all the samples
with an average MW: 200KD). After an incubation period of
1h at 37°C, the culture medium was substituted with fresh
medium only containing the samples to be tested. The cy-
topathogenicity of the virus was tested on the 2nd day of
25 incubation. The measurement of the inhibition of the vi-

ral cytopathogenicity was evaluated by determining the inhibition of the synthesis of DNA and RNA in the "infected" cells and subjected to treatment as indicated above: the cells were seeded in micro-wells in a culture medium containing different concentrations of the samples to be tested with 2.5 μ Ci of 3H-thymidine and 3H-uridine per ml. After 16 h at 37°C, the cells were treated with Trichloroacetic acid, washed in ethanol, left to dry and counted in 7.5 ml of liquid by scintillation. The antiviral activity of the samples tested is expressed as the minimum concentration required for inhibiting the cytopathogenicity of the virus by 50%: IC50. Furthermore, in order to also evaluate the cytotoxicity of the samples tested, the minimum concentrations necessary for causing morphological damage (observable with an optical microscope) to the cells used, were determined. The comparison was effected vs dextran-sulfate and the drug Acyclovir (both molecules with a known antiviral effectiveness, therefore used as positive control).

20 The results obtained are expressed in Figure 11:

 The experimental data confirm what was seen for HIV, i.e. the powerful antiviral action of both HAS1 and HAS3: a comparison between HA-NS1 and HAS1 shows that not all sulfated hyaluronic acids are equivalent as HA-NS1 did not prove to be active, and this difference in efficacy

does not depend on the molecular weight or the sulfation degree of hyaluronic acid, it therefore lies in the very structure of HA-NS1 vs HAS1, i.e. on the sulfation. HAS in fact shows an efficacy equal to that of dextran-
5 sulfate and comparable to that of acyclovir, a reference drug for the treatment of Herpes Simplex. Furthermore, it should be pointed out that acyclovir is inactive vs VSV, whereas HAS degree 1 and 3 has a very powerful antiviral activity vs VSV.

10 All the samples tested are non-cytotoxic towards the host cell, the minimum cytotoxic concentration obtained is in fact equal to that of the reference drugs normally used in clinical practice for the treatment of Herpes, and proved to be on an average 100 times higher than that
15 proved active in the inhibition of the viral replication.

Cytomegalovirus

The activity of the samples tested was determined by evaluating the inhibition of the cytopathogenicity determined by Cytomegalovirus (CMV: AD-169 and Davis strain)
20 using the previous protocol. The antiviral activity was tested vs HEL cells (pulmonary embryonic cells) and was expressed as the concentration required for inhibiting the number of plaques formed by the above virus by 50%.

The results obtained are expressed in Figure 12:

25 the table indicates the clear and significant positive

result obtained for both HAS1 and HAS3 which again confirms them as effective antiviral agents. Also in this case, HA-NS1 did not prove to be active in inhibiting the proliferation of the virus, confirming the absolute diversity between the two types of degree 1 sulfated product in having an antiviral capacity.

Example 14

Preparation of a formulation in the form of a solution for inhalations containing sulfated hyaluronic acid degree 1

40 mg (or 20 mg if the HAS has a MW of 500-730KD) of sulfated hyaluronic acid degree 1, having a low or medium MW, are introduced into a 50 ml glass flask, after which 15 ml of PBS 0.2M at sterile pH 7.4, are added. The mixture is subjected to stirring for about 30 minutes, until the complete dissolution of the powder. When the complete dissolution has been obtained, 2 ml of glycol propylene and further PBS 0.2M at sterile pH 7.4 are added until the total volume of 20 ml is reached. The solution is maintained under stirring for a few minutes.

Example 15

Preparation of a formulation in the form of a SOLUTION for inhalations containing sulfated hyaluronic acid degree 3

100 mg of sulfated hyaluronic acid degree 3 obtained

from HA 200KD are introduced into a 50 ml glass flask, after which 15 ml of PBS 0.2M at sterile pH 7.4, are added. The mixture is subjected to stirring for about 30 minutes, until the complete dissolution of the powder.

- 5 When the complete dissolution has been obtained, 2 ml of glycol propylene and further PBS 0.2M at sterile pH 7.4 are added until the total volume of 20 ml is reached. The solution is maintained under stirring for a few minutes.

Example 16

- 10 Preparation of a formulation in the form of an injectable SOLUTION for intra-articular use containing sulfated hyaluronic acid degree 1

500 mg of sulfated hyaluronic acid degree 1 obtained from HA with a MW of 500-730KD are introduced into a 50 ml glass flask, and PBS 0.2M at sterile pH 7.4, is then added until the total volume of 20 ml is reached. The mixture is subjected to stirring for about 60 minutes, until the complete dissolution of the powder.

Example 17

- 20 Preparation of a formulation in the form of a HYDROPHILIC GEL containing HAS, HA and CMC

Methyl- and propyl-parabene are dissolved in purified water at 80°C. After cooling the solution to room temperature, sodium hyaluronate is added under stirring until dissolution followed by HAS1 (or HAS3), maintaining

the stirring until complete dissolution. Glycerol and propylene glycol are then added under stirring until complete dissolution. Sodium carboxymethylcellulose (CMC) is finally added and the mixture is mixed until a gelified
5 solution is obtained.

Example 18

Preparation of a formulation in the form of a HYDROPHILIC GEL for mucosal use (without preservatives) containing HAS and HA

10 Sodium hyaluronate is dissolved under stirring, and then HAS1 (or HAS3) in a quantity of water of about 90% of that envisaged in the formula. Propylene glycol, Symdiol 68 are added followed by MP Diol Glycol and the mixture is mixed until the complete dissolution of the various
15 components. Carbomer 974P is subsequently added and the stirring is maintained until the homogeneous dispersion of the latter. Beads of sodium hydroxide are dissolved in the remaining 10% of water and this solution is added to the that previously obtained, to obtain the
20 gelification of the aqueous phase.

Example 19

Preparation of a formulation in the form of a LIPSTICK containing HAS and HA

The correct quantity of liquid paraffin indicated in
25 the manufacturing formula is charged into a suitable con-

tainer. It is heated to 88-92°C and white soft paraffin, hard paraffin, white beeswax, ceresin, arlacel are then added under stirring, the stirring is maintained until the complete melting of the various components. All-rac-
5 a-tocopheryl acetate, allantoin, butylhydroxytoluene, propyl p-hydroxybenzoate are then incorporated and the mixture is mixed until complete dissolution, maintaining the mass at 88-92°C.

The quantity of purified water envisaged in the formula is charged separately into a suitable container, sodium hyaluronate, HAS1 (or HAS3) are then added under stirring until complete dissolution, followed by Disodium Edetate maintaining the stirring until dissolution.
10

The aqueous phase is transferred under stirring to the container containing the molten mass, maintaining the system at 88-92°C and the stirring until a limpid solution is obtained. The two aromatizing agents are then added under stirring and the mixture is mixed for 10'. The molten mass is poured into moulds and is immediately
15 cooled to $T < 0^{\circ}\text{C}$ until solid sticks are obtained.
20

Example 20

Preparation of a formulation in the form of VAGINAL OVULES containing HAS and HA

The gelatine is allowed to swell in 70% of purified water at 85°C; sodium hyaluronate and HAS1 (or HAS3) are
25

dissolved in the remaining quantity of water and this solution is mixed with the glycerine brought to the same temperature. The glycerine solution is added to the swollen gelatine solution and the stirring is maintained until the complete dissolution of the gelatine. The mass is poured into moulds and cooled to $T < 0^{\circ}\text{C}$ until solid ovules are obtained.

Example 21

Preparation of a formulation in the form of a HYDROPHILIC CREAM containing HAS and HA

The oil phase is prepared by melting liquid paraffin, stearic acid and Tefose 1500 under stirring at 50°C . The aqueous phase is prepared separately by the initial dissolution at 80°C of methyl-parabene and subsequent cooling to room temperature and the incorporation of glycerol, sodium hyaluronate and subsequently HAS1 (or HAS3) under stirring until the complete dissolution of the various components.

The aqueous phase is joined to the oil phase and emulsification is effected, the emulsion O/A obtained is cooled under stirring to room temperature.

Example 22

Preparation of a formulation in the form of an OINTMENT containing HAS

The base ointment is prepared by melting light liq-

uid paraffin and white Vaseline under stirring at 70°C. After cooling to room temperature, HAS1 (or HAS3) is incorporated under stirring and the mixture is mixed until a homogeneous suspension is obtained.

5 **Example 23**

Preparation of a formulation in the form of CAPSULES (hard gelatine) containing HAS

HAS1 (HAS3) is mixed with calcium phosphate, magnesium stearate and silica by progressive dilution. The capsules are then filled.

Example 24

Preparation of a formulation in the form of TABLETS containing HAS

HAS1 (or HAS3) is subjected to wet granulation in a fluid bed with a ligand solution consisting of water and about 70% of the total quantity of sodium CMC. The granulate obtained is subjected to sieving on an 0.8 mm net.

The remaining ingredients (calcium phosphate, microcrystalline cellulose, sodium CMC and silica) are mixed and sieved on an 0.8 mm net.

The granulate previously obtained is mixed with the mixture consisting of the remaining ingredients (extragranules), and is finally mixed with magnesium stearate (previously sieved on an 0.8 mm net), followed by compression.

Example 25Preparation of a formulation in the form of an INJECTIVE SOLUTION containing HAS

After the preparation (at room temperature) of a
 5 physiological solution buffered at pH 6.4-7.2, lactose is dissolved under stirring and finally HAS1 or HAS3. The solution thus obtained is filtered on 0.22 microns.

Table

HYDROPHILIC GEL (Example 17)

Components	Quantity (mg/1g of hydrogel)
HAS1 (HAS3)	40 mg (10 mg)
CMC	20 mg
Glycerol	100 mg
Propylene Glycol	66,75 mg
Sodium Hyaluronate	2 mg
Methyl p-hydroxybenzoate	2 mg
Propyl p-hydroxybenzoate	0,2 mg
Purified Water	q.s to 1g

HYDROPHILIC GEL for mucosal use (Example 18)

Components	Quantity (mg/1g of hydrogel)
HAS1 (HAS3)	10 mg

Carbomer 974P	15 mg
Propylene Glycol	100 mg
Sodium Hydroxyde	0,33 mg
Sodium Hyaluronate	2 mg
MP-Diol Glycol	37,5 mg
SymDiol 68	90 mg
Purified Water	q.s to 1g

LIPSTICK (Example 19)	
Components	Quantity (mg/1g lipstick)
HAS1 (HAS3)	30 mg (10 mg)
Liquid Paraffin	253,2 mg
White Soft Paraffin	326,2 mg
Hard Paraffin	144,3 mg
Beeswax white	96 mg
Ceresin	28,2 mg
Arlacel 582	95,8 mg
Sodium Hyaluronate	2 mg
Allantoin	1,1 mg
all-rac-a Tocopheryl Acetate	1,1 mg
Propyl p-hydroxybenzoate	0,4 mg
Butylhydroxytoluene	0,4 mg
Purified Water	19,2 mg

Disodium Edetate	1,1 mg
Vanilla Flavour	0,5 mg
Sweet Flavour	0,5 mg
VAGINAL OVULES (Example 20)	
Components	Quantity (mg/1g of ovule)
HAS1 (HAS3)	10 mg
Glycerin	580
Gelatin	200
Sodium Hyaluronate	2
Purified Water	q.s to 1g
HYDROPHILIC CREAM (Emulsion O/A) (Example 21)	
Components	Quantity mg/1g of cream)
HAS1 (HAS3)	10 mg
Tefose 1500	110 mg
Glycerol	80 mg
Stearic Acid	33 mg
Sodium Hyaluronate	2 mg
Liquid Paraffin	40 mg
Methyl p-hydroxybenzoate	1 mg
Purified Water	q.s to 1g

OINTMENT (Example 22)	
Components	Quantity (mg/1g of ointment)
HAS1 (HAS3)	20 mg
Light Liquid Paraffin	200 mg
White Petrolatum	q.s to 1g

CAPSULES (Hard Gelatine) (Example 23)	
Components	Quantity (mg/capsule)
HAS-1 (HAS3)	80 mg (40mg)
Calcium Phosphate	256 mg
Magnesium Stearate	7 mg
Silica	7 mg
TABLETS (Example 24)	
Components	Quantity (mg/tablet)
HAS-1 (HAS-3)	120 mg
Calcium Phosphate	200
Microcrystalline cellulose	185
Sodium Carboxymethylcellulose	10
Magnesium Stearate	10
Silica	15

INJECTIVE FORMULATIONS (Example 25)	
Components	Quantity (mg/ml of sol)
HAS-3	50 mg
Lactose	0,93 mg
Potassium Phosphate Dibasic	0,36
Potassium Phosphate Monobasic	0,23
Sodium Chloride	9

The description being thus described, these methods can obviously be modified in various ways. These modifications should not be considered as diverging from the spirit and perspectives of the invention and all the modifications which can appear evident to a skilled person in the field are included in the scope of the following claims.

CLAIMS

1. Sulphated hyaluronic acid for use in the prevention and/or treatment of herpes genitalis, said sulphated hyaluronic acid (HA) being obtained by a sulfation involving
5 the alcohol hydroxyls present in the HA polysaccharide chain, starting from a HA having a molecular weight ranging from 10,000 to 50,000 Da, 150,000 to 250,000 Da and 500,000 to 750,000 Da, said sulphated hyaluronic acid having a sulphation degree equal to 1 or 3.
- 10 2. Sulphated hyaluronic acid for use according to claim 1 for systemic administration.
3. Sulphated hyaluronic acid according to any one of claims 1 to 2 for topic or inhalation administration.

1/6

Fig. 1





Fig. 2



2/6

Fig. 3





Fig. 4



3/6

Fig. 5





Fig. 6



4/6

Fig. 7




Fig. 8




Fig. 9




Fig. 10

Product tested	Minimum inhibitory concentration IC 50 (µg/ml)	
	HIV-1 (III _s Strain)	HIV-2 (ROD Strain)
HA -NS1	>125	>250
HAS1	2,3	14,4
HAS3	0,15	0,66
DS	0,135	0,025

6/6

Fig. 11

Product tested	Minimum Cytotox. concentration (µg/ml)	Minimum inhibitory concentration IC 50 (µg/ml)						
		Herpes simplex virus-1 (KOS)	Herpes simplex virus-1 (F)	Herpes simplex virus-1 (McIntyre)	Herpes simplex virus-2 (G)	Herpes simplex virus-2 (196)	Herpes simplex virus-2 (Lyons)	Vesicular stomatitis virus
HA-NS1	>400	>400	>400	>400	>400	>400	>400	>400
HAS1	>400	48	16	9,6	9,6	3,2	3,2	1,92
HAS3	>400	16	3,2	1,92	1,92	1,92	1,92	0,128
DS	>400	9,6	9,6	9,6	3,2	3,2	1,92	16
Acyclovir	>400	0,384	0,384	0,64	0,384	0,384	0,384	>80

Fig. 12

Product tested	Minimum inhibitory concentration IC 50 (µg/ml)	
	Cytomegalovirus AD-169 strain	Cytomegalovirus Davis strain
HA-NS1	>50	>50
HAS1	5	3,4
HAS3	0,2	0,2
DS	2,7	0,5