

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 December 2006 (28.12.2006)

PCT

(10) International Publication Number
WO 2006/138656 A2

(51) International Patent Classification:

C07D 487/04 (2006.01) A61P 3/04 (2006.01)
A61K 31/5025 (2006.01)

(74) Agents: DUNCAN, Sammy G., Ph.D. et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(21) International Application Number:

PCT/US2006/023637

(22) International Filing Date: 19 June 2006 (19.06.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/692,042 17 June 2005 (17.06.2005) US
11/455,045 16 June 2006 (16.06.2006) US

(71) Applicant (for all designated States except US): BRISTOL-MYERS SQUIBB COMPANY [US/US]; Route 206 and Province Line Road, Princeton, New Jersey 08543-4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SUN, Chongqing [US/US]; 527 Dutch Neck Road, East Windsor, New Jersey 08520 (US). HUANG, Yanting [CA/US]; 42 Manley Road, Pennington, New Jersey 08534 (US). SITKOFF, Doree, F. [US/US]; 1537 Lexington Drive, Drescher, Pennsylvania 19025 (US). MIKKILINENI, Amarendra, B. [US/US]; 1230 Sycamore Avenue, Easton, Pennsylvania 18040 (US). YU, Guixue [US/US]; 8 Greene Drive, Princeton Junction, New Jersey 08550 (US). EWING, William, R. [US/US]; 1193 Waterwheel Drive, Princeton Junction, New Jersey 19067 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]

(54) Title: BICYCLIC HETEROCYCLES AS CANNABINOID-1 RECEPTOR MODULATORS



(I)

(57) Abstract: The present application describes compounds according to Formula (I), pharmaceutical compositions comprising at least one compound according to Formula (I) and optionally one or more additional therapeutic agents and methods of treatment using the compounds according to Formulas (I) both alone and in combination with one or more additional therapeutic agents. The compounds have the following general formulas: including all prodrugs, solvates, pharmaceutically acceptable salts and stereoisomers, wherein B, R¹, R², R³ and x are described herein.



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

BICYCLIC HETEROCYCLES AS CANNABINOIND-1 RECEPTOR MODULATORS


BACKGROUND

- 5 [0001] Delta-9-tetrahydrocannabinol or Delta-9 THC, the principle active component of *Cannabis sativa* (marijuana), is a member of a large family of lipophilic compounds (i.e., cannabinoids) that mediate physiological and psychotropic effects including regulation of appetite, immunosuppression, analgesia, inflammation, emesis, anti-nociception, sedation, and intraocular pressure. Other members of the
- 10 cannabinoid family include the endogenous (arachidonic acid-derived) ligands, anandamide, 2-arachidonyl glycerol, and 2-arachidonyl glycerol ether. Cannabinoids work through selective binding to and activation of G-protein coupled cannabinoid receptors. Two types of cannabinoid receptors have been cloned including CB-1 (L. A. Matsuda, et al., *Nature*, 346, 561-564 (1990)), and CB-2 (S. Munro, et al., *Nature*,
- 15 365, 61-65 (1993)). The CB-1 receptor is highly expressed in the central and peripheral nervous systems (M. Glass, et al., *Neuroscience*, 77, 299-318 (1997)), while the CB-2 receptor is highly expressed in immune tissue, particularly in spleen and tonsils. The CB-2 receptor is also expressed on other immune system cells, such as lymphoid cells (S. Galiege, et al., *Eur J Biochem*, 232, 54-61 (1995)). Agonist
- 20 activation of cannabinoid receptors results in inhibition of cAMP accumulation, stimulation of MAP kinase activity, and closure of calcium channels.
- [0002] There exists substantial evidence that cannabinoids regulate appetitive behavior. Stimulation of CB-1 activity by anandamide or Delta-9 THC results in increased food intake and weight gain in multiple species including humans (Williams and Kirkham, *Psychopharm.*, 143, 315-317 (1999)). Genetic knock-out of CB-1 result
- 25 in mice that were hypophagic and lean relative to wild-type litter mates (DiMarzo, et al., *Nature*, 410, 822- 825 (2001)). Published studies with CB-1 small molecule antagonists have demonstrated decreased food intake and body weight in rats (Trillou, et. al., *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, R345-R353, (2003)). Chronic
- 30 administration of the CB-1 antagonist AM-251 for two weeks resulted in substantial body weight reduction and decreased adipose tissue mass (Hildebrandt, et. al., *Eur. J. Pharm*, 462, 125-132 (2003)). There are multiple studies that have assessed the

anorexic effect of the Sanofi CB-1 antagonist, SR-141716 (Rowland, et. al., *Psychopharm.*, 159, 111-116 (2001); Colombo, et. al., *Life Sci.*, 63, 113-117 (1998)). There are at least two CB-1 antagonists in clinical trials for regulation of appetite, Sanofi's SR-141716 and Solvay's SLV-319. Published Phase IIb data reveal that SR-141716 dose-dependently reduced body weight in human subjects over a 16 week trial period. CB-1 antagonists have also been shown to promote cessation of smoking behavior. Phase II clinical data on smoking cessation were presented in September of 2002 at Sanofi-Synthelabo's Information meeting. This data showed that 30.2% of patients treated with the highest dose of SR-141716 stayed abstinent from cigarette smoke relative to 14.8% for placebo.

DETAILED DESCRIPTION

[0003] The present application describes compounds according to Formula I, pharmaceutical compositions comprising at least one compound according to Formula I and optionally one or more additional therapeutic agents and methods of treatment using the compounds according to Formula I both alone and in combination with one or more additional therapeutic agents. The compounds claimed have the general Formula I



I

20 including all prodrugs, solvates, pharmaceutically acceptable salts and stereoisomers,
wherein B, R¹, R², R³ and x are described herein.

DEFINITIONS

[0004] The following definitions apply to the terms as used throughout this
25 specification, unless otherwise limited in specific instances.

[0005] As used herein, the term "alkyl" denotes branched or unbranched hydrocarbon chains containing 1 to 20 carbons, preferably 1 to 12 carbons, and more preferably 1 to 8 carbons, in the normal chain, such as, methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, iso-butyl, tert-butyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl and the like. Further, alkyl groups, as defined herein, may optionally be substituted on any available carbon atom with one or more functional groups commonly attached to such chains, such as, but not limited to hydroxyl, halo, haloalkyl, mercapto or thio, cyano, alkylthio, cycloalkyl, heterocyclyl, aryl, heteroaryl, carboxyl, carbalkoyl, carboxamido, carbonyl, alkenyl, alkynyl, nitro, amino, alkoxy, aryloxy, arylalkyloxy, heteroaryloxy, amido, -OC(O)NR⁸R⁹, -OC(O)R⁸, -OPO₃H, -OSO₃H, and the like to form alkyl groups such as trifluoromethyl, 3-hydroxyhexyl, 2-carboxypropyl, 2-fluoroethyl, carboxymethyl, cyanobutyl and the like.

[0006] Unless otherwise indicated, the term "alkenyl" as used herein by itself or as part of another group refers to straight or branched chains of 2 to 20 carbons, preferably 2 to 12 carbons, and more preferably 2 to 8 carbons with one or more double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like. Further, alkenyl groups, as defined herein, may optionally be substituted on any available carbon atom with one or more functional groups commonly attached to such chains, such as, but not limited to halo, haloalkyl, alkyl, alkoxy, alkynyl, aryl, arylalkyl, cycloalkyl, amino, hydroxyl, heteroaryl, cycloheteroalkyl, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, alkylthio and/or any of the alkyl substituents set out herein.

[0007] Unless otherwise indicated, the term "alkynyl" as used herein by itself or as part of another group refers to straight or branched chains of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons with one or more triple bonds in the normal chain, such as 2-propynyl, 3-butylnyl, 2-butylnyl, 4-pentylnyl, 3-pentylnyl, 2-hexynyl, 3-hexynyl, 2-heptylnyl, 3-heptylnyl, 4-heptylnyl, 3-octynyl, 3-nonylnyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like. Further, alkynyl groups, as defined herein, may optionally be substituted on any available carbon atom with one

or more functional groups commonly attached to such chains, such as, but not limited to halo, haloalkyl, alkyl, alkoxy, alkenyl, aryl, arylalkyl, cycloalkyl, amino, hydroxyl, heteroaryl, cycloheteroalkyl, alkanoylamino, alkylamido, arylcarbonylamino, nitro, cyano, thiol, alkylthio and/or any of the alkyl substituents set out herein.


- 5 [0008] Unless otherwise indicated, the term “cycloalkyl” as employed herein alone or as part of another group includes saturated or partially unsaturated (containing one or more double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, appended or fused, including monocyclicalkyl, bicyclicalkyl and tricyclicalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 3 to 10 carbons, forming the ring and which may be fused to 1 or 2 aromatic rings as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl,



- 15 [0009] Further, any cycloalkyl may be optionally substituted through any available carbon atoms with one or more groups selected from hydrogen, halo, haloalkyl, alkyl, alkoxy, haloalkyloxy, hydroxyl, alkenyl, alkynyl, aryl, aryloxy, heteroaryl, heteroaryloxy, arylalkyl, heteroarylalkyl, alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the alkyl substituents.

- 20 [0010] The term “cycloalkylalkyl” as used herein alone or as part of another group refers to alkyl groups as defined above having a cycloalkyl substituent, wherein said “cycloalkyl” and/or “alkyl” groups may optionally be substituted as defined above.


- [0011] Unless otherwise indicated, the term “aryl” as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including 1-naphthyl and 2-naphthyl) and may optionally include one to three additional rings fused to a carbocyclic ring or a heterocyclic ring, for example



- 5 **[0012]** Further, “aryl”, as defined herein, may optionally be substituted with one or more functional groups, such as halo, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocycloalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, alkoxycarbonyl, arylcarbonyl, arylalkenyl, aminocarbonylaryl, arylthio, arylsulfinyl, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxyl, nitro, cyano, amino, substituted amino
- 10 wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl,
- 15 alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino or arylsulfonaminocarbonyl and/or any of the alkyl substituents set out herein.

- [0013]** Unless otherwise indicated, the term “heteroaryl” as used herein alone or as part of another group refers to a 5- or 6- membered aromatic ring which includes 1,
- 20 2, 3 or 4 hetero atoms such as nitrogen, oxygen or sulfur. Such rings may be fused to an aryl, cycloalkyl, heteroaryl or heterocyclyl and include possible N-oxides as described in Katritzky, A. R. and Rees, C. W., eds. *Comprehensive Heterocyclic Chemistry: The Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds* 1984, Pergamon Press, New York, NY; and Katritzky, A. R., Rees, C. W., Scriven, E.

F., eds. *Comprehensive Heterocyclic Chemistry II: A Review of the Literature 1982-1995* 1996, Elsevier Science, Inc., Tarrytown, NY; and references therein. Further, “heteroaryl”, as defined herein, may optionally be substituted with one or more substituents such as the substituents included above in the definition of “substituted alkyl” and “substituted aryl”. Examples of heteroaryl groups include the following:



and the like.

[0014] The term “heteroarylalkyl” as used herein alone or as part of another group refers to alkyl groups as defined above having a heteroaryl substituent, wherein said heteroaryl and/or alkyl groups may optionally be substituted as defined above.

[0015] The term “heterocyclo”, “heterocycle”, “heterocyclyl” or “heterocyclic ring”, as used herein, represents an unsubstituted or substituted stable 4 to 7-membered monocyclic ring system which may be saturated or unsaturated, and which consists of carbon atoms, with one to four heteroatoms selected from nitrogen, oxygen or sulfur, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in

the creation of a stable structure. Examples of such heterocyclic groups include, but is not limited to, piperidinyl, piperazinyl, oxopiperazinyl, oxopiperidinyl, oxopyrrolidinyl, oxoazepinyl, azepinyl, pyrrolyl, pyrrolidinyl, furanyl, thienyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isooxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, thiadiazolyl, tetrahydropyranyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, oxadiazolyl and other heterocycles described in Katritzky, A. R. and Rees, C. W., eds. *Comprehensive Heterocyclic Chemistry: The Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds* 1984, Pergamon Press, New York, NY; and Katritzky, A. R., Rees, C. W., Scriven, E. F., eds. *Comprehensive Heterocyclic Chemistry II: A Review of the Literature 1982-1995* 1996, Elsevier Science, Inc., Tarrytown, NY; and references therein.

[0016] The term "heterocycloalkyl" as used herein alone or as part of another group refers to alkyl groups as defined above having a heterocyclyl substituent, wherein said heterocyclyl and/or alkyl groups may optionally be substituted as defined above.

[0017] The terms "arylalkyl", "arylalkenyl" and "arylalkynyl" as used alone or as part of another group refer to alkyl, alkenyl and alkynyl groups as described above having an aryl substituent. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, phenethyl, benzhydryl and naphthylmethyl and the like.

[0018] The term "alkoxy", "aryloxy", "heteroaryloxy" "arylalkyloxy", or "heteroarylalkyloxy" as employed herein alone or as part of another group includes an alkyl or aryl group as defined above linked through an oxygen atom.

[0019] The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine, with bromine, chlorine or fluorine being preferred.

[0020] The term "cyano," as used herein, refers to a -CN group.

[0021] The term "methylene," as used herein, refers to a -CH₂- group.

[0022] The term "nitro," as used herein, refers to a -NO₂ group.

- [0023] The compounds of formula I can be present as salts, which are also within the scope of this invention. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred. If the compounds of formula I have, for example, at least one basic center, they can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, for example sulfuric acid, phosphoric acid or a hydrohalic acid, with organic carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms, for example acetic acid, which are unsubstituted or substituted, for example, by halogen as chloroacetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, (for example aspartic or glutamic acid or lysine or arginine), or benzoic acid, or with organic sulfonic acids, such as (C₁-C₄) alkyl or arylsulfonic acids which are unsubstituted or substituted, for example by halogen, for example methyl- or *p*-toluene- sulfonic acid. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center. The compounds of formula I having at least one acid group (for example COOH) can also form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono, di or tri-lower alkylamine, for example ethyl, *tert*-butyl, diethyl, diisopropyl, triethyl, tributyl or dimethyl-propylamine, or a mono, di or trihydroxy lower alkylamine, for example mono, di or triethanolamine. Corresponding internal salts may furthermore be formed. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds of formula I or their pharmaceutically acceptable salts, are also included.
- [0024] Preferred salts of the compounds of formula I which contain a basic group include monohydrochloride, hydrogensulfate, methanesulfonate, phosphate, nitrate or acetate.
- [0025] Preferred salts of the compounds of formula I which contain an acid group include sodium, potassium and magnesium salts and pharmaceutically acceptable organic amines.

[0026] The term “modulator” refers to a chemical compound with capacity to either enhance (e.g., “agonist” activity) or partially enhance (e.g., “partial agonist” activity) or inhibit (e.g., “antagonist” activity or “inverse agonist” activity) a functional property of biological activity or process (e.g., enzyme activity or receptor binding); such enhancement or inhibition may be contingent on the occurrence of a specific event, such as activation of a signal transduction pathway, and/or may be manifest only in particular cell types.

[0027] The term “bioactive metabolite” as employed herein refers to any functional group contained in a compound of formula I with an open valence for further substitution wherein such substitution can, upon biotransformation, generate a compound of formula I. Examples of such functional groups of bioactive metabolites include, but are not limited to, -OH, -NH or functional groups wherein the hydrogen can be replaced with a functional group such as -PO₃H₂ for example, which, upon biotransformation generates an -OH or -NH functional group of a compound of formula I.

[0028] The term “prodrug esters” as employed herein includes esters and carbonates formed by reacting one or more hydroxyls of compounds of formula I with alkyl, alkoxy, or aryl substituted acylating agents employing procedures known to those skilled in the art to generate acetates, pivalates, methylcarbonates, benzoates and the like. Prodrug esters may also include- but are not limited to groups such as phosphate esters, phosphonate esters, phosphoramidate esters, sulfate esters, sulfonate esters, and sulfonamidate esters wherein the ester may be further substituted with groups that confer a pharmaceutical advantage such as-but not limited to-favorable aqueous solubility or in vivo exposure to the bioactive component formula I.

[0029] The term “prodrug” as employed herein includes functionalization of bioactive amine- or hydroxyl-containing compounds of formula I to form alkyl-, acyl-, sulfonyl-, phosphoryl-, or carbohydrate-substituted derivatives. Such derivatives are formed by reacting compounds of formula I with alkylating-, acylating-, sulfonylating-, or phosphorylating reagents employing procedures known to those skilled in the art. Alkylation of amines of formula I may result in- but are not limited to- derivatives that include spacer units to other prodrug moieties such as substituted alkyloxymethyl-, acyloxymethyl-, phosphoryloxymethyl-, or sulfonyloxymethyl-

groups. Alkylation of amines of formula I may result in the generation of quarternary amine salts that act in vivo to provide the bioactive agent (i.e., the compound of formula I).

- 5 [0030] Preferred prodrugs consist of a compound of formula I where a pendant hydroxyl is phosphorylated to generate a phosphate derivative. Such a prodrug may also include a spacer group between the compound of formula I and the phosphate group, such as a methyleneoxy-group. Methods to generate such a prodrug from a compound of formula I are known to those skilled in the art, and are listed in the references below.
- 10 [0031] Preferred prodrugs also consist of a compound of formula I where a pendant amine, such as a pyridine group, is alkylated with a group, such as methyl, to form a quarternary ammonium ion salt. Methods to generate such a prodrug from a compound of formula I are known to those skilled in the art, and are listed in the references below.
- 15 [0032] Any compound that can be converted in vivo to provide the bioactive agent (i.e., the compound of formula I) is a prodrug within the scope and spirit of the invention.
- [0033] Various forms of prodrugs are well known in the art. A comprehensive description of prodrugs and prodrug derivatives are described in:
- 20 a) *The Practice of Medicinal Chemistry*, Camille G. Wermuth et al., Ch 31, (Academic Press, 1996);
- b) *Design of Prodrugs*, edited by H. Bundgaard, (Elsevier, 1985);
- c) *A Textbook of Drug Design and Development*, P. Krogsgaard-Larson and H. Bundgaard, eds. Ch 5, pgs 113 – 191 (Harwood Academic Publishers, 1991);
- 25 d) *Hydrolysis in Drug and Prodrug Metabolism*, B. Testa and J. M. Mayer, (Verlag Helvetica Chimica Acta AG, Zurich, Switzerland; Wiley-VCH, Weinheim, Federal Republic of Germany, 2003);
- e) Etmayer, P.; Amidon, G. L.; Clement, B.; Testa, B. "Lessons Learned from Marketed and Investigational Prodrugs" *J. Med. Chem.* 2004, 47 (10), 2393-
- 30 2404; and

f) Davidsen, S. K. et al. "N-(Acyloxyalkyl)pyridinium Salts as Soluble Prodrugs of a Potent Platelet Activating Factor Antagonist" *J. Med. Chem.* 1994, 37 (26), 4423-4429.

Said references are incorporated herein by reference.

5

[0034] An administration of a therapeutic agent of the invention includes administration of a therapeutically effective amount of the agent of the invention. The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent to treat or prevent a condition treatable by administration of a composition of the invention. That amount is the amount sufficient to exhibit a detectable therapeutic or preventative or ameliorative effect. The effect may include, for example, treatment or prevention of the conditions listed herein. The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition being treated, recommendations of the treating physician, and the therapeutics or combination of therapeutics selected for administration.

[0035] All stereoisomers of the compounds of the instant invention are contemplated, either in mixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of the carbon atoms including any one of the R substituents. Consequently, compounds of formula I can exist in enantiomeric or diastereomeric forms or in mixtures thereof. The processes for preparation can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric or enantiomeric products are prepared, they can be separated by conventional methods for example, chromatographic techniques, chiral HPLC or fractional crystallization.

[0036] The compounds of formula I of the invention can be prepared as shown in the following reaction schemes and description thereof, as well as relevant published literature procedures that may be used by one skilled in the art. Exemplary reagents and procedures for these reactions appear hereinafter and in the working Examples.

30

ABBREVIATIONS

[0037] The following abbreviations are employed in the Schemes, Examples and elsewhere herein:

- AcOH = acetic acid
- 5 AlCl₃ = aluminum chloride
- Boc = *tert*-butoxycarbonyl
- DMF = N,N-dimethylformamide
- DMSO = dimethyl sulfoxide
- EtOAc = ethyl acetate
- 10 Et₃N = triethylamine
- Et₂O = diethyl ether
- HCl = hydrochloric acid
- HPLC or LC = high performance liquid chromatography
- K₃PO₄ = potassium phosphate tribasic
- 15 K₂CO₃ = potassium carbonate
- MeOH = methanol
- MOM = methoxymethyl
- Ms = methanesulfonyl
- MS or Mass Spec = mass spectrometry
- 20 NaCl = sodium chloride
- Na₂CO₃ = sodium carbonate
- NaHCO₃ = sodium bicarbonate
- Na₂SO₄ = sodium sulfate
- NaOH = sodium hydroxide
- 25 Pd(Ph₃P)₄ = tetrakis(triphenylphosphine)palladium(0)
- PG = protecting group
- Pd(dppf)Cl₂ = [1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium(II)
- POCl₃ = phosphorus oxychloride
- TFA = trifluoroacetic acid
- 30 THF = tetrahydrofuran
- Ts = *p*-toluenesulfonyl
- min = minute(s)

- h = hour(s)
L = liter(s)
mL = milliliter(s)
 μ L = microliter(s)
- 5 g = gram(s)
mg = milligram(s)
mol = mole(s)
mmol = millimole(s)
M = molar
- 10 nM = nanomolar
[M+H]⁺ = parent plus a proton
[M+Na]⁺ = parent plus a sodium ion
[M-H]⁻ = parent minus a proton
- 15 **[0038]** Compounds of the present invention may be prepared by procedures illustrated in the accompanying schemes.


METHODS OF PREPARATION

- [0039]** The compounds of the present invention may be prepared by methods such
20 as those illustrated in the following Scheme 1-3. Solvents, temperatures, pressures, and other reaction conditions may readily be selected by one of ordinary skill in the art. Starting materials are either commercially available or can be readily prepared by one of ordinary skill in the art using known methods. For all of the schemes and compounds described below, B, R¹, R², R³ and x are as described for a compound of
25 formula I.
- [0040]** The following are the definitions of symbols used throughout Schemes 1 to 3:
- PG suitable nitrogen or oxygen protecting group, exemplified by benzyl, methoxymethyl- [MOM], benzyloxymethyl- [BOM], 2-
30 (trimethylsilyl)ethoxymethyl- [SEM], methoxyethoxymethyl- [MEM], or *t*-butyl groups;

EE leaving group exemplified by halogen (F, Cl, Br, I) and sulfonates (-OSO₂-aryl (e.g., -OSO₂Ph or -OSO₂PhCH₃), or -OSO₂-alkyl (e.g., -OSO₂CH₃ or -OSO₂CF₃));

5

SCHEME 1



[0041] Compounds of formula **II** are either commercially available or available by means known to one skilled in the art. Compounds of formula **III** can be prepared by reacting compounds of formula **II** with an appropriately protected hydrazine such as benzylhydrazine. Exemplary nitrogen protecting groups and methods of protecting the nitrogen are similar to those for protecting amines, such as those described in T. W.

Greene and P. G. M. Wuts, *Protecting Groups in Organic Synthesis*, John Wiley & Sons, Inc, New York, 1991. Preferred nitrogen protecting groups are benzyl, *tert*-butyl, methoxymethyl (MOM) groups.

5 [0042] Compounds of formula **IV** can be prepared by reacting compounds of formula **III** with a chlorinating agent such as POCl₃, either neat or in an inert solvent such as toluene at elevated temperature.

[0043] Compounds of formula **V** can be prepared by the reactions of compounds of formula **IV** with activated R², such as activated by boronic acids, tin, Grignard reagents, Zinc, Cu, etc in the presence of an appropriate catalyst if needed such as
10 Pd(dppf)Cl₂ complex with CH₂Cl₂.

[0044] Compounds of formula **VI** can be prepared by removing the protecting group (PG) in compound **V** under catalytic hydrogenation (for benzyl), or Lewis acid (e.g. AlCl₃ for benzyl or MOM), acidic (e.g. TFA for *t*-butyl or Boc) conditions.

[0045] Compounds of formula **VII**, where EE = Cl, can be prepared by reacting
15 compounds of formula **VI** with a chlorinating agent such as POCl₃, either neat or in an inert solvent such as toluene at elevated temperature.


[0046] Compounds of formula **VIII** can be prepared by reacting compounds of formula **VII** with hydrazine in an inert solvent such as pyridine at elevated temperature.

20 [0047] Compounds of formula **IX** can be prepared by reacting compounds of formula **VIII** with a carbonylating agent such as carbonyldiimidazole, phosgene, triphosgene or urea in an inert solvent such as tetrahydrofuran.

[0048] Compounds of formula **Ia** can be prepared by reacting compounds of formula **IX** with R³-X (X is a leaving group such as Cl, Br, I, -OMs, -OTs), or
25 epoxides at elevated temperatures. **Ia** can also be prepared by reacting compounds of formula **IX** with R³-OH under Mitsunobu conditions.

[0049] Compounds of formula **Ib** can be prepared by reacting compounds of formula **VIII** with R³COCO₂R (R³ substituted 2-oxoacetic acid, or its ester, which are either commercially available or available by means known to one skilled in the art) in
30 an acidic solvent, such as acetic acid at elevated temperature.

SCHEME 2



[0050] Compounds of formula **X** can be prepared by reacting compounds of formula **II** with a hydrazine mineral acid salt, such as hydrazine dihydrochloride in an aqueous or alcoholic solvent at elevated temperature.

[0051] Compounds of formula **XI**, where EE = Cl, can be prepared by reacting compounds of formula **X** with a chlorinating agent such as POCl₃, either neat or in an inert solvent such as toluene at elevated temperature.

[0052] Compounds of formula **XII** can be prepared by reacting compounds of formula **XI** with hydrazine in an inert solvent such as pyridine at elevated temperature.

[0053] Compounds of formula **XIIIa** can be prepared by reacting compounds of formula **XII** with a carbonylating agent such as carbonyldiimidazole, phosgene, triphosgene or urea in an inert solvent such as tetrahydrofuran.

[0054] Compounds of formula **XIV** can be prepared by reacting compounds of formula **XIIIa** with R³-X (X is a leaving group such as Cl, Br, I, -OMs, -OTs), or

epoxides at elevated temperatures. Compounds of formula **XIV** can also be prepared by reacting compounds of formula **XIIIa** with R^3 -OH under Mitsunobu conditions.


[0055] Compounds of formula **Ia** can be prepared by the reactions of compounds of formula **XIV** with activated R^2 , such as activated by boronic acids, tin, Grignard reagents, Zinc, Cu, etc in the presence of an appropriate catalyst if needed such as Pd(dppf)Cl₂ complex with CH₂Cl₂.

[0056] Compounds of formula **XIIIb** can be prepared by reacting compounds of formula **XII** with R^3 COCO₂R (R^3 substituted 2-oxoacetic acid or its ester, which are either commercially available or available by means known to one skilled in the art) in an acidic solvent, such as acetic acid at elevated temperature.

[0057] Compounds of formula **Ib** can be prepared by the reactions of compounds of formula **XIIIb** with activated R^2 , such as activated by boronic acids, tin, Grignard reagents, Zinc, Cu, etc in the presence of an appropriate catalyst if needed such as Pd(dppf)Cl₂ complex with CH₂Cl₂.

15

SCHEME 3



[0058] Compounds of formula **XV** are either commercially available or available by means known to one skilled in the art. Compounds of formula **XVI** can be

5 prepared by reacting compounds of formula **XV** with a hydrazine hydrate in a dipolar protic solvent, such as ethanol at elevated temperature.

[0059] Compounds of formula **XVII**, where EE' = Cl, can be prepared by reacting compounds of formula **XVI** with a chlorinating agent such as POCl₃, either neat or in an inert solvent such as toluene at elevated temperature.

10 [0060] Compounds of formula **XVIII** can be prepared by reacting compounds of formula **XVII** with hydrazine in an inert solvent such as pyridine at elevated temperature.

[0061] Compounds of formula **XIXa** can be prepared by reacting compounds of formula **XVIII** with a carbonylating agent such as carbonyldiimidazole, phosgene, 15 triphosgene or urea in an inert solvent such as tetrahydrofuran.

[0062] Compounds of formula **XX** can be prepared by reacting compounds of formula **XIXa** with R^3-X (X is a leaving group such as Cl, Br, I, -OMs, -OTs), or epoxides at elevated temperatures. Compounds of formula **XIV** can also be prepared by reacting compounds of formula **XIIIa** with R^3-OH under Mitsunobu conditions.

- 5 [0063] Compounds of formula **Ia** can be prepared by the reactions of compounds of formula **XX** with activated R^1 , such as activated by boronic acids, tin, Grignard reagents, Zinc, Cu, etc in the presence of an appropriate catalyst if needed such as $Pd(dppf)Cl_2$ complex with CH_2Cl_2 . Compounds of formula **Ia** may also be prepared from a compound of formula **XX** via displacement of the leaving group (EE) by the
- 10 conjugate base of a compound R^1-H , wherein R^1 is as previously defined, using a base in an inert solvent. Exemplary bases include sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, or alkyl lithiums.

- [0064] Compounds of formula **XIXb** can be prepared by reacting compounds of formula **XVIII** with R^3COCO_2R (R^3 substituted 2-oxoacetic acid, or its ester, which
- 15 are either commercially available or available by means known to one skilled in the art) in an acidic solvent, such as acetic acid at elevated temperature.

- [0065] Compounds of formula **Ib** can be prepared by the reactions of compounds of formula **XIXb** with activated R^1 , such as activated by boronic acids, tin, Grignard reagents, Zinc, Cu, etc in the presence of an appropriate catalyst if needed such as
- 20 $Pd(dppf)Cl_2$ complex with CH_2Cl_2 . Compounds of formula **Ib** may also be prepared from a compound of formula **XIXb** via displacement of the leaving group (EE) by the conjugate base of a compound R^1-H , wherein R^1 is as previously defined, using a base in an inert solvent. Exemplary bases include sodium carbonate, potassium carbonate, cesium carbonate, sodium hydride, potassium hydride, or alkyl lithiums.

25

EXAMPLES

[0066] The following examples serve to better illustrate, but not limit, some of the preferred embodiments of the invention.

Analytical HPLC and HPLC/MS Methods Employed in Characterization of Examples

[0067] Analytical HPLC was performed on Shimadzu LC10AS liquid chromatographs. Analytical HPLC/MS was performed on Shimadzu LC10AS liquid chromatographs and Waters ZMD Mass Spectrometers using the following methods:

Method A. Linear gradient of 0 to 100% solvent B over 4.0 min, with 1 min hold at 100% B;

UV visualization at 220 nm

10 Column: Phenomenex Lura C18, 4.6 X 50 mm

Flow rate: 4 ml/min

Solvent A: 0.1% trifluoroacetic acid, 90% water, 10% methanol

Solvent B: 0.1% trifluoroacetic acid, 10% water, 90% methanol.

15 **Method B.** Linear gradient of 0 to 100% solvent B over 8.0 min, with 3 min hold at 100% B;

UV visualization at 220 nm

Column: Zorbax SB C18, 4.6 X 75 mm

Flow rate: 2.5 ml/min

20 Solvent A: 0.2% phosphoric acid, 90% water, 10% methanol

Solvent B: 0.2% phosphoric acid, 10% water, 90% methanol.

Method C. Linear gradient of 0 to 100% solvent B over 4.0 min, with 1 min hold at 100% B;

25 UV visualization at 220 nm

Column: Phenomenex Lura C18, 4.6 X 50 mm

Flow rate: 4 ml/min



Solvent A: 10 mM NH₄OAc, 90% water, 10% methanol

Solvent B: 10 mM NH₄OAc, 10% water, 90% methanol.

30

NMR Employed in Characterization of Examples


[0068] ^1H NMR spectra were obtained with Bruker or JOEL fourier transform spectrometers operating at the following frequencies: ^1H NMR: 400 MHz (Bruker), 400 MHz (JOEL), or 500 MHz (JOEL); ^{13}C NMR: 100 MHz (Bruker), 100 MHz (JOEL) or 125 MHz (JOEL). Spectra data are reported as Chemical shift (multiplicity, coupling constants in Hz, number of hydrogens) and are reported in ppm(δ units) relative to either an internal standard (tetramethylsilane = 0 ppm) for ^1H NMR spectra, or are referenced to the residual solvent peak (2.49 ppm for $\text{CD}_2\text{HSOCD}_3$, 3.30 ppm for CD_2HOD , 7.24 ppm for CHCl_3 , 39.7 ppm for CD_3SOCD_3 , 49.0 ppm for CD_3OD , 77.0 ppm for CDCl_3). All ^{13}C NMR spectra were proton decoupled.

EXAMPLE 1**Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one****1A. Preparation of 2-benzyl-5-(4-chlorophenyl)-6-hydroxypyridazin-3(2*H*)-one**

[0069] To a suspension of 3-(4-chlorophenyl)-2,5-furandione (10.24 g, 66.9 mmol) in acetic acid (300 mL) was added benzylhydrazine dihydrochloride in one portion (19.6 g, 100 mmol). The reaction mixture was stirred at reflux for 14 h. Analysis by HPLC/MS indicated the reaction was complete. After cooling to room temperature, the resulting white suspension was filtered. The collected solid was washed with EtOAc (20 mL x 2), and dried in a 50°C vacuum oven for 16 h to yield


6.5 g of the desired product as a white solid. The filtrate was concentrated under reduced pressure, and the obtained residue was crystallized in EtOAc to afford additional 1.6 g of the title compound (8.1 g in total, 39% yield). LC/MS (method A): retention time = 3.30 min, $(M+H)^+ = 313$. 1H NMR (DMSO- d_6 , 500 MHz): δ 11.62 (s, 1H), 7.65 (d, $J = 10$ Hz, 2H), 7.51 (d, $J = 10$ Hz, 2H), 7.23-7.38 (m, 5H), 7.04 (s, 1H), 5.11 (s, 2H).

1B. Preparation of 2-benzyl-6-chloro-5-(4-chlorophenyl)pyridazin-3(2H)-one




10 **[0070]** A suspension of 2-benzyl-5-(4-chlorophenyl)-6-hydroxypyridazin-3(2H)-one (8.5 g, 27.2 mmol) in phosphorus oxychloride (50 mL) was stirred at 95°C for 1 h. Analysis by HPLC/MS indicated the starting material had been consumed. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure. The obtained residue was diluted with EtOAc. The resulting EtOAc solution
15 was washed with saturated aqueous $NaHCO_3$, saturated aqueous NaCl, dried (Na_2SO_4), filtered and concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (330 g) eluting with a gradient of EtOAc (0-40%) in hexanes to afford 6.2 g (69%) of the title compound as a white solid. LC/MS (method A): retention time = 3.81 min, $(M+H)^+ = 331$. 1H NMR ($CDCl_3$, 400 MHz): δ
20 7.27-7.80 (m, 9H), 6.88 (s, 1H), 5.36 (s, 2H).

1C. Preparation of 2-benzyl-6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3(2H)-one




[0071] To a flame-dried round bottom flask was placed 2-benzyl-6-chloro-5-(4-chlorophenyl)pyridazin-3(2H)-one (6.0 g, 18.1 mmol), 2-chlorophenylboronic acid (5.7 g, 36.2 mmol), K_3PO_4 (7.7 g, 36.2 mmol), and [1,1'-bis(diphenylphosphino)-ferrocene]dichloropalladium(II), complex with CH_2Cl_2 (1:1) (2.96 g, 3.6 mmol). The flask was purged with argon for 15 min before degassed THF (120 mL) was added. The reaction mixture was stirred in a 70°C oil bath under argon for 16 h. After cooling to room temperature, the reaction mixture was diluted with EtOAc (200 mL), washed with 1N aqueous NaOH (50 mL), H_2O , saturated aqueous NaCl, dried (Na_2SO_4), filtered and concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (330 g) eluting with a gradient of EtOAc (0-40%) in hexanes to afford 6.5 g (88%) of the title compound as a white solid. LC/MS (method A): retention time = 3.97 min, $(M+H)^+ = 407$. 1H NMR ($CDCl_3$, 400 MHz): δ 7.51 (d, 2H, $J = 8.0$ Hz), 7.24-7.40 (m, 9H), 6.95-7.03 (m, 3H), 5.33 (s, 2H).


15 **1D. Preparation of 6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3(2H)-one**



[0072] To a solution of 2-benzyl-6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3(2H)-one (6.4 g, 15.7 mmol) in toluene (100 mL) at 90°C was added aluminum chloride (5.23 g, 39.3 mmol) in one portion. After addition, the reaction mixture was stirred at 90°C for 20 min. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to reduce the volume to about 50 mL, then water (150 mL) was added. The resulting mixture was sonicated for 5 min, then stirred at room temperature for 10 min. The resulting suspension was filtered, the collected solid washed with water (20 mL x 3), hexanes (20 mL x 3) and dried in a 50°C vacuum oven for 16 h to afford 4.2 g (76%) of the title compound as a white solid. LC/MS (method A): retention time = 3.27 min, $(M+H)^+ = 317$. 1H NMR ($CDCl_3$, 400 MHz): δ 11.6 (s, 1H), 7.17-7.40 (m, 6H), 6.90-7.10 (m, 3H).

1E. Preparation of 6-chloro-3-(2-chlorophenyl)-4-(4-chlorophenyl)pyridazine

[0073] A suspension of 6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3(2*H*)-one (4.1 g, 12.9 mmol) in phosphorus oxychloride (15 mL) was stirred at 95°C for 1 h. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure. The residue was partitioned between EtOAc and saturated aqueous NaHCO₃. The separated aqueous layer was extracted with EtOAc (40 mL x 2). The combined organic was washed with saturated aqueous NaHCO₃, saturated aqueous NaCl, dried (Na₂SO₄) and concentrated under reduced pressure to afford 4.2 g (97%) of the title compound as a pink foam. LC/MS (method A): retention time = 3.50 min, (M+H)⁺ = 335.


1F. Preparation of 1-(6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3)hydrazine

15

[0074] To a solution of 6-chloro-3-(2-chlorophenyl)-4-(4-chlorophenyl)pyridazine (2.1 g, 6.26 mmol) in pyridine (20 mL) at 90°C was added anhydrous hydrazine (1 mL), and the resulting mixture was stirred at 90°C for 1 h. Analysis by HPLC/MS indicated the reaction was not complete. Additional hydrazine (0.5 mL) was added, and the reaction continued at 90°C for 1 h more. After cooling to room temperature, the reaction was concentrated under reduced pressure. The obtained residue was treated with water (50 mL), and the resulting suspension stirred at room temperature for 10 min, then filtered. The collected solid was washed with water (20 mL x 3),


dried in a 50°C vacuum oven for 16 h to afford 2.1 g (100%) of the title compound as a pink solid. LC/MS (method A): retention time = 2.67 min, $(M+H)^+ = 331$.

1G. Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-
5 *b*]pyridazin-3(2*H*)-one




[0075] To a suspension of 1,1'-carbonyl diimidazole (2.84 g, 17.5 mmol) in anhydrous THF (15 mL) at 65°C was added a suspension of 1-(6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3)hydrazine (1.16 g, 3.5 mmol) in THF (20 mL). After
10 addition, the reaction mixture was stirred at 65°C for 20 min. Analysis by HPLC/MS indicated the starting material had been consumed. After cooling to room temperature, the reaction was concentrated under reduced pressure. The obtained residue was dissolved in EtOAc (20 mL), and the resulting solution was adjusted to pH 6-8 by slow addition of 1N aqueous HCl. The resulting yellow suspension was filtered. The
15 collected solid was washed with water (20 mL x 2), EtOAc (10 mL x 2), and dried in air to obtain 850 mg of the title compound as a yellow solid. The filtrate was extracted with EtOAc (20 mL x 2). The combined EtOAc extracts were washed with saturated aqueous NaCl, dried (Na_2SO_4), filtered and concentrated under reduced pressure. The obtained residue was triturated with EtOAc (10 mL), and the resulting yellow
20 precipitate was collected by filtration. The collected solid was washed with EtOAc/hexanes (1:2) (10 mL x 2) and dried in air to afford additional 150 mg of the title compound (1.0 g in total, 80% yield). LC/MS (method A): retention time = 3.24 min, $(M+H)^+ = 357$. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz): δ 12.92 (s, 1H), 7.92 (s, 1H), 7.62-7.70 (m, 1H), 7.35-7.50 (m, 3H), 7.33 (d, $J = 8.0$ Hz, 2H), 7.20 (d, $J = 8.0$ Hz,
25 2H).


EXAMPLE 2

Preparation of 2-(4-(trifluoromethyl)benzyl)-6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one

- 5 **[0076]** To a solution of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-
b]pyridazin-3(2H)-one (18 mg, 0.05 mmol) and 4-trifluoromethyl benzylbromide (18
mg, 0.075 mmol) in anhydrous DMF (0.5 mL)) at 70°C was added solid K₂CO₃ (13.8
mg, 0.1 mmol). After addition, the reaction mixture was stirred at 70°C for 30 min.
Analysis by HPLC/MS indicated the starting material had been consumed. After
10 cooling to room temperature, the reaction was diluted with water, and extracted with
EtOAc (5 mL x 3). The combined EtOAc extracts were washed with saturated
aqueous NaCl, dried (Na₂SO₄), filtered and concentrated under reduced pressure. The
obtained residue was purified using reverse phase preparative HPLC (Conditions:
Phenomenex Luna 5μ C18 21.2 x 100 mm; Eluted with 70% to 100% B, 8 min
15 gradient, hold at 100% B for 6 min (A = 90% H₂O, 10% MeOH and B = 10% H₂O,
90% MeOH); Flow rate at 20 mL/min, UV detection at 220 nm) to afford 18 mg
(70%) of the title compound as a yellow powder. LC/MS (method A): retention time =
4.05 min, (M+H)⁺ = 515. ¹H NMR (CDCl₃, 400 MHz): δ 7.57-7.70 (m, 4H), 7.42-
7.48 (m, 1H), 7.41 (s, 1H), 7.30-7.38 (m, 2H), 7.17-7.25 (m, 3H), 7.02 (d, J = 8.0 Hz,
20 2H), 5.30 (s, 2H).

EXAMPLE 3**Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**

- 5 [0077] The title compound was prepared from 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (79 mg, 0.22 mmol) and 5-(chloromethyl)-2-(trifluoromethyl)pyridine (74 mg, 0.38 mmol) according to the procedures described for 2-(4-(trifluoromethyl)benzyl)-6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (Example 2). The crude product was purified using a silica gel cartridge (12 g) eluting with a gradient of EtOAc (0-80%) in hexanes to obtain the desired product as a yellow foam, which was lyophilized in acetonitrile to afford 87 mg (76%) of the title compound as a yellow powder. LC/MS (method A): retention time = 3.76 min, $(M+H)^+ = 516$. 1H NMR (CDCl₃, 400 MHz): δ 8.86 (s, 1H), 8.02 (d, J = 8.3 Hz, 1H), 7.47 (m, 1H), 7.42 (s, 1H), 7.33 (m, 2H), 7.25 (m, 2H), 7.20 (d, J = 8.8 Hz, 2H), 7.02 (d, J = 8.8 Hz, 2H), 5.34 (s, 2H).


EXAMPLE 4**Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-2-cyclohexyl-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**

[0078] A suspension of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one (36 mg, 0.1 mmol), bromocyclohexane (40 μ L, 0.3 mmol) and solid K_2CO_3 (42 mg, 0.3 mmol) in anhydrous DMF (1 mL) was stirred at 75°C for 17 h. Analysis by HPLC/MS indicated the reaction was not
5 complete. Additional bromocyclohexane (50 μ L, 0.375 mmol) was added, the reaction was allowed to continue at 75°C for 24 h more. After cooling to room temperature, the reaction mixture was partitioned between water (20 mL) and EtOAc (20 mL). The separated EtOAc layer was washed with saturated aqueous NaCl, dried (Na_2SO_4) and concentrated under reduced pressure. The crude product was purified using a silica gel
10 cartridge (12 g) eluting with a gradient of EtOAc (0-60%) in hexanes to yield a yellow glassy product, which was lyophilized in acetonitrile to afford 15 mg (34%) of the title compound as a yellow powder. LC/MS (method A): retention time = 4.06 min, $(M+H)^+ = 439$. 1H NMR ($CDCl_3$, 400 MHz): δ 7.18-7.50 (m, 5H), 7.19 (d, $J = 8.8$ Hz, 2H), 7.03 (d, $J = 8.8$ Hz, 2H), 4.47 (m, 1H), 1.26-2.04 (m, 10H).

15

EXAMPLE 5

Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-2-cyclohexyl-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one




20 [0079] A suspension of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one (20 mg, 0.056 mmol), 2,2,2-trifluoroethyl trifluoromethane sulfonate (16 mg, 0.073 mmol) and cesium carbonate (20 mg, 0.056 mmol) in anhydrous DMF (0.6 mL) was stirred at room temperature for 1 h. Analysis by HPLC/MS indicated the reaction was complete. The reaction mixture was
25 partitioned between water and EtOAc. The separated EtOAc layer was washed with water, saturated aqueous NaCl, dried (Na_2SO_4) and concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (4 g) eluting with

a gradient of EtOAc (0-25%) in hexanes to afford 20 mg (81%) of the title compound as a yellow solid. LC/MS (method A): retention time = 3.66 min, $(M+H)^+ = 439$. 1H NMR ($CDCl_3$, 400 MHz): δ 7.47 (m, 2H), 7.35 (m, 2H), 7.25 (m, 1H), 7.21 (d, $J = 8.8$ Hz, 2H), 7.05 (d, $J = 8.8$ Hz, 2H), 4.68 (m, 2H).

5

EXAMPLE 6


Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-2-(3,3,3-trifluoropropyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one



- 10 **[0080]** A suspension of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2H)-one (36 mg, 0.1 mmol), 1,1,1-trifluoro-3-iodopropane (67 mg, 0.3 mmol) and solid K_2CO_3 (42 mg, 0.3 mmol) in anhydrous DMF (0.6 mL) was stirred at room temperature for 2 h. Analysis by HPLC/MS indicated the reaction was not complete. Additional 1,1,1-trifluoro-3-iodopropane (67
- 15 mg, 0.3 mmol) was added, the reaction was allowed to continue at room temperature for 16 h, then at 45°C for 30 min. After cooling to room temperature, the reaction mixture was partitioned between water and EtOAc. The separated EtOAc layer was washed with water, saturated aqueous NaCl, dried (Na_2SO_4) and concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (12 g)
- 20 eluting with a gradient of EtOAc (0-60%) in hexanes to yield a yellow oily product, which was lyophilized in acetonitrile to afford 38 mg (84%) of the title compound as a yellow powder. LC/MS (method A): retention time = 3.69 min, $(M+H)^+ = 453$. 1H NMR ($CDCl_3$, 400 MHz): δ 7.44 (m, 2H), 7.35 (m, 2H), 7.25 (m, 1H), 7.21 (d, $J = 8.8$ Hz, 2H), 7.05 (d, $J = 8.8$ Hz, 2H), 4.36 (m, 2H), 2.76 (m, 2H).


25

EXAMPLE 7


Preparation of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-2-(4,4,4-trifluorobutyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one

- 5 **[0081]** A suspension of 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one (20 mg, 0.056 mmol), 1,1,1-trifluoro-4-iodobutane (20 mg, 0.084 mmol) and solid K₂CO₃ (12 mg, 0.084 mmol) in anhydrous DMF (0.6 mL) was stirred at 80°C for 1 h. Analysis by HPLC/MS indicated the reaction was not complete. Additional 1,1,1-trifluoro-4-iodobutane (12 mg, 0.084
- 10 mmol) was added, the reaction was allowed to continue at 80°C for 30 min. After cooling to room temperature, the reaction mixture was diluted with water, and extracted with EtOAc (20 mL x 2). The combined EtOAc extracts were washed with water, saturated aqueous NaCl, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (12 g) eluting
- 15 with a gradient of EtOAc (0-60%) in hexanes to yield a yellow oily product, which was lyophilized in acetonitrile to afford 19 mg (73%) of the title compound as a yellow powder. HPLC (method B): retention time = 7.70 min. ¹H NMR (CDCl₃, 400 MHz): δ 7.45 (m, 2H), 7.35 (m, 2H), 7.25 (m, 1H), 7.20 (d, J = 8.8 Hz, 2H), 7.04 (d, J = 8.8 Hz, 2H), 4.17 (t, J = 6.1 Hz, 2H), 2.17-2.25 (m, 4H).

20


EXAMPLE 8**Preparation of 6,7-bis(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**

5

8A. Preparation of 4-(4-chlorophenyl)-1,2-dihydropyridazine-3,6-dione

[0082] To a solution of hydrazine dihydrochloride (5.6 g, 53.3 mmol) in H₂O (60 mL) at 120°C was added 3-(4-chlorophenyl)-2,5-furandione (10 g, 48 mmol) in one portion. After addition, the reaction mixture was stirred at 120°C for 40 min. After cooling to room temperature, the reaction became a pale yellow suspension. The precipitate was collected by filtration, washed with water (30 mL x 3), hexanes (30 mL x 3), and dried in a 50°C vacuum oven for 16 h to afford 8.7 g (81%) of the title compound as a pale yellow solid. LC/MS (method A): retention time = 2.57 min, (M+H)⁺ = 223.

15


8B. Preparation of 3,6-dichloro-4-(4-chlorophenyl)pyridazine

[0083] A suspension of 4-(4-chlorophenyl)-1,2-dihydropyridazine-3,6-dione (7.9 g, 35.5 mmol) in phosphorus oxychloride (25 mL) was heated at 120°C for 30 min,

20

then allowed to cool to room temperature. After removing most of the remaining phosphorus oxychloride under reduced pressure, icy water (200 mL) was carefully added to the residue. The resulting mixture was extracted with EtOAc (150 mL x 2), and the combined EtOAc extracts washed with saturated aqueous NaHCO₃, saturated aqueous NaCl, dried (Na₂SO₄) and concentrated to afford 7.1 g (77%) of the title compound as an off-white solid. LC/MS (method A): retention time = 3.16 min, (M+H)⁺ = 259.

8C. Preparation of 1-(6-chloro-5-(4-chlorophenyl)pyridazin-3-yl)hydrazine




10

[0084] Anhydrous hydrazine (8 mL) was added to a solution of 3,6-dichloro-4-(4-chlorophenyl)pyridazine (7.1 g, 27.4 mmol) in pyridine (30 mL) at 100°C. The resulting mixture was stirred at 100°C for 30 min. Analysis by HPLC/MS indicated the reaction was complete. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to remove most of the pyridine. To the residue was added isopropanol (200 mL), the resulting suspension was refluxed for 20 min. After cooling to room temperature, the suspension was filtered. The collected solid was washed with isopropanol, dried in a 50°C vacuum oven overnight to afford 4.0 g (57%) of the title compound as an off-white solid. LC/MS (method A): retention time = 2.24 min, (M+H)⁺ = 255.

20

8D. Preparation of 6-chloro-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one




[0085] The title compound (900 mg, 80%) as a yellow solid was prepared from 1-(6-chloro-5-(4-chlorophenyl)pyridazin-3-yl)hydrazine (1.02 g, 4.0 mmol) and 1,1'-

25


carbonyl diimidazole (3.24 g, 20.0 mmol) according to the procedures described for 6-(2-chlorophenyl)-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one (Example 1G). LC/MS (method A): retention time = 2.77 min, (M+H)⁺ = 281.

5 **8E. Preparation of 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one**



[0086] A suspension of 6-chloro-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-
10 *b*]pyridazin-3(2*H*)-one (843 mg, 3 mmol), 5-chloromethyl-2-(trifluoromethyl)pyridine (704 mg, 3.6 mmol) and solid K₂CO₃ (830 mg, 6 mmol) in anhydrous DMF (10 mL) was stirred at 70 °C for 20 min. Analysis by HPLC/MS indicated the reaction was complete. After cooling to room temperature, the reaction was diluted with water and extracted with EtOAc (40 mL x 3). The combined EtOAc extracts were washed with
15 saturated aqueous NaCl, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified by trituration with CH₂Cl₂-hexanes to afford 1.10 g (84%) of the title compound as a yellow solid. LC/MS (method A): retention time = 3.50 min, (M+H)⁺ = 440.


20 **8F. Preparation of 6,7-bis(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-*b*]pyridazin-3(2*H*)-one**




[0087] To a flame-dried round bottom flask was placed 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (44 mg, 0.1 mmol), 4-chlorophenylboronic acid (31 mg, 0.2 mmol), K₃PO₄ (42.5 mg, 0.2 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with CH₂Cl₂ (1:1) (16 mg, 0.02 mmol). The flask was purged with argon for 15 min before addition of degassed anhydrous THF (1 mL). The reaction mixture was stirred at 70°C under argon for 16 h. After cooling to room temperature, the reaction mixture was diluted with EtOAc (20 mL), washed with 1 N aqueous NaOH, water, saturated aqueous NaCl, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified using reverse phase preparative HPLC (Conditions: Phenomenex Luna 5µ C18 21.2 x 100 mm; Eluted with 60% to 100% B, 8 min gradient, hold at 100% B for 6 min (A = 90% H₂O, 10% MeOH, 0.1% trifluoroacetic acid and B = 10% H₂O, 90% MeOH, 0.1% trifluoroacetic acid); Flow rate at 20 mL/min, UV detection at 220 nm). The desired fractions were concentrated under reduced pressure. The residue was dissolved in EtOAc, washed with saturated aqueous NaHCO₃ once, water twice, then saturated aqueous NaCl, dried over Na₂SO₄ and evaporated. The yellow residue was dissolved in acetonitrile, frozen and lyophilized to afford 18.2 mg (35%) of the title compound as a yellow powder. LC/MS (method A): retention time = 3.94 min, (M+H)⁺ = 516. ¹H NMR (CDCl₃, 400 MHz): δ 8.83 (d, J=1.71 Hz, 1H), 7.98 (dd, J=7.95, 2.08 Hz, 1H), 7.69 (d, J=8.07 Hz, 1H), 7.41 (s, 1H), 7.25-7.32 (m, 4H), 7.20-7.23 (m, 2H), 7.00-7.04 (m, 2H), 5.33 (s, 2H).

EXAMPLE 9

Preparation of 7-(4-chlorophenyl)-6-phenyl-2-((6-(trifluoromethyl) pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one




- 5 **[0088]** To a flame-dried tube was placed 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (44 mg, 0.1 mmol), phenylboronic acid (24.4 mg, 0.2 mmol), K_3PO_4 (42.5 mg, 0.2 mmol) and [1,1'-bis(diphenyl-phosphino) ferrocene]dichloropalladium(II), complex with CH_2Cl_2 (1:1) (16 mg, 0.02 mmol). The vessel was purged with argon for 10 min
- 10 before addition of degassed anhydrous THF (1 mL), and sealed. The reaction mixture was stirred at 75°C for 16 h. After cooling to room temperature, the reaction mixture was diluted with EtOAc (20 mL), washed with 1 N aqueous NaOH, water, saturated aqueous NaCl, dried (Na_2SO_4) and concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (12 g) eluting with a gradient of
- 15 EtOAc (0-60%) in hexanes to obtain 23 mg of the desired product as an oil. The oily residue was triturated with CH_2Cl_2 -hexanes to afford 20 mg (42%) of the title compound as a yellow powder. LC/MS (method A): retention time = 3.76 min, $(M+H)^+ = 482$. 1H NMR ($CDCl_3$, 400 MHz): δ 8.84 (d, $J=1.71$ Hz, 1H), 7.99 (dd, $J=8.07$, 1.96 Hz, 1H), 7.68 (d, $J=8.07$ Hz, 1H), 7.41 (s, 1H), 7.33-7.38 (m, 1H), 7.24-
- 20 7.30 (m, 6H), 6.99-7.06 (m, 2H), 5.33 (s, 2H).

EXAMPLE 10**Preparation of 7-(4-chlorophenyl)-6-(2-fluoropyridin-3-yl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**

- 5 **[0089]** The title compound (41 mg, 42%) as a yellow solid was prepared from 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (88 mg, 0.2 mmol) and 2-fluoropyridin-3-ylboronic acid (56 mg, 0.4 mmol) according to the procedures described for 7-(4-chlorophenyl)-6-phenyl-2-((6-(trifluoromethyl) pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (Example 9). LC/MS (method A): retention time= 3.38 min, $(M+H)^+ = 501$. 1H NMR ($CDCl_3$, 400 MHz): δ 8.84 (s, 1H), 8.14 (d, $J=1.96$ Hz, 1H), 7.97-8.01 (m, 1H), 7.68-7.76 (m, 2H), 7.46 (s, 1H), 7.34 (d, $J=8.56$ Hz, 2H), 7.04 (d, $J=8.56$ Hz, 2H), 6.89 (dd, $J=8.56$, 2.93 Hz, 1H), 5.34 (s, 2H).

15

EXAMPLE 11**Preparation of 7-(4-chlorophenyl)-6-(2,4-dichlorophenyl-3-yl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**


- 20 **[0090]** The title compound was prepared from 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (44 mg, 0.1 mmol) and 2,4-dichlorophenylboronic acid (23 mg, 0.12 mmol) according to

the procedures described for 7-(4-chlorophenyl)-6-phenyl-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (Example 9). The crude product was purified using reverse phase preparative HPLC (Conditions: Phenomenex Luna 5 μ C18 21.2 x 100 mm; Eluted with 65% to 100% B, 8 min gradient, hold at 100% B for 7 min (A = 90% H₂O, 10% MeOH, 0.1% trifluoroacetic acid and B = 10% H₂O, 90% MeOH, 0.1% trifluoroacetic acid); Flow rate at 20 mL/min, UV detection at 220 nm). The desired fractions were concentrated under reduced pressure. The residue was dissolved in EtOAc, washed with saturated aqueous NaHCO₃ once, water twice, then saturated aqueous NaCl, dried over Na₂SO₄ and evaporated. The yellow residue was lyophilized in acetonitrile to afford 3.5 mg (6.4%) of the title compound as a yellow powder. LC/MS (method A): retention time= 4.03 min, (M+H)⁺ = 550. ¹H NMR (CDCl₃, 400 MHz): δ 8.85 (d, J=1.71 Hz, 1H), 8.00 (dd, J=7.95, 1.83 Hz, 1H), 7.70 (d, J=8.07 Hz, 1H), 7.38-7.44 (m, 2H), 7.31-7.36 (m, 1H), 7.22-7.29 (m, 3H), 6.99-7.05 (m, 2H), 5.34 (s, 2 H).

15

EXAMPLE 12

Preparation of 7-(4-chlorophenyl)-6-(2-fluorophenyl-3-yl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one




[0091] The title compound (8.5 mg, 17%) as a yellow powder was prepared from 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (44 mg, 0.1 mmol) and 2-fluorophenylbromic acid (28 mg, 0.2 mmol) according to the procedures described for 7-(4-chlorophenyl)-6-phenyl-2-((6-(trifluoromethyl) pyridin-3-yl)methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (Example 9). LC/MS (method A): retention time= 3.66 min, (M+H)⁺ = 500. ¹H NMR (CDCl₃, 400 MHz): δ 8.84 (s, 1 H), 7.95-


25

8.01 (m, 1H), 7.69 (d, $J=8.07$ Hz, 1H), 7.51 (td, $J=7.34, 1.71$ Hz, 1H), 7.40 (s, 1H), 7.35-7.40 (m, 1H), 7.18-7.27 (m, 3H), 6.98-7.05 (m, 2H), 6.87 (t, $J=9.17$ Hz, 1H), 5.32 (s, 2H).

5

EXAMPLE 13**Preparation of 2-(4-(trifluoromethyl)benzyl)-6,7-bis(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**


10 **13A. Preparation of 2-(4-(trifluoromethyl)benzyl)-6-chloro-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one**




[0092] The title compound (469 mg, 60%) was prepared from 6-chloro-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (500 mg, 1.78 mmol) and 1-chloromethyl-4-trifluoromethyl benzene (425 mg, 1.78 mmol) according to the procedures described for 6-chloro-7-(4-chlorophenyl)-2-((6-(trifluoromethyl)pyridin-3-yl) methyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (Example 8E). LC/MS (Method C): retention time = 3.91 min, (M+H)⁺ = 439.

15


13B. Preparation of 2-(4-(trifluoromethyl)benzyl)-6,7-bis(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one



- [0093]** To a suspension of 2-(4-(trifluoromethyl)benzyl)-6-chloro-7-(4-chlorophenyl)-[1,2,4]triazolo[4,3-b]pyridazin-3(2H)-one (150 mg, 0.34 mmol), 4-chlorophenylboronic acid (60 mg, 0.38 mmol), Na₂CO₃ (90 mg, 0.85 mmol) dissolved in water (0.2 mL) in toluene (3 mL) was added Pd(PPh₃)₄ (25 mg, 0.02 mmol). The resulting reaction mixture was bubbled with argon for 10 min before the flask was placed in an oil bath preheated at 125°C. After stirring at 125°C for 10 h, the reaction was then allowed to cool to room temperature. To the reaction was added water (10 mL), the aqueous mixture extracted with EtOAc (10 mL x 3). The combined EtOAc extracts were washed with water (10 mL x 2), followed by saturated aqueous NaCl (10 mL x 2) and concentrated under reduced pressure. The crude product was purified by reverse phase preparative HPLC (Conditions: Phenomenex Luna 5μ C18 21.2 x 100 mm; Eluted with 17% to 100% B, 25 min gradient, hold at 100% B for 5 min (A = 90% H₂O, 10% acetonitrile and B = 10% H₂O, 90% acetonitrile); Flow rate at 20 mL/min, UV detection at 220 nm) to obtain 22 mg (13%) the title compound as a yellow solid. LC/MS (Method C): retention time = 4.26 min; (M+H)⁺ = 515. ¹H NMR (CDCl₃, 500MHz): δ 7.61 (d, *J* = 5.0 Hz, 2H), 7.56 (d, *J* = 5.0 Hz, 2H), 7.41(s, 1H), 7.24-7.30 (m, 4H), 7.21 (d, *J* = 5.0 Hz, 2H), 7.01 (d, *J* = 5.0 Hz, 2H), 5.29 (s, 2H).


EXAMPLE 14**Preparation of 7-(2-chlorophenyl)-8-(4-chlorophenyl)-3-cyclohexyl-4H-pyridazino[6,1-c][1,2,4]triazin-4-one**

5

14A. Preparation of methyl 2-cyclohexyl-2-oxoacetate

[0094] The title compound (1.1 g, 65%) as a colorless oil was prepared from cyclohexyl-magnesium bromide and methyl chlorooxoacetate by the procedures described in F. Babudri *et al.*, *Tetrahedron*, **52** (42), pp.13512-20 (1996).

10

14B. Preparation of 7-(2-chlorophenyl)-8-(4-chlorophenyl)-3-cyclohexyl-4H-pyridazino[6,1-c][1,2,4]triazin-4-one


[0095] To a suspension of 1-(6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3-yl)hydrazine (50 mg, 0.15 mmol) in acetic acid (1 mL) was added methyl 2-cyclohexyl-2-oxoacetate (51 mg, 0.3 mmol). The reaction mixture was heated at 120°C for 1 h. After cooling to room temperature, the reaction was concentrated under reduced pressure. The residue was diluted with EtOAc (30 mL), washed with water, saturated aqueous NaHCO₃, saturated aqueous NaCl, dried (Na₂SO₄), filtered and

20


concentrated under reduced pressure. The crude product was purified using a silica gel cartridge (12g) eluting with a gradient of EtOAc (0-40%) in hexanes to afford 50 mg (74%) of the title compound as a pale yellow solid. LC/MS (method A): retention time = 3.93 min, $(M+H)^+ = 451$. 1H NMR ($CDCl_3$, 400 MHz): δ 7.89 (s, 1H), 7.50-7.60 (m, 1H), 7.35-7.43 (m, 2H), 7.20-7.32 (m, 3H), 7.10 (d, 2H, $J = 12.0$ Hz), 3.30-3.43 (m, 1H), 2.07 (d, 2H, $J = 12.0$ Hz), 1.90 (d, 2H, $J = 12.0$ Hz), 1.65-1.72 (m, 3H), 1.40-1.62 (m, 3H).

EXAMPLE 15

10 **Preparation of 3-(4-(trifluoromethyl)benzyl)-7-(2-chlorophenyl)-8-(4-chlorophenyl)-4H-pyridazino[6,1-c][1,2,4]triazin-4-one**




15A. **Preparation of (Z)-2-hydroxy-3-(4-trifluoromethyl)phenyl)acrylic acid**



[0096] The title compound (300 mg, 65%) as a pale yellow powder was prepared from 4-(trifluoromethyl)phenylacetonitrile and diethyl oxalate by the procedures described in M. Brana *et al.*, *Org. Biomol. Chem.*, **2**, pp.1864-1871, (2004).

15B. Preparation of 3-(4-(trifluoromethyl)benzyl)-7-(2-chlorophenyl)-8-(4-chlorophenyl)-4H-pyridazino[6,1-c][1,2,4]triazin-4-one



[0097] The title compound (55 mg, 70%) as a pale yellow solid was prepared from 1-(6-(2-chlorophenyl)-5-(4-chlorophenyl)pyridazin-3)hydrazine (50 mg, 0.15 mmol) and (Z)-2-hydroxy-3-(4-trifluoromethyl)phenyl)acrylic acid (52.2 mg, 0.225 mmol) according to the procedures described for 7-(2-chlorophenyl)-8-(4-chlorophenyl)-3-cyclohexyl-4H-pyridazino[6,1-c][1,2,4]triazin-4-one (Example 14B). LC/MS (method A): retention time = 3.95 min, (M+H)⁺ = 527. ¹H NMR (CDCl₃, 400 MHz): δ 7.91 (s, 1H), 7.64 (d, 2H, *J* = 8.0 Hz), 7.55 (d, 2H, *J* = 8.0 Hz), 7.48-7.53 (m, 1H), 7.35-7.45 (m, 2H), 7.21-7.32 (m, 3H), 7.09 (d, 2H, *J* = 8.0 Hz), 4.46 (s, 2H).

BIOLOGICAL EVALUATION

Cannabinoid Receptor Binding Assay

[0098] Radioligand binding studies were conducted in membranes prepared from Chinese Hamster Ovary (CHO) cells that over-express recombinant human CB-1 (CHO-CB-1 cells). Total assay volume for the binding studies was 100 μl. 5 μg of membranes were brought up to a final volume of 95 μl with Binding Buffer (25 mM HEPES, 150 mM NaCl, 2.5 mM CaCl₂, 1 mM MgCl₂, 0.25% BSA). The diluted membranes were preincubated with a compound or DMSO vehicle. The binding reaction was initiated by the addition of 2 nM final ³H-CP-55,940 (120 Ci/mmol) and proceeded for 2.5 hours at room temperature. The binding reaction was terminated by transferring the reaction to GF/B 96 well plates (presoaked with 0.3% polyethylenimine) using a Packard Cell Harvester. The filter was washed with 0.25x PBS, 30 μl MicroScint was added per well, and the bound radiolabel was quantitated by scintillation counting on a Packard TopCount Scintillation Counter. The CB-2

radioligand binding assay was conducted identically except that the membranes from CHO-CB-2 cells were used.

[0099] For a compound to be considered a CB-1 antagonist, the compound must possess a CB-1 receptor binding affinity K_i less than 13000 nM. As determined by the assay described above, the CB-1 receptor binding K_i values of working Examples 1-63 fall within the range of 0.01 nM to 10000 nM.

Cannabinoid Receptor Functional Activity Assay

[00100] Functional CB-1 inverse agonist activity of test compounds was determined in CHO-CB-1 cells using a cAMP accumulation assay. CHO-CB-1 cells were grown in 96 well plates to near confluence. On the day of the functional assay, growth medium was aspirated and 100 of Assay Buffer (PBS plus 25 mM HEPES / 0.1 mM 3-isobutyl-1-methylxanthine/ 0.1% BSA) was added. Compounds were added to the Assay buffer diluted 1:100 from 100% DMSO and allowed to preincubate for 10 minutes prior to addition of 5 uM forskolin. The mixture was allowed to proceed for 15 minutes at room temperature and was terminated by the addition of 0.1 N HCl. The total intracellular cAMP concentration was quantitated using the Amersham cAMP SPA kit.

20

UTILITIES AND COMBINATIONS

Utilities

[00101] The compounds of the present invention are cannabinoid receptor modulators, and include compounds which are, for example, selective agonists, partial agonists, inverse agonists, antagonists or partial antagonists of the cannabinoid receptor. Accordingly, the compounds of the present invention may be useful for the treatment or prevention of diseases and disorders associated with G-protein coupled cannabinoid receptor activity. Preferably, compounds of the present invention possess activity as antagonists or inverse agonists of the CB-1 receptor, and may be used in the treatment of diseases or disorders associated with the activity of the CB-1 receptor.

[00102] Accordingly, the compounds of the present invention can be administered to mammals, preferably humans, for the treatment of a variety of conditions and

disorders, including, but not limited to metabolic and eating disorders as well as conditions associated with metabolic disorders, (e.g., obesity, diabetes, arteriosclerosis, hypertension, polycystic ovary disease, cardiovascular disease, osteoarthritis, dermatological disorders, hypertension, insulin resistance, hypercholesterolemia, hypertriglyceridemia, cholelithiasis and sleep disorders, hyperlipidemic conditions, bulimia nervosa and compulsive eating disorders) or psychiatric disorders, such as substance abuse, depression, anxiety, mania and schizophrenia. These compounds could also be used for the improvement of cognitive function (e.g., the treatment of dementia, including Alzheimer's disease, short term memory loss and attention deficit disorders); neurodegenerative disorders (e.g., Parkinson's Disease, cerebral apoplexy and craniocerebral trauma) and hypotension (e.g., hemorrhagic and endotoxin-induced hypotension). These compounds could also be used for treatment of catabolism in connection with pulmonary dysfunction and ventilator dependency; treatment of cardiac dysfunction (e.g., associated with valvular disease, myocardial infarction, cardiac hypertrophy or congestive heart failure); and improvement of the overall pulmonary function; transplant rejection; rheumatoid arthritis; multiple sclerosis; inflammatory bowel disease; lupus; graft vs. host disease; T-cell mediated hypersensitivity disease; psoriasis; asthma; Hashimoto's thyroiditis; Guillain-Barre syndrome; cancer; contact dermatitis; allergic rhinitis; and ischemic or reperfusion injury.

[00103] Compounds useful in the treatment of appetitive or motivational disorders regulate desires to consume sugars, carbohydrates, alcohol or drugs and more generally to regulate the consumption of ingredients with hedonic value. In the present description and in the claims, appetitive disorders are understood as meaning: disorders associated with a substance and especially abuse of a substance and/or dependency on a substance, disorders of eating behaviors, especially those liable to cause excess weight, irrespective of its origin, for example: bulimia nervosa, craving for sugars. The present invention therefore further relates to the use of a CB-1 receptor antagonist or inverse agonist for the treatment of bulimia and obesity, including obesity associated with type II diabetes (non-insulin-dependent diabetes), or more generally any disease resulting in the patient becoming overweight. Obesity, as described herein, is defined by a body mass index (kg/m^2) of at least 26. It may be

due to any cause, whether genetic or environmental, including overeating and
bulemia, polycystic ovary disease, craniopharyngeoma, Prader-Willi Syndrome,
Frohlich's Syndrome, Type II diabetes, growth hormone deficiency, Turner's
Syndrome and other pathological states characterized by reduced metabolic activity or
5 reduced energy expenditure. As used with reference to the utilities described herein,
the term "treating" or "treatment" encompasses prevention, partial alleviation, or cure
of the disease or disorder. Further, treatment of obesity is expected to prevent
progression of medical covariants of obesity, such as arteriosclerosis, Type II diabetes,
polycystic ovary disease, cardiovascular disease, osteoarthritis, dermatological
10 disorders, hypertension, insulin resistance, hypercholesterolemia,
hypertriglyceridemia, cholelithiasis and sleep disorders.

[00104] Compounds in the present invention may also be useful in treating
substance abuse disorders, including substance dependence or abuse without
physiological dependence. Substances of abuse include alcohol, amphetamines (or
15 amphetamine-like substances), caffeine, cannabis, cocaine, hallucinogens, inhalants,
nicotine, opioids, phencyclidine (or phencyclidine-like compounds), sedative-
hypnotics or benzodiazepines, and other (or unknown) substances and combinations
of the above. The terms "substance abuse disorders" also includes drug or alcohol
withdrawal syndromes and substance-induced anxiety or mood disorder with onset
20 during withdrawal.

[00105] Compounds in the present invention may be useful in treating memory
impairment and cognitive disorders. The condition of memory impairment is
manifested by impairment of the ability to learn new information and/or the inability
to recall previously learned information. Memory impairment is a primary symptom
25 of dementia and can also be a symptom associated with such diseases as Alzheimer's
disease, schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease,
Creutzfeld-Jakob disease, HIV, cardiovascular disease, and head trauma as well as
age-related cognitive decline. Dementias are diseases that include memory loss and
additional intellectual impairment separate from memory. Cannabinoid receptor
30 modulators may also be useful in treating cognitive impairments related to attentional
deficits, such as attention deficit disorder.

[00106] Compounds in the present invention may also be useful in treating diseases associated with dysfunction of brain dopaminergic systems, such as Parkinson's Disease and substance abuse disorders. Parkinson's Disease is a neurodegenerative movement disorder characterized by bradykinesia and tremor.

5 [00107] As modulators of the cannabinoid receptor, the compounds of the present invention are further useful for the treatment and prevention of respiratory diseases and disorders. Respiratory diseases for which cannabinoid receptor modulators are useful include, but are not limited to, chronic pulmonary obstructive disorder, emphysema, asthma, and bronchitis. In addition, cannabinoid receptor modulators
10 block the activation of lung epithelial cells by moieties such as allergic agents, inflammatory cytokines or smoke, thereby limiting release of mucin, cytokines, and chemokines, or selectively inhibiting lung epithelial cell activation.

[00108] Moreover, the compounds employed in the present invention may stimulate inhibitory pathways in cells, particularly in leukocytes, lung epithelial cells,
15 or both, and are thus useful in treating such diseases. "Leukocyte activation" is defined herein as any or all of cell proliferation, cytokine production, adhesion protein expression, and production of inflammatory mediators. "Epithelial cell activation" is defined herein as the production of any or all of mucins, cytokines, chemokines, and adhesion protein expression.

20 [00109] Use of the compounds of the present invention for treating leukocyte activation-associated disorders is exemplified by, but is not limited to, treating a range of disorders such as: transplant (such as organ transplant, acute transplant, xenotransplant or heterograft or homograft (such as is employed in burn treatment)) rejection; protection from ischemic or reperfusion injury such as ischemic or
25 reperfusion injury incurred during organ transplantation, myocardial infarction, stroke or other causes; transplantation tolerance induction; arthritis (such as rheumatoid arthritis, psoriatic arthritis or osteoarthritis); multiple sclerosis; respiratory and pulmonary diseases including but not limited to chronic obstructive pulmonary disease (COPD), emphysema, bronchitis, and acute respiratory distress syndrome
30 (ARDS); inflammatory bowel disease, including ulcerative colitis and Crohn's disease; lupus (systemic lupus erythematosus); graft vs. host disease; T-cell mediated hypersensitivity diseases, including contact hypersensitivity, delayed-type

- hypersensitivity, and gluten-sensitive enteropathy (Celiac disease); psoriasis; contact dermatitis (including that due to poison ivy); Hashimoto's thyroiditis; Sjogren's syndrome; Autoimmune Hyperthyroidism, such as Graves' Disease; Addison's disease (autoimmune disease of the adrenal glands); Autoimmune polyglandular disease (also known as autoimmune polyglandular syndrome); autoimmune alopecia; pernicious anemia; vitiligo; autoimmune hypopituitarism; Guillain-Barre syndrome; other autoimmune diseases; glomerulonephritis; serum sickness; urticaria; allergic diseases such as respiratory allergies (asthma, hayfever, allergic rhinitis) or skin allergies; scleroderma; mycosis fungoides; acute inflammatory and respiratory responses (such as acute respiratory distress syndrome and ischemia/reperfusion injury); dermatomyositis; alopecia areata; chronic actinic dermatitis; eczema; Behcet's disease; Pustulosis palmoplantis; Pyoderma gangrenosum; Sezary's syndrome; atopic dermatitis; systemic sclerosis; and morphea. The term "leukocyte activation-associated" or "leukocyte-activation mediated" disease as used herein includes each of the above referenced diseases or disorders. In a particular embodiment, the compounds of the present invention are useful for treating the aforementioned exemplary disorders irrespective of their etiology. The combined activity of the present compounds towards monocytes, macrophages, T-cells, etc. may be useful in treating any of the above-mentioned disorders.
- 20 **[00110]** Cannabinoid receptors are important in the regulation of Fc gamma receptor responses of monocytes and macrophages. Compounds of the present invention inhibit the Fc gamma dependent production of TNF alpha in human monocytes/macrophages. The ability to inhibit Fc gamma receptor dependent monocyte and macrophage responses results in additional anti-inflammatory activity
- 25 for the present compounds. This activity is especially of value, for example, in treating inflammatory diseases such as arthritis or inflammatory bowel disease. In particular, the present compounds are useful for treating autoimmune glomerulonephritis and other instances of glomerulonephritis induced by deposition of immune complexes in the kidney that trigger Fc gamma receptor responses leading to
- 30 kidney damage.
- [00111]** Cannabinoid receptors are expressed on lung epithelial cells. These cells are responsible for the secretion of mucins and inflammatory cytokines/chemokines in

the lung and are thus intricately involved in the generation and progression of respiratory diseases. Cannabinoid receptor modulators regulate both the spontaneous and the stimulated production of both mucins and cytokines. Thus, such compounds are useful in treating respiratory and pulmonary diseases including, COPD, ARDS, and bronchitis.

[00112] Further, cannabinoid receptors may be expressed on gut epithelial cells and hence regulate cytokine and mucin production and may be of clinical use in treating inflammatory diseases related to the gut. Cannabinoid receptors are also expressed on lymphocytes, a subset of leukocytes. Thus, cannabinoid receptor modulators will inhibit B and T-cell activation, proliferation and differentiation. Thus, such compounds will be useful in treating autoimmune diseases that involve either antibody or cell mediated responses such as multiple sclerosis and lupus.

[00113] In addition, cannabinoid receptors regulate the Fc epsilon receptor and chemokine induced degranulation of mast cells and basophils. These play important roles in asthma, allergic rhinitis, and other allergic disease. Fc epsilon receptors are stimulated by IgE-antigen complexes. Compounds of the present invention inhibit the Fc epsilon induced degranulation responses, including the basophil cell line, RBL. The ability to inhibit Fc epsilon receptor dependent mast cell and basophil responses results in additional anti-inflammatory and anti-allergic activity for the present compounds. In particular, the present compounds are useful for treating asthma, allergic rhinitis, and other instances of allergic disease.

Combinations

[00114] The present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, a therapeutically effective amount of at least one of the compounds of formula I, alone or in combination with a pharmaceutical carrier or diluent. Optionally, compounds of the present invention can be used alone, in combination with other suitable therapeutic agents useful in the treatment of the aforementioned disorders including: anti-obesity agents; anti-diabetic agents, appetite suppressants; cholesterol/lipid-lowering agents, HDL-raising agents, cognition enhancing agents, agents used to treat neurodegeneration, agents used to treat respiratory conditions, agents used to treat bowel disorders, anti-inflammatory

agents; anti-anxiety agents; anti-depressants; anti-hypertensive agents; cardiac glycosides; and anti-tumor agents.

[00115] Such other therapeutic agent(s) may be administered prior to, simultaneously with, or following the administration of the cannabinoid receptor modulators in accordance with the invention.

[00116] Examples of suitable anti-obesity agents for use in combination with the compounds of the present invention include melanocortin receptor (MC4R) agonists, melanin-concentrating hormone receptor (MCHR) antagonists, growth hormone secretagogue receptor (GHSR) antagonists, galanin receptor modulators, orexin antagonists, CCK agonists, GLP-1 agonists, and other Pre-proglucagon-derived peptides; NPY1 or NPY5 antagonists, NPY2 and NPY4 modulators, corticotropin releasing factor agonists, histamine receptor-3 (H3) modulators, α 2 inhibitors, PPAR gamma modulators, PPAR delta modulators, acetyl-CoA carboxylase (ACC) inhibitors, 11- β -HSD-1 inhibitors, adipopectin receptor modulators; beta 3 adrenergic agonists, such as AJ9677 (Takeda/Dainippon), L750355 (Merck), or CP331648 (Pfizer) or other known beta 3 agonists as disclosed in U.S. Patent Nos. 5,541,204, 5,770,615, 5,491,134, 5,776,983 and 5,488,064, a thyroid receptor beta modulator, such as a thyroid receptor ligand as disclosed in WO 97/21993 (U. Cal SF), WO 99/00353 (KaroBio) and WO00/039077 (KaroBio), a lipase inhibitor, such as orlistat or ATL-962 (Alizyme), serotonin receptor agonists, (e.g., BVT-933 (Biovitrum)), monoamine reuptake inhibitors or releasing agents, such as fenfluramine, dexfenfluramine, fluvoxamine, fluoxetine, paroxetine, sertraline, chlorphentermine, cloforex, clortermine, picilorex, sibutramine, dexamphetamine, phentermine, phenylpropanolamine or mazindol, anorectic agents such as topiramate (Johnson & Johnson), CNTF (ciliary neurotrophic factor) /Axokine[®] (Regeneron), BDNF (brain-derived neurotrophic factor), leptin and leptin receptor modulators, or cannabinoid-1 receptor antagonists, such as SR-141716 (Sanofi) or SLV-319 (Solvay).

[00117] Examples of suitable anti-diabetic agents for use in combination with the compounds of the present invention include: insulin secretagogues or insulin sensitizers, which may include biguanides, sulfonyl ureas, glucosidase inhibitors, aldose reductase inhibitors, PPAR γ agonists such as thiazolidinediones, PPAR α

agonists (such as fibric acid derivatives), PPAR δ antagonists or agonists, PPAR α/γ dual agonists, 11- β -HSD-1 inhibitors, dipeptidyl peptidase IV (DP4) inhibitors, SGLT2 inhibitors, glycogen phosphorylase inhibitors, and/or meglitinides, as well as insulin, and/or glucagon-like peptide-1 (GLP-1), GLP-1 agonist, and/or a PTP-1B inhibitor (protein tyrosine phosphatase-1B inhibitor).

[00118] The antidiabetic agent may be an oral antihyperglycemic agent preferably a biguanide such as metformin or phenformin or salts thereof, preferably metformin HCl. Where the antidiabetic agent is a biguanide, the compounds of the present invention will be employed in a weight ratio to biguanide within the range from about 0.001:1 to about 10:1, preferably from about 0.01:1 to about 5:1.

[00119] The antidiabetic agent may also preferably be a sulfonyl urea such as glyburide (also known as glibenclamide), glimepiride (disclosed in U.S. Patent No. 4,379,785), glipizide, gliclazide or chlorpropamide, other known sulfonylureas or other antihyperglycemic agents which act on the ATP-dependent channel of the beta-cells, with glyburide and glipizide being preferred, which may be administered in the same or in separate oral dosage forms. The oral antidiabetic agent may also be a glucosidase inhibitor such as acarbose (disclosed in U.S. Patent No. 4,904,769) or miglitol (disclosed in U.S. Patent No. 4,639,436), which may be administered in the same or in a separate oral dosage forms.

[00120] The compounds of the present invention may be employed in combination with a PPAR γ agonist such as a thiazolidinedione oral anti-diabetic agent or other insulin sensitizers (which has an insulin sensitivity effect in NIDDM patients) such as rosiglitazone (SKB), pioglitazone (Takeda), Mitsubishi's MCC-555 (disclosed in U.S. Patent No. 5,594,016), Glaxo-Wellcome's GL-262570, englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer, isaglitazone (MIT/J&J), JTT-501 (JPNT/P&U), L-895645 (Merck), R-119702 (Sankyo/WL), NN-2344 (Dr. Reddy/NN), or YM-440 (Yamanouchi), preferably rosiglitazone and pioglitazone.

[00121] The compounds of the present invention may be employed with a PPAR α/γ dual agonist such as MK-767/KRP-297 (Merck/Kyorin; as described in , K. Yajima, et. al., *Am. J. Physiol. Endocrinol. Metab.*, 284: E966-E971 (2003)), AZ-242

(tesaglitazar; Astra-Zeneca; as described in B. Ljung, et. al., *J. Lipid Res.*, 43, 1855-1863 (2002)); muraglitazar; or the compounds described in US patent 6,414,002.

[00122] The compounds of the present invention may be employed in combination with anti-hyperlipidemia agents, or agents used to treat arteriosclerosis. An example of an hypolipidemic agent would be an HMG CoA reductase inhibitor which includes, but is not limited to, mevastatin and related compounds as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related compounds as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds such as disclosed in U.S. Patent No. 4,346,227, simvastatin and related compounds as disclosed in U.S. Patent Nos. 4,448,784 and 4,450,171. Other HMG CoA reductase inhibitors which may be employed herein include, but are not limited to, fluvastatin, disclosed in U.S. Patent No. 5,354,772, cerivastatin disclosed in U.S. Patent Nos. 5,006,530 and 5,177,080, atorvastatin disclosed in U.S. Patent Nos. 4,681,893, 5,273,995, 5,385,929 and 5,686,104, pitavastatin (Nissan/Sankyo's nisvastatin (NK-104) or itavastatin), disclosed in U.S. Patent No. 5,011,930, Shionogi-Astra/Zeneca rosuvastatin (visastatin (ZD-4522)) disclosed in U.S. Patent No. 5,260,440, and related statin compounds disclosed in U.S. Patent No. 5,753,675, pyrazole analogs of mevalonolactone derivatives as disclosed in U.S. Patent No. 4,613,610, indene analogs of mevalonolactone derivatives as disclosed in PCT application WO 86/03488, 6-[2-(substituted-pyrrol-1-yl)-alkyl]pyran-2-ones and derivatives thereof as disclosed in U.S. Patent No. 4,647,576, Searle's SC-45355 (a 3-substituted pentanedioic acid derivative) dichloroacetate, imidazole analogs of mevalonolactone as disclosed in PCT application WO 86/07054, 3-carboxy-2-hydroxy-propane-phosphonic acid derivatives as disclosed in French Patent No. 2,596,393, 2,3-disubstituted pyrrole, furan and thiophene derivatives as disclosed in European Patent Application No. 0221025, naphthyl analogs of mevalonolactone as disclosed in U.S. Patent No. 4,686,237, octahydronaphthalenes such as disclosed in U.S. Patent No. 4,499,289, keto analogs of mevinolin (lovastatin) as disclosed in European Patent Application No. 0,142,146 A2, and quinoline and pyridine derivatives disclosed in U.S. Patent Nos. 5,506,219 and 5,691,322. In addition, phosphinic acid compounds useful in inhibiting HMG CoA reductase suitable for use herein are disclosed in GB 2205837.

[00123] The squalene synthetase inhibitors suitable for use herein include, but are not limited to, α -phosphono-sulfonates disclosed in U.S. Patent No. 5,712,396, those disclosed by Biller, et al., *J. Med. Chem.*, 31, 1869-1871 (1998) including isoprenoid (phosphinyl-methyl)phosphonates as well as other known squalene synthetase inhibitors, for example, as disclosed in U.S. Patent No. 4,871,721 and 4,924,024 and in Biller, S.A., Neuenschwander, K., Ponpipom, M.M., and Poulter, C.D., *Current Pharmaceutical Design*, 2, 1-40 (1996).

[00124] In addition, other squalene synthetase inhibitors suitable for use herein include the terpenoid pyrophosphates disclosed by P. Ortiz de Montellano, et al., *J. Med. Chem.*, 20, 243-249 (1977), the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-PP) analogs as disclosed by Corey and Volante, *J. Am. Chem. Soc.*, 98, 1291-1293 (1976), phosphinylphosphonates reported by McClard, R.W. et al., *J. Am. Chem. Soc.*, 109, 5544 (1987) and cyclopropanes reported by Capson, T.L., PhD dissertation, June, 1987, Dept. Med. Chem. U of Utah, Abstract, Table of Contents, pp 16, 17, 40-43, 48-51, Summary.

[00125] Other hypolipidemic agents suitable for use herein include, but are not limited to, fibric acid derivatives, such as fenofibrate, gemfibrozil, clofibrate, bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds as disclosed in U.S. Patent No. 3,674,836, probucol and gemfibrozil being preferred, bile acid sequestrants such as cholestyramine, colestipol and DEAE-Sephadex (SECHOLEX, POLICEXIDE) and cholestagel (Sankyo/Geltex), as well as lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphosphorylcholine (SPC, Roche), aminocyclodextrin (Tanabe Seiyoku), Ajinomoto AJ-814 (azulene derivative), melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid (niacin), acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin, poly(diallylmethylamine) derivatives such as disclosed in U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents.

[00126] The other hypolipidemic agent may be an ACAT inhibitor (which also has anti-atherosclerosis activity) such as disclosed in, *Drugs of the Future*, 24, 9-15

- (1999), (Avasimibe); "The ACAT inhibitor, CI-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters", Nicolosi et al., *Atherosclerosis* (Shannon, Irel), 137 (1), 77-85 (1998); "The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by selective suppression of the hepatic secretion of ApoB100-containing lipoprotein", Ghiselli, Giancarlo, *Cardiovasc. Drug Rev.*, 16 (1), 16-30 (1998); "RP 73163: a bioavailable alkylsulfinyl-diphenylimidazole ACAT inhibitor", Smith, C., et al., *Bioorg. Med. Chem. Lett*, 6 (1), 47-50 (1996); "ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic activities in experimental animals", Krause et al., Editor(s): Ruffolo, Robert R., Jr.; Hollinger, Manfred A., *Inflammation: Mediators Pathways*, 173-98 (1995), Publisher: CRC, Boca Raton, Fla.; "ACAT inhibitors: potential anti-atherosclerotic agents", Sliskovic et al., *Curr. Med. Chem.*, 1 (3), 204-25 (1994); "Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. 6. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). 7. Development of a series of substituted N-phenyl-N'-[(1-phenylcyclopentyl)-methyl]ureas with enhanced hypocholesterolemic activity", Stout et al., *Chemtracts: Org. Chem.*, 8 (6), 359-62 (1995), or TS-962 (Taisho Pharmaceutical Co. Ltd), as well as F-1394, CS-505, F-12511, HL-004, K-10085 and YIC-C8-434.
- 20 [00127] The hypolipidemic agent may be an upregulator of LDL receptor activity such as MD-700 (Taisho Pharmaceutical Co. Ltd) and LY295427 (Eli Lilly). The hypolipidemic agent may be a cholesterol absorption inhibitor preferably Schering-Plough's SCH48461 (ezetimibe) as well as those disclosed in *Atherosclerosis* 115, 45-63 (1995) and *J. Med. Chem.* 41, 973 (1998).
- 25 [00128] The other lipid agent or lipid-modulating agent may be a cholesteryl transfer protein inhibitor (CETP) such as Pfizer's CP-529,414 as well as those disclosed in WO/0038722 and in EP 818448 (Bayer) and EP 992496, and Pharmacia's SC-744 and SC-795, as well as CETi-1 and JTT-705.
- [00129] The hypolipidemic agent may be an ileal Na⁺/bile acid cotransporter inhibitor such as disclosed in *Drugs of the Future*, 24, 425-430 (1999). The ATP citrate lyase inhibitor which may be employed in the combination of the invention may include, for example, those disclosed in U.S. Patent No. 5,447,954.
- 30

- [00130] The other lipid agent also includes a phytoestrogen compound such as disclosed in WO 00/30665 including isolated soy bean protein, soy protein concentrate or soy flour as well as an isoflavone such as genistein, daidzein, glycitein or equol, or phytosterols, phytostanol or tocotrienol as disclosed in WO 2000/015201;
- 5 a beta-lactam cholesterol absorption inhibitor such as disclosed in EP 675714; an HDL upregulator such as an LXR agonist, a PPAR α -agonist and/or an FXR agonist; an LDL catabolism promoter such as disclosed in EP 1022272; a sodium-proton exchange inhibitor such as disclosed in DE 19622222; an LDL-receptor inducer or a steroidal glycoside such as disclosed in U.S. Patent No. 5,698,527 and GB 2304106;
- 10 an anti-oxidant such as beta-carotene, ascorbic acid, α -tocopherol or retinol as disclosed in WO 94/15592 as well as Vitamin C and an antihomocysteine agent such as folic acid, a folate, Vitamin B6, Vitamin B12 and Vitamin E; isoniazid as disclosed in WO 97/35576; a cholesterol absorption inhibitor, an HMG-CoA synthase inhibitor, or a lanosterol demethylase inhibitor as disclosed in WO 97/48701; a PPAR δ agonist
- 15 for treating dyslipidemia; or a sterol regulating element binding protein-I (SREBP-1) as disclosed in WO 2000/050574, for example, a sphingolipid, such as ceramide, or neutral sphingomyelinase (N-SMase) or fragment thereof. Preferred hypolipidemic agents are pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, pitavastatin and rosuvastatin, as well as niacin and/or cholestagel.
- 20 [00131] The compounds of the present invention may be employed in combination with anti-hypertensive agents. Examples of suitable anti-hypertensive agents for use in combination with the compounds of the present invention include beta adrenergic blockers, calcium channel blockers (L-type and/or T-type; e.g. diltiazem, verapamil, nifedipine, amlodipine and mybefradil), diuretics (e.g., chlorothiazide,
- 25 hydrochlorothiazide, flumethiazide, hydroflumethiazide, bendroflumethiazide, methylchlorothiazide, trichloromethiazide, polythiazide, benzthiazide, ethacrynic acid, tricynafene, chlorthalidone, furosemide, bumetanide, triamterene, amiloride, spironolactone), renin inhibitors, ACE inhibitors (e.g., captopril, zofenopril, fosinopril, enalapril, ceranopril, cilazapril, delapril, pentopril, quinapril,
- 30 ramipril, lisinopril), AT-1 receptor antagonists (e.g., losartan, irbesartan, valsartan), ET receptor antagonists (e.g., sitaxsentan, atrisentan and compounds disclosed in U.S.

Patent Nos. 5,612,359 and 6,043,265), Dual ET/AII antagonist (e.g., compounds disclosed in WO 00/01389), neutral endopeptidase (NEP) inhibitors, vasopeptidase inhibitors (dual NEP-ACE inhibitors) (e.g., omapatrilat and gemopatrilat), and nitrates.

5 **[00132]** Cannabinoid receptor modulators could be useful in treating other diseases associated with obesity, including sleep disorders. Therefore, the compounds described in the present invention could be used in combination with therapeutics for treating sleep disorders. Examples of suitable therapies for treatment of sleeping disorders for use in combination with the compounds of the present invention include
10 melatonin analogs, melatonin receptor antagonists, ML 1 B agonists, GABA receptor modulators; NMDA receptor modulators, histamine-3 (H3) receptor modulators, dopamine agonists and orexin receptor modulators.

[00133] Cannabinoid receptor modulators may reduce or ameliorate substance abuse or addictive disorders. Therefore, combination of cannabinoid receptor
15 modulators with agents used to treat addictive disorders may reduce the dose requirement or improve the efficacy of current addictive disorder therapeutics. Examples of agents used to treat substance abuse or addictive disorders are: selective serotonin reuptake inhibitors (SSRI), methadone, buprenorphine, nicotine and bupropion.

20 **[00134]** Cannabinoid receptor modulators may reduce anxiety or depression; therefore, the compounds described in this application may be used in combination with anti-anxiety agents or antidepressants. Examples of suitable anti-anxiety agents for use in combination with the compounds of the present invention include benzodiazepines (e.g., diazepam, lorazepam, oxazepam, alprazolam,
25 chlordiazepoxide, clonazepam, chlorazepate, halazepam and prazepam), 5HT1A receptor agonists (e.g., buspirone, flesinoxan, gepirone and ipsapirone), and corticotropin releasing factor (CRF) antagonists.

[00135] Examples of suitable classes of anti-depressants for use in combination with the compounds of the present invention include norepinephrine reuptake
30 inhibitors (tertiary and secondary amine tricyclics), selective serotonin reuptake inhibitors (SSRIs) (fluoxetine, fluvoxamine, paroxetine and sertraline), monoamine oxidase inhibitors (MAOIs) (isocarboxazid, phenelzine, tranlycypromine, selegiline),

reversible inhibitors of monoamine oxidase (RIMAs) (moclobemide), serotonin and norepinephrine reuptake inhibitors (SNRIs) (venlafaxine), corticotropin releasing factor (CRF) receptor antagonists, alpha-adrenoreceptor antagonists, and atypical antidepressants (bupropion, lithium, nefazodone, trazodone and viloxazine).

5 [00136] The combination of a conventional antipsychotic drug with a CB-1 receptor antagonist could also enhance symptom reduction in the treatment of psychosis or mania. Further, such a combination could enable rapid symptom reduction, reducing the need for chronic treatment with antipsychotic agents. Such a combination could also reduce the effective antipsychotic dose requirement, resulting
10 in reduced probability of developing the motor dysfunction typical of chronic antipsychotic treatment.

[00137] Examples of suitable antipsychotic agents for use in combination with the compounds of the present invention include the phenothiazine (chlorpromazine, mesoridazine, thioridazine, acetophenazine, fluphenazine, perphenazine and
15 trifluoperazine), thioxanthine (chlorprothixene, thiothixene), heterocyclic dibenzazepine (clozapine, olanzepine and aripiprazole), butyrophenone (haloperidol), dipheylbutylpiperidine (pimozide) and indolone (molindolone) classes of antipsychotic agents. Other antipsychotic agents with potential therapeutic value in combination with the compounds in the present invention include loxapine, sulpiride
20 and risperidone.

[00138] Combination of the compounds in the present invention with conventional antipsychotic drugs could also provide an enhanced therapeutic effect for the treatment of schizophrenic disorders, as described above for manic disorders. As used here, schizophrenic disorders include paranoid, disorganized, catatonic,
25 undifferentiated and residual schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder and psychotic disorder not specified. Examples of suitable antipsychotic drugs for combination with the compounds in the present invention include the antipsychotics mentioned above, as well as dopamine receptor antagonists, muscarinic receptor agonists, 5HT_{2A}
30 receptor antagonists and 5HT_{2A}/dopamine receptor antagonists or partial agonists (e.g., olanzepine, aripiprazole, risperidone, ziprasidone).

- [00139] The compounds described in the present invention could be used to enhance the effects of cognition-enhancing agents, such as acetylcholinesterase inhibitors (e.g., tacrine), muscarinic receptor-1 agonists (e.g., milameline), nicotinic agonists, glutamic acid receptor (AMPA and NMDA) modulators, and nootropic agents (e.g., piracetam, levetiracetam). Examples of suitable therapies for treatment of Alzheimer's disease and cognitive disorders for use in combination with the compounds of the present invention include donepezil, tacrine, revastigraïne, 5HT₆, gamma secretase inhibitors, beta secretase inhibitors, SK channel blockers, Maxi-K blockers, and KCNQs blockers.
- 10 [00140] The compounds described in the present invention could be used to enhance the effects of agents used in the treatment of Parkinson's Disease. Examples of agents used to treat Parkinson's Disease include: levodopa with or without a COMT inhibitor, antihlutamatergic drugs (amantadine, riluzole), alpha-2 adrenergic antagonists such as idazoxan, opiate antagonists, such as naltrexone, other dopamine agonists or transporter modulators, such as ropinirole, or pramipexole or neurotrophic factors such as glial derived neurotrophic factor (GDNF).
- 15 [00141] The compounds described in the present invention could be used in combination with suitable anti-inflammatory agents. Examples of suitable anti-inflammatory agents for use in combination with the compounds of the present invention include prednisone, dexamethasone, cyclooxygenase inhibitors (i.e., COX-1 and/or COX-2 inhibitors such as NSAIDs, aspirin, indomethacin, ibuprofen, piroxicam, Naproxen[®], Celebrex[®], Vioxx[®]), CTLA4-Ig agonists/antagonists, CD40 ligand antagonists, IMPDH inhibitors, such as mycophenolate (CellCept[®]), integrin antagonists, alpha-4 beta-7 integrin antagonists, cell adhesion inhibitors, interferon gamma antagonists, ICAM-1, tumor necrosis factor (TNF) antagonists (e.g., infliximab, OR1384, including TNF-alpha inhibitors, such as tenidap, anti-TNF antibodies or soluble TNF receptor such as etanercept (Enbrel[®]), rapamycin (sirolimus or Rapamune) and leflunomide (Arava)), prostaglandin synthesis inhibitors, budesonide, clofazimine, CNI-1493, CD4 antagonists (e.g., priliximab), p38 mitogen-activated protein kinase inhibitors, protein tyrosine kinase (PTK) inhibitors, IKK
- 20
25
30

inhibitors, and therapies for the treatment of irritable bowel syndrome (e.g., Zelnorm[®] and Maxi-K[®] openers such as those disclosed in U.S. Patent No. 6,184,231 B1).

[00142] Exemplary of such other therapeutic agents which may be used in combination with cannabinoid receptor modulators include the following:

- 5 cyclosporins (e.g., cyclosporin A), anti-IL-2 receptor (Anti-Tac), anti-CD45RB, anti-CD2, anti-CD3 (OKT-3), anti-CD4, anti-CD80, anti-CD86, monoclonal antibody OKT3, agents blocking the interaction between CD40 and gp39, such as antibodies specific for CD40 and/or gp39 (i.e., CD154), fusion proteins constructed from CD40 and gp39 (CD40Ig and CD8gp39), inhibitors, such as nuclear translocation inhibitors, 10 of NF-kappa B function, such as deoxyspergualin (DSG), gold compounds, antiproliferative agents such as methotrexate, FK506 (tacrolimus, Prograf), mycophenolate mofetil, cytotoxic drugs such as azathioprine and cyclophosphamide, anticytokines such as antiIL-4 or IL-4 receptor fusion proteins and PDE 4 inhibitors such as Ariflo, and the PTK inhibitors disclosed in the following U.S. patent 15 applications, incorporated herein by reference in their entirety: Ser. No. 09/097,338, filed Jun. 15, 1998; Ser. No. 09/094,797, filed Jun. 15, 1998; Ser. No. 09/173,413, filed Oct. 15, 1998; and Ser. No. 09/262,525, filed Mar. 4, 1999. See also the following documents and references cited therein and incorporated herein by reference: Hollenbaugh, D., et al., "Cleavable CD40Ig Fusion Proteins and the 20 Binding to Sgp39", *J. Immunol. Methods* (Netherlands), 188 (1), pp. 1-7 (Dec. 15, 1995); Hollenbaugh, D., et al., "The Human T Cell Antigen Gp39, A Member of the TNF Gene Family, Is a Ligand for the CD40 Receptor: Expression of a Soluble Form of Gp39 with B Cell Co-Stimulatory Activity", *EMBO J* (England), 11 (12), pp. 4313-4321 (December 1992); and Moreland, L. W. et al., "Treatment of Rheumatoid 25 Arthritis with a Recombinant Human Tumor Necrosis Factor Receptor (P75)-Fc Fusion Protein," *New England J. of Medicine*, 337 (3), pp. 141-147 (1997).

[00143] The above other therapeutic agents, when employed in combination with the compounds of the present invention, may be used, for example, in those amounts indicated in the Physicians' Desk Reference (PDR) or as otherwise determined by one 30 of ordinary skill in the art.

[00144] The compounds of formula (I) of the invention can be administered orally or parenterally, such as subcutaneously or intravenously, as well as by nasal

application, rectally or sublingually to various mammalian species known to be subject to such maladies, e.g., humans, in an effective amount up to 1 gram, preferably up to 200 mg, more preferably up to 100 mg in a regimen of single, two or four divided daily doses.

- 5 **[00145]** The compounds of the formula (I) can be administered for any of the uses described herein by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or
- 10 suspensions); nasally, including administration to the nasal membranes, such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The present compounds can, for example, be administered in a form suitable for immediate release or extended
- 15 release. Immediate release or extended release can be achieved by the use of suitable pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps. The present compounds can also be administered liposomally.
- [00146]** Exemplary compositions for oral administration include suspensions
- 20 which can contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which can contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other
- 25 excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The compounds of formula I can also be delivered through the oral cavity by sublingual and/or buccal administration. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms which may be used. Exemplary compositions include those formulating the present compound(s) with fast dissolving
- 30 diluents such as mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (avicel) or polyethylene glycols (PEG). Such formulations can also include an excipient to aid

mucosal adhesion such as hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (e.g., Gantrez), and agents to control release such as polyacrylic copolymer (e.g. Carbopol 934). Lubricants, glidants, flavors, coloring agents and stabilizers may
5 also be added for ease of fabrication and use.

[00147] Exemplary compositions for nasal aerosol or inhalation administration include solutions in saline which can contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

10 [00148] Exemplary compositions for parenteral administration include injectable solutions or suspensions which can contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty
15 acids, including oleic acid, or Cremaphor.

[00149] Exemplary compositions for rectal administration include suppositories which can contain, for example, a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols, which are solid at ordinary temperatures, but liquify and/or dissolve in the rectal cavity to release the drug.

20 [00150] Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene).


[00151] It will be understood that the specific dose level and frequency of dosage for any particular subject can be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and
25 length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the particular condition.

[00152] It should be understood that while this invention has been described herein in terms of specific embodiments set forth in detail, such embodiments are presented
30 by way of illustration of the general principles of the invention, and the invention is not necessarily limited thereto. Certain modifications and variations in any given material, process step or chemical formula will be readily apparent to those skilled in

the art without departing from the true spirit and scope of the present invention, and all such modifications and variations should be considered within the scope of the claims that follow.

WHAT IS CLAIMED IS:

1. A compound or a prodrug or a solvate or a pharmaceutically acceptable salt or a stereoisomer thereof according to Formula I



I

wherein:

B is selected from the group consisting of CR⁴ and a direct bond;

x is a double bond when B is CR⁴;

x is a single bond when B is a direct bond;

10 R¹ is selected from the group consisting of halogen, cyano, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, OR⁶, OCONR⁷R⁸, OSO₂NR⁷R⁸, NR⁷R⁸, NR⁷COR⁸, NR⁷CO₂R⁶, NR⁷CONR⁸R⁹, NR⁷SO₂R⁶, NR⁷SO₃R⁶, NR⁷SO₂NR⁸R⁹, SR⁶, SO₂R⁶, SO₂NR⁷R⁸, COR⁶, CO₂R⁶ and CONR⁷R⁸, any of which may optionally be substituted with 1 to 3 R¹⁰;

15 R² is selected from the group consisting of halogen, cyano, cycloalkyl, aryl, heteroaryl, wherein aryl and heteroaryl may optionally be substituted with 1 to 3 R¹⁰;

R³, when B is a direct bond, is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, arylalkyl and heteroarylalkyl, any of which may be optionally be substituted with 1 to 3 R¹⁰;

20 R³ is absent when B is CR⁴;

R⁴ is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, aryl, heteroaryl, arylalkyl and heteroarylalkyl, any of which may optionally be substituted with 1 to 3 R¹⁰;

25 R⁶ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl and heteroaryl, any of which may optionally be substituted with 1 to 3 R¹⁰;

R^7 , R^8 and R^9 are independently selected from the group consisting of hydrogen, aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl and heterocyclyl, any of which may optionally be substituted with 1 to 3 R^{10} ;

any two of R^6 , R^7 , R^8 and R^9 may optionally together form a 4 to 7-membered
 5 heterocyclyl ring, or a 5 to 6-membered heteroaryl ring;

R^{10} selected from the group consisting of aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, OR^6 , $OCOR^6$, $OCONR^7R^8$, $OSO_2NR^7R^8$, NR^7R^8 , NR^7COR^8 , $NR^7CO_2R^6$, $NR^7CONR^8R^9$, $NR^7SO_2R^6$, $NR^7SO_3R^6$, $NR^7SO_2NR^8R^9$, SR^6 , SO_2R^6 , $SO_2NR^7R^8$, COR^6 , CO_2R^6 , $CONR^7R^8$, halogen, cyano, and the divalent
 10 groups oxo ($=O$) and oximato ($=NOR^7$), any of which may optionally be substituted with 1 to 3 R^{11} ;

R^{11} selected from the group consisting of aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, OR^6 , $OCONR^7R^8$, $OSO_2NR^7R^8$, NR^7R^8 , NR^7COR^8 , $NR^7CO_2R^6$, $NR^7CONR^8R^9$, $NR^7SO_2R^6$, $NR^7SO_3R^6$, $NR^7SO_2NR^8R^9$, SR^6 , SO_2R^6 ,
 15 $SO_2NR^7R^8$, COR^6 , CO_2R^6 , $CONR^7R^8$, halogen, cyano, and the divalent groups oxo ($=O$) and oximato ($=NOR^7$), any of which may optionally be substituted with 1 to 3 R^{12} ; and

R^{12} selected from the group consisting of aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, OR^6 , $OCONR^7R^8$, $OSO_2NR^7R^8$, NR^7R^8 , NR^7COR^8 ,
 20 $NR^7CO_2R^6$, $NR^7CONR^8R^9$, $NR^7SO_2R^6$, $NR^7SO_3R^6$, $NR^7SO_2NR^8R^9$, SR^6 , SO_2R^6 , $SO_2NR^7R^8$, COR^6 , CO_2R^6 , $CONR^7R^8$, halogen, cyano, and the divalent groups oxo ($=O$) and oximato ($=NOR^7$).

2. The compound according to Claim 1, wherein R^1 is selected from the
 25 group consisting of aryl and heteroaryl, either of which may be optionally substituted with 1 to 3 R^{10} .

3. The compound according to Claim 1, wherein R^2 is selected from the
 group consisting of aryl and heteroaryl, either of which may be optionally substituted
 30 with 1 to 3 R^{10} .

4. The compound according to Claim 1, wherein:

B is a direct bond;

x is a single bond;

R^1 is selected from the group consisting of aryl and heteroaryl, either of which may be optionally substituted with 1 to 3 R^{10} ;

5 R^2 is selected from the group consisting of aryl and heteroaryl, either of which may be optionally substituted with 1 to 3 R^{10} ; and

R^3 is selected from the group consisting of alkyl, cycloalkyl, heterocyclyl; cycloalkylalkyl, heterocyclalkyl, arylalkyl and heteroarylalkyl, any of which may be optionally be substituted with 1 to 3 R^{10} .

10

5. The compound according to Claim 1, wherein:

B is CR^4 ;


x is a double bond;


15 R^1 is selected from the group consisting of aryl and heteroaryl, either of which may be optionally substituted with 1 to 3 R^{10} ;

R^2 is selected from the group consisting of aryl and heteroaryl, either of which may be optionally substituted with 1 to 3 R^{10} ; and


20 R^4 is selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclalkyl, arylalkyl and heteroarylalkyl, any of which may be optionally be substituted with 1 to 3 R^{10} .

6. The compound according to Claim 1, wherein the compound is selected from the group consisted of:







,




,



,




,




5


,







,



,







,







, and



- 5 7. The compound according to Claim 1, wherein the compound is selected from the group consisted of:



10



5



, and

8. A pharmaceutical composition, comprising: at least one compound according to Claim 1; and at least one pharmaceutically acceptable carrier or diluent.

10

9. The pharmaceutical composition according to Claim 7, further comprising: at least one additional therapeutic agent.

10. A pharmaceutical combination comprising a pharmaceutical composition of Claim 7 and a therapeutic agent selected from anti-obesity agents;

15

appetite suppressants; anti-diabetic agents; anti-hyperlipidemia agents; hypolipidemic agents; hypocholesterolemic agents; lipid-modulating agents; cholesterol-lowering agents; lipid-lowering agents; HDL-raising agent, anti-hypertensive agents; agents used to treat sleep disorders; agents used to treat substance abuse and addictive disorders; anti-anxiety agents; anti-depressants; anti-psychotic agents; cognition enhancing agents; agents used to treat cognitive disorders; agents used to treat Alzheimer's disease; agents used to treat Parkinson's disease; anti-inflammatory agents; agents used to treat neurodegeneration; agents used to treat arteriosclerosis; agents used to treat respiratory conditions; agents used to treat bowel disorders; cardiac glycosides; and anti-tumor agents.

11. The pharmaceutical combination according to Claim 9 wherein the other therapeutic agent may be administered prior to, simultaneously with, or following the administration of the pharmaceutical composition of Claim 7.

15

12. A method for treating obesity, comprising: administering a therapeutically effective amount of at least one cannabinoid receptor 1 antagonist according to Claim 1 to a patient in need.

13. A method for smoking cessation, comprising: administering a therapeutically effective amount of at least one cannabinoid receptor 1 antagonist according to Claim 1 to a patient in need.