




(86) Date de dépôt PCT/PCT Filing Date: 2012/04/12
(87) Date publication PCT/PCT Publication Date: 2012/10/18
(45) Date de délivrance/Issue Date: 2019/09/03
(85) Entrée phase nationale/National Entry: 2013/10/10
(86) N° demande PCT/PCT Application No.: EP 2012/056649
(87) N° publication PCT/PCT Publication No.: 2012/140120
(30) Priorité/Priority: 2011/04/13 (US61/474,936)

(51) Cl.Int./Int.Cl. *C07D 409/10* (2006.01),
C07D 409/14 (2006.01)
(72) Inventeurs/Inventors:
HOUPIS, IOANNIS NICOLAOS, BE;
LEMAIRE, SEBASTIEN FRANCOIS EMMANUEL, BE
(73) Propriétaire/Owner:
JANSSEN PHARMACEUTICA NV, BE
(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : PROCÉDE DE PRÉPARATION DE COMPOSÉS UTILES EN TANT QU'INHIBITEURS DE SGLT2
(54) Title: PROCESS FOR THE PREPARATION OF COMPOUNDS USEFUL AS INHIBITORS OF SGLT2



(57) Abrégé/Abstract:

The present invention is directed to a novel process for the preparation of compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof; comprising: reacting a compound of formula (V) wherein LG¹ is a leaving group, with a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about -78°C to about room temperature; to yield a mixture of the corresponding compound of formula (VI), wherein M¹ is lithium, and the zinc salt; admixing to the mixture of the compound of formula (VI) and the zinc salt, a first ether solvent; to yield the corresponding compound of formula (VII), wherein M² is a reactive zinc species.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2012/140120 A1(43) International Publication Date
18 October 2012 (18.10.2012)

(51) International Patent Classification:

C07D 409/10 (2006.01) C07D 409/14 (2006.01)

(21) International Application Number:

PCT/EP2012/056649

(22) International Filing Date:

12 April 2012 (12.04.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/474,936 13 April 2011 (13.04.2011) US

(71) Applicant (for all designated States except US):

JANSSEN PHARMACEUTICA NV [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HOUPIS, Ioannis, Nicolaos [GR/BE]; c/o Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE). LEMAIRE, Sebastien, François, Emmanuel [FR/BE]; c/o Janssen Pharmaceutica NV, Turnhoutseweg 30, B-2340 Beerse (BE).




(74) Common Representative: JANSSEN PHARMACEUTICA NV; Turnhoutseweg 30, B-2340 Beerse (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: PROCESS FOR THE PREPARATION OF COMPOUNDS USEFUL AS INHIBITORS OF SGLT2



(57) **Abstract:** The present invention is directed to a novel process for the preparation of compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof; comprising: reacting a compound of formula (V) wherein LG^1 is a leaving group, with a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about -78°C to about room temperature; to yield a mixture of the corresponding compound of formula (VI), wherein M^1 is lithium, and the zinc salt; admixing to the mixture of the compound of formula (VI) and the zinc salt, a first ether solvent; to yield the corresponding compound of formula (VII), wherein M^2 is a reactive zinc species.

WO 2012/140120 A1

- GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

 - *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
 - *of inventorship (Rule 4.17(iv))*

Published:

 - *with international search report (Art. 21(3))*

PROCESS FOR THE PREPARATION OF COMPOUNDS USEFUL AS
INHIBITORS OF SGLT2

5 This application claims priority to U. S. Provisional Application 61/474,936, filed
on April 13, 2011.

FIELD OF THE INVENTION

10 The present invention is directed to a novel process for the preparation of
compounds having inhibitory activity against sodium-dependent glucose transporter
(SGLT) being present in the intestine or kidney.

BACKGROUND OF THE INVENTION

15 Diet therapy and exercise therapy are essential in the treatment of diabetes
mellitus. When these therapies do not sufficiently control the conditions of patients,
insulin or an oral antidiabetic agent is additionally used for the treatment of diabetes. At
the present, there have been used as an antidiabetic agent biguanide compounds,
sulfonylurea compounds, insulin resistance improving agents and a-glucosidase
inhibitors. However, these antidiabetic agents have various side effects. For example,
biguanide compounds cause lactic acidosis, sulfonylurea compounds cause significant
20 hypoglycemia, insulin resistance improving agents cause edema and heart failure, and
a-glucosidase inhibitors cause abdominal bloating and diarrhea. Under such
circumstances, it has been desired to develop novel drugs for treatment of diabetes
mellitus having no such side effects.

25 Recently, it has been reported that hyperglycemia participates in the onset and
progressive impairment of diabetes mellitus, i.e., glucose toxicity theory. Namely,
chronic hyperglycemia leads to decrease of insulin secretion and further to decrease of
insulin sensitivity, and as a result, the blood glucose concentration is increased so that
diabetes mellitus is self-exacerbated (UNGER, R.H., et al., "Hyperglycemia as an
Inducer as well as a Consequence of Impaired Islet Cell Function and Insulin
30 Resistance: Implications for the

Management of Diabetes”, Diabetologia, 1985, pp119-121, Vol. 28, Issue 3;
ROSSETTI, L. et al., “Glucose Toxicity”, Diabetes Care, 1990, pp 610-630, Vol.
13, Issue 6). Therefore, by treating hyperglycemia, the aforementioned self-
exacerbating cycle is interrupted so that the prophylaxis or treatment of
5 diabetes mellitus is made possible.

As one of the methods for treating hyperglycemia, it is considered to
excrete an excess amount of glucose directly into urine so that the blood
glucose concentration is normalized. For example, by inhibiting sodium-
dependent glucose transporter being present at the proximal convoluted tubule
10 of kidney, the re-absorption of glucose at the kidney is inhibited, by which the
excretion of glucose into urine is promoted so that the blood glucose level is
decreased. In fact, it is confirmed that by continuous subcutaneous
administration of phlorizin having SGLT inhibitory activity to diabetic animal
models, hyperglycemia is normalized and the blood glucose level thereof can
15 be kept normal for a long time so that the insulin secretion and insulin
resistance are improved (ROSSETTI, L., et al., “Correction of Hyperglycemia
with Phlorizin Normalizes Tissue sensitivity to Insulin in Diabetic Rats”, Journal
of Clinical Investigation, 1987, pp 1510-1515, Vol. 79, Issue 5, pp. 1510-1515;
ROSSETTI, L., et al., “Effect of Chronic Hyperglycemia on in vivo Insulin
20 Secretion in Partially Pancreatectomized Rats”, Journal of Clinical
Investigation, 1987, pp 1037-1044, Vol. 80, Issue 4; KAHN, B.B., et al.,
“Normalization of blood glucose in diabetic rats with phlorizin treatment
reverses insulin-resistant glucose transport in adipose cells without restoring
glucose transporter gene expression”, J. Clin. Invest., 1991, pp 561-570, Vol.
25 87).


In addition, by treating diabetic animal models with SGLT inhibitory
agents for a long time, insulin secretion response and insulin sensitivity of the
animals are improved without incurring any adverse affects on the kidney or
imbalance in blood levels of electrolytes, and as a result, the onset and
30 progress of diabetic nephropathy and diabetic neuropathy are prevented
(KENJI, T., et al., “ Na⁺-Glucose Co-transporter (SGLT) Inhibitors as
Antidiabetic Agents. 4. Synthesis and Pharmacological Properties of 4’-

- Dehydroxyphlorizin Derivatives Substituted on the B Ring", J. Med. Chem., 1999, pp 5311-5324, Vol. 42; KENJI, A., et al., "Improved diabetic syndrome in C57BL/KsJ-db/db mice by oral administration of the Na⁺-glucose cotransporter inhibitor T-1095", British Journal of Pharmacology, 2001, pp 578-586, Vol. 132, Issue 2; UETA, K., et al., "Long Term Treatment with the Na⁺ Glucose Co-transporter Inhibitor T-1095 causes Sustained Improvement in Hyperglycemia and Prevents Diabetic Neuropathy in Goto-Kakizaki Rats", Life Sci., 2005, pp 2655-2668, Vol. 76, Issue 23)

From the above, SGLT inhibitors may be expected to improve insulin secretion and insulin resistance by decreasing the blood glucose level in diabetic patients and further prevent the onset and progress of diabetes mellitus and diabetic complications.

SUMMARY OF THE INVENTION

The present invention is directed to a process for the preparation of compounds of formula (I)



wherein Ring A and Ring B are one of the following:

- (1) Ring A is an optionally substituted unsaturated monocyclic heterocyclic ring, and Ring B is an optionally substituted unsaturated monocyclic heterocyclic ring, an optionally substituted unsaturated fused heterobicyclic ring, or an optionally substituted benzene ring; or
- (2) Ring A is an optionally substituted benzene ring, and Ring B is an optionally substituted unsaturated monocyclic heterocyclic ring, or an optionally substituted unsaturated fused heterobicyclic ring wherein Y is linked to the heterocyclic ring of the fused heterobicyclic ring; or


- (3) Ring A is an optionally substituted unsaturated fused heterobicyclic ring, wherein the sugar moiety X-(sugar) and the moiety -Y-(Ring B) are both on the same heterocyclic ring of the fused heterobicyclic ring, and Ring B is an optionally substituted unsaturated monocyclic heterocyclic ring, an optionally substituted unsaturated fused heterobicyclic ring, or an optionally substituted benzene ring;

X is a carbon atom;


Y is $-(CH_2)_n-$; wherein n is 1 or 2;

provided that in Ring A, X is part of an unsaturated bond;


- and pharmaceutically acceptable salts and solvates thereof; comprising




- reacting a compound of formula (V) wherein LG^1 is a leaving group, with a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about $-78^\circ C$ to about room temperature; to yield a mixture of the corresponding compound of formula (VI), wherein M^1 is lithium, and the zinc salt;



- admixing to the mixture of the compound of formula (VI) and the zinc salt, a first ether solvent; to yield the corresponding compound of formula (VII), wherein M^2 is a reactive zinc species;




reacting the compound of formula (VII) with a compound of formula (VIII), wherein each Z is an independently selected oxygen protecting group and wherein LG² is a leaving group; optionally in a mixture of a second ether solvent and a second hydrocarbon solvent; to yield the corresponding compound of formula (IX);



de-protecting the compound of formula (IX); to yield the corresponding compound of formula (I).

In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (I), as herein defined, comprising




reacting a compound of formula (V), wherein LG¹ is a leaving group, with

(a) a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about -78°C to about room temperature;


(b) then admixing a first ether solvent; and

5 (c) then reacting the resulting mixture with a compound of formula (VIII), wherein each Z is an independently selected oxygen protecting group and wherein LG² is a leaving group; optionally in a mixture of a second ether solvent and a second hydrocarbon solvent; to yield the corresponding compound of formula (IX);




de-protecting the compound of formula (IX); to yield the corresponding compound of formula (I).


15 In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-S)




or solvate thereof; (also known as 1-(β-D-glucopyranosyl)-4-methyl-3-[5-(4-fluorophenyl)-2-thienylmethyl]benzene); comprising




reacting a compound of formula (V-S) wherein LG¹ is a leaving group,
 with a mixture of a zinc salt and an organo-lithium reagent; in a first
 hydrocarbon solvent; at a temperature in the range of from about -78°C to
 5 about room temperature; to yield a mixture of the corresponding compound of
 formula (VI-S), wherein M¹ is lithium, and the zinc salt;



admixing to the mixture of the compound of formula (VI-S) and the zinc
 salt a first ether solvent; to yield the corresponding compound of formula (VII-
 10 S), wherein M² is a reactive zinc species;




reacting a compound of formula (VII-S), with a compound of formula (VIII-S), wherein each Z is an independently selected oxygen protecting group and wherein LG² is a leaving group; optionally in a mixture of a second ether solvent and a second hydrocarbon solvent; to yield the corresponding compound of formula (IX-S);



de-protecting the compound of formula (IX-S); to yield the corresponding compound of formula (I-S).

In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-S), as herein defined, comprising




5 reacting a compound of formula (V-S), wherein LG¹ is a leaving group,
 with

(a) a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about -78°C to about room temperature;


(b) then admixing a first ether solvent; and

10 (c) then reacting the resulting mixture with a compound of formula (VIII-S), wherein each Z is an independently selected oxygen protecting group and wherein LG² is a leaving group; optionally in a mixture of a second ether solvent and a second hydrocarbon solvent; to yield the corresponding compound of formula (IX-S);




de-protecting the compound of formula (IX-S); to yield the corresponding compound of formula (I-S).

- 5 In another embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-K)




or pharmaceutically acceptable salt or solvate thereof; (also known as 1-(β -D-glucopyranosyl)-4-chloro-3-[5-(4-fluoro-3-pyridyl)-2-thienylmethyl]benzene); comprising

10




reacting a compound of formula (V-K) wherein LG^1 is a leaving group, with a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about -78°C to

about room temperature; to yield a mixture of the corresponding compound of formula (VI-K), wherein M^1 is lithium, and the zinc salt;




admixing to the mixture of the compound of formula (VI-K) and the zinc salt a first ether solvent; to yield the corresponding compound of formula (VII-K), wherein M^2 is a reactive zinc species;



reacting the compound of formula (VII-K), with a compound of formula (VIII-S), wherein each Z is an independently selected oxygen protecting group and wherein LG^2 is a leaving group; optionally in a mixture of a second ether

Chemical reaction scheme showing the conversion of compound (IX-K) to compound (I-K). Compound (IX-K) is a substituted tetrahydropyran with four Z-protected groups (ZO) and a side chain containing a thiophene ring and a 2-fluoropyridine ring. Compound (I-K) is the deprotected form, where the ZO groups are replaced by hydroxyl (OH) groups. The reaction is indicated by a right-pointing arrow.




12

(a) a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon solvent; at a temperature in the range of from about -78°C to about room temperature;

(b) then admixing a first ether solvent; and

5 (c) then reacting the resulting mixture with a compound of formula (VIII-K), wherein each Z is an independently selected oxygen protecting group and wherein LG² is a leaving group; optionally in a mixture of a second ether solvent and a second hydrocarbon solvent; to yield the corresponding compound of formula (IX-K);



de-protecting the compound of formula (IX-K); to yield the corresponding compound of formula (I-K).

15 The present invention is further directed to a product prepared according to any of the processes described herein.

Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and the product prepared according to any of the processes described herein. An illustration of the invention is a pharmaceutical composition made by mixing the product prepared according to

20 any of the processes described herein and a pharmaceutically acceptable carrier. Illustrating the invention is a process for making a pharmaceutical composition comprising mixing the product prepared according to any of the processes described herein and a pharmaceutically acceptable carrier.

Exemplifying the invention are methods of treating a disorder mediated


25 by SGLT (including treating or delaying the progression or onset of diabetes mellitus, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy,

delayed wound healing, insulin resistance, hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids, elevated blood levels of glycerol, hyperlipidemia, obesity, hypertriglyceridemia, Syndrome X, diabetic complications, atherosclerosis, or hypertension,) comprising administering to
 5 the subject in need thereof a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

Further exemplifying the invention are methods of treating type 1 and type 2 diabetes mellitus, comprising administering to a subject in need of treatment a therapeutically effective amount of a therapeutically effective
 10 amount of any of the compounds or pharmaceutical compositions described above, alone or in combination with at least one antidiabetic agent, agent for treating diabetic complications, anti-obesity agent, antihypertensive agent, antiplatelet agent, anti-atherosclerotic agent and/or hypolipidemic agent.

15 DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a process for the preparation of compounds of formula (I)



wherein X, Y, Ring A and Ring B are as herein defined; and
 20 pharmaceutically acceptable salts and solvates thereof; as described in more detail herein. The compounds of the formula (I) exhibits an inhibitory activity against sodium-dependent glucose transporter being present in the intestine and the kidney of mammalian species, and is useful in the treatment of diabetes mellitus or diabetic complications such as diabetic retinopathy,
 25 diabetic neuropathy, diabetic nephropathy, obesity, and delayed wound healing. In an embodiment, the present invention is directed to a process for

the preparation of a compound of formula (I-S), as described in more detail herein. In another embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-K), as described in more detail herein.

5

The term “**halogen**”, shall include chlorine, bromine, fluorine and iodine. When referring to substituents on the compound of formula (I), the term “**halogen atom**” or “**halo**” shall mean chlorine, bromine and fluorine, and chlorine and fluorine are preferable.

10

The term “**alkyl group**” means a straight or branched saturated monovalent hydrocarbon chain having 1 to 12 carbon atoms. The straight chain or branched chain alkyl group having 1 to 6 carbon atoms is preferable, and the straight chain or branched chain alkyl group having 1 to 4 carbon atoms is more preferable. Examples thereof are methyl group, ethyl group, propyl group, isopropyl group, butyl group, t-butyl group, isobutyl group, pentyl group, hexyl group, isohexyl group, heptyl group, 4,4-dimethylpentyl group, octyl group, 2,2,4-trimethylpentyl group, nonyl group, decyl group, and various branched chain isomers thereof. Further, the alkyl group may optionally and independently be substituted by 1 to 4 substituents as listed below, if

15

20

necessary.

The term “**alkylene group**” or “**alkylene**” means a straight or branched divalent saturated hydrocarbon chain having 1 to 12 carbon atoms. The straight chain or branched chain alkylene group having 1 to 6 carbon atoms is preferable, and the straight chain or branched chain alkylene group having 1 to 4 carbon atoms is more preferable. Examples thereof are methylene group, ethylene group, propylene group, trimethylene group, etc. If necessary, the alkylene group may optionally be substituted in the same manner as the above-mentioned “alkyl group”. Where alkylene groups as defined above attach at two different carbon atoms of the benzene ring, they form an annelated five, six or seven membered carbocycle together with the carbon atoms to which they are attached, and may optionally be substituted by one or more substituents defined below.

25

30

The term “**alkenyl group**” means a straight or branched monovalent hydrocarbon chain having 2 to 12 carbon atoms and having at least one double bond. Preferable alkenyl group is a straight chain or branched chain alkenyl group having 2 to 6 carbon atoms, and the straight chain or branched chain alkenyl group having 2 to 4 carbon atoms is more preferable. Examples thereof are vinyl group, 2-propenyl group, 3-butenyl group, 2-butenyl group, 4-pentenyl group, 3-pentenyl group, 2-hexenyl group, 3-hexenyl group, 2-heptenyl group, 3-heptenyl group, 4-heptenyl group, 3-octenyl group, 3-nonenyl group, 4-decenyl group, 3-undecenyl group, 4-dodecenyl group, 4,8,12-tetradecatrienyl group, etc. The alkenyl group may optionally and independently be substituted by 1 to 4 substituents as mentioned below, if necessary.

The term “**alkenylene group**” means a straight or branched divalent hydrocarbon chain having 2 to 12 carbon atoms and having at least one double bond. The straight chain or branched chain alkenylene group having 2 to 6 carbon atoms is preferable, and the straight chain or branched chain alkenylene group having 2 to 4 carbon atoms is more preferable. Examples thereof are vinylene group, propenylene group, butadienylene group, etc. If necessary, the alkylene group may optionally be substituted by 1 to 4 substituents as mentioned below, if necessary. Where alkenylene groups as defined above attach at two different carbon atoms of the benzene ring, they form an annelated five, six or seven membered carbocycle (e.g., a fused benzene ring) together with the carbon atoms to which they are attached, and may optionally be substituted by one or more substituents defined below.

The term “**alkynyl group**” means a straight or branched monovalent hydrocarbon chain having at least one triple bond. The preferable alkynyl group is a straight chain or branched chain alkynyl group having 2 to 6 carbon atoms, and the straight chain or branched chain alkynyl group having 2 to 4 carbon atoms is more preferable. Examples thereof are 2-propynyl group, 3-butynyl group, 2-butynyl group, 4-pentynyl group, 3-pentynyl group, 2-hexynyl group, 3-hexynyl group, 2-heptynyl group, 3-heptynyl group, 4-heptynyl group, 3-octynyl group, 3-nonyl group, 4-decynyl group, 3-undecynyl group, 4-

dodecynyl group, etc. The alkynyl group may optionally and independently be substituted by 1 to 4 substituents as mentioned below, if necessary.

The term “**cycloalkyl group**” means a monocyclic or bicyclic monovalent saturated hydrocarbon ring having 3 to 12 carbon atoms, and the
5 monocyclic saturated hydrocarbon group having 3 to 7 carbon atoms is more preferable. Examples thereof are a monocyclic alkyl group and a bicyclic alkyl group such as cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group, cyclooctyl group, cyclodecyl group, etc. These groups may optionally and independently be substituted by 1 to 4
10 substituents as mentioned below, if necessary. The cycloalkyl group may optionally be condensed with a saturated hydrocarbon ring or an unsaturated hydrocarbon ring (said saturated hydrocarbon ring and unsaturated hydrocarbon ring may optionally contain an oxygen atom, a nitrogen atom, a sulfur atom, SO or SO₂ within the ring, if necessary), and the condensed
15 saturated hydrocarbon ring and the condensed unsaturated hydrocarbon ring may be optionally and independently be substituted by 1 to 4 substituents as mentioned below.

The term “**cycloalkylidene group**” means a monocyclic or bicyclic divalent saturated hydrocarbon ring having 3 to 12 carbon atoms, and the
20 monocyclic saturated hydrocarbon group having 3 to 6 carbon atoms is preferable. Examples thereof are a monocyclic alkylidene group and a bicyclic alkylidene group such as cyclopropylidene group, cyclobutylidene group, cyclopentylidene group, cyclohexylidene group, etc. These groups may optionally and independently be substituted by 1 to 4 substituents as mentioned
25 below, if necessary. Besides, the cycloalkylidene group may optionally be condensed with a saturated hydrocarbon ring or an unsaturated hydrocarbon ring (said saturated hydrocarbon ring and unsaturated hydrocarbon ring may optionally contain an oxygen atom, a nitrogen atom, a sulfur atom, SO or SO₂ within the ring, if necessary), and the condensed saturated hydrocarbon ring
30 and the unsaturated hydrocarbon ring may be optionally and independently be substituted by 1 to 4 substituents as mentioned below.

The term “**cycloalkenyl group**” means a monocyclic or bicyclic monovalent unsaturated hydrocarbon ring having 4 to 12 carbon atoms and having at least one double bond. The preferable cycloalkenyl group is a monocyclic unsaturated hydrocarbon group having 4 to 7 carbon atoms.

- 5 Examples thereof are monocyclic alkenyl groups such as cyclopentenyl group, cyclopentadienyl group, cyclohexenyl group, etc. These groups may optionally and independently be substituted by 1 to 4 substituents as mentioned below, if necessary. Besides, the cycloalkenyl group may optionally be condensed with a saturated hydrocarbon ring or an unsaturated hydrocarbon ring (said
- 10 saturated hydrocarbon ring and unsaturated hydrocarbon ring may optionally contain an oxygen atom, a nitrogen atom, a sulfur atom, SO or SO₂ within the ring, if necessary), and the condensed saturated hydrocarbon ring and the unsaturated hydrocarbon ring may be optionally and independently be substituted by 1 to 4 substituents as mentioned below.

- 15 The term “**cycloalkynyl group**” means a monocyclic or bicyclic unsaturated hydrocarbon ring having 6 to 12 carbon atoms, and having at least one triple bond. The preferable cycloalkynyl group is a monocyclic unsaturated hydrocarbon group having 6 to 8 carbon atoms. Examples thereof are monocyclic alkynyl groups such as cyclooctynyl group, cyclodecynyl group.
- 20 These groups may optionally be substituted by 1 to 4 substituents as mentioned below, if necessary. Besides, the cycloalkynyl group may optionally and independently be condensed with a saturated hydrocarbon ring or an unsaturated hydrocarbon ring (said saturated hydrocarbon ring and unsaturated hydrocarbon ring may optionally contain an oxygen atom, a
- 25 nitrogen atom, a sulfur atom, SO or SO₂ within the ring, if necessary), and the condensed saturated hydrocarbon ring or the unsaturated hydrocarbon ring may be optionally and independently be substituted by 1 to 4 substituents as mentioned below.

- The term “**aryl group**” means a monocyclic or bicyclic monovalent aromatic hydrocarbon group having 6 to 10 carbon atoms. Examples thereof
- 30 are phenyl group, naphthyl group (including 1-naphthyl group and 2-naphthyl group). These groups may optionally and independently be substituted by 1 to

4 substituents as mentioned below, if necessary. Besides, the aryl group may optionally be condensed with a saturated hydrocarbon ring or an unsaturated hydrocarbon ring (said saturated hydrocarbon ring and unsaturated hydrocarbon ring may optionally contain an oxygen atom, a nitrogen atom, a sulfur atom, SO or SO₂ within the ring, if necessary), and the condensed saturated hydrocarbon ring or the unsaturated hydrocarbon ring may be optionally and independently be substituted by 1 to 4 substituents as mentioned below.

The term “**unsaturated monocyclic heterocyclic ring**” means an unsaturated hydrocarbon ring containing 1-4 heteroatoms independently selected from a nitrogen atom, an oxygen atom and a sulfur atom, and the preferable one is a 4- to 7-membered saturated or unsaturated hydrocarbon ring containing 1-4 heteroatoms independently selected from a nitrogen atom, an oxygen atom and a sulfur atom. Examples thereof are pyridine, pyrimidine, pyrazine, furan, thiophene, pyrrole, imidazole, pyrazole, oxazole, isoxazole, 4,5-dihydrooxazole, thiazole, isothiazole, thiadiazole, triazole, tetrazole, etc. Among them, pyridine, pyrimidine, pyrazine, furan, thiophene, pyrrole, imidazole, oxazole, and thiazole can be preferably used. The “unsaturated monocyclic heterocyclic ring” may optionally and independently be substituted by 1-4 substituents as mentioned below, if necessary.

The term “**unsaturated fused heterobicyclic ring**” means hydrocarbon ring comprised of a saturated or a unsaturated hydrocarbon ring condensed with the above mentioned unsaturated monocyclic heterocyclic ring where said saturated hydrocarbon ring and said unsaturated hydrocarbon ring may optionally contain an oxygen atom, a nitrogen atom, a sulfur atom, SO, or SO₂ within the ring, if necessary. The “unsaturated fused heterobicyclic ring” includes, for example, benzothiophene, indole, tetrahydrobenzothiophene, benzofuran, isoquinoline, thienothiophene, thienopyridine, quinoline, indoline, isoindoline, benzothiazole, benzoxazole, indazole, dihydroisoquinoline, etc. Further, the “heterocyclic ring” also includes possible N- or S-oxides thereof.

The term “**heterocyclyl**” means a monovalent group of the above-mentioned unsaturated monocyclic heterocyclic ring or unsaturated fused

heterobicyclic ring and a monovalent group of the saturated version of the above-mentioned unsaturated monocyclic heterocyclic or unsaturated fused heterobicyclic ring. If necessary, the heterocyclyl may optionally and independently be substituted by 1 to 4 substituents as mentioned below.

5 The term “**alkanoyl group**” means a formyl group and ones formed by binding an “alkyl group” to a carbonyl group.

 The term “**alkoxy group**” means ones formed by binding an “alkyl group” to an oxygen atom.

 The **substituent** for the above each group includes, for example, a
 10 halogen atom (fluorine, chlorine, bromine), a nitro group, a cyano group, an oxo group, a hydroxy group, a mercapto group, a carboxyl group, a sulfo group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkynyl group, an aryl group, a heterocyclyl group, an alkoxy group, an alkenyloxy group, an
 15 alkynyloxy group, a cycloalkyloxy group, a cycloalkenyloxy group, a cycloalkynyloxy group, an aryloxy group, a heterocyclioxy group, an alkanoyl group, an alkenylcarbonyl group, an alkynylcarbonyl group, a cycloalkylcarbonyl group, a cycloalkenylcarbonyl group, a cycloalkynylcarbonyl group, an arylcarbonyl group, a hetero-cyclylcarbonyl group, an alkoxy-
 20 carbonyl group, an alkenyloxy-carbonyl group, an alkynyloxy-carbonyl group, a cycloalkyloxy-carbonyl group, a cycloalkenyl-oxy-carbonyl group, a cycloalkynyl-oxycarbonyl group, an aryloxycarbonyl group, a hetero-cyclioxycarbonyl group, an alkanoyloxy group, an alkenyl-carbonyloxy group, an alkynyl-carbonyloxy group, a cycloalkyl-carbonyloxy group, a cycloalkenyl-carbonyloxy group, a cycloalkynyl-carbonyloxy group, an arylcarbonyloxy group, a hetero-cyclylcarbonyloxy group, an alkylthio group, an alkenyl-thio group, an alkynylthio group, a cycloalkylthio group, a cycloalkenyl-thio group, a cycloalkynylthio group, an arylthio group, a heterocyclylthio group, an amino group, a mono- or di-alkyl-amino group, a mono- or di-alkanoylamino group, a
 25 mono- or di-alkoxy-carbonyl-amino group, a mono- or di-arylcarbonyl-amino group, an alkylsulfinylamino group, an alkyl-sulfonyl-amino group, an arylsulfinylamino group, an arylsulfonylamino group, a carbamoyl group, a
 30

mono- or di-alkyl-carbamoyl group, a mono- or di-arylcarbamoyl group, an alkylsulfinyl group, an alkenyl-sulfinyl group, an alkynylsulfinyl group, a cycloalkyl-sulfinyl group, a cycloalkenylsulfinyl group, a cycloalkynyl-sulfinyl group, an arylsulfinyl group, a heterocyclyl-sulfinyl group, an alkyl-sulfonyl group, an alkenylsulfonyl group, an alkynylsulfonyl group, a cycloalkylsulfonyl group, a cycloalkenyl-sulfonyl group, a cycloalkynylsulfonyl group, an aryl-sulfonyl group, and a heterocyclylsulfonyl group. Each group as mentioned above may optionally be substituted by these substituents.

Further, the terms such as a haloalkyl group, a halo-lower alkyl group, a haloalkoxy group, a halo-lower alkoxy group, a halophenyl group, or a haloheterocyclyl group mean an alkyl group, a lower alkyl group, an alkoxy group, a lower alkoxy group, a phenyl group or a heterocyclyl group (hereinafter, referred to as an alkyl group, etc.) being substituted by one or more halogen atoms, respectively. Preferable ones are an alkyl group, etc. being substituted by 1 to 7 halogen atoms, and more preferable ones are an alkyl group, etc. being substituted by 1 to 5 halogen atoms. Similarly, the terms such as a hydroxyalkyl group, a hydroxy-lower alkyl group, a hydroxyalkoxy group, a hydroxy-lower alkoxy group and a hydroxyphenyl group mean an alkyl group, etc., being substituted by one or more hydroxy groups. Preferable ones are an alkyl group, etc., being substituted by 1 to 4 hydroxy groups, and more preferable ones are an alkyl group, etc., being substituted by 1 to 2 hydroxy groups. Further, the terms such as an alkoxyalkyl group, a lower alkoxyalkyl group, an alkoxy-lower alkyl group, a lower alkoxy-lower alkyl group, an alkoxyalkoxy group, a lower alkoxyalkoxy group, an alkoxy-lower alkoxy group, a lower alkoxy-lower alkoxy group, an alkoxyphenyl group, and a lower alkoxyphenyl group means an alkyl group, etc., being substituted by one or more alkoxy groups. Preferable ones are an alkyl group, etc., being substituted by 1 to 4 alkoxy groups, and more preferable ones are an alkyl group, etc., being substituted by 1 to 2 alkoxy groups.

The terms "**arylalkyl**" and "**arylalkoxy**" as used alone or as part of another group refer to alkyl and alkoxy groups as described above having an aryl substituent.

The term “**lower**” used in the definitions for the formulae in the present specification means a straight or branched carbon chain having 1 to 6 carbon atoms, unless defined otherwise. More preferably, it means a straight or branched carbon chain having 1 to 4 carbon atoms.

5

The term “**prodrug**” means an ester or carbonate, which is formed by reacting one or more hydroxy groups of the compound of the formula I with an acylating agent substituted by an alkyl, an alkoxy or an aryl by a conventional method to produce acetate, pivalate, methylcarbonate, benzoate, etc. Further, the prodrug includes also an ester or amide, which is similarly formed by reacting one or more hydroxy groups of the compound of the formula I with an α -amino acid or a β -amino acid, etc. using a condensing agent by a conventional method.

The **pharmaceutically acceptable salt** of the compound of the formula I includes, for example, a salt with an alkali metal such as lithium, sodium, potassium, etc.; a salt with an alkaline earth metal such as calcium, magnesium, etc.; a salt with zinc or aluminum; a salt with an organic base such as ammonium, choline, diethanolamine, lysine, ethylenediamine, t-butylamine, t-octylamine, tris(hydroxymethyl)aminomethane, N-methyl glucosamine, triethanolamine and dehydroabietylamine; a salt with an inorganic acid such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, nitric acid, phosphoric acid, etc.; or a salt with an organic acid such as formic acid, acetic acid, propionic acid, oxalic acid, malonic acid, succinic acid, fumaric acid, maleic acid, lactic acid, malic acid, tartaric acid, citric acid, ethanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, etc.; or a salt with an acidic amino acid such as aspartic acid, glutamic acid, etc.

The compound of the present invention also includes a mixture of stereoisomers, or each pure or substantially pure isomer. For example, the present compound may optionally have one or more asymmetric centers at a carbon atom containing any one of substituents. Therefore, the compound of the formula I may exist in the form of enantiomer or diastereomer, or a mixture thereof. When the present compound (I) contains a double bond, the present

compound may exist in the form of geometric isomerism (cis-compound, trans-compound), and when the present compound (I) contains an unsaturated bond such as carbonyl, then the present compound may exist in the form of a tautomer, and the present compound also includes these isomers or a mixture thereof. The starting compound in the form of a racemic mixture, enantiomer or diastereomer may be used in the processes for preparing the present compound. When the present compound is obtained in the form of a diastereomer or enantiomer, they can be separated by a conventional method such as chromatography or fractional crystallization.

10 In addition, the present compound (I) includes an intramolecular salt, hydrate, solvate or polymorph thereof.

Examples of the optionally substituted unsaturated monocyclic heterocyclic ring of the present invention include an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-5 substituents selected from the group consisting of a halogen atom, a nitro group, a cyano group, an oxo group, a hydroxyl group, a mercapto group, a carboxyl group, a sulfo group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkynyl group, an aryl group, a heterocyclyl group, an alkoxy group, an alkenyloxy group, an alkynyloxy group, a cycloalkyloxy group, a cycloalkenyloxy group, a cycloalkynyloxy group, an aryloxy group, a heterocyclyloxy group, an alkanoyl group, an alkenylcarbonyl group, an alkynylcarbonyl group, a cycloalkylcarbonyl group, a cycloalkenylcarbonyl group, a cycloalkynylcarbonyl group, an arylcarbonyl group, a heterocyclylcarbonyl group, an alkoxycarbonyl group, an alkenyloxycarbonyl group, an alkynyloxycarbonyl group, a cycloalkyloxycarbonyl group, a cycloalkenyloxycarbonyl group, a cycloalkynyloxycarbonyl group, an aryloxycarbonyl group, a heterocyclyloxycarbonyl group, an alkanoyloxy group, an alkenylcarbonyloxy group, an alkynylcarbonyloxy group, a cycloalkylcarbonyloxy group, a cycloalkenylcarbonyloxy group, a cycloalkynylcarbonyloxy group, an arylcarbonyloxy group, a heterocyclylcarbonyloxy group, an alkylthio group, an alkenylthio group, an alkynylthio group, a cycloalkylthio group, a

cycloalkenylthio group, a cycloalkynylthio group, an arylthio group, a heterocyclylthio group, an amino group, a mono- or di-alkylamino group, a mono- or di-alkanoylamino group, a mono- or di-alkoxycarbonylamino group, a mono- or di-arylcarbonylamino group, an alkylsulfinylamino group, an
 5 alkylsulfonylamino group, an arylsulfinylamino group, an arylsulfonylamino group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, a mono- or di-arylcarbamoyl group, an alkylsulfinyl group, an alkenylsulfinyl group, an alkynylsulfinyl group, a cycloalkylsulfinyl group, a cycloalkenylsulfinyl group, a cycloalkynylsulfinyl group, an arylsulfinyl group, a heterocyclylsulfinyl group, an
 10 alkylsulfonyl group, an alkenylsulfonyl group, an alkynylsulfonyl group, a cycloalkylsulfonyl group, a cycloalkenylsulfonyl group, a cycloalkynylsulfonyl group, an arylsulfonyl group, and a heterocyclylsulfonyl group wherein each substituent may optionally be further substituted by these substituents.

Examples of the optionally substituted unsaturated fused heterobicyclic
 15 ring of the present invention include an unsaturated fused heterobicyclic ring which may optionally be substituted by 1-5 substituents selected from the group consisting of a halogen atom, a nitro group, a cyano group, an oxo group, a hydroxy group, a mercapto group, a carboxyl group, a sulfo group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a
 20 cycloalkylidene- methyl group, a cycloalkenyl group, a cycloalkynyl group, an aryl group, a heterocyclyl group, an alkoxy group, an alkenyloxy group, an alkynyloxy group, a cycloalkyloxy group, a cycloalkenyloxy group, a cycloalkynyloxy group, an aryloxy group, a heterocycliloxy group, an alkanoyl group, an alkenylcarbonyl group, an alkynylcarbonyl group, a
 25 cycloalkylcarbonyl group, a cycloalkenyl- carbonyl group, a cycloalkynyl- carbonyl group, an arylcarbonyl group, a heterocyclylcarbonyl group, an alkoxycarbonyl group, an alkenyloxycarbonyl group, an alkynyloxy- carbonyl group, a cycloalkyloxycarbonyl group, a cycloalkenyloxy- carbonyl group, a cycloalkynyloxycarbonyl group, an aryloxycarbonyl group, a
 30 heterocycliloxy carbonyl group, an alkanoyloxy group, an alkenylcarbonyloxy group, an alkynylcarbonyloxy group, a cyclo- alkylcarbonyloxy group, a cycloalkenylcarbonyloxy group, a cyclo- alkynylcarbonyloxy group, an

arylcarbonyloxy group, a heterocyclyl- carbonyloxy group, an alkylthio group,
 an alkenylthio group, an alkynylthio group, a cycloalkylthio group, a
 cycloalkenylthio group, a cycloalkynylthio group, an arylthio group, a
 heterocyclylthio group, an amino group, a mono- or di-alkylamino group, a
 5 mono- or di-alkanoyl- amino group, a mono- or di-alkoxycarbonylamino group,
 a mono- or di-arylcarbonylamino group, an alkylsulfinylamino group, an alkyl-
 sulfonylamino group, an arylsulfinylamino group, an arylsulfonylamino group, a
 carbamoyl group, a mono- or di-alkylcarbamoyl group, a mono- or di-
 arylcarbamoyl group, an alkylsulfinyl group, an alkenylsulfinyl group, an
 10 alkynylsulfinyl group, a cycloalkylsulfinyl group, a cyclo- alkenylsulfinyl group, a
 cycloalkynylsulfinyl group, an arylsulfinyl group, a heterocyclylsulfinyl group, an
 alkylsulfonyl group, an alkenylsulfonyl group, an alkynylsulfonyl group, a
 cycloalkylsulfonyl group, a cyclo- alkenylsulfonyl group, a cycloalkynylsulfonyl
 group, an arylsulfonyl group, and a heterocyclylsulfonyl group, wherein each
 15 substituent may optionally be further substituted by these substituents.

Examples of the optionally substituted benzene ring of the present
 invention include a benzene ring which may optionally be substituted by 1-5
 substituents selected from the group consisting of a halogen atom, a nitro
 group, a cyano group, a hydroxy group, a mercapto group, a carboxyl group, a
 20 sulfo group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl
 group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkynyl
 group, an aryl group, a heterocyclyl group, an alkoxy group, an alkenyloxy
 group, an alkynyloxy group, a cycloalkyloxy group, a cycloalkenyloxy group, a
 cycloalkynyloxy group, an aryloxy group, a heterocycliloxy group, an alkanoyl
 25 group, an alkenylcarbonyl group, an alkynylcarbonyl group, a
 cycloalkylcarbonyl group, a cycloalkenylcarbonyl group, a cycloalkynylcarbonyl
 group, an arylcarbonyl group, a heterocyclylcarbonyl group, an alkoxycarbonyl
 group, an alkenyloxycarbonyl group, an alkynyloxycarbonyl group, a
 cycloalkyloxycarbonyl group, a cycloalkenyloxycarbonyl group, a cycloalkynyl-
 30 oxy carbonyl group, an aryloxycarbonyl group, a heterocycliloxy carbonyl group,
 an alkanoyloxy group, an alkenylcarbonyloxy group, an alkynylcarbonyloxy
 group, a cycloalkylcarbonyloxy group, a cycloalkenylcarbonyloxy group, a

cycloalkynylcarbonyloxy group, an arylcarbonyloxy group, a heterocyclylcarbonyloxy group, an alkylthio group, an alkenylthio group, an alkynylthio group, a cycloalkylthio group, a cycloalkenylthio group, a cycloalkynylthio group, an arylthio group, a heterocyclylthio group, an amino group, a mono- or di-alkylamino group, a mono- or di-alkanoylamino group, a mono- or di-alkoxycarbonylamino group, a mono- or di-arylcarbonylamino group, an alkylsulfinylamino group, an alkylsulfonylamino group, an arylsulfinylamino group, an arylsulfonylamino group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, a mono- or di-arylcarbamoyl group, an alkylsulfinyl group, an alkenylsulfinyl group, an alkynylsulfinyl group, a cycloalkylsulfinyl group, a cycloalkenylsulfinyl group, a cycloalkynylsulfinyl group, an arylsulfinyl group, a heterocyclylsulfinyl group, an alkylsulfonyl group, an alkenylsulfonyl group, an alkynylsulfonyl group, a cycloalkylsulfonyl group, a cycloalkenylsulfonyl group, a cycloalkynylsulfonyl group, an arylsulfonyl group, a heterocyclylsulfonyl group, an alkylene group, an alkyleneoxy group, an alkylenedioxy group, and an alkenylene group wherein each substituent may optionally be further substituted by these substituents.

Moreover, examples of the optionally substituted benzene ring include a benzene ring substituted with an alkylene group to form an annelated carbocycle together with the carbon atoms to which they are attached, and also includes a benzene ring substituted with an alkenylene group to form an annelated carbocycle such as a fused benzene ring together with the carbon atoms to which they are attached.

Preferable examples of the optionally substituted unsaturated monocyclic heterocyclic ring include an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents selected from the group consisting of a halogen atom, a hydroxy group, an alkoxy group, an alkyl group, a haloalkyl group, a haloalkoxy group, a hydroxyalkyl group, an alkoxyalkyl group, an alkoxyalkoxy group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkyloxy group, an aryl group, an aryloxy group, an arylalkoxy group, a cyano group, a nitro group, an amino group, a mono- or di-alkylamino group, an

alkanoylamino group, an alkoxycarbonylamino group, a carboxyl group, an
alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group,
an alkanoyl group, an alkylsulfonylamino group, an arylsulfonylamino group, an
alkylsulfinyl group, an alkylsulfonyl group, an arylsulfonyl group, a heterocycl
5 group, and an oxo group.

Preferable examples of the optionally substituted unsaturated fused
heterobicyclic ring include an unsaturated fused heterobicyclic ring which may
optionally be substituted by 1-3 substituents independently selected from the
group consisting of a halogen atom, a hydroxy group, an alkoxy group, an alkyl
10 group, a haloalkyl group, a haloalkoxy group, a hydroxyalkyl group, an
alkoxyalkyl group, an alkoxyalkoxy group, an alkenyl group, an alkynyl group, a
cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cyclo-
alkyloxy group, an aryl group, an aryloxy group, an arylalkoxy group, a cyano
group, a nitro group, an amino group, a mono- or di-alkylamino group, an
15 alkanoylamino group, an alkoxycarbonylamino group, a carboxyl group, an
alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group,
an alkanoyl group, an alkylsulfonylamino group, an arylsulfonylamino group, an
alkylsulfinyl group, an alkylsulfonyl group, an arylsulfonyl group, a heterocycl
group, and an oxo group.

Preferable examples of the optionally substituted benzene ring include a
benzene ring which may optionally be substituted by 1-3 substituents selected
from the group consisting of a halogen atom, a hydroxy group, an alkoxy group,
an alkyl group, a haloalkyl group, a haloalkoxy group, a hydroxyalkyl group, an
alkoxyalkyl group, an alkoxyalkoxy group, an alkenyl group, an alkynyl group, a
25 cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a
cycloalkyloxy group, an aryl group, an aryloxy group, an arylalkoxy group, a
cyano group, a nitro group, an amino group, a mono- or di-alkylamino group, an
alkanoylamino group, an alkoxycarbonylamino group, a carboxyl group, an
alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group,
30 an alkanoyl group, an alkylsulfonylamino group, an arylsulfonylamino group, an
alkylsulfinyl group, an alkylsulfonyl group, an arylsulfonyl group, a heterocycl

group, an alkylene group, an alkyleneoxy group, an alkylendioxy group, and an alkenylene group.

In another preferable embodiment of the present invention, the optionally substituted unsaturated monocyclic heterocyclic ring is an

5 unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl

10 group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, an alkanoylamino group, an alkoxycarbonylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsulfonylamino group, a

15 phenylsulfonyl group, a heterocyclyl group, and an oxo group;

the optionally substituted unsaturated fused heterobicyclic ring is an unsaturated fused heterobicyclic ring which may optionally be substituted by 1-3 substituents selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an

20 alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, an alkanoylamino group, an alkoxycarbonylamino group, a sulfamoyl group, a mono- or di-alkyl- sulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono-

25 or di-alkylcarbamoyl group, an alkanoyl group, an alkylsulfonylamino group, a phenyl group, a phenoxy group, a phenylsulfonylamino group, phenylsulfonyl group, a heterocyclyl group, and an oxo group; and

the optionally substituted benzene ring is a benzene ring which may optionally be substituted by 1-3 substituents, independently selected from the

30 group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio

group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, an alkanoylamino group, an alkoxycarbonylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsulfonylamino group, a phenylsulfonyl group, a heterocyclyl group, an alkylene group, and an alkenylene group;

wherein each of the above-mentioned substituents on the unsaturated monocyclic heterocyclic ring, the unsaturated fused heterobicyclic ring and the benzene ring may further be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, an alkyl group, a haloalkyl group, an alkoxy group, a haloalkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, a mono- or di-alkylamino group, a carboxyl group, an alkoxycarbonyl group, a phenyl group, an alkyleneoxy group, an alkylenedioxy group, an oxo group, a carbamoyl group, and a mono- or di-alkylcarbamoyl group.

In a preferable embodiment, the optionally substituted unsaturated monocyclic heterocyclic ring is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a cyano group, an alkyl group, an alkoxy group, an alkanoyl group, a mono- or di-alkylamino group, an alkanoylamino group, an alkoxycarbonylamino group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, a phenyl group, a heterocyclyl group, and an oxo group;

the optionally substituted unsaturated fused heterobicyclic ring is an unsaturated fused heterobicyclic ring which may optionally be substituted by 1-3 substituents independently selected from the group consisting of a halogen atom, a cyano group, an alkyl group, an alkoxy group, an alkanoyl group, a mono- or di-alkylamino group, an alkanoylamino group, an alkoxycarbonylamino group, a carboxy group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, a phenyl group, a heterocyclyl group, and an oxo group; and

the optionally substituted benzene ring is a benzene ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a cyano group, an alkyl group, an alkoxy group, an alkanoyl group, a mono- or di-alkylamino group, an alkanoylamino group, an alkoxycarbonylamino group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, a phenyl group, a heterocyclyl group, an alkylene group, and an alkenylene group;

wherein each of the above-mentioned substituents on the unsaturated monocyclic heterocyclic ring, the unsaturated fused heterobicyclic ring and the benzene ring may further be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a cyano group, an alkyl group, a haloalkyl group, an alkoxy group, a haloalkoxy group, an alkanoyl group, a mono- or di-alkylamino group, a carboxyl group, a hydroxy group, a phenyl group, an alkylendioxy group, an alkyleneoxy group, an alkoxycarbonyl group, a carbamoyl group and a mono- or di-alkylcarbamoyl group.

In another preferable embodiment,

(1) Ring A is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsulfonylamino group, a phenylsulfonyl group, a heterocyclyl group, and an oxo group, and

Ring B is an unsaturated monocyclic heterocyclic ring, an unsaturated fused heterobicyclic ring, or a benzene ring, each of which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a

cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsufonylamino group, a phenylsulfonyl group, a heterocyclyl group, an alkylene group, and an alkenylene group;

(2) Ring A is a benzene ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, an alkanoylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsufonylamino group, a phenylsulfonyl group, a heterocyclyl group, an alkylene group, and an alkenylene group, and

Ring B is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring, each of which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsufonylamino group, a phenylsulfonyl group, a heterocyclyl group, an alkylene group and an oxo group; or

(3) Ring A is an unsaturated fused heterobicyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the

group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono-
 5 or di-alkylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy group, a phenylsulfonylamino group, a phenylsulfonyl group, a heterocyclyl group, and an oxo group, and

10 Ring B is an unsaturated monocyclic heterocyclic ring, an unsaturated fused heterobicyclic ring, or a benzene ring, each of which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a hydroxy group, a cyano group, a nitro group, an alkyl group, an alkenyl group, an alkynyl group, a cycloalkyl group, a
 15 cycloalkylidenemethyl group, an alkoxy group, an alkanoyl group, an alkylthio group, an alkylsulfonyl group, an alkylsulfinyl group, an amino group, a mono- or di-alkylamino group, a sulfamoyl group, a mono- or di-alkylsulfamoyl group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkylsufonylamino group, a phenyl group, a phenoxy
 20 group, a phenylsulfonylamino group, a phenylsulfonyl group, a heterocyclyl group, an alkylene group and an oxo group;

wherein each of the above-mentioned substituents on Ring A and Ring B may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a cyano group, an alkyl group, a
 25 haloalkyl group, an alkoxy group, a haloalkoxy group, an alkanoyl group, a mono- or di-alkylamino group, a carboxyl group, a hydroxy group, a phenyl group, an alkylenedioxy group, an alkyleneoxy group, an alkoxycarbonyl group, a carbamoyl group and a mono- or di-alkylcarbamoyl group.

In a more preferable embodiment of the present invention, Ring A and
 30 Ring B are

(1) Ring A is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by a halogen atom, a lower alkyl group, a halo-lower

alkyl group, a lower alkoxy group, or an oxo group, and Ring B is (a) a benzene ring which may optionally be substituted by a halogen atom; a cyano group; a lower alkyl group; a halo-lower alkyl group; a lower alkoxy group; a halo-lower alkoxy group; a mono- or di-lower alkylamino group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; (b) an unsaturated monocyclic heterocyclic ring which may optionally be substituted by a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a phenyl group which may be substituted with a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; and a heterocyclyl group which may optionally be substituted with a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; or (c) an unsaturated fused heterobicyclic ring which may optionally be substituted by a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a phenyl group which may be substituted with a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; and a heterocyclyl group which may optionally be substituted with a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group;

(2) Ring A is a benzene ring which may optionally be substituted by a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a phenyl group, or a lower alkenylene group, and Ring B is (a) an unsaturated monocyclic heterocyclic ring which may optionally be substituted by a halogen atom; a cyano group; a lower alkyl group; a halo-lower alkyl

group; a phenyl-lower alkyl group; a lower alkoxy group; a halo-lower alkoxy group; a mono- or di-lower alkylamino group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a mono- or di-lower alkylamino group, or a carbamoyl group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a mono- or di-lower alkylamino group or a carbamoyl group; (b) an unsaturated fused heterobicyclic ring which may optionally be substituted by a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a phenyl-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a phenyl group which may be substituted with a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; and a heterocyclyl group which may optionally be substituted with a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; or

(3) Ring A is an unsaturated fused heterobicyclic ring which may optionally be substituted by a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or an oxo group, and Ring B is (a) a benzene ring which may optionally be substituted by a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a phenyl group which may be substituted with a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; and a heterocyclyl group which may optionally be substituted with a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; (b) an unsaturated monocyclic heterocyclic ring which may optionally be substituted by a halogen atom; a cyano group; a lower alkyl group; a halo-lower alkyl group; a lower alkoxy group; a halo-lower alkoxy group; a mono- or di-lower alkylamino group; a phenyl group optionally

substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; or (c) an unsaturated fused heterobicyclic ring which may optionally be substituted by a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a phenyl group which may be substituted with a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group; and a heterocyclyl group which may optionally be substituted with a group selected from a halogen atom, cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a mono- or di-lower alkylamino group.

15 In another more preferable embodiment, Y is $-\text{CH}_2-$ and is linked at the 3-position of Ring A, with respect to X being the 1-position, Ring A is a benzene ring which is substituted by 1-3 substituents selected from the group consisting of a lower alkyl group, a halo-lower alkyl group, a halogen atom, a lower alkoxy group, a phenyl group, and a lower alkenylene group, and Ring B is an
20 unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring, each of which may be substituted by 1-3 substituents selected from the group consisting of a lower alkyl group, a halo-lower alkyl group, a phenyl-lower alkyl group, a halogen atom, a lower alkoxy group, a halo-lower alkoxy group, a phenyl group, a halophenyl group, a cyanophenyl group, a lower alkylphenyl group, a halo-lower alkylphenyl group, a lower alkoxyphenyl group, a halo-lower alkoxy phenyl group, a lower alkylendioxyphenyl group, a lower
25 alkyleneoxy phenyl group, a mono- or di-lower alkylaminophenyl group, a carbamoyl phenyl group, a mono- or di-lower alkylcarbamoylphenyl group, a heterocyclyl group, a haloheterocyclyl group, a cyanoheterocyclyl group, a
30 lower alkylheterocyclyl group, a lower alkoxyheterocyclyl group, a mono- or di-lower alkylaminoheterocyclyl group, a carbamoylheterocyclyl group, and a mono- or di-lower alkylcarbamoyl group.

In another more preferable embodiment, Y is $-\text{CH}_2-$ and is linked at the 3-position of Ring A, with respect to X being the 1-position, Ring A is an unsaturated monocyclic heterocyclic ring which may be substituted by 1-3 substituents selected from the group consisting of a lower alkyl group, a halogen atom, a lower alkoxy group, and an oxo group, and Ring B is a benzene ring which may be substituted by 1-3 substituents selected from the group consisting of a lower alkyl group, a halo-lower alkyl group, a halogen atom, a lower alkoxy group, a halo-lower alkoxy group, a phenyl group, a halophenyl group, a cyanophenyl group, a lower alkylphenyl group, a halo-lower alkylphenyl group, a lower alkoxyphenyl group, a heterocyclyl group, a haloheterocyclyl group, a cyanoheterocyclyl group, a lower alkylheterocyclyl group, and a lower alkoxyheterocyclyl group.

Further, in another preferable embodiment, Y is $-\text{CH}_2-$ and is linked at the 3-position of Ring A, with respect to X being the 1-position, Ring A is an unsaturated monocyclic heterocyclic ring which may be substituted by 1-3 substituents selected from the group consisting of a lower alkyl group, a halogen atom, a lower alkoxy group, and an oxo group, and Ring B is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring, each of which may be substituted by 1-3 substituents selected from the group consisting of a lower alkyl group, a halo-lower alkyl group, a halogen atom, a lower alkoxy group, a halo-lower alkoxy group, a phenyl group, a halophenyl group, a cyanophenyl group, a lower alkylphenyl group, a halo-lower alkylphenyl group, a lower alkoxyphenyl group, a halo-lower alkoxyphenyl group, a heterocyclyl group, a haloheterocyclyl group, a cyanoheterocyclyl group, a lower alkylheterocyclyl group, and a lower alkoxyheterocyclyl group.

In a more preferable embodiment of the present invention, X is a carbon atom and Y is $-\text{CH}_2-$.

Further, in another preferable embodiment, Ring A and Ring B are:

(1) Ring A is a benzene ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a lower alkyl group optionally substituted by a halogen atom or a lower alkoxy group, a lower alkoxy group optionally substituted by a halogen atom or

a lower alkoxy group, a cycloalkyl group, a cycloalkoxy group, a phenyl group, and a lower alkenylene group, and

Ring B is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring, each of which may optionally be substituted by 1-3
5 substituents, independently selected from the group consisting of a halogen atom; a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group or a phenyl group; a lower alkoxy group optionally substituted by a halogen atom or a lower alkoxy group; a cycloalkyl group; a cycloalkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group,
10 a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, or a carbamoyl group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group or a carbamoyl group; and an oxo group,

15 (2) Ring A is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a lower alkyl group optionally substituted by a lower alkoxy group, a lower alkoxy group optionally substituted by a halogen atom or a lower alkoxy group, a cycloalkyl group, a cycloalkoxy group,
20 and an oxo group, and

Ring B is a benzene ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom; a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group or a phenyl group; a lower alkoxy group optionally substituted by
25 a halogen atom or a lower alkoxy group; a cycloalkyl group; a cycloalkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group; a lower alkenylene group,
30

(3) Ring A is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the

group consisting of a halogen atom, a lower alkyl group optionally substituted by a halogen atom or a lower alkoxy group, a lower alkoxy group optionally substituted by a halogen atom or a lower alkoxy group, a cycloalkyl group, a cycloalkoxy group, and an oxo group,

5 Ring B is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring, each of which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom; a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group or a phenyl group; a lower alkoxy group optionally substituted by
10 a halogen atom or a lower alkoxy group; a cycloalkyl group; a cycloalkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower
15 alkoxy group or a halo-lower alkoxy group; and an oxo group;

(4) Ring A is an unsaturated fused heterobicyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a lower alkyl group optionally substituted by a lower alkoxy group, a lower alkoxy group optionally substituted by a
20 halogen atom or a lower alkoxy group, a cycloalkyl group, a cycloalkoxy group, and an oxo group,

Ring B is a benzene ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom; a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group or a phenyl group; a lower alkoxy group optionally substituted by
25 a halogen atom or a lower alkoxy group; a cycloalkyl group; a cycloalkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower
30 alkoxy group or a halo-lower alkoxy group; and a lower alkylene group, or

(5) Ring A is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen atom, a lower alkyl group optionally substituted by a lower alkoxy group, a lower alkoxy group optionally substituted by a
5 halogen atom or a lower alkoxy group, a cycloalkyl group, a cycloalkoxy group, and an oxo group,

Ring B is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring, each of which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen
10 atom; a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group or a phenyl group; a lower alkoxy group optionally substituted by a halogen atom or a lower alkoxy group; a cycloalkyl group; a cycloalkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-
15 lower alkoxy group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group; and an oxo group.

In another preferable embodiment of the present invention, Y is linked at the 3-position of Ring A, with respect to X being the 1-position, Ring A is a
20 benzene ring which may optionally be substituted by a halogen atom, a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group, or a phenyl group, and Ring B is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring which may optionally be substituted by 1-3 substituents, independently selected from the group consisting of a halogen
25 atom; a lower alkyl group optionally substituted by a halogen atom or a phenyl group; a lower alkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, or a lower alkoxy group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, or a lower alkoxy
30 group; and an oxo group.

In another more preferable embodiment of the present invention, Y is linked at the 3-position of Ring A, with respect to X being the 1-position, Ring A

is an unsaturated monocyclic heterocyclic ring which may optionally be substituted by a substituent selected from a halogen atom, a lower alkyl group, and an oxo group, and Ring B is a benzene ring which may optionally be substituted by a substituent selected from the group consisting of a halogen atom; a lower alkyl group optionally substituted by a halogen atom or a phenyl group; a lower alkoxy group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, or a lower alkoxy group; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, or a lower alkoxy group; and a lower alkylene group.

Preferable examples of unsaturated monocyclic heterocyclic ring include a 5- or 6-membered unsaturated heterocyclic ring containing 1 or 2 hetero atoms independently selected from a nitrogen atom, an oxygen atom, and a sulfur atom. More specifically, preferred are furan, thiophene, oxazole, isoxazole, triazole, tetrazole, pyrazole, pyridine, pyrimidine, pyrazine, dihydroisoxazole, dihydropyridine, and thiazole. Preferable unsaturated fused heterobicyclic ring includes a 9- or 10-membered unsaturated fused heterocyclic ring containing 1 to 4 hetero atoms independently selected from a nitrogen atom, an oxygen atom, and a sulfur atom. More specifically, preferred are indoline, isoindoline, benzothiazole, benzoxazole, indole, indazole, quinoline, isoquinoline, benzothiophene, benzofuran, thienothiophene, and dihydroisoquinoline.


In a more preferred embodiment of the present invention, Ring A is a benzene ring which may optionally be substituted by a substituent selected from the group consisting of a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, and a phenyl group, and Ring B is a heterocyclic ring selected from the group consisting of thiophene, furan, benzofuran, benzothiophene, and benzothiazole, wherein the heterocyclic ring may optionally be substituted by a substituent selected from the following group: a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a phenyl-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a phenyl group, a halophenyl group, a lower alkylphenyl group, a lower

alkoxyphenyl group, a thienyl group, a halothienyl group, a pyridyl group, a halopyridyl group, and a thiazolyl group.

In yet another preferred embodiment, Y is $-\text{CH}_2-$, Ring A is an unsaturated monocyclic heterocyclic ring or an unsaturated fused heterobicyclic ring selected from the group consisting of thiophene, dihydroisoquinoline, dihydroisoxazole, triazole, pyrazole, dihydropyridine, dihydroindole, indole, indazole, pyridine, pyrimidine, pyrazine, quinoline, and a isoindoline, wherein the heterocyclic ring may optionally substituted by a substituent selected from the following group: a halogen atom, a lower alkyl group, and an oxo group, and Ring B is a benzene ring which may optionally be substituted by a substituent selected from the following group: a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, and a halo-lower alkoxy group.

In a further preferred embodiment of the present invention, Ring A is a benzene ring which is substituted by a halogen atom or a lower alkyl group, and Ring B is thienyl group which is substituted by phenyl group or a heterocyclyl group in which said phenyl group and heterocyclyl group is substituted by 1-3 substituents selected from a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, and a halo-lower alkoxy group.


Further, in another aspect of the present invention, preferable examples of the compound of the formula I include a compound wherein Ring A is



wherein R^{1a} , R^{2a} , R^{3a} , R^{1b} , R^{2b} , and R^{3b} are each independently a hydrogen atom, a halogen atom, a hydroxy group, an alkoxy group, an alkyl group, a haloalkyl group, a haloalkoxy group, a hydroxyalkyl group, an alkoxyalkyl group, an alkoxyalkoxy group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkyloxy group, a phenyl group, a phenylalkoxy group, a cyano group, a

nitro group, an amino group, a mono- or di-alkylamino group, an alkanoylamino group, a carboxyl group, an alkoxy carbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkanoyl group, an alkylsulfonylamino group, a phenylsulfonylamino group, an alkylsulfinyl group, an alkylsulfonyl group, or a
 5 phenylsulfonyl group, and

Ring B is



wherein R^{4a} and R^{5a} are each independently a hydrogen atom; a halogen atom; a hydroxy group; an alkoxy group; an alkyl group; a haloalkyl group; a haloalkoxy group; a hydroxyalkyl group; an alkoxyalkyl group; a
 10 phenylalkyl group; an alkoxyalkoxy group; a hydroxyalkoxy group; an alkenyl group; an alkynyl group; a cycloalkyl group; a cycloalkylidenemethyl group; a cycloalkenyl group; a cycloalkyloxy group; a phenyloxy group; a phenylalkoxy group; a cyano group; a nitro group; an amino group; a mono- or di-alkylamino group; an alkanoylamino group; a carboxyl group; an alkoxy carbonyl group; a
 15 carbamoyl group; a mono- or di-alkylcarbamoyl group; an alkanoyl group; an alkylsulfonylamino group; a phenylsulfonylamino group; an alkylsulfinyl group; an alkylsulfonyl group; a phenylsulfonyl group; a phenyl group optionally substituted by a halogen atom, a cyano group, an alkyl group, a haloalkyl group, an alkoxy group, a haloalkoxy group, an alkylenedioxy group, an
 20 alkyleneoxy group, a mono- or di-alkylamino group, a carbamoyl group, or a mono- or di-alkylcarbamoyl group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, an alkyl group, a haloalkyl group, an alkoxy group, a haloalkoxy group, a carbamoyl group, or a mono- or di-alkylcarbamoyl group, or R^{4a} and R^{5a} are bonded to each other at the terminals thereof to form an alkylene group; and

R^{4b} , R^{5b} , R^{4c} and R^{5c} are each independently a hydrogen atom; a halogen atom; a hydroxy group; an alkoxy group; an alkyl group; a haloalkyl group; a haloalkoxy group; a hydroxyalkyl group; an alkoxyalkyl group; a
 30 phenylalkyl group; an alkoxyalkoxy group; a hydroxyalkoxy group; an alkenyl


group; an alkynyl group; a cycloalkyl group; a cycloalkylidenemethyl group; a cycloalkenyl group; a cycloalkyloxy group; a phenyloxy group; a phenylalkoxy group; a cyano group; a nitro group; an amino group; a mono- or di-alkylamino group; an alkanoylamino group; a carboxyl group; an alkoxycarbonyl group; a carbamoyl group; a mono- or di-alkylcarbamoyl group; an alkanoyl group; an alkylsulfonylamino group; a phenylsulfonylamino group; an alkylsulfinyl group; an alkylsulfonyl group; a phenylsulfonyl group; a phenyl group optionally substituted by a halogen atom, a cyano group, an alkyl group, a haloalkyl group, an alkoxy group, a haloalkoxy group, a methylenedioxy group, an ethyleneoxy group, or a mono- or di-alkylamino group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, an alkyl group, a haloalkyl group, an alkoxy group or a haloalkoxy group.

More preferred is a compound wherein R^{1a} , R^{2a} , R^{3a} , R^{1b} , R^{2b} , and R^{3b} are each independently a hydrogen atom, a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a phenyl group;

R^{4a} and R^{5a} are each independently a hydrogen atom; a halogen atom; a lower alkyl group; a halo-lower alkyl group; a phenyl-lower alkyl group; a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a methylenedioxy group, an ethyleneoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a lower alkoxy group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group, or R^{4a} and R^{5a} are bonded to each other at the terminals thereof to form a lower alkylene group; and

R^{4b} , R^{5b} , R^{4c} and R^{5c} are each independently a hydrogen atom, a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a halo-lower alkoxy group.

Further preferred is a compound in which Ring B is




30

wherein R^{4a} is a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a methylenedioxy group, an ethyleneoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a lower alkoxy group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group, and


R^{5a} is a hydrogen atom, or

R^{4a} and R^{5a} are bonded to each other at the terminals thereof to form a lower alkylene group.

Further more preferred is a compound in which Ring A is



wherein R^{1a} is a halogen atom, a lower alkyl group, or a lower alkoxy group, and R^{2a} and R^{3a} are hydrogen atoms; and Ring B is




wherein R^{4a} is a phenyl group optionally substituted by a substituent selected from the group consisting of a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group; or a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a lower alkoxy group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group, and R^{5a} is a hydrogen atom, and Y is $-CH_2-$.

In more preferable embodiment, R^{4a} is a phenyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a halo-lower alkoxy group; or a

heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group, or a lower alkoxy group.

In another preferable embodiment of the present invention, a preferable compound can be represented by the following formula IA:




5

wherein R^A is a halogen atom, a lower alkyl group or a lower alkoxy group; R^B is a phenyl group optionally substituted by 1-3 substituents selected from a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a methylenedioxy group, an ethyleneoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group; or a heterocyclyl group optionally substituted by 1-3 substituents selected from a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group; and R^C is hydrogen atom; or R^B and R^C taken together are a fused benzene ring which may be substituted by a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group.

In a preferable embodiment, R^A is a halogen atom or a lower alkyl group, R^C is hydrogen atom, and R^B is phenyl group substituted by 1-3 substituents selected from a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a methylenedioxy group, an ethyleneoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group; or a heterocyclyl group substituted by 1-3 substituents selected from the group consisting of a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-

lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group. The chemical structure of such compounds are represented by the following formula (IA'):



5 wherein R^A is a halogen atom, or a lower alkyl group, Ring C is a phenyl group substituted by 1-3 substituents selected from the group consisting of a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a methylenedioxy group, an ethyleneoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group; or a heterocyclyl group substituted by 1-3 substituents selected from the group consisting of a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, a mono- or di-lower alkylamino group, a carbamoyl group, and a mono- or di-lower alkylcarbamoyl group.


15 In a more preferable embodiment, Ring C is a phenyl group substituted by 1-3 substituents selected from the group consisting of a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, a halo-lower alkoxy group, and a mono- or di-lower alkylamino group; or a heterocyclyl group substituted by a substituent selected from the group consisting of a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, and a halo-lower alkoxy group.

25 Among them, a compound in which Ring C is a phenyl group substituted by a halogen atom, a cyano group, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group or a halo-lower alkoxy group; or a heterocyclyl group substituted by a halogen atom, a cyano group, a lower alkyl group, or a lower alkoxy group is preferred.


A preferred heterocyclyl group includes a 5- or 6-membered heterocyclyl group containing 1 or 2 hetero atoms independently selected from the group consisting of a nitrogen atom, an oxygen atom, and a sulfur atom, or a 9- or 10-membered heterocyclyl group containing 1 to 4 hetero atoms independently
 5 selected from the group consisting of a nitrogen atom, an oxygen atom, and a sulfur atom. Specifically, a thienyl group, a pyridyl group, a pyrimidyl group, a pyrazinyl group, pyrazolyl group, a thiazolyl group, a quinolyl group, a tetrazolyl group and an oxazolyl group are preferred.

In a further preferable embodiment, Ring C is a phenyl group substituted
 10 by a halogen atom or a cyano group, or a pyridyl group substituted by a halogen atom.

In another preferable embodiment of the present invention, preferred is a compound in which Ring A is




15 wherein R^{1a} is a halogen atom, a lower alkyl group, or a lower alkoxy group, and R^{2a} and R^{3a} are hydrogen atoms; and Ring B is




20 wherein R^{4b} and R^{5b} are each independently a hydrogen atom, a halogen atom, a lower alkyl group, a halo-lower alkyl group, a lower alkoxy group, or a halo-lower alkoxy group.

In another aspect of the present invention, preferable examples of the compound I include a compound represented by the following formula IB:




wherein R^8 , R^9 and R^{10} are each independently a hydrogen atom, a halogen atom, a hydroxy group, an alkoxy group, an alkyl group, a haloalkyl group, a haloalkoxy group, a hydroxyalkyl group, an alkoxyalkyl group, an alkoxyalkoxy group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkyloxy group, an aryloxy group, an arylalkoxy group, a cyano group, a nitro group, an amino group, a mono- or di-alkylamino group, an alkylcarbonylamino group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkanoyl group, an alkylsulfonylamino group, an arylsulfonylamino group, an alkylsulfinyl group, an alkylsulfonyl group, or an arylsulfonyl group; and

a group represented by:




is



wherein R^{6a} and R^{7a} are each independently a hydrogen atom, a halogen atom, a hydroxy group, an alkoxy group, an alkyl group, a haloalkyl group, a haloalkoxy group, a hydroxyalkyl group, an alkoxyalkyl group, an alkoxyalkoxy group, an alkenyl group, an alkynyl group, a cycloalkyl group, a cycloalkylidenemethyl group, a cycloalkenyl group, a cycloalkyloxy group, an


aryloxy group, an arylalkoxy group, a cyano group, a nitro group, an amino group, a mono- or di-alkylamino group, an alkylcarbonylamino group, a carboxyl group, an alkoxycarbonyl group, a carbamoyl group, a mono- or di-alkylcarbamoyl group, an alkanoyl group, an alkylsulfonylamino group, an
 5 arylsulfonylamino group, an alkylsulfinyl group, an alkylsulfonyl group, or an arylsulfonyl group and R^{6b} and R^{7b} are each independently a hydrogen atom, a halogen atom, an alkyl group, a haloalkyl group, or an alkoxy group.

Among the compounds represented by the formula IB, more preferred is a compound in which R^8 , R^9 and R^{10} are each independently a hydrogen atom,
 10 a halogen atom, a lower alkyl group, a cycloalkyl group, a hydroxy-lower alkyl group, a halo-lower alkyl group, a lower alkoxy-lower alkyl group, a lower alkoxy group, a cycloalkoxy group, a halo-lower alkoxy group, or a lower alkoxy-lower alkoxy group, and
 a group represented by:




15


is



wherein R^{6a} , R^{7a} are each independently a hydrogen atom, a halogen atom, a lower alkyl group, a cycloalkyl group, a hydroxy-lower alkyl group, a
 20 halo-lower alkyl group, a lower alkoxy-lower alkyl group, a lower alkoxy group, a cycloalkoxy group, a halo-lower alkoxy group, or a lower alkoxy-lower alkoxy group, or a group represented by:




is



wherein R^{6b} and R^{7b} are each independently a hydrogen atom, a halogen atom, a lower alkyl group, a halo-lower alkyl group, or a lower alkoxy group.

- 5 In another aspect of the present invention, preferable examples of the compound I include a compound represented by the following formula IC:



- wherein Ring B' is an optionally substituted benzene ring, an optionally substituted unsaturated monocyclic heterocyclic ring, or an optionally substituted unsaturated fused heterobicyclic ring.
- 10

- Preferable examples of Ring B' include a benzene ring and a heterocyclic ring, both of which may have a substituent(s) selected from the group consisting of a halogen atom; a cyano group; a lower alkyl group optionally substituted by a halogen atom; a lower alkoxy group optionally substituted by a halogen atom; a lower alkanoyl group; a mono- or di-lower alkylamino group; a lower alkoxycarbonyl group; a carbamoyl group; a mono- or di-lower alkylcarbamoyl group; a phenyl group optionally substituted by a substituent(s) selected from a halogen atom, a cyano group, a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group optionally substituted by a halogen atom, a lower alkanoyl group, a mono- or di-lower alkylamino group, a lower alkoxycarbonyl group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group; a heterocyclyl group optionally substituted by a substituent(s) selected from a halogen atom, a cyano group, a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy
- 15
- 20

group optionally substituted by a halogen atom, a lower alkanoyl group, a mono- or di-lower alkylamino group, a lower alkoxy carbonyl group, a carbamoyl group, or a mono- or di-lower alkylcarbamoyl group; an alkylene group; and an oxo group.

5 More preferable examples of Ring B' include a benzene ring which may be substituted by a substituent selected from the group consisting of a halogen atom; a cyano group; a lower alkyl group optionally substituted by a halogen atom; a lower alkoxy group optionally substituted by a halogen atom; a mono- or di-lower alkylamino group; a phenyl group optionally substituted by a
 10 halogen atom, a cyano group, a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group optionally substituted by a halogen atom; a heterocyclyl group optionally substituted by a halogen atom, a cyano group, a lower alkyl group optionally substituted by a halogen atom, a lower alkoxy group optionally substituted by a halogen atom.

15 Preferred compound of the present invention may be selected from the following group:

1-(β -D-glucopyranosyl)-4-chloro-3-(6-ethylbenzo[b]thiophen-2-ylmethyl)benzene;

20 1-(β -D-glucopyranosyl)-4-chloro-3-[5-(5-thiazolyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-chloro-3-(5-phenyl-2-thienyl-methyl)benzene;

1-(β -D-glucopyranosyl)-4-methyl-3-[5-(4-fluorophenyl)-2-thienylmethyl]benzene;

25 1-(β -D-glucopyranosyl)-4-chloro-3-[5-(2-pyrimidinyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-methyl-3-[5-(2-pyrimidinyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-chloro-3-[5-(3-cyanophenyl)-2-thienylmethyl]benzene;

30 1-(β -D-glucopyranosyl)-4-chloro-3-[5-(4-cyanophenyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-methyl-3-[5-(6-fluoro-2-pyridyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-chloro-3-[5-(6-fluoro-2-pyridyl)-2-thienylmethyl]benzene;

5 1-(β -D-glucopyranosyl)-4-methyl-3-[5-(3-difluoromethyl-phenyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-methyl-3-[5-(3-cyanophenyl)-2-thienylmethyl]benzene;

10 1-(β -D-glucopyranosyl)-4-methyl-3-[5-(4-cyanophenyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-chloro-3-[5-(6-fluoro-3-pyridyl)-2-thienylmethyl]benzene;

1-(β -D-glucopyranosyl)-4-fluoro-3-[5-(3-cyanophenyl)-2-thienylmethyl]benzene;

15 the pharmaceutically acceptable salt thereof; and the prodrug thereof.

Particularly preferred compounds of the present invention include:

1-(β -D-glucopyranosyl)-4-methyl-3-[5-(3-cyano-phenyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

20 1-(β -D-glucopyranosyl)-4-methyl-3-[5-(4-cyano-phenyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

1-(β -D-glucopyranosyl)-4-methyl-3-[5-(4-fluoro-phenyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

25 1-(β -D-glucopyranosyl)-4-chloro-3-[5-(3-cyano-phenyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

30 1-(β -D-glucopyranosyl)-4-methyl-3-[5-(6-fluoro-2-pyridyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

1-(β -D-glucopyranosyl)-4-chloro-3-[5-(6-fluoro-2-pyridyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof;

5 1-(β -D-glucopyranosyl)-4-chloro-3-[5-(6-fluoro-3-pyridyl)-2-thienylmethyl]benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof; and

1-(β -D-glucopyranosyl)-4-fluoro-3-(5-(3-cyanophenyl)-2-thienylmethyl)benzene, or a pharmaceutically acceptable salt thereof, or a prodrug thereof.

10

Abbreviations used in the specification, particularly the Schemes and Examples, are as follows:

AcOEt	=	Ethyl acetate
CPME	=	Cyclopentyl methyl ether
DI (water)	=	Deionized (water)
DMAP	=	4-Dimethylaminopyridine
HPLC	=	High Pressure Liquid Chromatography
IPA	=	Isopropyl Alcohol
2-Me-THF	=	2-Methyl-tetrahydrofuran
MPLC	=	Medium Pressure Liquid Chromatography
MTBE	=	Methyl-t-butyl Ether
n-BuLi	=	n-Butyl lithium
n-Bu ₂ O	=	di-(n-butyl) ether
Pd/C	=	Palladium on carbon
Pd(OAc) ₂ / Et ₃ SiH	=	Palladium acetate and triethylsilane
RaNi	=	RANEY [®] nickel (aluminum nickel alloy)
RBF	=	Round Bottom Flask
TEA	=	Triethylamine
THF	=	Tetrahydrofuran
TMEDA	=	Tetramethylethylenediamine
TMS	=	Trimethylsilyl
TMSBr	=	Trimethylsilyl bromide

TMSCH₂ = Trimethylsilyl-CH₂-

As used herein, unless otherwise noted, the term “**isolated form**” shall mean that the compound is present in a form which is separate from any solid mixture with another compound(s), solvent system or biological environment.

5 In an embodiment, the product prepared according to the process described herein (more particularly, a compound of formula (I), preferably a compound of formula (I-S) or compound of formula (I-K)) is prepared as an isolated form.

As used herein, unless otherwise noted, the term “**substantially pure**” shall mean that the mole percent of impurities in the isolated compound is less
10 than about 5 mole percent, preferably less than about 2 mole percent, more preferably, less than about 0.5 mole percent, most preferably, less than about 0.1 mole percent.

In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (I), wherein the compound of formula (I)
15 is substantially pure. In another embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-S), wherein the compound of formula (I-S) is substantially pure. In another embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-K), wherein the compound of formula (I-K) is substantially pure.

20 As used herein, unless otherwise noted, the term “**substantially free of a corresponding salt form(s)**” when used to described the compound of formula (I) shall mean that mole percent of the corresponding salt form(s) in the isolated base of formula (I) is less than about 5 mole percent, preferably less than about 2 mole percent, more preferably, less than about 0.5 mole percent,
25 most preferably less than about 0.1 mole percent.

In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (I), wherein the compound of formula (I) is substantially free of corresponding salt forms. In another embodiment, the present invention is directed to a process for the preparation of a compound of
30 formula (I-S), wherein the compound of formula (I-S) is substantially free of corresponding salt forms. In another embodiment, the present invention is

directed to a process for the preparation of a compound of formula (I-K), wherein the compound of formula (I-K) is substantially free of corresponding salt forms.

5 As used herein, unless otherwise noted, the terms “**treating**”, “**treatment**” and the like, shall include the management and care of a subject or patient (preferably mammal, more preferably human) for the purpose of combating a disease, condition, or disorder and includes the administration of a compound of the present invention to prevent the onset of the symptoms or
10 complications, alleviate the symptoms or complications, or eliminate the disease, condition, or disorder.

 As used herein, unless otherwise noted, the term “**prevention**” shall include (a) reduction in the frequency of one or more symptoms; (b) reduction in the severity of one or more symptoms; (c) the delay or avoidance of the
15 development of additional symptoms; and / or (d) delay or avoidance of the development of the disorder or condition.

 One skilled in the art will recognize that wherein the present invention is directed to methods of prevention, a subject in need of thereof (i.e. a subject in need of prevention) shall include any subject or patient (preferably a mammal,
20 more preferably a human) who has experienced or exhibited at least one symptom of the disorder, disease or condition to be prevented. Further, a subject in need thereof may additionally be a subject (preferably a mammal, more preferably a human) who has not exhibited any symptoms of the disorder, disease or condition to be prevented, but who has been deemed by a
25 physician, clinician or other medical profession to be at risk of developing said disorder, disease or condition. For example, the subject may be deemed at risk of developing a disorder, disease or condition (and therefore in need of prevention or preventive treatment) as a consequence of the subject's medical history, including, but not limited to, family history, pre-disposition, co-existing
30 (comorbid) disorders or conditions, genetic testing, and the like.

 The term “**subject**” as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment,

observation or experiment. Preferably, the subject has experienced and / or exhibited at least one symptom of the disease or disorder to be treated and / or prevented.

The term “**therapeutically effective amount**” as used herein, means that
5 amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term “**composition**” is intended to encompass a
10 product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

The compound of formula (I) of the present invention exhibits an
15 excellent inhibitory activity against sodium-dependent glucose transporter, and an excellent blood glucose lowering effect. Therefore, the compound of the present invention is useful for treating or delaying the progression or onset of diabetes mellitus, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, delayed wound healing, insulin resistance, hyperglycemia,
20 hyperinsulinemia, elevated blood levels of fatty acids, elevated blood levels of glycerol, hyperlipidemia, obesity, hypertriglyceridemia, Syndrome X, diabetic complications, atherosclerosis, or hypertension. In particular, the compound of the present invention is useful in the treatment or the prophylaxis of diabetes mellitus (type 1 and type 2 diabetes mellitus, etc.), diabetic complications (such
25 as diabetic retinopathy, diabetic neuropathy, diabetic nephropathy) or obesity, or is useful in the treatment of postprandial hyperglycemia.

The compound of formula (I) of the present invention or a
pharmaceutically acceptable salt thereof may be administered either orally or parenterally, and can be used in the form of a suitable pharmaceutical
30 preparation. Suitable pharmaceutical preparation for oral administration includes, for example, solid preparation such as tablets, granules, capsules, powders, etc., or solution preparations, suspension preparations, or emulsion

preparations, etc. Suitable pharmaceutical preparation for parenteral administration includes, for example, suppositories; injection preparations and intravenous drip preparations using distilled water for injection, physiological saline solution or aqueous glucose solution; or inhalant preparations.

5 The dosage of the present compound of formula (I) or a pharmaceutically acceptable salt thereof may vary according to the administration routes, ages, body weight, conditions of a patient, or kinds and severity of a disease to be treated, and it is usually in the range of about 0.01 to 300 mg/kg/day, or any amount or range therein, preferably in the range of
10 about 0.1 to 50 mg/kg/day, or any amount or range therein, preferably in the range of about 0.1 to 30 mg/kg/day, or any amount or range therein.

 The compound of the formula I may be used, if necessary, in combination with one or more of other antidiabetic agents, one or more agents for treating diabetic complications, and/or one or more agents for treatment of
15 other diseases. The present compound and these other agents may be administered in the same dosage form, or in a separate oral dosage form or by injection.

 The other antidiabetic agents include, for example, antidiabetic or antihyperglycemic agents including insulin, insulin secretagogues, or insulin
20 sensitizers, or other antidiabetic agents having an action mechanism different from SGLT inhibition, and 1, 2, 3 or 4 of these other antidiabetic agents may preferably be used. Concrete examples thereof are biguanide compounds, sulfonylurea compounds, α -glucosidase inhibitors, PPAR γ agonists (e.g., thiazolidinedione compounds), PPAR α/γ dual agonists, dipeptidyl peptidase IV
25 (DPP4) inhibitors, mitiglinide compounds, and/or nateglinide compounds, and insulin, glucagon-like peptide-1 (GLP-1), PTP1B inhibitors, glycogen phosphorylase inhibitors, RXR modulators, and/or glucose 6-phosphatase inhibitors.

 The agents for treatment of other diseases include, for example, an anti-
30 obesity agent, an antihypertensive agent, an antiplatelet agent, an anti-atherosclerotic agent and/or a hypolipidemic agent.

The SGLT inhibitors of the formula I may be used in combination with agents for treatment of diabetic complications, if necessary. These agents include, for example, PKC inhibitors and/or ACE inhibitors.

5 The dosage of those agents may vary according to ages, body weight, and conditions of patients, and administration routes, dosage forms, etc.

These pharmaceutical compositions may be orally administered to mammalian species including human beings, apes, dogs, etc., for example, in the dosage form of tablet, capsule, granule or powder, or parenterally administered in the form of injection preparation, or intranasally, or in the form
10 of transdermal patch.

One skilled in the art will recognize that, where not otherwise specified, the reaction step(s) is performed under suitable conditions, according to known methods, to provide the desired product.

15 One skilled in the art will further recognize that, in the specification and claims as presented herein, wherein a reagent or reagent class/type (e.g. base, solvent, etc.) is recited in more than one step of a process, the individual reagents are independently selected for each reaction step and may be the same or different from each other. For example wherein two steps of a process
20 recite an organic or inorganic base as a reagent, the organic or inorganic base selected for the first step may be the same or different than the organic or inorganic base of the second step. Further, one skilled in the art will recognize that wherein a reaction step of the present invention may be carried out in a variety of solvents or solvent systems, said reaction step may also be carried
25 out in a mixture of the suitable solvents or solvent systems.

Examples of suitable solvents, bases, reaction temperatures, and other reaction parameters and components are provided in the detailed descriptions which follows herein. One skilled in the art will recognize that the listing of said examples is not intended, and should not be construed, as limiting in any way
30 the invention set forth in the claims which follow thereafter.

To provide a more concise description, some of the quantitative expressions herein are recited as a range from about amount X to about

amount Y. It is understood that wherein a range is recited, the range is not limited to the recited upper and lower bounds, but rather includes the full range from about amount X through about amount Y, or any amount or range therein.

To provide a more concise description, some of the quantitative
5 expressions given herein are not qualified with the term “**about**”. It is understood that whether the term “about” is used explicitly or not, every quantity given herein is meant to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including approximations due
10 to the experimental and/or measurement conditions for such given value.

As used herein, unless otherwise noted, the term “**leaving group**” shall mean a charged or uncharged atom or group which departs during a substitution or displacement reaction. Suitable examples include, but are not limited to, Br, Cl, I, , tosylate, and the like, preferably the leaving group is a
15 halogen, such as Br, Cl or I.

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective
20 Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

As used herein, unless otherwise noted, the term “**nitrogen protecting**
25 **group**” shall mean a group which may be attached to a nitrogen atom to protect said nitrogen atom from participating in a reaction and which may be readily removed following the reaction. Suitable nitrogen protecting groups include, but are not limited to carbamates –containing groups of the formula –C(O)O-R wherein R is for example methyl, ethyl, t-butyl, benzyl, phenylethyl,
30 CH₂=CH-CH₂-, and the like; amides – containing groups of the formula –C(O)-R’ wherein R’ is for example methyl, phenyl, trifluoromethyl, and the like; N-sulfonyl derivatives – containing groups of the formula –SO₂-R” wherein R” is


for example tolyl, phenyl, trifluoromethyl, 2,2,5,7,8-pentamethylchroman-6-yl-, 2,3,6-trimethyl-4-methoxybenzene, and the like. Other suitable nitrogen protecting groups may be found in texts such as T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991.

5 As used herein, unless otherwise noted, the term “**oxygen protecting group**” shall mean a group which may be attached to a oxygen atom to protect said oxygen atom from participating in a reaction and which may be readily removed following the reaction. Suitable oxygen protecting groups include, but are not limited to, acetyl, benzoyl, pivaloyl, t-butyl-dimethylsilyl, trimethylsilyl
10 (TMS), MOM, THP, and the like. Other suitable oxygen protecting groups may be found in texts such as T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991.

Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be
15 separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt
20 formation with an optically active acid, such as (-)-di-p-toluoyl-D-tartaric acid and/or (+)-di-p-toluoyl-L-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds
25 may be resolved using a chiral HPLC column.

One skilled in the art will recognize that in any of the processes described herein, reactive substituents on the compounds of formula (I), such as hydroxy groups, oxo groups, carboxy groups, and the like, are preferably protected and subsequently de-protected, according to known methods, at
30 suitable points along the synthesis route.

The present invention is directed to a process for the preparation of compounds of formula (I) as outlined in Scheme 1, below.



Scheme 1

Accordingly, a suitably substituted compound of formula (V), wherein LG¹ is a suitably selected leaving group such as bromo, iodo, and the like, preferably LG¹ is bromo or iodo, a known compound or compound prepared by known methods, is reacted with a

mixture of a suitably selected zinc salt such as zinc dibromide (ZnBr₂), zinc diiodide (ZnI₂), zinc ditriflate, and the like, preferably ZnBr₂; or with an amine complex of zinc halide such as pyridine zinc bromide complex, N-methylmorpholine zinc bromide complex, and the like; wherein the zinc salt or amine complex of zinc halide is preferably present in an amount in the range of

from about 0.33 to about 3.0 molar equivalents, more preferably in an amount in the range of from about 0.33 to about 1.0 molar equivalents, more preferably in an amount of about 0.5 molar equivalents; and a suitably selected organo-lithium reagent such as trimethylsilylmethyl lithium, n-hexyl lithium, sec-butyl lithium, n-butyllithium, t-butyllithium, methyl lithium, and the like, preferably n-hexyl lithium or n-butyl lithium; wherein the organo-lithium reagent is preferably present in an amount in the range of from about 0.5 to about 2.0 molar equivalents, preferably in an amount in the range of from about 1.0 to about 1.2 molar equivalents;

10 in a suitably selected first hydrocarbon solvent, such as toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, preferably toluene; preferably at a temperature less than about room temperature, more preferably at a temperature in the range of from about -78°C to about room temperature; more preferably at about 0°C; to yield a mixture of the

15 corresponding compound of formula (VI), wherein M¹ is lithium and the zinc salt. Preferably, the compound of formula (VI) is not isolated.

In an embodiment of the present invention, the zinc salt is ZnBr₂ and the organo-lithium reagent is n-butyl-lithium. In another embodiment of the present invention, the molar ratio of the zinc salt to the organo-lithium reagent is about


20 1:2. In another embodiment of the present invention, the zinc salt and the organo-lithium reagent are pre-mixed; preferably for a period of time in the range of from about 1 to about 2 hours.

It has been discovered that (a) pre-mixing of the zinc salt and organo-lithium reagent and (b) adding the compound of formula (V) to the pre-mixed zinc salt and organo-lithium reagent, improves the reaction conditions for making the compound of formula (VII). More particularly, the use of the pre-mixed zinc salt and organo-lithium reagent mixture permits for the preparation of the compound of formula (VI) at a temperature of about 0°C. The higher

25 temperature is preferred, particularly for large scale manufacture. Additionally, the use of the pre-mixed zinc salt and organo-lithium reagent mixture results in improved yield of the compound of formula (IX).

30

To the mixture of the compound of formula (VI) and the zinc salt is admixed a suitably selected first ether solvent such as diethyl ether, diisopropyl ether, di-n-butyl ether, MTBE, cyclopentylmethyl ether, and the like, preferably
 5 di-n-butyl ether or cyclopentyl methyl ether; wherein the first ether solvent is preferably present in an amount in the range of from about 5% to about 15% by volume (relative to the total volume of the first hydrocarbon solvent and the first ether solvent), or any amount or range therein; preferably in an amount in the range of from about 7% to about 10%, or any amount or range therein; more
 10 preferably, at about amount in the range of from about 8% to about 9%, or any amount or range therein; (to minimize the formation of undesired by-product, more particularly to minimize formation of the compound of formula (U))



to yield the corresponding compound of formula (VII), wherein M^2 is a
 15 corresponding reactive zinc species, which compound is not isolated.

It is theorized that, when the zinc salt used in the previous reaction step is $ZnBr_2$, then in the compound of formula (VII), M^2 may include $ZnBr$; when the zinc salt used in the previous reaction step may include ZnI_2 , then in the compound of formula (VII), M^2 is ZnI ; when the zinc salt used in the previous
 20 reaction step is zinc ditriflate, then in the compound of formula (VII), M^2 may include zinc triflate. At this time, the intermediate of formula (VII) has not been isolated or fully characterized (particularly, as to the identity of the M^2 group).

It is further theorized that admixing the suitably selected first ether solvent with the mixture of the compound of formula (VI) and the zinc salt,
 25 preferably zinc bromide, results in chelation of the zinc salt. The chelated zinc bromide then reacts with the compound of formula (VI) to yield the corresponding compound of formula (VII).

The compound of formula (VII) is reacted with a suitably substituted
 30 compound of formula (VIII), wherein LG^2 is a suitably selected leaving group

such as bromo, chloro, iodo, and the like, preferably bromo; and wherein each Z is independently a suitably selected oxygen protecting group, for example Z may selected from the group consisting of benzyl, benzoyl, pivaloyl, isobutyryl, p-methoxy-benzyl, acetyl, propionyl, and the like, preferably, each Z protecting
5 group is the same, more preferably each Z is pivaloyl, a known compound or compound prepared by known methods; wherein the compound of formula (VIII) is preferably present in an amount in the range of from about 0.5 to about 3.0 molar equivalents, or any amount or range therein, more preferably in an amount in the range of from about 0.8 to about 1.25 molar equivalents, or any
10 amount or range therein, more preferably in an amount of about 1.0 to about 1.1 molar equivalents;

optionally in a mixture of a suitably selected second ether solvent and a suitably selected second hydrocarbon solvent, wherein the second ether solvent is for example, diethyl ether, di-n-butyl ether, MTBE, 2-methyl-THF, di-
15 isopropyl ether, cyclopentylmethyl ether, and the like, preferably di-n-butyl ether or cyclopentyl methyl ether; and wherein the second hydrocarbon solvent is for example toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, preferably toluene; (in an embodiment, the second ether solvent and the second hydrocarbon solvent are the same as the first ether solvent and the
20 first hydrocarbon solvent, respectively);

at a temperature in the range of from about room temperature to about reflux temperature, more preferably at a temperature in the range of from about 60°C to about 95°C; to yield the corresponding compound of formula (IX).

Preferably, the compound of formula (VIII), as a solution in a suitably
25 selected second hydrocarbon solvent, more preferably a suitably selected second aromatic hydrocarbon, such as toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, more preferably, toluene; is added to a solution of the compound of formula (VII) in a suitably selected second ether solvent other than THF, such as diisopropyl ether, 1,4-dioxane, 2-
30 methyl-THF, MTBE, cyclopentyl methyl ether (CPME), di-n-butyl ether, and the like, more preferably CPME or di-(n-butyl) ether, more preferably, di-(n-butyl) ether. Preferably, the final solvent mixture is present in a volume ratio of


second ether solvent : second hydrocarbon solvent of from about 1: 1 to about 1:3.


The compound of formula (IX) is de-protected according to known
 5 methods, to yield the corresponding compound of formula (I). For example,
 wherein each Z is pivaloyl, the compound of formula (IX) may be de-protected
 by reacting with a suitably selected alkoxide or hydroxide base such as sodium
 methoxide, sodium ethoxide, lithium hydroxide, and the like, in a suitably
 selected solvent such as methanol, ethanol, and the like, to yield the
 10 corresponding compound of formula (I).

One skilled in the art will recognize that, depending on the particular
 protecting group Z, other reagents may be used in the de-protection step
 including, but not limited to, Pd/C, Pd(OH)₂, PdCl₂, Pd(OAc)₂/Et₃SiH, RaNi, a
 suitably selected acid, a suitably selected base, fluoride, and the like.

15 The compound of formula (I) is preferably isolated according to known
 methods, for example by extraction, filtration or column chromatography. The
 compound of formula (I) is further, preferably purified according to known
 methods, for example by recrystallization.

20 In an embodiment, the present invention is directed to a process for the
 preparation of a compound of formula (I-S), as outlined in Scheme 2, below.





5 Accordingly, a suitably substituted compound of formula (V-S), wherein
LG¹ is a suitably selected leaving group such as bromo, iodo, and the like,
preferably LG¹ is bromo or iodo, a known compound or compound prepared by
known methods, is reacted with
a mixture of a suitably selected zinc salt such as zinc dibromide (ZnBr₂),
10 zinc diiodide (ZnI₂), zinc ditriflate, and the like, preferably ZnBr₂; or with an
amine complex of zinc halide such as pyridine zinc bromide complex, N-
methylmorpholine zinc bromide complex, and the like; wherein the zinc salt or
amine complex of zinc halide is preferably present in an amount in the range of
from about 0.33 to about 3.0 molar equivalents, more preferably in an amount
15 in the range of from about 0.33 to about 1.0 molar equivalents, more preferably
in an amount of about 0.5 molar equivalents; and a suitably selected organo-
lithium reagent such as trimethylsilylmethyl lithium, n-hexyl lithium, sec-butyl
lithium, n-butyllithium, t-butyllithium, methyl lithium, and the like, preferably n-

hexyl lithium or n-butyl lithium; wherein the organo-lithium reagent is preferably present in an amount in the range of from about 0.5 to about 2.0 molar equivalents, preferably in an amount in the range of from about 1.0 to about 1.2 molar equivalents;


- 5 in a suitably selected first hydrocarbon solvent, such as toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, preferably toluene; preferably at a temperature less than about room temperature, more preferably at a temperature in the range of from about -78°C to about room temperature; more preferably at about 0°C; to yield a mixture of the
- 10 corresponding compound of formula (VI-S), wherein M¹ is lithium, and the zinc salt. Preferably, the compound of formula (VI-S) is not isolated.

- In an embodiment of the present invention, the zinc salt is ZnBr₂ and the organo-lithium reagent is n-butyl-lithium. In another embodiment of the present invention, the molar ratio of the zinc salt to the organo-lithium reagent is about
- 15 1:2. In another embodiment of the present invention, the zinc salt and the organo-lithium reagent are pre-mixed; preferably for a period of time in the range of from about 1 to about 2 hours.

- It has been discovered that (a) pre-mixing of the zinc salt and organo-
- 20 lithium reagent and (b) adding the compound of formula (V-S) to the pre-mixed zinc salt and organo-lithium reagent, improves the reaction conditions for making the compound of formula (VII-S). More particularly, the use of the pre-mixed zinc salt and organo-lithium reagent solution permits for the preparation of the compound of formula (VI-S) at a temperature of about 0°C. The higher
- 25 temperature is preferred, particularly for large scale manufacture. Additionally, the use of the pre-mixed zinc salt and organo-lithium reagent mixture results in improved yield of the compound of formula (IX-S).

- To the mixture of the compound of formula (VI-S) and the zinc salt is
- 30 admixed a suitably selected first ether solvent such as diethyl ether, diisopropyl ether, di-n-butyl ether, MTBE, cyclopentylmethyl ether, and the like, preferably di-n-butyl ether or cyclopentyl methyl ether; wherein the first ether solvent is

preferably present in an amount in the range of from about 5% to about 15% by volume (relative to the total volume of the first hydrocarbon solvent and the first ether solvent), or any amount or range therein; preferably in an amount in the range of from about 7% to about 10%, or any amount or range therein; more preferably, at about amount in the range of from about 8% to about 9%, or any amount or range therein; (to minimize the formation of undesired by-product, more particularly to minimize formation of the compound of formula (U-S))



also known as 2-(4-fluorophenyl)-5-(2-methylbenzyl)thiophene);
 10 to yield the corresponding compound of formula (VII-S), wherein M^2 is a corresponding reactive zinc species, which compound is not isolated.

It is theorized that, when the zinc salt used in the previous reaction step is $ZnBr_2$, then in the compound of formula (VII-S), M^2 may include $ZnBr$; when the zinc salt used in the previous reaction step is ZnI_2 , then in the compound of
 15 formula (VII-S), M^2 may include ZnI ; when the zinc salt used in the previous reaction step is zinc ditriflate, then in the compound of formula (VII-S), M^2 may include zinc triflate. At this time, the intermediate of formula (VII) has not been isolated or fully characterized (particularly, as to the identity of the M^2 group).

It is further theorized that admixing the suitably selected first ether
 20 solvent with the mixture of the compound of formula (VI-S) and the zinc salt, preferably zinc bromide, results in chelation of the zinc salt. The chelated zinc bromide then reacts with the compound of formula (VI-S) to yield the corresponding compound of formula (VII-S).

25 The compound of formula (VII-S) is reacted with a suitably substituted compound of formula (VIII-S), wherein LG^2 is a suitably selected leaving group such as bromo, chloro, iodo, and the like, preferably bromo; and wherein each Z is independently a suitably selected oxygen protecting group, for example Z may selected from the group consisting of benzyl, benzoyl, pivaloyl, isobutyryl,

p-methoxy-benzyl, acetyl, propionyl, and the like; preferably, each Z protecting group is the same, more preferably each Z is pivaloyl, a known compound or compound prepared by known methods; wherein the compound of formula (VIII-S) is preferably present in an amount in the range of from about 0.5 to
5 about 3.0 molar equivalents, or any amount or range therein, more preferably in an amount in the range of from about 0.8 to about 1.25 molar equivalents, or any amount or range therein, more preferably in an amount of about 1.0 to about 1.1 molar equivalents;

optionally in a mixture of a suitably selected second ether solvent and a
10 suitably selected second hydrocarbon solvent, wherein the second ether solvent is for example, diethyl ether, di-n-butyl ether, MTBE, 2-Me-THF, cyclopentylmethyl ether, di-isopropyl ether, and the like, preferably di-n-butyl ether or cyclopentyl methyl ether; and wherein the second hydrocarbon solvent is for example toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride,
15 and the like, preferably toluene; (in an embodiment, the second ether solvent and the second hydrocarbon solvent are the same as the first ether solvent and the first hydrocarbon solvent, respectively);

at a temperature in the range of from about room temperature to about reflux temperature, more preferably at a temperature in the range of from about
20 60°C to about 95°C; to yield the corresponding compound of formula (IX-S).


Preferably, the compound of formula (VIII-S), as a solution in a suitably selected second hydrocarbon solvent, more preferably a suitably selected second aromatic hydrocarbon, such as toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, more preferably, toluene; is
25 added to a solution of the compound of formula (VII) in a suitably selected second ether solvent other than THF, such as diisopropyl ether, 1,4-dioxane, 2-methyl-THF, MTBE, cyclopentyl methyl ether (CPME), di-n-butyl ether, and the like, more preferably CPME or di-(n-butyl) ether, more preferably di-(n-butyl) ether. Preferably, the final solvent mixture is present in a volume ratio of
30 second ether solvent : second hydrocarbon solvent of from about 1:1 to about 1:3.


The compound of formula (IX-S) is de-protected according to known methods, to yield the corresponding compound of formula (I-S). For example, wherein each Z is pivaloyl, the compound of formula (IX-S) may be de-protected by reacting with a suitably selected alkoxide or hydroxide base such as sodium methoxide, sodium ethoxide, lithium hydroxide, and the like, in a suitably selected solvent such as methanol, ethanol, and the like, to yield the corresponding compound of formula (I-S).

One skilled in the art will recognize that, depending on the particular protecting group Z, other reagents may be used in the de-protection step including, but not limited to, Pd/C, Pd(OH)₂, PdCl₂, Pd(OAc)₂/Et₃SiH, RaNi, a suitably selected acid, a suitably selected base, fluoride, and the like.

The compound of formula (I-S) is preferably isolated according to known methods, for example by extraction, filtration or column chromatography. The compound of formula (I-S) is further, preferably purified according to known methods, for example by recrystallization.

In another embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-K), as outlined in Scheme 3, below.





5

Scheme 3

Accordingly, a suitably substituted compound of formula (V-K), wherein LG^1 is a suitably selected leaving group such as bromo, iodo, and the like, preferably LG^1 is bromo or iodo, a known compound or compound prepared by known methods, is reacted with

10

a mixture of a suitably selected zinc salt such as zinc dibromide ($ZnBr_2$), zinc diiodide (ZnI_2), zinc ditriflate, and the like, preferably $ZnBr_2$; or with an amine complex of zinc halide such as pyridine zinc bromide complex, N-methylmorpholine zinc bromide complex, and the like; wherein the zinc salt or amine complex of zinc halide is preferably present in an amount in the range of from about 0.33 to about 3.0 molar equivalents, more preferably in an amount in the range of from about 0.33 to about 1.0 molar equivalents, more preferably in an amount of about 0.5 molar equivalents; and a suitably selected organo-lithium reagent such as trimethylsilylmethyl lithium, n-hexyl lithium, sec-butyl

15

lithium, n-butyllithium, t-butyllithium, methyl lithium, and the like, preferably n-hexyl lithium or n-butyl lithium; wherein the organo-lithium reagent is preferably present in an amount in the range of from about 0.5 to about 2.0 molar equivalents, preferably in an amount in the range of from about 1.0 to about 1.2
5 molar equivalents;


in a suitably selected first hydrocarbon solvent, such as toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, preferably toluene; preferably at a temperature less than about room temperature, more preferably at a temperature in the range of from about -78°C to about room
10 temperature; more preferably at about 0°C; to yield a mixture of the corresponding compound of formula (VI-K), wherein M¹ is lithium, and the zinc salt. Preferably, the compound of formula (VI-K) is not isolated.

In an embodiment of the present invention, the zinc salt is ZnBr₂ and the organo-lithium reagent is n-butyl-lithium. In another embodiment of the present
15 invention, the molar ratio of the zinc salt to the organo-lithium reagent is about 1:2. In another embodiment of the present invention, the zinc salt and the organo-lithium reagent are pre-mixed; preferably for a period of time in the range of from about 1 to about 2 hours.

20 It has been discovered that (a) pre-mixing of the zinc salt and organo-lithium reagent and (b) adding the compound of formula (V-K) to the pre-mixed zinc salt and organo-lithium reagent, improves the reaction conditions for making the compound of formula (VII-K). More particularly, the use of the pre-mixed zinc salt and organo-lithium reagent solution permits allows for the
25 preparation of the compound of formula (VI-K) at a temperature of about 0°C. The higher temperature is preferred, particularly for large scale manufacture. Additionally, the use of the pre-mixed zinc salt and organo-lithium reagent mixture results in improved yield of the compound of formula (IX-K).

30 To the mixture of the compound of formula (VI-K) and the zinc salt is admixed a suitably selected first ether solvent such as diethyl ether, diisopropyl ether, di-n-butyl ether, MTBE, cyclopentylmethyl ether, and the like, preferably

di-n-butyl ether or cyclopentyl methyl ether; wherein the first ether solvent is preferably present in an amount in the range of from about 5% to about 15% by volume (relative to the total volume of the first hydrocarbon solvent and the first ether solvent), or any amount or range therein; preferably in an amount in the range of from about 7% to about 10%, or any amount or range therein; more preferably, at about amount in the range of from about 8% to about 9%, or any amount or range therein; (to minimize the formation of undesired by-product, more particularly to minimize formation of the compound of formula (U-K)



also known as 5-(5-(2-chlorobenzyl)thiophen-2-yl)-2-fluoropyridine); to yield the corresponding compound of formula (VII-K), wherein M^2 is a corresponding reactive zinc species, which compound is not isolated.

It is theorized that, when the zinc salt used in the previous reaction step is $ZnBr_2$, then in the compound of formula (VII-K), M^2 may include $ZnBr$; when the zinc salt used in the previous reaction step is ZnI_2 , then in the compound of formula (VII-K), M^2 may include ZnI ; when the zinc salt used in the previous reaction step is zinc ditriflate, then in the compound of formula (VII-K), M^2 may include zinc triflate. At this time, the intermediate of formula (VII-K) has not been isolated or fully characterized (particularly, as to the identity of the M^2 group).

It is further theorized that admixing the suitably selected first ether solvent with the mixture of the compound of formula (VI-K) and the zinc salt, preferably zinc bromide, results in chelation of the zinc salt. The chelated zinc bromide then reacts with the compound of formula (VI-K) to yield the corresponding compound of formula (VII-K).

The compound of formula (VII-K) is reacted with a suitably substituted compound of formula (VIII-S), wherein LG^2 is a suitably selected leaving group such as bromo, chloro, iodo, and the like, preferably bromo; and wherein each

Z is independently a suitably selected oxygen protecting group, for example Z may selected from the group consisting of benzyl, benzoyl, pivaloyl, isobutyryl, p-methoxy-benzyl, acetyl, propionyl, and the like; preferably, each Z protecting group is the same, more preferably each Z is pivaloyl, a known compound or
5 compound prepared by known methods; wherein the compound of formula (VIII) is preferably present in an amount in the range of from about 0.5 to about 3.0 molar equivalents, or any amount or range therein, more preferably in an amount in the range of from about 0.8 to about 1.25 molar equivalents, or any amount or range therein, more preferably in an amount of about 1.0 to about
10 1.1 molar equivalents;

optionally in a mixture of a suitably selected second ether solvent and a suitably selected second hydrocarbon solvent, wherein the second ether solvent is for example, diethyl ether, di-n-butyl ether, MTBE, 2-Me-THF, cyclopentylmethyl ether, and the like, preferably di-n-butyl ether or cyclopentyl
15 methyl ether; and wherein the second hydrocarbon solvent is for example toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, preferably toluene; (in an embodiment, the second ether solvent and the second hydrocarbon solvent are the same as the first ether solvent and the first hydrocarbon solvent, respectively);

20 at a temperature in the range of from about room temperature to about reflux temperature, more preferably at a temperature in the range of from about 60°C to about 95°C; to yield the corresponding compound of formula (IX-K).

Preferably, the compound of formula (VIII-S), as a solution in a suitably selected second hydrocarbon solvent, more preferably a suitably selected
25 second aromatic hydrocarbon, such as toluene, xylene, fluorobenzene, chlorobenzene, benzotrifluoride, and the like, more preferably, toluene; is added to a solution of the compound of formula (VII-K) in a suitably selected second ether solvent other than THF, such as diisopropyl ether, 1,4-dioxane, 2-methyl-THF, MTBE, cyclopentyl methyl ether (CPME), di-n-butyl ether, and the
30 like, more preferably CPME or di-(n-butyl) ether, more preferably di-(n-butyl) ether. Preferably, the final solvent mixture is present in a volume ratio of

second ether solvent : second hydrocarbon solvent of from about 1:1 to about 1:3.

The compound of formula (IX-K) is de-protected according to known
5 methods, to yield the corresponding compound of formula (I-K). For example, wherein each Z is pivaloyl, the compound of formula (IX-K) may be de-protected by reacting with a suitably selected alkoxide or hydroxide base such as sodium methoxide, sodium ethoxide, lithium hydroxide, and the like, in a suitably selected solvent such as methanol, ethanol, and the like, to yield the
10 corresponding compound of formula (I-K).

One skilled in the art will recognize that, depending on the particular protecting group Z, other reagents may be used in the de-protection step including, but not limited to, Pd/C, Pd(OH)₂, PdCl₂, Pd(OAc)₂/Et₃SiH, RaNi, a suitably selected acid, a suitably selected base, fluoride, and the like.

15 The compound of formula (I-K) is preferably isolated according to known methods, for example by extraction, filtration or column chromatography. The compound of formula (I-K) is further, preferably purified according to known methods, for example by recrystallization.

20 The present invention further comprises pharmaceutical compositions containing a compound prepared according to any of the processes described herein with a pharmaceutically acceptable carrier. Pharmaceutical compositions containing one or more of the compounds of the invention described herein as the active ingredient can be prepared by intimately mixing
25 the compound or compounds with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral). Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils,
30 alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents,

lubricants, binders, disintegrating agents and the like. Solid oral preparations may also be coated with substances such as sugars or be enteric-coated so as to modulate major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other ingredients may be added to
5 increase solubility or preservation. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives.

To prepare the pharmaceutical compositions of this invention, one or more compounds of the present invention as the active ingredient is intimately admixed with a pharmaceutical carrier according to conventional
10 pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and
15 solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like.
20 Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes
25 such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient
30 necessary to deliver an effective dose as described above. The pharmaceutical compositions herein may contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from

about 0.01 to about 1,000 mg or any amount or range therein, and may be given at a dosage of from about 0.01 to about 300 mg/kg/day, or any amount or range therein, preferably from about 0.1 to about 50 mg/kg/day, or any amount or range therein. The dosages, however, may be varied depending
5 upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or
10 suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound,
15 such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums,
20 and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the
25 composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 1,000 mg, or any amount or range therein, of the active ingredient of the present invention. The tablets or pills of the novel composition
30 can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of

an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

The methods of treating described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.01 mg and about 1000 mg of the compound, or any amount or range therein; preferably about 10 to about 500 mg of the compound, or any amount or range therein, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixirs, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present

invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than
5 intermittent throughout the dosage regimen.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and
10 coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without
15 limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable
20 preservatives are employed when intravenous administration is desired.

To prepare a pharmaceutical composition of the present invention, a compound prepared according to any of the processes described herein as the active ingredient is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may
25 take a wide variety of forms depending of the form of preparation desired for administration (e.g. oral or parenteral). Suitable pharmaceutically acceptable carriers are well known in the art. Descriptions of some of these pharmaceutically acceptable carriers may be found in The Handbook of Pharmaceutical Excipients, published by the American Pharmaceutical
30 Association and the Pharmaceutical Society of Great Britain.

Methods of formulating pharmaceutical compositions have been described in numerous publications such as Pharmaceutical Dosage Forms:

Tablets, Second Edition, Revised and Expanded, Volumes 1-3, edited by Lieberman et al; Pharmaceutical Dosage Forms: Parenteral Medications, Volumes 1-2, edited by Avis et al; and Pharmaceutical Dosage Forms: Disperse Systems, Volumes 1-2, edited by Lieberman et al; published by
5 Marcel Dekker, Inc.

Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of disorders as described herein is required.

Optimal dosages to be administered may be readily determined by those
10 skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.


15 One skilled in the art will recognize that, both *in vivo* and *in vitro* trials using suitable, known and generally accepted cell and / or animal models are predictive of the ability of a test compound to treat or prevent a given disorder.

One skilled in the art will further recognize that human clinical trials including first-in-human, dose ranging and efficacy trials, in healthy patients
20 and / or those suffering from a given disorder, may be completed according to methods well known in the clinical and medical arts.

The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way
25 the invention set forth in the claims which follow thereafter. In the Examples which follow, some synthesis products are listed as having been isolated as a residue. It will be understood by one of ordinary skill in the art that the term "**residue**" does not limit the physical state in which the product was isolated and may include, for example, a solid, an oil, a foam, a gum, a syrup, and the
30 like.

Example 1

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)




- 5 In a 250mL RBF with mechanical stirrer, dried and under argon atmosphere, 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (22.20 mmoles; 9.06 g) was dissolved in a mixture of dried and degassed toluene (37.00 mL; 32.23 g) / diethyl ether (37.00 mL; 26.24 g) at room temperature. After cooling to -50°C (isopropanol + dry ice bath) under vigorous stirring,
- 10 (trimethylsilyl)methyl lithium (1M in pentane, 37.00 mL) was added dropwise to the heterogeneous mixture. 30 min after the end of the addition, the conversion was checked by sampling and extra (trimethylsilyl)methyl lithium was added if needed. After 15min., zinc dibromide (22.20 mmoles; 5.00 g) (solid extra dry from Aldrich) was added in one portion and the resulting mixture
- 15 was allowed to warm up to 25°C over 1 hour. After 1 hour stirring at room temperature, diethyl ether and pentane were evaporated under reduced pressure (400mmHg) at 15°C. Finally α -D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (10.72 g, 18.50 mmoles) dissolved in degassed toluene (18.50 mL) was added dropwise over 10 min and the
- 20 resulting mixture was heated at 75°C for 21 hours. After cooling to room temperature, aqueous solution of ammonium chloride (1M, 100 mL) and ethyl acetate (150 mL) were added. After 10 min. stirring, the 2 phases were

separated and the organic layer was washed twice with water (100 mL) and once with brine (100mL). The organic layer was thereafter dried over sodium sulfate and the solvent was evaporated under reduced pressure to yield a clear brown oil. The oil was purified by MPLC (cartridge: 330g SiO₂, solvent system :
 5 95/5 to 85/15 heptane / AcOEt) to yield the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a single isomer. The ¹H NMR spectrum was consistent with the previously measured ¹H NMR spectra for the title compound.

10

Example 2

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)



15


In a 25mL Schlenk reactor, dried and under argon atmosphere, 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (1.99 mmoles; 813.71 mg) was dissolved in dry cyclopentylmethyl ether (CPME) (7.2 mL) at room temperature. After cooling to -50°C (acetonitrile + dry ice) under vigorous
 20 stirring, n-hexyllithium (2.3M in hexane, 966.31 µL) was added dropwise to the mixture. After 15min, zinc dibromide (996.50 µL; 2M solution in CPME) was added and the resulting mixture was allowed to warm up to 15°C over 1.5 hour.

Then α -D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (1.05 g, 1.81 mmoles) dissolved in degassed CPME (1.81 mL) was added dropwise over 10 min and the resulting mixture was heated at 85°C overnight. After cooling to room temperature, an aqueous solution of ammonium chloride (1M, 10mL) and ethyl acetate (15mL) were added. After 10 min. stirring, the 2 phases were separated, and the organic layer was washed twice with water (10mL) and once with brine (10mL). The organic layer was dried over sodium sulfate and the solvent was evaporated under reduced pressure to yield a clear brown oil, which was determined by quantitative HPLC to contain the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a single isomer. The ^1H NMR spectrum was consistent with the previously measured ^1H NMR spectra for the title compound.

15

Example 3

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)




20

In a 25 mL Schlenk reactor, dried and under argon atmosphere, 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (1.90 mmoles; 775 mg) was dissolved in toluene (3.45 mL) / diethyl ether (3.45 mL) at room temperature.

After cooling to -50°C (acetonitrile + dry ice) under vigorous stirring, n-hexyllithium (2.3M in hexane, 920.29 µL) was added dropwise to the mixture. After 15min., zinc dibromide (2.07 mmoles; 466 mg) was added in one portion and the resulting mixture was allowed to warm up to 15°C over 1.5 hours. The
5 resulting mixture was then cooled to 0°C and (trimethylsilyl)methylolithium (1M in pentane, 1.9mL) was added dropwise. After 1 hour, diethyl ether and hexane were evaporated under reduced pressure (400 mmHg) at 15°C. Then α-D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (1.73 mmoles; 1.00 g) dissolved in degassed toluene (1.73 mL) was added dropwise
10 over 10 min and the resulting mixture was heated at 85°C overnight. After cooling to room temperature, aqueous solution of ammonium chloride (1M, 10mL) and ethyl acetate (15mL) were added. After 10 min. stirring, the 2 phases were separated and the organic layer was washed twice with water (10mL) and once with brine (10mL). The organic layer was dried over sodium
15 sulfate and the solvent was evaporated under reduced pressure to yield a clear brown oil, which was determined by quantitative HPLC to contain the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl)tetrahydro-2H-pyran 3,4,5 -triyl tris(2,2-dimethylpropanoate) as a single isomer. The ¹H NMR spectrum was consistent
20 with the previously measured ¹H NMR spectra for the title compound.

Example 4

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-
25 dimethylpropanoate)




In a 25 mL Schlenk reactor, dried and under argon atmosphere, 2-(4-fluorophenyl)-5-(5-(5-iodo-2-methylbenzyl)thiophene (1.58 mmol; 643 mg) was dissolved in toluene (2.86 mL) / 2-methyltetrahydrofuran (2.86 mL) at room temperature. After cooling to -50°C (acetonitrile+ dry ice) under vigorous stirring, n-hexyllithium (2.3M in hexane; 764 µL) was added dropwise to the mixture. After 15min., zinc dibromide (1.72 mmol; 387 mg) dissolved in 2-methyltetrahydrofuran (859 µL) was added in one portion and the resulting mixture was allowed to warm up to 15°C over 1.5 hours. Then α-D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (1.43 mmol; 830 mg) dissolved in degassed toluene (1.43 mL) was added dropwise over 10 min and the resulting mixture was heated at 85°C overnight. After cooling to room temperature, an aqueous solution of ammonium chloride (1M, 10mL) and ethyl acetate (15mL) were added. After 10 min stirring, the phases were separated and the organic layer was washed twice with water (10mL) and once with brine (10mL). The organic layer was dried over sodium sulfate and the solvent was evaporated under reduced pressure to yield a clear brown oil, which was determined by quantitative HPLC to contain the title compound (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a single isomer. The ¹H NMR spectrum was consistent with the previously measured ¹H NMR spectra for the title compound.

Example 5

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-
dimethylpropanoate)

5




- In a 25 mL Schlenk reactor, dried and under argon atmosphere, 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (1.90 mmoles; 775 mg) was dissolved in toluene (3.45 mL) / diethyl ether (3.45 mL) at room temperature.
- 10 After cooling to -50°C (acetonitrile+ dry ice) under vigorous stirring, n-hexyllithium (2.3M in hexane, 920 µL) was added dropwise to the mixture. After 15 min, zinc dibromide (2.07 mmoles; 466 mg) was added in one portion and the resulting mixture was allowed to warm up to 15°C over 1.5 hours. After 1 hour, diethyl ether and hexane were evaporated under reduced pressure
- 15 (400 mmHg) at 15°C. Then α-D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (1.73 mmoles; 1.00 g) dissolved in degassed toluene (1.73 mL) was added dropwise over 10 min and the resulting mixture was heated at 50°C for 2 days. After cooling to room temperature, aqueous solution of ammonium chloride (1M, 10 mL) and ethyl acetate (15 mL) were added.
- 20 After 10 min stirring, the phases were separated and the organic layer was washed twice with water (10mL) and once with brine (10 mL). The organic layer was dried over sodium sulfate and the solvent was evaporated under

reduced pressure to yield a clear brown oil, which was determined by quantitative HPLC to contain the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a single isomer.

- 5 The ^1H NMR spectrum was consistent with the previously measured ^1H NMR spectra for the title compound.

Example 6

10 (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)




- 15 In a 25 mL Schlenk reactor, dried and under argon atmosphere, 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (2.60 mmoles; 1.06 g) was dissolved in toluene (4.73 mL) / Methoxy-cyclopentane (4.73 mL) at room temperature. After cooling to -50°C (acetonitrile + dry ice) under vigorous stirring, n-hexyllithium (2.3M in hexane, 1.26 mL) was added dropwise to the mixture. After 15min., zinc dibromide (2.84 mmoles; 639 mg) dissolved in dry methoxy-cyclopentane (1.40 mL) was added dropwise and the resulting mixture
- 20 was allowed to warm up to 15°C over 1 hour. Then α -D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (2.36 mmoles; 1.37 g) dissolved in degassed toluene (2.36 mL) was added dropwise over 10 min and

the resulting mixture was heated at 75°C for 2 days. After cooling to room temperature, aqueous solution of ammonium chloride (1M, 10mL) and ethyl acetate (15 mL) were added. After 10 mi. stirring, the phases were separated and the organic layer was washed twice with water (10 mL) and once with brine (10 mL). The organic layer was dried over sodium sulfate and the solvent was evaporated under reduced pressure to yield a clear brown oil, which was determined by quantitative HPLC to contain the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a single isomer. The ¹H NMR spectrum was consistent with the previously measured ¹H NMR spectra for the title compound.

Example 7

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)




In a 50mL Schlenk reactor under argon atmosphere at room temperature, 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (2.45 mmols; 1.00 g) was dissolved in n-butyl ether (980 µL) / toluene (8.8 mL). The temperature was then decreased to -60°C. N-hexyllithium (2.3M in hexane, 1.20 mL) was added dropwise. After 2 hours, zinc dibromide (607 mg)

was added in one portion at -60°C. The resulting mixture was allowed to warm up slowly to 10°C over 2 hours. At 10°C, α -D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) (2.69 mmol; 1.56 g) dissolved in toluene (2.69 mL) was added over 1 min. and the temperature was increased to 50°C overnight. The temperature of the mixture was increased to 60°C for 1 hour and finally for 2 days at 70°C. After cooling to room temperature, aqueous solution of ammonium chloride (1M, 10 mL) and ethyl acetate (15 mL) were added. After 10 min stirring, the phases were separated and the organic layer was washed twice with water (10 mL) and once with brine (10 mL). The organic layer was dried over sodium sulfate and the solvent was evaporated under reduced pressure to yield a clear brown oil, which was determined by quantitative HPLC to contain the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a single isomer. The ^1H NMR spectrum was consistent with the previously measured ^1H NMR spectra for the title compound.

Example 8

(2R,3R,4S,5R,6R)-6-(pivaloyloxymethyl)tetrahydro-2H-pyran-2,3,4,5-tetrayl
tetrakis(2,2-dimethylpropanoate)




D-glucose (25.0g, 0.139 mol) was suspended in anhydrous dichloromethane (416 mL) under nitrogen and the resulting mixture was stirred

for 5 minutes at room temperature, then cooled to 0°C and stirred for 10 minutes. To the resulting mixture was then added TEA (154.7 mL), dropwise over about 10-15 min, with stirring; then DMAP (1.25 g, 0.0102 mol) in one portion. To the resulting mixture was added pivaloyl chloride (136 mL) diluted
 5 with dichloromethane (83 mL) at 0°C, over 30 min. The ice bath was removed and the resulting mixture stirred at room temperature for 20 hours. The resulting mixture was then poured into dichloromethane (500 mL) and hydrochloric acid (1.5M, 375 mL) and the resulting phases separated. The organic layer was washed with sodium bicarbonate solution (550 g in 500mL DI
 10 water, 1N) and then evaporated to a small volume. To the resulting residue was added ethanol (95%, 240 mL) and the mixture heated to reflux temperature to yield a homogeneous mixture. The resulting mixture was cooled to 0°C, resulting in the formation of white crystals, which were filtered and dried in vacuo at room temperature, overnight, to yield the title compound.

15

Example 9

(2R,3R,4S,5R,6R)-2-bromo-6-(pivaloyloxymethyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)




20 (2R,3R,4S,5R,6R)-6-(pivaloyloxymethyl)tetrahydro-2H-pyran-2,3,4,5-tetrayl tetrakis(2,2-dimethylpropanoate) (10.0 g, 16.65 mmol) was dissolved in anhydrous dichloromethane (100 mL) under nitrogen and stirred for 5 min at room temperature. To the mixture was then added zinc bromide (0.76 g, 3.33 mmol) and the resulting yellow solution stirred for 5 min at room temperature.
 25 To the mixture was then added TMS bromide (10.2 g, 66.58 mmol) diluted with

dichloromethane (10 mL) over about 15-20 min and the resulting mixture stirred at room temperature for 24 hours. The resulting mixture was filtered to remove the solids and the filtrate cooled to 0°C. To the cooled filtrate was then added sodium bicarbonate solution (132 g in 120 mL water) to a final pH in the range of 7-8. The resulting phases were separated, the organic layer washed with water (120 mL) and the combined aqueous layers evaporated to a small volume. To the resulting residue was added IPA (39.3 g) and the mixture heated to dissolve. The resulting mixture was cooled to 0°C, resulting in the formation of white crystals, which were filtered and dried in vacuo at room temperature, overnight, to yield the title compound.

Example 10

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)



STEP A: Preparation of Aryllithium Mixture

2-(4-Fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene (12.81 g, 31.37 mmol) was placed in a dry Schlenk tube under an argon atmosphere. Anhydrous toluene (15.7 mL) and anhydrous CPME (9.4 mL) were added by syringe, without stirring and the resulting mixture cooled to -45°C and then stirred. To the resulting cooled mixture was then added n-hexyllithium (14.3 g,

32.94 mmol), as a 2.5M solution in hexane (14.3 mL) over about 5-10 min; and the mixture warmed to -25°C over 1 hour.


STEP B: Preparation of Title Compound

Zinc bromide (3.88 g, 17.25 mmol) and lithium bromide (2.72 g, 34.50
5 mmol) were dried at 200°C in vacuo, in anhydrous CPME (18.6 mL) in a Schlenk tube. The mixture was then added by cannula, at -25°C to the aryllithium mixture (prepared as described in STEP A above) and the resulting mixture was warmed to 0°C over 1 hour. To the resulting mixture was then added (2R,3R,4S,5R,6R)-2-bromo-6-(pivaloyloxymethyl)tetrahydro-2H-pyran-
10 3,4,5-triyl tris(2,2-dimethylpropanoate) (20.0g, 34.50 mmol) in anhydrous toluene (31.4 mL). The ice bath was removed and the resulting mixture stirred at room temperature for 30 min; then heated to 65°C for 48 hours. The resulting suspension was filtered through a glass frit, rinsed with toluene (20 mL) and the filtrate washed with 1N ammonium chloride solution (100 mL) and
15 water (100mL). The toluene was distilled off to a small volume. Methanol (157 mL) was added to the resulting residue and the mixture cooled to 0°C, resulting in the formation of crystals, which were filtered and dried in vacuo at 40°C, overnight, to yield the title compound. Yield: 18.30 g, 75%.

20

Example 11

(2S,3R,4R,5S,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol




25 (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-

dimethylpropanoate) (39.0 g, 50.0 mmol) was suspended in methanol (150 mL) at room temperature. Sodium methoxide solution (9.3 mL) was added and the resulting suspension was stirred at room temperature, heated to 60°C for 16 hours and then cooled. To the resulting yellow solution was then added water (50 mL) and seeds to the title compound. An additional portion of water (50 mL) was added, and the mixture stirred at 0°C for 1 hour, resulting in the formation of a precipitate, which was collected by filtration to yield the title compound. Yield: 20.00 g, 90%.

10

Example 12

(2S,3R,4R,5S,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol



15 STEP A: Coupling to (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-((pivaloyloxy)methyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)

A jacketed reactor with overhead stirrer, under dry nitrogen flow was heated at 100°C for 16h, then cooled to ~20°C under nitrogen flow. To the reactor was then charged ZnBr₂ (10g, 44.45mmol, 0.55eq) and toluene (264mL, 8V), under nitrogen; and the resulting mixture stirred for 5 minutes. *n*-BuLi 2.3M in heptane (84.87mmol, 1.05eq.) was then added at ~20°C, under nitrogen in one portion. The resulting mixture was stirred for 2h, then cooled to 0°C within 30min. 2-(4-Fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene, as a solid (1eq., 33g, 80.83mmol), was added to the reactor, under nitrogen within 10min, and the resulting mixture was stirred for 1h at 0°C. *n*-Bu₂O (26.4mL,

0.8V, 10 v%) was added to the reactor, under nitrogen. The resulting mixture was warmed to ~25°C and then stirred for 3 h.

To the resulting mixture was then added (2R,3R,4S,5R,6R)-2-bromo-6-((pivaloyloxy)methyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as a solid (1eq., 47.75g, 80.83mmol)), at ~25°C, in one portion. The resulting mixture was heated to ~95°C within ~15min, then stirred at ~95°C for 1h.

The resulting mixture was cooled to room temperature, quenched with aqueous ammonia 50 w/w% (132mL, 4V) and stirred for 1h at 30°C. Water (4V, 132mL) was added and the resulting mixture was stirred for 1 extra hour at 30°C. The resulting phases were separated at 30°C, and the organic layer collected. The organic layer was then placed under reduced pressure (90mbar), and 300 mL of solvent was distilled, at a maximum temperature of 50°C. To the resulting residue was added 1-butanol (150mL, 4.5V) and 150 mL was distilled under reduced pressure (90mbar). To the resulting residue was again added 1-butanol (150mL, 4.5V) and 150mL distilled under reduced pressure (90mbar) at a maximum temperature of 80°C. To the resulting residue was added 1-butanol (150mL, 4.5V) a third time. The resulting mixture was cooled to room temperature, then to 0°C, and after 1hour at 0°C, the mixture was filtered. The filtercake was washed with 1-butanol (1V, 33mL), then dried under reduced pressure at 50°C to yield (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-((pivaloyloxy)methyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) as clear yellow solution, with 84% in situ yield, as determined against a qualified standard.

STEP B: (2S,3R,4R,5S,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol

A reactor was charged with methanol (13mL, .2.56V) and (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-((pivaloyloxy)methyl)tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate) (5.08g, 6.50mmoles, 1eq.). NaOCH₃ 30w/w% in


methanol (0.233g, 0.2eq.) was added and the resulting mixture was heated at reflux (65°C) and stirred for 5 hours.

The resulting mixture was heated to distill methanol (1.3V, 6.5mL) at 66°C under atmospheric pressure. To the resulting residue was added
 5 methanol (1.3V, 6.5ml), and the mixture heated to distill methanol (1.3V, 6.5mL) at 66°C under atmospheric pressure. Methanol (1.3V, 6.5ml) was added a second time, and the mixture heated to again distill methanol (1.3V, 6.5mL) at 66°C under atmospheric pressure. The resulting mixture was then cooled to 60°C. Acetic acid (78mg, 0.2eq.) and water (0.96V, 4.88mL) were added and
 10 the resulting mixture cooled to 26°C, seeded with the desired product (14.5mg, 0.005mol/mol) and stirred for 6h (min 4h). Water (0.77V, 3.9mL) was added over 2h and the resulting mixture stirred for at least 1h. The resulting suspension was then cooled to 20°C (15-25°C) and stirred for at least 5h, then filtered. The filtercake was washed with a mixture of water / methanol (1/1 v/v,
 15 0.63V, 3.24mL), then dried under reduced pressure at 50°C overnight to yield (2S,3R,4R,5S,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(hydroxymethyl)tetrahydro-2H-pyran-3,4,5-triol as a hemi-hydrate in 95% yield as off-white solid.

20

Example 13

(2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate)



In a jacket reactor with overhead stirrer (dried nitrogen flow at 100°C for 16h) at 20°C under nitrogen flow, were charged ZnBr₂ (15.17g, 0.55eq) and toluene (90mL). After 5 min. stirring at 20°C under nitrogen atmosphere, *n*-butyl lithium (46mL, 2.8M in heptane, 1.05eq.) was added in one portion and the resulting mixture was stirred for 2 hours. After cooling down to -10°C over 30min., a solution of 2-(4-fluorophenyl)-5-(5-iodo-2-methylbenzyl)thiophene in toluene (50g, 1eq. dissolved in 325mL of toluene) was added dropwise, and the resulting mixture maintained at -10°C for 1h. *n*-Butyl ether (45mL) was then added in one portion, under nitrogen and the resulting mixture was warmed to 25°C. After 1.5h at 25°C, a solution of α-D-glucopyranosyl bromide, 2,3,4,6-tetrakis(2,2-dimethyl propanoate) in toluene (71g, 1.0eq, 1M in toluene) was added, at 25°C, in one portion. The resulting mixture was heated to 95°C over 90 min and then stirred at 95°C for 5h. After cooling to 25°C, an aqueous solution of ammonium chloride (500mL) was added. The resulting mixture was then stirred for 30min., the two phases were separated and the organic layer collected and assayed at 86.6% of the title compound, (2S,3S,4R,5R,6R)-2-(3-((5-(4-fluorophenyl)thiophen-2-yl)methyl)-4-methylphenyl)-6-(pivaloyloxymethyl) tetrahydro-2H-pyran-3,4,5-triyl tris(2,2-dimethylpropanoate).

20

Formulation Example

Solid, Oral Formulation - Prophetic Example

As a specific embodiment of an oral composition, 100 mg of the compound prepared as in Example 11 above, is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.


5

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the

10 following claims and their equivalents.


We Claim

1. A process for the preparation of compounds of formula (I)




wherein Ring A and Ring B are one of the following:


- 5 (1) Ring A is an optionally substituted unsaturated monocyclic heterocyclic ring, and Ring B is an optionally substituted unsaturated monocyclic heterocyclic ring, an optionally substituted unsaturated fused heterobicyclic ring, or an optionally substituted benzene ring; or
- (2) Ring A is an optionally substituted benzene ring, and Ring B is an
- 10 optionally substituted unsaturated monocyclic heterocyclic ring, or an optionally substituted unsaturated fused heterobicyclic ring wherein Y is linked to the heterocyclic ring of the fused heterobicyclic ring; or
- (3) Ring A is an optionally substituted unsaturated fused heterobicyclic ring, wherein the sugar moiety X-(sugar) and the moiety -Y-(Ring B) are both
- 15 on the same heterocyclic ring of the fused heterobicyclic ring, and Ring B is an optionally substituted unsaturated monocyclic heterocyclic ring, an optionally substituted unsaturated fused heterobicyclic ring, or an optionally substituted benzene ring;
- X is a carbon atom;
- 20 Y is $-(CH_2)_n-$; wherein n is 1 or 2;
- provided that in Ring A, X is part of an unsaturated bond;
- or a pharmaceutically acceptable salt or solvate thereof; comprising




reacting a compound of formula (V) wherein LG^1 is a leaving group, with
a mixture of a zinc salt and an organo-lithium reagent; in a first hydrocarbon
solvent; at a temperature in the range of from about -78°C to about room
5 temperature; to yield a mixture of the corresponding compound of formula (VI),
wherein M^1 is lithium, and the zinc salt;



admixing to the mixture of the compound of formula (VI) and the zinc salt
a first ether solvent; to yield the corresponding compound of formula (VII),
10 wherein M^2 is a reactive zinc species;



reacting the compound of formula (VII), with a compound of formula
(VIII), wherein each Z is an independently selected oxygen protecting group
and wherein LG^2 is a leaving group; to yield the corresponding compound of
15 formula (IX);



de-protecting the compound of formula (IX); to yield the corresponding compound of formula (I).

- 5 2. A process as in Claim 1, where in the zinc salt is ZnBr_2 and wherein the organo-lithium reagent is n-butyl lithium.

3. A process as in Claim 1, wherein the zinc salt and the organo-lithium reagent are present in a molar ratio of about 1:2.

- 10 4. A process as in Claim 1, wherein the zinc salt and the organo-lithium reagent are pre-mixed for a period of about 1 to about 2 hours.


5. A process as in Claims 1, wherein the first hydrocarbon solvent is
- 15 toluene.

6. A process as in Claim 1, wherein LG^1 is iodo, the zinc salt is ZnBr_2 , the organolithium reagent is n-butyl lithium, the zinc salt and the organo-lithium reagent are pre-mixed, the first hydrocarbon solvent is toluene, and wherein the
- 20 compound of formula (V) is reacted with the mixture of zinc salt and organo-lithium reagent at a temperature of about 0°C .


7. A process as in Claim 1, wherein the first ether solvent is di-n-butyl ether.

25

8. A process as in Claim 1, wherein the first ether solvent is present in an amount in the range of about 7% to about 10% by volume.
9. A process as in Claim 1, wherein LG¹ is iodo, Z is pivaloyl and LG² is
5 bromo.
10. A process as in Claim 1, wherein
X is a carbon atom;
Ring A is selected from the group consisting of 4-methylphenyl and 4-
10 chlorophenyl;
Y is -CH₂- and is bound at the 3-position of Ring A; and
Ring B is selected from the group consisting of 2-(5-(4-fluorophenyl)-
thienyl) and 2-(5-(6-fluoro-pyrid-3-yl)-thienyl).
11. A process for the preparation of a compound of formula (I-S)




or solvate thereof; comprising




- reacting a compound of formula (V-S) wherein LG¹ is a leaving group,
20 with a mixture of a zinc salt and an organo-lithium reagent; in a first


hydrocarbon solvent; at a temperature in the range of from about -78°C to about room temperature; to yield a mixture of the corresponding compound of formula (VI-S), wherein M^1 is lithium, and the zinc salt;



- 5 admixing to the mixture of the compound of formula (VI-S) and the zinc salt a first ether solvent; to yield the corresponding compound of formula (VII-S), wherein M^2 is a reactive zinc species;



- 10 reacting a compound of formula (VII-S), wherein M^2 is a zinc species, with a compound of formula (VIII-S), wherein each Z is an independently selected oxygen protecting group and wherein LG^2 is a leaving group; to yield the corresponding compound of formula (IX-S);



de-protecting the compound of formula (IX-S); to yield the corresponding compound of formula (I-S).

5

12. A process as in Claim 11, where in the zinc salt is ZnBr_2 and wherein the organo-lithium reagent is n-butyl lithium.

13. A process as in Claim 11, wherein the zinc salt and the organo-lithium reagent are present in a molar ratio of about 1:2.

14. A process as in Claim 11, wherein the zinc salt and the organo-lithium reagent are pre-mixed for a period of about 1 to about 2 hours.

15. A process as in Claims 11, wherein the first hydrocarbon solvent is toluene.


16. A process as in Claim 1, wherein LG^1 is iodo, the zinc salt is ZnBr_2 , the organolithium reagent is n-butyl lithium, the zinc salt and the organo-lithium reagent are pre-mixed, the first hydrocarbon solvent is toluene, and wherein the compound of formula (V-S) is reacted with the mixture of zinc salt and organo-lithium reagent at a temperature of about 0°C .

17. A process as in Claim 11, wherein the first ether solvent is di-n-butyl ether.

18. A process as in Claim 11, wherein the first ether solvent is present in an amount in the range of about 7% to about 10% by volume.

19. A process as in Claim 11, wherein LG¹ is iodo, Z is pivaloyl and LG² is bromo.

5



(I)