



(12)

Oversættelse af europæisk patent

Patent- og
Varemærkestyrelsen

(51) Int.Cl.: **C 12 N 1/20 (2006.01)** **A 61 K 31/22 (2006.01)** **A 61 K 35/74 (2015.01)**
C 12 R 1/225 (2006.01)

(45) Oversættelsen bekendtgjort den: **2016-02-01**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2015-11-11**

(86) Europæisk ansøgning nr.: **07748532.4**

(86) Europæisk indleveringsdag: **2007-05-30**

(87) Den europæiske ansøgnings publiceringsdag: **2009-04-01**

(86) International ansøgning nr.: **SE2007050371**

(87) Internationalt publikationsnr.: **WO2007142596**

(30) Prioritet: **2006-06-05 US 446628**

(84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**

(73) Patenthaver: **BIOGAIA AB, Box 3242, 103 64 Stockholm, Sverige**

(72) Opfinder: **CONNOLLY, Eamonn, Motionsvägen 3, S-181 30 LIDINGÖ, Sverige**
MÖLLSTAM, Bo, Cederflychtsvagen 12, S-443 31 LERUM, Sverige

(74) Fuldmægtig i Danmark: **Synergon AB, Box 32 S-443 21 Lerum, Sverige**

(54) Benævnelse: **ANVENDELSE AF UDVALGTE MÆLKESYREBAKTERIER TIL REDUKTION AF SPÆDBARNSKOLIK**

(56) Fremdragne publikationer:
EP-A1- 1 384 483
WO-A1-2007/142597
WO-A2-2005/117532
US-A1- 2004 185 032
US-A1- 2005 271 641
US-A1- 2006 251 634
SAVINO F ET AL: "Lactobacillus reuteri ATCC 55730 versus simethicone in the treatment colic: A perspective randomised study: 328" PEDIATRIC RESEARCH, WILLIAMS AND WILKINS, BALTIMORE, MD, US LNKD-DOI:10.1203/00006450-200508000-00356, vol. 58, no. 2, 1 August 2005 (2005-08-01) , page 411, XP003018394 ISSN: 0031-3998
KASTNER S ET AL: "Antibiotic susceptibility patterns and resistance genes of starter cultures and probiotic bacteria used in food" SYSTEMATIC AND APPLIED MICROBIOLOGY, URBAN UND FISCHER VERLAG, DE LNKD- DOI:10.1016/J.SYAPM.2005.07.009, vol. 29, no. 2, 1 March 2006 (2006-03-01), pages 145-155, XP025163627 ISSN: 0723-2020 [retrieved on 2006-03-01]
HUYG G ET AL: "Genetic Basis of Tetracycline and Minocycline Resistance in Potentially Probiotic Lactobacillus plantarum Strain CCUG 43738" ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, AMERICAN SOCIETY FOR MICROBIOLOGY, WASHINGTON, DC, US LNKD- DOI:10.1128/AAC.50.4.1550-1551.2006, vol. 50,

Fortsættes ...

no. 4, 1 April 2006 (2006-04-01), pages 1550-1551, XP003018395 ISSN: 0066-4804

ROSANDER ANNA ET AL: "Removal of antibiotic resistance gene-carrying plasmids from *Lactobacillus reuteri* ATCC 55730 and characterization of the resulting daughter strain, *L. reuteri* DSM 17938" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 74, no. 19, October 2008 (2008-10), pages 6032-6040, XP002578248 ISSN: 0099-2240

GOSWAMI A.G. ET AL.: 'Role of lactobacilli in the treatment of lactose intolerance in children with diarrhoea' INDIAN JOURNAL OF PHYSIOLOGY AND ALLIED SCIENCES vol. 48, no. 2, 1994, pages 83 - 87, XP009038713

WEIZMAN Z. ET AL.: 'Effect of a Probiotic Infant Formula on Infections in Child Care Centers: Comparison of Two Probiotic Agents' PEDIATRICS vol. 115, 2005, pages 5 - 9, XP003018393

SAVINO F. ET AL.: 'Lactobacillus reuteri ATCC 55730 versus simethicone in the treatment colic: A perspective randomised study: 328' PEDIATRIC RESEARCH vol. 58, no. 2, August 2005, page 411, XP003018394

DESCRIPTION

BACKGROUND OF INVENTION

Field of the Invention

[0001] The invention herein provides a method of selecting of lactic acid bacteria, selected for their ability of increasing cytokine IL-10 levels, useful for reducing infantile colic.

Description of the Related Art

[0002] Despite its salience in terms of both prevalence and distress the nature and causes of infantile colic have remained poorly understood. A mother's description of this condition is a baby who has been happy during the day, begins to frown, his face becomes red, he draws up his legs, screams and continues to cry for about 2-20 min after which the attack ends suddenly. Controversy arises even in the terms used to describe the condition. These terms include, "infantile colic", "evening colic" because the pain is mainly confined to evening, and "three months colic" under the pretext that it disappears after about three months from birth (Illingworth RS. Difficulties in breastfeeding. In Ronald S. Illingworth, ed. The Normal Child. 10th edn. Harcourt (India) Pvt. Ltd. 1997; 39-44). Different authors have used different definitions. The definition of Wessels that colic is paroxysms of crying for three or more hours per days for three days or more per week during a period of at least three weeks, is mostly accepted in the literature (Sondergaard C, Skajaa E, Henriksen TB. Fetal growth and infantile colic. Arch Dis Child Fetal-Neonatal Ed 2000; 83 (1): F44-47). To date, the main possible causative factors has been divided into three groups: psychosocial, gastrointestinal and neuro-developmental disorders.

[0003] Psychosocial factors include: variant of normal crying, behavioral effects of atypical parenting, and manifestation of problems in parent-infant interaction.

[0004] Gastrointestinal disorders have been implicated in colic because of the infant's leg position and grimacing during a crying spell. The gastrointestinal factors are briefly reviewed below:

Improper feeding techniques like bottle feeding, feeding in a horizontal position and lack of burping post-feeding have been considered as causative factors. Breast-feeding in the first six months has been found to be a sole protective factor. The risk of infantile colic is 1.86 times higher among non-breast fed infants (Saavedra MA, Dacosta JS, Garcias G, Horta BL, Tomasi E, Mendoca R. Infantile colic incidence and associated risk factors: a cohort study. Pediatr (Rio J) 2003; 79(2): 115-122). Lothe et al. showed that colic is from sensitivity to cow's milk whey protein (Lothe L, Lindberg T. Cow's milk whey protein elicits symptoms of infantile colic in colicky formula -fed infants: A double -blind cross over study. Pediatrics 1989; 83: 262).

[0005] In a recent study, it was concluded that food hypersensitivity could be defined as causative in only a minority of cases of infantile colic (Hill DJ, Hosking CS. Infantile colic and food hypersensitivity. J Pediatr Gastroenterol Nutr 2000; 30 (Suppl): S67-76). Recently Buchanan has shown that a trial of hypoallergenic milk in infantile colic is not supported by enough evidence (Buchanan P. Effectiveness of treatment for infantile colic. Trial of hypoallergenic milk is not supported by strong enough evidence. BMJ 1998; 317(7170): 1451-1452). Babies with infantile colic are thought to be at higher risk of allergic diseases. However, in a recent study, markers of atopy, allergic rhinitis, asthma, wheezing and peak flow variability, are shown to be comparable in infants with or without infantile colic (Castro- Rodriguez JA, Stern DA, Halonen M, Wright AL, Holberg CJ et al. Relationship between infantile colic and Asthma/ atopy: A prospective study in an unselected population. Pediatrics 2001; 108(4): 878-882).

[0006] Lactose malabsorption has been reported by authors based on breath hydrogen tests (Hyams J, Geerstama M, Etienne N, Treem W. Colonic hydrogen production in infants with colic. J Pediatr 1989; 115: 592). No differences have been found in stool hydrogen concentrations between infants with or without infantile colic. However, infants producing higher methane levels have been found to have decreased colic suggesting a role of methane production in its alleviation (Belson A, Shetty AK, Yorgin PD, Bujanover Y, Peled Y, Dar MH, Riaf S. Colonic hydrogen elimination and methane production in infants with and without infantile colic syndrome. Dig Dis Sci 2003; 48(9): 1762-1766).

[0007] Gastrointestinal (GIT) hormones such as motilin, vasoactive intestinal peptide have been shown to be abnormally high in babies with colic. Lothe et al. have shown elevated levels from first day of life in babies who later developed colic suggesting an abnormal GIT physiology in infantile colic (Lothe L, Ivansson SA, Ekman R, Lindberg T. Motilin and infantile colic: A prospective study. *Acta Pediatr Scand* 1990; 79(4): 410-416).

[0008] Neuro-developmental disorders that have been put forward suggest that abdominal cramping and colic may be a result of hyperperistalsis. The theory is supported by evidence that the use of anticholinergic agents decreases colic symptoms (Gupta SK. Is colic a gastrointestinal disorder? *Curr Opin Pediatr* 2002;14:588-92).

[0009] The fact that most infants outgrow colic by four months of age lends support to a neurodevelopmental cause of colic (Barr RG. Colic and crying syndromes in infants. *Pediatrics* 1998; 102(5 suppl E):1282-6).

[0010] A probiotic, by the generally accepted definition, is a live microbial feed supplement which beneficially affects the host animal by improving its intestinal microbial balance. Although originally referring to the supplementation of animal feeds for farm animals, the definition is easily applied to the human situation. The major consumption of probiotics by humans is in the form of dairy-based foods containing intestinal species of lactobacilli and bifidobacteria. It is implicit in the definition that consumption of the probiotic affects the composition of the intestinal microflora.

[0011] It is proposed that this effect of the probiotic on the intestinal ecosystem impacts in some beneficial way on the consumer. A number of potential benefits arising from changes to the intestinal milieu through the agency of probiotics have been documented including: increased resistance to infectious diseases, particularly of the intestine, decreased duration of diarrhea, reduction in blood pressure, reduction in serum cholesterol concentration, reduction in allergy, stimulation of phagocytosis by peripheral blood leucocytes, modulation of cytokine gene expression, adjuvant effect, regression of tumors, and reduction in carcinogen or co-carcinogen production.

[0012] Christensen et al. was one of the first reporting that probiotic lactobacilli exerted their immunomodulatory effects by modulating the Th1/Th2/Th3/Tr1/Treg-promoting capacity of dendritic cells (DCs). They showed that when murine DCs were exposed to co-cultures of different *Lactobacillus* species, including *Lactoacillus reuteri*, they were differentially modulated for production of cytokines IL-6, IL-10, IL-12, and TNF- α , and for up-regulation of MHC class II and CD86 surface markers in a concentration dependent manner. All lactobacilli upregulated surface MHC class II and CD86 markers - indicative of DC maturation. Particularly notable in these studies was that *L. reuteri* (strain 12246) was a poor IL-12 inducer, but when co-cultured with *L. johnsonii* or *L. casei*, it differentially inhibited production of the proinflammatory cytokine signals IL-12, IL-6 and TNF- α , which were stimulated by the latter two species. IL-10 production remained unaltered under these conditions.

[0013] These findings led to their conclusions that "*L. reuteri* may contribute to an environmental modulation of the intestinal dendritic cell generation favoring tolerance toward antigens bearing no 'danger signal' while at the same time keeping intact the capacity to respond against pathogens recognized via a danger signal like LPS," that "*L. reuteri* might be a potential fine-targeted treatment effective for downregulating production of IL-12 and TNF- α (and IL-6) while inducing the antiinflammatory IL-10, thus representing an alternative therapeutic approach to counterbalance the pro-inflammatory intestinal milieu," and thus "the potential exists for Th1/Th2/Th3 driving capacities of the gut to be modulated according to composition of gut microflora, including ingested probiotics." (Christensen HR, Frokiaer H, Pestka JJ (2002) Lactobacilli differentially modulate expression of cytokines and maturation surface markers in murine dendritic cells. *J Immunol* 168 171-178).

[0014] Smits et al. extended these observations and showed that *L. reuteri* has the ability to prime DCs to stimulate T regulatory (TR) cell production. They used three different *Lactobacillus* species co-cultured in vitro with human monocyte-derived DCs. Two of the lactobacilli, a human *L. reuteri* strain (ATCC 53609) and *L. casei*, but not a *L. plantarum* strain, primed these DCs to stimulate development of TR cells. These TR cells were shown to produce increased levels of IL-10 and were able to inhibit proliferation of bystander T cells in an IL-10-dependent fashion (Smits HH, Engering A, van der Kleij D, de Jong EC, Schipper K, van Capel TMM, Zaat BAJ, Yazdanbakhsh M, Wierenga EA, van Kooyk Y, Kapsenberg L (2005) Selective probiotic bacteria induce IL-10-producing regulatory T cells in vitro by modulating dendritic cell function through dendritic cell-specific intercellular adhesion molecule 3-grabbing nonintegrin. *J Allergy Clin Immunol* 115 1260-1267).

[0015] One of the first pieces of evidence concerning an effect of orally administered *L. reuteri* on the host per se was an increased CD4+/CD8+ T cell ratio observed in the chick ileum (Walter J. Dobrogosz, NUTRAfoods 2005 4(2/3) 15-28). Valeur et al. recently extended this observation to include the human ileum. They obtained direct *in situ* evidence in human subjects showing (a) distribution of individual human-specific *L. reuteri* ATCC 55730 cells throughout the human gut and (b) apparent involvement of *L. reuteri* ATCC 55730 in the recruitment and/or proliferation of CD4+ T cells specifically in the ileal regions of the intestines. They concluded that "*L. reuteri* administrations elicited a recruitment of CD4+ T-helper cells to the human epithelium.

This recruitment may be one factor in explaining the probiotic effect of this *L. reuteri* strain in man" (Valeur N, Engel P, Carbalal N, Connolly E, Ladefoged K (2004) Colonization and immunomodulation by *Lactobacillus reuteri* ATCC 55730 in the human gastrointestinal tract, *Appl Environ Microbiol* 70 1176-1181).

[0016] Toll like receptors (TLRs) recognize microbial motifs and activate a set of genes that lead to cytokine production. Traditionally, TLRs have been regarded as sensors of microbial infections, and their role is to induce an inflammatory response. However, the motifs recognized by TLRs are not unique to pathogens but are general motifs shared by entire classes of microorganisms, and it is not fully understood how the immune system differentiates between commensal and pathogenic bacteria via the TLRs. Recently, data have shown that TLRs, despite their role in induction of the inflammatory response, also play a role in maintaining intestinal homeostasis by recognizing the commensal microflora (Rakoff-Nahoum S, Paglino J, Eslami-Varzaneh F, Edberg S, Medzhitov R. Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis. *Cell*. 2004 Jul 23;118(2):229-41).

[0017] Strains of a wide variety of *Lactobacillus* species, including *L. reuteri* have been used in probiotic formulations. *L. reuteri* is one of the naturally occurring inhabitants of the gastrointestinal tract of animals, and is routinely found in the intestines of healthy animals, including humans. It is known to have antimicrobial activity. See, for example U.S. Patent Nos. 5,439,678, 5,458,875, 5,534,253, 5,837,238, and 5,849,289. When *L. reuteri* cells are grown under anaerobic conditions in the presence of glycerol, they produce the antimicrobial substance known as β -hydroxy-propionaldehyde (3-HPA).

[0018] There is a clear and complex relationship between the intestinal immune system and the commensal flora. Recently it has been demonstrated that the luminal endogenous flora can initiate the key process of bacteria-induced innate and adaptive host response through the activation of toll-like receptors (TLRs) and NOD-receptors, located on the intestinal epithelial cells (Haller D, Jobin C. Interaction between resident luminal bacteria and the host: can a healthy relationship turn sour? *J Pediatr Gastroenterol Nutr* 2004; 38: 123-36. Rakoff-Naholm S, Paglino J, Eslami-Varzaneh F, Edberg S, Medzhitov R. Recognition of commensal microflora by Toll-like receptors is required for intestinal homeostasis. *Cell* 2004; 118: 229-241.). In animal models, cytokines can initiate a hyper-reflex response of the enteric neuromusculature through neuro- and myo-immune interactions (Milla PJ. Inflammatory cells and the regulation of gut motility. *J Pediatr Gastroenterol Nutr* 2004; 39: S750.).

[0019] In experimental and clinical studies it has been demonstrated that specific probiotic strains inhibit proliferation of the T cells and reduce secretion of both Th1 and Th2 cytokines while preferentially generating suppressive cytokine such as IL-10 and TGF- β (Rautava S, Kalliomaki M, Isolauri E. Probiotics during pregnancy and breast-feeding might confer immunomodulatory protection against atopic disease in the infants. *J Allergy Clin Immunol* 2002; 109: 119-121.). Further, in human volunteers, *L. reuteri* colonizes the human gastrointestinal tract and is able to exert immunomodulatory activity, including recruitment of CD4+ T-helper cells at the human ileum epithelium (Valeur N, Engel P, Carbalal N, Connolly E, Ladefoged K. Colonization and immunomodulation by *Lactobacillus reuteri* ATCC 55730 in the human gastrointestinal tract. *Appl Environ Microbiol* 2004; 70: 1176-81). Maturation of DCs is the process that converts immature DCs to mature antigen presenting cells that migrate to lymph nodes. This process results in the loss of the powerful antigen uptake capacity that characterizes immature DC and in the upregulation of co-stimulatory molecule expression and of various cytokines (Mellman I, Steinman RM: Dendritic cells: specialized and regulated antigen processing machines. *Cell* 2001, 106:255-8. Banchereau J, Briere F, Caux C, Davoust J, Lebecque S, Liu YJ, Pulendran B, Palucka K: Immunobiology of dendritic cells. *Annu Rev Immunol* 2000, 18:767-811).

[0020] Known maturation protocols are based on the environment that DCs are believed to encounter after or during exposure to antigens. The best example of this approach is the use of monocyte-conditioned media (MCM). MCM is generated in vitro by culturing monocytes and then using the culture supernatant fluid as a source of maturation factors. The major components in MCM responsible for maturation are the pro-inflammatory cytokines interleukin-1 beta (IL-1 β), IL-6 and TNF- α (Reddy A, Sapp M, Feldman M, Subdewe M, Bhardwaj N: A monocyte conditioned medium is more effective than defined cytokines in mediating the terminal maturation of human dendritic cells. *Blood* 1997, 90:3640-6.). Mature DCs produce a variety of cytokines, which stimulate and direct the T cell response. Two of these cytokines are IL-10 and IL-12. These cytokines have opposing effects on the direction of the induced T cell response: IL-12 induces a Th1 type response whereas IL-10 inhibits such response.

[0021] Thus, the activation state of APCs (antigen presenting cells), including DCs, determines the type and magnitude of the CD4 $^{+}$ T cell response. Resting APCs (including epithelial cells in the thymus) may promote the development of CD4 $^{+}$ CD25 $^{+}$ TR cells. During infection by pathogens, recognition of microbial molecules by TLRs results in activation of APCs. The APCs then produce IL-6 and additional soluble factors that together override the suppressive effects of TR cells, allowing efficient generation of TE (T effector cells) cells against the pathogen. The dynamic equilibrium between resting and activated APCs will also be influenced by the actions of both TR and TE cells (Figure 1).

[0022] Pessi et al. (2000) describe IL-10 generation in atopic children following oral *Lactobacillus rhamnosus* GG. But unlike the invention herein the study does not describe specifically strains with the ability of producing high amounts of IL-10, selected for their efficiency against motility disorders and colic (Pessi T, Sutas Y, Hurme M, Isolauri E. Interleukin-10 generation in atopic children following oral *Lactobacillus rhamnosus*. GG. *Clin Exp Allergy*. 2000 Dec;30(12):1804-8).

[0023] A study by Hermelijn et al. (2005) shows *L. reuteri* and *Lactobacillus casei*, but not *Lactobacillus plantarum*, priming monocyte-derived DCs to drive the development of TR cells. These TR cells produced increased levels of IL-10. Unlike the invention herein, the authors do not connect the IL-10 elevation to gut motility or colic. Even when it comes to strains, the authors mention two different bacterial species being efficient in increasing IL-10 levels. This is in contrast to the invention herein where it is showed that the probiotic to be most effective reducing colic has to be determined and selected at the strain level as the different strains of the same species have a different ability to be effective in increasing IL-10 levels.

[0024] At present there is no cure for colic. The current treatment paradigm for colic consists of either pharmacological and/or non- pharmacological methods, providing at best marginal reduction of symptoms. Typical therapeutic interventions for colic offered to parents fall within four categories, including, dietary, physical, behavioral and pharmacological. Dietary manipulations include professional advice on various feeding techniques, or the use of hypoallergenic milk, soy or lactose free formulas, and an early introduction to solids (Lothe, L., et al. cow's milk formula as a cause of infantile colic: a double-blind study. *Pediatrics* 1982; 70:7-10; Forsyth B W C. Colic and the effect of changing formulas: a double- blind multiple-crossover study. *J Pediatr* 1989; 115,521- 6; Treem, W R, et al. Evaluation of the effect of a fiber-enriched formula on infant colic. *J Pediatr* 1991; 119:695-701). However, neither the use of soy formulas, or changes in feeding techniques works effectively for every case of colic. A review of the data studying these recommendations showed that use of hypoallergenic formulas, such as partially hydrolyzed or amino acid-based, may benefit only about 25% of infants (Lucassen, P L B J, et al. Infantile colic: crying time reduction with a whey hydrolysate: a double-blind, randomized placebo-controlled trial. *Pediatrics* 2000;106:1349-54; Estep, D C, et al. Treatment of infant colic with amino acid-based infant formula: a preliminary study. *Acta Paediatr* 2000;89:22- 7).

[0025] Physical strategies for the management of colic include physical movement of body positions to alleviate gas production/reflux, carrying, swaddling, applying abdominal pressure, or massaging the baby. Other methods include creating a sense of distraction to minimize infant awareness of colic such as taking an infant for a car ride, use of a car ride simulator, crib vibrator, or infant swings (Lipton E L. Swaddling and child care practice: historical, cultural and experimental observations. *Pediatrics* 1965;35:521-67; Byrne J M, Horowitz F D. Rocking as a soothing intervention: the influence of direction and type of movement. *Infant Behav Dev* 1981;4:207-18).

[0026] Another approach is to play recordings of sounds that supposedly soothe the baby. However, there is evidence in the medical literature that these methods do not work (Parkin P C, Schwartz C J, Manuel B A. Randomized controlled trial of three interventions in the management of persistent crying of infancy. *Pediatrics* 1993;92(2): 197- 201). These strategies, at best, are only marginally effective in abatement of colic symptoms.

[0027] Recommendations for behavioral interventions for the treatment of colic are the most inconsistent therapies available. Some authors advocate increasing sensory stimulation, while others advocate decreasing such stimulation (Balon A J. Management of infantile colic. *Amer Pham Physician* 1997; 55:235-242; Lucassen P L B J, Assendelft W J J, Gubbels J W, van Eijk T M, van Geldrop W J, Effectiveness of treatments for infantile colic: systematic review. *BMJ* 1998;316(5): 1563-9; and Carey W B, "Colic"--primary excessive crying as an infant-environmental interaction. *Pediatr Clin North Am* 1984;31:993-1005). Other recommendations include early response to crying, or allowing the infant to cry, offering a pacifier, implementation of a routine feeding schedule, using eye contact and interactive playing.

[0028] Pharmacologic intervention for the treatment of colic has led to the use of prescription and non-prescription medications. Currently employed prescription medications include belladonna alkaloids and opiates (paregoric), which may provide relief, but are fraught with risks including extra pyramidal symptoms, respiratory depression, and constipation. For example, anticholinergic drugs, similar in their effect to atropine, such as, Hyoscyamine (LEVISINE®, or GASTROSED®) and Dicyclomine dilate pupils, increase heart rate, decrease production of saliva, relieve spasms of gastrointestinal and urinary tracts, as well as bronchi. Although the anticholinergic drugs are the only prescription medications on the U.S. market that consistently have been shown to effectively treat infantile colic, unfortunately, up to 5% of treated infants may develop side effects, including breathing difficulties, apnea, seizures, syncope, asphyxia, coma and muscular hypotonia (Williams J, Watkin-Jones R. Dicyclomine: worrying symptoms associated with its use in some small babies (BMJ 1984;288:901; Myers J H, Moro-Sutherland D, Shook J E. Anticholinergic poisoning in colicky infants treated with hyoscyamine sulfate. *Am J Emerg Med* 1997;15:532-5). In addition, several cases of death have been reported in infants taking dicyclomine (Garriott J C, Rodriguez R, Norton L E. Two cases of death involving dicyclomine in infants. *Clinical Toxicol* 1984;22(5) :455-462).

[0029] Non-prescription medications that have been reported as effective treatment for infantile colic include several sedative or sleep- inducing drugs, including supraphysiologic (high dose) diphenhydramine (BENADRYL®), phenobarbital, chloral hydrate, and even alcohol. However, there is the potential for serious side effects associated with several of these agents in children with respiratory disease, thus limiting their widespread use in treating colic (Balon A J. Management of infantile colic. Amer Pham Physician 1997;55:235-242; Gurry D. Infantile colic. Australian Pham Phys 1994;23(3):337-34632).

[0030] A safer non-prescription medication for treatment of colic has largely included the administration of simethicone or dimethylpolysiloxane, a non-absorbable, over-the-counter drug, which reduces the size of intestinal gas bubbles. Simethicone has a very safe profile and is frequently recommended, despite several studies demonstrating that effectiveness of simethicone on infantile colic is no better than placebo (Metcalf, T J, et al., Pediatrics 1994 July; 94(1) :29-34. Sferra, T J, et al., Pediatr Clin North Am 1996 April; 43(2):489-510. Danielson, B. et al., Acta Paediatr Scand 1985 May; 74(3):446-50. Colon, A R, et al., Am Fam Physician 1989 December; 40(6):122-4.). As a result, the most common treatment for colic today is to simply wait for the baby to grow out of the condition.

[0031] Therefore, there currently is a need for safe and effective compounds and compositions and techniques that prove useful for treating colic in infants and young children. The methods of the present invention, respond to this need, providing product that can safely and effectively treat the symptoms associated with colic in infants. Patent application EP1384483 describes probiotics for treatment of gut- neuromuscular abnormalities, such as colic in babies. As a probiotic the applicants mention several different bacterial species. This is in contrast to the invention herein where the probiotic to be most effective reducing colic is a specific lactic acid bacteria strain selected to be effective in increasing IL-10 levels and not a whole bacterial species, as the inventors of the present invention has showed that there are substantial differences in stimulation IL-10 production between strains also of the same species.

[0032] In 2005 Savino demonstrated that supplementation with *L. reuteri* ATCC 55730 significantly improved colicky symptoms in breast-fed infants compared to the standard therapy with Simethicone within 7 days of treatment. The response rate to the treatment with *L. reuteri* was 95% while only 7% of infants responded to Simethicone. He presented results at the European Society for Pediatric Research (ESPR), Meeting August 31, 2005 - September 3, 2005, Siena, Italy (Paediatr Res. 2005;58(2): 411). Though Savino's results showed beneficial effects, he was not aware of the connection between certain strains promoting the production of IL-10 and the reduction of gut motility and consequently colic. The invention herein provides a method for selecting such best strains.

[0033] *L. reuteri* DSM 17938 is a daughter strain of ATCC 55730, after removal of the antibiotic resistance gene carrying plasmids (Rosander et al, Appl. Environ. Microbiol. 2008, 74(19):6032). Such plasmids may cause potential problems in probiotic cultures. Unless eliminated from the final product, they reach the consumers' intestinal tract in high viable numbers, probiotics being specifically designed for this purpose. Their safety should be verified not only with respect to virulence factors and other potential disease-causing traits, but also concerning their capability of acquiring and disseminating resistance determinants (Kastner et al., Systematic and applied microbiology, 2006, 29(2): 145-155).

[0034] WO 2005/117532 A2 describes that selected strains of Lactobacillus have the capability to increase the level of IL-10 in breast milk.

[0035] Collins described that disturbances of motor function produce a variety of symptoms that can arise in the context of inflammation or immune activation in various gut regions and embrace such common entities as esophagitis, gastritis, and idiopathic inflammatory bowel disease (IBD). These observations suggest that motility of the gut is also subject to the influence of the immune system. In this context, the motor system may play an important role in defending the gut against noxious stimuli present in the lumen (Collins S., The Immunomodulation of Enteric Neuromuscular Function: Implications for Motility and Inflammatory Disorders. Gastroenterology 1996; 111:1683-1699). This notion is reflected, for example, by the observation of Vantrappen et al., who showed that bacterial overgrowth in the small intestine is accompanied by disruption of the normal cyclic pattern of interdigestive motility in the small intestine (Vantrappen G, Jannssens J, Hellermans J, Ghoos Y. The interdigestive motor complex of normal subjects and patients with bacterial overgrowth of the small intestine. J Clin Invest 1977; 9:1158-1168).

[0036] It has been well known for many years that elevated IL-10 levels suppress over-activated immune systems. It has also previously been demonstrated that gut motility is controlled by neurological signals, that are connected to the intestinal immune system and that colic is a consequence of increased gut motility, for example, due to bacterial overgrowth.

[0037] In industrialized countries, hygienic measures begin as early as the birthing process, which disrupts the neonates capacity to uptake the mother's gut microflora. As a consequence, different microflora are established in the infant. Instead of harboring

for example *Escherichia coli* and Lactobacilli the newborns are more often colonized with *Staphylococcus aureus* and other skin bacteria.

[0038] The inventors of the present invention has made the unexpected finding that while over-growth of skin bacteria over-activates the baby's immune system, leading to over-activated gut motility and colic as a consequence, a higher number of specific gut bacteria, such as *L. reuteri* DSM 17938, with capacity of promoting IL-10 production, leads to maturation of the TR system i.e. proliferation of CD4+CD25+TR. cells. Up-regulation of CD4+CD25+ cells leads to calmed gut motility and consequently to beneficial effects on babies with colic.

[0039] On account of these findings, nonpathogenic bacterial strains were selected for IL-10 increasing properties, and surprisingly, this property of the strains were found to correlate with a decrease in infantile colic.

SUMMARY OF THE INVENTION .

[0040] The invention herein provides a method of selecting of lactic acid bacteria, selected for their capability of promoting production of IL-10 and consequently proliferation of CD4+CD25+ TR cells, for prophylaxis and/or treatment of colic.

[0041] Other objects and features of the invention will be more fully apparent from the following disclosure and appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0042]

Figure 1. Proposed role for certain selected lactic acid bacteria strains in promoting the development of CD4⁺CD25⁺TR cells.

Figure 2a. IL10 production by DC cells in bar-graph form; Figure 2b. IL10 production by DC cells in a table.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS THEREOF

[0043] Colonization of skin bacteria in the gut and deficiency of TR-cells over-activate the immune system of the new-born baby, leading to over-activated gut motility and consequently colic. A larger number of specific gut bacteria cells, such as *L. reuteri* DSM 17938, with capacity of promoting IL-10 production, leads to maturation of the TR system i.e. proliferation of CD4+CD25+TR cells. Other lactobacilli strains have previously been reported by many researchers to induce production of IL-10 for example Rautava et al (above). Up-regulation of CD4+CD25+ cells leads to calmed gut motility and consequently beneficial effects on colic. Surprisingly the strains that bring about an increase in cytokine IL-10 levels were found to be the same strains being able to decrease the average crying time (Example 2). The present invention herein comprises a method for selecting strains of lactic acid bacteria which have been selected for their capability of reducing colic, including *L. reuteri* DSM 17938. Products such as foods, nutritional additives and formulations, pharmaceuticals or medical devices containing whole cells or components derived from these strains may be formulated as is known in the art, and generally include an ingestible support as known plus the *Lactobacillus*-strain, or its derived component.

[0044] An *in vitro* study may be used as a method for selecting strains of lactobacilli for their capability of stimulating the production of IL10 by monocyte-derived DCs and thereby capacity to induce the development of CD4+CD25+TR cells (Example 1).

[0045] Data disclose an indication of a powerful stimulation of IL-10 by the specific strains *L. reuteri* ATCC 55730 and *L. reuteri* DSM 17938, and that this regulation is mediated by a substance released into the growth medium by these two specific strains during late log/stationary growth phase. On the contrary, the two other strains of *L. reuteri* were unable to stimulate relevant IL-10 production.

[0046] To confirm the clinical relevance of the selected strains for preventing or treating colic further studies are performed in breast-fed infants with a diagnosis of infantile colic.

[0047] The features of the present invention will be more clearly understood by reference to the following examples, which are not to be construed as limiting the invention.

EXAMPLE 1.

[0048] Study of probiotic strains ability to promote IL-10 expression by monocyte-derived DC:

Immature DCs are generated from peripheral blood monocytes (Hilkens, C. M. U., P. Kalinski, M. de Boer, and Kapsenberg. 1997. Human dendritic cells require exogenous interleukin-12-inducing factors to direct the development of naive T-helper cells toward the Th1 phenotype. *Blood* 90:1920) cultured in IMDM (Life Technologies, Paisley, U.K.) containing 10% FCS HyClone, Logan, UT), recombinant human (rh)GM-CSF (500 U/ml; Schering-Plough, Uden, The Netherlands), and rhIL-4 (250 U/ml; Pharma Biotechnologie Hannover, Hannover, Germany) (Kalinski, P., J. H. N. Schuitemaker, C. M. U. Hilkens, E. A. Wierenga, and M. L. Kapsenberg. 1999. Final maturation of dendritic cells is associated with impaired responsiveness to IFN- γ and to bacterial IL-12 inducers: decreased ability of mature dendritic cells to produce IL-12 during the interaction with Th cells. *J. Immunol.* 162:3231).

[0049] Strains to be tested in this example are *Lactobacillus reuteri* ATCC 55730, *Lactobacillus reuteri* DSM 17938, *Lactobacillus reuteri* ATCC PTA 4660 and *Lactobacillus reuteri* ATCC PTA 4964 obtainable from ATCC (Manassas, VA, USA and DSMZ, Braunschweig, Germany). Strains are cultured on Columbia agar (Oxoid, Basingstoke, United Kingdom) containing 6.25% sheep blood. Lactobacilli are incubated at 37°C in a 5% CO₂ atmosphere. After 3 days, the number of bacteria is determined by measuring the optical density at 620 nm (OD620). Typically, an OD620 of 0.35 corresponded to 1x10⁸ colony-forming units (cfu) for all test strains.

[0050] On day 6, maturation of immature DCs is induced by LPS (E coli; Sigma-Aldrich, St Louis, Mo) and the test *Lactobacilli* in the presence of the combination of the cytokines IL-1 β (25 ng/mL) and TNF- α (50 ng/mL), together all used as maturation factors (cytokines purchased from Peprotech, Rocky Hill, NJ).

[0051] IL-12, IL-10, and IL-6 cytokine production by mature DCs is determined by means of a 24-hour stimulation with CD40 ligand-expressing mouse plasmacytoma cells (J558), as described by Vieira PL, et. al. (Vieira PL, de Jong EC, Wierenga EA, Kapsenberg ML, Kalinski P. Development of Th1-inducing capacity in myeloid dendritic cells requires environmental instruction. *J Immunol* 2000;164:4507-12). Supernatants are harvested after 24 h, and the concentrations of IL-10 are measured by ELISA. For results see Figure 2. As can be seen from the results, there is a substantial difference in influence by different *Lactobacillus* strains in their ability to promote IL-10 production by the DCs.

EXAMPLE 2

Selected probiotic strains versus Simethicone in the treatment of infantile colic

[0052] Breast-fed infants with a diagnosis of infantile colic are recruited in the Department of Pediatric and Adolescence Science (Drottning Silvias University Hospital, Göteborg). Patients aged 21-90 days, adequate for gestational age with a birth weight between 2500 and 4000g, with colic symptoms satisfying the Wessel's criteria, and arisen almost 6 ± 1 days before the enrollment, are considered for inclusion in the study (Wessel MA, Cobb JC, Jackson EB, Harris GS, Detwiler AC. Paroxysmal fussing in infancy, sometimes called "colic". *Pediatrics* 1954; 14: 421-35). All infants enrolled are exclusively breast-fed in order to reduce variability in the intestinal microflora due to variations in the diet, which may in turn have influenced the response to the probiotic. Infants are excluded if they had clinical evidence of chronic illness or gastrointestinal disorders or if they had received either antibiotics or probiotics in the week preceding recruitment.

[0053] In this study, colicky infants are randomized to receive the probiotic or Simethicone in the following treatments:

(P1) *Lactobacillus reuteri* strain ATCC 55730 also called SD2112,

(P2) *Lactobacillus reuteri* strain DSM 17938 (deposited at DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen

GmbH (Mascheroder Weg 1b, D-38124 Braunschweig) on February 6, 2006, under the Budapest Treaty)

(P3) *Lactobacillus reuteri* strain ATCC PTA 4660 and

(S) Simethicone

[0054] *L. reuteri* is administered at a dose of 10^8 colony forming units (CFU) in 5 drops of a commercially available MCT-oil (medium chain tri-glyceride oil) suspension 30 min after feeding, once a day for 28 days. MCT oils occur naturally, and the most abundant source is coconut oil. Most MCT oil is refined from coconut oil. MCT oil is a clear light colored liquid with no flavor and low viscosity. This oil suspension is stable for 21 months at 2-8°C (as documented by the manufacturer, BioGaia AB, Stockholm, Sweden) (alternative source of MCT oil is Akomed R, by Karlshamns AB, Karlshamn, Sweden). During the study, parents are instructed to keep the product in the refrigerator when not in use. Simethicone (S) was given at a dose of 60 mg/day in 3mL of a commercially available solution (Mylicon Infants Gas Relief Drops, J&J, 7050 Camp Hill Rd., Fort Washington, Pennsylvania, 19034-2210, USA) after feeding, twice a day for 28 days. At the enrollment, all mothers are asked to follow a cows' milk-free diet as reported: avoidance of milk, yogurt, fresh and strong cheese, cream, butter, biscuits. Adherence to the diet is monitored with a diet diary maintained for all the treatment period. At day 7, 14, 21 and 28 one of the researchers examined the compliance to this diet.

[0055] The Ethical Committee at the Institution approved the study protocol and infants are enrolled in the study only after written informed consent was obtained from the parents.

Follow-up visits

[0056] The day on which the pediatrician saw the infant for the first time is defined as Day -1. On this occasion each infant underwent a medical examination and the parents were interviewed in order to obtain background data concerning type of delivery, birth weight and gestational age, family history of gastrointestinal disease and atopy. In particular, the last one is considered positive if the infants had one or more family members (mother, father and /or older sibling) with atopic eczema, allergic rhinitis or asthma. Moreover, any signs and symptoms of atopic disease during the study period were recorded. Parents were also invited to record data concerning daily average crying time and number of colic episodes on the day after the recruitment (Day 0). The doctor randomly assigned the child to any of the study groups. Administration of study products began on Day 1.

[0057] Parents are given written information about the study and are asked to record daily number of inconsolable crying episodes and their duration, stool consistency and frequency as well as any observed side effects (constipation, vomiting, cutaneous reactions, etc) starting from Day 0 up to Day 28, using a structured diary. In order to ensure that all the parents noted crying time in a uniform way and to ensure that the infants are given the medication correctly, one of the researchers is always available by phone to help parents.

[0058] Each patient is re-examined by the same pediatrician on Days 1, 7, 14, 21 and 28.

Results

[0059] Of the 120 breast-fed colicky infants enrolled, 30 are randomly assigned to treatment with *L. reuteri* ATCC 55730 (P1), 30 are randomly assigned to treatment with *L. reuteri* DSM 17938 (P2), 30 are randomly assigned to treatment with *L. reuteri* ATCC PTA 4660 (P3) and 30 to Simethicone (S). No babies are withdrawn because of any side effect related to the trial. The groups are similar with respect to age, birth weight, gender, type of delivery, family history of atopy or gastrointestinal diseases, and exposure to smoking.

[0060] The average crying time per day is similar in the treatment groups on Day 0, and on Day 1. Infants receiving *L. reuteri* ATCC 55730 (P1) and DSM 17938 (P2) showed a significant reduction in daily crying time by Day 7 compared to infants treated with ATCC PTA 4660 and Simethicone(S). On Day 14, 21 and 28 crying time is significantly different between the four treatment groups. The difference between (P1) and (S) in average crying time per day from the beginning to the end of the study is 82 minutes, and the difference between (P2) and (S) in average crying time per day from the beginning to the end of the study is 83 minutes on Day 28 (Table 1). As can be seen from the results are the four treatment groups fairly even before the administration

of the test products. Already after 7 days there is a clear difference in favor of *L.reuteri* strains ATCC 55730 and DSM 17938 over the two other alternatives. After 28 days the difference has grown larger.

Table 1. Crying time (mean minutes per day) in *L. reuteri* and Simethicone groups.

Colicky infants N=120	<i>L. reuteri</i> ATCC55730 N= 30	<i>L. reuteri</i> DSM 17938 N= 30	<i>L. reuteri</i> ATCC PTA 4660 N= 30	Simethicone N=30
Day 0	206	203	204	205
Day 1	191	190	198	194
Day 7	147	148	171	172
Day 14	98	100	156	155
Day 21	75	77	150	148
Day 28	58	59	145	141

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US5439678A \[0017\]](#)
- [US5458875A \[0017\]](#)
- [US5534253A \[0017\]](#)
- [US5837238A \[0017\]](#)
- [US5849289A \[0017\]](#)
- [EP1384483A \[0031\]](#)
- [WO2005117632A2 \[0034\]](#)

Non-patent literature cited in the description

- **ILLINGWORTH RS.** Difficulties in breastfeedingPvt. Ltd. 1997000039-44 [\[0002\]](#)
- **SONDERGAARD CSKAJAA EHENRIKSEN TB.** Fetal growth and infantile colic20000000vol. 83, F44-47 [\[0002\]](#)
- **SAAVEDRA MADACOSTA JS GARCIAS GHORTA BL TOM ASI EMENDOCA R.** Infantile colic incidence and associated risk factors: a cohort study.Pediatr (Rio J), 2003, vol. 79, 2115-122 [\[0004\]](#)
- **LOTHE LLINDBERG T.** Cow's milk whey protein elicits symptoms of infantile colic in colicky formula -fed infants: A double - blind cross over studyPediatrics, 1989, vol. 83, 262- [\[0004\]](#)
- **HILL DJHOSKING CS.** Infantile colic and food hypersensitivityJ Pediatr Gastroenterol Nutr, 2000, vol. 30, 67-76 [\[0005\]](#)
- **BUCHANAN P.** Effectiveness of treatment for infantile colic. Trial of hypoallergenic milk is not supported by strong enough evidenceBMJ, 1998, vol. 317, 71701451-1452 [\[0005\]](#)
- **CASTRO- RODRIGUEZ JASTERN DAHALONEN MWRIGHT ALHOLBERG CJ et al.** Relationship between infantile colic and Asthma/ atopy: A prospective study in an unselected populationPediatrics, 2001, vol. 108, 4878-882 [\[0005\]](#)
- **HYAMS JGEERSTAMA METIENNE NTREEM W.** Colonic hydrogen production in infants with colicJ Pediatr, 1989, vol. 115, 592- [\[0006\]](#)
- **BELSON ASHETTY AKYORGIN PDBUJANOVER YPELED YDAR MHRIAF S.** Colonic hydrogen elimination and methane production in infants with and without infantile colic syndrome.Dig Dis Sci, 2003, vol. 48, 91762-1766 [\[0006\]](#)
- **LOTHE LIVASSON SAEKMAN RLINDBERG T.** Motilin and infantile colic: A prospective study.Acta Pediatr Scand, 1990, vol. 79, 4410-416 [\[0007\]](#)

- **GUPTA SK.** Is colic a gastrointestinal disorder? *Curr Opin Pediatr*, 2002, vol. 14, 588-92 [0008]
- **BARR RG.** Colic and crying syndromes in infants *Pediatrics*, 1998, vol. 102, 51282-6 [0009]
- **CHRISTENSEN HRFROKIAER HPESTKA JJ.** Lactobacilli differentially modulate expression of cytokines and maturation surface markers in murine dendritic cells *J Immunol*, 2002, vol. 168, 171-178 [0010]
- **SMITS HHENGERING AVAN DER KLEIJ DDE JONG ECSCHIPPER KVAN CAPEL TMM ZAAT BAJ. YAZDANBAKHSH MWIERENGA EAVAN KOOYK Y.** Selective probiotic bacteria induce IL-10-producing regulatory T cells in vitro by modulating dendritic cell function through dendritic cell-specific intercellular adhesion molecule 3-grabbing nonintegrin *J Allergy Clin Immunol*, 2005, vol. 115, 1260-1267 [0014]
- **WALTER J. DOBROGOSENUTRA** foods, 2005, vol. 4, 2/315-28 [0015]
- **VALEUR NENGEL PCARBAJAL NCONNOLLY ELADEFOGED K.** Colonization and immunomodulation by *Lactobacillus reuteri* ATCC 55730 in the human gastrointestinal tract *Appl Environ Microbiol*, 2004, vol. 70, 1176-1181 [0015]
- **RAKOFF-NAHOUM SPAGLINO JESLAMI-VARZANEH FEDBERG SMEDZHITOVA R.** Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis *Cell*, 2004, vol. 118, 2229-41 [0016]
- **HALLER DJOBIN C.** Interaction between resident luminal bacteria and the host: can a healthy relationship turn sour? *J Pediatr Gastroenterol Nutr*, 2004, vol. 38, 123-36 [0018]
- **RAKOFF-NAHOLM SPAGLINO JESLAMI-VARZANEH FEDBERG SMEDZHITOVA R.** Recognition of commensal microflora by Toll-like receptors is required for intestinal homeostasis *Cell*, 2004, vol. 118, 229-241 [0018]
- **MILLA PJ.** Inflammatory cells and the regulation of gut motility *J Pediatr Gastroenterol Nutr*, 2004, vol. 39, S750- [0018]
- **RAUTAVA SKALLIOMAKI MISOLAUER E.** Probiotics during pregnancy and breast-feeding might confer immunomodulatory protection against atopic disease in the infants *J Allergy Clin Immunol*, 2002, vol. 109, 119-121 [0019]
- **VALEUR NENGEL PCARBAJAL NCONNOLLY ELADEFOGED K.** Colonization and immunomodulation by *Lactobacillus reuteri* ATCC 55730 in the human gastrointestinal tract *Appl Environ Microbiol*, 2004, vol. 70, 1176-81 [0019]
- **MELLMAN ISTEINMAN RM** Dendritic cells: specialized and regulated antigen processing machines *Cell*, 2001, vol. 106, 255-8 [0019]
- **BANCHEREAU JBRIERE FCAUX CDAVOUST JLEBECQUE SLIU YJPULENDRAN BPALUCKA K.** Immunobiology of dendritic cells *Annu Rev Immunol*, 2000, vol. 18, 767-811 [0019]
- **REDDY ASAPP MFELDMAN MSUBLDEWE MBHARDWAJ NA** monocyte conditioned medium is more effective than defined cytokines in mediating the terminal maturation of human dendritic cells *Blood*, 1997, vol. 90, 3640-6 [0020]
- **PESSI TSUTAS YHURME MISOLAUER E.** Interleukin-10 generation in atopic children following oral *Lactobacillus rhamnosus* GG *Clin Exp Allergy*, 2000, vol. 30, 121804-8 [0022]
- **LOTHE, L. et al.** cow's milk formula as a cause of infantile colic: a double-blind study *Pediatrics*, 1982, vol. 70, 7-10 [0024]
- **FORSYTH B W C.** Colic and the effect of changing formulas: a double-blind multiple-crossover study *J Pediatr*, 1989, vol. 115, 521-6 [0024]
- **TREEM, W R et al.** Evaluation of the effect of a fiber-enriched formula on infant colic *J Pediatr*, 1991, 119695-701 [0024]
- **LUCASSEN, P L B J et al.** Infantile colic: crying time reduction with a whey hydrolysate: a double-blind, randomized placebo-controlled trial *Pediatrics*, 2000, vol. 106, 1349-54 [0024]
- **ESTEP, D C et al.** Treatment of infant colic with amino acid-based infant formula: a preliminary study *Acta Paediatr*, 2000, vol. 89, 22-7 [0024]
- **LIPTON E L.** Swaddling and child care practice: historical, cultural and experimental observations *Pediatrics*, 1965, vol. 35, 521-67 [0025]
- **BYRNE J MHOROWITZ F D.** Rocking as a soothing intervention: the influence of direction and type of movement *Infant Behav Dev*, 1981, vol. 4, 207-18 [0025]
- **PARKIN P CSCHWARTZ C JMANUEL B A.** Randomized controlled trial of three interventions in the management of persistent crying of infancy *Pediatrics*, 1993, vol. 92, 2197-201 [0026]
- **BALON A J.** Management of infantile colic *Amer Pham Physician*, 1997, vol. 55, 235-242 [0027]
- **LUCASSEN P L B JASSENDELFT W J JGUBBELS J W VAN EIJK T M VAN GELDROP W J.** Effectiveness of treatments for infantile colic: systematic review *BMJ*, 1998, vol. 316, 51563-9 [0027]
- **CAREY W B.** "Colic"--primary excessive crying as an infant-environmental interaction *Pediatr Clin North Am*, 1984, vol. 31, 993-1005 [0027]
- **WILLIAMS JWATKIN-JONES R.** Dicyclomine: worrying symptoms associated with its use in some small babies *BMJ*, 1984, vol. 288, 901- [0028]
- **MYERS J HMORO-SUTHERLAND DSHOOK J E.** Anticholinergic poisoning in colicky infants treated with hyoscymamine sulfate *Am J Emerg Med*, 1997, vol. 15, 532-5 [0028]
- **GARIOTT J CRODRIGUEZ RNORTON L E.** Two cases of death involving dicyclomine in infants *Clinical Toxicol*, 1984, vol. 22, 5455-462 [0028]
- **BALON A J.** Management of infantile colic *Amer Pham Physician*, 1997, vol. 55, 235-242 [0029]
- **GURRY D.** Infantile colic *Australian Pham Phys*, 1994, vol. 23, 3337-34632 [0029]
- **METCALF, T J et al.** *Pediatrics*, 1994, vol. 94, 129-34 [0029]

- **SFERRA, T J et al.** Pediatr Clin North Am, 1996, vol. 43, 2489-510 [0039]
- **DANIELSON, B. et al.** Acta Paediatr Scand, 1985, vol. 74, 3446-50 [0020]
- **COLON, A R et al.** Am Fam Physician, 1989, vol. 40, 6122-4 [0030]
- Paediatr Res., 2005, vol. 58, 2411- [0032]
- **ROSANDER et al.** Appl. Environ. Microbiol., 2008, vol. 74, 196032- [0033]
- **CASTNER et al.** Systematic and applied microbiology, 2006, vol. 29, 2145-155 [0033]
- **COLLINS S.** The Immunomodulation of Enteric Neuromuscular Function: Implications for Motility and Inflammatory Disorders Gastroenterology, 1996, vol. 111, 1683-1699 [0035]
- **VANTRAPPEN GJANNSENS JHELEMANS JGHOOS Y.** The interdigestive motor complex of normal subjects and patients with bacterial overgrowth of the small intestine J Clin Invest, 1977, vol. 9, 1158-1168 [0035]
- **HILKENS, C. M. U.P. KALINSKIM. DE BOERKAPSENBERG.** Human dendritic cells require exogenous interleukin-12-inducing factors to direct the development of naive T-helper cells toward the Th1 phenotype. Blood, 1997, vol. 90, 1920- [0048]
- **KALINSKI, P. J. H. N. SCHUITEMAKER C. M. U. HILKENSE. A. WIERENGAM. L. KAPSENBERG** Final maturation of dendritic cells is associated with impaired responsiveness to IFN- γ and to bacterial IL-12 inducers: decreased ability of mature dendritic cells to produce IL-12 during the interaction with Th cells J. Immunol., 1999, vol. 162, 3231- [0048]
- **VIEIRA PLDE JONG ECWIERENGA EAKAPSENBERG M L KALINSKI P.** Development of Th1-inducing capacity in myeloid dendritic cells requires environmental instruction J. Immunol., 2000, vol. 164, 4507-12 [0051]
- **WESSEL MACOBB JCJACKSON EBHARRIS GSDETWILER AC.** Paroxysmal fussing in infancy, sometimes called "colic" Pediatrics, 1954, vol. 14, 421-35 [0052]

Patentkrav

1. Fremgangsmåde til udvælgelse af en mælkesyrebakteriestamme, som kan anvendes til reduktion af spædbarnskolik, hvilken fremgangsmåde omfatter
5 bestemmelse af en mælkesyrebakteriestammes evne til at øge IL-10-niveauer.
2. Fremgangsmåde ifølge krav 1, hvor mælkesyrebakteriestammen kan anvendes til profylakse og/eller behandling af kolik.
- 10 3. Fremgangsmåde ifølge krav 1, hvilken fremgangsmåde omfatter *in vitro*-bestemmelse af mælkesyrebakteriestammens evne til at stimulere fremstillingen af IL-10 ved hjælp af monocytafledte dendritiske celler (DC) og derved deres evne til at fremkalde udvikling af CD4+CD25+ TR-celler.

DRAWINGS

Figure 1




Figure 2a.




Figure 2b.

	IL-10 production (pg/ml)
MRS	0
ATCC 55730	1500
DSM 17938	1450
ATCC PTA 4660	35
ATCC PTA 4964	47