




(86) Date de dépôt PCT/PCT Filing Date: 2006/01/13
(87) Date publication PCT/PCT Publication Date: 2006/07/20
(85) Entrée phase nationale/National Entry: 2007/07/06
(86) N° demande PCT/PCT Application No.: US 2006/001308
(87) N° publication PCT/PCT Publication No.: 2006/076620
(30) Priorités/Priorities: 2005/01/14 (US60/644,354);
2006/01/12 (US11/330,868)

(51) Cl.Int./Int.Cl. *A61K 9/19* (2006.01),
A61K 31/4184 (2006.01), *A61P 35/00* (2006.01),
A61P 35/02 (2006.01)
(71) Demandeur/Applicant:
CEPHALON, INC., US
(72) Inventeurs/Inventors:
BRITTAIN, JASON EDWARD, US;
FRANKLIN, JOE CRAIG, US
(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSITIONS PHARMACEUTIQUES DE BENDAMUSTINE
(54) Title: BENDAMUSTINE PHARMACEUTICAL COMPOSITIONS FOR LYOPHILISATION



(57) **Abrégé/Abstract:**

The present invention provides pharmaceutical formulations of lyophilized bendamustine suitable for pharmaceutical use. The present invention further provides methods of producing lyophilized bendamustine. The pharmaceutical formulations can be used for any disease that is sensitive to treatment with bendamustine, such as neoplastic diseases. Preferred solvent used in a pre-lyophilized composition of bendamustine is tertiary-butyl alcohol.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 July 2006 (20.07.2006)

PCT

(10) International Publication Number
WO 2006/076620 A3

(51) International Patent Classification:

A61K 9/19 (2006.01) A61P 35/02 (2006.01)
A61K 31/4184 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2006/001308

(22) International Filing Date: 13 January 2006 (13.01.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/644,354 14 January 2005 (14.01.2005) US
11/330,868 12 January 2006 (12.01.2006) US

(71) Applicant (for all designated States except US):
CEPHALON, INC. [US/US]; 41 Moores Road, P.O.
Box 4011, Frazer, Pennsylvania 19355 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRITTAIN, Jason, Edward** [US/US]; 1580 Chiswick Court, El Cajon, California 92020 (US). **FRANKLIN, Joe, Craig** [US/US]; 3708 45th Street, Tulsa, Oklahoma 74135 (US).

(74) Agent: **MILLER, Suzanne, E.**; Woodcock Washburn LLP, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania 19103 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report

(88) Date of publication of the international search report:

28 December 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BENDAMUSTINE PHARMACEUTICAL COMPOSITIONS FOR LYOPHILISATION



(57) Abstract: The present invention provides pharmaceutical formulations of lyophilized bendamustine suitable for pharmaceutical use. The present invention further provides methods of producing lyophilized bendamustine. The pharmaceutical formulations can be used for any disease that is sensitive to treatment with bendamustine, such as neoplastic diseases. Preferred solvent used in a pre-lyophilized composition of bendamustine is tertiary-butyl alcohol.

WO 2006/076620 A3

BENDAMUSTINE PHARMACEUTICAL COMPOSITIONS

FIELD OF THE INVENTION

5 The present invention pertains to the field of pharmaceutical compositions for the treatment of various disease states, especially neoplastic diseases and autoimmune diseases. Particularly, it relates to pharmaceutical formulations comprising nitrogen mustards, particularly the nitrogen mustard bendamustine, e.g., bendamustine HCl.


10 BACKGROUND OF THE INVENTION

 The present invention claims the benefit of and priority to US Serial No. 60/644,354, filed January 14, 2005, entitled, "Bendamustine Pharmaceutical Compositions," which is incorporated herein by reference in its entirety, including figures and claims.

15 The following description includes information that may be useful in understanding the present invention. It is not an admission that any such information is prior art, or relevant, to the presently claimed inventions, or that any publication specifically or implicitly referenced is prior art.

 Because of their high reactivity in aqueous solutions, nitrogen mustards are
20 difficult to formulate as pharmaceuticals and are often supplied for administration in a lyophilized form that requires reconstitution, usually in water, by skilled hospital personnel prior to administration. Once in aqueous solution, nitrogen mustards are subject to degradation by hydrolysis, thus, the reconstituted product should be administered to a patient as soon as possible after its reconstitution.

25 Bendamustine, (4-{5-[Bis(2-chloroethyl)amino]-1-methyl-2-benzimidazolyl} butyric acid, is an atypical structure with a benzimidazole ring, whose structure includes an active nitrogen mustard (see Formula I, which shows bendamustine hydrochloride).



Formula I

Bendamustine was initially synthesized in 1963 in the German Democratic Republic (GDR) and was available from 1971 to 1992 in that location under the name Cytostasan®. Since that time, it has been marketed in Germany under the tradename
5 Ribomustin®. It has been widely used in Germany to treat chronic lymphocytic leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, multiple myeloma, and breast cancer.

Due to its degradation in aqueous solutions (like other nitrogen mustards), bendamustine is supplied as a lyophilized product. The current lyophilized
10 formulation of bendamustine (Ribomustin®) contains bendamustine hydrochloride and mannitol in a sterile lyophilized form as a white powder for intravenous use following reconstitution. The finished lyophilisate is unstable when exposed to light. Therefore, the product is stored in brown or amber-colored glass bottles. The current lyophilized formulation of bendamustine contains degradation products that may occur during
15 manufacturing of the drug substance and/or during the lyophilization process to make the finished drug product.

Currently bendamustine is formulated as a lyophilized powder for injection with 100 mg of drug per 50 mL vial or 25 mg of drug per 20 mL vial. The vials are opened and reconstituted as close to the time of patient administration as possible. The
20 product is reconstituted with 40 mL (for the 100 mg presentation) or 10 mL (for the 25 mg presentation) of Sterile Water for Injection. The reconstituted product is further diluted into 500 mL, q.s., 0.9% Sodium Chloride for Injection. The route of administration is by intravenous infusion over 30 to 60 minutes.

Following reconstitution with 40 mL Sterile Water for Injection, vials of
25 bendamustine are stable for a period of 7 hours under room temperature storage or for 6 days upon storage at 2-8°C. The 500 mL admixture solution must be administered to the patient within 7 hours of vial reconstitution (assuming room temperature storage of the admixture).

The reconstitution of the present bendamustine lyophilized powder is difficult.
30 Reports from the clinic indicate that reconstitution can require at least fifteen minutes and may require as long as thirty minutes. Besides being burdensome and time-consuming for the healthcare professional responsible for reconstituting the product, the lengthy exposure of bendamustine to water during the reconstitution process

increases the potential for loss of potency and impurity formation due to the hydrolysis of the product by water.

Thus, a need exists for lyophilized formulations of bendamustine that are easier to reconstitute and which have a better impurity profile than the current lyophilate
5 (lyophilized powder) formulations of bendamustine.

German (GDR) Patent No. 34727 discloses a method of preparing ω -[5-bis-(β -chloroethyl)-amino-benzimidazolyl-(2)]-alkane carboxylic acids substituted in the 1-position.

German (GDR) Patent No. 80967 discloses an injectable preparation of γ -[1-methyl-5-bis-(β -chloroethyl)-amino-benzimidazolyl-(2)]-butric acid hydrochloride.
10

German (GDR) Patent No. 159877 discloses a method for preparing 4-[1-methyl-5-bis (2-chloroethyl) amino-benzimidazolyl-2)-butyric acid.

German (GDR) Patent No. 159289 discloses an injectable solution of bendamustine.

15 Ribomustin® bendamustine Product monograph (updated 1/2002)
http://www.ribosepharm.de/pdf/ribomustin_bendamustin/productmonograph.pdf
provides information about Ribomustin® including product description.

Ni et al. report that the nitrosourea SarCNU was more stable in pure tertiary butanol than in pure acetic acid, dimethyl sulfoxide, methylhydroxy, water or in
20 TBA/water mixtures (Ni et al. (2001) *Intl. J. Phamaceutics* 226:39-46).

Lyophilized cyclophosphamide is known in the art see e.g., US Patent Nos. 5,418,223; 5,413,995; 5,268,368; 5,227,374; 5,130,305; 4,659,699; 4,537,883; and 5,066,647.

The lyophilized nitrogen mustard Ifosfamide is disclosed in International
25 Publication No. WO 2003/066027; US Pat. Nos. 6,613,927; 5,750,131; 5,972,912; 5,227,373; and 5,204,335.


Teagarden et al. disclose lyophilized formulations of prostaglandin E-1 made by dissolving PGE-1 in a solution of lactose and tertiary butyl alcohol (US Pat. No. 5,770,230).

30

SUMMARY OF THE INVENTION

The present invention is directed to stable pharmaceutical compositions of nitrogen mustards, in particular lyophilized bendamustine and its use in treatment of various disease states, especially neoplastic diseases and autoimmune diseases.


An embodiment of the invention is a pharmaceutical composition of bendamustine containing not more than about 0.5% to about 0.9% (area percent of bendamustine) HP1, as shown in Formula II,



Formula II


at the time of release or where the HP1 is the amount of HP1 present at time zero after reconstitution of a lyophilized pharmaceutical composition of bendamustine as described herein. In a preferred embodiment is a pharmaceutical composition of bendamustine containing not more than about 0.5% (area percent of bendamustine) HP1, preferably not more than about 0.45%, more preferably not more than about 0.40%, more preferably not more than about 0.35%, even more preferably not more than 0.30%.

Another embodiment of the invention is a lyophilized preparation of bendamustine containing not more than about 0.1 % to about 0.3 % bendamustine dimer as shown in Formula III at release or at time zero after reconstitution



Formula III.

Yet another embodiment of the invention is a lyophilized preparation of bendamustine containing not more than about 0.5%, preferably 0.15% to about 0.5%, bendamustine ethylester, as shown in Formula IV at release or at time zero after reconstitution



Formula IV.

Yet another embodiment of the invention is a lyophilized preparation of bendamustine wherein the concentration of bendamustine ethylester (Formula IV) is no more than 0.2%, preferably 0.1%, greater than the concentration of bendamustine ethylester as found in the drug substance used to make the lyophilized preparation.

In another embodiment of the invention is a lyophilized preparation of bendamustine containing not more than about 0.5% to about 0.9% (area percent of bendamustine) HP1 at the time of drug product release. In a preferred embodiment is a lyophilized preparation of bendamustine containing not more than about 0.50% (area percent of bendamustine) HP1, preferably not more than about 0.45%, more preferably not more than about 0.40%, more preferably not more than about 0.35%, even more preferably not more than 0.30%. An aspect of this embodiment is lyophilized preparations of bendamustine containing not more than about 0.5% to about 0.9%, preferably 0.5%, (area percent of bendamustine) HP1 at the time of release of drug product where the lyophilized preparation is packaged in a vial or other pharmaceutically acceptable container.

In yet another aspect of the invention, the lyophilized preparations of bendamustine are stable with respect to the amount of HP1 for at least about 6 months, preferably 12 months, preferably 24 months, to about 36 months or greater when stored at about 2° to about 30°. Preferred temperatures for storage are about 5° C and about room temperature.

Another embodiment of the invention is a pharmaceutical dosage form that includes a pharmaceutical composition of bendamustine containing not more than about 0.5% to about 0.9% HP1, preferably not more than about 0.50%, preferably not more than about 0.45%, more preferably not more than about 0.40%, more preferably not more than about 0.35%, even more preferably not more than 0.30%, where the HP1 is the amount of HP1 present at release or at time zero after reconstitution of a lyophilized preparation of bendamustine of the present invention. In preferred aspects

of the invention, the dosage form can be about 5 to about 500 mg of bendamustine, about 10 to about 300 mg of bendamustine, about 25 mg of bendamustine, about 100 mg of bendamustine, and about 200 mg of bendamustine.

Yet another embodiment of the invention is a pharmaceutical dosage form that includes a lyophilized preparation of bendamustine containing not more than about 0.5% to about 0.9%, preferably 0.5%, HP1. Preferred dosage forms can be about 5 to about 500 mg of bendamustine, about 10 to about 300 mg of bendamustine, about 25 mg of bendamustine, about 100 mg of bendamustine, and about 200 mg of bendamustine.

In still another embodiment, the invention includes a pharmaceutical composition of bendamustine including bendamustine containing not more than about 0.5% to about 0.9% (area percent of bendamustine), preferably not more than about 0.50%, preferably not more than about 0.45%, more preferably not more than about 0.40%, more preferably not more than about 0.35%, even more preferably not more than 0.30%, and a trace amount of one or more organic solvents, wherein said HP1 is the amount of HP1 present at release or time zero after reconstitution of a lyophilized pharmaceutical composition of bendamustine as disclosed herein. In different aspects of this embodiment, the organic solvent is selected from one or more of tertiary butanol, n-propanol, n-butanol, isopropanol, ethanol, methanol, acetone, ethyl acetate, dimethyl carbonate, acetonitrile, dichloromethane, methyl ethyl ketone, methyl isobutyl ketone, 1-pentanol, methyl acetate, carbon tetrachloride, dimethyl sulfoxide, hexafluoroacetone, chlorobutanol, dimethyl sulfone, acetic acid, and cyclohexane. Preferred organic solvents include one or more of ethanol, methanol, propanol, butanol, isopropanol, and tertiary butanol. A more preferred organic solvent is tertiary butanol, also known as TBA, t-butanol, tert-butyl alcohol or tertiary butyl alcohol.

The present invention involves a method for obtaining agency approval for a bendamustine product, the improvement which includes setting a release specification for bendamustine degradants at less than about 4.0%, preferably about 2.0 % to about 4.0 %, (area percent bendamustine) or otherwise to achieve the pharmaceutical compositions described herein. An aspect of this embodiment is a method for obtaining agency approval for a bendamustine product which includes setting a release specification for HP1 to be less than or equal to 1.5% (area percent Bendamustine).

The bendamustine product herein contains not more than about 0.5% (area percent of bendamustine) HP1 at release.

Another embodiment is a method for obtaining agency approval for a bendamustine product, the improvement which includes setting a shelf-life
5 specification for bendamustine degradants at less than about 7.0%, preferably about 5.0% to about 7.0%, (area percent bendamustine) where the product is stored at about 2°C to about 30°C. Preferred temperatures for storage are about 5°C and about room temperature. The bendamustine product herein contains not more than about 0.5% (area percent of bendamustine) HP1 at release.

10 Another embodiment of the invention is a process for manufacturing a lyophilized preparation of bendamustine which includes controlling for the concentration of bendamustine degradants in the final product, such that the concentration of bendamustine degradants is less than about 4.0%, preferably no more than about 2.0 % to about 4.0 %, (area percent of bendamustine) at release or
15 otherwise to achieve the pharmaceutical compositions described herein. The bendamustine product herein contains not more than about 0.5% to about 0.9%, preferably about 0.5%, (area percent of bendamustine) HP1 at release.

The present invention discloses a process for manufacturing a lyophilized preparation of bendamustine which comprises controlling for the concentration of
20 bendamustine degradants in the final product, such that, at release, the concentration of HP1 is less than 0.9%, preferably 0.5%, (area percent of bendamustine) and, at the time of product expiration, the concentration of bendamustine degradants is less than about 7.0%, preferably no more than about 5.0% to about 7.0%; wherein said product is stored at about 2°C to about 30°C.

25 Another embodiment of the invention is a bendamustine pre-lyophilization solution or dispersion comprising one or more organic solvents where the solution or dispersions include at least one stabilizing concentration of an organic solvent which reduces the level of degradation of bendamustine so that the amount of HP1 produced during lyophilization from about 0 to 24 hours does not exceed about 0.5% to about
30 0.9% (area percent of bendamustine) preferably 0.50%, preferably 0.45%, more preferably 0.40%, more preferably 0.35%, even more preferably 0.30%. An aspect of this embodiment is the lyophilized powder produced from the pre-lyophilization solution or dispersion.

Still another embodiment of the invention is a bendamustine pre-lyophilization solution or dispersion comprising one or more organic solvents where the solution or dispersions include at least one stabilizing concentration of an organic solvent which reduces the level of degradation of bendamustine so that the amount of bendamustine ethylester produced during lyophilization from about 0 to 24 hours does not exceed about 0.5% (area percent bendamustine). An aspect of this embodiment is the lyophilized powder produced from the pre-lyophilization solution or dispersion.

Still another embodiment of the invention is a bendamustine pre-lyophilization solution or dispersion comprising one or more organic solvents where the solution or dispersions include at least one stabilizing concentration of an organic solvent which reduces the level of degradation of bendamustine so that the amount of bendamustine ethylester (as shown in Formula IV) produced during lyophilization from about 0 to 24 hours is no more than 0.2%, preferably 0.1%, greater than the concentration of bendamustine ethylester as found in the drug substance used to make the pre-lyophilization solution. A preferred organic solvent is tertiary butanol.

The invention also discloses methods for preparing a bendamustine lyophilized preparation that includes dissolving bendamustine in a stabilizing concentration of an alcohol solvent of between about 5% to about 100% (v/v alcohol to form a pre-lyophilization solution; and lyophilizing the pre-lyophilization solution; wherein the bendamustine lyophilized preparation made from such methods contains not more than about 0.5% to about 0.9%, preferably 0.5%, (area percent of bendamustine) HP1 as shown in Formula II, wherein said HP1 is the amount of HP1 present at release or at time zero after reconstitution of the lyophilized pharmaceutical composition of bendamustine. Other alcohol concentrations include about 5% to about 99.9%, about 5% to about 70%, about 5% to about 60%, about 5% to about 50%, about 5% to about 40%, about 20% to about 35%. Preferred concentrations of alcohol are from about 20% to about 30%. Preferred alcohols include one or more of methanol, ethanol, propanol, iso-propanol, butanol, and tertiary-butanol. A more preferred alcohol is tertiary-butanol. A preferred concentration of tertiary-butanol is about 20% to about 30%, preferably about 30%. An aspect of this embodiment is the addition of an excipient before lyophilization. A preferred excipient is mannitol. Preferred pre-lyophilized concentrations of bendamustine are from about 2 mg/mL to about 50 mg/mL.

In a preferred method for preparing a bendamustine lyophilized preparation, lyophilizing the pre-lyophilization solution comprises i) freezing the pre-lyophilization solution to a temperature below about -40°C , preferably -50°C , to form a frozen solution; ii) holding the frozen solution at or below -40°C , preferably -50°C , for at least 2 hours; iii) ramping the frozen solution to a primary drying temperature between about -40°C and about -10°C to form a dried solution; iv) holding for about 10 to about 70 hours; v) ramping the dried solution to a secondary drying temperature between about 25°C and about 40°C ; and vii) holding for about 5 to about 40 hours to form a bendamustine lyophilized preparation. In a more preferred method lyophilizing the pre-lyophilization solution comprises i) freezing the pre-lyophilization solution to about -50°C to form a frozen solution; ii) holding the frozen solution at about -50°C for at least 2 hours to about 4 hours; iii) ramping to a primary drying temperature between about -20°C and about -12°C to form a dried solution; iv) holding at a primary drying temperature for about 10 to about 48 hours; v) ramping the dried solution to a secondary drying temperature between about 25°C and about 40°C ; and vi) holding at a secondary drying temperature for at least 5 hours up to about 20 hours. A preferred alcohol is tertiary-butanol. A preferred concentration of tertiary-butanol is about 20% to about 30%, preferably about 30%. An aspect of this embodiment is the addition of an excipient before lyophilization. A preferred excipient is mannitol. Preferred pre-lyophilized concentrations of bendamustine are from about 2 mg/mL to about 50 mg/mL.

Another embodiment of the invention is the lyophilized powder or preparation obtained from the methods of preparing a bendamustine lyophilized preparation disclosed herein.

The invention also involves bendamustine formulations for lyophilization that include an excipient and a stabilizing concentration of an organic solvent. A preferred formulation includes bendamustine at a concentration of about 15 mg/mL, mannitol at a concentration of about 25.5 mg/mL, tertiary-butyl alcohol at a concentration of about 30% (v/v) and water. Included in this embodiment of the invention are the lyophilized preparations made from such bendamustine formulations.

Included in the inventions are methods of treating a medical condition in a patient that involve administering a therapeutically effective amount of a pharmaceutical composition of the invention where the condition is amenable to

treatment with said pharmaceutical composition. Some conditions amenable to treatment with the compositions of the invention include chronic lymphocytic leukemia (CLL), Hodgkin's disease, non-Hodgkin's lymphoma (NHL), multiple myeloma (MM), breast cancer, small cell lung cancer, hyperproliferative disorders, and an autoimmune disease. Preferred conditions include NHL, CLL, breast cancer, and MM. Preferred autoimmune diseases include rheumatoid arthritis, multiple sclerosis or lupus.

Included in the inventions are the use of the pharmaceutical compositions or pharmaceutical preparations of the invention in the manufacture of a medicament for the treatment of a medical condition, as defined herein, in a patient that involve administering a therapeutically effective amount of a pharmaceutical composition of the invention where the condition is amenable to treatment with said pharmaceutical composition.

Also included in the invention are methods of treating in which the pharmaceutical compositions of the invention are in combination with one or more anti-neoplastic agents where the antineoplastic agent is given prior, concurrently, or subsequent to the administration of the pharmaceutical composition of the invention. Preferred antineoplastic agents are antibodies specific for CD20.

Another embodiment of the invention is a lyophilization cycle for producing lyophilized bendamustine preparations of the invention. A preferred lyophilization cycle includes a) freezing to about -50°C over about 8 hours; b) holding at -50°C for about 4 hours; c) ramping to -25°C over about 3 hours; d) holding at -10°C for 30 hours; e) ramping to between about 25°C and about 40°C or higher for about 3 hours; f) holding between about 25°C and about 40°C for about 25 hours; g) ramping to about 20°C in 1 hour; h) unloading at about 20°C, at a pressure of 13.5 psi in a pharmaceutically acceptable container that is hermetically sealed; wherein the pressure is about 150 microns throughout primary drying and 50 microns throughout secondary drying. An aspect of this cycle involves step (e) which is ramped to about 30-35°C for 3 hours and then ramped to 40°C for 5 hours. Another aspect of this embodiment is the lyophilized prepared prepared from such lyophilization cycles. A more preferred lyophilization cycle includes i) starting with a shelf temperature of about 5°C for loading; ii) freezing to about -50°C over about 8 hours; iii) holding at -50°C for about 4 hours; iv) ramping to about -20°C over about 3 hours; v) holding at about -20°C for 6

hours; ramping to about -15°C over about 1 hour; vi) holding at -15°C for about 20 hours; vii) ramping to about -15°C over about 1 hour; viii) holding at about -15°C for about 20 hours; ix) ramping to about -12°C over about 0.5 hours; x) holding at about -12°C for about 15.5 hours; xi) ramping to between about 25°C and about 40°C or higher for about 15 hours; xii) holding between about 25°C and about 40°C for about 10 hours; xiii) ramping to about 40°C over about 1 hour; and xiv) holding at about 40°C for about 5 hours; unloading at about 5°C, at a pressure of about 13.5 psi in a pharmaceutically acceptable container that is hermetically sealed; wherein the pressure is about 150 microns throughout primary drying and 50 microns throughout secondary drying. In a preferred embodiment step (xi) is ramped to about 30-35°C for about 15 hours.

The invention also encompasses a pharmaceutical dosage form of bendamustine containing not more than about 0.5% to about 0.9%, preferably 0.5%, HP1 (area percent of bendamustine) wherein said dosage form comprises a vial or other pharmaceutically acceptable container, wherein said HP1 is the amount of HP1 present pre-reconstitution or at time zero after reconstitution of said dosage form. Preferred concentrations of bendamustine include about 10 to about 500 mg/container, about 100 mg/container, about 5 mg to about 2 g/container and about 170 mg/container.

The present invention also includes pre-lyophilized pharmaceutical compositions of bendamustine. A preferred pre-lyophilized composition includes bendamustine HCl about 15 mg/mL, mannitol about 25.5 mg/mL, about 30% (v/v) tertiary-butyl alcohol, and water.

These and other embodiments of the invention are described hereinbelow or are evident to persons of ordinary skill in the art based on the following disclosures.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the solubility of bendamustine at various temperatures for two different solutions of bendamustine in tertiary butanol.

Fig. 2 shows the purity results of an HPLC analysis after incubating bendamustine in various alcohols for 24 hours at 5°C. Results are presented as the area percent of the bendamustine peak.

Fig. 3 shows HP1 (Formula II) formation after 24 hours in various alcohol/water co-solvents at 5°C

Fig 4 shows dimer (Formula III) formation after 24 hours in various alcohol/water co-solvents at 5°C

5 Fig. 5- shows a lyophilization cycle for bendamustine using a TBA/water co-solvent.

Fig. 6 shows a chromatogram for Ribomustin® using HPLC method No. 1.

DETAILED DESCRIPTION OF THE INVENTION

10 As used herein, the terms “formulate” refers to the preparation of a drug, e.g., bendamustine, in a form suitable for administration to a mammalian patient, preferably a human. Thus, “formulation” can include the addition of pharmaceutically acceptable excipients, diluents, or carriers.

As used herein, the term “lyophilized powder” or “lyophilized preparation”
15 refers to any solid material obtained by lyophilization, i.e., freeze-drying of an aqueous solution. The aqueous solution may contain a non-aqueous solvent, i.e. a solution composed of aqueous and one or more non-aqueous solvent(s). Preferably, a lyophilized preparation is one in which the solid material is obtained by freeze-drying a solution composed of aqueous and one or more non-aqueous solvents, more
20 preferably the non-aqueous solvent is an alcohol.

By “stable pharmaceutical composition” is meant any pharmaceutical composition having sufficient stability to have utility as a pharmaceutical product. Preferably, a stable pharmaceutical composition has sufficient stability to allow storage at a convenient temperature, preferably between -20°C and 40°C, more
25 preferably about 2°C to about 30°C, for a reasonable period of time, e.g., the shelf-life of the product which can be as short as one month but is typically six months or longer, more preferably one year or longer even more preferably twenty-four months or longer, and even more preferably thirty-six months or longer. The shelf-life or expiration can be that amount of time where the active ingredient degrades to a point
30 below 90% purity. For purposes of the present invention stable pharmaceutical composition includes reference to pharmaceutical compositions with specific ranges of impurities as described herein. Preferably, a stable pharmaceutical composition is one which has minimal degradation of the active ingredient, e.g., it retains at least about 85

% of un-degraded active, preferably at least about 90 %, and more preferably at least about 95%, after storage at 2-30°C for a 2-3 year period of time.

By “stable lyophilized preparation” is meant any lyophilized preparation having sufficient stability, such characteristics as similarly defined herein for a stable pharmaceutical composition, to have utility as a pharmaceutical product

By “degraded” is meant that the active has undergone a change in chemical structure.

The term "therapeutically effective amount" as used herein refers to that amount of the compound being administered that will relieve to some extent one or more of the symptoms of the disorder being treated. In reference to the treatment of neoplasms, a therapeutically effective amount refers to that amount which has the effect of (1) reducing the size of the tumor, (2) inhibiting (that is, slowing to some extent, preferably stopping) tumor metastasis, (3) inhibiting to some extent (that is, slowing to some extent, preferably stopping) tumor growth, and/or, (4) relieving to some extent (or, preferably, eliminating) one or more symptoms associated with the cancer. Therapeutically effective amount can also mean preventing the disease from occurring in an animal that may be predisposed to the disease but does not yet experience or exhibit symptoms of the disease (prophylactic treatment). Further, therapeutically effective amount can be that amount that increases the life expectancy of a patient afflicted with a terminal disorder. Typical therapeutically effective doses for bendamustine for the treatment of non-Hodgkin's lymphoma can be from about 60-120 mg/m² given as a single dose on two consecutive days. The cycle can be repeated about every three to four weeks. For the treatment of chronic lymphocytic leukemia (CLL) bendamustine can be given at about 80-100 mg/m² on days 1 and 2. The cycle can be repeated after about 4 weeks. For the treatment of Hodgkin's disease (stages II-IV), bendamustine can be given in the “DBVBe regimen” with daunorubicin 25 mg/m² on days 1 and 15, bleomycin 10 mg/m² on days 1 and 15, vincristine 1.4 mg/m² on days 1 and 15, and bendamustine 50 mg/m² on days 1-5 with repetition of the cycle about every 4 weeks. For breast cancer, bendamustine (120 mg/m²) on days 1 and 8 can be given in combination with methotrexate 40 mg/m² on days 1 and 8, and 5-fluorouracil 600 mg/m² on days 1 and 8 with repetition of the cycle about every 4 weeks. As a second-line of therapy for breast cancer, bendamustine can be given at

about 100-150 mg/m² on days 1 and 2 with repetition of the cycle about every 4 weeks.

As used herein "neoplastic" refers to a neoplasm, which is an abnormal growth, such growth occurring because of a proliferation of cells not subject to the usual limitations of growth. As used herein, "anti-neoplastic agent" is any compound,
5 composition, admixture, co-mixture, or blend which inhibits, eliminates, retards, or reverses the neoplastic phenotype of a cell.

As used herein "hyperproliferation" is the overproduction of cells in response to a particular growth factor. "Hyperproliferative disorders" are diseases in which the
10 cells overproduce in response to a particular growth factor. Examples of such "hyperproliferative disorders" include diabetic retinopathy, psoriasis, endometriosis, cancer, macular degenerative disorders and benign growth disorders such as prostate enlargement.

As used herein, the term "vial" refers to any walled container, whether rigid or
15 flexible.

"Controlling" as used herein means putting process controls in place to facilitate achievement of the thing being controlled. For example, in a given case, "controlling" can mean testing samples of each lot or a number of lots regularly or randomly; setting the concentration of degradants as a release specification; selecting
20 process conditions, e.g., use of alcohols and/or other organic solvents in the pre-lyophilization solution or dispersion, so as to assure that the concentration of degradants of the active ingredient is not unacceptably high; etc. Controlling for degradants by setting release specifications for the amount of degradants can be used to facilitate regulatory approval of a pharmaceutical product by a regulatory agency,
25 such as the U.S. Food and Drug Administration and similar agencies in other countries or regions ("agency").

The term "pharmaceutically acceptable" as used herein means that the thing that is pharmaceutically acceptable, e.g., components, including containers, of a pharmaceutical composition, does not cause unacceptable loss of pharmacological
30 activity or unacceptable adverse side effects. Examples of pharmaceutically acceptable components are provided in The United States Pharmacopeia (USP), The National Formulary (NF), adopted at the United States Pharmacopeial Convention, held in Rockville, Md. in 1990 and FDA Inactive Ingredient Guide 1990, 1996 issued

by the U.S. Food and Drug Administration (both are hereby incorporated by reference herein, including any drawings). Other grades of solutions or components that meet necessary limits and/or specifications that are outside of the USP/NF may also be used.

The term "pharmaceutical composition" as used herein shall mean a
5 composition that is made under conditions such that it is suitable for administration to humans, e.g., it is made under GMP conditions and contains pharmaceutically acceptable excipients, e.g., without limitation, stabilizers, bulking agents, buffers, carriers, diluents, vehicles, solubilizers, and binders. As used herein pharmaceutical composition includes but is not limited to a pre-lyophilization solution or dispersion as
10 well as a liquid form ready for injection or infusion after reconstitution of a lyophilized preparation.

A "pharmaceutical dosage form" as used herein means the pharmaceutical compositions disclosed herein being in a container and in an amount suitable for reconstitution and administration of one or more doses, typically about 1-2, 1-3, 1-4, 1-
15 5, 1-6, 1-10, or about 1-20 doses. Preferably, a "pharmaceutical dosage form" as used herein means a lyophilized pharmaceutical composition disclosed herein in a container and in an amount suitable for reconstitution and delivery of one or more doses, typically about 1-2, 1-3, 1-4, 1-5, 1-6, 1-10, or about 1-20 doses. The pharmaceutical dosage form can comprise a vial or syringe or other suitable pharmaceutically
20 acceptable container. The pharmaceutical dosage form suitable for injection or infusion use can include sterile aqueous solutions or dispersions or sterile powders comprising an active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and
25 storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol such as glycerol, propylene glycol, or liquid polyethylene glycols and the like, vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The prevention of the growth of microorganisms can be accomplished by various antibacterial and antifungal agents, for example, parabens,
30 chlorobutanol, phenol, sorbic acid, thimerosal, and the like.

As used herein, the term "excipient" means the substances used to formulate active pharmaceutical ingredients (API) into pharmaceutical formulations; in a preferred embodiment, an excipient does not lower or interfere with the primary

therapeutic effect of the API. Preferably, an excipient is therapeutically inert. The term "excipient" encompasses carriers, diluents, vehicles, solubilizers, stabilizers, bulking agents, and binders. Excipients can also be those substances present in a pharmaceutical formulation as an indirect or unintended result of the manufacturing process. Preferably, excipients are approved for or considered to be safe for human and animal administration, i.e., GRAS substances (generally regarded as safe). GRAS substances are listed by the Food and Drug administration in the Code of Federal Regulations (CFR) at 21 CFR § 182 and 21 CFR § 184, incorporated herein by reference. Preferred excipients include, but are not limited to, hexitols, including mannitol and the like.

As used herein "a stabilizing concentration of an organic solvent" or "a stabilizing concentration of an alcohol" means that amount of an organic solvent or alcohol that reduces the level of degradation of bendamustine to achieve a specified level of degradants in the final drug product. For example, with respect to the degradant HP1, a stabilizing concentration of an organic solvent is that amount which results in an HP1 concentration (area percent of bendamustine) of less than about 0.5%, preferably less than 0.45 %, preferably less than 0.40 %, more preferably less than 0.35%, more preferably less than 0.30%, and even more preferably less than 0.25%. With respect to the overall or total degradant concentration of the final drug product, a stabilizing concentration of an organic solvent is that amount that results in a total degradant concentration (at the time of drug product release) of less than about 7% (area percent bendamustine), preferably less than about 6%, more preferably less than about 5%, and even more preferably less than about 4.0%. By "area percent of bendamustine" is meant the amount of a specified degradant, e.g., HP1, relative to the amount of bendamustine as determined, e.g., by HPLC.

The term "organic solvent" means an organic material, usually a liquid, capable of dissolving other substances.

As used herein, "trace amount of an organic solvent" means an amount of solvent that is equal to or below recommended levels for pharmaceutical products, for example, as recommended by ICH guidelines (International Conferences on Harmonization, Impurities-- Guidelines for Residual Solvents. Q3C. Federal Register. 1997;62(247):67377). The lower limit is the lowest amount that can be detected.

The term “release” or “at release” means the drug product has met the release specifications and can be used for its intended pharmaceutical purpose.

A. General

The invention provides stable, pharmaceutically acceptable compositions prepared from bendamustine. In particular, the invention provides formulations for the lyophilization of bendamustine HCl. The lyophilized powder obtained from such formulations is more easily reconstituted than the presently available lyophilized powder of bendamustine. Further, the lyophilized products of the present invention have a better impurity profile than Ribomustin® with respect to certain impurities, in particular HP1, bendamustine dimer, and bendamustine ethylester, prior to reconstitution, upon storage of the lyophilate, or following reconstitution and admixture.

The present invention further provides formulations of bendamustine useful for treating neoplastic diseases. The formulations described herein can be administered alone or in combination with at least one additional anti-neoplastic agent and/or radioactive therapy.

An aspect of the invention is conditions and means for enhancing the stability of bendamustine prior to and during the lyophilization process, upon shelf storage or upon reconstitution.

Anti-neoplastic agents which may be utilized in combination with the formulations of the invention include those provided in the Merck Index 11, pp 16-17, Merck & Co., Inc. (1989) and The Chemotherapy Source Book (1997). Both books are widely recognized and readily available to the skilled artisan.

There are large numbers of antineoplastic agents available in commercial use, in clinical evaluation and in pre-clinical development, which could be selected for treatment of neoplasia by combination drug chemotherapy. Such antineoplastic agents fall into several major categories, namely, antibiotic-type agents, covalent DNA-binding drugs, antimetabolite agents, hormonal agents, including glucocorticoids such as prednisone and dexamethasone, immunological agents, interferon-type agents, differentiating agents such as the retinoids, pro-apoptotic agents, and a category of miscellaneous agents, including compounds such as antisense, small interfering RNA, and the like. Alternatively, other anti-neoplastic agents, such as metallomatrix proteases (MMP) inhibitors, SOD mimics or $\alpha_v\beta_3$ inhibitors may be used.

One family of antineoplastic agents which may be used in combination with the compounds of the inventions consists of antimetabolite-type antineoplastic agents. Suitable antimetabolite antineoplastic agents may be selected from the group consisting of alanosine, AG2037 (Pfizer), 5-FU-fibrinogen, acanthifolic acid, aminothiadiazole, brequinar sodium, carmofur, Ciba-Geigy CGP-30694, cyclopentyl cytosine, cytarabine phosphate stearate, cytarabine conjugates, Lilly DATHF, Merrel Dow DDFC, dezaguanine, dideoxycytidine, dideoxyguanosine, didox, Yoshitomi DMDC, doxifluridine, Wellcome EHNA, Merck & Co. EX-015, fazarabine, floxuridine, fludarabine phosphate, 5-fluorouracil, N-(2'-furanidyl)-5-fluorouracil, Daiichi Seiyaku FO-152, isopropyl pyrrolizine, Lilly LY-188011, Lilly LY-264618, methobenzaprim, methotrexate, Wellcome MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI NSC-39661, NCI NSC-612567, Warner-Lambert PALA, pentostatin, piritrexim, plicamycin, Asahi Chemical PL-AC, Takeda TAC-788, thioguanine, tiazofurin, Erbamont TIF, trimetrexate, tyrosine kinase inhibitors, tyrosine protein kinase inhibitors, Taiho UFT and uricytin.

A second family of antineoplastic agents which may be used in combination with the compounds of the invention consists of covalent DNA-binding agents. Suitable alkylating-type antineoplastic agents may be selected from the group consisting of Shionogi 254-S, aldo-phosphamide analogues, altretamine, anaxirone, Boehringer Mannheim BBR-2207, bestrabucil, budotitane, Wakunaga CA-102, carboplatin, carmustine, Chinoin-139, Chinoin-153, chlorambucil, cisplatin, cyclophosphamide, American Cyanamid CL-286558, Sanofi CY-233, cyplatate, Degussa D-19-384, Sumimoto DACHP(Myrr)2, diphenylspiromustine, diplatinum cytostatic, Erba distamycin derivatives, Chugai DWA-2114R, ITI E09, elmustine, Erbamont FCE-24517, estramustine phosphate sodium, fotemustine, Unimed G-6-M, Chinoin GYKI-17230, hepsul-fam, ifosfamide, iproplatin, lomustine, mafosfamide, melphalan, mitolactol, Nippon Kayaku NK-121, NCI NSC-264395, NCI NSC-342215, oxaliplatin, Upjohn PCNU, prednimustine, Proter PTT-119, ranimustine, semustine, SmithKline SK&F-101772, Yakult Honsha SN-22, spiromustine, Tanabe Seiyaku TA-077, tauromustine, temozolomide, teroxirone, tetraplatin and trimelamol.

Another family of antineoplastic agents which may be used in combination with the compounds disclosed herein consists of antibiotic-type antineoplastic agents. Suitable antibiotic-type antineoplastic agents may be selected from the group

consisting of Taiho 4181-A, aclarubicin, actinomycin D, actinoplanone, alanosine, Erbamont ADR-456, aeroplysinin derivative, Ajinomoto AN-201-II, Ajinomoto AN-3, Nippon Soda anisomycins, anthracycline, azino-mycin-A, bisucaberin, Bristol-Myers BL-6859, Bristol-Myers BMY-25067, Bristol-Myers BMY-25551, Bristol-Myers 5 BMY-26605, Bristol-Myers BMY-27557, Bristol-Myers BMY-28438, bleomycin sulfate, bryostatin-1, Taiho C-1027, caliche mycin, chromoximycin, dactinomycin, daunorubicin, Kyowa Hakko DC-102, Kyowa Hakko DC-79, Kyowa Hakko DC-88A, Kyowa Hakko DC89-A1, Kyowa Hakko DC92-B, ditrisarubicin B, Shionogi DOB-41, doxorubicin, doxorubicin-fibrinogen, elsamicin-A, epirubicin, erbstatin, esorubicin, 10 esperamicin-A1, esperamicin-Alb, Erbamont FCE-21954, Fujisawa FK-973, fostriecin, Fujisawa FR-900482, glidobactin, gregatin-A, grincamycin, herbimycin, idarubicin, illudins, kazusamycin, kesarirhodins, Kyowa Hakko KM-5539, Kirin Brewery KRN-8602, Kyowa Hakko KT-5432, Kyowa Hakko KT-5594, Kyowa Hakko KT-6149, American Cyanamid LL-D49194, Meiji Seika ME 2303, menogaril, mitomycin, 15 mitoxantrone, SmithKline M-TAG, neoenactin, Nippon Kayaku NK-313, Nippon Kayaku NKT-01, SRI International NSC-357704, oxalysine, oxaunomycin, peplomycin, pilatin, pirarubicin, porothramycin, pyrindamycin A, Tobishi RA-I, rapamycin, rhizoxin, rodorubicin, sibanomicin, siwenmycin, Sumitomo SM-5887, Snow Brand SN-706, Snow Brand SN-07, sorangicin-A, sparsomycin, SS 20 Pharmaceutical SS-21020, SS Pharmaceutical SS-7313B, SS Pharmaceutical SS-9816B, steffimycin B, Taiho 4181-2, talisomycin, Takeda TAN-868A, terpentecin, thiazine, tricrozarin A, Upjohn U-73975, Kyowa Hakko UCN-10028A, Fujisawa WF-3405, Yoshitomi Y-25024 and zorubicin.

A fourth family of antineoplastic agents which may be used in combination 25 with the compounds of the invention include a miscellaneous family of antineoplastic agents selected from the group consisting of alpha-carotene, alpha-difluoromethyl-arginine, acitretin, arsenic trioxide, Avastin® (bevacizumab), Biotec AD-5, Kyorin AHC-52, alstonine, amonafide, amphetinile, amsacrine, Angiostat, ankinomycin, anti-neoplaston A10, antineoplaston A2, antineoplaston A3, antineoplaston A5, 30 antineoplaston AS2-1, Henkel APD, aphidicolin glycinate, asparaginase, Avarol, baccharin, batracylin, benfluron, benzotript, Ipsen-Beaufour BIM-23015, bisantrene, Bristo-Myers BMY-40481, Vestar boron-10, bromofosfamide, Wellcome BW-502, Wellcome BW-773, caracemide, carmethizole hydrochloride, Ajinomoto CDAF,

chlorsulfaquinoxalone, Chemes CHX-2053, Chemex CHX-100, Warner-Lambert CI-921, Warner-Lambert CI-937, Warner-Lambert CI-941, Warner-Lambert CI-958, clanfenur, claviridenone, ICN compound 1259, ICN compound 4711, Contracan, Yakult Honsha CPT-11, crisnatol, curaderm, cytochalasin B, cytarabine, cytocytin,

5 Merz D-609, DABIS maleate, dacarbazine, datelliptinium, didemnin- B, dihaematoporphyrin ether, dihydrolenperone, dinaline, distamycin, Toyo Pharmar DM-341, Toyo Pharmar DM-75, Daiichi Seiyaku DN-9693, elliprabin, elliptinium acetate, epothionesTsumura EPMTC, erbitux, ergotamine, erlotnib, etoposide,

10 etretinate, fenretinide, Fujisawa FR-57704, gallium nitrate, genkwadaphnin, Gleevec® (imatnib), Chugai GLA-43, Glaxo GR-63178, gefitinib, grifolan NMF-5N, hexadecylphosphocholine, Green Cross HO-221, homoharringtonine, hydroxyurea, BTG ICRF-187, indanocine, ilmofosine, isoglutamine, isotretinoin, Otsuka JI-36, Ramot K-477, Otsuak K-76COONa, Kureha Chemical K-AM, MECT Corp KI-8110, American Cyanamid L-623, leukoregulin, lonidamine, Lundbeck LU-23-112, Lilly

15 LY-186641, NCI (US) MAP, marycin, mefloquine, Merrel Dow MDL-27048, Medco MEDR-340, merbarone, merocyanine derivatives, methylanilinoacridine, Molecular Genetics MGI-136, minactivin, mitonafide, mitoquidone, mopidamol, motretinide, Zenyaku Kogyo MST-16, N-(retinoyl)amino acids, Nisshin Flour Milling N-021, N-acylated-dehydroalanines, nafazatrom, Taisho NCU-190, nocodazole derivative,

20 Normosang, NCI NSC-145813, NCI NSC-361456, NCI NSC-604782, NCI NSC-95580, octreotide, Ono ONO-112, oquizanocine, Akzo Org-10172, paclitaxel, pancratistatin, pazelliptine, Warner-Lambert PD-111707, Warner-Lambert PD-115934, Warner-Lambert PD-131141, Pierre Fabre PE-1001, ICRT peptide D, piroxantrone, polyhaematoporphyrin, polypreic acid, Efamol porphyrin, probimane,

25 procarbazine, proglumide, Invitron protease nexin I, Tobishi RA-700, razoxane, Sapporo Breweries RBS, restrictin-P, retelliptine, retinoic acid, Rhone-Poulenc RP-49532, Rhone-Poulenc RP-56976, Rituxan® (and other anti CD20 antibodies, e.g. Bexxar®, Zevalin®), SmithKline SK&F-104864, statins (Lipitor® etc.), Sumitomo SM-108, Kuraray SMANCS, SeaPharm SP-10094, spatol, spirocyclopropane

30 derivatives, spirogermanium, Unimed, SS Pharmaceutical SS-554, strypoldinone, Stypoldione, Suntory SUN 0237, Suntory SUN 2071, superoxide dismutase, Thalidomide, Thalidomide analogs, Toyama T-506, Toyama T-680, taxol, Teijin TEI-0303, teniposide, thaliblastine, Eastman Kodak TJB-29, tocotrienol, Topostin, Teijin

Examples of radioprotective agents which may be used in the combination chemotherapy of this invention are AD-5, adchnon, amifostine analogues, detox, dimesna, 1-102, MM-159, N-acylated-dehydroalanines, TGF-Genentech, tiprotimod, amifostine, WR-151327, FUT-187, ketoprofen transdermal, nabumetone, superoxide dismutase (Chiron and Enzon).

Preferred anti-neoplastic agents include, without limitation, one or more of
15 daunorubicin, bleomycin, vincristine, doxorubicin, dacarbazine, prednisolone,
mitoxantrone, prednisone, methotrexate, 5-fluorouracil, dexamethasone, thalidomide,
thalidomide derivatives, 2ME2, Neovastat, R 11 5777, arsenic trioxide, bortezomib,
tamoxifen, G3139 (antisense), and SU5416, mitomycin, anti-CD20 antibodies, such as
Rituxan® and R-etodolac.

Preferred drug regimens for which the present formulation may be used in conjunction with or as a replacement for one or more of the components includes, without limitation, ABVD (doxorubicin, bleomycin, vincristine, dacarbazine), DBV (daunorubicin, belomycin, vincristine), CVPP (cyclophosphamide, vinblastine, procarbazine, prednisolone), COP (cyclophosphamide, vincristine, prednisolone), CHOP (cyclophosphamide, doxorubicin,

Table A- Cancer Therapeutic Regimens

Abbreviation	Drugs Used	Disease
AC	Doxorubicin & Cyclophosphamide	Breast cancer
CFM (CF, FNC)	Cyclophosphamide,	Breast cancer

	Fluorouracil, Mitoxantrone	
CMF	Cyclophosphamide, Methotrexate, Fluorouracil	Breast cancer
NFL	Mitoxantrone, Fluorouracil, Leucovorin	Breast cancer
Sequential Dox-CMF	Doxorubicin	Breast cancer
VATH	Vinblastine, Doxorubicin, Thiotepa, Fluoxymesterone	Breast cancer
EMA-86	Etoposide, Mitoxantrone, Ctyarabine	AML (induction)
7 + 3	Cytarabine WITH Daunorubicin OR Idarubicin OR Mitoxantrone	AML (induction)
5 + 2	Cytarabine WITH Daunorubicin OR Mitoxantrone	AML (induction)
HiDAC	Cytarabine	AML (post- remission)
ABVD	Doxorubicin, Bleomycin, Vinblastine, Dacarbazine	Hodgkin's
ChlVPP	Chlorambucil, Vinblastine, Procarbazine, Prednisone	Hodgkin's
EVA	Etoposide, Vinblastine, Doxorubicin	Hodgkin's
MOPP	Mechlorethamine, Vincristine, Procarbazine, Prednisone	Hodgkin's
MOPP/ABV Hybrid	Mechlorethamine, Vincristine, Procarbazine, Prednisone, Doxorubicin, Bleomycin, Vinblastine	Hodgkin's
MOPP/ABVD	Mechlorethamine,	Hodgkin's

	Doxorubicin, Vinblastine, Bleomycin, Etoposide, Prednisone	
CNOP	Cyclophosphamide, Mitoxantrone, Vincristine, Prednisone	Non-Hodgkin's
COMLA	Cyclophosphamide, Vincristine, Methotrexate, Leucovorin, Cytarabine	Non-Hodgkin's
DHAP	Dexamethasone, Cisplatin, Cytarabine	Non-Hodgkin's
ESHAP	Etoposide, Methylprednisilone, Cisplatin, Cytarabine	Non-Hodgkin's
MACOP-B	Methotrexate, Leucovorin, Doxorubicin, Cyclophosphamide, Vincristine, Prednisone, Bleomycin, Septra, Ketoconazole	Non-Hodgkin's
m-BACOD	Methotrexate, Leucovorin, Bleomycin, Doxorubicin, Cyclophosphamide, Vincristine, Dexamethasone	Non-Hodgkin's
MINE-ESHAP	Mesna, Ifosfamide, Mitoxantrone, Etoposide	Non-Hodgkin's
NOVP	Mitoxantrone, Vinblastine, Prednisone, Vincristine	Non-Hodgkin's
ProMACE/cytaBOM	Prednisone, Doxorubicin,	Non-Hodgkin's

	Cyclophosphamide, Etoposide, Cytarabine, Bleomycin, Vincristine, Methotrexate, Leucovorin, Septra	
M2	Vincristine, Carmustine, Cyclophosphamide, Melphalan, Prednisone	Multiple Myeloma
MP	Melphalan, Prednisone	Multiple Myeloma
VAD	Vincristine, Doxorubicin, Dexamethasone	Multiple Myeloma
VBMCP	Vincristine, Carmustine, Melphalan, Cyclophosphamide, Prednisone	Multiple Myeloma

As described herein, a lyophilized formulation of bendamustine is achieved following removal of an organic solvent in water. The most typical example of the solvent used to prepare this formulation is tertiary butanol (TBA). Other organic solvents can be used including ethanol, n-propanol, n-butanol, isopropanol, ethyl acetate, dimethyl carbonate, acetonitrile, dichloromethane, methyl ethyl ketone, methyl isobutyl ketone, acetone, 1-pentanol, methyl acetate, methanol, carbon tetrachloride, dimethyl sulfoxide, hexafluoroacetone, chlorobutanol, dimethyl sulfone, acetic acid, cyclohexane. These preceding solvents may be used individually or in combination.

Useful solvents must form stable solutions with bendamustine and must not appreciably degrade or deactivate the API. The solubility of bendamustine in the selected solvent must be high enough to form commercially useful concentrations of the drug in solvent. Additionally, the solvent should be capable of being removed easily from an aqueous dispersion or solution of the drug product, e.g., through lyophilization or vacuum drying. Preferably, a solution having a concentration of

about 2-80 mg/mL, preferably about 5 to 40 mg/mL, more preferably 5-20 mg/mL and even more preferably 12 to 17 mg/mL bendamustine is used.

A pharmaceutically acceptable lyophilization excipient can be dissolved in the aqueous phase. Examples of excipients useful for the present invention include, without limitation, sodium or potassium phosphate, citric acid, tartaric acid, gelatin, glycine, and carbohydrates such as lactose, sucrose, maltose, glycerin, dextrose, dextran, trehalose and hetastarch. Mannitol is a preferred excipient. Other excipients that may be used if desired include antioxidants, such as, without limitation, ascorbic acid, acetylcysteine, cysteine, sodium hydrogen sulfite, butyl-hydroxyanisole, butyl-hydroxytoluene or alpha-tocopherol acetate, or chelators.

A typical formulation and lyophilization cycle useful in accordance with the present invention is provided below. Lyophilization can be carried out using standard equipment as used for lyophilization or vacuum drying. The cycle may be varied depending upon the equipment and facilities used for the fill/finish.

In accordance with a typical embodiment of the present invention, an aqueous pre-lyophilization solution or dispersion is first formulated in a pharmaceutically acceptable compounding vessel. The solution is aseptically filtered into a sterile container, filled into an appropriate sized vial, partially stoppered and loaded into the lyophilizer. Using lyophilization techniques described herein the solution is lyophilized until a moisture content in the range of about 0.1 to about 8.0 percent is achieved. The resulting lyophilization powder is stable as a lyophilized powder for about six months to greater than about 2 years, preferably greater than about 3 years at about 5°C to about 25° C and can be readily reconstituted with Sterile Water for Injection, or other suitable carrier, to provide liquid formulations of bendamustine, suitable for internal administration e.g., by parenteral injection. For intravenous administration, the reconstituted liquid formulation, i.e., the pharmaceutical composition, is preferably a solution.

The pre-lyophilization solution or dispersion normally is first formulated in a pharmaceutically acceptable container by: 1) adding an excipient, such as mannitol (about 0 to about 50 mg/mL) with mixing to water (about 65% of the total volume) at ambient temperature, 2) adding an organic solvent (0.5- 99.9% v/v), such as TBA to the aqueous solution with mixing at about 20°-35°C, 4) adding bendamustine HCl to the desired concentration with mixing, 5) adding water to achieve the final volume,

and 6) cooling the solution to about 1°C to about 30°C, preferably about 5°C.

Although the preceding steps are shown in a certain order, it is understood that one skilled in the art can change the order of the steps and quantities as needed. Quantities can be prepared on a weight basis also.

5 The pre-lyophilization solution or dispersion can be sterilized prior to lyophilization, sterilization is generally performed by aseptic filtration, e.g., through a 0.22 micron or less filter. Multiple sterilization filters can be used. Sterilization of the solution or dispersion can be achieved by other methods known in the art, e.g., radiation.

10 In this case, after sterilization, the solution or dispersion is ready for lyophilization. Generally, the filtered solution will be introduced into a sterile receiving vessel, and then transferred to any suitable container or containers in which the formulation may be effectively lyophilized. Usually the formulation is effectively and efficiently lyophilized in the containers in which the product is to be marketed,
15 such as, without limitation, a vial, as described herein and as known in the art.

 A typical procedure for use in lyophilizing the pre-lyophilization solutions or dispersions is set forth below. However, a person skilled in the art would understand that modifications to the procedure or process may be made depending on such things as, but not limited to, the pre-lyophilization solution or dispersion and lyophilization
20 equipment.

 Initially, the product is placed in a lyophilization chamber under a range of temperatures and then subjected to temperatures well below the product's freezing point, generally for several hours. Preferably, the temperature will be at or below about -40°C for at least 2 hours. After freezing is complete, the chamber and the
25 condenser are evacuated through vacuum pumps, the condenser surface having been previously chilled by circulating refrigerant. Preferably, the condenser will have been chilled below the freezing point of the solution preferably to about -40°, more preferably to about -50°C or lower, even more preferably to about -60°C or lower. Additionally, evacuation of the chamber should continue until a pressure of about 10
30 to about 600 microns, preferably about 50 to about 150 microns is obtained.

 The product composition is then warmed under vacuum in the chamber and condenser. This usually will be carried out by warming the shelves within the lyophilizer on which the product rests during the lyophilization process at a pressure

ranging from about 10 to about 600 microns. The warming process will optimally take place very gradually, over the course of several hours. For example, the product temperature should initially be increased from about -30°C to about -10°C and maintained for about 10-70 hours. Additionally, the product temperature can be
5 increased from the freezing temperature to about 25°C-40°C over a period of 30-192 hours. To prevent powder ejection of the lyophilate from vials, complete removal of the organic solvent and water should be done during the initial drying phase. Complete drying can be confirmed by stabilization of vacuum, condenser temperature and product shelf temperature. After the initial drying, the product temperature should
10 be increased to about 25°C-40°C and maintained for about 5-40 hours.

Once the drying cycle is completed, the pressure in the chamber can be slowly released to atmospheric pressure (or slightly below) with sterile, dry-nitrogen gas (or equivalent gas). If the product composition has been lyophilized in containers such as vials, the vials can be stoppered, removed and sealed. Several representative samples
15 can be removed for purposes of performing various physical, chemical, and microbiological tests to analyze the quality of the product.

The lyophilized bendamustine formulation is typically marketed in pharmaceutical dosage form. The pharmaceutical dosage form of the present invention, although typically in the form of a vial, may be any suitable container, such
20 as ampoules, syringes, co-vials, which are capable of maintaining a sterile environment. Such containers can be glass or plastic, provided that the material does not interact with the bendamustine formulation. The closure is typically a stopper, most typically a sterile rubber stopper, preferably a bromobutyl rubber stopper, which affords a hermetic seal.

25 After lyophilization, the bendamustine lyophilization powder may be filled into containers, such as vials, or alternatively the pre-lyophilization solution can be filled into such vials and lyophilized therein, resulting in vials which directly contain the lyophilized bendamustine formulation. Such vials are, after filling or lyophilization of the solution therein, sealed, as with a stopper, to provide a sealed, sterile,
30 pharmaceutical dosage form. Typically, a vial will contain a lyophilized powder including about 10-500 mg/vial, preferably about 100 mg/vial, bendamustine and about 5mg-2g/vial, preferably about 170 mg/vial, mannitol.

The lyophilized formulations of the present invention may be reconstituted with water, preferably Sterile Water for Injection, or other sterile fluid such as co-solvents, to provide an appropriate solution of bendamustine for administration, as through parenteral injection following further dilution into an appropriate intravenous admixture container, for example, normal saline.

B. Solubility

The solubility of bendamustine HCl (bendamustine) in water (alone) and with varying amounts of alcohols commonly used in lyophilization, e.g., methanol, ethanol, propanol, isopropanol, butanol and tertiary-butyl alcohol (TBA) was determined by visual inspection. Amounts of bendamustine at 15 mg/mL, combined with mannitol at 25.5 mg/mL were prepared in 10 mL of the indicated alcohol solutions at room temperature (see Table 1). Samples were then refrigerated at 5°C and inspected after 0, 3, 6 and 24 hours for particulates and/or precipitates.

The results shown in Table 1 indicate that bendamustine solubility is dependant on temperature and the amount of alcohol in aqueous solutions. For the alcohols tested, the solubility of bendamustine increased as the concentration of alcohol increased. The formation of a precipitant was also dependent on the temperature and time. Bendamustine did not precipitate immediately with any alcohol, but crystallized after storage at 5°C. Alcohols varied in their effect on solubility. Without wishing to be bound to any particular theory, smaller alcohols such as methanol and ethanol have less of an effect on solubility as compared with larger alcohols (tertiary-butanol and n-butanol). However, the shape of the alcohol is also important. For example n-propanol was found to be better than iso-propanol in preventing precipitation in this system. The two alcohols with the greatest effect on solubility were n-propanol and tertiary-butanol.

Table 1. Bendamustine solubility over a 24 hour period in various alcohols when stored at 5°C.

	Zero Time	3 Hours	6 Hours	24 Hours
Methanol (v/v)				
0% (Water Only)	CCS	CCS	Precipitate	Precipitate
5%	CCS	CCS	Precipitate	Precipitate
10%	CCS	CCS	CCS	Precipitate
20%	CCS	CCS	CCS	Precipitate
30%	CCS	CCS	CCS	CCS
Ethanol (v/v)				
1.9%	CCS	CCS	Precipitate	Precipitate
5%	CCS	CCS	Precipitate	Precipitate
10%	CCS	CCS	CCS	Precipitate

20%	CCS	CCS	CCS	CCS
30%	CCS	CCS	CCS	CCS
n-Propanol (v/v)				
5%	CCS	CCS	CCS	Precipitate
10%	CCS	CCS	CCS	CCS
20%	CCS	CCS	CCS	CCS
30%	CCS	CCS	CCS	CCS
Iso-propanol (v/v)				
5%	CCS	Precipitate	Precipitate	Precipitate
10%	CCS	CCS	CCS	CCS
20%	CCS	CCS	CCS	CCS
30%	CCS	CCS	CCS	CCS
n-Butanol (v/v)				
5%	CCS	CCS	CCS	CCS
10%	CCS	CCS	CCS	CCS
20%	2 layers	2 layers	2 layers	2 layers
30%	2 layers	2 layers	2 layers	2 layers
Tert-Butanol (v/v)				
5%	CCS	CCS	CCS	Precipitate
10%	CCS	CCS	CCS	Precipitate
20%	CCS	CCS	CCS	CCS
30%	CCS	CCS	CCS	CCS

CCS stands for clear colorless solution

Experiments to quantitatively determine the solubility of bendamustine at various temperatures for three different solutions are summarized in Figure 1 and Table 2. The amount of TBA, 20% (v/v) and 30% (v/v), used in the experiment was based on stability studies (results described below). For both solutions tested, the solubility of bendamustine decreased linearly with temperatures from 25°C to 0°C. This experiment confirmed the data shown in Table 1 and highlights the difference in bendamustine solubility for 20% and 30% TBA solutions.

Table 2. Solubility of bendamustine in TBA

	-8°C	0°C	5°C	25°C
20% (v/v) TBA 25.5 mg/mL mannitol Water, q.s. to desired volume	14 mg/mL	11 mg/mL	17 mg/mL	47 mg/mL
30% (v/v) TBA 25.5 mg/mL mannitol Water, q.s. to desired volume	20 mg/mL	18 mg/mL	27 mg/mL	65 mg/mL

C. Stability


Because of its instability in aqueous solutions due to hydrolysis with water, bendamustine requires lyophilization in order to make a product suitable for

pharmaceutical use. However, during the manufacturing of lyophilized drug products, aqueous solutions are commonly needed for filling, prior to lyophilization. Thus, the use of aqueous solutions during the compounding and fill processes for bendamustine and other nitrogen mustards can result in degradation of the drug product.

- 5 Consequently, the effect of various alcohols on the degradation of bendamustine was evaluated to determine if formulations could be found that would allow longer fill-finish times, provide lyophilate powders that could be reconstituted more quickly than the current Ribomustin® formulation, and/or provide lyophilized preparations of bendamustine with a better impurity profile with respect to certain impurities, e.g.,
 10 HP1, and BM1 dimer than Ribomustin®.

Preferably, a lyophilized preparation of the invention is stable with respect to HP1, i.e., the amount of HP1 does not increase appreciably (does not exceed the shelf-life specifications), for 6 months, more preferably 12 months, and even more preferably greater than 24 months, e.g., 36 months, when stored at about 2°C to about
 15 30°C, preferably 5°C.

Table 3 shows the stability results of bendamustine in water with no addition of alcohol over a 24 hour period at 5°C. Bendamustine degrades rapidly in water alone and forms predominantly the hydrolysis product, HP1 (monohydroxy bendamustine).




Monohydroxy bendamustine (HP1)

Formula II

Table 3. Stability of bendamustine in water

	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
0% Alcohol, i.e., Water Alone	0 hours	99.11	0.60	0.11
	3 hours	98.83	0.86	0.13
	6 hours	98.44	1.22	0.17
	24 hours	95.67	3.81	0.29


The other major degradant observed during this study and other long term stability studies was the dimer of bendamustine.



Bendamustine Dimer (BM1 Dimer)


Formula III

5 Other degradants contained in the Ribomustin lyophilized product are bendamustine ethylester (BM1EE) (Formula IV) and BM1DCE (Formula V). BM1EE is formed when bendamustine reacts with ethyl alcohol.



Bendamustine ethylester (BM1EE)

Formula IV



BM1DCE

Formula V

15 Figure 2 summarizes the purity results of an HPLC analysis after incubating bendamustine in various alcohols for 24 hours at 5°C. Results are presented as the area percent of the total peak area. The numerical values for Figure 2 are provided in
20 Tables 3-9. The purity was highest in solutions containing higher concentration of alcohols, regardless of the alcohol. Of the alcohols evaluated, bendamustine degraded the least in a solution containing about 30% (v/v) TBA. In about 10% and about 20 % alcohol solutions, n-butanol was superior in preventing degradation of bendamustine. At 20% and 30% (v/v), n-butanol in water resulted in a biphasic system due to the
25 insolubility of n-butanol in water at these concentrations.

Figures 3 and 4 show the amount of degradation of bendamustine as measured by HP1 and dimer formation quantified by HPLC (as described herein). HP1 and dimer formation increased as the amount of alcohol concentration decreased regardless of the alcohol. This increase in impurities occurred with an anticipated time dependence (see Tables 3-9). Tert-butanol and n-butanol appeared superior to other alcohols in preventing degradation of the product. As seen in Table 10, mannitol had no effect on the stabilization of bendamustine with TBA.

10 Table 4. HPLC stability results for the stability of bendamustine in various ethyl alcohol concentrations over a 24 hour period. HP1 and Dimer were impurities that increased in this study.

V/V alcohol	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
1.9% Ethanol	0 hours	99.11	0.64	0.12
	3 hours	98.83	0.90	0.14
	6 hours	98.60	1.12	0.15
	24 hours	96.16	3.41	0.27
5% Ethanol	0 hours	99.31	0.44	0.12
	3 hours	99.10	0.64	0.13
	6 hours	98.87	0.86	0.14
	24 hours	96.89	2.68	0.25
10% Ethanol	0 hours	99.44	0.33	0.11
	3 hours	99.28	0.48	0.12
	6 hours	99.10	0.65	0.12
	24 hours	98.03	1.57	0.18
20% Ethanol	0 hours	99.54	0.22	0.10
	3 hours	99.45	0.30	0.11
	6 hours	99.36	0.39	0.11
	24 hours	98.61	0.96	0.15
30% Ethanol	0 hours	99.62	0.15	0.10
	3 hours	99.56	0.21	0.11
	6 hours	99.52	0.24	0.12
	24 hours	99.21	0.45	0.12

Table 5. HPLC stability results for bendamustine in various Tert-butanol concentrations over a 24 hour period. HP1 and Dimer were impurities that increased in this study.

Concentration alcohol (v/v)	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
5% Tert-butanol	0 hours	99.34	0.41	0.12
	3 hours	99.10	0.64	0.14
	6 hours	98.85	0.88	0.13
	24 hours	97.58	2.09	0.20
10% Tert-butanol	0 hours	99.46	0.30	0.11
	3 hours	99.26	0.48	0.12
	6 hours	99.05	0.69	0.13
	24 hours	98.04	1.64	0.19
20% Tert-butanol	0 hours	99.59	0.17	0.11
	3 hours	99.48	0.29	0.11
	6 hours	99.35	0.40	0.12
	24 hours	98.35	1.27	0.20
30% Tert-butanol	0 hours	99.63	0.13	0.10
	3 hours	99.60	0.16	0.10
	6 hours	99.58	0.18	0.11
	24 hours	99.42	0.34	0.12

Table 6. HPLC stability results for various n-propyl alcohol concentrations over a 24 hour period. HP1 and Dimer were impurities that increased in this study.

Concentration alcohol (v/v)	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
5% n-Propanol	0 hours	99.25	0.43	0.13
	3 hours	99.00	0.66	0.15
	6 hours	98.72	0.94	0.16
	24 hours	97.24	2.33	0.26
10% n-Propanol	0 hours	99.34	0.33	0.15
	3 hours	99.17	0.48	0.14
	6 hours	98.92	0.70	0.16
	24 hours	97.67	1.83	0.28
20% n-Propanol	0 hours	99.45	0.33	0.13
	3 hours	99.42	0.26	0.13
	6 hours	99.29	0.39	0.14
	24 hours	98.60	0.97	0.24
30% n-Propanol	0 hours	99.53	0.15	0.13
	3 hours	99.51	0.15	0.15
	6 hours	99.44	0.20	0.11
	24 hours	99.27	0.36	0.17

5 Table 7. HPLC stability results for bendamustine in various iso-propyl alcohol concentrations over a 24 hour period. HP1 and Dimer were impurities that increased in this study.

Concentration alcohol (v/v)	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
5% Iso-propanol	0 hours	99.21	0.48	0.13
	3 hours	98.65	0.72	0.14
	6 hours	98.56	1.02	0.14
	24 hours	96.14	3.35	0.26
10% Iso-propanol	0 hours	99.32	0.37	0.12
	3 hours	99.11	0.55	0.14
	6 hours	98.85	0.75	0.16
	24 hours	97.68	1.92	0.21
20% Iso-propanol	0 hours	99.49	0.21	0.11
	3 hours	99.39	0.31	0.12
	6 hours	99.22	0.42	0.13
	24 hours	98.61	1.04	0.17
30% Iso-propanol	0 hours	99.56	0.15	0.10
	3 hours	99.47	0.20	0.12
	6 hours	99.40	0.24	0.11
	24 hours	99.15	0.52	0.14

10 Table 8. HPLC stability results for bendamustine in various methyl alcohol concentrations over a 24 hour period. HP1 and Dimer were impurities that increased in this study.

Concentration alcohol (v/v)	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
5% Methanol	0 hours	99.35	0.40	0.12
	3 hours	98.97	0.70	0.14
	6 hours	98.66	0.95	0.14
	24 hours	96.65	2.83	0.23
10% Methanol	0 hours	99.42	0.34	0.11
	3 hours	99.01	0.59	0.12
	6 hours	98.86	0.80	0.12

	24 hours	97.65	1.85	0.18
20% Methanol	0 hours	99.56	0.22	0.11
	3 hours	99.31	0.38	0.11
	6 hours	98.99	0.50	0.12
	24 hours	98.31	1.15	0.16
30% Methanol	0 hours	99.59	0.18	0.10
	3 hours	99.43	0.27	0.11
	6 hours	99.25	0.34	0.11
	24 hours	98.65	0.76	0.13

Table 9. HPLC stability results for bendamustine in various n-butyl alcohol concentrations over a 24 hour period. HP1 and Dimer were impurities that increased in this study.

Concentration alcohol (v/v)	Hold Time	Purity (%Area)	HP1 (%)	Dimer (%)
5% Butanol	0 hours	99.25	0.49	0.13
	3 hours	98.94	0.73	0.14
	6 hours	98.76	0.91	0.14
	24 hours	97.46	2.20	0.21
10% Butanol	0 hours	99.44	0.30	0.11
	3 hours	99.18	0.49	0.12
	6 hours	99.03	0.64	0.12
	24 hours	98.13	1.55	0.17
20% Butanol ^a	0 hours	99.54	0.23	0.10
	3 hours	99.45	0.31	0.11
	6 hours	99.30	0.40	0.11
	24 hours	98.81	0.91	0.14
30% Butanol ^a	0 hours	99.55	0.24	0.10
	3 hours	99.40	0.29	0.10
	6 hours	99.40	0.37	0.11
	24 hours	99.00	0.74	0.12

5 a – Both solutions had 2 layers/phases of liquids in the vial. Solutions were vortexed prior to sample preparation.

The results in Tables 1-9 indicate that the stability of bendamustine HCl with respect to HP1 and dimer improves with increasing alcohol concentration.

10

Table 10. HPLC stability results for bendamustine in TBA with and without mannitol over a 24 hour period.

Sample	Purity (%Area)	HP1 (%)
TBA 20% (v/v) with Mannitol		
0 hours	99.59	0.17
24 hours @ 5°C	99.35	1.27
TBA 20% (v/v) without Mannitol		
0 hours	100.0	0.00
24 hours @ 5°C	98.80	1.21

NOTE: The samples analyzed without mannitol were analyzed by HPLC using a normal phase method while the samples analyzed with mannitol used a reverse phase HPLC method. Slight variability may be seen in other samples analyzed between the two methods.

15

D. Lyophilization Cycle Development

Different pre-lyophilization formulations were prepared at various concentrations of bendamustine, mannitol, and alcohols in water. The cycle

development was changed and optimized at each step for freezing (fast vs. slow), primary drying (both temperature and pressure), and secondary drying as described herein.

Based upon all of the information detailed above on solubility, stability, and ease of lyophilization, preferred formulations include the following:

	Ingredients	Concentration
	Bendamustine	about 2-40 mg/mL
	Mannitol	about 0-50 mg/mL
10	Alcohol	about 0.5%-40% (v/v)
	Water, q.s. to	desired volume
	wherein the alcohol is selected from methanol, n-propanol, or isopropanol	

	Ingredients	Concentration
15	Bendamustine	about 5-20 mg/mL
	Mannitol	10-30 mg/mL
	Alcohol	1-20% (v/v)
	Water, q.s. to	desired volume
	wherein the alcohol is selected from methanol, n-propanol, or isopropanol	

20	Ingredients	Concentration
	Bendamustine	about 5-20 mg/mL
	Mannitol	10-30 mg/mL
	Alcohol	5-40% (v/v)
25	Water, q.s. to	desired volume

	Ingredients	Concentration
	Bendamustine HCl	about 12-17 mg/mL
	Mannitol	about 20-30 mg/mL
30	Alcohol	about 5-15% (v/v)
	Water, q.s. to	desired volume

	Ingredients	Concentration
	Bendamustine HCl	about 15 mg/mL
35	Mannitol	about 25.5 mg/mL
	Alcohol	about 10% (v/v)
	Water, q.s. to	desired volume

	Ingredients	Concentration
	Bendamustine HCl	about 2-40 mg/mL
	Mannitol	about 0-50 mg/mL
5	Butanol	about 0.5-20% (v/v)
	Water, q.s. to	desired volume
	Ingredients	Concentration
	Bendamustine HCl	about 5-20 mg/mL
10	Mannitol	about 10-30 mg/mL
	Butanol	about 1-10 % (v/v)
	Water, q.s. to	desired volume
	Ingredients	Concentration
15	Bendamustine HCl	about 12-17 mg/mL
	Mannitol	about 20-30 mg/mL
	Butanol	about 1-10% (v/v)
	Water, q.s. to	desired volume
20	Ingredients	Concentration
	Bendamustine HCl	about 15 mg/mL
	Mannitol	about 25.5 mg/mL
	Butanol	about 10% (v/v)
	Water, q.s. to	desired volume
25	Ingredients	Concentration
	Bendamustine HCl	about 2-50 mg/mL
	Mannitol	about 0-50 mg/mL
	Tertiary butanol	about 0.5-100 % (v/v)
30	Water, q.s. to	desired volume
	Ingredients	Concentration
	Bendamustine HCl	about 2-50 mg/mL
	Mannitol	about 0-50 mg/mL
35	Tertiary butanol	about 0.5-99.9 % (v/v)
	Water, q.s. to	desired volume

Ingredients	Concentration
-------------	---------------

	Bendamustine HCl	about 2-50 mg/mL
	Mannitol	about 0-50 mg/mL
	Tertiary butanol	about 0.5-99 % (v/v)
	Water, q.s. to	desired volume
5	Ingredients	Concentration
	Bendamustine HCl	about 2-50 mg/mL
	Mannitol	about 0-50 mg/mL
	Tertiary butanol	about 90-99 % (v/v)
10	Water, q.s. to	desired volume
	Ingredients	Concentration
	Bendamustine HCl	about 5-20 mg/mL
	Mannitol	about 10-30 mg/mL
15	Tertiary butanol	about 5-80 % (v/v)
	Water, q.s. to	desired volume
	Ingredients	Concentration
	Bendamustine HCl	about 12-17 mg/mL
20	Mannitol	about 20-30 mg/mL
	Tertiary butanol	about 10-50 % (v/v)
	Water, q.s. to	desired volume
	Ingredients	Concentration
25	Bendamustine HCl	about 12.5-15 mg/mL
	Mannitol	about 0-30 mg/mL
	Ethanol	about 20-30 % (v/v)
	Water, q.s. to	desired volume
30	Ingredients	Concentration
	Bendamustine HCl	about 15 mg/mL
	Mannitol	about 25.5 mg/mL
	Tertiary butanol	about 30 % (v/v)
35	Water, q.s. to	desired volume

EXAMPLES

The following Examples are provided to illustrate certain aspects of the present invention and to aid those of skill in the art in practicing the invention. These

Examples are in no way to be considered to limit the scope of the invention in any manner.

Materials:

Bendamustine HCl, (Degussa, Lot #s 0206005 and 0206007)

5 Mannitol, NF or equivalent (Mallinckrodt)

Ethyl Alcohol Dehydrated (200 proof), USP or equivalent (Spectrum)

Tertiary-butyl alcohol, ACS (EM Science)

Methanol (Spectrum and EMD)

Propanol (Spectrum)

10 Iso-propanol (Spectrum)

Butanol (Spectrum)

Water, HPLC grade or equivalent (EMD)

Acetonitrile, HPLC grade or equivalent (EMD)

Trifluoroacetic Acid, J.T. Baker

15 Methanol, HPLC grade or equivalent (EM Science, Cat # MX0488P-1)

Trifluoroacetic Acid, HPLC grade or equivalent (JT Baker, Cat# JT9470-01)

Equipment:

Waters 2695 Alliance HPLC system with photodiode array detector

20 Waters 2795 Alliance HPLC system with dual wavelength detector

Analytical Balance (Mettler AG285, ID #1028) and (Mettler XS205)

VirTis Lyophilizer AdVantage

Agilent Zorbax SB-C18 5 μ m 80 Å 4.6 $\times$ 250 mm column, Cat# 880975-902

25 Example 1- HPLC Procedures

Method 1

Mobile Phase A: 0.1% TFA; H₂O

Mobile Phase B: 0.1% TFA; 50% ACN:50% H₂O

UV: 230 nm

30 Flow rate: 1.0 mL/min

Column temp.: 30 °C

Column: Zorbax SB-C18 5 μ m 80 Å 4.6 $\times$ 250 mm

Sample temp.: 5 °C

Injection Volume: 10 μ L

35 Sample Concentration: 0.25 mg/mL in MeOH

Gradient: 20%B for 1 min

20 – 90%B in 23 min

90%B for 6 min

back to 20%B in 1 min

hold at 20%B for 4 min

Run time: 30 min

Post run time: 5 min

5

Method 2

Mobile Phase A: 0.1% TFA; H₂O:ACN (9:1)

Mobile Phase B: 0.1% TFA; H₂O:ACN (5:5)

UV: 230 nm

10 Flow rate: 1.0 mL/min

Column: Zorbax SB-C18 5 µm 80 Å 4.6 × 250 mm

Column temp.: 30 °C

Sample temp.: 5 °C

Injection Volume: 10 µL

15 Sample Concentration: 0.25 mg/mL in MeOH

Gradient: 0%B for 3 min

0 – 50%B in 13 min

50 – 70%B in 17 min

70 – 90%B in 2 min

20 90%B for 5 min

back to 0%B in 1 min

hold at 0%B for 4 min

Run time: 40 min

Post run time: 5 min

25

Method 3

Phase A: HPLC grade water with 0.1 % TFA(v/v)

Phase B: HPLC grade ACN / water(1:1 v/v) with 0.1%TFA(v/v)

UV: 254 nm

30 Flow rate: 1.0 mL/min

Column: Zorbax SB-C18 5 µm 80 Å 4.6 × 250 mm

Column temp.: 30 °C

Sample temp.: 5 °C

Injection Volume: 5 µL

35 Acquisition time: 30 min

Post time: 9 min

Diluent: methanol

Gradient:

Time (min.)	% Phase A	% Phase B
0.0	82	18
7.0	60	40
11.0	60	40
15.0	20	80
30.0	20	80
31.0	82	18

Sample preparation- dissolve the drug product with 200 mL MeOH. Sonicate 6 minutes. The solution can be injected directly into the HPLC (ca. 0.5 mg/mL)

5

Method 4

Phase A: HPLC grade water with 0.1 % TFA(v/v)

Phase B: HPLC grade ACN with 0.1%TFA(v/v)

UV: 254 nm

10 Flow rate: 1.0 mL/min

Column: Zorbax Bonus RP-C14 5 μ m 4.6 $\times$ 150 mm

Column temp.: 30°C

Sample temp.: 5°C

Injection Volume: 2 μ L

15 Acquisition time:31 min

Post time: 5 min

Diluent: NMP/0.1% TFA in water (50:50 v/v)

Gradient:

Time (min.)	% Phase A	% Phase B
0.0	93	7
5	93	7
13	73	27
16	73	27
25	10	90
31	10	90

20

Sample preparation for method 4- dissolve the drug product with a known amount of diluent to prepare a concentration of 4.2 mg/mL for injection directly into the HPLC. It may be necessary to perform a second dilution (the 100 mg/vial dosage form) to obtain a 4.2 mg/mL sample concentration.

25

Results

The retention times for some Bendamustine impurities using HPLC Method 1 described above are shown in Table 11. An HPLC chromatograph for Ribomustin® using the HPLC procedure described herein is shown in Fig. 6.

5 Table 11: Retention Time for Bendamustine and some of its Impurities using HPLC Method 1

Sample Name	Retention Time (min)
HP1	14.110
Bendamustine	22.182
BM1 Dimer	24.824
BM1EE	26.968

Although HPLC Method 1 was capable of resolving impurities found in bendamustine it was not capable of separating a potential impurity formed during
 10 analysis, the methyl ester of bendamustine (BM1ME). The retention time difference between BM1ME and BM1 Dimer was only 0.3 minutes. In order to resolve BM1 Dimer, another HPLC method (# 2) was developed. HPLC method #2 was capable of separating all the impurities but required a longer run time of 45 minutes (Table 12).

15 Table 12: Retention Time for bendamustine and impurities using HPLC Method 2.

Sample Name	Retention Time (min)
HP1	15.694
BM1	25.420
BM1ME	31.065
BM1 Dimer	32.467
BM1EE	36.038

20 The impurity profile of various lots of Ribomustin using HPLC Method 3 are shown in Table 13.

Table13- Ribomustine Impuirty Profile using HPLC Method 3

% Area					
Batch	Bendamustine(HCl)	HP1	BM1EE	BM1 Dimer	BM1DCE
03H08	98.14	1.07	0.21	0.34	0.03
03H07	97.67	1.5	0.2	0.33	0.04
02K27	96.93	0.93	0.29	1.18	0.08

03C08	97.61	1.24	0.19	0.46	0.02
-------	-------	------	------	------	------

Example 2- Solubility

The solubility of bendamustine HCl (bendamustine) in water (alone) and with varying amounts of methanol, ethanol, propanol, isopropanol, butanol and tertiary-butyl alcohol (TBA) was determined by visual inspection. Amounts of bendamustine at 15 mg/mL, mannitol at 25.5 mg/mL were prepared in 10 mL of the indicated alcohol solutions (Table 1) at room temperature. Samples were then refrigerated at 5°C and inspected after 0, 3, 6 and 24 hours for particulates and/or precipitates.

Results summarized in Table 1 indicate that bendamustine solubility is dependant on temperature and the amount of alcohol in aqueous solutions. For all alcohols the solubility of bendamustine increased as the concentration of alcohol increased. The formation of a precipitant was also dependent on the temperature and time.

The solubility of bendamustine was also determined in 20% (v/v) TBA containing 25.5 mg/mL mannitol in water, and 30% (v/v) TBA containing 25.5 mg/mL mannitol in water (Fig 1). Bendamustine was added to 4 mL of each solution while mixing until it would no longer dissolve. The saturated solutions were allowed to mix for 1 hour at -8°C, 0°C, 5°C, or 25°C. The samples were centrifuged and placed back at the original temperature for a minimum of 30 minutes. The -8°C sample was placed into an ice bath containing sodium chloride, which lowers the temperature of the ice bath, and the temperature was measured when the sample was pulled for analysis. An aliquot of each sample was taken and prepared for HPLC analysis.

The results of these experiments are shown in Figure 1 and Table 2. The amount of TBA, 20% (v/v) and 30% (v/v), used in the experiment (Fig. 1) was based on stability studies described herein.

As indicated in Fig. 1, the solubility of bendamustine decreased linearly with temperature (25°C to 0°C). The solubility of bendamustine was temperature dependant whether it was dissolved in water alone or with an alcohol. The 20% (v/v) TBA may likely be the lower limit required for efficient and robust pharmaceutical manufacturing due to the stability and solubility of bendamustine. A filling solution of 15 mg/mL bendamustine is close to the saturation limit of 17.2 mg/mL bendamustine at 5°C but higher than the limit at 0°C. The 30% (v/v) TBA is the recommended

concentration of TBA for the final formulation and is well within the solubility limit regardless of temperature.

Example 3-Stability

A. Stability in Water

5 Solutions of bendamustine (15 mg/mL), and mannitol (25.5 mg/mL) were prepared in water at room temperature and immediately placed in an ice bath (to lower the temperature quickly to about 5°C) for 10 minutes and then refrigerated at 5°C. A sample of each formulation was analyzed by HPLC using the methods described herein after 0, 3, 6 and 24 hours when stored at 5°C.

10 B. Stability in Alcohols

Solutions containing 15 mg/mL bendamustine, 25.5 mg/mL mannitol, and 1.9%, 5%, 10%, 20% or 30% (v/v) ethyl alcohol in water or 5%, 10%, 20% or 30% (v/v) TBA, methanol, propanol, iso-propanol, or butanol in water were prepared at room temperature, placed into an ice bath for 10 minutes and then refrigerated at 5°C. 15 A sample of each formulation was analyzed by HPLC after 0, 3, 6 and 24 hours when stored at 5°C.

C. Stability Results

Table 3 shows the stability results of bendamustine in water with no addition of alcohol over a 24 hour period at 5°C. Bendamustine degrades quickly in water but the 20 stability of bendamustine increases with increasing alcohol concentrations (Figs. 2, 3 and 4). Although alcohols are frequently used in lyophilization to aid in solubility problems, the effect of alcohols on bendamustine stability is unique, unexpected and useful in manufacturing bendamustine with fewer impurities since an aqueous solution can be used while maintaining the stability of bendamustine. TBA was found to be the 25 best stabilizer of the six alcohols tested (Figs. 2, 3, and 4). All alcohols at 30% (v/v) reduced the formation of impurities HP1 and Dimer at 5°C for up to 24 hours. With respect to TBA, HP1 reaches only about 0.4% when stored at 5°C for up to 24 hours. Lower concentrations of alcohol may not be efficient, when formulated at 15 mg/mL bendamustine and stored at 5°C due to bendamustine precipitation and impurity 30 formation.

Example 4- Formulation Optimization

After the solubility and stability of bendamustine were determined, the formulation was optimized for lyophilization. Since the concentration of

bendamustine is higher in a 30% TBA/water saturated solution as compared with other alcohol solutions, it is anticipated that the vial size required to fill 100 mg of bendamustine can be decreased from the current Ribomustin® presentation. Although a saturated solution of bendamustine contains 18 mg/mL at 0°C, a concentration of 15 mg/mL was selected for the formulation to compensate for slight differences in API solubility due to differences in bulk API purity as a result of batch differences. A concentration of 15 mg/mL bendamustine requires 6.67 mL to fill 100 mg of bendamustine HCl per vial.

The surface (sublimation) area to volume ratio is critical to producing a lyophilized product with good appearance that freeze dries quickly. Generally, lyophilized products occupy between 30% to 50% of the vial volume. A 20 mL vial with 6.67 mL contains about 30% of its capacity and has a surface area ratio of 0.796 cm²/mL.

Mannitol was selected as the bulking agent in order to maintain a formulation similar to Ribomustin®. Studies were performed to evaluate the effect of mannitol on bendamustine solubility and appearance of the product. Mannitol decreases the solubility of bendamustine (at 15 mg/mL) in both ethanol and TBA aqueous solutions. For example, solutions containing 5% and 10% ethanol and TBA without mannitol did not precipitate over 24 hours. However, for samples with mannitol (Table 1) precipitate was observed within 24 hours. There was no precipitate with aqueous solutions containing 30% (v/v) TBA, 15 mg/mL bendamustine, and 25.5 mg/mL mannitol. In order to maintain a well formed cake resistant to breakage during handling, a minimum of 134 mg/vial of mannitol was required with no difference observed in vials up to 200 mg/vial of mannitol.

All alcohols tested increased the stability and solubility of bendamustine. However, a significant mole fraction was required to affect the stability of the filling solution and the ease of manufacturing. Smaller alcohols have the undesirable effect of lowering the freezing point of the bulk solution and thus requiring long lyophilization cycles at lower temperatures. Higher concentrations of methanol and ethanol produced unattractive cakes that were difficult to reconstitute. 10% ethanol, 20% ethanol, 10% iso-propanol, 20% iso-propanol, or 30% TBA aqueous solutions containing bendamustine (15 mg/mL), mannitol (25.5 mg/mL) were prepared and lyophilized. The lyophilized vials filled from solutions of 10% ethanol, 20% ethanol,

10% iso-propanol, 20% iso-propanol produced either a collapsed cake or a film residue. The only solvent system producing an acceptable cake was 30% TBA. Additionally, reconstitution of 10% ethanol, 20% ethanol, 10% iso-propanol, 20% iso-propanol lyophilized vials were difficult and did not fully dissolve until >45 minutes.

5 The ability to utilize a smaller vial is constrained by the concentration or solubility of bendamustine in the aqueous/organic solution. At lower concentrations of ethanol, methanol, isopropanol and n-propanol, which produced acceptable cake appearance, a more dilute solution of bendamustine is required due to solubility limitations. To maintain a presentation with 100 mg of bendamustine per vial, a vial
10 larger than 50 mL would be required. Also, stability studies herein indicated that at the lower alcohol concentration, the chemical stability was not sufficient to allow for acceptable filling times.

One of the factors affecting the ease of reconstitution is the porosity of the lyophilate. In general, amorphously precipitated solids with little surface area are
15 more difficult to solubilize. Most lyophilates containing mannitol will reconstitute within 3-5 minutes as long as there is no precipitate formed during lyophilization, frequently caused by evaporation of a liquid (melt back). Based on our experience with several lyophilization solvent systems and not wishing to be bound to any particular theory, the problems associated with Ribomustin® reconstitution may be
20 associated with precipitation caused by melt back during lyophilization. Most organic solvents do not lyophilize efficiently and cause melt back because of their low melting point. TBA (tertiary butyl alcohol) has a high melting point and a similar vapor pressure as compared to water. TBA is removed by sublimation, not evaporation, at about the same rate as water. Lyophilates produced with 30% (v/v) TBA according to
25 the invention reconstitute within 3-10 minutes as compare to commercially available Ribomustin which may take 30-45 minutes.

Based upon the solubility, stability, ease of reconstitution and manufacturing considerations, the following is a preferred pre-lyophilization formulation of the present invention: bendamustine HCl about 15 mg/mL, mannitol about 25.5 mg/mL,
30 about 30% (v/v) tertiary-butyl alcohol, and q.s. using water for Injection. The formulation is then filled at 5°C using 6.67 mL in an amber 20 mL, 20 mm vial and partially stoppered with a bromobutyl stopper and loaded into a pre-chilled lyophilizer.

Example 5- Impurity assessment

Major impurities introduced during Ribomustin® manufacturing, compounding, fill, and lyophilization procedure, as determined by HPLC analysis (Fig. 6), are the hydrolysis product HP1, the Dimer, and the ethyl ester of bendamustine, BM1EE. BM1EE can be formed during drug substance manufacturing, e.g., during recrystallization and/or purification processes. BM1EE is known to be a more potent cytotoxic drug than bendamustine. Experiments were undertaken to determine if the use of a 30% TBA aqueous filling solution would lead to the formation of bendamustine t-butyl ester.

Experiments were performed using traditional Fisher esterification reaction conditions required for the formation of t-butyl ester of bendamustine. Bendamustine was heated in 60°C TBA with HCl for 20 hours. No reaction was observed. This result indicated that it would be very difficult to form the tert-butyl ester of bendamustine during the fill/finish process. No new impurities in drug product manufactured from TBA have been observed in stability studies to date.

To aid in the testing of the drug product, synthetic routes using more reactive sources of the t-butyl moiety were developed. Another attempt to make tert-butyl ester was carried out by formation of the acyl chloride of bendamustine. A suspension of bendamustine in methylene chloride was treated with oxalyl chloride and N,N-dimethylformamide. After acyl chloride was formed, the solvent was concentrated. The residue was added to methylene chloride, tert-butanol, triethylamine, and 4-dimethylaminopyridine and the mixture was stirred at room temperature overnight. After adding all solvents and purification, an unknown compound was given. The LC-MS did not match the molecular weight of bendamustine tert-butyl ester and the proton NMR did not showed the peak for tert-butyl. Therefore, this attempt also failed to produce the bendamustine tert-butyl ester. Thus, using TBA as the co-solvent has an additional benefit of not forming the ester from the alcohol.

Example 6- Lyophilization Cycle Development

Numerous lyophilization cycles were performed to evaluate the critical stages of lyophilization and achieve the most efficient drying cycle. Experiments were performed to evaluate the effect of the freezing rate, primary drying temperature, time, and pressure on the product.

A. Freezing Rate

The literature reports that TBA adopts different crystal forms depending on the freeze rate. In some TBA solutions, the slower the product froze, the quicker it dried. Larger crystals formed during slow freezing producing bigger pores allowing more efficient sublimation. However, during studies with bendamustine, the freezing rate
5 was not found to be a critical processing parameter when evaluated at 2 and 8 hours.

B. Primary and Secondary Drying

During the first attempts to lyophilize from 30% TBA solutions, the lyophilized cake fractured and powder was ejected from the vial. These cakes appeared to contain amorphous particles within the lyophilate, an indication of melt
10 back. This phenomenon was reproducible and occurred when the product reached about -10°C (refer to Fig. 5) independent of the warming rate. Several variables were tested to determine the cause and solution to the problem of the powder ejection. The pressure was raised from 50 µm to 150 µm during primary drying, but powder ejection was still observed but to a lesser extent. This experiment was then repeated except the
15 freezing rate was extended to 8 hours from 2 hours. This change had no effect.

The length of primary drying was next evaluated. For example, the following very slow drying cycle was evaluated: freezing from +25°C to -50°C in eight hours; holding at -50°C for 5 hours, warming and drying from -50°C to -25°C in seven hours; holding for twenty hours at -25°C, warming and drying from -25°C to -15°C in two
20 hours and holding for twenty hours at -15°C, warming and drying from -15°C to 40°C in six hours and holding for twenty hours at 40°C while maintaining a chamber pressure of 150 µm throughout drying. No powder ejection (Fig 5) was observed. This cycle resulted in a well-formed cake without fracture that reconstituted readily. Without wishing to be bound to a particular theory, the problems with powder ejection
25 and difficulty with reconstitution may be the result of drying the lyophilate too quickly, thus resulting in strong vapor flow out of the cake as well as melt back. With the use of a less aggressive drying cycle an aesthetic, stable, and easy to reconstitute cake was reproducibly formed. Thus, removing all unbound water and tertiary-butyl alcohol prior to secondary drying may prevent melt back as well as powder ejection.
30 The lyophilization cycle was further optimized under these gentle conditions (Fig. 5). There were no immediate degradation products as a result of drying at 40°C for up to 20 hours.

Example 7- Lyophilization cycle

Step	Description	Time (Hour)	Temperature (°C)	Pressure (Microns)
1	Hold	0.25	5°C	-
2	Ramp	8	-50°C	-
3	Hold	4	-50°C	-
4	Ramp	3	-20°C	150
5	Hold	6	-20°C	150
6	Ramp	1	-15°C	150
7	Hold	20	-15°C	150
8	Ramp	0.5	-12°C	150
9	Hold	15.5	-12°C	150
10	Ramp	15	35°C	50
11	Hold	10	35°C	50
12	Ramp	1	40°C	50
	Hold	5	40°C	50
Total		89.25	-	-

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods and in the steps or in the sequence of steps of the method described herein without departing from the spirit and scope of the invention. More specifically, it will be apparent that certain solvents which are both chemically and physiologically related to the solvents disclosed herein may be substituted for the solvents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit and scope of the invention as defined by the appended claims.

All patents, patent applications, and publications mentioned in the specification are indicative of the levels of those of ordinary skill in the art to which the invention pertains. All patents, patent applications, and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.


The invention illustratively described herein suitably may be practiced in the absence of any element(s) not specifically disclosed herein. Thus, for example, in

each instance herein any of the terms “comprising”, “consisting essentially of”, and “consisting of” may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of
5 excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those
10 skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

What is claimed is:

1. A pharmaceutical composition of bendamustine containing not more than about 0.9% (area percent of bendamustine) HP1 as shown in Formula II,

5



Formula II

wherein said HP1 is the amount of HP1 present at time zero after reconstitution of a lyophilized preparation of bendamustine.

10

2. The composition according to claim 1, wherein the amount of HP1 is not more than 0.5% (area percent of bendamustine) at time zero after reconstitution of a lyophilized preparation of bendamustine.

15

3. The composition according to claim 1, wherein the amount of HP1 is not more than 0.4% (area percent of bendamustine) at time zero after reconstitution of a lyophilized preparation of bendamustine.

20

4. The composition according to claim 1, wherein the amount of HP1 is not more than 0.3% (area percent of bendamustine) at time zero after reconstitution of a lyophilized preparation of bendamustine.

25

5. A lyophilized preparation of bendamustine containing not more than about 0.9% (area percent of bendamustine) HP1 at release.

30


6. A lyophilized preparation of bendamustine containing not more than about 0.5% (area percent of bendamustine) HP1 at release.

7. The lyophilized preparation according to claim 5, wherein the preparation is packaged in a vial or other pharmaceutically acceptable container.

8. The lyophilized preparation according to claim 6, wherein said preparation is stable with respect to the amount of HP1 for at least about six months when stored at 5° C.
9. The lyophilized preparation according to claim 6, wherein said preparation is stable with respect to the amount of HP1 for at least about 12 months when stored at 5° C.
10. The lyophilized preparation according to claim 6, wherein said preparation is stable with respect to the amount of HP1 for at least about 24 months when stored at 5° C.
11. A pharmaceutical dosage form comprising a pharmaceutical composition of bendamustine containing not more than about 0.9% HP1, wherein said HP1 is the amount of HP1 present at release.
12. A pharmaceutical dosage form comprising a pharmaceutical composition of bendamustine containing not more than about 0.5% HP1, wherein said HP1 is the amount of HP1 present at release.
13. A pharmaceutical dosage form of claim 11, wherein the pharmaceutical dosage form comprises about 5 mg to about 500 mg of bendamustine.
14. A pharmaceutical dosage form of claim 11, wherein the pharmaceutical dosage form comprises about 10 mg to about 300 mg of bendamustine.
15. A pharmaceutical dosage form of claim 11, wherein the pharmaceutical dosage form comprises about 25 mg of bendamustine.
16. A pharmaceutical dosage form of claim 11, wherein the pharmaceutical dosage form comprises about 100 mg of bendamustine.
17. A pharmaceutical dosage form of claim 11, wherein the pharmaceutical dosage form comprises about 200 mg of bendamustine.

18. A pharmaceutical dosage form comprising the lyophilized preparation of claim 5.
19. A pharmaceutical composition of bendamustine comprising bendamustine containing not more than about 0.5% (area percent of bendamustine) HP1 and a trace
5 amount of one or more organic solvents, wherein said HP1 is the amount of HP1 present at release.
20. A pharmaceutical composition of bendamustine according to claim 19 wherein the organic solvent is selected from the group consisting of one or more of tertiary
10 butanol, n-propanol, n-butanol, isopropanol, ethanol, methanol, acetone, ethyl acetate, dimethyl carbonate, acetonitrile, dichloromethane, methyl ethyl ketone, methyl isobutyl ketone, 1-pentanol, methyl acetate, carbon tetrachloride, dimethyl sulfoxide, hexafluoroacetone, chlorobutanol, dimethyl sulfone, acetic acid, and cyclohexane.
- 15 21. A pharmaceutical composition according to claim 20, wherein the organic solvent is selected from the group consisting of one or more of ethanol, methanol, propanol, butanol, isopropanol, and tertiary butanol.
22. A pharmaceutical composition according to claim 19, wherein the organic solvent
20 is tertiary butanol.
23. A lyophilized preparation according to claim 5 further comprising a trace amount of an organic solvent.
- 25 24. A lyophilized preparation according to claim 23 wherein said organic solvent is tertiary butanol.
25. In a method for obtaining agency approval for a bendamustine product, the improvement which comprises setting a release specification for bendamustine
30 degradants at less than 4.0 % (area percent bendamustine) for a bendamustine product containing not more than about 0.5% (area percent of bendamustine) HP1 at release.

26. In a method for obtaining agency approval for a bendamustine product, the improvement which comprises setting a release specification for bendamustine of HP1 at less than or equal to 1.5% for a bendamustine product containing not more than about 0.5% (area percent of bendamustine) HP1 at release.
- 5
27. In a method for obtaining agency approval for a bendamustine product, the improvement which comprises setting a shelf-life specification for bendamustine degradants at less than 7.0% (area percent bendamustine) for a bendamustine product containing not more than about 0.5% (area percent of bendamustine) HP1 at release.
- 10
28. A process for manufacturing a lyophilized preparation of bendamustine which comprises controlling for the concentration of bendamustine degradants in the final product, such that, at release, the concentration of bendamustine degradants is less than 4.0 % (area percent of bendamustine) and the concentration of HP1 is less than 0.5% (area percent of bendamustine).
- 15
29. A process for manufacturing a lyophilized preparation of bendamustine which comprises controlling for the concentration of bendamustine degradants in the final product, such that the concentration of HP1 is less than 0.9% (area percent of bendamustine) at release and the concentration of bendamustine degradants is less than 7.0% at the time of product expiration; wherein said product is stored at 5°C.
- 20
30. A process for manufacturing a lyophilized preparation of bendamustine which comprises controlling for the concentration of bendamustine degradants in the final product, such that the concentration of HP1 is less than 0.5% (area percent of bendamustine) at release and the concentration of bendamustine degradants is less than 7.0% at the time of product expiration; wherein said product is stored at 5°C.
- 25
31. A lyophilized preparation of bendamustine wherein the concentration of bendamustine ethylester (as shown in Formula IV)
- 30




Formula IV

is no more than 0.2% greater than the concentration of bendamustine ethylester as found in the drug substance used to make the lyophilized preparation.

5

32. A lyophilized preparation of bendamustine according to claim 5 containing not more than about 0.5% bendamustine ethylester as shown in Formula IV



Formula IV.

10

33. A bendamustine pre-lyophilization solution or dispersion comprising one or more organic solvents, wherein said solution or dispersion comprises at least one stabilizing concentration of an organic solvent which reduces the level of degradation of bendamustine so that the amount of HP1 produced during lyophilization, from about 0 to 24 hours, does not exceed 0.9% (area percent bendamustine).

34. A bendamustine pre-lyophilization solution or dispersion comprising one or more organic solvents, wherein said solution or dispersion comprises at least one stabilizing concentration of an organic solvent which reduces the level of degradation of bendamustine so that the amount of HP1 produced during lyophilization, from about 0 to 24 hours, does not exceed 0.5% (area percent bendamustine).

35. The lyophilized powder produced from the pre-lyophilization solution or dispersion according to claim 33.

36. A method of preparing a bendamustine lyophilized preparation comprising,
a) dissolving bendamustine in a stabilizing concentration of an alcohol solvent comprising between about 5% to about 100% (v/v) alcohol to form a pre-lyophilization solution; and
5 b) lyophilizing the pre-lyophilization solution;
wherein said bendamustine lyophilized preparation contains not more than about 0.9% (area percent of bendamustine) HP1 as shown in Formula II, wherein said HP1 is the amount of HP1 present at release.
- 10 37. A method of preparing a bendamustine lyophilized preparation comprising,
a) dissolving bendamustine in a stabilizing concentration of an alcohol solvent comprising between about 5% to about 100% (v/v) alcohol to form a pre-lyophilization solution; and
b) lyophilizing the pre-lyophilization solution;
15 wherein said bendamustine lyophilized preparation contains not more than about 0.5% (area percent of bendamustine) HP1 as shown in Formula II, wherein said HP1 is the amount of HP1 present at release.
38. A method according to claim 36, wherein the alcohol concentration is between
20 about 5% to about 99.9%.
39. A method according to claim 36, wherein said alcohol is selected from one or more of methanol, ethanol, propanol, iso-propanol, butanol, and tertiary-butanol.
- 25 40. A method according to claim 39, wherein said alcohol is tertiary-butanol.
41. A method according to claim 40, wherein said tertiary butanol is at a concentration of about 20% to 30%.
- 30 42. A method according to claim 40, wherein said tertiary butanol is at a concentration of about 30%.

43. A method according to claim 36, wherein an excipient is added before lyophilization.
44. A method according to claim 43, wherein the excipient is mannitol.
- 5 45. A method according to claim 36, wherein the bendamustine concentration is about 2 to 50 mg/mL.
- 10 46. The lyophilized powder obtained from the method according to one of claims 36 to 45.
47. A method according to claim 36 wherein step b) comprises:
- i) freezing the pre-lyophilization solution to a temperature below about -40°C to form a frozen solution;
 - 15 ii) holding the frozen solution at or below -40°C for at least 2 hours;
 - iii) ramping the frozen solution to a primary drying temperature between about -40°C and about -10°C to form a dried solution;
 - iv) holding for about 10 to about 70 hours;
 - v) ramping the dried solution to a secondary drying temperature between about 20 25°C and about 40°C; and
 - vii) holding for about 5 to about 40 hours to form a bendamustine lyophilized preparation.
- 25 48. A method according to claim 47, wherein said alcohol is tertiary-butanol.
49. A method according to claim 48, wherein said tertiary butanol is at a concentration of about 20% to 30%.
- 30 50. A method according to 49, wherein said tertiary butanol is at a concentration of about 30%.
51. The lyophilized powder obtained from the method according to one of claims 47 to 50.

52. A method according to claim 36 wherein step b) comprises:
- i) freezing the pre-lyophilization solution to about -50°C to form a frozen solution;
 - ii) holding the frozen solution at about -50°C for at least 2 hours to about 4 hours;
 - 5 iii) ramping to a primary drying temperature between about -20°C and about -12°C to form a dried solution;
 - iv) holding at a primary drying temperature for about 10 to about 48 hours;
 - v) ramping the dried solution to a secondary drying temperature between about 25°C and about 40°C; and
 - 10 vi) holding at a secondary drying temperature for at least 5 hours up to about 20 hours.
53. A method according to claim 52, wherein said alcohol is tertiary-butanol.
- 15 54. A method according to claim 53, wherein said tertiary butanol is at a concentration of about 20% to 30%.
55. A method according to 54, wherein said tertiary butanol is at a concentration of about 30%.
- 20 56. The lyophilized powder obtained from the method according to one of claims 52 to 55.
57. A method according to claim 36 wherein step b) comprises: i) starting with a shelf temperature of about 5°C for loading; ii) freezing to about -50°C over about 8 hours; iii) holding at -50°C for about 4 hours; iv) ramping to about -20°C over about 3 hours; v) holding at about -20°C for 6 hours; ramping to about -15°C over about 1 hour; vi) holding at -15°C for about 20 hours; vii) ramping to about -15°C over about 1 hour; viii) holding at about -15°C for about 20 hours; ix) ramping to about -12°C over about 0.5 hours; x) holding at about -12°C for about 15.5 hours; xi) ramping to between about 25°C and about 40°C or higher for about 15 hours; xii) holding between about 25°C and about 40°C for about 10 hours; xiii) ramping to about 40°C over about 1 hour; andxiv) holding at about 40°C for about 5 hours; unloading at about 5°C, at a
- 25
- 30

pressure of about 13.5 psi in a pharmaceutically acceptable container that is hermetically sealed; wherein the pressure is about 150 microns throughout primary drying and 50 microns throughout secondary drying.

5 58. A lyophilization cycle according to claim 57, wherein step (xi) is ramped to about 30-35°C for about 15 hours.

59. The lyophilized powder prepared from the lyophilization cycle of claim 57 or 58.

10 60. A formulation for lyophilization comprising bendamustine at a concentration of about 15 mg/mL, mannitol at a concentration of about 25.5 mg/mL, tertiary-butyl alcohol at a concentration of about 30% (v/v) and water.

61. A lyophilized preparation made from the formulation according to claim 60.

15

62. A method of treating a medical condition in a patient comprising dissolving the preparation of claim 5 in a pharmaceutically acceptable solvent to produce a pharmaceutically acceptable solution and administering to said patient a therapeutically effective amount of said solution, wherein said condition is amenable to treatment with said preparation.

20

63. A method of treating according to claim 62, wherein said condition is selected from chronic lymphocytic leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, multiple myeloma, breast cancer, small cell lung cancer, and an autoimmune disease.

25

64. A method of treating according to claim 63, wherein said condition is non-Hodgkin's lymphoma.

65. A method of treating according to claim 63, wherein said condition is chronic lymphocytic leukemia.

30

66. A method of treating according to claim 63, wherein said condition is multiple myeloma.

67. A method of treating according to claim 62 further comprising administering the dissolved preparation of claim 5 in combination with one or more anti-neoplastic agents wherein said antineoplastic agent is given prior, concurrently, or subsequent to the administration of the dissolved preparation of claim 5.

68. A method of treating according to claim 67 wherein the antineoplastic agent is an antibody specific for CD20, wherein said antibody is given prior, concurrently or subsequent to the administration of the dissolved preparation of claim 5.

10

69. A method of treating according to claim 62 wherein the autoimmune disease is rheumatoid arthritis, multiple sclerosis or lupus.

70. A method of treating according to claim 62, wherein the medical condition is a hyperproliferative disorder.

15

71. A pharmaceutical dosage form of bendamustine containing not more than about 0.9% HP1 (area percent of bendamustine) wherein said dosage form comprises a vial or other pharmaceutically acceptable container, wherein said HP1 is the amount of HP1 present pre-reconstitution or at time zero after reconstitution of said dosage form.

20

72. A pharmaceutical dosage form of bendamustine containing not more than about 0.5% HP1 (area percent of bendamustine) wherein said dosage form comprises a vial or other pharmaceutically acceptable container, wherein said HP1 is the amount of HP1 present pre-reconstitution or at time zero after reconstitution of said dosage form.

25

73. A pharmaceutical dosage form according to claim 72, wherein the vial or other pharmaceutically acceptable container contains bendamustine at a concentration of about 10 to about 500 mg/container.

30

74. A pharmaceutical dosage form according to claim 72, wherein the vial or other pharmaceutically acceptable container contains bendamustine at a concentration of about 100 mg/container.

75. A pharmaceutical dosage form according to claim 72, wherein the vial or other pharmaceutically acceptable container further comprises mannitol at a concentration of about 5 mg to about 2 g/container.

5

76. A pharmaceutical dosage form according to claim 72, wherein the vial or other pharmaceutically acceptable container further comprises mannitol at a concentration of about 170 mg/container.

10 77. A pre-lyophilized pharmaceutical composition of bendamustine comprising about 15 mg/mL bendamustine HCl, about 25.5 mg/mL mannitol, about 30% (v/v) tertiary-butyl alcohol, and water.

78. The preparation of claim 5 which is a pharmaceutical composition.

15

1/6




FIG. 1

Bendamustine Purity after 24 hours at 5°C in Various
Alcohol/Water Co-Solvents




FIG. 2

3/6

HP1 information after 24 hours stored at 5°C in Various
Alcohol/Water Co-Solvents




FIG. 3

4/6




FIG. 4

5/6




FIG. 5





FIG. 6

9/6

HP1 information after 24 hours stored at 5°C in Various
Alcohol/Water Co-Solvents

