

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2009203350 B2

(54) Title
Humanized anti-alpha9 integrin antibodies and the uses thereof

(51) International Patent Classification(s)

C12N 15/09 (2006.01)	C12N 1/21 (2006.01)
A61K 39/395 (2006.01)	C12N 5/10 (2006.01)
C07K 16/28 (2006.01)	C12P 21/08 (2006.01)
C12N 1/15 (2006.01)	G01N 33/563 (2006.01)
C12N 1/19 (2006.01)	G01N 33/577 (2006.01)

(21) Application No: **2009203350** **(22) Date of Filing:** **2009.01.13**

(87) WIPO No: **WO09/088105**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/020,527	2008.01.11	US

(43) Publication Date: **2009.07.16**

(44) Accepted Journal Date: **2014.03.13**

(71) Applicant(s)

Gene Techno Science Co., Ltd.;Kaken Pharmaceutical Co., Ltd.

(72) Inventor(s)

Tsurushita, Naoya;Kumar, Shankar;Kon, Shigeyuki;Tso, J. Yun

(74) Agent / Attorney

FB Rice, Level 23 44 Market Street, Sydney, NSW, 2000

(56) Related Art

AU 2007273467

EP 1840135

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

**(19) World Intellectual Property Organization
International Bureau**



**(43) International Publication Date
16 July 2009 (16.07.2009)**

PCT

(10) International Publication Number
WO 2009/088105 A1

(51) International Patent Classification:

<i>C12N 15/09</i> (2006.01)	<i>C12N 1/21</i> (2006.01)
<i>A61K 39/395</i> (2006.01)	<i>C12N 5/10</i> (2006.01)
<i>C07K 16/28</i> (2006.01)	<i>C12P 21/08</i> (2006.01)
<i>C12N 1/15</i> (2006.01)	<i>G01N 33/563</i> (2006.01)
<i>C12N 1/19</i> (2006.01)	<i>G01N 33/577</i> (2006.01)

(74) **Agents:** KOBAYASHI, Hiroshi et al.; ABE, IKUBO & KATAYAMA, Fukuoka Bldg. 9th Fl., 8-7, Yaesu 2-chome, Chuo-ku, Tokyo 1040028 (JP).

(21) International Application Number:

PCT/JP2009/050606

(22) International Filing Date: 13 January 2009 (13.01.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

61/020,527 11 January 2008 (11.01.2008) US

(71) **Applicants (for all designated States except US):**
GENE TECHNO SCIENCE CO., LTD. [JP/JP]; 1, Kita-2jo-Nishi 9-chome, Chuo-ku, Sapporo-shi, Hokkaido, 0600002 (JP). **KAKEN PHARMACEUTICAL CO., LTD.** [JP/JP]; 28-8, Honkomagome 2-chome, Bunkyo-ku, Tokyo, 1138650 (JP).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with *international search report*
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

WO 2009/088105 A1

(54) Title: HUMANIZED ANTI- α 9 INTEGRIN ANTIBODIES AND THE USES THEREOF

(57) Abstract: The present invention provides humanized antibodies that immunospecifically recognize human $\alpha 9$ integrin. Some of these antibodies inhibit the biological functions of the $\alpha 9$ integrin, thereby exhibiting therapeutic effects on various disorders or diseases that are associated with $\alpha 9$ integrin, including cancer, e.g., the growth and metastasis of a cancer cell, and inflammatory diseases, e.g., rheumatoid arthritis, osteoarthritis, hepatitis, bronchial asthma, fibrosis, diabetes, arteriosclerosis, multiple sclerosis, granuloma, an inflammatory bowel disease (ulcerative colitis and Crohn's disease), an autoimmune disease, and so forth.

DESCRIPTION

HUMANIZED ANTI- α 9 INTEGRIN ANTIBODIES AND THE USES THEREOF

5

1. FIELD OF THE INVENTION

The present invention relates to humanized antibodies that immunospecifically recognize human α 9 integrin and to their therapeutic and diagnostic uses for various diseases or disorders that are associated with or involve 10 α 9 integrin, including cancer, inflammatory diseases, autoimmune diseases, and the like.

2. BACKGROUND OF THE INVENTION

Cells adhere to extracellular matrix (hereinafter abbreviated as ECM) 15 mediated by a group of cell surface receptors which are termed integrins. Integrins perform their functions by forming 1 : 1 heterodimers of α and β chains. At least 18 types of α chain, 8 types of β chain and 24 types of $\alpha\beta$ heterodimer have been identified and confirmed so far. It is known that each integrin recognizes a specific ligand. Integrins are classified into subfamilies depending upon their ligand 20 specificities or functions, and divided into collagen receptors, laminin receptors, RGD receptors recognizing an Arg-Gly-Asp (RGD) sequence present in fibronectin, vitronectin, etc., and leukocyte-specific receptors present only in leukocytes (Hynes, R. O., 2002, Integrins: Bidirectional, Allosteric Signaling Machines. *Cell* 110: 673-87; Miyasaka, M., 2000, New edition of *Adhesion Molecule Handbook*, 25 Shujunsya). The α 4 and α 9 integrins are members of a subfamily that does not belong to any of these types and called the α 4 integrin subfamily (Elise L. Palmer, Curzio Riegg, Ronald Ferrando, Robert Pytela, Sheppard D., 1993, Sequence and Tissue Distribution of the Integrin α 9 Subunit, a Novel Partner of β 1 That Is Widely Distributed in Epithelia and Muscle. *The Journal of Cell Biology*, 123: 1289-97). 30 Meanwhile, ECM used to be considered so far to serve as a mere cementing substance between cells. It has now become clear that the integrin-mediated ECM-cell interaction is significantly involved in regulating the growth, adhesion, movement, etc. of cells and associated with the onset of diseases including a

progression of cancer, an exacerbation of inflammation, and the like.

For example, osteopontin (hereinafter abbreviated as OPN) which is one of the ECMs is a secreted, acidic phosphorylated glycoprotein with a molecular weight of about 41 kDa and is a molecule, whose expression is widely observed in breast 5 milk, urine, renal tubules, osteoclasts, osteoblasts, macrophages, activated T cells, tumor tissues, and so forth. OPN has the adhesion sequences, GRGDS (SEQ ID NO:1) at the center of its molecule, the SVVYGLR (SEQ ID NO:2) sequence in human OPN or the SLAYGLR (SEQ ID NO:3) sequence in mouse OPN, and a thrombin-cleavage site in close proximity thereto, and binds through the GRGDS 10 (SEQ ID NO:1) sequence to the RGD integrin or to the $\alpha 4$ ($\alpha 4\beta 1$) and $\alpha 9$ ($\alpha 9\beta 1$) integrins through the SVVYGLR (SEQ ID NO:2) sequence or the SLAYGLR (SEQ ID NO:3) sequence.

WO 02/081522 discloses a therapeutic effect on rheumatoid arthritis or hepatitis by inhibiting the OPN functions using OPN knockout mice or neutralizing 15 antibodies against OPN. Moreover, this publication discloses that the SVVYGLR (SEQ ID NO:2) sequence is essential as recognizing the $\alpha 9$ and $\alpha 4$ integrins for pathogenesis of an inflammatory disease and that receptors for OPN are expressed in immunocompetent cells or the like and associated with an inflammatory disease.

Differences in binding profile have been found in that $\alpha 4\beta 1$ binds both to 20 OPN not cleaved with thrombin (uncleaved OPN) and to the N-terminal fragment of thrombin-cleaved OPN (cleaved OPN), whereas $\alpha 9\beta 1$ binds only to the cleaved OPN (Y. Yokosaki, et al., (1999) *The Journal of Biological Chemistry*, 274: 36328-36334; P. M. Green, et al., (2001) *FEBS Letters*, 503: 75-79; S. T. Barry, et al., (2000) *Experimental Cell Research*, 258: 342-351).

25 The $\alpha 4$ and $\alpha 9$ integrins share many common ligands other than OPN. Known ligands are the EDA domain of fibronectin, propeptide-von Willebrand factor (pp-vWF), tissue transglutaminase (tTG), blood coagulation factor XIII, vascular cell adhesion molecule-1(VCAM-1), etc. In addition, the CS-1 domain of fibronectin, MadCAM-1 ($\alpha 4\beta 7$), etc. are known as the ligands specifically recognized by the $\alpha 4$ 30 integrin. Tenascin-C, plasmin, etc. are known as the ligands specifically recognized by the $\alpha 9$ integrin.

The amino acid sequences for the integrin subunits $\alpha 9$, $\alpha 4$ and $\beta 1$ are

publicly known. For instance, human $\alpha 9$ is registered as NM_002207, mouse $\alpha 9$ as NM_133721, human $\alpha 4$ as NM_000885, mouse $\alpha 4$ as NM_010576, human $\beta 1$ as X07979, and mouse $\beta 1$ as NM_010578, at the GenBank. These integrins are also known to have high similarities between species in amino acid sequence.

5

3. SUMMARY OF THE INVENTION

While a variety of drugs are known at present for the treatment of cancer, inflammatory diseases and autoimmune diseases, it has been desired to develop a preventive and/or therapeutic agent, etc. having more improved therapeutic effects on 10 cancer, inflammatory diseases and autoimmune diseases. The present invention is based, in part, on the discovery by the present inventors that a specific inhibitory antibody against the $\alpha 9$ integrin has cancer-suppressing and anti-inflammatory effects.

Previously, the present inventors isolated mouse monoclonal antibodies that 15 immunospecifically recognize human $\alpha 9$ integrin and are produced by hybridoma clones, 1K11, 21C5, 24I11, 25B6 and 28S1 (Depository Accession Nos. FERM BP-10510, FERM BP-10511, FERM BP-10512, FERM BP-10513 and FERM BP-10832, respectively), and those that immunospecifically recognize mouse $\alpha 9$ integrin and are produced by hybridoma clones, 18R18D, 12C4'58, 11L2B and 20 55A2C (Depository Accession Nos. FERM ABP-10195, FERM ABP-10196, FERM ABP-10197 and FERM ABP-10198, respectively). Herein, the hybridoma clone designations are interchangeably used as the designations of the monoclonal antibodies produced by the clones. All of these mouse anti-human $\alpha 9$ integrin antibodies were of IgG1 isotype. Some of these monoclonal antibodies inhibit the 25 binding between human and/or mouse $\alpha 9$ integrin and a ligand of $\alpha 9$ integrin, such as osteopontin. Thus, these anti- $\alpha 9$ integrin antibodies inhibit the $\alpha 9$ integrin functions and exhibit therapeutic effects on cancer, e.g., the growth or metastasis of cancer cells, and on inflammatory diseases, e.g., rheumatoid arthritis, osteoarthritis, hepatitis, bronchial asthma, fibrosis, diabetes mellitus, arteriosclerosis, multiple 30 sclerosis, granuloma, an inflammatory bowel disease (ulcerative colitis and Crohn's disease), an autoimmune disease, and the like.

Furthermore, the anti- $\alpha 9$ integrin antibodies of the present invention can be

used as an *in vivo* diagnostic agent to detect the presence and the level of $\alpha 9$ integrin expression in a subject, thereby diagnosing a disorder or a disease involving $\alpha 9$ integrin.

However, since these monoclonal antibodies are of mouse origin, possible 5 adverse effects due to their immunogenicity in humans have hampered their direct applications to diagnostic or therapeutic uses in humans. In order to reduce the immunogenicity, the present inventors have prepared a humanized antibody that have biological activities corresponding to those exhibited by the original mouse anti- $\alpha 9$ integrin antibody from which said humanized antibody was derived.

10 Accordingly, the present invention provides a humanized antibody or an antigen-binding fragment thereof, which immunospecifically recognizes human $\alpha 9$ integrin, said antibody comprising an antigen-binding region partially derived from a non-human origin and partially derived from a human origin. In a specific embodiment, the humanized antibody or an antigen-binding fragment thereof of the 15 present invention comprises a complementarity determining region (CDR) derived from a non-human source (donor), such as 1K11, 21C5, 24I11, 25B6 and 28S1 monoclonal antibodies, and a framework region (FR) derived from a human source (acceptor). In one embodiment, said humanized antibody or an antigen-binding fragment thereof inhibits the binding between human $\alpha 9$ integrin and a ligand of 20 human $\alpha 9$ integrin.

In a specific embodiment, said humanized antibody or an antigen-binding fragment thereof that immunospecifically recognizes human $\alpha 9$ integrin comprises: (i) a heavy chain (H-chain) comprising at least one H-chain FR (FRH) derived from a variable region (V-region) of a human H-chain, and at least one H-chain 25 complementarity determining region (CDRH) derived from at least one of the CDRHs of a non-human antibody that immunospecifically recognizes human $\alpha 9$ integrin; or (ii) a light chain (L-chain) comprising at least one L-chain FR (FRL) derived from a V-region of a human L-chain, and at least one L-chain complementarity determining region (CDRL) derived from at least one of the 30 CDRLs of a non-human antibody that immunospecifically recognizes human $\alpha 9$ integrin; or both (i) and (ii) above. For example, said non-human antibody, from which at least one of the CDRHs and/or at least one of the CDRLs of the humanized

antibody of the invention is derived, is a monoclonal antibody produced by a hybridoma selected from the group consisting of Accession Nos. FERM BP-10510, FERM BP-10511, FERM BP-10512, FERM BP-10513 and FERM BP-10832.

In a preferred specific embodiment, the humanized antibody or an

5 antigen-binding fragment thereof, of the present invention comprises: (i) at least one FRH derived from a human FRH, and at least one CDRH comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:4, 5 and 6; or (ii) at least one FRL derived from a human FRL, and at least one CDRL comprising an amino acid sequence selected from the group consisting of the

10 amino acid sequences of SEQ ID NOS:11, 12 and 13; or (iii) both (i) and (ii) above. Said humanized antibody or an antigen-binding fragment thereof, of the present invention may comprise CDRH1, CDRH2 and CDRH3, which comprise the amino acid sequences of SEQ ID NOS:4, 5 and 6, respectively. In the alternative, said humanized antibody or an antigen-binding fragment thereof, of the present invention

15 comprises CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS: 11, 12 and 13, respectively. In a preferred embodiment, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and CDRL3, which

20 comprise the amino acid sequences of SEQ ID NOS:4, 5, 6, 11, 12 and 13, respectively. In another alternative, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises a FRH derived from a variable region of a human H-chain encoded by GenBank Accession No. X65891 (SEQ ID NO:18), or a FRL derived from a variable region of a human κ-L-chain encoded by GenBank Accession No. X72441 (SEQ ID NO:23). In a preferred embodiment, the

25 FRH of the humanized antibody of the present invention comprises at least one amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:19, 20, 21 and 22 (FRH1, FRH2, FRH3 and FRH4, respectively, encoded by the corresponding portions of X65891). In another preferred embodiment, the FRL of the humanized antibody of the present invention comprises

30 at least one amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:24, 25, 26 and 27 (FRL1, FRL2, FRL3 and FRL4, respectively, encoded by the corresponding portions of X72441). In a most preferred embodiment, the humanized antibody or an antigen-binding fragment

thereof, of the present invention comprises: (i) a H-chain variable region (VH region) comprising the amino acid sequence of SEQ ID NO:29; or (ii) a L-chain variable region (VL region) comprising the amino acid sequence of SEQ ID NO:31; or (iii) both (i) and (ii) above.

5 In another preferred specific embodiment, the humanized antibody or an antigen-binding fragment thereof, of the present invention comprises: (i) at least one FRH derived from a human FRH, and at least one CDRH comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:32, 33 and 34; or (ii) at least one FRL derived from a human FRL, and
10 at least one CDRL comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:37, 38 and 39; or (iii) both (i) and (ii) above. Said humanized antibody or an antigen-binding fragment thereof, of the present invention may comprise CDRH1, CDRH2 and CDRH3, which comprise the amino acid sequences of SEQ ID NOS:32, 33 and 34, respectively. In
15 the alternative, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS:37, 38 and 39, respectively. In a preferred embodiment, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and
20 CDRL3, which comprise the amino acid sequences of SEQ ID NOS:32, 33, 34, 37, 38 and 39, respectively.

 In another preferred specific embodiment, the humanized antibody or an antigen-binding fragment thereof, of the present invention comprises: (i) at least one FRH derived from a human FRH, and at least one CDRH comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:42, 43 and 44; or (ii) at least one FRL derived from a human FRL, and at least one CDRL comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:47, 48 and 49; or (iii) both (i) and (ii) above. Said humanized antibody or an antigen-binding fragment thereof, of the present invention may comprise CDRH1, CDRH2 and CDRH3, which comprise the amino acid sequences of SEQ ID NOS:42, 43 and 44, respectively. In the alternative, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRL1, CDRL2 and CDRL3, which comprise the

amino acid sequences of SEQ ID NOS:47, 48 and 49, respectively. In a preferred embodiment, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS: 42, 43, 44, 47, 5 48 and 49, respectively.

In another preferred specific embodiment, the humanized antibody or an antigen-binding fragment thereof, of the present invention comprises: (i) at least one FRH derived from a human FRH, and at least one CDRH comprising an amino acid sequence selected from the group consisting of the amino acid sequences of 10 SEQ ID NOS:52, 53 and 54; or (ii) at least one FRL derived from a human FRL, and at least one CDRL comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:57, 58 and 59; or (iii) both (i) and (ii) above. Said humanized antibody or an antigen-binding fragment thereof, of the present invention may comprise CDRH1, CDRH2 and CDRH3, which 15 comprise the amino acid sequences of SEQ ID NOS:52, 53 and 54, respectively. In the alternative, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS:57, 58 and 59, respectively. In a preferred embodiment, said humanized antibody or an antigen-binding fragment thereof, of the 20 present invention comprises CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS:52, 53, 54, 57, 58 and 59, respectively.

In another preferred specific embodiment, the humanized antibody or an antigen-binding fragment thereof, of the present invention comprises: (i) at least 25 one FRH derived from a human FRH, and at least one CDRH comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:62, 63 and 64; or (ii) at least one FRL derived from a human FRL, and at least one CDRL comprising an amino acid sequence selected from the group consisting of the amino acid sequences of SEQ ID NOS:67, 68 and 69; or (iii) both (i) and (ii) above. Said humanized antibody or an antigen-binding fragment thereof, of the present invention may comprise CDRH1, CDRH2 and CDRH3, which 30 comprise the amino acid sequences of SEQ ID NOS:62, 63 and 64, respectively. In the alternative, said humanized antibody or an antigen-binding fragment thereof, of

the present invention comprises CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS:67, 68 and 69, respectively. In a preferred embodiment, said humanized antibody or an antigen-binding fragment thereof, of the present invention comprises CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and CDRL3, which comprise the amino acid sequences of SEQ ID NOS:62, 63, 64, 67, 68 and 69, respectively.

5 The present invention further provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding the humanized antibody or an antigen-binding fragment thereof of the present invention which immunospecifically
10 recognizes human $\alpha 9$ integrin. Specifically, the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a humanized H-chain comprising at least one amino acid sequence selected from the group consisting of SEQ ID NOS:4, 5, 6, 32, 33, 34, 42, 43, 44, 52, 53, 54, 62, 63 and 64, or a humanized L-chain comprising at least one amino acid sequence
15 selected from the group consisting of SEQ ID NOS:11, 12, 13, 37, 38, 39, 47, 48, 49, 57, 58, 59, 67, 68 and 69, or both said humanized H-chain and said humanized L-chain. In a preferred specific embodiment, such an isolated nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO:28, which encodes a VH region, or a nucleotide sequence encoding the amino acid sequence of SEQ ID
20 NO:29. In another preferred specific embodiment, such an isolated nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO:30, which encodes a VL region, or a nucleotide sequence encoding the amino acid sequence of SEQ ID NO:31. In yet another preferred specific embodiment, the isolated nucleic acid molecule of the present invention comprises the nucleotide sequences of both SEQ
25 ID NO:28 and 30. In yet another preferred specific embodiment, the isolated nucleic acid molecule of the present invention further comprises a nucleotide sequence encoding a signal peptide of donor origin, such as the amino acid sequences of SEQ ID NOS:10 and 17, respectively, or of heterologous origin.

30 The present invention further provides a vector, *e.g.*, an expression vector, comprising a nucleotide sequence encoding a H-chain or a L-chain, or both, of the humanized antibody or an antigen-binding fragment thereof of the present invention that immunospecifically recognizes human $\alpha 9$ integrin. In such a vector, the nucleotide sequence of the present invention may be operably linked to one or more

regulatory elements. The nucleotide sequence of the present invention may include a nucleotide sequence encoding a signal peptide native to a non-human donor antibody from which a CDR is derived, or a signal peptide of heterologous origin.

Furthermore, the present invention provides a host cell comprising the nucleic acid molecule of the present invention, including a vector comprising the nucleic acid molecule of the present invention. In one embodiment, the present invention provides an isolated host cell comprising a first nucleic acid molecule encoding a humanized H-chain of the present invention and a second nucleic acid molecule encoding a humanized L-chain of the present invention, said first and second nucleic acid molecules are each operably linked to a regulatory element in such a way that the biologically functional humanized antibody or antigen-binding fragment thereof of the present invention is expressed.

Accordingly, the present invention further provides a method for preparing the humanized antibody of the present invention, comprising culturing the host cell of the invention under conditions so that the humanized antibody is expressed; and collecting the produced humanized antibody.

The present invention further provides a composition comprising at least one of the humanized antibodies of the present invention. In addition, the present invention provides a pharmaceutical composition for preventing or treating a disorder or disease that is associated with $\alpha 9$ integrin, comprising at least one of the humanized antibodies of the present invention, and a pharmaceutically acceptable carrier. Either of said compositions can further comprise another active compound that can additively or synergistically ameliorate the disorder or disease. Such an active compound includes, but not by way of limitation, anti-inflammatory compounds, chemotherapeutic compounds, and the like, as well as an antibody or an antigen-binding fragment thereof, such as an antibody that can immunospecifically bind human $\alpha 4$ integrin.

In another aspect, the present invention provides a method for preventing or treating a disorder or disease that is associated with or involves $\alpha 9$ integrin, said method comprising administering a prophylactically or therapeutically effective amount of at least one of the humanized antibodies of the present invention to a subject in need thereof. For such uses, the humanized antibody of the present

invention may be conjugated to a therapeutic moiety that enhances the biological effect of the humanized antibody. Examples of such a therapeutic moiety include another antibody, such as anti- α 4 antibody (e.g., to form a bispecific antibody), cytotoxins that are cytostatic or cytoidal, radioactive elements, and/or other 5 therapeutic agents, including anti-inflammatory agents, antibiotics, and the like.

In yet another aspect, the present invention provides a method for diagnosing a disorder or disease, in a subject, that is associated with or involves α 9 integrin, said method comprising administering a diagnostically effective amount of the humanized antibody of the present invention to a subject to be examined. For such diagnostic 10 uses, the humanized antibody of the present invention may be labeled with detectable markers, such as radioactive elements.

3.1. Definitions

As used herein, the term "antibody" refers to an antibody molecule capable of immunospecifically binding to a desired antigen, such as the α 9 integrin, and 15 encompasses an antibody molecule as a whole or a fragment thereof, including an antigen-binding fragment.

The term "immunospecifically recognize" used herein refers to an ability of an antibody or an antigen-binding fragment thereof to bind specifically to a target polypeptide or protein, in particular, human α 9 integrin. Such an antibody does not 20 non-specifically bind to other polypeptides or proteins. However, an antibody or an antigen-binding fragment thereof that immunospecifically binds to the target polypeptide or protein (e.g., human α 9 integrin) may cross-react with other antigens. For example, the humanized antibody or an antigen-binding fragment of the present invention that immunospecifically recognizes human α 9 integrin may cross-react 25 with, for example, murine α 9 integrin. Preferably, an antibody or an antigen-binding fragment thereof that immunospecifically binds to human α 9 integrin does not cross-react with other antigens.

The term "an antigen-binding fragment" used herein refers to any fragment of an antibody that retains an ability to immunospecifically bind to a target polypeptide 30 or protein, in particular, human α 9 integrin and/or mouse α 9 integrin, and includes single chain antibodies, Fab fragments, $F(ab')_2$ fragments, disulfide-linked Fvs, and fragments containing either a variable region of a light chain (VL) and/or a variable

region of a heavy chain (VH) or even a complementary determining region (CDR) that specifically binds to a target polypeptide or protein. Thus, such antigen-binding fragments of humanized antibody may or may not include partial or full-length human constant regions. Various methods for obtaining the antibody fragments described above are well known in the art.

5 The term “derived from a human source” or “derived from a non-human source” used herein refers to an antibody portion whose amino acid sequence is derived from a corresponding portion of a human antibody or of a non-human antibody.

10 The term “an acceptor sequence” used herein refers to a nucleotide sequence or an amino acid sequence of framework regions from a human antibody VH or VL region that serves as an acceptor for CDRs from a donor antibody, which is usually a non-human antibody.

The present invention also provides a humanized antibody or an antigen-binding fragment thereof that immunospecifically recognizes human $\alpha 9$ integrin, comprising:

15 a H-chain comprising the amino acid sequence of SEQ ID NO:29, and at least one CDRH derived from at least one of CDRHs of a non-human antibody that immunospecifically recognizes human $\alpha 9$ integrin; and

20 a L-chain comprising the amino acid sequence of SEQ ID NO:31, and at least one CDRL derived from at least one of CDRLs of a non-human antibody that immunospecifically recognizes human $\alpha 9$ integrin.

The present invention also provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence of SEQ ID NO:29.

25 The present invention also provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence of SEQ ID NO:31.

4. BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows the nucleotide sequence (SEQ ID NO:7) of mouse 24I11 VH cDNA along with the deduced amino acid sequence (SEQ ID NO:8). Amino acid

residues are shown in single letter code. The signal peptide sequence (SEQ ID NO:10) is in italic. The N-terminal amino acid residue (E) of the mature VH is double-underlined. CDR sequences according to the definition of Kabat *et al.* (Sequences of Proteins of Immunological Interests, Fifth edition, NIH Publication No. 91-3242, U.S.

5 Department of Health and Human Services, 1991) are underlined.

Fig. 2 shows the nucleotide sequence (SEQ ID NO:14) of mouse 24I11 VL cDNA along with the deduced amino acid sequence (SEQ ID NO:15). Amino acid residues are shown in single letter code. The signal peptide sequence (SEQ ID NO:17) is in italic. The N-terminal amino acid residue (D) of the mature VL is double-underlined. CDR sequences according to the definition of Kabat et al. (1991) are 10 underlined.

Fig. 3 shows the nucleotide sequence (SEQ ID NO:72) of the designed 24I11 VH gene flanked by SpeI and HindIII sites (underlined), along with the deduced amino acid sequence (SEQ ID NO:8). Amino acid residues are shown in single letter code.

15 The signal peptide sequence (SEQ ID NO:10) is in italic. The

N-terminal amino acid residue (E) of the mature VH is double-underlined. CDR sequences according to the definition of Kabat et al. (1991) are underlined. The intron sequence is in italic.

Fig. 4 shows the nucleotide sequence (SEQ ID NO:73) of the designed 24I11 VL gene flanked by NheI and EcoRI sites (underlined), along with the deduced amino acid sequence (SEQ ID NO:15). Amino acid residues are shown in single letter code. The signal peptide sequence (SEQ ID NO:17) is in italic. The N-terminal amino acid residue (D) of the mature VL is double-underlined. CDR sequences according to the definition of Kabat et al. (1991) are underlined. The 10 intron sequence is in italic.

Fig. 5 shows the schematic structure of pCh24I11 and pHu24I11 (collectively Expression Vector). Proceeding clockwise from the Sall site at the top, the plasmid contains the heavy chain transcription unit starting with the human cytomegalovirus (CMV) major immediate early promoter and enhancer (CMV promoter) to initiate 15 transcription of the antibody heavy chain gene. The CMV promoter is followed by the VH exon, a genomic sequence containing the human gamma-1 heavy chain constant region including the CH1, hinge, CH2 and CH3 exons with the intervening introns, and a polyadenylation site of the gamma-1 gene for mRNA processing following CH3. After the heavy chain gene sequence, the light chain transcription 20 unit begins with the CMV promoter, followed by the VL exon and a genomic sequence containing the human kappa chain constant region exon (CL) with part of the intron preceding it, and a poly A signal of the kappa gene. The light chain gene is then followed by the SV40 early promoter (SV40 promoter), the *E. coli* xanthine 25 guanine phosphoribosyl transferase gene (gpt), and a segment containing the SV40 polyadenylation site (SV40 poly(A) site). Finally, the plasmid contains a part of the plasmid pUC19, comprising the bacterial origin of replication (pUC ori) and beta-lactamase gene (beta lactamase).

Fig. 6 shows the alignment of the amino acid sequences of 24I11 VH (SEQ ID NO:9), humanized 24I11 (Hu24I11) VH (SEQ ID NO:29) and FRH1 (SEQ ID 30 NO:19), FRH2 (SEQ ID NO:20), FRH3 (SEQ ID NO:21) and FRH4 (SEQ ID NO:22) of human acceptor sequences, derived from the amino acid sequence encoded by the nucleotide sequence of GenBank Accession No. X65891. Amino acid residues are shown in single letter code. Numbers above the sequences

indicate the positions according to Kabat *et al.* (1991). CDR sequences defined by Kabat *et al.* (1991) are underlined. Double-underlined residues were predicted to contact with the CDRs and the mouse residues were retained at these locations in the humanized form. CDR residues in X65891 are omitted in the figure.

5 **Fig. 7** shows the alignment of the amino acid sequences of 24I11 VL (SEQ ID NO:16), humanized 24I11 (Hu24I11) VL (SEQ ID NO:31) and FRL1 (SEQ ID NO:24), FRL2 (SEQ ID NO:25), FRL3 (SEQ ID NO:26) and FRL4 (SEQ ID NO:27) of human acceptor sequences, derived from the amino acid sequence encoded by the nucleotide sequence of GenBank Accession No. X72441. Amino acid residues are
10 shown in single letter code. Numbers above the sequences indicate the positions according to Kabat *et al.* (1991). CDR sequences defined by Kabat *et al.* (1991) are underlined. Double-underlined residues were predicted to contact with the CDRs and the mouse residues were retained at these locations in the humanized form. CDR residues in X72441 are omitted in the figure.

15 **Fig. 8** shows the oligonucleotides used for construction of the Hu24I11 VH gene.

Fig. 9 shows the oligonucleotides used for construction of the Hu24I11 VL gene.

20 **Fig. 10** shows the oligonucleotides used for construction of the Hu24I11 VH gene flanked by SpeI and HindIII sites (SEQ ID NO:74 with 5'-GGG tail at 5'-terminal and CCC-3' tail at 3'-terminal). An arrow denotes the position and orientation (5' to 3') of each oligonucleotide. Amino acid residues of the signal peptide (SEQ ID NO:10) and the VH region (SEQ ID NO:29) are shown in single letter code.

25 **Fig. 11** shows oligonucleotides used for construction of the Hu24I11 VL gene flanked by NheI and EcoRI sites (SEQ ID NO:75 with 5'-GGG tail at 5'-terminal and CCC-3' tail at 3'-terminal). An arrow denotes the position and orientation (5' to 3') of each oligonucleotide. Amino acid residues of the signal peptide (SEQ ID NO:17) and the VL region (SEQ ID NO:31) are shown in single letter code.

30 **Fig. 12** shows the nucleotide sequence (SEQ ID NO:74) of the Hu24I11 VH gene flanked by SpeI and HindIII sites (underlined), along with the deduced amino acid sequence of the signal peptide (SEQ ID NO:10; shown in italic) and the VH

region (SEQ ID NO:29). Amino acid residues are shown in single letter code. The N-terminal amino acid residue (Q) of the mature VH is double-underlined. CDR sequences according to the definition of Kabat et al. (1991) are underlined. The intron sequence is in italic.

5 **Fig. 13** shows the nucleotide sequence (SEQ ID NO:75) of the Hu24I11 VL gene flanked by NheI and EcoRI sites (underlined), along with the deduced amino acid sequence of the signal peptide (SEQ ID NO:17; shown in italic) and the VL region (SEQ ID NO:31). Amino acid residues are shown in single letter code. The N-terminal amino acid residue (D) of the mature VL is double-underlined.

10 10 CDR sequences according to the definition of Kabat et al. (1991) are underlined. The intron sequence is in italic.

Fig. 14 shows the comparison of the affinity of chimeric and humanized 24I11 antibodies to human $\alpha 9$ integrin. The binding of chimeric and humanized 24I11 at 1 and 0.5 μ g/ml to CHO/ $\alpha 9$ cells was examined by cell ELISA.

15 15 Experiments were carried out in triplicate. The mean absorbance value with SEM is shown in the figure.

Fig. 15 shows the result of the FACS analysis of the binding of mouse, chimeric and humanized 24I11 antibodies to human $\alpha 9$ integrin. Each antibody was tested at 1, 0.33, 0.11, 0.037 and 0.012 μ g/ml for binding to CHO/hu $\alpha 9$ cells.

20 20 Geometric mean channel fluorescence values (MCF; Y-axis) are plotted at each antibody concentration tested (X-axis) in the figure.

Fig. 16 shows the result of cell adhesion inhibitory activity of anti-human $\alpha 9$ integrin antibodies, between Human $\alpha 9$ /CHO-K1 cells and hOPN(RAA)N-half, Tenascin-C, VCAM-1, or human fibronectin.

25 25 **Fig. 17** shows the result of cell adhesion inhibitory effects of anti-human $\alpha 9$ integrin antibodies on human melanoma cells in the presence of anti-human $\alpha 4$ integrin.

Fig. 18 shows therapeutic effects on hepatitis by the anti- $\alpha 4$ integrin antibodies and the anti- $\alpha 9$ -integrin antibodies. In the Figure, NHG indicates

30 30 normal hamster antibody and NRG indicates normal rat antibody.

Fig. 19 shows that growth of the B16-BL6 cells was inhibited by the anti- $\alpha 9$

integrin antibodies.

Fig. 20 shows the results of FACS analysis using anti-human $\alpha 9$ integrin antibodies for Human $\alpha 9$ /CHO-K1 cells (Fig. 20a), Human $\alpha 4$ /CHO-K1 (Fig. 20b) and human neutrophils (Fig. 20c).

5 **Fig. 21** shows the cell growth inhibition of the B16-BL6 cells by the anti- $\alpha 9$ integrin antibodies using immobilized VCAM-1 as an ECM.

Fig. 22 shows the therapeutic effect of anti- $\alpha 9$ integrin in a mouse rheumatoid arthritis model.

10

5. DETAILED DESCRIPTION OF THE INVENTION

5.1. Preparation of Antibodies Against Human $\alpha 9$ Integrin

Antibodies that immunospecifically recognize human $\alpha 9$ integrin or any epitope thereof may be generated by any suitable method known in the art.

15 The $\alpha 9$ integrin used as an antigen in the present invention may be (1) proteins derived from all cells from human that express $\alpha 9$ integrin, or all tissues where these cells are present, (2) recombinant proteins in which the $\alpha 9$ integrin-encoding gene DNA, preferably cDNA, is transfected into bacteria, yeast, cell lines including animal cells, etc. and expressed, or (3) synthetic proteins.

20 The $\alpha 9$ integrin includes polypeptides comprising substantially the same amino acid sequences as the amino acid sequences of human $\alpha 9$ integrins (SEQ ID NO:76, wherein 1-29 residues are the signal peptide).

25 Herein, the term "polypeptides comprising substantially the same amino acid sequence" means variant polypeptides comprising an amino acid sequence, in which multiple amino acids, preferably 1 to 10 amino acids and more preferably 1 to several (e.g., 1 to 5) amino acids are substituted, deleted and/or modified, as long as these variant polypeptides have biological properties substantially equivalent to the naturally occurring human $\alpha 9$ integrin; and variant polypeptides comprising an amino acid sequence, wherein multiple amino acids, preferably 1 to 10 amino acids and more preferably 1 to several (e.g., 1 to 5) amino acids are added to the amino acid sequence of naturally occurring human $\alpha 9$ integrin. Furthermore, the variant

polypeptides may be those having a plurality of these substitutions, deletions, modifications and additions of amino acids.

The human $\alpha 9$ integrin as an antigen in the present invention can be produced by methods well known in the art, such as chemical synthesis method, cell culture method, etc., or their modifications, in addition to the gene recombinant techniques.

Examples of the methods for producing variant polypeptides include a synthetic oligonucleotide site-directed mutagenesis (gapped duplex method), a point mutagenesis method which involves introducing a point mutation at random by treatment with nitrite or sulfite, a method which involves preparing a deletion mutant with Bal31 enzyme, or other enzymes, a cassette mutagenesis, a linker scanning method, a miss incorporation method, a mismatch primer method, a DNA segment synthesis method, and the like.

The human $\alpha 9$ integrin to be used as an antigen in the present invention also includes a "part" of said $\alpha 9$ integrin. As used herein, the "part" refers to a part comprising a region required for binding to a ligand of the $\alpha 9$ integrin, for example, OPN, VCAM-1, tenascin-C, etc.; specifically, a part comprising the 14th-980th amino acid residues, and a part comprising the 11th-981st amino acid residues, of the mature human $\alpha 9$ integrin (the 30th-1035th amino acid residues of SEQ ID NO:76). The "part" of said $\alpha 9$ integrin can be produced by gene recombination or chemical synthesis according to methods known in the art described below, or modifications thereof, or can be produced by appropriately digesting the human $\alpha 9$ integrin isolated by the cell culture method with a proteolytic enzyme or the like.

As an antigen, a cell per se that overexpresses the $\alpha 9$ integrin on the cell membrane, or a membrane fraction thereof, can be also used. Cells overexpressing human $\alpha 9$ integrin can be prepared by recombinant DNA technologies well known in the art.

Using appropriate antigens prepared as described above, antibodies specific for human $\alpha 9$ integrin or any epitope thereof may be prepared by various methods well known in the art. Polyclonal antibodies to human $\alpha 9$ integrin can be produced by various procedures well known in the art. For example, an antigen of interest can be administered to various host animals including, but not limited to, rabbits,

mice, rats, etc., to induce the production of antisera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete) adjuvant, mineral gels such as

5 aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful adjuvants for humans such as BCG (Bacille Calmette-Guerin) and Corynebacterium parvum. Such adjuvants are also well known in the art.

10 Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring

15 Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas*, pp. 563-681 (Elsevier, N.Y., 1981) (both of which are incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived

20 from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art. In a non-limiting example, mice can be immunized with an antigen of interest or a cell expressing such an antigen.

25 Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells (e.g., P3U1, P3X63-Ag8, P3X63-Ag8-U1, P3NS1-Ag4, SP2/0-Ag14, P3X63-Ag8-653, etc.). Hybridomas are selected and cloned by limiting dilution.

30 The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding the antigen. Ascites fluid, which generally contains high levels of antibodies, can be generated by inoculating mice intraperitoneally with positive hybridoma clones.

Antibody fragments what recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the complete light chain, and the variable region, the CH1 region and the hinge region of the heavy chain.

5 The antibodies of the invention or an antigen-binding fragment thereof can be also produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression 10 techniques.

The nucleotide sequence encoding an antibody may be obtained from any information available to those skilled in the art (i.e., from Genbank, the literature, or by routine cloning and sequence analysis). If a clone containing a nucleic acid encoding a particular antibody or an epitope-binding fragment thereof is not 15 available, but the sequence of the antibody molecule or epitope-binding fragment thereof is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from any tissue or cells expressing the antibody, such as hybridoma cells selected to 20 express an antibody) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in 25 the art.

5.2. Preparation of Recombinant Antibodies

Once the nucleotide sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site 30 directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., *supra*; and Ausubel et al., eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in

their entireties), to generate antibodies having a different amino acid sequence by, for example, introducing amino acid substitutions, deletions, and/or insertions into the epitope-binding domain regions of the antibodies or any portion of antibodies which may enhance or reduce biological activities of the antibodies.

5 Recombinant expression of an antibody requires construction of an expression vector containing a nucleotide sequence that encodes the antibody. Once a nucleotide sequence encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using
10 techniques well known in the art as discussed in the previous sections. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic
15 recombination. The nucleotide sequence encoding the heavy-chain variable region, light-chain variable region, both the heavy-chain and light-chain variable regions, an epitope-binding fragment of the heavy- and/or light-chain variable region, or one or more complementarity determining regions (CDRs) of an antibody may be cloned into such a vector for expression. Such a sequence may be fused with a
20 polynucleotide encoding a signal peptide native to the original antibody or a heterologous signal peptide. Thus-prepared expression vector can be then introduced into appropriate host cells for the expression of the antibody. Accordingly, the invention includes host cells containing a polynucleotide encoding a humanized antibody or an antigen-binding fragment thereof that immunospecifically
25 recognizes human $\alpha 9$ integrin.

30 The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides or different selectable markers to ensure maintenance of both plasmids. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

In another embodiment, antibodies can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage 5 can be utilized to display antigen binding domains, such as Fab and Fv or disulfide-bond stabilized Fv, expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used 10 in these methods are typically filamentous phage, including fd and M13. The antigen binding domains are expressed as a recombinantly fused protein to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the immunoglobulins, or fragments thereof, of the present invention include those disclosed in Brinkman et al., *J. Immunol. Methods*, 182:41-50, 1995; 15 Ames et al., *J. Immunol. Methods*, 184:177-186, 1995; Kettleborough et al., *Eur. J. Immunol.*, 24:952-958, 1994; Persic et al., *Gene*, 187:9-18, 1997; Burton et al., *Advances in Immunology*, 57:191-280, 1994; PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 20 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, 25 including human antibodies, or any other desired fragments, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')2 fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., 30 BioTechniques, 12(6):864-869, 1992; and Sawai et al., AJRI, 34:26-34, 1995; and Better et al., *Science*, 240:1041-1043, 1988 (each of which is incorporated by reference in its entirety). Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patent Nos.

4,946,778 and 5,258,498; Huston et al., *Methods in Enzymology*, 203:46-88, 1991; Shu et al., *PNAS*, 90:7995-7999, 1993; and Skerra et al., *Science*, 240:1038-1040, 1988.

Once an antibody molecule of the invention has been produced by any

5 methods described above, it may then be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A or Protein G purification, and sizing column chromatography), centrifugation, differential solubility, or by any other standard techniques for the purification of

10 proteins. Further, the antibodies of the present invention or fragments thereof may be fused to heterologous polypeptide sequences described herein or otherwise known in the art to facilitate purification.

For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human

15 antibodies. Chimeric antibodies and humanized antibodies are discussed in details in Section 5.3, *infra*.

Antibodies fused or conjugated to other compounds or heterologous polypeptides may be used in *in vitro* immunoassays, in purification methods (e.g., affinity chromatography), as well as *in vivo* therapeutic or diagnostic uses. See e.g.,

20 PCT publication Number WO 93/21232; EP 439,095; Naramura et al., *Immunol. Lett.*, 39:91-99, 1994; U.S. Patent 5,474,981; Gillies et al., *PNAS*, 89:1428-1432, 1992; and Fell et al., *J. Immunol.*, 146:2446-2452, 1991, which are incorporated herein by reference in their entireties. For example, antibodies can be labeled in various ways using a known method or commercially available kit (e.g., biotin

25 labeling, FITC labeling, APC labeling). As another example, antibodies may be conjugated to a therapeutic moiety that enhances the biological effect of the antibodies *in vivo*. Examples of such a therapeutic moiety include another antibody, cytotoxins that are cytostatic or cytoidal, radioactive element, and/or other therapeutic agents, including anti-inflammatory agents, antibiotics, and the like. In

30 the present invention, the humanized anti-human α 9 integrin may be conjugated to another antibody, such as anti- α 4 antibody (e.g., to form a bispecific antibody). As another example, the humanized antibody of the present invention may be labeled with detectable markers, such as radioactive elements, for *in vivo* diagnostic uses.

5.3. Chimeric and Humanized Antibodies

A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a constant region

5 derived from a human immunoglobulin. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, *Science*, 229:1202, 1985; Oi et al., *BioTechniques*, 4:214 1986; Gillies et al., *J. Immunol. Methods*, 125:191-202, 1989; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entireties.

10 A humanized antibody is a molecule that binds a desired antigen and comprises a variable region containing one or more complementarity determining regions (CDRs) derived from a non-human species and one or more framework regions derived from a human immunoglobulin molecule. The typical methods for humanizing non-human antibodies have been described in various references, such as
15 those: by Queen et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:10029-10033 and U.S. Patent Nos. 5,585,089 and 5,693,762; by Riechmann et al., *Nature*, 332:323, 1988; and by Tsurushita et al., *Methods* 36:69-83, 2005, all of which are incorporated herein by reference in their entireties). For example, the reference by Tsurushita et al. (2005, *supra*; hereafter “*Tsurushita*”) provides a practical and instructive protocol
20 for the humanization of mouse monoclonal antibodies based on the antibody-humanization method originally developed by Queen et al. (1989, *supra*). The general protocol disclosed in *Tsurushita* is briefly summarized below.

5.3.1. General protocol for preparing humanized antibodies

Cloning and sequencing of mouse V genes

25 Various methods are available for cloning cDNAs encoding the VH and VL regions of a target mouse monoclonal antibody. For example, 5' RACE (rapid amplification of cDNA ends) method using SMART RACE cDNA Amplification Kit (BD Biosciences, CA) or the GeneRacer Kit (Invitrogen, CA) has been commonly used. A gene-specific primer for 5' RACE can be prepared based on the isotypes of
30 the H-chain and the L-chain of the target monoclonal antibody so that it can bind immediately downstream of the variable region for each of the H-chain and L-chain. Thus, 5' RACE primer may be designed to be specific for each subtype in mouse,

such as $\gamma 1$, $\gamma 2a$, $\gamma 2b$ or $\gamma 3$. Alternatively, a common primer for all subtypes may be designed based on the consensus or highly homologous region among the subtypes.

In *Tsurushita*, the following 5' RACE primers are disclosed as examples:

5 (i) 5'-GCCAGTGGATAGACTGATGG- (SEQ ID NO:129) (for cloning of mouse $\gamma 1$, $\gamma 2a$, $\gamma 2b$ and $\gamma 3$ H-chains)

15 (ii) 5'-GATGGATACAGTTGGTGCAGC- (SEQ ID NO:130) (for cloning of mouse κ light chains).

PCR-amplified V gene fragments can be directly cloned into a plasmid vector, for example, using the Zero Blunt TOPO PCR Cloning Kit (Invitrogen), and 10 their DNA sequences determined. The obtained sequences should be confirmed by, for example, comparing their encoding amino acid sequences with those of the target monoclonal antibody determined by the N-terminal amino acid sequencing, using, for example a Model 241 Protein Sequencer (Hewlett-Packard, CA). Typically, the determination of at least 15-20 amino acid residues at the N-terminus of the target 15 antibody, for example, by Edman degradation, is sufficient to confirm the authenticity of the cloned DNA sequences. *Tsurushita* cautions that when glutamine, which is one of the two most common N-terminal amino acid in mouse, is the N-terminal amino acid, it might have been converted to pyroglutamine and blocks the sequencing at the N-terminus. In that case, it is necessary to deblock the 20 N-terminus to obtain the sequence.

Three-dimensional modeling of V regions

Based on the sequences of the VH and VL regions, the framework residues of the target antibody that are potentially important for maintaining the conformational structure of the CDRs, are first identified by the method, for 25 example, described by R. Levy *et al.*, 1989, *Biochemistry* 28:7168-7175; and by B. Zilber *et al.*, 1990, *Biochemistry* 29:10032-10041. Typically, each of the VH and VL regions is divided into 14 structurally meaningful segments, which are β strands and loop-like structures comprising the domain structure of the immunoglobulin 30 superfamily. The amino acid sequence of each of the segments from the target antibody is aligned with the corresponding segments of antibodies of known structures, in the PDB database (see H.M. Berman *et al.*, 2000, *Nucleic Acids Res.* 28:235-342). By multiple sequence alignment, a corresponding segment having the

highest sequence homology to each of the target segment is selected and the three-dimensional model of the V-region is constructed. In order to optimize the structure, the model is subjected to multiple cycles of conjugate gradient energy minimization (e.g., using ENCAD, or as described by Press *et al.*, 1990, in

5 "Numerical Recipes, Cambridge University Press, Cambridge; AMBER by Weiner *et al.*, 1981, *J. Comp. Chem.* 2:287-303; 3D-JIG-SAW available at BioMolecularModelling or "BMM" web site run by Cancer Research UK; or SWISS-MODEL available at ExPASy Proteomics Server web site run by Swiss Institute of Bioinformatics, Geneva).

10 **Selection of human frameworks**

In parallel with modeling the structure of the V regions, the amino acid sequences deduced from the cDNA cloning of the mouse VH and VL regions, respectively, are compared to human V region sequences in the databases, for example, the Kabat database (see Johnson *et al.*, 2000, *Nucleic Acids Res.*

15 28:214-218.), GenBank, and so forth. Human framework regions that have overall sequence identity of at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, or at least 95% identity, with the mouse sequence, can be searched using, for example, the Smith-Waterman algorithm (by Gusfield, 1997, in "Algorithms on Strings, Trees, and Sequences",
20 Cambridge University Press, Cambridge), or BLAST (by Karlin *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87:2264-2268), and the like. These human sequences may be based on cDNA-based and protein-derived sequences; however, the use of germline is often preferable as it may be useful in eliminating potential immunogenicity associated with somatic hypermutations in cDNA-based, protein-derived sequences.
25 In the alternative, as described in Queen *et al.* (1989, *supra*), the use of a consensus framework sequence can also identify and remove such hypermutated residues in the framework obtained from cDNA-based or protein-derived sequences. In the case where a germline VH segment is used as an acceptor framework, VH segments encoded on chromosome 14, rather than 15 and 16, should be used as only those on
30 chromosome 14 produce functional VH regions.

Design of humanized V regions

According to Queen *et al.* (1989, *supra*), it is necessary to identify framework

amino acids within about 4-6 Å of the CDRs as these residues are considered to be potential key framework residues that support the correct CDR structures. Such a process can be achieved using a computer program, such as RASMOL available at Molecular Visualization Freeware web site supported by National Science

5 Foundation (NSF), that calculates interatomic distances from the atomic coordinates or, through manual inspection of a computer model. If amino acids at key framework positions are different between mouse donor and human acceptor sequences, those of mouse donor usually replace the human residues. However, if such residues have minimal contribution to support the CDR structures, the

10 corresponding human residues are typically used. Also, if the selected human acceptor contains “atypical” amino acids, which occur in less than about 10-20% of the V region sequences, they may be the result of somatic hypermutation during affinity maturation and should be replaced with the donor residues in order to avoid potential immunogenicity in humans.

15 In addition, other factors, such as residues of potential N-linked glycosylation signals, need to be carefully considered in order to design humanized V regions (*see Tsurushita* for details).

Humanized antibodies may contain a human constant region or a portion thereof from the human κ or λ light chain, and/or the γ1, γ2, γ3, γ4, μ, α1, α2, δ, or ε

20 heavy chain of human antibodies, or variants thereof, depending on the effector functions required or to be eliminated for therapeutic uses. For example, a Fc portion of the constant region containing a mutation may be fused to the variable region of the chimeric or humanized antibody of the present invention so as to reduce the binding of the antibody to Fc receptors and/or to reduce its ability to fix

25 complement (*see*, for example, Winter *et al.*, GB 2,209,757 B; Morrison *et al.*, WO 89/07142, Morgan *et al.*, WO 94/29351). Such manipulations of antibody molecules can be carried out by recombinant DNA technology as described in Section 5.2.

30 Preferably the resulting chimeric or humanized antibody has the same specificity as the non-human donor antibody and an affinity similar to or at least about 1/3, at least about 1/2, or at least about 2/3, of that of the non-human donor antibody. In another aspect, the resulting chimeric or humanized antibody has an affinity constant of at least about $1 \times 10^7 \text{ M}^{-1}$, preferably at least about $1 \times 10^8 \text{ M}^{-1}$,

and most preferably at least about $1 \times 10^9 \text{ M}^{-1}$.

In addition to the general protocol described above, antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101 and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology, 28(4/5):489-498, 1991; Studnicka et al., Protein Engineering, 7(6):805-814, 1994; Roguska et al., Proc Natl. Acad. Sci. USA, 91:969-973, 1994), and chain shuffling (U.S. Patent No. 5,565,332), all of which are hereby incorporated by reference in their entireties.

10 **5.3.2. Additional considerations for preparing humanized antibodies as pharmaceuticals**

To offer humanized antibodies as pharmaceuticals, an efficient and consistent production system therefor needs to be prepared. For example, an appropriate expression vector for humanized antibodies is prepared by inserting H- and L-chain 15 sequences, and a high-productivity cell line transfected with the expression vector can be obtained as a seed cell for a master cell bank (MCB), which serves as a stable and semi-permanent source for a working cell bank (WCB). Humanized antibodies can be then prepared by culturing working cells from the WCB and collecting the culture medium.

20 Various expression vectors with appropriate regulatory genes can be used for the preparation of such a production cell line. As a host cell, those commonly used for expressing mammalian proteins can be used for the expression of humanized antibodies. Examples of such host cells include, but are not limited to, Chinese Hamster Ovary (CHO) cells, SP2/0-Ag14.19 cells, NSO cells, and the like. The 25 productivity of humanized antibodies can be maximized by selecting the best combination of an expression vector and a host cell. Furthermore, the composition of culture media should be explored in order to select suitable media, from various serum-free culture media and supplements, so that the expression of humanized antibodies by the host cell can be optimized.

30 Based on the efficiency and the final yield, the humanized antibodies produced by the host cell can be purified from the culture supernatant using various methods well known in the art, including affinity chromatography, ion-exchange

chromatography, hydrophobic interaction chromatography, and the like.

5.4. Pharmaceutical Composition and Therapeutic Uses

The present invention provides a pharmaceutical composition comprising the humanized antibody or an antigen-binding fragment thereof, described above, that

5 immunospecifically recognizes human $\alpha 9$ integrin. The pharmaceutical composition comprising the humanized antibody of the present invention as an active ingredient can be used as an agent for preventing and/or treating a disorder or disease that is associated with $\alpha 9$ integrin, including, but not limited to, cancer, *e.g.*, the growth or metastasis of cancer cells, and an inflammatory disease, *e.g.*, rheumatoid

10 arthritis, osteoarthritis, hepatitis, bronchial asthma, fibrosis, diabetes mellitus, arteriosclerosis, multiple sclerosis, granuloma, an inflammatory bowel disease (ulcerative colitis and Crohn's disease), an autoimmune disease, and the like.

The pharmaceutical composition comprising the humanized antibody of the present invention can also be used to treat chronic rejection after organ

15 transplantation, and an autoimmune disease such as systemic autoimmune disease, erythematosus, uveitis, Behcet's disease, polymyositis, glomerular proliferative nephritis, sarcoidosis, and the like.

The preventive and/or therapeutic agent for preventing or treating the disorders or diseases described above, comprising the humanized antibody of the

20 present invention, has low toxicity and can be administered to humans orally or parenterally, directly as a liquid preparation by mixing in a suitable solvent, or as a pharmaceutical composition in an appropriate dosage form.

The pharmaceutical composition used for the administration described above contains the aforesaid antibody or salts thereof and pharmaceutically acceptable

25 carriers, diluents or excipients. Such a composition is provided in a dosage form suitable for oral or parenteral administration.

The dose may vary depending upon the age and the size of a subject to be administered, target disease, conditions, route of administration, and the like. When the antibody is used for preventing and/or treating, for example, rheumatoid arthritis

30 in an adult patient, it is advantageous to intravenously administer the antibody of the present invention normally at a single dose of about 0.01 to about 20 mg/kg body weight, preferably about 0.1 to about 10 mg/kg body weight, and more preferably

about 0.1 to about 5 mg/kg body weight, approximately 1 to 5 times per day, preferably approximately 1 to 3 times per day. In other parenteral administration and oral administration, the antibody can be administered in a dose corresponding to the dose given above. When the condition is especially severe, the dose may be 5 increased according to the condition.

Various delivery systems are known and can be used to administer the pharmaceutical composition of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the mutant viruses, receptor mediated endocytosis (see, e.g., Wu and Wu, 1987, *J. Biol. Chem.* 10 262:4429 4432). Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may 15 be administered together with other biologically active agents. Administration can be systemic or local. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; 20 this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, by means of nasal spray, or by means of an implant, said implant being of a porous, non porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. 25 In one embodiment, administration can be by direct injection at the site (or former site) infected tissues.

In another embodiment, the pharmaceutical composition can be delivered in a vesicle, in particular a liposome (see Langer, 1990, *Science* 249:1527-1533; Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez Berestein 30 and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.* , pp. 317-327; see generally *ibid.*).

In yet another embodiment, the pharmaceutical composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see

Langer, *supra*; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:201; Buchwald et al., 1980, *Surgery* 88:507; and Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); 5 Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., 1985, *Science* 228:190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105). In yet another embodiment, a controlled release system can be placed in proximity of 10 the composition's target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)). Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

Examples of the composition for oral administration include solid or liquid 15 dosage forms, specifically, tablets (including dragees and film-coated tablets), pills, granules, powdery preparations, capsules (including soft capsules), syrup, emulsions, suspensions, etc. Such a composition is manufactured by publicly known methods and contains a vehicle, a diluent or an excipient conventionally used in the field of pharmaceutical preparations. Examples of the vehicle or excipient for tablets are 20 lactose, starch, sucrose, magnesium stearate, and the like.

The injectable preparations may include dosage forms for intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, etc. These injectable preparations may be prepared by methods publicly known. The injectable preparations may be prepared, e.g., by dissolving, suspending or 25 emulsifying the antibody or its salt described above in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), 30 a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl

benzoate, benzyl alcohol, etc. The injection thus prepared is preferably filled in an appropriate ampoule. The suppository used for rectal administration may be prepared by blending the aforesaid antibody or its salt with conventional bases for suppositories.

5 Advantageously, the pharmaceutical compositions for oral or parenteral use described above are prepared into dosage forms in a unit dose suited to fit a dose of the active ingredients. Such dosage forms in a unit dose include, for example, tablets, pills, capsules, injections (ampoules), suppositories, etc. The amount of the aforesaid antibody contained is generally about 5 to 500 mg per dosage form in a unit
10 dose; especially in the form of injection, it is preferred that the aforesaid antibody is contained in about 5 to 100 mg and in about 10 to 250 mg for the other dosage forms.

Each composition described above may further contain other active components unless formulation causes any adverse interaction with the antibodies described above.

15 The present invention also relates to an inhibitor and/or promoter for cell and/or tissue remodeling, which comprises an $\alpha 9$ integrin-binding functional molecule (e.g., OPN, VCAM-1, tenascin-C, fibronectin, pp-vWF, tTG, etc.) as an active ingredient; and a method for inhibiting and/or promoting cell and/or tissue remodeling, which comprises contacting the $\alpha 9$ integrin-expressing cell and/or tissue
20 (e.g., a tumor cell, neutrophil, smooth muscle, etc.) with the $\alpha 9$ integrin-binding functional molecule. The dose, method for administration, pharmaceutical preparation, etc. of the active ingredient in such a therapeutic agent can be appropriately determined by referring to the foregoing description of medicaments comprising the humanized antibodies of the present invention.

25 As described above, the present invention further provides a method for preventing or treating a disorder or disease that is associated with or involves $\alpha 9$ integrin, said method comprising administering an effective amount of at least one of the humanized antibodies of the present invention to a subject in need thereof.

5.5. Diagnostic Uses

30 The pharmaceutical composition comprising the humanized antibody of the present invention can be used as a diagnostic agent for cancer, e.g., the growth or

metastasis of cancer cells, and an inflammatory disease, e.g., rheumatoid arthritis, osteoarthritis, hepatitis, bronchial asthma, fibrosis, diabetes mellitus, cancer metastasis, arteriosclerosis, multiple sclerosis, granuloma, etc., or as a diagnostic agent for chronic rejection after organ transplantation, an autoimmune disease such as systemic autoimmune disease, erythematosus, uveitis, Behcet's disease, polymyositis, glomerular proliferative nephritis, sarcoidosis, and so forth. The humanized antibodies of the present invention are capable of specifically recognizing the $\alpha 9$ integrin and hence can be used to quantify the $\alpha 9$ integrin in a test fluid, especially for quantification by the sandwich immunoassay, competitive assay, immunometry, nephrometry, etc., immunostaining, or the like. In applying these immunological methods to the assay methods of the present invention, it is not required to set forth any particular conditions, procedures, etc. It is sufficient to construct assay systems by adding ordinary technical consideration in the art to conventional conditions and procedures. For details of these general technical means, reference can be made to reviews, texts or the like.

As described above, the $\alpha 9$ integrin can be quantified with high sensitivity by using the antibodies of the present invention. The humanized antibodies of the present inventions are particularly useful for diagnosing various diseases associated with the $\alpha 9$ integrin by applying the method for quantifying the $\alpha 9$ integrin *in vivo*.
20 For instance, where an increase or decrease in the expression level of the $\alpha 9$ integrin is detected, it can be diagnosed that it is highly likely that one now suffers from diseases associated with the $\alpha 9$ integrin, e.g., cancer or an inflammatory disease, or it is highly likely that one will suffer from these diseases in the future. Thus, the present invention also provides a method for diagnosing a disorder or disease associated with or involve $\alpha 9$ integrin in a subject, said method comprising administering an effective amount of at least one of the humanized antibodies of the present invention or both to a subject in need thereof. Required dosages for such an *in vivo* diagnosis may be less than those required for therapeutic uses and can be determined by one skilled in the art according to routine procedures.

30 The humanized antibodies of the present invention can also be used for specifically detecting the $\alpha 9$ integrin present in a test fluid such as a body fluid, a tissue, etc. The humanized antibodies can also be used for preparation of antibody columns for purification of the $\alpha 9$ integrin, for detection of the $\alpha 9$ integrin contained

in each fraction upon purification or for analysis of behaviors of the $\alpha 9$ integrin in cells to be tested.

6. EXAMPLES

5 The following examples illustrate the preparation of monoclonal antibodies that immunospecifically recognize human and/or mouse $\alpha 9$ integrin, the sequencing of the variable regions of the monoclonal antibodies, the epitope mapping and other characterization of the antibodies and the chimerization and the humanization of such antibodies, as well as the characterization of the resulting chimeric and humanized 10 antibodies. These examples should not be construed as limiting.

6.1. Preparation of Mouse Antibody Against Human $\alpha 9$ Integrin

Mouse monoclonal antibodies against human $\alpha 9$ integrin were prepared according to the subtractive immunization method (by Williams C.V., *et al.*, 1992, *Biotechniques* 12:842-847). Briefly, three Balb/c mice were injected 15 intraperitoneally with CHO-K1 cells at 4×10^6 per mouse. In the following two days, the mice received 4 mg/mouse of cyclophosphamide intraperitoneally. At two weeks after the cyclophosphamide injection, the mice were injected intraperitoneally with 2×10^6 cells/mouse of CHO-K1 cells expressing human $\alpha 9$ integrin (Human $\alpha 9$ /CHO-K1 cells), followed by another intraperitoneal injection of the same cells at 20 3×10^6 cell/mouse two weeks later. Hybridomas were prepared by the methods well known in the art (see, for example, Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, *et al.*, in: *Monoclonal Antibodies and T-Cell Hybridomas*, pp. 563-681 (Elsevier, N.Y., 1981). Hybridoma clones producing monoclonal antibodies that were immunospecifically 25 reactive with Human $\alpha 9$ /CHO-K1 cells but not with CHO K1 cells expressing human $\alpha 4$ integrin were established and five hybridoma clones (*i.e.*, 1K11, 21C5, 24I11, 25B6 and 28S1) producing monoclonal antibodies immunospecifically recognizing human $\alpha 9$ integrin were isolated.

6.2. Epitope Analysis for Anti-Human $\alpha 9$ Integrin Monoclonal Antibodies

30 Twelve-residue polypeptides, starting from N-terminal of human $\alpha 9$ integrin

and every three residues thereafter (*i.e.*, amino acid residues 1-12, 4-15, 7-18, and so forth) were prepared and coupled to a cellulose membrane via C6-spacer and 2 β Ala residues at 5 nmol/spot. The membrane was blocked with a blocking buffer (milk/0.05% Tween20 in PBS) and reacted with 10 ml of solution containing 1.0 5 μ g/ml of each of the anti-human α 9 integrin monoclonal antibodies (*i.e.*, 1K11, 21C5, 24I11 and 25 B6, respectively) labeled with peroxidase for three hours at room temperature. After washing with T-TBS, the membrane was reacted with enhanced 10 chemiluminescence (ECL) detecting reagent for 1 minute at room temperature. The luminescence emitted as a result of the enzymatic reaction is measured and epitopes 15 of the antibodies were determined based on the luminescence intensities. As a control, Y9A2 (see Wang *et al.*, 1996, *Am J Respir Cell Mol Biol* 15, 664-672), a commercially available monoclonal antibody against human α 9 integrin, was used.

Table 1 below shows the result of the epitope mapping, which indicated that the monoclonal antibodies isolated by the present inventors have epitopes that are 15 distinct from that of Y9A2.

Table 1

Human $\alpha 9$ integrin sequences	1K11	21C5	24I11	25B6	Y9A2
FQGPADSFFGYA (SEQ ID NO:77)	-	-	-	++	-
KSPGAVFKCRVHTNPDRR (SEQ ID NO:78)	++	+++	++++	-	+
WMGVSLARQPKADGRVLA (SEQ ID NO:79)	+	+++	+	-	+
CAHRWKNIYYEADHI (SEQ ID NO:80)	+	+	+++	+	+
GFCYIIPSNLQAKGRTLI (SEQ ID NO:81)	+++	+++++	+++++	+++++	+++++
VMGAPGSFYWAGTIKVNL (SEQ ID NO:82)	+	+	+	+++	+
VIMNRRYTYLGYAVT (SEQ ID NO:83)	+++++	++	+++++	+++++	+++
VYIFRADRRSGTLIKIFQ (SEQ ID NO:84)	++++	+++++	+++	-	+
QYSMKLSGQKINPVLRMFGQSISG (SEQ ID NO:85)	+	++++	-	-	+
VVLLRARPVITVDVSIFL (SEQ ID NO:86)	++	++	-	-	+++++
RHYVAHVKRQDVISPI (SEQ ID NO:87)	+++	+++	+++++	+	++
ELPPLTPVLRWKKGQKIAQKNQTVFERNCR (SEQ ID NO:88)	+	+++++	+	-	++
YLALGAVKNISL (SEQ ID NO:89)	+	+	-	++	+++++
CSVGFPFMRSKSKYEFPSV (SEQ ID NO:90)	+	++	++	-	+++
SSSVIQFMSRAKVKVDPALRV (SEQ ID NO:91)	+	++	+++	-	+

6.3. CDR Analysis of Anti-Human $\alpha 9$ Integrin Antibodies

The amino acid sequences of CDRs of the monoclonal antibodies (*i.e.*, 1K11, 21C5, 24I11, 25B6 and 28S1) were determined by reverse transcription of the mRNA extracted from the corresponding hybridomas to prepare cDNAs. Using the cDNAs as templates, the variable regions of the H-chains and L-chains were extended and amplified by PCR using ScFv-cloning primers (Light Primer Mix and Heavy Primer Mix; by Amersham Biosciences Corp., IL). The PCR products were cloned into pCRII TOPO vector, sequenced and the amino acid sequences were determined. This process was repeated three times for each antibody. The results are shown in Table 2.

Table 2

CDRs	1K11	21C5	24I11	25B6	28S1
CDRH1	DYNNMD (SEQ ID NO:32)	DYYMY (SEQ ID NO:42)	DTYVH (SEQ ID NO:4)	SYGVH (SEQ ID NO:32)	GYGVN (SEQ ID NO:62)
CDRH2	DINPNNGGTYNQKEFQG (SEQ ID NO:33)	TISDGGGNYTYPDSVKG (SEQ ID NO:43)	NDPANGNTKYDPKFQG (SEQ ID NO:5)	VTWSSGGSTYNTNSALMS (SEQ ID NO:53)	MWGDGITEYNSALKS (SEQ ID NO:63)
CDRH3	SGVISTDY (SEQ ID NO:34)	DRDGSSLFAY (SEQ ID NO:44)	WLRHFYYAMDY (SEQ ID NO:6)	DYGNYPWFAY (SEQ ID NO:54)	RDASSGYGFA (SEQ ID NO:64)
CDRL1	RASQEISGYLI (SEQ ID NO:37)	KASQDVNIAVA (SEQ ID NO:47)	RASENTYSLA (SEQ ID NO:11)	KASQDVNTAVA (SEQ ID NO:57)	TASSVSSSYLH (SEQ ID NO:67)
CDRL2	AASLILDS (SEQ ID NO:38)	WASTRHT (SEQ ID NO:48)	NANSLED (SEQ ID NO:12)	SASYRYT (SEQ ID NO:58)	STSNLAS (SEQ ID NO:68)
CDRL3	LQYANTPP1 (SEQ ID NO:39)	QQHYNTPW (SEQ ID NO:49)	KQAYDVPYT (SEQ ID NO:13)	QQHYSTPCA (SEQ ID NO:59)	HQYHRSPTYT (SEQ ID NO:69)

6.4. Cell Adhesion Inhibitory Activity

(1) Since it is known that cell adhesion involves the binding of $\alpha 9$ integrin to

its ligands, *i.e.*, various ECMs, including OPN, fibronectin, Tenascin-C, VCAM-1, and the like, the isolated anti-human α 9 integrin antibodies were examined for their cell adhesion inhibitory activity.

Briefly, hOPN(RAA)N-half was prepared as a Glutathione S-transferase (GST)-fusion protein by isolating from *E. coli* host cells an N-terminal portion down to the thrombin-cleavage site of OPN, in which the GRD sequence has been replaced with the RAA sequence, and cleaving the GST portion with Precision protease (Amersham Biosciences). VCAM-1 was purchased from R&D Systems, Inc. (Minneapolis, MN). Tenascin-C and human fibronectin were prepared by synthesizing polypeptides containing AEIDGIEL (SEQ ID NO:92); the α 9 integrin-binding region of Tenascin-C) and CPEDGIHEFP (SEQ ID NO:93); the α 9 integrin-binding region of human fibronectin), respectively, and subsequently attaching them to bovine serum albumin (BSA). For human α 9 integrin, CHO-K1 cells that abundantly expressed human α 9 integrin (Human α 9/CHO-K1) were used.

15 Fifty microliters of Tenascin-C, fibronectin, VCAM-1 or hOPN(RAA)N-half were added to a 96-well plate at 1.25-5.0 μ g/ml and incubated at 37°C for 1 hour to coat the plate. After blocking the plate with a blocking solution (0.5% BSA/PBS) and washing it with PBS once, the mixture of human α 9/CHO-K1 cells (1.0×10^5 cells/ml) and the isolated monoclonal antibodies (10 g/ml) in 0.25% BSA-Minimum 20 Essential Media (MEM) was added to the plate at 200 μ l/well and incubated at 37°C for 1 hour under 5% CO₂. Non-adherent cells were rinsed off with PBS and adherent cells were fixed and stained with 0.5% Crystal Violet (by WAKO, Osaka, Japan)/20% methanol. The stained cells were allowed to stand at room temperature for 30 minutes and 20% acetic acid solution was added thereto to effect dissolution.

25 The adhesion activity was quantified by measuring OD at 590 nm wavelength.

As shown in Fig. 16, the cell adhesion involving Tenascin-C was inhibited by 21C5, 24I11, 25B6 and 28S1, but not by 1K11. The cell adhesion involving fibronectin was inhibited by 21C5, 25B6 and 28S1 and by 24I11 to a less degree, but no inhibition was observed with 1K11. The cell adhesion involving VICAM-1 was inhibited by 21C5, 24I11, 25B6 and 28S1, but not by 1K11. Likewise, the cell adhesion involving hOPN(RAA)N-half was inhibited by 21C5, 24I11, 25B6 and 28S1, but not by 1K11.

(2) Since $\alpha 4$ integrin and $\alpha 9$ integrin have many common ECM ligands, the presence of both anti- $\alpha 4$ integrin and anti- $\alpha 9$ integrin antibodies is expected to enhance the cell adhesion inhibitory activity. Thus, the effect of the both types of antibodies in combination was examined *in vitro* for a possible inhibitory effect on 5 metastatic cancer, based on the cell adhesion between human melanoma cells (G361) expressing $\alpha 4$ integrin as well as $\alpha 9$ integrin and VCAM-1 (1.25 μ g/ml) as an ECM; a rat monoclonal antibody, P1H4 (Cat. No. MAB16983Z, Chemicon International Inc., CA) was used as an anti-human $\alpha 4$ integrin antibody.

As shown in Fig. 17, the adhesion involving VCAM-1 was not inhibited by 10 any of the anti-human $\alpha 9$ integrin antibodies alone, but was inhibited by the positive control (Y9A2), 21C5 and 24I11, in the co-presence of the anti-human $\alpha 4$ integrin antibody. Since cells expressing many $\alpha 9$ integrin molecules usually express $\alpha 4$ integrin molecules also, this result indicates that the inhibition of cell adhesion can 15 be effectively achieved by the combination use of anti-human $\alpha 9$ integrin antibody and anti-human $\alpha 4$ integrin antibody, thereby enhancing the suppression of various disorders and diseases, including metastatic cancer involving these integrin molecules.

6.5. Use of Anti-Human $\alpha 9$ Integrin Antibodies in FACS Analysis

Whether the anti-human $\alpha 9$ integrin antibodies were usable for FACS was 20 examined using Human $\alpha 9$ /CHO-K1 cells, CHO-K1 cells and human neutrophils endogenously expressing the $\alpha 9$ integrin. In human neutrophils, FACS analysis was conducted at the cell count of 1.0×10^5 and antibodies were reacted on ice. Non-specific reaction with the Fc receptor was blocked with 50% goat serum. FITC-labeled anti-mouse IgG antibody was used as a secondary antibody. As a 25 result (see Figs. 20a, 20b and 20c), all of the anti-human $\alpha 9$ integrin antibodies could detect the $\alpha 9$ integrin on Human $\alpha 9$ /CHO-K1 and human neutrophils. None of the antibodies reacted with Human $\alpha 4$ /CHO-K1 cells (see Figs. 20a, 20b and 20c). These results revealed that all of the anti-human $\alpha 9$ integrin antibodies could detect the human $\alpha 9$ integrin proteins expressed on cells using FACS.

6.6. Therapeutic Effects of Anti- α 9 Integrin Antibody

Therapeutic effects of anti- α 9 integrin antibodies were examined in a mouse system.

The anti-mouse α 9 integrin monoclonal antibodies (11L2B, 12C4'58, 5 18R18D and 55A2C) were prepared substantially in the same manner as described for mouse anti-human α 9 integrin antibodies (see Section 6.1, *supra*), except that hamsters were immunized with CHO-K1 cells expressing mouse α 9 integrin (mouse α 9/CHO-K1 cells) and the resulting monoclonal antibodies that reacted with mouse α 9/NIH3T3 cells but not with mouse α 4/NIH3T3 were selected.

10 6.6.1. Therapeutic effect on hepatitis

WO 02/081522 discloses that hepatitis can be treated by inhibiting the OPN functions. Accordingly, therapeutic effects of anti- α 9 integrin antibody was studied in a mouse hepatitis model using a hamster anti-mouse α 9 integrin antibody, 11L2B, and a rat anti-mouse α 4 integrin antibody, R1-2 (Pharmingen). The blood AST and 15 ALT levels in the mice were measured using GPT/ALT-PIII and GOT/AST-PIII (Fuji Film), 12 hours after an intravenous injection of 200 μ g of concanavalin A (Con A) (Vector). Three hours before the Con A injection, 200 μ g of the antibody were administered. As shown in Fig. 18, the AST and ALT levels were found to be decreased by the anti- α 9 integrin antibody, and the therapeutic effects could be 20 noted. In addition, the therapeutic effects could be boosted by concomitant use with the anti- α 4 integrin antibody. The results revealed that hepatitis could be treated by the anti- α 9 integrin antibody.

6.6.2. Effect of anti- α 9 integrin antibodies on growth of mouse cancer cell 25 line

Murine melanoma cell line B16-BL6 expresses abundant α 9 integrin. Accordingly, cell growth inhibitory activities of the established anti-mouse α 9 integrin antibodies against cancer cells were examined.

The B16-BL6 cells were prepared on a 96-well plate for cell culture (Becton Dickinson) at 5×10^4 cells/mL in 10% FCS/DMEM. After 10 μ g/ml of the 30 anti-mouse α 9 integrin antibody and anti-mouse α 4 integrin antibody were added,

100 µL each of the cell-antibody suspension was added to each well. Incubation was conducted at 37°C for 24 hours under 5% CO₂, and 10 µL each of Cell Counting Kit 8 (Dojin Kagaku Kenkyu-sho) was added, followed by incubation at 37°C for an hour under 5% CO₂. Absorbance at O.D. 450 was measured and the cell count was 5 quantitatively analyzed. As shown in Fig. 19, 12C4'58 gave the highest inhibitory activity and inhibited the growth of B16-BL6 cells by about 35%. Both 55A2C and R1-2 could inhibit the growth by about 20%.

Next, for analysis of inhibitory effects against cell growth under conditions closer to the in vivo conditions, VCAM-1 was immobilized on a solid phase and 10 assayed in a similar fashion. VCAM-1 is a ligand for $\alpha 9$ integrin and a recombinant soluble form of VCAM-1 protein, rhVCAM-1-Fc chimera (Roche), was used. Using the rhVCAM-1-Fc chimera immobilized on a solid phase with 10 µg/mL, non-specific reaction was blocked with 0.5% BSA/PBS. The chimera was added in a concentration of 10 µg/ml in single use of the antibody, and in 5 µg 15 each/ml in concomitant use. Thereafter, the same procedures as in Fig. 19 were followed. As a result, the effect was not obtained at all or only an imperceptible effect was obtained by single administration of 12C4'58 and by single use of the $\alpha 4$ inhibitory antibody clone R1-2, whereas in simultaneous administration of 12C4'58 with R1-2 the cell growth inhibitory effect showed a marked increase by about 20%, 20 as shown in Fig. 21.

6.6.3. Therapeutic effect of anti- $\alpha 9$ integrin in mouse rheumatoid arthritis model

Seven-week old female mice (Balb/c) (3 mice per group) were injected intraperitoneally with the hamster anti-mouse $\alpha 9$ integrin antibody (55A2C), or 25 normal hamster IgG (NHG) at 400 µg/mouse. After 24 hours, 2 mg/mouse of the arthritis-inducing cocktail of type II collagen-specific monoclonal antibody (Chondrex Inc.) was injected intravenously. After 72 hours, 400 µg/mouse of 55A2C or NHG as well as 50 µg/mouse of LPS were injected intraperitoneally. The mice were observed from 3 days before the LPS injection until 6 days after the LPS 30 injection and the levels of arthritis were scored according to the method by Wood *et al.* (1969, *Int. Arch. Allergy Appl. Immunol.* 35:456). The result is shown in Fig. 22. The mice injected with control NHG had a high score and developed rheumatoid

arthritis, whereas in those injected with anti-mouse $\alpha 9$ integrin antibody, the development of rheumatoid arthritis was completely blocked. Thus, anti- $\alpha 9$ integrin antibodies were indicated to have prophylactic and therapeutic effects on rheumatoid arthritis.

5 6.7. Humanization of Non-Human Antibodies

6.7.1. Cloning and sequencing of mouse 24I11 V genes

Mouse 24I11 hybridoma cells were grown in TIL Media I (Immuno-Biological Laboratories, Gunma, Japan) containing 10% fetal bovine serum (FBS; HyClone, Logan, UT) at 37°C in a 7.5% CO₂ incubator. Total RNA 10 was extracted from approximately 3 x 10⁶ hybridoma cells using TRIzol reagent (Invitrogen, Carlsbad, CA) according to the supplier's protocol. Oligo dT-primed cDNA was synthesized using the GeneRacer Kit (Invitrogen) following the supplier's protocol. The variable region cDNAs for 24I11 heavy and light chains were 15 amplified by polymerase chain reaction (PCR) with Phusion DNA polymerase (New England Biolabs, Beverly, MA) using primers that anneal respectively to the mouse gamma-1 and kappa chain constant regions, and a GeneRacer 5' primer (5'-CGACTGGAGCACGAGGACACTGA-) (SEQ ID NO:94) provided in the GeneRacer Kit. For PCR amplification of heavy chain variable region (VH), the primer has the sequence 5'-GCCAGTGGATAGACAGATGG- (SEQ ID NO:95). 20 For PCR amplification of light chain variable region (VL), the primer has the sequence 5'-GATGGATACAGTTGGTGCAGC- (SEQ ID NO:96). The amplified VH and VL cDNAs were subcloned into the pCR4Blunt-TOPO vector (Invitrogen) for sequence determination. DNA sequencing of the variable regions was carried 25 out at Tocore (Menlo Park, CA). Several heavy and light chain clones were sequenced and unique sequences homologous to typical mouse heavy and light chain variable regions were identified. The consensus cDNA sequences along with deduced amino acid sequences of 24I11 VH and VL are shown in Figs. 1 and 2, respectively.

6.7.2. Construction of chimeric 24I11 IgG1/k antibody

30 A gene encoding 24I11 VH was generated as an exon including a splice donor signal and appropriate flanking restriction enzyme sites by PCR using 24I11 VH

cDNA as a template,
5'-GGGACTAGTACCACCATGAAATGCAGCTGGTTATCTTC- (SEQ ID NO:97) (SpeI site is underlined) as a 5' primer, and
5'-GGGAAGCTTAGAGGCCATTCTTACCTGAGGAGACGGTGA
5 C- (SEQ ID NO:98) (HindIII site is underlined) as a primer (Fig. 3). Likewise, a gene encoding 24I11 VL was generated as an exon including a splice donor signal and appropriate flanking restriction enzyme sites by PCR using 24I11 VL cDNA as a template, 5'-GGGGCTAGACCACCATGAGTGTGCCACTCAACTCCTG- (SEQ ID NO:99) (NheI site is underlined) as a 5' primer, and 5'-
10 GGGGAATTCTGAGAAGACTACTTACGTTTATTCCAGCTGGTCCCCC- (SEQ ID NO:100) (EcoRI site is underlined) as a primer (Fig. 4). The splice donor signals of the 24I11 VH and VL exons were derived from the mouse germline JH4 and Jκ2 sequences, respectively. PCR-amplified fragments were gel-purified using QIAquick Gel Extraction Kit (Qiagen, Valencia, CA), digested with SpeI and
15 HindIII (for VH) or NheI and EcoRI (for VL), and cloned into a mammalian expression vector carrying human gamma-1 and kappa constant regions for production of chimeric 24I11 IgG1/κ antibody. The schematic structure of the resulting expression vector, pCh24I11, is shown in Fig. 5.

6.7.3. Generation of humanized 24I11 V genes

20 Humanization of the 24I11 variable regions was carried out as outlined by Queen et al. (Proc. Natl. Acad. Sci. USA 86: 10029-10033, 1989). First, a molecular model of the 24I11 variable regions was constructed with the aid of computer programs. Next, based on a homology search against human variable region sequences, the human amino acid sequence encoded by the nucleotide sequence of GenBank Accession No. X65891, which has a high homology [72.4% (63/87) amino acid identity in the FRHs] to 24I11 VH, was chosen as an acceptor to provide the frameworks for humanized 24I11 VH. Likewise, the human amino acid sequence encoded by the nucleotide sequence of GenBank Accession No. X72441 [77.5% (62/80) amino acid identity in the FRLs] was chosen as an acceptor for
25 humanization of 24I11 VL.
30

At framework positions where the computer model suggested significant contact with the complementarity determining regions (CDRs), the amino acids from the 24I11 variable regions were substituted for the human framework amino acids.

This was done at positions 27, 28, 29, 30, 48, 66, 67 and 71 (according to Kabat numbering system; *see, Kabat et al., Sequences of Proteins of Immunological Interests, Fifth edition, NIH Publication No. 91-3242, U.S. Department of Health and Human Services, 1991*) to generate humanized 24I11 (Hu24I11) VH (Fig. 6). For 5 the light chain, replacements were made at residues 70 and 71 to generate humanized 24I11 (Hu24I11) VL (Fig. 7). The alignments of 24I11, designed Hu24I11, and the human acceptor amino acid sequence are shown for VH in Fig. 6 and for VL in Fig. 7.

A gene encoding each of Hu24I11 VH and VL was designed as an exon 10 including a signal peptide, a splice donor signal, and appropriate restriction enzyme sites for subsequent cloning into a mammalian expression vector. The splice donor signals of the Hu24I11 VH and VL exons were derived from the human germline JH4 and Jκ1 sequences, respectively. The signal peptide sequences in the Hu24I11 VH and VL exons were derived from the corresponding mouse 24I11 VH and VL 15 sequences, respectively. The Hu24I11 VH and VL genes were constructed by extension and PCR amplification of several overlapping synthetic oligonucleotide primers using ThermalAce DNA polymerase (Invitrogen) as outlined by He et al. (J. Immunol. 160: 1029-1035, 1998). The oligonucleotides used for construction of Hu24I11 VH and VL genes are listed in Figs. 8 and 9, respectively. The location of 20 the oligonucleotides in the Hu24I11 VH and VL genes is shown in Figs. 10 and 11, respectively. PCR-amplified fragments were gel-purified using QIAquick Gel Extraction Kit (Qiagen) and cloned into pCR4Blunt-TOPO vector for sequence determination. After digestion with SpeI and HindIII (for VH) or NheI and EcoRI (for VL), Hu24I11 VH and VL genes were subcloned into corresponding sites in a 25 mammalian expression vector for production in the human IgG1/κ form. The schematic structure of the resulting expression vector, pHu24I11, is shown in Fig. 5. The nucleotide sequences of the obtained Hu24I11 VH and VL genes along with deduced amino acid sequences are shown in Figs. 12 and 13, respectively.

6.7.4. Transient expression of chimeric and humanized 24I11 IgG1/κ

30 Chimeric and humanized 24I11 IgG1/κ antibodies were transiently expressed by transfecting pCh24I11 and pHu24I11 plasmid DNA, respectively, to HEK293 cells using polyethylenimine according to Durocher et al. (Nucl. Acids Res. 30: e9,

2002). Transiently transfected HEK293 cells were maintained for four days in DMEM containing 10% FBS at 37°C in a 7.5% CO₂ incubator. The expression level of each of Ch24I11 and Hu24I11 IgG1/κ antibodies in culture supernatant was measured by sandwich ELISA. An ELISA plate was coated overnight at 4°C with 5 100 μl/well of 1/2,000-diluted goat anti-human IgG Fcγ-chain-specific polyclonal antibodies (SouthernBiotech, Birmingham, AL) in PBS, washed with Wash Buffer (PBS containing 0.05% Tween 20), and blocked for 1 hr at room temperature with 300 μl/well of Blocking Buffer (PBS containing 2% Skim Milk and 0.05% Tween 20). After washing with Wash Buffer, 100 μl/well of samples appropriately diluted 10 in ELISA Buffer (PBS containing 1% Skim Milk and 0.025% Tween 20) were applied to the ELISA plate. Human IgG1/κ antibody purified from human myeloma serum (SouthernBiotech) was used as a standard. After incubating the ELISA plate for 2 hr at room temperature and washing with Wash Buffer, bound antibodies were detected using 100 μl/well of 1/2,000-diluted horse radish 15 peroxidase (HRP)-conjugated goat anti-human kappa chain polyclonal antibodies (SouthernBiotech). After incubating for 1 hr at room temperature and washing with Wash Buffer, color development was performed by adding 100 μl/well of ABTS substrate (bioWORLD, Dublin, OH). Color development was stopped by adding 100 μl/well of 2% oxalic acid. Absorbance was read at 405 nm.

20 **6.7.5. Characterization of humanized 24I11**

Binding of chimeric and humanized 24I11 antibodies to human α9 integrin was examined by cell ELISA. CHO-K1 stable transfectants expressing recombinant human α9 integrin on the surface (CHO/huα9; provided by Gene Techno Science) were seeded at 2 x 10⁵ cells/well in 50 μl of F12/DMEM (HyClone) 25 containing 10% FBS in a 96-well tissue culture plate and grown overnight at 37°C in a 7.5% CO₂ incubator. For testing of binding to human α9 integrin, 50 μl of chimeric 24I11, humanized 24I11 or irrelevant human IgG1/κ myeloma antibody (SouthernBiotech) in F12/DMEM containing 10% FBS was added to each well. After incubating for 1 hr at 4°C and washing cells twice with ice-cold PBS, 100 μl of 30 1/1,000-diluted HRP-conjugated goat anti-human IgG polyclonal antibodies (SouthernBiotech) was added to each well. After incubating for 1 hr at 4°C, cells were washed three times with ice-cold PBS. For color development, 100 μl of

ABTS substrate was added. Color development was stopped by adding 100 μ l of 2% oxalic acid. Absorbance was read at 405 nm. The result showed that the binding of chimeric 24I11 antibody to human α 9 integrin was almost same as that of humanized 24I11 antibody at both 0.5 and 1 μ g/ml (Fig. 14).

5 Antigen binding of mouse, chimeric and humanized 24I11 monoclonal antibodies was also examined in a FACS binding assay using CHO/hu α 9 cells. Purified mouse 24I11 monoclonal antibody was provided by Gene Techno Sciences. Approximately 8×10^5 CHO/hu α 9 cells/test were washed with FACS Binding Buffer (PBS containing 0.5% BSA and 0.05% NaN₃) and suspended in 200 μ l of FACS

10 10 Binding Buffer containing various amounts of test antibody. After 30 min on ice, the cells were washed twice with FACS Binding Buffer. The cells stained with mouse 24I11 were then suspended in 200 μ l of 1/200-diluted FITC-labeled goat anti-mouse IgG polyclonal antibody (SouthernBiotech) in FACS Binding Buffer. The cells stained with chimeric or humanized 24I11 were suspended in 200 μ l of

15 15 1/200-diluted FITC-labeled goat anti-human IgG polyclonal antibody (SouthernBiotech) in FACS Binding Buffer. After 30 min on ice, the cells were washed with FACS Binding Buffer, suspended in 200 μ l of FACS Binding Buffer, and analyzed using a FACSCan flow cytometer (BD Biosciences, Franklin Lakes, NJ). The binding of chimeric and humanized 24I11 antibodies to CHO/hu α 9 cells

20 20 was very similar to each other in this analysis (Fig. 15).

The results of the cell ELISA and FACS experiments using transiently expressed antibodies suggest that humanization of mouse 24I11 antibody is successful.

Humanization of the other mouse anti-human α 9 antibodies disclosed herein

25 25 (i.e., 1K11, 21C5, 25B6 and 28S1) can be also carried out by employing the same procedure described herein. The DNA sequences and the amino acid sequences of the VH and VL regions, respectively, of these mouse monoclonal antibodies are summarized below.

Mouse monoclonal antibodies	DNA sequence of VH region (SEQ ID NO:) ¹	Deduced amino acid sequence of mature VH (SEQ ID NO:) ²	DNA sequence of VL region (SEQ ID NO:) ¹	Deduced amino acid sequence of mature VL (SEQ ID NO:)
1K11	35	36	40	41
21C5	45	46	50	51
25B6	55	56	60	61
28S1	65	66	70	71

¹ The V genes of each antibody were cloned by a method using Amersham's degenerate primers.

² The deduced amino acid sequence starts from the 2nd residue of VH region (according to the Kabat numbering system) for each clone.

DEPOSITION

The hybridomas designated herein as 1K11, 21C5, 24I11, 25B6 and 28S1 producing mouse anti-human α 9 integrin monoclonal antibodies were deposited on 10 February 15, 2006 with International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology, located at AIST Tsukuba Central 6, 1-1, Higashi 1-chome Tsukuba-shi, Ibaraki-ken 305-8566 Japan in accordance with the Budapest Treaty on the Deposit of Microorganisms, and accorded Accession Nos. FERM BP-10510, FERM BP-10511, FERM BP-10512, FERM BP-10513 and FERM 15 BP-10832, respectively, all of which are incorporated herein by reference in their entirieties.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of 20 any other element, integer or step, or group of elements, integers or steps.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed before the priority 25 date of each claim of this application.

INDUSTRIAL APPLICABILITY

The humanized monoclonal antibodies of the present invention inhibit the function of $\alpha 9$ integrin to exhibit therapeutic effects on cancer, e.g., the growth or metastasis of cancer cells, and an inflammatory disease, e.g., rheumatoid arthritis, 5 osteoarthritis, hepatitis, bronchial asthma, fibrosis, diabetes mellitus, cancer metastasis, arteriosclerosis, multiple sclerosis, granuloma, an inflammatory bowel disease (ulcerative colitis and Crohn's disease), an autoimmune disease, and the like. The pharmaceutical composition comprising both the anti- $\alpha 9$ integrin antibody and anti- $\alpha 4$ integrin antibody of the present invention exerts more improved therapeutic effects on 10 cancer and an inflammatory disease.

9. LIST OF SEQUENCES

The sequences referenced throughout the specification are summarized below.

SEQ ID NO.	TYPE	DESCRIPTION	SEQUENCE
1	AA	OPN adhesion sequence	GRGDS
2	AA	HuOPN's $\alpha 4\beta 1/\alpha 9\beta 1$ -binding site	SVVYGLR
3	AA	MuOPN's $\alpha 4\beta 1/\alpha 9\beta 1$ -binding site	SLAYGLR
4	AA	CDRH1 of 24I11 (FERM BP-10512)	DTYVH
5	AA	CDRH2 of 24I11 (FERM BP-10512)	NIDPANGNTKYDPKFQG
6	AA	CDRH3 of 24I11 (FERM BP-10512)	WLRHFYYAMDY
7	DNA	VH of 24I11 (FERM BP-10512) including sequence encoding signal peptide (1-57)	ATGAAATGCAGCTGGTTATCTTCTCCTGATG GCAGTGGTTACAGGGTCAATTAGAGGTTCA GCTGCAGCAGTCTGGGGCAGAGCTTGTGAAG CCAGGGGCCTCAGTCAGTTGTCCCTGCACAG CTTCTGGCTTCAACATTAAAGACACCTATGTG CACTGGGTGAAGCAGAGGCCTGAACAGGGCC TGGAGTGGATTGGAAATATTGATCCTGCGAAT GGTAATACTAAATATGACCCGAAGTTCCAGGG CAAGGCCACTATAACAGCAGACACATCCTCCA ACACAGCCTACCTGCACCTCAGCAGCCTGACA TCTGAGGACACTGCCGTCTATTACTGTGCTAG ATGGTTACGACATTAACTATGCTATGGACTA CTGGGGTCAAGGAACCTCAGTCACCGTCTCCT CA
8	AA	VH of 24I11 (FERM BP-10512) including signal peptide (1-19)	MKCSWVIFFLMAAVVTGVNSEVQLQQSGAELVKP GASVJKLSCASGFNIKDTYVHWVKQRPEQGLE WIGNIDPANGNTKYDPKFQGKATITADTSNTAY LHLSSLTSEDTAVYYCARWLRHFYYAMDYWGQ GTSVTVSS

9	AA	Mature VH of 24I11 (FERM BP-10512)	EVQLQQSGAELVKPGASVQLSCTASGFNIKDTY VHVKQRPEQGLEWIGNIDPANGNTKYDPKFQ GKATITADTSSNTAYLHLSSLTSEDTAVYYCARW LRHFYYAMDYWGQGTSVTVSS
10	AA	Signal peptide of 24I11 H-chain	MKCSWVIVFLMAVVTGVNS
11	AA	CDRL1 of 24I11 (FERM BP-10512)	RASENIYYSLA
12	AA	CDRL2 of 24I11 (FERM BP-10512)	NANSLED
13	AA	CDRL3 of 24I11 (FERM BP-10512)	KQAYDVPYT
14	DNA	VL of 24I11 (FERM BP-10512) including sequence encoding signal peptide (1-60)	ATGAGTGTGCCACTCAACTCCTGGGGTTGCT GCTGCTGTGGCTTACAGACGCAGGATGTGACA TCCAGATGACTCAGTCTCCAGCCTCCCTGGCT GCATCTGTGGGAGAAACTGTCACCGACATCCA GATGACTCAGTCTCCAGCCTCCCTGGCTGCAT CTGTGGGAGAAACTGTCACCGGAAATTCCT CAGCTCCTGATCTATAATGCAAACAGCTTGA AGATGGTGTCCCATCGAGGTTCAAGTGGCAGTG GATCTGGGACACAGTATTCTATGAAGATCAAC AGCATGCAGCCTGAAGATACCGCAACTTATT CTGTAAACAGGCTTATGACGTTCCGTACACGT TCGGAGGGGGGACCAAGCTGGAAATAAAA
15	AA	VL of 24I11 (FERM BP-10512) including signal peptide (1-20)	MSVPTQLLGLLLWLTDAGCDIQMTQSPASLAAS VGETVTITCRASENIYYSLAWYQQKQGKSPQLL IYNANSLEDGVPSRFSGSGSGTQYSMKINSMQP EDTATYFCKQAYDVPYTGGGTKLEIK
16	AA	Mature VL of 24I11 (FERM BP-10512)	DIQMTQSPASLAASVGETVTITCRASENIYYSLA WYQQKQGKSPQLLIYNANSLEDGVPSRFSGSGS GTQYSMKINSMQPEDTATYFCKQAYDVPYTFG GGTKLEIK
17	AA	Signal peptide of 24I11 L-chain	MSVPTQLLGLLLWLTDAGC
18	DNA	X65891	ATGGACTGGACCTGGAGGGTCTCTTTTGTT GGCAGCAGCCACAGGTGCCACTCCCAGGTC CAGCTTGTGCAGTCTGGGGCTGAGGTGAAGA AGCCTGGGGCCTCAGTGAAGGTTCTGCAGA GGCTTCTGGATAACACCTTCACTAGCTATGCTAT GCATTGGGTGCGCCAGGCCCCGGACAAAGG

			CTTGAGTGGATGGATGGATCAACGCTGGCAA TGGTAACACAAAATATTACAGAAGTTCCAGG GCAGAGTCACCATTACCAGGGACACATCCGCG AGCACAGCCTACATGGAGCTGAGCAGCCTGA GATCTGAAGACACGGCTGTGTATTACTGTGCG AGAATAACCCGTATTAGCAGTGGCTGGTTGGG GGACTACTTGACTACTGGGCCAGGGAACCC TGGTCACCGTCTCCTCA
19	AA	FRH1 of X65891	QVQLVQSGAEVKKPGASVKVSCKASGYTFT
20	AA	FRH2 of X65891	WVRQAPGQRLEWMG
21	AA	FRH3 of X65891	RVTITRDTSASTAYMELSSLRSEDTAVYYCAR
22	AA	FRH4 of X65891	WGQGTLTVSS
23	DNA	X72441	CGCTCAGCTCCTGGGGCTCCTGCTACTCTGGC TCCGAGGTGCCAGATGTGACATCCAGATGACC CAGTCTCCATCCTCCCTGCTGCACTGTAGGA GACAGAGTCACCACACTTGCCGGCAAGTC AGAGCATTAGCAGCTATTAAATTGGTATCAGC AGAAACCAGGGAAAGCCCCTAACGCTCCTGAT CTATGCTGCATCCAGTTGCAAAGTGGGTCC CATCAAGGTTCAGTGGCAGTGGATCTGGGACA GATTCACTCTCACCATCAGCAGTCTGCAACC TGAAGATTTGCAACTACTACTGTCAACAGA GTTACAGTACCCCTCGGACGTTGGCCAAGGG ACCAAGGTGGAAATCAA
24	AA	FRL1 of X72441	DIQMTQSPSSLSASVGDRVTITC
25	AA	FRL2 of X72441	WYQQKPGKAPKLLIY
26	AA	FRL3 of X72441	GVPSRFSGSGSGTDFLTISLQPEDFATYYC
27	AA	FRL4 of X72441	FGQGTKVEIK
28	DNA	VH of Hu24I11	CAGGTTCAGCTGGTGCAGTCTGGGGCAGA GGTGAAGAAGCCAGGGGCCTCAGTCAAGG TTCCCTGCAAGGCTCTGGCTTCACATTA AAGACACCTATGTGCACTGGGTGCGCCAGG CCCCTGGACAGAGGCTGGAGTGGATTGGA AATATTGATCCTGCGAATGGTAATACTAAATAT GACCCGAAGTCCAGGGCAAGGCCACTATAA CAGCAGACACATCCGCGAGCACAGCCTACA TGGAGCTCAGCAGCCTGAGATCTGAGGAC

			ACTGCCGTCTATTACTGTGCTAGATGGTTAC GACATTTTACTATGCTATGGACTACTGGGGTC AAGGAACCCTGGTCACCGTCTCCTCA
29	AA	VH of Hu24I11	QVQLVQSGAEVKKPGASVKVSCKASGFNIKDT YVHWVRQAPGQRLEWIGNIDPANGNTKYDPKF QGKATITADTSASTAYMELSSLRSEDTAVYYCAR WLRHFYYAMDYWGQGTLVTVSS
30	DNA	VL of Hu24I11	GACATCCAGATGACTCAGTCTCCATCCTCCCT GTCTGCATCTGTGGGAGACAGAGTCACCATCA CATGTCGAGCAAGTGAGAAACATTACTACAGT TTAGCATGGTATCAGCAGAAGCCAGGGAAAG CCCCTAACGCTCCTGATCTATAATGCAAACAGCT TGGAAAGATGGTGTCCCATCGAGGTTCAGTGGC AGTGGATCTGGGACACAGTATACTCTCACCAT CAGCAGCCTGCAGCCTGAAGATTGCAACTT ATTACTGAAACAGGCTTATGACGTTCCGTAC ACGTTCGGACAAGGGACCAAGGTGGAAATCA AA
31	AA	VL of Hu24I11	DIQMTQSPSSLSASVGDRVITCRASENIYVSLA WYQQKPGKAPKLLIYNANSLEDGVPSRFSGSGS GTQYTLTISSLQPEDFATYYCKQAYDVPYTFGQ GTKVEIK
32	AA	CDRH1 of 1K11 (FERM BP-10510)	DYNMD
33	AA	CDRH2 of 1K11 (FERM BP-10510)	DINPNNGGTIYNQKFQG
34	AA	CDRH3 of 1K11 (FERM BP-10510)	SGVISTDY
35	DNA	VH of 1K11 (FERM BP-10510)	GTGCAGCTGCAGGAGTCAGGACCTGAGCTGG TGAAGCCTGGGGCTTCAGTGAAGATACCCCTGC AAGGCTTCTGGATACACATTCACTGACTACAA CATGGACTGGGTGAAGCAGAGCCATGGAAAG AGCCTTGAGTGGATTGGAGATATTAACTCTAAC AACGGTGGTACAATCTACAACCAGAAGTCCA GGGCAAGGCCACATTGACTGTAGACAAGTCC TCCAGCACAGCCTACATGGAGCTCCGCAGCCT GACATCTGAGGACACTGCAGTCTATTACTGTG CAAGATCGGGGGTTATTAGTACGGACTACTGG GGCCAAGGGACCACGGTCACCGTCTCCTCA
36	AA	Mature VH of 1K11 (FERM BP-10510) starting from	?VQLQESGPELVKPGASVKIPCKASGYTFTDYN MDWVKQSHGKSLEWIGDINPNNGGTIYNQKFQ GKATLTVDKSSSTAYMELRSLTSEDTAVYYCARS GVISTDYWGQGTTVTVSS

		2nd residue according to Kabat numbering	
37	AA	CDRL1 of 1K11 (FERM BP-10510)	RASQEISGYLI
38	AA	CDRL2 of 1K11 (FERM BP-10510)	AASTLDS
39	AA	CDRL3 of 1K11 (FERM BP-10510)	LQYANYPPT
40	DNA	VL of 1K11 (FERM BP-10510)	GACATCCAGATGACACAGTCTCCACCCTCCCTATCTGCCTCTCTGGGAGAAAGAGTCAGTCTCACCTGTCGGGCAAGTCAGGAAATTAGTGGTTACTTAACATCTGGCTTCAACAGAAACCAGATGGAACATTCAACGCCTGATCTACGCCGCATCCACTTAGATTCTGGTGTCCAAAAAGGTTCACTGGCAAGGGTCTGGGTCAAGATTATTCTCTCACCATCAGAGCCTTGAGTCTGAAGATTTCAGACTATTACTGTCTACAATATGCTAATTATCCTCCGACGTTCGGTGGAGGCACCAAGCTGGAAATCAAACGG
41	AA	Mature VL of 1K11 (FERM BP-10510)	DIQMTQSPPSLSASLGERVSLTCRASQEISGYLIWLQQKPDGTIQRLIYAASTLDSGVPKRFSGSRSGSDYSLTISSLESEDFADYYCLQYANYPPTFGGGTKLEIKR
42	AA	CDRH1 of 21C5 (FERM BP-10511)	DYYMY
43	AA	CDRH2 of 21C5 (FERM BP-10511)	TISDGGNYTYYPDSVKG
44	AA	CDRH3 of 21C5 (FERM BP-10511)	DRDGSSLFAY
45	DNA	VH of 21C5 (FERM BP-10511)	GTGCAGCTGCAGGAGTCTGGGGGAGGCTTAGTGAAGCCTGGAGGGTCCCTGAAACTCTCCTGTGCAGCCTCTGGATTCACTTCAGTGACTATTACATGTATTGGGTTGCCAGACTCCGGAAAAGAGGCTGGAGTGGGTCGCAACCATTAGTGATGGTGTAATTACACCTACTATCCAGACAGTGTGAAGGGCGATTCAACCATTCCAGAGACAATGCCAAGAATAACCTGTACCTGCAAATGAGCAGTCTGAGTCTGAGGAACACAGCCATGTATTACTGTGCA

			AGAGATCGGGACGGTAGTAGCCTGTTGCTTA CTGGGGCCAAGGGACCACGGTCACCGTCTCC TCA
46	AA	Mature VH of 21C5 (FERM BP-10511) starting from 2nd residue according to Kabat numbering	?VQLQESGGGLVKPGGSLKLSCAASGFTFSDYY MYWVRQTPEKRLEWVATISDGNYTYPDSVK GRFTISRDNAKNNLQMQSSLKSEDTAMYCA RDRDGSSLFAYWGQGTTVTVSS
47	AA	CDRL1 of 21C5 (FERM BP-10511)	KASQDVNIAVA
48	AA	CDRL2 of 21C5 (FERM BP-10511)	WASTRHT
49	AA	CDRL3 of 21C5 (FERM BP-10511)	QQHYNTPW
50	DNA	VL of 21C5 (FERM BP-10511)	CATCCAGATGACACAGTCTCCAATTATGTC CACATCAGTAGGAGACAGGGTCAGCATCACCT GCAAGGCCAGTCAGGATGTGAATATTGCTGTA GCCTGGTATCAACAAAGACCAAGGGCAATCTCC TAAACTACTGATTACTGGGCATCCACCCGGC ACACTGGAGTCCTGATCGCTTCACAGGCAGT GGATCTGGACAGATTACTCTCACCATCAG CAGTGTGCAGGCTGAAGACCTGGCACTTATT ACTGTCAACATCATAACACTCCGTGGACG TTCGGTGGAGGCACCAAGCTGGAAATCAAAC GG
51	AA	Mature VL of 21C5 (FERM BP-10511)	HPDDTVSKFMSTSVGDRVSITCKASQDVNIAVA WYQQRPGQSPKLLIYWASTRHTGVPDRFTGSGS GTDYTLTISSVQAEDLALYYCQQHYNTPWTFGG GTKLEIKR
52	AA	CDRH1 of 25B6 (FERM BP-10513)	SYGVH
53	AA	CDRH2 of 25B6 (FERM BP-10513)	VIWSGGSTNYNSALMS
54	AA	CDRH3 of 25B6 (FERM BP-10513)	DYGNYPWFAY

55	DNA	VH of 25B6 (FERM BP-10513)	GTCAAGCTGCAGCAGTCTGGACCTGGCCTGG TGGCGCCCTCACAGAGCCTGTCCATCACTTGC ACTGTCTCTGGGTTTCATTAACCAGTTAGGT GTACACTGGGTTGCCAGCCTCCAGGAAAGG GTCTGGAGTGGCTGGAGTAATATGGTCTGGT GGAAGCACAAATTATAATTGGCTCATGTCC AGACTGAGCATCAGTAAGACAAATTAAAGAG CCAAGTTTCTTAAAAATGAACAGTCTGCCAAA CTGATGACACAGCCATATACTACTGTGCCAGA GACTATGGTAACTACCCCTGGTTGCTTACTGG GGCCAAGGGACCACGGTCACCGTCTCCTCA
56	AA	Mature VH of 25B6 (FERM BP-10513) starting from 2nd residue according to Kabat numbering	?VKLQQSGPGLVAPSQSLSITCTVSGFSLTSYGV HWVRQPPGKGLEWLGVIVSGGSTNYNSALMS RLSISKDNFKSQVFLKMNSLQTDDTAIYYCARD YGNYPWFAYWGQGTTVTVS
57	AA	CDRL1 of 25B6 (FERM BP-10513)	KASQDVNTAVA
58	AA	CDRL2 of 25B6 (FERM BP-10513)	SASYRYT
59	AA	CDRL3 of 25B6 (FERM BP-10513)	QQHYSTPCA
60	DNA	VL of 25B6 (FERM BP-10513)	CATCCAGATGACACAGTCTCCAAATTCTATGTC CACATCAGTAGGAGACAGGGTCAGCATCACCT GCAAGGCCAGTCAGGATGTGAATACTGCTGTG GCCTGGTATCAACAGAAACCAGGACAATCCCC TAAACTACTGATTACTCGGCATCCTACCGGTA CACTGGAGTCCCTGATCGCTCACTGGCAGTG GATCTGGGACGGATTCACTTCACCATCAGC AGTGTGCAGGCTGAAGACCTGGCAGTTATTA CTGTCAGCAACATTATAGTACTCCGTGCGCGTT CGGAGGGGGGACAAAGTTGAAATAAACGG
61	AA	Mature VL of 25B6 (FERM BP-10513)	HPDDTVSKFMSTVGDRVSITCKASQDVNTAVA WYQQKPGQSPKLLIYSASYRYTGVPDRFTGSGS GTDFTFTISSVQAEDLAVYYCQQHYSTPCAFGG GTKLEIKR
62	AA	CDRH1 of 28S1 (FERM BP- 10832)	GYGVN

63	AA	CDRH2 of 28S1 (FERM BP- 10832)	MIWGDGITEYNSALKS
64	AA	CDRH3 of 28S1 (FERM BP- 10832)	RDASSGYGFA
65	DNA	VH of 28S1 (FERM BP- 10832)	AGGTGAAGCTGCAGGAGTCAGGACCTGGCCTGGTCGCCCCCACAGAGCCTGTCCATCACATGCACCGTCTCAGGGTTCTCATTAACCGGCTATGGTAAACTGGGTTGCCAGCCTCCAGGAAAGGGTCTGGAGTGGCTGGGAATGATATGGGGTGATGGAATCACAGAGTATAATTAGCTCTCAAATCCAGACTGAGCATCAGCAAGGACAACCTCAAGAGCCAAGTTTCTAAAAATGAACAGTCTGCAAACGTGATGACACAGCCAGGTACTACTGTGCCAGAGATGCCAGCTCGGGCTACGGGTTGCTTACTGGGCCAAGGGACCACGGTCACCGTCTCCTCA
66	AA	Mature VH of 28S1 (FERM BP- 10832) starting from 2nd residue according to Kabat numbering	?VKLQESGPLVAPSQSLSITCTVSGFSLTGYGVNWVRQPPGKGLEWLGMIWGDGITEYNSALKSRLSISKDNNSKSQVFLKMNSLQTDDTARYYCARDASSGYGFAYWGQGTTVTVSS
67	AA	CDRL1 of 28S1 (FERM BP- 10832)	TASSSVSSSYLH
68	AA	CDRL2 of 28S1 (FERM BP- 10832)	STSNLAS
69	AA	CDRL3 of 28S1 (FERM BP- 10832)	HQYHRSPYT
70	DNA	VL of 28S1 (FERM BP- 10832)	TACATTGTGCTGACCCAGTCTCCAGCAATCATGTCTGCATCTCTAGGGGAACGGGTACCATGACCTGCACTGCCAGCTCAAGTGTAAAGTCCAGTTACTGCACAGTGGTACCAAGCAGAAGCCAGGATCCTCCCCAAACTCTGGATTATAGCACATCCAACTGGCTCTGGAGTCCAGCTCGCTTCAGTGCAGTGGGTCTGGGACCTCTTACTCTCACATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTATTACTGCCACCAAGTATCATCGTCCCCGTAACGTTCGGAGGGGGACAAAGTTGGAAATAAACGG

71	AA	Mature VL of 28S1 (FERM BP- 10832)	YIVLTQSPAAMSASLGERVTMTCTASSSVSSYL HWYQQKPGSSPKLWIYSTSNLASGVPARFSGSG SGTSYSLTISSMEAEDAATYYCHQYHRSPYTFG GGTKLEIKR
72	DNA	VH of 24I11 (FERM BP-10512) including sequence encoding signal peptide (1-57), flanked by SpeI and HindIII sites	ACTAGTACCAACCATGAAATGCAGCTGGTTAT CTTCTTCCTGATGGCAGTGGTTACAGGGTCA ATTCAGAGGTTCAGCTGCAGCAGTCTGGGC AGAGCTTGTGAAGCCAGGGGCCTCAGTCAAG TTGTCCTGCACAGCTCTGGCTTCAACATTAA AGACACCTATGTGCACTGGGTGAAGCAGAGG CCTGAACAGGGCCTGGAGTGGATTGGAAATAT TGATCCTGCGAATGGTAATACTAAATATGACCC GAAGTTCCAGGGCAAGGCCACTATAACAGCA GACACATCCTCCAACACAGCCTACCTGCACCT CAGCAGCCTGACATCTGAGGACACTGCCGTCT ATTACTGTGCTAGATGGTACGACATTACT ATGCTATGGACTACTGGGGTCAAGGAACCTCA GTCACCGTCTCCTCAGGTAAAGAATGGCCTCTA AGCTT
73	DNA	VL of 24I11 (FERM BP-10512) including sequence encoding signal peptide (1-60), flanked by NheI and EcoRI sites	GCTAGCACCACCATGAGTGTGCCACTCAACT CCTGGGGTTGCTGCTGCTGGCTTACAGACG CAGGATGTGACATCCAGATGACTCAGTCTCCA GCCTCCCTGGCTGCATCTGTGGGAGAAACTGT CACCGACATCCAGATGACTCAGTCTCCAGCCT CCCTGGCTGCATCTGTGGGAGAAACTGTCA GGGAAATCTCCTCAGCTCCTGATCTATAATGCA AACAGCTTGGAAAGATGGTGTCCCATCGAGGTT CAGTGGCAGTGGATCTGGGACACAGTATTCTA TGAAGATCAACAGCATGCAGCCTGAAGATACC GCAACTTATTCTGAAACAGGCTATGACGTT CCGTACACGTTGGAGGGGGACCAAGCTGG AAATAAAACGTAAGTAGTCTCTCAGAATT
74	DNA	Fig. 10 (w/ 5'-GGG & CCC-) and Fig. 12 Hu24I11 VH gene flanked by SpeI and HindIII sites	ACTAGTACCAACCATGAAATGCAGCTGGTTAT CTTCTTCCTGATGGCAGTGGTTACAGGGTCA ATTCACAGGTTCAGCTGGTGCAGTCTGGGC GAGGTGAAGAAGCCAGGGGCCTCAGTCAAGG TTTCCTGCAAGGCTTCTGGCTTCAACATTAAA GACACCTATGTGCACTGGGTGCCAGGCC TGGACAGAGGCTGGAGTGGATTGGAAATATTG ATCCTGCGAATGGTAATACTAAATATGACCCGA AGTTCCAGGGCAAGGCCACTATAACAGCAGA CACATCCCGAGCACAGCCTACATGGAGCTCA GCAGCCTGAGATCTGAGGACACTGCCGTCTAT TACTGTGCTAGATGGTACGACATTACTAT GCTATGGACTACTGGGGTCAAGGAACCTGGT CACCGTCTCCTCAGGTGAGTCCTCACAAAAGC TT
75	DNA	Fig. 11 (w/	GCTAGCACCACCATGAGTGTGCCACTCAACT

		5'-GGG & CCC-) and Fig. 13 Hu24I11 VL gene flanked by NheI and EcoRI sites	CCTGGGGTTGCTGCTGCTGTGGCTTACAGACG CACGATGTGACATCCAGATGACTCAGTCTCCA TCCTCCCTGTCTGCATCTGTGGGAGACAGAGT CACCATCACATGTCGAGCAAGTGAGAACATT ACTACAGTTAGCATGGTATCAGCAGAACGCCA GGGAAAGCCCCCTAACGCTCCTGATCTATAATGC AAACAGCTTGGAAAGATGGTGTCCCATCGAGGT TCAGTGGCAGTGGATCTGGGACACAGTATACT CTCACCATCAGCAGCCTGCAGCCTGAAGATT TGCAACTTATTACTGTAACACAGGCTTATGACGT TCCGTACACGTTGGACAAGGGACCAAGGTG GAAATCAAACGTGAGTAGAATTAAAGAATT
76	AA	Human α 9 integrin (signal peptide; 1-29 residues; in italic)	<i>MGGPAAPRGAGRLALLALVVAGIPAGAYNLDpq</i> RPVHFQGPADSFFGYAVLEHFHDNTRWVLVGAP KADSKYSPSVKSPGAVFKCRVHTNPDRRCTELD MARGKNRGTSCGKTCREDRDEWMGVSLARQ PKADGRVLACAHWRKNIYYEADHILPHGFCYII PSNLQAKGRTLIPCYEEYKKKYGEEHGSCQAGI AGFFTEELVVMGAPGSFYWAGTIKVNLTDNTY LKLNDEVIMNRRYTYLGYAVTAGHFSHPSTIDV VGGAPQDKGIGKVYIFRADRRSGTLIKIFQASGK KMGSYFGSSLCAVDLNGDGLSDLLVGAPMFSEI RDEGQVTVYINRGNGALEEQLALTGDAYNAH FGESIASLDDLDNDGFPDVAIGAPKEDDFAGAV YIYHGDAGGIVPQYSMKGQKINPVRMFGQS ISGGIDMDGNGYPDVTVGAFMSDSVVLLARP VITVDVSIFLPGSINITAPQCHDGQQPVNCLNTV TCFSFHGKHVPGEIGLNYVLMADVAKKEKGQM PRVYFVLLGETMGQVTEKLQLTYMEETCRHYV AHVKRRVQDVSPIVFEAAYSLSEHVTGEEEREL PPLTPVLRWKKGQKIAQKNQTVFERNCRSEDCA ADLQLQGKLLLSSMDEKTLYLALGAVKNISLNISI SISNLGDDAYDANVSFNVSRELFFINMWQKEEM GISCELLESDFLKCSVGFPMRSKSKYEFSVIFDT SHLSGEEEVLSFIVTAQSGNTERSESLHDNTLVL MVPLMHEVDT SITGIMSPTSFVYGESVDAANFI QLDDLECHFQPINITLQVYNTGPSTLPGSSVSISF PNRLSSGGAEMFHVQEMVVGQEKGNCSFQKNP TPCIIPQEKENIFHTIAFFTKSGRKVLDCEKPGIS CLTAHCNFSALAKEEVRTIDYMLLNTIELKKDS SSVIQFMSRAKVVDPALRVVEIAHGNPEEVTV VFEALHNLEPRGYVVGWIAISLLVGILIFLLAV LLWKMGGFFRRRYKEIIAEKNRKENEDSWDWV QKNQ
77	AA	Partial aa sequence of hu α 9 integrin	FQGPADSFFGYA
78	AA	Partial aa sequence of	KSPGAVFKCRVHTNPDRR

		hu α 9 integrin	
79	AA	Partial aa sequence of hu α 9 integrin	WMGVSLARQPKADGRVLA
80	aa	Partial aa sequence of hu α 9 integrin	CAHRWKNIYYEADHI
81	aa	Partial aa sequence of hu α 9 integrin	GFCYIIPSNLQAKGRTLI
82	aa	Partial aa sequence of hu α 9 integrin	VMGAPGSFYWAGTIKVLN
83	aa	Partial aa sequence of hu α 9 integrin	VIMNRRYTYLGYAVT
84	aa	Partial aa sequence of hu α 9 integrin	VYIFRADRRSGTLIKIFQ
85	aa	Partial aa sequence of hu α 9 integrin	QYSMKLSGQKINPVLRMFGQSISG
86	aa	Partial aa sequence of hu α 9 integrin	VVLLRARPVITVDVSIFL
87	aa	Partial aa sequence of hu α 9 integrin	RHYVAHVKKRQDVVISPI
88	aa	Partial aa sequence of hu α 9 integrin	ELPPLTPVLRWKKGQKIAQKNQTVFERNCR
89	aa	Partial aa sequence of hu α 9 integrin	YLALGAVKNISL
90	aa	Partial aa sequence of hu α 9 integrin	CSVGFPFMRSKSKYEFSV
91	aa	Partial aa sequence of hu α 9 integrin	SSSVIQFMSRAKVKVDPALRV
92	aa	Tenascin-C hu α 9-binding site	AEIDGIEL

93	aa	Human fibronectin hu α 9-binding site	CPEDGIHELF
94	DNA	GeneRacer 5' primer	CGACTGGAGCACGAGGACACTGA
95	DNA	VH primer	GCCAGTGGATAGACAGATGG
96	DNA	VL primer	GATGGATACAGTTGGTGCAGC
97	DNA	VH 5' primer w/ SpeI site	GGGACTAGTACCACCATGAAATGCAGCTGGGT TATCTTC
98	DNA	VH primer w/ HindIII site	GGGAAGCTTAGAGGCCATTCTTACCTGAGGA GACGGTGAUTGAGGTTCC3
99		VL 5' primer w/ NheI site	GGGGCTAGCACCACCATGAGTGTGCCACTCA ACTCCTG
100		VL 5' primer w/ EcpRI site	GGGGAATTCTGAGAAGACTACTTACGTTTAT TTCCAGCTTGGTCCCCCCC
101	DNA	JNJ120	GGGACTAGTACCACCATGAAATGCAGC
102	DNA	JNJ137	GGGACTAGTACCACCATGAAATGCAGCTGGGT TATCTTCTCCTGATGGCAGTGGTT
103	DNA	JNJ138	AGACTGCACCAGCTGAACCTGTGAATTGACCC CTGTAACCACTGCCATCAGGAAGAA
104	DNA	JNJ139	CAGGTTCAGCTGGTGCAGTCTGGGGCAGAGG TGAAGAACGCCAGGGCCTCAGTCAAG
105	DNA	JNJ140	GTCTTAATGTTGAAGCCAGAAGCCTGCAGG AAACCTTGACTGAGGCCCTGGCTT
106	DNA	JNJ141	TCTGGCTCAACATTAAAGACACCTATGTGCA CTGGGTGCGCCAGGCCCTGGACAGAGG
107	DNA	JNJ142	ACCATTCGCAGGATCAATATTCCAATCCACTC CAGCCTCTGTCCAGGGCCTGGCG
108	DNA	JNJ143	AATATTGATCCTGCGAATGGTAATACTAAATAT GACCCGAAGTCCAGGGCAAGGCCACT
109	DNA	JNJ144	CATGTAGGCTGTGCTCGGGATGTGTCTGCTG TTATAGTGGCCTGCCCTGGAACCTT
110	DNA	JNJ145	TCCCGCAGCACAGCCTACATGGAGCTCAGCA GCCTGAGATCTGAGGACACTGCCGTC
111	DNA	JNJ146	ATAGAAAAATGTCGTAAACCATCTAGCACAGT AATAGACGGCAGTGTCTCAGA
112	DNA	JNJ147	TGGTTACGACATTACTATGCTATGGACTAC TGGGGTCAAGGAACCCCTGGTCACC
113	DNA	JNJ148	GGGAAGCTTTGTGAGGACTCACCTGAGGAG

			ACGGTGACCAGGGTTCTTGACC
114	DNA	JNJ149	GGGAAGCTTGTGAGGACTC
115	DNA	JNJ150	GGGGCTAGCACCACCATGAGTGTGCCACTCA
116	DNA	JNJ126	GGGGCTAGCACCACCATGAGTGTGCCACTCA ACTCCTGGGTTGCTGCTGCTGTGG
117	DNA	JNJ127	AGACTGAGTCATCTGGATGTCACATCGTGCCT CTGTAAGCCACAGCAGCAGCAACCCCAG
118	DNA	JNJ128	GACATCCAGATGACTCAGTCTCCATCCTCCCT GTCTGCATCTGTGGGAGACAGA
119	DNA	JNJ129	GTAAATGTTCTCACTGCTCGACATGTGATGGT GACTCTGTCTCCCACAGATGCAGA
120	DNA	JNJ130	CGAGCAAGTGAGAACATTACTACAGTTAGC ATGGTATCAGCAGAACGCCAGGGAAA
121	DNA	JNJ131	CAAGCTTTGCATTATAGATCAGGAGCTTAG GGGCTTCCCTGGCTCTGCTGATA
122	DNA	JNJ132	ATCTATAATGCAAACAGCTTGGAAAGATGGTGT CCCATCGAGGTTCACTGGCAGTGG
123	DNA	JNJ133	CAGGCTGCTGATGGTGAGAGTATACTGTGTCC CAGATCCACTGCCACTGAACCTCGA
124	DNA	JNJ134	ACTCTCACCATCAGCAGCCTGCAGCCTGAAGA TTTGCAACTTATTACTGTAAACAG
125	DNA	JNJ135	GGTCCCTTGTCCGAACGTGTACGGAACGTCAT AAGCCTTTACAGTAATAAGTTGC
126	DNA	JNJ136	TACACGTTGGACAAGGGACCAAGGTGGAAA TCAAACGTGAGTAG
127	DNA	JNJ101	GGGAATTCTTAAATTCTACTCACGTTGATT TCCA
128	DNA	JNJ117	GGGAATTCTTAAATTCTA
129	DNA	Primer for mouse γ 1, γ 2a, γ 2b and γ 3 H-chains	GCCAGTGGATAGACTGATGG
130	DNA	Primer for mouse κ L-chain primer	GATGGATACAGTTGGTGCAGC

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A humanized antibody or an antigen-binding fragment thereof that immunospecifically recognizes human $\alpha 9$ integrin, comprising:
 - 5 a H-chain comprising the amino acid sequence of SEQ ID NO:29, and at least one CDRH derived from at least one of CDRHs of a non-human antibody that immunospecifically recognizes human $\alpha 9$ integrin; and
 - a L-chain comprising the amino acid sequence of SEQ ID NO:31, and at least one CDRL derived from at least one of CDRLs of a non-human antibody that
- 10 immunospecifically recognizes human $\alpha 9$ integrin.
2. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence of SEQ ID NO:29.
- 15 3. The nucleic acid molecule of claim 2, wherein said nucleotide sequence has the nucleotide sequence of SEQ ID NO:28.
4. The nucleic acid molecule of claim 2 further comprising a nucleotide sequence encoding a signal peptide.
- 20 5. The nucleic acid molecule of claim 4, wherein said signal peptide comprises the amino acid sequence of SEQ ID NO:10.
6. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the amino acid sequence of SEQ ID NO:31.
- 25 7. The nucleic acid molecule of claim 6, wherein said nucleotide sequence has the nucleotide sequence of SEQ ID NO:30.
- 30 8. The nucleic acid molecule of claim 6 further comprising a nucleotide sequence encoding a signal peptide.
9. The nucleic acid molecule of claim 8, wherein said signal peptide comprises the

amino acid sequence of SEQ ID NO:17.

10. A vector comprising the nucleic acid molecule of claim 2 or 6, or both, wherein said nucleic acid molecule is operably linked to one or more regulatory elements.

5

11. An isolated host cell comprising the vector of claim 10.

12. A method for preparing a humanized antibody or an antigen-binding fragment thereof, comprising culturing the host cell of claim 11 under conditions so that the 10 humanized antibody or an antigen-binding fragment thereof is expressed, and collecting the expressed humanized antibody.

13. A pharmaceutical composition comprising the humanized antibody or an antigen-binding fragment thereof of claim 1, and a pharmaceutically acceptable carrier.

15

14. A method for preventing or treating a disorder or disease associated with $\alpha 9$ integrin, said method comprising administering an effective amount of the humanized antibody or an antigen-binding fragment thereof of claim 1 to a subject in need thereof.

20

15. A method for diagnosing a disorder or disease associated with $\alpha 9$ integrin *in vivo*, said method comprising administering an effective amount of the humanized antibody or an antigen-binding fragment thereof of claim 1 to a subject to be examined.

25

20. The humanized antibody or an antigen-binding fragment thereof of claim 1 or the nucleic acid molecule of claims 2 or 6, substantially as hereinbefore described with reference to any of the Examples and/or Figures.

Fig. 1

ATGAAATGCAGCTGGGTTATCTTCCTCCGATGGCAGTGGTACAGGGGTCATTCAAGAG
M K C S W V I F F L M A V V T G V N S E

GTTCAGCTGCAGCAGTCTGGGGCAGAGCTTGTGAAGCCAGGGGCTCAGTCAGTTGTCC
V Q L Q Q S G A E L V K P G A S V K L S

TGCACAGCTCTGGCTTCAACATTAAGACACCTATGTGCACTGGGTGAAGCAGAGCCT
C T A S G F N I K D T Y V H W V K Q R P

GAACAGGGCTGGAGTGGATTGGAAATATTGATCCTGCGAATGGTAATACTAAATATGAC
E Q G L E W I G N I D P A N G N T K Y D

CCGAAGTTCCAGGGCAAGGCCACTATAACAGCAGACACATCCTCCAACACAGCCTACCTG
P K F Q G K A T I T A D T S S N T A Y L

CACCTCAGCAGCCTGACATCTGAGGACACTGCCGTCTATTACTGTGCTAGATGGTTACGA
H L S S L T S E D T A V Y Y C A R W L R

CATTTTTACTATGCTATGGACTACTGGGGTCAAGGAACCTCAGTCACCGTCTCCTCA
H F Y Y A M D Y W G Q G T S V T V S S

Fig. 2

ATGAGTGTGCCACTCAACTCCTGGGTTGCTGCTGTGGCTTACAGACGCAGGATGT
M S V P T Q L L G L L L W L T D A G C

GACATCCAGATGACTCAGTCTCCAGCCTCCCTGGCTGCATCTGTGGAGAACTGTCACC
D I Q M T Q S P A S L A A S V G E T V T

ATCACATGTGGAGCAAGTGAGAACATTACTACAGTTAGCATGGTATCAGCAGAACAA
I T C R A S E N I Y Y S L A W Y Q Q K Q

GGGAAATCTCCTCAGCTCCTGATCTATAATGCAAACAGCTTGGAGATGGTGTCCATCG
G K S P Q L L I Y N A N S L E D G V P S

AGGTTCACTGGCAGTGGATCTGGGACACAGTATTCTATGAAGATCAACAGCATGCAGCCT
R F S G S G S G T Q Y S M K I N S M Q P

GAAGATAACCGCAACTTATTCGTAAACAGGCTTATGACGTTCCGTACACGTTGGAGGG
E D T A T Y F C K Q A Y D V P Y T F G G

GGGACCAAGCTGGAAATAAAA
G T K L E I K

Fig. 3

SpcI

ACTAGTACCACCATGAAATGCAGCTGGGTTATCTTCTTCTGATGGCAGTGGTTACAGGG
 M K C S W V I F F L M A V V T G

GTCAATTCAAGAGGTTCAAGCTGCAGCAGTCTGGGGCAGAGCTTGTGAAGCCAGGGGCTCA
 V N S E V Q L Q Q S G A E L V K P G A S

GTCAAGTTGTCTGCACAGCTTCTGGCTTCAACATTAAAGACACCTATGTGCACTGGGTG
 V K L S C T A S G F N I K D T Y V H W V

AAGCAGAGGCCTGAACAGGGCCTGGAGTGGATTGGAAATATGATCCTGCGAATGGTAAT
 K Q R P E Q G L E W I G N I D P A N G N

ACTAAATATGACCCGAAGTTCCAGGGCAAGGCCACTATAACAGCAGACACATCCTCCAAC
 T K Y D P K F Q G K A T I T A D T S S N

ACAGCCTACCTGCACCTCAGCAGCCTGACATCTGAGGACACTGCCGTCTATTACTGTGCT
 T A Y L H L S S L T S E D T A V Y Y C A

AGATGGTTACGACATTTTACTATGCTATGGACTACTGGGGTCAAGGAACCTCAGTCACC
 R W L R H F Y Y A M D Y W G Q G T S V T

HindIII

GTCTCCTCAGGTAAGAATGGCCTCTAAGCTT
 V S S

Fig. 4

NheI

GCTAGCACCACCATGAGTGTGCCACTCAACTCTGGGTTGCTGCTGCTGTGGCTTACA
 M S V P T Q L L G L L L W L T

GACGCAGGATGTGACATCCAGATGACTCAGTCTCCAGCCTCCCTGGCTGCATCTGTGGGA
 D A G C D I Q M T Q S P A S L A A S V G

GAAACTGTCAACCACATGTCGAGCAAGTGAGAACATTACTACAGTTACCATGGTAT
 E T V T I T C R A S E N I Y Y S L A W Y

CAGCAGAAGCAAGGGAAATCTCCTCAGCTCCTGATCTATAATGCAAACAGCTTGGAAAGAT
 Q Q K Q G K S P Q L L I Y N A N S L E D

GGTGTCCCATCGAGGTTCACTGGCAGTGGATCTGGACACAGTATTCTATGAAGATCAAC
 G V P S R F S G S G S G T Q Y S M K I N

AGCATGCAGCCTGAAGATAACCGCAACTTATTTCTGTAAACAGGCTTATGACGTTCCGTAC
 S M Q P E D T A T Y F C K Q A Y D V P Y

EcoRI

ACGTTCGGAGGGGGGACCAAGCTGAAATAAAACGTAAGTAGTCTTCAGAATT
T F G G G T K L E I K


Fig. 5

Fig. 6

	1	2	3	
24I11 VH	123456789	0123456789	0123456789	0123456789
Hu24I11 VH	EVQLQQSGA	ELVKPGASVK	LSCTASGFNI	<u>KDTYVHWVKQ</u>
X65891	QVQLVQSGA	EVKKPGASVK	VSCKASGFNI	<u>KDTYVHWVRQ</u>
	QVQLVQSGA	EVKKPGASVK	VSCKASGYTF	T-----WVRQ
	4	5	6	7
	0123456789	01223456789	0123456789	0123456789
	a			
24I11 VH	RPEQGLEWIG	<u>NIDPANGNTKY</u>	<u>DPKFQGKATI</u>	TADTSSNTAY
Hu24I11 VH	APGQRLEWIG	<u>NIDPANGNTKY</u>	<u>DPKFQGKATI</u>	<u>TADTSASTAY</u>
X65891	APGQRLEWMG	-----	-----	RVTI TRDTSASTAY
	8	9	0	1
	0122223456789	0123456789	0000123456789	0123
	abc			
24I11 VH	LHLSSLTSEDTAV	<u>YYCARWLRHF</u>	<u>YYAMDYWGQGTSV</u>	TVSS
Hu24I11 VH	MELSSLRSEDTAV	<u>YYCARWLRHF</u>	<u>YYAMDYWGQGTLV</u>	TVSS
X65891	MELSSLRSEDTAV	YYCAR-----	-----WGQGTLV	TVSS

Fig. 7

	1	2	3	
24I11 VL	123456789	0123456789	0123456789	0123456789
Hu24I11 VL	DIQMTQSPA	SLAASVGETV	TITCRASENI	YYSLAWYQQK
X72441	DIQMTQSPS	SLSASVGDRV	TITCRASENI	YYSLAWYQQK
	DIQMTQSPS	SLSASVGDRV	TITC-----	-----WYQQK
	4	5	6	7
24I11 VL	0123456789	0123456789	0123456789	0123456789
Hu24I11 VL	QGKSPQLLIY	<u>NANSLEDGVP</u>	SRFSGSGSGT	QYSMKINSMQ
X72441	PGKAPKLLIY	<u>NANSLEDGVP</u>	SRFSGSGSGT	<u>QYTLTISSLQ</u>
	PGKAPKLLIY	-----GVP	SRFSGSGSGT	DFTLTISSLQ
		8	9	1
		0123456789	0123456789	01234567
24I11 VL	PEDTATYFCK	<u>QAYDVPYTFG</u>	GGTKLEIK	
Hu24I11 VL	PEDPATYYCK	<u>QAYDVPYTFG</u>	QGTKVEIK	
X72441	PEDPATYYC-	-----FG	QGTKVEIK	

Fig. 8

OLIGO-NUCLEOTIDES	SEQUENCE	SEQ ID NO.
JNJ120	GGGACTAGTACCAACCATGAAATGCAGC	101
JNJ137	GGGACTAGTACCAACCATGAAATGCAGCTGGGTTATCTTCTTCTGATGGCAGTGGTT	102
JNJ138	AGACTGCACCAAGCTGAAACCTGTGAATTGACCCCTGTAACCACTGCCATCAGGAAGAA	103
JNJ139	CAGGTTCACTGGTGCAGTCTGGGCAGAGGTGAAGAAGCCAGGGCCCTCAGTCAAG	104
JNJ140	GTCTTAATGTTGAAGCCAGAACGCTTGAGGAAACCTTGACTGAGGCCCTGGCTT	105
JNJ141	TCTGGCTTCAACATTAAGACACCTATGTGCACTGGTGCAGGCCAGGGCCCTGGACAGAGG	106
JNJ142	ACCATTGCAAGGATCAATATTCACACTCCAGCCTCTGCCAGGGCCCTGGCG	107
JNJ143	AATATTGATCCTGCAATGGTAATACTAAATATGACCCGAAGTCCAGGGCAAGGCCACT	108
JNJ144	CATGTAGGCTGTGCTCGGGATGTGCTGTTATAGTGGCCTTGCCCTGGAACCTT	109
JNJ145	TCCGCGAGCACAGCCTACATGGAGCTCAGCAGCCTGAGATCTGAGGACACTGCCGTC	110
JNJ146	ATAGTAAAATGTCGAACCATCTAGCACAGTAATAGACGGCAGTGTCCCTCAGA	111
JNJ147	TGGTTACGACATTTTACTATGCTATGGACTACTGGGTCAAGGAACCCCTGGTCACC	112
JNJ148	GGGAAGCTTTGTGAGGACTCACCTGAGGAGACGGTGACCAGGGTTCCCTGACC	113
JNJ149	GGGAAGCTTTGTGAGGACTC	114

Fig. 9

OLIGO-NUCLEOTIDES	SEQUENCE	SEQ ID NO.
JNJ150	GGGGCTAGCACCACCATGAGT	115
JNJ126	GGGGCTAGCACCACCATGAGTGTGCCACTCAACTCCTGGGGTTGCTGCTGTGG	116
JNJ127	AGACTGAGTCATCTGGATGTCACATCGTGCCTGTAAGCCACAGCAGCAGCAACCCAG	117
JNJ128	GACATCCAGATGACTCAGTCTCCATCCTCCCTGCTGCATCTGGGAGACAGA	118
JNJ129	GTAAATGTTCTCACTGCTCGACATGTGATGGTACTCTGTCTCCCACAGATGCAGA	119
JNJ130	CGAGCAAGTGAGAACATTACTACAGTTAGCATGGTATCAGCAGAAGCCAGGGAAA	120
JNJ131	CAAGCTGTTGCATTATAGATCAGGAGCTTAGGGGCTTCCCTGGCTTGCTGATA	121
JNJ132	ATCTATAATGCAAACAGCTTGGAGATGGTGTCCCATCGAGGTTCACTGGCAGTGG	122
JNJ133	CAGGCTGCTGATGGTAGAGTATACTGTGTCCAGATCCACTGCCACTGAACCTCGA	123
JNJ134	ACTCTCACCATCAGCAGCCTGCAGCCTGAAGATTTGCAACTTATTACTGAAACAG	124
JNJ135	GGTCCCTTGTCCGAACGTGTACGGAACGTCTAAAGCCTGTTACAGTAATAAGTGC	125
JNJ136	TACACGTTGGACAAGGGACCAAGGTGAAATCAAACGTGAGTAG	126
JNJ101	GGGGATTCTTAAATTCTACTCACGTTGATTCCA	127
JNJ117	GGGGATTCTTAAATTCTA	128

Fig. 10

Spel

1 GGGACTAGTACCAACC ATG AAA TGC AGC TGG GTT ATC TTC TCC CTG ATG GCA GTG GTT ACA ACA GGG
 17 ▶ M K C S W V I F F L M A V V T G
 JNJ120 → ← JNJ137

64 GTC AAT TCA CAG GTT CAG CTG GTG CAG TCT GGG GCA GAG GTG AAG AAG CCA GGG GCC TCA
 17 ▶ V N S Q V Q L V Q S G A E V K K P G A S
 JNJ138 → ← JNJ139

124 GTC AAG GTT TCC TGC AAG GCT TCT GGC TTC AAG ATT AAA GAC ACC TAT GTG CAC TGG GTG
 37 ▶ V K V S C K A S G F N I K D T Y V H W V
 JNJ140 → ← JNJ141

184 CGC CAG GCC CCT GGA CAG AGG CTG GAG TGG ATT GGA AAT ATT GAT CCT GCG AAT GGT AAT
 57 ▶ R Q A P G Q R L E W I G N I D P A N G N
 ← JNJ142

244 ACT AAA TAT GAC CCG AAG TTC CAG GGC AAG GCC ACT ATA ACA GCA GAC ACA TCC GCG AGC
 77 ▶ T K Y D P K F Q G K A T I T A D T S A S
 JNJ143 → ← JNJ144

BgIII

304 ACA GCC TAC ATG GAG CTC AGC AGC CTG AGA TCT GAG GAC ACT GCC GTC TAT TAC TGT GCT
 97 ▶ T A Y M E L S S L R S E D T A V Y Y C A
 JNJ145 → ← JNJ146

364 AGA TGG TTA CGA CAT TTT TAC TAT GCT ATG GAC TAC TGG GGT CAA GGA ACC CTG GTC ACC
 117 ▶ R W L R H F Y Y A M D Y W G Q G T L V T
 JNJ147 → ← JNJ148

HindIII

424 GTC TCC TCA GGTGAGTCCTCACAAAAGCTTCCC
 137 ▶ V S S ← JNJ149

JNJ148

Fig. 11

NheI

GGGGCTAGCACCACC ATG AGT GTG CCC ACT CAA CTC CTG GGG TTG CTG CTG CTG TGG CTT ACA
 ▶ M S V P T Q L L G L L L W L T

██████████ → JNJ126 ← ██████████

GAC GCA CGA TGT GAC ATC CAG ATG ACT CAG TCT CCA TCC TCC CTG TCT GCA TCT GTG GGA
 ▶ D A R C D I Q M T Q S P S S L S A S V G

██████████ → JNJ128 ← ██████████

JNJ127

GAC AGA GTC ACC ATC ACA TGT CGA GCA AGT GAG AAC ATT TAC TAC AGT TTA GCA TGG TAT
 ▶ D R V T I T C R A S E N I Y Y S L A W Y

██████████ → JNJ130 ← ██████████

JNJ129

CAG CAG AAG CCA GGG AAA GCC CCT AAG CTC CTG ATC TAT AAT GCA AAC AGC TTG GAA GAT
 ▶ Q Q K P G K A P K L L I Y N A N S L E D

██████████ → JNJ131 ← ██████████

JNJ132

GCT GTC CCA TCG AGG TTC AGT GGC AGT GGA TCT GGG ACA CAG TAT ACT CTC ACC ATC AGC
 ▶ G V P S R F S G S G S G T Q Y T L T I S

██████████ → JNJ133 ← ██████████

PstI

AGC CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC TGT AAA CAG GCT TAT GAC GTT CCG TAC
 ▶ S L Q P E D F A T Y Y C K Q A Y D V P Y

██████████ → JNJ134 ← ██████████

JNJ135

EcORI

ACG TTC GGA CAA GGG ACC AAG GTG GAA ATC AAA CGTGAGTAGAATTAAAGAATTCCCC
 ▶ T F G Q G T K V E I K

██████████ → JNJ117 ← ██████████

JNJ136

██████████ → JNJ101 ← ██████████

Fig. 12

SpeI

ACTAGTACCACCATGAAATGCAGCTGGTTATCTTCTTCCTGATGGCAGTGGTTACAGGG
 M K C S W V I F F L M A V V T G

GTCAATTACAGGTTCAAGCTGGTGCACTCTGGGCAGAGGTGAAGAAGCCAGGGCCTCA
 V N S Q V Q L V Q S G A E V K K P G A S

GTCAAGGTTCTGCAAGGCTCTGGCTTCAACATTAAAGACACCTATGTGCACTGGTG
 V K V S C K A S G F N I K D T Y V H W V

CGCCAGGCCCTGGACAGAGGCTGGAGTGGATTGAAATATTGATCCTGCGAATGGTAAT
 R Q A P G Q R L E W I G N I D P A N G N

ACTAAATATGACCCGAAGTTCCAGGGCAAGGCCACTATAACAGCAGACACATCCGCGAGC
 T K Y D P K F Q G K A T I T A D T S A S

ACAGCCTACATGGAGCTCAGCAGCCTGAGATCTGAGGACACTGCCGTCTATTACTGTGCT
 T A Y M E L S S L R S E D T A V Y Y C A

AGATGGTTACGACATTTTACTATGCTATGGACTACTGGGTCAAGGAACCTGGTCACC
 R W L R H F Y Y A M D Y W G Q G T L V T

HindIII

GTCTCCTCAGGTGAGTCCTCACAAAGCTT
 V S S

Fig. 13

NheI

GCTAGCACCACCATGAGTGTGCCACTCAACTCCTGGGTTGCTGCTGTGGCTTACA
 M S V P T Q L L G L L L W L T

GACGCACGATGTGACATCCAGATGACTCAGTCTCCATCCTCCCTGTCTGCATCTGGGA
 D A R C D I Q M T Q S P S S L S A S V G

GACAGAGTCACCACATCACATGTCGAGCAAGTGAGAACATTTACTACAGTTAGCATGGTAT
 D R V T I T C R A S E N I Y Y S L A W Y

CAGCAGAAGCCAGGGAAAGCCCTAAGCTCCTGATCTATAATGCAAACAGCTTGGAAAGAT
 Q Q K P G K A P K L L I Y N A N S L E D

GGTGTCCCCTCGAGGTTCACTGGCAGTGGATCTGGACACAGTATACTCTCACCATCAGC
 G V P S R F S G S G S G T Q Y T L T I S

AGCCTGCAGCCTGAAGATTTGCAACTTATTACTGTAAACAGGCTTATGACGTTCCGTAC
 S L Q P E D F A T Y Y C K Q A Y D V P Y

ECORI

ACGTTGGACAAGGGACCAAGGTGAAATCAAACGTGAGTAGAATTAAAGAATTTC
T F G Q G T K V E I K


Fig. 14


Fig. 15



Fig. 16

Fig. 17

Cells : 1.0×10^6 /ml
 Ab : P1H4(10 μ g/ml)
 5A1, Y9A2, 1K11, 21C5, 24I11, 25B6(20 μ g/ml)

VCAM-1(1.25 μ g/ml)

[IU/L]





Fig. 19


Fig. 20aHuman $\alpha 9/\text{CHO-K1}$ 


Fig. 20bHuman α 4/CHO-K1


Fig. 20cNeutrophils


Fig. 21

Fig. 22

