

(19) **DANMARK**



Patent- og  
Varemærkestyrelsen

(12)

## Oversættelse af europæisk patentskrift

(10) **DK/EP 2100959 T3**

- 
- (51) Int.Cl.: *C 12 N 15/24 (2006.01)* *C 07 K 14/54 (2006.01)* *C 07 K 16/24 (2006.01)*  
*C 12 Q 1/68 (2006.01)* *G 01 N 33/68 (2006.01)*
- (45) Oversættelsen bekendtgjort den: **2017-01-30**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-12-07**
- (86) Europæisk ansøgning nr.: **09156327.0**
- (86) Europæisk indleveringsdag: **1998-07-24**
- (87) Den europæiske ansøgnings publiceringsdag: **2009-09-16**
- (30) Prioritet: **1997-07-25 US 900905**
- (62) Stamansøgningsnr: **98937105.9**
- (84) Designerede stater: **AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL PT SE**
- (73) Patenthaver: **Merck Sharp & Dohme Corp., 126 East Lincoln Avenue, Rahway NJ 07065-0907, USA**
- (72) Opfinder: **BAZAN, J. Fernando, 775 University Drive, Menlo Park, CA 94025, USA**
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Pattedyrscytokin: Interleukin-B30 og tilhørende reagenser**
- (56) Fremdragne publikationer:  
**EP-A1- 1 072 610**  
**WO-A-96/24676**  
**WO-A1-99/54357**  
**Y. ITO ET AL: "Pig STAT2 cDNA" EMBL DATABASE ENTRY SSAB4061, ACCESSION NUMBER AB004061, 3 June 1997 (1997-06-03), XP002085722**  
**J. TAKEDA: EMBL DATABASE ENTRY HSC3686, ACCESSION NUMBER C06368, 25 August 1996 (1996-08-25), XP002085721**  
**L. HILLIER ET AL: "WashU-Merck EST project 1997" EMBL DATABASE ENTRY HS1214377, ACCESSION NUMBER AA418955, 14 May 1997 (1997-05-14), XP002085720**  
**OPPMANN B ET AL: "Novel p19 Protein Engages IL-12p40 to Form a Cytokine, IL-23, with Biological Activities Similar as Well as Distinct from IL-12" IMMUNITY, CELL PRESS, US, vol. 13, no. 5, 1 November 2000 (2000-11-01), pages 715-725, XP002903120 ISSN: 1074-7613**  
**CUA D J ET AL: "Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain" NATURE, NATURE PUBLISHING GROUP, LONDON, UK, vol. 421, 13 February 2003 (2003-02-13), pages 744-748, XP002971969 ISSN: 0028-0836**  
**MURPHY C A ET AL: "Divergent pro- and antiinflammatory roles for IL-23 and IL-12 in joint autoimmune inflammation" JOURNAL OF EXPERIMENTAL MEDICINE, ROCKEFELLER UNIVERSITY PRESS, JP, vol. 198, no. 12, 15 December 2003 (2003-12-15), pages 1951-1957, XP002294202 ISSN: 0022-1007**  
**PITTI R M ET AL: "Induction of apoptosis by Apo-2 ligand, a new member of the tumor necrosis factor cytokine family", JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, US, vol. 271, no. 22, 31 May 1996 (1996-05-31), pages 12687-12690, XP003024616,**

Fortsættes ...

ISSN: 0021-9258, DOI: 10.1074/JBC.271.22.12687

DATABASE EMBL [Online] 27 January 1996 'Human clone JkA3 mRNA induced upon T-cell activation, 3' end.'  
Retrieved from EBI, accession no. EMBL:U38443 Database accession no. U38443

DATABASE EMBL [Online] 14 May 1997 'zw01c10.s1 Soares\_NhHMPu\_S1 Homo sapiens cDNA clone  
IMAGE:768018 3', mRNA sequence.' Retrieved from EBI, accession no. EMBL:AA418747 Database accession  
no. AA418747

DATABASE EMBL [Online] 09 January 1998 'nx39f06.s1 NCI\_CGAP\_GC4 Homo sapiens cDNA clone  
IMAGE:1258499 3', mRNA sequence.' Retrieved from EBI, accession no. EMBL:AA729815 Database accession  
no. AA729815

DATABASE EMBL [Online] 02 May 1994 'hbc2125 Human pancreatic islet Homo sapiens cDNA clone hbc2125  
5', mRNA sequence.' Retrieved from EBI, accession no. EMBL:T18577 Database accession no. T18577

## DESCRIPTION

### FIELD OF THE INVENTION

**[0001]** The present invention pertains to compositions related to proteins which function in controlling biology and physiology of mammalian cells, e.g., cells of a mammalian immune system. In particular, it provides purified genes, proteins, antibodies, and related reagents useful, e.g., to regulate activation, development, differentiation, and function of various cell types, including hematopoietic cells.

### BACKGROUND OF THE INVENTION

**[0002]** Recombinant DNA technology refers generally to the technique of integrating genetic information from a donor source into vectors for subsequent processing, such as through introduction into a host, whereby the transferred genetic information is copied and/or expressed in the new environment. Commonly, the genetic information exists in the form of complementary DNA (cDNA) derived from messenger RNA (mRNA) coding for a desired protein product. The carrier is frequently a plasmid having the capacity to incorporate cDNA for later replication in a host and, in some cases, actually to control expression of the cDNA and thereby direct synthesis of the encoded product in the host.

**[0003]** For some time, it has been known that the mammalian immune response is based on a series of complex cellular interactions, called the "immune network". Recent research has provided new insights into the inner workings of this network. While it remains clear that much of the response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines, play a critical role in controlling these cellular interactions. Thus, there is considerable interest in the isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which will lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system disorders. Some of these factors are hematopoietic growth factors, e.g., granulocyte colony stimulating factor (G-CSF). See, e.g., Thomson (1994; ed.) The Cytokine Handbook (2d ed.) Academic Press, San Diego; Metcalf and Nicola (1995) The Hematopoietic Colony Stimulating Factors Cambridge University Press; and Aggarwal and Gutterman (1991) Human Cytokines Blackwell Pub. Ito et al. (Pig STAT2 cDNA EMBL database entry SSAB4061, Accession No. AB004061, 1997) discloses a sequence for pig STAT2.

**[0004]** Lymphokines apparently mediate cellular activities in a variety of ways. They have been shown to support the proliferation, growth, and differentiation of pluripotential hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular lineages making up a complex immune system. Proper and balanced interactions between the cellular components are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

**[0005]** Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These lymphocytes interact with many other cell types.

**[0006]** Another important cell lineage is the mast cell (which has not been positively identified in all mammalian species), which is a granule-containing connective tissue cell located proximal to capillaries throughout the body. These cells are found in especially high concentrations in the lungs, skin, and gastrointestinal and genitourinary tracts. Mast cells play a central role in allergy-related disorders, particularly anaphylaxis as follows: when selected antigens crosslink one class of immunoglobulins bound to receptors on the mast cell surface, the mast cell degranulates and releases mediators, e.g., histamine, serotonin, heparin, and prostaglandins, which cause allergic reactions, e.g., anaphylaxis.

**[0007]** Research to better understand and treat various immune disorders has been hampered by the general inability to maintain cells of the immune system in vitro. Immunologists have discovered that culturing these cells can be accomplished through the use of T-cell and other cell supernatants, which contain various growth factors, including many of the lymphokines.

**[0008]** From the foregoing, it is evident that the discovery and development of new lymphokines, e.g., related to G-CSF and/or IL-6, could contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or indirectly involve the immune system and/or hematopoietic cells. In particular, the discovery and development of lymphokines which enhance or

potentiate the beneficial activities of known lymphokines would be highly advantageous. The present invention provides new interleukin compositions and related compounds, and methods for their use.

## **SUMMARY OF THE INVENTION**

**[0009]** The present invention is directed to mammalian, e.g., rodent, canine, feline, primate, interleukin-B30 (IL-B30) and its biological activities. Disclosed herein are nucleic acids coding for polypeptides themselves and methods for their production and use. The nucleic acids disclosed herein are characterized, in part, by their homology to cloned complementary DNA (cDNA) sequences enclosed herein, and/or by functional assays for growth factor- or cytokine-like activities, e.g., G-CSF (see Nagata (1994) in Thomson The Cytokine Handbook 2d ed., Academic Press, San Diego) and/or IL-6 (see Hirano (1994) in Thomson The Cytokine Handbook 2d ed., Academic Press, San Diego), applied to the polypeptides, which are typically encoded by these nucleic acids. Methods for modulating or intervening in the control of a growth factor dependent physiology or an immune response are disclosed.

**[0010]** The present invention is based, in part, upon the discovery of a new cytokine sequence exhibiting significant sequence and structural similarity to G-CSF and IL-6. In particular, disclosed herein is primate, e.g., human, gene encoding a protein whose mature size is about 168 amino acids, and pig and murine, e.g., mouse, sequences. Functional equivalents exhibiting significant sequence homology will be available from other mammalian, e.g., cow, horse, and rat, and non-mammalian species. the IL-B30. may comprises a mature sequence of Table 1; or may exhibits a post-translational modification pattern distinct from natural IL-B30; or the protein or peptide: may be from a warm blooded animal selected from a mammal, including a primate; may comprises at least one polypeptide segment of SEQ ID NO: 2; may exhibits a plurality of portions exhibiting the identity; may be a natural allelic variant of IL-B30; may have a length at least about 30 amino acids; may exhibits at least two non-overlapping epitopes which are specific for a mammalian IL-B30; may exhibits a sequence identity at least about 90% over a length of at least about 20 amino acids to mammalian IL-B30; may be glycosylated; may have a molecular weight of at least 10 kD with natural glycosylation; may be a synthetic polypeptide; may be attached to a solid substrate; is conjugated to another chemical moiety; may be a 5-fold or less substitution from natural sequence; or may be a deletion or insertion variant from a natural sequence. Disclosed herein is a composition comprising: a sterile IL-B30 protein or peptide; or the IL-B30 protein or peptide and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration. Disclosed herein are fusion proteins in which, the protein can have: mature protein sequence of Table 1; a detection or purification tag, including a FLAG, His6, or Ig sequence; and/or sequence of another cytokine or chemokine.

**[0011]** Disclosed herein are binding compounds in which the compound may have an antigen binding site from an antibody, which specifically binds to a natural IL-B30 protein, wherein: the IL-B30 may be a mammalian protein; the binding compound may be an Fv, Fab, or Fab2 fragment; the binding compound may be conjugated to another chemical moiety; or the antibody: may be raised against a peptide sequence of a mature polypeptide of Table 1; may be raised against a mature IL-B30; may be raised to a purified rodent IL-B30; may be immunoselected; may be a polyclonal antibody; may binds to a denatured IL-B30, may exhibits a Kd of at least 30  $\mu$ M; may be attached to a solid substrate, including a bead or plastic membrane; may be in a sterile composition; or may be detectably labeled, including a radioactive or fluorescent label. Disclosed herein are kits containing binding compounds including those with: a compartment comprising the binding compound; and/or instructions for use or disposal of reagents in the kit. Often the kit may be capable of making a qualitative or quantitative analysis. Compositions may comprise: a sterile binding compound; or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

**[0012]** The invention is has defined in that claims.

## **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

**[0013]** All references cited herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

### **I. General**

**[0014]** Disclosed herein are amino acid sequences and DNA sequences encoding various mammalian proteins which are cytokines, e.g., which are secreted molecules which can mediate a signal between immune or other cells. See, e.g., Paul (1994)

Fundamental Immunology (3d ed.) Raven Press, N.Y. The full length cytokines, and fragments, or antagonists will be useful in physiological modulation of cells expressing a receptor. It is likely that IL-B30 has either stimulatory or inhibitory effects on hematopoietic cells, including, e.g., lymphoid cells, such as T-cells, B-cells, natural killer (NK) cells, macrophages, dendritic cells, hematopoietic progenitors, etc. The proteins will also be useful as antigens, e.g., immunogens, for raising antibodies to various epitopes on the protein, both linear and conformational epitopes.

[0015] A cDNA encoding IL-B30 was identified from a human cell line. The molecule was designated huIL-B30. A related gene corresponding to a pig sequence was also identified. A rodent sequence, e.g., from mouse, is also described.

[0016] The human gene encodes a small soluble cytokine-like protein, of about 168 amino acids. The signal sequence probably is about 21 residues, and would run from the Met to about Ala. See Table 1 and SEQ. ID. NO: 1 and 2. IL-B30 exhibits structural motifs characteristic of a member of the long chain cytokines. Compare, e.g., IL-B30, G-CSF, and IL-6, sequences available from GenBank. See also Table 2.

**Table 1: Nucleic acid (SEQ ID NO: 1) encoding IL-B30 from a primate, e.g., human. Translated amino acid sequence is SEQ ID NO: 2.**

[0017]

```

ATG CTG GGG AGC AGA GCT GTA ATG CTG CTG TTG CTG CTG CCC TGG ACA
48
Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr
-21 -20 -15 -10

GCT CAG GGC AGA GCT GTG CCT GGG GGC AGC AGC CCT GCC TGG ACT CAG
96
Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln
-5 1 5 10

TGC CAG CAG CTT TCA CAG AAG CTC TGC ACA CTG GCC TGG AGT GCA CAT
144
Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His
15 20 25

CCA CTA GTG GGA CAC ATG GAT CTA AGA GAA GAG GGA GAT GAA GAG ACT
192
Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr
30 35 40

ACA AAT GAT GTT CCC CAT ATC CAG TGT GGA GAT GGC TGT GAC CCC CAA
240
Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln
45 50 55

GGA CTC AGG GAC AAC AGT CAG TTC TGC TTG CAA AGG ATC CAC CAG GGT
288
Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly
60 65 70 75

CTG ATT TTT TAT GAG AAG CTG CTA GGA TCG GAT ATT TTC ACA GGG GAG
336
Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu
80 85 90

CCT TCT CTG CTC CCT GAT AGC CCT GTG GCG CAG CTT CAT GCC TCC CTA
384
Pro Ser Leu Leu Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu
95 100 105

CTG GGC CTC AGC CAA CTC CTG CAG CCT GAG GGT CAC CAC TGG GAG ACT
432
Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr
110 115 120

CAG CAG ATT CCA AGC CTC AGT CCC AGC CAG CCA TGG CAG CGT CTC CTT
480
Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu
125 130 135

CTC CGC TTC AAA ATC CTT CGC AGC CTC CAG GCC TTT GTG GCT GTA GCC
528
Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala
140 145 150 155

GCC CGG GTC TTT GCC CAT GGA GCA GCA ACC CTG AGT CCC TAA
570
Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro
160 165

```

coding sequence:

ATGCTGGGGA GCAGAGCTGT AATGCTGCTG TTGCTGCTGC CCTGGACAGC  
 TCAGGGCAGA GCTGTGCCTG GGGGCAGCAG CCTGCTCTGG ACTCAGTGCC  
 AGCAGCTTTC ACAGAAGCTC TGCACACTGG CCTGGAGTGC ACATCCACTA  
 GTGGGACACA TGGATCTAAG AGAAGAGGGA GATGAAGAGA CTACAAATGA  
 TGTTCCCAT ATCCAGTGTG GAGATGGCTG TGACCCCAAA GGACTCAGGG  
 ACAACAGTCA GTTCTGCTTG CAAAGGATCC ACCAGGGTCT GATTTTTTAT  
 GAGAAGCTGC TAGGATCGGA TATTTTCACA GGGGAGCCTT CTCGTCTCCC  
 TGATAGCCCT GTGGCGCAGC TTCATGCCCT CCTACTGGGC CTCAGCCAAC  
 TCCTGCAGCC TGAGGGTCAC CACTGGGAGA CTCAGCAGAT TCCAAGCCTC  
 AGTCCCAGCC AGCCATGGCA GCGTCTCCTT CTCGCTTCA AAATCCCTCG  
 CAGCCTCCAG GCCTTTGTGG CTGTAGCCGC CCGGCTCTTT GCCCATGGAG  
 CAGCAACCTT GAGTCCCTAA

Rodent, e.g., mouse, IL-B30 (SEQ ID NO: 3 and 4):

CGCTTAGAAG TCGGACTACA GAGTTAGACT CAGAACCAA GGAGGTGGAT AGGGGGTCCA 60  
 CAGGCCTGCT GCAGATCACA GAGCCAGCCA GATCTGAGAA GCAGGGAACA AG ATG 115  
 Met  
 -21  
 CTG GAT TGC AGA GCA GTA ATA ATG CTA TGG CTG TTG CCC TGG GTC ACT 163  
 Leu Asp Cys Arg Ala Val Ile Met Leu Trp Leu Leu Pro Trp Val Thr  
 -20 -15 -10 -5  
 CAG GGC CTG GCT GTG CCT AGG AGT AGC AGT CCT GAC TGG GCT CAG TGC 211  
 Gln Gly Leu Ala Val Pro Arg Ser Ser Ser Pro Asp Trp Ala Gln Cys  
 1 5 10  
 CAG CAG CTC TCT CGG AAT CTC TGC ATG CTA GCC TGG AAC GCA CAT GCA 259  
 Gln Gln Leu Ser Arg Asn Leu Cys Met Leu Ala Trp Asn Ala His Ala  
 15 20 25  
 CCA GCG GGA CAT ATG AAT CTA CTA AGA GAA GAA GAG GAT GAA GAG ACT 307  
 Pro Ala Gly His Met Asn Leu Leu Arg Glu Glu Glu Asp Glu Glu Thr  
 30 35 40  
 AAA AAT AAT GTG CCC CGT ATC CAG TGT GAA GAT GGT TGT GAC CCA CAA 355  
 Lys Asn Asn Val Pro Arg Ile Gln Cys Glu Asp Gly Cys Asp Pro Gln  
 45 50 55 60  
 GGA CTC AAG GAC AAC AGC CAG TTC TGC TTG CAA AGG ATC CGC CAA GGT 403  
 Gly Leu Lys Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile Arg Gln Gly  
 65 70 75  
 CTG GCT TTT TAT AAG CAC CTG CTT GAC TCT GAC ATC TTC AAA GGG GAG 451  
 Leu Ala Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys Gly Glu  
 80 85 90  
 CCT GCT CTA CTC CCT GAT AGC CCC ATG GAG CAA CTT CAC ACC TCC CTA 499  
 Pro Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr Ser Leu  
 95 100 105  
 CTA GGA CTC AGC CAA CTC CTC CAG CCA GAG GAT CAC CCC CGG GAG ACC 547  
 Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg Glu Thr  
 110 115 120  
 CAA CAG ATG CCC AGC CTG AGT TCT AGT CAG CAG TGG CAG CGC CCC CTT 595  
 Gln Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg Pro Leu  
 125 130 135 140  
 CTC CGT TCC AAG ATC CTT CGA AGC CTC CAG GCC TTT TTG GCC ATA GCT 643  
 Leu Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala Ile Ala  
 145 150 155  
 GCC CGG GTC TTT GCC CAC GGA GCA GCA ACT CTG ACT GAG CCC TTA GTG 691  
 Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro Leu Val  
 160 165 170  
 CCA ACA GCT TAAGGATGCC CAGGTTCCCA TGGCTACCAT GATAAGACTA 740  
 Pro Thr Ala  
 175  
 ATCTATCAGC CCAGACATCT ACCAGTTAAT TAACCCATTA GGACTTGTGC TGTCTTGTGTT 800  
 TCGTTTGTGTT TGCCTGAAGG GCAAGGACAC CATTATTAAA GAGAAAAGAA ACAAAACCCA 860  
 GAGCAGGCAG CTGGCTAGAG AAAGGAGCTG GAGAAGAAGA ATAAAGTCTC GAGCCCTTGG 920  
 CCTTGAAGC GGGCAAGCAG CTGCGTGGCC TGAGGGGAAG GGGGCGGTGG CATCGAGAAA 980  
 CTGTGAGAAA ACCCAGAGCA TCAGAAAAAG TGAGCCCAGG CTTTGGCCAT TATCTGTAAG 1040  
 AAAAACAAGA AAAGGGGAAC ATTATACCTT CCTGGGTGGC TCAGGGAAAT GTGCAGATGC 1100  
 ACAGTACTCC AGACAGCAGC TCTGTACCTG CCTGCTCTGT CCCTCAGTTC TAACAGAATC 1160  
 TAGTCACTAA GAACCTAACAG GACTACCAAT ACGAAGTGC AAA 1203

Table 2: Comparison of various IL-6 and G-CSF variants compared to IL-B30. Human IL-B30 is SEQ ID NO: 2; mouse IL-B30 is SEQ ID NO: 4; pig IL-B30 is SEQ ID NO: 5; bovine G-CSF is SEQ ID NO: 6; feline G-CSF is SEQ ID NO: 7; human G-CSF is SEQ ID NO: 8; mouse G-CSF is SEQ ID NO: 9; otter IL-6 is SEQ ID NO: 10; feline IL-6 is SEQ ID NO: 11; human IL-6 is SEQ ID NO: 12; sheep IL-6 is SEQ ID NO: 13; mouse IL-6 is SEQ ID NO: 14; chicken MGF is SEQ ID NO: 15; and KSHV, kaposi's sarcoma herpes virus, a viral IL-6, is SEQ ID NO: 16.

|             |             |            |            |            |
|-------------|-------------|------------|------------|------------|
| il130_human | .....VPGG   | SSPVWTQCQQ | LSQKLCT.LA | WSAHPLVG.. |
| il130_mouse | .....VPRS   | SSPDWAQCQQ | LSRNLCM.LA | WNAHAPAG.. |
| il130_pig   | .....       | .....      | .....      | .....      |
| gcsf_bovin  | .....TPLG   | P.....AR   | SLPQSFLKLC | LEQVRKIQAD |
| gcsf_felca  | .....TPLG   | P.....TS   | SLPQSFLKLC | LEQVRKIQAD |
| gcsf_human  | .....TPLG   | P.....AS   | SLPQSFLKLC | LEQVRKIQGD |
| gcsf_mouse  | VPLVTVSALP  | P.....SL   | PLPRSFLKLS | LEQVRKIQAS |
| il6_otter   | .AFPTPGPLG  | GDSKDDATSN | RPPLTSADRM | EDFIKFIKLG |
| il6_felca   | .AFPTPGPLG  | G....DATSN | RLPLTPADRM | EELIKYILGK |
| il6_human   | .AFPAPVPPG  | EDSKDVAAPH | RQPLTSSERI | DKQIRYILDG |
| il6_sheep   | .AFPTPGPLG  | EDFKNDTPS  | RLLLTTPKPT | EALIKHIVDK |
| il6_mouse   | .AFPTSQVRR  | GDFTEDTTPN | R.PVYTTSQV | GGLITHVLWE |
| mgf_chick   | .....       | .APLAELSGD | HDFQLFLHKN | LEFTRKIRGD |
| il6_khsv    | .....       | .....TRG   | KLPDAPEFEK | DLIIQRLNWM |
| il130_human | .HMD.LREEG  | DEETNDVPH  | I...QCGDGC | DPQGLRDNQS |
| il130_mouse | .HMNLLREEE  | DEETKNNVPR | I...QCEDGC | DPQGLKDNQS |
| il130_pig   | .....       | .....      | .....      | .....      |
| gcsf_bovin  | .CAA.HKLCH  | PEELMLLRHS | LGIP.QAPLS | SCSSQSLLQR |
| gcsf_felca  | .CAA.HKLCH  | PEELVLLGHA | LGIP.QAPLS | SCSSQALQLT |
| gcsf_human  | .ECAT.YKLCH | PEELVLLGHS | LGIP.WAPLS | SCPSQALQLA |
| gcsf_mouse  | .CAT.YKLCH  | PEELVLLGHS | LGIP.KASLS | GCSSQALQQT |
| il6_otter   | .CDK.YNKCE  | DSKEVLAENN | LNLPLKAEKD | RCFQSGFNQE |
| il6_felca   | .CDN.YNKCE  | DSKEALAENN | LNLPLKAEKD | RCFQSGFNQE |
| il6_human   | .CNK.SNMCE  | SSKEALAENN | LNLPKMAEKD | RCFQSGFNQE |
| il6_sheep   | .CEK.NDECE  | NSKETLAENK | LKLPMKEEKD | RCFQSGFNQA |
| il6_mouse   | .CNG.NSDCM  | NNDDALAENN | LKLPEIQRND | GCYQTGYNQE |
| mgf_chick   | .CDT.FQLCT  | EEELQLVQPD | PHLV.QAPLD | QCHKRGFQAE |
| il6_khsv    | LCYR.TGICK  | GILEPAATFH | LKLPAINDTD | HCGLIGFNET |
| il130_human | IFYEKLLGSD  | IFTGE..... | .PSLLPDSFV | AQLHASLLGL |
| il130_mouse | AFYKHLLDSD  | IFKGE..... | .PALLPDSFM | EQHLSLLGL  |
| il130_pig   | VFYEKLLGSD  | IFTGE..... | .PSLHPDGSV | GQLHASLLGL |
| gcsf_bovin  | FLYQGLLQAL  | AGIS.....  | .PELAPTLT  | LQLDVDFAT  |
| gcsf_felca  | FLYQGLLQAL  | AGIS.....  | .PELAPTLT  | LQLDITDFAI |
| gcsf_human  | FLYQGLLQAL  | EGIS.....  | .PELAPTLT  | LQLDVDFAT  |
| gcsf_mouse  | CLYQGLLQAL  | SGIS.....  | .PALAPTLT  | LQLDVANFAT |
| il6_otter   | QEFQIHLKYL  | ESNYEG...N | KDNAHSVYIS | TKHLIQLTRP |
| il6_felca   | QEFQIYLKFL  | QDKYEG...D | KENAKSVYTS | TNVLLQMLKR |
| il6_human   | LEFEVYLEYL  | QNRFE...S  | EEQARAVQMS | TKVLIQFLQK |
| il6_sheep   | LEYQIYLDLFL | QNEFEG...N | QETVMEQSS  | IRTLIQILKE |
| il6_mouse   | LEYHSYLEYM  | KNNLKDN..K | KDKARVLQED | TETLIHIFNQ |
| mgf_chick   | HAYHDSLGAV  | LRLLP....  | ..NHTTLVET | LQLDANLSS  |
| il6_khsv    | FEFEVLKFL   | TTEFGKSVIN | VDVMELLTKT | LGWDIQEELN |

Table 2 (continued)

|             |            |             |            |             |
|-------------|------------|-------------|------------|-------------|
| il130_human | HHWETQIP.  | .SLSPSQ..P  | WQRLRLRFKI | LRSLQAFVAV  |
| il130_mouse | HPRETQOMP. | .SLSSSQ..Q  | WQRLPLRSKI | LRSLQAFVAV  |
| il130_pig   | HHWETQTP.  | .SPSPSQ..P  | WQRLRLRLKI | LRSLQAFVAV  |
| gcsf_bovin  | AAPAVQPTQ. | .GAMPTTSA   | FQRRAGGVLV | ASQLHRFLEL  |
| gcsf_felca  | MAPAVPPTQ. | .GTMPPTSA   | FQRRAGGTLV | ASNLSQSFLEV |
| gcsf_human  | MAPALQPTQ. | .GAMPFASA   | FQRRAGGVLV | ASHLSQSFLEV |
| gcsf_mouse  | VAPTVPPTQ. | .SAMPAPTSA  | FQRRAGGVLA | ISYLGQFLET  |
| il6_otter   | PDPTTASL.  | .QALFKSQDK  | WLKHTTIHLI | LRRLLEDLQF  |
| il6_felca   | PVPTVEVGL. | .QLSCSHR.R  | VAEAHNNHLT | LRRLLEDLQF  |
| il6_human   | PDPTTNASL. | .LTKLQAQNO  | WLQDMTTHLI | LRSLFKEFLQS |
| il6_sheep   | TPPATHTDM. | .LEKMQSSNE  | WVKNAKVIII | LRSLNENFLQF |
| il6_mouse   | PTPISNALL. | .TDKLESQKE  | WLRTKTIQFI | LKSLEEFKLV  |
| mgf_chick   | LDTVTLPAEQ | RSPPPTFSGP  | FQQQVGGFPI | LANFQRFLET  |
| il6_khsv    | P.PKFDGR.  | .LLGRLQGLKY | WVRHFASFYV | LSAMEKFAGQ  |
| il130_human | TLSP....   | .....       | .....      | .....       |
| il130_mouse | TLTEPLVPTA | .....       | .....      | .....       |
| il130_pig   | TLSP....   | .....       | .....      | .....       |
| gcsf_bovin  | P.....     | .....       | .....      | .....       |
| gcsf_felca  | P.....     | .....       | .....      | .....       |
| gcsf_human  | P.....     | .....       | .....      | .....       |
| gcsf_mouse  | .....      | .....       | .....      | .....       |
| il6_otter   | .....      | .....       | .....      | .....       |
| il6_felca   | .....      | .....       | .....      | .....       |
| il6_human   | .....      | .....       | .....      | .....       |
| il6_sheep   | .....      | .....       | .....      | .....       |
| il6_mouse   | .....      | .....       | .....      | .....       |
| mgf_chick   | L.....     | .....       | .....      | .....       |
| il6_khsv    | VTPDVHDK   | .....       | .....      | .....       |

[0018] The structural homology of IL-B30 to related cytokine proteins suggests related function of this molecule. IL-B30 is a long chain cytokine exhibiting sequence similarity to IL-6 and G-CSF.

**[0019]** IL-B30 agonists, or antagonists, may also act as functional or receptor antagonists, e.g., which block IL-6 or G-CSF binding to their respective receptors, or mediating the opposite actions. Thus, IL-B30, or its antagonists, may be useful in the treatment of abnormal medical conditions, including immune disorders, e.g., T cell immune deficiencies, chronic inflammation, or tissue rejection, or in cardiovascular or neurophysiological conditions.

**[0020]** The natural antigens are capable of mediating various biochemical responses which lead to biological or physiological responses in target cells. The variant characterized herein is from human, but other primate, or other species counterparts exist in nature. Additional sequences for proteins in other mammalian species, e.g., primates, canines, felines, and rodents, should also be available. See below. The descriptions below are directed, for exemplary purposes, to a human IL-B30, but are likewise applicable to related variants from other species.

## **II. Purified IL-B30**

**[0021]** Human IL-B30 amino acid sequence, is disclosed herein within SEQ ID NO: 2. Other naturally occurring nucleic acids which encode the protein can be isolated by standard procedures using the provided sequence, e.g., PCR techniques, or by hybridization. These amino acid sequences, provided amino to carboxy, are important in providing sequence information for the cytokine allowing for distinguishing the protein antigen from other proteins and exemplifying numerous variants. Moreover, the peptide sequences allow preparation of peptides to generate antibodies to recognize such segments, and nucleotide sequences allow preparation of oligonucleotide probes, both of which are strategies for detection or isolation, e.g., cloning, of genes encoding such sequences.

**[0022]** As used herein, the term "human soluble IL-B30" shall encompass, when used in a protein context, a protein having amino acid sequence corresponding to a soluble polypeptide shown in SEQ ID NO: 2, or significant fragments thereof. Disclosed herein are a plurality of distinct, e.g., nonoverlapping, segments of the specified length. Typically, the plurality will be at least two, more usually at least three, and preferably 5, 7, or even more. While the length minima are provided, longer lengths, of various sizes, may be appropriate, e.g., one of length 7, and two of length 12.

**[0023]** Binding components, e.g., antibodies, typically bind to an IL-B30 with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. Counterpart proteins will be found in mammalian species other than human, e.g., other primates, ungulates, or rodents. Non-mammalian species should also possess structurally or functionally related genes and proteins, e.g., birds or amphibians.

**[0024]** The term "polypeptide" as used herein includes a significant fragment or segment, and encompasses a stretch of amino acid residues of at least about 8 amino acids, generally at least about 12 amino acids, typically at least about 16 amino acids, preferably at least about 20 amino acids, and, particularly at least about 30 or more amino acids, e.g., 35, 40, 45, 50, etc. Such fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at residues 1, 2, 3, etc., and ending at, e.g., 150, 149, 148, etc., in all practical combinations. Particularly interesting peptides have ends corresponding to structural domain boundaries, e.g., helices A, B, C, and/or D. See Table 1.

**[0025]** The term "binding composition" refers to molecules that bind with specificity to IL-B30, e.g., in an antibody-antigen interaction. The specificity may be more or less inclusive, e.g., specific to a particular variant, or to groups of related variants, e.g., primate, rodent, etc. It also includes compounds, e.g., proteins, which specifically associate with IL-B30, including in a natural physiologically relevant protein-protein interaction, either covalent or non-covalent. The molecule may be a polymer, or chemical reagent. A functional analog may be a protein with structural modifications, or it may be a molecule which has a molecular shape which interacts with the appropriate binding determinants. The compounds may serve as agonists or antagonists of a receptor binding interaction, see, e.g., Goodman, et al. (eds.) Goodman & Gilman's: The Pharmacological Bases of Therapeutics (current ed.) Pergamon Press.

**[0026]** Substantially pure, e.g., in a protein context, typically means that the protein is free from other contaminating proteins, nucleic acids, or other biologicals derived from the original source organism. Purity may be assayed by standard methods, typically by weight, and will ordinarily be at least about 40% pure, generally at least about 50% pure, often at least about 60% pure, typically at least about 80% pure, preferably at least about 90% pure, and in particular, at least about 95% pure. Carriers or excipients will often be added.

**[0027]** Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters affect

polypeptide solubility, including temperature, electrolyte environment, size and molecular characteristics of the polypeptide, and nature of the solvent. Typically, the temperature at which the polypeptide is used ranges from about 4° C to about 65° C. Usually the temperature at use is greater than about 18° C. For diagnostic purposes, the temperature will usually be about room temperature or warmer, but less than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C for humans and mice, though under certain situations the temperature may be raised or lowered in situ or in vitro.

**[0028]** The size and structure of the polypeptide should generally be in a substantially stable state, and usually not in a denatured state. The polypeptide may be associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents.

**[0029]** The solvent and electrolytes will usually be a biologically compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological aqueous solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, one or more detergents will be added, typically a mild non-denaturing one, e.g., CHS (cholesteryl hemisuccinate) or CHAPS (3-[3-cholamidopropyl]dimethylammonio]-1-propane sulfonate), or a low enough concentration as to avoid significant disruption of structural or physiological properties of the protein. In other instances, a harsh detergent may be used to effect significant denaturation.

### III. Physical Variants

**[0030]** Disclosed herein are proteins or peptides having substantial amino acid sequence identity with the amino acid sequence of the IL-B30 antigen. The variants include species, polymorphic, or allelic variants.

**[0031]** Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. See also Needleham, et al. (1970) J. Mol. Biol. 48:443-453; Sankoff, et al. (1983) Chapter One in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group, Madison, WI. Sequence identity changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. The conservation may apply to biological features, functional features, or structural features. Homologous amino acid sequences are typically intended to include natural polymorphic or allelic and interspecies variations of a protein sequence. Typical homologous proteins or peptides will have from 25-100% identity (if gaps can be introduced), to 50-100% identity (if conservative substitutions are included) with the amino acid sequence of the IL-B30. Identity measures will be at least about 35%, generally at least about 40%, often at least about 50%, typically at least about 60%, usually at least about 70%, preferably at least about 80%, and more preferably at least about 90%.

**[0032]** The isolated IL-B30 DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of short nucleotide stretches. These modifications result in novel DNA sequences which encode these antigens, their derivatives, or proteins having similar physiological, immunogenic, antigenic, or other functional activity. These modified sequences can be used to produce mutant antigens or to enhance expression. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. "Mutant IL-B30" encompasses a polypeptide otherwise falling within the sequence identity definition of the IL-B30 as set forth above, but having an amino acid sequence which differs from that of IL-B30 as normally found in nature, whether by way of deletion, substitution, or insertion. This generally includes proteins having significant identity with a protein having sequence of SEQ ID NO: 2, and as sharing various biological activities, e.g., antigenic or immunogenic, with those sequences, and in particular contain most of the natural full length disclosed sequences. Full length sequences will typically be preferred, though truncated versions will also be useful, likewise, genes or proteins found from natural sources are typically most desired. Similar concepts apply to different IL-B30 proteins, particularly those found in various warm blooded animals, e.g., mammals and birds. These descriptions are generally meant to encompass all IL-B30 proteins, not limited to the particular primate variants specifically discussed.

**[0033]** IL-B30 mutagenesis can also be conducted by making amino acid insertions or deletions. Substitutions, deletions, insertions, or any combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy- terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or polymerase chain reaction (PCR) techniques. See, e.g., Sambrook, et al. (1989); Ausubel, et al. (1987 and Supplements); and Kunkel, et al. (1987) Methods in Enzymol. 154:367-382. Disclosed herein

are e.g., 1-fold, 2-fold, 3-fold, 5-fold, 7-fold, etc., preferably conservative substitutions at the nucleotide or amino acid levels. Preferably the substitutions will be away from the conserved cysteines, and often will be in the regions away from the helical structural domains. Such variants may be useful to produce specific antibodies, and often will share many or all biological properties.

**[0034]** Disclosed herein are recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. A similar concept applies to heterologous nucleic acid sequences.

**[0035]** In addition, new constructs may be made from combining similar functional domains from other proteins. For example, target-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) *Science* 243:1330-1336; and O'Dowd, et al. (1988) *J. Biol. Chem.* 263:15985-15992.

**[0036]** The phosphoramidite method described by Beaucage and Carruthers (1981) *Tetra. Letts.* 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence, e.g., PCR techniques.

**[0037]** Structural analysis can be applied to this gene, in comparison to the IL-6 family of cytokines. The family includes, e.g., IL-6, IL-11, IL-12, G-CSF, LIF, OSM, CNTF, and Ob. Alignment of the human, pig, and mouse IL-B30 sequences with other members of the IL-6 family should allow definition of structural features. In particular,  $\beta$ -sheet and  $\alpha$ -helix residues can be determined using, e.g., RASMOL program, see Bazan, et al. (1996) *Nature* 379:591; Lodi, et al. (1994) *Science* 263:1762-1766; Sayle and Milner-White (1995) *TIBS* 20:374-376; and Gronenberg, et al. (1991) *Protein Engineering* 4:263-269. Preferred residues for substitutions include the surface exposed residues which would be predicted to interact with receptor. Other residues which should conserve function will be conservative substitutions, particularly at position far from the surface exposed residues.

#### **IV. Functional Variants**

**[0038]** The blocking of physiological response to IL-B30s may result from the competitive inhibition of binding of the ligand to its receptor.

**[0039]** In vitro assays disclosed herein will often use isolated protein, soluble fragments comprising receptor binding segments of these proteins, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or cytokine mutations and modifications, e.g., IL-B30 analogs.

**[0040]** Disclosed herein is the use of competitive drug screening assays, e.g., where neutralizing antibodies to the cytokine, or receptor binding fragments compete with a test compound.

**[0041]** "Derivatives" of IL-B30 antigens include amino acid sequence mutants from naturally occurring forms, glycosylation variants, and covalent or aggregate conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in IL-B30 amino acid side chains or at the N- or C- termini, e.g., by standard means. See, e.g., Lundblad and Noyes (1988) *Chemical Reagents for Protein Modification*, vols. 1-2, CRC Press, Inc., Boca Raton, FL; Hugli (ed. 1989) *Techniques in Protein Chemistry*, Academic Press, San Diego, CA; and Wong (1991) *Chemistry of Protein Conjugation and Cross Linking*, CRC Press, Boca Raton, FL.

**[0042]** In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. See, e.g., Elbein (1987) *Ann. Rev. Biochem.* 56:497-534. Also embraced are versions of the peptides with the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

**[0043]** Fusion polypeptides between IL-B30s and other homologous or heterologous proteins are also disclosed. Many cytokine receptors or other surface proteins are multimeric, e.g., homodimeric entities, and a repeat construct may have various advantages, including lessened susceptibility to proteolytic cleavage. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a protein, e.g., a receptor-binding segment, so that the presence or location of the fused ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609. Other gene fusion partners include bacterial  $\beta$ -galactosidase, trpE, Protein A,  $\beta$ -lactamase, alpha amylase, alcohol dehydrogenase, yeast alpha mating factor, and detection or

purification tags such as a FLAG sequence or His6 sequence. See, e.g., Godowski, et al. (1988) *Science* 241:812-816.

**[0044]** Fusion peptides will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, e.g., in Sambrook, et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed.), vols. 1-3, Cold Spring Harbor Laboratory; and Ausubel, et al. (eds. 1993) *Current Protocols in Molecular Biology*, Greene and Wiley, NY. Techniques for synthesis of polypeptides are described, e.g., in Merrifield (1963) *J. Amer. Chem. Soc.* 85:2149-2156; Merrifield (1986) *Science* 232: 341-347; Atherton, et al. (1989) *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, Oxford; and Grant (1992) *Synthetic Peptides: A User's Guide*, W.H. Freeman, NY. Refolding methods may be applicable to synthetic proteins.

**[0045]** Disclosed herein is the use of derivatives of IL-B30 proteins other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties or protein carriers. Covalent or aggregative derivatives will be useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of binding partners, e.g., other antigens. An IL-B30 can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated SEPHAROSE, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-IL-B30 antibodies or an alternative binding composition. The IL-B30 proteins can also be labeled with a detectable group, e.g., for use in diagnostic assays. Purification of IL-B30 may be effected by an immobilized antibody or complementary binding partner, e.g., binding portion of a receptor.

**[0046]** A solubilized IL-B30 or fragment disclosed herein can be used as an immunogen for the production of antisera or antibodies specific for binding. Purified antigen can be used to screen monoclonal antibodies or antigen-binding fragments, encompassing antigen binding fragments of natural antibodies, e.g., Fab, Fab', F(ab)<sub>2</sub>, etc. Purified IL-B30 antigens can also be used as a reagent to detect antibodies generated in response to the presence of elevated levels of the cytokine, which may be diagnostic of an abnormal or specific physiological or disease condition. Disclosed herein are antibodies raised against amino acid sequences encoded by nucleotide sequence shown in SEQ ID NO: 1, or fragments of proteins containing it. In particular, disclosed herein are antibodies having binding affinity to or being raised against specific domains, e.g., helices A, B, C, or D.

**[0047]** Disclosed herein is the isolation of additional closely related species variants. Southern and Northern blot analysis will establish that similar genetic entities exist in other mammals. It is likely that IL-B30s are widespread in species variants, e.g., rodents, lagomorphs, carnivores, artiodactyla, perissodactyla, and primates.

**[0048]** Disclosed herein are means to isolate a group of related antigens displaying both distinctness and similarities in structure, expression, and function. Elucidation of many of the physiological effects of the molecules will be greatly accelerated by the isolation and characterization of additional distinct species or polymorphic variants of them. In particular, disclosed herein are useful probes for identifying additional homologous genetic entities in different species.

**[0049]** The isolated genes will allow transformation of cells lacking expression of an IL-B30, e.g., either species types or cells which lack corresponding proteins and exhibit negative background activity. This should allow analysis of the function of IL-B30 in comparison to untransformed control cells.

**[0050]** Dissection of critical structural elements which effect the various physiological functions mediated through these antigens is possible using standard techniques of modern molecular biology, particularly in comparing members of the related class. See, e.g., the homolog-scanning mutagenesis technique described in Cunningham, et al. (1989) *Science* 243:1339-1336; and approaches used in O'Dowd, et al. (1988) *J. Biol. Chem.* 263:15985-15992; and Lechleiter, et al. (1990) *EMBO J.* 9:4381-4390.

**[0051]** Intracellular functions would probably involve receptor signaling. However, protein internalization may occur under certain circumstances, and interaction between intracellular components and cytokine may occur. Specific segments of interaction of IL-B30 with interacting components may be identified by mutagenesis or direct biochemical means, e.g., cross-linking or affinity methods. Structural analysis by crystallographic or other physical methods will also be applicable. Further investigation of the mechanism of signal transduction will include study of associated components which may be isolatable by affinity methods or by genetic means, e.g., complementation analysis of mutants.

**[0052]** Further study of the expression and control of IL-B30 will be pursued. The controlling elements associated with the antigens should exhibit differential physiological, developmental, tissue specific, or other expression patterns. Upstream or downstream genetic regions, e.g., control elements, are of interest.

[0053] Structural studies of the IL-B30 antigens will lead to design of new antigens, particularly analogs exhibiting agonist or antagonist properties on the molecule. This can be combined with previously described screening methods to isolate antigens exhibiting desired spectra of activities.

## V. Antibodies

[0054] Antibodies can be raised to various epitopes of the IL-B30 proteins, including species, polymorphic, or allelic variants, and fragments thereof, both in their naturally occurring forms and in their recombinant forms. Additionally, antibodies can be raised to IL-B30s in either their active forms or in their inactive forms, including native or denatured versions. Anti-idiotypic antibodies are also contemplated.

[0055] Antibodies, including binding fragments and single chain versions, against predetermined fragments of the antigens can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective IL-B30s, or screened for agonistic or antagonistic activity, e.g., mediated through a receptor. Antibodies may be agonistic or antagonistic, e.g., by sterically blocking binding to a receptor. These monoclonal antibodies will usually bind with at least a  $K_D$  of about 1 mM, more usually at least about 300  $\mu$ M, typically at least about 100  $\mu$ M, more typically at least about 30  $\mu$ M, preferably at least about 10  $\mu$ M, and more preferably at least about 3  $\mu$ M or better.

[0056] The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they can be screened for ability to bind to the antigens without inhibiting binding to a receptor. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying IL-B30 protein or its receptors. See, e.g., Chan (ed. 1987) *Immunology: A Practical Guide*, Academic Press, Orlando, FL; Price and Newman (eds. 1991) *Principles and Practice of Immunoassay*, Stockton Press, N.Y.; and Ngo (ed. 1988) *Nonisotopic Immunoassay*, Plenum Press, N.Y. Cross absorptions or other tests will identify antibodies which exhibit various spectra of specificities, e.g., unique or shared species specificities.

[0057] Further, the antibodies, including antigen binding fragments, of this invention can be potent antagonists that bind to the antigen and inhibit functional binding, e.g., to a receptor which may elicit a biological response. They also can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides so that when the antibody binds to antigen, a cell expressing it, e.g., on its surface, is killed. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker, and may effect drug targeting.

[0058] Antigen fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. An antigen and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) *Specificity of Serological Reactions*, Dover Publications, New York; Williams, et al. (1967) *Methods in Immunology and Immunochemistry*, vol. 1, Academic Press, New York; and Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, CSH Press, NY, for descriptions of methods of preparing polyclonal antisera.

[0059] In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) *Basic and Clinical Immunology* (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, CSH Press; Goding (1986) *Monoclonal Antibodies: Principles and Practice* (2d ed.), Academic Press, New York; and particularly in Kohler and Milstein (1975) in *Nature* 256:495-497, which discusses one method of generating monoclonal antibodies.

[0060] Other suitable techniques involve *in vitro* exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," *Science* 246:1275-1281; and Ward, et al. (1989) *Nature* 341:544-546. The polypeptides and antibodies disclosed herein may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also,

recombinant immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; Moore, et al., U.S. Patent No. 4,642,334; and Queen, et al. (1989) Proc. Nat'l Acad. Sci. USA 86:10029-10033.

**[0061]** The antibodies of this invention can also be used for affinity chromatography in isolating the protein. Columns can be prepared where the antibodies are linked to a solid support. See, e.g., Wilchek et al. (1984) Meth. Enzymol. 104:3-55.

**[0062]** Antibodies raised against each IL-B30 will also be useful to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the respective antigens.

## **VI. Nucleic Acids**

**[0063]** The described peptide sequences and the related reagents are useful in detecting, isolating, or identifying a DNA clone encoding IL-B30, e.g., from a natural source. Typically, it will be useful in isolating a gene from mammal, and similar procedures will be applied to isolate genes from other species, e.g., warm blooded animals, such as birds and mammals. Cross hybridization will allow isolation of IL-B30 from the same, e.g., polymorphic variants, or other species. A number of different approaches will be available to successfully isolate a suitable nucleic acid clone.

**[0064]** The purified protein or defined peptides are useful for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein can be presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual, Cold Spring Harbor Press.

**[0065]** For example, the specific binding composition could be used for screening of an expression library made from a cell line which expresses an IL-B30. Screening of intracellular expression can be performed by various staining or immunofluorescence procedures. Binding compositions could be used to affinity purify or sort out cells expressing a surface fusion protein.

**[0066]** The peptide segments can also be used to predict appropriate oligonucleotides to screen a library. The genetic code can be used to select appropriate oligonucleotides useful as probes for screening. See, e.g., SEQ ID NO: 1. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting correct clones from a library. Complementary sequences will also be used as probes, primers, or antisense strands. Various fragments should be particularly useful, e.g., coupled with anchored vector or poly-A complementary PCR techniques or with complementary of other peptides.

**[0067]** Disclosed herein is use of isolated DNA or fragments to encode a biologically active corresponding IL-B30 polypeptide, particularly lacking the portion coding the untranslated 5' portion of the described sequence. In addition, disclosed herein is isolated or recombinant DNA which encodes a biologically active protein or polypeptide and which is capable of hybridizing under appropriate conditions with the DNA sequences described herein. Said biologically active protein or polypeptide can be an intact antigen, or fragment, and have an amino acid sequence disclosed in, e.g., SEQ ID NO: 2, particularly a mature, secreted polypeptide. Further, disclosed herein is the use of isolated or recombinant DNA, or fragments thereof, which encode proteins which exhibit high identity to a secreted IL-B30. The isolated DNA can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others. Alternatively, expression may be effected by operably linking a coding segment to a heterologous promoter, e.g., by inserting a promoter upstream from an endogenous gene.

**[0068]** An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially separated from other components which naturally accompany a native sequence, e.g., ribosomes, polymerases, and/or flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule. Generally, the nucleic acid will be in a vector or fragment less than about 50 kb, usually less than about 30 kb, typically less than about 10 kb, and preferably less than about 6 kb.

**[0069]** An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some instances, contain minor heterogeneity. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

**[0070]** A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human

intervention in the nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants. Thus, e.g., products made by transforming cells with any unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site.

**[0071]** Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species or polymorphic variants.

**[0072]** A significant "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least about 22 nucleotides, ordinarily at least about 29 nucleotides, more often at least about 35 nucleotides, typically at least about 41 nucleotides, usually at least about 47 nucleotides, preferably at least about 55 nucleotides, and particularly will be at least about 60 or more nucleotides, e.g., 67, 73, 81, 89, 95, etc.

**[0073]** A DNA which codes for an IL-B30 protein will be particularly useful to identify genes, mRNA, and cDNA species which code for related or similar proteins, as well as DNAs which code for homologous proteins from different species. There will be homologs in other species, including primates, rodents, canines, felines, and birds. Various IL-B30 proteins should be homologous and are encompassed herein. However, even proteins that have a more distant evolutionary relationship to the antigen can readily be isolated under appropriate conditions using these sequences if they are sufficiently homologous. Primate IL-B30 proteins are of particular interest.

**[0074]** Recombinant clones derived from the genomic sequences, e.g., containing introns, will be useful for transgenic studies, including, e.g., transgenic cells and organisms, and for gene therapy. See, e.g., Goodnow (1992) "Transgenic Animals" in Roitt (ed.) *Encyclopedia of Immunology*, Academic Press, San Diego, pp. 1502-1504; Travis (1992) *Science* 256:1392-1394; Kuhn, et al. (1991) *Science* 254:707-710; Capecchi (1989) *Science* 244:1288; Robertson (1987)(ed.) *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, IRL Press, Oxford; and Rosenberg (1992) *J. Clinical Oncology* 10:180-199.

**[0075]** Substantial homology, e.g., identity, in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least about 58%, ordinarily at least about 65%, often at least about 71%, typically at least about 77%, usually at least about 85%, preferably at least about 95 to 98% or more, and in particular, as high as about 99% or more of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to a strand, or its complement, typically using a sequence of IL-B30, e.g., in SEQ ID NO: 1. Typically, selective hybridization will occur when there is at least about 55% identity over a stretch of at least about 30 nucleotides, preferably at least about 75% over a stretch of about 25 nucleotides, and most preferably at least about 90% over about 20 nucleotides. See, Kanehisa (1984) *Nuc. Acids Res.* 12:203-213. The length of identity comparison, as described, may be over longer stretches, and in certain instances will be over a stretch of at least about 17 nucleotides, usually at least about 28 nucleotides, typically at least about 40 nucleotides, and preferably at least about 75 to 100 or more nucleotides.

**[0076]** Stringent conditions, in referring to homology in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters, typically those controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30° C, usually in excess of about 37° C, typically in excess of about 55° C, preferably in excess of about 70° C. Stringent salt conditions will ordinarily be less than about 1000 mM, usually less than about 400 mM, typically less than about 250 mM, preferably less than about 150 mM, including about 100, 50, or even 20 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) *J. Mol. Biol.* 31:349-370. Hybridization under stringent conditions should give a background of at least 2-fold over background, preferably at least 3-5 or more.

**[0077]** For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence,

based on the designated program parameters.

**[0078]** Optical alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1981) *Adv. Appl. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Nat'l Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel et al., *supra*).

**[0079]** One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng and Doolittle (1987) *J. Mol. Evol.* 35:351-360. The method used is similar to the method described by Higgins and Sharp (1989) *CABIOS* 5:151-153. The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. For example, a reference sequence can be compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

**[0080]** Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described Altschul, et al. (1990) *J. Mol. Biol.* 215:403-410. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul, et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a wordlength (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Nat'l Acad. Sci. USA* 89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

**[0081]** In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) *Proc. Nat'l Acad. Sci. USA* 90:5873-5877). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

**[0082]** A further indication that two nucleic acid sequences of polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions, as described below.

**[0083]** IL-B30 from other mammalian species can be cloned and isolated by cross-species hybridization of closely related species. Homology may be relatively low between distantly related species, and thus hybridization of relatively closely related species is advisable. Alternatively, preparation of an antibody preparation which exhibits less species specificity may be useful in expression cloning approaches.

## VII. Making IL-B30; Mimetics

**[0084]** DNA which encodes the IL-B30 or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or screening genomic libraries prepared from a wide variety of cell lines or tissue samples. See, e.g., Okayama and Berg (1982) *Mol. Cell. Biol.* 2:161-170; Gubler and Hoffman (1983) *Gene* 25:263-269; and Glover (ed. 1984) *DNA Cloning: A Practical Approach*, IRL Press, Oxford. Alternatively, the sequences provided herein provide useful PCR primers or allow synthetic or other preparation of suitable genes encoding an IL-B30; including naturally occurring variants.

**[0085]** This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length IL-B30 or fragments which can in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified molecules; and for structure/function studies.

**[0086]** Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. See, e.g., Pouwels, et al. (1985 and Supplements) *Cloning Vectors: A Laboratory Manual*, Elsevier, N.Y.; and Rodriguez, et al. (1988)(eds.) *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworth, Boston, MA.

**[0087]** For purposes of this disclosure, DNA sequences are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression. See, e.g., Rodriguez, et al., Chapter 10, pp. 205-236; Balbas and Bolivar (1990) *Methods in Enzymology* 185:14-37; and Ausubel, et al. (1993) *Current Protocols in Molecular Biology*, Greene and Wiley, NY.

**[0088]** Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) *Mol. Cell Biol.* 5:1136-1142; pMCIneo Poly-A, see Thomas, et al. (1987) *Cell* 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610. See, e.g., Miller (1988) *Ann. Rev. Microbiol.* 42:177-199.

**[0089]** It will often be desired to express an IL-B30 polypeptide in a system which provides a specific or defined glycosylation pattern. See, e.g., Luckow and Summers (1988) *Bio/Technology* 6:47-55; and Kaufman (1990) *Meth. Enzymol.* 185:487-511.

**[0090]** The IL-B30, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by standard procedures of protein chemistry. See, e.g., Low (1989) *Biochim. Biophys. Acta* 988:427-454; Tse, et al. (1985) *Science* 230:1003-1008; and Brunner, et al. (1991) *J. Cell Biol.* 114:1275-1283.

**[0091]** Now that the IL-B30 has been characterized, fragments or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) *Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) *The Practice of Peptide Synthesis*, Springer-Verlag, New York; Bodanszky (1984) *The Principles of Peptide Synthesis*, Springer-Verlag, New York; and Villafranca (ed. 1991) *Techniques in Protein Chemistry II*, Academic Press, San Diego, Ca.

## **VIII.Uses**

**[0092]** Disclosed herein are reagents which will find use in diagnostic applications as described elsewhere herein, e.g., in IL-B30 mediated conditions, or below in the description of kits for diagnosis. The gene may be useful in forensic sciences, e.g., to distinguish rodent from human, or as a marker to distinguish between different cells exhibiting differential expression or modification patterns.

**[0093]** Disclosed herein are reagents with significant commercial and/or therapeutic potential. The IL-B30 (naturally occurring or recombinant), fragments thereof, and antibodies thereto, along with compounds identified as having binding affinity to IL-B30, should be useful as reagents for teaching techniques of molecular biology, immunology, or physiology. Appropriate kits may be prepared with the reagents, e.g., in practical laboratory exercises in production or use of proteins, antibodies, cloning methods, histology, etc.

**[0094]** The reagents will also be useful in the treatment of conditions associated with abnormal physiology or development, including inflammatory conditions. They may be useful in vitro tests for presence or absence of interacting components, which may correlate with success of particular treatment strategies. In particular, modulation of physiology of various, e.g., hematopoietic or lymphoid, cells will be achieved by appropriate methods for treatment using the compositions provided herein. See, e.g., Thomson (1994; ed.) *The Cytokine Handbook* (2d ed.) Academic Press, San Diego; Metcalf and Nicola (1995) *The Hematopoietic Colony Stimulating Factors* Cambridge University Press; and Aggarwal and Gutterman (1991) *Human Cytokines* Blackwell Pub.

**[0095]** For example, a disease or disorder associated with abnormal expression or abnormal signaling by an IL-B30 should be a likely target for an agonist or antagonist. The new cytokine should play a role in regulation or development of hematopoietic cells, e.g., lymphoid cells, which affect immunological responses, e.g., inflammation and/or autoimmune disorders. Alternatively, it may affect vascular physiology or development, or neuronal effects.

**[0096]** In particular, the cytokine should mediate, in various contexts, cytokine synthesis by the cells, proliferation, etc. Antagonists of IL-B30, such as mutein variants of a naturally occurring form of IL-B30 or blocking antibodies, may provide a selective and powerful way to block immune responses, e.g., in situations as inflammatory or autoimmune responses. See also Samter, et al. (eds.) *Immunological Diseases* vols. 1 and 2, Little, Brown and Co.

**[0097]** In addition, certain combination compositions would be useful, e.g., with other modulators of inflammation. Such other molecules may include steroids, other versions of IL-6 and/or G-CSF, including species variants, or viral homologs, and their respective antagonists.

**[0098]** Various abnormal conditions are known in each of the cell types shown to produce IL-B30 mRNA by Northern blot analysis. See Berkow (ed.) *The Merck Manual of Diagnosis and Therapy*, Merck & Co., Rahway, N.J.; Thorn, et al. *Harrison's Principles of Internal Medicine*, McGraw-Hill, N.Y.; and Weatherall, et al. (eds.) *Oxford Textbook of Medicine*, Oxford University Press, Oxford. Many other medical conditions and diseases involve activation by macrophages or monocytes, and many of these will be responsive to treatment by an agonist or antagonist provided herein. See, e.g., Stites and Terr (eds.; 1991) *Basic and Clinical Immunology* Appleton and Lange, Norwalk, Connecticut; and Samter, et al. (eds.) *Immunological Diseases* Little, Brown and Co. These problems should be susceptible to prevention or treatment using compositions provided herein. The pancreatic islet localization suggests a possible relevance to diabetes.

**[0099]** IL-B30, antagonists, antibodies, etc., can be purified and then administered to a patient, veterinary or human. These reagents can be combined for therapeutic use with additional active or inert ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, e.g., immunogenic adjuvants, along with physiologically innocuous stabilizers, excipients, or preservatives. These combinations can be sterile filtered and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof, including forms which are not complement binding.

**[0100]** Drug screening using IL-B30 or fragments thereof can be performed to identify compounds having binding affinity to or other relevant biological effects on IL-B30 functions, including isolation of associated components. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is therefore a blocker or antagonist in that it blocks the activity of the cytokine. Likewise, a compound having intrinsic stimulating activity can activate the signal pathway and is thus an agonist in that it simulates the activity of IL-B30. This invention further contemplates the therapeutic use of blocking antibodies to IL-B30 as antagonists and of stimulatory antibodies as agonists. This approach should be particularly useful with other IL-B30 species variants.

**[0101]** The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicaments administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds. 1990) *Goodman and Gilman's: The Pharmacological Bases of Therapeutics*, 8th Ed., Pergamon Press; and Remington's *Pharmaceutical Sciences*, 17th ed. (1990), Mack Publishing Co., Easton, Penn. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10  $\mu$ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or a slow release apparatus will often be

utilized for continuous or long term administration. See, e.g., Langer (1990) *Science* 249:1527-1533.

**[0102]** IL-B30, fragments thereof, and antibodies to it or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations typically comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier should be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, topical, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) *Goodman and Gilman's: The Pharmacological Bases of Therapeutics*, 8th Ed., Pergamon Press; and Remington's *Pharmaceutical Sciences*, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) *Pharmaceutical Dosage Forms: Parenteral Medications*, Dekker, New York; Lieberman, et al. (eds. 1990) *Pharmaceutical Dosage Forms: Tablets*, Dekker, New York; and Lieberman, et al. (eds. 1990) *Pharmaceutical Dosage Forms: Disperse Systems*, Dekker, New York. The therapy of this invention may be combined with or used in association with other agents, e.g., other cytokines, including IL-6 or G-CSF, or their respective antagonists.

**[0103]** Both naturally occurring and recombinant forms of the IL-B30s disclosed herein are particularly useful in kits and assay methods which are capable of screening compounds for binding activity to the proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) *Science* 251:767-773, which describes means for testing of binding affinity by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays can be greatly facilitated by the availability of large amounts of purified, soluble IL-B30 as disclosed herein.

**[0104]** Other methods can be used to determine the critical residues in IL-B30-IL-B30 receptor interactions. Mutational analysis can be performed, e.g., see Somoza, et al. (1993) *J. Exp. Med.* 178:549-558, to determine specific residues critical in the interaction and/or signaling. PHD (Rost and Sander (1994) *Proteins* 19:55-72) and DSC (King and Sternberg (1996) *Protein Sci.* 5:2298-2310) can provide secondary structure predictions of  $\alpha$ -helix (H),  $\beta$ -strand (E), or coil (L). Helices A and D are most important in receptor interaction, with the D helix being the more important region. Helix A would run in the human from about pro(7) to his(27), while helix D would run from about trp(135) to gly(162). Surface exposed residues would affect receptor binding, while embedded residues would affect general structure. Predicted residues of particular importance would likely correspond to arg(146), ser(147), gln(149), ala(150), ala(153), val(154), ala(156), arg(157), ala(160), and his(161).

**[0105]** For example, antagonists can normally be found once the antigen has been structurally defined, e.g., by tertiary structure data. Testing of potential interacting analogs is now possible upon the development of highly automated assay methods using a purified IL-B30. In particular, new agonists and antagonists will be discovered by using screening techniques described herein. Of particular importance are compounds found to have a combined binding affinity for a spectrum of IL-B30 molecules, e.g., compounds which can serve as antagonists for species variants of IL-B30.

**[0106]** One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing an IL-B30. Cells may be isolated which express an IL-B30 in isolation from other molecules. Such cells, either in viable or fixed form, can be used for standard binding partner binding assays. See also, Parce, et al. (1989) *Science* 246:243-247; and Owicki, et al. (1990) *Proc. Nat'l Acad. Sci. USA* 87:4007-4011, which describe sensitive methods to detect cellular responses.

**[0107]** Another technique for drug screening involves an approach which provides high throughput screening for compounds having suitable binding affinity to an IL-B30 and is described in detail in Geysen, European Patent Application 84/03564, published on September 13, 1984. First, large numbers of different small peptide test compounds are synthesized on a solid substrate, e.g., plastic pins or some other appropriate surface, see Fodor, et al. (1991). Then all the pins are reacted with solubilized, unpurified or solubilized, purified IL-B30, and washed. The next step involves detecting bound IL-B30.

**[0108]** Rational drug design may also be based upon structural studies of the molecular shapes of the IL-B30 and other effectors or analogs. Effectors may be other proteins which mediate other functions in response to binding, or other proteins which normally interact with IL-B30, e.g., a receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions, as modeled, e.g., against other cytokine-receptor models. For a

detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York.

## **IX. Kits**

**[0109]** Disclosed herein is use of IL-B30 proteins, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of another IL-B30 or binding partner. Typically the kit will have a compartment containing either a defined IL-B30 peptide or gene segment or a reagent which recognizes one or the other, e.g., IL-B30 fragments or antibodies.

**[0110]** A kit for determining the binding affinity of a test compound to an IL-B30 would typically comprise a test compound; a labeled compound, for example a binding partner or antibody having known binding affinity for IL-B30; a source of IL-B30 (naturally occurring or recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the molecule. Once compounds are screened, those having suitable binding affinity to the antigen can be evaluated in suitable biological assays, as are well known in the art, to determine whether they act as agonists or antagonists to the IL-B30 signaling pathway. The availability of recombinant IL-B30 polypeptides also provide well defined standards for calibrating such assays.

**[0111]** A preferred kit for determining the concentration of, e.g., an IL-B30 in a sample would typically comprise a labeled compound, e.g., binding partner or antibody, having known binding affinity for the antigen, a source of cytokine (naturally occurring or recombinant) and a means for separating the bound from free labeled compound, e.g., a solid phase for immobilizing the IL-B30. Compartments containing reagents, and instructions, will normally be provided.

**[0112]** Antibodies, including antigen binding fragments, specific for the IL-B30 or fragments are useful in diagnostic applications to detect the presence of elevated levels of IL-B30 and/or its fragments. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, body fluids, and further can involve the detection of antigens related to the antigen in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent and antigen-binding partner complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. See, e.g., Van Vunakis, et al. (1980) *Meth Enzymol.*, 70:1-525; Harlow and Lane (1980) *Antibodies: A Laboratory Manual*, CSH Press, NY; and Coligan, et al. (eds. 1993) *Current Protocols in Immunology*, Greene and Wiley, NY.

**[0113]** Anti-idiotypic antibodies may have similar use to diagnose presence of antibodies against an IL-B30, as such may be diagnostic of various abnormal states. For example, overproduction of IL-B30 may result in production of various immunological reactions which may be diagnostic of abnormal physiological states, particularly in proliferative cell conditions such as cancer or abnormal activation or differentiation. Moreover, the distribution pattern available provides information that the cytokine is expressed in pancreatic islets, suggesting the possibility that the cytokine may be involved in function of that organ, e.g., in a diabetes relevant medical condition.

**[0114]** Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. Depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody or binding partner, or labeled IL-B30 may be provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

**[0115]** Many of the aforementioned constituents of the drug screening and the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the binding partner, test compound, IL-B30, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as <sup>125</sup>I, enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

[0116] There are also numerous methods of separating the bound from the free IL-B30, or alternatively the bound from the free test compound. The IL-B30 can be immobilized on various matrixes followed by washing. Suitable matrixes include plastic such as an ELISA plate, filters, and beads. See, e.g., Coligan, et al. (eds. 1993) *Current Protocols in Immunology*, Vol. 1, Chapter 2, Greene and Wiley, NY. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) *Clin. Chem.* 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678.

[0117] Methods for linking proteins or their fragments to the various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

[0118] Disclosed herein is use of oligonucleotide or polynucleotide sequences taken from the sequence of an IL-B30. These sequences can be used as probes for detecting levels of the IL-B30 message in samples from patients suspected of having an abnormal condition, e.g., inflammatory or autoimmune. Since the cytokine may be a marker or mediator for activation, it may be useful to determine the numbers of activated cells to determine, e.g., when additional therapy may be called for, e.g., in a preventative fashion before the effects become and progress to significance. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. See, e.g., Langer-Safer, et al. (1982) *Proc. Nat'l Acad. Sci.* 79:4381-4385; Caskey (1987) *Science* 236:962-967; and Wilchek et al. (1988) *Anal. Biochem.* 171:1-32.

[0119] Diagnostic kits which also test for the qualitative or quantitative expression of other molecules are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) *Progress in Growth Factor Res.* 1:89-97. Other kits may be used to evaluate other cell subsets.

## **X. Isolating a IL-B30 Receptor**

[0120] Having isolated a ligand of a specific ligand-receptor interaction, methods exist for isolating the receptor. See, Gearing, et al. (1989) *EMBO J.* 8:3667-3676. For example, means to label the IL-B30 cytokine without interfering with the binding to its receptor can be determined. For example, an affinity label can be fused to either the amino- or carboxyl-terminus of the ligand. Such label may be a FLAG epitope tag, or, e.g., an Ig or Fc domain. An expression library can be screened for specific binding of the cytokine, e.g., by cell sorting, or other screening to detect subpopulations which express such a binding component. See, e.g., Ho, et al. (1993) *Proc. Nat'l Acad. Sci. USA* 90:11267-11271; and Liu, et al. (1994) *J. Immunol.* 152:1821-29. Alternatively, a panning method may be used. See, e.g., Seed and Aruffo (1987) *Proc. Nat'l Acad. Sci. USA* 84:3365-3369.

[0121] Protein cross-linking techniques with label can be applied to isolate binding partners of the IL-B30 cytokine. This would allow identification of proteins which specifically interact with the cytokine, e.g., in a ligand-receptor like manner.

[0122] Early experiments will be performed to determine whether the known IL-6 or G-CSF receptor components are involved in response(s) to IL-B30. It is also quite possible that these functional receptor complexes may share many or all components with an IL-B30 receptor complex, either a specific receptor subunit or an accessory receptor subunit.

## **EXAMPLES**

### **I. General Methods**

[0123] Many of the standard methods below are described or referenced, e.g., in Maniatis, et al. (1982) *Molecular Cloning*, A Laboratory Manual Cold Spring Harbor Laboratory, Cold Spring Harbor Press, NY; Sambrook, et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed.) Vols. 1-3, CSH Press, NY; Ausubel, et al., *Biology* Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) *Current Protocols in Molecular Biology* Wiley/Greene, NY; Innis, et al. (eds. 1990) *PCR Protocols: A Guide to Methods and Application* Academic Press, NY. Methods for protein purification include such methods as

ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification," Methods in Enzymology vol. 182, and other volumes in this series; Coligan, et al. (1995 and supplements) Current Protocols in Protein Science John Wiley and Sons, New York, NY; P. Matsudaira (ed. 1993) A Practical Guide to Protein and Peptide Purification for Microsequencing, Academic Press, San Diego, CA; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, NJ, or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments (epitope tags), e.g., to a FLAG sequence or an equivalent which can be fused, e.g., via a protease-removable sequence. See, e.g., Hochuli (1989) *Chemische Industrie* 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) *Genetic Engineering, Principle and Methods* 12:87-98, Plenum Press, NY; and Crowe, et al. (1992) *OIAexpress: The High Level Expression & Protein Purification System* QUIAGEN, Inc., Chatsworth, CA.

**[0124]** Standard immunological techniques are described, e.g., in Hertzzenberg, et al. (eds. 1996) *Weir's Handbook of Experimental Immunology* vols. 1-4, Blackwell Science; Coligan (1991) *Current Protocols in Immunology* Wiley/Greene, NY; and *Methods in Enzymology* vols. 70, 73, 74, 84, 92, 93, 108, 116, 121, 132, 150, 162, and 163. Cytokine assays are described, e.g., in Thomson (ed. 1994) *The Cytokine Handbook* (2d ed.) Academic Press, San Diego; Metcalf and Nicola (1995) *The Hematopoietic Colony Stimulating Factors* Cambridge University Press; and Aggarwal and Gutterman (1991) *Human Cytokines* Blackwell Pub.

**[0125]** Assays for vascular biological activities are well known in the art. They will cover angiogenic and angiostatic activities in tumor, or other tissues, e.g., arterial smooth muscle proliferation (see, e.g., Koyoma, et al. (1996) *Cell* 87:1069-1078), monocyte adhesion to vascular epithelium (see McEvoy, et al. (1997) *J. Exp. Med.* 185:2069-2077), etc. See also Ross (1993) *Nature* 362:801-809; Rekhter and Gordon (1995) *Am. J. Pathol.* 147:668-677; Thyberg, et al. (1990) *Atherosclerosis* 10:966-990; and Gumbiner (1996) *Cell* 84:345-357.

**[0126]** Assays for neural cell biological activities are described, e.g., in Wouterlood (ed. 1995) *Neuroscience Protocols* modules 10, Elsevier; *Methods in Neurosciences* Academic Press; and *Neuromethods* Humana Press, Totowa, NJ. Methodology of developmental systems is described, e.g., in Meisami (ed.) *Handbook of Human Growth and Developmental Biology* CRC Press; and Chrispeels (ed.) *Molecular Techniques and Approaches in Developmental Biology* Interscience.

**[0127]** FACS analyses are described in Melamed, et al. (1990) *Flow Cytometry and Sorting* Wiley-Liss, Inc., New York, NY; Shapiro (1988) *Practical Flow Cytometry* Liss, New York, NY; and Robinson, et al. (1993) *Handbook of Flow Cytometry Methods* Wiley-Liss, New York, NY.

## **II. Cloning of Human IL-B30**

**[0128]** The sequence of the gene is provided in Table 1. The sequence is derived from a cDNA library made from melanocyte, fetal heart, and pregnant uterus. It is also found from a cDNA library sequence derived from a pancreatic islet. These sequences allow preparation of PCR primers, or probes, to determine cellular distribution of the gene. The sequences allow isolation of genomic DNA which encode the message.

**[0129]** Using the probe or PCR primers, various tissues or cell types are probed to determine cellular distribution. PCR products are cloned using, e.g., a TA cloning kit (Invitrogen). The resulting cDNA plasmids are sequenced from both termini on an automated sequencer (Applied Biosystems).

## **III. Cellular Expression of IL-B30**

**[0130]** An appropriate probe or primers specific for cDNA encoding primate IL-B30 are prepared. Typically, the probe is labeled, e.g., by random priming. The expression is probably in the cell types described, and perhaps also in pancreatic islets. Southern Analysis: DNA (5 µg) from a primary amplified cDNA library was digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and transferred to a nylon membrane (Schleicher and Schuell, Keene, NH).

**[0131]** Samples for human mRNA isolation include: peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); T cell, TH1 clone HY06,

resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13, Tc783.58, Tc782.69, resting (T118); T cell random  $\gamma\delta$  T cell clones, resting (T119); CD28- T cell clone; Splenocytes, resting (B100); Splenocytes, activated with anti-CD40 and IL-4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); NK 20 clones pooled, resting (K100); NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101); NK1 clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101); elutriated monocytes, activated with LPS, IFN $\gamma$ , anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFN $\gamma$ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFN $\gamma$ , anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFN $\gamma$ , IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated monocytes, activated LPS for 6 h (M109); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D104); DC 95% CD14+, ex CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNF $\alpha$ , monocyte supe for 4, 16 h pooled (D110); epithelial cells, unstimulated; epithelial cells, IL-1 $\beta$  activated; lung fibroblast sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102). Expression of IL-B30 transcript was very high in elutriated monocytes, activated with LPS, IFN $\gamma$ , anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFN $\gamma$ , anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated LPS for 6 h (M109); and elutriated monocytes, activated LPS for 1 h (M108). Expression was high in DC 95% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D104); and NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101). Lesser expression was detected in DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, activated with PMA and ionomycin for 6 hr (D103); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, activated with PMA and ionomycin for 1 hr (D102); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T103); Splenocytes, activated with anti-CD40 and IL-4 (B101); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); Splenocytes, resting (B100); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); epithelial cells, IL-1 $\beta$  activated; elutriated monocytes, activated with LPS, IFN $\gamma$ , IL-10 for 4, 16 h pooled (M107); and B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103). Detectable expression was observed in DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); T cell, TH0 clone Mot 72, resting (T102); peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti IFN- $\gamma$ , TH2 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell clones, pooled AD130.2, Tc783.12, Tc783.13, Tc783.58, Tc782.69, resting (T118); U937 premonocytic line, resting (M100); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D106); elutriated monocytes, activated with LPS, IFN $\gamma$ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); DC from monocytes GM-CSF, IL-4 5 days, activated TNF $\alpha$ , monocyte supe for 4, 16 h pooled (D110); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101); T cell random  $\gamma\delta$  T cell clones, resting (T119); and T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108). No signal was detected in the other samples.

**[0132]** In summary, the distribution shows IL-B30 elevated in activated macrophages, suggesting a role in inflammation; activated Th1 cells, suggesting a regulation or effector role in T helper subsets, particularly Th1 immune responses; and activated dendritic cells, suggesting a role in antigen presentation or germinal center T or B cell interactions with DC.

**[0133]** Samples for mouse mRNA isolation include: resting mouse fibroblastic L cell line (C200); Braf:ER (Braf fusion to estrogen receptor) transfected cells, control (C201); Me114 naive T cells from spleen, resting (T209); Me114+ naive T cells from spleen, stimulated with IFN $\gamma$ , IL-12, and anti IL-4 to polarize to TH1 cells, exposed to IFN $\gamma$  and IL-4 for 6, 12, 24 h, pooled (T210); Me114+ naive T cells from spleen, stimulated with IL-4 and anti IFN $\gamma$  to polarize to Th2 cells, exposed to IL-4 and anti IFN $\gamma$  for 6, 13, 24 h, pooled (T211); T cells, TH1 polarized (Me114 bright, CD4+ cells from spleen, polarized for 7 days with IFN- $\gamma$  and anti IL-4; T200); T cells, TH2 polarized (Me114 bright, CD4+ cells from spleen, polarized for 7 days with IL-4 and anti-IFN- $\gamma$ ; T201); T cells, highly TH1 polarized 3x from transgenic Balb/C (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367; activated with anti-CD3 for 2,

6, 24 h pooled; T202); T cells, highly TH2 polarized 3x from transgenic Balb/C (activated with anti-CD3 for 2, 6, 24 h pooled (T203); T cells, highly TH1 polarized 3x from transgenic C57 bl/6 (activated with anti-CD3 for 2, 6, 24 h pooled; T212); T cells, highly TH2 polarized 3x from transgenic C57 bl/6 (activated with anti-CD3 for 2, 6, 24 h pooled; T213); T cells, highly TH1 polarized (naive CD4+ T cells from transgenic Balb/C, polarized 3x with IFN $\gamma$ , IL-12, and anti-IL-4; stimulated with IGF, IL-12, and anti IL-4 for 6, 12, 24 h, pooled); CD44- CD25+ pre T cells, sorted from thymus (T204); TH1 T cell clone D1.1, resting for 3 weeks after last stimulation with antigen (T205); TH1 T cell clone D1.1, 10  $\mu$ g/ml ConA stimulated 15 h (T206); TH2 T cell clone CDC35, resting for 3 weeks after last stimulation with antigen (T207); TH2 T cell clone CDC35, 10  $\mu$ g/ml ConA stimulated 15 h (T208); unstimulated B cell line CH12 (B201); unstimulated mature B cell leukemia cell line A20 (B200); unstimulated large B cells from spleen (B202); B cells from total spleen, LPS activated (B203); metrizamide enriched dendritic cells from spleen, resting (D200); dendritic cells from bone marrow, resting (D201); unstimulated bone marrow derived dendritic cells depleted with anti B220, anti CD3, and anti Class II, cultured in GM-CSF and IL-4 (D202); bone marrow derived dendritic cells depleted with anti B220, anti CD3, and anti Class II, cultured in GM-CSF and IL-4, stimulated with anti CD40 for 1, 5 d, pooled (D203); monocyte cell line RAW 264.7 activated with LPS 4 h (M200); bone-marrow macrophages derived with GM and M-CSF (M201); bone-marrow macrophages derived with GM-CSF, stimulated with LPS, IFN $\gamma$ , and IL-10 for 24 h (M205); bone-marrow macrophages derived with GM-CSF, stimulated with LPS, IFN $\gamma$ , and anti IL-10 for 24 h (M206); peritoneal macrophages (M207); macrophage cell line J774, resting (M202); macrophage cell line J774 + LPS + anti-IL-10 at 0.5, 1, 3, 6, 12 h pooled (M203); macrophage cell line J774 + LPS + IL-10 at 0.5, 1, 3, 5, 12 h pooled (M204); unstimulated mast cell lines MC-9 and MCP-12 (M208); immortalized endothelial cell line derived from brain microvascular endothelial cells, unstimulated (E200); immortalized endothelial cell line derived from brain microvascular endothelial cells, stimulated overnight with TNF $\alpha$  (E201); immortalized endothelial cell line derived from brain microvascular endothelial cells, stimulated overnight with TNF $\alpha$  and IL-10 (E203); total aorta from wt C57 bl/6 mouse; total aorta from 5 month ApoE KO mouse (X207); total aorta from 12 month ApoE KO mouse (X207); wt thymus (0214); total thymus, rag-1 (0208); total kidney, rag-1 (0209); total kidney, NZ B/W mouse; and total heart, rag-1 (0202). High signal was detected in the monocyte cell line RAW 264.7 activated with LPS 4 h (M200); T cells, highly TH1 polarized 3x from transgenic C57 bl/6 (activated with anti-CD3 for 2, 6, 24 h pooled; T212); and T cells, highly TH1 polarized (naive CD4+ T cells from transgenic Balb/C, polarized 3x with IFN $\gamma$ , IL-12, and anti-IL-4; stimulated with IGF, IL-12, and anti IL-4 for 6, 12, 24 h, pooled). Detectable signals were detected in T cells, highly TH1 polarized 3x from transgenic Balb/C (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367; activated with anti-CD3 for 2, 6, 24 h pooled; T202); T cells, TH2 polarized (Mel14 bright, CD4+ cells from spleen, polarized for 7 days with IL-4 and anti-IFN- $\gamma$ ; T201); T cells, TH1 polarized (Mel14 bright, CD4+ cells from spleen, polarized for 7 days with IFN- $\gamma$  and anti IL-4; T200); macrophage cell line J774 + LPS + anti-IL-10 at 0.5, 1, 3, 6, 12 h pooled (M203); macrophage cell line J774, resting (M202); macrophage cell line J774 + LPS + IL-10 at 0.5, 1, 3, 5, 12 h pooled (M204); immortalized endothelial cell line derived from brain microvascular endothelial cells, stimulated overnight with TNF $\alpha$  (E201); and bone-marrow macrophages derived with GM-CSF, stimulated with LPS, IFN $\gamma$ , and anti IL-10 for 24 h (M206). Other samples showed no signal. The expression in the RAW 264.7 mouse monocyte cell line suggests a natural source for protein.

#### **IV. Chromosome mapping of IL-B30**

**[0134]** An isolated cDNA encoding the IL-B30 is used. Chromosome mapping is a standard technique. See, e.g., BIOS Laboratories (New Haven, CT) and methods for using a mouse somatic cell hybrid panel with PCR. Circumstantial evidence suggests that the mouse IL-B30 gene maps to chromosome 10.

#### **V. Purification of IL-B30 Protein**

**[0135]** Multiple transfected cell lines are screened for one which expresses the cytokine at a high level compared with other cells. Various cell lines are screened and selected for their favorable properties in handling. Natural IL-B30 can be isolated from natural sources, or by expression from a transformed cell using an appropriate expression vector. Purification of the expressed protein is achieved by standard procedures, or may be combined with engineered means for effective purification at high efficiency from cell lysates or supernatants. FLAG or His<sub>6</sub> segments can be used for such purification features. Alternatively, affinity chromatography may be used with specific antibodies, see below.

**[0136]** Protein is produced in coli, insect cell, or mammalian expression systems, as desired.

#### **VI. Isolation of Homologous IL-B30 Genes**

**[0137]** The IL-B30 cDNA, or other species counterpart sequence, can be used as a hybridization probe to screen a library from a desired source, e.g., a primate cell cDNA library. Many different species can be screened both for stringency necessary for easy hybridization, and for presence using a probe. Appropriate hybridization conditions will be used to select for clones exhibiting specificity of cross hybridization.

**[0138]** Screening by hybridization using degenerate probes based upon the peptide sequences will also allow isolation of appropriate clones. Alternatively, use of appropriate primers for PCR screening will yield enrichment of appropriate nucleic acid clones.

**[0139]** Similar methods are applicable to isolate either species, polymorphic, or allelic variants. Species variants are isolated using cross-species hybridization techniques based upon isolation of a full length isolate or fragment from one species as a probe.

**[0140]** Alternatively, antibodies raised against human IL-B30 will be used to screen for cells which express cross-reactive proteins from an appropriate, e.g., cDNA library. The purified protein or defined peptides are useful for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) *Current Protocols in Immunology* Wiley/Greene; and Harlow and Lane (1989) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press. The resulting antibodies are used for screening, purification, or diagnosis, as described.

## **VII. Preparation of antibodies specific for IL-B30**

**[0141]** Synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) *Current Protocols in Immunology* Wiley/Greene; and Harlow and Lane (1989) *Antibodies: A Laboratory Manual* Cold Spring Harbor Press. Polyclonal serum, or hybridomas may be prepared. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods. Immunoselection and related techniques are available to prepare selective reagents, as desired.

## **VIII. Evaluation of Breadth of Biological Functions**

**[0142]** Biological activities of IL-B30 were tested based on the sequence and structural homology between IL-B30 and IL-6 and G-CSF. Initially, assays that had shown biological activities of IL-6 or G-CSF are examined.

### **A. Effects on proliferation of cells**

**[0143]** The effect on proliferation of various cell types are evaluated with various concentrations of cytokine. A dose response analysis is performed, in combinations with the related cytokines IL-6, G-CSF, etc.

### **B. Effects on the expression of cell surface molecules on human monocytes**

**[0144]** Monocytes are purified by negative selection from peripheral blood mononuclear cells of normal healthy donors. Briefly,  $3 \times 10^8$  ficoll banded mononuclear cells are incubated on ice with a cocktail of monoclonal antibodies (Becton-Dickinson; Mountain View, CA) consisting, e.g., of 200  $\mu$ l of  $\alpha$ CD2 (Leu-5A), 200  $\mu$ l of  $\alpha$ CD3 (Leu-4), 100  $\mu$ l of  $\alpha$ CD8 (Leu 2a), 100  $\mu$ l of  $\alpha$ CD19 (Leu-12), 100  $\mu$ l of  $\alpha$ CD20 (Leu-16), 100  $\mu$ l of  $\alpha$ CD56 (Leu-19), 100  $\mu$ l of  $\alpha$ CD67 (IOM 67; Immunotech, Westbrook, ME), and anti-glycophorin antibody (10F7MN, ATCC, Rockville, MD). Antibody bound cells are washed and then incubated with sheep anti-mouse IgG coupled magnetic beads (Dynal, Oslo, Norway) at a bead to cell ratio of 20:1. Antibody bound cells are separated from monocytes by application of a magnetic field. Subsequently, human monocytes are cultured in Yssel's medium (Gemini Bioproducts, Calabasas, CA) containing 1% human AB serum in the absence or presence of IL-B30, IL-6, G-CSF or combinations.

**[0145]** Analyses of the expression of cell surface molecules can be performed by direct immunofluorescence. For example,  $2 \times$

10<sup>5</sup> purified human monocytes are incubated in phosphate buffered saline (PBS) containing 1% human serum on ice for 20 minutes. Cells are pelleted at 200 x g. Cells are resuspended in 20 ml PE or FITC labeled mAb. Following an additional 20 minute incubation on ice, cells are washed in PBS containing 1% human serum followed by two washes in PBS alone. Cells are fixed in PBS containing 1% paraformaldehyde and analyzed on FACScan flow cytometer (Becton Dickinson; Mountain View, CA). Exemplary mAbs are used, e.g.: CD11b (anti-mac1), CD11c (a gp150/95), CD14 (Leu-M3), CD54 (Leu 54), CD80 (anti-BB1/B7), HLA-DR (L243) from Becton-Dickinson and CD86 (FUN 1; Pharmingen), CD64 (32.2; Medarex), CD40 (mAb89; Schering-Plough France).

**C. Effects of IL-B30 on cytokine production by human monocytes**

**[0146]** Human monocytes are isolated as described and cultured in Yssel's medium (Gemini Bioproducts, Calabasas, CA) containing 1% human AB serum in the absence or presence of IL-B30 (1/100 dilution baculovirus expressed material). In addition, monocytes are stimulated with LPS (E. coli 0127:B8 Difco) in the absence or presence of IL-B30 and the concentration of cytokines (IL-1 $\beta$ , IL-6, TNF $\alpha$ , GM-CSF, and IL-10) in the cell culture supernatant determined by ELISA.

**[0147]** For intracytoplasmic staining for cytokines, monocytes are cultured (1 million/ml) in Yssel's medium in the absence or presence of IL-B30 and LPS (E. coli 0127:B8 Difco) and 10 mg/ml Brefeldin A (Epicentre technologies Madison WI) for 12 hrs. Cells are washed in PBS and incubated in 2% formaldehyde/PBS solution for 20 minutes at RT. Subsequently cells are washed, resuspended in permeabilization buffer (0.5% saponin (Sigma) in PBS/BSA (0.5%) /Azide (1 mM)) and incubated for 20 minutes at RT. Cells (2 x 10<sup>5</sup>) are centrifuged and resuspended in 20 ml directly conjugated anti-cytokine mAbs diluted 1:10 in permeabilization buffer for 20 minutes at RT. The following antibodies can be used: IL-1 $\alpha$ -PE (364-3B3-14); IL-6-PE (MQ2-13A5); TNF $\alpha$ -PE (MAb11); GM-CSF-PE (BVD2-21C11); and IL-12-PE (C11.5.14; Pharmingen San Diego, CA). Subsequently, cells are washed twice in permeabilization buffer and once in PBS/BSA/Azide and analyzed on FACScan flow cytometer (Becton Dickinson; Mountain View, CA).

**D. Effects of IL-B30 on proliferation of human peripheral blood mononuclear cells (PBMC).**

**[0148]** Total PBMC are isolated from buffy coats of normal healthy donors by centrifugation through ficoll-hypaque as described (Boyum, et al.). PBMC are cultured in 200  $\mu$ l Yssel's medium (Gemini Bioproducts, Calabasas, CA) containing 1% human AB serum in 96 well plates (Falcon, Becton-Dickinson, NJ) in the absence or presence of IL-B30. Cells are cultured in medium alone or in combination with 100 U/ml IL-2 (R&D Systems) for 120 hours. 3H-Thymidine (0.1 mCi) is added during the last six hours of culture and 3H-Thymidine incorporation determined by liquid scintillation counting.

**[0149]** The native, recombinant, and fusion proteins would be tested for agonist and antagonist activity in many other biological assay systems, e.g., on T-cells, B-cells, NK, macrophages, dendritic cells, hematopoietic progenitors, etc. Because of the IL-6 and G-CSF structural relationship, assays related to those activities should be analyzed

**[0150]** IL-B30 is evaluated for agonist or antagonist activity on transfected cells expressing IL-6 or G-CSF receptor and controls. See, e.g., Ho, et al. (1993) Proc. Nat'l Acad. Sci. USA 90, 11267-11271 ; Ho, et al. (1995) Mol. Cell. Biol. 15:5043-5053; and Liu, et al. (1994). J. Immunol. 152:1821-1829.

**[0151]** IL-B30 is evaluated for effect in macrophage/dendritic cell activation and antigen presentation assays, T cell cytokine production and proliferation in response to antigen or allogeneic stimulus. See, e.g., de Waal Malefyt et al. (1991) J. Exp. Med. 174:1209-1220; de Waal Malefyt et al. (1991) J. Exp. Med. 174:915-924; Fiorentino, et al. (1991) J. Immunol. 147, 3815-3822; Fiorentino, et al. (1991) J. Immunol. 146:3444-3451; and Groux, et al. (1996) J. Exp. Med. 184:19-29.

**[0152]** IL-B30 will also be evaluated for effects on NK cell stimulation. Assays may be based, e.g., on Hsu, et al. (1992) Internat. Immunol. 4:563-569; and Schwarz, et al. (1994) J. Immunother. 16:95-104.

**[0153]** B cell growth and differentiation effects will be analyzed, e.g., by the methodology described, e.g., in Defrance, et al. (1992). J. Exp. Med. 175:671-682; Rousset, et al. (1992) Proc. Nat'l Acad. Sci. USA 89:1890-1893; including IgG2 and IgA2 switch factor assays. Note that, unlike COS7 supernatants, NIH3T3 and COP supernatants apparently do not interfere with human B cell assays.

**IX. Generation and Analysis of Genetically Altered Animals**

**[0154]** Transgenic mice can be generated by standard methods. Such animals are useful to determine the effects of deletion of the gene, in specific tissues, or completely throughout the organism. Such may provide interesting insight into development of the animal or particular tissues in various stages. Moreover, the effect on various responses to biological stress can be evaluated. See, e.g., Hogan, et al. (1995) *Manipulating the Mouse Embryo: A Laboratory Manual* (2d ed.) Cold Spring Harbor Laboratory Press.

**[0155]** A transgenic mouse has been generated, and while the animal seems to survive birth, it fails to thrive, and typically dies within a few weeks. The construct is based upon an actin promoter with a CMV enhancer, which should lead to broad and high expression. The mice, like IL-6 transgenic mice, are runted. Moreover, they exhibit a bloated abdomen, inflammation of the stomach and intestines, infiltration of cells into the liver, and typically die before day 50. These mice do not breed. A second subset of the transgenic mice have a less severe phenotype, and attempts to breed them are taking place.

**[0156]** The genomic structure for the mouse IL-B30 has been determined. A strategy for the production of IL-B30 knock-out mice has been developed, and constructs have been started.

#### **SEQUENCE LISTING**

##### **[0157]**

SEQ ID NO: 1 is a primate IL-B30 natural nucleic acid sequence.

SEQ ID NO: 2 is a primate IL-B30 natural amino acid sequence.

SEQ ID NO: 3 is a rodent IL-B30 natural nucleic acid sequence.

SEQ ID NO: 4 is a rodent IL-B30 natural amino acid sequence.

SEQ ID NO: 5 is a pig IL-B30.

SEQ ID NO: 6 is bovine G-CSF.

SEQ ID NO: 7 is feline G-CSF.

SEQ ID NO: 8 is human G-CSF.

SEQ ID NO: 9 is mouse G-CSF.

SEQ ID NO: 10 is otter IL-6.

SEQ ID NO: 11 is feline IL-6.

SEQ ID NO: 12 is human IL-6.

SEQ ID NO: 13 is sheep IL-6.

SEQ ID NO: 14 is mouse IL-6.

SEQ ID NO: 15 is chicken MGF.

SEQ ID NO: 16 is KSHV, kaposi's sarcoma herpes virus, a viral IL-6.

##### **(1) GENERAL INFORMATION:**

1. (i) APPLICANT: Schering Corporation
2. (ii) TITLE OF INVENTION: MAMMALIAN CYTOKINE; RELATED REAGENTS
3. (iii) NUMBER OF SEQUENCES: 16
4. (iv) CORRESPONDENCE ADDRESS:
  1. (A) ADDRESSEE: Schering Corporation
  2. (B) STREET: 2000 Galloping Hill Road
  3. (C) CITY: Kenilworth

4. (D) STATE: New Jersey
5. (E) COUNTRY: USA
6. (F) ZIP: 07033-0530
5. (v) COMPUTER READABLE FORM:
  1. (A) MEDIUM TYPE: Floppy disk
  2. (B) COMPUTER: Power Macintosh 7600/120
  3. (C) OPERATING SYSTEM: Windows
  4. (D) SOFTWARE: MS Word
6. (vi) CURRENT APPLICATION DATA:
  1. (A) APPLICATION NUMBER:
  2. (B) FILING DATE: July 24, 1998
  3. (C) CLASSIFICATION:
7. (vii) PRIOR APPLICATION DATA:
  1. (A) APPLICATION NUMBER: US 08/900,905
  2. (B) FILING DATE: 25-JUL-1997
8. (viii) ATTORNEY/AGENT INFORMATION:
  1. (A) NAME: Thampoe, Immac J.
  2. (B) REGISTRATION NUMBER: 36,322
  3. (C) REFERENCE/DOCKET NUMBER: DX0758K1
9. (ix) TELECOMMUNICATION INFORMATION:
  1. (A) TELEPHONE: (908) 298-5061
  2. (B) TELEFAX: (908) 298-5388

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 570 base pairs
2. (B) TYPE: nucleic acid
3. (C) STRANDEDNESS: single
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

1. (A) NAME/KEY: CDS
2. (B) LOCATION: 1..567

(ix) FEATURE:

1. (A) NAME/KEY: mat\_peptide
2. (B) LOCATION: 64..567

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

|                                                                                                                                                       |     |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| ATG CTG GGG AGC AGA GCT GTA ATG CTG CTG TTG CTG CTG CCC TGG ACA<br>Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr<br>-21 -20 -15 -10 | 48  |
| GCT CAG GGC AGA GCT GTG CCT GGG GGC AGC AGC CCT GCC TGG ACT CAG<br>Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln<br>-5 1 5 10       | 96  |
| TGC CAG CAG CTT TCA CAG AAG CTC TGC ACA CTG GCC TGG AGT GCA CAT<br>Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His<br>15 20 25        | 144 |
| CCA CTA GTG GGA CAC ATG GAT CTA AGA GAA GAG GGA GAT GAA GAG ACT<br>Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr<br>30 35 40        | 192 |
| ACA AAT GAT GTT CCC CAT ATC CAG TGT GGA GAT GGC TGT GAC CCC CAA<br>Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln<br>45 50 55        | 240 |
| GGA CTC AGG GAC AAC AGT CAG TTC TGC TTG CAA AGG ATC CAC CAG GGT<br>Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly<br>60 65 70 75     | 288 |
| CTG ATT TTT TAT GAG AAG CTG CTA GGA TCG GAT ATT TTC ACA GGG GAG<br>Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu<br>80 85 90        | 336 |
| CCT TCT CTG CTC CCT GAT AGC CCT GTG GCG CAG CTT CAT GCC TCC CTA<br>Pro Ser Leu Leu Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu<br>95 100 105      | 384 |
| CTG GGC CTC AGC CAA CTC CTG CAG CCT GAG GGT CAC CAC TGG GAG ACT<br>Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr<br>110 115 120     | 432 |
| CAG CAG ATT CCA AGC CTC AGT CCC AGC CAG CCA TGG CAG CGT CTC CTT<br>Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu<br>125 130 135     | 480 |
| CTC CGC TTC AAA ATC CTT CGC AGC CTC CAG GCC TTT GTG GCT GTA GCC<br>Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala<br>140 145 150 155 | 528 |
| GCC CGG GTC TTT GCC CAT GGA GCA GCA ACC CTG AGT CCC TAA<br>Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro<br>160 165                             | 570 |

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 189 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr
-21 -20                      -15                      -10

Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln
-5                      1                      5                      10

Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His
15                      20                      25

Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr
30                      35                      40

Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln
45                      50                      55

Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly
60                      65                      70                      75

Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu
80                      85                      90

Pro Ser Leu Leu Pro Asp Ser Pro Val Ala Gln Leu His Ala Ser Leu
95                      100                     105

Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr
110                     115                     120

Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu
125                     130                     135

Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala
140                     145                     150                     155

Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro
160                     165

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 1203 base pairs
2. (B) TYPE: nucleic acid
3. (C) STRANDEDNESS: single
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE: (A) NAME/KEY: CDS (B) LOCATION: 113..700

(ix) FEATURE: (A) NAME/KEY: mat\_peptide (B) LOCATION: 176..700

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

|                                                                                                                                                       |      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| CGCTTAGAAG TCGGACTACA GAGTTAGACT CAGAACCAAA GGAGGTGGAT AGGGGGTCCA                                                                                     | 60   |
| CAGCCCTGGT GCAGATCACA GAGCCAGCCA GATCTGAGAA GCAGGGAACA AG ATG<br>Met<br>-21                                                                           | 115  |
| CTG GAT TGC AGA GCA GTA ATA ATG CTA TGG CTG TTG CCC TGG GTC ACT<br>Leu Asp Cys Arg Ala Val Ile Met Leu Trp Leu Leu Pro Trp Val Thr<br>-20 -15 -10 -5  | 163  |
| CAG GGC CTG GCT GTG CCT AGG AGT AGC AGT CCT GAC TGG GCT CAG TGC<br>Gln Gly Leu Ala Val Pro Arg Ser Ser Ser Pro Asp Trp Ala Gln Cys<br>1 5 10          | 211  |
| CAG CAG CTC TCT CGG AAT CTC TGC ATG CTA GCC TGG AAC GCA CAT GCA<br>Gln Gln Leu Ser Arg Asn Leu Cys Met Leu Ala Trp Asn Ala His Ala<br>15 20 25        | 259  |
| CCA GCG GGA CAT ATG AAT CTA CTA AGA GAA GAA GAG GAT GAA GAG ACT<br>Pro Ala Gly His Met Asn Leu Leu Arg Glu Glu Glu Asp Glu Glu Thr<br>30 35 40        | 307  |
| AAA AAT AAT GTG CCC CGT ATC CAG TGT GAA GAT GGT TGT GAC CCA CAA<br>Lys Asn Asn Val Pro Arg Ile Gln Cys Glu Asp Gly Cys Asp Pro Gln<br>45 50 55 60     | 355  |
| GGA CTC AAG GAC AAC AGC CAG TTC TGC TTG CAA AGG ATC CGC CAA GGT<br>Gly Leu Lys Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile Arg Gln Gly<br>65 70 75        | 403  |
| CTG GCT TTT TAT AAG CAC CTG CTT GAC TCT GAC ATC TTC AAA GGG GAG<br>Leu Ala Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys Gly Glu<br>80 85 90        | 451  |
| CCT GCT CTA CTC CCT GAT AGC CCC ATG GAG CAA CTT CAC ACC TCC CTA<br>Pro Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr Ser Leu<br>95 100 105      | 499  |
| CTA GGA CTC AGC CAA CTC CTC CAG CCA GAG GAT CAC CCC CGG GAG ACC<br>Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg Glu Thr<br>110 115 120     | 547  |
| CAA CAG ATG CCC AGC CTG AGT TCT AGT CAG CAG TGG CAG CGC CCC CTT<br>Gln Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg Pro Leu<br>125 130 135 140 | 595  |
| CTC CGT TCC AAG ATC CTT CGA AGC CTC CAG GCC TTT TTG GCC ATA GCT<br>Leu Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala Ile Ala<br>145 150 155     | 643  |
| GCC CGG GTC TTT GCC CAC GGA GCA GCA ACT CTG ACT GAG CCC TTA GTG<br>Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro Leu Val<br>160 165 170     | 691  |
| CCA ACA GCT TAAGGATGCC CAGGTTCCCA TGGCTACCAT GATAAGACTA<br>Pro Thr Ala<br>175                                                                         | 740  |
| ATCTATCAGC CCAGACATCT ACCAGTTAAT TAACCCATTA GGACTTGTGC TGTCTTGT                                                                                       | 800  |
| TCGTTTGTGTT TGCCTGAAGG GCAAGGACAC CATTATTAAA GAGAAAAGAA ACAAACCCCA                                                                                    | 860  |
| GAGCAGGCAG CTGGCTAGAG AAAGGAGCTG GAGAAGAAGA ATAAAGTCTC GAGCCCTTGG                                                                                     | 920  |
| CCTTGGAAGC GGGCAAGCAG CTGCGTGGCC TGAGGGGAAG GGGGCGGTGG CATCGAGAAA                                                                                     | 980  |
| CTGTGAGAAA ACCCAGAGCA TCAGAAAAAG TGAGCCCAGG CTTTGGCCAT TATCTGTAAG                                                                                     | 1040 |
| AAAAACAAGA AAAGGGGAAC ATTATACCTT CCTGGGTGGC TCAGGGAAAT GTGCAGATGC                                                                                     | 1100 |
| ACAGTACTCC AGACAGCAGC TCTGTACCTG CCTGCTCTGT CCCTCAGTTC TAACAGAATC                                                                                     | 1160 |
| TAGTCACTAA GAACTAACAG GACTACCAAT ACGAACTGAC AAA                                                                                                       | 1203 |

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 196 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Leu Asp Cys Arg Ala Val Ile Met Leu Trp Leu Leu Pro Trp Val
-21 -20 -15 -10
Thr Gln Gly Leu Ala Val Pro Arg Ser Ser Ser Pro Asp Trp Ala Gln
-5 1 5 10
Cys Gln Gln Leu Ser Arg Asn Leu Cys Met Leu Ala Trp Asn Ala His
15 20 25
Ala Pro Ala Gly His Met Asn Leu Leu Arg Glu Glu Glu Asp Glu Glu
30 35 40
Thr Lys Asn Asn Val Pro Arg Ile Gln Cys Glu Asp Gly Cys Asp Pro
45 50 55
Gln Gly Leu Lys Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile Arg Gln
60 65 70 75
Gly Leu Ala Phe Tyr Lys His Leu Leu Asp Ser Asp Ile Phe Lys Gly
80 85 90
Glu Pro Ala Leu Leu Pro Asp Ser Pro Met Glu Gln Leu His Thr Ser
95 100 105
Leu Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Asp His Pro Arg Glu
110 115 120
Thr Gln Gln Met Pro Ser Leu Ser Ser Ser Gln Gln Trp Gln Arg Pro
125 130 135
Leu Leu Arg Ser Lys Ile Leu Arg Ser Leu Gln Ala Phe Leu Ala Ile
140 145 150 155
Ala Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Thr Glu Pro Leu
160 165 170
Val Pro Thr Ala
175

```

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 102 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

Ser Cys Leu Gln Arg Ile His Gln Gly Leu Val Phe Tyr Glu Lys Leu
1 5 10 15
Leu Gly Ser Asp Ile Phe Thr Gly Glu Pro Ser Leu His Pro Asp Gly
20 25 30
Ser Val Gly Gln Leu His Ala Ser Leu Leu Gly Leu Arg Gln Leu Leu
35 40 45
Gln Pro Glu Gly His His Trp Glu Thr Glu Gln Thr Pro Ser Pro Ser
50 55 60
Pro Ser Gln Pro Trp Gln Arg Leu Leu Leu Arg Leu Lys Ile Leu Arg
65 70 75 80
Ser Leu Gln Ala Phe Val Ala Val Ala Ala Arg Val Phe Ala His Gly
85 90 95
Ala Ala Thr Leu Ser Gln
100

```

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 174 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Thr Pro Leu Gly Pro Ala Arg Ser Leu Pro Gln Ser Phe Leu Leu Lys
1      5      10      15
Cys Leu Glu Gln Val Arg Lys Ile Gln Ala Asp Gly Ala Glu Leu Gln
20      25      30
Glu Arg Leu Cys Ala Ala His Lys Leu Cys His Pro Glu Glu Leu Met
35      40      45
Leu Leu Arg His Ser Leu Gly Ile Pro Gln Ala Pro Leu Ser Ser Cys
50      55      60
Ser Ser Gln Ser Leu Gln Leu Arg Gly Cys Leu Asn Gln Leu His Gly
65      70      75      80
Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Ala Gly Ile Ser
85      90      95
Pro Glu Leu Ala Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Thr Asp
100     105     110
Phe Ala Thr Asn Ile Trp Leu Gln Met Glu Asp Leu Gly Ala Ala Pro
115     120     125
Ala Val Gln Pro Thr Gln Gly Ala Met Pro Thr Phe Thr Ser Ala Phe
130     135     140
Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser Gln Leu His Arg Phe
145     150     155     160
Leu Glu Leu Ala Tyr Arg Gly Leu Arg Tyr Leu Ala Glu Pro
165     170

```

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 174 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

Thr Pro Leu Gly Pro Thr Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
1      5      10      15
Cys Leu Glu Gln Val Arg Lys Val Gln Ala Asp Gly Thr Ala Leu Gln
20      25      30
Glu Arg Leu Cys Ala Ala His Lys Leu Cys His Pro Glu Glu Leu Val
35      40      45
Leu Leu Gly His Ala Leu Gly Ile Pro Gln Ala Pro Leu Ser Ser Cys
50      55      60
Ser Ser Gln Ala Leu Gln Leu Thr Gly Cys Leu Arg Gln Leu His Ser
65      70      75      80
Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Ala Gly Ile Ser
85      90      95
Pro Glu Leu Ala Pro Thr Leu Asp Met Leu Gln Leu Asp Ile Thr Asp
100     105     110
Phe Ala Ile Asn Ile Trp Gln Gln Met Glu Asp Val Gly Met Ala Pro
115     120     125
Ala Val Pro Pro Thr Gln Gly Thr Met Pro Thr Phe Thr Ser Ala Phe
130     135     140
Gln Arg Arg Ala Gly Gly Thr Leu Val Ala Ser Asn Leu Gln Ser Phe
145     150     155     160
Leu Glu Val Ala Tyr Arg Ala Leu Arg His Phe Thr Lys Pro
165     170

```

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 177 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant

## 4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
1      5      10      15
Cys Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln
20      25      30
Glu Lys Leu Val Ser Glu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu
35      40      45
Glu Leu Val Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu
50      55      60
Ser Ser Cys Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln
65      70      75      80
Leu His Ser Gly Leu Met Ala Pro Ala Leu Gln Pro Thr Gln Gly Ala
85      90      95
Met Pro Ala Phe Ala Ser Ala Phe Gln Arg Arg Ala Gly Gly Val Leu
100      105      110
Val Ala Ser His Leu Gln Ser Phe Leu Glu Val Ser Tyr Arg Val Leu
115      120      125
Arg His Leu Ala Gln Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Glu
130      135      140
Gly Ile Ser Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp
145      150      155      160
Val Ala Asp Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly
165      170      175
Pro

```

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 178 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Val Pro Leu Val Thr Val Ser Ala Leu Pro Pro Ser Leu Pro Leu Pro
1      5      10      15
Arg Ser Phe Leu Leu Lys Ser Leu Glu Gln Val Arg Lys Ile Gln Ala
20      25      30
Ser Gly Ser Val Leu Leu Glu Gln Leu Cys Ala Thr Tyr Lys Leu Cys
35      40      45
His Pro Glu Glu Leu Val Leu Leu Gly His Ser Leu Gly Ile Pro Lys
50      55      60
Ala Ser Leu Ser Gly Cys Ser Ser Gln Ala Leu Gln Gln Thr Gln Cys
65      70      75      80
Leu Ser Gln Leu His Ser Gly Leu Cys Leu Tyr Gln Gly Leu Leu Gln
85      90      95
Ala Leu Ser Gly Ile Ser Pro Ala Leu Ala Pro Thr Leu Asp Leu Leu
100      105      110
Gln Leu Asp Val Ala Asn Phe Ala Thr Thr Ile Trp Gln Gln Met Glu
115      120      125
Asn Leu Gly Val Ala Pro Thr Val Gln Pro Thr Gln Ser Ala Met Pro
130      135      140
Ala Phe Thr Ser Ala Phe Gln Arg Arg Ala Gly Gly Val Leu Ala Ile
145      150      155      160
Ser Tyr Leu Gln Gly Phe Leu Glu Thr Ala Arg Leu Ala Leu His His
165      170      175
Leu Ala

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 186 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

Ala Phe Pro Thr Pro Gly Pro Leu Gly Gly Asp Ser Lys Asp Asp Ala
 1         5         10        15
Thr Ser Asn Arg Pro Pro Leu Thr Ser Ala Asp Lys Met Glu Asp Phe
 20        25        30
Ile Lys Phe Ile Leu Gly Lys Ile Ser Ala Leu Arg Asn Glu Met Cys
 35        40        45
Asp Lys Tyr Asn Lys Cys Glu Asp Ser Lys Glu Val Leu Ala Glu Asn
 50        55        60
Asn Leu Asn Leu Pro Lys Leu Ala Glu Lys Asp Arg Cys Phe Gln Ser
 65        70        75        80
Arg Phe Asn Gln Glu Thr Cys Leu Thr Arg Ile Thr Thr Gly Leu Gln
 85        90        95
Glu Phe Gln Ile His Leu Lys Tyr Leu Glu Ser Asn Tyr Glu Gly Asn
100       105       110
Lys Asp Asn Ala His Ser Val Tyr Ile Ser Thr Lys His Leu Leu Gln
115       120       125
Thr Leu Arg Pro Met Asn Gln Ile Glu Val Thr Thr Pro Asp Pro Thr
130       135       140
Thr Asp Ala Ser Leu Gln Ala Leu Phe Lys Ser Gln Asp Lys Trp Leu
145       150       155       160
Lys His Thr Thr Ile His Leu Ile Leu Arg Arg Leu Glu Asp Phe Leu
165       170       175
Gln Phe Ser Leu Arg Ala Ile Arg Ile Met
180       185

```

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 183 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Ala Phe Pro Thr Pro Gly Pro Leu Gly Gly Asp Ala Thr Ser Asn Arg  
 1 5 10 15  
 Leu Pro Leu Thr Pro Ala Asp Lys Met Glu Glu Leu Ile Lys Tyr Ile  
 20 25 30  
 Leu Gly Lys Ile Ser Ala Leu Lys Lys Glu Met Cys Asp Asn Tyr Asn  
 35 40 45  
 Lys Cys Glu Asp Ser Lys Glu Ala Leu Ala Glu Asn Asn Leu Asn Leu  
 50 55 60  
 Pro Lys Leu Ala Glu Lys Asp Gly Cys Phe Gln Ser Gly Phe Asn Gln  
 65 70 75 80  
 Glu Thr Cys Leu Thr Arg Ile Thr Thr Gly Leu Gln Glu Phe Gln Ile  
 85 90 95  
 Tyr Leu Lys Phe Leu Gln Asp Lys Tyr Glu Gly Asp Lys Glu Asn Ala  
 100 105 110  
 Lys Ser Val Tyr Thr Ser Thr Asn Val Leu Leu Gln Met Leu Lys Arg  
 115 120 125  
 Lys Gly Lys Asn Gln Asp Glu Val Thr Ile Pro Val Pro Thr Val Glu  
 130 135 140  
 Val Gly Leu Gln Leu Ser Cys Ser His Arg Arg Val Ala Glu Ala His  
 145 150 155 160  
 Asn Asn His Leu Thr Leu Arg Arg Leu Glu Asp Phe Leu Gln Leu Arg  
 165 170 175  
 Leu Arg Ala Val Arg Ile Met  
 180

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 188 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: single
4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ala Phe Pro Ala Pro Val Pro Pro Gly Glu Asp Ser Lys Asp Val Ala  
 1 5 10 15  
 Ala Pro His Arg Gln Pro Leu Thr Ser Ser Glu Arg Ile Asp Lys Gln  
 20 25 30  
 Ile Arg Tyr Ile Leu Asp Gly Ile Ser Ala Leu Arg Lys Glu Thr Cys  
 35 40 45  
 Asn Lys Ser Asn Met Cys Glu Ser Ser Lys Glu Ala Leu Ala Glu Asn  
 50 55 60  
 Asn Leu Asn Leu Pro Lys Met Ala Glu Lys Asp Gly Cys Phe Gln Ser  
 65 70 75 80  
 Gly Phe Asn Glu Glu Thr Cys Leu Val Lys Ile Ile Thr Gly Leu Leu  
 85 90 95  
 Glu Phe Glu Val Tyr Leu Glu Tyr Leu Gln Asn Arg Phe Glu Ser Ser  
 100 105 110  
 Glu Glu Gln Ala Arg Ala Val Gln Met Ser Thr Lys Val Leu Ile Gln  
 115 120 125  
 Phe Leu Gln Lys Lys Ala Lys Asn Leu Asp Ala Ile Thr Thr Pro Asp  
 130 135 140  
 Pro Thr Thr Asn Ala Ser Leu Leu Thr Lys Leu Gln Ala Gln Asn Gln  
 145 150 155 160  
 Trp Leu Gln Asp Met Thr Thr His Leu Ile Leu Arg Ser Phe Lys Glu  
 165 170 175  
 Phe Leu Gln Ser Ser Leu Arg Ala Leu Arg Gln Met  
 180 185

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 184 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

Ala Phe Pro Thr Pro Gly Pro Leu Gly Glu Asp Phe Lys Asn Asp Thr
1      5      10      15
Thr Pro Ser Arg Leu Leu Leu Thr Thr Pro Glu Lys Thr Glu Ala Leu
20      25      30
Ile Lys His Ile Val Asp Lys Ile Ser Ala Ile Arg Lys Glu Ile Cys
35      40      45
Glu Lys Asn Asp Glu Cys Glu Asn Ser Lys Glu Thr Leu Ala Glu Asn
50      55      60
Lys Leu Lys Leu Pro Lys Met Glu Glu Lys Asp Gly Cys Phe Gln Ser
65      70      75      80
Gly Phe Asn Gln Ala Ile Cys Leu Ile Lys Thr Thr Ala Gly Leu Leu
85      90      95
Glu Tyr Gln Ile Tyr Leu Asp Phe Leu Gln Asn Glu Phe Glu Gly Asn
100     105     110
Gln Glu Thr Val Met Glu Leu Gln Ser Ser Ile Arg Thr Leu Ile Gln
115     120     125
Ile Leu Lys Glu Lys Ile Ala Gly Leu Ile Thr Thr Pro Ala Thr His
130     135     140
Thr Asp Met Leu Glu Lys Met Gln Ser Ser Asn Glu Trp Val Lys Asn
145     150     155     160
Ala Lys Val Ile Ile Ile Leu Arg Ser Leu Glu Asn Phe Leu Gln Phe
165     170     175
Ser Leu Arg Ala Ile Arg Met Lys
180

```

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 188 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

Ala Phe Pro Thr Ser Gln Val Arg Arg Gly Asp Phe Thr Glu Asp Thr
1      5      10      15
Thr Pro Asn Arg Pro Val Tyr Thr Thr Ser Gln Val Gly Gly Leu Ile
20      25      30
Thr His Val Leu Trp Glu Ile Val Glu Met Arg Lys Glu Leu Cys Asn
35      40      45
Gly Asn Ser Asp Cys Met Asn Asn Asp Asp Ala Leu Ala Glu Asn Asn
50      55      60
Leu Lys Leu Pro Glu Ile Gln Arg Asn Asp Gly Cys Tyr Gln Thr Gly
65      70      75      80
Tyr Asn Gln Glu Ile Cys Leu Leu Lys Ile Ser Ser Gly Leu Leu Glu
85      90      95
Tyr His Ser Tyr Leu Glu Tyr Met Lys Asn Asn Leu Lys Asp Asn Lys
100     105     110
Lys Asp Lys Ala Arg Val Leu Gln Arg Asp Thr Glu Thr Leu Ile His
115     120     125
Ile Phe Asn Gln Glu Val Lys Asp Leu His Lys Ile Val Leu Pro Thr
130     135     140
Pro Ile Ser Asn Ala Leu Leu Thr Asp Lys Leu Glu Ser Gln Lys Glu
145     150     155     160
Trp Leu Arg Thr Lys Thr Ile Gln Phe Ile Leu Lys Ser Leu Glu Glu
165     170     175
Phe Leu Lys Val Thr Leu Arg Ser Thr Arg Gln Thr
180     185

```

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 178 amino acids
2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

```

Ala Pro Leu Ala Glu Leu Ser Gly Asp His Asp Phe Gln Leu Phe Leu
1      5      10      15
His Lys Asn Leu Glu Phe Thr Arg Lys Ile Arg Gly Asp Val Ala Ala
20      25      30
Leu Gln Arg Ala Val Cys Asp Thr Phe Gln Leu Cys Thr Glu Glu Glu
35      40      45
Leu Gln Leu Val Gln Pro Asp Pro His Leu Val Gln Ala Pro Leu Asp
50      55      60
Gln Cys His Lys Arg Gly Phe Gln Ala Glu Val Cys Phe Thr Gln Ile
65      70      75      80
Arg Ala Gly Leu His Ala Tyr His Asp Ser Leu Gly Ala Val Leu Arg
85      90      95
Leu Leu Pro Asn His Thr Thr Leu Val Glu Thr Leu Gln Leu Asp Ala
100     105     110
Ala Asn Leu Ser Ser Asn Ile Gln Gln Gln Met Glu Asp Leu Gly Leu
115     120     125
Asp Thr Val Thr Leu Pro Ala Glu Gln Arg Ser Pro Pro Pro Thr Phe
130     135     140
Ser Gly Pro Phe Gln Gln Gln Val Gly Gly Phe Phe Ile Leu Ala Asn
145     150     155     160
Phe Gln Arg Phe Leu Glu Thr Ala Tyr Arg Ala Leu Arg His Leu Ala
165     170     175
Arg Leu

```

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

1. (A) LENGTH: 185 amino acids

2. (B) TYPE: amino acid
3. (C) STRANDEDNESS: not relevant
4. (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

```

Thr Arg Gly Lys Leu Pro Asp Ala Pro Glu Phe Glu Lys Asp Leu Leu
1      5      10      15
Ile Gln Arg Leu Asn Trp Met Leu Trp Val Ile Asp Glu Cys Phe Arg
20      25      30
Asp Leu Cys Tyr Arg Thr Gly Ile Cys Lys Gly Ile Leu Glu Pro Ala
35      40      45
Ala Ile Phe His Leu Lys Leu Pro Ala Ile Asn Asp Thr Asp His Cys
50      55      60
Gly Leu Ile Gly Phe Asn Glu Thr Ser Cys Leu Lys Lys Leu Ala Asp
65      70      75      80
Gly Phe Phe Glu Phe Glu Val Leu Phe Lys Phe Leu Thr Thr Glu Phe
85      90      95
Gly Lys Ser Val Ile Asn Val Asp Val Met Glu Leu Leu Thr Lys Thr
100     105     110
Leu Gly Trp Asp Ile Gln Glu Glu Leu Asn Lys Leu Thr Lys Thr His
115     120     125
Tyr Ser Pro Pro Lys Phe Asp Arg Gly Leu Leu Gly Arg Leu Gln Gly
130     135     140
Leu Lys Tyr Trp Val Arg His Phe Ala Ser Phe Tyr Val Leu Ser Ala
145     150     155     160
Met Glu Lys Phe Ala Gly Gln Ala Val Arg Val Leu Asp Ser Ile Pro
165     170     175
Asp Val Thr Pro Asp Val His Asp Lys
180     185

```

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- [US4859609A \[0043\]](#)
- [US3617837A \[0060\]](#)
- [US3850752A \[0060\]](#)
- [US3939350A \[0060\]](#)
- [US3996345A \[0060\]](#)
- [US4277437A \[0060\]](#)
- [US4275149A \[0060\]](#)
- [US4366241A \[0060\]](#)
- [US4816567A \[0060\]](#)
- [US4642334A \[0060\]](#)
- [EP6403564A \[0107\]](#)
- [US3646090A \[0115\]](#)
- [US3940475A \[0115\]](#)
- [US4659678A \[0116\]](#)
- [US08900905B \[0157\]](#)

## Non-patent literature cited in the description

- **THOMSON**The Cytokine HandbookAcademic Press19940000 [0003]
- **METCALFNICOLA**The Hematopoietic Colony Stimulating FactorsCambridge University Press19950000 [0003] [0094] [0124]
- **AGGARWALGUTTERMAN**Human CytokinesBlackwell Pub19910000 [0003] [0094] [0124]
- **ITO et al.**Pig STAT2 cDNA EMBL, 1997, [0003]
- **NAGATA**Thomson The Cytokine HandbookAcademic Press19940000 [0009]
- **HIRANO**Thomson The Cytokine HandbookAcademic Press19940000 [0009]
- **PAUL**Fundamental ImmunologyRaven Press19940000 [0014]
- Goodman & Gilman's: The Pharmacological Bases of TherapeuticsPergamon Press [0025]
- **NEEDLEHAM et al.**J. Mol. Biol., 1970, vol. 48, 443-453 [0031]
- **SANKOFF et al.**Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence ComparisonAddison-Wesley19830000 [0031]
- **KUNKEL et al.**Methods in Enzymol., 1987, vol. 154, 367-382 [0033]
- **CUNNINGHAM et al.**Science, 1989, vol. 243, 1330-1336 [0035]
- **O'DOWD et al.**J. Biol. Chem., 1988, vol. 263, 15985-15992 [0035] [0050]
- **BEAUCAGECARRUTHERS**Tetra. Letts., 1981, vol. 22, 1859-1862 [0036]
- **BAZAN et al.**Nature, 1996, vol. 379, 591- [0037]
- **LODI et al.**Science, 1994, vol. 263, 1762-1766 [0037]
- **SAYLEMILNER-WHITE**TIBS, 1995, vol. 20, 374-376 [0037]
- **GRONENBERG et al.**Protein Engineering, 1991, vol. 4, 263-269 [0037]
- **LUNDBLADNOYES**Chemical Reagents for Protein ModificationCRC Press, Inc.19880000vol. 1-2, [0041]
- Techniques in Protein ChemistryAcademic Press19890000 [0041]
- **WONG**Chemistry of Protein Conjugation and Cross LinkingCRC Press19910000 [0041]
- **ELBEIN**Ann. Rev. Biochem., 1987, vol. 56, 497-534 [0042]
- **GODOWSKI et al.**Science, 1988, vol. 241, 812-816 [0043]
- **SAMBROOK et al.**Molecular Cloning: A Laboratory ManualCold Spring Harbor Laboratory19890000vol. 1-3, [0044]
- Current Protocols in Molecular BiologyGreene and Wiley19930000 [0044]
- **MERRIFIELD**J. Amer. Chem. Soc., 1963, vol. 85, 2149-2156 [0044]
- **MERRIFIELD**Science, 1986, vol. 232, 341-347 [0044]
- **ATHERTON et al.**Solid Phase Peptide Synthesis: A Practical ApproachIRL Press19890000 [0044]
- **GRANTS**Synthetic Peptides: A User's GuideW.H. Freeman19920000 [0044]
- **CUNNINGHAM et al.**Science, 1989, vol. 243, 1339-1336 [0050]
- **LECHLEITER et al.**EMBO J., 1990, vol. 9, 4381-4390 [0050]
- Immunology: A Practical GuideAcademic Press19870000 [0056]
- Principles and Practice of ImmunoassayStockton Press19910000 [0056]
- Nonisotopic ImmunoassayPlenum Press19880000 [0056]
- **HARPERROW**Hoeber Medical DivisionMicrobiology, 1969, [0058]
- **LANDSTEINER**Specificity of Serological ReactionsDover Publications19620000 [0058]
- **WILLIAMS et al.**Methods in Immunology and ImmunochemistryAcademic Press19670000vol. 1, [0058]
- **HARLOWLANE**Antibodies: A Laboratory ManualCSH Press19880000 [0058] [0059]
- Basic and Clinical ImmunologyLange Medical Publications [0059]
- **GODING**Monoclonal Antibodies: Principles and PracticeAcademic Press19860000 [0059]
- **KOHLERMILSTEIN**Nature, 1975, vol. 256, 495-497 [0059]
- **HUSE et al.**Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage LambdaScience, 1989, vol. 246, 1275-1281 [0060]
- **WARD et al.**Nature, 1989, vol. 341, 544-546 [0060]
- **QUEEN et al.**Proc. Nat'l Acad. Sci. USA, 1989, vol. 86, 10029-10033 [0060]
- **WILCHEK et al.**Meth. Enzymol., 1984, vol. 104, 3-55 [0061]
- **COLIGAN**Current Protocols in ImmunologyWiley/Greene19910000 [0064] [0124] [0140] [0141]
- **HARLOWLANE**Antibodies: A Laboratory ManualCold Spring Harbor Press19890000 [0064] [0140] [0141]
- Transgenic Animals**GOODNOW**Encyclopedia of ImmunologyAcademic Press199200001502-1504 [0074]
- **TRAVIS**Science, 1992, vol. 256, 1392-1394 [0074]

- **KUHN et al.** Science, 1991, vol. 254, 707-710 [0074]
- **CAPECCHI** Science, 1989, vol. 244, 1288- [0074]
- **Teratocarcinomas and Embryonic Stem Cells: A Practical Approach** IRL Press [0074]
- **ROSENBERG J.** Clinical Oncology, 1992, vol. 10, 180-199 [0074]
- **KANEHISA** Nuc. Acids Res., 1984, vol. 12, 203-213 [0075]
- **WETMURDAVIDSON J.** Mol. Biol., 1968, vol. 31, 349-370 [0076]
- **SMITHWATERMAN** Adv. Appl. Math., 1981, vol. 2, 482- [0078]
- **NEEDLEMANWUNSCH J.** Mol. Biol., 1970, vol. 48, 443- [0078]
- **PEARSONLIPMAN** Proc. Nat'l Acad. Sci. USA, 1988, vol. 85, 2444- [0078]
- **FENGDOOLITTLE J.** Mol. Evol., 1987, vol. 35, 351-360 [0079]
- **HIGGINSHARP** CABIOS, 1989, vol. 5, 151-153 [0079]
- **ALTSCHUL et al.** J. Mol. Biol., 1990, vol. 215, 403-410 [0080]
- **HENIKOFFHENIKOFF** Proc. Nat'l Acad. Sci. USA, 1989, vol. 89, 10915- [0080]
- **KARLINALTSCHUL** Proc. Nat'l Acad. Sci. USA, 1993, vol. 90, 5873-5787 [0081]
- **OKAYAMABERG** Mol. Cell. Biol., 1982, vol. 2, 161-170 [0084]
- **GUBLERHOFFMAN** Gene, 1983, vol. 25, 263-269 [0084]
- **DNA Cloning: A Practical Approach** IRL Press 19840000 [0084]
- **POUWELS et al.** Cloning Vectors: A Laboratory Manual Elsevier 19850000 [0086]
- **Vectors: A Survey of Molecular Cloning Vectors and Their Uses** Butterworth 19880000 [0086]
- **BALBASBOLIVAR** Methods in Enzymology, 1990, vol. 185, 14-37 [0087]
- **AUSUBEL et al.** Current Protocols in Molecular Biology Greene and Wiley 19930000 [0087]
- **OKAYAMA et al.** Mol. Cell Biol., 1985, vol. 5, 1136-1142 [0088]
- **THOMAS et al.** Cell, 1987, vol. 51, 503-512 [0088]
- **MILLER** Ann. Rev. Microbiol., 1988, vol. 42, 177-199 [0088]
- **LUCKOWSUMMERS** Bio/Technology, 1988, vol. 6, 47-55 [0089]
- **KAUFMAN** Meth. Enzymol., 1990, vol. 185, 487-511 [0089]
- **LOW** Biochim. Biophys. Acta, 1989, vol. 988, 427-454 [0090]
- **TSE et al.** Science, 1985, vol. 230, 1003-1008 [0090]
- **BRUNNER et al.** J. Cell Biol., 1991, vol. 114, 1275-1283 [0090]
- **STEWARTYOUNG** Solid Phase Peptide Synthesis Pierce Chemical Co. 19840000 [0091]
- **BODANSZKYBODANSZKY** The Practice of Peptide Synthesis Springer-Verlag 19840000 [0091]
- **BODANSZKY** The Principles of Peptide Synthesis Springer-Verlag 19840000 [0091]
- **Techniques in Protein Chemistry II** Academic Press 19910000 [0091]
- **The Cytokine Handbook** Academic Press 19940000 [0094] [0124]
- **Immunological Diseases** Little, Brown and Co. vol. 1 and 2, [0096]
- **The Merck Manual of Diagnosis and Therapy** Merck & Co. [0098]
- **THORN et al.** Harrison's Principles of Internal Medicine McGraw-Hill [0098]
- **Oxford Textbook of Medicine** Oxford University Press [0098]
- **Appleton and Lange, Norwalk, Connecticut** Basic and Clinical Immunology 19910000 [0098]
- **Immunological Diseases** Little, Brown and Co. [0098]
- **Goodman and Gilman's: The Pharmacological Bases of Therapeutics** Pergamon Press 19900000 [0101] [0102]
- **Remington's Pharmaceutical Sciences** Mack Publishing Co. 19900000 [0101] [0102]
- **Merck Index** Merck & Co. [0101]
- **LANGER** Science, 1990, vol. 249, 1527-1533 [0101]
- **Pharmaceutical Dosage Forms: Parenteral Medications** Dekker 19930000 [0102]
- **Pharmaceutical Dosage Forms: Tablets** Dekker 19900000 [0102]
- **Pharmaceutical Dosage Forms: Disperse Systems** Dekker 19900000 [0102]
- **FODOR et al.** Science, 1991, vol. 251, 767-773 [0103]
- **SOMOZA et al.** J. Exp. Med., 1993, vol. 178, 549-558 [0104]
- **ROSTSANDER** Proteins, 1994, vol. 19, 55-72 [0104]
- **KINGSTERNBERG** Protein Sci., 1996, vol. 5, 2298-2310 [0104]
- **PARCE et al.** Science, 1989, vol. 246, 243-247 [0106]
- **OWICKI et al.** Proc. Nat'l Acad. Sci. USA, 1990, vol. 87, 4007-4011 [0106]
- **BLUNDELLJOHNSON** Protein Crystallography Academic Press 19760000 [0108]
- **VAN VUNAKIS et al.** Meth. Enzymol., 1980, vol. 70, 1-525 [0112]
- **HARLOWLANE** Antibodies: A Laboratory Manual CSH Press 19800000 [0112]
- **Current Protocols in Immunology** Greene and Wiley 19930000 [0112]

- Current Protocols in Immunology Greene and Wiley 19930000 vol. 1, [0116]
- **RATTLE et al.** Clin. Chem., 1984, vol. 30, 1457-1461 [0116]
- **LANGER-SAFER et al.** Proc. Nat'l Acad. Sci., 1982, vol. 79, 4381-4385 [0118]
- **CASKEY** Science, 1987, vol. 236, 962-967 [0118]
- **WILCHEK et al.** Anal. Biochem., 1988, vol. 171, 1-32 [0118]
- **VIALLET et al.** Progress in Growth Factor Res., 1989, vol. 1, 89-97 [0119]
- **GEARING et al.** EMBO J., 1989, vol. 8, 3667-3676 [0120]
- **HO et al.** Proc. Nat'l Acad. Sci. USA, 1993, vol. 90, 11267-11271 [0120]
- **LIU et al.** J. Immunol., 1994, vol. 152, 1821-29 [0120]
- **SEEDARUFFO** Proc. Nat'l Acad. Sci. USA, 1987, vol. 84, 3365-3369 [0120]
- **MANIATIS et al.** Molecular Cloning, A Laboratory Manual Cold Spring Harbor Laboratory 19820000 [0123]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual CSH Press 19890000 vol. 1-3, [0123]
- **AUSUBEL et al.** Biology Greene Publishing Associates [0123]
- **AUSUBEL et al.** Current Protocols in Molecular Biology Wiley/Greene 19870000 [0123]
- PCR Protocols: A Guide to Methods and Application Academic Press 19900000 [0123]
- Guide to Protein Purification **DEUTSCHER** Methods in Enzymology 19900000 vol. 182, [0123]
- **COLIGAN et al.** Current Protocols in Protein Science John Wiley and Sons 19950000 [0123]
- A Practical Guide to Protein and Peptide Purification for Microsequencing Academic Press 19930000 [0123]
- **HOCHULI** Chemische Industrie, 1989, vol. 12, 69-70 [0123]
- Purification of Recombinant Proteins with Metal Chelate Absorbent **HOCHULI** Genetic Engineering, Principle and Methods Plenum Press 19900000 vol. 12, 87-98 [0123]
- **CROWE et al.** OIAexpress: The High Level Expression & Protein Purification System QUIAGEN, Inc. 19920000 [0123]
- Weir's Handbook of Experimental Immunology Blackwell Science 19960000 vol. 1-4, [0124]
- Methods in Enzymology, vol. 70,73,74,84,92,93,108,116,121,132,150,162,163, [0124]
- **KOYOMA et al.** Cell, 1996, vol. 87, 1069-1078 [0125]
- **MCEVOY et al.** J. Exp. Med., 1997, vol. 185, 2069-2077 [0125]
- **ROSS** Nature, 1993, vol. 362, 801-809 [0125]
- **REKHTERGORDON** Am. J. Pathol., 1995, vol. 147, 668-677 [0125]
- **THYBERG et al.** Atherosclerosis, 1990, vol. 10, 966-990 [0125]
- **GUMBINER** Cell, 1996, vol. 84, 345-357 [0125]
- Neuroscience Protocols Elsevier 19950000 [0126]
- Methods in Neurosciences Academic Press [0126]
- Neuromethods Humana Press [0126]
- Handbook of Human Growth and Developmental Biology CRC Press [0126]
- Molecular Techniques and Approaches in Developmental Biology Interscience [0126]
- **MELAMED et al.** Flow Cytometry and Sorting Wiley-Liss, Inc. 19900000 [0127]
- **SHAPIRO** Practical Flow Cytometry Liss 19880000 [0127]
- **ROBINSON et al.** Handbook of Flow Cytometry Methods Wiley-Liss 19930000 [0127]
- **OPENSHAW et al.** J. Exp. Med., 1995, vol. 182, 1357-1367 [0133] [0133]
- **HO et al.** Proc. Nat'l Acad. Sci. USA, 1993, vol. 90, 11267-11271 [0150]
- **HO et al.** Mol. Cell. Biol., 1995, vol. 15, 5043-5053 [0150]
- **LIU et al.** J. Immunol., 1994, vol. 152, 1821-1829 [0150]
- **WAAL MALEFYT et al.** J. Exp. Med., 1991, vol. 174, 1209-1220 [0151]
- **WAAL MALEFYT et al.** J. Exp. Med., 1991, vol. 174, 915-924 [0151]
- **FIORENTINO et al.** J. Immunol., 1991, vol. 147, 3815-3822 [0151]
- **FIORENTINO et al.** J. Immunol., 1991, vol. 146, 3444-3451 [0151]
- **GROUX et al.** J. Exp. Med., 1996, vol. 184, 19-29 [0151]
- **HSU et al.** Internat. Immunol., 1992, vol. 4, 563-569 [0152]
- **SCHWARZ et al.** J. Immunother, 1994, vol. 16, 95-104 [0152]
- **DEFRANCE et al.** J. Exp. Med., 1992, vol. 175, 671-682 [0153]
- **ROUSSET et al.** Proc. Nat'l Acad. Sci. USA, 1992, vol. 89, 1890-1893 [0153]
- **HOGAN et al.** Manipulating the Mouse Embryo: A Laboratory Manual Cold Spring Harbor Laboratory Press 19950000 [0154]

**Patentkrav**

1. Antistof, eller antigenbindende fragment deraf, som specifikt binder til den modne aminosyresekvens SEQ ID NO.: 2 eller 4 eller en variant deraf med 1, 2, 3, 5 eller 7 konservative aminosyresubstitutioner.
2. Antistof, eller antigenbindende fragment deraf, ifølge krav 1, hvor antistoffet specifikt binder til den modne aminosyresekvens SEQ ID NO.: 2 eller en variant deraf med 1, 2, 3, 5 eller 7 konservative aminosyresubstitutioner.
3. Antistof, eller antigenbindende fragment deraf, ifølge krav 2, hvor antistoffet, eller det antigenbindende fragment deraf, er et monoklonalt antistof eller fragment deraf.
4. Antistof, eller antigenbindende fragment deraf, ifølge krav 2, hvor antistoffet eller det antigenbindende fragment er:
- a) et Fab-fragment;
  - b) et Fab'-fragment;
  - c) et F(ab)<sub>2</sub>-fragment; eller
  - d) et enkelt kædeantistof.
5. Antistof, eller antigenbindende fragment deraf, ifølge krav 2, hvor antistoffet, eller det antigenbindende fragment deraf, er et kimærisk eller humaniseret antistof, eller et antigenbindende fragment deraf.
6. Steril sammensætning omfattende antistoffet, eller det antigenbindende fragment deraf, ifølge et af kravene 1-5.
7. Farmaceutisk formulering omfattende:
- a) antistoffet, eller det antigenbindende fragment deraf, ifølge et af kravene 1-5; og
  - b) en eller flere farmaceutisk acceptable bærere.