

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46374 A1

(51) International Patent Classification⁷: **C11D 3/50**, D06M 13/00

(74) Agents: **REED, T., David** et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).

(21) International Application Number: PCT/US00/34833

(22) International Filing Date: 20 December 2000 (20.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

99870277.3	22 December 1999 (22.12.1999)	EP
00870070.0	13 April 2000 (13.04.2000)	EP
00202168.1	22 June 2000 (22.06.2000)	EP

(71) Applicant (for all designated States except US): **THE PROCTER & GAMBLE COMPANY** [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SMETS, Johan** [BE/BE]; Bollenberg 79, B-3210 Lubbeek (BE). **WEVERS, Jean** [BE/BE]; Heide 17, B-1840 Steenhuffel (BE). **SAINI, Guarav** [IN/JP]; 1-4-127-506, Koyocho-Naka, Higashinada-ku, Kobe 658-0032 (JP). **ROSALDO, Rafael, Trujillo** [MX/BE]; Ave. Princesse Paola #39, B-Waterloo (BE).

(81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/46374 A1

(54) Title: LAUNDRY AND CLEANING AND/OR FABRIC CARE COMPOSITIONS

(57) Abstract: There is provided a laundry and/or cleaning and/or fabric care composition comprising a benefit agent whereby said benefit agent is carried with a carrier material, thereby providing an enhanced deposition on the treated fabric of the benefit agent.

Laundry and cleaning and/or fabric care compositions

5

10

Technical field of the Invention

15

The present invention relates to a laundry and/or cleaning and/or fabric care composition comprising a benefit agent, for imparting sustained release of the benefit agent on the treated surfaces like fabrics, in particular dry fabrics.

20

Background of the invention

25

Perfumed products are well-known in the art. However, consumer acceptance of such perfumed products like laundry and cleaning products is determined not only by the performance achieved with these products but also by the aesthetics associated therewith. The perfume components are therefore an important aspect of the successful formulation of such commercial products.

30

It is also desired by consumers for treated surfaces like fabrics to maintain the pleasing fragrance over time. Indeed, perfume additives make such compositions more aesthetically pleasing to the consumer, and in some cases the perfume imparts a pleasant fragrance to surfaces, like fabrics, treated therewith. However, the amount of perfume 5 carried-over from an aqueous laundry or cleaning bath onto fabrics is often marginal and does not last long on the surface. Furthermore, fragrance materials are often very costly and their inefficient use in laundry and cleaning compositions and ineffective delivery to surfaces like fabrics results in a very high cost to both consumers and laundry and cleaning manufacturers. Industry, therefore, continues to seek with 10 urgency for more efficient and effective fragrance delivery in laundry and cleaning products, especially for improvement in the provision of long-lasting fragrance to the surfaces like fabrics.

Further, after drying fabrics under the sun, fabrics obtain a “sun-dried type” of odor. 15 Consumers often prefer this to a standard perfume odor. Also they often consider fabrics with these odors to be cleaner. Because consumers like the odor, they like to dry fabrics under the sun. In some countries, however, consumer cannot dry their fabrics outside because the air is not clean, or there is too much rain. As a result, they have to dry their fabrics indoors and cannot expect to enjoy this benefit of having a 20 “sun-dried type” of odor on their fabrics.

Recently, a new class of materials, namely the amine reaction product of a compound containing a primary amine functional group and an active ketone or aldehyde containing component, have found increasing use in the domestic treatment of fabrics in 25 order to provide long lasting perfume release on the laundered fabric. Disclosure of such compounds can be found in recently filed applications EP 98870227.0, EP 98870226.2, EP 99870026.4, and EP 99870025.6, all incorporated herein by reference.

Still, the above citations are limited to deposit only one or two type of perfume 30 ingredients on the treated surfaces, whereas there is a need for a deposition of a more

complete perfume formulation so that the various "aspects" of a perfume scent are represented, thereby increasing the consumer's acceptance.

Further, there is also a need for a process for making such composition that is
5 economical and simple.

It has now been found that a laundry and/or cleaning composition which incorporates a benefit agent like a perfume composition with a carrier, wherein the carried composition has a viscosity of at least 400 cps, preferably 1.500 cps, more preferably
10 10.000 at 20°C fulfills such a need.

Perfume which is combined with polymeric component is known in the art. Hence, JP-56075159 discloses the combination of methacrylonitrilebutadiene-styrene tertiary polymer with a liquid perfume so as to yield a semi-solid visco-elastic material for use in the adhesive industry. GB2141726 discloses perfumes which are mixed with adhesives glues for use in the adhesive industry to mask the odor of the adhesive. Finally, DE 3247709 discloses perfumed adhesive cardboard for paper package by using a polymer with a viscosity of 800 to 2500mPa.s.

20 Perfume which is combined with solid carrier in laundry composition is also known in the art. Hence, WO 97/34982 uses zeolites particles as solid carrier, WO 94/19449 uses starch, whilst WO 98/28398 uses organic polymers.

25 Surprisingly, it has been found that when the combination of a benefit agent (e.g. perfume) with a carrier (e.g. polymer) is incorporated in a laundry and/or cleaning and/or fabric care product, the perfume composition is sufficiently protected from the wash oxidative solution and effectively deposited on the fabric whilst still providing efficient release of the perfume on the fabric, in particular dry fabric.

Summary of the invention

5 The present invention is a laundry and/or cleaning and/or fabric care composition comprising a detergent and/or cleaning and/or surfactant and/or fabric care ingredient and a benefit agent, said benefit agent being carried with a carrier, characterised in that the carried benefit agent has a viscosity of at least 400 cps at 20°C.

10 In another aspect of the invention, there is provided a process for the perfume composition.

15 Still in a further aspect of the invention, there is provided a method for providing an enhanced deposition of the benefit agent treated surfaces which comprises the steps of contacting the surface with a composition of the invention, or carried benefit agent as defined herein.

Detailed description of the invention

20

Benefit Agent

The benefit agent is a component that will provide a beneficial effect on the treated surface like fabric. Hence, the benefit agent may be selected from a flavour ingredient, 25 a pharmaceutical ingredient, a biocontrol ingredient, a perfume composition, a refreshing cooling ingredient and mixtures thereof.

Of course, various other features like the one you may want to deposit on the surface may be incorporated in this system, i.e. fabric softener, photobleaching agent,

brightener, bleaching agents, enzymes, lubricants, bleach quenchers, anti-abrasion agents, crystal growth inhibitors, etc...

Typically, the benefit agent comprises from 0.01 to 25%, more preferably from 0.02 to 5% 10%, and most preferably from 0.05 to 5% by weight of the invention composition.

Flavour ingredients include spices, flavor enhancers that contribute to the overall flavour perception.

10 Pharmaceutical ingredients include drugs.

Biocontrol ingredients include biocides, antimicrobials, bactericides, bacteriostatics, fungicides, algaecides, mildewcides, disinfectants, antiseptics, insecticides, vermicides, plant growth hormones.

15 Typical antimicrobials or antibacterials or bacteriostatics which can be carried by the carrier material include amine oxide surfactants, photo-activated bleaches, chlorhexidine diacetate, glutaraldehyde, cinnamon oil and cinnamaldehyde, citric acid, decanoic acid, lactic acid, maleic acid, nonanoic acid, polybiguanide, propylene glycol, cumene sulfonate, eugenol, thymol, benzalkonium chloride, geraniol, and mixtures thereof. Preferred are compounds which can react with the carrier material.

20 Preferably, the carrier material is a polymer, preferably a polymer which is reacted with another benefit agent, such as for example perfumes described herein, and this polymer or polymer reaction product functions as a carrier for the biocide. Preferred carried compositions for use in fabric care and cleaning compositions have a viscosity 25 of at least 500cps, or even at least 1000cps, or even at least 10,000cps or even more more than 100,000 or even more than 500,000cps, as described hereinafter. Preferred polymers are also described in more detail hereinafter.

Typical insect and/or moth repellants are perfume ingredients, such as citronellal, citral, N, N diethyl meta toluamide, Rotundial, 8-acetoxycarvotanacetone, ethyl-3-[N-butyl-N-acetyl]aminopropionate, allethrin, permethrin and mixtures thereof. Other examples of insect and/or moth repellant for use herein are disclosed in US 4,449,987, 4,693,890, 5 4,696,676, 4,933,371, 5,030,660, 5,196,200, and "Semio Activity of Flavor and Fragrance molecules on various Insect Species", B.D. Mookherjee et al., published in Bioactive Volatile Compounds from Plants, ASC Symposium Series 525, R. Teranishi, R.G. Buttery, and H. Sugisawa, 1993, pp. 35-48.

10 One preferred benefit agent is a perfume composition.

Perfume composition

Perfume compositions are typically comprised of one or a mixture of perfumes ingredients.

15 One typical perfume ingredient is a aldehyde perfume ingredient. Preferably, the perfume aldehyde is selected from adoxal; anisic aldehyde; cymal; ethyl vanillin; florhydral; helional; heliotropin; hydroxycitronellal; koavone; lauric aldehyde; lyral; methyl nonyl acetaldehyde; P. T. bucinal; phenyl acetaldehyde; undecylenic aldehyde; 20 vanillin; 2,6,10-trimethyl-9-undecenal, 3-dodecen-1-al, alpha-n-amyl cinnamic aldehyde, 4-methoxybenzaldehyde, benzaldehyde, 3-(4-tert butylphenyl)-propanal, 2-methyl-3-(para-methoxyphenyl propanal, 2-methyl-4-(2,6,6-trimethyl-2(1)-cyclohexen-1-yl) butanal, 3-phenyl-2-propenal, cis-/trans-3,7-dimethyl-2,6-octadien-1-al, 3,7-dimethyl-6-octen-1-al, [(3,7-dimethyl-6-octenyl)oxy] acetaldehyde, 4- 25 isopropylbenzaldehyde, 1,2,3,4,5,6,7,8-octahydro-8,8-dimethyl-2-naphthaldehyde, 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde, 2-methyl-3-(isopropylphenyl)propanal, 1-decanal; decyl aldehyde, 2,6-dimethyl-5-heptenal, 4-(tricyclo[5.2.1.0(2,6)]-decylidene-8)-butanal, octahydro-4,7-methano-1H-indenecarboxaldehyde, 3-ethoxy-4-hydroxy benzaldehyde, para-ethyl-alpha, alpha-dimethyl hydrocinnamaldehyde, alpha-methyl-30 3,4-(methylenedioxy)-hydrocinnamaldehyde, 3,4-methylenedioxybenzaldehyde, alpha-n-

hexyl cinnamic aldehyde, m-cymene-7-carboxaldehyde, alpha-methyl phenyl acetaldehyde, 7-hydroxy-3,7-dimethyl octanal, Undecenal, 2,4,6-trimethyl-3-cyclohexene-1-carboxaldehyde, 4-(3)(4-methyl-3-pentenyl)-3-cyclohexen-carboxaldehyde, 1-dodecanal, 2,4-dimethyl cyclohexene-3-carboxaldehyde, 4-(4-5 hydroxy-4-methyl pentyl)-3-cyclohexene-1-carboxaldehyde, 7-methoxy-3,7-dimethyloctan-1-al, 2-methyl undecanal, 2-methyl decanal, 1-nonanal, 1-octanal, 2,6,10-trimethyl-5,9-undecadienal, 2-methyl-3-(4-tertbutyl)propanal, dihydrocinnamic aldehyde, 1-methyl-4-(4-methyl-3-pentenyl)-3-cyclohexene-1-carboxaldehyde, 5 or 6 methoxyhexahydro-4,7-methanoindan-1 or 2- carboxaldehyde, 3,7-dimethyloctan-1-al, 10-1-undecanal, 10-undecen-1-al, 4-hydroxy-3-methoxy benzaldehyde, 1-methyl-3-(4-methylpentyl)-3-cyclhexenecarboxaldehyde, 7-hydroxy-3,7-dimethyl-octanal, trans-4-decenal, 2,6-nonadienal, para-tolylacetaldehyde; 4-methylphenylacetaldehyde, 2-methyl-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-butenal, ortho-methoxycinnamic aldehyde, 3,5,6-trimethyl-3-cyclohexene carboxaldehyde, 3,7-dimethyl-2-methylene-6-octenal, 15 phenoxyacetaldehyde, 5,9-dimethyl-4,8-decadienal, peony aldehyde (6,10-dimethyl-3-oxa-5,9-undecadien-1-al), hexahydro-4,7-methanoindan-1-carboxaldehyde, 2-methyl octanal, alpha-methyl-4-(1-methyl ethyl) benzene acetaldehyde, 6,6-dimethyl-2-norpinene-2-propionaldehyde, para methyl phenoxy acetaldehyde, 2-methyl-3-phenyl-2-propen-1-al, 3,5,5-trimethyl hexanal, Hexahydro-8,8-dimethyl-2-naphthaldehyde, 3-20 propyl-bicyclo[2.2.1]-hept-5-ene-2-carbaldehyde, 9-decenal, 3-methyl-5-phenyl-1-pentanal, methylnonyl acetaldehyde, 1-p-menthene-q-carboxaldehyde, citral, lilial, florhydral, mefloral, and mixtures thereof.

More preferred aldehydes are selected from citral, 1-decanal, benzaldehyde, florhydral, 25 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde; cis/trans-3,7-dimethyl-2,6-octadien-1-al; heliotropin; 2,4,6-trimethyl-3-cyclohexene-1-carboxaldehyde; 2,6-nonadienal; alpha-n-amyl cinnamic aldehyde, alpha-n-hexyl cinnamic aldehyde, P.T. Bucinal, lyral, cymal, methyl nonyl acetaldehyde, trans-2-nonenal, lilial, trans-2-nonenal, lauric aldehyde, undecylenic aldehyde, mefloral and mixture thereof.

Another typical perfume ingredient is a ketone perfume ingredient. Preferably, the perfume ketone is selected from bucoxime; iso jasmone; methyl beta naphthyl ketone; musk indanone; tonalid/musk plus; Alpha-Damascone, Beta-Damascone, Delta-Damascone, Iso-Damascone, Damascenone, Damarose, Methyl-Dihydrojasmonate, 5 Menthone, Carvone, Camphor, Fenchone, Alpha-Ionone, Beta-Ionone, Gamma-Methyl so-called Ionone, Fleuramone, Dihydrojasmone, Cis-Jasmone, Iso-E-Super, Methyl-Cedrenyl-ketone or Methyl- Cedrylone, Acetophenone, Methyl-Acetophenone, Para-Methoxy-Acetophenone, Methyl-Beta-Naphtyl-Ketone, Benzyl-Acetone, Benzophenone, Para-Hydroxy-Phenyl-Butanone, Celery Ketone or Livescone, 6-Isopropyldecahydro-2-naphthone, Dimethyl-Octenone, Freskomenthe, 4-(1-Ethoxyvinyl)-3,3,5,5,-tetramethyl-Cyclohexanone, Methyl-Heptenone, 2-(2-(4-Methyl-3-cyclohexen-1-yl)propyl)-cyclopentanone, 1-(p-Menthene-6(2)-yl)-1-propanone, 4-(4-Hydroxy-3-methoxyphenyl)-2-butanone, 2-Acetyl-3,3-Dimethyl-Norbornane, 6,7-Dihydro-1,1,2,3,3-Pentamethyl-4(5H)-Indanone, 4-Damascol, Dulcinal or Cassione, Gelsone, Hexalon, Isocyclemon E, Methyl Cyclocitron, Methyl-Lavender-Ketone, Orivon, Para-tertiary-Butyl-Cyclohexanone, Verdone, Delphone, Muscone, Neobutenone, Plicatone, Veloutone, 2,4,4,7-Tetramethyl-oct-6-en-3-one, Tetrameran, hedione, and mixtures thereof. The number of different perfume raw materials in the mixtures can be higher than 5, higher than 10 and even higher than 20.

20

More preferably, for the above mentioned compounds, the preferred ketones are selected from Alpha Damascone, Delta Damascone, Iso Damascone, Carvone, Gamma-Methyl-Ionone, Iso-E-Super, 2,4,4,7-Tetramethyl-oct-6-en-3-one, Benzyl Acetone, Beta Damascone, Damascenone, methyl dihydrojasmonate, methyl cedrylone, hedione, and mixtures thereof.

25 Still, the perfume composition may also be mixture of perfume ingredients including or not the above mentioned aldehyde or ketone.

Typical of these ingredients include fragrant substance or mixture of substances including natural (i.e., obtained by extraction of flowers, herbs, leaves, roots, barks, wood, blossoms or plants), artificial (i.e., a mixture of different nature oils or oil constituents) and synthetic (i.e., synthetically produced) odoriferous substances. Such 5 materials are often accompanied by auxiliary materials, such as fixatives, extenders, stabilizers and solvents. These auxiliaries are also included within the meaning of "perfume", as used herein. Typically, perfumes are complex mixtures of a plurality of organic compounds.

10 Suitable perfumes are disclosed in U.S. Pat. 5,500,138, said patent being incorporated herein by reference.

Examples of perfume ingredients useful in the perfume compositions include, but are not limited to, amyl salicylate; hexyl salicylate; terpineol; 3,7-dimethyl-*cis*-2,6-octadien-1-ol; 2,6-dimethyl-2-octanol; 2,6-dimethyl-7-octen-2-ol; 3,7-dimethyl-3-octanol; 3,7-dimethyl-*trans*-2,6-octadien-1-ol; 3,7-dimethyl-6-octen-1-ol; 3,7-dimethyl-1-octanol; 2-methyl-3-(para-tert-butylphenyl)-propionaldehyde; 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde; tricyclodeceny1 propionate; tricyclodeceny1 acetate; anisaldehyde; 2-methyl-2-(para-iso-propylphenyl)-propionaldehyde; ethyl-3-methyl-3-phenyl glycidate; 4-(para-hydroxyphenyl)-butan-2-one; 1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-2-buten-1-one; para-methoxyacetophenone; para-methoxy-alpha-phenylpropene; methyl-2-n-hexyl-3-oxo-cyclopentane carboxylate; undecalactone gamma.

Additional examples of fragrance materials include, but are not limited to, orange oil; 25 lemon oil; grapefruit oil; bergamot oil; clove oil; dodecalactone gamma; methyl-2-(2-pentyl-3-oxo-cyclopentyl) acetate; beta-naphthol methylether; methyl-beta-naphthylketone; coumarin; 4-tert-butylcyclohexyl acetate; alpha, alpha-dimethylphenethyl acetate; methylphenylcarbonyl acetate; cyclic ethyleneglycol diester of tridecanoic acid; 3,7-dimethyl-2,6-octadiene-1-nitrile; ionone gamma methyl; 30 ionone alpha; ionone beta; petitgrain; methyl cedrylone; 7-acetyl-1,2,3,4,5,6,7,8-

octahydro-1,1,6,7-tetramethyl-naphthalene; ionone methyl; methyl-1,6,10-trimethyl-2,5,9-cyclododecatrien-1-yl ketone; 7-acetyl-1,1,3,4,4,6-hexamethyl tetralin; 4-acetyl-6-tert-butyl-1,1-dimethyl indane; benzophenone; 6-acetyl-1,1,2,3,3,5-hexamethyl indane; 5-acetyl-3-isopropyl-1,1,2,6-tetramethyl indane; 1-dodecanal; 7-hydroxy-3,7-dimethyl 5 octanal; 10-undecen-1-al; iso-hexenyl cyclohexyl carboxaldehyde; formyl tricyclodecan; cyclopentadecanolide; 16-hydroxy-9-hexadecenoic acid lactone; 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta-gamma-2-benzopyrane; ambroxane; dodecahydro-3a,6,6,9a-tetramethylnaphtho-[2,1b]furan; cedrol; 5-(2,2,3-trimethylcyclopent-3-enyl)-3-methylpentan-2-ol; 2-ethyl-4-(2,2,3-trimethyl-3-cyclopenten-1-yl)-2-buten-1-ol; 10 caryophyllene alcohol; cedryl acetate; para-tert-butylcyclohexyl acetate; patchouli; olibanum resinoid; labdanum; vetivert; copaiba balsam; fir balsam; hydroxycitronellal and indol; phenyl acetaldehyde and indol;

More examples of perfume components are geraniol; geranyl acetate; linalool; linalyl acetate; tetrahydrolinalool; citronellol; citronellyl acetate; dihydromyrcenol; dihydromyrcenyl acetate; tetrahydromyrcenol; terpinyl acetate; nopol; nopyl acetate; 2-phenylethanol; 2-phenylethyl acetate; benzyl alcohol; benzyl acetate; benzyl salicylate; benzyl benzoate; styrallyl acetate; dimethylbenzylcarbinol; trichloromethylphenylcarbonyl methylphenylcarbonyl acetate; isononyl acetate; vetiveryl acetate; vetiverol; 2-methyl-3-(p-tert-butylphenyl)-propanal; 2-methyl-3-(p-isopropylphenyl)-propanal; 3-(p-tert-butylphenyl)-propanal; 4-(4-methyl-3-pentenyl)-3-cyclohexenecarbaldehyde; 4-acetoxy-3-pentyltetrahydropyran; methyl dihydrojasmonate; 2-n-heptylcyclopentanone; 3-methyl-2-pentyl-cyclopentanone; n-decanal; n-dodecanal; 9-decenol-1; phenoxyethyl isobutyrate; phenylacetaldehyde 25 dimethylacetal; phenylacetaldehyde diethylacetal; geranonitrile; citronellonitrile; cedryl acetal; 3-isocamphylcyclohexanol; cedryl methylether; isolongifolanone; aubepine nitrile; aubepine; heliotropine; eugenol; vanillin; diphenyl oxide; hydroxycitronellal ionones; methyl ionones; isomethyl ionomes; irones; cis-3-hexenol and esters thereof; indane musk fragrances; tetralin musk fragrances; isochroman musk fragrances; 30 macrocyclic ketones; macrolactone musk fragrances; ethylene brassylate. Also suitable

herein as perfume ingredients of the perfume composition are the so-called Schiff base. Schiff-Bases are the condensation of an aldehyde perfume ingredient with an anthranilate. A typical description can be found in US 4853369. Typical of Schiff bases are selected from Schiff's base of 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde and methyl anthranilate; condensation products of: hydroxycitronellal and methyl anthranilate; 4-(4-hydroxy-4-methyl pentyl)-3-cyclohexene-1-carboxaldehyde and methyl anthranilate; Methyl Anthranilate and HydroxyCitronellal commercially available under the tradename Aurantiol; Methyl Anthranilate and Methyl Nonyl Acetaldehyde commercially available under the tradename Agrumea; Methyl Anthranilate and PT Bucinal commercially available under the tradename Verdantiol; Methyl anthranilate and Lyral commercially available under the tradename Lyrame; Methyl Anthranilate and Ligustral commercially available under the tradename Ligantral; and mixtures thereof.

Preferably, the perfume compositions useful in the present invention compositions are substantially free of halogenated materials and nitromusks.

More preferably, the perfume compounds are characterised by having a low Odor Detection Threshold. Such Odor Detection Threshold (ODT) should be lower than 1ppm, preferably lower than 10ppb - measured at controlled Gas Chromatography (GC) conditions such as described here below. This parameter refers to the value commonly used in the perfumery arts and which is the lowest concentration at which significant detection takes place that some odorous material is present. Please refer for example in "Compilation of Odor and Taste Threshold Value Data (ASTM DS 48 A)", edited by F. A. Fazzalari, International Business Machines, Hopwell Junction, NY and in Calkin et al., Perfumery, Practice and Principles, John Wiley & Sons, Inc., page 243 et seq (1994). For the purpose of the present invention, the Odor Detection Threshold is measured according to the following method :

The gas chromatograph is characterized to determine the exact volume of material injected by the syringe, the precise split ratio, and the hydrocarbon response using a

hydrocarbon standard of known concentration and chain-length distribution. The air flow rate is accurately measured and, assuming the duration of a human inhalation to last 0.02 minutes, the sampled volume is calculated. Since the precise concentration at the detector at any point in time is known, the mass per volume inhaled is known and hence the concentration of material. To determine the ODT of a perfume material, 5 solutions are delivered to the sniff port at the back-calculated concentration. A panelist sniffs the GC effluent and identifies the retention time when odor is noticed. The average over all panelists determines the threshold of noticeability. The necessary amount of analyte is injected onto the column to achieve a certain concentration, such as 10 ppb, at the detector. Typical gas chromatograph parameters for determining odor 10 detection thresholds are listed below.

GC: 5890 Series II with FID detector

7673 Autosampler

Column: J&W Scientific DB-1

15 Length 30 meters ID 0.25 mm film thickness 1 micrometer

Method:

Split Injection: 17/1 split ratio

Autosampler: 1.13 microliters per injection

Column Flow: 1.10 mL/minute

20 Air Flow: 345 mL/minute

Inlet Temp. 245°C

Detector Temp. 285°C

Temperature Information

Initial Temperature: 50°C

25 Rate: 5C/minute

Final Temperature: 280°C

Final Time: 6 minutes

Leading assumptions: 0.02 minutes per sniff

GC air adds to sample dilution

Examples of such preferred perfume components are those selected from : 2-methyl-2-(para-iso-propylphenyl)-propionaldehyde, 1-(2,6,6-trimethyl-2-cyclohexan-1-yl)-2-buten-1-one and/or para-methoxy-acetophenone. Even more preferred are the following compounds having an ODT \leq 10ppb measured with the method described above :

5 undecylenic aldehyde, undecalactone gamma, heliotropin, dodecalactone gamma, p-anisic aldehyde, para hydroxy-phenyl-butanone, cymal, benzyl acetone, ionone alpha, p.t.bucinal, damascenone, ionone beta, methyl-nonyl ketone, methyl heptine carbonate, linalool, indol, cis-3-hexenyl salicylate, vanillin, methyl isobutetyl tetrahydropyran, ethylvanillin, coumarin, ethyl methyl phenyl glycidate, eugenol, methylanthranilate, iso 10 eugenol, beta naphtol methyl ester, herbavert, lyral, allyl amyl glycolate, dihydro iso jasmonate, ethyl-2-methylbutyrate, nerol, and phenylacetaldehyde. Most preferably the perfume composition comprises at least 5%, more preferably at least 10% of such components

15 Most preferably, the perfume ingredients are those as described in WO 96/12785 on page 12-14. Even most preferred are those perfume compositions comprising at least 10%, preferably 25%, by weight of perfume ingredient with an ClogP of at least 2.0, preferably at least 3.0, and boiling point of at least 250°C. still another preferred perfume composition is a composition comprising at least 20%, preferably 35%, by 20 weight of perfume ingredient with an ClogP at least 2.0, preferably at least 3.0, and boiling point of less than or equal to 250°C.

Clog P is a commonly known calculated measure as defined in the following references “Calculating log P_{oct} from Structures”; Albert Leo (Medicinal Chemistry Project, 25 Pomona College, Claremont, CA USA. Chemical Reviews, Vol. 93, number 4, June 1993; as well as from Comprehensive Medicinal Chemistry, Albert Leo, C. Hansch, Ed. Pergamon Press: Oxford, 1990, Vol. 4, p.315; and Calculation Procedures for molecular lipophilicity: a comparative Study, Quant. Struct. Act. Realt. 15, 403-409 (1996), Raymund Mannhold and Karl Dross.

Carrier

A carrier is another essential component of the invention. Indeed, the carrier will serve for the deposition of the benefit agent onto the surface as well as protecting the benefit agent from oxidation from the wash liquor as well as from diffusing in the aqueous environment.

Preferably, for the purpose of the invention, the carrier or even the carried composition is water-insoluble, preferably the carrier is a water-insoluble polymer. Carrier to be used herein are selected from polymers which have chemically reacted with a benefit agent like perfume ingredient, components which have chemically reacted with a benefit agent like perfume ingredient to make the carrier as above mentioned, polymers which are not capable of chemically reacting with a benefit agent like a perfume ingredient above mentioned, i.e. chemically inert, and mixtures thereof.

These carrier components are selected so as to provide the required viscosity of at least 400 cps for the resulting carried composition. Preferably, these components will also provide the water-insolubility of the carried composition.

In the present invention, if using polymers as the carrier and aldehyde mixtures as the benefit agent, it is possible to adjust the ratio of the carrier and the benefit agent. If the amount of polymers are low, some aldehydes remain unreacted. In this case, these unreacted-aldehydes can also perform as perfume in the final product in the present invention.

a)- Compounds which have chemically reacted with a benefit agent

Examples of compounds which have chemically reacted with a benefit agent are the so-called "amines which form amine reaction products", i.e. a product of reaction between a compound containing a primary amine functional group and/or secondary amine functional group and an active ketone or aldehyde containing component. Preferred compounds for use herein are polymers which have been previously reacted with an

aldehyde and/or ketone perfume ingredient, thereby imparting a more effective scent to the fabrics.

5 A typical disclosure of amine reaction product suitable for use herein can be found in recently filed applications EP 98870227.0, EP 98870226.2, EP 99870026.4, and EP 99870025.6, all incorporated herein by reference.

A-Primary amine and/or secondary amine

10 By "primary and/or secondary amine", it is meant a component which carries at least one primary and/or secondary amine and/or amide function.

Of course, one amine compound may carry both primary and secondary amine compound, thereby enabling the reaction with several aldehydes and/or ketones.

15 Preferably, the primary amine and/or secondary amine compound is also characterized by an Odour Intensity Index of less than that of a 1% solution of methylanthranilate in dipropylene glycol.

Odour Intensity Index method

20 By Odour Intensity Index, it meant that the pure chemicals were diluted at 1% in Dipropylene Glycol, odor-free solvent used in perfumery. This percentage is more representative of usage levels. Smelling strips, or so called "blotters", were dipped and presented to the expert panellist for evaluation. Expert panellists are assessors trained for at least six months in odor grading and whose gradings are checked for accuracy and reproducibility versus a reference on an on-going basis. For each amine compound, the panellist was presented two blotters: one reference (Me Anthranilate, unknown from the panellist) and the sample. The panellist was asked to rank both smelling strips on the 0-5 odor intensity scale, 0 being no odor detected, 5 being very strong odor present.

Results:

The following represents Odour Intensity Index of an amine compound suitable for use in the present invention and according to the above procedure. In each case, numbers 5 are arithmetic averages among 5 expert panellists and the results are statistically significantly different at 95% confidence level:

Methylantranilate 1% (reference)	3.4
Ethyl-4-aminobenzoate (EAB) 1%	0.9
10 1,4-bis-(3-aminopropyl)-piperazine (BNPP) 1%	1.0

A general structure for the primary amine compound of the invention is as follows:

15 B-(NH₂)_n;
wherein B is a carrier material, and n is an index of value of at least 1.

Compounds containing a secondary amine group have a structure similar to the above excepted that the compound comprises one or more -NH- groups instead of -NH₂.
20 Further, the compound structure may also have one or more of both -NH₂ and -NH- groups.

Preferred B carriers are inorganic or organic carriers.

25 By “inorganic carrier”, it is meant carrier which are non-or substantially non carbon based backbones.

Preferred primary and/or secondary amines, among the inorganic carriers, are those selected from mono or polymers or organic-organosilicon copolymers of amino 30 derivatised organo silane, siloxane, silazane, alumane, aluminum siloxane, or aluminum

silicate compounds. Typical examples of such carriers are: organosiloxanes with at least one primary amine moiety like the diaminoalkylsiloxane [H2NCH2(CH3) 2Si]O, or the organoaminosilane (C6H5) 3SiNH2 described in: Chemistry and Technology of Silicone, W. Noll, Academic Press Inc. 1998, London, pp 209, 106).

5

Preferred primary and/or secondary amines, among the organic carriers, are those selected from aminoaryl derivatives, polyamines, amino acids and derivatives thereof, substituted amines and amides, glucamines, dendrimers, polyvinylamines and derivatives thereof, and/or copolymer thereof, alkylene polyamine, polyaminoacid and 10 copolymer thereof, cross-linked polyaminoacids, amino substituted polyvinylalcohol, polyoxyethylene bis amine or bis aminoalkyl, aminoalkyl piperazine and derivatives thereof, bis (amino alkyl) alkyl diamine linear or branched, and mixtures thereof.

Preferred aminoaryl derivatives are the amino-benzene derivatives including the alkyl 15 esters of 4-amino benzoate compounds, and more preferably selected from ethyl-4-amino benzoate, phenylethyl-4-aminobenzoate, phenyl-4-aminobenzoate, 4-amino-N'-(3-aminopropyl)-benzamide, and mixtures thereof.

Polyamines suitable for use in the present invention are polyethyleneimines polymers, 20 poly[oxy(methyl-1,2-ethanediyl)], α -(2-aminomethylethyl)- ω -(2-aminomethyl-ethoxy)- (= C.A.S No. 9046-10-0); poly[oxy(methyl-1,2-ethanediyl)], α -hydro-)- ω -(2-aminomethylethoxy)-, ether with 2-ethyl-2-(hydroxymethyl)-1,3-propanediol (= C.A.S. No. 39423-51-3); commercially available under the tradename Jeffamines T-403, D-230, D-400, D-2000; 2,2',2"-triaminotriethylamine; 2,2'-diamino-diethylamine; 3,3'- 25 diamino-dipropylamine, 1,3 bis aminoethyl-cyclohexane commercially available from Mitsubishi and the C12 Sternamines commercially available from Clariant like the C12 Sternamin(propylenamine)_n with n=3/4, and mixtures thereof. Preferred polyamines are polyethyleneimines commercially available under the tradename Lupasol like Lupasol HF (MW 25000), P (MW 750000), PS (MW 750000), SK (MW 2000000),

SNA (MW 1000000), G20 (MW1300), G35 (MW2000), G100, PR8515 (MW2000), FG (MW800).

Preferred amino acids for use herein are selected from tyrosine, tryptophane, lysine, 5 glutamic acid, glutamine, aspartic acid, arginine, asparagine, phenylalanine, proline, glycine, serine, histidine, threonine, methionine, and mixture thereof, most preferably selected from tyrosine, tryptophane, and mixture thereof. Preferred amino acid derivatives are selected from tyrosine ethylate, glycine methylate, tryptophane ethylate, and mixture thereof.

10

Preferred substituted amines and amides for use herein are selected from nipecotamide, N-coco-1,3-propenediamine; N-oleyl-1,3-propenediamine; N-(tallow alkyl)-1,3-propenediamine; 1,4-diamino cyclohexane; 1,2-diamino-cyclohexane; 1,12-diaminododecane, and mixtures thereof.

15

Other primary amine compounds suitable for use herein are the glucamines, preferably selected from 2,3,4,5,6-pentamethoxy-glucamine; 6-acetylglucamine, glucamine, and mixture thereof.

20

Also preferred compounds are the polyethylenimine and/or polypropylenimine dendrimers and the commercially available Starburst[®] polyamidoamines (PAMAM) dendrimers, generation G0-G10 from Dendritech and the dendrimers Astromols[®], generation 1-5 from DSM being DiAminoButane PolyAmine DAB (PA)x dendrimers with x = 2ⁿ x 4 and n being generally comprised between 0 and 4.

25

Polyamino acid is one suitable and preferred class of amino-functional polymer. Polyaminoacids are compounds which are made up of amino acids or chemically modified amino acids. They can contain alanine, serine, aspartic acid, arginine, valine, threonine, glutamic acid, leucine, cysteine, histidine, lysine, isoleucine, tyrosine, 30 asparagine, methionine, proline, tryptophan, phenylalanine, glutamine, glycine or

mixtures thereof. In chemically modified amino acids, the amine or acidic function of the amino acid has reacted with a chemical reagent. This is often done to protect these chemical amine and acid functions of the amino acid in a subsequent reaction or to give special properties to the amino acids, like improved solubility. Examples of such 5 chemical modifications are benzyloxycarbonyl, aminobutyric acid, butyl ester, pyroglutamic acid. More examples of common modifications of amino acids and small amino acid fragments can be found in the Bachem, 1996, Peptides and Biochemicals Catalog.

10 Preferred polyamino acids are polylysines, polyarginine, polyglutamine, polyasparagine, polyhistidine, polytryptophane or mixtures thereof. Most preferred are polylysines or polyamino acids where more than 50% of the amino acids are lysine, since the primary amine function in the side chain of the lysine is the most reactive amine of all amino acids.

15

The preferred polyamino acid has a molecular weight of 500 to 10.000.000, more preferably between 2.000 and 25.000.

20 The polyamino acid can be cross linked. The cross linking can be obtained for example by condensation of the amine group in the side chain of the amino acid like lysine with the carboxyl function on the amino acid or with protein cross linkers like PEG derivatives. Other examples are described herein below. The cross linked polyamino acids still need to have free primary and/or secondary amino groups left for reaction with the active ingredient.

25

The preferred cross linked polyamino acid has a molecular weight of 20.000 to 10.000.000, more preferably between 200.000 and 2.000.000.

30 The polyamino acid or the amino acid can be co-polymerized with other reagents like for instance with acids, amides, acyl chlorides. More specifically with aminocaproic

acid, adipic acid, ethylhexanoic acid, caprolactam or mixture thereof. The molar ratio used in these copolymers ranges from 1:1 (reagent/ amino acid (lysine)) to 1:20, more preferably from 1:1 to 1:10.

5 The polyamino acid like polylysine can also be partially ethoxylated.

Examples and supply of polyaminoacids containing lysine, arginine, glutamine, asparagine are given in the Bachem 1996, Peptides and Biochemicals catalog.

10 The polyaminoacid can be obtained before reaction with the active ingredient, under a salt form. For example polylysine can be supplied as polylysine hydrobromide. Polylysine hydrobromide is commercially available from Sigma, Applichem, Bachem and Fluka.

15 Examples of suitable amino functional polymers containing at least one primary and/or secondary amine group for the purpose of the present invention are :

- Polyvinylamine with a MW of about 300-2.10E6;
- Polyvinylamine alkoxylated with a MW of about 600, 1200 or 3000 and an ethoxylation degree of 0.5;

20 - Polyvinylamine vinylalcohol - molar ratio 2:1, polyvinylaminevinylformamide - molar ratio 1:2 and polyvinylamine vinylformamide-molar ratio 2:1;

- Triethylenetetramine, diethylenetriamine, tetraethylenepentamine;
- Bis-aminopropylpiperazine;

25 - Polyamino acid (L-lysine / lauric acid in a molar ratio of 10/1), Polyamino acid (L-lysine / aminocaproic acid / adipic acid in a molar ratio of 5/5/1), Polyamino acid (L-lysine / aminocaproic acid /ethylhexanoic acid in a molar ratio of 5/3/1) Polyamino acid (polylysine-cocaprolactam); Polylysine; Polylysine hydrobromide; cross-linked polylysine,

- amino substituted polyvinylalcohol with a MW ranging from 400-300,000;

30 - polyoxyethylene bis [amine] available from e.g. Sigma;

- polyoxyethylene bis [6-aminohexyl] available from e.g. Sigma;
- N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched (TPTA); and
- 1,4-bis-(3-aminopropyl) piperazine (BNPP).

5 The more preferred compounds are selected from ethyl-4-amino benzoate, polyethyleneimine polymers commercially available under the tradename Lupasol like Lupasol WFG20 waterfree, PR8515, HF, P, PS, SK, SNA; the diaminobutane dendrimers Astramol®, polylysine, cross-linked polylysine, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched; 1,4-bis-(3-aminopropyl) piperazine, and mixtures thereof. Even most preferred compounds are those selected from ethyl-4-amino benzoate, polyethyleneimine polymers commercially available under the tradename Lupasol like Lupasol WF, G20 waterfree, PR8515, HF, P, PS, SK, SNA; polylysine, cross-linked polylysine, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched, 1,4-bis-(3-aminopropyl) piperazine, and mixtures thereof.

15

Advantageously, such most preferred primary and/or secondary amine compounds also provide fabric appearance benefit, in particular colour appearance benefit, thus providing a resulting amine reaction product with the properties of fabric appearance benefit, deposition onto the surface to be treated, and delayed release of the active as well as release of the perfume composition. Further, when the primary and/or secondary amine compound has more than one free primary and/or secondary amine group, several different active ingredients (aldehyde and/or ketone) can be linked to the amine compound.

25 Of course, the primary and/or secondary amine compound may also be used as is, i.e. without having been reacted with the above benefit agent like aldehyde and/or ketone perfume ingredient. Moreover, the primary and/or secondary amine compound may also be reacted with compounds other than the benefit agent mentioned above like acyl halides, like acetylchloride, palmytoyl chloride or myristoyl chloride, acid anhydrides like acetic anhydride, alkylhalides or arylhalides to do alkylation or arylation, aldehydes

or ketones not used as perfume ingredients like formaldehyde, glutaraldehyde, unsaturated ketones, aldehydes or carboxylic acids like 2-decylpropenoic acid, propenal, propenone to form reaction products with the required viscosity.

5 The carrier mix can be treated (during the mix is formed or after) with plasticisers like phtalates, with tactifiers like rosin acids or rosin esters, cross-linking agents like bifunctional aldehydes, or with thickeners, as described herein after. These agents can give the polymer the proper carrier characteristics like the required viscosity if the viscosity is not high enough. Of course, other known viscosity enhancer may be used
10 herein for that purpose. Preferably the ratio of carried benefit agent to thickening and/or cross-linking agent being from 100:1 to 10:1.

b)- Polymers which are not capable of chemically reacting with a benefit agent

Polymers which are not capable of chemically reacting with a benefit agent include
15 block copolymers like block copolymer of styrene and butadiene, polyisoprene, polyacrylate, acrylic emulsion polymers using preferably ethylacrylate butyl acrylate, 2-ethylhexylacrylate, methylacrylate, acrylic acid, methacrylic acid as monomers, acrylic emulsion polymers copolymerized with vinyl acetate, vinyl chloride or maleic acid, styrene polymers, polyurethane, polybutadiene, polyepichlorohydrin, neoprene or
20 chloroprene, natural latex rubbers, polyvinylpyrrolidine, polyvinylpyridine N oxide, vinylpyrrolodone vinyl imidazole copolymer, chlorosulfonyl polyethylene, ethylene propylene copolymer, ethylene polysulfide, polyvinylacetate, polyamide, polyvinylacetate-ethylene copolymers, urea-formaldehyde resins, cyanoacrylates, polysulphides, polyvinylalcohol, styrene-butadiene polymers, polyolefines based on
25 polyethylene or polypropylene, polyester, nitrile rubber polymers based on butadiene and acrylonitrile, as well as silicone rubbers having methyl, phenyl and vinyl groups or mixtures thereof or copolymers (random, block or grafted) of the above mentioned polymers or the above polymers further cross linked with cross linking agents like zinc oxide.

The polymers can be treated with plasticisers like phtalates, with tactifiers like rosin acids or rosin esters, or with thickeners. These agents can give the polymer the proper carrier characteristics like the required viscosity.

5 . Preferred polymers from this class are polymers used in the adhesive industry, more preferably polyisobutylene polymers supplied by BASF under the commercial name of Oppanol.

10 It is most preferred that the benefit agent and the carrier are present in weight ratios of from 0.05:1 to 5:1, preferably of from 1:1 to 4:1. Indeed, not to be bound by theory, it is speculated that below a ratio of 0.05:1 or even 0.5:1, the amount of polymer that would be required to form the carried composition would be too high while above a ratio of 5:1, the system would be too liquid and therefore not provide its purpose of deposition onto the treated surface.

15

Viscosity

20 Viscosity of the carried perfume composition, i.e. the perfume composition which is carried by the carrier material, is an essential feature of the invention. Indeed, with the viscosity characteristic, the perfume composition is ensured to be protected from its oxidative environment present in the wash liquor, effectively deposited on the surface to be treated and thereafter to deliver its release on the treated surface.

25 To achieve these benefits, the viscosity of the carried composition is between 400 cps, preferably between 1.500 cps and 100.000.000 cps, preferably between 5.000 and 10.000.000 cps, more preferably between 10.000 and 1.000.000 cps, most preferably between 10.000 and 100.000

The viscosity is measured on a rheometer, TA Instrument CSL²₁₀₀ at a temperature of 20C with a gap setting of 500 micrometers.

30

Process

The carried composition comprising the benefit agent is obtained by mixing the benefit agent with the carrier in such a way that a very viscous homogeneous fluid is obtained with the desired viscosity.

5

One convenient way for making the carried composition in industrial quantities is via a continuous process like by means of a twin Screw Extruder (TSE). Suitable TSE include the TX-57 MAG, TX-85 MAG, TX-110 MAG, TX-144 MAG, or TX-178 MAG twin screw extruder from Wenger. One preferred for use herein is the TX-57 MAG. TSE suitable for use herein comprise at one of their extremities so called herein after "first part of the TSE" two distinct inlet: one for the active and the other for the amine, and at about the middle of the TSE, so called hereinafter "second part of the TSE" another inlet for the carrier. Temperature controllers are also distributed along the TSE.

15 Preferred is that the carried composition is made into suspendable particels or solid particles by dispersing it into a carrier dispersing agent (or below referred to as 'carrier'), preferably a liquid carrier, which is preferably a material which is solid at room temperature, e.g. below 25°C or even below 30°C, and is liquid due to the temperature of the equipment wherein the mixing takes place and/ or the temperature of the product or mixture of step a). Thus, the carrier material has preferably a melting point above 30°C. Preferably, the temperature of the product of step a) and/or the carrier material is such that the carrier material is in its molten state, preferably the temperature of the carrier material and/ or the reaction product/ mixture of step a) is between 30°C and 100°C, preferably between 40°C and 80°C or even between 50°C
20 and 80°C. Preferably, for the purpose of the invention, when the resulting carried composition is to be a suspendable material, the carrier also has a viscosity from 500 or even from 700 to 100,000 or even 70,000cps.
25

30 Highly preferred are carrier materials which do not react with the carried composition of the invention.

Highly preferred are organic nonionic material, including nonionic surfactants.

Preferred carrier material include liquids conventionally used in cleaning products as solvents, such as alcohols, glycerols.

5

Preferred are nonionic surfactants. Essentially any nonionic surfactants useful for detergents purposes can be included in the compositions provided it has a melting point between 30°C and 135°C.

10 Exemplary, non-limiting classes of useful nonionic surfactants are:

Polyhydroxy fatty acid amides suitable for use herein are those having the structural formula R₂CONR₁Z wherein : R₁ is H, C₁-C₄ hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl, or a mixture thereof, preferable C₁-C₄ alkyl, more preferably C₁ or C₂ alkyl, most preferably C₁ alkyl (i.e., methyl); and R₂ is a C₅-C₃₁ hydrocarbyl, preferably straight-chain C₅-C₁₉ alkyl or alkenyl, more preferably straight-chain C₉-C₁₇ alkyl or alkenyl, most preferably straight-chain C₁₁-C₁₇ alkyl or alkenyl, or mixture thereof; and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably will be derived from a reducing sugar in a reductive amination reaction; more preferably Z is a glycetyl.

25 The alkyl ethoxylate condensation products of aliphatic alcohols with from about 1 to about 150 moles of ethylene oxide are suitable for use herein. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 6 to 22 carbon atoms. Particularly preferred are the condensation products of alcohols having an alkyl group containing from 8 to 20 carbon atoms with from about 25 to about 150 moles of ethylene oxide per mole of alcohol, preferably 50 to 100, more preferably 80 moles of ethylene oxide per mole of alcohol.

Preferred nonionic ethoxylated alcohol surfactants are selected from tallow (C₁₆-C₁₈) alcohol ethoxylated with 25, 50, 80, or 100 moles of ethylene oxide commercially available from under the tradename of Lutensol from BASF, Empilan from Albright and Wilson, and Genapol from Clariant. The most preferred nonionic ethoxylated alcohol surfactant is tallow (C₁₆-C₁₈) alcohol ethoxylated with 80moles of ethylene oxide and commercially available under the tradename of Lutensol 80/80 from BASF, Empilan KM 80 from Albright and Wilson, or Genapol T800 from Clariant.


The ethoxylated C₆-C₂₂ fatty alcohols and C₆-C₂₂ mixed ethoxylated/propoxylated fatty alcohols are suitable surfactants for use herein, particularly where water soluble. Preferably the ethoxylated fatty alcohols are the C₁₀-C₂₂ ethoxylated fatty alcohols with a degree of ethoxylation of from 25 to 150, most preferably these are the C₁₂-C₁₈ ethoxylated fatty alcohols with a degree of ethoxylation from 50 to 80. Preferably the mixed ethoxylated/propoxylated fatty alcohols have an alkyl chain length of from 10 to 18 carbon atoms, a degree of ethoxylation of from 3 to 30 and a degree of propoxylation of from 1 to 30.

The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol are suitable for use herein. The hydrophobic portion of these compounds preferably has a molecular weight of from about 1500 to about 1800 and exhibits water insolubility. Examples of compounds of this type include certain of the commercially-available PluronicTM surfactants, marketed by BASF.

The condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine are suitable for use herein. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. Examples of this type of nonionic surfactant include certain of the commercially available TetronicTM compounds, marketed by BASF.

Suitable alkylpolysaccharides for use herein are disclosed in U.S. Patent 4,565,647, Llenado, issued January 21, 1986, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g., a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties. (Optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside.) The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units.

15 The preferred alkylpolyglycosides have the formula



wherein R₂ is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from 10 to 18, preferably from 12 to 14, carbon atoms; n is 2 or 3; t is from 0 to 10, preferably 0, and X is from 1.3 to 8, preferably from 1.3 to 3, most preferably from 1.3 to 2.7. The glycosyl is preferably derived from glucose.

25

Fatty acid amide surfactants suitable for use herein are those having the formula:

$R^6CON(R^7)_2$ wherein R⁶ is an alkyl group containing from 7 to 21, preferably from 9 to 17 carbon atoms and each R⁷ is selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, and -(C₂H₄O)_xH, where x is in the range of from 1 to 30 3.

Preferred carrier materials are selected from nonionic ethoxylated alcohol surfactants, polyalkoxylated compounds, such as polyalkoxylated esters, polyalkoxylated amines, polyalkoxylated amides, polyalkoxylated alcohols, preferably poly ethoxylated
5 compounds. Preferred average alkoxylation degrees are at least 25, or even at least 40 or even at least 70. Also useful herein as carrier are quaternary oligoamine oligomers, preferably alkoxylated quaternary oligoamines, more preferable polyethoxylated quaternary diamines, preferably ahveing an alkoxylation degree of 10 to 40, or even 16 to 26, preferably the quaternary amine groups being spaced apart by 2 carbon atoms or
10 more, preferably by 4 carbon atoms or more, pr even at 6 or more carbon atoms, preferably this being alkylene moieties.

Highly preferred carrier materials include polyalkoxylated alcohols. Tallow alcohol polyethoxylates, such as TAE80, and PEG and cationically modified PEG are preferred
15 examples. Also preferred are polyethylene glycols, preferably with an weight average molecular weight of more than 400, preferably more than 1000 or even mor than 2000 or even more than 3000, for example PEG 4000, preferably up to 10,000.

Also useful as carrier can be anionic surfactants, cationic surfactants, amphoteric
20 surfactants, zwitterionic surfactants and/ or amphiphilic surfactants.

Also useful are hydrotropes, such as salts (sodium) of xylenen sulphonate, toluene sulphonate and/ or cumene sulphonate.

25 Other preferred carriers include silicone materials. Preferred are non-volatile silicone fluids such as polydimethyl siloxane gums and fluids, such as linear silicone polymer fluids having the formula $(CH_3)_3SiO[(CH_3)_2SiO]_mSi(CH_3)_3$ where m is 0 or more and whereby m has an average value such that the viscosity at 25°C of the silicone fluid is preferably 5 centistokes or more, more preferably 500 centistokes or more, the silicone
30 fluid preferably having a weight average molecular weight of 800 or more, preferably

25,000 or more; or such as volatile silicone fluid which can be a cyclic silicone fluid of the formula $[(CH_3)_2SiO]n$ where n ranges between about 3 to about 7, preferably about 5 or 6; or such as silicone surfactants, such as silicone-glycolethers; other suitable silicone surfactants are described in 'Silicone Surfactants', by R.M.Hill, ISBN 0-8247-5 0010-4, 1999, Marcel Dekker Inc. New York, Basel. These silicone surfactants can be ABA type copolymers, grafted copolymers or ter- or tri-siloxane polymers. The silicone surfactants can be silicone polyether copolymers and can have ethylene oxide, propylene oxide or butylene oxide based chains and/or mixtures thereof. More preferably the silicone surfactant has a weight average molecular weight of more than 1000, more preferably more than 5000. The silicone or silicone surfactants can be a fluorosilicone as well, preferably a fluorosilicone with a viscosity of at least 1000 cps.

10 Suitable examples of silicone for use herein as carrier include silicones commercially available from Dow Corning Corporation like the DC 3225C; DC5225C and DC246 for cyclic silicones; silicone glycols ethers like DC5200, DC1248, DC190; the DC 244 Fluids, DC 245 Fluids, DC 344 Fluids, or DC 345 Fluids, or ABIL K4, ABIL B 8839 for the cyclomethicone, or the DC 200 fluids, ABIL K 520 (hexamethyl disiloxane), ABIL 10 to ABIL 100000 (dimethicone), ABIL AV 8853 (Phenyl dimethicone) for the linear silicones; Dow Corning's FS1265 fluorosilicones.

20

Still another suitable carrier is a combination of various silicone materials and/ or other carriers, such as those described herein before. Preferably, one or more silicone material(s) is (are) emulsified or microemulsified in one or more other silicone materials. Preferably, at least 80% or even 90% of such a silicone mixture is formed by 2 silicone materials. Then, the weight ratio of a first silicone material to a second silicone material in such a mixture is preferably from 1:50 to 2:1, more preferably 1:19 to 3:2, or even 1:9 to 1:1. Preferably, the carrier for suspendable components formed by the process herein is selected from glycols and/ or silicones, described above, most preferably is selected from silicones. Preferably, the carrier for solid components formed by the process herein is selected from nonionic surfactants, alkoxylated

compounds, including alkoxylated alcohol nonionic surfactants, alcohols, glycols and/or (polyalkylene) glycols.

When solid compositions are preferred, a solid granulation agent is preferably added to the carried composition or the carried composition which is mixed with the carriers described above. The solid granulation aid can be any material which is solid under the reaction conditions, other than the compounds reacted with another. Preferred are inorganic or organic acids or salts. The granulation agent should be such that it does not react with the reaction product of step a). Preferred are anhydrous materials.

10

Highly preferred are solid granulation agents in powder form having a weight mean particle size of from 1 to 200 microns, preferably up to 150 microns or even up to 100 microns.

15

Preferred are organic carboxylic acid or salts thereof, conventional chelating agents, including phosphonate chelating agents are suitable herein, or preferred are inorganic materials such as inorganic salts, including bicarbonates, carbonates, sulphates, phosphates, amorphous and crystalline (layered) silicates, including aluminosilicates. Preferred salts are salts of sodium, potassium or magnesium. Highly preferred is the use of at least a carbonate salt or an aluminosilicate or mixtures thereof.

Suitable water-soluble solid granulation agents as organic acid carriers include monocarboxylic acids, monomeric polycarboxylic acids, homo or copolymeric polycarboxylic acids, inorganic acids, and mixtures thereof.

Suitable example of monocarboxylic acids containing one carboxy group include formic acid, acetic acid, propionic acid, butyric acid, valeric acid, caproic acid, enanthic acid, caprylic acid, pelargonic acid, capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, and mixtures thereof.

25

Still other suitable monocarboxylic acids are the monocarboxylic acids substituted by any of the following groups: CH₃- (CH₂)_n, wherein n is an integer of value of at least 1, CH₃, OH, NH₂, Cl, Br, F, I, OR'', NHR'', NR''², NO₂, SO₃, cyclic rings like cyclopentane, cyclohexane, phenyl, benzyl, or a mixture of these substituents; wherein 5 R'' is selected from saturated or unsaturated alkyl chains. Preferred examples are 1-methylcyclohexanecarboxylic acid, glycolic acid, mandelic acid, lactic acid, salicylic acid, benzoic acid, and derivatives thereof. The substituents may also be anywhere in the alkyl chain attached to the acidic function. The alkyl chain can be saturated or non saturated.

10

Other dicarboxylic acids suitable for use herein are the dicarboxylic acids substituted by CH₃- (CH₂)_n, wherein n is an integer of value of at least 1, CH₃, OH, NH₂, Cl, Br, F, I, OR'', NHR'', NR''², NO₂, SO₃, cyclic rings like cyclopentane, cyclohexane, phenyl, benzyl, or a mixture of these substituents; wherein R'' is selected from 15 saturated or unsaturated alkyl chain. Preferred examples of such substituted dicarboxylic acids are phthalic acid, isophthalic acid, terephthalic acid, malic acid, fumaric acid, tartaric acid, or mixtures thereof. The substituents may also be anywhere in the alkyl chain attached to the acidic functions. The alkyl chains can be saturated or non saturated.

20

Other polycarboxylic acids suitable for use herein are the polycarboxylic acids containing three carboxy groups and include, in particular, water-soluble citric acid, aconitic and citraconic acid as well as succinic derivatives such as the carboxymethyloxysuccinic described in British Patent No. 1,379,241, lactoxysuccinic described in British Patent No. 1,389,732, and aminosuccinic described in Netherlands Application 7205873, and the oxypolycarboxylic materials such as 2-oxa-1,1,3-propane tricarboxylic described in British Patent No. 1,387,447.

25

Other polycarboxylic acids suitable for use herein are the polycarboxylic acids containing four carboxy groups and include oxydisuccinic disclosed in British Patent 30 No. 1,261,829, 1,1,2,2-ethane tetracarboxylic, 1,1,3,3-propane tetracarboxylic and

1,1,2,3-propane tetracarboxylic. Polycarboxylic containing sulfo substituents include the sulfosuccinic derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in U.S. Patent No. 3,936,448, and the sulfonated pyrolysed citric acid described in British Patent No. 1,439,000.

5

Other suitable granulation agents are amino acids like, glycine, lysine, alanine, valine, leucine, isoleucine, proline, phenylalanine, tyrosine, tryptophan, serine, threonine, cysteine, methionine, asparagine, glutamine, aspartate, glutamate, arginine, histidine, and mixtures thereof.

10

Other suitable solid granulation agents are acid anhydrides and acyl halides. Acid anhydrides react in the presence of water to acids. Sometimes, the production of the amine reaction product is followed by the incomplete removal of the water in the amine samples. It may then be desired to remove the remaining water by reacting it with the acid anhydrides to form acids which in turn make the salt with the amine reaction product.

15 Preferably, to avoid possible hydrolysis of the product in-situ due to the eventual additional water coming from the acid carrier, the acid carrier is used in its anhydrous forms. For example, citric acid is available under anhydrous form or as a monohydrate.

20 Of the above, the preferred are polycarboxylic acids selected from citric acid, tartaric acid, malonic acid, succinic acid, oxalic acid, crotonic acid, adipic acid, maleic acid, malic acid, phthalic acid, succinic acid, hydroxysuccinic acid, polyacrylic acid, and mixtures thereof.

Preferred are also mixtures of granulation agents, for examples mixtures of inorganic salts or mixtures of organic acids and inorganic salts, including effervescing mixtures such as carboxylic acids and (bi)carbonates.

Suitable organic polymeric compounds suitable as solid granulation agents include cellulose derivatives such as methylcellulose, carboxymethylcellulose, hydroxypropylcellulose and hydroxyethylcellulose, as well as carbohydrates like pectins, and gums. Further compounds are carbohydrates and derivatives such as 5 fructose, xylose, galactose, galacturonic acid or glucose based polymers like inuline, dextran, xyloglucan, pectin or gums

Examples of carbonates are the alkaline earth and alkali metal carbonates, including 10 sodium carbonate and sesqui-carbonate and mixtures thereof with ultra-fine calcium carbonate as disclosed in German Patent Application No. 2,321,001 published on November 15, 1973.

Specific examples of water-soluble phosphates are the alkali metal tripolyphosphates, 15 sodium, potassium and ammonium pyrophosphate, sodium and potassium and ammonium pyrophosphate, sodium and potassium orthophosphate, sodium polymeta/phosphate in which the degree of polymerization ranges from about 6 to 21, and salts of phytic acid.


Typical disclosure of cyclodextrin derivatives are disclosed in WO96/05358, U.S. Pat. 20 Nos: 3,426,011, Parmerter et al., issued Feb. 4, 1969; 3,453,257; 3,453,258; 3,453,259; and 3,453,260, all in the names of Parmerter et al., and all issued July 1, 1969; 3,459,731, Gramera et al., issued Aug. 5, 1969; 3,553,191, Parmerter et al., issued Jan. 5, 1971; 3,565,887, Parmerter et al., issued Feb. 23, 1971; 4,535,152, Szejtli et al., issued Aug. 13, 1985; 4,616,008, Hirai et al., issued Oct. 7, 1986; 25 4,678,598, Ogino et al., issued Jul. 7, 1987; 4,638,058, Brandt et al., issued Jan. 20, 1987; and 4,746,734, Tsuchiyama et al., issued May 24, 1988; all of said patents being incorporated herein by reference.

30 Although less preferred for use herein because of their lower solubility, the solid granulation agents may also comprise silicates and aluminosilicates.

Suitable silicates include the water soluble sodium silicates with an SiO₂: Na₂O ratio of from 1.0 to 2.8, with ratios of from 1.6 to 2.4 being preferred, and 2.0 ratio being most preferred. The silicates may be in the form of either the anhydrous salt or a hydrated salt. Sodium silicate with an SiO₂: Na₂O ratio of 2.0 is the most preferred silicate.

5

Crystalline layered sodium silicates have the general formula



10

wherein M is sodium or hydrogen, x is a number from 1.9 to 4 and y is a number from 0 to 20. Crystalline layered sodium silicates of this type are disclosed in EP-A-0164514 and methods for their preparation are disclosed in DE-A-3417649 and DE-A-3742043. For the purpose of the present invention, x in the general formula above has a value of 15 2, 3 or 4 and is preferably 2. The most preferred material is δ -Na₂Si₂O₅, available from Hoechst AG as NaSKS-6.

Suitable aluminosilicate zeolites have the unit cell formula Na_z[(AlO₂)_z(SiO₂)_y].

20

XH₂O wherein z and y are at least 6; the molar ratio of z to y is from 1.0 to 0.5 and x is at least 5, preferably from 7.5 to 276, more preferably from 10 to 264. The aluminosilicate material are in hydrated form and are preferably crystalline, containing from 10% to 28%, more preferably from 18% to 22% water in bound form.

25

Typically when the amine reaction product is mixed with a solid granulation agent and further processed to form a particle, the amine reaction product will be present in an amount of from 1 to 75%, preferably 5 to 30%, more preferably 6 to 25% by weight of the processed reaction product in the produced particle.

Typically the granulation agent will be present in an amount of from 10% to 95%, preferably from 30 to 90%, more preferably, 50 to 75% by weight of the resulting granule.

5 One method for applying mixing with the solid granulation agent involves agglomeration. Any conventional agglomerator/mixer may be used including, but not limited to pan, rotary drum and vertical blender types. Molten coating compositions may also be applied either by being poured onto, or spray atomized onto a moving bed of the mixture of amine reaction product with carrier.

10

One typical method involves:

In the first part of the TSE, the active brought at a temperature between 5 and 40°C and the amine together with the perfume mix brought at a temperature between 5 and 40°C are incorporated into the TSE via their respective inlet and mixed together at a screw speed between 50 and 200, preferably 150 rpm, to make the resulting amine reaction product with perfume mix. Typical weight rate of material which is introduced in the TSE are of 5 to 200kg/hour for each of the active and of the amine. The temperature within the reaction mixture is preferably within the range of 20 to 40°C with a residence time between 10 and 45 seconds. Thereafter, the resulting product is brought along the TSE for dispersion into a carrier, preferably a carrier having a melting point between 30°C and 135°C, the carrier having been previously brought to a temperature between 20 and 150°C at a rate of between 50 and 200, preferably 150 kg/hour. The dispersion temperature at the end of the TSE was about 80°C and the total residence time of the mixture within the TSE is preferably between 10 seconds to 2 minutes. The resulting dispersion is then collected for optional agglomeration and/or coating process as outlined thereafter.

Specifically, in the first part of the TSE, the Damascone brought at a temperature of 20°C and Lupasol P (water free) with a perfume mixture brought at a temperature of 20°C are mixed at a screw speed of 150 rpm to make the resulting amine reaction

product with the perfume mixture, at a weight ratio of 20 to 40kg/hour, preferably about 25kg/hour, Damascone and 56kg/hour of Lupasol P (water free) with perfume mix of which the Lupasol P (water free) is 16kg/hour and. In the second part of the TSE, the amine reaction product is dispersed into TAE80 brought at a temperature of 5 70°C at a rate of 120kg/hour. The total production rate was thus 200kg/hour.

Still, an alternative process for making the amine reaction product in a carrier is by a batch process using a mixing tank in which pre-or melted therein carrier, e.g. TAE80 is placed into the mixing tank before incorporation of the amine component and 10 subsequently of the active both incorporated at room temperature.

The carried composition, or the carried composition mixed with the liquid carrier and/ or the granulation aid, may comprise further compounds to control the viscosity, as described herein. Examples are asticisers like phtalates, with tactifiers like rosin acids 15 or rosin esters, cross linking agents such as polyethylene oxide/ polypropylene oxide (co)polymers, polyethylene- or propylene-glycols, pentanal, nonanal, hexanal, heptanal, octanal, or bifunctional aldehydes, such as glutaraldehyde, and/ or with thickeners, such as for xample used in liquid detergent compositions. These agents can give the polymer the proper carrier characteristics like the required viscosity if the 20 viscosity is not high enough. Of course, other known viscosity enhancer may be used herein for that purpose.

Particle Size

For ease of handling and incorporation into the laundry and cleaning composition of the invention, it might be preferred to further process the carried composition. Typically, this involves making agglomerates of the above obtained viscous mix by first making a dispersion in a dispersing carrier like a water-soluble material having a melting point of from 30°C to 135°C like a nonionic ethoxylated alcohol surfactant and then agglomerating it with a coating material having a melting point between 35 and 135°C, like carbonate, starch, cyclodextrin, and mixtures thereof. Typical description of such process can respectively be found in co-pending application EP 99401736.6 at page 19 lines 11 to page 22 line 36 and page 28 line 31 to page 32 line 20.

Typically, the particle size of the carried composition in the dispersed carrier is from 0.1 micrometers to 150 micrometers, more preferably from 1 micrometer to 100 micrometers, and most preferably from 3 or even from 10 to 70 micrometers. When further processed, it has been found that in order for these processed carried benefit agent to impart their beneficial deposition and release on the surface, the agglomerated carried benefit agent preferably has an average particle diameter of from about 1 to about 2000 micrometers, preferably from about 150 to about 1700 micrometers, more preferably from about 250 to about 1000 micrometers. The term "average particle diameter" represents the mean particle size diameter of the actual particles of a given material. The mean is calculated on a weight percent basis. The mean is determined by conventional analytical techniques such as, for example, laser light diffraction or microscopic determination utilizing a scanning electron microscope. Preferably, greater than 50% by weight and more preferably greater than 60% by weight and most preferably greater than 70% by weight, of the particles have actual diameters which fall within the range of from about 250 to about 1000 micrometers, preferably from about 250 to about 850 micrometers.

The desired particle sizes can be achieved by, for example, mechanically grinding the resulting carried perfume composition in blenders (e.g., an Oster® blender) or in large

scale mills (e.g., a Wiley® Mill) to the desired particle size range or by prilling in a conventional manner (e.g., forcing the well-circulated co-melt through a heated nozzle into cooled atmospheric temperatures).

5 Laundry and cleaning and/or fabric care products

The carried perfume composition is then incorporated in a laundry or cleaning and/or fabric composition. Means of incorporation into the laundry and/or cleaning and/or fabric composition are conventionally known in the art, and is typically made depending on its end form by either spraying when in sprayable liquid form, or dry-addition.

10 Preferably, the carried composition is in processed form as mentioned above and incorporated by dry-addition.

15 Preferably, the carried composition which is incorporated into such laundry or cleaning and/or fabric composition provides a dry surface Odor Index of more than 5 preferably at least 10.

20 By Dry Surface Odor Index, it is meant that the carried composition provides a Delta of more than 5, wherein Delta is the difference between the Odor Index of the dry surface treated with the carried composition and the Odor Index of the dry surface treated with only the perfume raw material.

Measurement method of Dry Surface Odor Index:

For the above Dry Surface Odor Index, the carried composition suitable for use in the present invention needs to fulfill the following test.

25

Product preparation:

The carried composition is added to the unperfumed product base. Levels of carried composition are selected so as to obtain an odor grade on the dry fabric of at least 20. After careful mixing, by shaking the container in case of a liquid, with a spatula in case 30 of a powder, the product is allowed to sit for 24 hrs.

Washing process:

The resulting product is added into the washing machine in the dosage and in the dispenser appropriate for its category. The quantity corresponds to recommended dosages made for the corresponding market products: typically between 70 and 150 g for a detergent powder or liquid via current dosing device like granulette, or ariellette. The load is composed of four bath towels (170g) using a Miele W830 washing machine at 40°C short cycle, water input :15°Hardness at a temperature of 10-18°C, and full spin of 1200rpm.

10

The same process is applied for the corresponding free perfume ingredient in consideration and is used as the reference. Dosages, fabric loads and washing cycles for the reference and the sample are identical.

15

Drying Process:

Within two hours after the end of the washing cycle, the spinned but still wet fabrics are assessed for their odors using the scale mentioned below. Afterwards, half of the fabric pieces are hung on a line for 24 hr drying, away from any possible contaminations. Unless specified, this drying takes place indoor. Ambient conditions are at temperature between 18-25C and air moisture between 50-80%. The other half is placed in a tumble drier and undergoes a full "very dry" cycle, i.e. in a Miele, Novotronic T430 set on program white-extra dry (full cycle). Tumble dry fabrics are also assessed on the next day. Fabrics are then stored in opened aluminum bags in an odor free room, and assessed again after 7 days.

25

Odor Evaluations:

Odor is assessed by expert panellist smelling the fabrics. A 0-100 scale is used for all fabric odor gradings. The grading scale is as follows :

100 = extremely strong perfume odor

30 75 = very strong perfume odor

50 = strong odor
40 = moderate perfume odor
30 = slight perfume odor
20 = weak perfume odor
10 = very weak perfume odor
0 = no odor

5

A difference of more than 5 grades after one day and/or 7 days between the carried composition and the benefit agent, e.g. perfume is statistically significant. A difference 10 of 10 grades or more after one day and/or 7 days represents a step-change. In other words, when a difference of grade of more than 5, preferably at least 10 is observed between the amine reaction product and the perfume raw material, after either 1 day or 7 days or both 1 day and 7 days, it can be concluded that the carried composition is suitable for use in the present invention.

15

The laundry or cleaning composition typically comprises one or more detergent and/or cleaning and/or surfactant ingredient, whilst the fabric care composition will typically comprises a fabric care ingredient. By "fabric care ingredient", it is meant an ingredient which provide care to the fiber integrity of the treated fabric like a color 20 protecting agent, e.g DTI, crystal growth inhibitor, bleach quencher-scavenger, anti abrasive agent, etc....

Preferably, the invention composition is a laundry and/or cleaning composition.

25 Laundry compositions also encompass compositions providing color care, or composition counteracting malodours, as well as compositions suitable for use in any steps of the domestic treatment, that is as a pre-treatment composition, as a wash additive as a composition suitable for use in the rinse-cycle of the laundry cycle or applied on a dryer-sheet. Obviously, multiple applications can be made such as treating 30 the fabric with a pre-treatment composition of the invention and also thereafter with a

composition of the invention suitable for use in the rinse cycle and/or suitable for use as a dryer-sheet.

5 The liquid finished compositions of the invention may also be in a spray, foam, or aerosol form which for example can be suitable for use while ironing, or applied on the surfaces of the tumble dryer.

Laundry compositions encompass laundry detergent compositions, including liquid, solid form like powdered, tablets as well as softening compositions including rinse added softening composition as well as dryer added softening compositions.

10

A conventional disclosure of softening ingredients to be used in the softening composition of the invention can be found in EP 98870227.0, incorporated herein by reference, which typically include components selected from a surfactant like a quaternary ammonium softening component, a stabilising agent like a nonionic ethoxylated surfactant, a chelating agent, a crystal growth inhibitor, a soil release agent, 15 a polyalkyleneimine component, brighteners, preservatives, antibacterials, cyclodextrins, and mixtures thereof.

A conventional disclosure of a laundry or cleaning composition can be found in EP-A-20 0,659,876 and European patent application No. 98870226.2 which are both incorporated herein by reference.

25 Typical laundry or cleaning composition comprises a detergent and/or cleaning ingredient. By detergent or cleaning ingredient, it is meant ingredient which are respectively conventional to the detergent composition or cleaning composition. Typical of conventionals in detergent compositions includes one or more of surfactants, or organic and inorganic builders. The preferred laundry or cleaning composition, embodiment of the invention, will also preferably contain a bleaching system and/or other components conventional in detergent compositions. Typical of bleaching systems 30 include a peroxyacid, a hypohalite, or a bleach precursor with a source of alkaline

hydrogen peroxide necessary to form a peroxyacid bleaching species in the wash solution. Other optionals include soil suspending and anti-redeposition agents, suds suppressors, enzymes, fluorescent whitening agents, photoactivated bleaches, perfumes, colours, and mixtures thereof.

5

Preferably, the finished composition is a detergent composition, more preferably in solid form.

10 In addition, when the composition is a laundry composition, it is preferred that the detergent composition comprises a clay.

Clay

15 The compositions of the invention may preferably contain a clay, preferably present at a level of from 0.05% to 40%, more preferably from 0.5% to 30%, most preferably from 2% to 20% by weight of the composition. For clarity, it is noted that the term clay mineral compound, as used herein, excludes sodium aluminosilicate zeolite builder compounds, which however, may be included in the compositions of the invention as optional components.

20 One preferred clay may be a bentonite clay. Highly preferred are smectite clays, as for example disclosed in the US Patents No.s 3,862,058 3,948,790, 3,954,632 and 4,062,647 and European Patents No.s EP-A-299,575 and EP-A-313,146 all in the name of the Procter and Gamble Company.

25 The term smectite clays herein includes both the clays in which aluminium oxide is present in a silicate lattice and the clays in which magnesium oxide is present in a silicate lattice. Smectite clays tend to adopt an expandable three layer structure.

30 Specific examples of suitable smectite clays include those selected from the classes of the montmorillonites, hectorites, volchonskoites, nontronites, saponites and sauconites,

particularly those having an alkali or alkaline earth metal ion within the crystal lattice structure. Sodium or calcium montmorillonite are particularly preferred.

Suitable smectite clays, particularly montmorillonites, are sold by various suppliers
5 including English China Clays, Laviosa, Georgia Kaolin and Colin Stewart Minerals.

Clays for use herein preferably have a particle dimension of from 10nm to 800nm more
preferably from 20nm to 500 nm, most preferably from 50nm to 200 nm.

10 Particles of the clay mineral compound may be included as components of agglomerate
particles containing other detergent compounds. Where present as such components,
the term "largest particle dimension" of the clay mineral compound refers to the largest
dimension of the clay mineral component as such, and not to the agglomerated particle
as a whole.

15 Substitution of small cations, such as protons, sodium ions, potassium ions, magnesium
ions and calcium ions, and of certain organic molecules including those having
positively charged functional groups can typically take place within the crystal lattice
structure of the smectite clays. A clay may be chosen for its ability to preferentially
20 absorb one cation type, such ability being assessed by measurements of relative ion
exchange capacity. The smectite clays suitable herein typically have a cation exchange
capacity of at least 50 meq/100g. U.S. Patent No. 3,954,632 describes a method for
measurement of cation exchange capacity.

25 The crystal lattice structure of the clay mineral compounds may have, in a preferred
execution, a cationic fabric softening agent substituted therein. Such substituted clays
have been termed 'hydrophobically activated' clays. The cationic fabric softening agents
are typically present at a weight ratio, cationic fabric softening agent to clay, of from
1:200 to 1:10, preferably from 1:100 to 1:20. Suitable cationic fabric softening agents

include the water insoluble tertiary amines or dilong chain amide materials as disclosed in GB-A-1 514 276 and EP-B-0 011 340.

5 A preferred commercially available "hydrophobically activated" clay is a bentonite clay containing approximately 40% by weight of a dimethyl ditallow quaternary ammonium salt sold under the tradename Claytone EM by English China Clays International.

10 In a highly preferred embodiment of the invention, the clay is present in an intimate mixture or in a particle with a humectant and a hydrophobic compound, preferably a wax or oil, such as paraffin oil. Preferred humectants are organic compounds, including propylene glycol, ethylene glycol, dimers or trimers of glycol, most preferably glycerol. The particle is preferably an agglomerate. Alternatively, the particle may be such that the wax or oil and optionally the humectant form an encapsulate on the clay or alternatively, the clay be a encapsulate for the wax or oil and the humectant. It may be 15 preferred that the particle comprises an organic salt or silica or silicate.

20 However, in another embodiment of the invention, the clay is preferably mixed with one or more surfactants and optionally builders and optionally water, in which case the mixture is preferably subsequently dried. Preferably, such a mixture is further processed in a spray-drying method to obtain a spray dried particle comprising the clay.

It may be preferred that the flocculating agent is also comprised in the particle or granule comprising the clay.

25 It may also be preferred that the intimate mixture comprises a chelating agent.

Flocculating agent

30 The compositions of the invention may contain a clay flocculating agent, preferably present at a level of from 0.005% to 10%, more preferably from 0.05% to 5%, most preferably from 0.1% to 2% by weight of the composition.

The clay flocculating agent functions such as to bring together the particles of clay compound in the wash solution and hence to aid their deposition onto the surface of the fabrics in the wash. This functional requirement is hence different from that of clay dispersant compounds which are commonly added to laundry detergent compositions to aid the removal of clay soils from fabrics and enable their dispersion within the wash solution.

Preferred as clay flocculating agents herein are organic polymeric materials having an average weight of from 100,000 to 10,000,000, preferably from 150,000 to 5,000,000, more preferably from 200,000 to 2,000,000.

Suitable organic polymeric materials comprise homopolymers or copolymers containing monomeric units selected from alkylene oxide, particularly ethylene oxide, acrylamide, acrylic acid, vinyl alcohol, vinyl pyrrolidone, and ethylene imine. Homopolymers of, on particular, ethylene oxide, but also acrylamide and acrylic acid are preferred.

European Patents No.s EP-A-299,575 and EP-A-313,146 in the name of the Procter and Gamble Company describe preferred organic polymeric clay flocculating agents for use herein.

The weight ratio of clay to the flocculating polymer is preferably from 1000:1 to 1:1, more preferably from 500:1 to 1:1, most preferably from 300:1 to 1:1, or even more preferably from 80:1 to 10:1, or in certain applications even from 60:1 to 20:1.

25


Inorganic clay flocculating agents are also suitable herein, typical examples of which include lime and alum.

The flocculating agent is preferably present in a detergent base granule such as a detergent agglomerate, extrudate or spray-dried particle, comprising generally one or more surfactants and builders.

5 Effervescent

Effervescent means may also be optionally used in the compositions of the invention.

Effervescency as defined herein means the evolution of bubbles of gas from a liquid, as the result of a chemical reaction between a soluble acid source and an alkali metal 10 carbonate, to produce carbon dioxide gas,



Further examples of acid and carbonate sources and other effervescent systems may be found in : (Pharmaceutical Dosage Forms : Tablets Volume 1 Page 287 to 291).

15

Carbonate salts

Suitable alkali and/ or earth alkali inorganic carbonate salts herein include carbonate and hydrogen carbonate of potassium, lithium, sodium, and the like amongst which sodium and potassium carbonate are preferred. Suitable bicarbonates to be used herein include 20 any alkali metal salt of bicarbonate like lithium, sodium, potassium and the like, amongst which sodium and potassium bicarbonate are preferred. However, the choice of carbonate or bicarbonate or mixtures thereof may be made depending on the pH desired in the aqueous medium wherein the granules are dissolved. For example where a relative high pH is desired in the aqueous medium (e.g., above pH 9.5) it may be 25 preferred to use carbonate alone or to use a combination of carbonate and bicarbonate wherein the level of carbonate is higher than the level of bicarbonate. The inorganic alkali and/ or earth alkali carbonate salt of the compositions of the invention comprises preferably a potassium or more preferably a sodium salt of carbonate and/ or bicarbonate. Preferably, the carbonate salt comprises sodium carbonate, optionally also 30 a sodium bicarbonate.

The inorganic carbonate salts herein are preferably present at a level of at least 20% by weight of the composition. Preferably they are present at a level of at least 23% or even 25% or even 30% by weight, preferably up to about 60% by weight or more 5 preferably up to 55% or even 50% by weight.

They may be added completely or partially as separate powdered or granular component, as co-granules with other detergent ingredients, for example other salts or surfactants. In solid detergent compositions of the invention, they may also completely 10 or partially be present in detergent granules such as agglomerates or spray dried granules.

In one embodiment of the invention, an effervescence source is present, preferably comprising an organic acid, such as carboxylic acids or aminoacids, and a carbonate. 15 Then it may be preferred that part or all of the carbonate salt herein is premixed with the organic acid, and thus present in an separate granular component.

Preferred effervescent source are selected from compressed particles of citric acid and carbonate optionally with a binder; and particle of carbonate, bicarbonate and malic or maleic acid in weight ratios of 4:2:4. The dry add form of citric acid and carbonate are preferably used.

The carbonate may have any particle size. In one embodiment, in particular when the carbonate salt is present in a granule and not as separately added compound, the carbonate salt has preferably a volume median particle size from 5 to 375 micrometers, whereby preferably at least 60%, preferably at least 70% or even at least 80% or even at least 90% by volume, has a particle size of from 1 to 425 micrometers. More preferably, the carbon dioxide source has a volume median particle size of 10 to 250, whereby preferably at least 60 %, or even at least 70% or even at least 80% or even at least 90% by volume, has a particle size of from 1 to 375 micrometers; or even

preferably a volume median particle size from 10 to 200 micrometers, whereby preferably at least 60 %, preferably at least 70% or even at least 80% or even at least 90% by volume, has a particle size of from 1 to 250 micrometers.

In particular when the carbonate salt is added as separate component, so to say 'dry-added' or admixed to the other detergent ingredients, the carbonate may have any particle size, including the above specified particle sizes, but preferably at least an 5 volume average particle size of 200 micrometers or even 250 micrometers or even 300 micrometers.

It may be preferred that the carbon dioxide source of the required particle size is obtained by grinding a larger particle size material, optionally followed by selecting the 10 material with the required particle size by any suitable method.

Whilst percarbonate salts may be present in the compositions of the invention as a bleaching agent, they are not included in the carbonate salts as defined herein

15 Form of the composition

The composition of the invention may take a variety of physical form including liquid, gel, foam in either aqueous or non-aqueous form, or solid form, including bar, beads, granular and tablet forms.

20 Still in another aspect of the invention, there is provided a packaged composition comprising the processed product of the invention or composition of the invention. Preferably, the packaged composition is a closed packaging system having a moisture vapour transmission rate of less than 20g/m²/24 hours. Typical disclosure of such a package can be found in WO 98/40464.

25 Still another preferred package is a spray dispenser.

Spray Dispenser

The present invention also relates to such compositions incorporated into a spray dispenser to create an article of manufacture that can facilitate treatment of fabric articles and/or surfaces with said compositions containing the amine reaction product and other ingredients (examples are cyclodextrins, polysaccharides, polymers, surfactant, perfume, softener) at a level that is effective, yet is not discernible when dried on the surfaces. The spray dispenser comprises manually activated and non-manual powered (operated) spray means and a container containing the treating composition. Typical disclosure of such spray dispenser can be found in WO 96/04940 page 19 line 21 to page 22 line 27. The articles of manufacture preferably are in association with instructions for use to ensure that the consumer applies sufficient ingredient of the composition to provide the desired benefit. Typical compositions to be dispensed from a sprayer contain a level of amine reaction product of from about 0.01% to about 5%, preferably from about 0.05% to about 2%, more preferably from about 0.1% to about 1%, by weight of the usage composition.

Method

Also provided herein is a method for providing an enhanced deposition as well as a delayed release of the benefit agent, preferably a perfume composition, which comprises the step of contacting the surface to be treated with a composition of the invention, and preferably thereafter contacting the treated surface with a material, preferably an aqueous medium like moisture or any other means susceptible of releasing the perfume from the composition.

By "surface", it is meant any surface onto which the compound can deposit. Typical examples of such material are fabrics, hard surfaces such as dishware, floors, bathrooms, toilet, kitchen and other surfaces in need of a delayed release of a perfume such as that with litter like animal litter. Preferably, the surface is selected from a fabric, a tile, a ceramic; more preferably is a fabric.

By "enhanced deposition", it is meant a better deposition of the benefit agent (e.g. perfume) on the treated surface than by the use of the benefit agent (e.g. perfume) itself.

5 By "delayed release" is meant release of the benefit agent (e.g perfume) over a longer period of time than by the use of the benefit agent (e.g., perfume) itself.

10 Where the carrier is a polymer or component which has been chemically reacted with a benefit agent like perfume, the release of the benefit agent which is entrapped or embedded within the reacted carrier, i.e. not chemically reacted, is released from the carried benefit agent composition by hydrolysis of the reacted carrier. Indeed, the hydrolysis of the "protective shell" made by the reacted carrier into the respective aldehyde and/or ketone on the one hand and the polymer on the other will gradually open the shell, thereby enabling release of the entrapped benefit agent.

15 Still in another aspect of the invention, there is provided the use of the product of the invention for the manufacture of a laundry and cleaning composition for delivering residual fragrance onto the fabrics on which it is applied.

20 For the purposes of the present invention the term "contacting" is defined as "intimate contact of a surface with an aqueous solution of the hereinabove described composition." Contacting typically occurs by soaking, washing, rinsing the composition onto fabric, but can also include contact of a substrate *inter alia* a material onto which the composition has been absorbed, with the fabric.

25 EXAMPLES

I-Synthesis Example of a carrier and a perfume mix benefit agent

30 In a reaction vessel of 2l, placed on a rotary evaporator, 100g of δ -Damascone and 150g of LupasolP(about 50% of water) and 175g of a perfume mixture are mixed together for 4 hours at 42°C. The temperature of the reaction mixture, during the

mixing, is controlled via a thermostat and not allowed to go higher than 42°C. 335g of product is obtained and only traces of unreacted δ-Damascone remain. The viscosity of the synthesised product is 55000 cps.

5 II-Synthesis Example of a carrier and a perfume mix benefit agent

In a reaction vessel of 250ml, 20g of Lilial and 16g of water-free Lupasol P (water-free Lupasol P is taken from the commercial Lupasol sample from which the water has been removed by vacuum distillation) and 83g of a perfume mixture are mixed together for 4 hours at 42°C. The temperature of the reaction mixture, during the mixing, is controlled via a thermostat and not allowed to go higher than 42°C. 118g of product is obtained and only traces of unreacted Lilial remain. The viscosity of the synthesised product is 1600cps.

III-Synthesis Example of a carrier and a perfume mix benefit agent

15 In a reaction vessel of 250ml, 12g of Carvone and 10g of waterfree Lupasol P and 49g of a perfume mixture are mixed together for 4 hours at 42°C. The temperature of the reaction mixture, during the mixing, is controlled via a thermostat and not allowed to go higher than 42°C. 71g of product is obtained and most of the Carvone has reacted . The viscosity of the synthesised product is 2300cps.

20

IV-Synthesis Example of a carrier and a perfume mix benefit agent

25 In a reaction vessel of 250ml, 12g of Triplal and 10g of waterfree Lupasol P and 22g of a perfume mixture are mixed together for 4 hours at 42°C. The temperature of the reaction mixture, during the mixing, is controlled via a thermostat and not allowed to go higher than 42°C. 42g of product is obtained and only traces of unreacted Triplal remain. The viscosity of the synthesised product is 9764cps.

V-Synthesis Example of a benefit agent with a carrier

30 Lupasol WF was reacted with palmitoylchloride. 32g of LupasolWF was dissolved in 250ml of dry dichloromethane. The solution was cooled to 0C and 76ml of palmitoyl

chloride dissolved in 50ml dry dichloromethane with a dropping funnel. The solution was stirred 1hour under N2 atmosphere. The reaction products were washed with a saturated water solution of potassiumcarbonate. After washing, the reaction product is dried by vaccum distillation. 88g of reaction product is obtained.

5 8 g of the above product is mixed with 24 g of perfume mix at 40C till a yellow viscous and homogeneous product is obtained.

Any type of perfume mixture may be used. One preferred composition of the perfume mix is as follows:

10

Citronellol	7
Geraniol	7
Linalool	7
Para Tertiary Butyl Cyclohexyl Acetate	10
Phenyl Ethyl Alcohol	19
Habanolide	4.5
Para Methoxy Acetophenone	1.5
Benzyl Acetate	4
Eugenol	2
Phenyl Ethyl Acetate	5
Verdyl Acetate	6
Verdyl Propionate	4
Hexyl Cinnamic Aldehyde	3
Ionone Gamma Methyl	2
Methyl Cedrylone	10
P.T. Bucinal	7
Para Cresyl Methyl Ether	1

The synthesised "carried composition" may be used as is or may be further processed to enable easy incorporation into finished product.

VI-Synthesis Example of a carrier and a perfume mix benefit agent

In a reaction vessel of 250ml, 8 gms of perfume mix FC1 and 2.5 g of waterfree Lupasol WF is mixed together for 30 minutes hours at room temperature C. The 5 temperature of the reaction mixture, during the mixing, is controlled via a thermostat and not allowed to go higher than 80 C. After mixing the mixture is kept overnight in a waterbath at 60C. The product thus obtained is a mixture of Lupasol completely reacted with molar proportions of each of the aldehydes of FC1, and unreacted aldehydes of FC1. All of the Lupasol WF is assumed to be reacted. The viscosity of the synthesised 10 product is 190.000 cps.

FC1:-

Methyl Nonyl Acetaldehyde :	15
Undecylenic Aldehyde :	30
Triplal :	35
15 Lauric Aldehyde :	19.5
Iris Aldehyde :	0.5

Processing method

Processing of the carried composition is done as follows: 80g of one of the carried 20 composition as above synthesised is mixed in an Ultra Turrax containing 120g of dispersing carrier, e.g. TAE80 or PEG 1000 to 10,000, for 5 minutes, the temperature of mixing being of about 70°C (melting temperature of the carrier), and the speed of the mixer being sufficient so as to maintain such temperature substantially constant. Temperature and time will depend on the nature of the dispersing carrier but are 25 conventional steps to the skilled man. The resulting mixture is maintained at a temperature substantially equal to the melting point of the carrier material. Once the mixture is at a suitable temperature, it is poured onto the coating material i.e. carbonate and agglomerated in an electrical mixer like a Braun Mixer. Care is also taken that the temperature during the mixing does not substantially exceed the melting point of the 30 carrier material. For example, 150g of a mixture containing 90g TAE80 and 60g of the

carried composition is poured at 60°C into a Braun Mixer containing 300g of carbonate. The mixing of the ingredients is carried out for about 5 minutes. Care is also taken that the temperature during the mixing does not exceed 65°C. Again, temperature and time will depend on the nature of the coating agent but are conventional steps to the skilled man.

Abbreviations used in the following laundry and cleaning composition Examples

In the laundry and cleaning compositions, the abbreviated component identifications have the following meanings:

10

DEQA	:	Di-(tallowyl-oxy-ethyl) dimethyl ammonium chloride
DTDMAC	:	Ditallow dimethylammonium chloride
DEQA (2)	:	Di-(soft-tallowyloxyethyl) hydroxyethyl methyl ammonium methylsulfate.
DTDMAMS	:	Ditallow dimethyl ammonium methylsulfate.
SDASA	:	1:2 ratio of stearyldimethyl amine:triple-pressed stearic acid.
Fatty acid	:	Stearic acid of IV =0
Electrolyte	:	Calcium chloride
PEG	:	Polyethylene Glycol 4000
Neodol 45-13	:	C14-C15 linear primary alcohol ethoxylate, sold by Shell Chemical CO.
Silicone antifoam	:	Polydimethylsiloxane foam controller with siloxane-oxyalkylene copolymer as dispersing agent with a ratio of said foam controller to said dispersing agent of 10:1 to 100:1.
PEI	:	Polyethyleneimine with an average molecular weight of 1800 and an average ethoxylation degree of 7 ethyleneoxy residues per nitrogen
HEDP	:	1,1-hydroxyethane diphosphonic acid
15 LAS	:	Sodium linear C ₁₁₋₁₃ alkyl benzene sulfonate
TAS	:	Sodium tallow alkyl sulfate

	CxyAS	:	Sodium C _{1x} - C _{1y} alkyl sulfate
	C46SAS	:	Sodium C ₁₄ - C ₁₆ secondary (2,3) alkyl sulfate
	CxyEzS	:	Sodium C _{1x} -C _{1y} alkyl sulfate condensed with z moles of ethylene oxide
5	CxyEz	:	C _{1x} -C _{1y} predominantly linear primary alcohol condensed with an average of z moles of ethylene oxide
	QAS	:	R ₂ .N ⁺ (CH ₃) ₂ (C ₂ H ₄ OH) with R ₂ = C ₁₂ - C ₁₄
	QAS 1	:	R ₂ .N ⁺ (CH ₃) ₂ (C ₂ H ₄ OH) with R ₂ = C ₈ - C ₁₁
	APA	:	C ₈ - C ₁₀ amido propyl dimethyl amine
10	Soap	:	Sodium linear alkyl carboxylate derived from an 80/20 mixture of tallow and coconut fatty acids
	STS	:	Sodium toluene sulphonate
	CFAA	:	C ₁₂ -C ₁₄ (coco) alkyl N-methyl glucamide
	TFAA	:	C ₁₆ -C ₁₈ alkyl N-methyl glucamide
15	TPKFA	:	C ₁₂ -C ₁₄ topped whole cut fatty acids
	STPP	:	Anhydrous sodium tripolyphosphate
	TSPP	:	Tetrasodium pyrophosphate
	Zeolite A	:	Hydrated sodium aluminosilicate of formula Na ₁₂ (AlO ₂ SiO ₂) ₁₂ .27H ₂ O having a primary particle size in the range from 0.1 to 10 micrometers (weight expressed on an anhydrous basis)
20	NaSKS-6	:	Crystalline layered silicate of formula δ- Na ₂ Si ₂ O ₅
	Citric acid	:	Anhydrous citric acid
	Borate	:	Sodium borate
25	Carbonate	:	Anhydrous sodium carbonate with a particle size between 200μm and 900μm
	Bicarbonate	:	Anhydrous sodium bicarbonate with a particle size distribution between 400μm and 1200μm

Silicate	:	Amorphous sodium silicate (SiO ₂ :Na ₂ O = 2.0:1)
Sulfate	:	Anhydrous sodium sulfate
Mg sulfate	:	Anhydrous magnesium sulfate
Citrate	:	Tri-sodium citrate dihydrate of activity 86.4% with a 5 particle size distribution between 425μm and 850μm
MA/AA	:	Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 70,000
MA/AA (1)	:	Copolymer of 4:6 maleic/acrylic acid, average molecular weight about 10,000
10 AA	:	Sodium polyacrylate polymer of average molecular weight 4,500
CMC	:	Sodium carboxymethyl cellulose
Cellulose ether	:	Methyl cellulose ether with a degree of polymerization of 650 available from Shin Etsu Chemicals
15 Protease	:	Proteolytic enzyme, having 3.3% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Savinase
Protease I	:	Proteolytic enzyme, having 4% by weight of active enzyme, as described in WO 95/10591, sold by Genencor 20 Int. Inc.
Alcalase	:	Proteolytic enzyme, having 5.3% by weight of active enzyme, sold by NOVO Industries A/S
Cellulase	:	Cellulytic enzyme, having 0.23% by weight of active enzyme, sold by NOVO Industries A/S under the 25 tradename Carezyme
Amylase	:	Amylolytic enzyme, having 1.6% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Termamyl 120T

	Lipase	:	Lipolytic enzyme, having 2.0% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Lipolase
5	Lipase (1)	:	Lipolytic enzyme, having 2.0% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Lipolase Ultra
	Endolase	:	Endoglucanase enzyme, having 1.5% by weight of active enzyme, sold by NOVO Industries A/S
10	PB4	:	Sodium perborate tetrahydrate of nominal formula $\text{NaBO}_2 \cdot 3\text{H}_2\text{O} \cdot \text{H}_2\text{O}_2$
	PB1	:	Anhydrous sodium perborate bleach of nominal formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$
	Percarbonate	:	Sodium percarbonate of nominal formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$
15	NOBS	:	Nonanoyloxybenzene sulfonate in the form of the sodium salt
	NAC-OBS	:	(6-nonamidocaproyl) oxybenzene sulfonate
	TAED	:	Tetraacetyl ethylenediamine
	DTPA	:	Diethylene triamine pentaacetic acid
20	DTPMP	:	Diethylene triamine penta (methylene phosphonate), marketed by Monsanto under the Tradename Dequest 2060
	EDDS	:	Ethylenediamine-N,N'-disuccinic acid, (S,S) isomer in the form of its sodium salt.
25	Photoactivated bleach (1)	:	Sulfonated zinc phthlocyanine encapsulated in dextrin soluble polymer
	Photoactivated bleach (2)	:	Sulfonated alumino phthlocyanine encapsulated in dextrin soluble polymer
	Brightener 1	:	Disodium 4,4'-bis(2-sulphostyryl)biphenyl

Brightener 2	:	Disodium 4,4'-bis(4-anilino-6-morpholino-1,3,5-triazin-2-yl)amino) stilbene-2:2'-disulfonate
HEDP	:	1,1-hydroxyethane diphosphonic acid
PEGx	:	Polyethylene glycol, with a molecular weight of x 5 (typically 4,000)
PEO	:	Polyethylene oxide, with an average molecular weight of 200000 to 400000
TEPAE	:	Tetraethylenepentaamine ethoxylate
PVI	:	Polyvinyl imidosole, with an average molecular weight of 20,000 10
PVP	:	Polyvinylpyrrolidone polymer, with an average molecular weight of 60,000
PVNO	:	Polyvinylpyridine N-oxide polymer, with an average molecular weight of 50,000
15 PVPVI	:	Copolymer of polyvinylpyrrolidone and vinylimidazole, with an average molecular weight of 20,000
QEA	:	bis((C ₂ H ₅ O)(C ₂ H ₄ O) _n)(CH ₃) -N ⁺ -C ₆ H ₁₂ -N ⁺ -(CH ₃) bis((C ₂ H ₅ O)-(C ₂ H ₄ O)) _n , wherein n = from 20 to 30
SRP 1	:	Anionically end capped poly esters
20 SRP 2	:	Diethoxylated poly (1, 2 propylene terephthalate) short block polymer
PEI	:	Polyethyleneimine with an average molecular weight of 1800 and an average ethoxylation degree of 7 ethyleneoxy residues per nitrogen
25 Silicone antifoam	:	Polydimethylsiloxane foam controller with siloxane-oxyalkylene copolymer as dispersing agent with a ratio of said foam controller to said dispersing agent of 10:1 to 100:1
Opacifier	:	Water based monostyrene latex mixture, sold by BASF 30 Aktiengesellschaft under the tradename Lytron 621

Wax	:	Paraffin wax
PA30	:	Polyacrylic acid of average molecular weight of between about 4,500 - 8,000.
480N	:	Random copolymer of 7:3 acrylate/methacrylate, average molecular weight about 3,500.
Polygel/carbopol	:	High molecular weight crosslinked polyacrylates.
Metasilicate	:	Sodium metasilicate (SiO ₂ :Na ₂ O ratio = 1.0).
Nonionic	:	C ₁₃ -C ₁₅ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5.
Neodol 45-13	:	C ₁₄ -C ₁₅ linear primary alcohol ethoxylate, sold by Shell Chemical CO.
MnTACN	:	Manganese 1,4,7-trimethyl-1,4,7-triazacyclononane.
PAAC	:	Pentaamine acetate cobalt(III) salt.
Paraffin	:	Paraffin oil sold under the tradename Winog 70 by Wintershall.
NaBz	:	Sodium benzoate.
BzP	:	Benzoyl Peroxide.
SCS	:	Sodium cumene sulphonate.
BT	:	Benzotriazole.
PH	:	Measured as a 1% solution in distilled water at 20°C.
CaP1	:	Processed amine reaction product of d-Damascone with Lupasol P and perfume mix as made from Synthesis example I, mixed with a carrier and agglomerated with TAE80 coating agent according to processing method above described.

CaP2 : Processed amine reaction product of Lupasol P with Lilial and perfume mix as made from Synthesis example II, and agglomerated with PEG4000 and carbonate coating agent according to processing method above described.

CaP 3 : Processed amine reaction product of Lupasol P with Carvone and perfume mix as made from Synthesis example III, mixed with a carrier and agglomerated with TAE80 coating agent according to processing method above described.

CaP 4 : Processed amine reaction product of Lupasol P with Triplal and perfume mix as made from Synthesis example IV, mixed with a carrier and agglomerated with PEG4000 coating agent according to processing method above described.

CaP 5 : Processed amine reaction product of Lupasol WF with palmitoylchloride and perfume mix. as made from Synthesis example V, mixed with a carrier and agglomerated with TAE80 coating agent according to processing method above described.

CAP6 Processed amine reaction product of Lupasol P with Lilial and perfume mix as made from Synthesis example II,

CAP7 Processed amine reaction product of Lupasol P with Carvone and perfume mix as made from Synthesis example III,

CAP8 Processed amine reaction product of Lupasol P with Triplal and perfume mix as made from Synthesis example IV,

CAP9	Processed amine reaction product of Lupasol WF and perfume mix as made in Synthesis example VI
Clay I	Bentonite clay
Clay II	Smectite clay
Flocculating agent I	polyethylene oxide of average molecular weight of between 200,000 and 400,000
5 Flocculating agent II	polyethylene oxide of average molecular weight of between 400,000 and 1,000,000
Flocculating agent III	polymer of acrylamide and/ or acrylic acid of average molecular weight of 200,000 and 400,000
DOBS	Decanoyl oxybenzene sulfonate in the form of the sodium salt
10 SRP 3	Polysaccharide soil release polymer
SRP 4	Nonionically end capped poly esters
Polymer	Polyvinylpyrrolidone K90 available from BASF under the tradename Luviskol K90
15 Dye fixative	Dye fixative commercially available from Clariant under the tradename Cartafix CB
Polyamine	1,4-Bis-(3-aminopropyl)piperazine
Bayhibit AM	2-Phosphonobutane-1,2,4-tricarboxylic acid commercially available from Bayer
20 Fabric softener active	Di-(canoloyl-oxy-ethyl)hydroxyethyl methyl ammonium methylsulfate
HPBDC	Hydroxypropyl beta-cyclodextrin
RAMEB	Randomly methylated beta-cyclodextrin
Bardac 2050	Diethyl dimethyl ammonium chloride, 50% solution
25 Bardac 22250	Didecyl dimethyl ammonium chloride, 50% solution
Genamin C100	Coco fatty amine ethoxylated with 10 moles

ethylene oxide and commercially available
from Clariant

Genapol V4463 : Coco alcohol ethoxylated with 10 moles
ethylene oxide and commercially available
from Clariant

5 Silwet 7604: Polyalkyleneoxide polysiloxanes of MW 4000 of formula

$$R-(CH_3)_2SiO-[(CH_3)_2SiO]_a-[(CH_3)(R)SiO]_b-Si(CH_3)_2-R,$$
 wherein average $a+b$ is 21, and commercially available
from Osi Specialties, Inc., Danbury, Connecticut

10 Silwet 7600: Polyalkyleneoxide polysiloxanes of MW 4000, of formula

$$R-(CH_3)_2SiO-[(CH_3)_2SiO]_a-[(CH_3)(R)SiO]_b-Si(CH_3)_2-R,$$
 wherein average $a+b$ is 11, and commercially available
from Osi Specialties, Inc., Danbury, Connecticut

15 In the following formulation examples all levels are quoted as % by weight of the composition unless otherwise stated, and incorporation of the carried perfume composition so called herein after "CAP" in the fully formulated composition is carried out as is.

20 Example 1

The following high density granular laundry detergent compositions are in accord with the invention:

	A	B	C	D	E
LAS	6.0	6.0	8.0	8.0	8.0
TAS	1.0	0.1	-	0.5	-
C46(S)AS	-	-	2.0	2.5	-
C25AS	4.5	5.5	-	-	-
C68AS	-	-	2.0	5.0	7.0
C25E5	4.6	4.6	-	-	3.4

C25E7	-	-	3.4	3.4	1.0
C25E3S	5.0	4.5	-	-	-
QAS	-	-	-	0.8	-
QAS (I)	0.5	1.0	-	-	-
Zeolite A	20.0	18.1	18.1	18.0	14.1
Citric acid	-	2.5	-	-	-
Carbonate	10.0	13.0	13.0	13.0	25.0
SKS-6	-	10.0	-	-	-
Silicate	0.5	0.3	1.4	1.4	3.0
Citrate	-	-	-	1.0	-
Sulfate	-	-	26.1	26.1	26.1
Mg sulfate	-	0.2	0.3	-	-
MA/AA	1.0	1.0	0.3	0.3	0.3
CMC	0.4	0.4	0.2	0.2	0.2
PB4	-	-	9.0	9.0	5.0
Percarbonate	18.0	18.0	-	-	-
TAED	3.9	4.2	1.5	0.4	1.5
NAC-OBS	-	-	-	2.0	1.0
DTPMP	-	-	0.25	0.25	0.25
SRP 2	-	0.2	-	-	-
EDDS	0.5	0.5	-	0.25	0.4
CFAA	-	-	-	1.0	-
HEDP	0.4	0.4	0.3	0.3	0.3
QEA	-	0.5	-	-	-
Protease I	-	-	-	-	0.26
Protease	1.5	1.0	0.26	0.26	-
Cellulase	0.3	0.3	0.3	-	-
Amylase	0.5	0.5	0.1	0.1	0.1
Lipase (1)	0.5	0.5	0.3	-	-

Photoactivated bleach (ppm)	20 ppm	20 ppm	15 ppm	15 ppm	15 ppm
Brightener 1	0.09	0.09	0.09	0.09	0.09
Perfume spray on	0.4	0.08	0.05	0.3	0.3
CAP 1	2.0	1.0	0.05	0.1	0.5
CAP 3	-	0.5	-	-	-
Silicone antifoam	0.3	0.3	0.5	0.5	0.5
Misc/minors to 100%					
Density in g/litre	850	850	850	850	850

	F	G	H	I
LAS	2.0	6.0	6.0	5.0
TAS	0.5	1.0	0.1	1.5
C25AS	7.0	4.5	5.5	2.5
C68AS	-	-	-	0.2
C25E5	10.0	4.6	4.6	2.6
C25E3S	2.0	5.0	4.5	0.5
QAS (I)	0.8	0.5	1.0	1.5
Zeolite A	18.1	20.0	18.1	16.2
Citric acid	2.5	-	2.5	1.5
Carbonate	10.0	10.0	13.0	20.6
SKS-6	10.0	-	10.0	4.3
Silicate	0.3	0.5	0.3	-
Citrate	3.0	-	-	1.4
Sulfate	6.0	-	-	-
Mg sulfate	0.2	-	0.2	0.03
MA/AA	4.0	1.0	1.0	0.6

CMC	0.2	0.4	0.4	0.3
Percarbonate	-	18.0	18.0	9.0
TAED	-	3.9	4.2	3.2
DTPMP	0.25	-	-	-
SRP 2	0.2	-	0.2	-
EDDS	-	0.5	0.5	0.1
CFAA	2.0	-	-	-
TFAA	-	-	-	1.1
HEDP	0.3	0.4	0.4	0.3
QEA	0.2	-	0.5	-
Protease I	1.0	-	-	0.3
Protease	-	1.5	1.0	-
Cellulase	0.3	0.3	0.3	0.3
Amylase	0.4	0.5	0.5	0.1
Lipase (1)	0.5	0.5	0.5	0.1
Photoactivated bleach (ppm)	-	20 ppm	20 ppm	20 ppm
PVNO/PVPVI	0.1	-	-	-
Brightener 1	-	0.09	0.09	0.01
Brightener 2	-	-	-	0.09
Perfume spray on	0.4	0.4	0.04	-
CAP 2	2.0	1.0	0.1	0.8
Silicone antifoam	-	0.3	0.3	0.3
Clay II	-	-	-	12.0
Flocculating agent I	-	-	-	0.3
Glycerol	-	-	-	0.6
Wax	-	-	-	0.4
Misc/minors to				

100%				
Density in g/litre	850	850	850	850

Example 2

The following granular laundry detergent compositions of particular utility under European machine wash conditions are in accord with the invention:

5

	A	B	C	D	E	F
LAS	5.5	7.5	5.0	5.0	6.0	7.0
TAS	1.25	1.86	-	0.8	0.4	0.3
C24AS/C25AS	-	2.24	5.0	5.0	5.0	2.2
C25E3S	-	0.76	1.0	1.5	3.0	1.0
C45E7	3.25	-	-	-	-	3.0
TFAA	-	-	2.0	-	-	-
C25E5	-	5.5	-	-	-	-
QAS	0.8	-	-	-	-	-
QAS II	-	0.7	1.0	0.5	1.0	0.7
STPP	19.7	-	-	-	-	-
Zeolite A	-	19.5	25.0	19.5	20.0	17.0
NaSKS-6/citric acid (79:21)	-	10.6	-	10.6	-	-
NaSKS-6	-	-	9.0	-	10.0	10.0
Carbonate	6.1	10.0	9.0	10.0	10.0	18.0
Bicarbonate	-	2.0	7.0	5.0	-	2.0
Silicate	6.8	-	-	0.3	0.5	-
Citrate	-	-	4.0	4.0	-	-
Sulfate	39.8	-	-	5.0	-	12.0
Mg sulfate	-	-	0.1	0.2	0.2	-
MA/AA	0.5	1.6	3.0	4.0	1.0	1.0

CMC	0.2	0.4	1.0	1.0	0.4	0.4
PB4	5.0	12.7	-	-	-	-
Percarbonate	-	-	-	-	18.0	15.0
TAED	0.5	3.1	-	-	5.0	-
NAC-OBS	1.0	3.5	-	-	-	2.5
DTPMP	0.25	0.2	0.3	0.4	-	0.2
HEDP	-	0.3	-	0.3	0.3	0.3
QEA	-	-	1.0	1.0	1.0	-
Protease I	-	-	-	0.5	1.2	-
Protease	0.26	0.85	0.9	1.0	-	0.7
Lipase (1)	0.15	0.15	0.3	0.3	0.3	0.2
Cellulase	0.28	0.28	0.2	0.2	0.3	0.3
Amylase	0.1	0.1	0.4	0.4	0.6	0.2
PVNO/PVPVI	-	-	0.2	0.2	-	-
PVP	0.9	1.3	-	-	-	0.9
SRP 1	-	-	0.2	0.2	0.2	-
Photoactivated bleach (1) (ppm)	15 ppm	27 ppm	-	-	20 ppm	20 ppm
Photoactivated bleach (2) (ppm)	15 ppm	-	-	-	-	-
Brightener 1	0.08	0.19	-	-	0.09	0.15
Brightener 2	-	0.04	-	-	-	-
Perfume	0.3	0.3	0.04	0.0	0.0	0.3
CAP4	2.0	1.0	4.0	-	-	0.1
CAP3	-	-	-	2.0	1.5	0.4
Silicone antifoam	0.5	2.4	0.3	0.5	0.3	2.0
Minors/misc to 100%						
Density in g/litre	750	750	750	750	750	750

Example 3

The following detergent formulations of particular utility under European machine wash conditions were prepared in accord with the invention.

5

	A	B	C	D
Blown powder				
LAS	6.0	5.0	11.0	6.0
TAS	2.0	-	-	2.0
Zeolite A	24.0	-	-	20.0
STPP	-	27.0	24.0	-
Sulfate	4.0	6.0	13.0	-
MA/AA	1.0	4.0	6.0	2.0
Silicate	1.0	7.0	3.0	3.0
CMC	1.0	1.0	0.5	0.6
Brightener 1	0.2	0.2	0.2	0.2
Silicone antifoam	1.0	1.0	1.0	0.3
DTPMP	0.4	0.4	0.2	0.4
Spray on				
Brightener	0.02	-	-	0.02
C45E7	-	-	-	5.0
C45E2	2.5	2.5	2.0	-
C45E3	2.6	2.5	2.0	-
Perfume	0.5	0.3	0.1	-
Silicone antifoam	0.3	0.3	0.3	-
Dry additives				
QEA	-	-	-	1.0
EDDS	0.3	-	-	-
Sulfate	2.0	3.0	5.0	10.0

Carbonate	6.0	13.0	11.0	14.0
Citric acid	2.5	-	-	2.0
QAS II	0.5	-	-	0.5
SKS-6	10.0	-	-	-
Percarbonate	18.5	-	-	-
PB4	-	18.0	10.0	21.5
TAED	2.0	2.0	-	2.0
NAC-OBS	3.0	2.0	4.0	-
Protease	1.0	1.0	1.0	1.0
Lipase	-	0.4	-	0.2
Lipase (1)	0.4	-	0.4	-
Amylase	0.2	0.2	0.2	0.4
Brightener 1	0.05	-	-	0.05
CAP3	1.2	1.5	2.0	0.1
Misc/minor to 100%				

Example 4

The following granular detergent formulations were prepared in accord with the invention.

5

	A	B	C	D	E	F
Blown powder						
LAS	23.0	8.0	7.0	9.0	7.0	7.0
TAS	-	-	-	-	1.0	-
C45AS	6.0	6.0	5.0	8.0	-	-
C45AES	-	1.0	1.0	1.0	-	-
C45E35	-	-	-	-	2.0	4.0
Zeolite A	10.0	18.0	14.0	12.0	10.0	10.0
MA/AA	-	0.5	-	-	-	2.0

MA/AA (1)	7.0	-	-	-	-	-
AA	-	3.0	3.0	2.0	3.0	3.0
Sulfate	5.0	6.3	14.3	11.0	15.0	19.3
Silicate	10.0	1.0	1.0	1.0	1.0	1.0
Carbonate	13.0	19.0	8.0	20.7	8.0	6.0
PEG 4000	0.4	1.5	1.5	1.0	1.0	1.0
DTPA	-	0.9	0.5	-	-	0.5
Brightener 2	0.3	0.2	0.3	-	0.1	0.3
Spray on						
C45E7	-	2.0	-	-	2.0	2.0
C25E9	3.0	-	-	-	-	-
C23E9	-	-	1.5	2.0	-	2.0
Perfume	0.3	0.3	0.3	2.0	0.03	-
Agglomerates						
C45AS	-	5.0	5.0	2.0	-	5.0
LAS	-	2.0	2.0	-	-	2.0
Zeolite A	-	7.5	7.5	8.0	-	7.5
Carbonate	-	4.0	4.0	5.0	-	4.0
PEG 4000	-	0.5	0.5	-	-	0.5
Misc (water etc)	-	2.0	2.0	2.0	-	2.0
Dry additives						
QAS (I)	-	-	-	-	1.0	-
Citric acid	-	-	-	-	2.0	-
PB4	-	-	-	-	12.0	1.0
PB1	4.0	1.0	3.0	2.0	-	-
Percarbonate	-	-	-	-	2.0	10.0
Carbonate	-	5.3	1.8	-	4.0	4.0
NOBS	4.0	-	6.0	-	-	0.6
Methyl cellulose	0.2	-	-	-	-	-

SKS-6	8.0	-	-	-	-	-
STS	-	-	2.0	-	1.0	-
Cumene sulfonic acid	-	1.0	-	-	-	2.0
Lipase	0.2	-	0.2	-	0.2	0.4
Cellulase	0.2	0.2	0.2	0.3	0.2	0.2
Amylase	0.2	-	0.1	-	0.2	-
Protease	0.5	0.5	0.5	0.3	0.5	0.5
PVPVI	-	-	-	-	0.5	0.1
PVP	-	-	-	-	0.5	-
PVNO	-	-	0.5	0.3	-	-
QEA	-	-	-	-	1.0	-
SRP1	0.2	0.5	0.3	-	0.2	-
CAP3	1.2	1.0	3.0	1.5	0.06	0.1
Silicone antifoam	0.2	0.4	0.2	0.4	0.1	-
Mg sulfate	-	-	0.2	-	0.2	-
Misc/minors to 100%						

	G	H	I	J
Blown powder				
Clay I or II	7.0	10.0	6.0	2.0
LAS	16.0	5.0	11.0	6.0
TAS	-	5.0	-	2.0
Zeolite A	-	20.0	-	10.0
STPP	24.0	-	14.0	-
Sulfate	-	2.0	-	-
MA/AA	-	2.0	1.0	1.0
Silicate	4.0	7.0	3.0	-

CMC	1.0	-	0.5	0.6
Brightener 1	0.2	0.2	0.2	0.2
Carbonate	10.0	10.0	20.0	-
DTPMP	0.4	0.4	0.2	-
Spray on				
Brightener 1	0.02	-	-	0.02
C45E7 or E9	-	-	2.0	1.0
C45E3 or E4	-	-	2.0	4.0
Perfume	0.5	-	0.5	0.2
Silicone antifoam	0.3	-	-	-
Dry additives				
Flocculating agent I or II	0.3	1.0	1.0	0.5
QEA	-	-	-	1.0
HEDP/ EDDS	0.3	-	-	-
Sulfate	2.0	-	-	-
Carbonate	20.0	13.0	15.0	24.0
Citric acid	2.5	-	-	2.0
QAS	-	-	0.5	0.5
NaSKS-6	3.5	-	-	5.0
Percarbonate	-	-	-	9.0
PB4	-	-	5.0	
NOBS	-	-	-	1.3
TAED	-	-	2.0	1.5
Protease	1.0	1.0	1.0	1.0
Lipase	-	0.4	-	0.2
Amylase	0.2	0.2	0.2	0.4
Brightener 2	0.05	-	-	0.05
Perfume	-	0.2	0.5	0.3

Speckle	1.2	0.5	2.0	-
CAP1	1.0	0.5	1.4	2.0
Misc/minor to 100%				

Example 5

5 The following nil bleach-containing detergent formulations of particular use in the washing of coloured clothing, according to the present invention were prepared:

	A	B	C
Blown Powder			
Zeolite A	15.0	15.0	-
Sulfate	0.0	5.0	-
LAS	3.0	3.0	-
DTPMP	0.4	0.5	-
CMC	0.4	0.4	-
MA/AA	4.0	4.0	-
Agglomerates			
C45AS	-	-	11.0
LAS	6.0	5.0	-
TAS	3.0	2.0	-
Silicate	4.0	4.0	-
Zeolite A	10.0	15.0	13.0
CMC	-	-	0.5
MA/AA	-	-	2.0
Carbonate	9.0	7.0	7.0
Spray On			
Perfume	0.3	0.3	0.5

C45E7	4.0	4.0	4.0
C25E3	2.0	2.0	2.0
Dry additives			
MA/AA	-	-	3.0
NaSKS-6	-	-	12.0
Citrate	10.0	-	8.0
Bicarbonate	7.0	3.0	5.0
Carbonate	6.0	-	7.0
PVPVI/PVNO	0.5	0.5	0.5
Alcalase	0.5	0.3	0.9
Lipase	0.4	0.4	0.4
Amylase	0.6	0.6	0.6
Cellulase	0.6	0.6	0.6
CAP2	3.0	2.0	0.45
Silicone antifoam	5.0	5.0	5.0
Sulfate	0.0	9.0	0.0
Misc/minors to 100%	100.0	100.0	100.0
Density (g/litre)	700	700	700

Example 6

The following granular detergent formulations were prepared in accord with the invention.

5

	A	B	C	D
Base granule				
Zeolite A	30.0	22.0	24.0	10.0
Sulfate	10.0	5.0	10.0	7.0
MA/AA	3.0	-	-	-
AA	-	1.6	2.0	-

MA/AA (1)	-	12.0	-	6.0
LAS	14.0	10.0	9.0	20.0
C45AS	8.0	7.0	9.0	7.0
C45AES	-	1.0	1.0	-
Silicate	-	1.0	0.5	10.0
Soap	-	2.0	-	-
Brightener 1	0.2	0.2	0.2	0.2
Carbonate	6.0	9.0	10.0	10.0
PEG 4000	-	1.0	1.5	-
DTPA	-	0.4	-	-
Spray on				
C25E9	-	-	-	5.0
C45E7	1.0	1.0	-	-
C23E9	-	1.0	2.5	-
Perfume	0.2	0.3	0.3	-
Dry additives				
Carbonate	5.0	5.0	15.0	6.0
PVPVI/PVNO	0.5	-	0.3	-
Protease	1.0	1.0	1.0	0.5
Lipase	0.4	-	-	0.4
Amylase	0.1	-	-	0.1
Cellulase	0.1	0.2	0.2	0.1
NOBS	-	4.0	-	4.5
PB1	1.0	5.0	1.5	6.0
Sulfate	4.0	5.0	-	5.0
SRPI	-	0.4	-	-
CAP2	5.0	2.0	0.4	0.1
CAP3	-	-	-	0.1
Sud suppressor	-	0.5	0.5	-

Misc/minor to 100%				
--------------------	--	--	--	--

Example 7

The following granular detergent compositions were prepared in accord with the invention.

5

	A	B	C
Blown powder			
Zeolite A	20.0	-	15.0
STPP	-	20.0	-
Sulphate	-	-	5.0
Carbonate	-	-	5.0
TAS	-	-	1.0
LAS	6.0	6.0	6.0
C68AS	2.0	2.0	-
Silicate	3.0	8.0	-
MA/AA	4.0	2.0	2.0
CMC	0.6	0.6	0.2
Brightener 1	0.2	0.2	0.1
DTPMP	0.4	0.4	0.1
STS	-	-	1.0
Spray on			
C45E7	5.0	5.0	4.0
Silicone antifoam	0.3	0.3	0.1
Perfume	0.2	0.2	0.3
Dry additives			
QEA	-	-	1.0
Carbonate	14.0	9.0	10.0
PB1	1.5	2.0	-

PB4	18.5	13.0	13.0
TAED	2.0	2.0	2.0
QAS (I)	-	-	1.0
Photoactivated bleach	15 ppm	15 ppm	15ppm
SKS-6	-	-	3.0
Protease	1.0	1.0	0.2
Lipase	0.2	0.2	0.2
Amylase	0.4	0.4	0.2
Cellulase	0.1	0.1	0.2
Sulfate	10.0	20.0	5.0
CAP1	1.2	2.0	0.5
Misc/minors to 100%			
Density (g/litre)	700	700	700

Example 8

The following detergent compositions, according to the present invention were prepared:

5

	A	B	C
Blown Powder			
Zeolite A	15.0	15.0	15.0
Sulfate	0.0	5.0	0.0
LAS	3.0	3.0	3.0
QAS	-	1.5	1.5
DTPMP	0.4	0.2	0.4
EDDS	-	0.4	0.2
CMC	0.4	0.4	0.4
MA/AA	4.0	2.0	2.0
Agglomerates			
LAS	5.0	5.0	5.0
TAS	2.0	2.0	1.0
Silicate	3.0	3.0	4.0
Zeolite A	8.0	8.0	8.0
Carbonate	8.0	8.0	4.0
Spray On			
Perfume	0.3	0.3	0.3
C45E7	2.0	2.0	2.0
C25E3	2.0	-	-
Dry additives			
Citrate	5.0	-	2.0
Bicarbonate	-	3.0	-

Carbonate	8.0	15.0	10.0
TAED	6.0	2.0	5.0
PB1	14.0	7.0	10.0
PEO	-	-	0.2
CAP1	1.2	1.0	0.75
Bentonite clay	-	-	10.0
Protease	1.0	1.0	1.0
Lipase	0.4	0.4	0.4
Amylase	0.6	0.6	0.6
Cellulase	0.6	0.6	0.6
Silicone antifoam	5.0	5.0	5.0
Sodium sulfate	0.0	3.0	0.0
Misc/minors to 100%	100.0	100.0	100.0
Density (g/litre)	850	850	850

	D	E	F	G	H
Blown Powder					
STPP/ Zeolite A	9.0	15.0	15.0	9.0	9.0
Flocculating agent II or III	0.5	0.2	0.9	1.5	-
LAS	7.5	23.0	3.0	7.5	7.5
QAS	2.5	1.5	-	-	-
DTPMP	0.4	0.2	0.4	0.4	0.4
HEDP or EDDS	-	0.4	0.2	-	-
CMC	0.1	0.4	0.4	0.1	0.1
Sodium carbonate	5.0	20.0	20.0	10.0	-
Brightener	0.05	-	-	0.05	0.05
Clay I or II	-	10.0	-	-	-
STS	0.5	-	-	0.5	0.5

MA/AA	1.5	2.0	2.0	1.5	1.5
Agglomerates					
Suds suppresser (silicon)	1.0	1.0	-	2.0	0.5
Agglomerate					
Clay	9.0	-	-	4.0	10.0
Wax	0.5	-	-	0.5	1.5
Glycerol	0.5	-	-	0.5	0.5
Agglomerate					
LAS	-	5.0	5.0	-	-
TAS	-	2.0	1.0	-	-
Silicate	-	3.0	4.0	-	-
Zeolite A	-	8.0	8.0	-	-
Carbonate	-	8.0	4.0	-	-
Spray On					
Perfume	0.3	-	-	0.3	0.3
C45E7 or E9	2.0	-	-	2.0	2.0
C25E3 or E4	2.0	-	-	2.0	2.0
Dry additives					
Citrate or citric acid	2.5	-	2.0	2.5	2.5
Clay I or II	-	5.0	5.0	-	-
Flocculating agent I or II	-	-	-	-	0.2
Bicarbonate	-	3.0	-	-	-
Carbonate	15.0	-	-	25.0	31.0
TAED	1.0	2.0	5.0	1.0	-
Sodium perborate or percarbonate	6.0	7.0	10.0	6.0	-
SRP1, 2, 3 or 4	0.2	0.1	0.2	0.5	0.3
CMC or nonionic cellulose ether	1.0	1.5	0.5	-	-

Protease	0.3	1.0	1.0	0.3	0.3
Lipase	-	0.4	0.4	-	-
Amylase	0.2	0.6	0.6	0.2	0.2
Cellulase	0.2	0.6	0.6	0.2	0.2
Silicone antifoam	-	5.0	5.0	-	-
Perfume (starch)	0.2	0.3	1.0	0.2	0.2
Speckle	0.5	0.5	0.1	-	1.0
NaSKS-6 (silicate 2R)	3.5	-	-	-	3.5
Photobleach	0.1	-	-	0.1	0.1
Soap	0.5	2.5	-	0.5	0.5
Sodium sulfate	-	3.0	-	-	-
CAP5	0.7	1.0	2.0	0.4	1.5
Misc/minors to 100%	100.0	100.0	100.0	100.0	100.0
Density (g/litre)	850	850	850	850	850

Example 9

The following detergent formulations, according to the present invention were prepared:

	A	B	C	D
LAS	18.0	14.0	24.0	20.0
QAS	0.7	1.0	-	0.7
TFAA	-	1.0	-	-
C23E56.5	-	-	1.0	-
C45E7	-	1.0	-	-
C45E3S	1.0	2.5	1.0	-
STPP	32.0	18.0	30.0	22.0
Silicate	9.0	5.0	9.0	8.0
Carbonate	9.0	7.5	-	5.0
Bicarbonate	-	7.5	-	-
PB1	3.0	1.0	-	-
PB4	-	1.0	-	-
NOBS	2.0	1.0	-	-
DTPMP	-	1.0	-	-
DTPA	0.5	-	0.2	0.3
SRP 1	0.3	0.2	-	0.1
MA/AA	1.0	1.5	2.0	0.5
CMC	0.8	0.4	0.4	0.2
PEI	-	-	0.4	-
Sodium sulfate	20.0	10.0	20.0	30.0
Mg sulfate	0.2	-	0.4	0.9
Protease	0.8	1.0	0.5	0.5
Amylase	0.5	0.4	-	0.25

Lipase	0.2	-	0.1	-
Cellulase	0.15	-	-	0.05
Photoactivated bleach (ppm)	30ppm	20ppm	-	10ppm
CAP2	2.0	1	0.8	2
Perfume spray on	0.3	0.3	0.1	-
Brightener 1/2	0.05	0.2	0.08	0.1
Misc/minors to 100%				

Example 10

The following is a composition in the form of a tablet, bar, extrudate or granule in accord with the invention

5

		A	B	C	D	E	F	G
Sodium alkylbenzenesulfonate	C ₁₁ -C ₁₃	12.0	16.0	23.0	19.0	18.0	20.0	16.0
Sodium sulfate	C ₁₄ -C ₁₅ alcohol		4.5	-		-	-	4.0
C ₁₄ -C ₁₅ alcohol sulfate	ethoxylate (3)	-	-	2.0	-	1.0	1.0	1.0
Sodium ethoxylate	C ₁₄ -C ₁₅ alcohol	2.0	2.0	-	1.3	-	-	5.0
C ₉ -C ₁₄ alkyl dimethyl hydroxy ethyl quaternary ammonium salt				-	-	1.0	0.5	2.0
Tallow fatty acid				-	-	-	-	1.0
Sodium Zeolite	tripolyphosphate /	23.0	25.0	14.0	22.0	20.0	10.0	20.0
Sodium carbonate		25.0	22.0	35.0	20.0	28.0	41.0	30.0

Sodium Polyacrylate (45%)	0.5	0.5	0.5	0.5	-	-	-
Sodium polyacrylate/maleate polymer	-	-	1.0	1.0	1.0	2.0	0.5
Sodium silicate (1:6 ratio NaO/SiO ₂)(46%)	3.0	6.0	9.0	8.0	9.0	6.0	8.0
Sodium sulfate	-	-	-	-	-	2.0	3.0
Sodium perborate/ percarbonate	5.0	5.0	10.0	-	3.0	1.0	-
Poly(ethyleneglycol), MW ~4000 (50%)	1.5	1.5	1.0	1.0	-	-	0.5
Sodium carboxy methyl cellulose	1.0	1.0	1.0	-	0.5	0.5	0.5
NOBS/ DOBS	-	1.0	-	-	1.0	0.7	-
TAED	1.5	1.0	2.5	-	3.0	0.7	-
SRP 1	1.5	1.5	1.0	1.0	-	1.0	-
Clay I or II	5.0	6.0	12.0	7.0	10.0	4.0	3.0
Flocculating agent I or III	0.2	0.2	3.0	2.0	0.1	1.0	0.5
Humectant	0.5	1.0	0.5	1.0	0.5	0.5	-
Wax	0.5	0.5	1.0	-	-	0.5	0.5
Moisture	7.5	7.5	6.0	7.0	5.0	3.0	5.0
Magnesium sulphate	-	-	-	-	-	0.5	1.5
Chelant	-	-	-	-	0.8	0.6	1.0
Enzymes, including amylase, cellulase, protease and lipase	-	-	-	-	2.0	1.5	2.0
Speckle	2.5	4.1	4.2	4.4	5.6	5.0	5.2
minors, e.g. perfume, PVP, PVPVI/PVNO, brightener, photo-bleach,	2.0	1.0	1.0	1.0	2.5	1.5	1.0
CAP2	1.6	2.0	0.4	2.0	1.0	1.6	0.5

	H	I	J	K
Sodium C ₁₁ -C ₁₃ alkylbenzenesulfonate	23.0	13.0	20.0	18.0
Sodium C ₁₄ -C ₁₅ alcohol sulfate	-	4.0	-	-
Clay I or II	5.0	10.0	14.0	6.0
Flocculating agent I or II	0.2	0.3	0.1	0.9
Wax	0.5	0.5	1.0	-
Humectant (glycerol/ silica)	0.5	2.0	1.5	-
C ₁₄ -C ₁₅ alcohol ethoxylate sulfate	-		-	2.0
Sodium C ₁₄ -C ₁₅ alcohol ethoxylate (2.5	3.5	-	-
C ₉ -C ₁₄ alkyl dimethyl hydroxy ethyl quaternary ammonium salt		-	-	0.5
Tallow fatty acid	0.5	-	-	-
Tallow alcohol ethoxylate (50)	-	-		1.3
Sodium tripolyphosphate	-	41.0	-	20.0
Zeolite A, hydrate (0.1-10 micrometer size)	26.3	-	21.3	-
Sodium carbonate	24.0	22.0	35.0	27.0
Sodium Polyacrylate (45 %)	2.4	-	2.7	-
Sodium polyacrylate/maleate polymer	-	-	1.0	2.5
Sodium silicate (1.6 or 2 or 2.2 ratio NaO/SiO ₂)(46%)	4.0	7.0	2.0	6.0
Sodium sulfate	-	6.0	2.0	-
Sodium perborate/ percarbonate	8.0	4.0	-	12.0
Poly(ethyleneglycol), MW ~4000 (50%)	1.7	0.4	1.0	-
Sodium carboxy methyl cellulose	1.0	-	-	0.3
Citric acid	-	-	3.0	-
NOBS/ DOBS	1.2	-	-	1.0
TAED	0.6	1.5	-	3.0
Perfume	0.5	1.0	0.3	0.4

SRP 1	-	1.5	1.0	1.0
Moisture	7.5	3.1	6.1	7.3
Magnesium sulphate	-	-	-	1.0
Chelant	-	-	-	0.5
speckle	1.0	0.5	0.2	2.7
Enzymes, including amylase, cellulase, protease and lipase	-	1.0	-	1.5
minors, e.g. brightener, photo-bleach	1.0	1.0	1.0	1.0
CAP2	1.2	0.4	1.6	2.0

Example 11

The following liquid detergent formulations were prepared in accord with the invention (levels are given as parts per weight).

5

	A	B	C	D	E
LAS	11.5	8.8	-	3.9	-
C25E2.5S	-	3.0	18.0	-	16.0
C45E2.25S	11.5	3.0	-	15.7	-
C23E9	-	2.7	1.8	2.0	1.0
C23E7	3.2	-	-	-	-
CFAA	-	-	5.2	-	3.1
TPKFA	1.6	-	2.0	0.5	2.0
Citric acid (50%)	6.5	1.2	2.5	4.4	2.5
Calcium formate	0.1	0.06	0.1	-	-
Sodium formate	0.5	0.06	0.1	0.05	0.05
Sodium cumene sulfonate	4.0	1.0	3.0	1.18	-
Borate	0.6	-	3.0	2.0	2.9
Sodium hydroxide	5.8	2.0	3.5	3.7	2.7

Ethanol	1.75	1.0	3.6	4.2	2.9
1, 2 propanediol	3.3	2.0	8.0	7.9	5.3
Monoethanolamine	3.0	1.5	1.3	2.5	0.8
TEPAE	1.6	-	1.3	1.2	1.2
Protease	1.0	0.3	1.0	0.5	0.7
Lipase	-	-	0.1	-	-
Cellulase	-	-	0.1	0.2	0.05
Amylase	-	-	-	0.1	-
SRP1	0.2	-	0.1	-	-
DTPA	-	-	0.3	-	-
PVNO	-	-	0.3	-	0.2
CAP1	2.0	-	0.1	-	-
CAP6	-	0.4	-	-	-
CAP7	-	-	0.2	-	0.1
CAP8	-	-	-	0.5	-
Brightener 1	0.2	0.07	0.1	-	-
Silicone antifoam	0.04	0.02	0.1	0.1	0.1
Water/minors up to 100%					

Example 12

The following liquid detergent formulations were prepared in accord with the invention (levels are given in parts per weight):

	A	B	C	D	E	F	G	H
LAS	10.0	13.0	9.0	-	25.0	-	-	-
C25AS	4.0	1.0	2.0	10.0	-	13.0	18.0	15.0
C25E3S	1.0	-	-	3.0	-	2.0	2.0	4.0
C25E7	6.0	8.0	13.0	2.5	-	-	4.0	4.0

TFAA	-	-	-	4.5	-	6.0	8.0	8.0
APA	-	1.4	-	-	3.0	1.0	2.0	-
TPKFA	2.0	-	13.0	7.0	-	15.0	11.0	11.0
Citric acid	2.0	3.0	1.0	1.5	1.0	1.0	1.0	1.0
Dodecetyl/tetradecenyl succinic acid	12.0	10.0	-	-	15.0	-	-	-
Rape seed fatty acid	4.0	2.0	1.0	-	1.0	-	3.5	-
Ethanol	4.0	4.0	7.0	2.0	7.0	2.0	3.0	2.0
1,2 Propanediol	4.0	4.0	2.0	7.0	6.0	8.0	10.0	13.0
Monoethanolamine	-	-	-	5.0	-	-	9.0	9.0
Triethanolamine	-	-	8.0	-	-	-	-	-
TEPAE	0.5	-	0.5	0.2	-	-	0.4	0.3
DTPMP	1.0	1.0	0.5	1.0	2.0	1.2	1.0	-
Protease	0.5	0.5	0.4	0.25	-	0.5	0.3	0.6
Alcalase	-	-	-	-	1.5	-	-	-
Lipase	-	0.10	-	0.01	-	-	0.15	0.15
Amylase	0.25	0.25	0.6	0.5	0.25	0.9	0.6	0.6
Cellulase	-	-	-	0.05	-	-	0.15	0.15
Endolase	-	-	-	0.10	-	-	0.07	-
SRP2	0.3	-	0.3	0.1	-	-	0.2	0.1
Boric acid	0.1	0.2	1.0	2.0	1.0	1.5	2.5	2.5
Calcium chloride	-	0.02	-	0.01	-	-	-	-
Bentonite clay	-	-	-	-	4.0	4.0	-	-
Brightener 1	-	0.4	-	-	0.1	0.2	0.3	-
Sud suppressor	0.1	0.3	-	0.1	0.4	-	-	-
Opacifier	0.5	0.4	-	0.3	0.8	0.7	-	-
CAP6	0.2	0.1	0.05	0.1	3.3	-	-	-
CAP7	-	-	-	0.1	-	0.1	0.2	0.05

Water/minors up to 100%								
NaOH up to pH	8.0	8.0	7.6	7.7	8.0	7.5	8.0	8.2

Example 13

The following liquid detergent compositions were prepared in accord with the invention (levels are given in parts per weight).

5

	A	B
LAS	27.6	18.9
C45AS	13.8	5.9
C13E8	3.0	3.1
Oleic acid	3.4	2.5
Citric acid	5.4	5.4
Sodium hydroxide	0.4	3.6
Calcium formate	0.2	0.1
Sodium formate	-	0.5
Ethanol	7.0	-
Monoethanolamine	16.5	8.0
1,2 propanediol	5.9	5.5
Xylene sulfonic acid	-	2.4
TEPAE	1.5	0.8
Protease	1.5	0.6
PEG	-	0.7
Brightener 2	0.4	0.1
Perfume	0.5	0.3
CAP6	0.2	-
CAP8		
Silicone-based suspensing agent	2.0	-

Water/minors up to 100%		
-------------------------	--	--

Example 14

The following laundry bar detergent compositions were prepared in accord with the invention (levels are given in parts per weight).

5

	A	B	C	D	E	F	G	H
LAS	-	-	19.0	15.0	21.0	6.75	8.8	-
C28AS	30.0	13.5	-	-	-	15.75	11.2	22.5
Sodium laurate	2.5	9.0	-	-	-	-	-	-
Zeolite A	2.0	1.25	-	-	-	1.25	1.25	1.25
Carbonate	10.0	-	11.0	5.0	2.0	7.0	13.0	9.0
Calcium carbonate	27.5	39.0	35.0	-	-	40.0	-	40.0
Sulfate	5.0	5.0	3.0	5.0	3.0	-	-	5.0
TSPP	5.0	-	-	-	-	5.0	2.5	-
STPP	5.0	15.0	10.0	-	-	7.0	8.0	10.0
Bentonite clay	-	10.0	-	-	5.0	-	-	-
DTPMP	-	0.7	0.6	-	0.6	0.7	0.7	0.7
CMC	-	1.0	1.0	1.0	1.0	-	-	1.0
Talc	-	-	10.0	15.0	10.0	-	-	-
Silicate	-	-	4.0	5.0	3.0	-	-	-
PVNO	0.02	0.03	-	0.01	-	0.02	-	-
MA/AA	0.4	1.0	-	-	0.2	0.4	0.5	0.4
SRP1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Protease	-	0.12	-	0.08	0.08	-	-	0.1
Lipase	-	0.1	-	0.1	-	-	-	-
Amylase	-	-	0.8	-	-	-	0.1	-
Cellulase	-	0.15	-	-	0.15	0.1	-	-

PEO	-	0.2	-	0.2	0.3	-	-	0.3
Perfume	1.0	0.5	0.3	0.2	0.4	-	-	0.4
Mg sulfate	-	-	3.0	3.0	3.0	-	-	-
CAP1	3.0	1.4	0.8	0.4	0.1	2.0	2.0	1.0
CAP2	-	1.4	-	-	-	-	2.0	-
CAP3	-	-	0.8	-	-	-	-	1.0
CAP4	-	-	-	0.4	0.1	-	-	0.05
Brightener	0.15	0.10	0.15	-	-	-	-	0.1
Photoactivated bleach (ppm)	-	15.0	15.0	15.0	15.0	-	-	15.0

Example 15

The following detergent additive compositions were prepared according to the present invention :

5

	A	B	C
LAS	-	5.0	5.0
STPP	30.0	-	20.0
Zeolite A	-	35.0	20.0
PB1	20.0	15.0	-
TAED	10.0	8.0	-
CAP1	3.1	0.4	1.1
CAP2	-	0.4	0.2
Protease	-	0.3	0.3
Amylase	-	0.06	0.06
Minors, water and miscellaneous			Up to 100%

Example 16

The following compact high density (0.96Kg/l) dishwashing detergent compositions were prepared according to the present invention :

	A	B	C	D	E	F	G	H
STPP	-	-	54.3	51.4	51.4	-	-	50.9
Citrate	35.0	17.0	-	-	-	46.1	40.2	-
Carbonate	-	15.0	12.0	14.0	4.0	-	7.0	31.1
Bicarbonate	-	-	-	-	-	25.4	-	-
Silicate	32.0	14.8	14.8	10.0	10.0	1.0	25.0	3.1
Metasilicate	-	2.5	-	9.0	9.0	-	-	-
PB1	1.9	9.7	7.8	7.8	7.8	-	-	-
PB4	8.6	-	-	-	-	-	-	-
Percarbonate	-	-	-	-	-	6.7	11.8	4.8
Nonionic	1.5	2.0	1.5	1.7	1.5	2.6	1.9	5.3
TAED	5.2	2.4	-	-	-	2.2	-	1.4
HEDP	-	1.0	-	-	-	-	-	-
DTPMP	-	0.6	-	-	-	-	-	-
MnTACN	-	-	-	-	-	-	0.008	-
PAAC	-	-	0.00	0.01	0.00	-	-	-
			8		7			
BzP	-	-	-	-	1.4	-	-	-
Paraffin	0.5	0.5	0.5	0.5	0.5	0.6	-	-
CAP1	1.2	1.4	1.2	1.1	-	0.1	-	0.5
CAP2	-	-	-	-	2.1	2.3	4.2	-
Protease	0.07	0.07	0.02	0.053	0.04	0.026	0.059	0.06
	2	2	9		6			
Amylase	0.01	0.01	0.00	0.012	0.01	0.009	0.017	0.03
	2	2	6		3			

Lipase	-	0.00	-	0.005	-	-	-	-	-
		1							
BTA	0.3	0.3	0.3	0.3	0.3	-	0.3	0.3	
MA/AA	-	-	-	-	-	-	4.2	-	
480N	3.3	6.0	-	-	-	-	-	0.9	
Perfume	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	
Sulphate	7.0	20.0	5.0	2.2	0.8	12.0	4.6	-	
pH	10.8	11.0	10.8	11.3	11.3	9.6	10.8	10.9	
Miscellaneous and water						Up to 100%			

Example 17

The following granular dishwashing detergent compositions of bulk density 1.02Kg/L were prepared according to the present invention :

	A	B	C	D	E	F	G	H
STPP	30.0	30.0	33.0	34.2	29.6	31.1	26.6	17.6
Carbonate	29.5	30.0	29.0	24.0	15.0	36.0	2.1	38.0
Silicate	7.4	7.4	7.5	7.2	13.3	3.4	43.7	12.4
Metasilicate	-	-	4.5	5.1	-	-	-	-
Percarbonate	-	-	-	-	-	4.0	-	-
PB1	4.4	4.2	4.5	4.5	-	-	-	-
NADCC	-	-	-	-	2.0	-	1.6	1.0
Nonionic	1.2	1.0	0.7	0.8	1.9	0.7	0.6	0.3
TAED	1.0	-	-	-	-	0.8	-	-
PAAC	-	0.004	0.004	0.004	-	-	-	-
BzP	-	-	-	1.4	-	-	-	-
Paraffin	0.25	0.25	0.25	0.25	-	-	-	-
CAP1	1.0	0.5	1.4	1.8	0.1	-	-	-
CAP2	-	-	-	-	0.1	0.15	0.2	0.1
Protease	0.036	0.015	0.03	0.028	-	0.03	-	-
Amylase	0.003	0.003	0.01	0.006	-	0.01	-	-
Lipase	0.005	-	0.001	-	-	-	-	-
BTA	0.15	0.15	0.15	0.15	-	-	-	-
Perfume	0.2	0.2	0.2	0.2	0.1	0.2	0.2	-
Sulphate	23.4	25.0	22.0	18.5	30.1	19.3	23.1	23.6
pH	10.8	10.8	11.3	11.3	10.7	11.5	12.7	10.9
5 Miscellaneous and water					Up to 100%			

Example 18

The following tablet detergent compositions were prepared according to the present invention by compression of a granular dishwashing detergent composition at a pressure of 13KN/cm² using a standard 12 head rotary press:

	A	B	C	D	E	F
STPP	-	48.8	49.2	38.0	-	46.8
Citrate	26.4	-	-	-	31.1	-
Carbonate	-	4.0	12.0	14.4	10.0	20.0
Silicate	26.4	14.8	15.0	12.6	17.7	2.4
CAP1	2.3	-	-	-	2.5	0.2
CAP2	-	0.8	-	-	-	0.2
CAP3	-	-	1.2	1	-	-
Protease	0.058	0.072	0.041	0.033	0.052	0.013
Amylase	0.01	0.03	0.012	0.007	0.016	0.002
Lipase	0.005	-	-	-	-	-
PB1	1.6	7.7	12.2	10.6	15.7	-
PB4	6.9	-	-	-	-	14.4
Nonionic	1.5	2.0	1.5	1.65	0.8	6.3
PAAC	-	-	0.02	0.009	-	-
MnTACN	-	-	-	-	0.007	-
TAED	4.3	2.5	-	-	1.3	1.8
HEDP	0.7	-	-	0.7	-	0.4
DTPMP	0.65	-	-	-	-	-
Paraffin	0.4	0.5	0.5	0.55	-	-
BTA	0.2	0.3	0.3	0.3	-	-
PA30	3.2	-	-	-	-	-
MA/AA	-	-	-	-	4.5	0.55
Perfume	-	-	0.05	0.05	0.2	0.2
Sulphate	24.0	13.0	2.3	-	10.7	3.4

Weight of tablet	25g	25g	20g	30g	18g	20g
pH	10.6	10.6	10.7	10.7	10.9	11.2
Miscellaneous and water	Up to 100%					

Example 19

5 The following liquid dishwashing detergent compositions of density 1.40Kg/L were prepared according to the present invention :

	A	B	C	D
STPP	17.5	17.5	17.2	16.0
Carbonate	2.0	-	2.4	-
Silicate	5.3	6.1	14.6	15.7
NaOCl	1.15	1.15	1.15	1.25
Polygen/carbopol	1.1	1.0	1.1	1.25
Nonionic	-	-	0.1	-
NaBz	0.75	0.75	-	-
CAP6	0.4	0.8	0.1	0.5
NaOH	-	1.9	-	3.5
KOH	2.8	3.5	3.0	-
pH	11.0	11.7	10.9	11.0
Sulphate, miscellaneous and water	up to 100%			

Example 20

10 The following liquid rinse aid compositions were prepared according to the present invention :

	A	B	C
Nonionic	12.0	-	14.5
Nonionic blend	-	64.0	-

Citric	3.2	-	6.5	
HEDP	0.5	-	-	
PEG	-	5.0	-	
SCS	4.8	-	7.0	
Ethanol	6.0	8.0	-	
CAP7	3	-	1	
CAP8	3.0	0.2	0.1	
pH of the liquid	2.0	7.5	/	
Miscellaneous and water			Up to 100%	

Example 21

The following liquid dishwashing compositions were prepared according to the present
5 invention :

	A	B	C	D	E
C17ES	28.5	27.4	19.2	34.1	34.1
Amine oxide	2.6	5.0	2.0	3.0	3.0
C12 glucose amide	-	-	6.0	-	-
Betaine	0.9	-	-	2.0	2.0
Xylene sulfonate	2.0	4.0	-	2.0	-
Neodol C11E9	-	-	5.0	-	-
Polyhydroxy fatty acid amide	-	-	-	6.5	6.5
Sodium diethylene penta acetate (40%)	-	-	0.03	-	-
TAED	-	-	-	0.06	0.06
Sucrose	-	-	-	1.5	1.5
Ethanol	4.0	5.5	5.5	9.1	9.1
Alkyl diphenyl oxide disulfonate	-	-	-	-	2.3

Ca formate	-	-	-	0.5	1.1
Ammonium citrate	0.06	0.1	-	-	-
Na chloride	-	1.0	-	-	-
Mg chloride	3.3	-	0.7	-	-
Ca chloride	-	-	0.4	-	-
Na sulfate	-	-	0.06	-	-
Mg sulfate	0.08	-	-	-	-
Mg hydroxide	-	-	-	2.2	2.2
Na hydroxide	-	-	-	1.1	1.1
Hydrogen peroxide	200pp	0.16	0.006	-	-
		m			
CAP6	0.4	1.6	1.2	-	0.1
CAP7	-	-	-	1	1
Protease	0.017	0.005	.0035	0.003	0.002
Perfume	0.18	0.09	0.09	0.2	0.2
Water and minors				Up to 100%	

Example 22

The following liquid hard surface cleaning compositions were prepared according to the

5 present invention :

	A	B	C	D	E
CAP6	2.8	-	1.6	1.0	0.4
CAP7	-	1.2	-	1.0	0.5
Amylase	0.01	0.002	0.005	-	-
Protease	0.05	0.01	0.02	-	-
Hydrogen peroxide	-	-	-	6.0	6.8
Acetyl triethyl citrate	-	-	-	2.5	-
DTPA	-	-	-	0.2	-
Butyl hydroxy toluene	-	-	-	0.05	-

EDTA*	0.05	0.05	0.05	-	-
Citric / Citrate	2.9	2.9	2.9	1.0	-
LAS	0.5	0.5	0.5	-	-
C12 AS	0.5	0.5	0.5	-	-
C10AS	-	-	-	-	1.7
C12(E)S	0.5	0.5	0.5	-	-
C12,13 E6.5 nonionic	7.0	7.0	7.0	-	-
Neodol 23-6.5	-	-	-	12.0	-
Dobanol 23-3	-	-	-	-	1.5
Dobanol 91-10	-	-	-	-	1.6
C25AE1.8S	-	-	-	6.0	-
Na paraffin sulphonate	-	-	-	6.0	-
Perfume	1.0	1.0	1.0	0.5	0.2
Propanediol	-	-	-	1.5	-
Ethoxylated tetraethylene pentaimine	-	-	-	1.0	-
2, Butyl octanol	-	-	-	-	0.5
Hexyl carbitol**	1.0	1.0	1.0	-	-
SCS	1.3	1.3	1.3	-	-
pH adjusted to	7-12	7-12	7-12	4	-
Miscellaneous and water				Up to 100%	

*Na4 ethylenediamine diacetic acid

**Diethylene glycol monohexyl ether

5 Example 23

The following spray composition for cleaning of hard surfaces and removing household mildew was prepared according to the present invention :

CAP6	1
Amylase	0.01

100

Protease	0.01
Na octyl sulfate	2.0
Na dodecyl sulfate	4.0
Na hydroxide	0.8
Silicate	0.04
Butyl carbitol*	4.0
Perfume	0.35
Water/minors	up to 100%

*Diethylene glycol monobutyl ether

Example 24

5 The following lavatory cleansing block compositions were prepared according to the present invention.

	A	B	C
C16-18 fatty alcohol/50EO	70.0	-	-
LAS	-	-	80.0
Nonionic	-	1.0	-
Oleoamide surfactant	-	25.0	-
Partially esterified copolymer of vinylmethyl ether and maleic anhydride, viscosity 0.1-0.5	5.0	-	-
Polyethylene glycol MW 8000	-	38.0	-
Water-soluble K-polyacrylate MW 4000-8000	-	12.0	-
Water-soluble Na-copolymer of acrylamide (70%) and acryclic acid (30%) low MW	-	19.0	-
Na triphosphate	10.0	-	-
Carbonate	-	-	-
CAP6	1.0	1.2	-
CAP7	-	-	0.5

101

Dye	2.5	1.0	1.0
Perfume	3.0	-	7.0
KOH / HCL solution	pH 6-11		

Example 25

The following toilet bowl cleaning composition was prepared according to the present invention.

5

	A	B
C14-15 linear alcohol 7EO	2.0	10.0
Citric acid	10.0	5.0
CAP1	2.0	-
CAP7	2.0	4.0
DTPMP	-	1.0
Dye	2.0	1.0
Perfume	3.0	3.0
NaOH	pH 6-11	
Water and minors	Up to 100%	

Example 26

10 The following fabric softening compositions are in accordance with the present invention

Component	A	B	C	D	E	F
DTDMAC	-	-	-	-	4.5	15.0
DEQA	2.6	2.9	18.0	19.0	-	-
Fatty acid	0.3	-	1.0	-	-	-
HCl	0.02	0.02	0.02	0.02	0.02	0.02
PEG	-	-	0.6	0.6	-	0.6
Perfume	1.0	1.0	1.0	1.0	1.0	1.0

Silicone antifoam	0.01	0.01	0.01	0.01	0.01	0.01
CAP 6	0.4	0.1	0.8	0.2	1.0	0.6
Electrolyte (ppm)	-	-	600	1200	-	1200
Silicone suspending agent	1.0	0.5				
Dye (ppm)	10	10	50	50	10	50
Water and minors to balance to 100%						

Example 27

The following dryer added fabric conditioner compositions were prepared according to the present invention :

5

	A	B	C	D
DEQA(2)	-	-	-	50.0
DTMAMS	-	-	26.0	-
SDASA	70.0	70.0	42.0	35.0
Neodol 45-13	13.0	13.0	-	-
Ethanol	1.0	1.0	-	-
CAP 6	1.5	-	1.5	3.0
CAP 7	1.5	0.2	5.0	1.0
Perfume	0.75	0.75	1.0	1.5
Glycoperse S-20	-	-	-	10.0
Glycerol monostearate	-	-	26.0	-
Digeranyl Succinate	0.38	0.38	-	-
Clay	-	-	3.0	-
Dye	0.01	0.01	-	-
Minors to balance to 100%				

Example 28

The following are non-limiting examples of pre-soak fabric conditioning and/or fabric enhancement compositions according to the present invention which can be suitably used in the laundry rinse cycle.

Ingredients	A	B	C	D	E	F
Polymer	3.5	3.5	3.5	3.5	3.5	3.5
Dye fixative	2.3	2.3	2.4	2.4	2.5	2.5
Polyamine	15.0	15.0	17.5	17.5	20.0	20.0
Bayhibit AM	1.0	1.0	1.0	1.0	1.0	1.0
C ₁₂ -C ₁₄ dimethyl hydroxyethyl quaternary ammonium chloride	-	5.0	5.0	-	-	-
Fabric softener active	-	-	2.5	2.5	-	-
Genamin C100	0.33	-	0.33	0.33	0.33	-
Genapol V4463	0.2	-	0.2	0.2	0.2	-
CAP6	2.0	4.0	0.2	1.0	0.1	0.16
Water & minors	balance	balance	balance	balance	balance	balance

5

Example 29

The following are non-limiting examples of odor-absorbing compositions suitable for spray-on applications:

Examples	A	B	C	D	E
Ingredients	Wt. %				
HPBCD	1.0	-	1.0	-	1.2

RAMEB	-	1.0	-	0.8	-
Tetronic 901	-	-	0.1	-	-
Silwet L-7604	-	-	-	0.1	-
Silwet L-7600	0.1	-	-	-	0.1
Bardac 2050	-	-	-	0.03	-
Bardac 2250	-	0.2	-	-	0.1
Diethylene glycol	-	1.0	-	-	0.2
Triethylene glycol	-	-	0.1	-	-
Ethanol	-	-	-	-	2.5
Perfume 1	0.1	-	-	-	-
Perfume 2	-	0.05	-	0.1	-
Perfume 3	-	-	0.1	-	0.1
Kathon	3 ppm	3 ppm	3 ppm	3 ppm	-
HCl	to pH 4.5	to pH 4.5	to pH 3.5	to pH 3.5	to pH 3.5
CAP6	5.0	2.0	1.0	0.2	0.16
Distilled water	Bal.	Bal.	Bal.	Bal.	Bal.

The perfume 1, 2, and 3 have the following compositions:

Perfume	1	2	3
Perfume Ingredients	Wt. %	Wt. %	Wt. %
Anisic aldehyde	-	-	2
Benzophenone	3	5	-
Benzyl acetate	10	15	5
Benzyl salicylate	5	20	5
Cedrol	2	-	-

Citronellol	10	-	5
Coumarin	-	-	5
Cymal	-	-	3
Dihydromyrcenol	10	-	5
Flor acetate	5	-	5
Galaxolide	10	-	-
Lilial	10	15	20
Linalyl acetate	4	-	5
Linalool	6	15	5
Methyl dihydro jasmonate	3	10	5
Phenyl ethyl acetate	2	5	1
Phenyl ethyl alcohol	15	15	20
alpha-Terpineol	5	-	8
Vanillin	-	-	1
Total	100	100	100

What is claimed is:

1. A laundry and/or cleaning and/or fabric care composition comprising a detergent and/or cleaning and/or surfactant and/or fabric care ingredient and a benefit agent, said benefit agent being carried with a carrier, characterised in that the carried benefit agent has a viscosity of at least 400 cps at 20°C.
2. A composition according to Claim 1, wherein the carried composition which is incorporated into the laundry and/or cleaning and/or fabric care compositions provides a dry surface Odor Index of more than 5, preferably at least 10.
3. A composition according to either one of Claim 1 or 2, wherein the carrier is selected from a liquid carrier, solid carrier, and mixtures thereof, preferably liquid carrier.
4. A composition according to any one of Claim 1-3, wherein the carrier is selected from polymers which have chemically reacted with a benefit agent, components which have chemically reacted with a benefit agent, polymers which are not capable of chemically reacting with a benefit agent, and mixtures thereof.
5. A composition according to any one of Claims 1-4, wherein the carrier is water-insoluble.
6. A composition according to any one of Claim 1-5, wherein the carrier is polymer which is reacted with a component selected from acyl halides, palmytoyl chloride, myristoyl chloride, acid anhydrides alkyl halides, arylhalides, aldehydes, ketones, anhydride, carboxylic acid, and mixtures thereof, preferably is a polymer which is reacted with a component selected from aldehydes, ketones, and mixtures thereof.

7. A composition according to Claim 6, wherein the aldehyde and/or ketone are perfume aldehyde and/or ketones, preferably selected from undecylenic aldehyde, undecalactone gamma, heliotropin, dodecalactone gamma, p-anisic aldehyde, para hydroxy-phenyl-butanone, cymal, benzyl acetone, ionone alpha, p.t.bucinal, damascenone, ionone beta, methyl-nonyl ketone, lyral, dihydro iso jasmonate, citral, 1-decanal, benzaldehyde, florhydral, 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde; cis/trans-3,7-dimethyl-2,6-octadien-1-al; heliotropin; 2,4,6-trimethyl-3-cyclohexene-1-carboxaldehyde; 2,6-nonadienal; alpha-n-amyl cinnamic aldehyde, alpha-n-hexyl cinnamic aldehyde, P.T. Bucinal, lyral, cymal, methyl nonyl acetaldehyde, trans-2-nonenal, lilial, trans-2-nonenal, lauric aldehyde, undecylenic aldehyde, mefloral, phenylacetaldehyde, Alpha Damascone, Delta Damascone, Iso Damascone, Carvone, Gamma-Methyl-Ionone, Iso-E-Super, 2,4,4,7-Tetramethyl-oct-6-en-3-one, Benzyl Acetone, Beta Damascone, Damascenone, methyl dihydrojasmonate, methyl cedrylone, hedione and mixtures thereof.

8. A composition according to any one of Claims 4-6, wherein the polymers or components which have chemically reacted with a benefit agent is an amino-functional component, preferably selected from ethyl-4-amino benzoate, polyethyleneimine polymers; diaminobutane dendrimers Astramol®, polylysine, cross-linked polylysine, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched; 1,4-bis-(3-aminopropyl) piperazine, and mixtures thereof.

9 .A composition according to any one of Claims 4 or 5, wherein the polymers which are not capable of chemically reacting with a benefit agent are polyisobutylene polymers.

10. A composition according to any one of Claims 1-8, wherein the carrier of the carried composition has a molecular weight ranging from 2.000 to 10.000.000.

11. A composition according to any one of Claims 1-10, wherein the benefit agent comprises or is a perfume composition, preferably comprising at least 5%, preferably 10% by weight of perfume ingredient with an ODT of less than 1ppm, and whereby preferably the ratio of the benefit agent to the carrier is from 10:1 to 1:10, or even from 1:3 to 3:1.
12. A composition according to any one of claims 1-10, wherein the benefit agent is a perfume composition, preferably comprising at least 10%, preferably 25%, by weight of perfume ingredient with a Clog P of at least 2.0, preferably of at least 3.0, and boiling point of at least 250°C.
13. A composition according to any one of claims 1-10, wherein the benefit agent is a perfume composition, preferably comprising at least 20%, preferably 35%, by weight of perfume ingredient with a Clog P at least 2.0, preferably of at least 3.0, and boiling point of less than or equal to 250°C.
14. A composition to any of Claims 1 to 13 whereby the benefit agent comprises or is a biocide, preferably an antimicrobial, insect-repellent or a pharmaceutical agent, and whereby preferably the ratio of the benefit agent to the carrier is from 10:1 to 1:10, or even from 1:3 to 3:1.
15. A composition according to any one of Claims 1-14, wherein the carried benefit agent is further processed to result in an agglomerate having a particle size of from 150 to 850 micrometers.
16. A composition according to any one of Claim 1-15, wherein the carried benefit agent is further mixed with a liquid carrier dispersing material, preferably a liquid carrier selected from nonionic alkoxylated compounds and glycol derivatives, preferably TAE80 and/ or PEG with an weight average molecular width of 1000 to 10,000.

17. A composition according to any one of Claim 1-16 whereby the carrier benefit agent is further mixed with a solid granulation agent, preferably organic acids or salts, inorganic acids or salts, silicates, preferably selected from zeolite, carbonate clay, sulfate, organic acid or acid salt, silicate or mixture thereof.
17. A composition according to any of Claims 1 to 16 whereby the carried benefit agent also comprises a thickening agent and/ or the carried benefit agent comprises as carrier a polymer and it also comprises a cross-linking agent, preferably aldehydes or bifunctional aldehydes, preferably the ratio of carried benefit agent to thickening and/ or cross-linking agent being from 100:1 to 10:1.
18. A composition according to any one of Claim 1-15, wherein the composition is a laundry solid, preferably granular or tablet composition.
19. A composition according to any one of Claims 1-18, wherein said laundry and cleaning composition is in liquid form.
20. A method for providing an enhanced deposition of the benefit agent onto treated surfaces which comprises the steps of contacting the surface with a laundry or cleaning or carried benefit agent as defined in any one of Claims 1-9.
21. A method for providing a delayed release of the benefit agent onto treated surfaces which comprises the steps of contacting the surface with a laundry or cleaning or carried benefit agent as defined in any one of Claims 1-19.
22. A method according to either one of Claim 20 or 21, wherein the surface is a fabric.

23. A method according to any one of Claim 20-22, wherein the release of the benefit agent from the carried benefit agent composition is achieved by hydrolysis of the carrier which has been reacted with a benefit agent.

INTERNATIONAL SEARCH REPORT

Int	nal Application No
PCT/US 00/34833	

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D3/50 D06M13/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D D06M

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 273 775 A (ALBRIGHT & WILSON) 6 July 1988 (1988-07-06) claims examples page 2, line 63 -page 4, line 8 page 4, line 23 - line 28 page 5, line 65 -page 6, line 26 ---	1-5, 11-15, 19-23
A	page 2, line 63 -page 4, line 8 page 4, line 23 - line 28 page 5, line 65 -page 6, line 26 ---	10, 16
X	US 5 780 527 A (O'LEARY NICHOLAS) 14 July 1998 (1998-07-14) claims 1-8, 13 example 15 column 2, line 40 - line 62 column 3, line 60 - line 67 ---	1-5, 10-13
		-/-

 Further documents are listed in the continuation of box C. Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

12 February 2001

Date of mailing of the international search report

19/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
Fax: (+31-70) 340-3016

Authorized officer

Neys, P

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/US 00/34833

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KLECAN V: "STABILISIERTE PARFUMOLE IN DETERGENTIEN UND SEIFEN" SOFW-JOURNAL SEIFEN, OLE, FETTE, WACHSE, DE, VERLAG FUR CHEMISCHE INDUSTRIE, H. ZIOLKOWSKY K.G. AUGSBURG, vol. 116, no. 1, 18 January 1990 (1990-01-18), pages 11-16, XP000114338 ISSN: 0942-7694 Page 12: 'the crosslinking of the substance with a polymer', ... Page 14-15: 'stabilization of perfume oils in toilette soaps' ----	1-4, 11-13
P, X	EP 0 971 025 A (PROCTER & GAMBLE) 12 January 2000 (2000-01-12) cited in the application the whole document ----	1-8, 10-14, 19-23
A	EP 0 816 484 A (GIVAUDAN ROURE INT) 7 January 1998 (1998-01-07) claims examples page 2, line 52 - line 56 page 3, line 16 - line 20 ----	1-5, 10-13, 15, 19-23
A	KAMOGAWA H ET AL: "Chemical release control- Schiff bases of perfume aldehydes and aminostyrenes" JOURNAL OF POLYMER SCIENCE, POLYMER CHEMISTRY EDITION, US, JOHN WILEY AND SONS. NEW YORK, vol. 20, 1 January 1982 (1982-01-01), pages 3121-3129, XP002084908 ISSN: 0887-624X the whole document ----	1, 5-8, 21-24
A	EP 0 392 619 A (QUEST INT) 17 October 1990 (1990-10-17) the whole document -----	1, 5-8, 21, 24

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/US 00/34833

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
EP 0273775	A 06-07-1988	AU 8318487 A	07-07-1988	BR 8707185 A	09-08-1988
		DK 690187 A	01-07-1988	FI 875790 A	01-07-1988
		GB 2200377 A	03-08-1988	JP 1096296 A	14-04-1989
		NO 875493 A	01-07-1988	PT 86485 A, B	01-01-1988
		ZA 8709757 A	23-06-1988	CN 1035124 A	30-08-1989
		PT 88721 B	31-12-1992		
US 5780527	A 14-07-1998	AU 685968 B	29-01-1998	AU 2989995 A	14-03-1996
		BR 9506319 A	05-08-1997	CA 2173153 A	29-02-1996
		CN 1134115 A	23-10-1996	EP 0728021 A	28-08-1996
		WO 9605870 A	29-02-1996	JP 9504221 T	28-04-1997
		NZ 289918 A	26-11-1996	ZA 9506802 A	01-04-1996
EP 0971025	A 12-01-2000	AU 4870399 A	01-02-2000	AU 4870499 A	01-02-2000
		AU 4984699 A	01-02-2000	EP 0971027 A	12-01-2000
		EP 0971021 A	12-01-2000	WO 0002991 A	20-01-2000
		WO 0002986 A	20-01-2000	WO 0002987 A	20-01-2000
EP 0816484	A 07-01-1998	AU 2616697 A	15-01-1998	BR 9703689 A	27-10-1998
		JP 10060482 A	03-03-1998	SG 55330 A	21-12-1998
		ZA 9705339 A	10-09-1998		
EP 0392619	A 17-10-1990	AT 124078 T	15-07-1995	DE 69020219 D	27-07-1995
		DE 69020219 T	02-11-1995	JP 2649983 B	03-09-1997
		JP 3205495 A	06-09-1991	US 5204023 A	20-04-1993
		US 5789010 A	04-08-1998	ZA 9002789 A	24-12-1991