

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number
WO 03/093242 A2

(51) International Patent Classification⁷: **C07D 231/00**

(21) International Application Number: PCT/US02/31770

(22) International Filing Date: 3 October 2002 (03.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/326,721	3 October 2001 (03.10.2001)	US
60/338,623	24 October 2001 (24.10.2001)	US
60/332,857	6 November 2001 (06.11.2001)	US
60/344,957	7 November 2001 (07.11.2001)	US
60/332,107	21 November 2001 (21.11.2001)	US
60/332,104	21 November 2001 (21.11.2001)	US

(71) Applicant (for all designated States except US): **PHARMACIA CORPORATION** [US/US]; Global Patent Department, 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SOUTH, Michael, S.** [US/US]; 11671 Chieftain Drive, St. Louis, MO 63146 (US). **WEBBER, Ronald, K.** [US/US]; 5379 Wolfrum Way, St. Charles, MO 63304 (US). **HUANG, Horng-Chih** [US/US]; 15481 Duxbury Way, Chesterfield, MO 63017 (US). **TOTH, Mihaly, V.** [US/US]; 1031 Claridge Place, St. Louis, MO 63122 (US). **MOORMANN, Alan, E.** [US/US]; 5578 Hennsley Circle, Weldon Springs, MO 63304 (US). **SNYDER, Jeffrey, S.** [US/US]; 478 Corteqway Court, Manchester, MO 63021 (US). **SCHOLTEN, Jeffrey, A.** [US/US]; 246 Cheval Square, Chesterfield, MO 63005 (US). **GARLAND, Danny, J.** [US/US]; 656 Crowsnet Drive, Ballwin, MO 63021 (US). **RUEPPEL, Melvin, L.** [US/US]; 1904 Grassy Ridge Road, St. Louis, MO 63122 (US). **NEUMANN, William, L.** [US/US];

212 West Monroe, St. Louis, MO 63146 (US). **LONG, Scott** [US/US]; 212 Oakbriar Farm Drive, Ballwin, MO 63021 (US). **WEI, Huang** [CN/US]; 919 Crown Pointe Estates Drive, Wildwood, MO 63021 (US). **TRUJILLO, John** [US/US]; 15 Eagles Landing Drive, St. Peters, MO 63376 (US). **PARLOW, John, J.** [US/US]; 4841 Placid Hills Road, Arnold, MO 63010 (US). **JONES, Darin, E.** [US/US]; 408 Johanna Place Drive, Ballwin, MO 63021 (US). **CASE, Brenda** [US/US]; 8 South Tealbrook, St. Louis, MO 63141 (US). **HAYES, Michael, J.** [US/US]; 668 Chelsea Avenue, St. Louis, MO 63122 (US).

(74) Agents: **DOTY, Kathryn, E.** et al.; Senniger, Powers, Leavitt & Roedel, One Metropolitan Square, 16th Floor, St. Louis, MO 63102 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED 5-MEMBERED POLYCYCLIC COMPOUNDS USEFUL FOR SELECTIVE INHIBITION OF THE COAGULATION CASCADE

(57) Abstract: The present invention relates to compounds, and prodrugs thereof, compositions and methods useful for preventing and treating thrombotic conditions in mammals. The compounds of the present invention, and prodrugs thereof, selectively inhibit certain proteases of the coagulation cascade.



WO 03/093242 A2

**Substituted 5-membered Polycyclic Compounds Useful for
Selective Inhibition of the Coagulation Cascade**

Field of the Invention

5 The present invention relates to compounds,
compositions and methods for preventing and treating
thrombotic conditions such as coronary artery and
cerebrovascular disease. More particularly, the invention
relates to 5-membered polycyclic compounds, and prodrugs
thereof that inhibit serine proteases of the coagulation
10 cascade.

Background of the Invention

15 Hemorrhage, intravascular thrombosis, and embolism are
common clinical manifestations of many diseases (see R. I.
Handin *in* Harrison's Principles of Internal Medicine (J.D.
Wilson, et al. eds., 12th ed. 1991) New York, McGraw-Hill
Book Co., pp. 348-351). The normal hemostatic system
limits blood loss by precisely regulated interactions
between components of the vessel wall, circulating blood
platelets, and plasma proteins. Unregulated activation of
20 the of the hemostatic system, however, may cause
thrombosis, which can reduce blood flow to critical organs
like the brain and myocardium.

25 Physiological systems control the fluidity of blood in
mammals (see P.W. Majerus, et al. *in* Goodman & Gilman's The
Pharmacological Basis of Therapeutics (J.G. Hardman & L.E.
Limbird, eds., 9th ed. 1996) New York, McGraw-Hill Book

Co., pp. 1341-1343). Blood must remain fluid within the vascular systems and yet quickly be able to undergo hemostasis. Hemostasis, or clotting, begins when platelets first adhere to macromolecules in subendothelial regions of injured and/or damaged blood vessels. These platelets aggregate to form the primary hemostatic plug and stimulate local activation of plasma coagulation factors leading to generation of a fibrin clot that reinforces the aggregated platelets.

Plasma coagulation factors, also referred to as protease zymogens, include factors II, V, VII, VIII, IX, X, XI, and XII. These coagulation factors or protease zymogens are activated by serine proteases leading to coagulation in a so called "coagulation cascade" or chain reaction.

Coagulation or clotting occurs in two ways through different pathways. An intrinsic or contact pathway leads from XII to XIIa to XIa to IXa and to the conversion of X to Xa. Xa with factor Va converts prothrombin (II) to thrombin (IIa) leading to conversion of fibrinogen to fibrin. Polymerization of fibrin leads to a fibrin clot. An extrinsic pathway is initiated by the conversion of coagulation factor VII to VIIa by Xa. Factor VIIa, a plasma protease, is exposed to, and combines with its essential cofactor tissue factor (TF) which resides constitutively beneath the endothelium. The resulting factor VIIa/TF complex proteolytically activates its substrates, factors IX and X, triggering a cascade of reactions that leads to the generation of thrombin and a fibrin clot as described above.

While clotting as a result of an injury to a blood vessel is a critical physiological process for mammals, clotting can also lead to disease states. A pathological process called thrombosis results when platelet aggregation and/or a fibrin clot blocks (i.e., occludes) a blood vessel. Arterial thrombosis may result in ischemic

necrosis of the tissue supplied by the artery. When the thrombosis occurs in a coronary artery, a myocardial infarction or heart attack can result. A thrombosis occurring in a vein may cause tissues drained by the vein to become edematous and inflamed. Thrombosis of a deep vein may be complicated by a pulmonary embolism. Preventing or treating clots in a blood vessel may be therapeutically useful by inhibiting formation of blood platelet aggregates, inhibiting formation of fibrin, inhibiting thrombus formation, inhibiting embolus formation, and for treating or preventing unstable angina, refractory angina, myocardial infarction, transient ischemic attacks, atrial fibrillation, thrombotic stroke, embolic stroke, deep vein thrombosis, disseminated intravascular coagulation, ocular build up of fibrin, and reocclusion or restenosis of recanalized vessels.

In order to treat such conditions, researchers have sought to discover chemical compounds that efficaciously and selectively control the clotting process. In addition, such compounds may provide a better understanding of the pathways involved in the coagulation process.

Thus far, the compounds that have been discovered often possess a polar or basic functional group which is integrally responsible for the desired biological activity. Frequently, this polar functional group is a nitrogen atom of, for example, a guanidine, alkyl-amidine or aryl-amidine group. Because these functionalities are highly basic, they remain protonated at physiologically relevant pH's. The ionic nature of such protonated species hinders their permeability across lipophilic membranes, which can reduce bioavailability when the pharmaceutical agent is administered orally.

In order to circumvent such a problem, it is often advantageous to perform a derivatization or chemical modification of the polar functionality such that the pharmaceutical agent becomes neutrally charged and more


lipophilic, thereby facilitating absorption of the drug. However, for the derivatization to be useful, the derivatization must be bioconvertible at the target site or sites of desired pharmacological activity and cleaved under normal physiological conditions to yield the biologically active drug. The term "prodrug" has been used to denote such a chemically modified intermediate.

Summary of the Invention

Among the several aspects of the present invention, therefore, is the provision of compounds useful for selective inhibition of certain enzymes that act upon the coagulation cascade thereby preventing and treating thrombotic conditions in mammals.

Another aspect of the present invention is the provision of prodrug compounds useful for selective inhibition of certain enzymes that act upon the coagulation cascade thereby preventing and treating thrombotic conditions in mammals. In general, these prodrug compounds undergo hydrolysis, oxidation, reduction or elimination at a derivatized amidine group to yield the active compound.

Briefly, therefore, the present invention is directed to the compound, per se, to the prodrug of the compound, to pharmaceutical compositions comprising the compound or prodrug and a pharmaceutically acceptable carrier, and to methods of use. In one embodiment, the invention embraces compounds having formula (I):



(I)

wherein:

X comprises a 5-membered heterocyclic or carbocyclic ring, the ring atoms being X_1 , X_2 , X_3 , X_4 , and X_5 , wherein X_1 ,

X_3 , and X_4 , are independently carbon or nitrogen and X_2 and X_5 are independently carbon, nitrogen, oxygen or sulfur;


L_1 , L_3 and L_4 are linkages through which Z_1 , Z_3 , and Z_4 , respectively, are covalently bonded to different ring atoms of the 5-membered heterocyclic or carbocyclic ring of X, wherein Z_1 is covalently bonded to X_1 , Z_3 is covalently bonded to X_3 , and Z_4 is covalently bonded to X_4 , each of L_1 , L_3 and L_4 independently being a covalent bond or comprising one or more atoms through which Z_1 , Z_3 , and Z_4 are covalently bonded to X_1 , X_3 and X_4 , respectively;

Z_1 is hydrocarbyl or substituted hydrocarbyl.

Z_3 comprises a 5- or 6-membered heterocyclic or aromatic ring substituted with an amidine group, the ring atoms of the 5- or 6-membered heterocyclic or aromatic ring of Z_3 being carbon, sulfur, nitrogen, or oxygen, wherein the 5- or 6-membered ring is optionally substituted at any position with halogen, alkyl or hydroxy; and

Z_4 comprises a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of the 5- or 6-membered heterocyclic or carbocyclic ring of Z_4 being carbon, nitrogen, oxygen, or sulfur.

Another aspect of the invention provides compounds corresponding to formula (II)



(II)

wherein:

X_2 and X_5 are as defined for compounds having formula (I);

X_1 , X_3 and X_4 are independently carbon or nitrogen;

X_9 is a direct bond or $-(CH_2)_m-$ where m is 1 or 2; and

Z₁, Z₃ and Z₄ are as defined for compounds having formula (I).

Other aspects and features of this invention will be in part apparent and in part pointed out hereafter.

5 **Abbreviations and Definitions**

The term "elimination" is generally meant to encompass any one or more of the following reactions: (1) a reaction that results in a compound fragmenting into two or more compounds; and (2) a reaction that results in one or more
10 groups being removed from a compound without being replaced by other groups.

The term "oxidation" is generally meant to encompass any one or more of the following reactions: (1) a reaction that results in an increase in the oxidation number of an
15 atom in a compound, whether the atom is uncharged or charged and whether free or covalently bound; (2) a reaction that results in the loss of hydrogen from a compound; (3) a reaction that results in the loss or removal of one or more electrons from a compound, with or
20 without concomitant loss or removal of a proton or protons; (4) the action or process of reacting a compound with oxygen; and (5) a reaction that results in the addition of one or more oxygen atoms to a compound.

The term "reduction" is generally meant to encompass
25 any one or more of the following reactions: (1) any reaction which results in a decrease in the oxidation number of an atom in a compound; and (2) any reaction that results in oxygen being withdrawn from, hydrogen being added to, or an electron being added to (with or without
30 the addition of a proton) a compound.

The term "hydrolysis" is generally meant to encompass any one or more of the following reactions: (1) any
reaction which results in the addition of a nucleophile to a compound to form a new bond with concurrent loss of a
35 group from the compound; (2) any reaction which results in

the addition of water to a compound; and (3) any reaction that results in the rupture of one or more chemical bonds by reaction with, and involving the addition of, the elements of water.

5 The term "physiological conditions" are those conditions characteristic to an organism's (to a human beings) healthy or normal functioning.

 The terms "hydrocarbon" and "hydrocarbyl" as used herein describe organic compounds or radicals consisting
10 exclusively of the elements carbon and hydrogen. These moieties include alkyl, alkenyl, alkynyl, and aryl moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl moieties substituted with other aliphatic or cyclic hydrocarbon groups, such as alkaryl, alkenaryl
15 and alkynaryl. Unless otherwise indicated, these moieties preferably comprise 1 to 20 carbon atoms.

 The "substituted hydrocarbyl" moieties described herein are hydrocarbyl moieties which are substituted with at least one atom other than carbon, including moieties in
20 which a carbon chain atom is substituted with a heteroatom such as nitrogen, oxygen, silicon, phosphorus, boron, sulfur, or a halogen atom. Exemplary substituted hydrocarbyl moieties include, heterocyclo, alkoxyalkyl, alkenyloxyalkyl, alkynyloxyalkyl, aryloxyalkyl,
25 hydroxyalkyl, protected hydroxyalkyl, keto, acyl, nitroalkyl, aminoalkyl, cyano, alkylthioalkyl, arylthioalkyl, ketals, acetals, amides, acids, esters and the like.

 The term "heteroatom" shall mean atoms other than
30 carbon and hydrogen.

 Unless otherwise indicated, the alkyl groups described herein are preferably lower alkyl containing from one to eight carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain or
35 cyclic and include methyl, ethyl, propyl, isopropyl, butyl, hexyl and the like.

Unless otherwise indicated, the alkenyl groups described herein are preferably lower alkenyl containing from two to eight carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain or cyclic and include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like.

Unless otherwise indicated, the alkynyl groups described herein are preferably lower alkynyl containing from two to eight carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and include ethynyl, propynyl, butynyl, isobutynyl, hexynyl, and the like.

The terms "aryl" or "ar" as used herein alone or as part of another group denote optionally substituted homocyclic aromatic groups, preferably monocyclic or bicyclic groups containing from 6 to 12 carbons in the ring portion, such as phenyl, biphenyl, naphthyl, substituted phenyl, substituted biphenyl or substituted naphthyl. Phenyl and substituted phenyl are the more preferred aryl.

The terms "halogen" or "halo" as used herein alone or as part of another group refer to chlorine, bromine, fluorine, and iodine.

The terms "heterocyclo" or "heterocyclic" as used herein alone or as part of another group denote optionally substituted, fully saturated or unsaturated, monocyclic or bicyclic, aromatic or nonaromatic groups having at least one heteroatom in at least one ring, and preferably 5 or 6 atoms in each ring. The heterocyclo group preferably has 1 or 2 oxygen atoms, 1 or 2 sulfur atoms, and/or 1 to 4 nitrogen atoms in the ring, and may be bonded to the remainder of the molecule through a carbon or heteroatom. Exemplary heterocyclo include heteroaromatics such as furyl, thienyl, pyridyl, oxazolyl, pyrrolyl, indolyl, quinolinyl, or isoquinolinyl and the like. Exemplary substituents include one or more of the following groups: hydrocarbyl, substituted hydrocarbyl, keto, hydroxy,

protected hydroxy, acyl, acyloxy, alkoxy, alkenoxy, alkynoxy, aryloxy, halogen, amido, amino, nitro, cyano, thiol, ketals, acetals, esters and ethers.

5 The term "heteroaromatic" as used herein alone or as part of another group denote optionally substituted aromatic groups having at least one heteroatom in at least one ring, and preferably 5 or 6 atoms in each ring. The heteroaromatic group preferably has 1 or 2 oxygen atoms, 1 or 2 sulfur atoms, and/or 1 to 4 nitrogen atoms in the
10 ring, and may be bonded to the remainder of the molecule through a carbon or heteroatom. Exemplary heteroaromatics include furyl, thienyl, pyridyl, oxazolyl, pyrrolyl, indolyl, quinolinyl, or isoquinolinyl and the like. Exemplary substituents include one or more of the following
15 groups: hydrocarbyl, substituted hydrocarbyl, keto, hydroxy, protected hydroxy, acyl, acyloxy, alkoxy, alkenoxy, alkynoxy, aryloxy, halogen, amido, amino, nitro, cyano, thiol, ketals, acetals, esters and ethers.

20 The term "acyl," as used herein alone or as part of another group, denotes the moiety formed by removal of the hydroxyl group from the group -COOH of an organic carboxylic acid, e.g., $RC(O)-$, wherein R is hydrogen, R^1 , R^1O- , R^1R^2N- , or R^1S- , R^1 is hydrocarbyl, heterosubstituted hydrocarbyl, or heterocyclo, and R^2 is hydrogen, hydrocarbyl
25 or substituted hydrocarbyl.

The term "acyloxy," as used herein alone or as part of another group, denotes an acyl group as described above bonded through an oxygen linkage (-O-), e.g., $RC(O)O-$ wherein R is as defined in connection with the term "acyl."

30 The term "acetamidyl" as used herein describes a chemical moiety represented by the formula $NR_1C(O)R_2$.

The term "carboxamido" as used herein, describes a chemical moiety represented by the formula $C(O)NR_1R_2$.

35 The term "alkoxycarbonyl" as used herein describes a chemical moiety represented by the formula $C(O)OR$.

The term "sulfonamido" as used herein describes a chemical moiety represented by the formula $\text{SO}_2\text{NR}_1\text{R}_2$.


The term "alkylsulfonyl" as used herein describes a chemical moiety represented by the formula SO_2R .

5 The term "sulfonamidyl" as used herein describes a chemical moiety represented by the formula NRSO_2R .

As described herein for the terms "acetamidyl", "carboxamido", "alkocycarbonyl", "sulfonamido", "alkylsulfonyl", and "sulfonamidyl", R, R_1 and R_2 are
 10 independently hydrogen, alkyl, aryl, and arylakyl, optionally substituted with halogen, hydroxy or alkoxy.

Description of the Preferred Embodiment

One aspect of the invention embraces compounds that
 15 correspond to formula (I)



(I)

wherein:

X comprises a 5-membered heterocyclic or carbocyclic ring, the ring atoms being X_1 , X_2 , X_3 , X_4 , and X_5 , wherein X_1 , X_3 , and X_4 , are independently carbon or nitrogen and X_2 and X_5 are independently carbon, nitrogen, oxygen or sulfur;
 20

L_1 , L_3 and L_4 are linkages through which Z_1 , Z_3 , and Z_4 , respectively, are covalently bonded to different ring atoms of the 5-membered heterocyclic or carbocyclic ring of X, wherein Z_1 is covalently bonded to X_1 , Z_3 is covalently bonded to X_3 , and Z_4 is covalently bonded to X_4 , each of L_1 , L_3 and L_4 independently being a covalent bond or comprising one or more atoms through which Z_1 , Z_3 , and Z_4 are covalently bonded to X_1 , X_3 and X_4 , respectively;
 25

30 Z_1 is hydrocarbyl or substituted hydrocarbyl;

Z₃ comprises a 5- or 6-membered heterocyclic or aromatic ring substituted with an amidine group, the ring atoms of the 5- or 6-membered heterocyclic or aromatic ring of Z₃ being carbon, sulfur, nitrogen, or oxygen, wherein the 5- or 6-membered ring is optionally substituted at any position with halogen, alkyl or hydroxy; and

Z₄ comprises a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of the 5- or 6-membered heterocyclic or carbocyclic ring of Z₄ being carbon, nitrogen, oxygen, or sulfur.

In another embodiment for compounds having formula (I), each of X₁, X₂, X₃, X₄, and X₅ are as defined in the embodiment above;

L₁ is -X₉NH- wherein X₉ is covalently bonded directly to Z₁ and X₉ is a direct bond or -(CH₂)_m- wherein m is 1 to 5;

L₃ is a glycine derivative;

L₄ is a direct bond;

Z₁ is selected from the group consisting of C₁-C₈ alkyl, C₂-C₈ alkenyl, and C₂-C₈ alkynyl, the alkyl, alkenyl, or alkynyl being optionally substituted at any substitutable position with a halogen;

Z₃ comprises a phenyl, furanyl or thienyl ring, the phenyl, furanyl or thienyl ring being optionally further substituted with fluorine or hydroxy;

Z₄ comprises a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of Z₄ being Z₄₀, Z₄₁, Z₄₂, Z₄₄ and Z₄₅ when Z₄ is a 5-membered ring and Z₄₀, Z₄₁, Z₄₂, Z₄₃, Z₄₄ and Z₄₅ when Z₄ is a 6-membered ring, Z₄₀, Z₄₁, Z₄₂, Z₄₃, Z₄₄ and Z₄₅, being carbon, nitrogen, oxygen or sulfur, Z₄₀ being the ring atom through which Z₄ is attached to the heterocyclic ring, Z₄₁ and Z₄₅ each being in an alpha position relative to Z₄₀, Z₄₂ and Z₄₄ each being in a beta position relative to Z₄₀, Z₄₃ being in the gamma position relative to Z₄₀ when Z₄ is a 6-membered ring, Z₄ having a substituent R₄₂ covalently attached to Z₄₂, and a second

substituent bonded to one of Z_{41} , Z_{43} , Z_{44} , or Z_{45} , the substituent being R_{41} when bonded to Z_{41} , the substituent being R_{43} when bonded to Z_{43} , the substituent being R_{44} when bonded to Z_{44} , and the substituent being R_{45} when bonded to Z_{45} ;

R_{42} is amino; and

R_{41} , R_{43} , R_{44} and R_{45} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen, or an optionally substituted heteroatom selected from nitrogen, oxygen, sulfur and phosphorus, provided at least one of R_{41} , R_{43} , R_{44} or R_{45} is other than hydrogen.

Still a further embodiment provides compounds having formula (I), where L_1 is $-X_9NH-$ wherein X_9 is covalently bonded directly to Z_1 and X_9 is a direct bond;

L_3 is $-CH_2CONHCH_2-$ and Z_3 is covalently bonded to the methylene bonded to the amine nitrogen of L_3 ;

Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl;

Z_3 comprises a phenyl ring, the phenyl ring being optionally further substituted with fluorine or hydroxy;

Z_4 comprises a phenyl or thienyl ring having two substituents, R_{42} and R_{44} , and two ring atoms each of which is in the beta position relative to the ring atom of Z_4 through which Z_4 is covalently bonded to X , wherein one of R_{42} and R_{44} is covalently bonded to one of the beta positions and the other of R_{42} and R_{44} is covalently bonded to the other of the beta positions;

R_{42} is amino; and

R_{44} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen, and an optionally substituted heteroatom selected from nitrogen, oxygen, sulfur and phosphorus.

In one embodiment for compounds having formula (I), the L₁ linkage is -X₉NH- where X₉ is covalently bonded directly to Z₁ and is a direct bond or an alkylene chain having the formula (CH₂)_m wherein m is 0 to 5. In an
5 alternative embodiment, m is 0 to 2. In another embodiment, the L₁ linkage is a bond.


In another embodiment for compounds having formula (I), the L₃ linkage is a glycine derivative, an alanine derivative, an amino derivative, or a sulfonyl derivative.
10 In an alternative embodiment, the L₃ linkage is a glycine derivative. In still another embodiment, the L₃ linkage is -CH₂CONHCH₂- where Z₃ is covalently bonded to the methylene bonded to the amine nitrogen of L₃.

In a further embodiment for compounds having formula (I), the L₄ linkage is a direct bond, methylene, ethylene or
15 an optionally substituted heteroatom selected from the group nitrogen, oxygen, sulfur or phosphorus. In another embodiment, the L₄ linkage is a direct bond.

In yet another embodiment for compounds having formula (I), Z₁ is a C₁-C₅ alkyl optionally substituted at any
20 substitutable position with fluorine, hydroxy, carboxy or alkoxy carbonyl. Generally speaking, the C₁-C₅ alkyl is a cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted
25 at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl. In another embodiment, the C₁-C₅ alkyl is isopropyl or cyclobutyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl. In yet another alternative of
30 this embodiment, Z₁ is trifluoroethyl or carboxymethyl.

In one embodiment of compounds corresponding to formula (I), Z₃ is a 5- or 6-membered heterocyclic or aromatic ring substituted with an amidine group and may be
35 optionally substituted at any substitutable position with halogen, hydroxy, haloalkyl, alkyl, carboxy, alkoxy carbonyl, or hydrocarbyloxy, or any combination

thereof. A preferred halogen is fluorine. In another alternative of this embodiment, Z_3 is a 6-membered carbocyclic aromatic ring substituted with an amidine group. In yet another embodiment, Z_3 comprises a substituted phenyl, thienyl, or furanyl ring, the phenyl, thienyl or furanyl ring being substituted with an amidine group, and optionally further substituted at any substitutable position with fluorine, hydroxy, carboxy, alkoxy, alkoxy, or hydroxy. In yet another alternative of this embodiment, Z_3 corresponds to formula (a)



(a)

wherein:

R_{304} and R_{306} are independently selected from the group consisting of hydrogen, fluorine, hydroxy, carboxy, hydrocarbyloxy and alkoxy; and

R_{305} and R_{307} are independently selected from the group consisting of hydrogen, fluorine, methoxy, hydroxy and carboxy.

In one alternative of compounds wherein Z_3 corresponds to formula (a), Z_3 is selected from the group consisting of benzamidine-4-yl, 3-hydroxybenzamidine-4-yl, 3,5-dihydroxybenzamidine-4-yl, 2,5,6-trifluoro-3-hydroxybenzamidine-4-yl, 2-hydroxybenzamidine-4-yl and 3,5,6-trifluoro-2-hydroxybenzamidine-4-yl.

A further embodiment embraces prodrug compounds having formula (I) where Z_3 is $-R_{300}C(=NR_{301})NR_{302}R_{303}$, wherein R_{300} is a

6-membered carbocyclic aromatic ring, R_{301} , R_{302} , R_{303} are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur, provided at least one of R_{301} , R_{302} , R_{303} is other than hydrogen. In yet another embodiment, Z_3 is


$-R_{300}C(=NR_{301})NR_{302}R_{303}$, wherein R_{300} is a 6-membered carbocyclic aromatic ring, and at least two of R_{301} , R_{302} , R_{303} are ring atoms of a heterocyclic ring. In an alternative embodiment, Z_3 is $-R_{300}C(=NR_{301})NR_{302}R_{303}$, R_{300} is a 6-membered carbocyclic aromatic ring, and at least one of R_{301} , R_{302} , R_{303} are ring atoms of a heterocyclic ring fused to R_{300} .

Yet another embodiment encompasses prodrug compounds having formula (I) where Z_3 is a benzamide derivatized with one or more groups selected from carbonyl, thiocarbonyl, imino, enamino, phosphorus, and sulfur, where the benzamide derivative hydrolyzes under physiological conditions to form benzamide. In an alternative to this embodiment, Z_3 is a benzamide derivatized with one or more groups selected from optionally substituted hydrocarbyl, provided that the carbon atom directly bonded to the amidine is sp^3 hybridized and aryl, where the benzamide derivative is oxidized under physiological conditions to form benzamide. In yet another alternative for this embodiment, Z_3 is a benzamide derivatized with one or more heteroatoms selected from oxygen, nitrogen in its most reduced state, and sulfur in its most reduced state, where the benzamide derivative is reduced under physiological conditions to form benzamide. In still another alternative for this embodiment, Z_3 is a benzamide derivatized with one or more substituents selected from a hydrocarbyl substituted at the beta carbon with carbonyl, sulfonyl, sulfinyl, cyano, nitro and an alkyl, aryl, or heterocyclic group substituted with oxygen, nitrogen, or sulfur at the carbon directly bonded to the amidine group,

where the benzamidine derivative undergoes elimination at physiological conditions to form benzamidine.

In a further embodiment for prodrug compounds having formula (I), Z_3 corresponds to formula (b)

5



(b)

wherein:

R_{301} , R_{302} , and R_{303} are independently selected from the group consisting of:

10

(i) hydrogen, $-C(=O)R_a$, $-C(=O)OR_a$, $-S(=O)OR_a$, $-S(=O)SR_a$, $-S(=O)_2OR_a$, $-S(=O)_2SR_a$ and alkenyl, wherein R_a is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, and heterocylo, provided, however, that the carbon atom of R_{301} , R_{302} , and R_{303} directly bonded to the amidine is sp^2 hybridized when R_{301} , R_{302} , and R_{303} is alkenyl,

15

(ii) hydrogen, optionally substituted hydrocarbyl and aryl, provided, however, the carbon atom of R_{301} , R_{302} , and R_{303} directly bonded to the amidine is sp^3 hybridized when R_{301} , R_{302} , and R_{303} is optionally substituted hydrocarbyl,

20

(iii) hydrogen, $-OR_b$, $-SR_b$, $-NR_b$, or $-N(R_b)_2$, wherein each R_b is independently optionally substituted hydrocarbyl, and heterocylo, and

25

(iv) hydrogen, substituted hydrocarbyl wherein the carbon bonded to the amidine group is substituted with $-OR_c$, $-SR_c$, $-NR_c$, or $-N(R_c)_2$, wherein each R_c is independently $-C(O)R_d$, $-C(O)NR_d$, $-C(O)OR_d$, $-C(O)N(R_d)_2$ and each R_d is independently hydrocarbyl, substituted hydrocarbyl or heterocyclo, and substituted alkyl with the carbon atom

beta to the point of attachment to the amidine group being an unsaturated electron withdrawing group, provided, however, at least one of R_{301} , R_{302} , and R_{303} is other than hydrogen;

5 R_{304} is selected from the group consisting of halogen, hydrogen, hydroxyl, alkyl, sulfhydryl, alkoxy, and alkylthio;

R_{305} is selected from the group consisting of oxygen, sulfur, halogen, hydrogen, hydroxyl, alkyl, sulfhydryl, alkoxy, and alkylthio;

10 R_{306} is selected from the group consisting of halogen, hydrogen, hydroxyl, alkyl, sulfhydryl, alkoxy, and alkylthio; and

R_{307} is selected from the group consisting of oxygen, sulfur, halogen, hydrogen, hydroxyl, alkyl, sulfhydryl, alkoxy, and alkylthio.


In one embodiment for prodrug compounds having formula (I), the benzamidine derivative is hydrolyzed under physiological conditions to form benzamidine when Z_3 is a benzamidine derivative having formula (b) and R_{301} , R_{302} , and R_{303} are independently selected from hydrogen, $-C(=O)R_a$, $-C(=O)OR_a$, $-S(=O)OR_a$, $-S(=O)SR_a$, $-S(=O)_2OR_a$, $-S(=O)_2SR_a$ and alkenyl, wherein R_a is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, and heterocylo, provided, however, that the carbon atom of R_{301} , R_{302} , and R_{303} directly bonded to the amidine is sp^2 hybridized when R_{301} , R_{302} , and R_{303} is alkenyl.

In a further embodiment for prodrug compounds having formula (I), the benzamidine derivative is oxidized under physiological conditions to form benzamidine when Z_3 is a benzamidine derivative having formula (b) and R_{301} , R_{302} , and R_{303} are independently selected from hydrogen, optionally substituted hydrocarbyl and aryl, provided, however, the carbon atom of R_{301} , R_{302} , and R_{303} directly bonded to the amidine is sp^3 hybridized when R_{301} , R_{302} , and R_{303} is optionally substituted hydrocarbyl.

In still another embodiment for prodrug compounds having formula (I), the benzamidine derivative is reduced under physiological conditions to form benzamidine when Z_3 is a benzamidine derivative having formula (b) and R_{301} , R_{302} , and R_{303} are independently selected from hydrogen, $-OR_b$, $-SR_b$, $-NR_b$, or $-N(R_b)_2$, wherein each R_b is independently optionally substituted hydrocarbyl, and heterocyclo.

In an alternative embodiment for prodrug compounds having formula (I), the benzamidine derivative undergoes elimination at physiological conditions to form benzamidine when Z_3 is a benzamidine derivative having formula (b) and R_{301} , R_{302} , and R_{303} are independently selected from hydrogen, substituted hydrocarbyl wherein the carbon bonded to the amidine group is substituted with $-OR_c$, $-SR_c$, $-NR_c$, or $-N(R_c)_2$, wherein each R_c is independently $-C(O)R_d$, $-C(O)NR_d$, $-C(O)OR_d$, $-C(O)N(R_d)_2$ and each R_d is independently hydrocarbyl, substituted hydrocarbyl or heterocyclo, and substituted alkyl with the carbon atom beta to the point of attachment to the amidine group being an unsaturated electron withdrawing group.

A further embodiment embraces compounds having formula (I) where Z_4 is a substituted, 6-membered carbocyclic aromatic ring. In an alternative of this embodiment, Z_4 corresponds to formula (c)



(c)

wherein:

R_{42} is amino;

R_{44} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, halogen and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur; and

5 R_{41} , R_{43} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.

10 In another embodiment of compounds wherein Z_4 corresponds to formula (c) and R_{42} is amino, R_{44} is selected from hydrocarbyl, substituted hydrocarbyl, acetamido, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkyl, haloalkoxy, haloalkylthio, carboalkoxy, carboxy, 15 carboxamidoalkyl, and carboxamidoalkylaryl. In an alternative of this embodiment, R_{44} is selected from hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroaryl, heterocyclo, halogen, acetamido, guanidino, hydroxy, nitro, amino, amidosulfonyl, acylamido, hydrocarbyloxy, 20 substituted hydrocarbyloxy, hydrocarbylthio, substituted hydrocarbylthio, hydrocarbylsulfonyl, and substituted hydrocarbylsulfonyl. In yet another alternative of this embodiment, R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, 25 carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacylamine, 30 isobutylamide, *sec*-pentylamine, cyclopentylacylamine, 1-carboxyl-2-methylbutylamide, isobutylacylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and 35 carbmethoxy. In still another alternative of this embodiment, R_{41} , R_{43} and R_{45} are independently selected from

the group consisting of hydrogen, halogen, alkoxy, or hydroxy and R_{44} is as defined in any of the alternative embodiments above. In yet another alternative of this embodiment, R_{41} , R_{43} and R_{45} are independently selected from the group consisting of hydrogen and halogen and R_{44} is as defined in any of the alternative embodiments above.

In another embodiment for compounds wherein Z_4 corresponds to formula (c) and R_{42} is amino, R_{45} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamidyl, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkoxy, haloalkythio, alkoxy-carbonyl, carboxy, sulfonamido, carboxamido and sulfonamidyl, optionally substituted with fluorine. In another alternative of this embodiment (i.e., when Z_4 corresponds to formula (c) and R_{42} is amino), R_{45} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamidyl, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkoxy, haloalkythio, alkoxy-carbonyl, sulfonamido, carboxamido and sulfonamidyl, optionally substituted with fluorine. In still another alternative of this embodiment, R_{45} is selected from the group consisting of hydroxy, carboxy, carboxamido, alkoxy, alkylsulfonyl, sulfonamido, and alkoxy-carbonyl. In yet another alternative of this embodiment, R_{45} is selected from the group consisting of sec-butylamide, carboxy, ethoxy-carbonyl, isopropoxy-carbonyl, butoxy-carbonyl, isopropylamide and hydroxy. In still another alternative of this embodiment, R_{41} , R_{43} and R_{44} are independently selected from the group consisting of hydrogen, halogen, alkoxy, or hydroxy and R_{45} is as defined in any of the alternative embodiments above. In yet another alternative of this embodiment, R_{41} , R_{43} and R_{44} are independently selected from the group consisting of hydrogen and halogen and R_{45} is as defined in any of the alternative embodiments above.


In yet another embodiment for compounds wherein Z_4 corresponds to formula (c) and R_{42} is amino, R_{43} is selected

from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamidyl, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkoxy, haloalkythio, alkoxycarbonyl, carboxy, sulfonamido, carboxamido and sulfonamidyl, optionally substituted with fluorine. In another alternative of this embodiment (i.e., when Z_4 corresponds to formula (c) and R_{42} is amino), R_{43} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamidyl, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkoxy, haloalkythio, alkoxycarbonyl, sulfonamido, carboxamido and sulfonamidyl, optionally substituted with fluorine. In still another alternative of this embodiment, R_{43} is selected from the group consisting of hydroxy, carboxy, carboxamido, alkoxy, alkylsulfonyl, sulfonamido, and alkoxycarbonyl. In yet another alternative of this embodiment, R_{43} is selected from the group consisting of sec-butylamide, carboxy, ethoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isopropylamide and hydroxy. In still another alternative of this embodiment, R_{41} , R_{44} and R_{45} are independently selected from the group consisting of hydrogen, halogen, alkoxy, or hydroxy and R_{43} is as defined in any of the alternative embodiments above. In yet another alternative of this embodiment, R_{41} , R_{44} and R_{45} are independently selected from the group consisting of hydrogen and halogen and R_{43} is as defined in any of the alternative embodiments above.

In still another embodiment for compounds wherein Z_4 corresponds to formula (c) and R_{42} is amino, R_{41} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamidyl, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkoxy, haloalkythio, alkoxycarbonyl, carboxy, sulfonamido, carboxamido and sulfonamidyl, optionally substituted with fluorine. In another alternative of this embodiment (i.e., when Z_4 corresponds to formula (c) and R_{42} is amino), R_{41} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl,

acetamidyl, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkoxy, haloalkythio, alkoxy-carbonyl, sulfonamido, carboxamido and sulfonamidyl, optionally substituted with fluorine. In still another alternative of this embodiment, 5 R_{41} is selected from the group consisting of hydroxy, carboxy, carboxamido, alkoxy, alkylsulfonyl, sulfonamido, and alkoxy-carbonyl. In yet another alternative of this embodiment, R_{41} is selected from the group consisting of sec-butylamide, carboxy, ethoxycarbonyl, 10 isopropoxycarbonyl, butoxycarbonyl, isopropylamide and hydroxy. In still another alternative of this embodiment, R_{43} , R_{44} and R_{45} are independently selected from the group consisting of hydrogen, halogen, alkoxy, or hydroxy and R_{41} is as defined in any of the alternative embodiments above. 15 In yet another alternative of this embodiment, R_{43} , R_{44} and R_{45} are independently selected from the group consisting of hydrogen and halogen and R_{41} is as defined in any of the alternative embodiments above.

In yet another embodiment, Z_4 is a five-membered ring having formula (d) 20



(d)

wherein:

Z_{40} , Z_{41} , Z_{42} , Z_{44} , and Z_{45} are independently selected from the group consisting of carbon, nitrogen, oxygen and sulfur; 25

R_{42} is amino;

R_{44} is selected from the group consisting of is selected from the group consisting of hydrocarbyl,

substituted hydrocarbyl, halogen and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur; and

5 R_{41} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.

10 In another embodiment of compounds wherein Z_4 corresponds to formula (d) and R_{42} is amino, R_{44} is selected from hydrocarbyl, substituted hydrocarbyl, acetamido, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkyl, haloalkoxy, haloalkylthio, carboalkoxy, carboxy, carboxamidoalkyl, and carboxamidoalkylaryl. In an
15 alternative of this embodiment, R_{44} is selected from hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroaryl, heterocyclo, halogen, acetamido, guanidino, hydroxy, nitro, amino, amidosulfonyl, acylamido, hydrocarbyloxy, substituted hydrocarbyloxy, hydrocarbylthio, substituted
20 hydrocarbylthio, hydrocarbylsulfonyl, and substituted hydrocarbylsulfonyl. In yet another alternative of this embodiment, R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-
25 aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-
30 carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy. In still another alternative of this
35 embodiment, R_{41} and R_{45} are independently selected from the group consisting of hydrogen, halogen, alkoxy, or hydroxy

and R₄₄ is as defined in any of the alternative embodiments above. In yet another alternative of this embodiment, R₄₁ and R₄₅ are independently selected from the group consisting of hydrogen and halogen and R₄₄ is as defined in any of the alternative embodiments above.


Still another embodiment provides compounds having formula (I) where X₂ or X₅ are hydrogen bond acceptors. In another embodiment, both X₂ and X₅ are hydrogen bond acceptors. Generally speaking, the phrase "hydrogen bond acceptor" encompasses heteroatoms having a lone pair of electrons available for hydrogen bonding. Suitable hydrogen bond acceptors, when taken with the carbon to which Z₂ is attached, are typically selected from the group consisting of C(O), C(S), C(Cl), C(Br), C(F), C(OH), COCH₃, COR, C(SH), CSR, and CNR₁R₂ wherein R, R₁ and R₂ are independently hydrogen, alkyl, aryl, and arylalkyl, optionally substituted with halogen, hydroxy or alkoxy. Suitable X₂ groups include carbon substituted with hydrogen, fluorine, oxygen, or sulfur, nitrogen optionally substituted with hydrogen or oxygen, oxygen, or sulfur. Suitable X₅ groups include oxygen, sulfur, nitrogen, carbonyl and carbon substituted with a halogen selected from fluorine, chlorine and bromine.

In one embodiment for compounds having formula (I), X₁, X₂, X₃, X₄, and X₅ are carbon or nitrogen. In an alternative of this embodiment, X₁, X₂ and X₅ are carbon, and X₃ and X₄ are nitrogen. In another alternative of this embodiment, X₁, X₃, X₄ and X₅ are carbon and X₂ is nitrogen. In yet another alternative of this embodiment, X₁, X₂, X₃, X₄ and X₅ are all carbon. In still a further alternative of this embodiment, X₁, X₄ and X₅ are carbon and X₂ and X₃ are nitrogen.

In another embodiment for compounds having formula (I), X₁, X₂, X₃, X₄, and X₅ form a 5-membered heterocyclic or carbocyclic ring selected from pyrazolinone, pyrrole, thiophene, pyrazole-N-oxide, 1-amino-pyrazole, 1,3,4-

triazole, 2-amino-4-aryl-thiazole, 2-amino-5-aryl-thiazole,
 pyrrolidine, 2-amino-5-aryl-oxazole, 3-amino-pyrazole, 2-
 amio-4-aryl-oxazole, tetrahydrofuran, cyclopentadienone,
 and N-hydroxypyrrolidine. In an alternative of this
 5 embodiment, X_1 , X_2 , X_3 , X_4 , and X_5 form a pyrazolinone,
 thiophene, pyrazole-N-oxide, 2-amino-5-aryl-thiazole,
 tetrahydrofuran, cyclopentadienone, or a N-
 hydroxypyrrolidine ring. In yet another embodiment, the
 ring is a pyrazolinone, pyrazole-N-oxide,
 10 cyclopentadienone, or a N-hydroxypyrrolidine. In still
 another embodiment, the ring is a pyrazolinone.

Another aspect of the invention embraces compounds
 that correspond to formula (II)



(II)

15 wherein:

X_2 and X_5 are as defined for compounds having formula
 (I);

X_1 , X_3 and X_4 are independently carbon or nitrogen;

X_9 is a direct bond or $-(CH_2)_m-$ where m is 1 or 2; and

20 Z_1 , Z_3 and Z_4 are as defined for compounds having
 formula (I).

In yet another embodiment for compounds having formula
 (II), each of X_1 , X_2 , X_3 , X_4 , and X_5 is carbon or nitrogen;

25 X_9 is selected from the group consisting of a direct
 bond, methylene, and ethylene;

Z_1 is selected from the group consisting of C_1-C_8
 alkyl, C_2-C_8 alkenyl, and C_2-C_8 alkynyl, the alkyl, alkenyl,
 or alkynyl being optionally substituted at any
 substitutable position with a halogen;

Z₃ comprises a phenyl, furanyl or thienyl ring, the phenyl, furanyl or thienyl ring being optionally further substituted with fluorine or hydroxy;

5 Z₄ comprises a phenyl or thienyl ring having two substituents, R₄₂ and R₄₄, and two ring atoms each of which is in the beta position relative to the ring atom of Z₄ through which Z₄ is covalently bonded to X, wherein one of R₄₂ and R₄₄ is covalently bonded to one of said beta positions and the other of R₄₂ and R₄₄ is covalently bonded
10 to the other of said beta positions;

R₄₂ is amino; and

R₄₄ is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen, and an optionally substituted heteroatom selected from
15 nitrogen, oxygen, sulfur and phosphorus.

In yet another embodiment for compounds having formula (II), each of X₁, X₂, X₃, X₄, and X₅ is carbon or nitrogen;

X₉ is a direct bond;

Z₁ is selected from the group consisting of
20 cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, tert-butyl and sec-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy-carbonyl;


Z₃ is a phenyl ring, the phenyl ring being optionally
25 further substituted with fluorine or hydroxy;

Z₄ is a substituted phenyl ring as described in other embodiments for compounds having formula (II); and

R₄₄ is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, p-fluorobenzylamide, cyclobutylamide, -fluorobenzylamide, 1-
35 methylbenzylamide, sec-butylamide, benzylacetylamine, isobutylamide, sec-pentylamine, cyclopentylacetylamine, 1-

carboxyl-2-methylbutylamide, isobutylacylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy.

5 A further embodiment provides compounds having formula (II) that are represented by formula (IIa)



(IIa)

wherein:

10 each of Z_1 , Z_3 , and Z_4 are as defined for compounds having formulas (I) and (II).

In another embodiment for compounds having formula (IIa), Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted
15 at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl;

Z_3 is $-R_{300}C(=NR_{301})NR_{302}R_{303}$, wherein R_{300} is a phenyl ring, R_{301} , R_{302} , R_{303} are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom
20 selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur; and


Z_4 is a phenyl ring having formula (c) wherein:
 R_{42} is amino;

25 R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine,
30 methoxyethylamide, 1-carboxylbenzylamide, *p*-

fluorobenzylamide, cyclobutylamide, -fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacylamine, isobutylamide, *sec*-pentylamine, cyclopentylacylamine, 1-carboxyl-2-methylbutylamide, isobutylacylamine,
 5 isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and

R_{41} , R_{43} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl,
 10 substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.

A further embodiment provides compounds having formula (II) that are represented by formula (IIb)



(IIb)

15 wherein:

each of Z₁, Z₃, and Z₄ are as defined for compounds having formulas (I) and (II).

In an alternative for compounds having formula (IIb),
 20 Z₁ is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl;

Z₃ is $-R_{300}C(=NR_{301})NR_{302}R_{303}$, wherein R_{300} is a phenyl ring,
 25 R_{301} , R_{302} , R_{303} are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen,
 30 phosphorus and sulfur; and


Z_4 is a phenyl ring having formula (c) wherein:

R_{42} is amino;

R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and

R_{41} , R_{43} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.

Yet a further embodiment provides compounds having formula (II) that are represented by formula (IIc)



(IIc)

wherein:

each of Z_1 , Z_3 , and Z_4 are as defined for compounds having formulas (I) and (II).

In an alternative for compounds having formula (IIc), Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted at any substitutable

position with fluorine, hydroxy, carboxy, or alkoxy-carbonyl;

Z_3 is $-R_{300}C(=NR_{301})NR_{302}R_{303}$, wherein R_{300} is a phenyl ring, R_{301} , R_{302} , R_{303} are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur; and


Z_4 is a phenyl ring having formula (c) wherein:

R_{42} is amino;

R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, -fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and

R_{41} , R_{43} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.

Still a further embodiment provides compounds having formula (II) that are represented by formula (IIId)



(IIId)

wherein:

each of Z_1 , Z_3 , and Z_4 are as defined for compounds having formulas (I) and (II).

5 In an alternative for compounds having formula (IIId), Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or
10 alkoxyacetyl;


Z_3 is $-R_{300}C(=NR_{301})NR_{302}R_{303}$, wherein R_{300} is a phenyl ring, R_{301} , R_{302} , R_{303} are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom
15 selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur; and

Z_4 is a phenyl ring having formula (c) wherein:
 R_{42} is amino;

R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-
20 fluorobenzylamide, cyclobutylamide, *p*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide,
25 isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and

R_{41} , R_{43} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen,
35 nitrogen, and sulfur.

A further aspect of the invention embraces compounds having formula (III)



(III)

wherein:

- 5 each of X_1 , X_2 , X_3 , X_4 , and X_5 form any of the 5-membered heterocyclic or carbocyclic rings indicated in Table 1 below and are selected from pyrazolinone, pyrrole, thiophene, pyrazole-N-oxide, 1-amino-pyrazole, 1,3,4-triazole, 2-amino-4-aryl-thiazole, 2-amino-5-aryl-thiazole,
- 10 pyrrolidine, 2-amino-5-aryl-oxazole, 3-amino-pyrazole, 2-amio-4-aryl-oxazole, tetrahydrofuran, cyclopentadienone, and N-hydroxypyrrolidine; and

Z_1 and R_{44} are any of the groups detailed in Table 1 below.

TABLE 1		
Core	Z ₁	R ₄₄
pyrazolinone	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
pyrrole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
thiophene	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
pyrazole-N-oxide	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
1-amino-pyrazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
1,3,4-triazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
2-amio-4-aryl-thiazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2-methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2-cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoromethyl, carboxyl, carbmethoxy, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, p-fluorobenzylamide, cyclobutylamide, 1-methylbenzylamide, sec-butylamide, benzylacylamine, isobutylamide, sec-pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
2-amino-5-aryl-thiazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2-methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2-cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoromethyl, carboxyl, carbmethoxy, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, p-fluorobenzylamide, cyclobutylamide, 1-methylbenzylamide, sec-butylamide, benzylacylamine, isobutylamide, sec-pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
pyrrolidine	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
2-amino-5-aryl-oxazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2-methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2-cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoromethyl, carboxyl, carbmethoxy, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, p-fluorobenzylamide, cyclobutylamide, 1-methylbenzylamide, sec-butylamide, benzylacylamine, isobutylamide, sec-pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
3-amino-pyrazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine


TABLE 1		
Core	Z ₁	R ₄₄
2-amino-4-aryl-oxazole	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2-methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2-cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoromethyl, carboxyl, carbmethoxy, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, p-fluorobenzylamide, cyclobutylamide, 1-methylbenzylamide, sec-butylamide, benzylacylamine, isobutylamide, sec-pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
tetrahydrofuran	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine


TABLE 1		
Core	Z ₁	R ₄₄
cyclopentadienone	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2- methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2- cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoro- methyl, carboxyl, carbmethoxy, amino, 3- aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1- carboxylbenzylamide, p- fluorobenzylamide, cyclobutylamide, 1- methylbenzylamide, sec- butylamide, benzylacylamine, isobutylamide, sec- pentylamine, or cyclopentylacylamine

TABLE 1		
Core	Z ₁	R ₄₄
N-hydroxypyrrolidine	alkyl, substituted alkyl, phenyl, substituted phenyl, cycloalkyl, or substituted cycloalkyl	hydroxy, 1-carboxyl-2-methylbutylamide, isobutylsulfonyl, isobutylacylamine, trifluoromethyl, isobutylsulfoxyl, carboxamidobenzyl, 2-cyclohexylamide, carboxamidobutyl-2-yl, methoxyl, isobutyramido, sulfonamide, isobutoxy, isobutylsulfonamide, carboethoxy, aminoacyltrifluoromethyl, carboxyl, carbmethoxy, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, p-fluorobenzylamide, cyclobutylamide, 1-methylbenzylamide, sec-butylamide, benzylacylamine, isobutylamide, sec-pentylamine, or cyclopentylacylamine


The structure for each ring formed by X₁, X₂, X₃, X₄, and X₅ identified in Table 1 is shown below. Additionally, the structure for each R₄₄ group identified in Table 1 is shown below.




PYRAZOLINONE




PYRROLE




THIOPHENE




PYRAZOLE-N-OXIDE




1-AMINO-PYRAZOLE




1,3,4-TRIAZOLE




2-AMINO-4-ARYL-THIAZOLE




2-AMINO-5-ARYL-THIAZOLE




PYRROLIDINE




2-AMINO-5-ARYL-OXAZOLE




3-AMINO-PYRAZOLE




2-AMINO-4-ARYL-OXAZOLE




TETRAHYDROFURAN




CYCLOPENTADIENONE




N-HYDROXYPYRROLIDINE




hydroxy




isobutylsulfonyl




trifluoromethyl




isobutoxy




carboethoxy




carboxyl




carboxamidobenzyl




carboxamidobutyl-2-yl




isobutyramido




phenethylamine




isobutylamine




methoxyethylamide




amino




3-aminomethylthiophene




benzylamine




1-carboxylbenzylamide
R or S ISOMERS




p-fluorobenzylamide




cyclobutylamide




benzylacetylamine




isobutylamide




sec-pentylamine




m-fluorobenzylamide




1-methylbenzylamide
RACEMIC or R or S




sec-butylamide
R or S ISOMERS




isobutylsulfoxyl




2-cyclohexylamide
R or S ISOMERS




methoxyl




cyclopentylacetylamine




1-carboxyl-2-methylbutylamine




isobutylacetylamine




sulfonamide



isobutylsulfonamide




aminoacyltrifluoromethyl



carbmethoxy

Another aspect of the invention embraces compounds that correspond to formula (IV)



(IV)


wherein:

Z₁ is isopropyl, cyclopropyl, cyclobutyl or cyclopentyl optionally substituted by fluorine, hydroxy, carboxy, or alkoxy carbonyl;

5 Z₃ is -R₃₀₀C(=NR₃₀₁)NR₃₀₂R₃₀₃, wherein R₃₀₀ is a 6-membered carbocyclic aromatic ring, R₃₀₁, R₃₀₂, R₃₀₃ are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of
10 oxygen, nitrogen, phosphorus and sulfur; and

R₄₄ is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene,
15 benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-
20 carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy.

A further embodiment provides compounds having formula
25 (IV) that are represented by formula (IVa)




wherein:

Z_1 is isopropyl, cyclopropyl, cyclobutyl or cyclopentyl optionally substituted by fluorine, hydroxy, carboxy, or alkoxy carbonyl;

5 R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine,
 10 methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine,
 15 isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and

R_{310} and R_{311} are independently selected from the group consisting of hydrogen, fluorine, hydroxy, alkoxy, and
 20 carboxy, provided at least one of R_{310} and R_{311} is other than fluorine and hydrogen.

Yet another embodiment provides compounds having formula (IV) that are represented by formula (IVb)



(IVb)


wherein:

Z_1 is isopropyl, cyclopropyl, cyclobutyl or cyclopentyl optionally substituted by fluorine, hydroxy, carboxy, or alkoxy carbonyl;

5 Z_3 corresponds to formula (a) detailed above; and

R_{440} is C_1 - C_6 alkyl, aryl, aralkyl, carboxy, hydroxy or carboxyalkyl, wherein the alkyl, aryl, aralkyl, carboxy, hydroxy or carboxyalkyl is optionally further substituted by fluorine.

10 Still another embodiment provides compounds having formula (IV) that are represented by formula (IVc)



(IVc)


wherein:

15 Z_1 is isopropyl, cyclopropyl, cyclobutyl or cyclopentyl optionally substituted by fluorine, hydroxy, carboxy, or alkoxy carbonyl;

Z_3 corresponds to formula (a) detailed above ; and

20 R_{440} is C_1 - C_6 alkyl, aryl, aralkyl, carboxy, or carboxyalkyl, wherein the alkyl, aryl, aralkyl, carboxy, or carboxyalkyl is optionally further substituted by fluorine.

A further aspect of the invention embraces compounds having formula (V)



(V)

wherein:

Z₁, Z₃, Z₄, L₃, X₂, X₃, and X₄ are as defined for any of formulas (I) - (IV);

5 X₁ and X₅ are independently carbon and nitrogen;
 X₇ and X₈ are independently a covalent bond, carbon, nitrogen, oxygen or sulfur;


X₉ is carbon substituted with a methylene group or carbon substituted with an ethylene group wherein said
 10 methylene or ethylene group covalently links X₉ and Z₁;

Z₂, when present, is a hydrogen bond acceptor covalently or datively bonded to X₂;

n is 0 to 2; and

R₇₀ and R₈₀ are independently selected from the group
 15 consisting of hydrogen, halogen, amino, hydrocarbyl, substituted hydrocarbyl, aryl, wherein aryl is phenyl either unsubstituted or substituted with hydroxy, amino, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or halogen provided that R₇₀ is not present when X₇ is a bond and R₈₀ is not present when X₈ is a
 20 bond; or R₇₀ and R₈₀, along with the ring atoms to which each is attached, form a 5 or 6 membered saturated ring.

Yet another aspect of the invention provides compounds that are intermediates in the synthesis of compounds having formulas (I) - (V). These intermediate compounds correspond
 25 to formula (VI)



(VI)

wherein:

X_1 , X_2 , X_3 , and X_5 are members of a heterocyclic or carbocyclic ring;

5 X_1 , X_2 , and X_3 are independently carbon or nitrogen;

X_5 is selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, provided that when X_5 is carbon, it is $-CH=$, $-C(F)=$, or $-C(Br)=$;


10 T_3 is selected from the group consisting of hydroxy, alkoxy, substituted alkoxy, and substituted amino;

T_4 is selected from the group consisting of $-Cl$, $-Br$, $-I$, $-S(CH_3)$, and $-OSO_2(CF_3)$;

Z_1 is selected from the group consisting of hydrocarbyl, and substituted hydrocarbyl; and

15 Z_2 , when present, is a hydrogen bond acceptor covalently or datively bonded to X_2 .

Yet another aspect of the invention provides compounds that are intermediates in the synthesis of compounds having formulas (I)-(V). These intermediate compounds correspond to formula (VII)



(VII)

wherein:

X_1 , X_2 , X_3 , and X_5 are members of a heterocyclic or aromatic core ring;

X_1 , X_2 , and X_3 are independently carbon or nitrogen;

5 X_5 is selected from the group consisting of carbon, nitrogen, oxygen, and sulfur, provided that when X_5 is carbon, it is $-CH=$, $-C(F)=$, or $-C(Br)=$;

Z_1 is hydrocarbyl, or substituted hydrocarbyl;

10 Z_2 , when present, is a hydrogen bond acceptor covalently or datively bonded to X_2 ; and

Z_4 is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl and a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of the 5- or 6-membered heterocyclic or carbocyclic ring of Z_4 being
15 carbon, nitrogen, oxygen, or sulfur.

In yet a further embodiment, compounds having any of formulas (I)-(IV) are selected from the compounds in Table 2 below.

20

TABLE 2	
Compound No.	Compound
1	

TABLE 2	
Compound No.	Compound
2	 <chem>CC(C)Nc1nc(Cc2ccc(N)cc2C(F)(F)F)c3c1CC(=O)Nc4ccc(C(=N)N)cc4</chem>
3	 <chem>CC(C)Nc1sc(Cc2ccc(N)cc2C(F)(F)F)c3c1CC(=O)Nc4ccc(C(=N)N)cc4</chem>
4	 <chem>CC(C)Nc1nc(Cc2ccc(N)cc2S(=O)(=O)CC)nc1=OCC(=O)Nc3ccc(C(=N)N)cc3</chem>




TABLE 2	
Compound No.	Compound
5	 <chem>CC(C)Nc1nc(Cc2cc(C(=O)N)cc2N)cn1CC(=O)Nc3ccc(C(=N)N)cc3</chem>
6	 <chem>CC(C)Nc1nc(Cc2cc(N)cc2S(=O)(=O)C)cn1CC(=O)Nc3ccc(C(=N)N)cc3</chem>
7	 <chem>CC(C)Nc1nc(Cc2cc(N)cc2OC)s1CC(=O)Nc3ccc(C(=N)N)cc3</chem>





TABLE 2	
Compound No.	Compound
8	 <chem>CC(C)Nc1nc(s1)CC(=O)NCCc2ccc(cc2)C(=N)N</chem>
9	 <chem>CC(C)Nc1c[nH]c1CC(=O)NCCc2ccc(cc2)C(=N)N</chem>
10	 <chem>CC(C)Nc1nc2oc1c2CC(=O)NCCc3ccc(cc3)C(=N)N</chem>
11	 <chem>CC(C)Nc1c[nH]n1CC(=O)NCCc2ccc(cc2)C(=N)N</chem>

TABLE 2	
Compound No.	Compound
12	 <chem>CC(C)Nc1nc(Cc2cc(C(=O)Nc3ccc(C(=N)N)cc3)cc2)c4cc(N)c(OC)cc41</chem>
13	 <chem>CC1OC(CC1c2cc(N)c(C(=O)OC)cc2)C(=O)Nc3ccc(C(=N)N)cc3N4CC4</chem>
14	 <chem>CC1OC(CC1c2cc(N)c(C(=O)OC)cc2)C(=O)Nc3ccc(C(=N)N)cc3N(O)C4CC4</chem>

TABLE 2	
Compound No.	Compound
15	

Any compound corresponding to any of formulas (I) - (V), having one or more prodrug moieties as part of the molecule, can be converted under physiological conditions to the biologically active drug by a number of chemical and biological mechanisms. In general terms, these prodrug conversion mechanisms are hydrolysis, reduction, oxidation, and elimination. For illustrative purposes, the following paragraphs detail prodrugs in which the prodrug moiety is covalently bonded to the amidine group on Z₃, as depicted embodiments for each of formulas (I) - (V) above.

In one embodiment, conversion of the prodrug to the biologically active drug can be accomplished by hydrolysis of the prodrug moiety provided the prodrug moiety is chemically or enzymatically hydrolyzable with water. The reaction with water typically results in removal of the prodrug moiety and liberation of the biologically active drug. By way of example, a hydrolyzable prodrug derivative at the amidine group may be a carbonyl derivative such as N-acyl. Hydrolysis results in freeing the amidine group of the drug by removal of the acyl as carbon acid. Other suitable hydrolyzable prodrug derivatives include carbonyl, thiocarbonyl, imine, enamine, and oxygenated sulfur.

Yet another aspect of the invention provides conversion of the prodrug to the biologically active drug by reduction of the prodrug moiety. Typically in this embodiment, the

prodrug moiety is reducible under physiological conditions in the presence of a reducing enzymatic process. The reduction preferably results in removal of the prodrug moiety and liberation of the biologically active drug. An
5 example of a reducible prodrug derivative at the amidine group is an oxygen containing group in which an oxygen is directly attached to the amidine. Reduction results in freeing the amidine group of the drug by removal of oxygen as water or an alcohol. Generally speaking, other suitable
10 reducible prodrug derivatives include a nitrogen containing group, and a sulfur containing group, provided both nitrogen and sulfur are each preferably in their most reduced state.

In another embodiment, conversion of the prodrug to the biologically active drug can also be accomplished by
15 oxidation of the prodrug moiety. Typically in this embodiment, the prodrug moiety is oxidizable under physiological conditions in the presence of an oxidative enzymatic process. The oxidation preferably results in removal of the prodrug moiety and liberation of the
20 biologically active drug. An example of an oxidizable prodrug derivative at the amidine group is a hydrocarbyl containing unsaturation in the carbon beta to the carbon directly connected to the amidine group. Oxidation results in forming an oxygenated intermediate that breaks down,
25 thereby freeing the amidine group of the drug with concurrent hydrolysis of the oxygenated hydrocarbyl residue. Other suitable oxidizable prodrug derivatives of the amidine include saturated hydrocarbyl, unsaturated substituted hydrocarbyl, aryl, and aralkyl.

30 A further aspect of the invention encompasses conversion of the prodrug to the biologically active drug by elimination of the prodrug moiety. Generally speaking, in this embodiment the prodrug moiety is removed under physiological conditions with a chemical or biological
35 reaction. The elimination results in removal of the prodrug moiety and liberation of the biologically active drug. By way of example, an eliminateable prodrug derivative at the

amidine group is a hydrocarbyl containing an unsaturated electron withdrawing group bonded to the carbon beta to the carbon directly connected to the amidine. More specifically, for illustration purposes and exemplification, the hydrocarbyl group could have a cyano group beta to the carbon directly bonded to the amidino group. Elimination results in the freeing of the amidine group of the drug with concurrent removal of the unsaturated hydrocarbyl residue derived from the prodrug moiety. Other suitable eliminateable prodrug derivatives of the amidine include a hydrocarbyl substituted at the beta carbon with carbonyl, alkoxy carbonyl, amidocarbonyl, nitro, or sulfonyl or an alkyl group substituted with oxygen, nitrogen or sulfur at the carbon directly bonded to the amidine group.

Any compound of the present invention corresponding to formulas (I)-(V) may undergo any combination of the above detailed mechanisms to convert the prodrug to the biologically active compound. For example, a particular compound may undergo hydrolysis, oxidation, elimination, and reduction to convert the prodrug to the biologically active compound. Equally, a particular compound may undergo only one of these mechanisms to convert the prodrug to the biologically active compound.

The compounds of the present invention can exist in tautomeric, geometric or stereoisomeric forms. The present invention contemplates all such compounds, including cis- and trans-geometric isomers, E- and Z-geometric isomers, R- and S-enantiomers, diastereomers, d-isomers, l-isomers, the racemic mixtures thereof and other mixtures thereof, as falling within the scope of compounds having any of formulas (I)-(V). Pharmaceutically acceptable salts of such tautomeric, geometric or stereoisomeric forms are also included within the invention. The terms "cis" and "trans", as used herein, denote a form of geometric isomerism in which two carbon atoms connected by a double bond will each have a hydrogen atom on the same side of the double bond ("cis") or on opposite sides of the double bond ("trans").

Some of the compounds described contain alkenyl groups, and are meant to include both cis and trans or "E" and "Z" geometric forms. Furthermore, some of the compounds described contain one or more stereocenters and are meant to include R, S, and mixtures or R and S forms for each stereocenter present.

Also included in the family of compounds having any of formulas (I)-(V) are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salt" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that it is pharmaceutically acceptable. Suitable pharmaceutically-acceptable acid addition salts of the compounds may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethylsulfonic, benzenesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of the compounds include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, choline, chlorprocaine, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procain. All of these salts may be prepared by conventional means from the corresponding compound by reacting, for example, the appropriate acid or base with the selected compound of any of formulas (I)-(V).

The present invention also comprises a pharmaceutical composition comprising a therapeutically-effective amount of the compound of the invention in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent. Pharmaceutical compositions of the present invention can comprise the active compounds of formulas (I)-(V) in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended.

The active compounds and composition may, for example, be administered orally, intravascularly, intraperitoneally, subcutaneously, intramuscularly, ocularly, or topically. For treating ocular build up of fibrin, the compounds may be administered intraocularly or topically as well as orally or parenterally.

The compounds can be administered in the form of a depot injection or implant preparation which may be formulated in such a manner as to permit a sustained release of the active ingredient. The active ingredient can be compressed into pellets or small cylinders and implanted subcutaneously or intramuscularly as depot injections or implants. Implants may employ inert materials such as biodegradable polymers or synthetic silicones, for example, silastic, silicone rubber or other silicon containing polymers.

Moreover, the compounds can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds may also be delivered by the use of

monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran

5

copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds may be coupled to a class of biodegradable polymers useful in achieving controlled

10

15

release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihdropyrans, polycyanoacrylates and cross linked or amphiphathic block copolymers of hydrogels.

For oral administration, the pharmaceutical composition may be in the form of, for example, tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixers, tinctures, suspensions, liquids including syrups, and emulsions. The pharmaceutical composition is preferably made in the form of a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by

20

25

injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier.

The amount of therapeutically active compounds which are administered and the dosage regimen for treating a disease condition with the compounds and/or compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely.

30

35

The pharmaceutical compositions may contain active ingredients in the range of about 0.1 to 2000 mg, and preferably in the range of about 0.5 to 500 mg. A daily

dose of about 0.01 to 100 mg/kg body weight, and preferably between about 0.5 and about 20 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day.

5 The compounds may be formulated in topical ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active
10 ingredients may be employed with either paraffinic or a water-miscible ointment base.

 Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at
15 least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through
20 the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be accomplished using a patch either of
25 the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the
30 active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane.

 The oily phase of the emulsions of this invention may
35 be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or

an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as diisoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

For therapeutic purposes, the active compounds of the present invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered *per os*, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin,

acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided
5 in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules
10 having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or
15 various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

As a further embodiment, compounds having formula (I)-(V) or a pharmaceutically-acceptable salt thereof comprise a
20 treatment and prophylaxis for thrombotic events resulting from coronary artery disease, cerebrovascular disease and other coagulation cascade related disorders in a subject. The treatment comprises administering to the subject having such disorder a therapeutically-effective amount of
25 compounds having formulas (I)-(V) or a pharmaceutically-acceptable salt thereof.

In another aspect of the invention, the compounds or a pharmaceutically-acceptable salt thereof can also be used whenever inhibition of blood coagulation is required such as
30 to prevent coagulation of stored whole blood and to prevent coagulation in other biological samples for testing or storage. Thus coagulation inhibitors of the present invention can be added to or contacted with stored whole blood and any medium containing or suspected of containing
35 plasma coagulation factors and in which it is desired that blood coagulation be inhibited, e.g. when contacting the mammal's blood with material selected from the group

consisting of vascular grafts, stents, orthopedic prothesis, cardiac prothesis, and extracorporeal circulation systems.

Compounds of Formula (I)-(V) are capable of inhibiting activity of serine proteases related to the coagulation cascade. Thus, these compounds could be used in the manufacture of a medicament as a method for the prophylactic or therapeutic treatment of diseases mediated by coagulation cascade serine proteases, such as inhibiting the formation of blood platelet aggregates, inhibiting the formation of fibrin, inhibiting thrombus formation, and inhibiting embolus formation in a mammal, in blood, in blood products, and in mammalian organs. The compounds also can be used for treating or preventing unstable angina, refractory angina, myocardial infarction, transient ischemic attacks, atrial fibrillation, thrombotic stroke, embolic stroke, deep vein thrombosis, disseminated intravascular coagulation, ocular build up of fibrin, and reocclusion or restenosis of recanalized vessels in a mammal. Moreover, the compounds also can be used to study the mechanism of action of coagulation cascade serine proteases to enable the design of better inhibitors and development of better assay methods. The compounds would be also useful in prevention of cerebral vascular accident (CVA) or stroke.

In practicing the methods of the present invention for the treatment and prevention of a variety of thrombotic conditions including coronary artery and cerebrovascular disease, the compounds and pharmaceutical compositions are administered alone or in combination with one another, or in combination with other therapeutics or *in vivo* diagnostic agents. In another aspect, the compounds can also be co-administered with suitable anti-platelet aggregation agents, including, but not limited to aspirin, ticlopidine, or clopidrogel, fibrinogen receptor antagonists (e.g. to treat or prevent unstable angina or to prevent reocclusion after angioplasty and restenosis), anti-coagulants such as aspirin, warfarin or heparins, thrombolytic agents such as plasminogen activators or streptokinase to achieve

synergistic effects in the treatment of various pathologies, lipid lowering agents including antihypercholesterolemics (e.g. HMG CoA reductase inhibitors such as mevastatin, lovastatin, simvastatin, pravastatin, and fluvastatin, HMG CoA synthetase inhibitors, etc.), anti-diabetic drugs, or other cardiovascular agents (e.g. loop diuretics, thiazide type diuretics, nitrates, aldosterone antagonists (e.g. spironolactone and epoxy-mexlerenone), angiotensin converting enzyme (e.g. ACE) inhibitors, angiotensin II receptor antagonists, beta-blockers, antiarrhythmics, anti-hypertension agents, and calcium channel blockers to treat or prevent atherosclerosis. By way of example, patients suffering from coronary artery disease, and patients subjected to angioplasty procedures, would benefit from coadministration of fibrinogen receptor antagonists and coagulation cascade inhibitors of the present invention. Also, coagulation cascade inhibitors could enhance the efficiency of tissue plasminogen activator-mediated thrombolytic reperfusion.

Typical doses of compounds of the present invention with other suitable anti-platelet agents, anticoagulation agents, cardiovascular therapeutic agents, or thrombolytic agents may be the same as those doses of coagulation cascade inhibitors administered without coadministration of additional anti-platelet agents, anticoagulation agents, cardiovascular therapeutic agents, or thrombolytic agents, or may be substantially less than those doses of coagulation cascade inhibitors administered without coadministration of additional anti-platelet agents, anticoagulation agents, cardiovascular therapeutic agents, or thrombolytic agents, depending on a patient's therapeutic needs.

The present methods preferably employ prodrug compounds that when converted to the biologically active compound selectively inhibit human TF-VIIA over the inhibition of both human Thrombin II and human factor Xa. Preferably, the compounds have a human TF-VIIA IC_{50} of less than 0.5 mM and also have a selectivity ratio of TF-VIIA inhibition over

both human Thrombin II and human factor Xa inhibition of at least 10, and more preferably at least 100. Even more preferably, the compounds have a human TF-VIIA IC₅₀ of less than 0.1 mM and also have a selectivity ratio of TF-VIIA inhibition over both human Thrombin II and human factor Xa inhibition of at least 1000, and most preferably at least 10,000.

All mentioned references are incorporated by reference as if here written.

Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations. The following examples are provided to illustrate the present invention and are not intended to limit the scope thereof. Without further elaboration, it is believed that one skilled in the art can, using the preceding descriptions, utilize the present invention to its fullest extent. Therefore, the following preferred specific embodiments are to be construed as merely illustrative and not limitative of the remainder of the disclosure in any way whatsoever. Compounds containing multiple variations of the structural modifications illustrated in the schemes or the following Examples are also contemplated. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds.

One skilled in the art may use these generic methods to prepare the following specific examples, which have been or may be properly characterized by ¹H NMR, mass spectrometry, elemental composition, and similar procedures. These compounds also may be formed *in vivo*. The following examples contain detailed descriptions of the methods of preparation of compounds having each of formulas (I)-(V). These detailed descriptions fall within the scope and are presented for illustrative purposes only and are not intended as a restriction on the scope of the invention. All parts are by weight and temperatures are Degrees

centigrade unless otherwise indicated.

GENERAL SYNTHETIC PROCEDURES AND SPECIFIC EXAMPLES

5 The compounds of the present invention can be synthesized, for example, according to the following procedures and Schemes given below.

Abbreviations used in the schemes and tables include:
"AA" represents amino acids, "AcCN" represents acetonitrile,
"AcOH" represents acetic acid, "BINAP" represents 2,2'-
10 bis(diphenylphosphino)-1,1'-binaphthyl, "BnOH" represents benzyl alcohol, "BnCHO" represents 2-phenylethanal, "BnSO₂Cl" represents benzylsulfonyl chloride, "Boc" represents tert-butyloxycarbonyl, "BOP" represents benzotriazol-1-yl-oxy-tris-(dimethylamino), "bu" represents
15 butyl, "dba" represents dibenzylidene-acetone, "DCC" represents 1,3-dicyclohexylcarbodiimide, "DCM" represents dichloromethane or methylene chloride, "DIBAH" or "DIBAL" represents diisobutylaluminum hydride, "DMF" represents dimethylformamide, "DMSO" represents dimethylsulfoxide,
20 "DPPA" represents diphenylphosphoryl azide", "EDC" represents 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride, "Ex. No." represents Example Number, "Fmoc" represents 9-fluorenylmethoxycarbonyl, "HOBT" represents hydroxybenzotriazole", "LDA" represents lithium
25 diisopropylamide, "MW" represents molecular weight, "NMM" represents N-methylmorpholine, "Ph" represents phenyl or aryl, "PHTH" represents a phthaloyl group, "pnZ" represents 4-nitrobenzyloxy-carbonyl, "PTC" represents a phase transfer catalyst, "py" represents pyridine, "R" represents a
30 hydrocarbyl or a substituted hydrocarbyl, "RNH₂" represents a primary organic amine, "SEM" represents 2-(trimethylsilyl)ethoxy-methyl chloride, "p-TsOH" represents paratoluenesulfonic acid, "TBAF" represents tetrabutylammonium fluoride, "TBTU" represents 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyl uronium
35 tetrafluoroborate, "TEA" represents triethylamine, "TFA" represents trifluoroacetic acid, "THF" represents

tetrahydrofuran, "TMS" represents trimethylsilyl, "TMSCN" represents trimethylsilyl cyanide, and "Cbz" or "Z" represents benzyloxycarbonyl "X" represents a halogen and typically is a bromine or chlorine. As used in the schemes and examples, Z₁, Z₃, Z₄, R₄₄, R₈₀, R₃₀₁, R₃₀₂, R₃₀₃, R₃₀₄, R₃₀₅, R₃₀₆, R₃₀₇, R₃₀₈, R₃₀₉, R₃₁₀, R₃₁₁, and X₇, along with any other variable depicted, encompasses every group described for each particular variable for each embodiment of compounds having formulas (I)-(V) as detailed herein. .

A specific synthetic process, useful in the preparation of many of the heterocyclic compounds of the present invention, is the arylation or heteroarylation of an intermediate compound characterized by having a suitable leaving group on a sp² hybridized carbon of a heterocyclic ring. In the product of the reaction, the leaving group is replaced by an aryl group or a heteroaryl group. Suitable leaving groups for the reaction include chloro, bromo, iodo, methylthio, and triflates. The heterocyclic ring with the leaving group will preferably have an acetic acid group or a derivative thereof bonded to a ring atom alpha to the bromo and a substituted or unsubstituted amino group bonded to a ring atom that is both beta to the carbon having the acetic acid group and gamma to the bromo substituted ring carbon. The aryl group that is reacted at the sp² hybridized carbon is generally an aryl boronic acid or an ester of the aryl boronic acid; similarly, heteroaryl boronic acids or esters of these boronic acids can be used in the same manner as aryl boronates. The aryl and heteroaryl boronates may be substituted or unsubstituted. The aryl or heteroaryl becomes bonded to the sp² hybridized carbon at the point at which the boron was attached to the aryl or heteroaryl ring. Aryl and heteroaryl organoSn compounds can also be used instead of the corresponding boronates.

Suitable reaction conditions for carrying out this transformation include:

1. Pd[P(phenyl)₃]₄, 2M Na₂CO₃, 60-75°C, dimethoxyethane (DME), H₂O, N₂;


2. Pd[P(phenyl)₃]₄, Cs₂CO₃, dioxane, 100°C;
3. Pd[P(phenyl)₃]₄, Cu(I)-2-thiophenecarboxylate, 70-75°C, anhydrous THF, argon; and
4. Z4-Sn(n-butyl)₃, Pd[P(phenyl)₃]₄, LiCl, anhydrous
5 dioxane, 85°C, argon or N₂.

The organo palladium (Pd[P(phenyl)₃]₄) compound is used catalytically in a ratio of 1 to 40 mole %. The carbonate base is normally used in an excess of 1.2 to 2 molar equivalents. Suitable solvents include dimethoxyethane
10 (DME), dioxane, 1-propanol, and tetrahydrofuran. The temperature of the reaction is normally in the range of from about 50 to 100°C. Cu(I)-2-thiophenecarboxylate (Cu(I)-TC) is normally used in a mole % of 110-150.


15 Scheme G shows a specific application of this specific synthetic process. Procedures for preparing the intermediate heterocyclic or cycloalkenyl ring compounds having a suitable leaving group on sp² hybridized carbon and useful as suitable intermediates in this specific synthetic process are given in the schemes and examples listed above.

20 Scheme J illustrates a general synthetic process for substitution at a nitrogen of the heterocyclic ring. The synthetic process applies whether the ring is 5- or 6-membered.



Scheme A:





Scheme A: (continued from page 77)



Scheme B:




Deprotection: H₂, Pd/C
or LiOH, H₂O, MeOH



Z = benzyloxycarbonyl


TBTU
DMF
DIEA




H₂, Pd/C or
LiOH, H₂O,
and MeOH

See Page 80




Scheme B: (continued from page 79)




Scheme C:




Z = benzyloxycarbonyl




Scheme C: (continued from page 81)




Scheme D:




Scheme D: (continued from page 83)




Scheme E:



Scheme E: (continued from page 85)




Scheme F:






See Page 88


Scheme F: (continued from page 87)




Z = benzyloxycarbonyl

5







Scheme G:


Scheme G: (continued from page 89)



Z = benzyloxycarbonyl



Scheme H:




5





See Page 92

TBTU
DMF
DIEA

Z = benzyloxycarbonyl


Scheme H: (continued from page 91)



Scheme I:

See Page 94


Scheme I: (continued from page 93)



Claims:

WHAT IS CLAIMED IS:

1. A compound having the structure



wherein:

X comprises a 5-membered heterocyclic or carbocyclic ring, the ring atoms being X_1 , X_2 , X_3 , X_4 , and X_5 , wherein X_1 , X_3 , and X_4 , are independently carbon or nitrogen and X_2 and X_5 are independently carbon, nitrogen, oxygen or sulfur;

L_1 , L_3 and L_4 are linkages through which Z_1 , Z_3 , and Z_4 , respectively, are covalently bonded to different ring atoms of the 5-membered heterocyclic or carbocyclic ring of X, wherein Z_1 is covalently bonded to X_1 , Z_3 is covalently bonded to X_3 , and Z_4 is covalently bonded to X_4 , each of L_1 , L_3 and L_4 independently being a covalent bond or comprising one or more atoms through which Z_1 , Z_3 , and Z_4 are covalently bonded to X_1 , X_3 and X_4 , respectively;

Z_1 is hydrocarbyl or substituted hydrocarbyl;

Z_3 comprises a 5- or 6-membered heterocyclic or aromatic ring substituted with an amidine group, the ring atoms of the 5- or 6-membered heterocyclic or aromatic ring of Z_3 being carbon, sulfur, nitrogen, or oxygen, wherein the 5- or 6-membered ring is optionally substituted at any position with halogen, alkyl or hydroxy; and

Z_4 comprises a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of the 5- or 6-membered heterocyclic or carbocyclic ring of Z_4 being carbon, nitrogen, oxygen, or sulfur.

2. The compound of claim 1 wherein:

each of X_1 , X_2 , X_3 , X_4 , and X_5 are as defined in claim 1;

L_1 is $-X_9NH-$ wherein X_9 is covalently bonded directly to Z_1 and X_9 is a direct bond or $-(CH_2)_m-$ wherein m is 1 to 5;

5 L_3 is a glycine derivative;

L_4 is a direct bond;

Z_1 is selected from the group consisting of C_1 - C_8 alkyl, C_2 - C_8 alkenyl, and C_2 - C_8 alkynyl, the alkyl, alkenyl, or alkynyl being optionally substituted at any substitutable position with a halogen;

10

Z_3 comprises a phenyl, furanyl or thienyl ring, the phenyl, furanyl or thienyl ring being optionally further substituted with fluorine or hydroxy;

Z_4 comprises a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of Z_4 being Z_{40} , Z_{41} , Z_{42} , Z_{44} and Z_{45} when Z_4 is a 5-membered ring and Z_{40} , Z_{41} , Z_{42} , Z_{43} , Z_{44} and Z_{45} when Z_4 is a 6-membered ring, Z_{40} , Z_{41} , Z_{42} , Z_{43} , Z_{44} and Z_{45} , being carbon, nitrogen, oxygen or sulfur, Z_{40} being the ring atom through which Z_4 is attached to the heterocyclic ring, Z_{41} and Z_{45} each being in an alpha position relative to Z_{40} , Z_{42} and Z_{44} each being in a beta position relative to Z_{40} , Z_{43} being in the gamma position relative to Z_{40} when Z_4 is a 6-membered ring, Z_4 having a substituent R_{42} covalently attached to Z_{42} , and a second substituent bonded to one of Z_{41} , Z_{43} , Z_{44} , or Z_{45} , the substituent being R_{41} when bonded to Z_{41} , the substituent being R_{43} when bonded to Z_{43} , the substituent being R_{44} when bonded to Z_{44} , and the substituent being R_{45} when bonded to Z_{45} ;

15

20

25

R_{42} is amino; and

R_{41} , R_{43} , R_{44} and R_{45} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen, or an optionally substituted heteroatom selected from nitrogen, oxygen, sulfur and phosphorus, provided at least one of R_{41} , R_{43} , R_{44} or R_{45} is other than hydrogen.

3. The compound of claim 2 wherein:

X_9 is a direct bond;

L_3 is $-\text{CH}_2\text{CONHCH}_2-$ and Z_3 is covalently bonded to the methylene bonded to the amine nitrogen of L_3 ;

5 Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, tert-butyl and sec-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxycarbonyl;

10 Z_3 comprises a phenyl ring, the phenyl ring being optionally further substituted with fluorine or hydroxy;

Z_4 comprises a phenyl or thienyl ring having two substituents, R_{42} and R_{44} , and two ring atoms each of which is in the beta position relative to the ring atom of Z_4 through which Z_4 is covalently bonded to X , wherein one of R_{42} and R_{44} is covalently bonded to one of the beta positions and the other of R_{42} and R_{44} is covalently bonded to the other of the beta positions;

R_{42} is amino; and

20 R_{44} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen, and an optionally substituted heteroatom selected from nitrogen, oxygen, sulfur and phosphorus.

4. The compound of claim 1 wherein L_1 is $-\text{X}_9\text{NH}-$ wherein X_9 is covalently bonded directly to Z_1 and X_9 is a direct bond or $-(\text{CH}_2)_m-$ wherein m is 1 to 5.

5. The compound of claim 1 wherein L_3 is selected from the group consisting of a glycine derivative, an alanine derivative, an amino derivative, and a sulfonyl derivative.


6. The compound of claim 5 wherein L_3 is a glycine derivative.

7. The compound of claim 6 wherein L_3 is $-\text{CH}_2\text{CONHCH}_2-$ and Z_3 is covalently bonded to the methylene bonded to the amine nitrogen of L_3 .

8. The compound of claim 1 wherein L_4 is selected from the group consisting of a direct bond, methylene, ethylene and an optionally substituted heteroatom selected from the group consisting of nitrogen, oxygen, sulfur and phosphorus.

9. The compound of claim 8 wherein L_4 is a direct bond.

10. The compound of claim 1 having the structure



wherein:

X_2 and X_5 are as defined in claim 1;

X_1 , X_3 and X_4 are independently carbon or nitrogen;

X_9 is a direct bond or $-(\text{CH}_2)_m-$ where m is 1 or 2; and

Z_1 , Z_3 and Z_4 are as defined in claim 1.

11. The compound of claim 10 wherein:

X_9 is selected from the group consisting of a direct bond, methylene, and ethylene;

Z_1 is selected from the group consisting of C_1 - C_8 alkyl, C_2 - C_8 alkenyl, and C_2 - C_8 alkynyl, the alkyl, alkenyl, or alkynyl being optionally substituted at any substitutable position with a halogen;

Z_3 comprises a phenyl, furanyl or thienyl ring, the phenyl, furanyl or thienyl ring being optionally further substituted with fluorine or hydroxy;

Z_4 comprises a phenyl or thienyl ring having two

substituents, R_{42} and R_{44} , and two ring atoms each of which is in the beta position relative to the ring atom of Z_4 through which Z_4 is covalently bonded to X, wherein one of R_{42} and R_{44} is covalently bonded to one of said beta positions and the other of R_{42} and R_{44} is covalently bonded to the other of said beta positions;

R_{42} is amino; and

R_{44} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen, and an optionally substituted heteroatom selected from nitrogen, oxygen, sulfur and phosphorus.

12. The compound of claim 11 wherein:

X_9 is a direct bond;

Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl;

Z_3 is a phenyl ring, the phenyl ring being optionally further substituted with fluorine or hydroxy;

Z_4 is a substituted phenyl ring; and

R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, -fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy.

13. The compound of claim 1 or 10 wherein:
X₁, X₂ and X₅ are carbon; and
X₃ and X₄ are nitrogen.
14. The compound of claim 1 or 10 wherein:
X₁, X₃, X₄ and X₅ are carbon; and
X₂ is nitrogen.
15. The compound of claim 1 or 10 wherein:
X₁, X₂, X₃, X₄ and X₅ are carbon.
16. The compound of claim 1 or 10 wherein:
X₁, X₄ and X₅ are carbon; and
X₂ and X₃ are nitrogen.
17. The compound of claim 1 or 10 wherein X₁, X₂, X₃, X₄, and X₅ are selected to provide a heterocyclic or carbocyclic ring selected from the group consisting of pyrazolinone, pyrrole, thiophene, pyrazole-N-oxide, 1-amino, pyrazole, 1,3,4-triazole, 2-amino-4-aryl-thiazole, 2-amino-5-aryl-thiazole, pyrrolidine, 2-amino-5-aryl-oxazole, 3-amino-pyrazole, 2-amio-4-aryl-oxazole, tetrahydrofuran, cyclopentadienone, and N-hydroxypyrrolidine.
18. The compound of 17 wherein the heterocyclic or carbocyclic ring is selected from the group consisting of cyclopentadienone, pyrazole-N-oxide, pyrazolinone and N-hydroxypyrrolidine.
19. The compound of claim 18 wherein the heterocyclic ring is pyrazolinone.
20. The compound of claim 2 or 10 wherein X₉ is a direct bond.
21. The compound of claim 2 or 10 wherein X₉ is methylene or ethylene.

22. The compound of claim 1 or 10 wherein Z_1 is selected from the group consisting of C_1-C_8 alkyl, C_2-C_8 alkenyl, and C_2-C_8 alkynyl, the alkyl, alkenyl, or alkynyl being optionally substituted at any substitutable position with a halogen.

23. The compound of claim 22 wherein Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, *tert*-butyl and *sec*-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl.


24. The compound of claim 23 wherein Z_1 is selected from the group consisting of cyclopropyl, isopropyl, cyclobutyl, isobutyl, and *sec*-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl.

25. The compound of claim 24 wherein Z_1 is isopropyl or cyclobutyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy carbonyl.

26. The compound of claim 1 or 10 wherein Z_3 comprises a phenyl, furanyl or thienyl ring, the phenyl, furanyl or thienyl ring being optionally further substituted with fluorine or hydroxy.

27. The compound of claim 26 wherein Z_3 is $-R_{300}C(=NH)NH_2$, wherein R_{300} is a 6-membered carbocyclic aromatic ring, the 6-membered ring being optionally substituted at any substitutable position with a halogen or hydroxy.

28. The compound of claim 27 wherein Z₃ is



wherein:

R₃₀₄ and R₃₀₆ are independently selected from the group consisting of hydrogen, fluorine, hydroxy, carboxy, hydrocarbyloxy, and alkoxy carbonyl; and

R₃₀₅ and R₃₀₇ are independently selected from the group consisting of hydrogen, fluorine, methoxy, hydroxy, and carboxy.

29. The compound of claim 1 or 10 wherein Z₃ is -R₃₀₀C(=NR₃₀₁)NR₃₀₂R₃₀₃, wherein R₃₀₀ is a 6-membered carbocyclic aromatic ring, R₃₀₁, R₃₀₂, R₃₀₃ are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur, provided at least one of R₃₀₁, R₃₀₂, R₃₀₃ is other than hydrogen.

30. The compound of claim 29 wherein Z₃ is a benzamidine derivative, which upon hydrolysis, oxidation, reduction or elimination forms a benzamidine.

31. The compound of claim 1 or 10 wherein Z₄ comprises a 5- or 6-membered heterocyclic or carbocyclic ring, the ring atoms of Z₄ being Z₄₀, Z₄₁, Z₄₂, Z₄₄ and Z₄₅ when Z₄ is a 5-membered ring and Z₄₀, Z₄₁, Z₄₂, Z₄₃, Z₄₄ and Z₄₅ when Z₄ is a 6-membered ring, Z₄₀, Z₄₁, Z₄₂, Z₄₃, Z₄₄ and Z₄₅, being carbon, nitrogen, oxygen or sulfur, Z₄₀ being the ring atom through which Z₄ is attached to the heterocyclic ring, Z₄₁ and Z₄₅ each being in an alpha position relative to Z₄₀, Z₄₂ and Z₄₄


each being in a beta position relative to Z_{40} , Z_{43} being in
 10 the gamma position relative to Z_{40} , when Z_4 is a 6-membered
 ring, Z_4 having a substituent R_{42} covalently attached to Z_{42} ,
 and a second substituent bonded to one of Z_{41} , Z_{43} , Z_{44} , or
 Z_{45} , the substituent being R_{41} when bonded to Z_{41} , the
 15 substituent being R_{43} when bonded to Z_{43} , the substituent
 being R_{44} when bonded to Z_{44} , and the substituent being R_{45}
 when bonded to Z_{45} ;

R_{42} is amino; and

R_{41} , R_{43} , R_{44} and R_{45} are independently hydrogen,
 hydrocarbyl, substituted hydrocarbyl, heterocyclo, halogen,
 or a substituted or unsubstituted heteroatom selected from
 nitrogen, oxygen, sulfur and phosphorus, provided at least
 one of R_{41} , R_{43} , R_{44} or R_{45} is other than hydrogen.

32. The compound of claim 31 wherein Z_4 is a
 substituted, 6-membered, carbocyclic aromatic ring.

33. The compound of claim 32 wherein Z_4 has the
 structure



wherein

R_{42} is amino;

5 R_{44} is selected from the group consisting of
 hydrocarbyl, substituted hydrocarbyl, halogen and an
 optionally substituted heteroatom selected from the group
 consisting of oxygen, nitrogen, and sulfur; and

10 R_{41} , R_{43} and R_{45} are independently selected from the
 group consisting of hydrogen, halogen, hydrocarbyl,
 substituted hydrocarbyl, and an optionally substituted

heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.


34. The compound of claim 33 wherein R_{44} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamido, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkyl, haloalkoxy, haloalkylthio, carboalkoxy, carboxy, carboxamidoalkyl, and carboxamidoalkylaryl.

35. The compound of claim 33 wherein R_{44} is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroaryl, heterocyclo, halogen, acetamido, guanidino, hydroxy, nitro, amino, amidosulfonyl, acylamido, hydrocarbyloxy, substituted hydrocarbyloxy, hydrocarbylthio, substituted hydrocarbylthio, hydrocarbylsulfonyl, and substituted hydrocarbylsulfonyl.

36. The compound of claim 33 wherein R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacylamine, isobutylamide, *sec*-pentylamine, cyclopentylacylamine, 1-carboxyl-2-methylbutylamide, isobutylacylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy.

37. The compound of claim 33 wherein each of R_{41} , R_{43} and R_{45} is hydrogen.

38. The compound of claim 31 wherein Z_4 has the structure



wherein:

5 Z_{40} , Z_{41} , Z_{42} , Z_{44} , and Z_{45} are independently selected from the group consisting of carbon, nitrogen, oxygen and sulfur;

R_{42} is amino;

10 R_{44} is selected from the group consisting of is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, halogen and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur; and

15 R_{41} and R_{45} are independently selected from the group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.


39. The compound of claim 38 wherein R_{44} is selected from the group consisting of hydrocarbyl, substituted hydrocarbyl, acetamido, alkoxy, hydroxy, amino, alkylsulfonyl, haloalkyl, haloalkoxy, haloalkylthio, 5 carboalkoxy, carboxy, carboxamidoalkyl, and carboxamidoalkylaryl.

40. The compound of claim 38 wherein R_{44} is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroaryl, heterocyclo, halogen, acetamido, guanidino, hydroxy, nitro, amino, amidosulfonyl, 5 acylamido, hydrocarbyloxy, substituted hydrocarbyloxy, hydrocarbylthio, substituted hydrocarbylthio, hydrocarbylsulfonyl, and substituted hydrocarbylsulfonyl.

41. The compound of claim 38 wherein R_{44} is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy.

42. The compound of claim 38 wherein each of R_{41} , R_{42} and R_{45} is hydrogen.


43. The compound of claim 10 having the structure



wherein:

each of Z_1 , Z_3 , and Z_4 are as defined in claim 10.


44. The compound of claim 10 having the structure



wherein:

each of Z_1 , Z_3 , and Z_4 are as defined in claim 10.


45. The compound of claim 10 having the structure



wherein:

each of Z_1 , Z_3 , and Z_4 are as defined in claim 10.

46. The compound of claim 10 having the structure



wherein:


each of Z_1 , Z_3 , and Z_4 are as defined in claim 10.

47. The compound of each of claims 43 to 46 wherein:

Z_1 is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, tert-butyl and sec-butyl optionally substituted at any substitutable position with fluorine, hydroxy, carboxy, or alkoxy-carbonyl;

Z_3 is $-\text{R}_{300}\text{C}(=\text{NR}_{301})\text{NR}_{302}\text{R}_{303}$, wherein R_{300} is a phenyl ring, R_{301} , R_{302} , R_{303} are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur; and

Z₄ is a phenyl ring having the structure



wherein:


- 15 R₄₂ is amino;
- R₄₄ is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene,
- 20 benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, -fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacylamine, isobutylamide, *sec*-pentylamine, cyclopentylacylamine, 1-
- 25 carboxyl-2-methylbutylamide, isobutylacylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and
- R₄₁, R₄₃ and R₄₅ are independently selected from the
- 30 group consisting of hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, and sulfur.

48. The compound of claim 47 wherein:

- Z₁ is selected from the group consisting of cyclopropyl, isopropyl, methyl, ethyl, cyclobutyl, isobutyl, and *sec*-butyl optionally substituted at any substitutable
- 5 position with fluorine, hydroxy, carboxy, or alkoxy carbonyl; and

 Z₃ is as defined in claim 28.

49. The compound of claim 10 having the structure



wherein:

Z₁ is isopropyl, cyclopropyl, cyclobutyl or cyclopentyl optionally substituted by fluorine, hydroxy, carboxy, or alkoxy carbonyl;


5

Z₃ is -R₃₀₀C(=NR₃₀₁)NR₃₀₂R₃₀₃, wherein R₃₀₀ is a 6-membered carbocyclic aromatic ring, R₃₀₁, R₃₀₂, R₃₀₃ are independently selected from the group consisting of hydrogen, halogen, optionally substituted hydrocarbyl, and an optionally substituted heteroatom selected from the group consisting of oxygen, nitrogen, phosphorus and sulfur; and

10

R₄₄ is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy.

50. The compound of claim 10 having the structure




wherein:

Z₁ is isopropyl, cyclopropyl, cyclobutyl or cyclopentyl optionally substituted by fluorine, hydroxy, carboxy, or alkoxy carbonyl;

R₄₄ is selected from the group consisting of hydroxy, isobutylsulfonyl, trifluoromethyl, carboxamidobenzyl, carboxamidobutyl-2-yl, isobutyramido, isobutoxy, carboethoxy, carboxyl, amino, 3-aminomethylthiophene, benzylamine, phenethylamine, isobutylamine, methoxyethylamide, 1-carboxylbenzylamide, *p*-fluorobenzylamide, cyclobutylamide, *m*-fluorobenzylamide, 1-methylbenzylamide, *sec*-butylamide, benzylacetylamine, isobutylamide, *sec*-pentylamine, cyclopentylacetylamine, 1-carboxyl-2-methylbutylamide, isobutylacetylamine, isobutylsulfoxyl, 2-cyclohexylamide, methoxy, sulfonamide, isobutylsulfonamide, aminoacyltrifluoromethyl, and carbmethoxy; and

R₃₁₀ and R₃₁₁ are independently selected from the group consisting of hydrogen, fluorine, hydroxy, alkoxy, and carboxy, provided at least one of R₃₁₀ and R₃₁₁ is other than fluorine and hydrogen.

51. The compound of claim 50 having the structure




wherein:

Z₁ is isopropyl, cyclopropyl, cyclobutyl or
cyclopentyl optionally substituted by fluorine, hydroxy,
5 carboxy, or alkoxy-carbonyl;

Z₃ is as defined in claim 28; and

R₄₄₀ is C₁-C₆ alkyl, aryl, aralkyl, carboxy, hydroxy or
carboxyalkyl, wherein the alkyl, aryl, aralkyl, carboxy,
hydroxy or carboxyalkyl is optionally further substituted by
10 fluorine.

52. The compound of claim 50 having the structure



wherein:

Z₁ is isopropyl, cyclopropyl, cyclobutyl or
cyclopentyl optionally substituted by fluorine, hydroxy,
5 carboxy, or alkoxy-carbonyl;

Z₃ is as defined in claim 28; and

R₄₄₀ is C₁-C₆ alkyl, aryl, aralkyl, carboxy, or carboxyalkyl, wherein the alkyl, aryl, aralkyl, carboxy, or carboxyalkyl is optionally further substituted by fluorine.

53. A composition for substantially inhibiting thrombotic conditions in blood comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

54. A method for substantially inhibiting thrombotic conditions in blood comprising adding to blood the composition of claim 53.

55. A method for substantially inhibiting formation of platelet aggregates in blood comprising adding to blood the composition of claim 53.

56. A method for substantially inhibiting thrombus formation in blood comprising adding to blood the composition of claim 53.

57. A method for treating or preventing venous thromboembolism and pulmonary embolism in a mammal comprising administering to the mammal the composition of claim 53.

58. A method for treating or preventing deep vein thrombosis in a mammal comprising administering to the mammal the composition of claim 53.

59. A method for treating or preventing cardiogenic thromboembolism in a mammal comprising administering to the mammal the composition of claim 53.

60. A method for treating or preventing thromboembolic stroke in a mammal comprising administering to the mammal the composition of claim 53.

61. A method for treating or preventing thrombosis associated with cancer and cancer chemotherapy in a mammal comprising administering to the mammal the composition of claim 53.

62. A method for treating or preventing unstable angina in a mammal comprising administering to the mammal the composition of claim 53.

63. A method for inhibiting thrombus formation in blood comprising adding to blood the composition of claim 53 along with a fibrinogen receptor antagonist.