

Boehringer Ingelheim International GmbH

BENZODIOXANE INHIBITORS OF LEUKOTRIENE PRODUCTION**Field of the Invention**

This invention relates to benzodioxanes that are useful as inhibitors of leukotriene A₄ hydrolase (LTA₄H) and are thus useful for treating a variety of diseases and disorders that are mediated or sustained through the activity of leukotrienes including asthma, allergy and cardiovascular diseases including atherosclerosis, myocardial infarction and stroke. This invention also relates to pharmaceutical compositions comprising these compounds, methods of using these compounds in the treatment of various diseases and disorders, processes for preparing these compounds and intermediates useful in these processes.


Background of the Invention

Leukotrienes (LT) are oxidized lipids that are produced by several cell types including neutrophils, mast cells, eosinophils, basophils, monocytes and macrophages. The first committed step in the intracellular synthesis of LTs involves oxidation of arachidonic acid by 5-lipoxygenase (5-LO) to leukotriene A₄ (LTA₄), a process requiring the 5-lipoxygenase-activating protein (FLAP). Leukotriene A₄ hydrolase (LTA₄H) catalyzes the hydrolysis of LTA₄ to produce leukotriene B₄ (LTB₄). Through the engagement of the LTB₄ receptors (BLT1, BLT2), LTB₄ stimulates an array of pro-inflammatory responses (leukocyte chemotaxis, cytokine release, etc.). The leukotriene pathway has been implicated in diseases in which inflammation is a critical component of the pathology; these include cancer, asthma, atherosclerosis, colitis, glomerulonephritis, and pain (for a review, see M. Peters-Golden and W.R. Henderson, Jr., M.D., N. Engl. J. Med., 2007, 357, 1841-1854).

Brief Summary of the Invention

The present invention provides novel compounds which inhibit leukotriene A₄ hydrolase (LTA₄H) and are thus useful for treating a variety of diseases and disorders that are mediated or sustained through the activity of leukotrienes, including allergic, pulmonary, fibrotic, inflammatory and cardiovascular diseases and cancer. ✓

In one embodiment, the invention relates to a compound of formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein:

5 X is N or CH;

n is an integer from 0 to 3;

R¹ is selected from halo, -OH, -CN, -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, and -(C₃-C₆)cycloalkyl;


10 R² and R³ are each independently selected from -H and -(C₁-C₆)alkyl; wherein R² and R³ may join to form a 3- to 6-membered ring optionally comprising one to three heteroatoms, and further optionally substituted with one to three groups selected from halo, -OH, (=O), -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, -C(O)OH, -C(O)(C₁-C₆)alkyl, and -C(O)NH₂;

A is a group of formula -NR⁴R⁵, wherein

15 R⁴ and R⁵ are each independently selected from -H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl; wherein each of the foregoing -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl of said R⁴ and R⁵ groups is optionally independently substituted by one to three R⁶ groups; wherein two R⁶ groups when attached to the
20 same carbon atom of said -(C₁-C₆)alkyl may join to form a 3- to 6-membered ring optionally comprising one to three heteroatoms, and further optionally substituted

with one to three groups selected from halo, -OH, (=O), -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, -C(O)OH, -C(O)(C₁-C₆)alkyl, and -C(O)NH₂;

or A is a (4- to 11-membered)N-heterocyclic ring of formula B:



5

wherein said ring B may be a non-aromatic 4-8 membered monocyclic radical; a bridged bicyclic radical; a spirocyclic radical; or a 6 to 11-membered fused bicyclic radical which may be non-aromatic or have one aromatic ring provided that the aromatic ring of the bicyclic radical, when present, is not attached to methylene carbon atom 1 of the compound of formula (I);

10

wherein said ring B may additionally comprise one to three additional ring heteroatoms independently selected from N, O and S;

wherein said ring B may be further optionally substituted by one to three groups selected from halo, -OH, (=O), -C(O)OH, -C(O)O-(C₁-C₆)alkyl, and -(C₁-C₆)alkyl; and

15

wherein L is absent or a linker selected from -(C₁-C₆)alkylene;

each R⁶ is independently selected from halo, -OR⁷, -CF₃, -CN, -(C₁-C₆)alkyl, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁷)₂, -N(R⁷)₂, -NHC(O)R⁷, -NHC(O)N(R⁷)₂, -S(O)₂R⁷, -NH-S(O)₂-R⁷, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl; wherein each of said, -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl of said R⁶ group is optionally substituted where possible with one to three groups selected from halo, -OH, -CF₃, -CN, (=O), -(C₁-C₆)alkyl, -C(O)OH, -C(O)O-(C₁-C₆)alkyl, -NH₂, -NH(C₁-C₆)alkyl, -N((C₁-C₆)alkyl)₂, -S(O)₂(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl; and

20

25

each R⁷ is independently selected from -H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₃-C₆)cycloalkyl-OH, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl.

- 5 This invention also relates to pharmaceutical compositions comprising the compounds of formula (I), methods of using these compounds in the treatment of various diseases and disorders, processes for preparing these compounds and intermediates useful in these processes.

10

Detailed Description of the Invention

Definitions:

DCE = dichloroethane

DCM = dichloromethane

DEA = diethylamine

- 15 DIBAL-H = diisobutylaluminum hydride

DIPEA = diisopropylethylamine

DMA = dimethylacetamide

DMAP = 4-dimethylaminopyridine

DME = dimethyl ether

- 20 DMF = dimethylformamide

DMSO = dimethylsulfoxide

Et₂O = ethylether

EtOAc = ethyl acetate

EtOH = ethanol

- 25 IPA = isopropyl alcohol

KHMDS = potassium bis(trimethylsilyl)amide

MeCN = acetonitrile

MeOH = methanol

TBTU = 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate

- 30 TEA = triethylamine

TFA = trifluoroacetic acid 

THF = tetrahydrofuran


TMSCF₃ = (trifluoromethyl)trimethylsilane

It will be understood that the terms “compounds of formula (I)” and “compounds of the invention” have the same meaning unless indicated otherwise.

In its broadest embodiment (“the first embodiment of the invention”), the invention relates to compounds of formula (I) as described above, and pharmaceutically acceptable salts thereof, as described above in the summary of the invention.

In another embodiment (“the second embodiment of the invention”), the invention relates to a compound of formula (I) as described in the first embodiment immediately of the invention, or a pharmaceutically acceptable salt thereof, wherein group A is a group of formula -NR⁴R⁵.

In another embodiment (“the third embodiment of the invention”), the invention relates to a compound of formula (I) as described in the first embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein group A is a (4- to 11-membered)N-heterocyclic ring of formula B:



In another embodiment, the invention relates to a compound of formula (I) as described in the second embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein R⁴ is -H or -(C₁-C₆)alkyl, and R⁵ is -(C₁-C₆)alkyl; wherein each -(C₁-C₆)alkyl of said R⁴ and R⁵ groups, when present, is optionally independently substituted by one to three R⁶ groups.

In another embodiment, the invention relates to a compound of formula (I) as described in the embodiment immediately above, or a pharmaceutically acceptable salt thereof, wherein R⁴ is -H or -(C₁-C₆)alkyl, and R⁵ is -(C₁-C₆)alkyl; wherein said -(C₁-C₆)alkyl of said R⁵

group is substituted by $-(C_3-C_6)$ cycloalkyl, $-(4- \text{ to } 11\text{-membered})$ heterocycloalkyl, $-(C_6-C_{10})$ aryl, or $-(5- \text{ to } 11\text{-membered})$ heteroaryl; wherein each of said, $-(C_3-C_6)$ cycloalkyl, $-(4- \text{ to } 11\text{-membered})$ heterocycloalkyl, $-(C_6-C_{10})$ aryl, and $-(5- \text{ to } 11\text{-membered})$ heteroaryl is optionally substituted with one to three groups independently selected from $-(C_1-C_6)$ alkyl, $-\text{CF}_3$, and $-\text{C}(\text{O})\text{OR}^8$.

In another embodiment, the invention relates to a compound of formula (I) as described in the second embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein R^4 is $-\text{H}$ or $-(C_1-C_6)$ alkyl, and R^5 is $-(C_1-C_6)$ alkyl; wherein said $-(C_1-C_6)$ alkyl of said R^5 group is independently substituted by one to three groups selected from $-(C_1-C_6)$ alkyl, $-\text{O}(\text{C}_1-C_6)$ alkyl, $-\text{C}(\text{O})\text{R}^8$, $-\text{C}(\text{O})\text{OR}^8$, $-\text{S}(\text{O})_2\text{R}^8$, and $-\text{NHC}(\text{O})\text{R}^8$.

In another embodiment, the invention relates to a compound of formula (I) as described in the second embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein R^4 and R^5 are each independently selected from $-\text{H}$ or $-(C_1-C_6)$ alkyl.

In another embodiment, the invention relates to a compound of formula (I) as described in the second embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein R^4 is $-\text{H}$ or $-(C_1-C_6)$ alkyl, and R^5 is $-(C_3-C_6)$ cycloalkyl, $-(4- \text{ to } 11\text{-membered})$ heterocycloalkyl, $-(C_6-C_{10})$ aryl, and $-(5- \text{ to } 11\text{-membered})$ heteroaryl; wherein each of the foregoing $-(C_3-C_6)$ cycloalkyl, $-(4- \text{ to } 11\text{-membered})$ heterocycloalkyl, $-(C_6-C_{10})$ aryl, and $-(5- \text{ to } 11\text{-membered})$ heteroaryl groups of said R^5 is optionally independently substituted by one to three groups selected from $-(C_1-C_6)$ alkyl, $-\text{O}(\text{C}_1-C_6)$ alkyl, $-\text{C}(\text{O})\text{R}^8$, $-\text{C}(\text{O})\text{OR}^8$, $-\text{S}(\text{O})_2\text{R}^8$, and $-\text{NHC}(\text{O})\text{R}^8$.


In another embodiment, the invention relates to a compound of formula (I) as described in the third embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein said ring B is 4-8 membered monocyclic radical.


In another embodiment, the invention relates to a compound of formula (I) as described in the embodiment immediately above, or a pharmaceutically acceptable salt thereof, wherein said 4-8 membered monocyclic radical is selected from the group consisting of azetidine, tetrahydropyrrole, piperidine, hexamethyleneimine, 1,2-diazetidine, pyrazolidine, 

imidazolidine, piperazine, hexahydrodiazepine, isoxazolidine, oxazolidine, tetrahydro-2H-1,3-oxazine, morpholine, and hexahydro-1,4-oxazepine; wherein said monocyclic ring may be further optionally substituted by one to three groups selected from halo, -OH, (=O), -C(O)OH, -C(O)O-(C₁-C₆)alkyl, and -(C₁-C₆)alkyl.

- 5 In another embodiment, the invention relates to a compound of formula (I) as described in the third embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein said ring B is a spirocyclic heterocyclic radical.


In another embodiment, the invention relates to a compound of formula (I) as described in the embodiment immediately above, or a pharmaceutically acceptable salt thereof, wherein
10 said spirocyclic heterocyclic radical is selected from:






In another embodiment, the invention relates to a compound of formula (I) as described in the third embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein said ring B is a bridged bicyclic radical; or a 6 to 11-membered fused bicyclic radical which may be non-aromatic or have one aromatic ring provided that the aromatic ring of the bicyclic radical, when present, is not attached to methylene carbon atom 1 of the compound of formula (I).

In another embodiment, the invention relates to a compound of formula (I) as described in the embodiment immediately above, or a pharmaceutically acceptable salt thereof, wherein said 6 to 11-membered fused bicyclic radical or bridged bicyclic radical is selected from:





In another embodiment, the invention relates to a compound of formula (I) as described in the third embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein L is $-\text{CH}_2-$.

- 5 In another embodiment, the invention relates to a compound of formula (I) as described in the third embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein L is absent. *w*

In another embodiment, the invention relates to a compound of formula (I) as described in the third embodiment of the invention, or a pharmaceutically acceptable salt thereof, wherein said 4-8-membered heterocyclic ring B is a selected from azetidiny, pyrrolidinyl, piperidinyl and azepanyl; wherein each of the foregoing azetidiny, pyrrolidinyl, piperidinyl and azepanyl rings is optionally substituted by one to three groups selected from halo, -OH, (=O), -C(O)OH, C(O)O-(C₁-C₆)alkyl, and -(C₁-C₆)alkyl; and wherein

L is absent or a linker selected from -(C₁-C₆)alkylene; and wherein
 10 R⁶ is elected from halo, -OR⁷, -CF₃, -CN, -(C₁-C₆)alkyl, -C(O)R⁷, -C(O)₂R⁷, -C(O)N(R⁷)₂, -N(R⁷)₂, -NHC(O)R⁷, -NHC(O)N(R⁷)₂, -S(O)₂R⁷, -NH-S(O)₂-R⁷, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl; wherein each of said, -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl of said R⁶ group is optionally substituted where
 15 possible with one to three groups selected from halo, -OH, -CF₃, -CN, (=O), -(C₁-C₆)alkyl, -C(O)OH, -C(O)O-(C₁-C₆)alkyl, -NH₂, -NH(C₁-C₆)alkyl, -N((C₁-C₆)alkyl)₂, -S(O)₂(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl.

20










In another embodiment, the invention relates to a compound of formula (I) as described in any of the embodiments above, or a pharmaceutically acceptable salt thereof, wherein X is N.

In another embodiment, the invention relates to a compound of formula (I) as described in
 25 any of the embodiments above except the embodiment immediately above, or a pharmaceutically acceptable salt thereof, wherein X is CH.










The following are representative compounds of the invention which were made by the general synthetic schemes, the examples, and known methods in the art. 

Table 1. Exemplary compounds of the invention.

Cpd No.	Structure	Name
1		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]pyrrolidine
2		4-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]morpholine
3		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-4,4-dimethylpiperidine
4		8-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-2,8-diazaspiro[4.5]decan-1-one
5		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-4-fluoropiperidine
6		(1s,4s)-7-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-7-azabicyclo[2.2.1]heptane
7		4-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]thiomorpholine 1,1-dioxide
8		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,N-dimethylpiperidine-4-carboxamide
9		(3S)-1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]pyrrolidin-3-ol
10		1-({1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-3-yl}methyl)pyrrolidin-2-one

11		1-{4-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperazin-1-yl}ethanone
12		2-{[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]amino}-1-(pyrrolidin-1-yl)ethanone
13		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-methyl-1-(methylsulfonyl)piperidin-4-amine
14		1-{4-[{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}(methyl)amino]piperidin-1-yl}ethanone
15		3-[4-(pyrrolidin-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
16		7-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-5,6,7,8-tetrahydro[1,2,4]triazolo[4,3-a]pyrazine
17		3-{4-[(1,1-dioxidothiomorpholin-4-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
18		3-[4-(morpholin-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
19		(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-3-carboxylic acid

20		(3S)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-3-carboxylic acid
21		1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2,2,2-trifluoroethanol
22		2-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-1,1,1,3,3,3-hexafluoropropan-2-ol
23		N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-2-methylpropan-2-amine
24		(2R)-N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]butan-2-amine
25		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-N-methylpiperidine-4-carboxamide
26		4-{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}butanoic acid
27		{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}methanol
28		2-{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}propan-2-ol
29		3-{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}propan-1-ol










30		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-4-methyl-1,4-diazepane
31		1-{4-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-1,4-diazepan-1-yl} ethanone
32		4-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-1,4-oxazepane
33		N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-2-methoxy-N-methylethanamine
34		(3R)-1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]pyrrolidin-3-ol
35		8-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-1,3,8-triazaspiro[4.5]decane-2,4-dione
36		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-3-methoxyazetidine
37		{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}(morpholin-4-yl)methanone
38		2-{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}-N,N-dimethylacetamide

39		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-4-(methylsulfonyl)piperidine
40		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]azepane
41		N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]cyclopentanamine
42		N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-N-methyl-2-(pyridin-2-yl)ethanamine
43		1-cyclopropyl-N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]methanamine
44		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-4-phenylpiperidin-4-ol
45		N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-N-ethylethanamine
46		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]azetidine-3-carbonitrile
47		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-3-methoxypyrrolidine
48		N-{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}methanesulfonamide











49		N-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-2-methyl-1-(pyrrolidin-1-yl)propan-2-amine
50		1-({1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}methyl)pyrrolidin-2-one
51		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-N,N-dimethylpiperidine-4-carboxamide
52		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-N-(2-hydroxyethyl)piperidine-4-carboxamide
53		1-{1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidin-4-yl}urea
54		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)phenyl]-N-(pyridin-3-ylmethyl)methanamine
55		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)phenyl]-N-[(1-methyl-1H-imidazol-4-yl)methyl]methanamine
56		2-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,2,3,4-tetrahydroisoquinoline-4-carboxylic acid

57		(1R,3S)-3-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)cyclopentanecarboxylic acid
58		3-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)-4,4-dimethylpentanoic acid
59		1-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)cyclopentanecarboxylic acid
60		N-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)-N-methylglycine
61		1-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)pyrrolidine-3-carboxylic acid
62		trans-4-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)cyclohexanecarboxylic acid
63		cis-4-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)cyclohexanecarboxylic acid
64		1-((3R)-3-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)pyrrolidin-1-yl)ethanone
65		1-((3S)-3-((4-((2S)-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl)amino)pyrrolidin-1-yl)ethanone

66		trans-4-((4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)amino)cyclohexanecarboxamide
67		N-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)-N-methylcyclohexanamine
68		1-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)-2-methylpiperidine
69		(1-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)piperidin-3-yl)methanol
70		2-(1-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)piperidin-4-yl)ethanol
71		N-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)propan-2-amine
72		N-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)-1-methoxypropan-2-amine
73		N-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)propan-1-amine
74		N-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)-N-methylethanamine
75		1-(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]phenyl)-N,N-dimethylmethanamine
76		trans-4-((4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)amino)cyclohexanol








77		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpyrrolidine
78		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-ol
79		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,N',N'-trimethylethane-1,2-diamine
80		2-(cyclohexyl{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)ethanol
81		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,2-dimethylpropan-2-amine
82		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)acetamide
83		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)-N-methylacetamide
84		(1R,2R,4S)-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}bicyclo[2.2.1]heptan-2-amine
85		(4aR,8aS)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}decahydroquinoline









86		(1S,2R)-2-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclohexanecarboxamide
87		[(1S,2R)-2-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclohexyl]methanol
88		(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-ol
89		[(1R,2R)-2-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclohexyl]methanol
90		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methanol
91		(3S)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-ol
92		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}imidazolidin-4-one
93		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,N-dimethylpyrrolidin-3-amine
94		1'-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,4'-bipiperidin-2-one










95		N-(cyclopropylmethyl)-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}cyclohexanamine
96		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-carboxamide
97		(1R,2R)-2-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclohexanol
98		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-methoxypiperidine
99		1-[(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methyl]pyrrolidin-2-one
100		trans-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-methylcyclohexanamine
101		(1S,2R)-2-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclopentanol
102		(1S,2S)-2-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclopentanol
103		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}tetrahydro-2H-pyran-3-amine
104		N-cyclohexyl-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N',N'-dimethylethane-1,2-diamine









105		(1S,2S)-2-[(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)(methyl)amino]cyclohexanol
106		(1R,2S)-2-[(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)(methyl)amino]cyclohexanol
107		4-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-3-methylmorpholine
108		5-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)-1-methylpiperidin-2-one
109		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-ethylcyclopentanamine
110		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,1-dimethylpiperidin-4-amine
111		4-[(4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl)amino)methyl]phenol
112		2-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,2,3,4-tetrahydroisoquinolin-6-ol
113		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidine-3-carboxylic acid
114		1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]piperidine-3-carboxamide

115		(3S)-1-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]-3-fluoropyrrolidine
116		9-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2,9-diazaspiro[5.5]undecan-1-one
117		7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one
118		1-(7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]non-1-yl)ethanone
119		7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonane-1-carboxamide
120		9-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methyl-2,9-diazaspiro[5.5]undecan-1-one
121		8-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methyl-2,8-diazaspiro[4.5]decan-1-one
122		7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1-(methylsulfonyl)-1,7-diazaspiro[4.4]nonane

123		2-(7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]non-1-yl)acetamide
124		(7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]non-1-yl)acetonitrile
125		8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one
126		(3S)-3-[4-(pyrrolidin-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
127		7-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one
128		1-(7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]non-1-yl)-2-methoxyethanone
129		8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-2,8-diazaspiro[4.5]decan-1-one







130		9-[(S)-4-(2,3-Dihydrobenzo[1,4]dioxin-2-yl)-benzyl]-2-methyl-2,9-diazaspiro[5.5]undecan-1-one
131		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,4-diazepan-5-one
132		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,4-diazepan-5-one
133		N-[2-({4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}amino)ethyl]acetamide
134		3-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)propanoic acid
135		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}cyclopentanamine
136		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-3-carboxamide
137		(3S)-3-{4-[(4-methylpiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine

138		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methyl-2-(pyridin-2-yl)ethanamine
139		(3S)-3-[4-(azepan-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
140		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylethanamine
141		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-ethylethanamine
142		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylcyclopentanamine
143		(3S)-3-{4-[(4-methyl-1,4-diazepan-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
144		(3R)-1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}pyrrolidin-3-ol
145		(3S)-3-{4-[(1s,4s)-7-azabicyclo[2.2.1]hept-7-ylmethyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
146		(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanol

147		(3S)-1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}pyrrolidin-3-ol
148		1-(4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,4-diazepan-1-yl)ethanone
149		3-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)propan-1-ol
150		(3S)-3-[4-(1,4-oxazepan-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
151		4-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)butanoic acid
152		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylpiperidine-4-carboxamide
153		1-{4-[(4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}amino)piperidin-1-yl]ethanone
154		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-carboxamide

155		(3S)-3-{4-[(4-fluoropiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
156		(3S)-3-[4-(5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
157		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-1-(pyrrolidin-1-yl)propan-2-amine
158		(3S)-3-{4-[(3-methoxypiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
159		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carbonitrile
160		N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)acetamide
161		(3S)-3-{4-[(1,1-dioxidothiomorpholin-4-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
162		(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone
163		1-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-3-yl)methyl]pyrrolidin-2-one











164		4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperazine-1-carboxamide
165		8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,3,8-triazaspiro[4.5]decane-2,4-dione
166		(3S)-3-{4-[(3-methoxyazetidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
167		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1-(methylsulfonyl)piperidin-4-amine
168		(3S)-3-{4-[(3-methoxypyrrolidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
169		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methyl-1-(methylsulfonyl)piperidin-4-amine
170		(3S)-3-(4-{4-(2-methoxyethoxy)piperidin-1-yl}methyl)phenyl)-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
171		2-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-N,N-dimethylacetamide










172		(3S)-3-(4-{{4-(methylsulfonyl)piperidin-1-yl}methyl}phenyl)-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
173		N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}cyclobutanamine
174		N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1-(methylsulfonyl)piperidin-4-amine
175		1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)urea
176		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methanesulfonamide
177		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-4-carbonitrile

178		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)acetamide
179		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-methylpiperidine-4-carboxamide
180		(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone
181		4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)butanoic acid
182		[(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-yl]acetic acid









183		[(3S)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-yl]acetic acid
184		[(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl]acetic acid
185		1-(4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperazin-1-yl)ethanone
186		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-ol
187		1-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)urea

188		(3S)-3-(4-{{4-(methylsulfonyl)piperazin-1-yl}methyl}phenyl)-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
189		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxylic acid
190		N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanesulfonamide
191		(1S,3R)-3-({4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}amino)cyclopentanecarboxylic acid
192		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-ol
193		1-{4-[(2R)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-ol
194		8-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one

195		8-{4-[(2R)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one
196		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidine
197		1-{4-[(2R)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidine
198		4-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}morpholine
199		4-{4-[(2R)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}morpholine
200		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-4-carboxylic acid
201		1-{4-[(2R)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-4-carboxylic acid
202		4-[4-(7-fluoro-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]morpholine
203		1-[4-(7-fluoro-2,3-dihydro-1,4-benzodioxin-2-yl)benzyl]pyrrolidine
204		(3S)-3-[4-(morpholin-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine

205		(3R)-3-[4-(morpholin-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine
206		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide
207		1-{4-[(3R)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide
208		1-[4-(2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl)benzyl]pyrrolidin-2-one
209		3-[4-(2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl)benzyl]-1,3-oxazolidin-2-one
210		1-[4-(2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl)phenyl]methanamine
211		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-methylpiperidine-4-carboxylic acid
212		(3R,4R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-3-methylpiperidine-4-carboxylic acid
213		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-fluoropiperidine-4-carboxylic acid

214		(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidine-3-carboxylic acid
215		(3S)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidine-3-carboxylic acid
216		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-(1H-tetrazol-5-yl)piperidine
217		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-amine
218		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2-hydroxyacetamide
219		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2-methoxyacetamide

220		N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2-hydroxy-2-methylpropanamide
221		N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxy-2-methylpropanamide
222		N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxyacetamide
223		N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-1-hydroxycyclopropanecarboxamide
224		1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-(1,1-dioxido-1,2-thiazolidin-2-yl)piperidine
225		1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]phenyl}ethyl)pyrrolidine
226		4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]phenyl}ethyl)morpholine
227		1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]phenyl}ethyl)piperidine-4-carboxylic acid

228		1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-4-methylpiperidine-4-carboxylic acid
229		2-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2-methylpropanoic acid
230		2-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-methylpropanoic acid
231		4-[(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methyl]benzoic acid
232		2-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid
233		4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)benzoic acid
234		4-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methyl]benzoic acid
235		4-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)benzoic acid
236		4-{[4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl](ethylamino)methyl}benzoic acid

237		4-[(butyl {4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl} amino)methyl]benzoic acid
238		3-[[{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl} (ethyl)amino]methyl]benzoic acid
239		3-[(4-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl} piperazin-1-yl)methyl]benzoic acid

In one embodiment, the invention relates to any of the compounds depicted in Table 1, and pharmaceutically acceptable salts thereof.

5

In another embodiment, the invention relates to a compound selected from the group consisting of:

4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl} piperidin-4-yl)butanoic acid;

4-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl} piperidin-4-yl)benzoic

10 acid;

(3S)-3-{4-[(1s,4s)-7-azabicyclo[2.2.1]hept-7-ylmethyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;

N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl} piperidin-4-yl)methanesulfonamide;

15 (3S)-3-[4-(azepan-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;

1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpiperidine;

7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonane-1-carboxamide;

7-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,7-diazaspiro[4.4]nonan-

20 2-one;

1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl} piperidine-4-carboxylic acid; ✓

- (1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone;
 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,3,8-triazaspiro[4.5]decane-2,4-dione;
- 5 (3S)-3-{4-[(3-methoxypiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)-N-methylacetamide;
 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-(1,1-dioxido-1,2-thiazolidin-2-yl)piperidine;
- 10 (3R)-1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}pyrrolidin-3-ol;
 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxyacetamide;
 4-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methyl]benzoic acid;
- 15 (1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone;
 (3S)-3-[4-(morpholin-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 8-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one;
- 20 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carbonitrile;
 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-methylpiperidine-4-carboxamide;
 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one;
 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-
- 25 hydroxy-2-methylpropanamide;
 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-1-hydroxycyclopropanecarboxamide;
 N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-ethylcyclopentanamine;
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylpiperidine-4-
- 30 carboxamide; 

- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylcyclopentanamine;
- 1-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-3-yl)methyl]pyrrolidin-2-one;
- 5 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpyrrolidine;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-1-(pyrrolidin-1-yl)propan-2-amine;
- N-cyclohexyl-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N',N'-dimethylethane-1,2-diamine;
- 10 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)acetamide;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methyl-2-(pyridin-2-yl)ethanamine;
- (3S)-3-[4-(pyrrolidin-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- 15 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-3-carboxamide;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide;
- N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)acetamide;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-carboxamide;
- 20 (3S)-3-[4-(1,4-oxazepan-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-carboxamide;
- 4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)benzoic acid;
- 1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)urea;
- 25 7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one;
- 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-2,8-diazaspiro[4.5]decan-1-one;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-ol;
- N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methanesulfonamide; 
- 30

- 3-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)propan-1-ol;
- (3S)-3-{4-[(4-methylpiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- 5 N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-ethylethanamine; N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1-(methylsulfonyl)piperidin-4-amine;
- (3S)-3-{4-[(4-fluoropiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- 10 1-(4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,4-diazepan-1-yl)ethanone;
- [(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-yl]acetic acid;
- (1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanol;
- 4-[(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methyl]benzoic acid;
- 15 (3S)-3-{4-[(4-methyl-1,4-diazepan-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- (3S)-3-{4-[(3-methoxypyrrolidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine; and
- 20 N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,2-dimethylpropan-2-amine; or a pharmaceutically salt thereof of each of the foregoing.

In another embodiment, the invention relates to a compound selected from the group consisting of:

- 25 (3S)-3-[4-(azepan-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylcyclopentanamine;
- N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanesulfonamide;
- 30 (3S)-3-{4-[(3-methoxypiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine; 

- (3S)-3-{4-[(4-methylpiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)acetamide;
- 5 (3S)-3-{4-[(1s,4s)-7-azabicyclo[2.2.1]hept-7-ylmethyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 (3S)-3-[4-(pyrrolidin-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-ethylethanamine;
 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)-N-methylacetamide;
- 10 N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-1-(pyrrolidin-1-yl)propan-2-amine;
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-ol;
 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one;
- 15 (1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone;
 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)acetamide;
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylpiperidine-4-carboxamide;
- 20 7-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one;
 (3S)-3-[4-(1,4-oxazepan-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 3-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)propan-1-ol;
- 25 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-2,8-diazaspiro[4.5]decan-1-one;
 (3S)-3-{4-[(4-methyl-1,4-diazepan-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- 30 4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)benzoic acid;
 (3R)-1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}pyrrolidin-3-ol; 

- 1-(4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,4-diazepan-1-yl)ethanone;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carbonitrile;
- N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxy-2-methylpropanamide;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-3-carboxamide;
- (1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanol;
- 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,3,8-triazaspiro[4.5]decane-2,4-dione;
- 10 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-1-hydroxycyclopropanecarboxamide;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-carboxamide;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1-(methylsulfonyl)piperidin-4-amine;
- 15 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide;
- 4-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)benzoic acid;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpyrrolidine;
- 20 1-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-3-yl)methyl]pyrrolidin-2-one;
- 7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one;
- 1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)urea;
- N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-ethylcyclopentanamine;
- 25 N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methyl-2-(pyridin-2-yl)ethanamine;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpiperidine;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-4-carboxylic acid;
- 4-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methyl]benzoic acid;
- 30 (3S)-3-[4-(morpholin-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine; 

- (3S)-3-{4-[(4-fluoropiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 (1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone;
- 5 8-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one;
 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxyacetamide;
 4-[(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methyl]benzoic acid;
- 10 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methanesulfonamide;
 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-methylpiperidine-4-carboxamide;
 4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)butanoic acid;
 7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonane-1-
- 15 carboxamide;
 N-cyclohexyl-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N',N'-dimethylethane-1,2-diamine;
 [(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-yl]acetic acid;
 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-(1,1-dioxido-1,2-thiazolidin-2-
- 20 yl)piperidine; and
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-carboxamide; or
 a pharmaceutically acceptable salt thereof of each of the foregoing.
- 25 In another embodiment, the invention relates a pharmaceutical composition comprising one or more compounds of formula (I) as defined in any of the embodiments above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient. ✓

All terms as used herein in this specification, unless otherwise stated, shall be understood in their ordinary meaning as known in the art. Other more specific definitions are as follows:

- 5 The term “(C₁-C₆)alkyl” refers to branched and unbranched alkyl groups having from 1 to 6 carbon atoms. Examples of -(C₁-C₆)alkyls include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentane, iso-pentyl, neopentyl, n-hexane, iso-hexanes (e.g., 2-methylpentyl, 3-methylpentyl, 2,3-dimethylbutyl, and 2,2-dimethylbutyl). It will be understood that any chemically feasible carbon atom of the (C₁-C₆)alkyl group
10 can be the point of attachment to another group or moiety.

The term “(C₃-C₆)cycloalkyl” refers to a nonaromatic 3- to 6-membered monocyclic carbocyclic radical. Examples of “(C₃-C₆)cycloalkyls” include cyclopropyl, cyclobutyl, cyclohexyl, cyclopentyl and cyclohexyl.

- 15 As used herein, the term “(C₆-C₁₀)aryl” refers to an aromatic hydrocarbon rings containing from six to ten carbon ring and includes monocyclic rings and bicyclic rings where at least one of the rings is aromatic. Non-limiting examples of C₆₋₁₀ aryls include phenyl, indanyl, indenyl, benzocyclobutanyl, dihydronaphthyl, tetrahydronaphthyl, naphthyl,
20 benzocycloheptanyl and benzocycloheptenyl.

- As used herein, the term “4 to 11-membered heterocycle” includes stable nonaromatic 4-8 membered monocyclic heterocyclic radical or a stable nonaromatic 6 to 11-membered fused bicyclic, bridged bicyclic or spirocyclic heterocyclic radical. The 4 to 11-membered
25 heterocycle consists of carbon atoms and one or more, preferably from one to four heteroatoms chosen from nitrogen, oxygen and sulfur. The heterocycle may be either saturated or partially unsaturated. Non-limiting examples of nonaromatic 4-8 membered monocyclic heterocyclic radicals include tetrahydrofuranyl, azetidiny, pyrrolidinyl, pyranyl, tetrahydropyranyl, dioxanyl, thiomorpholinyl, 1,1-dioxo-1λ⁶-thiomorpholinyl, morpholinyl, piperidinyl, piperazinyl, and azepinyl. Non-limiting examples of
30 nonaromatic 6 to 11-membered fused bicyclic radicals include octahydroindolyl, 

octahydrobenzofuranyl, and octahydrobenzothiophenyl. Non-limiting examples of nonaromatic 6 to 11-membered bridged bicyclic radicals include 2-azabicyclo[2.2.1]heptanyl, 3-azabicyclo[3.1.0]hexanyl, and 3-azabicyclo[3.2.1]octanyl. Non-limiting examples of nonaromatic 6 to 11-membered spirocyclic heterocyclic radicals include 7-aza-spiro[3,3]heptanyl, 7-spiro[3,4]octanyl, and 7-aza-spiro[3,4]octanyl. As used herein, the term "5 to 11-membered heteroaryl" includes aromatic 5 to 6-membered monocyclic heteroaryls and aromatic 7 to 11-membered heteroaryl bicyclic rings where at least one of the rings is aromatic, wherein the heteroaryl ring contains 1-4 heteroatoms such as N, O and S. Non-limiting examples of 5 to 6-membered monocyclic heteroaryl rings include furanyl, oxazolyl, isoxazolyl, oxadiazolyl, pyranyl, thiazolyl, pyrazolyl, pyrrolyl, imidazolyl, tetrazolyl, triazolyl, thienyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, and purinyl. Non-limiting examples of 7 to 11-membered heteroaryl bicyclic rings include benzimidazolyl, 1,3-dihydrobenzoimidazol-2-one, quinolinyl, dihydro-2*H*-quinolinyl, isoquinolinyl, quinazolinyl, indazolyl, thieno[2,3-*d*]pyrimidinyl, indolyl, isoindolyl, indazolyl, benzotriazolyl, benzofuranyl, benzopyranyl, benzodioxolyl, benzoxazolyl, benzothiazolyl, pyrrolo[2,3-*b*]pyridinyl, and imidazo[4,5-*b*]pyridinyl.


It will be understood that when a heterocyclyl or heteroaryl contains a S ring atom, such S ring atom can be present in the ring in its divalent, tetravalent, or hexavalent form, i.e., -S-, -S(O)- or -S(O)₂-.

Each aryl or heteroaryl unless otherwise specified includes its partially or fully hydrogenated derivatives. For example, quinolinyl may include decahydroquinolinyl and tetrahydroquinolinyl, naphthyl may include its hydrogenated derivatives such as tetrahydronaphthyl. Other partially or fully hydrogenated derivatives of the aryl and heteroaryl compounds described herein will be apparent to one of ordinary skill in the art.

The term "heteroatom" as used herein shall be understood to mean atoms other than carbon such as O, N, and S.

The term “halo” or “halogen” refers to fluoro, chloro, bromo or iodo.

The symbol



5 means point of attachment of a group R to a moiety.

In all alkyl groups or carbon chains one or more carbon atoms can be optionally replaced by heteroatoms: O, S or N, it shall be understood that if N is not substituted then it is NH, it shall also be understood that the heteroatoms may replace either terminal carbon atoms
 10 or internal carbon atoms within a branched or unbranched carbon chain. Such groups can be substituted as herein above described by groups such as oxo to result in definitions such as but not limited to: alkoxycarbonyl, acyl, amido and thioxo.

For all compounds disclosed in this application, in the event the nomenclature is in conflict
 15 with the structure, it shall be understood that the compound is defined by the structure.

The invention also relates to pharmaceutical preparations, containing as active substance one or more compounds of the invention, or the pharmaceutically acceptable derivatives thereof, optionally combined with conventional excipients and/or carriers.

20

Compounds of the invention also include their isotopically-labelled forms. An isotopically-labelled form of an active agent of a combination of the present invention is identical to said active agent but for the fact that one or more atoms of said active agent have been replaced by an atom or atoms having an atomic mass or mass number different
 25 from the atomic mass or mass number of said atom which is usually found in nature.

Examples of isotopes which are readily available commercially and which can be incorporated into an active agent of a combination of the present invention in accordance with well established procedures, include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, *e.g.*, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F ,
 30 and ^{36}Cl , respectively. An active agent of a combination of the present invention, a 

prodrug thereof, or a pharmaceutically acceptable salt of either which contains one or more of the above-mentioned isotopes and/or other isotopes of other atoms is contemplated to be within the scope of the present invention.

- 5 The invention includes the use of any compounds of described above containing one or more asymmetric carbon atoms may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. Isomers shall be defined as being enantiomers and diastereomers. All such isomeric forms of these compounds are expressly included in the present invention. Each stereogenic carbon may
10 be in the R or S configuration, or a combination of configurations.

Some of the compounds of the invention can exist in more than one tautomeric form. The invention includes methods using all such tautomers.

- 15 The compounds of the invention are only those which are contemplated to be 'chemically stable' as will be appreciated by those skilled in the art. For example, a compound which would have a 'dangling valency', or a 'carbanion' is not compounds contemplated by the inventive methods disclosed herein.
- 20 The invention includes pharmaceutically acceptable derivatives of compounds of formula (I). A "pharmaceutically acceptable derivative" refers to any pharmaceutically acceptable salt or ester, or any other compound which, upon administration to a patient, is capable of providing (directly or indirectly) a compound useful for the invention, or a pharmacologically active metabolite or pharmacologically active residue thereof. A
25 pharmacologically active metabolite shall be understood to mean any compound of the invention capable of being metabolized enzymatically or chemically. This includes, for example, hydroxylated or oxidized derivative compounds of the invention.

- Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable
30 inorganic and organic acids and bases. Examples of suitable acids include hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycolic, lactic, 

salicylic, succinic, toluene-p-sulfuric, tartaric, acetic, citric, methanesulfonic, formic, benzoic, malonic, naphthalene-2-sulfuric and benzenesulfonic acids. Other acids, such as oxalic acid, while not themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds and their

5 pharmaceutically acceptable acid addition salts. Salts derived from appropriate bases include alkali metal (*e.g.*, sodium), alkaline earth metal (*e.g.*, magnesium), ammonium and N-(C₁-C₄)alkyl)⁴⁺ salts.

In addition, within the scope of the invention is use of prodrugs of compounds of the

10 invention. Prodrugs include those compounds that, upon simple chemical transformation, are modified to produce compounds of the invention. Simple chemical transformations include hydrolysis, oxidation and reduction. Specifically, when a prodrug is administered to a patient, the prodrug may be transformed into a compound disclosed hereinabove, thereby imparting the desired pharmacological effect.

15


GENERAL SYNTHETIC METHODS

The compounds of the invention may be prepared by the general methods, examples presented below, and methods known to those of ordinary skill in the art and reported in the chemical literature. In each of the schemes below, the groups R¹ to R³ and A are as


20 defined above for the compound of formula (I), unless noted otherwise. Optimum reaction conditions and reaction times may vary depending on the particular reactants used. Unless otherwise specified, solvents, temperatures, pressures and other reaction conditions may be readily selected by one of ordinary skill in the art. Specific procedures are provided in the Synthetic Examples section.

25

Scheme 1 below depicts the general synthetic procedure for making the compounds of formula (I) wherein X is CH ("the benzodioxane LTAH₄ inhibitors"). 

Scheme 1: General synthetic scheme for making benzodioxane LTA₄H inhibitors

Scheme 2 below depicts the general synthetic procedure for making the compounds of
 5 formula (I) wherein X is N (“the 8-azabenzodioxane LTA₄H inhibitors”).

Scheme 2: General synthetic scheme for 8-azabenzodioxane LTA₄H inhibitors

Starting materials and reagents used in Schemes 1 and 2 are commercially available or may
 10 be prepared by one of ordinary skill in the art using methods described in the chemical
 literature and in the Synthetic Examples section below.

The examples which follow are illustrative and, as recognized by one skilled in the art,
 particular reagents or conditions could be modified as needed for individual compounds
 15 without undue experimentation.

Synthetic Examples

General Methods: Unless noted otherwise, all reactions are run at room temperature
 (about 25°C), under inert atmosphere (e.g., Argon, N₂), and under anhydrous conditions.

All compounds are characterized by at least one of the following methods: ^1H NMR, HPLC, HPLC-MS, and melting point.

Typically, reaction progress is monitored by thin layer chromatography (TLC) or HPLC-MS. Intermediates and products are purified using at least one of the following methods:

Flash chromatography on silica gel,

Recrystallization,

Chiral HPLC using a 20 x 500 mm Chiralpak AD-H column, or 20 x 500 mm Chiralpak OD-H column, and eluting with an isocratic mixture of isopropanol in heptanes with 0.1% diethylamine (DEA) at 7.5 mL/min,

20 x 250 mm Chiralcel OD-H column, and eluting with an isocratic mixture of isopropanol in heptanes at 7.5 mL/min,

Super Critical Fluid (SCF) Chiral HPLC using a 3.0 x 25.0 cm RegisPack column, eluting with an isocratic mixture of MeOH, isopropylamine (IPA), and super critical carbon dioxide at 125 bar; 80 mL/min, and/or

Reversed phase HPLC using a C18 semi-preparative column eluting with a gradient of MeCN+0.1% TFA /H₂O+0.1% TFA, or MeCN+0.1% formic acid /H₂O+0.1% formic acid.

The reported MS data is for observed $[\text{M}+\text{H}]^+$. For bromine containing compounds, the $[\text{M}+\text{H}]^+$ is either reported for one or both of the bromine isotopes (i.e., ^{79}Br and ^{81}Br).

LC/MS methods used in to characterize and isolate the compounds of the inventions are described in Tables 2a and 2b below.

Table 2a. LC/MS Methods and retention times (RT).

LC/MS Method	Time (min)	Mobile Phase		Flow (mL/min)	Column
		H ₂ O (0.1%FA)	CH ₃ CN (0.1%FA)		
1	0	95	5	2.5	Agilent Zorbax C18 SB 3.5um 4.6x30mm cartridge
	1.7	5	95	2.5	

	2	5	95	2.5	
	2.1	95	5	2.5	
	2.3	95	5	2.5	
2	0	70	30	2.5	Agilent Zorbax C18 SB 3.5um 4.6x30mm cartridge
	1.7	5	95	2.5	
	2	5	95	2.5	
	2.1	70	30	2.5	
	2.3	70	30	2.5	
3	0	99	1	2.5	Agilent Zorbax C18 SB 3.5um 4.6x30mm cartridge
	1.7	50	50	2.5	
	2	5	95	2.5	
	2.1	5	95	2.5	
	2.3	99	1	2.5	
4	0	95	5	1.5	Agilent Zorbax Eclipse XDB-C8 5um 4.6x150mm
	7	5	95	1.5	
	9	5	95	1.5	
	9.3	95	5	1.5	
	10	95	5	1.5	
5	0	99	1	2.5	Agilent Zorbax C18 SB 3.5um 4.6x30mm cartridge
	1.6	80	20	2.5	
	1.7	5	95	2.5	
	2	5	95	2.5	
	2.1	99	1	2.5	
	2.3	99	1	2.5	
6	0	99	1	1.5	Agilent Zorbax Eclipse XDB-C8 5um 4.6x150mm column
	2	80	20	1.5	
	7	5	95	1.5	
	9	5	95	1.5	
	9.3	99	1	1.5	
	10	99	1	1.5	
7	0	88	12	1.5	Agilent SB-C18 1.8um 3x50mm column
	0.25	70	30	1.5	
	0.3	60	40	1.5	
	1.19	5	95	1.5	
	1.75	0	100	1.5	
8	0	60	40	1.5	Agilent Eclipse C8 1.8um 3x50mm column
	1.19	15	85	1.5	
	1.75	0	100	1.5	
9	0	95	5	1.5	Agilent SB-AQ 1.8um 3x50mm column
	0.25	50	50	1.5	
	0.3	70	30	1.5	
	1.3	10	90	1.5	


	1.7	0	100	1.5	
10	0	95	5	1.5	Agilent SB-C18 1.8um 3x50mm column
	3.8	10	90	1.5	
	4.5	0	100	1.5	

Table 2b. LC/MS Methods and retention times (RT).

LC/MS Method	Time (min)	Mobile Phase		Flow (mL/min)	Column
		95% H ₂ O 2+ 5% CH ₃ CN (0.05% Formic Acid)	CH ₃ CN (0.05% Formic Acid)		
11	0	90	10	0.8	BEH 2.1x50mm C18, 1.7um particle diameter
	1.19	5	95	0.8	
	1.7	5	95	0.8	
12	0	90	10	0.8	BEH 2.1x50mm C18, 1.7um particle diameter
	1.19	0	100	0.8	
	1.7	0	100	0.8	
13	0	95	5	0.6	Waters HSS T3 2.1X100mm 18 um column
	4.45	0	100	0.6	
	5	0	100	0.6	
14	0	100	0	0.6	Waters HSS T3 2.1X100mm 18 um column
	1	100	0	0.6	
	4.45	0	100	0.6	
	5	0	100		
15	0	90	10	0.6	BEH 2.1x50mm C18, 1.7um particle diameter
	4.45	0	100	0.6	
	4.58	0	100	0.6	

5

Synthesis of IntermediatesPreparation of (S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzaldehyde (A) 



To a stirred solution of pyrocatechol (23.8 g, 216 mmol) in acetone (300mL) is added cesium carbonate (84.4 g, 259 mmol) and 2-Bromo-1-(4-bromo-phenyl)-ethanone (60 g, 216 mmol) at room temperature. The reaction is stirred at room temperature for 1 hour then
 5 water (200 mL) is added. The precipitate is filtered and triturated with EtOAc (150 mL) to give A-1 as a solid.

To a solution of A-1 (50.0 g, 163 mmol) in anhydrous THF (375 mL) is added acetic anhydride (23.0 mL, 244 mmol), TEA (34.0 mL, 244 mmol), and DMAP (199 mg, 1.63
 10 mmol). The reaction mixture is stirred at 40°C for 45 min, cooled to room temperature and diluted with EtOAc (250 mL). The organic solution is washed with water (2 x 100mL), 0.25N HCl (100mL), saturated sodium bicarbonate solution (100mL), and brine (100mL), and dried over Na₂SO₄. After removal of volatile solvent, the residue is triturated with 5% EtOAc in heptane (1500 mL). The solid is filtered and air dried to give A-2.

To degassed DMF (500 mL) is added A-2 (41.0 g, 117 mmol), (1S,2S)-(+)-N-(4-Toluenesulfonyl)-1,2-diphenylethylenediamine (756 mg, 2.10 mmol) and Pentamethylcyclopentadienylrhodium(III)dichloride (Cp*RhCl₂) dimer (319 mg, 0.520 mmol). The resulting mixture is stirred at 0°C for 20 minutes under argon sparging and treated dropwise with formic acid/triethylamine complex (5:2, 31 mL, 72 mmol). The
 20 reaction mixture is stirred under argon at 0°C for 2 hours, diluted with EtOAc (600mL),

and washed with half-saturated sodium bicarbonate solution, saturated sodium bicarbonate, and brine. The organic layer is dried over Na_2SO_4 and concentrated. The residue is purified through a pad of silica gel (400mL), eluting with EtOAc/heptane (1:1, 3 L) to give A-3 as a solid.

5


To a MeOH solution (125 mL) of A-3 (24.6 g, 69.0 mmol) is added a solution of $\text{LiOH}\cdot\text{H}_2\text{O}$ (5.8 g, 137 mmol) in water (125 mL). The mixture is stirred at 60°C for 30 min, cooled to room temperature and concentrated. The residue is diluted with water and neutralized with 1N aqueous HCl to a pH of 6. The resulting mixture is extracted with
 10 EtOAc (3 x 150 mL). The combined organic extracts are washed by saturated sodium bicarbonate solution, brine, dried over Na_2SO_4 , filtered and concentrated to give A-4 as an oil.

To a 0°C solution of triphenylphosphine (32.7 g, 125 mmol) and diisopropyl
 15 azodicarboxylate (24.7mL, 125 mmol) in THF (anhydrous, 400 mL) is added a solution of A-4 (35 g, 113 mmol) in THF (anhydrous, 200 mL) over 30 min. The resulting solution is warmed to room temperature, stirred for 1 hour, and concentrated. The residue is vigorously stirred in heptane (1.8 L) for 2 hours. The precipitate is filtered, and rinsed with heptane. The filtrate is concentrated and purified by flash column chromatography on silica
 20 gel (0-10% EtOAc in heptane) to give A-5 as a solid.

To an argon-degassed solution of A-5 (30.7 g, 105 mmol) in DMF (anhydrous, 400 mL) is added $\text{Zn}(\text{CN})_2$ (12.4 g, 105 mmol), tris(dibenzylideneacetone)dipalladium(0) ($\text{Pd}_2(\text{dba})_3$) (2.9 g, 3.2 mmol), and 1,1'-bis(diphenylphosphino)ferrocene (dppf) (3.5 g, 6.3 mmol). The
 25 resulting mixture is sparged with argon and stirred at 80°C overnight. The reaction is cooled to room temperature and filtered through a pad of Diatomaceous earth, and rinsed with EtOAc. The filtrate is diluted with water (400mL) and extracted with EtOAc (2 x 400mL). The combined organic extracts are washed with brine and stirred with activated carbon (80 g). After 30 min, the mixture is filtered through a pad of Diatomaceous earth
 30 and concentrated. The residue is triturated with 2% EtOAc in heptane (1L), and filtered to give A-6 as a solid. ✓

A solution of A-6 (11.1 g, 46.7 mmol) in THF (anhydrous, 400 mL) at 0°C is treated dropwise with DIBAL-H (25 wt% in toluene, 77.8 mL, 117 mmol). The reaction is stirred at 0°C for 30 min, warmed to room temperature and stirred for 2 hours. The reaction is cooled down to 0°C and quenched with EtOAc (250 mL) followed by saturated potassium sodium tartrate solution (400mL). The mixture is diluted with EtOAc (300 mL) and water (300 mL) and stirred for 30 min. The organic layer is separated and washed with water, 1N HCl solution, and brine, and dried over Na₂SO₄. After filtering through a pad of Diatomaceous earth, the filtrate is concentrated and purified by flash column chromatography on silica gel (0-30% EtOAc in heptane) to give the title product as a solid.


Preparation of (±)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzaldehyde (B)



To a stirred solution of A-1 (1.2 g, 3.9 mmol) in EtOH (40 mL) is added sodium borohydride (295 mg, 7.80 mmol). The reaction is stirred for 14 h, quenched with 1N HCl (10 mL) and concentrated to remove the EtOH. The solid residue is filtered, washed with water and dried in vacuo to give B-1 as a solid.

The title product is synthesized from B-1 according to the procedure described for the synthesis of A from A-4.

Preparation of (S)-4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzaldehyde (C)



To a solution of 2-chloro-3-hydroxy-pyridine (25.0 g, 193 mmol) and 2,4'-dibromo-
acetophenone (53.6 g, 193 mmol) in acetone (400 mL) is added Cs_2CO_3 (75.4 g, 232
5 mmol), and the suspension is stirred at room temperature for 1 h. The reaction is poured
into 1 L of water with stirring. Filtration of the mixture gives C-1 as a solid.

A solution of C-1 (30.0 g, 91.9 mmol), Cp^*RhCl_2 dimer (0.57 g, 0.92 mmol) and N-
((1R,2R)-2-Amino-1,2-diphenyl-ethyl)-4-methyl-benzenesulfonamide (1.0 g, 2.8 mmol) in
anhydrous DMF (400 mL) is cooled to 0 °C and sparged with argon for 20 minutes before
10 the dropwise addition of formic acid: TEA mixture (5:2 mixture; 28.2 mL). The reaction is
stirred at 0 °C with Argon sparging for 1 hr. The reaction mixture is slowly added to 1.5L
of vigorously stirred water. Filtration gives C-2 as a solid.

A solution of C-2 (10.0 g, 30.4 mmol) in DME (350 mL) is heated to 60°C, KHMDS (61.5
15 mL, 0.5M in toluene) is added slowly and the resulting solution is stirred for 30 minutes.
The reaction is cooled to room temperature, quenched with water, concentrated in vacuo
and extracted with EtOAc. The combined organics are washed with brine, dried over
 Na_2SO_4 , filtered and concentrated. The residue is purified by flash column
chromatography (0-40% EtOAc in heptanes) to give C-3 as a solid.

20


To a degassed solution of C-3 (5.50 g, 18.8 mmol) in anhydrous DMF (100 mL) is added
 $\text{Zn}(\text{CN})_2$ (2.2 g, 18.8 mmol) and dppf (1.0 g, 1.9 mmol) followed by $\text{Pd}_2(\text{dba})_3$ (0.86 g,
0.90 mmol), and the reaction is warmed to 80°C overnight. The reaction is then cooled to
room temperature and stirred for 48h. The mixture is filtered through a bed of

25 Diatomaceous earth, and the filtrate slowly poured into 1 L of vigorously stirred water. $\sqrt{\quad}$


The resulting solid is isolated by filtration and purified by flash chromatography on silica gel (0-40% EtOAc in heptanes) to give C-4 as a solid.

A solution of C-4 (3.5 g, 14.7 mmol) in 125 mL of THF is cooled down to 0°C in a ice bath. 25 mL of 1.5M DIBAL-H (36.7 mmol, 2.5 eq) solution in toluene is added dropwise via addition funnel (over 15 min). The reaction is stirred at 0°C for 30 min and then allowed to warm to room temperature. The reaction mixture is stirred for 2h at room temperature. The reaction is cooled to 0 °C and carefully quenched with EtOAc (200 mL total), followed by 100 mL of water and 400 mL of saturated aqueous Rochelle's salt solution, and the mixture is stirred for 5 minutes. The entire mixture is transferred to a separatory funnel and the layers are separated. The aqueous layer is extracted with 100 mL of EtOA twice, and the extracts are combined and washed with 0.5 N HCl (100mL). Some product is observed in the acid layer. Acid layer is cooled to 0°C, neutralized with saturated NaHCO₃, and extracted with EtOAc twice. The organic layers are combined, washed with brine, and dried over anhydrous Na₂SO₄, and evaporated. The resulting residue is purified by flash chromatography eluting with 0-80 % EtOAc/Heptane to give the title compound as a solid.

Preparation of (±)-4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzaldehyde (D)



Compound D-1 is synthesized from C-1 according to the procedure described for the synthesis of B-1. ~




A solution of piperidine-1,4-dicarboxylic acid monobenzyl ester (1.0 g, 3.80 mmol), 2,3,4,5,6-pentafluoro-phenol (0.77 g, 4.18 mmol) and dicyclohexyl-carbodiimide (0.86 g, 4.18 mmol) in dioxane (12 mL) is stirred at room temperature for 16 h. The mixture is filtered and concentrated in vacuo. The residue is purified by flash chromatography (EtOAc/heptane) to give F-2.

To a solution of F-2 (200 mg, 0.47 mmol) in DME (1.0 mL) is added TMSCF_3 (139 mg, 0.98 mol) and tetramethylammonium fluoride (43 mg, 0.47 mmol) at -50°C . The resulting mixture is allowed to warm to room temperature and stirred for 16h. The mixture is concentrated in vacuo and the residue is purified by reverse HPLC (30-95%, MeCN/Water) to give F-3.

A mixture of F-3 (670 mg, 1.74 mmol) and 10% palladium on carbon (210 mg) in MeOH (17 mL) is stirred under an atmosphere of H_2 at room temperature for 15h. The mixture is filtered through Diatomaceous earth and the filter pad is washed with MeOH. The filtrate is concentrated to provide the title product (F).

20 Preparation of 4-methyl-piperidine-carboxylic acid methyl ester hydrochloride (I-1)



To a stirred solution of 4-methyl-piperidine-1,4-dicarboxylic acid mono-tert-butyl ester (1.00 g, 4.10 mmol) in MeOH (2 mL) is added HCl (5 ml, 4 M in dioxane). After 18h, the mixture is evaporated to dryness, the residue is dissolved in MeOH (3mL), and the stirred solution is treated with Et₂O (45 ml). The resulting solid is filtered and dried to give the title compound.

The following intermediates are also prepared according to the procedure described for the synthesis of I-1:

Intermediate #	Structure
I-2	HCl
I-3	HCl
I-4	HCl
I-5	HCl
I-6	HCl


10

Synthesis of Compounds of Formula I

General Method A through E (protocols for reductive amination).

Example of General Method A: ✓


Preparation of 8-[(S)-4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzyl]-2,8-diaza-spiro[4.5]decan-1-one (Example 125)



TEA (0.12mL, 0.83 mmol) is added to a mixture of C (100 mg, 0.42 mmol) and
 5 2,8-Diaza-spiro[4.5]decan-1-one; hydrochloride (158 mg, 0.83 mmol) in 2 mL of DCM.
 One drop of acetic acid is added, and the mixture is stirred for 10 min,
 sodiumacetoxylborohydride (132mg, 0.83 mmol) is added, and the resulting mixture is
 stirred for 24 h. The solvent is evaporated and the crude mixture is dissolved in 2 ml of
 MeCN/H₂O (1:1). The mixture is purified on a reverse phase C18 semi-preparative HPLC
 10 column eluting with a gradient of 0-95% MeCN/H₂O to give the title product.

Example of General Method B:

Preparation of (±)-4-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazine-1-carboxylic acid tert-butyl ester




15 To a solution of B (100 mg, 0.420 mmol), and piperazine-1-carboxylic acid tert-butyl ester
 (93 mg, 0.50 mmol) in DCE (4 mL) is added acetic acid (50 mg, 0.83mmol). The mixture
 is stirred at room temperature for 10 min, treated with sodium triacetoxylborohydride (141
 mg, 0.67 mmol), and stirred at room temperature for 16 hours. The reaction is diluted with
 20 saturated aqueous sodium bicarbonate (5 mL) and extracted with EtOAc (5 mL x 3). The
 combined organic layers is washed with brine, dried over Na₂SO₄, filtered and
 concentrated. The residue is purified on a reversed phase C18 semi-preparative HPLC
 column eluting with a gradient of 5-85% MeCN+0.1%TFA /H₂O+0.1%TFA). The

combined fractions are concentrated and basified by saturated aqueous sodium bicarbonate (5 mL) and extracted with EtOAc (5 mL x 3). The combined organic phases is washed by brine, dried over Na₂SO₄, filtered, and concentrated to give the title product.

5 Example of General Method C:

Preparation of 4-{1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-ylmethyl}-benzoic acid methyl ester




A solution of A (100 mg, 0.42 mmol), 4-piperidin-4-ylmethyl-benzoic acid methyl ester hydrochloride (146 mg, 0.54 mmol), sodium cyanoborohydride (52 mg, 0.83 mmol), and TEA (0.08 mL, 0.54 mmol) in THF (5 mL) is treated with 2 drops of acetic acid, and stirred at room temperature for 16 h. The mixture is concentrated, and the residue is purified by flash chromatography eluting with a gradient of 0-10% MeOH in DCM to give the title compound.

15

Example of General Method D:

Preparation of 1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid methylamide (Example 25) :




A solution of B (40mg, 0.17mmol) and piperidine-4-carboxylic acid methylamide (47.2mg, 0.332mmol) is treated with acetic acid (0.01mL). After shaking for 1 hour, a solution of sodium triacetoxyborohydride (70.6mg, 0.33mmol) in DMA (0.5mL) is added and the

resulting mixture is shaken overnight. The mixture is concentrated, diluted with DMSO (0.8 mL), filtered and purified on a C18 semi-preparative HPLC column eluting with a gradient of 5-85% MeCN+0.1%TFA /H₂O+0.1%TFA) to provide the title compound.

5 **Example of General Method E:**

Preparation of 4-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzylamino]-methyl}-benzoic acid methyl ester



- A solution of A (310 mg), methyl 4-(aminomethyl)benzoate hydrochloride (338 mg),
 10 sodium cyanoborohydride (162 mg), and DIPEA (0.3 mL) in MeOH (5 mL) is treated with
 2 drops of acetic acid, and the resulting mixture is stirred at room temperature for 16h.
 The mixture is concentrated, and the residue is purified by flash chromatography eluting
 with a gradient of 0-10% MeOH in DCM to give the title compound.

- 15 Table 3 provides a summary of the key reagents used to prepare Examples 1-191 according
 to general methods A, B, C, D, E, or F as depicted in the reaction below.






























































Table 3: Examples synthesized by General Method A, B, C, D, E, or F

Ex #	X	Chirality at *	---A	Synthesis Method	LC/MS Method	[M+H] ⁺	Rt (min)
1	CH	racemic		B	10	296.2	1.20
2	CH	racemic		B	10	312.2	1.20
3	CH	racemic		B	10	338.4	1.20
4	CH	racemic		B	10	379.4	1.10
5	CH	racemic		B	10	328.4	1.11
6	CH	racemic		B	10	322.4	1.13
7	CH	racemic		B	10	360.4	1.40
8	CH	S		B	10	381.3	0.67
9	CH	racemic		B	10	312.4	1.04
10	CH	racemic		B	10	407.4	1.14
11	CH	racemic		B	10	353.40	1.47
12	CH	racemic		B	10	353.40	1.57

13	C H	S		A	10	417.40	1.62
14	C H	S		A	10	381.40	1.57
15	N	racemic		B	10	297.40	0.97
16	C H	racemic		B	10	349.40	2.14
17	N	racemic		B	10	361.20	1.66
18	N	racemic		B	10	313.40	0.89
19	C H	S		C	1	354.52	0.55
20	C H	S		C	1	354.24	0.56
21	C H	S		C	1	408.26	0.71
22	C H	S		C	1	476.23	0.77
23	C H	racemic		D	11	298.2	0.7









24	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	298.2	0.73
25	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	367.3	0.66
26	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	396.3	0.73
27	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	340.2	0.66
28	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	368.3	0.72
29	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	368.5	0.7
30	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	339.2	0.56
31	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	367.2	0.65
32	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	326.2	0.67
33	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	314.2	0.71
34	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	312.2	0.65
35	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	394.2	0.65














36	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	312.4	0.68
37	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	423.3	0.69
38	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	395.3	0.70
39	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	388.2	0.66
40	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	324.3	0.79
41	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	310.2	0.78
42	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	361.2	0.79
43	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	296.2	0.75
44	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	402.3	0.82
45	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	298.2	0.75
46	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	306.8	0.73
47	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	326.2	0.74
48	$\begin{smallmatrix} \text{C} \\ \text{H} \end{smallmatrix}$	racemic		D	11	403.2	0.71













49	C H	racemic		D	11	367.2	0.68
50	C H	racemic		D	11	407.3	0.73
51	C H	racemic		D	11	381.3	0.73
52	C H	racemic		D	11	397.3	0.67
53	C H	racemic		D	11	368.2	0.67
54	C H	racemic		D	11	333.2	0.66
55	C H	racemic		D	11	336.2	0.59
56	C H	S		D	11	402.3	0.75
57	C H	S		D	11	354.1	0.63
58	C H	S		D	11	370.2	0.70
59	C H	S		D	11	354.2	0.66
60	C H	S		D	11	314.3	0.63










61	C H	S		D	11	340.1	0.61
62	C H	S		D	11	367.9	0.61
63	C H	S		D	11	368.2	0.64
64	C H	S		D	11	353.8	0.63
65	C H	S		D	11	353.1	0.63
66	C H	S		D	11	367.1	0.64
67	C H	S		D	11	337.8	0.78
68	C H	S		D	11	323.9	0.73
69	C H	S		D	11	339.8	0.66
70	C H	S		D	11	353.9	0.67
71	C H	S		D	11	284.3	0.68
72	C H	S		D	11	313.6	0.70














73	C H	S		D	11	283.8	0.7
74	C H	S		D	11	283.9	0.67
75	C H	S		D	11	269.8	0.65
76	C H	S		D	11	340.2	0.65
77	C H	S		D	11	310.2	0.70
78	C H	S		D	11	325.9	0.65
79	C H	S		D	11	327.1	0.61
80	C H	S		D	11	367.9	0.76
81	C H	S		D	11	311.7	0.71
82	C H	S		D	11	352.9	0.64
83	C H	S		D	11	366.9	0.67
84	C H	S		D	11	336.2	0.73
85	C H	S		D	11	363.8	0.76














86	C H	S		D	11	366.9	0.63
87	C H	S		D	11	354.4	0.68
88	C H	S		D	11	313.3	0.64
89	C H	S		D	11	353.9	0.70
90	C H	S		D	11	340.8	0.59
91	C H	S		D	11	311.8	0.57
92	C H	S		D	11	311.1	0.75
93	C H	S		D	11	339.1	0.49
94	C H	S		D	11	407.2	0.63














95	C H	S		D	11	377.8	0.78
96	C H	S		D	11	397.3	0.56
97	C H	S		D	11	339.8	0.65
98	C H	S		D	11	340.2	0.63
99	C H	S		D	11	407.4	0.62
100	C H	S		D	11	338.2	0.76
101	C H	S		D	11	325.8	0.62
102	C H	S		D	11	325.7	0.6
103	C H	S		D	11	326.3	0.61
104	C H	S		D	11	395.2	0.8
105	C H	S		D	11	353.9	0.66
106	C H	S		D	11	353.9	0.67
107	C H	S		D	11	325.9	0.61














108	CH	S		D	11	352.9	0.56
109	CH	S		D	11	338.4	0.71
110	CH	S		D	11	353.1	0.44
111	CH	S		D	11	348.1	0.64
112	CH	S		D	11	374.1	0.66
113	CH	Racemic		A	13	354.4	1.1
114	CH	Racemic		A	13	353.4	1.0
115	CH	Racemic		A	13	314.4	1.0
116	CH	S		A	12	393.1	1.3
117	CH	S		A	13	365.4	1.3
118	CH	S		A	13	393.4	1.2
119	CH	S		A	13	393.4	1.1










12 0	C H	S		A	7	407.3	0.7
12 1	C H	S		A	13	393.4	1.1
12 2	C H	S		A	7	429.5	0.7
12 3	C H	S		A	7	408.3	0.6
12 4	C H	S		A	7	390.3	0.7
12 5	N	S		A	11	380.4	0.4
12 6	N	S		A	11	297.2	0.4
12 7	N	S		A	11	366.2	0.4
12 8	C H	S		A	11	423.3	0.6
12 9	N	S		A	11	394.2	0.4

13 0	N	S		A	11	408.3	0.5
13 1	C H	S		A	11	339.2	0.6
13 2	N	S		A	13	340.2	0.9
13 3	N	S		A	11	328.2	0.4
13 4	C H	S		A	13	382.4	0.6
13 5	N	S		D	11	311.1	0.52
13 6	N	S		D	11	354.1	0.42
13 7	N	S		D	11	325.1	0.53
13 8	N	S		D	11	362.1	0.51
13 9	N	S		D	11	325.1	0.52
14 0	N	S		D	11	285.3	0.43
14 1	N	S		D	11	299.1	0.47
14 2	N	S		D	11	325.1	0.53


14 3	N	S		D	11	340.1	0.31
14 4	N	S		D	11	313.1	0.41
14 5	N	S		D	11	323.1	0.49
14 6	N	S		D	11	341.1	0.43
14 7	N	S		D	11	313.1	0.41
14 8	N	S		D	11	368.1	0.41
14 9	N	S		D	11	369.1	0.47
15 0	N	S		D	11	327.1	0.44
15 1	N	S		D	11	397.1	0.51
15 2	N	S		D	11	368.1	0.42
15 3	N	S		D	11	368.1	0.42
15 4	N	S		D	11	398.1	0.40
15 5	N	S		D	11	329.1	0.47

15 6	N	S		D	11	350.1	0.56
15 7	N	S		D	11	368.1	0.40
15 8	N	S		D	11	341.1	0.48
15 9	N	S		D	11	336.1	0.44
16 0	N	S		D	11	368.1	0.41
16 1	N	S		D	11	361.0	0.65
16 2	N	S		D	11	424.1	0.46
16 3	N	S		D	11	408.1	0.47
16 4	N	S		D	11	355.1	0.38
16 5	N	S		D	11	395.1	0.41
16 6	N	S		D	11	313.1	0.45
16 7	N	S		D	11	404.1	0.45
16 8	N	S		D	11	327.1	0.46


16 9	N	S		D	11	418.1	0.47
17 0	N	S		D	11	385.1	0.50
17 1	N	S		D	11	396.1	0.47
17 2	N	S		D	11	389.1	0.43
17 3	N	S		D	11	297.1	0.48
17 4	C H	S		A	15	403.2	0.69
17 5	C H	S		A	15	368.4	0.55
17 6	C H	S		A	15	403.2	0.90
17 7	C H	S		A	15	336.2	0.85
17 8	C H	S		A	15	368.0	0.98
17 9	C H	S		A	15	367.2	1.02
18 0	C H	S		A	15	423.3	1.03
18 1	C H	S		A	15	395.9	1.13
18 2	C H	S		A	15	368.2	1.04

183	C H	S		A	15	368.2	1.05
184	C H	S		A	15	354.2	0.99
185	N	S		A	14	354.4	2.09
186	N	S		A	14	327.1	2.13
187	N	S		A	14	369.2	2.09
188	N	S		A	14	390.4	2.22
189	N	S		A	14	355.1	2.16
190	N	S		A	14	404.2	2.16
191	C H	S		A	14	353.8	0.63

Examples 192 and 193: Preparation of (S)-1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-ol (192), and (R)-1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-ol (193).



192





193

5

A racemic mixture of 192 and 193 is prepared from intermediate B and 4-hydroxypiperidine according to the General Method B, and resolved by SCF Chiral HPLC using 20% MeOH, 1% IPA, and super critical carbon dioxide to give 192 as the first- ✓



eluting peak, and 193 as the second-eluting peak. 192: LC/MS Method 10; $R_t = 0.98$ min.; $[M+H]^+ = 326.4$. 193: LC/MS Method 10; $R_t = 0.98$ min.; $[M+H]^+ = 326.4$.

Examples 194 and 195: Preparation of 8-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-2,8-diaza-spiro[4.5]decan-1-one (194) and (8-[(R)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-2,8-diaza-spiro[4.5]decan-1-one (195)





Compound 4 (racemate) is resolved by SCF Chiral HPLC using 55% methanol, 1% isopropylamine, and super critical carbon dioxide to give 194 as the first-eluting peak, and 195 as the second-eluting peak. 194: LC/MS Method 10; $R_t = 1.10$ min.; $[M+H]^+ = 379.4$. 195: LC/MS Method 10; $R_t = 1.09$ min.; $[M+H]^+ = 379.4$.

Examples 196 and 197: Preparation of 1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-pyrrolidine (196) and 1-[(R)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-pyrrolidine (197).




Compound 1 (racemate) is resolved by HPLC using a Chiralpak AD-H column, and eluting with 7% IPA in heptanes with 0.1% DEA to give 196 as the first-eluting peak, and 197 as the second-eluting peak. 196: LC/MS Method 10; $R_t = 1.21$ min.; $[M+H]^+ = 296.2$. 197: LC/MS Method 10; $R_t = 1.21$ min.; $[M+H]^+ = 296.2$.

Examples 198 and 199: Preparation of 4-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-morpholine (198) and 4-[(R)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-morpholine (199).



- 5 Compound 2 (racemate) is resolved by HPLC using a Chiralpak OD-H column and eluting with 7% IPA in heptanes with 0.1% DEA to give 198 as the first-eluting peak and 199 as the second-eluting peak. 198: LC/MS Method 10; Rt = 1.20 min.; $[M+H]^+ = 312.4$. 199: LC/MS Method 10; Rt = 1.21 min.; $[M+H]^+ = 312.4$.
- 10 **Examples 200 and 201: Preparation of (S)-1-[4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid (200) and (R)-1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid (201).** ✓




1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid ethyl ester is prepared from intermediate B and ethyl isonipecotate according to the procedure described in General Method B, and resolved by HPLC using a Chiralpak OD-H column, and eluting with 12% IPA in heptanes with 0.1% DEA to give (S)-1-[4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid ethyl ester as the first-eluting peak, and (R)-1-[4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid ethyl ester as the second-eluting peak.

(S)-1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid ethyl ester (145 mg, 0.380 mmol) and lithium hydroxide monohydrate (48 mg, 1.1 mmol) are heated in 1:1 mixture of MeOH/water (2 mL) at 75°C for 2h. The reaction mixture is acidified with TFA (300 μ L). The resulting white precipitate is filtered off, washed with water, and dried to give compound 200. LC/MS Method 10; Rt = 1.14 min.; $[M+H]^+ = 382.4$. $\checkmark$

Compound 201 is prepared from (R)-1-[4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidine-4-carboxylic acid ethyl ester according to the procedure described for the synthesis of compound 201. LC/MS Method 10; $R_t = 1.13$ min.; $[M+H]^+ = 382.4$.

5 Example: 202

Preparation of 4-[4-(7-Fluoro-2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-morpholine (202)



A solution of 4-fluoro-2-methoxy-phenol (3.0 g, 21.1 mmol) in acetone (250 mL) is treated with cesium carbonate (8.3 g, 25.3 mmol) followed by 2-bromo-1-(4-bromo-phenyl)-ethanone (5.9 g, 21.1 mmol). The resulting mixture is stirred at room temperature for 2 h. Water (600 mL) is added slowly to the vigorously stirring solution. After stirring for 30 minutes, the precipitate is filtered off and washed with copious water to give 1-(4-bromo-phenyl)-2-(4-fluoro-2-methoxy-phenoxy)-ethanone (G-1).

15

G-1 (3.0 g, 8.85 mmol) is dissolved in DCM (30 mL) and cooled to 0°C . Aluminum chloride (2.9 g, 22.1 mmol) is added in one portion and the reaction is stirred at 0°C for 10 minutes. Ethanethiol (1.6 mL, 22.1 mmol) is added and the reaction is stirred at 0°C for 30 minutes. The reaction mixture is poured onto ice and the resulting slurry is stirred for 30 minutes. The product is extracted with EtOAc (3 x 50 mL). The combined organic extracts are dried over sodium sulfate, concentrated, and purified by flash column.

20


chromatography on silica gel (0 to 50% EtOAc in heptane) to provide 1-(4-bromo-phenyl)-2-(4-fluoro-2-hydroxy-phenoxy)-ethanone (G-2).

To a solution of G-2 (1.25g, 3.85 mmol) in EtOH (25 mL) is added sodium borohydride (291 mg, 7.69 mmol), and the mixture is stirred at room temperature for 2 h. Water (5 mL) is added, and the resulting mixture is stirred at room temperature for 1 h. The reaction mixture is concentrated, and the residue is dissolved in 1N HCl and extracted with EtOAc. The organic layer is washed with brine, dried over sodium sulfate, and concentrated. The residue is purified by flash column chromatography on silica gel (0 to 40% EtOAc in heptane) to provide 2-[2-(4-bromo-phenyl)-2-hydroxy-ethoxy]-5-fluoro-phenol (G-3).

Triphenylphosphine (918 mg, 3.5 mmol) is dissolved in THF (25 mL) and cooled to 0°C. Diisopropyl azodicarboxylate (0.7 mL, 3.5 mmol) is added to the mixture and stirred at 0°C for 20 minutes. The mixture is then treated dropwise over 5 minutes with a solution of G-3 (1.1 g, 3.33 mmol) in THF (10 mL), and the resulting mixture is stirred at 0°C for 30 minutes and at room temperature for 30 minutes. The reaction mixture is concentrated, and the residue is purified by flash column chromatography on silica gel (0 to 40% EtOAc in heptane) to give 2-(4-bromo-phenyl)-7-fluoro-2,3-dihydro-benzo[1,4]dioxine (G-4).

A solution of G-4 (200 mg, 0.65 mmol), potassium (morpholin-4-yl)methyltrifluoroborate (134 mg, 0.65 mmol), palladium(II) acetate (4.3 mg, 0.019 mmol), 2-dicyclohexylphosphino-2',4',6'-tri-isopropyl-1,1'-biphenyl (19 mg, 0.039 mmol), and cesium carbonate (632 mg, 1.9 mmol) in 10:1 THF/water (2 mL) is stirred at 95°C for 18 h under an atmosphere of nitrogen. The mixture is taken up in EtOAc, and the organic layer is washed with water, brine, dried over Na₂SO₄, and concentrated. The residue is purified by preparative C18 reversed phase HPLC (MeCN/water; 0.1% TFA) to give the title compound. LC/MS Method 10; Rt = 1.09 min.; [M+H]⁺ = 354.4.


Example 203: Preparation of 1-[4-(7-Fluoro-2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-pyrrolidine (203) 



The title compound is prepared from G-4 and Potassium 1-trifluoroboratomethylpyrrolidine according to the procedure described for the synthesis of compound 202. 203: LC/MS Method 10; $R_t = 1.07$ min.; $[M+H]^+ = 354.4$.


5

Examples 204 and 205: Preparation of (S)-3-(4-Morpholin-4-ylmethyl-phenyl)-2,3-dihydro-[1,4]dioxino[2,3-b]pyridine (204) and (R)-3-(4-Morpholin-4-ylmethyl-phenyl)-2,3-dihydro-[1,4]dioxino[2,3-b]pyridine (205).




10 Compound 18 (racemate) is resolved by HPLC using a Chiralcel OD-H column eluting with 28% isopropanol in heptane to give compound 204 (LCMS method 15: ES^+ m/z 313.2 $[M+H]^+$, $rt = 0.47$ min) and compound 205 (LCMS method 15: ES^+ m/z 313.2 $[M+H]^+$, $rt = 0.50$ min).

15 **Examples 206 and 207: Preparation of 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide (206) and 1-{4-[(3R)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide (207).** *u*



The racemic form of 1-[4-(2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzyl]-piperidine-4-carboxylic acid amide is prepared from compound 4-(2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzaldehyde and piperidine-4-carboxylic acid amide according to the general method B. Compounds 206 and 207 are resolved from the corresponding racemic compound by chiral HPLC according to the procedure described for Examples 204 and 205:




10

Table 4. Preparation of compounds 206 and 207.

Ex #	Chirality at *	MS Method	[M+H] ⁺	rt (min)
206	S	15	354.2	0.47
207	R	15	354.2	0.45

Example 208: Preparation of 11-[4-(2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzyl]-pyrrolidin-2-one (208) ✓




A solution of D (1.0g, 4.15 mmol) in THF (50mL) is treated with sodium borohydride (188 mg, 5.00 mmol) at 0°C. The resulting mixture is allowed to warm to room temperature and stirred at room temperature for 1h. The reaction mixture is concentrated and the residue
 5 dissolved in EtOAc. The organic solution is washed with water and brine, dried over sodium sulfate, and concentrated. The residue is purified by flash column chromatography (silica gel) with MeOH in DCM (from 2% to 8%) to give [4-(2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-phenyl]-methanol H-1.

A solution of H-1 (400 mg, 1.64 mmol) in THF (10 mL) is treated with triphenylphosphine
 10 dibromide (1.39g, 3.29 mmol) and imidazole (224 mg, 3.29 mmol) at room temperature, and the resulting mixture is stirred at room temperature for 72h. The mixture is diluted with water and extracted with EtOAc (25mL, 3X). The combined organic layers are washed with brine, dried over sodium sulfate, and concentrated. The residue is purified by flash column chromatography (silica gel) with EtOAc in heptane (from 15% to 50%) to
 15 give 3-(4-Bromomethyl-phenyl)-2,3-dihydro-[1,4]dioxino[2,3-b]pyridine H-2.

A solution of pyrrolidinone (18 mg, 0.21 mmol) in anhydrous DMF (2 mL) is treated with sodium hydride (60% dispersion in mineral oil, 7.8 mg, 0.2 mmol), and the mixture is stirred at room temperature for 15 minutes. Intermediate H-2 (50 mg, 0.16 mmol) is added, and the mixture is stirred at 50°C. After 15 minutes, the mixture is quenched with
 20 water and extracted with EtOAc. The organic layer is concentrated, and the residue is purified by reversed phase HPLC eluting with a gradient of 5-85% of MeCN in H₂O


(+0.1%TFA). The desired fractions are concentrated. The residue is dissolved in EtOAc, washed with saturated aqueous NaHCO₃, brine, and dried over Na₂SO₄. The solution is then filtered and concentrated to give the title compound as a solid (LCMS method 10: ES⁺ m/z 311.4 [M+H]⁺, Rt = 1.84 min).

5 **Example 209: 3-[4-(2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl)benzyl]-1,3-oxazolidin-2-one (209)**



Compound 209 is prepared from intermediate H-2 according to the procedure described for the synthesis of 208. (LCMS method 10: ES⁺ m/z 313.4 [M+H]⁺, Rt = 1.72 min)

10 **Example 210: Preparation of 4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzylamine (210)**




A solution of H-1 (340 mg, 1.4 mmol), triphenylphosphine (550 mg, 2.1 mmol) and diphenylphosphophyl azide (0.45 mL, 2.1 mmol) in anhydrous THF (30 mL) is treated with diisopropyl azodicarboxylate (0.41 mL, 2.1 mmol). The reaction is stirred at room temperature for 24 hours, diluted with water (50 mL), and extracted with EtOAc (3 x 50 mL). The combined organic solution is washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue is purified by flash chromatography eluting with a gradient of 10-50% EtOAc in Heptane to give H-3 as an oil.

20 A solution of H-3 (390mg, 78% pure, 1.1 mmol) and triphenylphosphine (446 mg, 1.7 mmol) in THF (20 mL) is treated with water (0.2 mL, 11.3 mmol). The mixture is stirred at ~~~~~

40°C for 24 hours, cooled to room temperature, diluted with water (25 mL), and extracted with EtOAc (3 x 25 mL). The combined organic solution is washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue is purified by reversed phase HPLC eluting with a gradient of 5-85% MeCN in H₂O (+0.1%TFA). The combined fractions is concentrated, basified with saturated aqueous NaHCO₃ (10 mL), and extracted with EtOAc (10 mL x 3). The combined organic phase is washed with brine, dried over Na₂SO₄, and concentrated to give the title compound as a solid (LCMS method 10: ES⁺ m/z 243.4 [M+H]⁺, Rt = 0.57 min).

Example 211: Preparation of 1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-4-methyl-piperidine-4-carboxylic acid (211)



Intermediate A (100 mg, 0.42 mmol), methyl-piperidine-4-carboxylic acid methyl ester hydrochloride (105 mg, 0.54 mmol), and TEA (75 uL, 0.54 mmol) are stirred in dry THF (3 mL) for 10 minutes. Sodium triacetoxyborohydride (176 mg) is added and stirred for 4h. The mixture is diluted with saturated NaHCO₃ and extracted with EtOAc. The organic layer is washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-3% MeOH in DCM to give 1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-4-methyl-piperidine-4-carboxylic acid methyl ester. ✓

A solution of 1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-4-methyl-piperidine-4-carboxylic acid methyl ester in MeOH (2 mL) is treated with a solution of LiOH·H₂O (52 mg, 1.23mmol) in water (2 mL). The mixture is heated to 70°C for 2h, concentrated, and treated with TFA (96 uL, 1.23 mmol). The mixture is diluted with water and extracted with EtOAc/THF. The organic layer is dried over Na₂SO₄, filtered through Diatomaceous earth, and concentrated. The residue is purified by reversed phase HPLC eluting with a gradient of 5-80% MeCN in water (+0.1%TFA) to provide the title compound as the TFA salt (LC/MS method 1: ES⁺ m/z 368.23 [M+H]⁺; Rt = 0.62 min).

10 Examples 212-215: Preparation of Compounds 212-215

Compounds 212-215 are prepared from intermediates I-2, I-3, I-5 and I-6 according to the procedure described for the synthesis of compound 211 and shown in Table 5.






Table 5. Preparation of compounds 212-215.

Ex #	--A	MS Method	[M+H] ⁺	Rt (min)
212		1	368.24	0.62
213		1	372.20	0.61
214		1	341.20	0.61

215		1	341.23	0.58
-----	---	---	--------	------


Example 216: Preparation of, 1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-4-(1H-tetrazol-5-yl)-piperidine (216).



To a solution of 177 (115 mg, 0.34 mmol) in DMF (2 mL) is added NaN₃ (89.0 mg, 1.38 mmol) and NH₄Cl (147 mg, 2.75 mmol). The mixture is heated at 120°C for 18 h. Additional NaN₃ (89.0 mg, 1.38 mmol) is added, and the reaction is stirred at 120°C for an additional 72 h. The reaction is filtered, and the filtrate is purified by reversed phase HPLC eluting with a gradient of 5-80% MeCN in water (+0.1%TFA to provide the title compound (LC/MS method 1: ES⁺ m/z 378.2 [M+H]⁺; Rt = 0.54 min).

10


Example 217: Preparation of 1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-ylamine (217).



A solution of intermediate A (300 mg, 1.25 mmol) and piperidin-4-yl-carbamic acid tert-butyl ester (300 mg, 1.5 mmol, 1.2 equiv.) is stirred in dry THF (3 mL) for 10 minutes. Sodium triacetoxyborohydride (316 mg, 1.49 mmol) is added, and the reaction is stirred for

18h. The reaction is concentrated and partitioned between EtOAc and saturated aqueous NaHCO₃. The organic layer is dried over Na₂SO₄, filtered, and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-5% MeOH in DCM. The residue is dissolved in MeOH (1 mL), treated with HCl (10 mL, 4M in dioxane), and stirred for 18h. The reaction is diluted with Et₂O (40 mL) and filtered to provide the title compound as the HCl salt (LC/MS method 1: ES+ m/z 325.2 [M+H]⁺, Rt = 0.35 min).

Example 218: Preparation of N-{1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-yl}-2-hydroxy-acetamide (218).



A solution of compound 217 (80 mg, 0.22 mmol), TEA (0.09 mL, 0.67 mmol), hydroxyacetic acid (22 mg, 0.29 mmol) and TBTU (93 mg, 0.29 mmol) in DMF (2 mL) is stirred for 2h. The reaction is filtered and purified by reversed phase HPLC eluting with a gradient of 0-80% MeCN in water (+0.1%TFA) to provide the title compound as a TFA salt (LC/MS method 1: ES+ m/z 383.2 [M+H]⁺, Rt = 0.59 min).

Example 219-220: Preparation of N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2-methoxyacetamide (219) and N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)-2-hydroxy-2-methylpropanamide (220).

20 Compounds 219 through 223 are prepared and according to the procedure described for compound 218 and shown in Table 6. The products are purified by reversed phase HPLC or flash chromatography eluting with a gradient of 0-10% MeOH in DCM.





Table 6. Preparation of compounds 219-223.

Ex #	X	---R ⁶	MS Method	[M+H] ⁺	Rt (min)
219	CH		1	397.08	0.61
220	CH		1	411.30	0.55
221	N		1	412.27	0.48
222	N		1	384.22	0.43
223	N		1	410.26	0.47

Example 224: Preparation of 1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-4-(1,1-dioxo-1λ⁶-isothiazolidin-2-yl)-piperidine (224).




5

To a stirred solution of compound 217 (535 mg, 1.65 mmol) in THF (10 mL) is added 3-chloro-propane-1-sulfonyl chloride (0.40 mL, 3.3 mmol) and pyridine (0.27 mL). After 18 h

h, the mixture is diluted with saturated NaHCO_3 and extracted with EtOAc. The organic layer is dried over Na_2SO_4 , filtered, and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-10% MeOH in DCM to provide 3-chloro-propane-1-sulfonic acid {1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-yl}-amide. LC/MS method 1: ES+ m/z 465.2 $[\text{M}]^+$, R_t = 0.68 min).

To a solution of 3-chloro-propane-1-sulfonic acid {1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-yl}-amide (410 mg, 0.88 mmol) in DMF (5 mL) is added NaH (60% dispersion in mineral oil, 71 mg, 1.8 mmol). The reaction is heated to 80 °C for 1h, diluted with EtOAc, washed with water and brine, dried over Na_2SO_4 , filtered, and concentrated. The residue is purified by reversed phase HPLC eluting with a gradient of 0-80% MeCN in water (+0.1%TFA). The desired fractions are lyophilized, partitioned between saturated aqueous NaHCO_3 and EtOAc. The organic layer is dried over Na_2SO_4 , filtered and concentrated to provide the title compound (LC/MS method 1: ES+ m/z 429.4 $[\text{M}+\text{H}]^+$, R_t = 0.63 min).

Example 225: Preparation of 1-{1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-phenyl]-ethyl}-pyrrolidine (225).



A solution of A (1.0 g, 4.16 mmol) in THF (10mL) is treated with 1.4M methylmagnesium bromide solution in toluene at 0°C. The resulting mixture is stirred at 0°C for 1h. The

mixture is then quenched with saturated ammonium chloride solution and extracted with EtOAc. The organic solution is dried over Na₂SO₄, filtered, and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-30% EtOAc in heptane to give 1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-phenyl]-ethanol (J-1).


- 5 A solution of J-1 (500 mg, 1.95 mmol) in THF (10 mL) is treated with triphenylphosphine dibromide (1.65 g, 3.90 mmol) and imidazole (265 mg, 3.90 mmol) at room temperature, and the resulting mixture is stirred at room temperature for 72h. The mixture is diluted with water and extracted with EtOAc (25mL, 3X). The combined organic layers are washed with brine, dried over sodium sulfate, and concentrated. The residue is purified by
10 flash column chromatography (silica gel) with EtOAc in Heptane (from 0% to 30%) to give (S)-2-[4-(1-bromo-ethyl)-phenyl]-2,3-dihydro-benzo[1,4]dioxine J.

- A mixture of intermediate J (560 mg, 90% pure, 1.58 mmol) in pyrrolidine (0.5 mL) is heated at 60°C for 18h. The reaction is diluted with MeOH and purified by reversed phase HPLC eluting with a gradient of 5-80% MeCN in water (+0.1%TFA). The desired
15 fractions are combined, diluted with EtOAc, and washed with saturated aqueous NaHCO₃. The organic layer is dried over Na₂SO₄, filtered, and concentrated. The residue is dissolved in Et₂O (2 mL), treated with HCl (2 mL, 2M in Et₂O), and concentrated to provide the title product as the HCl salt (LC/MS method 1: ES+ m/z 311.2 [M+H]⁺, Rt = 0.63 min).

20

Example 226: 4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]phenyl}ethyl)morpholine (226)

Compound 226 is prepared from intermediate J and morpholine according to the procedure described for the synthesis of compound 225.




25

Ex #	MS Method	[M+H] ⁺	Rt (min)
------	-----------	--------------------	----------

226	1	327.20	0.89
-----	---	--------	------


Example 227: Preparation of 1-{1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-phenyl]-ethyl}-piperidine-4-carboxylic acid (227)



- 5 A mixture of intermediate J (188 mg, 0.59 mmol) and piperidine-4-carboxylic acid ethyl ester (0.5 mL, 3.24 mmol) is heated at 60°C for 18h. The reaction is diluted with MeOH and purified by reversed phase HPLC eluting with a gradient of 5-80% CH₃CN in water (+0.1%TFA). The desired fractions are combined, diluted with EtOAc, and washed with saturated aqueous NaHCO₃. The organic layer is dried over Na₂SO₄, filtered, and
- 10 concentrated. The residue is dissolved in a mixture of MeOH (4 mL) and water (4 mL) containing KOH (110 mg, 2 mmol) and heated at 50°C for 18h. The mixture is concentrated, treated with TFA (0.15 mL, 2 mmol), and extracted with EtOAc. The organic layer is dried over Na₂SO₄, filtered, and concentrated to provide the title compound as the TFA salt (LCMS method 7: ES+ m/z 369.2 [M+H]⁺, Rt = 0.56 min).

15


Example 228: Preparation of 1-[(S)-4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzyl]-4-methyl-piperidine-4-carboxylic acid formate salt (228)



- 20 A mixture of 1-[(S)-4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzyl]-4-methyl-piperidine-4-carboxylic acid methyl ester (prepared according to the General Method A) ~~~~~


(43 mg, 0.10 mmol), LiOH·H₂O (21 mg, 0.5 mmol), MeOH (3 mL), and water (1 mL) is warmed to 50 °C overnight. The reaction is concentrated, neutralized with 1 N aqueous HCl, and purified by reversed phase HPLC eluting with a gradient of 0-70% MeCN in water (+0.1% formic acid) to afford the title compound as the formate salt (LCMS method 15: ES+ m/z 382.8 [M+H]⁺, Rt = 0.54 min).

Example 229: Preparation of 2-{1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-yl}-2-methyl-propionic acid formate salt (229)




2-{1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-yl}-2-methyl-propionic acid ethyl ester (prepared according to General Method A) (226 mg, 0.430 mmol) is treated with HCl (1.5 mL, 4M in dioxane, 6 mmol) and 1 mL of water. The mixture is warmed to 140 °C for 1 hour, concentrated, diluted with water, and neutralized with 2N aqueous Na₂CO₃. The aqueous layer is decanted and the remaining residue is purified by reversed phase HPLC eluting with a gradient of 0-70% MeCN in water (+0.1% formic acid) to afford the title compound as the formate salt (LCMS method 15: ES+ m/z 395.8 [M+H]⁺, Rt = 1.25 min).

Example 230: Preparation of 2-{1-[(S)-4-(2,3-Dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl)-benzyl]-piperidin-4-yl}-2-methyl-propionic acid formate salt (230)



Compound 230 is prepared according to the procedure described for the synthesis of compound 229.

Example 231: Preparation of 4-{1-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-ylmethyl}-benzoic acid (231).




- 5 A mixture of 4-{1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperidin-4-ylmethyl}-benzoic acid methyl ester (prepared according to General Method C) (80 mg, 0.17 mmol), LiOH·H₂O (15 mg, 0.36 mmol), MeOH (3 mL) and water (0.5 mL) is stirred at room temperature for 16h. The reaction mixture is neutralized with acetic acid and concentrated. The residue is triturated with water to give the title compound.

10

Examples 231-235: Preparation of Compounds 231-235





Compounds 231-235 are prepared according to the procedure described for the synthesis of compound 231 as shown in Table 7 below.




15

Table 7. Preparation of compounds 231-235.

Ex #	X	---A	MS Method	[M+H] ⁺	Rt (min)
231	CH		4	444.30	1.42

232	CH		4	402.25	1.28
233	CH		4	430.26	1.21
234	N		4	445.29	0.81
235	N		3	431.25	1.59

Example 236: Preparation of 4-({[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-ethyl-amino}-methyl)-benzoic acid (236).



5

A mixture of 4-{{[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzylamino]-methyl}-benzoic acid methyl ester (prepared according to General Method E) (130 mg, 0.33 mmol), acetaldehyde (0.03 mL, 0.50 mmol), and sodium cyanoborohydride (42 mg, 0.67 mmol) in MeOH (15 mL) is treated with 2 drops of acetic acid. The mixture is stirred at room temperature for 16h, and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-10% MeOH in DCM to give 4-({[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-ethyl-amino}-methyl)-benzoic acid methyl ester .

A mixture of 4-({[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-ethyl-amino}-methyl)-benzoic acid methyl ester (65 mg, 0.16 mmol), LiOH•H₂O (23 mg, 0.55 mmol), MeOH (5 mL) and water (0.5 mL) is stirred at room temperature for 16h. The reaction mixture is neutralized with acetic acid and concentrated. The residue is diluted with water and DCM, phases are separated, the organic layer is dried over Na₂SO₄, filtered and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-10% MeOH in DCM to give the title compound.

Examples 236-238: Preparation of Compounds 236-238

Compounds 236-238 are prepared according to the procedure described for the synthesis of compound 236 and as shown in Table 8 below.





Table 8. Preparation of compounds 236-238.

Ex #	X	---A	MS Method	[M+H] ⁺	Rt (min)
236	CH		3	404.40	1.86
237	CH		3	432.29	2.29
238	CH		3	404.26	1.98

Example 239: Preparation of 3-{4-[(S)-4-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazin-1-ylmethyl}-benzoic acid (239) ✓



Methanol (30mL) is added dropwise to acetyl chloride (1.4 mL) at 0°C. The solution is added to 4-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazine-1-carboxylic acid tert-butyl ester (408 mg, 0.99 mmol) (prepared according to the General Method E).
 5 The resulting mixture is stirred at room temperature for 16 h and concentrated. The residue is suspended in a mixture of heptane and EtOAc, and the precipitate is collected and dried under vacuum to give 1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazine dihydrochloride.

10

A solution of 1-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazine dihydrochloride (80 mg, 0.21 mmol), 4-formyl-benzoic acid methyl ester (41 mg, 0.25 mmol), sodium cyanoborohydride (26 mg, 0.42 mmol), and DIPEA (0.07 mL, 0.42 mmol) in MeOH (5 mL) is treated with 2 drops of acetic acid. The resulting mixture is stirred at
 15 room temperature for 16 h, concentrated, diluted with water, and extracted with ethyl acetate. The organic layer is washed with brine, dried over Na₂SO₄, filtered, and concentrated. The residue is purified by flash chromatography eluting with a gradient of 0-10% MeOH in DCM to give 4-{4-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazin-1-ylmethyl}-benzoic acid methyl ester.

20

A mixture of 4-{4-[(S)-4-(2,3-dihydro-benzo[1,4]dioxin-2-yl)-benzyl]-piperazin-1-ylmethyl}-benzoic acid methyl ester (48 mg, 0.11 mmol), LiOH·H₂O (15 mg, 0.37 mmol), and

dioxane (5 mL), and water (0.5 mL) is stirred at room temperature for 16h. The reaction mixture is neutralized with acetic acid and concentrated. The residue is triturated with water to give the title compound (LCMS method 4: ES+ m/z 445.2 $[M+H]^+$, Rt = 1.31 min).

5

Assessment of Biological Properties

The compounds of the invention are assessed for the ability to interact with human LTA₄ hydrolase in an enzymatic assay that measures the ability of the enzyme to cleave the peptide bond of arginyl-aminomethylcoumarin (Arg-AMC). LTA₄H Enzyme (1nM final),
 10 Arg-AMC substrate (50 μ M final), and compound are combined in a reaction buffer (50 mM Tris-HCl (pH 7.5), 100 mM KCl, 0.5% bovine serum albumin) at room temperature for 1h. The formation of product is assessed by measuring the fluorescence of aminomethylcoumarin product (excitation wavelength 380nm/emission wavelength 460nm). In general, the preferred potency range (IC₅₀) of compounds in the LTA₄H
 15 Enzyme assay is between 0.1 nM to 10 μ M, the more preferred potency range is 0.1 nM to 0.1 μ M, and the most preferred potency range is 0.1 nM to 10 nM.

Table 9. IC₅₀ values of LTA₄H Enzyme assay.

Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)
1	0.38	61	0.60	121	0.37	181	0.042
2	2.45	62	1.79	122	0.91	182	0.29
3	2.57	63	7.90	123	0.73	183	0.48
4	0.74	64	0.83	124	2.45	184	0.11
5	2.96	65	1.15	125	0.16	185	0.59
6	0.46	66	1.79	126	0.18	186	0.24
7	2.79	67	0.61	127	0.12	187	0.07
8	0.32	68	0.10	128	0.65	188	0.87
9	1.49	69	0.60	129	0.23	189	0.16
10	0.75	70	0.57	130	0.51	190	0.09
11	2.95	71	1.88	131	1.73	191	1.62
12	10.19	72	1.80	132	0.91	192	0.43
13	0.36	73	3.65	133	1.75	193	5.35

Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)
14	0.27	74	1.00	134	0.47	194	0.15
15	0.36	75	4.51	135	0.47	195	1.59
16	2.32	76	1.90	136	0.19	196	0.39
17	0.77	77	0.18	137	0.26	197	2.69
18	1.14	78	1.40	138	0.18	198	2.28
19	0.73	79	0.51	139	0.10	199	40.12
20	1.30	80	0.71	140	0.38	200	0.12
21	4.43	81	0.31	141	0.26	201	1.59
22	200.00	82	0.20	142	0.17	202	23.37
23	5.20	83	0.13	143	0.30	203	2.94
24	5.90	84	2.69	144	0.14	204	0.15
25	0.76	85	0.45	145	0.09	205	27.50
26	0.43	86	0.92	146	0.29	206	0.19
27	1.20	87	0.69	147	0.35	207	0.86
28	3.40	88	0.54	148	0.28	208	21.45
29	2.04	89	1.40	149	0.24	209	12.41
30	1.77	90	0.77	150	0.21	210	19.00
31	1.54	91	0.54	151	0.10	211	0.69
32	1.80	92	35.99	152	0.17	212	0.49
33	3.19	93	1.98	153	0.82	213	0.81
34	1.89	94	0.45	154	0.20	214	0.47
35	0.26	95	0.49	155	0.28	215	0.70
36	4.45	96	0.22	156	0.91	216	0.13
37	1.05	97	2.87	157	0.18	217	2.28
38	1.14	98	0.61	158	0.13	218	0.37
39	2.14	99	0.37	159	0.16	219	0.49
40	0.82	100	2.36	160	0.18	220	0.47
41	3.71	101	1.90	161	0.41	221	0.16
42	0.69	102	2.68	162	0.14	222	0.14
43	4.42	103	2.40	163	0.17	223	0.16
44	0.69	104	0.18	164	0.84	224	0.13
45	0.90	105	0.51	165	0.13	225	5.30
46	24.82	106	0.46	166	0.68	226	42.95
47	1.73	107	1.35	167	0.27	227	1.40

Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)
48	0.16	108	0.87	168	0.31	228	0.61
49	0.32	109	0.17	169	0.33	229	3.85
50	0.60	110	2.15	170	0.47	230	1.24
51	0.82	111	2.25	171	0.45	231	0.29
52	0.75	112	1.07	172	0.53	232	2.75
53	0.42	113	2.49	173	0.73	233	0.22
54	5.93	114	0.77	174	0.60	234	0.14
55	3.63	115	3.03	175	0.22	235	0.08
56	6.08	116	0.82	176	0.24	236	6.04
57	13.66	117	0.23	177	1.45	237	0.81
58	1.36	118	0.45	178	0.35	238	0.55
59	89.24	119	0.10	179	0.16	239	0.15
60	31.02	120	0.51	180	0.12		

The compounds of the invention are additionally tested in a human whole blood (HWB) assay to determine their ability to inhibit the synthesis of LTB₄ in a cellular system.

Compounds are combined with heparinized human whole blood and incubated for 15

- 5 minutes at 37°C. Calcimycin (20μM final, prepared in phosphate-buffered saline, pH 7.4) is then added and the mixture is incubated for another 30 minutes at 37°C. The samples are centrifuged for 5 min at low speed (1500 x g) and the plasma layer is removed. Plasma LTB₄ concentrations are then measured using an antibody-based homogenous time-resolved fluorescence method (CisBio, Bedford, MA). In general, the preferred potency range (IC₅₀) of compounds in the HWB assay is between 10 nM to 10 μM, the more preferred potency range is 10 nM to 1 μM, and the most preferred potency range is 10 nM to 100 nM. The potencies of representative compounds of the invention in the WHB assays are shown in Table 10.

- 15 Table 10. IC₅₀ values of LTB₄ production inhibition assay in human whole blood.

Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)	Example	IC ₅₀ (nM)
139	13.18	175	72.56	52	145.12	61	250.398

142	24.37	173	72.75	219	148.90	118	255.61
190	26.03	109	75.52	183	151.13	174	268.33
158	27.50	138	76.03	14	154.92	87	268.33
137	27.64	68	80.65	53	154.92	9	284.85
160	27.98	81	80.74	64	157.95	229	289.83
145	28.39	200	82.95	156	158.75	25	294.12
126	32.30	220	91.13	121	159.37	85	304.96
141	32.81	185	92.28	178	159.61	227	307.44
140	33.44	237	94.53	69	164.89	26	308.47
83	33.49	134	94.60	214	168.70	116	310.32
157	35.00	234	94.77	18	176.14	123	312.41
186	36.46	204	94.98	187	176.77	37	323.66
168	38.99	155	95.39	1	179.33	67	345.74
125	39.18	99	95.39	98	180.00	10	352.40

Methods of Use

The compounds of the invention are effective inhibitors of leukotriene A₄ hydrolase (LTA₄H) and thus inhibit leukotriene production. Therefore, in one embodiment of the invention, there is provided methods of treating leukotriene-mediated disorders using compounds of the invention. In another embodiment, there is provided methods of treating cardiovascular, inflammatory, allergic, pulmonary and fibrotic diseases, renal diseases and cancer using compounds of the invention.

Without wishing to be bound by theory, by inhibiting the activity of LTA₄H, the compounds of the invention block the production of LTB₄ resulting from the oxidation of arachidonic acid by 5-LO and subsequent metabolism. Thus, the inhibition of LTA₄H activity is an attractive means for preventing and treating a variety of diseases mediated by LTB₄. These include:

- Cardiovascular diseases including atherosclerosis, myocardial infarction, stroke, aortic aneurysm, sickle cell crisis, ischemia-reperfusion injury, pulmonary arterial hypertension and sepsis;
- Allergic diseases including asthma, allergic rhinitis, rhinosinusitis, atopic dermatitis and urticaria;

Fibrotic diseases including airway remodeling in asthma, idiopathic pulmonary fibrosis, scleroderma, asbestosis;

Pulmonary syndromes including adult respiratory distress syndrome, viral bronchiolitis, obstructive sleep apnea, chronic obstructive pulmonary disease, cystic fibrosis, and bronchopulmonary dysplasia;

Inflammatory diseases including rheumatoid arthritis, osteoarthritis, gout, glomerulonephritis, interstitial cystitis, psoriasis, inflammatory bowel disease systemic lupus erythematosus, transplant rejection, inflammatory and allergic ocular diseases;

Cancer including solid tumors, leukemias and lymphomas; and Renal diseases such as glomerulonephritis.

For treatment of the above-described diseases and conditions, a therapeutically effective dose will generally be in the range from about 0.01 mg to about 100 mg/kg of body weight per dosage of a compound of the invention; preferably, from about 0.1 mg to about 20 mg/kg of body weight per dosage. For example, for administration to a 70 kg person, the dosage range would be from about 0.7 mg to about 7000 mg per dosage of a compound of the invention, preferably from about 7.0 mg to about 1400 mg per dosage. Some degree of routine dose optimization may be required to determine an optimal dosing level and pattern. The active ingredient may be administered from 1 to 6 times a day.

General Administration and Pharmaceutical Compositions

When used as pharmaceuticals, the compounds of the invention are typically administered in the form of a pharmaceutical composition. Such compositions can be prepared using procedures well known in the pharmaceutical art and comprise at least one compound of the invention. The compounds of the invention may also be administered alone or in combination with adjuvants that enhance stability of the compounds of the invention, facilitate administration of pharmaceutical compositions containing them in certain embodiments, provide increased dissolution or dispersion, increased antagonist activity, provide adjunct therapy, and the like. The compounds according to the invention may be used on their own or in conjunction with other active substances according to the

invention, optionally also in conjunction with other pharmacologically active substances. In general, the compounds of this invention are administered in a therapeutically or pharmaceutically effective amount, but may be administered in lower amounts for diagnostic or other purposes.

5


Administration of the compounds of the invention, in pure form or in an appropriate pharmaceutical composition, can be carried out using any of the accepted modes of administration of pharmaceutical compositions. Thus, administration can be, for example, orally, buccally (e.g., sublingually), nasally, parenterally, topically, transdermally, 10 vaginally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as, for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages. The pharmaceutical compositions will generally include a conventional pharmaceutical carrier or excipient and 15 a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, vehicles, or combinations thereof. Such pharmaceutically acceptable excipients, carriers, or additives as well as methods of making pharmaceutical compositions for various modes or administration are well-known to those of skill in the art. The state of the art is evidenced, 20 e.g., by *Remington: The Science and Practice of Pharmacy*, 20th Edition, A. Gennaro (ed.), Lippincott Williams & Wilkins, 2000; *Handbook of Pharmaceutical Additives*, Michael & Irene Ash (eds.), Gower, 1995; *Handbook of Pharmaceutical Excipients*, A.H. Kibbe (ed.), American Pharmaceutical Ass'n, 2000; H.C. Ansel and N.G. Popovich, *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 5th ed., Lea and Febiger, 25 1990; each of which is incorporated herein by reference in their entireties to better describe the state of the art.

As one of skill in the art would expect, the forms of the compounds of the invention utilized in a particular pharmaceutical formulation will be selected (e.g., salts) that possess 30 suitable physical characteristics (e.g., water solubility) that are required for the formulation to be efficacious. ~~~~~

Claims

What is claimed is:

1. A compound of formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein:

X is N or CH;

n is an integer from 0 to 3;

- 10 R^1 is selected from halo, -OH, -CN, -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, and -(C₃-C₆)cycloalkyl;

- R^2 and R^3 are each independently selected from -H and -(C₁-C₆)alkyl; wherein R^2 and R^3 may join to form a 3- to 6-membered ring optionally comprising one to three heteroatoms, and further optionally substituted with one to three groups selected from halo, -OH, (=O), -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, -C(O)OH, -C(O)(C₁-C₆)alkyl, and -C(O)NH₂;
- 15


A is a group of formula -NR⁴R⁵, wherein

- R^4 and R^5 are each independently selected from -H, -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl; wherein each of the foregoing -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl of said R^4 and R^5 groups is optionally independently substituted by one to three R^6 groups; wherein two R^6 groups when attached to the
- 20

same carbon atom of said $-(C_1-C_6)\text{alkyl}$ may join to form a 3- to 6-membered ring optionally comprising one to three heteroatoms, and further optionally substituted with one to three groups selected from halo, $-OH$, $(=O)$, $-(C_1-C_6)\text{alkyl}$, $-O(C_1-C_6)\text{alkyl}$, $-C(O)OH$, $-C(O)(C_1-C_6)\text{alkyl}$, and $-C(O)NH_2$;

5

or A is a (4- to 11-membered)N-heterocyclic ring of formula B:



10

wherein said ring B may be a non-aromatic 4-8 membered monocyclic radical; a bridged bicyclic radical; a spirocyclic radical; or a 6 to 11-membered fused bicyclic radical which may be non-aromatic or have one aromatic ring provided that the aromatic ring of the bicyclic radical, when present, is not attached to methylene carbon atom 1 of the compound of formula (I);

wherein said ring B may additionally comprise one to three additional ring heteroatoms independently selected from N, O and S;

15

wherein said ring B may be further optionally substituted by one to three groups selected from halo, $-OH$, $(=O)$, $-C(O)OH$, $-C(O)O-(C_1-C_6)\text{alkyl}$, and $-(C_1-C_6)\text{alkyl}$; and

wherein L is absent or a linker selected from $-(C_1-C_6)\text{alkylene}$;

each R^6 is independently selected from halo, $-OR^7$, $-CF_3$, $-CN$, $-(C_1-C_6)\text{alkyl}$, $-C(O)R^7$, $-C(O)_2R^7$, $-C(O)N(R^7)_2$, $-N(R^7)_2$, $-NHC(O)R^7$, $-NHC(O)N(R^7)_2$, $-S(O)_2R^7$, $-NH-S(O)_2-R^7$, $-(C_3-C_6)\text{cycloalkyl}$, $-(4\text{- to }11\text{-membered})\text{heterocycloalkyl}$, $-(C_6-C_{10})\text{aryl}$, and $-(5\text{- to }11\text{-membered})\text{heteroaryl}$; wherein each of said, $-(C_1-C_6)\text{alkyl}$, $-O(C_1-C_6)\text{alkyl}$, $-(C_3-C_6)\text{cycloalkyl}$, $-(4\text{- to }11\text{-membered})\text{heterocycloalkyl}$, $-(C_6-C_{10})\text{aryl}$, and $-(5\text{- to }11\text{-membered})\text{heteroaryl}$ of said R^6 group is optionally substituted where possible with one to three groups selected from halo, $-OH$, $-CF_3$, $-CN$, $(=O)$, $-(C_1-C_6)\text{alkyl}$, $-C(O)OH$, $-C(O)O-(C_1-C_6)\text{alkyl}$, $-NH_2$, $-NH(C_1-C_6)\text{alkyl}$, $-N((C_1-C_6)\text{alkyl})_2$, $-S(O)_2(C_1-C_6)\text{alkyl}$, $-(C_3-$

C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl; and

each R⁷ is independently selected from -H, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl, -(C₁-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -(C₃-C₆)cycloalkyl-OH, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein group A is a group of formula -NR⁴R⁵.

10 3. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R⁴ is -H or -(C₁-C₆)alkyl, and R⁵ is -(C₁-C₆)alkyl; wherein each -(C₁-C₆)alkyl of said R⁴ and R⁵ groups, when present, is optionally independently substituted by one to three R⁶ groups.


15 4. The compound of claim 1 or 3, or a pharmaceutically acceptable salt thereof, wherein R⁴ is -H or -(C₁-C₆)alkyl, and R⁵ is -(C₁-C₆)alkyl; wherein said -(C₁-C₆)alkyl of said R⁵ group is substituted by -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, or -(5- to 11-membered)heteroaryl; wherein each of said, -(C₃-C₆)cycloalkyl, -(4- to 11-membered)heterocycloalkyl, -(C₆-C₁₀)aryl, and -(5- to 11-membered)heteroaryl
20 is optionally substituted with one to three groups independently selected from -(C₁-C₆)alkyl, -CF₃, and -C(O)OR⁸.

5. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R⁴ is -H or -(C₁-C₆)alkyl, and R⁵ is -(C₁-C₆)alkyl; wherein said -(C₁-C₆)alkyl of
25 said R⁵ group is independently substituted by one to three groups selected from -(C₁-C₆)alkyl, -O(C₁-C₆)alkyl, -C(O)R⁸, -C(O)OR⁸, -S(O)₂R⁸, and -NHC(O)R⁸.

6. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R⁴ and R⁵ are each independently selected from -H or -(C₁-C₆)alkyl. 

7. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^4 is $-H$ or $-(C_1-C_6)alkyl$, and R^5 is $-(C_3-C_6)cycloalkyl$, $-(4- \text{ to } 11- \text{ membered})heterocycloalkyl$, $-(C_6-C_{10})aryl$, and $-(5- \text{ to } 11- \text{ membered})heteroaryl$; wherein each of the foregoing $-(C_3-C_6)cycloalkyl$, $-(4- \text{ to } 11- \text{ membered})heterocycloalkyl$, $-(C_6-C_{10})aryl$, and $-(5- \text{ to } 11- \text{ membered})heteroaryl$ groups of said R^5 is optionally independently substituted by one to three groups selected from $-(C_1-C_6)alkyl$, $-O(C_1-C_6)alkyl$, $-C(O)R^8$, $-C(O)OR^8$, $-S(O)_2R^8$, and $-NHC(O)R^8$.

8. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein group A is a $(4- \text{ to } 11- \text{ membered})N$ -heterocyclic ring of formula B:



9. The compound of claim 1 or 8, or a pharmaceutically acceptable salt thereof, wherein said ring B is 4-8 membered monocyclic radical.

10. The compound of claim 9, or a pharmaceutically acceptable salt thereof, wherein said 4-8 membered monocyclic radical is selected from the group consisting of azetidine, tetrahydropyrrole, piperidine, hexamethyleneimine, 1,2-diazetidine, pyrazolidine, imidazolidine, piperazine, hexahydrodiazepine, isoxazolidine, oxazolidine, tetrahydro-2H-1,3-oxazine, morpholine, and hexahydro-1,4-oxazepine; wherein said monocyclic ring may be further optionally substituted by one to three groups selected from halo, $-OH$, $(=O)$, $-C(O)OH$, $-C(O)O-(C_1-C_6)alkyl$, and $-(C_1-C_6)alkyl$.

11. The compound of claim 1 or 8, or a pharmaceutically acceptable salt thereof, wherein said ring B is a spirocyclic heterocyclic radical.

12. The compound of claim 1 or 8, or a pharmaceutically acceptable salt thereof, wherein said ring B is a bridged bicyclic radical; or a 6 to 11-membered fused bicyclic radical which may be non-aromatic or have one aromatic ring provided that the aromatic

ring of the bicyclic radical, when present, is not attached to methylene carbon atom 1 of the compound of formula (I).

13. The compound of claim 1 or 8, or a pharmaceutically acceptable salt thereof,
5 wherein L is $-\text{CH}_2-$.

14. The compound of claim 1 or 8, or a pharmaceutically acceptable salt thereof,
wherein L is absent.

10 15. The compound of claim 1 or 8, or a pharmaceutically acceptable salt thereof,
wherein said 4-8-membered heterocyclic ring B is a selected from azetidiny, pyrrolidiny,
piperidiny and azepany; wherein each of the foregoing azetidiny, pyrrolidiny,
piperidiny and azepany rings is optionally substituted by one to three groups selected
from halo, $-\text{OH}$, $(=\text{O})$, $-\text{C}(\text{O})\text{OH}$, $\text{C}(\text{O})\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl}$, and $-(\text{C}_1-\text{C}_6)\text{alkyl}$; and
15 wherein

L is absent or a linker selected from $-(\text{C}_1-\text{C}_6)\text{alkylene}$; and wherein
 R^6 is elected from halo, $-\text{OR}^7$, $-\text{CF}_3$, $-\text{CN}$, $-(\text{C}_1-\text{C}_6)\text{alkyl}$, $-\text{C}(\text{O})\text{R}^7$, $-\text{C}(\text{O})_2\text{R}^7$,
 $-\text{C}(\text{O})\text{N}(\text{R}^7)_2$, $-\text{N}(\text{R}^7)_2$, $-\text{NHC}(\text{O})\text{R}^7$, $-\text{NHC}(\text{O})\text{N}(\text{R}^7)_2$, $-\text{S}(\text{O})_2\text{R}^7$, $-\text{NH}-\text{S}(\text{O})_2-\text{R}^7$,
 $-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$, $-(4\text{- to }11\text{-membered})\text{heterocycloalkyl}$, $-(\text{C}_6-\text{C}_{10})\text{aryl}$, and $-(5\text{-}$
20 $\text{to }11\text{-membered})\text{heteroaryl}$; wherein each of said, $-(\text{C}_1-\text{C}_6)\text{alkyl}$, $-\text{O}(\text{C}_1-\text{C}_6)\text{alkyl}$,
 $-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$, $-(4\text{- to }11\text{-membered})\text{heterocycloalkyl}$, $-(\text{C}_6-\text{C}_{10})\text{aryl}$, and $-(5\text{-}$
 $\text{to }11\text{-membered})\text{heteroaryl}$ of said R^6 group is optionally substituted where
possible with one to three groups selected from halo, $-\text{OH}$, $-\text{CF}_3$, $-\text{CN}$, $(=\text{O})$, $-(\text{C}_1-\text{C}_6)\text{alkyl}$,
 $-\text{C}(\text{O})\text{OH}$, $-\text{C}(\text{O})\text{O}-(\text{C}_1-\text{C}_6)\text{alkyl}$, $-\text{NH}_2$, $-\text{NH}(\text{C}_1-\text{C}_6)\text{alkyl}$, $-\text{N}((\text{C}_1-\text{C}_6)\text{alkyl})_2$,
25 $-\text{S}(\text{O})_2(\text{C}_1-\text{C}_6)\text{alkyl}$, $-(\text{C}_3-\text{C}_6)\text{cycloalkyl}$, $-(4\text{- to }11\text{-membered})\text{heterocycloalkyl}$, $-(\text{C}_6-\text{C}_{10})\text{aryl}$, and $-(5\text{- to }11\text{-membered})\text{heteroaryl}$.

16. The compound of any of the preceeding claims, or a pharmaceutically acceptable
salt thereof, wherein X is N. 

17. The compound of any of claims 1 to 15, or a pharmaceutically acceptable salt thereof, wherein X is CH.

18. A compound selected from the group consisting of:

4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)butanoic acid;

5 4-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)benzoic acid;

(3S)-3-{4-[(1s,4s)-7-azabicyclo[2.2.1]hept-7-ylmethyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;

10 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanesulfonamide;

(3S)-3-[4-(azepan-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;

1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpiperidine;

7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonane-1-carboxamide;

15 7-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one;

1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-4-carboxylic acid;

(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone;

20 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,3,8-triazaspiro[4.5]decane-2,4-dione;

(3S)-3-{4-[(3-methoxypiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;

25 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)-N-methylacetamide;

1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-4-(1,1-dioxido-1,2-thiazolidin-2-yl)piperidine;

(3R)-1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}pyrrolidin-3-ol;

30 N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxyacetamide; 

- 4-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methyl]benzoic acid;
- (1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)(morpholin-4-yl)methanone;
- 5 (3S)-3-[4-(morpholin-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
- 8-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2,8-diazaspiro[4.5]decan-1-one;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carbonitrile;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-methylpiperidine-4-carboxamide;
- 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2,8-diazaspiro[4.5]decan-
- 10 1-one;
- N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-2-hydroxy-2-methylpropanamide;
- N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)-1-hydroxycyclopropanecarboxamide;
- 15 N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-ethylcyclopentanamine;
- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylpiperidine-4-carboxamide;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methylcyclopentanamine;
- 20 1-[(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-3-yl)methyl]pyrrolidin-2-one;
- 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-2-methylpyrrolidine;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-1-(pyrrolidin-1-yl)propan-2-amine;
- 25 N-cyclohexyl-N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N',N'-dimethylethane-1,2-diamine;
- N-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)acetamide;
- N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-methyl-2-(pyridin-2-
- 30 yl)ethanamine;
- (3S)-3-[4-(pyrrolidin-1-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine; ✓

- 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-3-carboxamide;
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidine-4-carboxamide;
 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}pyrrolidin-3-yl)acetamide;
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-(2-
 5 hydroxyethyl)piperidine-4-carboxamide;
 (3S)-3-[4-(1,4-oxazepan-4-ylmethyl)phenyl]-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;
 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N-(2-hydroxyethyl)piperidine-4-
 carboxamide;
 4-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)benzoic acid;
 10 1-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)urea;
 7-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-1,7-diazaspiro[4.4]nonan-2-one;
 8-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-2-methyl-2,8-
 diazaspiro[4.5]decan-1-one;
 1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-ol;
 15 N-(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-
 yl)methanesulfonamide;
 3-(1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)propan-1-
 ol;
 (3S)-3-{4-[(4-methylpiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-
 20 b]pyridine;
 N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-N-ethylethanamine;
 N-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1-
 (methylsulfonyl)piperidin-4-amine;
 (3S)-3-{4-[(4-fluoropiperidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-
 25 b]pyridine;
 1-(4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}-1,4-diazepan-1-
 yl)ethanone;
 [(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-yl]acetic acid;
 (1-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperidin-4-yl)methanol;
 30 4-[(1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-4-yl)methyl]benzoic
 acid; 

(3S)-3-{4-[(4-methyl-1,4-diazepan-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine;

(3S)-3-{4-[(3-methoxypyrrolidin-1-yl)methyl]phenyl}-2,3-dihydro[1,4]dioxino[2,3-b]pyridine; and

- 5 N-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}-N,2-dimethylpropan-2-amine; or a pharmaceutically salt thereof of each of the foregoing.

19. The compound 4-{4-[(3S)-2,3-dihydro[1,4]dioxino[2,3-b]pyridin-3-yl]benzyl}piperazine-1-carboxamide, or a pharmaceutically acceptable salt thereof.

10

20. The compound 1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidine-4-carboxylic acid, or a pharmaceutically acceptable salt thereof.

21. The compound [(3R)-1-{4-[(2S)-2,3-dihydro-1,4-benzodioxin-2-yl]benzyl}piperidin-3-yl]acetic acid, or a pharmaceutically acceptable salt thereof.

15

22. A pharmaceutical composition comprising the compound of any of claims 1 to 21, or a pharmaceutically acceptable salt thereof.

23. Use of a compound of any of claims 1 to 21, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating leukotriene-mediated disorders.

24. Use of a compound of any of claims 1 to 21, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a cardiovascular disease.

25

25. The use of claim 24, wherein the cardiovascular disease is selected from the group consisting of atherosclerosis, myocardial infarction, stroke, aortic aneurysm, sickle cell crisis, ischemia-reperfusion injury, pulmonary arterial hypertension and sepsis. ✓

30

26. Use of a compound of any of claims 1 to 21, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment of atherosclerosis. ✓

5

21 AOUT 2013
EKEME LYSAGHT Sarl
B.P. 6370
YAOUNDE - CAMEROUN
Tél.- Fax: 22 31 67 53

