



US006869778B2

(12) **United States Patent**
Madison

(10) **Patent No.:** **US 6,869,778 B2**
(45) **Date of Patent:** **Mar. 22, 2005**

(54) **TISSUE TYPE PLASMINOGEN ACTIVATOR (T-PA) VARIANTS: COMPOSITIONS AND METHODS OF USE**

WO	WO 88/05081	7/1988
WO	WO 88/10119	12/1988
WO	WO 89/00191	1/1989
WO	WO 90/02798	3/1990

(75) Inventor: **Edwin L. Madison**, San Diego, CA (US)

OTHER PUBLICATIONS

(73) Assignee: **The Scripps Research Institute**, La Jolla, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/705,633**

(22) Filed: **Nov. 10, 2003**

(65) **Prior Publication Data**

US 2004/0147009 A1 Jul. 29, 2004

Related U.S. Application Data

(62) Division of application No. 09/600,985, filed as application No. PCT/US97/20226 on Nov. 12, 1997, now Pat. No. 6,706,504.

(60) Provisional application No. 60/030,655, filed on Nov. 12, 1996.

(51) **Int. Cl.⁷** **C12P 21/06**

(52) **U.S. Cl.** **435/69.1**

(58) **Field of Search** 435/69.1; 530/300, 530/350

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,246,850 A	9/1993	Bennett et al.
5,520,913 A	5/1996	Anderson et al.
5,525,477 A	6/1996	Hassouna
5,616,486 A	4/1997	Anderson et al.

FOREIGN PATENT DOCUMENTS

EP	0 199 574	10/1986
EP	0 201 153	11/1986
EP	0 225 286	6/1987
EP	0 231 624	8/1987
EP	0 233 013	8/1987
EP	0 238 304	9/1987
EP	0 240 334	10/1987
EP	0 241 209	10/1987
EP	0 290 118	11/1988
EP	0 292 009	11/1988
EP	0 293 934	12/1988
EP	0 293 936	12/1988
EP	0 297 066	12/1988
EP	0 299 706	1/1989
EP	0 351 246	1/1990
WO	WO 87/04722	8/1987

Petersen, et al., Quenching of the amidolytic activity of one-chain tissue-type plasminogen activator by mutation of lysine-416, 1990, *Biochemistry*, 29(14):3451-3457.

Tachias, et al., Variants of tissue-type plasminogen activator which display substantially enhanced stimulation by fibrin, 1995, *J. Biol. Chem.*, 270(31):18319-18322.

Krause, et al., Properties of molecular variants of tissue-type plasminogen activator, 1989, *Arzneim.-Forsch.*, 39:632-637.

van Zonneveld, et al., Mapping of epitopes on human tissue-type plasminogen activator with recombinant deletion mutant proteins, 1987, *Thrombosis & Haemostasis*, 57(1):82-86.

Browne, et al., A tissue-type plasminogen activator mutant with prolonged clearance in vivo. Effect of removal of the growth factor domain, 1988, *J. Biol. Chem.*, 263:1599-1602.

Madison, et al., Serpin resistant mutants of human tissue-type plasminogen activator, 1989, *Nature*, 339:721-724.

Higgins, et al., The effect of the one-chain to two-chain conversion in tissue plasminogen activator: Characterization of mutations at position 275, 1990, *Thrombosis Res.*, 57:527-539.

Lamba, et al., "The 2.3 Å Crystal Structure of the Catalytic Domain of Recombinant Two-chain Human Tissue-type Plasminogen Activator", *J. Mol. Biol.* 258: 117-135 (1996).

Strandberg, et al., "Variants of Tissue-type Plasminogen Activator with Substantially Enhanced Response and Selectivity toward Fibrin Co-factors", *J. Biol. Chem.* 270(40): 23444-23449 (1995).

Tate, et al., "Functional Role of Proteolytic Cleavage at Arginine-275 of Human Tissue Plasminogen Activator as Assessed by Site-Directed Mutagenesis", *Biochemistry* 26: 338-343 (1987).

Primary Examiner—Jon Weber

Assistant Examiner—Sheridan Snedden

(74) *Attorney, Agent, or Firm*—Thomas Fitting; Michael J. McCarthy

(57) **ABSTRACT**

Variants of tissue plasminogen factor exhibit significantly enhanced fibrin stimulation, dramatically increased discrimination among fibrin co-factors, marked resistance to inhibition by PAI-1, and substantially increased zymogenicity, a combination of properties that enhance the therapeutic utility of the enzyme.

8 Claims, 5 Drawing Sheets

GTTC TGAGCACAGGGCTGGAGAGAAAACCTCTGCGAGGAAAGGGAAGGAGCAAGCCGTGA

ATTTAAGGGACGCTGTGAAGCAATC -35 met asp ala met lys -30 arg gly leu
ATG GAT GCA ATG AAG AGA GGG CTC

cys cys val leu leu leu cys gly ala val phe val ser pro ser
TGC TGT GTG CTG CTG CTG TGT GGA GCA GTC TTC GTT TCG CCC AGC

gln glu ⁻¹⁰ ile his ala arg phe arg arg gly ala arg ¹ SER TYR GLN
CAG GAA ATC CAT GCC CGA TTC AGA AGA GGA GCC AGA TCT TAC CAA

10
VAL ILE CYS ARG ASP GLU LYS THR GLN MET ILE TYR GLN GLN HIS
GTG ATC TGC AGA GAT GAA AAA ACG CAG ATG ATA TAC CAG CAA CAT

GLN²⁰ SER TRP LEU ARG PRO VAL LEU ARG SER ASN³⁰ ARG VAL GLU TYR
CAG TCA TGG CTG CGC CCT GTG CTC AGA AGC AAC CGG GTG GAA TAT

40
CYS TRP CYS ASN SER GLY ARG ALA GLN CYS HIS SER VAL PRO VAL
TGC TGG TGC AAC AGT GGC AGG GCA CAG TGC CAC TCA GTG CCT GTC

50
60
 LYS SER CYS SER GLU PRO ARG CYS PHE ASN GLY GLY THR CYS GLN
 AAA AGT TGC AGC GAG CCA AGG TGT TTC AAC GGG GGC ACC TGC CAG

70
GLN ALA LEU TYR PHE SER ASP PHE VAL CYS GLN CYS PRO GLU GLY
CAG GCC CTG TAC TTC TCA GAT TTC GTG TGC CAG TGC CCC GAA GGA

80 90
PHE ALA GLY LYS CYS CYS GLU ILE ASP THR ARG ALA THR CYS TYR
TTT GCT GGG AAG TGC TGT GAA ATA GAT ACC AGG GCC ACG TGC TAC

GLU ASP GLN GLY ILE SER ¹⁰⁰ TYR ARG GLY THR TRP SER THR ALA GLU
GAG GAC CAG GGC ATC AGC TAC AGG GGC ACG TGG AGC ACA GCG GAG

SER¹¹⁰ GLY ALA GLU CYS THR ASN TRP ASN SER SER¹²⁰ ALA LEU ALA GLN
AGT GGC GCC GAG TGC ACC AAC TGG AAC AGC AGC GCG TTG GCC CAG

LYS PRO TYR SER GLY ARG ¹³⁰ ARG PRO ASP ALA ILE ARG LEU GLY LEU
 AAG CCC TAC AGC GGG CGG AGG CCA GAC GCC ATC AGG CTG GGC CTG

140
 GLY ASN HIS ASN TYR CYS ARG ASN PRO ASP ARG 150
 GGG AAC CAC AAC TAC TGC AGA AAC CCA GAT CGA GAC SER LYS PRO
 TCA AAG CCC

FIG. 1A

160
TRP CYS TYR VAL PHE LYS ALA GLY LYS TYR SER SER GLU PHE CYS
TGG TGC TAC GTC TTT AAG GCG GGG AAG TAC AGC TCA GAG TTC TGC

170
SER THR PRO ALA CYS SER GLU GLY ASN SER ASP CYS TYR PHE GLY
AGC ACC CCT GCC TGC TCT GAG GGA AAC AGT GAC TGC TAC TTT GGG

180
ASN GLY SER ALA TYR ARG GLY THR HIS SER LEU THR GLU SER GLY
AAT GGG TCA GCC TAC CGT GGC ACG CAC AGC CTC ACC GAG TCG GGT

190
ALA SER CYS LEU PRO TRP ASN SER MET ILE LEU ILE GLY LYS VAL
GCC TCC TGC CTC CCG TGG AAT TCC ATG ATC CTG ATA GGC AAG GTT

200
TYR THR ALA GLN ASN PRO SER ALA GLN ALA LEU GLY LEU GLY LYS
TAC ACA GCA CAG AAC CCC AGT GCC CAG GCA CTG GGC CTG GGC AAA

210
HIS ASN TYR CYS ARG ASN PRO ASP GLY ASP ALA LYS PRO TRP CYS
CAT AAT TAC TGC CGG AAT CCT GAT GGG GAT GCC AAG CCC TGG TGC

220
HIS VAL LEU LYS ASN ARG ARG LEU THR TRP GLU TYR CYS ASP VAL
CAC GTG CTG AAG AAC CGC AGG CTG ACG TGG GAG TAC TGT GAT GTG

230
PRO SER CYS SER THR CYS GLY LEU ARG GLN TYR SER GLN PRO GLN
CCC TCC TGC TCC ACC TGC GGC CTG AGA CAG TAC AGC CAG CCT CAG

240
PHE ARG ILE LYS GLY GLY LEU PHE ALA ASP ILE ALA SER HIS PRO
TTT CGC ATC AAA GGA GGG CTC TTC GCC GAC ATC GCC TCC CAC CCC

250
TRP GLN ALA ALA ILE PHE ALA LYS HIS ARG ARG SER PRO GLY GLU
TGG CAG GCT GCC ATC TTT GCC AAG CAC AGG AGG TCG CCC GGA GAG

260
ARG PHE LEU CYS GLY GLY ILE LEU ILE SER SER CYS TRP ILE LEU
CGG TTC CTG TGC GGG GGC ATA CTC ATC AGC TCC TGC TGG ATT CTC

270
SER ALA ALA HIS CYS PHE GLN GLU ARG PHE PRO PRO HIS HIS LEU
TCT GCC GCC CAC TGC TTC CAG GAG AGG TTT CCG CCC CAC CAC CTG

280
THR VAL ILE LEU GLY ARG THR TYR ARG VAL VAL PRO GLY GLU GLU
ACG GTG ATC TTG GGC AGA ACA TAC CGG GTG GTC CCT GGC GAG GAG

290
300
310
320
330
340

FIG. 1B

350
GLU GLN LYS PHE GLU VAL GLU LYS TYR ILE VAL 360
GAG CAG AAA TTT GAA GTC GAA AAA TAC ATT GTC CAT AAG GAA TTC

370
ASP ASP ASP THR TYR ASP ASN ASP ILE ALA LEU LEU GLN LEU LYS
GAT GAT GAC ACT TAC GAC AAT GAC ATT GCG CTG CTG CAG CTG AAA

380
SER ASP SER SER ARG CYS ALA GLN GLU SER SER 390
TCG GAT TCG TCC CGC TGT GCC CAG GAG AGC AGC GTG GTC CGC ACT

400
VAL CYS LEU PRO PRO ALA ASP LEU GLN LEU PRO ASP TRP THR GLU
GTG TGC CTT CCC CCG GCG GAC CTG CAG CTG CCG GAC TGG ACG GAG

410
CYS GLU LEU SER GLY TYR GLY LYS HIS GLU ALA 420
TGT GAG CTC TCC GGC TAC GGC AAG CAT GAG GCC TTG TCT CCT TTC

430
TYR SER GLU ARG LEU LYS GLU ALA HIS VAL ARG LEU TYR PRO SER
TAT TCG GAG CGG CTG AAG GAG GCT CAT GTC AGA CTG TAC CCA TTC

440
SER ARG CYS THR SER GLN HIS LEU LEU ASN ARG 450
AGC CGC TGC ACA TCA CAA CAT TTA CTT AAC AGA ACA GTC ACC GAC

460
ASN MET LEU CYS ALA GLY ASP THR ARG SER GLY GLY PRO GLN ALA
AAC ATG CTG TGT GCT GGA GAC ACT CGG AGC GGC GGG CCC CAG GCA

470
ASN LEU HIS ASP ALA CYS GLN GLY ASP SER GLY 480
AAC TTG CAC GAC GCC TGC CAG GGC GAT TCG GGA GGC CCC CTG GTG

490
CYS LEU ASN ASP GLY ARG MET THR LEU VAL GLY ILE ILE SER TRP
TGT CTG AAC GAT GGC CGC ATG ACT TTG GTG GGC ATC ATC AGC TGG

500
GLY LEU GLY CYS GLY GLN LYS ASP VAL PRO GLY 510
GGC CTG GGC TGT GGA CAG AAG GAT GTC CCG GGT GTG VAL TYR THR LYS

520
VAL THR ASN TYR LEU ASP TRP ILE ARG ASP ASN MET ARG 527
GTT ACC AAC TAC CTA GAC TGG ATT CGT GAC AAC ATG CGA CCG TGA

FIG. 1C

CCAGGAACACCCGACTCCTCAAAAGCAAATGAGATCCCGCCTCTTCTTCTTCAGAAGACA
CTGCAAAGGCGCAGTGCTTCTCTACAGACTTCTCCAGACCCACCACCCGCAGAAGCGGG
ACGAGACCCTACAGGAGAGGGAAGAGTGCATTTTCCAGATACTTCCCATTTTGGAAGT
TTTCAGGACTTGGTCTGATTTTCAGGATACTCTGTCAGATGGGAAGACATGAATGCACACT
AGCCTCTCCAGGAATGCCTCCTCCCTGGGCAGAAAGTGGCCATGCCACCCTGTTTTTCAGCTA
AAGCCCAACCTCCTGACCTGTCACCGTGAGCAGCTTTGGAAACAGGACCACAAAAATGAA
AGCATGTCTCAATAGTAAAAGATAACAAGATCTTTCAGGAAAGACGGATTGCATTAGAA
ATAGACAGTATATTTATAGTCACAAGAGCCAGCAGGGCCTCAAAGTTGGGGCAGGCTGGC
TGGCCCGTCATGTTCTCAAAAGCACCCCTTGACGTCAAGTCTCCTTCCCCTTCCCCACT
CCCTGGCTCTCAGAAGGTATTCCTTTTGTGTACAGTGTGTAAAGTGTAATCCTTTTTCT
TTATAAACTTTAGAGTAGCATGAGAGAATTGTATCATTGAACAAGTGGCTTCAGCATA
TTTATAGCAATCCATGTTAGTTTTTACTTTCTGTTGCCACAACCCTGTTTTTATACTGTA
CTTAATAAATTCAGATATATTTTTTCACAGTTTTTCCAAAAAAAAAAAAAA

FIG. 1D

FIG. 2A




FIG. 2B




FIG. 2C





FIG. 2D



TISSUE TYPE PLASMINOGEN ACTIVATOR (T-PA) VARIANTS: COMPOSITIONS AND METHODS OF USE

REFERENCE TO RELATED APPLICATION

This application is a divisional of U.S. patent application Ser. No. 09/600,985 filed Nov. 13, 2000, now U.S. Pat. No. 6,706,504, pending, which is a national phase application of International Patent Application Serial No. PCT/US97/20226, filed Nov. 12, 1997 which claims the benefit of U.S. Provisional Application Ser. No. 60/030,655, filed Nov. 12, 1996, abandoned, which is hereby incorporated by reference.

GOVERNMENTAL RIGHTS

This invention was made with government support under Contract Nos. R01 52475 and P01 HL 31950 by the National Institutes of Health. The government has certain rights in the invention.

This invention was made with governmental support from the United States Government, National Institutes of Health, Grants HL52475 and HL31950; the United States Government has certain rights in the invention

The invention comprises protein single chain variants of tissue type plasminogen activator, also referred to as t-PA as well as nucleic acids encoding such protein single chain variants of tissue type plasminogen activator. The t-PA protein variants have higher zymogenicity than the wild-type single chain t-PA form. Methods of making and using the t-PA variant compositions are also described.

BACKGROUND

Tissue-type plasminogen activator (t-PA) is a serine protease that plays a critical role in the process of fibrinolysis, the dissolution of clots, by activating plasminogen to the protease plasmin. t-PA has been fully identified and characterized by underlying DNA sequence and deduced amino acid sequence. See Pennica et al., *Nature*, 301: 214 (1983) and U.S. Pat. No. 4,853,330, issued Aug. 1, 1989, the teachings of both of which are incorporated by reference. The nucleotide sequence and deduced primary amino acid sequence of human t-PA is depicted in FIG. 1A, FIG. 1B, FIG. 1C and FIG. 1D.

The group of amino acid residues from -35 to -1 preceding the sequence of the mature t-PA is the "pro" sequence. The mature t-PA molecule (amino acid residues 1-527) contains five domains that have been defined with reference to homologous or otherwise similar structures identified in various other proteins such as trypsin, chymotrypsin, plasminogen, prothrombin, fibronectin, and epidermal growth factor (EGF). These domains have been designated, starting at the N-terminus of the amino acid sequence of mature t-PA, as 1) the finger region (F) that has variously been defined as including amino acid residues 1 to about 44, 2) the growth factor region (G) that has been variously defined as stretching from about amino acid residues 45 to 91 (based upon its homology with EGF), 3) kringle one (K1) that has been defined as stretching from about amino acid residue 92 to about amino acid residue 173, 4) kringle two (K2) that has been defined as stretching from about amino acid residue 180 to about amino acid residue 261, and 5) the so-called serine protease domain (P) that generally has been defined as stretching from about amino acid residue 264 to the C-terminal end of the molecule at amino acid residue 527. These domains, which are

situated generally adjacent to one another, or are separated by short "linker" regions, account for the entire amino acid sequence of from 1 to 527 amino acid residues of the mature form of t-PA.

Each domain has been described variously as contributing certain specific biologically significant properties. The finger domain has been characterized as containing a sequence of at least major importance for high binding affinity to fibrin. (This activity is thought important for the high specificity that t-PA displays with respect to clot lysis at the locus of a fibrin-rich thrombus.) The growth factor-like region likewise has been associated with cell surface binding activity. The kringle 2 region also has been strongly associated with fibrin binding and with the ability of fibrin to stimulate the activity of t-PA. The serine protease domain is responsible for the enzymatic cleavage of plasminogen to produce plasmin.

t-PA is unusual among proteases in the level of the enzymatic activity of its precursor. In general, proteases are synthesized as zymogens, inactive precursors that must either be proteolytically processed or bind to a specific co-factor to develop substantial catalytic activity. The increase in catalytic efficiency after zymogen activation, or zymogenicity, is dramatic in almost all cases, although varying widely among individual members of the chymotrypsin family. For example, strong zymogens, i.e., those having high zymogenicity, such as trypsinogen, chymotrypsinogen, or plasminogen are almost completely inactive, with measured zymogenicities of 10^4 to 10^6 (Robinson, N. C., Neurath, H., and Walsh, K. A. (1973) *Biochemistry* 12, 420-426; Gertler, A., Walsh, K. A., and Neurath, H. (1974) *Biochemistry* 13, 1302-1310). Other serine proteases exhibit intermediate zymogenicity. For example, the enzymatic activity of Factor XIIa is 4000-fold greater than that of its corresponding zymogen, Factor XII (Silverberg, M., and Kaplan, A. P. (1982) *Blood* 60, 64), and the catalytic efficiency of urokinase is 250-fold greater than that of pro-urokinase (Lijnen, H. R., Van Hoef, B., Nelles, L., and Collen, D. (1990) *J. Biol. Chem.* 265, 5232-5236). By contrast, the catalytic activities of single and two chain t-PA vary by a factor of only 5-10.

The zymogenicity, expressed as the ratio of the activity of the mature cleaved two-chain enzyme to that of the single chain precursor form, is only 5-10 for wild-type t-PA, in contrast to other precursors of other proteases that have little or no catalytic activity. Thus, the single chain form of wild-type t-PA is not a true zymogen.

There have been many attempts to improve the usefulness of t-PA by genetic engineering. These efforts have been only partially successful. The state of the art has been reviewed by Krause, J., & Tanswell, P. *Arzneim.-Forsch.* 39: 632-637 (1989) and in U.S. Pat. No. 5,616,486, the teachings of both of which are incorporated by reference.

Despite the profound advantages associated with natural t-PA as a clot-dissolving agent, it is not believed that the natural protein necessarily represents the optimal t-PA agent under all circumstances. Therefore, several variants have been proposed or devised to enhance specific properties of t-PA. Certain of those variants address disadvantages associated with the use of natural t-PA in situations where an agent with a longer half-life or slower clearance rate would be preferred, e.g., in the treatment of deep-vein thrombosis and following reperfusion of an infarct victim, or where a single-chain agent is preferred.

For example, removal of a substantial portion or all of the finger domain results in a molecule with substantially dimin-

ished fibrin binding characteristics, albeit in return there is a decrease in the overall rate of clearance of the resultant entity—See WO 89/00197 published Jan. 12, 1989.

Variants are described in EPO Pat. Publ. No. 199,574 that have amino acid substitutions at the proteolytic cleavage sites at positions 275, 276, and 277. These variants, characterized preferentially as t-PA variants having an amino acid other than arginine at position 275, are referred to as protease-resistant one-chain t-PA variants in that, unlike natural t-PA, which can exist in either a one-chain or two-chain form, they are resistant to protease cleavage at position 275 and are therefore not converted metabolically in vivo into a two-chain form. This form is thought to have certain advantages biologically and commercially, in that it is more stable while the fibrin binding and fibrin stimulation are increased relative to two-chain t-PA. Furthermore, plasminogen activators are described that comprise one domain capable of interacting with fibrin and the protease domain of urokinase, with one embodiment being urokinase altered to make it less susceptible to forming two-chain urokinase. See WO 88/05081 published Jul. 14, 1988.

For further patent literature regarding modification of the protease cleavage site of t-PA, see, for example, EPO Pat. Nos. 241,209; EP 201,153 published Nov. 12, 1986; EP 233,013 published Aug. 19, 1987; EP 292,009 published Nov. 23, 1988; EP 293,936 published Dec. 7, 1988; and EP 293,934 published Dec. 7, 1988; and WO 88/10119.

Glycosylation mutants at positions 117–119, 184–186, and 448–450 exhibited higher specific activity as the mole percent carbohydrate was reduced. See EPO Pub. No. 227,462 published Jul. 1, 1987. This patent application additionally discloses using an assay of fibrin/fibrin degradation products and teaches that one may also modify the t-PA molecule at positions 272–280 or delete up to 25 amino acids from the C-terminus. Further, the t-PA mutants with Asn 119, Ala 186 and Asn 450, which have the N-glycosylation sites selectively removed by DNA modification but contain residual O-linked carbohydrate, were found to be about two-fold as potent as melanoma t-PA in an in vitro lysis assay. See EPO Publ. No. 225,286 published Jun. 10, 1987.

Replacement of the amino acid at position 449 of t-PA with any amino acid except arginine to modify the glycosylation site, as well as modification of Arg 275 or deletion of the –3 to 91 region, is also taught. See WO 87/04722 published Aug. 13, 1987. An amino acid substitution at position 448 of t-PA is disclosed as desirable to remove glycosylation. See EPO Pub. No. 297,066 published Dec. 28, 1988. The combination of modifications at positions 448–450 and deletion of the N-terminal 1–82 amino acids is disclosed by WO 89/00191 published Jan. 12, 1989. Additionally, urokinase has been modified in the region of Asp 302-Ser 303-Thr 304 to prevent glycosylation. See EPO Pub. No. 299,706 published Jan. 18, 1989.

However, alteration of the glycosylation sites, and in particular that at amino acid 117, seems invariably to result in a molecule having affected solubility characteristics that may result additionally in an altered circulating half-life pattern and/or fibrin binding characteristics. See EPO Pat. Publ. No. 238,304, published Sep. 23, 1987.

When the growth factor domain of t-PA is deleted, the resultant variant is still active and binds to fibrin, as reported by A. J. van Zonneveld et al., *Thrombos. Haemostas.* 54 (1): 4 (1985). Various deletions in the growth factor domain have also been reported in the patent literature. See EPO Publ. No. 241,209 (del-51–87), EPO Publ. No. 241,208 (del-

51–87 and del-51–173), PCT 87/04722 (deletion of all or part of the N-terminal 1–91), EPO Publ. No. 231,624 (all of growth factor domain deleted), and EPO Publ. No. 242,836 and Jap. Pat. Appl. Kokai No. 62-269688 (some or all of the growth factor domain deleted).

It has further been shown that t-PA can be modified both in the region of the first kringle domain and in the growth factor domain, resulting in increased circulatory half-life. See EPO Pat. Publ. No. 241,208 published Oct. 14, 1987. The region between amino acids 51 and 87, inclusive, can be deleted from t-PA to result in a variant having slower clearance from plasma. Browne et al., *J. Biol. Chem.*, 263: 1599–1602 (1988). Also, t-PA can be modified, without adverse biological effects, in the region of amino acids 67 to 69 of the mature, native t-PA, by deletion of certain amino acid residues or replacement of one or more amino acids with different amino acids. See EPO Pat. Publ. No. 240,334 published Oct. 7, 1987.

A hybrid of t-PA/urokinase using the region of t-PA encompassing amino acids 273–527 is also disclosed. See EPO 290,118 published Nov. 9, 1988. Serpin-resistant mutants of human t-PA with alterations in the protease domain, including del296–302 t-PA, R304S t-PA, and R304E t-PA, are disclosed in Madison et al., *Nature*, 339: 721–724 (1989). The above list is not an exhaustive review of the numerous variants of t-PA that have described.

As a result of the catalytic activity of precursor t-PA, despite effective clot lysis at targeted sites, undesirable proteolysis occurs systemically resulting in the deleterious depletion of circulating fibrinogen, α_2 -anti-plasmin and plasminogen. What is needed are more zymogenic t-PA variant proteins that provide effective local clot lysis is achieved with diminished systemic proteolytic effects.

SUMMARY OF THE INVENTION

The present invention provides single chain variant t-PA proteins having at least two substitutions of basic amino acid residues by neutral or acidic amino acid residues, compared to the wild-type human t-PA, as well as polynucleotides encoding such single chain variant t-PA proteins. The single chain variant t-PA proteins of the present invention have the R275 amino acid residue substituted by an amino acid residue chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid, and glutamic acid. Preferably the single chain variant t-PA proteins of the present invention have the R275 amino acid residue substituted by an amino acid residue chosen from the group consisting of an aspartic acid residue and a glutamic acid residue, and most preferably by a glutamic acid residue.

The single chain variant t-PA proteins of the present invention have additionally at least one other basic amino acid residue in the serine protease region residue substituted by a non-basic amino acid such that the salt bridge interaction normally found in wildtype single chain t-PA between aspartate 477 and lysine 429 is disrupted. Preferably, basic amino acids are replaced with polar or acidic amino acids, and more preferably, amino acid residues chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid and glutamic acid.

The salt bridge interaction between aspartate 477 and lysine 429 can be disrupted by a substitution at position 477 or 429, or by an appropriate substitution at a position within the serine protease region that provides an alternative salt bridge interaction partner for at least one of aspartate 477 and lysine 429. In one preferred embodiment, the H417 amino acid residue is substituted by an amino acid residue

5

chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid, and glutamic acid. More preferably the single chain variant t-PA proteins of the present invention have both the R275 amino acid residue and the H417 amino acid residue substituted by an amino acid residue chosen from the group consisting of an aspartic acid residue and a glutamic acid residue. Two exemplary preferred single chain variant t-PA proteins are the t-PA variants designated as R275E, H417E and R275E, H417D.

In another preferred embodiment, the K429 amino acid residue is substituted by an amino acid residue chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid, and glutamic acid. More preferably the single chain variant t-PA proteins of the present invention have both the R275 amino acid residue and the K429 amino acid residue substituted by an amino acid residue chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid, and glutamic acid. One preferred single chain variant t-PA protein is the t-PA variant designated as R275E, K429Y.

The single chain variant t-PA proteins of the present invention exhibit greater zymogenicity, expressed as the ratio of the activity of the mature cleaved two-chain enzyme to that of the single chain precursor form, than that of the wild type single chain t-PA protein. The single chain variant t-PA proteins of the present invention have zymogenicity of at least 10, preferably about 50 to about 200.

The single chain variant t-PA proteins of the present invention exhibit a greater fibrin stimulation factor, expressed as the ratio of the catalytic efficiencies in the presence and absence of fibrin, compared to the wild type single chain t-PA protein. The single chain variant t-PA proteins of the present invention have a fibrin stimulation factor of at least 7,000, preferably about 20,000 to about 50,000.

The single chain variant t-PA proteins of the present invention exhibit a reduced inhibition by plasminogen activator inhibitor 1 (PAI-1) to the wild type single chain t-PA protein. The single chain variant t-PA proteins of the present invention are at least a factor of 5, preferably at least a factor of about 9, most preferably at least a factor of about 200 less inhibited by PAI-1 compared to the wild type single chain t-PA protein.

The single chain variant t-PA proteins of the present invention exhibit a greater fibrin selectivity factor, expressed as the ratio of the catalytic efficiencies in the presence fibrin to that in the presence of fibrinogen, compared to the wild type single chain t-PA protein. Preferred embodiments of the single chain variant t-PA proteins of the present invention have a fibrin selectivity factor of at least 10, preferably at least 50, more preferably at least 100.

BRIEF DESCRIPTION OF THE DRAWINGS

In the drawings,

FIGS. 1A, 1B, 1C, and 1D show the nucleotide sequence (SEQ ID NO:11) and deduced amino acid sequence (SEQ ID NO:12) of the full-length human t-PA cDNA; and

FIG. 2 is a graphical representation of the results of standard chromogenic assays of plasminogen activation in the presence of buffer (open squares), DESAFIB (open diamonds), fibrinogen (open circles), cyanogen bromide fragments of fibrinogen (open triangles) or the stimulatory peptide P368 (hatched squares).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

As used herein, "wild-type t-PA" refers to the t-PA protein naturally occurring in humans. While this human t-PA is

6

exemplified by the amino acid sequence depicted in FIGS. 1A, 1B and 1C, the term wild-type t-PA should be understood to encompass naturally occurring allelic variations. t-PA Variant Compositions

The t-PA variant cDNAs and the corresponding expressed recombinant proteins of this invention are useful compounds that function in the serine protease-mediated control of fibrinolysis as described herein.

The t-PA variant cDNAs of the present invention contain at least one nucleotide substitution to generate a t-PA cDNA that encodes a noncleavable single chain t-PA variant, i.e., not cleavable by plasmin under normal conditions. The nucleotide substitution results in a substitution of a glutamic acid (E) for an arginine (R) at amino acid residue 275 (or position 15 using the chymotrypsin numbering system) in the t-PA precursor that is responsible for creating a non-cleavable variant. Positions 15, 144, 156, and 194 of the chymotrypsin numbering system correspond to positions 275, 417, 429, and 477, respectively, in the t-PA numbering system as depicted in FIG. 1.

The variants, which are substitution mutants, are designated by the single letter code of the wild type human t-PA amino acid residue, the position of the residue relative to the amino terminus of the mature human t-PA as depicted in FIG. 1, followed by the single letter code of the amino acid residue substituted for the amino acid residue in mature human t-PA. The substitution of glutamic acid for arginine at position 275 is designated as R275E. Equivalent substitutions generating noncleavable single chain t-PA are known in the art (Higgins, D. L., et al., (1990) *Thrombosis Res.* 57: 527-539).

In addition to the R275E substitution, the variant cDNAs of the present invention further comprise at least one other nucleotide substitution at a separate site to create a t-PA variant having at least two amino acid substitutions. Preferred cDNA variants include at least one nucleotide substitution that results in an amino acid substitution of an amino acid residue chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid, and glutamic acid for a histidine at amino acid residue position 417. Preferred embodiments are designated R275E, H417D and R275E, H417E. A further cDNA variant comprises at least one nucleotide substitution resulting in the substitution of an amino acid residue chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid, and glutamic acid for the lysine (K) at amino acid residue position 429. One such preferred embodiment is designated R275E, K429Y.

The variant t-PA cDNAs of the present invention are useful for generating the recombinant expressed variant t-PAs described above. In a further embodiment, the variant t-PA cDNAs have therapeutic uses in gene therapy as described below.

The invention includes embodiments such as expression vectors or plasmids in which the cDNAs for encoding variant t-PAs are operably linked to provide for the expression of recombinant variant t-PAs for use in the methods as described below. One preferred embodiment is the expression of a variant t-PA protein by COS 1 cells comprising pSVT7 expression vector operably linked to a polynucleotide encoding the variant protein. Constitutive and inducible expression vectors are disclosed. In a further embodiment, transiently and stably transfected cells contain cDNA encoding variant t-PAs.

The resultant recombinant expressed t-PA variants described herein are characterized as having one or more of the following structural and functional properties: 1) The

t-PA variant is in the form of a noncleavable single chain protein containing an R275E amino acid substitution or equivalents thereof that prevent cleavage by t-PA activating enzymes; 2) The t-PA variant exhibits increased resistance to inhibition by the serpin plasminogen activator inhibitor, type I (PAI-1); 3) The t-PA variants has diminished catalytic activity on substrates, such as plasminogen, in the absence of co-factors, such as fibrin; 4) The t-PA variants exhibit enhanced stimulation by fibrin; 5) The t-PA variants exhibit comparable catalytic activity to substrates, such as plasminogen, in the presence of co-factors, such as fibrin; and 6) In view of the proceeding properties, the t-PA variants thus are effective at local fibrinolysis function without extensive systemic proteolysis thereby negating the depletion of circulating fibrinogen, α 2-anti-plasmin and plasminogen, as is seen with wild type human single chain t-PA precursor.

Preferred recombinant expressed t-PA variants thus include R275E,H417D, R275E,H417E and R275E,K429Y, and conservative substitutions thereof. In general examples of conservative substitutions include the substitution of one non-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another. For further discussion of the classifications of amino acids see Lehninger, A. L., Biochemistry, 2nd Edition, Worth Publishers, New York, 1975, pp.71-94.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that such protein displays the requisite binding activity. "Chemical derivative" refers to a subject protein having one or more residues chemically derivatized by reaction of a functional side group. Such derivatized molecules include for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl-groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-im-benzylhistidine. Also included as chemical derivatives are those peptides which contain one or more naturally occurring amino acid derivatives of the twenty standard amino acids. For example, 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine. D-amino acids may also be included in place of one or more L-amino acids.

In the specific case of the present invention, basic amino acids, i.e., arginine, lysine and histidine are replaced with non-basic amino acids. Preferably basic amino acids are replaced with polar or acidic amino acids, i.e. amino acid residues chosen from the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid and glutamic acid. Conservative substitutions are thus defined, for the purpose of the present invention, as meaning that non-basic amino acids replacing particular basic amino acids in mature wild type human t-PA may be chosen from the group of non-basic amino acids generally, preferably from

the group consisting of glycine, serine, threonine, asparagine, tyrosine, glutamine, aspartic acid and glutamic acid, and more preferably from the group consisting of tyrosine, aspartic acid and glutamic acid. For example, the use of aspartic acid instead of glutamic acid to replace an histidine residue is a conservative substitution. Preferred variants are R275E,H417D and R275E,H417E, described in Example 1 and the R275E,K429Y variant, described in Example 2.

The expressed recombinant t-PA variants having at least two amino acid substitutions, e.g., R275E,H417D, R275E,H417E and R275E,K429Y, further exhibit unique properties. R275E and R275E,H417E are activated by both fibrinogen and fibrin while R275E,K429Y is activated primarily by fibrin and is not sensitive to fibrinogen. The latter is also more resistant than the R275E,H417D and R275E,H417E variants to inhibition by PAI-1. These characteristics provide additional advantages in administering the compounds as therapeutic thrombolytic compositions as further described below. In addition, the t-PA variants described herein are useful in diagnostic applications as described below.

Methods of Making and Using t-PA Variant Compositions

Methods of Making

The t-PA variant cDNA and recombinant expressed variant proteins described above are useful in a number of methodological aspects as described in Examples 1 and 2. In particular, the isolated cDNA clones are useful in an expression vector system to produce encoded t-PA variant proteins of this invention. Thus, expression vector systems having a t-PA variant cDNA operably linked therein, including cells containing the expression vectors, are contemplated for generating the recombinant expressed variant proteins of this invention.

Diagnostic Applications

Preferred diagnostic methodological aspects are described herein. In particular, the recombinant expressed t-PA variants of the present invention are useful in diagnostic assays to detect fibrin and fibrin degradation products that have altered activities. The assays are thus indicated in thrombotic conditions. Other diagnostic applications, including kits comprising antibodies against the t-PA variants are familiar to one of ordinary skill in the art.

Therapeutic Applications

The t-PA variant cDNAs of the present invention are useful in genetic therapeutic applications for use in ameliorating thrombotic disorders including both acute and chronic conditions. Acute conditions include among others both heart attack and stroke while chronic situations include those of arterial and deep vein thrombosis and restenosis. Preferred therapeutic compositions thus include the cDNA compounds themselves as naked DNA, presented as part of a viral vector delivery system or other vector-based gene expression delivery system, presented in a liposome delivery system and the like.

The recombinant expressed t-PA variant proteins of the present invention are contemplated as thrombolytic therapeutic agents for ameliorating the same conditions outlined above. Based on the individual structural and functional properties of various t-PA variant proteins described above, the selection of the particular t-PA variant is determined by the desired therapeutic outcome. For example, the fibrinogen-mediated activation of endogenous human t-PA is activated by bleeding which then results in a widespread undesired systemic response. Thus, to mediate fibrinolytic processes locally in either an acute or chronic thrombotic condition while simultaneously preventing proteolytic activation systemically, one would therefore use the t-PA

variant, namely R275E,K429Y, that is primarily activated by fibrin and not fibrinogen. A composition for use as thrombolytic therapeutic agents generally a physiologically effective amount of the t-PA variant protein in a pharmaceutically suitable excipient. Depending on the mode of administration and the condition to be treated, the thrombolytic therapeutic agents are administered in single or multiple doses. If "bolus" doses are administered, doses of about 0.01 to about 0.6 mg/kg will typically be administered, preferably doses of about 0.05 to about 0.2 mg/kg, with subsequent administrations of about 0.1 to about 0.2 mg/kg to maintain a t-PA blood level of about 3 microgram/ml. One skilled in the art will appreciate that variations in dosage depend on the condition to be treated. In other applications, a composition of variant t-PA in a gel composition is useful in preventing the formation of adhesions.

Other variations and uses of the present invention will be apparent to one skilled in the art.

EXAMPLE 1

Site Directed Mutagenesis and Construction of Expression Vectors Encoding Variants of T-PA

Oligonucleotide directed site specific mutagenesis was performed by the method of Zoller and Smith (Zoller, M. I., and Smith, M. (1984) DNA 3, 479488) as modified by Kunkel (Kunkel; T. A. (1985) Proc. Natl. Acad. Sci. U.S.A. 82, 488492). Mutations were introduced into the 290 bp SacI-SmaI fragment of cDNA encoding t-PA that had been previously subcloned into bacteriophage M13mp18. The mutagenic primers had the following nucleotide sequences:

H417D:
5'-CTACGGCAAGGACGAGGCCTTGT-3' (SEQ ID NO: 8)

H417E:
5'-CTACGGCAAGGAGGAGGCCTTGT-3' (SEQ ID NO: 9)

Following mutagenesis, ssDNA corresponding to the entire 290 bp SacI-SmaI fragment was fully sequenced to assure the presence of the desired mutation and the absence of any additional mutations. The sequence corresponding to the 290 bp SacI-SmaI fragment of the H417D mutation is shown in SEQ ID NO: 5; the corresponding sequence of the H417E mutation is shown in SEQ ID NO: 6. Replicative form (RF) DNA was prepared for appropriate phage, and the mutated 290 bp SacI-SmaI fragments were recovered after digestion of RF DNA with SacI and SmaI and electrophoresis of the digestion products on an agarose gel. The isolated, mutated SacI-SmaI fragments were used to replace the corresponding fragment in full length cDNAs encoding wild type human t-PA or t-PA/R275E to yield new, full length cDNAs encoding t-PA/H417D; t-PA/H417E; t-PA/R275E, H417D (SEQ ID NO: 1); and t-PA/R275E, H417E (SEQ ID NO: 2).

Expression of Enzymes by Transient Transfection of COS Cells.

cDNAs encoding t-PA; t-PA/R275E; t-PA/H417D; t-PA/H417E; t-PA/R275E, H417D; and t-PA/R275E were ligated into the transient expression vector pSVT7 which is described in Madison, E. L., et al. (1989) *Nature* 339, 721-724; Bird, P. M., et al., (1987) *J. Cell Biol.* 105: 2905-2914; and Sambrook, J., et al., (1986) *Mol. Biol. Med.* 3: 459481. See also U.S. Pat. No. 5,550,042, incorporated herein by reference, which describes the construction and use of pSVT7 as well as the deposit with American Type Culture Collection, 12301 Parklawn Dr., Rockville, Md.

20852 of cultures comprising other pSVT7 t-PA constructs. Vectors with ligated cDNA inserts were then introduced into COS 1 cells by electroporation using a Bio Rad Gene Pulser. An aliquot containing 20 μ g of cDNA, 100 μ g of carrier DNA and approximately 10^7 COS cells were placed into a 0.4 cm cuvette, and electroporation was performed at 320 V, 960 μ FD, and $\Omega n = \infty$. Following electroporation, cells were incubated overnight at 37 degrees Celsius in DMEM containing 10% fetal calf serum and 5 mM sodium butyrate. Cells were then washed with serum free medium and incubated in DMEM for 48 hours at 37 degrees Celsius. After the incubation with serum free media, conditioned media were collected. Enzyme concentrations in aliquots of the collected conditioned media were determined by ELISA.

Kinetic Analysis of Plasminogen Activation Using Indirect Chromogenic Assays.

Indirect chromogenic assays of t-PA utilized the substrates lys-plasminogen (American Diagnostica, Greenwich, Conn.) and Spectrozyme PL (American Diagnostica) and were performed as previously described (Madison, E. L., Goldsmith, E. J., Gerard, R. D., Gething, M.-J., and Sambrook, J. F. (1989) *Nature* 339, 721-724; Madison, E. L., Goldsmith, E. J., Gerard, R. D., Gething, M. J., Sambrook, J. F., and Bassel-Duby, R. S. (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87, 3530-3533; Madison, E. L., Goldsmith, E. J., Gething, M. J., Sambrook, J. F., and Gerard, R. D. (1990) *J. Biol. Chem.* 265, 21423-21426.). Assays were performed both in the presence and absence of the co-factor DESAFIB (American Diagnostica). The concentration of lys-plasminogen was varied from 0.0125-0.2 μ M in the presence of DESAFIB and from 0.9-15 μ M in the absence of the co-factor.

Kinetic Analysis of t-PA Activity Using a Small, Synthetic Substrate

The direct chromogenic assay utilized the substrate methylsulfonyl-D-cyclohexyltyrosyl-glycyl-arginine-p-nitroaniline (Spectrozyme t-PA, American Diagnostica) and was performed as previously described (Strandberg, L., and Madison, E. L. (1995) *J. Biol. Chem.* 270, 23444-23449; Smith, J. W., Tachias, K., and Madison, E. L. (1995) *J. Biol. Chem.* 270, 30486-30490).

Measurement of Second Order Rate Constants for Inhibition by PAI-1

Second order rate constants for the inhibition of wild type human t-PA and variant t-PA were measured under pseudo-first order conditions as previously described. Briefly, enzyme and inhibitor were preincubated at 23 degrees Celsius for periods of time varying from 0-30 minutes. Following preincubation, the mixtures were diluted, and the residual enzymatic activity was measured in a standard indirect chromogenic assay. For each enzyme, the concentrations of enzyme and inhibitor and the times of preincubation were chosen to yield several data points for which the residual enzymatic activity varied between 20% and 80% of the initial activity. Data were analyzed by plotting the natural logarithm of the ratio (residual activity/initial activity) versus time of preincubation and measuring the resulting slopes. Division of this slope by $-[I]$ yielded the second order rate constants shown.

It was found that replacement of histidine 417 of t-PA with an acidic residue selectively suppresses the catalytic activity of single chain t-PA. Histidine 417 was replaced by either an aspartate or glutamate residue to yield two variants: t-PA/H417D and t-PA/H417E. Accurate measurement of the enzymatic activity toward plasminogen of the single chain form of these two variants proved difficult, however,

because the plasmin produced in this assay rapidly converted the single chain enzymes into their mature, two-chain form by cleaving the R275–1276 peptide bond. Consequently, to overcome this technical difficulty, we also constructed noncleavable forms of the two mutated enzymes by introducing the additional mutation R275E into the existing mutants.

Wild type human t-PA, t-PA/R275E, and all four variants containing mutations at position 417 were expressed by transient expression of COS-1 cells. Since this procedure yielded predominantly single chain enzymes, two-chain t-PAs were generated by treating the enzyme preparations with plasmin-Sepharose (Strandberg, L., and Madison, E. L. (1995) *J. Biol. Chem.* 270, 23444–23449). Quantitative conversion of the enzymes into their mature, two-chain form was confirmed by SDS-PAGE. As expected, variants containing the mutation R275E were synthesized and secreted exclusively as single chain enzymes and were not cleaved by plasmin-Sepharose.

The enzymatic activity of the single and two-chain forms of wild type human t-PA and each variant toward a small synthetic substrate is listed in Table I below. Mutation of histidine 417 had only very modest effects on the activity of the two-chain enzymes. Two-chain t-PA/H417D and t-PA/H417E displayed 67% or 80%, respectively, the activity of the two-chain, wild type human t-PA enzyme in this assay. The H417D and H417E mutations, however, had more significant effects on the activities of the single chain enzymes. Compared to single chain t-PA/R275E, single chain t-PA/R275E,H417D (SEQ ID NO: 1) and t-PA/R275E, H417E (SEQ ID NO: 2) exhibited approximately 16% or 25%, respectively, the activity of single chain t-PA/R275E.

TABLE 1

Kinetic constants for cleavage of the chromogenic substrate Spectrozyme t-PA by single- and two-chain t-PA variants			
Enzyme	K_{cat} (s^{-1})	K_m (mM)	K_{cat}/K_m ($M^{-1}s^{-1}$)
<u>Two-chain form</u>			
t-PA	59	0.4	1.5×10^5
t-PA/H417D	41	0.4	1.0×10^5
t-PA/H417E	58	0.5	1.2×10^5
<u>Single-chain form</u>			
t-PA/R275E	26	0.7	3.7×10^4
t-PA/R275E, H417D	5.9	1.0	5.9×10^3
t-PA/R275E, H417E	12	1.3	9.2×10^3

All of the variants analyzed maintained high enzymatic activity towards the natural substrate, plasminogen, in the presence of the co-factor fibrin (Table II below). The catalytic activity of the two-chain form of wild type human t-PA, t-PA/H417D, and t-PA/H417E varied by a factor of only 1.4. Similarly, the activities of single chain t-PA/R275E, t-PA/R275E,H417D, and t-PA/R275E,H417E differed by a factor of less than 1.8.

TABLE II

Kinetic constants for activation of plasminogen by single- and two-chain t-PA variants in the presence of fibrin			
Enzyme	K_{cat} (s^{-1})	K_m (μM)	K_{cat}/K_m ($M^{-1}s^{-1}$)
<u>Two-chain form</u>			
t-PA	0.11	0.017	6.5×10^6
t-PA/H417D	0.11	0.024	4.6×10^6
t-PA/H417E	0.10	0.022	4.5×10^6

TABLE II-continued

Kinetic constants for activation of plasminogen by single- and two-chain t-PA variants in the presence of fibrin			
Enzyme	K_{cat} (s^{-1})	K_m (μM)	K_{cat}/K_m ($M^{-1}s^{-1}$)
<u>Single-chain form</u>			
t-PA/R275E	0.16	0.017	9.4×10^6
t-PA/R275E, H417D	0.23	0.043	5.3×10^6
t-PA/R275E, H417E	0.17	0.028	6.1×10^6

In the absence of a co-factor, the mutations at position 417 had little effect on the activity of two-chain t-PA toward plasminogen; however, these mutations significantly reduced the catalytic efficiency of single chain t-PA (Table III below). Compared to that of single chain t-PA/R275E, the activity of t-PA/R275E,H417D and t-PA/R275E,H417E was reduced by a factor of approximately 14 or 6, respectively. In this assay, the “zymogenicity”, or ratio of the activities of the two-chain and single chain form of a particular enzyme, were approximately 9 for wild type t-PA. By contrast, for variants containing the H417D or H417E mutation, this ratio increased to approximately 150 or 50, respectively (Table III).

TABLE III

Kinetic constants for activation of plasminogen by single- and two-chain variants of t-PA in the absence of a cofactor			
Enzyme	K_{cat} (s^{-1})	K_m (μM)	K_{cat}/K_m ($M^{-1}s^{-1}$)
<u>Two-chain form</u>			
t-PA	0.093	6.7	1.4×10^4
t-PA/H417D	0.110	6.8	1.6×10^4
t-PA/H417E	0.099	8.7	1.1×10^4
<u>Single-chain form</u>			
t-PA/R275E	0.014	9.5	1.5×10^3
t-PA/R275E, H417D	0.001	9.4	1.1×10^2
t-PA/R275E, H417E	0.002	8.5	2.4×10^2

Molecular details of the stimulation of t-PA by fibrin, a complex process that almost certainly involves multiple points of contact between the two proteins, remain unclear. While fibrin stimulation of two-chain t-PA may occur through a single mechanism; stimulation of single chain t-PA by fibrin co-factors, however, appears to utilize at least two distinct mechanisms. First, fibrin apparently stimulates both single- and two-chain t-PA through a templating mechanism resulting in formation of a ternary complex which greatly augments the local concentration of enzyme and substrate. Second, because single- and two-chain t-PA have equivalent activity in the presence but not the absence of fibrin, it seems likely that binding to fibrin induces a conformational change in the activation domain of single chain t-PA. Similar activation of plasminogen upon binding to streptokinase as well as activation of prothrombin by binding to staphylocoagulase have been described previously. Although the mechanism of this nonclassical, nonproteolytic activation of serine protease zymogens remains completely unclear, the behavior of single chain t-PA/R275E,H417D and t-PA/R275E,H417E indicates that His 417 does not play an essential role in this process. In addition, the properties of two-chain t-PA/H417D and t-PA/H417E indicate that His 417 does not play an essential role during zymogen activation of t-PA through the classical, proteolytic mechanism.

The primary effect of the H417D and H417E mutations was a selective reduction of the activity of single chain t-PA

13

in the absence of fibrin and, consequently, an increase in the zymogenicity of the enzyme. At the molecular level this effect could be mediated either by stabilizing a less active, new conformation of single chain t-PA or by shifting the equilibrium between one or more existing conformations, with distinct activities, towards the less active conformation. Without being held to a single hypothesis, based on structural studies of trypsinogen, trypsin, chymotrypsinogen, and chymotrypsin, that the existence of an equilibrium among multiple conformations of the activation domain is likely to be a general feature of chymotrypsinogen family zymogens.

It is believed that the effect produced by converting His 417 to an acidic residue is mediated by disrupting the important salt bridge between Asp 477 and Lys 429 by providing an alternative, electrostatic interaction for Lys 429. The observation of an electrostatic interaction between K429 and E417 in the recently reported structure of the protease domain of two-chain u-PA, although the distance and geometry of this interaction vary somewhat in the two members of the unit cell in this study, lends credence to this hypothesis. Moreover, as observed in this study, formation of a new salt bridge between Lys 429 and Asp/Glu 417 would be expected to selectively suppress the activity of single chain t-PA because Lys 429 does not interact with Asp 477 in the two-chain enzyme. Instead, in two-chain t-PA, as in other mature chymotrypsin like enzymes, the mature amino terminus inserts into the activation pocket and plays this role. Consequently, as observed, two-chain t-PA/H417D and t-PA/H417E are expected to maintain high catalytic activity. Variants of t-PA containing an acidic residue at position 417, therefore, exhibit significantly enhanced zymogenicity.

TABLE IV

Stimulatory effect of fibrin on the catalytic efficiencies for variants of t-PA	
Enzyme	Fold stimulation of k_{cat}/K_m
<u>Two-Chain form</u>	
t-PA	460
t-PA/H417D	290
t-PA/H417E	410
<u>Single-chain form</u>	
t-PA/R275E	6300
t-PA/R275E, H417D	48,200
t-PA/R275E, H417E	25,400

The extent of fibrin stimulation displayed by the single chain form of the mutated enzymes examined in this study is significantly greater than that displayed by wild type t-PA. Wild type, two-chain t-PA possesses a fibrin stimulation factor, defined as the ratio of the catalytic efficiencies in the presence and absence of fibrin, of approximately 460 (Table IV above). The two-chain variants display similar stimulation factors of 290 (t-PA/H417D) and 410 (t-PA/H417E). Single chain wild type t-PA, with a fibrin stimulation factor of 6300, is stimulated to a substantially greater degree than the two-chain enzymes, presumable reflecting the ability of fibrin to stimulate the single chain enzymes not only through a templating mechanism but also by inducing nonproteolytic zymogen activation. Stimulation of single chain t-PA is further enhanced by the H417D or H417E mutations. The fibrin stimulation factors for single chain t-PA/R275E, H417D and t-PA/H417E are 48,200 and 25,400, respectively (Table IV above). Enhanced fibrin stimulation of the variants did not result from increased activity in the presence of fibrin

14

but rather from decreased activity in the absence of a stimulator, an observation consistent with the belief that the effects of these mutations are mediated by disruption of a salt bridge between Lys 429 and Asp 477 in single chain t-PA.

The single chain form of a zymogen-like variant of t-PA is expected to exhibit reduced activity not only towards substrates (Tables I and III, above) but also towards specific inhibitors. To demonstrate this, we measured the second order rate constant for inhibition of single chain t-PA/R275E, t-PA/R275E,H417D, and t-PA/R275E,H417E by the serpin plasminogen activator inhibitor, type 1 (PAI-1) (Table V below). As expected, both variants containing mutations at position 417 exhibited resistance to inhibition by PAI-1. The second order rate constant for inhibition by PAI-1 of t-PA/R275E,H417D or t-PA/R275E,H417E was reduced by factors of approximately 12 or 9, respectively, compared with t-PA/R275E.

TABLE V

Inhibition of wild type and variants of t-PA by PAI-1	
Enzyme	Second Order Rate Constant ($M^{-1}s^{-1}$)
t-PA/R275E	1.8×10^6
t-PA/R275E, H417D	1.5×10^5
t-PA/R275E, H417E	2.1×10^5

t-PA exhibits unusually high catalytic activity as a single chain enzyme and consequently very low zymogenicity. By contrast, a closely related enzyme urokinase (u-PA) exhibits much lower catalytic activity as a single chain enzyme and substantially higher zymogenicity. An important finding of this study is that the presence or absence of a favorable electrostatic interaction between residues at positions 417 and 429 appears to be the major determinant of this key functional distinction between the two human plasminogen activators. The zymogenicity of wild type t-PA, u-PA, and t-PA containing an aspartate at position 417 are approximately 9,250, and 150, respectively.

These studies demonstrated structure/function relationships within the activation domain of t-PA, and elucidated the molecular basis of important functional distinctions between t-PA and u-PA. These results can also aid the design of improved thrombolytic agents. For example t-PA/R275E, H417D, exhibits substantially enhanced fibrin stimulation, resistance to inhibition by PAI-1, and significantly increased zymogenicity, a useful combination of properties that enhances the therapeutic utility of the enzyme.

EXAMPLE 2

Site Directed Mutagenesis and Construction of Expression Vectors Encoding Variants of T-PA

Oligonucleotide directed site specific mutagenesis was performed as described in Example 1. The K429Y mutation was introduced into the 290 bp SacI-SmaI fragment of cDNA encoding t-PA that had been previously subcloned into bacteriophage M13 mp18. The mutagenic primer had the following nucleotide sequence:

5'-CGGAGCGGCTGTATGAGGCTCATGT-3' (SEQ ID NO: 10).

Following mutagenesis, ssDNA corresponding to the entire 290 bp SacI-SmaI fragment was fully sequenced to assure the presence of the desired mutation and the absence of any additional mutations. The sequence corresponding to the 290 bp SacI-SmaI fragment of the K429Y mutation is

15

shown in SEQ ID NO: 7. Replicative form (RF) DNA was prepared for appropriate phage, and the mutated 290 bp SacI-SmaI fragment was recovered after digestion of RF DNA with SacI and SmaI and electrophoresis of the digestion products on an agarose gel. The isolated, mutated SacI-SmaI fragment was used to replace the corresponding fragment in full length cDNAs encoding wild type t-PA or t-PA/R275E to yield new, full length cDNAs encoding t-PA/K429Y and t-PA/R275E,K429Y.

Expression of Enzymes by Transient Transfection of COS Cells.

cDNAs encoding t-PA, t-PA/R275E, t-PA/K429Y, and t-PA/R275E,K429Y were ligated into the transient expression vector pSVT7 and then introduced into COS cells by electroporation using a Bio Rad Gene pulser as described in Example 1. Following electroporation, cells were incubated overnight at 37 degrees Celsius in DMEM containing 10% fetal calf serum and 5 mM sodium butyrate. Cells were then washed with serum free medium and incubated in DMEM for 48 hours at 37 degrees Celsius. After the incubation with serum free media, conditioned media were collected and enzyme concentrations were determined by ELISA.

Purification of Wild Type and Mutated Variants of t-PA.

Wild type and mutated variants of t-PA were purified using an FPLC system and a 1 ml HiTrap chelating column (Pharmacia Biotech). The column was charged with 0.1 M CuSO₄ solution, washed with 5–10 ml distilled water, and equilibrated with start buffer (0.02 M NaHPO₄, pH 7.2, 1 M NaCl and 1 mM Imidazole). Conditioned medium containing wild type or variants of t-PA was adjusted to 1 M NaCl and injected into the column with a 50 ml superloop (Pharmacia Biotech). The column was then washed with 10 column volumes of start buffer and eluted using a 0–0.32 M linear gradient of imidazole in the same buffer. Peak fractions were collected and pooled. Purified t-PA samples were concentrated, and buffer was exchanged to 25 mM Tris (pH=7.5), 50 mM NaCl, 1 mM EDTA, using a Centrplus 30 concentrator (Amicon).

Kinetic Analysis of t-PA Activity Using a Small, Synthetic Substrate.

The direct chromogenic assay utilized the substrate methylsulfonyl-D-cyclohexyltyrosyl-glycyl-arginine-p-nitroaniline (Spectrozyme t-PA, American Diagnostica) and was performed as described in Example 1.

Kinetic Analysis of Plasminogen Activation Using Indirect Chromogenic Assays.

Indirect chromogenic assays of t-PA utilized the substrates lys-plasminogen (American Diagnostica) and Spectrozyme PL (American Diagnostica) and were performed as previously described in Example 1. Assays were performed both in the presence and absence of the co-factor DESAFIB (American Diagnostica).

Indirect Chromogenic Assays in the Presence of Various Fibrin Co-Factors.

Standard indirect chromogenic assays were performed. Briefly, 0.25–1 ng of enzyme, 0.2 μ M lys-plasminogen and 0.62 mM Spectrozyme PL were present in a total volume of 100 μ l. Assays were performed either in the presence of buffer, 25 μ g/ml DESAFIB, 100 μ g/ml fibrinogen, 100 μ g/ml cyanogen bromide fragments of fibrinogen (American Diagnostica), or 100 μ g/ml of the stimulatory, thirteen amino acid peptide P368. P368 was kindly provided by Marshall Runge (University of Texas Medical Center, Galveston, Tex.). Assays were performed in microtiter plates, and the optical density at 405 nm was measured every 30 seconds for one hour in a Molecular Devices Thermomax. Reactions were performed at 37 degrees Celsius.

16

Measurement of Second Order Rate Constants for Inhibition by PAI-1.

Second order rate constants for the inhibition of wild type and mutated t-PA were measured under pseudo-first order conditions as described in Example 1.

Oligonucleotide directed site specific mutagenesis was used to produce a mutation of Lys 429 of t-PA that selectively suppressed the catalytic activity of single chain t-PA. Lysine 429 was replaced by a tyrosine residue to yield t-PA/K429Y. In addition, to permit accurate measurement of the enzymatic activity toward plasminogen of the single chain form of this variant, a noncleavable form of the mutated enzyme was constructed by introducing the additional mutation R275E into the existing mutant to yield the R275E,K429Y variant.

Wild type t-PA, t-PA/R275E, t-PA/K429Y, and t-PA/R275E,K429Y were expressed by transient expression in COS 1 cells as described in Example 1. Since this procedure yielded predominantly single chain enzymes, two-chain t-PAs were generated by treating the enzyme preparations with plasmin-Sepharose. Quantitative conversion of the enzymes into their mature, two-chain form was confirmed by SDS-PAGE. As previously demonstrated, variants containing the mutation R275E were synthesized and secreted exclusively as single chain enzymes and were not cleaved by plasmin-Sepharose.

TABLE VI

Kinetic constants for cleavage of the chromogenic substrate
Spectrozyme t-PA by single- and two-chain t-PA variants

Enzyme	$K_{cat}(s^{-1})$	$K_m(mM)$	$K_{cat}/K_m(M^{-1}s^{-1})$
<u>Two-chain form</u>			
t-PA	40	0.5	8.0×10^4
t-PA/K429Y	35	0.5	7.0×10^4
<u>Single-chain form</u>			
t-PA/R275E	24	0.7	3.4×10^4
t-PA/R275E, K429Y	0.3	0.5	6.0×10^2

The enzymatic activity of the single and two-chain forms of wild type and t-PAs toward a small synthetic substrate is listed in Table VI above. Mutation of lysine 429 had little effect on the activity of two-chain t-PA. Two-chain t-PA/K429Y displayed approximately 90% of the activity of the two-chain, wild type enzyme in this assay. By contrast, the K429Y mutation had a very substantial effect on the activity of single chain t-PA. Single chain t-PA/R275E,K429Y exhibited approximately 2% of the activity of single chain t-PA/R275E. In this assay, the zymogenicity, defined as the ratio of the activities of the two-chain to that of the single chain form of a particular enzyme, was approximately 2.5 for wild type t-PA. By contrast, for variants containing the K429Y mutation, this ratio increased to approximately 117 (Table VI).

In the absence of a co-factor, the K429Y mutation had little effect on the activity of two-chain t-PA toward plasminogen; however, this mutation significantly reduced the catalytic efficiency of single chain t-PA (Table VII below). Compared with that of single chain t-PA/R275E, the activity of single chain t-PA/R275E,K429Y was reduced by a factor

of 17. By contrast, the activities of two-chain t-PA and t-PA/K429Y differed by a factor of only 1.2.

TABLE VII

Kinetic constants for activation of plasminogen by single- and two-chain variants of t-PA in the absence of a cofactor			
Enzyme	$K_{cat}(s^{-1})$	$K_m(\mu M)$	$K_{cat}/K_m(M^{-1}s^{-1})$
<u>Two-chain form</u>			
t-PA	0.16	10	1.6×10^4
t-PA/K429Y	0.18	14	1.3×10^4
<u>Single-chain form</u>			
t-PA/R275E	[0.038]	[30]	1.3×10^3
t-PA/R275E, K429Y	0.00046	5.9	7.8×10^1

All of the variants analyzed in this study maintained reasonably high enzymatic activity towards the natural substrate plasminogen in the presence of the co-factor fibrin (Table VIII below). The single chain form of variants containing the K429Y mutation were, however, affected to a slightly greater extent than the corresponding mature, two-chain enzymes. Two-chain t-PA/K429Y possessed approximately 75% of the activity of two-chain t-PA while single chain t-PA/R275E,K429Y exhibited approximately 40% of the activity of single chain t-PA/R275E.

TABLE VIII

Kinetic constants for activation of plasminogen by single- and two-chain t-PA variants in the presence of fibrin			
Enzyme	$K_{cat}(s^{-1})$	$K_m(\mu M)$	$K_{cat}/K_m(M^{-1}s^{-1})$
<u>Two-chain form</u>			
t-PA	0.08	0.02	4.0×10^6
t-PA/K429Y	0.08	0.03	3.0×10^6
<u>Single-chain form</u>			
t-PA/R275E	0.10	0.02	5.0×10^6
t-PA/R275E, K429Y	0.10	0.07	2.0×10^6

The extent of fibrin stimulation displayed by the single chain form of t-PA/R275E,K429Y is significantly greater than that displayed by wild type t-PA. Wild type, two-chain t-PA possesses a fibrin stimulation factor, defined as the ratio of the catalytic efficiencies in the presence and absence of fibrin, of approximately 250 (Table IX below). The two-chain t-PA/K429Y variant displays a similar stimulation factor of 230. Single chain wild type t-PA, with a fibrin stimulation factor of 3800, is stimulated to a substantially greater degree than the two-chain enzymes, presumable reflecting the ability of fibrin to stimulate the single chain enzymes not only through a templating mechanism but also by inducing nonproteolytic zymogen activation. Stimulation of single chain t-PA is further enhanced by the K429Y mutation. The fibrin stimulation factor for single chain t-PA/R275E,K429Y is approximately 26,000. Enhanced fibrin stimulation of the variant did not result from increased activity in the presence of fibrin but rather from decreased activity in the absence of a stimulator, an observation consistent with our proposal that the effects of these muta-

tions are mediated by disruption of a salt bridge between Lys 429 and Asp 477 in single chain t-PA.

TABLE IX

Stimulatory effect of fibrin on the catalytic efficiencies for variants of t-PA	
Enzyme	Fold stimulation of k_{cat}/K_m
<u>Two-chain form</u>	
t-PA	250
t-PA/K429Y	230
<u>Single-chain form</u>	
t-PA/R275E	3800
t-PA/R275E, K429Y	26,000

The mutated enzyme t-PA/R275E,K429Y is not only stimulated to a significantly greater extent by soluble fibrin than t-PA (Table IX above), but it is also substantially more discriminating among fibrin co-factors than the wild type enzyme (FIG. 2). The two-chain form of both wild type t-PA and t-PA/K429Y are strongly stimulated by soluble fibrin monomers (DESAFIB), fibrinogen, CNBr fragments of fibrinogen, and a 13 amino acid peptide (P368). Single chain t-PA/R275E, on the other hand, is stimulated strongly by soluble fibrin and fibrinogen and moderately by the CNBr fragments and peptide P368. In striking contrast to these enzymes, single chain t-PA/R275E,K429Y, although dramatically stimulated by fibrin monomers, is virtually non-responsive to fibrinogen, CNBr fragments of fibrinogen, peptide P368.

The ratio of the specific activity of a plasminogen activator in the presence of fibrin to that in the presence of fibrinogen, or "fibrin selectivity factor", is one indication of the "clot selectivity" an enzyme will demonstrate in vivo. An enzyme with enhanced fibrin selectivity can accomplish efficient thrombolysis while displaying decreased systemic activity. Under the conditions of the assays reported here, the fibrin selectivity is 1.5 for two-chain t-PA, 1.5 for two-chain t-PA/K429Y, and 1.0 for single chain t-PA/R275E. The fibrin selectivity factor for single chain t-PA/R275E,K429Y, however, is 146. This double mutant, therefore, is approximately two orders of magnitude more discriminating between fibrin and fibrinogen than either single or two-chain wild type t-PA.

The single chain form of a zymogen-like variant of t-PA is expected to exhibit reduced activity not only towards substrates (Tables VI and VIII above) but also towards specific inhibitors. The second order rate constant for inhibition of the single chain form of both t-PA/R275E and t-PA/R275E,K429Y by the serpin plasminogen activator inhibitor, type 1 (PAI-1), the primary physiological inhibitor of t-PA is shown in Table X below. As expected, t-PA/R275E,K429Y exhibited resistance to inhibition by PAI-1. The second order compared with t-PA/R275E.

TABLE X

Inhibition of wild type and variants of t-PA by PAI-1	
Enzyme	Second order rate constant ($M^{-1}s^{-1}$)
t-PA/R275E	1.8×10^6
t-PA/R275E,K429Y	7.7×10^3

An important finding of this study is that conversion of lysine 429 to tyrosine residue selectively suppresses the activity of single chain t-PA and thereby substantially

19

enhances the zymogenicity of the enzyme. We have demonstrated, in addition, that single chain t-PA/R275E, K429Y is significantly more fibrin stimulated and substantially more fibrin selective than either single or two-chain, wild type t-PA. Single chain t-PA/R275E,K429Y also exhibits marked resistance to inhibition by PAI-1. It is believed that the effects of this mutation are mediated by disruption of a critical salt bridge formed by Lys 429 and Asp 477 that has been predicted to be present in single- but not two-chain t-PA. The primary role of this putative salt bridge is believed to be stabilization of the active conformation of single chain t-PA. Two-chain t-PA/K429Y, therefore, as demonstrated in this study, is expected to maintain high enzymatic activity.

20

These results aid in the design of improved thrombolytic agents. For Example t-PA/R275E,K429Y, exhibits significantly enhanced fibrin stimulation, dramatically increased discrimination among fibrin co-factors, marked resistance to inhibition by PAI-1, and substantially increased zymogenicity, a combination of properties that enhance the therapeutic utility of the enzyme.

The foregoing is intended to be illustrative of the present invention, but not limiting. Numerous variations and modifications of the present invention may be effected without departing from the true spirit and scope of the invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 12

<210> SEQ ID NO 1

<211> LENGTH: 527

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Ser Tyr Gln Val Ile Cys Arg Asp Glu Lys Thr Gln Met Ile Tyr Gln
 1           5           10           15

Gln His Gln Ser Trp Leu Arg Pro Val Leu Arg Ser Asn Arg Val Glu
 20           25           30

Tyr Cys Trp Cys Asn Ser Gly Arg Ala Gln Cys His Ser Val Pro Val
 35           40           45

Lys Ser Cys Ser Glu Pro Arg Cys Phe Asn Gly Gly Thr Cys Gln Gln
 50           55           60

Ala Leu Tyr Phe Ser Asp Phe Val Cys Gln Cys Pro Glu Gly Phe Ala
 65           70           75           80

Gly Lys Cys Cys Glu Ile Asp Thr Arg Ala Thr Cys Tyr Glu Asp Gln
 85           90           95

Gly Ile Ser Tyr Arg Gly Thr Trp Ser Thr Ala Glu Ser Gly Ala Glu
100           105           110

Cys Thr Asn Trp Asn Ser Ser Ala Leu Ala Gln Lys Pro Tyr Ser Gly
115           120           125

Arg Arg Pro Asp Ala Ile Arg Leu Gly Leu Gly Asn His Asn Tyr Cys
130           135           140

Arg Asn Pro Asp Arg Asp Ser Lys Pro Trp Cys Tyr Val Phe Lys Ala
145           150           155           160

Gly Lys Tyr Ser Ser Glu Phe Cys Ser Thr Pro Ala Cys Ser Glu Gly
165           170           175

Asn Ser Asp Cys Tyr Phe Gly Asn Gly Ser Ala Tyr Arg Gly Thr His
180           185           190

Ser Leu Thr Glu Ser Gly Ala Ser Cys Leu Pro Trp Asn Ser Met Ile
195           200           205

Leu Ile Gly Lys Val Tyr Thr Ala Gln Asn Pro Ser Ala Gln Ala Leu
210           215           220

Gly Leu Gly Lys His Asn Tyr Cys Arg Asn Pro Asp Gly Asp Ala Lys
225           230           235           240

Pro Trp Cys His Val Leu Lys Asn Arg Arg Leu Thr Trp Glu Tyr Cys
245           250           255

Asp Val Pro Ser Cys Ser Thr Cys Gly Leu Arg Gln Tyr Ser Gln Pro
260           265           270

```

-continued

Gln Phe Glu Ile Lys Gly Gly Leu Phe Ala Asp Ile Ala Ser His Pro
 275 280 285
 Trp Gln Ala Ala Ile Phe Ala Lys His Arg Arg Ser Pro Gly Glu Arg
 290 295 300
 Phe Leu Cys Gly Gly Ile Leu Ile Ser Ser Cys Trp Ile Leu Ser Ala
 305 310 315 320
 Ala His Cys Phe Gln Glu Arg Phe Pro Pro His His Leu Thr Val Ile
 325 330 335
 Leu Gly Arg Thr Tyr Arg Val Val Pro Gly Glu Glu Glu Gln Lys Phe
 340 345 350
 Glu Val Glu Lys Tyr Ile Val His Lys Glu Phe Asp Asp Asp Thr Tyr
 355 360 365
 Asp Asn Asp Ile Ala Leu Leu Gln Leu Lys Ser Asp Ser Ser Arg Cys
 370 375 380
 Ala Gln Glu Ser Ser Val Val Arg Thr Val Cys Leu Pro Pro Ala Asp
 385 390 395 400
 Leu Gln Leu Pro Asp Trp Thr Glu Cys Glu Leu Ser Gly Tyr Gly Lys
 405 410 415
 Asp Glu Ala Leu Ser Pro Phe Tyr Ser Glu Arg Leu Lys Glu Ala His
 420 425 430
 Val Arg Leu Tyr Pro Ser Ser Arg Cys Thr Ser Gln His Leu Leu Asn
 435 440 445
 Arg Thr Val Thr Asp Asn Met Leu Cys Ala Gly Asp Thr Arg Ser Gly
 450 455 460
 Gly Pro Gln Ala Asn Leu His Asp Ala Cys Gln Gly Asp Ser Gly Gly
 465 470 475 480
 Pro Leu Val Cys Leu Asn Asp Gly Arg Met Thr Leu Val Gly Ile Ile
 485 490 495
 Ser Trp Gly Leu Gly Cys Gly Gln Lys Asp Val Pro Gly Val Tyr Thr
 500 505 510
 Lys Val Thr Asn Tyr Leu Asp Trp Ile Arg Asp Asn Met Arg Pro
 515 520 525

<210> SEQ ID NO 2

<211> LENGTH: 527

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Ser Tyr Gln Val Ile Cys Arg Asp Glu Lys Thr Gln Met Ile Tyr Gln
 1 5 10 15
 Gln His Gln Ser Trp Leu Arg Pro Val Leu Arg Ser Asn Arg Val Glu
 20 25 30
 Tyr Cys Trp Cys Asn Ser Gly Arg Ala Gln Cys His Ser Val Pro Val
 35 40 45
 Lys Ser Cys Ser Glu Pro Arg Cys Phe Asn Gly Gly Thr Cys Gln Gln
 50 55 60
 Ala Leu Tyr Phe Ser Asp Phe Val Cys Gln Cys Pro Glu Gly Phe Ala
 65 70 75 80
 Gly Lys Cys Cys Glu Ile Asp Thr Arg Ala Thr Cys Tyr Glu Asp Gln
 85 90 95
 Gly Ile Ser Tyr Arg Gly Thr Trp Ser Thr Ala Glu Ser Gly Ala Glu
 100 105 110
 Cys Thr Asn Trp Asn Ser Ser Ala Leu Ala Gln Lys Pro Tyr Ser Gly

-continued

115	120	125
Arg Arg Pro Asp Ala Ile Arg Leu Gly Leu Gly Asn His Asn Tyr Cys 130 135 140		
Arg Asn Pro Asp Arg Asp Ser Lys Pro Trp Cys Tyr Val Phe Lys Ala 145 150 155 160		
Gly Lys Tyr Ser Ser Glu Phe Cys Ser Thr Pro Ala Cys Ser Glu Gly 165 170 175		
Asn Ser Asp Cys Tyr Phe Gly Asn Gly Ser Ala Tyr Arg Gly Thr His 180 185 190		
Ser Leu Thr Glu Ser Gly Ala Ser Cys Leu Pro Trp Asn Ser Met Ile 195 200 205		
Leu Ile Gly Lys Val Tyr Thr Ala Gln Asn Pro Ser Ala Gln Ala Leu 210 215 220		
Gly Leu Gly Lys His Asn Tyr Cys Arg Asn Pro Asp Gly Asp Ala Lys 225 230 235 240		
Pro Trp Cys His Val Leu Lys Asn Arg Arg Leu Thr Trp Glu Tyr Cys 245 250 255		
Asp Val Pro Ser Cys Ser Thr Cys Gly Leu Arg Gln Tyr Ser Gln Pro 260 265 270		
Gln Phe Glu Ile Lys Gly Gly Leu Phe Ala Asp Ile Ala Ser His Pro 275 280 285		
Trp Gln Ala Ala Ile Phe Ala Lys His Arg Arg Ser Pro Gly Glu Arg 290 295 300		
Phe Leu Cys Gly Gly Ile Leu Ile Ser Ser Cys Trp Ile Leu Ser Ala 305 310 315 320		
Ala His Cys Phe Gln Glu Arg Phe Pro Pro His His Leu Thr Val Ile 325 330 335		
Leu Gly Arg Thr Tyr Arg Val Val Pro Gly Glu Glu Glu Gln Lys Phe 340 345 350		
Glu Val Glu Lys Tyr Ile Val His Lys Glu Phe Asp Asp Asp Thr Tyr 355 360 365		
Asp Asn Asp Ile Ala Leu Leu Gln Leu Lys Ser Asp Ser Ser Arg Cys 370 375 380		
Ala Gln Glu Ser Ser Val Val Arg Thr Val Cys Leu Pro Pro Ala Asp 385 390 395 400		
Leu Gln Leu Pro Asp Trp Thr Glu Cys Glu Leu Ser Gly Tyr Gly Lys 405 410 415		
Glu Glu Ala Leu Ser Pro Phe Tyr Ser Glu Arg Leu Lys Glu Ala His 420 425 430		
Val Arg Leu Tyr Pro Ser Ser Arg Cys Thr Ser Gln His Leu Leu Asn 435 440 445		
Arg Thr Val Thr Asp Asn Met Leu Cys Ala Gly Asp Thr Arg Ser Gly 450 455 460		
Gly Pro Gln Ala Asn Leu His Asp Ala Cys Gln Gly Asp Ser Gly Gly 465 470 475 480		
Pro Leu Val Cys Leu Asn Asp Gly Arg Met Thr Leu Val Gly Ile Ile 485 490 495		
Ser Trp Gly Leu Gly Cys Gly Gln Lys Asp Val Pro Gly Val Tyr Thr 500 505 510		
Lys Val Thr Asn Tyr Leu Asp Trp Ile Arg Asp Asn Met Arg Pro 515 520 525		

-continued

```

<211> LENGTH: 527
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Ser Tyr Gln Val Ile Cys Arg Asp Glu Lys Thr Gln Met Ile Tyr Gln
 1             5             10             15

Gln His Gln Ser Trp Leu Arg Pro Val Leu Arg Ser Asn Arg Val Glu
          20             25             30

Tyr Cys Trp Cys Asn Ser Gly Arg Ala Gln Cys His Ser Val Pro Val
          35             40             45

Lys Ser Cys Ser Glu Pro Arg Cys Phe Asn Gly Gly Thr Cys Gln Gln
          50             55             60

Ala Leu Tyr Phe Ser Asp Phe Val Cys Gln Cys Pro Glu Gly Phe Ala
          65             70             75             80

Gly Lys Cys Cys Glu Ile Asp Thr Arg Ala Thr Cys Tyr Glu Asp Gln
          85             90             95

Gly Ile Ser Tyr Arg Gly Thr Trp Ser Thr Ala Glu Ser Gly Ala Glu
          100            105            110

Cys Thr Asn Trp Asn Ser Ser Ala Leu Ala Gln Lys Pro Tyr Ser Gly
          115            120            125

Arg Arg Pro Asp Ala Ile Arg Leu Gly Leu Gly Asn His Asn Tyr Cys
          130            135            140

Arg Asn Pro Asp Arg Asp Ser Lys Pro Trp Cys Tyr Val Phe Lys Ala
          145            150            155            160

Gly Lys Tyr Ser Ser Glu Phe Cys Ser Thr Pro Ala Cys Ser Glu Gly
          165            170            175

Asn Ser Asp Cys Tyr Phe Gly Asn Gly Ser Ala Tyr Arg Gly Thr His
          180            185            190

Ser Leu Thr Glu Ser Gly Ala Ser Cys Leu Pro Trp Asn Ser Met Ile
          195            200            205

Leu Ile Gly Lys Val Tyr Thr Ala Gln Asn Pro Ser Ala Gln Ala Leu
          210            215            220

Gly Leu Gly Lys His Asn Tyr Cys Arg Asn Pro Asp Gly Asp Ala Lys
          225            230            235            240

Pro Trp Cys His Val Leu Lys Asn Arg Arg Leu Thr Trp Glu Tyr Cys
          245            250            255

Asp Val Pro Ser Cys Ser Thr Cys Gly Leu Arg Gln Tyr Ser Gln Pro
          260            265            270

Gln Phe Glu Ile Lys Gly Gly Leu Phe Ala Asp Ile Ala Ser His Pro
          275            280            285

Trp Gln Ala Ala Ile Phe Ala Lys His Arg Arg Ser Pro Gly Glu Arg
          290            295            300

Phe Leu Cys Gly Gly Ile Leu Ile Ser Ser Cys Trp Ile Leu Ser Ala
          305            310            315            320

Ala His Cys Phe Gln Glu Arg Phe Pro Pro His His Leu Thr Val Ile
          325            330            335

Leu Gly Arg Thr Tyr Arg Val Val Pro Gly Glu Glu Glu Gln Lys Phe
          340            345            350

Glu Val Glu Lys Tyr Ile Val His Lys Glu Phe Asp Asp Asp Thr Tyr
          355            360            365

Asp Asn Asp Ile Ala Leu Leu Gln Leu Lys Ser Asp Ser Ser Arg Cys
          370            375            380

Ala Gln Glu Ser Ser Val Val Arg Thr Val Cys Leu Pro Pro Ala Asp

```

-continued

385	390	395	400
Leu Gln Leu Pro Asp	Trp Thr Glu Cys	Glu Leu Ser Gly Tyr	Gly Lys
405		410	415
His Glu Ala Leu Ser Pro Phe Tyr	Ser Glu Arg Leu Tyr	Glu Ala His	
420	425	430	
Val Arg Leu Tyr Pro Ser Ser Arg Cys Thr Ser Gln His Leu Leu Asn			
435	440	445	
Arg Thr Val Thr Asp Asn Met Leu Cys Ala Gly Asp Thr Arg Ser Gly			
450	455	460	
Gly Pro Gln Ala Asn Leu His Asp Ala Cys Gln Gly Asp Ser Gly Gly			
465	470	475	480
Pro Leu Val Cys Leu Asn Asp Gly Arg Met Thr Leu Val Gly Ile Ile			
485	490	495	
Ser Trp Gly Leu Gly Cys Gly Gln Lys Asp Val Pro Gly Val Tyr Thr			
500	505	510	
Lys Val Thr Asn Tyr Leu Asp Trp Ile Arg Asp Asn Met Arg Pro			
515	520	525	

<210> SEQ ID NO 4
 <211> LENGTH: 290
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

ctacggcaag catgaggcct tgtctccttt ctattcggag cggctgaagg aggctcatgt	60
cagactgtac ccatccagcc gctgcacatc acaacattta cttaacagaa cagtcaccga	120
caacatgctg tgtgctggag aactcggag cggcggggccc caggcaaact tgcacgacgc	180
ctgccagggc gattcgggag gcccctggt gtgtctgaac gatggccgca tgactttggt	240
gggcatcatc agctggggcc tgggctgtgg acagaaggat gtcccgggtg	290

<210> SEQ ID NO 5
 <211> LENGTH: 290
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

ctacggcaag gacgaggcct tgtctccttt ctattcggag cggctgaagg aggctcatgt	60
cagactgtac ccatccagcc gctgcacatc acaacattta cttaacagaa cagtcaccga	120
caacatgctg tgtgctggag aactcggag cggcggggccc caggcaaact tgcacgacgc	180
ctgccagggc gattcgggag gcccctggt gtgtctgaac gatggccgca tgactttggt	240
gggcatcatc agctggggcc tgggctgtgg acagaaggat gtcccgggtg	290

<210> SEQ ID NO 6
 <211> LENGTH: 290
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

ctacggcaag gaggaggcct tgtctccttt ctattcggag cggctgaagg aggctcatgt	60
cagactgtac ccatccagcc gctgcacatc acaacattta cttaacagaa cagtcaccga	120
caacatgctg tgtgctggag aactcggag cggcggggccc caggcaaact tgcacgacgc	180
ctgccagggc gattcgggag gcccctggt gtgtctgaac gatggccgca tgactttggt	240
gggcatcatc agctggggcc tgggctgtgg acagaaggat gtcccgggtg	290

-continued

<210> SEQ ID NO 7
 <211> LENGTH: 290
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```
ctacggcaag catgaggcct tgtctccttt ctattcggag cggctgtatg aggetcatgt      60
cagactgtac ccatccagcc gctgcacatc acaacattta cttaacagaa cagtcaccga    120
caacatgctg tgtgtcggag acactcggag cggcggggccc caggcaaact tgcacgacgc    180
ctgccagggc gattcgggag gccccctggg gtgtctgaac gatggccgca tgactttggg    240
gggcatcatc agctggggcc tgggctgtgg acagaaggat gtcccgggtg                290
```

<210> SEQ ID NO 8
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```
ctacggcaag gacgaggcct tgt                                             23
```

<210> SEQ ID NO 9
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```
ctacggcaag gaggaggcct tgt                                             23
```

<210> SEQ ID NO 10
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

```
cggagcggct gtatgaggct mcatgt                                         26
```

<210> SEQ ID NO 11
 <211> LENGTH: 2544
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (86)...(1774)
 <220> FEATURE:
 <221> NAME/KEY: sig_peptide
 <222> LOCATION: (86)...(190)

<400> SEQUENCE: 11

```
gttctgagca cagggtgga gagaaaacct ctgcgaggaa agggaaggag caagccgtga      60
atttaaggga cgctgtgaag caatc atg gat gca atg aag aga ggg ctc tgc      112
                                     Met Asp Ala Met Lys Arg Gly Leu Cys
                                     -35                      -30
tgt gtg ctg ctg ctg tgt gga gca gtc ttc gtt tcg ccc agc cag gaa      160
Cys Val Leu Leu Leu Cys Gly Ala Val Phe Val Ser Pro Ser Gln Glu
-25                      -20                      -15
atc cat gcc cga ttc aga aga gga gcc aga tct tac caa gtg atc tgc      208
Ile His Ala Arg Phe Arg Arg Gly Ala Arg Ser Tyr Gln Val Ile Cys
-10                      -5                      1                      5
```

-continued

aga gat gaa aaa acg cag atg ata tac cag caa cat cag tca tgg ctg	256
Arg Asp Glu Lys Thr Gln Met Ile Tyr Gln Gln His Gln Ser Trp Leu	
10 15 20	
cgc cct gtg ctc aga agc aac cgg gtg gaa tat tgc tgg tgc aac agt	304
Arg Pro Val Leu Arg Ser Asn Arg Val Glu Tyr Cys Trp Cys Asn Ser	
25 30 35	
ggc agg gca cag tgc cac tca gtg cct gtc aaa agt tgc agc gag cca	352
Gly Arg Ala Gln Cys His Ser Val Pro Val Lys Ser Cys Ser Glu Pro	
40 45 50	
agg tgt ttc aac ggg ggc acc tgc cag cag gcc ctg tac ttc tca gat	400
Arg Cys Phe Asn Gly Gly Thr Cys Gln Gln Ala Leu Tyr Phe Ser Asp	
55 60 65 70	
ttc gtg tgc cag tgc ccc gaa gga ttt gct ggg aag tgc tgt gaa ata	448
Phe Val Cys Gln Cys Pro Glu Gly Phe Ala Gly Lys Cys Cys Glu Ile	
75 80 85	
gat acc agg gcc acg tgc tac gag gac cag ggc atc agc tac agg ggc	496
Asp Thr Arg Ala Thr Cys Tyr Glu Asp Gln Gly Ile Ser Tyr Arg Gly	
90 95 100	
acg tgg agc aca gcg gag agt ggc gcc gag tgc acc aac tgg aac agc	544
Thr Trp Ser Thr Ala Glu Ser Gly Ala Glu Cys Thr Asn Trp Asn Ser	
105 110 115	
agc gcg ttg gcc cag aag ccc tac agc ggg cgg agg cca gac gcc atc	592
Ser Ala Leu Ala Gln Lys Pro Tyr Ser Gly Arg Arg Pro Asp Ala Ile	
120 125 130	
agg ctg ggc ctg ggg aac cac aac tac tgc aga aac cca gat cga gac	640
Arg Leu Gly Leu Gly Asn His Asn Tyr Cys Arg Asn Pro Asp Arg Asp	
135 140 145 150	
tca aag ccc tgg tgc tac gtc ttt aag gcg ggg aag tac agc tca gag	688
Ser Lys Pro Trp Cys Tyr Val Phe Lys Ala Gly Lys Tyr Ser Ser Glu	
155 160 165	
ttc tgc agc acc cct gcc tgc tct gag gga aac agt gac tgc tac ttt	736
Phe Cys Ser Thr Pro Ala Cys Ser Glu Gly Asn Ser Asp Cys Tyr Phe	
170 175 180	
ggg aat ggg tca gcc tac cgt ggc acg cac agc ctc acc gag tcg ggt	784
Gly Asn Gly Ser Ala Tyr Arg Gly Thr His Ser Leu Thr Glu Ser Gly	
185 190 195	
gcc tcc tgc ctc ccg tgg aat tcc atg atc ctg ata ggc aag gtt tac	832
Ala Ser Cys Leu Pro Trp Asn Ser Met Ile Leu Ile Gly Lys Val Tyr	
200 205 210	
aca gca cag aac ccc agt gcc cag gca ctg ggc ctg ggc aaa cat aat	880
Thr Ala Gln Asn Pro Ser Ala Gln Ala Leu Gly Leu Gly Lys His Asn	
215 220 225 230	
tac tgc cgg aat cct gat ggg gat gcc aag ccc tgg tgc cac gtg ctg	928
Tyr Cys Arg Asn Pro Asp Gly Asp Ala Lys Pro Trp Cys His Val Leu	
235 240 245	
aag aac cgc agg ctg acg tgg gag tac tgt gat gtg ccc tcc tgc tcc	976
Lys Asn Arg Arg Leu Thr Trp Glu Tyr Cys Asp Val Pro Ser Cys Ser	

-continued

250	255	260	
acc tgc ggc ctg aga cag tac agc cag cct cag ttt cgc atc aaa gga			1024
Thr Cys Gly Leu Arg Gln Tyr Ser Gln Pro Gln Phe Arg Ile Lys Gly			
265	270	275	
ggg ctc ttc gcc gac atc gcc tcc cac ccc tgg cag gct gcc atc ttt			1072
Gly Leu Phe Ala Asp Ile Ala Ser His Pro Trp Gln Ala Ala Ile Phe			
280	285	290	
gcc aag cac agg agg tcg ccc gga gag cgg ttc ctg tgc ggg ggc ata			1120
Ala Lys His Arg Arg Ser Pro Gly Glu Arg Phe Leu Cys Gly Gly Ile			
295	300	305	310
ctc atc agc tcc tgc tgg att ctc tct gcc gcc cac tgc ttc cag gag			1168
Leu Ile Ser Ser Cys Trp Ile Leu Ser Ala Ala His Cys Phe Gln Glu			
315	320	325	
agg ttt ccg ccc cac cac ctg acg gtg atc ttg ggc aga aca tac cgg			1216
Arg Phe Pro Pro His His Leu Thr Val Ile Leu Gly Arg Thr Tyr Arg			
330	335	340	
gtg gtc cct ggc gag gag gag cag aaa ttt gaa gtc gaa aaa tac att			1264
Val Val Pro Gly Glu Glu Glu Gln Lys Phe Glu Val Glu Lys Tyr Ile			
345	350	355	
gtc cat aag gaa ttc gat gat gac act tac gac aat gac att gcg ctg			1312
Val His Lys Glu Phe Asp Asp Asp Thr Tyr Asp Asn Asp Ile Ala Leu			
360	365	370	
ctg cag ctg aaa tcg gat tcg tcc cgc tgt gcc cag gag agc agc gtg			1360
Leu Gln Leu Lys Ser Asp Ser Ser Arg Cys Ala Gln Glu Ser Ser Val			
375	380	385	390
gtc cgc act gtg tgc ctt ccc ccg gcg gac ctg cag ctg ccg gac tgg			1408
Val Arg Thr Val Cys Leu Pro Pro Ala Asp Leu Gln Leu Pro Asp Trp			
395	400	405	
acg gag tgt gag ctc tcc ggc tac ggc aag cat gag gcc ttg tct cct			1456
Thr Glu Cys Glu Leu Ser Gly Tyr Gly Lys His Glu Ala Leu Ser Pro			
410	415	420	
ttc tat tcg gag cgg ctg aag gag gct cat gtc aga ctg tac cca tcc			1504
Phe Tyr Ser Glu Arg Leu Lys Glu Ala His Val Arg Leu Tyr Pro Ser			
425	430	435	
agc cgc tgc aca tca caa cat tta ctt aac aga aca gtc acc gac aac			1552
Ser Arg Cys Thr Ser Gln His Leu Leu Asn Arg Thr Val Thr Asp Asn			
440	445	450	
atg ctg tgt gct gga gac act cgg agc ggc ggg ccc cag gca aac ttg			1600
Met Leu Cys Ala Gly Asp Thr Arg Ser Gly Gly Pro Gln Ala Asn Leu			
455	460	465	470
cac gac gcc tgc cag ggc gat tcg gga ggc ccc ctg gtg tgt ctg aac			1648
His Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Leu Asn			
475	480	485	
gat ggc cgc atg act ttg gtg ggc atc atc agc tgg ggc ctg ggc tgt			1696
Asp Gly Arg Met Thr Leu Val Gly Ile Ile Ser Trp Gly Leu Gly Cys			
490	495	500	
gga cag aag gat gtc ccg ggt gtg tac acc aag gtt acc aac tac cta			1744

-continued

Gly	Gln	Lys	Asp	Val	Pro	Gly	Val	Tyr	Thr	Lys	Val	Thr	Asn	Tyr	Leu			
		505					510						515					
gac tgg att cgt gac aac atg cga ccg tga ccaggaacac ccgactcctc 1794																		
Asp	Trp	Ile	Arg	Asp	Asn	Met	Arg	Pro	*									
	520					525												
aaaagcaaat gagatcccg ctcctctctt tcagaagaca ctgcaaaggc gcagtgcctc 1854																		
tctacagact tctccagacc caccacaccg cagaagcggg acgagaccct acaggagagg 1914																		
gaagagtga ttttccaga tacttcccat ttggaagt ttcaggactt ggtctgattt 1974																		
caggatactc tgtcagatgg gaagacatga atgcacacta gcctctccag gaatgcctcc 2034																		
tccctgggca gaaagtggc atgccaccct gttttcagct aaagcccaac ctccctgacct 2094																		
gtcaccgtga gcagctttgg aaacaggacc acaaaaatga aagcatgtct caatagtaaa 2154																		
agataacaag atctttcagg aaagacggat tgcattagaa atagacagta tatttatagt 2214																		
cacaagagcc cagcagggcc tcaaagtgg ggcaggctgg ctggcccgtc atgttcctca 2274																		
aaagcaccct tgacgtcaag tctccttccc ctttccccac tccctggctc tcagaaggta 2334																		
ttccttttgt gtacagtgtg taaagtgtaa atcctttttc tttataaact ttagagtagc 2394																		
atgagagaat tgtatcattt gaacaactag gcttcagcat atttatagca atccatgtta 2454																		
gtttttactt tctgttgcca caaccctgtt ttatactgta cttaataaat tcagatatat 2514																		
ttttcacagt ttttcaaaa aaaaaaaaaa 2544																		

<210> SEQ ID NO 12

<211> LENGTH: 562

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: SIGNAL

<222> LOCATION: (1)...(35)

<400> SEQUENCE: 12

Met	Asp	Ala	Met	Lys	Arg	Gly	Leu	Cys	Cys	Val	Leu	Leu	Leu	Cys	Gly		
-35					-30					-25					-20		
Ala	Val	Phe	Val	Ser	Pro	Ser	Gln	Glu	Ile	His	Ala	Arg	Phe	Arg	Arg		
			-15				-10						-5				
Gly	Ala	Arg	Ser	Tyr	Gln	Val	Ile	Cys	Arg	Asp	Glu	Lys	Thr	Gln	Met		
			1			5						10					
Ile	Tyr	Gln	Gln	His	Gln	Ser	Trp	Leu	Arg	Pro	Val	Leu	Arg	Ser	Asn		
	15				20					25							
Arg	Val	Glu	Tyr	Cys	Trp	Cys	Asn	Ser	Gly	Arg	Ala	Gln	Cys	His	Ser		
30				35					40				45				
Val	Pro	Val	Lys	Ser	Cys	Ser	Glu	Pro	Arg	Cys	Phe	Asn	Gly	Gly	Thr		
			50				55						60				
Cys	Gln	Gln	Ala	Leu	Tyr	Phe	Ser	Asp	Phe	Val	Cys	Gln	Cys	Pro	Glu		
			65			70					75						
Gly	Phe	Ala	Gly	Lys	Cys	Cys	Glu	Ile	Asp	Thr	Arg	Ala	Thr	Cys	Tyr		
	80				85					90							
Glu	Asp	Gln	Gly	Ile	Ser	Tyr	Arg	Gly	Thr	Trp	Ser	Thr	Ala	Glu	Ser		
	95				100				105								
Gly	Ala	Glu	Cys	Thr	Asn	Trp	Asn	Ser	Ser	Ala	Leu	Ala	Gln	Lys	Pro		
110				115					120					125			
Tyr	Ser	Gly	Arg	Arg	Pro	Asp	Ala	Ile	Arg	Leu	Gly	Leu	Gly	Asn	His		
			130				135							140			
Asn	Tyr	Cys	Arg	Asn	Pro	Asp	Arg	Asp	Ser	Lys	Pro	Trp	Cys	Tyr	Val		

[illegible]

39

I claim:

1. A polynucleotide encoding a variant of human tissue-type plasminogen activator (t-PA) protein; wherein said human t-PA protein comprises residues 1–527 of SEQ ID NO:12; and wherein said variant comprises a single chain human t-PA protein having an R275 and at least one other basic amino acid residue in a serine protease region substituted by a non-basic amino acid residue thereby disrupting a salt bridge interaction between an aspartate 477 and a lysine 429.

2. An expression vector comprising the polynucleotide of claim 1.

3. A cell comprising the expression vector of claim 2.

4. A polynucleotide encoding a variant of human tissue-type plasminogen activator t-PA) protein; wherein said

40

human t-PA protein comprises residues 1–527 of SEQ ID NO:12; and wherein said variant comprises a single ham human t-PA protein selected from the group consisting of R275E,H417D, R275E,H417E and R275E K429Y.

5. An expression vector comprising the polynucleotide of claim 4.

6. A cell comprising the expression vector of claim 5.

7. A method of making variant single chain human tissue-type plasminogen activator protein comprising the step of culturing the cell of claim 6.

8. The method of claim 7 further comprising the additional step of purifying the protein.

* * * * *