



US 20060218673A9

(19) **United States**
(12) **Patent Application Publication**
Aukerman

(10) **Pub. No.: US 2006/0218673 A9**
(48) **Pub. Date: Sep. 28, 2006**
CORRECTED PUBLICATION

(54) **GENE SILENCING**

(75) Inventor: **Milo J. Aukerman**, Newark, DE (US)

Correspondence Address:

PIONEER HI-BRED INTERNATIONAL, INC.
7250 N.W. 62ND AVENUE
P.O. BOX 552
JOHNSTON, IA 50131-0552 (US)

[0195], [0196], [0198], [0199], [0200], [0202],
[0206], [0208], [0210], [0211], [0212], Claims 1, 9,
17, 18, 19, 20, 21, 22, 23, 24, 25, 52, 54, 56, 62, 65
and 68.

(65) US 2005/0138689 A1 Jun. 23, 2005

Related U.S. Application Data

(60) Provisional application No. 60/509,958, filed on Oct.
9, 2003.

(73) Assignee: **E.I. du Pont de Nemours and Company**

(21) Appl. No.: **10/963,394**

(22) Filed: **Oct. 12, 2004**

Publication Classification

(51) **Int. Cl.**

A01H 1/00 (2006.01)

C12N 15/82 (2006.01)

C07H 21/04 (2006.01)

A01H 5/00 (2006.01)

C12N 5/04 (2006.01)

(52) **U.S. Cl.** **800/286**; 800/320.1; 435/468;
435/412; 435/419; 536/23.6

(57)

ABSTRACT

The invention provides methods and compositions useful in
target sequence suppression and target sequence validation.
The invention provides polynucleotide constructs useful for
gene silencing, as well as cells, plants and seeds comprising
the polynucleotides. The invention also provides a method
for using microRNA to silence a target sequence.

Prior Publication Data

(15) Correction of US 2005/0138689 A1 Jun. 23, 2005
See Paragraphs [0006], [0007], [0013], [0016],
[0017], [0018], [0020], [0030], [0031], [0045],
[0046], [0047], [0054], [0057], [0064], [0067],
[0068], [0069], [0070], [0071], [0072], [0074],
[0075], [0077], [0078], [0080], [0081], [0082],
[0083], [0084], [0085], [0086], [0087], [0088],
[0091], [0092], [0094], [0098], [0099], [0100],
[0101], [0102], [0103], [0106], [0108], [0132],
[0135], [0139], [0141], [0143], [0144], [0145],
[0148], [0152], [0153], [0154], [0155], [0156],
[0157], [0187], [0189], [0190], [0192], [0194],

A
A U
A U
UA
CG
CG
CG
AU
U
AU
AU
AU
UG
A
G A
U U
UA
AU
A G
A U
G A
GU
UA
AU
CG
A
C G
UA
UA
AU
CG
AU
UG
UA
UA
CG
C
AU
CG
CG
AU
CG
GC
C
G
GC
A
UA
GC
UA
UA
U U
GC
UG
UA
GC
CG
UG
GC

A
 A U
 AU
 UA
 CG
 CG
 CG
 AU
 U
 AU
 AU
 AU
 UG
 A
 G A
 U U
 UA
 AU
 A G
 A U
 G A
 GU
 UA
 AU
 CG
 A A
 CG
 UA
 UA
 AU
 GC
 AU
 AU
 UG
 UA
 AU
 CG
 C A
 AU
 CG
 GC
 AU
 CG
 GC
 C A
 G U
 GC
 A A
 UA
 GC
 UA
 UA
 U U
 GC
 UG
 UA
 GC
 CG
 UG
 GC

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5

A



B




FIG. 6





FIG. 7




FIG. 8





FIG. 9

a



b




FIG. 10

GENE SILENCING

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 60/509,958, filed Oct. 9, 2003. The entire contents of the above application is herein incorporated by reference.

FIELD OF THE INVENTION

[0002] The field of the present invention relates generally to plant molecular biology. More specifically it relates to constructs and methods to suppress the expression of targeted genes.

BACKGROUND

[0003] Reduction of the activity of specific genes (also known as gene silencing, or gene suppression) is desirable for several aspects of genetic engineering in plants. There is still a need for methods and constructs that induce gene suppression against a wide selection of target genes, and that result in effective silencing of the target gene at high efficiency.

BRIEF DESCRIPTION OF THE FIGURES

[0004] **FIG. 1.** Predicted hairpin structure formed by the sequence surrounding miR172a-2. The mature microRNA is indicated by a grey box.

[0005] **FIG. 2.** miR172a-2 overexpression phenotype. a, Wild type (Columbia ecotype) plant, 3.5 weeks old. b, EAT-D plant, 3.5 weeks old. c, Wild type flower. d, EAT-D flower. Note absence of second whorl organs (petals). Arrow indicates sepal with ovules along the margins and stigmatic papillae at the tip. e, Cauline leaf margin from a 35S-EAT plant. Arrows indicate bundles of stigmatic papillae projecting from the margin. f, Solitary gynoecium (arrow) emerging from the axil of a cauline leaf of a 35S-EAT plant.

[0006] **FIG. 3.** The EAT gene contains a miRNA that is complementary to APETALA2 (AP2). a, Location of the EAT gene on chromosome 5. The T-DNA insertion and orientation of the 35S enhancers is indicated. The 21-nt sequence corresponding to miR172a-2 is shown below the EAT gene (SEQ ID NO: 86). b, Putative 21-nt miR172a-2/AP2 RNA duplex is shown below the gene structure of AP2. The GU wobble in the duplex is underlined. c, Alignment of AP221-nt region (black bar) and surrounding sequence with three other *Arabidopsis* AP2 family members, and with two maize AP2 genes (IDS1 and GL15). d, Alignment of miR172a-2 miRNA (black bar) and surrounding sequence with miR172-like sequences from *Arabidopsis*, tomato, soybean, potato and rice.

[0007] **FIG. 4.** miR172a-2 miRNA expression. a, Northern blot of total RNA from wild type (lanes 3 and 7) and EAT-D (lanes 4 and 8). Blots were probed with sense (lanes 1-4) or antisense (lanes 5-8) oligo to miR172a-2 miRNA. 100 pg of sense oligo (lanes 2 and 6) and antisense oligo (lanes 1 and 5) were loaded as hybridization controls. Nucleotide size markers are indicated on the left. b, S1 nuclease mapping of miR172a-2 miRNA. A 5'-end-labeled probe undigested (lane 1) or digested after hybridization to total RNA from wild-type (lane 2), EAT-D (lane 3), or tRNA (lane 4).

[0008] **FIG. 5.** Developmental expression pattern of miR172 family members. a, RT-PCR of total RNA from wild type seedlings harvested at 2, 5, 12, and 21 days after germination (lanes 1-4, respectively), or from mature leaves (lane 5) and floral buds (lane 6). Primers for PCR are indicated on the left. b, Northern analysis of mirR172 expression in the indicated mutants, relative to wild type (Col). Blot was probed with an oligo to miR172a-2; however, all miR172 members should cross hybridize.

[0009] **FIG. 6.** Expression analysis of putative EAT target genes. a, Northern blot analysis of polyA+ RNA isolated from wild type (Col) or EAT-D floral buds. Probes for hybridization are indicated on the right. b, Western blot of proteins from wild type or EAT-D floral buds, probed with AP2 antibody. RbcL, large subunit of ribulose 1,5-bisphosphate carboxylase as loading control.

[0010] **FIG. 7.** Identification of LAT-D. a, Location of the T-DNA insert in LAT-D, in between At2g28550 and At2g28560. The 4×35S enhancers are approximately 5 kb from At2g28550. b, RT-PCR analysis of At2g28550 expression in wild type versus LAT-D plants.

[0011] **FIG. 8.** EAT-D is epistatic to LAT-D. Genetic cross between EAT-D and LAT-D plants, with the resultant F1 plants shown, along with their flowering time (measured as rosette leaf number).

[0012] **FIG. 9.** Loss-of-function At2g28550 (2-28550) and At5g60120 (6-60120) mutants. Location of T-DNA in each line is indicated, along with intron/exon structure.

[0013] **FIG. 10.** Potential function of the miR172 miRNA family. a, Temporal expression of miR172a-2 and its relatives may cause temporal downregulation of AP2 targets (e.g. At2g28550 and At5g60120), which may trigger flowering once the target proteins drop below a critical threshold (dotted line). b, Dicer cleavage at various positions may generate at least four distinct miRNAs from the miR172 family (indicated as a single hairpin with a miRNA consensus sequence). Sequences at the 5' and 3' ends of each miRNA are indicated, with the invariant middle 15 nt shown as ellipses. The putative targets recognized by the individual miRNAs are in parentheses below each.

SUMMARY OF THE INVENTION

[0014] The invention provides methods and compositions useful in target sequence suppression and target sequence validation. The invention provides polynucleotide constructs useful for gene silencing, as well as cells, plants and seeds comprising the polynucleotides. The invention also provides a method for using microRNA to silence a target sequence.

DETAILED DESCRIPTION

[0015] Recently discovered small RNAs play an important role in controlling gene expression. Regulation of many developmental processes including flowering is controlled by small RNAs. It is now possible to engineer changes in gene expression of plant genes by using transgenic constructs which produce small RNAs in the plant.

[0016] The invention provides methods and compositions useful for suppressing targeted sequences. The compositions can be employed in any type of plant cell, and in other cells which comprise the appropriate processing components

(e.g., RNA interference components), including invertebrate and vertebrate animal cells. The compositions and methods are based on an endogenous miRNA silencing process discovered in *Arabidopsis*, a similar strategy can be used to extend the number of compositions and the organisms in which the methods are used. The methods can be adapted to work in any eukaryotic cell system. Additionally, the compositions and methods described herein can be used in individual cells, cells or tissue in culture, or in vivo in organisms, or in organs or other portions of organisms.

[0017] The compositions selectively suppress the target sequence by encoding a miRNA having substantial complementarity to a region of the target sequence. The miRNA is provided in a nucleic acid construct which, when transcribed into RNA, is predicted to form a hairpin structure which is processed by the cell to generate the miRNA, which then suppresses expression of the target sequence.

[0018] A nucleic acid construct is provided to encode the miRNA for any specific target sequence. Any miRNA can be inserted into the construct, such that the encoded miRNA selectively targets and suppresses the target sequence. The construct is modeled on the EAT (mir-172a) miRNA precursor from *Arabidopsis*.

[0019] A method for suppressing a target sequence is provided. The method employs the constructs above, in which a miRNA is designed to a region of the target sequence, and inserted into the construct. Upon introduction into a cell, the miRNA produced suppresses expression of the targeted sequence. The target sequence can be an endogenous plant sequence, or a heterologous transgene in the plant. The target gene may also be a gene from a plant pathogen, such as a pathogenic virus, nematode, insect, or mold or fungus.

[0020] A plant, cell, and seed comprising the construct and/or the miRNA is provided. Typically, the cell will be a cell from a plant, but other prokaryotic or eukaryotic cells are also contemplated, including but not limited to viral, bacterial, yeast, insect, nematode, or animal cells. Plant cells include cells from monocots and dicots. The invention also provides plants and seeds comprising the construct and/or the miRNA.

[0021] Units, prefixes, and symbols may be denoted in their SI accepted form. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxyl orientation, respectively. Numeric ranges recited within the specification are inclusive of the numbers defining the range and include each integer within the defined range. Amino acids may be referred to herein by either commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes. Unless otherwise provided for, software, electrical, and electronics terms as used herein are as defined in The New IEEE Standard Dictionary of Electrical and Electronics Terms (5th edition, 1993). The terms defined below are more fully defined by reference to the specification as a whole.

[0022] As used herein, “nucleic acid construct” or “construct” refers to an isolated polynucleotide which is introduced into a host cell. This construct may comprise any

combination of deoxyribonucleotides, ribonucleotides, and/or modified nucleotides. The construct may be transcribed to form an RNA, wherein the RNA may be capable of forming a double-stranded RNA and/or hairpin structure. This construct may be expressed in the cell, or isolated or synthetically produced. The construct may further comprise a promoter, or other sequences which facilitate manipulation or expression of the construct.

[0023] As used here “suppression” or “silencing” or “inhibition” are used interchangeably to denote the down-regulation of the expression of the product of a target sequence relative to its normal expression level in a wild type organism. Suppression includes expression that is decreased by about 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% relative to the wild type expression level.

[0024] As used herein, “encodes” or “encoding” refers to a DNA sequence which can be processed to generate an RNA and/or polypeptide.

[0025] As used herein, “expression” or “expressing” refers to the generation of an RNA transcript from an introduced construct, an endogenous DNA sequence, or a stably incorporated heterologous DNA sequence. The term may also refer to a polypeptide produced from an miRNA generated from any of the above DNA precursors.

[0026] As used herein, “heterologous” in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or is synthetically designed, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. A heterologous protein may originate from a foreign species or, if from the same species, is substantially modified from its original form by deliberate human intervention.

[0027] By “host cell” is meant a cell which contains an introduced nucleic acid construct and supports the replication and/or expression of the construct. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as fungi, yeast, insect, amphibian, nematode, or mammalian cells. Alternatively, the host cells are monocotyledonous or dicotyledonous plant cells. An example of a monocotyledonous host cell is a maize cell.

[0028] The term “introduced” means providing a nucleic acid or protein into a cell. Introduced includes reference to the incorporation of a nucleic acid into a eukaryotic or prokaryotic cell where the nucleic acid may be incorporated into the genome of the cell, and includes reference to the transient provision of a nucleic acid or protein to the cell. Introduced includes reference to stable or transient transformation methods, as well as sexually crossing.

[0029] The term “isolated” refers to material, such as a nucleic acid or a protein, which is: (1) substantially free from components which normally accompany or interact with the material as found in its naturally occurring environment or (2) if the material is in its natural environment, the material has been altered by deliberate human intervention to a composition and/or placed at a locus in the cell other than the locus native to the material.

[0030] As used herein, “miRNA” refers to an oligoribonucleic acid, which suppresses expression of a polynucle-

otide comprising the target sequence transcript. A “miRNA precursor” refers to a larger polynucleotide which is processed to produce a mature miRNA, and includes a DNA which encodes an RNA precursor, and an RNA transcript comprising the miRNA. A “mature miRNA” refers to the miRNA generated from the processing of a miRNA precursor. A “miRNA template” is an oligonucleotide region, or regions, in a nucleic acid construct which encodes the miRNA. The “backside” region of a miRNA is a portion of a polynucleotide construct which is substantially complementary to the miRNA template and is predicted to base pair with the miRNA template. The miRNA template and backside may form a double-stranded polynucleotide, including a hairpin structure.

[0031] As used herein, the phrases “target sequence” and “sequence of interest” are used interchangeably. Target sequence is used to mean the nucleic acid sequence that is selected for suppression of expression, and is not limited to polynucleotides encoding polypeptides. The target sequence comprises a sequence that is substantially or completely complementary to the miRNA. The target sequence can be RNA or DNA, and may also refer to a polynucleotide comprising the target sequence.

[0032] As used herein, “nucleic acid” means a polynucleotide and includes single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases. Nucleic acids may also include fragments and modified nucleotides.

[0033] By “nucleic acid library” is meant a collection of isolated DNA or RNA molecules which comprise and substantially represent the entire transcribed fraction of a genome of a specified organism or of a tissue from that organism. Construction of exemplary nucleic acid libraries, such as genomic and cDNA libraries, is taught in standard molecular biology references such as Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology*, Vol. 152, Academic Press, Inc., San Diego, Calif. (Berger); Sambrook et al., *Molecular Cloning—A Laboratory Manual*, 2nd ed., Vol. 1-3 (1989); and *Current Protocols in Molecular Biology*, F. M. Ausubel et al., Eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc. (1994).

[0034] As used herein “operably linked” includes reference to a functional linkage of at least two sequences. Operably linked includes linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence.

[0035] As used herein, “plant” includes plants and plant parts including but not limited to plant cells, plant tissue such as leaves, stems, roots, flowers, and seeds.

[0036] As used herein, “polypeptide” means proteins, protein fragments, modified proteins, amino acid sequences and synthetic amino acid sequences. The polypeptide can be glycosylated or not.

[0037] As used herein, “promoter” includes reference to a region of DNA that is involved in recognition and binding of an RNA polymerase and other proteins to initiate transcription.

[0038] The term “selectively hybridizes” includes reference to hybridization, under stringent hybridization condi-

tions, of a nucleic acid sequence to a specified nucleic acid target sequence to a detectably greater degree (e.g., at least 2-fold over background) than its hybridization to non-target nucleic acid sequences and to the substantial exclusion of non-target nucleic acids. Selectively hybridizing sequences typically have about at least 80% sequence identity, or 90% sequence identity, up to and including 100% sequence identity (i.e., fully complementary) with each other.

[0039] The term “stringent conditions” or “stringent hybridization conditions” includes reference to conditions under which a probe will selectively hybridize to its target sequence. Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences can be identified which are 100% complementary to the probe (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, optionally less than 500 nucleotides in length.

[0040] Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55° C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60 to 65° C.

[0041] Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl, *Anal. Biochem.*, 138:267-284 (1984): $T_m = 81.5^\circ \text{C.} + 16.6 (\log M) + 0.41 (\% \text{ GC}) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1° C. for each 1% of mismatching; thus, T_m , hybridization and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $\geq 90\%$ identity are sought, the T_m can be decreased 10° C. Generally, stringent conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4° C. lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash

at 6, 7, 8, 9, or 10° C. lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20° C. lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45° C. (aqueous solution) or 32° C. (formamide solution) it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 “Overview of principles of hybridization and the strategy of nucleic acid probe assays”, Elsevier, New York (1993); and *Current Protocols in Molecular Biology*, Chapter 2, Ausubel et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). Hybridization and/or wash conditions can be applied for at least 10, 30, 60, 90, 120, or 240 minutes.

[0042] As used herein, “transgenic” includes reference to a plant or a cell which comprises a heterologous polynucleotide. Generally, the heterologous polynucleotide is stably integrated within the genome such that the polynucleotide is passed on to successive generations. Transgenic is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term “transgenic” as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

[0043] As used herein, “vector” includes reference to a nucleic acid used in introduction of a polynucleotide of the invention into a host cell. Expression vectors permit transcription of a nucleic acid inserted therein.

[0044] Polynucleotide sequences may have substantial identity, substantial homology, or substantial complementarity to the selected region of the target gene. As used herein “substantial identity”, and “substantial homology” indicate sequences that have sequence identity or homology to each other. Generally, sequences that are substantially identical or substantially homologous will have about 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity wherein the percent sequence identity is based on the entire sequence and is determined by GAP alignment using default parameters (GCG, GAP version 10, Accelrys, San Diego, Calif.). GAP uses the algorithm of Needleman and Wunsch (*J. Mol. Biol.* 48:443-453, 1970) to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of sequence gaps. Sequences which have 100% identity are identical. “Substantial complementarity” refers to sequences that are complementary to each other, and are able to base pair with each other. In describing complementary

sequences, if all the nucleotides in the first sequence will base pair to the second sequence, these sequences are fully complementary.

[0045] Through a forward genetics approach, a microRNA that confers a developmental phenotype in *Arabidopsis* was identified. This miRNA, miR172a-2 (Park et al., *Curr. Biol.* 12:1484-1495 2002), causes early flowering and defects in floral organ identity when overexpressed. The predicted target of miR172a-2 is a small subfamily of APETALA2-like transcription factors (Okamuro et al. 1997). Overexpression of miR172a-2 downregulates at least one member of this family. In addition, overexpression of one of the AP2-like target genes, At2g28550, causes late flowering. This result, in conjunction with loss-of-function analyses of At2g28550 and another target gene, At5g60120, indicates that at least some of the AP2-like genes targeted by miR172a-2 normally function as floral repressors. The EAT-D line overexpressing miR172a-2 has a wild-type response to photoperiod. The genomic region encoding the miRNA was also identified (SEQ ID NO: 1) and used to produce a cassette into which other miRNAs to target sequences can be inserted (SEQ ID NO: 3), and to produce an expression vector (SEQ ID NO: 44) useful for cloning the cassettes and expressing the miRNA. The expression vector comprises the 1.4 kb region encoding the miRNA. Expression of this region is processed in the cell to produce the miRNA which suppresses expression of the target gene. Alternatively, the miRNA may be synthetically produced and introduced to the cell directly.

[0046] In one embodiment, there is provided a method for the suppression of a target sequence comprising introducing into a cell a nucleic acid construct encoding a miRNA substantially complementary to the target. In some embodiments the miRNA comprises about 10-200 nucleotides, about 10-15, 15-20, 19, 20, 21, 22, 23, 24, 25, 26, 27, 25-30, 30-50, 50-100, 100-150, or about 150-200 nucleotides. In some embodiments the nucleic acid construct encodes the miRNA. In some embodiments the nucleic acid construct encodes a polynucleotide precursor which may form a double-stranded RNA, or hairpin structure comprising the miRNA. In some embodiments, nucleotides 39-59 and 107-127 of SEQ ID NO: 3 are replaced by the backside of the miRNA template and the miRNA template respectively. In some embodiments, this new sequence replaces the equivalent region of SEQ ID NO: 1. In further embodiments, this new sequence replaces the equivalent region of SEQ ID NO: 44.

[0047] In some embodiments, the nucleic acid construct comprises a modified endogenous plant miRNA precursor, wherein the precursor has been modified to replace the endogenous miRNA encoding regions with sequences designed to produce a miRNA directed to the target sequence. In some embodiments the miRNA precursor template is a miR172a miRNA precursor. In some embodiments, the miR172a precursor is from a dicot or a monocot. In some embodiments the miRNA precursor is SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 44.

[0048] In another embodiment the method comprises:

[0049] A method of inhibiting expression of a target sequence in a cell comprising:

[0050] (a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide, wherein the polynucleotide comprises in the following order:

[0051] (i) at least about 20 contiguous nucleotides in the region of nucleotides 1-38 of SEQ ID NO: 3,

[0052] (ii) a first oligonucleotide of 10 to about 50 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide,

[0053] (iii) at least about 20 contiguous nucleotides in the region of nucleotides 60-106 of SEQ ID NO: 3,

[0054] (iv) the second oligonucleotide of about 10 to about 50 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and the second oligonucleotide is substantially complementary to the target sequence, and

[0055] (v) at least about 20 contiguous nucleotides in the region of nucleotides 128-159 of SEQ ID NO:3;

wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin, and

[0056] (b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

[0057] In another embodiment the method comprises:

[0058] A method of inhibiting expression of a gene comprising a target sequence in a cell comprising:

[0059] (a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide, wherein the polynucleotide comprises in the following order:

[0060] (i) nucleotides 1-38 of SEQ ID NO: 3,

[0061] (ii) a first oligonucleotide of 21 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide,

[0062] (iii) nucleotides 60-106 of SEQ ID NO: 3,

[0063] (iv) the second oligonucleotide of 21 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and wherein the second oligonucleotide is substantially complementary to the target sequence, and

[0064] (v) nucleotides 128-159 of SEQ ID NO:3;

wherein polynucleotide encodes an RNA precursor capable of forming a hairpin, and

[0065] (b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

[0066] In another embodiment, the method comprises selecting a target sequence of a gene, and designing a nucleic

acid construct comprising polynucleotide encoding a miRNA substantially complementary to the target sequence. In some embodiments, the target sequence is selected from any region of the gene. In some embodiments, the target sequence is selected from an untranslated region. In some embodiments, the target sequence is selected from a coding region of the gene. In some embodiments, the target sequence is selected from a region about 50 to about 200 nucleotides upstream from the stop codon, including regions from about 50-75, 75-100, 100-125, 125-150, or 150-200 upstream from the stop codon. In further embodiments, the target sequence and/or the miRNA is based on the polynucleotides and process of EAT suppression of *Apetela2*-like genes in *Arabidopsis thaliana*. In some embodiments, nucleotides 39-59 and 107-127 of SEQ ID NO: 3 are replaced by the backside of the miRNA template (first oligonucleotide) and the miRNA template (second oligonucleotide) respectively. In some embodiments, this new sequence replaces the equivalent region of SEQ ID NO: 1. In further embodiments, this new sequence replaces the equivalent region of SEQ ID NO: 44.

[0067] In some embodiments, the miRNA template, (i.e. the polynucleotide encoding the miRNA), and thereby the miRNA, may comprise some mismatches relative to the target sequence. In some embodiments the miRNA template has ≥ 1 nucleotide mismatch as compared to the target sequence, for example, the miRNA template can have 1, 2, 3, 4, 5, or more mismatches as compared to the target sequence. This degree of mismatch may also be described by determining the percent identity of the miRNA template to the complement of the target sequence. For example, the miRNA template may have a percent identity including about at least 70%, 75%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% as compared to the complement of the target sequence.

[0068] In some embodiments, the miRNA template, (i.e. the polynucleotide encoding the miRNA) and thereby the miRNA, may comprise some mismatches relative to the miRNA backside. In some embodiments the miRNA template has ≥ 1 nucleotide mismatch as compared to the miRNA backside, for example, the miRNA template can have 1, 2, 3, 4, 5, or more mismatches as compared to the miRNA backside. This degree of mismatch may also be described by determining the percent identity of the miRNA template to the complement of the miRNA backside. For example, the miRNA template may have a percent identity including about at least 70%, 75%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% as compared to the complement of the miRNA backside.

[0069] In some embodiments, the target sequence is selected from a plant pathogen. Plants or cells comprising a miRNA directed to the target sequence of the pathogen are expected to have decreased sensitivity and/or increased resistance to the pathogen. In some embodiments, the miRNA is encoded by a nucleic acid construct further comprising an operably linked promoter. In some embodiments, the promoter is a pathogen-inducible promoter.

[0070] In another embodiment, the method comprises replacing the miRNA encoding sequence in the polynucle-

otide of SEQ ID NO: 3 with a sequence encoding a miRNA substantially complementary to the target region of the target gene.

[0071] In another embodiment a method is provided comprising a method of inhibiting expression of a target sequence in a cell comprising:

[0072] (a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide encoding a modified plant miRNA precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide encodes a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin; and

[0073] (b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

[0074] In another embodiment a method is provided comprising a method of inhibiting expression of a target sequence in a cell comprising:

[0075] (a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide encoding a modified plant miR172 miRNA precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide encodes a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin; and

[0076] (b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

[0077] In another embodiment a method is provided comprising a method of inhibiting expression of a target sequence in a cell comprising:

[0078] (a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide encoding a modified *Arabidopsis* miR172 miRNA precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide encodes a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin; and

[0079] (b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

[0080] In another embodiment, there is provided a nucleic acid construct for suppressing a target sequence. The nucleic acid construct encodes a miRNA substantially complementary to the target. In some embodiments, the nucleic acid

construct further comprises a promoter operably linked to the polynucleotide encoding the miRNA. In some embodiments, the nucleic acid construct lacking a promoter is designed and introduced in such a way that it becomes operably linked to a promoter upon integration in the host genome. In some embodiments, the nucleic acid construct is integrated using recombination, including site-specific recombination. See, for example, WO 99/25821, herein incorporated by reference. In some embodiments, the nucleic acid construct is an RNA. In some embodiments, the nucleic acid construct comprises at least one recombination site, including site-specific recombination sites.

[0081] In some embodiments, the nucleic acid construct comprises a modified endogenous plant miRNA precursor, wherein the precursor has been modified to replace the miRNA encoding region with a sequence designed to produce a miRNA directed to the target sequence. In some embodiments the miRNA precursor template is a miR172a miRNA precursor. In some embodiments, the miR172a precursor is from a dicot or a monocot. In some embodiments the miR172a precursor is from *Arabidopsis thaliana*, tomato, soybean, rice, or corn. In some embodiments the miRNA precursor is SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 44.

[0082] In another embodiment, the nucleic acid construct comprises an isolated polynucleotide comprising a polynucleotide which encodes a modified plant miRNA precursor, the modified precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide comprises a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin.

[0083] In another embodiment, the nucleic acid construct comprises an isolated polynucleotide comprising a polynucleotide which encodes a modified plant miR172 miRNA precursor, the modified precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide comprises a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin.

[0084] In another embodiment, the nucleic acid construct comprises an isolated polynucleotide comprising a polynucleotide which encodes a modified *Arabidopsis* miR172 miRNA precursor, the modified precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide comprises a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin.

[0085] In some embodiments the miRNA comprises about 10-200 nucleotides, about 10-15, 15-20, 19, 20, 21, 22, 23, 24, 25, 26, 27, 25-30, 30-50, 50-100, 100-150, or about 150-200 nucleotides. In some embodiments the nucleic acid construct encodes the miRNA. In some embodiments the

nucleic acid construct encodes a polynucleotide precursor which may form a double-stranded RNA, or hairpin structure comprising the miRNA. In some embodiments, nucleotides 39-59 and/or 107-127 of SEQ ID NO: 3 are replaced by the backside of the miRNA template and the miRNA template respectively. In some embodiments, this new sequence replaces the equivalent region of SEQ ID NO: 1. In further embodiments, this new sequence replaces the equivalent region of SEQ ID NO: 44. In some embodiments, the target region is selected from any region of the target sequence. In some embodiments, the target region is selected from a untranslated region. In some embodiments, the target region is selected from a coding region of the target sequence. In some embodiments, the target region is selected from a region about 50 to about 200 nucleotides upstream from the stop codon, including regions from about 50-75, 75-100, 100-125, 125-150, or 150-200 upstream from the stop codon. In further embodiments, the target region and/or the miRNA is based on the polynucleotides and process of EAT suppression of *Apetela2*-like sequences in *Arabidopsis thaliana*.

[0086] In another embodiment the nucleic acid construct comprises an isolated polynucleotide comprising in the following order at least 20 contiguous nucleotides in the region from nucleotides 1-38 of SEQ ID NO: 3, a first oligonucleotide of about 10 to about 50 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide, at least about 20 contiguous nucleotides in the region from nucleotides 60-106 of SEQ ID NO: 3, a second oligonucleotide of about 10 to about 50 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and the second oligonucleotide is substantially complementary to the target sequence, and at least about 20 contiguous nucleotides in the region from nucleotides 128-159 of SEQ ID NO: 3, wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin structure.

[0087] In another embodiment the nucleic acid construct comprises an isolated polynucleotide comprising in the following order nucleotides 1-38 of SEQ ID NO: 3, a first oligonucleotide of 21 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide, nucleotides 60-106 of SEQ ID NO: 3, a second oligonucleotide of 21 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and the second oligonucleotide is substantially complementary to the target sequence, and nucleotides 128-159 of SEQ ID NO: 3, wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin.

[0088] In some embodiments there are provided cells, plants, and seeds comprising the introduced polynucleotides, and/or produced by the methods of the invention. The cells include prokaryotic and eukaryotic cells, including but not limited to bacteria, yeast, fungi, viral, invertebrate, vertebrate, and plant cells. Plants, plant cells, and seeds of the invention include gymnosperms, monocots and dicots, including but not limited to, for example, rice, wheat, oats, barley, millet, sorghum, soy, sunflower, safflower, canola, alfalfa, cotton, *Arabidopsis*, and tobacco.

[0089] In some embodiments, the cells, plants, and/or seeds comprise a nucleic acid construct comprising a modified plant miRNA precursor, wherein the precursor has been

modified to replace the endogenous miRNA encoding regions with sequences designed to produce a miRNA directed to the target sequence. In some embodiments the miRNA precursor template is a miR172a miRNA precursor. In some embodiments, the miR172a precursor is from a dicot or a monocot. In some embodiments the miR172a precursor is from *Arabidopsis thaliana*. In some embodiments the miRNA precursor is SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 44. In some embodiments the miRNA precursor is encoded by SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 44. In some embodiments, the nucleic acid construct comprises at least one recombination site, including site-specific recombination sites.

[0090] The present invention concerns methods and compositions useful in suppression of a target sequence and/or validation of function. The invention also relates to a method for using microRNA (miRNA) mediated RNA interference (RNAi) to silence or suppress a target sequence to evaluate function, or to validate a target sequence for phenotypic effect and/or trait development. Specifically, the invention relates to constructs comprising small nucleic acid molecules, miRNAs, capable of inducing silencing, and methods of using these miRNAs to selectively silence target sequences.

[0091] RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Fire et al., *Nature* 391:806 1998). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing (PTGS) or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla (Fire et al., *Trends Genet.* 15:358 1999). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA of viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized.

[0092] The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Berstein et al., *Nature* 409:363 2001). Short interfering RNAs derived from dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes (Elbashir et al., *Genes Dev.* 15:188 2001). Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner et al., 2001, *Science* 293:834). The RNAi response also features an endonuclease complex, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence complementarity to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir et al., *Genes Dev.* 15:188 2001). In addition, RNA interference can

also involve small RNA (e.g., microRNA, or miRNA) mediated gene silencing, presumably through cellular mechanisms that regulate chromatin structure and thereby prevent transcription of target gene sequences (see, e.g., Allshire, *Science* 297:1818-1819 2002; Volpe et al., *Science* 297:1833-1837 2002; Jenuwein, *Science* 297:2215-2218 2002; and Hall et al., *Science* 297:2232-2237 2002). As such, miRNA molecules of the invention can be used to mediate gene silencing via interaction with RNA transcripts or alternately by interaction with particular gene sequences, wherein such interaction results in gene silencing either at the transcriptional or post-transcriptional level.

[0093] RNAi has been studied in a variety of systems. Fire et al. (*Nature* 391:806 1998) were the first to observe RNAi in *C. elegans*. Wianny and Goetz (*Nature Cell Biol.* 2:70 1999) describe RNAi mediated by dsRNA in mouse embryos. Hammond et al. (*Nature* 404:293 2000) describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir et al., (*Nature* 411:494 2001) describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa cells.

[0094] Small RNAs play an important role in controlling gene expression. Regulation of many developmental processes, including flowering, is controlled by small RNAs. It is now possible to engineer changes in gene expression of plant genes by using transgenic constructs which produce small RNAs in the plant.

[0095] Small RNAs appear to function by base-pairing to complementary RNA or DNA target sequences. When bound to RNA, small RNAs trigger either RNA cleavage or translational inhibition of the target sequence. When bound to DNA target sequences, it is thought that small RNAs can mediate DNA methylation of the target sequence. The consequence of these events, regardless of the specific mechanism, is that gene expression is inhibited.

[0096] It is thought that sequence complementarity between small RNAs and their RNA targets helps to determine which mechanism, RNA cleavage or translational inhibition, is employed. It is believed that siRNAs, which are perfectly complementary with their targets, work by RNA cleavage. Some miRNAs have perfect or near-perfect complementarity with their targets, and RNA cleavage has been demonstrated for at least a few of these miRNAs. Other miRNAs have several mismatches with their targets, and apparently inhibit their targets at the translational level. Again, without being held to a particular theory on the mechanism of action, a general rule is emerging that perfect or near-perfect complementarity favors RNA cleavage, whereas translational inhibition is favored when the miRNA/target duplex contains many mismatches. The apparent exception to this is microRNA 172 (miR172) in plants. One of the targets of miR172 is APETALA2 (AP2), and although miR172 shares near-perfect complementarity with AP2 it appears to cause translational inhibition of AP2 rather than RNA cleavage.

[0097] MicroRNAs (miRNAs) are noncoding RNAs of about 19 to about 24 nucleotides (nt) in length that have been identified in both animals and plants (Lagos-Quintana et al., *Science* 294:853-858 2001, Lagos-Quintana et al., *Curr. Biol.* 12:735-739 2002; Lau et al., *Science* 294:858-862 2001; Lee and Ambros, *Science* 294:862-864 2001; Llave et

al., *Plant Cell* 14:1605-1619 2002; Mourelatos et al., *Genes. Dev.* 16:720-728 2002; Park et al., *Curr. Biol.* 12:1484-1495 2002; Reinhart et al., *Genes. Dev.* 16:1616-1626 2002). They are processed from longer precursor transcripts that range in size from approximately 70 to 200 nt, and these precursor transcripts have the ability to form stable hairpin structures. In animals, the enzyme involved in processing miRNA precursors is called Dicer, an RNase III-like protein (Grishok et al., *Cell* 106:23-34 2001; Hutvagner et al., *Science* 293:834-838 2001; Ketting et al., *Genes. Dev.* 15:2654-2659 2001). Plants also have a Dicer-like enzyme, DCL1 (previously named CARPEL FACTORY/SHORT INTEGUMENTS1/SUSPENSOR1), and recent evidence indicates that it, like Dicer, is involved in processing the hairpin precursors to generate mature miRNAs (Park et al., *Curr. Biol.* 12:1484-1495 2002; Reinhart et al., *Genes. Dev.* 16:1616-1626 2002). Furthermore, it is becoming clear from recent work that at least some miRNA hairpin precursors originate as longer polyadenylated transcripts, and several different miRNAs and associated hairpins can be present in a single transcript (Lagos-Quintana et al., *Science* 294:853-858 2001; Lee et al., *EMBO J.* 21:4663-4670 2002).

[0098] In animals, there is direct evidence indicating a role for specific miRNAs in development. The lin-4 and let-7 miRNAs in *C. elegans* have been found to control temporal development, based on the phenotypes generated when the genes producing the lin-4 and let-7 miRNAs are mutated (Lee et al., *Cell* 75:843-854 1993; Reinhart et al., *Nature* 403:901-906 2000). In addition, both miRNAs display a temporal expression pattern consistent with their roles in developmental timing. Other animal miRNAs display developmentally regulated patterns of expression, both temporal and tissue-specific (Lagos-Quintana et al., *Science* 294:853-858 2001, Lagos-Quintana et al., *Curr. Biol.* 12:735-739 2002; Lau et al., *Science* 294:858-862 2001; Lee and Ambros, *Science* 294:862-864 2001), leading to the hypothesis that miRNAs may, in many cases, be involved in the regulation of important developmental processes. Likewise, in plants, the differential expression patterns of many miRNAs suggests a role in development (Llave et al., *Plant Cell* 14:1605-1619 2002; Park et al., *Curr. Biol.* 12:1484-1495 2002; Reinhart et al., *Genes. Dev.* 16:1616-1626 2002). However, a developmental role for miRNAs has not been directly proven in plants, because to date there has been no report of a developmental phenotype associated with a specific plant miRNA.

[0099] MicroRNAs appear to regulate target genes by binding to complementary sequences located in the transcripts produced by these genes. In the case of lin-4 and let-7, the target sites are located in the 3' UTRs of the target miRNAs (Lee et al., *Cell* 75:843-854 1993; Wightman et al., *Cell* 75:855-862 1993; Reinhart et al., *Nature* 403:901-906 2000; Slack et al., *Mol. Cell* 5:659-669 2000), and there are several mismatches between the lin-4 and let-7 miRNAs and their target sites. Binding of the lin-4 or let-7 miRNA appears to cause downregulation of steady-state levels of the protein encoded by the target miRNA without affecting the transcript itself (Olsen and Ambros, *Dev. Biol.* 216:671-680 1999). On the other hand, recent evidence suggests that miRNAs can, in some cases, cause specific RNA cleavage of the target transcript within the target site, and this cleavage step appears to require 100% complementarity between the miRNA and the target transcript (Hutvagner and Zamore, *Science* 297:2056-2060 2002; Llave et al., *Plant Cell*

14:1605-1619 2002). It seems likely that miRNAs can enter at least two pathways of target gene regulation: Protein downregulation when target complementarity is <100%, and RNA cleavage when target complementarity is 100%. MicroRNAs entering the RNA cleavage pathway are analogous to the 21-25 nt short interfering RNAs (siRNAs) generated during RNA interference (RNAi) in animals and posttranscriptional gene silencing (PTGS) in plants (Hamilton and Baulcombe 1999; Hammond et al., 2000; Zamore et al., 2000; Elbashir et al., 2001), and likely are incorporated into an RNA-induced silencing complex (RISC) that is similar or identical to that seen for RNAi.

[0100] Identifying the targets of miRNAs with bioinformatics has not been successful in animals, and this is probably due to the fact that animal miRNAs have a low degree of complementarity with their targets. On the other hand, bioinformatic approaches have been successfully used to predict targets for plant miRNAs (Llave et al., *Plant Cell* 14:1605-1619 2002; Park et al., *Curr. Biol.* 12:1484-1495 2002; Rhoades et al., *Cell* 110:513-520 2002), and thus it appears that plant miRNAs have higher overall complementarity with their putative targets than do animal miRNAs. Most of these predicted target transcripts of plant miRNAs encode members of transcription factor families implicated in plant developmental patterning or cell differentiation. Nonetheless, biological function has not been directly demonstrated for any plant miRNA. Although Llave et al. (*Science* 297:2053-2056 2002) have shown that a transcript for a SCARECROW-like transcription factor is a target of the *Arabidopsis* miRNA mir171, these studies were performed in a heterologous species and no plant phenotype associated with mir171 was reported.

[0101] The methods provided can be practiced in any organism in which a method of transformation is available, and for which there is at least some sequence information for the target sequence, or for a region flanking the target sequence of interest. It is also understood that two or more sequences could be targeted by sequential transformation, co-transformation with more than one targeting vector, or the construction of a DNA construct comprising more than one miRNA sequence. The methods of the invention may also be implemented by a combinatorial nucleic acid library construction in order to generate a library of miRNAs directed to random target sequences. The library of miRNAs could be used for high-throughput screening for gene function validation.

[0102] General categories of sequences of interest include, for example, those genes involved in regulation or information, such as zinc fingers, transcription factors, homeotic genes, or cell cycle and cell death modulators, those involved in communication, such as kinases, and those involved in housekeeping, such as heat shock proteins.

[0103] Target sequences further include coding regions and non-coding regions such as promoters, enhancers, terminators, introns and the like, which may be modified in order to alter the expression of a gene of interest. For example, an intron sequence can be added to the 5' region to increase the amount of mature message that accumulates (see for example Buchman and Berg, *Mol. Cell Biol.* 8:4395-4405 (1988); and Callis et al., *Genes Dev.* 1:1183-1200 (1987)).

[0104] The target sequence may be an endogenous sequence, or may be an introduced heterologous sequence,

or transgene. For example, the methods may be used to alter the regulation or expression of a transgene, or to remove a transgene or other introduced sequence such as an introduced site-specific recombination site. The target sequence may also be a sequence from a pathogen, for example, the target sequence may be from a plant pathogen such as a virus, a mold or fungus, an insect, or a nematode. A miRNA could be expressed in a plant which, upon infection or infestation, would target the pathogen and confer some degree of resistance to the plant.

[0105] In plants, other categories of target sequences include genes affecting agronomic traits, insect resistance, disease resistance, herbicide resistance, sterility, grain characteristics, and commercial products. Genes of interest also included those involved in oil, starch, carbohydrate, or nutrient metabolism as well as those affecting, for example, kernel size, sucrose loading, and the like. The quality of grain is reflected in traits such as levels and types of oils, saturated and unsaturated, quality and quantity of essential amino acids, and levels of cellulose. For example, genes of the phytic acid biosynthetic pathway could be suppressed to generate a high available phosphorous phenotype. See, for example, phytic acid biosynthetic enzymes including inositol polyphosphate kinase-2 polynucleotides, disclosed in WO 02/059324, inositol 1,3,4-trisphosphate 5/6-kinase polynucleotides, disclosed in WO 03/027243, and myo-inositol 1-phosphate synthase and other phytate biosynthetic polynucleotides, disclosed in WO 99/05298, all of which are herein incorporated by reference. Genes in the lignification pathway could be suppressed to enhance digestibility or energy availability. Genes affecting cell cycle or cell death could be suppressed to affect growth or stress response. Genes affecting DNA repair and/or recombination could be suppressed to increase genetic variability. Genes affecting flowering time could be suppressed, as well as genes affecting fertility. Any target sequence could be suppressed in order to evaluate or confirm its role in a particular trait or phenotype, or to dissect a molecular, regulatory, biochemical, or proteomic pathway or network.

[0106] A number of promoters can be used, these promoters can be selected based on the desired outcome. It is recognized that different applications will be enhanced by the use of different promoters in plant expression cassettes to modulate the timing, location and/or level of expression of the miRNA. Such plant expression cassettes may also contain, if desired, a promoter regulatory region (e.g., one conferring inducible, constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific/selective expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

[0107] Constitutive, tissue-preferred or inducible promoters can be employed. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'-promoter derived from T-DNA of *Agrobacterium tumefaciens*, the ubiquitin 1 promoter, the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (U.S. Pat. No. 5,683,439), the Nos promoter, the pEmu promoter, the rubisco promoter, the GRP1-8 promoter and other transcription initiation regions from various plant genes known to those of skill. If low level expression is desired, weak promoter(s) may be used. Weak constitutive promoters include, for example, the core pro-

motor of the Rsyn7 promoter (WO 99/43838 and U.S. Pat. No. 6,072,050), the core 35S CaMV promoter, and the like. Other constitutive promoters include, for example, U.S. Pat. Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; and 5,608,142. See also, U.S. Pat. No. 6,177,611, herein incorporated by reference.

[0108] Examples of inducible promoters are the Adh1 promoter which is inducible by hypoxia or cold stress, the Hsp70 promoter which is inducible by heat stress, the PPKK promoter and the pepcarboxylase promoter which are both inducible by light. Also useful are promoters which are chemically inducible, such as the In2-2 promoter which is safer induced (U.S. Pat. No. 5,364,780), the ERE promoter which is estrogen induced, and the Axig1 promoter which is auxin induced and tapetum specific but also active in callus (PCT US01/22169).

[0109] Examples of promoters under developmental control include promoters that initiate transcription preferentially in certain tissues, such as leaves, roots, fruit, seeds, or flowers. An exemplary promoter is the anther specific promoter 5126 (U.S. Pat. Nos. 5,689,049 and 5,689,051). Examples of seed-preferred promoters include, but are not limited to, 27 kD gamma zein promoter and waxy promoter, Boronat, A. et al. (1986) *Plant Sci.* 47:95-102; Reina, M. et al. *Nucl. Acids Res.* 18(21):6426; and Kloesgen, R. B. et al. (1986) *Mol. Gen. Genet.* 203:237-244. Promoters that express in the embryo, pericarp, and endosperm are disclosed in U.S. Pat. No. 6,225,529 and PCT publication WO 00/12733. The disclosures each of these are incorporated herein by reference in their entirety.

[0110] In some embodiments it will be beneficial to express the gene from an inducible promoter, particularly from a pathogen-inducible promoter. Such promoters include those from pathogenesis-related proteins (PR proteins), which are induced following infection by a pathogen; e.g., PR proteins, SAR proteins, beta-1,3-glucanase, chitinase, etc. See, for example, Redolfi et al. (1983) *Neth. J. Plant Pathol.* 89:245-254; Uknes et al. (1992) *Plant Cell* 4:645-656; and Van Loon (1985) *Plant Mol. Virol.* 4:111-116. See also WO 99/43819, herein incorporated by reference.

[0111] Of interest are promoters that are expressed locally at or near the site of pathogen infection. See, for example, Marineau et al. (1987) *Plant Mol. Biol.* 9:335-342; Matton et al. (1989) *Molecular Plant-Microbe Interactions* 2:325-331; Somsisch et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:2427-2430; Somsisch et al. (1988) *Mol. Gen. Genet.* 2:93-98; and Yang (1996) *Proc. Natl. Acad. Sci. USA* 93:14972-14977. See also, Chen et al. (1996) *Plant J.* 10:955-966; Zhang et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:2507-2511; Warner et al. (1993) *Plant J.* 3:191-201; Siebertz et al. (1989) *Plant Cell* 1:961-968; U.S. Pat. No. 5,750,386 (nematode-inducible); and the references cited therein. Of particular interest is the inducible promoter for the maize PRms gene, whose expression is induced by the pathogen *Fusarium moniliforme* (see, for example, Cordero et al. (1992) *Physiol. Mol. Plant Pathol.* 41:189-200).

[0112] Additionally, as pathogens find entry into plants through wounds or insect damage, a wound-inducible promoter may be used in the constructions of the polynucleotides. Such wound-inducible promoters include potato proteinase inhibitor (pin II) gene (Ryan (1990) *Ann. Rev.*

Phytopath. 28:425-449; Duan et al. (1996) *Nature Biotech.* 14:494-498; wun1 and wun2, U.S. Pat. No. 5,428,148; win1 and win2 (Stanford et al. (1989) *Mol. Gen. Genet.* 215:200-208); systemin (McGurl et al. (1992) *Science* 225:1570-1573); WIP1 (Rohmeier et al. (1993) *Plant Mol. Biol.* 22:783-792; Eckelkamp et al. (1993) *FEBS Lett.* 323:73-76); MPI gene (Corderok et al. (1994) *Plant J.* 6(2):141-150); and the like, herein incorporated by reference.

[0113] Chemical-regulated promoters can be used to modulate the expression of a gene in a plant through the application of an exogenous chemical regulator. Depending upon the objective, the promoter may be a chemical-inducible promoter, where application of the chemical induces gene expression, or a chemical-repressible promoter, where application of the chemical represses gene expression. Chemical-inducible promoters are known in the art and include, but are not limited to, the maize In2-2 promoter, which is activated by benzenesulfonamide herbicide safeners, the maize GST promoter, which is activated by hydrophobic electrophilic compounds that are used as pre-emergent herbicides, and the tobacco PR-1a promoter, which is activated by salicylic acid. Other chemical-regulated promoters of interest include steroid-responsive promoters (see, for example, the glucocorticoid-inducible promoter in Schena et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:10421-10425 and McNellis et al. (1998) *Plant J.* 14(2):247-257) and tetracycline-inducible and tetracycline-repressible promoters (see, for example, Gatz et al. (1991) *Mol. Gen. Genet.* 227:229-237, and U.S. Pat. Nos. 5,814,618 and 5,789,156), herein incorporated by reference.

[0114] Tissue-preferred promoters can be utilized to target enhanced expression of a sequence of interest within a particular plant tissue. Tissue-preferred promoters include Yamamoto et al. (1997) *Plant J.* 12(2):255-265; Kawamata et al. (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen et al. (1997) *Mol. Gen. Genet.* 254(3):337-343; Russell et al. (1997) *Transgenic Res.* 6(2):157-168; Rinehart et al. (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp et al. (1996) *Plant Physiol.* 112(2):525-535; Canevascini et al. (1996) *Plant Physiol.* 112(2):513-524; Yamamoto et al. (1994) *Plant Cell Physiol.* 35(5):773-778; Lam (1994) *Results Probl. Cell Differ.* 20:181-196; Orozco et al. (1993) *Plant Mol. Biol.* 23(6):1129-1138; Matsuoka et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(20):9586-9590; and Guevara-Garcia et al. (1993) *Plant J.* 4(3):495-505. Such promoters can be modified, if necessary, for weak expression.

[0115] Leaf-preferred promoters are known in the art. See, for example, Yamamoto et al. (1997) *Plant J.* 12(2):255-265; Kwon et al. (1994) *Plant Physiol.* 105:357-67; Yamamoto et al. (1994) *Plant Cell Physiol.* 35(5):773-778; Gotor et al. (1993) *Plant J.* 3:509-18; Orozco et al. (1993) *Plant Mol. Biol.* 23(6):1129-1138; and Matsuoka et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(20):9586-9590. In addition, the promoters of cab and ribisco can also be used. See, for example, Simpson et al. (1958) *EMBO J* 4:2723-2729 and Timko et al. (1988) *Nature* 318:57-58.

[0116] Root-preferred promoters are known and can be selected from the many available from the literature or isolated de novo from various compatible species. See, for example, Hire et al. (1992) *Plant Mol. Biol.* 20(2):207-218 (soybean root-specific glutamine synthetase gene); Keller and Baumgartner (1991) *Plant Cell* 3(10):1051-1061 (root-

specific control element in the GRP 1.8 gene of French bean); Sanger et al. (1990) *Plant Mol. Biol.* 14(3):433-443 (root-specific promoter of the mannopine synthase (MAS) gene of *Agrobacterium tumefaciens*); and Miao et al. (1991) *Plant Cell* 3(1):11-22 (full-length cDNA clone encoding cytosolic glutamine synthetase (GS), which is expressed in roots and root nodules of soybean). See also Bogusz et al. (1990) *Plant Cell* 2(7):633-641, where two root-specific promoters isolated from hemoglobin genes from the nitrogen-fixing nonlegume *Parasponia andersonii* and the related non-nitrogen-fixing nonlegume *Trema tomentosa* are described. The promoters of these genes were linked to a β -glucuronidase reporter gene and introduced into both the nonlegume *Nicotiana tabacum* and the legume *Lotus corniculatus*, and in both instances root-specific promoter activity was preserved. Leach and Aoyagi (1991) describe their analysis of the promoters of the highly expressed roIC and roID root-inducing genes of *Agrobacterium rhizogenes* (see *Plant Science* (Limerick) 79(1):69-76). They concluded that enhancer and tissue-preferred DNA determinants are dissociated in those promoters. Teeri et al. (1989) used gene fusion to lacZ to show that the *Agrobacterium* T-DNA gene encoding octopine synthase is especially active in the epidermis of the root tip and that the TR2' gene is root specific in the intact plant and stimulated by wounding in leaf tissue, an especially desirable combination of characteristics for use with an insecticidal or larvicidal gene (see *EMBO J.* 8(2):343-350). The TR1' gene, fused to nptII (neomycin phosphotransferase II) showed similar characteristics. Additional root-preferred promoters include the VtENOD-GRP3 gene promoter (Kuster et al. (1995) *Plant Mol. Biol.* 29(4):759-772); and roIB promoter (Capana et al. (1994) *Plant Mol. Biol.* 25(4):681-691. See also U.S. Pat. Nos. 5,837,876; 5,750,386; 5,633,363; 5,459,252; 5,401,836; 5,110,732; and 5,023,179. The phaseolin gene (Murai et al. (1983) *Science* 23:476-482 and Sengopta-Gopalén et al. (1988) *PNAS* 82:3320-3324).

[0117] Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. Suitable methods of introducing the DNA construct include microinjection (Crossway et al. (1986) *Biotechniques* 4:320-334; and U.S. Pat. No. 6,300,543), sexual crossing, electroporation (Riggs et al. (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606), *Agrobacterium*-mediated transformation (Townsend et al., U.S. Pat. No. 5,563,055; and U.S. Pat. No. 5,981,840), direct gene transfer (Paszowski et al. (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford et al., U.S. Pat. No. 4,945,050; Tomes et al., U.S. Pat. No. 5,879,918; Tomes et al., U.S. Pat. No. 5,886,244; Bidney et al., U.S. Pat. No. 5,932,782; Tomes et al. (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe et al. (1988) *Biotechnology* 6:923-926). Also see Weissinger et al. (1988) *Ann. Rev. Genet.* 22:421-477; Sanford et al. (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou et al. (1988) *Plant Physiol.* 87:671-674 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh et al. (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta et al. (1990) *Biotechnology* 8:736-740 (rice); Klein et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize);

Klein et al. (1988) *Biotechnology* 6:559-563 (maize); Tomes, U.S. Pat. No. 5,240,855; Buising et al., U.S. Pat. Nos. 5,322,783 and 5,324,646; Klein et al. (1988) *Plant Physiol.* 91:440-444 (maize); Fromm et al. (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren et al. (1984) *Nature* (London) 311:763-764; Bowen et al., U.S. Pat. No. 5,736,369 (cereals); Bytebier et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet et al. (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman et al. (Longman, New York), pp. 197-209 (pollen); Kaeppler et al. (1990) *Plant Cell Reports* 9:415-418 and Kaeppler et al. (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin et al. (1992) *Plant Cell* 4:1495-1505 (electroporation); Li et al. (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda et al. (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); and U.S. Pat. No. 5,736,369 (meristem transformation), all of which are herein incorporated by reference.

[0118] The nucleotide constructs may be introduced into plants by contacting plants with a virus or viral nucleic acids. Generally, such methods involve incorporating a nucleotide construct of the invention within a viral DNA or RNA molecule. Further, it is recognized that useful promoters encompass promoters utilized for transcription by viral RNA polymerases. Methods for introducing nucleotide constructs into plants and expressing a protein encoded therein, involving viral DNA or RNA molecules, are known in the art. See, for example, U.S. Pat. Nos. 5,889,191, 5,889,190, 5,866,785, 5,589,367 and 5,316,931; herein incorporated by reference.

[0119] In some embodiments, transient expression may be desired. In those cases, standard transient transformation techniques may be used. Such methods include, but are not limited to viral transformation methods, and microinjection of DNA or RNA, as well other methods well known in the art.

[0120] The cells from the plants that have stably incorporated the nucleotide sequence may be grown into plants in accordance with conventional ways. See, for example, McCormick et al. (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic imparted by the nucleotide sequence of interest and/or the genetic markers contained within the target site or transfer cassette. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the desired phenotypic characteristic has been achieved.

[0121] Initial identification and selection of cells and/or plants comprising the DNA constructs may be facilitated by the use of marker genes. Gene targeting can be performed without selection if there is a sensitive method for identifying recombinants, for example if the targeted gene modification can be easily detected by PCR analysis, or if it results in a certain phenotype. However, in most cases, identification of gene targeting events will be facilitated by the use of markers. Useful markers include positive and negative selectable markers as well as markers that facilitate

screening, such as visual markers. Selectable markers include genes carrying resistance to an antibiotic such as spectinomycin (e.g. the *aada* gene, Svab et al. 1990 *Plant Mol. Biol.* 14:197), streptomycin (e.g., *aada*, or SPT, Svab et al. 1990 *Plant Mol. Biol.* 14:197; Jones et al. 1987 *Mol. Gen. Genet.* 210:86), kanamycin (e.g., *nptII*, Fraley et al. 1983 *PNAS* 80:4803), hygromycin (e.g., HPT, Vanden Elzen et al. 1985 *Plant Mol. Biol.* 5:299), gentamycin (Hayford et al. 1988 *Plant Physiol.* 86:1216), phleomycin, zeocin, or bleomycin (Hille et al. 1986 *Plant Mol. Biol.* 7:171), or resistance to a herbicide such as phosphinothricin (*bar* gene), or sulfonylurea (acetolactate synthase (*ALS*)) (Charest et al. (1990) *Plant Cell Rep.* 8:643), genes that fulfill a growth requirement on an incomplete media such as *HIS3*, *LEU2*, *URA3*, *LYS2*, and *TRP1* genes in yeast, and other such genes known in the art. Negative selectable markers include cytosine deaminase (*codA*) (Stougaard 1993 *Plant J.* 3:755-761), *tms2* (DePicker et al. 1988 *Plant Cell Rep.* 7:63-66), nitrate reductase (Nussame et al. 1991 *Plant J.* 1:267-274), *SU1* (O'Keefe et al. 1994 *Plant Physiol.* 105:473-482), *aux-2* from the Ti plasmid of *Agrobacterium*, and thymidine kinase. Screenable markers include fluorescent proteins such as green fluorescent protein (*GFP*) (Chalfie et al., 1994 *Science* 263:802; U.S. Pat. No. 6,146,826; U.S. Pat. No. 5,491,084; and WO 97/41228), reporter enzymes such as β -glucuronidase (*GUS*) (Jefferson R. A. 1987 *Plant Mol. Biol. Rep.* 5:387; U.S. Pat. No. 5,599,670; and U.S. Pat. No. 5,432,081), β -galactosidase (*lacZ*), alkaline phosphatase (*AP*), glutathione S-transferase (*GST*) and luciferase (U.S. Pat. No. 5,674,713; and Ow et al. 1986 *Science* 234(4778):856-859), visual markers like anthocyanins such as *CRC* (Ludwig et al. (1990) *Science* 247(4841):449-450) *R* gene family (e.g. *Lc*, *P*, *S*), *A*, *C*, *R-nj*, *body* and/or eye color genes in *Drosophila*, coat color genes in mammalian systems, and others known in the art.

[0122] One or more markers may be used in order to select and screen for gene targeting events. One common strategy for gene disruption involves using a target modifying polynucleotide in which the target is disrupted by a promoterless selectable marker. Since the selectable marker lacks a promoter, random integration events are unlikely to lead to transcription of the gene. Gene targeting events will put the selectable marker under control of the promoter for the target gene. Gene targeting events are identified by selection for expression of the selectable marker. Another common strategy utilizes a positive-negative selection scheme. This scheme utilizes two selectable markers, one that confers resistance (*R+*) coupled with one that confers a sensitivity (*S+*), each with a promoter. When this polynucleotide is randomly inserted, the resulting phenotype is *R+/S+*. When a gene targeting event is generated, the two markers are uncoupled and the resulting phenotype is *R+/S-*. Examples of using positive-negative selection are found in Thykjaer et al. (1997) *Plant Mol. Biol.* 35:523-530; and WO 01/66717, which are herein incorporated by reference.

EXAMPLES

[0123] The following are non-limiting examples intended to illustrate the invention. Although the present invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

Example 1

The Example Describes the Identification of a microRNA

[0124] The following experiments were carried out on the *Arabidopsis thaliana* Col-0 ecotype. Plants were grown in long days (16 h light, 8 h dark) under cool white light at 22° C.

[0125] *Arabidopsis* plants were transformed by a modified version of the floral dip method, in which *Agrobacterium* cell suspension was applied to plants by direct watering from above. The T-DNA vector used, pHSbarENDs, contained four copies of the CAMV 35S enhancer adjacent to the right border, an arrangement similar to that described by Weigel et al. (*Plant Physiol.* 122:1003-1013, 2000). Transformed plants were selected with glufosinate (BASTA) and screened for flowering time, which resulted in the identification of the early-flowering EAT-D mutant. A single T-DNA cosegregating with early flowering was identified in EAT-D, and TAIL-PCR was performed to amplify sequences adjacent to the left and right borders of the T-DNA. To identify transcripts upregulated in the EAT-D mutant, we probed Northern blots containing RNA extracted from wild type (Col-0) and EAT-D plants. Probes for the genes At5g04270 and At5g04280 (GenBank NC_003076) did not detect any difference between wild type and EAT-D, whereas a probe from the intergenic region identified an ~1.4 kb transcript that was expressed at significantly higher levels in EAT-D than in wild type.

[0126] To isolate the full-length EAT cDNA, we performed 5'- and 3'-RACE-PCR with a GeneRacer kit (Invitrogen) that selects for 5'-capped miRNAs. Reverse transcription was carried out using an oligo-dt primer, and PCR utilized a gene-specific primer (SEQ ID NO: 45 5'-CTGT-GCTCAGATCTTGTGTTCTTGTATC-3') paired with the 5' kit primer, or a second gene-specific primer (SEQ ID NO: 46 5'-GTCGGCGGATCCATGGMGAAGCTCATC-5') paired with the 3' kit primer.

[0127] The *Arabidopsis* EAT-D (Early Activation Tagged-Dominant) mutant was identified in an activation tagging population (Weigel et al., *Plant Physiol.* 122:1003-1013, 2000). As evidenced by visual inspection and by measuring rosette leaf number (Table 1), the EAT-D mutant flowers extremely early. In addition, EAT-D displays floral defects that are virtually identical to those observed for strong *apetala2* (*ap2*) mutant alleles (Bowman et al., *Development* 112:1-20, 1991), including the complete absence of petals and the transformation of sepals to carpels. This *ap2*-like phenotype is only observed in EAT-D homozygotes, whereas both EAT-D heterozygotes and homozygotes are early flowering, indicating that the flowering time phenotype is more sensitive to EAT-D dosage than the *ap2*-like floral phenotype.

TABLE 1

Rosette leaf numbers for <i>Arabidopsis</i> lines		
Genotype	rosette leaf no.	floral phenotype
Col-0	11.4 +/- 1.2	wild type
EAT-D	3.1 +/- 0.8	<i>ap2</i>
EAT-OX	2.0 +/- 0.2	<i>ap2</i> + additional
eatdel	11.1 +/- 1.1	wild type
miR172a1-OX	2.1 +/- 0.3	<i>ap2</i> + additional

TABLE 1-continued

Rosette leaf numbers for <i>Arabidopsis</i> lines		
Genotype	rosette leaf no.	floral phenotype
LAT-D	22.5 +/- 2.1	wild type
At2g28550-OX	28.6 +/- 3.6	wild type
5-60120	10.2 +/- 1.4	wild type
2-28550	8.7 +/- 0.6	wild type
5-60120; 2-28550	6.0 +/- 0.8	wild type

[0128] We mapped the activation-tagged T-DNA insert in EAT-D to chromosome 5, in between the annotated genes At5g04270 and At5g04280. We then used 5'- and 3'-RACE PCR with primers located within this region to identify a 1.4 kb transcript (SEQ ID NO: 1), which we named EAT, that is upregulated in EAT-D. When the 1.4 kb EAT cDNA was fused to the constitutive CAMV 35S promoter and the resultant 35S::EAT construct was introduced into wild type (Col-0) plants by *Agrobacterium*-mediated transformation (Clough and Bent, *Plant J.* 16:735-743 1998), the 35S::EAT transformants displayed the identical early-flowering and ap2-like phenotypes seen for EAT-D (Table 1). Many of the 35S::EAT transformants occasionally displayed additional defects, including stigmatic papillae on cauline leaf margins and the formation of a complete or partial flower rather than a secondary inflorescence in the axils of cauline leaves. Ectopic expression of the EAT gene in 35S::EAT plants, therefore, affects both flowering time and the specification of floral organ identity.

[0129] The EAT gene produces a 1417-nucleotide non-coding RNA that is predicted to be 5'-capped and polyadenylated, based on our RACE-PCR methodology. BLASTN and BLASTX searches of several databases with the EAT cDNA did not reveal extensive nucleotide or predicted amino acid sequence identity between EAT and any other gene. We did, however, identify a 21-nucleotide (nt) (SEQ ID NO: 4) stretch in the middle of the EAT transcript that is identical to miR172a-2, a recently identified miRNA (Park et al., *Curr. Biol.* 12:1484-1495, 2002). To confirm the

wild-type and had normal flowers (Table 1), indicating that the miR172a-2 sequence is necessary to confer both the flowering time and floral organ identity phenotypes seen in EAT-overexpressing lines.

[0130] As noted by Park et al. (*Curr. Biol.* 12:1484-1495, 2002), the 21-nt miR172a-2 miRNA has the potential to form an RNA duplex with a sequence near the 3' end of the coding region of AP2 (Table 2).

TABLE 2

Putative 21-nt miR172a-2/AP2 RNA duplex		
Sequence	Duplex	SEQ ID NO:
AP2 RNA	5' -CUGCAGCAUCA <u>CAGGAU</u> UCU-3'	47
EAT miRNA	3' -UACGUCGUAGUAGU <u>CUA</u> AGA-5'	48

The GU wobble in the duplex is underlined.

[0131] This particular region of the AP2 gene is poorly conserved at the nucleotide level among the AP2 family; nevertheless, the AP2 sequence (SEQ ID NO: 49) that is complementary to miR172a-2 is found in a similar location in three other *Arabidopsis* AP2 family members, At5g60120 (SEQ ID NO: 50), At2g28550 (SEQ ID NO: 51), At5g67180 (SEQ ID NO: 52). In addition, the sequence can be found at the corresponding positions of the maize AP2 genes indeterminate spikelet1 (Chuck et al., *Genes. Dev.* 12:1145-1154 1998) (IDS1 (SEQ ID NO: 53)) and glossy15 (Moose and Sisco, *Genes. Dev.* 10:3018-3027 1996) (GL15 (SEQ ID NO: 54)), and in AP2 family members from many other plant species, including soybean, rice, wheat, tomato and pea (not shown). The alignment of three *Arabidopsis* and two maize AP2 family members is shown in Table 3 below.

TABLE 3

Alignment of AP2 21-nt region (black bar) and surrounding sequence	
AP2	ACCAAGTGTGAGAAATGCTGCAGCATCATCAGGATTCTCTGCTCATCATCACAATCAG
At5g60120	CACCGCCACTGTTTTCAAATGCAGCATCATCAGGATTCTCACTGTCTAGCTACACGCCCT
At2g28550	CACCATTTGTTCTCAGTTGCAGCAGCATCATGAGGATTCTCATTTCGCGCCACAACCT
At5g67180	GAAATCGAGTGGTGGGAATGGCAGGATGATCAGGATTCTCTCTCAACCTTCCCGTTAC
IDS1	ACGTGCCGTTGCACCACTCTGCAGCATGATCAGGATTCTCTACCGCCGCGGGGCCAAC
GL15	ACGCCAGCAGCGCCGCGCTGCAGCATGATCAGGATTCTCCACTGTGGCAGCTGGGTGCG

functional importance of the miR172a-2 sequence within the EAT cDNA, we generated a mutant form of EAT in which the miR172a-2 sequence was deleted, and made a construct consisting of this mutant EAT cDNA, eatdel, driven by the 35S promoter. Transgenic plants carrying this 35S::eatdel construct flowered with the same number of leaves as

[0132] There is an additional copy of the miR172a-2 miRNA in the *Arabidopsis* genome on chromosome 2 (miR172a-1, FIG. 2d), and miR172a-2 is highly similar to three other *Arabidopsis* loci. Like the miR172a-2 miRNA, all four reiterations of the sequence are in intergenic regions, i.e. in between the *Arabidopsis* genes currently annotated in

GenBank. In addition, the sequence is found in ESTs from tomato, potato and soybean, and four copies were found in the genomic sequence of rice.

Example 2

This Example Describes the Construction of Expression Vectors

[0133] To overexpress the EAT gene, we designed primers containing XhoI sites (SEQ ID NO: 55 5'-GACTACTC-GAGCACCTCTCACTCCCTTTCTCTAAC-3' and SEQ ID NO: 56 5'-GACTACTCGAGGTTCTCAAGTTGAC-CACTTGAAAAC-3') to amplify the entire EAT gene from Col-0 DNA. The PCR product was digested with XhoI and inserted into a modified pBluescriptSK+ vector (Stratagene, La Jolla, Calif.) that lacked BamHI and HindIII sites, to generate EATX4 (SEQ ID NO: 44). To generate the 35S::EAT transformants, the XhoI-cut EAT gene was inserted into the binary vector pBE851 in between a CAMV 35S promoter and b-phaseolin terminator, and Col-0 was transformed by floral dip. To generate the eatdel construct, two oligonucleotides were synthesized (SEQ ID NO: 57 5'-GATCCATGGAAGAAAGCTCATCTGTCGT-TGTTTGTAGGCGCAGCACCATTAAAGA TTCACATG-GAAATTGATAAATAC-3' and SEQ ID NO: 58 5'-CCTAAATTAGGGTTTGTATATGATATAT-TCAACAATCGACGGCTACAAATACCTAA-3') that completely recreated the BamHI/HindIII fragment of the EAT cDNA except that it lacked the 21 nt miR172a-2 sequence located within the fragment. These two oligos were annealed to their synthesized complementary strands (SEQ ID NO: 59 5'-TAGGGTATTTATCAATTTCATGT-GAATCTTAATGGTGCTGCGCTACAAACAAC GACAGATGAGCTTTCTTCCATG-3' and SEQ ID NO: 60 5'-AGCTTTAGGTATTTGTAGCCGTCGAT-TGTTGAATATACATATCAAAACCCTAATT-3') and ligated to EATX4 that had been digested with BamHI and HindIII, in a trimolecular ligation reaction. This resulted in the replacement of 159 bp of wild-type EAT sequence with the 138 bp mutant sequence. The eatdel cDNA was then subcloned into pBE851 and transformed as described above. BASTA was used to select in plants for both the EAT and eatdel overexpression constructs.

[0134] To test whether another member of the miR172 family, miR172a-1, would confer a phenotype similar to that of miR172a-2, we generated a construct containing the 35S promoter fused to the genomic region surrounding miR172a-1. Plants containing the 35S::miR172a-1 construct flowered early and displayed an ap2 phenotype (Table 1), indicating that miR172a-1 behaves in an identical manner to miR172a-2 when overexpressed.

[0135] All of the miR172 miRNA family members are located within a sequence context that allows an RNA hairpin to form (FIG. 1). Presumably this hairpin is the substrate which is subsequently cleaved by a plant Dicer homolog to generate the mature miRNA. The location of the miRNA within the hairpin, i.e. on the 3' side of the stem, is conserved amongst all the members of the miR172 family, and this may reflect a structural requirement for processing of this particular miRNA family. The 21-nt miR172a-2 miRNA, therefore, is predicted to be a member of a family of miRNAs that have the capacity to regulate a subset of AP2 genes by forming an RNA duplex with a 21-nt cognate sequence in these genes.

Example 3

The Example Describes the Analysis of microRNA Expression and AP2 Expression

[0136] Total RNA was isolated from wild type and EAT-D whole plants that had already flowered, using TRIzol reagent (Sigma). 50 mg of each RNA was subjected to electrophoresis on a 15% TBE-Urea Criterion gel (BioRad), electroblotted onto Hybond-N+ filter paper (Amersham) using a TransBlot-SD apparatus (BioRad). The filter was then hybridized at 37° C. overnight in UltraHyb-Oligo buffer (Ambion) with 32P-labeled oligos. The oligos were 30-mers that corresponded to either the sense or antisense strands of the miR172a-2 miRNA, with 4-5 nt of flanking sequence on each side. The filter was washed twice at 37° C., in buffer containing 2×SSC and 0.5% SDS. For S1 analysis, probe was made by end-labeling an oligo (SEQ ID NO: 61) (5'-ATGCAGCATCATCMGATTCTCATATACAT-3') with T4 polynucleotide kinase and 32P. Hybridization and processing of S1 reactions were carried out using standard protocols. For developmental analysis of miR172a-2 and miR172a-1, total RNA was isolated from plants at the various stages and tissues indicated in Example 4, using an Rneasy kit (Qiagen). RT-PCR was carried out using standard protocols, and utilized oligos specific for sequences adjacent to miR172a-2 (SEQ ID NO: 62) (5'-GTCGGCGGATC-CATGGAAGAAAGCTCATC-3' and (SEQ ID NO: 63) 5'-CAAAGATCGATCCAGACTTCAATCAATATC-3') or sequences adjacent to miR172a-1 (SEQ ID NO: 64) (5'-TAATTTCCGGAGCCACGGTCGTTGTTG-3' and (SEQ ID NO: 65) 5'-AATAGTCGTTGATTGCCGATGCAG-CATC-3'). Oligos used to amplify the ACT11 (Actin) transcript were: (SEQ ID NO: 66) 5'-ATGGCAGATGGTGAA-GACATTCAG-3', and (SEQ ID NO: 67) 5'-GAAGCACTTCCTGTGGACTATTGATG-3'. RT-PCR analysis of AP2 was performed on RNA from floral buds, and utilized the following oligos: (SEQ ID NO: 68) 5'-TTTCCGGGCAGCAGCAACATTGGTAG-3', and (SEQ ID NO: 69) 5'-GTTTCGCTAAGTTAACAAGAGGATT-TAGG-3'. Oligos used to amplify the ANT transcript were: (SEQ ID NO: 70) 5'-GATCAACTTCAATGAC-TAACTCTGGTTTTC-3', and (SEQ ID NO: 71) 5'-GT-TATAGAGAGATTCAATCTGTTTCACATG-3'.

[0137] Immunoblot analysis of AP2 was performed on proteins extracted from floral buds. Following electrophoresis on a 10% SDS-PAGE gel, proteins were transferred to a Hybond-P membrane (Amersham) and incubated with an antibody specific for AP2 protein (aA-20, Santa Cruz Biotechnology). The blot was processed using an ECL-plus kit (Amersham).

[0138] Northern analysis using probes both sense and antisense to the miR172a-2 miRNA identified a small single-stranded RNA of 21-25 nucleotides accumulating too much higher levels in EAT-D mutant plants relative to wild type. The small amount of transcript seen in wild type presumably represents endogenous levels of not only the miR172a-2 miRNA but also its family members, which are similar enough to cross-hybridize with the probe. The predicted miR172a-2 hairpin is 117 nt in length (FIG. 1), a small amount of an ~100 nt transcript accumulating is detected in EAT-D, this likely represents partially processed miR172a-2 hairpin precursor. S1 nuclease mapping of the miR172a-2

miRNA provides independent confirmation of the 5' end of miR172a-2 reported by Park et al. (*Curr. Biol.* 12:1484-1495, 2002).

Example 4

The Example Describes the Developmental Pattern of EAT miRNA Expression.

[0139] To address the wild-type expression pattern of miR172a-2 separate from its other *Arabidopsis* family members, RT-PCR was used to specifically detect a fragment of the 1.4 kb EAT full-length precursor transcript containing miR172a-2. EAT precursor transcript expression is temporally regulated, with little or no transcript detected two days after germination, and progressively more steady-state transcript accumulation seen as the plant approaches flowering. The precursor transcript of miR172a-1 showed a similar temporal pattern of expression. Both miR172a-2 and miR172a-1 precursor transcripts continue to be expressed after flowering has occurred, and accumulate in both leaves and floral buds. We were unable to detect expression of the precursors for the other miR172 family members, perhaps due to their exclusive expression in tissue types not included in this analysis, or because their precursor transcripts are too transient to detect. The temporal expression pattern seen for miR172a-2 and miR172a-1 is reminiscent of that observed for let-7 and lin-4, two miRNAs that control developmental timing in *C. elegans* (Feinbaum and Ambros, *Dev. Biol.* 210:87-95 1999; Reinhart et al., *Nature* 403:901-906 2000).

Example 5

[0140] We assessed the levels of miR172 in various flowering time mutants, in an attempt to position miR172 within the known flowering time pathways. The levels of miR172 were not altered in any of the mutants tested, and the levels of the EAT transcript were identical in plants grown in long days versus plants grown in short days.

Example 6

The Example Describes Evaluation of Protein Expression

[0141] Immunoblot analysis indicates that AP2 protein is reduced 3.5-fold in the EAT-D mutant relative to wild type, whereas the AP2 transcript is unaffected. This data suggests that the miR172a-2 miRNA negatively regulates AP2 by translational inhibition. The predicted near-perfect complementarity between the miR172a-2 miRNA and the AP2 target site would be predicted to trigger AP2 miRNA cleavage by the RNA interference (RNAi) pathway (Llave et al., *Plant Cell* 14:1605-1619 2002; Hutvagner and Zamore, *Science* 297:2056-2060 2002). Indeed, others have proposed that many plant miRNAs enter the RNAi pathway exclusively due to their near-perfect complementarity to putative targets (Rhoades et al., *Cell* 110:513-520 2002). While there is no evidence regarding the GU wobble base pair in the predicted miR172a-2/AP2 RNA duplex, it is conserved in all predicted duplexes between miR172 family members and their AP2 targets. Regardless of the mechanism, it is apparent from the AP2 expression data and the observed phenotype of EAT-D that AP2 is a target of negative regulation by miR172a-2, at least when miR172a-2 is overexpressed.

Example 7

[0142] In the same genetic screen that identified the early-flowering EAT-D mutant, we identified an activation-tagged

late-flowering mutant, called LAT-D. The LAT-D mutant displays no additional phenotypes besides late flowering (Table 1), and the late-flowering phenotype cosegregated with a single T-DNA insertion. Sequence analysis of the T-DNA insert in LAT-D indicated that the 4x35S enhancer was located approximately 5 kb upstream of At2g28550, which is one of the AP2-like target genes that are potentially regulated by miR172. RT-PCR analysis using primers specific for At2g28550 indicates that the transcript corresponding to this gene is indeed expressed at higher levels in the LAT-D mutant relative to wild type. To confirm that overexpression of At2g28550 causes late flowering, we fused a genomic region containing the entire At2g28550 coding region (from start to stop codon) to the 35S promoter, and created transgenic plants containing this construct. Transgenic 35S::At2g28550 plants flowered later than wild type plants, and were slightly later than the LAT-D mutant (Table 1). This late flowering phenotype was observed in multiple independent transformants.

[0143] The fact that overexpression of At2g28550 causes late flowering suggests that miR172 promotes flowering in part by downregulating At2g28550. However, because miR172 appears to affect protein rather than transcript accumulation of its target genes, and because we do not have an antibody to the At2g28550 gene product, we decided to test this regulation indirectly via a genetic cross. A plant heterozygous for LAT-D was crossed to a plant homozygous for EAT-D, such that all F1 progeny would contain one copy of EAT-D and 50% of the F1 progeny would also have one copy of LAT-D. F1 progeny were scored for the presence or absence of the LAT-D allele by PCR, and also were scored for flowering time. All of the F1 plants were early flowering, regardless of whether or not they contained a copy of the LAT-D allele, indicating that EAT-D is epistatic to LAT-D. This result is consistent with the idea that miR172a-2, which is overexpressed in EAT-D, directly downregulates At2g28550, which is overexpressed in LAT-D.

Example 8

[0144] To assess the effects of reducing At2g28550 function, we identified plants containing a T-DNA insertion in the At2g28550 gene. In addition, we identified a T-DNA mutant for At2g60120, a closely related AP2-like gene that also contains the miR172 target sequence. Plants homozygous for either the At2g28550 insert or the At5g60120 insert were slightly early flowering relative to wild type (Table 1). The two mutants were crossed, and the double mutant was isolated by PCR genotyping. The At2g28550/At5g60120 double mutant was earlier flowering than either individual mutant (Table 1), suggesting that the genes have overlapping function. The early flowering phenotype of the At2g28550/At5g60120 double mutant is consistent with the idea that the early flowering phenotype of miR172-overexpressing lines is due to downregulation of several AP2-like genes, including At2g28550 and At5g60120. Interestingly, the At2g28550/At5g60120 double mutant is not as early as miR172-overexpressing lines (c.f. EAT-OX, Table 1), which suggests that other AP2-like targets of miR172, for example AP2 itself or At5g67180, also contribute to flowering time control. Because ap2 mutants are not early flowering, any potential negative regulation of flowering by AP2 must be normally masked by genetic redundancy.

Example 9

[0145] This Example Describes a Method of Target Selection and Method to Design DNA Constructs to Generate miRNAs Using the Constructs of SEQ ID NOS: 3 and 44. Any Sequence of Interest can be Selected for Silencing by miRNA Generated Using the Following Method:

[0146] 1. Choose a region from the coding strand in a gene of interest to be the target sequence. Typically, choose a region of about 10-50 nucleotides found in a similar location to the region targeted by EAT in AP2-like genes, which are regions about 100 nt upstream of the stop codon. The exact location of the target, however, does not appear to be critical. It is recommended to choose a region that has ~50% GC and is of high sequence complexity, i.e. no repeats or long polynucleotide tracts. It is also recommended that the chosen region ends with a T or A, such that the complementary miRNA will start with an A or U. This is to help ensure a lower stability at the 5' end of the miRNA in its double-stranded Dicer product form (Schwartz, et al. 2003 Cell 115:199-208). For example, in the miR172a-2 precursor, the miRNA sequence starts with an A, and many other miRNAs start with a U.

[0147] 2. To use the construct of SEQ ID NO: 3, create a 21 nucleotide sequence complementary to the 21 nt target region (miRNA). Optionally, change a C in the miRNA to a T, which will generate a GU wobble with the target sequence, which mimics the GU wobble seen in EAT.

[0148] 3. Create the 21 nucleotide "backside" sequence of the hairpin. This will be substantially complementary to the miRNA from step 2. Note, this backside sequence will also be substantially identical to the target sequence. Typically, introduce a few mismatches to make some bulges in the stem of the hairpin that are similar to the bulges in the original EAT hairpin. Optionally, introduce an A at the 3' end of the backside, to create mismatch at the 5' end of the miRNA. This last step may help ensure lower stability at the 5' end of the miRNA in its double-stranded Dicer product form (Schwartz, et al. 2003 Cell 115:199-208).

[0149] 4. Replace the 21 nucleotide miRNA sequence and the 21 nucleotide "backside" sequence in the EAT BamHI/HindIII DNA construct (SEQ ID NO: 3) with the new miRNA and "backside" sequences from steps 2 and 3.

[0150] 5. Use MFOLD (GCG, Accelrys, San Diego, Calif.), or an equivalent program, to compare the new hairpin from Step 4 with the original hairpin. Generally, the sequence substantially replicate the structure of the original hairpin (FIG. 1). It is predicted that the introduced bulges need not be exactly identical in length, sequence or position to the original. Examine the miRNA sequence in the hairpin for the relative stability of the 5' and 3' ends of the predicted dsRNA product of Dicer.

[0151] 6. Generate four synthetic oligonucleotides of 76-77 nucleotides in length to produce two double-stranded fragments which comprise the BamHI and HindIII restriction sites, and a 4 nucleotide overhang to facilitate directional ligation which will recreate the BamHI/HindIII fragment. Design of the overhang can be done by one of skill in the art, the current example uses the 4 nucleotide region of

positions 79-82 (CCTA) of SEQ ID NO: 3. Hence, for example:

[0152] Oligo 1 will have an unpaired BamHI site at the 5' end, and will end with the nucleotide at position 78 of SEQ ID NO: 3.

[0153] Oligo 2 will have the nucleotides of position 79-82 (CCTA) unpaired at the 5' end, and will terminate just before the HindIII site (or positions 151-154 in SEQ ID NO: 3).

[0154] Oligo 3 will be essentially complementary to Oligo 1, (nucleotides 5-78 of SEQ ID NO: 3), and will terminate with 4 nucleotides complementary to nucleotides 1-4 (CCTA) of Oligo 2.

[0155] Oligo 4 will be essentially complementary to Oligo 2 beginning at the nucleotide of position 5, and will terminate with the HindIII site at the 3' end.

[0156] Anneal the oligonucleotides to generate two fragments to be used in a subsequent ligation reaction with the plasmid sequence.

[0157] 7. Ligate the two DNA fragments from Step 6 in a trimolecular ligation reaction with a plasmid cut with BamHI/HindIII. The current example uses the modified pBluescript SK+ plasmid of SEQ ID NO: 44, which comprises the 1.4 kb EAT sequence of SEQ ID NO: 1, digested with BamHI/HindIII and gel purified away from the small fragment using standard molecular biological techniques. The new designed miRNA to the gene of interest has replaced the previous miRNA.

[0158] 8. The plasmid from Step 7, subject to any other preparations or modifications as needed, is used to transform the target organism using techniques appropriate for the target.

[0159] 9. Silencing of the target gene can be assessed using techniques well-known in the art, for example, Northern blot analysis, immunoblot analysis if the target gene of interest encodes a polypeptide, and any phenotypic screens relevant to the target gene, for example flowering time, or floral morphology.

Example 10

[0160] Described in this example are methods one may use for introduction of a polynucleotide or polypeptide into a plant cell.

A. Maize Particle-Mediated DNA Delivery

[0161] A DNA construct can be introduced into maize cells capable of growth on suitable maize culture medium. Such competent cells can be from maize suspension culture, callus culture on solid medium, freshly isolated immature embryos or meristem cells. Immature embryos of the Hi-II genotype can be used as the target cells. Ears are harvested at approximately 10 days post-pollination, and 1.2-1.5 mm immature embryos are isolated from the kernels, and placed scutellum-side down on maize culture medium.

[0162] The immature embryos are bombarded from 18-72 hours after being harvested from the ear. Between 6 and 18 hours prior to bombardment, the immature embryos are placed on medium with additional osmoticum (MS basal medium, Musashige and Skoog, 1962, *Physiol. Plant* 15:473-497, with 0.25 M sorbitol). The embryos on the

high-osmotic medium are used as the bombardment target, and are left on this medium for an additional 18 hours after bombardment.

[0163] For particle bombardment, plasmid DNA (described above) is precipitated onto 1.8 mm tungsten particles using standard CaCl_2 -spermidine chemistry (see, for example, Klein et al., 1987, *Nature* 327:70-73). Each plate is bombarded once at 600 PSI, using a DuPont Helium Gun (Lowe et al., 1995, *Bio/Technol* 13:677-682). For typical media formulations used for maize immature embryo isolation, callus initiation, callus proliferation and regeneration of plants, see Armstrong, C., 1994, In "The Maize Handbook", M. Freeling and V. Walbot, eds. Springer Verlag, NY, pp 663-671.

[0164] Within 1-7 days after particle bombardment, the embryos are moved onto N6-based culture medium containing 3 mg/l of the selective agent bialaphos. Embryos, and later callus, are transferred to fresh selection plates every 2 weeks. The calli developing from the immature embryos are screened for the desired phenotype. After 6-8 weeks, transformed calli are recovered.

B. Soybean Transformation

[0165] Soybean embryogenic suspension cultures are maintained in 35 ml liquid media SB196 or SB172 in 250 ml Erlenmeyer flasks on a rotary shaker, 150 rpm, 26°C with cool white fluorescent lights on 16:8 hr day/night photoperiod at light intensity of 30-35 $\mu\text{E}/\text{m}^2\text{s}$. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 ml of fresh liquid media. Alternatively, cultures are initiated and maintained in 6-well Costar plates.

[0166] SB 172 media is prepared as follows: (per liter), 1 bottle Murashige and Skoog Medium (Duchefa # M 0240), 1 ml B5 vitamins 1000 $\times$ stock, 1 ml 2,4-D stock (Gibco 11215-019), 60 g sucrose, 2 g MES, 0.667 g L-Asparagine anhydrous (GibcoBRL 11013-026), pH 5.7. SB 196 media is prepared as follows: (per liter) 1.0 ml MS FeEDTA, 10 ml MS Sulfate, 10 ml FN-Lite Halides, 10 ml FN-Lite P, B, Mo, 1 ml B5 vitamins 1000 $\times$ stock, 1 ml 2,4-D, (Gibco 11215-019), 2.83g KNO_3 , 0.463 g $(\text{NH}_4)_2\text{SO}_4$, 2g MES, 1 g Asparagine Anhydrous, Powder (Gibco 11013-026), 10 g Sucrose, pH 5.8. 2,4-D stock concentration 10 mg/ml is prepared as follows: 2,4-D is solubilized in 0.1 N NaOH, filter-sterilized, and stored at -20°C . B5 vitamins 1000 $\times$ stock is prepared as follows: (per 100 ml)-store aliquots at -20°C , 10 g myo-inositol, 100 mg nicotinic acid, 100 mg pyridoxine HCl, 1 g thiamin.

[0167] Soybean embryogenic suspension cultures are transformed with various plasmids by the method of particle gun bombardment (Klein et al., 1987 *Nature* 327:70). To prepare tissue for bombardment, approximately two flasks of suspension culture tissue that has had approximately 1 to 2 weeks to recover since its most recent subculture is placed in a sterile 60 $\times$ 20 mm petri dish containing 1 sterile filter paper in the bottom to help absorb moisture. Tissue (i.e. suspension clusters approximately 3-5 mm in size) is spread evenly across each petri plate. Residual liquid is removed from the tissue with a pipette, or allowed to evaporate to remove excess moisture prior to bombardment. Per experiment, 4-6 plates of tissue are bombarded. Each plate is made from two flasks.

[0168] To prepare gold particles for bombardment, 30 mg gold is washed in ethanol, centrifuged and resuspended in

0.5 ml of sterile water. For each plasmid combination (treatments) to be used for bombardment, a separate microcentrifuge tube is prepared, starting with 50 μl of the gold particles prepared above. Into each tube, the following are also added; 5 μl of plasmid DNA (at 1 $\mu\text{g}/\mu\text{l}$), 50 μl CaCl_2 , and 20 μl 0.1 M spermidine. This mixture is agitated on a vortex shaker for 3 minutes, and then centrifuged using a microcentrifuge set at 14,000 RPM for 10 seconds. The supernatant is decanted and the gold particles with attached, precipitated DNA are washed twice with 400 μl aliquots of ethanol (with a brief centrifugation as above between each washing). The final volume of 100% ethanol per each tube is adjusted to 40 μl , and this particle/DNA suspension is kept on ice until being used for bombardment.

[0169] Immediately before applying the particle/DNA suspension, the tube is briefly dipped into a sonicator bath to disperse the particles, and then 5 μl of DNA prep is pipetted onto each flying disk and allowed to dry. The flying disk is then placed into the DuPont Biolistics PDS1000/HE. Using the DuPont Biolistic PDS1000/HE instrument for particle-mediated DNA delivery into soybean suspension clusters, the following settings are used. The membrane rupture pressure is 1100 psi. The chamber is evacuated to a vacuum of 27-28 inches of mercury. The tissue is placed approximately 3.5 inches from the retaining/stopping screen (3rd shelf from the bottom). Each plate is bombarded twice, and the tissue clusters are rearranged using a sterile spatula between shots.

[0170] Following bombardment, the tissue is re-suspended in liquid culture medium, each plate being divided between 2 flasks with fresh SB196 or SB172 media and cultured as described above. Four to seven days post-bombardment, the medium is replaced with fresh medium containing a selection agent. The selection media is refreshed weekly for 4 weeks and once again at 6 weeks. Weekly replacement after 4 weeks may be necessary if cell density and media turbidity is high.

[0171] Four to eight weeks post-bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated, green tissue is removed and inoculated into 6-well microtiter plates with liquid medium to generate clonally-propagated, transformed embryogenic suspension cultures.

[0172] Each embryogenic cluster is placed into one well of a Costar 6-well plate with 5 mls fresh SB196 media with selection agent. Cultures are maintained for 2-6 weeks with fresh media changes every 2 weeks. When enough tissue is available, a portion of surviving transformed clones are subcultured to a second 6-well plate as a back-up to protect against contamination.

[0173] To promote in vitro maturation, transformed embryogenic clusters are removed from liquid SB196 and placed on solid agar media, SB 166, for 2 weeks. Tissue clumps of 2-4 mm size are plated at a tissue density of 10 to 15 clusters per plate. Plates are incubated in diffuse, low light ($<10 \mu\text{E}$) at $26\pm 1^\circ\text{C}$. After two weeks, clusters are subcultured to SB 103 media for 3-4 weeks.

[0174] SB 166 is prepared as follows: (per liter), 1 pkg. MS salts (Gibco/BRL-Cat# 111117-017), 1 ml B5 vitamins 1000 $\times$ stock, 60 g maltose, 750 mg MgCl_2 hexahydrate, 5 g activated charcoal, pH 5.7, 2 g gelrite. SB 103 media is

prepared as follows: (per liter), 1 pkg. MS salts (Gibco/BRL-Cat# 11117-017), 1 ml B5 vitamins 1000× stock, 60 g maltose, 750 mg MgCl₂ hexahydrate, pH 5.7, 2 g gelrite. After 5-6 week maturation, individual embryos are desiccated by placing embryos into a 100×15 petri dish with a 1 cm² portion of the SB103 media to create a chamber with enough humidity to promote partial desiccation, but not death.

[0175] Approximately 25 embryos are desiccated per plate. Plates are sealed with several layers of parafilm and again are placed in a lower light condition. The duration of the desiccation step is best determined empirically, and depends on size and quantity of embryos placed per plate. For example, small embryos or few embryos/plate require a shorter drying period, while large embryos or many embryos/plate require a longer drying period. It is best to check on the embryos after about 3 days, but proper desiccation will most likely take 5 to 7 days. Embryos will decrease in size during this process.

[0176] Desiccated embryos are planted in SB 71-1 or MSO medium where they are left to germinate under the same culture conditions described for the suspension cultures. When the plantlets have two fully-expanded trifoliate leaves, germinated and rooted embryos are transferred to sterile soil and watered with MS fertilizer. Plants are grown to maturity for seed collection and analysis. Healthy, fertile transgenic plants are grown in the greenhouse.

[0177] SB 71-1 is prepared as follows: 1 bottle Gamborg's B5 salts w/sucrose (Gibco/BRL-Cat# 21153-036), 10 g sucrose, 750 mg MgCl₂ hexahydrate, pH 5.7, 2 g gelrite. MSO media is prepared as follows: 1 pkg Murashige and Skoog salts (Gibco 11117-066), 1 ml B5 vitamins 1000× stock, 30 g sucrose, pH 5.8, 2g Gelrite.

Example 11

[0178] This example describes the design and synthesis of miRNA targets and hairpins directed to various gene targets found in maize, soy, and/or *Arabidopsis*, using the method described in Example 9.

A. Targeting *Arabidopsis* AGAMOUS, At4g18960

[0179] The miRNA sequence of SEQ ID NO: 4 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 12-15, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

[0180] *Arabidopsis thaliana* Col-0 was transformed and grown as described in Example 1. After transformation with a vector comprising the miRNA of SEQ ID NO: 4, 88% of the transformants exhibited a mutant AGAMOUS (ag) floral phenotype, characterized by the conversion of stamens to petals in whorl 3, and carpels to another ag flower in whorl 4 (Bowman, et al. (1991) *The Plant Cell* 3:749-758). The mutant phenotype varied between transformants, with approximately 1/3 exhibiting a strong ag phenotype, 1/3 exhibiting an intermediate ag phenotype, and 1/3 exhibiting a weak ag phenotype. Gel electrophoresis and Northern Blot analysis of small RNAs isolated from the transformants demonstrated that the degree of the mutant ag phenotype was directly related to the level of antiAG miRNA, with the strongest phenotype having the highest accumulation of the processed miRNA (~21 nt).

B. Targeting *Arabidopsis* Apetela3 (AP3), At3g54340

Two miRNA Targets from AP3 were Selected and Oligonucleotides Designed.

[0181] The miRNA sequence of SEQ ID NO: 5 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 16-19, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

[0182] The miRNA sequence of SEQ ID NO: 6 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 20-23, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

[0183] *Arabidopsis thaliana* Col-0 was transformed and grown as described in Example 1. After transformation with a vector comprising the miRNA of SEQ ID NO: 5, the transformants had novel leaf and floral phenotypes, but did not exhibit any mutant AP3 phenotype. Gel electrophoresis and Northern analysis of RNA isolated from 2 week old rosette leaf tissue from the transformants demonstrated that the highest accumulation of the processed miRNA (~21 nt) corresponded to the "backside" strand of the precursor, which evidently silenced a different target sequence to produce the novel leaf and floral phenotypes.

[0184] A new target sequence was selected, with the correct asymmetry in order for the miRNA target strand to be selected during incorporation into RISC (Schwartz et al., 2003, *Cell* 115:199-208). The miRNA sequence of SEQ ID NO: 6 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 20-23, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44. Greater than 90% of the transformants showed silencing for the AP3 gene, as demonstrated by floral phenotype and electrophoretic analysis. An approximately 21 nt miRNA (antiAP3b) was detected at high levels in the transgenic plants, and not in wild type control plants. RT-PCR analysis confirmed that the amount of AP3 transcript was reduced in the transformants, as compared to wild type control plants.

C. Targeting Maize Phytoene Desaturase

Two miRNA Targets from Phytoene Desaturase (PDS) were Selected and Oligonucleotides Designed.

[0185] The miRNA sequence of SEQ ID NO: 7 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 24-27, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

[0186] The miRNA sequence of SEQ ID NO: 8 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 28-31, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

D. Targeting Maize Phytic Acid Biosynthetic Enzymes

[0187] Three maize phytic acid biosynthetic enzyme gene targets were selected and miRNA and oligonucleotides designed. Inositol polyphosphate kinase-2 polynucleotides are disclosed in WO 02/059324, herein incorporated by reference. Inositol 1,3,4-trisphosphate 5/6-kinase polynucle-

otides are disclosed in WO 03/027243, herein incorporated by reference. Myo-inositol 1-phosphate synthase polynucleotides are disclosed in WO 99/05298, herein incorporated by reference.

Inositol Polyphosphate Kinase-2 (IPPK2)

[0188] The miRNA sequence of SEQ ID NO: 9 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 32-35, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

Inositol 1,3,4-trisphosphate 5/6-kinase-5 (ITPK5)

[0189] The miRNA sequence of SEQ ID NO: 10 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 36-39, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

Myo-Inositol 1-Phosphate Synthase (mi1 ps)

[0190] The miRNA sequence of SEQ ID NO: 11 was selected and designed. The sequence is put into the BamHI/HindIII hairpin cassette by annealing the synthetic oligonucleotides of SEQ ID NOS: 40-43, and ligating them into the BamHI/HindIII backbone fragment of SEQ ID NO: 44.

E. Targeting Soy Apetela2-Like Sequences (AP2)

[0191] The same EAT (miR172a-2) construct, comprising SEQ ID NO: 1, used for *Arabidopsis* transformation was used to transform soybean. This construct has a miRNA template sequence which encodes the miRNA of SEQ ID NO: 48. The construct was created using a PCR amplification of miR172a-2 precursor sequence from *Arabidopsis*, restriction digestion, and ligation as described in Example 2.

[0192] Soybean tissue was transformed and grown essentially as described in Example 10. After transformation, 42% of the transformants exhibited a mutant phenotype, characterized by the conversion of sepals to leaves. Plants exhibiting the strongest phenotypes were sterile, and produced no seed. Both the homeotic conversion of the organs and the effects on fertility are similar to that seen for ap2 mutant alleles in *Arabidopsis*. Small RNA gel electrophoresis and Northern analysis, probed with an oligonucleotide probe antisense to miR172, showed accumulation of miR172 in

the transgenic lines. A small amount of endogenous soy miR172 is also detected in the soy control line. The degree of the mutant phenotype was directly related to the level of miRNA, with the strongest phenotype having the highest accumulation of the processed miRNA (~21 nt).

Example 12

[0193] This example summarizes the target sequences and oligos used for miRNA silencing constructs as described in the examples.

TABLE 4

Organism	Target gene	miRNA name	miRNA/ template	Precursor oligos SEQ ID NOS
<i>Arabidopsis</i>	AP2-like	miR172-a2	SEQ ID NO: 47	55-56 (PCR)
	none	EATdel	none	57-60
	AGAMOUS	antiAG	SEQ ID NO: 4	12-15
	APETELA3	antiAP3a	SEQ ID NO: 5	16-19
	(a)			
Corn	APETELA3	antiAP3b	SEQ ID NO: 6	20-23
	(b)			
	PDS1	antiPDS1	SEQ ID NO: 7	24-27
	PDS2	antiPDS1	SEQ ID NO: 8	28-31
	IPPK2	antiIPPK2	SEQ ID NO: 9	32-35
Soybean	ITPK5	antiITPK5	SEQ ID NO: 10	36-39
	MI1PS	antiMI1PS	SEQ ID NO: 11	40-43
	AP2-like	miR172a-2	SEQ ID NO: 47	55-56 (PCR)

[0194] The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, patents, patent applications, and computer programs cited herein are hereby incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 71

<210> SEQ ID NO 1

<211> LENGTH: 1417

<212> TYPE: DNA

<213> ORGANISM: *Arabidopsis thaliana*

<400> SEQUENCE: 1

```
gcacctctca ctccctttct ctaactagtc ttgtgtgcac ccatttatgt gtacgtacta      60
ttatctcata aataaatatt tttaaaatta gatgcattta ttgatatgaa aaagttacaa      120
gattagtttg ttgtgtgtga gactttggat cgacagatcg aaaaattaac taaccggtca      180
gtattgaata tcaactatta tatgtccat gcattcgctt atagtttcac acaatttggt      240
```

-continued

```

ttcttcacgg tctaaatca gaagattcca tatatcttct tatgacgtaa aaggaccact 300
tataagttga cagctcagcc cttggattcg tgagggtttt ctctctactt cacctatcta 360
cttttcctca tatcccactg cttttctcct tcttggttctt gtttttctcg ttttttctt 420
cttcttctcc aagaaaaatag agatcgaaaa gattagatct attttgtgta gcaagaaatt 480
atcattttcg tttcttcatt catatattgt tctattatgt tgtacaataa tagatactcg 540
atctcttggt cgtgcgtaaa ttttatacaa gttgtcggcg gatccatgga agaaagctca 600
tctgtcgttg tttgtaggcg cagcaccatt aagattcaca tggaaattga taaataccct 660
aaattagggg tttgatattg atatgagaat cttgatgatg ctgcatcaac aatcgacggc 720
tacaataacc taaagcttga gaaagaaact tgaagatatt gattgaagtc tggatcgatc 780
tttggtaaat ctctctcttg attagtttta agaatactt ttttttttct gtgtttgaac 840
atgtttacat atatcatcta tgtctcaata tatatatttt cttaacttag ggtcaatgac 900
ggattagggc gtaattaca atgaatatgg aaaaactatt ttgccttga tcttgacttg 960
agtgttgatg aacagatgta taatgttatg tagtatgtac tgtatttttt ctagaatcat 1020
tctttagtct ccaactctcc attaatcaaa tgaggtcctt ataggtaatg ctatgatcaa 1080
gaacaacaag atcgtgagca cagatcggcc agttcgggtca ctttttaaaa gagagatggt 1140
atattgttaa tttgttatta tcaggataaa taaatacaga atagttcgtc cagagaccag 1200
acattttata gtttcaattt tatgacagtc ttgtaataat atttgtttaa tagtgtgtca 1260
ccttctatct ctgggttatt acttggtccc gaaattttct tattgttcta attttgtaat 1320
attagaaatt tggttttctt gccaaatcaa atcaaacatt acggtgtggt gtacattgta 1380
ccagaacttt tgttttcaag tgctcaactt gagaacc 1417

```

```

<210> SEQ ID NO 2
<211> LENGTH: 95
<212> TYPE: DNA
<213> ORGANISM: Arabidopsis thaliana

```

```

<400> SEQUENCE: 2

```

```

aggcgagca ccattaagat tcacatggaa attgataaat accctaaatt agggttttga 60
tatgtatatg agaactttga tgatgctgca tcaac 95

```

```

<210> SEQ ID NO 3
<211> LENGTH: 159
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template cassette

```

```

<400> SEQUENCE: 3

```

```

ggatccatgg aagaaagctc atctgtcgtt gtttgtaggc gcagcaccat taagattcac 60
atggaaattg ataaataccc taaattaggg ttttgatatg tatatgagaa tcttgatgat 120
gtgcatcaa caatcgacgg ctacaaatc ctaaagctt 159

```

```

<210> SEQ ID NO 4
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target At4g18960

```

-continued

<400> SEQUENCE: 4
taggttgtaa tgccgcgact t 21

<210> SEQ ID NO 5
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target At3g54340

<400> SEQUENCE: 5
ggtggaaatg aagagcgtaa g 21

<210> SEQ ID NO 6
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target At3g54340

<400> SEQUENCE: 6
agagcgtaag cacgtgaccc t 21

<210> SEQ ID NO 7
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target maize phytoene
desaturase

<400> SEQUENCE: 7
tgctggcaga agtccgattg c 21

<210> SEQ ID NO 8
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target maize phytoene
desaturase

<400> SEQUENCE: 8
agcttcctgg ataggactgc a 21

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target is maize IPPK2

<400> SEQUENCE: 9
aagttgtggt taatcacccc a 21

<210> SEQ ID NO 10
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target is maize ITPK5

<400> SEQUENCE: 10

-continued

gaggacagtt tcgtatcctg g 21

<210> SEQ ID NO 11
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: miRNA template to target is maize Milps3

<400> SEQUENCE: 11

gagcgtttac caccggtgtg c 21

<210> SEQ ID NO 12
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo1/4 for At4g18960 target

<400> SEQUENCE: 12

gatccatgga agaaagctca tctgtcgttg tttgtaggca gtcgcggcac tacaacaaa 60

tggaattga taaatac 77

<210> SEQ ID NO 13
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo2/4 for At4g18960 target

<400> SEQUENCE: 13

tagggtatatt atcaatttcc atttggttgt agtgccgcga ctgcctacaa acaacgacag 60

atgagctttc ttccatg 77

<210> SEQ ID NO 14
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo3/4 for At4g18960 target

<400> SEQUENCE: 14

cctaaattag ggttttgata tgtatattag gttgtaatgc cgcgactttc aacaatcgac 60

ggctacaaat acctaa 76

<210> SEQ ID NO 15
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo4/4 for At4g18960 target

<400> SEQUENCE: 15

agcttttaggt atttgtagcc gtcgattgtt gaaagtcgcg gcattacaac ctaatatata 60

tatcaaaaacc ctaatt 76

<210> SEQ ID NO 16
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo1/4 for At3g54340 target

<400> SEQUENCE: 16

gatccatgga agaaagctca tctgtcgttg tttgtaggat tacgcccttc attaccacca      60
tggaattga taaatac                                                    77

<210> SEQ ID NO 17
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo2/4 for At3g54340 target

<400> SEQUENCE: 17

tagggatattt atcaatttcc atggtggtaa tgaagggcgt aatcctacaa acaacgacag      60
atgagctttc ttccatg                                                    77

<210> SEQ ID NO 18
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo3/4 for At3g54340 target

<400> SEQUENCE: 18

cctaatttag ggttttgata tgtatatggt ggaaatgaag agcgtaagtc aacaatcgac      60
ggctacaaat acctaa                                                    76

<210> SEQ ID NO 19
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo4/4 for At3g54340 target

<400> SEQUENCE: 19

agcttttaggt attttagtcc gtcgattgtt gacttacgct cttcatttcc accatataca      60
tatcaaaacc ctaatt                                                    76

<210> SEQ ID NO 20
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo1/4 for At3g54340 target

<400> SEQUENCE: 20

gatccatgga agaaagctca tctgtcgttg tttgtaggag gtcacgcgct tacgctcaca      60
tggaattga taaatac                                                    77

<210> SEQ ID NO 21
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo2/4 for At3g54340 target

<400> SEQUENCE: 21

tagggatattt atcaatttcc atgtgagcgt aagcgcgtga ccgcctacaa acaacgacag      60

```

-continued

atgagctttc ttccatg 77

<210> SEQ ID NO 22
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligo3/4 for At3g54340 target
 <400> SEQUENCE: 22

cctaaattag ggttttgata tgtatatgag agcgtaagca cgtgaccctc aacaatcgac 60

ggctacaaat acctaa 76

<210> SEQ ID NO 23
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligo4/4 for At3g54340 target
 <400> SEQUENCE: 23

agcttttaggt atttgtagcc gtcgattgtt gagggtcacg tgcttacgct ctcatataca 60

tatcaaaaacc ctaatt 76

<210> SEQ ID NO 24
 <211> LENGTH: 77
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligo1/4 for phytoene desaturase target
 <400> SEQUENCE: 24

gatccatgga agaaagctca tctgtcgttg tttgtaggca atcggacttc tgccagcaca 60

tggaattga taaatac 77

<210> SEQ ID NO 25
 <211> LENGTH: 77
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligo2/4 for phytoene desaturase target
 <400> SEQUENCE: 25

tagggatattt atcaatttcc atgtgctggc agaagtcgga ttgcctacaa acaacgacag 60

atgagctttc ttccatg 77

<210> SEQ ID NO 26
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligo3/4 for phytoene desaturase target
 <400> SEQUENCE: 26

cctaaattag ggttttgata tgtatatgtg ctggcagaag tccgattgcc aacaatcgac 60

ggctacaaat acctaa 76

```
<210> SEQ ID NO 31
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo4/4 for phytoene desaturase
        target

<400> SEQUENCE: 31

agcttttaggt atttgtagcc gtcgattggt gtgcagtcct atccaggaag ctcatatata
60

tatcaaaaacc ctaatt
76
```

-continued

<210> SEQ ID NO 32
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo1/4 for IPPK2 target

<400> SEQUENCE: 32

gatccatgga agaaagctca tctgtcgttg tttgtaggcg gggtgataaa ccacaacata 60
tggaattga taaatac 77

<210> SEQ ID NO 33
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo2/4 for IPPK2 target

<400> SEQUENCE: 33

tagggtatatt atcaatttcc atatgttgtg gtttatcacc ccgcctacaa acaacgacag 60
atgagctttc ttccatg 77

<210> SEQ ID NO 34
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo3/4 for IPPK2 target

<400> SEQUENCE: 34

cctaaattag ggttttgata tgtatataag ttgtggttaa tcaccccatc acaaatcgac 60
ggctacaaat acctaa 76

<210> SEQ ID NO 35
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo4/4 for IPPK2 target

<400> SEQUENCE: 35

agcttttaggt attttagtacc gtcgattgtt gatggggtga ttaaccacaa cttatatata 60
tatcaaaacc ctaatt 76

<210> SEQ ID NO 36
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo1/4 for ITPK5 target

<400> SEQUENCE: 36

gatccatgga agaaagctca tctgtcgttg tttgtaggac aggatacgta actgtccaca 60
tggaattga taaatac 77

<210> SEQ ID NO 37
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo2/4 for ITPK5 target

-continued

<400> SEQUENCE: 37

taggggtatatt atcaattttcc atgtggacag ttacgtatcc tgtcctacaa acaacgacag 60

atgagctttc ttccatg 77

<210> SEQ ID NO 38

<211> LENGTH: 76

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligo3/4 for ITPK5 target

<400> SEQUENCE: 38

cctaaattag ggttttgata tgtatatgag gacagtttcg taccctggtc aacaatcgac 60

ggctacaaat acctaa 76

<210> SEQ ID NO 39

<211> LENGTH: 76

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligo4/4 for ITPK5 target

<400> SEQUENCE: 39

agcttttaggt attttagtgc gtcgattgtt gaccaggata cgaaactgtc ctcatatata 60

tatcaaaaacc ctaatt 76

<210> SEQ ID NO 40

<211> LENGTH: 77

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligo1/4 for milps target

<400> SEQUENCE: 40

gatccatgga agaaagctca tctgtcgttg ttgttaggac acaccggcgg taaacgcaca 60

tggaattga taaatac 77

<210> SEQ ID NO 41

<211> LENGTH: 77

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligo2/4 for milps target

<400> SEQUENCE: 41

taggggtatatt atcaattttcc atgtgcgttt accgccggtg tgtcctacaa acaacgacag 60

atgagctttc ttccatg 77

<210> SEQ ID NO 42

<211> LENGTH: 76

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligo3/4 for milps target

<400> SEQUENCE: 42

cctaaattag ggttttgata tgtatatgag cgtttaccac cgggtgtgctc aacaatcgac 60

ggctacaaat acctaa 76

-continued

<210> SEQ ID NO 43
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligo4/4 for milps target

<400> SEQUENCE: 43

agcttttaggt attttagtacc gtcgattgtt gagcacaccg gtggtaaacg ctcatataca 60
tatcaaaacc ctaatt 76

<210> SEQ ID NO 44
<211> LENGTH: 4426
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Plasmid

<400> SEQUENCE: 44

ctaaattgta agcgtaata ttttgttaaa attcgcgtta aatttttgtt aaatcagctc 60
attttttaac caataggccg aaatcggcaa aatcccttat aaatcaaaag aatagaccga 120
gatagggttg agtgtgttgc cagtttggaa caagagtcca ctattaaaga acgtggactc 180
caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg aaccatcacc 240
ctaatcaagt tttttggggt cgaggtgccg taaagcacta aatcggaacc cttaaaggag 300
cccccgaatt agagcttgac ggggaaagcc ggcgaacgtg gcgagaaagg aagggaagaa 360
agcgaaagga gcggcgctga gggcgctggc aagtgtagcg gtcacgctgc gcgtaaccac 420
cacaccgcc gcgcttaatg cgccgctaca gggcgctgcc cattcgccat tcaggctgcg 480
caactgttgg gaaggcgcat cgggtcgggc ctcttcgcta ttacgccagc tggcgaaagg 540
gggatgtgct gcaaggcgat taagttgggt aacgccaggg ttttcccagt cagcagcttg 600
taaaacgacg gccagtgcgc gcgcgtaata cgactcacta tagggcgaat tgggtaccgg 660
gccctctaga tgcattgctg agcggccgcc agtgtgatgg atatctgcag aattcgcctt 720
tgactactcg agcacctctc actcccttct tctaactagt cttgtgtgca cccatttatg 780
tgtactgtat attatctcat aaataaatat ttttaaaatt agatgcattt attgatatga 840
aaaagttaca agattagttt gttgtgtgtg agactttgga tcgacagatc gaaaaattaa 900
ctaaccggtc agtattgaat atcaactatt atatgctcca tgcattcgct tatagtttca 960
cacattttgt tttcttcacg gtctaaaatc agaagattcc atatattttc ttatgacgta 1020
aaaggaccac ttataagttg acacgtcagc ccttgatttc gtgaggtttt tctctctact 1080
tcacctatct acttttcttc atatccactt gcttttctcc ttcttgttct tgtttttctc 1140
gtttttttct tcttcttctc caagaaaata gagatcgaaa agattagatc tattttgtgt 1200
agcaagaaat tatcattttc gtttcttcat tcataatttg ttctattatg ttgtacaata 1260
atagatactc gatctcttgt gcgtgcgtaa attttatata agttgtcggc ggatccatgg 1320
aagaaagctc atctgtcgtt gttttagagg gcagcaccat taagattcac atggaaattg 1380
ataaatacc ctaaataggg ttttgatatg tatatgagaa tcttgatgat gctgcatcaa 1440
caatcgacgg ctacaaatcc cttaaagctt agaaagaaac ttgaagatat tgattgaagt 1500
ctggatcgat ctttggtaaa tctctctctt gattagtttt aagaatcact tttttttttc 1560

-continued

tgtgtttgaa catgtttaca tatatcatct atgtctcaat atatataatt tcttaattcta	1620
gggtcaatga cggattaggg cgtaattac aatgaatatg gaaaaactat ttgaccttg	1680
atcttgactt gagtgtgat gaacagatgt ataattgtat gtagtatgta ctgtattttt	1740
tctagaatca ttcttttagtc tccaactctc cattaatcaa atgaggtcct tataggtaat	1800
gctatgatca agaacaacaa gatcgtgagc acagatcggc cagttcggtc actttttaaa	1860
agagagatgt tatattgtta atttggtatt atcaggtata ataaatacag aatagttcgt	1920
ccagagacca gacattttat agtttcaatt ttatgacagt cttgtaataa tattttgtta	1980
atagtgtgtc accttctatt tctgggttat tacttggtcc cgaaattttc ttattgttct	2040
aattttgtaa tattagaaat ttggttttct tgccaaatca aatcaaact tacggtgtgt	2100
tgtacattgt accagaactt ttgttttcaa gtgctcaact tgagaacctc gagtagtcaa	2160
ggcggaattc cagcacactg gcggccgtta ctagtcttag agcgccgcc accgcggtg	2220
agctccagct ttgttccct ttagtgaggg ttaattgcgc gcttgcgta atcatggtca	2280
tagctgtttc ctgtgtgaaa ttgttatccg ctcaaatc cacaacat acgagccgga	2340
agcataaagt gtaaagcctg gggtgcctaa tgagttagct aactcacatt aattgcgtt	2400
cgctcactgc ccgctttcca gtcgggaaac ctgctgtgcc agctgcatta atgaatcggc	2460
caacgcgcgg gagagggcg ttgctgtatt gggcgctctt ccgcttctc gctcactgac	2520
tcgtgcgct cggtcgttcg gctgcggcga gcggtatcag ctactcaaa ggcggtaata	2580
cggttatcca cagaatcagg ggataacgca ggaagaaca tgtgagcaa aggccagcaa	2640
aaggccagga accgtaaaaa ggcgcggtt ctggcggttt tccataggct ccgccccct	2700
gacgagcatc aaaaaatcg acgctcaagt cagaggtggc gaaacccgac aggactataa	2760
agataccagg cgtttccccc tggaagctcc ctctgcgct ctctgttcc gacctgccc	2820
cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgcttc tcatagctca	2880
cgctgtagggt atctcagttc ggtgtaggtc gttcgtcca agctgggctg tgtgcacgaa	2940
cccccggtc agcccgaccg ctgcgcctta tccgtaact atcgtcttga gtccaaccg	3000
gtaagacacg acttatcgcc actggcagca gccactggt acaggattag cagagcgagg	3060
tatgtaggcg gtgtacaga gttcttgaag tgggtgccta actacggcta cactagaagg	3120
acagtatttg gtatctgcgc tctgtgaag ccagttacct tcggaaaaag agttggtagc	3180
tcttgatccg gaaaaaaac caccgctggt agcgggtggt tttttgttg caagcagcag	3240
attacgcga gaaaaaagg atctcaagaa gatcctttga tctttctac ggggtctgac	3300
gctcagtga acgaaaactc acgttaaggg attttggta tgagattatc aaaaaggatc	3360
ttcacctaga tcctttttaa ttaaaatga agttttaaat caatctaaag tatatatgag	3420
taaacttggt ctgacagtta ccaatgctta atcagttagg cacctatctc agcgatctgt	3480
ctatttcgtt catccatagt tgcctgactc ccgctcgtg agataactac gatacgggag	3540
ggcttaccat ctggccccag tgctgcaatg ataccgcgag acccacgctc accggctcca	3600
gatttatcag caataaacca gccagccgga agggccgagc gcagaagtgg tcctgcaact	3660
ttatccgcct ccatccagtc tattaattgt tgccgggaag ctagagtaag tagttcgcca	3720
gttaatagtt tgcgcaacgt tgttgccatt gctacaggca tcgtggtgtc acgctcgtcg	3780
tttggtatgg cttcattcag ctccggttcc caacgatcaa ggcgagttac atgatcccc	3840

-continued

```

atgtttgtgca aaaaagcggg tagctccttc ggtcctccga tcgttgcag aagtaagttg 3900
gccgcagtggt tatcactcat gggtatggca gcactgcata attctcttac tgcatgcca 3960
tccgtaagat gcttttctgt gactggtgag tactcaacca agtcattctg agaatagtgt 4020
atgcgcgcgac cgagttgctc ttgcccgcg tcaatacggg ataataccgc gccacatagc 4080
agaactttaa aagtgtcat cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc 4140
ttaccgctgt tgagatccag ttcgatgtaa cccactcgtg cacccaactg atcttcagca 4200
tcttttactt tcaccagcgt ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa 4260
aagggaataa gggcgacacg gaaatgttga atactcatac tcttcctttt tcaatattat 4320
tgaagcattt atcagggtta ttgtctcatg agcggataca tatttgaatg tatttagaaa 4380
aataaaca aa taggggttcc gcgcacattt ccccgaaaag tgccac 4426

```

```

<210> SEQ ID NO 45
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EAT 3' PCR primer

```

<400> SEQUENCE: 45

```

ctgtgctcac gatcttgttg ttcttgatc 29

```

```

<210> SEQ ID NO 46
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EAT 5' PCR primer

```

<400> SEQUENCE: 46

```

gtcggcggat ccatggaaga aagctcatc 29

```

```

<210> SEQ ID NO 47
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Arabidopsis thaliana
<220> FEATURE:
<223> OTHER INFORMATION: AP2 RNA

```

<400> SEQUENCE: 47

```

cugcagcauc aucaggauuc u 21

```

```

<210> SEQ ID NO 48
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Arabidopsis thaliana
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (0)...(0)
<223> OTHER INFORMATION: EAT miRNA

```

<400> SEQUENCE: 48

```

agaaucuuga ugaugcugca u 21

```

```

<210> SEQ ID NO 49
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Arabidopsis thaliana

```

-continued

<400> SEQUENCE: 49

accaagtgtt gacaaatgct gcagcatcat caggattctc tcctcatcat cacaatcag 59

<210> SEQ ID NO 50

<211> LENGTH: 59

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 50

caccgccact gttttcaaat gcagcatcat caggattctc actctcagct acacgccct 59

<210> SEQ ID NO 51

<211> LENGTH: 59

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 51

caccattgtt ctcatgtgca gcagcatcat caggattctc acatttcggt ccacaacct 59

<210> SEQ ID NO 52

<211> LENGTH: 59

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 52

gaaatcgagt ggtgggaatg gcagcatcat caggattctc tcctcaacct tccccttac 59

<210> SEQ ID NO 53

<211> LENGTH: 59

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<400> SEQUENCE: 53

acgtgccgtt gcaccactct gcagcatcat caggattctc taccgccgcc ggggccaac 59

<210> SEQ ID NO 54

<211> LENGTH: 59

<212> TYPE: DNA

<213> ORGANISM: Zea mays

<400> SEQUENCE: 54

acgccagcag cgccgccgct gcagcatcat caggattccc actgtggcag ctgggtgcg 59

<210> SEQ ID NO 55

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: EAT PCR primer

<400> SEQUENCE: 55

gactactcga gcacctotca ctccctttct ctaac 35

<210> SEQ ID NO 56

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: EAT PCR primer

<400> SEQUENCE: 56

-continued

gactactcga gggtctcaag ttgagcactt gaaaac 36

<210> SEQ ID NO 57
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EAT deletion oligonucleotide

<400> SEQUENCE: 57

gatccatgga agaaagctca tctgtcgttg tttgtaggcg cagcaccatt aagattcaca 60

tggaattga taaatac 77

<210> SEQ ID NO 58
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EAT deletion oligonucleotide

<400> SEQUENCE: 58

cctaaattag gggtttgata tgtatattca acaatcgacg gctacaaata cctaa 55

<210> SEQ ID NO 59
<211> LENGTH: 77
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EAT deletion oligonucleotide

<400> SEQUENCE: 59

tagggatatt atcaatttcc atgtgaatct taatgggtgct ggcctacaa acaacgacag 60

atgagctttc ttccatg 77

<210> SEQ ID NO 60
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EAT deletion oligonucleotide

<400> SEQUENCE: 60

agcttttaggt attttagacc gtcgattgtt gaatatacat atcaaaaccc taatt 55

<210> SEQ ID NO 61
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: S1 probe

<400> SEQUENCE: 61

atgcagcatc atcaagattc tcatatacat 30

<210> SEQ ID NO 62
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mir172a-2 PCR primer

<400> SEQUENCE: 62

-continued

gtcggcggat ccatggaaga aagtcatac 29

<210> SEQ ID NO 63
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mir172a-2 PCR primer

<400> SEQUENCE: 63

caaagatcga tccagacttc aatcaatatac 30

<210> SEQ ID NO 64
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mir172a-1 PCR primer

<400> SEQUENCE: 64

taatttccgg agccacggtc gttgttg 27

<210> SEQ ID NO 65
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mir172a-1 PCR primer

<400> SEQUENCE: 65

aatagtcggt gattgccgat gcagcatc 28

<210> SEQ ID NO 66
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Actin PCR primer

<400> SEQUENCE: 66

atggcagatg gtgaagacat tcag 24

<210> SEQ ID NO 67
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Actin PCR primer

<400> SEQUENCE: 67

gaagcacttc ctgtggacta ttgatg 26

<210> SEQ ID NO 68
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AP2 PCR primer

<400> SEQUENCE: 68

tttccgggca gcagcaacat tggtag 26

-continued

```

<210> SEQ ID NO 69
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AP2 PCR primer

```

```

<400> SEQUENCE: 69

```

```

gttcgcctaa gttaacaaga ggatttagg

```

29

```

<210> SEQ ID NO 70
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: ANT PCR primer

```

```

<400> SEQUENCE: 70

```

```

gatcaacttc aatgactaac tctggttttc

```

30

```

<210> SEQ ID NO 71
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: ANT PCR primer

```

```

<400> SEQUENCE: 71

```

```

gttatagaga gattcattct gtttcacatg

```

30

What is claimed:

1. A method of inhibiting expression of a target sequence in a cell comprising:

(a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide, the polynucleotide comprising in the following order:

- (i) at least about 20 contiguous nucleotides in the region from nucleotides 1-38 of SEQ ID NO: 3,
- (ii) a first oligonucleotide of 10 to about 50 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide,
- (iii) at least about 20 contiguous nucleotides in the region from nucleotides 60-106 of SEQ ID NO: 3,
- (iv) the second oligonucleotide of about 10 to about 50 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and wherein the second oligonucleotide is substantially complementary to the target sequence, and
- (v) at least about 20 contiguous nucleotides in the region from nucleotides 128-159 of SEQ ID NO: 3;

wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin, and

(b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

2. The method of claim 1, wherein the nucleic acid construct further comprises a promoter operably linked to the polynucleotide

3. The method of claim 1, wherein the cell is a plant cell.

4. A cell comprising the nucleic acid construct of claim 1.

5. The cell of claim 4, wherein the cell is a plant cell.

6. The method of claim 1, wherein target sequence expression is inhibited by at least 10%.

7. The method of claim 1, wherein inhibition of the target sequence generates a loss-of-function phenotype.

8. The method of claim 2, wherein the promoter is a pathogen-inducible promoter and inhibition of the target sequence confers resistance to a pathogen.

9. An isolated polynucleotide comprising in the following order at least 20 contiguous nucleotides in the region from nucleotides 1-38 of SEQ ID NO: 3, a first oligonucleotide of about 10 to about 50 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide, at least about 20 contiguous nucleotides in the region from nucleotides 60-106 of SEQ ID NO: 3, a second oligonucleotide of about 10 to about 50 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and the second oligonucleotide is substantially complementary to the target sequence, and at least about 20 contiguous nucleotides in the region from nucleotides 128-159 of SEQ ID NO: 3, wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin.

10. The isolated polynucleotide of claim 9, further comprising an operably linked promoter.

11. A cell comprising the isolated polynucleotide of claim 9.

12. The cell of claim 11, wherein the cell is a plant cell.
 13. A transgenic plant comprising the isolated polynucleotide of claim 9.

14. A transgenic seed comprising the isolated polynucleotide of claim 9.

15. The isolated polynucleotide of claim 10, wherein the promoter is a pathogen-inducible promoter.

16. The isolated polynucleotide of claim 9, wherein the polynucleotide suppresses expression of a target sequence.

17. A method of inhibiting expression of a target sequence in a cell comprising:

(a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide, wherein the polynucleotide comprises in the following order:

(i) nucleotides 1-38 of SEQ ID NO: 3,

(ii) a first oligonucleotide of 21 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide,

(iii) nucleotides 60-106 of SEQ ID NO: 3,

(iv) a second oligonucleotide of 21 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and wherein the second oligonucleotide is substantially complementary to the target sequence, and

(v) nucleotides 128-159 of SEQ ID NO:3;

wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin, and

(b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

18. An isolated polynucleotide comprising in the following order nucleotides 1-38 of SEQ ID NO: 3, a first oligonucleotide of 21 contiguous nucleotides, wherein the first oligonucleotide is substantially complementary to a second oligonucleotide, nucleotides 60-106 of SEQ ID NO: 3, a second oligonucleotide of 21 contiguous nucleotides, wherein the second oligonucleotide encodes a miRNA, and the second oligonucleotide is substantially complementary to the target sequence, and nucleotides 128-159 of SEQ ID NO:3, wherein the polynucleotide encodes an RNA precursor capable of forming a hairpin.

19. A method of inhibiting expression of a target sequence in a cell comprising:

(a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide encoding a modified plant miRNA precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide encodes a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin; and

(b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

20. A method of inhibiting expression of a target sequence in a cell comprising:

(a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide encoding a modified plant miR172 miRNA precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide encodes a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin; and

(b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

21. The method of claim 20, wherein the plant miR172 miRNA precursor is from a dicot or a monocot.

22. The method of claim 20, wherein the plant miR172 miRNA precursor is from *Arabidopsis*.

23. The method of claim 20, wherein the plant miR172 miRNA precursor comprises SEQ ID NO: 3.

24. A method of inhibiting expression of a target sequence in a cell comprising:

(a) introducing into the cell a nucleic acid construct comprising a promoter operably linked to a polynucleotide encoding a modified *Arabidopsis* miR172 miRNA precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide encodes a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin; and

(b) expressing the nucleic acid construct for a time sufficient to produce the miRNA, wherein the miRNA inhibits expression of the target sequence.

25. The method of claim 24, wherein the *Arabidopsis* miR172 miRNA precursor comprises SEQ ID NO: 3.

26. A cell produced by the method of claim 1.

27. The cell of claim 26, wherein the cell is a plant cell.

28. The cell of claim 27, wherein the plant cell is from a monocot or a dicot.

29. The cell of claim 28, wherein the cell is selected from the group consisting of corn, wheat, rice, barley, oats, sorghum, millet, sunflower, safflower, cotton, soy, canola, alfalfa, *Arabidopsis*, and tobacco.

30. The cell of claim 29, wherein the cell is from *Arabidopsis*.

31. The cell of claim 29, wherein the cell is from corn.

32. A cell produced by the method of claim 19.

33. The cell of claim 32, wherein the cell is a plant cell.

34. The cell of claim 33, wherein the plant cell is from a monocot or a dicot.

35. The cell of claim 34, wherein the cell is selected from the group consisting of corn, wheat, rice, barley, oats, sorghum, millet, sunflower, safflower, cotton, soy, canola, alfalfa, *Arabidopsis*, and tobacco.

36. The cell of claim 35, wherein the cell is from *Arabidopsis*.

37. The cell of claim 35, wherein the cell is from corn.

38. A cell produced by the method of claim 20.
39. The cell of claim 38, wherein the cell is a plant cell.
40. The cell of claim 39, wherein the plant cell is from a monocot or a dicot.
41. The cell of claim 40, wherein the cell is selected from the group consisting of corn, wheat, rice, barley, oats, sorghum, millet, sunflower, safflower, cotton, soy, canola, alfalfa, *Arabidopsis*, and tobacco.
42. The cell of claim 41, wherein the cell is from *Arabidopsis*.
43. The cell of claim 41, wherein the cell is from corn.
44. A cell produced by the method of claim 24.
45. The cell of claim 44, wherein the cell is a plant cell.
46. The cell of claim 45, wherein the plant cell is from a monocot or a dicot.
47. The cell of claim 46, wherein the cell is selected from the group consisting of corn, wheat, rice, barley, oats, sorghum, millet, sunflower, safflower, cotton, soy, canola, alfalfa, *Arabidopsis*, and tobacco.
48. The cell of claim 47, wherein the cell is from *Arabidopsis*.
49. The cell of claim 47, wherein the cell is from corn.
50. The method of claim 3, the method further comprising producing a transformed plant, wherein the plant comprises the nucleic acid construct which encodes the miRNA.
51. A plant produced by the method of claim 50.
52. The method of claim 19, wherein the cell is a plant cell, the method further comprising producing a transformed plant, wherein the plant comprises the nucleic acid construct which encodes the miRNA.
53. A plant produced by the method of claim 52.
54. The method of claim 20, wherein the cell is a plant cell, the method further comprising producing a transformed plant, wherein the plant comprises the nucleic acid construct which encodes the miRNA.
55. A plant produced by the method of claim 50.
56. The method of claim 24, wherein the cell is a plant cell, the method further comprising producing a transformed plant, wherein the plant comprises the nucleic acid construct which encodes the miRNA.
57. A plant produced by the method of claim 56.
58. The plant of claim 57, wherein the plant is a monocot or a dicot.
59. The plant of claim 58, wherein the plant is selected from the group consisting of corn, wheat, rice, barley, oats, sorghum, millet, sunflower, safflower, cotton, soy, canola, alfalfa, *Arabidopsis*, and tobacco.
60. The plant of claim 59, wherein the plant is *Arabidopsis*.
61. The plant of claim 59, wherein the plant is corn.
62. An isolated polynucleotide comprising a polynucleotide which encodes a modified plant miRNA precursor, the modified precursor comprising a first and a second oligo-

nucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide comprises a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin.

63. A cell comprising the isolated polynucleotide of claim 62.
64. The cell of claim 63, wherein the cell is a plant cell.
65. An isolated polynucleotide comprising a polynucleotide encoding a modified plant miR172 miRNA precursor, the modified precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide comprises a miRNA substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin.
66. A cell comprising the isolated polynucleotide of claim 65.
67. The cell of claim 66, wherein the cell is a plant cell.
68. An isolated polynucleotide comprising polynucleotide encoding a modified *Arabidopsis* miR172 miRNA precursor, the modified precursor comprising a first and a second oligonucleotide, wherein at least one of the first or the second oligonucleotides is heterologous to the precursor, wherein the first oligonucleotide is substantially complementary to the second oligonucleotide, and the second oligonucleotide comprises a miRNA which is substantially complementary to the target sequence, wherein the precursor is capable of forming a hairpin.
69. A cell comprising the isolated polynucleotide of claim 68.
70. The cell of claim 69, wherein the cell is a plant cell.
71. The cell of claim 70, wherein the plant cell is from a dicot or a monocot.
72. The cell of claim 71, wherein the cell is from *Arabidopsis*.
73. The cell of claim 71, wherein the cell is from corn.
74. The isolated polynucleotide of claim 62, wherein the first and the second oligonucleotides are heterologous to the precursor.
75. The isolated polynucleotide of claim 65, wherein the first and the second oligonucleotides are heterologous to the precursor.
76. The isolated polynucleotide of claim 68, wherein the first and the second oligonucleotides are heterologous to the precursor.

* * * * *