

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2016311429 C1**

(54) Title
Modified anthrax toxin protective antigen

(51) International Patent Classification(s)
C07K 14/325 (2006.01) **A61K 39/07** (2006.01)

(21) Application No: **2016311429** (22) Date of Filing: **2016.08.25**

(87) WIPO No: **WO17/035359**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/210,771	2015.08.27	US
62/323,218	2016.04.15	US

(43) Publication Date: **2017.03.02**

(44) Accepted Journal Date: **2021.07.08**

(44) Amended Journal Date: **2021.10.07**

(71) Applicant(s)
The United States of America, as represented by the Secretary Department of Health and Human Services

(72) Inventor(s)
Liu, Shi-Hui;Leppla, Stephen H.;Bugge, Thomas H.;Wein, Alexander N.;Peters, Diane E.;Liu, Jie;Chen, Kuang-Hua

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
WO 2014/134187 A1
CN 101215568 A
Mourez, Michael, et al. "Mapping dominant-negative mutations of anthrax protective antigen by scanning mutagenesis." Proceedings of the National Academy of Sciences 100.24 (2003): 13803-13808.
US 7601351 B1



- (51) **International Patent Classification:**
C07K 14/325 (2006.01) *A61K 39/07* (2006.01)
- (21) **International Application Number:**
PCT/US2016/048706
- (22) **International Filing Date:**
25 August 2016 (25.08.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/210,771 27 August 2015 (27.08.2015) US
62/323,218 15 April 2016 (15.04.2016) US
- (71) **Applicant:** THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Office of Technology Transfer, National Institutes of Health, 6011 Executive Boulevard, Suite 325, MSC 7660, Bethesda, Maryland 20892-7660 (US).
- (72) **Inventors:** LIU, Shi-Hui; 14516 Settlers Landing Way, Gaithersburg, Maryland 20878 (US). LEPLA, Stephen H.; 9501 Starmont Road, Bethesda, Maryland 20817 (US). BUGGE, Thomas H.; 5270 POOK Hills Road, Bethesda, Maryland 20814 (US). WEIN, Alexander N.; 783 Sunnybrook Drive, Decatur, Georgia 30033 (US). PETERS, Diane E.; 126 Millbury Street, Grafton, Massachusetts 01519 (US). LIU, Jie; 14516 Settlers Landing Way, Gaithersburg, Maryland 20878 (US). CHEN, Kuang-Hua; 14403 Sylvan Glade Drive, Gaithersburg, Maryland 20878 (US).
- (74) **Agents:** LAWLEY, Stephanie M. et al.; Leydig, Voit & Mayer, Two Prudential Plaza, Suite 4900, 180 North Stetson Avenue, Chicago, Illinois 60601-6731 (US).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) **Title:** MODIFIED ANTHRAX TOXIN PROTECTIVE ANTIGEN

(57) **Abstract:** Disclosed is a *Bacillus anthracis* protective antigen (PA) comprising a PA amino acid sequence, wherein one or more of amino acid residues I207, I210, E654, I656, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine. Related compositions, nucleic acids, recombinant expression vectors, host cells, populations of cells, methods of treating or preventing cancer in a mammal, and methods of inhibiting the growth of a target cell are also disclosed.



MODIFIED ANTHRAX TOXIN PROTECTIVE ANTIGEN

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This patent application claims the benefit of U.S. Provisional Patent Application No. 62/323,218, filed April 15, 2016, and U.S. Provisional Patent Application No. 62/210,771, filed August 27, 2015, each of which is incorporated by reference in its entirety herein.

STATEMENT REGARDING
FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] This invention was made with Government support under project numbers HL006185-01, ZIA A10009829-12, 1 ZIA AI000929-12, and Z01DE0699 by the National Institutes of Health, National Institute of Allergy and Infectious Diseases. The Government has certain rights in this invention.

INCORPORATION-BY-REFERENCE OF MATERIAL SUBMITTED
ELECTRONICALLY

[0003] Incorporated by reference in its entirety herein is a computer-readable nucleotide/amino acid sequence listing submitted concurrently herewith and identified as follows: One 49,007 Byte ASCII (Text) file named "726173_ST25.TXT," dated August 25, 2016.

BACKGROUND OF THE INVENTION

[0004] *Bacillus anthracis* anthrax toxin is a bacterial toxin with cytotoxic activity that may be effective for destroying or inhibiting the growth of undesirable cells, e.g., cancer cells. Accordingly, one or more components of the anthrax toxin may be useful for treating or preventing diseases such as, e.g., cancer. However, anthrax toxin or components thereof may be toxic to normal cells. Accordingly, there exists a need for improved anthrax toxins.

BRIEF SUMMARY OF THE INVENTION

[0005] An embodiment of the invention provides a *Bacillus anthracis* protective antigen (PA) comprising a PA amino acid sequence, wherein one or more of amino acid residues

I207, I210, E654, I656, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine.

[0006] Another embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or both of amino acid residues I207 and I210, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine.

[0007] Another embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or more of amino acid residues I656, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted.

[0008] Another embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or more of amino acid residues E654, R659, and M662, as defined by reference to SEQ ID NO: 1, are, independently, substituted.

[0009] Another embodiment of the invention provides a method of treating or preventing a solid tumor in a mammal, the method comprising administering to the mammal pentostatin and cyclophosphamide in an amount effective to treat or prevent the solid tumor in the mammal.

[0010] Further embodiments of the invention provide related compositions, nucleic acids, recombinant expression vectors, host cells, and methods of treating or preventing cancer.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWING(S)

[0011] Figure 1 is a schematic representation of PA oligomers, with each LF-binding site including three subsites (circles) located on adjacent PA monomers. While a heptamer is shown, the same interactions occur within octamers.

[0012] Figure 2 is a graph showing the cell viability (%) of RAW264.7 macrophage cells treated with the indicated PA variants at about 500 ng/mL in the presence of 100 ng/mL LF for 20 h.

[0013] Figure 3A is a schematic representation of intermolecular complementation of the PA variants.

[0014] Figure 3B is a graph showing the cell viability (%) of RAW264.7 macrophage cells treated with the indicated PA variants at 500 ng/mL in the presence of 100 ng/mL LF

(single protein, unshaded bars) or at 250 ng/mL plus 250 ng/mL PA-R200A (shaded bars) or PA-R200C (striped bars) in the presence of 100 ng/mL LF for 6 h.

[0015] Figure 4A is a graph showing the viability (%) of B16-BL6 cells treated with various concentrations (ng/mL) of PA variants PA-U2-R200A (closed circles), PA-L1-I207R (closed squares), PA-L1-I210A (diamonds), PA-L1-I207R+PA-U2-R200A (open squares), PA-L1-I210A+PA-U2-R200A (open circles).

[0016] Figure 4B is a graph showing the survival of C57BL/6J mice injected intraperitoneally with three doses of 20 μ g PA-L1-I207R (squares) or PA-L1-I210A (diamonds) plus 10 μ g FP59 at days 0, 1 and 2.

[0017] Figure 5 is a graph showing the tumor weight (mg) in B16-BL6 tumor-bearing mice treated intraperitoneally with phosphate buffered saline (PBS) (squares), L1-I210A/U2-R200A/LF (7.5 μ g/7.5 μ g/5 μ g) (circles), L1-I210A/U2-R200A/LF (22.5 μ g/22.5 μ g/15 μ g) (diamonds), L1-I207R/U2-R200A/LF (7.5 μ g/7.5 μ g/5 μ g) ($\blacktriangle$), or L1-I207R/U2-R200A/LF (22.5 μ g/22.5 μ g/15 μ g) ($\blacktriangledown$). Doses were administered on each of the five days shown by the arrows.

[0018] Figures 6A-6C are graphs showing the tumor volume (mm^3) (mean $\pm$ SE) in *Cmg2*^{+/+} (squares), *Cmg2*^{+/-} (diamonds), or *Cmg2*^{-/-} (circles) mice bearing A549 (A), LL3 (B), or B16 (C) tumors at various time points (days) after inoculation.

[0019] Figures 6D-6E are graphs showing the tumor volume (mm^3) (mean $\pm$ SE) of *Tem8*^{-/-} mice (circles) and littermate *Tem8*^{+/+} mice (squares) bearing LL3 (D) or B16 (E) tumors at various time points (days) after inoculation.

[0020] Figures 6F-6H are graphs showing the body weight (g) (mean $\pm$ SD) in *Cmg2*^{+/+} (squares), *Cmg2*^{+/-} (diamonds), or *Cmg2*^{-/-} (circles) mice bearing A549 (F), LL3 (G), or B16 (H) tumors at various time points (days) after inoculation.

[0021] Figures 6I is a graph showing the body weight (g) (mean $\pm$ SD) of *Tem8*^{-/-} mice (grey bars) and littermate *Tem8*^{+/+} mice (black bars) bearing LL3 tumors the day when tumor cells were inoculated.

[0022] Figure 6J is a graph showing the body weight (g) (mean $\pm$ SD) of *Tem8*^{-/-} mice (circles) and littermate *Tem8*^{+/+} mice (squares) bearing B16 tumors at various time points (days) after inoculation.

[0023] Figures 7A and 7B are graphs showing the tumor volume (mm^3) (mean $\pm$ SE) of LL3 tumor-bearing mice treated with PBS (squares), PA-L1 (circles) (A), or IC2-PA/LF

(diamonds) (B) at various time points (days) after first treatment. The treatment was administered on the days indicated by arrows.

[0024] Figure 7C is a graph showing the tumor volume (mm^3) (mean $\pm$ SE) of B16-BL6 tumor-bearing mice treated with PBS (squares), PA-L1 (circles), or IC2-PA/LF (diamonds) at various time points (days) after first treatment. The treatment was administered on the days indicated by arrows.

[0025] Figure 7D is a graph showing the cell viability (%) of LL3 cells treated with PA (open squares) or PA-L1 (closed squares) or B16-BL6 cells treated with PA (open circles) or PA-L1 (closed circles) in the presence of LF (mean $\pm$ SD).

[0026] Figures 7E and 7F are graphs showing the tumor volume (mm^3) (mean $\pm$ SE) (E) and body weight (g) (mean $\pm$ SD) (F) of *Cmg2^{+/+}* (squares), *Cmg2^{+/-}* (diamonds), and *Cmg2^{-/-}* (circles) mice treated with 15 μg PA-L1 plus 7.5 μg LF on days after the first treatment. The treatment was administered on the days indicated by arrows.

[0027] Figures 7G and 7H are graphs showing the tumor volume (mm^3) (mean $\pm$ SE) (G) and body weight (g) (mean $\pm$ SD) (H) of A549 tumor-bearing littermate *Tem8^{+/+}* (squares), *Tem8^{+/-}* (diamonds), and *Tem8^{-/-}* (circles) mice treated with 15 μg PA-L1 plus 7.5 μg LF on days after the first treatment. The treatment was administered on the days indicated by arrows.

[0028] Figures 8A-8C are graphs showing the tumor volume (mm^3) (mean $\pm$ SE) of B16-BL6 melanoma-bearing mice with the indicated CMG2 genotypes that were treated with 30 μg PA-L1 plus 15 μg LF on days after the first treatment. The treatment was administered on the days indicated by arrows.

[0029] Figure 8D is a graph showing the tumor volume (mm^3) (mean $\pm$ SE) of B16-BL6 melanoma-bearing endothelial cell-specific CMG2-null mice and myeloid-specific CMG2-null mice (*Cmg2(Myel)^{-/-}*) treated with 30 μg IC2-PA (15 μg PA-L1-I207R + 15 μg PA-U2-R200A) plus 10 μg LF or PBS at the days indicated by the arrows.

[0030] Figure 8E is a graph showing the survival (%) of WT mice and *Cmg2^{EC}* mice that were treated with PA-L1/LF on the days indicated by arrows.

[0031] Figure 8F is a graph showing the number of blood vessels per 10 mm^2 of mice treated with PBS (left bar) or 3 doses of 30 μg PA-L1 plus 15 μg LF (right bar) on days 0, 2, and 4.

[0032] Figure 9A is a graph showing the cell density (%) of tumor endothelial cells incubated with various concentrations of PA-L1 in the presence of LF (500 ng/mL). Data are shown as mean $\pm$ SD.

[0033] Figure 9B is a graph showing the ECAR (mpH/min) measured at various time points (min) under basal conditions or conditions following sequential additions of the mitochondrial inhibitors (oligomycin, FCCP, and antimycin A) of tumor endothelial cells treated with or without PA-L1/LF (1 μ g/mL each) for 24 h. The ECAR readings were normalized to amounts of cells having 50 mg total protein, mean $\pm$ SD. (n=3 independent experiments).

[0034] Figure 9C is a graph showing the OCR (pM/min) measured at various time points (min) under basal conditions or conditions following sequential additions of the mitochondrial inhibitors (oligomycin, FCCP, and antimycin A) of tumor endothelial cells treated with or without PA-L1/LF (1 μ g/mL each) for 24 h. The OCR readings were normalized to amounts of cells having 50 mg total protein, mean $\pm$ SD. (n=3 independent experiments).

[0035] Figure 9D is a graph showing (i) the OCR_{ATP} obtained by subtracting the OCRs after addition of oligomycin from the basal OCRs of the endothelial cells, (ii) spare respiratory capacity (SRC), and (iii) the maximal respiration (MR) values of cells treated with (dark grey bars) or without (light grey bars) PA-L1/LF (1 μ g/mL each). Data are shown as mean $\pm$ SD.

[0036] Figure 9E is a graph showing the relative cellular ATP levels of tumor endothelial cells treated with PA-L1/LF (0.5 μ g/mL or 2 μ g/mL each) for 24 h vs. untreated cells. Data are shown as mean $\pm$ SD.

[0037] Figure 9F is a graph showing the change in expression (%) of the indicated genes in tumor endothelial cells treated with or without PA-L1/LF (1 μ g/mL each) for 24 h. Eukaryotic translation initiation factor *EIF35S* was used as an internal normalization control. *, $P < 0.05$; ***, $P < 0.01$.

[0038] Figures 10A and 10B are graphs showing the tumor volume (mm³) (A) and body weight (g) (B) in LL3 lung carcinomas-bearing immunocompetent C57BL/6 mice that were treated with PBS (closed squares), IC2-PA/LF (20 μ g/6.7 μ g) (* and open circles), PC regimen (1 mg pentostatin and 50 mg cyclophosphamide) (∇), high (25 μ g/8.1 μ g) (Δ), medium (20 μ g/6.7 μ g) (open squares), or low (15 μ g/5 μ g) (diamonds and $\blacktriangle$) doses of IC2-PA/LF combined with PC regimen at various time points (days) after tumor inoculation. Schedules for PC and the toxin treatments are indicated by the arrows. Tumor weights, mean $\pm$ SE. Body weights, mean $\pm$ SD.

[0039] Figure 10C is a graph showing the viability (%) of RAW264.7 cells that were incubated with IC2-PA/LF (100 ng/mL each) for 5 h in the presence of various dilutions of sera obtained from representative mice in Figures 10A and 10B after the 1st (closed circles) or 2nd (open circles) round of therapy. The viability of cells treated with PC and IC2-PA/LF after the 2nd (diamonds) or 3rd (Δ) round of therapy, PC ($\blacktriangledown$), or PBS (squares) are also shown.

[0040] Figure 10D is a graph showing the viability (%) of RAW264.7 cells that were incubated with IC2-PA/LF (100 ng/mL each) for 5 h in the presence of various dilutions of sera obtained from representative mice in Figures 10A and 10B after the 1st round of therapy (circles). The viability of cells treated with PC and IC2-PA/LF after the 4th round of therapy (squares) is also shown.

[0041] Figure 10E is a graph showing the viability (%) of cells treated as described in Figures 10A-10D using various concentrations of 14B7 anti-PA monoclonal antibodies as a positive control for neutralizing antibodies.

[0042] Figures 11A-11D are graphs showing the number of cells (millions) isolated from B16-BL6 melanoma-bearing mice that were positive for the indicated B-cell (A), T-cell (B and C), or granulocyte markers (D) after the 2nd cycle of therapy with PBS (grey bars), PC (black bars), IC2-PA/LF (right striped bars), or the combined PC and the toxin (left striped bars) as measured by flow cytometry. Naïve, untreated cells (white bars) were measured as a control.

[0043] Figure 11E is a graph showing the total splenocyte count (millions) of the mice treated in Figures 11A-11D.

[0044] Figure 12A is a graph showing the tumor volume (mm^3) (mean $\pm$ SE) of *Tem8*^{-/-} mice (circles), *Tem8*^{+/-} mice (diamonds), *Tem8*^{+/+} mice (squares) bearing A549 tumors at various time points (days) after inoculation.

[0045] Figure 12B is a graph showing the tumor volume (mm^3) (mean $\pm$ SE) of *Tem8*^{-/-} mice (grey bars) and *Tem8*^{+/+} mice (black bars) bearing LL3 tumors at various time points (days) after inoculation.

[0046] Figure 12C is a graph showing the body weight (g) (mean $\pm$ SD) of *Tem8*^{-/-} mice (circles), *Tem8*^{+/-} mice (diamonds), *Tem8*^{+/+} mice (squares) bearing A549 tumors at various time points (days) after inoculation.

[0047] Figure 12D is a graph showing the body weight (g) (mean $\pm$ SD) of *Tem8^{-/-}* mice (grey bars) and *Tem8^{+/+}* mice (black bars) bearing LL3 tumors at various time points (days) after inoculation.

[0048] Figure 12E is a graph showing the percentage of mice with misaligned teeth having the following genotypes and fed the following foods: *Tem8^{+/+}* (hard food) (n=46) (closed squares); *Tem8^{+/+}* (soft food) (n=26) (open squares); *Tem8^{+/-}* (hard food) (n=91) (closed diamonds); *Tem8^{+/-}* (hard food) (n=61) (open diamonds); *Tem8^{+/-}* (soft food) (n=151) (closed circles); and *Tem8^{-/-}* (soft food) (n=74) (open circles).

[0049] Figures 12F and 12G are graphs showing the body weight of female (F) and male (G) mice that were fed hard food and which had the following genotypes: *Tem8^{+/+}* (squares); *Tem8^{+/-}* (diamonds); and *Tem8^{-/-}* (circles).

[0050] Figures 12H and 12I are graphs showing the body weight of female (H) and male (I) mice that were fed soft food and which had the following genotypes: *Tem8^{+/+}* (squares); *Tem8^{+/-}* (diamonds); and *Tem8^{-/-}* (circles).

[0051] Figure 13A is a graph showing the viability (%) of cells treated with LF and various concentrations of PA. The cell types and treatments were as follows: A549 cells treated with PA/LF (open squares); A549 cells treated with PA-L1/LF (closed squares); Colo205 cells treated with PA/LF (open circles); and Colo205 cells treated with PA-L1/LF (closed circles).

[0052] Figure 13B is a graph showing the viability (%) of cells treated with FP59 and various concentrations of PA. The cell types and treatments were as follows: A549 cells treated with PA/FP59 (striped bars right of dotted line); A549 cells treated with PA-L1/FP59 (shaded bars right of dotted line); Colo205 cells treated with PA/FP59 (striped bars left of dotted line); and Colo205 cells treated with PA-L1/FP59 (shaded bars left of dotted line).

[0053] Figure 14 is a graph showing the Colo205 tumor volume (mm³) in mice at various time points (days) after first treatment. Treatments were administered on the days indicated by arrows. The genotypes and treatments were as follows: *Cmg2^{+/+}* mice treated with PBS (open squares); *Cmg2^{+/+}* mice treated with PA-L1/LF (closed squares); *Cmg2^{+/-}* treated with PBS (open diamonds); and *Cmg2^{+/-}* mice treated with PA-L1/LF (closed diamonds).

[0054] Figures 15A and 15B are graphs showing the tumor volume (mm³) (A) and body weight (g) (B) in mice bearing B16-BL6 melanoma tumors receiving the indicated treatments at various time points (days) after tumor inoculation. The treatments were given on the days

indicated by arrows. The treatments were as follows: PBS (closed squares); IC2-PA/LF (circles); PC (triangles); and PC + IC2-PA/LF (open squares).

[0055] Figure 15C is a graph showing the viability (%) of cells that were incubated with the indicated treatments in the presence of various dilutions of sera obtained from representative mice in Figures 15A and 15B after the indicated round of therapy. The treatments and the rounds of therapy were as follows: IC2-PA/LF (2nd) (circles); PC (▼); PC+IC2-PA/LF (2nd) (△); PC+IC2-PA/LF (5th) (▲); PBS (closed squares); and naïve (untreated) (open squares).

[0056] Figure 15D is a graph showing the viability of cells (%) that were treated with the LF and the indicated concentrations of PA. The EC₅₀ of PA is 6 ng/mL.

[0057] Figures 16A and 16B are graphs showing the viability (%) of LL3 (circles) and B16-BL cells (squares) that were treated with various concentrations of cyclophosphamide (A) or pentostatin (B).

[0058] Figure 16C is a graph showing the viability (%) of LL3 (circles) or B16-BL cells (squares) (%) that were treated with various concentrations of pentostatin and cyclophosphamide.

[0059] Figure 16D is a graph showing the viability (%) of endothelial cells that were treated with various concentrations of pentostatin and cyclophosphamide.

[0060] Figure 17A is a graph showing the survival (percentage) of wild-type (WT) (squares), CMG2^{-/-} (circles), and TEM8^{-/-} (diamonds) mice at various time points (hours) following injection of native PA and FP59. Group sizes are indicated for each toxin.

[0061] Figure 17B is a graph showing the survival (percentage) of WT (squares), CMG2^{-/-} (circles), and TEM8^{-/-} (diamonds) mice at various time points (hours) following injection of PA I656Q and FP59. Group sizes are indicated for each toxin.

[0062] Figure 17C is a graph showing the survival (percentage) of WT (squares), CMG2^{-/-} (circles), and TEM8^{-/-} (diamonds) mice at various time points (hours) following injection of PA I656V and FP59. Group sizes are indicated for each toxin.

[0063] Figure 17D is a graph showing the survival (percentage) of (i) WT (open squares) and CMG2^{-/-} (open circles) mice at various time points (hours) following injection of the PA E654T/R659S/M662R protein (abbreviated PA TSR) and FP59 and (ii) WT (closed squares) and CMG2^{-/-} (closed circles) mice at various time points (hours) after injection of native PA and FP59. Group sizes are indicated for each toxin.

[0064] Figure 18A is a graph showing the body weight (g) of mice bearing a HeLa xenograft which were injected intraperitoneally with PBS (■), 15 µg PA-L1 + 5 µg LF (▲), 15 µg PA-L1 I656Q + 5 µg LF (●) or 30 µg PA-L1 TSR + 10 µg LF (◆). Group sizes are indicated for each toxin.

[0065] Figure 18B is a graph showing the weight (g) of intradermal tumor nodules (expressed as mean of tumor weight) in mice bearing a HeLa xenograft which were injected intraperitoneally with PBS (■), 15 µg PA-L1 + 5 µg LF (▲), 15 µg PA-L1 I656Q + 5 µg LF (●) or 30 µg PA-L1 TSR + 10 µg LF (◆). Group sizes are indicated for each toxin. Arrows indicate days on which toxin was injected. PBS vs. PA-L1/LF = $P < 0.01$. PBS vs. PA-L1-I656Q/LF = $P < 0.01$. PA-L1-TSR/LF vs. PA-L1/LF = $P < 0.01$. PA-L1-TSR/LF vs PA-L1-I656Q/LF = $P < 0.01$.

[0066] Figures 19A and 19C are graphs showing oral melanoma tumor size (cm³) in Dog 1 (A) and Dog 2 (C) treated with 750 µg IC2-PA (PA-L1-I207R+PA-U2-R200A) plus 250 µg LF at various time points (days) after first treatment. The rightmost arrow indicates surgical removal of the tumor. The remainder of the arrows indicate the days on which the dogs were treated with IC2-PA plus LF.

[0067] Figures 19B and 19D are graphs showing the body weight (kg) of Dog 1 (B) and Dog 2 (D) at various time points after first treatment with IC2-PA plus LF.

DETAILED DESCRIPTION OF THE INVENTION

[0068] Anthrax toxin is a protein toxin produced by *Bacillus anthracis*. Anthrax toxin includes three components: the protective antigen (PrAg or PA), the lethal factor (LF), and the edema factor (EF). PA binds to a cell surface receptor for PA. The native, wild-type PA is cleaved after the sequence RKKR (SEQ ID NO: 4) by cell-surface furin or furin-like proteases into two fragments: PrAg63 and PrAg20. PrAg63 is a 63 kDa, C-terminal fragment which remains receptor-bound and forms a homo-oligomeric heptamer. PrAg20 is a 20 kDa, N-terminal fragment which is released into the cell medium. The PrAg63 homo-oligomeric heptamer forms the binding site for effector components LF and/or EF. Each LF/EF binding site is comprised of three subsites on two adjacent PA monomers. Upon the binding of LF and/or EF to PA, the resulting complex is internalized in a target cell where the LF and/or EF exert toxic effects.

[0069] The premature native, wild-type PA amino acid sequence is set forth in Genbank Accession No. NP_052806 (SEQ ID NO: 2). A coding sequence for the premature native,

wild-type PA amino acid sequence is the nucleotide sequence of SEQ ID NO: 33. Another premature PA amino acid sequence is set forth in SEQ ID NO: 5. Premature PA contains an N-terminal signal peptide (SEQ ID NO: 3, which corresponds to amino acid residues 1-29 of SEQ ID NO: 2) which is removed in the mature form of PA. The mature, wild-type PA amino acid sequence (without the N-terminal signal peptide) is set forth in SEQ ID NO: 1.

[0070] Unless specified otherwise, amino acid residue position numbers of PA are defined herein by reference to the amino acid sequence of the mature, wild-type PA amino acid sequence of SEQ ID NO: 1. Thus, substitutions of PA are described herein by reference to the amino acid residue present at a particular position, followed by the amino acid with which that residue has been replaced in the particular substitution under discussion. In this regard, the positions of the amino acid sequence of a particular embodiment of a PA are referred to herein as the positions as defined by SEQ ID NO: 1. The actual positions of the amino acid sequence of a particular embodiment of a PA are defined relative to the corresponding positions of SEQ ID NO: 1 and may represent different residue position numbers than the residue position numbers of SEQ ID NO: 1. Thus, for example, substitutions refer to a replacement of an amino acid residue in the amino acid sequence of a particular embodiment of a PA corresponding to the indicated position of the 735-amino acid sequence of SEQ ID NO: 1 with the understanding that the actual positions in the respective amino acid sequences may be different. For example, when the positions are as defined by SEQ ID NO: 1, the term "R200" refers to the arginine normally present at position 200 of SEQ ID NO: 1, "R200A" indicates that the arginine normally present at position 200 of SEQ ID NO: 1 is replaced by an alanine, while "I207R" indicates that the isoleucine normally present at position 207 of SEQ ID NO: 1 has been replaced with an arginine. In the event of multiple substitutions at two or more positions, the two or more substitutions may be the same or different, i.e., each amino acid residue of the two or more amino acid residues being substituted can be substituted with the same or different amino acid residue unless explicitly indicated otherwise.

[0071] Variants of PA, and combinations thereof, may provide cancer-cell specific cytotoxicity. For example, the PA variant PA-U2-R200A is a urokinase plasminogen activator-activated PA variant with LF-binding subsite II residue R200 mutated to Ala, and the PA variant PA-L1-I210A is a matrix metalloproteinase-activated PA variant with LF-binding subsite III residue I210 mutated to Ala. PA-U2-R200A and PA-L1-I210A provide reduced cytotoxicity when used singly. However, when combined, PA-U2-R200A and PA-

L1-I210A (collectively referred to as IC-PA) form LF-binding competent heterogeneous oligomers by intermolecular complementation, and achieve high specificity in tumor targeting. Nevertheless, IC-PA may provide undesirable non-specific toxicity to normal tissues.

[0072] It has been discovered that replacing the native amino acid residue at one or both of positions 207 and 210 of *Bacillus anthracis* PA with an amino acid residue other than alanine advantageously reduces non-specific toxicity to normal tissues and provides potent anti-tumor activity. Accordingly, an embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or both of amino acid residues I207 and I210, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine.

[0073] In a preferred embodiment, the PA comprises a PA amino acid sequence, wherein amino acid residue I207, as defined by reference to SEQ ID NO: 1, is substituted, with the proviso that amino acid residue I207 is not substituted with alanine.

[0074] Amino acid residues I207 and I210 may be independently substituted with any amino acid residue other than alanine. Preferably, amino acid residue I210 is substituted with aspartic acid, glutamic acid, lysine, glutamine, arginine, or serine. Preferably, amino acid residue I207 is substituted with arginine, tryptophan, or tyrosine. In an especially preferred embodiment, amino acid residue I207 is substituted with arginine.

[0075] It has also been discovered that replacing the native amino acid residue at one or more of positions 656, 681, and 687 of *Bacillus anthracis* PA advantageously improves selectivity for CMG2 (which is expressed by tumor endothelial cells), reduces non-specific toxicity to normal tissues, and provides potent anti-tumor activity. Accordingly, an embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or more of amino acid residues I656, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted.

[0076] Amino acid residues I656, Y681, and L687 may be independently substituted with any amino acid residue. Preferably, amino acid residue I656 is substituted with glutamine, valine, alanine, cysteine, or glutamic acid. Preferably, amino acid residue Y681 is substituted with alanine. Preferably, amino acid residue L687 is substituted with alanine.

[0077] It has also been discovered that replacing the native amino acid residue at one or more of positions 654, 659, and 662 of *Bacillus anthracis* PA advantageously improves

selectivity for TEM8. Accordingly, an embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or more of amino acid residues E654, R659, and M662, as defined by reference to SEQ ID NO: 1, are, independently, substituted. A PA wherein one or more of amino acid residues E654, R659, and M662 is substituted may be useful as, for example, a research tool and/or may provide anti-tumor activity for tumors that have a high expression of TEM8.

[0078] Amino acid residues E654, R659, and M662 may be independently substituted with any amino acid residue. Preferably, amino acid residue E654 is substituted with threonine. Preferably, amino acid residue R659 is substituted with serine. Preferably, amino acid residue M662 is substituted with arginine.

[0079] The inventive PA may comprise any of the amino acid residue substitutions described herein in any combination. Accordingly, an embodiment of the invention provides a *Bacillus anthracis* PA comprising a PA amino acid sequence, wherein one or more of amino acid residues I207, I210, E654, I656, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine.

[0080] The terms “protective antigen,” “PA,” and “PrAg,” as used herein, include PA that includes any one or more of the following modifications: (a) substitution of a native cell-recognition domain for a non-native cell-recognition domain; (b) substitution of a native proteolytic activation site for a non-native proteolytic activation site; (c) modification of a first PA to generate a first modified PA, whereby the first modified PA can pair only with a second PA; or (d) modification of the first PA and the second PA, whereby an effector component (e.g., LF and/or EF) can bind only at a site formed by the interaction of the first PA and the second PA.

[0081] For example, the PA may be modified by substitution of the native cell recognition domain for a non-native cell recognition domain as described in, e.g., U.S. Patent 7,947,289, which is incorporated herein by reference. The non-native cell recognition domain recognizes and/or binds to a molecule on the surface of a target cell population (e.g., a cancer cell), thus specifically targeting the modified PA to the target cells. Substitution of a native cell recognition domain may comprise, e.g., substitution of the native cell recognition domain with a recognition domain for another cell surface molecule. The PA may be modified such that it no longer binds to the PA receptors, but binds to other cell surface

molecules (e.g., receptors for cytokines). The PA may also be modified such that it binds to the PA receptor in addition to other cell surface molecules. Exemplary non-native cell recognition domains include, e.g., an antibody, a cytokine, or a cell surface receptor ligand. Suitable non-native cell recognition domains include, e.g., VEGF, C-CSF, GM-CSF, EPO, EGF, IL-1, IL-2, IL-4, IL-5, IL-6, interferon α , interferon γ , growth hormone, prolactin, thrombopoietin, and TGF- β . Exemplary antibodies may include, e.g., an antibody that specifically binds a protein that is overexpressed on cancer cells such as, for example, Her2/Neu, CD19, CD276, CD25, CD30, CA19-9, CA-125, VEGF receptors, C-CSF receptors, GM-CSF receptors, EPO receptors, EGF receptors, interleukin receptors (e.g., IL-1R, IL-2R, IL-4R, IL-5R, or IL-6R), interferon receptors (e.g., interferon α or interferon γ), growth hormone receptors, prolactin receptors, thrombopoietin receptors, and TGF- β receptors.

[0082] Alternatively or additionally, the PAs may be modified by substitution of the native proteolytic cleavage site(s) with nonnative proteolytic cleavage site(s) as described in, e.g., U.S. Patent 7,947,289, which is incorporated herein by reference. Thus, the PA can be activated via cleavage by proteases present on the surface of specific target cell types (e.g., cancer cells). Examples of proteolytic cleavage sites that can be substituted for native cleavage sites on the PA include cleavage sites for any proteases known in the art including, e.g., metalloproteinase, a cysteine protease, an aspartic acid protease, a plasminogen activator, a kallikrein, a type 1 transmembrane serine protease, a type 2 transmembrane serine protease, or a GPI anchored serine protease. Proteases include, e.g., plasminogen activators (uPA and tPA), matrix metalloproteinases, (e.g., MMP-1; MMP-2; MMP-3; MMP-7; MMP-8; MMP-9; MMP-10; MMP-11; MMP-12; MMP-13; MMP-14; MMP-15; MMP-16; MMP-17; MMP-19; MMP-20; MMP-21; MMP-23A; MMP-23B; MMP-24; MMP-25; MMP-26; MMP-27; MMP-28; and MT2-MMP); metalloproteases (e.g., Meprin a; Meprin b; Decysin; ADAM1a; ADAM2; ADAM3B; ADAM4; ADAM4B; ADAM5; ADAM6; ADAM7; ADAM8; ADAM9; ADAM10; ADAM11; ADAM12; ADAM15; ADAM17; ADAM18; ADAM19; ADAM20; ADAM21; ADAM22; ADAM23; ADAM28; ADAM29; ADAM30; ADAM32; ADAM33; ADAMTS1; ADAMTS2; ADAMTS3; ADAMTS4; ADAMTS5/11; ADAMTS6; ADAMTS7; ADAMTS8; ADAMTS9; ADAMTS10; ADAMTS12; ADAMTS13; ADAMTS14; ADAMTS15; ADAMTS16; ADAMTS17; ADAMTS18; ADAMTS19; ADAMTS20); serine proteases (e.g., Kallikrein hK1; Kallikrein hK2; Kallikrein hK3/PSA; Kallikrein hK4; Kallikrein hK5; Kallikrein hK6; Kallikrein hK7;

Kallikrein hK8; Kallikrein hK9; Kallikrein hK10; Kallikrein hK11; Kallikrein hK12; Kallikrein hK13; Kallikrein hK14; Kallikrein hK15; Thrombin; Coagulation factor VIIa; Coagulation factor IXa; Coagulation factor Xa; Coagulation factor XIa; Coagulation factor XIIa; Protein C; Protein Z; Mastin; Trypsin- α 1; Trypsin- α 2; Trypsin- β 31; Trypsin- δ 1; Trypsin- γ 1; Marapsin; Marapsin-2; Testisin; Brain serine protease-2; Proctasin; Proctasin-like 1; Proctasin-like 2; Chymase; Cathepsin G; Neutrophil elastase; Azurocidin; Hepsin; HAT-related protease; HAT (Human Airway Trypsin-like Protease); Type I transmembrane serine proteases, type II transmembrane serine proteases, cysteine proteases, aspartic acid proteases, HAT-like 1; HAT-like 2; HAT-like 3; HAT-like 4; HAT-like 5; DESC1; Corin; Matriptase; Matriptase-2; Matriptase-3; TMPRSS3; TMPRSS4; Spinesin; Polyserase; MSPL; Neurotrypsin; Urokinase plasminogen activator; Tissue plasminogen activator; Plasminogen; Acrosin; Plasma-kallikrein-like 1; Plasma-kallikrein-like 2; Plasma-kallikrein-like 3; Plasma-kallikrein-like 4; and Seprase); and aspartic proteases (e.g., Cathepsin D; and Cathepsin E); Cysteine Proteases (e.g., Cathepsin B; Cathepsin C; Cathepsin F; Cathepsin H; Cathepsin K; Cathepsin L; Cathepsin L2; Cathepsin S; Cathepsin W; Cathepsin Z and Cathepsin J).

[0083] In a preferred embodiment, the PA is modified by substitution of the native proteolytic cleavage site(s) with an MMP cleavage site or a plasminogen activator cleavage site. Both MMP and plasminogen activator proteases are overproduced by tumor tissues and are implicated in cancer cell growth and metastasis (Dano et al., *APMIS*, 107: 120-127 (1999)). In an embodiment of the invention, the native furin cleavage site (e.g., SEQ ID NO: 4) of the PA is replaced with an MMP cleavage site or a plasminogen activator cleavage site. Examples of MMP cleavage sites that may be useful in the inventive PAs include GPLGMLSQ (SEQ ID NO: 6) and GPLGLWAQ (SEQ ID NO: 7). In this regard, the PA may comprise an MMP cleavage site. Examples of plasminogen activator cleavage sites that may be useful in the inventive PAs include PCPGRVVGG (SEQ ID NO: 8), PGSGRSA (SEQ ID NO: 9), PGSGKSA (SEQ ID NO: 10), PQRGRSA (SEQ ID NO: 11), PCPGRVVGG (SEQ ID NO: 12), GSGRSA (SEQ ID NO: 13), GSGKSA (SEQ ID NO: 14), QRGRSA (SEQ ID NO: 15). In this regard, the PA may comprise a plasminogen activator cleavage site.

[0084] Alternatively or additionally, the PA may be modified so that each individual PA molecule (PA monomer) can only form hetero-oligomers (i.e., each PA monomer can pair only with a PA monomer of a different type (i.e., having a different amino acid sequence)) as described in, e.g., U.S. Patent 7,947,289, which is incorporated herein by reference. The

modified PA monomers may comprise PA monomer binding sites which have been modified such that the PA monomer can bind only to a PA monomer of a different type. In some embodiments, the PA monomer binding sites have been modified such that the PA monomer can bind two PA monomers of two different types. For example, a PA monomer that forms a heptameric complex can be modified so that each PA monomer can bind only to two different, but complementary PA monomers, each of which has a modified PA monomer binding site. For example, one or more of the following amino acid substitutions may provide a PA monomer that can only form a hetero-oligomeric heptamer: aspartic acid at position 512 for alanine; aspartic acid at position 512 for lysine; lysine at position 199 for glutamic acid; arginine at position 468 for alanine, and arginine at position 470 for aspartic acid. For example, a modified PA monomer comprising an alanine at position 512 is unable to homo-oligomerize, but can form functional hetero-oligomers with a modified PA with a glutamic acid at position 199, an alanine at position 468, and an aspartic acid at position 470, as defined by reference to SEQ ID NO: 1.

[0085] Alternatively or additionally, the PA may be modified so that at least two PA monomers (each having a different amino acid sequence) are needed to bind an effector component (e.g., LF and/or EF) so that the effector component can be delivered to a target cell and exert a biological effect (e.g., target cell killing or inhibition of target cell proliferation) as described in, e.g., U.S. Patent 7,947,289, which is incorporated herein by reference. The portion of a first PA monomer that binds to the effector component may be modified so that a second PA monomer of a different type (i.e., having a different amino acid sequence) is required to effectively bind the effector molecule.

[0086] For example, the native lethal factor (LF) binding site of the PA may be mutated so that at least two different PA monomer types are required to bind LF. In an embodiment of the invention, a substitution of the native amino acid residue at one or more of the following positions may provide a PA monomer that cannot homo-oligomerize to form a functional LF binding site: 178, 197, 200, 207, 210, and 214, as defined by reference to SEQ ID NO: 1. However, in one embodiment of the invention, the combination of (i) a PA monomer comprising a substitution of the native amino acid residue at position 200 and (ii) a PA monomer comprising a substitution of the amino acid residue at position 207 may form functional PA heptamers that bind LF. In another embodiment of the invention, the combination of (i) a PA monomer comprising a substitution of the native amino acid residue at position 200 and (ii) a PA monomer comprising a substitution of the amino acid residue at

position 210 may form functional PA heptamers that bind LF. For example, one or more of the following substitutions may provide a PA monomer that cannot homo-oligomerize to form a functional LF binding site: arginine at position 178 for alanine; lysine at position 197 for alanine; arginine at position 200 for alanine; isoleucine at position 207 for arginine; isoleucine at position 210 for alanine; lysine at position 214 for alanine, as defined by reference to SEQ ID NO: 1. Modified PA monomers comprising alanine at position 200 (e.g., R200A), modified PA monomers comprising alanine at position 210 (e.g., I210A), modified PA monomers comprising arginine at position 210 (e.g., I207R), and modified PA monomers comprising alanine at position 214 (e.g., K214A) are unable to homo-oligomerize and form a functional PA heptamer that binds LF. However, the combination of any one or more of (i) a PA monomer having the substitution of R200A or R200C with (ii) any one or more of a PA monomer having the substitution of I207R, I207W, I210D, I210E, I210K, I210Q, I210R, or I210S may form functional PA heptamers that bind LF. For example, the combination of a PA monomer comprising R200A and a PA monomer comprising I207R may form functional PA heptamers that bind LF.

[0087] Accordingly, an embodiment of the invention also provides compositions comprising combinations of any of the PA monomers described herein which, when employed singly, are unable to homo-oligomerize and form a functional PA heptamer that binds LF, but when combined, may form functional PA heptamers that bind LF.

Accordingly, an embodiment of the invention provides a composition comprising a first PA comprising a first PA amino acid sequence and a second PA comprising a second PA amino acid sequence. The first PA comprises a first PA amino acid sequence, wherein (i) one or both of amino acid residues I207 and I210, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine; (ii) one or more of amino acid residues I207, I210, E654, I656, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine; (iii) one or more of amino acid residues I656, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted; or (iv) one or more of amino acid residues E654, R659, and M662, as defined by reference to SEQ ID NO: 1, are, independently, substituted. The second PA comprises a second PA amino acid sequence, wherein amino acid residue R200 of the second PA amino acid sequence, as defined by

reference to SEQ ID NO: 1, is substituted. The second PA amino acid sequence is different from the first PA amino acid sequence.

[0088] Amino acid residue R200 of the second PA amino acid sequence may be substituted with any amino acid residue. In an embodiment of the invention, amino acid residue R200 of the second PA amino acid sequence is substituted with alanine, cysteine, aspartic acid, glutamic acid, glycine, isoleucine, methionine, proline, serine, valine, or tryptophan. In a preferred embodiment, amino acid residue R200 of the second PA amino acid sequence is substituted with alanine.

[0089] The first PA and the second PA may include any of the modifications described herein with respect to other aspects of the invention. In an embodiment of the invention, the second PA amino acid sequence is modified by substitution of the native proteolytic cleavage site(s) with any of the nonnative proteolytic cleavage site(s) described herein. In an embodiment of the invention, the first PA comprises an MMP cleavage site and the second PA comprises a plasminogen activator cleavage site. The plasminogen activator cleavage site may be selected from the group consisting of SEQ ID NOs: 8-15.

[0090] The inventive compositions may further comprise an effector component which binds to a heptamer formed by the modified PAs and exerts a biological effect (e.g., killing of a target cell or inhibition of target cell proliferation). Suitable effector components include, e.g., anthrax lethal factor, anthrax edema factor, truncated anthrax lethal factor (e.g., LFn or amino acids 1-254 of anthrax lethal factor), FP59 (LFn fused to the ADP-ribosylation domain of *Pseudomonas* exotoxin A as described in, e.g., Arora et al., *J. Biol. Chem.*, 268:3334-3341 (1993) and WO 01/21656), and cytolethal distending toxin subunit B (CdtB) conjugated or fused to *Bacillus anthracis* toxin LF as described in International Publication No. WO 2014/205187.

[0091] Toxin proteins such as, for example, the PAs described herein, may be highly immunogenic upon administration to a mammal. Such immunogenicity may reduce the amount of PA that can be given to a mammal which may, in turn, reduce the effectiveness of the PA for treating or preventing the disease, e.g., cancer. It has been discovered that the combination of pentostatin and cyclophosphamide advantageously reduces or prevents the induction of antibodies against the inventive PAs described herein. Accordingly, an embodiment of the invention provides any of the compositions described herein, wherein the composition further comprises an immunosuppressive agent. In an embodiment of the

invention, the immunosuppressive agent is the combination of pentostatin and cyclophosphamide.

[0092] An embodiment of the invention provides a nucleic acid comprising a nucleotide sequence encoding any of the inventive PAs described herein. In an embodiment, the nucleic acid comprises a nucleotide sequence encoding a PA amino acid sequence including the I207R substitution, wherein the nucleotide sequence comprises SEQ ID NO: 29.

[0093] The term “nucleic acid,” as used herein, includes “polynucleotide,” “oligonucleotide,” and “nucleic acid molecule,” and generally means a polymer of DNA or RNA, which can be single-stranded or double-stranded, which can be synthesized or obtained (e.g., isolated, purified, or both isolated and purified) from natural sources, which can contain natural, non-natural or altered nucleotides, and which can contain a natural, non-natural, or altered internucleotide linkage, such as a phosphoroamidate linkage or a phosphorothioate linkage, instead of the phosphodiester found between the nucleotides of an unmodified oligonucleotide. It is generally preferred that the nucleic acid does not comprise any insertions, deletions, inversions, substitutions, or combinations thereof. However, it may be suitable in some instances, as discussed herein, for the nucleic acid to comprise one or more insertions, deletions, inversions, substitutions, or combinations thereof.

[0094] Preferably, the nucleic acids of the invention are recombinant. As used herein, the term “recombinant” refers to (i) molecules that are constructed outside living cells by joining natural or synthetic nucleic acid segments, or (ii) molecules that result from the replication of those described in (i) above. For purposes herein, the replication can be *in vitro* replication or *in vivo* replication.

[0095] The nucleic acids can be constructed based on chemical synthesis, enzymatic ligation reactions, or combinations thereof using procedures known in the art. See, for example, Green and Sambrook, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 4th Ed. (2012). For example, a nucleic acid can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed upon hybridization (e.g., phosphorothioate derivatives and acridine substituted nucleotides). Examples of modified nucleotides that can be used to generate the nucleic acids include, but are not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil,

dihydrouracil, beta-D-galactosylqueosine, inosine, N⁶-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N⁶-substituted adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N⁶-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, 3-(3-amino-3-N-2-carboxypropyl) uracil, and 2,6-diaminopurine. Alternatively, one or more of the nucleic acids of the invention can be purchased from companies, such as Macromolecular Resources (Fort Collins, CO) and Synthegen (Houston, TX).

[0096] The invention also provides a nucleic acid comprising a nucleotide sequence which is complementary to the nucleotide sequence of any of the nucleic acids described herein or a nucleotide sequence which hybridizes under stringent conditions to the nucleotide sequence of any of the nucleic acids described herein.

[0097] The nucleotide sequence which hybridizes under stringent conditions preferably hybridizes under high stringency conditions. By "high stringency conditions" is meant that the nucleotide sequence specifically hybridizes to a target sequence (the nucleotide sequence of any of the nucleic acids described herein) in an amount that is detectably stronger than non-specific hybridization. High stringency conditions include conditions which would distinguish a polynucleotide with an exact complementary sequence, or one containing only a few scattered mismatches, from a random sequence that happened to have only a few small regions (e.g., 3-10 bases) that matched the nucleotide sequence. Such small regions of complementarity are more easily melted than a full-length complement of 14-17 or more bases, and high stringency hybridization makes them easily distinguishable. Relatively high stringency conditions would include, for example, low salt conditions, high temperature conditions, or combinations thereof, such as provided by about 0.02-0.1 M NaCl or the equivalent, at temperatures of about 50-70 °C. Such high stringency conditions tolerate little, if any, mismatch between the nucleotide sequence and the template or target strand, and are particularly suitable for detecting expression of any of the inventive proteins or chimeric molecules. It is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide.

[0098] The invention also provides a nucleic acid comprising a nucleotide sequence that is about 70% or more, e.g., about 80% or more, about 90% or more, about 91% or more,

about 92% or more, about 93% or more, about 94% or more, about 95% or more, about 96% or more, about 97% or more, about 98% or more, or about 99% or more identical to any of the nucleic acids described herein.

[0099] The nucleic acids of the invention can be incorporated into a recombinant expression vector. In this regard, the invention provides recombinant expression vectors comprising any of the nucleic acids of the invention. For purposes herein, the term “recombinant expression vector” means a genetically-modified oligonucleotide or polynucleotide construct that permits the expression of an mRNA, protein, polypeptide, or peptide by a host cell, when the construct comprises a nucleotide sequence encoding the mRNA, protein, polypeptide, or peptide, and the vector is contacted with the cell under conditions sufficient to have the mRNA, protein, polypeptide, or peptide expressed within the cell. The vectors of the invention are not naturally-occurring as a whole. However, parts of the vectors can be naturally-occurring. The inventive recombinant expression vectors can comprise any type of nucleotides, including, but not limited to DNA and RNA, which can be single-stranded or double-stranded, which can be synthesized or obtained in part from natural sources, and which can contain natural, non-natural or altered nucleotides. The recombinant expression vectors can comprise naturally-occurring, non-naturally-occurring internucleotide linkages, or both types of linkages. Preferably, the non-naturally occurring or altered nucleotides or internucleotide linkages do not hinder the transcription or replication of the vector.

[0100] The recombinant expression vector of the invention can be any suitable recombinant expression vector, and can be used to transform or transfect any suitable host cell. Suitable vectors include those designed for propagation and expansion or for expression or for both, such as plasmids and viruses. The vector can be selected from the group consisting of the pUC series (Fermentas Life Sciences), the pBluescript series (Stratagene, LaJolla, CA), the pET series (Novagen, Madison, WI), the pGEX series (Pharmacia Biotech, Uppsala, Sweden), and the pEX series (Clontech, Palo Alto, CA). Bacteriophage vectors, such as λ GT10, λ GT11, λ ZapII (Stratagene), λ EMBL4, and λ NM1149, also can be used. Examples of plant expression vectors include pBI01, pBI101.2, pBI101.3, pBI121 and pBIN19 (Clontech). Examples of animal expression vectors include pEUK-CI, pMAM, and pMAMneo (Clontech). Preferably, the recombinant expression vector is a viral vector, e.g., a retroviral vector.

[0101] The recombinant expression vectors of the invention can be prepared using standard recombinant DNA techniques described in, for example, Green and Sambrook, *supra*. Constructs of expression vectors, which are circular or linear, can be prepared to contain a replication system functional in a prokaryotic or eukaryotic host cell. Replication systems can be derived, e.g., from ColEI, 2 μ plasmid, λ , SV40, bovine papilloma virus, and the like.

[0102] Desirably, the recombinant expression vector comprises regulatory sequences, such as transcription and translation initiation and termination codons, which are specific to the type of host (e.g., bacterium, fungus, plant, or animal) into which the vector is to be introduced, as appropriate and taking into consideration whether the vector is DNA- or RNA-based.

[0103] The recombinant expression vector can include one or more marker genes, which allow for selection of transformed or transfected hosts. Marker genes include biocide resistance, e.g., resistance to antibiotics, heavy metals, etc., complementation in an auxotrophic host to provide prototrophy, and the like. Suitable marker genes for the inventive expression vectors include, for instance, neomycin/G418 resistance genes, hygromycin resistance genes, histidinol resistance genes, tetracycline resistance genes, and ampicillin resistance genes.

[0104] The recombinant expression vector can comprise a native or nonnative promoter operably linked to the nucleotide sequence encoding the inventive protein or chimeric molecule (including functional portions and functional variants), or to the nucleotide sequence which is complementary to or which hybridizes to the nucleotide sequence encoding the protein or chimeric molecule. The selection of promoters, e.g., strong, weak, inducible, tissue-specific, and developmental-specific, is within the ordinary skill of the artisan. Similarly, the combining of a nucleotide sequence with a promoter is also within the ordinary skill of the artisan. The promoter can be a non-viral promoter or a viral promoter, e.g., a cytomegalovirus (CMV) promoter, an SV40 promoter, an RSV promoter, or a promoter found in the long-terminal repeat of the murine stem cell virus.

[0105] The inventive recombinant expression vectors can be designed for either transient expression, for stable expression, or for both. Also, the recombinant expression vectors can be made for constitutive expression or for inducible expression.

[0106] Another embodiment of the invention further provides a host cell comprising any of the recombinant expression vectors described herein. As used herein, the term "host cell"

refers to any type of cell that can contain the inventive recombinant expression vector. The host cell can be a eukaryotic cell, e.g., plant, animal, fungi, or algae, or can be a prokaryotic cell, e.g., bacteria or protozoa. The host cell can be a cultured cell, an adherent cell or a suspended cell, i.e., a cell that grows in suspension. For purposes of producing a recombinant inventive protein or chimeric molecule, the host cell is preferably a prokaryotic cell, e.g., an *E. coli* cell.

[0107] Also provided by the invention is a population of cells comprising at least one host cell described herein. The population of cells can be a heterogeneous population comprising the host cell comprising any of the recombinant expression vectors described, in addition to at least one other cell, e.g., a host cell which does not comprise any of the recombinant expression vectors. Alternatively, the population of cells can be a substantially homogeneous population, in which the population comprises mainly (e.g., consisting essentially of) host cells comprising the recombinant expression vector. The population also can be a clonal population of cells, in which all cells of the population are clones of a single host cell comprising a recombinant expression vector, such that all cells of the population comprise the recombinant expression vector. In one embodiment of the invention, the population of cells is a clonal population of host cells comprising a recombinant expression vector as described herein.

[0108] The inventive proteins, chimeric molecules (including functional portions and functional variants), nucleic acids, recombinant expression vectors, host cells, and populations of cells can be isolated, purified, or both isolated or purified. The term “isolated” as used herein means having been removed from its natural environment. The term “purified” as used herein means having been increased in purity, wherein “purity” is a relative term, and not to be necessarily construed as absolute purity. For example, the purity can be about 50% or more, about 60% or more, about 70% or more, about 80% or more, about 90% or more, or about 100%. The purity preferably is about 90% or more (e.g., about 90% to about 95%) and more preferably about 98% or more (e.g., about 98% to about 99%).

[0109] The inventive PAs, nucleic acids, recombinant expression vectors, host cells, populations of cells, and compositions, all of which are collectively referred to as “inventive PA materials” hereinafter, can be formulated into a pharmaceutical composition. In this regard, the invention provides a pharmaceutical composition comprising any of the PAs, nucleic acids, recombinant expression vectors, host cells, populations of cells, or compositions described herein, and a pharmaceutically acceptable carrier.

[0110] Preferably, the carrier is a pharmaceutically acceptable carrier. With respect to pharmaceutical compositions, the carrier can be any of those conventionally used and is limited only by chemico-physical considerations, such as solubility and lack of reactivity with the active compound(s), and by the route of administration. The pharmaceutically acceptable carriers described herein, for example, vehicles, adjuvants, excipients, and diluents, are well-known to those skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one which is chemically inert to the active agent(s) and one which has no detrimental side effects or toxicity under the conditions of use.

[0111] The choice of carrier will be determined in part by the particular inventive PA material, as well as by the particular method used to administer the inventive PA material. Accordingly, there are a variety of suitable formulations of the pharmaceutical composition of the invention. The following formulations for parenteral (e.g., subcutaneous, intravenous, intraarterial, intramuscular, intradermal, interperitoneal, intratumoral, and intrathecal) and oral administration are exemplary and are in no way limiting. More than one route can be used to administer the inventive PA materials, and in certain instances, a particular route can provide a more immediate and more effective response than another route.

[0112] Formulations suitable for oral administration can include (a) liquid solutions, such as an effective amount of the inventive PA material dissolved in diluents, such as water or saline; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant. Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and other pharmacologically compatible excipients. Lozenge forms can comprise the inventive PA material in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising

the inventive PA material in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like additionally containing such excipients as are known in the art.

[0113] Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The inventive PA material can be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol or hexadecyl alcohol, a glycol, such as propylene glycol or polyethylene glycol, dimethylsulfoxide, glycerol, ketals such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, poly(ethyleneglycol) 400, oils, fatty acids, fatty acid esters or glycerides, or acetylated fatty acid glycerides with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

[0114] Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters.

[0115] Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl- β -aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (e) mixtures thereof.

[0116] The parenteral formulations will typically contain from about 0.5% to about 2% by weight per volume of the inventive PA material in solution. Preservatives and buffers may be used. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants having a hydrophile-lipophile

balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulations will typically range from about 5% to about 15% by weight. Suitable surfactants include polyethylene glycol sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use.

Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. The requirements for effective pharmaceutical carriers for parenteral compositions are well-known to those of ordinary skill in the art (see, e.g., Lloyd et al. (Eds.) *Remington: The Science and Practice of Pharmacy*, Pharmaceutical Press, 22ND Ed. (2012)).

[0117] For purposes of the invention, the amount or dose of the inventive PA material administered should be sufficient to effect a desired response, e.g., a therapeutic or prophylactic response, in the mammal over a reasonable time frame. For example, the dose of the inventive PA material should be sufficient to inhibit growth of a target cell or treat or prevent cancer in a period of from about 2 hours or longer, e.g., 12 to 24 or more hours, from the time of administration. In certain embodiments, the time period could be even longer. The dose will be determined by the efficacy of the particular inventive PA material and the condition of the mammal (e.g., human), as well as the body weight of the mammal (e.g., human) to be treated.

[0118] Many assays for determining an administered dose are known in the art. An administered dose may be determined *in vitro* (e.g., cell cultures) or *in vivo* (e.g., animal studies). For example, an administered dose may be determined by determining the IC₅₀ (the dose that achieves a half-maximal inhibition of symptoms), LD₅₀ (the dose lethal to 50% of the population), the ED₅₀ (the dose therapeutically effective in 50% of the population), and the therapeutic index in cell culture, animal studies, or combinations thereof. The therapeutic index is the ratio of LD₅₀ to ED₅₀ (i.e., LD₅₀/ED₅₀).

[0119] The dose of the inventive PA material also may be determined by the existence, nature, and extent of any adverse side effects that might accompany the administration of a particular inventive PA material. Typically, the attending physician will decide the dosage of the inventive PA material with which to treat each individual patient, taking into

consideration a variety of factors, such as age, body weight, general health, diet, sex, inventive PA material to be administered, route of administration, and the severity of the condition being treated. By way of example and not intending to limit the invention, the dose of the inventive PA material can be about 0.001 to about 1000 mg/kg body weight of the subject being treated/day, from about 0.01 to about 10 mg/kg body weight/day, about 0.01 mg to about 1 mg/kg body weight/day, from about 1 to about to about 1000 mg/kg body weight/day, from about 5 to about 500 mg/kg body weight/day, from about 10 to about 250 mg/kg body weight/day, about 25 to about 150 mg/kg body weight/day, about 10 mg/kg body weight/day, about 2 mg/kg body weight/day to about 5 mg/kg body weight/day, or about 4 mg/kg body weight/day.

[0120] The inventive PA materials may be assayed for cytotoxicity by assays known in the art. Examples of cytotoxicity assays include a MTT (3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide) assay, as described in Liu et al., *J. Biol. Chem.*, 278: 5227-5234 (2003).

[0121] It is contemplated that the inventive PA materials can be used in methods of treating or preventing a condition such as, for example, cancer. Without being bound to a particular theory or mechanism, it is believed that compositions comprising a combination of the first PA and the second PA as described herein form LF-binding competent heterogeneous oligomers by intermolecular complementation, and specifically kill cancer cells with no or minimal cross-reactivity with normal, non-cancerous cells. In this regard, the invention provides a method of treating or preventing cancer in a mammal, the method comprising administering to the mammal any of the inventive PA materials described herein in an amount effective to treat or prevent cancer in the mammal.

[0122] In an embodiment of the invention, the method further comprises administering to the mammal one or more of any of the effector components described herein. The first PA, the second PA, and the effector component may be administered simultaneously or in any suitable sequence. For example, the method may comprise administering the first PA prior to administering the second PA and administering the effector component after administering the second PA. Alternatively, the method may comprise administering the first PA and the second PA simultaneously and prior to administering the effector component. In a preferred embodiment, the method comprises administering the first PA, the second PA, and the effector component simultaneously.

[0123] In an embodiment, the method further comprises administering one or more immunosuppressive agents to the mammal in an amount effective to decrease the immune response to the inventive PA material. Administering one or more immunosuppressive agents to the mammal may, advantageously, eliminate or reduce an anti-PA material immune response including, for example, the production of anti-PA material antibodies and/or T-cells, that may otherwise undesirably neutralize the cytotoxic activity of the inventive PA material. In this regard, administering one or more immunosuppressive agents to the mammal may, advantageously, increase the amount of inventive PA material that can be given to the mammal which may, in turn, increase the effectiveness of the inventive PA material for treating or preventing a condition, such as, for example, cancer.

[0124] The method may comprise administering the inventive PA material and the one or more immunosuppressive agents simultaneously or sequentially to the mammal. While the method may comprise administering the one or more immunosuppressive agents to the mammal after administering the inventive PA material, preferably the method comprises administering the one or more immunosuppressive agents to the mammal prior to administering the inventive PA material or sequentially with the inventive PA material. Administering the one or more immunosuppressive agents to the mammal prior to administering the inventive PA material advantageously may eliminate or reduce an undesirable immune response to the inventive PA material before the immune system is exposed to the inventive PA material. For example, the method may comprise administering the one or more immunosuppressive agents to the mammal about one to about seven days prior to administering the inventive PA material to the mammal.

[0125] The one or more immunosuppressive agents may comprise any agent suitable for suppressing an immune response to the inventive PA material. In an embodiment, the one or more immunosuppressive agents are selected from the group consisting of pentostatin, cyclophosphamide, and tofacitinib (Hassan et al., *Sci. Transl. Med.*, 5(208):208ra147 (2013); Mossoba et al., *Clin. Cancer Res.*, 17: 3697-3705 (2011); and Onda et al., *J. Immunol.*, 193(1): 48-55 (2014). In an embodiment of the invention, the one or more immunosuppressive agents comprise a combination of pentostatin and cyclophosphamide. Pentostatin and cyclophosphamide may be administered simultaneously or in sequence. For example, pentostatin may be administered before administering cyclophosphamide. Alternatively, cyclophosphamide may be administered prior to administering pentostatin.

[0126] It has also been discovered that the combination of pentostatin and cyclophosphamide provides anti-solid tumor effects when used in the absence of any other pharmaceutically active agents. Accordingly, an embodiment of the invention provides a method of treating or preventing a solid tumor in a mammal, the method comprising administering to the mammal pentostatin and cyclophosphamide in an amount effective to treat or prevent the solid tumor in the mammal. In a preferred embodiment, the method comprises administering pentostatin and cyclophosphamide in the absence of any other pharmaceutically active agents (e.g., in the absence of any other anti-cancer agents). For example, the method comprises administering pentostatin and cyclophosphamide in the absence of any of the inventive PA materials described herein. In an especially preferred embodiment, the method comprises administering pentostatin and cyclophosphamide (and a pharmaceutically inactive carrier) alone. Pentostatin and cyclophosphamide may be administered simultaneously or in sequence as described herein with respect to other aspects of the invention.

[0127] The terms “treat” and “prevent” as well as words stemming therefrom, as used herein, do not necessarily imply 100% or complete treatment or prevention. Rather, there are varying degrees of treatment or prevention of which one of ordinary skill in the art recognizes as having a potential benefit or therapeutic effect. In this respect, the inventive methods can provide any amount or any level of treatment or prevention of cancer in a mammal. Furthermore, the treatment or prevention provided by the inventive method can include treatment or prevention of one or more conditions or symptoms of the disease, e.g., cancer, being treated or prevented. Also, for purposes herein, “prevention” can encompass delaying the onset of the disease, or a symptom or condition thereof.

[0128] With respect to the inventive methods, the cancer can be any cancer, including any of adrenal gland cancer, sarcomas (e.g., synovial sarcoma, osteogenic sarcoma, leiomyosarcoma uteri, angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma, myxoma, rhabdomyoma, fibroma, lipoma, and teratoma), lymphomas (e.g., small lymphocytic lymphoma, Hodgkin lymphoma, and non-Hodgkin lymphoma), hepatocellular carcinoma, glioma, head cancers (e.g., squamous cell carcinoma, e.g., oral squamous cell carcinoma), neck cancers (e.g., squamous cell carcinoma), acute lymphocytic cancer, leukemias (e.g., hairy cell leukemia, myeloid leukemia (acute and chronic), lymphatic leukemia (acute and chronic), prolymphocytic leukemia (PLL), myelomonocytic leukemia (acute and chronic), and lymphocytic leukemia (acute and chronic)), bone cancer (osteogenic

sarcoma, fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor, chordoma, osteochondroma (osteocartilaginous exostoses), benign chondroma, chondroblastoma, chondromyxoid fibroma, osteoid osteoma, and giant cell tumors), brain cancer (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma (pinealoma), glioblastoma multiforme, oligodendroglioma, schwannoma, and retinoblastoma), fallopian tube cancer, breast cancer, cancer of the anus, anal canal, or anorectum, cancer of the eye, cancer of the intrahepatic bile duct, cancer of the joints, cancer of the neck, gallbladder, or pleura, cancer of the nose, nasal cavity, or middle ear, cancer of the oral cavity, cancer of the vulva (e.g., squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, and fibrosarcoma), myeloproliferative disorders (e.g., chronic myeloid cancer), colon cancers (e.g., colon carcinoma), esophageal cancer (e.g., squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, and lymphoma), cervical cancer (cervical carcinoma and pre-invasive cervical dysplasia), gastric cancer, gastrointestinal carcinoid tumor, hypopharynx cancer, larynx cancer, liver cancers (e.g., hepatocellular carcinoma, cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, and hemangioma), lung cancers (e.g., bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, and adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, chondromatous hamartoma, small cell lung cancer, non-small cell lung cancer, and lung adenocarcinoma), malignant mesothelioma, skin cancer (e.g., melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, nevi, dysplastic nevi, lipoma, angioma, dermatofibroma, and keloids), multiple myeloma, nasopharynx cancer, ovarian cancer (e.g., ovarian carcinoma (serous cystadenocarcinoma, mucinous cystadenocarcinoma, endometrioid carcinoma, and clear cell adenocarcinoma), granulosa-theca cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, and malignant teratoma), pancreatic cancer (e.g., ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, and VIPoma), peritoneum, omentum, mesentery cancer, pharynx cancer, prostate cancer (e.g., adenocarcinoma and sarcoma), rectal cancer, kidney cancer (e.g., adenocarcinoma, Wilms tumor (nephroblastoma), and renal cell carcinoma), small intestine cancer (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, and fibroma), soft tissue cancer, stomach cancer (e.g., carcinoma, lymphoma, and leiomyosarcoma), testicular cancer (e.g., seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, Leydig cell tumor, fibroma, fibroadenoma,

adenomatoid tumors, and lipoma), cancer of the uterus (e.g., endometrial carcinoma), thyroid cancer, and urothelial cancers (e.g., squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma, ureter cancer, and urinary bladder cancer). In an embodiment, the cancer is a solid tumor. In a preferred embodiment, the solid tumor is an oral squamous carcinoma tumor, a melanoma tumor, a lung tumor, or an ovarian tumor. Without being bound to a particular theory or mechanism, it is believed that the inventive PA materials, wherein one or more of amino acid residues I207, I210, I656, Y681, and L687 are substituted as described herein, are toxic to tumor endothelial cells that express CMG2. Accordingly, in an embodiment of the invention, the endothelial cells of the solid tumor express CMG2. Without being bound to a particular theory or mechanism, it is believed that the inventive PA materials, wherein one or more of amino acid residues E654, R659, and M662 are substituted as described herein, are toxic to cells that express TEM8. Accordingly, in an embodiment of the invention, the cells of the tumor express TEM8.

[0129] As used herein, the term “mammal” refers to any mammal, including, but not limited to, mammals of the orders Rodentia, such as mice and hamsters; Logomorpha, such as rabbits; Carnivora, including Felines (cats) and Canines (dogs); Artiodactyla, including Bovines (cows) and Swines (pigs); Perssodactyla, including Equines (horses); Primates, Ceboids, or Simoids (monkeys) or Anthropoids (humans and apes). In a preferred embodiment, the mammal is a human, mouse, cat or dog.

[0130] Also provided is a method of inhibiting the growth of a target cell comprising contacting the cell with any of the inventive PA materials described herein, in an amount effective to inhibit growth of the target cell. The growth of the target cell may be inhibited by any amount, e.g., by about 10% or more, about 15% or more, about 20% or more, about 25% or more, about 30% or more, about 35% or more, about 40% or more, about 45% or more, about 50% or more, about 55% or more, about 60% or more, about 65% or more, about 70% or more, about 75% or more, about 80% or more, about 85% or more, about 90% or more, about 95% or more, or about 100%. The target cell may be provided in a biological sample. A biological sample may be obtained from a mammal in any suitable manner and from any suitable source. The biological sample may, for example, be obtained by a blood draw, leukapheresis, tumor biopsy necropsy, or combinations thereof. The contacting step can take place *in vitro* or *in vivo* with respect to the mammal. Preferably, the contacting is *in vitro*.

[0131] In an embodiment of the invention, the target cell is a cancer cell. Preferably, the target cell is a cancer cell. The target cell may be a cancer cell of any of the cancers described herein. In an embodiment of the invention, the target may express a cell surface molecule. The cell surface marker may be any cell surface molecule described herein with respect to other aspects of the invention.

[0132] The method may comprise contacting the cell with the inventive PA material and the effector component simultaneously or sequentially. In an embodiment of the invention, the method comprises contacting the cell with the inventive PA material prior to contacting the cell with the effector component. In another embodiment, the method comprises contacting the cell with the effector component prior to administering the inventive PA material. Preferably, the method comprises contacting the cell with the effector component and the inventive PA simultaneously.

[0133] The following examples further illustrate the invention but, of course, should not be construed as in any way limiting its scope.

EXAMPLES 1-3

[0134] The following materials and methods were employed in the experiments described in Examples 1-3.

Mutagenic PCR

[0135] Mutations were introduced by overlap polymerase chain reaction (PCR) with NNS codons into the plasmid pYS52 (a derivative of *B. anthracis* expression vector pYS5 (Singh et al., *J. Biol. Chem.*, 264: 19103-19107 (1989)), which contains a synthetic DNA sequence coding for domain II of PA with additional unique restriction sites. PHUSION High-Fidelity DNA polymerase (New England Biologicals, Ipswich, MA) was used for mutagenic PCR reactions. Flanking primers used for all reactions were SEQ ID NO: 16 (PA-250) and SEQ ID NO: 17 (Pa-SwaI-Rev). Specific primers were SEQ ID NO: 18 (R178X mut), SEQ ID NO: 19 (R178X rev), SEQ ID NO: 20 (R200X mut), SEQ ID NO: 21 (R200X rev), SEQ ID NO: 22 (I207X mut), SEQ ID NO: 23 (I207/210X rev), SEQ ID NO: 24 (I210X mut), SEQ ID NO: 25 (K214X mut), and SEQ ID NO: 26 (K214X rev). Mutagenic primers were combined with PA-SwaI-Rev and reverse primers with PA-250 for the first round of PCR. The products were gel-purified and the complementary PCR fragments were extended to full length by 10 cycles of PCR before addition of PA-250 and PA-SwaI-rev and an additional 35

cycles. These mutagenic inserts were gel-purified and the inserts, along with pYS52, were digested with PstI and HindIII. The digestion products were ligated overnight and transformed into chemically competent *E. coli* MC1061, which were plated on lysogeny broth (LB)-agar plates containing 100 µg/mL ampicillin. Colony PCR using PA-250 and PA-SwaI-Rev primers was performed to identify positive clones. Colonies were screened using primer PA-250 to sequence the plasmid and identify clones having all possible substitutions. Selected clones were then grown overnight in LB containing 100 µg/mL ampicillin and the plasmids were extracted by mini-scale preparation.

Expression of protein library

[0136] Mini-prepared plasmids from *E. coli* MC1061 were transformed into chemically competent *E. coli* SCS110, which is *dam*⁻ and *dcm*⁻. The purified, non-methylated plasmids from SCS110 were then transformed into an electrocompetent *B. anthracis* BH480 strain, which was plated on LB-agar containing 20 µg/mL kanamycin. BH480 is an avirulent large plasmids-cured, sporulation-defective *B. anthracis* strain with eight proteases deleted, serving as an efficient host for recombinant protein expression (Pomerantsev et al., *Protein Expr. Purif.*, 80: 80-90 (2011)). Single colonies were grown overnight in 5 mL FA medium containing 20 µg/mL kanamycin (Pomerantsev et al., *supra*). The supernatants containing the mutant PA proteins were sterilized by centrifugation and concentrated ~10 fold using AMICON ULTRA-4 (30K) Centrifugal Filter Devices (Millipore Corp., Billerica, MA). The supernatants were analyzed by native gel electrophoresis, stained with coomassie blue dye, and the protein concentrations were estimated by densitometry to compare the supernatant bands to a sample of purified PA. This was performed twice on each protein.

PA variant screen

[0137] RAW264.7 macrophages and murine melanoma B16-BL6 cells were grown in Dulbecco's Modified Eagle Medium (Life Technologies, Grand Island, NY) supplemented with fetal bovine serum to 10% (Invitrogen) and gentamycin at 50 µg/mL (Invitrogen) at 37 °C in a tissue culture incubator with 5% CO₂.

[0138] To test the PA variants for a loss of function, RAW 264.7 macrophages were plated in a 96-well plate at 10⁵ cells/well and grown overnight. The next day, the PA variants were added at a concentration of 500 ng/mL and LF at a concentration of 100 ng/mL. The cells were incubated for 20 hours (h) and MTT (3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-

2H-tetrazolium bromide, Sigma, St. Louis, MO) was added at 500 µg/mL for the final hour. The medium was aspirated and the oxidized MTT was solubilized in 91% isopropanol containing 0.5% SDS and 0.038 M hydrochloric acid, then read at 570 nm using a SPECTRAMAX 190 plate reader (Molecular Devices, Sunnyvale, CA). The absorbance was used to determine percent survival compared to an untreated control. PA variants that showed less toxicity than the original constructs PA-R200A and PA-I210A were subjected to a double-agent gain-of-function test. RAW 264.7 macrophages were seeded as above and respective PA variants (250 ng/mL) combined with PA-R200A or PA-I210A (250 ng/mL) and 100 ng/mL LF were added. The cells were incubated for 6 h and viability was measured as above.

Creation and cytotoxicity of PA-L1-I207R

[0139] Site-directed mutagenesis was used to introduce mutation I207R into pYS5-L1 ff, an overexpression plasmid for PA-L1, in which the furin cleavage sequence RKKR (SEQ ID NO: 4) (residues 164-167) was replaced with a MMP substrate sequence GPLGMLSQ (SEQ ID NO: 6) (Liu et al., *Cancer Res.*, 60: 6061-6067 (2000)). Primers used were SEQ ID NO: 27 (I207R sense), SEQ ID NO: 28 (I207R antisense). Sense and anti-sense primers were used with the QUIKCHANGE LIGHTNING Kit (Agilent, Santa Clara, CA) according to manufacturer's recommendations and transformed into chemically competent *E. coli* XL-10 GOLD cells. The plasmids were mini-prepared, sequenced, and transformed into *E. coli* SCS110 before transformation into BH480 for expression and purification. The coding sequence for PA-L1-I207R was the nucleotide sequence of SEQ ID NO: 29. Cytotoxicities of PA-L1-I207R and PA-L1-I210A were tested singly and in combination with PA-U2-R200A (with the furin cleavage sequence replaced with an uPA substrate sequence PGSGRSA (SEQ ID NO: 9) (Liu et al., *J. Biol. Chem.*, 276, 17976-17984 (2001))) in the presence of 30 ng/mL FP59. B16-BL6 cells were incubated with the indicated toxins in 96-well plates for 48 h, and cell viabilities were determined by MTT assay as described above.

Protein purification

[0140] PA-L1-I210A, PA-L1-I207R, PA-U2-R200A, LF and FP59 were expressed using pYS5-based expression plasmids from *B. anthracis* BH480 strain. The recombinant proteins secreted into culture supernatants were purified as described previously (Pomerantsev et al., *supra*; Liu et al., *Cell. Microbiol.*, 9: 977-987 (2007)). In brief, the expression plasmid

transformed BH480 strains were grown in FA medium with 10 µg/ml kanamycin for 12 h at 37 °C. The proteins secreted into the culture supernatants were precipitated on PHENYL-SEPHAROSE 6 FAST FLOW resin (low substitution, GE Healthcare Life Sciences, Pittsburg, PA) (30 ml per liter supernatant) in the presence of 2 M ammonium sulfate in rotating bottles. The resin was collected on a porous plastic funnel and washed with wash buffer (1.5 M ammonium sulfate, 10 mM Tris-HCl (pH 8.0), 1 mM EDTA) and the proteins were eluted using elution buffer (0.3 M ammonium sulfate, 10 mM Tris-HCl (pH 8.0), 1 mM EDTA). The eluted proteins were precipitated by adding 2 M ammonium sulfate. The precipitate was collected by centrifugation, resuspended and dialyzed in 10 mM Tris-HCl (pH 8.0), 1 mM EDTA. The toxin proteins were further purified by chromatography on Q-SEPHAROSE FAST FLOW column (GE Healthcare Life Sciences) and eluted with a 0–0.5 M NaCl gradient in 20 mM Tris-HCl, 0.5 mM EDTA (pH 8.0). The toxin proteins were further purified by SEPHACRYL S-200 high resolution gel filtration (GE Healthcare Life Sciences) using 10 mM Tris-HCl (pH 8.0), 100 mM NaCl, and 0.5 mM ethylenediaminetetraacetic acid (EDTA) to one prominent band at the expected molecular mass.

In vivo toxicity of PA mutants

[0141] C57BL/6J male and female mice (10 to 12-week-old) were injected intraperitoneally with 20 µg PA-L1-I207R or PA-L1-I210A along with 10 µg FP59 at 0, 24, and 48 hours and checked twice daily for signs of malaise and mortality for two weeks following the first injection. Mice were euthanized at the end of experiment. All animal studies were carried out in accordance with protocols approved by the National Institute of Allergy and Infectious Diseases Animal Care and Use Committee.

In vivo anti-tumor study

[0142] Twelve-week-old female C57BL/6J mice (Jackson Laboratory, Bar Harbor, Maine) were injected with 5×10^5 B16-BL6 cells in the mid-scapular subcutis. B16-BL6 melanoma cells were authenticated by continual assessment of cellular morphology at both low and high magnifications. Eight days after injection, established tumors were measured with digital calipers (FV Fowler Company, Inc., Newton, MA). Tumor weights were estimated with the longest and shortest tumor dimensions in the formula: tumor weight (mg) = (length in mm $\times$ width in mm²) $\times$ 0.5 (Geran et al., *Cancer Chemother. Rep.*, 3: 1-103

(1972)). Tumor-bearing mice were randomized into groups and injected intraperitoneally on study days 0 (eight days after tumor cell injection), 2, 4, 7, and 9 with a 200 μ L PBS solution containing the PA variant proteins and LF, at doses shown in Figure 5. Mice were weighed and the tumors were measured before each injection. The study was ended on day 10 when tumors in the PBS alone control group reached 10% of the body weights, a condition requiring euthanasia according to the animal study protocol.

Statistical analysis

[0143] Statistical analysis was done with unpaired student's t test using EXCEL software. Survival curves were analyzed using log-rank test (Mantel-Cox) using GRAPHPAD PRISM software.

EXAMPLE 1

[0144] This example demonstrates the mutagenesis and expression of PA mutants.

[0145] To select PA variants with activity strictly dependent on intermolecular complementation for LF-binding, a PA variant library with NNS (N = any nucleotide, S = C or G) codons at LF-binding subsite residue R200 on the counterclockwise-side monomer was constructed, and the residues R178, I207, I210, and K214 on the clockwise-side monomer (Figure 1). While DNA coding sequences for all of the possible PA variants were isolated, some proteins were not expressed, or were expressed only at very low levels, from the corresponding transformed avirulent *B. anthracis* BH480 strains. Out of a theoretical library of 95 mutants, 79 were successfully expressed and supernatants containing the secreted PA variants were prepared for initial testing. PA proteins were usually expressed at high levels by the correspondingly transformed BH480 strains, often reaching levels above 50% of the total protein in the culture supernatant. Of the sixteen proteins that were not expressed, nine were mutants of K214 (C, D, E, F, G, P, S, W, Y), four were mutants at I207 (D, K, P, S), two were mutants of I210 (G, N), and one was an R178 mutant (P). K214 lies at the end of an α -helix and the side chain makes no close contact with any other residues on PA. It is not clear why so many mutants at this position could not be expressed.

EXAMPLE 2

[0146] This example demonstrates that PA-I207R is an improved clockwise-side monomer variant.

[0147] The sterilized supernatants containing PA proteins were screened for activity relative to the original versions of the intermolecular complementing, PA-R200A or PA-I210A, as appropriate using cytotoxicity assays. In the screen of counterclockwise-side monomer mutants, ten mutant PA-R200X proteins (C, D, E, G, I, M, P, S, V, and W) were initially identified as being less intrinsically toxic than PA-R200A. However, later detailed characterization found that these PA variants also showed reduced intermolecular complementation with PA-I210, resulting in less LF-induced cytotoxicity. As an example, although PA-R200C was slightly less toxic than PA-R200A, it was also slightly less effective in complementing with PA-I210A to promote killing of RAW264.7 cells. Therefore, none of the counterclockwise-side monomer PA variants were found to be significantly superior to the original PA-R200A.

[0148] In screens for the clockwise-side monomer variants PA-R178X, PA-I207X, PA-I210X, and PA-K214X, eight proteins with greatly reduced inherent cytotoxicity to RAW264.7 macrophages were identified: I207R, I207W, I210D, I210E, I210K, I210Q, I210R, and I210S (Figure 2). These were tested in combination with R200A and R200C for their ability in intermolecular complementation (Figure 3B). All of these clockwise-side mutants could be complemented by PA-R200A and to a lesser extent by PA-R200C. Among these PA variants, PA-I207R behaved the best in complementing with both PA-R200A and PA-R200C to achieve killing of RAW264.7 cells. Therefore, PA-I207R was identified as an improved clockwise-side monomer variant for the intermolecular complementation PA system, displaying very low cytotoxicity when used singly.

[0149] To further characterize PA-I207R, the MMP-activated variant, i.e., PA-L1-I207R (SEQ ID NO: 30) was generated and the protein purified. The new combination of PA-L1-I207R and PA-U2-R200A was compared with the original combination of PA-L1-I210A and PA-U2-R200A for cytotoxicity towards mouse melanoma B16-BL6 cells, which express high levels of both MMPs and uPA. PA-L1-I207R showed similar activity as PA-L1-I207A in complementing with PA-U2-R200A to kill B16-BL6 cells in the presence of FP59, a LF fusion effector protein that kills cells in a PA-dependent manner (Arora et al., *Infect. Immun.*, 62: 4955-4961 (1994)) (Figure 4A). Remarkably, the single component PA-L1-I207R showed no cytotoxicity, whereas the original counterpart PA-L1-I210A displayed moderate

cytotoxicity to B16-BL6 cells when used singly ($IC_{50} = 200$ ng/mL) in these assays (Figure 4A). PA-U2-R200A also showed no cytotoxicity when used singly in these settings (Figure 4A).

[0150] The toxicity of PA-L1-I207R and PA-L1-I210A was further compared when administered with FP59 to C57BL/6J mice. It was found that PA-L1-I207R was much less toxic than PA-L1-I210A in that all of the mice challenged with 3 doses of 20 μ g PA-L1-I207R and 10 μ g FP59 survived whereas all of the mice challenged with three doses of 20 μ g PA-L1-I210A and 10 μ g FP59 succumbed to the challenges within a week ($P = 0.0007$, log-rank test (Mantel-Cox)) (Figure 4B).

EXAMPLE 3

[0151] This example demonstrates that the combination of PA-L1-I207R and PA-U2-R200A provides high efficacy in antitumor activity.

[0152] To further evaluate the anti-tumor activity of the new PA variant, a side-by-side comparison of the new combination, i.e., PA-L1-I207R (SEQ ID NO: 30) plus PA-U2-R200A (SEQ ID NO: 32), and the original combination of PA-L1-I210A and PA-U2-R200A in the treatment of B16-BL6 syngeneic tumors in C57BL/6J mice was performed. PA-U2-R200A was encoded by the nucleotide sequence of SEQ ID NO: 31. The tumor-bearing mice were treated with either PBS, low doses of the new combination or the original combination of the PA variants plus LF (7.5 μ g/7.5 μ g/5 μ g), or high doses of each combination plus LF (22.5 μ g/22.5 μ g/15 μ g). All of the toxin-treated groups showed significant anti-tumor activities compared to the PBS control group (Figure 5) ($P < 0.0001$ for all toxin-treated groups, Student's *t* test). The new combination of PA-L1-I207R and PA-U2-R200A showed significantly higher anti-tumor activity than the original combination of PA-L1-I210A and PA-U2-R200A (Figure 5) ($P = 0.0326$ for the two high dose groups at day 10, Student's *t* test). A mortality of 20% was observed in the PBS group before termination of the experiment at day 10, with deaths apparently due to the high tumor burden. The groups receiving the low doses of both combinations had 10% mortality, whereas in the high dose groups the new combination appeared to be safer, with a 10% mortality versus 30% mortality in the original combination group (although this difference was not statistically significant).

EXAMPLES 4-10

[0153] The following materials and methods were employed in the experiments described in Examples 4-10.

Proteins and reagents

[0154] Recombinant PA variants and LF proteins were purified from supernatants of BH480, an avirulent, sporulation-defective, protease-deficient *B. anthracis* strain, as described previously (Gupta et al., *PLoS ONE*, 3: e3130 (2008); Pomerantsev et al., *supra*). FP59 is a fusion protein of LF amino acids 1–254 and the catalytic domain of *Pseudomonas aeruginosa* exotoxin A that kills cells by ADP-ribosylation of eukaryotic elongation factor-2 after delivery to cytosol by PA. The LF and FP59 used here contain the native amino-terminal sequence AGG (Gupta et al., *supra*). MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) and pentostatin (SML0508-25 mg) were from Sigma (Atlanta, GA). Cyclophosphamide (NDC10019-957-01) was from Baxter Healthcare (Deerfield, IL).

Cells and cytotoxicity assay

[0155] All cultured cells were grown at 37 °C in a 5% CO₂ atmosphere. Murine B16-BL6 melanoma cells, LL3 Lewis lung carcinoma cells, human lung carcinoma A549 cells, and colorectal carcinoma Colo205 cells were cultured in DMEM (Dulbecco's Modified Eagle Medium) supplemented with 10% fetal bovine serum. Mouse lung endothelial cells and tumor endothelial cells from B16-BL6 melanomas were isolated following the protocol for lung endothelial cell isolation (Reynolds et al., *Methods Mol. Med.*, 120: 503-509 (2006)). Briefly, mouse lungs and B16-BL6 tumors were digested with type I collagenase and plated on gelatin and collagen-coated flasks. The cells were then subjected to sequential negative sorting by magnetic beads coated with a sheep anti-rat antibody using a Fc Blocker (rat anti-mouse CD16/CD32, Cat. 553142, BD Pharmingen, San Diego, CA) to remove macrophages and positive sorting by magnetic beads using an anti-intermolecular adhesion molecule 2 (ICAM2 or CD102) antibody (Cat. 553326, rat anti-mouse CD102, BD Pharmingen) to isolate endothelial cells. Endothelial cells were cultured in DMEM supplemented with 20% fetal bovine serum, endothelial cell growth supplement (30 mg in 500 mL DMEM) (E2759-15 mg, Sigma), heparin (50 mg in 500 mL DMEM) (H3149-100 KU, Sigma).

[0156] For cytotoxicity assays, cells grown in 96-well plates (50% confluence) were incubated with various concentrations of PA or PA variant proteins combined with 500 ng/mL LF for 72 h. In some experiments, cells were incubated with various concentrations of pentostatin or cyclophosphamide or their combination for 72 h. Cell viabilities were then assayed by MTT as described previously (Liu et al., *J. Biol. Chem.*, 278: 5227-5234 (2003)), expressed as % of MTT signals of untreated cells.

MEK cleavage assay

[0157] Tumor endothelial cells grown in 12-well plate were incubated with indicated concentrations of PA-L1 plus LF for 3 h at 37 °C, then washed three times with Hank's Balanced Salt Solution (Biofluids, Rockville, MD) to remove unbound toxins. Cells were then lysed in modified radioimmunoprecipitation assay (RIPA) lysis buffer containing protease inhibitors (Liu et al., *J. Biol. Chem.*, 278: 5227-5234 (2003)) and lysates were subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and Western blotting to detect MEK2 cleavage using anti-MEK2 (N-terminus) antibody (sc-524, Santa Cruz Biotechnology, Santa Cruz, CA).

Gene expression

[0158] Endothelial cells cultured in 12-well plates were treated with or without PA-L1/LF for 24 h, total RNA was then prepared using TRIZOL reagent (Invitrogen, Carlsbad, CA). Single-strand cDNA was synthesized using reverse transcriptase reaction kit following the manufacturer's manual (Invitrogen). Expression changes of the selected key genes involved in glucose uptake, glycolysis, tricarboxylic acid cycle, glutaminolysis, and lipid synthesis were analyzed by real-time quantitative PCR using a SYBR Green PCR Mastermix kit.

Oxygen consumption rates and extracellular acidification rates

[0159] Metabolic activities of tumor endothelial cells were assessed in an XF24 Extracellular Flux analyzer (Seahorse Bioscience, North Billerica, MA). Tumor cells and tumor endothelial cells grown to confluence in 24-well XF24 tissue culture plates were incubated with or without PA-L1/LF (1 µg/mL each) in pentaplicates for 24 h. Cells were changed into fresh unbuffered serum-free DMEM with 200 mM GLUTAMAX-1 medium, 25

mM D-glucose, pH 7.4, and equilibrated in the medium for 1 h. Real-time extracellular acidification rates (ECARs) and oxygen consumption rates (OCRs) were then measured at 37 °C under basal conditions and conditions following sequential additions of oligomycin (0.5 μ M), FCCP (carbonyl cyanide p-trifluoromethoxyphenylhydrazone) (0.5 μ M), and antimycin A (1 μ M). ECARs and OCRs were normalized to 50 mg total protein in cell lysates. Adenosine triphosphate (ATP) production-coupled OCR is calculated as the difference between basal OCR and OCR after addition of oligomycin. Spare respiratory capacity (SRC) is defined as the difference between the OCR following FCCP addition and the OCR under basal condition. Maximal respiration (MR) is defined as the difference between the OCR following FCCP addition and the OCR following oligomycin addition. Cellular ATP levels were measured using ATPLITE 1STEP kit (PerkinElmer, Boston, MA).

Mice and tumor studies

[0160] TEM8- and CMG2-null mice were generated previously (Liu et al., *Proc. Natl. Acad. Sci. USA*, 106: 12424-12429 (2009)). TEM8- and CMG2-null mice were also crossed with athymic nude (*Foxn1^{nu/nu}*) mice (Jackson Laboratory, Bar Harbor, Maine) to generate athymic nude TEM8- and CMG2-null mice, which were used to establish human tumor xenografts. Various tissue-specific CMG2-null mice, including *Cmg2(EC)^{-/-}*, *Cmg2(SM)^{-/-}*, *Cmg2(Mye)^{-/-}* mice, and the tissue-specific CMG2-expressing mice, including *Cmg2^{EC}* and *Cmg2SM* mice were generated as described previously (Liu et al., *Proc. Natl. Acad. Sci. USA*, 106: 12424-12429 (2009); Liu et al., *Cell Host Microbe*, 8: 455-462 (2010); Liu et al., *Nature*, 501: 63-68 (2013)). (Full descriptions of the genotypes are provided in the legends of Figures 8A-8E. For tumor studies, 10-14-week-old male and female mice were used. To grow syngeneic tumors, 5×10^5 cells/mouse B16-BL6 melanoma cells or LL3 lung carcinoma cells were injected in the mid-scapular subcutis of the pre-shaved mice with indicated genotypes. For human tumor xenografts, 1×10^7 cells/mouse Colo205 colorectal carcinoma cells or A549 lung carcinoma cells were injected intradermally into athymic nude mice having the indicated TEM8 or CMG2 genotypes. Tumors were measured with digital calipers (FV Fowler Company, Inc., Newton, MA) and tumor weights were estimated with the length, width, and height tumor dimensions using formulas: tumor weight (mg) = $\frac{1}{2}(\text{length in mm} \times \text{width in mm}^2)$ or $\frac{1}{2}(\text{length in mm} \times \text{width in mm} \times \text{height in mm})$. Tumor-bearing mice were randomized into groups and injected intraperitoneally following schedules indicated in the figures, with PBS, the engineered toxins, a PC regimen, or a

combined therapy. Mice were weighed and the tumors measured before each injection. All animal studies were carried out in accordance with protocols approved by the National Institute of Allergy and Infectious Diseases Animal Care and Use Committee.

Visualization of blood vessels with a lipophilic carbocyanine dye DiI

[0161] The procedure was previously described (Li et al., *Nat. Protoc.*, 3: 1703-1708 (2008)). In brief, B16-BL6 tumor-bearing mice treated with three doses of 30 μ g PA-L1 plus 15 μ g LF or PBS were euthanized by CO₂ inhalation, followed immediately by sequential cardiac perfusion using PBS, DiI dye (Sigma), and 4% paraformaldehyde. Frozen tissue sections were then prepared for fluorescent microscopy to visualize vasculatures of tumors and various normal tissues. For tumor blood vessel quantifications, blood vessels were counted in five random views (11 mm²/view) from each tumor sample (n=3 for each treatment group).

Measurement of toxin-neutralizing antibodies

[0162] B16-BL6 or LL3 tumor-bearing mice from various treatment groups were terminally bled and sera prepared. To titrate toxin neutralizing antibodies in the sera, RAW264.7 cells grown in 96-well plates were incubated with 100 ng/mL PA plus 100 ng/mL LF (amounts that kill >95% of the cells) in the presence of various dilutions of the sera for 5 h, followed by MTT assay to determine cell viabilities as described above.

Flow Cytometry

[0163] Spleens from naïve mice and the B16-BL6 melanoma-bearing mice from the groups treated with PBS, PC regimen, IC2-PA/LF, or the combined PC and the toxin were dissected and weighed after the second round treatments as shown in Figures 15A-15D. Splenocytes were isolated, counted, and stained with fluorochrome-conjugated mAbs anti-CD45R APC-Cy7 (Cat. No. 552094, BD Pharmingen), anti-CD4 APC (Cat. 553051, BD Pharmingen), anti-CD8 PE (Cat. 553033, BD Pharmingen), anti-CD11b PerCP-Cy5 (Cat. 550993, BD Pharmingen), and anti-Gr-1 FITC (Cat. 553127, BD Pharmingen), or anti-IgD FITC (Cat. 553439, BD Pharmingen), anti-IgM PE (Cat. 553409, BD Pharmingen), and anti-CD27 PerCP-Cy5 (Cat. 563603, BD Pharmingen). The cells were analyzed using a BD FACSCanto Flow Cytometer and percentages of each cell population positive for indicated

immune cell markers were obtained. Cell numbers positive for each immune cell marker were obtained by: total splenocytes $\times$ % of the marker positive cells.

Statistical analysis

[0164] Statistical analysis was done with unpaired Student's t test using EXCEL software. Survival curves were analyzed using log-rank test (Mantel-Cox) using GRAPHPAD PRISM software.

EXAMPLE 4

[0165] This example demonstrates that CMG2 and TEM8 in tumor stromal compartments are not important for tumor growth.

[0166] Without being bound to a particular theory or mechanism, it is believed that the angiogenic process is necessary for tumor growth. To directly assess the roles of the anthrax toxin receptors CMG2 and TEM8 in tumor angiogenesis, the growth rates of three different solid tumors in previously described TEM8- and CMG2-null mice (Liu et al., *Proc. Natl. Acad. Sci. USA*, 106: 12424-12429 (2009)) were measured. The tumors evaluated were human lung carcinoma A549 xenografts and the syngeneic mouse LL3 Lewis lung carcinoma and B16-BL6 melanoma (Figures 6A, 6B, 6C, 6F, 6G, and 6H). Consistently, all three tumors grew as rapidly in CMG2-null mice as in their littermate control mice, indicating that CMG2 expression in tumor stromal compartments (e.g., endothelial cells and inflammatory cells) is not required for tumor growth (Figures 6A, 6B, and 6C). No differences in body weight were observed between the tumor-bearing littermate *Cmg2*^{+/+}, *Cmg2*^{+/-}, and *Cmg2*^{-/-} mice (Figures 6F-6H).

[0167] In preliminary studies, it was observed that tumors grew more slowly in TEM8-null mice than in their littermate controls (Figures 12A, 12B, 12C, and 12D). However, it was found that nearly all *Tem8*^{-/-} mice progressively developed misaligned overgrown incisor teeth (malocclusion) beginning at the time of weaning (3-weeks old), causing these mice to have difficulty in chewing the hard food that was routinely provided (Figure 12D). Consequently, the *Tem8*^{-/-} mice became malnourished, reflected in lower body weights associated with lack of subcutaneous fat deposition (Figures 12E, 12F, 12G, 12H, and 12I). The malnourished phenotypes became more pronounced as the *Tem8*^{-/-} mice aged, leading to about 50% mortality at one-year of age. It was found that the *Tem8*^{-/-} mice were completely rescued from malnourishment after providing soft food (Nutra-Gel from Bio-Serv,

Frenchtown, NJ) (Figures 12E, 12F, 12G, 12H, and 12I). Thus, the cause of the malnourished phenotype of *Tem8*^{-/-} mice was their inability to obtain sufficient amounts of nutrients. Furthermore, the frequency of malocclusion seen in *Tem8*^{-/-} mice was greatly decreased by soft food feeding (Figure 12E). Therefore, TEM8 is essential in maintaining mouse tooth function required for the gnawing and chewing of hard food.

[0168] To investigate whether the decreases in tumor growth in *Tem8*^{-/-} mice observed above were due to the malnourished phenotype, the syngeneic LL3 lung carcinomas and B16-BL6 melanomas were then grown in the soft food-fed, body weight-corrected *Tem8*^{-/-} mice. Notably, no significant differences in tumor growth were now observed between littermate *Tem8*^{-/-} and *Tem8*^{+/+} mice (Figures 6D, 6E, 6I, and 6J). Taken together, the results above demonstrate that expression of neither CMG2 nor TEM8 in stromal compartments is important for tumor growth and, by implication, tumor angiogenesis.

EXAMPLE 5

[0169] This example demonstrates that engineered anthrax lethal toxins block tumor growth through host CMG2.

[0170] To investigate the anti-tumor mechanisms of engineered lethal toxins, LL3 carcinoma-bearing mice and B16-BL6 melanoma-bearing mice were treated systemically with PA-L1 plus LF or IC2-PA plus LF. "IC2-PA" refers to the combination of PA-L1-I207R and PA-U2-R200A. These types of tumors were highly and equally sensitive to these engineered lethal toxins *in vivo* (Figures 7A, 7B, and 7C). LL3 cells were sensitive to the lethal toxins in *in vitro* cytotoxicity assay, whereas B16-BL6 cells were highly resistant (Figure 7D). These results suggested that targeting certain cell-types in tumor stromal compartments may play a role in tumor responses to the toxins.

[0171] Because both CMG2- and TEM8-null mice are able to support normal tumor growth, these mice provide powerful genetic tools to dissect the mechanisms by which the engineered anthrax toxins control tumor growth. To determine the role of stromal compartments in the potent anti-tumor activities of the engineered anthrax lethal toxins, A549 tumor-bearing *Cmg2*^{-/-} and *Tem8*^{-/-} mice and their littermate control mice were treated with PA-L1/LF after tumors had grown to about one gram. A549 cells contain wild-type BRAF and are resistant to PA-L1/LF in *in vitro* cytotoxicity assays (Figures 13A and 13B). While A549 tumors in *Cmg2*^{+/+} and *Cmg2*^{+/-} mice were very sensitive to the toxin, the tumors growing in *Cmg2*^{-/-} mice were much less sensitive (Figures 7E and 7F). In contrast, the A549

tumors growing in *Tem8^{-/-}* mice, as well as in their littermate control mice, all were sensitive to the toxin treatments (Figures 7G and 7H). These results clearly demonstrate that the anti-tumor activities of the engineered toxin involve targeting certain tumor stromal compartments. The results also reveal that CMG2 rather than TEM8 is the major toxin receptor mediating the anti-tumor activities of the toxin. In the presence of stromal CMG2 expression, the engineered toxin was highly potent, showing efficacy even for tumors that were very large in size ($\approx 5\%$ of total body weight) (Figures 7E, 7F, 7G, and 7H).

EXAMPLE 6A

[0172] This example demonstrates that targeting tumor endothelial cells is responsible for the anti-tumor activities of the engineered anthrax lethal toxins.

[0173] To determine which cell-type in tumor stromal compartments is responsible for the anti-tumor action of PA-L1/LF, B16-BL6 tumors were inoculated into three types of mice: *Cmg2^{-/-}* mice, *Cmg2^{-/-}* mice with a CMG2-transgene expressed only in endothelial cells (named *Cmg2^{EC}* hereafter, see Liu et al., *Nature*, 501: 63-68 (2013) for detailed description), and *Cmg2^{-/-}* mice with a CMG2-transgene expressed only in vascular smooth muscle cells (*Cmg2SM*) (see Liu et al., *Nature*, 501: 63-68 (2013) for detailed description). B16-BL6 cells are insensitive to PA-L1/LF in *in vitro* cytotoxicity assays (Figure 7D). As seen above for A549 tumors (Figure 7E), B16-BL6 melanomas in *Cmg2^{-/-}* mice were insensitive to PA-L1/LF, whereas the same tumors in *Cmg2^{+/+}* mice were exquisitely sensitive (Figures 8A and 8B). While the B16-BL6 tumors in *Cmg2SM* mice were, like in *Cmg2^{-/-}* mice, insensitive to the toxin, the tumors in *Cmg2^{EC}* mice were fully sensitive (Figures 8A and 8B). Thus, CMG2 endothelial expression is sufficient to mediate the anti-tumor activities of the toxin. To further evaluate the role of targeting tumor endothelial cells in cancer targeted therapy, B16-BL6 tumors were grown in endothelial cell-specific CMG2-null (termed *Cmg2(EC)^{-/-}* hereafter, see Liu et al., *Nature*, 501: 63-68 (2013) for detailed description) mice. The tumors in *Cmg2(EC)^{-/-}* mice completely lost sensitivity to PA-L1/LF as well as to PA-IC2/LF (Figures 8C and 8D), whereas the tumors in myeloid CMG2-specific CMG2-null (*Cmg2(Myel)^{-/-}*, see reference Liu et al., *Cell Host Microbe*, 8: 455-462 (2010) for detailed description) mice remained sensitive to PA-IC2/LF (Figure 8D). No antitumor activity was observed when PA-L1 was used alone, confirming that the antitumor activity of these toxins requires the action of LF, the enzymatic moiety of the toxins.

[00174] Taken together, the above results clearly demonstrate that the potent anti-tumor activities of the engineered anthrax toxins are due to the toxins' unique toxicities to the host-derived tumor endothelial cells rather than to other cell-types in the tumor stromal compartments, e.g., vascular smooth muscle cells and myeloid lineage cells. An additional observation was that the *Cmg2^{EC}* mice used in the tumor trials were themselves (when tumor-free) highly resistant to challenge with PA-L1/LF, in that 100% of the *Cmg2^{EC}* mice survived 5 doses of 50 µg PA-L1 plus 50 µg LF, a regimen that killed 50% of wild-type mice (Figure 8E). These results indicate that the engineered toxins are selectively toxic to proliferating tumor endothelial cells rather than to the mostly quiescent normal endothelial cells that do not express MMPs or uPA. To test this hypothesis, blood vessels of B16-BL6 tumor-bearing mice treated with PBS or PA-L1/LF were labeled with the fluorescent lipophilic carbocyanine dye DiI by cardiac perfusion (Li et al., *Nat. Protoc.*, 3: 1703-1708 (2008)). During perfusion, DiI directly incorporates into endothelial cell membranes upon contact, allowing visualization by fluorescence microscopy of vasculature structures within tumors and normal tissues. Notably, no differences were detected in vasculature structures of various normal tissues, including the spleens, kidneys, livers, and hearts of the B16-BL6 tumor-bearing mice treated with PBS and the toxin. Remarkably, while blood vessels were abundant in the tumors treated with PBS, vessels in the tumors treated with PA-L1/LF were rarely detected (Figure 8F). B16-BL6 melanomas and LLC carcinomas were also sectioned and histologically analyzed after the tumor-bearing mice were treated with PA-L1/LF or PBS. Extensive tumor necrosis (H&E staining) and decreases in cell proliferation (Ki67 staining) accompanied by loss of tumor vascular structures were readily detected in the toxin-treated B16-BL6 and LLC tumors. These results support the view that the engineered lethal toxins selectively damage tumor endothelial cells while sparing normal endothelial cells. CD31 and TUNEL co-staining was also performed on B16-BL6 tumors. Although extensive apoptotic tumor cell death was detected in PA-L1/LF-treated tumors, no apoptotic cell death was identified among the rarely detected tumor endothelial cells in the toxin-treated tumors, suggesting that the toxin may exert the antitumor effects through affecting endothelial cell proliferation rather than by inducing apoptosis.

EXAMPLE 6B

[0174] This example demonstrates that engineered anthrax lethal toxins inhibit the proliferation of tumor endothelial cells.

[0176] Tumor endothelial cells were isolated from B16-BL6 tumors through intermolecular adhesion molecule 2 (ICAM2) sorting to investigate the toxic effects of the engineered toxins on tumor endothelial cells. The purity of the isolated endothelial cells was confirmed by another endothelial marker, CD31. Delivery of LF into the cytosol of endothelial cells by PAL1 was evidenced by the cleavage of MEK1 and MEK2, accompanied by a dramatic decrease in phosphorylation of ERK1/2, as measured by Western blotting using MEK1, MEK2, ERK, and Phospho-ERK antibodies. Expression of the toxin-activating proteases by endothelial cells was also confirmed by the cells' susceptibilities to the protease-activated PA variants (PA, PA-L1 (MMP-activated PA variant), and PA-U2 (urokinase-activated PA variant)) in the presence of FP59. FP59 is a fusion protein of LF amino acids 1–254 and the catalytic domain of *Pseudomonas aeruginosa* exotoxin A that kills all cells by ADP ribosylation of eEF2 after delivery to cytosol by PA. To examine the cytotoxic effects of the toxin on tumor endothelial cells, the cells were treated with PA-L1/LF for 48 h and 72 h, respectively, followed by annexin V plus propidium iodide (PI) staining to identify apoptotic cells by flow cytometry. Although PA-L1 plus FP59 could induce dramatic apoptotic cell death 24 h after incubation, PA-L1 plus LF could not do so even after 72 h incubation. Although the engineered lethal toxin did not directly kill endothelial cells, the toxin displayed potent inhibitory effects on endothelial cell proliferation. Thus, Ki67 staining revealed that tumor endothelial cell proliferation nearly completely ceased after 72 h incubation with the toxin. The toxin's effects on endothelial cells could be fully replicated by trametinib (although much higher molar concentrations were required), a small molecule inhibitor of MEK1/2 approved by the Food and Drug Administration for treating patients having metastatic melanoma with BRAF^{V600E} mutation. These data suggest that the inhibitory effects of the engineered toxins were through disruption of the MEK-ERK pathway.

EXAMPLE 7

[0177] This example demonstrates the additional benefit of the engineered toxin in targeting tumors having the BRAF mutation.

[0178] Due to their unique action on tumor endothelial cells, the tumor-associated protease-activated anthrax lethal toxins exhibit potent anti-tumor activities even for the tumors composed of cancer cells that are insensitive to the toxins. However, a subset of human cancer cells have oncogenic BRAF mutations such as BRAF^{V600E} that make the tumor

cells dependent on the RAF-MEK-ERK pathway for survival while also making them exquisitely sensitive to anthrax lethal toxin (Abi-Habib et al., *Mol. Cancer Ther.*, 4: 1303-1310 (2005)). It was hypothesized that engineered anthrax lethal toxins would have additional benefit in treatment of solid tumors having the BRAF^{V600E} mutation. To test this, human colorectal carcinoma Colo205 cells, which contain the oncogenic BRAF^{V600E} mutation and are sensitive to PA-L1/LF *in vitro* (Figure 13A and 13B), were inoculated into littermate *Cmg2*^{-/-} and *Cmg2*^{+/+} mice and treated with PA-L1/LF. Significantly, Colo205 tumors in *Cmg2*^{-/-} mice were sensitive to the toxin treatment, although the response to the toxin treatment was lower than the strong response of the tumors growing in *Cmg2*^{+/+} mice (Figure 14). These results suggest that in the 'toxin-sensitive' tumors, the anti-tumor activity of the toxin depends on targeting both tumor endothelial cells as well as the cancer cells.

EXAMPLE 8

[0179] This example demonstrates the effects of anthrax lethal toxin on metabolism of tumor endothelial cells.

[0180] Tumor endothelial cells were isolated from B16-BL6 tumors to investigate the underlying mechanisms of the toxic effects of the engineered toxins on tumor endothelial cells. Notably, proliferation of the endothelial cells cultured in endothelial cell growth media was decreased significantly after the cells were incubated with PA-L1/LF for 72 h (Figure 9A). The delivery of LF into the cytosol of endothelial cells by PA-L1 was evidenced by the cleavage of MEK2 after a 3-h incubation with the toxin. Images were taken of the live, MTT-stained endothelial cells after incubation with or without 5 µg/mL PA-L1 plus 0.5 µg/mL LF for 72 h. The relative lower density of the cells treated with the toxin was noted. Because cellular metabolism is crucial for all cellular processes, it was examined whether lethal toxin affected the bioenergetics of tumor endothelial cells. After incubation with PA-L1/LF for 24 h, the extracellular acidification rates (ECARs) and oxygen consumption rates (OCRs) of tumor endothelial cells were measured under basal conditions, and following additions of the mitochondrial inhibitors oligomycin (ATP synthase inhibitor), FCCP (mitochondrial oxidative phosphorylation uncoupler), and antimycin A (Complex III inhibitor). ECAR reflects cytosolic glycolytic activity whereas OCR reflects mitochondrial oxidative phosphorylation. The engineered toxin significantly inhibited endothelial cell glycolytic activity under basal conditions, as well as when mitochondria were inhibited (Figure 9B). The up-regulation of glycolytic activity, which attempts to compensate for

diminished energy production during mitochondria inhibition (by oligomycin and FCCP), was also compromised by the toxin (Figure 9B). Furthermore, the basal OCRs, ATP production-coupled OCRs (OCR_{ATPS}), maximal respiration (MR), spare respiratory capacity (SRC), and cellular ATP levels of the endothelial cells were also remarkably decreased by the toxin (Figures 9C, 9D, and 9E). These results demonstrate that the toxin profoundly affects tumor endothelial cell metabolism through affecting glycolysis, as well as mitochondrial oxidative phosphorylation.

[0181] To explore the underlying mechanisms of the toxin's effects on metabolism of endothelial cells, the expression levels of key genes in central metabolism by real-time PCR analyses were surveyed. Surprisingly, many genes key to glucose uptake, glycolysis, tricarboxylic acid cycle, glutamine usage, as well as lipid synthesis were significantly down-regulated by PA-L1/LF (Figure 9F). Because glucose and glutamine are two major carbon sources for energetic metabolism and macromolecule syntheses, and lipid synthesis provides essential plasma membrane building blocks for cell proliferation, the profound effects of the toxin on tumor growth can be attributed at least in part to the transcriptional repression of key metabolic genes demonstrated here.

EXAMPLE 9

[0182] This example demonstrates that preventing antibody responses to the engineered toxin allows repeated courses of treatment.

[0183] The tumor-associated protease-activated anthrax toxins provide high tumor specificity and high anti-tumor efficacy. However, these bacterial proteins are foreign antigens to mammalian hosts and may induce neutralizing antibodies that prevent long-term use. Therefore, strategies for preventing an immune response may be useful.

[0184] To examine whether a combination of pentostatin and cyclophosphamide (PC) blocks production of antibodies that neutralize engineered anthrax toxins, a trial was performed using the highly metastatic LL3 (mouse) carcinomas established in syngeneic immunocompetent C57BL/6 mice. The tumor-bearing mice were treated with PBS, a PC regimen, IC2-PA/LF, or the combined therapy of the PC regimen and IC2-PA/LF, following the schedule shown in Figures 10A and 10B. For the combined treatment groups, the tumor-bearing mice were prepared with doses of PC 3 and 4 days prior to the first toxin treatment. The combined treatment groups were treated with a total of 4 cycles of toxin and PC, with intervals of 5-7 days between cycles. IC2-PA/LF alone showed strong anti-tumor effects

(Figures 10A and 10B). Surprisingly, all of the combined treatments showed much higher anti-tumor efficacy at both early and late times, with the tumors remaining responsive to the treatments even after the 4th cycle of the therapy (Figures 10A-10B). Importantly, no mortality was observed in the low (15 μ g IC2-PA + 5 μ g LF) and the medium (20 μ g IC2-PA + 6.7 μ g LF) dose groups. In fact, the mice receiving the combined treatments were alive after 42 days, well after mice in the other groups had to be euthanized due to their high tumor burdens (Figure 10A-10B). The PC regimen alone exhibited potent anti-tumor activities (Figure 10A-10B). As expected, neutralizing antibodies were detected in all of the mice treated with the toxin alone. Antibodies were detected as early as 10 days after the first treatment, the time at which the tumors began to show decreases in response to the toxin (only) treatment. Strikingly, no neutralizing antibodies were detected in the tumor-bearing mice of the combined therapy group even after the 4th round of therapy (Figures 10C, 10D, and 10E).

[0185] This study was next extended to include therapy of another highly malignant syngeneic tumor, the B16-BL6 melanomas implanted in immunocompetent C57BL/6 mice. This experiment used a modified toxin and PC regimen as shown in Figures 15A-15B. The B16-BL6 melanoma-bearing mice were treated with PBS, IC2-PA/LF (30 μ g /10 μ g), a PC regimen, or the combined therapy of the PC regimen and IC2-PA/LF twice in the first week and weekly in the following weeks (Figures 15A-15B). Again, the PC alone regimen had a significant anti-tumor effect (Figures 15A-15B), and the combined treatment showed remarkable efficacy, with the tumors remaining responsive to the treatments even after the 5th cycle of the therapy (Figures 15A-15B). Consistently, no neutralizing antibodies against the engineered toxin were detected in mice treated with the combined PC and the toxin even after the five cycles of therapy (Figures 15C-15D).

[0186] Taken together, the above results reveal that the combined toxin and PC therapy has remarkable and prolonged anti-tumor effects. It was demonstrated that the PC regimen contributes to the antitumor efficacy of the combined therapy not only by preventing the induction of neutralizing antibodies, thereby allowing multiple cycles of therapy, but also via its previously unrecognized anti-solid tumor activity. *In vitro* cytotoxicity assays showed that pentostatin and cyclophosphamide were not toxic to B16-BL6, LL3, and endothelial cells when used either alone or in combination (Figures 16A, 16B, 16C, and 16D), suggesting that the anti-solid tumor effects of PC discovered in this work might occur through effects on immune cell compartments that support tumor cell proliferation and survival.

EXAMPLE 10

[0187] This example demonstrates the effects of pentostatin and cyclophosphamide on immune cells.

[0188] To investigate the effects of the PC regimen on immune cells, splenocytes were isolated from naïve mice and B16-BL6 melanoma-bearing mice from various treatment groups after the second round of treatments as shown in Figures 15A-15B. Flow cytometry analyses revealed that B-cell populations ($CD45R^+$, IgM^+ , IgD^+ cells) were nearly completely depleted in the PC as well as in the combined therapy groups (Figures 11A). To a lesser extent, T-cell populations ($CD4^+$, $CD8^+$, and $CD27^+$ cells) were also reduced in these treatment groups (Figures 11B and 11C). The PC regimen and the combined PC and toxin treatments did not affect granulocyte populations ($CD11b^+$ and $Gr-1^+$ cells). It was found that the numbers of $CD11b^+$ and $Gr-1^+$ granulocytes were significantly increased in the tumor-bearing mice when compared to the numbers in naïve mice regardless of treatment type, suggesting the existence of innate immune responses to the tumors (Figure 11E). The IC2-PA/LF alone did not significantly affect these cell populations (Figures 11A, 11B, 11C, 11D, and 11E). Therefore, the above results clearly demonstrate that the PC regimen efficiently depletes lymphocytes, in particular B-cells, while sparing innate immunity. In agreement with PC's effects on lymphocytes, the total splenocytes of the mice treated with PC regimen alone or in combination with the toxin were also significantly decreased (Figure 11E). Without being bound to a particular theory or mechanism, it is believed that the absence of a humoral immune response to the engineered toxins was due to the B-cell depletion caused by the PC regimen.

EXAMPLES 11-18

[0189] The following materials and methods were employed in the experiments of Examples 11-18.

Reagents

[0190] Enzymes for DNA manipulations and modifications were purchased from New England Biolabs (Beverly, MA). Dynabeads TALON was obtained from Invitrogen (Carlsbad, CA). PA and FP59 were prepared as described in Pomerantsev et al., *Protein Expr. Purif.*, 80: 80-90 (2011); Liu et al., *Cell. Microbiol.*, 9: 977-987 (2007); and Gupta et

al., *PLoS One*, 3: e3130 (2008). The extracellular domain (ECD) of CMG2 (amino acids 40–218) was produced in *Escherichia coli* with an N-terminal His6 tag (Santelli et al., *Nature*, 430: 905-908 (2004); Chen et al., *J. Biol. Chem.*, 282: 9834-9846 (2007)). The ECD of TEM8 (amino acids 35–227) with a C-terminal His6 sequence was expressed in CHO cells. The latter protein was concentrated and purified from the culture medium with a nickel-nitrilotriacetic acid column (Qiagen, Valencia, CA) and further purified by MONOQ column chromatography (GE Healthcare, Pittsburgh, PA). Minimum essential medium alpha, HEPES buffer, hygromycin B, Hanks' balanced salt solution (HBSS), and fetal bovine serum (FBS) were from Invitrogen. The mutagenic nucleoside triphosphates 8-oxo-2'-deoxyguanosine triphosphate (8-Oxo-dGTP) and 6-(2-deoxy- β -D-ribofuranosyl)-3,4-dihydro-8H-pyrimido-[4,5-c][1,2]oxazin-7-one triphosphate (dPTP) were purchased from Trilink Biotechnologies (San Diego, CA). Other chemical reagents were obtained from Sigma (St. Louis, MO).

Construction of a Randomly Mutated PAD4 Phage Display Library

[0191] Random mutagenesis of PAD4 was accomplished by PCR amplification with 10 ng of pYS5 template DNA and nucleotide analogues 8-oxo-dGTP and dPTP (Chen et al., *J. Biol. Chem.*, 282: 9834-9846 (2007); Zaccolo et al., *J. Mol. Biol.*, 255: 589-603 (1996)). The PCR reaction (a total volume of 25 μ l) contained 2.5 units of Takara Ex TaqDNA polymerase (Clontech, Mountain View, CA), 1 μ M each of forward primer 5'GCTTGAATTCATTTCATTATGATAGAAATAAC3' (SEQ ID NO: 34) and reverse primer 5'AATTCAAGCTTTCCTATCTCATAGCCTTTTTT 3' (SEQ ID NO: 35) (with EcoRI and HindIII sites underlined and bolded, respectively), 2 mM MgCl₂, 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 400 μ M each of dATP, dCTP, dGTP, dTTP, 50 μ M of 8-oxo-dGTP and 10 μ M of dPTP. After forty-two cycles (94 °C for 1 min, 60 °C for 1.5 min, and 72 °C for 5 min), 1 μ l of the amplified DNA was used in a secondary PCR in which the above conditions were used except that 8-oxo-dGTP and dPTP were omitted. The secondary PCR product was digested with EcoRI and HindIII, cloned into the T7Select10-3 phage display system, and packaged into bacteriophage according to the manufacturer's instructions (Novagen, Madison, WI).

Selection of CMG2- and TEM8-Specific PAD4 Variants using magnetic beads

[0192] Bacteriophage libraries were selected for CMG2-specific clones by four rounds of panning with Dynabeads TALON. Each round of panning was begun by washing 1 mg Dynabeads TALON with TALON binding and washing buffer (50 mM Na Phosphate, pH 8.0, 300 mM NaCl, 5 mM MgCl₂, 0.01% Tween-20, and 100 µg/ml of bovine serum albumin), and then adding 10 µg of extracellular domain of CMG2. After incubation on a roller at 4 °C for 30 min, the beads were washed four times with the same buffer. Bacteriophage (1 x 10⁹ particles) were then added to the mixture and incubated at 4 °C for 90 min. The beads were washed five times with the same buffer over total time periods of 10 min, 2 h, 5 h and 20 h in the four successive panning cycles. Following the final wash in each cycle, bacteriophage were eluted from the beads with 100 µl of TALON elution buffer (150 mM imidazole, 50 mM Na Phosphate, pH 8.0, 300 mM NaCl, 5 mM MgCl₂, 0.01% Tween-20, and 100 µg/ml of bovine serum albumin) by incubating the mixture on a roller for 10 min at 4° C. Phage were amplified by growth in *E. coli* BLT5403 and used in the next cycle of panning. To select TEM8-specific PAD4 variants, the experimental procedures described above were conducted except that 10 µg of extracellular domain of TEM8 was used for binding to the Dynabeads.

Sequencing of DNA Inserts in T7 Bacteriophage

[0193] Bacteriophage from the fourth round of panning were picked from individual bacteriophage plaques, suspended in M9LB broth (Novagen, Madison, WI) and set at room temperature for 2 h to allow bacteriophage particles to diffuse from the agar. EDTA was then added to the bacteriophage suspension to give a final concentration of 10 mM, and the suspension was heated at 65 °C for 10 min. One microliter of the sample was used for PCR amplification in reactions containing 2.5 units of Takara Ex TaqDNA polymerase, forward primer 5'TAAGTACGCAATGGGCCACG3' (SEQ ID NO: 36), and reverse primer 5'AACTCAGCGGCAGTCTCAAC3' (SEQ ID NO: 37), with PCR conditions as described above, except the annealing temperature was 50 °C. The PCR product was sequenced by dideoxy-mediated sequencing reactions.

Construction of PA I656 and PA E654 Mutants

[0194] Cloning of mutated PA domain 4 sequences was done with plasmid pYS54, an expression vector having the silent restrictions sites *Stu*I and *Xho*I at positions corresponding to amino acids 650-652 and 665-667, respectively (Chen et al., *J. Biol. Chem.*, 282: 9834-9846 (2007)). To obtain all 19 amino acid substitutions of either I656 or E654, oligonucleotides were synthesized having the sense strand sequence of SEQ ID NO: 38 and antisense strand sequence of SEQ ID NO: 39, where either (but not both) the E654 codon gaa or the I656 codon ata in the sense strand and the corresponding codons in the antisense strands were fully randomized. Additionally, mutants having R659S/M662R substitutions plus all 19 amino acid substitutions of E654 were made in a similar way by including the 659S and 662R codons in the oligonucleotides described above. In each case, double strand cassettes were obtained by heating the mixture of sense and antisense strands at 100 °C for 10 min and allowing the mixture to cool slowly to room temperature. The resulting products were digested with *Stu*I and *Xho*I and ligated into pYS54 digested with the same restriction endonucleases, and the ligation mixtures were transformed into *E. coli* XL-1 Blue. Plasmid DNAs isolated from individual clones were sequenced to select all 20 amino acids substitutions.

Construction of PA Proteins with PAD4 substitutions and having a Modified Furin Site

[0195] To prepare PA proteins having substitutions in residues identified as being involved in binding to CMG2 and TEM8 (I656, etc.) and also an uncleavable furin site, the pYS54 constructs described above were digested with *Pst*I and *Hind*III, and the intervening region was replaced with the corresponding *Pst*I to *Hind*III fragment isolated from a plasmid encoding PA-U7 (Liu et al., *J. Biol. Chem.*, 276: 17976-17984 (2001)), in which the ¹⁶⁴RKKR¹⁶⁷ sequence is replaced by PGG. The doubly-substituted PA proteins derived from PA I656V and PA I656Q were accordingly named PA-U7 I656V and PA-U7 I656Q, respectively. In a similar way, altered furin site sequences were inserted into constructs having multiple substitutions, including the PA E654T/R659S/M662R protein (abbreviated PA TSR). In this case, alternative furin site sequences used included those from PA-U7 (as above) and PA-L1, the latter being one cleaved by matrix metalloproteases (Liu et al., *Cancer Res.*, 60: 6061-6067 (2000)). The doubly-mutated PA proteins derived from PA TSR were accordingly named as PA-U7 TSR and PA-L1 TSR.

Expression and Purification of PA Proteins

[0196] Plasmids encoding individual constructs described above were transformed into the nonvirulent *B. anthracis* strains BH450 or BH460 and transformants were grown in FA medium with 10 µg/ml kanamycin for 15 h at 37 °C (Pomerantsev et al., *Protein Expr. Purif.*, 80: 80-90 (2011)). PA proteins were concentrated from culture supernatants and purified by chromatography on a MONOQ column (GE Healthcare) by methods described previously (Pomerantsev et al., *Protein Expr. Purif.*, 80: 80-90 (2011); Park et al., *Protein Expr. Purif.*, 18, 293-302 (2000)).

Measurement of PA Affinity for Receptors

[0197] Schild plot analyses were used as described previously (Liu et al., *Proc. Natl. Acad. Sci. U. S. A.*, 106: 12424-12429 (2009); Ittelson et al., *Nature*, 242: 330-332 (1973)) to determine the affinity (apparent K_d) of PA variant proteins for cellular receptors. Briefly, the nontoxic PA-U7, PA-U7 I656Q, PA-U7 I656V, and PA-U7 TSR proteins were used as competitors to block the toxicity of wild-type PA to CHO cells. Multiple PA dose-response cytotoxicity assays were done, each in the presence of a different fixed concentration of the competitor. The midpoints on the dose-response curves (T_i) were plotted against the competitor concentration. T_i is the midpoint of the dose-response curve obtained at a particular fixed competitor concentration. T_o is the value of T_i with no competitor added. The intercept of the resulting line at the point where $\log((T_i/T_o) - 1) = 0$ identifies the competitor concentration equal to the apparent K_d value, a measure of the affinity of the competitor for the receptor used by the toxin to produce toxicity.

Cells and Culture Media

[0198] Parental wild type Chinese hamster ovary (CHO) cells (CHO WTP4) and the PA receptor-expressing CHO CMG2-C4 and CHO TEM8-T4 cells, which overexpress CMG2 and TEM8 respectively, are as described previously (Liu et al., *Cell. Microbiol.* 9: 977-987 (2007); Liu et al., *J. Biol. Chem.* 278: 5227-5234 (2003)). CHO cells were grown in Minimum essential medium alpha with 5% FBS, 2 mM glutamine, 5 mM HEPES, pH 7.4, and 50 µg/ml gentamicin, with or without 300 µg/ml hygromycin B. HeLa and SN12C cells were cultured in Dulbecco's modified Eagle's medium with 10% FBS, 2 mM glutamine, and 50 µg/ml gentamicin. Mouse embryonic fibroblast (MEF) cells were isolated from E13.5

embryos of the wild type, CMG2^{-/-}, TEM8^{-/-} and CMG2^{-/-}/TEM8^{-/-} mice (to be described below) as described previously (Liu et al., *Proc. Natl. Acad. Sci. U. S. A.*, 106: 12424-12429 (2009)) and cultured in Dulbecco's modified Eagle's medium with 10% FBS, 2 mM glutamine, and 50 µg/ml gentamicin. MEF cells were passaged at 85–90% confluence by splitting 1:4 and used within the first six passages.

Cytotoxicity Assays of Cultured Cells

[0199] Cells were plated at 10,000 cells/well in 96-well plates and cultured for 24 h before treatment. PA proteins, combined with 100 ng/ml of FP59 for CHO cells and with 400 ng/ml of FP59 for MEF cells, were added to cells to a final volume of 200 µl/well. Cell viability was assayed 48 h after treatment by replacing the medium with 50 µl of medium containing 2.5 mg/ml MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide). Medium was removed after 1 h of incubation at 37 °C and the blue pigment produced by viable cells was dissolved in 50 µl/well of 0.5% (w/v) SDS, 25 mM HCl, in 90% (v/v) isopropyl alcohol. Plates were vortexed and oxidized MTT was measured as A_{570} using a microplate reader (Spectra Max 190, Molecular Devices, Sunnyvale, CA). Results were analyzed with PRISM software (GraphPad Software Inc., San Diego, CA) as percentage viability of control wells containing FP59 without PA. EC₅₀ values were determined by nonlinear regression sigmoidal dose-response analysis with variable slopes. Each assay was performed three times, and data from a representative assay are shown.

Lethality of PA Variants for Receptor-deficient Mice

[0200] The CMG2^{-/-}, TEM8^{-/-}, and CMG2^{-/-}/TEM8^{-/-} (double null) C57BL/6 mice used here were described previously (Liu et al., *Proc. Natl. Acad. Sci. U. S. A.*, 106: 12424-12429 (2009); Liu et al., *Nature*, 501: 63-68 (2013)). These mice and littermate control mice (all 8–10 weeks old) were challenged intraperitoneally with two doses of PA or PA variants along with FP59 (in 0.5 ml PBS, doses indicated in tables and figures) with a 2-day interval. Mice were observed for signs of malaise twice daily for one week following injection. All mice studies were carried out in accordance with protocols approved by the National Institute of Allergy and Infectious Diseases Animal Care and Use Committee.

Tumoricidal Activity of PA variants

[0201] Athymic nude mice were injected intradermally with 1×10^7 HeLa cells/mouse, and tumor xenografts were allowed to grow to 0.05% of body mass. Tumor-bearing mice were randomized into groups and injected intraperitoneally at intervals of 2 or 3 days, as indicated in the figures, with six doses of PBS or the engineered toxins. Body weights and tumor masses were measured at 2 or 3 day intervals. Tumors were measured with digital calipers (FV Fowler Company, Inc., Newton, MA) and tumor weights were estimated with the length, width, and height tumor dimensions using formulas: tumor weight (mg) = $\frac{1}{2}(\text{length in mm} \times \text{width in mm}^2)$ or $\frac{1}{2}(\text{length in mm} \times \text{width in mm} \times \text{height in mm})$.

EXAMPLE 11

[0202] This example demonstrates the selection of PAD4 mutants that bind to CMG2.

[0203] An improved panning procedure was developed using soluble forms of the extracellular domains of CMG2 and TEM8 bound to magnetic beads (Dynabeads TALON). The specific binding of the phage-displayed PAD4 to each type of beads was verified by showing that preincubation of the beads with 100 ng PA reduced phage binding by 90% (Table 1).

TABLE 1

Treatments	Phage bound to Dynabeads
Exp #1 – CMG2 magnetic beads	
Control ^a	7.5×10^2
CMG2-mag beads	6.6×10^4
CMG2-mag beads + PA ^b	5.6×10^3
Exp #2 – TEM8 magnetic beads	
Control ^a	7.4×10^2
TEM8-mag beads	7.0×10^4
TEM8-mag beads + PA ^b	5.6×10^3

^abinding of bacteriophage to Dynabeads without addition of an extracellular domain.

^b100 ng of PA was mixed with bacteriophage lysate before adding to the Dynabeads.

[0204] *Table 1:* After binding of bacteriophage, the mixture was washed five times with binding and washing buffer for a total of 10 min and bacteriophage were eluted with elution buffer and titered.

[0205] To enrich for PAD4 variants having high affinity, the stringency of washing in each successive amplification round was increased, using washing times of 10 min, 2 h, 5 h and 20 h. The number of phage increased 600-fold by the third round of panning. The fourth round of panning on CMG2 resulted in a 65% decrease in phage numbers (Table 2), possibly because the highest possible affinity had already been reached by the third round, and the prolonged 20-h washing removed some of these tightly binding phage. No similar effect was seen when panning on TEM8 beads. After the fourth round of panning, individual bacteriophage plaques were chosen and sequenced.

TABLE 2

Round of panning	Phage bound to Dynabeads
Expt. #1 – CMG2-magnetic beads	
1 st	3.8×10^4
2 nd	3.8×10^5
3 rd	2.3×10^7
4 th	8.1×10^6
Expt. #2 – TEM8-magnetic beads	
1 st	7.0×10^4
2 nd	1.2×10^7
3 rd	1.1×10^7
4 th	4.5×10^7

Table 2: Experimental procedures were the same as described in Table 1, except the magnetic beads were washed for 10 min, 2 h, 5 h, and 20 h in 1st, 2nd, 3rd and 4th rounds of panning, respectively.

[0206] A total of 104 phage clones were sequenced over the 420-base pair (bp) region corresponding to PA amino acids 596-735. These DNA sequences were aligned. The

resulting amino acid sequences of the clones were determined. Among the resulting total of 43,680 bp examined, a total of 616 changes were found, for a mutation rate of 1.4%. However, this value and the subsequent analyses were for the phage library after selection for binding to CMG2. This selection will both eliminate the large fraction of sequences coding for misfolded or unstable proteins as well as those with specific surface residue changes that decrease receptor recognition. The selection is also expected to enrich for silent mutations. It was found that the number of substitutions in codon positions 1-3 were 207, 165, and 244, respectively. Thus, substitutions in the 3rd position, which are less likely to change the amino acid, were enriched, consistent with the expected elimination of clones having the most common mutagenic events - amino acid changes that decrease structure and/or function.

[0207] While mutation frequencies and other numerical values were calculated, these can be viewed only as estimates. Comparison of the DNA and amino acid sequences shows that many clones were related. The bp changes seen in multiple clones must have occurred in an early PCR cycle and then expanded into many progeny sequences. This illustrates that the clones are not independent, as would be needed to achieve an accurate statistical analysis.

[0208] Because 8-oxo-GTP causes A to C and T to G transversions and dPTP causes A to G and T to C transitions (Zaccolo et al., *J.Mol.Biol.*, 255: 589-603 (1996)), a broad spectrum of amino acid substitutions should be generated, although codons containing G and C will be less affected. Although there were multiple mutations in nearly every clone from the CMG2-selected library, strikingly, it was found that mutation of Ile656 to Val, in the region between 486 and 487 (amino acids 656-665), occurred in 85 of the 86 clones sequenced, indicating that residue Ile656 has a role in the binding of PA to the CMG2 receptor. In contrast, the TEM8-selected library showed a greater diversity of substitutions. However, residue Glu654, in the same region as Ile656, was very frequently mutated. Thus, of the 95 random clones sequenced, 27 have Glu654 substituted by either Gly or Ala, indicating that this residue may have a role in the binding of PA to TEM8.

EXAMPLE 12

[0209] This example demonstrates PAD4 variants having substitutions at Ile656.

[0210] To further characterize the role of Ile656 in binding of PA to TEM8, Ile656 was mutated to all other 19 amino acids. Substitutions of Ile656 with Leu, Met, Gln, Ser, Thr, Val, and Tyr were well tolerated, because these proteins retained cytotoxicity to the wild-type CHO WTP4 cells (Table 3). While these Ile656 mutants also retained activity to CHO

CMG2-C4 cells, the CHO cell line that over-expresses only CMG2, all these PA variants displayed greatly decreased cytotoxicity to CHO TEM8-T4 cells, which over-express only TEM8. In particular, PA I656Q and I656S showed no cytotoxicity to CHO TEM8-T4 cells at the highest concentrations tested, 80 ng/ml. Therefore, PA I656Q and I656S are highly selective for CMG2. All other variants showed decreased cytotoxicity to all three types of CHO cells. PA I656W showed no cytotoxicity to the cell lines even at the highest concentration (80 ng/ml) tested (Table 3). Nearly all the I656 substitution variants showed a much greater loss of cytotoxicity toward the CHO TEM8-T4 cells than toward CHO CMG2-C4 and CHO WTP4 cells.

TABLE 3

PA variant	EC ₅₀ (ng/ml) on CHO cells [#]			Selectivity for CMG2 vs. TEM8 [§]
	WTP4	TEM8-T4	CMG2-C4	
PA (native)	0.14*	0.21	0.05	4
PA I656Q	0.48*	>83.3	0.05	>1667
PA I656S	0.42	>83.3	0.07	>1190
PA I656N	1.10	>83.3	0.12	>694
PA I656H	4.11	>83.3	0.17	>490
PA I656E	1.71	>83.3	0.17	>490
PA I656A	3.87	>83.3	0.26	>320
PA I656G	1.83	>83.3	0.29	>287
PA I656K	9.27	>83.3	0.31	>269
PA I656C	0.63*	17.33	0.10	173
PA I656M	0.24	4.70	0.05	99
PA I656Y	2.13	13.61	0.14	99
PA I656L	0.22	3.71	0.04	92
PA I656F	1.76	7.29	0.12	61
PA I656T	0.28	1.59	0.03	53
PA I656D	15.33	>83.3	1.91	>44
PA I656P	2.76	5.25	0.22	24
PA I656V	0.35	1.34	0.06	22
PA I656R	>83.3	>83.3	5.33	>16
PA I656W	>83.3	>83.3	>83.3	nc [#]

[§]Ratio: (EC₅₀ for TEM8-T4)/(EC₅₀ for CMG2-C4). Variants are sorted according to this value.

[#] nc is for not able to be calculated.

[#]In EC₅₀ values, standard deviations (SD) are indicated as follows:

No asterisk indicates that SD = <0.5 x the absolute value or cannot be calculated and one asterisk indicates that SD = 0.5-1.0 x the absolute value.

[0211] *Table 3:* PA I656 mutants containing all 20 amino acids at position 656 were tested for toxicity to CHO cells. Values in the table are EC₅₀, the concentrations of the PA proteins that killed 50% of cells when combined with 100 ng/ml of FP59. The analyses were performed three times, and the averages are listed.

EXAMPLE 13

[0212] This example demonstrates PAD4 variants having substitutions at Glu654.

[0213] To further characterize the role of position 654 in binding to CMG2, E654 was mutated to all other 19 amino acids. In many cases, the substitutions led to a greater reduction in potency toward CHO-CMG2-C4 than toward CHO-TEM8-T4 cells (Table 4), although the losses in potency were smaller than those caused by substitutions at I656 (Table 3). Only the substitution of Val was fully tolerated, as shown by the retention of toxicity toward both cell types. In contrast, substitution of Ala, Cys, Gly, His, Asn, Gln, Ser, Thr, Trp and Tyr caused decreases in potency toward CHO CMG2-C4 cells that were not paralleled by decreases in potency toward CHO TEM8-T4. In fact, many of these substitutions appeared to cause a slight increase in potency toward the CHO-TEM8-T4 cells, as seen particularly with the E654Q, E654V and E654W variants. These variants have a 2- to 3-fold higher toxicity toward CHO-TEM8-T4 than does native PA, indicating that the E654 residue plays a role in determining the relative affinities for the two receptors.

TABLE 4

PA variant	EC ₅₀ (ng/ml) on CHO cells [#]		Selectivity for TEM8 vs. CMG2 ^s
	TEM8-T4	CMG2-C4	
PA (native)	0.52	0.39*	0.7
PA E654C	0.16	0.81*	4.9
PA E654N	0.16	0.68	4.3
PA E654Q	0.13	0.39	2.9
PA E654S	0.10	0.26*	2.6
PA E654H	0.22	0.53	2.4
PA E654F	0.63	1.43	2.3
PA E654Y	0.16	0.36	2.3
PA E654W	0.15*	0.34	2.2
PA E654T	0.42	0.89	2.1
PA E654I	0.73	1.27	1.7
PA E654M	0.35	0.58	1.7
PA E654A	0.54	0.88	1.6

PA variant	EC ₅₀ (ng/ml) on CHO cells [#]		Selectivity for TEM8 vs. CMG2 ^{\$}
	TEM8-T4	CMG2-C4	
PA E654L	0.52	0.70	1.3
PA E654G	0.58	0.59	1.0
PA E654V	0.19*	0.20	1.0
PA E654R	2.40	1.12	0.5
PA E654P	7.20	2.80	0.4
PA E654D	2.40	0.83	0.3
PA E654K	2.50	0.50	0.2

^{\$}Ratio: (EC₅₀ for CMG2-C4)/(EC₅₀ for TEM8-T4). Variants are sorted according to this value.

[#]In EC₅₀ values, standard deviations (SD) are indicated as follows:

No asterisk indicates that SD = <0.5 x the absolute value or cannot be calculated, and one asterisk indicates that SD = 0.5-1.0 x the absolute value.

[0214] *Table 4:* PA E654 variants containing all 20 amino acids at position 654 were tested for toxicity to CHO cells. Values in the table are EC₅₀, the concentrations of the PA proteins that killed 50% of cells when combined with 100 ng/ml of FP59. The analyses were performed three times, and the averages are listed.

EXAMPLE 14

[0215] This example demonstrates PAD4 variants having substitutions in multiple residues.

[0216] To examine the effects of combining substitutions, PA variants were selected combining all 19 amino acids replacing E654 with the R659S/M662R substitutions. All 19 substitution mutants showed decreased toxicity toward CHO-CMG2-C4 cells (Table 5). Particularly, the PA variants KSR, PSR and RSR were nontoxic to CHO-CMG2-C4 cells when added at 80 ng/ml, the highest concentration tested. In contrast, variants PA ASR, GSR, HSR, MSR, NSR, QSR, SSR, TSR and WSR had potencies equal or slightly greater than that of native PA toward CHO-TEM8-T4 cells (Table 5). For example, PA TSR was 2-fold-more toxic to CHO-TEM8-T4 cells than native PA but 23-fold less toxic to CHO-CMG2-C4 cells. The PA QSR, MSR and GSR variants were similar to PA TSR, being only slightly less selective. Based on these data, PA MSR, QSR and TSR were selected for further study.

TABLE 5

PA variant		EC ₅₀ (ng/ml) on CHO cells*		Selectivity for TEM8 vs. CMG2 [§]
designation	Substitution (in addition to R659S/ M662R)	TEM8-T4	CMG2-C4	
PA (native)		0.40	0.16	0.4
MSR	E654M	0.60	6.83	11.4
TSR	E654T	0.35	3.00	8.5
QSR	E654Q	0.44	3.20	7.3
GSR	E654G	0.49	3.47	7.1
YSR	E654Y	0.63	4.33	6.8
VSR	E654V	0.71	3.93	5.5
RSR	E654R	15.67	>83	>5.3
LSR	E654L	3.53	17.3	4.9
FSR	E654F	0.62	2.97	4.8
ISR	E654I	2.00	9.33	4.7
NSR	E654N	0.43	1.73	4.1
CSR	E654C	6.40	17.0	2.7
SSR	E654S	0.35	0.91*	2.6
KSR	E654K	40.2	>83	>2
HSR	E654H	0.21	0.38	1.8
ASR	E654A	0.36	0.60	1.7
WSR	E654W	0.47	0.63	1.3
DSR	E654D	8.00	5.00	0.6
PSR	E654P	>83	>83	nc [#]

[§]Ratio: (EC₅₀ for CMG2-C4)/(EC₅₀ for TEM8-T4).

[#]nc is for not able to be calculated.

^{*}In EC₅₀ values, standard deviations (SD) are indicated as follows:

No asterisk indicates that SD = <0.5 x the absolute value or cannot be calculated, and one asterisk indicates that SD = 0.5-1.0 x the absolute value.

[0217] Table 5: PA variants having R659S and M662R substitutions along with substitution of each of the other 19 amino acids at position 654 were tested for toxicity to CHO cells. The 3-letter designation identifies the non-native substitutions at positions 654, 659, and 662. Values in the table are EC₅₀, the concentration of the PA protein that produces 50% killing when combined with 100 ng/ml of FP59. The analyses were performed three times, and the averages are listed.

EXAMPLE 15

[0218] This example demonstrates the cytotoxicity of PA variants to receptor-deficient mouse embryonic fibroblasts (MEFs).

[0219] While the CHO cell lines expressing a single receptor were essential in determining the preference of PA variants for CMG2 and TEM8, it was sought to verify the behavior of the PA proteins on other cell types. Mice having each receptor deleted, and cells isolated from them, provide a well-defined genetic system for examining the receptor specificity of the PA variants (Liu et al. *Cell Host Microbe*, 8: 455-462 (2010); Liu et al., *Nature*, 501: 63-68 (2013); Liu et al., *Toxins (Basel)*, 5: 1-8 (2012)). Cytotoxicity tests of the PA variants having substitutions at I656 (from Table 3) on MEFs isolated from these mice clearly demonstrated the high specificity of the PA I656 variants for CMG2 (Expt. 1, Table 6). The wild-type (WT) MEF cells appear to express both receptors, whereas the CMG2^{-/-} MEFs are highly resistant to PA I656Q, while remaining sensitive to native PA and other variants which can use the remaining TEM8 receptor. The MEF cells from mice deleted for both receptors were highly resistant to all the PA variants, consistent with prior evidence that there are no physiologically significant receptors other than CMG2 and TEM8 (Liu et al., *Nature*, 501: 63-68 (2013); Liu et al., *Toxins (Basel)*, 5: 1-8 (2012)).

[0220] In a parallel experiment (Table 6, Expt. #2), the MEF was used again, and it was observed that the PA triple substitution variants such as PA TSR exert their toxicity through the TEM8 receptor, since absence of the CMG2 receptor has little effect on their potency, while absence of the TEM8 receptor decreases potency >100-fold. Taken together, these data using MEF cells confirm the selectivity of the PA variants that was seen in Tables 3 and 5.

TABLE 6

PA variant	EC ₅₀ (ng/ml) on MEFs			
	WT	CMG2 ^{-/-}	TEM8 ^{-/-}	CMG2 ^{-/-} / TEM8 ^{-/-}
Expt. #1				
PA (native)	0.8	4.8	0.9	>1000
PA I656Q	0.9	>1000	0.9	>1000
PA I656V	1.0	7.0	1.1	>1000
PA I656W	>1000	>1000	>1000	>1000
Expt. #2				
PA (native)	0.9	17	1.0	
PA MSR	9.5	19	>1,000	
PA QSR	9.0	16	700	
PA TSR	3.5	7	130	

[0221] Table 6: PA variants were tested for cytotoxicity to the indicated MEF cells. Values in the table are EC₅₀, the concentrations of the PA proteins that killed 50% of cells

when combined with 400 ng/ml of FP59. The analyses were performed two additional times with results similar to those presented.

EXAMPLE 16

[0222] This example demonstrates the affinity of PA variants for receptors as determined by Schild plot analyses.

[0223] It was expected that the differences in toxicity of the PA variants as described above result principally from differences in receptor-binding affinities. To examine this hypothesis, apparent affinities were measured by competitive Schild plot analyses. These analyses require that the protein ligand be a nontoxic variant of the protein of interest. Nontoxic variants of the PA I656Q, PA I656V, and PA TSR proteins were constructed by replacing their furin cleavage sites, ¹⁶⁴RKKR¹⁶⁷ (SEQ ID NO: 4), with the uncleavable sequence PGG of the previously characterized PA-U7 protein (Liu et al., *J. Biol. Chem.*, 276: 17976-17984 (2001)). The resulting PA-U7 I656Q, PA-U7 I656V, and PA-U7 TSR were used as nontoxic receptor binding competitors. Reciprocal plots of the midpoints of the dose-response curves performed in varying amounts of the competitors yielded apparent dissociation constants (K_d) for the apparent affinity of the competitors to the PA receptors. These analyses showed that PA I656Q retained affinity for CMG2 receptor but greatly lost affinity for the TEM8 receptor (Table 7, Expt. #1). Although PA I656V retained affinity for CMG2, it had moderately decreased affinity for TEM8. Conversely, PA-U7 TSR showed a decrease in affinity to CMG2 (Table 7, Expt. #2), consistent with its preference for TEM8.

TABLE 7

Competitor	K _d (nM) on cell lines	
	CHO CMG2-C4	CHO TEM8-T4
Expt. #1		
PA-U7	8	25
PA-U7 I656V	8	46
PA-U7 I656Q	8	>240
Expt. #2		
PA-U7	12	35
PA-U7 TSR	48	30

[0224] Table 7: Analyses were performed as described in the Methods section. PA-U7, PA-U7 I656V, and PA-U7 I656Q, and PA-U7 TSR were used as competitors for PA in

cytotoxicity assays with the two cell lines listed. Schild plot analyses determined the apparent affinities of the competitors for the receptors on the two cell lines.

EXAMPLE 17

[0225] This example demonstrates the toxicity of PA variants to CMG2 and TEM8 knockout mice.

[0226] To explore whether the differing receptor specificities of the PA variants described above applied *in vivo*, CMG2^{-/-} and/or TEM8^{-/-} mice and their littermate control mice were challenged using 20 µg PA variants plus 20 µg FP59 (Figs. 17A-17D). All the WT and TEM8^{-/-} mice were killed by native PA + FP59 within 24 h (Fig. 17A), whereas most of the CMG2^{-/-} mice survived to 48 h but succumbed to the second dose given at 48 h within the following 24 h, consistent with CMG2 being the major toxin receptor mediating mouse lethality. All the CMG2^{-/-} mice survived two doses of the CMG2-selective PA I656Q + FP59 (Fig. 17B), while all their littermate control mice and TEM8^{-/-} mice died, mostly within 24 h after the toxin challenge, demonstrating that the high specificity of PA I656Q for CMG2 extends to the case of intact mice. The moderate increase in specificity of PA I656V to CMG2 was also confirmed, as 50% the CMG2^{-/-} mice survived to 75 h after receiving two doses of PA I656V + FP59 (Fig. 17C), while all CMG2^{-/-} mice had succumbed by this time after challenge with the native PA + FP59 (Fig. 17A and 17C, compared).

[0227] Comparison of the potency of the putatively TEM8-specific PA TSR variant toward wild type and CMG2-null mice produced the results shown in Fig. 17D. The PA TSR variant was much less toxic toward wild type mice than native PA, consistent with the CMG2 receptor being the main determinant of toxin sensitivity. However, PA TSR was more potent toward the CMG2-null mice than toward wild type mice. In the CMG2-null mice, death must result from targeting the TEM8 receptor in certain (unidentified) tissues, and PA TSR appeared to do this more efficiently than native PA.

EXAMPLE 18

[0228] This example demonstrates the tumoricidal activity of PA variants to human HeLa xenografts in mice.

[0229] To examine the effects of altering PA's receptor specificity on its tumor-targeting ability, two of the PA variants were tested against HeLa cell xenografts in mice, in comparison to the previously described PA-L1 variant. The PA variants used here contained

the MMP-specific L1 sequence at the furin site so as to achieve tumor specificity, and the effector used was LF rather than FP59. Mice bearing solid intradermal tumor nodules constituting $\approx 0.05\%$ of the total body mass were challenged by intraperitoneal injection of six doses of the toxins at 2- or 3-day intervals. At the doses used, mouse body weights did not differ between the groups (Fig. 18A). In addition, the mice showed no outward sign of illness or gross abnormalities. However, large differences were seen in the effect of the PA variants on tumor size. Both PA-L1 and PA-L1 I656Q had strong and similar anti-tumor activities, reducing tumor size by $>80\%$ compared to PBS-treated tumors (Fig. 18B). In contrast, the tumors showed less response to treatment with PA-L1 TSR, with only a 50% reduction of tumor size occurring (Fig. 18B). These data are consistent with other evidence that targeting of tumors is most effective when PA acts through CMG2 rather than TEM8.

EXAMPLE 19

[0230] This example demonstrates a method of treating oral melanoma in cats by administering IC2-PA and LF.

[0231] In a phase 0 trial with a goal to establish preliminary safety and efficacy data, five cats were enrolled and treated with a microdose (15 μg PA-U2-R200A + 15 μg PA-L1-I207R + 10 μg LF, defined as less than 1/100th of an observed safe dose in mice) three times in a week by intratumoral injection. All of the cats tolerated the treatments well with no significant side effects. Two cats had an increase in grooming behavior and ability to eat. One cat had a 31% reduction in tumor volume after three injections of IC2-PA + LF, as measured by computerized tomography (CT) scan. Four cats had stable disease. Histology analyses showed the rate of tumor cell apoptosis was significantly increased in three cats.

[0232] Encouraged by the results of the phase 0 trial, a combined phase I/II dose dose-finding and efficacy trial was initiated. The first and second cohorts of this trial received six injections of a dose 5x or 25x greater than in the phase 0 trial, respectively. A reduction in tumor volume has been observed in three of the five cats with no adverse effects. The trial is currently ongoing.

EXAMPLE 20

[0233] This example demonstrates a method of treating oral melanoma in dogs by administering IC2-PA and LF.

[0234] In this ongoing phase 1 trial, two dogs with oral melanoma were enrolled and treated with 375 µg PA-U2-R200A + 375 µg PA-L1-I207R + 250 µg LF, three times a week for two weeks by intratumoral injection, followed by surgical removal of the tumors. The two dogs tolerated the treatments well with no significant side effects. The dogs' body weight and tumor sizes were measured throughout the study. The results are shown in Figures 19A-19D. One dog had a 50% reduction in tumor volume (Figure 19A) and the other experienced a modest reduction (Figure 19C). The trial is currently ongoing.

[0235] All references, including publications, patent applications, and patents, cited herein are hereby incorporated by reference to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein.

[0236] The use of the terms “a” and “an” and “the” and “at least one” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The use of the term “at least one” followed by a list of one or more items (for example, “at least one of A and B”) is to be construed to mean one item selected from the listed items (A or B) or any combination of two or more of the listed items (A and B), unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (i.e., meaning “including, but not limited to,”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[0237] Preferred embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those preferred

embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

[0238] In a first aspect, the invention relates to a *Bacillus anthracis* protective antigen (PA) comprising a PA amino acid sequence, wherein one or more of amino acid residues I207, I210, and I656, as defined by reference to SEQ ID NO: 1, are, independently, substituted,

wherein the one or more substitutions are selected from

(a) amino acid residue I207 substituted with the proviso that amino acid residue I207 is not substituted with alanine;

(b) amino acid residue I210 substituted with aspartic acid, glutamic acid, lysine, glutamine, arginine, or serine; and

(c) amino acid residue I656 substituted with glutamine, valine, alanine, or glutamic acid;
and

wherein amino acid residues E654, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted or wild-type.

[0238a] In a second aspect, the invention relates to a nucleic acid comprising a nucleotide sequence encoding the PA according to the first aspect.

[0238b] In a third aspect, the invention relates to a recombinant expression vector comprising the nucleic acid according to the second aspect.

[0238c] In a fourth aspect, the invention relates to a host cell comprising the recombinant expression vector according to the third aspect.

[0239] In a fifth aspect, the invention relates to a composition comprising a first *Bacillus anthracis* protective antigen (PA) comprising a first PA amino acid sequence and a second PA

comprising a second PA amino acid sequence, wherein the second PA amino acid sequence is different from the first PA amino acid sequence,

wherein the first PA has one or more of amino acid residues I207, I210, E654, I656, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, independently substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine, and

wherein amino acid residue R200 of the second PA amino acid sequence, as defined by reference to SEQ ID NO: 1, is substituted.

[0239a] In a sixth aspect, the invention relates to a composition comprising the PA of the first aspect, the nucleic acid of the second aspect, the recombinant expression vector of the third aspect, the host cell of the fourth aspect, or the composition according to the fifth aspect, further comprising a pharmaceutically acceptable carrier formulated for parenteral administration.

[0239b] In a seventh aspect, the invention relates to a method of treating or preventing cancer in a mammal, the method comprising administering to the mammal the PA of the first aspect, the nucleic acid of the second aspect, the recombinant expression vector of the third aspect, the host cell of the fourth aspect, or the composition according to the fifth aspect or the sixth aspect and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF, in an amount effective to treat or prevent cancer in the mammal.

[0239c] In an eighth aspect, the invention relates to use of the PA of the first aspect, the nucleic acid of the second aspect, the recombinant expression vector of the third aspect, the host cell of the fourth aspect, or the composition according to the fifth aspect or the sixth aspect and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in the manufacture of a medicament to treat or prevent cancer in a mammal.

[0239d] In a ninth aspect, the invention relates to a method for inhibiting growth of a target cell, the method comprising contacting the cell with the PA of the first aspect, the nucleic acid of the second aspect, the recombinant expression vector of the third aspect, the host cell of the fourth aspect, or the composition according to the fifth aspect or the sixth aspect and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in an amount effective to inhibit growth of the target cell.

[0239e] In a tenth aspect, the invention relates to use of the PA of the first aspect, the nucleic acid of the second aspect, the recombinant expression vector of the third aspect, the host cell of the fourth aspect, or the composition according to the fifth aspect or the sixth aspect and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in the manufacture of a medicament to inhibit growth of a target cell.

[0240] The term “comprise” and variants of the term such as “comprises” or “comprising” are used herein to denote the inclusion of a stated integer or stated integers but not to exclude any other integer or any other integers, unless in the context or usage an exclusive interpretation of the term is required.

[0241] Any reference to any prior art in this specification is not, and should not be taken as an acknowledgement or any form of suggestion that the prior art forms part of the common general knowledge.

CLAIMS:

1. A *Bacillus anthracis* protective antigen (PA) comprising a PA amino acid sequence, wherein one or more of amino acid residues I207, I210, and I656, as defined by reference to SEQ ID NO: 1, are, independently, substituted,

wherein the one or more substitutions are selected from

(a) amino acid residue I207 substituted with the proviso that amino acid residue I207 is not substituted with alanine;

(b) amino acid residue I210 substituted with aspartic acid, glutamic acid, lysine, glutamine, arginine, or serine; and

(c) amino acid residue I656 substituted with glutamine, valine, alanine, or glutamic acid;

and

wherein amino acid residues E654, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, are, independently, substituted or wild-type.

2. The PA according to claim 1, wherein one or both of amino acid residues I207 and I210 are, independently, substituted,

wherein the one or both substitutions are selected from

(a) amino acid residue I207 substituted with the proviso that amino acid residue I207 is not substituted with alanine; and

(b) amino acid residue I210 substituted with aspartic acid, glutamic acid, lysine, glutamine, arginine, or serine.

3. The PA according to claim 1 or 2, wherein amino acid residue I210 is substituted with aspartic acid, glutamic acid, lysine, glutamine, arginine, or serine.

4. The PA according to claim 1 or 2, wherein amino acid residue I207 is substituted with arginine, tryptophan, or tyrosine.

5. The PA according to claim 1 or 2, wherein amino acid residue I207 is substituted with arginine.
6. The PA according to any one of claims 1 and 3-5, wherein one or both of amino acid residues Y681 and L687 are, independently, substituted.
7. The PA according to any one of claims 1 and 3-6, wherein amino acid residue I656 is substituted with glutamine, valine, alanine, or glutamic acid.
8. The PA according to any one of claims 1 and 3-7, wherein amino acid residue Y681 is substituted with alanine.
9. The PA according to any one of claims 1 and 3-8, wherein amino acid residue L687 is substituted with alanine.
10. The PA according to any one of claims 1, 3-5, and 7-9, wherein one or more of amino acid residues E654, R659, and M662 are, independently, substituted.
11. The PA according to any one of claims 1, 3-5, and 7-10, wherein amino acid residue E654 is substituted with threonine.
12. The PA according to any one of claims 1, 3-5, and 7-11, wherein amino acid residue R659 is substituted with serine.
13. The PA according to any one of claims 1, 3-5, and 7-12, wherein amino acid residue M662 is substituted with arginine.

14. The PA according to any one of claims 1-13, comprising a matrix metalloproteinase (MMP) cleavage site.
15. The PA according to claim 14, wherein the MMP cleavage site is SEQ ID NO: 6 or 7.
16. A nucleic acid comprising a nucleotide sequence encoding the PA according to any one of claims 1-15.
17. A recombinant expression vector comprising the nucleic acid according to claim 16.
18. A host cell comprising the recombinant expression vector according to claim 17.
19. A composition comprising a first *Bacillus anthracis* protective antigen (PA) comprising a first PA amino acid sequence and a second PA comprising a second PA amino acid sequence, wherein the second PA amino acid sequence is different from the first PA amino acid sequence,
- wherein the first PA has one or more of amino acid residues I207, I210, E654, I656, R659, M662, Y681, and L687, as defined by reference to SEQ ID NO: 1, independently substituted, with the proviso that amino acid residue I207 is not substituted with alanine and amino acid residue I210 is not substituted with alanine, and
- wherein amino acid residue R200 of the second PA amino acid sequence, as defined by reference to SEQ ID NO: 1, is substituted.
20. The composition according to claim 19, wherein the one or more substitutions within the first PA are selected from
- (a) amino acid residue I207 substituted with the proviso that amino acid residue I207 is not substituted with alanine;

- (b) amino acid residue I210 substituted with aspartic acid, glutamic acid, lysine, glutamine, arginine, or serine;
- (c) amino acid residue I656 substituted with glutamine, valine, alanine, cysteine, or glutamic acid;
- (d) amino acid residue R659 substituted; and
- (e) amino acid residue M662 substituted with arginine.

21. The composition according to claim 19 or 20, wherein within the first PA amino acid residue I207 is substituted with arginine.

22. A composition comprising a first *Bacillus anthracis* protective antigen (PA) comprising a first PA amino acid sequence and a second PA comprising a second PA amino acid sequence, wherein the second PA amino acid sequence is different from the first PA amino acid sequence,

wherein the first PA is the PA according to any one of claims 1-15, and

wherein amino acid residue R200 of the second PA amino acid sequence, as defined by reference to SEQ ID NO: 1, is substituted.

23. The composition according to any one of claims 19-22, wherein amino acid residue R200 of the second PA amino acid sequence is substituted with alanine, cysteine, aspartic acid, glutamic acid, glycine, isoleucine, methionine, proline, serine, valine, or tryptophan.

24. The composition according to any one of claims 19-22, wherein amino acid residue R200 of the second PA amino acid sequence is substituted with alanine.

25. The composition according to any one of claims 19-24, wherein amino acid residue I656 of the first PA amino acid sequence, the second PA amino acid sequence, or both

the first and second PA amino acid sequence, as defined by reference to SEQ ID NO: 1, is substituted.

26. The composition according to claim 25, wherein amino acid residue I656 of the first PA amino acid sequence, the second PA amino acid sequence, or both the first and second PA amino acid sequence, is substituted with glutamine, valine, alanine, cysteine, or glutamic acid.

27. The composition according to claim 26, wherein amino acid residue I656 of the first PA amino acid sequence is substituted with glutamine, and wherein amino acid residue I656 of the second PA amino acid sequence is substituted with glutamine.

28. The composition according to any one of claims 19-27, wherein the second PA comprises a plasminogen activator cleavage site.

29. The composition according to claim 28, wherein the plasminogen activator cleavage site is selected from the group consisting of SEQ ID NOs: 8-15.

30. The composition according to any one of claims 19-29, further comprising one or more of (i) *Bacillus anthracis* lethal factor (LF), (ii) *Bacillus anthracis* edema factor (EF), (iii) FP59, and (iv) cytolethal distending toxin subunit B (CdtB) conjugated or fused to *Bacillus anthracis* toxin LF.

31. A composition comprising the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-30, further comprising a pharmaceutically acceptable carrier formulated for parenteral administration.

32. The composition according to any one of claims 19-31, further comprising pentostatin and cyclophosphamide.

33. A method of treating or preventing cancer in a mammal, the method comprising administering to the mammal the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-32 and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF, in an amount effective to treat or prevent cancer in the mammal.

34. Use of the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-32 and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in the manufacture of a medicament to treat or prevent cancer in a mammal.

35. A method of treating or preventing cancer in a mammal, the method comprising administering to the mammal a set comprising (a) pentostatin and cyclophosphamide, (b) the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-32, and (c) one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in an amount effective to treat or prevent cancer in the mammal.

36. Use of a set comprising (a) pentostatin and cyclophosphamide, (b) the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-32, and (c) one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in the manufacture of a medicament to treat or prevent cancer in a mammal.

37. The method according to claim 35 or use according to claim 36, wherein (a) is to be administered prior to or sequentially with the administration of (b).

38. The method according to any one of claims 33, 35, and 37 or the use according to any one of claims 34, 36, and 37, wherein the cancer is a solid tumor.

39. The method according to claim 38 or the use according to claim 38, wherein endothelial cells of the solid tumor express CMG2.

40. The method according to claim 38 or 39 or the use according to claim 38 or 39, wherein the solid tumor is an oral squamous carcinoma tumor, a melanoma tumor, a lung tumor, or an ovarian tumor.

41. The method according to any one of claims 33, 35, 37, 38, 39, and 40 or the use according to any one of claims 34, 36, 37, 38, 39, and 40, wherein the mammal is a human, a mouse, a cat, or a dog.

42. A method for inhibiting growth of a target cell, the method comprising contacting the cell with the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-32 and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in an amount effective to inhibit growth of the target cell.

43. Use of the PA of any one of claims 1-15, the nucleic acid of claim 16, the recombinant expression vector of claim 17, the host cell of claim 18, or the composition according to any one of claims 19-32 and one or more of (i) *Bacillus anthracis* LF, (ii) *Bacillus anthracis* EF, (iii) FP59, and (iv) CdtB conjugated or fused to *Bacillus anthracis* toxin LF in the manufacture of a medicament to inhibit growth of a target cell.

**The United States of America, as represented by the Secretary, Department of Health
and Human Services**

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 6




FIG. 7




FIG. 7




FIG. 8





FIG. 8

12/27




FIG. 9




FIG. 9




FIG. 10




FIG. 10




FIG. 11




FIG. 12




FIG. 12




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19

726173_ST25
SEQUENCE LISTING

<110> THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE
SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES

<120> MODIFIED ANTHRAX TOXIN PROTECTIVE ANTIGEN

<130> 726173

<150> US 62/323,218

<151> 2016-04-15

<150> US 62/210,771

<151> 2015-08-27

<160> 39

<170> PatentIn version 3.5

<210> 1

<211> 735

<212> PRT

<213> Bacillus anthracis

<400> 1

Glu Val Lys Gln Glu Asn Arg Leu Leu Asn Glu Ser Glu Ser Ser Ser
1 5 10 15

Gln Gly Leu Leu Gly Tyr Tyr Phe Ser Asp Leu Asn Phe Gln Ala Pro
20 25 30

Met Val Val Thr Ser Ser Thr Thr Gly Asp Leu Ser Ile Pro Ser Ser
35 40 45

Glu Leu Glu Asn Ile Pro Ser Glu Asn Gln Tyr Phe Gln Ser Ala Ile
50 55 60

Trp Ser Gly Phe Ile Lys Val Lys Lys Ser Asp Glu Tyr Thr Phe Ala
65 70 75 80

Thr Ser Ala Asp Asn His Val Thr Met Trp Val Asp Asp Gln Glu Val
85 90 95

Ile Asn Lys Ala Ser Asn Ser Asn Lys Ile Arg Leu Glu Lys Gly Arg
100 105 110

Leu Tyr Gln Ile Lys Ile Gln Tyr Gln Arg Glu Asn Pro Thr Glu Lys
115 120 125

Gly Leu Asp Phe Lys Leu Tyr Trp Thr Asp Ser Gln Asn Lys Lys Glu
130 135 140

Val Ile Ser Ser Asp Asn Leu Gln Leu Pro Glu Leu Lys Gln Lys Ser
145 150 155 160

Ser Asn Ser Arg Lys Lys Arg Ser Thr Ser Ala Gly Pro Thr Val Pro
165 170 175

726173_ST25

Asp Arg Asp Asn Asp Gly Ile Pro Asp Ser Leu Glu Val Glu Gly Tyr
 180 185 190
 Thr Val Asp Val Lys Asn Lys Arg Thr Phe Leu Ser Pro Trp Ile Ser
 195 200 205
 Asn Ile His Glu Lys Lys Gly Leu Thr Lys Tyr Lys Ser Ser Pro Glu
 210 215 220
 Lys Trp Ser Thr Ala Ser Asp Pro Tyr Ser Asp Phe Glu Lys Val Thr
 225 230 235 240
 Gly Arg Ile Asp Lys Asn Val Ser Pro Glu Ala Arg His Pro Leu Val
 245 250 255
 Ala Ala Tyr Pro Ile Val His Val Asp Met Glu Asn Ile Ile Leu Ser
 260 265 270
 Lys Asn Glu Asp Gln Ser Thr Gln Asn Thr Asp Ser Gln Thr Arg Thr
 275 280 285
 Ile Ser Lys Asn Thr Ser Thr Ser Arg Thr His Thr Ser Glu Val His
 290 295 300
 Gly Asn Ala Glu Val His Ala Ser Phe Phe Asp Ile Gly Gly Ser Val
 305 310 315 320
 Ser Ala Gly Phe Ser Asn Ser Asn Ser Ser Thr Val Ala Ile Asp His
 325 330 335
 Ser Leu Ser Leu Ala Gly Glu Arg Thr Trp Ala Glu Thr Met Gly Leu
 340 345 350
 Asn Thr Ala Asp Thr Ala Arg Leu Asn Ala Asn Ile Arg Tyr Val Asn
 355 360 365
 Thr Gly Thr Ala Pro Ile Tyr Asn Val Leu Pro Thr Thr Ser Leu Val
 370 375 380
 Leu Gly Lys Asn Gln Thr Leu Ala Thr Ile Lys Ala Lys Glu Asn Gln
 385 390 395 400
 Leu Ser Gln Ile Leu Ala Pro Asn Asn Tyr Tyr Pro Ser Lys Asn Leu
 405 410 415
 Ala Pro Ile Ala Leu Asn Ala Gln Asp Asp Phe Ser Ser Thr Pro Ile
 420 425 430
 Thr Met Asn Tyr Asn Gln Phe Leu Glu Leu Glu Lys Thr Lys Gln Leu
 435 440 445

726173_ST25

Arg Leu Asp Thr Asp Gln Val Tyr Gly Asn Ile Ala Thr Tyr Asn Phe
 450 455 460
 Glu Asn Gly Arg Val Arg Val Asp Thr Gly Ser Asn Trp Ser Glu Val
 465 470 475 480
 Leu Pro Gln Ile Gln Glu Thr Thr Ala Arg Ile Ile Phe Asn Gly Lys
 485 490 495
 Asp Leu Asn Leu Val Glu Arg Arg Ile Ala Ala Val Asn Pro Ser Asp
 500 505 510
 Pro Leu Glu Thr Thr Lys Pro Asp Met Thr Leu Lys Glu Ala Leu Lys
 515 520 525
 Ile Ala Phe Gly Phe Asn Glu Pro Asn Gly Asn Leu Gln Tyr Gln Gly
 530 535 540
 Lys Asp Ile Thr Glu Phe Asp Phe Asn Phe Asp Gln Gln Thr Ser Gln
 545 550 555 560
 Asn Ile Lys Asn Gln Leu Ala Glu Leu Asn Ala Thr Asn Ile Tyr Thr
 565 570 575
 Val Leu Asp Lys Ile Lys Leu Asn Ala Lys Met Asn Ile Leu Ile Arg
 580 585 590
 Asp Lys Arg Phe His Tyr Asp Arg Asn Asn Ile Ala Val Gly Ala Asp
 595 600 605
 Glu Ser Val Val Lys Glu Ala His Arg Glu Val Ile Asn Ser Ser Thr
 610 615 620
 Glu Gly Leu Leu Leu Asn Ile Asp Lys Asp Ile Arg Lys Ile Leu Ser
 625 630 635 640
 Gly Tyr Ile Val Glu Ile Glu Asp Thr Glu Gly Leu Lys Glu Val Ile
 645 650 655
 Asn Asp Arg Tyr Asp Met Leu Asn Ile Ser Ser Leu Arg Gln Asp Gly
 660 665 670
 Lys Thr Phe Ile Asp Phe Lys Lys Tyr Asn Asp Lys Leu Pro Leu Tyr
 675 680 685
 Ile Ser Asn Pro Asn Tyr Lys Val Asn Val Tyr Ala Val Thr Lys Glu
 690 695 700
 Asn Thr Ile Ile Asn Pro Ser Glu Asn Gly Asp Thr Ser Thr Asn Gly
 705 710 715 720

726173_ST25

Ile Lys Lys Ile Leu Ile Phe Ser Lys Lys Gly Tyr Glu Ile Gly
 725 730 735

<210> 2
 <211> 764
 <212> PRT
 <213> Bacillus anthracis

<400> 2

Met Lys Lys Arg Lys Val Leu Ile Pro Leu Met Ala Leu Ser Thr Ile
 1 5 10 15

Leu Val Ser Ser Thr Gly Asn Leu Glu Val Ile Gln Ala Glu Val Lys
 20 25 30

Gln Glu Asn Arg Leu Leu Asn Glu Ser Glu Ser Ser Ser Gln Gly Leu
 35 40 45

Leu Gly Tyr Tyr Phe Ser Asp Leu Asn Phe Gln Ala Pro Met Val Val
 50 55 60

Thr Ser Ser Thr Thr Gly Asp Leu Ser Ile Pro Ser Ser Glu Leu Glu
 65 70 75 80

Asn Ile Pro Ser Glu Asn Gln Tyr Phe Gln Ser Ala Ile Trp Ser Gly
 85 90 95

Phe Ile Lys Val Lys Lys Ser Asp Glu Tyr Thr Phe Ala Thr Ser Ala
 100 105 110

Asp Asn His Val Thr Met Trp Val Asp Asp Gln Glu Val Ile Asn Lys
 115 120 125

Ala Ser Asn Ser Asn Lys Ile Arg Leu Glu Lys Gly Arg Leu Tyr Gln
 130 135 140

Ile Lys Ile Gln Tyr Gln Arg Glu Asn Pro Thr Glu Lys Gly Leu Asp
 145 150 155 160

Phe Lys Leu Tyr Trp Thr Asp Ser Gln Asn Lys Lys Glu Val Ile Ser
 165 170 175

Ser Asp Asn Leu Gln Leu Pro Glu Leu Lys Gln Lys Ser Ser Asn Ser
 180 185 190

Arg Lys Lys Arg Ser Thr Ser Ala Gly Pro Thr Val Pro Asp Arg Asp
 195 200 205

Asn Asp Gly Ile Pro Asp Ser Leu Glu Val Glu Gly Tyr Thr Val Asp
 210 215 220

726173_ST25

Val Lys Asn Lys Arg Thr Phe Leu Ser Pro Trp Ile Ser Asn Ile His
 225 230 235 240
 Glu Lys Lys Gly Leu Thr Lys Tyr Lys Ser Ser Pro Glu Lys Trp Ser
 245 250 255
 Thr Ala Ser Asp Pro Tyr Ser Asp Phe Glu Lys Val Thr Gly Arg Ile
 260 265 270
 Asp Lys Asn Val Ser Pro Glu Ala Arg His Pro Leu Val Ala Ala Tyr
 275 280 285
 Pro Ile Val His Val Asp Met Glu Asn Ile Ile Leu Ser Lys Asn Glu
 290 295 300
 Asp Gln Ser Thr Gln Asn Thr Asp Ser Gln Thr Arg Thr Ile Ser Lys
 305 310 315 320
 Asn Thr Ser Thr Ser Arg Thr His Thr Ser Glu Val His Gly Asn Ala
 325 330 335
 Glu Val His Ala Ser Phe Phe Asp Ile Gly Gly Ser Val Ser Ala Gly
 340 345 350
 Phe Ser Asn Ser Asn Ser Ser Thr Val Ala Ile Asp His Ser Leu Ser
 355 360 365
 Leu Ala Gly Glu Arg Thr Trp Ala Glu Thr Met Gly Leu Asn Thr Ala
 370 375 380
 Asp Thr Ala Arg Leu Asn Ala Asn Ile Arg Tyr Val Asn Thr Gly Thr
 385 390 395 400
 Ala Pro Ile Tyr Asn Val Leu Pro Thr Thr Ser Leu Val Leu Gly Lys
 405 410 415
 Asn Gln Thr Leu Ala Thr Ile Lys Ala Lys Glu Asn Gln Leu Ser Gln
 420 425 430
 Ile Leu Ala Pro Asn Asn Tyr Tyr Pro Ser Lys Asn Leu Ala Pro Ile
 435 440 445
 Ala Leu Asn Ala Gln Asp Asp Phe Ser Ser Thr Pro Ile Thr Met Asn
 450 455 460
 Tyr Asn Gln Phe Leu Glu Leu Glu Lys Thr Lys Gln Leu Arg Leu Asp
 465 470 475 480
 Thr Asp Gln Val Tyr Gly Asn Ile Ala Thr Tyr Asn Phe Glu Asn Gly
 485 490 495

726173_ST25

Arg Val Arg Val Asp Thr Gly Ser Asn Trp Ser Glu Val Leu Pro Gln
 500 505 510
 Ile Gln Glu Thr Thr Ala Arg Ile Ile Phe Asn Gly Lys Asp Leu Asn
 515 520 525
 Leu Val Glu Arg Arg Ile Ala Ala Val Asn Pro Ser Asp Pro Leu Glu
 530 535 540
 Thr Thr Lys Pro Asp Met Thr Leu Lys Glu Ala Leu Lys Ile Ala Phe
 545 550 555 560
 Gly Phe Asn Glu Pro Asn Gly Asn Leu Gln Tyr Gln Gly Lys Asp Ile
 565 570 575
 Thr Glu Phe Asp Phe Asn Phe Asp Gln Gln Thr Ser Gln Asn Ile Lys
 580 585 590
 Asn Gln Leu Ala Glu Leu Asn Ala Thr Asn Ile Tyr Thr Val Leu Asp
 595 600 605
 Lys Ile Lys Leu Asn Ala Lys Met Asn Ile Leu Ile Arg Asp Lys Arg
 610 615 620
 Phe His Tyr Asp Arg Asn Asn Ile Ala Val Gly Ala Asp Glu Ser Val
 625 630 635 640
 Val Lys Glu Ala His Arg Glu Val Ile Asn Ser Ser Thr Glu Gly Leu
 645 650 655
 Leu Leu Asn Ile Asp Lys Asp Ile Arg Lys Ile Leu Ser Gly Tyr Ile
 660 665 670
 Val Glu Ile Glu Asp Thr Glu Gly Leu Lys Glu Val Ile Asn Asp Arg
 675 680 685
 Tyr Asp Met Leu Asn Ile Ser Ser Leu Arg Gln Asp Gly Lys Thr Phe
 690 695 700
 Ile Asp Phe Lys Lys Tyr Asn Asp Lys Leu Pro Leu Tyr Ile Ser Asn
 705 710 715 720
 Pro Asn Tyr Lys Val Asn Val Tyr Ala Val Thr Lys Glu Asn Thr Ile
 725 730 735
 Ile Asn Pro Ser Glu Asn Gly Asp Thr Ser Thr Asn Gly Ile Lys Lys
 740 745 750
 Ile Leu Ile Phe Ser Lys Lys Gly Tyr Glu Ile Gly
 755 760

726173_ST25

<210> 3
 <211> 29
 <212> PRT
 <213> Bacillus anthracis

<400> 3

Met Lys Lys Arg Lys Val Leu Ile Pro Leu Met Ala Leu Ser Thr Ile
 1 5 10 15

Leu Val Ser Ser Thr Gly Asn Leu Glu Val Ile Gln Ala
 20 25

<210> 4
 <211> 4
 <212> PRT
 <213> Bacillus anthracis

<400> 4

Arg Lys Lys Arg
 1

<210> 5
 <211> 764
 <212> PRT
 <213> Bacillus anthracis

<400> 5

Met Lys Lys Arg Lys Val Leu Ile Pro Leu Met Ala Leu Ser Thr Ile
 1 5 10 15

Leu Val Ser Ser Thr Gly Asn Leu Glu Val Ile Gln Ala Glu Val Lys
 20 25 30

Gln Glu Asn Arg Leu Leu Asn Glu Ser Glu Ser Ser Ser Gln Gly Leu
 35 40 45

Leu Gly Tyr Tyr Phe Ser Asp Leu Asn Phe Gln Ala Pro Met Val Val
 50 55 60

Thr Ser Ser Thr Thr Gly Asp Leu Ser Ile Pro Ser Ser Glu Leu Glu
 65 70 75 80

Asn Ile Pro Ser Glu Asn Gln Tyr Phe Gln Ser Ala Ile Trp Ser Gly
 85 90 95

Phe Ile Lys Val Lys Lys Ser Asp Glu Tyr Thr Phe Ala Thr Ser Ala
 100 105 110

Asp Asn His Val Thr Met Trp Val Asp Asp Gln Glu Val Ile Asn Lys
 115 120 125

Ala Ser Asn Ser Asn Lys Ile Arg Leu Glu Lys Gly Arg Leu Tyr Gln
 130 135 140

726173_ST25

Ile Lys Ile Gln Tyr Gln Arg Glu Asn Pro Thr Glu Lys Gly Leu Asp
 145 150 155 160
 Phe Lys Leu Tyr Trp Thr Asp Ser Gln Asn Lys Lys Glu Val Ile Ser
 165 170 175
 Ser Asp Asn Leu Gln Leu Pro Glu Leu Lys Gln Lys Ser Ser Asn Ser
 180 185 190
 Arg Lys Lys Arg Ser Thr Ser Ala Gly Pro Thr Val Pro Asp Arg Asp
 195 200 205
 Asn Asp Gly Ile Pro Asp Ser Leu Glu Val Glu Gly Tyr Thr Val Asp
 210 215 220
 Val Lys Asn Lys Arg Thr Phe Leu Ser Pro Trp Ile Ser Asn Ile His
 225 230 235 240
 Glu Lys Lys Gly Leu Thr Lys Tyr Lys Ser Ser Pro Glu Lys Trp Ser
 245 250 255
 Thr Ala Ser Asp Pro Tyr Ser Asp Phe Glu Lys Val Thr Gly Arg Ile
 260 265 270
 Asp Lys Asn Val Ser Pro Glu Ala Arg His Pro Leu Val Ala Ala Tyr
 275 280 285
 Pro Ile Val His Val Asp Met Glu Asn Ile Ile Leu Ser Lys Asn Glu
 290 295 300
 Asp Gln Ser Thr Gln Asn Thr Asp Ser Gln Thr Arg Thr Ile Ser Lys
 305 310 315 320
 Asn Thr Ser Thr Ser Arg Thr His Thr Ser Glu Val His Gly Asn Ala
 325 330 335
 Glu Val His Ala Ser Phe Phe Asp Ile Gly Gly Ser Val Ser Ala Gly
 340 345 350
 Phe Ser Asn Ser Asn Ser Ser Thr Val Ala Ile Asp His Ser Leu Ser
 355 360 365
 Leu Ala Gly Glu Arg Thr Trp Ala Glu Thr Met Gly Leu Asn Thr Ala
 370 375 380
 Asp Thr Ala Arg Leu Asn Ala Asn Ile Arg Tyr Val Asn Thr Gly Thr
 385 390 395 400
 Ala Pro Ile Tyr Asn Val Leu Pro Thr Thr Ser Leu Val Leu Gly Lys
 405 410 415

726173_ST25

Asn Gln Thr Leu Ala Thr Ile Lys Ala Lys Glu Asn Gln Leu Ser Gln
 420 425 430
 Ile Leu Ala Pro Asn Asn Tyr Tyr Pro Ser Lys Asn Leu Ala Pro Ile
 435 440 445
 Ala Leu Asn Ala Gln Asp Asp Phe Ser Ser Thr Pro Ile Thr Met Asn
 450 455 460
 Tyr Asn Gln Phe Leu Glu Leu Glu Lys Thr Lys Gln Leu Arg Leu Asp
 465 470 475 480
 Thr Asp Gln Val Tyr Gly Asn Ile Ala Thr Tyr Asn Phe Glu Asn Gly
 485 490 495
 Arg Val Arg Val Asp Thr Gly Ser Asn Trp Ser Glu Val Leu Pro Gln
 500 505 510
 Ile Gln Glu Thr Thr Ala Arg Ile Ile Phe Asn Gly Lys Asp Leu Asn
 515 520 525
 Leu Val Glu Arg Arg Ile Ala Ala Val Asn Pro Ser Asp Pro Leu Glu
 530 535 540
 Thr Thr Lys Pro Asp Met Thr Leu Lys Glu Ala Leu Lys Ile Ala Phe
 545 550 555 560
 Gly Phe Asn Glu Pro Asn Gly Asn Leu Gln Tyr Gln Gly Lys Asp Ile
 565 570 575
 Thr Glu Phe Asp Phe Asn Phe Asp Gln Gln Thr Ser Gln Asn Ile Lys
 580 585 590
 Asn Gln Leu Ala Glu Leu Asn Val Thr Asn Ile Tyr Thr Val Leu Asp
 595 600 605
 Lys Ile Lys Leu Asn Ala Lys Met Asn Ile Leu Ile Arg Asp Lys Arg
 610 615 620
 Phe His Tyr Asp Arg Asn Asn Ile Ala Val Gly Ala Asp Glu Ser Val
 625 630 635 640
 Val Lys Glu Ala His Arg Glu Val Ile Asn Ser Ser Thr Glu Gly Leu
 645 650 655
 Leu Leu Asn Ile Asp Lys Asp Ile Arg Lys Ile Leu Ser Gly Tyr Ile
 660 665 670
 Val Glu Ile Glu Asp Thr Glu Gly Leu Lys Glu Val Ile Asn Asp Arg
 675 680 685

726173_ST25

Tyr Asp Met Leu Asn Ile Ser Ser Leu Arg Gln Asp Gly Lys Thr Phe
690 695 700

Ile Asp Phe Lys Lys Tyr Asn Asp Lys Leu Pro Leu Tyr Ile Ser Asn
705 710 715 720

Pro Asn Tyr Lys Val Asn Val Tyr Ala Val Thr Lys Glu Asn Thr Ile
725 730 735

Ile Asn Pro Ser Glu Asn Gly Asp Thr Ser Thr Asn Gly Ile Lys Lys
740 745 750

Ile Leu Ile Phe Ser Lys Lys Gly Tyr Glu Ile Gly
755 760

<210> 6
<211> 8
<212> PRT
<213> Artificial sequence

<220>
<223> synthetic

<400> 6

Gly Pro Leu Gly Met Leu Ser Gln
1 5

<210> 7
<211> 8
<212> PRT
<213> Artificial sequence

<220>
<223> synthetic

<400> 7

Gly Pro Leu Gly Leu Trp Ala Gln
1 5

<210> 8
<211> 9
<212> PRT
<213> Artificial sequence

<220>
<223> synthetic

<400> 8

Pro Cys Pro Gly Arg Val Val Gly Gly
1 5

<210> 9
<211> 7
<212> PRT
<213> Artificial sequence

<220>

<223> synthetic

<400> 9

Pro Gly Ser Gly Arg Ser Ala
1 5

<210> 10

<211> 7

<212> PRT

<213> Artificial sequence

<220>

<223> synthetic

<400> 10

Pro Gly Ser Gly Lys Ser Ala
1 5

<210> 11

<211> 7

<212> PRT

<213> Artificial sequence

<220>

<223> synthetic

<400> 11

Pro Gln Arg Gly Arg Ser Ala
1 5

<210> 12

<211> 9

<212> PRT

<213> Artificial sequence

<220>

<223> synthetic

<400> 12

Pro Cys Pro Gly Arg Val Val Gly Gly
1 5

<210> 13

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic

<400> 13

Gly Ser Gly Arg Ser Ala
1 5

<210> 14

<211> 6

<212> PRT
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 14

Gly Ser Gly Lys Ser Ala
 1 5

<210> 15
 <211> 6
 <212> PRT
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 15

Gln Arg Gly Arg Ser Ala
 1 5

<210> 16
 <211> 21
 <212> DNA
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 16
 gtttagacgat aaaccagtcc t

21

<210> 17
 <211> 31
 <212> DNA
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 17
 ctaccagatt taaatccttt ccattaataaa t

31

<210> 18
 <211> 48
 <212> DNA
 <213> Artificial sequence

<220>
 <223> synthetic

<220>
 <221> misc_feature
 <222> (27)..(28)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (29)..(29)
 <223> n is g or c

726173_ST25

<400> 18
caagtgcctgg acctacggtt ccagacnng acaatgatgg aatcccta 48

<210> 19
<211> 24
<212> DNA
<213> Artificial sequence

<220>
<223> synthetic

<400> 19
ctggaaccgt aggtccagca cttg 24

<210> 20
<211> 58
<212> DNA
<213> Artificial sequence

<220>
<223> synthetic

<220>
<221> misc_feature
<222> (38)..(39)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (40)..(40)
<223> n is g or c

<400> 20
cgaggtagaa ggatatacgg ttgatgtcaa aaataaannn acctttcttt caccatgg 58

<210> 21
<211> 29
<212> DNA
<213> Artificial sequence

<220>
<223> synthetic

<400> 21
tgacatcaac cgtatatacct tctacctcg 29

<210> 22
<211> 58
<212> DNA
<213> Artificial sequence

<220>
<223> synthetic

<220>
<221> misc_feature
<222> (32)..(33)
<223> n is a, c, g, or t

<220>
<221> misc_feature

<222> (34)..(34)
 <223> n is g or c

 <400> 22
 caaaaataaa aggacctttc tttcaacatg gnnntctaata attcatgaaa agaaaggg 58

 <210> 23
 <211> 30
 <212> DNA
 <213> Artificial sequence

 <220>
 <223> synthetic

 <400> 23
 catggtgaaa gaaaggctct tttatttttg 30

 <210> 24
 <211> 58
 <212> DNA
 <213> Artificial sequence

 <220>
 <223> synthetic

 <220>
 <221> misc_feature
 <222> (41)..(42)
 <223> n is a, c, g, or t

 <220>
 <221> misc_feature
 <222> (43)..(43)
 <223> n is g or c

 <400> 24
 caaaaataaa aggacctttc tttcaccatg gatatactaata nncatgaaa agaaaggg 58

 <210> 25
 <211> 59
 <212> DNA
 <213> Artificial sequence

 <220>
 <223> synthetic

 <220>
 <221> misc_feature
 <222> (34)..(35)
 <223> n is a, c, g, or t

 <220>
 <221> misc_feature
 <222> (36)..(36)
 <223> n is g or c

 <400> 25
 ctttcacat ggatatctaa tattcatgaa aagnnngggt taaccaaata taaatcatc 59

 <210> 26
 <211> 33
 <212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 26

cttttcatga atattagata tccatggtga aag

33

<210> 27

<211> 59

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 27

aaaataaaaag aacttttctt tcaccatggc gttctaatat tcatgaaaag aaaggatta

59

<210> 28

<211> 59

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 28

taatcctttc ttttcatgaa tattagaacg ccatggtgaa agaaaagttc ttttatttt

59

<210> 29

<211> 2307

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 29

atgaaaaaac gaaaagtgtt aataccatta atggcattgt ctacgatatt agtttcaagc

60

acaggttaatt tagaggtgat tcaggcagaa gttaaacagg agaaccggtt attaaatgaa

120

tcagaatcaa gttcccagg gttactagga tactatttta gtgatttgaa ttttcaagca

180

cccatggtgg ttacctcttc tactacagg gatttatcta ttcctagttc tgagttagaa

240

aatattccat cggaaaacca atattttcaa tctgctattt ggtcaggatt tatcaaagtt

300

aagaagagtg atgaatatac atttgctact tccgctgata atcatgtaac aatgtgggta

360

gatgaccaag aagtgattaa taaagcttct aattctaaca aaatcagatt agaaaaagga

420

agattatatac aaataaaaat tcaatatcaa cgagaaaatc ctactgaaaa aggattggat

480

ttcaagttgt actggaccga ttctcaaaat aaaaaagaag tgatttctag tgataactta

540

caattgccag aattaaaaca aaaatcttcg aactcaggac cattaggaat gttgagtcaa

600

agtacaagtg ctggacctac ggttcagac cgtgacaatg atggaatccc tgattcatta

660

gaggtagaag gatatacggg tgatgtcaaa aataaaagaa cttttctttc accatggcgc

720

tctaataattc atgaaaagaa aggattaacc aaatataaat catctcctga aaaatggagc

780

726173_ST25

```

acggcttctg atccgtacag tgatttcgaa aagggttacag gacggattga taagaatgta      840
tcaccagagg caagacaccc ccttgtggca gcttatccga ttgtacatgt agatatggag      900
aatattattc tctcaaaaaa tgaggatcaa tccacacaga atactgatag tcaaacgaga      960
acaataagta aaaatacttc tacaagtagg acacatacta gtgaagtaca tggaaatgca     1020
gaagtgcatt cgtcgttctt tgatattggg gggagtgtat ctgcaggatt tagtaattcg     1080
aattcaagta cggtcgcaat tgatcattca ctatctctag caggggaaag aacttgggct     1140
gaaacaatgg gtttaaatac cgctgataca gcaagattaa atgccaatat tagatatgta     1200
aatactggga cggctccaat ctacaacgtg ttaccaacga cttcgttagt gttaggaaaa     1260
aatcaaacac tcgcgacaat taaagctaag gaaaaccaat taagtcaaat acttgcacct     1320
aataattatt atccttctaa aaacttggcg ccaatcgcat taaatgcaca agacgatttc     1380
agttctactc caattacaat gaattacaat caatttcttg agttagaaaa aacgaaacaa     1440
ttaagattag atacggatca agtatatggg aatatagcaa catacaattt tgaaaatgga     1500
agagtgaggg tggatacagg ctcgaactgg agtgaagtgt taccgcaaat tcaagaaaca     1560
actgcacgta tcatttttaa tggaaaagat ttaaatctgg tagaaaggcg gatagcggcg     1620
gttaatccta gtgatccatt agaaacgact aaaccggata tgacattaaa agaagccctt     1680
aaaatagcat ttggatttaa cgaaccgaat ggaaacttac aatatcaagg gaaagacata     1740
accgaatttg attttaattt cgatcaacaa acatctcaaa atatcaagaa tcagttagcg     1800
gaattaaacg caactaacat atatactgta ttagataaaa tcaaattaaa tgcaaaaatg     1860
aatattttta taagagataa acgttttcat tatgatagaa ataacatagc agttggggcg     1920
gatgagtcag tagttaagga ggctcataga gaagtaatta attcgtcaac agagggatta     1980
ttgttaaata ttgataagga tataagaaaa atattatcag gttatattgt agaaattgaa     2040
gatactgaag ggcttaaaga agttataaat gacagatatg atatgttgaa tatttctagt     2100
ttacggcaag atggaaaaac atttatagat tttaaaaaat ataatagata attaccgtta     2160
tatataagta atcccaatta taaggtaa atgtatgctg ttactaaaga aaacactatt     2220
attaatccta gtgagaatgg ggatactagt accaacggga tcaagaaaat tttaatcttt     2280
tctaaaaaag gctatgagat aggataa                                     2307

```

<210> 30
 <211> 768
 <212> PRT
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 30

Met Lys Lys Arg Lys Val Leu Ile Pro Leu Met Ala Leu Ser Thr Ile
 1 5 10 15

Leu Val Ser Ser Thr Gly Asn Leu Glu Val Ile Gln Ala Glu Val Lys

726173_ST25

20

25

30

Gln Glu Asn Arg Leu Leu Asn Glu Ser Glu Ser Ser Ser Gln Gly Leu
 35 40 45

Leu Gly Tyr Tyr Phe Ser Asp Leu Asn Phe Gln Ala Pro Met Val Val
 50 55 60

Thr Ser Ser Thr Thr Gly Asp Leu Ser Ile Pro Ser Ser Glu Leu Glu
 65 70 75 80

Asn Ile Pro Ser Glu Asn Gln Tyr Phe Gln Ser Ala Ile Trp Ser Gly
 85 90 95

Phe Ile Lys Val Lys Lys Ser Asp Glu Tyr Thr Phe Ala Thr Ser Ala
 100 105 110

Asp Asn His Val Thr Met Trp Val Asp Asp Gln Glu Val Ile Asn Lys
 115 120 125

Ala Ser Asn Ser Asn Lys Ile Arg Leu Glu Lys Gly Arg Leu Tyr Gln
 130 135 140

Ile Lys Ile Gln Tyr Gln Arg Glu Asn Pro Thr Glu Lys Gly Leu Asp
 145 150 155 160

Phe Lys Leu Tyr Trp Thr Asp Ser Gln Asn Lys Lys Glu Val Ile Ser
 165 170 175

Ser Asp Asn Leu Gln Leu Pro Glu Leu Lys Gln Lys Ser Ser Asn Ser
 180 185 190

Gly Pro Leu Gly Met Leu Ser Gln Ser Thr Ser Ala Gly Pro Thr Val
 195 200 205

Pro Asp Arg Asp Asn Asp Gly Ile Pro Asp Ser Leu Glu Val Glu Gly
 210 215 220

Tyr Thr Val Asp Val Lys Asn Lys Arg Thr Phe Leu Ser Pro Trp Arg
 225 230 235 240

Ser Asn Ile His Glu Lys Lys Gly Leu Thr Lys Tyr Lys Ser Ser Pro
 245 250 255

Glu Lys Trp Ser Thr Ala Ser Asp Pro Tyr Ser Asp Phe Glu Lys Val
 260 265 270

Thr Gly Arg Ile Asp Lys Asn Val Ser Pro Glu Ala Arg His Pro Leu
 275 280 285

Val Ala Ala Tyr Pro Ile Val His Val Asp Met Glu Asn Ile Ile Leu

290

295

Ser Lys Asn Glu Asp Gln Ser Thr Gln Asn Thr Asp Ser Gln Thr Arg
305 310 315 320

Thr Ile Ser Lys Asn Thr Ser Thr Ser Arg Thr His Thr Ser Glu Val
325 330 335

His Gly Asn Ala Glu Val His Ala Ser Phe Phe Asp Ile Gly Gly Ser
340 345 350

Val Ser Ala Gly Phe Ser Asn Ser Asn Ser Thr Val Ala Ile Asp
355 360 365

His Ser Leu Ser Leu Ala Gly Glu Arg Thr Trp Ala Glu Thr Met Gly
370 375 380

Leu Asn Thr Ala Asp Thr Ala Arg Leu Asn Ala Asn Ile Arg Tyr Val
385 390 395 400

Asn Thr Gly Thr Ala Pro Ile Tyr Asn Val Leu Pro Thr Thr Ser Leu
405 410 415

Val Leu Gly Lys Asn Gln Thr Leu Ala Thr Ile Lys Ala Lys Glu Asn
420 425 430

Gln Leu Ser Gln Ile Leu Ala Pro Asn Asn Tyr Tyr Pro Ser Lys Asn
435 440 445

Leu Ala Pro Ile Ala Leu Asn Ala Gln Asp Asp Phe Ser Ser Thr Pro
450 455 460

Ile Thr Met Asn Tyr Asn Gln Phe Leu Glu Leu Glu Lys Thr Lys Gln
465 470 475 480

Leu Arg Leu Asp Thr Asp Gln Val Tyr Gly Asn Ile Ala Thr Tyr Asn
485 490 495

Phe Glu Asn Gly Arg Val Arg Val Asp Thr Gly Ser Asn Trp Ser Glu
500 505 510

Val Leu Pro Gln Ile Gln Glu Thr Thr Ala Arg Ile Ile Phe Asn Gly
515 520 525

Lys Asp Leu Asn Leu Val Glu Arg Arg Ile Ala Ala Val Asn Pro Ser
530 535 540

Asp Pro Leu Glu Thr Thr Lys Pro Asp Met Thr Leu Lys Glu Ala Leu
545 550 555 560

Lys Ile Ala Phe Gly Phe Asn Glu Pro Asn Gly Asn Leu Gln Tyr Gln

Gly Lys Asp Ile Thr Glu Phe Asp Phe Asn Phe Asp Gln Gln Thr Ser
580 585 590

Gln Asn Ile Lys Asn Gln Leu Ala Glu Leu Asn Ala Thr Asn Ile Tyr
595 600 605

Thr Val Leu Asp Lys Ile Lys Leu Asn Ala Lys Met Asn Ile Leu Ile
610 615 620

Arg Asp Lys Arg Phe His Tyr Asp Arg Asn Asn Ile Ala Val Gly Ala
625 630 635 640

Asp Glu Ser Val Val Lys Glu Ala His Arg Glu Val Ile Asn Ser Ser
645 650 655

Thr Glu Gly Leu Leu Leu Asn Ile Asp Lys Asp Ile Arg Lys Ile Leu
660 665 670

Ser Gly Tyr Ile Val Glu Ile Glu Asp Thr Glu Gly Leu Lys Glu Val
675 680 685

Ile Asn Asp Arg Tyr Asp Met Leu Asn Ile Ser Ser Leu Arg Gln Asp
690 695 700

Gly Lys Thr Phe Ile Asp Phe Lys Lys Tyr Asn Asp Lys Leu Pro Leu
705 710 715 720

Tyr Ile Ser Asn Pro Asn Tyr Lys Val Asn Val Tyr Ala Val Thr Lys
725 730 735

Glu Asn Thr Ile Ile Asn Pro Ser Glu Asn Gly Asp Thr Ser Thr Asn
740 745 750

Gly Ile Lys Lys Ile Leu Ile Phe Ser Lys Lys Gly Tyr Glu Ile Gly
755 760 765

<210> 31
<211> 2304
<212> DNA
<213> Artificial sequence

<220>
<223> synthetic

<400> 31
atgaaaaaac gaaaagtgtt aataccatta atggcattgt ctacgatatt agtttcaagc 60
acaggttaatt tagaggtgat tcaggcagaa gttaaacagg agaaccgggtt attaatgaa 120
tcagaatcaa gttcccaggg gttactagga tactatttta gtgatttgaa ttttcaagca 180
cccatggtgg ttacctcttc tactacaggg gatttatcta ttcctagttc tgagttagaa 240

726173_ST25

aatattccat	cggaaaacca	atattttcaa	tctgctat	ggtcaggatt	tatcaaagtt	300
aagaagagt	atgaatatac	atttgctact	tccgctgata	atcatgtaac	aatgtgggta	360
gatgaccaag	aagtgattaa	taaagcttct	aatttctaaca	aaatcagatt	agaaaaagga	420
agattatatac	aaataaaaat	tcaatatcaa	cgagaaaatc	ctactgaaaa	aggattggat	480
ttcaagttgt	actggaccga	ttctcaaaaat	aaaaaagaag	tgattttctag	tgataactta	540
caattgccag	aattaaaaca	aaaatcttcg	aactcaccag	gaagtggaag	atcagcaagt	600
acaagtgctg	gacctacggt	tccagaccgt	gacaatgatg	gaatccctga	ttcattagag	660
gtagaaggat	atacggttga	tgtcaaaaat	aaagcaactt	ttctttcacc	atggattttct	720
aatattcatg	aaaagaaagg	attaaccaa	tataaatcat	ctcctgaaaa	atggagcacg	780
gcttctgatc	cgtacagtga	tttcgaaaag	gttacaggac	ggattgataa	gaatgtatca	840
ccagaggcaa	gacacccct	tgtggcagct	tatccgattg	tacatgtaga	tatggagaat	900
attattctct	caaaaaatga	ggatcaatcc	acacagaata	ctgatagtca	aacgagaaca	960
ataagtaaaa	atacttctac	aagtaggaca	catactagt	aagtacatgg	aaatgcagaa	1020
gtgcatgcgt	cgttctttga	tattgggtgg	agtgtatctg	caggatttag	taattcgaat	1080
tcaagtacgg	tcgcaattga	tcattcacta	tctctagcag	gggaaagaac	ttgggctgaa	1140
acaatgggtt	taaataccgc	tgatacagca	agattaaatg	ccaatattag	atatgtaaat	1200
actgggacgg	ctccaatcta	caacgtgtta	ccaacgactt	cgttagtggt	aggaaaaaat	1260
caaacactcg	cgacaattaa	agctaaggaa	aaccaattaa	gtcaaatact	tgcacctaata	1320
aattattatc	cttctaaaaa	cttggcgcca	atcgcattaa	atgcacaaga	cgatttcagt	1380
tctactccaa	ttacaatgaa	ttacaatcaa	tttcttgagt	tagaaaaaac	gaaacaatta	1440
agattagata	cggatcaagt	atatgggaat	atagcaacat	acaattttga	aaatggaaga	1500
gtgaggggtg	atacaggctc	gaactggagt	gaagtgttac	cgcaaattca	agaaacaact	1560
gcacgtatca	tttttaatgg	aaaagattta	aatctggtag	aaaggcggat	agcggcggtt	1620
aatcctagt	atccattaga	aacgactaaa	ccggatatga	cattaaaaga	agcccttaaa	1680
atagcatttg	gatttaacga	accgaatgga	aacttacaat	atcaaggga	agacataacc	1740
gaatttgatt	ttaatttcga	tcaacaaaca	tctcaaaaata	tcaagaatca	gttagcggaa	1800
ttaaacgcaa	ctaacaatata	tactgtatta	gataaaatca	aattaaatgc	aaaaatgaat	1860
attttaataa	gagataaacg	ttttcattat	gatagaaata	acatagcagt	tggggcggat	1920
gagtcagtag	ttaaggaggc	tcatagagaa	gtaattaatt	cgtcaacaga	gggattattg	1980
ttaaatattg	ataaggatat	aagaaaaata	ttatcagggt	atattgtaga	aattgaagat	2040
actgaagggc	ttaaagaagt	tataaatgac	agatatgata	tgttgaatat	ttctagttta	2100
cggcaagatg	gaaaaacatt	tatagatttt	aaaaaatata	atgataaatt	accgttatat	2160
ataagtaatc	ccaattataa	ggtaaatgta	tatgctgtta	ctaaagaaaa	cactattatt	2220
aatcctagt	agaatgggga	tactagtacc	aacgggatca	agaaaatttt	aatcttttct	2280

aaaaaaggct atgagatagg ataa

<210> 32
 <211> 767
 <212> PRT
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 32

Met Lys Lys Arg Lys Val Leu Ile Pro Leu Met Ala Leu Ser Thr Ile
 1 5 10 15

Leu Val Ser Ser Thr Gly Asn Leu Glu Val Ile Gln Ala Glu Val Lys
 20 25 30

Gln Glu Asn Arg Leu Leu Asn Glu Ser Glu Ser Ser Ser Gln Gly Leu
 35 40 45

Leu Gly Tyr Tyr Phe Ser Asp Leu Asn Phe Gln Ala Pro Met Val Val
 50 55 60

Thr Ser Ser Thr Thr Gly Asp Leu Ser Ile Pro Ser Ser Glu Leu Glu
 65 70 75 80

Asn Ile Pro Ser Glu Asn Gln Tyr Phe Gln Ser Ala Ile Trp Ser Gly
 85 90 95

Phe Ile Lys Val Lys Lys Ser Asp Glu Tyr Thr Phe Ala Thr Ser Ala
 100 105 110

Asp Asn His Val Thr Met Trp Val Asp Asp Gln Glu Val Ile Asn Lys
 115 120 125

Ala Ser Asn Ser Asn Lys Ile Arg Leu Glu Lys Gly Arg Leu Tyr Gln
 130 135 140

Ile Lys Ile Gln Tyr Gln Arg Glu Asn Pro Thr Glu Lys Gly Leu Asp
 145 150 155 160

Phe Lys Leu Tyr Trp Thr Asp Ser Gln Asn Lys Lys Glu Val Ile Ser
 165 170 175

Ser Asp Asn Leu Gln Leu Pro Glu Leu Lys Gln Lys Ser Ser Asn Ser
 180 185 190

Pro Gly Ser Gly Arg Ser Ala Ser Thr Ser Ala Gly Pro Thr Val Pro
 195 200 205

Asp Arg Asp Asn Asp Gly Ile Pro Asp Ser Leu Glu Val Glu Gly Tyr
 210 215 220

726173_ST25

Thr Val Asp Val Lys Asn Lys Ala Thr Phe Leu Ser Pro Trp Ile Ser
 225 230 235 240
 Asn Ile His Glu Lys Lys Gly Leu Thr Lys Tyr Lys Ser Ser Pro Glu
 245 250 255
 Lys Trp Ser Thr Ala Ser Asp Pro Tyr Ser Asp Phe Glu Lys Val Thr
 260 265 270
 Gly Arg Ile Asp Lys Asn Val Ser Pro Glu Ala Arg His Pro Leu Val
 275 280 285
 Ala Ala Tyr Pro Ile Val His Val Asp Met Glu Asn Ile Ile Leu Ser
 290 295 300
 Lys Asn Glu Asp Gln Ser Thr Gln Asn Thr Asp Ser Gln Thr Arg Thr
 305 310 315 320
 Ile Ser Lys Asn Thr Ser Thr Ser Arg Thr His Thr Ser Glu Val His
 325 330 335
 Gly Asn Ala Glu Val His Ala Ser Phe Phe Asp Ile Gly Gly Ser Val
 340 345 350
 Ser Ala Gly Phe Ser Asn Ser Asn Ser Ser Thr Val Ala Ile Asp His
 355 360 365
 Ser Leu Ser Leu Ala Gly Glu Arg Thr Trp Ala Glu Thr Met Gly Leu
 370 375 380
 Asn Thr Ala Asp Thr Ala Arg Leu Asn Ala Asn Ile Arg Tyr Val Asn
 385 390 395 400
 Thr Gly Thr Ala Pro Ile Tyr Asn Val Leu Pro Thr Thr Ser Leu Val
 405 410 415
 Leu Gly Lys Asn Gln Thr Leu Ala Thr Ile Lys Ala Lys Glu Asn Gln
 420 425 430
 Leu Ser Gln Ile Leu Ala Pro Asn Asn Tyr Tyr Pro Ser Lys Asn Leu
 435 440 445
 Ala Pro Ile Ala Leu Asn Ala Gln Asp Asp Phe Ser Ser Thr Pro Ile
 450 455 460
 Thr Met Asn Tyr Asn Gln Phe Leu Glu Leu Glu Lys Thr Lys Gln Leu
 465 470 475 480
 Arg Leu Asp Thr Asp Gln Val Tyr Gly Asn Ile Ala Thr Tyr Asn Phe
 485 490 495

726173_ST25

Glu Asn Gly Arg Val Arg Val Asp Thr Gly Ser Asn Trp Ser Glu Val
 500 505 510
 Leu Pro Gln Ile Gln Glu Thr Thr Ala Arg Ile Ile Phe Asn Gly Lys
 515 520 525
 Asp Leu Asn Leu Val Glu Arg Arg Ile Ala Ala Val Asn Pro Ser Asp
 530 535 540
 Pro Leu Glu Thr Thr Lys Pro Asp Met Thr Leu Lys Glu Ala Leu Lys
 545 550 555 560
 Ile Ala Phe Gly Phe Asn Glu Pro Asn Gly Asn Leu Gln Tyr Gln Gly
 565 570 575
 Lys Asp Ile Thr Glu Phe Asp Phe Asn Phe Asp Gln Gln Thr Ser Gln
 580 585 590
 Asn Ile Lys Asn Gln Leu Ala Glu Leu Asn Ala Thr Asn Ile Tyr Thr
 595 600 605
 Val Leu Asp Lys Ile Lys Leu Asn Ala Lys Met Asn Ile Leu Ile Arg
 610 615 620
 Asp Lys Arg Phe His Tyr Asp Arg Asn Asn Ile Ala Val Gly Ala Asp
 625 630 635 640
 Glu Ser Val Val Lys Glu Ala His Arg Glu Val Ile Asn Ser Ser Thr
 645 650 655
 Glu Gly Leu Leu Leu Asn Ile Asp Lys Asp Ile Arg Lys Ile Leu Ser
 660 665 670
 Gly Tyr Ile Val Glu Ile Glu Asp Thr Glu Gly Leu Lys Glu Val Ile
 675 680 685
 Asn Asp Arg Tyr Asp Met Leu Asn Ile Ser Ser Leu Arg Gln Asp Gly
 690 695 700
 Lys Thr Phe Ile Asp Phe Lys Lys Tyr Asn Asp Lys Leu Pro Leu Tyr
 705 710 715 720
 Ile Ser Asn Pro Asn Tyr Lys Val Asn Val Tyr Ala Val Thr Lys Glu
 725 730 735
 Asn Thr Ile Ile Asn Pro Ser Glu Asn Gly Asp Thr Ser Thr Asn Gly
 740 745 750
 Ile Lys Lys Ile Leu Ile Phe Ser Lys Lys Gly Tyr Glu Ile Gly
 755 760 765

726173_ST25

<210> 33
 <211> 2295
 <212> DNA
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 33
 atgaaaaaac gaaaagtgtt aataccatta atggcattgt ctacgatatt agtttcaagc 60
 acaggtaatt tagaggatgat tcaggcagaa gttaaacagg agaaccgggtt attaaatgaa 120
 tcagaatcaa gttcccagggtt gttactagga tactattttta gtgattttgaa ttttcaagca 180
 cccatgggtg ttacctcttc tactacagggtt gatttatctt ttcctagttc tgagtttagaa 240
 aatattccat cggaaaacca atattttcaa tctgctatct gggtcaggatt tatcaaagtt 300
 aagaagagtg atgaatatac atttgctact tccgctgata atcatgtaac aatgtgggta 360
 gatgaccaag aagtgattaa taaagcttct aattctaaca aaatcagatt agaaaaagga 420
 agattatatc aaataaaaaat tcaatatcaa cgagaaaatc ctactgaaaa aggattggat 480
 ttcaagttgt actggaccga ttctcaaaat aaaaaagaag tgattttctag tgataactta 540
 caattgccag aattaaaaca aaaatcttcg aactcaagaa aaaagcgaag tacaagtgtc 600
 ggacctacgg ttccagaccg tgacaatgat ggaatccctg attcattaga ggtagaagga 660
 tatacggttg atgtcaaaaa taaaagaact tttctttcac catggatttc taatattcat 720
 gaaaagaaag gattaaccaa atataaatca tctcctgaaa aatggagcac ggcttctgat 780
 ccgtacagtg atttcgaaaa gggtacagga cggattgata agaatgtatc accagaggca 840
 agacaccccc ttgtggcagc ttatccgatt gtacatgtag atatggagaa tattattctc 900
 tcaaaaaatg aggatcaatc cacacagaat actgatagtc aaacgagaac aataagtaaa 960
 aatacttcta caagtaggac acatactagt gaagtacatg gaaatgcaga agtgcattgcg 1020
 tcgttctttg atattgggtg gagtgtatct gcaggattta gtaattcgaa ttcaagtacg 1080
 gtcgcaattg atcattcact atctctagca ggggaaagaa cttgggctga aacaatgggt 1140
 ttaaataccg ctgatacagc aagattaaat gccaatatta gatattgtaa tactgggacg 1200
 gctccaatct acaacgtgtt accaacgact tcgttagtgt taggaaaaaa tcaaacactc 1260
 gcgacaatta aagctaagga aaaccaatta agtcaaatac ttgcacctaa taattattat 1320
 ccttctaaaa acttggcgcc aatcgcatc aatgcacaag acgatttcag ttctactcca 1380
 attacaatga attacaatca atttcttgag ttagaaaaaa cgaaacaatt aagattagat 1440
 acggatcaag tatatgggaa tatagcaaca tacaattttg aaaatggaag agtgaggggtg 1500
 gatacaggct cgaactggag tgaagtgtta ccgcaaattc aagaaacaac tgcacgtatc 1560
 atttttaatg gaaaagattt aaatctggta gaaaggcgga tagcggcggt taatcctagt 1620
 gatccattag aaacgactaa accggatatg acattaaaag aagcccttaa aatagcattt 1680
 ggatttaacg aaccgaatgg aaacttacia tatcaaggga aagacataac cgaatttgat 1740

726173_ST25

tttaatttcg atcaacaaac atctcaaaat atcaagaatc agttagcgga attaaacgca	1800
actaacatat atactgtatt agataaaatc aaattaaatg caaaaatgaa tattttaata	1860
agagataaac gttttcatta tgatagaaat aacatagcag ttggggcgga tgagtcagta	1920
gttaaggagg ctcatagaga agtaattaat tcgtcaacag agggattatt gttaaataatt	1980
gataaggata taagaaaaat attatcaggt tatattgtag aaattgaaga tactgaaggg	2040
cttaaagaag ttataaatga cagatatgat atgttgaata tttctagttt acggcaagat	2100
ggaaaaacat ttatagattt taaaaaatat aatgataaat taccgttata tataagtaat	2160
cccaattata aggtaaatgt atatgctgtt actaaagaaa acactattat taatcctagt	2220
gagaatgggg atactagtag caacgggatc aagaaaattt taatcttttc taaaaaaggc	2280
tatgagatag gataa	2295

<210> 34
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 34	
gcttgaattc atttcattat gatagaaata ac	32

<210> 35
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 35	
aattcaagct ttcctatctc atagcctttt tt	32

<210> 36
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 36	
taagtacgca atgggccacg	20

<210> 37
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 37	
aactcagcgg cagtctcaac	20

726173_ST25

<210> 38
 <211> 75
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 38
 tgaagatact gaaggcctta aagaagttat aaatgacaga tatgatatgt tgaatatctc 60
 gagtttacgg caaga 75

<210> 39
 <211> 75
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> synthetic

<400> 39
 tcttgccgta aactcgagat attcaacata tcatatctgt catttataac ttctttaagg 60
 ccttcagtat cttca 75