



US 20200093874A1

(19) **United States**

(12) **Patent Application Publication**
AGARWAL et al.

(10) **Pub. No.: US 2020/0093874 A1**

(43) **Pub. Date: Mar. 26, 2020**

(54) **COMPOSITIONS AND METHODS FOR THE TREATMENT OF NETHERTON SYNDROME**

(71) Applicant: **Krystal Biotech, Inc.**, Pittsburgh, PA (US)

(72) Inventors: **Pooja AGARWAL**, Mars, PA (US); **Suma KRISHNAN**, San Francisco, CA (US); **John C. FREEDMAN**, Pittsburgh, PA (US)

(73) Assignee: **Krystal Biotech, Inc.**, Pittsburgh, PA (US)

(21) Appl. No.: **16/581,150**

(22) Filed: **Sep. 24, 2019**

Related U.S. Application Data

(60) Provisional application No. 62/735,582, filed on Sep. 24, 2018.

Publication Classification

(51) **Int. Cl.**
A61K 35/763 (2006.01)
A61K 38/57 (2006.01)
(52) **U.S. Cl.**
CPC *A61K 35/763* (2013.01); *A61K 38/57* (2013.01)

(57) **ABSTRACT**

The present disclosure provides recombinant nucleic acids comprising one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide (e.g., a SPINK5 polypeptide); viruses comprising the recombinant nucleic acids; compositions and formulations comprising the recombinant nucleic acids and/or viruses; methods of their use (e.g., for the treatment of Netherton Syndrome); and articles of manufacture or kits thereof.

Specification includes a Sequence Listing.




FIG. 1A





FIG. 1B



→ ≡ heterologous promoter

▨ ≡ Serine Protease Inhibitor Kazal-type 5 (SPINK5) coding sequence

■ ≡ regulatory elements

FIG. 1C





FIG. 1D



↑ == heterologous promoter

▨ == Serine Protease Inhibitor Kazal-type 5 (SPINK5) coding sequence

▩ == regulatory elements

FIG. 1E





FIG. 1F



↑ = heterologous promoter

▨ = Serine Protease Inhibitor Kazal-type 5 (SPINK5) coding sequence

▣ = regulatory elements

FIG. 1G





FIG. 1H




↑ = heterologous promoter

▨ = Serine Protease Inhibitor Kazal-type 5 (SPINK5) coding sequence

▭ = regulatory elements

FIG. 11



↑ = heterologous promoter

▨ = Serine Protease Inhibitor Kazal-type 5 (SPINK5) coding sequence

▣ = regulatory elements

FIG. 2A

SPINK5 DNA Copies




FIG. 2B

SPINK5 Transcripts




FIG. 2C




FIG. 2D




FIG. 4




FIG. 3




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E





FIG. 5F



COMPOSITIONS AND METHODS FOR THE TREATMENT OF NETHERTON SYNDROME

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of U.S. Provisional Application Ser. No. 62/735,582, filed Sep. 24, 2018, which is incorporated herein by reference in its entirety.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

[0002] The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: 761342000400SEQLIST.txt, date recorded: Sep. 23, 2019, size: 60 KB).

FIELD OF THE INVENTION

[0003] The present disclosure relates, in part, to recombinant nucleic acids comprising one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide (e.g., a human SPINK5 polypeptide), viruses comprising the same, pharmaceutical compositions, formulations, and medicaments thereof, and methods of their use (e.g., for the treatment of Netherton Syndrome).

BACKGROUND

[0004] Netherton Syndrome (NS), also referred to as Comèl-Netherton Syndrome, is a debilitating autosomal recessive skin disorder that causes defective keratinization, severe skin barrier defects, and recurrent infections. Patients present shortly after birth with generalized rashes that develop into severe ichthyosis (Comèl, 1949. *Dermatologica*. 98(3):133-6). Infants with more severe NS symptoms are associated with failure to thrive, hypernatremic dehydration secondary to excess fluid loss, delayed growth, short stature, and recurrent infections (Jones et al., 1986. *Br J Dermatol*. 114(6):741-3; Hausser et al., 1996. *Pediatr Dermatol*. 13(3):183-99). Postnatal mortality rates for Netherton Syndrome are extremely high, with up to 20% of patients failing to reach their first birthday (Renner et al., 2009. *J Allergy Clin Immunol*. 124(3):536-43).

[0005] Clinically, Netherton Syndrome is characterized by congenital ichthyosiform erythroderma, hair shaft defects, recurrent infections, and a defective skin barrier (Judge et al., 1994. *Br J Dermatol*. 131(5):615-21; Schmuth et al., 2013. *Eur J Hum Genet*. 21(2): 123-33). Hair shafts are fragile and break easily due to trichorrhexis invaginata, or "bamboo hair", resulting in short sparse hair. A predisposition to allergies, asthma, and eczema is also characteristic of NS. Ultimately, those afflicted by Netherton Syndrome often experience chronic skin inflammation, severe dehydration, and stunted growth.

[0006] The disease arises due to mutations in the Serine Protease Inhibitor Kazal-type 5 (SPINK5) gene, resulting in loss of activity of its encoded serine protease inhibitor protein SPINK5 (also known as Lympho-Epithelial Kazal-type-related Inhibitor (LEKTI)) (Bitoun et al., 2002. *J Invest Dermatol*. 118(2):352-61). In healthy individuals, SPINK5 is one of the serine protease inhibitors expressed in the outermost layers of the skin, and it plays a critical role in the regulation of serine proteases which hydrolyze extracellular

proteins that hold corneocytes together. In patients suffering from Netherton Syndrome, the suppressive effects of SPINK5 on these serine proteases is abolished due to underlying genetic mutations in the SPINK5 gene (Komatsu et al., 2008. *J Invest Dermatol*. 128(5):1148-59). Consequently, hyperactivated serine proteases in the skin cause uncontrolled desquamation, leading to a defective skin barrier (Descargues et al., 2005. *Nat Genet*. 37(1):56-65).

[0007] Presently, there is no known cure for Netherton Syndrome, and treatment options for patients are limited. Current care focuses on managing the symptoms of the disease, including using moisturizing products to minimize scaling/cracking of the skin, and providing anti-infective treatments when appropriate. Additionally, intravenous administration of immunoglobulin to reduce skin infections has gained in popularity. Unfortunately, while steroid and retinoid products have shown some success in treating other ichthyosis-related disorders, these products have proven ineffective against NS, and may in fact make things worse for affected individuals (Braun et al., 1997. *Dermatology*. 195(1):75). Thus, there exists a clear need for novel treatment options targeting molecular correction of SPINK5 deficiencies observed in this sensitive patient population.

[0008] All references cited herein, including patent applications, patent publications, non-patent literature, and NCBI/UniProtKB/Swiss-Prot accession numbers are herein incorporated by reference in their entirety, as if each individual reference were specifically and individually indicated to be incorporated by reference.

BRIEF SUMMARY

[0009] In order to meet these and other needs, provided herein are recombinant nucleic acids (e.g., recombinant herpes simplex virus genomes) encoding Serine Protease Inhibitor Kazal-type (SPINK) polypeptides (e.g., SPINK5 polypeptides) for use in viruses (e.g., herpes viruses), pharmaceutical compositions and formulations, medicaments, and/or methods useful for remedying SPINK deficiencies and/or for treating an individual having, or at risk of developing, a SPINK-associated disorder (e.g., Netherton Syndrome, atopic dermatitis, hereditary pancreatitis, tropical calcific pancreatitis, spermatogenic failure 29, etc). The present inventors have shown that the recombinant viruses described herein were capable of transducing human epidermal cells, successfully expressing the encoded exogenous SPINK (RNA and protein), and that the exogenous polypeptide was appropriately secreted and was fully functional (see e.g., Example 2). Moreover, the present inventors have shown that the viruses described herein may be successfully administered either topically or intradermally in vivo without significant cytotoxicity, allowing for the encoded human SPINK polypeptide to be expressed in and localized to the appropriate region of the skin (see e.g., Example 3). Without wishing to be bound by theory, it is believed that increasing, augmenting, and/or supplementing the levels of SPINK polypeptide (e.g., human SPINK5) in a subject in need thereof by administering one or more of the recombinant nucleic acids, viruses, compositions, and/or medicaments described herein will: 1) enhance anti-inflammatory and/or anti-microbial protection of mucous epithelia; 2) reduce transepidermal water loss (TEWL); 3) inhibit desquamation; and 4) reduce or treat skin barrier defects in the subject. In addition, without wishing to be bound by theory, it is believed that increasing, augmenting, and/or

supplementing the levels of a SPINK polypeptide (e.g., human SPINK5) in one or more cells of a subject (by administering any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical composition described herein) will lead to the treatment of existing skin abnormalities in individuals suffering from a SPINK deficiency (e.g., Netherton Syndrome, atopic dermatitis, etc.), as well as will prevent or delay reformation of skin abnormalities in the treated areas.

[0010] Accordingly, certain aspects of the present disclosure relate to a recombinant herpes virus genome comprising one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide. In some embodiments, the recombinant herpes virus genome comprises two or more polynucleotides encoding a SPINK polypeptide. In some embodiments, the recombinant herpes virus genome is replication competent. In some embodiments, the recombinant herpes virus genome is replication defective. In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes virus genome is selected from a recombinant herpes simplex virus genome, a recombinant varicella zoster virus genome, a recombinant human cytomegalovirus genome, a recombinant herpesvirus 6A genome, a recombinant herpesvirus 6B genome, a recombinant herpesvirus 7 genome, a recombinant Kaposi's sarcoma-associated herpesvirus genome, and any combinations or derivatives thereof. In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes virus genome is a recombinant herpes simplex virus genome. In some embodiments, the recombinant herpes simplex virus genome is a recombinant type 1 herpes simplex virus (HSV-1) genome, a recombinant type 2 herpes simplex virus (HSV-2) genome, or any derivatives thereof. In some embodiments, the recombinant herpes simplex virus genome is a recombinant HSV-1 genome.

[0011] In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation. In some embodiments, the inactivating mutation is in a herpes simplex virus gene. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the herpes simplex virus gene. In some embodiments, the herpes simplex virus gene is selected from Infected Cell Protein (ICP) 0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27, ICP47, thymidine kinase (tk), Long Unique Region (UL) 41, and UL55. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in one or both copies of the ICP4 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP22 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL41 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in one or both copies of the ICP0 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP27 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL55 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the Joint region. In some embodiments, the recombinant herpes simplex virus genome comprises a deletion of the Joint region.

[0012] In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within one or more viral gene loci. In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes simplex virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within one or both of the ICP4 viral gene loci. In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes simplex virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within the ICP22 viral gene locus. In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes simplex virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within the UL41 viral gene locus.

[0013] In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide is a human SPINK polypeptide. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide is a Serine Protease Inhibitor Kazal-type 5 (SPINK5) polypeptide. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide is a human SPINK5 polypeptide. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to an amino acid sequence selected from SEQ ID NOS: 7-25. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to an amino acid sequence selected from SEQ ID NOS: 7-9. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 7. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 8. In some embodiments that may be combined with any of the preceding embodiments, the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 9.

[0014] In some embodiments that may be combined with any of the preceding embodiments, the recombinant herpes virus genome has reduced cytotoxicity when introduced into a target cell as compared to a corresponding wild-type herpes virus genome. In some embodiments, the target cell is a human cell. In some embodiments, the target cell is a cell

of the epidermis and/or dermis. In some embodiments, the target cell is a keratinocyte or fibroblast.

[0015] Other aspects of the present disclosure relate to a herpes virus comprising any of the recombinant herpes virus genomes described herein. In some embodiments, the herpes virus is replication competent. In some embodiments, the herpes virus is replication defective. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus has reduced cytotoxicity as compared to a corresponding wild-type herpes virus. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus is selected from a herpes simplex virus, a varicella zoster virus, a human cytomegalovirus, a herpesvirus 6A, a herpesvirus 6B, a herpesvirus 7, a Kaposi's sarcoma-associated herpesvirus, and any combinations or derivatives thereof. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus is a herpes simplex virus. In some embodiments, the herpes simplex virus is an HSV-1, an HSV-2, or any derivatives thereof. In some embodiments, the herpes simplex virus is an HSV-1.

[0016] Other aspects of the present disclosure relate to a pharmaceutical composition comprising any of the recombinant herpes virus genomes described herein and/or any of the herpes viruses described herein and a pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition is suitable for topical, transdermal, subcutaneous, intradermal, oral, sublingual, buccal, rectal, vaginal, inhaled, intravenous, intraarterial, intramuscular, intracardiac, intraosseous, intraperitoneal, transmucosal, intravitreal, subretinal, intraarticular, peri-articular, local, epicutaneous administration, or any combinations thereof. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition is suitable for topical, transdermal, subcutaneous, intradermal, and/or transmucosal administration. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition is suitable for topical, transdermal, and/or intradermal administration. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition is suitable for topical and/or intradermal administration. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition is suitable for topical administration. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition comprises a methylcellulose gel (e.g., a carboxy methylcellulose gel, a hydroxypropyl methylcellulose gel, etc.). In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition comprises a phosphate buffer. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition comprises glycerol. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition comprises a lipid carrier. In some embodiments that may be combined with any of the preceding embodiments, the pharmaceutical composition comprises a nanoparticle carrier.

[0017] Other aspects of the present disclosure relate to the use of any of the recombinant nucleic acids, herpes viruses, and/or pharmaceutical compositions described herein as a medicament.

[0018] Other aspects of the present disclosure relate to the use of any of the recombinant nucleic acids, herpes viruses, and/or pharmaceutical compositions described herein in a therapy.

[0019] Other aspects of the present disclosure relate to the use of any of the recombinant nucleic acids, herpes viruses, and/or pharmaceutical compositions described herein in the production or manufacture of a medicament for treating one or more signs or symptoms of SPINK polypeptide deficiency (e.g., Netherton Syndrome, atopic dermatitis, etc.).

[0020] Other aspects of the present disclosure relate to a method of enhancing, increasing, augmenting, and/or supplementing the levels of a SPINK polypeptide in one or more cells of a subject comprising administering to the subject an effective amount of any of the herpes viruses described herein and/or any of the pharmaceutical compositions described herein. In some embodiments, the SPINK polypeptide is a SPINK5 polypeptide (e.g., a human SPINK5 polypeptide). In some embodiments that may be combined with any of the preceding embodiments, the subject is a human. In some embodiments that may be combined with any of the preceding embodiments, the subject's genome comprises a loss-of-function mutation in a SPINK5 gene. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically to the subject. In some embodiments that may be combined with any of the preceding embodiments, the skin of the subject is abraded prior to administration.

[0021] Other aspects of the present disclosure relate to a method of enhancing, increasing, augmenting, and/or supplementing anti-inflammatory and/or anti-microbial protection of mucous epithelia in a subject in need thereof comprising administering to the subject an effective amount of any of the herpes viruses described herein and/or any of the pharmaceutical compositions described herein. In some embodiments, the subject is a human. In some embodiments that may be combined with any of the preceding embodiments, the subject's genome comprises a loss-of-function mutation in a SPINK5 gene. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, sub-

cutaneously, intradermally, or transmucosally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically to the subject. In some embodiments that may be combined with any of the preceding embodiments, the skin of the subject is abraded prior to administration.

[0022] Other aspects of the present disclosure relate to a method of repressing desquamation in a subject in need thereof comprising administering to the subject an effective amount of any of the herpes viruses described herein and/or any of the pharmaceutical compositions described herein. In some embodiments, the subject is a human. In some embodiments that may be combined with any of the preceding embodiments, the subject's genome comprises a loss-of-function mutation in a SPINK5 gene. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically to the subject. In some embodiments that may be combined with any of the preceding embodiments, the skin of the subject is abraded prior to administration.

[0023] Other aspects of the present disclosure relate to a method of reducing or treating a skin barrier defect in a subject in need thereof comprising administering to the subject an effective amount of any of the herpes viruses described herein and/or any of the pharmaceutical compositions described herein. In some embodiments, the skin barrier defect is transepithelial water loss (TEWL). In some embodiments that may be combined with any of the preceding embodiments, the subject is a human. In some embodiments that may be combined with any of the preceding embodiments, the subject's genome comprises a loss-of-function mutation in a SPINK5 gene. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically to the subject. In some embodiments that may be combined with any of the preceding embodiments, the skin of the subject is abraded prior to administration.

[0024] Other aspects of the present disclosure relate to a method of providing prophylactic, palliative, or therapeutic relief of one or more signs or symptoms of Netherton Syndrome (NS) in a subject in need thereof comprising administering to the subject an effective amount of any of the herpes viruses described herein and/or any of the pharmaceutical compositions described herein. In some embodiments, the one or more signs or symptoms of NS are selected from defective keratinization, a defective skin barrier, chronic skin inflammation, universal pruritus, severe dehydration, stunted growth, trichorrhexis invaginata and/or trichorrhexis nodosa, leaking fluid from the skin, development of ring-like lesions on the skin, eczema, increased susceptibility to infection, recurrent skin infections, increased susceptibility to allergy, development of scaly/reddish skin, development of ichthyosis linearis circumflexa and/or ichthyosiform erythroderma, altered immunoglobulin levels, immature natural killer cells having reduced lytic function, difficulty regulating body temperature, and any combinations thereof. In some embodiments that may be combined with any of the preceding embodiments, the one or more sign or symptoms of NS are selected from defective keratinization, a defective skin barrier, recurrent skin infections, congenital ichthyosiform erythroderma, ichthyosis linearis circumflexa, trichorrhexis invaginata, chronic skin inflammation, and any combinations thereof. In some embodiments that may be combined with any of the preceding embodiments, the subject is a human. In some embodiments that may be combined with any of the preceding embodiments, the subject's genome comprises a loss-of-function mutation in a SPINK5 gene. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically or intradermally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered intradermally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically to the subject. In some embodiments that may be combined with any of the preceding embodiments, the skin of the subject is abraded prior to administration.

[0025] Other aspects of the present disclosure relate to a method of providing prophylactic, palliative, or therapeutic relief of one or more signs or symptoms of atopic dermatitis in a subject in need thereof comprising administering to the

subject an effective amount of any of the herpes viruses described herein and/or any of the pharmaceutical compositions described herein. In some embodiments, the one or more signs or symptoms of atopic dermatitis are selected from itchy skin, dry skin, red to brownish-grey patches on the skin, small raised bumps on the skin, thickened skin, cracked skin, scaly skin, swollen skin, weeping sores, skin infections, eyelid dermatitis, cataracts, and any combinations thereof. In some embodiments that may be combined with any of the preceding embodiments, the subject is a human. In some embodiments that may be combined with any of the preceding embodiments, the subject's genome comprises a loss-of-function mutation in a SPINK5 gene. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically or intradermally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered intradermally to the subject. In some embodiments that may be combined with any of the preceding embodiments, the herpes virus and/or pharmaceutical composition is administered topically to the subject. In some embodiments that may be combined with any of the preceding embodiments, the skin of the subject is abraded prior to administration.

[0026] Other aspects of the present disclosure relate to an article of manufacture or kit comprising any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein and instructions for administration thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0027] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

[0028] FIGS. 1A-I show schematics of wild-type and modified herpes simplex virus genomes. FIG. 1A shows a wild-type herpes simplex virus genome. FIG. 1B shows a modified herpes simplex virus genome comprising deletions of the coding sequence of ICP4 (both copies), with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at each of the ICP4 loci. FIG. 1C shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies) and ICP22, with an expression cassette containing a nucleic acid encoding a human SPINK5 poly-

peptide integrated at each of the ICP4 loci. FIG. 1D shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies) and ICP22, with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at the ICP22 locus. FIG. 1E shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies) and UL41, with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at each of the ICP4 loci. FIG. 1F shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies) and UL41, with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at the UL41 locus. FIG. 1G shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies), ICP22, and UL41, with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at each of the ICP4 loci. FIG. 1H shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies), ICP22, and UL41, with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at the UL41 locus. FIG. 1I shows a modified herpes simplex virus genome comprising deletions of the coding sequences of ICP4 (both copies), ICP22, and UL41, with an expression cassette containing a nucleic acid encoding a human SPINK5 polypeptide integrated at the ICP22 locus.

[0029] FIGS. 2A-D show human SPINK5 nucleic acid and protein analyses in immortalized normal keratinocytes infected with HSV-S5. FIG. 2A shows the levels of codon-optimized human SPINK5 DNA present in immortalized normal keratinocytes 48 hours after infection with HSV-S5 at the indicated MOIs, as determined by qPCR analysis. FIG. 2B shows the levels of codon-optimized human SPINK5 transcripts present in immortalized normal keratinocytes 48 hours after infection with HSV-S5 at the indicated MOIs, as determined by qRT-PCR analysis. FIG. 2C shows western blot analysis of human SPINK5 protein expression in immortalized normal keratinocytes 48 hours after infection with HSV-S5 at the indicated MOIs. GAPDH was used as a loading control. FIG. 2D shows representative immunofluorescence images of human SPINK5 protein expression in immortalized normal keratinocytes 48 hours after infection with HSV-S5 at the indicated MOIs. DAPI staining was used to visualize nuclei. For all experiments, uninfected (mock) cells were used as a negative control.

[0030] FIG. 3 shows the concentration of human SPINK5 secreted into the supernatant of cultured immortalized normal keratinocytes 48 hours after infection with HSV-S5 at the indicated MOIs, as assessed by ELISA. Cell supernatant collected from uninfected (mock) immortalized normal keratinocytes was used as a negative control.

[0031] FIG. 4 shows the percent inhibition in proteolytic activity of recombinant human Kallikrein 5 (KLK5) after incubation with cell culture supernatants collected from immortalized normal keratinocytes infected with HSV-S5 at the indicated MOIs, as assessed in a fluorometric assay employing a synthetic, non-natural peptide substrate of human KLK5. Cell supernatant collected from uninfected (mock) immortalized normal keratinocytes was used as a negative control. Data is presented as the average \pm SEM.

[0032] FIGS. 5A-F show nucleic acid and protein analyses, as well as histology, of skin biopsies harvested from control- or HSV-S5-treated BALB/c mice. FIG. 5A shows the levels of human SPINK5 DNA present in the skin of immunocompetent animals harvested 48 hours after topical application of HSV-S5 or vehicle control, as assessed by qPCR analysis. FIG. 5B shows the levels of human SPINK5 transcripts present in the skin of immunocompetent animals harvested 48 hours after topical application of HSV-S5 or vehicle control, as assessed by qRT-PCR analysis. FIG. 5C shows the levels of human SPINK5 DNA present in the skin of immunocompetent animals harvested 48 hours after intradermal injection of HSV-S5 or vehicle control, as assessed by qPCR analysis. FIG. 5D shows the levels of human SPINK5 transcripts present in the skin of immunocompetent animals harvested 48 hours after intradermal injection of HSV-S5 or vehicle control, as assessed by qRT-PCR analysis. For qPCR and qRT-PCR analyses, vehicle control data is presented as the average of two tissue samples (two replicates/tissue sample) \pm SEM, and HSV-S5 data is presented as the average of six tissue samples (two replicates/tissue sample) \pm SEM. FIG. 5E shows representative immunofluorescence images of human SPINK5 and mouse filaggrin protein expression in skin biopsies harvested from BALB/c mice 48 hours after topical application of HSV-S5 or vehicle control. DAPI staining was used to visualize nuclei. FIG. 5F shows representative histology of mouse skin biopsies harvested from BALB/c mice 48 hours after topical application of HSV-S5 or vehicle control.

DETAILED DESCRIPTION

[0033] In some embodiments, the present disclosure relates to recombinant nucleic acids (e.g., recombinant herpes viral genomes) comprising one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide (e.g., a human SPINK5 polypeptide), and/or use of these recombinant nucleic acids in viruses (e.g., herpes viruses), compositions, formulations, medicaments, and/or methods in order to supplement or treat SPINK gene deficiencies (e.g., in a subject whose genome naturally harbors a loss-of-function and/or pathogenic variant of a SPINK gene), and/or provide medical intervention to a subject in need thereof (e.g., to provide prophylactic, palliative, and/or therapeutic relief to one or more diseases or disorders arising from a SPINK gene deficiency (e.g., Netherton Syndrome, atopic dermatitis, etc.). Without wishing to be bound by theory, it is believed that the recombinant nucleic acids, viruses, compositions, formulations, medicaments, and/or methods described herein will help to treat the existing skin abnormalities in individuals suffering from Netherton Syndrome and/or atopic dermatitis, as well as prevent or delay reformation of skin abnormalities in treated subjects.

[0034] The following description sets forth exemplary methods, parameters, and the like. It should be recognized, however, that such description is not intended as a limitation on the scope of the present disclosure but is instead provided as a description of exemplary embodiments.

I. GENERAL TECHNIQUES

[0035] The techniques and procedures described or referenced herein are generally well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized meth-

odologies described in Sambrook et al., 2001. *Molecular Cloning: A Laboratory Manual* 3d edition Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; *Current Protocols in Molecular Biology* (F. M. Ausubel, et al. eds., (2003)); the series *Methods in Enzymology* (Academic Press, Inc.); *PCR 2: A Practical Approach* (M. J. MacPherson, B. D. Hames and G. R. Taylor eds. (1995)), Harlow and Lane, eds. (1988); *Oligonucleotide Synthesis* (M. J. Gait, ed., 1984); *Methods in Molecular Biology*, Humana Press; *Cell Biology: A Laboratory Notebook* (J. E. Cellis, ed., 1998) Academic Press; *Animal Cell Culture* (R. I. Freshney, ed., 1987); *Introduction to Cell and Tissue Culture* (J. P. Mather and P. E. Roberts, 1998) Plenum Press; *Cell and Tissue Culture: Laboratory Procedures* (A. Doyle, J. B. Griffiths, and D. G. Newell, eds., 1993-8) J. Wiley and Sons; *Gene Transfer Vectors for Mammalian Cells* (J. M. Miller and M. P. Calos, eds., 1987); *PCR: The Polymerase Chain Reaction*, (Mullis et al., eds., 1994); *Short Protocols in Molecular Biology* (Wiley and Sons, 1999).

II. DEFINITIONS

[0036] Before describing the present disclosure in detail, it is to be understood that the present disclosure is not limited to particular compositions or biological systems, which can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

[0037] As used herein, the singular forms “a”, “an” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a molecule” optionally includes a combination of two or more such molecules, and the like.

[0038] As used herein, the term “and/or” may include any and all combinations of one or more of the associated listed items. For example, the term “a and/or b” may refer to “a alone”, “b alone”, “a or b”, or “a and b”; the term “a, b, and/or c” may refer to “a alone”, “b alone”, “c alone”, “a or b”, “a or c”, “b or c”, “a, b, or c”, “a and b”, “a and c”, or “a, b, and c”; etc.

[0039] As used herein, the term “about” refers to the usual error range for the respective value readily known to the skilled person in this technical field. Reference to “about” a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter per se.

[0040] It is understood that aspects and embodiments of the present disclosure include “comprising”, “consisting”, and “consisting essentially of” aspects and embodiments

[0041] As used herein, the terms “polynucleotide”, “nucleic acid sequence”, “nucleic acid”, and variations thereof shall be generic to polydeoxyribonucleotides (containing 2-deoxy-D-ribose), to polyribonucleotides (containing D-ribose), to any other type of polynucleotide that is an N-glycoside of a purine or pyrimidine base, and to other polymers containing non-nucleotidic backbones, provided that the polymers contain nucleobases in a configuration that allows for base pairing and base stacking, as found in DNA and RNA. Thus, these terms include known types of nucleic acid sequence modifications, for example, substitution of one or more of the naturally occurring nucleotides with an analog, and inter-nucleotide modifications.

[0042] As used herein, a nucleic acid is “operatively linked” or “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, a promoter or enhancer is operably linked to

a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operatively linked” or “operably linked” means that the DNA sequences being linked are contiguous.

[0043] As used herein, the term “vector” refers to discrete elements that are used to introduce heterologous nucleic acids into cells for either expression or replication thereof. An expression vector includes vectors capable of expressing nucleic acids that are operatively linked with regulatory sequences, such as promoter regions, that are capable of effecting expression of such nucleic acids. Thus, an expression vector may refer to a DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector that, upon introduction into an appropriate host cell, results in expression of the nucleic acids. Appropriate expression vectors are well known to those of skill in the art and include those that are replicable in eukaryotic cells and those that remain episomal or those which integrate into the host cell genome.

[0044] As used herein, an “open reading frame” or “ORF” refers to a contiguous stretch of nucleic acids, either DNA or RNA, that encode a protein or polypeptide. Typically, the nucleic acid comprises a translation start signal or initiation codon, such as ATG or AUG, and a termination codon.

[0045] As used herein, an “untranslated region” or “UTR” refers to untranslated nucleic acids at the 5' and/or 3' ends of an open reading frame. The inclusion of one or more UTRs in a polynucleotide may affect post-transcriptional regulation, mRNA stability, and/or translation of the polynucleotide.

[0046] As used herein, the term “transgene” refers to a polynucleotide that is capable of being transcribed into RNA and translated and/or expressed under appropriate conditions, after being introduced into a cell. In some embodiments, it confers a desired property to a cell into which it was introduced, or otherwise leads to a desired therapeutic or diagnostic outcome.

[0047] As used herein, the terms “polypeptide,” “protein,” and “peptide” are used interchangeably and may refer to a polymer of two or more amino acids.

[0048] As used herein, a “subject”, “host”, or an “individual” refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, as well as animals used in research, such as mice, rats, hamsters, rabbits, and non-human primates, etc. In some embodiments, the mammal is human.

[0049] As used herein, the terms “pharmaceutical formulation” or “pharmaceutical composition” refer to a preparation which is in such a form as to permit the biological activity of the active ingredient(s) to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the composition or formulation would be administered. “Pharmaceutically acceptable” excipients (e.g., vehicles, additives) are those which can reasonably be administered to a subject mammal to provide an effective dose of the active ingredient(s) employed.

[0050] As used herein, an “effective amount” is at least the minimum amount required to affect a measurable improvement or prevention of one or more symptoms of a particular disorder. An “effective amount” may vary according to factors such as the disease state, age, sex, and weight of the patient. An effective amount is also one in which any toxic

or detrimental effects of the treatment are outweighed by the therapeutically beneficial effects. For prophylactic use, beneficial or desired results include results such as eliminating or reducing the risk, lessening the severity, or delaying the onset of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. For therapeutic use, beneficial or desired results include clinical results such as decreasing one or more symptoms resulting from the disease, increasing the quality of life of those suffering from the disease, decreasing the dose of other medications used to treat symptoms of the disease, delaying the progression of the disease, and/or prolonging survival. An effective amount can be administered in one or more administrations. For purposes of the present disclosure, an effective amount of a recombinant nucleic acid, virus, and/or pharmaceutical composition is an amount sufficient to accomplish prophylactic or therapeutic treatment either directly or indirectly. As is understood in the clinical context, an effective amount of a recombinant nucleic acid, virus, and/or pharmaceutical composition may or may not be achieved in conjunction with another drug, compound, or pharmaceutical composition. Thus, an “effective amount” may be considered in the context of administering one or more therapeutic agents, and a single agent may be considered to be given in an effective amount if, in conjunction with one or more other agents, a desirable result may be or is achieved.

[0051] As used herein, “treatment” refers to clinical intervention designed to alter the natural course of the individual or cell being treated during the course of clinical pathology. Desirable effects of treatment include decreasing the rate of disease/disorder/defect progression, ameliorating or palliating the disease/disorder/defect state, and remission or improved prognosis. For example, an individual is successfully “treated” if one or more symptoms associated with Netherton Syndrome and/or atopic dermatitis are mitigated or eliminated.

[0052] As used herein, the term “delaying progression of” a disease/disorder/defect refers to deferring, hindering, slowing, retarding, stabilizing, and/or postponing development of the disease/disorder/defect. This delay can be of varying lengths or time, depending on the history of the disease/disorder/defect and/or the individual being treated. As is evident to one of ordinary skill in the art, a sufficient or significant delay can, in effect, encompass prevention, in that the individual does not develop the disease.

III. RECOMBINANT NUCLEIC ACIDS

[0053] Certain aspects of the present disclosure relate to recombinant nucleic acids (e.g., isolated recombinant nucleic acids) comprising one or more (e.g., one or more, two or more, three or more, four or more, five or more, ten or more, etc.) polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide. In some embodiments, the recombinant nucleic acid comprises one polynucleotide encoding a SPINK polypeptide. In some embodiments, the recombinant nucleic acid comprises two polynucleotides encoding a SPINK polypeptide. In some embodiments, the SPINK polypeptide is a human SPINK polypeptide. In some embodiments, the SPINK polypeptide is a Serine Protease Inhibitor Kazal-type 5 (SPINK5) polypeptide. In some embodiments, the SPINK5 polypeptide is a human SPINK5 polypeptide.

[0054] In some embodiments, the recombinant nucleic acid is a vector. In some embodiments, the recombinant nucleic acid is a viral vector. In some embodiments, the recombinant nucleic acid is a herpes viral vector. In some embodiments, the recombinant nucleic acid is a herpes simplex virus amplicon. In some embodiments, the recombinant nucleic acid is a recombinant herpes virus genome. In some embodiments, the recombinant nucleic acid is a recombinant herpes simplex virus genome. In some embodiments, the recombinant nucleic acid is a recombinant type 1 herpes simplex virus (HSV-1) genome.

[0055] Polynucleotides Encoding SPINK Polypeptides

[0056] In some embodiments, the present disclosure relates to a recombinant nucleic acid comprising one or more polynucleotides comprising the coding sequence of a SPINK gene, or portions thereof (e.g., the coding sequence corresponding to one or more Kazal-type domains resulting from proteolytic cleavage (e.g., via furin cleavage) of a full-length SPINK polypeptide). Any suitable SPINK gene (including any isoform thereof) known in the art may be encoded by a polynucleotide of the present disclosure, including, for example, a SPINK1 gene (e.g., a human SPINK1 gene (see e.g., NCBI Gene ID: 6690)), a SPINK2 gene (e.g., a human SPINK2 gene (see e.g., NCBI Gene ID: 6691)), a SPINK4 gene (e.g., a human SPINK4 gene (see e.g., NCBI Gene ID: 27920)), a SPINK5 gene (e.g., a human SPINK5 gene (see e.g., NCBI Gene ID: 11005)), a SPINK6 gene (e.g., a human SPINK6 gene (see e.g., NCBI Gene ID: 404203)), a SPINK7 gene (e.g., a human SPINK7 gene (see e.g., NCBI Gene ID: 84651)), a SPINK8 gene (e.g., a human SPINK8 gene (see e.g., NCBI Gene ID: 646424)), a SPINK9 gene (e.g., a human SPINK9 gene (see e.g., NCBI Gene ID: 643394)), a SPINK13 gene (e.g., a human SPINK13 gene (see e.g., NCBI Gene ID: 153218)), a SPINK14 gene (e.g., a human SPINK14 gene (see e.g., NCBI Gene ID: 408187)), etc. In some embodiments, a polynucleotide of the present disclosure comprises a sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the sequence of any SPINK gene (and/or coding sequences thereof) described herein or known in the art. Methods of identifying SPINK gene homologs/orthologs from additional species are known to one of ordinary skill in the art, including, for example, using a nucleic acid sequence alignment program such as the BLAST® blastn suite. In some embodiments, one or more polynucleotides of the present disclosure comprises the coding sequence of a human SPINK gene.

[0057] In some embodiments, a polynucleotide of the present disclosure comprises a codon-optimized variant of any SPINK gene described herein or known in the art. In some embodiments, use of a codon-optimized variant of a SPINK gene increases stability and/or yield of heterologous expression (RNA and/or protein) of the encoded SPINK polypeptide in a target cell (e.g., a target human cell such as a human keratinocyte or fibroblast), as compared to the stability and/or yield of heterologous expression of a corresponding non-codon-optimized, wild-type sequence. Any suitable method known in the art for performing codon optimization of a sequence for expression in one or more target cells (e.g., one or more human cells) may be used,

including, for example, by the methods described by Fath et al. (PLoS One. 2011 Mar. 3; 6(3): e17596).

[0058] In some embodiments, the present disclosure relates to a recombinant nucleic acid comprising one or more polynucleotides comprising the coding sequence of a SPINK5 gene (e.g., a human SPINK5 gene), or portions thereof (e.g., the coding sequence corresponding to one or more Kazal-type domains resulting from proteolytic cleavage (e.g., via furin cleavage) of a full-length SPINK5 polypeptide). Any suitable SPINK5 gene (including any isoform thereof) known in the art may be encoded by a polynucleotide of the present disclosure, including, for example, a human SPINK5 gene (see e.g., NCBI Gene ID: 11005; SEQ ID NOS: 1, 3, or 5), a chimpanzee SPINK5 gene (see e.g., NCBI Gene ID: 462173), a mouse SPINK5 gene (see e.g., NCBI Gene ID: 72432), a rat SPINK5 gene (see e.g., NCBI Gene ID: 361319), a dog SPINK5 gene (see e.g., NCBI Gene ID: 478055), a cow SPINK5 gene (see e.g., NCBI Gene ID: 526637), a horse SPINK5 gene (see e.g., NCBI Gene ID: 100071873), a pig SPINK5 gene (see e.g., NCBI Gene ID: 100512160), etc. In some embodiments, a polynucleotide of the present disclosure comprises a sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the sequence of any SPINK5 gene (and/or the coding sequences thereof) described herein or known in the art. Methods of identifying SPINK5 gene homologs/orthologs from additional species are known to one of ordinary skill in the art, including, for example, using a nucleic acid sequence alignment program such as the BLAST® blastn suite.

[0059] In some embodiments, a polynucleotide of the present disclosure comprises a codon-optimized variant of any SPINK5 gene described herein or known in the art (see e.g., SEQ ID NOS: 2, 4, 6, or 26). In some embodiments, use of a codon-optimized variant of a SPINK5 gene (e.g., a codon-optimized variant of a human SPINK5 gene) increases stability and/or yield of heterologous expression (RNA and/or protein) of the encoded SPINK5 polypeptide in a target cell (e.g., a target human cell such as a human keratinocyte or fibroblast), as compared to the stability and/or yield of heterologous expression of a corresponding non-codon-optimized, wild-type sequence. Any suitable method known in the art for performing codon optimization of a sequence for expression in one or more target cells (e.g., one or more human cells) may be used, including, for example, by the methods described by Fath et al. (PLoS One. 2011 Mar. 3; 6(3): e17596).

[0060] In some embodiments, one or more polynucleotides of the present disclosure comprises the coding sequence of a human SPINK5 gene (or a codon-optimized variant thereof), or portions thereof (e.g., the coding sequence corresponding to one or more Kazal-type domains resulting from proteolytic cleavage (e.g., via human furin cleavage) of a full-length human SPINK5 polypeptide). In some embodiments, a polynucleotide of the present disclosure comprises a sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a sequence selected from SEQ ID NOS: 1-6 and 26. In some embodi-

or a fragment of the sequence of SEQ ID NO: 5, SEQ ID NO: 6, or SEQ ID NO: 26. In some embodiments, the 5' truncation, 3' truncation, or fragment of the sequence of SEQ ID NO: 5, SEQ ID NO: 6, or SEQ ID NO: 26 is a polynucleotide that has at least 25, at least 50, at least 75, at least 100, at least 125, at least 150, at least 175, at least 200, at least 250, at least 300, or at least 350, at least 400, at least 450, at least 500, at least 750, at least 1000, at least 1250, at least 1500, at least 1750, at least 2000, at least 2250, at least 2500, at least 2750, at least 3000, but fewer than 3195 consecutive nucleotides of SEQ ID NO: 5, SEQ ID NO: 6, or SEQ ID NO: 26. In some embodiments, a polynucleotide of the present disclosure comprises a sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the sequence of nucleic acids 1-3192 of SEQ ID NO: 5, SEQ ID NO: 6, or SEQ ID NO: 26. In some embodiments, a polynucleotide of the present disclosure comprises the sequence of nucleic acids 1-3192 of SEQ ID NO: 5, SEQ ID NO: 6, or SEQ ID NO: 26.

[0082] A polynucleotide of the present disclosure encoding a SPINK polypeptide (e.g., a human SPINK5 polypeptide) may further encode additional coding and non-coding sequences. Examples of additional coding and non-coding sequences may include, but are not limited to, sequences encoding additional polypeptide tags (e.g., encoded in-frame with the SPINK protein in order to produce a fusion protein), introns (e.g., native, modified, or heterologous introns), 5' and/or 3' UTRs (e.g., native, modified, or heterologous 5' and/or 3' UTRs), and the like. Examples of suitable polypeptide tags may include, but are not limited, to any combination of purification tags, such as his-tags, flag-tags, maltose binding protein and glutathione-S-transferase tags, detection tags, such as tags that may be detected photometrically (e.g., green fluorescent protein, red fluorescent protein, etc.) and tags that have a detectable enzymatic activity (e.g., alkaline phosphatase, etc.), tags containing secretory sequences, signal sequences, leader sequences, and/or stabilizing sequences, protease cleavage sites (e.g., furin cleavage sites, TEV cleavage sites, Thrombin cleavage sites, etc.), and the like. In some embodiments, the 5' and/or 3'UTRs increase the stability, localization, and/or translational efficiency of the polynucleotides. In some embodiments, the 5' and/or 3'UTRs improve the level and/or duration of protein expression. In some embodiments, the 5' and/or 3'UTRs include elements (e.g., one or more miRNA binding sites, etc.) that may block or reduce off-target expression (e.g., inhibiting expression in specific cell types (e.g., neuronal cells), at specific times in the cell cycle, at specific developmental stages, etc.). In some embodiments, the 5' and/or 3'UTRs include elements (e.g., one or more miRNA binding sites, etc.) that may enhance effector protein expression in specific cell types (such as human keratinocytes and/or fibroblasts).

[0083] In some embodiments, a polynucleotide of the present disclosure is operably linked to one or more (e.g., one or more, two or more, three or more, four or more, five or more, ten or more, etc.) regulatory sequences. The term "regulatory sequence" may include enhancers, insulators, promoters, and other expression control elements (e.g., polyadenylation signals). Any suitable enhancer(s) known in the art may be used, including, for example, enhancer

sequences from mammalian genes (such as globin, elastase, albumin, α -fetoprotein, insulin and the like), enhancer sequences from a eukaryotic cell virus (such as SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, adenovirus enhancers, and the like), and any combinations thereof. Any suitable insulator(s) known in the art may be used, including, for example, herpes simplex virus (HSV) chromatin boundary (CTRL/CTCF-binding/insulator) elements CTRL1 and/or CTRL2, chicken hypersensitive site 4 insulator (cHS4), human HNRPA2B1-CBX3 ubiquitous chromatin opening element (UCOE), the scaffold/matrix attachment region (S/MAR) from the human interferon beta gene (IFNB1), and any combinations thereof. Any suitable promoter (e.g., suitable for transcription in mammalian host cells) known in the art may be used, including, for example, promoters obtained from the genomes of viruses (such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus, Simian Virus 40 (SV40), and the like), promoters from heterologous mammalian genes (such as the actin promoter (e.g., the β -actin promoter), a ubiquitin promoter (e.g., a ubiquitin C (UbC) promoter), a phosphoglycerate kinase (PGK) promoter, an immunoglobulin promoter, from heat-shock protein promoters, and the like), promoters from native and/or homologous mammalian genes (e.g., human SPINK gene promoters), synthetic promoters (such as the CAGG promoter), and any combinations thereof, provided such promoters are compatible with the host cells. Regulatory sequences may include those which direct constitutive expression of a nucleic acid, as well as tissue-specific regulatory and/or inducible or repressible sequences.

[0084] In some embodiments, a polynucleotide of the present disclosure is operably linked to one or more heterologous promoters. In some embodiments, the one or more heterologous promoters are one or more of constitutive promoters, tissue-specific promoters, temporal promoters, spatial promoters, inducible promoters and repressible promoters. In some embodiments, the one or more heterologous promoters are one or more of the human cytomegalovirus (HCMV) immediate early promoter, the human elongation factor-1 (EF1) promoter, the human β -actin promoter, the human UbC promoter, the human PGK promoter, the synthetic CAGG promoter, and any combinations thereof. In some embodiments, a polynucleotide of the present disclosure is operably linked to an HCMV promoter.

[0085] In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a Collagen alpha-1 (VII) chain polypeptide (COLT). In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a Lysyl hydroxylase 3 polypeptide (LH3). In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a Keratin type I cytoskeletal 17 polypeptide (KRT17). In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a transglutaminase (TGM) polypeptide (e.g., a human transglutaminase poly-

peptide such as a human TGM1 polypeptide). In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a cosmetic protein (e.g., collagen proteins, fibronectins, elastins, lumicans, vitronectins/vitronectin receptors, laminins, neuromodulators, fibrillins, additional dermal extracellular matrix proteins, etc.). In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) an antibody (e.g., a full-length antibody, an antibody fragments, etc.). In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a Collagen alpha-1 (VII) chain polypeptide, a Lysyl hydroxylase 3 polypeptide, a Keratin type I cytoskeletal 17 polypeptide, and/or any chimeric polypeptides thereof. In some embodiments, a recombinant nucleic acid of the present disclosure does not comprise a polynucleotide comprising the coding sequence of (e.g., a transgene encoding) a Collagen alpha-1 (VII) chain polypeptide, a Lysyl hydroxylase 3 polypeptide, a Keratin type I cytoskeletal 17 polypeptide, a transglutaminase (TGM) polypeptide, a cosmetic protein, an antibody, and/or any chimeric polypeptides thereof.

[0086] SPINK Polypeptides

[0087] In some embodiments, the present disclosure relates to one or more polynucleotides encoding a SPINK polypeptide, or any portions thereof (e.g., the amino acid sequence of one or more Kazal-type domains resulting from proteolytic cleavage (e.g., via furin cleavage) of a full-length SPINK polypeptide). Any suitable SPINK polypeptide (or portions thereof) known in the art may be encoded by a polynucleotide of the present disclosure, including, for example, a SPINK1 polypeptide (e.g., a human SPINK1 polypeptide (see e.g., UniProt accession number: P00995)), a SPINK2 polypeptide (e.g., a human SPINK2 polypeptide (see e.g., UniProt accession number: P20155)), a SPINK4 polypeptide (e.g., a human SPINK4 polypeptide (see e.g., UniProt accession number: O60575)), a SPINK5 polypeptide (e.g., a human SPINK5 polypeptide (see e.g., UniProt accession number: Q9NQ38)), a SPINK6 polypeptide (e.g., a human SPINK6 polypeptide (see e.g., UniProt accession number: Q6UWN8)), a SPINK7 polypeptide (e.g., a human SPINK7 polypeptide (see e.g., UniProt accession number: P58062)), a SPINK8 polypeptide (e.g., a human SPINK8 polypeptide (see e.g., UniProt accession number: P0C7L1)), a SPINK9 polypeptide (e.g., a human SPINK9 polypeptide (see e.g., UniProt accession number: Q5DT21)), a SPINK13 polypeptide (e.g., a human SPINK13 polypeptide (see e.g., UniProt accession number: Q1W4C9)), a SPINK14 polypeptide (e.g., a human SPINK14 polypeptide (see e.g., UniProt accession number: Q6IE38)), etc. In some embodiments, a SPINK polypeptide of the present disclosure comprises a sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the amino acid sequence of any SPINK polypeptide described herein or known in the art. Methods of identifying SPINK polypeptide homologs/orthologs from additional species are known to one of ordinary skill in the art, including, for example, using an amino acid sequence alignment program such as the

BLAST® blastp suite or OrthoDB. In some embodiments, the SPINK polypeptide is a human SPINK polypeptide.

[0088] In some embodiments, the present disclosure relates to one or more polynucleotides encoding a SPINK5 polypeptide (e.g., SEQ ID NOS: 7-9), or any portions thereof (e.g., the amino acid sequence of one or more Kazal-type domains resulting from proteolytic cleavage (e.g., via furin cleavage) of a full-length SPINK5 polypeptide (e.g., SEQ ID NOS: 10-25)). Any suitable SPINK5 polypeptide (or portions thereof) known in the art may be encoded by a polynucleotide of the present disclosure, including, for example, a human SPINK5 polypeptide (see e.g., UniProt accession number: 11005), a chimpanzee SPINK5 polypeptide (see e.g., UniProt accession number: 462173), a mouse SPINK5 polypeptide (see e.g., UniProt accession number: Q5K5D4), a rat SPINK5 polypeptide (see e.g., UniProt accession number: D3ZET2), a dog SPINK5 polypeptide (see e.g., UniProt accession number: F1PR80), a cow SPINK5 polypeptide (see e.g., UniProt accession number: F1MJH0), a horse SPINK5 polypeptide (see e.g., UniProt accession number: B9VJ40), a pig SPINK5 polypeptide (see e.g., UniProt accession number: F1RLZ2), etc. In some embodiments, a SPINK5 polypeptide of the present disclosure comprises a sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the amino acid sequence of any SPINK5 polypeptide described herein or known in the art. Methods of identifying SPINK5 polypeptide homologs/orthologs from additional species are known to one of ordinary skill in the art, including, for example, using an amino acid sequence alignment program such as the BLAST® blastp suite or OrthoDB.

[0089] In some embodiments, a SPINK5 polypeptide of the present disclosure is a human SPINK5 polypeptide, or portions thereof (e.g., a polypeptide comprising the amino acid sequence corresponding to one or more Kazal-type domains resulting from proteolytic cleavage (e.g., via human furin cleavage) of a full-length human SPINK5 polypeptide). In some embodiments, a polynucleotide encoding a human SPINK5 polypeptide is a polynucleotide that encodes a polypeptide comprising an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a sequence selected from SEQ ID NOS: 7-25. In some embodiments, a polynucleotide encoding a human SPINK5 polypeptide is a polynucleotide that encodes a polypeptide comprising an amino acid sequence selected from SEQ ID NOS: 7-25. In some embodiments, a polynucleotide encoding a human SPINK5 polypeptide is a polynucleotide that encodes a polypeptide comprising an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a sequence selected from SEQ ID NOS: 7-9. In some embodiments, a polynucleotide encoding a human SPINK5 polypeptide is a polynucleotide that encodes a polypeptide comprising an amino acid sequence selected from SEQ ID NOS: 7-9.

cells of a subject. In some embodiments, expression of the SPINK polypeptide (e.g., a SPINK5 polypeptide) enhances, increases, augments, and/or supplements the levels, function, and/or activity of a SPINK polypeptide in one or more target cells of a subject (e.g., as compared to prior to expression of the exogenous SPINK polypeptide). In some embodiments, expression of the SPINK polypeptide (e.g., a SPINK5 polypeptide) enhances, increases, augments, and/or supplements the anti-inflammatory and/or anti-microbial protection of mucous epithelia in the subject. In some embodiments, expression of the SPINK polypeptide (e.g., a SPINK5 polypeptide) enhances, increases, augments, and/or supplements the integrity and/or protective barrier function of the skin of the subject. In some embodiments, expression of the SPINK polypeptide (e.g., a SPINK5 polypeptide) decreases, augments, and/or inhibits one or more of Kallikrein-5 (KLK5), Kallikrein-7 (KLK7), Kallikrein-14 (KLK14), Caspase-14 (CASP14) and/or trypsin in the subject (e.g., decreases, augments, and/or inhibits one or more activities of KLK5, KLK7, KLK14, CASP14, and/or trypsin).

[0115] Recombinant Nucleic Acids

[0116] In some embodiments, the present disclosure relates to recombinant nucleic acids comprising any one or more of the polynucleotides described herein. In some embodiments, the recombinant nucleic acid is a vector (e.g., an expression vector, a display vector, etc.). In some embodiments, the vector is a DNA vector or an RNA vector. Generally, vectors suitable to maintain, propagate, and/or express polynucleotides to produce one or more polypeptides in a subject may be used. Examples of suitable vectors may include, for example, plasmids, cosmids, episomes, transposons, and viral vectors (e.g., adenoviral vectors, adeno-associated viral vectors, vaccinia viral vectors, Sindbis-viral vectors, measles vectors, herpes viral vectors, lentiviral vectors, retroviral vectors, etc.). In some embodiments, the vector is a herpes viral vector. In some embodiments, the vector is capable of autonomous replication in a host cell. In some embodiments, the vector is incapable of autonomous replication in a host cell. In some embodiments, the vector can integrate into a host DNA. In some embodiments, the vector cannot integrate into a host DNA (e.g., is episomal). Methods of making vectors containing one or more polynucleotides of interest are well known to one of ordinary skill in the art, including, for example, by chemical synthesis or by artificial manipulation of isolated segments of nucleic acids (e.g., by genetic engineering techniques).

[0117] In some embodiments, a recombinant nucleic acid of the present disclosure is a herpes simplex virus (HSV) amplicon. Herpes virus amplicons, including the structural features and methods of making the same, are generally known to one of ordinary skill in the art (see e.g., de Silva S. and Bowers W. "Herpes Virus Amplicon Vectors". *Viruses* 2009, 1, 594-629). In some embodiments, the herpes simplex virus amplicon is an HSV-1 amplicon. In some embodiments, the herpes simplex virus amplicon is an HSV-1 hybrid amplicon. Examples of HSV-1 hybrid amplicons may include, but are not limited to, HSV/AAV hybrid amplicons, HSV/EBV hybrid amplicons, HSV/EBV/RV hybrid amplicons, and/or HSV/Sleeping Beauty hybrid amplicons. In some embodiments, the amplicon is an HSV/AAV hybrid amplicon. In some embodiments, the amplicon is an HSV/Sleeping Beauty hybrid amplicon.

[0118] In some embodiments, a recombinant nucleic acid of the present disclosure is a recombinant herpes virus genome. The recombinant herpes virus genome may be a recombinant genome from any member of the Herpesviridae family of DNA viruses known in the art, including, for example, a recombinant herpes simplex virus genome, a recombinant varicella zoster virus genome, a recombinant human cytomegalovirus genome, a recombinant herpesvirus 6A genome, a recombinant herpesvirus 6B genome, a recombinant herpesvirus 7 genome, a recombinant Kaposi's sarcoma-associated herpesvirus genome, and any combinations or derivatives thereof. In some embodiments, the recombinant herpes virus genome comprises one or more (e.g., one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, etc.) inactivating mutations. In some embodiments, the one or more inactivating mutations are in one or more (e.g., one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, etc.) herpes virus genes. In some embodiments, the recombinant herpes virus genome is attenuated (e.g., as compared to a corresponding, wild-type herpes virus genome). In some embodiments, the recombinant herpes virus genome is replication competent. In some embodiments, the recombinant herpes virus genome is replication defective.

[0119] In some embodiments, the recombinant nucleic acid is a recombinant herpes simplex virus (HSV) genome. In some embodiments, the recombinant herpes simplex virus genome is a recombinant type 1 herpes simplex virus (HSV-1) genome, a recombinant type 2 herpes simplex virus (HSV-2) genome, or any derivatives thereof. In some embodiments, the recombinant herpes simplex virus genome is a recombinant HSV-1 genome. In some embodiments, the recombinant HSV-1 genome may be from any HSV-1 strain known in the art, including, for example, strains 17, Ty25, R62, S25, Ku86, S23, R11, Ty148, Ku47, H166syn, 1319-2005, F-13, M-12, 90237, F-17, KOS, 3083-2008, F12g, L2, CD38, H193, M-15, India 2011, 0116209, F-111, 66-207, 2762, 369-2007, 3355, MacIntyre, McKrae, 7862, 7-hse, HF10, 1394,2005, 270-2007, OD4, SC16, M-19, 4J1037, 5J1060, J1060, KOS79, 132-1988, 160-1982, H166, 2158-2007, RE, 78326, F18g, F11, 172-2010, H129, F, E4, CJ994, F14g, E03, E22, E10, E06, E11, E25, E23, E35, E15, E07, E12, E14, E08, E19, E13, ATCC 2011, etc. (see e.g., Bowen et al. *J Virol.* 2019 Apr. 3; 93(8)). In some embodiments, the recombinant HSV-1 genome is from the KOS strain. In some embodiments, the recombinant HSV-1 genome is not from the McKrae strain. In some embodiments, the recombinant herpes simplex virus genome is attenuated. In some embodiments, the recombinant herpes simplex virus genome is replication competent. In some embodiments, the recombinant herpes simplex virus genome is replication defective. In some embodiments, the recombinant herpes simplex virus genome comprises one or more (e.g., one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, etc.) inactivating mutations. In some embodiments, the one or more inactivating mutations are in one or more (e.g., one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, etc.) herpes simplex virus genes. As used herein, an "inactivating mutation" may refer to any mutation that results in a gene or regulon product (RNA or protein) having

reduced, undetectable, or eliminated quantity and/or function (e.g., as compared to a corresponding sequence lacking the inactivating mutation). Examples of inactivating mutations may include, but are not limited to, deletions, insertions, point mutations, and rearrangements in transcriptional control sequences (promoters, enhancers, insulators, etc.) and/or coding sequences of a given gene or regulon. Any suitable method of measuring the quantity of a gene or regulon product known in the art may be used, including, for example, qPCR, Northern blots, RNAseq, western blots, ELISAs, etc.

[0120] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in at least one, at least two, at least three, at least four, at least five, at least six, at least seven, or all eight of the Infected Cell Protein (or Infected Cell Polypeptide) (ICP) 0, ICP4, ICP22, ICP27, ICP47, thymidine kinase (tk), Long Unique Region (UL) 41 and/or UL55 herpes simplex virus genes. In some embodiments, the recombinant herpes simplex virus genome does not comprise an inactivating mutation in the ICP34.5 (one or both copies) and/or ICP47 herpes simplex virus genes (e.g., to avoid production of an immune-stimulating virus). In some embodiments, the recombinant herpes simplex virus genome does not comprise an inactivating mutation in the ICP34.5 (one or both copies) herpes simplex virus gene. In some embodiments, the recombinant herpes simplex virus genome does not comprise an inactivating mutation in the ICP47 herpes simplex virus gene. In some embodiments, the recombinant herpes simplex virus genome does not comprise an inactivating mutation in the ICP34.5 (one or both copies) and ICP47 herpes simplex virus genes. In some embodiments, the recombinant herpes simplex virus genome is not oncolytic.

[0121] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies). In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies), and further comprises an inactivating mutation in the ICP4 (one or both copies), ICP22, ICP27, ICP47, UL41, and/or UL55 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies), and an inactivating mutation in the ICP4 gene (one or both copies). In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies), and an inactivating mutation in the ICP22 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies), and an inactivating mutation in the ICP4 gene (one or both copies), and an inactivating mutation in the ICP22 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies), an inactivating mutation in the ICP4 gene (one or both copies), and an inactivating mutation in the UL41 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 gene (one or both copies), an inactivating mutation in the ICP22 gene, and an inactivating mutation in the UL41 gene. In some embodiments, the recombinant herpes simplex virus

genome comprises an inactivating mutation in the ICP0 gene (one or both copies), an inactivating mutation in the ICP4 gene (one or both copies), an inactivating mutation in the ICP22 gene, and an inactivating mutation in the UL41 gene. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, and/or UL41 genes. In some embodiments, the recombinant herpes simplex virus genome further comprises an inactivating mutation in the ICP27, ICP47, and/or UL55 genes.

[0122] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 gene (one or both copies). In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 gene (one or both copies), and further comprises an inactivating mutation in the ICP0 (one or both copies), ICP22, ICP27, ICP47, UL41, and/or UL55 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 gene (one or both copies), and an inactivating mutation in the ICP22 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 gene (one or both copies), and an inactivating mutation in the UL41 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 gene (one or both copies), an inactivating mutation in the ICP22 gene, and an inactivating mutation in the UL41 gene. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the ICP4 (one or both copies), ICP22, and/or UL41 genes. In some embodiments, the recombinant herpes simplex virus genome further comprises an inactivating mutation in the ICP0 (one or both copies), ICP27, ICP47, and/or UL55 genes.

[0123] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP22 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP22 gene, and further comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP27, ICP47, UL41, and/or UL55 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP22 gene, and an inactivating mutation UL41 gene. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the ICP22 and/or UL41 genes. In some embodiments, the recombinant herpes simplex virus genome further comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP27, ICP47, and/or UL55 genes.

[0124] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP27 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP27 gene, and further comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP47, UL41, and/or UL55 genes. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the ICP27 gene.

[0125] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP47 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP47 gene, and further comprises an

inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27, UL41, and/or UL55 genes. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the ICP47 gene.

[0126] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL41 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL41 gene, and further comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27, ICP47, and/or UL55 genes. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the UL41 gene.

[0127] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL55 gene. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL55 gene, and further comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27, ICP47, and/or UL41 genes. In some embodiments, the inactivating mutation is a deletion of the coding sequence of the UL55 gene.

[0128] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in (e.g., a deletion of) the internal repeat (Joint) region comprising the internal repeat long (IR_L) and internal repeat short (IR_S) regions. In some embodiments, inactivation (e.g., deletion) of the Joint region eliminates one copy each of the ICP4 and ICP0 genes. In some embodiments, inactivation (e.g., deletion) of the Joint region further inactivates (e.g., deletes) the promoter for the ICP22 and ICP47 genes. If desired, expression of one or both of these genes can be restored by insertion of an immediate early promoter into the recombinant herpes simplex virus genome (see e.g., Hill et al. (1995). *Nature* 375(6530): 411-415; Goldsmith et al. (1998). *J Exp Med* 187(3): 341-348). Without wishing to be bound by theory, it is believed that inactivating (e.g., deleting) the Joint region may contribute to the stability of the recombinant herpes simplex virus genome and/or allow for the recombinant herpes simplex virus genome to accommodate more and/or larger transgenes.

[0129] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 (one or both copies), ICP22, and ICP27 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 (one or both copies), ICP27, and UL55 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP4 (one or both copies), ICP22, ICP27, ICP47, and UL55 genes. In some embodiments, the inactivating mutation in the ICP4 (one or both copies), ICP27, and/or UL55 genes is a deletion of the coding sequence of the ICP4 (one or both copies), ICP27, and/or UL55 genes. In some embodiments, the inactivating mutation in the ICP22 and ICP47 genes is a deletion in the promoter region of the ICP22 and ICP47 genes (e.g., the ICP22 and ICP47 coding sequences are intact but are not transcriptionally active). In some embodiments, the recombinant herpes simplex virus genome comprises a deletion in the coding sequence of the ICP4 (one or both copies), ICP27, and UL55 genes, and a deletion in the promoter region of the ICP22 and ICP47 genes. In some embodi-

ments, the recombinant herpes simplex virus genome further comprises an inactivating mutation in the ICP0 (one or both copies) and/or UL41 genes.

[0130] In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 (one or both copies) and ICP4 (one or both copies) genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), and ICP22 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, and ICP27 genes. In some embodiments, the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27 and UL55 genes. In some embodiments, the inactivating mutation in the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27 and/or UL55 genes comprises a deletion of the coding sequence of the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27 and/or UL55 genes. In some embodiments, the recombinant herpes simplex virus genome further comprises an inactivating mutation in the ICP47 and/or the UL41 genes.

[0131] In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within one, two, three, four, five, six, seven or more viral gene loci. Examples of suitable viral loci may include, without limitation, the ICP0 (one or both copies), ICP4 (one or both copies), ICP22, ICP27, ICP47, tk, UL41 and/or UL55 herpes simplex viral gene loci. In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within one or both of the viral ICP4 gene loci (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in one or both of the ICP4 loci). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within the viral ICP22 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in the ICP22 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within the viral ICP47 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in the ICP47 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within the viral UL41 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in the UL41 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within the viral ICP47 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in the ICP47 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within one or both of the viral ICP4 gene loci, and one or more polynucleotides of the present disclosure within the viral ICP22 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in one or both of the ICP4 loci, and a polynucleotide encoding the same or a different SPINK polypeptide in the ICP22 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within one or both of the viral ICP4 gene loci, and one or more polynucleotides of the present disclosure within the viral UL41 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK poly-

peptide in one or both of the ICP4 loci, and a polynucleotide encoding the same or a different SPINK polypeptide in the UL41 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within the viral ICP22 gene locus, and one or more polynucleotides of the present disclosure within the viral UL41 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in the ICP22 locus, and a polynucleotide encoding the same or a different SPINK polypeptide in the UL41 locus). In some embodiments, a recombinant herpes simplex virus genome comprises one or more polynucleotides of the present disclosure within one or both of the viral ICP4 gene loci, one or more polynucleotides of the present disclosure within the viral ICP22 gene locus, and one or more polynucleotides of the present disclosure within the viral UL41 gene locus (e.g., a recombinant virus comprising a polynucleotide encoding a SPINK polypeptide in one or both of the ICP4 loci, a polynucleotide encoding the same or a different SPINK polypeptide in the ICP22 locus, and a polynucleotide encoding a SPINK polypeptide in the UL41 locus (which may be the same SPINK polypeptide encoded in the ICP4 and/or ICP22 loci or may be a different SPINK polypeptide than those encoded in the ICP4 and/or ICP22 loci).

[0132] In some embodiments, the recombinant herpes virus genome (e.g., a recombinant herpes simplex virus genome) has been engineered to decrease or eliminate expression of one or more herpes virus genes (e.g., one or more toxic herpes virus genes), such as one or both copies of the HSV ICP0 gene, one or both copies of the HSV ICP4 gene, the HSV ICP22 gene, the HSV UL41 gene, the HSV ICP27 gene, etc. In some embodiments, the recombinant herpes virus genome (e.g., recombinant herpes simplex virus genome) has been engineered to reduce cytotoxicity of the recombinant genome (e.g., when introduced into a target cell) as compared to a corresponding wild-type herpes virus genome (e.g., a wild-type herpes simplex virus genome). In some embodiments, the target cell is a human cell. In some embodiments, the target cell is a cell of the epidermis and/or dermis (e.g., a cell of the human epidermis and/or dermis). In some embodiments, the target cell is a keratinocyte or fibroblast (e.g., a human keratinocyte or human fibroblast). In some embodiments, cytotoxicity (e.g., in human keratinocytes and/or fibroblasts) of the recombinant genome (e.g., a recombinant herpes simplex virus genome) is reduced by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% as compared to a corresponding wild-type herpes virus genome (e.g., measuring the relative cytotoxicity of a recombinant Δ ICP4 (one or both copies) herpes simplex virus genome vs. a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); measuring the relative cytotoxicity of a recombinant Δ ICP4 (one or both copies)/ Δ ICP22 herpes simplex virus genome vs. a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); etc.). In some embodiments, cytotoxicity (e.g., in human keratinocytes and/or fibroblasts) of the recombinant genome (e.g., a

recombinant herpes simplex virus genome) is reduced by at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 15-fold, at least about 20-fold, at least about 25-fold, at least about 50-fold, at least about 75-fold, at least about 100-fold, at least about 250-fold, at least about 500-fold, at least about 750-fold, at least about 1000-fold, or more as compared to a corresponding wild-type herpes virus genome (e.g., measuring the relative cytotoxicity of a recombinant Δ ICP4 (one or both copies) herpes simplex virus genome vs. a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); measuring the relative cytotoxicity of a recombinant Δ ICP4 (one or both copies)/ Δ ICP22 herpes simplex virus genome vs. a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); etc.). Methods of measuring cytotoxicity are known to one of ordinary skill in the art, including, for example, through the use of vital dyes (formazan dyes), protease biomarkers, an MTT assay (or an assay using related tetrazolium salts such as XTT, MTS, water-soluble tetrazolium salts, etc.), measuring ATP content, etc.

[0133] In some embodiments, the recombinant herpes virus genome (e.g., a recombinant herpes simplex virus genome) has been engineered to reduce its impact on host cell proliferation after exposure of a target cell to the recombinant genome, as compared to a corresponding wild-type herpes virus genome (e.g., a wild-type herpes simplex virus genome). In some embodiments, the target cell is a human cell. In some embodiments, the target cell is a cell of the epidermis and/or dermis (e.g., a cell of the human epidermis and/or dermis). In some embodiments, the target cell is a keratinocyte or fibroblast (e.g., a human keratinocyte or human fibroblast). In some embodiments, host cell proliferation (e.g., of human keratinocytes and/or fibroblasts) after exposure to the recombinant genome is at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% faster as compared to host cell proliferation after exposure to a corresponding wild-type herpes virus genome (e.g., measuring the relative cellular proliferation after exposure to a recombinant Δ ICP4 (one or both copies) herpes simplex virus genome vs. cellular proliferation after exposure to a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); measuring the relative cellular proliferation after exposure to a recombinant Δ ICP4 (one or both copies)/ Δ ICP22 herpes simplex virus genome vs. cellular proliferation after exposure to a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); etc.). In some embodiments, host cell proliferation (e.g., of human keratinocytes and/or fibroblasts) after exposure to the recombinant genome is at least about 1.5-fold, at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 15-fold, at least about 20-fold, at least about 25-fold, at least

about 50-fold, at least about 75-fold, at least about 100-fold, at least about 250-fold, at least about 500-fold, at least about 750-fold, or at least about 1000-fold faster as compared to host cell proliferation after exposure to a corresponding wild-type herpes virus genome (e.g., measuring the relative cellular proliferation after exposure to a recombinant Δ ICP4 (one or both copies) herpes simplex virus genome vs. cellular proliferation after exposure to a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); measuring the relative cellular proliferation after exposure to a recombinant Δ ICP4 (one or both copies)/ Δ ICP22 herpes simplex virus genome vs. cellular proliferation after exposure to a wild-type herpes simplex virus genome in human keratinocytes or fibroblasts (primary cells or cell lines); etc.). Methods of measuring cellular proliferation are known to one of ordinary skill in the art, including, for example, through the use of a Ki67 cell proliferation assay, a BrdU cell proliferation assay, etc.

[0134] A vector (e.g., herpes viral vector) may include one or more polynucleotides of the present disclosure in a form suitable for expression of the polynucleotide in a host cell. Vectors may include one or more regulatory sequences operatively linked to the polynucleotide to be expressed (e.g., as described above).

[0135] In some embodiments, a recombinant nucleic acid (e.g., a recombinant herpes virus genome, such as a recombinant herpes simplex virus genome) of the present disclosure comprises one or more of the polynucleotides described herein inserted in any orientation in the recombinant nucleic acid. If the recombinant nucleic acid comprises two or more polynucleotides described herein (e.g., two or more, three or more, etc.), the polynucleotides may be inserted in the same orientation or opposite orientations to one another. Without wishing to be bound by theory, incorporating two polynucleotides (e.g., two transgenes) into a recombinant nucleic acid (e.g., a vector) in an antisense orientation may help to avoid read-through and ensure proper expression of each polynucleotide.

[0136] In some embodiments, the present disclosure relates to one or more heterologous polynucleotides (e.g., a bacterial artificial chromosome (BAC)) comprising any of the recombinant nucleic acids described herein.

IV. Viruses

[0137] Certain aspects of the present disclosure relate to viruses comprising any of the polynucleotides and/or recombinant nucleic acids described herein. In some embodiments, the virus is capable of infecting one or more target cells of a subject (e.g., a human). In some embodiments, the virus is suitable for delivering the polynucleotides and/or recombinant nucleic acids into one or more target cells of a subject (e.g., a human). In some embodiments, the present disclosure relates to one or more viral particles comprising any of the polynucleotides and/or recombinant nucleic acids described herein. In some embodiments, the one or more target cells are one or more human cells. In some embodiments, the one or more target cells are one or more cells with a SPINK deficiency (e.g., one or more cells comprising a loss-of-function mutation in, or a pathogenic variant of, a native SPINK gene, such as a SPINK5 gene). In some embodiments, the one or more target cells are one or more cells of the skin (e.g., one or more cells of the epidermis, dermis, and/or subcutis). In some embodiments, the one or more target cells are cells of the epidermis and/or dermis

(e.g., cells of the human epidermis and/or dermis). In some embodiments, the one or more target cells are selected from keratinocytes, melanocytes, Langerhans cells, Merkel cells, mast cells, fibroblasts, and/or adipocytes. In some embodiments, the one or more target cells are keratinocytes. In some embodiments, the one or more target cells reside in the stratum corneum, stratum granulosum, stratum spinulosum, stratum basale, and/or basement membrane. In some embodiments, the one or more target cells are one or more epidermal cells. In some embodiments, the one or more target cells are one or more dermal cells.

[0138] Any suitable virus known in the art may be used, including, for example, adenovirus, adeno-associated virus, retrovirus, lentivirus, sendai virus, papillomavirus, herpes virus (e.g., a herpes simplex virus), vaccinia virus, and/or any hybrid or derivative viruses thereof. In some embodiments, the virus is attenuated. In some embodiments, the virus is replication defective. In some embodiments, the virus is replication competent. In some embodiments, the virus has been modified to alter its tissue tropism relative to the tissue tropism of a corresponding unmodified, wild-type virus. In some embodiments, the virus has reduced cytotoxicity as compared to a corresponding wild-type virus. Methods of producing a virus comprising recombinant nucleic acids are well known to one of ordinary skill in the art.

[0139] In some embodiments, the virus is a member of the Herpesviridae family of DNA viruses, including, for example, a herpes simplex virus, a varicella zoster virus, a human cytomegalovirus, a herpesvirus 6A, a herpesvirus 6B, a herpesvirus 7, and a Kaposi's sarcoma-associated herpesvirus, etc. In some embodiments, the herpes virus is attenuated. In some embodiments, the herpes virus is replication defective. In some embodiments, the herpes virus is replication competent. In some embodiments, the herpes virus has reduced cytotoxicity as compared to a corresponding wild-type herpes virus. In some embodiments, the herpes virus is not oncolytic.

[0140] In some embodiments, the herpes virus is a herpes simplex virus. Herpes simplex viruses comprising recombinant nucleic acids may be produced by a process disclosed, for example, in WO2015/009952 and/or WO2017/176336. In some embodiments, the herpes simplex virus is attenuated. In some embodiments, the herpes simplex virus is replication defective. In some embodiments, the herpes simplex virus is replication competent. In some embodiments, the herpes simplex virus is an HSV-1 virus, an HSV-2, or any derivatives thereof. In some embodiments, the herpes simplex virus is an HSV-1 virus. In some embodiments, the HSV-1 is attenuated. In some embodiments, the HSV-1 is replication defective. In some embodiments, the HSV-1 is replication competent. In some embodiments, the herpes simplex virus (e.g., the HSV-1) has reduced cytotoxicity as compared to a corresponding wild-type herpes simplex virus (e.g., a wild-type HSV-1). In some embodiments, the herpes simplex virus (e.g., the HSV-1) is not oncolytic.

[0141] In some embodiments, the herpes simplex virus has been modified to alter its tissue tropism relative to the tissue tropism of an unmodified, wild-type herpes simplex virus. In some embodiments, the herpes simplex virus comprises a modified envelope. In some embodiments, the modified envelope comprises one or more (e.g., one or more, two or more, three or more, four or more, etc.) mutant herpes simplex virus glycoproteins. Examples of herpes simplex

virus glycoproteins may include, but are not limited to, the glycoproteins gB, gC, gD, gH, and gL. In some embodiments, the modified envelope alters the herpes simplex virus tissue tropism relative to a wild-type herpes simplex virus.

[0142] In some embodiments, the transduction efficiency (in vitro and/or in vivo) of a virus of the present disclosure (e.g., a herpes virus such as a herpes simplex virus) for one or more target cells (e.g., one or more human keratinocytes and/or fibroblasts) is at least about 25%. For example, the transduction efficiency of the virus for one or more target cells may be at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99%, at least about 99.5%, or more. In some embodiments, the virus is a herpes simplex virus and the transduction efficiency of the virus for one or more target cells (e.g., one or more human keratinocytes and/or fibroblasts) is about 85% to about 100%. In some embodiments, the virus is a herpes simplex virus and the transduction efficiency of the virus for one or more target cells (e.g., one or more human keratinocytes and/or fibroblasts) is at least about 85%, at least about 86%, at least about 87%, at least about 88%, at least about 89%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100%. Methods of measuring viral transduction efficiency in vitro or in vivo are well known to one of ordinary skill in the art, including, for example, qPCR analysis, deep sequencing, western blotting, fluorometric analysis (such as fluorescent in situ hybridization (FISH), fluorescent reporter gene expression, immunofluorescence, FACS), etc.

V. PHARMACEUTICAL COMPOSITIONS AND FORMULATIONS

[0143] Certain aspects of the present disclosure relate to pharmaceutical compositions and formulations comprising any of the recombinant nucleic acids (e.g., recombinant herpes virus genomes) and/or viruses (e.g., herpes viruses comprising the recombinant genomes) described herein (such as a herpes simplex virus comprising a recombinant herpes simplex virus genome), and a pharmaceutically acceptable excipient or carrier.

[0144] In some embodiments, the pharmaceutical composition or formulation comprises any one or more of the viruses (e.g., herpes viruses) described herein. In some embodiments, the pharmaceutical composition or formulation comprises from about 10^4 to about 10^{12} plaque forming units (PFU)/mL of the virus. For example, the pharmaceutical composition or formulation may comprise from about 10^4 to about 10^{12} , about 10^5 to about 10^{12} , about 10^6 to about 10^{12} , about 10^{11} to about 10^{12} , about 10^8 to about 10^{12} , about 10^9 to about 10^{12} , about 10^{10} to about 10^{12} , about 10^{11} to about 10^{12} , about 10^4 to about 10^{11} , about 10^5 to about 10^{11} , about 10^6 to about 10^{11} , about 10^7 to about 10^{11} , about 10^8 to about 10^{11} , about 10^9 to about 10^{11} , about 10^{10} to about 10^{11} , about 10^4 to about 10^{10} about 10^5 to about 10^{10} about 10^6 to about 10^{10} , about 10^7 to about 10^{10} about 10^8 to about 10^{10} about 10^9 to about 10^{10} about 10^4 to about 10^9 about 10^5 to about 10^9 , about 10^6 to about 10^9 , about 10^7 to about 10^9 , about 10^8 to about 10^9 , about 10^4 to about 10^8 , about 10^5 to

about 10^8 , about 10^6 to about 10^8 , about 10^7 to about 10^8 , about 10^4 to about 10^7 , about 10^5 to about 10^7 , about 10^6 to about 10^7 , about 10^4 to about 10^6 , about 10^5 to about 10^6 , or about 10^4 to about 10^5 PFU/mL of the virus. In some embodiments, the pharmaceutical composition or formulation comprises about 10^4 , about 10^5 , about 10^6 , about 10^7 , about 10^8 , about 10^9 , about 10^{10} , about 10^{11} , or about 10^{12} PFU/mL of the virus.

[0145] Pharmaceutical compositions and formulations can be prepared by mixing the active ingredient(s) (such as a recombinant nucleic acid and/or a virus) having the desired degree of purity with one or more pharmaceutically acceptable carriers or excipients. Pharmaceutically acceptable carriers or excipients are generally nontoxic to recipients at the dosages and concentrations employed, and may include, but are not limited to: buffers (such as phosphate, citrate, acetate, and other organic acids); antioxidants (such as ascorbic acid and methionine); preservatives (such as octadecyldimethylbenzyl ammonium chloride, benzalkonium chloride, benzethonium chloride, phenol, butyl or benzyl alcohol, alkyl parabens, catechol, resorcinol, cyclohexanol, 3-pentanol, and m-cresol); amino acids (such as glycine, glutamine, asparagine, histidine, arginine, or lysine); low molecular weight (less than about 10 residues) polypeptides; proteins (such as serum albumin, gelatin, or immunoglobulins); polyols (such as glycerol, e.g., formulations including 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, etc. glycerol); hydrophilic polymers (such as polyvinylpyrrolidone); monosaccharides, disaccharides, and other carbohydrates (including glucose, mannose, or dextrans); chelating agents (such as EDTA); sugars (such as sucrose, mannitol, trehalose, or sorbitol); salt-forming counter-ions (such as sodium); metal complexes (such as Zn-protein complexes); and/or non-ionic surfactants (such as polyethylene glycol (PEG)). A thorough discussion of pharmaceutically acceptable carriers is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991).

[0146] In some embodiments, the pharmaceutical composition or formulation comprises one or more lipid (e.g., cationic lipid) carriers. In some embodiments, the pharmaceutical composition or formulation comprises one or more nanoparticle carriers. Nanoparticles are submicron (less than about 1000 nm) sized drug delivery vehicles that can carry encapsulated drugs (such as synthetic small molecules, proteins, peptides, cells, viruses, and nucleic acid-based biotherapeutics) for rapid or controlled release. A variety of molecules (e.g., proteins, peptides, recombinant nucleic acids, etc.) can be efficiently encapsulated in nanoparticles using processes well known in the art. In some embodiments, a molecule "encapsulated" in a nanoparticle may refer to a molecule (such as a virus) that is contained within the nanoparticle or attached to and/or associated with the surface of the nanoparticle, or any combination thereof. Nanoparticles for use in the compositions or formulations described herein may be any type of biocompatible nanoparticle known in the art, including, for example, nanoparticles comprising poly(lactic acid), poly(glycolic acid), PLGA, PLA, PGA, and any combinations thereof (see e.g., Vauthier et al. Adv Drug Del Rev. (2003) 55: 519-48; US2007/0148074; US2007/0092575; US2006/0246139; U.S. Pat. Nos. 5,753,234; 7,081,483; and WO2006/052285).

[0147] In some embodiments, the pharmaceutically acceptable carrier or excipient may be adapted for or suit-

able for any administration route known in the art, including, for example, intravenous, intramuscular, subcutaneous, cutaneous, oral, nasal, intratracheal, sublingual, buccal, topical, transdermal, intradermal, intraperitoneal, intraorbital, intravitreal, subretinal, transmucosal, intraarticular, by implantation, by inhalation, intrathecal, intraventricular, and/or intranasal administration. In some embodiments, the pharmaceutical composition or formulation is adapted for or suitable for any administration route known in the art, including, for example, intravenous, intramuscular, subcutaneous, cutaneous, oral, nasal, intratracheal, sublingual, buccal, topical, transdermal, intradermal, intraperitoneal, intraorbital, intravitreal, subretinal, transmucosal, intraarticular, by implantation, by inhalation, intrathecal, intraventricular, and/or intranasal administration. In some embodiments, the pharmaceutically acceptable carrier or excipient is adapted for or suitable for topical, transdermal, subcutaneous, intradermal, and/or transmucosal administration. In some embodiments, the pharmaceutical composition or formulation is adapted for or suitable for topical, transdermal, subcutaneous, intradermal, and/or transmucosal administration. In some embodiments, the pharmaceutically acceptable carrier or excipient is adapted for or suitable for topical, transdermal, and/or intradermal administration. In some embodiments, the pharmaceutical composition or formulation is adapted for or suitable for topical, transdermal, and/or intradermal administration. In some embodiments, the pharmaceutically acceptable carrier or excipient is adapted for or suitable for topical administration. In some embodiments, the pharmaceutical composition or formulation is adapted for or suitable for topical administration.

[0148] Examples of carriers or excipients adapted for or suitable for use in pharmaceutical compositions or formulations of the present disclosure may include, but are not limited to, ointments, oils, pastes, creams, aerosols, suspensions, emulsions, fatty ointments, gels (e.g., methylcellulose gels, such as carboxy methylcellulose, hydroxypropyl methylcellulose, etc.), powders, liquids, lotions, solutions, sprays, patches (e.g., transdermal patches or microneedle patches), adhesive strips, a microneedle or microneedle arrays, and inhalants. In some embodiments, the carrier or excipient (e.g., the pharmaceutically acceptable carrier or excipient) comprises one or more (e.g., one or more, two or more, three or more, four or more, five or more, etc.) of an ointment, oil, paste, cream, aerosol, suspension, emulsion, fatty ointment, gel, powder, liquid, lotion, solution, spray, patch, adhesive strip, and an inhalant. In some embodiments, the carrier comprises a patch (e.g. a patch that adheres to the skin), such as a transdermal patch or microneedle patch. In some embodiments, the carrier comprises a microneedle or microneedle array. Methods for making and using microneedle arrays suitable for composition delivery are generally known in the art (see e.g., Kim Y. et al. "Microneedles for drug and vaccine delivery". *Advanced Drug Delivery Reviews* 2012, 64 (14): 1547-68).

[0149] In some embodiments, the pharmaceutical composition or formulation further comprises one or more additional components. Examples of additional components may include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc,

silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); wetting agents (e.g., sodium lauryl sulphate, etc.); salt solutions; alcohols; polyethylene glycols; gelatin; lactose; amylase; magnesium stearate; talc; silicic acid; viscous paraffin; methylcellulose; polyvinylpyrrolidone; sweetenings; flavorings; perfuming agents; colorants; moisturizers; sunscreens; antibacterial agents; agents able to stabilize polynucleotides or prevent their degradation, and the like. In some embodiments, the pharmaceutical composition or formulation comprises a methylcellulose gel, such as a carboxy methylcellulose gel, a hydroxypropyl methylcellulose gel, etc. (e.g., at about 0.5%, at about 1%, at about 1.5%, at about 2%, at about 2.5%, at about 3%, at about 3.5%, at about 4%, at about 4.5%, at about 5%, at about 5.5%, at about 6%, at about 6.5%, at about 7%, at about 7.5%, at about 8%, at about 8.5%, at about 9%, at about 9.5%, at about 10%, at about 10.5%, at about 11%, at about 11.5%, at about 12%, etc.). In some embodiments, the pharmaceutical composition or formulation comprises a phosphate buffer. In some embodiments, the pharmaceutical composition or formulation comprises glycerol (e.g., at about 1%, about 2%, about 3%, about 4%, about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, etc.). In some embodiments, the pharmaceutical composition or formulation comprises a methylcellulose gel (a carboxy methylcellulose gel, a hydroxypropyl methylcellulose gel, etc.), a phosphate buffer, and/or glycerol.

[0150] Compositions and formulations (e.g., pharmaceutical compositions and formulations) to be used for in vivo administration are generally sterile. Sterility may be readily accomplished, e.g., by filtration through sterile filtration membranes.

[0151] In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used to deliver one or more polynucleotides encoding a SPINK polypeptide into one or more cells of a subject (e.g., one or more SPINK-deficient cells, one or more cells harboring a SPINK gene mutation, etc.). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations may be used in a therapy. In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the treatment of a disease or condition that would benefit from the expression of a SPINK polypeptide (e.g., a disease associated with a SPINK deficiency and/or a disease associated with a SPINK gene mutation). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used for providing prophylactic, palliative, or therapeutic relief to one or more signs or symptoms of Netherton Syndrome (e.g., via delivery of a SPINK5 polypeptide), atopic dermatitis (e.g., via delivery of a SPINK5 polypeptide), hereditary pancreatitis (PCTT) (e.g., via delivery of a SPINK1 polypeptide), tropical calcific pancreatitis (e.g., via delivery of a SPINK1 polypeptide), and/or spermatogenic failure 29 (SPGF29) (e.g., via delivery of a SPINK2 polypeptide). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations

described herein may be used to treat Netherton Syndrome (e.g., via delivery of a human SPINK5 polypeptide). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used to treat atopic dermatitis (e.g., via delivery of a human SPINK5 polypeptide).

[0152] In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the preparation or manufacture of a medicament. In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the preparation or manufacture of a medicament useful for delivering one or more polynucleotides encoding a SPINK polypeptide into one or more cells of a subject (e.g., one or more SPINK-deficient cells, one or more cells harboring a SPINK gene mutation, etc.). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the preparation or manufacture of a medicament useful for the treatment of a disease or condition that would benefit from the expression of a SPINK polypeptide (e.g., a disease associated with a SPINK deficiency and/or a disease associated with a SPINK gene mutation). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the preparation or manufacture of a medicament useful for the treatment of Netherton Syndrome (e.g., via delivery of a SPINK5 polypeptide), atopic dermatitis (e.g., via delivery of a SPINK5 polypeptide), hereditary pancreatitis (PCTT) (e.g., via delivery of a SPINK1 polypeptide), tropical calcific pancreatitis (e.g., via delivery of a SPINK1 polypeptide), and/or spermatogenic failure 29 (SPGF29) (e.g., via delivery of a SPINK2 polypeptide). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the preparation or manufacture of a medicament useful for the treatment of Netherton Syndrome (e.g., via delivery of a human SPINK5 polypeptide). In some embodiments, any of the recombinant nucleic acids, viruses, and/or pharmaceutical compositions or formulations described herein may be used in the preparation or manufacture of a medicament useful for the treatment of atopic dermatitis (e.g., via delivery of a human SPINK5 polypeptide).

VI. METHODS

[0153] Certain aspects of the present disclosure relate to enhancing, increasing, augmenting, and/or supplementing the levels of a SPINK polypeptide in one or more cells of a subject comprising administering to the subject an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein. In some embodiments, the SPINK polypeptide is a human SPINK polypeptide. In some embodiments, the SPINK polypeptide is a SPINK5 polypeptide. In some embodiments, the SPINK5 polypeptide is a human SPINK5 polypeptide. In some embodiments, the subject is a human. In some embodiments, the subject's genome comprises a mutation (e.g., a loss-of-function mutation, a pathogenic variant) in an endogenous SPINK gene (such as an endogenous SPINK5 gene). In some embodi-

ments, the subject suffers from Netherton Syndrome. In some embodiments, the subject suffers from atopic dermatitis.

[0154] In some embodiments, administration of the recombinant nucleic acid, virus, medicament, and/or pharmaceutical composition or formulation to the subject increases SPINK (e.g., SPINK5) levels (transcript or protein levels) by at least about 2-fold in one or more contacted or treated cells of the subject, as compared to the endogenous levels of the SPINK in one or more corresponding untreated cells of the subject. For example, administration of the recombinant nucleic acid, virus, medicament, and/or pharmaceutical composition or formulation may increase SPINK (e.g., SPINK5) levels (transcript or protein levels) by at least about 2-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, at least about 10-fold, at least about 15-fold, at least about 20-fold, at least about 25-fold, at least about 50-fold, at least about 75-fold, at least about 100-fold, at least about 250-fold, at least about 500-fold, at least about 750-fold, at least about 1000-fold, or more in one or more contacted or treated cells of the subject, as compared to the endogenous levels of the SPINK in one or more corresponding untreated cells of the subject. In some embodiments, the one or more contacted or treated cells are one or more cells of the epidermis and/or dermis (e.g., a keratinocyte). Methods of measuring transcript or protein levels from a sample are well known to one of ordinary skill in the art, including, for example, by qPCR, western blot, mass spectrometry, etc.

[0155] In some embodiments, administering to an individual an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulation described herein enhances, increases, augments, and/or supplements the integrity and/or protective barrier function of the skin of the subject. In some embodiments, administering to an individual an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein decreases, augments, and/or inhibits one or more of Kallikrein-5 (KLK5), Kallikrein-7 (KLK7), Kallikrein-14 (KLK14), Caspase-14 (CASP14) and trypsin in the subject (e.g., decreases, augments, and/or inhibits one or more activities (e.g., proteolytic activities) of KLK5, KLK7, KLK14, CASP14, and/or trypsin).

[0156] Other aspects of the present disclosure relate to enhancing, increasing, augmenting, and/or supplementing anti-inflammatory and/or anti-microbial protection of mucous epithelia in a subject in need thereof comprising administering to the subject an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein. In some embodiments, administration of the recombinant nucleic acid, virus, medicament, and/or pharmaceutical composition reduces the susceptibility to skin infections of the subject.

[0157] Other aspects of the present disclosure relate to repressing or inhibiting desquamation in a subject in need thereof comprising administering to the subject an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein.

[0158] Other aspects of the present disclosure relate to reducing or treating a skin barrier defect in a subject in need

thereof comprising administering to the subject an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein. In some embodiments, the skin barrier defect is transepidermal water loss (TEWL), also called transepithelial water loss. In some embodiments, the methods of the present disclosure reduce transepidermal water loss in a subject in need thereof. Methods of measuring skin barrier function, including TEWL, are well known to one of ordinary skill in the art, including, for example, by any of the methods described by Antonov et al. (Curr Probl Dermatol. 2016; 49:61-70).

[0159] Other aspects of the present disclosure relate to providing prophylactic, palliative, or therapeutic relief of one or more signs or symptoms of Netherton Syndrome in a subject in need thereof comprising administering to the subject an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein. In some embodiments, the subject's genome comprises a mutation (e.g., a loss-of-function mutation, a pathogenic variant) in an endogenous SPINK5 gene. In some embodiments, the recombinant nucleic acid, virus, medicament, and/or pharmaceutical composition or formulation comprises one or more polynucleotides encoding a human SPINK5 polypeptide.

[0160] Signs and/or symptoms of Netherton Syndrome may include, but are not limited to defective keratinization, a defective skin barrier, chronic skin inflammation, universal pruritus (itch), severe dehydration, stunted growth, trichorrhexis invaginata and/or trichorrhexis *nodosa* (hair shaft defect, also known as bamboo hair), leaking fluid from the skin, development of ring-like lesions on the skin, eczema, increased susceptibility to infection (particularly of the skin, including recurrent skin infections with *Staphylococcus*), increased susceptibility to allergy, development of scaly/reddish skin (similar to atopic dermatitis), development of ichthyosis linearis circumflexa and/or ichthyosiform erythroderma, altered immunoglobulin levels (typically high IgE and low to normal IgG immunoglobulins), immature natural killer cells (having reduced lytic function), and difficulty regulating body temperature.

[0161] Other aspects of the present disclosure relate to providing prophylactic, palliative, or therapeutic relief of one or more signs or symptoms of atopic dermatitis in a subject in need thereof comprising administering to the subject an effective amount of any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein. In some embodiments, the subject's genome comprises a mutation (e.g., a loss-of-function mutation, a pathogenic variant) in an endogenous SPINK5 gene. In some embodiments, the recombinant nucleic acid, virus, medicament, and/or pharmaceutical composition or formulation comprises one or more polynucleotides encoding a human SPINK5 polypeptide.

[0162] Signs and symptoms of atopic dermatitis may include, without limitation: dry skin; itching, which may be severe, especially at night; red to brownish-gray patches, especially on the hands, feet, ankles, wrists, neck, upper chest, eyelids, inside the bend of the elbows and knees, and in infants, the face and scalp; small, raised bumps on the skin which may be weeping; skin infections; eyelid dermatitis;

cataracts; increased IgE levels; thickened, cracked, or scaly skin; and raw, sensitive, swollen skin from scratching.

[0163] The recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein may be administered by any suitable method or route known in the art, including, without limitation, orally, sublingually, buccally, topically, rectally, via inhalation, transdermally, subcutaneously, intradermally, intravenously, intraarterially, intramuscularly, intracardially, intraosseously, intraperitoneally, transmucosally, vaginally, intravitreally, intraorbitally, subretinally, intraarticularly, peri-articularly, locally, epicutaneously, or any combinations thereof. The present disclosure thus encompasses methods of delivering any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein to an individual (e.g., an individual having, or at risk of developing, Netherton Syndrome and/or atopic dermatitis).

[0164] In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein are administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject. In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered topically, transdermally, or intradermally to the subject. In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered topically or intradermally to the subject. In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered topically to the subject.

[0165] In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered once to the subject. In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered at least twice (e.g., at least 2 times, at least 3 times, at least 4 times, at least 5 times, at least 10 times, etc.) to the subject. In some embodiments, at least about 1 hour (e.g., at least about 1 hour, at least about 6 hours, at least about 12 hours, at least about 18 hours, at least about 1 day, at least about 2 days, at least about 3 days, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 15 days, at least about 20 days, at least about 30 days, at least about 40 days, at least about 50 days, at least about 60 days, at least about 70 days, at least about 80 days, at least about 90 days, at least about 100 days, at least about 120 days, etc.) pass between administrations (e.g., between the first and second administrations, between the second and third administrations, etc.). In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered one, two, three, four, five or more times per day to the subject. In some embodiments, the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations are administered to one or more affected (e.g., one or more regions of the skin displaying one or more signs or symptoms of Netherton Syndrome and/or atopic dermatitis) and/or unaffected areas of the subject.

[0166] In some embodiments, one or more portions of the skin of the subject is abraded or made more permeable prior

to treatment. Any suitable method of abrading the skin or increasing skin permeability known in the art may be used, including, for example, use of a dermal roller, repeated use of adhesive strips to remove layers of skin cells (tape stripping), scraping with a scalpel or blade, use of sandpaper, use of chemical permeation enhancers or electrical energy, use of sonic or ultrasonic energy, use of light (e.g., laser) energy, use of micron-sized needles or blades with a length suitable to pierce but not completely pass through the epidermis, etc.

VII. HOST CELLS

[0167] Certain aspects of the present disclosure relate to one or more host cells comprising any of the recombinant nucleic acids described herein. Any suitable host cell (prokaryotic or eukaryotic) known in the art may be used, including, for example: prokaryotic cells including eubacteria, such as Gram-negative or Gram-positive organisms, for example Enterobacteriaceae such as *Escherichia* (e.g., *E. coli*), *Enterobacter*, *Erminia*, *Klebsiella*, *Proteus*, *Salmonella* (e.g., *S. typhimurium*), *Serratia* (e.g., *S. marcescans*), and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis*; fungal cells (e.g., *S. cerevisiae*); insect cells (e.g., S2 cells, etc.); and mammalian cells, including monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651), human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture), baby hamster kidney cells (BHK, ATCC CCL 10), mouse Sertoli cells (TM4), monkey kidney cells (CV1 ATCC CCL 70), African green monkey kidney cells (VERO-76, ATCC CRL-1587), human cervical carcinoma cells (HELA, ATCC CCL 2), canine kidney cells (MDCK, ATCC CCL 34), buffalo rat liver cells (BRL 3A, ATCC CRL 1442), human lung cells (W138, ATCC CCL 75), human liver cells (Hep G2, HB 8065), mouse mammary tumor (MMT 060562, ATCC CCL51), TRI cells, MRC 5 cells, FS4 cells, human hepatoma line (Hep G2), Chinese hamster ovary (CHO) cells, including DHFR^r CHO cells, and myeloma cell lines such as NS0 and Sp2/0. In some embodiments, the host cell is a human or non-human primate cell. In some embodiments, the host cells are cells from a cell line. Examples of suitable host cells or cell lines may include, but are not limited to, 293, HeLa, SH-Sy5y, Hep G2, CACO-2, A549, L929, 3T3, K562, CHO-K1, MDCK, HUVEC, Vero, N20, COS-7, PSN1, VCaP, CHO cells, and the like.

[0168] In some embodiments, the recombinant nucleic acid is a herpes simplex viral vector. In some embodiments, the recombinant nucleic acid is a herpes simplex virus amplicon. In some embodiments, the recombinant nucleic acid is an HSV-1 amplicon or HSV-1 hybrid amplicon. In some embodiments, a host cell comprising a helper virus is contacted with an HSV-1 amplicon or HSV-1 hybrid amplicon described herein, resulting in the production of a virus comprising one or more recombinant nucleic acids described herein. In some embodiments, the virus is collected from the supernatant of the contacted host cell. Methods of generating virus by contacting host cells comprising a helper virus with an HSV-1 amplicon or HSV-1 hybrid amplicon are known in the art.

[0169] In some embodiments, the host cell is a complementing host cell. In some embodiments, the complementing host cell expresses one or more genes that are inactivated in any of the viral vectors described herein. In some embodiments, the complementing host cell is contacted with a

recombinant herpes virus genome (e.g., a recombinant herpes simplex virus genome) described herein. In some embodiments, contacting a complementing host cell with a recombinant herpes virus genome results in the production of a herpes virus comprising one or more recombinant nucleic acids described herein. In some embodiments, the virus is collected from the supernatant of the contacted host cell. Methods of generating virus by contacting complementing host cells with a recombinant herpes simplex virus are generally described in WO2015/009952 and/or WO2017/176336.

VIII. ARTICLES OF MANUFACTURE OR KITS

[0170] Certain aspects of the present disclosure relate to an article of manufacture or a kit comprising any of the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations described herein. In some embodiments, the article of manufacture or kit comprises a package insert comprising instructions for administering the recombinant nucleic acid, virus, medicament, and/or pharmaceutical composition or formulation to treat a SPINK deficiency (e.g., in a subject harboring a SPINK gene mutation) and/or to provide prophylactic, palliative, or therapeutic relief of one or more signs or symptoms of a disease associated with a SPINK deficiency (such as Netherton Syndrome and/or atopic dermatitis).

[0171] Suitable containers for the recombinant nucleic acids, viruses, medicaments, and/or pharmaceutical compositions or formulations may include, for example, bottles, vials, bags, tubes, and syringes. The container may be formed from a variety of materials such as glass, plastic (such as polyvinyl chloride or polyolefin), or metal alloy (such as stainless steel or hastelloy). In some embodiments, the container comprises a label on, or associated with the container, wherein the label indicates directions for use. The article of manufacture or kit may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, package inserts, and the like.

IX. ENUMERATED EMBODIMENTS

[0172] Embodiment 1: a recombinant herpes virus genome comprising one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide.

[0173] Embodiment 2: the recombinant herpes virus genome of embodiment 1, wherein the recombinant herpes virus genome is replication competent.

[0174] Embodiment 3: the recombinant herpes virus genome of embodiment 1, wherein the recombinant herpes virus genome is replication defective.

[0175] Embodiment 4: the recombinant herpes virus genome of any one of embodiments 1-3, wherein the recombinant herpes virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within one or more viral gene loci.

[0176] Embodiment 5: the recombinant herpes virus genome of any one of embodiments 1-4, wherein the recombinant herpes virus genome is selected from a recombinant herpes simplex virus genome, a recombinant varicella zoster virus genome, a recombinant human cytomegalovirus genome, a recombinant herpesvirus 6A genome, a recombinant herpesvirus 6B genome, a recombinant herpesvirus 7

genome, a recombinant Kaposi's sarcoma-associated herpesvirus genome, and any derivatives thereof.

[0177] Embodiment 6: the recombinant herpes virus genome of any one of embodiments 1-5, wherein the recombinant herpes virus genome is a recombinant herpes simplex virus genome.

[0178] Embodiment 7: the recombinant herpes virus genome of embodiment 5 or embodiment 6, wherein the recombinant herpes simplex virus genome is a recombinant type 1 herpes simplex virus (HSV-1) genome, a recombinant type 2 herpes simplex virus (HSV-2) genome, or any derivatives thereof.

[0179] Embodiment 8: the recombinant herpes virus genome of any one of embodiments 1-7, wherein the SPINK polypeptide is a human SPINK polypeptide.

[0180] Embodiment 9: the recombinant herpes virus genome of any one of embodiments 1-7, wherein the SPINK polypeptide is a Serine Protease Inhibitor Kazal-type 5 (SPINK5) polypeptide.

[0181] Embodiment 10: the recombinant herpes virus genome of any one of embodiments 1-9, wherein the SPINK polypeptide is a human SPINK5 polypeptide.

[0182] Embodiment 11: the recombinant herpes virus genome of embodiment 9 or embodiment 10, wherein the SPINK5 polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to an amino acid sequence selected from SEQ ID NOS: 7-25.

[0183] Embodiment 12: the recombinant herpes virus genome of any one of embodiments 9-11, wherein the SPINK5 polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to an amino acid sequence selected from SEQ ID NOS: 7-9.

[0184] Embodiment 13: the recombinant herpes virus genome of any one of embodiments 9-12, wherein the SPINK5 polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 7.

[0185] Embodiment 14: the recombinant herpes virus genome of any one of embodiments 5-13, wherein the recombinant herpes simplex virus genome comprises an inactivating mutation.

[0186] Embodiment 15: the recombinant herpes virus genome of embodiment 14, wherein the inactivating mutation is in a herpes simplex virus gene.

[0187] Embodiment 16: the recombinant herpes virus genome of embodiment 15, wherein the inactivating mutation is a deletion of the coding sequence of the herpes simplex virus gene.

[0188] Embodiment 17: the recombinant herpes virus genome of embodiment 15 or embodiment 16, wherein the herpes simplex virus gene is selected from Infected Cell Protein (ICP) 0, ICP4, ICP22, ICP27, ICP47, thymidine kinase (tk), Long Unique Region (UL) 41, and UL55.

[0189] Embodiment 18: the recombinant herpes virus genome of embodiment 17, wherein the recombinant herpes simplex virus genome comprises an inactivating mutation in one or both copies of the ICP4 gene.

[0190] Embodiment 19: the recombinant herpes virus genome of embodiment 17 or embodiment 18, wherein the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP22 gene.

[0191] Embodiment 20: the recombinant herpes virus genome of any one of embodiments 17-19, wherein the recombinant herpes simplex virus genome comprises an inactivating mutation in the UL41 gene.

[0192] Embodiment 21: the recombinant herpes virus genome of any one of embodiments 17-20, wherein the recombinant herpes simplex virus genome comprises an inactivating mutation in one or both copies of the ICP0 gene.

[0193] Embodiment 22: the recombinant herpes virus genome of any one of embodiments 17-21, wherein the recombinant herpes simplex virus genome comprises an inactivating mutation in the ICP27 gene.

[0194] Embodiment 23: the recombinant herpes virus genome of any one of embodiments 5-22, wherein the recombinant herpes simplex virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within one or both of the ICP4 viral gene loci.

[0195] Embodiment 24: the recombinant herpes virus genome of any one or embodiments 5-23, wherein the recombinant herpes simplex virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within the ICP22 viral gene locus.

[0196] Embodiment 25: the recombinant herpes virus genome of any one of embodiments 5-24, wherein the recombinant herpes simplex virus genome comprises the one or more polynucleotides encoding the SPINK polypeptide within the UL41 viral gene locus.

[0197] Embodiment 26: the recombinant herpes virus genome of any one of embodiments 1-25, wherein the recombinant herpes virus genome has reduced cytotoxicity when introduced into a target cell as compared to a corresponding wild-type herpes virus genome.

[0198] Embodiment 27: the recombinant herpes virus genome of embodiment 26, wherein the target cell is a human cell.

[0199] Embodiment 28: the recombinant herpes virus genome of embodiment 26 or embodiment 27, wherein the target cell is a cell of the epidermis and/or dermis.

[0200] Embodiment 29: the recombinant herpes virus genome of any one of embodiments 26-28, wherein the target cell is a keratinocyte or fibroblast.

[0201] Embodiment 30: a herpes virus comprising the recombinant herpes virus genome of any one of embodiments 1-29.

[0202] Embodiment 31: the herpes virus of embodiment 30, wherein the herpes virus is replication competent.

[0203] Embodiment 32: the herpes virus of embodiment 30, wherein the herpes virus is replication defective.

[0204] Embodiment 33: the herpes virus of any one of embodiments 30-32, wherein the herpes virus has reduced cytotoxicity as compared to a corresponding wild-type herpes virus.

[0205] Embodiment 34: the herpes virus of any one of embodiments 30-33, wherein the herpes virus is selected from a herpes simplex virus, a varicella zoster virus, a human cytomegalovirus, a herpesvirus 6A, a herpesvirus 6B, a herpesvirus 7, a Kaposi's sarcoma-associated herpesvirus, and any derivatives thereof.

[0206] Embodiment 35: the herpes virus of any one of embodiments 30-34, wherein the herpes virus is a herpes simplex virus.

[0207] Embodiment 36: the herpes virus of embodiment 34 or embodiment 35, wherein the herpes simplex virus is an HSV-1 virus, an HSV-2 virus, or any derivatives thereof.

[0208] Embodiment 37: a pharmaceutical composition comprising the recombinant herpes virus genome of any one of embodiments 1-29 and/or the herpes virus of any one of embodiments 30-36 and a pharmaceutically acceptable excipient.

[0209] Embodiment 38: the pharmaceutical composition of embodiment 37, wherein the pharmaceutical composition is suitable for topical, transdermal, subcutaneous, intradermal, oral, sublingual, buccal, rectal, vaginal, inhaled, intravenous, intraarterial, intramuscular, intracardiac, intraosseous, intraperitoneal, transmucosal, intravitreal, subretinal, intraarticular, peri-articular, local, or epicutaneous administration.

[0210] Embodiment 39: the pharmaceutical composition of embodiment 37 or embodiment 38, wherein the pharmaceutical composition is suitable for topical, transdermal, subcutaneous, intradermal, or transmucosal administration.

[0211] Embodiment 40: the pharmaceutical composition of any one of embodiments 37-39, wherein the pharmaceutical composition is suitable for topical administration.

[0212] Embodiment 41: the pharmaceutical composition of any one of embodiments 37-40, wherein the pharmaceutical composition comprises a hydroxypropyl methylcellulose gel.

[0213] Embodiment 42: the pharmaceutical composition of any one of embodiments 37-41, wherein the pharmaceutical composition comprises a phosphate buffer.

[0214] Embodiment 43: the pharmaceutical composition of any one of embodiments 37-42, wherein the pharmaceutical composition comprises glycerol.

[0215] Embodiment 44: the pharmaceutical composition of any one of embodiments 37-43, wherein the pharmaceutical composition comprises a lipid carrier.

[0216] Embodiment 45: the pharmaceutical composition of any one of embodiments 37-44, wherein the pharmaceutical composition comprises a nanoparticle carrier.

[0217] Embodiment 46: a method of enhancing, increasing, augmenting, and/or supplementing the levels of a SPINK polypeptide in one or more cells of a subject, the method comprising administering to the subject an effective amount of the herpes virus of any one of embodiments 30-36 or the pharmaceutical composition of any one of embodiments 37-45.

[0218] Embodiment 47: the method of embodiment 46, wherein the SPINK polypeptide is a SPINK5 polypeptide.

[0219] Embodiment 48: a method of enhancing, increasing, augmenting, and/or supplementing anti-inflammatory and/or anti-microbial protection of mucous epithelia in a subject in need thereof, the method comprising administering to the subject an effective amount of the herpes virus of any one of embodiments 30-36 or the pharmaceutical composition of any one of embodiments 37-45.

[0220] Embodiment 49: a method of repressing desquamation in a subject in need thereof, the method comprising administering to the subject an effective amount of the herpes virus of any one of embodiments 30-36 or the pharmaceutical composition of any one of embodiments 37-45.

[0221] Embodiment 50: a method of reducing or treating a skin barrier defect in a subject in need thereof, the method comprising administering to the subject an effective amount of the herpes virus of any one of embodiments 30-36 or the pharmaceutical composition of any one of embodiments 37-45.

[0222] Embodiment 51: the method of embodiment 50, wherein the skin barrier defect is transepithelial water loss (TEWL).

[0223] Embodiment 52: a method of providing prophylactic, palliative, or therapeutic relief of one or more signs or symptoms of Netherton Syndrome (NS) in a subject in need thereof, the method comprising administering to the subject an effective amount of the herpes virus of any one of embodiments 30-36 or the pharmaceutical composition of any one of embodiments 37-45.

[0224] Embodiment 53: the method of embodiment 52, wherein the one or more signs or symptoms of NS are selected from the group consisting of defective keratinization, a defective skin barrier, recurrent skin infections, congenital ichthyosiform erythroderma, ichthyosis linearis circumflexa, trichorrhexis invaginata, chronic skin inflammation, and any combinations thereof.

[0225] Embodiment 54: the method of any one of embodiments 46-53, wherein the subject is a human.

[0226] Embodiment 55: the method of any one of embodiments 46-54, wherein the subject's genome comprises a loss-of-function mutation in a SPINK5 gene.

[0227] Embodiment 56: the method of any one of embodiments 46-55, wherein the herpes virus or pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, periarticularly, locally, or via inhalation to the subject.

[0228] Embodiment 57: the method of any one of embodiments 46-56, wherein the herpes virus or pharmaceutical composition is administered topically, transdermally, subcutaneously, intradermally, or transmucosally to the subject.

[0229] Embodiment 58: the method of any one of embodiments 46-57, wherein the herpes virus or pharmaceutical composition is administered topically to the subject.

[0230] Embodiment 59: the method of any one of embodiments 46-58, wherein the skin of the subject is abraded prior to administration.

[0231] The specification is considered to be sufficient to enable one skilled in the art to practice the present disclosure. Various modifications of the present disclosure in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

EXAMPLES

[0232] The present disclosure will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the present disclosure. It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

Example 1: Modified Herpes Simplex Virus Vectors Encoding a Human SPINK5 Protein

[0233] To make modified, recombinant herpes simplex virus genome vectors capable of expressing SPINK5 polypeptides in a target mammalian cell (such as a human keratinocyte or fibroblast), a herpes simplex virus genome (FIG. 1A) is first modified to inactivate one or more herpes simplex virus genes. Such modifications may decrease the toxicity of the genome in mammalian cells. Next, variants of these modified/attenuated recombinant viral constructs are generated such that they carry one or more polynucleotides encoding the desired SPINK5 polypeptide. These variants include: 1) a recombinant Δ ICP4-modified HSV-1 genome comprising expression cassettes containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 7) under the control of a heterologous promoter integrated at each ICP4 locus (FIG. 1B); 2) a recombinant Δ ICP4/ Δ ICP22-modified HSV-1 genome comprising expression cassettes containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at each ICP4 locus (FIG. 1C); 3) a recombinant Δ ICP4/ Δ ICP22-modified HSV-1 genome comprising an expression cassette containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at the ICP22 locus (FIG. 1D); 4) a recombinant Δ ICP4/ Δ UL41-modified HSV-1 genome comprising an expression cassette containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at each ICP4 locus (FIG. 1E); 5) a recombinant Δ ICP4/ Δ UL41-modified HSV-1 genome comprising an expression cassette containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at the UL41 locus (FIG. 1F); 6) a recombinant Δ ICP4/ Δ ICP22/ Δ UL41-modified HSV-1 genome comprising an expression cassette containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at each ICP4 locus (FIG. 1G); 7) a recombinant Δ ICP4/ Δ ICP22/ Δ UL41-modified HSV-1 genome comprising an expression cassette containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at the UL41 locus (FIG. 1H); and 8) a recombinant Δ ICP4/ Δ ICP22/ Δ UL41-modified HSV-1 genome comprising an expression cassette containing the coding sequence (e.g., SEQ ID NO: 2) of a human SPINK5 polypeptide (e.g., SEQ ID NO: 4) under the control of a heterologous promoter integrated at the ICP22 locus (FIG. 1I).

[0234] These modified herpes simplex virus genome vectors are transfected into engineered Vero cells that are modified to express one or more herpes virus genes. These engineered Vero cells subsequently secrete into the supernatant of the cell culture a replication-defective herpes simplex virus with the modified genomes packaged therein. The supernatant is then collected, concentrated, and sterile filtered through a 5 μ m filter.

Example 2: Construction and In Vitro Analysis of an HSV Candidate Encoding Human SPINK5

[0235] The following example describes the construction of a recombinant herpes simplex virus type 1 (HSV-1) modified to express human SPINK5, and further, provides experiments showing that the recombinant virus was capable of expression functional human SPINK5 in vitro in infected human cells.

[0236] Materials and Methods

Virus Construction

[0237] The “HSV-S5” vector was generated as follows: a recombinant HSV-1 was engineered to incorporate a codon-optimized human SPINK5 expression cassette containing a heterologous promoter and polyA sequence into each of the ICP4 loci. Multiple plaques of the engineered virus putatively containing the human SPINK5 cassette were picked and screened by infection in Vero cells to test for SPINK5 expression (data not shown). The highest SPINK5 producing virus, termed “HSV-S5”, was isolated and purified.

Cell Culture

[0238] All cells were cultured at 37° C. in 5% CO₂. Immortalized normal keratinocytes were cultured in DMEM supplemented with 10% FBS and 1% HEPES buffer. Cells were grown in 6-well plates for qPCR/qRT-PCR analyses, western blotting, and the effector secretion and functionality assays. Cells were grown in 8 chamber slides for immunofluorescence.

Viral Infections

[0239] Viral aliquots were removed from the -80° C. freezer and were left to defrost under the tissue culture laminar flow hood. Multiplicity of infection was calculated from the virus titer and target cell number, the appropriate volume of virus stock was diluted in cell culture medium to a final volume of 100 μ L, and the prepared virus was incubated with the target cells for 1 hour at 37° C. After the hour-long infection, 2 mL of the appropriate cell culture medium was added to well of the 6-well plate, or in the case of 8 chamber slides, 4004, of medium was added.

qPCR/qRT-PCR Analysis

[0240] Immortalized keratinocytes were plated in 6-well plates at 7×10^5 cells/well to achieve 90-100% confluence the following day. 48 hours after mock infection or infection with HSV-S5, cells were resuspended in 3504, RLT buffer containing DTT and homogenized by passing through a QiaShredder column according to the manufacturer's protocol. DNA and RNA were isolated using an AllPrep DNA/RNA Mini kit (Qiagen) according to the manufacturer's protocol. For qPCR/qRT-PCR analysis, 50 ng of DNA or RNA were used per reaction in a total reaction volume of 25 μ L. DNA quantification was determined by qPCR analysis using a Taqman® Fast Advanced Master Mix (Applied Biosystems); RNA quantification was determined by qRT-PCR analysis using Quantabio 1-Step RT-qPCR ToughMix. All samples were run in duplicate.

Western Blotting

[0241] Immortalized keratinocytes were plated in 6-well plates at 7×10^5 cells/well to achieve 90-100% confluence the following day. 48 hours after mock infection or infection

with HSV-S5, media was aspirated, cells were recovered with gentle scraping, and cell pellets were generated by centrifugation. Cells were lysed with RIPA buffer and incubated with intermittent agitation. Next, benzonase (10U) was added to each lysate for 10 minutes at room temperature. Lysates were then mixed with 4×LDS sample buffer (Invitrogen) containing 5% 2-mercaptoethanol. Prior to loading, samples were first boiled for 10 minutes at 95° C., allowed to cool to room temperature, and then briefly centrifuged. Lysates were then run on a 4-20% Tris-Glycine gel and then transferred onto PVDF. Primary antibodies (rabbit anti-human SPINK5, R&D Systems cat. no. AF8515; rabbit anti-human GAPDH, Abcam cat. no. ab9485) were incubated with the blocked blots. Blots were then incubated with secondary antibody (goat anti-rabbit IgG-AP conjugated, Sigma) for 1 hour at room temperature, the blots were washed three times with TBS for five minutes each, and the washed blots were then incubated with AP Chromogenic substrate (Invitrogen) to allow for development of the bands.

Immunocytochemistry

[0242] Immortalized keratinocytes were plated in 8-chamber slides at 1.3×10^5 cells/well to achieve 90-100% confluence the following day. 48 hours after mock infection or infection with HSV-S5, cells were fixed with 2% paraformaldehyde, and were permeabilized with 0.2% Triton™ X-100 for five minutes. The cells were incubated with Powerblock (BioGenex) for 30 minutes, then were incubated with rabbit anti-human SPINK5 primary antibody (Novus Biologicals cat. no. NBP1-90509) overnight. Slides were subsequently incubated with Alexa Fluor® 488-conjugated secondary antibody, and mounted with ProLong Gold antifade reagent containing DAPI.

SPINK5 ELISA

[0243] The expression and secretion of human SPINK5 protein was assessed using a commercially available ELISA kit according to the manufacturer's protocol (LSBio). Briefly, standards or samples (media from cell culture) were added to the wells, unbound standard or sample was washed away, and a biotin-conjugated detection antibody was then added. An avidin-horseradish peroxidase conjugate was incubated in the wells to bind to the biotin, and a TMB substrate was then added resulting in quantitative colorimetric development. A sulfuric acid stop solution was added to terminate color development, and optical density of each well was measured at a wavelength of 450 nm using a Synergy H1 Hybrid Multi-Mode Reader.

SPINK5 Activity Assay

[0244] The inhibition of recombinant human Kallikrein 5 (R&D Systems) protease activity by HSV-S5-infected immortalized keratinocyte cell supernatants was evaluated using a modified method supplied by R&D Systems. Briefly, rhKLK5 (2 ng/μL) was preincubated with culture supernatants from HSV-S5-infected (or mock infected) cells for 5 minutes at room temperature. Following preincubation, the fluorogenic peptide Boc-VPR-AMC was added to a final concentration of 20 μM. Immediately after the addition of substrate, the cleavage of substrate (relative fluorescence units, RFU) was measured at Excitation 380 nm/Emission 460 nm using a Synergy H1 Hybrid Multi-Mode Reader every 70 seconds for 15 minutes.

Results

[0245] Immortalized normal human keratinocytes (HaCaTs) were infected with various multiplicities of infection (MOIs) of HSV-S5 ranging from 0.3 to 3.0. SPINK5 expression was evaluated by quantitative polymerase chain reaction (qPCR), quantitative reverse transcription PCR (qRT-PCR), western blot analysis, and an enzyme-linked immunosorbent assay (ELISA) 48 hours post-infection.

[0246] SPINK5 was detected in infected normal keratinocytes at an MOI as low as 0.3, and appeared to show a dose-dependent increase at both the DNA (FIG. 2A) and transcript (FIG. 2B) levels up to an MOI of 3.0. A concomitant dose-dependent increase in SPINK5 protein expression was observed in the cytoplasm of infected normal keratinocytes, as assessed by western blot (FIG. 2C).

[0247] SPINK5 expression in HSV-S5-infected keratinocytes was also evaluated by immunocytochemistry (ICC). In agreement with the nucleic acid and western blot analyses, a dose-dependent increase in SPINK5 protein expression was observed by ICC analysis (FIG. 2D). 7.36% of mock infected cells were SPINK5-positive, while 27.3%, 68.48%, and 95.20% of the HSV-S5-transduced cells were SPINK5-positive when infected at a MOI of 0.3, 1.0, and 3.0, respectively. Notably, there was no significant effect of infection on either cell morphology or viability, even at high doses of HSV-S5.

[0248] Because endogenous SPINK5 is a naturally secreted protein, it was important to show that the HSV-S5-expressed exogenous human SPINK5 could also be effectively secreted from infected cells, thus supporting HSV-S5's use in SPINK5-deficient patients. As such, SPINK5 protein levels were quantitated in cell culture medium taken from infected HaCaTs. SPINK5 was successfully secreted from infected normal keratinocytes in a dose-dependent manner (FIG. 3).

[0249] Finally, functionality of the HSV-S5-expressed human SPINK5 secreted from infected keratinocytes was confirmed using an enzymatic inhibition assay that measured SPINK5's ability to inhibit a native target of the SPINK5, the human serine protease Kallikrein 5 (KLK5). Briefly, recombinant human KLK5 was preincubated with cell culture supernatants collected from HSV-S5-infected HaCaTs (supernatant from mock-infected cells was used as a control). Following this preincubation, a synthetic, non-natural peptide substrate of human KLK5 (Boc-VPR-AMC) was added to the samples, and KLK5-mediated cleavage of the substrate was assessed in each sample. Using this experimental approach, the proteolytic activity of KLK5 (and its inhibition by SPINK5) could be directly quantitated by determining fluorescence resulting from the liberation of the previously quenched fluorescent AMC moiety of Boc-VPR-AMC after hydrolysis of the Arg-AMC amide bond by KLK5.

[0250] As expected, no inhibition of KLK5 activity was observed when preincubating the protease with cell culture supernatant obtained from mock-infected keratinocytes. In contrast, infection with HSV-S5 resulted in a dose-dependent increase in functional SPINK5 expression, as observed by increased KLK5 inhibition when the protease was pre-treated with cell culture supernatant collected from keratinocytes infected with increasing MOIs of HSV-S5 (FIG. 4).

[0251] Taken together, the data presented in this example demonstrate that an engineered herpes simplex virus can

efficiently transduce normal human keratinocytes and produce high levels of exogenous SPINK5 after transduction. Importantly, while HSV-S5 was capable of rescuing/supplementing SPINK5 expression in human cells, the vector did not induce any major toxicity in any of the studies. In addition, the data presented herein indicates that HSV-S5 is capable of inducing secretion of functional SPINK5 at therapeutically relevant levels.

Example 3: In Vivo Characterization of HSV-S5

[0252] The following example describes in vivo experiments establishing multiple routes of delivery for HSV-S5 in healthy immunocompetent animals. These studies were conducted in BALB/c mice because there are no practical disease animal models for SPINK5 deficiency, as homozygous deletion of SPINK5 is neonatal-lethal in animals.

Materials and Methods

[0253] All procedures conducted in this example were in compliance with applicable animal welfare acts and were approved by the local Institutional Animal Care and Use Committee (IACUC).

Topical HSV-S5 Administration

[0254] The backs of mice were shaved, and hair follicles were removed with a hair removal product before further manipulations. Tape stripping (using Tegaderm™) was carried out as described previously (Ekanayake-Mudiyanselage et al. *J Invest Dermatol* (1998), 111(3):517-23). After tape stripping, two sites on the backs of each mouse were treated topically with the appropriate test article. To contain the topical formulation at the treatment site, sterile plastic wells covered by a transparent adhesive dressing were adhered to the skin of the animals using surgical glue. HSV-S5 (or vehicle control) formulated in a methylcellulose gel carrier was then applied to the treatment sites via injection through the transparent adhesive dressing.

Intradermal HSV-S5 Injection

[0255] The backs of mice were shaved before further manipulations. HSV-S5 (or vehicle control) was then injected intradermally to two sites in the backs of each mouse.

Tissue Harvest

[0256] After infection and the subsequent recovery period, the animals were euthanized by CO₂ inhalation followed by cervical dislocation, and the treatment sites were removed using an 8 mm punch biopsy. One half of each biopsy was quick-frozen in liquid nitrogen for qPCR/qRT-PCR analysis, while the other half was processed for immunofluorescence analysis and H&E staining.

qPCR/qRT-PCR Analyses

[0257] Quick-frozen biopsy halves were stored at -80° C. until analysis. For processing and analysis, samples were resuspended in 350 µL RLT buffer prepared with fresh DTT according to the manufacturer's protocol (Qiagen) and were sonicated three times at 25% amplitude with intermittent incubation for 1 minute on ice. DNA and RNA extractions were performed using the Qiagen AllPrep DNA/RNA extraction kit according to the manufacturer's protocol (with the inclusion of the optional DNase treatment step for the

RNA samples). Both DNA and RNA samples were resuspended in distilled deionized RNase free water and quantified spectrophotometrically on a Synergy™ H1 microplate reader (BioTek).

[0258] Absolute quantification of SPINK5 DNA and RNA copies was performed by TaqMan Real Time PCR analysis using custom, transgene-specific primer/probe pairs. Taqman® Fast Advanced Master Mix (Applied Biosystems) was used for DNA quantification (qPCR) and Quantabio 1-Step RT-qPCR ToughMix was used for RNA quantification (qRT-PCR). All samples were run in duplicate, and copy number was determined using a standard curve derived from a dilution series of plasmid standard containing a known copy number of the SPINK5 transgene.

Immunofluorescence Staining

[0259] 5 µm sections were taken from OCT frozen tissue, mounted on slides, and air dried for up to 1 hour. The slides were then dipped in 100% methanol (MeOH) for 10 minutes at -20° C. and left to air dry. The methanol-fixed sections were rehydrated by washing 3 times in PBS (5 minutes each) at room temperature, followed by an incubation in 3% H₂O₂ for 10 minutes at room temperature, and 3 washes with PBS. The samples were then incubated with a blocking solution (Power Block) for 10 minutes at room temperature in a humidified chamber. Excess blocking solution was removed, and the sections were stained with a drop of primary antibody (Ab) solution (1:200 final dilution) prepared in antibody diluent buffer (30 to 50 µL primary Ab solution/section). The sections were incubated with the primary antibody for 16 hours at 4° C. or 1 hour at room temperature, washed three times in TBST (TBS+0.025% Triton X-100) for 5 minutes at room temperature, and secondary Ab was applied at a 1:200 dilution in antibody diluent buffer for 30 minutes at room temperature in a humidified chamber. Slides were once again washed three times with TBST, and the stained sections were mounted with mounting media (ProLong™ Gold Antifade Mountant with DAPI, Thermo-Fisher, cat. no. P36931) and covered with a coverslip. The sections were imaged after dehydration (approximately 24 hours) using an ECHO Fluorescence Microscope. The primary and secondary antibodies used in this study are presented in Table 1.

TABLE 1

antibodies used for immunofluorescence			
Antibody:	Primary/Secondary:	Vendor:	Cat. No.:
Rabbit anti-SPINK5	Primary	R&D Systems	AF8515
Mouse anti-Filaggrin	Primary	BOSTER	M01063
Anti-rabbit Alexa Fluor® 488	Secondary	Invitrogen	A11034
Anti-mouse Alexa Fluor® 594	Secondary	Abcam	Ab50120

Hematoxylin and Eosin (H&E) Staining

[0260] 5 µm sections were taken from cryopreserved tissue, mounted on slides, and air dried for up to 1 hour. The dried slides were rehydrated by soaking in double-distilled water for 2 minutes at room temperature. Sections were then incubated in Hematoxylin Gill 2x(VWR) for 2 minutes at

room temperature, followed by being dipped 2 to 3 times in acid alcohol, dipped 3 to 4 times in Blue in Ammonia water, and incubated in eosin (Eosin Y Solution 1%, VWR) for 2 minutes. Samples were rinsed 3 to 4 times with tap water between each step. The stained and rinsed sections were gradually dehydrated with ethanol (EtOH) by first rinsing twice with 95% EtOH for 2 minutes each, then twice with 100% EtOH for 2 minutes each. Sections were then cleared through three rinses with Histo-Clear for 2 minutes each, mounted with mounting media (Permount™ Mounting Medium), and covered with a coverslip. The sections were imaged approximately 24 hours after dehydration using a brightfield microscope

[0261] Results

[0262] A single dose pharmacology study was conducted in immunocompetent BALB/c mice to determine feasibility of administering human SPINK5 via topical and/or intradermal administration of HSV-S5. A total of 4 BALB/c mice were used for this study. Prior to test article administration, the backs of the mice were shaved, and hair follicles were removed using a chemical hair removal product. Next, at the sites of topical treatment, the exposed skin was tape stripped nine times to disrupt/remove the stratum corneum and 1×10^8 plaque forming units (PFUs) of HSV-S5 (or vehicle control) formulated in a gel carrier were topically administered to two regions of the tape stripped skin on each mouse. For intradermal administration, 1×10^8 PFUs of HSV-S5 (or vehicle control) were injected at two parallel sites on the back of each animal. Table 2 below provides a synopsis of the experimental design.

TABLE 2

study design and test article administration					
Group No.	N	Test Article	Route of Administration	Location; No. of Sites	Termination
1	1	Vehicle	Topical & Intradermal	Back; 2 topical, 2 intradermal	48 hours
2	3	HSV-S5	Topical & Intradermal	Back; 2 topical, 2 intradermal	48 hours

[0263] 48 hours post-administration, a full thickness 8 mm biopsy was taken from each treatment site and split in half. One half of each section was flash frozen in liquid nitrogen and subsequently processed for qPCR and qRT-PCR analysis in order to quantify SPINK5 DNA copy numbers and

transcript levels, respectively. The remaining half of each biopsy was embedded in OCT for immunofluorescence (IF). [0264] qPCR analysis of the topically treated, tape-stripped skin indicated that the engineered HSV genomes encoding the human SPINK5 transgene efficiently transduced disrupted skin of immunocompetent animals (FIG. 5A). Not only did the genomes efficiently enter the targeted tissues, but the recombinant human SPINK5 was robustly expressed after infection, as assessed by qRT-PCR analysis (FIG. 5B). Similar results were found after intradermal injection of HSV-S5 into fully intact skin at both the DNA (FIG. 5C) and RNA (FIG. 5D) levels. No SPINK5 DNA or RNA was detected in the vehicle treated tissues, indicating specificity of the assay for the HSV-S5 test article.

[0265] SPINK5 expression in cryosections harvested from topically treated animals was determined by immunofluorescent analysis using an anti-human SPINK5 antibody. To confirm that the SPINK5 expressed from HSV-S5 was correctly localized to the cornified layer of the epidermis, the samples were also counterstained for mouse Filaggrin, another structural protein localized to this region of the skin (FIG. 5E). This data demonstrated that topical application of HSV-S5 led to successful transduction of mouse skin, inducing robust expression of the encoded human transgene in the correct layer of the epidermis.

[0266] Histological evaluation of topically treated skin showed no inflammatory infiltration at the treated site, and the HSV-S5 treated skin appeared morphologically normal (comparable to vehicle-treated skin) (FIG. 5F), demonstrating the safety of this therapy.

[0267] Taken together, the in vivo data presented herein clearly demonstrates that (1) HSV-S5 successfully expresses human SPINK5 in vivo when administered topically and intradermally to immunocompetent mice, and (2) the recombinant SPINK5 is expressed in, and localizes to, the appropriate region of the epidermis. In addition, no inflammatory infiltration or gross structural changes to the skin were observed at the HSV-S5-treated sites, confirming that HSV-S5 is well tolerated. Without wishing to be bound by theory, it is believed that these data, paired with the results of the in vitro testing, provide strong support for the safe and effective use of topical or intradermal HSV-S5 for the transient and repeated delivery of human SPINK5. In addition, without wishing to be bound by theory, it is believed that HSV-S5 has the potential to be a clinically beneficial, non-invasive gene therapy candidate for the treatment of Netherton Syndrome.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 26

<210> SEQ ID NO 1

<211> LENGTH: 3285

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

atgaagatag ccacagtgtc agtgcttctg cccttggctc ttgacctcat acaagatgct      60
gccagtaaga atgaagatca ggaaatgtgc catgaatttc aggcatttat gaaaaatgga      120
aaactgttct gtccccagga taagaaattt ttccaagtc ttgatggaat aatgttcac      180
aataaatgtg ccacgtgcaa aatgatactg gaaaaagaag caaaatcaca gaagagggcc      240

```

-continued

aggcatttag caagagctcc caaggctact gcccacacag agctgaattg tgatgatttt	300
aaaaaaggag aaagagatgg ggattttatc tgcctgatt attatgaagc tgtttgaggc	360
acagatggga aaacatatga caacagatgt gcaactgtgtg ctgagaatgc gaaaaccggg	420
tcccaaatg gtgtaaaaag tgaaggggaa tgtaagagca gtaatccaga gcaggatgta	480
tgcaagtgtt ttcggccctt tgtagagat ggaagacttg gatgcacaag ggaaaatgat	540
cctgttcttg gtcctgatgg gaagacgcat ggcaataagt gtgcaatgtg tgctgagctg	600
tttttaaaag aagctgaaaa tgccaagcga gagggtgaaa ctagaattcg acgaaatgct	660
gaaaaggatt tttgcaagga atatgaaaa caagtgagaa atggaaggct tttttgtaca	720
cgggagagtg atccagtcoc tggccctgac ggcaggatgc atggcaacaa atgtgccctg	780
tgtgtgaaa ttttcaagca gcgtttttca gaggaaaaa gtaaaacaga tcaaaatttg	840
ggaaaagctg aagaaaaaac taaagttaa agagaaattg tgaaactctg cagtcaatat	900
caaatcagg caaagaatgg aatacttttc tgtaccagag aaaatgaccc tattcgtggt	960
ccagatggga aaatgcatgg caacttgtgt tccatgtgtc aagcctactt ccaagcagaa	1020
aatgaagaaa agaaaaaggc tgaagcacga gctagaaaa aaagagaatc tggaaaagca	1080
acctcatatg cagagctttg cagtgaatat cgaaaagcttg tgaggaacgg aaaacttgct	1140
tgaccagag agaacgatcc tatccagggc ccagatggga aagtgcattg caacacctgc	1200
tccatgtgtg aggtcttctt ccaagcagaa gaagaagaaa agaaaaagaa ggaaggtaaa	1260
tcaagaaaca aaagacaatc taagagtaca gcttctttg aggagttgtg tagtgaatac	1320
cgcaaatcca ggaaaaaacg acggttttt tgcaccagag agaatgaccc catccagggc	1380
ccagatggaa aaatgcatgg caacacctgc tccatgtgtg aggccttctt tcaacaagaa	1440
gaaagagcaa gagcaaaggc taaaagagaa gctgcaaagg aatctgcag tgaatttcgg	1500
gaccaagtga ggaatggaac acttatatgc accagggagc ataactcgtt ccgtggccca	1560
gatggcaaaa tgcattgaaa caagtgtgcc atgtgtgcca gtgtgttcaa acttgaagaa	1620
gaagagaaga aaaatgataa agaagaaaaa gggaaagtgc aggctgaaaa agttaagaga	1680
gaagcagttc aggagctgtg cagtgaatat cgtcattatg tgaggaatgg acgactcccc	1740
tgtaccagag agaatgatcc tattgagggt ctatagggga aaatccacgg caacacctgc	1800
tccatgtgtg aagccttctt ccagcaagaa gcaaaagaaa aagaaagagc tgaaccacga	1860
gcaaaagtca aaagagaagc tgaagaggag acatgcatg aatttcggag acttttgcaa	1920
aatggaaaac ttttctgcac aagagaaaat gatcctgtgc gtggcccaga tggcaagacc	1980
catggcaaca agtgtgcat gtgtaaggca gtcttccaga aagaaaatga ggaagaaaag	2040
aggaagaag aggaagatca gagaaatgct gcaggacatg gttccagtgg tggtaggaga	2100
ggaaacactc aggacgaatg tgctgagat cggaacaaa tgaaaaatgg aagactcagc	2160
tgtactcggg agagtgatcc tgtacgtgat gctgatggca aatcgtacaa caatcagtg	2220
accatgtgta aagcaaaatt ggaagagaa gcagagagaa aaaatgagta ttctcgtcc	2280
agatcaaatg ggactggatc agaatcaggg aaggatacat gtgatgagtt tagaagccaa	2340
atgaaaaatg gaaaactcat ctgcaactga gaaagtgacc ctgtccgggg tccagatggc	2400
aagacacatg gcaataagtg tactatgtgt aagaaaaaac tggaaaagga agcagctgaa	2460
aaaaaaaaga aagaggatga agacaggagc aatacaggag aaaggagcaa tacaggagaa	2520

-continued

```

aggagcaatg acaaagagga tctgtgtcgt gaatttcgaa gcatgcagag aaatggaaaag 2580
cttatctgca ccagagaaaa taacctgtt cgaggccat atggcaagat gcacatcaat 2640
aaatgtgcta tgtgtcagag catctttgat cgagaagcta atgaaagaaa aaagaaagat 2700
gaagagaaat caagtagcaa gccctcaaat aatgcaaagg accagtgcag acaggttcag 2760
aatgaagcgg aggatgcaaa atttagacia cctgggctt ccttggcctc tgttgccagg 2820
atgagtacag atgagtgcag tgaatttcga aactataaa ggaacaatga actcatctgc 2880
cctagagaga atgaccagc gcacgggtgct gatggaaagt tctatacaaa caagtgtac 2940
atgtgcagag ctgtctttct aacagaagct ttgaaaggg caaagcttca agaaaagcca 3000
tcccatgtta gagcttctca agaggaagac agcccagact ctttcagttc tctggattct 3060
gagatgtgca aagactaccg agtattgcc aggataggtt atctttgtcc aaaggattta 3120
aagcctgtct gtggtgacga tggccaaacc tacaacaatc cttgcatgct ctgtcatgaa 3180
aacctgatac gccaaacaaa tacacacatc cgcagtacag ggaagtgtga ggagagcagc 3240
acccagga ccaccgcagc cagcatgcc cegtctgacg aatga 3285

```

```

<210> SEQ ID NO 2
<211> LENGTH: 3285
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 2

```

```

atgaagatcg ccaccgtgtc tgtgtgtcgt cctctggctc tgtgctgat tcaggatgcc 60
gccagcaaga acgaggacca agagatgtgc cagcagttcc aggcctttat gaagaacggc 120
aagctgttct gccctcaaga caagaagttc ttccagagcc tggacggcat catgttcatc 180
aacaagtgcg ccacatgcaa gatgatcctg gaaaaagagg ccaagagcca gaagcgggcc 240
agacaccttg ctagagcccc taaagccacc gctcctaccg agctgaactg cgacgaactc 300
aagaaaggcg agcgcgacgg cgacttcatc tgccccgatt attacgaggc cgtgtgcggc 360
accgatggca agacctacga taacagatgc gccctgtgcy ccgagaatgc caagacagga 420
tctcagatcg gctgaaagtc tgaggcgag tgcaagagca gcaacccca gcaggatgtg 480
tgcagcgctc tcagaccctt cgtcagagat ggcagactgg gctgcaccag agaaaacgat 540
cctgtgctgg gccctgacgg caagacacac gaaaacaagt gtgctatgtg tgccgagctg 600
ttcctgaaag aggccgaaaa cgccaagaga gaggcgaga cacggatcag aagaaacgcc 660
gagaaggact tctgcaaaga atacgagaag caagtgcgga acggacgctt gttctgtacc 720
cgggaaagcg atccagtcag aggccccgat ggaagaatgc acggcaacaa atgcgcactg 780
tgtgtgaga tcttcaagca gcggttcagc gaggaaaaca gcaagaccga ccagaacctg 840
ggcaaagcgg aagaaaagac caaagtgaag cgcgagatcg tgaagctgtg cagccagtac 900
cagaaccagg ccaagaatgg catcctgttc tgcacacgcy agaacgaccc catcagagga 960
cccgacggaa agatgcatgg aaacctgtgc agcatgtgcc aggcctactt ccaggccgag 1020
aacgaagaga agaagaaggc cgaagccagg gccagaaaca agagagaaag cggcaaggcc 1080
accagctacg ccgaactgtg tagcagtagc agaaagctcg tgccgaaatg caagctggcc 1140
tgcactaggg aaaatgaccc tattcaggga ccagacggca aggtgcacgg gaatacctgt 1200

```

-continued

tccatgtgcg aggtgttctt tcaggcggag gaagaggaaa aaaagaagaa agagggcaag	1260
tcccggaaaca agcggcagag caagagcacc gccagcttcg aggaactgtg ctccaggtac	1320
cggaaagtcca gaaagaatgg ccggcttttc tgcacgagag agaatgatcc catccagggg	1380
cctgatggca aatgcatgg caatacttgc tctatgtgcg aggcattttt ccagcaagag	1440
gaacgggcca gagccaaggc caaaagagag gccgccaag agatctgcag cgagttagg	1500
gaccaagtcc gcaacggcac cctgatctgc accagggaac acaaccctgt cagagggcca	1560
gacgggaaaa tgcacgggaa caagtgtgcc atgtgcgcct ccgtgttcaa gctggaagaa	1620
gaagaaaaga aaaacgacaa agaagagaag ggaaaagtcg aagccgagaa agtgaagagg	1680
gaagccgtgc aagaactctg ttctgagtac cggcactacg tccggaacgg cagactgcct	1740
tgtactcggg aaaacgaccc aatcgaagga ctggacggga agatccatgg caacacctgt	1800
agtatgtgcg aggcattctt tcaacaagaa gccaaagaaa aagagcgggc cgagcctcgg	1860
gccaaagtca aaagagaagc tgagaagag acatgcgacg agttccggcg gctgctgcag	1920
aatgaaaac tcttctgtac acgtgaaaat gatcccgtgc ggggaccaga tggaaagacc	1980
catgggaaca aatgcgctat gtgcaaggcc gtgttccaaa aagagaatga ggaacgcaag	2040
cggaaagagg aagaagatca gcggaacgcc gctggccacg gatctagtgg cggaggtggt	2100
ggaaataccc aggacgagtg tgccgaatac cgggaacaga tgaagaatgg aaggctgagc	2160
tgcacacggg aatctgacct cgttagagat gccgacggca agtcctacaa caaccagtgc	2220
accatgtgta aagccaagct cgagcgcgag gccgagcggg agaatgagta cagcagaagc	2280
cggctctaacg gcaccggctc tgagtctggc aaggatacct gtgacgagtt cagatcccag	2340
atgaagaacg gaaactcat ctgcacaaga gaatcagacc cagtcggggg acctgacggg	2400
aaaactcatg gcaacaagtg cacaatgtgc aaagagaaac tggaaacgca ggctgccgag	2460
aagaaaaaga aagaggacga ggatcgggag aacaccggcg agagatctaa taccggcgag	2520
cggagcaatg acaagagga cctctgccgg gaattccggt ccatgcagag aaacggaaaag	2580
ctgatatgta cgcgcgagaa caatcccgtt cggggccctt atgggaagat gcacattaac	2640
aaatgtgcaa tgtgccagag catcttcgac agagaggcca acgaaacgca aaagaaggat	2700
gaagagaagt ccagcagcaa gccacgcaac aacgccaagg accagtgcag acaggtgcag	2760
aacgaggctg aggacgcaa gtttagacag cccggtagaa gcctggcctc tgtggccaga	2820
atgagcaccg atgagtgcag cgaattccgg aactatatcc ggaacaacga gctgatctgt	2880
ccgcgagaga atgaccctgt gcatggcgcc gatggaagt tctacaccaa caagtgttat	2940
atgtgcagag ccgtctttct gacagaggcc ctggaaagag ccaagctgca agagaagcct	3000
tctcacgtgc gggcctctca agaagaggac agccctgata gcttcagcag cctggacagc	3060
gagatgtgca aggactacag agtgcgtgcc cggatcggct atctgtgccc caaggatctg	3120
aagcctgtgt gtggcgacga cggccagaca tacaacaatc cctgcatgct gtgtcaagag	3180
aacctgatcc ggcagaccaa cacacacatc cggtcaccgg gcaagtgcga agagtctagc	3240
acacctggaa caaccgccc cctctatgcct cctagcagc aatga	3285

<210> SEQ ID NO 3

<211> LENGTH: 2751

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 3

```

atgaagatag ccacagtgtc agtgcctctg cccttggtc tttgectcat acaagatgct    60
gccagtaaga atgaagatca ggaatgtgc catgaatttc aggcatztat gaaaaatgga    120
aaactgttct gtccccagga taagaaattt tttcaaagtc ttgatggaat aatgttcate    180
aataaatgtg ccacgtgcaa aatgatactg gaaaaagaag caaaatcaca gaagagggcc    240
aggcatntag caagagctcc caaggctact gcccacacag agctgaattg tgatgatttt    300
aaaaaaggag aaagagatgg ggattttatc tgtcctgatt attatgaagc tgtttgtggc    360
acagatggga aacatatga caacagatgt gcaactgtgtg ctgagaatgc gaaaaccggg    420
tcccaaatgt gtgtaaaaag tgaaggggaa tgtaagagca gtaatccaga gcaggatgta    480
tgcagtgtct ttcggccctt tgttagagat ggaagacttg gatgcacaag ggaaaatgat    540
cctgttcttg gtcctgatgg gaagacgcat ggcaataagt gtgcaatgtg tgctgagctg    600
tttttaaaag aagctgaaaa tgccaagcga gagggtgaaa ctagaattcg acgaaatgct    660
gaaaaggatt tttgcaagga atatgaaaaa caagtgagaa atggaaggct tttttgtaca    720
cgggagagtg atccagtcog tggccctgac ggcaggatgc atggcaacaa atgtgccttg    780
tgtgtctgaa tttcaagca gcgtttttca gaggaaaaca gtaaaacaga tcaaaatttg    840
ggaaaagctg aagaaaaaac taaagttaaa agagaaattg tgaactctg cagtcaatat    900
caaaatcagg caaagaatgg aatacttttc tgtaccagag aaaatgacct tattcgtggt    960
ccagatggga aaatgcatgg caacttgtgt tccatgtgtc aagcctactt ccaagcagaa   1020
aatgaagaaa agaaaaaggc tgaagcacga gctagaaaca aaagagaatc tggaaaagca   1080
acctcatatg cagagctttg cagtgaatat cgaaaagcttg tgaggaacgg aaaacttgct   1140
tgcaccagag agaacgatcc tatccagggc ccagatggga aagtgcattg caacacctgc   1200
tccatgtgtg aggtcttctt ccaagcagaa gaagaagaaa agaaaaagaa ggaaggtaaa   1260
tcaagaaaca aaagacaatc taagagtaca gcttcctttg aggagttgtg tagtgaatac   1320
cgcaaatcca ggaaaaacgg acggcttttt tgcaccagag agaatgacct catccagggc   1380
ccagatggaa aaatgcatgg caacacctgc tccatgtgtg aggccttctt tcaacaagaa   1440
gaaagagcaa gagcaaaggc taaaagagaa gctgcaaagg aaatctgcag tgaatttcgg   1500
gaccaagtga ggaatggaac acttatatgc accagggagc ataatcctgt ccgtggccca   1560
gatggcaaaa tgcattgaaa caagtgtgcc atgtgtgcca gtgtgttcaa acttgaagaa   1620
gaagagaaga aaaatgataa agaagaaaaa gggaaagtgc aggctgaaaa agttaagaga   1680
gaagcagttc aggagctgtg cagtgaatat cgtcattatg tgaggaatgg acgactcccc   1740
tgtaccagag agaatgatcc tattgagggt ctagatggga aaatccacgg caacacctgc   1800
tccatgtgtg aagccttctt ccagcaagaa gcaaaagaaa aagaaagagc tgaaccagaa   1860
gcaaaagtca aaagagaagc tgaaaaggag acatgcatg aatttcggag acttttgcaa   1920
aatggaaaac ttttctgcac aagagaaaat gatcctgtgc gtggcccaga tggcaagacc   1980
catggcaaca agtgtgcat gtgtaaggca gtcttcagaa aagaaaatga ggaagaaaag   2040
aggaaagaag aggaagatca gagaaatgct gcaggacatg gttccagtgg tggaggagga   2100
ggaaacactc aggacgaatg tgctgagat cggaacaaa tgaaaaatgg aagactcagc   2160
tgtactcggg agagtgatcc tgtacgtgat gctgatggca aatcgtacaa caatcagttg   2220

```

-continued

```

accatgtgta aagcaaaatt ggaaagagaa gcagagagaa aaaatgagta ttctcgctcc 2280
agatcaaatg ggactggatc agaatcaggg aaggatacat gtgatgagtt tagaagccaa 2340
atgaaaaatg gaaaactcat ctgcaactcga gaaagtgacc ctgtccgggg tccagatggc 2400
aagacacatg gcaataagtg tactatgtgt aaggaaaaac tggaaagga agcagctgaa 2460
aaaaaaaaga aagaggatga agacaggagc aatacaggag aaaggagcaa tacaggagaa 2520
aggagcaatg acaagagga tctgtgtcgt gaatttcgaa gcatgcagag aaatggaaag 2580
cttatctgca ccagagaaaa taacctgtt cgaggcccat atggcaagat gcacatcaat 2640
aaatgtgcta tgtgtcagag catctttgat cgagaagcta atgaaagaa aaagaaagat 2700
gaagaaaaat caagtagcaa gccctcaaat aatgcaaagg ttatttatta a 2751

```

```

<210> SEQ ID NO 4
<211> LENGTH: 2751
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 4

```

```

atgaagatcg ccaccgtgtc tgtgctgctg cctctggctc tgtgcctgat tcaggatgcc 60
gccagcaaga acgaggacca agagatgtgc cagcagttcc aggcctttat gaagaacggc 120
aagctgttct gccctcaaga caagaagttc ttccagagcc tggacggcat catgttcatc 180
aacaagtgcg ccacatgcaa gatgatcctg gaaaaagagg ccaagagcca gaagcgggcc 240
agacaccttg ctagagcccc taaagccacc gctcctaccg agctgaactg cgacgacttc 300
aagaaaggcg agcgcgacgg cgacttcatc tgccccgatt attacgaggc cgtgtgcggc 360
accgatggca agacctacga taacagatgc gccctgtgcg ccgagaatgc caagacagga 420
tctcagatcg gcgtgaagtc tgagggcgag tgcaagagca gcaaccccga gcaggatgtg 480
tgcagcgctc tcagaccctt cgtcagagat ggcagactgg gctgcaccag agaaaacgat 540
cctgtgctgg gccctgacgg caagacacac gaaacaagt gtgctatgtg tgccgagctg 600
ttcctgaaag aggccgaaaa cgccaagaga gagggcgaga cacggatcag aagaaacgcc 660
gagaaggact tctgcaaaga atacgagaag caagtgcgga acggacgctt gttctgtacc 720
cgggaaagcg atccagtcag aggcctccgat ggaagaatgc acggcaacaa atgcgactg 780
tgtgtgagga tcttcaagca gcggttcagc gaggaaaaa gcaagaccga ccagaacctg 840
ggcaaaagcc aagaaaagac caaagtgaag cgcgagatcg tgaagctgtg cagccagtac 900
cagaaccagg ccaagaatgg catcctgttc tgcacacgcg agaacgacct catcagagga 960
cccagcgaaa agatgcattg aaacctgtgc agcatgtgcc aggcctactt ccaggccgag 1020
aacgaagaga agaagaaggc cgaagccagg gccagaaaca agagagaaag cggcaaggcc 1080
accagctacg ccgaactgtg tagcgagtac agaaagctcg tgcggaatgg caagctggcc 1140
tgcactaggg aaaatgacct tattcaggga ccagacggca aggtgcacgg gaatacctgt 1200
tccatgtgcg aggtgttctt tcaggccgag gaagaggaaa aaaagaagaa agagggcaag 1260
tcccggaaaca agcggcagag caagagcacc gccagcttcg aggaactgtg ctccgagtac 1320
cggaagtcca gaaagaatgg ccgcttttc tgcacgagag agaatgatcc catccagggg 1380
cctgatggca aaatgcatgg caatactgct tctatgtgcg aggcattttt ccagcaagag 1440

```

-continued

gaacgggcca gagccaaggc caaaagagag gccgccaag agatctgcag cgagtttagg	1500
gaccaagtcc gcaacggcac cctgatctgc accagggaac acaaccctgt cagagggcca	1560
gacgggaaaa tgcacgggaa caagtgtgcc atgtgcgcct ccgtgttcaa gctggaagaa	1620
gaagaaaaga aaaacgacaa agaagagaag ggaaaagtgc aagccgagaa agtgaagagg	1680
gaagccgtgc aagaactctg ttctgagtac cggcactacg tccggaacgg cagactgcct	1740
tgtactcggg aaaacgacct aatcgaagga ctggacggga agatccatgg caaacctgt	1800
agtatgtgcg aggattctt tcaacaagaa gccaaagaaa aagagcgggc cgagcctcgg	1860
gccaaagtca aaagagaagc tgagaaagag acatgcgcag agttccggcg gctgctgcag	1920
aatggaaaac tcttctgtac acgtgaaaat gatcccgtgc ggggaccaga tggaaagacc	1980
catgggaaca aatcgcctat gtgcaaggcc gtgttccaaa aagagaatga ggaacgcaag	2040
cggaaagagg aagaagatca gcggaacgcc gctggccacg gatctagtgg cggaggtggt	2100
ggaaataccc aggacgagtg tgccaatac cgggaacaga tgaagaatgg aaggctgagc	2160
tgcacacggg aatctgacct cgttagagat gccgacggca agtcctacaa caaccagtgc	2220
accatgtgta aagccaagct cgagcgcgag gccgagcggga agaatagta cagcagaagc	2280
cggctaacg gcaccggctc tgagtctggc aaggatacct gtgacgagtt cagatccag	2340
atgaagaacg ggaactcat ctgcacaaga gaatcagacc cagtcgggg acctgacggg	2400
aaaactcatg gcaacaagtg cacaatgtgc aaagagaaac tggaaacgca ggctgcccag	2460
aagaaaaaga aagaggacga ggtcgggac aacaccggcg agagatctaa taccggcgag	2520
cggagcaatg acaaaagga cctctgcccg gaattccggt ccatgcagag aaacggaaa	2580
ctgatatgta cgcgcgagaa caatccggtt cggggccctt atgggaagat gcacattaac	2640
aatgtgcaa tgtccagag catcttcgac agagaggcca acgaacgcaa aaagaaggat	2700
gaagagaagt ccagcagcaa gccacgcaac aacgctaaag tgatctactg a	2751

<210> SEQ ID NO 5

<211> LENGTH: 3195

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

atgaagatag ccacagtgtc agtgcctctg cccttggctc tttgcctcat acaagatgct	60
gccagtaaga atgaagatca ggaaatgtgc catgaatttc aggcatttat gaaaaatgga	120
aaactgttct gtccccagga taagaaattt tttcaaagtc ttgatggaat aatgttcatc	180
aataaatgtg ccacgtgcaa aatgatactg gaaaaagaag caaaatcaca gaagagggcc	240
aggcatttag caagagctcc caaggctact gcccacacag agctgaattg tgatgatttt	300
aaaaaaggag aaagagatgg ggattttatc tgtcctgatt attatgaagc tgtttgtggc	360
acagatggga aaacatatga caacagatgt gcaactgtgtg ctgagaatgc gaaaaccggg	420
tcccaaatgt gtgtaaaaag tgaaggggaa tgtaagagca gtaatccaga gcaggatgta	480
tgcagtgtct ttcggccctt tgttagagat ggaagacttg gatgcacaag ggaaaatgat	540
cctgttcttg gtccctgatg gaagacgcat ggcaataagt gtgcaatgtg tgctgagctg	600
tttttaaaag aagctgaaaa tgccaagcga gagggtgaaa ctagaattcg acgaaatgct	660
gaaaaggatt tttgcaagga atatgaaaaa caagtgagaa atggaaggct tttttgtaca	720

-continued

cgggagagtg atccagtcg tggccctgac ggcaggatgc atggcaacaa atgtgcctcg	780
tgtgctgaaa ttttcaagca gcgtttttca gaggaaaaca gtaaacaga tcaaaatttg	840
ggaaaagctg aagaaaaaac taaagttaa agagaaattg tgaactctg cagtcaatat	900
caaaatcagg caaagaatgg aatacttttc tgtaccagag aaaatgacct tattcgtggt	960
ccagatggga aaatgcatgg caacttgtgt tccatgtgtc aagcctactt ccaagcagaa	1020
aatgaagaaa agaaaaaggc tgaagcacga gctagaaaca aaagagaatc tggaaaagca	1080
acctcatatg cagagctttg cagtgaatat cgaaagcttg tgaggaaagg aaaacttget	1140
tgcaccagag agaacgatcc tatccagggc ccagatggga aagtgcattg caacacctgc	1200
tccatgtgtg aggtcttctt ccaagcagaa gaagaagaaa agaaaaagaa ggaaggtaaa	1260
tcaagaaaca aaagacaatc taagagtaca gcttctttg aggagttgtg tagtgaatac	1320
cgcaaatcca gaaaaaacgg acggttttt tgcaccagag agaatgacct catccagggc	1380
ccagatggaa aaatgcatgg caacacctgc tccatgtgtg aggccttctt tcaacaagaa	1440
gaaagagcaa gagcaaaggc taaaagagaa gctgcaaagg aaatctgcag tgaatttcgg	1500
gaccaagtga ggaatggaac acttatatgc accaggagc ataactctgt ccgtggccca	1560
gatggcaaaa tgcattgaaa caagtgtgcc atgtgtgcca gtgtgttcaa acttgaagaa	1620
gaagagaaga aaaatgataa agaagaaaaa gggaaagtgc aggtgaaaa agttaagaga	1680
gaagcagttc aggagctgtg cagtgaatat cgtcattatg tgaggaaagg acgactcccc	1740
tgtaccagag agaatgatcc tattgagggt ctgatggga aaatccacgg caacacctgc	1800
tccatgtgtg aagccttctt ccagcaagaa gcaaaagaaa aagaaagagc tgaaccaga	1860
gcaaaagtca aaagagaagc tgaagaggag acatgcatg aatttcggag acttttgcaa	1920
aatgaaaaac tttctgcac aagagaaaat gatcctgtgc gtggcccaga tggcaagacc	1980
catggcaaca agtgtgcat gtgtaaggca gtcttcaga aagaaatga ggaagaaaag	2040
agaaagaag aggaagatca gagaaatgct gcaggacatg gttccagtgg tggaggagga	2100
ggaaacactc aggacgaatg tctgagatc cgggaacaaa tgaaaaatgg aagactcagc	2160
tgtactcggg agagtgatcc tgtacgtgat gctgatggca aatcgtacaa caatcagttg	2220
accatgtgta aagcaaaatt ggaagagaa gcagagagaa aaaatgagta ttctcgtccc	2280
agatcaaatg ggactggatc agaatcaggg aaggatacat gtgatgagtt tagaagccaa	2340
atgaaaaatg gaaaactcat ctgcaactga gaaagtgacc ctgtccgggg tccagatggc	2400
aagacacatg gcaataagtg tactatgtgt aagaaaaaac tggaaaggga agcagctgaa	2460
aaaaaaaaaga aagaggatga agcaggagc aatacaggag aaaggagcaa tacaggagaa	2520
aggagcaatg acaaaagga tctgtgtcgt gaatttcgaa gcatgcagag aaatggaaag	2580
cttatctgca ccagagaaaa taacctgtt cgaggccat atggcaagat gcacatcaat	2640
aaatgtgcta tgtgtcagag catctttgat cgagaagcta atgaaagaaa aaagaaagat	2700
gaagagaaat caagtagcaa gccctcaaat aatgcaaagg atgagtgagc tgaatttcga	2760
aactatataa ggaacaatga actcatctgc cctagagaga atgacctagt gcacggtgct	2820
gatgaaagt tctatacaaa caagtgtac atgtgcagag ctgtcttctt aacagaagct	2880
ttgaaaggg caaagcttca agaaaagcca tccatgtta gagcttctca agaggaagac	2940
agcccagact ctttcagttc tctggattct gagatgtgca aagactaccg agtattgccc	3000

-continued

```

aggataggtt atctttgtcc aaaggattta aagcctgtct gtggtgacga tggccaaacc 3060
tacaacaatc cttgcatgct ctgtcatgaa aacctgatac gccaaacaaa tacacacatc 3120
cgcagtacag ggaagtgtga ggagagcagc accccaggaa ccaccgcagc cagcatgccc 3180
ccgtctgacg aatga 3195

```

```

<210> SEQ ID NO 6
<211> LENGTH: 3195
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 6

```

```

atgaagatcg ccaccgtgtc tgtgctgctg cctctggctc tgtgcctgat tcaggatgcc 60
gccagcaaga acgaggacca agagatgtgc cacgagttcc aggcctttat gaagaacggc 120
aagctgttct gccctcaaga caagaagttc ttccagagcc tggacggcat catgttcatc 180
aacaagtgcg ccacatgcaa gatgatcctg gaaaagagg ccaagagcca gaagcgggcc 240
agacaccttg ctagagcccc taaagccacc gctcctaccg agctgaactg cgacgacttc 300
aagaaaggcg agcgcgacgg cgacttcatc tgccccgatt attacgaggc cgtgtgcggc 360
accgatggca agacctacga taacagatgc gccctgtgcg ccgagaatgc caagacagga 420
tctcagatcg gcgtgaagtc tgagggcgag tgcaagagca gcaacccca gaggatgtg 480
tgcagcgctc tcagaccctt cgtcagagat ggcagactgg gctgcaccag agaaaacgat 540
cctgtgctgg gccctgacgg caagacacac gaaaacaagt gtgctatgtg tgccgagctg 600
ttcctgaaag aggccgaaaa cgccaagaga gagggcgaga cacggatcag aagaaacgcc 660
gagaaggact tctgcaaaga atacgagaag caagtgcgga acggacgctt gttctgtacc 720
cgggaaagcg atccagtcag aggccccgat ggaagaatgc acggcaacaa atgcgcactg 780
tgtgctgaga tcttcaagca gcggttcagc gaggaaaaca gcaagaccga ccagaacctg 840
ggcaaaagccg aagaaaagac caaagtgaag cgcgagatcg tgaagctgtg cagccagtac 900
cagaaccagg ccaagaatgg catcctgttc tgcacacgcy agaacgaccc catcagagga 960
cccagcggaa agatgcatgg aaacctgtgc agcatgtgcc aggcctactt ccaggccgag 1020
aacgaagaga agaagaaggc cgaagccagg gccagaaaaca agagagaaag cggcaaggcc 1080
accagctacg ccgaactgtg tagcaggtac agaaagctcg tgcggaatgg caagctggcc 1140
tgcactaggg aaaatgaccc tattcaggga ccagacggca aggtgcacgg gaatacctgt 1200
tccatgtgcy aggtgttctt tcaggccgag gaagaggaaa aaaagaagaa agagggcaag 1260
tcccggaaaca agcggcagag caagagcacc gccagcttcg aggaactgtg ctccgagtac 1320
cggaagtcca gaaagaatgg ccggcttttc tgcacgagag agaatgatcc catccagggg 1380
cctgatggca aaatgcatgg caatacttgc tctatgtgcy aggcattttt ccagcaagag 1440
gaacggggcca gagccaaggc caaaagagag gccgccaaag agatctgcag cgagtttagg 1500
gaccaagtcc gcaacggcac cctgatctgc accagggaac acaaccctgt cagagggcca 1560
gacgggaaaa tgcacgggaa caagtgtgcc atgtgcccct ccgtgttcaa gctggaagaa 1620
gaagaaaaga aaaacgacaa agaagagaag ggaaaagtcg aagccgagaa agtgaagagg 1680
gaagccgtgc aagaactctg ttctgagtac cggcactacg tccggaacgg cagactgcct 1740

```

-continued

```

tgtactcggg aaaacgacct aatcgaagga ctggacggga agatccatgg caacacctgt 1800
agtatgtgag aggcattctt tcaacaagaa gccaaagaaa aagagcgggc cgagcctcgg 1860
gccaaagtca aaagagaagc tgagaagag acatgacgacg agttccggcg gctgctgcag 1920
aatggaaaac tcttctgtac acgtgaaaat gatcccgtgc ggggaccaga tggaaagacc 1980
catgggaaca aatgcgctat gtgcaaggcc gtgttccaaa aagagaatga ggaacgcaag 2040
cggaaagagg aagaagatca gcggaacgcc gctggccacg gatctagtgg cggaggtggt 2100
ggaaataccc aggacgagtg tgccgaatac cgggaacaga tgaagaatgg aaggctgagc 2160
tgcacacggg aatctgacct cgttagagat gccgacggca agtcctacaa caaccagtgc 2220
accatgtgta aagccaagct cgagcgcgag gccgagcggc agaagtagta cagcagaagc 2280
cgggtctaacg gcaccggctc tgagtctggc aaggatacct gtgacgagtt cagatcccag 2340
atgaagaacg gaaactcat ctgcacaaga gaatcagacc cagtcggggg acctgacggg 2400
aaaactcatg gcaacaagtg cacaatgtgc aaagagaaac tggaaacgca ggctgccgag 2460
aagaaaaaga aagaggacga ggatcgggag aacaccggcg agagatctaa taccggcgag 2520
cggagcaatg acaagagga cctctgccgg gaattccggt ccatgcagag aaacggaaaag 2580
ctgatatgta cgcgcgagaa caatcccgtt cggggcccctt atgggaagat gcacattaac 2640
aaatgtgcaa tgtgccagag catcttcgac agagaggcca acgaacgcaa aaagaaggat 2700
gaagagaagt ccagcagcaa gccagcaac aacgccaagg acgaatgcag cgaattccgg 2760
aactatatcc ggaacaacga gctgatctgt ccgagagaga atgaccctgt gcatggcgcc 2820
gatggaaagt tctacaccaa caagtgttat atgtgcagag ccgtctttct gacagaggcc 2880
ctggaaagag ccaagctgca agagaagcct tctcacgtgc gggcctctca agaagaggac 2940
agccctgata gcttcagcag cctggacagc gagatgtgca aggactacag agtgcctgcc 3000
cggatcggct atctgtgccc caaggatctg aagcctgtgt gtggcgacga cggccagaca 3060
tacaacaatc cctgcatgct gtgtcacgag aacctgatcc ggcagaccaa cacacacatc 3120
cggtcaccag gcaagtgcga agagtctagc acacctggaa caaccgccgc ctctatgcct 3180
cctagcgcag aatga 3195

```

<210> SEQ ID NO 7

<211> LENGTH: 1094

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

Met Lys Ile Ala Thr Val Ser Val Leu Leu Pro Leu Ala Leu Cys Leu
1           5           10          15
Ile Gln Asp Ala Ala Ser Lys Asn Glu Asp Gln Glu Met Cys His Glu
20          25          30
Phe Gln Ala Phe Met Lys Asn Gly Lys Leu Phe Cys Pro Gln Asp Lys
35          40          45
Lys Phe Phe Gln Ser Leu Asp Gly Ile Met Phe Ile Asn Lys Cys Ala
50          55          60
Thr Cys Lys Met Ile Leu Glu Lys Glu Ala Lys Ser Gln Lys Arg Ala
65          70          75          80
Arg His Leu Ala Arg Ala Pro Lys Ala Thr Ala Pro Thr Glu Leu Asn
85          90          95

```

-continued

Cys Asp Asp Phe Lys Lys Gly Glu Arg Asp Gly Asp Phe Ile Cys Pro
 100 105 110
 Asp Tyr Tyr Glu Ala Val Cys Gly Thr Asp Gly Lys Thr Tyr Asp Asn
 115 120 125
 Arg Cys Ala Leu Cys Ala Glu Asn Ala Lys Thr Gly Ser Gln Ile Gly
 130 135 140
 Val Lys Ser Glu Gly Glu Cys Lys Ser Ser Asn Pro Glu Gln Asp Val
 145 150 155 160
 Cys Ser Ala Phe Arg Pro Phe Val Arg Asp Gly Arg Leu Gly Cys Thr
 165 170 175
 Arg Glu Asn Asp Pro Val Leu Gly Pro Asp Gly Lys Thr His Gly Asn
 180 185 190
 Lys Cys Ala Met Cys Ala Glu Leu Phe Leu Lys Glu Ala Glu Asn Ala
 195 200 205
 Lys Arg Glu Gly Glu Thr Arg Ile Arg Arg Asn Ala Glu Lys Asp Phe
 210 215 220
 Cys Lys Glu Tyr Glu Lys Gln Val Arg Asn Gly Arg Leu Phe Cys Thr
 225 230 235 240
 Arg Glu Ser Asp Pro Val Arg Gly Pro Asp Gly Arg Met His Gly Asn
 245 250 255
 Lys Cys Ala Leu Cys Ala Glu Ile Phe Lys Gln Arg Phe Ser Glu Glu
 260 265 270
 Asn Ser Lys Thr Asp Gln Asn Leu Gly Lys Ala Glu Glu Lys Thr Lys
 275 280 285
 Val Lys Arg Glu Ile Val Lys Leu Cys Ser Gln Tyr Gln Asn Gln Ala
 290 295 300
 Lys Asn Gly Ile Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Arg Gly
 305 310 315 320
 Pro Asp Gly Lys Met His Gly Asn Leu Cys Ser Met Cys Gln Ala Tyr
 325 330 335
 Phe Gln Ala Glu Asn Glu Glu Lys Lys Lys Ala Glu Ala Arg Ala Arg
 340 345 350
 Asn Lys Arg Glu Ser Gly Lys Ala Thr Ser Tyr Ala Glu Leu Cys Ser
 355 360 365
 Glu Tyr Arg Lys Leu Val Arg Asn Gly Lys Leu Ala Cys Thr Arg Glu
 370 375 380
 Asn Asp Pro Ile Gln Gly Pro Asp Gly Lys Val His Gly Asn Thr Cys
 385 390 395 400
 Ser Met Cys Glu Val Phe Phe Gln Ala Glu Glu Glu Glu Lys Lys Lys
 405 410 415
 Lys Glu Gly Lys Ser Arg Asn Lys Arg Gln Ser Lys Ser Thr Ala Ser
 420 425 430
 Phe Glu Glu Leu Cys Ser Glu Tyr Arg Lys Ser Arg Lys Asn Gly Arg
 435 440 445
 Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Gln Gly Pro Asp Gly Lys
 450 455 460
 Met His Gly Asn Thr Cys Ser Met Cys Glu Ala Phe Phe Gln Gln Glu
 465 470 475 480
 Glu Arg Ala Arg Ala Lys Ala Lys Arg Glu Ala Ala Lys Glu Ile Cys
 485 490 495

-continued

Ser Glu Phe Arg Asp Gln Val Arg Asn Gly Thr Leu Ile Cys Thr Arg
 500 505 510

Glu His Asn Pro Val Arg Gly Pro Asp Gly Lys Met His Gly Asn Lys
 515 520 525

Cys Ala Met Cys Ala Ser Val Phe Lys Leu Glu Glu Glu Lys Lys
 530 535 540

Asn Asp Lys Glu Glu Lys Gly Lys Val Glu Ala Glu Lys Val Lys Arg
 545 550 555 560

Glu Ala Val Gln Glu Leu Cys Ser Glu Tyr Arg His Tyr Val Arg Asn
 565 570 575

Gly Arg Leu Pro Cys Thr Arg Glu Asn Asp Pro Ile Glu Gly Leu Asp
 580 585 590

Gly Lys Ile His Gly Asn Thr Cys Ser Met Cys Glu Ala Phe Phe Gln
 595 600 605

Gln Glu Ala Lys Glu Lys Glu Arg Ala Glu Pro Arg Ala Lys Val Lys
 610 615 620

Arg Glu Ala Glu Lys Glu Thr Cys Asp Glu Phe Arg Arg Leu Leu Gln
 625 630 635 640

Asn Gly Lys Leu Phe Cys Thr Arg Glu Asn Asp Pro Val Arg Gly Pro
 645 650 655

Asp Gly Lys Thr His Gly Asn Lys Cys Ala Met Cys Lys Ala Val Phe
 660 665 670

Gln Lys Glu Asn Glu Glu Arg Lys Arg Lys Glu Glu Glu Asp Gln Arg
 675 680 685

Asn Ala Ala Gly His Gly Ser Ser Gly Gly Gly Gly Gly Asn Thr Gln
 690 695 700

Asp Glu Cys Ala Glu Tyr Arg Glu Gln Met Lys Asn Gly Arg Leu Ser
 705 710 715 720

Cys Thr Arg Glu Ser Asp Pro Val Arg Asp Ala Asp Gly Lys Ser Tyr
 725 730 735

Asn Asn Gln Cys Thr Met Cys Lys Ala Lys Leu Glu Arg Glu Ala Glu
 740 745 750

Arg Lys Asn Glu Tyr Ser Arg Ser Arg Ser Asn Gly Thr Gly Ser Glu
 755 760 765

Ser Gly Lys Asp Thr Cys Asp Glu Phe Arg Ser Gln Met Lys Asn Gly
 770 775 780

Lys Leu Ile Cys Thr Arg Glu Ser Asp Pro Val Arg Gly Pro Asp Gly
 785 790 795 800

Lys Thr His Gly Asn Lys Cys Thr Met Cys Lys Glu Lys Leu Glu Arg
 805 810 815

Glu Ala Ala Glu Lys Lys Lys Lys Glu Asp Glu Asp Arg Ser Asn Thr
 820 825 830

Gly Glu Arg Ser Asn Thr Gly Glu Arg Ser Asn Asp Lys Glu Asp Leu
 835 840 845

Cys Arg Glu Phe Arg Ser Met Gln Arg Asn Gly Lys Leu Ile Cys Thr
 850 855 860

Arg Glu Asn Asn Pro Val Arg Gly Pro Tyr Gly Lys Met His Ile Asn
 865 870 875 880

Lys Cys Ala Met Cys Gln Ser Ile Phe Asp Arg Glu Ala Asn Glu Arg
 885 890 895

Lys Lys Lys Asp Glu Glu Lys Ser Ser Ser Lys Pro Ser Asn Asn Ala

-continued

900					905					910					
Lys	Asp	Gln	Cys	Arg	Gln	Val	Gln	Asn	Glu	Ala	Glu	Asp	Ala	Lys	Phe
	915						920					925			
Arg	Gln	Pro	Gly	Arg	Ser	Leu	Ala	Ser	Val	Ala	Arg	Met	Ser	Thr	Asp
	930					935					940				
Glu	Cys	Ser	Glu	Phe	Arg	Asn	Tyr	Ile	Arg	Asn	Asn	Glu	Leu	Ile	Cys
	945					950					955				960
Pro	Arg	Glu	Asn	Asp	Pro	Val	His	Gly	Ala	Asp	Gly	Lys	Phe	Tyr	Thr
				965					970					975	
Asn	Lys	Cys	Tyr	Met	Cys	Arg	Ala	Val	Phe	Leu	Thr	Glu	Ala	Leu	Glu
			980					985						990	
Arg	Ala	Lys	Leu	Gln	Glu	Lys	Pro	Ser	His	Val	Arg	Ala	Ser	Gln	Glu
		995					1000						1005		
Glu	Asp	Ser	Pro	Asp	Ser	Phe	Ser	Ser	Leu	Asp	Ser	Glu	Met	Cys	Lys
	1010					1015						1020			
Asp	Tyr	Arg	Val	Leu	Pro	Arg	Ile	Gly	Tyr	Leu	Cys	Pro	Lys	Asp	Leu
	1025					1030					1035				1040
Lys	Pro	Val	Cys	Gly	Asp	Asp	Gly	Gln	Thr	Tyr	Asn	Asn	Pro	Cys	Met
				1045					1050					1055	
Leu	Cys	His	Glu	Asn	Leu	Ile	Arg	Gln	Thr	Asn	Thr	His	Ile	Arg	Ser
			1060					1065					1070		
Thr	Gly	Lys	Cys	Glu	Glu	Ser	Ser	Thr	Pro	Gly	Thr	Thr	Ala	Ala	Ser
		1075					1080						1085		
Met	Pro	Pro	Ser	Asp	Glu										
	1090														
<210> SEQ ID NO 8															
<211> LENGTH: 916															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 8															
Met	Lys	Ile	Ala	Thr	Val	Ser	Val	Leu	Leu	Pro	Leu	Ala	Leu	Cys	Leu
1				5					10					15	
Ile	Gln	Asp	Ala	Ala	Ser	Lys	Asn	Glu	Asp	Gln	Glu	Met	Cys	His	Glu
			20					25					30		
Phe	Gln	Ala	Phe	Met	Lys	Asn	Gly	Lys	Leu	Phe	Cys	Pro	Gln	Asp	Lys
		35					40					45			
Lys	Phe	Phe	Gln	Ser	Leu	Asp	Gly	Ile	Met	Phe	Ile	Asn	Lys	Cys	Ala
	50					55					60				
Thr	Cys	Lys	Met	Ile	Leu	Glu	Lys	Glu	Ala	Lys	Ser	Gln	Lys	Arg	Ala
	65			70					75					80	
Arg	His	Leu	Ala	Arg	Ala	Pro	Lys	Ala	Thr	Ala	Pro	Thr	Glu	Leu	Asn
				85					90					95	
Cys	Asp	Asp	Phe	Lys	Lys	Gly	Glu	Arg	Asp	Gly	Asp	Phe	Ile	Cys	Pro
			100					105					110		
Asp	Tyr	Tyr	Glu	Ala	Val	Cys	Gly	Thr	Asp	Gly	Lys	Thr	Tyr	Asp	Asn
		115					120					125			
Arg	Cys	Ala	Leu	Cys	Ala	Glu	Asn	Ala	Lys	Thr	Gly	Ser	Gln	Ile	Gly
	130					135					140				
Val	Lys	Ser	Glu	Gly	Glu	Cys	Lys	Ser	Ser	Asn	Pro	Glu	Gln	Asp	Val
	145			150						155				160	

-continued

Cys Ser Ala Phe Arg Pro Phe Val Arg Asp Gly Arg Leu Gly Cys Thr
 165 170 175

Arg Glu Asn Asp Pro Val Leu Gly Pro Asp Gly Lys Thr His Gly Asn
 180 185 190

Lys Cys Ala Met Cys Ala Glu Leu Phe Leu Lys Glu Ala Glu Asn Ala
 195 200 205

Lys Arg Glu Gly Glu Thr Arg Ile Arg Arg Asn Ala Glu Lys Asp Phe
 210 215 220

Cys Lys Glu Tyr Glu Lys Gln Val Arg Asn Gly Arg Leu Phe Cys Thr
 225 230 235 240

Arg Glu Ser Asp Pro Val Arg Gly Pro Asp Gly Arg Met His Gly Asn
 245 250 255

Lys Cys Ala Leu Cys Ala Glu Ile Phe Lys Gln Arg Phe Ser Glu Glu
 260 265 270

Asn Ser Lys Thr Asp Gln Asn Leu Gly Lys Ala Glu Glu Lys Thr Lys
 275 280 285

Val Lys Arg Glu Ile Val Lys Leu Cys Ser Gln Tyr Gln Asn Gln Ala
 290 295 300

Lys Asn Gly Ile Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Arg Gly
 305 310 315 320

Pro Asp Gly Lys Met His Gly Asn Leu Cys Ser Met Cys Gln Ala Tyr
 325 330 335

Phe Gln Ala Glu Asn Glu Glu Lys Lys Lys Ala Glu Ala Arg Ala Arg
 340 345 350

Asn Lys Arg Glu Ser Gly Lys Ala Thr Ser Tyr Ala Glu Leu Cys Ser
 355 360 365

Glu Tyr Arg Lys Leu Val Arg Asn Gly Lys Leu Ala Cys Thr Arg Glu
 370 375 380

Asn Asp Pro Ile Gln Gly Pro Asp Gly Lys Val His Gly Asn Thr Cys
 385 390 395 400

Ser Met Cys Glu Val Phe Phe Gln Ala Glu Glu Glu Glu Lys Lys Lys
 405 410 415

Lys Glu Gly Lys Ser Arg Asn Lys Arg Gln Ser Lys Ser Thr Ala Ser
 420 425 430

Phe Glu Glu Leu Cys Ser Glu Tyr Arg Lys Ser Arg Lys Asn Gly Arg
 435 440 445

Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Gln Gly Pro Asp Gly Lys
 450 455 460

Met His Gly Asn Thr Cys Ser Met Cys Glu Ala Phe Phe Gln Gln Glu
 465 470 475 480

Glu Arg Ala Arg Ala Lys Ala Lys Arg Glu Ala Ala Lys Glu Ile Cys
 485 490 495

Ser Glu Phe Arg Asp Gln Val Arg Asn Gly Thr Leu Ile Cys Thr Arg
 500 505 510

Glu His Asn Pro Val Arg Gly Pro Asp Gly Lys Met His Gly Asn Lys
 515 520 525

Cys Ala Met Cys Ala Ser Val Phe Lys Leu Glu Glu Glu Lys Lys
 530 535 540

Asn Asp Lys Glu Glu Lys Gly Lys Val Glu Ala Glu Lys Val Lys Arg
 545 550 555 560

Glu Ala Val Gln Glu Leu Cys Ser Glu Tyr Arg His Tyr Val Arg Asn

-continued

Met Lys Ile Ala Thr Val Ser Val Leu Leu Pro Leu Ala Leu Cys Leu
 1 5 10 15
 Ile Gln Asp Ala Ala Ser Lys Asn Glu Asp Gln Glu Met Cys His Glu
 20 25 30
 Phe Gln Ala Phe Met Lys Asn Gly Lys Leu Phe Cys Pro Gln Asp Lys
 35 40 45
 Lys Phe Phe Gln Ser Leu Asp Gly Ile Met Phe Ile Asn Lys Cys Ala
 50 55 60
 Thr Cys Lys Met Ile Leu Glu Lys Glu Ala Lys Ser Gln Lys Arg Ala
 65 70 75 80
 Arg His Leu Ala Arg Ala Pro Lys Ala Thr Ala Pro Thr Glu Leu Asn
 85 90 95
 Cys Asp Asp Phe Lys Lys Gly Glu Arg Asp Gly Asp Phe Ile Cys Pro
 100 105 110
 Asp Tyr Tyr Glu Ala Val Cys Gly Thr Asp Gly Lys Thr Tyr Asp Asn
 115 120 125
 Arg Cys Ala Leu Cys Ala Glu Asn Ala Lys Thr Gly Ser Gln Ile Gly
 130 135 140
 Val Lys Ser Glu Gly Glu Cys Lys Ser Ser Asn Pro Glu Gln Asp Val
 145 150 155 160
 Cys Ser Ala Phe Arg Pro Phe Val Arg Asp Gly Arg Leu Gly Cys Thr
 165 170 175
 Arg Glu Asn Asp Pro Val Leu Gly Pro Asp Gly Lys Thr His Gly Asn
 180 185 190
 Lys Cys Ala Met Cys Ala Glu Leu Phe Leu Lys Glu Ala Glu Asn Ala
 195 200 205
 Lys Arg Glu Gly Glu Thr Arg Ile Arg Arg Asn Ala Glu Lys Asp Phe
 210 215 220
 Cys Lys Glu Tyr Glu Lys Gln Val Arg Asn Gly Arg Leu Phe Cys Thr
 225 230 235 240
 Arg Glu Ser Asp Pro Val Arg Gly Pro Asp Gly Arg Met His Gly Asn
 245 250 255
 Lys Cys Ala Leu Cys Ala Glu Ile Phe Lys Gln Arg Phe Ser Glu Glu
 260 265 270
 Asn Ser Lys Thr Asp Gln Asn Leu Gly Lys Ala Glu Glu Lys Thr Lys
 275 280 285
 Val Lys Arg Glu Ile Val Lys Leu Cys Ser Gln Tyr Gln Asn Gln Ala
 290 295 300
 Lys Asn Gly Ile Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Arg Gly
 305 310 315 320
 Pro Asp Gly Lys Met His Gly Asn Leu Cys Ser Met Cys Gln Ala Tyr
 325 330 335
 Phe Gln Ala Glu Asn Glu Glu Lys Lys Lys Ala Glu Ala Arg Ala Arg
 340 345 350
 Asn Lys Arg Glu Ser Gly Lys Ala Thr Ser Tyr Ala Glu Leu Cys Ser
 355 360 365
 Glu Tyr Arg Lys Leu Val Arg Asn Gly Lys Leu Ala Cys Thr Arg Glu
 370 375 380
 Asn Asp Pro Ile Gln Gly Pro Asp Gly Lys Val His Gly Asn Thr Cys
 385 390 395 400
 Ser Met Cys Glu Val Phe Phe Gln Ala Glu Glu Glu Glu Lys Lys Lys

-continued

<400> SEQUENCE: 11

```

Ala Pro Thr Glu Leu Asn Cys Asp Asp Phe Lys Lys Gly Glu Arg Asp
          5              10              15
Gly Asp Phe Ile Cys Pro Asp Tyr Tyr Glu Ala Val Cys Gly Thr Asp
          20              25              30
Gly Lys Thr Tyr Asp Asn Arg Cys Ala Leu Cys Ala Glu Asn Ala Lys
          35              40              45
Thr Gly Ser Gln Ile Gly Val Lys Ser Glu Gly Glu Cys Lys Ser
          50              55              60

```

<210> SEQ ID NO 12

<211> LENGTH: 62

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 12

```

Asn Pro Glu Gln Asp Val Cys Ser Ala Phe Arg Pro Phe Val Arg Asp
          5              10              15
Gly Arg Leu Gly Cys Thr Arg Glu Asn Asp Pro Val Leu Gly Pro Asp
          20              25              30
Gly Lys Thr His Gly Asn Lys Cys Ala Met Cys Ala Glu Leu Phe Leu
          35              40              45
Lys Glu Ala Glu Asn Ala Lys Arg Glu Gly Glu Thr Arg Ile
          50              55              60

```

<210> SEQ ID NO 13

<211> LENGTH: 67

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 13

```

Asn Ala Glu Lys Asp Phe Cys Lys Glu Tyr Glu Lys Gln Val Arg Asn
          5              10              15
Gly Arg Leu Phe Cys Thr Arg Glu Ser Asp Pro Val Arg Gly Pro Asp
          20              25              30
Gly Arg Met His Gly Asn Lys Cys Ala Leu Cys Ala Glu Ile Phe Lys
          35              40              45
Gln Arg Phe Ser Glu Glu Asn Ser Lys Thr Asp Gln Asn Leu Gly Lys
          50              55              60
Ala Glu Glu
          65

```

<210> SEQ ID NO 14

<211> LENGTH: 62

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 14

```

Arg Glu Ile Val Lys Leu Cys Ser Gln Tyr Gln Asn Gln Ala Lys Asn
          5              10              15
Gly Ile Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Arg Gly Pro Asp
          20              25              30

```

-continued

Gly Lys Met His Gly Asn Leu Cys Ser Met Cys Gln Ala Tyr Phe Gln
 35 40 45

Ala Glu Asn Glu Glu Lys Lys Lys Ala Glu Ala Arg Ala Arg
 50 55 60

<210> SEQ ID NO 15
 <211> LENGTH: 63
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 15

Thr Ser Tyr Ala Glu Leu Cys Ser Glu Tyr Arg Lys Leu Val Arg Asn
 5 10 15

Gly Lys Leu Ala Cys Thr Arg Glu Asn Asp Pro Ile Gln Gly Pro Asp
 20 25 30

Gly Lys Val His Gly Asn Thr Cys Ser Met Cys Glu Val Phe Phe Gln
 35 40 45

Ala Glu Glu Glu Glu Lys Lys Lys Lys Glu Gly Lys Ser Arg Asn
 50 55 60

<210> SEQ ID NO 16
 <211> LENGTH: 59
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 16

Ala Ser Phe Glu Glu Leu Cys Ser Glu Tyr Arg Lys Ser Arg Lys Asn
 5 10 15

Gly Arg Leu Phe Cys Thr Arg Glu Asn Asp Pro Ile Gln Gly Pro Asp
 20 25 30

Gly Lys Met His Gly Asn Thr Cys Ser Met Cys Glu Ala Phe Phe Gln
 35 40 45

Gln Glu Glu Arg Ala Arg Ala Lys Ala Lys Arg
 50 55

<210> SEQ ID NO 17
 <211> LENGTH: 62
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 17

Glu Ala Ala Lys Glu Ile Cys Ser Glu Phe Arg Asp Gln Val Arg Asn
 5 10 15

Gly Thr Leu Ile Cys Thr Arg Glu His Asn Pro Val Arg Gly Pro Asp
 20 25 30

Gly Lys Met His Gly Asn Lys Cys Ala Met Cys Ala Ser Val Phe Lys
 35 40 45

Leu Glu Glu Glu Glu Lys Lys Asn Asp Lys Glu Glu Lys Gly
 50 55 60

<210> SEQ ID NO 18
 <211> LENGTH: 62

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 18

Glu Ala Val Gln Glu Leu Cys Ser Glu Tyr Arg His Tyr Val Arg Asn
      5              10              15
Gly Arg Leu Pro Cys Thr Arg Glu Asn Asp Pro Ile Glu Gly Leu Asp
      20              25              30
Gly Lys Ile His Gly Asn Thr Cys Ser Met Cys Glu Ala Phe Phe Gln
      35              40              45
Gln Glu Ala Lys Glu Lys Glu Arg Ala Glu Pro Arg Ala Lys
      50              55              60

```

```

<210> SEQ ID NO 19
<211> LENGTH: 63
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 19

Glu Ala Glu Lys Glu Thr Cys Asp Glu Phe Arg Arg Leu Leu Gln Asn
      5              10              15
Gly Lys Leu Phe Cys Thr Arg Glu Asn Asp Pro Val Arg Gly Pro Asp
      20              25              30
Gly Lys Thr His Gly Asn Lys Cys Ala Met Cys Lys Ala Val Phe Gln
      35              40              45
Lys Glu Asn Glu Glu Arg Lys Arg Lys Glu Glu Glu Asp Gln Arg
      50              55              60

```

```

<210> SEQ ID NO 20
<211> LENGTH: 57
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 20

Gly Asn Thr Gln Asp Glu Cys Ala Glu Tyr Arg Glu Gln Met Lys Asn
      5              10              15
Gly Arg Leu Ser Cys Thr Arg Glu Ser Asp Pro Val Arg Asp Ala Asp
      20              25              30
Gly Lys Ser Tyr Asn Asn Gln Cys Thr Met Cys Lys Ala Lys Leu Glu
      35              40              45
Arg Glu Ala Glu Arg Lys Asn Glu Tyr
      50              55

```

```

<210> SEQ ID NO 21
<211> LENGTH: 63
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 21

Glu Ser Gly Lys Asp Thr Cys Asp Glu Phe Arg Ser Gln Met Lys Asn
      5              10              15

```

-continued

Gly Lys Leu Ile Cys Thr Arg Glu Ser Asp Pro Val Arg Gly Pro Asp
 20 25 30

Gly Lys Thr His Gly Asn Lys Cys Thr Met Cys Lys Glu Lys Leu Glu
 35 40 45

Arg Glu Ala Ala Glu Lys Lys Lys Lys Glu Asp Glu Asp Arg Ser
 50 55 60

<210> SEQ ID NO 22
 <211> LENGTH: 63
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 22

Asn Asp Lys Glu Asp Leu Cys Arg Glu Phe Arg Ser Met Gln Arg Asn
 5 10 15

Gly Lys Leu Ile Cys Thr Arg Glu Asn Asn Pro Val Arg Gly Pro Tyr
 20 25 30

Gly Lys Met His Ile Asn Lys Cys Ala Met Cys Gln Ser Ile Phe Asp
 35 40 45

Arg Glu Ala Asn Glu Arg Lys Lys Lys Asp Glu Glu Lys Ser Ser
 50 55 60

<210> SEQ ID NO 23
 <211> LENGTH: 92
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 23

Asn Asn Ala Lys Asp Gln Cys Arg Gln Val Gln Asn Glu Ala Glu Asp
 5 10 15

Ala Lys Phe Arg Gln Pro Gly Arg Ser Leu Ala Ser Val Ala Arg Met
 20 25 30

Ser Thr Asp Glu Cys Ser Glu Phe Arg Asn Tyr Ile Arg Asn Asn Glu
 35 40 45

Leu Ile Cys Pro Arg Glu Asn Asp Pro Val His Gly Ala Asp Gly Lys
 50 55 60

Phe Tyr Thr Asn Lys Cys Tyr Met Cys Arg Ala Val Phe Leu Thr Glu
 65 70 75 80

Ala Leu Glu Arg Ala Lys Leu Gln Glu Lys Pro Ser
 85 90

<210> SEQ ID NO 24
 <211> LENGTH: 62
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 24

Asn Asn Ala Lys Asp Glu Cys Ser Glu Phe Arg Asn Tyr Ile Arg Asn
 5 10 15

Asn Glu Leu Ile Cys Pro Arg Glu Asn Asp Pro Val His Gly Ala Asp
 20 25 30

Gly Lys Phe Tyr Thr Asn Lys Cys Tyr Met Cys Arg Ala Val Phe Leu

-continued

tccatgtgcg aggtgttctt tcaggccgag gaagaggaaa aaaagaagaa agaggccaag	1260
tcccggaaca agcggcagag caagagcacc gccagcttcg aggaactgtg ctccgagtac	1320
cggaagtcca gaaagaatgg ccggcttttc tgcacgagag agaatgatcc catccagggg	1380
cctgatggca aaatgcatgg caatacttgc tctatgtgcg aggcattttt ccagcaagag	1440
gaacgggcca gagccaaggc caaaagagag gccgccaag agatctgcag cgagttagg	1500
gaccaagtcc gcaacggcac cctgatctgc accagggaac acaaccctgt cagagggcca	1560
gacgggaaaa tgcacgggaa caagtgtgcc atgtgcgcct ccgtgttcaa gctggaagaa	1620
gaagaaaaga aaaacgacaa agaagagaag ggaaaagtgc aagccgagaa agtgaagagg	1680
gaagccgtgc aagaactctg ttctgagtac cggcactacg tccggaacgg cagactgcct	1740
tgtactcggg aaaacgaccc aatcgaagga ctggacggga agatccatgg caacacctgt	1800
agtatgtgcg aggcattctt tcaacaagaa gccaaagaaa aagagcgggc cgagcctcgg	1860
gccaaagtca aaagagaagc tgagaagag acatgcgacg agttccggcg gctgctgcag	1920
aatggaaaac tcttctgtac acgtgaaaat gatcccgtgc ggggaccaga tggaaagacc	1980
catgggaaca aatcgctat gtgcaaggcc gtgttccaaa aagagaatga ggaacgcaag	2040
cggaagagg aagaagatca gcggaacgcc gctggccacg gatctagtgg cggaggtggt	2100
ggaaataccc aggcagagtg tgcgaatac cgggaacaga tgaagaatgg aaggctgagc	2160
tgcacacggg aatctgaccc cgttagagat gccgacggca agtcctacaa caaccagtgc	2220
accatgtgta aagccaagct cgagcgcgag gccgagcggga agaattgagta cagcagaagc	2280
cggcttaacg gcaccggctc tgagtctggc aaggatacct gtgacgagtt cagatcccag	2340
atgaagaacg gaaaactcat ctgcacaaga gaatcagacc cagtcocggg acctgacggg	2400
aaaactcatg gcaacaagtg cacaatgtgc aaagagaac tggaacgcga ggctgccgag	2460
aagaaaaaga aagaggacga ggatcggagc aacaccggcg agagatctaa taccggcgag	2520
cggagcaatg acaaaagga cctctgccgg gaatttcgga gcatgcagag aaacggaaaag	2580
ctcatatgta cgcgcgagaa caatccggtt cggggccctt atgggaagat gcacattaac	2640
aaatgtgcaa tgtgccagag catcttcgac agagaggcca acgaacgcaa aaagaaggat	2700
gaagagaagt ccagcagcaa gccagcaac aacccaagg acgaatgctc tgagttccgc	2760
aattacatcc ggaacaacga gctgatctgt ccgcgagaga acgatccagt tcatggcgcc	2820
gatgggaagt tctacaccaa caagtgttat atgtgcagag ccgtctttct gacagaggcc	2880
ctggaagag ccaagctgca agagaagcct tctcacgtgc gggcctctca agaagaggac	2940
agccctgata gcttcagcag cctggacagc gagatgtgca aggactacag agtgcgtgcc	3000
cggatcggct atctgtgccc caaggatctg aagcctgtgt gtggcgacga cggccagaca	3060
tacaacaatc cctgcatgct gtgtcacgag aacctgatcc ggcagaccaa cacacacatc	3120
cgggccaccg gcaagtgcga agagtctagc acacctggaa caaccggcgc ctctatgcct	3180
cctagcgacg aatga	3195

What is claimed is:

1. A pharmaceutical composition comprising:
 - (a) a herpes virus comprising a recombinant herpes virus genome, wherein the recombinant herpes virus genome comprises one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide; and
 - (b) a pharmaceutically acceptable excipient.
2. The pharmaceutical composition of claim 1, wherein the recombinant herpes virus genome is replication competent.
3. The pharmaceutical composition of claim 1, wherein the recombinant herpes virus genome is replication defective.
4. The pharmaceutical composition of claim 1, wherein the recombinant herpes virus genome is selected from the group consisting of a recombinant herpes simplex virus genome, a recombinant varicella zoster virus genome, a recombinant human cytomegalovirus genome, a recombinant herpesvirus 6A genome, a recombinant herpesvirus 6B genome, a recombinant herpesvirus 7 genome, a recombinant Kaposi's sarcoma-associated herpesvirus genome, and any derivatives thereof.
5. The pharmaceutical composition of claim 1, wherein the recombinant herpes virus genome is a recombinant herpes simplex virus type 1 (HSV-1) genome.
6. The pharmaceutical composition of claim 5, wherein the recombinant HSV-1 genome comprises an inactivating mutation in a herpes simplex virus gene selected from the group consisting of Infected Cell Protein (ICP) 0, ICP4, ICP22, ICP27, ICP47, thymidine kinase (tk), Long Unique Region (UL) 41, and UL55.
7. The pharmaceutical composition of claim 1, wherein the SPINK polypeptide is a human SPINK polypeptide.
8. The pharmaceutical composition of claim 1, wherein the SPINK polypeptide is a Serine Protease Inhibitor Kazal-type 5 (SPINK5) polypeptide.
9. The pharmaceutical composition of claim 1, wherein the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NOS: 7-25.
10. The pharmaceutical composition of claim 1, wherein the herpes virus has reduced cytotoxicity as compared to a corresponding wild-type herpes virus.
11. The pharmaceutical composition of claim 1, wherein the herpes virus is selected from the group consisting of a herpes simplex virus, a varicella zoster virus, a human cytomegalovirus, a herpesvirus 6A, a herpesvirus 6B, a herpesvirus 7, and a Kaposi's sarcoma-associated herpesvirus.
12. The pharmaceutical composition of claim 1, wherein the herpes virus is a herpes simplex virus type 1 (HSV-1).
13. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is suitable for topical, transdermal, subcutaneous, intradermal, oral, intranasal, intratracheal, sublingual, buccal, rectal, vaginal, inhaled, intravenous, intraarterial, intramuscular, intracardiac, intraosseous, intraperitoneal, transmucosal, intravitreal, subretinal, intraarticular, peri-articular, local, or epicutaneous administration.
14. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is suitable for topical, transdermal, or intradermal administration.
15. A method of providing prophylactic, palliative, or therapeutic relief to one or more signs or symptoms of Netherton Syndrome (NS) or atopic dermatitis (AD) in a subject in need thereof, the method comprising administering to the subject an effective amount of a pharmaceutical composition comprising:
 - (a) a herpes virus comprising a recombinant herpes virus genome, wherein the recombinant herpes virus genome comprises one or more polynucleotides encoding a Serine Protease Inhibitor Kazal-type (SPINK) polypeptide; and
 - (b) a pharmaceutically acceptable excipient.
16. The method of claim 15, wherein the subject is a human.
17. The method of claim 15, wherein the one or more signs or symptoms of NS are selected from the group consisting of defective keratinization, a defective skin barrier, recurrent skin infections, congenital ichthyosiform erythroderma, ichthyosis linearis circumflexa, trichorrhexis invaginata, chronic skin inflammation, and any combinations thereof.
18. The method of claim 15, wherein the one or more signs or symptoms of atopic dermatitis are selected from the group consisting of itchy skin, dry skin, red to brownish-grey patches on the skin, small raised bumps on the skin, thickened skin, cracked skin, scaly skin, swollen skin, weeping sores, skin infections, eyelid dermatitis, cataracts, and any combinations thereof.
19. The method of claim 15, wherein the pharmaceutical composition is administered topically, transdermally, subcutaneously, epicutaneously, intradermally, orally, sublingually, buccally, rectally, vaginally, intravenously, intraarterially, intramuscularly, intraosseously, intracardially, intraperitoneally, transmucosally, intravitreally, subretinally, intraarticularly, peri-articularly, locally, or via inhalation to the subject.
20. The method of claim 15, wherein the pharmaceutical composition is administered topically, transdermally, or intradermally to the subject.
21. The method of claim 20, wherein the skin of the subject is abraded prior to administration.
22. The method of claim 15, wherein the recombinant herpes virus genome is selected from the group consisting of a recombinant herpes simplex virus genome, a recombinant varicella zoster virus genome, a recombinant human cytomegalovirus genome, a recombinant herpesvirus 6A genome, a recombinant herpesvirus 6B genome, a recombinant herpesvirus 7 genome, a recombinant Kaposi's sarcoma-associated herpesvirus genome, and any derivatives thereof.
23. The method of claim 15, wherein the recombinant herpes virus genome is a recombinant herpes simplex virus type 1 (HSV-1) genome.
24. The method of claim 23, wherein the recombinant HSV-1 genome comprises an inactivating mutation in a herpes simplex virus gene selected from the group consisting of Infected Cell Protein (ICP) 0, ICP4, ICP22, ICP27, ICP47, thymidine kinase (tk), Long Unique Region (UL) 41, and UL55.
25. The method of claim 15, wherein the SPINK polypeptide is a human SPINK polypeptide.

26. The method of claim **15**, wherein the wherein the SPINK polypeptide is a Serine Protease Inhibitor Kazal-type 5 (SPINK5) polypeptide.

27. The method of claim **15**, wherein the SPINK polypeptide comprises a sequence having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NOS: 7-25.

28. The method of claim **15**, wherein the herpes virus has reduced cytotoxicity as compared to a corresponding wild-type herpes virus.

29. The method of claim **15**, wherein the herpes virus is selected from the group consisting of a herpes simplex virus, a varicella zoster virus, a human cytomegalovirus, a herpesvirus 6A, a herpesvirus 6B, a herpesvirus 7, and a Kaposi's sarcoma-associated herpesvirus.

30. The method of claim **15**, wherein the herpes virus is a herpes simplex virus type 1 (HSV-1).

* * * * *