Innovation, Sciences et Développement économique Canada
Office de la Propriété Intellectuelle du Canada

Innovation, Science and Economic Development Canada
Canadian Intellectual Property Office

CA 2787383 C 2018/05/22

(11)(21) **2 787 383**

(12) BREVET CANADIEN / CANADIAN PATENT

(13) **C**

(86) **Date de dépôt PCT/PCT Filing Date:** 2011/01/27
(87) **Date publication PCT/PCT Publication Date:** 2011/08/04
(45) **Date de délivrance/Issue Date:** 2018/05/22
(85) **Entrée phase nationale/National Entry:** 2012/07/16
(86) **N° demande PCT/PCT Application No.:** US 2011/022690
(87) **N° publication PCT/PCT Publication No.:** 2011/094388
(30) **Priorité/Priority:** 2010/01/29 (US61/299,517)

(51) **Cl.Int./Int.Cl.** ***A01N 43/40*** (2006.01), ***A01N 39/04*** (2006.01), ***A01P 13/00*** (2006.01)

(72) **Inventeur/Inventor:**
CARRANZA GARZON, NELSON M., CO

(73) **Propriétaire/Owner:**
DOW AGROSCIENCES LLC, US

(74) **Agent:** SMART & BIGGAR

(54) **Titre : COMPOSITION HERBICIDE SYNERGIQUE CONTENANT DE L'AMINOPYRALIDE ET DE L'ACIDE 2,4-DICHLOROPHENOXYACETIQUE**
(54) **Title: SYNERGISTIC HERBICIDAL COMPOSITION CONTAINING AMINOPYRALID AND 2,4-DICHLOROPHENOXYACETIC ACID**

(57) **Abrégé/Abstract:**
An herbicidal composition containing (a) aminopyralid and (b) 2,4-D provides synergistic control of selected broadleaf weeds.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
4 August 2011 (04.08.2011)

PCT

(10) International Publication Number
WO 2011/094388 A3

(51) International Patent Classification:
A01N 43/40 (2006.01) *A01P 13/00* (2006.01)
A01N 39/04 (2006.01)

(21) International Application Number:
PCT/US2011/022690

(22) International Filing Date:
27 January 2011 (27.01.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/299,517 29 January 2010 (29.01.2010) US

(71) Applicant *(for all designated States except US)*: **DOW AGROSCIENCES LLC** [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268 (US).

(72) Inventor; and
(75) Inventor/Applicant *(for US only)*: **CARRANZA GARZON, Nelson** [CO/CO]; Girasol Mz. F Casa 1718 # 7 Floor, Ibague (CO).

(74) Agent: **MIXAN, Craig**; Dow AgroSciences LLC, 9330 Zionsville Rd, Indianapolis, Indiana 46268 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(88) Date of publication of the international search report:
29 March 2012

WO 2011/094388 A3

(54) Title: SYNERGISTIC HERBICIDAL COMPOSITION CONTAINING AMINOPYRALID AND 2,4-DICHLOROPHENOXYACETIC ACID

(57) Abstract: An herbicidal composition containing (a) aminopyralid and (b) 2,4-D provides synergistic control of selected broadleaf weeds.

SYNERGISTIC HERBICIDAL COMPOSITION CONTAINING AMINOPYRALID AND 2,4-DICHLOROPHENOXYACETIC ACID

The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem, researchers in the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use.

In some cases, herbicidal active ingredients have been shown to be more effective in combination than when applied individually and this is referred to as "synergism." As described in the *Herbicide Handbook* of the Weed Science Society of America, Eighth Edition, 2002, p. 462, "'synergism' [is] an interaction of two or more factors such that the effect when combined is greater than the predicted effect based on the response to each factor applied separately." The present invention is based on the discovery that 2,4-D and aminopyralid, already known individually for their herbicidal efficacy, display a synergistic effect when applied in combination.

The herbicidal compounds forming the synergistic composition of this invention are independently known in the art for their effects on plant growth.

The present invention concerns a synergistic herbicidal mixture comprising an herbicidally effective amount of (a) aminopyralid and (b) 2,4-D. The compositions may also contain an agriculturally acceptable adjuvant or carrier.

The present invention also concerns a method of controlling the growth of undesirable vegetation in crops, rangeland and pastures, and non-croplands, particularly in rice, and the use of this synergistic composition.

Aminopyralid is the common name for 4-amino-3,6-dichloro-2-pyridinecarboxylic acid. Its herbicidal activity is described in *The Pesticide Manual*, Fourteenth Edition, 2006. Aminopyralid controls annual and perennial broadleaf weeds in grassland.

2,4-D is the common name for 2,4-dichlorophenoxyacetic acid. Its herbicidal activity is described in *The Pesticide Manual*, Fourteenth Edition, 2006. 2,4-D controls both annual and perennial broadleaf weeds in a variety of grassy crops.

The term herbicide is used herein to mean an active ingredient that kills, controls or otherwise adversely modifies the growth of plants. An herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect and includes deviations from natural development, killing, regulation, desiccation, retardation, and the like. The terms plants and vegetation include germinant seeds, emerging seedlings and established vegetation.

Herbicidal activity is exhibited by the compounds of the synergistic mixture when they are applied directly to the plant or to the locus of the plant at any stage of growth or before planting or emergence. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted as is known in the art to promote non-selective or selective herbicidal action. Generally, it is preferred to apply the composition of the present invention postemergence to relatively immature undesirable vegetation to achieve the maximum control of weeds.

Both aminopyralid and 2,4-D are carboxylic acids and can be applied either as an ester or a salt. Preferred esters of 2,4-D include the ethyl, isopropyl, butyl, isobutyl, isooctyl, 2-ethylhexyl and 2-butoxyethyl esters. Preferred salts of 2,4-D include the sodium, isopropylammonium, dimethylammonium, diethanolammonium, di-isopropylammonium, triethanolammonium, tri-isopropylammonium, tri-isopropanolammonium and choline salts. Preferred esters of aminopyralid include the butyl ester. Preferred salts of aminopyralid include the potassium, dimethylammonium, and tri-isopropanolammonium salts.

In the composition of this invention, the active ingredient ratio (wt:wt) of 2,4-D to aminopyralid on an acid equivalent (ae) basis at which the herbicidal effect is synergistic lies within the range of between 10:1 and 280:1 with a ratio of 20:1 being preferred.

The rate at which the synergistic composition is applied will depend upon the particular type of weed to be controlled, the degree of control required, and the timing and

method of application. In general, the composition of the invention can be applied at an application rate of between 104 grams acid equivalent per hectare (gae/ha) and 1180 gae/ha based on the total amount of active ingredients in the composition. An application rate between 180 gae/ha and 840 gae/ha is preferred. In an especially preferred embodiment of the invention, 2,4-D is applied at a rate between 180 gae/ha and 240 gae/ha, and aminopyralid is applied at a rate between 9 gae/ha and 12 gae/ha.

The components of the synergistic mixture of the present invention can be applied either separately or as part of a multipart herbicidal system.

The synergistic mixture of the present invention can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the synergistic composition of the present invention include: acetochlor, acifluorfen, aclonifen, AE0172747, alachlor, amidosulfuron, aminocyclopyraclor, aminotriazole, ammonium thiocyanate, anilifos, atrazine, AVH 301, azimsulfuron, benfuresate, bensulfuron-methyl, bentazone, benthiocarb, benzobicyclon, bifenox, bispyribac-sodium, bromacil, bromoxynil, butachlor, butafenacil, butralin, cafenstrole, carbetamide, carfentrazone-ethyl, chlorflurenol, chlorimuron, chlorpropham, cinosulfuron, clethodim, clomazone, clopyralid, cloransulam-methyl, cyclosulfamuron, cycloxydim, cyhalofop-butyl, dicamba, dichlobenil, dichlorprop-P, diclosulam, diflufenican, diflufenzopyr, dimethenamid, dimethenamid-p, diquat, dithiopyr, diuron, EK2612, EPTC, esprocarb, ET-751, ethoxysulfuron, ethbenzanid, F7967, fenoxaprop, fenoxaprop-ethyl, fenoxaprop-ethyl + isoxadifen-ethyl, fentrazamide, flazasulfuron, florasulam, fluazifop, fluazifop-P-butyl, flucetosulfuron (LGC-421530, flufenacet, flufenpyr-Ethyl, flumetsulam, flumiclorac-pentyl, flumioxazin, fluometuron, flupyrsulfuron, fluroxypyr, fomesafen, foramsulfuron, fumiclorac, glufosinate, glufosinate-ammonium, glufosinate-P, glyphosate, halosulfuron, haloxyfop-methyl, haloxyfop-R, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, imazosulfuron, indanofan, iodosulfuron, ioxynil, ipfencarbazone, IR 5790, isoproturon, isoxaben, isoxaflutole, KUH-021, lactofen, linuron, MCPA, mecoprop-P, mefenacet, mesosulfuron, mesotrione, metamifop, metazosulfuron, metolachlor, metosulam, metribuzin, metsulfuron, molinate, MSMA, napropamide, nicosulfuron, norflurazon, OK-9701, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon,

oxazichlomefone, oxyfluorfen, paraquat, pendimethalin, penoxsulam, pentoxazone, pethoxamid, picloram, picolinafen, piperophos, pretilachlor, profoxydim, propachlor, propanil, propyzamide, prosulfocarb, prosulfuron, pyraclonil, pyrazogyl, pyrazosulfuron, pyribenzoxim (LGC-40863), pyriftalid, pyriminobac-methyl, pyroxasulfone, pyroxsulam, primisulfuron, quinclorac, quizalofop-ethyl-D, S-3252, saflufenacil, sethoxydim, simazine, SL-0401, SL-0402, s-metolachlor, sulcotrione, sulfentrazone, sulfosate, tebuthiuron, terbacil, TH-547, thiazopyr, thiobencarb, triclopyr, trifluralin and tritosulfuron.

The synergistic composition of the present invention can, further, be used in conjunction with glyphosate, glufosinate, dicamba or imidazolinones on glyphosate-tolerant, glufosinate-tolerant, dicamba-tolerant or imidazolinone-tolerant crops. The synergistic composition of the present invention is particularly well-suited for use on 2,4-D-tolerant crops.

It is generally preferred to use the synergistic composition of the present invention in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. It is further generally preferred to apply the synergistic composition of the present invention and other complementary herbicides at the same time, either as a combination formulation or as a tank mix.

The synergistic composition of the present invention can generally be employed in combination with known herbicide safeners, such as benoxacor, benthiocarb, brassinolide, cloquintocet (mexyl), cyometrinil, cyprosulfamate, daimuron, dichlormid, dicyclonon, dietholate, dimepiperate, disulfoton, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, mefenpyr-diethyl, mephenate, MG 191, MON 4660, naphthalic anhydride (NA), oxabetrinil, R29148 and *N*-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity.

In practice, it is preferable to use the synergistic composition of the present invention in mixtures containing an herbicidally effective amount of the herbicidal components along with at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such

mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions.

Suitable agricultural adjuvants and carriers that are useful in preparing the herbicidal mixtures of the invention are well known to those skilled in the art. Some of these adjuvants include, but are not limited to, crop oil concentrate (mineral oil (85%) + emulsifiers (15%)); nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C_9-C_{11} alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C_{12}-C_{16}) ethoxylate; di-*sec*-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

Liquid carriers that can be employed include water and organic solvents. The organic solvents typically used include, but are not limited to, petroleum fractions or hydrocarbons such as mineral oil, aromatic solvents, paraffinic oils, and the like; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; esters of the above vegetable oils; esters of monoalcohols or dihydric, trihydric, or other lower polyalcohols (4-6 hydroxy containing), such as 2-ethyl hexyl stearate, *n*-butyl oleate, isopropyl myristate, propylene glycol dioleate, di-octyl succinate, di-butyl adipate, di-octyl phthalate and the like; esters of mono, di and polycarboxylic acids and the like. Specific organic solvents include toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, *N*-methyl-2-pyrrolidinone, *N,N*-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. Water is generally the carrier of choice for the dilution of concentrates.

Suitable solid carriers include talc, pyrophyllite clay, silica, attapulgus clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, and the like.

It is usually desirable to incorporate one or more surface-active agents into the compositions of the present invention. Such surface-active agents are advantageously employed in both solid and liquid compositions, especially those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Surfactants conventionally used in the art of formulation and which may also be used in the present formulations are described, *inter alia*, in "McCutcheon's Detergents and Emulsifiers Annual," MC Publishing Corp., Ridgewood, New Jersey, 1998 and in "Encyclopedia of Surfactants," Vol. I-III, Chemical Publishing Co., New York, 1980-81. Typical surface-active agents include salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C_{18} ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C_{16} ethoxylate; soaps, such as sodium stearate; alkylnaphthalene-sulfonate salts, such as sodium dibutyl-naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono- and dialkyl phosphate esters; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cotton seed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils, particularly methyl esters.

Oftentimes, some of these materials, such as vegetable or seed oils and their esters, can be used interchangeably as an agricultural adjuvant, as a liquid carrier or as a surface active agent.

Other adjuvants commonly used in agricultural compositions include compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing

agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

The concentration of the active ingredients in the synergistic composition of the present invention is generally from 0.001 to 98 percent by weight. Concentrations from 0.01 to 90 percent by weight are often employed. In compositions designed to be employed as concentrates, the active ingredients are generally present in a concentration from 5 to 98 weight percent, preferably 10 to 90 weight percent. Such compositions are typically diluted with an inert carrier, such as water, before application. The diluted compositions usually applied to weeds or the locus of weeds generally contain 0.0001 to 1 weight percent active ingredient and preferably contain 0.001 to 0.05 weight percent.

The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation water, and by other conventional means known to those skilled in the art.

The following examples illustrate the present invention.

Examples

Evaluation of Postemergence Herbicidal Activity of Mixtures under Field Conditions

Methodology

These trials were conducted under field conditions in Tolima, Colombia. Trial sites were located in commercially grown fields of common rice (*Oryza sativa*). The rice crop was grown using normal cultural practices for fertilization, seeding, and maintenance to ensure good growth of the crop and the weeds. The trials were conducted using normal research methodology. Trial plots were 3 meters (m) wide by 6 m long. All treatments were applied using a randomized complete block trial design with 4 replications per treatment. The trial sites had naturally occurring populations of weeds. The weed spectrum included, but was not limited to, rice flatsedge (*Cyperus iria*, CYPIR); purple nutsedge (*Cyperus rotundus*, CYPRO); and common purslane, (*Portulaca oleracea*, POROL). The plots were treated with a postemergence foliar application 15 to 20 days after emergence of the rice.

Treatments consisted of tank mixes of soluble granules of aminopyralid tri-isopropanolammonium salt and commercially available formulations of 2,4-D (DMA 6 herbicide, water dispersible granules). The application volume was 160 liters per hectare (L/ha) of water. All application were made using precision gas hand sprayers using a 3 m boom using flat fan (80°) nozzles to broadcast the treatments over the top of the rice.

Evaluation

The treated plots and control plots were rated blind at various intervals after application. Ratings were based of Percent (%) Visual weed control, where 0 corresponds to no injury and 100 corresponds to complete kill.

Data were collected for all trials and analyzed using various statistical methods.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. *Weeds* **1967** *15*, 20-22).

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture;

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

The results are summarized in Table 1.

Table 1. Synergistic Herbicidal Activity 27 days after application

Application Rate		**% Control**					
Aminopyralid	**2,4-D**	**CYPRO**		**POROL**		**CYPIR**	
(rate in grams ae/ha)		Obs	Expected*	Obs	Expected*	Obs	Expected*
9	0	0	-	41	-	0	-
0	180	38	-	20	-	45	-
9	180	84	38	97	53	65	45
12	0	0	-	53	-	13	-
0	240	60	-	39	-	46	-
12	240	88	60	95	71	79	53

CYPRO – purple nutsedge (*Cyperus rotundus*)
POROL – common purslane (*Portulaca oleracea*)
CYPIR – rice flatsedge (*Cyperus iria*)
grams ae/ha – grams of acid equivalent per hectare
Obs – percent control observed
Expected* – percent control expected by Colby equation

CLAIMS:

1. A synergistic herbicidal mixture comprising an herbicidally effective amount of (a) aminopyralid and (b) 2,4-D, in which the weight ratio of 2,4-D to aminopyralid on an acid equivalent (ae) basis is between 10:1 and 20:1 or is 20:1.

2. The synergistic mixture of Claim 1, in which the aminopyralid is applied as an ester or a salt.

3. The synergistic mixture of Claim 1 or 2, in which 2,4-D is applied as an ester or a salt.

4. The synergistic mixture of Claim 1 or 3, wherein 2,4-D is a salt selected from the group consisting of sodium, isopropylammonium, dimethylammonium, diethanolammonium, di-isopropylammonium, triethanolammonium, tri-isopropylammonium, tri-isopropanolammonium and choline salts.

5. The synergistic mixture of Claim 1 or 2, wherein aminopyralid is a salt selected from the group consisting of potassium, dimethylammonium and tri-isopropanolammonium salts.

6. An herbicidal composition comprising an herbicidally effective amount of the synergistic herbicidal mixture of Claim 1 and an agriculturally acceptable adjuvant or carrier.

7. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with an herbicidally effective amount the synergistic herbicidal mixture of Claim 1 and in which the undesirable vegetation is in rice.

8. The method of Claim 7 for use on 2,4-D-tolerant crops.

9. The method of Claim 7, wherein the synergistic herbicidal mixture is applied at an application rate of between 104 grams acid equivalent per hectare (gae/ha) and 1180 gae/ha based on the total amount of active ingredients in the composition.

10. The method of Claim 7, wherein the synergistic herbicidal mixture is applied at an application rate of between 180 grams acid equivalent per hectare (gae/ha) and 840 gae/ha based on the total amount of active ingredients in the composition.

11. The method of Claim 7, wherein 2,4-D is applied at a rate between 180 gae/ha and 240 gae/ha, and aminopyralid is applied at a rate between 9 gae/ha and 12 gae/ha.

12. The method of Claim 7, wherein 2,4-D is a salt selected from the group consisting of sodium, isopropylammonium, dimethylammonium, diethanolammonium, di-isopropylammonium, triethanolammonium, tri-isopropylammonium, tri-isopropanolammonium and choline salts.

13. The method of Claim 7, wherein aminopyralid is a salt selected from the group consisting of potassium, dimethyl-ammonium and tri-isopropanolammonium salts.