


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 471/04, C07C 309/04 A61K 31/435 // (C07D 471/04 C07D 221:00, 209:00)	A1	(11) International Publication Number: WO 94/03452 (43) International Publication Date: 17 February 1994 (17.02.94)
(21) International Application Number: PCT/EP93/02014 (22) International Filing Date: 29 July 1993 (29.07.93) (30) Priority data: 9216154.6 30 July 1992 (30.07.92) GB (71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : HARTLEY, Peter, Leslie [GB/GB]; Glaxo Group Research Limited, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). ROBERTS, Tony, Gordon [GB/GB]; Glaxo Group Research Limited, Park Road, Ware, Hertfordshire SG12 0DP (GB). WHITESELL, Leonard, Graham [US/US]; Glaxo Inc., Five Moore Drive, Research Triangle Park, NC 27709 (US).		(74) Agents: FILLER, Wendy, Anne et al.; Glaxo Holdings plc, Glaxo House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). (81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: METHANESULPHONATE 		
(57) Abstract This invention relates to the novel salt 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrindo[4,3-b]indole-1-one methane sulphonate, to solvates of this salt, to pharmaceutical compositions containing it and to its use in medicine as 5-HT₃ receptor antagonists.		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

METHANESULPHONATE

This invention relates to a novel salt of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one, to solvates of this salt, to pharmaceutical compositions containing it and to its use in medicine.

- 5 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one, which may be represented by the formula (I)



- and its physiologically acceptable salts and solvates are described in European Patent Application publication No. 0353983. The compound of formula (I) is a potent and selective antagonist of 5-hydroxytryptamine (5-HT) at 5-HT₃ receptors and is useful in the treatment of, for example, emesis (i.e. nausea and vomiting), irritable bowel syndrome and gastrointestinal dysfunction such as dyspepsia. Physiologically acceptable salts of the compounds of formula (I) specifically disclosed in European Patent Application publication No. 0353983 are the maleate, benzoate and hydrochloride salts.
- 10
- 15

We have now found, surprisingly, that a novel salt of the compound of formula (I), and solvates, particularly hydrates of this salt, more particularly its dihydrate, is advantageous for the preparation of pharmaceutical compositions by virtue of its solubility and stability.

- 20 The present invention therefore provides in a first aspect 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate and physiologically acceptable solvates thereof.

Preferred solvates of the methanesulphonate salt of the compound of formula (I) are hydrates, in particular the dihydrate.

The preferred compound of the invention is 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate dihydrate.

5 The potent and selective antagonism of 5-HT at 5-HT₃ receptors by the compound of formula (I) has been demonstrated by its ability to inhibit 3-(5-methyl-1H-imidazol-4-yl)-1-[1-(methyl-t₃)-1H-indol-3-yl]-1-propanone binding in rat entorhinal cortex homogenates (following the general procedure described by G. Kilpatrick et al. in Nature, 1987, 330, 746).

10 In view of its 5-HT₃ antagonist activity, 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a physiologically acceptable solvate is useful in the treatment of conditions ameliorated by antagonism of 5-HT at 5-HT₃ receptors, such as emesis (i.e. nausea and vomiting), particularly that associated with cancer chemotherapy and radiotherapy, that occurring post operatively, and also that
15 induced by opioids; cognitive disorders such as dementia, particularly degenerative dementia (including senile dementia, Alzheimer's disease, Pick's disease, Huntington's chorea, Parkinson's disease and Creutzfeldt-Jakob disease), and vascular dementia (including multi-infarct dementia), as well as dementia associated with intracranial space occupying lesions, trauma,
20 infections and related conditions (including HIV infection), metabolism, toxins, anoxia and vitamin deficiency; and mild cognitive impairment associated with ageing, particularly Age Associated Memory Impairment; psychotic disorders, such as schizophrenia and mania; anxiety disorders, including panic disorder, agoraphobia, social phobia, simple phobia, obsessive compulsive disorders,
25 post traumatic stress disorder, mixed anxiety and depression, and generalized anxiety disorder; irritable bowel syndrome; gastric stasis; symptoms of gastrointestinal dysfunction such as occur with dyspepsia, peptic ulcer, reflux oesophagitis and flatulence; migraine; obesity and conditions such as bulimia; pain; dependency on drugs and substances of abuse; and depression.

30 There is also provided as a further aspect of the invention 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a physiologically acceptable solvate thereof for use in therapy, in particular in human medicine. It will be appreciated that use in

3.

therapy embraces but is not necessarily limited to use of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a physiologically acceptable solvate thereof as an active therapeutic substance.

- 5 It will be appreciated that reference to treatment is intended to include prophylaxis as well as the alleviation of established symptoms.

In an alternative aspect of the invention there is provided the use of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a physiologically acceptable
10 solvate thereof in the preparation of a medicament for use in the treatment of a human or animal subject suffering from a condition ameliorated by antagonism of 5-HT at 5-HT₃ receptors.

According to another aspect of the invention there is provided a method of treating a human or animal subject suffering from a condition ameliorated by
15 antagonism of 5-HT at 5-HT₃ receptors, which method comprises administering to said subject an effective amount of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a physiologically acceptable solvate thereof.

The anhydrous methanesulphonate salt of the compound of formula (I) is also of
20 use as an intermediate in the preparation of solvates thereof.

While it is possible that, for use in therapy, the compounds of the invention may be administered as the raw chemical it is preferable to present combinations as a pharmaceutical composition.

There is thus provided in a further aspect of the invention a pharmaceutical
25 composition comprising 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a physiologically acceptable solvate, for example a hydrate, such as a dihydrate thereof as active ingredient, together with a physiologically acceptable carrier, for use in human or veterinary medicine, and formulated for administration by
30 any convenient route.

Such compositions may be formulated in conventional manner using one or more physiologically acceptable carriers and/or excipients.

Thus the compound according to the invention or a physiologically acceptable solvate thereof may be formulated for oral, buccal, parenteral, transdermal or rectal administration or in a form suitable for administration by inhalation or insufflation (either through the mouth or nose).

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. Avicel PH102, lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); non-aqueous vehicles (e.g. almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g. methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavouring, colouring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

Tablets for oral administration may be suitably formulated to provide very fast release of the active compound, for example by formulation as a lyophilised tablet.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

5.

5 The compound of the invention or a physiologically acceptable solvate thereof may be formulated for parenteral administration by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form e.g. in ampoules, vials, syringes or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilising and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

10 A preferred composition for injection is a solution, particularly an aqueous solution. Such solutions may additionally contain other excipients, such as preservatives (for example methyl p-hydroxybenzoate or propyl p-hydroxybenzoate), buffering agents or isotonicity-adjusting agents (for example dextrose, sodium chloride or mannitol).

15 Preferably solutions according to the invention will be sterile, free from particulates and free from preservatives. Sterile formulations may be prepared by methods well known in the art, for example by aseptic manufacture or sterilisation of bulk products.

20 Aqueous solutions of the methanesulphonate dihydrate of the present invention adapted for injection will have a pH in the range 2 to 7. Most preferably the pH of aqueous solutions of the methanesulphonate dihydrate of the present invention will be 3 to 5, such as 3.9 to 4.1. Adjustment of the pH of aqueous solutions of the methanesulphonate dihydrate of the compound of formula (I) to within the desired range is conveniently effected by addition of an acid and/or
25 base. Suitable acids include inorganic acids, for example hydrochloric acid; suitable bases include inorganic bases, such as alkali metal hydroxides, for example sodium hydroxide.

For transdermal administration the compounds of the invention may be formulated as creams, gels, ointments or lotions or as transdermal patches.
30 Such compositions may, for example, be formulated with an aqueous or oily base with the addition of suitable thickening, gelling, emulsifying, stabilising, dispersing, suspending, and/or colouring agents.

The compound of the invention or a physiologically acceptable solvate thereof may also be formulated in rectal compositions such as suppositories or retention enemas, e.g. containing conventional suppository bases such as cocoa butter or other glycerides.

- 5 In addition to the formulations described previously, the compound of the invention or a physiologically acceptable solvate thereof may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compound of the invention or a
10 physiologically acceptable solvate thereof may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives.

- For administration by inhalation the compound according to the invention or a physiologically acceptable solvate thereof is conveniently delivered in the form
15 of an aerosol spray presentation from pressurised packs or a nebuliser, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurised aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and
20 cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound of the invention or a physiologically acceptable solvate thereof and a suitable powder base such as lactose or starch.

- For intranasal administration, the compound according to the invention or a
25 physiologically acceptable solvate thereof may be formulated as solutions for administration via a suitable metered or unit dose device or alternatively as a powder mix with a suitable carrier for administration using a suitable delivery device.

- The compound of the invention or a physiologically acceptable solvate thereof
30 may also be administered in combination with other therapeutic agents. Thus, for example, in the treatment of gastric stasis, symptoms of gastrointestinal dysfunction and emesis (i.e. nausea and vomiting), the compound of the

invention or a physiologically acceptable salt thereof may be administered in combination with antisecretory agents such as histamine H₂-receptor antagonists (e.g. ranitidine, sufofodine, cimetidine, famotidine, nizatidine or roxatidine) or H⁺K⁺ATPase inhibitors (e.g. omeprazole). In the treatment of emesis (i.e. nausea and vomiting), the compound of the invention or a physiologically acceptable solvate thereof may also be administered in combination with dexamethasone or a cyclo-oxygenase inhibitor such as piroxicam.

A proposed dose of the compounds of the invention for administration to man (of approximately 70kg body weight) is 0.001 to 100mg, preferably 0.01 to 50mg, more preferably 0.1 to 20mg of the active ingredient per unit dose expressed as the weight of free base, which could be administered, for example, 1 to 4 times per day. It will be appreciated that it may be necessary to make routine variations to the dosage, depending on the age and condition of the patient and the nature of the condition to be treated, and will be at the ultimate discretion of the attendant physician. The dosage will also depend on the route of administration.

The compound of the invention and physiologically acceptable solvates thereof may be prepared by the general methods outlined hereinafter.

According to one general process (A), the compound of the invention or a solvate thereof may be prepared by reaction of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one or a solvate thereof with methanesulphonic acid. The reaction may be carried out in the presence of a solvent, such as an alcohol (for example Industrial Methylated Spirits (IMS) or ethanol), or an aqueous alcohol (for example aqueous propan-1-ol) and preferably at elevated temperature (for example reflux).

According to one embodiment of process (A), where the compound of the invention is desired in anhydrous form, it is preferably prepared by admixture of the free base and methanesulphonic acid in IMS, at reflux. Preferably, the free base of the compound of the invention is recrystallised from methanol prior to use as a starting material.

According to another embodiment of process (A), where the compound of the invention is desired in hydrated form, in particular as its dihydrate, it is preferably prepared by admixture of the free base and methanesulphonic acid in aqueous propan-1-ol at a temperature preferably in the range 20°C-90°C. Most preferably the reaction will be carried out at a temperature of between 40°C-80°C, such as 60°-70°C.

According to another general process (B), a solvate of the compound of the invention may be prepared by reaction of the compound of the invention in anhydrous form or a solvate thereof with an appropriate solvent, if necessary at elevated temperature.

For example the hydrate, particularly the dihydrate of the compound of the invention may be prepared by reaction of the compound of the invention or a solvate thereof with water, if necessary at elevated temperature.

Such processes (A), and (B) form further aspects of the invention.

The following non-limiting examples further illustrate the invention.

Intermediate 1

6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one

6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one was dissolved in methanol (30ml) at reflux. Water (15ml) was added to the hot solution at 60° maintaining a clear solution. The solution was cooled for 1 hour at 0° and the solid was filtered off, washed with methanol:water (2:1, 15ml) at room temperature. The solid was dried in vacuo at 60° to constant weight to give a second polymorphic form of the title compound as a white solid (3.7g).

IR (Nujol Mull): 3208cm⁻¹, 1633cm⁻¹

Water Analysis Found: 0.5% w/w

Example 1

6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate

5 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one (224.9g) was suspended in IMS (1125ml) and heated to reflux. A solution of methanesulphonic acid (69.2g) in IMS (675ml) was added with stirring over 15 minutes. The mixture was subsequently allowed to cool over 3 hours, then seeded and left overnight at ambient temperature before being cooled to 4°C for 5 hours, with stirring. The mixture was filtered, the solid
10 washed with cold IMS (2 x 450ml), and then dried between 40 and 76° C in vacuo to give the title compound (226g) as a crystalline solid.

Water Analysis Found: 0.8% w/w=0.20mol H₂O
Analysis Found: C, 52.5; H, 5.2; N, 13.5;
C₁₇H₁₇FN₄O.CH₄O₃S Requires: C, 52.9; H, 5.2; N, 13.7%.

15

Example 2

6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate dihydrate

To 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-
20 1H-pyrido[4,3-b]indol-1-one methanesulphonate (33.0g) was added water (112ml) and the mixture heated to 45°C to give a clear yellow solution. This solution was allowed to reach ambient temperature, then cooled to 10°C, whereupon the mixture was filtered and the solid washed with water (20ml, at 10°C). The solid was dried at 48°C in vacuo for 16 hours to give the title
25 compound (20.3g).

Water Analysis Found: 8.2% w/w=2.02 mol H₂O
Analysis Found: C, 48.8; H, 5.7; N, 12.6; S, 7.2;
C₁₇H₁₇FN₄O.CH₄O₃S.2H₂O Requires: C, 48.6; H, 5.7; N, 12.6; S, 7.2%

Example 36-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate dihydrate

Water (0.29l) and methanesulphonic acid (0.23l) was added to a suspension of
 5 6-fluoro- 2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-
 pyrido[4,3-b]indol-1-one (1.04kg) in propan-1-ol (7.4l). The mixture was heated
 to about 65°C to give a clear solution, which was filtered, and the filter rinsed
 with propan-1-ol. The combined filtrate and wash were cooled to about 15°C
 and diisopropyl ether (7.9l) was added to the resulting suspension. The mixture
 10 was cooled to 5°C, and the solid isolated by filtration, washed with diisopropyl
 ether:propan-1-ol (3:1, 2x1.8l) and diisopropyl ether (2x2l) and finally dried in vacuo
 at about 35°C to yield the title compound (1.39kg).

The analysis and water analysis for this material were consistent with those
 obtained for the product of Example 2.

15

Example 4STERILE FORMULATION

	<u>mg/ml</u>
Compound of formula (I),	
methanesulphonate dihydrate	1.42mg
Dextrose Monohydrate USP	55.00mg
Hydrochloric Acid NF	adjust to pH4
Sodium Hydroxide NF	adjust to pH4
Water for Injection USP qs to	1ml

The compound of formula (I) methanesulphonate dihydrate was dissolved in
 water and the solution made up to approximately 85% of volume. Dextrose
 20 Monohydrate USP was added and the pH of the solution was adjusted to 4.0
 using hydrochloric acid and/or sodium hydroxide. The solution was made up to
 final volume with water and the pH remeasured and adjusted if necessary, to
 provide 1mg/ml of the compound of formula (I).

The solution may be packaged for injection, for example by filling and sealing in ampoules, vials or syringes. The ampoules, vials or syringes may be aseptically filled and/or terminally sterilised by, for example, autoclaving at 121°C.

- 5 Further sterile formulations were prepared, in a similar manner, containing 0.142mg and 14.23 mg of the compound of formula (I) methanesulphonate dihydrate, so as to provide 0.1mg/ml and 10mg/ml respectively of the compound of formula (I)

TABLETS FOR ORAL ADMINISTRATION

- 10 Tablets may be prepared by the normal methods such as direct compression or wet granulation.

The tablets may be film coated with suitable film forming materials, such as Opadry White, type YS-1-7027, using standard techniques. Alternatively the tablets may be sugar coated.

- 15 Example 5

Direct Compression

<u>Tablet</u>	<u>mg/Tablet</u>
Compound of formula (I),	
methanesulphonate dihydrate	0.71mg
Magnesium Stearate	0.75mg
Avicel PH102	qs 150.00mg

- 20 The compound of formula (I) methanesulphonate dihydrate is passed through a 30 mesh sieve and blended with Avicel PH102 and magnesium stearate. The resultant mix is compressed into tablets using a suitable tablet machine fitted with 9/32" diameter punches.

Tablets of other strengths, containing for example 0.142, 2.85 or 11.38 mg/tablet of the compound of formula (I) methanesulphonate dihydrate, may be

prepared in a similar manner, so as to provide 0.1, 2 or 8mg/tablet of the compound of formula (I).

Example 6

5 WET GRANULATION

A formulation as described in Example 5 may be used. The compound of formula (I) methanesulphonate dihydrate is dissolved in a suitable volume of granulating solution (purified water or 10% PVP K29/32 in water). After drying, the granules are screened, for example through 20 mesh screen, and blended with magnesium shearate. The granules are then compressed into tablets as described in Example 5.

Tablets of other strengths, such as those described in Example 5, may be prepared in a similar manner.

Example 7

15 SUPPOSITORY

Compound of formula (I),	14.23mg
methanesulphonate dihydrate	
Witepsol W32, hard fat	2000mg
qs	

Blend micronized drug in a portion of the melted Witepsol W32 at approximately 36°C for approximately 15 minutes in a high speed mixer. Incorporate the homogenized slurry into the remaining portion of the melted Witepsol W32 and blend at approximately 36°C until satisfactory dispersion is achieved. Fill moulds with 2000mg formulation, to provide 10mg/suppository of compound of formula (I).

Example 8CAPSULE

	<u>mg/capsule</u>
Compound of formula (I), methanesulphonate dihydrate	14.23mg
Polyethylene glycol	92.89mg
Propylene glycol qs	200mg

- 5 Blend together polyethylene glycol and propylene glycol using heat as necessary. Stir until homogeneous. Add micronised compound of formula (I) methanesulphonate dihydrate to blend. Mix until homogenous. Fill into an appropriate gelatin mass to give soft gelatin capsules containing 200mg of the formulation, to provide 10mg/capsule of compound of formula (I).

Example 910 ORAL SYRUP

	<u>mg/ml</u>
Compound of formula (I), methanesulphonate dihydrate	2.85mg
Sucrose	200mg
Methylparaben	1.2mg
Propylparaben	0.15mg
Flavouring	1.5mg
Citric Acid	0.1mg
Purified Water qs	1 ml

Dissolve the parabens in a small portion of the water that has been heated to approximately 90°C. Add the paraben solution to a large portion of the remaining water with mixing. Add and dissolve the other components. Bring the

formulation to final volume and mix until homogenous. Fill the formulation into a containing, such as a unit dose cup or a bottle for multiple-dose use, to provide 2mg/5ml.

Example 10

5 TRANSDERMAL SYSTEM

Compound of formula (I), methanesulphonate dihydrate	5% (of compound of formula (I))
Silicone fluid	90%
Colloidal silicone dioxide	5%

10 The silicone fluid and drug are mixed together and the colloidal silicone dioxide is added to increase the viscosity. The material is then dosed into a subsequently heat sealed polymeric laminate comprised of the following: polyester release liner, skin contact adhesive composed of silicone or acrylic polymers, a control membrane which is a polyolefin (e.g. polyethylene or polyvinyl acetate) or polyurethane, and an impermeable backing membrane of a polyester multilaminate.

Example 11

15 LYOPHILIZED PRODUCT

Compound of formula (I), methanesulphonate dihydrate	14.23mg
Mannitol	50.00mg
Citrate buffer	0.75mg
Water for injection	qs 1ml

20 Dissolve components in a portion of the water for injection. Make formulation up to final volume and mix until homogenous. Filter formulation through a sterilising filter and fill into glass vials. Lyophilize and seal vials. Reconstitute with appropriate solvent prior to use.

Claims

1. 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate, and solvates thereof.
- 5 2. 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate in the form of a hydrate.
3. 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate dihydrate.
- 10 4. A pharmaceutical composition comprising 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a solvate thereof as defined in any one of Claims 1 to 3 together with at least one physiologically acceptable carrier or excipient.
- 15 5. A pharmaceutical composition according to Claim 4 in a form adapted for oral or parenteral administration.
- 20 6. A process for the preparation of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate and solvates thereof, which comprises:
 - 25 (A) reaction of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate indol-1-one or a solvate thereof with methanesulphonic acid; or
 - (B) for the preparation of a solvate of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate, reaction of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a solvate thereof with an appropriate solvent.
- 30

7. 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a solvate thereof as defined in any of Claims 1 to 3 for use in therapy.
- 5 8. The use of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a solvate thereof as defined in any of Claims 1 to 3 for the manufacture of a medicament for the treatment or prophylaxis of a condition ameliorated by antagonism of 5-HT at 5-HT₃ receptors.
- 10 9. The use of 6-Fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a solvate thereof according to Claim 8 wherein the condition which may be treated is nausea or vomiting.
- 15 10. A method of treating a human or animal subject suffering from a condition ameliorated by antagonism of 5-HT at 5-HT₃ receptors, which method comprises administering to said subject an effective amount of 6-fluoro-2,3,4,5-tetrahydro-5-methyl-2-[(5-methyl-1H-imidazol-4-yl)methyl]-1H-pyrido[4,3-b]indol-1-one methanesulphonate or a solvate thereof as defined
- 20 in any of Claims 1 to 3.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 93/02014

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D471/04 C07C309/04 A61K31/435 //(C07D471/04,221:00,
209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 353 983 (GLAXO) 7 February 1990 cited in the application see page 3, line 54 - line 57; claims 1,9,14 -----	1,2,4

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 November 1993

Date of mailing of the international search report

26. 11. 93

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

ALFARO FAUS, I

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 93/02014

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 10 is directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/EP 93/02014

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0353983	07-02-90	AU-B- 631131	19-11-92
		AU-A- 3918689	08-02-90
		JP-A- 2160786	20-06-90
		US-A- 5183820	02-02-93
