



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/02/28
(87) Date publication PCT/PCT Publication Date: 2018/09/07
(85) Entrée phase nationale/National Entry: 2019/08/27
(86) N° demande PCT/PCT Application No.: US 2018/020335
(87) N° publication PCT/PCT Publication No.: 2018/160774
(30) Priorités/Priorities: 2017/02/28 (US62/464,675);
2017/03/16 (US62/472,468)

(51) Cl.Int./Int.Cl. *A61K 31/395* (2006.01),
A61K 31/505 (2006.01), *A61K 31/506* (2006.01)


(71) Demandeurs/Applicants:
THE GENERAL HOSPITAL CORPORATION, US;
DANA-FARBER CANCER INSTITUTE, INC., US

(72) Inventeurs/Inventors:
MURAKAMI, RYO, JP;
FISHER, DAVID E., US;
MUJAHID, NISMA, US;
GRAY, NATHANAEL S., US;
CHOI, HWAN GEUN, US;
LIANG, YANKE, US

(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : UTILISATIONS DE PYRIMIDOPYRIMIDINONES COMME INHIBITEURS DE SIK

(54) Title: USES OF PYRIMIDOPYRIMIDINONES AS SIK INHIBITORS



(57) Abrégé/Abstract:

The present disclosure provides methods of increasing skin pigmentation in a subject in need thereof using salt-inducible kinase (SIK) inhibitors, such as macrocyclic compounds of Formula (I), bicyclic urea compounds of Formula (II), (III), and (IV), and compounds of Formula (V), (VI), (VI-A), or (VII). Also provided are pharmaceutical compositions, methods, and uses that include or involve a compound described herein.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2018/160774 A1(43) International Publication Date
07 September 2018 (07.09.2018)

(51) International Patent Classification:

A61K 31/395 (2006.01) A61K 31/506 (2006.01)
A61K 31/505 (2006.01)

(21) International Application Number:

PCT/US2018/020335

(22) International Filing Date:

28 February 2018 (28.02.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/464,675 28 February 2017 (28.02.2017) US
62/472,468 16 March 2017 (16.03.2017) US

(71) Applicants: **THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US). **DANA-FARBER CANCER INSTITUTE, INC.** [US/US]; 450 Brookline Avenue, Boston, MA 02215 (US).

(72) Inventor; and

(71) Applicant: **MURAKAMI, Ryo** [JP/JP]; 3-16-1-301, Sen-goku, Bunkyo-ku, Tokyo, 112-0011 (JP).


(72) Inventors: **FISHER, David, E.**; 510 Ward Street, Newton, MA 02459 (US). **MUJAHID, Nisma**; 672 Ogden Parma Townline, Spencerport, NY 14559 (US). **GRAY, Nathanael, S.**; 26 Greenview Avenue, Boston, MA 02130 (US).

(74) Agent: **ZHANG, Wei**; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

(54) Title: USES OF PYRIMIDOPYRIMIDINONES AS SIK INHIBITORS



(57) Abstract: The present disclosure provides methods of increasing skin pigmentation in a subject in need thereof using salt-inducible kinase (SIK) inhibitors, such as macrocyclic compounds of Formula (I), bicyclic urea compounds of Formula (II), (III), and (IV), and compounds of Formula (V), (VI), (VI-A), or (VII). Also provided are pharmaceutical compositions, methods, and uses that include or involve a compound described herein.

[Continued on next page]

WO 2018/160774 A1

UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report (Art. 21(3))*

USES OF PYRIMIDOPYRIMIDINONES AS SIK INHIBITORS

RELATED APPLICATIONS

[0001] This application claims priority under 35 U.S.C. § 119(e) to U.S. Provisional Applications, U.S.S.N. 62/464,675, filed February 28, 2017, and U.S.S.N. 62/472,468, filed March 16, 2017, each of which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] A protein kinase inhibitor is an enzyme inhibitor that blocks the action of a protein kinase. A protein kinase is an enzyme that adds a phosphate group to a protein or other organic molecule. Phosphorylation is involved in a wide range of diseases, such as cancer. One protein kinase is a salt-inducible kinase (SIK), *e.g.*, SIK1, SIK2 or SIK3). SIK's (*e.g.*, SIK1, SIK2 or SIK3) are serine/threonine kinases in the adenosine monophosphate-activated protein kinase (AMPK) family. (*See Altarejos et al.* 2011, Patel *et al.* 2014, Park *et al.* 2014, and Henriksson *et al.* 2015).

[0003] The incidence of non-melanoma and melanoma skin cancers has been increasing in the United States over recent decades (Rogers *et al.* 2015; Ryerson *et al.* 2016; Watson *et al.* 2016). Epidemiological evidence suggests that there is a causal relationship between sun/UV exposure and the three major histologic forms of skin cancer: squamous cell carcinoma, basal cell carcinoma, and cutaneous melanoma (Wu *et al.* 2014; Kennedy *et al.* 2003; Gandini *et al.* 2005). Individuals with fair skin and/or poor tanning ability are at higher risk for developing these malignancies (Armstrong and Kricker 2001), which are uncommon in darkly pigmented individuals (Pennello, Devesa, and Gail 2000). During UV induced tanning, DNA damage in keratinocytes triggers p53 mediated transcription of the pro-opiomelanocortin (POMC) gene (Cui *et al.* 2007). Proteolytic cleavage of POMC, produces alpha-MSH, which binds to the melanocortin-receptor-1 (MC1R) on melanocytes, activating adenylate cyclase. Elevated cAMP activates protein kinase A (PKA), which phosphorylates the cAMP-responsive-element-binding protein (CREB) (Tsatmalia *et al.* 1999; Newton *et al.* 2005), resulting in stimulated transcription of the microphthalmia-associated transcription factor (MITF) gene (Price *et al.* 1998; Bertolotto *et al.* 1998). *MC1R* non-signaling variants are associated with lighter skin tones and red hair, and are linked to poor tanning responses (Valverde *et al.* 1995). Previously, topical application of the cAMP agonist forskolin was shown to rescue the cAMP-MITF-eumelanin pathway in *Mclr* deficient mice (D'Orazio *et al.* 2006). Subsequent studies identified the phosphodiesterase PDE4d3 as a key regulator of

melanocytic cAMP homeostasis, and its suppression produced similar hyperpigmentation as forskolin treatment in redhaired mice (Khaled, Levy, and Fisher 2010). However attempts to apply both of these small molecule approaches to human skin have been unsuccessful, likely related to poor skin penetration of the active species.


[0004] Genetic data in mice have suggested the presence of a pathway in which CREB-regulated transcription co-activator (CRTC) positively and salt-inducible kinase 2 (SIK2) negatively regulate MITF and pigment synthesis independently of CREB phosphorylation by PKA (Horike *et al.* 2010). In macrophages, the small molecule SIK inhibitor HG 9-91-01 has been shown to regulate CREB-dependent gene transcription by suppressing phosphorylation of CRTC (Clark *et al.* 2012), thereby inhibiting cytoplasmic sequestration and permitting its nuclear translocation. It was hypothesized that small molecule SIK inhibitors could be generated and optimized as topical agents capable of inducing cutaneous pigmentation independently of UV irradiation in human skin. There is a need for SIK inhibitors with good skin penetration for such a use in a subject.

SUMMARY OF THE INVENTION


[0005] The presence of dark melanin (eumelanin) within human epidermis represents one of the strongest predictors of protection from skin cancer and UV photodamage (Armstrong and Kricker 2001; Pennello, Devesa, and Gail 2000). Non-genotoxic topical rescue of eumelanin synthesis has been previously achieved in “redhaired” *Mclr* deficient mice and exhibited significant protection against UV damage and skin carcinogenesis (D'Orazio *et al.* 2006). However, application of this topical strategy to human skin has not been achieved, in large part due to the profoundly greater barrier function of human stratum corneum and epidermis. Salt Inducible Kinase (SIK) is a serine threonine kinase (Dentin *et al.* 2007) that has been demonstrated to regulate MITF, the master regulator of pigment gene expression through its effects on CRTC and CREB activity (Horike *et al.* 2010). A flavonoid inhibitor of SIK2 has been reported to promote melanogenesis in B16F10 melanoma cells (*See Kumagai et al.*). Here, the development of small molecule SIK inhibitors that are useful for human skin penetration are described, resulting in MITF up-regulation and induction of melanogenesis. When topically applied, pigment production was induced in redhead mice harboring an inactivating mutation in *Mclr*, as well as in normal human skin. These findings provide a realistic pathway towards topical modulation of human skin pigmentation without a requirement for UV irradiation, potentially impacting UV protection and skin cancer risk.








[0006] Described herein are compounds of Formula (I), (II), (III), (IV), (V), (VI), (VI-A), or (VII) that are salt-inducible kinase (SIK) inhibitors. The compounds include macrocyclic compounds of Formula (I), bicyclic urea compounds of Formulae (II), (III), and (IV), and compounds of Formulae (V), (VI), (VI-A), and (VII), and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. The described compounds are able to inhibit the activity of a salt-inducible kinase (SIK, *e.g.*, SIK1, SIK2, SIK3). Also provided in the present disclosure are methods of increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof using the described compounds (*e.g.*, via topical administration of the described compounds). In certain embodiments, provided herein are methods of increasing the appearance of skin darkening in a subject in need thereof using the described compounds (*e.g.*, via topical administration of the described compounds).

[0007] In one aspect, the compound utilized in the present disclosure is of Formula (I):






and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds of Formula (I) useful in the present invention include, but are not limited to:








and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.


[0008] Exemplary compounds of Formula (I) useful in the present invention include, but are not limited to:



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.


[0009] Exemplary compounds of Formula (I) useful in the present invention include, but are not limited to:







and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.


[0010] In another aspect, the compound utilized in the present disclosure is of Formula **(II)**:




and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds of Formula **(II)** useful in the present invention include, but are not limited to:











and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.


[0011] Exemplary compounds of Formula (II) useful in the present invention include, but are not limited to:




(HG-11-139-01),



(HG-11-143-01),




(YKL-04-136-10),



(YKL-04-136-11),


and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0012] In another aspect, the compound utilized in the present disclosure is of Formula **(III)**:

**(III)**,


and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds

of Formula (III) useful in the present invention include, but are not limited to:








and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0013] In another aspect, the compound utilized in the present disclosure is of Formula (IV):




and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds of Formula (IV) useful in the present invention include, but are not limited to:





and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.


[0014] Exemplary compounds of Formula (IV) useful in the present invention include, but are not limited to:




(HG-11-23-01),

and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0015] In another aspect, the compound utilized in the present disclosure is of Formula (V):


 $(\mathbf{V}),$

and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds of Formula (V) useful in the present invention include, but are not limited to:




and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0016] In another aspect, the compound utilized in the present disclosure is of Formula (VI):






and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0017] In another aspect, the compound utilized in the present disclosure is of Formula (VI-A):




and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds of Formula (VI) and (VI-A) useful in the present invention include, but are not limited to:






(MRT199665),



(KIN112),


and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0018] In another aspect, the compound utilized in the present disclosure is of Formula (VII):




(VII),







and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Exemplary compounds of Formula (VII) useful in the present invention include, but are not limited to:



(VII-5),



(VII-6),



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[0019] The pharmaceutical compositions including a compound described herein, and optionally a pharmaceutically acceptable excipient may be useful in increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof. In certain

embodiments, a pharmaceutical composition described herein includes an effective amount (*e.g.*, a prophylactically effective) amount of a compound described herein. In certain embodiments, the compound being administered or used selectively inhibits the activity of a salt-inducible protein kinase (SIK) (*e.g.*, SIK1, SIK2, or SIK3). In certain embodiments, the compound selectively inhibits SIK1. In certain embodiments, the compound selectively inhibits SIK2.

[0020] In certain embodiments, the subject being administered a compound or pharmaceutical composition described herein is a human. In certain embodiments, the subject being administered a compound or pharmaceutical composition described herein is a non-human animal.

[0021] In another aspect, the present disclosure provides methods of increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof. In another aspect, the present disclosure provides methods of inhibiting SIK in the skin of a subject. In another aspect, the present disclosure provides methods of inducing eumelanin synthesis. In certain embodiments, the present disclosure provides methods of inducing melanosomal maturation, export, and localization. In certain embodiments, the present disclosure provides methods of inducing microphthalmia-associated transcription factor (MITF) expression.

[0022] In still another aspect, described herein are kits including a container with a compound or pharmaceutical composition described herein for use in a method described herein. A kit described herein may include a single dose or multiple doses of the compound or pharmaceutical composition. The described kits may be useful in increasing skin pigmentation in a subject in need thereof. In certain embodiments, a kit described herein further includes instructions for using the kit.

[0023] In another aspect, the present disclosure provides methods of inhibiting the activity of a SIK (*e.g.*, SIK1, SIK2, or SIK3).

[0024] The methods of the present disclosure include administering to the subject an effective amount of a compound or pharmaceutical composition described herein for use in a method of the disclosure (*e.g.*, a method of increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof. In certain embodiments, the method comprises administering topically to the subject's skin an effective amount of a compound or pharmaceutical composition described herein.

[0025] In yet another aspect, the present disclosure provides compounds and pharmaceutical compositions described herein for use in a method of the disclosure (*e.g.*, a

method of increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof.

[0026] The present application refers to various issued patent, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. The details of one or more embodiments of the invention are set forth herein. Other features, objects, and advantages of the invention will be apparent from the Detailed Description, the Examples, and the Claims.

DEFINITIONS

[0027] Definitions of specific functional groups and chemical terms are described in more detail below. The chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, *Handbook of Chemistry and Physics*, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in *Organic Chemistry*, Thomas Sorrell, University Science Books, Sausalito, 1999; Smith and March *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; and Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987.

[0028] Compounds described herein can comprise one or more asymmetric centers, and thus can exist in various stereoisomeric forms, *e.g.*, enantiomers and/or diastereomers. For example, the compounds described herein can be in the form of an individual enantiomer, diastereomer or geometric isomer, or can be in the form of a mixture of stereoisomers, including racemic mixtures and mixtures enriched in one or more stereoisomer. Isomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred isomers can be prepared by asymmetric syntheses. See, for example, Jacques *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, E.L. *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S.H. *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972). The disclosure additionally encompasses compounds as individual isomers substantially free of other isomers, and alternatively, as mixtures of various isomers.

[0029] In a formula, --- is absent or a single bond, and == or == is a single or double bond.

[0030] The term “heteroatom” refers to an atom that is not hydrogen or carbon. In certain embodiments, the heteroatom is nitrogen. In certain embodiments, the heteroatom is oxygen. In certain embodiments, the heteroatom is sulfur.

[0031] When a range of values is listed, it is intended to encompass each value and sub-range within the range. For example “C₁₋₆ alkyl” is intended to encompass, C₁, C₂, C₃, C₄, C₅, C₆, C₁₋₆, C₁₋₅, C₁₋₄, C₁₋₃, C₁₋₂, C₂₋₆, C₂₋₅, C₂₋₄, C₂₋₃, C₃₋₆, C₃₋₅, C₃₋₄, C₄₋₆, C₄₋₅, and C₅₋₆ alkyl.

[0032] The term “aliphatic” refers to alkyl, alkenyl, alkynyl, and carbocyclic groups. Likewise, the term “heteroaliphatic” refers to heteroalkyl, heteroalkenyl, heteroalkynyl, and heterocyclic groups.


[0033] The term “alkyl” refers to a radical of a straight-chain or branched saturated hydrocarbon group having from 1 to 10 carbon atoms (“C₁₋₁₀ alkyl”). In some embodiments, an alkyl group has 1 to 9 carbon atoms (“C₁₋₉ alkyl”). In some embodiments, an alkyl group has 1 to 8 carbon atoms (“C₁₋₈ alkyl”). In some embodiments, an alkyl group has 1 to 7 carbon atoms (“C₁₋₇ alkyl”). In some embodiments, an alkyl group has 1 to 6 carbon atoms (“C₁₋₆ alkyl”). In some embodiments, an alkyl group has 1 to 5 carbon atoms (“C₁₋₅ alkyl”). In some embodiments, an alkyl group has 1 to 4 carbon atoms (“C₁₋₄ alkyl”). In some embodiments, an alkyl group has 1 to 3 carbon atoms (“C₁₋₃ alkyl”). In some embodiments, an alkyl group has 1 to 2 carbon atoms (“C₁₋₂ alkyl”). In some embodiments, an alkyl group has 1 carbon atom (“C₁ alkyl”). In some embodiments, an alkyl group has 2 to 6 carbon atoms (“C₂₋₆ alkyl”). Examples of C₁₋₆ alkyl groups include methyl (C₁), ethyl (C₂), n-propyl (C₃), isopropyl (C₃), n-butyl (C₄), tert-butyl (C₄), sec-butyl (C₄), iso-butyl (C₄), n-pentyl (C₅), 3-pentanyl (C₅), amyl (C₅), neopentyl (C₅), 3-methyl-2-butanyl (C₅), tertiary amyl (C₅), and n-hexyl (C₆). Additional examples of alkyl groups include n-heptyl (C₇), n-octyl (C₈) and the like. Unless otherwise specified, each instance of an alkyl group is independently unsubstituted (an “unsubstituted alkyl”) or substituted (a “substituted alkyl”) with one or more substituents. In certain embodiments, the alkyl group is an unsubstituted C₁₋₁₀ alkyl (*e.g.*, -CH₃). In certain embodiments, the alkyl group is a substituted C₁₋₁₀ alkyl.

[0034] The term “haloalkyl” is a substituted alkyl group, wherein one or more of the hydrogen atoms are independently replaced by a halogen, *e.g.*, fluoro, bromo, chloro, or iodo. In some embodiments, the haloalkyl moiety has 1 to 8 carbon atoms (“C₁₋₈ haloalkyl”). In some embodiments, the haloalkyl moiety has 1 to 6 carbon atoms (“C₁₋₆ haloalkyl”). In some

embodiments, the haloalkyl moiety has 1 to 4 carbon atoms (“C₁₋₄ haloalkyl”). In some embodiments, the haloalkyl moiety has 1 to 3 carbon atoms (“C₁₋₃ haloalkyl”). In some embodiments, the haloalkyl moiety has 1 to 2 carbon atoms (“C₁₋₂ haloalkyl”). Examples of haloalkyl groups include –CHF₂, –CH₂F, –CF₃, –CH₂CF₃, –CF₂CF₃, –CF₂CF₂CF₃, –CCl₃, –CFCl₂, –CF₂Cl, and the like.

[0035] The term “heteroalkyl” refers to an alkyl group, which further includes at least one heteroatom (*e.g.*, 1, 2, 3, or 4 heteroatoms) selected from oxygen, nitrogen, or sulfur within (*i.e.*, inserted between adjacent carbon atoms of) and/or placed at one or more terminal position(s) of the parent chain. In certain embodiments, a heteroalkyl group refers to a saturated group having from 1 to 10 carbon atoms and 1 or more heteroatoms within the parent chain (“heteroC₁₋₁₀ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 9 carbon atoms and 1 or more heteroatoms within the parent chain (“heteroC₁₋₉ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 8 carbon atoms and 1 or more heteroatoms within the parent chain (“heteroC₁₋₈ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 7 carbon atoms and 1 or more heteroatoms within the parent chain (“heteroC₁₋₇ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 6 carbon atoms and 1 or more heteroatoms within the parent chain (“heteroC₁₋₆ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 5 carbon atoms and 1 or 2 heteroatoms within the parent chain (“heteroC₁₋₅ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 4 carbon atoms and 1 or 2 heteroatoms within the parent chain (“heteroC₁₋₄ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 3 carbon atoms and 1 heteroatom within the parent chain (“heteroC₁₋₃ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 to 2 carbon atoms and 1 heteroatom within the parent chain (“heteroC₁₋₂ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 1 carbon atom and 1 heteroatom (“heteroC₁ alkyl”). In some embodiments, a heteroalkyl group is a saturated group having 2 to 6 carbon atoms and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₆ alkyl”). Unless otherwise specified, each instance of a heteroalkyl group is independently unsubstituted (an “unsubstituted heteroalkyl”) or substituted (a “substituted heteroalkyl”) with one or more substituents. In certain embodiments, the heteroalkyl group is an unsubstituted heteroC₁₋₁₀ alkyl. In certain embodiments, the heteroalkyl group is a substituted heteroC₁₋₁₀ alkyl.

[0036] The term “alkenyl” refers to a radical of a straight-chain or branched hydrocarbon group having from 2 to 10 carbon atoms and one or more carbon-carbon double bonds (*e.g.*,

1, 2, 3, or 4 double bonds). In some embodiments, an alkenyl group has 2 to 9 carbon atoms (“C₂₋₉ alkenyl”). In some embodiments, an alkenyl group has 2 to 8 carbon atoms (“C₂₋₈ alkenyl”). In some embodiments, an alkenyl group has 2 to 7 carbon atoms (“C₂₋₇ alkenyl”). In some embodiments, an alkenyl group has 2 to 6 carbon atoms (“C₂₋₆ alkenyl”). In some embodiments, an alkenyl group has 2 to 5 carbon atoms (“C₂₋₅ alkenyl”). In some embodiments, an alkenyl group has 2 to 4 carbon atoms (“C₂₋₄ alkenyl”). In some embodiments, an alkenyl group has 2 to 3 carbon atoms (“C₂₋₃ alkenyl”). In some embodiments, an alkenyl group has 2 carbon atoms (“C₂ alkenyl”). The one or more carbon-carbon double bonds can be internal (such as in 2-butenyl) or terminal (such as in 1-butenyl). Examples of C₂₋₄ alkenyl groups include ethenyl (C₂), 1-propenyl (C₃), 2-propenyl (C₃), 1-butenyl (C₄), 2-butenyl (C₄), butadienyl (C₄), and the like. Examples of C₂₋₆ alkenyl groups include the aforementioned C₂₋₄ alkenyl groups as well as pentenyl (C₅), pentadienyl (C₅), hexenyl (C₆), and the like. Additional examples of alkenyl include heptenyl (C₇), octenyl (C₈), octatrienyl (C₈), and the like. Unless otherwise specified, each instance of an alkenyl group is independently unsubstituted (an “unsubstituted alkenyl”) or substituted (a “substituted alkenyl”) with one or more substituents. In certain embodiments, the alkenyl group is an unsubstituted C₂₋₁₀ alkenyl. In certain embodiments, the alkenyl group is a substituted C₂₋₁₀ alkenyl. In an alkenyl group, a C=C double bond for which the stereochemistry is unspecified (*e.g.*, –CH=CHCH₃ or ) may be an (*E*)- or (*Z*)-double bond.

[0037] The term “heteroalkenyl” refers to an alkenyl group, which further includes at least one heteroatom (*e.g.*, 1, 2, 3, or 4 heteroatoms) selected from oxygen, nitrogen, or sulfur within (*i.e.*, inserted between adjacent carbon atoms of) and/or placed at one or more terminal position(s) of the parent chain. In certain embodiments, a heteroalkenyl group refers to a group having from 2 to 10 carbon atoms, at least one double bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₁₀ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 9 carbon atoms at least one double bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₉ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 8 carbon atoms, at least one double bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₈ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 7 carbon atoms, at least one double bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₇ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 6 carbon atoms, at least one double bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₆ alkenyl”). In

some embodiments, a heteroalkenyl group has 2 to 5 carbon atoms, at least one double bond, and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₅ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 4 carbon atoms, at least one double bond, and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₄ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 3 carbon atoms, at least one double bond, and 1 heteroatom within the parent chain (“heteroC₂₋₃ alkenyl”). In some embodiments, a heteroalkenyl group has 2 to 6 carbon atoms, at least one double bond, and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₆ alkenyl”). Unless otherwise specified, each instance of a heteroalkenyl group is independently unsubstituted (an “unsubstituted heteroalkenyl”) or substituted (a “substituted heteroalkenyl”) with one or more substituents. In certain embodiments, the heteroalkenyl group is an unsubstituted heteroC₂₋₁₀ alkenyl. In certain embodiments, the heteroalkenyl group is a substituted heteroC₂₋₁₀ alkenyl.

[0038] The term “alkynyl” refers to a radical of a straight-chain or branched hydrocarbon group having from 2 to 10 carbon atoms and one or more carbon-carbon triple bonds (*e.g.*, 1, 2, 3, or 4 triple bonds) (“C₂₋₁₀ alkynyl”). In some embodiments, an alkynyl group has 2 to 9 carbon atoms (“C₂₋₉ alkynyl”). In some embodiments, an alkynyl group has 2 to 8 carbon atoms (“C₂₋₈ alkynyl”). In some embodiments, an alkynyl group has 2 to 7 carbon atoms (“C₂₋₇ alkynyl”). In some embodiments, an alkynyl group has 2 to 6 carbon atoms (“C₂₋₆ alkynyl”). In some embodiments, an alkynyl group has 2 to 5 carbon atoms (“C₂₋₅ alkynyl”). In some embodiments, an alkynyl group has 2 to 4 carbon atoms (“C₂₋₄ alkynyl”). In some embodiments, an alkynyl group has 2 to 3 carbon atoms (“C₂₋₃ alkynyl”). In some embodiments, an alkynyl group has 2 carbon atoms (“C₂ alkynyl”). The one or more carbon-carbon triple bonds can be internal (such as in 2-butyne) or terminal (such as in 1-butyne). Examples of C₂₋₄ alkynyl groups include, without limitation, ethynyl (C₂), 1-propynyl (C₃), 2-propynyl (C₃), 1-butyne (C₄), 2-butyne (C₄), and the like. Examples of C₂₋₆ alkenyl groups include the aforementioned C₂₋₄ alkynyl groups as well as pentynyl (C₅), hexynyl (C₆), and the like. Additional examples of alkynyl include heptynyl (C₇), octynyl (C₈), and the like. Unless otherwise specified, each instance of an alkynyl group is independently unsubstituted (an “unsubstituted alkynyl”) or substituted (a “substituted alkynyl”) with one or more substituents. In certain embodiments, the alkynyl group is an unsubstituted C₂₋₁₀ alkynyl. In certain embodiments, the alkynyl group is a substituted C₂₋₁₀ alkynyl.

[0039] The term “heteroalkynyl” refers to an alkynyl group, which further includes at least one heteroatom (*e.g.*, 1, 2, 3, or 4 heteroatoms) selected from oxygen, nitrogen, or sulfur within (*i.e.*, inserted between adjacent carbon atoms of) and/or placed at one or more terminal

position(s) of the parent chain. In certain embodiments, a heteroalkynyl group refers to a group having from 2 to 10 carbon atoms, at least one triple bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₁₀ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 9 carbon atoms, at least one triple bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₉ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 8 carbon atoms, at least one triple bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₈ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 7 carbon atoms, at least one triple bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₇ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 6 carbon atoms, at least one triple bond, and 1 or more heteroatoms within the parent chain (“heteroC₂₋₆ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 5 carbon atoms, at least one triple bond, and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₅ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 4 carbon atoms, at least one triple bond, and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₄ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 3 carbon atoms, at least one triple bond, and 1 heteroatom within the parent chain (“heteroC₂₋₃ alkynyl”). In some embodiments, a heteroalkynyl group has 2 to 6 carbon atoms, at least one triple bond, and 1 or 2 heteroatoms within the parent chain (“heteroC₂₋₆ alkynyl”). Unless otherwise specified, each instance of a heteroalkynyl group is independently unsubstituted (an “unsubstituted heteroalkynyl”) or substituted (a “substituted heteroalkynyl”) with one or more substituents. In certain embodiments, the heteroalkynyl group is an unsubstituted heteroC₂₋₁₀ alkynyl. In certain embodiments, the heteroalkynyl group is a substituted heteroC₂₋₁₀ alkynyl.

[0040] The term “carbocyclyl” or “carbocyclic” refers to a radical of a non-aromatic cyclic hydrocarbon group having from 3 to 14 ring carbon atoms (“C₃₋₁₄ carbocyclyl”) and zero heteroatoms in the non-aromatic ring system. In some embodiments, a carbocyclyl group has 3 to 10 ring carbon atoms (“C₃₋₁₀ carbocyclyl”). In some embodiments, a carbocyclyl group has 3 to 8 ring carbon atoms (“C₃₋₈ carbocyclyl”). In some embodiments, a carbocyclyl group has 3 to 7 ring carbon atoms (“C₃₋₇ carbocyclyl”). In some embodiments, a carbocyclyl group has 3 to 6 ring carbon atoms (“C₃₋₆ carbocyclyl”). In some embodiments, a carbocyclyl group has 4 to 6 ring carbon atoms (“C₄₋₆ carbocyclyl”). In some embodiments, a carbocyclyl group has 5 to 6 ring carbon atoms (“C₅₋₆ carbocyclyl”). In some embodiments, a carbocyclyl group has 5 to 10 ring carbon atoms (“C₅₋₁₀ carbocyclyl”). Exemplary C₃₋₆ carbocyclyl groups include, without limitation, cyclopropyl (C₃), cyclopropenyl (C₃), cyclobutyl (C₄), cyclobutenyl (C₄), cyclopentyl (C₅), cyclopentenyl (C₅), cyclohexyl (C₆),

cyclohexenyl (C₆), cyclohexadienyl (C₆), and the like. Exemplary C₃₋₈ carbocyclyl groups include, without limitation, the aforementioned C₃₋₆ carbocyclyl groups as well as cycloheptyl (C₇), cycloheptenyl (C₇), cycloheptadienyl (C₇), cycloheptatrienyl (C₇), cyclooctyl (C₈), cyclooctenyl (C₈), bicyclo[2.2.1]heptanyl (C₇), bicyclo[2.2.2]octanyl (C₈), and the like. Exemplary C₃₋₁₀ carbocyclyl groups include, without limitation, the aforementioned C₃₋₈ carbocyclyl groups as well as cyclononyl (C₉), cyclononenyl (C₉), cyclodecyl (C₁₀), cyclodecenyl (C₁₀), octahydro-1*H*-indenyl (C₉), decahydronaphthalenyl (C₁₀), spiro[4.5]decanyl (C₁₀), and the like. As the foregoing examples illustrate, in certain embodiments, the carbocyclyl group is either monocyclic (“monocyclic carbocyclyl”) or polycyclic (*e.g.*, containing a fused, bridged or spiro ring system such as a bicyclic system (“bicyclic carbocyclyl”) or tricyclic system (“tricyclic carbocyclyl”)) and can be saturated or can contain one or more carbon-carbon double or triple bonds. “Carbocyclyl” also includes ring systems wherein the carbocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups wherein the point of attachment is on the carbocyclyl ring, and in such instances, the number of carbons continue to designate the number of carbons in the carbocyclic ring system. Unless otherwise specified, each instance of a carbocyclyl group is independently unsubstituted (an “unsubstituted carbocyclyl”) or substituted (a “substituted carbocyclyl”) with one or more substituents. In certain embodiments, the carbocyclyl group is an unsubstituted C₃₋₁₄ carbocyclyl. In certain embodiments, the carbocyclyl group is a substituted C₃₋₁₄ carbocyclyl.

[0041] In some embodiments, “carbocyclyl” is a monocyclic, saturated carbocyclyl group having from 3 to 14 ring carbon atoms (“C₃₋₁₄ cycloalkyl”). In some embodiments, a cycloalkyl group has 3 to 10 ring carbon atoms (“C₃₋₁₀ cycloalkyl”). In some embodiments, a cycloalkyl group has 3 to 8 ring carbon atoms (“C₃₋₈ cycloalkyl”). In some embodiments, a cycloalkyl group has 3 to 6 ring carbon atoms (“C₃₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group has 4 to 6 ring carbon atoms (“C₄₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group has 5 to 6 ring carbon atoms (“C₅₋₆ cycloalkyl”). In some embodiments, a cycloalkyl group has 5 to 10 ring carbon atoms (“C₅₋₁₀ cycloalkyl”). Examples of C₅₋₆ cycloalkyl groups include cyclopentyl (C₅) and cyclohexyl (C₆). Examples of C₃₋₆ cycloalkyl groups include the aforementioned C₅₋₆ cycloalkyl groups as well as cyclopropyl (C₃) and cyclobutyl (C₄). Examples of C₃₋₈ cycloalkyl groups include the aforementioned C₃₋₆ cycloalkyl groups as well as cycloheptyl (C₇) and cyclooctyl (C₈). Unless otherwise specified, each instance of a cycloalkyl group is independently unsubstituted (an “unsubstituted cycloalkyl”) or substituted (a “substituted cycloalkyl”) with one or more substituents. In

certain embodiments, the cycloalkyl group is an unsubstituted C₃₋₁₄ cycloalkyl. In certain embodiments, the cycloalkyl group is a substituted C₃₋₁₄ cycloalkyl.

[0042] The term “heterocyclyl” or “heterocyclic” refers to a radical of a 3- to 14-membered non-aromatic ring system having ring carbon atoms and 1 to 4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“3-14 membered heterocyclyl”). In heterocyclyl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. A heterocyclyl group can either be monocyclic (“monocyclic heterocyclyl”) or polycyclic (*e.g.*, a fused, bridged or spiro ring system such as a bicyclic system (“bicyclic heterocyclyl”) or tricyclic system (“tricyclic heterocyclyl”)), and can be saturated or can contain one or more carbon-carbon double or triple bonds. Heterocyclyl polycyclic ring systems can include one or more heteroatoms in one or both rings. “Heterocyclyl” also includes ring systems wherein the heterocyclyl ring, as defined above, is fused with one or more carbocyclyl groups wherein the point of attachment is either on the carbocyclyl or heterocyclyl ring, or ring systems wherein the heterocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups, wherein the point of attachment is on the heterocyclyl ring, and in such instances, the number of ring members continue to designate the number of ring members in the heterocyclyl ring system. Unless otherwise specified, each instance of heterocyclyl is independently unsubstituted (an “unsubstituted heterocyclyl”) or substituted (a “substituted heterocyclyl”) with one or more substituents. In certain embodiments, the heterocyclyl group is an unsubstituted 3-14 membered heterocyclyl. In certain embodiments, the heterocyclyl group is a substituted 3-14 membered heterocyclyl.

[0043] In some embodiments, a heterocyclyl group is a 5-10 membered non-aromatic ring system having ring carbon atoms and 1-4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5-10 membered heterocyclyl”). In some embodiments, a heterocyclyl group is a 5-8 membered non-aromatic ring system having ring carbon atoms and 1-4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5-8 membered heterocyclyl”). In some embodiments, a heterocyclyl group is a 5-6 membered non-aromatic ring system having ring carbon atoms and 1-4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5-6 membered heterocyclyl”). In some embodiments, the 5-6 membered heterocyclyl has 1-3 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5-6 membered heterocyclyl has 1-2

ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heterocyclyl has 1 ring heteroatom selected from nitrogen, oxygen, and sulfur.

[0044] Exemplary 3-membered heterocyclyl groups containing 1 heteroatom include, without limitation, azirdinyl, oxiranyl, and thiiranyl. Exemplary 4-membered heterocyclyl groups containing 1 heteroatom include, without limitation, azetidiny, oxetanyl, and thietanyl. Exemplary 5-membered heterocyclyl groups containing 1 heteroatom include, without limitation, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothiophenyl, dihydrothiophenyl, pyrrolidinyl, dihydropyrrolyl, and pyrrolyl-2,5-dione. Exemplary 5-membered heterocyclyl groups containing 2 heteroatoms include, without limitation, dioxolanyl, oxathiolanyl and dithiolanyl. Exemplary 5-membered heterocyclyl groups containing 3 heteroatoms include, without limitation, triazolinyl, oxadiazolinyl, and thiadiazolinyl. Exemplary 6-membered heterocyclyl groups containing 1 heteroatom include, without limitation, piperidinyl, tetrahydropyranyl, dihydropyridinyl, and thianyl. Exemplary 6-membered heterocyclyl groups containing 2 heteroatoms include, without limitation, piperazinyl, morpholinyl, dithianyl, and dioxanyl. Exemplary 6-membered heterocyclyl groups containing 3 heteroatoms include, without limitation, triazinyl. Exemplary 7-membered heterocyclyl groups containing 1 heteroatom include, without limitation, azepanyl, oxepanyl and thiepanyl. Exemplary 8-membered heterocyclyl groups containing 1 heteroatom include, without limitation, azocanyl, oxecanyl and thiocanyl. Exemplary bicyclic heterocyclyl groups include, without limitation, indolinyl, isoindolinyl, dihydrobenzofuranyl, dihydrobenzothienyl, tetrahydrobenzothienyl, tetrahydrobenzofuranyl, tetrahydroindolyl, tetrahydroquinoliny, tetrahydroisoquinoliny, decahydroquinoliny, decahydroisoquinoliny, octahydrochromenyl, octahydroisochromenyl, decahydronaphthyridinyl, decahydro-1,8-naphthyridinyl, octahydropyrrolo[3,2-b]pyrrole, phthalimidyl, naphthalimidyl, chromanyl, chromenyl, , and the like.

[0045] The term “aryl” refers to a radical of a monocyclic or polycyclic (*e.g.*, bicyclic or tricyclic) $4n+2$ aromatic ring system (*e.g.*, having 6, 10, or 14 π electrons shared in a cyclic array) having 6–14 ring carbon atoms and zero heteroatoms provided in the aromatic ring system (“C_{6–14} aryl”). In some embodiments, an aryl group has 6 ring carbon atoms (“C₆ aryl”; *e.g.*, phenyl). In some embodiments, an aryl group has 10 ring carbon atoms (“C₁₀ aryl”; *e.g.*, naphthyl such as 1–naphthyl and 2–naphthyl). In some embodiments, an aryl group has 14 ring carbon atoms (“C₁₄ aryl”; *e.g.*, anthracyl). “Aryl” also includes ring systems wherein the aryl ring, as defined above, is fused with one or more carbocyclyl or heterocyclyl groups wherein the radical or point of attachment is on the aryl ring, and in such

instances, the number of carbon atoms continue to designate the number of carbon atoms in the aryl ring system. Unless otherwise specified, each instance of an aryl group is independently unsubstituted (an “unsubstituted aryl”) or substituted (a “substituted aryl”) with one or more substituents. In certain embodiments, the aryl group is an unsubstituted C₆₋₁₄ aryl. In certain embodiments, the aryl group is a substituted C₆₋₁₄ aryl.

[0046] “Aralkyl” is a subset of “alkyl” and refers to an alkyl group substituted by an aryl group, wherein the point of attachment is on the alkyl moiety.

[0047] The term “heteroaryl” refers to a radical of a 5–14 membered monocyclic or polycyclic (*e.g.*, bicyclic, tricyclic) 4n+2 aromatic ring system (*e.g.*, having 6, 10, or 14 π electrons shared in a cyclic array) having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5–14 membered heteroaryl”). In heteroaryl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. Heteroaryl polycyclic ring systems can include one or more heteroatoms in one or both rings. “Heteroaryl” includes ring systems wherein the heteroaryl ring, as defined above, is fused with one or more carbocyclyl or heterocyclyl groups wherein the point of attachment is on the heteroaryl ring, and in such instances, the number of ring members continue to designate the number of ring members in the heteroaryl ring system. “Heteroaryl” also includes ring systems wherein the heteroaryl ring, as defined above, is fused with one or more aryl groups wherein the point of attachment is either on the aryl or heteroaryl ring, and in such instances, the number of ring members designates the number of ring members in the fused polycyclic (aryl/heteroaryl) ring system. Polycyclic heteroaryl groups wherein one ring does not contain a heteroatom (*e.g.*, indolyl, quinoliny, carbazolyl, and the like) the point of attachment can be on either ring, *i.e.*, either the ring bearing a heteroatom (*e.g.*, 2-indolyl) or the ring that does not contain a heteroatom (*e.g.*, 5-indolyl).

[0048] In some embodiments, a heteroaryl group is a 5–10 membered aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5–10 membered heteroaryl”). In some embodiments, a heteroaryl group is a 5–8 membered aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5–8 membered heteroaryl”). In some embodiments, a heteroaryl group is a 5–6 membered aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected

from nitrogen, oxygen, and sulfur (“5–6 membered heteroaryl”). In some embodiments, the 5–6 membered heteroaryl has 1–3 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heteroaryl has 1–2 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heteroaryl has 1 ring heteroatom selected from nitrogen, oxygen, and sulfur. Unless otherwise specified, each instance of a heteroaryl group is independently unsubstituted (an “unsubstituted heteroaryl”) or substituted (a “substituted heteroaryl”) with one or more substituents. In certain embodiments, the heteroaryl group is an unsubstituted 5–14 membered heteroaryl. In certain embodiments, the heteroaryl group is a substituted 5–14 membered heteroaryl.

[0049] Exemplary 5–membered heteroaryl groups containing 1 heteroatom include, without limitation, pyrrolyl, furanyl, and thiophenyl. Exemplary 5–membered heteroaryl groups containing 2 heteroatoms include, without limitation, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, and isothiazolyl. Exemplary 5–membered heteroaryl groups containing 3 heteroatoms include, without limitation, triazolyl, oxadiazolyl, and thiadiazolyl. Exemplary 5–membered heteroaryl groups containing 4 heteroatoms include, without limitation, tetrazolyl. Exemplary 6–membered heteroaryl groups containing 1 heteroatom include, without limitation, pyridinyl. Exemplary 6–membered heteroaryl groups containing 2 heteroatoms include, without limitation, pyridazinyl, pyrimidinyl, and pyrazinyl. Exemplary 6–membered heteroaryl groups containing 3 or 4 heteroatoms include, without limitation, triazinyl and tetrazinyl, respectively. Exemplary 7–membered heteroaryl groups containing 1 heteroatom include, without limitation, azepinyl, oxepinyl, and thiepinyl. Exemplary 5,6–bicyclic heteroaryl groups include, without limitation, indolyl, isoindolyl, indazolyl, benzotriazolyl, benzothiophenyl, isobenzothiophenyl, benzofuranyl, benzoisofuranyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzoxadiazolyl, benzthiazolyl, benzisothiazolyl, benzthiadiazolyl, indolizinyl, and purinyl. Exemplary 6,6–bicyclic heteroaryl groups include, without limitation, naphthyridinyl, pteridinyl, quinolinyl, isoquinolinyl, cinnolinyl, quinoxalinyl, phthalazinyl, and quinazolinyl. Exemplary tricyclic heteroaryl groups include, without limitation, phenanthridinyl, dibenzofuranyl, carbazolyl, acridinyl, phenothiazinyl, phenoxazinyl, and phenazinyl.

[0050] “Heteroaralkyl” is a subset of “alkyl” and refers to an alkyl group substituted by a heteroaryl group, wherein the point of attachment is on the alkyl moiety.

[0051] The term “unsaturated bond” refers to a double or triple bond.

[0052] The term “unsaturated” or “partially unsaturated” refers to a moiety that includes at least one double or triple bond.

[0053] The term “saturated” refers to a moiety that does not contain a double or triple bond, *i.e.*, the moiety only contains single bonds.

[0054] Affixing the suffix “-ene” to a group indicates the group is a divalent moiety, *e.g.*, alkylene is the divalent moiety of alkyl, alkenylene is the divalent moiety of alkenyl, alkynylene is the divalent moiety of alkynyl, heteroalkylene is the divalent moiety of heteroalkyl, heteroalkenylene is the divalent moiety of heteroalkenyl, heteroalkynylene is the divalent moiety of heteroalkynyl, carbocyclylene is the divalent moiety of carbocyclyl, heterocyclylene is the divalent moiety of heterocyclyl, arylene is the divalent moiety of aryl, and heteroarylene is the divalent moiety of heteroaryl.

[0055] A group is optionally substituted unless expressly provided otherwise. In certain embodiments, alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl groups are optionally substituted. “Optionally substituted” refers to a group which may be substituted or unsubstituted (*e.g.*, “substituted” or “unsubstituted” alkyl, “substituted” or “unsubstituted” alkenyl, “substituted” or “unsubstituted” alkynyl, “substituted” or “unsubstituted” heteroalkyl, “substituted” or “unsubstituted” heteroalkenyl, “substituted” or “unsubstituted” heteroalkynyl, “substituted” or “unsubstituted” carbocyclyl, “substituted” or “unsubstituted” heterocyclyl, “substituted” or “unsubstituted” aryl or “substituted” or “unsubstituted” heteroaryl group). In general, the term “substituted” means that at least one hydrogen present on a group is replaced with a permissible substituent, *e.g.*, a substituent which upon substitution results in a stable compound, *e.g.*, a compound which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, or other reaction. Unless otherwise indicated, a “substituted” group has a substituent at one or more substitutable positions of the group, and when more than one position in any given structure is substituted, the substituent is either the same or different at each position. The term “substituted” is contemplated to include substitution with all permissible substituents of organic compounds, and includes any of the substituents described herein that results in the formation of a stable compound. The present disclosure contemplates any and all such combinations in order to arrive at a stable compound. For purposes of this disclosure, heteroatoms such as nitrogen may have hydrogen substituents and/or any suitable substituent as described herein which satisfy the valencies of the heteroatoms and results in the formation of a stable moiety.

[0056] Exemplary carbon atom substituents include, but are not limited to, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{ON}(\text{R}^{\text{bb}})_2$, $-\text{N}(\text{R}^{\text{bb}})_2$, $-\text{N}(\text{R}^{\text{bb}})_3^+\text{X}^-$, $-\text{N}(\text{OR}^{\text{cc}})\text{R}^{\text{bb}}$, $-\text{SH}$, $-\text{SR}^{\text{aa}}$, $-\text{SSR}^{\text{cc}}$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{CO}_2\text{H}$, $-\text{CHO}$, $-\text{C}(\text{OR}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{OC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OCO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{CO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{OR}^{\text{aa}}$, $-\text{OSO}_2\text{R}^{\text{aa}}$, $-\text{S}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OS}(=\text{O})\text{R}^{\text{aa}}$, $-\text{Si}(\text{R}^{\text{aa}})_3$, $-\text{OSi}(\text{R}^{\text{aa}})_3$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{OC}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{OR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{OP}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{OP}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{P}(=\text{O})(\text{N}(\text{R}^{\text{bb}})_2)_2$, $-\text{OP}(=\text{O})(\text{N}(\text{R}^{\text{bb}})_2)_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{N}(\text{R}^{\text{bb}})_2)_2$, $-\text{P}(\text{R}^{\text{cc}})_2$, $-\text{P}(\text{OR}^{\text{cc}})_2$, $-\text{P}(\text{R}^{\text{cc}})_3^+\text{X}^-$, $-\text{P}(\text{OR}^{\text{cc}})_3^+\text{X}^-$, $-\text{P}(\text{R}^{\text{cc}})_4$, $-\text{P}(\text{OR}^{\text{cc}})_4$, $-\text{OP}(\text{R}^{\text{cc}})_2$, $-\text{OP}(\text{R}^{\text{cc}})_3^+\text{X}^-$, $-\text{OP}(\text{OR}^{\text{cc}})_2$, $-\text{OP}(\text{OR}^{\text{cc}})_3^+\text{X}^-$, $-\text{OP}(\text{R}^{\text{cc}})_4$, $-\text{OP}(\text{OR}^{\text{cc}})_4$, $-\text{B}(\text{R}^{\text{aa}})_2$, $-\text{B}(\text{OR}^{\text{cc}})_2$, $-\text{BR}^{\text{aa}}(\text{OR}^{\text{cc}})$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, hetero C_{1-10} alkyl, hetero C_{2-10} alkenyl, hetero C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3-14 membered heterocyclyl, C_{6-14} aryl, and 5-14 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups; wherein X^- is a counterion;

or two geminal hydrogens on a carbon atom are replaced with the group $=\text{O}$, $=\text{S}$, $=\text{NN}(\text{R}^{\text{bb}})_2$, $=\text{NNR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $=\text{NNR}^{\text{bb}}\text{C}(=\text{O})\text{OR}^{\text{aa}}$, $=\text{NNR}^{\text{bb}}\text{S}(=\text{O})_2\text{R}^{\text{aa}}$, $=\text{NR}^{\text{bb}}$, or $=\text{NOR}^{\text{cc}}$;

each instance of R^{aa} is, independently, selected from C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, hetero C_{1-10} alkyl, hetero C_{2-10} alkenyl, hetero C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3-14 membered heterocyclyl, C_{6-14} aryl, and 5-14 membered heteroaryl, or two R^{aa} groups are joined to form a 3-14 membered heterocyclyl or 5-14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{bb} is, independently, selected from hydrogen, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CN}$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{R}^{\text{cc}}$, $-\text{SO}_2\text{OR}^{\text{cc}}$, $-\text{SOR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{cc}}$, $-\text{C}(=\text{S})\text{SR}^{\text{cc}}$, $-\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{P}(=\text{O})(\text{N}(\text{R}^{\text{cc}})_2)_2$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, hetero C_{1-10} alkyl, hetero C_{2-10} alkenyl, hetero C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3-14 membered heterocyclyl, C_{6-14} aryl, and 5-14 membered heteroaryl, or two R^{bb} groups are joined to form a 3-14 membered heterocyclyl or 5-14 membered heteroaryl

ring, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups; wherein X⁻ is a counterion;

each instance of R^{cc} is, independently, selected from hydrogen, C₁₋₁₀ alkyl, C₁₋₁₀ perhaloalkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, heteroC₁₋₁₀ alkyl, heteroC₂₋₁₀ alkenyl, heteroC₂₋₁₀ alkynyl, C₃₋₁₀ carbocyclyl, 3-14 membered heterocyclyl, C₆₋₁₄ aryl, and 5-14 membered heteroaryl, or two R^{cc} groups are joined to form a 3-14 membered heterocyclyl or 5-14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

each instance of R^{dd} is, independently, selected from halogen, -CN, -NO₂, -N₃, -SO₂H, -SO₃H, -OH, -OR^{ee}, -ON(R^{ff})₂, -N(R^{ff})₂, -N(R^{ff})₃⁺X⁻, -N(OR^{ee})R^{ff}, -SH, -SR^{ee}, -SSR^{ee}, -C(=O)R^{ee}, -CO₂H, -CO₂R^{ee}, -OC(=O)R^{ee}, -OCO₂R^{ee}, -C(=O)N(R^{ff})₂, -OC(=O)N(R^{ff})₂, -NR^{ff}C(=O)R^{ee}, -NR^{ff}CO₂R^{ee}, -NR^{ff}C(=O)N(R^{ff})₂, -C(=NR^{ff})OR^{ee}, -OC(=NR^{ff})R^{ee}, -OC(=NR^{ff})OR^{ee}, -C(=NR^{ff})N(R^{ff})₂, -OC(=NR^{ff})N(R^{ff})₂, -NR^{ff}C(=NR^{ff})N(R^{ff})₂, -NR^{ff}SO₂R^{ee}, -SO₂N(R^{ff})₂, -SO₂R^{ee}, -SO₂OR^{ee}, -OSO₂R^{ee}, -S(=O)R^{ee}, -Si(R^{ee})₃, -OSi(R^{ee})₃, -C(=S)N(R^{ff})₂, -C(=O)SR^{ee}, -C(=S)SR^{ee}, -SC(=S)SR^{ee}, -P(=O)(OR^{ee})₂, -P(=O)(R^{ee})₂, -OP(=O)(R^{ee})₂, -OP(=O)(OR^{ee})₂, C₁₋₆ alkyl, C₁₋₆ perhaloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, heteroC₁₋₆alkyl, heteroC₂₋₆alkenyl, heteroC₂₋₆alkynyl, C₃₋₁₀ carbocyclyl, 3-10 membered heterocyclyl, C₆₋₁₀ aryl, 5-10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups, or two geminal R^{dd} substituents can be joined to form =O or =S; wherein X⁻ is a counterion;

each instance of R^{ee} is, independently, selected from C₁₋₆ alkyl, C₁₋₆ perhaloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, heteroC₁₋₆ alkyl, heteroC₂₋₆alkenyl, heteroC₂₋₆ alkynyl, C₃₋₁₀ carbocyclyl, C₆₋₁₀ aryl, 3-10 membered heterocyclyl, and 5-10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups;

each instance of R^{ff} is, independently, selected from hydrogen, C₁₋₆ alkyl, C₁₋₆ perhaloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, heteroC₁₋₆alkyl, heteroC₂₋₆alkenyl, heteroC₂₋₆alkynyl, C₃₋₁₀ carbocyclyl, 3-10 membered heterocyclyl, C₆₋₁₀ aryl and 5-10 membered heteroaryl, or two R^{ff} groups are joined to form a 3-10 membered heterocyclyl or 5-10 membered

heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups; and

each instance of R^{gg} is, independently, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OC}_{1-6}\text{ alkyl}$, $-\text{ON}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{N}(\text{C}_{1-6}\text{ alkyl})_3^+\text{X}^-$, $-\text{NH}(\text{C}_{1-6}\text{ alkyl})_2^+\text{X}^-$, $-\text{NH}_2(\text{C}_{1-6}\text{ alkyl})^+\text{X}^-$, $-\text{NH}_3^+\text{X}^-$, $-\text{N}(\text{OC}_{1-6}\text{ alkyl})(\text{C}_{1-6}\text{ alkyl})$, $-\text{N}(\text{OH})(\text{C}_{1-6}\text{ alkyl})$, $-\text{NH}(\text{OH})$, $-\text{SH}$, $-\text{SC}_{1-6}\text{ alkyl}$, $-\text{SS}(\text{C}_{1-6}\text{ alkyl})$, $-\text{C}(=\text{O})(\text{C}_{1-6}\text{ alkyl})$, $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_{1-6}\text{ alkyl})$, $-\text{OC}(=\text{O})(\text{C}_{1-6}\text{ alkyl})$, $-\text{OCO}_2(\text{C}_{1-6}\text{ alkyl})$, $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{OC}(=\text{O})\text{NH}(\text{C}_{1-6}\text{ alkyl})$, $-\text{NHC}(=\text{O})(\text{C}_{1-6}\text{ alkyl})$, $-\text{N}(\text{C}_{1-6}\text{ alkyl})\text{C}(=\text{O})(\text{C}_{1-6}\text{ alkyl})$, $-\text{NHCO}_2(\text{C}_{1-6}\text{ alkyl})$, $-\text{NHC}(=\text{O})\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{NHC}(=\text{O})\text{NH}(\text{C}_{1-6}\text{ alkyl})$, $-\text{NHC}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{NH})\text{O}(\text{C}_{1-6}\text{ alkyl})$, $-\text{OC}(=\text{NH})(\text{C}_{1-6}\text{ alkyl})$, $-\text{OC}(=\text{NH})\text{OC}_{1-6}\text{ alkyl}$, $-\text{C}(=\text{NH})\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{C}(=\text{NH})\text{NH}(\text{C}_{1-6}\text{ alkyl})$, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{OC}(=\text{NH})\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{OC}(\text{NH})\text{NH}(\text{C}_{1-6}\text{ alkyl})$, $-\text{OC}(\text{NH})\text{NH}_2$, $-\text{NHC}(\text{NH})\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{NHC}(=\text{NH})\text{NH}_2$, $-\text{NHSO}_2(\text{C}_{1-6}\text{ alkyl})$, $-\text{SO}_2\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{SO}_2\text{NH}(\text{C}_{1-6}\text{ alkyl})$, $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{C}_{1-6}\text{ alkyl}$, $-\text{SO}_2\text{OC}_{1-6}\text{ alkyl}$, $-\text{OSO}_2\text{C}_{1-6}\text{ alkyl}$, $-\text{SOC}_{1-6}\text{ alkyl}$, $-\text{Si}(\text{C}_{1-6}\text{ alkyl})_3$, $-\text{OSi}(\text{C}_{1-6}\text{ alkyl})_3$, $-\text{C}(=\text{S})\text{N}(\text{C}_{1-6}\text{ alkyl})_2$, $\text{C}(=\text{S})\text{NH}(\text{C}_{1-6}\text{ alkyl})$, $\text{C}(=\text{S})\text{NH}_2$, $-\text{C}(=\text{O})\text{S}(\text{C}_{1-6}\text{ alkyl})$, $-\text{C}(=\text{S})\text{SC}_{1-6}\text{ alkyl}$, $-\text{SC}(=\text{S})\text{SC}_{1-6}\text{ alkyl}$, $-\text{P}(=\text{O})(\text{OC}_{1-6}\text{ alkyl})_2$, $-\text{P}(=\text{O})(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{OP}(=\text{O})(\text{C}_{1-6}\text{ alkyl})_2$, $-\text{OP}(=\text{O})(\text{OC}_{1-6}\text{ alkyl})_2$, $\text{C}_{1-6}\text{ alkyl}$, $\text{C}_{1-6}\text{ perhaloalkyl}$, $\text{C}_{2-6}\text{ alkenyl}$, $\text{C}_{2-6}\text{ alkynyl}$, $\text{heteroC}_{1-6}\text{ alkyl}$, $\text{heteroC}_{2-6}\text{ alkenyl}$, $\text{heteroC}_{2-6}\text{ alkynyl}$, $\text{C}_{3-10}\text{ carbocyclyl}$, $\text{C}_{6-10}\text{ aryl}$, 3-10 membered heterocyclyl, 5-10 membered heteroaryl; or two geminal R^{gg} substituents can be joined to form $=\text{O}$ or $=\text{S}$; wherein X^- is a counterion.

[0057] Exemplary carbon atom substituents include, but are not limited to, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OR}^{aa}$, $-\text{ON}(\text{R}^{bb})_2$, $-\text{N}(\text{R}^{bb})_2$, $-\text{N}(\text{OR}^{cc})\text{R}^{bb}$, $-\text{SH}$, $-\text{SR}^{aa}$, $-\text{C}(=\text{O})\text{R}^{aa}$, $-\text{CO}_2\text{H}$, $-\text{CHO}$, $-\text{C}(\text{OR}^{cc})_2$, $-\text{CO}_2\text{R}^{aa}$, $-\text{OC}(=\text{O})\text{R}^{aa}$, $-\text{OCO}_2\text{R}^{aa}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{bb})_2$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{bb})_2$, $-\text{NR}^{bb}\text{C}(=\text{O})\text{R}^{aa}$, $-\text{NR}^{bb}\text{CO}_2\text{R}^{aa}$, $-\text{NR}^{bb}\text{C}(=\text{O})\text{N}(\text{R}^{bb})_2$, $-\text{C}(=\text{NR}^{bb})\text{R}^{aa}$, $-\text{C}(=\text{NR}^{bb})\text{OR}^{aa}$, $-\text{OC}(=\text{NR}^{bb})\text{R}^{aa}$, $-\text{OC}(=\text{NR}^{bb})\text{OR}^{aa}$, $-\text{C}(=\text{NR}^{bb})\text{N}(\text{R}^{bb})_2$, $-\text{OC}(=\text{NR}^{bb})\text{N}(\text{R}^{bb})_2$, $-\text{NR}^{bb}\text{C}(=\text{NR}^{bb})\text{N}(\text{R}^{bb})_2$, $-\text{C}(=\text{O})\text{NR}^{bb}\text{SO}_2\text{R}^{aa}$, $-\text{SO}_2\text{R}^{aa}$, $-\text{SO}_2\text{OR}^{aa}$, $-\text{OSO}_2\text{R}^{aa}$, $-\text{S}(=\text{O})\text{R}^{aa}$, or $-\text{OS}(=\text{O})\text{R}^{aa}$.

[0058] The term “halo” or “halogen” refers to fluorine (fluoro, $-\text{F}$), chlorine (chloro, $-\text{Cl}$), bromine (bromo, $-\text{Br}$), or iodine (iodo, $-\text{I}$).

[0059] The term “hydroxyl” or “hydroxy” refers to the group $-\text{OH}$. The term “substituted hydroxyl” or “substituted hydroxyl,” by extension, refers to a hydroxyl group wherein the oxygen atom directly attached to the parent molecule is substituted with a group other than hydrogen, and includes groups selected from $-\text{OR}^{aa}$, $-\text{ON}(\text{R}^{bb})_2$, $-\text{OC}(=\text{O})\text{SR}^{aa}$,

$-\text{OC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OCO}_2\text{R}^{\text{aa}}$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$,
 $-\text{OC}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OS}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OSO}_2\text{R}^{\text{aa}}$, $-\text{OSi}(\text{R}^{\text{aa}})_3$, $-\text{OP}(\text{R}^{\text{cc}})_2$, $-\text{OP}(\text{R}^{\text{cc}})_3^+\text{X}^-$,
 $-\text{OP}(\text{OR}^{\text{cc}})_2$, $-\text{OP}(\text{OR}^{\text{cc}})_3^+\text{X}^-$, $-\text{OP}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{OP}(=\text{O})(\text{OR}^{\text{cc}})_2$, and $-\text{OP}(=\text{O})(\text{N}(\text{R}^{\text{bb}}))_2$,
 wherein X^- , R^{aa} , R^{bb} , and R^{cc} are as defined herein.

[0060] The term “thiol” or “thio” refers to the group $-\text{SH}$. The term “substituted thiol” or “substituted thio,” by extension, refers to a thiol group wherein the sulfur atom directly attached to the parent molecule is substituted with a group other than hydrogen, and includes groups selected from $-\text{SR}^{\text{aa}}$, $-\text{S}=\text{SR}^{\text{cc}}$, $-\text{SC}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{OR}^{\text{aa}}$, and $-\text{SC}(=\text{O})\text{R}^{\text{aa}}$, wherein R^{aa} and R^{cc} are as defined herein.

[0061] The term “amino” refers to the group $-\text{NH}_2$. The term “substituted amino,” by extension, refers to a monosubstituted amino, a disubstituted amino, or a trisubstituted amino. In certain embodiments, the “substituted amino” is a monosubstituted amino or a disubstituted amino group.

[0062] The term “monosubstituted amino” refers to an amino group wherein the nitrogen atom directly attached to the parent molecule is substituted with one hydrogen and one group other than hydrogen, and includes groups selected from $-\text{NH}(\text{R}^{\text{bb}})$, $-\text{NHC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{NHCO}_2\text{R}^{\text{aa}}$, $-\text{NHC}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NHC}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NHSO}_2\text{R}^{\text{aa}}$, $-\text{NHP}(=\text{O})(\text{OR}^{\text{cc}})_2$, and $-\text{NHP}(=\text{O})(\text{N}(\text{R}^{\text{bb}}))_2$, wherein R^{aa} , R^{bb} and R^{cc} are as defined herein, and wherein R^{bb} of the group $-\text{NH}(\text{R}^{\text{bb}})$ is not hydrogen.

[0063] The term “disubstituted amino” refers to an amino group wherein the nitrogen atom directly attached to the parent molecule is substituted with two groups other than hydrogen, and includes groups selected from $-\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{CO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, and $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{N}(\text{R}^{\text{bb}}))_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein, with the proviso that the nitrogen atom directly attached to the parent molecule is not substituted with hydrogen.

[0064] The term “trisubstituted amino” refers to an amino group wherein the nitrogen atom directly attached to the parent molecule is substituted with three groups, and includes groups selected from $-\text{N}(\text{R}^{\text{bb}})_3$ and $-\text{N}(\text{R}^{\text{bb}})_3^+\text{X}^-$, wherein R^{bb} and X^- are as defined herein.

[0065] The term “sulfonyl” refers to a group selected from $-\text{SO}_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{SO}_2\text{R}^{\text{aa}}$, and $-\text{SO}_2\text{OR}^{\text{aa}}$, wherein R^{aa} and R^{bb} are as defined herein.

[0066] The term “sulfinyl” refers to the group $-\text{S}(=\text{O})\text{R}^{\text{aa}}$, wherein R^{aa} is as defined herein.

[0067] The term “carbonyl” refers a group wherein the carbon directly attached to the parent molecule is sp^2 hybridized, and is substituted with an oxygen, nitrogen or sulfur atom,

e.g., a group selected from ketones ($-\text{C}(=\text{O})\text{R}^{\text{aa}}$), carboxylic acids ($-\text{CO}_2\text{H}$), aldehydes ($-\text{CHO}$), esters ($-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{SR}^{\text{aa}}$), amides ($-\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{bb}})_2$), and imines ($-\text{C}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$), wherein R^{aa} and R^{bb} are as defined herein.

[0068] The term “silyl” refers to the group $-\text{Si}(\text{R}^{\text{aa}})_3$, wherein R^{aa} is as defined herein.

[0069] The term “oxo” refers to the group $=\text{O}$, and the term “thiooxo” refers to the group $=\text{S}$.

[0070] Nitrogen atoms can be substituted or unsubstituted as valency permits, and include primary, secondary, tertiary, and quaternary nitrogen atoms. Exemplary nitrogen atom substituents include, but are not limited to, hydrogen, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CN}$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{R}^{\text{cc}}$, $-\text{SO}_2\text{OR}^{\text{cc}}$, $-\text{SOR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{cc}}$, $-\text{C}(=\text{S})\text{SR}^{\text{cc}}$, $-\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{P}(=\text{O})(\text{N}(\text{R}^{\text{cc}})_2)_2$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, hetero C_{1-10} alkyl, hetero C_{2-10} alkenyl, hetero C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, or two R^{cc} groups attached to an N atom are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa} , R^{bb} , R^{cc} and R^{dd} are as defined above.

[0071] In certain embodiments, the substituent present on the nitrogen atom is a nitrogen protecting group (also referred to herein as an “amino protecting group”). Nitrogen protecting groups include, but are not limited to, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{cc}})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{N}(\text{R}^{\text{cc}})_2$, $-\text{SO}_2\text{R}^{\text{cc}}$, $-\text{SO}_2\text{OR}^{\text{cc}}$, $-\text{SOR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{cc}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{cc}}$, $-\text{C}(=\text{S})\text{SR}^{\text{cc}}$, C_{1-10} alkyl (*e.g.*, aralkyl, heteroaralkyl), C_{2-10} alkenyl, C_{2-10} alkynyl, hetero C_{1-10} alkyl, hetero C_{2-10} alkenyl, hetero C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl groups, wherein each alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl, heterocyclyl, aralkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa} , R^{bb} , R^{cc} and R^{dd} are as defined herein. Nitrogen protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[0072] For example, nitrogen protecting groups such as amide groups (*e.g.*, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$) include, but are not limited to, formamide, acetamide, chloroacetamide, trichloroacetamide,

trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, picolinamide, 3-pyridylcarboxamide, *N*-benzoylphenylalanyl derivative, benzamide, *p*-phenylbenzamide, *o*-nitrophenylacetamide, *o*-nitrophenoxyacetamide, acetoacetamide, (*N*'-dithiobenzyloxyacylamino)acetamide, 3-(*p*-hydroxyphenyl)propanamide, 3-(*o*-nitrophenyl)propanamide, 2-methyl-2-(*o*-nitrophenoxy)propanamide, 2-methyl-2-(*o*-phenylazophenoxy)propanamide, 4-chlorobutanamide, 3-methyl-3-nitrobutanamide, *o*-nitrocinnamide, *N*-acetylmethionine derivative, *o*-nitrobenzamide, and *o*-(benzyloxymethyl)benzamide.

[0073] Nitrogen protecting groups such as carbamate groups (*e.g.*, $-\text{C}(=\text{O})\text{OR}^{\text{aa}}$) include, but are not limited to, methyl carbamate, ethyl carbamate, 9-fluorenylmethyl carbamate (Fmoc), 9-(2-sulfo)fluorenylmethyl carbamate, 9-(2,7-dibromo)fluorenylmethyl carbamate, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl carbamate (DBD-Tmoc), 4-methoxyphenacyl carbamate (Phenoc), 2,2,2-trichloroethyl carbamate (Troc), 2-trimethylsilylethyl carbamate (Teoc), 2-phenylethyl carbamate (hZ), 1-(1-adamantyl)-1-methylethyl carbamate (Adpoc), 1,1-dimethyl-2-haloethyl carbamate, 1,1-dimethyl-2,2-dibromoethyl carbamate (DB-*t*-BOC), 1,1-dimethyl-2,2,2-trichloroethyl carbamate (TCBOC), 1-methyl-1-(4-biphenyl)ethyl carbamate (Bpoc), 1-(3,5-di-*t*-butylphenyl)-1-methylethyl carbamate (*t*-Bumeoc), 2-(2'- and 4'-pyridyl)ethyl carbamate (Pyoc), 2-(*N,N*-dicyclohexylcarboxamido)ethyl carbamate, *t*-butyl carbamate (BOC or Boc), 1-adamantyl carbamate (Adoc), vinyl carbamate (Voc), allyl carbamate (Alloc), 1-isopropylallyl carbamate (Ipaoc), cinnamyl carbamate (Coc), 4-nitrocinnamyl carbamate (Noc), 8-quinolyl carbamate, *N*-hydroxypiperidinyl carbamate, alkylthio carbamate, benzyl carbamate (Cbz), *p*-methoxybenzyl carbamate (Moz), *p*-nitrobenzyl carbamate, *p*-bromobenzyl carbamate, *p*-chlorobenzyl carbamate, 2,4-dichlorobenzyl carbamate, 4-methylsulfinylbenzyl carbamate (MsZ), 9-anthrylmethyl carbamate, diphenylmethyl carbamate, 2-methylthioethyl carbamate, 2-methylsulfonylethyl carbamate, 2-(*p*-toluenesulfonyl)ethyl carbamate, [2-(1,3-dithianyl)]methyl carbamate (Dmoc), 4-methylthiophenyl carbamate (Mtpc), 2,4-dimethylthiophenyl carbamate (Bmpc), 2-phosphonioethyl carbamate (Peoc), 2-triphenylphosphonioisopropyl carbamate (Ppoc), 1,1-dimethyl-2-cyanoethyl carbamate, *m*-chloro-*p*-acyloxybenzyl carbamate, *p*-(dihydroxyboryl)benzyl carbamate, 5-benzisoxazolylmethyl carbamate, 2-(trifluoromethyl)-6-chromonylmethyl carbamate (Tcroc), *m*-nitrophenyl carbamate, 3,5-dimethoxybenzyl carbamate, *o*-nitrobenzyl carbamate, 3,4-dimethoxy-6-nitrobenzyl carbamate, phenyl(*o*-nitrophenyl)methyl carbamate, *t*-amyl carbamate, *S*-benzyl thiocarbamate, *p*-cyanobenzyl

carbamate, cyclobutyl carbamate, cyclohexyl carbamate, cyclopentyl carbamate, cyclopropylmethyl carbamate, *p*-decyloxybenzyl carbamate, 2,2-dimethoxyacetylvinyl carbamate, *o*-(*N,N*-dimethylcarboxamido)benzyl carbamate, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl carbamate, 1,1-dimethylpropynyl carbamate, di(2-pyridyl)methyl carbamate, 2-furanylmethyl carbamate, 2-iodoethyl carbamate, isoborynl carbamate, isobutyl carbamate, isonicotinyl carbamate, *p*-(*p'*-methoxyphenylazo)benzyl carbamate, 1-methylcyclobutyl carbamate, 1-methylcyclohexyl carbamate, 1-methyl-1-cyclopropylmethyl carbamate, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl carbamate, 1-methyl-1-(*p*-phenylazophenyl)ethyl carbamate, 1-methyl-1-phenylethyl carbamate, 1-methyl-1-(4-pyridyl)ethyl carbamate, phenyl carbamate, *p*-(phenylazo)benzyl carbamate, 2,4,6-tri-*t*-butylphenyl carbamate, 4-(trimethylammonium)benzyl carbamate, and 2,4,6-trimethylbenzyl carbamate.

[0074] Nitrogen protecting groups such as sulfonamide groups (*e.g.*, $-S(=O)_2R^{aa}$) include, but are not limited to, *p*-toluenesulfonamide (Ts), benzenesulfonamide, 2,3,6-trimethyl-4-methoxybenzenesulfonamide (Mtr), 2,4,6-trimethoxybenzenesulfonamide (Mtb), 2,6-dimethyl-4-methoxybenzenesulfonamide (Pme), 2,3,5,6-tetramethyl-4-methoxybenzenesulfonamide (Mte), 4-methoxybenzenesulfonamide (Mbs), 2,4,6-trimethylbenzenesulfonamide (Mts), 2,6-dimethoxy-4-methylbenzenesulfonamide (iMds), 2,2,5,7,8-pentamethylchroman-6-sulfonamide (Pmc), methanesulfonamide (Ms), β -trimethylsilyl ethanesulfonamide (SES), 9-anthracenesulfonamide, 4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonamide (DNMBS), benzylsulfonamide, trifluoromethylsulfonamide, and phenacysulfonamide.

[0075] Other nitrogen protecting groups include, but are not limited to, phenothiazinyl-(10)-acyl derivative, *N'*-*p*-toluenesulfonylaminoacyl derivative, *N'*-phenylaminothioacyl derivative, *N*-benzoylphenylalanyl derivative, *N*-acetylmethionine derivative, 4,5-diphenyl-3-oxazolin-2-one, *N*-phthalimide, *N*-dithiasuccinimide (Dts), *N*-2,3-diphenylmaleimide, *N*-2,5-dimethylpyrrole, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct (STABASE), 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridone, *N*-methylamine, *N*-allylamine, *N*-[2-(trimethylsilyl)ethoxy]methylamine (SEM), *N*-3-acetoxypyrrolamine, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrroline-3-yl)amine, quaternary ammonium salts, *N*-benzylamine, *N*-di(4-methoxyphenyl)methylamine, *N*-5-dibenzosuberylamine, *N*-triphenylmethylamine (Tr), *N*-[(4-methoxyphenyl)diphenylmethyl]amine (MMTr), *N*-9-phenylfluorenylamine (PhF), *N*-2,7-dichloro-9-fluorenylmethyleneamine, *N*-

ferrocenylmethylamino (Fcm), *N*-2-picolylamino *N'*-oxide, *N*-1,1-dimethylthiomethyleneamine, *N*-benzylideneamine, *N*-*p*-methoxybenzylideneamine, *N*-diphenylmethyleneamine, *N*-[(2-pyridyl)mesityl]methyleneamine, *N*-(*N'*,*N'*-dimethylaminomethylene)amine, *N,N'*-isopropylidenediamine, *N*-*p*-nitrobenzylideneamine, *N*-salicylideneamine, *N*-5-chlorosalicylideneamine, *N*-(5-chloro-2-hydroxyphenyl)phenylmethyleneamine, *N*-cyclohexylideneamine, *N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl)amine, *N*-borane derivative, *N*-diphenylborinic acid derivative, *N*-[phenyl(pentaacylchromium- or tungsten)acyl]amine, *N*-copper chelate, *N*-zinc chelate, *N*-nitroamine, *N*-nitrosoamine, amine *N*-oxide, diphenylphosphinamide (Dpp), dimethylthiophosphinamide (Mpt), diphenylthiophosphinamide (Ppt), dialkyl phosphoramidates, dibenzyl phosphoramidate, diphenyl phosphoramidate, benzenesulfenamide, *o*-nitrobenzenesulfenamide (Nps), 2,4-dinitrobenzenesulfenamide, pentachlorobenzenesulfenamide, 2-nitro-4-methoxybenzenesulfenamide, triphenylmethylsulfenamide, and 3-nitropyridinesulfenamide (Npys).

[0076] In certain embodiments, the substituent present on an oxygen atom is an oxygen protecting group (also referred to herein as an “hydroxyl protecting group”). Oxygen protecting groups include, but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3^+X^-$, $-P(OR^{cc})_2$, $-P(OR^{cc})_3^+X^-$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, and $-P(=O)(N(R^{bb})_2)_2$, wherein X^- , R^{aa} , R^{bb} , and R^{cc} are as defined herein. Oxygen protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[0077] Exemplary oxygen protecting groups include, but are not limited to, methyl, methoxymethyl (MOM), methylthiomethyl (MTM), *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl (SMOM), benzyloxymethyl (BOM), *p*-methoxybenzyloxymethyl (PMBM), (4-methoxyphenoxy)methyl (*p*-AOM), guaiacolmethyl (GUM), *t*-butoxymethyl, 4-pentenylloxymethyl (POM), siloxymethyl, 2-methoxyethoxymethyl (MEM), 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl (SEMOR), tetrahydropyranyl (THP), 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl (MTHP), 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxide, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl (CTMP), 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl,

2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, benzyl (Bn), *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p,p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl, α -naphthyldiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxyphenyl)diphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(imidazol-1-yl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-2-yl, benzisothiazolyl S,S-dioxido, trimethylsilyl (TMS), triethylsilyl (TES), triisopropylsilyl (TIPS), dimethylisopropylsilyl (IPDMS), diethylisopropylsilyl (DEIPS), dimethylhexylsilyl, *t*-butyldimethylsilyl (TBDMS), *t*-butyldiphenylsilyl (TBDPS), tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl (DPMS), *t*-butylmethoxyphenylsilyl (TBMPS), formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-(ethylenedithio)pentanoate (levulinoyldithioacetal), pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate (mesitoate), methyl carbonate, 9-fluorenylmethyl carbonate (Fmoc), ethyl carbonate, 2,2,2-trichloroethyl carbonate (Troc), 2-(trimethylsilyl)ethyl carbonate (TMSEC), 2-(phenylsulfonyl) ethyl carbonate (Psec), 2-(triphenylphosphonio) ethyl carbonate (Peoc), isobutyl carbonate, vinyl carbonate, allyl carbonate, *t*-butyl carbonate (BOC or Boc), *p*-nitrophenyl carbonate, benzyl carbonate, *p*-methoxybenzyl carbonate, 3,4-dimethoxybenzyl carbonate, *o*-nitrobenzyl carbonate, *p*-nitrobenzyl carbonate, *S*-benzyl thiocarbonate, 4-ethoxy-1-naphthyl carbonate, methyl dithiocarbonate, 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl, 4-(methylthiomethoxy)butyrate, 2-(methylthiomethoxymethyl)benzoate, 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate,


(*E*)-2-methyl-2-butenolate, *o*-(methoxyacyl)benzoate, α -naphthoate, nitrate, alkyl *N,N,N',N'*-tetramethylphosphorodiamidate, alkyl *N*-phenylcarbamate, borate, dimethylphosphinothioyl, alkyl 2,4-dinitrophenylsulfenate, sulfate, methanesulfonate (mesylate), benzyisulfonate, and tosylate (Ts).









[0078] In certain embodiments, the substituent present on a sulfur atom is a sulfur protecting group (also referred to as a “thiol protecting group”). Sulfur protecting groups include, but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3^+X^-$, $-P(OR^{cc})_2$, $-P(OR^{cc})_3^+X^-$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, and $-P(=O)(N(R^{bb})_2)_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein; wherein X^- is a counterion. Sulfur protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

[0079] A “counterion” or “anionic counterion” is a negatively charged group associated with a positively charged group in order to maintain electronic neutrality. An anionic counterion may be monovalent (*i.e.*, including one formal negative charge). An anionic counterion may also be multivalent (*i.e.*, including more than one formal negative charge), such as divalent or trivalent. Exemplary counterions include halide ions (*e.g.*, F^- , Cl^- , Br^- , I^-), NO_3^- , ClO_4^- , OH^- , $H_2PO_4^-$, HCO_3^- , HSO_4^- , sulfonate ions (*e.g.*, methanesulfonate, trifluoromethanesulfonate, *p*-toluenesulfonate, benzenesulfonate, 10-camphor sulfonate, naphthalene-2-sulfonate, naphthalene-1-sulfonic acid-5-sulfonate, ethan-1-sulfonic acid-2-sulfonate, and the like), carboxylate ions (*e.g.*, acetate, propanoate, benzoate, glycerate, lactate, tartrate, glycolate, gluconate, and the like), BF_4^- , PF_4^- , PF_6^- , AsF_6^- , SbF_6^- , $B[3,5-(CF_3)_2C_6H_3]_4^-$, $B(C_6F_5)_4^-$, BPh_4^- , $Al(OC(CF_3)_3)_4^-$, and carborane anions (*e.g.*, $CB_{11}H_{12}^-$ or $(HCB_{11}Me_5Br_6)^-$). Exemplary counterions which may be multivalent include CO_3^{2-} , HPO_4^{2-} , PO_4^{3-} , $B_4O_7^{2-}$, SO_4^{2-} , $S_2O_3^{2-}$, carboxylate anions (*e.g.*, tartrate, citrate, fumarate, maleate, malate, malonate, gluconate, succinate, glutarate, adipate, pimelate, suberate, azelate, sebacate, salicylate, phthalates, aspartate, glutamate, and the like), and carboranes.


[0080] A “hydrocarbon chain” refers to a substituted or unsubstituted divalent alkyl, alkenyl, or alkynyl group. A hydrocarbon chain includes (1) one or more chains of carbon atoms immediately between the two radicals of the hydrocarbon chain; (2) optionally one or more hydrogen atoms on the chain(s) of carbon atoms; and (3) optionally one or more substituents (“non-chain substituents,” which are not hydrogen) on the chain(s) of carbon atoms. A chain of carbon atoms consists of consecutively connected carbon atoms (“chain

atoms”) and does not include hydrogen atoms or heteroatoms. However, a non-chain substituent of a hydrocarbon chain may include any atoms, including hydrogen atoms, carbon atoms, and heteroatoms. For example, hydrocarbon chain $-C^A H(C^B H_2 C^C H_3)-$ includes one chain atom C^A , one hydrogen atom on C^A , and non-chain substituent $-(C^B H_2 C^C H_3)$. The term “ C_x hydrocarbon chain,” wherein x is a positive integer, refers to a hydrocarbon chain that includes x number of chain atom(s) between the two radicals of the hydrocarbon chain. If there is more than one possible value of x , the smallest possible value of x is used for the definition of the hydrocarbon chain. For example, $-CH(C_2H_5)-$ is a C_1 hydrocarbon chain,

and  is a C_3 hydrocarbon chain. When a range of values is used, the meaning of the range is as described herein. For example, a C_{3-10} hydrocarbon chain refers to a hydrocarbon chain where the number of chain atoms of the shortest chain of carbon atoms immediately between the two radicals of the hydrocarbon chain is 3, 4, 5, 6, 7, 8, 9, or 10. A hydrocarbon chain may be saturated (*e.g.*, $-(CH_2)_4-$). A hydrocarbon chain may also be unsaturated and include one or more $C=C$ and/or $C\equiv C$ bonds anywhere in the hydrocarbon chain. For instance, $-CH=CH-(CH_2)_2-$, $-CH_2-C\equiv C-CH_2-$, and $-C\equiv C-CH=CH-$ are all examples of a unsubstituted and unsaturated hydrocarbon chain. In certain embodiments, the hydrocarbon chain is unsubstituted (*e.g.*, $-C\equiv C-$ or $-(CH_2)_4-$). In certain embodiments, the hydrocarbon chain is substituted (*e.g.*, $-CH(C_2H_5)-$ and $-CF_2-$). Any two substituents on the hydrocarbon chain may be joined to form an optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl ring.

For instance, , , , , , and  are all examples of a hydrocarbon chain. In contrast, in certain embodiments,  and  are not within the scope of the hydrocarbon chains described herein.

When a chain atom of a C_x hydrocarbon chain is replaced with a heteroatom, the resulting group is referred to as a C_x hydrocarbon chain wherein a chain atom is replaced with

a heteroatom, as opposed to a C_{x-1} hydrocarbon chain. For example,  is a C_3 hydrocarbon chain wherein one chain atom is replaced with an oxygen atom.

[0081] As used herein, a “leaving group” (LG) is an art-understood term referring to a molecular fragment that departs with a pair of electrons in heterolytic bond cleavage, wherein the molecular fragment is an anion or neutral molecule. As used herein, a leaving group can be an atom or a group capable of being displaced by a nucleophile. See, for example, Smith, March Advanced Organic Chemistry 6th ed. (501-502). Exemplary leaving groups include, but are not limited to, halo (*e.g.*, chloro, bromo, iodo) and activated substituted hydroxyl groups (*e.g.*, $-OC(=O)SR^{aa}$, $-OC(=O)R^{aa}$, $-OCO_2R^{aa}$, $-OC(=O)N(R^{bb})_2$, $-OC(=NR^{bb})R^{aa}$, $-OC(=NR^{bb})OR^{aa}$, $-OC(=NR^{bb})N(R^{bb})_2$, $-OS(=O)R^{aa}$, $-OSO_2R^{aa}$, $-OP(R^{cc})_2$, $-OP(R^{cc})_3$, $-OP(=O)_2R^{aa}$, $-OP(=O)(R^{aa})_2$, $-OP(=O)(OR^{cc})_2$, $-OP(=O)_2N(R^{bb})_2$, and $-OP(=O)(NR^{bb})_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein).

[0082] A “counterion” or “anionic counterion” is a negatively charged group associated with a positively charged group in order to maintain electronic neutrality. An anionic counterion may be monovalent (*i.e.*, including one formal negative charge). An anionic counterion may also be multivalent (*i.e.*, including more than one formal negative charge), such as divalent or trivalent. Exemplary counterions include halide ions (*e.g.*, F^- , Cl^- , Br^- , I^-), NO_3^- , ClO_4^- , OH^- , $H_2PO_4^-$, HCO_3^- , HSO_4^- , sulfonate ions (*e.g.*, methanesulfonate, trifluoromethanesulfonate, *p*-toluenesulfonate, benzenesulfonate, 10-camphor sulfonate, naphthalene-2-sulfonate, naphthalene-1-sulfonic acid-5-sulfonate, ethan-1-sulfonic acid-2-sulfonate, and the like), carboxylate ions (*e.g.*, acetate, propanoate, benzoate, glycerate, lactate, tartrate, glycolate, gluconate, and the like), BF_4^- , PF_4^- , PF_6^- , AsF_6^- , SbF_6^- , $B[3,5-(CF_3)_2C_6H_3]_4^-$, $B(C_6F_5)_4^-$, BPh_4^- , $Al(OC(CF_3)_3)_4^-$, and carborane anions (*e.g.*, $CB_{11}H_{12}^-$ or $(HCB_{11}Me_5Br_6)^-$). Exemplary counterions which may be multivalent include CO_3^{2-} , HPO_4^{2-} , PO_4^{3-} , $B_4O_7^{2-}$, SO_4^{2-} , $S_2O_3^{2-}$, carboxylate anions (*e.g.*, tartrate, citrate, fumarate, maleate, malate, malonate, gluconate, succinate, glutarate, adipate, pimelate, suberate, azelate, sebacate, salicylate, phthalates, aspartate, glutamate, and the like), and carboranes.

[0083] The term “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, Berge *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1–19, incorporated herein by reference.

Pharmaceutically acceptable salts of the compounds of this disclosure include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid, or malonic acid or by using other methods known in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium, and $N^+(C_{1-4} \text{ alkyl})_4^-$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate, and aryl sulfonate. In certain embodiments, a “pharmaceutically acceptable salt” is a “cosmetically acceptable salt.”

[0084] The term “solvate” refers to forms of the compound, or a salt thereof, that are associated with a solvent, usually by a solvolysis reaction. This physical association may include hydrogen bonding. Conventional solvents include water, methanol, ethanol, acetic acid, DMSO, THF, diethyl ether, and the like. The compounds described herein may be prepared, *e.g.*, in crystalline form, and may be solvated. Suitable solvates include pharmaceutically acceptable solvates and further include both stoichiometric solvates and non-stoichiometric solvates. In certain instances, the solvate will be capable of isolation, for example, when one or more solvent molecules are incorporated in the crystal lattice of a crystalline solid. “Solvate” encompasses both solution-phase and isolatable solvates. Representative solvates include hydrates, ethanolates, and methanolates.

[0085] The term “hydrate” refers to a compound that is associated with water. Typically, the number of the water molecules contained in a hydrate of a compound is in a definite ratio

to the number of the compound molecules in the hydrate. Therefore, a hydrate of a compound may be represented, for example, by the general formula $R \cdot x H_2O$, wherein R is the compound, and x is a number greater than 0. A given compound may form more than one type of hydrate, including, *e.g.*, monohydrates (x is 1), lower hydrates (x is a number greater than 0 and smaller than 1, *e.g.*, hemihydrates ($R \cdot 0.5 H_2O$)), and polyhydrates (x is a number greater than 1, *e.g.*, dihydrates ($R \cdot 2 H_2O$) and hexahydrates ($R \cdot 6 H_2O$)).

[0086] The term “tautomers” or “tautomeric” refers to two or more interconvertible compounds resulting from at least one formal migration of a hydrogen atom and at least one change in valency (*e.g.*, a single bond to a double bond, a triple bond to a single bond, or *vice versa*). The exact ratio of the tautomers depends on several factors, including temperature, solvent, and pH. Tautomerizations (*i.e.*, the reaction providing a tautomeric pair) may catalyzed by acid or base. Exemplary tautomerizations include keto-to-enol, amide-to-imide, lactam-to-lactim, enamine-to-imine, and enamine-to-(a different enamine) tautomerizations.

[0087] It is also to be understood that compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed “isomers”. Isomers that differ in the arrangement of their atoms in space are termed “stereoisomers”.

[0088] Stereoisomers that are not mirror images of one another are termed “diastereomers” and those that are non-superimposable mirror images of each other are termed “enantiomers”. When a compound has an asymmetric center, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (*i.e.*, as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a “racemic mixture”.

[0089] The term “polymorphs” refers to a crystalline form of a compound (or a salt, hydrate, or solvate thereof). All polymorphs have the same elemental composition. Different crystalline forms usually have different X-ray diffraction patterns, infrared spectra, melting points, density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Recrystallization solvent, rate of crystallization, storage temperature, and other

factors may cause one crystal form to dominate. Various polymorphs of a compound can be prepared by crystallization under different conditions.

[0090] The term “prodrugs” refers to compounds that have cleavable groups and become by solvolysis or under physiological conditions the compounds described herein, which are pharmaceutically active *in vivo*. Such examples include, but are not limited to, choline ester derivatives and the like, N-alkylmorpholine esters and the like. Other derivatives of the compounds described herein have activity in both their acid and acid derivative forms, but in the acid sensitive form often offer advantages of solubility, tissue compatibility, or delayed release in the mammalian organism (see, Bundgard, H., *Design of Prodrugs*, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acid with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a substituted or unsubstituted amine, or acid anhydrides, or mixed anhydrides. Simple aliphatic or aromatic esters, amides, and anhydrides derived from acidic groups pendant on the compounds described herein are particular prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy)alkyl esters or ((alkoxycarbonyl)oxy)alkylesters. C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, aryl, C₇-C₁₂ substituted aryl, and C₇-C₁₂ arylalkyl esters of the compounds described herein may be preferred.

[0091] The “molecular weight” of a monovalent moiety –R is calculated by subtracting 1 from the molecular weight of the compound R–H. The “molecular weight” of a divalent moiety –L– is calculated by subtracting 2 from the molecular weight of the compound H–L–H.

[0092] The terms “composition” and “formulation” are used interchangeably.

[0093] A “subject” to which administration is contemplated includes, but is not limited to, humans (*i.e.*, a male or female of any age group, *e.g.*, a pediatric subject (*e.g.*, infant, child, adolescent) or adult subject (*e.g.*, young adult, middle-aged adult, or senior adult)) and/or other non-human animals, for example, mammals (*e.g.*, primates (*e.g.*, cynomolgus monkeys, rhesus monkeys); commercially relevant mammals such as cattle, pigs, horses, sheep, goats, cats, and/or dogs) and birds (*e.g.*, commercially relevant birds such as chickens, ducks, geese, and/or turkeys). In certain embodiments, the animal is a mammal. The animal may be a male or female at any stage of development. The animal may be a transgenic animal or genetically engineered animal. In certain embodiments, the subject is a non-human animal. A “patient” refers to a human subject in need of treatment of a disease.

[0094] The term “topical” refers to local application of the compounds described herein to a body surface of a human or non-human animal. In certain embodiments, the body surface is skin. In certain embodiments, the skin is on a body part. In certain embodiments, the skin is on the face. In certain embodiments, the skin is on the neck. In certain embodiments, the skin is on the torso. In certain embodiments, the skin is on the chest. In certain embodiments, the skin is on the back. In certain embodiments, the skin is on the arms. In certain embodiments, the skin is on the legs.

[0095] The term “skin condition” or “skin disease” refers to a condition related to the skin. Exemplary skin conditions or skin diseases include, but are not limited to, skin cancers (*e.g.*, non-melanoma and melanoma skin cancers, basal or squamous cell skin cancer, Merkel cell carcinoma, skin lymphoma, or Kaposi sarcoma), pruritus (itch), psoriasis, eczema, burns, or dermatitis.

[0096] The term “administer,” “administering,” or “administration” refers to implanting, absorbing, ingesting, injecting, inhaling, or otherwise introducing a compound described herein, or a composition thereof, in or on a subject.

[0097] The terms “treatment,” “treat,” and “treating” refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease described herein. In other embodiments, treatment may be administered in the absence of signs or symptoms of the disease. For example, treatment may be administered to a susceptible subject prior to the onset of symptoms (*e.g.*, in light of a history of symptoms and/or in light of exposure to a pathogen). Treatment may also be continued after symptoms have resolved, for example, to delay and/or prevent recurrence.

[0098] The term “prevent” refers to a prophylactic treatment of a subject who is not and did not have a disease but is at risk of developing the disease or who had a disease, does not have the disease, but is at risk of regression of the disease.

[0099] The terms “condition,” “disease,” and “disorder” are used interchangeably.

[00100] An “effective amount” of a compound described herein refers to an amount sufficient to elicit the desired biological response. An effective amount of a compound described herein may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the condition being treated, the mode of administration, and the age and health of the subject. In certain embodiments, an effective amount is a therapeutically effective amount. In certain embodiments, an effective amount is a prophylactic treatment. In certain embodiments, an effective amount is the amount of a

compound described herein in a single dose. In certain embodiments, an effective amount is the combined amounts of a compound described herein in multiple doses.

[00101] A “prophylactically effective amount” of a compound described herein is an amount sufficient to prevent a condition, or one or more symptoms associated with the condition or prevent its recurrence. A prophylactically effective amount of a compound means an amount of a therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of skin cancer. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

[00102] A “kinase” is a type of enzyme that transfers phosphate groups from high energy donor molecules, such as ATP, to specific substrates, referred to as phosphorylation. Kinases are part of the larger family of phosphotransferases. One of the largest groups of kinases are protein kinases, which act on and modify the activity of specific proteins. Kinases are used extensively to transmit signals and control complex processes in cells. Various other kinases act on small molecules such as lipids, carbohydrates, amino acids, and nucleotides, either for signaling or to prime them for metabolic pathways. Kinases are often named after their substrates. More than 500 different protein kinases have been identified in humans. Salt-inducible kinase (SIK) is one exemplary human protein kinase (*e.g.*, SIK1, SIK2, or SIK3).

[00103] The term “salt-inducible kinase” or “SIK” refers to a subfamily of serine/threonine protein kinases including SIK1, SIK2, and SIK3 that belong to an AMP-activated protein kinase family. In certain embodiments, the SIK is SIK1. In certain embodiments, the SIK is SIK2. In certain embodiments, the SIK is SIK3.

[00104] The term “inhibition,” “inhibiting,” “inhibit,” or “inhibitor” refer to the ability of a compound to reduce, slow, halt, and/or prevent activity of a particular biological process (*e.g.*, SIK kinase activity) in a cell relative to vehicle.

[00105] When a compound, pharmaceutical composition, method, use, or kit is referred to as “selectively” or “specifically” modulating (*e.g.*, increasing or inhibiting) the activity of a first protein kinase, the compound, pharmaceutical composition, method, use, or kit modulates the activity of the first protein kinase (*e.g.*, SIK) to a greater extent (*e.g.*, not less than about 2-fold, not less than about 5-fold, not less than about 10-fold, not less than about 30-fold, not less than about 100-fold, not less than about 1,000-fold, or not less than about 10,000-fold) than the activity of a second protein kinase that is different from the first protein kinase.

[00106] The term “cancer” refers to a class of diseases characterized by the development of abnormal cells that proliferate uncontrollably and have the ability to infiltrate and destroy normal body tissues. See, *e.g.*, *Stedman’s Medical Dictionary*, 25th ed.; Hensyl ed.; Williams & Wilkins: Philadelphia, 1990. Exemplary cancers include, but are not limited to, acoustic neuroma; adenocarcinoma; adrenal gland cancer; anal cancer; angiosarcoma (*e.g.*, lymphangiosarcoma, lymphangioendotheliosarcoma, hemangiosarcoma); appendix cancer; benign monoclonal gammopathy; biliary cancer (*e.g.*, cholangiocarcinoma); bladder cancer; breast cancer (*e.g.*, adenocarcinoma of the breast, papillary carcinoma of the breast, mammary cancer, medullary carcinoma of the breast); brain cancer (*e.g.*, meningioma, glioblastomas, glioma (*e.g.*, astrocytoma, oligodendroglioma), medulloblastoma); bronchus cancer; carcinoid tumor; cervical cancer (*e.g.*, cervical adenocarcinoma); choriocarcinoma; chordoma; craniopharyngioma; colorectal cancer (*e.g.*, colon cancer, rectal cancer, colorectal adenocarcinoma); connective tissue cancer; epithelial carcinoma; ependymoma; endotheliosarcoma (*e.g.*, Kaposi’s sarcoma, multiple idiopathic hemorrhagic sarcoma); endometrial cancer (*e.g.*, uterine cancer, uterine sarcoma); esophageal cancer (*e.g.*, adenocarcinoma of the esophagus, Barrett’s adenocarcinoma); Ewing’s sarcoma; ocular cancer (*e.g.*, intraocular melanoma, retinoblastoma); familial hypereosinophilia; gall bladder cancer; gastric cancer (*e.g.*, stomach adenocarcinoma); gastrointestinal stromal tumor (GIST); germ cell cancer; head and neck cancer (*e.g.*, head and neck squamous cell carcinoma, oral cancer (*e.g.*, oral squamous cell carcinoma), throat cancer (*e.g.*, laryngeal cancer, pharyngeal cancer, nasopharyngeal cancer, oropharyngeal cancer)); hematopoietic cancers (*e.g.*, leukemia such as acute lymphocytic leukemia (ALL) (*e.g.*, B-cell ALL, T-cell ALL), acute myelocytic leukemia (AML) (*e.g.*, B-cell AML, T-cell AML), chronic myelocytic leukemia (CML) (*e.g.*, B-cell CML, T-cell CML), and chronic lymphocytic leukemia (CLL) (*e.g.*, B-cell CLL, T-cell CLL)); lymphoma such as Hodgkin lymphoma (HL) (*e.g.*, B-cell HL, T-cell HL) and non-Hodgkin lymphoma (NHL) (*e.g.*, B-cell NHL such as diffuse large cell lymphoma (DLCL) (*e.g.*, diffuse large B-cell lymphoma), follicular lymphoma, chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma (MCL), marginal zone B-cell lymphomas (*e.g.*, mucosa-associated lymphoid tissue (MALT) lymphomas, nodal marginal zone B-cell lymphoma, splenic marginal zone B-cell lymphoma), primary mediastinal B-cell lymphoma, Burkitt lymphoma, lymphoplasmacytic lymphoma (*i.e.*, Waldenström’s macroglobulinemia), hairy cell leukemia (HCL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma and primary central nervous system (CNS) lymphoma; and T-cell NHL such as precursor T-lymphoblastic

lymphoma/leukemia, peripheral T-cell lymphoma (PTCL) (*e.g.*, cutaneous T-cell lymphoma (CTCL) (*e.g.*, mycosis fungoides, Sezary syndrome), angioimmunoblastic T-cell lymphoma, extranodal natural killer T-cell lymphoma, enteropathy type T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, and anaplastic large cell lymphoma); a mixture of one or more leukemia/lymphoma as described above; and multiple myeloma (MM)), heavy chain disease (*e.g.*, alpha chain disease, gamma chain disease, mu chain disease); hemangioblastoma; hypopharynx cancer; inflammatory myofibroblastic tumors; immunocytic amyloidosis; kidney cancer (*e.g.*, nephroblastoma *a.k.a.* Wilms' tumor, renal cell carcinoma); liver cancer (*e.g.*, hepatocellular cancer (HCC), malignant hepatoma); lung cancer (*e.g.*, bronchogenic carcinoma, small cell lung cancer (SCLC), non-small cell lung cancer (NSCLC), adenocarcinoma of the lung); leiomyosarcoma (LMS); mastocytosis (*e.g.*, systemic mastocytosis); muscle cancer; myelodysplastic syndrome (MDS); mesothelioma; myeloproliferative disorder (MPD) (*e.g.*, polycythemia vera (PV), essential thrombocytosis (ET), agnogenic myeloid metaplasia (AMM) *a.k.a.* myelofibrosis (MF), chronic idiopathic myelofibrosis, chronic myelocytic leukemia (CML), chronic neutrophilic leukemia (CNL), hypereosinophilic syndrome (HES)); neuroblastoma; neurofibroma (*e.g.*, neurofibromatosis (NF) type 1 or type 2, schwannomatosis); neuroendocrine cancer (*e.g.*, gastroenteropancreatic neuroendocrine tumor (GEP-NET), carcinoid tumor); osteosarcoma (*e.g.*, bone cancer); ovarian cancer (*e.g.*, cystadenocarcinoma, ovarian embryonal carcinoma, ovarian adenocarcinoma); papillary adenocarcinoma; pancreatic cancer (*e.g.*, pancreatic adenocarcinoma, intraductal papillary mucinous neoplasm (IPMN), Islet cell tumors); penile cancer (*e.g.*, Paget's disease of the penis and scrotum); pinealoma; primitive neuroectodermal tumor (PNT); plasma cell neoplasia; paraneoplastic syndromes; intraepithelial neoplasms; prostate cancer (*e.g.*, prostate adenocarcinoma); rectal cancer; rhabdomyosarcoma; salivary gland cancer; skin cancer (*e.g.*, squamous cell carcinoma (SCC), keratoacanthoma (KA), melanoma, basal cell carcinoma (BCC)); small bowel cancer (*e.g.*, appendix cancer); soft tissue sarcoma (*e.g.*, malignant fibrous histiocytoma (MFH), liposarcoma, malignant peripheral nerve sheath tumor (MPNST), chondrosarcoma, fibrosarcoma, myxosarcoma); sebaceous gland carcinoma; small intestine cancer; sweat gland carcinoma; synovioma; testicular cancer (*e.g.*, seminoma, testicular embryonal carcinoma); and thyroid cancer (*e.g.*, papillary carcinoma of the thyroid, papillary thyroid carcinoma (PTC), medullary thyroid cancer) ; urethral cancer; vaginal cancer; and vulvar cancer (*e.g.*, Paget's disease of the vulva).

BRIEF DESCRIPTION OF THE DRAWINGS

[00107] *Figure 1* shows that inhibition of SIK by HG 9-91-01 promotes *Mitf* transcription and pigmentation *in vitro*. Panel A shows mRNA expression of *Mitf* relative to *Rpl11* (control ribosomal gene) mRNA quantified by QPCR in normal human melanocytes 3 hours after HG 9-91-01 or vehicle control (70% ethanol; 30% propylene glycol) treatment (n=3, mean $\pm$ SD). Panel B shows mRNA expression of *Mitf* and Panel C shows *Mitf*-dependent gene, *TRPM1* over 24 hours after 4 μ M HG 9-91-01 treatment, relative to *Rpl11* and vehicle control (70% ethanol; 30% propylene glycol) *Mitf* or *TRPM1* mRNA expression, respectively, in normal human melanocytes quantified by QPCR (n=3, mean $\pm$ SD). Panel D shows cell pellets of UACC257 melanoma cells after 3 days of treatment with vehicle control (70% ethanol; 30% propylene glycol) or 4 μ M of SIK inhibitor HG 9-91-01 (n=3). For all graphs, statistical significance is reported as follows: ***P < 0.001, ****P < 0.0001.

[00108] *Figure 2* shows that topical treatment with HG 9-91-01 caused robust darkening in MC1R^{e/e};K14-Scf mice. Panel A shows MC1R^{e/e};K14-Scf mice and Tyr^{c/c};K14-Scf mice (albinos, which serve as negative controls) before treatment (Day 0) and after 7 days of treatment (Day 7) with 30 μ L of vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM HG 9-91-01 (n=4). Panel B shows reflective colorimetry measurements (CIE L* white-black color axis), and Panel D shows melanin extraction of MC1R^{e/e};K14-Scf mice and Tyr^{c/c};K14-Scf mice described in Panel A (n=4, mean $\pm$ SD). Fontana-Masson (eumelanin) (top two panels of C) and hematoxylin and eosin (bottom two panels of C) stained skin sections of MC1R^{e/e};K14-Scf mice (x400 magnification) are described in Panel A. For all graphs, statistical significance is reported as follows: ****P < 0.0001.

[00109] *Figure 3* shows that darkening induced by topical application of SIK inhibitor HG 9-91-01 (labeled “SIKi” in Figure 3) to MC1R^{e/e};K14-Scf mice is progressive and reversible. Panel A shows MC1R^{e/e};K14-Scf mice and Tyr^{c/c};K14-Scf mice after treatment with 30 μ L of vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM HG 9-91-01: before treatment (Day 0) and after 6 days of treatment (Day 6), and 40 days post-treatment (Day 46) (vehicle mouse in Day 46 photo is different from Day 0 and Day 6 photo). In Panel A, for Tyr^{c/c};K14-Scf (albino) mice vehicle: n=3 (Day 0-10), n=2 (Day 11-20); for Tyr^{c/c};K14-Scf mice HG 9-91-01: n=3; for MC1R^{e/e};K14-Scf mice vehicle: n=5 (Day 0-19), n=4 (Day 24-34). In Panel A, for MC1R^{e/e};K14-Scf mice HG 9-91-01: n=3 mean $\pm$ SD. Panels B and C show reflective colorimetry measurements (CIE L* white-black color axis) of MC1R^{e/e};K14-Scf mice (Panel B) and Tyr^{c/c};K14-Scf mice (Panel C) described in Panel A above. For all graphs, statistical significance is reported as follows: *P < 0.05, **P

< 0.01 , *** $P < 0.001$, **** $P < 0.0001$.

[00110] *Figure 4* shows that second generation SIK inhibitors (YKL 06-061 and YKL 06-062) are as efficacious in inducing the pigmentation pathway as HG 9-91-01. Panels A and B show mRNA expression of *Mitf* relative to *Rpl11* mRNA quantified by QPCR in normal human melanocytes, treated with YKL 06-061 (Panel A) YKL 06-062 (Panel B) ($n=3$, mean $\pm$ SD). Panels C and D show mRNA expression of *Mitf* (Panel C) and *Mitf*-dependent gene, *TRPM1* (Panel D), relative to *Rpl11* and vehicle control (70% ethanol; 30% propylene glycol) treatment's *Mitf* or *TRPM1* mRNA expression, respectively, in normal human melanocytes treated with 4 μ M of SIK inhibitor quantified by QPCR. For all graphs, statistical significance is reported as follows: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

[00111] *Figure 5* shows that treatment of human skin explants with 37.5 mM of SIK inhibitor induces pigmentation. Human breast skin explants were treated with passive application of vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM of SIK inhibitors YKL 06-061, YKL 06-062, or HG 9-91-01 for 8 days (10 μ L; 1x/day). Panel A shows an image taken 2 days after the end of treatment ($n=3$). Panel B shows hematoxylin and eosin (bottom panel) and Fontana-masson (top panel) staining (x400 magnification) (Panel B) of breast skin described in Panel A. Human breast skin explants were treated with a passive application of vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM of SIK inhibitors YKL 06-061, YKL 06-062, or HG 9-91-01 for 5 days (10 μ L; 2x/day). Panel C shows an image taken 1 day after end of treatment ($n=1$). Panel D shows human breast skin explants treated with passive application of control (70% ethanol; 30% propylene glycol) or 37.5 mM of SIK inhibitors YKL 06-061, YKL 06-062, or HG 9-91-01 for 6 days (10 μ L; 2x/day). The image was taken 1 day after end of treatment ($n=1$). Panel E shows human breast skin explants treated with mechanical application of vehicle control (70% ethanol; 30% propylene glycol) or 50 mM (50 μ L for 1 day; 1x/day) or 25 mM (50 μ L for 3 days; 3x/day)) of HG 9-91-01 ($n=1$). The image was taken 4 days after start of treatment. Panel F shows Fontana-masson (top panel) and hematoxylin and eosin (bottom panel) and staining (400x magnification) of human skin explants described in Panel E.

[00112] *Figure 6* shows characterization of SIK inhibitors. Panel A shows structures of HG-9-91-01, YKL-06-061 and YKL-06-062 and their biochemical IC_{50} against SIKs. Panel B shows KinomeScan kinase selectivity profiles for YKL-06-061. YKL-06-061 was profiled at a concentration of 1 μ M against a diverse panel of 468 kinases by DiscoverX. Kinases that exhibited a score of 1 or below (Score is percent relative to DMSO control. Smaller numbers

indicate stronger binding) are marked in circles. Panel C shows biochemical kinase IC₅₀'s of YKL-06-061 top hits as shown in Panel B.

[00113] *Figure 7* shows that inhibition of SIK by HG 9-91-01, YKL 06-061, and YKL 06-062 in UACC62 and UACC257 melanoma cells induces *Mitf* expression. Panel A shows mRNA expression of *Mitf* relative to *Rpl11* mRNA quantified by QPCR in melanocytes, UACC62 melanoma cells (Panels A, C and E), and UACC257 melanoma cells (Panels B, D, and E) treated with HG 9-91-01 (Panels A and B), YKL 06-061 (Panels C and D), or YKL 06-062 (Panels E and F) (n=3, mean $\pm$ SD). Panels G and H show mRNA expression of *Mitf* (Panel G) and *Mitf*-dependent gene, *TRPM1* (Panel H), relative to *Rpl11* and vehicle control (70% ethanol; 30% propylene glycol) treatment's *Mitf* or *TRPM1* mRNA expression, respectively, in normal human melanocytes treated with 4 μ M of SIK inhibitor quantified by QPCR.

[00114] *Figure 8* shows MC1R^{e/e};K14-Scf mice (Panel A) and *Tyr*^{c/c};K14-Scf mice (Panel B) after treatment with vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM HG 9-91-01 before treatment (Day 0) and after 7 days of treatment (Day 7). Panel C shows fontana-masson (eumelanin) (top panel) and hematoxylin and eosin stain (bottom panel) of *Tyr*^{c/c};K14-Scf mice (as described in Figure 2A) after treatment with vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM HG 9-91-01 before treatment (Day 0) and after 7 days of treatment (Day 7) (n=4) (x630 magnification) (Panel C). Panel D shows fontana-masson (eumelanin) (left two panels) stained skin sections of MC1R^{e/e};K14-Scf mice treated 37.5 mM for 7 days. Image is at the margin of treated and untreated area (x100 magnification) (Panel D). Panel E shows fontana-masson (eumelanin) (left panels) and hematoxylin and eosin stain (right panels) of MC1R^{e/e};K14-Scf mice and Albino;K14-Scf mice with 6 days of treatment of vehicle control (70% ethanol; 30% propylene glycol) or 37.5 mM HG 9-91-01 after 46 days from start of treatment (200x magnification).

[00115] *Figure 9* shows induction of microphthalmia-associated transcription factor (MITF) by exemplary SIK inhibitor compounds in mouse melanoma and human melanoma cell lines.

[00116] *Figure 10* shows a schematic of SIK inhibitor treatment regime for the skin experiments in *Figure 11* and in *Figure 12*.

[00117] *Figure 11* shows skin darkening 9 days post treatment with exemplary compounds YKL-05-120, YKL-05-200-01, YKL-05-200-02, YKL-05-201-1, YKL-05-203-1, YKL-05-203-2, YKL-05-204-1, YKL-05-204-2, YKL-06-203-029, YKL-06-030, YKL-06-031, YKL-06-033, YKL-06-046, YKL-06-050, YKL-06-058, YKL-06-059, YKL-06-060, YKL-06-061,

YKL-06-062, HG 9-91-01, HG 11-139-02, and HG 11-137-01, compared with DMSO treatment. Treatment of human skin explants 9 days after start of treatment with vehicle control (DMSO) or 10 mM SIK inhibitor solution for all SIK inhibitor compounds except HG 9-91-01, which was applied at a 37.5 mM concentration. Skin was treated with passive application as displayed in the schematic in *Figure 10*: 1x/day for the first day, and 2x/day for the next three days with a 2 day break and another 2x/day treatment for the next three days. Image and Colorimeter analysis were conducted 24 hours after the last treatment.


[00118] *Figure 12* shows reflective colorimetry measurements of human skin darkening 9 days after treatment with exemplary compounds under the same conditions as for *Figure 11*, with exemplary compounds YKL-05-120, YKL-05-200-01, YKL-05-200-02, YKL-05-201-1, YKL-05-203-1, YKL-05-203-2, YKL-05-204-1, YKL-05-204-2, YKL-06-203-029, YKL-06-030, YKL-06-031, YKL-06-033, YKL-06-046, YKL-06-050, YKL-06-058, YKL-06-059, YKL-06-060, YKL-06-061, YKL-06-062, HG 9-91-01, HG 11-139-02, and HG 11-137-01, compared with DMSO treatment.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

[00119] Described herein are compounds of Formulas (I), (II), (III), (IV), (V), (VI), (VI-A), and (VII), and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof. Certain compounds described herein are useful in inhibiting the activity of SIK kinase, and are useful in increasing skin pigmentation in a subject in need thereof. Increasing the skin pigmentation in a subject may reduce the risk of developing skin cancer in the subject. Also provided are pharmaceutical compositions and kits including a compound described herein. In certain embodiments, compounds for use in this application are the compounds described in U.S. Patent Applications, U.S.S.N. 14/385,077, filed September 12, 2014, U.S.S.N. 15/606,970, filed May 26, 2017, U.S.S.N. 62/027,099, filed July 21, 2014, U.S.S.N. 15/327,690, filed January 20, 2017, U.S.S.N. 62/358,524, filed July 5, 2016, U.S.S.N. 62/027,122, filed July 21, 2014, U.S.S.N. 15/327,705, filed January 20, 2017, each of which is incorporated herein by reference.

*Compounds Useful in the Invention***Compounds of Formula (I)**

[00120] In one aspect, the compound utilized in the present disclosure is of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is a substituted or unsubstituted phenyl ring or a substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur;

each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{A1}$, $-N(R^{A1})_2$, $-SR^{A1}$, $-CN$, $-SCN$, $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, $-C(=NR^{A1})N(R^{A1})_2$, $-C(=O)R^{A1}$, $-C(=O)OR^{A1}$, $-C(=O)N(R^{A1})_2$, $-NO_2$, $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$, $-NR^{A1}C(=O)N(R^{A1})_2$, $-OC(=O)R^{A1}$, $-OC(=O)OR^{A1}$, or $-OC(=O)N(R^{A1})_2$, wherein each instance of R^{A1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^{A1} groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, or 4;

L is a substituted or unsubstituted, saturated or unsaturated C_{3-10} hydrocarbon chain, optionally wherein one or more chain atoms of the hydrocarbon chain are independently replaced with $-O-$, $-S-$, $-NR^N-$, $-N=$, or $=N-$, wherein each instance of R^N is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

R^B is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each of X^A , X^B , and X^C is independently N or CR^X , wherein R^X is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{X1}$, $-N(R^{X1})_2$, $-SR^{X1}$, $-CN$, $-SCN$, $-C(=NR^{X1})R^{X1}$, $-C(=NR^{X1})OR^{X1}$, $-C(=NR^{X1})N(R^{X1})_2$, $-C(=O)R^{X1}$, $-C(=O)OR^{X1}$, $-C(=O)N(R^{X1})_2$, $-NO_2$, $-NR^{X1}C(=O)R^{X1}$, $-NR^{X1}C(=O)OR^{X1}$, $-NR^{X1}C(=O)N(R^{X1})_2$, $-OC(=O)R^{X1}$, $-OC(=O)OR^{X1}$, or $-OC(=O)N(R^{X1})_2$, wherein each instance of R^{X1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^{X1} groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

Y is $-O-$ or $-NR^Y-$, wherein R^Y is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

or when Y is $-NR^Y-$, and X^A is CR^X , R^Y and R^X of X^A are joined to form a substituted or unsubstituted, monocyclic, 5- to 7-membered heterocyclic ring that is fused with Ring B;


each instance of R^C is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{C1}$, $-N(R^{C1})_2$, $-SR^{C1}$, $-CN$, $-SCN$, $-C(=NR^{C1})R^{C1}$, $-C(=NR^{C1})OR^{C1}$, $-C(=NR^{C1})N(R^{C1})_2$, $-C(=O)R^{C1}$, $-C(=O)OR^{C1}$, $-C(=O)N(R^{C1})_2$, $-NO_2$, $-NR^{C1}C(=O)R^{C1}$, $-NR^{C1}C(=O)OR^{C1}$, $-NR^{C1}C(=O)N(R^{C1})_2$, $-OC(=O)R^{C1}$, $-OC(=O)OR^{C1}$, or $-OC(=O)N(R^{C1})_2$, wherein each instance of R^{C1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or

a sulfur protecting group when attached to a sulfur atom, or two R^{C1} groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

m is 0, 1, 2, 3, 4, or 5; and

R^D is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{D1}$, $-N(R^{D1})_2$, $-SR^{D1}$, $-CN$, $-SCN$, $-C(=NR^{D1})R^{D1}$, $-C(=NR^{D1})OR^{D1}$, $-C(=NR^{D1})N(R^{D1})_2$, $-C(=O)R^{D1}$, $-C(=O)OR^{D1}$, $-C(=O)N(R^{D1})_2$, $-NO_2$, $-NR^{D1}C(=O)R^{D1}$, $-NR^{D1}C(=O)OR^{D1}$, $-NR^{D1}C(=O)N(R^{D1})_2$, $-OC(=O)R^{D1}$, $-OC(=O)OR^{D1}$, or $-OC(=O)N(R^{D1})_2$, wherein each instance of R^{D1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.


[00121] Formula (I) includes Ring A that is unsubstituted (*e.g.*, when k is 0) or substituted with one or more substituents R^A (*e.g.*, when k is 1, 2, 3, or 4). In certain embodiments, Ring A is an unsubstituted phenyl ring. In certain embodiments, Ring A is a substituted phenyl ring. In certain embodiments, Ring A is a substituted or unsubstituted, monocyclic, 5-membered heteroaryl ring (*e.g.*, furanyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, or isothiazolyl ring), wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur. In certain embodiments, Ring A is substituted or unsubstituted pyrazolyl. In certain embodiments, Ring A is a substituted or unsubstituted, monocyclic, 6-membered heteroaryl ring (*e.g.*, a pyridyl, pyrazinyl, pyrimidinyl, or pyridazinyl ring), wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur. In certain embodiments, Ring A is substituted or unsubstituted pyridinyl. In certain embodiments, Ring A is of the formula:





wherein the radical marked with “*” is directly attached to N(R^B), and the radical marked with “**” is directly attached to L.

[00122] In Formula (I), Ring A may include one or more substituents R^A. In certain embodiments, at least two instances of R^A are different. In certain embodiments, all instances of R^A are the same. In certain embodiments, at least one instance of R^A is halogen. In certain embodiments, at least one instance of R^A is F. In certain embodiments, at least one instance of R^A is Cl. In certain embodiments, at least one instance of R^A is Br. In certain embodiments, at least one instance of R^A is I (iodine). In certain embodiments, at least one instance of R^A is substituted alkyl. In certain embodiments, at least one instance of R^A is unsubstituted alkyl. In certain embodiments, at least one instance of R^A is unsubstituted C₁₋₆ alkyl. In certain embodiments, all instances of R^A are unsubstituted C₁₋₆ alkyl. In certain embodiments, at least one instance of R^A is substituted C₁₋₆ alkyl. In certain embodiments, at least one instance of R^A is C₁₋₆ alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^A is –CH₃. In certain embodiments, all instances of R^A are –CH₃. In certain embodiments, at least one instance of R^A is substituted methyl. In certain embodiments, at least one instance of R^A is –CH₂F. In certain embodiments, at least one instance of R^A is –CHF₂. In certain embodiments, at least one instance of R^A is –CF₃. In certain embodiments, at least one instance of R^A is ethyl. In certain embodiments, at least one instance of R^A is propyl. In certain embodiments, at least one instance of R^A is butyl. In certain embodiments, at least one instance of R^A is pentyl. In certain embodiments, at least one instance of R^A is hexyl. In certain embodiments, at least one instance of R^A is substituted alkenyl. In certain embodiments, at least one instance of R^A is unsubstituted alkenyl. In certain embodiments, at least one instance of R^A is substituted alkynyl. In certain embodiments, at least one instance of R^A is unsubstituted alkynyl. In certain embodiments, at least one instance of R^A is substituted carbocyclyl. In certain embodiments, at least one instance of R^A is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^A is saturated carbocyclyl. In certain embodiments, at least one instance of R^A is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^A is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^A is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^A is substituted heterocyclyl.

In certain embodiments, at least one instance of R^A is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^A is saturated heterocyclyl. In certain embodiments, at least one instance of R^A is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^A is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^A is monocyclic heterocyclyl. In certain embodiments, at least one instance of R^A is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^A is substituted or unsubstituted, monocyclic, 3- to 7-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^A is substituted or unsubstituted oxetanyl, substituted or unsubstituted tetrahydrofuranyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl, or substituted or unsubstituted piperazinyl. In certain embodiments, at least one instance of R^A is of the formula:



(e.g., ). In certain embodiments, at least one instance of R^A is of the formula



. In certain embodiments, at least one instance of R^A is substituted aryl. In certain embodiments, at least one instance of R^A is unsubstituted aryl. In certain embodiments, at least one instance of R^A is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^A is substituted phenyl. In certain embodiments, at least one instance of R^A is unsubstituted phenyl. In certain embodiments, at least one instance of R^A is substituted heteroaryl. In certain embodiments, at least one instance of R^A is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^A is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^A is monocyclic heteroaryl. In certain embodiments, at least one instance of R^A is 5-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^A is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^A is bicyclic


heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^A is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^A is $-OR^{A1}$. In certain embodiments, at least one instance of R^A is $-OH$. In certain embodiments, at least one instance of R^A is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^A is $-OMe$. In certain embodiments, at least one instance of R^A is $-OEt$. In certain embodiments, at least one instance of R^A is $-OPr$. In certain embodiments, at least one instance of R^A is $-OBu$. In certain embodiments, at least one instance of R^A is $-OBn$. In certain embodiments, at least one instance of R^A is $-OPh$. In certain embodiments, at least one instance of R^A is $-SR^{A1}$. In certain embodiments, at least one instance of R^A is $-SH$. In certain embodiments, at least one instance of R^A is $-SMe$. In certain embodiments, at least one instance of R^A is $-N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-NH_2$. In certain embodiments, at least one instance of R^A is $-NHMe$. In certain embodiments, at least one instance of R^A is $-NMe_2$. In certain embodiments, at least one instance of R^A is $-CN$. In certain embodiments, at least one instance of R^A is $-SCN$. In certain embodiments, at least one instance of R^A is $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, or $-C(=NR^{A1})N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-C(=O)R^{A1}$ or $-C(=O)OR^{A1}$. In certain embodiments, at least one instance of R^A is $-C(=O)N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, at least one instance of R^A is $-NO_2$. In certain embodiments, at least one instance of R^A is $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$, or $-NR^{A1}C(=O)N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-OC(=O)R^{A1}$, $-OC(=O)OR^{A1}$, or $-OC(=O)N(R^{A1})_2$.

[00123] In certain embodiments, at least one instance of R^{A1} is H. In certain embodiments, at least one instance of R^{A1} is substituted acyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{A1} is acetyl. In certain embodiments, at least one instance of R^{A1} is substituted alkyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{A1} is methyl. In certain embodiments, at least one instance of R^{A1} is ethyl. In certain embodiments, at least one instance of R^{A1} is propyl. In certain embodiments, at least one instance of R^{A1} is butyl. In certain embodiments, at least one instance of R^{A1} is pentyl. In certain embodiments, at least one instance of R^{A1} is hexyl. In certain embodiments, at least one instance of R^{A1} is substituted alkenyl. In certain embodiments, at least one instance of

R^{A1} is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{A1} is substituted alkynyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{A1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{A1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{A1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{A1} is saturated heterocyclyl. In certain embodiments, at least one instance of R^{A1} is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^{A1} is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{A1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{A1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{A1} is monocyclic aryl. In certain embodiments, at least one instance of R^{A1} is substituted phenyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{A1} is bicyclic aryl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{A1} is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{A1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{A1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{A1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{A1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{A1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{A1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{A1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{A1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{A1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of


R^{A1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{A1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{A1} are joined to form a heterocyclic ring, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{A1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{A1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{A1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.


[00124] In certain embodiments, k is 0. In certain embodiments, k is 1. In certain embodiments, k is 2. In certain embodiments, k is 3. In certain embodiments, k is 4. In certain


embodiments, k is 1 and R^A is  unsubstituted C_{1-6} alkyl.

[00125] Formula (I) includes divalent linker L. L consists of a chain, and optionally one or more hydrogen atoms and/or one or more substituents (*e.g.*, =O) on the chain, where any two substituents may optionally be joined to form a ring. In certain embodiments, the molecular weight of L is not more than about 300 g/mol, not more than about 200 g/mol, not more than about 150 g/mol, not more than about 100 g/mol, or not more than 80 g/mol. In certain embodiments, the molecular weight of L is between 50 and 150 g/mol, inclusive. In certain embodiments, L consists of not more than about 70 atoms, not more than about 50 atoms, not more than about 30 atoms, not more than about 20 atoms, or not more than 15 atoms. In certain embodiments, L consists of between 10 and 30 atoms, inclusive. In certain embodiments, L does not include unsaturated bonds in the chain. In certain embodiments, L consists of one unsaturated bond in the chain. In certain embodiments, L consists of 2, 3, or 4 unsaturated bonds in the chain. In certain embodiments, L is a substituted or unsubstituted, saturated or unsaturated C_{3-10} hydrocarbon chain (*e.g.*, a C_{5-6} hydrocarbon chain). In certain embodiments, L is a substituted or unsubstituted, saturated or unsaturated C_{3-10} hydrocarbon chain (*e.g.*, a C_{5-6} hydrocarbon chain), wherein one chain atom of the hydrocarbon chain is replaced with $-O-$, $-S-$, $-NR^N-$, $-N=$, or $=N-$. In certain embodiments, L is a substituted or unsubstituted, saturated or unsaturated, C_{3-10} hydrocarbon chain (*e.g.*, a C_{5-6} hydrocarbon chain), wherein 2, 3, 4, or 5 chain atoms of the hydrocarbon chain are independently replaced with $-O-$, $-S-$, $-NR^N-$, $-N=$, or $=N-$. In certain embodiments, L is a substituted or


unsubstituted, saturated or unsaturated C₅₋₆ hydrocarbon chain, wherein one or two chain atoms of the hydrocarbon chain are independently replaced with -O-, -S-, or -NR^N-. In

certain embodiments, L is of the formula: . In certain embodiments, L


is of the formula: . In certain embodiments, L is of the formula:




In certain embodiments, L is of the formula:




In certain embodiments, L is of the formula:




In certain embodiments, L is of the formula:







In certain embodiments, L is of the formula:



In certain embodiments, L is of the formula:




certain embodiments, L is of the formula:



certain embodiments, L is of the formula:



of the formula:



[00126] In certain embodiments, at least two instances of R^N are different. In certain embodiments, all instances of R^N are the same. In certain embodiments, at least one instance of R^N is H. In certain embodiments, each instance of R^N is H. In certain embodiments, at least one instance of R^N is substituted acyl. In certain embodiments, at least one instance of R^N is unsubstituted acyl. In certain embodiments, at least one instance of R^N is acetyl. In certain embodiments, at least one instance of R^N is unsubstituted C_{1-6} alkyl. In certain embodiments, each instance of R^N is independently unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^N is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^N is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^N is unsubstituted methyl. In certain embodiments, at least one instance of R^N is substituted methyl. In certain embodiments, at least one instance of R^N is $-CH_2F$. In certain embodiments, at least one instance of R^N is $-CHF_2$. In certain embodiments, at least one instance of R^N is $-CF_3$. In certain embodiments, at least one instance of R^N is ethyl. In certain embodiments, at least one instance of R^N is propyl. In certain embodiments, at least one

instance of R^N is butyl. In certain embodiments, at least one instance of R^N is pentyl. In certain embodiments, at least one instance of R^N is hexyl. In certain embodiments, at least one instance of R^N is a nitrogen protecting group. In certain embodiments, at least one instance of R^N is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts.

[00127] Formula (I) includes substituent R^B on a nitrogen atom. In certain embodiments, R^B is H. In certain embodiments, R^B is substituted acyl. In certain embodiments, R^B is unsubstituted acyl. In certain embodiments, R^B is acetyl. In certain embodiments, R^B is unsubstituted C_{1-6} alkyl. In certain embodiments, R^B is substituted C_{1-6} alkyl. In certain embodiments, R^B is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, R^B is unsubstituted methyl. In certain embodiments, R^B is substituted methyl. In certain embodiments, R^B is $-CH_2F$. In certain embodiments, R^B is $-CHF_2$. In certain embodiments, R^B is $-CF_3$. In certain embodiments, R^B is ethyl. In certain embodiments, R^B is propyl. In certain embodiments, R^B is butyl. In certain embodiments, R^B is pentyl. In certain embodiments, R^B is hexyl. In certain embodiments, R^B is a nitrogen protecting group. In certain embodiments, R^B is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts.

[00128] Formula (I) includes Ring B that includes moieties X^A , X^B , and X^C in the ring system. In certain embodiments, X^A is CR^X , and each of X^B and X^C is N. In certain embodiments, X^A is CH, and each of X^B and X^C is N. In certain embodiments, X^B is CR^X , and each of X^A and X^C is N. In certain embodiments, X^B is CH, and each of X^A and X^C is N. In certain embodiments, X^C is CR^X , and each of X^A and X^B is N. In certain embodiments, X^C is CH, and each of X^A and X^B is N. In certain embodiments, X^A is N, and each of X^B and X^C is independently CR^X . In certain embodiments, X^A is N, and each of X^B and X^C is CH. In certain embodiments, X^B is N, and each of X^A and X^C is independently CR^X . In certain embodiments, X^B is N, and each of X^A and X^C is CH. In certain embodiments, X^C is N, and each of X^A and X^B is independently CR^X . In certain embodiments, X^C is N, and each of X^A and X^B is CH. In certain embodiments, each of X^A , X^B , and X^C is independently CR^X . In certain embodiments, each of X^A , X^B , and X^C is CH.

[00129] In certain embodiments, when X^A , X^B , or X^C is CR^X , R^X is H. In certain embodiments, R^X is halogen. In certain embodiments, R^X is F. In certain embodiments, R^X is Cl. In certain embodiments, R^X is Br. In certain embodiments, R^X is I (iodine). In certain embodiments, R^X is substituted alkyl. In certain embodiments, R^X is unsubstituted alkyl. In certain embodiments, R^X is unsubstituted C_{1-6} alkyl. In certain embodiments, R^X is substituted C_{1-6} alkyl. In certain embodiments, R^X is C_{1-6} alkyl substituted with at least one

halogen. In certain embodiments, R^X is $-\text{CH}_3$. In certain embodiments, R^X is substituted methyl. In certain embodiments, R^X is $-\text{CH}_2\text{F}$. In certain embodiments, R^X is $-\text{CHF}_2$. In certain embodiments, R^X is $-\text{CF}_3$. In certain embodiments, R^X is ethyl. In certain embodiments, R^X is propyl. In certain embodiments, R^X is butyl. In certain embodiments, R^X is pentyl. In certain embodiments, R^X is hexyl. In certain embodiments, R^X is substituted alkenyl. In certain embodiments, R^X is unsubstituted alkenyl. In certain embodiments, R^X is substituted alkynyl. In certain embodiments, R^X is unsubstituted alkynyl. In certain embodiments, R^X is substituted carbocyclyl. In certain embodiments, R^X is unsubstituted carbocyclyl. In certain embodiments, R^X is saturated carbocyclyl. In certain embodiments, R^X is unsaturated carbocyclyl. In certain embodiments, R^X is monocyclic carbocyclyl. In certain embodiments, R^X is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^X is substituted heterocyclyl. In certain embodiments, R^X is unsubstituted heterocyclyl. In certain embodiments, R^X is saturated heterocyclyl. In certain embodiments, R^X is unsaturated heterocyclyl. In certain embodiments, R^X is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^X is monocyclic heterocyclyl. In certain embodiments, R^X is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^X is substituted aryl. In certain embodiments, R^X is unsubstituted aryl. In certain embodiments, R^X is 6- to 10-membered aryl. In certain embodiments, R^X is substituted phenyl. In certain embodiments, R^X is unsubstituted phenyl. In certain embodiments, R^X is substituted heteroaryl. In certain embodiments, R^X is unsubstituted heteroaryl. In certain embodiments, R^X is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^X is monocyclic heteroaryl. In certain embodiments, R^X is 5-membered, monocyclic heteroaryl. In certain embodiments, R^X is 6-membered, monocyclic heteroaryl. In certain embodiments, R^X is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, R^X is $-\text{OR}^{X1}$. In certain embodiments, R^X is $-\text{OH}$. In certain embodiments, R^X is $-\text{O}(\text{substituted or unsubstituted } \text{C}_{1-6} \text{ alkyl})$. In certain embodiments, R^X is $-\text{OMe}$. In certain embodiments, R^X is $-\text{OEt}$. In certain embodiments, R^X is $-\text{OPr}$. In certain embodiments, R^X is $-\text{OBu}$. In certain embodiments, R^X is $-\text{OBn}$. In certain embodiments, R^X is $-\text{OPh}$. In certain embodiments, R^X is $-\text{SR}^{X1}$. In certain embodiments, R^X is $-\text{SH}$. In certain embodiments, R^X is $-\text{SMe}$. In certain embodiments, R^X is $-\text{N}(\text{R}^{X1})_2$. In certain embodiments, R^X is $-\text{NH}_2$. In certain embodiments, R^X is $-\text{NHMe}$. In certain embodiments, R^X is $-\text{NMe}_2$. In

certain embodiments, R^X is $-\text{CN}$. In certain embodiments, R^X is $-\text{SCN}$. In certain embodiments, R^X is $-\text{C}(=\text{NR}^{X1})\text{R}^{X1}$, $-\text{C}(=\text{NR}^{X1})\text{OR}^{X1}$, or $-\text{C}(=\text{NR}^{X1})\text{N}(\text{R}^{X1})_2$. In certain embodiments, R^X is $-\text{C}(=\text{O})\text{R}^{X1}$ or $-\text{C}(=\text{O})\text{OR}^{X1}$. In certain embodiments, R^X is $-\text{C}(=\text{O})\text{N}(\text{R}^{X1})_2$. In certain embodiments, R^X is $-\text{C}(=\text{O})\text{NMe}_2$, $-\text{C}(=\text{O})\text{NHMe}$, or $-\text{C}(=\text{O})\text{NH}_2$. In certain embodiments, R^X is $-\text{NO}_2$. In certain embodiments, R^X is $-\text{NR}^{X1}\text{C}(=\text{O})\text{R}^{X1}$, $-\text{NR}^{X1}\text{C}(=\text{O})\text{OR}^{X1}$, or $-\text{NR}^{X1}\text{C}(=\text{O})\text{N}(\text{R}^{X1})_2$. In certain embodiments, R^X is $-\text{OC}(=\text{O})\text{R}^{X1}$, $-\text{OC}(=\text{O})\text{OR}^{X1}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^{X1})_2$.

[00130] In certain embodiments, R^{X1} is H. In certain embodiments, R^{X1} is substituted acyl. In certain embodiments, R^{X1} is unsubstituted acyl. In certain embodiments, R^{X1} is acetyl. In certain embodiments, R^{X1} is substituted alkyl. In certain embodiments, R^{X1} is unsubstituted alkyl. In certain embodiments, R^{X1} is unsubstituted C_{1-6} alkyl. In certain embodiments, R^{X1} is methyl. In certain embodiments, R^{X1} is ethyl. In certain embodiments, R^{X1} is propyl. In certain embodiments, R^{X1} is butyl. In certain embodiments, R^{X1} is pentyl. In certain embodiments, R^{X1} is hexyl. In certain embodiments, R^{X1} is substituted alkenyl. In certain embodiments, R^{X1} is unsubstituted alkenyl. In certain embodiments, R^{X1} is substituted alkynyl. In certain embodiments, R^{X1} is unsubstituted alkynyl. In certain embodiments, R^{X1} is substituted or unsubstituted carbocyclyl. In certain embodiments, R^{X1} is saturated carbocyclyl. In certain embodiments, R^{X1} is unsaturated carbocyclyl. In certain embodiments, R^{X1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^{X1} is substituted or unsubstituted heterocyclyl. In certain embodiments, R^{X1} is saturated heterocyclyl. In certain embodiments, R^{X1} is unsaturated heterocyclyl. In certain embodiments, R^{X1} is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^{X1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^{X1} is substituted or unsubstituted aryl. In certain embodiments, R^{X1} is 6- to 10-membered aryl. In certain embodiments, R^{X1} is monocyclic aryl. In certain embodiments, R^{X1} is substituted phenyl. In certain embodiments, R^{X1} is unsubstituted phenyl. In certain embodiments, R^{X1} is bicyclic aryl. In certain embodiments, R^{X1} is substituted or unsubstituted heteroaryl. In certain embodiments, R^{X1} is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^{X1} is monocyclic heteroaryl. In certain embodiments, R^{X1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, R^{X1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, R^{X1} is a

nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, R^{X1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{X1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{X1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{X1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{X1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulphenyl, 2-pyridine-sulphenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{X1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{X1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{X1} are joined to form a heterocyclic ring, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{X1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{X1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{X1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.

[00131] Formula (I) includes divalent moiety Y. In certain embodiments, Y is $-O-$. In certain embodiments, Y is $-NR^Y-$. In certain embodiments, Y is $-NH-$. In certain embodiments, Y is $-NR^Y-$; X^A is CR^X ; and R^Y and R^X of X^A are joined to form a substituted or unsubstituted, monocyclic, 5- to 7-membered heterocyclic ring that is fused with Ring B, optionally wherein there are 2 or 3 nitrogen atoms, 0 or 1 oxygen atom, and 0 or 1 sulfur atom, in the monocyclic heterocyclic ring system. The monocyclic heterocyclic ring formed by joining R^Y and R^X of X^A is fused with Ring B to form a substituted or unsubstituted, bicyclic, 9- to 11-membered ring. In certain embodiments, Y is $-NR^Y-$; X^A is CR^X ; and R^Y and R^X of X^A are joined to form a substituted or unsubstituted, monocyclic, 6-membered heterocyclic ring that is fused with Ring B.

[00132] In certain embodiments, when Y is $-NR^Y-$, R^Y is H. In certain embodiments, R^Y is substituted acyl. In certain embodiments, R^Y is unsubstituted acyl. In certain embodiments, R^Y is acetyl. In certain embodiments, R^Y is unsubstituted C_{1-6} alkyl. In certain embodiments, R^Y is substituted C_{1-6} alkyl. In certain embodiments, R^Y is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, R^Y is unsubstituted methyl. In certain embodiments, R^Y is substituted methyl. In certain embodiments, R^Y is $-CH_2F$. In certain embodiments, R^Y is $-$

CHF₂. In certain embodiments, R^Y is –CF₃. In certain embodiments, R^Y is ethyl. In certain embodiments, R^Y is propyl. In certain embodiments, R^Y is butyl. In certain embodiments, R^Y is pentyl. In certain embodiments, R^Y is hexyl. In certain embodiments, R^Y is a nitrogen protecting group. In certain embodiments, R^Y is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts.


[00133] In Formula (I), Ring B includes substituent R^D. In certain embodiments, R^D is H. In certain embodiments, R^D is halogen. In certain embodiments, R^D is F. In certain embodiments, R^D is Cl. In certain embodiments, R^D is Br. In certain embodiments, R^D is I (iodine). In certain embodiments, R^D is substituted alkyl. In certain embodiments, R^D is unsubstituted alkyl. In certain embodiments, R^D is unsubstituted C₁₋₆ alkyl. In certain embodiments, R^D is substituted C₁₋₆ alkyl. In certain embodiments, R^D is C₁₋₆ alkyl substituted with at least one halogen. In certain embodiments, R^D is –CH₃. In certain embodiments, R^D is substituted methyl. In certain embodiments, R^D is –CH₂F. In certain embodiments, R^D is –CHF₂. In certain embodiments, R^D is –CF₃. In certain embodiments, R^D is ethyl. In certain embodiments, R^D is propyl. In certain embodiments, R^D is butyl. In certain embodiments, R^D is pentyl. In certain embodiments, R^D is hexyl. In certain embodiments, R^D is substituted alkenyl. In certain embodiments, R^D is unsubstituted alkenyl. In certain embodiments, R^D is substituted alkynyl. In certain embodiments, R^D is unsubstituted alkynyl. In certain embodiments, R^D is substituted carbocyclyl. In certain embodiments, R^D is unsubstituted carbocyclyl. In certain embodiments, R^D is saturated carbocyclyl. In certain embodiments, R^D is unsaturated carbocyclyl. In certain embodiments, R^D is monocyclic carbocyclyl. In certain embodiments, R^D is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^D is substituted heterocyclyl. In certain embodiments, R^D is unsubstituted heterocyclyl. In certain embodiments, R^D is saturated heterocyclyl. In certain embodiments, R^D is unsaturated heterocyclyl. In certain embodiments, R^D is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^D is monocyclic heterocyclyl. In certain embodiments, R^D is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^D is substituted aryl. In certain embodiments, R^D is unsubstituted aryl. In certain embodiments, R^D is 6- to 10-membered aryl. In certain embodiments, R^D is substituted phenyl. In certain embodiments, R^D is unsubstituted phenyl. In certain embodiments, R^D is substituted heteroaryl. In certain embodiments, R^D is unsubstituted heteroaryl. In certain embodiments, R^D is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group



consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^D is monocyclic heteroaryl. In certain embodiments, R^D is 5-membered, monocyclic heteroaryl. In certain embodiments, R^D is 6-membered, monocyclic heteroaryl. In certain embodiments, R^D is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, R^D is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, R^D is $-OR^{D1}$. In certain embodiments, R^D is $-OH$. In certain embodiments, R^D is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, R^D is $-OMe$. In certain embodiments, R^D is $-OEt$. In certain embodiments, R^D is $-OPr$. In certain embodiments, R^D is $-OBu$. In certain embodiments, R^D is $-OBn$. In certain embodiments, R^D is $-OPh$. In certain embodiments, R^D is $-SR^{D1}$. In certain embodiments, R^D is $-SH$. In certain embodiments, R^D is $-SMe$. In certain embodiments, R^D is $-N(R^{D1})_2$. In certain embodiments, R^D is $-NH_2$. In certain embodiments, R^D is $-NHMe$. In certain embodiments, R^D is $-NMe_2$. In certain embodiments, R^D is $-CN$. In certain embodiments, R^D is $-SCN$. In certain embodiments, R^D is $-C(=NR^{D1})R^{D1}$, $-C(=NR^{D1})OR^{D1}$, or $-C(=NR^{D1})N(R^{D1})_2$. In certain embodiments, R^D is $-C(=O)R^{D1}$ or $-C(=O)OR^{D1}$. In certain embodiments, R^D is $-C(=O)N(R^{D1})_2$. In certain embodiments, R^D is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, R^D is $-NO_2$. In certain embodiments, R^D is $-NR^{D1}C(=O)R^{D1}$, $-NR^{D1}C(=O)OR^{D1}$, or $-NR^{D1}C(=O)N(R^{D1})_2$. In certain embodiments, R^D is $-OC(=O)R^{D1}$, $-OC(=O)OR^{D1}$, or $-OC(=O)N(R^{D1})_2$.


[00134] In certain embodiments, R^{D1} is H. In certain embodiments, R^{D1} is substituted acyl. In certain embodiments, R^{D1} is unsubstituted acyl. In certain embodiments, R^{D1} is acetyl. In certain embodiments, R^{D1} is substituted alkyl. In certain embodiments, R^{D1} is unsubstituted alkyl. In certain embodiments, R^{D1} is unsubstituted C_{1-6} alkyl. In certain embodiments, R^{D1} is methyl. In certain embodiments, R^{D1} is ethyl. In certain embodiments, R^{D1} is propyl. In certain embodiments, R^{D1} is butyl. In certain embodiments, R^{D1} is pentyl. In certain embodiments, R^{D1} is hexyl. In certain embodiments, R^{D1} is substituted alkenyl. In certain embodiments, R^{D1} is unsubstituted alkenyl. In certain embodiments, R^{D1} is substituted alkynyl. In certain embodiments, R^{D1} is unsubstituted alkynyl. In certain embodiments, R^{D1} is substituted or unsubstituted carbocyclyl. In certain embodiments, R^{D1} is saturated carbocyclyl. In certain embodiments, R^{D1} is unsaturated carbocyclyl. In certain embodiments, R^{D1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^{D1} is substituted or unsubstituted heterocyclyl. In certain embodiments, R^{D1} is saturated heterocyclyl. In certain embodiments, R^{D1} is unsaturated heterocyclyl. In certain embodiments, R^{D1} is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are


independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^{D1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^{D1} is substituted or unsubstituted aryl. In certain embodiments, R^{D1} is 6- to 10-membered aryl. In certain embodiments, R^{D1} is monocyclic aryl. In certain embodiments, R^{D1} is substituted phenyl. In certain embodiments, R^{D1} is unsubstituted phenyl. In certain embodiments, R^{D1} is bicyclic aryl. In certain embodiments, R^{D1} is substituted or unsubstituted heteroaryl. In certain embodiments, R^{D1} is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^{D1} is monocyclic heteroaryl. In certain embodiments, R^{D1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, R^{D1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, R^{D1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, R^{D1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{D1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{D1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{D1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{D1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{D1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{D1} are joined to form a heterocyclic ring, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{D1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{D1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{D1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.


[00135] Formula (I) includes Ring C that is unsubstituted (*e.g.*, when *m* is 0) or substituted with one or more substituents R^C (*e.g.*, when *m* is 1, 2, 3, 4, or 5). In certain embodiments,


Ring C is of the formula: . In certain embodiments, Ring C is of the formula:



. In certain embodiments, Ring C is of the formula: . In certain





embodiments, Ring C is of the formula: . In certain embodiments, Ring C is of the

formula: , wherein each instance of R^C is independently substituted or unsubstituted

C_{1-6} alkyl. In certain embodiments, Ring C is of the formula: . In certain

embodiments, Ring C is of the formula: . In certain embodiments, Ring C is of the

formula: . In certain embodiments, Ring C is of the formula: ,

, ,
, or . In certain embodiments, at least two instances of R^C are

different. In certain embodiments, all instances of R^C are the same. In certain embodiments, at least one instance of R^C is halogen. In certain embodiments, at least one instance of R^C is F. In certain embodiments, at least one instance of R^C is Cl. In certain embodiments, at least one instance of R^C is Br. In certain embodiments, at least one instance of R^C is I (iodine). In certain embodiments, at least one instance of R^C is substituted alkyl. In certain embodiments, at least one instance of R^C is unsubstituted alkyl. In certain embodiments, at least one

instance of R^C is unsubstituted C_{1-6} alkyl. In certain embodiments, all instances of R^C are unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^C is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^C is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^C is $-CH_3$. In certain embodiments, all instances of R^C are $-CH_3$. In certain embodiments, at least one instance of R^C is substituted methyl. In certain embodiments, at least one instance of R^C is $-CH_2F$. In certain embodiments, at least one instance of R^C is $-CHF_2$. In certain embodiments, at least one instance of R^C is $-CF_3$. In certain embodiments, at least one instance of R^C is ethyl. In certain embodiments, at least one instance of R^C is propyl. In certain embodiments, at least one instance of R^C is butyl. In certain embodiments, at least one instance of R^C is pentyl. In certain embodiments, at least one instance of R^C is hexyl. In certain embodiments, each instance of R^C is independently halogen (*e.g.*, Cl) or substituted or unsubstituted C_{1-6} alkyl (*e.g.*, unsubstituted C_{1-6} alkyl (*e.g.*, Me)). In certain embodiments, at least one instance of R^C is substituted alkenyl. In certain embodiments, at least one instance of R^C is unsubstituted alkenyl. In certain embodiments, at least one instance of R^C is substituted alkynyl. In certain embodiments, at least one instance of R^C is unsubstituted alkynyl. In certain embodiments, at least one instance of R^C is substituted carbocyclyl. In certain embodiments, at least one instance of R^C is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^C is saturated carbocyclyl. In certain embodiments, at least one instance of R^C is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^C is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^C is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^C is substituted heterocyclyl. In certain embodiments, at least one instance of R^C is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^C is saturated heterocyclyl. In certain embodiments, at least one instance of R^C is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^C is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^C is monocyclic heterocyclyl. In certain embodiments, at least one instance of R^C is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^C is substituted aryl. In certain embodiments, at least one instance of R^C is unsubstituted aryl. In certain embodiments, at least one instance of R^C is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^C is substituted phenyl. In certain embodiments, at least one instance of R^C is unsubstituted phenyl. In certain embodiments, at least one instance of R^C is substituted heteroaryl. In

certain embodiments, at least one instance of R^C is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^C is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^C is monocyclic heteroaryl. In certain embodiments, at least one instance of R^C is 5-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^C is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^C is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^C is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^C is $-OR^{C1}$. In certain embodiments, at least one instance of R^C is $-OH$. In certain embodiments, at least one instance of R^C is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^C is $-OMe$. In certain embodiments, at least one instance of R^C is $-OEt$. In certain embodiments, at least one instance of R^C is $-OPr$. In certain embodiments, at least one instance of R^C is $-OBu$. In certain embodiments, at least one instance of R^C is $-OBn$. In certain embodiments, at least one instance of R^C is $-OPh$. In certain embodiments, at least one instance of R^C is $-SR^{C1}$. In certain embodiments, at least one instance of R^C is $-SH$. In certain embodiments, at least one instance of R^C is $-SMe$. In certain embodiments, at least one instance of R^C is $-N(R^{C1})_2$. In certain embodiments, at least one instance of R^C is $-NH_2$. In certain embodiments, at least one instance of R^C is $-NHMe$. In certain embodiments, at least one instance of R^C is $-NMe_2$. In certain embodiments, at least one instance of R^C is $-CN$. In certain embodiments, at least one instance of R^C is $-SCN$. In certain embodiments, at least one instance of R^C is $-C(=NR^{C1})R^{C1}$, $-C(=NR^{C1})OR^{C1}$, or $-C(=NR^{C1})N(R^{C1})_2$. In certain embodiments, at least one instance of R^C is $-C(=O)R^{C1}$ or $-C(=O)OR^{C1}$. In certain embodiments, at least one instance of R^C is $-C(=O)N(R^{C1})_2$. In certain embodiments, at least one instance of R^C is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, at least one instance of R^C is $-NO_2$. In certain embodiments, at least one instance of R^C is $-NR^{C1}C(=O)R^{C1}$, $-NR^{C1}C(=O)OR^{C1}$, or $-NR^{C1}C(=O)N(R^{C1})_2$. In certain embodiments, at least one instance of R^C is $-OC(=O)R^{C1}$, $-OC(=O)OR^{C1}$, or $-OC(=O)N(R^{C1})_2$.

[00136] In certain embodiments, at least one instance of R^{C1} is H. In certain embodiments, at least one instance of R^{C1} is substituted acyl. In certain embodiments, at least one instance of R^{C1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{C1} is acetyl. In certain embodiments, at least one instance of R^{C1} is substituted alkyl. In certain embodiments,


at least one instance of R^{C1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{C1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{C1} is methyl. In certain embodiments, at least one instance of R^{C1} is ethyl. In certain embodiments, at least one instance of R^{C1} is propyl. In certain embodiments, at least one instance of R^{C1} is butyl. In certain embodiments, at least one instance of R^{C1} is pentyl. In certain embodiments, at least one instance of R^{C1} is hexyl. In certain embodiments, at least one instance of R^{C1} is substituted alkenyl. In certain embodiments, at least one instance of R^{C1} is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{C1} is substituted alkynyl. In certain embodiments, at least one instance of R^{C1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{C1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{C1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{C1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{C1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{C1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{C1} is saturated heterocyclyl. In certain embodiments, at least one instance of R^{C1} is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^{C1} is heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{C1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{C1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{C1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{C1} is monocyclic aryl. In certain embodiments, at least one instance of R^{C1} is substituted phenyl. In certain embodiments, at least one instance of R^{C1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{C1} is bicyclic aryl. In certain embodiments, at least one instance of R^{C1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{C1} is heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{C1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{C1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{C1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{C1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{C1} is Bn, Boc,

Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{C1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{C1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{C1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{C1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{C1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{C1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{C1} are joined to form a heterocyclic ring, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{C1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{C1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{C1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.

[00137] In certain embodiments, *m* is 0. In certain embodiments, *m* is 1. In certain embodiments, *m* is 2. In certain embodiments, *m* is 3. In certain embodiments, *m* is 4. In certain embodiments, *m* is 5.


[00138] In certain embodiments, *m* is 2; and each instance of R^C is halogen (*e.g.*, Cl). In certain embodiments, *m* is 2; and each instance of R^C is substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, *m* is 2; and each instance of R^C is methyl. In certain embodiments, *m* is 2; and each instance of R^C is independently halogen (*e.g.*, Cl) or substituted or unsubstituted C_{1-6} alkyl (*e.g.*, unsubstituted C_{1-6} alkyl (*e.g.*, Me)).

[00139] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00140] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00141] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00142] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00143] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00144] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00145] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00146] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00147] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof, wherein each instance of R^E is independently hydrogen, halogen, or substituted or unsubstituted C₁₋₆ alkyl. In certain embodiments, the two instances of R^E are the same. In certain embodiments, the two instances of R^E are not the same. In certain embodiments, at least one instance of R^E is hydrogen. In certain embodiments, each instance of R^E is hydrogen. In certain embodiments, at least one instance of R^E is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^E is substituted or unsubstituted C₁₋₆ alkyl. In certain embodiments, at least


one instance of R^E is Me. In certain embodiments, at least one instance of R^E is substituted methyl (*e.g.*, $-\text{CF}_3$ or Bn), Et, substituted ethyl (*e.g.*, perfluoroethyl), Pr, substituted propyl (*e.g.*, perfluoropropyl), Bu, or substituted butyl (*e.g.*, perfluorobutyl).

[00148] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00149] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00150] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00151] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00152] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00153] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00154] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:



[00155] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00157] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00158] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:






or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00159] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00160] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00161] In certain embodiments, the compound of Formula (I) useful in the present invention is of the formula:






or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

Compounds of Formula (II)

[00162] In another aspect, the compound utilized in the present disclosure is of Formula (II):



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:

R^A is substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl,

, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocyclyl;

each instance of R^{A1} is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or

unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^a are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

each instance of R^b is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^b are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, or 4;

each instance of R^B is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

m is 0, 1, 2, 3, 4, or 5;

R^C is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^D is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^E is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^F is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Ring A is substituted or unsubstituted phenyl; substituted or unsubstituted, polycyclic aryl; substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl; or substituted or unsubstituted, polycyclic heteroaryl;

each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or



unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^b)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^b)\text{R}^a$, $-\text{C}(=\text{NR}^b)\text{OR}^a$, $-\text{C}(=\text{NR}^b)\text{N}(\text{R}^b)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^b)_2$, $-\text{NO}_2$, $-\text{NR}^b\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^b\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^b\text{C}(=\text{O})\text{N}(\text{R}^b)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^b)_2$;



n is 0, 1, 2, 3, or 4, as valency permits;




L is a bond or a substituted or unsubstituted C_{1-6} hydrocarbon chain, optionally wherein one or more chain atoms of the hydrocarbon chain are independently replaced with $-\text{C}(=\text{O})-$, $-\text{O}-$, $-\text{S}-$, $-\text{NR}^b-$, $-\text{N}=\text{}$, or $=\text{N}-$; and



R^H is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted heterocyclyl, $-\text{OH}$, or $-\text{N}(\text{R}^c)_2$, wherein each instance of R^c is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.



[00163] Formula (II) includes substituent R^A . In certain embodiments, R^A is substituted alkenyl. In certain embodiments, R^A is unsubstituted alkenyl. In certain embodiments, R^A is substituted alkynyl. In certain embodiments, R^A is unsubstituted alkynyl. In certain embodiments, R^A is substituted phenyl. In certain embodiments, R^A is of the


formula:  or . In certain embodiments, k is 0. In certain






embodiments, R^A is of the formula:  or . In certain embodiments, k is 1. In


certain embodiments, R^A is of the formula: , , , or



. In certain embodiments, R^A is of the formula: . In certain


embodiments, R^A is . In certain embodiments, R^A is . In certain

embodiments, k is 2. In certain embodiments, R^A is of the formula: ,

, , , , or . In certain

embodiments, k is 3. In certain embodiments, R^A is of the formula: ,

, or . In certain embodiments, k is 4. In certain embodiments,

R^A is of the formula: .

[00164] In certain embodiments, when R^A is substituted phenyl, R^A includes one or more R^{A1} substituents. In certain embodiments, at least one instance of R^{A1} is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one R^{A1} is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted methyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted ethyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted propyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{2-6} alkenyl). In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted alkynyl (*e.g.*,


substituted or unsubstituted C₂₋₆ alkynyl). In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclyl comprising zero, one, or two double bonds in the carbocyclic ring system). In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted piperazinyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted oxetanyl, substituted or unsubstituted tetrahydrofuranyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl, or substituted or unsubstituted piperazinyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of R^{A1} is benzyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur; or substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^{A1} is -OR^a (*e.g.*, -OH or -OMe). In certain embodiments, at least one instance of R^{A1} is -N(R^b)₂, -SR^a, -CN, -SCN, -C(=NR^b)R^a, -C(=NR^b)OR^a, -C(=NR^b)N(R^b)₂, -C(=O)R^a, -C(=O)OR^a, -C(=O)N(R^b)₂, -NO₂, -NR^bC(=O)R^a, -NR^bC(=O)OR^a, -NR^bC(=O)N(R^b)₂, -OC(=O)R^a, -OC(=O)OR^a, or -OC(=O)N(R^b)₂.

[00165] In certain embodiments, at least one instance of R^a is hydrogen. In certain embodiments, at least one instance of R^a is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^a is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C₁₋₆ alkyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted methyl. In certain embodiments, at least one instance of R^a is substituted or unsubstituted ethyl. In certain embodiments, at least one instance of R^a is substituted or unsubstituted propyl. In certain embodiments, at least one instance of R^a is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C₂₋₆ alkenyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted alkynyl (*e.g.*,

substituted or unsubstituted C₂₋₆ alkynyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclyl comprising zero, one, or two double bonds in the carbocyclic ring system). In certain embodiments, at least one instance of R^a is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^a is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of R^{A1} is benzyl. In certain embodiments, at least one instance of R^a is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^a is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur; or substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^a is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, at least one instance of R^a is a sulfur protecting group when attached to a sulfur atom.


[00166] In certain embodiments, at least one instance of R^b is hydrogen. In certain embodiments, at least one instance of R^b is substituted or unsubstituted C₁₋₆ alkyl (*e.g.*, substituted or unsubstituted methyl, ethyl, or propyl). In certain embodiments, at least one instance of R^b is a nitrogen protecting group (*e.g.*, benzyl (Bn), t-butyl carbonate (BOC or Boc), benzyl carbamate (Cbz), 9-fluorenylmethyl carbonate (Fmoc), trifluoroacetyl, triphenylmethyl, acetyl, or p-toluenesulfonamide (Ts)). In certain embodiments, two instances of R^b are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur; or substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, two instances of R^b are taken together with their intervening

atoms to form substituted or unsubstituted piperazinyl. In certain embodiments, two instances



of R^b are taken together with their intervening atoms to form or .

[00167] In certain embodiments, R^A is substituted or unsubstituted heteroaryl. In certain embodiments, R^A is substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur (*e.g.*, furanyl, thiophenyl, pyridinyl, or pyrimidinyl, *etc.*) In certain embodiments, R^A is substituted or unsubstituted furanyl. In certain embodiments, R^A is substituted or unsubstituted thiophenyl. In certain embodiments, R^A is substituted or unsubstituted pyridinyl. In certain embodiments, R^A is of the formula:



embodiments, R^A is of the formula: , , or . In certain embodiments, k is 1.


In certain embodiments, R^A is of the formula: , , , or .

In certain embodiments, R^A is of the formula: or . In certain


embodiments, R^A is of the formula: , , , or . In certain


embodiments, R^A is of the formula: or . In certain embodiments, k is

2. In certain embodiments, R^A is of the formula:







, or


 . In certain embodiments, R^A is of the formula:




,


 ,  ,  , or  . In certain embodiments, R^A is of

the formula:









, or  . In certain

embodiments, k is 3. In certain embodiments, R^A is of the formula:





,

 , or  . In certain embodiments, R^A is of the formula:


 ,  ,  , or  . In certain embodiments, R^A


is of the formula:



or  . In certain embodiments, k is 4. In certain

embodiments, R^A is of the formula:



, or  . In certain embodiments, R^A is not substituted or unsubstituted pyridinyl. In certain


embodiments, R^A is not substituted or unsubstituted 2-pyridinyl. In certain embodiments, R^A is not substituted 2-pyridinyl. In certain embodiments, R^A is substituted or unsubstituted pyrimidinyl. In certain embodiments, R^A is substituted or unsubstituted pyrazinyl. In certain embodiments, R^A is substituted or unsubstituted triazinyl. In certain embodiments, R^A is substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^A is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur), In certain embodiments, R^A is not substituted or unsubstituted 3-pyrrolidinyl. In certain embodiments, R^A is substituted or unsubstituted tetrahydropyranyl. In certain embodiments, R^A is unsubstituted tetrahydropyranyl. In certain embodiments, R^A is piperidinyl. In certain embodiments, R^A is substituted or unsubstituted morpholinyl. In certain embodiments, R^A is substituted or unsubstituted piperazinyl.


[00168] Formula (II) includes Ring B. Ring B is described in the Detailed Description for Formula (IV) below.


[00169] Formula (II) includes substituents R^C , R^D , R^E , and R^F . Substituents R^C , R^D , R^E , and R^F are described in the Detailed Description for Formula (IV) below.

[00170] Formula (II) includes Ring A and one or more instances of substituent R^G . Ring A and substituent R^G are described in the Detailed Description for Formula (IV) below.


[00171] Formula (II) includes linker L that connects Ring A to substituent R^H . In certain embodiments, L is a substituted or unsubstituted C_{1-6} hydrocarbon chain. In certain embodiments, one or more chain atoms of the hydrocarbon chain of L are independently replaced with $-C(=O)-$, $-O-$, $-S-$, $-NR^b-$, $-N=$, or $=N-$. In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain.


[00172] In certain embodiments, L is of the formula: , wherein a is 0, 1, 2, 3, 4, 5, or 6. In certain embodiments, a is 0. In certain embodiments, L is a bond. In certain embodiments, a is 1. In certain embodiments, a is 2. In certain embodiments, a is 3. In certain embodiments, a is 4. In certain embodiments, a is 5. In certain embodiments, a is 6. In


certain embodiments, L is of the formula: ,



, wherein l^A indicates the point of attachment to Ring A, and l^R indicates the point of attachment to R^H .


[00173] In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one or more chain atoms of the hydrocarbon chain are independently replaced with $-O-$ or $-NR^b-$. In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $-O-$. In certain embodiments, L is of

the formula: , wherein l^A indicates the point of attachment to Ring A, and l^R indicates the point of attachment to R^A . In certain embodiments, L is of the formula:


. In certain embodiments, L is a substituted C_{1-6} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $-N-$. In certain embodiments, L is





of the formula: , wherein l^A indicates the point of attachment to Ring A, and l^R indicates the point of attachment to R^H . In certain embodiments, L is of the formula:


. In certain embodiments, L is of the formula: . In certain








embodiments, L is of the formula: . In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $-C(=O)-$. In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $-S-$. In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $-NR^b-$. In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $-N=$. In certain embodiments, L is an unsubstituted C_{1-3} hydrocarbon chain, wherein one chain atom of the hydrocarbon chain is replaced with $=N-$.

[00174] Formula (II) includes substituent R^H . In certain embodiments, R^H is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, methyl, ethyl, or propyl). In certain embodiments, R^H is methyl. In certain embodiments, R^H is substituted or unsubstituted methyl. In certain

embodiments, R^H is substituted methyl. In certain embodiments, R^H is . In certain embodiments, R^H is unsubstituted methyl. In certain embodiments, R^H is ethyl. In certain embodiments, R^H is propyl. In certain embodiments, R^H is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^H is substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl, or substituted or unsubstituted piperazinyl. In certain embodiments, R^H is of the

formula: , , , , or


, wherein R^1 is substituted or unsubstituted C_{1-6} alkyl or $-OR^{x1}$, wherein R is hydrogen, substituted or unsubstituted C_{1-6} alkyl or nitrogen protecting group; R^{x1} is hydrogen or substituted or unsubstituted C_{1-6} alkyl; and x is 0, 1, 2, or 3. In certain

embodiments, R^H is of the formula: , , , , , , or . In certain embodiments, R^H


is $-OH$. In certain embodiments, R^H is $-N(R^c)_2$. As generally described herein, R^H may include substituent R^c . In certain embodiments, R^c is hydrogen. In certain embodiments, R^c is substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, R^c is substituted or unsubstituted C_{1-3} alkyl. In certain embodiments, R^c is substituted or unsubstituted methyl. In certain embodiments, R^c is methyl. In certain embodiments, R^c is substituted or unsubstituted ethyl. In certain embodiments, R^c is substituted or unsubstituted methyl. In certain embodiments, R^c is a nitrogen protecting group. In certain embodiments, R^H is $-NMe_2$. In certain embodiments, two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic ring (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heteroaryl ring (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system

are independently nitrogen, oxygen, or sulfur; or substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur).


[00175] In certain embodiments, Ring A with linker L and substituent R^H is of the




In certain embodiments, Ring A with linker L and substituent R^H is of the formula:




and substituent R^H is of the formula:




In certain embodiments, Ring A with linker L and substituent R^H is not of the formula:




L and substituent R^H is of the formula:






. In certain embodiments,


Ring A with linker L and substituent R^H is not of the formula:



[00176] In certain embodiments, Ring A with linker L and substituent R^H is of the




[00177] In certain embodiments, the compound of Formula (II) is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00178] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00179] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00180] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00181] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00182] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:






or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00183] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00184] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:





or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00185] In certain embodiments, the compound of Formula (II) useful in the present invention is of the formula:











or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00186] In certain embodiments, the compounds of Formula (II) useful in the present invention include, but are not limited to:




(HG-11-139-01),



(HG-11-143-01),




(YKL-04-136-10),




(YKL-04-136-11),

and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[00187] In certain embodiments, the compounds of Formula (II) useful in the present invention include, but are not limited to:




(HG-11-139-02),



(HG-11-143-01),


and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[00188] In certain embodiments, the compounds of Formula (II) useful in the present invention include, but are not limited to:




and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.


[00189] In certain embodiments, the compounds of Formula (II) useful in the present invention include, but are not limited to:




(YKL-05-57)




(YKL-05-58)




(YKL-05-59)




(YKL-05-60)




(YKL-05-68)




(YKL-05-69)




(YKL-05-70)




(YKL-05-74)




(YKL-05-76)




(YKL-05-77)




(YKL-05-88)




(YKL-05-89)




(YKL-05-90)




(YKL-05-91)




(YKL-05-92)




(YKL-05-93)




(YKL-05-94)




(YKL-05-95)




(YKL-05-96)




(YKL-05-97)




(YKL-05-98)




(YKL-05-99)




(YKL-05-100)




(YKL-05-151)




(YKL-05-152)




(YKL-05-153)




(YKL-05-154)




(YKL-05-155)




(YKL-05-156)




(YKL-05-163)




(YKL-05-164)




(YKL-05-165)




(YKL-05-166)




(YKL-05-178)




(YKL-05-179)




(YKL-05-180)




(YKL-05-181)




(YKL-05-182)




(YKL-05-183)




(Example 2)



(YKL-04-136-1; SB1-D-01)




(SB1-D-09; YKL-04-136-6)




(SB1-D-10)

(YKL-04-136-10)




(SB1-D-11)

(YKL-04-136-8)




(YKL-06-038)

(SB1-D-40)




(YKL-06-039)


(SB1-D-42)




(YKL-06-040)




(YKL-06-044)




(YKL-06-045)




(YKL-06-051)




(YKL-06-054)



(YKL-06-055)




(YKL-06-056)



(YKL-06-057)

(SB1-D-43)




(YKL-06-077)


(YKL-06-078)

(SB1-D-57)

(SB1-D-58)




(YKL-06-080-1)



(YKL-06-081-1)

(SB1-D-60)

(SB1-D-61)




(YKL-06-082)

(SB1-D-62)


and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof.

[00190] In certain embodiments, the compound of Formula (II) useful in the present invention is not of the formula:



(YKL-05-95)


(YKL-05-96)




(YKL-05-99),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00191] In certain embodiments, the compound of Formula (II) useful in the present invention is not of the formula:



(HG-11-139-02),




(HG-11-143-01),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

Compounds of Formula (III)

[00192] In another aspect, the compound utilized in the present disclosure is of Formula (III):



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:

R^J is substituted or unsubstituted carbocyclyl;

each instance of R^B is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or

unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

each instance of R^b is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^b are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

m is 0, 1, 2, 3, 4, or 5;

R^C is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^D is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^E is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^F is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Ring A is substituted or unsubstituted phenyl; substituted or unsubstituted, polycyclic aryl; substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl; or substituted or unsubstituted, polycyclic heteroaryl;

each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;


n is 0, 1, 2, 3, or 4, as valency permits;



R^K is unsubstituted methyl, substituted or unsubstituted heterocyclyl, $-OR^a$, or $-N(R^c)_2$, wherein each instance of R^c is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.




[00193] Formula (III) includes substituent R^J . In certain embodiments, R^J is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclyl comprising zero, one, or two double bonds in the carbocyclic ring system). In certain embodiments, R^J is substituted or unsubstituted, C_{3-6} carbocyclyl. In certain embodiments, R^J is substituted or unsubstituted cyclopropyl. In certain embodiments, R^J is


substituted or unsubstituted cyclobutyl. In certain embodiments, R^J is cyclobutyl. In certain embodiments, R^J is substituted or unsubstituted cyclopentyl. In certain embodiments, R^J is cyclopentyl. In certain embodiments, R^J is substituted or unsubstituted cyclohexyl. In certain embodiments, R^J is cyclohexyl.


[00194] As generally defined herein, as applicable to Formula (II) and (III), Ring B is an unsubstituted phenyl ring (*e.g.*, when m is 0) or a phenyl ring substituted with one or more substituents R^B (*e.g.*, when m is 1, 2, 3, 4, or 5). In certain embodiments, at least two instances of R^B are different. In certain embodiments, all instances of R^B are the same. In certain embodiments, m is 0. In certain embodiments, m is 1. In certain embodiments, m is 2. In certain embodiments, m is 3. In certain embodiments, m is 4. In certain embodiments, m is



5. In certain embodiments, Ring B is of the formula: . In certain embodiments, Ring

B is of the formula: . In certain embodiments, Ring B is of the formula: .

In certain embodiments, Ring B is of the formula: , , or . In

certain embodiments, Ring B is not of the formula: . In certain embodiments, Ring

B is of the formula: . In certain embodiments, Ring B is of the formula:

. In certain embodiments, Ring B is of the formula: .

[00195] In certain embodiments, at least one instance of R^B is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^B is F. In certain embodiments, at least one instance of R^B is Cl. In certain embodiments, at least one instance of R^B is Br. In certain embodiments, at least one instance of R^B is I. In certain embodiments, at least one R^B is


substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C₁₋₆ alkyl). In certain embodiments, at least one instance of R^B is substituted or unsubstituted methyl. In certain embodiments, at least one instance of R^B is methyl. In certain embodiments, m is 2, and both instances of R^B are methyl. In certain embodiments, m is 2, and one instance of R^B is halogen, and the other instance of R^B is methyl. In certain embodiments, m is 2, and one instance of R^B is Cl, and the other instance of R^B is methyl. In certain embodiments, at least one instance of R^B is substituted or unsubstituted ethyl. In certain embodiments, at least one instance of R^B is substituted or unsubstituted propyl. In certain embodiments, at least one instance of R^B is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C₂₋₆ alkenyl). In certain embodiments, at least one instance of R^B is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C₂₋₆ alkynyl). In certain embodiments, at least one instance of R^B is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclyl comprising zero, one, or two double bonds in the carbocyclic ring system). In certain embodiments, at least one instance of R^B is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^B is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of R^B is benzyl. In certain embodiments, at least one instance of R^B is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^B is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur; or substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^B is -OR^a (*e.g.*, -OH or -OMe). In certain embodiments, at least one instance of R^B is -N(R^b)₂, -SR^a, -CN, -SCN, -C(=NR^b)R^a, -C(=NR^b)OR^a, -C(=NR^b)N(R^b)₂, -C(=O)R^a, -C(=O)OR^a, -C(=O)N(R^b)₂, -NO₂, -NR^bC(=O)R^a, -NR^bC(=O)OR^a, -NR^bC(=O)N(R^b)₂, -OC(=O)R^a, -OC(=O)OR^a, or -OC(=O)N(R^b)₂.

[00196] Formula (III) includes substituents R^C, R^D, R^E, and R^F. Substituents R^C, R^D, R^E, and R^F are described in the Detailed Description for Formula (IV) below.


[00197] Formula (III) includes Ring A and one or more instances of substituent R^G. Ring A and substituent R^G are described in the Detailed Description for Formula (IV) below.

[00198] As generally defined herein, Formula (III) includes substituent R^K attached to Ring A. Substituent R^K is described in the Detailed Description for Formula (IV) below.

[00199] In certain embodiments, Ring A with substituent R^K is of the formula:




[00200] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00201] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00202] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:






or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00203] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00204] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00205] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00206] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00207] In certain embodiments, the compound of Formula (III) useful in the present invention is of the formula:




(YKL-06-063)




(YKL-06-064)




(YKL-06-075)




(YKL-06-076)




(YKL-06-088)



(YKL-06-089)



(YKL-06-090)




(YKL-06-091)

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

Compounds of Formula (IV)

[00208] In another aspect, the compound utilized in the present disclosure is of Formula (IV):



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:

R^L is substituted or unsubstituted alkyl;

Ring C is unsubstituted phenyl or of the formula: or ;

each instance of R^{B1} is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^d)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^d)R^a$, $-C(=NR^d)OR^a$, $-C(=NR^d)N(R^d)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^d)_2$, $-NO_2$, $-NR^dC(=O)R^a$, $-NR^dC(=O)OR^a$, $-NR^dC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^d)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

each instance of R^d is independently hydrogen, $-C(=O)R^a$, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^d are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

R^C is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^D is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^E is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^F is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Ring A is substituted or unsubstituted phenyl; substituted or unsubstituted, polycyclic aryl; substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl; or substituted or unsubstituted, polycyclic heteroaryl;


each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;


n is 0, 1, 2, 3, or 4, as valency permits;

R^K is unsubstituted methyl, substituted or unsubstituted heterocyclyl, $-OR^a$, or $-N(R^c)_2$, wherein each instance of R^c is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

R^Y is substituted phenyl.


[00209] Formula (IV) includes Ring C. In certain embodiments, Ring C is unsubstituted




phenyl. In certain embodiments, Ring C is of the formula: . In certain embodiments, at least one instance of R^{B1} is halogen. In certain embodiments, at least one instance of R^{B1} is halogen. In certain embodiments, at least one instance of R^{B1} is F. In certain embodiments, at least one instance of R^{B1} is Cl. In certain embodiments, at least one instance of R^{B1} is Br. In certain embodiments, at least one instance of R^{B1} is I (iodine). In certain embodiments, at least one instance of R^{B1} is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, methyl, ethyl, or propyl). In certain embodiments, at least one instance of R^{B1} is substituted or unsubstituted methyl. In certain embodiments, at least one instance of R^{B1} is methyl. In certain embodiments, at least one instance of R^{B1} is $-N(R^d)_2$, wherein each instance of R^d is hydrogen or $-C(=O)R^a$. In certain embodiments, at least one instance of R^{B1} is $-NH(C(=O)R^a)$. In certain embodiments, at least one instance of R^{B1} is $-OR^a$ (*e.g.*, $-OH$ or


–OMe). In certain embodiments, at least one instance of R^{B1} is $-SR^a$, $-CN$, $-SCN$, $-C(=NR^d)R^a$, $-C(=NR^d)OR^a$, $-C(=NR^d)N(R^d)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^d)_2$, $-NO_2$, $-NR^dC(=O)R^a$, $-NR^dC(=O)OR^a$, $-NR^dC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or –


$OC(=O)N(R^d)_2$. In certain embodiments, Ring C is of the formula:



embodiments, Ring C is of the formula:




formula: . In certain embodiments, R^Y is substituted phenyl. In certain embodiments, R^Y is substituted phenyl. In certain embodiments, R^Y is of the formula:




, wherein R^{y1} is halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, R^{y1} is halogen (*e.g.*, Br, Cl, F). In certain embodiments, R^{y1} is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, substituted or unsubstituted, methyl, ethyl, or propyl). In certain embodiments, R^{y1} is substituted or unsubstituted methyl. In certain embodiments, R^{y1} is substituted methyl. In certain embodiments, R^{y1} is methyl. In certain embodiments, R^{y1} is –

CF_3 . In certain embodiments, Ring C is of the formula:



embodiments, Ring C is of the formula:




[00210] Formula (IV) includes substituent R^L . In certain embodiments, R^L is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, R^L is substituted or unsubstituted methyl. In certain embodiments, R^L is methyl. In certain

embodiments, R^L is unsubstituted methyl. In certain embodiments, R^L is substituted methyl. In certain embodiments, R^L is substituted or unsubstituted ethyl. In certain embodiments, R^L is ethyl. In certain embodiments, R^L is unsubstituted ethyl. In certain embodiments, R^L is substituted ethyl. In certain embodiments, R^L is substituted or unsubstituted propyl. In certain embodiments, R^L is propyl. In certain embodiments, R^L is isopropyl. In certain embodiments, R^L is substituted or unsubstituted butyl.


[00211] As generally defined herein, Formulae (III) and (IV) include substituent R^K attached to Ring A. In certain embodiments, R^K is unsubstituted methyl. In certain embodiments, R^K is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^K is substituted or unsubstituted tetrahydropyranyl. In certain embodiments, R^K is substituted or unsubstituted piperidinyl. In certain embodiments, R^K is substituted or unsubstituted morpholinyl. In certain embodiments, R^K is substituted or unsubstituted

piperazinyl. In certain embodiments, R^K is of the formula:



wherein R^1 is substituted or unsubstituted C_{1-6} alkyl or $-OR^{x1}$, wherein R is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or nitrogen protecting group; R^{x1} is hydrogen or substituted or unsubstituted C_{1-6} alkyl;

and x is 0, 1, 2, or 3. In certain embodiments, R^K is of the formula:



. In certain embodiments, R^K is $-OR^a$ (*e.g.*, $-OH$ or $-OMe$). In certain embodiments, R^K is $-N(R^c)_2$. In certain embodiments, two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic ring (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heteroaryl ring (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur; or substituted or

unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^K is $-NMe_2$. In certain embodiments, R^K is $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$.

[00212] As generally defined herein, Formulae (II), (III), and (IV) include substituent R^C . In certain embodiments, R^C is hydrogen. In certain embodiments, R^C is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, R^C is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, methyl, ethyl, or propyl). In certain embodiments, R^C is substituted or unsubstituted methyl. In certain embodiments, R^C is methyl. In certain embodiments, R^C is substituted or unsubstituted ethyl. In certain embodiments, R^C is ethyl. In certain embodiments, R^C is substituted or unsubstituted propyl. In certain embodiments, R^C is unsubstituted isopropyl.

[00213] As generally defined herein, Formulae (II), (III), and (IV) include substituent R^D . In certain embodiments, R^D is hydrogen. In certain embodiments, R^D is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, R^D is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, methyl, ethyl, or propyl). In certain embodiments, R^D is substituted or unsubstituted methyl. In certain embodiments, R^D is methyl. In certain embodiments, R^D is substituted or unsubstituted ethyl. In certain embodiments, R^D is ethyl. In certain embodiments, R^D is substituted or unsubstituted propyl. In certain embodiments, R^D is isopropyl.

[00214] As generally defined herein, Formulae (II), (III), and (IV) include substituent R^E . In certain embodiments, R^E is hydrogen. In certain embodiments, R^E is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, R^E is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, methyl, ethyl, or propyl). In certain embodiments, R^E is substituted or unsubstituted methyl. In certain embodiments, R^E is methyl. In certain embodiments, R^E is substituted or unsubstituted ethyl. In certain embodiments, R^E is ethyl. In certain embodiments, R^E is substituted or unsubstituted propyl. In certain embodiments, R^E is isopropyl.

[00215] As generally defined herein, Formulae (II), (III), and (IV) include substituent R^F . In certain embodiments, R^F is hydrogen. In certain embodiments, R^F is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, methyl, ethyl, or propyl). In certain embodiments, R^F is substituted or unsubstituted methyl. In certain embodiments, R^F is methyl. In certain embodiments, R^F is substituted or unsubstituted ethyl. In certain embodiments, R^F is ethyl. In certain embodiments, R^F is substituted or unsubstituted propyl. In certain embodiments, R^F is isopropyl. In certain embodiments, R^F is a nitrogen protecting group (*e.g.*, a nitrogen


protecting group (*e.g.*, benzyl (Bn), t-butyl carbonate (BOC or Boc), benzyl carbamate (Cbz), 9-fluorenylmethyl carbonate (Fmoc), trifluoroacetyl, triphenylmethyl, acetyl, or *p*-toluenesulfonamide (Ts)).


[00216] In certain embodiments, R^C , R^D , R^E , and R^F are each hydrogen. In certain embodiments, at least one substituent selected from the group consisting of R^C , R^D , R^E , and R^F is substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, R^C is substituted or unsubstituted C_{1-6} alkyl; and R^D , R^E , and R^F are each hydrogen. In certain embodiments, R^C is unsubstituted methyl; and R^D , R^E , and R^F are each hydrogen. In certain embodiments, R^C is unsubstituted isopropyl; and R^D , R^E , and R^F are each hydrogen. In certain embodiments, R^D is substituted or unsubstituted C_{1-6} alkyl; and R^C , R^E , and R^F are each hydrogen. In certain embodiments, R^E is substituted or unsubstituted C_{1-6} alkyl; and R^C , R^D , and R^F are each hydrogen. In certain embodiments, R^F is substituted or unsubstituted C_{1-6} alkyl; and R^C , R^D , and R^E are each hydrogen.


[00217] As generally defined herein, Formulae (II), (III), and (IV) include Ring A. In certain embodiments, Ring A is substituted or unsubstituted phenyl. In certain embodiments, Ring A is not substituted or unsubstituted phenyl. In certain embodiments, Ring A is not substituted phenyl. In certain embodiments, Ring A is not unsubstituted phenyl. In certain embodiments, Ring A is unsubstituted phenyl. In certain embodiments, Ring A is phenyl, and includes one or more R^G substituents. In certain embodiments, Ring A includes one R^G substituent. In certain embodiments, Ring A includes two R^G substituents. In certain embodiments, Ring A is substituted or unsubstituted polycyclic aryl (*e.g.*, naphthalene or anthracene). In certain embodiments, Ring A is substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, Ring A is substituted or unsubstituted furan. In certain embodiments, Ring A is substituted or unsubstituted thiophene. In certain embodiments, Ring A is substituted or unsubstituted pyrrole. In certain embodiments, Ring A is substituted or unsubstituted pyrazole. In certain embodiments, Ring A is pyrazole. In certain embodiments, Ring A is substituted or unsubstituted pyridinyl. In certain embodiments, Ring A is pyridinyl. In certain embodiments, Ring A is substituted or unsubstituted polycyclic heteroaryl (*e.g.*, substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur).

[00218] As generally defined herein, Formulae (II), (III), and (IV) include one or more

instances of substituent R^G . In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2. In certain embodiments, n is 3. In certain embodiments, n is 4. In certain embodiments, at least one instance of R^G is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^G is F. In certain embodiments, at least one instance of R^G is Cl. In certain embodiments, at least one instance of R^G is Br. In certain embodiments, at least one instance of R^G is I. In certain embodiments, at least one R^G is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^G is substituted or unsubstituted C_{1-3} alkyl. In certain embodiments, at least one instance of R^G is substituted or unsubstituted methyl. In certain embodiments, at least one instance of R^G is unsubstituted methyl. In certain embodiments, at least one instance of R^G is substituted methyl. In certain embodiments, at least one instance of R^G is $-CF_3$. In certain embodiments, at least one instance of R^G is substituted or unsubstituted ethyl. In certain embodiments, at least one instance of R^G is substituted ethyl. In certain embodiments, at least one instance of R^G is unsubstituted ethyl. In certain embodiments, at least one instance of R^G is substituted or unsubstituted propyl. In certain embodiments, at least one instance of R^G is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{2-6} alkenyl). In certain embodiments, at least one instance of R^G is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C_{2-6} alkynyl). In certain embodiments, at least one instance of R^G is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclyl comprising zero, one, or two double bonds in the carbocyclic ring system). In certain embodiments, at least one instance of R^G is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, 5- to 10-membered monocyclic or bicyclic heterocyclic ring, wherein one or two atoms in the heterocyclic ring are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^G is substituted or unsubstituted morpholinyl. In

certain embodiments, at least one instance of R^G is of the formula: . In certain embodiments, at least one instance of R^G is substituted or unsubstituted piperazinyl. In


certain embodiments, at least one instance of R^G is of the formula: . In certain

embodiments, at least one instance of R^G is of the formula: . In certain

embodiments, at least one instance of R^G is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of


R^G is benzyl. In certain embodiments, at least one instance of R^G is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^B is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur; or substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^G is $-OR^a$, wherein R^a is hydrogen or substituted or unsubstituted C_{1-6} alkyl (*e.g.*, $-OH$ or $-OMe$). In certain embodiments, at least one instance of R^G is $-OMe$. In certain embodiments, at least one instance of R^G is $-OEt$. In certain embodiments, at least one instance of R^G is $-O(Pr)$. In certain embodiments, at least one instance of R^G is $-O(iPr)$. In certain embodiments, at least one instance of R^G is $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$.

[00219] In certain embodiments, Ring A with substituent R^K is of the formula:




. In certain embodiments, Ring A with

substituent R^K is of the formula:




[00220] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00221] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00222] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00223] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00224] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00225] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00226] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00227] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:





or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00228] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:



(HG-11-23-01),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00229] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




(HG-11-139-02),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00230] In certain embodiments, the compound of Formula (IV) useful in the present invention is of the formula:




(YKL-06-084)




(YKL-06-085)



(YKL-06-086)



(YKL-06-087)




(HG-4-34-01),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.


[00231] In certain embodiments, the compound of Formula (IV) useful in the present

invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00232] In certain embodiments, the compound of Formula (IV) useful in the present invention is not of the formula:




(HG-11-139-02),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

Compounds of Formula (V)


[00233] In another aspect, the compound utilized in the present disclosure is of Formula (V):



(V),

and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:

R^G is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heteroaryl, or of the formula:



each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or

unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$, or two R^a groups are joined to form a substituted or unsubstituted carbocyclic ring, substituted or unsubstituted heterocyclic ring, substituted or unsubstituted aryl ring, or substituted or unsubstituted heteroaryl ring;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^a groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, 4, or 5;

R^b is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each of X^A , X^B , and X^C is independently N or CR^X , wherein each instance of R^X is independently hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;

or: X^B is CR^X , and R^G and R^X of X^B are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

R^C is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;

$\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;

R^D is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or a nitrogen protecting group;


Y is $-\text{O}-$ or $-\text{NR}^Y-$, wherein R^Y is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Z is a bond or $-\text{C}(\text{R}^Z)_2-$, wherein each instance of R^Z is independently hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

each instance of R^E is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NR}^a\text{S}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{S}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{S}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NR}^a\text{S}(=\text{O})_2\text{R}^a$, $-\text{NR}^a\text{S}(=\text{O})_2\text{OR}^a$, $-\text{NR}^a\text{S}(=\text{O})_2\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$; and

m is 0, 1, 2, 3, 4, or 5.

[00234] In certain embodiments, a compound of Formula (V) useful in the present invention is of Formula (V-A):



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof, wherein:

each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NR}^a\text{S}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{S}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{S}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NR}^a\text{S}(=\text{O})_2\text{R}^a$, $-\text{NR}^a\text{S}(=\text{O})_2\text{OR}^a$, $-\text{NR}^a\text{S}(=\text{O})_2\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;

NO_2 , $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^a groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, 4, or 5;

R^b is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each of X^A , X^B , and X^C is independently N or CR^X , wherein each instance of R^X is independently hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;

R^c is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^a$, $-\text{N}(\text{R}^a)_2$, $-\text{SR}^a$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^a)\text{R}^a$, $-\text{C}(=\text{NR}^a)\text{OR}^a$, $-\text{C}(=\text{NR}^a)\text{N}(\text{R}^a)_2$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^a$, $-\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{NO}_2$, $-\text{NR}^a\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^a\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^a)_2$;


R^d is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or a nitrogen protecting group;


Y is $-\text{O}-$ or $-\text{NR}^Y-$, wherein R^Y is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;


each instance of R^E is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^a)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$; and

m is 0, 1, 2, 3, 4, or 5.

[00235] Formula (V) includes substituent R^G . In certain embodiments, R^G is hydrogen. In certain embodiments, R^G is substituted or unsubstituted alkyl. In certain embodiments, R^G is substituted C_{1-6} alkyl (*e.g.*, $-CF_3$, perfluoroethyl, perfluoropropyl, perfluorobutyl, Bn, or C_{1-6} alkyl substituted with at least one instance of halogen and/or $-OR^a$). In certain embodiments, R^G is C_{1-6} alkyl substituted with at least one instance of $-OR^a$, optionally wherein R^a is hydrogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, R^G is of the


formula: . In certain embodiments, R^G is unsubstituted C_{1-6} alkyl (*e.g.*, Me, Et, Pr, or Bu). In certain embodiments, R^G is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{1-6} alkenyl). In certain embodiments, R^G is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C_{1-6} alkynyl). In certain embodiments, R^G is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3- to 7-membered carbocyclyl). In certain embodiments, R^G is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^G is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^G is substituted or unsubstituted 2-pyridyl. In certain embodiments, R^G is substituted or unsubstituted 3-pyridyl.





In certain embodiments, R^G is of the formula: , wherein R^a is hydrogen, halogen, substituted or unsubstituted C_{1-6} alkyl, $-OH$, $-O$ (substituted or unsubstituted C_{1-6} alkyl), or substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein

one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur; and each instance of R^b is independently hydrogen, halogen, substituted or unsubstituted C_{1-6} alkyl, $-OH$, or $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain


embodiments, R^G is of the formula:



, , or






. In certain embodiments, R^G is substituted or unsubstituted 4-pyridyl. In certain embodiments, R^G is substituted or unsubstituted 1-pyrazolyl. In certain embodiments, R^G is substituted or unsubstituted 3-pyrazolyl. In certain embodiments, R^G is substituted or unsubstituted 4-pyrazolyl. In certain embodiments, R^G is of the formula:




, wherein R^a is hydrogen, substituted or unsubstituted C_{1-6} alkyl, a nitrogen protecting group, or $-(\text{substituted or unsubstituted } C_{1-6} \text{ alkylene})-(\text{substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur})$; and each instance of R^b is independently hydrogen, halogen, substituted or unsubstituted C_{1-6} alkyl, $-OH$, or $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, R^G is of the


formula:











, , or . In certain embodiments,


R^G is substituted or unsubstituted furanyl, substituted or unsubstituted thienyl, substituted or unsubstituted pyrrolyl, substituted or unsubstituted imidazolyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted isoxazolyl, substituted or unsubstituted thiazolyl, substituted or unsubstituted isothiazolyl, or substituted or unsubstituted tetrazolyl. In certain embodiments, R^G is substituted or unsubstituted pyrazinyl, substituted or unsubstituted pyrimidinyl, or substituted or unsubstituted pyridazinyl. In certain embodiments, R^G is substituted or unsubstituted, bicyclic, 9- to 10-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.


[00236] In certain embodiments, R^G is of the formula: . Ring A is unsubstituted (*e.g.*, when k is 0) or substituted with one or more substituents R^A (*e.g.*, when k is 1, 2, 3, 4, or 5). In certain embodiments, Ring A is an unsubstituted phenyl ring. In certain embodiments, Ring A is a substituted phenyl ring. In certain embodiments, Ring A is of the


formula: , optionally wherein R^A is substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur. In certain embodiments, Ring A is of the formula:


, *e.g.*,  or . In certain embodiments, R^G is . In certain embodiments, Ring A is of the formula: , , or


, In certain embodiments, Ring A is of the formula: , optionally wherein R^A is halogen, substituted or unsubstituted C_{1-6} alkyl, or $-OR^a$. In certain

embodiments, Ring A is of the formula: . In certain embodiments,


Ring A is of the formula: . In certain embodiments, Ring A is of the formula:

, optionally wherein R^A is halogen, substituted or unsubstituted C_{1-6} alkyl, or $-OR^a$.


In certain embodiments, Ring A is of the formula: . In certain





embodiments, Ring A is of the formula: , optionally wherein each instance of R^A is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, or substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen,



oxygen, or sulfur. In certain embodiments, Ring A is of the formula: ,




certain embodiments, Ring A is of the formula:




. In certain embodiments, Ring A is of the formula: , optionally wherein each instance of R^A is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, or substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen, oxygen,


or sulfur. In certain embodiments, Ring A is of the formula: or




. In certain embodiments, Ring A is of the formula:




, wherein the two instances of R^A are joined to form a substituted or unsubstituted carbocyclic ring (*e.g.*, substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclic ring), substituted or unsubstituted heterocyclic ring (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur), substituted or unsubstituted aryl ring (*e.g.*, substituted or unsubstituted phenyl ring), or substituted or unsubstituted heteroaryl ring (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl ring, wherein one or two atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In




certain embodiments, Ring A is of the formula:




of the formula: , optionally wherein each instance of R^A is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, or substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur. In certain



embodiments, Ring A is of the formula:




Ring A is of the formula: , wherein each instance of R^A is independently $-OR^a$.





In certain embodiments, Ring A is of the formula: , wherein each instance of R^A


is independently –O(substituted or unsubstituted alkyl). In certain embodiments, Ring A is of




[00237] In Formula (V), Ring A may include one or more substituents R^A . In certain embodiments, all instances of R^A are the same. In certain embodiments, at least two instances of R^A are different. In certain embodiments, at least one instance of R^A is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^A is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^A is $-CH_3$. In certain embodiments, at least one instance of R^A is $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn. In certain embodiments, at least one instance of R^A is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{1-6} alkenyl). In certain embodiments, at least one instance of R^A is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C_{1-6} alkynyl). In certain embodiments, at least one instance of R^A is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3- to 7-membered carbocyclyl). In certain embodiments, at least one instance of R^A is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance



unsubstituted phenyl. In certain embodiments, at least one instance of R^A is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^A is $-OR^a$. In certain embodiments, at least one instance of R^A is $-OH$. In certain embodiments, at least one instance of R^A is $-O$ (substituted or unsubstituted alkyl), such as $-O$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-OMe$, $-OEt$, $-OPr$, $-OBu$, $-OBn$, or



). In certain embodiments, at least one instance of R^A is $-O$ (substituted or unsubstituted phenyl) (*e.g.*, $-OPh$). In certain embodiments, at least one instance of R^A is $-SR^a$ (*e.g.*, $-SH$, $-S$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-SMe$, $-SEt$, $-SPr$, $-SBu$, or $-SBn$), or $-S$ (substituted or unsubstituted phenyl) (*e.g.*, $-SPh$). In certain embodiments, at least one instance of R^A is $-N(R^a)_2$ (*e.g.*, $-NH_2$, $-NH$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-NHMe$), or $-N$ (substituted or unsubstituted C_{1-6} alkyl)–(substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-NMe_2$). In certain embodiments, at least one instance of R^A is $-CN$, $-SCN$, or $-NO_2$. In certain embodiments, at least one instance of R^A is $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, or $-C(=NR^a)N(R^a)_2$. In certain embodiments, at least one instance of R^A is $-C(=O)R^a$ or $-C(=O)OR^a$. In certain embodiments, at least one instance of R^A is $-C(=O)N(R^a)_2$ (*e.g.*, $-$



$C(=O)NH_2$, $-C(=O)NHMe$, $-C(=O)NMe_2$, or). In certain

embodiments, at least one instance of R^A is $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, or $-NR^aC(=O)N(R^a)_2$. In certain embodiments, at least one instance of R^A is $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$.

[00238] Each instance of R^A , R^C , R^E , and R^X may independently include one or more substituents R^a . In certain embodiments, all instances of R^a are the same. In certain embodiments, at least two instances of R^a are different. In certain embodiments, at least one instance of R^a is H. In certain embodiments, each instance of R^a is H. In certain embodiments, at least one instance of R^a is substituted or unsubstituted acyl (*e.g.*, acetyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^a is $-CH_3$. In certain embodiments, at least one instance of R^a is $-CF_3$, unsubstituted ethyl,

perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn. In certain embodiments, at least one instance of R^a is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{1-6} alkenyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C_{1-6} alkynyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3- to 7-membered carbocyclyl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^a is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of R^a is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^a is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^a is a nitrogen protecting group (*e.g.*, Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts) when attached to a nitrogen atom. In certain embodiments, at least one instance of R^a is an oxygen protecting group (*e.g.*, silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl) when attached to an oxygen atom. In certain embodiments, at least one instance of R^a is a sulfur protecting group (*e.g.*, acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl) when attached to a sulfur atom. In certain embodiments, two instances of R^a are joined to form a substituted or unsubstituted heterocyclic ring (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, two instances of R^a are joined to form a substituted or unsubstituted heteroaryl ring (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur).

[00239] In certain embodiments, k is 0. In certain embodiments, k is 1. In certain embodiments, k is 2. In certain embodiments, k is 3. In certain embodiments, k is 4. In certain embodiments, k is 5.

[00240] Formula (V) includes substituent R^B on the nitrogen atom that connects Rings A and B. In certain embodiments, R^B is hydrogen. In certain embodiments, R^B is substituted or

unsubstituted acyl (*e.g.*, acetyl). In certain embodiments, R^B is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, $-CH_3$, $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn). In certain embodiments, R^B is a nitrogen protecting group (*e.g.*, Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts).

[00241] Formula (V) includes a heteroaryl ring as Ring B that includes moieties X^A , X^B , and X^C in the heteroaryl ring system. In certain embodiments, X^A is CR^X , and each of X^B and X^C is N. In certain embodiments, X^A is CH, and each of X^B and X^C is N. In certain embodiments, X^B is CR^X , and each of X^A and X^C is N. In certain embodiments, X^B is CH, and each of X^A and X^C is N. In certain embodiments, X^C is CR^X , and each of X^A and X^B is N. In certain embodiments, X^C is CH, and each of X^A and X^B is N. In certain embodiments, X^A is N, and each of X^B and X^C is independently CR^X . In certain embodiments, X^A is N, and each of X^B and X^C is CH. In certain embodiments, X^B is N, and each of X^A and X^C is independently CR^X . In certain embodiments, X^B is N, and each of X^A and X^C is CH. In certain embodiments, X^C is N, and each of X^A and X^B is independently CR^X . In certain embodiments, X^C is N, and each of X^A and X^B is CH. In certain embodiments, each of X^A , X^B , and X^C is independently CR^X . In certain embodiments, each of X^A , X^B , and X^C is CH.

[00242] In certain embodiments, X^B is CR^X , and R^G and R^X of X^B are joined to form a substituted or unsubstituted heterocyclic ring (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclic ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur, further wherein at least one atom in the heterocyclic ring system is nitrogen). In certain embodiments, X^B is CR^X , and R^G and R^X of X^B are joined to form a substituted or unsubstituted heteroaryl ring (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl ring, wherein one or two atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur, further wherein at least one atom in the heteroaryl ring system is nitrogen). In certain embodiments, X^B is CR^X , and R^G and R^X of X^B are joined to form substituted or unsubstituted pyrrolyl ring.

[00243] In certain embodiments, all instances of R^X are the same. In certain embodiments, at least two instances of R^X are different. In certain embodiments, at least one instance of R^X is hydrogen. In certain embodiments, at least one instance of R^X is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^X is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^X is $-CH_3$. In certain embodiments, at least one instance of R^X is $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or


Bn. In certain embodiments, at least one instance of R^X is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{1-6} alkenyl). In certain embodiments, at least one instance of R^X is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C_{1-6} alkynyl). In certain embodiments, at least one instance of R^X is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3- to 7-membered carbocyclyl). In certain embodiments, at least one instance of R^X is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^X is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of R^X is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^X is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^X is $-OR^a$ (*e.g.*, $-OH$, $-O$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-OMe$, $-OEt$, $-OPr$, $-OBu$, or $-OBn$), or $-O$ (substituted or unsubstituted phenyl) (*e.g.*, $-OPh$)). In certain embodiments, at least one instance of R^X is $-SR^a$ (*e.g.*, $-SH$, $-S$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-SMe$, $-SEt$, $-SPr$, $-SBu$, or $-SBn$), or $-S$ (substituted or unsubstituted phenyl) (*e.g.*, $-SPh$)). In certain embodiments, at least one instance of R^X is $-N(R^a)_2$ (*e.g.*, $-NH_2$, $-NH$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-NHMe$), or $-N$ (substituted or unsubstituted C_{1-6} alkyl)–(substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-NMe_2$)). In certain embodiments, at least one instance of R^X is $-CN$, $-SCN$, or $-NO_2$. In certain embodiments, at least one instance of R^X is $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, or $-C(=NR^a)N(R^a)_2$. In certain embodiments, at least one instance of R^X is $-C(=O)R^a$, $-C(=O)OR^a$, or $-C(=O)N(R^a)_2$ (*e.g.*, $-C(=O)NH_2$, $-C(=O)NHMe$, or $-C(=O)NMe_2$). In certain embodiments, at least one instance of R^X is $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, or $-NR^aC(=O)N(R^a)_2$. In certain embodiments, at least one instance of R^X is $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$.

[00244] Formula (V) includes substituent R^C on Ring B. In certain embodiments, R^C is hydrogen. In certain embodiments, R^C is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, R^C is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, R^C is $-CH_3$. In certain embodiments, R^C is $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn. In certain embodiments, R^C is substituted or unsubstituted


alkenyl (*e.g.*, substituted or unsubstituted C₁₋₆ alkenyl). In certain embodiments, R^C is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C₁₋₆ alkynyl). In certain embodiments, R^C is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3- to 7-membered carbocyclyl). In certain embodiments, R^C is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^C is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, R^C is substituted or unsubstituted phenyl. In certain embodiments, R^C is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^C is -OR^a (*e.g.*, -OH, -O(substituted or unsubstituted C₁₋₆ alkyl) (*e.g.*, -OMe, -OEt, -OPr, -OBu, or -OBn), or -O(substituted or unsubstituted phenyl) (*e.g.*, -OPh)). In certain embodiments, R^C is -SR^a (*e.g.*, -SH, -S(substituted or unsubstituted C₁₋₆ alkyl) (*e.g.*, -SMe, -SEt, -SPr, -SBu, or -SBn), or -S(substituted or unsubstituted phenyl) (*e.g.*, -SPh)). In certain embodiments, R^C is -N(R^a)₂ (*e.g.*, -NH₂, -NH(substituted or unsubstituted C₁₋₆ alkyl) (*e.g.*, -NHMe), or -N(substituted or unsubstituted C₁₋₆ alkyl)-(substituted or unsubstituted C₁₋₆ alkyl) (*e.g.*, -NMe₂)). In certain embodiments, R^C is -CN, -SCN, or -NO₂. In certain embodiments, R^C is -C(=NR^a)R^a, -C(=NR^a)OR^a, or -C(=NR^a)N(R^a)₂. In certain embodiments, R^C is -C(=O)R^a, -C(=O)OR^a, or -C(=O)N(R^a)₂ (*e.g.*, -C(=O)NH₂, -C(=O)NHMe, or -C(=O)NMe₂). In certain embodiments, R^C is -NR^aC(=O)R^a, -NR^aC(=O)OR^a, or -NR^aC(=O)N(R^a)₂. In certain embodiments, R^C is -OC(=O)R^a, -OC(=O)OR^a, or -OC(=O)N(R^a)₂.

[00245] Formula (V) includes substituent R^D on a nitrogen atom of the urea or carbamate moiety. In certain embodiments, R^D is hydrogen. In certain embodiments, R^D is substituted or unsubstituted alkyl, such as substituted or unsubstituted C₁₋₆ alkyl (*e.g.*, -CH₃, -CF₃, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn). In certain embodiments, R^D is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C₁₋₆ alkenyl). In certain embodiments, R^D is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C₁₋₆ alkynyl). In certain embodiments, R^D is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3- to 7-membered carbocyclyl). In certain embodiments, R^D is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system


are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^D is substituted or unsubstituted oxetanyl, substituted or unsubstituted azetidiny, substituted or unsubstituted tetrahydrofuranyl, substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted piperazinyl, or substituted or unsubstituted morpholinyl. In certain embodiments, R^D is of the formula:



. In certain embodiments, R^D is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, R^D is substituted or




unsubstituted phenyl. In certain embodiments, R^D is of the formula: $(R^F)_n$, wherein each instance of R^F is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^a)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$; and n is 0, 1, 2, 3, 4, or 5. In certain embodiments, R^D is of the formula:





(*e.g.*, R^F). In certain embodiments, R^D is of the


formula: (*e.g.*, R^F), or R^F .

In certain embodiments, R^D is of the formula: (*e.g.*, R^F), or R^F (*e.g.*, OR^a).



In certain embodiments, R^D is of the formula . In certain embodiments, R^D is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, R^D is substituted or unsubstituted 1-pyrazolyl, substituted or unsubstituted 3-pyrazolyl, or substituted or



unsubstituted 4-pyrazolyl (*e.g.*, ). In certain embodiments, R^D is substituted or unsubstituted furanyl, substituted or unsubstituted thienyl, substituted or unsubstituted pyrrolyl, substituted or unsubstituted imidazolyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted isoxazolyl, substituted or unsubstituted thiazolyl, substituted or unsubstituted isothiazolyl, or substituted or unsubstituted tetrazolyl. In certain embodiments, R^D is substituted or unsubstituted pyridyl, substituted or unsubstituted pyrazinyl, substituted or unsubstituted pyrimidinyl, or substituted or unsubstituted pyridazinyl.

[00246] In certain embodiments, R^D is a nitrogen protecting group (*e.g.*, Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts).

[00247] Formula (V) includes divalent moiety Y. In certain embodiments, Y is $-O-$. In certain embodiments, Y is $-NR^Y-$. In certain embodiments, Y is $-NH-$.

[00248] In certain embodiments, R^Y is hydrogen. In certain embodiments, R^Y is substituted or unsubstituted acyl (*e.g.*, acetyl). In certain embodiments, R^Y is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, $-CH_3$, $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn). In certain embodiments, R^Y is a nitrogen protecting group (*e.g.*, Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts).


[00249] Formula (V) includes divalent moiety Z. In certain embodiments, Z is a bond. In certain embodiments, Z is $-C(R^Z)_2-$. In certain embodiments, Z is $-CH_2-$. In certain embodiments, Z is $-CHF-$ or $-CF_2-$.


[00250] In certain embodiments, the two instances of R^Z are the same. In certain embodiments, the two instances of R^Z are not the same. In certain embodiments, at least one instance of R^Z is hydrogen. In certain embodiments, each instance of R^Z is hydrogen. In


certain embodiments, at least one instance of R^Z is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^Z is substituted or unsubstituted C_{1-6} alkyl (*e.g.*, $-CH_3$, $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn).



[00251] In certain embodiments, $-Y-Z-$ is $-N(R^Y)-$. In certain embodiments, $-Y-Z-$ is $-NH-$. In certain embodiments, $-Y-Z-$ is $-N(Me)-$. In certain embodiments, $-Y-Z-$ is $-O-$. In certain embodiments, $-Y-Z-$ is $-N(R^Y)-C(R^Z)_2-$ (*e.g.*, $-N(R^Y)-CH_2-$). In certain embodiments, $-Y-Z-$ is $-NH-CH_2-$. In certain embodiments, $-Y-Z-$ is $-N(Me)-CH_2-$. In certain embodiments, $-Y-Z-$ is $-O-C(R^Z)_2-$ (*e.g.*, $-O-CH_2-$).


[00252] Formula (V) includes a phenyl ring as Ring C, which is unsubstituted (*e.g.*, when m is 0) or substituted with one or more substituents R^E (*e.g.*, when m is 1, 2, 3, 4, or 5). In certain embodiments, Ring C is an unsubstituted phenyl ring. In certain embodiments, Ring C





is a substituted phenyl ring. In certain embodiments, Ring C is of the formula: , optionally wherein R^E is halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-SCN$, or $-NO_2$. In





certain embodiments, Ring C is of the formula:  or . In certain








embodiments, Ring C is of the formula: , optionally wherein R^E is halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-SCN$, or $-NO_2$. In certain embodiments,





Ring C is of the formula:  F. In certain embodiments, Ring C is of the formula: , optionally wherein R^E is halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-$


$C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-SCN$, or $-NO_2$. In




certain embodiments, Ring C is of the formula:  or . In certain embodiments, Ring


C is of the formula: , , or . In certain embodiments, Ring C is of

the formula: , optionally wherein each instance of R^E is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-SCN$, or $-NO_2$. In certain embodiments,


Ring C is of the formula: . In certain embodiments, Ring C is of

the formula: , optionally wherein each instance of R^E is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-SCN$, or $-NO_2$. In certain embodiments,

Ring C is of the formula: , , .




embodiments, Ring C is of the formula:




. In certain embodiments, Ring C is of

the formula:



, wherein each instance of R^E is independently substituted or


unsubstituted alkyl. In certain embodiments, Ring C is of the formula:



, wherein


each instance of R^E is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-$

SCN , or $-NO_2$. In certain embodiments, Ring C is of the formula:




. In certain

embodiments, Ring C is of the formula:




or




. In certain embodiments,


Ring C is of the formula:





. In certain embodiments, Ring C is of the formula:





. In certain embodiments, Ring C is of the formula:



, R^E





formula: , optionally wherein each instance of R^E is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^a$, $-N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, $-OC(=O)N(R^a)_2$, $-CN$, $-SCN$, or $-NO_2$. In certain embodiments,




[00253] In Formula (V), Ring C may include one or more substituents R^E . In certain embodiments, all instances of R^E are the same. In certain embodiments, at least two instances of R^E are different. In certain embodiments, at least one instance of R^E is halogen (*e.g.*, F, Cl, Br, or I). In certain embodiments, at least one instance of R^E is substituted or unsubstituted alkyl (*e.g.*, substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^E is $-CH_3$. In certain embodiments, at least one instance of R^E is $-CF_3$, unsubstituted ethyl, perfluoroethyl, unsubstituted propyl, perfluoropropyl, unsubstituted butyl, perfluorobutyl, or Bn. In certain embodiments, at least one instance of R^E is substituted or unsubstituted alkenyl (*e.g.*, substituted or unsubstituted C_{1-6} alkenyl). In certain embodiments, at least one instance of R^E is substituted or unsubstituted alkynyl (*e.g.*, substituted or unsubstituted C_{1-6} alkynyl). In certain embodiments, at least one instance of R^E is substituted or unsubstituted carbocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 3-

to 7-membered carbocyclyl). In certain embodiments, at least one instance of R^E is substituted or unsubstituted heterocyclyl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^E is substituted or unsubstituted aryl (*e.g.*, substituted or unsubstituted, 6- to 10-membered aryl). In certain embodiments, at least one instance of R^E is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^E is substituted or unsubstituted heteroaryl (*e.g.*, substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur). In certain embodiments, at least one instance of R^E is $-OR^a$ (*e.g.*, $-OH$, $-O$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-OMe$, $-OEt$, $-OPr$, $-OBu$, or $-OBn$), or $-O$ (substituted or unsubstituted phenyl) (*e.g.*, $-OPh$)). In certain embodiments, at least one instance of R^E is $-SR^a$ (*e.g.*, $-SH$, $-S$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-SMe$, $-SEt$, $-SPr$, $-SBu$, or $-SBn$), or $-S$ (substituted or unsubstituted phenyl) (*e.g.*, $-SPh$)). In certain embodiments, at least one instance of R^E is $-N(R^a)_2$ (*e.g.*, $-NH_2$, $-NH$ (substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-NHMe$), or $-N$ (substituted or unsubstituted C_{1-6} alkyl)–(substituted or unsubstituted C_{1-6} alkyl) (*e.g.*, $-NMe_2$)). In certain embodiments, at least one instance of R^E is $-CN$, $-SCN$, or $-NO_2$. In certain embodiments, at least one instance of R^E is $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, or $-C(=NR^a)N(R^a)_2$. In certain embodiments, at least one instance of R^E is $-C(=O)R^a$, $-C(=O)OR^a$, or $-C(=O)N(R^a)_2$ (*e.g.*, $-C(=O)NH_2$, $-C(=O)NHMe$, or $-C(=O)NMe_2$). In certain embodiments, at least one instance of R^E is $-NR^aC(=O)R^a$, optionally wherein each instance of R^a is independently H, substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted phenyl, or a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^E is $-NHC(=O)R^a$, wherein R^a is substituted or unsubstituted phenyl. In certain




embodiments, at least one instance of R^E is of the formula: , optionally wherein each instance of R^c is independently H, halogen, substituted or unsubstituted C_{1-6} alkyl, $-OH$, or $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least

one instance of R^E is of the formula: . In certain embodiments, at least one

instance of R^E is $-NR^aC(=O)OR^a$ or $-NR^aC(=O)N(R^a)_2$. In certain embodiments, at least one instance of R^E is $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$. In certain embodiments, at least one instance of R^E is $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, or $-NR^aS(=O)N(R^a)_2$, optionally wherein each instance of R^a is independently H, substituted or unsubstituted C_{1-6} alkyl, nitrogen protecting group when attached to a nitrogen atom, or an oxygen protecting group when attached to an oxygen atom. In certain embodiments, at least one instance of R^E is $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, or $-NR^aS(=O)_2N(R^a)_2$, optionally wherein each instance of R^a is independently H, substituted or unsubstituted C_{1-6} alkyl, nitrogen protecting group when attached to a nitrogen atom, or an oxygen protecting group when attached to an oxygen atom. In certain embodiments, at least one instance of R^E is $-NHS(=O)_2R^a$, optionally wherein R^a is substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^E is $-NHS(=O)_2Me$.


[00254] In certain embodiments, m is 0. In certain embodiments, m is 1. In certain embodiments, m is 2. In certain embodiments, m is 3. In certain embodiments, m is 4. In certain embodiments, m is 5.

[00255] In certain embodiments, the compound of Formula (V) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00256] In certain embodiments, the compound of Formula (V) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00257] In certain embodiments, the compound of Formula (V) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00258] In certain embodiments, the compound of Formula (V) useful in the present invention is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00259] In certain embodiments, the compound of Formula (V) useful in the present invention is not of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00260] In certain embodiments, the compound of Formula (V) useful in the present invention is of the formula:




(HG-9-91-01),

or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

Compounds of Formula (VI)

[00261] In another aspect, the compound utilized in the present disclosure is of Formula (VI):





(VI),

and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:


Ar₁ is a 5- or 6-membered hetero- or homo-cyclic aromatic ring optionally having a C₁-C₄ alkyl, or saturated heterocyclic or methyl-heterocyclic substituent;

X is separately N or CH;







[00262] In another aspect, the compound utilized in the present disclosure is of Formula (VI-A):



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:

Ar₁ is a 5- or 6-membered hetero- or homo-cyclic aromatic ring optionally having a C₁-C₄ alkyl, or saturated heterocyclic or methyl-heterocyclic substituent;







R₃ and R₄ are joined together with the intervening atoms to form a pyrrolidine ring where one or both of the free carbons are substituted with an alkyl or oxygen-containing substituent.





[00263] In certain embodiments, in Formula (VI) or (VI-A), Ar₁ is a 5- or 6-membered hetero- or homo-cyclic aromatic ring optionally having a C₁-C₄ alkyl, or saturated heterocyclic or methyl-heterocyclic substituent. In certain embodiments, a hetero-cyclic aromatic ring is a heteroaryl ring. In certain embodiments, a homo-cyclic aromatic ring is an aryl ring. In certain embodiments, in Formula (VI) or (VI-A), Ar₁ is optionally substituted 5-



or 6-membered heteroaryl or aryl. In certain embodiments, Ar₁ is optionally substituted with a C₁-C₄ alkyl, or saturated heterocyclic or methyl-heterocyclic substituent. In certain embodiments, Ar₁ is optionally substituted aryl. In certain embodiments, Ar₁ is optionally substituted phenyl. In certain embodiments, Ar₁ is unsubstituted phenyl. In certain embodiments, Ar₁ is optionally substituted 5- or 6-membered heteroaryl. In certain embodiments, Ar₁ is optionally substituted 5- membered heteroaryl. In certain embodiments, Ar₁ is optionally substituted pyrazole. In certain embodiments, Ar₁ is optionally substituted 6-membered heteroaryl. In certain embodiments, Ar₁ is optionally substituted heterocyclyl. In certain embodiments, Ar₁ is optionally substituted 5-10 membered heterocyclyl with 1–4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur. In certain embodiments, Ar₁ is a saturated heterocyclic or methyl-heterocyclic substituent.


[00264] In certain embodiments, X is separately N or CH. In certain embodiments, at least one instance of X is N. In certain embodiments, at least one instance of X is CH. In certain embodiments, one instance of X is N and the other instance of X is CH.


[00265] In certain embodiments, R₂ is  or , wherein R^{Z1} is hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted heterocyclyl, or


optionally substituted heteroaryl. In certain embodiments, R₂ is  or , wherein R^{Z1} is optionally substituted alkyl, optionally substituted aryl. In certain

embodiments, R₂ is , , , or . In certain


embodiments, R₂ is . In certain embodiments, R₂ is . In certain



embodiments, R₂ is , where R^{Z2} is optionally substituted alkyl, optionally substituted aryl, optionally substituted carbocyclyl, or optionally substituted heterocyclyl. In

certain embodiments, R₂ is , where R^{Z2} is optionally substituted carbocyclyl.


In certain embodiments, R₂ is . In certain embodiments, R₂ is optionally


substituted carbocyclyl. In certain embodiments, R_2 is optionally substituted C_{3-14} carbocyclyl. In certain embodiments, R_2 is optionally substituted C_{5-10} carbocyclyl. In certain

embodiments, R_2 is .



[00266] In certain embodiments, in Formula (VI-A), R_3 is hydrogen or . In certain embodiments, in Formula (VI-A), R_3 is hydrogen. In certain embodiments, R_3 is optionally substituted carbocyclyl. In certain embodiments, R_3 is optionally substituted C_{3-14} carbocyclyl. In certain embodiments, R_3 is optionally substituted C_{3-10} carbocyclyl. In certain embodiments, R_3 is optionally substituted C_{3-8} carbocyclyl. In certain embodiments, R_3 is .



[00267] In certain embodiments, R_4 is optionally substituted aryl (*e.g.*, optionally substituted phenyl or benzyl). In certain embodiments, R_4 is optionally substituted phenyl. In

certain embodiments, R_4 is hydrogen. In certain embodiments, R_4 is  or



. In certain embodiments, R_5 is H or optionally substituted alkyl. In certain


embodiments, R_4 is  or , where R_5 is H or optionally substituted

alkyl. In certain embodiments, R_4 is hydrogen, , or , where R_5 is H or a C_1 - C_4 alkyl.

[00268] In certain embodiments, R_3 and R_4 are joined together with the intervening atoms to form a pyrrolidine ring where at least one the free carbons is substituted with optionally

substituted alkyl, optionally substituted heterocyclyl, optionally substituted heteroaryl, or $-OR^{Z1}$, wherein R^{Z1} is hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted heterocyclyl, or optionally substituted heteroaryl. In certain embodiments, R_3 and R_4 are joined together with the intervening atoms to form a pyrrolidine ring where one or both of the free carbons are substituted with an alkyl or oxygen-containing substituent. In certain embodiments, R_3 and R_4 are joined together with the intervening atoms to form a pyrrolidine ring where both of the free carbons are substituted with optionally substituted alkyl. In certain embodiments, R_3 and R_4 are joined together with the intervening atoms to form a pyrrolidine ring where both of the free carbons are substituted with optionally substituted C_{1-6} alkyl. In certain embodiments, R_3 and R_4 are joined together with the intervening atoms to form a pyrrolidine ring where both of the free carbons are substituted with unsubstituted methyl.


[00269] In certain embodiments, the compound of Formula (VI) or (VI-A) useful in the present invention is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

Compounds of Formula (VII)

[00270] In another aspect, the compound utilized in the present disclosure is of Formula (VII):



and pharmaceutically acceptable salts, solvates, hydrates, polymorphs, co-crystals, tautomers, stereoisomers, isotopically labeled derivatives, and prodrugs thereof, wherein:

each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{A1}$, $-N(R^{A1})_2$, $-SR^{A1}$, $-CN$, $-SCN$, $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, $-C(=NR^{A1})N(R^{A1})_2$, $-C(=O)R^{A1}$, $-C(=O)OR^{A1}$, $-C(=O)N(R^{A1})_2$, $-NO_2$, $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$, $-NR^{A1}C(=O)N(R^{A1})_2$, $-OC(=O)R^{A1}$, $-OC(=O)OR^{A1}$, or $-OC(=O)N(R^{A1})_2$, wherein each instance of R^{A1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{A1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

j is 0, 1, or 2;

R^B is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or a nitrogen protecting group;

R^C is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^D is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or

unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^{\text{D1}}$, $-\text{N}(\text{R}^{\text{D1}})_2$, $-\text{SR}^{\text{D1}}$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^{\text{D1}})\text{R}^{\text{D1}}$, $-\text{C}(=\text{NR}^{\text{D1}})\text{OR}^{\text{D1}}$, $-\text{C}(=\text{NR}^{\text{D1}})\text{N}(\text{R}^{\text{D1}})_2$, $-\text{C}(=\text{O})\text{R}^{\text{D1}}$, $-\text{C}(=\text{O})\text{OR}^{\text{D1}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{D1}})_2$, $-\text{NO}_2$, $-\text{NR}^{\text{D1}}\text{C}(=\text{O})\text{R}^{\text{D1}}$, $-\text{NR}^{\text{D1}}\text{C}(=\text{O})\text{OR}^{\text{D1}}$, $-\text{NR}^{\text{D1}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{D1}})_2$, $-\text{OC}(=\text{O})\text{R}^{\text{D1}}$, $-\text{OC}(=\text{O})\text{OR}^{\text{D1}}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{D1}})_2$, wherein each instance of R^{D1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

m is 0, 1, or 2;

R^{E} is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^{F} is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-\text{OR}^{\text{F1}}$, $-\text{N}(\text{R}^{\text{F1}})_2$, $-\text{SR}^{\text{F1}}$, $-\text{CN}$, $-\text{SCN}$, $-\text{C}(=\text{NR}^{\text{F1}})\text{R}^{\text{F1}}$, $-\text{C}(=\text{NR}^{\text{F1}})\text{OR}^{\text{F1}}$, $-\text{C}(=\text{NR}^{\text{F1}})\text{N}(\text{R}^{\text{F1}})_2$, $-\text{C}(=\text{O})\text{R}^{\text{F1}}$, $-\text{C}(=\text{O})\text{OR}^{\text{F1}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{F1}})_2$, $-\text{NO}_2$, $-\text{NR}^{\text{F1}}\text{C}(=\text{O})\text{R}^{\text{F1}}$, $-\text{NR}^{\text{F1}}\text{C}(=\text{O})\text{OR}^{\text{F1}}$, $-\text{NR}^{\text{F1}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{F1}})_2$, $-\text{OC}(=\text{O})\text{R}^{\text{F1}}$, $-\text{OC}(=\text{O})\text{OR}^{\text{F1}}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{F1}})_2$, wherein each instance of R^{F1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{F1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring; and

n is 0, 1, 2, 3, 4, or 5.

[00271] Formula (VII) includes as Ring A a pyrimidinyl ring that is unsubstituted (*e.g.*, when j is 0) or substituted with one or two substituents R^{A} (*e.g.*, when j is 1 or 2). In certain

embodiments, the two instances of R^A are different. In certain embodiments, both instances of R^A are the same. In certain embodiments, at least one instance of R^A is halogen. In certain embodiments, at least one instance of R^A is F. In certain embodiments, at least one instance of R^A is Cl. In certain embodiments, at least one instance of R^A is Br. In certain embodiments, at least one instance of R^A is I (iodine). In certain embodiments, at least one instance of R^A is substituted alkyl. In certain embodiments, at least one instance of R^A is unsubstituted alkyl. In certain embodiments, at least one instance of R^A is unsubstituted C_{1-6} alkyl. In certain embodiments, both instances of R^A are unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^A is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^A is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^A is $-CH_3$. In certain embodiments, at least one instance of R^A is substituted methyl. In certain embodiments, at least one instance of R^A is $-CH_2F$. In certain embodiments, at least one instance of R^A is $-CHF_2$. In certain embodiments, at least one instance of R^A is $-CF_3$. In certain embodiments, at least one instance of R^A is ethyl. In certain embodiments, at least one instance of R^A is propyl. In certain embodiments, at least one instance of R^A is butyl. In certain embodiments, at least one instance of R^A is pentyl. In certain embodiments, at least one instance of R^A is hexyl. In certain embodiments, at least one instance of R^A is halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^A is substituted alkenyl. In certain embodiments, at least one instance of R^A is unsubstituted alkenyl. In certain embodiments, at least one instance of R^A is substituted alkynyl. In certain embodiments, at least one instance of R^A is unsubstituted alkynyl. In certain embodiments, at least one instance of R^A is substituted carbocyclyl. In certain embodiments, at least one instance of R^A is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^A is saturated carbocyclyl. In certain embodiments, at least one instance of R^A is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^A is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^A is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^A is substituted heterocyclyl. In certain embodiments, at least one instance of R^A is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^A is saturated heterocyclyl. In certain embodiments, at least one instance of R^A is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^A is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^A is monocyclic heterocyclyl. In certain embodiments, at least one instance of

R^A is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^A is substituted aryl. In certain embodiments, at least one instance of R^A is unsubstituted aryl. In certain embodiments, at least one instance of R^A is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^A is substituted phenyl. In certain embodiments, at least one instance of R^A is unsubstituted phenyl. In certain embodiments, at least one instance of R^A is substituted heteroaryl. In certain embodiments, at least one instance of R^A is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^A is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^A is monocyclic heteroaryl. In certain embodiments, at least one instance of R^A is 5-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^A is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^A is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^A is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^A is $-OR^{A1}$. In certain embodiments, at least one instance of R^A is $-OH$. In certain embodiments, at least one instance of R^A is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^A is $-OMe$. In certain embodiments, at least one instance of R^A is $-OEt$. In certain embodiments, at least one instance of R^A is $-OPr$. In certain embodiments, at least one instance of R^A is $-OBu$. In certain embodiments, at least one instance of R^A is $-OBn$. In certain embodiments, at least one instance of R^A is $-OPh$. In certain embodiments, at least one instance of R^A is $-SR^{A1}$. In certain embodiments, at least one instance of R^A is $-SH$. In certain embodiments, at least one instance of R^A is $-SMe$. In certain embodiments, at least one instance of R^A is $-N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-NH_2$. In certain embodiments, at least one instance of R^A is $-NHMe$. In certain embodiments, at least one instance of R^A is $-NMe_2$. In certain embodiments, at least one instance of R^A is $-CN$. In certain embodiments, at least one instance of R^A is $-SCN$. In certain embodiments, at least one instance of R^A is $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, or $-C(=NR^{A1})N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-C(=O)R^{A1}$ or $-C(=O)OR^{A1}$. In certain embodiments, at least one instance of R^A is $-C(=O)N(R^{A1})_2$. In certain embodiments, at least one instance of R^A is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, at least one instance of R^A is $-NO_2$. In certain embodiments, at least one instance of R^A is $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$,

or $-\text{NR}^{\text{A1}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{A1}})_2$. In certain embodiments, at least one instance of R^{A} is $-\text{OC}(=\text{O})\text{R}^{\text{A1}}$, $-\text{OC}(=\text{O})\text{OR}^{\text{A1}}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{A1}})_2$.


[00272] In certain embodiments, at least one instance of R^{A1} is H. In certain embodiments, at least one instance of R^{A1} is substituted acyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{A1} is acetyl. In certain embodiments, at least one instance of R^{A1} is substituted alkyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{A1} is methyl. In certain embodiments, at least one instance of R^{A1} is ethyl. In certain embodiments, at least one instance of R^{A1} is propyl. In certain embodiments, at least one instance of R^{A1} is butyl. In certain embodiments, at least one instance of R^{A1} is pentyl. In certain embodiments, at least one instance of R^{A1} is hexyl. In certain embodiments, at least one instance of R^{A1} is substituted alkenyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{A1} is substituted alkynyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{A1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{A1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{A1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{A1} is saturated heterocyclyl. In certain embodiments, at least one instance of R^{A1} is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^{A1} is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{A1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{A1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{A1} is monocyclic aryl. In certain embodiments, at least one instance of R^{A1} is substituted phenyl. In certain embodiments, at least one instance of R^{A1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{A1} is bicyclic aryl. In certain embodiments, at least one instance of R^{A1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{A1} is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain


embodiments, at least one instance of R^{A1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{A1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{A1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{A1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{A1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{A1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{A1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{A1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{A1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{A1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{A1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{A1} are joined to form a heterocyclic ring, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{A1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{A1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{A1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.

[00273] In certain embodiments, *j* is 0. In certain embodiments, *j* is 1. In certain embodiments, *j* is 2.


[00274] Formula (VII) includes substituent R^B on a nitrogen atom. In certain embodiments, R^B is H. In certain embodiments, R^B is substituted acyl. In certain embodiments, R^B is unsubstituted acyl. In certain embodiments, R^B is acetyl. In certain embodiments, R^B is substituted alkyl. In certain embodiments, R^B is unsubstituted alkyl. In certain embodiments, R^B is unsubstituted C_{1-6} alkyl. In certain embodiments, R^B is substituted C_{1-6} alkyl. In certain embodiments, R^B is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, R^B is $-CH_3$. In certain embodiments, R^B is substituted methyl. In certain embodiments, R^B is $-CH_2F$. In certain embodiments, R^B is $-CHF_2$. In certain embodiments, R^B is $-CF_3$. In certain embodiments, R^B is ethyl. In certain embodiments, R^B is


propyl. In certain embodiments, R^B is butyl. In certain embodiments, R^B is pentyl. In certain embodiments, R^B is hexyl. In certain embodiments, R^B is $-(CH_2)_{1-4}-(\text{Ring F})$, wherein Ring F is a substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, R^B is $-(CH_2)_{1-4}-(\text{substituted or unsubstituted pyrrolidinyl})$. In certain

embodiments, R^B is of the formula: , wherein R^4 is H, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group. In certain embodiments, R^B is of the formula:



In certain embodiments, R^B is $-(CH_2)_{1-4}-(\text{Ring F})$, wherein Ring F is a substituted or unsubstituted oxetanyl, substituted or unsubstituted azetidiny, substituted or unsubstituted tetrahydrofuranyl, substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidiny, substituted or unsubstituted piperazinyl, or substituted or unsubstituted morpholinyl ring. In certain embodiments, R^B is substituted alkenyl. In certain embodiments, R^B is unsubstituted alkenyl. In certain embodiments, R^B is substituted alkynyl. In certain embodiments, R^B is unsubstituted alkynyl. In certain embodiments, R^B is substituted carbocyclyl. In certain embodiments, R^B is unsubstituted carbocyclyl. In certain embodiments, R^B is saturated carbocyclyl. In certain embodiments, R^B is unsaturated carbocyclyl. In certain embodiments, R^B is monocyclic carbocyclyl. In certain embodiments, R^B is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^B is unsubstituted cyclopropyl. In certain embodiments, R^B is substituted cyclopropyl. In certain embodiments, R^B is substituted heterocyclyl. In certain embodiments, R^B is unsubstituted heterocyclyl. In certain embodiments, R^B is saturated heterocyclyl. In certain embodiments, R^B is unsaturated heterocyclyl. In certain embodiments, R^B is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^B is monocyclic heterocyclyl. In certain embodiments, R^B is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^B is substituted aryl. In certain embodiments, R^B is unsubstituted aryl. In certain embodiments, R^B is 6- to 10-membered aryl. In certain embodiments, R^B is substituted phenyl. In certain embodiments, R^B is unsubstituted phenyl. In certain embodiments, R^B is substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur. In certain embodiments, R^B is substituted or

unsubstituted pyrazolyl. In certain embodiments, R^B is of the formula: , wherein R^1 is H, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group. In

certain embodiments, R^B is of the formula: . In certain embodiments, R^B is substituted or unsubstituted furanyl, substituted or unsubstituted thienyl, substituted or unsubstituted pyrrolyl, substituted or unsubstituted imidazolyl, substituted or unsubstituted oxazolyl, substituted or unsubstituted isoxazolyl, substituted or unsubstituted thiazolyl, or substituted or unsubstituted isothiazolyl. In certain embodiments, R^B is substituted or unsubstituted pyridyl, substituted or unsubstituted pyrazinyl, substituted or unsubstituted pyrimidinyl, or substituted or unsubstituted pyridazinyl. In certain embodiments, R^B is substituted or unsubstituted, 9- to 10-membered, bicyclic heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur. In certain embodiments, R^B is a nitrogen protecting group. In certain embodiments, R^B is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts.

[00275] Formula (VII) includes substituent R^C on a nitrogen atom. In certain embodiments, R^C is H. In certain embodiments, R^C is substituted acyl. In certain embodiments, R^C is unsubstituted acyl. In certain embodiments, R^C is acetyl. In certain embodiments, R^C is unsubstituted C_{1-6} alkyl. In certain embodiments, R^C is substituted C_{1-6} alkyl. In certain embodiments, R^C is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, R^C is unsubstituted methyl. In certain embodiments, R^C is substituted methyl. In certain embodiments, R^C is $-CH_2F$. In certain embodiments, R^C is $-CHF_2$. In certain embodiments, R^C is $-CF_3$. In certain embodiments, R^C is ethyl. In certain embodiments, R^C is propyl. In certain embodiments, R^C is butyl. In certain embodiments, R^C is pentyl. In certain embodiments, R^C is hexyl. In certain embodiments, R^C is a nitrogen protecting group. In certain embodiments, R^C is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts.

[00276] In certain embodiments, R^B is substituted or unsubstituted C_{1-6} alkyl, and R^C is H. In certain embodiments, R^B is $-(CH_2)_{1-4}-(\text{Ring F})$, wherein Ring F is a substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclic ring; and R^C is H. In certain embodiments, R^B is substituted or unsubstituted phenyl (*e.g.*, *para*-substituted phenyl), and R^C is H. In certain embodiments, R^B is substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system

are independently nitrogen, oxygen, or sulfur; and R^C is H. In certain embodiments, R^B is substituted or unsubstituted pyrazolyl, and R^C is H. In certain embodiments, R^B is substituted or unsubstituted, 3- to 7-membered, monocyclic carbocyclyl; and R^C is H. In certain embodiments, R^B is substituted or unsubstituted cyclopropyl, and R^C is H.

[00277] Formula (VII) includes as Ring B an imidazolyl ring that is unsubstituted (*e.g.*, when m is 0) or substituted with one or two substituents R^D (*e.g.*, when m is 1 or 2). In certain embodiments, Ring B does not include substituents R^D , that is, m is 0. In certain embodiments, the two instances of R^D are different. In certain embodiments, both instances of R^D are the same. In certain embodiments, at least one instance of R^D is halogen. In certain embodiments, at least one instance of R^D is F. In certain embodiments, at least one instance of R^D is Cl. In certain embodiments, at least one instance of R^D is Br. In certain embodiments, at least one instance of R^D is I (iodine). In certain embodiments, at least one instance of R^D is substituted alkyl. In certain embodiments, at least one instance of R^D is unsubstituted alkyl. In certain embodiments, at least one instance of R^D is unsubstituted C_{1-6} alkyl. In certain embodiments, both instances of R^D are unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^D is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^D is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^D is $-CH_3$. In certain embodiments, at least one instance of R^D is substituted methyl. In certain embodiments, at least one instance of R^D is $-CH_2F$. In certain embodiments, at least one instance of R^D is $-CHF_2$. In certain embodiments, at least one instance of R^D is $-CF_3$. In certain embodiments, at least one instance of R^D is ethyl. In certain embodiments, at least one instance of R^D is propyl. In certain embodiments, at least one instance of R^D is butyl. In certain embodiments, at least one instance of R^D is pentyl. In certain embodiments, at least one instance of R^D is hexyl. In certain embodiments, at least one instance of R^D is halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^D is substituted alkenyl. In certain embodiments, at least one instance of R^D is unsubstituted alkenyl. In certain embodiments, at least one instance of R^D is substituted alkynyl. In certain embodiments, at least one instance of R^D is unsubstituted alkynyl. In certain embodiments, at least one instance of R^D is substituted carbocyclyl. In certain embodiments, at least one instance of R^D is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^D is saturated carbocyclyl. In certain embodiments, at least one instance of R^D is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^D is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^D is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least

one instance of R^D is substituted heterocyclyl. In certain embodiments, at least one instance of R^D is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^D is saturated heterocyclyl. In certain embodiments, at least one instance of R^D is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^D is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^D is monocyclic heterocyclyl. In certain embodiments, at least one instance of R^D is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^D is substituted aryl. In certain embodiments, at least one instance of R^D is unsubstituted aryl. In certain embodiments, at least one instance of R^D is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^D is substituted phenyl. In certain embodiments, at least one instance of R^D is unsubstituted phenyl. In certain embodiments, at least one instance of R^D is substituted heteroaryl. In certain embodiments, at least one instance of R^D is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^D is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^D is monocyclic heteroaryl. In certain embodiments, at least one instance of R^D is 5-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^D is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^D is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^D is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^D is $-OR^{D1}$. In certain embodiments, at least one instance of R^D is $-OH$. In certain embodiments, at least one instance of R^D is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^D is $-OMe$. In certain embodiments, at least one instance of R^D is $-OEt$. In certain embodiments, at least one instance of R^D is $-OPr$. In certain embodiments, at least one instance of R^D is $-OBu$. In certain embodiments, at least one instance of R^D is $-OBn$. In certain embodiments, at least one instance of R^D is $-OPh$. In certain embodiments, at least one instance of R^D is $-SR^{D1}$. In certain embodiments, at least one instance of R^D is $-SH$. In certain embodiments, at least one instance of R^D is $-SMe$. In certain embodiments, at least one instance of R^D is $-N(R^{D1})_2$. In certain embodiments, at least one instance of R^D is $-NH_2$. In certain embodiments, at least one instance of R^D is $-NHMe$. In certain embodiments, at least one instance of R^D is $-NMe_2$. In certain embodiments, at least one instance of R^D is $-CN$. In certain embodiments, at least

one instance of R^D is $-\text{SCN}$. In certain embodiments, at least one instance of R^D is $-\text{C}(=\text{NR}^{D1})\text{R}^{D1}$, $-\text{C}(=\text{NR}^{D1})\text{OR}^{D1}$, or $-\text{C}(=\text{NR}^{D1})\text{N}(\text{R}^{D1})_2$. In certain embodiments, at least one instance of R^D is $-\text{C}(=\text{O})\text{R}^{D1}$ or $-\text{C}(=\text{O})\text{OR}^{D1}$. In certain embodiments, at least one instance of R^D is $-\text{C}(=\text{O})\text{N}(\text{R}^{D1})_2$. In certain embodiments, at least one instance of R^D is $-\text{C}(=\text{O})\text{NMe}_2$, $-\text{C}(=\text{O})\text{NHMe}$, or $-\text{C}(=\text{O})\text{NH}_2$. In certain embodiments, at least one instance of R^D is $-\text{NO}_2$. In certain embodiments, at least one instance of R^D is $-\text{NR}^{D1}\text{C}(=\text{O})\text{R}^{D1}$, $-\text{NR}^{D1}\text{C}(=\text{O})\text{OR}^{D1}$, or $-\text{NR}^{D1}\text{C}(=\text{O})\text{N}(\text{R}^{D1})_2$. In certain embodiments, at least one instance of R^D is $-\text{OC}(=\text{O})\text{R}^{D1}$, $-\text{OC}(=\text{O})\text{OR}^{D1}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^{D1})_2$.

[00278] In certain embodiments, at least one instance of R^{D1} is H. In certain embodiments, at least one instance of R^{D1} is substituted acyl. In certain embodiments, at least one instance of R^{D1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{D1} is acetyl. In certain embodiments, at least one instance of R^{D1} is substituted alkyl. In certain embodiments, at least one instance of R^{D1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{D1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{D1} is methyl. In certain embodiments, at least one instance of R^{D1} is ethyl. In certain embodiments, at least one instance of R^{D1} is propyl. In certain embodiments, at least one instance of R^{D1} is butyl. In certain embodiments, at least one instance of R^{D1} is pentyl. In certain embodiments, at least one instance of R^{D1} is hexyl. In certain embodiments, at least one instance of R^{D1} is substituted alkenyl. In certain embodiments, at least one instance of R^{D1} is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{D1} is substituted alkynyl. In certain embodiments, at least one instance of R^{D1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{D1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{D1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{D1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{D1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{D1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{D1} is saturated heterocyclyl. In certain embodiments, at least one instance of R^{D1} is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^{D1} is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{D1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{D1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{D1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{D1} is monocyclic aryl. In

certain embodiments, at least one instance of R^{D1} is substituted phenyl. In certain embodiments, at least one instance of R^{D1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{D1} is bicyclic aryl. In certain embodiments, at least one instance of R^{D1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{D1} is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{D1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{D1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{D1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{D1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{D1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{D1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{D1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{D1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{D1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{D1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{D1} are joined to form a heterocyclic ring, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{D1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{D1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{D1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.


[00279] In certain embodiments, *m* is 0. In certain embodiments, *m* is 1. In certain embodiments, *m* is 2.


[00280] Formula (VII) includes substituent R^E on a nitrogen atom. In certain embodiments, R^E is H. In certain embodiments, R^E is substituted acyl. In certain embodiments, R^E is unsubstituted acyl. In certain embodiments, R^E is acetyl. In certain



embodiments, R^E is unsubstituted C_{1-6} alkyl. In certain embodiments, R^E is substituted C_{1-6} alkyl. In certain embodiments, R^E is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, R^E is unsubstituted methyl. In certain embodiments, R^E is substituted methyl. In certain embodiments, R^E is $-CH_2F$. In certain embodiments, R^E is $-CHF_2$. In certain embodiments, R^E is $-CF_3$. In certain embodiments, R^E is ethyl. In certain embodiments, R^E is propyl. In certain embodiments, R^E is butyl. In certain embodiments, R^E is pentyl. In certain embodiments, R^E is hexyl. In certain embodiments, R^E is a nitrogen protecting group. In certain embodiments, R^E is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts.


[00281] In certain embodiments, each of R^C and R^E is H.



[00282] Formula (VII) includes as Ring C a phenyl ring that is unsubstituted (*e.g.*, when n is 0) or substituted with one or more substituents R^F (*e.g.*, when n is 1, 2, 3, 4, or 5). In



certain embodiments, Ring C is of the formula: . In certain embodiments, Ring C is


of the formula: . In certain embodiments, Ring C is of the formula:

. In certain embodiments, Ring C is of the formula: . In certain

embodiments, Ring C is of the formula: , wherein each instance of R^F is independently halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments,

Ring C is of the formula:  or . In certain embodiments, Ring C is of the

formula: . In certain embodiments, Ring C is of the formula: . In



certain embodiments, Ring C is of the formula:

. In certain embodiments, at least two instances of R^F are different. In certain embodiments, all instances of R^F are the same. In certain embodiments, at least one instance of R^F is halogen. In certain embodiments, at least one instance of R^F is F. In certain embodiments, at least one instance of R^F is Cl. In certain embodiments, at least one instance of R^F is Br. In certain embodiments, at least one instance of R^F is I (iodine). In certain embodiments, at least one instance of R^F is substituted alkyl. In certain embodiments, at least one instance of R^F is unsubstituted alkyl. In certain embodiments, at least one instance of R^F is unsubstituted C_{1-6} alkyl. In certain embodiments, all instances of R^F are unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^F is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^F is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^F is $-CH_3$. In certain embodiments, all instances of R^F are $-CH_3$. In certain embodiments, at least one instance of R^F is substituted methyl. In certain embodiments, at least one instance of R^F is $-CH_2F$. In certain embodiments, at least one instance of R^F is $-CHF_2$. In certain embodiments, at least one instance of R^F is $-CF_3$. In certain embodiments, at least one instance of R^F is ethyl. In certain embodiments, at least one instance of R^F is propyl. In certain embodiments, at least one instance of R^F is butyl. In certain embodiments, at least one instance of R^F is pentyl. In certain embodiments, at least one instance of R^F is hexyl. In certain embodiments, at least one instance of R^F is substituted alkenyl. In certain embodiments, at least one instance of R^F is unsubstituted alkenyl. In certain embodiments, at least one instance of R^F is substituted alkynyl. In certain embodiments, at least one instance of R^F is unsubstituted alkynyl. In certain embodiments, at least one instance of R^F is substituted carbocyclyl. In certain embodiments, at least one instance of R^F is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^F is saturated carbocyclyl. In certain embodiments, at least one instance of R^F is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^F is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^F is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^F is substituted heterocyclyl. In certain embodiments, at least one instance of R^F is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^F is saturated heterocyclyl. In certain embodiments, at least one instance of R^F is unsaturated heterocyclyl. In certain embodiments, at least one

instance of R^F is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^F is monocyclic heterocyclyl. In certain embodiments, at least one instance of R^F is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^F is substituted aryl. In certain embodiments, at least one instance of R^F is unsubstituted aryl. In certain embodiments, at least one instance of R^F is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^F is substituted phenyl. In certain embodiments, at least one instance of R^F is unsubstituted phenyl. In certain embodiments, at least one instance of R^F is substituted heteroaryl. In certain embodiments, at least one instance of R^F is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^F is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^F is monocyclic heteroaryl. In certain embodiments, at least one instance of R^F is 5-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^F is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^F is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^F is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^F is $-OR^{F1}$. In certain embodiments, at least one instance of R^F is $-OH$. In certain embodiments, at least one instance of R^F is $-O(\text{substituted or unsubstituted } C_{1-6} \text{ alkyl})$. In certain embodiments, at least one instance of R^F is $-OMe$. In certain embodiments, at least one instance of R^F is $-OEt$. In certain embodiments, at least one instance of R^F is $-OPr$. In certain embodiments, at least one instance of R^F is $-OBu$. In certain embodiments, at least one instance of R^F is $-OBn$. In certain embodiments, at least one instance of R^F is $-OPh$. In certain embodiments, at least one instance of R^F is $-SR^{F1}$. In certain embodiments, at least one instance of R^F is $-SH$. In certain embodiments, at least one instance of R^F is $-SMe$. In certain embodiments, at least one instance of R^F is $-N(R^{F1})_2$. In certain embodiments, at least one instance of R^F is $-NH_2$. In certain embodiments, at least one instance of R^F is $-NHMe$. In certain embodiments, at least one instance of R^F is $-NMe_2$. In certain embodiments, at least one instance of R^F is $-CN$. In certain embodiments, at least one instance of R^F is $-SCN$. In certain embodiments, at least one instance of R^F is $-C(=NR^{F1})R^{F1}$, $-C(=NR^{F1})OR^{F1}$, or $-C(=NR^{F1})N(R^{F1})_2$. In certain embodiments, at least one instance of R^F is $-C(=O)R^{F1}$ or $-C(=O)OR^{F1}$. In certain embodiments, at least one instance of R^F is $-C(=O)N(R^{F1})_2$.

[00283] In certain embodiments, at least one instance of R^F is $-C(=O)N(R^{F1})_2$, wherein each instance of R^{F1} is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted phenyl, or a nitrogen protecting group. In certain embodiments, at least one instance of R^F is $-C(=O)NHR^{F1}$, wherein R^{F1} is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^F is $-C(=O)NHR^{F1}$, wherein R^{F1} is phenyl substituted with one, two, three, four, or five substituents independently selected from the group consisting of halogen and substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^F is $-C(=O)NMeR^{F1}$, wherein R^{F1} is substituted or unsubstituted phenyl. In certain embodiments, at least one instance of R^F is $-C(=O)NMeR^{F1}$, wherein R^{F1} is phenyl substituted with one, two, three, four, or five substituents independently selected from the group consisting of halogen and substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^F is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, at least one instance of R^F is $-NO_2$. In certain embodiments, at least one instance of R^F is $-NR^{F1}C(=O)R^{F1}$, $-NR^{F1}C(=O)OR^{F1}$, or $-NR^{F1}C(=O)N(R^{F1})_2$. In certain embodiments, at least one instance of R^F is $-OC(=O)R^{F1}$, $-OC(=O)OR^{F1}$, or $-OC(=O)N(R^{F1})_2$.

[00284] In certain embodiments, at least one instance of R^F is halogen, substituted or unsubstituted C_{1-6} alkyl, or $-OR^{F1}$. In certain embodiments, at least one instance of R^F is halogen, substituted or unsubstituted C_{1-6} alkyl, or $-OR^{F1}$, wherein R^{F1} is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or an oxygen protecting group. In certain embodiments, at least one instance of R^F is halogen, unsubstituted C_{1-6} alkyl, or $-OR^{F1}$, wherein R^{F1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^F is $-CH_3$ or Cl .

[00285] In certain embodiments, at least one instance of R^{F1} is H . In certain embodiments, at least one instance of R^{F1} is substituted acyl. In certain embodiments, at least one instance of R^{F1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{F1} is acetyl. In certain embodiments, at least one instance of R^{F1} is substituted alkyl. In certain embodiments, at least one instance of R^{F1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{F1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{F1} is methyl. In certain embodiments, at least one instance of R^{F1} is ethyl. In certain embodiments, at least one instance of R^{F1} is propyl. In certain embodiments, at least one instance of R^{F1} is butyl. In certain embodiments, at least one instance of R^{F1} is pentyl. In certain embodiments, at least one instance of R^{F1} is hexyl. In certain embodiments, at least one instance of R^{F1} is substituted alkenyl. In certain embodiments, at least one instance of R^{F1}


is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{F1} is substituted alkynyl. In certain embodiments, at least one instance of R^{F1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{F1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{F1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{F1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{F1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{F1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{F1} is saturated heterocyclyl. In certain embodiments, at least one instance of R^{F1} is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^{F1} is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{F1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{F1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{F1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{F1} is monocyclic aryl. In certain embodiments, at least one instance of R^{F1} is substituted phenyl. In certain embodiments, at least one instance of R^{F1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{F1} is bicyclic aryl. In certain embodiments, at least one instance of R^{F1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{F1} is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{F1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{F1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{F1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{F1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{F1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{F1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{F1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{F1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{F1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of

R^{F1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{F1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{F1} are joined to form a heterocyclic ring, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{F1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{F1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{F1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.

[00286] In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2. In certain embodiments, n is 3. In certain embodiments, n is 4. In certain embodiments, n is 5.

[00287] In certain embodiments, n is 1; and R^F is $-C(=O)N(R^{F1})_2$. In certain embodiments, n is 1; and R^F is $-C(=O)N(R^{F1})_2$, wherein each instance of R^{F1} is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted phenyl, or a nitrogen protecting group. In certain embodiments, n is 2; and each instance of R^F is independently halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, n is 2; and each instance of R^F is independently halogen or unsubstituted C_{1-6} alkyl (*e.g.*, $-CH_3$). In certain embodiments, n is 2; and each instance of R^F is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^{F1}$, or $-C(=O)N(R^{F1})_2$. In certain embodiments, n is 2; and each instance of R^F is independently halogen, substituted or unsubstituted C_{1-6} alkyl, $-OR^{F1}$, or $-C(=O)N(R^{F1})_2$, wherein each instance of R^{F1} is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted phenyl, an oxygen protecting group when attached to an oxygen atom, or a nitrogen protecting group when attached to a nitrogen atom.

[00288] In certain embodiments, the compound of Formula (VII) is of Formula (VII-A):




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof, wherein:


each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{G1}$, $-N(R^{G1})_2$, $-SR^{G1}$, $-CN$, $-SCN$, $-C(=NR^{G1})R^{G1}$, $-C(=NR^{G1})OR^{G1}$, $-C(=NR^{G1})N(R^{G1})_2$, $-C(=O)R^{G1}$, $-C(=O)OR^{G1}$, $-C(=O)N(R^{G1})_2$, $-NO_2$, $-NR^{G1}C(=O)R^{G1}$, $-NR^{G1}C(=O)OR^{G1}$, $-NR^{G1}C(=O)N(R^{G1})_2$, $-OC(=O)R^{G1}$, $-OC(=O)OR^{G1}$, or $-OC(=O)N(R^{G1})_2$, wherein each instance of R^{G1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{G1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring; and



k is 0, 1, 2, 3, 4, or 5.

[00289] Formula (VII-A) includes as Ring D a phenyl ring that is unsubstituted (*e.g.*, when k is 0) or substituted with one or more substituents R^G (*e.g.*, when k is 1, 2, 3, 4, or 5).




In certain embodiments, Ring D is of the formula: . In certain embodiments, Ring D


is of the formula: . In certain embodiments, Ring D is of the formula:



instances of R^G are different. In certain embodiments, all instances of R^G are the same. In certain embodiments, at least one instance of R^G is halogen. In certain embodiments, at least one instance of R^G is F. In certain embodiments, at least one instance of R^G is Cl. In certain

embodiments, at least one instance of R^G is Br. In certain embodiments, at least one instance of R^G is I (iodine). In certain embodiments, at least one instance of R^G is substituted alkyl. In certain embodiments, at least one instance of R^G is unsubstituted alkyl. In certain embodiments, at least one instance of R^G is unsubstituted C_{1-6} alkyl. In certain embodiments, all instances of R^G are unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^G is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^G is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^G is $-CH_3$. In certain embodiments, all instances of R^G are $-CH_3$. In certain embodiments, at least one instance of R^G is substituted methyl. In certain embodiments, at least one instance of R^G is $-CH_2F$. In certain embodiments, at least one instance of R^G is $-CHF_2$. In certain embodiments, at least one instance of R^G is $-CF_3$. In certain embodiments, at least one instance of R^G is ethyl. In certain embodiments, at least one instance of R^G is propyl. In certain embodiments, at least one instance of R^G is butyl. In certain embodiments, at least one instance of R^G is pentyl. In certain embodiments, at least one instance of R^G is hexyl. In certain embodiments, at least one instance of R^G is substituted alkenyl. In certain embodiments, at least one instance of R^G is unsubstituted alkenyl. In certain embodiments, at least one instance of R^G is substituted alkynyl. In certain embodiments, at least one instance of R^G is unsubstituted alkynyl. In certain embodiments, at least one instance of R^G is substituted carbocyclyl. In certain embodiments, at least one instance of R^G is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^G is saturated carbocyclyl. In certain embodiments, at least one instance of R^G is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^G is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^G is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^G is substituted heterocyclyl. In certain embodiments, at least one instance of R^G is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^G is saturated heterocyclyl. In certain embodiments, at least one instance of R^G is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^G is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^G is monocyclic heterocyclyl. In certain embodiments, at least one instance of R^G is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^G is substituted or unsubstituted piperazinyl. In certain embodiments, at least one

instance of R^G is of the formula: , wherein R^2 is H, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group. In certain embodiments, at least one instance of R^G

is of the formula: . In certain embodiments, at least one instance of R^G is substituted aryl. In certain embodiments, at least one instance of R^G is unsubstituted aryl. In certain embodiments, at least one instance of R^G is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^G is substituted phenyl. In certain embodiments, at least one instance of R^G is unsubstituted phenyl. In certain embodiments, at least one instance of R^G is substituted heteroaryl. In certain embodiments, at least one instance of R^G is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^G is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^G is monocyclic heteroaryl. In certain embodiments, at least one instance of R^G is 5-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^G is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^G is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^G is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^G is $-OR^{G1}$. In certain embodiments, at least one instance of R^G is $-OH$. In certain embodiments, at least one instance of R^G is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^G is $-O-(CH_2)_{2-4}-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, at least one instance of R^G is $-O-(CH_2)_2-OMe$. In certain embodiments, at least one instance of R^G is $-OMe$. In certain embodiments, at least one instance of R^G is $-OEt$. In certain embodiments, at least one instance of R^G is $-OPr$. In certain embodiments, at least one instance of R^G is $-OBu$. In certain embodiments, at least one instance of R^G is $-OBn$. In certain embodiments, at least one instance of R^G is $-OPh$. In certain embodiments, at least one instance of R^G is $-SR^{G1}$. In certain embodiments, at least one instance of R^G is $-SH$. In certain embodiments, at least one instance of R^G is $-SMe$. In certain embodiments, at least one instance of R^G is $-N(R^{G1})_2$. In certain embodiments, at least one instance of R^G is $-N(R^{G1})_2$, wherein each instance of R^{G1} is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or two

instances of R^{G1} are joined to form a substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, at least one instance of R^G is $-NH_2$. In certain embodiments, at least one instance of R^G is $-NHMe$. In certain embodiments, at least one instance of R^G is $-NMe_2$. In certain embodiments, at least one instance of R^G is $-CN$. In certain embodiments, at least one instance of R^G is $-SCN$. In certain embodiments, at least one instance of R^G is $-C(=NR^{G1})R^{G1}$, $-C(=NR^{G1})OR^{G1}$, or $-C(=NR^{G1})N(R^{G1})_2$. In certain embodiments, at least one instance of R^G is $-C(=O)R^{G1}$ or $-C(=O)OR^{G1}$. In certain embodiments, at least one instance of R^G is $-C(=O)N(R^{G1})_2$. In certain embodiments, at least one instance of R^G is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, at least one instance of R^G is $-NO_2$. In certain embodiments, at least one instance of R^G is $-NR^{G1}C(=O)R^{G1}$, $-NR^{G1}C(=O)OR^{G1}$, or $-NR^{G1}C(=O)N(R^{G1})_2$. In certain embodiments, at least one instance of R^G is $-OC(=O)R^{G1}$, $-OC(=O)OR^{G1}$, or $-OC(=O)N(R^{G1})_2$.

[00290] In certain embodiments, at least one instance of R^G is $-OR^{G1}$, $-N(R^{G1})_2$, or substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently nitrogen, oxygen, or sulfur.

[00291] In certain embodiments, at least one instance of R^{G1} is H. In certain embodiments, at least one instance of R^{G1} is substituted acyl. In certain embodiments, at least one instance of R^{G1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{G1} is acetyl. In certain embodiments, at least one instance of R^{G1} is substituted alkyl. In certain embodiments, at least one instance of R^{G1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{G1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{G1} is methyl. In certain embodiments, at least one instance of R^{G1} is ethyl. In certain embodiments, at least one instance of R^{G1} is propyl. In certain embodiments, at least one instance of R^{G1} is butyl. In certain embodiments, at least one instance of R^{G1} is pentyl. In certain embodiments, at least one instance of R^{G1} is hexyl. In certain embodiments, at least one instance of R^{G1} is substituted alkenyl. In certain embodiments, at least one instance of R^{G1} is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{G1} is substituted alkynyl. In certain embodiments, at least one instance of R^{G1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{G1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{G1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{G1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{G1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{G1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{G1} is saturated heterocyclyl. In certain


embodiments, at least one instance of R^{G1} is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^{G1} is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{G1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{G1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{G1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{G1} is monocyclic aryl. In certain embodiments, at least one instance of R^{G1} is substituted phenyl. In certain embodiments, at least one instance of R^{G1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{G1} is bicyclic aryl. In certain embodiments, at least one instance of R^{G1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{G1} is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{G1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{G1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{G1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{G1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{G1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{G1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{G1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{G1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{G1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{G1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{G1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{G1} are joined to form a heterocyclic ring, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{G1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{G1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{G1} are joined to form a substituted

or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.

[00292] In certain embodiments, k is 0. In certain embodiments, k is 1. In certain embodiments, k is 2. In certain embodiments, k is 3. In certain embodiments, k is 4. In certain embodiments, k is 5.


[00293] In certain embodiments, k is 1; and R^G is $-OR^{G1}$, $-N(R^{G1})_2$, or substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently nitrogen, oxygen, or sulfur. In certain embodiments, k is 1; and R^G is $-OR^{G1}$, $-N(R^{G1})_2$, or substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently nitrogen, oxygen, or sulfur; and each instance of R^{G1} is independently H, substituted or unsubstituted C_{1-6} alkyl, an oxygen protecting group when attached to an oxygen atom, or a nitrogen protecting group when attached to a nitrogen group.

[00294] In certain embodiments, the compound of Formula (VII) is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00295] In certain embodiments, the compound of Formula (VII) is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00296] In certain embodiments, the compound of Formula (VII) is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00297] In certain embodiments, the compound of Formula (VII) is of the formula:




or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00298] In certain embodiments, the compound of Formula (VII) is of the formula:



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00299] In certain embodiments, the compound of Formula (VII) is of Formula (VII-B):



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof, wherein:


R^H is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^J is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^J$, $-N(R^J)_2$, $-SR^J$, $-CN$, $-SCN$, $-C(=NR^J)R^J$, $-C(=NR^J)OR^J$, $-C(=NR^J)N(R^J)_2$, $-C(=O)R^J$, $-C(=O)OR^J$, $-C(=O)N(R^J)_2$, $-NO_2$, $-NR^J C(=O)R^J$, $-NR^J C(=O)OR^J$, $-NR^J C(=O)N(R^J)_2$, $-OC(=O)R^J$, $-OC(=O)OR^J$, or $-OC(=O)N(R^J)_2$, wherein each instance of R^J is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^J are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

q is 0, 1, 2, 3, or 4; and

R^K is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^K$, $-N(R^K)_2$, $-SR^K$, $-CN$, $-SCN$, $-C(=NR^K)R^K$, $-C(=NR^K)OR^K$, $-C(=NR^K)N(R^K)_2$, $-C(=O)R^K$, $-C(=O)OR^K$, $-C(=O)N(R^K)_2$, $-NO_2$, $-NR^K C(=O)R^K$, $-NR^K C(=O)OR^K$, $-NR^K C(=O)N(R^K)_2$, $-OC(=O)R^K$, $-OC(=O)OR^K$, or $-OC(=O)N(R^K)_2$, wherein each instance of R^K is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^K are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.


[00300] In certain embodiments, a compound of Formula (VII) is of Formula (VII-B),



wherein when R^C is hydrogen, R^B is not unsubstituted cyclopropyl or .


[00301] Formula (VII-B) includes substituent R^H on a nitrogen atom. In certain embodiments, R^H is H. In certain embodiments, R^H is not H. In certain embodiments, R^H is substituted acyl. In certain embodiments, R^H is unsubstituted acyl. In certain embodiments, R^H is acetyl. In certain embodiments, R^H is unsubstituted C_{1-6} alkyl. In certain embodiments, R^H is substituted C_{1-6} alkyl. In certain embodiments, R^H is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, R^H is unsubstituted methyl. In certain embodiments, R^H is substituted methyl. In certain embodiments, R^H is $-CH_2F$. In certain embodiments, R^H is $-CHF_2$. In certain embodiments, R^H is $-CF_3$. In certain embodiments, R^H is ethyl. In certain embodiments, R^H is propyl. In certain embodiments, R^H is butyl. In certain embodiments, R^H is pentyl. In certain embodiments, R^H is hexyl. In certain embodiments, R^H is a nitrogen protecting group. In certain embodiments, R^H is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts. In certain embodiments, R^H is hydrogen or unsubstituted C_{1-6} alkyl.


[00302] Formula (VII-B) includes as Ring E a phenyl ring that is substituted with R^K and optionally one or more substituents R^J . In certain embodiments, Ring E is of the formula:




embodiments, Ring E is of the formula: . In certain embodiments, Ring E is of

the formula: . In certain embodiments, Ring E is of the formula: .


In certain embodiments, Ring E is of the formula: . In certain embodiments,





Ring E is of the formula: . In certain embodiments, R^K is H. In certain embodiments, R^K is halogen. In certain embodiments, R^K is F. In certain embodiments, R^K is Cl. In certain embodiments, R^K is Br. In certain embodiments, R^K is I (iodine). In certain embodiments, R^K is substituted alkyl. In certain embodiments, R^K is unsubstituted alkyl. In certain embodiments, R^K is unsubstituted C₁₋₆ alkyl. In certain embodiments, R^K is substituted C₁₋₆ alkyl. In certain embodiments, R^K is C₁₋₆ alkyl substituted with at least one halogen. In certain embodiments, R^K is –CH₃. In certain embodiments, R^K is substituted methyl. In certain embodiments, R^K is –CH₂F. In certain embodiments, R^K is –CHF₂. In certain embodiments, R^K is –CF₃. In certain embodiments, R^K is ethyl. In certain embodiments, R^K is propyl. In certain embodiments, R^K is butyl. In certain embodiments, R^K is pentyl. In certain embodiments, R^K is hexyl. In certain embodiments, R^K is substituted alkenyl. In certain embodiments, R^K is unsubstituted alkenyl. In certain embodiments, R^K is substituted alkynyl. In certain embodiments, R^K is unsubstituted alkynyl. In certain embodiments, R^K is substituted carbocyclyl. In certain embodiments, R^K is unsubstituted carbocyclyl. In certain embodiments, R^K is saturated carbocyclyl. In certain embodiments, R^K is unsaturated carbocyclyl. In certain embodiments, R^K is monocyclic carbocyclyl. In certain embodiments, R^K is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^K is substituted heterocyclyl. In certain embodiments, R^K is unsubstituted heterocyclyl. In certain embodiments, R^K is saturated heterocyclyl. In certain embodiments, R^K is unsaturated heterocyclyl. In certain embodiments, R^K is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^K is monocyclic heterocyclyl. In certain embodiments, R^K is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^K is substituted aryl. In certain embodiments, R^K is unsubstituted aryl. In certain embodiments, R^K is 6- to 10-membered aryl. In certain embodiments, R^K is substituted phenyl. In certain embodiments, R^K is unsubstituted phenyl. In certain embodiments, R^K is substituted heteroaryl. In certain embodiments, R^K is unsubstituted heteroaryl. In certain embodiments, R^K is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^K is monocyclic heteroaryl. In certain embodiments, R^K is 5-membered, monocyclic heteroaryl. In certain embodiments, R^K is not substituted imidazolyl. In certain

embodiments, R^K is 6-membered, monocyclic heteroaryl. In certain embodiments, R^K is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, R^K is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, R^K is $-OR^{K1}$. In certain embodiments, R^K is $-OH$. In certain embodiments, R^K is $-O$ (substituted or unsubstituted C_{1-6} alkyl). In certain embodiments, R^K is $-OMe$. In certain embodiments, R^K is $-OEt$. In certain embodiments, R^K is $-OPr$. In certain embodiments, R^K is $-OBu$. In certain embodiments, R^K is $-OBn$. In certain embodiments, R^K is $-OPh$. In certain embodiments, R^K is $-SR^{K1}$. In certain embodiments, R^K is $-SH$. In certain embodiments, R^K is $-SMe$. In certain embodiments, R^K is $-N(R^{K1})_2$. In certain embodiments, R^K is $-NH_2$. In certain embodiments, R^K is $-NHMe$. In certain embodiments, R^K is $-NMe_2$. In certain embodiments, R^K is $-CN$. In certain embodiments, R^K is $-SCN$. In certain embodiments, R^K is $-C(=NR^{K1})R^{K1}$, $-C(=NR^{K1})OR^{K1}$, or $-C(=NR^{K1})N(R^{K1})_2$. In certain embodiments, R^K is $-C(=O)R^{K1}$ or $-C(=O)OR^{K1}$. In certain embodiments, R^K is $-C(=O)N(R^{K1})_2$. In certain embodiments, R^K is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, R^K is $-NO_2$. In certain embodiments, R^K is $-NR^{K1}C(=O)R^{K1}$, $-NR^{K1}C(=O)OR^{K1}$, or $-NR^{K1}C(=O)N(R^{K1})_2$. In certain embodiments, R^K is $-OC(=O)R^{K1}$, $-OC(=O)OR^{K1}$, or $-OC(=O)N(R^{K1})_2$.

[00303] In certain embodiments, R^K is $-(CH_2)_{1-3}-(\text{Ring G})$, wherein Ring G is a substituted or unsubstituted, 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, R^K is $-(CH_2)_{1-3}-(\text{substituted or unsubstituted piperazinyl})$. In certain

embodiments, R^K is of the formula: , wherein R^3 is H, substituted or unsubstituted C_{2-6} alkyl, substituted methyl, or a nitrogen protecting group. In certain

embodiments, R^K is of the formula: . In certain embodiments, R^K is not of

the formula: . In certain embodiments, R^K is $-(CH_2)_{1-3}-(\text{Ring G})$, wherein Ring G is a substituted or unsubstituted oxetanyl, substituted or unsubstituted azetidiny, substituted or unsubstituted tetrahydrofuranyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidinyl, or substituted or unsubstituted morpholinyl ring.

[00304] In certain embodiments, R^{K1} is H. In certain embodiments, R^{K1} is substituted acyl. In certain embodiments, R^{K1} is unsubstituted acyl. In certain embodiments, R^{K1} is acetyl. In

certain embodiments, R^{K1} is substituted alkyl. In certain embodiments, R^{K1} is unsubstituted alkyl. In certain embodiments, R^{K1} is unsubstituted C_{1-6} alkyl. In certain embodiments, R^{K1} is methyl. In certain embodiments, R^{K1} is ethyl. In certain embodiments, R^{K1} is propyl. In certain embodiments, R^{K1} is butyl. In certain embodiments, R^{K1} is pentyl. In certain embodiments, R^{K1} is hexyl. In certain embodiments, R^{K1} is substituted alkenyl. In certain embodiments, R^{K1} is unsubstituted alkenyl. In certain embodiments, R^{K1} is substituted alkynyl. In certain embodiments, R^{K1} is unsubstituted alkynyl. In certain embodiments, R^{K1} is substituted or unsubstituted carbocyclyl. In certain embodiments, R^{K1} is saturated carbocyclyl. In certain embodiments, R^{K1} is unsaturated carbocyclyl. In certain embodiments, R^{K1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, R^{K1} is substituted or unsubstituted heterocyclyl. In certain embodiments, R^{K1} is saturated heterocyclyl. In certain embodiments, R^{K1} is unsaturated heterocyclyl. In certain embodiments, R^{K1} is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^{K1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, R^{K1} is substituted or unsubstituted aryl. In certain embodiments, R^{K1} is 6- to 10-membered aryl. In certain embodiments, R^{K1} is monocyclic aryl. In certain embodiments, R^{K1} is substituted phenyl. In certain embodiments, R^{K1} is unsubstituted phenyl. In certain embodiments, R^{K1} is bicyclic aryl. In certain embodiments, R^{K1} is substituted or unsubstituted heteroaryl. In certain embodiments, R^{K1} is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, R^{K1} is monocyclic heteroaryl. In certain embodiments, R^{K1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, R^{K1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, R^{K1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, R^{K1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{K1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{K1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{K1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{K1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulphenyl, 2-pyridine-sulphenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{K1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain

embodiments, two instances of R^{K1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{K1} are joined to form a heterocyclic ring, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{K1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{K1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{K1} are joined to form a substituted or unsubstituted, 5- to 6-membered, monocyclic heteroaryl ring, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.

[00305] Ring E of Formula (VII-B) may include one or more substituents R^J . In certain embodiments, at least two instances of R^J are different. In certain embodiments, all instances of R^J are the same. In certain embodiments, at least one instance of R^J is halogen. In certain embodiments, at least one instance of R^J is F. In certain embodiments, at least one instance of R^J is Cl. In certain embodiments, at least one instance of R^J is Br. In certain embodiments, at least one instance of R^J is I (iodine). In certain embodiments, at least one instance of R^J is substituted alkyl. In certain embodiments, at least one instance of R^J is unsubstituted alkyl. In certain embodiments, at least one instance of R^J is unsubstituted C_{1-6} alkyl. In certain embodiments, all instances of R^J are unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^J is substituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^J is C_{1-6} alkyl substituted with at least one halogen. In certain embodiments, at least one instance of R^J is $-CH_3$. In certain embodiments, all instances of R^J are $-CH_3$. In certain embodiments, at least one instance of R^J is substituted methyl. In certain embodiments, at least one instance of R^J is $-CH_2F$. In certain embodiments, at least one instance of R^J is $-CHF_2$. In certain embodiments, at least one instance of R^J is $-CF_3$. In certain embodiments, at least one instance of R^J is ethyl. In certain embodiments, at least one instance of R^J is propyl. In certain embodiments, at least one instance of R^J is butyl. In certain embodiments, at least one instance of R^J is pentyl. In certain embodiments, at least one instance of R^J is hexyl. In certain embodiments, at least one instance of R^J is halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^J is substituted alkenyl. In certain embodiments, at least one instance of R^J is unsubstituted alkenyl. In certain embodiments, at least one instance of R^J is substituted alkynyl. In certain embodiments, at least one instance of R^J is unsubstituted alkynyl. In certain embodiments, at least one instance of R^J is substituted carbocyclyl. In certain embodiments, at least one instance of R^J is unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^J is saturated carbocyclyl. In certain

embodiments, at least one instance of R^J is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^J is monocyclic carbocyclyl. In certain embodiments, at least one instance of R^J is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^J is substituted heterocyclyl. In certain embodiments, at least one instance of R^J is unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^J is saturated heterocyclyl. In certain embodiments, at least one instance of R^J is unsaturated heterocyclyl. In certain embodiments, at least one instance of R^J is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^J is monocyclic heterocyclyl. In certain embodiments, at least one instance of R^J is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^J is substituted aryl. In certain embodiments, at least one instance of R^J is unsubstituted aryl. In certain embodiments, at least one instance of R^J is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^J is substituted phenyl. In certain embodiments, at least one instance of R^J is unsubstituted phenyl. In certain embodiments, at least one instance of R^J is substituted heteroaryl. In certain embodiments, at least one instance of R^J is unsubstituted heteroaryl. In certain embodiments, at least one instance of R^J is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^J is monocyclic heteroaryl. In certain embodiments, at least one instance of R^J is 5-membered, monocyclic heteroaryl. In certain embodiments, no instance of R^J is substituted imidazolyl. In certain embodiments, at least one instance of R^J is 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^J is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^J is 9- or 10-membered, bicyclic heteroaryl. In certain embodiments, at least one instance of R^J is $-OR^{J1}$. In certain embodiments, at least one instance of R^J is $-OH$. In certain embodiments, at least one instance of R^J is $-O(\text{substituted or unsubstituted } C_{1-6} \text{ alkyl})$. In certain embodiments, at least one instance of R^J is $-OMe$. In certain embodiments, at least one instance of R^J is $-OEt$. In certain embodiments, at least one instance of R^J is $-OPr$. In certain embodiments, at least one instance of R^J is $-OBu$. In certain embodiments, at least one instance of R^J is $-OBn$. In certain embodiments, at least one instance of R^J is $-OPh$. In certain embodiments, at least one instance of R^J is $-SR^{J1}$. In certain embodiments, at least one instance of R^J is $-SH$. In certain embodiments, at least one instance of R^J is $-SMe$. In certain

embodiments, at least one instance of R^J is $-N(R^{J1})_2$. In certain embodiments, at least one instance of R^J is $-NH_2$. In certain embodiments, at least one instance of R^J is $-NHMe$. In certain embodiments, at least one instance of R^J is $-NMe_2$. In certain embodiments, at least one instance of R^J is $-CN$. In certain embodiments, at least one instance of R^J is $-SCN$. In certain embodiments, at least one instance of R^J is $-C(=NR^{J1})R^{J1}$, $-C(=NR^{J1})OR^{J1}$, or $-C(=NR^{J1})N(R^{J1})_2$. In certain embodiments, at least one instance of R^J is $-C(=O)R^{J1}$ or $-C(=O)OR^{J1}$. In certain embodiments, at least one instance of R^J is $-C(=O)N(R^{J1})_2$. In certain embodiments, at least one instance of R^J is $-C(=O)NMe_2$, $-C(=O)NHMe$, or $-C(=O)NH_2$. In certain embodiments, at least one instance of R^J is $-NO_2$. In certain embodiments, at least one instance of R^J is $-NR^{J1}C(=O)R^{J1}$, $-NR^{J1}C(=O)OR^{J1}$, or $-NR^{J1}C(=O)N(R^{J1})_2$. In certain embodiments, at least one instance of R^J is $-OC(=O)R^{J1}$, $-OC(=O)OR^{J1}$, or $-OC(=O)N(R^{J1})_2$.

[00306] In certain embodiments, at least one instance of R^J is halogen or substituted or unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^J is halogen, unsubstituted C_{1-6} alkyl, or C_{1-6} alkyl substituted with at least one halogen.

[00307] In certain embodiments, at least one instance of R^{J1} is H. In certain embodiments, at least one instance of R^{J1} is substituted acyl. In certain embodiments, at least one instance of R^{J1} is unsubstituted acyl. In certain embodiments, at least one instance of R^{J1} is acetyl. In certain embodiments, at least one instance of R^{J1} is substituted alkyl. In certain embodiments, at least one instance of R^{J1} is unsubstituted alkyl. In certain embodiments, at least one instance of R^{J1} is unsubstituted C_{1-6} alkyl. In certain embodiments, at least one instance of R^{J1} is methyl. In certain embodiments, at least one instance of R^{J1} is ethyl. In certain embodiments, at least one instance of R^{J1} is propyl. In certain embodiments, at least one instance of R^{J1} is butyl. In certain embodiments, at least one instance of R^{J1} is pentyl. In certain embodiments, at least one instance of R^{J1} is hexyl. In certain embodiments, at least one instance of R^{J1} is substituted alkenyl. In certain embodiments, at least one instance of R^{J1} is unsubstituted alkenyl. In certain embodiments, at least one instance of R^{J1} is substituted alkynyl. In certain embodiments, at least one instance of R^{J1} is unsubstituted alkynyl. In certain embodiments, at least one instance of R^{J1} is substituted or unsubstituted carbocyclyl. In certain embodiments, at least one instance of R^{J1} is saturated carbocyclyl. In certain embodiments, at least one instance of R^{J1} is unsaturated carbocyclyl. In certain embodiments, at least one instance of R^{J1} is 3- to 7-membered, monocyclic carbocyclyl. In certain embodiments, at least one instance of R^{J1} is substituted or unsubstituted heterocyclyl. In certain embodiments, at least one instance of R^{J1} is saturated heterocyclyl. In certain embodiments, at least one instance of R^{J1} is unsaturated heterocyclyl. In certain

embodiments, at least one instance of R^{J1} is heterocyclyl, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{J1} is 3- to 7-membered, monocyclic heterocyclyl. In certain embodiments, at least one instance of R^{J1} is substituted or unsubstituted aryl. In certain embodiments, at least one instance of R^{J1} is 6- to 10-membered aryl. In certain embodiments, at least one instance of R^{J1} is monocyclic aryl. In certain embodiments, at least one instance of R^{J1} is substituted phenyl. In certain embodiments, at least one instance of R^{J1} is unsubstituted phenyl. In certain embodiments, at least one instance of R^{J1} is bicyclic aryl. In certain embodiments, at least one instance of R^{J1} is substituted or unsubstituted heteroaryl. In certain embodiments, at least one instance of R^{J1} is heteroaryl, wherein one, two, three, or four atoms of the heteroaryl ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, at least one instance of R^{J1} is monocyclic heteroaryl. In certain embodiments, at least one instance of R^{J1} is 5- or 6-membered, monocyclic heteroaryl. In certain embodiments, at least one instance of R^{J1} is bicyclic heteroaryl, wherein the point of attachment may be on any atom of the bicyclic heteroaryl ring system, as valency permits. In certain embodiments, at least one instance of R^{J1} is a nitrogen protecting group when attached to a nitrogen atom. In certain embodiments, at least one instance of R^{J1} is Bn, Boc, Cbz, Fmoc, trifluoroacetyl, triphenylmethyl, acetyl, or Ts when attached to a nitrogen atom. In certain embodiments, R^{J1} is an oxygen protecting group when attached to an oxygen atom. In certain embodiments, R^{J1} is silyl, TBDPS, TBDMS, TIPS, TES, TMS, MOM, THP, *t*-Bu, Bn, allyl, acetyl, pivaloyl, or benzoyl when attached to an oxygen atom. In certain embodiments, R^{J1} is a sulfur protecting group when attached to a sulfur atom. In certain embodiments, R^{J1} is acetamidomethyl, *t*-Bu, 3-nitro-2-pyridine sulfenyl, 2-pyridine-sulfenyl, or triphenylmethyl when attached to a sulfur atom. In certain embodiments, two instances of R^{J1} are joined to form a substituted or unsubstituted heterocyclic ring. In certain embodiments, two instances of R^{J1} are joined to form a saturated or unsaturated heterocyclic ring. In certain embodiments, two instances of R^{J1} are joined to form a heterocyclic ring, wherein one, two, or three atoms of the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur. In certain embodiments, two instances of R^{J1} are joined to form a 3- to 7-membered, monocyclic heterocyclic ring. In certain embodiments, two instances of R^{J1} are joined to form a substituted or unsubstituted heteroaryl ring. In certain embodiments, two instances of R^{J1} are joined to form a substituted or unsubstituted, 5- to 6-membered,





monocyclic heteroaryl ring, wherein one, two, three, or four atoms of the heteroaryl ring system are independently nitrogen, oxygen, or sulfur.







[00308] In certain embodiments, q is 0. In certain embodiments, q is 1. In certain embodiments, q is 2. In certain embodiments, q is 3. In certain embodiments, q is 4.

[00309] In certain embodiments, no instance of R^J and R^K is substituted or unsubstituted heteroaryl. In certain embodiments, no instance of R^J and R^K is substituted or unsubstituted imidazolyl. In certain embodiments, no instance of R^J and R^K is substituted imidazolyl.

[00310] In certain embodiments, the compound of Formula (VII) is of the formula:


[00311] In certain embodiments, the compound of Formula (VII) is of the formula:







or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof.

[00312] In certain embodiments, a compound of Formula (VII) is not of the formula:





[00313] In certain embodiments, the compound utilized in the present disclosure is of Formula (VII):



or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, or tautomer thereof. each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{A1}$, $-N(R^{A1})_2$, $-SR^{A1}$, $-CN$, $-SCN$, $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, $-C(=NR^{A1})N(R^{A1})_2$, $-C(=O)R^{A1}$, $-C(=O)OR^{A1}$, $-C(=O)N(R^{A1})_2$, $-NO_2$, $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$, $-NR^{A1}C(=O)N(R^{A1})_2$, $-OC(=O)R^{A1}$, $-OC(=O)OR^{A1}$, or $-OC(=O)N(R^{A1})_2$, wherein each instance of R^{A1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when

attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{A1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

j is 0, 1, or 2;

R^B is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or a nitrogen protecting group;

R^C is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^D is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{D1}$, $-N(R^{D1})_2$, $-SR^{D1}$, $-CN$, $-SCN$, $-C(=NR^{D1})R^{D1}$, $-C(=NR^{D1})OR^{D1}$, $-C(=NR^{D1})N(R^{D1})_2$, $-C(=O)R^{D1}$, $-C(=O)OR^{D1}$, $-C(=O)N(R^{D1})_2$, $-NO_2$, $-NR^{D1}C(=O)R^{D1}$, $-NR^{D1}C(=O)OR^{D1}$, $-NR^{D1}C(=O)N(R^{D1})_2$, $-OC(=O)R^{D1}$, $-OC(=O)OR^{D1}$, or $-OC(=O)N(R^{D1})_2$, wherein each instance of R^{D1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

m is 0, 1, or 2;


R^E is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;


each instance of R^F is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{F1}$, $-N(R^{F1})_2$, $-SR^{F1}$, $-CN$, $-SCN$, $-C(=NR^{F1})R^{F1}$, $-C(=NR^{F1})OR^{F1}$, $-C(=NR^{F1})N(R^{F1})_2$, $-C(=O)R^{F1}$, $-C(=O)OR^{F1}$,

$-\text{C}(=\text{O})\text{N}(\text{R}^{\text{F1}})_2$, $-\text{NO}_2$, $-\text{NR}^{\text{F1}}\text{C}(=\text{O})\text{R}^{\text{F1}}$, $-\text{NR}^{\text{F1}}\text{C}(=\text{O})\text{OR}^{\text{F1}}$, $-\text{NR}^{\text{F1}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{F1}})_2$, $-\text{OC}(=\text{O})\text{R}^{\text{F1}}$, $-\text{OC}(=\text{O})\text{OR}^{\text{F1}}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{F1}})_2$, wherein each instance of R^{F1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{F1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring; and

n is 0, 1, 2, 3, 4, or 5;

provided that the compound is not of the formula:





or a pharmaceutically acceptable salt thereof.

Pharmaceutical Compositions and Administration

[00314] In certain embodiments, a SIK inhibitor for use in the invention described herein is a compound of Formula (I), (II), (III), (IV), (V), (VI), (VI-A), or (VII), or a pharmaceutically acceptable salt, solvate, hydrate, polymorph, co-crystal, tautomer, stereoisomer, isotopically labeled derivative, or prodrug thereof. In certain embodiments, the compound described herein is provided in an effective amount in the pharmaceutical composition. In certain embodiments, the effective amount is a prophylactically effective amount. In certain embodiments, the effective amount is an amount effective for increasing skin pigmentation in a subject in need thereof. In certain embodiments, the effective amount is an amount effective for increasing skin pigmentation in a subject in need thereof for

cosmetic purposes. In certain embodiments, the effective amount is an amount effective for increasing skin pigmentation in a subject in need thereof for treating polymorphic light eruption (*e.g.*, sun hypersensitivity). In certain embodiments, the effective amount is an amount effective for reducing the risk of skin cancer in a subject in need thereof. In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK, (*e.g.*, SIK1, SIK2, or SIK3)) in a subject or cell (*e.g.*, the skin of a subject or in skin cells).

[00315] In certain embodiments, the subject being administered a compound or composition described herein is an animal. The animal may be of either sex and may be at any stage of development. In certain embodiments, the subject described herein is a human. In certain embodiments, the subject is a non-human animal. In certain embodiments, the subject is a mammal.

[00316] An effective amount of a compound described herein may vary from about 0.001 mg/kg to about 1000 mg/kg in one or more dose administrations for one or several days (depending on the mode of administration), wherein mg/kg is mg of compound to kg weight of the subject. In certain embodiments, the effective amount per dose varies from about 0.001 mg/kg to about 1000 mg/kg, from about 0.01 mg/kg to about 750 mg/kg, from about 0.1 mg/kg to about 500 mg/kg, from about 1.0 mg/kg to about 250 mg/kg, and from about 10.0 mg/kg to about 150 mg/kg.

[00317] In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK by at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 98%. In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK by not more than 10%, not more than 20%, not more than 30%, not more than 40%, not more than 50%, not more than 60%, not more than 70%, not more than 80%, not more than 90%, not more than 95%, or not more than 98%. In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK by a range between a percentage described in this paragraph and another percentage described in this paragraph, inclusive.

[00318] Pharmaceutical compositions described herein can be prepared by any method known in the art of pharmacology. In general, such preparatory methods include bringing the compound described herein (*i.e.*, the “active ingredient”) into association with a carrier or excipient, and/or one or more other accessory ingredients, and then, if necessary and/or desirable, shaping, and/or packaging the product into a desired single- or multi-dose unit.

[00319] Pharmaceutical compositions can be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. A “unit dose” is a discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage, such as one-half or one-third of such a dosage.

[00320] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition described herein will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. The composition may comprise between 0.1% and 100% (w/w) active ingredient.

[00321] Pharmaceutically acceptable excipients used in the manufacture of provided pharmaceutical compositions include inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and perfuming agents may also be present in the composition.

[00322] Exemplary diluents include calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, and mixtures thereof.

[00323] Exemplary granulating and/or dispersing agents include potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose, and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum), sodium lauryl sulfate, quaternary ammonium compounds, and mixtures thereof.

[00324] Exemplary surface active agents and/or emulsifiers include natural emulsifiers (*e.g.*, acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (*e.g.*, bentonite (aluminum silicate) and Veegum (magnesium aluminum silicate)), long chain

amino acid derivatives, high molecular weight alcohols (*e.g.*, stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (*e.g.*, carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (*e.g.*, carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (*e.g.*, polyoxyethylene sorbitan monolaurate (Tween[®] 20), polyoxyethylene sorbitan (Tween[®] 60), polyoxyethylene sorbitan monooleate (Tween[®] 80), sorbitan monopalmitate (Span[®] 40), sorbitan monostearate (Span[®] 60), sorbitan tristearate (Span[®] 65), glyceryl monooleate, sorbitan monooleate (Span[®] 80), polyoxyethylene esters (*e.g.*, polyoxyethylene monostearate (Myrj[®] 45), polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and Solutol[®]), sucrose fatty acid esters, polyethylene glycol fatty acid esters (*e.g.*, Cremophor[®]), polyoxyethylene ethers, (*e.g.*, polyoxyethylene lauryl ether (Brij[®] 30)), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, Pluronic[®] F-68, poloxamer P-188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, and/or mixtures thereof.

[00325] Exemplary binding agents include starch (*e.g.*, cornstarch and starch paste), gelatin, sugars (*e.g.*, sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol, *etc.*), natural and synthetic gums (*e.g.*, acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum[®]), and larch arabogalactan), alginates, polyethylene oxide, polyethylene glycol, inorganic calcium salts, silicic acid, polymethacrylates, waxes, water, alcohol, and/or mixtures thereof.

[00326] Exemplary preservatives include antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, antiprotozoan preservatives, alcohol preservatives, acidic preservatives, and other preservatives. In certain embodiments, the preservative is an antioxidant. In other embodiments, the preservative is a chelating agent.

[00327] Exemplary antioxidants include alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and sodium sulfite.

[00328] Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA) and salts and hydrates thereof (*e.g.*, sodium edetate, disodium edetate, trisodium edetate, calcium disodium edetate, dipotassium edetate, and the like), citric acid and salts and hydrates thereof (*e.g.*, citric acid monohydrate), fumaric acid and salts and hydrates thereof, malic acid and salts and hydrates thereof, phosphoric acid and salts and hydrates thereof, and tartaric acid and salts and hydrates thereof. Exemplary antimicrobial preservatives include benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and thimerosal.

[00329] Exemplary antifungal preservatives include butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and sorbic acid.

[00330] Exemplary alcohol preservatives include ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and phenylethyl alcohol.

[00331] Exemplary acidic preservatives include vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and phytic acid.

[00332] Other preservatives include tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, Glydant[®] Plus, Phenonip[®], methylparaben, Germall[®] 115, Germaben[®] II, Neolone[®], Kathon[®], and Euxyl[®].

[00333] Exemplary buffering agents include citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, and mixtures thereof.

[00334] Exemplary lubricating agents include magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behanate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, and mixtures thereof.

[00335] Exemplary natural oils include almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary synthetic oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and mixtures thereof.

[00336] Liquid dosage forms for oral and parenteral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredients, the liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (*e.g.*, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents. In certain embodiments for parenteral administration, the conjugates described herein are mixed with solubilizing agents such as Cremophor[®], alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and mixtures thereof.

[00337] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or (a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders such as, for example, carboxymethylcellulose,

alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, (c) humectants such as glycerol, (d) disintegrating agents such as agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, (e) solution retarding agents such as paraffin, (f) absorption accelerators such as quaternary ammonium compounds, (g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, (h) absorbents such as kaolin and bentonite clay, and (i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets, and pills, the dosage form may include a buffering agent.

[00338] Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the art of pharmacology. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of encapsulating compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00339] The active ingredient can be in a micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings, and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active ingredient can be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may comprise buffering agents. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of encapsulating agents which can be used include polymeric substances and waxes.

[00340] Dosage forms for topical and/or transdermal administration of a compound described herein may include ointments, pastes, creams, lotions, gels, powders, solutions,

sprays, inhalants, and/or patches. Generally, the active ingredient is admixed under sterile conditions with a pharmaceutically acceptable carrier or excipient and/or any needed preservatives and/or buffers as can be required. Additionally, the present disclosure contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of an active ingredient to the body. Such dosage forms can be prepared, for example, by dissolving and/or dispensing the active ingredient in the proper medium. Alternatively or additionally, the rate can be controlled by either providing a rate controlling membrane and/or by dispersing the active ingredient in a polymer matrix and/or gel.

[00341] Suitable devices for use in delivering intradermal pharmaceutical compositions described herein include short needle devices. Intradermal compositions can be administered by devices which limit the effective penetration length of a needle into the skin. Alternatively or additionally, conventional syringes can be used in the classical mantoux method of intradermal administration. Jet injection devices which deliver liquid formulations to the dermis *via* a liquid jet injector and/or *via* a needle which pierces the stratum corneum and produces a jet which reaches the dermis are suitable. Ballistic powder/particle delivery devices which use compressed gas to accelerate the compound in powder form through the outer layers of the skin to the dermis are suitable.

[00342] Formulations suitable for topical administration include, but are not limited to, liquid and/or semi-liquid preparations such as liniments, lotions, oil-in-water and/or water-in-oil emulsions such as creams, ointments, and/or pastes, and/or solutions and/or suspensions. Topically administrable formulations may, for example, comprise from about 1% to about 10% (w/w) active ingredient, although the concentration of the active ingredient can be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical administration may further comprise one or more of the additional ingredients described herein.

[00343] A pharmaceutical composition described herein can be prepared, packaged, and/or sold in a formulation suitable for pulmonary administration *via* the buccal cavity. Such a formulation may comprise dry particles which comprise the active ingredient and which have a diameter in the range from about 0.5 to about 7 nanometers, or from about 1 to about 6 nanometers. Such compositions are conveniently in the form of dry powders for administration using a device comprising a dry powder reservoir to which a stream of propellant can be directed to disperse the powder and/or using a self-propelling solvent/powder dispensing container such as a device comprising the active ingredient

dissolved and/or suspended in a low-boiling propellant in a sealed container. Such powders comprise particles wherein at least 98% of the particles by weight have a diameter greater than 0.5 nanometers and at least 95% of the particles by number have a diameter less than 7 nanometers. Alternatively, at least 95% of the particles by weight have a diameter greater than 1 nanometer and at least 90% of the particles by number have a diameter less than 6 nanometers. Dry powder compositions may include a solid fine powder diluent such as sugar and are conveniently provided in a unit dose form.

[00344] Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant may constitute 50 to 99.9% (w/w) of the composition, and the active ingredient may constitute 0.1 to 20% (w/w) of the composition. The propellant may further comprise additional ingredients such as a liquid non-ionic and/or solid anionic surfactant and/or a solid diluent (which may have a particle size of the same order as particles comprising the active ingredient).

[00345] Although the descriptions of pharmaceutical compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to animals of all sorts. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with ordinary experimentation.

[00346] Compounds provided herein are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions described herein will be decided by a physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject or organism will depend upon a variety of factors including the disease being treated and the severity of the disorder; the activity of the specific active ingredient employed; the specific composition employed; the age, body weight, general health, sex, and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific active ingredient employed; the duration of the treatment; drugs used in combination or coincidental with the specific active ingredient employed; and like factors well known in the medical arts.

[00347] The compounds and compositions provided herein can be administered by any route, including enteral (*e.g.*, oral), parenteral, intravenous, intramuscular, intra-arterial,

intramedullary, intrathecal, subcutaneous, intraventricular, transdermal, interdermal, rectal, intravaginal, intraperitoneal, topical (as by powders, ointments, creams, and/or drops), mucosal, nasal, bucal, sublingual; by intratracheal instillation, bronchial instillation, and/or inhalation; and/or as an oral spray, nasal spray, and/or aerosol. Specifically contemplated routes are oral administration, intravenous administration (*e.g.*, systemic intravenous injection), regional administration *via* blood and/or lymph supply, and/or direct administration to an affected site. In general, the most appropriate route of administration will depend upon a variety of factors including the nature of the agent (*e.g.*, its stability in the environment of the gastrointestinal tract), and/or the condition of the subject (*e.g.*, whether the subject is able to tolerate oral administration). In certain embodiments, the compound or pharmaceutical composition described herein is suitable for topical administration to the skin of a subject. In certain embodiments, the compound or pharmaceutical composition described herein is suitable for transdermal administration to the skin of a subject. In certain embodiments, the compound or pharmaceutical composition described herein is suitable for administration to the skin cells of a subject.

[00348] The exact amount of a compound required to achieve an effective amount will vary from subject to subject, depending, for example, on species, age, and general condition of a subject, severity of the side effects or disorder, identity of the particular compound, mode of administration, and the like. An effective amount may be included in a single dose (*e.g.*, single oral dose) or multiple doses (*e.g.*, multiple oral doses). In certain embodiments, when multiple doses are administered to a subject or applied to a tissue or cell, any two doses of the multiple doses include different or substantially the same amounts of a compound described herein. In certain embodiments, when multiple doses are administered to a subject or applied to a tissue or cell, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is three doses a day, two doses a day, one dose a day, one dose every other day, one dose every third day, one dose every week, one dose every two weeks, one dose every three weeks, or one dose every four weeks. In certain embodiments, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is one dose per day. In certain embodiments, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is two doses per day. In certain embodiments, the frequency of administering the multiple doses to the subject or applying the multiple doses to the tissue or cell is three doses per day. In certain embodiments, when multiple doses are administered to a subject or applied to a tissue or cell, the duration between the first dose and last dose of the multiple doses is one

day, two days, four days, one week, two weeks, three weeks, one month, two months, three months, four months, six months, nine months, one year, two years, three years, four years, five years, seven years, ten years, fifteen years, twenty years, or the lifetime of the subject, tissue, or cell. In certain embodiments, the duration between the first dose and last dose of the multiple doses is three months, six months, or one year. In certain embodiments, the duration between the first dose and last dose of the multiple doses is the lifetime of the subject, tissue, or cell. In certain embodiments, a dose (*e.g.*, a single dose, or any dose of multiple doses) described herein includes independently between 0.1 μg and 1 μg , between 0.001 mg and 0.01 mg, between 0.01 mg and 0.1 mg, between 0.1 mg and 1 mg, between 1 mg and 3 mg, between 3 mg and 10 mg, between 10 mg and 30 mg, between 30 mg and 100 mg, between 100 mg and 300 mg, between 300 mg and 1,000 mg, or between 1 g and 10 g, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 1 mg and 3 mg, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 3 mg and 10 mg, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 10 mg and 30 mg, inclusive, of a compound described herein. In certain embodiments, a dose described herein includes independently between 30 mg and 100 mg, inclusive, of a compound described herein.

[00349] Dose ranges as described herein provide guidance for the administration of provided pharmaceutical compositions to an adult. The amount to be administered to, for example, a child or an adolescent can be determined by a medical practitioner or person skilled in the art and can be lower or the same as that administered to an adult. In certain embodiments, a dose described herein is a dose to an adult human whose body weight is 70 kg.

[00350] A SIK inhibitor or composition, as described herein, can be administered in combination with one or more additional pharmaceutical agents (*e.g.*, therapeutically and/or prophylactically active agents) useful in increasing skin pigmentation and/or reducing the risk of skin cancer. The compounds or compositions can be administered in combination with additional pharmaceutical agents that improve their activity (*e.g.*, activity (*e.g.*, potency and/or efficacy) in increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof, and/or in inhibiting the activity of SIK in a subject), improve bioavailability, improve safety, reduce drug resistance, reduce and/or modify metabolism, inhibit excretion, and/or modify distribution in a subject or cell. It will also be appreciated that the therapy employed may achieve a desired effect for the same disorder, and/or it may

achieve different effects. In certain embodiments, a pharmaceutical composition described herein including a SIK inhibitor described herein and an additional pharmaceutical agent shows a synergistic effect that is absent in a pharmaceutical composition including one of the SIK inhibitor and the additional pharmaceutical agent, but not both.

[00351] The SIK inhibitor or composition can be administered concurrently with, prior to, or subsequent to one or more additional pharmaceutical agents, which are different from the compound or composition and may be useful as, *e.g.*, combination therapies in increasing skin pigmentation and/or reducing the risk of skin cancer. Pharmaceutical agents include therapeutically active agents. Pharmaceutical agents also include prophylactically active agents. Pharmaceutical agents include small organic molecules such as drug compounds (*e.g.*, compounds approved for human or veterinary use by the U.S. Food and Drug Administration as provided in the Code of Federal Regulations (CFR)), peptides, proteins, carbohydrates, monosaccharides, oligosaccharides, polysaccharides, nucleoproteins, mucoproteins, lipoproteins, synthetic polypeptides or proteins, small molecules linked to proteins, glycoproteins, steroids, nucleic acids, DNAs, RNAs, nucleotides, nucleosides, oligonucleotides, antisense oligonucleotides, lipids, hormones, vitamins, and cells. In certain embodiments, the additional pharmaceutical agent is a pharmaceutical agent useful for increasing skin pigmentation and/or reducing the risk of skin cancer. Each additional pharmaceutical agent may be administered at a dose and/or on a time schedule determined for that pharmaceutical agent. The additional pharmaceutical agents may also be administered together with each other and/or with the SIK inhibitor or composition described herein in a single dose or administered separately in different doses. The particular combination to employ in a regimen will take into account compatibility of the SIK inhibitor described herein with the additional pharmaceutical agent(s) and/or the desired therapeutic and/or prophylactic effect to be achieved. In general, it is expected that the additional pharmaceutical agent(s) in combination be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

[00352] The additional pharmaceutical agents include, but are not limited to, anti-proliferative agents, anti-musculoskeletal disease agents, anti-cancer agents, cytotoxic agents, anti-angiogenesis agents, anti-inflammatory agents, immunosuppressants, anti-bacterial agents, anti-viral agents, cardiovascular agents, cholesterol-lowering agents, anti-diabetic agents, anti-allergic agents, contraceptive agents, and pain-relieving agents. In certain embodiments, the additional pharmaceutical agent is an anti-proliferative agent. In certain

embodiments, the additional pharmaceutical agent is an anti- musculoskeletal disease agent. In certain embodiments, the additional pharmaceutical agent is an anti-cancer agent. In certain embodiments, the additional pharmaceutical agent is an anti-viral agent. In certain embodiments, the additional pharmaceutical agent is an binder or inhibitor of a protein kinase. In certain embodiments, the additional pharmaceutical agent is selected from the group consisting of epigenetic or transcriptional modulators (*e.g.*, DNA methyltransferase inhibitors, histone deacetylase inhibitors (HDAC inhibitors), lysine methyltransferase inhibitors), antimitotic drugs (*e.g.*, taxanes and vinca alkaloids), hormone receptor modulators (*e.g.*, estrogen receptor modulators and androgen receptor modulators), cell signaling pathway inhibitors (*e.g.*, tyrosine protein kinase inhibitors), modulators of protein stability (*e.g.*, proteasome inhibitors), Hsp90 inhibitors, glucocorticoids, all-*trans* retinoic acids, and other agents that promote differentiation. In certain embodiments, the compounds described herein or pharmaceutical compositions can be administered in combination with an anti-cancer therapy including, but not limited to, surgery, radiation therapy, transplantation (*e.g.*, stem cell transplantation, bone marrow transplantation), immunotherapy, and chemotherapy. In certain embodiments, the additional pharmaceutical agent is an agent for increasing skin pigmentation in a subject. In certain embodiments, the additional pharmaceutical agent is an agent for reducing the risk of skin cancer in a subject. In certain embodiments, the additional pharmaceutical agent is an agent for inducing eumelanin synthesis, inducing melanosomal maturation, export, and localization, or inducing microphthalmia-associated transcription factor (MITF) expression.

Methods of Treatment and Uses

[00353] The present disclosure provides methods of inhibiting the activity of SIK (*e.g.*, SIK1, SIK2, or SIK3) in a subject or cell. The present disclosure also provides methods for increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof. The present disclosure provides methods for increasing skin pigmentation for cosmetic purposes. In certain embodiments, provided herein are methods of increasing the appearance of skin darkening in a subject in need thereof using the described compounds (*e.g.*, via topical administration of the described compounds). In certain embodiments, the present disclosure provides methods of increasing the appearance of skin pigmentation in a subject, the methods comprising administering topically to the subject's skin an effective amount of a salt inducible kinase (SIK) inhibitor of any one of Formulae (I), (II), (III), (IV),

(V), (VI), (VI-A), and (VII), or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, or stereoisomer thereof, or a pharmaceutical composition thereof.

[00354] In certain embodiments, the present disclosure provides methods of treating polymorphic light eruption (*e.g.*, sun hypersensitivity). The present disclosure provides methods of inducing eumelanin synthesis. The present disclosure provides methods of inducing melanosomal maturation, export, and localization. In certain embodiments, the present disclosure provides methods of inducing (*e.g.*, temporarily and reversibly inducing) microphthalmia-associated transcription factor (MITF) expression. In certain embodiments, the present disclosure provides methods of inducing increased microphthalmia-associated transcription factor (MITF) expression. In certain embodiments, inducing MITF expression comprises temporarily and reversibly inducing MITF expression.

[00355] In another aspect, the present disclosure provides methods of inhibiting the activity of SIK, (*e.g.*, SIK1, SIK2, or SIK3) in a subject. In certain embodiments, the activity of a SIK, (*e.g.*, SIK1, SIK2, or SIK3) in a subject is inhibited by a method described herein by at least about 1%, at least about 3%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90%. In some embodiments, the activity of SIK, (*e.g.*, SIK1, SIK2, or SIK3) in a subject is selectively inhibited by the method. In some embodiments, the activity of SIK2 in a subject is selectively inhibited by the method. In another aspect, the present disclosure provides methods of inhibiting SIK in the skin of a subject.

[00356] Another aspect of the present disclosure relates to methods of treating a disease associated with SIK, (*e.g.*, SIK1, SIK2, or SIK3).

[00357] In still another aspect, the present disclosure provides methods of preventing skin cancer described herein in a subject in need thereof.

[00358] In another aspect, the present disclosure provides methods of increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof, the method comprising administering topically to the subject's skin an effective amount of a salt inducible kinase (SIK) inhibitor of any one of Formulae (I), (II), (III), (IV), (V), (VI), (VI-A), and (VII), or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, or stereoisomer thereof, or a pharmaceutical composition thereof. In certain embodiments, provided are methods of reversibly increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof, the method comprising administering topically to the subject's skin an effective amount of a SIK inhibitor of compounds described herein. In

certain embodiments, increasing skin pigmentation comprises reversibly increasing skin pigmentation.

[00359] In certain embodiments, the methods of the disclosure include administering to a subject in need thereof an effective amount of a compound or pharmaceutical composition described herein. In certain embodiments, the subject being administered a compound or pharmaceutical composition described herein is a human. In certain embodiments, the subject being administered a compound or pharmaceutical composition described herein is a non-human animal. In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK by at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 98%. In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK by not more than 10%, not more than 20%, not more than 30%, not more than 40%, not more than 50%, not more than 60%, not more than 70%, not more than 80%, not more than 90%, not more than 95%, or not more than 98%. In certain embodiments, the effective amount is an amount effective for inhibiting the activity of a SIK by a range between a percentage described in this paragraph and another percentage described in this paragraph, inclusive.

[00360] In certain embodiments, the methods of the disclosure include administering to a subject in need thereof an effective amount of a compound or pharmaceutical composition described herein. In certain embodiments, the methods of the disclosure include topically administering to a subject in need thereof an effective amount of a compound or pharmaceutical composition described herein. In certain embodiments, the methods of the disclosure include administering to a subject in need thereof a prophylactically effective amount of a compound or pharmaceutical composition described herein. In certain embodiments, the methods of the disclosure include topically administering to a subject in need thereof a prophylactically effective amount of a compound or pharmaceutical composition described herein.

[00361] In another aspect, the present disclosure provides the compounds described herein for use in a method described herein (*e.g.*, method of inhibiting SIK, (*e.g.*, SIK1, SIK2, or SIK3)), or a method of increasing skin pigmentation and/or reducing the risk of skin cancer in a subject in need thereof. The compounds provided may be used in a method of inducing a skin condition in a subject in need thereof. In certain embodiments, provided is a method of increasing skin pigmentation by administering the compounds described herein. In certain embodiments, provided is a method of increasing skin pigmentation and/or reducing the risk

of skin cancer by topically administering the compounds described herein to a subject's skin on a body part. In certain embodiments, the body part is the face of the subject. In certain embodiments, the body part is the neck of the subject. In certain embodiments, the body part is the chest of the subject. In certain embodiments, the body part is the back of the subject. In certain embodiments, the skin on the body part is skin on the arms of the subject. In certain embodiments, the skin on the body part is skin on the legs of the subject. In certain embodiments, the skin is on the torso of the subject. In another aspect, the present disclosure provides methods of inhibiting SIK in the skin of a subject. In another aspect, the present disclosure provides methods of inducing eumelanin synthesis. In certain embodiments, the present disclosure provides methods of inducing melanosomal maturation, export, and localization. In certain embodiments, the present disclosure provides methods of inducing microphthalmia-associated transcription factor (MITF) expression. In certain embodiments, the present disclosure provides methods of inducing increased microphthalmia-associated transcription factor (MITF) expression.

[00362] In still another aspect, the present disclosure provides the pharmaceutical compositions described herein for use in a method described herein (*e.g.*, a method of inhibiting SIK, (*e.g.*, SIK1, SIK2, or SIK3)), or method of increasing skin pigmentation in a subject in need thereof. In still another aspect, the present disclosure provides the compounds described herein, or a topical formulation thereof, for use in a method of inducing a condition (*e.g.*, increasing skin pigmentation) in a subject in need thereof.

EXAMPLES

[00363] In order that the disclosure may be more fully understood, the following examples are set forth. The examples described in this application are offered to illustrate the compounds, pharmaceutical compositions, uses, and methods provided herein and are not to be construed in any way as limiting their scope.

Methods

[00364] *Materials.* Salt inducible kinase inhibitors were dissolved in 30% propylene glycol + 70% ethanol.

[00365] *Kinome Profiling.* Kinome profiling was performed using KinomeScan ScanMAX at compound concentration of 1 μ M. Protocols are available from DiscoverX®.

[00366] *Kinase activity in vitro assay.* The biochemical activities against SIK2 were measured by Caliper-based mobility shift assay (PerkinElmer®). For these experiments, full

length His6-MBP-tagged hSIK2 (4 nM) was incubated with HG-9-91-01 derivatives in buffer containing 100 mM HEPES 7.5, 10 mM MgCl₂, 2.5 mM DTT, 0.004% Tween20, 0.003% Brij-35, 30 μM ATP and 1.5 μM ProfilerPro FL-Peptide 10 (5-FAMKKKVSRSGLYRSPSPENLNRPR-COOH (SEQ ID NO: 1), PerkinElmer®, Catalog No. 760354) at room temperature. Reactions were quenched by adding 20 mM EDTA (pH 8) after 1 hour, and percentage of substrate conversion was measured by LabChip EZ Reader II (PerkinElmer®). IC₅₀'s for SIK2 inhibition were calculated using SmartFit nonlinear regression in Genedata® Screener software suite (Genedata®).

[00367] All other *in vitro* kinase assay were conducted using the SelectScreen Kinase Profiling Service at Thermo Fisher Scientific® (Madison, WI). The protocols are available from Thermo Fisher Scientific® website.

[00368] *Cell Culture.* UACC257 and UACC62 cells were grown in RPMI + 1% penicillin and streptomycin + 5% fetal bovine serum. Normal human melanocytes were grown in TIVA, and starved 24 hours in HAM's F-12 + 1% penicillin and streptomycin before all molecular experiments.

[00369] *qPCR.* mRNA was extracted from cells using the RNeasy Mini Kit (Qiagen®). KAPA SYBR® FAST Universal One-Step qRT-PCR Kit (KAPA BIOSYSTEMS®) was used to prepare mRNA samples for qPCR and samples were analyzed using the 7500 Fast Real Time PCR System (Applied Biosystems®). The relative expression of each gene was calculated by 7500 Fast System Software program, which utilizes *Ct* normalized to mRNA levels of RPL11 to calculate relative expression. Results are reported relative to control cells.

[00370] *Cell pelleting experiments.* 1x10⁵ UACC257 cells were plated per well in a 6-well plate. Cells were treated daily (1x/day) with 4 μM of SIK inhibitors HG 9-91-01, YKL 06-061, or YKL 06-062 or vehicle control. After 3 days, cells were trypsinized with 0.25% trypsin, resuspended in RPMI media, and pelleted. Pellets were washed 1x with PBS and allowed to dry before imaging.

[00371] *Mice.* C57BL/6J *Mc1r^{ele}* mice were crossed with K14-Scf transgenic mice. C57BL/6J *Tyr^{c2j/c2j}* were crossed with K14-Scf transgenic mice. (Kunisada *et al.* 1998; D'Orazio *et al.* 2006).

[00372] *Photos.* Photos were taken using a Nikon® D50 DSLR with a Nikon® Nikkor 40 mm f/2.8 DX G AF-S. Shutter speed ranged from 1/40 to 1/250 and aperture ranged from F3-F8. Ott-Lite Model L139AB were used to create uniform lighting for photos.

[00373] *Mouse pigmentation experiments.* Animals were treated with vehicle control or

37.5 mM HG 9-9-01 until uniform gross darkening was visible as stated in figure legends. Daily differences in darkening of the skin were measured by reflective colorimetry. Skin was harvested, fixed, and processed for paraffin embedding. Sections were cut from paraffin blocks, and sections were stained utilizing hematoxylin and eosin (morphology) and Fontana-Masson (melanin). Melanin dissolution was conducted utilizing NaOH lysis method (Wakamatsu and Ito 2002).

[00374] *Human pigmentation experiments.* Full thickness human breast skin explants were cultured in petri dishes with a solid phase and liquid phase phenol red free DMEM medium with 20% penicillin streptomycin, 5% fungizone (Gibco®), and 10% FBS. Explants were treated daily with vehicle or SIK inhibitor as indicated in figure legends. Passive application refers to simply applying the treatment to skin without further rubbing or manipulation. Mechanical application refers to application of agents to skin with further rubbing of treatment with 10 clockwise turns of a gloved cotton swab applicator. Skin was harvested, fixed, and processed for paraffin embedding. Sections were cut from paraffin blocks, and sections were stained utilizing hematoxylin and eosin (morphology) and Fontana-Masson (melanin).


[00375] *Colorimeter Measurements:* Differences in darkening of the skin were measured by reflective colorimetry (CIE L* white-black color axis) utilizing a CR-400 Colorimeter (Minolta®) calibrated by a calibration plate (C: Y 93.1, x .3133, y .3194) before each set of measurements.

[00376] *Statistical Analysis.* Data are presented as the mean $\pm$ SD. Statistical significance of differences between experimental groups for dose curve experiments were assessed by one-way ANOVA with Dunnet post test. Statistical significance of differences between experimental groups for time course experiments were assessed by two-way ANOVA with Dunnet post test. Multiplicity adjusted P values were reported for each comparison, and differences of means were considered significant if $P < 0.05$.

Example I. Preparation and Characterization of Exemplary Compounds

[00377] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Proton nuclear magnetic resonance spectra (^1H NMR) were obtained on Bruker AVANCE spectrometer at 400 MHz for proton. Spectra are given in ppm (δ) and coupling constants, J , are reported in Hertz. The solvent peak was used as the reference peak for proton spectra. LC-MS spectra were obtained on Agilent 1100 HPLC LC-MS ion trap electrospray ionization (ESI) mass spectrometer.

Synthesis of YKL-04-114 and YKL-05-093



2,4-dichloro-5-(iodomethyl)pyrimidine (2)

[00378] A mixture of 2,4-dichloro-5-(chloromethyl)pyrimidine (15.0 g, 76.0 mmol), NaI (13.7 g, 91.4 mmol) in acetone was stirred at 60 °C for 45 min. The resulting precipitate (NaCl) was removed by filtration and washed with acetone. The combined filtrate was concentrated to give light yellow solid, which was purified by column chromatography on silica gel (eluting with DCM) to obtain 2,4-dichloro-5-(iodomethyl)pyrimidine **2** as a light yellow solid (30.8 g, yield 96%). LCMS (m/z): 289.3 [M + H]⁺.

N-((2,4-dichloropyrimidin-5-yl)methyl)-2,6-dimethylaniline (3)

[00379] A mixture of 2,4-dichloro-5-(iodomethyl)pyrimidine **2** (7.0 g, 24.2 mmol), 2,6-dimethylaniline (3.8 g, 31.4 mmol), K₂CO₃ (5.0 g, 36.2 mmol) in acetone (60 mL) was stirred at 55 °C overnight. The solvent was removed and the residue was extracted with EtOAc (150 mL × 3). The combined organic phase was washed with brine (80 mL × 3), dried with Na₂SO₄, filtered, and concentrated. The residue was purified by column chromatography on silica gel (petroleum ether / EtOAc = 8/1, 4/1, 1/1) to get N-((2,4-dichloropyrimidin-5-yl)methyl)-2,6-dimethylaniline **3** as a light brown solid (4.5 g, yield 66%). LCMS (m/z): 282.3 [M + H]⁺.

1-((2,4-dichloropyrimidin-5-yl)methyl)-3-(2,4-dimethoxyphenyl)-1-(2,6-dimethylphenyl)urea (4)


[00380] A round bottomed flask with a Dean-Stark apparatus was charged with N-((2,4-dichloropyrimidin-5-yl)methyl)-2,6-dimethylaniline **3** (3.0 g, 10.6 mmol), 1-isocyanato-2,4-

dimethoxybenzene (2.5 g, 14.0 mmol), toluene (3 mL). The mixture was stirred at 130 °C for 2 d, cooled to rt, and concentrated. The residue was purified by column chromatography on silica gel (petroleum ether / EtOAc = 4/1, 2/1, 1/1, EA) to get 1-((2,4-dichloropyrimidin-5-yl)methyl)-3-(2,4-dimethoxyphenyl)-1-(2,6-dimethylphenyl)urea **4** as a light brown solid (4.1 g, yield 84%). LCMS (m/z): 461.4 [M + H]⁺.

7-chloro-1-(2,4-dimethoxyphenyl)-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (5)

[00381] To the solution of 1-((2,4-dichloropyrimidin-5-yl)methyl)-3-(2,4-dimethoxyphenyl)-1-(2,6-dimethylphenyl) urea **4** (3.1 g, 6.7 mmol) in DCM (20 mL) was added Bu₄NOH (174 mg, 0.67 mmol), NaOH (474 mg, in 2 mL H₂O, 11.8 mmol). The mixture was stirred at rt for 4 h. The final mixture was diluted with H₂O (20 mL), extracted with DCM (80 mL × 3). The combined organic phase was washed with brine (50 mL × 2), dried with Na₂SO₄, filtered, and concentrated. The residue was purified by column chromatography on silica gel (eluting with DCM/MeOH = 20/1) to give 7-chloro-1-(2,4-dimethoxyphenyl)-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one **5** as an off-white solid (2.4 g, yield 85%). ¹H NMR (DMSO-*d*₆, 400 MHz): δ 8.37 (s, 1H), 7.16-7.19 (m, 4H), 6.68 (d, *J* = 2.4 Hz, 1H), 6.58 (dd, *J* = 8.8, 2.4 Hz, 1H), 4.74 (dd, *J* = 5.5, 1.6 Hz, 2H), 3.81 (s, 3H), 3.70 (s, 3H), 2.25 (s, 3H), 2.20 (s, 3H); LCMS (m/z): 425.4 [M + H]⁺.


YKL-04-114



[00382] A mixture of 7-chloro-1-(2,4-dimethoxyphenyl)-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one **5** (10 mg, 0.024 mmol), 3-methoxy-4-(4-methylpiperazin-1-yl)aniline (7.8 mg, 0.035 mmol), and TFA (5.5 mg, 0.048 mmol) in 2-BuOH (0.5 mL) was stirred at 100 °C overnight. The reaction was cooled and concentrated. The residue was purified by prep-HPLC (MeOH/H₂O 5:95 – 100:0), followed by column


chromatography on silica gel (0-10% MeOH in DCM) to afford YKL-04-114 as a white solid (8.0 mg, 56%). ^1H NMR (DMSO- d_6 , 400 MHz): δ 9.21 (s, 1H), 8.20 (s, 1H), 7.25-7.22 (m, 4H), 7.03 (d, $J = 8.4$ Hz, 1H), 6.98 (s, 1H), 6.77 (d, $J = 2.8$ Hz, 1H), 6.68 (dd, $J = 8.8, 2.8$ Hz, 1H), 6.51 (d, $J = 8.4$ Hz, 1H), 4.73 (d, $J = 14.4$ Hz, 1H), 4.59 (d, $J = 14.4$ Hz, 1H), 3.91 (s, 3H), 3.73 (s, 3H), 3.68 (s, 3H), 2.94 (m, 4H), 2.58 (m, 4H), 2.34 (s, 3H), 2.32 (s, 3H), 2.29 (s, 3H); LCMS (m/z): 610.7 [M + H] $^+$.

YKL-05-093



[00383] A mixture of 7-chloro-1-(2,4-dimethoxyphenyl)-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-*d*]pyrimidin-2(1*H*)-one **5** (100 mg, 0.24 mmol), 3-methoxy-4-(1-methylpiperidin-4-yl)aniline (78 mg, 0.35 mmol), and TFA (55 mg, 0.48 mmol) in 2-BuOH (5 mL) was stirred at 100 °C overnight. The reaction was cooled and concentrated. The residue was purified by prep-HPLC (MeOH/H₂O 5:95 – 100:0), followed by column chromatography on silica gel (0-10% MeOH in DCM) to afford YKL-05-093 as a white solid (127 mg, 89%). ^1H NMR (DMSO- d_6 , 400 MHz): δ 9.16 (s, 1H), 8.09 (s, 1H), 7.12-7.09 (m, 4H), 6.95 (d, $J = 8.0$ Hz, 1H), 6.86 (s, 1H), 6.65-6.62 (m, 2H), 6.55 (dd, $J = 8.4, 2.4$ Hz, 1H), 4.60 (d, $J = 14.8$ Hz, 1H), 4.47 (d, $J = 14.8$ Hz, 1H), 3.78 (s, 3H), 3.60 (s, 3H), 3.54 (s, 3H), 2.81 (m, 2H), 2.66-2.57 (m, 1H), 2.19 (s, 3H), 2.16 (s, 3H), 2.15 (s, 3H), 1.95-1.90 (m, 2H), 1.56-1.46 (m, 4H); LCMS (m/z): 609.7 [M + H] $^+$.

Synthesis of YKL-06-061



2-chloro-N-cyclobutyl-5-((2,6-dimethylphenylamino)methyl)pyrimidin-4-amine (2)

[00384] A mixture of N-((2,4-dichloropyrimidin-5-yl)methyl)-2,6-dimethylaniline (**1**) (1.10 g, 3.91 mmol), cyclobutanamine (1.67 g, 23.49 mmol) and DIPEA (6 mL) in THF (100 mL) was stirred at room temperature overnight. Then the mixture was concentrated, the residue was purified by flash column (eluting with ethyl acetate /PE = 0-15%) to give 2-chloro-N-cyclobutyl-5-((2,6-dimethylphenylamino)methyl)pyrimidin-4-amine (**2**) as white solid (0.90 g, yield 72%). LCMS (m/z): 317 [M + H]⁺.


7-chloro-1-cyclobutyl-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (3)

[00385] A mixture of 2-chloro-N-cyclobutyl-5-((2,6-dimethylphenylamino)methyl)pyrimidin-4-amine (**2**) (0.75 g, 2.37 mmol) and triphosgene (1.05 g, 3.56 mmol) in DCM (100 mL) was stirred at room temperature for 1 hour, the a solution of NaOH (1.90 g, 47.5 mmol) and *n*-Bu₄NOH (78 mg, 0.301 mmol) in H₂O (24 mL) was added, the resulting mixture was stirred at room temperature overnight. The organic layer was washed with H₂O (50 mL × 2), dried with Na₂SO₄, filtered and concentrated, the residue was purified by flash column (eluting with ethyl acetate /PE = 0-15%) to give 7-chloro-1-cyclobutyl-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d] pyrimidin-2(1H)-one (**3**) as white solid (750 mg, yield 92%). LCMS (m/z): 343 [M + H]⁺. ¹H-NMR (CDCl₃, 400 MHz): δ 8.11 (s, 1H), 7.11-7.18 (m, 3H), 4.83-4.92 (m, 1H), 4.47 (d, *J* = 0.8Hz, 2H), 2.52-2.59 (m, 4H), 2.22 (s, 6H), 1.74-1.92 (m, 2H).

1-cyclobutyl-3-(2,6-dimethylphenyl)-7-(2-methoxy-4-(4-methylpiperazin-1-yl)phenylamino)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (YKL-06-061)

[00386] A mixture of 7-chloro-1-cyclobutyl-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (**3**) (70 mg, 0.204 mmol), 2-methoxy-4-(4-methylpiperazin-1-yl)aniline (66 mg, 0.298 mmol) and TFA (0.5 mL) in *n*-BuOH (5 mL) was stirred at 110°C overnight, the mixture was concentrated, the residue was purified by prep-HPLC (0.05% NH₄HCO₃ in CH₃CN-H₂O) to give **YKL-06-061** as white solid (50 mg, yield 47%). LCMS (m/z): 528 [M + H]⁺. ¹H-NMR (CDCl₃, 400 MHz): δ 8.22 (d, *J* = 8.4 Hz, 1H), 7.95 (s, 1H), 7.36 (s, 1H), 7.09 - 7.16 (m, 3H), 6.56 - 6.60 (m, 2H), 4.90 - 4.98 (m, 1H), 4.37 (s, 2H), 3.90 (s, 3H), 3.20 (t, *J* = 5.2 Hz, 4H), 2.58 - 2.68 (m, 6H), 2.46-2.54 (m, 2H), 2.38 (s, 3H), 2.23 (s, 6H), 1.74-1.89 (m, 2H).

Synthesis of YKL-06-062



2-chloro-N-cyclohexyl-5-((2,6-dimethylphenylamino)methyl)pyrimidin-4-amine (4)

[00387] A mixture of A mixture of N-((2,4-dichloropyrimidin-5-yl)methyl)-2,6-dimethylaniline (**1**) (1.5g, 5.34 mmol), cyclohexylamine (3.17 g, 32.0 mmol) and DIPEA (6 mL) was stirred at room temperature overnight. Then the mixture was concentrated, the residue was purified by flash column (eluting with ethyl acetate /PE = 0-12%) to give 2-chloro-N-cyclohexyl-5-((2,6-dimethylphenylamino)methyl)pyrimidin-4-amine (**4**) as white solid (1.5 g, yield 81%). LCMS (m/z): 345 [M + H]⁺.

7-chloro-1-cyclohexyl-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (5)






[00388] A mixture of 2-chloro-N-cyclohexyl-5-((2,6-dimethylphenylamino)methyl)pyrimidin-4-amine (**4**) (1.50 g, 4.36 mmol) and triphosgene (1.94 g, 6.54 mmol) in DCM (100 mL) was stirred at room temperature for 30 minutes, then a solution of NaOH (3.49 g, 87.3 mmol) and n-Bu₄NOH (78 mg, 0.301 mmol) in H₂O (43 mL) was added, the resulting mixture was stirred at room temperature overnight. The organic layer was washed with H₂O (50 mL × 2), dried with Na₂SO₄, filtered and concentrated to give crude product, then purified by flash column (eluting with ethyl acetate /PE = 0-20%) to give 7-chloro-1-cyclohexyl-3-(2,6-dimethylphenyl)-3,4-dihydro-pyrimido[4,5-d]pyrimidin-2(1H)-one (**5**) as white solid (1.1 g, yield 68%). LCMS (m/z): 371 [M + H]⁺. ¹H-NMR (CDCl₃, 400 MHz): δ 8.08 (s, 1H), 7.11-7.18 (m, 3H), 4.63-4.71 (m, 1H), 4.49 (d, J = 0.8Hz, 2H), 2.45-2.56 (m, 2H), 2.22 (s, 6H), 1.65-1.86 (m, 5H), 1.34-1.43 (m, 2H), 1.17-1.28 (m, 1H).






1-cyclohexyl-3-(2,6-dimethylphenyl)-7-(4-(4-methylpiperazin-1-yl)phenylamino)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (YKL-06-062)






[00389] A mixture of 7-chloro-1-cyclohexyl-3-(2,6-dimethylphenyl)-3,4-dihydropyrimido[4,5-d]pyrimidin-2(1H)-one (**5**) (94 mg, 0.254 mmol), 4-(4-methylpiperazin-1-yl)aniline (97 mg, 0.508 mmol) and TFA (0.5 mL) in n-BuOH (8 mL) was stirred at 110°C






overnight. The mixture was concentrated, the residue was purified by prep-HPLC (0.05% NH_4HCO_3 in $\text{CH}_3\text{CN}-\text{H}_2\text{O}$) to give **YKL-06-062** as light yellow solid (94 mg, yield 70%). LCMS (m/z): 526 $[\text{M} + \text{H}]^+$. ^1H -NMR (CDCl_3 , 400 MHz): δ 7.92 (s, 1H), 7.46 - 7.50 (m, 2H), 7.09 - 7.15 (m, 3H), 7.02 (s, 1H), 6.93 - 6.96 (m, 2H), 4.61 - 4.69 (m, 1H), 4.38 (s, 2H), 3.19 (t, $J = 5.2$ Hz, 4H), 2.61 (t, $J = 5.2$ Hz, 4H), 2.43-2.53 (m, 2H), 2.36 (s, 3H), 2.23 (s, 6H), 1.76 - 1.86 (m, 4H), 1.66 (d, $J = 12.4$ Hz, 1H), 1.32 - 1.43 (m, 2H), 1.14-1.23 (m, 1H).





Table 1. Exemplary Compounds of Formula (I), (II), (III), (IV), and (V)





Compound Number	Compound Structure
YKL 05-120	
YKL 05-200-01	
YKL 05-200-02	
YKL 05-201-1	
YKL 05-201-2	






Compound Number	Compound Structure
YKL 05-203-1	
YKL 05-203-2	
YKL 05-204-1	
YKL 05-204-2	
YKL 06-029	






Compound Number	Compound Structure
YKL 06-030	
YKL 06-031	
YKL 06-033	
YKL 06-046	
YKL 06-050	






Compound Number	Compound Structure
YKL 06-058	
YKL 06-059	
YKL 06-059	
YKL 06-060	
YKL 06-061	



Compound Number	Compound Structure
YKL 06-062	
YKL-04-136-8	
YKL-04-136-7	
YKL-04-136-6	

Compound Number	Compound Structure
YKL-04-136-5	
YKL-04-136-4	
YKL-04-136-3	
YKL-04-136-2	

Compound Number	Compound Structure
YKL-04-136-1	
YKL-04-125	
YKL-04-118	
YKL-04-115	
YKL-04-114	

Compound Number	Compound Structure
YKL-04-113	
YKL-04-112	
YKL-04-108	
YKL-04-107	
YKL-04-106	

Compound Number	Compound Structure
YKL-04-105	
YKL-04-104	
YKL-04-103	
HG-11-136-01	
HG 9-91-01	

Compound Number	Compound Structure
HG 11-139-02	
HG 11-137-01	

Example 2. A UV independent topical small molecule approach for melanin production in human skin

Small molecule inhibition of SIK induces Mitf expression in vitro.

[00390] To test regulation of the pigmentation pathway by the SIK inhibitor HG 9-91-01 (HG) *in vitro*, normal human melanocytes, UACC62 melanoma cells, and UACC257 melanoma cells were treated. Dose-dependent increases in expression of MITF were observed in these cells (See Figure 1A; Figure 7A, 7B, 7G). RNA levels of the MITF gene target TRPM1 (Miller et al) were also increased and followed the anticipated delayed kinetics relative to MITF induction in normal human melanocytes (See Figure 1B, 1C) and UACC257 cells (Figure 7G, 7H). Gross pigmentation was observed in cell pellets of UACC257 cells after 3 days of HG 9-91-01 treatment (See Figure 1D). These data suggest that small molecule SIK inhibition may stimulate the pigmentation pathway *in vitro*.

[00391] Tables 2 to 5 below show the effects of exemplary SIK inhibitor compounds on skin darkening, including an *in vitro* test of MITF induction following the same protocol as described above in paragraph 333, wherein exemplary SIK inhibitor compounds were tested for their *in vitro* induction of MITF.

Table 2. Effects of exemplary SIK inhibitor compounds on skin darkening

Compound Name	Type of Test	Type of Cells	Results
HG-10-32-01	<i>In vitro</i> MITF induction	B16	
HG-10-88-02	<i>In vitro</i> MITF induction	B16	
HG-10-93-01	<i>In vitro</i> MITF induction	B16	
HG-11-123-01	<i>In vitro</i> MITF induction	B16	
HG-11-136-01	<i>In vitro</i> MITF induction	B16	
HG-11-137-01	<i>In vitro</i> MITF induction	B16	
HG-11-139-01	<i>In vitro</i> MITF induction	B16	
HG-11-139-02	<i>In vitro</i> MITF induction	B16	
HG-11-143-01	<i>In vitro</i> MITF induction	B16	
HG-9-91-01	<i>In vitro</i> MITF induction	B16	
YKL-04-103	<i>In vitro</i> MITF induction	B16	
YKL-04-104	<i>In vitro</i> MITF induction	B16	
YKL-04-105	<i>In vitro</i> MITF induction	B16	
YKL-04-106	<i>In vitro</i> MITF induction	B16	
YKL-04-107	<i>In vitro</i> MITF induction	B16	
YKL-04-108	<i>In vitro</i> MITF induction	B16	
YKL-04-112	<i>In vitro</i> MITF induction	B16	
YKL-04-113	<i>In vitro</i> MITF induction	B16	
YKL-04-114	<i>In vitro</i> MITF induction	B16	
YKL-04-115	<i>In vitro</i> MITF induction	B16	
YKL-04-118	<i>In vitro</i> MITF induction	B16	
YKL-04-125	<i>In vitro</i> MITF induction	B16	
YKL-04-136-1	<i>In vitro</i> MITF induction	B16	
YKL-04-136-10	<i>In vitro</i> MITF induction	B16	
YKL-04-136-11	<i>In vitro</i> MITF induction	B16	
YKL-04-136-2	<i>In vitro</i> MITF induction	B16	
YKL-04-136-3	<i>In vitro</i> MITF induction	B16	
YKL-04-136-4	<i>In vitro</i> MITF induction	B16	
YKL-04-136-5	<i>In vitro</i> MITF induction	B16	
YKL-05-120	Human Skin Explants		Slight Darkening
YKL-05-200-1	Human Skin Explants		Slight Darkening
YKL-05-200-2	Human Skin Explants		Darkens
YKL-05-201-1	Human Skin Explants		Darkens
YKL-05-201-2	Human Skin Explants		Slight Darkening
YKL-05-203-1	Human Skin Explants		Slight Darkening
YKL-05-203-2	Human Skin Explants		Slight Darkening
YKL-05-204-1	Human Skin Explants		Darkens
YKL-05-204-2	Human Skin Explants		Darkens

YKL-06-029	<i>In vitro</i> MITF induction	UACC62	
YKL-06-058	Human Skin Explants		Darkens
YKL-06-059	Human Skin Explants		Darkens
YKL-06-060	Human Skin Explants		Darkens
YKL-06-061	Human Skin Explants		Darkens
YKL-06-062	Human Skin Explants		Darkens
YKL-06-29	Human Skin Explants		Darkens
YKL-06-30	Human Skin Explants		Slight Darkening
YKL-06-31	Human Skin Explants		Slight Darkening
YKL-06-33	Human Skin Explants		Slight Darkening
YKL-06-46	Human Skin Explants		Slight Darkening
YKL-06-50	Human Skin Explants		Slight Darkening

Table 3: Effects of exemplary SIK inhibitor compounds on skin darkening

Compound Name	Type of Test	Type of Cells	Results
HG-11-137-01	Human Skin Explants		Darkens
HG-11-139-02	Human Skin Explants		No Darkening
HG-9-91-01	Human Skin Explants		Darkens
YKL-04-108	Human Skin Explants		Darkens
YKL-05-200-2	<i>In vitro</i> MITF induction	UACC62	
YKL-05-201-1	<i>In vitro</i> MITF induction	UACC62	
YKL-05-204-1	<i>In vitro</i> MITF induction	UACC62	
YKL-06-029	<i>In vitro</i> MITF induction	UACC62	
YKL-06-059	<i>In vitro</i> MITF induction	UACC62	
YKL-06-060	<i>In vitro</i> MITF induction	UACC62	
YKL-06-061	<i>In vitro</i> MITF induction	UACC62	
YKL-06-062	<i>In vitro</i> MITF induction	UACC62	

Table 4: Effects of exemplary SIK inhibitor compounds on skin darkening

Compound Name	Type of Test	Results
HG-11-137-01	Human Skin Explants	Slight Darkening
HG-11-139-02	Human Skin Explants	Slight Darkening
HG-9-91-01	Human Skin Explants	Darkens
YKL-05-120	Human Skin Explants	Slight Darkening
YKL-05-200-1	Human Skin Explants	Slight Darkening
YKL-05-200-2	Human Skin Explants	Darkens
YKL-05-201-1	Human Skin Explants	Darkens
YKL-05-201-2	Human Skin Explants	Slight Darkening
YKL-05-203-1	Human Skin Explants	Slight Darkening
YKL-05-203-2	Human Skin Explants	Slight Darkening

YKL-05-204-1	Human Skin Explants	Darkens
YKL-05-204-2	Human Skin Explants	Darkens
YKL-06-058	Human Skin Explants	Darkens
YKL-06-059	Human Skin Explants	Darkens
YKL-06-060	Human Skin Explants	Darkens
YKL-06-061	Human Skin Explants	Darkens
YKL-06-062	Human Skin Explants	Darkens
YKL-06-29	Human Skin Explants	Darkens
YKL-06-30	Human Skin Explants	Slight Darkening
YKL-06-31	Human Skin Explants	Slight Darkening
YKL-06-33	Human Skin Explants	Slight Darkening
YKL-06-46	Human Skin Explants	Slight Darkening
YKL-06-50	Human Skin Explants	Slight Darkening

[00392] Table 5 below additionally provides characteristics of exemplary SIK inhibitor compounds, MITF induction in B16 cells, EC₅₀, logP, and experimental conditions for skin darkening experiments in mice and humans.

Table 5. Effects of exemplary SIK inhibitor compounds on skin darkening; and characteristics of exemplary SIK inhibitor compounds

Compound Name	Molecular weight (g/mol)	Highest fold mitf induction in B16 cells	H-bond donors	H-bond acceptors	Concentration highest induction observed (nM)	EC50 (nM)	logP	Days Treated Mouse	□ Mouse	Darkening (Mouse)	Days Treated Human	□ Human	Darkening (Human)
YKL-04-118		1.1			0.08	1424.000	6.36						
YKL-04-136-07		2.4			0.4	0.127	3.38						
HG-9-91-01		2.5			0.06	0.016	6.01	2	25 mM	Darkens	5	25 mM	No darkening
YKL-04-107		3.6			0.4	0.015	6.43						
YKL-04-136-08		4.1			0.4	0.014	3.57						

Compound Name	Molecular weight (g/mol)	Highest fold mitf induction in B16 cells	H-bond donors	H-bond acceptors	Concentration highest induction observed (nM)	EC50 (uM)	logP	Days Treated Mouse	□ Mouse	Darkening (Mouse)	Days Treated Human	□ Human	Darkening (Human)
YKL-04-112		5.2			0.4	0.215	6.59						
HG-11-136-01		6.1			0.06	0.000	5.55	12	25 mM	Darkens	16	75 mM	No Darkening
YKL-04-136-03		6.2			0.08	0.025	5.8						
HG-11-143-01		6.8			0.4	0.057	5.6	12	25 mM (4 days) -130 mM (8 days)	No Darkening			No Darkening
YKL-04-104		8.0			0.4	0.021	6.52						
YKL-04-136-05		8.9			0.4	0.042	5.05						
YKL-04-106		9.0			0.4	0.074	5.4						
YKL-04-136-04		9.2			0.08	0.015	4.13						
YKL-04-136-01		9.7			0.4	0.053	5.55						
YKL-04-125		9.7			0.4	0.133	6.42						
YKL-04-105		9.9			0.08	0.005	4.15						
YKL-04-115		10.1			0.08	0.024	5.49						
YKL-04-103		10.2			0.08	0.001	6.78						

Compound Name	Molecular weight (g/mol)	Highest fold mitf induction in B16 cells	H-bond donors	H-bond acceptors	Concentration highest induction observed (nM)	EC50 (uM)	logP	Days Treated Mouse	□ Mouse	Darkening (Mouse)	Days Treated Human	□ Human	Darkening (Human)
YKL-04-114		10.4			0.4	0.067	5.55						
YKL-04-113		11.2			0.4	0.657	6.52						
YKL-04-108		13.9			0.08	0.014	6.21						
YKL-04-136-02		16.1			0.08	0.021	4.04						
YKL-04-136-09		16.5			0.4	0.007	5.64						
YKL-04-136-06		26.9			0.08	0.009	5.25						
HG-10-32-01	401.461	2.4	2	5			4.91						
YKL-04-136-10	564.681	16.28	1	8			5.08						
YKL-04-136-11	689.849	9.58	1	9			4.89						
YKL-04-193-1	465.593	2.02	2	7			5.92						
YKL-04-193-2	428.533	2.65	2	7			4.41						
HG-11-137-01	429.518	4.82	1	6			3.34						
HG-11-139-02	459.544	5.76	1	7			3.19						
Contr ol								2		No darkening	16		No darkenin g

[00393] Table 6 below provides characteristics of exemplary SIK inhibitor compounds, highest fold MITF induction, and logP. Table 7 below provides characteristics of exemplary SIK inhibitor compounds and logP

Table 6. Characteristics of exemplary SIK inhibitor compounds, and effects of exemplary SIK inhibitor compounds on MITF induction

Compound Name	Molecular weight (g/mol)	log P	H-bond donors	H-bond acceptors	Highest fold Mitf induction
HG-10-32-01	401.461	4.91	2	5	2.4
YKL-04-136-10	564.681	5.08	1	8	16.28
YKL-04-136-11	689.849	4.89	1	9	9.58
YKL-04-193-1	465.593	5.92	2	7	2.02
YKL-04-193-2	428.533	4.41	2	7	2.65
HG-11-137-01	429.518	3.34	1	6	4.82
HG-11-139-02	459.544	3.19	1	7	5.76

Table 7. Characteristics of exemplary SIK inhibitor compounds

Molecule Name	Molecular weight (g/mol)	log P	H-bond donors	H-bond acceptors
HG-9-91-01	567.681	6.01	2	8
HG-11-137-01	429.518	3.34	1	6
HG-11-139-02	459.544	3.19	1	7
YKL-05-200-2	427.498	4.65	1	5
YKL-05-201-1	447.917	4.74	1	5
YKL-05-204-1	511.618	4.57	1	7
YKL-06-029	457.571	4.37	1	6
YKL-06-059	515.65	4.99	1	7
YKL-06-060	497.634	5.28	1	6
YKL-06-061	527.66	5.12	1	7
YKL-06-062	525.688	6.17	1	6

HG 9-91-01 rescues melanogenesis in mice with inactive melanocortin 1 receptor.

[00394] Since the *in vitro* results demonstrated that inhibition of SIK by HG 9-91-01 positively regulated the CRTC-CREB-MITF pathway, it was next evaluated whether topical application of this compound could induce pigmentation independent of Mc1r *in vivo*. To test this, a previously described mouse “redhair” model was utilized which carries the inactivating *Mc1r^{ele}* mutant allele and a transgene, K14-SCF, in which Stem Cell Factor expression is driven by the Keratin-14 promoter allowing for epidermal homing of melanocytes (Kunisada *et al.* 1998; D'Orazio *et al.* 2006). Albino mice harboring a mutation

in the *tyrosinase* gene were combined with the K14-Scf transgene (*tyr^{cl/c}*;K14-Scf mice) and served as controls to evaluate whether the pigmentation afforded by topical SIK inhibitor was dependent upon the canonical tyrosinase-melanin pathway. Daily application of the SIK inhibitor HG 9-91-01 for 6 days, caused robust darkening in *Mc1r^{ee/e}*;K14-Scf mice (Figure 2A; Figure 8A). No visible change in skin pigmentation was observed in *Mc1r^{ee/e}*;K14-Scf mice treated with vehicle or *Tyr^{cl/c}*;K14-Scf mice treated with vehicle or HG 9-91-01 (Figure 2A; Figure 8A, 8B). Reflective colorimetry analysis (CIE L* white-black color axis (Park, Suh, and Youn 1999)) revealed significant darkening in *Mc1r^{ee/e}*;K14-Scf mice treated with SIK inhibitor, but not in vehicle treated *Mc1r^{ee/e}*;K14-Scf mice or *Tyr^{cl/c}*;K14-Scf mice treated with either SIK inhibitor or vehicle control (Figure 2B). Fontana-Masson staining, a specialized melanin stain, revealed strong induction of melanin production in *Mc1r^{ee/e}*;K14-Scf mice in areas treated with HG 9-91-01, (Figure 2C; Figure 8D) but no pigment induction in *Mc1r^{ee/e}*;K14-Scf mice treated with vehicle or in albino (*Tyr^{cl/c}*;K14-Scf) mice treated with either vehicle or SIK inhibitor (Figure 8C). Nuclear capping of melanin-laden melanosomes was observed within epidermal keratinocytes in *Mc1r^{ee/e}*;K14-Scf mice treated with HG 9-91-01 (indicated by white arrows) and represents a known subcellular localization typical of physiologic skin pigmentation (Kobayashi *et al.* 1998) (Figure 2C). This feature suggests that SIK inhibitor treatment not only stimulates melanocytic pigment synthesis, but also export of melanin in a fashion that closely mimics the known pathway of UV melanogenesis. Hematoxylin and eosin staining revealed normal morphology of HG 9-91-01 treated *Mc1r^{ee/e}*;K14-Scf mice (Figure 2C) and *Tyr^{cl/c}*;K14-Scf epidermis (Figure 8C). Additionally, NaOH lysis of skin samples (Wakamatsu and Ito 2002) revealed a visible increase in extractable eumelanin from *Mc1r^{ee/e}*;K14-Scf mice treated with HG 9-91-01 compared to all other treatment groups (Figure 2D). The darkening induced by topical application of HG 9-91-01 to *Mc1r^{ee/e}*;K14-Scf mice was progressive over 6 days of treatment and reversible by 10 days after treatment was stopped (Figure 3A, 3B). Skin pigmentation remained in its pretreatment state over the next 30 days (Figure 3A; 3B). No change was observed in *Tyr^{cl/c}*;K14-Scf mice during treatment or 14 days after treatment was stopped (Figure 3C). Fontana-Masson staining of skin sections revealed no differences in either treatment group for *Mc1r^{ee/e}*;K14-Scf mice or *Tyr^{cl/c}*;K14-Scf mice and hematoxylin and eosin stain illustrated normal morphology for all mice 40 days after treatment (Figure 8E). These findings combined with the small molecule and lipophilic nature of the SIK inhibitors led us to investigate the use of SIK inhibitors for topical eumelanization of human skin.

Second generation SIK inhibitors are as efficacious in inducing the pigmentation pathway as HG 9-91-01.

[00395] Since there are limitations to topical delivery of drugs into human skin epidermis, novel SIK inhibitors designed to enhance epidermal permeation by reducing the size and increasing the lipophilicity (cLogP) to develop second generation SIK inhibitors YKL 06-061 and YKL 06-062 were derived (Figure 6) (Bos and Meinardi 2000; Hadgraft and Pugh 1998). They had comparable IC₅₀ values for inhibition of SIKs (Figure 6). To assess the kinome selectivity information of new analogs, YKL-06-061 was screened across a panel of 468 human kinases at a concentration of 1 μ M using the KinomeScan methodology (DiscoverX). YKL-06-061 exhibited a S(1) score of 0.02 with 16 kinases displaying tight binding to it (Ambit scores of less than or equal to 1). As the KinomeScan assays measure binding, enzymatic assays for these targets were also performed either in-house or using the SelectScreen Kinase Profiling Service at Thermo Fisher Scientific (Madison, WI). YKL-06-061 only inhibited one kinase (FRK) stronger than SIKs, which demonstrates its high overall selectivity. It was anticipated that YKL-06-062 had similar kinase selectivity considering their high structural similarity.

[00396]— Similar to observations with HG 9-91-01, after 3 hours of treatment of normal human melanocytes (Figure 4A, B), UACC62, and UACC257 cells (Figure 7C-F), there was a dose dependent increase in MITF mRNA expression for YKL 06-061 and YKL 06-062. Levels of TRPM1 mRNA increased after MITF induction with agents these cells (Figure C-D; Figure 7G, 7H).

Topical SIK inhibitors induce human skin eumelanization

[00397] Treatment of human skin explants with passive topical application of the second generation SIK inhibitors, YKL 06-061 and YKL 06-062, induced significant pigmentation without any additional treatments after 8 days (1x/day) of treatment, but no significant gross pigmentation was observed in skin treated with HG 9-91-01 (Figure 5A). Fontanna Mason staining revealed increased melanin content in skin treated with YKL 06-061 and YKL 06-062 and marginally increased melanin in skin treated with HG 9-91-01 as compared to control (Figure 5B). This effect was reproducible with independent preparations of synthesized drugs applied passively to different human skin donors (Figure 5C; 5D). Mechanical application of the first generation SIK inhibitor HG 9-91-01 (by rubbing via an applicator) induced significant gross pigmentation (Figure 5E), and increased melanin content was observed upon Fontana Masson staining of skin sections (Figure 5F), suggesting

HG 9-91-01's human skin limited penetration can be at least partially overcome through mechanical application. YKL 06-061 and YKL 06-062 did not require mechanical application (rubbing) to induce significant human epidermal darkening. Human skin explants were treated with passive topical application of the exemplary SIK inhibitors shown in Tables 2-5 above. Whether the exemplary SIK inhibitors induced pigmentation without any additional treatments after 8 days (1x/day) of treatment is shown in Tables 2-5 above. Fontanna Mason staining further illustrated whether exemplary SIK inhibitors induced pigmentation in the human skin explants. *See* Tables 2-5.

[00398] These results illustrate, but are not limited to, the development and successful application of small molecule SIK inhibitors for topical induction of skin pigmentation independently of UV irradiation in human skin. SIK inhibitors were shown to induce enhanced expression of the MITF transcription factor which is known to regulate expression of numerous pigment enzymes that promote biosynthesis of eumelanin. A new generation of SIK inhibitors was developed, based upon strategies designed to enhance likelihood of skin penetration through increased lipophilicity. Two such SIK targeted inhibitors, YKL 06-061 and YKL 06-062, were shown to induce similar responses both *in vitro* and when applied to human skin explants. In addition to upregulating mRNA levels of MITF and TRPM1, topical SIK inhibitors were seen to trigger transfer of melanosomes into epidermal keratinocytes in a manner which recapitulates the peri-nuclear capping (subcellular localization) that is seen in normal human epidermal pigmentation. Thus SIK inhibitor treatments appear to induce not only synthesis of melanin, but also melanosomal maturation, export, and localization features, even after import into keratinocytes. These features closely resemble the previously observed behavior of forskolin treatment in redhaired mice (D'Orazio *et al.* 2006).

[00399] For example, the induction of dark pigmentation is associated with the lowest risk of most skin cancers in humans, and this pigment synthesis is believed to be dependent upon MITF. Yet fixed genomic mutation or amplification of the MITF gene can be oncogenic in certain contexts (Garraway *et al.*; Yokoyama *et al.*; Bertolotto *et al.*). Reversible upregulation of MITF as reported here, is also likely to occur in routine instances of UV-tanning, or to be constitutively elevated in skin of individuals with darker pigmentation levels, and would not be anticipated to trigger genomic mutation of the MITF gene. Analogously, transient administration of recombinant hematopoietic growth factors has not been associated with formation of oncogenic transformation or leukemias. In mice, topical forskolin's pigmentary rescue in "redheads" resulted in significant protection from UV carcinogenesis without apparent associated toxicities over many months of treatment (D'Orazio *et al.*). A recent

study has utilized injections of the synthetic alpha-MSH analogue, afamelanotide, for treatment of photosensitivity associated with erythropoietic protoporphyria (Langendonk et al 2015). In this work systemic elevations in MSH downstream signals were induced, and pigmented lesions/melanoma were carefully evaluated and reported not to occur at elevated risk.

[00400] The half-life of melanin in skin is thought to be several weeks and primarily diminishes only after superficial keratinocyte sloughing. Most epidermal melanin resides within keratinocytes after transfer of melanosomes from melanocytes. Therefore it is possible that small molecule approaches like that described here, but not limited to these exemplary approaches, might be achievable, or maintained, through intermittent dosing strategies. In conclusion, these studies describe an exemplary small molecule topical approach to the rescue of eumelanin synthesis in a UV-independent manner. The applications of the strategy could be significant in a number of UV-protection related settings.

REFERENCES

1. Armstrong, B. K., and A. Krickler. 2001. 'The epidemiology of UV induced skin cancer', *J Photochem Photobiol B*, 63: 8-18.
2. Bertolotto, C., P. Abbe, T. J. Hemesath, K. Bille, D. E. Fisher, J. P. Ortonne, and R. Ballotti. 1998. 'Microphthalmia gene product as a signal transducer in cAMP-induced differentiation of melanocytes', *J Cell Biol*, 142: 827-35.
3. Bos, J. D., and M. M. Meinardi. 2000. 'The 500 Dalton rule for the skin penetration of chemical compounds and drugs', *Exp Dermatol*, 9: 165-9.
4. Clark, K., K. F. MacKenzie, K. Petkevicius, Y. Kristariyanto, J. Zhang, H. G. Choi, M. Pegg, L. Plater, P. G. Pedrioli, E. McIver, N. S. Gray, J. S. Arthur, and P. Cohen. 2012. 'Phosphorylation of CRT3 by the salt-inducible kinases controls the interconversion of classically activated and regulatory macrophages', *Proc Natl Acad Sci U S A*, 109: 16986-91.
5. Cui, R., H. R. Widlund, E. Feige, J. Y. Lin, D. L. Wilensky, V. E. Igras, J. D'Orazio, C. Y. Fung, C. F. Schanbacher, S. R. Granter, and D. E. Fisher. 2007. 'Central role of p53 in the suntan response and pathologic hyperpigmentation', *Cell*, 128: 853-64.
6. D'Orazio, J. A., T. Nobuhisa, R. Cui, M. Arya, M. Spry, K. Wakamatsu, V. Igras, T. Kunisada, S. R. Granter, E. K. Nishimura, S. Ito, and D. E. Fisher. 2006. 'Topical drug rescue strategy and skin protection based on the role of Mc1r in UV-induced tanning', *Nature*, 443: 340-4.
7. Dentin, R., Y. Liu, S. H. Koo, S. Hedrick, T. Vargas, J. Heredia, J. Yates, 3rd, and M.

- Montminy. 2007. 'Insulin modulates gluconeogenesis by inhibition of the coactivator TORC2', *Nature*, 449: 366-9.
8. Gandini, S., F. Sera, M. S. Cattaruzza, P. Pasquini, O. Picconi, P. Boyle, and C. F. Melchi. 2005. 'Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure', *Eur J Cancer*, 41: 45-60.
9. Hadgraft, J., and W. J. Pugh. 1998. 'The selection and design of topical and transdermal agents: a review', *J Investig Dermatol Symp Proc*, 3: 131-5.
10. Harms, J. H., S. Lautenschlager, C. E. Minder, and E. I. Minder. 2009. 'Mitigating photosensitivity of erythropoietic protoporphyria patients by an agonistic analog of alpha-melanocyte stimulating hormone', *Photochem Photobiol*, 85: 1434-9.
11. Horike, N., A. Kumagai, Y. Shimono, T. Onishi, Y. Itoh, T. Sasaki, K. Kitagawa, O. Hatano, H. Takagi, T. Susumu, H. Teraoka, K. Kusano, Y. Nagaoka, H. Kawahara, and H. Takemori. 2010. 'Downregulation of SIK2 expression promotes the melanogenic program in mice', *Pigment Cell Melanoma Res*, 23: 809-19.
12. Kennedy, C., C. D. Bajdik, R. Willemze, F. R. De Gruijl, and J. N. Bouwes Bavinck. 2003. 'The influence of painful sunburns and lifetime sun exposure on the risk of actinic keratoses, seborrheic warts, melanocytic nevi, atypical nevi, and skin cancer', *J Invest Dermatol*, 120: 1087-93.
13. Khaled, M., C. Levy, and D. E. Fisher. 2010. 'Control of melanocyte differentiation by a MITF-PDE4D3 homeostatic circuit', *Genes Dev*, 24: 2276-81.
14. Kobayashi, N., A. Nakagawa, T. Muramatsu, Y. Yamashina, T. Shirai, M. W. Hashimoto, Y. Ishigaki, T. Ohnishi, and T. Mori. 1998. 'Supranuclear melanin caps reduce ultraviolet induced DNA photoproducts in human epidermis', *J Invest Dermatol*, 110: 806-10.
15. Kumagai, A., N. Horike, Y. Satoh, T. Uebi, T. Sasaki, Y. Itoh, Y. Hirata, K. Uchio-Yamada, K. Kitagawa, S. Uesato, H. Kawahara, H. Takemori, and Y. Nagaoka. 2011. 'A potent inhibitor of SIK2, 3, 3', 7-trihydroxy-4'-methoxyflavon (4'-O-methylfisetin), promotes melanogenesis in B16F10 melanoma cells', *PLoS One*, 6: e26148.
16. Kunisada, T., S. Z. Lu, H. Yoshida, S. Nishikawa, S. Nishikawa, M. Mizoguchi, S. Hayashi, L. Tyrrell, D. A. Williams, X. Wang, and B. J. Longley. 1998. 'Murine cutaneous mastocytosis and epidermal melanocytosis induced by keratinocyte expression of transgenic stem cell factor', *J Exp Med*, 187: 1565-73.
17. Lim, H. W., W. D. James, D. S. Rigel, M. E. Maloney, J. M. Spencer, and R. Bhushan. 2011. 'Adverse effects of ultraviolet radiation from the use of indoor tanning equipment: time to ban the tan', *J Am Acad Dermatol*, 64: 893-902.

18. Newton, R. A., S. E. Smit, C. C. Barnes, J. Pedley, P. G. Parsons, and R. A. Sturm. 2005. 'Activation of the cAMP pathway by variant human MC1R alleles expressed in HEK and in melanoma cells', *Peptides*, 26: 1818-24.
19. Oancea, E., J. Vriens, S. Brauchi, J. Jun, I. Splawski, and D. E. Clapham. 2009. 'TRPM1 forms ion channels associated with melanin content in melanocytes', *Sci Signal*, 2: ra21.
20. Park, S. B., D. H. Suh, and J. I. Youn. 1999. 'A long-term time course of colorimetric evaluation of ultraviolet light-induced skin reactions', *Clin Exp Dermatol*, 24: 315-20.
21. Pennello, G., S. Devesa, and M. Gail. 2000. 'Association of surface ultraviolet B radiation levels with melanoma and nonmelanoma skin cancer in United States blacks', *Cancer Epidemiol Biomarkers Prev*, 9: 291-7.
22. Prevention, Centers for Disease Control and. 21 June 2016. 'Skin Cancer Statistics'. <https://www.cdc.gov/cancer/skin/statistics/>.
23. Price, E. R., M. A. Horstmann, A. G. Wells, K. N. Weilbaecher, C. M. Takemoto, M. W. Landis, and D. E. Fisher. 1998. 'alpha-Melanocyte-stimulating hormone signaling regulates expression of microphthalmia, a gene deficient in Waardenburg syndrome', *J Biol Chem*, 273: 33042-7.
24. Rogers, H. W., M. A. Weinstock, S. R. Feldman, and B. M. Coldiron. 2015. 'Incidence Estimate of Nonmelanoma Skin Cancer (Keratinocyte Carcinomas) in the U.S. Population, 2012', *JAMA Dermatol*, 151: 1081-6.
25. Ryerson, A. B., C. R. Ehemann, S. F. Altekruse, J. W. Ward, A. Jemal, R. L. Sherman, S. J. Henley, D. Holtzman, A. Lake, A. M. Noone, R. N. Anderson, J. Ma, K. N. Ly, K. A. Cronin, L. Penberthy, and B. A. Kohler. 2016. 'Annual Report to the Nation on the Status of Cancer, 1975-2012, featuring the increasing incidence of liver cancer', *Cancer*, 122: 1312-37.
26. Tsatmalia, M., K. Wakamatsu, A. J. Graham, and A. J. Thody. 1999. 'Skin POMC peptides. Their binding affinities and activation of the human MC1 receptor', *Ann N Y Acad Sci*, 885: 466-9.
27. Valverde, P., E. Healy, I. Jackson, J. L. Rees, and A. J. Thody. 1995. 'Variants of the melanocyte-stimulating hormone receptor gene are associated with red hair and fair skin in humans', *Nat Genet*, 11: 328-30.
28. Wakamatsu, K., and S. Ito. 2002. 'Advanced chemical methods in melanin determination', *Pigment Cell Res*, 15: 174-83.
29. Watson, M., A. C. Geller, M. A. Tucker, G. P. Guy, Jr., and M. A. Weinstock. 2016. 'Melanoma burden and recent trends among non-Hispanic whites aged 15-49years, United

States', *Prev Med*, 91: 294-98.

30. Wu, S., J. Han, R. A. Vleugels, R. Puett, F. Laden, D. J. Hunter, and A. A. Qureshi. 2014. 'Cumulative ultraviolet radiation flux in adulthood and risk of incident skin cancers in women', *Br J Cancer*, 110: 1855-61.

31. Altarejos, J. Y., and Montminy, M. (2011) CREB and the CRTC co-activators: sensors for hormonal and metabolic signals. *Nat. Rev. Mol. Cell Biol.* 12, 141–151.

32. Patel, K., Foretz, M., Marion, A., Campbell, D. G., Gourlay, R., Boudaba, N., Tournier, E., Titchenell, P., Pegg, M., Deak, M., Wan, M., Kaestner, K. H., Goransson, O., Viollet, B., Gray, N. S., Birnbaum, M. J., Sutherland, C., and Sakamoto, K. (2014) The LKB1-salt-inducible kinase pathway functions as a key gluconeogenic suppressor in the liver, *Nat. Commun.*, 5.

33. Park, J., Yoon, Y. S., Han, H. S., Kim, Y. H., Ogawa, Y., Park, K. G., Lee, C. H., Kim, S. T., and Koo, S. H. (2014) SIK2 Is Critical in the Regulation of Lipid Homeostasis and Adipogenesis In Vivo. *Diabetes*, 63, 3659–3673.

34. Henriksson, E., Sall, J., Gormand, A., Wasserstrom, S., Morrice, N. A., Fritzen, A. M., Foretz, M., Campbell, D. G., Sakamoto, K., Ekelund, M., Degerman, E., Stenkula, K. G., and Goransson, O. (2015) SIK2 regulates CRTC, HDAC4 and glucose uptake in adipocytes. *J. Cell Sci.*, 128, 472–486.

35. Kumagai *et al.*, PLOS One, Oct. 13, 2011.

36. Horike *et al.*, Pigment Cell Melanoma Res. 23; 809–819 (2010).

37. Langendonk JG *et al.*, (2015). Afamelanotide for Erythropoietic Protoporphyrria. *New Eng J Med*. 373 (1): 48-59.

38. Yasumoto *et al.*, Molecular Cell Biology. 1994, 8058-70.

39. Pfeifer *et al.*, Mutation Research. 2005, 571, 19-31.

EQUIVALENTS AND SCOPE

1. In the claims articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention

includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

2. Furthermore, the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims is introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, *e.g.*, in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements and/or features, certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth *in haec verba* herein. It is also noted that the terms “comprising” and “containing” are intended to be open and permits the inclusion of additional elements or steps. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

3. This application refers to various issued patents, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. If there is a conflict between any of the incorporated references and the instant specification, the specification shall control. In addition, any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Because such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the invention can be excluded from any claim, for any reason, whether or not related to the existence of prior art.


4. Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments described herein. The scope of the present embodiments described herein is not intended to be limited to the above Description, but rather is as set forth in the appended claims. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made

without departing from the spirit or scope of the present invention, as defined in the following claims.

CLAIMS

What is claimed is:

1. A method of increasing skin pigmentation in a subject, the method comprising administering topically to the subject's skin an effective amount of a salt inducible kinase (SIK) inhibitor of any one of Formulae (I), (II), (III), (IV), (V), (VI), (VI-A), and (VII), or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, or stereoisomer thereof, or a pharmaceutical composition thereof.
2. A method of increasing the appearance of skin pigmentation in a subject, the methods comprising administering topically to the subject's skin an effective amount of a salt inducible kinase (SIK) inhibitor of any one of Formulae (I), (II), (III), (IV), (V), (VI), (VI-A), and (VII), or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, or stereoisomer thereof, or a pharmaceutical composition thereof.
3. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is a substituted or unsubstituted phenyl ring or a substituted or unsubstituted, monocyclic, 5- to 6-membered heteroaryl ring, wherein one, two, three, or four atoms in the heteroaryl ring system are independently nitrogen, oxygen, or sulfur;

each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{A1}$, $-N(R^{A1})_2$, $-SR^{A1}$, $-CN$, $-SCN$, $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, $-C(=NR^{A1})N(R^{A1})_2$, $-C(=O)R^{A1}$, $-C(=O)OR^{A1}$, $-C(=O)N(R^{A1})_2$, $-NO_2$, $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$, $-NR^{A1}C(=O)N(R^{A1})_2$, $-OC(=O)R^{A1}$, $-OC(=O)OR^{A1}$, or $-OC(=O)N(R^{A1})_2$, wherein each instance of R^{A1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl,

substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^{A1} groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, or 4;

L is a substituted or unsubstituted, saturated or unsaturated C_{3-10} hydrocarbon chain, optionally wherein one or more chain atoms of the hydrocarbon chain are independently replaced with $-O-$, $-S-$, $-NR^N-$, $-N=$, or $=N-$, wherein each instance of R^N is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

R^B is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each of X^A , X^B , and X^C is independently N or CR^X , wherein R^X is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{X1}$, $-N(R^{X1})_2$, $-SR^{X1}$, $-CN$, $-SCN$, $-C(=NR^{X1})R^{X1}$, $-C(=NR^{X1})OR^{X1}$, $-C(=NR^{X1})N(R^{X1})_2$, $-C(=O)R^{X1}$, $-C(=O)OR^{X1}$, $-C(=O)N(R^{X1})_2$, $-NO_2$, $-NR^{X1}C(=O)R^{X1}$, $-NR^{X1}C(=O)OR^{X1}$, $-NR^{X1}C(=O)N(R^{X1})_2$, $-OC(=O)R^{X1}$, $-OC(=O)OR^{X1}$, or $-OC(=O)N(R^{X1})_2$, wherein each instance of R^{X1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^{X1} groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

Y is $-O-$ or $-NR^Y-$, wherein R^Y is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

or when Y is $-NR^Y-$ and X^A is CR^X , R^Y and R^X of X^A are joined to form a substituted or unsubstituted, monocyclic, 5- to 7-membered heterocyclic ring that is fused with Ring B;

each instance of R^C is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{C1}$, $-N(R^{C1})_2$, $-SR^{C1}$, $-CN$, $-SCN$, $-C(=NR^{C1})R^{C1}$, $-C(=NR^{C1})OR^{C1}$, $-C(=NR^{C1})N(R^{C1})_2$, $-C(=O)R^{C1}$, $-C(=O)OR^{C1}$, $-C(=O)N(R^{C1})_2$, $-NO_2$, $-NR^{C1}C(=O)R^{C1}$, $-NR^{C1}C(=O)OR^{C1}$, $-NR^{C1}C(=O)N(R^{C1})_2$, $-OC(=O)R^{C1}$, $-OC(=O)OR^{C1}$, or $-OC(=O)N(R^{C1})_2$, wherein each instance of R^{C1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^{C1} groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

m is 0, 1, 2, 3, 4, or 5; and


R^D is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{D1}$, $-N(R^{D1})_2$, $-SR^{D1}$, $-CN$, $-SCN$, $-C(=NR^{D1})R^{D1}$, $-C(=NR^{D1})OR^{D1}$, $-C(=NR^{D1})N(R^{D1})_2$, $-C(=O)R^{D1}$, $-C(=O)OR^{D1}$, $-C(=O)N(R^{D1})_2$, $-NO_2$, $-NR^{D1}C(=O)R^{D1}$, $-NR^{D1}C(=O)OR^{D1}$, $-NR^{D1}C(=O)N(R^{D1})_2$, $-OC(=O)R^{D1}$, $-OC(=O)OR^{D1}$, or $-OC(=O)N(R^{D1})_2$, wherein each instance of R^{D1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.

4. The method of claim 2, or a pharmaceutically acceptable salt thereof, wherein Ring A is a substituted or unsubstituted phenyl ring.

5. The method of any one of claims 2-4, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^A is substituted or unsubstituted, monocyclic, 3- to 7-membered heterocyclyl, wherein one, two, or three atoms in the heterocyclic ring system are independently selected from the group consisting of nitrogen, oxygen, and sulfur.


6. The method of claim 5, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^A is substituted or unsubstituted oxetanyl, substituted or unsubstituted tetrahydrofuranyl, substituted or unsubstituted pyrrolidinyl, substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl, or substituted or unsubstituted piperazinyl.

7. The method of claim 6, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^A is of the formula:



8. The method of claim 7, or a pharmaceutically acceptable salt thereof, wherein at least

one instance of R^A is of the formula:



The structure shows a six-membered piperidine ring. The nitrogen at the bottom-left position is connected to a wavy line, representing a point of attachment. The nitrogen at the top-right position is connected to a methyl group (represented by a single line).

9. The method of any one of claims 3-8, or a pharmaceutically acceptable salt thereof, wherein k is 0.


10. The method of any one of claims 3-8, or a pharmaceutically acceptable salt thereof, wherein k is 1.

11. The method of any one of claims 3-10, or a pharmaceutically acceptable salt thereof, wherein L consists of not more than about 30 atoms.


12. The method of any one of claims 3-11, or a pharmaceutically acceptable salt thereof, wherein the molecular weight of L is not more than about 150 g/mol.

13. The method of any one of claims 3-12, or a pharmaceutically acceptable salt thereof, wherein L is a substituted or unsubstituted, saturated or unsaturated C₅₋₆ hydrocarbon chain, wherein one or two chain atoms of the hydrocarbon chain are independently replaced with –O–, –S–, or –NR^N–.


14. The method of any one of claims 3-13, or a pharmaceutically acceptable salt thereof, wherein L is of the formula:



15. The method of any one of claims 3-14, or a pharmaceutically acceptable salt thereof, wherein L is of the formula:



16. The method of any one of claims 3-14, or a pharmaceutically acceptable salt thereof, wherein L is of the formula:




17. The method of any one of claims 3-15, or a pharmaceutically acceptable salt thereof, wherein R^B is hydrogen.

18. The method of any one of claims 3-17, or a pharmaceutically acceptable salt thereof, wherein X^A is CR^X , and each of X^B and X^C is N.

19. The method of any one of claims 3-17, or a pharmaceutically acceptable salt thereof, wherein Y is $-NR^Y-$.

20. The method of claim 19, wherein Y is $-NH-$.


21. The method of any one of claims 3-20, or a pharmaceutically acceptable salt thereof, wherein Ring C is of the formula:



22. The method of any one of claims 3-21, or a pharmaceutically acceptable salt thereof, wherein each instance of R^C is independently halogen or unsubstituted C_{1-6} alkyl.


23. The method of claim 22, wherein at least one instance of R^C is chlorine or methyl.
24. The method of any one of claims 3-23, or a pharmaceutically acceptable salt thereof, wherein m is 2.
25. The method of any one of claims 3-24, wherein:
 Y is $-NR^Y-$;
 X^A is CR^X ; and
 R^Y and R^X of X^A are joined to form a substituted or unsubstituted, monocyclic, 5- to 7-membered heterocyclic ring that is fused with Ring B.

26. The method of any one of claims 3-25, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.


27. The method of claim 26, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof.




28. The method of any one of claims 3-27, or a pharmaceutically acceptable salt thereof, wherein the compound is of the formula:





, or a pharmaceutically acceptable salt thereof.


29. The method of any one of claims 3-27, or a pharmaceutically acceptable salt thereof, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof.

wherein:

R^A is substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl,



each instance of R^{A1} is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

each instance of R^b is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^b are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, or 4;

each instance of R^B is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

m is 0, 1, 2, 3, 4, or 5;

R^C is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^D is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^E is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^F is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Ring A is substituted or unsubstituted phenyl; substituted or unsubstituted, polycyclic aryl; substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl; or substituted or unsubstituted, polycyclic heteroaryl;


each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;



n is 0, 1, 2, 3, or 4, as valency permits;



L is a bond or a substituted or unsubstituted C_{1-6} hydrocarbon chain, optionally wherein one or more chain atoms of the hydrocarbon chain are independently replaced with $-C(=O)-$, $-O-$, $-S-$, $-NR^b-$, $-N=$, or $=N-$; and


R^H is substituted or unsubstituted C_{1-6} alkyl, substituted or unsubstituted heterocyclyl, $-OH$, or $-N(R^c)_2$, wherein each instance of R^c is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.

32. The method of claim 31, or a pharmaceutically acceptable salt thereof, wherein R^A is




33. The method of claim 32, wherein R^A is  or .



34. The method of any one of claims 32-33, wherein R^A is , , or



35. The method of any one of claims 32-34, or a pharmaceutically acceptable salt thereof, wherein R^B is halogen or methyl.
36. The method of claim 35, or a pharmaceutically acceptable salt thereof, wherein R^B is chlorine or methyl.
37. The method of any one of claims 32-36, or a pharmaceutically acceptable salt thereof, wherein Ring A is phenyl.
38. The method of any one of claims 32-36, or a pharmaceutically acceptable salt thereof, wherein Ring A is pyrazolyl or pyridinyl.
39. The method of any one of claims 31-38, or a pharmaceutically acceptable salt thereof, wherein L is a bond.

40. The method of any one of claims 31-39, or a pharmaceutically acceptable salt thereof, wherein L is an unsubstituted C₁₋₃ hydrocarbon chain, optionally wherein one or more chain atoms of the hydrocarbon chain are independently replaced with -O- or -NR^b-.

41. The method of claim 40, wherein L is of the formula: , wherein l^A indicates the point of attachment to Ring A, and l^R indicates the point of attachment to R^H.

42. The method of claim 41, wherein L is of the formula: , , or , wherein l^A indicates the point of attachment to Ring A, and l^R indicates the point of attachment to R^H.

43. The method of any one of claims 31-42, or a pharmaceutically acceptable salt thereof, wherein R^H is substituted or unsubstituted heterocyclyl.

44. The method of claim 43, or a pharmaceutically acceptable salt thereof, wherein R^H is substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted morpholinyl, or substituted or unsubstituted piperazinyl.

45. The method of any one of claims 31-44, or a pharmaceutically acceptable salt thereof, wherein R^H is -N(R^c)₂, and each instance of R^c is substituted or unsubstituted C₁₋₆ alkyl.

46. The method of 45, or a pharmaceutically acceptable salt thereof, wherein R^H is -N(Me)₂.

47. The method of any one of claims 31-46, or a pharmaceutically acceptable salt thereof, wherein R^C is hydrogen.

48. The method of any one of claims 31-47, or a pharmaceutically acceptable salt thereof, wherein R^D is hydrogen.

49. The method of any one of claims 31-48, or a pharmaceutically acceptable salt thereof, wherein R^E is hydrogen.

50. The method of any one of claims 31-49, or a pharmaceutically acceptable salt thereof, wherein R^F is hydrogen.

51. The method of any one of claims 31-50, or a pharmaceutically acceptable salt thereof, wherein R^C , R^D , R^E , and R^F are each hydrogen.


52. The method of any one of claims 31-51, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^G is substituted or unsubstituted C_{1-3} alkyl.


53. The method of claim 52, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^G is methyl, ethyl, or $-CF_3$.


54. The method of any one of claims 31-53, wherein at least one instance of R^G is $-OMe$, $-OEt$, or $-O(iPr)$.

55. The method of any one of claims 31-54, or a pharmaceutically acceptable salt thereof, wherein n is 0, 1, or 2.

56. The method of any one of claims 1 or 31-55, wherein the compound is of the formula:









or a pharmaceutically acceptable salt thereof.


57. The method of any one of claims 1 or 31-55, wherein the compound is of the formula:




(HG-11-139-01),



(HG-11-143-01),




(YKL-04-136-10),




(YKL-04-136-11),

or a pharmaceutically acceptable salt thereof.


58. The method of any one of claims 1 or 31-55, wherein the compound is of the formula:




(YKL-05-57)




(YKL-05-58)




(YKL-05-59)




(YKL-05-60)




(YKL-05-68)




(YKL-05-69)




(YKL-05-70)



(YKL-05-74)



(YKL-05-76)




(YKL-05-77)




(YKL-05-88)







(YKL-05-89)




(YKL-05-90)




(YKL-05-91)




(YKL-05-92)




(YKL-05-93)




(YKL-05-94)




(YKL-05-95)




(YKL-05-96)




(YKL-05-97)




(YKL-05-98)




(YKL-05-99)











(YKL-05-100)




(YKL-05-151)




(YKL-05-152)







(YKL-05-178)




(YKL-05-179)




(YKL-05-180)




(YKL-05-181)




(YKL-05-182)



(YKL-05-183)




(Example 2)




(YKL-04-136-1)

(SB1-D-01)




(SB1-D-09; YKL-04-136-6)




(SB1-D-10)

(YKL-04-136-10)




(SB1-D-11)

(YKL-04-136-8)




(YKL-06-038)

(SB1-D-40)




(YKL-06-039)


(SB1-D-42)




(YKL-06-040)




(YKL-06-044)




(YKL-06-045)




(YKL-06-051)




(YKL-06-054)



(YKL-06-055)




(YKL-06-056)




(YKL-06-057)

(SB1-D-43)




(YKL-06-077)

(SB1-D-57)




(YKL-06-078)

(SB1-D-58)




(YKL-06-080-1)

(SB1-D-60)



(YKL-06-081-1)

(SB1-D-61)




(YKL-06-082)


(SB1-D-62)

or a pharmaceutically acceptable salt thereof.

59. The method of any one of claims 1 or 31-55, wherein the compound is not of the formula:




(HG-11-136-01),



(HG-11-143-01),

or a pharmaceutically acceptable salt thereof.

60. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (III):



or a pharmaceutically acceptable salt thereof,

wherein:

R^J is substituted or unsubstituted carbocyclyl;

each instance of R^B is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, an

oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

each instance of R^b is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^b are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

m is 0, 1, 2, 3, 4, or 5;

R^C is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^D is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^E is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^F is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Ring A is substituted or unsubstituted phenyl; substituted or unsubstituted, polycyclic aryl; substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl; or substituted or unsubstituted, polycyclic heteroaryl;

each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^b)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^b)_2$, $-NO_2$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^a$, $-NR^bC(=O)N(R^b)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^b)_2$;

n is 0, 1, 2, 3, or 4, as valency permits;

R^K is unsubstituted methyl, substituted or unsubstituted heterocyclyl, $-OR^a$, or $-N(R^c)_2$, wherein each instance of R^c is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.

61. The method of claim 60, or a pharmaceutically acceptable salt thereof, wherein R^J is substituted or unsubstituted, 4- to 6-membered carbocyclyl.

62. The method of claim 61, or a pharmaceutically acceptable salt thereof, wherein R^J is substituted or unsubstituted cyclobutyl, substituted or unsubstituted cyclopentyl, or substituted or unsubstituted cyclohexyl.

63. The method of any one of claims 60-62, or a pharmaceutically acceptable salt thereof, wherein R^K is substituted or unsubstituted heterocyclyl.

64. The method of claim 63, or a pharmaceutically acceptable salt thereof, wherein R^K is substituted or unsubstituted piperidinyl or substituted or unsubstituted piperazinyl.


65. The method of any one of claims 60-64, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^G is $-OR^a$, wherein R^a is substituted or unsubstituted C_{1-6} alkyl.

66. The method of claim 65, wherein at least one instance of R^G is $-OMe$.

67. The method of any one of claims 60-66, or a pharmaceutically acceptable salt thereof, wherein m is 2.

68. The method of any one of claims 60-67, or a pharmaceutically acceptable salt thereof,


wherein Ring B is of the formula:



69. The method of claim 68, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^B is halogen or substituted or unsubstituted C_{1-6} alkyl.

70. The method of any one of claims 62 or 67-69, or a pharmaceutically acceptable salt

thereof, wherein Ring B is of the formula:



71. The method of any one of claims 60-70, or a pharmaceutically acceptable salt thereof, wherein R^C is hydrogen.


72. The method of any one of claims 60-71, or a pharmaceutically acceptable salt thereof, wherein R^D is hydrogen.

73. The method of any one of claims 60-72, or a pharmaceutically acceptable salt thereof, wherein R^E is hydrogen.

74. The method of any one of claims 60-73, or a pharmaceutically acceptable salt thereof, wherein R^F is hydrogen.


75. The method of any one of claims 60-74, or a pharmaceutically acceptable salt thereof, wherein n is 0.

76. The method of any one of claims 60-69 or 71-75, wherein Ring B is of the formula:




, R^C , R^D , R^E , and R^F are each hydrogen, R^G is $-OR^a$, R^J is substituted or unsubstituted, 4- to 6-membered carbocyclyl, and R^K is substituted or unsubstituted piperidinyl, or substituted or unsubstituted piperazinyl.

77. The method of any one of claims 60-69 or 71-75, wherein Ring B is of the formula:




, R^C , R^D , R^E , and R^F are each hydrogen, n is 0, R^J is substituted or unsubstituted, 4- to 6-membered carbocyclyl, and R^K is substituted or unsubstituted piperidinyl, or substituted or unsubstituted piperazinyl.


78. The method of any one of claims 60-77, or a pharmaceutically acceptable salt thereof, wherein the compound is of the formula:




or a pharmaceutically acceptable salt thereof.

79. The method of any one of claims 60-77, or a pharmaceutically acceptable salt thereof, wherein the compound is of the formula:






(YKL-06-076)




(YKL-06-089)



(YKL-06-091)


80. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (IV):



287

R^L is substituted or unsubstituted alkyl;

Ring C is unsubstituted phenyl or of the formula:



each instance of R^{B1} is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^d)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^d)R^a$, $-C(=NR^d)OR^a$, $-C(=NR^d)N(R^d)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^d)_2$, $-NO_2$, $-NR^dC(=O)R^a$, $-NR^dC(=O)OR^a$, $-NR^dC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^d)_2$;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom;

each instance of R^d is independently hydrogen, $-C(=O)R^a$, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^d are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

R^C is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^D is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^E is hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

R^F is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Ring A is substituted or unsubstituted phenyl; substituted or unsubstituted, polycyclic aryl; substituted or unsubstituted, 5- or 6-membered, monocyclic heteroaryl; or substituted or unsubstituted, polycyclic heteroaryl;

each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^b)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^b)R^a$, $-C(=NR^b)OR^a$, $-C(=NR^b)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-$

NO_2 , $-\text{NR}^b\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^b\text{C}(=\text{O})\text{OR}^a$, $-\text{NR}^b\text{C}(=\text{O})\text{N}(\text{R}^a)_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{OR}^a$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^b)_2$;

n is 0, 1, 2, 3, or 4, as valency permits;

R^k is unsubstituted methyl, substituted or unsubstituted heterocyclyl, $-\text{OR}^a$, or $-\text{N}(\text{R}^c)_2$, wherein each instance of R^c is independently hydrogen, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group, or optionally two instances of R^c are taken together with their intervening atoms to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;


R^y is substituted phenyl.

81. The method of claim 80, or a pharmaceutically acceptable salt thereof, wherein R^L is substituted or unsubstituted C_{1-6} alkyl.

82. The method of any one of claims 80-81, or a pharmaceutically acceptable salt thereof, wherein R^L is Me, or iPr.

83. The method of any one of claims 80-82, or a pharmaceutically acceptable salt thereof, wherein Ring C is unsubstituted phenyl.


84. The method of any one of claims 80-83, or a pharmaceutically acceptable salt thereof,



wherein Ring C is of the formula:


85. The method of claim 84, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^{B1} is halogen or substituted or unsubstituted C_{1-6} alkyl, or $-\text{N}(\text{R}^d)_2$, wherein each instance of R^d is hydrogen or $-\text{C}(=\text{O})\text{R}^a$.

86. The method of any one of claims 80-81, or 84, or a pharmaceutically acceptable salt




thereof, wherein Ring C is of the formula:

87. The method of any one of claims 80-81, or a pharmaceutically acceptable salt thereof, wherein Ring C is of the formula:



88. The method of claim 87, or a pharmaceutically acceptable salt thereof, wherein Ring C is of the formula:



89. The method of any one of claims 80-88, or a pharmaceutically acceptable salt thereof, wherein R^K is substituted or unsubstituted heterocyclyl.

90. The method of claim 89, or a pharmaceutically acceptable salt thereof, wherein R^K is substituted or unsubstituted tetrahydropyranyl, substituted or unsubstituted piperidinyl, substituted or unsubstituted morpholinyl, or substituted or unsubstituted piperazinyl.

91. The method of any one of claims 80-90, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^G is -OR^a, wherein R^a is substituted or unsubstituted C₁₋₆ alkyl.

92. The method of any one of claims 80-90, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^G is substituted or unsubstituted C₁₋₆ alkyl.


93. The method of any one of claims 80-90, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^G is -OMe or ethyl.

94. The method of any one of claims 80-93, or a pharmaceutically acceptable salt thereof, wherein each one of R^C and R^D is hydrogen.

95. The method of any one of claims 80-94, or a pharmaceutically acceptable salt thereof, wherein R^E is hydrogen.
96. The method of any one of claims 80-95, or a pharmaceutically acceptable salt thereof, wherein R^F is hydrogen.
97. The method of any one of claims 80-96, or a pharmaceutically acceptable salt thereof, wherein Ring A is substituted or unsubstituted phenyl.
98. The method of any one of claims 80-97, or a pharmaceutically acceptable salt thereof, wherein n is 0 or 1.
99. The method of any one of claims 80-98, or a pharmaceutically acceptable salt thereof, wherein R^C is hydrogen.
100. The method of any one of claims 80-99, or a pharmaceutically acceptable salt thereof, wherein R^D is hydrogen.
101. The method of any one of claims 80-100, or a pharmaceutically acceptable salt thereof, wherein R^E is hydrogen.
102. The method of any one of claims 80-101, or a pharmaceutically acceptable salt thereof, wherein R^F is hydrogen.
103. The method of any one of claims 80-102, wherein R^C , R^D , R^E , and R^F are each hydrogen.

292


105. The method of any one of claims 80-103, wherein the compound is of the formula:




(HG-11-23-01),

or a pharmaceutically acceptable salt thereof.

106. The method of any one of claims 80-103, wherein the compound is of the formula:




(YKL-06-084)




(YKL-06-085)

(YKL-06-086)




(YKL-06-087)



(HG-4-34-01),

or a pharmaceutically acceptable salt thereof.


107. The method of any one of claims 80-103, wherein the compound is not of the formula:



(HG-11-139-02),


or a pharmaceutically acceptable salt thereof.

108. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (V):



or a pharmaceutically acceptable salt thereof, wherein:

R^G is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heteroaryl, or of the formula:



each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^a)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$, or two R^A groups are joined to form a substituted or unsubstituted carbocyclic ring, substituted or unsubstituted heterocyclic ring, substituted or unsubstituted aryl ring, or substituted or unsubstituted heteroaryl ring;

each instance of R^a is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two R^a groups are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

k is 0, 1, 2, 3, 4, or 5;

R^B is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each of X^A , X^B , and X^C is independently N or CR^X , wherein each instance of R^X is independently hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^a)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$;

or: X^B is CR^X , and R^G and R^X of X^B are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

R^C is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^a)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$;

R^D is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or a nitrogen protecting group;


Y is $-O-$ or $-NR^Y-$, wherein R^Y is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

Z is a bond or $-C(R^Z)_2-$, wherein each instance of R^Z is independently hydrogen, halogen, or substituted or unsubstituted C_{1-6} alkyl;

each instance of R^E is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^a$, $-N(R^a)_2$, $-SR^a$, $-CN$, $-SCN$, $-C(=NR^a)R^a$, $-C(=NR^a)OR^a$, $-C(=NR^a)N(R^a)_2$, $-C(=O)R^a$, $-C(=O)OR^a$, $-C(=O)N(R^a)_2$, $-NO_2$, $-NR^aC(=O)R^a$, $-NR^aC(=O)OR^a$, $-NR^aC(=O)N(R^a)_2$, $-NR^aS(=O)R^a$, $-NR^aS(=O)OR^a$, $-NR^aS(=O)N(R^a)_2$, $-NR^aS(=O)_2R^a$, $-NR^aS(=O)_2OR^a$, $-NR^aS(=O)_2N(R^a)_2$, $-OC(=O)R^a$, $-OC(=O)OR^a$, or $-OC(=O)N(R^a)_2$; and


m is 0, 1, 2, 3, 4, or 5.

109. The method of claim 108, wherein the SIK inhibitor is of the formula:




or a pharmaceutically acceptable salt thereof.

110. The method of any one of claims 108-109, wherein the SIK inhibitor is of the formula:



or a pharmaceutically acceptable salt thereof.


111. The method of any one of claims 108-110, wherein the SIK inhibitor is of the formula:



or a pharmaceutically acceptable salt thereof.

112. The method of any one of claims 108-110, or a pharmaceutically acceptable salt thereof, wherein R^B is hydrogen.


113. The method of any one of claims 108-110 or 112, or a pharmaceutically acceptable





salt thereof, wherein Ring A is of the formula:

114. The method of claim 113, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^A is substituted or unsubstituted, 5- to 6-membered, monocyclic heterocyclic ring, wherein one, two, or three atoms in the heterocyclic ring system are independently oxygen, nitrogen, or sulfur.

115. The method of claim 114, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^A is of the formula:



116. The method of any one of claims 114-115, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^A is of the formula:





117. The method of any one of claims 108-116, or a pharmaceutically acceptable salt thereof, wherein k is 1.


118. The method of any one of claims 108-110 or 112-117, or a pharmaceutically acceptable salt thereof, wherein R^C is hydrogen.

119. The method of any one of claims 108-110 or 112-1181, or a pharmaceutically acceptable salt thereof, wherein R^D is substituted or unsubstituted phenyl.

120. The method of 119, or a pharmaceutically acceptable salt thereof, wherein R^D is of the




121. The method of any one of claims 119-120, or a pharmaceutically acceptable salt



thereof, wherein R^D is of the formula: .

122. The method of any one of claims 108-110 or 112-121, or a pharmaceutically acceptable salt thereof, wherein $-Y-Z-$ is $-NR^Y-$.


123. The method of any one of claims 108-110 or 112-122, or a pharmaceutically



acceptable salt thereof, wherein Ring C is of the formula: .

124. The method of claim 123, or a pharmaceutically acceptable salt thereof, wherein at least one instance of R^E is substituted or unsubstituted alkyl.


125. The method of claim 124, or a pharmaceutically acceptable salt thereof, wherein Ring



C is of the formula: .


126. The method of any one of claims 108-125, or a pharmaceutically acceptable salt thereof, wherein m is 2.

127. The method of any one of claims 108-126, or a pharmaceutically acceptable salt thereof, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof.



128. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (VI):




or a pharmaceutically acceptable salt thereof, wherein:

Ar₁ is a 5- or 6-membered hetero- or homo-cyclic aromatic ring optionally having a C₁-C₄ alkyl, or saturated heterocyclic or methyl-heterocyclic substituent;

X is separately N or CH;





129. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (VI-A):



or a pharmaceutically acceptable salt thereof, wherein:

Ar₁ is a 5- or 6-membered hetero- or homo-cyclic aromatic ring optionally having a C₁-C₄ alkyl, or saturated heterocyclic or methyl-heterocyclic substituent;




R₅ is H or a C₁-C₄ alkyl; or

R₃ and R₄ are joined together with the intervening atoms to form a pyrrolidine ring where one or both of the free carbons are substituted with an alkyl or oxygen-containing substituent.

130. The method of claim 128 or 129, or a pharmaceutically acceptable salt thereof, wherein Ar₁ is phenyl or saturated heterocyclic.

131. The method of claim 129, or a pharmaceutically acceptable salt thereof, wherein one instance of X is N and the other instance of X is CH.


132. The method of any one of claims 128 or 129, or a pharmaceutically acceptable salt



133. The method of any one of claims 129-132, or a pharmaceutically acceptable salt




134. The method of any one of claims 127-131, or a pharmaceutically acceptable salt




135. The method of any one of claims 127-133, or a pharmaceutically acceptable salt thereof, wherein R_3 and R_4 are joined together with the intervening atoms to form a pyrrolidine ring where both of the free carbons are substituted with optionally substituted C_{1-6} alkyl.

136. The method of any one of claims 128-135, or a pharmaceutically acceptable salt thereof, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof.

137. The method of claim 1 or 2, wherein the SIK inhibitor is a compound of Formula (VII):



or a pharmaceutically acceptable salt thereof, wherein:

each instance of R^A is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{A1}$, $-N(R^{A1})_2$, $-SR^{A1}$, $-CN$, $-SCN$, $-C(=NR^{A1})R^{A1}$, $-C(=NR^{A1})OR^{A1}$, $-C(=NR^{A1})N(R^{A1})_2$, $-C(=O)R^{A1}$, $-C(=O)OR^{A1}$, $-C(=O)N(R^{A1})_2$, $-NO_2$, $-NR^{A1}C(=O)R^{A1}$, $-NR^{A1}C(=O)OR^{A1}$, $-NR^{A1}C(=O)N(R^{A1})_2$, $-OC(=O)R^{A1}$, $-OC(=O)OR^{A1}$, or $-OC(=O)N(R^{A1})_2$, wherein each instance of R^{A1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{A1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

j is 0, 1, or 2;

R^B is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or a nitrogen protecting group;

R^C is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^D is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{D1}$, $-N(R^{D1})_2$, $-SR^{D1}$, $-CN$, $-SCN$, $-C(=NR^{D1})R^{D1}$, $-C(=NR^{D1})OR^{D1}$, $-C(=NR^{D1})N(R^{D1})_2$, $-C(=O)R^{D1}$, $-C(=O)OR^{D1}$, $-C(=O)N(R^{D1})_2$, $-NO_2$, $-NR^{D1}C(=O)R^{D1}$, $-NR^{D1}C(=O)OR^{D1}$, $-NR^{D1}C(=O)N(R^{D1})_2$, $-OC(=O)R^{D1}$, $-OC(=O)OR^{D1}$, or $-OC(=O)N(R^{D1})_2$, wherein each instance of R^{D1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or

a sulfur protecting group when attached to a sulfur atom, or two instances of R^{D1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;


m is 0, 1, or 2;








R^E is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^F is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{F1}$, $-N(R^{F1})_2$, $-SR^{F1}$, $-CN$, $-SCN$, $-C(=NR^{F1})R^{F1}$, $-C(=NR^{F1})OR^{F1}$, $-C(=NR^{F1})N(R^{F1})_2$, $-C(=O)R^{F1}$, $-C(=O)OR^{F1}$, $-C(=O)N(R^{F1})_2$, $-NO_2$, $-NR^{F1}C(=O)R^{F1}$, $-NR^{F1}C(=O)OR^{F1}$, $-NR^{F1}C(=O)N(R^{F1})_2$, $-OC(=O)R^{F1}$, $-OC(=O)OR^{F1}$, or $-OC(=O)N(R^{F1})_2$, wherein each instance of R^{F1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{F1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring; and

n is 0, 1, 2, 3, 4, or 5;


provided that the compound is not of the formula:





or a pharmaceutically acceptable salt thereof.

138. The method of claim 137, wherein the compound is of Formula (VII-A):




or a pharmaceutically acceptable salt thereof, wherein:

each instance of R^G is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{G1}$, $-N(R^{G1})_2$, $-SR^{G1}$, $-CN$, $-SCN$, $-C(=NR^{G1})R^{G1}$, $-C(=NR^{G1})OR^{G1}$, $-C(=NR^{G1})N(R^{G1})_2$, $-C(=O)R^{G1}$, $-C(=O)OR^{G1}$, $-C(=O)N(R^{G1})_2$, $-NO_2$, $-NR^{G1}C(=O)R^{G1}$, $-NR^{G1}C(=O)OR^{G1}$, $-NR^{G1}C(=O)N(R^{G1})_2$, $-OC(=O)R^{G1}$, $-OC(=O)OR^{G1}$, or $-OC(=O)N(R^{G1})_2$, wherein each instance of R^{G1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{G1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring; and

k is 0, 1, 2, 3, 4, or 5.

139. The method of claim 137, wherein the compound is of Formula (VII-B):



or a pharmaceutically acceptable salt thereof, wherein:


R^H is hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted C_{1-6} alkyl, or a nitrogen protecting group;

each instance of R^J is independently halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{J1}$, $-N(R^{J1})_2$, $-SR^{J1}$, $-CN$, $-SCN$, $-C(=NR^{J1})R^{J1}$, $-C(=NR^{J1})OR^{J1}$, $-C(=NR^{J1})N(R^{J1})_2$, $-C(=O)R^{J1}$, $-C(=O)OR^{J1}$, $-C(=O)N(R^{J1})_2$, $-NO_2$, $-NR^{J1}C(=O)R^{J1}$, $-NR^{J1}C(=O)OR^{J1}$, $-NR^{J1}C(=O)N(R^{J1})_2$, $-OC(=O)R^{J1}$, $-OC(=O)OR^{J1}$, or $-OC(=O)N(R^{J1})_2$, wherein each instance of R^{J1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{J1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring;

q is 0, 1, 2, 3, or 4; and


R^K is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, $-OR^{K1}$, $-N(R^{K1})_2$, $-SR^{K1}$, $-CN$, $-SCN$, $-C(=NR^{K1})R^{K1}$, $-C(=NR^{K1})OR^{K1}$, $-C(=NR^{K1})N(R^{K1})_2$, $-C(=O)R^{K1}$, $-C(=O)OR^{K1}$, $-C(=O)N(R^{K1})_2$, $-NO_2$, $-NR^{K1}C(=O)R^{K1}$, $-NR^{K1}C(=O)OR^{K1}$, $-NR^{K1}C(=O)N(R^{K1})_2$, $-OC(=O)R^{K1}$, $-OC(=O)OR^{K1}$, or $-OC(=O)N(R^{K1})_2$, wherein each instance of R^{K1} is independently hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted carbocyclyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, a nitrogen protecting group when attached to a nitrogen atom, an oxygen protecting group when attached to an oxygen atom, or a sulfur protecting group when attached to a sulfur atom, or two instances of R^{K1} are joined to form a substituted or unsubstituted heterocyclic or substituted or unsubstituted heteroaryl ring.

140. The method of claim 139, wherein when R^C is hydrogen, R^B is not unsubstituted



141. The method of claim 139 or 140, wherein R^H is hydrogen.

142. The method of any one of claims 139-141, wherein Ring E is of the formula:



143. The method of any one of claims 139-142, wherein at least one instance of R^J is halogen or substituted or unsubstituted C_{1-6} alkyl.

144. The method of any one of claims 139-143, wherein R^K is hydrogen.

145. The method of any one of claims 139-144, wherein R^K is substituted or unsubstituted C_{1-6} alkyl.





146. The method of any one of claims 139-145, wherein at least one instance of R^A is halogen or substituted or unsubstituted C_{1-6} alkyl.







147. The method of any one of claims 139-146, wherein R^B is substituted or unsubstituted alkyl.

148. The method of any one of claims 139-147, wherein R^C is hydrogen.

149. The method of any one of claims 139-148, wherein at least one instance of R^D is halogen or substituted or unsubstituted C_{1-6} alkyl.

150. The method of any one of claims 139-149, wherein m is 0.
151. The method of any one of claims 139-150, wherein R^E is hydrogen.
152. The method of any one of claims 139-151, wherein at least one instance of R^F is halogen, substituted or unsubstituted C_{1-6} alkyl, or $-OR^{F1}$, wherein R^{F1} is hydrogen, substituted or unsubstituted C_{1-6} alkyl, or an oxygen protecting group.
153. The method of any one of claims 139-152, wherein n is 1.
154. The method of any one of claims 139-152, wherein n is 2.
155. The method of any one of claims 139-154, wherein the compound of Formula (VII) is of the formula:





or a pharmaceutically acceptable salt thereof.

156. The method of any one of claims 1-155, comprising administering to the subject a compound of Formulae (I)-(VII), a pharmaceutically acceptable salt thereof, and optionally a pharmaceutically acceptable excipient.

157. The method of any one of claims 1-155, comprising administering to the subject a compound of Formulae (I)-(VII), a pharmaceutically acceptable salt thereof, or a topical formulation thereof.

158. The method of any one of claims 1-157, or a pharmaceutically acceptable salt thereof, wherein the subject is a human.

159. The method of any one of claims 1-158, wherein the step of administering comprises administering the SIK inhibitor topically to the subject's skin on a body part.

160. The method of claim 159, wherein the body part comprises the face, neck, chest, back, arms, torso, or legs of the subject.

161. Use of a compound for increasing skin pigmentation, reducing the risk of skin cancer, and/or inducing microphthalmia-associated transcription factor expression in a subject, wherein the compound comprises: a compound of Formulae (I)-(VII), or a pharmaceutically acceptable salt thereof.

1/30




Figure 1A




Figure 1B

2/30




Figure 1C




Figure 1D

3/30




Figure 2A




Figure 2B

4/30




Figure 2C




Figure 2D

5/30




Figure 3A

6/30




Figure 3B




Figure 3C

7/30




Figure 4A




Figure 4B

8/30




Figure 4C






Figure 4D

9/30

Vehicle HG 9-91-01



YKL 06-061 YKL 06-062



Vehicle

HG 9-91-01

YKL 06-061

YKL 06-062

Figure 5A




Figure 5B

10/30




Figure 5C




Figure 5D




Figure 5E

11/30






Figure 5F


12/30

**YKL-06-062**

SIK1/2/3: 2.12/1.40/2.86

**YKL-06-061**

SIK1/2/3: 6.56/1.77/20.5 nM

**HG-9-91-01**

SIK1/2/3: 4.29/4.02/8.37 nM

Figure 6A

YKL-06-061
468 Assays Tested
16 Interactions Mapped
S-Score(1) = 0.04





Figure 6B

13/30

Kinase	% control at 1 μ M	Enzymatic IC ₅₀ (nM)
CSF1R	0	9.66
FRK	0	1.10
KIT	0	153
p38-alpha	0	10.1
p38-beta	0	9.64
SRC	0.15	58.8
BRK	0.2	24.1
PDGFRB	0.25	103
EPHB1	0.3	16.4
SIK1	0.3	6.56
TNK2	0.4	10.5
RSK4	0.6	>10000
SIK2	0.7	1.77
NLK	0.95	132
PRKR	1	>10000

Figure 6C



14/30

Figure 7C




Figure 7B




Figure 7A




Figure 7F




Figure 7E




Figure 7D

15/30




Figure 7G




Figure 7H

16/30




Figure 8A




Figure 8B

17/30




Figure 8C




Figure 8D




Figure 8E

19/30




Figure 9

20/30




Figure 9 (Continued)

21/30






Figure 9 (Continued)

22/30

Dose Response YKL04115



Dose Response YKL04118



Dose Response YKL04125






Figure 9 (Continued)

23/30

Dose Response YKL04136-01



Dose Response YKL04136-02



Dose Response YKL04136-03




Figure 9 (Continued)

24/30




Figure 9 (Continued)

25/30




Figure 9 (Continued)

26/30




Figure 9 (Continued)

27/30




Figure 9 (Continued)

28/30




Figure 10

29/30




Figure 11




Figure 12

