



- (51) **International Patent Classification:**
C07D 498/22 (2006.01) *C07D 489/08* (2006.01)
- (21) **International Application Number:**
PCT/CA2012/000384
- (22) **International Filing Date:**
2 May 2012 (02.05.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/481,359 2 May 2011 (02.05.2011) US
- (71) **Applicant (for all designated States except US):** **BROCK UNIVERSITY** [CA/CA]; 500 Glenridge Avenue, St. Catharines, Ontario L2S 3A1 (CA).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **HUDLICKY, Tomas** [US/CA]; 173 Lockhart Drive, St. Catharines, Ontario L2T 1W7 (CA). **WERNER, Lukas** [CZ/CZ]; Chomutovska 1229, 43201 Kadan (CZ). **WERNEROVA, Martina** [SK/SK]; Jurkova Vola 35, 08901 Svidnik (SK). **ENDO-MA-ARIAS, Mary Ann** [PH/PH]; c/o Institute of Chemistry, Quirino Street, University of Philippines, Diliman, Quezon City 1101 (PH). **MACHARA, Ales** [CZ/CZ]; Anastazova 21, Praha 6, 16900 Brevnov (CZ).
- (74) **Agent:** **BERESKIN & PARR LLP/S.E.N.C.R.L., s.r.l.**; 40 King Street West, 40th Floor, Toronto, Ontario M5H 3Y2 (CA).

- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** PROCESSES AND INTERMEDIATES IN THE PREPARATION OF MORPHINE ANALOGS VIA N-DEMETHYLATION OF N-OXIDES USING CYCLODEHYDRATION REAGENTS

(57) **Abstract:** A high-yielding method for the N-demethylation of oxycodone- and oxymorphone-N-oxides by the reaction of these compounds with cyclodehydration reagents has been performed. This method has been utilized to improve the synthesis of various morphine analogs, such as naltrexone, nalbuphine and naloxone.



TITLE: PROCESSES AND INTERMEDIATES IN THE PREPARATION OF MORPHINE ANALOGS VIA N-DEMETHYLATION OF N-OXIDES USING CYCLODEHYDRATION REAGENTS

CROSS-REFERENCE TO RELATED APPLICATIONS


[0001] The present application claims the benefit of priority from co-pending U.S. provisional application no. 61/481,359 filed on May 2, 2011, the contents of which are incorporated herein by reference in their entirety.

FIELD OF THE APPLICATION

[0002] The present application relates to new processes and intermediates useful in the preparation of morphine analogs, such as naltrexone, naloxone and nalbuphine. In a particular example, the process begins with oxymorphone or oxycodone N-oxides and includes the formation of an oxazolidine intermediate using a cyclodehydration reagent.

BACKGROUND OF THE APPLICATION

[0003] Various morphine antagonists such as naltrexone, naloxone, and nalbuphine are available by semi-synthesis from the natural opiates such as morphine, codeine, thebaine or oripavine, Scheme 1. These compounds are used extensively in medicine as antagonists (naltrexone and naloxone) and mixed agonist/antagonist (nalbuphine). Naltrexone has long been used for the treatment of alcoholism, and is the active ingredient in Vivitrol®, an extended release injectable suspension for the treatment of alcoholism and opioid dependence. Naloxone is the active ingredient in Narcan® for the reversal of opioid overdose and is used to mitigate side effects in combination with buprenorphine (Suboxone®) for the treatment of opioid addiction, with tilidine (Valoron N®) for the treatment of pain and with oxycodone (Targin®) for the prophylaxis and/or treatment of opioid-induced bowel dysfunction during the treatment of pain. Nalbuphine is the active ingredient in Nubain® and is used for the treatment of pain in very low doses particularly in women.




Scheme 1

[0004] The introduction of the C-14 hydroxyl into various natural morphinans to produce oxycodone and oxymorphone has been reduced to practice on large scales with a high degree of efficiency by oxidation of thebaine or oripavine. Methods for direct C-H oxidation at C-14 for compounds such as codeine, morphine, or hydrocodone have been reported but are not very efficient or practical at this time. On the other hand, N-demethylation of natural opiates still represents a challenge, especially in terms of efficiency or the focus on environmentally benign procedures and reagents. Many methods have been employed for the demethylation; these include the use of cyanogen bromide (von Braun reaction),ⁱ methyl or ethyl chloroformate,ⁱⁱ 1-chloroethyl chloroformate (ACE-Cl),ⁱⁱⁱ and microbial protocols,^{iv} including a recently published procedure employing fungal biotransformations.^v The biotransformations of several morphine alkaloids with the strain *Cunninghamella echinulata* and several others produced the free amines in reasonable yields and purity. Such processes, when scaled up and improved by the creation of a transgenic vector that would express the required fungal cytochrome in an *E. coli* carrier would have great potential as an environmentally benign N-demethylation protocol.

[0005] Recently, iron (II) as well as iron (0) catalyzed N-demethylation of several morphinan N-oxides was reported by Scammells.^{vi} Smith et al.^{vii} developed a method to convert N-methylated 6-oxo-14-hydroxymorphinanes to the corresponding nor compounds by treating the corresponding N-oxide with a Fe(II) based reducing agent in the presence of formic acid to form an

oxazolidine. The oxazolidine can be converted to the corresponding normorphinane by acid hydrolysis, as shown in Scheme 2. Conversion of the *N*-oxide to the corresponding oxazolidine works equally well whether the 7,8 carbon bond is unsaturated or saturated, as shown with oxymorphone, Scheme 2.



Scheme 2


[0006] The reactivity of the Burgess reagent, long associated only with the dehydration of alcohols, has been tested in a variety of ways with other functional groups. The synthesis of cis-fused sulfamidates was accomplished by the reaction of the Burgess reagent with epoxides^{viii} and 1,2-diols,^{ix} and the Burgess reagent was shown to oxidize thiols to disulfides in high yields.^x New applications^{xi} as well as more thermally stable forms of this reagent^{xii} are being reported, including its chiral auxiliary version,^{xiii} and the reagent has been used extensively in natural product syntheses.^{xiv}

SUMMARY OF THE APPLICATION

[0007] Because the conversion of natural opiates to their C-14 hydroxy derivatives is well established, it would be convenient to provide a direct conversion of oxymorphone to the corresponding analogs via *N*-demethylation and alkylation. The present application reports a rather unexpected reaction of a cyclodehydration reagent, such as Burgess reagent, with *N*-oxides derived from, for example, oxymorphone and oxycodone to provide the corresponding


oxazolidine, and an efficient conversion of these oxazolidines to naltrexone, nalbuphine, naloxone, and other analogs.

[0008] Accordingly, the present application includes a process for the preparation of a compound of Formula I:



comprising:

(a) reacting a compound of Formula II with an oxidizing agent under conditions to provide a compound of Formula III:



and

(b) reacting the compound of Formula III with a cyclodehydration reagent under conditions to provide the compound of Formula I,

wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;


R¹ and R² are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when --O represents =O, then R² is not present; and

PG is a protecting group; and

wherein in the compounds of Formulae I, II and III, one or more available hydrogens in R¹ and R² is/are optionally replaced with F and/or one or more of available atoms in R¹ and R² is/are optionally replaced with an isotopic label.

[0009] The compounds of Formula I are useful for the preparation of several different classes of morphine analogs.

[0010] For example, in one embodiment, the present application also includes a process for preparing compounds of Formula V:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R¹ and R² are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when ---O represents =O, then R² is not present;


PG is a protecting group;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl; and

X is a counteranion,

comprising

reacting a compound of Formula I:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when $=O$ represents $=O$, then R^2 is not present; and PG is a protecting group, with an alkylating reagent of Formula VI:




wherein R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl and LG is a leaving group, under conditions to provide the compound of Formula V,

wherein in the compounds of Formulae I, V and VI, one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label.

[0011] The oxazolidine ring in the compounds of Formula V can be cleaved under reducing or hydrolysis conditions to provide further morphine analogs. For example, the reduction of the compounds of Formula V provides the corresponding 14-O-methylated compounds or 14-OH compounds. Hydrolysis of the compounds of Formula V provides the corresponding 14-OH, 17-NH compounds.

[0012] As a further example of the use of the compounds of Formula I in the preparation of morphine analogs, hydrolysis of the compounds of Formula I under either acidic or basic conditions provides the free phenols of Formula X:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^8 and R^9 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents $=O$, then R^9 is not present; or


R^8 and R^9 are H, if in the compound of Formula I, R^1 and R^2 are a PG that is removed under the hydrolysis conditions; and

PG is a protecting group,

wherein in the compounds of Formula X, one or more available hydrogens in R^8 and R^9 is/are optionally replaced with F and/or one or more of available atoms in R^8 and R^9 is/are optionally replaced with an isotopic label.

[0013] Compounds of Formula X can be selectively alkylated at the 17-N to provide a wide variety of morphine analogs.

[0014] The present application includes compounds of Formula V:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents $=O$, then R^2 is not present;


PG is a protecting group;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label, or a salt or solvate thereof.

[0015] The present application also includes compounds of Formula VII:




wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other and when =O represents =O , then H is not present; R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion; and

one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label, or a salt or solvate thereof.

[0016] The present application also includes compounds of Formula VIII:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl;

R⁴ and R⁵ are independently selected from H, C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when =O represents =O, then R⁵ is not present;

PG is a protecting group; and

one or more available hydrogens in R³, R⁴ and R⁵ is/are optionally replaced with F and/or one or more of available atoms in R³, R⁴ and R⁵ is/are optionally replaced with an isotopic label, or
a salt or solvate thereof.

[0017] Other features and advantages of the present application will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating embodiments of the application are given by way of illustration only, since various changes and modifications within the spirit and scope of the application will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE APPLICATION

I. Definitions

[0018] Unless otherwise indicated, the definitions and embodiments described in this and other sections are intended to be applicable to all embodiments and aspects of the application herein described for which they are suitable as would be understood by a person skilled in the art.

[0019] As used in this application, the singular forms “a”, “an” and “the” include plural references unless the content clearly dictates otherwise. For example, an embodiment including “an oxidizing agent” should be understood to present certain aspects with one oxidizing agent, or two or more additional oxidizing agents.

[0020] In embodiments comprising an “additional” or “second” component, such as an additional or second oxidizing agent, the second

component as used herein is chemically different from the other components or first component. A "third" component is different from the other, first, and second components, and further enumerated or "additional" components are similarly different.

[0021] The term "suitable" as used herein means that the selection of the particular compound or conditions would depend on the specific synthetic manipulation to be performed, and the identity of the molecule(s) to be transformed, but the selection would be well within the skill of a person trained in the art. All process/method steps described herein are to be conducted under conditions sufficient to provide the product shown. A person skilled in the art would understand that all reaction conditions, including, for example, reaction solvent, reaction time, reaction temperature, reaction pressure, reactant ratio and whether or not the reaction should be performed under an anhydrous or inert atmosphere, can be varied to optimize the yield of the desired product and it is within their skill to do so.

[0022] In embodiments of the application, the compounds described herein have at least one asymmetric centre. Where compounds possess more than one asymmetric centre, they may exist as diastereomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present application. It is to be further understood that while the stereochemistry of the compounds may be as shown in any given compound listed herein, such compounds may also contain certain amounts (e.g. less than 20%, suitably less than 10%, more suitably less than 5%) of compounds of the application having alternate stereochemistry.


[0023] In understanding the scope of the present disclosure, the term "comprising" and its derivatives, as used herein, are intended to be open ended terms that specify the presence of the stated features, elements, components, groups, integers, and/or steps, but do not exclude the presence of other unstated features, elements, components, groups, integers and/or steps. The foregoing also applies to words having similar meanings such as

the terms, "including", "having" and their derivatives. The term "consisting" and its derivatives, as used herein, are intended to be closed terms that specify the presence of the stated features, elements, components, groups, integers, and/or steps, but exclude the presence of other unstated features, elements, components, groups, integers and/or steps. The term "consisting essentially of", as used herein, is intended to specify the presence of the stated features, elements, components, groups, integers, and/or steps as well as those that do not materially affect the basic and novel characteristic(s) of features, elements, components, groups, integers, and/or steps.

[0024] Terms of degree such as "substantially", "about" and "approximately" as used herein mean a reasonable amount of deviation of the modified term such that the end result is not significantly changed. These terms of degree should be construed as including a deviation of at least $\pm 5\%$ of the modified term if this deviation would not negate the meaning of the word it modifies.

[0025] The term "cyclodehydration reagent" as used herein refers to a reagent that facilitates the cyclization of a compound of Formula III to a compound of Formula I via the loss of one equivalent of H_2O under suitable conditions. The selection of a suitable cyclodehydration reagent can be made by a person skilled in the art. In an embodiment of the application, the cyclodehydration reagent is selected from Burgess reagent, TsCl, CrO_3 , DCC, XtalFluorTM and carbonyldiimidazole. It is an embodiment that the cyclodehydration reagent is Burgess reagent.

[0026] The term "Burgess reagent" as used herein refers to a reagent of the formula:



also known as methyl *N*-(triethylammoniumsulfonyl)carbamate. This reagent is commercially available (for example from Sigma Aldrich, St. Louis, MO,

USA) or may be prepared from chlorosulfonylisocyanate by treatment with methanol, followed by triethylamine in benzene.^{xv}

[0027] The term “counteranion” as used herein refers to a negatively charged species consisting of a single element, or a negatively charged species consisting of a group of elements connected by ionic and/or covalent bonds.

[0028] The term “acyl” as used herein, whether it is used alone or as part of another group, means straight or branched chain, saturated acyl groups. The term C₁₋₆acyl means an acyl group having 1, 2, 3, 4, 5 or 6 carbon atoms (i.e. C(O)C₁₋₅alkyl). It is an embodiment of the application that, in the acyl groups, one or more, including all of the available hydrogen atoms are optionally replaced with F or ²H and thus include, for example trifluoroacetyl and the like.

[0029] The term “alkyl” as used herein, whether it is used alone or as part of another group, means straight or branched chain, saturated alkyl groups. The term C₁₋₆alkyl means an alkyl group having 1, 2, 3, 4, 5, or 6 carbon atoms. It is an embodiment of the application that, in the alkyl groups, one or more, including all, of the hydrogen atoms are optionally replaced with F or ²H and thus include, for example trifluoromethyl, pentafluoroethyl and the like.

[0030] The term “alkylene” as used herein, whether it is used alone or as part of another group, refers to a bivalent alkyl group.

[0031] The term “alkenyl” as used herein, whether it is used alone or as part of another group, means straight or branched chain, unsaturated alkenyl groups. The term C₂₋₆alkenyl means an alkenyl group having 2, 3, 4, 5, or 6 carbon atoms and at least one double bond. It is an embodiment of the application that, in the alkenyl groups, one or more, including all, of the hydrogen atoms are optionally replaced with F or ²H and thus include, for example trifluoroethenyl, pentafluoropropenyl and the like.

[0032] The term "cycloalkyl" as used herein, whether it is used alone or as part of another group, means cyclic, saturated alkyl groups. The term C₃₋₁₀cycloalkyl means a cycloalkyl group having 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms. It is an embodiment of the application that, in the cycloalkyl groups, one or more, including all, of the hydrogen atoms are optionally replaced with F or ²H.

[0033] The term "cycloalkenyl" as used herein, whether it is used alone or as part of another group, means cyclic, unsaturated alkyl groups. The term C₃₋₁₀cycloalkenyl means a cycloalkenyl group having 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms and at least one double bond. It is an embodiment of the application that, in the cycloalkenyl groups, one or more, including all, of the hydrogen atoms are optionally replaced with F or ²H.

[0034] The term "aryl" as used herein refers to cyclic groups that contain at least one aromatic ring. In an embodiment of the application, the aryl group contains 6, 9 or 10 atoms, such as phenyl, naphthyl or indanyl. It is an embodiment of the application that, in the aryl groups, one or more, including all, of the hydrogen atoms are optionally replaced with F or ²H and thus include, for example pentafluorophenyl and the like.

[0035] The term "halo" as used herein refers to a halogen atom and includes F, Cl, Br and I.

[0036] The term "oxidizing agent" as used herein means any compound or combination of compounds that oxidizes a desired functional group(s) but does not otherwise react with or degrade the substrate comprising the functional group(s). An oxidizing agent results in the overall loss of electrons, or in the case of organic chemistry, hydrogen atoms from the functional group.

[0037] The term "reducing agent" as used herein means any compound or combination of compounds that reduces a desired functional group(s) but does not otherwise react with or degrade the substrate comprising the functional group(s). A reducing agent results in the overall gain of electrons, or in the case of organic chemistry, hydrogen atoms to the functional group. It

is an embodiment of the application that the reducing agent is a metal hydride reducing agent.

[0038] The term “inert solvent” as used herein means a solvent that does not interfere with or otherwise inhibit a reaction. Accordingly, the identity of the inert solvent will vary depending on the reaction being performed. The selection of inert solvent is within the skill of a person in the art. Examples of inert solvents include, but are not limited to, benzene, toluene, tetrahydrofuran, ethyl ether, ethyl acetate, dimethyl formamide (DMF), acetonitrile, C₁₋₆alkylOH (e.g. methanol, ethanol, n-propanol, 2-propanol, n-butanol, butan-2-ol and 2-methyl-1-propanol), diethylcarbonate, hexane and dimethylsulfoxide (DMSO). Further examples, can include aqueous solutions, such as water and dilute acids and bases, and ionic liquids, provided that such solvents do not interfere with the reaction.

[0039] The term “solvent” includes both a single solvent and a mixture comprising two or more solvents.

[0040] The term “available”, as in “available hydrogen atoms” or “available atoms” refers to atoms that would be known to a person skilled in the art to be capable of replacement by either a fluorine atom (in the case of hydrogen atoms) or isotopic labels (in the case of all atoms) using methods known in the art.

[0041] *t*-Boc as used herein refers to the group *t*-butoxycarbonyl.

[0042] Ac as used herein refers to the group acetyl.

[0043] Ts (tosyl) as used herein refers to the group *p*-toluenesulfonyl

[0044] Ms as used herein refers to the group methanesulfonyl

[0045] TBDMS as used herein refers to the group *t*-butyldimethylsilyl.

[0046] TBDPS as used herein refers to the group *t*-butyldiphenylsilyl.

[0047] TMS as used herein refers to the group trimethylsilyl.

[0048] Tf as used herein refers to the group trifluoromethanesulfonyl.

[0049] Ns as used herein refers to the group naphthalene sulphonyl.

[0050] Bn as used herein refers to the group benzyl.

[0051] Fmoc as used herein refers to the group fluorenylmethoxycarbonyl.

[0052] mCPBA as used herein refers to meta-chloroperbenzoic acid.

[0053] The term "leaving group" or "LG" as used herein refers to a group that is readily displaceable by a nucleophile, for example, under nucleophilic substitution reaction conditions. Examples of suitable leaving groups include, but are not limited to, halo, Ms, Ts, Ns, Tf, C₁₋₆acyl, and the like.


[0054] The terms "protective group" or "protecting group" or "PG" or the like as used herein refer to a chemical moiety which protects or masks a reactive portion of a molecule to prevent side reactions in those reactive portions of the molecule, while manipulating or reacting a different portion of the molecule. After the manipulation or reaction is complete, the protecting group is removed under conditions that do not degrade or decompose the remaining portions of the molecule. The selection of a suitable protecting group can be made by a person skilled in the art. Many conventional protecting groups are known in the art, for example as described in "Protective Groups in Organic Chemistry" McOmie, J.F.W. Ed., Plenum Press, 1973, in Greene, T.W. and Wuts, P.G.M., "Protective Groups in Organic Synthesis", John Wiley & Sons, 3rd Edition, 1999 and in Kocienski, P. Protecting Groups, 3rd Edition, 2003, Georg Thieme Verlag (The Americas). Examples of suitable protecting groups include, but are not limited to t-Boc, Ac, Ts, Ms, silyl ethers such as TMS, TBDMS, TBDPS, Tf, Ns, Bn, Fmoc, dimethoxytrityl, methoxyethoxymethyl ether, methoxymethyl ether, pivaloyl, p-methoxybenzyl ether, tetrahydropyranyl, trityl, ethoxyethyl ethers, carbobenzyloxy, benzoyl and the like.

[0055] The expression "proceed to a sufficient extent" as used herein with reference to the reactions or process steps disclosed herein means that

the reactions or process steps proceed to an extent that conversion of the starting material or substrate to product is maximized. Conversion may be maximized when greater than about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% of the starting material or substrate is converted to product.


II. Methods of the Application

[0056] The present application includes a process for the preparation of a compound of Formula I:



comprising:

(a) reacting a compound of Formula II with an oxidizing agent under conditions to provide a compound of Formula III:



and

(b) reacting the compound of Formula III with a cyclodehydration reagent under conditions to provide the compound of Formula I,

wherein

---- represents a single or double bond, provided that two double bonds are not adjacent to each other;


R¹ and R² are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when =O represents =O, then R² is not present; and

PG is a protecting group; and

wherein in the compounds of Formulae I, II and III, one or more available hydrogens in R^1 and R^2 is/are optionally replaced with F and/or one or more of available atoms in R^1 and R^2 is/are optionally replaced with an isotopic label.

[0057] In an embodiment of the application, R^1 and R^2 are independently selected from C_{1-6} alkyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{1-6} alkylene C_{6-10} aryl, C_{1-6} alkylene C_{3-6} cycloalkyl and PG. In a further embodiment of the application, R^1 and R^2 are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG. It is an embodiment of the application that PG is an alkyl acetate, such as acetyl.


[0058] In an embodiment, the compound of Formula II is selected from a compound of Formula II(a), II(b) and II(c):



wherein

--- represents a single or double bond;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, and PG is a protecting group, which provide, respectively, a compound of the Formula I(a), I(b) and I(c) using the process of the present application:



wherein

---- represents a single or double bond;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, and PG is a protecting group; and

wherein in the compounds of Formulae I(a), I(b), I(c), II(a), II(b) and II(c), one or more available hydrogens in R^1 and R^2 is/are optionally replaced with F and/or one or more of available atoms in R^1 and R^2 is/are optionally replaced with an isotopic label.

[0059] Oxidation of the compounds of Formula II to the compounds of Formula III is performed using any suitable oxidizing agent. In an embodiment of the application, the oxidizing agent is a peroxide or a peracid. In another embodiment, the peracid is *m*-chloroperbenzoic acid (*m*CPBA). Conditions to oxidize tertiary amines to the corresponding N-oxides are known in the art. Other exemplary oxidizing agents include hydrogen peroxide, peracetic acid, *t*-butylhydroperoxide and magnesium monoperoxyphthalate.


[0060] In an embodiment of the application, the cyclodehydration reagent is Burgess reagent. Other cyclodehydration agents were examined in the place of the Burgess reagent. Thus oxycodone-*N*-oxide also yielded oxazolidine on treatment with TsCl (30%), CrO_3 (44%) and DCC (50%). Treatment of the *N*-oxide with CS_2 or SeO_2 resulted only in its re-conversion to oxycodone. Other reagents that may be used in place of Burgess reagent are XtalFluorTM and carbonyldiimidazole, both of which are commercially available, for example from Sigma-Aldrich, USA.

[0061] In an embodiment of the application, the conditions to provide the compounds of Formula I from the compounds of Formula III using a cyclodehydration reagent comprise a temperature of about -50 °C to about 50 °C, in an inert solvent or mixture of solvents for a time for the conversion of the compound of Formula III to the compound of Formula I to proceed to a sufficient extent, for example from about 0.5 hours to about 48 hours, or about 2 hours to about 10 hours. In an embodiment, the molar ratio of

cyclodehydration reagent to the compound of Formula III is about 1.5:1 to about 1:1.

[0062] In a representative example of the process of the present application, the reaction of oxycodone *N*-oxide, derived from oxycodone (compound of Formula II(b), wherein $R^1 = \text{Me}$ and --- is a single bond), with the Burgess reagent was examined and a clean conversion to the corresponding oxazolidine (compound of Formula I(b), wherein $R^1 = \text{Me}$ and -- is a single bond) was obtained, providing significantly higher yields than that quoted in the Smith procedure shown in Scheme 2.

[0063] The compounds of Formula I, wherein R^1 and/or R^2 are PG, can be deprotected to provide the corresponding free hydroxy compounds, that is, compounds of Formula IV:




wherein --- represents a single or double bond, provided that two double bonds are not adjacent to each other and when =O represents =O , then the H is not present.

[0064] As noted above, the compounds of Formula I are useful for the preparation of a variety of different morphine analogs:

(i) Quaternary Salts of the Compounds of Formula I

[0065] Compounds of Formula I have been converted to the corresponding quaternary salts by reaction with an alkylating reagent. Therefore, the present application also includes a process for preparing compounds of Formula V:

20



wherein

— represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^2 is not present;


PG is a protecting group;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl; and

X is a counteranion,

comprising

reacting a compound of Formula I:



wherein

— represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^2 is not present; and

PG is a protecting group,

with an alkylating reagent of Formula VI:



wherein

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl, and

LG is a leaving group, under conditions to provide the compound of Formula V,


wherein in the compounds of Formulae I, V and VI, one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label.

[0066] In an embodiment, R^3 in the compounds of Formula V and VI is selected from C_{1-6} alkyl, C_{2-6} alkenyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkylene C_{6-10} aryl, and C_{1-6} alkylene C_{3-6} cycloalkyl. In a further embodiment of the application, R^3 is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

[0067] In an embodiment of the application, LG in the compounds of Formula VI is any suitable leaving group, for example, halo, Ms, Ts, Ns, Tf, C_{1-6} acyl, and the like. In a specific embodiment LG is halo, such as Br.

[0068] In another embodiment, X is the anion of LG, for example Br^- . In a further embodiment, X is LG^- and the process further comprises a hydrolysis step to convert LG^- to OH^- . Hydrolysis can be performed, for example, by treating the compound of Formula V with a base in an aqueous, alcoholic solvent system.

[0069] In an embodiment, the compound of Formula V is selected from a compound of Formula V(a), V(b) and V(c):



wherein

---- represents a single or double bond;


R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, and PG is a protecting group;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label.

[0070] The compounds of Formula V, wherein R^1 and/or R^2 are PG, may be deprotected to provide the corresponding free OH compounds, that is compounds of Formula VII:



wherein


---- represents a single or double bond, provided that two double bonds are not adjacent to each other and when $\equiv O$ represents $=O$, then H is not present;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion; and

one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label.

[0071] The oxazolidine ring in the compounds of Formula V can be cleaved using either reducing or hydrolysis (acidic or basic) conditions. Under reducing conditions, the compounds of the Formula V provide compounds of the Formula VIII:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;


R^4 and R^5 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^5 is not present, or

R^4 and R^5 are H if, in the compounds of Formula V, R^1 and R^2 are a PG that is removed under the reducing conditions;

PG is a protecting group that is not removed under the reducing conditions, and

one or more available hydrogens in R^3 , R^4 and R^5 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^4 and R^5 is/are optionally replaced with an isotopic label.

[0072] Under some reducing conditions, the compounds of the Formula V can provide compounds of the Formula VIII(d):



wherein

— represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

R^4 and R^5 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^5 is not present, or

R^4 and R^5 are H if, in the compounds of Formula V, R^1 and R^2 are a PG that is removed under the reducing conditions;


PG is a protecting group that is not removed under the reducing conditions, and

one or more available hydrogens in R^3 , R^4 and R^5 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^4 and R^5 is/are optionally replaced with an isotopic label.

[0073] In an embodiment of the application, R^4 and R^5 in the compounds of Formula VIII(d) are independently selected from C_{1-6} alkyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{1-6} alkylene C_{6-10} aryl, C_{1-6} alkylene C_{3-6} cycloalkyl and PG. In a further embodiment of the application, R^4 and R^5 are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG.

[0074] Under hydrolysis conditions, the compounds of the Formula V provide compounds of the Formula IX:

25



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl;

R⁶ and R⁷ are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when ---O represents =O, then R⁷ is not present, or

R⁶ and R⁷ are H if, in the compounds of Formula V, R¹ and R² are a PG that is removed under the hydrolysis conditions;

PG is a protecting group that is not removed under the hydrolysis conditions, and

one or more available hydrogens in R³, R⁶ and R⁷ is/are optionally replaced with F and/or one or more of available atoms in R³, R⁶ and R⁷ is/are optionally replaced with an isotopic label.

[0075] In an embodiment of the application, R⁶ and R⁷ in the compounds of Formula IX are independently selected from C₁₋₆alkyl, phenyl, naphthyl, indanyl, C₃₋₆cycloalkyl, C₁₋₆alkyleneC₆₋₁₀aryl, C₁₋₆alkyleneC₃₋₆cycloalkyl and PG. In a further embodiment of the application, R⁶ and R⁷ are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG.

[0076] In an embodiment, the reducing conditions to provide the compounds of Formula VIII comprise treating the compounds of Formula V with a suitable reducing agent, such as metal hydride reducing agents, optionally in the presence of a Lewis acid, for a time and temperature for the conversion of the compound of Formula V to the compound of Formula VIII to

proceed to a sufficient extent, for example at about -100 °C to about 100 °C for about 0.5 hours to about 48 hours.

[0077] In a further embodiment, the hydrolysis conditions to provide the compounds of Formula IX comprise treating the compounds of Formula V under suitable acidic (for example acetic acid/ammonia buffer) or basic (for example ammonium bicarbonate/ammonia) conditions for a time and temperature for the conversion of the compound of Formula V to the compound of Formula IX to proceed to a sufficient extent, for example at about -100 °C to about 100 °C for about 0.5 hours to about 48 hours.


[0078] In a particular embodiment, PG is a protecting group that is removed under conditions to hydrolyze the compound of Formula V to the compound of Formula IX. For example, when PG is an alkyl carbonate, hydrolysis under basic conditions hydrolyzes the oxazolidine and removes the protecting group simultaneously. In another embodiment, when PG is an alkyl acetate, hydrolysis under acidic conditions hydrolyzes the oxazolidine and removes the protecting group simultaneously. A person skilled in the art would appreciate that other protecting groups removable under reducing, acidic or basic conditions compatible with the compounds of Formula V, VIII and IX can also be used.

[0079] In an alternate embodiment, R¹ and R² in the compounds of Formula V are not a PG that is removed under the reducing or hydrolysis conditions, and the compounds of Formula VIII and IX are further treated under conditions to remove the PG group to provide the corresponding free hydroxy compounds (i.e. compounds of Formula VIII wherein R⁴ and R⁵ are H and compounds of Formula IX wherein R⁶ and R⁷ are H).

(ii) Hydrolysis of the Compounds of Formula I

[0080] Hydrolysis of the compounds of Formula I under either acidic or basic conditions provides the free phenols of Formula X:

27



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^8 and R^9 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^9 is not present; or

R^8 and R^9 are H, if in the compound of Formula I, R^1 and R^2 are a PG that is removed under the hydrolysis conditions; and

PG is a protecting group,


wherein in the compounds of Formula X, one or more available hydrogens in R^8 and R^9 is/are optionally replaced with F and/or one or more of available atoms in R^8 and R^9 is/are optionally replaced with an isotopic label.

[0081] In an embodiment of the application, R^8 and R^9 in the compounds of Formula X are independently selected from C_{1-6} alkyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{1-6} alkylene C_{6-10} aryl, C_{1-6} alkylene C_{3-6} cycloalkyl and PG. In a further embodiment of the application, R^8 and R^9 are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG.

[0082] In a particular embodiment, PG in the compounds of Formula I is a protecting group that is removed under conditions to hydrolyze the compound of Formula I to the compound of Formula X. For example, when PG is an alkyl carbonate, hydrolysis under basic conditions hydrolyzes the oxazolidine and removes the protecting group simultaneously. In another embodiment, when PG is an alkyl acetate, hydrolysis under acidic conditions hydrolyzes the oxazolidine and removes the protecting group simultaneously. A person skilled in the art would appreciate that other protecting groups

removable under acidic or basic conditions compatible with the compounds of Formula I and X can also be used. In an alternate embodiment, PG is a protecting group that is not removed under conditions to hydrolyze the compound of Formula I to the compound of Formula X and is optionally removed in a separate step after the preparation of the compound of Formula X.

[0083] The compounds of Formula X are selectively alkylated at N-17 by reaction with a compound of the Formula R^{10} -LG¹ (XI), wherein LG¹ is a leaving group and R^{10} is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl under standard alkylation conditions to provide compounds of Formula XII:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R⁸ and R⁹ are independently selected from H, C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when ---O represents =O, then R⁹ is not present;

PG is a protecting group;

R¹⁰ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl; and


one or more available hydrogens in R⁸, R⁹ and R¹⁰ is/are optionally replaced with F and/or one or more of available atoms in R⁸, R⁹ and R¹⁰ is/are optionally replaced with an isotopic label.

[0084] When R⁸ and/or R⁹ in the compounds of Formula XII is PG, it is an embodiment of the present application that the compounds of Formula XII

are further treated under conditions to remove the PG to provide the corresponding free hydroxy compounds (i.e. compounds of Formula XII wherein R⁸ and/or R⁹ are H).

[0085] In an embodiment, R¹⁰ in the compounds of Formula XII is selected from C₁₋₆alkyl, C₂₋₆alkenyl, phenyl, naphthyl, indanyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, C₁₋₆alkyleneC₆₋₁₀aryl, and C₁₋₆alkyleneC₃₋₆cycloalkyl. In a further embodiment of the application, R¹⁰ is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

[0086] In this embodiment, it is possible to prepare the known morphine analogs, naltrexone (R¹⁰ is cyclopropylmethyl), nalbuphine (R¹⁰ is cyclobutylmethyl) and naloxone (R¹⁰ is allyl). In each of these latter compounds, R¹ is H, R² is not present and ring C (i.e. the bottom ring) has the structure:




[0087] In a specific example of the present application, oxymorphone was converted to naltrexone or naloxone in just three operations in an overall yield of 55-65% using a process of the present application.

[0088] The processes of the present application may be performed using continuous or batch processes. For commercial scale preparations continuous processes are suitable. Methods of performing chemical processes in continuous or batch modes are known in the art. When continuous processes are used, the reaction temperature and/or pressure may be higher than those used in batch processes.

III. Compounds of the Application

[0089] The present application includes compounds of Formula V:

30



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R¹ and R² are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when ---O represents =O, then R² is not present;

PG is a protecting group;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R¹, R² and R³ is/are optionally replaced with F and/or one or more of available atoms in R¹, R² and R³ is/are optionally replaced with an isotopic label,

or a salt or solvate thereof.


[0090] In an embodiment of the application, R¹ and R² in the compounds of Formula V are independently selected from C₁₋₆alkyl, phenyl, naphthyl, indanyl, C₃₋₆cycloalkyl, C₁₋₆alkyleneC₆₋₁₀aryl, C₁₋₆alkyleneC₃₋₆cycloalkyl and PG. In a further embodiment of the application, R¹ and R² are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG. It is an embodiment of the application that PG is an alkyl acetate, such as acetyl.

[0091] In another embodiment, R³ in the compounds of Formula V is selected from C₁₋₆alkyl, C₂₋₆alkenyl, phenyl, naphthyl, indanyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, C₁₋₆alkyleneC₆₋₁₀aryl, and C₁₋₆alkyleneC₃₋₆cycloalkyl. In a further embodiment of the application, R³ is selected from Me, Et, allyl, Ph,

cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

[0092] In another embodiment of the present application, X in the compounds of Formula V is OH⁻, Br⁻ or Cl⁻.

[0093] In an embodiment, the compound of Formula V is selected from a compound of Formula V(a), V(b) and V(c):



wherein

--- represents a single or double bond;


R¹ and R² are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, and PG is a protecting group;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R¹, R² and R³ is/are optionally replaced with F and/or one or more of available atoms in R¹, R² and R³ is/are optionally replaced with an isotopic label, or a salt or solvate thereof.

[0094] The present application also includes compounds of Formula VII:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other and when $\equiv O$ represents $=O$, then H is not present; R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion; and


one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label, or

a salt or solvate thereof.

[0095] In another embodiment, R^3 in the compounds of Formula VII is selected from C_{1-6} alkyl, C_{2-6} alkenyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkylene C_{6-10} aryl, and C_{1-6} alkylene C_{3-6} cycloalkyl. In a further embodiment of the application, R^3 is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

[0096] In another embodiment of the present application, X in the compounds of Formula VII is OH^- , Br^- or Cl^- .

[0097] In a further embodiment, the compounds of Formula VII are selected from a compound of Formula VII(a), VII(b) and VII(c):



wherein


--- represents a single or double bond;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion; and

one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label, or
a salt or solvate thereof.

[0098] The present application also includes compounds of Formula VIII:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

R^4 and R^5 are independently selected from H, C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^5 is not present;


PG is a protecting group; and

one or more available hydrogens in R^3 , R^4 and R^5 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^4 and R^5 is/are optionally replaced with an isotopic label, or
a salt or solvate thereof.

[0099] In an embodiment, R^3 in the compounds of Formula VIII is selected from C_{1-6} alkyl, C_{2-6} alkenyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkylene C_{6-10} aryl, and C_{1-6} alkylene C_{3-6} cycloalkyl. In a further embodiment of the application, R^3 is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

[00100] In an embodiment of the application, R^4 and R^5 in the compounds of Formula VIII are independently selected from H, C_{1-6} alkyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{1-6} alkylene C_{6-10} aryl, C_{1-6} alkylene C_{3-6} cycloalkyl and PG. In a further embodiment of the application, R^4 and R^5 are independently selected from H, Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG. It is an embodiment of the application that PG is an alkyl acetate, such as acetyl.

[00101] In a further embodiment, the compounds of Formula VIII are selected from a compound of Formula VIII(a), VIII(b) and VIII(c):



wherein

--- represents a single or double bond;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

R^4 and R^5 are independently selected from H, C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG;

PG is a protecting group; and

one or more available hydrogens in R^3 , R^4 and R^5 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^4 and R^5 is/are optionally replaced with an isotopic label, or

a salt or solvate thereof.


[00102] The following non-limiting examples are illustrative of the present application:

EXAMPLES

Example 1: General procedure for N-oxidations


[00103] To a solution of oxycodone, oxymorphone, or 3-O-Ac-oxymorphone (1g scale) in dichloromethane (10 mL) cooled to 4 °C was added *m*CPBA (1 eq., 77% purity). The reaction mixture was stirred for 10 min and then added dropwise to vigorously stirred diethyl ether (100 mL). A white precipitate of product was filtered off to give nearly quantitative yield.

(a) Oxycodone N-oxide




[00104] $R_F = 0.26$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -167.6$ (c 1, CHCl_3); mp = 220 °C (decomposition, Et_2O); IR (KBr, cm^{-1}) ν 3426, 3018, 2997, 2956, 2932, 2862, 2832, 2312, 2243, 2160, 2133, 1900, 1723, 1632, 1606, 1533, 1499, 1461, 1436, 1342, 1312, 1256, 1160, 1017, 976, 933, 816; ^1H NMR (CDCl_3 , 300 MHz) δ 6.76 (d, 1H, $J=8.4$ Hz), 6.67 (d, 1H, $J=7.8$ Hz), 4.78 (s, 1H), 3.92 (s, 3H), 3.62 (d, 1H, $J=5.1$ Hz), 3.34 (s, 3H), 3.31-3.10 (m, 6H), 2.24 (ddd, 1H, $J=3.0, 3.0, 14.4$ Hz), 1.97 (ddd, 1H, $J=3.0, 4.8, 12.6$ Hz), 1.70 (dd, 1H, $J=3.3, 12.5$ Hz), 1.61 (ddd, 1H, $J=3.3, 12.9, 14.4$ Hz); ^{13}C NMR (CDCl_3 , 75 MHz) δ 207.74, 145.32, 143.96, 129.28, 120.33, 120.08, 115.75, 89.93, 75.78, 72.23, 61.69, 59.54, 59.85, 49.94, 34.93, 32.89, 28.68, 25.80; MS (+EI) m/z (%) 332 (100), 314 (29); HRMS (+FAB) calcd for $\text{C}_{18}\text{H}_{22}\text{NO}_5$: 332.14980. Found 332.14636.

(b) 3-Acetyl-oxymorphone N-oxide



[00105] Oxymorphone (600 mg; 1.99 mmol) was dissolved in tetrahydrofuran (8 mL). Solid K_2CO_3 (275 mg; 1.99 mmol) and acetic anhydride (188 μ L, 1.99 mmol) were then added and the reaction mixture was stirred at room temperature for 1.5 hours. TLC (95/5 dichloromethane:methanol, double development) showed only traces of starting material. The crude material was subjected to the oxidation protocol without isolation to furnish the *N*-oxide as a solid: $[\alpha]_D^{20} = -189.33$ (c 1, $CHCl_3$); mp = 160 °C ($CHCl_3$); IR (KBr, cm^{-1}) ν 3449, 2968, 2938, 1764, 1726, 1685, 1654, 1627, 1495, 1444, 1373, 1287, 1219, 1193, 1161, 1111, 1044, 931, 629; 1H NMR ($CDCl_3$, 300 MHz) δ 12.25 (bs, 1H), 6.91 (d, 1H, $J=8.4$ Hz), 6.72 (d, 1H, $J=8.1$ Hz), 4.80 (s, 1H), 3.60 (d, 1H, $J=5.1$ Hz), 3.36-3.03 (m, 9H), 2.31 (s, 3H), 2.23 (ddd, 1H, $J=3.0, 3.0, 14.7$ Hz), 1.96 (ddd, 1H, $J=3.0, 4.8, 12.6$ Hz), 1.71 (dd, 1H, $J=3.9, 11.4$ Hz), 1.58 (ddd, 1H, $J=3.6, 14.1, 14.1$ Hz); ^{13}C NMR ($CDCl_3$, 75 MHz) δ 206.97, 168.31, 148.23, 133.53, 129.93, 125.90, 124.10, 119.84, 90.15, 75.44, 72.14, 61.51, 59.49, 49.77, 34.84, 32.62, 28.94, 25.82, 20.74; MS (FAB+) m/z (%) 360 (100), 342 (20); HRMS (+FAB) calcd for $C_{19}H_{22}NO_6$: 360.14471. Found 360.14441.


(c) 3-Ethoxycarbonyl-oxymorphone *N*-oxide



[00106] Oxymorphone (100 mg, 0.33 mmol) was suspended in ethyl acetate (1 mL) and ethylchloroformate (32 μ L, 3.3 mmol) was added dropwise prior to addition of triethylamine (46 μ L, 3.3 mmol). The reaction mixture (white suspension) was stirred for 1 hour at room temperature. TLC analysis (dichloromethane/methanol/ammonia (90:8:2)) showed essentially clean conversion to the product (R_f = 0.70), which was immediately subjected without isolation to the oxidation protocol to yield the titled compound as a solid: R_F = 0.28 (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -50 \rightarrow -120$ (c 1, CH_2Cl_2) dynamic rotation; mp = 112-115 °C (*i*-PrOH);

IR (KBr, cm^{-1}) ν 3432, 3062, 2980, 2935, 2361, 2343, 1768, 1728, 1627, 1497, 1445, 1371, 1261, 1195, 1164, 1065, 1002, 931, 864, 738; ^1H NMR (CDCl_3 , 600 MHz) δ 7.01 (d, 1H, $J=8.4$ Hz), 6.74 (d, 1H, $J=7.8$ Hz), 4.86 (s, 1H), 4.32 (m, 2H), 3.69 (d, 1H, $J=6.0$ Hz), 3.38-3.35 (m, 5H), 3.27 (ddd, 1H, $J=4.2, 13.2, 13.2$ Hz), 3.22 (dd, 1H, $J=5.4, 19.86$ Hz), 3.16 (ddd, 1H, $J=4.8, 14.4, 14.4$ Hz), 3.13 (ddd, 1H, $J=4.2, 12.0, 12.0$ Hz), 2.26 (ddd, 1H, $J=3.0, 3.0, 14.4$ Hz), 2.02 (ddd, 1H, $J=3.0, 4.8, 13.2$ Hz), 1.74 (dd, 1H, $J=1.8, 13.2$ Hz), 1.61 (ddd, 1H, $J=3.0, 14.4, 14.4$ Hz), 1.39 (t, 3H, $J=7.2$ Hz); ^{13}C NMR (CDCl_3 , 125 MHz) δ 206.65, 152.66, 148.14, 134.09, 130.15, 126.16, 123.83, 119.90, 90.26, 75.47, 72.22, 65.36, 61.54, 59.44, 49.75, 34.85, 32.56, 28.98, 25.86, 14.14; MS (+EI) m/z (%) 390 (100); HRMS (+EI) calcd for $\text{C}_{20}\text{H}_{24}\text{NO}_7$: 390.15528. Found 390.15495.


Example 2: (5a*R*,8a*S*,11a*R*,11b*S*)-2-methoxy-5,5a,9,10-tetrahydro-6,11b-ethano-7*H*-furo[2',3',4',5':4,5]phenanthro[9,8a-*d*]oxazol-11(11a*H*)-one



[00107] Oxycodone *N*-oxide (Example 1a, 150 mg; 0.45 mmol) was dissolved in dichloromethane (10 mL). The reaction mixture was cooled to -20 to -25 °C in an acetone/ $\text{N}_2(l)$ bath and Burgess reagent (150 mg; 0.63 mmol) was added as a solid in one portion. The reaction mixture was stirred for 5 hours and allowed to warm to room temperature. At -5 °C the color of the reaction mixture changed from colorless to yellowish. The mixture was then diluted with dichloromethane (50 mL) and washed with NaHCO_3 (2 x 10 mL). The aqueous layer was re-extracted with dichloromethane (15 mL) and the combined organic layers were dried with MgSO_4 and concentrated to yield 157 mg of the titled oxazolidine as a yellow solid. The compound was not stable on silica, low-melting, hygroscopic solid; data was collected at 90% purity.

[00108] $R_F = 0.7$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -113.3$ (c 1, CHCl_3); IR (KBr, cm^{-1}) ν 2926, 2854, 1728, 1635, 1610, 1506, 1441, 1385, 1335, 1313, 1277, 1257, 1165, 1088, 1074, 1003, 951, 926, 895, 779; ^1H NMR (CDCl_3 , 600 MHz) δ 6.78 (d, 1H, $J=8.4$ Hz), 6.72 (d, 1H, $J=8.1$ Hz), 4.75 (d, 1H, $J=6.0$ Hz), 4.71 (d, 1H, $J=6.0$ Hz), 4.69 (s, 1H), 3.91 (s, 3H), 3.35 (d, 1H, $J=18.6$ Hz), 3.27 (d, 1H, $J=7.8$ Hz), 3.16 (dd, 1H, $J=7.8, 18.6$ Hz), 2.92 (ddd, 1H, $J=4.8, 14.4, 14.4$ Hz), 2.86-2.80 (m, 2H), 2.43 (ddd, 1H, $J=3.0, 3.0, 10.5$ Hz), 2.39 (m, 1H), 2.00 (ddd, 1H, $J=3.9, 3.9, 13.8$ Hz), 1.68 (ddd, 1H, $J=3.0, 3.0, 14.7$ Hz), 1.56 (m, 1H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 207.15, 144.74, 142.86, 129.10, 123.20, 120.03, 115.19, 91.01, 86.49, 77.21, 64.10, 56.78, 52.58, 44.35, 37.14, 34.12, 30.56, 26.80; MS (+EI) m/z (%) 313 (100), 257 (8), 229 (8); HRMS (+EI) calcd for $\text{C}_{18}\text{H}_{19}\text{NO}_4$: 313.13141. Found 313.13128.

Example 3: (5aR,8aS,11aR,11bS)-2-Acetoxy-5,5a,9,10-tetrahydro-6,11b-ethano-7H-furo[2',3',4',5':4,5]phenanthro[9,8a-d]oxazol-11(11aH)-one, one-pot protocol




[00109] Oxymorphone (600 mg; 1.99 mmol) was dissolved in tetrahydrofuran (8 mL). Solid K_2CO_3 (275 mg; 1.99 mmol) and acetic anhydride (188 μL , 1.99 mmol) were then added and the reaction mixture was stirred at room temperature for 1.5 hours. TLC (95/5 dichloromethane:methanol, double development) showed only traces of starting material. The reaction mixture was then cooled to $\sim 4^\circ\text{C}$ in an ice-bath, and a cold (4°C) solution of *m*CPBA (446 mg; 1.99 mmol; 77% purity) in dichloromethane (6 mL) was added dropwise over a period of 1 minute. [The solution of *m*CPBA was prepared by dissolving 669 mg *m*CPBA (77%) in dichloromethane (9 mL) and adding MgSO_4 (670 mg). The mixture was

agitated several times over a period of 30 min and cooled in an ice-bath to 4 °C]. After 1 hour stirring, a white precipitate of *N*-oxide formed and the reaction mixture was cooled to -20 °C. Burgess reagent (593 mg, 2.49 mmol) in dichloromethane (7 mL) was then cannulated at -20 °C into the reaction mixture over a period of 2 minutes. The reaction mixture was allowed to warm to 10 °C (3 h total reaction time) and then diluted with ethyl acetate (100 mL), and washed with NaHCO₃ solution (2 x 20 mL). The combined aqueous layer was reextracted with ethylacetate (2 x 20 mL) and the combined organic layer was dried with MgSO₄, filtered and concentrated to yield 636 mg (93%) of reasonably pure (92-95%) material. Crystallization of the product from a mixture of EtOH/ *i*-PrOH 1:1 (2 mL), temperature regime of 25 °C to 5-10 °C, afforded 520 mg (76%) of product. Repetition of the experiment on a scale of one gram yielded 78% of the titled compound as a solid.

[00110] $R_F = 0.7$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -84.1$ (c 1, CHCl₃); mp = 179-182 °C (*i*-PrOH); IR (KBr, cm⁻¹) ν 2954, 2892, 2864, 2834, 1765, 1722, 1624, 1494, 1445, 1370, 1339, 1317, 1216, 1201, 1185, 1158, 1073, 1009, 958, 930, 889, 781; ¹H NMR (CDCl₃, 600 MHz) δ 6.90 (d, 1H, *J*=8.2 Hz), 6.76 (d, 1H, *J*=8.4 Hz), 4.73 (d, 1H, *J*=6.0 Hz), 4.70 (d, 1H, *J*=6.0 Hz), 4.69 (s, 1H), 3.37 (d, 1H, *J*=19.2 Hz), 3.27 (d, 1H, *J*=7.8 Hz), 3.17 (dd, 1H, *J*=7.8, 19.2 Hz), 2.89 (ddd, 1H, *J*=4.8, 14.4, 14.4 Hz), 2.80 (m, 2H), 2.45-2.30 (m, 5H), 1.98 (ddd, 1H, *J*=3.3, 4.2, 13.8 Hz), 1.67 (ddd, 1H, *J*=3.1, 14.4, 14.4 Hz), 1.56 (m, 1H); ¹³C NMR (CDCl₃, 150 MHz) δ 206.36, 168.57, 147.49, 132.45, 129.76, 128.65, 123.49, 119.97, 91.25, 86.53, 77.00, 63.95, 52.40, 44.23, 37.09, 34.09, 30.28, 27.13, 20.83; MS (+EI) *m/z* (%) 341 (8), 299 (100), 243 (7); HRMS (+EI) calcd for C₁₉H₁₉NO₅: 341.12632. Found 341.12606.

Example 4: (5aR,8aS,11aR,11bS)-2-[(Ethoxycarbonyl)oxy]-5,5a,9,10-tetrahydro-6,11b-ethano-7H-furo[2',3',4',5':4,5]phenanthro[9,8a-d]oxazol-11(11aH)-one, one-pot protocol



[00111] Oxymorphone (100 mg, 0.33 mmol) was suspended in ethyl acetate (1 mL) and ethylchloroformate (32 μ L, 3.3 mmol) was added dropwise prior to addition of triethylamine (46 μ L, 3.3 mmol). The reaction mixture (white suspension) was stirred for one hour at room temperature. TLC analysis (dichloromethane/methanol/ammonia (90:8:2)) showed essentially a clean conversion to the ethyl carbonate-protected oxymorphone.

[00112] The crude reaction mixture containing the carbonate was then cooled to 4 °C in an ice-bath, and a 1 mL aliquot of a solution of *m*CPBA [The solution was prepared as follows: *m*CPBA (148 mg, 77% peroxide content, 0.66 mmol) was dissolved in ethyl acetate (2 mL) and MgSO_4 (140 mg) was added. The solution was dried 30 min and then cooled to 4 °C] was added dropwise. The reaction mixture was stirred for one hour at 4 °C. TLC analysis (dichloromethane/methanol/ammonia (90:8:2)) showed essentially clean conversion to the carbonate-protected product R_f =0.28. The reaction mixture was then cooled to -25 °C and Burgess reagent was added in one portion as a solid. The mixture was then allowed to reach room temperature in approximately 2-3 hours as the color changed from a white to a yellow suspension. The reaction mixture was then diluted with ethyl acetate (10 mL), washed with NaHCO_3 (2 x 4 mL), and the aqueous layer was re-extracted with ethyl acetate (5 mL). The combined organic layers were dried over MgSO_4 and concentrated to yield 125 mg of crude oxazolidine (85-90% purity) in ~84% yield as a low melting solid:

[00113] R_F = 0.70 (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20}$ = -64.19 (*c* 1, CHCl_3); mp = low-melting solid; IR (KBr, cm^{-1}) ν 3448, 2960, 2945, 2924, 2887, 1764, 1728, 1626, 1498, 1448, 1372, 1341, 1257, 1237, 1208, 1164, 1073, 1027, 931, 783; ^1H NMR (CDCl_3 , 600 MHz) δ 6.99 (d, 1H, $J=8.4$ Hz), 6.78 (d, 1H, $J=7.8$ Hz), 4.76 (d, 1H, $J=6.0$ Hz), 4.74 (s, 1H), 4.72 (d, 1H, $J=5.4$ Hz), 4.35-4.31 (m, 2H), 3.38 (d, 1H, $J=18.6$ Hz), 3.29 (d, 1H, $J=8.4$ Hz), 3.19 (dd, 1H, $J=7.8$, 19.2 Hz), 2.90 (ddd, 1H, $J=4.8$, 14.4, 14.4 Hz), 2.82 (bd, 2H, $J=8.4$ Hz), 2.42 (bd, 1H, $J=13.2$ Hz), 2.39 (m, 1H), 1.99 (ddd, 1H, $J=\sim 1.0$, ~ 1.0 , 13.8 Hz), 1.69 (ddd, 1H, $J=\sim 1.0$, 13.2, 13.2 Hz), 1.58 (d, 1H, $J=12.6$ Hz), 1.39 (t, 3H, $J=7.2$ Hz); ^{13}C NMR (CDCl_3 , 150 MHz) δ 206.19, 152.91, 147.38, 132.97, 130.00, 128.95, 123.15, 120.02, 91.35, 86.50, 65.15, 63.91, 52.37, 44.20, 37.07, 34.11, 30.23, 27.13, 14.16, 1.03; MS (+EI) m/z (%) 371 (13), 327 (14), 299 (100); HRMS (+EI) calcd for $\text{C}_{20}\text{H}_{21}\text{NO}_6$: 371.13689. Found 371.13697.


[00114] Analytical samples of intermediates (protected oxymorphone and protected oxymorphone *N*-oxide) were prepared in a stepwise manner and purified by column chromatography. An analytical sample of the oxazolidine from Example 4 was prepared from the protected oxymorphone. The *N*-oxidation and treatment with the Burgess reagent was performed in two steps in 95% yield (95% purity). Crystallization of the product from Example 4 was not possible because it is a low-melting hygroscopic solid).

Oxymorphone O-ethyl carbonate:

[00115] R_F = 0.75 (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20}$ = -146.84 (*c* 1, CH_2Cl_2); mp = 156-157 °C (*i*-PrOH); IR (KBr, cm^{-1}) ν 3433, 2982, 2936, 2907, 2870, 2811, 1755, 1727, 1627, 1498, 1447, 1373, 1348, 1321, 1197, 1165, 1032, 934, 782; ^1H NMR (CDCl_3 , 600 MHz) δ 6.92 (d, 1H, $J=8.4$ Hz), 6.71 (d, 1H, $J=7.8$ Hz), 5.09 (bs, 1H), 4.72 (s, 1H), 4.32 (m, 2H), 3.20 (d, 1H, $J=18.6$ Hz), 3.02 (ddd, 1H, $J=4.8$, 14.4, 14.4 Hz), 2.90 (d, 1H, $J=6.0$ Hz), 2.60 (dd, 1H, $J=6.0$, 18.6 Hz), 2.49 (dd, 1H, $J=4.8$, 12.0 Hz), 2.47 (m, 4H), 2.32 (ddd, 1H, $J=3.0$, 3.0, 14.4 Hz), 2.16 (ddd, 1H, $J=4.2$, 12.6, 12.6 Hz), 1.89 (ddd, 1H, $J=3.0$, 4.8, 13.2 Hz), 1.63 (ddd, 1H, $J=3.0$, 13.8, 13.8 Hz),

1.59 (dd, 1H, $J=3.6, 13.2$ Hz), 1.38 (t, 3H, $J=7.2$ Hz); ^{13}C NMR (CDCl_3 , 150 MHz) δ 207.51, 152.89, 147.59, 133.08, 130.59, 130.22, 122.59, 119.43, 90.69, 70.20, 65.09, 64.40, 50.05, 45.06, 42.71, 36.02, 31.11, 30.56, 22.27, 14.15; MS (+EI) m/z (%) 373 (100), 329 (21), 301 (99), 244 (34), 216 (38); HRMS (+EI) calcd for $\text{C}_{20}\text{H}_{23}\text{NO}_6$: 373.15254. Found 373.15284.


Example 5: Noroxymorphone



[00116] A. Acetic acid buffer: The oxazolidine from Example 3 (0.1 g, 0.29 mmol) was suspended in AcOH/NH_3 buffer (pH 9, 10% w/w, 1.5 mL) and heated for 16 hours at 50 °C. The reaction mixture was then cooled to room temperature and stirred for an additional two hours. A light-brown precipitate of product was filtered off and dried to yield 69 mg (82%) of noroxymorphone as a brownish solid. m.p. >300 °C (lit >300°C).^{xvi}

[00117] B. Ammonium carbonate buffer: The oxazolidine from Example 3 (0.2 g, 0.57 mmol) was suspended in $\text{NH}_4\text{HCO}_3/\text{NH}_3$ buffer (pH 9, 10% w/w, 1 mL) and heated for 16 hours at 50 °C. The reaction mixture was then cooled to room temperature and stirred for an additional two hours. The light brown precipitate of product was filtered off and dried to yield 131 mg (78%) of noroxymorphone as a brownish solid. m.p. >300 °C.

Example 6: Naltrexone




[00118] Cyclopropylmethyl bromide (64 mg; 0.479 mmol) and Et_3N (45 μL ; 0.327 mmol) were added to a suspension of noroxymorphone (Example 5,

100 mg; 0.348 mmol) in a mixture of N-Methyl-2-pyrrolidone (NMP)/H₂O (10:1; 0.35 mL). The reaction vessel was purged with argon and the reaction mixture was stirred at 70 °C for 2h. At that time, additional Et₃N (45 µl; 0.327 mmol) was added and the mixture was stirred for an additional 6 h at 70 °C. The reaction mixture was then cooled to room temperature, diluted with dichloromethane (15 mL) and washed with saturated NaHCO₃ (3 x 3 mL). The aqueous layer was re-extracted with dichloromethane (5 mL) and the combined organic layers were dried over MgSO₄. Column chromatography of the residue (dichloromethane/methanol 4:1) afforded 103 mg (87%) of naltrexone as a white solid: mp 173-175 °C (acetone), mp 159-161 °C (MeOH), [lit. mp 174-176 °C (acetone)]^{xvii} identical in all respects to the material described in the literature.^{xviii}


[00119] R_f 0.42 (ethyl acetate + 20% MeOH); [α]²⁰_D = -207.00 (c=1, CHCl₃); IR (CHCl₃) ν 3568, 3359, 3010, 2931, 2834, 1723, 1620, 1317, 1146, 1058, 943; ¹H NMR (600 MHz, CDCl₃) δ 6.74 (d, J=8.1 Hz, 1H), 6.60 (d, J=8.1 Hz, 1H), 5.82 (bs, 1H, OH), 4.74 (s, 1H), 3.21 (d, J=5.9 Hz, 1H), 3.11-3.03 (m, 2H), 2.72 (dd, J=12.0, 4.8 Hz, 1H), 2.58 (dd, J=18.4, 6.0 Hz, 1H), 2.49-2.39 (m, 3H), 2.34 (ddd, J=14.5, 3.0, 3.0 Hz, 1H), 2.18 (ddd, J=12.2, 3.8, 3.8 Hz, 1H), 1.91 (m, 1H), 1.66 (ddd, J=14.2, 14.2, 3.3 Hz, 1H), 1.59 (ddd, J=12.8, 2.7 Hz, 1H), 0.88 (m, 1H), 0.57 (m, 2H), 0.16 (m, 2H); ¹³C NMR (150 MHz, CDCl₃) δ 210.02, 142.51, 138.80, 129.05, 124.25, 119.90, 117.91, 90.60, 70.32, 62.01, 59.21, 51.07, 43.60, 36.21, 31.36, 30.65, 22.62, 9.42, 4.02, 3.81; MS (+EI) m/z (%): 47 (15), 55 (41), 84 (100), 110 (12), 202 (5), 256 (12), 286 (7), 300 (15), 341 (64); HRMS calcd for C₂₀H₂₃NO₄ 341.1627, found 341.16320.

Example 7: Naloxone



[00120] Allyl bromide (56 mg; 0.463 mmol) and Et₃N (45 μ l; 0.327 mmol) were added to a suspension of noroxymorphone (Example 5, 100 mg; 0.348 mmol) in a mixture of NMP/H₂O (10:1; 0.35 mL). The reaction vessel was purged with argon and the mixture was stirred at 70 °C for 2h. At that time, additional Et₃N (45 μ l; 0.327 mmol) was added and the mixture was stirred for an additional 7.5 h at 70 °C. The reaction mixture was then cooled to room temperature, diluted with dichloromethane (15 mL), and washed with saturated NaHCO₃ (3 x 3 mL). The aqueous layer was re-extracted with dichloromethane (5 mL) and the combined organic layers were dried over MgSO₄. Column chromatography of the residue (dichloromethane/methanol 4:1) afforded 96 mg (84%) of naloxone as a white solid mp: 181-182 °C (ethyl acetate), [lit. mp 173-175]^{xxix} [lit. 179.5 °C (toluene)]^{xx} identical in all respects to the material described in the literature.^{xxi}


Example 8: Nalbuphene



[00121] A slurry of noroxymorphone (Example 5, 220 mg; 0.766 mmol), sodium hydrogen carbonate (77 mg; 0.92 mmol), cyclobutylmethyl bromide (160 mg; 1.07 mmol) and NMP (1 mL) was stirred under a nitrogen atmosphere at 90 °C for 19 h. Then the reaction mixture was cooled and quenched with water (10 mL). After adjusting the pH to 9, the product was extracted with DCM (3x5 mL). The combined organic layers were washed with water, brine and dried over MgSO₄. Column chromatography afforded 180 mg (66%) of nalbuphene as a white solid; mp 170-172 °C (acetone), [lit. 173-174 °C (chloroform)]^{xxii}; R_f 0.64 (ethyl acetate + 20% methanol); [α]_D²⁰ = -180.44 (c = 1.0, MeOH); IR (CHCl₃) ν 3561, 3454, 2966, 2931, 2830, 1720, 1616, 1457, 1318, 1142, 1057, 944 cm⁻¹; ¹H NMR (600 MHz, CDCl₃) δ 6.74 (d, J=8.0 Hz, 1H), 6.61 (d, J=8.0 Hz, 1H), 5.64 (bs, 1H, OH), 4.72 (s, 1H), 3.11 (d, J=18.4 Hz, 1H), 3.04 (ddd, J=14.4, 14.4, 3.6 Hz, 1H), 2.92 (d, J=4.9 Hz, 1H),

2.57 (m, 5H), 2.42 (ddd, $J=12.4, 12.4, 4.4$ Hz, 1H), 2.33 (d, $J=14.4$ Hz, 1H), 2.20 (ddd, $J=12.0, 12.0, 2.2$ Hz, 1H), 2.11 (m, 2H), 1.95 (m, 1H), 1.90 (m, 2H), 1.87 (m, 2H), 1.66 (ddd, $J=13.6, 13.6, 2.2$ Hz, 1H), 1.56 (d, $J=12.6$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ 209.68, 143.45, 138.69, 129.02, 124.34, 119.87, 117.71, 90.58, 70.31, 62.74, 60.48, 50.93, 43.74, 36.18, 33.73, 31.32, 30.69, 27.00, 26.79, 22.96, 18.76; MS (FAB+) m/z (%): 41 (27), 69 (9), 98 (5), 300 (88), 355 (38), 356 (100); HRMS calcd for $\text{C}_{21}\text{H}_{26}\text{NO}_4$ 356.1856, found 356.18552.

Example 9: (5aR,8aS,11aR,11bS)-6-Allyl-2-methoxy-11-oxo-5,5a,9,10,11-pentahydro-6,11b-ethano-7H-furo[2',3',4',5':4,5]phenanthro[9,8a-d]oxazol-6-ium bromide




[00122] The compound from Example 2 (20 mg; 0.064 mmol) was dissolved in nitromethane (0.5 mL) and allyl bromide (77 mg; 0.63 mmol) was added. The reaction mixture was heated to 85 °C, stirred for 16 hours and then cooled to room temperature. The precipitated solid was filtered and dried in vacuo to yield essentially pure quaternary salt (22 mg, 80 %).

[00123] $R_f = 0.10-0.15$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -108.4$ (c 1, MeOH); IR (KBr, cm^{-1}) ν 3416, 2960, 2933, 2839, 1729, 1638, 1615, 1508, 1446, 1331, 1319, 1278, 1194, 1165, 1112, 1087, 1061, 1007, 948, 916, 801; ^1H NMR (CDCl_3 , 300 MHz) δ 6.97 (d, 1H, $J=8.4$ Hz), 6.89 (d, 1H, $J=8.4$ Hz), 6.09 (m, 1H), 5.83 (d, 1H, $J=15.6$ Hz), 5.78 (d, 1H, $J=9.3$ Hz), 5.50 (d, 1H, $J=5.1$ Hz), 5.24 (dd, 1H, $J=2.1, 5.1$ Hz), 5.12 (s, 1H), 4.41 (dd, 1H, $J=7.5, 13.2$ Hz), 4.33 (d, 1H, $J=7.5$ Hz), 4.17 (dd, 1H, $J=7.2, 13.2$ Hz), 3.93 (s, 3H), 3.77 (m, 1H), 3.72 (m, 1H), 3.40 (dddd, 1H, $J=3.3, 3.3, 7.2, 21.3$ Hz), 3.24 (ddd, 1H, $J=2.1, 4.8, 15.0$ Hz), 2.99 (ddd, 1H, $J=4.2, 14.1, 14.1$ Hz), 2.77 (ddd, 1H, $J=5.7, 14.4, 14.4$ Hz), 2.39-2.30 (m, 2H), 1.95 (m,

1H), 1.78 (ddd, 1H, $J=3.9, 15.6, 15.6$ Hz); ^{13}C NMR (CDCl_3 , 150 MHz): Major rotamer δ 205.7, 144.8, 143.5, 128.5, 126.9, 124.0, 120.6, 120.3, 117.1, 89.4, 87.9, 83.7, 70.3, 60.6, 56.3, 51.7, 50.5, 34.4, 30.8, 29.4, 22.5; MS (FAB+) m/z (%) 354 (100); HRMS (FAB+) calcd for $\text{C}_{21}\text{H}_{24}\text{NO}_4$: 354.17053. Found 354.17047.

[00124] Note: carbon signals at 600 MHz/150 MHz NMR indicate rotamers.

Example 10: (5aR,8aS,11aR,11bS)-2-Acetoxy-6-allyl-11-oxo-5,5a,9,10,11-pentahydro-6,11b-ethano-7H-furo[2',3',4',5':4,5]phenanthro[9,8a-d]oxazol-6-ium bromide




[00125] The compound of Example 3 (25.5 mg, 0.075 mmol) was stirred with 3 eq allyl bromide (19.0 μL , 0.224 mmol) in 0.3 mL nitromethane. After stirring for two hours, the solvents were evaporated to yield 36 mg of the title compound in essentially quantitative yield.

[00126] $R_F = 0.2$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -88$ to -92 (c 1, MeOH); IR (KBr) ν 3448, 2931, 1761, 1731, 1554, 1448, 1198 cm^{-1} ; ^1H NMR (300 MHz, MeOD) δ 7.06 (d, 1H, $J = 8.4$ Hz), 6.98 (d, 1H, $J = 8.1$ Hz), 6.21-6.06 (m, 1H), 5.87 (d, 1H, $J = 16.5$ Hz), 5.78 (d, 1H, $J = 9.9$ Hz), 5.55 (d, 1H, $J = 5.4$ Hz), 5.31 (dd, 1H, $J = 2.1, 5.4$ Hz), 5.20 (s, 1H), 4.54-4.46 (m, 2H), 4.29 (dd, 1H, $J = 7.2, 13.2$ Hz), 3.90 (d, 1H, $J = 20.7$ Hz), 3.80 (dd, 1H, $J = 5.4, 13.2$ Hz), 3.55-3.42 (m, 1H), 3.27 (dddd, $J = 1.4, 4.4, 6.0, 12.7$ Hz), 2.98 (ddd, 1H, $J = 4.2, 14.1, 14.1$ Hz), 2.81 (ddd, 1H, $J = 5.7, 13.5, 13.8$ Hz), 2.39-2.30 (m, 2H), 2.32 (s, 3H), 2.00-1.91 (m, 1H), 1.79 (ddd, 1H, $J = 3.3, 14.4, 14.4$ Hz) ppm; ^{13}C NMR (75 MHz, MeOD) δ 205.4, 168.8, 147.7, 132.9, 128.7, 127.6, 126.4, 124.6, 124.1, 120.6, 89.9, 87.9, 83.5,

70.1, 60.6, 51.5, 50.5, 34.5, 30.8, 29.2, 23.0, 19.2 ppm; MS (FAB+) m/z %: 414 ($M+CH_3OH$) (100), 382 (M^+) (83), 352 (13), 310 (7), 185 (6), 77 (7), 43 (11). HRMS Calcd for $C_{22}H_{24}NO_5$: 382.16545 found: 382.16100.


Example 11: (5aR,8aS,11aR,11bS)-6-Allyl-2-hydroxy-11-oxo-5,5a,9,10,11-pentahydro-6,11b-ethano-7H-furo[2',3',4',5':4,5]phenanthro[9,8a-d]oxazol-6-ium hydroxide



[00127] The compound of Example 3 (60 mg, 0.176 mmol) was dissolved in nitromethane (0.6 mL) and allyl bromide (0.15 mL, 1.759 mmol) was added to the mixture. The solution was stirred at room temperature for two hours, when TLC (DCM/MeOH/ NH_4OH 90/9/1) indicated only the formation of the product. No precipitate was observed at that time and the mixture was allowed to stir overnight. TLC after 12 hours was identical with the one from the day before. Solvents were evaporated under a stream of argon, and an NMR of the crude material was obtained. NMR showed 10% of "solvated" product. After leaving the compound in CD_3OD for a few hours, the ratio of "naked" to "solvated" product changes (40%, see NMR at 600 MHz). The CD_3OD was evaporated, the mixture was stirred in a saturated solution of $NaHCO_3$ (0.3 mL) and after concentration to dryness the residue was purified by chromatography on silica gel (7 mL) in DCM/MeOH/ H_2O 5/1/0.06) to furnish 52 mg (83%) of the title compound (as a zwitterion).

[00128] $R_F = 0.1$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); $[\alpha]_D^{20} = -106.971$ (c 2, MeOH); IR (KBr) ν 3422, 3258, 2969, 1728, 1627, 1504, 1464, 1319, 1087, 920 cm^{-1} ; ^1H NMR (300 MHz, MeOD) δ 6.80 (s, 2H), 6.15-6.04 (m, 1H), 5.83 (d, 1H, $J = 17.1$ Hz), 5.78 (d, 1H, $J = 9.9$ Hz), 5.50 (d, 1H, $J = 5.4$ Hz), 5.25 (dd, 1H, $J = 2.4, 5.4$ Hz), 5.10 (s, 1H), 4.42 (dd, 1H, $J = 7.5, 13.2$ Hz), 4.34 (d, 1H, $J = 7.2$ Hz), 4.19 (dd, 1H, $J = 6.9, 13.2$ Hz), 3.78-6.68 (m, 2H), 3.36 (s, 1H), 3.34-3.22 (m, 1H), 3.00 (ddd 1H, $J = 4.8, 14.1, 14.1$ Hz), 2.77 (ddd, 1H, $J = 6.0, 13.5, 13.5$ Hz), 2.40-2.29 (m, 2H), 2.00-1.91 (m, 1H), 1.79 (ddd, 1H, $J = 3.3, 15.6, 15.6$ Hz) ppm; ^{13}C NMR (75 MHz, CD_3OD) δ 206.6, 143.5, 140.3, 128.6, 126.6, 124.1, 120.6, 119.2, 118.8, 89.3, 87.9, 83.8, 70.4, 60.6, 51.7, 50.7, 34.5, 30.8, 29.4, 22.6 ppm; MS (FAB+) m/z %: 340 (M^+) (13), 176 (16), 149 (27), 136 (21), 95 (24), 83 (32), 69 (70), 55 (68), 43 (100). HRMS Calcd for $\text{C}_{20}\text{H}_{22}\text{NO}_4$: 340.15488 found: 340.15459.

Example 12: 3-Acetoxy-17-(2-nitroethyl)-noroxymorphone




[00129] The compound of Example 3 (59 mg, 0.173 mmol) was dissolved in nitromethane (0.6 mL) and cyclopropylmethyl bromide (50.0 μL , 0.519 mmol) was added to the mixture. The solution was stirred at room temperature for 7 hours at which time TLC analysis (DCM/MeOH/ NH_4OH 90/9/1) indicated no progress and the mixture was heated at 50 $^\circ\text{C}$ overnight. After this time, TLC analysis showed traces of starting material and additional cyclopropylmethyl bromide (50.0 μL , 0.519 mmol, 3 equiv) was added. After 6 hours of stirring at room temperature, the solvents were evaporated under a stream of argon. Chromatography of the residue (6 mL of silica gel) in DCM/MeOH 100/1 gradient elution to 25/1 gave as product, 27 mg of the titled compound (38%) as a colorless glassy material, in addition to a inseparable

mixture of compounds (14 mg). After trituration with MeOH, the titled compound was obtained as a white crystalline solid.

[00130] $R_F = 0.9$ (dichloromethane/methanol/ammonium hydroxide 90:8:2); mp = 145-174 °C, becomes brown (MeOH); $[\alpha]_D^{20} = -148.46$ (c 1, CHCl₃); IR (KBr) ν 3427, 2931, 2837, 1767, 1728, 1554, 1443, 1370, 1214, 1187 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 6.87 (d, 1H, $J = 8.1$ Hz), 6.71 (d, 1H, $J = 8.1$ Hz), 4.69 (s, 1H), 4.62-4.47 (m, 2H), 3.27 (ddd, 1H, $J = 4.8, 8.1, 14.4$ Hz), 3.11 (d, 1H, $J = 18.9$ Hz), 3.07 (dd, 1H, $J = 4.2, 10.2$ Hz), 3.01 (d, 1H, $J = 6.3$ Hz), 2.99 (dd, 1H, $J = 5.1, 12.9$ Hz), 2.77 (dd, 1H, $J = 5.7, 18.6$ Hz), 2.66-2.59 (m, 1H), 2.42-2.26 (m, 2H), 2.33 (s, 3H), 1.90 (ddd, 1H, $J = 3.0, 5.1, 13.2$ Hz), 1.67-1.55 (m, 2H) ppm; ¹³C NMR (75 MHz, CDCl₃) δ 207.4, 168.5, 147.8, 132.8, 129.8, 129.6, 123.2, 119.4, 90.4, 73.7, 70.1, 70.0, 63.9, 51.9, 50.2, 43.4, 35.9, 31.0, 30.3, 24.7, 20.8 ppm; MS (FAB+) m/z %: 403 (M+H⁺) (100), 402 (M⁺) (21), 385 (14), 360 (45), 342 (17), 300 (6), 214 (15), 187 (7), 129 (7), 84 (7), 56 (14), 43 (19). HRMS Calc'd for C₂₀H₂₃N₂O₇: 403.15053 found: 403.15129.

Example 13:



[00131] In preliminary experiments, the reduction of the above quaternary salt was attempted under a variety of conditions, including the activation of the C-14 oxygen with Lewis acids. The C-14 methyl ether was obtained as shown in the above scheme. Hydrolysis of the compound under acid buffer or basic conditions provided naltrexone.

[00132] While the present application has been described with reference to what are presently considered to be the preferred examples, it is to be understood that the application is not limited to the disclosed examples. To

the contrary, the application is intended to cover various modifications and equivalent arrangements included within the spirit and scope of the appended claims.

[00133] All publications, patents and patent applications are herein incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety. Where a term in the present application is found to be defined differently in a document incorporated herein by reference, the definition provided herein is to serve as the definition for the term.


**FULL CITATIONS FOR DOCUMENTS REFERRED TO IN THE
SPECIFICATION**

- ⁱ von Braun, J.; *Chem. Ber.*, **1900**, 33, 1438.
- ⁱⁱ Cooley, J.H.; Evain, E.J.; *Synthesis* **1989**, 1.
- ⁱⁱⁱ Olofson, R. A.; Martz, J. T.; Senet, J.-P.; Piteau, M.; Malfroot, T. *J. Org. Chem.* **1984**, 49, 2081.
- ^{iv} (a) K. M. Madyastha, *Proc. Indian Acad. Sci.* **1994**, 106, 1203; (b) K. M. Madyastha, G. V. B. Reddy, *J. Chem. Soc. Perkin Trans. 1* **1994**, 911.
- ^v Chaudhary, V.; Leisch, H.; Moudra, A.; Allen, B.; De Luca, V.; Cox, D. P.; Hudlicky, T. *Collect. Czech. Chem. Commun.* **2009**, 74, 1179-1193.
- ^{vi} (a) G. Kok, T. D. Asten, P. J. Scammells, *Adv. Synth. Catal.* **2009**, 351, 283; (b) Z. Dong, P. J. Scammells, *J. Org. Chem.* **2007**, 72, 9881.
- ^{vii} Smith, C.; Purcell, S.; Waddell, L.; Hayes, N.; Ritchie, J.; WO 2005/028483.
- ^{viii} Rinner, U.; Adams, D. R.; dos Santos, M. L.; Hudlicky, T. *Synlett* **2003**, 1247-1252.
- ^{ix} (a) Nicolaou, K. C.; Snyder, S. A.; Nalbandian, A. Z.; Longbottom, D. A. *J. Am. Chem. Soc.* **2004**, 126, 6234; (b) Nicolaou, K. C.; Huang, X.; Snyder, S. A.; Rao, P. B.; Bella, M.; Reddy, M. V. *Angew. Chem. Int. Ed. Engl.* **2002**, 41, 834; (c) Nicolaou, K. C.; Longbottom, D. A.; Snyder, S. A.; Nalbandian, A. Z.; Huang, X. *Angew. Chem. Int. Ed. Engl.* **2002**, 41, 3866.
- ^x Banfield, S. C.; Omori, A. T.; Leisch, H.; Hudlicky, T. *J. Org. Chem.* **2007**, 72, 4989-4992.
- ^{xi} (a) For a recent summary of new applications of the Burgess reagent see: Leisch, H.; Sullivan, B.; Fonovic, B.; Dudding, T.; Hudlicky, T. *Eur. J. Org. Chem.* **2009**, 2806-2819; Reviews of applications of the Burgess reagent: (b) Santra, S. *Synlett* **2009**, 852; (c) Nicolaou, K. C.; Snyder, S. A.; Longbottom, D. A.; Nalbandian, A. Z.; Huang, X. *Chem. Eur. J.* **2004**, 10, 5581; (d) Khapli, S.; Dey, S.; Mal, D. *J. Ind. Inst. Sci.* **2001**, 81, 461; (e) Burckhardt, S. *Synlett* **2000**, 559; (f) Lamberth, C. *J. Prakt. Chem.* **2000**, 342, 518; (g) Taibe, P.;

-
- Mobashery, S. in *Encyclopedia of Reagents in Organic Synthesis*, vol. 5, Paquette, L. A., Ed., Wiley, Chichester, 1995, p. 3345.
- ^{xii} Metcalf, T. A.; Simionescu, R.; Hudlicky, T. *J. Org. Chem.* **2010**, 75, 3447-3450.
- ^{xiii} Leisch, H.; Saxon, R.; Sullivan, B.; Hudlicky, T. *Synlett* **2006**, 445–449.
- ^{xiv} Use of the Burgess reagent in natural product synthesis: a) **cedrene**: Rigby, J. H.; Kirova-Snover, M. *Tetrahedron Lett.* **1997**, 38, 8153; b) **narciclasine**: Rigby, J. H.; Mateo, M. E. *J. Am. Chem. Soc.* **1997**, 119, 12655; c) **taxol**: Holton, R. A.; Kim, H. B.; Sonoza, C.; Liang, F.; Biediger, R. J.; Boatman, P. D.; Shindo, M.; Smith, C. C.; Kim, S.; Nadizadeh, H.; Suzuki, Y.; Tao, C.; Vu, P.; Tang, S.; Zhang, P.; Murthi, K. K.; Gentile, L. N.; Liu, J. H. *J. Am. Chem. Soc.* **1994**, 116, 1599; d) **efrotomycin**: Dolle, R. E.; Nicolaou, K. C. *J. Am. Chem. Soc.* **1985**, 107, 1691; e) **pravastatin**: Daniewski, A. R.; Wovkulich, P. M.; Uskokovic, M. R. *J. Org. Chem.* **1992**, 57, 7133; f) **balanol**: Sullivan, B.; Gilmet, J.; Leisch, H.; Hudlicky, T. *J. Nat. Prod.* **2008**, 71, 346–350.
- ^{xv} Edward M. Burgess, Harold R. Penton Jr., and E. A. Taylor. "Thermal reactions of alkyl N-carbomethoxysulfamate esters". *J. Org. Chem.* **1973**, 38(1):26–31.
- ^{xvi} Olofson, R. A.; Schnur, R. C.; Bunes, L.; Pepe, J. *Tetrahedron Lett.* **1977**, 18, 1567.
- ^{xvii} (a) Pillai, O.; Hamad, M.O.; Crooks, P.A.; Stinchcomb, A.L. *Pharm. Res.*, **2004**, 21, 1146; (b) Hamad, M.O.; Kiptoo, P.K.; Stinchcomb, A.L.; Crooks, P.A.; *Bioorg. Med. Chem.* **2006**, 14, 7051.
- ^{xviii} Naltrexone mp reference: Olofson et al. US Pat. No. 4,141,897 (1976).
- ^{xix} Ukrainets, I.V.; Tkach, A.A.; Gorokhova, O.V.; Turov, A.V.; Linsky, I.V.; in *Chemistry of Heterocyclic Compounds* (New York, NY, United States), 2009, vol. 45, #4, p. 405.
- ^{xx} Andre, J.; Dormoy, J.; Haymes, A. *Synth. Comm.*, **1992**, 22, 2313.
- ^{xxi} (a) Lewenstein et al. US Pat. No. 3,254,088 (1966); (b) Sankyo, Belg. Pat. No. 615,009 (1962).
- ^{xxii} Research Corp. Patent: US 4,161,597, 1979; *Chem. Abstr.* vol. 92, #22671.

We Claim:

1. A process for the preparation of a compound of Formula I:



wherein


---- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when $=O$ represents $=O$, then R^2 is not present; and

PG is a protecting group,

the process comprising:

- (a) reacting a compound of Formula II with an oxidizing agent under conditions to provide a compound of Formula III:



wherein

---- represents a single or double bond, provided that two double bonds are not adjacent to each other;


R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^2 is not present; and

PG is a protecting group; and

(b) reacting the compound of Formula III with a cyclodehydration reagent under conditions to provide the compound of Formula I,

wherein in the compounds of Formulae I, II and III, one or more available hydrogens in R^1 and R^2 is/are optionally replaced with F and/or one or more of available atoms in R^1 and R^2 is/are optionally replaced with an isotopic label.

2. The process of claim 1, wherein the compound of Formula I is selected from a compound of Formula I(a), I(b) and I(c):



wherein


--- represents a single or double bond;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, and PG is a protecting group; and

one or more available hydrogens in R^1 and R^2 is/are optionally replaced with F and/or one or more of available atoms in R^1 and R^2 is/are optionally replaced with an isotopic label.

3. The process of claim 1 or 2, wherein R^1 and R^2 are independently selected from C_{1-6} alkyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{1-6} alkylene C_{6-10} aryl, C_{1-6} alkylene C_{3-6} cycloalkyl and PG.

4. The process of claim 3, wherein R^1 and R^2 are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG.
5. The process of any one of claims 1 to 4, wherein the oxidizing agent is a peroxide or a peracid.
6. The process of claim 5, wherein the oxidizing agent is *m*-chloroperbenzoic acid (*m*CPBA).
7. The process of any one of claims 1 to 6, wherein the cyclodehydration reagent is selected from Burgess reagent, TsCl, CrO_3 , DCC, XtalFluorTM and carbonyldiimidazole, preferably Burgess reagent.
8. The process of claim 7, wherein the conditions to provide the compounds of Formula I from the compounds of Formula III using Burgess reagent comprise a temperature of about $-50\text{ }^{\circ}\text{C}$ to about $50\text{ }^{\circ}\text{C}$, in an inert organic solvent or mixture of organic solvents for a time for the conversion of the compound of Formula III to the compound of Formula I to proceed to a sufficient extent.
9. The process of claim 8, wherein the Burgess reagent and the compound of Formula III are used in a molar ratio of about 1.5:1 to about 1:1.
10. A process for preparing compounds of Formula V:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when $=O$ represents $=O$, then R^2 is not present;

PG is a protecting group;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl; and

X is a counteranion,

comprising

reacting a compound of Formula I as defined in any one of claims 1 to 4:

with an alkylating reagent of Formula VI:



wherein R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl and LG is a leaving group, under conditions to provide the compound of Formula V, wherein in the compounds of Formulae I, V and VI, one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label.


11. The process of claim 10, wherein R^3 is selected from C_{1-6} alkyl, C_{2-6} alkenyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkylene C_{6-10} aryl, and C_{1-6} alkylene C_{3-6} cycloalkyl.

12. The process of claim 11, wherein R^3 is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

13. The process of any one of claims 10 to 12, wherein LG is selected from halo, Ms, Ts, Ns, Tf and C₁₋₆acyl.

14. The process of any one of claims 10 to 13, wherein X is the anion of LG.

15. The process of any one of claims 10 to 14, wherein the compound of Formula V is selected from a compound of Formula V(a), V(b) and V(c):



wherein

--- represents a single or double bond; and


R¹ and R² are independently selected from C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, and PG is a protecting group;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R¹, R² and R³ is/are optionally replaced with F and/or one or more of available atoms in R¹, R² and R³ is/are optionally replaced with an isotopic label.

16. A process for preparing compounds of Formula VII:



wherein

---- represents a single or double bond, provided that two double bonds are not adjacent to each other and when $\equiv\text{O}$ represents $=\text{O}$, then H is not present;


R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion; and

one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label,

the process comprising deprotecting the compounds of Formula V as defined in any one of claims 10 to 15, wherein R^1 and/or R^2 are PG.

17. A process for preparing compounds of Formula VIII:



wherein

---- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;


R^4 and R^5 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when $\equiv\text{O}$ represents $=\text{O}$, then R^5 is not present, or

R^4 and R^5 are H if, in the compounds of Formula V, R^1 and R^2 are a PG that is removed under the reducing conditions;

PG is a protecting group that is not removed under the reducing conditions, and

one or more available hydrogens in R^3 , R^4 and R^5 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^4 and R^5 is/are optionally replaced with an isotopic label,
the process comprising treating compounds of Formula V as defined in any one of claims 10 to 15 under reducing conditions.

18. A process for preparing compounds of Formula IX:



wherein

— represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

R^6 and R^7 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^7 is not present, or

R^6 and R^7 are H if, in the compounds of Formula V, R^1 and R^2 are a PG that is removed under the hydrolysis conditions;


PG is a protecting group that is not removed under the hydrolysis conditions, and one or more available hydrogens in R^3 , R^6 and R^7 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^6 and R^7 is/are optionally replaced with an isotopic label,

the process comprising treating compounds of Formula V as defined in any one of claims 10 to 15 under hydrolysis conditions.

19. The process of claim 17, wherein the reducing conditions to provide the compounds of Formula VIII comprise treating the compounds of Formula V with a suitable reducing agent, such as metal hydride reducing agents, optionally in the presence of a Lewis acid, for a time and temperature for the conversion of the compound of Formula V to the compound of Formula VIII to proceed to a sufficient extent.

20. The process of claim 18, wherein the hydrolysis conditions to provide the compounds of Formula IX comprise treating the compounds of Formula V under suitable acidic (for example acetic acid/ammonia buffer) or basic (for example ammonium bicarbonate/ammonia) conditions for a time and temperature for the conversion of the compound of Formula V to the compound of Formula IX to proceed to a sufficient extent.

21. A process for preparing compounds of Formula X:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^8 and R^9 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when $=O$ represents $=O$, then R^9 is not present; or

R^8 and R^9 are H, if in the compound of Formula I, R^1 and R^2 are a PG that is removed under the hydrolysis conditions; and


PG is a protecting group,

wherein in the compounds of Formula X, one or more available hydrogens in R^8 and R^9 is/are optionally replaced with F and/or one or more of available atoms in R^8 and R^9 is/are optionally replaced with an isotopic label, the process comprising treating the compounds of Formula I as defined in any one of claims 1 to 9 under either acidic or basic conditions.

22. The process of claim 21, wherein PG in the compounds of Formula I is a protecting group that is removed under conditions to hydrolyze the compound of Formula I to the compound of Formula X.

23. The process of claim 21, wherein PG is a protecting group that is not removed under conditions to hydrolyze the compound of Formula I to the compound of Formula X and is optionally removed in a separate step after the preparation of the compound of Formula X.

24. A process for preparing compounds of Formula XII:



wherein

— represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^8 and R^9 are independently selected from H, C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when $=O$ represents $=O$, then R^9 is not present;

PG is a protecting group;


R^{10} is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl; and

one or more available hydrogens in R^8 , R^9 and R^{10} is/are optionally replaced with F and/or one or more of available atoms in R^8 , R^9 and R^{10} is/are optionally replaced with an isotopic label,

the process comprising selectively alkylating the N-17 of the compounds of Formula X as defined in any one of claims 21-23 by reaction with a compound of the Formula R^{10} -LG¹ (XI), wherein LG¹ is a leaving group and R^{10} is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl under alkylation conditions.

25. The process of claim 24, wherein R^8 and/or R^9 in the compounds of Formula XII is PG, and the compounds of Formula XII are further treated under conditions to remove the PG to provide the corresponding compounds of Formula XII wherein R^8 and/or R^9 are H.

26. A compound of Formula V:



wherein

--- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, except when ---O represents =O , then R^2 is not present;

PG is a protecting group;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label, or a salt or solvate thereof.

27. The compound of claim 26, wherein R^1 and R^2 are independently selected from C_{1-6} alkyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{1-6} alkylene C_{6-10} aryl, C_{1-6} alkylene C_{3-6} cycloalkyl and PG.


28. The compound of claim 27, wherein R^1 and R^2 are independently selected from Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG.

29. The compound of any one of claims 26 to 28, wherein R^3 is selected from C_{1-6} alkyl, C_{2-6} alkenyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkylene C_{6-10} aryl, and C_{1-6} alkylene C_{3-6} cycloalkyl.

30. The compound of claim 29, wherein R^3 is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

31. The compound of any one of claims 26 to 30, wherein X is OH^- , Br^- or Cl^- .

32. The compound of claim 26, selected from a compound of Formula V(a), V(b) and V(c):



wherein

— represents a single or double bond; and


R^1 and R^2 are independently selected from C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG, and PG is a protecting group;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion, and

one or more available hydrogens in R^1 , R^2 and R^3 is/are optionally replaced with F and/or one or more of available atoms in R^1 , R^2 and R^3 is/are optionally replaced with an isotopic label,
or a salt or solvate thereof.

33. A compound of Formula VII:



wherein

— represents a single or double bond, provided that two double bonds are not adjacent to each other and when $=O$ represents $=O$, then H is not present;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

X is a counteranion; and


one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label, or
a salt or solvate thereof.

34. The compound of claim 33, wherein R^3 is selected from C_{1-6} alkyl, C_{2-6} alkenyl, phenyl, naphthyl, indanyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, C_{1-6} alkylene C_{6-10} aryl, and C_{1-6} alkylene C_{3-6} cycloalkyl.

35. The compound of claim 34, wherein R^3 is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

36. The compound of any one of claims 33 to 35, wherein X is OH^- , Br^- or Cl^- .

37. The compound of claim 33, selected from a compound of Formula VII(a), VII(b) and VII(c):



wherein

— represents a single or double bond;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;


X is a counteranion; and

one or more available hydrogens in R^3 is/are optionally replaced with F and/or one or more of available atoms in R^3 is/are optionally replaced with an isotopic label, or

a salt or solvate thereof.

38. A compound of Formula VIII:

66



wherein

----- represents a single or double bond, provided that two double bonds are not adjacent to each other;

R³ is selected from C₃₋₁₀cycloalkyl, C₃₋₁₀cycloalkenyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₆₋₁₀aryl, C₁₋₁₀alkyleneC₆₋₁₀aryl and C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl;

R⁴ and R⁵ are independently selected from H, C₁₋₁₀alkyl, C₆₋₁₀aryl, C₃₋₁₀cycloalkyl, C₁₋₁₀alkyleneC₆₋₁₀aryl, C₁₋₁₀alkyleneC₃₋₁₀cycloalkyl and PG, except when =O represents =O, then R⁵ is not present;

PG is a protecting group; and

one or more available hydrogens in R³, R⁴ and R⁵ is/are optionally replaced with F and/or one or more of available atoms in R³, R⁴ and R⁵ is/are optionally replaced with an isotopic label, or
a salt or solvate thereof.


39. The compound of claim 38, wherein R³ is selected from C₁₋₆alkyl, C₂₋₆alkenyl, phenyl, naphthyl, indanyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, C₁₋₆alkyleneC₆₋₁₀aryl, and C₁₋₆alkyleneC₃₋₆cycloalkyl.

40. The compound of claim 39, wherein R³ is selected from Me, Et, allyl, Ph, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

41. The compound of any one of claims 38 to 40, wherein R⁴ and R⁵ are independently selected from H, C₁₋₆alkyl, phenyl, naphthyl, indanyl, C₃₋₆cycloalkyl, C₁₋₆alkyleneC₆₋₁₀aryl, C₁₋₆alkyleneC₃₋₆cycloalkyl and PG.

42. The compound of claim 41, wherein R^4 and R^5 are independently selected from H, Me, Et, Ph, cyclobutyl, cyclopentyl, cyclohexyl, Bn, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl and PG.

43. The compound of claim 38, selected from a compound of Formula VIII(a), VIII(b) and VIII(c):



wherein

---- represents a single or double bond;

R^3 is selected from C_{3-10} cycloalkyl, C_{3-10} cycloalkenyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{6-10} aryl, C_{1-10} alkylene C_{6-10} aryl and C_{1-10} alkylene C_{3-10} cycloalkyl;

R^4 and R^5 are independently selected from H, C_{1-10} alkyl, C_{6-10} aryl, C_{3-10} cycloalkyl, C_{1-10} alkylene C_{6-10} aryl, C_{1-10} alkylene C_{3-10} cycloalkyl and PG;

PG is a protecting group; and

one or more available hydrogens in R^3 , R^4 and R^5 is/are optionally replaced with F and/or one or more of available atoms in R^3 , R^4 and R^5 is/are optionally replaced with an isotopic label, or
a salt or solvate thereof.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2012/000384

A. CLASSIFICATION OF SUBJECT MATTER IPC: C07D 498/22 (2006.01) , C07D 489/08 (2006.01) According to International Patent Classification (IPC) or to both national classification and IPC																					
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC: C07D 498/00 (2006.01), C07D 489/00 (2006.01) Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used) STN (structure search in Registry for Formulae I, III, V, VII, VIII, IX, X, XII and VIII, CAPlus), EPOQUE (EPODOC: IPC and Inventor search), Canadian Patent Database (IPC and Inventor search)																					
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>WO 2005/028483 (SMITH et al) 31 March 2005 (31-03-2005) See the whole document.</td> <td>1-6, 21-25</td> </tr> <tr> <td>P, X</td> <td>WO 2012/013671 (SCHMIDHAMMER) 02 February 2012 (02-02-2012) See the whole document.</td> <td>38-43</td> </tr> <tr> <td>X</td> <td>DE 34 12 727 (SCHMIDHAMMER) 17 October 1985 (17-10-1985) See the whole document.</td> <td>38-43</td> </tr> <tr> <td>X</td> <td>R. KRASSNIG et al: "A New and Efficient Synthesis of the μ Opioid Receptor Antagonists 14-<i>O</i>-Methyl- and 14-<i>O</i>-Ethyl naloxone and -Naltrexone"; <i>Heterocycles</i>, 1998, 47(2), 1029-1032. See the whole document.</td> <td>38-43</td> </tr> <tr> <td>X</td> <td>R. KRASSNIG et al: "A Novel Method for the Introduction of a 5β-Methyl Group into 4,5α-Epoxy morphinan-6-ones <i>via</i> the Enol Ether"; <i>Helvetica Chimica Acta</i>, 2000, 83(2), 380-383. See the whole document.</td> <td>38-43</td> </tr> </tbody> </table>				Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	WO 2005/028483 (SMITH et al) 31 March 2005 (31-03-2005) See the whole document.	1-6, 21-25	P, X	WO 2012/013671 (SCHMIDHAMMER) 02 February 2012 (02-02-2012) See the whole document.	38-43	X	DE 34 12 727 (SCHMIDHAMMER) 17 October 1985 (17-10-1985) See the whole document.	38-43	X	R. KRASSNIG et al: "A New and Efficient Synthesis of the μ Opioid Receptor Antagonists 14- <i>O</i> -Methyl- and 14- <i>O</i> -Ethyl naloxone and -Naltrexone"; <i>Heterocycles</i> , 1998, 47 (2), 1029-1032. See the whole document.	38-43	X	R. KRASSNIG et al: "A Novel Method for the Introduction of a 5 β -Methyl Group into 4,5 α -Epoxy morphinan-6-ones <i>via</i> the Enol Ether"; <i>Helvetica Chimica Acta</i> , 2000, 83 (2), 380-383. See the whole document.	38-43
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																			
X	WO 2005/028483 (SMITH et al) 31 March 2005 (31-03-2005) See the whole document.	1-6, 21-25																			
P, X	WO 2012/013671 (SCHMIDHAMMER) 02 February 2012 (02-02-2012) See the whole document.	38-43																			
X	DE 34 12 727 (SCHMIDHAMMER) 17 October 1985 (17-10-1985) See the whole document.	38-43																			
X	R. KRASSNIG et al: "A New and Efficient Synthesis of the μ Opioid Receptor Antagonists 14- <i>O</i> -Methyl- and 14- <i>O</i> -Ethyl naloxone and -Naltrexone"; <i>Heterocycles</i> , 1998, 47 (2), 1029-1032. See the whole document.	38-43																			
X	R. KRASSNIG et al: "A Novel Method for the Introduction of a 5 β -Methyl Group into 4,5 α -Epoxy morphinan-6-ones <i>via</i> the Enol Ether"; <i>Helvetica Chimica Acta</i> , 2000, 83 (2), 380-383. See the whole document.	38-43																			
[X] Further documents are listed in the continuation of Box		[X] See patent family annex.																			
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family																				
Date of the actual completion of the international search 10 August 2012 (10-08-2012)		Date of mailing of the international search report 26 September 2012 (26-09-2012)																			
Name and mailing address of the ISA/CA Canadian Intellectual Property Office Place du Portage I, C114 - 1st Floor, Box PCT 50 Victoria Street Gatineau, Quebec K1A 0C9 Facsimile No.: 001-819-953-2476		Authorized officer Lu Jiang (819) 934-6738																			

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2012/000384

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	H. SCHMIDHAMMER et al.: "Synthesis and Biological Evaluation of 14-Alkoxymorphians. 1. Highly Potent Opioid Agonists in the Series of (-)-14-Methoxy-N-methylmorphinan-6-ones"; <i>J. Med. Chem.</i> , 1984, 27 , 1575-1579. See the whole document.	38-43
X	US 2010/0081819 (WANG et al) 01 April 2010 (01-04-2010) See the whole document.	24-25

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. ☐ Claim Nos. :
because they relate to subject matter not required to be searched by this Authority, namely :

2. ☐ Claim Nos. :
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :

3. ☐ Claim Nos. :
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

See the Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

- Remark on Protest** ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

The unique common concept linking the subject matter of claims 1-43 together seems to be compounds of Formula I or compounds of Formula X, which are already known in the art (*e.g.*, WO 2005/028483). Since the common linking feature is not novel, it cannot represent the single inventive concept which could have linked the different claimed subject matter together. Therefore, the claims are directed to a plurality of alleged inventions as follows:

Group A - Claims 1-9 are directed to a process for the preparation of a compound of Formula I from a compound of Formula II through a compound of Formula III reacting with a cyclodehydration reagent;

Group B - Claims 10-20 and 26-43 are directed to a process for preparing compounds of Formula V by alkylation of a compound of Formula I and subsequent processes of deprotection, reduction and hydrolysis of compounds of Formula V as well as the compounds produced by the above processes;

Group C - Claims 21-23 are directed to a process for the preparation of compounds of Formula X from a compound of Formula I by hydrolysis; and

Group D - Claims 24-25 are directed to a process for preparing compounds of Formula XII by alkylation of compounds of Formula X.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2012/000384

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
WO2005028483A1	31-03-2005	AU2004274038A1	31-03-2005
		AU2004274038B2	10-03-2011
		AU2004274038B8	04-08-2011
		AU2004274038C1	12-01-2012
		CA2539659A1	31-03-2005
		EP1664061A1	07-06-2006
		EP1664061A4	09-09-2009
		JP2007505935A	15-03-2007
		JP4879740B2	22-02-2012
		US2008045716A1	21-02-2008
		US7875718B2	25-01-2011
		US2011098474A1	28-04-2011
		US8106201B2	31-01-2012
WO2012013671A1	02-02-2012	EP2420503A1	22-02-2012
DE3412727A1	17-10-1985	None	
US2010081819A1	01-04-2010	AU2009300379A1	08-04-2010
		CA2738246A1	08-04-2010
		CN102227433A	26-10-2011
		EP2344506A2	20-07-2011
		US8080661B2	20-12-2011
		WO2010039209A2	08-04-2010
		WO2010039209A3	16-09-2010