

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2016/149370 A1

(43) International Publication Date
22 September 2016 (22.09.2016)

WIPO | PCT

(51) International Patent Classification:
C12N 15/86 (2006.01) C12N 15/113 (2010.01)

(21) International Application Number:
PCT/US2016/022645

(22) International Filing Date:
16 March 2016 (16.03.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/133,590 16 March 2015 (16.03.2015) US

(71) Applicant: MIRAGEN THERAPEUTICS, INC.
[US/US]; 6200 Lookout Road, Boulder, Colorado 80301
(US).

(72) Inventors: JACKSON, Aimee; c/o Miragen Therapeutics,
Inc., 6200 Lookout Road, Boulder, Colorado 80301 (US).
DALBY, Christina; 6200 Lookout Road, Boulder, Colorado
80301 (US).

(74) Agents: JOSHI, Trupti P. et al.; 1299 Pennsylvania Avenue, N.W., Suite 700, Washington, District of Columbia 20004-2400 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.





(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

[Continued on next page]

(54) Title: MIRNA MIMETICS AND THEIR USE IN TREATING SENSORY CONDITIONS



(57) Abstract: The present invention provides microRNA mimetic compounds that mimic the function or activity of miR-96, miR-182, and/or miR-183. The microRNA mimetic compounds of the invention comprise a first strand of about (22) to about (26) ribonucleotides comprising a mature miR-96, miR-182, or miR-183 sequence; and a second strand of about (20) to about (24) ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. The invention additionally provides expression vectors comprising a polynucleotide(s) encoding one or more of miR-96, miR-182, and miR-183. The invention also provides methods of treating ophthalmological or otic conditions by administering the microRNA mimetic compounds of miR-96, miR-182, and/or miR-183 and/or an expression vector encoding at least one of miR-96, miR-182, and miR-183 to a subject in need thereof.



Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

MIRNA MIMETICS AND THEIR USE IN TREATING SENSORY CONDITIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application claims the benefit of priority to U.S. Provisional Application No. 62/133,590, filed on March 16, 2015, the contents of which are hereby incorporated by reference herein in their entirety.

FIELD OF THE INVENTION

[0002] The present invention provides microRNA mimetic compounds that mimic the function or activity of miR-96, miR-182, and/or miR-183. The invention also provides expression vectors comprising a polynucleotide(s) encoding one or more of miR-96, miR-182, and miR-183. The invention further provides compositions comprising miR-96, miR-182, and/or miR-183 mimetic compounds or expression vectors encoding miR-96, miR-182, and/or miR-183, and methods of treating ophthalmological and otic conditions using the microRNA mimetic compounds or expression vectors encoding them.

BACKGROUND

[0003] MicroRNAs are small, endogenous, noncoding RNAs that act as posttranscriptional repressors of gene expression. MicroRNAs have unique expression profiles in the developing and adult retina and are involved in normal development and functions of the retina (Ryan *et al.*, Mol Vis 12:1175–1184, 2006; Xu *et al.*, J Biol Chem 282(34):25053–25066, 2007). MiRNAs are dysregulated in the retina of retinal degenerative mouse models, suggesting their potential involvement in retinal degeneration (Loscher *et al.*, Genome Biol 8(11):R248, 2007; Loscher *et al.*, Exp Eye Res 87(6):529–534 2008). Conditional inactivation of dicer, an RNase III endonuclease required for miRNA maturation in cytosol, in the mouse retina resulted in alteration of retinal differentiation and optic-cup patterning, increased cell death, and disorganization of axons of retinal ganglion cells (Pinter & Hindges, PLoS ONE 5(4):e10021, 2010; Davis & Ashery-Padan, Development 138(1):127–138, 2011; Georgi & Reh, J Neurosci 30(11):4048–4061, 2010), suggesting that miRNAs are important for normal development and functions of the mammalian retina. However, *in vivo* functions of individual miRNAs in the retina are still largely unknown.

[0004] The microRNA-183/96/182 cluster is expressed in the retina and other sensory organs. A knock-out mouse model of the miR-183/96/182 cluster showed that that the inactivation of the cluster during development results in the early-onset and progressive synaptic defects of the photoreceptors and progressive retinal degeneration (Lumayag *et al.*, Proc Natl Acad Sci, 110(6):E507-16, 2013). On the other hand, a transgenic anti-miRNA “sponge” mouse model that reduced the activities of all three miRNAs in the miR-183/96/182 cluster showed increased bright-light induced retinal degeneration; however, no histological or functional defects of the retina were observed under normal lighting conditions (Zhu *et al.*, J Biol Chem., 286(36):31749-60, 2011). Thus, the role of the miR-183/96/182 cluster in adult retina remains uncertain.

[0005] While many studies have shown therapeutic efficacy using single-stranded miRNA inhibitors called antimiRs, efforts to restore or increase the function of a miRNA have been lagging behind (van Rooij *et al.*, Cir Res, 110:496-507, 2012). Currently, miRNA function can be increased either by viral overexpression or by using synthetic double-stranded miRNAs. The use of adeno-associated viruses (AAV) to drive expression of a given miRNA for restoring its activity *in vivo* has shown to be effective in a mouse model of hepatocellular and lung carcinoma (Kasinski & Slack, Cancer Res, 72: 5576-5587, 2012; Kota *et al.*, Cell, 137: 1005-1017, 2009) and spinal and bulbar muscular atrophy (Miyazaki *et al.*, Nat Med., 18(7):1136-41, 2012). The use of synthetic oligonucleotide-based approaches to increase miRNA levels has not been well explored yet. The present invention provides synthetic oligonucleotides as well as virally expressed polynucleotides that mimic the activity of miR-96, miR-182 and/or miR-183 to restore eye and ear function.

SUMMARY OF THE INVENTION

[0006] The present invention provides microRNA mimetic compounds that mimic the function or activity of microRNAs, in particular, the function or activity of miR-96, miR-182, and/or miR-183. In some embodiments, the microRNA mimetic compounds of the invention include synthetic oligonucleotides. In other embodiments, the invention provides expression vectors comprising a polynucleotide(s) encoding one or more of miR-96, miR-182, and miR-183 for expression in a mammalian cell for treating eye or ear dysfunction.

[0007] In certain embodiments, the microRNA mimetic compounds of the invention comprise synthetic oligonucleotides comprising a first strand and a second strand, where the two strands form a double stranded region that is fully or partially complementary. In various embodiments, the first strand or the antisense strand of the microRNA mimetic compound comprises the sequence of a mature miR-96, miR-182, or miR-183 and the second strand or the sense strand comprises a sequence that is substantially complementary to the first strand and has at least one modified nucleotide.

[0008] The microRNA mimetic compounds of the invention mimic the function or activity of a mature, naturally-occurring miR-96, miR-182, or miR-183 microRNA and show enhanced resistance to the nuclease digestion of the antisense strand, improved ability to load the antisense strand into the miRNA-induced silencing complex (miRISC), and/or rapid degradation of the sense strand.

[0009] In some embodiments, the first strand of the miRNA mimetic compound is about 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 nucleotides in length and comprises the sequence of a mature miR-96, miR-182, or miR-183 and the second strand is about 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27 nucleotides in length and comprises a sequence that is substantially complementary to the first strand, wherein the second strand comprises at least one modified nucleotide. In certain embodiments, the first strand is about 22 to about 26 nucleotides in length and the second strand is about 20 to about 24 nucleotides in length. In some embodiments, the first strand is about 22 to about 26 nucleotides in length and the second strand is about 20 to about 26 nucleotides in length.

[0010] In some embodiments, the sequence of the first strand of the microRNA mimetic compounds of the invention is about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, inclusive of values therebetween, identical to the mature miR-96, miR-182, or miR-183 sequence. In other embodiments, the sequence of the first strand is completely (100%) identical to the mature miR-96, miR-182, or miR-183 sequence. In some embodiments, the first strand may comprise a 3' nucleotide overhang relative to the second strand. In other embodiments, the second strand may comprise a 3' nucleotide overhang relative to the first strand. The 3' overhang on the first or the second strand may comprise

about 1, 2, 3, or 4 nucleotides. In certain embodiments, the 3' overhang is about 1 or 2 nucleotides in length. In some embodiments, the first and the second strand contain the same number of nucleotides, *i.e.*, there is no overhang on either strand. In some embodiments, the overhang may be present on the 5' end of the first or the second strand.

[0011] In one embodiment, the sequence of the second strand is fully complementary (100%) to the sequence of the first strand. In another embodiment, the sequence of the second strand is substantially complementary, such as about 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%, inclusive of values therebetween, complementary to the first strand. In still another embodiment, the sequence of the second strand may contain about 1, 2, 3, 4, or 5 mismatches relative to the first strand.

[0012] In some embodiments, the first strand and/or the second strand may comprise one or more modified nucleotides. In some embodiments, the first strand comprising a mature miR-96, miR-182, or miR-183 sequence contains one or more modified nucleotides, such as one or more 2'-fluoro modified nucleotides. In certain embodiments, the first strand contains at least one 2'-fluoro nucleotide. In some embodiments, the second strand comprising a sequence that is substantially complementary to the sequence of the first strand comprises at least one modified nucleotide, such as a 2'-fluoro or 2'-O-methyl modified nucleotide. In certain embodiments, the at least one modified nucleotide in the second strand is a 2'-O-methyl modified nucleotide.

[0013] In some embodiments, the present invention provides a microRNA mimetic compound comprising a first strand of about 22 to about 26 ribonucleotides comprising a mature miR-96, miR-182, or miR-183 sequence; and a second strand of about 20 to about 24 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In some other embodiments, the present invention provides a microRNA mimetic compound comprising a first strand of about 22 to about 26 ribonucleotides comprising a mature miR-96, miR-182, or miR-183 sequence; and a second strand of about 20 to about 26 ribonucleotides comprising a sequence that is substantially

complementary to the first strand and having at least one modified nucleotide, wherein the first or the second strand has a 3' nucleotide overhang.

[0014] The invention further provides expression vectors comprising a polynucleotide encoding miR-96, miR-182, and/or miR-183 positioned for expression in a mammalian cell for use as a therapeutic agent for the treatment of ophthalmological or otic disorders. For example, the sequence encoding miR-96, miR-182, and/or miR-183 is positioned adjacent to an appropriate promoter for expression in an eye or ear cell and the promoter and coding sequence are flanked by inverted terminal repeats. The sequence is then further inserted into a vector sequence, which in certain embodiments can replicate in a human cell. The polynucleotide can be an isolated polynucleotide.

[0015] In certain embodiments, the vector is a viral expression vector. In certain embodiments, the viral expression vector is an adenoviral vector, an adeno-associated viral (AAV) vector, or a lentiviral vector. In some embodiments, the viral expression vector is a self-complementary adeno-associated viral vector. In certain embodiments, the AAV vector is based on a single serotype of AAV, such as AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, and AAV9. Modified AAV vectors based substantially on a single serotype are known in the art, for example serotype rh.74 which is AAV8-like and shares 93% amino acid identity with AAV8 can be used (see, e.g., Martin et al., Am. J. Cell. Physiol. 296:C476-C488, 2009, incorporated herein by reference). Alternatively, the adeno-associated viral vector is a chimeric adeno-associated viral vector based on multiple serotypes of AAV.

[0016] The expression vectors provided by the instant invention can include any sequence that encodes a functional miR-96, miR-182, and/or miR-183 for use in any of the methods of the instant invention. In certain embodiments, the expression vector includes a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 53-55.

[0017] The invention provides cells containing a polynucleotide encoding miR-96, miR-182, and/or miR-183 of the instant invention. In some embodiments, the cell is a bacterial cell or a mammalian cell. The cell can be a cell to which the polynucleotide has been delivered as a therapeutic intervention. Alternatively, the cell can be a cell in vitro used for

the preparation of a pharmaceutical composition for the treatment of ophthalmological or otic disorders.

[0018] In some embodiments, the invention provides compositions comprising at least one of the miR-96, miR-182, and miR-183 mimetic compounds or pharmaceutically acceptable salts thereof, and pharmaceutically acceptable carriers or excipients. In some other embodiments, the invention provides compositions comprising an expression vector encoding at least one of miR-96, miR-182, and miR-183, and pharmaceutically acceptable carriers or excipients.

[0019] The invention further provides methods of treating or preventing ophthalmological conditions such as retinal degeneration or retinitis pigmentosa comprising administering at least one of the miR-96, miR-182, and miR-183 mimetic compounds described herein to a subject in need thereof. The invention also encompasses methods for improving or restoring visual acuity in a subject in need thereof comprising administering at least one of the miR-96, miR-182, and miR-183 mimetic compounds described herein to a subject in need thereof.

[0020] The invention further provides methods of treating or preventing ophthalmological conditions such as retinal degeneration or retinitis pigmentosa comprising administering an expression vector encoding at least one of miR-96, miR-182, and miR-183 to a subject in need thereof. The invention also encompasses methods for improving or restoring visual acuity in a subject in need thereof comprising administering an expression vector encoding at least one of miR-96, miR-182, and miR-183 to a subject in need thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] FIGs. 1A-1D show quantitation of miRNA mimics and Rho siRNA in mouse retina using a sandwich ELISA assay. Values shown are pmol oligo/g of retina tissue and represent 4 retinas (microRNAs) or 2 retinas (siRNA). FIG. 1A shows the amount of miR-183 mimic, FIG. 1B shows the amount of miR-96 mimic, FIG. 1C shows the amount of miR-182 mimic, and FIG. 1D shows the amount of Rho siRNA.

[0022] Figure 2 shows the functional delivery of Rho siRNA to mouse retina and down-regulation of Rho mRNA. The solid line indicates the amount of rhodopsin siRNA detected (nmol/g retina tissue) and the dotted line indicates the fraction of rhodopsin mRNA detected in retina compared to untreated eyes (time 0).

[0023] FIGs. 3A-3B show the effect of passive transfection of a miRNA mimic in R-Ret cells. FIG. 3A shows R-Ret cells transfected with 10 μ M miR-206 mimic 72h post-transfection and FIG. 3B shows untreated R-Ret cells.

[0024] Figure 4 shows the results of an adenylate kinase assay of R-Ret cells transfected with miR-206 mimic.

[0025] Figure 5 shows a real time PCR analysis of Rho mRNA in R-Ret cells transfected with various concentrations of cholesterol-conjugated Rho siRNA.

[0026] FIGs. 6A-6B show a real time PCR analysis of mRNA expression profile of genes involved in the phototransduction pathway following exposure to 1 μ M oligonucleotides. FIG. 6A shows genes that were up-regulated following exposure to 1 μ M oligonucleotides. FIG. 6B shows genes that were down-regulated following exposure to 1 μ M oligonucleotides.

[0027] FIGs. 7A-7B show a real time PCR analysis of mRNA expression profile of genes involved in the phototransduction pathway following exposure to 3 μ M oligonucleotide. FIG. 7A shows genes that were up-regulated following exposure to 3 μ M oligonucleotides. FIG. 7B shows genes that were down-regulated following exposure to 3 μ M oligonucleotides.

[0028] Figure 8 shows a heat map of the log2-transformed average fold change values of the treatments shown in FIG. 6-7.

[0029] Figure 9 shows the effect of pooled microRNA mimics and Rho siRNA on visual acuity loss in the mouse model of retinitis pigmentosa.

[0030] Figure 10 shows the effect of pooled microRNA mimics and Rho siRNA on visual acuity loss in the mouse model of retinitis pigmentosa.

DETAILED DESCRIPTION OF THE INVENTION

[0031] MicroRNAs (miRNAs) are small, non-protein coding RNAs of about 18 to about 25 nucleotides in length that are derived from individual miRNA genes, from introns of protein coding genes, or from poly-cistronic transcripts that often encode multiple, closely related miRNAs. See review by Carrington *et al.* (*Science*, Vol. 301(5631):336-338, 2003). MiRNAs act as repressors of target mRNAs by promoting their degradation or by inhibiting translation.

[0032] MiRNAs are transcribed by RNA polymerase II (pol II) or RNA polymerase III (pol III; see Qi *et al.* (2006) *Cellular & Molecular Immunology*, Vol. 3:411-419) and arise from initial transcripts, termed primary miRNA transcripts (pri-miRNAs), that are generally several thousand bases long. Pri-miRNAs are processed in the nucleus by the RNase Drosha into about 70- to about 100-nucleotide hairpin-shaped precursors (pre-miRNAs). Following transport to the cytoplasm, the hairpin pre-miRNA is further processed by Dicer to produce a double-stranded miRNA. The mature miRNA strand is then incorporated into the RNA-induced silencing complex (RISC), where it associates with its target mRNAs by base-pair complementarity. In the relatively rare cases in which a miRNA base pairs perfectly with an mRNA target, it promotes mRNA degradation. More commonly, miRNAs form imperfect heteroduplexes with target mRNAs, affecting either mRNA stability or inhibiting mRNA translation.

[0033] The microRNA-183/96/182 cluster is expressed in the retina and other sensory organs. This cluster is located on chromosome 6 in mouse, chromosome 7 in human, and chromosome 4 in rat. The sequences for mature miR-96, miR-182, and miR-183 in mouse, human, and rat are given below.

Mature miR-96 sequence in mouse (SEQ ID NO: 1)

UUUGGCACUAGCACAUUUUGCU

Mature miR-96 sequence in human (SEQ ID NO: 2)

UUUGGCACUAGCACAUUUUGCU

Mature miR-96 sequence in rat (SEQ ID NO: 3)

UUUGGCACUAGCACAUUUUGCU

Mature miR-182 sequence in mouse (SEQ ID NO: 4)

UUUGGCAAUGGUAGAACUCACACCG

Mature miR-182 sequence in human (SEQ ID NO: 5)

UUUGGCAAUGGUAGAACUCACACU

Mature miR-182 sequence in rat (SEQ ID NO: 6)

UUUGGCAAUGGUAGAACUCACACCG

Mature miR-183 sequence in mouse (SEQ ID NO: 7)

UAUGGCACUGGUAGAAUUCACU

Mature miR-183 sequence in human (SEQ ID NO: 8)

UAUGGCACUGGUAGAAUUCACU

Mature miR-183 sequence in rat (SEQ ID NO: 9)

UAUGGCACUGGUAGAAUUCACU

[0034] A knockout of the microRNA-183/96/182 cluster in mice resulted in the early-onset and progressive synaptic defects of the photoreceptors and progressive retinal degeneration (Lumayag *et al.*, (2013) *Proc Nat Acad Sci*, 110(6) E507-E516). Although the microRNA-183/96/182 cluster has been shown to play a role in the development of the retina; it is not yet clear whether restoring or supplementing the activity of the microRNA-183/96/182 cluster in adults could ameliorate or prevent retinal disorders.

[0035] The present invention is based, in part, on the discovery that the administration of microRNA mimetic compounds that mimic the activity of miR-96, miR-182, and/or miR-183 regulates expression of phototransduction genes in retinal cells, prevents or reduces the death of photoreceptor cells, and prevents or reduces the loss of vision in a mouse model of retinitis pigmentosa. Accordingly, the present invention provides microRNA mimetic compounds,

expression vectors encoding miR-96, miR-182, and/or miR-183, compositions thereof, and methods thereof, for treating or preventing ophthalmological conditions. The invention also contemplates the use of miR-96, miR-182, and/or miR-183 mimetic compounds and expression vectors encoding miR-96, miR-182, and/or miR-183 for treating, improving, or preventing diseases or disorders of other sensory organs, for example, ear.

[0036] A microRNA mimetic compound according to the invention comprises a first strand and a second strand, wherein the first strand comprises a mature miR-96, miR-182, or miR-183 sequence and the second strand comprises a sequence that is substantially complementary to the first strand and has at least one modified nucleotide, wherein the microRNA mimetic compound mimics the activity of miR-96, miR-182, or miR-183 microRNA. The term “microRNA agonist” as used herein refers to a synthetic microRNA mimetic compound or an expression vector encoding miR-96, miR-182, and/or miR-183. Throughout the disclosure, the term “first strand” may be used interchangeably with the term “antisense strand” or “guide strand”; and the term “second strand” may be used interchangeably with the term “sense strand” or “passenger strand.”

Synthetic microRNA mimetic compounds

[0037] In one embodiment, the first strand of the microRNA mimetic compound comprises from about 20 to about 28 nucleotides and the second strand comprises from about 18 to about 26 nucleotides. In various embodiments, the first strand may comprise about 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 nucleotides and the second strand may comprise about 20, 21, 22, 23, 24, 25, or 26 nucleotides. In certain embodiments, the first strand comprises from about 22 to about 26 nucleotides comprising a sequence of mature miR-96, miR-182, or miR-183, and the second strand comprises from about 20 to about 24 nucleotides comprising a sequence that is fully or substantially complementary to the first strand. In other embodiments, the first strand comprises from about 22 to about 26 nucleotides comprising a sequence of mature miR-96, miR-182, or miR-183, and the second strand comprises from about 20 to about 26 nucleotides comprising a sequence that is fully or substantially complementary to the first strand.

[0038] The nucleotides that form the first and the second strand of the microRNA mimetic compounds may comprise ribonucleotides, deoxyribonucleotides, modified nucleotides, and combinations thereof. In certain embodiments, the first strand and the second strand of the microRNA mimetic compound comprise ribonucleotides and/or modified ribonucleotides. The term “modified nucleotide” means a nucleotide where the nucleobase and/or the sugar moiety is modified relative to unmodified nucleotides.

[0039] In certain embodiments, the microRNA mimetic compounds have a first strand or an antisense strand comprising a “miRNA region” whose sequence is identical to all or part of a mature miR-96, miR-182, or miR-183 sequence, and a second strand or a sense strand having a “complementary region” whose sequence is from between about 70% to about 100% complementary to the sequence of the miRNA region. The term “miRNA region” refers to a region on the first strand of the miRNA mimetic compound that is at least about 75, 80, 85, 90, 95, or 100% identical, including all integers there between, to the entire sequence of a mature, naturally occurring miR-96, miR-182, or miR-183 sequence. In certain embodiments, the miRNA region is about or is at least about 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identical to the sequence of a mature, naturally-occurring miRNA, such as the mouse, human, or rat miR-96, miR-182, or miR-183 sequence. For example, in some embodiments, the miRNA region is about or at least about 99, 99.1, 99.2, 99.3, 99.4, 99.5, 99.6, 99.7, 99.8, 99.9 or 100 % identical to the sequence of a mature, naturally-occurring miRNA, such as the mouse, human, or rat miR-96, miR-182, or miR-183 sequence. Alternatively, the miRNA region can comprise 18, 19, 20, 21, 22, 23, 24, 25 or more nucleotide positions in common with a mature, naturally-occurring miRNA as compared by sequence alignment algorithms and methods well known in the art. It is understood that the sequence of the miRNA region of the first strand may include modifications of the nucleotides compared to the sequence of a mature, naturally-occurring miRNA. For example, if a mature, naturally-occurring miRNA sequence comprises a cytidine nucleotide at a specific position, the miRNA region of the first strand of the mimetic compound may comprise a modified cytidine nucleotide, such as 2'-fluoro-cytidine, at the corresponding position or if a mature, naturally-occurring miRNA sequence comprises a uridine nucleotide at a specific position, the miRNA region of the first strand of the mimetic compound may comprise a modified uridine nucleotide, such as 2'-fluoro-uridine, 2'-O-methyl-uridine, 5-

fluorouracil, or 4-thiouracil at the corresponding position. Even if the sequence of the miRNA region of the first strand includes such modified nucleotides, the sequence is still considered identical to the sequence of the mature, naturally-occurring miRNA sequence as long as the nucleotide that is modified has the same base-pairing capability as the nucleotide present in the mature, naturally-occurring miRNA sequence. In some embodiments, the first strand may include a modification of the 5'-terminal residue. For example, the first strand may have a 5'-terminal monophosphate.

[0040] The term “complementary region” refers to a region on the second strand of the miRNA mimetic compound that is at least about 70% complementary to the sequence of the miRNA region on the first strand. For example, the complementary region is at least about 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%, inclusive of all values therebetween, complementary to the sequence of the miRNA region. In certain embodiments, about 18, 19, 20, 21, 22, or 23 nucleotides of the complementary region of the second strand may be complementary to the first strand. In some embodiments, the complementary region of the second strand comprises about 1, 2, 3, 4, or 5 mismatches relative to the miRNA region of the first strand. That is, up to 1, 2, 3, 4, or 5 nucleotides between the miRNA region of the first strand and the complementary region of the second strand may not be complementary. In one embodiment, the mismatches are consecutive and may create a bulge. In another embodiment, the mismatches are not consecutive and may be distributed throughout the complementary region. In yet another embodiment, up to 1, 2, 3, or 4 mismatches may be consecutive creating a bulge and the remaining mismatches may be distributed through the complementary region.

[0041] In some embodiments, the first and/or the second strand of the mimetic compound may comprise an overhang on the 5' or 3' end of the strands. In certain embodiments, the first strand comprises a 3' overhang, *i.e.*, a single-stranded region that extends beyond the duplex region, relative to the second strand. In some embodiments, the second strand comprises a 3' overhang relative to the first strand. The 3' overhang of the first or the second strand may range from about one nucleotide to about four nucleotides. In certain embodiments, the 3' overhang of the first or the second strand may comprise 1 or 2

nucleotides. In some embodiments, the nucleotides comprising the 3' overhang are linked by phosphorothioate linkages. The nucleotides comprising the 3' overhang may include ribonucleotides, deoxyribonucleotides, modified nucleotides, or combinations thereof. In certain embodiments, the 3' overhang in the first or the second strand comprises two ribonucleotides. In other embodiments, the 3' overhang in the first or the second strand comprises two modified ribonucleotides. In still other embodiments, the 3' overhang in the first or the second strand comprises one ribonucleotide and one modified ribonucleotide. In some embodiments, the overhang may be present on the 5' end of the first or the second strand. The 5' overhang may comprise from about one to four nucleotides. Similar to the 3' overhang, the 5' overhang may comprise ribonucleotides, deoxyribonucleotides, modified nucleotides, or combinations thereof. In some embodiments, the nucleotides comprising the 5' overhang may be linked by phosphorothioate linkages. In some embodiments, the miRNA mimetic compound may be a hairpin, *i.e.*, a single strand polynucleotide with a 5' and a 3' end, where one of the ends may generate an overhang when the single strand folds back on itself. In these embodiments, the single strand comprises the miRNA region and the complementary region that may be separated by a linker region. Such a single strand miRNA mimetic compound may have a greater range of length, for example, about 55 to about 100 nucleotides. The single stranded miRNA mimetic compound may contain an unpaired loop that would substantially correspond to the linker region.

[0042] It will be understood from the above description that the first or the antisense strand of the microRNA mimetic compound may comprise the entire sequence of a mature, naturally occurring microRNA or a part of it and may comprise additional nucleotides that are not part of the mature miRNA sequence. For example, the first strand of a mimetic compound according to the invention that mimics the activity of miR-96 may comprise the entire sequence of SEQ ID NOs: 1, 2, or 3 or a partial sequence, such as about, 15, 16, 17, 18, 19, 20, 21, or 22 nucleotides of SEQ ID NOs: 1, 2, or 3 and up to about 4 to 6 additional nucleotides that are not part of the mature miR-96 sequence. It will also be understood that the mimetic compound comprising the entire or partial sequence of the mature, naturally occurring microRNA and up to about 4 to 6 additional nucleotides still maintains the ability to mimic the activity or function of the microRNA. It will also be understood that the sequence of the first strand comprising the sequence of a mature, naturally occurring

microRNA may include modified nucleotides corresponding to the nucleotides present in the mature, naturally-occurring microRNA.

[0043] In one embodiment, the invention provides a microRNA mimetic compound comprising a first strand of about 22 to about 26 ribonucleotides comprising a mature miR-96, miR-182, or miR-183 sequence; and a second strand of about 20 to about 24 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In some embodiments, the invention provides a microRNA mimetic compound comprising a first strand of about 22 to about 26 ribonucleotides comprising a mature miR-96, miR-182, or miR-183 sequence; and a second strand of about 20 to about 26 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the second strand has a 3' nucleotide overhang relative to the second strand. The term “substantially complementary” means the sequence of the second strand is at least about 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%, inclusive of all values therebetween, complementary to the sequence of the first strand or the sequence of the second strand contains up to about 1, 2, 3, 4, 5, or 6 mismatches relative to the sequence of the first strand.

[0044] In certain embodiments, a miR-96 mimetic compound according to the invention comprises a first strand of about 22 to about 26 nucleotides comprising a sequence of SEQ ID NOs: 1, 2, or 3, and a second strand of about 20 to about 24 nucleotides that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In some embodiments, the second strand may comprise about 20 to about 26 nucleotides and may include a 3' or 5' nucleotide overhang relative to the first strand. Throughout this disclosure, the language reciting “a sequence of SEQ ID NOs. X, Y, or Z” encompasses all sequences where a naturally occurring nucleotide is replaced by a corresponding modified nucleotide. For example, the language encompasses sequences where adenosine is replaced with a modified adenosine; uridine is replaced with a modified uridine, thymidine, or modified thymidine;

guanosine is replaced with modified guanosine; or cytidine is replaced with modified cytidine.

[0045] In certain embodiments, a miR-96 mimetic compound according to the invention comprises a first strand comprising the sequence:

5'-rUrUfUrGrGfCrAfCfUrArGfCrAfCrAfUfUfUfUrGfCfUsrUsrU-3' (SEQ ID NO: 10) and a second strand that is substantially complementary, such as about 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%, inclusive of all values therebetween, complementary to the first strand. In one embodiment, the second strand of a miR-96 mimetic compound comprises the sequence:

5'-mAmGmCrArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArA-3' (SEQ ID NO: 11). In another embodiment, the second strand of a miR-96 mimetic compound comprises the sequence: 5'-mAmGmCrArArArArAmUrGmUrGmCmUrArGmUrGmCmCrArArA-3' (SEQ ID NO: 12). As used herein, an "m" preceding a base notation (e.g. A, U, G, C) indicates a 2'-O-methyl modified nucleotide, an "r" preceding a base notation indicates an unmodified ribonucleotide (*i.e.*, 2'-OH), an "f" preceding a base notation indicates a 2'-fluoronucleotide, and an "s" indicates a phosphorothioate linkage. Unless otherwise indicated, the nucleotides in the antisense and the sense strand are linked by phosphodiester linkages. In some embodiments, a miR-96 mimetic compound according to the invention comprises a first strand comprising a sequence selected from SEQ ID NOs: 10 and 26-29. In some embodiments, a miR-96 mimetic compound according to the invention comprises a second strand comprising a sequence selected from SEQ ID NOs: 11-14 and 30-34.

[0046] In certain embodiments, a miR-96 mimetic compound according to the invention comprises a first strand of about 22 to about 26 ribonucleotides comprising a sequence of SEQ ID NOs: 1, 2, 3, or 10 and a second strand of about 20 to about 24 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In certain embodiments, the second strand of a miR-96 mimetic compound comprises a sequence of SEQ ID NO: 11 or 12. In some embodiments, a miR-96 mimetic compound according to the invention comprises a first strand of about 22 to about 26 ribonucleotides comprising a sequence of SEQ ID NOs: 1, 2, 3, or 10 and a second strand of

about 20 to about 26 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the second strand has a 3' or 5' nucleotide overhang relative to the first strand.

[0047] In some embodiments, a miR-96 mimetic compound comprises a first strand of about 20 to about 26 nucleotides comprising the sequence of SEQ ID NO: 10 and a second strand of about 20 to about 24 nucleotides comprising the sequence of SEQ ID NO: 11, wherein the first strand has a 3' overhang relative to the second strand. In another embodiment, a miR-96 mimetic compound comprises a first strand of about 20 to about 26 nucleotides comprising the sequence of SEQ ID NO: 10 and a second strand of about 20 to about 24 nucleotides comprising the sequence of SEQ ID NO: 12, wherein the first strand has a 3' overhang relative to the second strand. In yet another embodiment, a miR-96 mimetic compound comprises a first strand of about 20 to about 26 nucleotides comprising the sequence of SEQ ID NO: 10 and a second strand of about 20 to about 24 nucleotides comprising the sequence of SEQ ID NO: 13 (5'-mA.mG.mC.rA.rA.rA.mU.mU.rG.rA.rG.mC.mU.rA.rG.mU.rG.mC.rG.rA.rA.chol6-3'), wherein the first strand has a 3' overhang relative to the second strand. In some embodiments, the second strand of a miR-96 mimetic compound is about 20 to about 24 nucleotides and comprises the sequence of SEQ ID NO: 14 (5'-mA.mG.mC.rA.rA.rA.mU.mU.rG.rA.rG.mC.mU.rA.rG.mU.rG.mC.rG.rA.rAs.chol6-3'). In this embodiment, the oligonucleotide sequence of the second strand is attached to cholesterol (carrier molecule) via a phosphorothioate linkage.

[0048] In some embodiments, a miR-96 mimetic compound comprises a first strand that is no more than 25, 26, 27 or 28 nucleotides long and comprises a sequence of SEQ ID NOs: 1, 2, 3, or 10. In some other embodiments, a miR-96 mimetic compound comprises a first strand that is no more than 26, 27 or 28 nucleotides long and comprises a sequence of SEQ ID NOs: 1, 2, or 3, where nucleotides at positions 3(U), 6(C), 8(C), 9(U), 12 (C), 14 (C), 16 (U), 17 (U), 18 (U), 19 (U), 20 (U), 22 (C), and/or 23 (U) in the 5' to 3' direction are modified relative to SEQ ID NOs. 1, 2, or 3. In certain embodiments, a miR-96 mimetic compound comprises a first strand having a sequence of SEQ ID NOs: 1, 2, 3, or 10, and a second strand that is complementary to the first strand except at nucleotide positions 4, 13,

and/or 16 from the 3' end of the second strand. In some embodiments, the second strand of a miR-96 mimetic compound is complementary to the sequence of SEQ ID NOs: 1, 2, 3, or 10 and comprises modified nucleotides at one or more positions selected from the group consisting of 1(A), 2(G), 3(C), 8(U), 9(U), 13(C), 14(U), 17(U), and 19(C), in the 5' to 3' direction.

[0049] In certain embodiments, a miR-182 mimetic compound according to the invention comprises a first strand of about 22 to about 26 nucleotides comprising a sequence of SEQ ID NOs: 4, 5, or 6, and a second strand that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In some embodiments, the second strand may comprise about 20 to about 26 nucleotides and may include a 3' or 5' nucleotide overhang relative to the first strand. In certain embodiments, a miR-182 mimetic compound according to the invention comprises a first strand comprising the sequence:

5'-rUrUfUrGrGfCrArAfUrGrGfUrArGrArAfCfUfCrAfCrAfCfUsrUsrU-3' (SEQ ID NO: 15) and a second strand that is substantially complementary, such as about 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%, inclusive of all values therebetween, complementary to the first strand. In one embodiment, the second strand of a miR-182 mimetic compound comprises the sequence:

5'-mAmGmUrGmUrGrArGrAmUmCrArAmCmCrAmUmUrGmCrGrArArA-3' (SEQ ID NO: 16). In another embodiment, the second strand of a miR-182 mimetic compound comprises the sequence: 5'-mAmGmUrGmUrGrArGmUmUmCmUrAmCmCrAmUmUrGmCmCrArArA-3' (SEQ ID NO: 17). In some embodiments, a miR-182 mimetic compound according to the invention comprises a first strand comprising a sequence selected from SEQ ID NOs: 15 and 35-38. In some embodiments, a miR-182 mimetic compound according to the invention comprises a second strand comprising a sequence selected from SEQ ID NOs: 16-19 and 39-43.

[0050] In some embodiments, a miR-182 mimetic compound comprises a first strand of about 22 to about 26 ribonucleotides containing a sequence of SEQ ID NOs: 4, 5, 6, or 15 and a second strand of about 20 to about 24 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide,

wherein the first strand has a 3' nucleotide overhang relative to the second strand. In certain embodiments, the second strand of a miR-182 mimetic compound comprises a sequence of SEQ ID NO: 16 or 17. In some embodiments, a miR-182 mimetic compound according to the invention comprises a first strand of about 22 to about 26 ribonucleotides comprising a sequence of SEQ ID NOs: 4, 5, 6, or 15 and a second strand of about 20 to about 26 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the second strand has a 3' or 5' nucleotide overhang relative to the first strand.

[0051] In some embodiments, a miR-182 mimetic compound comprises a first strand of about 22 to about 26 nucleotides comprising the sequence of SEQ ID NO: 15 and a second strand of about 20 to about 24 nucleotides comprising the sequence of SEQ ID NO: 16, wherein the first strand has a 3' overhang relative to the second strand. In another embodiment, a miR-182 mimetic compound comprises a first strand of about 22 to about 26 nucleotides comprising the sequence of SEQ ID NO: 15 and a second strand of about 20 to about 24 nucleotides comprising the sequence of SEQ ID NO: 17, wherein the first strand has a 3' overhang relative to the second strand. In yet another embodiment, a miR-182 mimetic compound comprises a first strand of about 22 to about 26 nucleotides comprising the sequence of SEQ ID NO: 15 and a second strand of about 20 to about 24 nucleotides comprising the sequence of SEQ ID NO: 18 (5'-mA.mG.mU.rG.mU.rG.rA.rG.rA.mU.mC.rA.rA.mC.mC.rA.mU.mU.rG.mC.rG.rA.rA.cho16-3'), wherein the first strand has a 3' overhang relative to the second strand. In some embodiments, the second strand of a miR-182 mimetic compound is about 20 to about 24 nucleotides and comprises the sequence of SEQ ID NO: 19 (5'-mA.mG.mU.rG.mU.rG.rA.rG.rA.mU.mC.rA.rA.mC.mC.rA.mU.mU.rG.mC.rG.rA.rA.rAs.cho16-3'). In this embodiment, the oligonucleotide sequence of the second strand is attached to cholesterol (carrier molecule) via a phosphorothioate linkage.

[0052] In some embodiments, a miR-182 mimetic compound comprises a first strand that is no more than 26, 27 or 28 nucleotides long and comprises a sequence of SEQ ID NOs: 4, 5, 6, or 15. In some other embodiments, a miR-182 mimetic compound comprises a first strand that is no more than 27 or 28 nucleotides long and comprises a sequence of SEQ ID NOs: 4,

5, or 6, where nucleotides at positions 3(U), 6(C), 9(U), 12 (U), 17 (C), 18 (U), 19 (C), 21 (C), 23 (C), and/or 24 (U) in the 5' to 3' direction are modified relative to SEQ ID NOs. 4, 5, or 6. In certain embodiments, a miR-182 mimetic compound comprises a first strand having a sequence of SEQ ID NOs: 4, 5, 6, or 15, and a second strand that is complementary to the first strand except at nucleotide positions 4, 13, and/or 16 from the 3' end of the second strand. In some embodiments, the second strand of a miR-182 mimetic compound is complementary to the sequence of SEQ ID NOs: 4, 5, 6, or 15 and comprises modified nucleotides at one or more positions selected from the group consisting of 1(A), 2 (G), 3(U), 5(U), 10(U), 11(C), 14(C), 15(C), 17(U), 18(U), and 20(C), in the 5' to 3' direction.

[0053] In certain embodiments, a miR-183 mimetic compound according to the invention comprises a first strand of about 22 to about 26 nucleotides comprising a sequence of SEQ ID NOs: 7, 8, or 9, and a second strand that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In some embodiments, the second strand may comprise about 20 to about 26 nucleotides and may include a 3' or 5' nucleotide overhang relative to the first strand. In certain embodiments, a miR-183 mimetic compound according to the invention comprises a first strand comprising the sequence:

5'- rUrAfUrGrGfCrAfCfUrGrGfUrArGrArAfUfUfCrAfCfUsrUsrU-3' (SEQ ID NO: 20) and a second strand that is substantially complementary, such as about 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100%, inclusive of all values therebetween, complementary to the first strand. In one embodiment, the second strand of a miR-183 mimetic compound comprises the sequence:

5'- mAmGmUrGrArArAmUmCrArAmCmCrArGmUrGmCrGrAmUrA-3' (SEQ ID NO: 21). In another embodiment, the second strand of a miR-183 mimetic compound comprises the sequence: 5'-mAmGmUrGrArArAmUmUmCmUrAmCmCrArGmUrGmCmCrAmUrA-3' (SEQ ID NO: 22). In some embodiments, a miR-183 mimetic compound according to the invention comprises a first strand comprising a sequence selected from SEQ ID NOs: 20 and 44-47. In some embodiments, a miR-183 mimetic compound according to the invention comprises a second strand comprising a sequence selected from SEQ ID NOs: 21-25 and 48-52.

[0054] In some embodiments, a miR-183 mimetic compound comprises a first strand of about 22 to about 26 ribonucleotides comprising a sequence of SEQ ID NOs: 7, 8, 9, or 20 and a second strand of about 20 to about 24 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the first strand has a 3' nucleotide overhang relative to the second strand. In certain embodiments, the second strand of a miR-183 mimetic compound comprises a sequence of SEQ ID NO: 21 or 22. In some embodiments, a miR-183 mimetic compound according to the invention comprises a first strand of about 22 to about 26 ribonucleotides comprising a sequence of SEQ ID NOs: 7, 8, 9, or 20 and a second strand of about 20 to about 26 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide, wherein the second strand has a 3' or 5' nucleotide overhang relative to the first strand.

[0055] In some embodiments, a miR-183 mimetic compound comprises a first strand of about 22 to about 26 nucleotides comprising the sequence of SEQ ID NO: 20 and a second strand of about 18 to about 22 nucleotides comprising the sequence of SEQ ID NO: 21, wherein the first strand has a 3' overhang relative to the second strand. In another embodiment, a miR-183 mimetic compound comprises a first strand of about 22 to about 26 nucleotides comprising the sequence of SEQ ID NO: 20 and a second strand of about 18 to about 22 nucleotides comprising the sequence of SEQ ID NO: 22, wherein the first strand has a 3' overhang relative to the second strand. In yet another embodiment, a miR-183 mimetic compound comprises a first strand of about 22 to about 26 nucleotides comprising the sequence of SEQ ID NO: 20 and a second strand of about 18 to about 22 nucleotides comprising the sequence of SEQ ID NO: 23 (5'-mAmGmUrGrArArAmUmCrArAmCmCrArGmUrGmCrGrAmUrA.chol6-3') or SEQ ID NO: 24 (5'-mAmGmUrGrArArAmUmUmCmUrAmCmCrArGmUrGmCmCrAmUrA.chol6-3'), wherein the first strand has a 3' overhang relative to the second strand. In one embodiment, the second strand of a miR-183 mimetic compound comprises the sequence of SEQ ID NO: 25 (5'-mAmGmUrGrArArAmUmCrArAmCmCrArGmUrGmCrGrAmUrAs.chol6-3'). In this embodiment, the oligonucleotide sequence of the second strand is attached to cholesterol (carrier molecule) via a phosphorothioate linkage.

[0056] In some embodiments, a miR-183 mimetic compound comprises a first strand that is no more than 25, 26, 27 or 28 nucleotides long and comprises a sequence of SEQ ID NOs: 7, 8, 9, or 20. In some other embodiments, a miR-183 mimetic compound comprises a first strand that is no more than 25, 26, 27 or 28 nucleotides long and comprises a sequence of SEQ ID NOs: 7, 8, or 9, where nucleotides at positions 3(U), 6(C), 8(C), 9(U), 12(U), 17(U), 18(U), 19(C), 21(C), and/or 22(U) in the 5' to 3' direction are modified relative to SEQ ID NOs. 7, 8, or 9. In certain embodiments, a miR-183 mimetic compound comprises a first strand having a sequence of SEQ ID NOs: 7, 8, 9, or 20, and a second strand that is complementary to the first strand except at nucleotide positions 4, 13, and/or 16 from the 3' end of the second strand. In some embodiments, the second strand of a miR-183 mimetic compound is complementary to the sequence of SEQ ID NOs: 7, 8, 9, or 20 and comprises modified nucleotides at one or more positions selected from the group consisting of 1(A), 2(G), 3(U), 7(U), 8(U), 9(C), 10(U), 12(C), 13(C), 16(U), 18(C), 19(C), and 21(U), in the 5' to 3' direction.

[0057] Specific microRNA mimetic compounds disclosed herein are summarized in Table 1 below. However, the invention is not limited to these specific mimetic compounds and other mimetic compounds that could be prepared based on the guidance provided throughout the specification are also encompassed by the invention.

Table 1: miR-96/182/183 mimics

SEQ ID NO.	Modified Sequence
<i>miR-96 First/antisense/guide strands</i>	
10	5'-rUrUfUrGrGfCrAfCfUrArGfCrAfCrAfUfUfUfUrGfCfUsrUsrU-sup-3'
26	5'-rUrUfUrGrGfCrAfCfUrArGfCrAfCrAfUfUfUfUrGfCfU-3'
27	5'-p.rUrUfUrGrGfCrAfCfUrArGfCrAfCrAfUfUfUfUrGfCfUsrUsrU-3'
28	5'-rUrUrUrGrGrCrArCrUrArGrCrArCrUrUrUrUrGrCrUrUrU-3'
29	5'-rUrUmUrGrGmCrAmCmUrArGmCrAmCmUmUmUmUmUrGmCmUsrUsrU-3'

<i>miR-96 Second/sense/pasenger strands</i>	
11	5'-mAmGmCrArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArA-3'
12	5'-mAmGmCrArArArArAmUrGmUrGmCmUrArGmUrGmCmCrArArA-3'
13	5'-mAmGmCrArArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArAcho6-3'
14	5'-mAmGmCrArArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArAscho6-3'
30	5'-mAmGmCrArArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArAsrU-3'
31	5'-C6Chol.mAmGmCrArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArA-3'
32	5'-mAmGrCrArArArArArUrGrUrGrCrUrArGrUrGrCrCrArArUrU-3'
33	5'-mAmGmCrArArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArAscho6-3'
34	5'-mAmGmCrArArArArAmUmUrGrArGmCmUrArGmUrGmCrGrArArAcho6-3'
<i>miR-182 First/antisense/guide strands</i>	
15	5'-rU.rU.flU.rG.rG.flC.rA.rA.flU.rG.rG.flU.rA.rG.rA.flC.flU.flC.rA.flC.flUs.rUs.rU-sup-3'
35	5'-rUrUrUrGrGfCrArAfUrGrGfUrArGrArAfCfUfCrAfCrAfCfU-3'
36	5'-p.rUrUrUrGrGfCrArAfUrGrGfUrArGrArAfCfUfCrAfCrAfCfUsrUsrU-3'
37	5'-rUrUrUrGrGrCrArArUrGrGrUrArGrArArCrUrCrArCrArCrUrUrU-3'
38	5'-rUrUmUrGrGmCrArAmUrGrGmUrArGrArAmCmUmCrAmCmUsrUsrU-3'
<i>miR-182 Second/sense/pasenger strands</i>	
16	5'-mA.mG.mU.rG.mU.rG.rA.rG.rA.mU.mC.rA.rA.mC.mC.rA.mU.mU.rG.mC.rG.rA.rA.rA-3'
17	5'-mA.mG.mU.rG.mU.rG.rA.rG.mU.mU.mC.mU.rA.mC.mC.rA.mU.mU.rG.mC.mC.rA.rA.rA-3'
18	5'-mA.mG.mU.rG.mU.rG.rA.rG.rA.mU.mC.rA.rA.mC.mC.rA.mU.mU.rG.mC.rG.rA.rA.rA.cho6-3'
19	5'-mA.mG.mU.rG.mU.rG.rA.rG.rA.mU.mC.rA.rA.mC.mC.rA.mU.mU.rG.mC.rG.rA.rA.rAs.cho6-3'
39	5'-mAmGmUrGmUrGrArGrAmUmCrArAmCmCrAmUmUrGmCrGrArArAsrU-3'
40	5'-C6cho1.mAmGmUrGmUrGrArGrAmUmCrArAmCmCrAmUmUrGmCrGrArArA-3'
41	5'-mAmGrUrGrUrGrArGrUrUrCrUrArCrArUrUrGrCrCrArArUrU-3'
42	5'-mAmGmUrGmUrGrArGrAmUmCrArAmCmCrAmUmUrGmCrGrArArAscho6-3'
43	5'-mAmGmUrGmUrGrArGrAmUmCrArAmCmCrAmUmUrGmCrGrArArAcho6-3'

<i>miR-183 First/antisense/guide strands</i>	
20	5'-rU.rA.fU.rG.rG.fC.rA.fC.fU.rG.rG.fU.rA.rG.rA.fU.fU.fC.rA.fC.fU.rU-sup-3'
44	5'-rUrAUrGrGfCrAfCfUrGrGfUrArGrArAfUfUfCrAfCfU-3'
45	5'-p.rUrAUrGrGfCrAfCfUrGrGfJrArGrArAfUfUfCrAfCfUsrUsrU-3'
46	5'-rUrArUrGrGrCrArCrUrGrGrUrArGrArUrCrArCrUrUrU-3'
47	5'-rUrAmUrGrGmCrAmCmUrGrGmUrArGrArAmUmUmCrAmCmUsrUsrU-3'
<i>miR-183 Second/sense/passenger strands</i>	
21	5'- mA.mG.mU.rG.rA.rA.mU.mC.rA.rA.mC.mC.rA.rG.mU.rG.mC.rG.rA.mU.rA-3'
22	5'-mA.mG.mU.rG.rA.rA.mU.mU.mC.mU.rA.mC.mC.rA.rG.mU.rG.mC.mC.rA.mU.rA-3'
23	5'- mA.mG.mU.rG.rA.rA.rA.mU.mC.rA.rA.mC.mC.rA.rG.mU.rG.mC.rG.rA.mU.rA.cho16-3'
24	5'-mA.mG.mU.rG.rA.rA.mU.mU.mC.mU.rA.mC.mC.rA.rG.mU.rG.mC.mC.rA.mU.rA.cho16-3'
25	5'- mA.mG.mU.rG.rA.rA.rA.mU.mC.rA.rA.mC.mC.rA.rG.mU.rG.mC.rG.rA.mU.rAs.cho16-3'
48	5'-mAmGmUrGrArAmUmUmCmUrAmCmCrArGmUrGmCmCrAmUrAsrUsrU-3'
49	5'-C6cho1.mAmGmUrGrArArAmUmCrArAmCmCrArGmUrGmCrGrAmUrA-3'
50	5'-mAmGrUrGrArArUrUrCrUrArCrCrArGrUrGrCrArUrArUrU-3'
51	5'-mAmGmUrGrArArArAmUmCrArAmCmCrArGmUrGmCrGrAmUrAChol6-3'
52	5'-mAmGmUrGrArArArAmUmCrArAmCmCrArGmUrGmCrGrAmUrAsChol6-3'

[0058] The modified nucleotides that may be used in the microRNA mimetic compounds of the invention can include nucleotides with a base modification or substitution. The natural or unmodified bases in RNA are the purine bases adenine (A) and guanine (G), and the pyrimidine bases cytosine (C) and uracil (U) (DNA has thymine (T)). In contrast, modified bases, also referred to as heterocyclic base moieties, include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo (including 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines), 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine,

8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine.

[0059] In some embodiments, the microRNA mimetic compounds can have nucleotides with modified sugar moieties. Representative modified sugars include carbocyclic or acyclic sugars, sugars having substituent groups at one or more of their 2', 3' or 4' positions and sugars having substituents in place of one or more hydrogen atoms of the sugar. In certain embodiments, the sugar is modified by having a substituent group at the 2' position. In additional embodiments, the sugar is modified by having a substituent group at the 3' position. In other embodiments, the sugar is modified by having a substituent group at the 4' position. It is also contemplated that a sugar may have a modification at more than one of those positions, or that an RNA molecule may have one or more nucleotides with a sugar modification at one position and also one or more nucleotides with a sugar modification at a different position.

[0060] Sugar modifications contemplated in the miRNA mimetic compounds include, but are not limited to, a substituent group selected from: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. In some embodiments, these groups may be chosen from: O(CH₂)_xOCH₃, O((CH₂)_xO)_yCH₃, O(CH₂)_xNH₂, O(CH₂)_xCH₃, O(CH₂)_xONH₂, and O(CH₂)_xON((CH₂)_xCH₃)₂, where x and y are from 1 to 10.

[0061] In some embodiments, miRNA mimetic compounds have a sugar substituent group selected from the following: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, Cl, Br, CN, OCN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, or similar substituents. In one embodiment, the modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, which is also known as 2'-O-(2-methoxyethyl) or 2'-MOE), that is, an alkoxyalkoxy group. Another modification includes 2'-dimethylaminoxyethoxy, that is, a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE and 2'-dimethylaminoethoxyethoxy

(also known in the art as 2'-O-dimethyl-amino-ethoxy-ethyl or 2'-DMAEOE), that is, 2'-O-CH₂-O-CH₂-N(CH₃)₂.

[0062] Additional sugar substituent groups include allyl (-CH₂-CH=CH₂), -O-allyl, methoxy (-O-CH₃), aminopropoxy (-OCH₂CH₂CH₂NH₂), and fluoro (F). Sugar substituent groups on the 2' position (2') may be in the arabino (up) position or ribo (down) position. One 2'-arabino modification is 2'-F. Other similar modifications may also be made at other positions on the sugar moiety, particularly the 3' position of the sugar on the 3' terminal nucleoside or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide.

[0063] In certain embodiments, the sugar modification is a 2'-O-alkyl (e.g. 2'-O-methyl, 2'-O-methoxyethyl), 2'-halo (e.g., 2'-fluoro, 2'-chloro, 2'-bromo), and 4' thio modifications. For instance, in some embodiments, the first strand of the miR-96, miR-182, or miR-183 mimetic compound comprises one or more 2' fluoro nucleotides. In another embodiment, the first strand of the mimetic compounds has no modified nucleotides. In yet another embodiment, the second strand of miR-96, miR-182, or miR-183 mimetic compound comprises at least one 2'-O-methyl modified nucleotide.

[0064] The first and the second strand of microRNA mimetic compounds of the invention can also include backbone modifications, such as one or more phosphorothioate, morpholino, or phosphonocarboxylate linkages (see, for example, U.S. Patent Nos. 6,693,187 and 7,067,641, which are herein incorporated by reference in their entireties). For example, in some embodiments, the nucleotides comprising the 3' overhang in the first strand are linked by phosphorothioate linkages.

[0065] In some embodiments, the microRNA mimetic compounds are conjugated to a carrier molecule such as a steroid (cholesterol), a vitamin, a fatty acid, a carbohydrate or glycoside, a peptide, or other small molecule ligand to facilitate *in vivo* delivery and stability. Preferably, the carrier molecule is attached to the second strand of the microRNA mimetic compound at its 3' or 5' end through a linker or a spacer group. In various embodiments, the carrier molecule is cholesterol, a cholesterol derivative, cholic acid or a cholic acid derivative. The use of carrier molecules disclosed in U.S. Patent No. 7,202,227, which is

incorporated by reference herein in its entirety, is also envisioned. In certain embodiments, the carrier molecule is cholesterol and it is attached to the 3' or 5' end of the second strand through at least a six carbon linker. In one embodiment, the carrier molecule is attached to the 3' end of the second strand through a linker. In various embodiments, the linker comprises a substantially linear hydrocarbon moiety. The hydrocarbon moiety may comprise from about 3 to about 15 carbon atoms and may be conjugated to cholesterol through a relatively non-polar group such as an ether or a thioether linkage. In certain embodiments, the hydrocarbon linker/spacer comprises an optionally substituted C2 to C15 saturated or unsaturated hydrocarbon chain (e.g. alkylene or alkenylene). A variety of linker/spacer groups described in U.S. Pre-grant Publication No. 2012/0128761, which is incorporated by reference herein in its entirety, can be used in the present invention.

Expression vectors encoding miR-96, miR-182, and/or miR-183

[0066] An expression vector comprising at least one gene encoding miR-96, miR-182, and/or miR-183 includes a sufficient portion of the miR-96, miR-182, and/or miR-183 native coding sequence, with or without flanking sequences present in the genomic context of miR-96, miR-182, and/or miR-183, to produce a mature miR-96, miR-182, and/or miR-183 to regulate expression of at least one miR-96, miR-182, and/or miR-183 target. For example, a sufficient portion can include about 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 nucleotides, including values and ranges thereof. In one embodiment, a sufficient portion contains at least the sequence of the mature miR. In certain embodiments, a sufficient portion includes at least the hairpin sequence of the miR. In certain embodiments, a sufficient portion includes the full length miR. A sufficient portion can be determined using methods routine in the art. It is understood that a sequence encoding a miR-96, miR-182, and/or miR-183 will be complementary to the RNA sequences provided and include T's rather than U's when the complementary DNA strand.

[0067] By "vector" is meant a nucleic acid molecule, for example, a plasmid, cosmid, or bacteriophage, that is capable of replication in a host cell. In one embodiment, a vector is an expression vector that is a nucleic acid construct, generated recombinantly or synthetically, bearing a series of specified nucleic acid elements that enable transcription of a nucleic acid molecule in a host cell. Typically, expression is placed under the control of certain regulatory

elements, including constitutive or inducible promoters, tissue-preferred regulatory elements, inverted terminal repeats, and enhancers.

[0068] In one embodiment, the expression vector is an AAV-based vector system which is an especially attractive platform for regulatory RNA delivery (Franich et al., Mol Ther 16, 947-956, 2008 and McCarty, Mol Ther 16, 1648-1656, 2008). When delivered in viral vectors, miRNAs are continually transcribed, allowing sustained high level expression in target tissues without the need for repeated dosing. Additionally, the use of tissue-specific promoters could restrict this expression to particular cell types of interest even with systemic delivery of the virus. Compared to retroviral delivery systems, DNA viruses such as AAV carry substantially diminished risk of insertional mutagenesis since viral genomes persist primarily as episomes (Schnepp et al., J Virol 77, 3495-3504, 2003). Further, the availability of multiple AAV serotypes allows efficient targeting of many tissues of interest (Gao et al., Proc Natl Acad Sci USA 99, 11854-11859, 2002; McCarty, Mol Ther 16, 1648-1656, 2008). Finally, the general safety of AAV has been well documented, with clinical trials using this platform already underway (Carter, Hum Gene Ther 16, 541-550, 2005; Maguire et al., N Engl J Med 358, 2240-2248, 2008; and Park et al., Front Biosci 13, 2653-2659, 2008). Recent advances in AAV vector technology include a self-complementary genome which enhances therapeutic gene expression and non-human primate AAV serotypes which facilitate efficient transduction following delivery. Due to their small size, regulatory RNAs are especially amenable to AAV-mediated delivery.

[0069] The expression vectors provided by the instant invention can include any sequence that encodes a functional miR-96, miR-182, and/or miR-183 for use in any of the methods of the instant invention. In some embodiments, the expression vector includes a nucleic acid sequence encoding partial or the entire sequence of pre-miRNA hairpin. In certain embodiments, the expression vector includes a nucleic acid sequence selected from SEQ ID NOS. 53-55 (Table 2). The nucleic acid sequence encoding a functional miR-96, miR-182, and/or miR-183 could be present as a single cluster or as two or three separate clusters.

Table 2

miR DNA Sequence	Sequence
miR-182	5'- GAGCTGCTGCCTCCCCCGTTTGGCAATGGTAGAACTCACACTGGT GAGGTAAACAGGATCCGGTGGTTAGACTTGCCAACTATGGGCGAGG ACTCAGCCGGCAC-3' (SEQ ID NO: 53)
miR-183	5'- CCGCAGAGTGTGACTCCTGTTCTGTATGGCACTGGTAGAATTCACTG TGAACAGTCTCAGTCAGTGAATTACCGAAGGGCCATAAACAGAGCAGA GACAGATCCACGA-3' (SEQ ID NO: 54)
miR-96	5'- TGGCCGATTTGGCACTAGCACATTGGCTGTCTCCGCTTGAG CAATCATGTGCAGTGCCAATATGGGAAA-3' (SEQ ID NO: 55)

[0070] The invention provides polynucleotide therapy useful for increasing the expression of miR-96, miR-182, or miR-183 microRNA, or any combination thereof for the treatment of ophthalmological and ear diseases. Expression vectors encoding a desired sequence (*e.g.* encoding a microRNA) can be delivered to cells of a subject having an ophthalmological or ear disease. The nucleic acid molecules must be delivered to the cells of a subject in a form in which they can be taken up and are advantageously expressed so that therapeutically effective levels can be achieved.

[0071] Methods for delivery of the polynucleotides to the cell according to the invention include using a delivery system such as liposomes, polymers, microspheres, gene therapy vectors, and naked DNA vectors.

[0072] Transducing viral (*e.g.*, retroviral, adenoviral, lentiviral and adeno-associated viral) vectors can be used for somatic cell gene therapy, especially because of their high efficiency of infection and stable integration and expression. For example, a polynucleotide encoding a nucleic acid molecule can be cloned into a retroviral vector and expression can be driven from its endogenous promoter, from the retroviral long terminal repeat, or from a promoter specific for a target cell type of interest. Other viral vectors that can be used include, for example, a vaccinia virus, a bovine papilloma virus, or a herpes virus, such as Epstein-Ban Virus. Retroviral vectors are particularly well developed and have been used in clinical settings (U.S. Pat. No.5,399,346). Viral vectors are preferably replication incompetent in the

cells to which they are delivered for therapeutic applications. However, replication competent viral vectors may also be used.

[0073] Preferred viral vectors for use in the invention include AAV vectors, *e.g.* AAV serotypes 1, 2, 3, 4, 5, 6, 7, 8, and/or 9, including chimeric AAV vectors. The availability of multiple AAV serotypes allows efficient targeting of many tissues of interest (Gao et al., 2002; McCarty, 2008; US Patent Publications 2008075737, 20080050343, 20070036760, 20050014262, 20040052764, 20030228282, 20030013189, 20030032613, and 20020019050 each incorporated herein by reference). In preferred embodiments, the invention includes the use of self-complementary (sc) AAV vectors which are described, for example, in US Patent Publications 20070110724 and 20040029106, and U.S. Pat. Nos. 7,465,583 and 7,186,699 (all of which are incorporated herein by reference). Exemplary methods for preparing AAVs for expressing microRNA are described in Knabel et al., PLoS One, 10(4):e0124411, 2015 and Xie et al., Semin Liver Dis, 35(1): 81-88, 2015 (both of which are incorporated herein by reference).

[0074] Non-viral approaches can also be employed for the introduction of a therapeutic nucleic acid molecule to a cell of a patient having an ophthalmological or ear disease. For example, an expression vector that encodes a miR-96, miR-182 and/or miR-183 microRNA can be introduced into a cell by administering the nucleic acid in the presence of lipofection, calcium phosphate co-precipitation, electroporation, microinjection, DEAE-dextran, transfection employing polyamine transfection reagents, cell sonication, gene bombardment using high velocity microprojectiles, and receptor-mediated transfection.

[0075] Nucleic acid molecule expression for use in polynucleotide therapy methods can be directed from any suitable promoter (*e.g.*, the human cytomegalovirus (CMV), simian virus 40 (SV40), metallothionein, U1a1 snRNA, U1b2 snRNA, histone H2, and histone H3 promoters), and regulated by any appropriate mammalian regulatory element. The expression may be directed using a tissue-specific or ubiquitously expressed promoter. Tissue specific promoters useful for the treatment of ophthalmological diseases include, but are not limited to, rhodopsin promoter, calcium binding protein 5 (CABP5) promoter, and cellular retinaldehyde binding protein (CRALBP) promoter. If desired, enhancers known to preferentially direct gene expression in specific cell types can be used to direct the expression

of a nucleic acid. The enhancers used can include, without limitation, those that are characterized as tissue- or cell-specific enhancers.

[0076] In one embodiment, an expression vector for expressing an agonist of miR-96, miR-182, or miR-183 comprises a promoter and a terminator operably linked to a polynucleotide sequence encoding an agonist, wherein the expressed agonist comprises a first strand comprising a mature sequence of miR-96 (SEQ ID NOs: 1, 2, or 3), miR-182 (SEQ ID NOs: 4, 5, or 6), or miR-183 (SEQ ID NOs: 7, 8, or 9) and a second strand that is substantially complementary to the first strand. In another embodiment, an expression vector for expressing an agonist of miR-96, miR-182, or miR-183 comprises a promoter and a terminator operably linked to a polynucleotide sequence encoding a pre-miRNA sequence, wherein the expressed agonist comprises a polynucleotide sequence in the form of a hairpin comprising a mature miRNA sequence. In yet another embodiment, an expression vector for expressing an agonist of miR-96, miR-182, or miR-183 comprises a first promoter and a first terminator operably linked to a first polynucleotide sequence encoding an antisense strand of miR-96, miR-182, or miR-183 mimetic compound and a second promoter and a second terminator operably linked to a second polynucleotide sequence encoding a sense strand of miR-96, miR-182, or miR-183 mimetic compound. The phrase 'operably linked' or 'under transcriptional control' as used herein means that the promoter and the terminator are in the correct location and orientation in relation to a polynucleotide to control the initiation and termination of transcription by RNA polymerase and expression of the polynucleotide.

[0077] As used herein, a "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. Suitable promoters include, but are not limited to RNA pol I, pol II, pol III, and viral promoters (e.g. human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter, and the Rous sarcoma virus long terminal repeat). In one embodiment, the promoter is a tissue-specific promoter. Of particular interest are retinal cell specific promoters, and more particularly, rods and cones specific promoters. These include the rhodopsin promoter, cone opsin promoter, calcium binding protein 5 (CABP5) promoter, cellular retinaldehyde binding protein (CRALBP) promoter, interphotoreceptor retinoid-binding protein (IRBP) promoter, arrestin promoter, and rhodopsin kinase promoter. As used

herein, a “terminator” refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, that is required to terminate the specific transcription of a gene. In one embodiment, a sequence comprising a polyadenylation signal/unit acts as a transcription terminator. The use of other transcription terminators is also contemplated.

[0078] In certain embodiments, the promoter operably linked to a polynucleotide encoding an agonist of miR-96, miR-182, or miR-183 can be an inducible promoter. Inducible promoters are known in the art and include, but are not limited to, tetracycline promoter, metallothionein II A promoter, heat shock promoter, steroid/thyroid hormone/retinoic acid response elements, the adenovirus late promoter, and the inducible mouse mammary tumor virus LTR.

Treatment methods

[0079] In various embodiments, the present invention provides methods of treating or preventing ophthalmological conditions in a subject in need thereof comprising administering to the subject a therapeutically effective amount of at least one agonist of miR-96, miR-182, and/or miR-183. In some embodiments, methods of treating or preventing ophthalmological conditions in a subject in need thereof comprise administering to the subject therapeutically effective amounts of at least two agonists, for example, agonists of miR-96 and miR-182, agonists of miR-96 and miR-183, or agonists of miR-182 and miR-183. In other embodiments, methods of treating or preventing ophthalmological conditions in a subject in need thereof comprise administering to the subject therapeutically effective amounts of all three agonists, *i.e.*, agonists of miR-96, miR-182, and miR-183. In some embodiments, the agonist of miR-96, miR-182, or miR-183 is a double-stranded oligonucleotide comprising a first strand containing a sequence of mature miR-96, miR-182, or miR-183 and a second strand comprising a sequence that is substantially complementary to the first strand, wherein at least one of the strands comprises one or more modified nucleotides. Any of the microRNA mimetic compounds described herein can be used in the methods of treating or preventing an ophthalmological condition in a subject in need thereof. In some embodiments, the agonist of miR-96, miR-182, or miR-183 is an expression vector comprising a polynucleotide sequence encoding a sufficient portion of the miR-96, miR-182,

or miR-183 native coding sequence to produce a mature miR-96, miR-182, or miR-183. In one embodiment, the expression vector is a recombinant AAV vector.

[0080] A “therapeutically effective amount or dose” is an amount sufficient to effect a beneficial or desired clinical result. For example, a therapeutically effective amount of microRNA mimetic compounds of miR-96, miR-182, and/or miR-183 includes an amount that is sufficient to maintain or improve visual acuity or an amount that is sufficient to reduce or prevent loss of vision or an amount that is sufficient to reduce or prevent photoreceptor cell death. In another embodiment, a therapeutically effective amount is an amount sufficient to increase expression of one or more phototransduction genes in the photoreceptor cells of the subject.

[0081] Ophthalmological conditions that may be treated by administering one or more agonists of miR-96, miR-182, and miR-183, include any condition caused by damage and/or death of photoreceptor cells. The term “photoreceptor cell” includes rods, cones, ganglion cells as well as other cells (*e.g.* retinal cells) present in the eye. In certain embodiments, ophthalmological conditions caused by damage or death of photoreceptor cells include retinal detachment, macular degeneration, Stargardt disease, retinal degeneration, retinitis pigmentosa, night blindness, retinal toxicity, uveal melanoma, sympathetic ophthalmia and retinopathies. In certain embodiments, the ophthalmological condition that may be treated according to the invention is retinal degeneration. In one embodiment, the ophthalmological condition that may be treated according to the invention is retinitis pigmentosa. In another embodiment, a subject in need of treatment with an agonist of miR-96, miR-182, and/or miR-183 may have signs of night blindness.

[0082] In various embodiments, administration of at least one agonist of miR-96, miR-182, or miR-183 to the subject prevents or slows the development and/or progression of one or more ophthalmological conditions and results in the improvement of one or more symptoms associated with these conditions. For instance, in one embodiment, administration of at least one agonist of miR-96, miR-182, or miR-183 results in the improvement of visual acuity. In another embodiment, administration of at least one agonist of miR-96, miR-182, or miR-183 reduces the signs of night blindness. In yet another embodiment, administration of at least

two agonists, for example, agonists of miR-96 and miR-182, agonists of miR-96 and miR-183, or agonists of miR-182 and miR-183, results in the improvement of visual acuity. In still another embodiment, administration of all three agonists, *i.e.*, agonists of miR-96, miR-182, and miR-183, results in the improvement of visual acuity.

[0083] In certain embodiments, the present invention provides methods for ameliorating or restoring visual acuity in a subject in need thereof comprising administering to the subject at least one agonist of miR-96, miR-182, and/or miR-183. According to the invention, systemic, local or topical administration of at least one agonist of miR-96, miR-182, or miR-183 to a subject in need thereof results in the increased activity of miR-96, miR-182, and/or miR-183 in various eye cells, such as rods, cones, Müller cells, horizontal cells, bipolar cells, amacrine cells, and/or ganglion cells of the subject.

[0084] In one embodiment, administration of at least one agonist of miR-96, miR-182, and/or miR-183 maintains or improves the function of photoreceptor cells such as rods and cones and/or other retinal cells, maintains or improves visual acuity, reduces or prevents the death of photoreceptor cells, and/or reduces or prevents vision loss. In certain embodiments, administration of at least one agonist of miR-96, miR-182, and/or miR-183 maintains or increases the expression of one or more phototransduction genes in the photoreceptor cells of the subject. The one or more phototransduction genes that may be upregulated upon administration of at least one agonist of miR-96, miR-182, and/or miR-183 include Recoverin (Revrn), NRL, Arrestin (Sag), Rhodopsin (Rho), Transducin (Gnat2), and Phosducin (PDC).

[0085] In one embodiment, vision loss, visual acuity, and/or retinal degeneration in a subject is measured using tests such as optokinetic tracking (OKT), electroretinography (ERG), mean spatial frequency threshold (SFT), and measurement of the thickness of the Inner and Outer Nuclear Cell layers of the retina. In another embodiment, a subject's visual acuity is determined using a protocol such as the Early Treatment for Diabetic Retinopathy Study ("ETDRS") or the Age-Related Eye Disease Study ("AREDS") protocol. In some embodiments, visual acuity is measured using a modified ETDRS and/or AREDS protocol, such as the measurement of visual acuity described in Ferris et al., Am J Ophthalmol 94:91-96, 1982. In one embodiment, a subject's visual acuity is determined by one or more of the following procedures: (1) measurement of best-corrected visual acuity (BCVA) with required

manifest refraction; (2) measurement of corrected visual acuity with conditional manifest refraction; or (3) measurement of corrected visual acuity without manifest refraction. In various aspects of the invention, administration of at least one agonist of miR-96, miR-182, or miR-183 to a subject in need thereof results in improved scores on one or more of these eye tests. In some embodiments, administration of at least two agonists, for example, agonists of miR-96 and miR-182, agonists of miR-96 and miR-183, or agonists of miR-182 and miR-183, to a subject results in improved scores on one or more of the eye tests. In some other embodiments, administration of all three agonists, *i.e.*, agonists of miR-96, miR-182 and miR-183, to a subject results in improved scores on one or more of the eye tests. For instance, in one embodiment, a subject upon administration with at least one, at least two, or all three agonists of miR-96, miR-182, and miR-183 has a greater than 3-line, 4-line or 5-line gain in visual acuity in a standardized chart of visual testing, *e.g.*, the ETDRS chart. In another embodiment, administration of at least one, at least two, or all three agonists of miR-96, miR-182, and miR-183 results in the subject's ability to read one or more additional, in some embodiments three or more additional, and in some embodiments 15 or more additional, letters of a standardized chart of vision testing, *e.g.*, the Early Treatment for Diabetic Retinopathy Study Chart ("ETDRS chart").

[0086] In some embodiments, the present invention provides methods of treating or preventing diseases or disorders of other sensory organs in a subject in need thereof comprising administering to the subject a therapeutically effective amount of at least one agonist of miR-96, miR-182, and/or miR-183. For example, in one embodiment, the present invention provides methods of treating or preventing diseases or disorders of ear such as hearing loss, tinnitus, Meniere's disease, ear infections, or damage caused to the ear by these conditions. In some embodiments, methods of treating or preventing ear disorders in a subject in need thereof comprise administering to the subject therapeutically effective amounts of at least two agonists, for example, agonists of miR-96 and miR-182, agonists of miR-96 and miR-183, or agonists of miR-182 and miR-183. In other embodiments, methods of treating or preventing ear disorders in a subject in need thereof comprise administering to the subject therapeutically effective amounts of all three agonists, *i.e.*, agonists of miR-96, miR-182, and miR-183. In some embodiments, the agonist of miR-96, miR-182, or miR-183 is a double-stranded oligonucleotide comprising a first strand containing a sequence of

mature miR-96, miR-182, or miR-183 and a second strand comprising a sequence that is substantially complementary to the first strand, wherein at least one of the strands comprises one or more modified nucleotides. Any of the microRNA mimetic compounds described herein can be used in the methods of treating or preventing an ear disorders in a subject in need thereof. In some embodiments, the agonist of miR-96, miR-182, or miR-183 is an expression vector comprising a polynucleotide sequence encoding a sufficient portion of the miR-96, miR-182, or miR-183 native coding sequence to produce a mature miR-96, miR-182, or miR-183. In one embodiment, the expression vector is a recombinant AAV vector.

[0087] In various embodiments, administration of at least one agonist of miR-96, miR-182, or miR-183 to the subject prevents or slows the development and/or progression of one or more ear disorders and results in the improvement of one or more symptoms associated with these conditions. For instance, in one embodiment, administration of at least one agonist of miR-96, miR-182, or miR-183 results in the improvement of hearing ability. In another embodiment, administration of at least one agonist of miR-96, miR-182, or miR-183 improves the function of sensory cells of the ear. In yet another embodiment, administration of at least two agonists, for example, agonists of miR-96 and miR-182, agonists of miR-96 and miR-183, or agonists of miR-182 and miR-183, results in the improvement of hearing ability and/or the function of ear cells. In still another embodiment, administration of all three agonists, *i.e.*, agonists of miR-96, miR-182, and miR-183, results in the improvement of hearing ability and/or the function of ear cells.

[0088] As used herein, the term “subject” or “patient” refers to any vertebrate including, without limitation, humans and other primates (*e.g.*, chimpanzees and other apes and monkey species), farm animals (*e.g.*, cattle, sheep, pigs, goats and horses), domestic mammals (*e.g.*, dogs and cats), laboratory animals (*e.g.*, rodents such as mice, rats, and guinea pigs), and birds (*e.g.*, domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like). In some embodiments, the subject is a mammal. In other embodiments, the subject is a human.

Pharmaceutical compositions

[0089] The present invention also provides pharmaceutical compositions comprising a therapeutically effective amount of one or more agonists of miR-96, miR-182, and/or miR-183 according to the invention and a pharmaceutically acceptable carrier or excipient. In one embodiment, the present invention provides pharmaceutical compositions comprising a therapeutically effective amount of one or more synthetic microRNA mimetic compounds of miR-96, miR-182, and/or miR-183 according to the invention or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient. In other embodiments, the invention provides pharmaceutical compositions comprising one or more expression vectors encoding miR-96, miR-182, and/or miR-183 and a pharmaceutically acceptable carrier or excipient, wherein the amount of the expression vectors provides therapeutically effective amount of miR-96, miR-182, and/or miR-183.

[0090] The term “pharmaceutically acceptable salt” refers to a salt prepared by combining a compound, such as the disclosed miRNA mimetic compounds, with an acid whose anion, or a base whose cation, is generally considered suitable for human consumption. Suitable pharmaceutically acceptable acid addition salts of the disclosed compounds include those derived from inorganic acids, such as hydrochloric, hydrobromic, hydrofluoric, boric, fluoroboric, phosphoric, metaphosphoric, nitric, carbonic, sulfonic, and sulfuric acids, and organic acids such as acetic, benzenesulfonic, benzoic, citric, ethanesulfonic, fumaric, gluconic, glycolic, isothionic, lactic, lactobionic, maleic, malic, methanesulfonic, trifluoromethanesulfonic, succinic, toluenesulfonic, tartaric, and trifluoroacetic acids.

[0091] Suitable organic acids generally include, for example, aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids. Specific examples of suitable organic acids include acetate, trifluoroacetate, formate, propionate, succinate, glycolate, gluconate, digluconate, lactate, malate, tartaric acid, citrate, ascorbate, glucuronate, maleate, fumarate, pyruvate, aspartate, glutamate, benzoate, anthranilic acid, mesylate, stearate, salicylate, p-hydroxybenzoate, phenylacetate, mandelate, embonate (pamoate), methanesulfonate, ethanesulfonate, benzenesulfonate, pantothenate, toluenesulfonate, 2-hydroxyethanesulfonate, sulfanilate, cyclohexylaminosulfonate, algenic acid, β -hydroxybutyric acid, galactarate, galacturonate, adipate, alginate, butyrate,

camphorate, camphorsulfonate, cyclopentanepropionate, dodecylsulfate, glycoheptanoate, glycerophosphate, heptanoate, hexanoate, nicotinate, 2-naphthalesulfonate, oxalate, palmoate, pectinate, 3-phenylpropionate, picrate, pivalate, thiocyanate, tosylate, and undecanoate.

[0092] Furthermore, where the disclosed compounds carry an acidic moiety, suitable pharmaceutically acceptable salts thereof can include alkali metal salts, *e.g.*, sodium or potassium salts; alkaline earth metal salts, *e.g.*, calcium or magnesium salts; and salts formed with suitable organic ligands, *e.g.*, quaternary ammonium salts. In some forms, base salts are formed from bases which form non-toxic salts, including aluminum, arginine, benzathine, choline, diethylamine, diolamine, glycine, lysine, meglumine, olamine, tromethamine and zinc salts.

[0093] Organic salts can be made from secondary, tertiary or quaternary amine salts, such as tromethamine, diethylamine, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), and procaine. Basic nitrogen-containing groups can be quaternized with agents such as lower alkyl (C1-C6) halides (*e.g.*, methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides), dialkyl sulfates (*e.g.*, dimethyl, diethyl, dibutyl, and diethyl sulfates), long chain halides (*e.g.*, decyl, lauryl, myristyl, and stearyl chlorides, bromides, and iodides), arylalkyl halides (*e.g.*, benzyl and phenethyl bromides), and others. In some forms, hemisalts of acids and bases can also be formed, for example, hemisulphate and hemicalcium salts. In certain embodiments, a pharmaceutically acceptable salt of the present mimetic compounds include a sodium salt.

[0094] In one embodiment, the pharmaceutical composition comprises a therapeutically effective amount of a miR-96 mimetic compound and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the mimetic compound comprises a mature miR-96 sequence and the second strand is substantially complementary to the first strand. In another embodiment, the pharmaceutical composition comprises a therapeutically effective amount of a miR-182 mimetic compound and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the mimetic compound comprises a mature miR-182 sequence and the second strand is substantially complementary to the first strand. In yet another embodiment, the pharmaceutical composition comprises a therapeutically effective amount of

a miR-183 mimetic compound and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the mimetic compound comprises a mature miR-183 sequence and the second strand is substantially complementary to the first strand.

[0095] In some embodiments, the pharmaceutical composition comprises a therapeutically effective amount of at least two microRNA mimetic compounds of the invention and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the first microRNA mimetic compound comprises a mature miR-96 sequence and the first strand of the second microRNA mimetic compound comprises a mature miR-182 sequence. In some other embodiments, the pharmaceutical composition comprises a therapeutically effective amount of at least two microRNA mimetic compounds of the invention and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the first microRNA mimetic compound comprises a mature miR-96 sequence and the first strand of the second microRNA mimetic compound comprises a mature miR-183 sequence. In still some other embodiments, the pharmaceutical composition comprises a therapeutically effective amount of at least two microRNA mimetic compounds of the invention and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the first microRNA mimetic compound comprises a mature miR-182 sequence and the first strand of the second microRNA mimetic compound comprises a mature miR-183 sequence. In yet some other embodiments, the invention provides pharmaceutical compositions comprising a therapeutically effective amount of three microRNA mimetic compounds of the invention and a pharmaceutically acceptable carrier or excipient, wherein the first strand of the first microRNA mimetic compound comprises a mature miR-96 sequence, the first strand of the second microRNA mimetic compound comprises a mature miR-182 sequence, and the first strand of the third microRNA mimetic compound comprises a mature miR-183 sequence.

[0096] Preferably, in the pharmaceutical compositions comprising at least two microRNA agonists according to the invention, the first and the second agonists or the first, second and the third agonists are present in equimolar concentrations. Other mixing ratios such as about 1:2, 1:3, 1:4, 1:5, 1:2:1, 1:3:1, 1:4:1, 1:2:3, 1:2:4 are also envisioned for preparing pharmaceutical compositions comprising at least two of the miR-96, miR-182, and miR-183 agonists.

[0097] In some embodiments, one or more microRNA agonists of the invention may be administered concurrently but in separate compositions, with concurrently referring to mimetic compounds given within short period, for instance, about 30 minutes of each other. In some other embodiments, miR-96, miR-182, and/or miR-183 agonists may be administered in separate compositions at different times.

[0098] The invention also encompasses embodiments where additional therapeutic agents may be administered along with miR-96, miR-182, and/or miR-183 agonists. The additional therapeutic agents may be administered concurrently but in separate formulations or sequentially. In other embodiments, additional therapeutic agents may be administered at different times prior to after administration of miR-96, miR-182, and/or miR-183 agonists. Prior administration includes, for instance, administration of the first agent within the range of about one week to up to 30 minutes prior to administration of the second agent. Prior administration may also include, for instance, administration of the first agent within the range of about 2 weeks to up to 30 minutes prior to administration of the second agent. After or later administration includes, for instance, administration of the second agent within the range of about one week to up to 30 minutes after administration of the first agent. After or later administration may also include, for instance, administration of the second agent within the range of about 2 weeks to up to 30 minutes after administration of the first agent. Where clinical applications are contemplated, pharmaceutical compositions will be prepared in a form appropriate for the intended application. Generally, this will entail preparing compositions that are essentially free of pyrogens, as well as other impurities that could be harmful to humans or animals.

[0099] In some embodiments, pharmaceutical compositions comprising the miR-96, miR-182, and/or miR-183 agonists can be formulated for topical ophthalmic application, for example, in the form of solutions, ointments, creams, lotions, eye ointments, eye drops or eye gels. In some other embodiments, pharmaceutical compositions comprising the miR-96, miR-182, and/or miR-183 agonists can be formulated for local ocular administration via injection. The routes for administering the miR-96, miR-182, and/or miR-183 agonists via

injection include intravitreal, peri-ocular, intracameral, subconjunctival, or transcleral administration.

[00100] In various embodiments, pharmaceutical compositions used for topical or local ocular administration can contain appropriate ophthalmological additives including, for example, buffers, isotonizing agents, preservatives, solubilizers (stabilizers), pH adjusting agents, thickeners and chelating agents, solvents to assist drug penetration, and emollients in ointments and creams. The buffers may be selected from but not limited by the group comprising a phosphate buffer, a borate buffer, a citrate buffer, a tartrate buffer, an acetate buffer (for example, sodium acetate) and an amino acid. The isotonizing agents may be selected from but not limited by the group comprising sugars such as sorbitol, glucose and mannitol, polyhydric alcohols such as glycerin, polyethylene glycol and polypropylene glycol, and salts such as sodium chloride. The preservatives may be selected from but not limited by the group comprising benzalkonium chloride, benzethonium chloride, alkyl paraoxybenzoates such as methyl paraoxybenzoate and ethyl paraoxybenzoate, benzyl alcohol, phenethyl alcohol, sorbic acid and salts thereof, thimerosal and chlorobutanol. The solubilizers (stabilizers) may be selected from but not limited by the group comprising cyclodextrin and derivatives thereof, water-soluble polymers such as poly(vinylpyrrolidone), and surfactants such as polysorbate 80 (trade name: Tween 80). The pH adjusting agents may be selected from but not limited by the group comprising hydrochloric acid, acetic acid, phosphoric acid, sodium hydroxide, potassium hydroxide and ammonium hydroxide. The thickeners may be selected from but not limited by the group comprising hydroxyethylcellulose, hydroxypropylcellulose, methylcellulose, hydroxypropylmethylcellulose and carboxymethylcellulose and salts thereof. The chelating agents may be selected from but not limited by the group comprising sodium edetate, sodium citrate and sodium condensed phosphate. Such topical formulations can further contain compatible ophthalmic carriers, for example cream or ointment bases, olive oil, arachis oil, castor oil, polyoxyethylated castor oil, mineral oil, petroleum jelly, dimethyl sulphoxide, an alcohol (ethanol or oleyl alcohol), liposome, silicone fluid and mixtures thereof as taught by U.S. Pat. No. 6,254,860.

[00101] Alternatively, the miR-96, miR-182, and miR-183 agonists may be applied to the eye via liposomes. In certain embodiments, liposomes used for delivery are amphoteric liposomes such SMARTICLES® (Marina Biotech, Inc.) which are described in detail in U.S. Pre-grant Publication No. 20110076322. The surface charge on the SMARTICLES® is fully reversible which make them particularly suitable for the delivery of nucleic acids. SMARTICLES® can be delivered via injection, remain stable, and aggregate free and cross cell membranes to deliver the nucleic acids.

[00102] Further, the agonists may be infused into the tear film via a pump-catheter system. Another embodiment of the present invention involves the mimetic compounds contained within a continuous or selective-release device, for example, membranes such as, but not limited to, those employed in the pilocarpine (OCUSERT™) System (Alza Corp., Palo Alto, Calif.). As an additional embodiment, the mimetic compounds can be contained within, carried by, or attached to contact lenses which are placed on the eye. Another embodiment of the present invention involves the mimetic compounds contained within a swab or sponge which can be applied to the ocular surface. Yet another embodiment of the present invention involves the mimetic compounds contained within a liquid spray which can be applied to the ocular surface.

[00103] This invention is further illustrated by the following additional examples that should not be construed as limiting. Those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made to the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

[00104] All patent and non-patent documents referenced throughout this disclosure are incorporated by reference herein in their entirety for all purposes.

EXAMPLES:

Example 1: Uptake and clearance of miRNA mimics in retina

[00105] Naked (not conjugated to cholesterol), miRNA mimics for mouse miR-96, miR-182, and miR-183 were resuspended in saline. The three duplexes were pooled at 3.3 µg/µl of

each duplex to obtain 10 µg/µl of pooled duplexes. Naked siRNA targeting mouse rhodopsin (NM_145383) was used as a positive control and was suspended in saline at a concentration of 10 µg/µl. To determine the uptake and clearance of miRNA mimics in retina, wild-type C57Bl/6 mice were injected intravitreally with 1 µl (10 µg) of pooled miRNA duplexes or rhodopsin siRNA per eye. Retina was isolated after each time point and the retinal distribution and clearance was measured by isolating RNA and measuring miRNA levels by sandwich ELISA. For the distribution and clearance of siRNA, RNA was isolated and target repression was measured by qRT-PCR.

[00106] Study Design:

2 study arms:

- 1) Pool of miR-96/182/183 mimics, 10 µg total (n=14)
- 2) siRNA targeting rhodopsin (n=14)

7 time points: (2 mice/study arm/timepoint)

- 1) Untreated (baseline)
- 2) 4h
- 3) 8h
- 4) 24h
- 5) 48h
- 6) 72h
- 7) 168h (day 7)

[00107] miRNA mimic/siRNA quantitation assay to assess oligonucleotide biodistribution

A sandwich hybridization assay was used to quantify miR-183, miR-96, miR-182, or rho siRNA in tissue samples as previously described by Efler, et al ("Quantitation of oligodeoxynucleotides in human plasma with a novel hybridization assay offers greatly enhanced sensitivity over capillary electrophoresis," *Oligonucleotides* 15(2), 119–131 (2005)). Briefly, probes for the hybridization assay were synthesized with 2'-O-methyl modified nucleotides and were tagged as 5' bTEG-sup-3' (capture probe) and 5'-6FAM-sup-3' (detection probe). Detection was accomplished using anti-fluorescence-peroxidase, Fab fragments (Roche), and TMB Peroxidase Substrate (KPL). Standard curves were generated with nonlinear logistic regression analysis with 4 parameters (4-PL). The working concentration range of the assay was 1 to 536 ng/mL. Tissue samples were prepared at 100 mg/mL by homogenizing in 3 mol/L GITC buffer (3 mol/L guanidineisothiocyanate, 0.5

mol/L NaCl, 0.1 mol/L Tris, pH 7.5, and 10 mmol/L EDTA) two times for 30 seconds with an MP FastPrep-24 at a speed setting of 6.0. Tissue homogenates were diluted in 1 mol/L GITC Buffer (1 mol/L guanidine isothiocyanate, 0.5 mol/L NaCl, 0.1 mol/L Tris, pH 7.5, and 10 mmol/L EDTA) for testing.

[00108] Quantitative Real-Time Polymerase Chain Reaction Analysis of mouse rhodopsin For in vivo real-time polymerase chain reaction (RT-PCR) analysis, RNA was extracted from retina tissue with Trizol (Invitrogen); then, 100 ng total RNA from each sample was used to generate cDNA with MultiScribe reverse transcriptase (Life Technologies) according to the manufacturer's specifications. Gene expression was measured with Life Technologies Taqman gene expression assays. Gene expression was normalized to a housekeeping gene such as GAPDH and calculated as relative expression compared to the average of the control group.

[00109] For animals treated with miRNA mimic pool, 2 retinas from each animal were pooled for biodistribution analysis. For animals treated with Rho siRNA, one retina per animal was used for biodistribution, and the other retina was used for RNA isolation to quantitate target knockdown.

[00110] All 3 miRNA mimics and the Rho siRNA distributed to the retina after a single intravitreal injection. The amount of each miRNA mimic is approximately one-third the amount of siRNA detected in retina, consistent with the fact that the miRNA mimic pool contained 3.3 μ g of each miRNA mimic (10 μ g total oligo), while the siRNA was dosed at a total concentration of 10 μ g. All three miRNA mimics were detected at 4h and 8h post-dose and were cleared rapidly after 8h (FIG. 1). Rho siRNA was detected at 4h and 8h post-injection and was cleared by 24h.

[00111] The Rho (rhodopsin) siRNA produced 60% silencing of the target gene at 72h post-injection, and silencing remained at ~50% at 7 days post-injection (FIG. 2). Rhodopsin is expressed in photoreceptors in the retina. The current data therefore demonstrate that siRNA duplexes distribute functionally to photoreceptors. Silencing is retained for ~ 1 week post-injection, even though the siRNA itself is not detectable in the retina after 24h.

Example 2: Administration of microRNA mimics (miR-96, miR-182, and miR-183) in rat retinal cell system

[00112] For this study, cholesterol-conjugated, mimics for miR-96, miR-182, and miR-183, and rat rhodopsin siRNA were used.

[00113] Table 3: miR-96/182/183 mimics used in the study:

SEQ ID NO.	Modified Sequence
13	5'-mA.mG.mC.rA.rA.rA.mU.mU.rG.rA.rG.mC.mU.rA.rG.mU.rG.mC.rG.rA.rA.rA.cho16-3'
18	5'-mA.mG.mU.rG.mU.rG.rA.rG.mU.mC.rA.rA.mC.mC.rA.mU.mU.rG.mC.rG.rA.rA.rA.cho16-3'
23	5'-mA.mG.mU.rG.rA.rA.rA.mU.mC.rA.rA.mC.mC.rA.rG.mU.rG.mC.rG.rA.mU.rA.cho16-3'
20	5'-rU.rA.fU.rG.rG.fC.rA.fC.fU.rG.rG.fU.rA.rG.rA.fU.fU.fC.rA.fC.fUs.rUs.rU-sup-3'
10	5'-rU.rU.fU.rG.rG.fC.rA.fC.fU.rA.rG.fC.rA.fC.rA.fU.fU.fU.rG.fC.fUs.rUs.rU-sup-3'
15	5'-rU.rU.fU.rG.rG.fC.rA.rA.fU.rG.rG.fU.rA.rG.rA.rA.fC.fU.fC.rA.fC.fUs.rUs.rU-sup-3'

[00114] Cholesterol-conjugated mimics of each of the 3 microRNAs were administered to rat retinal cells purchased from Lonza (R-RET-508). Lonza's rat retinal cell system isolated from neonatal (P3 or P4) Sprague-Dawley rats comprises 7 cell types normally found in the retina (rods, cones, Müller cells, horizontal cells, bipolar cells, amacrine cells, and ganglion cells). Although this is a mixed cell population, data on the proportions of cell types in the rat retina indicate that rods are the predominant cell type, and rods are also the cell type in which the miR-183 cluster is normally most highly expressed. Studying the effect of miRNA mimics in the context of other retinal neuronal cell types could provide more relevant, context-dependent, biology compared to a single cell type in isolation. Additionally, this mixed cell system could provide a more direct comparison for the gene expression changes identified in subsequent *in vivo* studies. At P3/P4 neonatal stage in the rats, the microRNAs in the miR-183/96/182 cluster are relatively lowly expressed (~1% of levels in the adult retina) thereby providing low background and a more robust signature for exogenously added mimic.

[00115] Rat retinal cells (R-Ret cells) in culture were passively transfected with cholesterol-conjugated microRNA mimics to determine various parameters: toxicity of miRNA mimics,

down-regulation of Rho mRNA using Rho siRNA, optimal plating density of R-Ret cells, maximum duration for culturing R-Ret cells, and expression profiling of genes involved in the phototransduction pathway using real-tile PCR.

[00116] Study Design:

- A. Determine the highest dose of cholesterol conjugated miRNA mimic tolerated by R-Ret cells with 10 μ M being the highest dose administered.
- B. Assess functional uptake by measuring Rho knockdown using cholesterol-conjugated siRNA pool
- C. Determine the optimal plating density of R-Ret cells that yields at least 800 ng of total RNA (the minimum required for microarray profiling on the Affymetrix platform)
- D. Determine maximal amount of time cells can be cultured
- E. Perform real time PCR on a selection of genes that are functionally involved in the phototransduction pathway

End points:

- A. Toxicity/viability assay and visual assessment
- B. qPCR of Rho
- C. UV/Visible RNA quantification
- D. miRNA qPCR
- E. qPCR of Rho, Sag, Arr3, Revm, Nrl, Pdc, Gnat1, Gnat2, Opnlmw

[00117] R-Ret cells were passively transfected with cholesterol-conjugated miR-206 mimic and the cells were observed 72 hours post-transfection. 10 μ M dose of cholesterol-conjugated miRNA mimic appeared to cause differentiation of the cells, 72 hours post-transfection and 1 week after start of the culture (FIG. 3). Toxicity of miRNA mimic on R-Ret cells was measured using an adenylate kinase assay. Toxicity was found to decrease over time in serum-free media (FIG. 4) and the treatment of cells with 10 μ M miRNA mimic was not found to be toxic.

[00118] R-Ret cells were passively transfected with various concentrations of cholesterol-conjugated Rho siRNA or non-targeting control (NTC) siRNA. RNA was isolated and a real-time PCR analysis of Rho mRNA was performed. FIG. 5 shows that 1, 5, and 10 μ M of cholesterol-conjugated Rho siRNA pool produced comparable Rho knockdown.

[00119] RNA was isolated from R-Ret cells cultured for one week and yield was determined using a UV/visible spectrophotometer (Table 4).

Table 4

Using miRNesay columns/qiazol					ng/uL	ng/uL
Cell	Day after plating	Density	Plate well	elution volume	1st elute	2nd elute
R-Ret	1 week	140K	12	30	60.7	39.4

[00120] To determine the maximum amount of time cells can be cultured, R-Ret cells were plated on poly-L-lysine. Cells survived for at least two weeks in culture where the cells were cultured in 5% serum media for first four days and then the media was changed to serum free media.

[00121] R-Ret cells were transfected with various concentrations of pooled or individual miR-183, miR-96, and miR-182 mimics and Rho siRNA. RNA was isolated 72 hours post-transfection and real time PCR analysis was performed to determine the mRNA expression profile of genes involved in the phototransduction pathway. FIGs. 6 and 7 show relative expression levels of genes that are expressed within the range of the PCR assay. P-values of difference were calculated by two-way ANOVA with Newman-Keuls test for multiple comparisons compared to the untreated group. FIG. 8 shows the heat map of the log2-transformed average fold change values of the treatments shown in FIG. 7.

Table 5: Genes examined by real time PCR

Gene name	Gene symbol	Cell type expression	Expression level (real time)
Rhodopsin	Rho	Rod-specific	Good
Arrestin	Sag	Rod-specific	Good

Transducin	Gnat1	Rod-specific	Low
Phosducin	PDC	Rod-specific	Good
NRL	NRL	Rod-specific	Good
Transducin	Gnat2	Cone-specific	Good
M-cone opsin	Opn1mw	Cone-specific	Not detectable
Arrestin	Arr3	Cone-specific	Low
Recoverin	Rcvn	Rod and cone	Good

[00122] Table 6 shows a comparison of the expression data of selected genes in the miR-183 cluster knockout mouse with that of rat retinal cells treated with cholesterol-conjugated miR-183 cluster miRNA mimics.

Table 6

Gene	miR-183 cluster Knock-out	Mimics of miR-183 cluster in vitro	Rho siRNA
Rcvn	-	↑	-
Nrl	↓	↑	-
Sag	-	↑	-
Rho	↓	↑	↓
Pdc	-	↑	-
Gnat2	↓	↑	-

Example 3: Identification of direct and downstream targets of microRNA mimics (miR-96, miR-182, miR-183) in retina

[00123] R-Ret cells in culture are passively transfected with cholesterol-conjugated microRNA mimics (miR-183, miR-96, miR-182) individually or pooled or with cholesterol-conjugated Rho siRNA. At various time points post-transfection, RNA is isolated and subjected to microarray profiling to identify significantly regulated transcripts.

Example 4: Administration of microRNA mimics prevents vision loss and retinal degeneration

[00124] The effect of microRNA mimetic compounds (pool of microRNA mimics of miR-96 and miR-182, and a control duplex) on vision loss and retinal degeneration in a mouse model of retinitis pigmentosa was examined. Rho siRNA was used as a positive control.

[00125] Rd10/rd10 mice were dark reared to P31, at which point they were moved to a 12 hour light/dark cycle to induce retinal degeneration. At P31, the test agents (microRNA pool or Rho siRNA) were administered via bilateral intravitreal (IVT) injection at a concentration of 10 µg (Rho siRNA), or 2 µg and 10 µg (microRNA pool of miR-96 and miR-182, and a control duplex). Control groups included rd10/rd10 mice receiving bilateral administration of vehicle at P31, or animals receiving intraperitoneal (i.p.) administration of phenyl-N-tert-butylnitron (PBN) daily from P29 to P35. For assessment of vision loss by optokinetic tracking (OKT) and electroretinography (ERG), an additional control group of untreated C57Bl/6J mice was included. Vision loss was assessed by both visual acuity measurements (P38 and P45) and ERG (P39 and P46). Visual acuity was assessed by determining the mean spatial frequency threshold (SFT) at which mice could distinguish visual stimuli presented in a virtual environment.

[00126] Administration of both 10 µg of pooled mimics (miR-96 and miR-182 mimics, and a control duplex) and PBN to rd10/rd10 mice resulted in a statistically significant retention of visual acuity as determined by OKT at P45.

[00127] Experimental Design

rd10/rd10 mice

- P1-P30: Animals dark-reared from birth
- P29-P35: Daily intraperitoneal administration of PBN (Arm 5)
- P31: Animals transferred to housing in normal cyclic light (~200 lux during daylight hours)
- P31: Bilateral intravitreal dosing of vehicle or test agents (Arms 1-4)
- P38: OKT analyses to quantify spatial frequency threshold
- P45: OKT analyses to quantify spatial frequency threshold

Animals

Strain: rd10/rd10 mice
 Sex: Male/Female
 Age Range: Newborn pups
 Weight Range: n/a
 Supplier: In-house breeding
 Number of Study Animals: 48
 Number of Spare Animals 0
 Number of Sentinel Animals: 0

Test compounds and Vehicle

Table 7

Compound	Preparation & Storage	Route of Administration
Pool of miR-96 mimic, miR-182 mimic, and control duplex	Store aliquots at -20°C until use. After thawing, store at 4°C.	Intravitreal
Rho siRNA	Store aliquots at -20°C until use. After thawing, store at 4°C.	Intravitreal
Phenyl-N-tert-butylnitronate (PBN)	Dry powder stored at RT. Working solutions prepared in 0.9% NaCl immediately prior to use.	Intraperitoneal

Study Arms

Table 8

Arm	Housing	Treatment	Treatment Details	Assessment
1	Dark-reared (P1-30); 12 h light/dark (P31-P46)	0.9% NaCl	Bilateral IVT (P31)	OKT (P38 and P45); ERG (P39 and P46); Retinal thickness quant
2	Dark-reared (P1-30); 12 h light/dark (P31-P46)	2 µg Pooled mimics	Bilateral IVT (P31)	OKT (P38 and P45); ERG (P39 and P46); Retinal thickness quant

3	Dark-reared (P1-30); 12 h light/dark (P31-P46)	10 µg Pooled mimics	Bilateral IVT (P31)	OKT (P38 and P45); ERG (P39 and P46); Retinal thickness quant
4	Dark-reared (P1-30); 12 h light/dark (P31-P46)	10 µg Rho siRNA	Bilateral IVT (P31)	OKT (P38 and P45); ERG (P39 and P46); Retinal thickness quant
5	Dark-reared (P1-30); 12 h light/dark (P31-P46)	100 mg/kg PBN	intraperitoneal administration (P29-P35)	OKT (P38 and P45); ERG (P39 and P46); Retinal thickness quant

[00128] Animal Housing

All animals were housed in groups of 3-5 in large cages kept in ventilated shelves under standard animal care conditions. Rd10/rd10 pregnant dams were housed in darkness upon observation of a mucus plug. Newborn pups were housed with mothers in complete darkness from postnatal day 1 through postnatal day 30. On postnatal day 31, animals were transitioned to maintenance under normal cyclical light conditions consisting of 12 hours of light (< 500 lux) followed by 12 hours of darkness.

[00129] Formulation preparation and storage

Test agents were supplied as aliquots at 10 µg/µl, and were ready to inject. For arm 2, test agent (pooled mimics) was diluted 1:5 in 0.9% NaCl to achieve a final concentration of 2 µg/µl. A total volume of 1 µl was delivered for all intravitreal injections. Aliquots were stored at -20°C until use. Following the initial thaw, all remaining material was stored at 4°C, and was not refrozen. PBN was prepared immediately prior to use as a 15 mg/ml solution in 0.9% NaCl (Cat #S4041, Teknova). PBN was delivered daily as indicated in Section 3.2 at a dose of 100 mg/kg in a total volume of 75-150 µl, depending on the animal's body weight.

[00130] Intraperitoneal (IP) Administration

Sedatives and the positive control test agent (PBN) were delivered by standard techniques for IP injection utilizing a 0.3 cc insulin syringe attached to an 8mm 31-gauge needle (BD#328438) with a total volume ≤ 150 µl.

[00131] Intravitreal Administration

Animals were anesthetized with ketamine/xylazine using a U-100 syringe utilizing Ketamine (85 mg/kg) and Xylazine (14 mg/kg). Pupils were then dilated with topical administration of Cyclogyl and Ak-Dilate. Following sedation and dilation, a total volume of 1 μ l per eye was injected into the vitreous at the pars plana using a Hamilton syringe and a 33 gauge needle.

[00132] Optokinetic tracking (OKT)

All optokinetic tracking experiments are performed using an Optomotry designed for rodent use (Cerebral Mechanics Inc.). In this non-invasive assessment, mice are placed onto a platform surrounded by 4 LCD screens which resides within a light-protected box. Visual stimuli are then presented to the mice via the LCD screens and a masked observer visualizes and scores optokinetic tracking reflexes from a digital camcorder which is mounted on the top of the box. For measurements of spatial frequency threshold, the mice were tested at a range of spatial frequencies from 0.034 to 0.514 cycles/degree. The Optomotry device employs a proprietary algorithm to accept the input from the masked observer and automatically adjust the testing stimuli based upon whether the animal exhibited the correct or incorrect tracking reflex.

[00133] Tissue collection

Following sedation with ketamine/xylazine, animals were euthanized with a lethal dose of pentobarbital. The right eye of all animals was scorched with a flamed needle to demarcate the superior portion of the eye, enucleated, fixed in Z-fix (zinc buffered neutral formalin), and processed for H&E histology. From the left eye of all animals, the retinas were individually isolated, and immediately snap-frozen in liquid N₂, and stored individually in a 2 mL screw cap

polypropylene tube at -70° C until further processing.

[00134] Data and Statistical Analyses

Statistical significance was determined using Prism software (Graphpad Inc.) to perform t-test (OKT and ERG) or one-way Analysis of Variance (ANOVA) calculations (retinal thickness), with a threshold of $p < 0.05$ to determine whether any changes are statistically significant.

[00135] FIG. 9 shows the effect of test agents on visual acuity loss in the mouse model of retinitis pigmentosa. Visual acuity was assessed by the mean SFT at which mice were able to distinguish visual stimuli. Visual acuity was lower in all groups of rd10 mice relative to C57Bl/6J control mice at both P38 and P45. At P38, visual acuity was comparable for the vehicle treated group and both groups receiving pooled mimics. The group receiving 10 μ g of Rho siRNA had statistically significant lower visual acuity relative to the vehicle treated group at P38 ($p = 0.0185$; unpaired t test), whereas the PBN treated positive control group displayed statistically significant higher visual acuity measurements relative to vehicle ($p = 0.0224$; unpaired t test). At P45, there is a dose-dependent amelioration in vision loss for the two groups receiving the pooled mimics, with a statistically significant difference between the 10 μ g group of the pooled mimics and the vehicle ($p = 0.0258$; unpaired t test). The PBN treated animals also display statistically significant better visual acuity relative to vehicle at P45 ($p = 0.0416$; unpaired t test). The visual acuity for the 10 μ g Rho siRNA group remained much lower than the age-matched vehicle group, but the difference was not statistically significant.

[00136] FIG. 10 shows the effect of test agents on visual acuity decline in the mouse model of retinitis pigmentosa. Visual acuity loss occurred across all groups from P38 to P45. Vehicle treated animals exhibit a 48% decrease in visual acuity from P38 to P45. The largest decline in visual acuity across the two time points occurred for the group receiving 10 μ g Rho siRNA test agent (59%). The group receiving 10 μ g of pooled mimics had the most negligible loss in visual acuity across the two time points (11%). Visual acuity loss was less dramatic for both the 2 μ g group of pooled mimics and the PBN group relative to vehicle (35% and 39%, respectively).

[00137] At both postnatal time points examined, vision loss assessed by OKT was statistically significantly lower in light exposed rd10/rd10 mice relative to both control C57Bl/6J mice, and rd10/rd10 mice prior to light exposure. The vehicle treated group was lower in its SFT than all groups except the 10 μ g Rho siRNA treatment group. Bilateral intravitreal administration of 10 μ g of pooled mimics (miR-96, miR-182, and control duplex) demonstrates a positive effect on visual acuity as demonstrated by a statistically significant retention in SFT measurements at P45, and the lowest percentage drop off in SFT from P38

to P45 amongst all groups (FIGs. 9 and 10). The PBN positive control treatment group also retained a statistically significant higher SFT at P45 relative to the vehicle control group, however, the loss in SFT from P38 to P45 was greater in the PBN group than the 10 μ g group of pooled mimics. The 2 μ g treatment group of pooled mimics had a higher SFT at P45 than the vehicle control group, but the difference was not statistically significant. The group receiving 10 μ g Rho siRNA had the largest loss in SFT measurements at both P38 and P45, and also had the largest percentage decline in SFT between the two time points. Based on the data, bilateral intravitreal administration of 10 μ g of pooled mimics ameliorates vision loss at the latter time point with a statistically significant difference in visual acuity measured by OKT.

CLAIMS:

1. A microRNA mimetic compound comprising:
 - a first strand of about 22 to about 26 ribonucleotides comprising a mature miR-96, miR-182, or miR-183 sequence; and
 - a second strand of about 20 to about 26 ribonucleotides comprising a sequence that is substantially complementary to the first strand and having at least one modified nucleotide.
2. The microRNA mimetic compound of claim 1, wherein the first strand has one or more 2' fluoro nucleotides.
3. The microRNA mimetic compound of claim 1, wherein the first strand has no modified nucleotides.
4. The microRNA mimetic compound of any one of claims 1 to 3, wherein the at least one modified nucleotide in the second strand is a 2'-O-methyl modified nucleotide.
5. The microRNA mimetic compound of any one of claims 1 to 4, wherein the second strand is not fully complementary to the first strand.
6. The microRNA mimetic compound of claim 5, wherein the second strand has 1, 2, or 3 mismatches relative to the first strand.
7. The microRNA mimetic compound of any one of claims 1 to 6, wherein the second strand is linked to a cholesterol molecule at its 3' or 5' terminus.
8. The microRNA mimetic compound of claim 7, wherein the cholesterol molecule is linked to the second strand through at least a six carbon linker.
9. The microRNA mimetic compound of any one of claims 1 to 8, wherein the first strand has a 5'-terminal monophosphate.

10. The microRNA mimetic compound of any one of claims 1 to 9, wherein the first or the second strand has a 3' nucleotide overhang relative to the other strand.
11. The microRNA mimetic compound of any one of claims 1 to 10, wherein the nucleotides comprising the 3' overhang are linked by phosphorothioate linkages.
12. The microRNA mimetic compound of any one of claims 1 to 11, wherein the 3' nucleotide overhang comprises two ribonucleotides.
13. The microRNA mimetic compound of any one of claims 1 to 12, wherein the first strand comprises a mature miR-96 sequence.
14. The microRNA mimetic compound of claim 13, wherein the first strand comprises a sequence of SEQ ID NO: 10.
15. The microRNA mimetic compound of claim 13 or 14, wherein the second strand comprises a sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, and SEQ ID NO: 13.
16. The microRNA mimetic compound of claim 13, wherein the first strand comprises a sequence selected from the group consisting of SEQ ID NOs: 10 and 26-29 and the second strand comprises a sequence selected from the group consisting of SEQ ID NOs: 11-14 and 30-34.
17. The microRNA mimetic compound of any one of claims 1 to 12, wherein the first strand comprises a mature miR-182 sequence.
18. The microRNA mimetic compound of claim 17, wherein the first strand comprises a sequence of SEQ ID NO: 15.

19. The microRNA mimetic compound of claim 17 or 18, wherein the second strand comprises a sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17, and SEQ ID NO: 18.
20. The microRNA mimetic compound of claim 17, wherein the first strand comprises a sequence selected from the group consisting of SEQ ID NOs: 15 and 35-38 and the second strand comprises a sequence selected from the group consisting of SEQ ID NOs: 16-19 and 39-43.
21. The microRNA mimetic compound of any one of claims 1 to 12, wherein the first strand comprises a mature miR-183 sequence.
22. The microRNA mimetic compound of claim 21, wherein the first strand comprises a sequence of SEQ ID NO: 20.
23. The microRNA mimetic compound of claim 21 or 22, wherein the second strand comprises a sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, and SEQ ID NO: 24.
24. The microRNA mimetic compound of claim 21, wherein the first strand comprises a sequence selected from the group consisting of SEQ ID NOs: 20 and 44-47 and the second strand comprises a sequence selected from the group consisting of SEQ ID NOs: 21-25 and 48-52.
25. A pharmaceutical composition comprising a therapeutically effective amount of the microRNA mimetic compound of any one of claims 1 to 24, or a pharmaceutically-acceptable salt thereof, and a pharmaceutically-acceptable carrier or diluent.
26. A pharmaceutical composition comprising a therapeutically effective amount of at least two microRNA mimetic compounds of claim 1, wherein the first strand of the first microRNA mimetic compound comprises a mature miR-96 sequence and the first strand of

the second microRNA mimetic compound comprises a mature miR-182 or miR-183 sequence.

27. The pharmaceutical composition of claim 26, wherein the first strand of the second microRNA mimetic compound comprises a mature miR-182 sequence.

28. The pharmaceutical composition of claim 27, further comprising a third microRNA mimetic compound, wherein the first strand of the third microRNA mimetic compound comprises a mature miR-183 sequence.

29. The pharmaceutical composition of claim 28, wherein the first, second, and third microRNA mimetic compounds are present in equimolar concentrations.

30. A method of treating or preventing an ophthalmological condition in a subject in need thereof comprising administering to the subject a therapeutically effective amount of at least one agonist of miR-96, miR-182, and/or miR-183, wherein the agonist is a double-stranded oligonucleotide comprising a first strand comprising a mature miR-96, miR-182, or miR-183 sequence and a second strand comprising a sequence that is substantially complementary to the first strand, wherein at least one of the strands comprises one or more modified nucleotides.

31. The method of claim 30, wherein the therapeutically effective amount is an amount sufficient to maintain or improve visual acuity in the subject.

32. The method of claim 30, wherein the therapeutically effective amount is an amount sufficient to reduce or prevent photoreceptor cell damage and/or death in the subject.

33. The method of any one of claims 30 to 32, wherein the first strand is about 22 to about 26 nucleotides in length and the second strand is about 20 to about 26 nucleotides in length.

34. The method of any one of claims 30 to 33, wherein the first strand has one or more 2' fluoro nucleotides.
35. The method of any one of claims 30 to 34, wherein the second strand has one or more 2'-O-methyl modified nucleotides.
36. The method of any one of claims 30 to 35, wherein the second strand has 1, 2, or 3 mismatches relative to the first strand.
37. The method of any one of claims 30 to 35, wherein the first or the second strand has a 3' nucleotide overhang relative to the other strand.
38. The method of claim 37, wherein the nucleotides comprising the 3' overhang are linked by phosphorothioate linkages.
39. The method of claim 37 or 38, wherein the 3' nucleotide overhang comprises two ribonucleotides.
40. The method of any one of claims 30 to 39, wherein the second strand is linked to a cholesterol molecule at its 3' or 5' terminus.
41. The method of claim 40, wherein the cholesterol molecule is linked to the second strand through at least a six carbon linker.
42. The method of any one of claims 30 to 41, wherein the agonist is a miR-96 agonist and the first strand of the double-stranded oligonucleotide comprises a mature miR-96 sequence.
43. The method of claim 42, wherein the first strand comprises a sequence selected from the group consisting of SEQ ID NOs: 10 and 26-29 and the second strand comprises a sequence selected from the group consisting of SEQ ID NOs: 11-14 and 30-34.

44. The method of any one of claims 30 to 41, wherein the agonist is a miR-182 agonist and the first strand of the double-stranded oligonucleotide comprises a mature miR-182 sequence.

45. The method of claim 44, wherein the first strand comprises a sequence selected from the group consisting of SEQ ID NOs: 15 and 35-38 and the second strand comprises a sequence selected from the group consisting of SEQ ID NOs: 16-19 and 39-43.

46. The method of any one of claims 30 to 41, wherein the agonist is a miR-183 agonist and the first strand of the double-stranded oligonucleotide comprises a mature miR-183 sequence.

47. The method of claim 46, wherein the first strand comprises a sequence selected from the group consisting of SEQ ID NOs: 20 and 44-47 and the second strand comprises a sequence selected from the group consisting of SEQ ID NOs: 21-25 and 48-52.

48. The method of claim 42, further comprising administering to the subject a miR-182 agonist, wherein the miR-182 agonist is a double-stranded oligonucleotide comprising a first strand comprising a mature miR-182 sequence and a second strand comprising a sequence that is substantially complementary to the first strand, wherein at least one of the strands comprises one or more modified nucleotides.

49. The method of claim 42 or 48, further comprising administering to the subject a miR-183 agonist, wherein the miR-183 agonist is a double-stranded oligonucleotide comprising a first strand comprising a mature miR-183 sequence and a second strand comprising a sequence that is substantially complementary to the first strand, wherein at least one of the strands comprises one or more modified nucleotides.

50. The method of claim 49, wherein the miR-96, miR-182, and miR-183 agonists are administered to the subject in separate compositions.

51. The method of claim 49, wherein the miR-96, miR-182, and miR-183 agonists are administered to the subject in the same composition.
52. The method of claim 51, wherein the miR-96, miR-182, and miR-183 agonists are present in the composition at equimolar concentrations.
53. The method of any one of claims 30 to 52, wherein the at least one agonist is administered to the subject ocularly.
54. The method of claim 53, wherein the ocular administration comprises intravitreal, peri-ocular, intracameral, subconjunctival, or transcleral administration.
55. The method of any one of claims 30 to 54, wherein the subject has retinitis pigmentosa.
56. The method of any one of claims 30 to 54, wherein the subject has signs of night blindness.
57. The method of any one of claims 30 to 54, wherein the subject has an ophthalmological condition selected from the group consisting of retinal detachment, retinal degeneration, macular degeneration, and Stargardt disease.
58. The method of any one of claims 30 to 57, wherein the therapeutically effective amount is an amount sufficient to increase expression of one or more phototransduction genes in the photoreceptor cells of the subject.
59. The method of claim 58, wherein the one or more phototransduction genes are selected from Recoverin (Revm), NRL, Arrestin (Sag), Rhodopsin (Rho), Transducin (Gnat2), and Phosducin (PDC).
60. A method of treating or preventing an ear disorder in a subject in need thereof comprising administering to the subject a therapeutically effective amount of at least one

agonist of miR-96, miR-182, and/or miR-183, wherein the agonist is a double-stranded oligonucleotide comprising a first strand comprising a mature miR-96, miR-182, or miR-183 sequence and a second strand comprising a sequence that is substantially complementary to the first strand, wherein at least one of the strands comprises one or more modified nucleotides.

61. The method of claim 60, wherein the ear disorder is selected from the group consisting of hearing loss, tinnitus, Meniere's disease, and ear infections.

62. An expression vector comprising a polynucleotide encoding miR-96, miR-182, or miR-183 for expression in a mammalian cell.

63. The expression vector of claim 62, wherein the vector is a viral expression vector.

64. The expression vector of claim 63, wherein the viral vector is an adeno-associated viral vector.

65. The expression vector of claim 63, wherein the adeno-associated viral vector is a self-complementary adeno-associated viral vector.

66. The expression vector of claim 64 or 65, wherein the adeno-associated viral vector is selected from the group consisting of AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, and AAV9.

67. The expression vector of claim 62, wherein the expression vector comprises a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 53-55.

68. A method of treating or preventing an ophthalmological condition or an ear condition in a subject in need thereof comprising administering to the subject an effective amount of an expression vector encoding a miR-96, miR-182, and/or miR-183.

1/9




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D

2/9




FIG. 2

3/9





FIG. 4

4/9




FIG. 5

5/9




FIG. 6A




FIG. 6B

6/9




FIG. 7A




FIG. 7B

7/9




FIG. 8




FIG. 9

9/9




FIG. 10

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US16/22645

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12N 15/86, 15/113 (2016.01)

CPC - A01K 67/0275; C12N 15/11, 15/86, 15/113, 15/111

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) Classifications: C12N 15/86; 15/13; CPC Classifications: A01K 67/0275; C12N 15/11; 15/86, 15/113, 15/111

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatSeer (US, EP, WO); USPTO Web Page; Google; Google Scholar; EBSCO; Entrez Pubmed; Science Direct; Search Terms -- microRNA, mimetics, miR-96, miR-182, miR-183, 'visual acuity', 'hearing loss', 'photoreceptor damage', 2-O nucleotide, 2-fluoro nucleotide

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2013/0225652 A1 (CHORN, G et al.) August 29, 2013; paragraphs [0005], [0007], [0013], [0015], [0016], [0079], [0084], [0096], [0098], [0119], [0186]	1-3, 4/1-3, 26-32, 33/30-32, 60-61
Y	US 2015/0051267 A1 (PURDUE RESEARCH FOUNDATION) February 19, 2015; abstract; paragraphs [0007], [0013], [0040], [0063], [0064], [0081], [0088], [0093]	1-3, 4/1-3, 26-29, 60, 61
Y	US 2012/0157514 A1 (ESAU, C et al.) June 21, 2012; paragraphs [0025], [0624]	29
Y	US 2013/0237587 A1 (THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES) September 12, 2013; paragraphs [0004], [0074], [0082], [0147], [0158]	30-32, 33/30-32
A	US 2005/0261218 A1 (ESAU, C et al.) November 24, 2005; entire document	1-3, 4/1-3, 26-32, 33/30-32, 60-61
A	WO 2014/100252 A1 (UNIVERSITY OF WASHINGTON THROUGH ITS CENTER FOR COMMERCIALIZATION) June 26, 2014; entire document	1-3, 4/1-3, 26-32, 33/30-32, 60-61
A	(WEI, Q et al.) Roles of miR-182 in Sensory Organ Development and Cancer. Thoracic Cancer. 7 January 2015, Vol. 6, No. 1; pages 2-9; DOI: 10.1111/1759-7714.	1-3, 4/1-3, 26-32, 33/30-32, 60-61

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:	
"A"	document defining the general state of the art which is not considered to be of particular relevance
"E"	earlier application or patent but published on or after the international filing date
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O"	document referring to an oral disclosure, use, exhibition or other means
"P"	document published prior to the international filing date but later than the priority date claimed
"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&"	document member of the same patent family

Date of the actual completion of the international search

19 July 2016 (19.07.2016)

Date of mailing of the international search report

19 AUG 2016

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450
Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US16/22645

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos..
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 5-25, 34-59
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

-***-Continued Within the Next Supplemental Box-***-

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-4, 26-33, 60, 61

Remark on Protest

The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/US16/22645

-***-Continued from Box III: Lack of Unity of Invention -***-

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, Claims 1-4, 26-33, 60 and 61 are directed toward a microRNA mimetic compound comprising a first strand, and a second strand that is substantially complementary to the first strand, and having at least one modified nucleotide; a composition comprising said compound and methods of treating or preventing an ophthalmological disorder or ear disorder with said compound

Groups II+, Claims 62-68 and SEQ ID NO: 53 are directed toward an expression vector comprising a polynucleotide encoding miR-96, miR-182, or miR-183 for expression in a mammalian cell; and a method of treating or preventing an ophthalmological condition or an ear condition, comprising administering said vector to a subject.

The expression vector can be searched to the extent that it comprises a sequence encoding an miRNA encompassing SEQ ID NO: 53 (first exemplary miRNA encoding sequence). Applicant is invited to elect additional miRNA-encoding sequence(s) with specified SEQ ID NO: for each, to be searched. Additional miRNA-encoding sequence(s) can be searched upon the payment of additional fees. It is believed that claims 62-66, 67 (in-part) and 68 encompass this first named invention and thus these claims can be searched with payment of a fee for the search of Groups II+ to the extent that they encompass SEQ ID NO: 53 (miRNA encoding sequence). Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched/examined. An exemplary election would be an miRNA-encoding sequence encompassing SEQ ID NO: 54 (first exemplary elected miRNA-encoding sequence).

The inventions listed as Groups I and II+ do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the special technical features of Group I include a modified nucleotide, not present in any of Groups II+; the special technical features of Groups II+ include an expression vector, not present in Group I.

No technical features are shared between the miRNA-encoding sequences of Groups II+ and, accordingly, these groups lack unity a priori.

Groups I and II+ share the technical features including: a nucleic acid encoding a miR-96, miR-182, or miR-183 sequence; and a method of treating or preventing an ophthalmological condition or an ear condition in a subject in need thereof, comprising administering to the subject an effective amount of a nucleic acid including a miR-96, miR-182, and/or miR-183 sequence. The special technical features of Groups II+ include: an expression vector comprising a polynucleotide encoding miR-96, miR-182, or miR-183 for expression in a mammalian cell; and a method of treating or preventing an ophthalmological condition or an ear condition in a subject in need thereof, comprising administering to the subject an effective amount of an expression vector encoding a miR-96, miR-182, and/or miR-183. However, these shared technical features are previously disclosed by WO 2014/100252 A1 to University of Washington Through Its Center for Commercialization (hereinafter 'Washington') in view of the article 'Roles of miR-182 in sensory organ development and cancer' by Wei et al. (hereinafter 'Wei').

Washington discloses a nucleic acid encoding (a vector (nucleic acid) encoding; paragraphs [00085], [00089], [000130]) a miR-96, miR-182, or miR-183 sequence (a miR-96, miR-182, or miR-183 sequence; paragraph [0035]); and a method of treating or preventing (therapeutic use of an miRNA for treatment (a method of treating or preventing); paragraph [00091]) an ophthalmological condition (age-related macular degeneration; age-related vision change (an ophthalmological condition); paragraph [00091]) or an ear condition (age-related hearing loss (an ear condition); paragraph [00091]) in a subject in need thereof (in a subject in need thereof; paragraphs [00091], [000124]), comprising administering to the subject an effective amount of a nucleic acid (comprising administering to the subject an effective amount of a vector (a nucleic acid); paragraphs [000124], [000130]) including a miR-96, miR-182, and/or miR-183 sequence (including a miR-96, miR-182, and/or miR-183 sequence; paragraphs [00035], [000124], [000130]); an expression vector comprising a polynucleotide (an expression vector comprising a polynucleotide; paragraphs [00085], [00089], [000130]) encoding miR-96, miR-182, or miR-183 (encoding miR-96, miR-182, or miR-183; paragraph [00035]) for expression in a mammalian cell (for expression in a mammalian cell; paragraphs [00085], [00089]); and a method of treating or preventing (therapeutic use of an miRNA for treatment (a method of treating or preventing); paragraph [00091]) an ophthalmological condition (age-related macular degeneration; age-related vision change (an ophthalmological condition); paragraph [00091]) or an ear condition (age-related hearing loss (an ear condition); paragraph [00091]) in a subject in need thereof (in a subject in need thereof; paragraphs [00091], [000124]), comprising administering to the subject an effective amount of an expression vector (comprising administering to the subject an effective amount of an expression vector; paragraphs [000124], [000130]) encoding a miR-96, miR-182, and/or miR-183 sequence (including (encoding) a miR-96, miR-182, and/or miR-183 sequence; paragraphs [00035], [000124], [000130]).

Washington does disclose limiting the encoded miRNA species to miR-96, miR-182, and/or miR-183.

Wei discloses wherein miR-182 is related to miR-183, and miR-96, sharing seed sequences (wherein miR-182 is related to miR-183, and miR-96, sharing seed sequences; page 2, first column, second paragraph - second column), and is abundantly expressed in the retina (wherein miR-182 is abundantly expressed in the retina; page 3, second column, third paragraph), and wherein miR-182 is expressed specifically in cochlear hair cells, and in the spiral ganglion (wherein miR-182 is expressed specifically in cochlear hair cells, and in the spiral ganglion; page 4, first column, second paragraph), where loss of expression correlated with progressive hair cell loss (where loss of expression correlated with progressive hair cell loss; page 4, first column, second paragraph).

It would have been obvious to a person of ordinary skill in the art at the time of the invention was made to have modified the disclosure of Washington to have specifically provided a vector encoding miR-182, or its related family members miR-96 and/or miR-183 for use in treatment of an ophthalmologic or ear condition, as disclosed by Washington, associated with reduced expression of these miRNA species in target tissues in the eye and ear, based on the disclosure of Wei, which indicates that a sufficient level of expression in said tissues is consistent with normal tissue.

Since none of the special technical features of the Groups I and II+ inventions is found in more than one of the inventions, and since all of the shared technical features are previously disclosed by a combination of the Washington and Wei references, unity of invention is lacking.