



US 20070224205A1

(19) **United States**(12) **Patent Application Publication**
Powell et al.(10) **Pub. No.: US 2007/0224205 A1**(43) **Pub. Date: Sep. 27, 2007**(54) **COMPOSITIONS THAT INCLUDE
HEMAGGLUTININ, METHODS OF MAKING
AND METHODS OF USE THEREOF**2006. Provisional application No. 60/856,451, filed
on Nov. 3, 2006.(76) Inventors: **Thomas J. Powell**, Madison, CT (US);
Albert E. Price, New Haven, CT (US);
Valerian Nakaar, Hamden, CT (US);
Robert K. Evans, Souderton, PA (US)**Publication Classification**(51) **Int. Cl.**
A61K 39/295 (2006.01)
(52) **U.S. Cl.** **424/159.1**; 424/209.1; 424/201.1;
424/272.1; 424/242.1

Correspondence Address:

**HAMILTON, BROOK, SMITH & REYNOLDS,
P.C.****530 VIRGINIA ROAD****P.O. BOX 9133****CONCORD, MA 01742-9133 (US)**(57) **ABSTRACT**(21) Appl. No.: **11/714,684**(22) Filed: **Mar. 6, 2007****Related U.S. Application Data**(60) Provisional application No. 60/779,854, filed on Mar.
7, 2006. Provisional application No. 60/784,497, filed
on Mar. 20, 2006. Provisional application No. 60/790,
457, filed on Apr. 7, 2006. Provisional application No.
60/814,292, filed on Jun. 16, 2006. Provisional appli-
cation No. 60/830,881, filed on Jul. 14, 2006. Provi-
sional application No. 60/838,007, filed on Aug. 16,

Compositions comprise a flagellin component or a Toll-like Receptor agonist component that is at least a portion of a flagellin or a Toll-like Receptor agonist, wherein the flagellin component or Toll-like Receptor agonist component includes at least one cysteine residue and whereby the flagellin component or Toll-like Receptor agonist component activates a Toll-like Receptor 5 or Toll-like Receptor. Compositions comprise a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, serine and histidine, whereby the flagellin component activates Toll-like Receptor 5. Compositions can further include an antigen, such as an influenza antigen, a flavivirus antigen, a pathogen-related antigen, a bacterial capsular antigen and a carrier protein. The compositions are used to stimulate an immune response and a protective immune response in a subject.

Design of Substrate Binding Domain as Influenza Vaccine (PR8)




Figure 1




Figure 2






Figure 3



Curve Fit Option - Fixed Weight Value: 1

Figure 4



4-P Fit: $y = (A - D) / (1 + (x/C)^B) + D$:

	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>R²</u>
○ Plot#1 (REF STF24xM2e@Experiment#1: Dilution vs ...	-11.075	0.501	10458.24	530.647	0.982
□ Plot#2 (Mal 1-2 Sup@Experiment#1: Dilution vs Mean...	9.743	1.712	8003.096	110.268	0.996
△ Plot#3 (SH 1-2 Sup@Experiment#1: Dilution vs Mean...	10.11	1.703	13620.02	128.77	0.997
◇ Plot#4 (Mal 1-1 Sup@Experiment#1: Dilution vs Mean...	10.599	1.047	5.86e5	1302.212	0.983

Curve Fit Option - Fixed Weight Value: 1

Figure 5




Figure 6




Figure 7




Figure 8A




Figure 8B




Figure 8C




Figure 9




Figure 10A




Figure 10B




Figure 11




Figure 12A




Figure 12B




Figure 13A




Figure 13B




Figure 13C




Figure 14




Figure 15




Figure 16




Figure 17

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSAKDDAAGQAIANRFTANIKG
LTQASRNANDGISIAQTTEGALNEINNNLQRVRELAVQSANSTNSQSDLDSIQAEITQRL
NEIDRVSGQTQFNGVVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
VKDTAVTTKAYANNGTTLDVSGLDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIG
GFTGADAAKNGDYEVNVATDGTVTLAAGATKTTMPAGATTKTEVQELKDTPAVVSADAKN
ALIAGGV DATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEATGAIKAKTTS
YTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGHDFKAQPELAEAAAKTTENPL
QKIDAALAQVDALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
QILQQAGTSVLAQANQVPQNVLSLLR

Figure 18





Figure 19



■ = STF2Δ construct

Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28

SEQ ID NO: 498 fljB/STF2 amino acid sequence (hinge region underlined)

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAIA NRFTANIKG
LTQASRNANDGISIAQTTEGALNEINNNLQRVRELAVQSANSTNSQSDLD SIQAEITQRL
NEIDRVSGQTQFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSL NVQKAYD
VKDTAVTTKAYANNGTTLDVSGLDDAAIKAATGGTNGTASVTGGAVKFDADNNKYFVTIG
GFTGADAAKNGDYEYVNVATDGTVTLAAGATKTTMPAGATTKTEVQELKDTPAVVSADAKN
ALIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEATGAIAKAKTTS
YTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGHDFKAQPELAEAAAKTTENPL
QKIDAALAQVDALRSDLGAVQNRFN SAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
QILQQAGTSVLAQANQVPQNVLSLLR

Figure 29

SEQ ID NO: 499 fljB/STF2 nucleic acid sequence (hinge region underlined)

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAACCTGAACAAA
TCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTTCTGGTCTGCGTATCAACAGC
GCGAAAGACGATGCGGCAGGTCAGGCGATTGCTAACCGTTTCACCGCGAACATCAAAGGT
CTGACTCAGGCTTCCCGTAACGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGC
GCGCTGAACGAAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTT CAGTCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCACCCAGCGCCTG
AACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACGGCGTGAAAGT CCTGGCGCAG
GACAACACCC TGACCATCCAGGTTGGCGCCAACGACGGTGAAACTATCGATATCGATCTG
AAGCAGATCAACTCTCAGACCC TGGGTCTGGACTCACTGAACGTGCAGAAAAGCGTATGAT
GTGAAAGATACAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGGTACGGCTTCT
GTAACCGGTGGTGCGGTTAAATTTGACGCAGATAATAACAAGTACTTTGTTACTATTGGT
GGCTTTACTGGTGCTGATGCCGCCAAAAATGGCGATTATGAAGTTAACGTTGCTACTGAC
GGTACAGTAACCCTTGCGGCTGGCGCAACTAAAACCACAATGCCTGCTGGTGCGACAAC T
AAAACAGAAGTACAGGAGTTAAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
GCCTTAATTGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTTGGTCAAAATG
TCTTATACCGATAAAAATGGTAAGACAATTGAAGGCGGTTATGCGCTTAAAGCTGGCGAT
AAGTATTACGCCGCAGATTACGATGAAGCGACAGGAGCAATTAAAGCTAAACTACAAGT
TATACTGCTGCTGACGGCACTACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGT
AAAACCGAAGTCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
GATTTCAAAGCACAAACCAGAGCTGGCGGAAGCAGCCGCTAAAACCACCGAAAACCCGCTG
CAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGCGCTCTGATCTGGGTGCGGTA
CAAAACCGTTTCAACTCTGCTATCACCAACCTGGGCAATACCGTAAACAATCTGTCTGAA
GCGCGTAGCCGTATCGAAGATTCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCG
CAGATTCTGCAGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
GTGCTGTCTCTGTTACGT

Figure 30

SEQ ID NO: 500 fljB/STF2Δ amino acid sequence

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAIANRFTANIKGLT
QASRNANDGISIAQTTEGALNEINNNLQRVRELAVQSANSTNSQSDLDSIQAEITQRLNEID
RVSGQTQFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSLNVHGAPVDPASPW
TENPLQKIDAALAQVDALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNM
SRAQILQQAGTSVLAQANQVPQNVLSLLR

Figure 31

SEQ ID NO: 501 fljB/STF2Δ nucleic acid sequence

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAACCTGAACAAATC
CCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTTCTGGTCTGCGTATCAACAGCGCGA
AAGACGATGCGGCAGGTCAGGCGATTGCTAACCGTTTCACCGCGAACATCAAAGGTCTGACT
CAGGCTTCCCGTAACGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAA
CGAAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTTCACTCTGCTAACAGCACCA
ACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCACCCAGCGCCTGAACGAAATCGAC
CGTGTATCCGGCCAGACTCAGTTCAACGGCGTGAAAGTCCTGGCGCAGGACAACACCCTGAC
CATCCAGGTTGGCGCCAACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTC
AGACCCTGGGTCTGGACTCACTGAACGTGCATGGAGCGCCGGTGGATCCTGCTAGCCCATGG
ACCGAAAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGCGCTCTGA
TCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAACCTGGGCAATACCGTAAACA
ATCTGTCTGAAGCGCGTAGCCGTATCGAAGATTCGACTACGCGACCGAAGTTTCCAACATG
TCTCGCGCGCAGATTTTGCAGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACCCAGTCCC
GCAGAACGTGCTGTCTCTGTTACGTG

Figure 32

SEQ ID NO: 502 E. coli fliC amino acid sequence (hinge region underlined)

MAQVINTNSLSLITQNNINKNQSA LSSSIERLSSGLRINSAKDDAAGQAIANRFTSNIKG
LTQAARNANDGISVAQTTEGALSEINN LQRIRELT VQASTGTNSDSDLDSIQDEIKSRL
DEIDRVSGQTQFNGVNV LAKDGSMKIQVGANDGQTITIDLKKIDSDTLGLNGFNVNGSGT
IANKAATISDLTAAKMDAATNTITTTNNALTASKALDQLKDGDVTIKADAAQTATVYTY
NASAGNFSLSNVSNNTSEKAGDVAASLLPPAGQTASGVYKAASGEVNFDVDANGKITIGG
QKAYLTSDGNLTTNDAGGATAATLDGLFKKAGDGQSIGFKKTASVTMGGTTYNFKTGADA
DAATANAGVSFTDTASKETVLNKVATAKQGKAAAADGDTSATITYKSGVQTYQAVFAAGD
GTASAKYADKADVSNATATYTDADGEMTTIGSYTTKYSIDANNGKVTVDSGTGTGKYAPK
VGAEVYVSANGTLTTDATSEGT VTKDPLKALDEAISSIDKFRSSLGAIQNRLDSAVTNLN
NTTTNLSEAQSRIQDADYATEVSNMSKAQIIQQAGNSVLAKANQVPQQVLSLLQG

Figure 33

SEQ ID NO: 503 E. coli fliC -nucleic acid sequence (hinge region underlined)

ATGGCACAAAGTCATTAATACCAACAGCCTCTCGCTGATCACTCAAAATAATATCAACAAG
AACCAGTCTGCGCTGTTCGAGTTCTATCGAGCGTCTGTCTTCTGGCTTGCGTATTAACAGC
GCGAAGGATGACGCCGAGGTGAGGCGATTGCTAACCGTTTACTTCTAACATTAAAGGC
CTGACTCAGGCTGCACGTAACGCCAACGACGGTATTTCCGTTGCGCAGACCACCGAAGGC
GCGCTGTCCGAAATCAACAACAACCTTACAGCGTATCCGTGAACTGACGGTTCAGGC'TTCT
ACCGGGACTAACTCCGATTGAGATCTGGACTCCATTGAGGACGAAATCAAATCCCGTCTG
GACGAAATTGACCGCGTATCTGGCCGAGCCAGTTCACGCGCGTGAACGTACTGGCGAAA
GACGGTTCAATGAAAATTGAGGTTGGTGCGAATGACGCGCCAGACTATCACGATTGATCTG
AAGAAAATTGACTCAGATACGCTGGGGCTGAATGGTTTAACTGAAATGGTTCCGGTACG
ATAGCCAATAAAGCGGCGACCATTAGCGACCTGACAGCAGCGAAAATGGATGCTGCAACT
AATACTATAACTACAACAAATAATGCGCTGACTGCATCAAAGGCGCTTGATCAACTGAAA
GATGGTGACACTGTTACTATCAAAGCAGATGCTGCTCAAACGCGGTTTATACATAC
AATGCATCAGCTGGTAACTTCTCACTCAGTAATGTATCGAATAATACTTCAGAAAAAGCA
GGTGATGTAGCAGCTAGCCTTCTCCGCGCGGCTGGGCAAACGCTAGTGGTGTTTATAAA
GCAGCAAGCGGTGAAGTGAAC'TTGATGTTGATGCGAATGGTAAAATCACAATCGGAGGA
CAGAAAGCATATTTAACTAGTGATGGTAACTTAACTACAAACGATGCTGGTGGTGCGACT
GCGGCTACGCTTGATGGTTTATTCAAGAAAGCTGGTGATGGTCAATCAATCGGGTTTAAAG
AAGACTGCATCAGTCACGATGGGGGGAACAAC'TTATAACTTTAAAACGGGTGCTGATGCT
GATGCTGCAACTGCTAACGAGGGGTATCGTTCAGTATACAGCTAGCAAAGAAACCGTT
TTAAATAAAGTGGCTACAGCTAAACAAGGCAAAGCAGCTGCAGCTGACGGTGATACATCC
GCAACAATTACCTATAAATCTGGCGTTGAGACGTATCAGGCTGTATTTGCCGAGGTGAC
GGTACTGCTAGCGCAAAATATGCCGATAAAGCTGACGTTTCTAATGCAACAGCAACATAC
ACTGATGCTGATGGTGAAATGACTACAATTGGTTCATACACCACGAAGTATTCAATCGAT
GCTAACAACGGCAAGGTAAC'TGTTGATTCTGGAAC'TGGTACGGGTAAATATGCGCCGAAA
GTAGGGGCTGAAGTATATGTTAGTGCTAATGGTACTTTAACAACAGATGCAACTAGCGAA
GGCACAGTAACAAAAGATCCACTGAAAGCTCTGGATGAAGCTATCAGCTCCATCGACAAA
TTCCGTTCTTCCCTGGGTGCTATCCAGAACCGTCTGGATTCCGCAGTCACCAACCTGAAC
AACACCACTACCAACCTGTCCGAAGCGCAGTCCCGTATTGAGGACGCCGACTATGCGACC
GAAGTGTCCAACATGTGAAAGCGCAGATCATTCAGCAGGCCGGTAACTCCGTGCTGGCA
AAAGCCAACCAGGTACCGCAGCAGGTTCTGTCTCTGCTGCAGGGTTAG

Figure 34

SEQ ID NO: 504 Salmonella muenchen flagellin fliC amino acid sequence (hinge region underlined)

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAIANRFTANIKGLT
QASRNANDGISIAQTTEGALNEINNNLQRVRELAVQSANGTNSQSDLSIQAEITQRLNEID
RVSGQTQFNGVKVLAQDNTLTIQVGANDGETIDIDLKEISSKTLGLDKLNVQDAYTPKETAV
TVDKTTYKNGTDTITAQSNTDIQTAIGGGATGVTGADIKFKDGQYYLDVKGGASAGVYKATY
DETTKKVNIDTTDKTPLATAEATAIRGTATITHNQIAEVTKEGVDTTVAQAALAAAGVTGAD
KDNTSLVKLSFEDKNGKVIDGGYAVKMGDDFYAATYDEKTGTITAKTTTYTDGAGVAQTGAV
KFGGANGKSEVVTATDGKTYLASDLDKHNFRTGGELKEVNTDKTENPLQKIDAALAQVDTLR
SDLGAVQNRFNSAITNLGNTVNNLSSARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQ
VPQNVLSLLR

Figure 35

SEQ ID NO: 505 Salmonella muenchen flagellin fliC nucleic acid sequence (hinge region underlined)

AATGGCACAAGTCATTAATACAAACAGCCTGTCGCTGTTGACCCAGAATAACCTGAACAAAT
CCCAGTCCGCTCTGGGCACCGCTATCGAGCGTCTGTCTTCCGGTCTGCGTATCAACAGCGCG
AAAGACGATGCGGCAGGTCAGGCGATTGCTAACCGTTTCACCGCGAACATCAAAGGTCTGAC
TCAGGCTTCCCGTAACGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGA
ACGAAATCAACAACAACCTGCAGCGTGTGCGTGAACTGGCGGTTCACTCTGCTAACGGTACT
AACTCCCAGTCTGACCTTGACTCTATCCAGGCTGAAATCAGGCGTCTGAACGAAATCGA
CCGTGTATCCGGTCAGACTCAGTTCAACGGCGTGAAAGTCTGGCGCAGGACAACACCCTGA
CCATCCAGGTTGGTGCCAACGACGGTGAAACTATTGATATTGATTTAAAAGAAATTAGCTCT
AAAACACTGGGACTTGATAAGCTTAATGTCCAGGATGCCTACACCCGAAAGAACTGCTGT
AACCGTTGATAAAACTACCTATAAAATGGTACAGATACTATTACAGCCCAGAGCAATACTG
ATATCCAACTGCAATTGGCGGTGGTGCAACGGGGTTACTGGGGCTGATATCAAATTTAAA
GATGGTCAATACTATTTAGATGTTAAAGGCGGTGCTTCTGCTGGTGTATATAAGCCACTTA
TGATGAAACTACAAAGAAAGTTAATATTGATACGACTGATAAACTCCGTTAGCAACTGCGG
AAGCTACAGCTATTCGGGGAACGGCCACTATAACCCACAACCAAATTGCTGAAGTAACAAAA
GAGGGTGTGATACGACCACAGTTGCGGCTCAACTTGCTGCTGCAGGGGTTACTGGTGCCGA
TAAGGACAATACTAGCCTTGTAAAACTATCGTTTGAGGATAAAAAACGGTAAGGTTATTGATG
GTGGCTATGCAGTGAAAATGGGCGACGATTTCTATGCCGCTACATATGATGAGAAAACAGGT
ACAATTACTGCTAAAAACAACCACTTATACAGATGGTGCTGGCGTTGCTCAAACCTGGAGCTGT
GAAATTTGGTGGCGCAAATGGTAAATCTGAAGTTGTTACTGCTACCGATGGTAAAACCTTACT
TAGCAAGCGACCTTGACAAACATAACTTCAGAACAGGCGGTGAGCTTAAAGAGGTTAATACA
GATAAGACTGAAAACCCACTGCAGAAAATTGATGCTGCCTTGGCACAGGTTGATACACTTCG
TTCTGACCTGGGTGCGGTACAGAACCGTTTCAACTCCGCTATCACCAACCTGGGCAATACCG
TAAATAACCTGTCTTCTGCCCCGTAGCCGTATCGAAGATTCCGACTACGCGACCGAAGTCTCC
AACATGTCTCGCGCGAGATTCTGCAGCAGGCCGGTACCTCCGTTCTGGCGCAGGCTAACCA
GGTTCGCAAAACGTCCTCTCTTTACTGCGTTAA

Figure 36

SEQ ID:506 Amino acid sequence of pMT/STF2 (Linker underlined)

MKLCILLAVVAFVGLSLGRSAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRI
NSAKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQRVREL
AVQSANSTNSQSDLDSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGAND
GETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTTLDVSGLDAAIK
AATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKNGDYEVNVATDGTVTLA
AGATKTTMPAGATTKTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYT
DKNGKTIEGGYALKAGDKYYAADYDEATGAIKAKTTSYTAADGTTKTAANQLGGVDG
KTEVV TIDGKTYNASKAAGHDFKAQPELAEEAAKT TENPLQKIDAALAQVDALRSDL
GAVQNRFN SAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQA
NQVPQNVLSLLRKGN SKLEGQLEFPRTSPVWWNSADIQHSGGRSSLEGPRFEGKPIP
NPLLGLDSTRTGHHHHHH

Figure 37

SEQ ID: 507 Amino acid sequence of pMT/STF2 (Linker underlined)

ATGAAGTTATGCATATTACTGGCCGTCGTGGCCTTTGTTGGCCTCTCGCTCGGGAGATCT
GCACAAGTAATCAACACTAACAGTCTGTCTGCTGCTGACCCAGAATAACCTGAACAAATCC
CAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTTCTGGTCTGCGTATCAACAGCGCG
AAAGACGATGCGGCAGGTCAGGCGATTGCTAACCGTTTCACCGCGAACATCAAAGTCTG
ACTCAGGCTTCCCGTAACGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCG
CTGAACGAAATCAACAACAACCTGCAGCGTGTGCGTGAACTGGCGGTTCACTCTGCTAAC
AGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCACCCAGCGCCTGAAC
GAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACGGCGTGAAAGTCCTGGCGCAGGAC
AACACCCTGACCATCCAGGTTGGCGCCAACGACGGTGAAACTATCGATATCGATCTGAAG
CAGATCAACTCTCAGACCCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTG
AAAGATACAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTATCG
GGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGGTACGGCTTCTGTA
ACCGGTGGTGCGGTTAAATTTGACGCAGATAATAACAAGTACTTTGTTACTATTGGTGGC
TTTACTGGTGCTGATGCCGCCAAAAATGGCGATTATGAAGTTAACGTTGCTACTGACGGT
ACAGTAACCCCTTGCGGCTGGCGCAACTAAAACCACAATGCCTGCTGGTGCGACAACATAA
ACAGAAGTACAGGAGTTAAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAATGCC
TTAATTGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTTGGTCAAAATGTCT
TATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTTATGCGCTTAAAGCTGGCGATAAG
TATTACGCCCGCAGATTACGATGAAGCGACAGGAGCAATTAAAGCTAAAACTACAAGTTAT
ACTGCTGCTGACGGCACTACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAA
ACCGAAGTCGTTACTATCGACGGTAAACCTACAATGCCAGCAAAGCCGCTGGTTCATGAT
TTCAAAGCACAACCAGAGCTGGCGGAAGCAGCCGCTAAAACCACCGAAAACCCGCTGCAG
AAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGCGCTCTGATCTGGGTGCGGTACAA
AACCGTTTCAACTCTGCTATCACCAACCTGGGCAATACCGTAAACAATCTGTCTGAAGCG
CGTAGCCGATATCGAAGATTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAG
ATTCTGCAGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACCAAGTCCCGCAGAACGTG
CTGTCTCTGTTACGTAAGGGCAATTCGAAGCTTGAAGGTCAATTGGAATTCCTAGGACT
AGTCCAGTGTGGTGGAATTCTGCAGATATCCAGCACAGTGGCGGCCGCTCGAGTCTAGAG
GGCCCCGCGGTTCAAGGTAAGCCTATCCCTAACCTCTCCTCGGTCTCGATTCTACGCGT
ACCGGTCATCATCACCATCACCAT

Figure 38



Figure 39




Figure 40




Figure 41A




Figure 41B




Figure 42

SEQ ID NO: 561

DTICIGYHANNSTDITVDTVLEKNVTVTHSVNLLED SHNGKLCRLKGIAPLQLGKCNIAGW
LLGNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEELREQLSSVSSFERFEIFPKE
SSWPNHNTNGVTAACSHEGKSSFYRNLLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHH
PPNSKEQQONLYQNENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLKPGDTII
FEANGNLIAPMYAFALS RGFSGIITSNASMHECNTK CQTPLGAINSSLPYQNIHPVTIG
ECPKYVRS AKLRMVTGLRNIPSIQSRGLFGAIAGFIEGGWTGMIDGWYGYHHQNEQGSY
AADQKSTQNAINGITNKVNTVIEKMNIQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTY
NAELLV LLENERTLDFHDSNVKNLYEKVKSQ LKNNAKEIGNGC FEFYHKCDNECMESVRN
GTYDYPKYSEESKLNREKVDGVKLESMGIYQ

Figure 43

SEQ ID NO: 562

GACCAGATCTGTATCGGTTATCATGCTAACAATTCTACTGAACAAGTAGATACTATCATG
GAGAAGAACGTTACAGTTACACATGCACAAGATATCCTGGAAAAGAAGCATAATGGAAAA
CTGTGTGACCTTGATGGTGTTAAACCACCTAATATTGCGTGACTGCTCAGTTGCTGGGTGG
TTGTTGGGGAATCCAATGTGCGACGAATTTATCAACGTTCCAGAATGGAGTTACATTGTT
GAAAAAGCTAACCCTGTTAACGACTTGTGTTACCCAGGCGATTTTAATGACTACGAGGAA
CTTAAGCATTTTGTGTCAAGAAATTAACCACTTCGAGAAAATTCAAATTATTCCAAAGTCA
TCTTGGTCCCTCCCATGAAGCATCCCTAGGAGTCTCTTCCGCTTGCCCTTACCAAGGCAAG
AGTTCCTTTTTTCGTAATGTCTGCTGGCTGATCAAAAAGAACTCCACCTATCCAACCTATA
AAGAGATCATACAACAACACAAATCAGGAGGATCTGCTAGTTCTGTGGGGCATTACCCAC
CCCAATGACGCAGCTGAGCAGACTAAATTGTACCAAACCCAACTACCTATATATCAGTT
GGTACCTCAACTCTTAACCAGCGACTAGTCCCCGTATTGCTACTAGGTCAAAGGTTAAT
GGTCAAAGTGAGCAATGGAGTTTTTCTGGACTATTTTGAAGCCCAACGATGCCATCAAC
TTCGAAAGTAATGGAAATTTTCATAGCCCCTGAGTACGCTTACAAAATCGTTAAAAAGGGT
GATTCCACTATCATGAAATCTGAACTGGAATACGGAACCTGTAACACCAAATGCCAGACG
CCAATGGGTGCCATCAACTCTTCTATGCCTTTTCAACAATTTCATCCTTTGACTATTGGT
GAATGCCCAAAGTACGTCAAATCTAACCGTTTGGTGTTGGCTACTGGTCTAAGGAACTCC
CCTCAGCGTGAAAGAAGAAGAAAGAAGAGGGGATTATTCGGTGCTATCGCTGGATTTATT
GAGGGAGGATGGCAGGGAATGGTCGATGGCTGGTATGGTTACCATCACTCAAATGAACAG
GGAAGTGGATACGCAGCTGATAAAGAATCTACTCAAAGGCTATCGACGGTGTTACAAAC
AAGGTCAATTCTATTATCGATAAGATGAATACACAGTTTGAGGCTGTTGGTAGAGAGTTC
AATAATCTTGAGAGAAGAATCGAAAACCTGAACAAGAAAATGGAAGACGGATTTTTTAGAT
GTATGGACTTACAATGCTGAGTTGTTGGTCTTGATGGAGAATGAACGAACGTTGGACTTC
CATGACTCCAATGTGAAGAACCTATATGACAAAGTGAGGCTGCAACTTAGAGACAACGCC
AAGGAATTGGGAAACGGGTGCTTCGAGTTTTTACCACAAATGCGACAACGAATGTATGGAA
TCAGTGAGAAACGGTACCTATGATTACCCCCAATATTCGAGGAGGCAAGACTGAAGAGA
GAAGAGATATCTGGTGTAAGTTGGAATCCATCGGTATTTATCAGATTCTATCTATATAT
TCTACCTAATAG

Figure 44

SEQ ID NO: 563 E. coli fliC Amino Acid sequence (without hinge region)

MAQVINTNSLSLITQNNINKNQSA LSSS IERLSSGLRINSAKDDAAGQA IANRFTSNIKG
LTQAARNANDGISVAQTTEGALSEINN NLQRIRELTVQASTGTNSDSDLDSIQDEIKSRL
DEIDRVSGQTQFNGVNV LAKDGSMKIQVGANDGQTITIDLKKIDSDLGTDPLKALDEA
ISSIDKFRSSLGAIQNRLDSAVTNLNN TTTNLSEAQSRIQDADYATEVSNMSKAQIIQQA
GNSVLAKANQVPQQVLSLLQG

Figure 45

SEQ ID: 585 Amino acid sequence of pMT/STF2Δ

MKLCILLAVVAFVGLSLGRSAQVINTNSLSLLTQNNLNKSQSALGTAIERLS
SGLRINSAKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEGALNEI
NNNLQRVRELAVQSANSTNSQSDLDISIQAETQRLNEIDRVSGQTQFNGVKV
LAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSLNVHGAPVDPASPWTEN
PLQKIDAALAQVDALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDY
ATEVSNMSRAQILQQAGTSVLAQANQVPQNVLSLLREFSRYP AQWRPLTRTG
HHHHHH

Figure 46

SEQ ID: 586 Nucleic acid sequence of pMT/STF2Δ

**ATGAAGTTATGCATATTACTGGCCGTCGTGGCCTTTGTTGGCCTCTCGCTCG
GGAGATCTGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAA
TAACCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCT
TCTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATTG
CTAACCGTTTCAACGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAACGC
TAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACGAAATC
AACAACAACCTGCAGCGTGTGCGTGAACTGGCGGTTTCAGTCTGCTAACAGCA
CCAACCTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCACCCAGCGCCT
GAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACGGCGTGAAAGTC
CTGGCGCAGGACAACACCCTGACCATCCAGGTTGGCGCCAACGACGGTGAAA
CTATCGATATCGATCTGAAGCAGATCAACTCTCAGACCCTGGGTCTGGACTC
ACTGAACGTGCATGGAGCGCCGGTGGATCCTGCTAGCCCATGGACCGAAAAC
CCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGCGCTCTG
ATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAACCTGGGCAA
TACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGATTCCGACTAC
GCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGCAGCAGGCCGGTA
CTTCCGTTCTGGCGCAGGCTAACCAGGTCCCGCAGAACGTGCTGTCTCTGTT
ACGTGAATTCTCTAGATATCCAGCACAGTGGCGGCCGCTCACGCGTACCGGT
CATCATCACCATCACCATTGA**

Figure 47

SEQ ID NO: 595 Salmonella muenchen fliC Amino Acid Sequence
(Hinge region deleted)

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSAKDDAAGQAIANRFTANIKG
LTQASRNANDGISIAQTTEGALNEINNNLQRVRELAVQSANGTNSQSDLDSIQAEITQRL
NEIDRVSGQTQFNGVKVLAQDNTLTIQVGANDGETIDIDLKEISSKTLDKHNFRGTGGELK
EVNTDKTENPLQKIDAALAQVDTLRSDLGAVQNRFNSAITNLGNTVNNLSSARSRIEDSD
YATEVSNMSRAQILQQAGTSVLAQANQVPQNVLSLLR

Figure 48

SEQ ID NO: 596 Salmonella Muenchen fliC Nucleic Acid Sequence
(Hinge region deleted)

ATGGCACAAGTCATTAATACAAACAGCCTGTCGCTGTTGACCCAGAATAACCTGAACAAA
TCCCAGTCCGCTCTGGGCACCGCTATCGAGCGTCTGTCTTCCGGTCTGCGTATCAACAGC
GCGAAAGACGATGCGGCAGGTCAGGCGATTGCTAACCGTTTCACCGCGAACATCAAAGGT
CTGACTCAGGCTTCCCGTAACGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGC
GCGCTGAACGAAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTT CAGTCTGCT
AACGGTACTAACTCCCAGTCTGACCTTGACTCTATCCAGGCTGAAATCACCCAGCGTCTG
AACGAAATCGACCGTGTATCCGGTCAGACTCAGTTCAACGGCGTGAAAGTCCTGGCGCAG
GACAACACCCCTGACCATCCAGGTTGGTGCCAACGACGGTGAACTATTGATATTGATTTA
AAAGAAATTAGCTCTAAAACACTGACAGATAAGACTGAAAACCCACTGCAGAAAATTGAT
GCTGCCTTGGCACAGGTTGATACACTTCGTTCTGACCTGGGTGCGGTACAGAACCGTTTC
AACTCCGCTATCACCAACCTGGGCAATACCGTAAATAACCTGTCTTCTGCCCCGTAGCCGT
ATCGAAGATTCCGACTACGCGACCGAAGTCTCCAACATGTCTCGCGCGCAGATTCTGCAG
CAGGCCGGTACCTCCGTTCTGGCGCAGGCTAACCAGGTTCCGCAAACGTCCTCTCTTTA
CTGCGTTAA

Figure 49




Figure 50




Figure 51




Figure 52




Figure 53




Figure 54




Figure 55




Figure 56




Figure 57

Rabbit IgG to M2e Individual Rabbits Day 49 (Day 7 Post Boost 2) 1:125 Dilution




Figure 58

STF2.4xM2e Rabbit Immunogenicity:
14 Day Post-prime




Figure 59




Figure 60




Figure 61




Figure 62A




Figure 62B




Figure 63




Figure 64A




Figure 64B




Figure 64C




Figure 64D




Figure 65




Figure 66




Figure 67




Figure 68




Figure 69




Figure 70




Figure 71




Figure 72

Flavivirus Envelope Protein Domain I/III junction
(domain I 280-297, domain III 298-406)

WN aa 281-307

WN LTSGHLKCRVKMEKLQLKGTTYGVCSK(SEQ ID NO: 691)

JE LTSGHLKCRLKMDKLALKGTTYGMCTE (SEQ ID NO: 692)

D1 IFAGHLKCRLKMDKLTLKGMSYVMCTG (SEQ ID NO: 639)

D2 LFTGHLKCRLRMDKLQLKGMSYSMCTG (SEQ ID NO: 694)

D3 IFAGHLKCRLKMDKLKLKGMSYAMCLN (SEQ ID NO: 695)

D4 MFAGHLKCKVRMEKLRIKGMSYTMCSG (SEQ ID NO: 696)

xxxGHLKCRxxMxKLxLKGxxYxxCxx (SEQ ID NO: 697)

GHLKCRMKLLKGYC (SEQ ID NO: 698)

Figure 73




Figure 74




Figure 75





Figure 76



Flagellin (1UCU)

Figure 77

Flagellin from *Pseudomonas aeruginosa* (accession number P21184)

```

      10      20      30      40 *      50      60
ALTVNTNIAS LNTQRNLNNS SASLNTSLQR LSTGSRINSA KDDAAGLQIA NRLTSQVNGL

      * 70      80      90      100      110      *  *
NVATKNANDG ISLAQTAEGA LQQSTNILQR MRDLSLQSAN GSNSDSERTA LNGEVKQLQK

      130      * 140      150      160      170      * 180
ELDRISNTTT FGGRKLLDGS FGVASFQVGS AANEIISVGI DEMSAESLNG TYFKADGGGA

      190      200      *  *      *      230      240
VTAATASGTV DIAIGITGGS AVNVKVDMMKG NETAEQAAAK IAAAVNDANV GIGAFSDGDT

      *  *      260      270      280      *290      * 300
ISYVSKAGKD GSGAITSavs GVVIADTGST GVGTAAGVTP SATAFAKTND TVAKIDISTA

      * 310      * 320      330      *      350      360
KGAQSAVLVI DEAIKQIDAQ RADLGAVQNR FDNTINNLKN IGENVSAARG RIEDTDFAAE

      * 370      380      390
TANLTKNQVL QQAGTAILAQ ANQLPQSVLS LLR

```

Figure 78

STF2 sequence

```

      10          *          30          40  *          50          *
MAQVINTNSL SLLTQNNLNK SQSALGTAIE RLSSGLRINS AKDDAAGQAI ANRFTANIKG

      70          80          90          100          110          120
LTQASRNAND GISIAQTTEG ALNEINNNLQ RVRELAVQSA NSTNSQSDLD SIQAEITQRL

      130          * 140          150          160 *          170          *180
NEIDRVSGQT QFNGVKVLAQ DNTLTIQVGA NDGETIDIDL KQINSQTLGL DSLNVQKAYD

      *          *          200          *          220          *          * 240
VKDTAVTTKA YANNGTTLDV SGLDDAAIKA ATGGTNGTAS VTGGAVKFDA DNNKYFVTIG

      *          260          270 *          280 *          *          *
GFTGADAAKN GDYEVNVATD GTVTLAAGAT KTTMPAGATT KTEVQELKDT PAVVSADAKN

      310          *          *          *          *340 *          350          * *360
ALIAGGV DAT DANGAELVKM SYTDKNGKTI EGGYALKAGD KYAAYDYDEA TGAIAKTTTS

      *          *          *          * 400          *          410          * 420
YTAADGTTKT AANQLGGVDG KTEVVTIDGK TYNASKAAGH DFKAQPELAE AAAKT TENPL

      *          430          440          450          460          470          480
QKIDAALAQV DALRSDLGAV QNRFNSAITN LGNTVNNLSE ARSRIEDSDY ATEVSNMSRA

      490
QILQQAGTSV LAQANQVPQN VLSLLR

```

Figure 79

Flagellin from *Listeria monocytogenes* (accession number Q92DW3)

```
*      10 *      *20      30      *      40      50      *      60
MKVNTNIISL KTQEYLRKNN EGMTQAQERL ASGKRINSSL DDAAGLAVVT RMNVKSTGLD

*      70      80      90      100      *      *
AASKNSSMGI DLLQTADSAL SSMSSILQRM RQLAVQSSNG SFSDEDRKQY TAEFGSLIKE

      130      *      140      150      *      160      *170      180
LDHVADTTNY NNIKLLDQTA TNAATQVSIQ ASDKANDLIN IDLFNAKGLS AGTITLGSGS

      190      200      210      220      230      *
TVAGYSALSV ADADSSQEAT EAIDELINNI SNGRALLGAG MSRLSYNVSN VNNQSIATKA

      250      *      *      270      280
SASSIEDADM AAEMSEMTKY KILTQTSISM LSQANQTPQM LTQLINS
```

Figure 80

COMPOSITIONS THAT INCLUDE HEMAGGLUTININ, METHODS OF MAKING AND METHODS OF USE THEREOF

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application Nos. 60/779,854, filed on Mar. 7, 2006; 60/784,497, filed on Mar. 20, 2006; 60/790,457, filed on Apr. 7, 2006; 60/814,292, filed on Jun. 16, 2006; 60/830,881, filed on Jul. 14, 2006; 60/838,007, filed on Aug. 16, 2006; and 60/856,451, filed on Nov. 3, 2006. The entire teachings of the above applications are incorporated herein by reference.

BACKGROUND OF THE INVENTION

[0002] Viral influenza infection can lead to disease. Strategies to prevent and manage disease associated with viral influenza infection can include vaccines with inactivated viruses and drugs. However, such strategies can be costly to maintain supply with demand and, thus, be limited in supply; may result in variable protection and less than satisfactory alleviation of symptoms, thereby ineffectively preventing or treating illness and, in some instances death, consequent to disease associated with viral influenza infection. Thus, there is a need to develop new, improved and effective methods of treatment for preventing and managing disease associated with viral influenza infection.

SUMMARY OF THE INVENTION

[0003] The present invention relates to compositions, such as compositions that stimulate a protective immune response, and methods of making proteins that stimulate a protective immune response in a subject.

[0004] In one embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transforming a nucleic acid sequence encoding the protein portion into a prokaryotic host cell; and culturing the prokaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0005] In another embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not a *Pichia pastoris* eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0006] In another embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not a *Drosophila melanogaster* eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0007] In a further embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not an insect eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0008] In a further embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not a stably transformed insect host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0009] In still another embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is neither a *Picia pastoris* eukaryotic host cell nor a stably transfected insect host cell;

and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0010] In yet another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein made by a method comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transforming a nucleic acid sequence encoding the portion into a prokaryotic host cell; and culturing the prokaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0011] In an additional embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein portion of a naturally occurring viral hemagglutinin, wherein the protein portion includes at least a portion of a globular head, and at least a portion of one secondary structure that causes the globular head to essentially retain its tertiary structure and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain.

[0012] In still another embodiment, the invention is a method of making a viral hemagglutinin protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the portion in a eukaryotic host cell, wherein the eukaryotic host cell is neither a *Pichia pastoris* eukaryotic host cell nor a stably transfected insect host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0013] In another embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby make a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; infecting a nucleic acid sequence encoding the protein portion into an insect cell host cell; and culturing the insect host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0014] In another embodiment, the invention is a method of making a viral hemagglutinin protein that stimulates a

protective immune response in a subject, comprising the steps of transforming a prokaryotic host cell with a nucleic acid sequence encoding at least one viral hemagglutinin that lacks a transmembrane domain and a cytoplasmic domain; and culturing the prokaryotic cell to thereby make the protein.

[0015] In a further embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein made by a method comprising the steps of transforming a prokaryotic host cell with a nucleic acid sequence encoding at least one viral hemagglutinin that lacks a transmembrane domain and a cytoplasmic domain; and culturing the prokaryotic host cell to thereby make the protein.

[0016] In another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein having at least one viral hemagglutinin that lacks a transmembrane domain and a cytoplasmic domain, wherein the protein was expressed in a prokaryotic cell.

[0017] In an additional embodiment, the invention is a composition comprising at least a portion of at least one pathogen-associated molecular pattern and a portion of a protein of a naturally occurring viral hemagglutinin, wherein the portion of the naturally occurring viral hemagglutinin includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the portion of the naturally occurring viral hemagglutinin lacks a membrane fusion domain, a transmembrane domain, and a cytoplasmic domain.

[0018] Another embodiment of the invention is a composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

[0019] In still another embodiment, the invention is a composition comprising a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor.

[0020] In an additional embodiment, the invention is a composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5.

[0021] In yet another embodiment, the invention is a composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0022] Another embodiment of the invention is a composition comprising a flagellin component that is at least a

portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0023] A further embodiment of the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

[0024] In still another embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5.

[0025] An additional embodiment of the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0026] In another embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0027] In yet another embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor agonist.

[0028] The methods and composition of the invention can be employed to stimulate an immune response, in particular, a protective immune response, in a subject. Advantages of the claimed invention include, for example, cost effective methods and compositions that can be produced in relatively large quantities for use in the prevention and treatment of disease associated with viral influenza infection. The claimed methods and compositions can be employed to prevent or treat viral influenza infection and, therefore, avoid serious illness and death consequent to viral influenza infection.

BRIEF DESCRIPTION OF THE FIGURES

[0029] FIG. 1 depicts a ribbon diagram of PR8 influenza A hemagglutinin (HA) crystal structure (1RU7). Schematic presentations from left to right are HA trimer, mono-

mer, the globular head domain and the three choices of vaccine candidates. Molecules or domains of interest are highlighted by dotted circle, such as the monomer in trimer structure and the globular head domain within the monomer structure. The domain boundaries of three vaccine candidates are marked by the cross signs. The residue numbers of each selection are also labeled in the individual vaccine presentation.

[0030] FIG. 2 depicts a hydrophobicity plot analysis of the B/Lee/40 HA (SEQ ID NO: 36) using ProtScale (<http://ca.expasy.org/tools>→Primary structure analysis→ProtScale) to confirm that the selected boundaries were in the hydrophilic regions of the protein.

[0031] FIG. 3 depicts a sandwich ELISA analysis of STF2.HA B strain proteins. ELISA plates were coated with antibody to flagellin, and the indicated proteins were incubated to allow capture. Proteins were detected using ferret antisera against B/Malaysia/2506/2004 and enzyme-labeled goat anti-ferret antibodies.

[0032] FIG. 4 depicts the TLR bioactivity of STF2.HA B strain proteins. HEK293 (TLR5+) cells were incubated with the indicated proteins overnight, and the cell culture supernatants were harvested and assessed for IL-8 by ELISA.

[0033] FIG. 5 depicts the TLR bioactivity of STF2.HA B strain proteins. HEK293 (TLR5+) cells were incubated with the indicated proteins overnight, and the cell culture supernatants were harvested and assessed for IL-8 by ELISA.

[0034] FIG. 6 depicts the dose-dependent antibody response to STF2.HA1-2(PR8) (SEQ ID NO: 90) in BALB/c mice. Mice (10/group) were immunized as indicated on days 0 and 14, and bled on days 12 and 21. Anti-HA IgG responses on Day 28 were examined by ELISA. Convalescent antisera (x) were included as a positive control. The data depict the mean±SD of 10 individual sera per group.

[0035] FIG. 7 depicts serum reactivity with influenza infected cells in vitro. The day 21 serum samples depicted in FIG. 6 were incubated with mock- and PR/8/34-infected MDCK cells. The data represent the OD₄₅₀ mean±SD of 10 individual sera/group.

[0036] FIG. 8A depicts survival of BALB/c mice immunized with STF2.HA1-2(PR8) (SEQ ID NO: 90). Mice from FIGS. 8 and 9 were challenged on day 28 with an LD₉₀ (8×10³ EID) of influenza A PR/8/34 administered intranasally (Day 0 post-challenge).

[0037] FIG. 8B depicts weights of BALB/c mice immunized with STF2.HA1-2(PR8) (SEQ ID NO: 90). Mice from FIGS. 6 and 7 were challenged on day 28 with an LD₉₀ (8×10³ EID) of influenza A PR/8/34 administered intranasally (Day 0 post-challenge). Graph reflects the average weight per group based on individual animals measured daily for 19 days.

[0038] FIG. 8C depicts clinical Score of BALB/c mice immunized with STF2.HA1-2(PR8) (SEQ ID: NO 90). Mice from FIGS. 6 and 7 were challenged on day 28 with an LD₉₀ (8×10³ EID) of influenza A PR/8/34 administered intranasally (Day 0 post-challenge). Graph reflects the average clinical score per group based on individual animals measured daily for 19 days. Clinical scores were assessed as

follows: 4 pts=healthy, 3 pts=reduced grooming, 2 pts=reduced physical activity and 1 pt=moribund.

[0039] FIG. 9 depicts neutralization activity of immune sera from BALB/c mice immunized with the indicated doses of STF2.HA1-2(PR8) (SEQ ID NO: 90) on days 0 and 14, and bled on day 21. The endpoint titer of each sample was calculated as the titer that inhibited virus-mediated lysis of the cells by at least 50%. Geometric mean titers $\pm$ SD were calculated for each dose group and plotted.

[0040] FIGS. 10A and 10B depict anti-HA antibody response of BALB/c mice immunized with 3 μ g of STF2.HA1-2(PR8) (SEQ ID NO: 90), or 3 or 0.3 μ g of STF2.HA1-2(VN) (SEQ ID NO: 95) s.c. on days 0 and 14. On day 21, sera were isolated and examined for reactivity with (FIG. 10A) HA purified from Influenza A/Viet Nam/1203/2004 (obtained from BEI Resources Cat# NR-660)) and (FIG. 10B) recombinant flagellin (STF2) (SEQ ID NO: 96) by ELISA.

[0041] FIG. 11 depicts a flow chart of the steps required to express pFastBac constructs using Bac-to-Bac Baculovirus Expression system.

[0042] FIGS. 12A and 12B depict an anti-HA specific IgG responses in BALB/c mice immunized with STF2.HA1-1 fusion proteins. Mice (10/group) were immunized with indicated STF2.HA1-1 fusion proteins on days 0 and 14, and bled on day 21. Anti-HA and anti-flagellin IgG responses were examined by ELISA. Convalescent antisera were included as a positive control. The data depict the mean $\pm$ SD of 10 individual sera per group.

[0043] FIG. 13A depicts survival of BALB/c mice immunized with recombinant STF2.HA1-1 proteins. On day 28, animals in FIG. 12 were challenged i.n. with an LD₉₀ (8×10^3 EID) of influenza A PR/8/34. The survival of individual mice was monitored for 21 days post challenge.

[0044] FIG. 13B depicts weight of BALB/c mice immunized with recombinant STF2.HA1-1 proteins. On day 28, animals in FIG. 12 were challenged i.n. with an LD₉₀ (8×10^3 EID) of influenza A PR/8/34. Graph reflects the average weight per group based on individual animals measured daily for 21 days.

[0045] FIG. 13C depicts clinical assessment of BALB/c mice immunized with recombinant STF2.HA1-1 proteins. On day 28, animals in FIG. 12 were challenged intranasally (i.n.) with an LD₉₀ (8×10^3 EID) of influenza A PR/8/34. Graph reflects the average weight per group based on individual animals measured daily for 21 days. Clinical scores were assessed as follows: 4 pts=healthy, 3 pts reduced grooming, 2 pts=reduced physical activity and 1 pt=moribund.

[0046] FIG. 14 depicts anti-HA antibody titers of mice following immunization with 10 μ g STF2.HA1-2(VN).

[0047] FIG. 15 depicts an anti-HA antibody titers of mice following immunization with 3 μ g STF2.HA1-2(VN).

[0048] FIG. 16 depicts an anti-HA antibody titers of mice following immunization with 1 μ g STF2.HA1-2(VN).

[0049] FIG. 17 depicts a survival of mice following challenge with 10 LD₉₀ of influenza A/Viet Nam/1203/04

[0050] FIG. 18 depicts the amino acid sequence of full-length flagellin (STF2) highlighting important domains and residues. Gray=experimentally-defined TLR5 binding site (Smith, et al., 2003); K=lysine residues; underline=flagellin sequences corresponding to STF2 Δ construct (linker not shown).

[0051] FIG. 19 depicts a space-filling model of flagellin (STF2) with the TLR5 activation site (left panel) and lysine residues (right panel) highlighted in gray.

[0052] FIG. 20 depicts a space-filling model of flagellin (STF2) with residues retained in STF2 Δ highlighted in black.

[0053] FIG. 21 depicts a schematic depiction of the two-step PCR cloning of pMT/STF2 Δ .

[0054] FIG. 22 depicts recognition of flagellin epitopes of STF Δ proteins in ELISA.

[0055] FIG. 23 depicts TLR5-dependent secretion of IL-8 following stimulation of HEK293 cells with the indicated STF2 Δ proteins.

[0056] FIG. 24 depicts the S200 chromatogram of STF2 Δ .HingeCys:CysH1C1. Void=column void volume; conjugate=elution peak of protein: peptide conjugate; A260, A280 and conductivity are indicated.

[0057] FIG. 25 depicts TLR5-dependent secretion of IL-8 following stimulation of RAWhTLR5 cells (closed symbols) or TLR5-negative RAW cells (open symbols) with STF2 Δ .HingeCys (squares), STF2 Δ .HingeCys:CysH1C1 conjugate (circles), or STF2.OVA (triangles).

[0058] FIG. 26 depicts the S200 chromatogram of STF2.4 $\times$ H1C1. Void=column void volume; conjugate=elution peak of protein: peptide conjugate; A260, A280 and conductivity are indicated.

[0059] FIG. 27 depicts an anti-H1C1 antibody response of BALB/c mice immunized with native H1C1 peptide or Pam3Cys.H1C1 peptide.

[0060] FIG. 28 depicts a survival plot of BALB/c mice immunized with native H1C1 peptide or Pam3Cys.H1C1 peptide then challenged with an LD₉₀ of influenza A/Puerto Rico/8/34 virus.

[0061] FIG. 29 depicts the amino acid sequence of *Salmonella typhimurium* flagellin type 2 (fljB/STF2) with the hinge region underlined (SEQ ID NO: 498).

[0062] FIG. 30 depicts the nucleic acid sequence (SEQ ID NO: 499) encoding SEQ ID NO: 498. The nucleic acid sequence encoding the hinge region is underlined.

[0063] FIG. 31 depicts the amino acid sequence of fljB/STF2 without the hinge region (also referred to herein as "fljB/STF2 Δ " or "STF2 Δ ") (SEQ ID NO: 500).

[0064] FIG. 32 depicts the nucleic acid sequence (SEQ ID NO: 501) encoding SEQ ID NO: 500.

[0065] FIG. 33 depicts the amino acid sequence of *E. coli* flagellin fliC (also referred to herein as "*E. coli* fliC") with the hinge region underlined (SEQ ID NO: 502).

[0066] FIG. 34 depicts the nucleic acid sequence (SEQ ID NO: 503) encoding SEQ ID NO: 502. The nucleic acid sequence encoding the hinge region is underlined.

[0067] FIG. 35 depicts the amino acid sequence of *S. muenchen* flagellin fliC (also referred to herein as “*S. muenchen* fliC”) with the hinge region underlined (SEQ ID NO: 504).

[0068] FIG. 36 depicts the nucleic acid sequence (SEQ ID NO: 505) encoding SEQ ID NO:504. The nucleic acid sequence encoding the hinge region is underlined.

[0069] FIG. 37 depicts the amino acid sequence of pMT/STF2. The linker is underlined and the sequence of the BiP secretion signal is bolded (SEQ ID NO: 506).

[0070] FIG. 38 depicts the nucleic acid sequence (SEQ ID NO: 507) of SEQ ID NO: 506. The nucleic acid sequence encoding the linker is underlined and the nucleic acid sequence encoding the BiP sequence is bolded.

[0071] FIG. 39 depicts a Pam3Cys.M2e fusion protein. The amino acid sequence (SEQ ID NO: 510) of M2e is shown in bold type.

[0072] FIG. 40 depicts the activation of an antigen-presenting cell (APC) by Toll-like Receptor (TLR) signaling.

[0073] FIGS. 41A and 41B depict plasmid constructs to express an amino-terminus of an M2 (e.g., SEQ ID NOS: 510, 554) of H1 and H5 (SEQ ID NO: 536) influenza A viral isolates. pMT: metallothionein promoter-based expression vector. BiP: secretion signal sequence of immunoglobulin-binding protein. STF2: full-length flagellin of *S. typhimurium*. STF2Δ: hinge region-deleted STF2. MCS: multiple cloning site.

[0074] FIG. 42 depicts plasmid constructs designed to express HA of H1 and H5 influenza A virus isolates. AOX1: AOX1 promoter of pPICZα expression vector (Invitrogen Corporation, Carlsbad, Calif.). α: secretion signal sequence of yeast. STF2: full-length flagellin of *S. typhimurium*. STF2Δ: hinge region-deleted STF2. MCS: multiple cloning site.

[0075] FIG. 43 depicts the amino acid sequence (SEQ ID NO: 561) of HA (PR8).

[0076] FIG. 44 depicts the nucleic acid sequence (SEQ ID NO: 562) encoding SEQ ID NO: 561.

[0077] FIG. 45 depicts the amino acid sequence (SEQ ID NO: 563) of *E. coli* fliC without the hinge region.

[0078] FIG. 46 depicts the amino acid sequence of pMT/STF2Δ (SEQ ID NO: 585). The linker sequence is underlined and the BiP secretion signal is bolded.

[0079] FIG. 47 depicts the nucleic acid sequence (SEQ ID NO: 586) encoding SEQ ID NO: 585. The nucleic acid sequence encoding the linker is underlined and the nucleic acid sequence encoding the BiP secretion signal is bolded.

[0080] FIG. 48 depicts the amino acid sequence (SEQ ID NO: 595) of the *Salmonella muenchen* fliC without the hinge region, which is also referred to herein as “*S. muenchen* fliCA.”

[0081] FIG. 49 depicts the nucleic acid sequence of *Salmonella muenchen* fliC (SEQ ID NO: 596) encoding SEQ ID NO: 595.

[0082] FIG. 50 depicts IL-8 secretion following stimulation of TLR5+ cells.

[0083] FIG. 51 depicts TNF secretion following stimulation of TLR2+ cells.

[0084] FIG. 52 depicts M2e-specific IgG.

[0085] FIG. 53 depicts the OVA-specific IgG.

[0086] FIG. 54 depicts the M2e-specific IgG serum titers.

[0087] FIG. 55 depicts the M2e-specific serum IgG titer post-boost.

[0088] FIG. 56 depicts the Pam3Cys.M2e dose response.

[0089] FIG. 57 depicts the M2e-specific serum IgG titer.

[0090] FIG. 58 depicts the rabbit IgG response to M2e.

[0091] FIG. 59 depicts the immunogenicity of STF2.4×M2e in a rabbit 14 days post-prime.

[0092] FIG. 60 depicts the survival of BALB1c mice following challenge. Mice were immunized on days 0 and 14 as indicated in the legend and challenged on day 28 by intranasal administration of an LD₅₀ of influenza A/Puerto Rico/8/34 virus (Day 0 post-challenge). Mice were monitored for survival for 21 days post-challenge.

[0093] FIG. 61 depicts fusion constructs in a pET24 vector. T7:T7 promoter; lacO: lac operator; STF2: *Salmonella typhimurium* flagellin; STF2Δ=STF2 with the hinge region deleted; EIII+ is domain III of a West Nile envelope protein with 6 amino acids of domain I amino acid.

[0094] FIGS. 62A and 62B depict TLR5 bioactivity of STF2.EIII+ (SEQ ID NOS: 657, 658) and STF2ΔEIII+ (SEQ ID NOS: 673, 674) fusion proteins. Serial dilutions of purified proteins were added to HEK293 (TLR5+) cells overnight and IL-8 content of the supernatants measured by ELISA. Purified STF2.OVA was used as a positive control (FIG. 129A). The TLR2 agonist Pam3CSK4 was used as a negative control (FIG. 129B).

[0095] FIG. 63 depicts STF2Δ.EIII+ antigenic epitopes assessed by ELISA. Plates were coated with full-length WNE (open bars) (SEQ ID NO: 642) or STF2Δ.EIII+ (SEQ ID NOS: 673,674) and probed with the indicated antibodies (mAb). Poly=polyclonal antiserum to WNE; 3D9 through 7H2=neutralizing monoclonal antibodies to WNE epitopes; anti-flagellin=monoclonal antibody to flagellin.

[0096] FIGS. 64A, 64B, 64C and 64D depict reactivity of STF2.E (SEQ ID NOS: 761, 762); STF2.EIII+ (SEQ ID NOS:657, 658) and STF2Δ.EIII+ (SEQ ID NOS:673,674) fusion proteins with antibodies to WNE and flagellin. Plates were coated with fusion proteins, blocked and incubated with antibodies to WNE or flagellin. Antibody reactivity was detected following incubation with HRP-labeled species specific IgG. Plates were developed in the presence of TMB substrate and O.D.450/650 using a TECAN plate reader and Magellian software.

[0097] FIG. 65 depicts IgG serum following injection with fusion proteins. Mice were immunized with either PBS, *Drosophila* conditioned medium containing STF2.E (CM, positive control), 25 μg of STF2Δ.EIII+ (SEQ ID NOS:673, 674) i.p., 25 μg STF2Δ.EIII+ s.c., 25 μg STF2.EIII+ (SEQ ID NOS:659,658) i.p., 25 μg STF2.EIII+ (SEQ ID NOS:657, 658) or 25 μg STF2.E (SEQ ID NOS: 761, 762). On day 35, immunized animals were challenged with WNV. Sera from individual mice (day 35) were characterized by direct

ELISA to determine IgG levels. Purified WNV-E protein (SEQ ID NO:642) was used as the antigen in this assay. This antigen (60) was produced in *Drosophila* as a his-tagged protein.

[0098] FIG. 66 depicts STF2Δ.EIII+ (SEQ ID NOS: 673, 674) and STF2Δ.EIII+ (SEQ ID NOS: 657, 658) protective immunity to WNV viral challenge. Mice were immunized and challenged with a lethal dose of WNV strain 2741 on day 35. Survival was monitored for 21 days.

[0099] FIG. 67 depicts IgG sera titers following immunization with fusion proteins. STF2Δ.EIII+ proteins induce WNV-specific IgG antibodies. Mice were immunized s.c. on days 0, 14 and 28 with PBS alone or about 25 μg of STF2Δ.EIII+ (SEQ ID NOS: 673, 674) (045 [positive control]), STF2Δ.EIII+ (067, trimer), STF2Δ.EIII+ (070, monomer) or STF2Δ.EIII+ (SEQ ID NOS: 675, 676) (069). On day 35 sera from individual mice were characterized by direct ELISA to determine IgG levels. Purified WNV-E protein (060, produced in *Drosophila* as a his-tagged protein) was used as the antigen in this assay.

[0100] FIG. 68 depicts STF2Δ.EIII+ (SEQ ID NOS: 673, 674) and STF2Δ.EIII+ (SEQ ID NOS: 675, 673) protective immunity in mice from WNV lethal challenge. On day 38 following immunization with fusion proteins, all groups were challenged with a lethal dose of WNV strain 2741 and survival was monitored for 21 days. Survival for each group (10 mice/group) is indicated as a percentage.

[0101] FIG. 69 depicts competition assays. Serial dilutions (five fold starting at 1:25) of antisera from immunized animals were incubated with biotinylated WNE protein (SEQ ID NO: 642) and then added to the wells of ELISA plates coated with mAb 7H2 at about 2 mg/ml. Wells were developed using avidin-HRP to determine inhibition of West Nile protein binding as a results of competition with mAb 7H2.

[0102] FIG. 70 depicts epitope mapping of the antibody response induced by STF2Δ.EIII+ (SEQ ID NOS: 675,676) fusion proteins. Immune sera from animals immunized with indicated STF2Δ-fusion proteins (E2-21, E27-E52, FIG. 142) were examined for the ability to recognize overlapping peptides corresponding to the junction of domains I and III of the WNV envelope protein.

[0103] FIG. 71 depicts epitope mapping of the antibody response induced by STFΔ.EIII+ (SEQ ID NOS: 675, 676) E-21 (envelope protein) epitope fusion proteins. Immune sera from animals immunized with the indicated STF2Δ-fusion proteins (E2-21, E2-21-1(S,C), E2-21-2(C,S), E2-21-2(C,S) and E2-21-4 through E2-21-24, see FIG. 139) were evaluated to identify the residues defining the E-21 epitope of West Nile envelope protein. Data reflects the response of sera to E-21 following the substitution of cysteine with serine (indicated by C,S); and the sequential replacement of amino acids with alanine. The peptides tested are listed in FIG. 139.

[0104] FIG. 72 depicts Pam3Cys.WNV001 (SEQ ID NO: 771) inducing EIII specific IgG antibodies. Mice were immunized s.c. on days 0, 14 and 28 with PBS alone, 22 mg of unmodified WNV001 (SEQ ID NO: 771) or 30 μg of Pam3Cys.WNV001. On day 35 sera from individual mice were characterized by direct ELISA to determine IgG levels to synthetic WNV001 peptide.

[0105] FIG. 73 depicts the amino acid sequences (SEQ ID NOS: 691-698) of the EI/EIII junction for West Nile, Japanese encephalitis and Dengue (serotypes 1 through 4) viruses. The West Nile epitope identified using antisera from STF2Δ.EIII+ immunized animals is underlined. This sequence corresponds to peptide E2-21 (SEQ ID NO: 728).

[0106] FIG. 74 depicts a tripalmitoylated peptide.

[0107] FIG. 75 depicts the D1 domain, D2 domain, TLR5 activation domain and hypervariable (D3 domain) of flagellin.

[0108] FIG. 76 depicts the D1 domain, D2 domain, TLR5 activation domain and hypervariable (D3 domain) of flagellin (Yonekura, et al. Nature 424, 643-650 (2003)).

[0109] FIG. 77 depicts the D0, D1, D2 and D3 domains of flagellin and regions in the D2 and D3 domains suitable for insertion or substitution with antigens (e.g., maturational cleavage site, HA, M2e).

[0110] FIG. 78 depicts the amino acid sequence of *Pseudomonas aeruginosa* flagellin (SEQ ID NO: 815). Lysine residues are indicated by the asterisks.

[0111] FIG. 79 depicts the amino acid sequence of a *S. typhimurium* flagellin (SEQ ID NO: 816). Lysine residues are indicated by the asterisks.

[0112] FIG. 80 depicts the amino acid sequence (SEQ ID NO: 820) of *Listeria monocytogenes* flagellin (GenBank Accession No: Q92DW3). Lysine residues are indicated by the asterisks.

DETAILED DESCRIPTION OF THE INVENTION

[0113] The features and other details of the invention, either as steps of the invention or as combinations of parts of the invention, will now be more particularly described and pointed out in the claims. It will be understood that the particular embodiments of the invention are shown by way of illustration and not as limitations of the invention. The principle features of this invention can be employed in various embodiments without departing from the scope of the invention.

[0114] The invention generally is directed to methods of making compositions, and to compositions, that stimulate an immune response in a subject, such as a protective immune response, and methods of treatment, such as by administering the composition to a subject.

[0115] "Stimulating an immune response," as used herein, refers to the generation of antibodies and/or T-cells to at least a portion of an antigen, such as the protein portions of hemagglutinin (HA) (e.g., HA1-1, HA1-2 proteins) described herein. The antibodies and/or T-cells can be generated to at least a portion of an influenza viral protein (e.g., HA and M2 proteins of influenza A, B and/or C).

[0116] Stimulating an immune response in a subject can include the production of humoral and/or cellular immune responses that are reactive against the antigen, such as a viral protein, in particular, an influenza viral protein.

[0117] The compositions of the invention for use in methods to stimulate immune responses in subjects, can be evaluated for the ability to stimulate an immune response in

a subject using well-established methods. Exemplary methods to determine whether the compositions of the invention stimulate an immune response in a subject, include measuring the production of antibodies specific to the antigen (e.g., IgG antibodies) by a suitable technique such as, ELISA assays; the potential to induce antibody-dependent enhancement (ADE) of a secondary infection; macrophage-like assays; neutralization assessed by using the Plaque Reduction Neutralization Test (PRNT₈₀); and the ability to generate serum antibodies in non-human models (e.g., mice, rabbits, monkeys) (Putnak, et al., *Vaccine* 23:4442-4452 (2005)).

[0118] "Stimulates a protective immune response," as used herein, means administration of the compositions of the invention, such as hemagglutinin (HA) proteins (e.g., HA1-1, HA1-2 proteins described herein), results in production of antibodies to the protein to thereby cause a subject to survive challenge by an otherwise lethal dose of a viral protein, such as viral HA. Techniques to determine a lethal dose of a virus (e.g., an influenza virus) are known to one of skill in the art (see, for example, WHO/CDS/CSR/NCS2002.5 "WHO Manual on Animal Influenza Diagnosis and Surveillance" World Health Organization, Dept of Communicable Disease Surveillance and Response, WHO Global Influenza Programme; Harmon, M. W., et al., *J. Clin. Microbiol.* 26:333-337 (1988); Reed, L. J., et al., *Am. J. Hyg.* 27:493-497 (1938); Rose, T., et al., *J. Clin. Microbiol.* 37:937-943 (1999); Walls, H. H. et al., *J. Clin. Microbiol.* 23:240-245 (1986); Current Protocols in Immunology, 19.11.1-19.11.32, Cottey, R., et al., John Wiley & Sons, Inc (2001)). Exemplary techniques for determining a lethal dose can include administration of varying doses of virus and a determination of the percent of subjects that survive following administration of the dose of virus (e.g., LD₁₀, LD₂₀, LD₄₀, LD₅₀, LD₆₀, LD₇₀, LD₈₀, LD₉₀). For example, a lethal dose of a virus that results in the death of 50% of a population of subjects is referred to as an "LD₅₀"; a lethal dose of a virus that results in the death of 80% of a population of subjects is referred to herein as "LD₈₀"; a lethal dose of a virus that results in death of 90% of a population of subjects is referred to herein as "LD₉₀."

[0119] For example, determination of the LD₉₀ can be conducted in subjects (e.g., mice) by administering intranasally varying doses (e.g., dilutions, such as log and half-log dilutions of 8×10³ egg-infectious doses (EID)) followed by an assessment of the survival of the subjects about 14 days to about 21 days after infection with the virus. Protective immunity can be assessed by physical appearance of the subject, general demeanor (active), weight (initial loss of weight followed by return to a weight about the weight of the subject prior to infection with the virus) and survival after about 14 to about 21 days following infection with the virus.

[0120] Assessment of stimulation of protective immunity can also be made by employing assays that assess the ability of the antibodies produced in response to the compositions of the invention (e.g., a portion of the protein of the naturally occurring virus, such as a protein portion of hemagglutinin) to neutralize binding of the viral protein (e.g., hemagglutinin protein) to a host cell (see, for example, Current Protocols in Immunology, 19.11.1-19.11.32, Cottey, R., et al., John Wiley & Sons, Inc (2001)). Assessment of stimulation of protective immunity can also be made by employing assays that

measure the ability of antibodies to inhibit hemagglutinin binding (see, for example, Burnett, F. M., et al., *J. exp. Biol. Med. Sci.* 25:227-233 (1947); Salk, J. E. *J. Immunol.* 49:87-98 (1944); Current Protocols in Immunology, 19.11.1-19.11.32, Cottey, R., et al., John Wiley & Sons, Inc (2001)).

[0121] It is believed that inhibition of hemagglutinin binding is indicative of the ability of antibodies, formed from the compositions and by the methods of the invention, to neutralize the sialic acid binding sites of the naturally occurring viral hemagglutinin ("neutralization of HA binding") and, thereby, prevent infection of the host cell as a consequence of stimulating a protective immune response. Inhibition or neutralization of hemagglutinin binding is believed to correlate with an ability of an immune response to protect against a lethal dose of virus.

[0122] Neutralization of HA binding can be assessed by in vitro assays (See, for example, Current Protocols in Immunology 19.11.1-19.11.32, Cottey, R., et al., Suppl. 42, John Wiley & Sons, Inc. (2001) and WHO Manual on Animal Influenza Diagnosis and Surveillance, Webster, R., et al., pages 28-36, 48-54, 82-92 (2002)). Exemplary viral neutralization assays rely on the ability of serum to specifically bind and prevent replication of influenza virus in culture, such as in the Madin-Darby Canine Kidney (MDCK) cell line. Briefly, cells are cultured in 96 well plates in the presence of a previously titrated virus and the cytopathic effect of the replicating virus is observed under a microscope. To test serum, serial dilutions of the serum are prepared and preincubated with the viral stock for 2 hours at 37° C. prior to infecting the MDCK cells. The mixture is incubated for an additional 2 hours after which the virus/serum mixture is removed and replaced with fresh media. The cells are grown for 4 days. Wells are scored as positive for viral growth if at least about 50% of the cells are dead in at least about half of the wells for a given serum dilution. The reciprocal of the highest dilution of serum which protects at least about half of the cells from death, in at least about half of the wells, is considered the neutralization titer.

[0123] Alternatively, a micro-neutralization in vitro assay can be performed to assess neutralization of HA binding. For example, serum is diluted and preincubated with a known titer of virus and mixed with MDCK cells, as described above. After 2 days of incubation, cells are washed and fixed with acetone. The plates are developed as an ELISA using a monoclonal antibody to the influenza nuclear antigen NP. A microneutralization titer is determined as the reciprocal of the highest dilution which yields less than about 50% of the anti-NP reading of the virus-only control wells.

[0124] The Hemagglutination Inhibition (HAI) assay is based on the HA antigen on the surface of the influenza virus agglutinating red blood cells (RBC) and preventing red blood cells from precipitating. Antibodies that specifically bind the sialic acid-binding regions of HA prevent agglutination allowing precipitation. The assay is performed in 96 well V bottom plates with fresh chicken RBC. A stock of viral antigen is titrated so that about a 4-fold excess of antigen is present relative to the minimum amount needed to prevent precipitation. The test serum, which can be from several species including mouse, ferret, poultry or human, is heated to about 56° C. to inactivate complement. Serial 2-fold dilutions of the inactivated serum are performed and mixed with the stock HA. After about 30 minutes at room

temperature, the RBCs are added and the plate is incubated for about 30 to about 45 minutes. Results are scored by observations: agglutination results in cloudy wells while inhibition results in a “button” of red cells precipitated at the bottom of the well. Controls include RBC with no HA, which forms a button, and HA and RBC with no serum, which remains cloudy. The HAI titer of a particular serum sample is the reciprocal of the last dilution which prevents agglutination (i.e., forms a button). For example, if about a 1:128 dilution reads as a button but the 1:256 dilution does not, the HAI titer is about 128.

[0125] In one embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby make a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain. The nucleic acid sequence encoding the protein portion is transformed into a prokaryotic cell and the prokaryotic host cell is cultured to thereby make the protein that stimulates a protective immune response in a subject.

[0126] “A portion of a protein,” or “protein portion,” as used herein in reference to a naturally occurring viral hemagglutinin, refers to any part of the naturally occurring viral hemagglutinin that is less than the entire naturally occurring viral hemagglutinin. “Naturally occurring viral hemagglutinin,” as used herein, refers to the entire viral hemagglutinin, as it occurs in nature.

[0127] The protein portion can further lack a signal sequence. The protein portion can further include a sialic acid binding site.

[0128] Portions of a protein of a naturally occurring viral hemagglutinin for use in the compositions and methods of the invention can be a portion of Orthomyxoviridae (influenza A, B, C), Paramyxovirus (parainfluenza, respiratory syncytial virus, Newcastle disease virus, Nipah, Measles, canine distemper, Sendai virus), Reoviridae (rotavirus), Parvoviridae (human parvovirus, porcine parvovirus), Orthopoxvirus (Monkeypox virus, Ectromelia virus), Flaviviridae (West Nile, Japanese Encephalitis, St. Louis, Murray Valley, Kunjin), Avipoxvirus (Chicken fowlpox), Nipah virus (Guillaume V., et al., *J. Virol.*, 80:7546-54 (2006)); Canine distemper virus (Singethan K., et al., *J. Gen. Virol.*, 87:1635-42 (2006)); Newcastle disease virus, (de Leeuw O. S., et al., *J. Gen. Virol.*, 86:1759-69 (2005); and Melanson V. R., et al., *J. Virol.*, 78:13053-61 (2004); Deng R., et al., *Virology*, 204:17-26; (1994)), Measles (Masse N., et al., *J. Virol.*, 78:9051-63 (2004)), Sendai virus (Tomasi M., et al., *FEBS Lett.*, 11:56-60 (2003)), Human parainfluenza (Porotto M., et al., *J. Virol.*, 79:2383-92 (2005); Tsurudome M., et al., *Virology*, 213:190-203 (1995); Bousse T., et al., *Virology*, 209:654-7 (1995); and Takimoto T., et al., *J. Virol.*, 66:7597-600 (1992)).

[0129] Portions of a viral hemagglutinin (“protein portions”) (e.g., an influenza A, an influenza B and an influenza C viral hemagglutinin) can include at least one member selected from the group consisting of protein portions referred to herein as “HA1-1,” “HA1-2” and “HA1-3.”

[0130] “HA1-1,” as used herein, refers to a protein portion of a viral hemagglutinin that includes at least about one β -sandwich that includes the substrate binding site, which includes at least about two β -sheets, at least about two to about three short α -helices, at least one small β -sheet and at least one additional small β -sandwich at the bottom of the molecule and at least about four disulfide bonds. The β -sandwich that includes the substrate binding site of the HA1-1 includes at least about four β -strands as the top sheet and at least about three to about four β -strands as the bottom sheet. At least about one α -helix of the HA1-1 portion is located by the side of β -sandwich that includes the substrate binding site and at least about one to about two are located at the bottom of the β -sandwich that includes the substrate binding site. The small β -sandwich of the HA1-1 can include at least about two to about three β -strands in each β -sheet; or about three to about four β -strands. Exemplary HA1-1 protein portions include SEQ ID NOS: 8, 11, 14, 17, 20, 38, 40, 45, 47, 49, 179, 180, 181 and 182.

[0131] “HA1-2,” as used herein, refers to a protein portion of a viral hemagglutinin that includes at least about one β -sandwich that includes the substrate binding site, at least about two to about three short α -helices, at least about one small β -sheet at the bottom of the molecule and at least about two disulfide bonds. A β -strand in a viral hemagglutinin can include between about two to about 15 amino acids. A small β -strand can include about two amino acids; or between about two to about three amino acids; or between about two to four amino acids or between about two to about five amino acids. A small β -sheet can include between about two to about three β -strands; or between about three to about four β -strands. The β -sandwich that includes the substrate binding site of HA1-2 can further include at least about four β -strands as the top sheet and at least about three to about four β -strands as the bottom sheet. At least about one α -helix of the HA1-2 portion is located by the side of the β -sandwich that includes the substrate binding site and at least about one to about two are located at the bottom of the β -sandwich that includes the substrate binding site. Exemplary HA1-2 protein portions include SEQ ID NOS: 9, 12, 15, 18, 21, 24, 26, 28, 30, 32, 39, 41, 46, 48 and 50.

[0132] “HA1-3,” as used herein, refers to a protein portion of a viral hemagglutinin that includes at least one β -sandwich that includes the substrate binding site, at least about two short α -helices and at least one disulfide bond. “ β -sandwich,” as used herein, refers to at least about two sets of beta-sheets that form at least about one interactive layer. “Substrate binding site,” as used herein in reference to the β -sandwich, means any part of the portion of the naturally occurring viral hemagglutinin that has the capacity to interact or bind to a molecule. For example, the β -sandwich that includes the substrate binding site of the portion can include a portion that binds sialic acid. The β -sandwich that includes the substrate binding site of HA1-3 can further include at least about four β -strands as the top sheet and at least about three β strands as the bottom sheet. At least about one α -helix of the HA1-1 portion is located by the side of the β -sandwich that includes the substrate binding site and at least one other α -helix is located at the bottom of the β -sandwich that includes the substrate binding site. A short α -helix can include less than about 5 turns (2, 3, 4, 5 turns) in an α -helix. An α -helix in a viral hemagglutinin can be between one to about 15 turns; or between about two to 15

turns. Exemplary HA1-3 protein portions include SEQ ID NOS: 10, 13, 16, 19, 22, 25, 27, 29, 31 and 33.

[0133] “A sialic acid binding site,” as that phrase is used herein in reference to the portion of the protein from the naturally occurring viral hemagglutinin, means a part of the protein portion that has the capacity to interact with sialic acid residues. “A sialic acid binding site” is also referred to herein as “a sialic acid binding domain.”

[0134] “At least a portion,” as used herein, refers to any part of a component (e.g., a globular head, a secondary structure) or molecule (e.g., a protein, antigen, Toll-like Receptor, a peptide, flagellin, HA, matrix 2 protein (M2), matrix 2 ectodomain (M2e)); or the entirety of the component or the molecule. “At least a portion,” is also referred to herein as a “fragment.”

[0135] “At least a portion,” as used herein in reference to a flagellin (e.g., fljB/STF2, *E. coli* fljC, *S. muenchen* fljC), refers to any part of the flagellin (e.g., motif C; motif N; domain 1, 2, 3) or the entirety of the flagellin that can initiate an intracellular signal transduction pathway for a Toll-like Receptor.

[0136] A single polypeptide can exhibit several types of secondary structure. Without any stabilizing interactions, a polypeptide can assume a random-coil conformation. However, secondary structures, such as alpha(α)-helices and beta(β)-strands, can stabilize a protein or a portion of a protein. Lateral association of β -strands form β -sheets (also referred to herein as “ β -pleated sheets”). Secondary structures can be located at the surfaces of the portion, the protein, or the naturally occurring protein (e.g., viral hemagglutinin, flagellin, M2e). A tertiary structure of a protein is the three-dimensional arrangement of amino acid residues. In contrast to secondary structure, which is stabilized by, for example, hydrogen bonds, α -helices, β -strands, tertiary structure results from hydrophobic interactions between non-polar side chains of the portion, protein or naturally occurring viral hemagglutinin. The hydrophobic interactions hold the helices strands in random coils in a compact internal scaffold. The size and shape of a protein can depend on its primary amino acid sequence, as well as the number, size and arrangement of secondary structures.

[0137] “A globular head,” as that phrase is used herein, refers to a portion of a protein of a naturally occurring viral hemagglutinin that includes the receptor or sialic acid binding regions. “Globular head,” is also referred to herein as a “globular domain.” The globular head of viral hemagglutinin proteins has been determined based on x-ray crystallography as described, for example, by Wilson I. A., et al. *Nature* 289:366-373 (1981); Chen, J., et al., *Cell* 95:409-417 (1998); Ha Y., et al., *The EMBO Journal* 21:865-875 (2002); Russell, R. J., et al., *Virology* 325:287-296 (2004); and Cox, N. J., et al., In: Topley and Wilson’s Microbiology and Microbial Infections, eds. B W J Mathy, et al., Vol. 1 (9th ed.) New York, N.Y., Oxford Univ. Press, Ch. 32, p. 634 (1998). The globular head of a naturally occurring viral hemagglutinin is a component of the HA1 subunit of, for example, influenza viral hemagglutinin. In addition to the receptor binding domain, the globular head can include the E subdomain and F subdomain as described, for example, by Ha, Y., et al. *The EMBO Journal* 21:865-875 (2002).

[0138] The phrase, “causes the globular head to essentially retain its tertiary structure,” as used herein, refers to main-

tenance of the tertiary structure of the globular head of the naturally occurring viral hemagglutinin sufficient to stimulate a protective immune response in a subject.

[0139] The membrane fusion domain of a viral hemagglutinin is that region of the viral hemagglutinin (involved in binding of the viral hemagglutinin) that binds a host cell. A transmembrane domain of the viral hemagglutinin is that portion of the viral hemagglutinin that spans the membrane of the virus. A cytoplasmic domain of a viral hemagglutinin is that portion of the viral hemagglutinin located on the cytoplasmic surface of the virus.

[0140] The portion of the protein of the naturally occurring viral hemagglutinin (also referred to herein as “protein portion”) can further lack a signal sequence. The portion of a globular head employed in the methods described herein can include at least a portion of at least one secondary structure that includes at least a portion of at least one β -pleated sheet; at least one alpha helix and/or at least one member selected from the group consisting of a salt bridge, a leucine zipper and a zinc finger. The portion of a globular head can further include at least about one disulfide bond, at least about two disulfide bonds, at least about three disulfide bonds, at least about four disulfide bonds, at least about five disulfide bonds and at least about six disulfide bonds.

[0141] The method of making a protein that stimulates a protective immune response in a subject can further include the step of substituting a nucleic acid sequence encoding at least one amino acid residue selected from the group consisting of a hydrophilic amino acid residue, a polar amino acid residue and a neutral amino acid residue for a nucleic acid sequence that encodes at least one hydrophobic amino acid residue in the protein portion. The hydrophobic amino acid residue substituted can include at least one member selected from the group consisting of a phenylalanine residue, a tryptophan residue and tyrosine residue. The polar amino acid residue substituted for the hydrophobic amino acid can include at least one member selected from the group consisting of an aspartic acid residue and a glutamic acid residue.

[0142] The portion of a protein of a naturally occurring viral hemagglutinin can be a portion of a naturally occurring influenza viral hemagglutinin protein (e.g., influenza A, B and C). The influenza A viral hemagglutinin protein can be at least one member selected from the group consisting of H1, H2, H3, H5, H7 and H9.

[0143] The host cell employed in the methods described herein can be a prokaryotic host cell. The prokaryotic host cell can be at least one member selected from the group consisting of an *E. coli* prokaryotic host cell, a *Pseudomonas* prokaryotic host cell, a *Bacillus* prokaryotic host cell, a *Salmonella* prokaryotic host cell and a *P. fluorescens* prokaryotic host cell.

[0144] The method of making a protein that stimulates a protective immune response in a subject can further include the step of transforming the prokaryotic host cell with a chaperone nucleic acid sequence. The chaperone nucleic acid sequence can be at least one member selected from the group consisting of a groES-groEL chaperone, a dnaK-dnaJ-grpE chaperone, a groES-groEL-tig chaperone and a tig chaperone.

[0145] The method of making a protein that stimulates a protective immune response in a subject can further include

the step of fusing at least a portion of a Toll-like Receptor (TLR) agonist to the protein. The nucleic acid sequence encoding at least a portion of a Toll-like Receptor agonist can be operably linked to the nucleic acid sequence encoding the protein portion of the viral hemagglutinin. A linker (e.g., peptide linker) can be between the Toll-like Receptor agonist and the portion of the viral hemagglutinin.

[0146] Toll-like Receptors refer to a family of receptor proteins that are homologous to the *Drosophila melano-gaster* Toll protein. Toll-like Receptors are type I transmembrane signaling receptor proteins characterized by an extracellular leucine-rich repeat domain and an intracellular domain homologous to an interleukin 1 receptor. Toll-like Receptors include TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR 8, TLR9, TLR10, TLR11 and TLR12.

[0147] "Agonist," as used herein in referring to a TLR, means a molecule that activates a TLR signaling pathway. A TLR signaling pathway is an intracellular signal transduction pathway employed by a particular TLR that can be activated by a TLR ligand or a TLR agonist. Common intracellular pathways are employed by TLRs and include, for example, NF- κ B, Jun N-terminal kinase and mitogen-activated protein kinase. The Toll-like Receptor agonist can include at least one member selected from the group consisting of a TLR1 agonist, a TLR2 agonist (e.g., Pam3Cys, Pam2Cys, bacterial lipoprotein), a TLR3 agonist (e.g., dsRNA), a TLR4 agonist (e.g., bacterial lipopolysaccharide), a TLR5 agonist (e.g., a flagellin), a TLR6 agonist, a TLR7 agonist, a TLR8 agonist, a TLR9 agonist (e.g., unmethylated DNA motifs), TLR10 agonist, a TLR11 agonist and a TLR12 agonist.

[0148] The Toll-like Receptor agonists for use in the methods and compositions of the invention can be a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor.

[0149] TLR4 ligands (e.g., TLR4 agonists) for use in the compositions and methods of the invention can include at least one member selected from the group consisting of

[0150] SEQ ID NOS: 359-406 (see, PCT/US 2006/002906/WO 2006/083706; PCT/US 2006/003285/WO 2006/083792; PCT/US 2006/041865; PCT/US 2006/042051).

GGKSGRTG	(SEQ ID NO: 359)
KGYDWLVVG	(SEQ ID NO: 360)
EDMVYRIGVP	(SEQ ID NO: 361)
VKLSGS	(SEQ ID NO: 362)
GMLSLALF	(SEQ ID NO: 363)
CVVGSVR	(SEQ ID NO: 364)
IVRGCLGW	(SEQ ID NO: 365)
AAEERTLG	(SEQ ID NO: 366)

-continued

WARVVGWLR	(SEQ ID NO: 367)
SEGYRLFEGG	(SEQ ID NO: 368)
LVGGVVRGGS	(SEQ ID NO: 369)
GRVNDLWLAA	(SEQ ID NO: 370)
SGWMLWREGS	(SEQ ID NO: 371)
ERMEDRGDDL	(SEQ ID NO: 372)
KLCCFTECM	(SEQ ID NO: 373)
AVGSMERGRG	(SEQ ID NO: 374)
RDWVGGLDV	(SEQ ID NO: 375)
FFEVAKISQQ	(SEQ ID NO: 376)
WWYWC	(SEQ ID NO: 377)
MHLCSHA	(SEQ ID NO: 378)
WLFRRIG	(SEQ ID NO: 379)
YFWFRIG	(SEQ ID NO: 380)
MHLYCIA	(SEQ ID NO: 381)
WPLFPWIV	(SEQ ID NO: 382)
DMRSHAR	(SEQ ID NO: 383)
MHLCTHA	(SEQ ID NO: 384)
NLFPFY	(SEQ ID NO: 385)
MHLCTRA	(SEQ ID NO: 386)
RHLWYHA	(SEQ ID NO: 387)
WPFSAYW	(SEQ ID NO: 388)
WYLRS	(SEQ ID NO: 389)
GKGTDLG	(SEQ ID NO: 390)
IFVRMR	(SEQ ID NO: 391)
WLFRPVF	(SEQ ID NO: 392)
FLGWLMG	(SEQ ID NO: 393)
MHLWHHA	(SEQ ID NO: 394)
WWFPWKA	(SEQ ID NO: 395)
WYLPWL	(SEQ ID NO: 396)
WPFPRTF	(SEQ ID NO: 397)
WPFPPAYW	(SEQ ID NO: 398)
FLGLRWL	(SEQ ID NO: 399)
SRTDVGVLV	(SEQ ID NO: 400)
REKVSRGDKG	(SEQ ID NO: 401)
DWDAVESEYM	(SEQ ID NO: 402)
VSSAQEVVVP	(SEQ ID NO: 403)
LYTGGLEALG	(SEQ ID NO: 404)

-continued

VEEYSSSGVS	(SEQ ID NO: 405)
VCEVSDSVMA	(SEQ ID NO: 406)

[0151] TLR2 ligands (e.g., TLR2 agonists) for use in the compositions and methods of the invention can also include at least one member selected from the group consisting of SEQ ID NOS: 455-494 (see, PCT/US 2006/002906/WO 2006/083706; PCT/US 2006/003285/WO 2006/083792; PCT/US 2006/041865; PCT/US 2006/042051).

NPPTT	(SEQ ID NO: 455)
MRRIL	(SEQ ID NO: 456)
MISS	(SEQ ID NO: 457)
RGGSK	(SEQ ID NO: 458)
RGGF	(SEQ ID NO: 459)
NRTVF	(SEQ ID NO: 460)
NRFLG	(SEQ ID NO: 461)
SRHGR	(SEQ ID NO: 462)
IMRHP	(SEQ ID NO: 463)
EVCAP	(SEQ ID NO: 464)
ACGVY	(SEQ ID NO: 465)
CGPKL	(SEQ ID NO: 466)
AGCFS	(SEQ ID NO: 467)
SGGLF	(SEQ ID NO: 468)
AVRLS	(SEQ ID NO: 469)
GGKLS	(SEQ ID NO: 470)
VSEGV	(SEQ ID NO: 471)
KCQSF	(SEQ ID NO: 472)
FCGLG	(SEQ ID NO: 473)
PESGV	(SEQ ID NO: 474)
DPDSG	(SEQ ID NO: 475)
IGRFR	(SEQ ID NO: 476)
MGTLF	(SEQ ID NO: 477)
ADTHQ	(SEQ ID NO: 478)
HLLPG	(SEQ ID NO: 479)
GPLLH	(SEQ ID NO: 480)
NYRRW	(SEQ ID NO: 481)
LRQGR	(SEQ ID NO: 482)
IMWFP	(SEQ ID NO: 483)
RVVAP	(SEQ ID NO: 484)
IHVVP	(SEQ ID NO: 485)

-continued

MFGVP	(SEQ ID NO: 486)
CVWLQ	(SEQ ID NO: 487)
IYKLA	(SEQ ID NO: 488)
KGWF	(SEQ ID NO: 489)
KYMPH	(SEQ ID NO: 490)
VGKND	(SEQ ID NO: 491)
THKPK	(SEQ ID NO: 492)
SHIAL	(SEQ ID NO: 493)
AWAGT	(SEQ ID NO: 494)

[0152] The TLR2 ligand (e.g., TLR2 agonist) can also include at least a portion of at least one member selected from the group consisting of flagellin modification protein FlmB of *Caulobacter crescentus*; Bacterial Type III secretion system protein; invasin protein of *Salmonella*; Type 4 fimbrial biogenesis protein (PilX) of *Pseudomonas*, *Salmonella* SciJ protein; putative integral membrane protein of *Streptomyces*; membrane protein of *Pseudomonas*; adhesin of *Bordetella pertussis*; peptidase B of *Vibrio cholerae*; virulence sensor protein of *Bordetella*; putative integral membrane protein of *Neisseria meningitidis*; fusion of flagellar biosynthesis proteins FlhR and FlhB of *Clostridium*; outer membrane protein (porin) of *Acinetobacter*; flagellar biosynthesis protein FlhF of *Helicobacter*; ompA related protein of *Xanthomonas*; omp2a porin of *Brucella*; putative porin/fimbrial assembly protein (LHrE) of *Salmonella*; wbdK of *Salmonella*; Glycosyltransferase involved in LPS biosynthesis; *Salmonella* putative permease.

[0153] The TLR2 ligand (e.g., TLR agonist) can include at least a portion of at least one member selected from the group consisting of lipoprotein/lipopeptides (a variety of pathogens); peptidoglycan (Gram-positive bacteria); lipoteichoic acid (Gram-positive bacteria); lipoarabinomannan (mycobacteria); a phenol-soluble modulin (*Staphylococcus epidermidis*); glycoinositolphospholipids (*Trypanosoma Cruzi*); glycolipids (*Treponema maltophilum*); porins (*Neisseria*); zymosan (fungi) and atypical LPS (*Leptospira interrogans* and *Porphyromonas gingivalis*).

[0154] The TLR2 ligand (e.g., TLR2 agonist) can also include at least one member selected from the group consisting of SEQ ID NOS: 495-497 (see, PCT/US 2006/002906/WO 2006/083706; PCT/US 2006/003285/WO 2006/083792; PCT/US 2006/041865; PCT/US 2006/042051).

KGGVGPVRRSSRLRRTTQPG	(SEQ ID NO: 495)
GRGLCRGCRTRGRIKQLQSAHK	(SEQ ID NO: 496)
RWGYHLRDRKYKGVRSKGVPR	(SEQ ID NO: 497)

[0155] The TLR2 agonist can include at least a portion of a bacterial lipoprotein (BLP).

[0156] The TLR2 agonist can be a bacterial lipoprotein, such as Pam2Cys (S-[2,3-bis(palmitoyloxy)propyl]cysteine), Pam3Cys ([Palmitoyl]-Cys((RS)-2,3-di(palmitoy-

loxy)-propyl cysteine) or *Pseudomonas aeruginosa* OprI lipoprotein (OprI). Exemplary OprI lipoproteins include SEQ ID NO: 782, encoded by SEQ ID NO: 783. An exemplary *E. coli* bacterial lipoprotein for use in the invention described herein is SEQ ID NO: 784 encoded by SEQ ID NO: 785. A bacterial lipoprotein that activates a TLR2 signaling pathway (a TLR2 agonist) is a bacterial protein that includes a palmitoleic acid (Omueti, K. O., et al., *J. Biol. Chem.* 280: 36616-36625 (2005)). For example, expression of SEQ ID NOS: 783 and 785 in bacterial expression systems (e.g., *E. coli*) results in the addition of a palmitoleic acid moiety to a cysteine residue of the resulting protein (e.g., SEQ ID NOS: 782 and 784) thereby generating a TLR2 agonist for use in the compositions, fusion proteins and polypeptides of the invention. Production of tripalmitoylated-lipoproteins (also referred to as triacyl-lipoproteins) in bacteria occurs through the addition of a diacylglycerol group to the sulfhydryl group of a cysteine (e.g., cysteine 21 of SEQ ID NO: 784) followed by cleavage of the signal sequence and addition of a third acyl chain to the free N-terminal group of the same cysteine (e.g., cysteine 21 of SEQ ID NO: 784) (Sankaran, K., et al., *J. Biol. Chem.* 269:19706 (1994)), to generate a tripalmitoylated peptide (a TLR2 agonist) as shown, for example, in FIG. 74.

[0157] The Toll-like Receptor agonist in the compositions of the invention can further include at least one cysteine residue at the terminal amino acid of the amino-terminus and/or the terminal amino acid of the carboxy-terminus of the Toll-like Receptor agonist. For example, SEQ ID NO: 359 can further include at least one cysteine residue in a peptide bond to the amino-terminal glycine residue and/or at least one cysteine residue in a peptide bond to the carboxy-terminal glycine residue; SEQ ID NO: 360 can further include at least one cysteine residue in a peptide bond to the amino-terminal lysine residue and/or at least one cysteine residue in a peptide bond to the carboxy-terminal glycine residue; SEQ ID NO: 361 can further include at least one cysteine residue in a peptide bond to the amino-terminal glutamic acid residue and/or at least one cysteine residue in a peptide bond to the carboxy-terminal proline residue.

[0158] TLR5 agonists for use in the methods of the invention can include at least a portion of a flagellin. Flagellin are pathogen-associated molecular patterns (PAMPs) that activate TLR5.

[0159] The flagellin in the compositions and methods described herein can be at least a portion of a *S. typhimurium* flagellin (Genbank Accession Number AF045151); at least a portion of the *S. typhimurium* flagellin selected from the group consisting of SEQ ID NO: 498, SEQ ID NO: 812, SEQ ID NO: 816 and SEQ ID NO: 500; at least a portion of an *S. muenchen* flagellin (Genbank Accession Number AB028476) that includes at least a portion of SEQ ID NO: 504 and SEQ ID NO: 813; at least a portion of *P. aeruginosa* flagellin that includes at least a portion of SEQ ID NO: 815; at least a portion of a *Listeria monocytogenes* flagellin that includes at least a portion of SEQ ID NO: 820; at least a portion of an *E. coli* flagellin that includes at least a portion of SEQ ID NO: 502 and SEQ ID NO: 814; at least a portion of a *Yersinia* flagellin; and at least a portion of a *Campylobacter* flagellin.

[0160] The flagellin employed in the compositions of the invention can also include the polypeptides of SEQ ID NO:

498, SEQ ID NO: 500, SEQ ID NO: 504 and SEQ ID NO: 502; at least a portion of SEQ ID NO: 498, at least a portion of SEQ ID NO: 500, at least a portion of SEQ ID NO: 504 and at least a portion of SEQ ID NO: 502; and a polypeptide encoded by SEQ ID NO: 499, SEQ ID NO: 501, SEQ ID NO: 505 and SEQ ID NO: 503; or at least a portion of a polypeptide encoded by SEQ ID NO: 499, SEQ ID NO: 501, SEQ ID NO: 505 and SEQ ID NO: 503.

[0161] The flagellin employed in the compositions and method of the invention can lack at least a portion of a hinge region. Hinge regions are the hypervariable regions of a flagellin. Hinge regions of a flagellin are also referred to herein as "D3 domain or region," "propellor domain or region," "hypervariable domain or region" and "variable domain or region." "Lack" of a hinge region of a flagellin, means that at least one amino acid or at least one nucleic acid codon encoding at least one amino acid that comprises the hinge region of a flagellin is absent in the flagellin. Examples of hinge regions include amino acids 176-415 of SEQ ID NO: 498, which are encoded by nucleic acids 528-1245 of SEQ ID NO: 499; amino acids 174-422 of SEQ ID NO: 502, which are encoded by nucleic acids 522-1266 of SEQ ID NO: 503; or amino acids 173-464 of SEQ ID NO: 504, which are encoded by nucleic acids 519-1392 of SEQ ID NO: 505. Thus, if amino acids 176-415 were absent from the flagellin of SEQ ID NO: 498, the flagellin would lack a hinge region. A flagellin lacking at least a portion of a hinge region is also referred to herein as a "truncated version" of a flagellin.

[0162] "At least a portion of a hinge region," as used herein, refers to any part of the hinge region of the flagellin, or the entirety of the hinge region. "At least a portion of a hinge region" is also referred to herein as a "fragment of a hinge region." At least a portion of the hinge region of fljB/STF2 can be, for example, amino acids 200-300 of SEQ ID NO: 498. Thus, if amino acids 200-300 were absent from SEQ ID NO: 498, the resulting amino acid sequence of STF2 would lack at least a portion of a hinge region.

[0163] Alternatively, at least a portion of a naturally occurring flagellin can be replaced with at least a portion of an artificial hinge region. "Naturally occurring," in reference to a flagellin amino acid sequence, means the amino acid sequence present in the native flagellin (e.g., *S. typhimurium* flagellin, *S. muenchen* flagellin, *E. coli* flagellin). The naturally occurring hinge region is the hinge region that is present in the native flagellin. For example, amino acids 176-415 of SEQ ID NO: 498, amino acids 174-422 of SEQ ID NO: 502 and amino acids 173-464 of SEQ ID NO: 504, are the amino acids corresponding to the natural hinge region of STF2, *E. coli* fljC and *S. muenchen* flagellins, fljC, respectively. "Artificial," as used herein in reference to a hinge region of a flagellin, means a hinge region that is inserted in the native flagellin in any region of the flagellin that contains or contained the native hinge region.

[0164] The hinge region of a flagellin be deleted and replaced with an antigen (e.g., HA1-1, HA1-2, the maturational cleavage site) and the resulting construct fused to another antigen (e.g., 4xM2e).

[0165] An artificial hinge region may be employed in a flagellin that lacks at least a portion of a hinge region, which may facilitate interaction of the carboxy- and amino-terminus of the flagellin for binding to TLR5 and, thus, activation

of the TLR5 innate signal transduction pathway. A flagellin lacking at least a portion of a hinge region is designated by the name of the flagellin followed by a "Δ." For example, an STF2 (e.g., SEQ ID NO: 498) that lacks at least a portion of a hinge region is referenced to as "STF2Δ" or "fljB/STF2Δ" (e.g., SEQ ID NO: 500).

[0166] The flagellin for use in the methods and compositions of the invention can be a at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5; a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5; a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5; a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5, as described herein.

[0167] A recombinant fusion protein can be generated by operably linking a TLR agonist to the protein portion (e.g., HA1-1, HA1-2).

[0168] "Fusion protein," as used herein, refers to a protein generated from at least two similar or distinct components (e.g., a protein portion of HA and a TLR agonist). Fusion proteins can be generated recombinantly or by chemical conjugation.

[0169] In an embodiment, fusion proteins of protein portions of HA and TLR agonists (e.g., SEQ ID NOS: 89-92, 95, 151-160, 177, 209, 210 and 211) can be admixed or coadministered with fusion proteins of TLR agonists and M2e proteins (e.g., SEQ ID NOS: 528, 587, 589 and 591) and administered to a subject to stimulate an immune response, such as a protective immune response.

[0170] Fusion proteins of the invention can be designated by components of the fusion proteins separated by a ".". For example, "STF2.HA1-2" refers to a fusion protein comprising one STF2 protein and one HA1-2 protein; and "STF2Δ.HA1-2" refers to a fusion protein comprising one STF2 protein without the hinge region and HA1-2. Exemplary fusion proteins of the invention include SEQ ID NOS: 89-92, 95, 151-160, 177, 209, 210 and 211.

[0171] The fusion proteins can include, for example, two, three, four, five, six or more Toll-like Receptor agonists (e.g., flagellin) and two, three, four, five, six or more antigens (e.g., protein portions, such as HA1-1, HA1-2). When two or more TLR agonists and/or two or more protein portions comprise fusion proteins of the invention, they are also referred to as "multimers." For example, a multimer of HA1-1 can be four HA1-1 sequences, which is referred to herein as 4×HA1-1. Likewise, "2×HA1-1" is a multimer of two HA1-1 sequences. (See, for example, SEQ ID NOS: 342-346 and 348).

[0172] The fusion proteins of the invention can further include a linker between at least one component of the fusion protein (e.g., TLR agonist) and at least one other component of the fusion protein (e.g., HA1-1, HA1-2) of the

composition, a linker between at least two of similar components of the fusion protein (e.g., HA1-1, HA1-2) or any combination thereof. "Linker," as used herein in reference to a fusion protein of the invention, refers to a connector between components of the fusion protein in a manner that the components of the fusion protein are not directly joined. For example, one part of the fusion protein (e.g., flagellin) can be linked to a distinct part (e.g., protein portion, HA1-1, HA1-2, an antigen) of the fusion protein. Likewise, at least two or more similar or like components of the fusion protein can be linked (e.g., two flagellin can further include a linker between each flagellin, or two HA proteins can further include a linker between each HA protein).

[0173] Additionally, or alternatively, the fusion proteins of the invention can include a combination of a linker between distinct components of the fusion protein and similar or like components of the fusion protein. For example, a fusion protein can comprise at least two TLR agonists that further includes a linker between, for example, two or more flagellin; at least two protein portions of HA that further include a linker between them; a linker between one component of the fusion protein (e.g., flagellin) and another distinct component of the fusion protein (e.g., protein portions of HA), or any combination thereof.

[0174] The linker can be an amino acid linker. The amino acid linker can include synthetic or naturally occurring amino acid residues. The amino acid linker employed in the fusion proteins of the invention can include at least one member selected from the group consisting of a lysine residue, a glutamic acid residue, a serine residue and an arginine residue. The amino acid linker can include, for example, SEQ ID NOS: 521, 523, 524 and 526, encoded by the nucleic acid sequences of SEQ ID NOS: 520, 522, 525 and 527 respectively.

[0175] The Toll-like Receptor agonist of the fusion proteins of the invention can be fused to a carboxy-terminus, the amino-terminus or both the carboxy- and amino-terminus of the protein portion of HA (e.g., HA1-1, HA1-2 or other antigens, such as the ectodomain of the Matrix 2 protein, a maturational cleavage site).

[0176] Fusion proteins can be generated by fusing the protein portions of HA (or other antigens, such as the maturational cleavage site of HA) to at least one of four regions (Regions 1, 2, 3 and 4) of flagellin, which have been identified based on the crystal structure of flagellin (PDB:1UCU) (see FIG. 77).

[0177] Region 1 is TIAL (SEQ ID NO: 823) GLG (95-111 of SEQ ID NO: 841). The corresponding residues for *Salmonella typhimurium* fljB construct are TTLD (SEQ ID NO: 824) GTN (196-216 of SEQ ID NO: 841). This region is an extended peptide sitting in a groove of two beta strands (GTDQKID (SEQ ID NO: 825) and NGEVTL (SEQ ID NO: 826) of (SEQ ID NO: 841). Substitution of this peptide with an antigen (e.g., HA maturational cleavage site) may mimic the conformation of wild type maturational cleavage site peptide in HA0. Exemplary amino acids that may be substituted include: flagellin residues SGLDDAAI-KAAT (SEQ ID NO: 827) (201-212 of SEQ ID NO: 841) substituted with maturational cleavage site residues: RGIF-GAIGFIE (SEQ ID NO: 828), which correspond to the A/H3N2 subtype, or RGLFGAIGFIE (SEQ ID NO: 803), which correspond to the maturational cleavage site residues

from the A/H2N1, A/H1N1, A/H5N1 subtypes or with RGFFGAIAGFLE (SEQ ID NO: 805), which correspond to Influenza B HA maturational cleavage site residues.

[0178] Region 2 of the *Salmonella* flagellin is a small loop GTG (238-240 of SEQ ID NO: 841) in 1UCU structure (see FIG. 77). The corresponding loop in *Salmonella* fljB is GADAA (SEQ ID NO: 829) (244-248 of SEQ ID NO: 841). Insertion of an antigen (e.g., protein portions of HA, a maturation cleavage site peptide) in this loop or replacement of the entire loop with an antigen (e.g., protein portions of HA, a maturational cleavage site peptide) should preserve the extended loop structure of the maturational cleavage site peptide that is associated with the native HA molecule.

[0179] Region 3 is a bigger loop that resides on the opposite side of the Region 1 peptide (see FIG. 77). This loop can be simultaneously substituted together with region 1 to create a double copy of the antigen (e.g., protein portions of HA, a maturational cleavage site peptide). The loop starts from AGGA (SEQ ID NO: 830) and ends at PATA (SEQ ID NO: 831) (259-274 of SEQ ID NO: 841). The corresponding *Salmonella* fljB sequence is AAGA (SEQ ID

NO: 832) ...-...ATTK (SEQ ID NO: 833) (266-281 of SEQ ID NO: 841). The sequence AGATKTTMPAGA (SEQ ID NO: 834) (267-278 of SEQ ID NO: 841) can be replaced with the antigens (e.g., protein portions of HA, a maturation cleavage site peptides).

[0180] Region 4 is the loop (VTGTG (SEQ ID NO: 835)) connecting a short α -helix (TEAKAALTA (SEQ ID NO: 836)) and a β -strand (ASVVKMSYTDN (SEQ ID NO: 837)) in 1UCU structure (see FIG. 77). The corresponding loop in *Salmonella* fljB is a longer loop VDATDANGA (SEQ ID NO: 838) (307-315 of SEQ ID NO: 841). An antigen (e.g., a protein portion of HA, a maturation cleavage site peptide) can be inserted into or replace this region.

[0181] One or more copies of antigens (e.g., maturation cleavage site) can be inserted or used to replace the peptides listed in the above four regions. Preferably, the replacements would be in Region 1 and Region 3.

[0182] Exemplary sequences of maturation cleavage site peptides of HA are listed below:

Sequence	Subtype
NVPEKQTRGIFGAIAGFIE	A/H3N2 (SEQ ID NO: 800)
NVPQIESRGLFGAIAGFIE	A/H2N1 (SEQ ID NO: 801)
NIPSIQSRGLFGAIAGFIE	A/H1N1 (SEQ ID NO: 802)
RERRRKKRGLFGAIAGFIE	A/H5N1 (SEQ ID NO: 839)
PAKLLKERGFFGAIAGFLE	B/HA (SEQ ID NO: 804)
Sequence alignment of X-ray model (1UCU) and SEQ ID NO: 841 75.1% identity in 506 residues overlap; Score: 1703.0; Gap frequency: 2.6%	
1UCU 1 AQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAIANRFTANIKGL	
Sfla 2 AQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAIANRFTANIKGL	

1UCU 61 TQASRNANDGISIAQTTEGALNEINNLRVRELAVQSANSTNSQSDLSIQAEITQRLN	
Sfla 62 TQASRNANDGISIAQTTEGALNEINNLRVRELAVQSANSTNSQSDLSIQAEITQRLN	

1UCU 121 EIDRVSGQTQFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDLSNVQKAYKV	
Sfla 122 EIDRVSGQTQFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDLSNVQKAYDV	

1UCU 181 SDTAATVTGYAD--TTI---ALDNSTFKASATGLGGTDQKIDGDLKFDTTGKYYAKVTV	
Sfla 182 KDTAVTTKAYANNGTTLDVSGLDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGG	
*** * * * * * * * * * * * * * * * *	
1UCU 236 TGGT-GKDGYYEVSVDKTNGEVTLAGGATSPLTGGLPATATEDVKNVQVANADLTEAKA	
Sfla 242 FTGADAANKGDYEVNV-ATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSALAKN	
* * * * * * * * * * * * * * * *	
1UCU 294 ALTAAGVTGT----ASVVKMSYTDNNGKTIDGGLAVKVGDDYYSATQNK-DGSISINTTK	
Sfla 301 ALIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYAAADYDEATGAIAKAKTTS	
* * * * * * * * * * * * * * * *	
1UCU 349 YTADGTSKTALNKIGGADGKTEVVSIGGKTYAASKAEGHNFKAQFDLAEAAATTENPL	
Sfla 361 YTAADGTTKTAANQLGGVDGKTEVVITDGKTYNASKAAGHDFKAQPELAEAAAKTTENPL	
*** * * * * * * * * * * * * * * * *	
1UCU 409 QKIDAALAQVDTLRSDLGAVQNRFNSAITNLGNTVNNLTSVRSRIEDSDYATEVSNMSRA	
Sfla 421 QKIDAALAQVDALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA	

-continued

Sequence	Subtype
1UCU 469 QILQQAGTSVLAQANQVPQNVL SLLR (SEQ ID NO: 840)	
Sfla 481 QILQQAGTSVLAQANQVPQNVL SLLR (SEQ ID NO: 841)	

[0183] The methods of making a protein that stimulates a protective immune response in a subject can further include the step of operably linking a nucleic acid sequence encoding a carrier protein to the nucleic acid sequence encoding a portion of the viral hemagglutinin.

[0184] "Carrier," as used herein, refers to a molecule (e.g., protein, peptide) that can enhance stimulation of a protective immune response. Carriers can be physically attached (e.g., linked by recombinant technology, peptide synthesis, chemical conjugation or chemical reaction) to a composition (e.g., a protein portion of a naturally occurring viral hemagglutinin) or admixed with the composition.

[0185] Carriers for use in the methods and compositions described herein can include, for example, at least one member selected from the group consisting of Tetanus toxoid (TT), *Vibrio cholerae* toxoid, Diphtheria toxoid (DT), a cross-reactive mutant (CRM) of diphtheria toxoid, *E. coli* enterotoxin, *E. coli* B subunit of heat labile enterotoxin (LTB), Tobacco mosaic virus (TMV) coat protein, protein Rabies virus (RV) envelope protein (glycoprotein), thyroglobulin (Thy), heat shock protein HSP 60 Kda, Keyhole limpet hemocyanin (KLH), an early secreted antigen tuberculosis-6 (ESAT-6), exotoxin A, cholera toxin, hepatitis B core antigen, and the outer membrane protein complex of *N. meningitidis* (OMP) (see, for example, Schneerson, R., et al., *Prog Clin Biol Res* 47:77-94 (1980); Schneerson, R., et al., *J Exp Med* 152:361-76 (1980); Chu, C., et al., *Infect Immun* 40: 245-56 (1983); Anderson, P., *Infect Immun* 39:233-238 (1983); Anderson, P., et al., *J Clin Invest* 76:52-59 (1985); Fenwick, B. W., et al., 54:583-586 (1986); Que, J. U., et al. *Infect Immun* 56:2645-9 (1988); Que, J. U., et al. *Infect Immun* 56:2645-9 (1988); (Que, J. U., et al. *Infect Immun* 56:2645-9 (1988); Murray, K., et al., *Biol Chem* 380:277-283 (1999); Fingerut, E., et al., *Vet Immunol Immunopathol* 112:253-263 (2006); and Granoff, D. M., et al., *Vaccine* 11:Suppl 1:S46-51 (1993)).

[0186] Exemplary carrier proteins for use in the methods and compositions described herein can include at least one member selected from the group consisting of SEQ ID NOS: 788-795:

Cross-reactive mutant (CRM) of diphtheria toxin including, CRM197 (SEQ ID NO: 788)
 GADDVVDSSKSFVMENFSSYHGTPKPGYVDSIQKGIQPKSGTQGNVDDW
 KGFYSTDNKYDAAGYSVDNENPLSGKAGGVVKVTPGLTKVLALKVDNAE
 TIKKELGLSLTEPLMEQVGTEEFIKRFGDGASRVVLSLPFAEGSSSVEYT
 NNWEQAKALSVELETNFETRGRGQDAMYEYMAQACAGNRVRRSVGSSLS
 CIINLDWDVIRDKTKTKIESLKEHGPIKNKMSSEPNKTVSEKAKQYLEE

-continued

FHQTALEHPSELSEKTVTGTNPVFAGANYAAWAVNVAQVIDSETADNLEK
 TTAALSILPGIGSVMIADGAVHHNTEIEVAQSIASSLMVAQAIPLVGE
 LVDIGFAAYNFVEsIINLFQVVHNSYNRPAYSPGHKTQPFLLHDGYAVSWN
 TVEDSIIRTGFQGESGHDIKITAENTPLPIAGVLLPTIPGKLDVNSKTH
 ISVNGRIDMRCPAIDGDVTFPCPYKSPVYVNGVHANLHVAFHRSSEKI
 HSNEISSDSIGVLGYQKTVDHDKVNSKLSLFFFEIKS
 Coat protein of Tobacco mosaic virus (TMV)
 coat protein (SEQ ID NO: 789)
 MMAYSIPTPSQLVYFTENYADYIPFVNRLINARSNSFQTQSGRDELREIL
 IKSQSVSVSPISRFPAPAYYIYLRDPSISTVYTALLQSTDTRNRVIEVE
 NSTNVTAEQLNAVRRDDASTAIHNNLEQLLSLLTNGTGCVFNRTSFESA
 SGLTWLVTTTPRTA
 Coat protein of alfalfa mosaic virus (AMV)
 (SEQ ID NO: 790)
 MSSSQKKAGGKAGKPTKRSQNYAALRKAQLPKPPALKVPVAKPTNTILPQ
 TGCWVQSLGTPLSLSSNGLGARFLYSFLKDFAAPRILEDLIFRMVFSI
 TPESHAGSFCLTDDVTEDGRAVHGNPMQEPFHGAFHANEKYGFELVFTA
 YTHAGMQNQNFKHSYAVALLCLDFDALPEGSRNPSYRTNEVWVERKAYPPA
 GPLRSLITVGLFDDADDLDRQ
 Coat protein of Potato virus X
 (SEQ ID NO: 791)
 MTTPTANTTQATGSTSTSTTKTAGATPATTSGLFTIPDGEFFSTAPAIVAS
 NAVATNEDLSKIEAIWKDMKVPTDTMAQAAWDLVRHCADVGSSAQTEMID
 TGPYSNGISRARLAAAIKEVCTLRQFCMKYAPVVWNWMLTNNSPANWQA
 QGFKPEHKFAAFDFNGVTPNPAAIMPKEGLIRPPSEAEENAAQTAAAFVK
 ITKARAQSNDFASLDAAVTRGRITGTTTAEAVVTLPPP
 Porins from *Neisseria* sp, e.g., class I outer
 membrane protein of *Neisseria meningitidis*
 (SEQ ID NO: 792)
 MRKKLTALVLSALPLAAVADVSLYGEIKAGVEGRNYQLQLTEAQAANGGA
 SGQVKVTVKTKAKSRIRTKISDFGSFIFGFKGSEDLGEGLKAVWQLEQDVS
 VAGGGATQWGNRESFIFGLAGEFGTLRAGRANQFDDASQAIDPWDSNNDV
 ASQLGIFKRFIDDMPVSVRYDSPEFSGFSGSVQFVPAQNSKSAKPAYWT
 TVNTGSATTTTFVPAVVGKPGSDVYAGLNYKIIGGFAGNYAFKYAAUIA
 NVGRDAFELFLGSGSDQAKGTDPLKNHQNHQVHRLTGGYEEGGLNLALA

-continued

AQLDLSSENGDKTNSTTEIAATASYRFGNAVPRISYAHGFDFIERGKKGE
 NTSYDQIIAGVDYDFSKRTSAIVSGAWLKRNTGIGNYTQINAAASVGLRHK
 F

Major fimbrial subunit protein type I (Fimbrillin)
 (SEQ ID NO: 793)
 MVLKTSNSNRAFGVGDDESKVAKLTVMVYNGEQQEAIKSAENATKVEDIK

CSAGQRTLVMANTGAMELVGKTLAEVKALTTELTAENQEAAGLIMTAEP
 KTIVLKAGKNYIGSYGTGEGNHIENDPLKIKRVHARMAFTEIKVQMSAAY
 DNIYTFVPEKIYGLIAKKQSNLFGATLVNADANYLTGSLTFNGAYTPAN
 YANVPWLSRNYVAPADAPQGFYVLENDYSANGGTIHTPILCVYGIQKN
 GADLAGADLAAAQANWVDAEGKTYYPVLVNFNSNNYTYDSNYTPKIERN
 HKYDIKLTITGPGTNNPENPITESAHLNVQCTVAEWLVGQNATW

Mycoplasma fermentans macrophage activating
 lipopeptide (MALP-2)

(SEQ ID NO: 794)
 MKKSKILLGLSPIAAVLPAVAVSCGNDESNISFKEKDISKYTTTNANG

KQVVKNAELLKLPVLITDEGKIDDKSFNQSAFEALKMNKQTGIEINSVE
 PSSNFESAYNSALSAGHKIWLNGFKHQQS IKQYIDAHREELERNQIKII
 GIDFDIETEKWFSYSLQFNIKESAFTTGAYIASWLSEQDESKRVVASFGV
 GAFFPGVITTFNEGFAKGILYYNQHKSSKIYHTSPVKLDSGFTAGEKMNTV
 INNVLSSSTPADVKYNPHVILSVAGPATFETVRLANKGQYVIGVDSQGMII
 QDKDRILTSVLKHKIQAVYETLLDLILEKEEGYKPYVVKDKKADKKWSHF
 GTQKEKWIGVAENHFSNTEEQAKINNKEIKEAIKMFRELPEDFVKYINS DK
 ALKDGKNKIDNVSERIEAII SATNKA AK

p19 protein of *Mycobacterium tuberculosis*

(SEQ ID NO: 795)
 ATTLPVQRHPRSLFPEFSELF AAFPSFAGLRPTFDTRLMRIEDEMKEGRY

EVRAELPGVDPDKVDIMVRDQGLT IKAERTEQKDFDGRSEFAYGSFVRT
 VSLPVGADED DIKATYDKGILTVSVAVSEGKPTKEIKHIQIRSTN

[0187] The compositions of the invention can further include at least one adjuvant. Adjuvants contain agents that can enhance the immune response against substances that are poorly immunogenic on their own (see, for example, Immunology Methods Manual, vol. 2, I. Lefkovits, ed., Academic Press, San Diego, Calif., 1997, ch. 13). Immunology Methods Manual is available as a four volume set, (Product Code Z37,435-0); on CD-ROM, (Product Code Z37,436-9); or both, (Product Code Z37,437-7). Adjuvants can be, for example, mixtures of natural or synthetic compounds that, when administered with compositions of the invention, such as proteins that stimulate a protective immune response made by the methods described herein, further enhance the immune response to the protein. Compositions that further include adjuvants may further increase the protective immune response stimulated by compositions of the invention by, for example, stimulating a cellular and/or a humoral response (i.e., protection from disease versus antibody production). Adjuvants can act by enhancing protein uptake and localization, extend or prolong protein release, macrophage activation, and T and B cell stimu-

lation. Adjuvants for use in the methods and compositions described herein can be mineral salts, oil emulsions, mycobacterial products, saponins, synthetic products and cytokines. Adjuvants can be physically attached (e.g., linked by recombinant technology, by peptide synthesis or chemical reaction) to a composition described herein or admixed with the compositions described herein.

[0188] In another embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not a *Pichia pastoris* eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein. In one embodiment, the protein portion can further lack a signal sequence. In another embodiment, the protein portion can further include a sialic acid binding site.

[0189] In a further embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not an insect eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0190] In a further embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head, and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the protein portion into a eukaryotic host cell, wherein the eukaryotic host cell is not a stably transformed insect cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0191] In another embodiment, the invention is a method of making a protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby make a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that

causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; infecting a nucleic acid sequence encoding the protein portion into an insect cell host cell (e.g., a baculovirus insect host cell, such as Sf9 or High5 cells); and culturing the insect host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0192] The methods of making a protein that stimulates a protective immune response in a subject can further include the step of deleting at least one glycosylation site in the nucleic acid sequence encoding the protein portion. The glycosylation site that is deleted can include an N-glycosylation site.

[0193] The eukaryotic host cells employed in the methods of the invention can include a *Saccharomyces* eukaryotic host cell, an insect eukaryotic host cell (e.g., at least one member selected from the group consisting of a Baculovirus infected insect cell, such as *Spodoptera frugiperda* (Sf9) or *Trichoplusia ni* (High5) cells; and a *Drosophila* insect cell, such as Dme12 cells), a fungal eukaryotic host cell, a parasite eukaryotic host cell (e.g., a *Leishmania tarentolae* eukaryotic host cell), CHO cells, yeast cells (e.g., *Pichia*) and a *Kluyveromyces lactis* host cell.

[0194] Suitable eukaryotic host cells and vectors can also include plant cells (e.g., tomato; chloroplast; mono- and dicotyledonous plant cells; *Arabidopsis thaliana*; *Hordeum vulgare*; *Zea mays*; potato, such as *Solanum tuberosum*; carrot, such as *Daucus carota* L.; and tobacco, such as *Nicotiana tabacum*, *Nicotiana benthamiana* (Gills, M., et al., *Plant Biotechnol J.* 3:613-20 (2005); He, D. M., et al., *Colloids Surf B Biointerfaces*, (2006); Huang, Z., et al., *Vaccine* 19:2163-71 (2001); Khandelwal, A., et al., *Virology* 308:207-15 (2003); Marquet-Blouin, E., et al., *Plant Mol Biol* 51:459-69 (2003); Sudarshana, M. R., et al., *Plant Biotechnol J.* 4:551-9 (2006); Varsani, A., et al., *Virus Res*, 120:91-6 (2006); Kamarajugadda S., et al., *Expert Rev Vaccines* 5:839-49 (2006); Koya V, et al., *Infect Immun*. 73:8266-74 (2005); Zhang, X., et al., *Plant Biotechnol J.* 4:419-32 (2006)).

[0195] The proteins made by the methods of the invention and the compositions of the invention can be purified and characterized employing well-known methods (e.g., gel chromatography, cation exchange chromatography, SDS-PAGE), as described herein.

[0196] For large scale production, fermentation techniques can be employed, as described herein. Additional exemplary fermentation techniques can include a proposed cycle that can start with a culture inoculated into 6 L of MRBR media, as described herein, held at about 30° C., about pH 7, and DO controlled to greater than about 30%. A 6 liter feed can then be started at least about 30 minutes after glucose exhaustion. The proposed 6 liter feed media, when combined with 6 L of MRBR media, can provide the necessary conditions for *E. coli* growth based on about 52% utilization of carbon for growth. The feed may or may not include IPTG. The batch can be induced with at least 2 mM IPTG, introduced as a bolus, shortly after the feed is started to initiate production. The feed rate can start at about 20 mL feed per hour per liter bioreactor volume and increase over time based on the ability of the culture to accept more glucose without glucose accumulation. The culture can be harvested when the feed is

complete. The 6 liter feed media, about pH 6.0, can include Glucose 180 g/L; KH₂PO₄ 2 g/L; NaH₂PO₄ (H₂O) 4 g/L; (NH₄)₂HPO₄ 12 g/L; (NH₄)₂HSO₄ 4 g/L; DL-Alanine 40 g/L; Citric Acid 4 g/L; MgSO₄(7H₂O) 5.5 g/L; Trace Metals 6 mL; CaCl₂ 2.5 g/L; FeSO₄ 7H₂O 1 g/L.

[0197] Cell disruption and clarification in a large scale production can include removal of Triton X-100 from the resuspension buffer; dissolution of insolubles by the addition of 50 mM Tris, 25 mM NaCl, 8 M urea, about pH 8 to the lysate; addition of PEI (poly ethylamine) and subsequent removal by centrifugation with one or more of the buffers to remove nucleic acids and/or aid in filtration; the addition of flocculants, such as Aerosil 380, Aerosil 200, Alkoxide Alu C, and Celpur; and subsequent removal by centrifugation to aid in filtration. Cation exchange chromatography can include the use of a process resin, adding a denaturing endotoxin removal step containing up to 8 M urea and up to about 2% Triton X-100, and a step gradient elution. The step elution gradient can include about 100 to about 200 mM NaCl.

[0198] In yet another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein made by a method comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby form a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transforming a nucleic acid sequence encoding the portion into a prokaryotic host cell; and culturing the prokaryotic host cell to thereby make the protein that stimulates protective immunity in a subject.

[0199] In still another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein portion of a naturally occurring viral hemagglutinin, wherein the protein portion includes at least a portion of a globular head and at least a portion of one secondary structure that causes the globular head to essentially retain its tertiary structure and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain.

[0200] In a further embodiment, the invention is a method of making a viral hemagglutinin protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby make a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the portion in a eukaryotic host cell, wherein the eukaryotic host cell is not a *Pichia pastoris* eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0201] In yet another embodiment, the invention is a method of making a viral hemagglutinin protein that stimulates a protective immune response in a subject, comprising the steps of separating a portion of a protein from a naturally occurring viral hemagglutinin to thereby make a protein portion, wherein the protein portion includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion lacks a membrane fusion domain, a transmembrane domain and a cytoplasmic domain; transfecting a nucleic acid sequence encoding the portion in a eukaryotic host cell, wherein the eukaryotic host cell is not a *Drosophila melanogaster* eukaryotic host cell; and culturing the eukaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0202] In still another embodiment, the invention is a method of making a viral hemagglutinin protein that stimulates a protective immune response in a subject, comprising the steps of transforming a prokaryotic host cell with a nucleic acid sequence encoding at least one viral hemagglutinin that lacks a transmembrane domain and a cytoplasmic domain; and culturing the prokaryotic cell to thereby make the protein that stimulates a protective immune response in a subject.

[0203] An additional embodiment of the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein made by a method comprising the steps of transforming a prokaryotic host cell with a nucleic acid sequence encoding at least one viral hemagglutinin that lacks a transmembrane domain and a cytoplasmic domain; and culturing the prokaryotic host cell to thereby make the protein that stimulates a protective immune response in a subject.

[0204] In another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a protein having at least one viral hemagglutinin that lacks a transmembrane domain and a cytoplasmic domain, wherein the protein was expressed in a prokaryotic cell.

[0205] In still another embodiment, the invention is a composition comprising at least a portion of at least one pathogen-associated molecular pattern and a protein portion of a naturally occurring viral hemagglutinin, wherein the protein portion of the naturally occurring viral hemagglutinin includes at least a portion of a globular head and at least a portion of at least one secondary structure that causes the globular head to essentially retain its tertiary structure, and wherein the protein portion of the naturally occurring viral hemagglutinin lacks a membrane fusion domain, a transmembrane domain, and a cytoplasmic domain.

[0206] Pathogen-associated molecular patterns (PAMPs), such as a flagellin or a bacterial lipoprotein, refer to a class of molecules (e.g., protein, peptide, carbohydrate, lipid, lipopeptide, nucleic acid) found in microorganisms that, when bound to a pattern recognition receptor (PRR), can trigger an innate immune response. The PRR can be a Toll-like Receptor (TLR).

[0207] TLRs are the best characterized type of Pattern Recognition Receptor (PRR) expressed on antigen-present-

ing cells (APC). APC utilize TLRs to survey the microenvironment and detect signals of pathogenic infection by engaging the cognate ligands of TLRs, PAMPs. PAMP and TLR interaction triggers the innate immune response, the first line of defense against pathogenic insult, manifested as release of cytokines, chemokines and other inflammatory mediators; recruitment of phagocytic cells; and important cellular mechanisms which lead to the expression of costimulatory molecules and efficient processing and presentation of antigens to T-cells. TLRs control both innate and the adaptive immune responses.

[0208] The binding of PAMPs to TLRs activates innate immune pathways. Target cells can result in the display of co-stimulatory molecules on the cell surface, as well as antigenic peptide in the context of major histocompatibility complex molecules (see FIG. 40). The compositions, fusion proteins or polypeptides of the invention can include a PAMP (e.g., a flagellin) that binds to a TLR (e.g., TLR5), promoting differentiation and maturation of the APC, including production and display of co-stimulatory signals (see FIG. 40). The compositions can be internalized by its interaction with the TLR and processed through the lysosomal pathway to generate antigenic peptides, which are displayed on the surface in the context of the major histocompatibility complex.

[0209] The compositions, fusion proteins and polypeptides of the invention employ pathogen-associated molecular patterns (TLR agonists) that trigger cellular events resulting in the expression of costimulatory molecules, secretion of critical cytokines and chemokines; and efficient processing and presentation of antigens to T-cells. As discussed above, TLRs recognize PAMPs including bacterial cell wall components (e.g., bacterial lipoproteins and lipopolysaccharides), bacterial DNA sequences that contain unmethylated CpG residues and bacterial flagellin. TLRs act as initiators of the innate immune response and gatekeepers of the adaptive immune response (Medzhitov, R., et al., *Cold Springs Harb. Symp. Quant. Biol.* 64:429 (1999); Pasare, C., et al., *Semin. Immunol* 16:23 (2004); Medzhitov, R., et al., *Nature* 388:394 (1997); Barton, G. M., et al., *Curr. Opin. Immunol* 14:380 (2002); Bendelac, A., et al., *J. Exp. Med.* 195:F19 (2002)).

[0210] As discussed above, the binding of PAMPs to TLRs activates immune pathways for use in the compositions, fusion proteins and polypeptides of the invention, which can be employed in stimulating the immune system in a subject. The compositions, fusion proteins and polypeptides of the invention can trigger an immune response to an antigen (e.g., a viral protein, such as an influenza viral) and trigger signal transduction pathways of the innate and adaptive immune system of the subject to thereby stimulate the immune system of a subject. Stimulation of the immune system of the subject may prevent infection by an antigen or a virus (e.g., an influenza virus) and thereby treat the subject or prevent the subject from disease, illness and, possibly, death.

[0211] In an additional embodiment, the invention is a composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

[0212] "Flagellin component," as used herein, means at least part of or the entirety of a flagellin.

[0213] “Activates,” when referring to a Toll-like Receptor (TLR), means that the component (e.g., a flagellin component or a Toll-like Receptor agonist component) stimulates a response associated with a TLR. For example, bacterial flagellin activates TLR5 and host inflammatory responses (Smith, K. D., et al., *Nature Immunology* 4:1247-1253 (2003)). Bacterial lipopeptide activates TLR1; Pam3Cys, Pam2Cys activate TLR2; dsRNA activates TLR3; LPS (LPS-binding protein) and LPS (lipopolysaccharide) activate TLR4; imidazoquinolines (anti-viral compounds and ssRNA) activate TLR7; and bacterial DNA (CpG DNA) activates TLR9. TLR1 and TLR6 require heterodimerization with TLR2 to recognize ligands (e.g., TLR agonists, TLR antagonists). TLR1/2 are activated by triacyl lipoprotein (or a lipopeptide, such as Pam3Cys), whereas TLR6/2 are activated by diacyl lipoproteins (e.g., Pam2Cys), although there may be some cross-recognition. In addition to the natural ligands, synthetic small molecules including the imidazoquinolines, with subclasses that are specific for TLR7 or TLR8 can activate both TLR7 and TLR8. There are also synthetic analogs of LPS that activate TLR4, such as monophosphoryl lipid A [MPL].

[0214] TLR activation can result in signaling through MyD88 and NF- κ B. There is some evidence that different TLRs induce different immune outcomes. For example, Hirschfeld, et al. *Infect Immun* 69:1477-1482 (2001)) and Re, et al. *J Biol Chem* 276:37692-37699 (2001) demonstrated that TLR2 and TLR4 activate different gene expression patterns in dendritic cells. Pulendran, et al *J Immunol* 167:5067-5076 (2001)) demonstrated that these divergent gene expression patterns were recapitulated at the protein level in an antigen-specific response, when lipopolysaccharides that signal through TLR2 or TLR4 were used to guide the response (TLR4 favored a Th1-like response with abundant IFN γ secretion, while TLR2 favored a Th2-line response with abundant IL-5, IL-10, and IL-13 with lower IFN γ levels). There is redundancy in the outcome of signaling through different TLRs.

[0215] Activation of TLRs can result in increased effector cell activity that can be detected, for example, by measuring IFN γ -secreting CD8 $^{+}$ cells (e.g., cytotoxic T-cell activity); increased antibody responses that can be detected by, for example, ELISA, virus neutralization, and flow cytometry (Schnare, M., et al., *Nat Immunol* 2:947 (2001); Alexopoulou, L., et al., *Nat Med* 8:878 (2002); Pasare, C., et al., *Science* 299:1033 (2003); Napolitani, G., et al., *Nat Immunol* 6:769 (2005); and Applequist, S. E., et al., *J Immunol* 175:3882 (2005)).

[0216] The composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 can further include at least a portion of at least one member selected from the group consisting of a Toll-like Receptor 1 agonist, a Toll-like Receptor 2 agonist (e.g., Pam3Cys, Pam2Cys), a Toll-like Receptor 3 agonist, a Toll-like Receptor 4 agonist, a Toll-like Receptor 6 agonist, a Toll-like Receptor 7 agonist, a Toll-like Receptor 8 agonist a Toll-like Receptor 9 agonist, a Toll-like Receptor 10 agonist, a Toll-like Receptor 11 agonist and a Toll-like Receptor 12 agonist.

[0217] The a Toll-like Receptor 1 agonist, a Toll-like Receptor 2 agonist, a Toll-like Receptor 3 agonist, a Toll-like

Receptor 4 agonist, a Toll-like Receptor 6 agonist, a Toll-like Receptor 7 agonist, a Toll-like Receptor 8 agonist a Toll-like Receptor 9 agonist a Toll-like Receptor 10 agonist, a Toll-like Receptor 11 agonist and a Toll-like Receptor 12 agonist employed in the compositions, such as compositions comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates TLR5, can further include at least one additional cysteine residue.

[0218] In one embodiment, at least one cysteine residue substitutes for at least one amino acid residue in a naturally occurring flagellin amino acid sequence of the flagellin component.

[0219] The cysteine residue that substitutes for at least one amino acid residue in a naturally occurring flagellin amino acid sequence of the flagellin component can be remote to at least one amino acid of the Toll-like Receptor 5 recognition site of the flagellin component. “Toll-like Receptor 5 recognition site,” means that part of the TLR5 ligand (e.g., TLR5 agonist) that interacts with TLR5 to mediate a cellular response. “Toll-like Receptor 5 recognition site” is also referred to as a “Toll-like Receptor 5 activation site” and a “Toll-like Receptor 5 activation domain.”

[0220] Likewise, “Toll-like Receptor recognition site,” means that part of the Toll-like Receptor ligand (e.g., a Toll-like Receptor agonist) that interacts with its respective TLR to mediate a cellular response. “Toll-like Receptor recognition site” is also referred to as a “Toll-like Receptor activation site” and a “Toll-like Receptor activation domain.”

[0221] The cysteine residue that substitutes for at least one amino acid residue in a naturally occurring flagellin amino acid sequence of the flagellin component can also be remote to at least one amino acid of the flagellin component involved in binding to the toll like receptor 5.

[0222] In another embodiment, the flagellin component includes at least a portion of a naturally occurring flagellin amino acid sequence in combination with the cysteine residue.

[0223] The cysteine residue used in combination (also referred to herein as “added cysteine”) with at least a portion of a naturally occurring flagellin amino acid sequence can be at least one member selected from the group consisting of the amino-terminal amino acid of the flagellin component or the Toll-like Receptor agonist component and the carboxy-terminal amino acid of the flagellin component or the Toll-like Receptor agonist component. The cysteine residue used in combination with at least a portion of a naturally occurring flagellin amino acid sequence can be remote to at least one amino acid of the Toll-like Receptor 5 recognition site of the flagellin component or remote to at least one amino acid of the Toll-like Receptor recognition site of the Toll-like Receptor agonist component.

[0224] The cysteine residue used in combination with at least a portion of a naturally occurring flagellin amino acid sequence can be remote to at least one amino acid of the flagellin component involved in binding to the Toll-like Receptor 5 or remote to at least one amino acid of a Toll-like Receptor agonist component involved in binding to the Toll-like Receptor.

[0225] In another embodiment, the flagellin component lacks at least a portion of a hinge region of a naturally occurring flagellin amino acid sequence.

[0226] The composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 can further include at least a portion of at least one antigen (e.g., an influenza antigen, such as an influenza A, B or C antigen).

[0227] The antigen can be an essentially hydrophobic antigen, such as a maturational cleavage site antigen of HA. The maturational cleavage site antigen can be at least one member selected from the group consisting of (SEQ ID NOS: 530, 532, 533, 534, 535, 600, 601, 602, 603 and NVPEKQTRGIFGAIAGFIE (H3) (SEQ ID NO: 796), NIPSIQSRGLFGAIAGFIE (H1) (SEQ ID NO: 797), PAKLLKERGFFGAIAGFIE (FLU B) (SEQ ID NO: 798), RERRRRKKRGLFGAIAGFIE (H5) (SEQ ID NO: 799), RGLXGAIAGFIE (SEQ ID NO: 821), RGLXGAIAGFIE (SEQ ID NO: 822).

[0228] "Essentially hydrophobic," as used herein, means that the antigen has limited solubility in an aqueous solution or environment. The hydrophobic nature of peptides or proteins can be determined and compared, for example, by the Kyte-Doolittle hydrophobicity scale (Kyte, J., et al., *Mol. Biol.* 157: 105-132 (1982), (1982)), which assigns a numerical value for each of the 20 amino acids according to relative hydrophobicity. A positive value indicates a hydrophobic amino acid. A negative value indicates a hydrophilic amino acid. The average of these values for an individual peptide or polypeptide (calculated by adding the individual hydrophobicity values for each amino acid of the polypeptide or protein and dividing the total value by the number of amino acids in the polypeptide or protein) provides an index of overall hydrophobicity known as the "grand average of hydropathicity," also referred to as "GRAVY." A GRAVY value greater than zero indicates the protein or peptide is essentially hydrophobic. A GRAVY value less than zero indicates the protein or peptide is essentially hydrophilic. The individual hydrophobicity values, for the 20 naturally occurring amino acids according to the Kyte-Doolittle scale (Kyte, J., et al., *Mol. Biol.* 157: 105-132 (1982)), are as follows:

Alanine	1.8
Arginine	-4.5
Asparagine	-3.5
Aspartic acid	-3.5
Cysteine	2.5
Glutamine	-3.5
Glutamic acid	-3.5
Glycine	-0.4
Histidine	-3.2
Isoleucine	4.5
Leucine	3.8
Lysine	-3.9
Methionine	1.9
Phenylalanine	2.8
Proline	-1.6
Serine	-0.8
Threonine	-0.7
Tryptophan	-0.9

-continued

Tyrosine	-1.3
Valine	4.2

[0229] The hydrophobicity of antigens, such as protein or peptide antigens, for use in the compositions and methods of the invention can be determined by calculating a GRAVY score based on the hydrophobicity values of the above Kyte-Doolittle scale. For example, GRAVY scores, which indicate essentially hydrophobic peptides, were calculated by addition of the hydrophobicity values (supra) of individual amino acids for the following maturational cleavage site amino peptides:

Maturational Cleavage Site	GRAVY
NVPEKQTRGIFGAIAGFIE (A/H3N2) (SEQ ID NO: 800)	0.053
NVPQIESRGLFGAIAGFIE (A/H2N1) (SEQ ID NO: 801)	0.453
NIPSIQSRGLFGAIAGFIE (A/H1N1) (SEQ ID NO: 802)	0.611
RGLFGAIAGFIE (Influenza A conserved region) (SEQ ID NO: 803)	1.067
PAKLLKERGFFGAIAGFIE (Influenza B) (SEQ ID NO: 804)	0.400
RGFFGAIAGFIE (Influenza B conserved region) (SEQ ID NO: 805)	0.925

[0230] Likewise, GRAVY scores, which indicate essentially hydrophilic peptides, were calculated for the following peptides:

Amino Acid Sequence	GRAVY
human influenza M2e (SEQ ID NO: 806) SLLTEVETPIRNEWGSRSDSDP	-1.129
Vietnam influenza M2e (SEQ ID NO: 807) GSGAGSLLTEVETPIRNEWECRSDSDP	-0.907
Hong Kong influenza M2e (SEQ ID NO: 808) GSGAGSLLTEVETLIRNGWGCRCSDSDP	-0.507
West Nile Virus E peptide 001 (SEQ ID NO: 809) LTSGLKCRVRMEKQLKGT	-0.475
Dengue 2 E peptide (SEQ ID NO: 810) EAEPFGDSYIIIGVEPGQLKLNWFKK	-0.333
BCRABL wt peptide (SEQ ID NO: 811) SLLTEVETPIRNEWGSRSDSDP	-1.129

[0231] An antigen can be any molecule (e.g., protein, peptide, glycoprotein, glycopeptide, carbohydrate, lipid, lipopeptide, polysaccharide) that can be recognized by the components of the immune system regardless of whether it can trigger activation of the immune system. The antigen can be a fragment or portion of a naturally occurring antigen

or a synthetic molecule that mimics the naturally occurring antigen or a portion of the naturally occurring antigen.

[0232] The antigen can be a viral antigen. A “viral antigen,” as used herein, refers to any portion of a virus (e.g., influenza virus, flavivirus) that generates an immune response in a subject either when employed in combination with a TLR agonist (e.g., a flagellin, Pam2Cys, Pam3Cys) or in the absence of a TLR agonist. The viral antigen can be a portion or a fragment of a naturally occurring virus or a synthetic molecule that mimics a naturally occurring virus, such as a recombinant or synthetic protein (e.g., influenza virus, flavivirus), peptide, lipid, carbohydrate, that generates an immune response in the subject. The influenza antigen can include at least one member selected from the group consisting of an influenza A antigen, influenza B antigen and an influenza C antigen. The influenza antigen can be an influenza virus integral membrane protein, such as HA, or a protein portion of HA, such as HA1-1 and HA1-2.

[0233] The antigen can be at least a portion of an influenza Matrix 2 (M2) protein, including at least a portion of the ectodomain of an influenza M2 protein (M2e).

[0234] Matrix protein 2 (M2 or M2 protein) is a proton-selective integral membrane ion channel protein of the influenza A virus. M2 is abundantly expressed at the plasma membrane of virus-infected cells, but is generally underexpressed by virions. For example, a portion of an M2 sequence of influenza A is SEQ ID NO: 508, which is encoded by SEQ ID NO: 509. The native form of the M2 protein is a homotetramer (i.e., four identical disulfide-linked M2 protein molecules). Each of the units are helices stabilized by two disulfide bonds. M2 is activated by low pH. Each of the M2 protein molecules in the homotetramer consists of three domains: a 24 amino acid outer or N (amino)-terminal domain (e.g., SEQ ID NO: 510; also referred to herein as a “human consensus sequence”), which is encoded by SEQ ID NO: 511; a 19 hydrophobic amino acid transmembrane region, and a 54 amino acid inner or C (carboxy)-terminal domain. The M2 protein can vary depending upon the influenza viral subtype (e.g., H1 and H5 subtypes of influenza A) and influenza viral source (e.g., Puerto Rico, Thailand, New York, Hong Kong), as shown, for example, in exemplary amino-terminal sequences of M2 proteins (SEQ ID NOS: 544-556 and 570-578) and as described in PCT/US2005/046662 (WO2006/069262).

[0235] The M2 protein has an important role in the life cycle of the influenza A virus. It is important in the uncoating stage where it permits the entry of protons into the viral particle, which lowers the pH inside the virus, resulting in dissociation of the viral matrix protein M1 from the ribonucleoprotein RNP. As a consequence, the virus coat is removed and the contents of the virus are released from the endosome into the cytoplasm of the host cell for infection.

[0236] The function of the M2 channel can be inhibited by antiviral drugs, such as amantadine and rimantadine, which prevent the virus from infecting the host cell. Such antiviral drugs usually bind the transmembrane region of the M2 protein and sterically block the ion channel created by the M2 protein, which prevents protons from entering and uncoating the virion.

[0237] The M2 protein for use in the compositions and methods of the invention can that include at least a portion

of SEQ ID NO: 510 encoded by SEQ ID NO: 511 or at least a portion of SEQ ID NO: 544, encoded by SEQ ID NO: 603. The M2 protein can further include at least one member selected from the group consisting of SEQ ID NO: 512, SEQ ID NO: 516, SEQ ID NO: 531; SEQ ID NO: 536 (Flu A H5N1 M2e, 2004 Viet Nam Isolate with serine replacing cysteine); SEQ ID NO: 537 (Flu A H5N1 M2e, 2004 Viet Nam Isolate); SEQ ID NO: 538 (Flu A H5N1 M2e, Hong Kong 97 Isolate with serine replacing cysteine); SEQ ID NO: 539 (Flu A H5N1 M2e, Hong Kong 97 Isolate); SEQ ID NO: 540 (Flu A H7N2 M2e Chicken/New York 95 Isolate with serine replacing cysteine); SEQ ID NO: 541 (Flu A H7N2 M2e, Chicken/New York 95 Isolate); SEQ ID NO: 542 (Flu A H9N2 M2e, Hong Kong 99 Isolate with serine replacing cysteine); and SEQ ID NO: 543 (Flu A, Hong Kong 99 Isolate). Certain cysteine residues, for example, amino acids 16 and 18 of SEQ ID NO: 537; amino acids 17 and 19 of SEQ ID NOS: 539, 541 and 543 in the naturally occurring sequence of at least a portion of M2 protein are replaced with a serine (see, SEQ ID NOS: 538, 540, 542 and 544, respectively).

[0238] The compositions comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 that further includes an antigen (e.g., a maturational cleavage site peptide, protein portion of HA, such as HA1-1, HA1-2) can be co-administered or admixed with another flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 that further includes a different, distinct antigen (e.g., M2e) or a fusion protein of a TLR agonist and an antigen (e.g., STF2.HA1-1, STF2.HA1-2, STF2Δ.HA1-1, STF2Δ.HA1-2, STF2.M2e, STF2.4×M2e, STF2Δ.M2e, STF2Δ.4×M2e).

[0239] The antigen included in the compositions and employed in the methods of the invention can be at least a portion of a microbial-related antigen, for example, antigens of pathogens, such as viruses, fungi or bacteria. Antigens can also be microorganism-related, and other disease-related antigens, such as antigens associated with allergies and cancer. The antigen can also be at least a portion of an antigen of a bacteria, a virus, a fungi, a yeast, a protozoa, a metazoa, a tumor, a malignant cell, a plant cell, an animal cell, a hormone and an amyloid-β peptide. The antigen can be a peptide, a polypeptide, a lipoprotein, a glycoprotein and a mucoprotein.

[0240] The antigen included in the compositions and employed in the methods of the invention can be a pathogen-related antigen. “Pathogen-related antigen,” as used herein, refers to any molecule (e.g., protein, peptide, carbohydrate, lipoprotein, polysaccharide) that is associated with a pathogen. Exemplary pathogen-related antigens include, for example, antigens of vaccinia, avipox virus, turkey influenza virus, bovine leukemia virus, feline leukemia virus, avian influenza, chicken pneumovirus, canine parvovirus, equine influenza, FHV, Newcastle Disease Virus (NDV), Chicken/Pennsylvania/1/83 influenza virus, infectious bronchitis virus; Dengue virus, measles virus, Rubella virus, pseudorabies, Epstein-Barr Virus, HIV, SIV, EHV, BHV, HCMV, Hantaan, *C. tetani*, mumps, Morbillivirus, Herpes Simplex Virus type 1, Herpes Simplex Virus type 2, Human

cytomegalovirus, Hepatitis A Virus, Hepatitis B Virus, Hepatitis C Virus, Hepatitis E Virus, Respiratory Syncytial Virus, Human Papilloma Virus, Influenza Virus, *Salmonella*, *Neisseria*, *Borrelia*, *Chlamydia*, *Bordetella*, *Plasmodium* (e.g., *Plasmodium falciparum* and *Plasmodium vivax*), *Toxoplasma*, *Cryptococcus*, *Streptococcus*, *Staphylococcus*, *Haemophilus*, *Diphtheria*, *Tetanus*, *Pertussis*, *Escherichia*, *Candida*, *Aspergillus*, *Entamoeba*, *Giardia* and *Trypanosoma*.

[0241] The antigen included in the compositions and employed in the methods of the invention can include bacterial capsular antigens. Exemplary bacterial capsular antigens include, for example, at least one member selected from the group consisting of a Group B *Streptococcus*, including *Streptococcus agalactiae*, capsular polysaccharides type Ia, Ib, Ia/c, II, III, IV, V and VI; *Streptococcus pneumoniae* capsular polysaccharides types 1, 2, 3, 4, 5, 6A, 6B, 7F, 9N, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 20, 22F, 23F, and 33F or any combination of these (typing according to the Danish system); *Staphylococcus aureus* capsular polysaccharides type 5, 8 and 336; *Bacillus anthracis* vegetative state capsular poly gamma D glutamic acid; *Mycobacteria tuberculosis* lipo-arabino mannan capsule; *Plasmodium falciparum* surface oligosaccharide; *Neisseria meningitidis* capsular polysaccharides of group A, B, C, X, Y and W135 or any combination thereof.

[0242] Compositions comprising the flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 can include an antigen that is produced by chemical modification of the side-chains of the antigen (e.g., capsular antigens, pathogen-related antigens, influenza antigens, flavivirus antigens) in a manner that generates an active group that is coupled (also referred to herein as chemical conjugation) to a compatible reactive group created on the substituted flagellins of the invention (see infra). Derivatization (modification) of the capsular material is carried out so that a small minority of the side-chains are modified (about 5%) to avoid destruction of the majority of immunoreactive structures on the capsular material.

[0243] The antigen included in the compositions and employed in the methods of the invention can be at least a portion of at least one member selected from the group consisting of a West Nile viral protein, a Langat viral protein, a Kunjin viral protein, a Murray Valley encephalitis viral protein, a Japanese encephalitis viral protein, a Tick-borne encephalitis viral protein, Dengue 1 viral protein, Dengue 2 viral protein, Dengue 3 viral protein, Dengue 4 viral protein, hepatitis C viral protein and a Yellow fever viral protein (see, for example, PCT/US2006/001623 (WO2006/078657)).

[0244] The genus flavivirus is in the virus family Flaviviridae and consists of about 70 viruses. Mosquito or ticks transmit most of these viruses. Several flaviviruses are significant human pathogens, including the four dengue viruses (Den1, Den2, Den3 and Den4), yellow fever (YF), Japanese encephalitis (JE), West Nile (WN, also referred to herein as "WNV") and Tick-borne encephalitis (TBE) (Weaver S. C., et al., *Nat Rev Microbiol* 10: 789-801 (2004)). The flavivirus genus is divided into a number of serogroups based on cross-neutralization tests, including the

dengue serogroup that contains four serologically and genetically distinct viruses termed. DEN-1, DEN-2, DEN-3 and DEN-4.

[0245] Flaviviruses are small, enveloped viruses with icosahedral capsids. The flavivirus genome is a single-stranded positive-sense RNA (about 11 kb) that is directly translated by the host cell machinery following infection. The viral genome is translated as a single polypeptide that undergoes co- and post-translational cleavage by viral and cellular enzymes to generate three structural proteins of the flavivirus (the capsid (C), the membrane (M) and the envelope (E) proteins); and seven nonstructural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B, and NS5) (Weaver, et al., *Annu Rev Microbiol* 1990:44-649 (2004)). The viral capsid is composed of the C-protein, while both the M- and envelope proteins are located on the envelope surface of the virion (Weaver, S. C., et al., *Nat. Rev. Microbiol.* 10:789-801 (2004); Chambers et al., *Annu Rev. Microbiol.* 44: 649-688 (1990)). A major immunogen for flaviviruses is the membrane envelope protein.

[0246] A flavivirus can enter a host cell when the viral envelope protein binds to a receptor and responds by conformational rearrangement to the reduced pH of an endosome. The conformational change induces fusion of viral and host-cell membranes.

[0247] The envelope of a flavivirus may function as a receptor binding protein and to facilitate fusion of the virus and host cell membrane. Envelope proteins of flaviviruses have common structural (domains I, II and III) and functional features (receptor binding of virus and host cell and fusion functions) and are class II fusion glycoproteins (Lescar et al., *Cell* 105:137-148 (2001)).

[0248] In the pre-fusion conformation, envelope proteins form homodimers on the outer surface of the virus particles (Rey, et al., *Nature* 375:291-298); Kuhn, et al., *Cell* 108:717-725 (2002); Mukhopadhyay, et al., *Science* 302:248 (2003)). Each envelope protein monomer folds into three structural domains (domains I, II and III) predominantly composed of β -strands. Domain I (also referred to herein as "I" or "DI") is centrally located in the structure and has an N-glycosylation site in glycosylated envelope proteins. Domain II (also referred to herein as "II" or "DII") of the envelope protein promotes dimerization and has a fusion loop that inserts into the target host membrane during the pH-dependent fusion of the virus (Modis, et al., *Nature* 427:313-319 (2004); Bressanelli, et al., *EMBO J* 23:728-738 (2004)). Domain III (also referred to herein as "III" or "DIII") is at the carboxy-terminus of the envelope protein. Domain III is also referred to as "domain B" in earlier antigenic mapping studies. Domain III has several epitopes that can elicit virus-neutralizing antibodies (Roehrig, *Adv Virus Res* 59:141-175 (2003)).

[0249] Domain I of the Tick-borne encephalitis envelope protein corresponds to amino acids 1-51, 137-189 and 285-302 of SEQ ID NO: 777; domain II of the Tick-borne encephalitis envelope protein of SEQ ID NO: 777 corresponds to amino acids 52-136 and 190-284; and domain III corresponds to amino acids 303-395 of SEQ ID NO: 777. (Rey, F. A., et al., *Nature* 375:291-298 (1995)). SEQ ID NO: 777 is encoded by SEQ ID NO: 778. Domain I of the Dengue 2 flavivirus envelope protein corresponds to amino acids 1-52, 132-193 and 280-296 of SEQ ID NO: 763;

domain II corresponds to amino acids 53-131 and 194-279 of SEQ ID NO: 763; and domain III corresponds to amino acids 297-495 of SEQ ID NO: 763 (Modis, Y., et al., *Nature* 427:313-319 (2004)). The location of domains I, II and III of other flavivirus (e.g., West Nile virus, Japanese encephalitis, Dengue 1 virus, Dengue 3 virus and Dengue 4 virus) is based on homology of the Tick-borne encephalitis envelope protein domains and the Dengue 2 envelope protein domains. Thus, reference herein to domains of flavivirus proteins, in particular, flaviviruses other than Tick-borne encephalitis flavivirus envelope proteins and Dengue 2 flavivirus envelope proteins, are based on homology to domains in the Tick-borne encephalitis flavivirus envelope protein and the Dengue 2 flavivirus envelope protein.

[0250] The domain III of the envelope protein of the DEN flavivirus encodes the majority of the flavivirus type-specific contiguous critical/dominant neutralizing epitopes (Roehring, J. T., *Adv. Virus Res.* 59:141 (2003)), including the four DEN (DEN1, DEN2, DEN3, DEN4) viruses. Flavivirus envelope proteins are highly homologous. Exemplary envelope protein sequences are SEQ ID NOS: 642, 763, 765, 767, 769 and 774.

[0251] West Nile virus (WNV) is a single-stranded positive sense RNA envelope virus. It was first isolated and identified in the West Nile region of Uganda in 1937 from a febrile female adult (Smithburn, et al., *Am J Trop Med Hyg* 3:9-18 (1954)).

[0252] Japanese encephalitis (JE) virus is localized in Asia and northern Australia (about 50,000 cases with about 10,000 deaths annually).

[0253] The Dengue (DEN) disease is caused by four mosquito-borne, serologically related flaviviruses known as DEN-1 (also referred to herein as "Den1" or Den 1"), DEN-2 (also referred to herein as "Den2" or "Den 2"), DEN-3 (also referred to herein as "Den3" or "Den 3"), and DEN-4 (also referred to herein as "Den4" or Den 4"). The compositions, fusion proteins and polypeptides of the invention can include Den 1 SEQ ID NO.: 623; Den 1 PR 94 (Puerto Rico, 1994) SEQ ID NO: 624; Den 3 SEQ ID NO: 626; and Den 4 SEQ ID NO: 627. SEQ ID NOS: 623, 624, 625, 626 and 627 are portions of domain III of Den1, Den2, Den3 and Den4 flaviviruses.

[0254] "EI," "EII," and "EIII," as used herein, refer to domains I, II and III, respectively, of the West Nile flavivirus envelope protein. "JEI," "JEEI," and "JEIII," as used herein, refer to domains I, II and III, respectively, of the Japanese encephalitis flavivirus envelope protein. "Den1 I," "Den1 II," and "Den1 III," as used herein refer to domains I, II and III, respectively, of the Dengue 1 flavivirus envelope protein. Likewise, designations for the domains of envelope proteins of other flaviviruses are referenced by the flavivirus name followed by the domain number (e.g., (Tick-borne) TBI (Tick-borne), TBII, TBIII, Den2 I, Den2 II, Den2 III).

[0255] The portion of an envelope protein of a flavivirus can include at least one member selected from the group consisting of at least a portion of domain I, at least a portion of domain II and at least a portion of domain III. When a domain is designated with a "+," for example "EIII+" or "JEIII+," the portion of the envelope protein referenced as "III" is one component of the total of that domain plus at least one of at least a portion of either or both of domains I

and II. For example, "EIII+," as used herein, means the compositions, fusion proteins and polypeptides of the invention include domain III and at least a portion of domain I. "EIII+" is also referred to as "EI/III." "JEIII+" is also referred to as "JEI/III." Similarly, when compositions include domains of envelope proteins of flavivirus, the domains can be any combination of domains I, II, and III and can be designated based on the domain. For example, EI/II includes domain I and II of the West Nile flavivirus. The absence of a "+" in reference to a domain (e.g., EIII, JEIII, Den1 III) of an envelope protein employed in the compositions, fusion proteins and polypeptides of the invention means that the composition, fusion protein and polypeptide includes the referenced domain. For example, "Den1 III" means the compositions, fusion proteins and compositions include domain III, not domains I and II, of the Dengue 1 virus.

[0256] The West Nile viral envelope protein can include at least a portion of at least one member selected from the group consisting of SEQ ID NO: 610, which is an EIII+ amino acid sequence, the italicized amino acids are domain I of the envelope protein and the remaining sequence is domain III of the envelope protein; SEQ ID NO: 611, West Nile virus, Stanford, Conn., also referred to as "West Nile S"; SEQ ID NO: 612, West Nile virus, New York, N.Y., also referred to as "West Nile NY"; and SEQ ID NO: 613, SEQ ID NO: 610 is encoded by SEQ ID NO: 614.

[0257] The Langat virus envelope protein for use in the compositions, fusion proteins and polypeptides of the invention can include at least a portion of SEQ ID NO: 615. The Kunjin virus envelope protein can include at least a portion of SEQ ID NO: 616. The Murray Valley encephalitis envelope protein can include at least a portion of SEQ ID NO: 617. The Japanese encephalitis envelope protein can include at least one member selected from the group consisting of at least a portion of SEQ ID NO: 618 and SEQ ID NO: 619. The Tick-borne encephalitis envelope protein can include at least a portion of SEQ ID NO: 620. The Yellow fever virus envelope protein can include at least a portion of SEQ ID NO: 621. The envelope protein of a flavivirus can include at least a portion of at least one member selected from the group consisting of SEQ ID NO: 622 and SEQ ID NO: 643. SEQ ID NOS: 615, 616, 617, 618, 619, 620, 621, 622 and 643 are portions of domain III of the viral envelope protein.

[0258] The antigen can be chemically conjugated to flagellin components and Toll-like Receptor agonist components. Chemical conjugation (also referred to herein as "chemical coupling") can include conjugation by a reactive group, such as a thiol group (e.g., a cysteine residue) or by derivatization of a primary (e.g., a amino-terminal) or secondary (e.g., lysine) group. Different crosslinkers can be used to chemically conjugate TLR ligands (e.g., TLR agonists) to proteins (e.g., antigens, compositions of the invention, HA and M2e constructs of the invention) or other molecules (e.g., nucleic acids, polysaccharides). Exemplary cross linking agents are commercially available, for example, from Pierce (Rockland, Ill.). Methods to chemically conjugate the antigen to the flagellin component are well-known and include the use of commercially available cross-linkers, such as those described herein.

[0259] For example, conjugation of peptide or protein antigens to a flagellin component or a Toll-like Receptor

agonist component of the invention can be through at least one cysteine residue of the flagellin component or the Toll-like Receptor component and at least one cysteine residue of a protein (e.g., an influenza antigen, such as HA, M2e) employing established techniques. The protein can be derivatized with a homobifunctional, sulfhydryl-specific crosslinker; the protein is then desalted to remove the unreacted crosslinker; and then the peptide or protein partner added and conjugated via at least one cysteine residue cysteine. Exemplary reagents for use in the conjugation methods can be purchased commercially from Pierce (Rockland, Ill.), for example, BMB (Catalog No: 22331), BMDB (Catalog No: 22332), BMH (Catalog No: 22330), BMOE (Catalog No: 22323), BM[PEO]₃ (Catalog No: 22336), BM[PEO]₄ (Catalog No: 22337), DPDPB (Catalog No: 21702), DTME (Catalog No: 22335), HBVS (Catalog No: 22334).

[0260] Alternatively, cysteine-containing proteins and antigens can also be conjugated to lysine residues on flagellin components, flagellin, Toll-like Receptor agonist components and Toll-like Receptor agonists of the invention. A protein containing no cysteine residues is derivatized with a heterobifunctional amine and sulfhydryl-specific crosslinker. After desalting, the cysteine-containing partner is added and conjugated. Exemplary reagents for use in the conjugation methods can be purchased from Pierce (Rockland, Ill.), for example, AMAS (Catalog No: 22295), BMPA (Catalog No: 22296), BMPS (Catalog No: 22298), EMCA (Catalog No: 22306), EMCS (Catalog No: 22308), GMBS (Catalog No: 22309), KMUA (Catalog No: 22211), LC-SMCC (Catalog No: 22362), LC-SPDP (Catalog No: 21651), MBS (Catalog No: 22311), SATA (Catalog No: 26102), SATP (Catalog No: 26100), SBAP (Catalog No: 22339), SIA (Catalog No: 22349), SIAB (Catalog No: 22329), SMCC (Catalog No: 22360), SMPB (Catalog No: 22416), SMPH (Catalog No: 22363), SMPT (Catalog No: 21558), SPDP (Catalog No: 21857), Sulfo-EMCS (Catalog No: 22307), Sulfo-GMBS (Catalog No: 22324), Sulfo-KMUS (Catalog No: 21111), Sulfo-LC-SPDP (Catalog No: 21650), Sulfo-MBS (Catalog No: 22312), Sulfo-SIAB (Catalog No: 22327), Sulfo-SMCC (Catalog No: 22322), Sulfo-SMPB (Catalog No: 22317), Sulfo-LC-SMPT (Catalog No: 21568).

[0261] Additionally, or alternatively, peptide or protein antigens can also be conjugated to flagellin components or Toll-like Receptor agonist components of the invention via at least one lysine residue on both conjugate partners. The two conjugate partners are combined along with a homobifunctional amine-specific crosslinker. The appropriate hetero-conjugate is then purified away from unwanted aggregates and homo-conjugates. Exemplary reagents for use in the conjugation methods can be purchased from Pierce (Rockland, Ill.), for example, BSOCOES (Catalog No: 21600), BS₃ (Catalog No: 21580), DFDNB (Catalog No: 21525), DMA (Catalog No: 20663), DMP (Catalog No: 21666), DMS (Catalog No: 20700), DSG (Catalog No: 20593), DSP (Catalog No: 22585), DSS (Catalog No: 21555), DST (Catalog No: 20589), DTBP (Catalog No: 20665), DTSSP (Catalog No: 21578), EGS (Catalog No: 21565), MSA (Catalog No: 22605), Sulfo-DST (Catalog No: 20591), Sulfo-EGS (Catalog No: 21566), THPP (Catalog No: 22607).

[0262] Similarly, peptide or protein antigens can be conjugated to flagellin components or Toll-like Receptor agonist components of the invention via at least one carboxyl group (e.g., glutamic acid, aspartic acid, or the carboxy-terminus of the peptide or protein) on one partner and amines on the other partner. The two conjugation partners are mixed together along with the appropriate heterobifunctional crosslinking reagent. The appropriate hetero-conjugate is then purified away from unwanted aggregates and homo-conjugates. Exemplary reagents for use in the conjugation methods can be purchased from Pierce (Rockland, Ill.), for example, AEDP (Catalog No: 22101), EDC (Catalog No: 22980) and TFCS (Catalog No: 22299).

[0263] In addition, carbohydrate antigens can be conjugated to proteins, flagellin components, flagellin, Toll-like Receptor agonist components and Toll-like Receptor agonists of the invention via at least one cysteine residue in the protein employing well-established techniques. For example, the protein is initially derivatized with a heterobifunctional crosslinker containing at least one sulfhydryl-specific group and at least one hydrazide group. The polysaccharide or oligosaccharide is then treated with an oxidizing agent such as sodium meta-periodate to generate terminal aldehyde groups. The oxidized carbohydrate is then added to the derivatized protein and conjugation to the aldehyde occurs via the hydrazide on the crosslinker. Exemplary reagents for use in the conjugation methods can be purchased from Pierce (Rockland, Ill.), for example, BMPH (Catalog No: 22297), EMCH (Catalog No: 22106), KMHU (Catalog No: 22111) and PDPH (Catalog No: 22301).

[0264] Further lipopeptides can be conjugated to protein antigens, flagellin components or Toll-like Receptor components or Toll-like Receptor agonists of the invention via amino-acid side chains on the peptide chain. Similar strategies and reagents, as described above for peptide conjugation to proteins, would be employed.

[0265] TLRs can be activated by nucleic acids. For example, TLR3 is activated by double-stranded (ds) RNA; TLR7 and TLR8 are activated by single-stranded (ss) RNA; and TLR9 is activated by CpG DNA sequences. Several different techniques can be employed to conjugate Toll-like Receptor agonist components of nucleic acid TLRs to protein antigens. For example, for un-modified DNA molecule, the 5' phosphate group can be modified with the water-soluble carbodiimide EDC (Pierce; Rockford, Ill., Catalog No: 22980) followed by imidazole to form a terminal phosphoylimidazolide. A terminal amine can then be substituted for this reactive group by the addition of ethylenediamine. The amine-modified nucleic acid can then be conjugated to a cysteine on a protein using a heterobifunctional maleimide (cysteine-specific) and NHS-ester (lysine-specific) crosslinker, as described above for peptide-protein conjugation.

[0266] Alternatively, or additionally, a sulfhydryl group can be incorporated at the 5' end by substituting cystamine for ethylenediamine in the second step. After reduction of the disulfide bond, the free sulfhydryl can be conjugated to cysteines on proteins, flagellin components, flagellin, Toll-like Receptor agonist components and Toll-like Receptor agonists employing a homo-bifunctional maleimide-based crosslinker, or to lysines using a heterobifunctional crosslinker, as described above. Alternately, or additionally,

synthetic oligonucleotides can be synthesized with modified bases on either the 5' or 3' end. These modified bases can include primary amine or sulfhydryl groups which can be conjugated to proteins using the methods described above. The 3' end of RNA molecules may be chemically modified to allow coupling with proteins or other macromolecules. The diol on the 3'-ribose residue may be oxidized using sodium meta-periodate to produce an aldehyde group. The aldehyde may then be conjugated to proteins using a hydrazide-containing crosslinker, such as MPBH (Pierce; Rockland, Ill., Catalog No: 22305), which covalently modifies the carbonyl group, and then conjugates to free thiols on proteins via a maleimide group. Alternatively, the 3' hydroxyl group may be derivatized directly with an isocyanate-containing crosslinker, such as PMPI (Pierce; Rockland, Ill., Catalog No: 28100), which also contains a maleimide group for conjugation to protein sulfhydryls.

[0267] Synthetic small-molecule TLR ligands (e.g., Toll-like Receptor agonists, Toll-like Receptor antagonists) have been identified. For example, imiquimod, which potently activates TLR7, and resiquimod, an activator of both TLR7 and TLR8. Analogs of these compounds with varying levels of potency and specificity have been synthesized. The ability to conjugate a small molecule TLR agonist to an antigen of interest can depend on the chemical nature of the TLR ligand. Some TLR ligands may have active groups that can be exploited for chemical conjugation. For example, imiquimod, as well as its analog gardiquimod (InvivoGen; San Diego, Calif.) and the TLR7-activating adenine analog CL087 (InvivoGen, San Diego, Calif.), have primary amine groups ($-\text{NH}_2$), which can be targets for derivatization by crosslinkers containing imidoesters or NHS esters. In another strategy, gardiquimod contains an exposed hydroxyl ($-\text{OH}$) group, which can be derivatized by an isocyanate-containing crosslinker, such as PMPI (Pierce; Rockland, ILL Catalog No: 28100) for subsequent crosslinking to protein sulfhydryl groups.

[0268] In addition, custom synthesis of derivatives of small-molecule TLR ligands can be arranged in order to attach novel functional groups at different positions in the molecule to facilitate crosslinking. Custom derivatives can also include groups, such as maleimides, which can then be used for direct linking to protein antigens.

[0269] Chemical conjugation of an antigen to the flagellin component can result in increased aqueous solubility of the antigen (e.g., an essentially hydrophobic antigen, such as a maturational cleavage site antigen) as a component of the composition.

[0270] The composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 can include a cysteine residue in the hypervariable region of the flagellin component.

[0271] At least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component. The cysteine residue can substitute for at least one amino acid selected from the group consisting of amino acid 1, 237, 238, 239, 240, 241 and 495 of SEQ ID NO: 812; at least one amino acid selected from the group consisting of amino acid 1, 240, 241, 242, 243, 244 and 505 of SEQ ID NO: 498; at least one amino acid

selected from the group consisting of amino acid 1, 237, 238, 239, 240, 241 and 504 of SEQ ID NO: 504; at least one amino acid selected from the group consisting of amino acid 1, 211, 212, 213 and 393 of SEQ ID NO: 815; at least one amino acid selected from the group consisting of amino acid 1, 151, 152, 153, 154 and 287 of SEQ ID NO: 820; at least one amino acid selected from the group consisting of amino acid 1, 238, 239, 240, 241, 242, 243 and 497 of SEQ ID NO: 502; at least one amino acid selected from the group consisting of amino acid 1, 237, 238, 239, 240, 241 and 495 of SEQ ID NO: 812.

[0272] The flagellin component or Toll-like Receptor agonist component can include at least a portion of a naturally occurring flagellin amino acid sequence in combination with the cysteine residue.

[0273] The composition of the invention wherein the cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component, or wherein the flagellin component includes at least a portion of a naturally occurring flagellin amino acid sequence in combination with the cysteine residue, can activate a Toll-like Receptor 5. For example, a cysteine residue can be placed within the D1/D2 domain proximate to the amino-terminus and carboxy-terminus, remote to the TLR5 recognition site (see, for example, FIGS. 75 and 76). Alternatively, or additionally, the cysteine residue can be placed at the distal point of the hypervariable domain (see, for example, FIGS. 75 and 76) at about amino acid 237, about 238, about 239, about 240 and about 241 of SEQ ID NO: 812. Substituting polar or charged amino acids is preferable to substituting hydrophobic amino acids with cysteine residues. Substitution within the TLR5 recognition site is least preferable.

[0274] Flagellin from *Salmonella typhimurium* STF1 (FliC) is depicted in SEQ ID NO: 812 (Accession No: P06179). The TLR5 recognition site is amino acid about 79 to about 117 and about 408 to about 439. Cysteine residues can substitute for or be included in combination with amino acid about 408 to about 439 of SEQ ID NO: 812; amino acids about 1 and about 495 of SEQ ID NO: 812; amino acids about 237 to about 241 of SEQ ID NO: 812; and/or amino acids about 79 to about 117 and about 408 to about 439 of SEQ ID NO: 812.

[0275] *Salmonella typhimurium* flagellin STF2 (FljB) is depicted in SEQ ID NO: 498. The TLR5 recognition site is amino acids about 80 to about 118 and about 420 to about 451 of SEQ ID NO: 498. Cysteine residues can substitute for or be included in combination with amino acids about 1 and about 505 of SEQ ID NO: 498; amino acids about 240 to about 244 of SEQ ID NO: 498; amino acids about 79 to about 117 and/or about 419 to about 450 of SEQ ID NO: 498.

[0276] *Salmonella muenchen* flagellin is depicted in SEQ ID NO: 504 (Accession No: #P06179). The TLR5 recognition site is amino acids about 79 to about 117 and about 418 to about 449 of SEQ ID NO: 504. Cysteine residues can substitute for or be included in combination with amino acids about 1 and about 504 of SEQ ID NO: 504; about 237 to about 241 of SEQ ID NO: 504; about 79 to about 117; and/or about 418 to about 449 of SEQ ID NO: 504.

[0277] *Escherichia coli* flagellin is depicted in SEQ ID NO: 502 (Accession No: P04949). The TLR5 recognition

site is amino acids about 79 to about 117 and about 410 to about 441 of SEQ ID NO: 502. Cysteine residues can substitute for or be included in combination with amino acids about 1 and about 497 of SEQ ID NO: 502; about 238 to about 243 of SEQ ID NO: 502; about 79 to about 117; and/or about 410 to about 441 of SEQ ID NO: 502.

[0278] *Pseudomonas aeruginosa* flagellin is depicted in SEQ ID NO: 815. The TLR5 recognition site is amino acids about 79 to about 114 and about 308 to about 338 of SEQ ID NO: 815. Cysteine residues can substitute for or be included in combination with amino acids about 1 and about 393 of SEQ ID NO: 815; about 211 to about 213 of SEQ ID NO: 815; about 79 to about 114; and/or about 308 to about 338 of SEQ ID NO: 815.

[0279] *Listeria monocytogenes* flagellin is depicted in SEQ ID NO: 820. The TLR5 recognition site is amino acids about 78 to about 116 and about 200 to about 231 of SEQ ID NO: 820. Cysteine residues can substitute for or be included in combination with amino acids about 1 and about 287 of SEQ ID NO: 820; about 151 to about 154 of SEQ ID NO: 820; about 78 to about 116; and/or about 200 to about 231 of SEQ ID NO: 820.

[0280] Experimentally defined TLR5 recognition sites on STF2 have been described (see, for example, Smith, K. D., et al., *Nature Immunology* 4:1247-1253 (2003) at amino acids about 79 to about 117 and about 420 to about 451. In addition, Smith, K. D., et al., *Nature Immunology* 4:1247-1253 (2003), based on sequence homology, identified TLR5 recognition sites on other flagellins, such as STF1 at amino acids about 79 to about 117, about 408 to about 439; *P. aeruginosa* at amino acids about 79 to about 117, about 308 to about 339; *L. pneumophila* at amino acids about 79 to about 117, about 381 to about 419; *E. coli* at amino acids about 79 to about 117, about 477, about 508; *S. marcescens* at amino acids about 79 to about 117, about 265-about 296; *B. subtilis* at amino acids about 77 to about 117, about 218 to about 249; and *L. monocytogenes* at amino acids about 77 to about 115, about 200 to about 231.

[0281] The high-resolution structure STF1 (FliC) (SEQ ID NO: 812) has been determined and can be a basis for analysis of TLR5 recognition by a flagellin and the location of cysteine substitutions/additions. Flagellin resembles a “boomerang,” with the amino- and carboxy-termini at the end of one arm (see, for example FIG. 77). The TLR5 recognition site is located roughly on the outer side of the boomerang just below the bend toward the amino- and carboxy-termini of the flagellin. The hinge region, which is not required for TLR5 recognition, is located above the bend. The region of greatest sequence homology of flagellins is in the TLR5 recognition site. The next region of sequence homology is in the D1 and D2 domains, which include the TLR5 recognition site and the amino- and carboxy-termini. The D1 and D2 domains, with or without a linker, is STF2Δ (SEQ ID NO: 500), which can activate TLR5. The region of least sequence homology between flagellins is the hypervariable region.

[0282] It is believed that the ability of the flagellin component or Toll-like Receptor agonist component to activate TLR5 can be accomplished by maintaining the conjugation sites (cysteine residues substituted for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component or at least a portion of a naturally

occurring flagellin amino acid sequence in combination with the cysteine residue) remote from the TLR5 or TLR recognition site. For example, for STF1 (SEQ ID NO: 812), for which a high resolution structural determination is available, this may be achieved in the D1 domain, D2 domain or in the hinge region. In the D1/D2 domain the amino- and carboxy-termini can be remote (also referred to herein as “distal”) from the TLR5 recognition site, and moving away from either the amino or carboxy terminus may bring the conjugation site closer to the recognition site and may interfere with TLR5 activity. In the hinge region amino acids about 237 to about 241 of SEQ ID NO: 812, are approximately at the other tip of the “boomerang” and are about the same distance from the TLR5 recognition site as the amino- and carboxy-termini. This site may also be a location that maintains TLR5 recognition.

[0283] Amino acid identity can be taken into consideration for the location of conjugation sites. Polar and charged amino acids (e.g., serine, aspartic acid, lysine) are more likely to be surface exposed and amenable to attachment of an antigen. Hydrophobic amino acids (e.g., valine, phenylalanine) are more likely to be buried and participate in structural interactions and should be avoided.

[0284] Compositions that include flagellin components with cysteine residues or Toll-like Receptor agonist components with cysteine residues activate TLR5 and can be chemically conjugated to antigens.

[0285] The compositions and methods of employing the compositions of the invention can further include a carrier protein. The carrier protein can be at least one member selected from the group consisting of a tetanus toxoid, a *Vibrio cholerae* toxoid, a diphtheria toxoid, a cross-reactive mutant of diphtheria toxoid, a *E. coli* B subunit of a heat labile enterotoxin, a tobacco mosaic virus coat protein, a rabies virus envelope protein, a rabies virus envelope glycoprotein, a thyroglobulin, a heat shock protein 60, a key-hole limpet hemocyanin and an early secreted antigen tuberculosis-6.

[0286] The composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5 can include at least one lysine of the flagellin component that has been substituted with at least one member selected from the group consisting of an arginine residue, a serine residue and a histidine residue.

[0287] In an additional embodiment, the invention is a composition comprising a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor. “Component,” as used herein in reference to a Toll-like Receptor agonist component, means at least part of or the entirety of a Toll-like Receptor agonist.

[0288] In one embodiment, the cysteine residue in the composition comprising a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component

includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor, substitutes for at least one amino acid in a naturally occurring amino acid sequence of a Toll-like Receptor agonist component. The cysteine can substitute for at least one amino acid remote to the Toll-like Receptor recognition site of the Toll-like Receptor agonist component.

[0289] In another embodiment, the Toll-like Receptor agonist component includes at least a portion of a naturally occurring Toll-like Receptor agonist amino acid sequence in combination with a cysteine residue. The cysteine residue in combination with the naturally occurring Toll-like Receptor agonist can be remote to the Toll-like Receptor recognition site of the Toll-like Receptor agonist component.

[0290] The composition comprising a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor can further include at least a portion of at least one antigen (e.g., an influenza antigen, such as an influenza integral membrane protein antigen, HA, HA1-1, HA1-2, M2, M2e).

[0291] In still another embodiment, the invention is a composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5.

[0292] "Substituted," as used herein in reference to the flagellin, flagellin component, Toll-like Receptor agonist or Toll-like Receptor agonist component, means that at least one amino acid, such as a lysine of the flagellin component, has been modified to another amino acid residue, for example, a conservative substitution (e.g., arginine, serine, histidine) to thereby form a substituted flagellin component or substituted Toll-like Receptor agonist component. The substituted flagellin component or substituted Toll-like Receptor agonist component can be made by generating recombinant constructs that encode flagellin with the substitutions, by chemical means, by the generation of proteins or peptides of at least a portion of the flagellin by protein synthesis techniques, or any combination thereof.

[0293] The lysine residue that is substituted with an amino acid (e.g., arginine, serine, histidine) can be at least one lysine residue selected from the group consisting of lysine 19, 41, 58, 135, 160, 177, 179, 203, 215, 221, 228, 232, 241, 251, 279, 292, 308, 317, 326, 338, 348, 357, 362, 369, 378, 384, 391 and 410 of SEQ ID NO: 812.

[0294] The flagellin can be a *S. typhimurium* flagellin that includes SEQ ID NO: 816. The lysine residue that is substituted with an amino acid (e.g., arginine, serine, histidine) can be at least one lysine residue selected from the group consisting of lysine 20, 42, 59, 136, 161, 177, 182, 189, 209, 227, 234, 249, 271, 281, 288, 299, 319, 325, 328, 337, 341, 355, 357, 369, 381, 390, 396, 403, 414 and 422 of SEQ ID NO: 816.

[0295] The flagellin can be an *E. coli* fliC that includes SEQ ID NO: 814. The flagellin can be a *S. muenchen* that

include the includes SEQ ID NO: 813. The flagellin can be a *P. aeruginosa* flagellin that includes SEQ ID NO: 815. The flagellin can be a *Listeria monocytogenes* flagellin that includes SEQ ID NO: 820.

[0296] The compositions of the invention can include a flagellin that has lysines substituted in a region adjacent to the motif C of the flagellin, the motif N of the flagellin, both the motif C and the motif N of the flagellin, domain 1 of the flagellin, domain 2 of the flagellin or any combination thereof. Motif C and motif N of flagellin and domain 1 and domain 2 of the flagellin can be involved in activation of TLR5 by the flagellin (Murthy, et al., *J. Biol. Chem.*, 279:5667-5675 (2004)).

[0297] Chemical conjugation of a protein, peptide, or polypeptide to another molecule can be by derivatization of a secondary group, such as a lysine. Certain lysine residues in flagellin are near or in domain 1, the motif C or motif N, motifs can be important in binding of the flagellin to TLR5. For example, lysine residues at amino acids 58, 135, 160 and 410 of SEQ ID NO: 812 may be substituted with at least one member selected from the group consisting of an arginine residue, a serine residue and a histidine residue. Derivatization of such lysine residues to, for example, chemically conjugated antigens to flagellins, may decrease the ability or the binding affinity of the flagellin to TLR5 and, thus, diminish an innate immune response mediated by TLR5. Substitution of at least one lysine residue in a flagellin that may be near to regions of the flagellin that are important in mediating interactions with TLR5 (e.g., motif C, motif N, domain 1) with another amino acid (e.g., arginine, serine, histidine) may preserve or enhance flagellin binding to TLR5. In a particular embodiment, the amino acid substitution is a conservative amino acid substitution with at least one member selected from the group consisting of arginine, serine and histidine. Exemplary commercially available reagents for chemical conjugation are described herein.

[0298] Certain lysine residues in flagellin are in the domain (domain 1) and can be important for activation of TLR5. For example, lysine residues at positions 58, 135, 160 and 410 of SEQ ID NO: 812 are in domain 1. Derivatization of such lysine residues to, for example, chemically conjugated antigens, may decrease TLR5 bioactivity and, thus, diminish an innate immune response mediated by TLR5.

[0299] Lysine residues that can be substituted can include lysine residues implicated in TLR5 activation. Lysine residues in motif N (amino acids 95-108 of SEQ ID NO: 812) and/or motif C (amino acids 441-449 of SEQ ID NO: 812) can be suitable for substitution. Substitution of certain lysine residue in the flagellin (e.g., lysine at amino acid position 19, 41) with, for example, an arginine, serine or histidine, can maintain binding of the flagellin to TLR5 and leave other lysines available for chemical conjugate to another molecule, such as an antigen (e.g., protein) or another molecule, such as another protein, peptide or polypeptide.

[0300] The X-ray crystal structure of the F41 fragment of flagellin from *Salmonella typhimurium* shows the domain structure of flagellin (Samatey, F. A., et al., *Nature* 410:321 (2001)). The full length flagellin protein contains 4 domains, designated as D0, D1, D2 and D3. Three of these domains are shown in the crystal structure because the structure was made with a proteolytic fragment of full length flagellin. The

amino acid sequences of *Salmonella typhimurium* flagellin for these regions, numbered relative to SEQ ID NO: 812 are as follows:

[0301] D0 contains the regions A1 through A55 and S451 through R494

[0302] D1 contains the regions N56 through Q176 and T402 through R450

[0303] D2 contains the regions K177 through GI 89 and A284 through A401

[0304] D3 contains the region Y190 though V283

[0305] Exemplary lysine residues of SEQ ID NO: 812 suitable for substitution with, for example, arginine, histidine, or serine, can include:

[0306] D0 contains 2 lysine residues; K19, K41

[0307] D1 contains 4 lysine residues; K58, K135, K160 and K410

[0308] D2 contains 14 lysine residues at positions 177, 179, 292, 308, 317, 326, 338, 348, 357, 362, 369, 378, 384, 391

[0309] D3 contains 8 lysine residues at positions 203, 215, 221, 228, 232, 241, 251, 279

[0310] Exemplary lysine residues suitable for substitution include lysines at positions 58, 135, 160 and 410 of SEQ ID NO: 812 (Jacchieri, S. G., et. al., *J. Bacteriol.* 185:4243 (2003); Donnelly, M. A., et al., *J. Biol. Chem.* 277:40456 (2002)). The sequences were obtained from the Swiss-Prot Protein Knowledge base located online at <http://us.expasy.org/sprot/>. Lysine residues that can be modified are indicated with a *.

[0311] Exemplary lysine residues of SEQ ID NO: 816 suitable for substitution can include:

[0312] D0—with two lysines at positions 20, 42;

[0313] D1—with five lysines at positions 59, 136, 161, 414, 422;

[0314] D2—with sixteen lysines at positions 177, 182, 189, 299, 319, 325, 328, 337, 341, 355, 357, 369, 381, 390, 396, 403 and

[0315] D3—with seven lysines at positions 209, 227, 234, 249, 271, 281, 288.

[0316] In a further embodiment, the invention is a composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0317] In yet another embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

[0318] In another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that

includes a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

[0319] In yet another embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5.

[0320] In still another embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0321] In another embodiment, the invention is a method of stimulating a protective immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0322] In an additional embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0323] In a further embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5.

[0324] Another embodiment of the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor agonist.

[0325] Another embodiment of the invention is a method of stimulating a protective immunity in a subject, comprising the step of administering to the subject a composition that includes a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does

not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor agonist.

[0326] In yet another embodiment, the invention is a composition comprising an antigen component that includes at least a portion of a hemagglutinin maturational cleavage site and an agonist component that includes a Toll-like Receptor agonist (e.g., at least one member selected from the group consisting of a Toll-like Receptor 1 agonist, a Toll-like Receptor 3 agonist, a Toll-like Receptor 5 agonist, a Toll-like Receptor 6 agonist, a Toll-like Receptor 7 agonist and a Toll-like Receptor 9 agonist) wherein the Toll-like Receptor agonist is not a Toll-like Receptor 2 agonist (e.g., the outer membrane protein complex (OMPC), such as OMPC of *Neisseria meningitidis*). The hemagglutinin maturational cleavage site can include at least one member selected from the group consisting of an influenza A hemagglutinin maturational cleavage site, an influenza B hemagglutinin maturational cleavage site and an influenza C hemagglutinin maturational cleavage site.

[0327] In one embodiment, the antigen component and the agonist component of the compositions of the invention can be components of a fusion protein. In another embodiment, the antigen component is chemically conjugated to the agonist component. The agonist component includes a composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5. The antigen component further includes at least a portion of a matrix 2 protein, which can further include a second agonist component that includes at least a portion of a second Toll-like Receptor agonist (e.g., at least a portion of at least one member selected from the group consisting of a Toll-like Receptor 2 agonist, a Toll-like Receptor 3 agonist, a Toll-like Receptor 4 agonist, a Toll-like Receptor 5 agonist, a Toll-like Receptor 6 agonist, a Toll-like Receptor 7 agonist and a Toll-like Receptor 9 agonist). The matrix-2 protein can be fused to the second agonist component or chemically conjugated to a second agonist component.

[0328] An "antigen component," as used herein, refers to a part or the entirety of an antigen.

[0329] An "agonist component" as used herein, refers to a part or the entirety of an agonist, such as a Toll-like Receptor agonist.

[0330] In an additional embodiment, the invention is a method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition comprising an antigen component that includes at least a portion of a hemagglutinin maturational cleavage site and an agonist component that includes a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist is not a Toll-like Receptor 2 agonist.

[0331] In a further embodiment, the invention is a method of stimulating protective immunity in a subject, comprising the step of administering to the subject a composition comprising an antigen component that includes at least a portion of a hemagglutinin maturational cleavage site and an agonist component that includes a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist is not a Toll-like Receptor 2 agonist.

[0332] In yet another embodiment, the invention is a composition that includes at least a portion of a HA antigen (e.g., the maturational cleavage site) that is fused (e.g., recombinantly) to or chemically conjugated to at least a portion of a hinge region of a flagellin, a flagellin component or in a flagellin in which the hinge region has been deleted; and at least a portion of an M2e protein that is fused (e.g., recombinantly) to or chemically conjugated to a flagellin or a flagellin component, for example, at the amino- or carboxy-terminus of the flagellin or the flagellin component (See, for example, FIG. 77).

[0333] In an additional embodiment, the invention includes a protein, peptide polypeptide having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98% and at least about 99% sequence identity to the proteins, polypeptides and peptides of the invention.

[0334] The percent identity of two amino acid sequences (or two nucleic acid sequences) can be determined by aligning the sequences for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first sequence). The amino acid sequence or nucleic acid sequences at corresponding positions are then compared, and the percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity=# of identical positions/total # of positions $\times$ 100). The length of the protein or nucleic acid encoding can be aligned for comparison purposes is at least 30%, preferably, at least 40%, more preferably, at least 60%, and even more preferably, at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100%, of the length of the reference sequence, for example, the nucleic acid sequence of a protein portion of HA (e.g., HA1-1, HA1-2), fusion protein, antigen, or polypeptide.

[0335] The actual comparison of the two sequences can be accomplished by well-known methods, for example, using a mathematical algorithm. A preferred, non-limiting example of such a mathematical algorithm is described in Karlin et al. (*Proc. Natl. Acad. Sci. USA*, 90:5873-5877 (1993), the teachings of which are hereby incorporated by reference in its entirety). Such an algorithm is incorporated into the BLASTN and BLASTX programs (version 2.2) as described in Schaffer et al. (*Nucleic Acids Res.*, 29:2994-3005 (2001), the teachings of which are hereby incorporated by reference in its entirety). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., BLASTN; available at the Internet site for the National Center for Biotechnology Information) can be used. In one embodiment, the database searched is a non-redundant (NR) database, and parameters for sequence comparison can be set at: no filters; Expect value of 10; Word Size of 3; the Matrix is BLOSUM62; and Gap Costs have an Existence of 11 and an Extension of 1.

[0336] Another mathematical algorithm employed for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989), the teachings of which are hereby incorporated by reference in its entirety. Such an algorithm is incorporated into the ALIGN program (version 2.0), which is part of the GCG (Accelrys, San Diego, Calif.) sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12,

and a gap penalty of 4 is used. Additional algorithms for sequence analysis are known in the art and include ADVANCE and ADAM as described in Torellis and Robotti (Comput. Appl. Biosci., 10: 3-5 (1994), the teachings of which are hereby incorporated by reference in its entirety); and FASTA described in Pearson and Lipman (*Proc. Natl. Acad. Sci. USA*, 85: 2444-2448 (1988), the teachings of which are hereby incorporated by reference in its entirety).

[0337] The percent identity between two amino acid sequences can also be accomplished using the GAP program in the GCG software package (Accelrys, San Diego, Calif.) using either a Blossum 63 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. In yet another embodiment, the percent identity between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package (Accelrys, San Diego, Calif.), using a gap weight of 50 and a length weight of 3.

[0338] The nucleic acid sequence encoding a protein portion of HA, polypeptide or fusion proteins of the invention and polypeptides of the invention can include nucleic acid sequences that hybridize to nucleic acid sequences or complements of nucleic acid sequences of the invention, for example, SEQ ID NOS: 53-58, 64, 68, 71, 72, 73, 76, 78, 80, 84, 87, 104, 107, 110, 111, 112, 115, 116, 117, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 137, 139, 142, 143, 146, 147, 150, 167, 184, 185, 186, 187, 188, 189, 190, 191, and 192); and nucleic acid sequences that encode amino acid sequences and fusion proteins of the invention (e.g., SEQ ID NOS: 89-92, 95, 151-160, 177, 209, 210 and 211) under selective hybridization conditions (e.g., highly stringent hybridization conditions). As used herein, the terms "hybridizes under low stringency," "hybridizes under medium stringency," "hybridizes under high stringency," or "hybridizes under very high stringency conditions," describe conditions for hybridization and washing of the nucleic acid sequences. Guidance for performing hybridization reactions, which can include aqueous and nonaqueous methods, can be found in Ausubel, F. M., et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (2001), the teachings of which are hereby incorporated herein in its entirety.

[0339] For applications that require high selectivity, relatively high stringency conditions to form hybrids can be employed. In solutions used for some membrane based hybridizations, addition of an organic solvent, such as formamide, allows the reaction to occur at a lower temperature. High stringency conditions are, for example, relatively low salt and/or high temperature conditions. High stringency are provided by about 0.02 M to about 0.10 M NaCl at temperatures of about 50° C. to about 70° C. High stringency conditions allow for limited numbers of mismatches between the two sequences. In order to achieve less stringent conditions, the salt concentration may be increased and/or the temperature may be decreased. Medium stringency conditions are achieved at a salt concentration of about 0.1 to 0.25 M NaCl and a temperature of about 37° C. to about 55° C., while low stringency conditions are achieved at a salt concentration of about 0.15 M to about 0.9 M NaCl, and a temperature ranging from about 20° C. to about 55° C. Selection of components and conditions for hybridization are well known to those skilled in the art and are reviewed

in Ausubel et al. (1997, *Short Protocols in Molecular Biology*, John Wiley & Sons, New York N.Y., Units 2.8-2.11, 3.18-3.19 and 4-64.9).

[0340] A "subject," as used herein, can be a mammal, such as a primate or rodent (e.g., rat, mouse). In a particular embodiment, the subject is a human.

[0341] An "effective amount," when referring to the amount of a composition and fusion protein of the invention, refers to that amount or dose of the composition and fusion protein, that, when administered to the subject is an amount sufficient for therapeutic efficacy (e.g., an amount sufficient to stimulate an immune response in the subject). The compositions and fusion proteins of the invention can be administered in a single dose or in multiple doses.

[0342] The methods of the present invention can be accomplished by the administration of the compositions and fusion proteins of the invention by enteral or parenteral means. Specifically, the route of administration is by oral ingestion (e.g., drink, tablet, capsule form) or intramuscular injection of the composition and fusion protein. Other routes of administration as also encompassed by the present invention including intravenous, intradermal, intraarterial, intraperitoneal, or subcutaneous routes, and nasal administration. Suppositories or transdermal patches can also be employed.

[0343] The compositions and fusion proteins of the invention can be administered ex vivo to a subject's autologous dendritic cells. Following exposure of the dendritic cells to the composition and fusion protein of the invention, the dendritic cells can be administered to the subject.

[0344] The compositions and fusion proteins of the invention can be administered alone or can be coadministered to the patient. Coadministration is meant to include simultaneous or sequential administration of the composition, fusion protein or polypeptide of the invention individually or in combination. Where the composition and fusion protein are administered individually, the mode of administration can be conducted sufficiently close in time to each other (for example, administration of the composition close in time to administration of the fusion protein) so that the effects on stimulating an immune response in a subject are maximal. It is also envisioned that multiple routes of administration (e.g., intramuscular, oral, transdermal) can be used to administer the compositions and fusion proteins of the invention.

[0345] The compositions and fusion proteins of the invention can be administered alone or as admixtures with conventional excipients, for example, pharmaceutically, or physiologically, acceptable organic, or inorganic carrier substances suitable for enteral or parenteral application which do not deleteriously react with the extract. Suitable pharmaceutically acceptable carriers include water, salt solutions (such as Ringer's solution), alcohols, oils, gelatins and carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, and polyvinyl pyrrolidone. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like which do not deleteriously react with the compositions, fusion proteins or polypeptides of the invention. The preparations can also be combined, when desired, with other active substances to reduce metabolic degrada-

tion. The compositions and fusion proteins of the invention can be administered by is oral administration, such as a drink, intramuscular or intraperitoneal injection or intranasal delivery. The compositions and fusion proteins alone, or when combined with an admixture, can be administered in a single or in more than one dose over a period of time to confer the desired effect (e.g., alleviate prevent viral infection, to alleviate symptoms of virus infection, such as influenza or flaviviral infection).

[0346] When parenteral application is needed or desired, particularly suitable admixtures for the compositions and fusion proteins are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. In particular, carriers for parenteral administration include aqueous solutions of dextrose, saline, pure water, ethanol, glycerol, propylene glycol, peanut oil, sesame oil, polyoxyethylene-block polymers, and the like. Ampules are convenient unit dosages. The compositions, fusion proteins or polypeptides can also be incorporated into liposomes or administered via transdermal pumps or patches. Pharmaceutical admixtures suitable for use in the present invention are well-known to those of skill in the art and are described, for example, in *Pharmaceutical Sciences* (17th Ed., Mack Pub. Co., Easton, Pa.) and WO 96/05309 the teachings of which are hereby incorporated by reference.

[0347] The compositions and fusion proteins of the invention can be administered to a subject on a support that presents the compositions, fusion proteins and polypeptides of the invention to the immune system of the subject to generate an immune response in the subject. The presentation of the compositions, fusion proteins and polypeptides of the invention would preferably include exposure of antigenic portions of the viral protein to generate antibodies. The components (e.g., PAMP and a viral protein) of the compositions, fusion proteins and polypeptides of the invention are in close physical proximity to one another on the support. The compositions and fusion proteins of the invention can be attached to the support by covalent or noncovalent attachment. Preferably, the support is biocompatible. "Biocompatible," as used herein, means that the support does not generate an immune response in the subject (e.g., the production of antibodies). The support can be a biodegradable substrate carrier, such as a polymer bead or a liposome. The support can further include alum or other suitable adjuvants. The support can be a virus (e.g., adenovirus, poxvirus, alphavirus), bacteria (e.g., *Salmonella*) or a nucleic acid (e.g., plasmid DNA).

[0348] The dosage and frequency (single or multiple doses) administered to a subject can vary depending upon a variety of factors, including prior exposure to an antigen, a viral protein, the duration of viral infection, prior treatment of the viral infection, the route of administration of the composition, fusion protein or polypeptide; size, age, sex, health, body weight, body mass index, and diet of the subject; nature and extent of symptoms of viral exposure, viral infection and the particular viral responsible for the infection (e.g., a flavivirus, influenza virus), or treatment or infection of another antigen, such as an influenza antigen, kind of concurrent treatment, complications from the viral exposure, viral infection or exposure or other health-related problems. Other therapeutic regimens or agents can be used in conjunction with the methods and compositions, fusion

proteins or polypeptides of the present invention. For example, the administration of the compositions and fusion proteins can be accompanied by other viral therapeutics or use of agents to treat the symptoms of a condition associated with or consequent to exposure to the antigen, such as flavivirus infection (e.g., high fever, numbness, DHF, meningoencephalitis) or influenza infection, for example. Adjustment and manipulation of established dosages (e.g., frequency and duration) are well within the ability of those skilled in the art.

[0349] Influenza viruses are single-stranded RNA viruses that belong to the viral family Orthomyxoviridae. Influenza viruses are divided into three types (A, B, C) determined by the antigenic differences in ribonucleoprotein (RNP) and matrix (M) antigens of the viruses. Influenza A virus naturally infects humans and several other mammalian species, including swine and horses, and a wide variety of avian species, and causes epidemics and pandemics in the human population. Influenza B virus appears to naturally infect only humans and can cause epidemics in humans. Influenza C virus has been isolated from humans and swine, but generally does not occur in epidemics and usually results in mild disease in humans.

[0350] Mature influenza virions are enveloped with a pleomorphic structure ranging in diameter from 80 to 120 nm. The single-stranded RNA genome is closely associated with a helical nucleoprotein and is present in seven (influenza C) or eight (influenza A and B) separate segments of ribonucleoprotein (RNP), each of which has to be present for successful replication of the virus. The segmented genome is enclosed within an outer lipoprotein envelope. Matrix protein 1 (MP1 or also referred to herein as "M1") lines the inside of the outer lipoprotein envelope and is bound to the RNP.

[0351] Hemagglutinin (HA) is a surface glycoprotein on a virus (e.g., an influenza virus) that is responsible for binding to N-AcetylNeuraminic Acid (NeuNAc; also referred to herein as "sialic acid") on host cells and subsequent fusion of viral and host membranes. HA acquired its name by virtue of its ability to cause red blood cells to clump, or agglutinate. Influenza HA is a trimer consisting of the three monomeric (HA0) subunits. HA performs two critical functions during the infection process: binding to a cell surface sialyloligosaccharide receptor and fusion of virus and host cell membrane. Following binding of the HA trimer to the plasma membrane of a host cell, the host cell membrane engulfs the virus in an endosome and attempts to digest the contents of the endosome by acidifying its interior and transferring it to a lysosome in the host cell. However, the acidic environment of the lysosome destabilizes HA, resulting in partial unfolding of HA0 which exposes a protease-sensitive site (the maturational cleavage site) that is cleaved by a host protease to form HA1 and HA2 subunits which are connected by a single disulfide bond (Wiley, D. C., et al., *Annu. Rev. Biochem.* 56:365-394 (1987)). Cleavage occurs at a specific amino acid residue and generates a hydrophobic amino terminus for the HA2 subunit. This hydrophobic terminus of HA2 mediates fusion between the viral envelope and the endosomal membrane of the host cell and releases the contents of the virion into the cytoplasm of the cell, a process known as uncoating. Thus, cleavage of the HA polypeptide is a requirement for infectivity.

[0352] The crystal structure of several viral hemagglutinins has been determined (see, for example, Wilson, I. A., et al., *Nature* 289:366-373 (1981); Chen, J., et al., *Cell* 95:409-417 (1998); Ha, Y., et al., *The EMBO Journal* 21: 865-875 (2002); Russell, R. J., et al., *Virology* 325:287-296 (2004); and Cox, N. J., et al., In: Topley and Wilson's Microbiology and Microbial Infections, eds. B. W. J. Mathy, et al., Vol. 1 (9th ed.) New York, N.Y., Oxford Univ. Press, Ch. 32, p. 634 (1998)). X-ray crystallographic structures show that HA is folded into two structural components or domains—a globular head and a fibrous stalk (see, for example, FIG. 1). The globular head includes HA1, including that part of HA1 that binds to sialic acid (also referred to as the “receptor binding site or domain” or “sialic acid binding site or domain”), and antiparallel β -sheets. The fibrous stalk is more proximal to the viral membrane and consists of all of HA2 and part of HA1, including the cleavage site between HA1 and HA2.

[0353] There are fifteen known subtypes of Influenza A HA (H1-H15) that share between about 40 to about 60% sequence identity (World Health Organization BULL. World Health Organ., 58:585-591 (1980)). Influenza viruses containing all 15 HA subtypes have been isolated from avian species (H5, H7, and H9), equine (H3 and H7), seals (H3, H4 and H7), whales (H1 and H13) and swine (H1, H3, and H9). Subtypes of influenza A virus are generally named according to the particular antigenic determinants of HA (H, 15 major types) and neuraminidase (N, about 9 major types). For example, subtypes include influenza A (H2N1), A(H3N2), A(H5N1), A(H7N2), A(H9N2), A(H1/H0), A(H3/H0) and A(H5/H0). In the last century, three subtypes of influenza A resulted in pandemics: H1 in 1918 and 1977; H2 in 1957 and H3 in 1968. In 1997, an H5 avian virus and in 1999, an H9 virus resulted in outbreaks of respiratory disease in Hong Kong. HA from influenza type B viruses have been isolated from humans and seals and are not divided into subtypes.

[0354] A host infected with influenza can mount an antibody response to the globular head of HA that protects that host from subsequent infection with the same strain of virus by blocking the interaction between HA and the host cell, i.e., neutralizing the infectivity of the virus. Due to the low fidelity and high rate of influenza RNA replication, the virus is constantly experiencing minor mutations in the HA gene that preserve the globular head structure and host cell interaction, but may allow progeny virus to escape immune surveillance. These point mutations are referred to as “antigenic drift.” In addition, if a single host is simultaneously infected with two different strains of influenza A, a new subtype of virus may emerge as a result of reassortment, or the exchange of the RNA segments, or genes, between different strains of influenza A viruses. The viruses emerging from reassortment present the human immune system with a new antigenic experience that usually results in high morbidity and mortality. This type of drastic antigenic change is known as “antigenic shift.” Since type B influenza viruses circulate almost exclusively in humans, these viruses cannot undergo reassortment with animal strains and, thus, are changed only by antigenic drift.

[0355] Immunity to HA can reduce the likelihood of infection and severity of disease if infection does occur. HA is an important antigenic target and the efficacy of vaccines depends on the antigenic match between the vaccine strain and the circulating strain. Since the hemagglutinin protein

readily undergoes antigenic shift and drift in order to evade the host's immune defense, traditional vaccines must be based on currently circulating influenza strains and annually updated. Annual updates of influenza vaccines are not only costly they also require significant amounts of production time and manufacturing infrastructure. A vaccine composition based on invariant regions of the virus may provide broadly cross-reactive protection.

[0356] In contrast to the globular head of HA, changes in amino acid residues surrounding the maturational cleavage site of HA are limited due to functional constraints. Amino acid residues surrounding the HA maturational cleavage site influence recognition and therefore cleavability of the site by the host protease. Since the virus does not code for the protease, changes in the amino acid residues surrounding the maturational cleavage site are restricted. As a consequence a peptide of about 20 amino acids spanning the maturational cleavage site remains genetically stable across influenza viruses of the same HA subtype (WO 2004/080403; Bianchi, et al. *J Virol* 79:7380-7388 (2005)) or as branched peptides (Horvath, et al. *Immunol Letters* 60:127-136 (1998), Nagy, et al. *Scand J Immunol* 40:281-291 (1994)).

[0357] A second highly conserved antigen of influenza A is the ectodomain of the matrix 2 protein (M2e). M2 is a 97-amino acid protein expressed at low levels in mature virions and much higher levels on infected cells. The M2 protein forms a homotetramer that functions as an ion channel which is critical to the replication of the virus, thus, mutations in M2e are not as well tolerated as mutations in HA. The 24-amino acid ectodomain (M2e) is highly conserved across multiple influenza A strains. In mammals, M2e is poorly immunogenic in its native form. Antibodies to M2e can confer passive protection in animal models of influenza A infection (Treanor, J. J., et al., *J Virol* 64:1375 (1990); Liu, W. P., et al., *Immunol Lett* 93:131 (2004)), not by neutralizing the virus and preventing infectivity, but by killing infected cells and disrupting the viral life cycle (Zebedee, S. L., et al., *J. Virol.* 62:2762 (1998); Jegerlehner, A. N., et al., *J Immunol* 172:5598 (2004)), which may be by antibody-dependent NK cell activity (Jegerlehner, A. N., et al., *J Immunol* 172:5598 (2004)). Composition that include M2e proteins may limit the severity of influenza A disease while allowing the host immune response to develop adaptive immunity to the dominant neutralizing influenza antigen, HA.

[0358] Strategies to manage infection and illness consequent to influenza viral infection have not changed significantly in the past four decades. Due to the seasonal nature of influenza disease, the distinct types of influenza virus (A and B) that threaten the human population, and the genetic instability of each type, it is necessary to reformulate a multivalent vaccine each year, based on epidemiological prediction of strains likely to be circulating in the human population in the upcoming flu season. The vaccine is produced from stocks of selected prototype viral strains grown in embryonated chicken eggs. Thus, the current strategy has several limitations that include: (a) dependence on uncertain prediction of circulating strains; (b) dependence on the ability to grow the appropriate strains in chicken eggs; (c) the egg-based production system carries risks of product contamination; (d) the product produced in eggs cannot be used in individuals with egg allergies; and (e) a significant risk that the typical multivalent vaccine will not

confer protection against a pandemic strain of virus to which the human population has no pre-existing immunity.

[0359] The dominant protective component of the currently-available influenza vaccine is the viral hemagglutinin (HA). A more effective vaccine may include not only strain-specific HA, but cross-protective antigens, such as M2e and maturational cleavage site. A vaccine production process that is more reliable, economical, and scaleable than the current egg-based method is also preferable. The compositions, fusion proteins and polypeptides of the invention provide compositions that include HA, M2e, and maturational cleavage site of influenza proteins, which can stimulate an immune response, specifically, a protective immune response to several influenza antigens in subject.

[0360] The teachings of all patents, published applications and references cited herein are incorporated by reference in their entirety.

EXEMPLIFICATION

Example 1

Design of Portions of a Naturally Occurring Hemagglutinin of Influenza A

Materials and Methods

[0361] Design of the HA1-1, 1-2 and 1-3 globular head constructs for A/Puerto Rico/8/34 (H1N1). Influenza strain A/Puerto Rico/8/34 (PR8) is a well-characterized mouse-adapted strain of influenza A virus. The published crystal structure of the mature PR8 HA (SEQ ID NO: 1) (Gamblin et al. 2005. *Science* 303:1838-1842; PDB accession number 1RU7) was used in combination with the Molecular Modeling Database to determine the boundaries for three PR8 globular head constructs. The complete list of solved crystal structures for influenza A hemagglutinin molecules can be found in the Protein Data Bank (PDB) or National Center for Biotechnology Information (NCBI) Structure website (<http://www.ncbi.nlm.nih.gov/Structure>).

[0362] The three dimensional crystal structure for the PR8 HA was viewed using the Cn3D program within the NCBI website. The trimeric HA molecule has the shape of a mushroom. The globular head refers to the portion of the hemagglutinin molecule at the top that resembles the mushroom head, and the coiled-coil stalk refers to the bottom part of the molecule that resembles the stalk of the mushroom (FIG. 1). The globular head contains the substrate binding site that binds to sialic acid on the cell surface, and therefore is important for viral entry. In addition, the majority of neutralizing antibodies epitopes are located near and around this receptor binding site, making the globular head a good target for protective vaccines. The globular head contains most of the HA1 peptide including residues numbered from E51 to K327 in PR8 for example.

[0363] The monomeric structure of PR8 chain A, which encompasses the globular head, was used to guide the design of the PR8 HA1-1, 1-2 and 1-3 constructs (FIG. 1). The HA constructs were designed with domain boundaries within the globular head so that when the molecule is expressed in a host cell, the encoded protein it may spontaneously fold or refolded in vitro to mimic the native conformation. A structural domain (also referred to herein as "domains")

within a protein is an element of overall structure that is self-stabilizing and often folds independently of the rest of the protein chain. Most domains can be classified into folds. Many domains are not unique to the proteins produced by one gene or one gene family but instead appear in a variety of proteins. Domains often are named and singled out because they play an important role in the biological function of the protein to which they belong; for example, the substrate binding domain of hemagglutinin can participate in binding to the substrate. Because domains can be self-stabilizing, domains can be swapped between one protein and another or associated to other carrier proteins by genetic engineering to make chimera proteins. A domain may be composed of none, one, or many structural motifs. In the cases of influenza hemagglutinin construct design, many structural motifs are preserved. The boundary selection is guided by known crystal structures. In the case of PR8, residues from E51 to K327 (SEQ ID NO: 1) fold into a compact structure that distinguishes from the rest of the HA molecules, thus is the region of focus to design HA constructs.

[0364] Design of the HA1-1, 1-2 and 1-3 globular head constructs for A/Viet Nam/1203/2004 (H5N1). The HA structure used as a reference in the design of the Viet Nam/1203/2004 (H5N1) globular head constructs is described in Stevens et al, 2006, *Science* 312:404-410 (MMDB number 38730; PDB accession number 2FK0). As described for the PR8 globular head constructs above, the published crystal structure of the mature A/Viet Nam/1203/2004 HA (SEQ ID NO: 2) was used in combination with the Molecular Modeling Database to determine the domain boundaries for three Viet Nam globular head constructs. The same structural criteria for preservation of secondary and tertiary structure for the structural domain were applied to the design of the Viet Nam constructs. While different hemagglutinin molecules can differ in the nature or number of residues that comprise the hemagglutinin molecule; the overall structure is remarkably similar. Thus, design of the domain boundaries for the Viet Nam globular head constructs required placement of the boundaries at structurally equivalent but numerically different positions within the HA molecule.

[0365] Design of the HA1-1, 1-2 and 1-3 globular head constructs for A/Indonesia/5/2005 (H5N1). The crystal structure for the Indonesia HA has not been solved. In general, when the crystal structure for a given HA molecule has not been solved, the available structure with the highest sequence identity can be used to guide the construct design. In the case of the design of the Indonesia globular head construct, the closest structure available is the A/Viet Nam/1203/2004 (Stevens et al. 2006. *Science* 312:404-410; MMDB number 38730; PDB Accession number 2FK0), which is of the same subtype as A/Indonesia/5/2005.

[0366] To design globular head constructs for this HA, the primary sequence of the Indonesia HA (SEQ ID NO: 3) was first aligned with A/Viet Nam/1203/2004 HA (H5VN; SEQ ID NO: 2) molecule (primary sequence identity: 96.13%). The primary sequence alignment for A/Viet Nam/1203/2004 HA (H5VN; SEQ ID NO: 2) and A/Indonesian/5/2005 (H5IN; SEQ ID NO: 3) was conducted using CLUSTALW and is shown below, where (*) asterisk=identity, (:) colon=conservative substitution; (.) period=weakly conservative substitution; and (space)=divergent substitution.

```

                    50
H5 IN  MEKIVLLLAIVSLVKSQICIGYHANNSTEQVDTIMEKNVTVTHAQDILE
H5 VN  MEKIVLLFAIVSLVKSQICIGYHANNSTEQVDTIMEKNVTVTHAQDILE
      *****;*****

                    100
H5 IN  KTHNGKLCDLGKPLILRDCSVAGWLLGNPMCDEFINVPESYIVEKAN
H5 VN  KKHNGKLCDLGKPLILRDCSVAGWLLGNPMCDEFINVPESYIVEKAN
      *.*****

                    150
H5 IN  PTNDLCYPGDFNDYEELKHLLSRINHFEKIQIIPKSSWSDHEASSGVSSA
H5 VN  PVNDLCYPGDFNDYEELKHLLSRINHFEKIQIIPKSSWSSHEASLGVSSA
      *.*****.*****.*****

                    200
H5 IN  CPYLGSPSFFRNVVWLIKKNSTYPTIKKSYNNTNQEDLLVLWGIHHPNDA
H5 VN  CPYQGKSSFFRNVVWLIKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPNDA
      *** *.*****;*****

                    250
H5 IN  AEQTRLYQNPTTYISIGTSTLNQRLVFKIATRISKVNGQSGRMEFFWTILK
H5 VN  AEQTKLYQNPTTYISVGTSTLNQRLVPRIATRISKVNGQSGRMEFFWTILK
      ****;*****;*****;*****

                    300
H5 IN  PNDAINFESNGNFIAPYAYKIVKKGDSAIMKSELEYGNCNTKCQTPMGA
H5 VN  PNDAINFESNGNFIAPYAYKIVKKGDSIMKSELEYGNCNTKCQTPMGA
      *****;*****

                    350
H5 IN  INSSMPFHNHPLTIGECPKYVKS NRLVLATGLRNSPQRESRRKKRGLFG
H5 VN  INSSMPFHNHPLTIGECPKYVKS NRLVLATGLRNSPQRESRRKKRGLFG
      *****

                    400
H5 IN  AIAGFIEGGWQGMVDGWYGYHHSNEQSGGYAADKESTQKAIDGVTNKVNS
H5 VN  AIAGFIEGGWQGMVDGWYGYHHSNEQSGGYAADKESTQKAIDGVTNKVNS
      *****

                    450
H5 IN  IIDKMTNQFEAVGREFNNLERRIENLNKKMEDGFLDVWTYNAELLVLMEN
H5 VN  IIDKMTNQFEAVGREFNNLERRIENLNKKMEDGFLDVWTYNAELLVLMEN
      *****

                    500
H5 IN  ERTLDFHDSNVKNLYDKVRLQLRDNKELGNGCFEFYHKCDNECMESIRN
H5 VN  ERTLDFHDSNVKNLYDKVRLQLRDNKELGNGCFEFYHKCDNECMESVRN
      *****;**

                    550
H5 IN  GTYNYPQYSEEARLKREEISGVKLESIGTYQILSIYSTVASSLALAIMMA
H5 VN  GTYDYPQYSEEARLKREEISGVKLESIGTYQILSIYSTVASSLALAIMVA
      ***;*****;*****;**

                    568
H5 IN  GLSLWMCNGLSLQCRICI (SEQ ID NO:3)
H5 VN  GLSLWMCNGLSLQCR--- (SEQ ID NO:2)
      *****

```

[0367] Design of the HA1-1, 1-2 and 1-3 globular head constructs for A/New Calcdonia/20/1999 (H1N1). The crystal structure for the New Calcdonia HA has not been solved. To design the globular head constructs for this HA, the primary sequence of the New Calcdonia HA (H1NC; SEQ ID NO: 4) was first aligned with two closely related HA molecules of the same subtype (primary sequence homology >85%) for which the structures have been resolved, specifically the H1N1 1918 virus (same as A/South Carolina/1/18)

(SEQ ID NO: 5) and the A/Puerto Rico/8/34 virus (H1PR8; SEQ ID NO: 1) (Gamblin, et al., *Science* 303, 1838-42 (2004). A/South Carolina/1/18: MMDB number 26943, PDB accession number: 1RUZ and PR8, MMDB number: 26941 and PDB accession number: 1RU7). The primary sequence alignment was conducted using CLUSTALW and is shown below, where (*) asterisk=identity, (:) colon=conservative substitution; (.) period=weakly conservative substitution; and (space)=divergent substitution.

```

H1NC      MKAKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLL
H1PR8      MKANLLVLLSALAAADADTICIGYHANNSTDTVDTVLEKTVTVTHSVNLL
          ***:****.::*: *****

H1NC      EDSHNGKLCLLKGIAPLQLGNCVAGWILGNPECELLISKESWSYIVETP
H1PR8      EDSHNGKLCRLKGIAPLQLGKCNLAGWLLGNPECDPLLPVRSWSYIVETP
          ***** :.::*:*****: :. .*****

H1NC      NPENGTCYPGYFADYEELREQLSSVSSFERFEIFPKESSWPNHNTVTGVSA
H1PR8      NSENGICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNHNTNGVTA
          *.*** ** * *****...*:

H1NC      SCSHNGKSSFYRNLLWLTGKNGLYPNLSKSYVNNKEKEVLVLWGVHHPPN
H1PR8      ACSHEGKSSFYRNLLWLTEKEGSYPKL2QSYVNKKGKEVLVLWGIHHPN
          :***:***** *: * :.::*: * *****:*****

H1NC      IGNQRALYHTENAYVSVVSSHYSRRFTPEIAKRPKVRDQEGRINYWTLL
H1PR8      SKEQQNIYQENAYVSVVTSNYNRRFTPEIAERPVRDQAGRMNYWTLL
          ::: :*.*****:*.*****:***** *:*****

H1NC      EPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNAPMDECDACKQTPQG
H1PR8      KPGDTIIFEANGNLIAPMYAFALSRGFGSGIITSNASHMHCNTKCQTPLG
          :***** *****..*::*:*****

H1NC      AINSSLFPQNVHPVTIGCEPKYVRSAKLRMVTGLRNIPSIQSRGLFGAIA
H1PR8      AINSSLFPYQNIHPVTIGCEPKYVRSAKLRMVTGLRNTPSIQSRGLFGAIA
          *****:*.***** *****

H1NC      GFIEGGWTGMVDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNSVIE
H1PR8      GFIEGGWTGMIDGWYGYHHQNEQGSYAADQKSTQNAINGITNKVNTVIE
          *****:*****:*****

H1NC      KMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENERT
H1PR8      KMNIQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTYNAELLVLENERT
          *** *****:*****

H1NC      LDFHDSNVKNLYEKVKSQLKNNAKEIGNC-----
H1PR8      LDFHDSNVKNLYEKVKSQLKNNAKEIGNCGFEFYHKCDNECMESVRNGTY
          *****

H1NC      -----
H1PR8      DYPKYSEESKLNREKVDGVKLESMGIYQILAIYSTVASSLVLLVSLGAIS

H1NC      ----- (SEQ ID NO: 4)
H1PR8      FWMCSNGSLQCRICI (SEQ ID NO: 1)

```

[0368] Design of the HA1-1, 1-2 and 1-3 globular head constructs for A/Wisconsin/67/2005 (H3N2). The crystal structure for the Wisconsin HA has not been solved. To design the globular head constructs for this HA, the primary sequence of the Wisconsin HA (H3W is; SEQ ID NO: 6) was first aligned with a closely related reference HA molecule of the same subtype (primary sequence identity: 81.16%) for which the structure had been resolved, specifically influenza A/X31 subtype H3N2 (H3X31; SEQ ID NO: 7) (PDB

accession number: 1VIU). The Wisconsin HA sequence was aligned with the X31 HA using CLUSTAL W and is shown below, where (*) asterisk=identity, (:) colon=conservative substitution; (.) period=weakly conservative substitution; and (space)=divergent substitution. Each amino acid in X31 was found to have a corresponding match in A/Wisconsin/67/2005 sequence. The domain boundaries the X31 structure were then used to identify the domain boundaries in the A/Wisconsin/67/2005.

```

                                     50
H3Wis  QKLPGNDNSTATLCLGHAVPNGTIVKTIITNDQIEVTNATELVQSSSTGG
H3X31  QDLPGNDNSTATLCLGNHAVPNGTLVKTITDDQIEVTNATELVQSSSTGK
      * .*****:*****:*****

                                     100
H3Wis  ICDSPHQILDGENCTLIDALLGDPQCDGFQNKKWLDFVERSKAYSNCYPY
H3X31  ICNNPHRILDGIDCTLIDALLGDPHCDVFNQNETWDLFVERSKAFSNCYPY
      * .*:*** :*****:* *:*****:*****

                                     150
H3Wis  DVDPYASLRSLVASSGTLEFNDESFNWTGVTQNGTSSACKRRSNNSFFSR
H3X31  DVDPYASLRSLVASSGTLEFITEGFTWTGVIQNGGSNACKPGPGSGFFSR
      ***** * .***** * .***** ...*****

                                     200
H3Wis  LNWLTHLKFYKYPALNVTMPNNEKFDKLYIWGVHHPGTDNDQIFLHAQASG
H3X31  LNWLTSGSTYPVLNVTMPNNDNFDKLYIWGIHHPSTNQEQTSLYVQASG
      *****: .*.*****:*****:***.*:***

                                     250
H3Wis  RITVSTKRSQQTIVIPNIGSRPRI RNIPSRISYWTIVKPGDILLINSTGN
H3X31  RVTVSTRRSQQTIIIPNIGSRPWVRLSSRISYWTIVKPGDVLVINSNGN
      *:*****:*****:*.*****:*****.*

                                     300
H3Wis  LIAPRGYFKIRSGKSSIMRSDAPIGKCNSECITPNGSIPNDKPFQNVNRI
H3X31  LIAPRGYFKMRTGKSSIMRSDAPIDTCISECITPNGSIPNDKPFQNVNKI
      *****:*.*****.* *****:*****.*

                                     329
H3Wis  TYGACPRYVKQNTLKLATGMRNVPEKQTR (SEQ ID NO: 6)
H3X31  TYGACPKYVKQNTLKLATGMRNVPEKQT- (SEQ ID NO: 7)
      *****

```

Results

[0369] Depiction of domain boundaries for PR8 HA constructs. The selected boundary domains are highlighted in the sequence below (SEQ ID NO: 1) as follows: HA1-1 boundaries are single-underlined (S53-R324 of SEQ ID NO: 1); HA1-2 boundaries are double-underlined (K62-S284 of SEQ ID NO: 1); HA1-3 boundaries are bold-underlined (N101-G276 of SEQ ID NO: 1). Detailed descriptions of each subunit design and boundary domains are given below.

```

                                     (SEQ ID NO:1)
MKANLLVLLSALAADADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLLEDSHNGKLCR 60

LKGIAPLQLQKGCNIAGWLLGNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEELRE 120

QLSSVSSFERFEIFPKESSWPNHNTNGVTAACSHGKSSFYRNLLWLTEKGSYPKLKNS 180

YVNKKGKEVLVLWGIHHPNSKEQQNLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQA 240

GRMNYWTLLKPGDTHIFEANGNLIAPMYAFALSRGFGSGIITSSNAMHECNTKCQTPLG 300

AINSSLPYQNIHPVTIGECPKYVRSAKLRMVTGLRNIPSIQSRGLFGAIAGFIEGGWTGM 360

IDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNTVIEKMNIQFTAVGKEFNKLEKRM 420

ENLNKKVDDGFLDIWYNAELLVLENERLDFHDSNVKNLYEKVKSQ LKNNAKEIGNGC 480

FEFYHKCDNECMESVRNGTYDYPKYSEESKLNREKVDGVKLESMGIYQILAIYSTVASSL 540

VLLVSLGAISFWMCSNGSLQCRICI 565

```

[0370] PR/8 HA1-1 construct (SEQ ID NO: 8). For this construct, four (4) conserved disulfide bonds were preserved by making the amino-terminal truncation before the Cysteine at position 59 of SEQ ID NO: 1 and the carboxyl-terminal truncation after the Cysteine at position 319 of SEQ ID NO: 1. The preserved disulfide bonds are listed as the following: C59-C291; C72-C84; C107-C152 and C295-C319 of SEQ ID NO: 1. Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the Serine at position 53 of SEQ ID NO: 1, which is immediately adjacent to the coil turn formed by residues 50-52 of SEQ ID NO: 1 (LED). The truncation completely eliminated the preceding β -strand formed by residues 44-49 of SEQ ID NO: 1 [THSVNL (SEQ ID NO: 214)]. The strand spanning residues 53-58 of SEQ ID NO: 1 [SHNGKL (SEQ ID NO: 215)] and the entire β -strand associated with residues 59-62 of SEQ ID NO: 1 [CRLK (SEQ ID NO: 216)] were preserved. Although in the structure residues 53-58 of SEQ ID NO: 1 [SHNGKL (SEQ ID NO: 215)] are defined as a random coil, these residues mimic a β -strand and thus serve to further stabilize the β -pleated sheets defined by residues 307-310 of SEQ ID NO: 1 [PYQN, SEQ ID NO: 217] and residues 320-323 of SEQ ID NO: 1 [PKYV, SEQ ID NO: 218]. The carboxyl-terminal truncation was made in the random coil sequence spanning residues 324-327 of SEQ ID NO: 1 [RSAK, SEQ ID NO: 219] at the Arginine at position 324 of SEQ ID NO: 1 such that the carboxyl-terminal tail which interacts with HA2 was completely eliminated.

[0371] PR/8 HA1-2 construct (SEQ ID NO: 9). For this construct two (2) conserved disulfide bonds (C72-C84 and C107-C152 of SEQ ID NO: 1) were preserved by making the amino-terminal truncation before the Cysteine at position 72 of SEQ ID NO: 1 and the carboxyl-terminal truncation after the Cysteine at position 152 of SEQ ID NO: 1. Using the crystal structure cited above and the MMD database to identify tertiary and secondary structure, the amino-terminal and the carboxyl-terminal truncations were placed in the loops [KGI, residues 62-64 of SEQ ID NO: 1 and NASMHE (SEQ ID NO: 220), residues 285-290, of SEQ ID NO: 1] that connect two distinct sets of β -pleated sheets. Truncation of the molecule within these loops preserves the surrounding secondary structure which in turn preserves the tertiary structure of the globular head.

[0372] The amino-terminal truncation was made at the Lysine at position 62 of SEQ ID NO: 1, such that the β -strand comprising APLQLG (SEQ ID NO: 221) at residues 65-70 of SEQ ID NO: 1 of the membrane-distal set of β -strands remained wholly intact, while the preceding β -strand [CRLK, SEQ ID NO: 216, residues 59-62 of SEQ ID NO: 1] that is more membrane-proximal was largely eliminated. The carboxyl-terminal truncation was made at

the Serine at position 284 of SEQ ID NO: 1 such that the β -strand [GIITS, SEQ ID NO: 222, residues 280-284 of SEQ ID NO: 1] of the distal set of β -strands remains wholly or largely intact and the subsequent β -strand [CNTK, SEQ ID NO: 223, residues 291-294 of SEQ ID NO: 1] of the proximal set of β -strands is wholly or largely eliminated. Preservation of the distal set of β -strands was the main determinant in selecting the domain boundaries for the HA1-2 construct. The distal β -pleated sheet, which is comprised of β -strands of APLQLG (SEQ ID NO: 221) [residues 65-70 of SEQ ID NO: 1], SYIVET (SEQ ID NO: 224) [residues 94-99 of SEQ ID NO: 1], and GIITS (SEQ ID NO: 222) [residues 280-284 of SEQ ID NO: 1], serves as the stabilizing secondary structural element that ensures a compact domain structure.

[0373] PR/8 HA1-3 construct (SEQ ID NO: 10). For this construct one (1) conserved disulfide bond (C107-C152 of SEQ ID NO: 1) was preserved by making the amino-terminal truncation before the Cysteine at position 107 of SEQ ID NO: 1 and the carboxyl terminal truncation after the Cysteine at position 152 of SEQ ID NO: 1. Using the crystal structure cited above and the MMDB database to identify tertiary and secondary structure the amino-terminal truncation was made at the Asparagine at position 101 of SEQ ID NO: 1 such that the preceding α -helix [NIAGWLLG, SEQ ID NO: 225, residues 73-80 of SEQ ID NO: 1] and β -strand [SYIVET, SEQ ID NO: 224, residues 94-99 of SEQ ID NO: 1] were completely eliminated while the subsequent β -strand [PGDFI, SEQ ID NO: 226, residues 109-113 of SEQ ID NO: 1] was preserved along with the strand NSENGICY, SEQ ID NO: 227 [residues 101-108 of SEQ ID NO: 1] that includes the conserved Cysteine at position 107 of SEQ ID NO: 1.

[0374] The carboxyl-terminal truncation was made near the end of the β -strand [AFALSRGF, SEQ ID NO: 228, residues 270-277 of SEQ ID NO: 1]. The last amino acid of this strand, position 277 of SEQ ID NO: 1, is a Phenylalanine, which was eliminated so as not to expose this hydrophobic residue. The carboxyl-terminal truncation was made at the Glycine at position 276 of SEQ ID NO: 1 which is in between two secondary structures. Truncation at this position keeps the preceding β -strand [AFALSRG, SEQ ID NO: 229, residues 270-276 of SEQ ID NO: 1] largely intact and completely eliminates the subsequent β -strand [GIITS, SEQ ID NO: 222, residues 280-284 of SEQ ID NO: 1].

[0375] Depiction of domain boundaries for VN04 HA construct. The selected boundary domains are highlighted in the sequence below (SEQ ID NO: 2) as follows: HA1-1 boundaries are single-underlined (E50-K323 of SEQ ID NO: 2); HA1-2 boundaries are double-underlined (G62-E284 of SEQ ID NO: 2); HA1-3 boundaries are bold-underlined (N103-G276 of SEQ ID NO: 2). Detailed descriptions of each subunit design and boundary domains are given below.

(SEQ ID NO:2)

MEKIVLLFAIVSLVKSQDQICIGYHANNSTEQVDTIMEKNVTVTHAQDILEKHHNGKLCDL 60

DGVKPLILRDCSVAGWLLGNPMCDFEINPEWSYIVEKANPVNDLCYPGDFNDYEELKHL 120

LSRINHFEKIQIIPKSSWSSEASLGVSACPYQKGSSFFRNVVWLIKKNSTYPTIKRSY 180

NNTNQEDLLVLWGIIHPNDAAEQTKLYQNPTTYISVGTSTLNQRLVPRIATRSKVNQSG 240

-continued

```

RMEFFWTILKPNDAINFESNGNFIAPYAYKIVKKGDSTIMKSELEYGNCNTKCQTPMGA 300
INSSMPFHNHPLTIGECPKYVKSRLVLATGLRNSPQRERRRKRGLFGAIAAGFIEGGW 360
QGMVDGWYGYHHSNEQGSGYAADKESTQKAIDGVTNKVNSIIDKMNTQFEAVGREFNNLE 420
RRIENLNKKMEDGFLDVWVTYNAELLVLMENERTLDFHDSNVKNLYDKVRLQLRDNAKELG 480
NGCFEFYHKCDNECMESVRNGTYDYPQYSEEARLKREEISGVYTESIGIYQILSIYSTVA 540
SSLALAIMVAGLSLWMCNSNGSLQCR 565

```

[0376] VN04 HA1-1 construct (SEQ ID NO: 11). For this construct four (4) conserved disulfide bonds were preserved by making the amino terminal truncation before the Cysteine at position 58 of SEQ ID NO: 2 and the carboxyl-terminal truncation after the Cysteine at position 318 of SEQ ID NO: 2. The preserved disulfide bonds are listed as the following: C58-C290; C71-C83; C106-C151 and C294-C318 of SEQ ID NO: 2. Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the Glutamic Acid found at position 50 of SEQ ID NO: 2 such that the preceding β -strand [THAQDI, SEQ ID NO: 230, residues 43-48 of SEQ ID NO: 2] is totally eliminated and the subsequent strand [EKKHNG, SEQ ID NO: 231, residues 50-55 of SEQ ID NO: 2] and the β -strand [KLCDDL, SEQ ID NO: 232] formed by residues 56-61 of SEQ ID NO: 2 were left wholly intact. Structurally the random peptide comprising residues 50-55 of SEQ ID NO: 2 [EKKHNG, SEQ ID NO: 231] mimics a β -strand that completes the membrane-distal β -pleated sheet to keep the domain structure intact. The carboxyl-terminal truncation was made in the loop region near residues 323-326 of SEQ ID NO: 2 [KSNR, SEQ ID NO: 233] that follow the Cysteine at position 318 of SEQ ID NO: 2. Truncation within this loop preserves secondary structure in that the preceding β -strand formed by residues 319-322 of SEQ ID NO: 2 [PKYV, SEQ ID NO: 234] remains wholly intact and the subsequent β -strand [LVLATG, SEQ ID NO: 235, residues 327-332 of SEQ ID NO: 2] is completely eliminated.

[0377] VN04 HA1-2 construct (SEQ ID NO: 12). For this construct two (2) conserved disulfide bonds (C71-C83 and C106-C151 of SEQ ID NO: 2) were preserved by making the amino-terminal truncation before the Cysteine at position 71 of SEQ ID NO: 2 and the carboxyl-terminal truncation after the Cysteine at position 151 of SEQ ID NO: 2. Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the glycine at position 62 of SEQ ID NO: 2 such that the subsequent β -strand comprising residues 64-68 of SEQ ID NO: 2 [KPLIL, SEQ ID NO: 236] remains wholly intact, while the preceding β -strand [KLCDDL, SEQ ID NO: 232, residues 56-61 of SEQ ID NO: 2] is completely eliminated.

[0378] To preserve secondary structure, the carboxyl-terminal truncation was made in the loop region at the Glutamic Acid at position 284 of SEQ ID NO: 2 such that the preceding β -strand [TIMKS, SEQ ID NO: 237, residues 279-283 of SEQ ID NO: 2] remained wholly or largely intact and the subsequent β -strand [YGNCN, SEQ ID NO: 238, residues 287-291 of SEQ ID NO: 2] was completely eliminated.

[0379] VN04 HA1-3 construct (SEQ ID NO: 13). For this construct one (1) conserved disulfide bond (C106-C151 of SEQ ID NO: 2) was preserved by making the amino-terminal truncation before the Cysteine at position 106 of SEQ ID NO: 2 and the carboxyl terminal truncation after the Cysteine at position 151 of SEQ ID NO: 2.

[0380] Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the Asparagine at position 103 of SEQ ID NO: 2 such that the preceding β -strand [SYIVEK, SEQ ID NO: 239, residues 93-98 of SEQ ID NO: 2] was completely eliminated while the subsequent β -strand sequence [PGDFN, SEQ ID NO: 240, residues 108-112 of SEQ ID NO: 2] remained wholly or largely intact. To preserve secondary structure, the carboxyl-terminal truncation was made in the random coil region at the Glycine at position 276 of SEQ ID NO: 2 such that the preceding β -strand formed by residues 102-105 of SEQ ID NO: 2 [KGDS, SEQ ID NO: 241] remains wholly or partially intact, the preceding β -strand [EYAYKIVK, SEQ ID NO: 242, residues 267-274 of SEQ ID NO: 2] is wholly preserved and the subsequent β -strand [TIMKS, SEQ ID NO: 237, residues 279-283 of SEQ ID NO: 2] is completely eliminated.

[0381] Depiction of domain boundaries for IND05 HA constructs. The A/Indonesia/5/2005 construct description is based on sequence alignment in reference to A/Viet Nam/1203/2004 structure. The selected boundary domains are highlighted in the sequence below (SEQ ID NO: 3) as follows: HA1-1 boundaries are single-underlined (E50-K323, of SEQ ID NO: 3); HA1-2 boundaries are double-underlined (G62-E284 of SEQ ID NO: 3); HA1-3 boundaries are bold-underlined (N103-G276 of SEQ ID NO: 3). Detailed descriptions of each subunit design and boundary domains are given below.

```

                                (SEQ ID NO:3)
MEKIVLLLLAIIVSLVKSDQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILEKTHNGKLCDL 60
DGVKPLILRDCSVAGWLLGNPMCDEFINVPESYIVEKANPTNDLCYPGSFNDYEELKHL 120
LSRINHFEKIQIIPKSSWSDEASSGVSSACPYLGSPSFFRNVLIKKNSTYPTIKKSY 180
NNTNQEDLLVLWGIHHPNDAAEQTRLYQNPTTYISIGTSTLNQRLVPKIATRSKVNQSG 240
RMEFFWTILKPNDAINFESNGNFIAPYAYKIVKKGDSAIMKSELEYGNCNTKCTPMGA 300
INSSMPFHNHPLTIGECPKYVKNRNLVLATGLRNSPQRESRRKRGLFGAAGFIEGGW 360
QGMVDGWYGYHHSNEQSGGYAADKESTQKAIDGVTNKVNSIIDKMNTQFEAVGREFNLE 420
RRIENLNKKMEDGFLDVWTYNAELLVLMENERTLDFHDSNVKNLYDKVRLQLRDNAKELG 480
NGCFEFYHKCDNECMESIRNGTYNYPQYSEEARLKREEISGVKLESIGTYQLSIYSTVA 540
SSLALAIMMAGLSLWMCNSNGSLQCRICI 560

```

[0382] IND05

HA1-1 construct (SEQ ID NO: 14). For this construct four (4) conserved disulfide bonds were preserved by making the amino-terminal truncation before the Cysteine at position 58 of SEQ ID NO: 3 and the carboxyl-terminal truncation after the Cysteine at position 318 of SEQ ID NO: 3. The preserved disulfide bonds are listed as the following: C58-C291; C71-C83; C106-C151 and C294-C318 of SEQ ID NO: 3. Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the Glutamic Acid found at position 50 of SEQ ID NO: 3 such that the preceding β -strand [THAQDI, SEQ ID NO: 243, residues 43-48 of SEQ ID NO: 3] was totally eliminated and the subsequent strand [EKTHNG, SEQ ID NO: 244, residues 50-55 of SEQ ID NO: 3] and β -strand [KLCDLD, SEQ ID NO: 245] formed by residues 56-61 of SEQ ID NO: 3 was left wholly intact. Structurally the random peptide comprising residues 50-55 of SEQ ID NO: 3 [EKTHNG, SEQ ID NO: 244] mimics a β -strand that completes the membrane-distal β -pleated sheet to keep the domain structure intact. The carboxyl-terminal truncation was made in the loop region near residues 323-326 of SEQ ID NO: 3 [KSNR, SEQ ID NO: 246] that follow the Cysteine at position 318 of SEQ ID NO: 3. Truncation within this loop preserves secondary structure in that the preceding β -strand formed by residues 319-322 of SEQ ID NO: 3 [PKYV, SEQ ID NO: 247] remains wholly intact and the subsequent β -strand [LVLATG, SEQ ID NO: 248, residues 327-332 of SEQ ID NO: 3] is completely eliminated.

[0383] IND05

HA1-2 construct (SEQ ID NO: 15). For this construct two (2) conserved disulfide bonds (C71-C83 and C106-C151 of SEQ ID NO: 3) were preserved by making the amino-terminal truncation before the Cysteine at position 71 of SEQ ID NO: 3 and the carboxyl-terminal truncation after the Cysteine at position 151 of SEQ ID NO: 3. Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the Glycine at position 62 of SEQ ID NO: 3 such that the subsequent α -strand comprising residues 64-68 of SEQ ID NO: 3 [KPLIL, SEQ ID NO: 249] remained wholly intact, while the preceding β -strand [KLCDLD, SEQ ID NO: 245, resi-

dues 56-61 of SEQ ID NO: 3] was completely eliminated. To preserve secondary structure, the carboxyl terminal truncation was made in the loop region at the Glutamic Acid at position 284 of SEQ ID NO: 3 in the sequence [ELE, residues 284-286 of SEQ ID NO: 3] such that the preceding β -strand [AIMKS, SEQ ID NO: 250, residues 279-283 of SEQ ID NO: 3] remained wholly or largely intact and the subsequent α -strand [YGNCN, SEQ ID NO: 251, residues 287-291 of SEQ ID NO: 3] was completely eliminated.

[0384] IND05

HA1-3 construct (SEQ ID NO: 16). For this construct one (1) conserved disulfide bond (C106-C151 of SEQ ID NO: 3) was preserved by making the amino-terminal truncation before the Cysteine at position 106 of SEQ ID NO: 3 and the carboxyl-terminal truncation after the Cysteine at position 151 of SEQ ID NO: 3. Using the crystal structure cited above and the Molecular Modeling Database (MMDB) to identify tertiary and secondary structure, the amino terminal truncation was placed at the asparagine at position 103 of SEQ ID NO: 3 such that the preceding α -strand [SYIVEK, SEQ ID NO: 252, residues 93-98 of SEQ ID NO: 3] was completely eliminated while the β -strand sequence [PGSFN, SEQ ID NO: 253, residues 108-112 of SEQ ID NO: 3] remained wholly or largely intact. To preserve secondary structure, the carboxyl-terminal truncation was made in the random coil region at the Glycine found at position 276 of SEQ ID NO: 3 such that the preceding β -strand [EYAYKIVK, SEQ ID NO: 254, residues 267-274 of SEQ ID NO: 3] was wholly preserved and the subsequent β -strand [AIMKS, SEQ ID NO: 250, residues 279-283 of SEQ ID NO: 3] was completely eliminated.

[0385] Depiction

of domain boundaries for New Calcdonia HA constructs. A/New Calcdonia/20/199 construct description is based on sequence alignment in reference of A/Puerto Rico/8/34 structure. The selected boundary domains are highlighted in the sequence below (SEQ ID NO: 4) as follows: HA1-1 boundaries are single-underlined (S53-R324 of SEQ ID NO: 4); HA1-2 boundaries are double-underlined (K62-S284 of SEQ ID NO: 4); HA1-3 boundaries are bold-underlined (N101-G276 of SEQ ID NO: 4). Detailed descriptions of each subunit design and boundary domains are given below.

(SEQ ID NO:4)
 MKAKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTTHSVNLLEDSHNGKLCL 60
 LKGIAPLQLGNCVAGWILGNPECELLISKESWSYIVETPNPENGTCYPGYFADYEELRE 120
 QLSSVSSFERFEIFPKESSWPNHTVTGVSASC SHNGKSSFYRNLLWLTGKNGLYPNLSKS 180
 YVNNKEKEVLVLGWVHPNIGNQRALYHTENAYVSVSSHYSRRFTPEIAKRPKVRDQE 240
 GRINYYWTLLLEPGDTIIFEANGNLIAPWYAFALSSRGFGSGIITSNAPMDECDACQTPQG 300
 AINSSLFPQNVHPVTIGECPKYVRSAKLRMVTGLRNIPSIQSRGLFGAIAGFIEGGWTGM 360
 VDGWGYHHQNEQSGYAADQKSTQNAINGITNKVNSVIEKMNTQFTAVGKEFNTERRN 420
 ENLNKKVDDGFLDIWTYNAELLVLLLENERTLDFHDSNVKNLYEKVKSQKNNAKEIGNGC 480
 FEFYHKCNNECMESVKNGTYDPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSL 540
 VLLVSLGAISFWMCSSNGSLQCRICI 565

[0386] New Caledonia HA1-1 construct design (SEQ ID NO: 17). For this construct, four (4) conserved disulfide bonds were preserved by making the amino terminal truncation before the Cysteine at position 59 of SEQ ID NO: 4 and the carboxyl-terminal truncation after the Cysteine at position 319 of SEQ ID NO: 4. The preserved disulfide bonds are listed as the following: C59-C291; C72-C84; C107-C152 and C295-C319 of SEQ ID NO: 4. Using the reference crystal structures and the MMDB to check secondary structures the amino terminal truncation was made at the Serine at position 53 of SEQ ID NO: 4 such that the preceding β -strand [THSVNLLED, SEQ ID NO: 255, residues 44-52 of SEQ ID NO: 4] was totally eliminated and the subsequent β -strands SHNGKL, SEQ ID NO: 256 [residues 53-58 of SEQ ID NO: 4] and APLQLG, SEQ ID NO: 257 [residues 65-70, of SEQ ID NO: 4] were left wholly intact. The amino-terminal truncation in the random coil sequence [SHNGKL, SEQ ID NO: 256, residues 53-58 of SEQ ID NO: 4] left the secondary structure largely intact because it mimics a β -strand that leaves the β -pleated sheet intact. The carboxyl-terminal truncation was made at the Arginine at position 324 of SEQ ID NO: 4 in the random coil sequence [RSAK, SEQ ID NO: 258, residues 324-327 of SEQ ID NO: 4] following the Cysteine at position 319 of SEQ ID NO: 4. The truncation in this random coil preserves secondary structure by leaving the preceding β -strand [PKYV, SEQ ID NO: 259, residues 320-323 of SEQ ID NO: 4] wholly intact while the subsequent β -strand [LRMVTG, SEQ ID NO: 260, residues 328-333 of SEQ ID NO: 4] is completely eliminated.

[0387] New Caledonia HA1-2 construct design (SEQ ID NO: 18). For this construct two (2) conserved disulfide bonds (C72-C84 and C107-C152 of SEQ ID NO: 4) were preserved by making the amino-terminal truncation before the Cysteine at position 72 of SEQ ID NO: 4 and the carboxyl-terminal truncation after the Cysteine at position 152 of SEQ ID NO: 4. Using the reference crystal structures and the MMDB to check secondary structures the amino-terminal truncation was made at the Lysine at position 62 of SEQ ID NO: 4 such that the random coil sequence comprising [SHNGKLCLLKGI, SEQ ID NO: 261, residues 53-64 of SEQ ID NO: 4] was partially or largely removed, the preceding β -strand [THSVNLLED, SEQ ID NO: 255, residues 44-52 of SEQ ID NO: 4] was completely eliminated

and the subsequent β -strand [APLQLG, SEQ ID NO: 257, residues 65-70 of SEQ ID NO: 4] remained wholly intact. The carboxyl-terminal truncation was made at the Serine at position 284 of SEQ ID NO: 4 such that the preceding β -strand [SGIITS, SEQ ID NO: 262, residues 279-284 of SEQ ID NO: 4] remained wholly intact and the subsequent random coil [NAPMDECDCA, SEQ ID NO: 263, residues 285-293 of SEQ ID NO: 4] was wholly or largely eliminated.

[0388] New Caledonia HA1-3 construct design (SEQ ID NO: 19). For this construct one (1) conserved disulfide bond (C107-C152 of SEQ ID NO: 4) was preserved by making the amino-terminal truncation before the Cysteine at position 107 of SEQ ID NO: 4 and the carboxyl terminal truncation after C152 of SEQ ID NO: 4. Using the reference crystal structures and the MMDB to check secondary structures the amino-terminal truncation was made at the Asparagine at position 101 of SEQ ID NO: 4 such that the receding β -strand [SYIVET, SEQ ID NO: 264, residues 94-99 of SEQ ID NO: 4] was completely eliminated and the subsequent β -strand [PGYFA, SEQ ID NO: 265, residues 109-113 of SEQ ID NO: 4] remained wholly intact. To preserve the secondary structure the carboxyl-terminal truncation was made at the Glycine at position 276 of SEQ ID NO: 4 such that the β -strand sequence [YAFALSRGF, SEQ ID NO: 266, residues 269-277 of SEQ ID NO: 4] remained largely intact, while the subsequent β -strand sequence [SGIITS, SEQ ID NO: 262, residues 279-284 of SEQ ID NO: 4] was eliminated.

[0389] Depiction of domain boundaries for A/Wisconsin/67/2005 HA constructs. A/Wisconsin/67/2005 construct description is based on sequence alignment in reference of A/X31 structure. The selected boundary domains are highlighted in the sequence below (SEQ ID NO: 6) as follows: HA1-1 boundaries are single-underlined (Q44-K310 of SEQ ID NO: 6); HA1-2 boundaries are double-underlined (S54-D271 of SEQ ID NO: 6); HA1-3 boundaries are bold-underlined (S95-G263 of SEQ ID NO: 6); Detailed descriptions of each subunit design and boundary domains are given below.

(SEQ ID NO:6)

QKLP GNDNSTATLCLGHVAVNGTIVKTIITNDQIEVTNATELVOSSTGGICDSPHOILD 60

GENCTLIDALLGDPQCDGFQNKWDLFVERSKAYSNCYPYDVPDYASLRSLVASSGTLEF 120

NDESFNWTGVTQNGTSSACKRRSNNSSFRLNWLTHLKFYKYPALNVTMPNNEKFDKLYIW 180

GVHHPGTDNDQIFLHAQASGRITVSTKRSQQTVIPNIGSRPRIRNIPSRISIIYWTIVKPG 240

DILLINSTGNLIAPRGYFKIRSGKSSIMRSDAPIGKCNSECITPNSIPNDKPFQNVNRI 300

TYGACPRYVKQNTLKLATGMNRNVEKQTR 329

[0390] A/Wisconsin/67/2005 HA1-1 construct design (SEQ ID NO: 20). For this construct, four (4) conserved disulfide bonds were preserved by making the amino terminal truncation before the Cysteine at position 52 of SEQ ID NO: 6 and the carboxyl-terminal truncation after the Cysteine at position 305 of SEQ ID NO: 6. The preserved disulfide bonds are listed as the following: C52-C277; C64-C76; C97-C139 and C281-C305 of SEQ ID NO: 6. Using the reference crystal structures and the MMDB to check secondary structures the amino terminal truncation was made at the Glutamine at position 44 of SEQ ID NO: 6, right after the tight turn formed by residues 42 and 43 of SEQ ID NO: 6 [LV], such that the preceding β -strand [TNATE, SEQ ID NO: 267, residues 37-41 of SEQ ID NO: 6] was totally eliminated while the entire short β -strand formed by residues QSS [44-46 of SEQ ID NO: 6] was preserved. The carboxyl-terminal truncation was made in the random coil sequence [PRYVK, SEQ ID NO: 268, residues 306-310 of SEQ ID NO: 6] following the Cysteine at position 305 of SEQ ID NO: 6 such that the carboxyl-terminal tail which interacts with HA2 was completely eliminated. Although the carboxyl-terminal sequence [PRYVK, SEQ ID NO: 268, residues 306-310 of SEQ ID NO: 6] is defined as a random coil, structurally it serves as an additional β -strand that further stabilizes the β -pleated sheets defined by residues 44-46 of SEQ ID NO: 6 [QSS] and residues 293-297 of SEQ ID NO: 6 [PFQNV, SEQ ID NO: 269]. This set of β -pleated encompasses the amino-terminus and carboxy-terminus within one stable secondary structure element to ensure a compact domain structure.

[0391] A/Wisconsin/67/2005 HA1-2 construct design (SEQ ID NO: 21). For this construct two (2) conserved disulfide bonds (C64-C76 and C97-C139 of SEQ ID NO: 6) were preserved by making the amino-terminal truncation before the Cysteine at position 64 of SEQ ID NO: 6 and the carboxyl-terminal truncation after the Cysteine at position 139 of SEQ ID NO: 6. Using the reference crystal structures and the MMDB to check secondary structures the amino-terminal truncation was made in the loops that connect two distinct sets of β -pleated sheets. The amino-terminal truncation was made at the Serine at position 54 of SEQ ID NO: 6, such that the β -strand comprising residues 57-62 of SEQ ID NO: 6 [QILDGE, SEQ ID NO: 270] of the membrane-distal set of β -strands remained wholly intact, while the membrane-proximal β -strand [GGICD, SEQ ID NO: 271, residues 49-53 of SEQ ID NO: 6] was completely eliminated. The carboxyl-terminal truncation was made at the Aspartic Acid at position 271 of SEQ ID NO: 6. Truncation at this position preserved secondary structure such that the membrane-distal β -strand [SSIMRS, SEQ ID NO: 272, residues 265-270 of SEQ ID NO: 6] remained intact while

the membrane-proximal β -strand [APIGK, SEQ ID NO: 273, residues 272-276 of SEQ ID NO: 6] was eliminated. The membrane-distal β -pleated sheets comprising β -strands QILDGE (SEQ ID NO: 270), [residues 57-62 of SEQ ID NO: 6] LFVER (SEQ ID NO: 274) [residues 86-90 of SEQ ID NO: 6], and SSIMRS, (SEQ ID NO: 272), [residues 265-270 of SEQ ID NO: 6] serve as the stabilizing secondary structure element that ensures a compact domain structure.

[0392] A/Wisconsin/67/2005 HA1-3 construct design (SEQ ID NO: 22). For this construct one (1) conserved disulfide bond (C97-C139 of SEQ ID NO: 6) was preserved by making the amino-terminal truncation before the Cysteine at position 97 of SEQ ID NO: 6 and the carboxy-terminal truncation after the Cysteine at position 139 of SEQ ID NO: 6. Using the reference crystal structure and the MMDB to check secondary structures the amino-terminal truncation was made at the Serine at position 95 of SEQ ID NO: 6 such that the preceding α -helix [TLIDALLG, SEQ ID NO: 275, residues 65-72 of SEQ ID NO: 6] was completely eliminated while the Serine at position 95 of SEQ ID NO: 6 and Asparagine at position 96 of SEQ ID NO: 6 were preserved along with the conserved Cysteine at position 97 of SEQ ID NO: 6. The carboxyl-terminal truncation was made in the random coil sequence at about the Glycine at position 263 of SEQ ID NO: 6 such that the random coil sequence containing residues 261-263 of SEQ ID NO: 6 [RSG] was kept wholly or partially intact, while the preceding β -strand [RGYFKI, SEQ ID NO: 276, residues 255-260 of SEQ ID NO: 6] was wholly preserved and the subsequent β -strand [SSIMRS, SEQ ID NO: 272, residues 265-270 of SEQ ID NO: 6] was completely eliminated.

Discussion

[0393] Parts of the globular head domain of the influenza virus hemagglutinin (HA) protein may generate neutralizing antibodies (Brand and Skehel; 1972; Eckert, 1973; Jackson et al., 1979; Russ et al., 1981) and the discrete, globular structure of this domain was thought to constitute a protein fragment which could be expressed in bacteria and refolded, or expressed and secreted from a eukaryotic host, thereby avoiding the expression and purification problems which have been encountered producing the full-length HA protein.

[0394] The largest HA construct (also referred to herein as "HA fragment") fused to STF2, HA1-1 (SEQ ID NO: 8, 11, 14, 17, 20), encompasses most of or likely the entire globular region of the naturally occurring (wild type) HA1 head domain. A naturally occurring fragment of HA1 very similar to the HA1-1 construct is released from the virus on

limited proteolysis (Bizeband et al., 1995). Thus, structural analysis and experimental data suggest that the HA1-1 construct should fold efficiently, whether produced in bacteria or eukaryotes, and maintain a stable native-like structure. Similarly, it is expected that the entire HA0s, the external portion of HA molecule outside the viral membrane (SEQ ID NO: 23) consisting of HA1 and HA2 but excluding the signal peptide (17 aa), internal peptide within the viral membrane (12 aa) and transmembrane domain (24 aa) should attain its native conformation when expressed in a eukaryotic host or can be refolded into a native state in vitro when expressed in bacteria. The two smaller HA fragments (HA1-2 (SEQ ID NO: 9, 12, 15, 18, 21) and HA1-3 (SEQ ID NO: 10, 13, 16, 19, 22)) truncate the region of the head domain distal to the receptor binding site in a manner which may either destabilize the resulting protein fragment (HA1-3) or expose the side chains of a few large hydrophobic residues (HA1-2). For this reason, experimental mutations were introduced into the HA1-2 and HA1-3 constructs, denoted as HA1-2mut (SEQ ID NO: 24, 26, 28, 30, 32) and HA1-3mut (SEQ ID NO: 25, 27, 29, 31, 33). The substitutions made in the HA1-3mut (SEQ ID NO: 25, 27, 29, 31, 33) construct introduce oppositely charged residues on the opposite strands which are left exposed by the HA1-3 truncations. "HA mut," as used herein, means an amino acid in the native or naturally occurring HA has been substituted with an amino acid that does not occur in the native or naturally occurring HA. Substituted residues may form a salt bridge and stabilize a structure which may otherwise fold poorly, or not at all. For example, in PR8, G105 was replaced by glutamate and Y115 was substituted with lysine (SEQ ID NO: 25). Negatively charged glutamate interacted with positively charged lysine residue so that N-terminal peptide 1-8 of SEQ ID NO: 25 [NSENEICY, SEQ ID NO: 277] was stabilized by charge-charge interaction with Lysine 15 located at the N-terminus of the short helix 15-22 of SEQ ID NO: 25 [KEELREQL, SEQ ID NO: 278] in the center of the molecule. For the HA1-2 construct, the truncations can lead to the exposure of hydrophobic side chains which would not be expected to impair expression of the protein. However, the exposed residues may lead to aggregation of expressed protein or instability of expressed protein. To avoid aggregation and enhance the stability of the expressed molecule the few large hydrophobic residues that become exposed were substituted with more neutral amino acids to generate HA1-2mut (SEQ ID NO: 24, 26, 28, 30, 32). These substitutions may result in a protein which is more amenable to a robust manufacturing process and/or long term stability.

Example 2

Design of Portions of a Naturally Occurring Hemagglutinin of Influenza B

Materials and Methods

[0395] Design of the HA1-1, 1-2 and 1-3 globular head constructs for B/Lee/40. Structural considerations in the design of an influenza B HA vaccine are similar to those for influenza A, i.e., the domain boundary of the globular head of HA must be identified so that the flagellin-HA fusion protein can fold correctly or be refolded correctly to expose appropriate antigenic epitopes. Unlike influenza A, well-defined X-ray crystallographic structures are not available for influenza B HA, thus it is more difficult to unambigu-

ously define the domain boundary of globular head. Therefore the influenza B HA model must be predicted based on bioinformatic and structural models (Tung et al. 2004. *J. Gen. Virol.* 85, 3249-3259). These investigators used a "knowledge-based" approach which depends on a high degree of sequence homology between the known structure from the protein data bank and the target unknown structure. In general, this approach benefits most from at least 35% sequence identity between the known and target proteins.

[0396] In the case of influenza B, the closest models come from A/Swine/Hong Kong/9/98 (SEQ ID NO: 34) (24% identity, PDB accession code 1JSD) and A/Aichi/2/68 (SEQ ID NO: 35) (21% identity, PDB accession code 1HGF). Although the sequence identities between the target model B/Lee/40 HA (SEQ ID NO: 36) and known template models are substantially below the desired minimum of 35%, the close similarity of the functions and tertiary folds of influenza A HA proteins in spite of their sequence divergence (H1, H3, H5 and H9 share only 18% sequence identity) suggest a possibly successful prediction of influenza B HA structure using the influenza A HA model. Moreover, influenza C HEF (Hemagglutinin-Esterase-Fusion) protein folds similarly to influenza A HA structure despite even lower sequence identity than any of the A/B comparison.

[0397] Since the crystal structure of influenza C HEF is known (PDB accession number 1FLC, MMDB accession number 12663), Tung et al included the knowledge of the structure similarity between C HEF and the known influenza A HA proteins to predict influenza B HA structure. Tung et al first aligned the HEF sequence from C/Johannesburg/1/66 (SEQ ID NO: 37) with one sequence from each of the 15 HA subtypes of influenza A virus (<http://flu.lanl.gov>) using CLUSTALW (Thompson et al. 1994. *Nucleic Acids Res* 22, 4673-4680) and compiled a profile based on structure-informed alignment.

[0398] The conserved secondary structural features were captured and assigned, as well as the variations among the types and subtypes by year and host species. They then further aligned the augmented A/C profile to an alignment of influenza B HA sequences, including B/Lee/40. The homology modeling techniques of Tung (1999) was used to construct B/Lee/40 model.

[0399] Briefly, they first matched the main-chain structures of the target to those of the template in the aligned regions. Insertions in the target relative to the template were treated as loops with known end-structure. Stretching the predicted structure accommodated insertions in the template relative to the target. The main chain structures of the loops were modeled by using an efficient Monte Carlo loop-sampling method (Tung, 1997; Ryu et al., 1998). Once the main-chain structure was modeled, the side-chain atoms were attached. As the head of the HA molecule is compact, limited space is available to place the side-chain atoms. Hence, in the analysis, side-chain torsional angles were initialized to equal or be close to those in the template structure. This consideration is particularly useful in avoiding clashes between side chains in the modeled structure. Finally, the all-atom models were subjected to a short run of energy minimization (1000 cycles) by using AMBER (Weiner et al., 1986) to relieve unfavourable steric interactions and to optimize the stereochemistry. The quality of the target model was checked by PROCHECK (Wilson et al.

1998. *J. Mol Biol.* 276, 417-436) and the functionality was checked by substrate docking to test whether the model substrate binding site can accommodate the natural receptor analogue sialyllactose as we have seen in the crystal structure of the HA of A/Aichi/2/68 (Weis et al. 1988. *Nature* 333, 426-431). The simulated B/Lee/40 model appears reasonable in a Ramachandran Plot (Wilson et al. 1998. *J. Mol Biol.* 276, 417-436) and shows no steric crash when docking the sialyl-2-3-lactose molecule into the substrate binding site of the B/Lee/40 model, indicating a correct stereochemistry. Therefore the B/Lee/40 model was used to guide the design of influenza B HA subunit vaccines.

Results and Discussion

[0400] Depiction of Domain Boundaries for B/Lee/40 HA constructs. The selected boundary domains are highlighted in the sequence below as follows: HA1-1 boundaries are single-underlined (T48-K340 of SEQ ID: 36), and HA1-2 boundaries are double-underlined (K60-G299 of SEQ ID: 36). Detailed descriptions of each subunit design and boundary domains are given below.

(SEQ ID NO:36)

```

MKAIIVLLMVVTSNADRICTGITSSNSPHVVKATQGEVNVITGVIPLTTTPTKSHFANLK 60
GTOTRGKLCPCNCFNCTDLVALGRPKCMGNTPSAKVSILHEVKPATSGCFPIIMHRTKIR 120
QLPNLLRGYENIRLSTNSVINTEAPGGPYKVGTSKSPNVANGNGFFNTMAWVPIKDN 180
KTAIPVPTVEVPYICSEGEDQITVWGFHSDDKTQMERLYGDSNPQKFTSSANGVTTHYVS 240
QIGGFNPQTEDEGLKQSGRIVVDYVMVQKPGKTGTIVYQRGILLPQKVWCASGRSKVIKGS 300
LPLIGEADCLHEKYGLNKSPPYTTGEHAKAIGNCPIWVKTPCLKANGTKYRPPAKLLKE 360
RGFFGAIAAGFLEGGWEGMIAGWHGYTSHGAHGVAADLKSTQEAINKITKNLNYLSESL 420
VKNLQRLSGAMNELHDEILELDEKVDLDRADTISSQIELAVLLSNEGIINSEDEHLLALE 480
RKLKMLGPSAVEIGNGCFETKHKCNQTCLDRIAAGTFNAGDFSLPTFDSLNTAASLND 540
DGLDNHTILLYSTAASSLAVTLMIAIFIVYMSRDNVSCSICL 584

```

[0401] The coordinates of the simulated B/Lee/40 model are available in Protein Data Bank with the accession code 1TX1. The pdb file was converted to the MMDB (Molecular Modeling Data Base) format by using VAST (Vector Alignment Search Tool) search (<http://www.ncbi.nlm.nih.gov/Structure/VAST/vastsearch.html>). The model of 1TX1 structure then was viewed by Cn3D and the structure was saved in MMDB format. Based on the same principal used to design influenza A HA1-1 and HA1-2 constructs, the domain boundaries of B/Lee/40 HA-1 and HA1-2 were pinpointed by examining the model structure. B/Lee/40 HA1-1 (SEQ ID NO: 38) includes the epitope-concentrated globular top, the membrane distal β -sheet and an additional β -sandwich located underneath the membrane distal β -sheet. The amino terminus of HA1-1 starts from residue 48 and ends at residue 340 (TTPTK (SEQ ID NO: 279) CPIWVK (SEQ ID NO: 280)) of SEQ ID NO: 36). The B/Lee/40 HA1-2 (SEQ ID NO: 39) was designed by removing the β -sandwich from HA1-1. HA1-2 starts from residue 60 and ends at residue 299 (KGTQTR (SEQ ID NO: 281) SKVIKG (SEQ ID NO: 282)) of SEQ ID NO: 36. In order to confirm the boundary selection, other independent methods were also employed. The first method used

was the primary sequence alignment. The sequence of A/Aichi/2/68 (SEQ ID NO: 35) was aligned with B/Lee/40 (SEQ ID NO: 36) sequence. The boundaries of A/Aichi/2/68 HA1-1 (SEQ ID NO: 40) (residues 60-326 QSSSTG (SEQ ID NO: 283) CPKYVK (SEQ ID NO: 284)) and HA1-2 (SEQ ID NO: 41) (residues 72-287; HRILDG (SEQ ID NO: 285) SIMRSD (SEQ ID NO: 286)) were aligned closely to those of B/Lee/40, supporting the boundary selections using simulated model. One residue adjustment was made to avoid exposing hydrophobic residues and either end of the construct. Thus, the 3-dimensional structure prediction matches very well with primary sequence alignment.

[0402] The second method used was the secondary structure prediction. The domain boundary is usually located in the loop or turn region without invading much of the secondary structure elements such as the α -helix and the β -sheet, especially the center of the α -helix bundle or the β -sheet. This criterion was used to double check if the boundary selection made from simulated 3-D structure is in agreement with independent secondary structure prediction.

The program PHD (<http://ca.expasy.org/tools>→Proteomics and sequence analysis tools→Secondary and tertiary structure tools→PredictProtein) was used to perform the secondary structure prediction. Other than two of the HA1-2 carboxy-terminal residues overlapping with a short α -helix (5 amino acids), all other boundary residues fall in the loop regions, indicating a reasonable boundary selection. PHD results is listed as the following, where AA is the amino acid sequence; OBS_sec is the observed secondary structure: H=helix, E=extended sheet, blank=other (loop); PROF_sec: PROF predicted secondary structure: H=helix, E=extended (sheet), blank=other (loop), PROF=PROF: Profile network prediction HeiDelberg; Rel_sec: reliability index for PROF-sec prediction (0=low to 9=high)

[0403] For the brief presentation strong predictions marked by '*'; SUB_sec: subset of the PROFsec prediction, for all residues with an expected average accuracy >82% NOTE: for this subset the following symbols are used: L: is loop (for which above '' is used) .: means that no prediction is made for this residue, as the reliability is: Rel<5;

O_3_acc: observed relative solvent accessibility (acc) in 3 states: b=0-9%, i=9-36%, e=36-100%; P_3_acc: PROF predicted relative solvent accessibility (acc) in 3 states: b=0-9%, i=9-36%, e=36-100%; Rel_acc: reliability index for PROFacc prediction (0=low to 9=high).

[0404] For the brief presentation strong predictions marked by '*'; SUB_acc: subset of the PROFacc prediction,

for all residues with an expected average correlation >0.69 (tables in header).

[0405] NOTE: for this subset the following symbols are used:

[0406] I: is intermediate (for which above " is used)

[0407] .: means that no prediction is made for this residue, as the reliability is: Rel<4.

```

.....1.....2.....3.....4.....5
AA      MKAIIVLLMVVTSNADRICTGITSSNSPHVVKATATQGEVNVTVGVIPLTTT
OBS_sec
PROF_sec EEEEEEEEEEE EEEEEEEE EEEEE EEEEEEEEE
Rel_sec  90456542111257762677454037753444441276147765653203
SUB_sec  L..EEE.....LLLL.EEE.E...LLL.....LL.EEEEEEE...

O_3_acc  bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc  ee bbbbbbbbeeb bbbbbbb eeee b bb eebeeb b bbe
Rel_acc  3316999752111103277892120122233322135234021321012
SUB_acc  ...bbbbbb.....bbbb.....e..b.....

.....6.....7.....8.....9.....10
AA      PTKSHFANLKGTRGKLCPCNCFNCTDLVALGRPKCMGNTPSAKVSILH
OBS_sec
PROF_sec EEEE EEEEE EEEE
Rel_sec  67752200245344675222033211232342043234337875203787
SUB_sec  LLLL.....L...LLL.....LLLL...EEE
O_3_acc  bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc  eee bbe eeeeeee eb bbbbbbbbbb b b ee eeeeb bbb
Rel_acc  23201011020002300210201031223343310100210221541265
SUB_acc  .....b.....eb..bb

.....11.1.,....12.1.,....13.1.,....14.1.,....15.1
AA      EVKPATSGCFPIHMDRTKIRQLPNLLRGYENIRLSTSNVINTETAPGGPY
OBS_sec
PROF_sec EE HHHHHHHHHH EEEEE EE
Rel_sec  50566441224778858887664221002304663021003444556531
SUB_sec  E.LLL.....LLLHHHHHHH.....EE.....LLLL..

O_3_acc  bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc  b eeee bbbbbb e b ebbeebbe eeb eeeee
Rel_acc  12210110221001337264350314531332301200011120132011
SUB_acc  .....e.bi.b...be.....

.....16.1.,....17.1.,....18.1.,....19.1.,....20.1
AA      KVGTSKSGCPNVANGNGFFNTMAWVTPKDNNTAINPVTVEVPYICSEGED
OBS_sec
PROF_sec HHHHHHHHHH EEEEE
Rel_sec  24431001255662036787888752378864236641012244157874
SUB_sec  .....LLLL...HHHHHHHHH..LLLL..LL.....LLLL..

O_3_acc  bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc  bbbbbbbe ee bbbbbbb bbeeeee ee ebeb b b eee
Rel_acc  11111012110301026314921522122321112112322203203212
SUB_acc  .....b..bb..b.....

.....21.1.,....22.1.,....23.1.,....24.1.,....25.1
AA      QITVWGFHSDDKTQMERLYGDSNPQKFTSSANGVTTHYVSQIGGFNPQTE
OBS_sec
PROF_sec EEEEEEEE HHHHHH EEEEE EEEEEEEE
Rel_sec  47897532588742544320368853775200122343201275331035
SUB_sec  .EEEE..LLLL..H.....LLL.EEE.....LL....L

O_3_acc  bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc  bbbbbbbb eeee b eeeeeeb b b e e e ebeb b e
Rel_acc  19696993033223122210203213224011122211311001202001
SUB_acc  .bbbbbb.....b.....

.....26.1.,....27.1.,....28.1.,....29.1.,....30.1
AA      DEGLKQSGRIIVVDMVQKPGKTGTIVYQRGILLPQKVWCASGRSKVIKGS
OBS_sec
PROF_sec EEEEEEEE EEEEE EEE EEEEE EEEE
Rel_sec  5675432037788898357724788643402302235520330245401
SUB_sec  LLLL.....EEEEEE.LLL.EEEE.....EE.....E...

```

-continued

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc eee bbbbbbb bbbb ee b b bbbbbbb bb b eeb bbe e
Rel_acc 17211105270744553301222473010244520230221020211103
SUB_acc .e.....b.b.bibbb.....ib.....bbb.....
.....,....31.1.,....32.1.,....33.1.,....34.1.,....35.1
AA LPLIGADCLHEKYGLNKSHPYYTGEHAKAIGNCPIWVKTPKLKLANGTK
OBS_sec
PROF_sec HHH HHHH HHHH
Rel_sec 32233333000113432246764334340342334133220003551573
SUB_sec LLL.....HH.LL.

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc e e e bb ee b bbbbbe be bb bbbbb
Rel_acc 10010011112110110111101210102102111021712452461321
SUB_acc b.bb.bb....
.....,....36.1.,....37.1.,....38.1.,....39.1.,....40.1
AA YRPPAKLLKERGFPGAIAAGFLEGGWEGMIAGWHGYTSHGAGVAVAADLK
OBS_sec
PROF_sec EEEEEHHH EEEE H
Rel_sec 56853211324203110113404531100120021023564325530623
SUB_sec LLLL.....L.....LL..EE..L..

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc b eee be bbbbbbbbbbbbbbb bbbbbbbbbbbbbbb bbbbbbb e
Rel_acc 0001141120235599999623520456353043010011315264032
SUB_acc bbbbbbbbb..b..bbb.b..b.....b.bb....
.....,....41.1.,....42.1.,....43.1.,....44.1.,....45.1
AA STQEANKITKKNLYLSELEVKNLQRLSGAMNELHDEILELDEKVDLRA
OBS_sec
PROF_sec HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH HHH
Rel_sec 47888989873212477733101234446677898888878756210111
SUB_sec .HHHHHHHHH....HHH.....HHHHHHHHHHHHHHHHHH.....

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc bbbbeebbbe b bbee beebbeb ebbbebbe beeb eebee
Rel_acc 2804982572534232231206312453406344467553254223101
SUB_acc .b.ebb.eb.e.b.....e..eb.e.b.ebbeibee.ee.....
.....,....46.1.,....47.1.,....48.1.,....49.1.,....50.1
AA DTISSQIELAVLLSNEGIINSEDEHLALERKCLKMLGPSAVEIGNGCFE
OBS_sec
PROF_sec HHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH EEE
Rel_sec 0364547777753464114550578879888887664424540663252
SUB_sec ..H.H.HHHHHHHH..L...LL.HHHHHHHHHHHHHHHHH..H..LL..E.

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc bbbb bb bbbbb eebb e e b eb e b e b e bee eebbbb
Rel_acc 04450592968651122001021323439443944173428232012340
SUB_acc .bbb.bb.bbbbb.....i.bie.bie.b.e.b.....b.
.....,....51.1.,....52.1.,....53.1.,....54.1.,....55.1
AA TKHKCNQTCLDRIAAGTFNAGDFSLPTFDSLNTAASLNDGDLNHTILL (SEQ ID NO: 36)
OBS_sec
PROF_sec EEEE HHHHHHHH HHHH EEEEEEEE EEEE
Rel_sec 22003402344311355466523223103101002476217776303665
SUB_sec LL.LLL.....EE.....EEE

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc b beb e bbbbb bbbb ee eebbe beb bbeb eeeeeeb bbb
Rel_acc 42120312620271012211330320211224131235303222133868
SUB_acc b.....b..b.....e.....b.....bbb
.....,....56.1.,....57.1.,....58.1. (SEQ ID NO:36)
AA YYSTAASSLAVTLMIATFIVYMVSRDNVSCSICL
OBS_sec
PROF_sec EEEEEHHHHHHHHHHHHHHHHH EEEEEEE
Rel_sec 4310021478888888776321006740346860
SUB_sec HHHHHHHHHH.....LL....EEE.

O_3_acc bbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbbb
P_3_acc bbbbbbbbbbbbbbbbbbbbbbbbee eb b bbb
Rel_acc 6362366679999996998517101010152400
SUB_acc b.b..bbbbbbbbbbbbbb.b.....b.b..

[0408] The third tool that used is the hydrophobicity analysis. In general, the structural core of a protein tends to be a hydrophobic cluster flanked by stretch of hydrophilic residues. Therefore the predicted boundaries should not locate in the center of hydrophobic core, rather should reside in the flanking hydrophilic regions. Hydrophobicity of B/Lee/40 HA sequence (SEQ ID NO: 36) was plotted using ProtScale (<http://ca.expasy.org/tools>→Primary structure analysis→ProtScale). The hydrophobicity analysis of B/Lee/40 sequence confirmed that the selected boundaries were in the hydrophilic regions of the protein (FIG. 2). The hydrophobicity strength used in the plot is listed in the following table:

SEQUENCE LENGTH: 584	
Using the scale Hphob./Kyte & Doolittle, the individual values for the 20 amino acids are:	
Ala:	1.800
Arg:	-4.500
Asn:	-3.500
Asp:	-3.500
Cys:	2.500
Gln:	-3.500
Glu:	-3.500
Gly:	-0.400
His:	-3.200
Ile:	4.500
Leu:	3.800
Lys:	-3.900
Met:	1.900
Phe:	2.800
Pro:	-1.600
Ser:	-0.800
Thr:	-0.700
Trp:	-0.900
Tyr:	-1.300
Val:	4.200
Asx:	-3.500
Glx:	-3.500
Xaa:	-0.490

results indicated that influenza B HA proteins are much more conserved compared to influenza A HA proteins, especially in the domain boundary sequences. Thus, the boundaries identified in B/Lee/40 were used to direct boundary selections of other influenza B strains. The resulting domain boundaries for each strain are summarized below.

[0410] B/Malaysia/2506/2004: HA1-1 44-337 (SEQ ID NO: 45) (TTTPTK (SEQ ID NO: 287)-CPIWVK (SEQ ID NO: 288)) HA1-2 56-296 (SEQ ID NO: 46) (KGTETR (SEQ ID NO: 289)-SKVIKG (SEQ ID NO: 290))

[0411] B/Ohio/1/2005: HA1-1 33-326 (SEQ ID NO: 47) (TTTPTK (SEQ ID NO: 291)-CPIWVK (SEQ ID NO: 292))

[0409] In order to identify the HA 1-1 and HA1-2 domain boundaries of other influenza B molecules, such as B/Malaysia/2506/2004 (SEQ ID NO: 42), B/Ohio/1/2005 (SEQ ID NO: 43), B/Victoria/2/87 lineage (SEQ ID NO: 787) and B/Shanghai/361/2002 (SEQ ID NO: 44), and B/Yamagata/16/88 lineage (SEQ ID NO: 213), each HA sequence was aligned with B/Lee/40 HA sequence (SEQ ID NO: 36). The

HA1-2 45-285 (SEQ ID NO: 48) (KGTKTR (SEQ ID NO: 293)-SKVIKG (SEQ ID NO: 294))

[0412] B/Shanghai/361/2002: HA1-1 33-325 (SEQ ID NO: 49) (TTTPIK (SEQ ID NO: 295)-CPIWVK (SEQ ID NO: 296)) HA1-2 45-284 (SEQ ID NO: 50) (KGTETR (SEQ ID NO: 297)-SKVIKG (SEQ ID NO: 298))

```

Bma1 ----IVLLMVVTSNADRICTGITSSNSPHVVKATQGEVNVTVGIPLTTT    50 (SEQ ID NO: 42)
Bohi -----DRICTGITSSNSPHVVKATQGEVNVTVGIPLTTT            (SEQ ID NO: 43)
BLee MKAIIVLLMVVTSNADRICTGITSSNSPHVVKATQGEVNVTVGIPLTTT    (SEQ ID NO: 36)
BSha -----DRICTGITSSNSPHVVKATQGEVNVTVGIPLTTT            (SEQ ID NO: 44)
*****

Bma1 PTKSHFANLKGTETRGKICPKCLNCTDLVALGRPKCTGNIPSARVSILH    100
Bohi PTKSHFANLKGTETRGKICPKCLNCTDLVALGRPKCTGNIPSAEVSILH
BLee PTKSHFANLKGTETRGKICPKCLNCTDLVALGRPKCMGNTPSAKVSILH
BSha PIKSHFANLKGTETRGKICPKCLNCTDLVALGRPMCVGTTTPSAKASILH
* *****.*:***** * *.***.***

Bma1 EVRPVTSGCCFPIMHDKIRQLPNLLRGYEHIRLSTHNVINAENAPGGSY    150
Bohi EVRPVTSGCCFPIMHDKIRQLPNLLRGYEHIRLSTHNVINAENAPGGSY
BLee EVKPATSGCCFPIMHDKIRQLPNLLRGYENIRLSTSNVINTETAPGGPY
BSha EVRPVTSGCCFPIMHDKIRQLPNLLRGYENIRLSTQNVIDAEKALGGPY
***.*:*****:***** ***:.*.***

Bma1 KIGTSGSCPNVTNGGFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGE    200
Bohi KIGTSGSCPNVTNGGFFATMAWAVPKMDNNKTATNSLTIEVPYICTEGE
BLee KVGTSKSCPNVANGGFFNTMAWVIPK-DNNKTAINPVTVEVPYICTEGE
BSha RLGTSGSCPNATSKSGFFATMAWAVPK-DNNKNATNPLTVEVPYICTEGE
*:*****.:.*** ***:.*:*****:***

Bma1 DQITVWGFHSDNEAQMALKYGDSPQKFTSSANGVTTHYVSQIGGFPNQ    250
Bohi DQITVWGFHSDSETQMAKLYGDSKPQKFTSSANGVTTHYVSQIGGFPNQ
BLee DQITVWGFHSDDKTQMERLYGDSNPQKFTSSANGVTTHYVSQIGGFPNQ
BSha DQITVWGFHSDDKTQMKNLGDSNPQKFTSSANGVTTHYVSQIGGFPDQT
***:*****.:**.*:*****:*****:***

```

-continued

```

BMal EDGGLPQSGRIVVDYVMVQKSGKTGTITYQRGILLPQKVWCASGRSKVIK 300
Bohi EDGGLPQSGRIVVDYVMVQKSGKTGTITYQRGILLPQKVWCASGRSKVIK
BLee EDEGLKQSGRIVVDYVMVQKPGKTGTIVYQRGILLPQKVWCASGRSKVIK
BSha EDGGLPQSGRIVVDYVMVQKPGKTGTIVYQRGVLLPQKVWCASGRSKVIK
      * * * * *
BMal SLPLIGEADCLHEKYGGLNKSPPYYTGEHAKAIGNCPIWVKTPCLKLANGT 350
Bohi SLPLIGEADCLHEKYGGLNKSPPYYTGEHAKAIGNCPIWVKTPLIQANGT
BLee SLPLIGEADCLHEKYGGLNKSPPYYTGEHAKAIGNCPIWVKTPCLKLANGT
BSha SLPLIGEADCLHEKYGGLNKSPPYYTGEHAKAIGHCPIWVKTPCLKLANGT
      * * * * *
BMal KYRPPAKLLKER----- 400
Bohi KYRPPAKLLKERGF-----
BLee KYRPPAKLLKERGFFGAIAGFLEGGWEGMIAGWHGYTSHGAHGVAVAADL
BSha KYRP-----
      * * *

```

[0413] B/Lee/40 HA1-1 construct design (SEQ ID NO: 38). For this construct, five (5) anticipated conserved disulfide bonds were preserved by making the amino terminal truncation before the Cysteine at position 69 and the carboxyl-terminal truncation after the Cysteine at position 335 of SEQ ID NO: 36. The likely disulfide pairings are as the following: C69 and C72, C75 and C87, C109 and C158, C195 and C289, C309 and C335 of SEQ ID NO: 36. Using the simulated structures and the MMDB to check secondary structures the amino terminal truncation was made at the Theonine at position 48, right after the β -strand kink introduced by Proline 46 followed by, Leucine 47 of SEQ ID NO: 36, such that the preceding β -strand [TGVI, SEQ ID NO: 299, residues 42-45 of SEQ ID NO: 36] was totally eliminated while the entire short β -strand formed by residues TTTP (SEQ ID NO: 300) [residues 48-51 of SEQ ID NO: 36] was preserved. The carboxyl-terminal truncation was made at the end of β -strand PIWV (SEQ ID NO: 301) [residues 336-340 of SEQ ID NO: 36] following the Cysteine at position 335 such that the carboxyl-terminal tail which interacts with HA2 was completely eliminated. Six α -strands including the C-terminal β -strand form a stable β -sandwich defined by β -strands PIWV (SEQ ID NO: 301) [residue 336-340 of SEQ ID NO: 36], PYYTG (SEQ ID NO: 302) [residues 322 to 326 of SEQ ID NO: 36] and TTTP (SEQ ID NO: 300) [residues 48-51 of SEQ ID NO: 36] (lower β -pleated sheet), and the β -strands DCLHE (SEQ ID NO: 303) [residues 308-312 of SEQ ID NO: 38], YGGLN (SEQ ID NO: 304) [residue 314-318 of SEQ ID NO: 36] and KAIGN (SEQ ID NO: 305) [residues 330-334 of SEQ ID NO: 36] (top β -pleated sheet). This set of β -pleated sheet includes both the amino-terminus and carboxyl-terminus within one stable secondary structure element. Such truncations were made to believe resulting a compact domain structure.

[0414] B/Lee/40 HA1-2 construct design (SEQ ID NO: 39). For this construct four (4) anticipated conserved disulfide bonds were preserved by making the amino-terminal truncation before the Cysteine at position 69 and the carboxyl-terminal truncation after the Cysteine at position 289 of SEQ ID NO: 36. Using the simulated structure and the MMDB to check secondary structures the amino-terminal truncation was made in the loops that connect two distinct sets of β -pleated sheets. The amino-terminal truncation was made at the Lysine at position 60, such that the β -strand comprising residues 62-66 of SEQ ID NO: 36 [TQTRG,

SEQ ID NO: 306] of the membrane-distal set of β -strands remained wholly intact, while the membrane-proximal β -strand [TTTP, SEQ ID NO: 300, residues 48-51 of SEQ ID NO: 36] that forms part of HA1-1 was completely eliminated. The carboxyl-terminal truncation was made at the Serine 300 of SEQ ID NO: 36. Truncation at this position preserved secondary structure such that the membrane-distal β -strand [KVIKG, SEQ ID NO: 307, residues 295-299 of SEQ ID NO: 36] remained intact while the membrane-proximal β -strand [DCLHE, SEQ ID NO: 308, residues 308-312 of SEQ ID NO: 36] was eliminated. The membrane-distal β -pleated sheets comprising β -strands TQTRG (SEQ ID NO: 306) [residues 62-66 of SEQ ID NO: 36], SILHEV (SEQ ID NO: 309) [residues 97-102 of SEQ ID NO: 36], and KVIKG (SEQ ID NO: 307) [residues 295-299 of SEQ ID NO: 36] serve as the stabilizing secondary structure element that is believed to conclude a compact domain structure.

[0415] B/Malaysia/2506/2004, B/Ohio/1/2005 and B/Shanghai/361/2002 follow the same reference structure as described above.

Example 3

Cloning and Expression of Recombinant
Flagellin-Hemagglutinin Fusion Proteins in *E. coli*

Materials and Methods

[0416] Cloning of HA influenza A subunits. Subunits of the HA globular head from several strains of influenza. A were cloned and expressed alone or as fusions with flagellin. We also expressed in fusion with the HA globular head domain, two proteins widely used as carrier proteins in conjugated vaccines. CRM-197 is a mutated diphtheria toxin (DTx) from *Corynebacterium diphtheriae* and LTb, is *E. coli* heat labile toxin B subunit.

[0417] These constructs were generated in one of four different methodologies:

[0418] Method #1: In this protocol, a fusion gene comprising flagellin (STF2) (SEQ ID NO: 212) and the HA subunit was codon-optimized for *E. coli* expression and obtained from a commercial vendor (DNA2.0 Inc., Menlo Park, Calif.) by chemical synthesis. The gene was excised with NdeI and BlnI enzymes, the insert fragment was gel purified and ligated to pET24a (Novagen, San Diego, Calif.)

which had been digested with NdeI and BlnI and treated with bacterial alkaline phosphatase (BAP).

[0419] Method #2: For facile cloning of genes in fusion with flagellin, a cassette plasmid containing a unique BlnI site at the 3' end of the flagellin (STF2) gene (SEQ ID NO: 212) was generated. This was done by introducing a silent mutation at nucleotides 5'-GTGCTGAGCCTGTTACGT-3' (SEQ ID NO: 310) [nt 1501 to 1518 of SEQ ID NO: 212] of STF2 creating the unique BlnI site in the plasmid cassette, pET24/STF2.blp (SEQ ID NO: 51). Synthetic genes for each target antigen were codon-optimized for *E. coli* expression and obtained from a commercial vendor (DNA2.0 Inc., Menlo Park, Calif.). The synthetic genes were excised with BlnI enzyme and ligated by compatible ends to pET24/STF2.blp which had been treated with BlnI and BAP.

[0420] Method #3: Using forward and reverse primers (Keck Foundation BRLK, Yale University, New Haven, Conn.; Midland Certified Reagent Company, Midland, Tex.) as indicated for each construct, PCR amplification was performed using the DNA templates shown in each table. The PCR product was subjected to BlnI digestion, gel-purified and ligated to pET24/STF2.blp (SEQ ID NO: 51) vector previously prepared by BlnI digestion and BAP treatment.

[0421] Method #4: A plasmid cassette, pET24/STF2.5G (SEQ ID NO: 52), was generated by introducing a flexible heptamer linker, Ser-Gly-Ser-Gly-Ser-Gly-Ser (S-G-S-G-S-G-S (SEQ ID NO: 311)) at the 3' end of the STF2 gene. A unique BamHI site was created in the linker to facilitate cloning of HA subunit fragments in fusion with flagellin. Synthetic genes codon-optimized for expression in *E. coli* were obtained from a commercial vendor (DNA2.0 Inc., Menlo Park, Calif.), excised from the parental plasmids with BamHI and BlnI restriction endonucleases, and ligated by compatible ends to the pET24/STF2 cassette that had previously been digested with BamHI and BAP-treated.

[0422] In each case, the constructed plasmids were used to transform competent *E. coli* TOP10 cells and putative recombinants were identified by PCR screening and restriction mapping analysis. The integrity of the constructs was verified by DNA sequencing and they were used to transform the expression host, BLR3 (DE3) (Novagen, San Diego, Calif.; Cat #69053). Transformants were selected on plates containing kanamycin (50 µg/mL), tetracycline (5 µg/mL) and glucose (0.5%). Colonies were picked and inoculated into 2 ml of LB medium supplemented with 25 µg/ml kanamycin, 12.5 µg/ml tetracycline and 0.5% glucose and grown overnight. Aliquots of these cultures were used to inoculate fresh cultures in the same medium formulation, which were cultured until they reached an OD₆₀₀=0.6, at which time protein expression was induced by the addition of 1 mM IPTG and culturing for 3 hours at 37° C. The cells were harvested and analyzed for protein expression.

[0423] SDS-PAGE and Western blot: Protein expression and identity were determined by gel electrophoresis and immunoblot analysis. Cells were harvested by centrifugation and lysed in Laemmli buffer. An aliquot of 10 µl of each lysate was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel and electrophoresed (SDS-PAGE). The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.)

to visualize protein bands. For western blot, 0.5 µl/lane cell lysate was electrophoresed and electrotransferred to a PVDF membrane and blocked with 5% (w/v) dry milk.

[0424] The membrane was then probed with either anti-flagellin antibody (Inotek; Beverly, Mass.) or influenza A PR/8/38 convalescent immune mouse serum. PR/8/34 immune serum was generated in BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) that received an experimentally determined sublethal challenge dose of 8×10⁴ egg infectious dosages (EID) of PR/8/34 influenza virus. Animals were then allowed to convalesce for >21 days post-infection at which time immune serum was isolated and clarified. After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockland, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega, Madison, Wis.). Bacterial clones which yielded protein bands of the correct molecular weight and reactive with the appropriate antibodies were selected for production of protein for use in biological assays.

[0425] The constructs derived from the HA of A/Puerto Rico/8/34 strain (PR8) (SEQ ID NO: 1) and listed in Table 1 were made by the synthetic gene route as described in Method #1 and Method #4. Similarly, the constructs derived from the A/Viet Nam/1203/2004 strain (SEQ ID NO: 2) are shown in Table 2, the constructs derived from the HA of A/Indonesia/2005 strain (IND) (SEQ ID NO: 3) are described in Table 3, and those from A/New Caledonia/12/99 strain (NC) (SEQ ID NO: 4) are described in Tables 4. "IND" as used herein, refers to "Indonesia." Where appropriate, the DNA primers and DNA templates used in the PCR amplification reaction are listed in the same table.

TABLE 1

PR8 HA constructs for expression in <i>E. coli</i>		
SEQ ID NO:	Construct	Method
53	STF2.HA1-1	#4
54	STF2.HA1-1.his	#4
55	STF2.HA1-2	#1
56	STF2.HA1-2mut	#1
57	STF2.HA1-3	#1
58	STF2.HA1-3mut	#1
59	HA1-1	#1
60	HA1-1.his	#1
61	HA1-2.his	#1
62	CRM.HA1-2	#1
63	LTB.HA1-2	#1

[0426]

TABLE 2

VN HA constructs for expression in <i>E. coli</i>					
SEQ ID NO:	Construct	Method	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:
64	STF2.HA1-1	#3	65	66	67
68	STF2.HA1-2	#3	69	70	67
71	STF2.HA1-2mut	#2	N/A	N/A	72

[0427]

TABLE 3

IND HA constructs for expression in <i>E. coli</i>					
SEQ ID NO:	Construct	Method	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:
73	STF2.HA1-1	#3	74	66	75
76	STF2.HA1-2	#3	69	77	75
78	STF2.HA1-2mut	#2	N/A	N/A	79

[0428]

TABLE 4

NC HA constructs for expression in <i>E. coli</i>					
SEQ ID NO:	Construct	Method	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:
80	STF2.HA1-1	#3	81	82	83
84	STF2.HA1-2	#3	85	86	83
87	STF2.HA1-2mut	#2	N/A	N/A	88

Results

[0429] Protein carriers have widespread application in human vaccines. The cross-reactive reactive material (CRM₁₉₇) of diphtheria toxin is considered to be advantageous as a carrier molecule in the formulation of several conjugate vaccines. Exemplary carriers include *E. coli* heat labile enterotoxin (LT) and its B subunit (LTB), Tetanus toxoid (TT) and cholera toxin (CT). Using CRM₁₉₇ and LTB as representatives of this group of carrier proteins, we have generated constructs in which the globular head of HA of (A/Puerto Rico/8/34 strain (PR8) (SEQ ID NO: 1) is fused to the 3' end of either CRM197 gene or LTB gene as described in Method #1 generating the constructs CRM.HA1-2 (SEQ ID NO: 62) and LTB.HA1-2 (SEQ ID NO: 63). The constructs were verified by DNA sequencing and used to transform the expression host, BLR3 (DE3) (Novagen, San Diego, Calif.; Cat #69053).

[0430] Transformants were selected on plates containing kanamycin (50 µg/mL), tetracycline (5 µg/mL) and glucose (0.5%). Several colonies were picked for an overnight culture which were used to inoculate a fresh LB medium supplemented with 25 µg/ml kanamycin, 12.5 µg/ml tetracycline and 0.5% glucose. At an OD₆₀₀=0.6 protein expression was induced with 1 mM IPTG for 3 h at 37° C. The cells were harvested and an aliquot of the lysate was analyzed on 10% SDS-PAGE by Coomassie blue staining and by immunoblot using PR/8/34 convalescent sera. In the case of CRM.HA1-2 construct several clones were picked and analyzed for expression by SDS-PAGE.

[0431] As assayed by Coomassie blue staining of the SDS-PAGE gel, all the clones displayed a band that migrated with an apparent MW 84 KDa and that corresponds to the predicted MW. The absence of this band in the control culture (without IPTG) indicates that it is specifically induced by IPTG. This observation was further confirmed

when the cell extracts of two clones #5 and #6 was fractionated in the presence or absence of a reducing agent (5 mM DTT). While the recombinant protein whose disulphide bonds have been disrupted by treatment with DTT is not recognized by the cognate antibodies, the native recombinant protein is. Similarly, a clone expressing construct LTB.HA1-2 displays a band corresponding to the predicted molecular weight of 39.8 KDa, protein when induced with IPTG. The identity of LTB-HA1-2 fusion protein is confirmed by western blot analysis using mouse convalescent serum. This band is diminished in intensity when a reductant (β-mercaptoethanol) was present. This latter observation suggests that insufficient amount of reductant was most likely employed in the experiment. Taken together the data presented herein support the notion that the globular head of HA can be successfully fused to carrier proteins to generate conformational sensitive proteins.

Cloning of Recombinant Flagellin-Hemagglutinin Fusion Proteins in *E. coli*

[0432] Cloning of HA Influenza B subunits. Subunits of the HA globular head from several strains of influenza B were cloned and expressed as fusions with flagellin. These constructs were generated by a two-step PCR.

[0433] The HA subunit was codon-optimized for *E. coli* expression and obtained from a commercial vendor (DNA2.0 Inc., Menlo Park, Calif.) by chemical synthesis. The HA1-1 or HA1-2 was PCR amplified using the synthesized DNA as templates. The flagellin (STF2) sequence (SEQ ID NO: 212) was derived from the plasmid pET24a-STF2.HA1-2. The STF2 DNA fragment was PCR amplified, and the C-terminal of the PCR product has a 28-30 bp overlap with the N-terminal sequence of the fusion HA subunit.

[0434] The STF2 and HA subunits were fused together by a 2nd PCR. Using respective forward and reverse primers (Integrated DNA Technologies, Inc, Coralville, Iowa 52241) listed below, fusion protein DNA was amplified from the DNA templates also shown below. The PCR product was subsequently subjected to XbaI digestion, gel-purification and ligation to pET24a-STF2.HA1-2 that was previously digested with XbaI and SnaBI.

[0435] The constructs are listed below, where appropriate, the DNA primers and DNA templates used in the PCR amplification reaction are also listed.

FluB STF2.HA constructs for expression in <i>E. coli</i>					
SEQ ID NO:	Construct	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:	
184	STF2.HA1-1 (MAL)	193, 195	194, 196	55, 183	
186	STF2.HA1-2 (MAL)	193, 198	197, 199	55, 185	
188	STF2.HA1-2 (SH)	193, 201	200, 202	55, 187	
190	STF2.HA1-2 (Lee)	193, 204	203, 205	55, 189	
192	STF2.HA1-2 (Ohio)	193, 207	206, 208	55, 191	

[0436] In each case, the constructed plasmids were used to transform competent *E. coli* DH5α cells and putative recombinants were identified by restriction mapping. The integrity

of the constructs was verified by DNA sequencing and was used to transform the expression host, BLR3 (DE3) (Novagen, San Diego, Calif.; Cat #69053). Transformants were selected on LB (Veggie peptone, EMD Bioscience, La Jolla, Calif.; Cat#71280; veggie yeast extract, EMD Bioscience, Cat# 71279;) plates containing kanamycin (35 µg/mL). Colonies were picked and inoculated into 2 ml of LB (Veggie) medium supplemented with 35 µg/ml kanamycin and grown overnight. Aliquots of these cultures were used to inoculate fresh cultures in the same medium formulation, which were cultured until OD₆₀₀=0.6 was reached. The protein expression was induced by the addition of 1 mM IPTG. The cells were harvested after 2 hours culturing at 37° C. and analyzed for protein expression.

[0437] SDS-PAGE and Western blot: Protein expression and identity were determined by gel electrophoresis and immunoblot analysis. Cells were harvested by centrifugation and lysed in Tris/NaCl (50 mM Tris, 200 mM NaCl, pH 8.0) buffer. Two aliquots of each lysate was diluted in SDS-PAGE sample buffer and boiled for 5 minutes. One aliquot sample was reduced by DTT (200 mM final concentration, EMD Bioscience, La Jolla, Calif. 92039; Cat#233155). The samples (1.5 µl/lane cell lysate) were loaded onto a 4-12% SDS polyacrylamide gel and electrophoresed (SDS-PAGE). The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands.

[0438] For western blot, the duplicated gel was electrophoresed and electrotransferred to a PVDF membrane and blocked with 5% (w/v) dry milk. The membrane was then probed with either anti-flagellin antibody (6H11, Inotek, Beverly, Mass.; Lot#1030B5) or ferret antiserum raised against B/Malaysia (2506/2004, CDC#2005741987). After probing with Horseradish Peroxidase (HRP)-conjugated secondary antibodies (1:10,000) (Goat anti-Mouse is from Jackson ImmunoResearch Laboratories; West Grove, Pa. Goat anti-Ferret is from Bethyl Laboratories; Montgomery, Tex.), protein bands were visualized after with addition of TMB, an HRP chromogenic substrate (Pierce; Rockland, Ill.; Cat#34018). The approximate molecular weight of STF2.HA1-1(MAL) is 84 kDa; and the approximate molecular weight of STF2.HA1-2 (MAL and SH) is ~78 kDa. Two micrograms of STF2.4×M2e fusion protein was loaded as a positive control for the anti-flagellin antibody and a negative control for the anti-HA sera. Virus lysate (B/Malaysia/2506/04, CDC# 2005756250) was included in the experiment as a positive control for the anti-HA sera and a negative control for the anti-flagellin antibody. Commassie blue staining of the gel showed low induction of STF2.HA1-1(MAL) protein; but much stronger bands for STF2.HA1-2(MAL) and STF2.HA1-2(SH) proteins. Western blot analysis using anti-flagellin monoclonal antibodies (1:4000) exhibited a strong positive band around 84 kDa for lanes loaded with STF2.HA1-1(MAL), and around 75 kDa in lanes loaded with STF2.HA1-2(MAL) or STF2.HA1-2(SH). The induction pattern of each fusion protein on the Western is very similar to the result for the commassie blue stained gel. The level of the non-reduced and the reduced protein recognized by the anti-flagellin was comparable. The positive control for anti-flagellin (STF2.4×M2e) showed a single strong band, and the negative control exhibited no signal.

[0439] Western blot analysis using anti-HA polyclonal anti-sera (1:1000) exhibited a similar pattern as the blot with anti-flagellin, except that the reduced STF2.HA1-2 protein

had a diminished signal compared to that of the non-reduced sample, indicating that a substantial portion of the anti-HA polyclonal antibodies in the antiserum recognize conformational epitopes formed by correct disulfide-bond formation while a small proportion of the antibodies in the anti-sera recognize linear epitopes that are not reliant on disulfide bound formation. Similar were observed using a capture ELISA assay. It is worth noting that anti-serum against B/Malaysia/2506/2004 (Victoria lineage) also recognizes B/Shanghai/361/2002, which belongs to Yamagata lineage. This result indicates that these two lineages not only share substantial primary sequence similarity, but also share tertiary structure.

[0440] The western blot and ELISA results confirm the expression of functional Influenza B HA1-1 and HA1-2 fusion proteins in *E. coli*.

Example 4

Purification and Biochemical Characterization of Recombinant Flagellin-Hemagglutinin Fusion Proteins

Materials and Methods

[0441] Bacterial growth and cell lysis: HA constructs were expressed in the *E. coli* host strain BLR (DE3). *E. coli* cells carrying a plasmid encoding STF2.HA1-1.his(PR8) (SEQ ID NO: 54) were cultured and harvested as described above. Individual strains were retrieved from glycerol stocks and grown in shake flasks to a final volume of 12 L. Cells were grown in LB medium containing 50 µg/ml kanamycin/12.5 µg/ml tetracycline/0.5% dextrose to OD₆₀₀=0.6 and induced with 1 mM IPTG for 3 h at 37° C. The cells were harvested by centrifugation (7000 rpm×7 minutes in a Sorvall RC5C centrifuge) and resuspended in 1×PBS, 1% glycerol, 1 µg/ml DNaseI, 1 mM PMSF, protease inhibitor cocktail and 1 mg/ml lysozyme. The cells were then lysed by two passes through a microfluidizer at 15,000 psi. The lysate was centrifuged at 45,000 g for one hour in a Beckman Optima L ultracentrifuge (Beckman Coulter; Fullerton, Calif.) to separate the soluble and insoluble fractions

[0442] Purification and refolding of STF2.HA1-1.his(PR8) (SEQ ID NO: 89) protein from *E. coli*. Following lysis and centrifugation, the insoluble (pellet) fraction was resuspended in 100 ml 1×TBS, pH 8.0+1 mM β-mercaptoethanol and solid urea was added to a final concentration of 6.2 M. After resuspension with a Dounce glass-ball homogenizer the mixture was centrifuged as described above. The resulting supernatant was loaded onto a chelating sepharose column (GE/Amersham Biosciences; Piscataway, N.J.) charged with NiSO₄ and equilibrated in Buffer A [20 mM Tris, pH 8.0/8 M urea/0.5 M NaCl].

[0443] After washing with 1 L buffer B [Buffer A+1% (w/v) TX-100] the column was eluted in a 5-column volume linear gradient from 100% Buffer A to 100% Buffer C [Buffer A+0.5 M imidazole]. Peak fractions were pooled, concentrated 3.5 fold on an Amicon 15 spin concentrator (Millipore; Billerica, Mass.) and dialyzed against 3×2 L of 8 M urea/20 mM Tris, pH 8.0/2 mM EDTA. Following dialysis, STF2.HA1-1.his(PR8) (SEQ ID NO: 89) was refolded by rapid dilution to a final concentration of 0.1 mg/ml protein in Refolding Buffer [0.1 M Tris, pH 8.0/0.1

M NaCl/1% (w/v) glycerol/5 mM reduced glutathione/1 mM oxidized glutathione] and incubated overnight at room temperature.

[0444] The refolded protein was then captured on a 5 ml chelating sepharose HiTrap column (GE/Amersham Biosciences; Piscataway, N.J.) charged with NiSO₄ and equilibrated in Buffer D [20 mM Tris, pH 8.0/0.5 M NaCl/1 mM reduced glutathione/0.2 mM oxidized glutathione] and eluted in 100% Buffer E [Buffer D+0.5 M imidazole]. Peak eluate fractions were pooled, concentrated using an Amicon 15 spin concentrator (Millipore; Billerica, Mass.) and fractionated on a Superdex 200 size-exclusion column (GE/Amersham; Piscataway, N.J.) equilibrated in 1×TBS, pH 8.0+0.5% (w/v) Na deoxycholate. S200 peak fractions were pooled, dialyzed against 3×2 L of 1×TBS, pH 8.0 (without deoxycholate), vialled and stored at -80° C.

[0445] Purification of STF2.HA1-2(PR8) (SEQ ID NO: 90) and STF2.HA1-2mut(PR8) (SEQ ID NO: 91): *E. coli* cells carrying a plasmid encoding STF2.HA1-2.his (PR8) (SEQ ID NO: 55) were cultured and harvested as described above. Following lysis and centrifugation, the insoluble (pellet) fraction containing STF2.HA1-2(PR8) was resuspended and homogenized in 50 mM Tris, pH 8.0+0.5% (w/v) Triton X-100 using a Dounce homogenizer. The homogenate was then centrifuged (19,000 rpm×10 min). The resulting pellet fraction was then re-homogenized in 50 mM Tris, pH 8.0+0.5% (w/v) Triton X-100+1.0 M NaCl and centrifuged. The inclusion body fraction (pellet) was then dissolved in buffer A [50 mM Na Acetate, pH 4.0+8.0 M urea] and centrifuged (19,000 rpm×10 min.) to remove insoluble debris.

[0446] The resulting supernatant was fractionated on a Source S column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated in Buffer A and eluted in a 5 column-volume linear gradient with Buffer B (50 mM Na Acetate, pH 4.0, 1.0 M NaCl). The protein was then refolded by rapid dilution into Buffer C (100 mM Tris-HCl, pH 8.0). The refolded protein was then fractionated on a Q sepharose HP column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated in Buffer C and eluted in a 0% to 60% linear gradient with buffer D (Buffer C+1 M NaCl). The Q HP eluate was then fractionated on a Superdex 200 size exclusion chromatography (SEC) column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated with 1× Tris-buffered saline, pH 8.0, to separate monomeric from aggregated protein. Monomeric fractions were then pooled, aliquoted and stored at -80° C.

[0447] Purification of STF2.HA1-3(PR8) (SEQ ID NO: 92): *E. coli* cells carrying a plasmid encoding STF2.HA1-3 (PR8) (SEQ ID NO: 57) were cultured and harvested as described above. Following lysis and centrifugation the soluble (supernatant) fraction containing STF2.HA1-3(PR8) was adjusted to 6.2 M urea, 100 mM DTT and incubated at room temperature overnight to completely reduce and denature the protein. The protein was then diluted five-fold in Buffer A (50 mM citric acid, pH 3.5, 8 M urea, 1 mM β-mercaptoethanol, 1 mM EDTA) and loaded onto an SP sepharose FF column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated with Buffer A. To remove endotoxin, the column washed with 10 column volumes of Buffer B (Buffer A+1% (w/v) Triton X-100) followed by 10 column volumes Buffer C (50 mM citric acid, pH 3.5, 1 mM β-mercaptoethanol, 1 mM EDTA, 70% (v/v) isopropanol).

[0448] The bound protein was then eluted in a five column-volume linear gradient of Buffer A:Buffer D (Buffer A+1M NaCl). The eluted protein was then dialyzed to Buffer E (20 mM Tris, pH 8.5, 8 M urea, 1 mM β-mercaptoethanol, 1 mM EDTA) and passed through a Source Q column (GE/Amersham Biosciences; Piscataway, N.J.). The STF2.HA1-3(PR8) (SEQ ID NO: 92) protein was collected in the flow-thru fraction. The protein was concentrated using Amicon spin-concentrators (Millipore; Billerica, Mass.) and dialyzed to Buffer F (20 mM Tris, pH 8.0, 8 M urea, 1 mM EDTA) to remove reductant. The denatured polypeptide was then refolded by rapid dilution to a final concentration of 0.1 mg/ml in Refolding Buffer (0.1 M Tris, pH 8.0, 0.1 M NaCl, 1% (w/v) glycerol, 5 mM reduced glutathione, 1 mM oxidized glutathione) and incubated overnight at room temperature. The refolded protein was then fractionated on a Superdex 200 size exclusion column equilibrated in 1× Tris-buffered saline (TBS), pH 8.0, plus 0.25% (w/v) sodium deoxycholate to separate monomeric STF2.HA1-3(PR8) from aggregated protein. Monomeric peak fractions were pooled, dialyzed to 1×TBS, pH 8.0, to remove deoxycholate, aliquoted and stored at -80° C.

[0449] Purification of HA1-2His(PR8) (SEQ ID NO: 93): *E. coli* cells carrying a plasmid encoding STF2.HA1-2.his (PR8) (SEQ ID NO: 61) were cultured and harvested as described above. Following cell lysis the insoluble fraction was homogenized and washed once with Buffer A (50 mM Tris, pH 8.0), followed by sequential washing with Buffer A+0.5% (w/v) Triton X-100 and Buffer A+0.5% (w/v) Triton X-100+1M NaCl. The insoluble material was then washed twice more with Buffer A alone. The washed inclusion bodies were then dissolved in Buffer B (1×PBS, pH 7.4, 8 M urea) and applied to a Ni-NTA (Sigma-Aldrich; St. Louis, Mo.) column charged with NiSO₄ and eluted in a linear gradient of Buffer C (50 mM Tris, pH 8.0, 8 M urea, 0.3 M NaCl, 0.5 M imidazole). In order to reduce endotoxin, the eluted protein was dialyzed overnight against 1×PBS, pH 7.4, 8 M urea, and reappplied to the Ni-NTA column.

[0450] The column was then washed with Buffer D (10 mM Tris, pH 8.8, 8 M urea, 60% (v/v) isopropanol) and eluted as described above. Ni-NTA purified protein was then refolded by 1:10 (v/v) dilution into Refolding Buffer (50 mM Tris, pH 8.8). The refolded HA1-2 His(PR8) (SEQ ID NO: 93) protein was then concentrated using an Amicon ultrafiltration unit (Millipore; Billerica, Mass.) and applied to a Superdex 200 size exclusion column (GE/Amersham; Piscataway, N.J.) equilibrated in 1×TBS, pH 8.0. Monomeric peak fractions were pooled, aliquoted and stored at -80° C.

[0451] SDS-PAGE and western blot analysis: Protein identity of all HA constructs was determined, and purity estimated, by SDS-PAGE. An aliquot of 5 μg of each sample was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel (LifeGels; French's Forrest, New South Wales, AUS) and electrophoresed. The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands. For western blot, 0.5 μg/lane total protein was electrophoresed as described above and the gels were then electro-transferred to a PVDF membrane and blocked with 5% (w/v) dry milk before probing with anti-flagellin antibody (Inotek; Beverly, Mass.) or influenza A PR/8/34 Convalescent immune serum

(described below under Protein Antigenicity ELISA). After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockland, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega; Madison, Wis.).

[0452] Protein assay: Total protein concentration for all proteins was determined using the Micro BCA (bicinchoninic acid) Assay (Pierce; Rockford Ill.) in the microplate format, using bovine serum albumin as a standard, according to the manufacturer's instructions.

[0453] Endotoxin assay: Endotoxin levels for all proteins were determined using the QCL-1000 Quantitative Chromogenic LAL test kit (Cambrex; E. Rutherford, N.J.), following the manufacturer's instructions for the microplate method.

[0454] TLR5 bioactivity assay: HEK293 cells constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96-well microplates (50,000 cells/well), and the following test proteins were added and incubated overnight: STF2.HA1-1His(PR8) (SEQ ID NO: 89); STF2.HA1-2(PR8) (SEQ ID NO: 90); STF2.HA1-2(PR8)mut (SEQ ID NO: 91); STF2.HA1-3(PR8) (SEQ ID NO: 92); HA1-2His(PR8) (SEQ ID NO: 93); STF2.HA1-2 (Mal) (SEQ ID NO: 211) supernatant; STF2. HA1-2 (Mal) (SEQ ID NO: 211) refolded; STF2.HA1-2 (SH) (SEQ ID NO: 211) supernatant; STF2.HA1-2 (SH) (SEQ ID NO: 211) refolded; STF2.HA1-1 (Mal) (SEQ ID NO: 209) supernatant; and STF2.HA1-1 (Mal) (SEQ ID NO: 209). The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20° C. After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce, Rockland, Ill.; #M801E and #M802B) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer (FAR-Cyte, GE/Amersham; Piscataway, N.J.).

[0455] Protein antigenicity ELISA: To determine whether the recombinant fusion proteins displayed correctly folded epitopes of HA, the antigenicity of individual HA-fusion proteins was evaluated by ELISA. 96-well ELISA plates were coated overnight at 4° C. with serial dilutions in PBS (100 µl/well) of each target protein starting at 5 µg/ml. Plates were blocked with 200 µl/well of Assay Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature, then washed three times in PBS-T. A fixed dose of primary antibody was then added to each well.

[0456] To assay HA reactivity, 100 µl/well of a 1:10,000 dilution of non-immune or PR/8/34 convalescent immune serum in ADB was added. PR/8/34 immune serum was generated in BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) that received an experimentally determined sublethal challenge dose of 8×10^1 egg infectious dosages (EID) of PR/8/34 influenza virus. Animals were then allowed to convalesce for >21 days post-infection at which time immune serum was isolated and clarified. For ELISA of flagellin or the 6× histidine tag, monoclonal antibody against 6×His (Invitrogen; Carlsbad, Calif.), or flagellin (Inotek; Beverly, Mass.) was added at 1 µg/ml in ADB (100 µl/well) and the plates were incubated for 1 hr at room temperature or overnight at 4° C. The plates were then washed three times with PBS-T. HRP-labeled goat anti-mouse IgG anti-

bodies (Jackson Immunochemical; West Grove, Pa.) diluted in ADB were added (100 µl/well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce; Rockford, Ill.) and monitoring color development, A_{450} was measured on a microplate spectrophotometer (FARCyte, GE/Amersham; Piscataway, N.J.).

Results and Discussion

[0457] Protein yield and purity: Results for the purification of recombinant HA and STF2.HA fusion proteins produced in *E. coli* are shown in Table 6. All four proteins were produced in high yield, with estimated purity exceeding 90% and endotoxin well below the standard acceptable level of 0.1 EU/µg. The three STF2 fusion proteins also had very high in vitro TLR5 bioactivity while the HA1-2His₆ (SEQ ID NO: 93) protein, as expected, had no TLR5 activity.

TABLE 6

protein	SEQ ID NO:	yield (mg)	purity est. (%)	endotoxin (EU/µg)	TLR5 activity (EC ₅₀ , ng/ml)
STF2.HA1-1His(PR8)	89	6.4	>90	0.04	0.2
STF2.HA1-2(PR8)	90	6.0	>98	0.02	0.15
STF2.HA1-2(PR8)mut	91	12	>95	0.01	1.2
STF2.HA1-3(PR8)	92	2.1	>90	0.016	1.2
HA1-2His(PR8)	93	10	>95	0.03	nd

[0458] Antigenicity of Influenza A HA proteins produced in *E. coli*: All four recombinant proteins were analyzed by western blotting with antibody against STF2 (flagellin) (Inotek; Beverly, Mass.) and immune antisera collected from PR/8/34 convalescent mice. The three STF2 fusion proteins appeared to react comparably with anti-flagellin antibody. STF2.HA1-1His(PR8) (SEQ ID NO: 89), STF2.HA1-2(PR8) (SEQ ID NO: 90) and HA1-2His(PR8) (SEQ ID NO: 93) also reacted with anti-PR/8/34 convalescent serum. This reactivity was only seen in non-reduced protein samples; the addition of DTT to the sample buffer greatly diminished the signal. This finding suggests that the majority of the anti-HA antibodies in the convalescent serum are conformationally dependent and that proper disulfide bonding in HA is necessary for reactivity by western blot. In contrast to these results, STF2.HA1-3(PR8) (SEQ ID NO: 92) did not react efficiently with the convalescent serum and what little signal was produced was not affected by reductant. This finding suggested that the STF2.HA1-3 protein may not be capable of folding properly and/or presenting native HA epitopes.

Capture ELISA for Influenza B Globular Head Proteins

[0459] The antigenicity of individual HA-fusion proteins was evaluated using capture ELISA. An anti-flagellin monoclonal antibody was used on the solid phase to capture these proteins. Ferret anti-sera raised on natural B/Malaysia/2506/2004 infection was used to detect the captured HA-fusion proteins. The results of this assay will determine whether the recombinant fusion proteins displayed correctly-folded epitopes of HA. Due to the considerable sequence homology between the different Influenza B HA globular head molecules, 84.6%, these antisera would be expected to react with both the Malaysian and the Shanghai globular head constructs.

Preparation of Samples

[0460] *E. coli* cells expressing the STF2.HA1-2(Malaysia (SEQ ID NO: 210)), STF2.HA1-2 (Shanghai (SEQ ID NO: 211)), and STF2.HA1-1 (Malaysia (SEQ ID NO: 209)) recombinant proteins were pelleted by centrifugation and resuspended in Tris buffer (50 mM Tris, 200 mM NaCl, pH 8). Cells were lysed by sonication. The cell lysate was centrifuged to separate the soluble proteins from the insoluble proteins. The insoluble protein pellet was resuspended in Tris buffer with 6M urea and the proteins were subjected to rapid refolding by quickly diluting the solution 1:10 fold in Tris buffer. The protein concentrations of both the soluble proteins and the pelleted samples containing refolded proteins were estimated by UV₂₈₀ (Spectrophotometer DU 800). Supernatants containing STF2.HA1-1(Mal) (SEQ ID NO: 209); STF2.HA1-2(Mal) (SEQ ID NO: 210), STF2.HA1-2(SH, Shanghai) (SEQ ID NO: 211) protein and solubilized pellets containing refolded STF2.HA1-1(Mal), STF2.HA1-2(Mal), STF2.HA1-2(SH) were evaluated by ELISA for reactivity with ferret antisera raised on natural infection against B/Malaysia/2506/2004. *E. coli* supernatant samples and solubilized, refolded samples containing STF2.4×M2e (SEQ ID NO: 94) protein were used as negative controls.

ELISA Method

[0461] The ELISA plate (Maxisorp, Nunc, Denmark) was coated with the flagellin-specific monoclonal antibody 6H11 (Inotek, Beverly, Mass.) at 0.5 µg/mL and incubated overnight at 2-5° C. The antibody coating solution was aspirated and the wells were blocked with 300 µL/well Super Block+ Tween-20 for 2 hours at 25° C. The plate was washed once with 1×PBS and blot-dried. Three-fold serial dilutions of the different protein solutions were performed in a dilution plate starting with a concentration of 5 µg/mL. 100 µL was transferred from the dilution plate to the ELISA plate.

[0462] The plate was incubated for 1 hr at 25° C. Unbound protein was removed by washing the plate 3 times with PBS+0.05% Tween-20. Ferret antiserum raised against B/Malaysia/2506/2004 (CDC, AL, Georgia) was diluted to 1:100 and 100 µL was added to each well. The plate was then incubated for 1 hr at 25° C. After this incubation step, the plates were washed 6 times with PBS containing 0.05% Tween-20. Goat anti-ferret IgG conjugated to horseradish peroxidase (HRP, Bethyl Labs Inc., IL) was diluted 1:10,000 and 100 µL was added to each well. The plate was incubated for 30 min. After this incubation step, the plate was washed 6 times with PBS+0.05% Tween-20. 100 µL of TMB Ultra containing the HRP substrate 3,3',5,5'-tetramethylbenzidine (Pierce, Rockford, Ill.) was added to each well. After the addition of this substrate, the color development was monitored and the reaction was stopped with the addition of 100 µL/well 1M H₂SO₄. A₄₅₀ was measured on a microplate reader (SpectraMax 190, Molecular devices, Sunnyvale, Calif.).

Antigenicity of Influenza B HA Proteins Produced in *E. coli*

[0463] The ELISA data (FIG. 3) indicated that the antisera recognized epitopes that were sensitive to reductant. Treatment of proteins with reducing agents, which alter the conformation of the proteins by the disruption of disulphide bonds and, diminished reactivity of the antisera with the proteins. The reactivity of STF2.HA1-2(Mal) total lysate

non-reduced, STF2.HA1-2(SH) total lysate non-reduced and STF2.HA1-1(Mal) total lysate non-reduced when maintained in the properly folded configuration was very good and indicates that the conformation of the individual HA-fusion proteins was comparable to that of the globular head in the native HA protein. While the antiserum contains conformationally sensitive antibodies and the reactivity depends on correct disulfide bonding, it also contains antibodies that are non-sensitive to disulfide bonding as demonstrated by the residual activity which was observed in both reduced and non-reduced conditions.

TLR5 Bioactivity of STF2.HA (Influenza B) Proteins

[0464] The recombinant fusion proteins showed potency which matched the results from the Capture ELISA indicating that refolded samples may be more active than the lysate in its native form (FIGS. 4 and 5). The samples in the native untreated form showed activity which were 2-fold less than the refolded samples. These proteins were misfolded in the native form and therefore require a refolding step to reinstall the TLR-5 bioactivity. This activity was consistent with HA activity in the Capture ELISA.

Example 5

Immunogenicity of Recombinant Flagellin-Hemagglutinin Fusion Proteins Representing Viral Strain A/Puerto Rico/8/34 in *E. coli*

Materials and Methods

[0465] Animal studies: Female BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) were used at the age of 6-8 weeks. Mice were divided into groups of 10 and received inguinal subcutaneous (s.c) immunizations on days 0 and 14 as follows:

- 1) PBS (phosphate buffered saline).
- 2) 3 µg of STF2.4×M2e (SEQ ID NO: 94) in saline buffer (10 mM Histidine, 10 mM Tris, 75 mM NaCl, 5% (vol/vol) sucrose, 0.02% (w/v) Polysorbate-80, 0.1 mM EDTA, 0.5% (v/v) ethanol, pH 7.2)
- 3) 30 µg of STF2.HA1-2(PR8) (SEQ ID NO: 90) in PBS
- 4) 3.0 µg of STF2.HA1-2(PR8) (SEQ ID NO: 90) in PBS
- 5) 0.3 µg of STF2.HA1-2(PR8) (SEQ ID NO: 90) in PBS
- 6) 0.03 µg of STF2.HA1-2(PR8) (SEQ ID NO: 90) in PBS

[0466] An additional group of five mice received an experimentally determined sublethal challenge with 8×10⁴ egg infectious dosages (EID) PR/8/34 and were allowed to convalesce for >21 days. These animals were then used as immune convalescent positive controls during the challenge studies (see below). Mice were bled on days 10 (primary) and 21 (boost), and sera were clarified by clotting and centrifugation and stored at -20° C.

[0467] Serum antibody determination: HA-specific IgG levels were determined by ELISA. 96-well ELISA plates (Costar (Cat #9018) Corning, N.Y.) were coated overnight at 4° C. with 100 µL/well HA0sHis protein (produced in *Drosophila*) (SEQ ID NO: 176) in PBS (5 µg/ml). Plates were blocked with 200 µL/well of Assay Diluent Buffer (ADB; BD Pharmingen, (Cat#: 555213)(San Diego, Calif.)

for one hour at room temperature. The plates were washed three times in PBS+0.05% (v/v) Tween 20 (PBS-T). Dilutions of the sera in ADB were added (100 μ l/well) and the plates were incubated overnight at 4° C. The plates were washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical, West Grove, Pa. (Cat#: 115-035-146)) diluted in ADB were added (100 μ l/well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce (Cat 34028), Rockford, Ill.)) and monitoring color development, A₄₅₀ was measured on a Tecan Farcyte (Durham, N.C.) microplate spectrophotometer.

[0468] MDCK whole cell ELISA: MDCK cells (ATCC (Cat# CCL-34) Manassas, Va.) were grown in 96-well culture plates (BD (Cat 353075), Corning, N.Y.) in DMEM complete medium containing 10% FCS at 37° C. for one to two days or until cells were near confluence. Wells were then incubated with 1 $\times$ 10⁶ EID of PR/8/34 virus (50 μ l) in DMEM medium without FCS or medium alone (for uninfected controls).

[0469] Following a 60 minute incubation at 37° C., 200 μ l of complete medium was added to each well and plates were incubated overnight at 37° C. The next day plates were washed with PBS and fixed with 10% formalin at room temperature for 10 minutes. Wells were washed three times with PBS/0.1% BSA and blocked with 200 μ l/well ADB (BD Pharmingen, Cat# 555213), San Diego, Calif.) for one hour at RT or overnight at 4° C. Serial dilutions of test sera were added to the wells and incubated for one to two hours at room temperature. Wells were washed and incubated with HRP-conjugated goat anti-mouse IgG (Jackson Immunochemical, Cat 115-035-146 (West Grove, Pa.) for 30 minutes at room temperature, followed by TMB Ultra substrate (Pierce (Cat# 34028), Rockford, Ill.) for two minutes at room temperature. The reaction was stopped with the addition of 25 μ l of 1N H₂SO₄ and the OD₄₅₀ was read using a microplate spectrophotometer (FARCyte, Amersham, Durham, N.C.).

Results and Discussion

[0470] Induction of HA-specific IgG response following immunization with STF2.HA1-2(PR8) (SEQ ID NO: 90): The immunogenicity of STF2.HA1-2 was examined by immunizing BALB/c mice (10/group) subcutaneously on day 0 and 14 with a dose range of STF2.HA1-2 (30, 3.0, 0.3, or 0.03 μ g). Control groups of mice were immunized with PBS (negative control), 3 μ g of STF2.4 $\times$ M2e (SEQ ID NO: 94) (negative control for HA immunogenicity, positive control for lethal challenge efficacy study), or a sub-lethal challenge with 8 $\times$ 10⁴ EID of the influenza isolate PR/8/34 to generate immune convalescent animals. HA-specific IgG responses were examined 7 days post boost (Day 21) by ELISA. The results demonstrate that immunization with 30, 3 or 0.3 μ g of STF2.HA1-2(PR8) induced consistent and significant HA0sHis-specific (SEQ ID NO: 176) IgG responses in a dose-dependent manner (FIG. 6).

[0471] Sera from BALB/c mice immunized with STF2.HA1-2 (SEQ ID NO: 90) react with MDCK cells infected with influenza virus: The direct ELISA results above demonstrated that the immune sera from STF2.HA1-2-immunized animals recognize recombinant *Drosophila* expressed HA0sHis (SEQ ID NO: 176) corresponding to the

PR/8/34 HA sequence. In order to demonstrate that the anti-HA antibodies recognize native viral HA, the same sera were examined for the ability to react with MDCK cells infected with PR/8/34. The results shown in FIG. 7 demonstrate that sera from mice immunized with 30, 3.0, or 0.30 μ g of STF2.HA1-2 bound specifically to PR/8/34-infected MDCK cells, indicating that the anti-HA antibodies elicited by immunization with STF2.HA1-2 recognize HA in its native conformation.

[0472] Collectively these results demonstrate that mice immunized with STF2.HA1-2(PR8) (SEQ ID NO: 90) in PBS, without conventional adjuvant or carrier, mounted vigorous anti-HA responses that recognized recombinant HA expressed in *Drosophila* and native viral HA expressed on the surface of PR/8/34 infected MDCK cells in vitro.

Example 6

Efficacy of Recombinant Flagellin-Hemagglutinin Fusion Proteins Representing Viral Strain A/Puerto Rico/8/34

Materials and Methods

[0473] Influenza virus challenge of mice. To assess efficacy, mice immunized as described in Example 5 were challenged on day 28 by intranasal administration of an LD₅₀ (dose lethal to 90% of mice) (8 $\times$ 10³ EID) of influenza A isolate PR/8/34. Animals were monitored daily for 21 days following the challenge for survival, weight loss and clinical presentation. The % weight loss was calculated based on the mean of ((Daily weight (g)/Initial weight (g) day 28) $\times$ 100) of each individual animal per group. Clinical scores were assigned as follows: 4 pts=healthy, 3 pts=reduced grooming, 2 pts=reduced physical activity and 1 pt=moribund. (Experimental results for clinical scores and weight loss reflect the results based on surviving animals on the day evaluated).

[0474] Virus Titration and Determination of TCID₅₀.

[0475] Cell preparation: MDCK cells (ATCC (Cat# CCL-34), Manassas, Va.) were cultured in 100 $\times$ 20 mm culture plates (BD (Cat# 353003) Corning, N.Y.) to 90-95% confluency and the monolayer was dislodged by incubation with trypsin-EDTA at 37° C./5% CO₂ for 20 minutes. The trypsin was inactivated by the addition of DMEM cell culture medium supplemented with 10% FBS, and the cell monolayer was scraped with a sterile spatula to complete detachment of cells. The cell suspension was harvested and washed twice with DMEM+10% FBS. Cells were resuspended in DMEM+10% FBS and counted. The cell concentration was adjusted to 4 $\times$ 10⁵ cells/ml and 100 μ l was added to each well of a 96-well tissue culture plate (BD (Cat# 353075), Corning N.Y.). Plates were incubated at 37° C./5% CO₂ until confluence reached 90-95%.

[0476] Viral titration: Influenza virus (strain A/Puerto Rico/8/34 [PR/8]) was diluted to 1 $\times$ 10⁸ EID in phenol red-free DMEM+0.1% BSA (fraction V (Rockland (Cat#BSA-50) Gilbertsville, Pa.), and serial 5-fold dilutions were prepared in 96-well plates using the same medium. Monolayers of MDCK cells in 96-well plates, prepared as described in herein, were washed by aspirating the culture medium, replacing with 200 μ l/well of 1 $\times$ PBS, and aspirating the PBS. Serial 5-fold dilutions of influenza virus prepared above were added to the washed monolayers in a

volume of 100 μ l/well. One row of wells was treated with medium only as a control. The cells were incubated for 2 hours at 37° C./5% CO₂ to allow viral attachment and entry. Wells were washed by aspiration, rinsing with 200 μ l/well PBS, and aspiration of the PBS. All wells received 100 μ l/well phenol red-free DMEM+0.1% BSA and the plates are incubated for 48 hours at 37° C./5% CO₂.

[0477] Determination of cell viability: Following incubation with virus as described above, the medium was aspirated from the wells and replaced with fresh medium containing 40 μ g/ml neutral red (Sigma Aldrich (Cat# N2889) St. Louis, Mo.). To determine maximum lysis, 2 μ l lysis solution (9% Triton X-100 in water, weight/vol) was added to triplicate wells that were incubated with medium only. Following a one hour incubation, the cells were fixed by the addition of 100 μ l/well 1% formaldehyde/1% CaCl₂ for 5 minutes at room temperature; this fix step was performed twice in succession. The fix solution was aspirated and the neutral red was released by the addition of 100 μ l/well of extraction medium (50% ethanol/1% acetic, acid).

[0478] The plate was incubated at room temperature for 20 minutes, with shaking for the final 2 minutes. The amount of dye released was determined by measuring absorbance at a wavelength of 540 nm using a microplate spectrophotometer. Cell death (and hence, viral infectivity) was measured as a decrease in the amount of dye released as compared to media control. At the end of the incubation, the neutral red assay was performed as described herein. The percentage lysis of each serum dilution was calculated as:

$$\% \text{ reduction} = 100 \times ((\text{sample} - \text{virus}) / (\text{med} - \text{virus}))$$

[0479] where sample, max, and med refer to the absorbance values in wells representing experimental samples, virus only, and medium only, respectively. The neutralizing titer of each sample was defined as the dilution of serum which resulted in a 50% reduction in viral infectivity.

[0480] Influenza A/PR/8/34 Neutralization Assay This assay was adapted with modifications from WHO Manual on Animal Influenza Diagnosis and Surveillance, p. 86-88 (WHO/CDS/CSR/NCS2002.5). The Neutral Red Assay is adapted from a public protocol of the Cell Lab at Gettysburg College (<http://www3.gettysburg.edu/~sorense/Cellab04/neutralred.htm>)

[0481] Specifically, MDCK (ATCC, Cat#(CCL 34), Manassas, Va.) cells were plated in 96-well tissue culture plates (BD (Cat 353075), Corning, N.Y.) as described above. Test reagents (experimental and control Sera) were heat-inactivated by incubating for 30 minutes in a water bath heated to 56° C. Sera were serially titrated in 96-well plates, in 3-fold dilutions in phenol red-free DMEM+0.1% BSA. An equal volume of PR/8/34 virus diluted to 5 $\times$ 10⁶/ml EID in the same medium was added to each serum dilution to achieve a final viral concentration of 2.5 $\times$ 10⁶ EID/ml (The pre-determined TCID₅₀ for our current stock of virus). Wells containing medium only and virus only were included as negative and positive controls, respectively. The plates were incubated for 30 minutes at 37° C./5% CO₂.

[0482] Cell monolayers prepared as described above (Cell preparation) were washed once with 200 μ l/well 1 $\times$ PBS, then 100 μ l/well of serum:virus mixtures and control reagents prepared as described above were added and incubated for 2 hours at 37° C./5% CO₂. Following incubation,

cell monolayers were washed once with PBS, and 100 μ l/well phenol red-free DMEM+0.1% BSA was added to each well and incubated at 37° C./5% CO₂ for 48 hours. At the end of the incubation, the neutral red assay was performed as described above (Determination of cell viability). The percentage lysis of each serum dilution was calculated as described above. The neutralizing titer of each sample was defined as the dilution of serum which results in a 50% reduction in viral infectivity

Results and Discussion

[0483] Immunization with STF2.HA1-2(PR8) (SEQ ID NO: 90) provides protection from a lethal challenge with influenza A: The results from Example 5 demonstrated that immunization of mice with STF2.HA1-2(PR8) (SEQ ID NO: 90) generated an antibody response that recognized native HA. In order to evaluate efficacy, the same mice were challenged on day 28 with an LD₉₀ (8 $\times$ 10³ EID) of PR/8/34 virus administered intra-nasally. Mice were monitored daily for 19 days following the challenge for survival, weight loss and clinical presentation. As shown in FIGS. 8A, 8B, and 8C, PBS-immunized mice showed signs of infection (weight loss and lower clinical scores) as early as three days post-challenge and all mice died by day 21 post-challenge. In contrast, mice immunized with 30, 3.0 or 0.3 μ g of STF2.HA1-2(PR8) (SEQ ID NO: 90) demonstrated markedly enhanced protection. These animals demonstrated little to no weight loss, significantly higher clinical scores and 100% survival that was similar to immune convalescent control animals over the 21 day period. These results demonstrate that *E. coli*-expressed STF2.HA1-2(PR8) (SEQ ID NO: 90) induces HA-specific immune responses that successfully protect BALB/c mice from a lethal challenge with virulent influenza A virus in vivo.

[0484] In order to evaluate the effectiveness of the antibody response in vitro a cell-based virus neutralization assay was developed to test the ability of immune sera to neutralize viral infectivity. In this study the viral inhibitory activity of serum from animals immunized with STF2.HA1-2(PR8) (SEQ ID NO: 90) and STF2.HA1-1(PR8) (SEQ ID NO: 151) was examined. Serial dilutions of non-immune and immune sera were pre-incubated with PR/8 and incubated with MDCK cells. Wells were washed to remove free virus and plates were incubated prior to staining with neutral red to detect live cells. The results indicate that immune sera from animals immunized with STF2.HA1-2(PR8) (SEQ ID NO: 90) demonstrated a tissue culture inhibition 50% dose (TCID) of >1:40 (FIG. 9).

Example 7

Immunogenicity of Recombinant
Flagellin-Hemagglutinin Fusion Proteins
Representing Viral Strain A/Viet Nam/1203/2004

Materials and Methods

[0485] Animal studies: Female BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) were used at the age of 6-8 weeks. Mice were divided into groups of 10 and received inguinal s.c immunizations on days 0 and 14 as follows:

[0486] 1) PBS (phosphate buffered saline).

[0487] 2) 3.0 μ g of STF2.HA1-2(PR8) (SEQ ID NO: 90) in PBS

[0488] 3) 3.0 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) in PBS

[0489] 4) 0.3 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) in PBS

[0490] Mice were bled on day 21 and sera were clarified by clotting and centrifugation and stored at -20° C.

[0491] Serum antibody determination: HA and STF2-specific IgG levels were determined by ELISA. 96-well ELISA plates (Costar (Cat#9018), Corning, N.Y.) were coated overnight at 4° C. with 100 µl/well of HA protein purified from Viet Nam/1203/2004 (BEI Resources (Cat #NR-660), Manassas, Va.) or recombinant flagellin (STF2) (SEQ ID NO: 96) in PBS (5 µg/ml). Plates were blocked with 200 µl/well of Assay Diluent Buffer (ADB; BD Pharmingen (Cat#555213) San Diego, Calif.) for one hour at room temperature. The plates were washed three times in PBS buffer containing 0.05% (v/v) Tween 20 (PBS-T). Dilutions of the sera in ADB were added (100 µl/well) and the plates were incubated overnight at 4° C. The plates were washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical (Cat#115-035-146), West Grove, Pa.) diluted in ADB were added (100 µl/well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce (Cat#34028), Rockford, Ill.)) and monitoring color development, A₄₅₀ was measured on a Tecan Farcyte (Durham, N.C.) microplate spectrophotometer.

Results and Discussion

[0492] The immunogenicity of STF2.HA1-2(VN) (SEQ ID NO: 95) was examined in BALB/c mice. Animals were immunized s.c. on days 0 and 14 with 3 µg STF2.HA1-2(PR8) (SEQ ID NO: 90), or 3 or 0.3 µg of STF2.HA1-2(VN) (SEQ ID NO: 95). On day 21 animals were bled and serum IgG responses to purified HA from A/Viet Nam/1203/2004 (obtained from BEI Resources Cat# NR-660) and recombinant flagellin (STF2; (SEQ ID NO: 96)) were examined by ELISA (FIGS. 10A and 10B). The results demonstrate that serum from mice immunized with STF2.HA1-2(VN) (SEQ ID NO: 95) exhibit antigen-specific reactivity with purified H5 protein from influenza A/Viet Nam/1203/2004.

Example 8

Cloning, Expression, and Biochemical Characterization of Recombinant Flagellin-Hemagglutinin Fusion Proteins Produced in Baculovirus

[0493] Cloning: Expression of HA and STF2.HA fusion proteins was carried out using the Bac-to-Bac Baculovirus Expression System (Invitrogen, Carlsbad, Calif.) according to the manufacturer's guidelines (FIG. 11). cDNAs encoding HA and STF2.HA fusion proteins were cloned into the pFastBac donor plasmid which was then used to transform DH10Bac cells containing the bacmid and helper DNA. Recombinant bacmid clones generated by homologous recombination in DH10Bac were then identified by blue-white screening on X-gal plates. The details of the cloning procedure are given in the following paragraphs.

[0494] Generation of vector cassettes: The pFastBac™1 vector is compatible with the Bac-to-Bac® Baculovirus Expression System (Invitrogen, Carlsbad, Calif.). This vector has a strong AcMNPV polyhedrin (PH) promoter for high level protein expression and a large multiple cloning site for simplified cloning. pFastBac™1 is a non-fusion vector (i.e. no fusion tags are present in the vector).

[0495] To ensure proper expression of recombinant protein, the insert must contain an ATG start codon for initiation of translation and a stop codon for termination of translation. In order to facilitate cloning of flagellin in fusion with several truncations of the HA gene, two plasmid cassettes were engineered: pFB-STF2Blp.wt (SEQ ID NO: 97) and pFB-STF2Blp.ng (SEQ ID NO: 98). In the latter, the putative glycosylation sites in the flagellin (STF2) gene were obliterated by substituting Gln (Q) for Asn (N). Potential glycosylation sites were determined using the consensus sequence for N-glycosylation N-X-S/T: N residue followed by any residue by S or T in the third position.

[0496] In the certain locations, residues 40, 122, 215, 237, 414, 478, and 497 of the STF2 gene (SEQ ID NO: 212) the N residue was incorporated as a Q substitution. Constructs which harbor these mutations and in which the proteins are not likely to be modified by sugars are designated as non-glycosylated (ng) mutants. Similarly, glycosylation mutations were introduced in HA0s gene at positions 11, 23, 268, 286 and 480 (SEQ ID NO: 23). Both vector constructs (pFB-STF2Blp.wt and pFB-STF2Blp.ng) harbor a silent mutation at nucleotides 5'-GTGCTGAGCCTGTTACGT-3' (SEQ ID NO: 310) (nt 1501 to 1518) of STF2 (SEQ ID NO: 212) creating a unique BlnI in the plasmid cassette. Each cassette contains the honey bee melittin (HBM) sequence (SEQ ID NO: 99) fused at the amino terminus of STF2 to provide a signal for secretion. Both plasmids were codon-optimized for expression in B Baculovirus (Midland Certified Reagent Co., Inc, Midland, Tex.; DNA2.0 Inc, Menlo Park, Calif.). The synthetic genes were excised with BamHI or BglII and SphI and cloned into pFastBac™ 1 vector by compatible end ligation generating the pFB-STF2Blp.wt and pFB-STF2Blp.ng cassettes.

[0497] Flagellin-HA fusion constructs: Subunits of the HA globular head from influenza strains A/Puerto Rico/8/34, A/Viet Nam/1203/2004, A/Indonesia/2005 and A/New Caledonia/12/99 were expressed alone or genetically fused to flagellin (STF2) (SEQ ID NO: 178) and expressed in Baculovirus. This was accomplished by one of three methods.

[0498] Method #1: To generate reagents for ELISA, HA1-1 subunits of PR8, VN, IND and NC strains (Tables 7 and 8) were cloned into pFASTBac™1, generating pFB.HA1-1 that harbors a hexa-his tag. This was achieved by employing PCR with a set of primers as indicated in the Tables below and with a synthetic codon-optimized HA0s gene (HA excluding the signal sequence and extracellular domain) as DNA template (DNA2.0 Inc., Menlo Park, Calif.). The PCR fragments were digested with BglII and SphI and inserted into pFastBac™1 that has previously been treated with BglII and SphI enzymes followed by BAP treatment. In order to generate a flagellin-HA subunit fusion, the PCR product was digested with BlnI, and ligated by compatible ends to pFB-STF2Blp.wt.

[0499] Method #2: In this protocol, codon-optimized HA subunit genes representing both the wt and non-glycosylated

forms were chemically synthesized (DNA2.0 Inc., Menlo Park, Calif.). The genes were excised with BspI and SphI enzymes and the fragment gel purified and ligated to pFB-STF2Blp.wt or the non-glycosylated version, pFB-STF2Blp.ng previously digested with BspI and SphI and BAP treated. In each case, the ligation mix was used to transform TOP10 cells and transformants were screened by PCR and DNA sequencing to confirm the presence and correct orientation of the inserts. The constructs were used to transform MAX Efficiency® DH10Bac™ competent *E. coli* (Invitrogen, Carlsbad, Calif.) to generate a recombinant bacmid. The colonies of bacteria were screened for positive bacmids by blue/white selection on plates containing 50 µg/mL kanamycin, 7 µg/mL gentamycin, 10 µg/mL tetracycline, 40 µg/mL IPTG and Blu-Gal (40 µg/mL). Recombinant bacmid DNA was prepared and used to transfect the insect cell line of choice (Sf9 or Sf21 cells) to generate a recombinant Baculovirus. The baculoviral stock was then amplified and titered and used to infect High Five insect cells to express the recombinant protein.

TABLE 7

PR8 HA constructs for expression in Baculovirus					
SEQ ID NO:	Construct (His tag)	Method	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:
100	STF2.HA0s	#1	101	102	103
104	STF2.HA1-1	#1	105	106	103
107	STF2.HA1-2	#1	108	109	103
110	STF2.HA1-2mut	#2	N/A	N/A	111
112	STF2.HA1-3	#1	113	114	103
115	STF2.HA1-3mut	#2	N/A	N/A	116
117	ngSTF2.HA0s	#2	N/A	N/A	118
119	ngSTF2.HA1-1	#2	N/A	N/A	120
121	ngSTF2.HA1-2	#2	N/A	N/A	122
123	ngSTF2.HA1-1ng	#2	N/A	N/A	124
125	ngSTF2.HA1-3	#2	N/A	N/A	126
127	ngSTF2.HA1-3mut	#2	N/A	N/A	128
129	wtSTF2.HA1-1ng	#2	N/A	N/A	130
131	ngSTF2.HA1-1wt	#2	N/A	N/A	132
133	HA1-1	#1	134	135	136
137	HA1-1 (no tag)	#1	134	138	136

[0500]

TABLE 8

NC, VN and NC HA constructs for expression in Baculovirus					
SEQ ID NO:	Construct (his tag)	Method	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:
139	HA1-1(NC)	#1	140	141	142
143	HA1-1(VN)	#1	144	145	146
147	HA1-1(IND)	#1	148	149	150

[0501] Protein Expression: *Spodoptera frugiperda* (Sf9) insect cells were cultured in Grace's Insect Cell Medium (Invitrogen; Carlsbad, Calif.). Recombinant bacmids were purified from DH10Bac clones and transfected into Sf9 cells using Cellfectin (Invitrogen; Carlsbad, Calif.) to generate passage one (P1) Baculovirus stocks which were then titered by traditional plaque assay according to the manufacturer's guidelines. Subsequent infections of Sf9 cells were per-

formed using P1 viral stocks at a multiplicity of infection (MOI) of 0.1 to produce larger volume P2 viral stocks with increased viral titers. Cultures expressing the following proteins were scaled up for protein purification: ngSTF2.HA(PR8)1-2mutHis (SEQ ID NO: 156); ngSTF2.HA(PR8)1-1His (SEQ ID NO: 152); STF2.HA(PR8)1-1His (SEQ ID NO: 151); STF2.HA(PR8)1-2His (SEQ ID NO: 153); STF2.HA1-2mutHis (SEQ ID NO: 155); HA(PR8)1-1His (SEQ ID NO: 179); HA(VN)1-1His (SEQ ID NO: 180); HA(IND)1-1His (SEQ ID NO: 181); HA(NC)1-1His (SEQ ID NO: 182). Protein expression was performed by infecting 2x1 L flasks of High-5 insect cells with P2 virus at an MOI of 2 and culturing for 24 hrs at 28° C. with slow shaking. Conditioned medium was harvested by centrifugation and clarified by passing through a 0.22 µm filter and stored at 4° C.

[0502] Protein Purification: NiSO₄ was added to each clarified supernatant to a final concentration of 0.5 mM and the pH adjusted to 8.0. The His₆-tagged protein was then captured on a chelating sepharose column (GE/Amersham Biosciences; Piscataway, N.J.) charged with NiSO₄ and equilibrated in Buffer A (20 mM Tris, pH 8.0, 0.5 M NaCl). After washing with Buffer A the bound protein was eluted with Buffer B (Buffer A+0.5 M imidazole). Peak fractions were pooled, dialyzed overnight into Buffer A and applied to a nickel NTA column (Sigma-Aldrich; St. Louis, Mo.). The column was eluted with a 5-column volume gradient from 100% Buffer A to 100% Buffer B. Peak fractions were pooled and dialyzed into 1xTBS, pH 8.0.

[0503] SDS-PAGE and western blot analysis: Protein identity was determined, and purity estimated, by SDS-PAGE. An aliquot of 5 µg of each sample was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel (LifeGels; French's Forrest, New South Wales, AUS) and electrophoresed. The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands. For western blot, 0.5 µg/lane total protein was electrophoresed as described above and the gels were then electro-transferred to a PVDF membrane and blocked with 5% (w/v) dry milk before probing with anti-flagellin antibody (Inotek; Beverly, Mass.), anti-His₆ antibody (Invitrogen; Carlsbad, Calif.), or influenza A PR/8/34 convalescent immune serum (described below under Protein Antigenicity ELISA). After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockford, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega; Madison, Wis.).

[0504] Protein assay: Total protein concentration was determined using the Micro BCA (bicinchoninic acid) Assay (Pierce; Rockford, Ill.) in the microplate format, using bovine serum albumin as a standard, according to the manufacturer's instructions.

[0505] Endotoxin assay: Endotoxin levels were determined using the QCL-1000 Quantitative Chromogenic LAL test kit (Cambrex; E. Rutherford, N.J.), following the manufacturer's instructions for the microplate method.

[0506] TLR5 bioactivity assay: HEK293 cells constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96-well microplates (50,000 cells/well), and the

following test proteins were added and incubated overnight: ngSTF2.HA(PR8)1-2mutHis (SEQ ID NO: 156); ngSTF2.HA(PR8)1-1His (SEQ ID NO: 152); STF2.HA(PR8)1-1His (SEQ ID NO: 151); STF2.HA(PR8)1-2His (SEQ ID NO: 153); STF2.HA1-2(PR8)mutHis (SEQ ID NO: 155). The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20°C . After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce; Rockford, Ill., #M801E and #M802B) following the manufacturer's instructions.

[0507] Protein antigenicity ELISA: The following constructs purified from Baculovirus culture were evaluated by ELISA to determine whether the proteins displayed correctly folded epitopes of HA: ngSTF2.HA(PR8)1-2mutHis (SEQ ID NO: 156); ngSTF2.HA(PR8)1-1His (SEQ ID NO: 152); STF2.HA(PR8)1-1His (SEQ ID NO: 151); STF2.HA(PR8)1-2His (SEQ ID NO: 153); STF2.HA1-2mutHis (SEQ ID NO: 155); and HA(PR8)1-1His (SEQ ID NO: 179). 96-well ELISA plates were coated overnight at 4°C with serial dilutions in PBS (100 μl /well) of each target protein starting at 5 $\mu\text{g}/\text{ml}$. Plates were blocked with 200 μl /well of Assay Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature, then washed three times in PBS-T. A fixed dose of primary antibody was then added to each well. To assay HA reactivity, 100 μl /well of a 1:10,000 dilution of non-immune or PR/8/34 convalescent immune serum in ADB was added. PR/8/34 immune serum was generated in BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) that received an experimentally determined sublethal challenge dose of 8×10^1 egg infectious dosages (EID) of PR/8/34 influenza virus. Animals were then allowed to convalesce for >21 days post-infection at which time immune serum was isolated and clarified.

[0508] For ELISA of flagellin or the 6 $\times$ histidine tag, monoclonal antibody against 6 $\times$ His (Invitrogen; Carlsbad, Calif.), or flagellin (Inotek; Beverly, Mass.) was added at 1 $\mu\text{g}/\text{ml}$ in ADB (100 μl /well) and the plates were incubated for 1 hr at room temperature or overnight at 4°C . The plates were then washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical; West Grove, Pa.) diluted in ADB were added (100 μl /well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce; Rockford, Ill.) and monitoring color development, A_{450} was measured on a microplate spectrophotometer (FARCyte, GE/Amersham; Piscataway, N.J.).

Results and Discussion

[0509] Expression and purification of recombinant HA proteins: The results of Baculovirus expression of recombinant STF2.HA fusion proteins are summarized in Table 9. All proteins were expressed in moderate to high yield. Purity after metal chelate chromatography was generally good and endotoxin levels were far below the acceptable limit of 0.1 EU/ μg . All STF2 fusion proteins demonstrated potent in vitro TLR5 activity.

TABLE 9

Protein	SEQ ID NO:	Yield (mg)	Purity est. (%)	Endotoxin (EU/ μg)	TLR activity (EC_{50} , ng/ml)
STF2.HA1-1His(PR8)	151	12	>80	0.01	3
ngSTF2.HA1-1His(PR8)	152	24	>90	<0.01	30
STF2.HA1-2His(PR8)	153	13.2	>90	<0.01	9
STF2.HA1-2mutHis(PR8)	155	25	>95	<0.01	900
ngSTF2.HA1-2mutHis(PR8)	156	6	>90	0.01	9

[0510] STF2.HA proteins for Baculovirus expression were engineered with and without N-linked glycosylation sites in order to determine the effect of this posttranslational modification on protein expression, folding, and biological activity. ngSTF2.HA1-2mutHis(PR8) (SEQ ID NO: 156), which is non-glycosylated, appears to be expressed at a lower level in Baculovirus supernatants than the corresponding glycosylated protein, suggesting that glycosylation may influence the folding or secretion of this protein. However, the glycosylated and non-glycosylated versions of this protein appear to react equally with convalescent antiserum from influenza PR8-infected mice when analyzed by western blot. Furthermore, ngSTF2.HA1-1His(PR8) (SEQ ID NO: 152) expressed at higher levels than its glycosylated counterpart, indicating that any effect of glycosylation on expression of these proteins may not be easily generalized.

[0511] In addition to the proteins expressed and purified at large scale, small-scale Baculovirus P1 supernatants of STF2.HA1-3His(PR8) (SEQ ID NO: 157) and STF2.HA1-3mutHis(PR8) (SEQ ID NO: 158) were analyzed for HA antibody reactivity. As seen with the STF2.HA1-3(PR8) protein (SEQ ID NO: 92) expressed in *E. coli* (See Example #4), these proteins showed very poor reactivity with PR8 convalescent mouse serum on western blots of conditioned medium. This result further confirms that STF2.HA1-3 proteins, whether expressed in a prokaryotic or eukaryotic host, are unable to fold properly and display native epitopes.

[0512] Antigenicity: Several STF2.HA proteins produced in Baculovirus and *E. coli* were analyzed by ELISA. All STF2 fusion proteins reacted equally well with anti-flagellin antibody (Inotek; Beverly, Mass.), demonstrating equal protein loading in the assay. Proper folding and display of antibody epitopes in STF2.HA fusion proteins was then examined by measuring reactivity of each protein with antisera from mice which had recovered from PR/8/34 influenza infection. Reactivity with PR/8/34 convalescent mouse serum was equivalent for Baculovirus-expressed STF2.HA1-1His(PR8) (SEQ ID NO: 151), ngSTF2.HA1-1His(PR8) (SEQ ID NO: 152), and ngSTF2.HA1-2mutHis(PR8) (SEQ ID NO: 156), and *E. coli*-expressed STF2.HA1-2(PR8) (SEQ ID NO: 90) (see Example #4). Thus, serum antibody recognition of the HA head domain may be focused on a region encompassed by the HA1-2 construct. Further, in vitro refolding of STF2.HA1-2 (SEQ ID NO: 90) produced in *E. coli* results in a protein with equivalent immune reactivity (and therefore similar folded conformation) to a protein processed and secreted from a eukaryotic host (Baculovirus). In addition, serum antibody recognition of the HA head domain may not appear to be

dependent on glycosylation and, at least for PR/8/34 HA, may not be negatively affected by glycosylation. HAs (hemagglutinins) from different influenza strains have variable patterns of N-linked glycosylation. (glycosylation may negatively affect immune recognition of other HAs. Serum antibody recognition of HA may be unaffected by N-glycosylation. It is possible glycosylation may otherwise influence the effectiveness, such as the half-life of a vaccine.

Example 9

Immunogenicity and Efficacy of Recombinant Flagellin-Hemagglutinin Fusion Proteins Produced in Baculovirus

Material and Methods

[0513] Animal studies: Female BALB/c mice (Jackson Laboratory, bar Harbor, Me.) were used at the age of 6-8 weeks. Mice were divided into groups of 10 and received inguinal subcutaneous immunizations on days 0 and 14 as follows:

- 1) PBS (phosphate buffered saline).
- 2) 3 μ g of STF2.HA1-2(PR8) *E. coli* (SEQ ID NO: 90) in PBS
- 3) 3 μ g of STF2.HA1-1(PR8) Baculovirus (SEQ ID NO: 151) in PBS
- 4) 3 μ g of ngSTF2.HA1-1(PR8) Baculovirus (SEQ ID NO: 152) in PBS
- 5) 0.3 μ g of ngSTF2.HA1-1(PR8) Baculovirus (SEQ ID NO: 152) in PBS
- 6) 3 μ g of STF2.HA1-1(PR8) *E. coli* (SEQ ID NO: 89) in PBS
- 7) 0.3 μ g of STF2.HA1-1(PR8) *E. coli* (SEQ ID NO: 89) in PBS

[0514] Mice were bled on days 10 (primary) and 21 (boost), and sera were clarified by clotting and centrifugation and stored at -20° C. Immunized animals were challenged on day 28 with an LD₅₀ (8×10^3 EID) of PR/8/34 virus administered intranasally (See Example 6). Mice were monitored daily for 21 days following the challenge for survival, weight loss and clinical presentation.

[0515] Serum antibody determination: HA-specific IgG levels were determined by ELISA. 96-well ELISA plates (Costar (Cat#9018), Corning, N.Y.) were coated overnight at 4° C. with 100 μ l/well HA0sHis protein produced in *Drosophila* (SEQ ID NO 176) in PBS (5 μ g/ml). Plates were blocked with 200 μ l/well of Assay Diluent Buffer (ADB; BD Pharmingen (Cat# 555213), San Diego, Calif.) for one hour at room temperature. The plates were washed three times in PBS containing 0.05% Tween-20 (PBS-T). Dilutions of the sera in ADB were added (100 μ l/well) and the plates were incubated overnight at 4° C. The plates were washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical (Cat# 115-035-146)) diluted in ADB were added (100 μ l/well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra

substrate (Pierce (Cat# 34028), Rockford, Pa.) and monitoring color development, A₄₅₀ was measured on a Tecan Farcyte (Durham, N.C.) microplate spectrophotometer.

Results and Discussion

[0516] Immunogenicity and efficacy of STF2.HA1-1(PR8) expressed in *E. coli* and Baculovirus: BALB/c mice were immunized s.c. with 3.0 or 0.3 μ g of indicated recombinant fusion proteins on days 0 and 14. On day 21 mice were bled and HA-specific IgG titers were examined by ELISA against HA0sHis (SEQ ID NO 176) expressed in *Drosophila*. FIGS. 12A, 12B shows the proteins induced varying levels of HA-specific IgG with ngSTF2.HA1-1(PR8) (SEQ ID NO: 152) inducing the strongest response similar to that observed in animals immunized with STF2.HA1-2(PR8) expressed in *E. coli* (SEQ ID NO: 90). Interestingly, animals immunized with STF2.HA1-1(PR8) containing the intact glycosylation sequences expressed in Baculovirus (SEQ ID NO: 151) elicited little to no detectable antibody responses to HA or flagellin, in marked contrast to that observed in animals similarly immunized with STF2.HA1-1(PR8) where the consensus glycosylation sequences had been removed (SEQ ID NO: 152).

[0517] The immunized mice were challenged on day 28 with an LD₅₀ (8×10^3 EID) of PR/8/34 virus administered intranasally. Mice were monitored daily for 21 days following the challenge for survival, weight loss and clinical presentation. As shown in FIGS. 13A, 13B, 13C, PBS-immunized mice showed signs of infection (weight loss and lower clinical scores) as early as three days after challenge and died by day 21, while mice immunized with 3.0 μ g of recombinant STF2.HA1-2(PR8) (SEQ ID NO: 90) fusion protein demonstrated enhanced protection with little to no observable clinical signs of infection or weight loss. Although animals immunized with STF2.HA1-1 fusion proteins demonstrated significantly higher efficacy than animals immunized with PBS alone, these animals demonstrated lower efficacy in terms of survival, clinical scores and overall weight loss, compared to animals receiving STF2.HA1-2(PR8) (SEQ ID NO: 90). These results demonstrate that HA subunits linked to STF2 in Baculovirus are immunogenic and efficacious, and suggest that STF2.HA1-1 (SEQ ID NO: 89) expressed in *E. coli* or Baculovirus is less immunogenic and efficacious than STF2.HA1-2 expressed in *E. coli* (SEQ ID NO: 90) or Baculovirus.

Example 10

Purification of STF2.HA1-2 (IND) (SEQ ID NO: 159)

Materials and Methods

[0518] Cell banking: *E. coli* cells engineered to express STF2.HA1-2(IND) (SEQ ID NO: 159) were adapted to culture in proprietary MRSF media, banked as glycerol stocks in 1 mL aliquots, and stored at -70° C.

MRSF media, pH 7.0

Composition	g/L
Glucose	10
KH ₂ PO ₄	7.8
(NH ₄) ₂ SO ₄	2.33
Citric Acid	1.0
MgSO ₄ (7H ₂ O)	1.0
CaCl ₂	0.04
Trace Metals	1 ml
Thiamine HCl	0.01
Kanamycin	0.0075

[0519]

Trace Metal Solution 1000x

Component	g/L
EDTA, disodium	5
FeSO ₄ (7H ₂ O)	10
ZnSO ₄ (7H ₂ O)	2
MnSO ₄ (H ₂ O)	2
CoCl ₂ (6H ₂ O)	0.2
CuSO ₄ (5H ₂ O)	0.1
Na ₂ MoO ₄ (2H ₂ O)	0.2
H ₃ BO ₃	0.1
H ₂ O	1000 ml

[0520] Cell scale-up: Two vials (2 ml, banked in MRSF media) of *E. coli* cells engineered to express STF2.HA1-2 (IND) (SEQ ID NO: 159) were retrieved from -70° C. and inoculated into 500 ml of MRSF medium. The cells were expanded by incubation at 37° C. for 11.5 hours, at which time the OD₆₀₀ was 3.2. The cell scale up provided biomass to inoculate the production bioreactor used in the fermentation.

[0521] Fermentation: A 250 mL aliquot of scaled up cells was used to inoculate a 12 L working volume bioreactor containing 10 L of proprietary MRBR media. The culture was incubated with constant stirring while maintaining pH at 7.0±0.1, temperature at 30±0.1° C., and dissolved oxygen (DO) at no less than 30%, and was run in batch mode until the glucose was exhausted. Glucose exhaustion is indicated by a sudden decrease in O₂ demand, and confirmed by measurement of glucose concentration on a YSI 2700 Select glucose meter. The OD₆₀₀ for the culture at this time was 20.1 AU.

[0522] Thirty minutes after glucose exhaustion, proprietary feed media was pumped into the bioreactor at a controlled rate until 2 L of feed was added. The incubation continued for 5.5 hours under glucose-limiting conditions. At an OD₆₀₀ of 34 AU, protein expression in the culture was induced by the addition of IPTG to a final concentration of 2 mM and incubation with constant stirring for 2 hours. At the conclusion of the induction period, cells were harvested by centrifugation in an Avanti JP-20 XP centrifuge for 20 minutes at 10,000 g. The final OD₆₀₀ for the culture was 58 AU and the total protein concentration as determined by BCA was 6.73 mg/mL. 1.16 kg of cell paste was recovered from 10.8 L harvest and frozen at -20° C. The total

bioreactor run time was 17 hours. STF2.HA1-2 (IND) (SEQ ID NO: 159) production was confirmed by SDS PAGE.

MRBR Media, pH 7.0

Composition	g/L
Glucose	20
KH ₂ PO ₄	2.2
(NH ₄) ₂ SO ₄	4.5
Citric Acid	1.0
MgSO ₄ (7H ₂ O)	1.0
CaCl ₂	0.04
Trace Metals	1 ml
Thiamine HCl	0.01
Antifoam	0.05
Kanamycin	0.0075

[0523]

Feed media, pH 6.0 - additions to
MRBR media (2 L final volume)

Composition	g/L
Glucose	160
KH ₂ PO ₄	5.6
DL-Alanine	50.0
Citric Acid	3.0
MgSO ₄ (7H ₂ O)	1.0 (added as 78 g/L solution)
CaCl ₂	2.5 (added as 80 g/L solution)
Trace Metals	24 ml
FeSO ₄ 7H ₂ O	0.75

[0524] Cell disruption and clarification: The cell paste was thawed and resuspended into a 15% suspension of 50 mM Tris, 25 mM NaCl, pH 8. The slurry was homogenized in an APV 1000 homogenizer three times at 12,000 psi. The lysate was adjusted to pH 4.0 by the addition of acetic acid and centrifuged to separate soluble from insoluble material. STF2.HA1-2 (IND) (SEQ ID NO: 159) along with most other proteins partition to the insoluble fraction. The soluble material was discarded and the insoluble material (pellet) was dissolved in two times the initial lysate volume of 50 mM acetate, 1% TritonX-100, 8 M Urea, pH 4. After mixing in the homogenizer at 0 psi, the sample was centrifuged and the supernatant was collected, filtered, and stored at 4° C. The presence of STF2.HA1-2 (IND) (SEQ ID NO: 159) in the proper phases (precipitate or supernatant) was confirmed by SDS PAGE.

[0525] Cation exchange chromatography (CEX): The clarified material was applied to a 50 ml Poros 50HS column. This step was designed to reduce the endotoxin concentration, reduce the nucleic acid concentration and increase the STF2.HA1-2 (IND) (SEQ ID NO: 159) purity of the eluate as compare to the load. After loading, the column washed with five column volumes of 50 mM acetate, 1% Triton X-100, 8 M Urea, pH 4 followed by 10 column volumes of 50 mM acetate, 8 M Urea, pH 4. The protein was eluted in a 0-100% gradient to 1 M NaCl. The target protein eluted at approximately 300 mM NaCl. The major peak (eluate) was pooled and stored at 4° C. The presence of STF2.HA1-2 (IND) (SEQ ID NO: 159) in the elution was confirmed by SDS PAGE. STF2.HA1-2 (IND) (SEQ ID NO:

159) purification across this step was determined by comparing the number and intensity of protein bands in the load material to those in the elution.

[0526] Endotoxin removal in cation exchange chromatography (CEX): Endotoxin is removed from eluting STF2.HA1-2 (IND) (SEQ ID NO: 159) with the 50 mM acetate, 1% Triton X-100, 8 M Urea, pH 4 wash and with proper elution conditions. This was tested using STF2.HA1-2 (VN) (SEQ ID NO: 95) Comparing two separate cation exchange runs (Toso SP650M resin) with and without the 50 mM acetate, 1% Triton X-100, 8 M Urea, pH 4 wash (ER wash vs. no ER wash), the endotoxin concentration in the ER wash elution at 150 mM NaCl shows significantly less endotoxin concentration than the 250 mM NaCl elution from the same run, or in either of the elutions without the 50 mM acetate, 1% Triton X-100, 8 M Urea, pH 4 wash.

Run	Elution (mM NaCl)	[Endotoxin] (EU/mL)	[Endotoxin] (EU/mg)	[protein] by BCA (g/L)	UV 280/260
Load	N/A	4.4×10^6	1.8×10^6	2.39	—
No	100	7.9×10^6	4.2×10^6	1.44	1.05
ER	250	8.0×10^6	4.1×10^6	1.12	2.76
wash					
ER	150	5,112	2,888	1.77	0.982
Wash	250	24,440	N/A	N/A	0.545

[0527] Refolding: The CEX eluate was dialyzed against 50 mM Tris, 8 M Urea, pH 8 buffer, and refolded by rapid dilution into 10 volumes of 50 mM Tris, pH 8.

[0528] Anion Exchange Chromatography (AEX): The refolded protein was applied to a 25 ml GE Source Q column. This step is designed to separate properly folded STF2.HA1-2 (IND) (SEQ ID NO: 159) from aggregates and host cell proteins and to reduce the endotoxin concentration of the eluate as compared to the load. After loading, the column washed with five column volumes of 50 mM Tris, 25 mM NaCl, pH 8. The protein was eluted in a 0-100% gradient to 1 M NaCl. A peak enriched for properly folded STF2.HA1-2(IND) (SEQ ID NO: 159) with low endotoxin (1.7 EU/mL) eluted at approximately 150 mM NaCl. A second peak eluting at approximately 170 to 270 mM NaCl contains other proteins including STF2.HA1-2 (IND) (SEQ ID NO: 159) aggregates. Fractions in first peak were pooled at stored at 4° C. Fractions from the first peak were kept for further processing. The presence of STF2.HA1-2 (IND) (SEQ ID NO: 159) in the first and second elutions was confirmed by SDS PAGE. STF2.HA1-2 (IND) (SEQ ID NO: 159) purification across this step was determined by comparing the number and intensity of protein bands in the load material to those in the elution.

[0529] Preparative size exclusion chromatography (SEC): The AEX eluate was run on a 320 ml SEC column in 2x5 mL fractions against TBS. This step is designed to separate correctly folded STF2.HA1-2 (IND) (SEQ ID NO: 159) from other proteins. Each run resulted in elution of a single major peak which was collected, pooled, and filtered through a 0.22 µm filter. The product was stored at -70° C.

[0530] The presence of STF2.HA1-2 (IND) (SEQ ID NO: 159) in the elution was confirmed by SDS PAGE.

STF2.HA1-2 (IND) (SEQ ID NO: 159) purification across this step was determined by comparing the number and intensity of protein bands in the load material to those in the elution. The final material profile on SDS PAGE is a strong single band.

Results and Discussion

[0531] Final characterization of protein: Recombinant STF2.HA1-2(IND) (SEQ ID NO: 159) protein was expressed in *E. coli* and purified to homogeneity. The final product was formulated in 1xTBS (27.7 mM Tris, 2.7 mM KCl, 137 mM NaCl, pH 7.4; made from 10x stock, Teknova catalog #T9530) and stored at -70° C. in aliquots of 1.0 ml. The final yield was 15.75 ml (45 ml at a concentration of 0.35 mg/ml), with an endotoxin level of 1.1 EU/mg protein as determined by the LAL method. The protein retained TLR5 biological activity as measured in a cell-based assay of cytokine release

[0532] SDS-PAGE: Protein identity was determined, and purity estimated, by SDS-PAGE. An aliquot sample was diluted in SDS-PAGE sample buffer and diluent. The samples were boiled for 5 minutes and loaded onto a 4 to 12% SDS polyacrylamide gel (Invitrogen NuPage) and electrophoresed. The gel was stained with Coomassie R-250 to visualize protein bands. Positive results for STF2.HA1-2 (IND) (SEQ ID NO: 159) were indicated by a band at about 75 kDa as compared to BioRad precision plus standards run on the same gel. Purity is estimated by visually comparing the intensity of the STF2.HA1-2 (IND) (SEQ ID NO: 159) band to all other bands.

[0533] TLR5 bioactivity assay: HEK293 cells constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96-well microplates (50,000 cells/well), and test proteins were added and incubated overnight. The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20° C. After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce, #M801E and #M802B) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer (FARCyte, Amersham).

[0534] Endotoxin assay: Endotoxin levels were determined using the QCL-1000 Quantitative Chromogenic LAL test kit (Cambrex), following the manufacturer's instructions for the microplate method.

[0535] Protein assay: Total protein concentration was determined using the Micro BCA (bicinchoninic acid) Assay (Pierce) in the microplate format, using bovine serum albumin as a standard, according to the manufacturer's instructions.

[0536] UV280/260 assay: This assay is a measure of the nucleic acid concentration in a sample as compared to the protein concentration. The absorbance maximum occurs at approximately 260 nm for nucleic acids, while the absorbance maximum occurs for proteins at approximately 280 nm. A higher ratio between the absorbance at 280 nm versus the absorbance at 260 nm indicates a sample enriched in protein over nucleic acids.

Example 11

Cloning, Expression, and Biochemical
Characterization of Recombinant
Flagellin-Hemagglutinin Fusion Proteins Produced
in *Drosophila*

Materials and Methods

[0537] Cloning and expression of HA in *Drosophila*: RNA was isolated from the influenza A strain A/Puerto Rico/8/34 using QIAamp MinElute Virus Spin Kit (Qiagen, Cat#57704) following the manufacturer's instructions. The HA RNA was reverse transcribed using SuperScript III First Strand Synthesis System for RT-PCR (Invitrogen; Cat #18080-051) or SuperScript III One-Step RT-PCR System with Platinum-Taq High Fidelity (Invitrogen; Cat# 12574-030) following the manufacturer's instructions, and the cDNA was cloned using the TOPO cloning vector PCR II Zero Blunt (Invitrogen). The HA0s fragment (HA gene without the signal sequence and trans-membrane domain) was then sub-cloned into pMT/STF2Δ at both the C-terminus and NH2 terminus of the cassette using PCR-based strategy to generate the constructs STF2Δ.HA0s (PR8) (SEQ ID NO: 160) and HA0s(PR8).STF2Δ (SEQ ID NO: 167) respectively. The HA gene was also cloned into the pMT/Bip/V5-his vector without STF2Δ to generate the construct HA0s.his (SEQ ID NO: 170). Table 10 list the constructs and the primers used in their construction.

TABLE 10

PR8 HA constructs for expression in <i>Drosophila</i>				
SEQ ID NO:	Construct	FOR Primer SEQ ID NO:	REV Primer SEQ ID NO:	DNA Template SEQ ID NO:
160	STF2Δ.HA0sHis	161	162	163
164	STF2Δ.HA0s	165	166	163
167	HA0s.STF2Δ	168	169	163
170	HA0sHis	171	172	163
173	HA0s	174	175	163

[0538] Recombinant protein expression from *Drosophila* Dmel-2 cells: The HA0s.His (SEQ ID NO: 170) and STF2Δ.HA0s (SEQ ID NO: 164) plasmids were co-transfected with the pCoBlast plasmid into *Drosophila* Dmel-2 cells to generate Dmel-2 HA0sHis(PR8) and Dmel-2 STF2Δ.HA0s(PR8) stable cells. Dmel-2 stable cells were expanded from adherent to shaker cultures in selection media [*Drosophila* Sfm medium (Invitrogen; Carlsbad, Calif.)+25 μg/ml blasticidin] and then expanded to a 12 L production scale.

[0539] Recombinant protein expression was induced by the addition of CuSO₄ to a final concentration of 0.5 mM. After incubation for 72 hours cells were removed from the conditioned medium by centrifugation. The conditioned medium was then clarified by passing through a 0.22 μm filter and stored at 4° C.

[0540] Purification of HA0sHis(PR8) (SEQ ID NO: 176): Conditioned medium was applied to a Chelating Sepharose column (GE/Amersham Biosciences; Piscataway, N.J.) charged with NiSO₄ and equilibrated in Buffer A [20 mM Tris, pH 8.0, 0.5 M NaCl] and eluted in a linear gradient with

Buffer B [Buffer A+0.5 M imidazole]. The protein was further purified by fractionation on a Superdex 200 SEC column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated in 1× Tris-buffered saline (TBS), pH 8.0. HA0s.His₆(PR8) fractions were pooled, aliquoted and stored at -80° C.

[0541] SDS-PAGE and western blot analysis: Protein identity was determined, and purity estimated, by SDS-PAGE. An aliquot of 5 μg of each sample was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel (LifeGels; French's Forrest, New South Wales, AUS) and electrophoresed. The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands. For western blot, 0.5 μg/lane total protein was electrophoresed as described above and the gels were then electro-transferred to a PVDF membrane and blocked with 5% (w/v) dry milk before probing with anti-flagellin antibody (Inotek; Beverly, Mass.), anti-His₆ antibody (Invitrogen; Carlsbad, Calif.) or influenza A PR/8/34 convalescent immune serum (described below under Protein Antigenicity ELISA). After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockford, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega; Madison, Wis.).

[0542] Protein assay: Total protein concentration was determined using the Micro BCA (bicinchoninic acid) Assay (Pierce; Rockford, Ill.) in the microplate format, using bovine serum albumin as a standard, according to the manufacturer's instructions.

[0543] Endotoxin assay: Endotoxin levels were determined using the QCL-1000 Quantitative Chromogenic LAL test kit (Cambrex; E. Rutherford, N.J.), following the manufacturer's instructions for the microplate method.

[0544] TLR5 bioactivity assay: HEK293 cells constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96-well microplates (50,000 cells/well), and recombinant *drosophila* conditioned medium containing either HA0sHis(PR8) (SEQ ID NO: 176) or STF2Δ.HA0s(PR8) (SEQ ID NO: 177) was added. The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20° C. After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce; Rockford, Ill., #M801E and #M802B) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer (FARCyte, GE/Amersham; Piscataway, N.J.).

[0545] Protein antigenicity ELISA: Purified HA0s.His₆(PR8) (SEQ ID NO: 176) was tested by ELISA to determine if the recombinant protein displayed correctly folded epitopes of HA. 96-well ELISA plates were coated overnight at 4° C. with serial dilutions in PBS (100 μl/well) of HA0s.His₆(PR8) (SEQ ID NO: 176) protein starting at 5 μg/ml. Plates were blocked with 200 μl/well of Assay Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature, then washed three times in PBS-T. A fixed dose of primary antibody was then added to each well. To assay HA reactivity, 100 μl/well of a 1:10,000 dilution of non-immune or PR/8/34 convalescent immune serum in ADB

was added. PR/8/34 immune serum was generated in BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) that received an experimentally determined sublethal challenge dose of 8×10^1 egg infectious dosages (EID) of PR/8/34 influenza virus.

[0546] Animals were then allowed to convalesce for >21 days post-infection at which time immune serum was isolated and clarified. For ELISA of the 6× histidine tag, monoclonal antibody against 6×His (Invitrogen; Carlsbad, Calif.), or flagellin (Inotek; Beverly, Mass.) was added at 1 µg/ml in ADB (100 µl/well) and the plates were incubated for 1 hr at room temperature or overnight at 4° C. The plates were then washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical; West Grove, Pa.) diluted in ADB were added (100 µl/well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce; Rockford, Ill.) and monitoring color development, A_{450} was measured on a microplate spectrophotometer (FARCyte, GE/Amersham; Piscataway, N.J.).

Results and Discussion

[0547] Characterization of *Drosophila*-expressed HA0s proteins: Western blot analysis of conditioned media with anti-His₆ antibody (Invitrogen; Carlsbad, Calif.) confirmed expression of HA0sHis(PR8) (SEQ ID NO: 176) and western blot with anti-flagellin antibody (Inotek; Beverly, Mass.) confirmed expression of STF2Δ.HA0s(PR8) (SEQ ID NO: 177) by *Drosophila* Dmel-2 cells transfected with the corresponding expression plasmid. Both proteins were recognized by western blot in non-reduced form with mouse PR/8/34 convalescent immune serum while reduction of the proteins with DTT abrogated recognition. This result indicates correct disulfide bonding of the two secreted proteins. The STF2Δ.HA0s(PR8) (SEQ ID NO: 177) conditioned medium showed significant in vitro TLR5 activity while the HA0sHis₆(PR8) (SEQ ID NO: 176) medium did not, as expected. Finally, purified HA0sHis₆(PR8) (SEQ ID NO: 176) protein showed significant reactivity with influenza A PR8/34 convalescent immune serum by ELISA. These results indicate that both HA0s proteins are secreted from *Drosophila* Dmel-2 cells in a properly folded form.

Example 12

Immunogenicity and Efficacy of Recombinant
Flagellin-Hemagglutinin Fusion Proteins
Representing Viral Strain A/Viet Nam/1203/04

Materials and Methods

[0548] Animal studies: Female BALB/c mice were used at the age of 6-8 weeks. Mice were divided into groups of 15 and received subcutaneous (s.c) immunizations on days 0 and 14 as follows:

[0549] TBS (phosphate buffered saline)

[0550] No immunization

[0551] 10 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) in TBS

[0552] 3.0 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) in PBS

[0553] 1 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) in TBS

[0554] Mice were bled on day 21 and sera were clarified by clotting and centrifugation and stored at -20° C.

[0555] Serum antibody determination: HA-specific IgG levels were determined by ELISA. 96-well ELISA plates (Costar (Cat #9018) Corning, N.Y.) were coated overnight at 4° C. with 100 µl/well recombinant HA protein from A/Vietnam/1203/04, produced in Baculovirus, (BEIR catalog number NR-660) in PBS (1 µg/ml). Plates were blocked with 300 µl/well of Assay Diluent Buffer (ADB; BD Pharmingen, (Cat#: 555213) (San Diego, Calif.) for two hours at 25° C. The plates were washed three times in PBS+0.05% (v/v) Tween 20 (PBS-T). Dilutions of the sera in ADB were added (100 µl/well) and the plates were incubated for 1.5 hours at 25° C. The plates were washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical, West Grove, Pa. (Cat#: 115-035-146)) diluted in ADB were added (100 µl/well) and the plates were incubated at 25° C. for 30 minutes. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce (Cat 34028), Rockford, Ill.)) and monitoring color development, A_{450} was measured on a SpectraMax 190 (Molecular Devices, Sunnyvale, Calif.) microplate spectrophotometer.

[0556] Influenza virus challenge of mice: To assess efficacy, mice were challenged on day 28 by intranasal administration of 10 LD₅₀ (10×dose lethal to 90% of mice) (6×10^3 EID) of influenza A/Viet Nam/1203/04. Animals were monitored daily for 21 days following the challenge for survival and clinical presentation.

[0557] Preparation of Influenza H5N1 Stocks: Influenza A/Vietnam/1203/04 (H5N1) was obtained from the Centers for Disease Control and Prevention (Atlanta, Ga.). Stocks of the virus were prepared in 10-day-old embryonated chicken eggs, and aliquots of 0.5 ml were stored at -80° C. until use.

[0558] Determination of Viral Titers: TCID₅₀ (50% tissue culture infectious dose) was determined utilizing MDCK cells seeded into the wells of 96-well plate and grown to confluency. A series of log₁₀ dilutions of the virus was inoculated onto the plate in quadruplicates. The plate was then incubated for 48-72 hours. The TCID₅₀ was determined by identifying the dilution at which quadruplicate wells were ½ positive and ½ negative for viral growth. The EID₅₀ was determined by inoculating log₁₀ dilutions of the stock into 2-4 eggs per dilution. Eggs were incubated for 40-48 hours, and 1 ml of the allantoic fluid was harvested from each egg. ED₅₀ was calculated as the dilution at which ½ of the eggs were negative and ½ of the eggs were positive for infectious virus.

Results and Discussion

[0559] Induction of HA-specific IgG response following immunization with STF2.HA1-2(VN) (SEQ ID NO: 95): The immunogenicity of STF2.HA1-2(VN) (SEQ ID NO: 95) was examined by immunizing BALB/c mice (15/group) subcutaneously on day 0 and 14 with a dose range of 10, 3, and 1 µg protein. Negative control groups of mice were immunized with TBS or not immunized. HA-specific IgG responses were examined 7 days post boost (Day 21) by ELISA. The results demonstrate that immunization with 10,

3 or 1 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) induced consistent and significant HA-specific IgG responses in a dose-dependent manner (FIGS. 14-16).

[0560] Immunization with STF2.HA1-2(VN) (SEQ ID NO: 95) provides protection from a lethal challenge with influenza A: The serological analysis described above demonstrated that immunization of mice with STF2.HA1-2(VN) (SEQ ID NO: 95) generated an antibody response that recognized native HA. In order to evaluate efficacy, the same mice were challenged on day 28 with 10 LD₅₀ (6×10⁵ EID) of A/Viet Nam/1203/04 virus administered intra-nasally. Mice were monitored daily for 21 days following the challenge for survival and clinical presentation. As shown in FIG. 17, TBS-immunized or naïve mice showed signs of infection (weight loss and lower clinical scores) as early as five days post-challenge and all mice died by day nine post-challenge. For the mice immunized with STF2.HA1-2(VN), there is a clear relationship between the dose received and efficacy. Mice immunized with 10 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) demonstrated markedly enhanced protection. Mice immunized with 3 or 1 µg of STF2.HA1-2(VN) (SEQ ID NO: 95) demonstrated a modest to a negligible impact on efficacy, respectively. Clinical parameters were similarly related to dose and animals receiving the highest dose of STF2.HA1-2(VN) (SEQ ID NO: 95) exhibited the mildest signs of disease. These results demonstrate that *E. coli*-expressed STF2.HA1-2(VN) (SEQ ID NO: 95) induces HA-specific immune responses that successfully protect BALB/c mice from a lethal challenge with virulent influenza A virus in vivo.

SEQ ID NO:1 A/Puerto Rico/8/34 HA
MKANLLVLLSALAADADTICIGYHANNSTDTVDTVLEKNVTVTHSVNL
EDSHNGKLCRLKGIAPLQLGKCNIAWLLGNPECDPLLPVRSWSYIVETP
NSENGICYPGDFIDYEELREQLSSVSSFERFEIPKSSWPHNHTNGVTA
ACSHGKSSFYRNLLWLTEKEGSYPKLKNSYVKKGKEVLVLWGVIHPPN
SKEQQLNYQENAYVSVVTSNYYNRRFTPEIAERPKVRDQAGRMNYYWTL
KPGDTIIFEANGNLIAPMYAFALSRGFGSGIITSNASMHECNTKQTPLG
AINSSLFPQNIHPVTIGCEPKYVRSALRMVTGLRNIPSIQSRGLFGAIA
GFIEGGWTGMDGWYGYHHQNEQSGGYAADQKSTQNAINGITNKVNTVIE
KMNIQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTYNAELLVLENER
LDFHDSNVKNLYEKVKSQKNNAKEIGNGCFEFYHKCDNECMESVRNGTY
DYPKYSEESKLNREKVDGVKLESMGIYQILAIYSTVASSLVLLVSLGAIS
FWMCSNGSLQCRICI

SEQ ID NO:2 A/Viet Nam/1203/2004 HA
MEKIVLLFAIVSLVKSDQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILE
KKHNGKLCDDLGVKPLILRDCSVAGWLLGNPMCEDEFINVPESYIVEKAN
PVNDLCYPGDFNDYEELKHLISRINHFEKIQIIPKSSWSSEASLGVSSA
CPYQKSSFFRNVVWLKKNSTYPTIKRSYNNNTQEDLLVLWGIIHPPNDA
AEQTKLYQNPTTYISVGTSTLNQRLVPRIATRSKVNQSGRMEFFWTILK
PNDAINFESNGNFIAPYAYKIVKKGDSIMKSELEYGNCNTKQTPMGA

-continued

INSSMPFHNIHPLTIGCEPKYVKSRLVLATGLRNSPQERRRRKKRGLFG
AIAGFIEGGWQGMVDGWYGYHHSNEQSGGYAADKESTQKAIDGVTNKVNS
IIDKMNTQFEAVGREFNNLERRIENLNKKMEDGFLDVWTYNAELLVLMEN
ERTLDFHDSNVKNLYDKVRLQLRDNALGNGCFEFYHKCDNECMESVRN
GTDYPQYSEEARLKREEISGVKLESIGIYQILSIYSTVASSLALAIMVAG
LSLWMCNSNGSLQCR

SEQ ID NO:3 A/Indonesia/5/2005 HA
MEKIVLLFAIVSLVKSDQICIGYHANNSTEQVDTIMEKNVTVTTHAQDILE
KTHNGKLCDDLGVKPLILRDCSVAGWLLGNPMCEDEFINVPESYIVEKAN
PTNDLCYPGDFNDYEELKHLISRINHFEKIQIIPKSSWSSEASLGVSSA
CPYLGSPSFFRNVVWLKKNSTYPTIKRSYNNNTQEDLLVLWGIIHPPNDA
AEQTRLYQNPTTYISIGTSTLNQRLVPKIATRSKVNQSGRMEFFWTILK
PNDAINFESNGNFIAPYAYKIVKKGDSAIMKSELEYGNCNTKQTPMGA
INSSMPFHNIHPLTIGCEPKYVKSRLVLATGLRNSPQERRRRKKRGLFG
AIAGFIEGGWQGMVDGWYGYHHSNEQSGGYAADKESTQKAIDGVTNKVNS
IIDKMNTQFEAVGREFNNLERRIENLNKKMEDGFLDVWTYNAELLVLMEN
ERTLDFHDSNVKNLYDKVRLQLRDNALGNGCFEFYHKCDNECMESVRN
GTYNYPQYSEEARLKREEISGVKLESIGTYQILSIYSTVASSLALAIMMA
GLSLWMCNSNGSLQCRICI

SEQ ID NO:4 A/NewCaledonia/20/1999
MKAKLLVLLCTFTATYADTICIGYHANNSTDTVDTVLEKNVTVTHSVNL
EDSHNGKLCCLLKGIAPLQLGNCVAGWILGNPECELLISKESWSYIVETP
NPENGTCPYGFADYEELREQLSSVSSFERFEIPKSSWPHNHTVTGVSA
SCSHNGKSSFYRNLLWLTKNGLYPNLSKSYVNNKEKEVLVLWGVIHPPN
IGNQRALYHTENAYVSVVSHYSRRFTPEIAKRPKVRDQEGRIYNYWTLL
EPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNAPMDECDKQTPQG
AINSSLFPQNVHPVTIGCEPKYVRSALRMVTGLRNIPSIQSRGLFGAIA
GFIEGGWTGMDGWYGYHHQNEQSGGYAADQKSTQNAINGITNKVNSVIE
KMNTQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTYNAELLVLENER
LDFHDSNVKNLYEKVKSQKNNAKEIGNGCFEFYHKCDNECMESVRNGTY
DYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAIS
FWMCSNGSLQCRICI

SEQ ID NO:5 A/South Carolina/1/18 HA
MEARLLVLLCAFAATNADTICIGYHANNSTDTVDTVLEKNVTVTHSVNL
EDSHNGKLCCLLKGIAPLQLGKCNIAWLLGNPECDLLLTASSWSYIVETS
NSENGTCYPGDFIDYEELREQLSSVSSFEKFEIPKSTSSWPHNETTKGVT
AACS YAGASSFYRNLLWLTKKGSSYPKLSKSYVNNKGKEVLVLWGVIHPPN
TGTDQQSLYQNADAYVSGSSKYNRRFTPEIAARPKVRDQAGRMNYYWTL
LEPGDTITFEATGNLIAPWYAFALNRGSGSGIITSAPVHDCNTKQTPH
GAINSSLFPQNIHPVTIGCEPKYVRSATKLRMATGLRNIPSIQSRGLFGAI

-continued

AGFIEGGWTGMIDGWYGYHHQNEQSGGYAADQKSTQNAIDGITNKVNSVI
 EKMNTQFTAVGKEFNLLERRIENLNKKVDDGFLDIWTYNAELLVLENER
 TLDPHDSNVRNLYEKVKSQALKNNAKEIGNGCFEFYHKCDDACMESVRNGT
 YDYPKYSEESKLNREEIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

SEQ ID NO:6 A/Wisconsin/67/2005 HA
 QKLPGNNDSTATLCLGHAVPNGTIVKTIITNDQIEVTNATELVQSSSTGG

ICDSPHQILDGENCTLIDALLGDPQCDGFQNKKWDLFVERS KAYSNCYPY
 DVDPDYASLRSLVASSGTLEFNDESFNWTGVTQNGTSSACKRRSNNSSF SR
 LNWLTHLKFKYPALNVMTMPNNEKFDKLYIWGVHHPGTDNDQIFLHAQASG
 RITVSTKRSQQTIVIPNIGSRPRI RNIPSRISYIWTIVKPGDILLINSTGN
 LIAPRGYFKIRSGKSSIMRSDAPIGKCNSECITPNGSIPNDKPFQNVNRI
 TYGACPRYVKQNTLKLATGM RNVP EKQTR

SEQ ID NO:7 A/X31 subtype H3N2 HA
 QDLPGNDNSTATLCLGHAVPNGTLVKTIITDDQIEVTNATELVQSSSTGK

ICNNPHRILDGIDCTLIDALLGDPHCDVFQNETWDLFVERSKAFSNCYPY
 DVDPDYASLRSLVASSGTLEFITEGFTWTGVIQNGGSNACKRGP GSGFFSR
 LNWLTSGSGSTYPVLNVMTMPNNDNFDKLYIWGIHHPSTNQEQTSLYVQASG
 RVTVSTRRSQQTIIIPNIGSRPWRGLSSRISYIWTIVKPGDVLVINSNGN
 LIAPRGYFKMRTGKSSIMRSDAPIDTCISECITPNGSIPNDKPFQNVNKI
 TYGACPKYVKQNTLKLATGM RNVP EKQT

SEQ ID NO:8 PR/8 HA1-1
 SHNGKLCRLKGIAPLQLGKCNIA GWLLGNPECDPLLPVRSWSYIVETPNS

ENGICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWP NHNTNGVTAAC
 SHEGKSSFYRNLLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHHPNSK
 EQQNLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYWTLLKP
 GDTIIFEANGNLIAPMYAFALSRGFGSGIITSNASMHECNTKCTPLGAI
 NSSLPYQNIHPVTIGEC PKYVR

SEQ ID NO:9 PR/8 HA1-2
 KGIAPLQLGKCNIA GWLLGNPECDPLLPVRSWSYIVETPNS ENGICYPGD

FIDYEELREQLSSVSSFERFEIFPKESSWP NHNTNGVTAACSHEGKSSFY
 RNLLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHHPNSKEQQNLYQNE
 NAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYWTLLKPGDTIIFEAN
 GNLIAPMYAFALSRGFGSGIITS

SEQ ID NO:10 PR/8 HA1-3
 NSENGICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWP NHNTNGVTA

ACSHGKSSFYRNLLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHHPN
 SKEQQNLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYWTLL
 KPGDTIIFEANGNLIAPMYAFALSRG

SEQ ID NO:11 VN HA1-1
 EKKHNGKLCDLGDKPLILRDCSVAGWLLGNPMCDEFIN VPEWSYIVEKA

NPVNDLCYPGDFNDYEELKHL LSRINHFEKIQIIPKSSWS SHEASLGVS

-continued

ACPYQGKSSFFRN VVWL IKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPND
 AAETKLYQNPTTYISVGTSTLNQRLVPRIATRSKVNGQSGRMEFFWTIL
 KPNDAINFESNGNFIAPEYAYKIVKKG DSTIMKSELEYGNCNTKCQTPMG
 AINSSMPFHNHPLTIGEC PKYVK

SEQ ID NO:12 VN HA1-2
 GVKPLILRDCSVAGWLLGNPMCDEFIN VPEWSYIVEKANPVNDLCYPGDF

NDYEELKHL LSRINHFEKIQIIPKSSWS SHEASLGVSACPYQGKSSFFR
 NVVWL IKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPNDAAETKLYQNPT
 TYISVGTSTLNQRLVPRIATRSKVNGQSGRMEFFWTILKPNDAINFESNG
 NFIAPEYAYKIVKKG DSTIMKSE

SEQ ID NO:13 VN HA1-3
 NDLCYPGDFNDYEELKHL LSRINHFEKIQIIPKSSWS SHEASLGVSAC

YQGKSSFFRN VVWL IKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPNDAAE
 QTKLYQNPTTYISVGTSTLNQRLVPRIATRSKVNGQSGRMEFFWTILKPN
 DAINFESNGNFIAPEYAYKIVKKG

SEQ ID NO:14 IND HA1-1
 EKTHNGKLCDLGDKPLILRDCSVAGWLLGNPMCDEFIN VPEWSYIVEKA

NPTNDLCYPGDFNDYEELKHL LSRINHFEKIQIIPKSSWS DHEASGVSS
 ACPYLGSPPFFRN VVWL IKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPND
 AAETRLYQNPTTYISIGTSTLNQRLVPK IATRSKVNGQSGRMEFFWTIL
 KPNDAINFESNGNFIAPEYAYKIVKKGSAIMKSELEYGNCNTKCQTPMG
 AINSSMPFHNHPLTIGEC PKYVK

SEQ ID NO:15 IND HA1-2
 GVKPLILRDCSVAGWLLGNPMCDEFIN VPEWSYIVEKANPTNDLCYPGSF

NDYEELKHL LSRINHFEKIQIIPKSSWS DHEASGVSSACPYLGSPSFFR
 NVVWL IKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPNDAAETRLYQNPT
 TYISIGTSTLNQRLVPK IATRSKVNGQSGRMEFFWTILKPNDAINFESNG
 NFIAPEYAYKIVKKGSAIMKSE

SEQ ID NO:16 IND HA1-3
 NDLCYPGSFNDYEELKHL LSRINHFEKIQIIPKSSWS DHEASGVSSAC

YLGSPSFFRN VVWL IKKNSTYPTIKRSYNNTNQEDLLVLWGIHHPNDAAE
 QTRLYQNPTTYISIGTSTLNQRLVPK IATRSKVNGQSGRMEFFWTILKPN
 DAINFESNGNFIAPEYAYKIVKKG

SEQ ID NO:17 NC HA1-1
 SHNGKLCLLKGIAPLQLGNC SVAGWILGNPECELLISKESWSYIVETPNP

ENGTCYPGYFADYEELREQLSSVSSFERFEIFPKESSWP NHNTVTVGSASC
 SHNGKSSFYRNLLWLTGKNGLYPNLSKSYVNKEKEVLVLWGVHHPNIG
 NQRALYHTENAYVSVSSHYSRRFTPEIAERP KVRDQEGRINYWTLLPE
 GDTIIFEANGNLIAPWYAFALSRGFGSGIITS NAPMDECDACKCTPQGAI
 NSSLPQNVHPVTIGEC PKYVR

-continued

SEQ ID NO:18 NC HA1-2
 KGIAPLQLGNCSVAGWILGNPECELLISKESWSYIVETPNPENGTCYPGY
 FADYEELREQLSSVSSFERFEIFPKESSWPNNHTVTGVSASCSHNGKSSFY
 RNLLWLTGKNGLYPNLSKSYVNNKEKEVLVLWGVHHPNIGNQRALYHTE
 NAYVSVVSSHYSRRFTPEIAKRPKVRDQEGRINYWTLLPEGDTIIFEAN
 GNLIAPWYAFALSRGFGSGIITS

SEQ ID NO:19 NC HA1-3
 NPENGTCYPGYFADYEELREQLSSVSSFERFEIFPKESSWPNNHTVTGVS
 A SC SHNGKSSFYRNLLWLTGKNGLYPNLSKSYVNNKEKEVLVLWGVHHPN
 IGNQRALYHTENAYVSVVSSHYSRRFTPEIAKRPKVRDQEGRINYWTLL
 EPGDTIIFEANGNLIAPWYAFALSRG

SEQ ID NO:20 WIS HA1-1
 QSSSTGGICDSPHQILDGENCTLIDALLGDPQCDGFQNKWDLFVERSKA
 YSNCYPYDVPDYASRLSLVASSGTLEFNDESFNWTGVTQNGTSSACKRRS
 NNSFFSRLNLWTLHKFKYPALNVTMPNNEKFDKLYIWGVHHPGTDNDQIF
 LHAQASGRITVSTKRSQQTIVIPNIGSRPRIRNIPSRISYWTIVKPGDIL
 LINSTGNLIAPRGYFKIRSGKSSIMRSDAPIGKCNSECITPNSIPNDKP
 FQNVNORITYGACPRYVK

SEQ ID NO:21 WIS HA1-2
 SPHQILDGENCTLIDALLGDPQCDGFQNKWDLFVERSKAYSNCYPYDVP
 DYASRLSLVASSGTLEFNDESFNWTGVTQNGTSSACKRRS NNSFFSRLNW
 LTHLKFYPALNVTMPNNEKFDKLYIWGVHHPGTDNDQIFLHAQASGRIT
 VSTKRSQQTIVIPNIGSRPRIRNIPSRISYWTIVKPGDILLINSTGNLIA
 PRGYFKIRSGKSSIMRSD

SEQ ID NO:22 WIS HA1-3
 SNCYPYDVPDYASRLSLVASSGTLEFNDESFNWTGVTQNGTSSACKRRSN
 NSFFSRLNLWTLHKFKYPALNVTMPNNEKFDKLYIWGVHHPGTDNDQIFL
 HAQASGRITVSTKRSQQTIVIPNIGSRPRIRNIPSRISYWTIVKPGDILL
 INSTGNLIAPRGYFKIRSG

SEQ ID NO:23 A/Puerto Rico/8/34 HA0s
 DTICIGYHANNSTDVDTVLEKNVTVTHSVNLED SHNGKLCRLKGIAPL
 QLGKCNIA GWLLGNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEE
 LREQLSSVSSFERFEIFPKESSWPNNHTNGVTAACSHGKSSFYRNLLWL
 TEKEGSYPKLKNSYVNNKGKEVLVLWGIHHPNSKEQQONLYQENAYVSV
 VTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLLKPGDTIIFEANGNLIAP
 MYAFALSRGFGSGIITSNASMHECNTKCQTPLGAINSSLPYQNIHPVTIG
 ECPKYVRS AKLRMVTGLRNIPSIQSRGLFGAIA GIEGGWTGMIDGWYGY
 HHQNEQGGSYAADQKSTQNAINGITNKVNTVIEKMNIQFTAVGKEFNKLE
 KRMENLNKKVDDGFLDIWTYNAELLV LLENERTLDFHDSNVKNLYEKVKS
 QLKNNAKE IGNGCFEFYHKCDNECMESVRNGTYDPKYSEESKLNREKVD
 GVKLESMGIYQ

-continued

SEQ ID NO:24 PR/8 HA1-2mut
 KGAAPLQLGKCNIA GWLLGNPECDPLLPVRSWSDIAETPNSENGICYPGD
 FIDYEELREQLSSVSSFERFEIFPKESSWPNNHTNGVTAACSHGKSSFY
 RNLLWLTEKEGSYPKLKNSYVNNKGKEVLVLWGIHHPNSKEQQONLYQNE
 NAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLLKPGDTIIFEAN
 GNLIAPMYAFALSRGFGSGIITS

SEQ ID NO:25 PR/8 HA1-3mut
 NSENEICYPGDFIDKEELREQLSSVSSFERFEIFPKESSWPNNHTNGVTA
 ACSHEGKSSFYRNLLWLTEKEGSYPKLKNSYVNNKGKEVLVLWGIHHPN
 SKEQQONLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLL
 KPGDTIIFEANGNLIAPMYAALSRG

SEQ ID NO:26 VN HA1-2mut
 GAKPLSLRDCSVAGWLLGNPMCDEFINVPESWDIAEKANPVNDLCYPGDF
 NDYEELKHLISRINHFEKIQIIPKSSWS SHEASLGVSACPYQGKSSFFR
 NVVWLICKNSTYPTIKRSYNNNTQEDLLVLWGIHHPNDAAEQTKLYQNPT
 TYISVGTSTLNQRLVPRIATRSKVNGQSGRMEFFWTILKPNDAINFESNG
 NFIAPEYAYKIVKKG DSTIMKSE

SEQ ID NO:27 VN HA1-3mut
 NDLCYPGDFNDKEELKHLISRINHFEKIQIIPKSSWS SHEASLGVSACP
 YQGKSSFFRN VVWLICKNSTYPTIKRSYNNNTQEDLLVLWGIHHPNDAAE
 QTKLYQNPTTYISVGTSTLNQRLVPRIATRSKVNGQSGRMEFFWTILKPN
 DAINFESNGNFIAPEYAYKIAKKG

SEQ ID NO:28 IND HA1-2mut
 GAKPLSLRDCSVAGWLLGNPMCDEFINVPESWDIAEKANPTNDLCYPGSF
 NDYEELKHLISRINHFEKIQIIPKSSWS DHEASSGVSSACPYLGSPSFFR
 NVVWLICKNSTYPTIKRSYNNNTQEDLLVLWGIHHPNDAAEQTRL YQNPT
 TYISIGTSTLNQRLVPKIATRSKVNGQSGRMEFFWTILKPNDAINFESNG
 NFIAPEYAYKIVKKGDSAIMKSE

SEQ ID NO:29 IND HA1-3mut
 NDLCYPGSFNDKEELKHLISRINHFEKIQIIPKSSWS DHEASSGVSSACP
 YLGSPSFFRN VVWLICKNSTYPTIKRSYNNNTQEDLLVLWGIHHPNDAAE
 QTRL YQNPTTYISIGTSTLNQRLVPKIATRSKVNGQSGRMEFFWTILKPN
 DAINFESNGNFIAPEYAYKIAKKG

SEQ ID NO:30 NC HA1-2mut
 KGAAPLQLGNCSVAGWILGNPECELLISKESWSDIAETPNPENGTCYPGY
 FADYEELREQLSSVSSFERFEIFPKESSWPNNHTVTGVSASCSHNGKSSFY
 RNLLWLTGKNGLYPNLSKSYVNNKEKEVLVLWGVHHPNIGNQRALYHTE
 NAYVSVVSSHYSRRFTPEIAKRPKVRDQEGRINYWTLLPEGDTTIFEAN
 GNLIAPWYAFALSRGFGSGIITS

SEQ ID NO:31 NC HA1-3mut
 NPENETCYPGYFADKEELREQLSSVSSFERFEIFPKESSWPNNHTVTGVS
 A SC SHNGKSSFYRNLLWLTGKNGLYPNLSKSYVNNKEKEVLVLWGVHHPN

-continued

IGNQRALYHTENAYVSVSSHYSRRFTPEIAKRPKVRDQEGRINYWTLL
 EPGDTIIFEANGNLIAPWYAAALSRG
 SEQ ID NO:32 WIS HA1-2mut
 SPHQALDGENCTLIDALLGDPQCDFGQNKKWDDFAERSKAYSNCYPYDVP
 DYASLRSLVASSGTLEFNDESFNWTGVTQNGTSSACKRRSNNSSFRLNW
 LTHLKFKYPALNVTMPNNEKFDKLYIWGVHHPGTDNDQIFLHAQASGRIT
 VSTKRSQQTVIPNIGSRPRIRNIPSRISYWTIVKPGDILLINSTGNLIA
 PRGYFKIRSGKSSIMRSD
 SEQ ID NO:33 WIS HA1-3mut
 SNCYPKDSPEASLRSLVASSGTDEFNDESFNWTGVTQNGTSSACKRRSN
 NSFFSRLNWLTHLKFKYPALNVTMPNNEKFDKLYIWGVHHPGTDNDQIFL
 HAQASGRITVSTKRSQQTVIPNTGSRPRIRNIPSRISYWTIVKPGDILL
 INSTGNLIAPRGYFKIRSG
 SEQ ID NO:34 A/Swine/Hong Kong/9/98 HA
 (ACCESSION BAB85618)
 DKICIGYQSTNSTETVDTLTETNPVPTHAKELLHTEHNGMLCATNLGHPL
 ILDTCTIEGLIYGNPSCDLLGGREWSYIVERPSAVNGMCYPGNVENLEE
 LRSLFSSASSYQRIQIFPDTIWNVSYSGETSKACSDSFYRSMRWLTQKNNNA
 YPIQDAQYTNNRGKSIILFMWGINHPPTDTVQTNLYTRDTTTSVTTEDIN
 RTFKPVIGRPLVNLGHGRIDYYSVLKPGQTLRVRSGNGLIAPWYGHIL
 SGESHGRILKTDLNSGNCVVQCQTERGGLNTTLPFHNVSKEYAFGNCPKYV
 GVKSLKAVGLRNVPARSSR
 SEQ ID NO:35 A/Aichi/2/1968 HA (ACCESSION
 BAF37221)
 MKTIIALS YTFCLALGQDLPGNDNSTATLCLGHHA VPNGTLVKTTITDDQI
 EVTNATELVQSSSTGKICNNPHRILDGIDCTLIDALLGDPHCDVFQNETW
 DLFVERSKAFSNCYPYDVPDYASLRSLVASSGTLEFITEGFTWTGVTQNG
 GSNACKRGP GSGFFSRLNWLTKSGSTYPVLNVTMPNNDNFDKLYIWGIHH
 PSTNQEQTSLYVQASGRVTVSTRRSQQTIIPNIGSRPWVRLSSRISYW
 TIVKPGDVLVINSNGNLIAPRGYFKMRTGKSSIMRSDAPIDTCISECITP
 NGSIPNDKPFQNVNKITYGACPKYVKQNTLKLATGMRNVPEKQTRGLFGA
 IAGFIENGWEGMIDGWYFRHQNSEGTQAADLKSTQAAIDQINGKLN RV
 IEKTNEKFHQIEKEFSEVEGRIQDLEKYVEDTKIDLWSYNAELLVALENQ
 HTIDLTDSEMKNLFKERRQLRENAEEMGNGCFKIYHKCDNACIESIRNG
 TYDHDVYRDEALNNRFQIKGVELKSGYKDWILWISFAISCFLLCVVLLGF
 IMWACQRGNIRCNICI
 SEQ ID NO:36 B/Lee/40 HA (Accession NP_056660)
 MKAIIVLLMVVTSNADRICTGITSSNSPHVVKATATQGEVNVTVGIPLTTT
 PTKSHFANLKGQTGRGKLCPCNFCTDL DVALGRPKCMGNTPSAKVSILH
 EVKPATSGCFPIIMHDKIRQLPNLLRGYENIRLSTSNVINTETAPGGPY
 KVGTS GSCPNVANGNGFFNTMAWVIPKDNNTAINPVTVEVPYICSEGED
 QITVWGFHSDDKTQMERLYGDSNPQKFTSSANGVTTHYVSQIGGFNPQTE

-continued

DEGLKQSGRIVVDYVMVQKPGKTGTIVYQRGILLPQKVWCASGRSKVIKGS
 LPLIGEADCLHEKYGGLNKS KPYTGEHAKAIGNCPIWVKTPCLKLANGTK
 YRPPAKLLKERGFFGAIAGFLEGGWEGMIAGWHGYTSHGAHGVAAADLK
 STQEAINKITKNLNYLSELEVKNLQRLSGAMNELHDEILELDEKNDDLRA
 DTISSQIELAVLLSNEGIINSEDEHLLALERKLLKMLGPSAVEIGNGCFE
 TKHKCNQTCLDRIAAGTFNAGDFSLPTFDSL NITASA SLNDDGLDNHTILL
 YYSTAASSLAVTLMIA TFIVYMVSRD NVSCSICL
 SEQ ID NO:37 C/Johannesburg/1/66 (ACCESSION
 AAW73083)
 MFFSLLLVLGLTEAEKIKICLQKQVNSSFSLHNGFGGNLYATEKRMFEL
 VKPKAGASVLNQSTWIGFGDSRTDKSNSAFPRSDVSAKTADKFRSLSGG
 SLMLSMFGPPGKVDYLYQCGCKHKVFYEGVNNWSPHAAINC YRKNWTDIKL
 NFQKN IYELASQSHCSMLVNALDKTIPLQVTTAGTAGNCNNSFLKNPALYT
 QEVKPSENKCGKENLAFFTLPTQFGTYECKLHLVASCYFIYDSKEVYNKR
 GCDN YFQVIYDSSGKVVGGLDN RVSPYTGNSGDTPTMQCDMLQLKPGRYS
 VRSSPRFLLPERSYCFDMKEKGPVTAVQS IWGKGRESDYAVDQACLSTP
 GCMLIQKQKPYIGEADDDHGDQEMRELLSGLDYEARCISQSGWVNETSPF
 TEKYLLPPKFGRCPLAAKEESIPKIPDGLL IPTSGTDTT VTTPKSRIFGI
 DDLIIGLLFVAIVEAGIGGYLLGSRKESGGGVTKESA EKGFEGIGNDIQI
 LKSSINIAIEKLNDRISHDEQAIRDLTLEIENARSEALLGELGIIRALLV
 GNISIGLQESLWELASEITNRAGDLAVEVSPGCWIIDNNICDQSCQNFI F
 KFNETAPVPTIPPLDTKIDLQSDPFYWGSSSLGLAITATISLAALVISGIA
 ICRTK
 SEQ ID NO:38 B/Lee/40 HA1-1
 TTTPTKSHFANLKGQTGRGKLCPCNFCTDL DVALGRPKCMGNTPSAKVS
 ILHEVKPATSGCFPIIMHDKIRQLPNLLRGYENIRLSTSNVINTETAPG
 GPYKVGTS GSCPNVANGNGFFNTMAWVIPKDNNTAINPVTVEVPYICSE
 GEDQITVWGFHSDDKTQMERLYGDSNPQKFTSSANGVTTHYVSQIGGFNP
 QTEDEGLKQSGRIVVDYVMVQKPGKTGTIVYQRGILLPQKVWCASGRSKVI
 KGS LPLIGEADCLHEKYGGLNKS KPYTGEHAKAIGNCPIWVK
 SEQ ID NO:39 B/Lee/40 HA1-2
 KGTQTRGKLCPCNFCTDL DVALGRPKCMGNTPSAKVSILHEVKPATSGC
 FPIIMHDKIRQLPNLLRGYENIRLSTSNVINTETAPGGPYKVGTS GSCP
 NVANGNGFFNTMAWVIPKDNNTAINPVTVEVPYICSEGEDQITVWGFHS
 DDKTQMERLYGDSNPQKFTSSANGVTTHYVSQIGGFNPQTEDEGLKQSGR
 IVVDYVMVQKPGKTGTIVYQRGILLPQKVWCASGRSKVIK
 SEQ ID NO:40 A/Aichi/2/68 HA1-1
 QSSSTGKICNNPHRILDGIDCTLIDALLGDPHCDVFQNETWDLFVERSKA
 FSNCYPYDVPDYASLRSLVASSGTLEFITEGFTWTGVTQNGGSNACKRGP
 GSGFFSRLNWLTKSGSTYPVLNVTMPNNDNFDKLYIWGIHHPSTNQEQTS
 LYVQASGRVTVSTRRSQQTIIPNIGSRPWVRLSSRISYWTIVKPGDVL

-continued

VINSNGNLIAPRGYFKMRTGKSSIMRSDAPIDTCISECITPNGSIPNDKP
FQNVNKITYGACPKYVK

SEQ ID NO:41 A/Aichi/2/68 HA1-2
HRILDGIDCTLIDALLGDPHCDVFQNETWDLFVERSKAFSNCYPYDVPDY
ASLRSLVASSGTLEFITEGFTWTGVTQNGGSNACKRGPGSGFFSRLNWL
KSGSTYPVLNVTMPNNDNFDKLYIWGIHHPSTNQEQTSLYVQASGRVTVS
TRRSQQTIIIPNIGSRPWRGLSSRISIYWTIVKPGDVLVINSNGNLIAPR
GYFKMRTGKSSIMRSD

SEQ ID NO:42 B/Malaysia/2506/2004 HA (Accession
ISDN126672)
IVLLMVVTSNADRICTGITSSNSPHVVKATQGEVNVTVGVIPLTTPTPKS
HFANLKGTETRGKLCPKCLNCTDLVALGRPKCTGNIPSARVSIHLEVRP
VTSGCFPIMHDRTKIRQLPNLLRGYEHIRLSTHNVINAENAPGGSYKIGT
SGSCPNTVNGNGFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQIT
VWGFHSDNEAQMAYLYGDSKPQKFTSSANGVTTHYVSQIGGFNPQTEDGG
LPQSGRIVVDYVMVQKSGKTGTITYQRGILLPQKVCASGRSKVIGSLPL
IGEADCLHEKYGGLNKSHPYTGGEHAKAIGNCPIWVKTPKLANGTKYRP
PAKLLKER

SEQ ID NO:43 B/Ohio/1/2005 HA (Accession
ISDN133312)
DRICTGITSSNSPHVVKATQGEVNVTVGVIPLTTPTPKSHFANLKGTR
GKLCPKCLNCTDLVALGRPKCTGNIPSAEVSILHEVRPVTSGCFPIMHD
RTKIRQLPNLLRGYEHIRLSTHNVINAENAPGGPYKIGTSGSCPNTVNGN
GFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQITVWGFHSDSETQ
MAKLYGDSKPQKFTSSANGVTTHYVSQIGGFNPQTEDGGLPQSGRIVVDY
MVQKSGKTGTITYQRGILLPQKVCASGRSKVIGSLPLIGEADCLHEKY
GGLNKSHPYTGGEHAKAIGNCPIWVKTPKLANGTKYRPPAKLLKERGF

SEQ ID NO:44 B/Shanghai/361/2002 HA (Accession
ISDN80784)
DRICTGITSSNSPHVVKATQGEVNVTVGVIPLTTPTPKSHFANLKGTR
GKLCPCDLNCTDLVALGRPMCVGTTPSAKASILHEVRPVTSGCFPIMHD
RTKIRQLPNLLRGYENIRLSTQNVDAEKALGGPYRLGTSGSCPNTATSKS
GFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQITVWGFHSDDKTQM
KNLYGDSNPQKFTSSANGVTTHYVSQIGGFDPQTEDGGLPQSGRIVVDY
VQKPGKTGTIVYQRGVLLPQKVCASGRSKVIGSLPLIGEADCLHEKY
GGLNKSHPYTGGEHAKAIGHCPIWVKTPKLANGTKYRP

SEQ ID NO:45 B/Malaysia/2506/2004 HA1-1
TTTTPTKSHFANLKGTETRGKLCPKCLNCTDLVALGRPKCTGNIPSARVS
ILHEVRPVTSGCFPIMHDRTKIRQLPNLLRGYEHIRLSTHNVINAENAPG
GSYKIGTSGSCPNTVNGNGFFATMAWAVPKNDNNKTATNSLTIEVPYICT
EGEDQITVWGFHSDNEAQMAYLYGDSKPQKFTSSANGVTTHYVSQIGGF
NQTEDGGLPQSGRIVVDYVMVQKSGKTGTITYQRGILLPQKVCASGRSKV
TKGSLPLIGEADCLHEKYGGLNKSHPYTGGEHAKAIGNCPIWVK

-continued

SEQ ID NO:46 B/Malaysia/2506/2004 HA1-2
KGTETRGKLCPKCLNCTDLVALGRPKCTGNIPSARVSIHLEVRPVTSGC
FPIMHDRTKIRQLPNLLRGYEHIRLSTHNVINAENAPGGSYKIGTSGSCP
NVTNGNGFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQITVWGFH
SDNEAQMAYLYGDSKPQKFTSSANGVTTHYVSQIGGFNPQTEDGGLPQSG
RIVVDYVMVQKSGKTGTITYQRGILLPQKVCASGRSKVIG

SEQ ID NO:47 B/Ohio/1/2005 HA1-1
TTTTPTKSHFANLKGTRGKLCPKCLNCTDLVALGRPKCTGNIPSAEVS
ILHEVRPVTSGCFPIMHDRTKIRQLPNLLRGYEHIRLSTHNVINAENAPG
GPYKIGTSGSCPNTVNGNGFFATMAWAVPKNDNNKTATNSLTIEVPYICT
EGEDQITVWGFHSDSETQMAKLYGDSKPQKFTSSANGVTTHYVSQIGGF
NQTEDGGLPQSGRIVVDYVMVQKSGKTGTITYQRGILLPQKVCASGRSKV
IKGSLPLIGEADCLHEKYGGLNKSHPYTGGEHAKAIGNCPIWVK

SEQ ID NO:48 B/Ohio/1/2005 HA1-2
KGTETRGKLCPKCLNCTDLVALGRPKCTGNIPSAEVSILHEVRPVTSGC
FPIMHDRTKIRQLPNLLRGYEHIRLSTHNVINAENAPGGPYKIGTSGSCP
NVTNGNGFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQITVWGFH
SDSETQMAKLYGDSKPQKFTSSANGVTTHYVSQIGGFNPQTEDGGLPQSG
RIVVDYVMVQKSGKTGTITYQRGILLPQKVCASGRSKVIG

SEQ ID NO:49 B/Shanghai/361/2002 HA1-1
TTTTPTKSHFANLKGTRGKLCPCDLNCTDLVALGRPMCVGTTPSAKAS
ILHEVRPVTSGCFPIMHDRTKIRQLPNLLRGYENIRLSTQNVDAEKALG
GPYRLGTSGSCPNTATSKSGFFATMAWAVPKNDNNKTATNSLTIEVPYICT
GEDQITVWGFHSDDKTQMKNLYGDSNPQKFTSSANGVTTHYVSQIGGF
QTEDGGLPQSGRIVVDYVMVQKPGKTGTIVYQRGVLLPQKVCASGRSKV
KGSPLPLIGEADCLHEKYGGLNKSHPYTGGEHAKAIGHCPIWVK

SEQ ID NO:50 B/Shanghai/361/2002 HA1-2
KGTETRGKLCPCDLNCTDLVALGRPMCVGTTPSAKASILHEVRPVTSGC
FPIMHDRTKIRQLPNLLRGYENIRLSTQNVDAEKALGGPYRLGTSGSCP
NATSKSGFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQITVWGFH
DDKTQMKNLYGDSNPQKFTSSANGVTTHYVSQIGGFDPQTEDGGLPQSGR
IVVDYVMVQKPGKTGTIVYQRGVLLPQKVCASGRSKVIG

SEQ ID NO:51 STF2.blp
ATGGCACAAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGCGGATT
GCTAACCGTTTCACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGCGCAGACTCAGTTCAACG

-continued

GCGTGAAAGTCCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTGGCGC
TGGCGCAACTAAAACCACAATGCCTGCTGGTGGCAGAACTAAAACAGAAG
TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
GCCTTAATTGCTGGCGCGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
ACAGGAGCAATTAAAGCTAAAACCTACAAGTTATACTGCTGTCAGCGCAC
TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
TCGTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA
AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
GCTCTGATCTGGGTGGGTACAAAACCGTTTCAACTCTGCTATCACCAC
CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
TTCCGACTACGCGACCGAAGTTTCCAACATGTCTGCGCGCAGATTCTGC
AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGAGAAC
GTGCTGAGCCTGTTACGT
SEQ ID NO:52 STF2.SG
ATGGCACAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
CCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAACGGCGGTTAGTCTGCT
AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
GCGTGAAAGTCCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT

-continued

GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
TGGCGCAACTAAAACCACAATGCCTGCTGGTGGCAGCAACTAAAACAGAAG
TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
GCCTTAATTGCTGGCGCGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
ACAGGAGCAATTAAAGCTAAAACCTACAAGTTATACTGCTGCTGACGGCAC
TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
TCGTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCGGCTGGTCAT
GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA
AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
GCTCTGATCTGGGTGGGTACAAAACCGTTTCAACTCTGCTATCACCAC
CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
TTCCGACTACGCGACCGAAGTTTCCAACATGTCTGCGCGCAGATTCTGC
AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGAGAAC
GTGCTGTCTCTGTTACGTAGCGGCAGTGGTAGCGGATCC
SEQ ID NO:53 STF2.HA1-1 PR8
ATGGCACAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
CCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAACGGCGGTTAGTCTGCT
AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
GCGTGAAAGTCCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
TGGCGCAACTAAAACCACAATGCCTGCTGGTGGCAGCAACTAAAACAGAAG
TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
GCCTTAATTGCTGGCGCGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
ACAGGAGCAATTAAAGCTAAAACCACAAGTTATACTGCTGCTGACGGCAC

-continued

TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTTCAT
 GATTTCAAAGCACAAACCAGAGCTGGCGGAAGCAGCCGCTAAAACCCGGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGCAGAAC
 GTGCTGTCTCTGTTACGTAGCGGAGTGGTAGCGGATCCTCTCATAACGG
 TAAACTGTGTCGTCTGAAAGGTATTGCACCACTGCAGCTGGGCAAAATGCA
 ACATCGCGGGTTGGCTGCTGGGTAAACCTGAATGTGACCCGCTGCTGCCG
 GTTCGTTCTCGGAGCTACATTGTTGAAACCCGAACTCCGAAAACCGTAT
 CTGCTACCCGGGCGACTTTATTGACTATGAAGAACTGCGTGAGCAGCTGT
 CTTCGCTGAGCAGCTTTGAACGCTTCGAAATCTTCCCGAAGGAAAGCTCC
 TGGCCGAACCAACACTAACGGCGTGACGGCGGCTTGCTCCCACGAAGG
 CAAATCTTCCTTTTATCGTAACCTGCTGTGGCTGACTGAAAAGGAAGGTT
 CCTACCCAAAACCTGAAAAACAGCTATGTTAACAAAAAGGGTAAAGAAGTC
 CTGGTGCTGTGGGGCATCCACCACCCGCCGAATCCAAGGAACAGCAGAA
 TCTGTATCAGAACGAAAACGCATACGTTTCTGTGCTTACTTCCAACATA
 ACCGTCGTTTCACTCCGGAATCGCGGAACGTCCGAAAGTACGCGACCAG
 GCTGGCCGTATGAACTACTACTGGACCTGCTGAAACCGGTGACACTAT
 TATCTTCGAAGCTAACGGTAACCTGATCGCACCAATGTACGCTTTCGCAC
 TGTCTCGTGGTTTCGGTTCCGGCATTATCACCAGCAACGCGTCTATGCAC
 GAATGCAACACGAAATGTCAGACGCCGCTGGGCGCAATTAATAGCAGCCT
 GCCGTACCAGAACATCCACCCGGTGACTATCGGCGAATGCCCGAAGTATG
 TTCGTTAATAG

SEQ ID NO:54 STF2.HA1-1.his PR8

ATGGCACAAGTAATCAACACTAACAGTCTGTCTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTTCAGTCTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA

-continued

TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGCGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGCTGATGCCGCCAAAAAT
 GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
 TGGCGCAACTAAAACCACAATGCCTGCTGGTGCGACAACATAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
 GCCTTAATGTGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGCGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCCGCAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCACAAGTTTACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTTCAT
 GATTTCAAAGCACAAACCAGAGCTGGCGGAAGCAGCCGCTAAAACCCGGA
 AAACCCGCTGCAAAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGCAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGCAGAAC
 GTGCTGTCTCTGTTACGTAGCGGCAGTGGTAGCGGATCCTCTCATAACGG
 TAAACTGTGCTGCTGTAAGAGGTATTGCACCACTGCAGCTGGGCAAAATGCA
 ACATCGCGGGTTGGCTGCTGGGTAACCTGAATGTGACCCGCTGCTGCCG
 GTTCGTTCTCGGAGCTACATTGTTGAAACCCCGAACTCCGAAAACGGTAT
 CTGCTACCCGGGCGACTTTATTGACTATGAAGAACTGCGTGAGCAGCTGT
 CTTCGCTGAGCAGCTTTGAACGCTTCGAAATCTTCCCGAAGGAAAGCTCC
 TGGCCGAACCAACACTAACGGCGTGACGGCGGCTTGCTCCCACGAAGG
 CAAATCTTCCTTTTATCGTAACCTGCTGTGGCTGACTGAAAAGGAAGGTT
 CCTACCCAAAACCTGAAAAACAGCTATGTTAACAAAAAGGGTAAAGAAGTC
 CTGGTGCTGTGGGGCATCCACCACCCGCCGAATCCAAGGAACAGCAGAA
 TCTGTATCAGAACGAAAACGCATACGTTTCTGTGCTTACTTCCAACATA
 ACCGTCGTTTCACTCCGGAATCGCGGAACGTCCGAAAGTACGCGACCAG
 GCTGGCCGTATGAACTACTACTGGACCTGCTGAAACCGGGTGACACTAT
 TATCTTCGAAGCTAACGGTAACCTGATCGCACCAATGTACGCTTTCGCAC
 TGTCTCGTGGTTTCGGTTCCGGCATTATCACCAGCAACGCGTCTATGCAC
 GAATGCAACACGAAATGTCAGACGCCGCTGGGCGCAATTAATAGCAGCCT
 GCCGTACCAGAACATCCACCCGGTGACTATCGGCGAATGCCCGAAGTATG
 TTCGTCATCACCATCATCACCATTAAATAG

SEQ ID NO:55 STF2.HA1-2 PR8

ATGGCACAAGTAATCAACACTAACAGTCTGTCTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT

-continued

CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTCACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTT CAGTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGTCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
 TGGCGCAACTAAAACCACAATGCCTGTGTGGTGCAGCAACTAAAACAGAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAAC TACAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAAACACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACC AAC
 CTGGGCAATACCGTTAAACAATCTGTCTGAAGCGGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTTACGTTCTGGTTCTGGTTCTGTTCTAAAGGTATTGC
 TCCACTGCAACTGGGTAAATGCAATATTGCGGGTTGGCTGTGGGCAACC
 CGGAATGCGATCCGCTGCTGCCGTTCCGTTCTGGAGCTATATTGTTGAA
 ACTCCGAACTCTGAGAACGGCATCTGCTATCCAGGTGATTTTATTGACTA
 TGAGGAACTGCGTGAACAAC TGTCTTCCGTGTCTTCTTTGAACGTTTCG
 AGATTTTTCCTAAAGAATCTTCTTGGCCGAACCATAACACTAATGGTGT
 ACCGCTGCGTGTCTCATGAAGGTAAATCTAGCTTTTACC GCAACCTGCT
 GTGGCTGACCGAGAAAGAAGGTTCTTACC GAAACTGAAAACAGCTACG
 TAAACAAAAAGGCAAGGAAGTTCTGTCTGTGGGTATCCACCATCCG
 CCGAACAGCAAGGAACAGCAGAACTCTGTATCAGAACGAAAACGCATACGT
 ATCTGTGTACTTCTAATAACACCGCGTTTACCCCTGAAATCGCGG

-continued

AACGTCCGAAAGTGCCTGACCAGGCAGGCCGCATGAACTATTACTGGACC
 CTGCTGAAGCCGGGTGATACTATCATCTTCGAAGCGAACGGTAACCTGAT
 CGCCCCGATGTACGCGTTCGCTCTGAGCCGTGGCTTCGGCTCTGCTATCA
 TTACGTCTTAATAA
 SEQ ID NO:56 STF2.HA1-2mut PR8
 ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTCACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTT CAGTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGTCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
 TGGCGCAACTAAAACCACAATGCCTGTGTGGTGCAGCAACTAAAACAGAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAAC CACAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAAACACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACC AAC
 CTGGGCAATACCGTTAAACAATCTGTCTGAAGCGGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTTACGTTCTGGTTCTGGTTCTGTTCTAAAGGTATTGC
 TCCACTGCAACTGGGTAAATGCAATATTGCGGGTTGGCTGTGGGCAACC
 CGGAATGCGATCCGCTGCTGCCGTTCCGTTCTGGAGCTATATTGTTGAA
 ACTCCGAACTCTGAGAACGGCATCTGCTATCCAGGTGATTTTATTGACTA
 TGAGGAACTGCGTGAACAAC TGTCTTCCGTGTCTTCTTTGAACGTTTCG
 AGATTTTTCCTAAAGAATCTTCTTGGCCGAACCATAACACTAATGGTGT
 ACCGCTGCGTGTCTCATGAAGGTAAATCTAGCTTTTACC GCAACCTGCT
 GTGGCTGACCGAGAAAGAAGGTTCTTACC GAAACTGAAAACAGCTACG
 TAAACAAAAAGGCAAGGAAGTTCTGTCTGTGGGTATCCACCATCCG
 CCGAACAGCAAGGAACAGCAGAACTCTGTATCAGAACGAAAACGCATACGT
 ATCTGTGTACTTCTAATAACACCGCGTTTACCCCTGAAATCGCGG

-continued

TGAGGAAC T GCGTGAACAAC TGTCTTCG TGTCTTCCTTTGAACGTTTCG
 AGATTTTTCCTAAAGAATCTTCTTGGCCGAACCATAACTAATGGTGT
 ACCGCTGCGTGTCTCATGAAGGTAAATCTAGCTTTTACCGCAACCTGCT
 GTGGCTGACCGAGAAAGAAGGTTCTTACCCGAAACTGAAAAACAGCTACG
 TAAACAAAAAGGGCAAGGAAGTTCTGGTCTGTGGGTATCCACCATCCG
 CCGAACAGCAAGGAACAGCAGAATCTGTATCAGAACGAAACGCATACGT
 ATCTGTTGTTACTTCTAATACAACCGCCGTTTACCCCTGAAATCGCGG
 AACGTCCGAAAGTGCGTGACCGAGCGAGCCGCATGAATATTACTGGACC
 CTGCTGAAGCCGGTGATACTATCATCTTCGAAGCGAACGGTAACCTGAT
 CGCCCGGATGTACGCGTCTGCTCTGAGCCGTGGCTTCGGCTCTGGTATCA
 TTACGTCTTAATAA

SEQ ID NO:57 STF2.HA1-3 PR8

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGTGACCCAGAATAA
 CCTGAACAAATCCCGATCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGGAAAGACGATGCGGCAGGTACGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTTCACTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTGGCGG
 TGGCGCAACTAAAACCAATGCCGTGCTGGTGGCACAAC TAAACAGAAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAAC TACAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTCAACTCTGCTATCACCAAC

-continued

CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTTACGTTCTGGTAGCGGTTCTGGCTCTAATCTGAAAA
 TGGTATCTGTTACCCGGGTGATTTTCATCGATTATGAGGAAC T GCGTGAAC
 AGCTGTCTAGCGTGAGCTCCTTTGAACGCTTCGAAATCTTCCCGAAGGAA
 TCTAGCTGGCCGAACCATAACACGAACGGTGTGACCGCTGCTTGCTCCCA
 CGAAGGTAAGAGCTCCTTCTACCGCAATCTGCTGTGGCTGACTGAAAAAG
 AAGGTAGCTACCCGAAACTGAAAAATCTTACGTCAACAAGAAAGGCAAG
 GAAGTGCTGGTTCTGTGGGGCATTACCAACCCACCGAACAGCAAAGAGCA
 ACAGAACCTGTACCAAATGAGAAGCTTACGTTTCTGTGTGACTTCTA
 ACTACAATCGTCGCTTTACCCCTGAAATCGCGGAGCGTCCAAAAGTGCCT
 GACCAGGCTGGTCTGATGAAC TACTATTGGACCTGCTGAAACCGGGCGA
 CACCATTATCTTTGAAGCGAACGGTAACCTGATCGCGCCTATGTACGCT
 TCGCTCTGTCTCGTGGCTAATAA

SEQ ID NO:58 STF2.HA1-3mut PR8

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTTCACTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTGGCGG
 TGGCGCAACTAAAACCAATGCCGTGCTGGTGGCACAAC TAAACAGAAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAAC TACAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTCAACTCTGCTATCACCAAC

-continued

TCGTTACTATCGACGGTAAAACTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACCAGAGCTGGCGGAAGCAGCCGCTAAAAACCACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACGTTTCAACTCTGCTATCACCAAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGCAAC
 GTGCTGTCTCTGTTACGTAGCGGAGTGGTAGCGGATCCAACCTGAAAA
 TGAATCTGTTACCCGGGTGATTTTCATCGATAAAGAGGAAGTGGTGAAC
 AGCTGTCTAGCGTGAGCTCCTTTGAACGCTTCGAAATCTTCCCAGGAA
 TCTAGCTGGCCGAACCATAACACGAACGGTGTGACCGCTGCTTGTCCCA
 CGAAGGTAAGAGCTCCTTCTACCGCAATCTGCTGTGGCTGACTGAAAAAG
 AAGGTAGCTACCCGAAACTGAAAAATCTTACGTCAACAAGAAAGGCAAG
 GAAGTGCTGGTTCTGTGGGCATTACACCCACCGAACAGCAAAGAGCA
 ACAGAACCTGTACCAAATGAGAACGCTTACGTTTCTGTTGTGACTTCTA
 ACTACAATCGTCGCTTTACCCCTGAAATCGCGGAGCGTCCAAAAGTGCCT
 GACCAGGCTGGTCGTATGAACACTATTTGGACCTGCTGAAACCGGGCGA
 CACCATTATCTTTGAAGCGAACGGTAACCTGATCGCGCTATGTACGCGG
 CGGCTCTGTCTCTGTGGCTAATAA

SEQ ID NO:59 HA1-1 PR8

ATGTCTCATAACGGTAAACTGTGTCGTCTGAAAGGTATTGCACCACTGCA
 GCTGGGCAAATGCAACATCGCGGTTGGCTGCTGGTAACCTGAATGTG
 ACCCGCTGCTGCCGGTTCTGTTCTGGAGCTACATTGTTGAAACCCGAAC
 TCCGAAAACGGTATCTGCTACCCGGGCGACTTTATTGACTATGAAGAACT
 GCGTGAGCAGCTGTCTTCCGTGAGCAGCTTGAACGCTTCGAAATCTTCC
 CGAAGGAAAGCTCCTGGCCGAACCACAACACTAACGGCGTGACGGCGGT
 TGCTCCACGAAGGCAAATCTTCTTTATCGTAACCTGCTGTGGCTGAC
 TGAAGGAAGGTTCTTACCCAAAAGTGAAGAAAGCTATGTTAACAAAA
 AGGGTAAAGAAGTCTGCTGTGGGCATCCACCACCCGCCGAACCTC
 AAGGAACAGCAGAATCTGTATCAGAACGAAACGCATACGTTTCTGTCTG
 TACTTCCAACATAAACCGTCGTTTCACTCCGAAATCGCGAACGTCCGA
 AAGTACGCGACCGGCTGGCCGTATGAACTACTACTGGACCTGCTGAAA
 CCGGGTGACACTATTATCTTCGAAGCTAACGGTAACCTGATCGACCAAT
 GTACGCTTTTCGACTGTCTGCTGGTTTCCGGTCCGGAATTATCACCAGCA
 ACGCGTCTATGCACGAATGCAACAGAAATGTGAGCGCCGCTGGGCGCA
 ATTAATAGCAGCCTGCCGTACCAGAACATCCACCGGTGACTATCGGCGA
 ATGCCCCGAAGTATGTTTCGTTCATCACCATCATCACCAT

-continued

SEQ ID NO:60 HA1-1.his PR8

ATGTCTCATAACGGTAAACTGTGTCGTCTGAAAGGTATTGCACCACTGCA
 GCTGGGCAAATGCAACATCGCGGTTGGCTGCTGGGTAACCTGAATGTG
 ACCCGCTGCTGCCGGTTCTGTTCTGGAGCTACATTGTTGAAACCCCGAAC
 TCCGAAAACGGTATCTGCTACCCGGGCGACTTTATTGACTATGAAGAACT
 GCGTGAGCAGCTGTCTTCCGTGAGCAGCTTTGAACGCTTCGAAATCTTCC
 CGAAGGAAAGCTCCTGGCCGAACCACAACACTAACGGCGTGACGGCGGCT
 TGCTCCACGAAGGCAATCTTCTTTTATCGTAACCTGCTGTGGCTGACT
 GAAAAAGGAAGGTTCTTACCCAAAAGTGAAGAAAGCTATGTTAACAAAA
 AGGGTAAAGAAGTCTGCTGTGTGGGGCATCCACCACCCGCCGAACCTC
 AAGGAACAGCAGAATCTGTATCAGAACGAAACGCATACGTTTCTGTCTG
 TACTTCCAACATAAACCGTCGTTTCACTCCGAAATCGCGAACGTCCGA
 AAGTACGCGACCGAGGCTGGCCGTATGAACTACTACTGGACCTGCTGAAA
 CCGGGTGACACTATTATCTTCAAGCTAACGGTAACCTGATCGCACCAAT
 GTACGCTTTCGCACTGTCTCGTGGTTTCGGTTCCGGCATTATCACCAGCA
 ACGCGTCTATGCACGAATGCAACACGAAATGTGACGCGCGCTGGGCGCA
 ATTAATAGCAGCCTGCCGTACCAGAACATCCACCCGGTACTATCGGCGA
 ATGCCCCGAAGTATGTTTCGTTCATCACCATCATCACCAT

SEQ ID NO:61 HA1-2.his PR8

ATGAAAGGTATTGCTCCACTGCAACTGGGTAAATGCAATATTGCGGGTTG
 GCTGTGGGCAACCCGGAATGCGATCCGCTGCTGCCGGTCCGTTCTGGA
 GCTATATTGTTGAAACTCCGAACCTCTGAGAACGGCATCTGCTATCCAGGT
 GATTTCAATGACTATGAGGAACTGCGTGAACAACCTGCTTCCGTGCTTTC
 CTTTGAACGTTTCGAGATTTTCTTAAAGAATCTTCTTGGCCGAACCATA
 ACACATAATGGTGTTACCGCTGCGTGCTCTCATGAAGGTAAATCTAGCTTT
 TACCGCAACCTGCTGTGGCTGACCGAGAAAGAGGTTCTTACCCGAAACT
 GAAAAACAGCTACGTAAACAAAAGGGCAAGGAAGTTCTGGTCTGTGGG
 GTATCCACCATCCGCCGAACAGCAAGGAACAGCAGAATCTGTATCAGAAC
 GAAAACGCATACGTATCTGTTGTTACTTCTAACTACAACCGCGTTTCAC
 CCCTGAAATCGCGGAACGTCCGAAAGTGCGTGACAGGCGAGCCGCATGA
 ACTATTACTGGACCTGCTGAAGCCGGTGATACTATCATCTTCGAAGCG
 AACGTAACCTGATCGCCCGATGTACGCGTTCTGCTGAGCCGTGGCTT
 CGGCTCTGGTATCATTACGTCTCATCACCATCATCACCAT

SEQ ID NO:62 CRM.HA1-2

ATGGGTGCTGATGATGCTGTTGATTCCTCCAAAAGCTTCGTTATGGAAAA
 TTTCTCTTCTTATCACGGCACCAACCGGGTTATGTGGATTCTATCCAAA
 AAGGCATCCAGAAACCGAAGTCCGGTACGACGGGTAACATGATGACGAT
 TGGAAAGGTTTTTACTCCACCGATAATAAATATGACGCGGCTGGCTACTC
 TGTTGACAACGAAAATCCACTGTCTGGTAAGGCTGGCGGTGCTGTAAAGG
 TAACGTATCCTGGCCTGACCAAGGTGCTGGCACTGAAGGTGGATAACGCT

-continued

GAAACCATCAAAAAGGAGCTGGGCCTGAGCCTGACTGAACCGCTGATGGA
GCAGGTCCGTACCGAGGAATTCATCAAACGCTTCGGTGATGGCGCTCCC
GTGTTGTGCTGTCCCTGCCGTTTCGAGAAGGCTCTTCTCTGTGCAATAT
ATCAACAACCTGGGAACAGGCTAAAGCTCTGAGCGTCGAACTGGAAATTAA
CTTTGAGACCCGTGGCAAGCGTGTCAGGACGCGATGTACGAATATATGG
CTCAGGCTTGTGCCGTAACCGTGTTCGCCGTTCCGTGGTTCCTCTCTG
TCTTGCATCAACCTGGATTGGGACGTTATCCGTGATAAGACCAAAACCAA
AATTGAAAGCCTGAAGGAACAGGTCCGATCAAAAACAAATGTCTGAAT
CTCCGAACAAAACCGTGTCCGAGGAAAAAGCGAAACAGTATCTGGAAGAA
TTCCACCAGACTGCCCTGGAACATCCTGAAGTGTCCGAACCTGAAGACTGT
AACCAGCACTAACCCGGTGTTCGAGGCGCAAACTACGCCGCGTGGGCGG
TAAAGCTTGGCGAGGTTATTGATAGCGAAACCGCAGATAACCTGGAAGAA
ACGACCGCAGCTCTGTCTATCCTGCCGGTATCGGTTCCGTTATGGGTAT
TGCGGACGGCGCTGTGCACCACAACACGGAAGAAATCGTCGCACAGTCTA
TCGCGCTGTCTTCTCTGATGGTTGCTCAGGCAATTCAGTGGTAGGTGAA
CTGGTGGACATTGGCTTTGCGGCGTACAACCTCGTCGAAAGCATATCAA
CCTGTTCCAGGTTGTACACAACCTTTACAACCGTCCGCGCTACAGCCCTG
GCCACAAAACCAACCGTTTCTGCACGACGGTTATGCGGTGTCTGGAAC
ACGGTTGAAGACTCTATCATTCGTACCGGCTTTCAGGGCGAGTCCGGCCA
CGATATCAAAATTACTGCAGAAAACACTCCGCTGCCGATCGCTGGCGTTC
TGCTGCCGACCATCCCGGTAAGCTGGATGTGAACAAAAGCAAAACCCAC
ATCTCTGTTAAGGTCGTAATAATTCGCATGCGCTGTCTGTATCGACGG
TGATGTTACCTTCTGCCGTCGAAATCTCCAGTCTACGTGGGCAACGGTG
TTCATGCCAACCTGCACGTGGCGTTCATCGTAGCTCTAGCGAAAAATC
CACTCCAACGAAATCTCTAGCGATTCTATCGGTGTTCTGGGTTATCAGAA
AACGGTGGATCATACGAAAGTCAATTCTAACTGAGCCTGTTCTTCGAAA
TCAAATCTAGCGGCTCTGGATCCGGTTCAAAGGCATCGCGCCGCTGCAG
CTGGGTAATGTAACATTGCGGGCTGGCTGCTGGGTAATCCGGAATGCGA
TCCGCTGCTGCCGGTCCGTAGCTGGTCTTACATTGTTGAAACTCCGAAC
CTGAGAAATGGCATCTGTACCCGGGCGATTTTATCGACTATGAAGAACTG
CGTGAACAGCTGTCTTCCGTTTCTTCTTTGAACGTTTCGAAATCTTCCC
GAAAGAAAGCAGCTGGCCGAATCACAATACGAACGGTGTACTGCTGCGT
GTTCTCATGAAGTAAATCCAGCTTCTACCGTAACCTGCTGTGGCTGACC
GAAAAAGAGGGTTCTTATCTCTAACTGAAAAACAGTACGTTAACAAGAA
GGGCAAGAAGTGTGGTGTGTGGGGTATCCATCACCCCTCCGAACCTTA
AAGAACAACAGAATCTGTATCAGAACGAAAACGCTTACGTTTCCGTGGTG
ACCTCTAACTATAACCGTCTGTTTACCCCGGAGATTGCTGAACGTCCGAA
AGTGCGCGATCAGGCTGGCCGTATGAAGTACTATTGGACCTGTGTAAC
CGGGCGATACCATCATTTTCAAGCTAACGGCAACCTGATTGCTCCGATG

-continued

TATGCGTTTGCTCTGTCTCGTGGCTTCGGCTCTGGTATTATTACGTCTTA
ATAA
SEQ ID NO:63 LTB.HA1-2
ATGAATAAGGTTAAGTTCTACGTACTGTTTACCGCGCTGCTGTCTTCTCT
GTGCGCGCATGGTGTCCGAGTCTATTACTGAACTGTGCTCTGAATACC
ACAACACCAGATCTATACTATCAACGATAAAATCCTGAGCTATACCGAA
TCTATGGCAGGCAACCGGAAATGGTTATCATTACCTTTAAAAGCGGCGC
CACCTTTCAAGTGGAAGTTCCGGGCTCTCAGCATATTGACTCTCAAAAAA
AAGCGATCGAACGTATGAAAGATACTCTGCGCATTACCTACCTGACCGAA
ACCAAAATCGATAAACTGTGCGTATGGAACAATAAGACTCCTAACTCTAT
CGCAGCTATTTCTATGGAAGTCTGGTAGCGGATCCGGTTCATAAGGCA
TCGCGCCGCTGCAGCTGGGTAAATGTAACATTGCGGGCTGGCTGTGGGT
AATCCGGAATGCGATCCGCTGCTGCCGGTCCGTAGCTGGTCTTACATTGT
TGAAACTCCGAACCTCTGAGAATGGCATCTGTACCCGGGCGATTTTATCG
ACTATGAAGAACTGCGTGAACAGCTGTCTTCCGTTTCTTCTTTGAACGT
TTCGAAATCTTCCGAAAGAAAGCAGCTGGCCGAATCACAATACGAACGG
TGTTACTGCTGCGTGTCTCATGAAGGTAATCCAGCTTCTACCGTAAC
TGCTGTGGCTGACCGAAAAAGAGGGTTCCTTATCCTAAACTGAAAAACAGC
TACGTTAACAAAAAGGGCAAGAGTGCTGGTGTGTTGGGTATCCATCA
CCCTCCGAACCTCTAAAGAACACAGAATCTGTATCAGAACGAAACGCTT
ACGTTTCCGTGGTGACCTCTAACTATAACCGTCTGTTTACCCCGGAGATT
GCTGAACGTCCGAAAGTGCGCATCAGGCTGGCCGTATGAACACTATTG
GACCTGTGTAACCGGGCGATACCATCATTTTCGAAGCTAACGGCAACC
TGATTGCTCCGATGTATGCGTTTGTCTGTCTGCTGGCTTCGGCTCTGGT
ATTATTACGTCTTAATAA
SEQ ID NO:64 STF2.HA1-1 VN
ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTGTCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGGTATCCGGCCAGACTCAGTTCAACG
GCGTGAAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAGTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG

-continued

TACGGCTTCTGTAACCGGTGGTGCAGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAACCACAATGCCTGCTGGTGCACAACTAAACAGAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAACCACAAGTTATACTGCTGCTGACGGCAC
 TACCAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAACCAGAAG
 TCGTTACTATCGACGGTAAACCTTACAATGCCAGCAAGCGCTGGTCAT
 GATTTCAAAGCACAACCAGAGCTGGCGGAAGCAGCGCTAAACCACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAC
 CTGGGAATACCGTAAACAATCTGTCTGAAGCGGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGCCCGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGAGCCTGTTAGCGGAGAAGAAACACAATGGCAACTGTGTGATCT
 GGATGGTGTGAAACCGCTGATTCTGCGCGATTGCTCTGTGGCAGGCTGGC
 TGCTGGGCAACCCATGTGTGACGAATTCATTAACGTTCCGGAATGGTCT
 TACATTGTTGAAAAAGCTAACCCGTGTCAACGATCTGTGTTACCCTGGTGA
 CTTTAACGATTACGAAGAACTGAAGCACCTGCTGTCTCGTATCAATCACT
 TCGAGAAAATCCAGATCATCCGAAATCCTCCTGGAGCTCCACGAAGCT
 TCTCTGGGCGTATCCTCCGCGTGCCGTACAGGGCAATCCTCTTTCTT
 TCGTAACGTTGTTTGGCTGATCAAGAAAAACTCCACCTACCCGACGATCA
 AGCGTAGCTATAATAACCAACCAGGAAGACCTGCTGGTTCTGTGGGGC
 ATCCACCATCCAAACGATGTGCGGAACAGACCAAGCTGTACCAGAACCC
 GACCACCTACATCAGCGTGGGCACCTCTACGCTGAACCAGCGTCTGGTAC
 CGCGTATCGCAACCCGACGCAAGGTAACGGTCAAAGCGGCGCATGGAA
 TTTTCTGGACCATCCTGAAACCGAACGACGCAATCAACTTCGAATCTAA
 CGGCAATTTTCATCGCTCCGGAGTATGCGTACAAAATCGTAAAGAAAGGTG
 ATAGCACTATCATGAAATCCGAGCTGGAATATGGCAACTGTAAACCCAAA
 TGCCAGACCCCGATGGGTGCAATCAACTCCTCCATGCCGTTTCACAACAT
 TCACCCGCTGACTATCGCGCAATGTCCGAAATACGTTAAATAGTAA

SEQ ID NO:65 FOR PRIMER

AGTGGCTGAGCCTGTTAGCGGAGAAGAAACACAATGGCAAACCTGTG

SEQ ID NO:66 REV PRIMER

AGTCGCTCAGCTTACTATTTAACGTATTTCCGACATTCGCCGATAGTCAG

-continued

SEQ ID NO:67 HA0s VN
 GTGCTGAGCCTGTTACGTCAGATTTGTATCGGCTACCACGCAAAACAACCTC
 TACCGAGCAAGTTGATACCATCATGGAGAAAAACGTGACCGTTACTCACG
 CGCAGGACATCCTGGAGAAGAAACACAATGGCAAACTGTGTGATCTGGAT
 GGTGTGAAACCGCTGATTCTGCGCGATTGCTCTGTGGCAGGCTGGCTGCT
 GGGCAACCCATGTGTGACGAATTCATTAACGTTCCGGAATGGTCTTACA
 TTGTTGAAAAAGCTAACCCGTGTCAACGATCTGTGTTACCCTGGTGACTTT
 AACGATTACGAAGAACTGAAGCACCTGCTGTCTCGTATCAATCACTTCGA
 GAAAATCCAGATCATCCCGAAATCCTCCTGGAGCTCCACGAAGCTTCTC
 TGGGCGTATCCTCCGCGTGCCCGTACCAGGGCAAAATCCTCTTTCTTTCGT
 AACGTTGTTTGGCTGATCAAGAAAAACTCCACCTACCCGACGATCAAGCG
 TAGCTATAATAACACCAACCAGGAAGACCTGCTGGTTCTGTGGGGCATCC
 ACCATCCAAACGATGTGCGGAACAGACCAAGCTGTACCAGAACCCGACC
 ACCTACATCAGCGTGGGCACCTCTACGCTGAACCAGCGTCTGGTACCGGC
 TATCGCAACCCGACGCAAGGTAACGGTCAAAGCGGCGCATGGAATTTT
 TCTGGACCATCCTGAAACCGAACGACGCAATCAACTTCGAATCTAACGGC
 AATTTTCATCGCTCCGGAGTATGCGTACAAAATCGTAAAGAAAGGTGATAG
 CACTATCATGAAATCCGAGCTGGAATATGGCAACTGTAAACCCAAATGCC
 AGACCCCGATGGGTGCAATCAACTCCTCCATGCCGTTTCACAACATTAC
 CCGCTGACTATCGCGCAATGTCCGAAATACGTTAAATCCAATCGTCTGGT
 TCTGGCTACCGGTCTGCGTAACTCCCCACAGCGTGAAACGTCGCTGAAGA
 AACGTGGTCTGTTTGGCGGATCGCTGGTTTCATCGAGGGCGGCTGGCAG
 GGTATGGTTGATGGCTGGTACGGTTATCATCATTCCAATGAACAGGGTTC
 CGGCTACGCCGACGATAAAGAAAGCACTCAGAAAGCAATTGATGGCGTAA
 CTAACAAAGTAAATCTATCATTGATAAAATGAACACCCAGTTTCGAGGGC
 GTTGGTCTGTGAGTTCAACAACCTGGAACCCGTATCGAAACCTGAACAA
 GAAAATGGAAGACGGTTTCTGGATGTGTGGACTTACAATGTGAACTGC
 TGGTGTGATGGAAAACGAGCGTACCCTGGACTTCCACGACAGCAACGTC
 AAAAATCTGTATGACAAAGTCCGCTGTCAGCTGCGTGATAACGCTAAAGA
 GCTGGGTAATGGCTGCTTTCGAGTTCTATCACAATGCGACAACGAATGCA
 TGGAATCTGTCCGCAACGGCACTTACGATTATCCGAGTACTCCGAAGAA
 GCGCGCTGAAACGCGAAGAGATCTCCGGTGTGAAGCTGGAGTCTATTGG
 CATCTACCAGATCCTGTCCATCTACAGCACCTAGTAAGCTGAGCGCCTAC
 GCAGC

SEQ ID NO:68 STF2.HA1-2 VN

ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
 GCTAACCGTTTCACCGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG

-continued

AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTCACTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCCTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTGCGGC
 TGGCGCAACTAAAACCAATGCTGCTGGTGCAGAACTAAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACCAACGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATACCAAC
 CTGGGCAATACCGTAAACAATCTGCTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGAGCCTGTTAGCAGGTGTGAAACCGCTGATTCTGCGCGATTGCTC
 TGTGGCAGGCTGGCTGCTGGGCAACCCATATGTGTGACGAATTCATTAACG
 TTCCGGAATGGTCTTACATTGTTGAAAAAGCTAACCTGTCAACGATCTG
 TGTACCCCTGGTGACTTTAACGATTACGAAGAACTGAAGCACCTGCTGTC
 TCGTATCAATCACTTCGAGALAATCCAGATCATCCCGAAATCCTCCTGGA
 GCTCCCACGAAGCTTCTCTGGCGGTATCCTCCGCGTGCCGTACCAAGGC
 AAATCCTCTTTCTTCGTGAACGTTGTTTGGCTGATCAAGAAAACTCCAC
 CTACCCGACGATCAAGCGTAGCTATAATAACCAACCAAGAGACCTGC
 TGGTCTGTGGGGCATCCACCATCCAAACGATGCTGCGGAACAGACCAAG
 CTGTACCAGAACCCGACCACTACATCAGCGTGGGCACCTCTACGCTGAA
 CCAGCGTCTGGTACCGGTATCGCAACCCGAGCAAGTAAACGGTCAAG
 CGGCCGATGGAATTTTCTGTGACCATCCTGAAACCGAACGAACGAATC

-continued

AACTTCGAATCTAACGGCAATTTTCATCGCTCCGGAGTATGCGTACAAAAT
 CGTAAAGAAAGGTGATAGCACTATCATGAAATCCGAGTAGTAA
 SEQ ID NO:69 FOR PRIMER
 AGTCGCTGAGCCTGTAGCAGGTGTGAAACCGTGATTCTGCGCGATTG
 SEQ ID NO:70 REV PRIMER
 TGACGCTCAGCTTACTACTCGGATTTCATGATAGTGCTATCACCTTTC
 SEQ ID NO:71 STF2.HA1-2mut VN
 ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTCACCGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTCACTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCCTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTGCGGC
 TGGCGCAACTAAAACCAATGCCTGCTGGTGCAGAACTAAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACCAACGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATACCAAC
 CTGGGCAATACCGTAAACAATCTGCTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGAGCCTGTTAGCAGGTGTGAAACCGCTGATTCTGCGCGATTGCTC
 TGTGGCAGGCTGGCTGCTGGGCAACCCATATGTGTGACGAATTCATTAACG
 TTCCGGAATGGTCTTACATTGTTGAAAAAGCTAACCTGTCAACGATCTG
 TGTACCCCTGGTGACTTTAACGATTACGAAGAACTGAAGCACCTGCTGTC
 TCGTATCAATCACTTCGAGALAATCCAGATCATCCCGAAATCCTCCTGGA
 GCTCCCACGAAGCTTCTCTGGCGGTATCCTCCGCGTGCCGTACCAAGGC
 AAATCCTCTTTCTTCGTGAACGTTGTTTGGCTGATCAAGAAAACTCCAC
 CTACCCGACGATCAAGCGTAGCTATAATAACCAACCAAGAGACCTGC
 TGGTCTGTGGGGCATCCACCATCCAAACGATGCTGCGGAACAGACCAAG
 CTGTACCAGAACCCGACCACTACATCAGCGTGGGCACCTCTACGCTGAA
 CCAGCGTCTGGTACCGGTATCGCAACCCGAGCAAGTAAACGGTCAAG
 CGGCCGATGGAATTTTCTGTGACCATCCTGAAACCGAACGAACGAATC

-continued

TGTTACCTTGGTGACTTTAACGATTACGAAGAACTGAAGCACCTGCTGTC
TCGTATCAATCACTTCGAGAAAAATCCAGATCATCCCGAAATCCTCCTGGA
GCTCCACGAAGCTTCTCTGGGCGTATCCTCCGCGTGCCCGTACCAGGGC
AAATCCTCTTTCTTTCGTAACGTTGTTTGGCTGATCAAGAAAACTCCAC
CTACCCGACGATCAAGCGTAGCTATAATAACACCAACCAGGAAGACCTGC
TGGTTCTGTGGGGCATCCACCATCCAAACGATGCTGCGGAACAGACCAAG
CTGTACCAGAACCCGACCACCTACATCAGCGTGGGCACCTCTACGCTGAA
CCAGCGTCTGGTACCGCGTATCGCAACCCGAGCAAGTAAACGGTCAAA
GCGGCGCATGGAATTTTCTGGACCATCTGAAACCGAACGACGCAATC
AACTTCGAATCTAACGCAATTTTCATCGCTCCGGAGTATGCGTACAAAAAT
CGTAAAGAAAGGTGATAGCACTATCATGAAATCCGAGTAGTAA

SEQ ID NO:72 HA1-2mut VN

GGTGCGAAACCGCTGCTCTGCGCGATTGCTCTGTGCGAGGCTGGCTGCT
GGGCAACCCATATGTGTGACGAATTCATTAACGTTCCGGAATGGTCTGATA
TTGCCGAAAAAGCTAACCCGTCAACGATCTGTGTTACCCGTGGTGACTTT
AACGATTACGAAGAACTGAAGCACCTGCTGCTCTCGTATCAATCACTTCGA
GAAAATCCAGATCATCCGAAATCCTCTGGAGCTCCACGAAGCTTCTC
TGGGCGTATCCTCCGCGTGCCCGTACCAGGGCAAATCCTCTTTCTTTCTG
AACGTTGTTTGGCTGATCAAGAAAAATCCACCTACCCGACGATCAAGCG
TAGCTATAATAACACCAACCAGGAAGACCTGCTGGTTCTGTGGGGCATCC
ACCATCCAAACGATGCTGCGGAACAGACCAAGCTGTACCAGAACCCGACC
ACCTACATCAGCGTGGGCACTCTACGCTGAACCAGCGTCTGGTACCGCG
TATCGCAACCCGAGCAAGGTAACCGGTCAAAGCGGCCGATGGAATTTT
TCTGGACCATCCTGAAACCGAACGACGCAATCAACTTCGAATCTAACGGC
AATTTTCATCGCTCCGGAGTATGCGTACAAAATCGTAAAGAAAGGTGATAG
CACTATCATGAAATCCGAGTAGTAA

SEQ ID NO:73 STF2.HA1-1 IND

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGATAAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
GCTAACCGTTTACCAGCAACATCAAAGGCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTTCACTCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
GCGTGAAAGTCCCTGGCGCAGGACAACCCCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA

-continued

TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAACCGGTGGTGCGGTTAAATTTGACGCAGATAATAACA
AGTACTTTGTACTATTGGTGGCTTTACTGGTGCTGATGCCGCCAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
TGGCGCAACTAAAACCACAATGCCTGCTGGTGCGACAACATAAACAGAAG
TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
GCCTTAATGTGCTGGCGCGGTTGACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGCGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
ACAGGAGCAATTAAAGCTAAAACCACAAGTTTACTGCTGCTGACGGCAC
TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCTAT
GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACCCCGA
AAACCCGCTGCAGAAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
TTCCGACTACGCGACCGAAGTTTCCAACATGCTCTCGCGCGCAGATTTTGC
AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
GTGCTGTCTCTGTTAGCGGAGAAAAACGCATAACCGTAAACTGTGCGACCT
GGACGGCGTGAAGCCGCTGATCCTGCGTGACTGTTCTGTTGCTGGCTGGC
TGCTGGGCAACCCGATGTGCGATGAGTTTCATCAACGTTCCGGAGTGGTCT
TATATTGTTGAAAAAGCGAACCAGCAATGACCTGTGCTACCCGGGCTC
TTTTAATGACTACGAGGAGCTGAAACATCTGCTGTCTCGCATCAACCATT
TCGAAAAAATTCAGATCATCCCGAAATCTTCTGGAGCGATCACGAAGCT
TCTTCCGCGTATCTTCTGCATGCCCTTACCTGGGTTCCTCCGCTTTTCTT
CCGTAACGTGGTTTGGCTGATCAAGAAGAACTCTACGTACCCGACCATCA
AGAAATCCTATAACAACACTAACAGGAAGATCTGCTGGTGCTGTGGGT
ATCCATCATCCAAACGATGCAGCCGAACAGACCCGCTGTACCAGAATCC
GACCACGTACATCTCTATCGGCACTTCTACCCCTGAATCAACGCTGGTGC
CGAAAATCGCAACTCGCAGCAAAGTCAACGCCAATCTGGTTCGATGGAG
TTTTTCTGGACCATCCTGAAACCGAACGACGCGATTAACTTCGAAAGCAA
CGGCAACTTCATTGACCAGGAATATGCTTACAAGATCGTTAAAAAGGGTG
ATTCTGCGATCATGAAAGCGAGCTGGAATACGGCAACTGCAACACTAAA
TGCCAGACCCCGATGGGTGCAATCAATTCAGCATGCCATTTTACAACAT
CCACCCGCTGACCATCGGCGAGTGTCCAAAATACGTTAAATAATAG

SEQ ID NO:74 FOR PRIMER

AGTGGCTGAGCCTGTTAGCGGAGAAAACGCATAACGGTAAACTGTG

SEQ ID NO:75 HA0s IND

GCTGAGCCTGTTACGTGATCAAAATCTGCATCGGTTACCACGCAAAACACT
CCACTGAACAAGTGGATACGATCATGGAAAGAACGTGACCGTGACCCAC

-continued

GCTCAAGATATCCTGGAGAAAACGCATAACGGTAAACTGTGCGACCTGGA
 CGGCGTGAAGCCGCTGATCTGCGTGAAGTCTTCTGTTGCTGGCTGGCTGC
 TGGGCAACCCGATGTGCGATGAGTTCATCAACGTTCCGGAGTGGTCTTAT
 ATTGTTGAAAAAGCGAACCCGACCAATGACCTGTGCTACCCGGGCTCTTT
 TAATGACTACGAGGAGCTGAAACATCTGCTGTCTCGCATCAACCATTTCTG
 AAAAAATTGAGATCATCCGAAATCTTCTGGAGCGATCAGGAAGCTTCT
 TCCGGCGTATCTTCTGCATGCCCTTACCTGGGTTCCCGCTTTTCTTCCG
 TAACGTGGTTTGGCTGATCAAGAAGAACTCTACGTACCCGACCATCAAGA
 AATCTATAACAACACTAACCAGGAAGATCTGCTGGTGTCTGTTGGGTATC
 CATCATCCAACGATGCGAGCCGAACAGACCCGCTGTACCAGAATCCGAC
 CACGTACATCTCTATCGGCACTTCTACCTGAATCAACGCTGCTGCGCA
 AATCGCAACTCGCAGCAAGTCAACGGCCAATCTGGTCGCATGGAGTTT
 TTCTGGACCATCTGAAACCGAACGACGCGATTAAC TTCGAAAGCAACGG
 CAAC TTCAATGACCCGGAATATGCTTACAAGATCGTTAAAAAGGGTGATT
 CTGCGATCATGAAAGCGAGCTGGAATACGGCAACTGCAACACTAAATGC
 CAGACCCGATGGGTGCAATCAATTCCAGCATGCCATTTCAACATCCA
 CCCGCTGACCATCGGCGAGTGTCCAAATACGTTAAATCCAATCGTCTGG
 TGCTGGCAACGGGCTGCGCAACTCTCCGCAACGTGAGTCTCGTCGCAAA
 AAGCGTGGCCTGTTTGGCGCAATTGCGGGCTTCATCGAGGCGGCTGGCA
 GGGTATGGTTGATGGTTGGTACGGTTATCACCATTCCAACGAGCAGGGCT
 CTGGTTACGCTGCGGACAAGAGTCCACGCAGAAAGCGATCGATGGCGTC
 ACCAACAAGTGAAGTCTATCATCGACAAGATGAACACTCAGTTCGAGGC
 TGTGGGCGGTGAATTCAACAATCTGGAACGTCGCATCGAAAACCTGAACA
 AAAAGATGGAAGACGGTTTCTGGACGTATGGACCTATAACGCAGAGCTG
 CTGGTTCTGATGGAAAACGAACGTACTCTGACTTCCACGATTCTAACGT
 TAAAAACCTGTACGACAAAGTTCGTCTGCAGCTGCGCGATAATGCAAAAAG
 AACTGGGTAACGGCTGCTTTGAAATTTTACCACAAATGTGACAACGAATGC
 ATGGAAGCATCCGTAACGGTACCTACAATTACCCACAGTACTCCGAAGA
 AGCGCGTCTGAAACGTGAAGAAATCTCCGGTGTAACCTGGAATCCATTG
 GCAGTATCAGATTCTGTCTATCTACTCCAGGTAGCGTCTCTCTGGCG
 CTGGCAATTATGATGGCGGCTGTCTCTGTGGATGTGCTCTAACGGCTC
 TCTGCAGTGGCGCATCTGCATCTAATAGGCTGAGC
 SEQ ID NO:76 STF2.HA1-2 IND
 ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAGACGATGCGGCAGGTACGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTTCACTCTGCT

-continued

AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGCGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAAACCACAATGCCTGCTGGTGCAGCAACTAAAACAGAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
 GCCTTAATGTCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCACAAGTTTACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCAT
 GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACCACCGA
 AAACCCGCTGCAGAAAATTGATGCCCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGAGATTTTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGAGCCCTGTAGCAGGTGTGAAACCGCTGATTCTGCGCGATTGTTT
 TGTTGCTGGCTGGCTGCTGGGCAACCCGATGTGCGATGAGTTTCATCAACG
 TTCCGGAGTGGTCTTATATTGTTGAAAAAGCGAACCCGACCAATGACCTG
 TGCTACCCGGGCTCTTTTAAATGACTACGAGGAGCTGAAACATCTGCTGTC
 TCGCATCAACCATTTTCGAAAAAATTCAGATCATCCGAAATCTTCTGGA
 GCGATCACGAAGCTTCTCCGCGTATCTTCTGCATGCCCTTACCTGGGT
 TCCCCGTCTTTCTTCCGTAACGTGGTTTGGCTGATCAAGAAGAACTCTAC
 GTACCCGACCATCAAGAAATCTATAACAACACTAACCGGAAGATCTGC
 TGGTGTGTGGGATATCCATCATCCAAACGATGCAGCCGAACAGACCCGC
 CTGTACCAGAATCCGACCACTACATCTCTATCGGCACTTCTACCTGAA
 TCAACGCTGGTGGCGAAAATCGCAACTCGCAGCAAGTCAACGGCCAAT
 CTGGTCTGCATGGAGTTTCTTCTGGACCATCCTGAAACCGAACGACGCGATT
 AACTTCGAAAGCAACGGCAACTTCATTGCACCGGAATATGCTTACAAGAT
 CGTTAAAAAGGGTGATTCTGCGATCATGAAAGCGAGTAATAG

-continued

SEQ ID NO:77 REV PRIMER
 AGTCGCTCAGCCTATTACTCGCTTTTCATGATCGCAGAATCAC

SEQ ID NO:78 STF2.HA1-2mut IND
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA

CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT

CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT

GCTAACCGTTTACC CGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA

CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG

AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTTTCAGTCTGCT

AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGTCGAAATCAC

CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG

GCGTGAAAGTCC TGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC

AACGACGGTGAACACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC

CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA

CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA

TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG

TACGGCTTCTGTAAACCGGTGGTGCAGTTAAATTTGACGCAGATAATAACA

AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT

GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC

TGGCGCAACTAAAACCAATGCCTGCTGGTGCAGCAACTAAAACAGAAG

TACAGGAGTTAAAGATACACCGGCAGTTGTTTTCAGCAGATGCTAAAAAT

GCCTTAATTGCTGGCGCGTGTACGCTACCGATGCTAATGGCGCTGAGTT

GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT

ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG

ACAGGAGCAATTAAAGCTAAAACCAAGTTATACTGCTGCTGACGGCAC

TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG

TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCAT

GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA

AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC

GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC

CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA

TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC

AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC

GTGCTGTCTCTGTAGCGGGCGCAAGCCGCTGTCTCTGCGTGACTGTTC

TGTTGCTGGCTGGCTGTGGGCAACCCGATGTGCGATGAGTTTCATCAACG

TTCCGGAGTGGTCTGATATTGCGGAAAAGCGAACCCGACCAATGACCTG

TGCTACCCGGGCTCTTTTAACTACTACGAGGAGCTGAACATCTGCTGTC

TCGCATCAACCATTTCGAAAAAATTCAGATCATCCCGAAATCTTCTCGGA

GCGATCACGAAGCTTCTTCCGGCGTATCTTCTGCATGCCCTTACCTGGGT

-continued

TCCCCGCTCTTTCTTCCGTAACGTGGTTTGGCTGATCAAGAAGAACTCTAC

GTACCCGACCATCAAGAAATCCTATAACAACACTAACCCAGGAAGATCTGC

TGGTGCTGTGGGGTATCCATCATCAAACGATGCAGCCGAACAGACCCGC

CTGTACCAGAATCCGACCACGTACATCTCTATCGGCACCTTCTACCCGTAA

TCAACGCCTGGTGCCGAAAATCGCAACTCGCAGCAAAGTCAACGGCCAAT

CTGGTCGATGGAGTTTTTCTGGACCATCTGAAACCGAACGACGCGATT

AACTTCGAAAGCAACGGCAACTTCATTGCACCGGAATATGCTTACAAGAT

CGTTAAAAAGGTGATTCTGCGATCATGAAAAGCGAGTAATAG

SEQ ID 79 HA1-2mut IND
 GGCGCAAAGCCGCTGTCTCTGCGTGACTGTTCTGTTGCTGGCTGGCTGCT

GGGCAACCCGATGTGCGATGAGTTTCATCAACGTTCCGGAGTGGTCTGATA

TTGCGGAAAAAGCGAACC CGACCAATGACCTGTGCTACCCGGGCTCTTTT

AATGACTACGAGGAGCTGAAACATCTGCTGTCTCGCATCAACCATTTTCGA

AAAAATTCAGATCATCCCGAAATCTTCTGAGCGATCACGAAGCTTCTT

CCGGCGTATCTTCTGCATGCCCTTACCTGGGTCCCGCTCTTCTTCCGT

AACGTGGTTTGGCTGATCAAGAAGAACTCTACGTACCCGACCATCAAGAA

ATCCTATAACAACACTAACCCAGGAAGATCTGCTGGTGTGTGGGGTATCC

ATCATCCAAACGATGCAGCCGAACAGACCCGCTGTACCGAATCCGACC

ACGTACATCTCTATCGGCACCTTCTACCCGTAATCAACGCCTGGTGCAGAA

AATCGCAACTCGCAGCAAAGTCAACGGCCAATCTGGTCGCATGGAGTTTT

TCTGGACCATCCTGAAACCGAACGACGCGATTAACTTCGAAAGCAACGGC

AACTTCATTGCACCGGAATATGCTTACAAGATCGTTAAAAAGGTGATTCT

TGCGATCATGAAAAGCGAGTAATAGGCTGAGC

SEQ ID NO:80 STF2.HA1-1 NC
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA

CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT

CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT

GCTAACCGTTTACC CGGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA

CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG

AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTTTCAGTCTGCT

AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC

CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG

GCGTGAAAGTCC TGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC

AACGACGGTGAACACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC

CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA

CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA

TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG

TACGGCTTCTGTAAACCGGTGGTGCAGTTAAATTTGACGCAGATAATAACA

AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT

GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC

-continued

TGGCGCAACTAAAACCACAATGCCTGCTGGTGCACAACTAAAACAGAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAGCTAAAACCACAAGTTATACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCGCTGGTCAT
 GATTTCAAAGCACAACCAGAGCTGGCGGAAGCAGCGCTAAAACCACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
 CTGGGAATACCGTAAACAATCTGCTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGCTCTCGCGCGAGATTTTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGAGCCTGTAGCGTCCCAACGGTAAACTGTGCTGCTGTAAGG
 CATCGCACCGCTGCAGCTGGGTAAGTGTAGCGTTGCAGGTTGGATCCTGG
 GTAACCCGGAGTGCAACTGCTGATTTCCAAGGAGAGCTGGTCTTACATT
 GTCGAAACGCCGAATCCGGAAGAACGGCACTTGTATCCGGGTTACTTCGC
 CGATTACGAAGAACTGCGTGAACAACCTGTCTTCCGTGAGCTCTTTCGAAC
 GTTTCGAGATCTTCCCGAAAGAGAGCAGCTGGCCAAACCACTGTTACC
 GGTGTGCTGCGAGCTGTTCTCACAACGGCAAGTCTTCTTTCTACCGCAA
 CCTGCTGTGGCTGACGGGTAAGAATGGCCTGTATCCGAACCTGTCTAAAT
 CTTACGTAAACAACAAAGAGAAAGAGGTGCTGGTCCGTGTTGGGCGTACAC
 CATCCACCAATATCGGCAACAGCGCGCCCTGTACCACACCGAAAACGC
 TTATGTGTCCGTGGTGAGCTCCCATACAGCCGCTGTTTACTCCGAGA
 TTGCCAAACGTCCGAAAGTTCTGTATCAGGAAGCCGTATTAATACTAC
 TGGACTCTGCTGGAGCCGGCGATACCATATTTTCGAGGCAACCGCAA
 CCTGATTGCGCCATGTTACGCGTTCGCCCTGAGCCGTGGTTTTGGCTCCG
 GTATTATCACCTCTAACCGCCCAATGGACGAATGCGACGCGAAATGCCAA
 ACGCCGAGGGCGCAATCAACAGCAGCCTGCCGTTCCAGAACGTTACCC
 GGTTACCATCGCGAATGCCCTAAATACGTGCGCTAATAG

 SEQ ID NO:81 FOR PRIMER
 AACCAAGTCCCGCAGAACGTGCTGAGCCTGTTAGCGTCCCAACGGTAA

 ACTGTGCTGCTGAAAGGCATCGCACCGCTGCAG

 SEQ ID NO:82 REV PRIMER
 GAACAGCTCAGTTCTCAGAGCTCAGCCTATTAGCGCAGTATTTAGGGC

 ATTCCCGATGTAACCGGGTGAACGTT

 SEQ ID NO:83 HA0s NC
 GACACGATCTGTATTGGTTATCATGCAACAACCTCTACTGACACTGTAGA
 TACTGTGCTGGAAAAGAACGTAACCGTTACCCACAGCGTTAACCTGCTGG

-continued

AAGATTCCCAACACGGTAACTGTGTCTGCTGAAAGGCATCGCACCGCTG
 CAGCTGGGTAACTGTAGCGTTGCAGGTTGGATCCTGGGTAAACCGGAGTG
 CGAAGTGTGATTTCCAAGGAGAGCTGGTCTTACATTGTGCAAAACGCCGA
 ATCCGGAACACGGCACTTGTATCCGGGTTACTTCGCCGATTACGAAGAA
 CTGCGTGAACAACCTGTCTTCCGTGAGCTCTTTCGAACGTTTCGAGATCTT
 CCCGAAAGAGAGCAGCTGGCCAAACCACTGTTACCGGTGTGCTGCGA
 GCTGTCTCACAACGGCAAGTCTTCTTTCTACCGCAACCTGTGTGGCTG
 ACGGGTAAGAATGGCCTGTATCCGAACCTGTCTAAATCTTACGTAAACAA
 CAAAGAGAAAGAGGTGCTGGTCTGTGGGGCGTACACCATCCACCAAATA
 TCGGCAACAGCGCGCCCTGTACCACACCGAAAACGCTTATGTGTCCGTG
 GTGAGCTCCCATACAGCCGTGTTTTACTCCGGAGATTGCCAAACGTCC
 GAAAGTTCGTGATCAGGAAGGCCGTATTAATACTACTACTGGACTCTGCTGG
 AGCCGGGCGATACCATCATTTTTCGAGGCAACGGCAACCTGATTGCGCCA
 TGGTACGCGTTCCGCCCTGAGCCGTGGTTTTGGCTCCGGTATTATCACCTC
 TAACGCGCAATGGACGAATGCGACGCGAAATGCCAAACGCCGAGGGCG
 CAATCAACAGCAGCCTGCCGTTCCAGAAGCTTACCCGGTTACCATCGGC
 GAATGCCCTAAATACGTGCGCTCCGCCAAACCTGCGCATGGTTACTGGTCT
 GCGTAACATCCCGAGCATTAGTCTCGCGGTCTGTTCCGGTGCATCGCGG
 GCTTCATTGAAGGCGGTTGGACCGGTATGGTGGATGGTTGGTACGGCTAC
 CATCACGAGAAGCAACAGGGTAGCGGTTACGCTGCCGACCAGAAATCCAC
 CCAGAAGCTATTAACGGTATCACCAACAAAGTTAACAGCGTAATTGAGA
 AGATGAACACGCGAGTTACCGCCGTAGGTAAGGAATTCAACAAGCTGGAA
 CGTCGCATGGAACCTGAACAAAAGGTGGACGACGGCTTTCTGGACAT
 CTGGACCTACAACGCTGAACTGCTGGTGTGCTGGAACCAACGTAACCC
 TGGATTTCACGACTCTAATGTTAAAAACCTGTACGAAAAGGTCAAGTCT
 CAACTGAAAAACAATGCGAAGGAATCGCAACCGCTGTTTCGAATTCTA
 CCATAAATGCAACAACGAATGCATGGAATCCGTTAAAAACGTTACCTATG
 ACTACCCCTAAATACTCCGAAGAAAGCAAACCTGAACCGCGAGAAAAATCGAT
 GGTGTAACCTGGAATCTATGGGTGTTTACCAGATCCTGGCGATCTACTC
 CACGGTAGCCAGCAGCCTGGTTCTGCTGGTTAGCCTGGGTGCAATTAGCT
 TCTGGATGTGCTCTAACGGCAGCCTGCAATGCCGCATCTGTAATAGGCTG

 AGC

 SEQ ID NO:84 STF2.HA1-2 NC
 ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACGGCGATT
 GCTAACCGTTTCACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTAGTCTGCT

-continued

AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGCAGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAAACCAATGCCTGCTGGTGCAGAACTAAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGGTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAC
 CTGGGCAATACCGTTAAACAATCTGTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGCAAG
 GTGCTGAGCCTGTTAGCGAAAGGCATCGCACCGCTGCAGCTGGGTAAGT
 TAGCGTTGCAGGTTGGATCCTGGTAACCCGGAGTGCGAAGTCTGATTT
 CCAAGGAGAGCTGGTCTTACATTGCTGAAACGCCGAATCCGGAAACGGC
 ACTTGTATCCGGTTACTTTCGCCGATTACGAAGAACTGCGTGAACAACT
 GTCTTCCGTGAGCTCTTTTGAACGTTTCGAGATCTTCCCGAAAGAGAGCA
 GCTGGCCAAACACACTGTTACCGGTGTGTCTGCGAGCTGTTCTCACAAC
 GGCAAGTCTTCTTCTACCGCAACCTGCTGTGGCTGACGGGTAAGAATGG
 CCTGTATCCGAACCTGTCTAAATCTTACGTAAACAACAAGAGAAAGAGG
 TGCTGGTCTGTGGGGCTACACCATCCACCAATATCGGCAACCGCGC
 GCCCTGTACCACACCGAAAACGCTTATGTGTCCGTGGTGAGCTCCCATTA
 CAGCCGTCGTTTTACTCCGAGATTGCCAAACGTCCGAAAGTTCGTGATC
 AGGAAGGCCGTATTAACTACTACTGGACTCTGTGGAGCCGGCGATACC
 ATCATTTTCGAGGCAAACGGCAACCTGATTGCGCCATGGTACGCGTTGCG
 CCTGAGCCGTGGTTTTGGCTCCGGTATTATCACCTCTTAATAA

-continued

SEQ ID NO:85 FOR PRIMER
 AACCAGGTCCCGCAGAAGCTGTGAGCCTGTTAGCGAAAGGCATCGCACC
 GCTGCAGCTGGGTAAGTGTAGCCTTGACAGTTGG
 SEQ ID NO:86 REV PRIMER
 GAACAGCTCAGTTCTTACGAGCTCAGCCTATTAAAGAGGTGATAATACCGG
 AGCCAAAACACGGCTCAGGGCGAAGCGGTACCA
 SEQ ID NO:87 STF2.HA1-2mut NC
 ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTCCAGCGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTGTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
 TGGCGCAACTAAAACCAATGCCTGCTGGTGCAGAACTAAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTACGAGATGCTAAAAAT
 GCCTTAATTGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGCAAG
 GTGCTGAGCCTGTTAGCGAAAGGCATCGCACCGCTGCAGCTGGGTAAGT
 TAGCGTTGCAGGTTGGATCCTGGTAACCCGGAGTGCGAAGTCTGATTT
 CCAAGGAGAGCTGGTCTTACATTGCTGAAACGCCGAATCCGGAAACGGC
 ACTTGTATCCGGTTACTTTCGCCGATTACGAAGAACTGCGTGAACAACT
 GTCTTCCGTGAGCTCTTTTGAACGTTTCGAGATCTTCCCGAAAGAGAGCA
 GCTGGCCAAACACACTGTTACCGGTGTGTCTGCGAGCTGTTCTCACAAC
 GGCAAGTCTTCTTCTACCGCAACCTGCTGTGGCTGACGGGTAAGAATGG
 CCTGTATCCGAACCTGTCTAAATCTTACGTAAACAACAAGAGAAAGAGG
 TGCTGGTCTGTGGGGCTACACCATCCACCAATATCGGCAACCGCGC
 GCCCTGTACCACACCGAAAACGCTTATGTGTCCGTGGTGAGCTCCCATTA
 CAGCCGTCGTTTTACTCCGAGATTGCCAAACGTCCGAAAGTTCGTGATC
 AGGAAGGCCGTATTAACTACTACTGGACTCTGTGGAGCCGGCGATACC
 ATCATTTTCGAGGCAAACGGCAACCTGATTGCGCCATGGTACGCGTTGCG
 CCTGAGCCGTGGTTTTGGCTCCGGTATTATCACCTCTTAATAA

-continued

ACTTGTATCCGGTTACTTCGCCGATTACGAAGAACTGCGTGAACT
 GTCTCCGTGAGCTCTTTCGAACGTTTCGAGATCTCCCGAAAGAGAGCA
 GCTGGCCAAACCACACTGTTACCGGTGTGTCTGCGAGCTGTTCTCACAAC
 GGCAAGTCTTCTTCTACCGCAACCTGCTGTGGCTGACGGGTAAGATGG
 CCTGTATCCGAACCTGTCTAAATCTTACGTAAACAACAAAGAGAAAGAGG
 TGCTGGTCTGTGGGGCTACACCATCCACCAATATCGGCAACCAGCGC
 GCCCTGTACCACACCGAAAACGCTTATGTGTCGTGAGCTCCCATTA
 CAGCCGTCGTTTTACTCCGAGATTGCCAAACGTCCGAAAGTTCGTGATC
 AGGAAGGCCGTATTAATACTACTGACTCTGCTGGAGCCGGCGATACC
 ATCATTTTCGAGGCAAACGGCAACCTGATTGCGCCATGGTACGCGTTCGC
 CCTGAGCCGTGGTTTTGGCTCCGTATTATCACCTCTTAATAG

SEQ ID NO:88 HA1-2mut NC
 GCTGAGCCTGTTAGCGAAAGCGCGGCACCGCTGCAGCTGGGTAACTGTA
 GCGTTGCAGGTTGGACTCTGGGTAACCCGAGTGCGAACTGCTGATTTCC
 AAGGAGAGCTGGTCTGATATTGCAGAAACGCCGAATCCGAAAACGGCAC
 TTGTTATCCGGGTTACTTCGCCGATTACGAAGAACTGCGTGAACAACTGT
 CTTCGCTGAGCTCTTTCGAACGTTTCGAGATCTCCCGAAAGAGAGCAGC
 TGGCCAAACCACACTGTTACCGGTGTGTCTGCGAGCTGTTCTCACAACGG
 CAAGTCTTCTTTCTACCGCAACCTGCTGTGGCTGACGGGTAAGATGGCC
 TGATATCCGAACCTGTCTAAATCTTACGTAAACAACAAAGAGAAAGAGGTG
 CTGGTCTGTGGGGCGTACACCATCCACCAATATCGGCAACCAGCGCGC
 CCTGTACCACACCGAAAACGCTTATGTGTCCGTGAGTCCCATTTACA
 GCCGTCGTTTTACTCCGAGATTGCCAAACGTCCGAAAGTTCGTGATCAG
 GAAGGCCGTATTAATACTACTGACTCTGCTGGAGCCGGCGATACCAT
 CATTTTCGAGGCAAACGGCAACCTGATTGCGCCATGGTACGCGTTCGCCC
 TGAGCCGTGGTTTTGGCTCCGGTATTATCACCTCTTAATAGCTGAGC

SEQ ID NO:89 STF2.HA1-1his (PR8)
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQVRRELAVQSA
 NSTNSQSDLDISQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLASHNGKLCRLKGIAPLQLGKCNIAWLLGNPECDPLLPVRSWSYI
 VETPNSENGICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNNHTN
 GVTAACSHGKSSFYRNLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIH

-continued

HPPNSKEQQNLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYY
 WTLKPGDTIIFEANGNLIAPMYAFALSRGFGSGIITSNASMHECNTKCQ
 TPLGAINSSLPYQNIHPVTIGECPKYVRHHHHHH

SEQ ID NO:90 STF2.HA1-2 (PR8)
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQVRRELAVQSA
 NSTNSQSDLDISQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLAKGIAPLQLGKCNIAWLLGNPECDPLLPVRSWSYIIVETPNSENG
 ICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNNHTNGVTAACSHE
 GKSSFYRNLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHPPNSKEQQ
 NLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLKPGDT
 IIFEANGNLIAPMYAFALSRGFGSGIITS

SEQ ID NO:91 STF2.RA1-2mut (PR8)
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQVRRELAVQSA
 NSTNSQSDLDISQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLAKGAAPLQLGKCNIAWLLGNPECDPLLPVRSWSDIAETPNSENG
 ICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNNHTNGVTAACSHE
 GKSSFYRNLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHPPNSKEQQ
 NLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLKPGDT
 IIFEANGNLIAPMYAFALSRGFGSGIITS

SEQ ID NO:92 STF2.HA1-3 (PR8)
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQVRRELAVQSA
 NSTNSQSDLDISQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV

-continued

SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGV DATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEEAAKT TENPLQKIDAALAQVDALRSDLGAVQNRFN SAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLANSENGICYPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNNHN
 TNGVTAACSHGKSSFYRNLLWLTEKEGSYPKLKNSYVNKKGKEVLVLWG
 IHHPNSKEQQNLYQENAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMN
 YYWTLKPGDTIIFEANGNLIAPMYAFALSRG

SEQ ID NO:93 HA1-2his (PR8)
 KGIAPLQLGKCNIAGWLLGNPECDPLLPVRSWSYIVETPNSENGICYPGD
 FIDYEELREQLSSVSSFERFEIFPKESSWPNNHNTNGVTAACSHGKSSFY
 RNLLWLTEKEGSYPKLKNSYVNKKGKEVLVLWGIHHPNSKEQQNLYQNE
 NAYVSVVTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLKPGDTIIFEAN
 GNLIAPMYAFALSRGFGSGIITSHHHHHH

SEQ ID NO:94 STF2.4xm2e
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINS AKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRRELAVQSA
 NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGV DATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEEAAKT TENPLQKIDAALAQVDALRSDLGAVQNRFN SAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLRLSLLTEVETPIRNEWGSRNSDSDPLESLLTEVETPIRNEWGSR
 SNDSDPGSSLLTEVETPIRNEWGSRNSDSDPELSLLTEVETPIRNEWG
 SRNSDSDPSR

SEQ ID NO:95 STF2.HA1-2 (VN)
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINS AKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRRELAVQSA
 NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGV DATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH

-continued

DFKAQPELAEEAAKT TENPLQKIDAALAQVDALRSDLGAVQNRFN SAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSPAQILQQAGTSVLAQANQVPQN
 VLSLLAGVKPLILRDCSVAGWLLGNPMCDEFINVPWEYSIVEKANPVNDL
 CYPGDFNDYEELKHLLSRINHFEKIQIIPKSSWSSHEASLGVS SACPYQG
 KSSFFRNVVWLKKNSTYPTIKRSYNNNTNQEDLLVLWGIHHPNDAAEQTK
 LYQNPTTYISVGTSTLNQRLVPRIATR SKVNGQSGRMEFFWTILKPNDAI
 NFESNGNFI APEYAYKIVKKGDSTIMKSE

SEQ ID NO:96 STF2
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINS AKDDAAGQAI
 ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRRELAVQSA
 NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
 ALIAGGV DATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
 TGAIKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEEAAKT TENPLQKIDAALAQVDALRSDLGAVQNRFN SAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLA

SEQ ID NO:97 STF2.Blp.wt
 ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTTC
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTCGTTGCTCA
 CCCAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACCGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCGCCGG
 TCAGGCCATCGCAAACCGCTTTACCGCCAATATCAAGGGACTGACGCAGG
 CTTTCGAGGAATGCTAACGATGGAATAAGCATCGCTCAAACCGAGGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCGTCAGGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCACCCAGCGCCTGAATGAGATTGACCGGGTGAGCGGTGAGACA
 CAGTTTAACGGCGTGAAGTACTTGCACAGGATAACACACTTACGATACA
 GGTGGGCGCCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAGCGGTACGAC
 GTGAAAGACACGCGGTCACTACCAAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGTACCGGGC
 GTACGAACGGAACCGCGTCCGTGACCGCGGTGCCGTGAAGTTCGATGCT
 GACAACAATAAGTATTTCTGCACCATTTGAGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGGTGCCACCAAGACCACCATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCGCGGTGGTTAGCGCAGA

-continued

TGCCAAAACGCGTTGATCGCCGGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAAATGTCGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTCGTTACGATCGATGGCAAGACATACAACGCCCTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGGCCGCGCGA
 AAACAACTGAGAACCCTTGCAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCCGTTCAAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCCGCGACTAGTGTCTGGCCAGGCCAATCAAGT
 CCGCAGAAATGTGCTGAGCCTACTACGAGAGTTCAGTCGTTACCCAGCCC
 AATGGCGGCCGCTC

SEQ ID NO:98 STF2.Blp.ng

ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTT
 TTACATCTATGGGCCCAAGTGATCAACACCAACTCCCTGTCCCTGTCTCA
 CTCAAAACAACCTCCAGAAATCTCAGTCTGCTCTCGGTACTGCTATCGAG
 CGTCTCTCTCCGGCCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
 ACAAGCTATCGTAAACGCTTTACTGCCAACATTAAGGCCCTGACCCAAG
 CCTCTAGGAACGCAATGATGGCATTCTATCGCTCAAAACCACGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCAACTCGCCGT
 CCAGTCCGCCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTCTGGTCAAAC
 CAGTTTAAACGGCGTCAAAGTGCTGGCTCAAGACAACACACTCACCATCCA
 AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGACTCGATGACGCCGCTATCAAGGCCCTACTGGCG
 GCACTCAGGGAACGCTCTGTGACCGGAGCGCTGTGAAGTTCGACGCC
 GATAACAACAAATACCTTCGTCACTATTGGTGGTTTCACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTGCCACTGACGGAACAGTGA
 CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGTGTGCTACTACT
 AAGACAGAGGTGCAGGAACCTCAAGGACACCCCTGCCGTGGTGTCCGCCGA
 TGCTAAGAACGCTCTCATTGTGTGGTGTGTCGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAAATGTCTACACCGACAAGAAGCGAAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGCTGATTA
 TGACGAAGCTACAGGCCCATTAAGCTAAACCACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGCTGCCAATCAGTGGGTGGAGTCGATGGA

-continued

AAAACCGAAGTGGTCACAATCGACGGAAGACCTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCCAACCCGAACCTGGCCGAGGCTGCTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG
 GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATTCTCCAACAAGCCGGCACCTCCGTCCTCGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGC

SEQ ID NO:99 Honey Bee mellitin

ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTT
 TTACATCTATGCG

SEQ ID NO:100 STF2.HA0s

ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTT
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTCTGTGCTCA
 CCCAAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACCGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCGCCGG
 TCAGGCCATCGCAAACCGCTTTACCGCCAATATCAAGGGACTGACGCAGG
 CTTCGAGGAATGCTAACGATGGAATAAGCATCGCTCAAACACGGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCGTCAAGGAGCTCGCAGT
 GCAGTCCGCCAATTTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCACCAGCGCCTGAATGAGATTGACCGGGTGAGCGGTGAGACA
 CAGTTTAAACGGCGTGAAGGTACTTGCACAGGATAACACACTTACGATACA
 GGTGGGCGCCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAAGCGTACGAC
 GTGAAAGACACGGCGGTCACTACCAAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGACGAATCAAGGCTGTACCGGCG
 GTACGAACGGAACCGGTCCTGACCGCGGTGCCGTGAAGTTGATGCT
 GACAACAATAAGTATTTCTGTCACCATTTGAGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGGTGCCACCAAGACCACCATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCGCGGTGGTTAGCGCAGA
 TGCCAAAACGCTTGATCGCCGGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAAATGTCGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTGCTTACGATCGATGGCAAGACATACAACGCTTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGGCCGCGCGA
 AAACAACTGAGAACCCTTGCAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCGTTCAAAATCGATTCAATCCGC

-continued

GATTACAAACCTGGGCAATACAGTAAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCCGCTAGTGTCTCGGCCAGGCCAATCAAGT
 CCCGCAGAATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCG
 ACACCATTTCATTTGATACCATGCAACAACCTCAACCGATCTGTTGAT
 ACCGTCTTGAGAAGAAGCTTACCGTACGCACTCGGTCAACCTATTAGA
 GGATAGCCACAACGGAAGCTGTGCCGTCTGAAAGGCATAGCGCCACTGC
 AGCTCGGCAATGTAACATCGCAGGTTGGCTCCTTGGTAACCCGAGTGC
 GACCCCTCTCCTGTATGATCTTGGAGTTATATCGTGGAGACCCCAA
 TAGCGAGAACGGAATTTGCTACCCAGGAGATTTATAGACTACGAGGAGT
 TGGCGAGCAGCTTTCGTCTGTGAGCAGCTTCGAAAGGTTTCGAGATATTC
 CCGAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCCGC
 CTGCAGTCATGAAGGAAAGAGTTTCTATCGCAACCTGCTGTGGTTGA
 CGGAGAAGAGGGCAGCTACCCCTAAGTTGAAGAACTCCTATGTGAACAAA
 AAAGGCAAGGAGGTTCTGGTGCTGTGGGGCATACACCAACCCCCCAATAG
 CAAGGAGCAGCAGAATCTGTACCAAAACGAGAATGCCTATGTGAGCGTGG
 TCACTAGTAACATAACCGTTCGGTTCACTCCCGAGATCGCCGAGCGTCCG
 AAGGTGAGGGACAGGCGAGCCGGATGAACACTACTGACCCCTATTGAA
 GCCAGGGGACACGATTATCTTCGAGGCAACGGAACCTCATAGCGCCGA
 TGTACGCCCTTCGCCCTGAGCCGTGGCTTTGGATCGGGATCATCACGTCT
 AACGCCCTCGATGCACGAATGTAATACCAATGCCAGACCCCACTGGGTGC
 TATCAACTCGTCCTTACCTATCAAAATATACATCCGGTCACCATAGGCG
 AGTGTCCCAAATATGTGATCCGCCAAGTTGCGGATGGTGACCGGCCTC
 CGCAATATTCCTAGTATTTCAGTCACGCGGCTTGTTCGCGCCATCGCTGG
 TTTTCATCGAAGCGGGTGACAGGCATGATTGATGGCTGGTATGGCTATC
 ACCACCAAGACGAGCAGGGCTCGGGCTACGCGGCTGACCAGAAGTCGACT
 CAGAATGCCATCAATGGCATCACGAATAAGGTGAACACGGTCATTGAAAA
 GATGAACATTCAATTTACAGCCGTAGGAAAAGAGTTTAAATAAACTGGAAA
 AAAGAATGGAGAATCTGAATAAGAAGGTGACGACGGATTTTGGACATC
 TGGACGTACAACGCCGAGCTGCTGGTTCTGCTGAAAAATGAGCGAACACT
 GGATTTTCATGATTCTAACGTAAAGAATCTGTACGAGAAGGTGAAGTCCC
 AACTAAAGAATAATGCCAAGGAAATCGGAAATGGATGCTTTGAGTTTAC
 CACAAGTGCAGATAATGAGTGCATGGAATCCGTGCGAAATGGTACATACGA
 TTACCCAAAGTACTCCGAAGAATCCAAGCTAAATCGCGAAAAGGTTGATG
 GTGTTAAACTTGAATCCATGGGTATTTACCAACACCATCATCACCACCAT
 TAATAG
 SEQ ID NO:101 FOR PRIMER
 AGTCGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCGACACCATT
 TGCATTGGATACCATGC

-continued

SEQ ID NO:102 REV PRIMER
 AGTCGCATGCCCTATTAATGGTGGTGATGATGGTGGTAAATACCCATG
 GATTG
 SEQ ID NO:103 HA0s PR8
 GACACCATTTCATTGGATACCATGCAAAACACTCAACCGATACTGTTGA
 TACCGTCCTTGAGAAGAAGCTTACCGTCACGCACTCGGTCAACCTATTAG
 AGGATAGCCACAACGGAAGCTGTGCCGTCTGAAAGGCATAGCGCCACTG
 CAGCTCGGCAATGTAACATCGCAGGTTGGCTCCTTGGTAACCCGGAGTG
 CGACCCCTCTCCTGTACGATCCTGGAGTTATATCGTGGAGACCCCCA
 ATAGCGAGAACGGAATTTGCTACCCAGGAGATTTATAGACTACGAGGAG
 TTGCGCGAGCAGCTTTCGTCTGTGAGCAGCTTCGAAAGGTTTCGAGATATT
 CCCGAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCCG
 CCTGCAGTCATGAAGGAAAGAGTTTCTATCGCAACCTGCTGTGGTTG
 ACGGAGAAAGAGGGCAGCTACCCCTAAGTTGAAGAACTCCTATGTGAACAA
 AAAAGGCAAGGAGGTTCTGGTGCTGTGGGGCATACACCAACCCCCCAATA
 GCAAGGAGCAGCAGAATCTGTACCAAAACGAGAATGCCTATGTGAGCGTG
 GTCCTAGTAACATAACCGTCGGTTCACTCCCGAGATCGCCGAGCGTCC
 GAAGGTGAGGGACAGGCAGGCGGATGAACACTACTGACCCCTATTGA
 AGCCAGGGGACACGATTATCTTCGAGGCAACGGAACCTCATAGCGCCG
 ATGTACGCCCTTCGCCCTGAGCCGTGGCTTTGGATCGGGATCATCACGTC
 TAACGCCCTCGATGCACGAATGTAATACCAATGCCAGACCCCACTGGGTG
 CTATCAACTCGTCCTTACCCTATCAAAATATACATCCGGTCACCATAGGC
 GAGTGTCCCAAATATGTGATCCGCCAAGTTGCGGATGGTGACCGGCCCT
 CCGCAATATTCCTAGTATTCAGTCACGCGGCTTGTTCGCGCCATCGCTG
 GTTTCATCGAAGGCGGGTGACAGGCATGATTGATGGCTGGTATGGCTAT
 CACCACCAAGACGAGCAGGGCTCGGGCTACGCGGCTGACCAGAAGTCGAC
 TCAGAATGCCATCAATGGCATCACGAATAAGGTGAACACGGTCATTGAAA
 AGATGAACATTCAATTTACAGCCGTAGGAAAAGAGTTTAAATAAACTGGAA
 AAAAGAATGGAGAATCTGAATAAGAAGGTGACGACGGATTTTGGACAT
 CTGGACGTACAACGCCGAGCTGCTGGTTCTGCTGAAAAATGAGCGAACAC
 TGGATTTTCATGATTCTAACGTAAAGAATCTGTACGAGAAGGTGAAGTCC
 CAACTAAAGAATAATGCCAAGGAAATCGGAAATGGATGCTTTGAGTTTAA
 CCACAAGTGCAGATAATGAGTGCATGGAATCCGTGCGAAATGGTACATACG
 ATTACCCAAAGTACTCCGAAGAATCCAAGCTAAATCGCGAAAAGGTTGAT
 GGTGTTAAACTTGAATCCATGGGTATTTACCAACACCATCATCACCACCA
 TTAATAG
 SEQ ID NO:104 STF2.HA1-1
 ATGAAATTCTTAGTCAACGTTGCCCTTGTGTTTATGGTCGTATACATTTT
 TTACATCTATGCGGGCCAGGTTATCAATACCAACTCCCTGCTGCTGCTCA
 CCCAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACCGCCATAGAA

-continued

CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCGCCGG
 TCAGGCCATCGCAAACCGCTTTACCGCAATATCAAGGACTGACGCAGG
 CTTTCGAGGAATGCTAACGATGGAATAAGCATCGCTCAAACCACGGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCGTCAGGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCAGCCAGCGCTGAATGAGATTGACCGGTCAGCGGTGAGACA
 CAGTTTAACGGCGTGAAGTACTTGCACAGGATAACACACTTACGATACA
 GGTGGGCGCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAAGGCGTACGAC
 GTGAAAGACACGGCGGTCACTACCAAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGCTACCGCGG
 GTACGAACGGAACCGCGTCCGTGACCGGCGGTGCCGTGAAGTTCGATGCT
 GACAACAATAAGTATTTTCGTCACCATTTGGAGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGGTGCCACCAAGACCACCATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCCGCGGTGGTTAGCGCAGA
 TGCCAAAACGCCTTGATCGCGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAATGTCGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTGCTTACGATCGATGGCAAGACATACAACGCTTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGCGCGCGGA
 AAACAACGTGAAACCGGTTGCGAGAAGATTGATCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCGGTTCAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCGGCACTAGTGTCTGGCCCAGGCCAATCAAGT
 CCCGACGAATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCA
 GCCACAACGGAAAGCTGTGCCGTCTGAAAGGCATAGCGCCACTGCAGCTC
 GGCAATGTAAACATCGCAGGTTGGCTCCTTGGTAACCCGAGTGCGACCC
 CCTCCTCCCTGTACGATCCTGGAGTTATATCGTGGAGACCCCCAATAGCG
 AGAACGGAATTTGCTACCCAGCAGATTTTCATAGACTACGAGGAGTTGCGC
 GAGCAGCTTTTCGTCTGTGAGCAGCTTCGAAAGGTTGAGATATTCCCGAA
 GGAGAGCAGTGCGCGAATCATACACTAACGCTGTGACAGCCGCTGCA
 GTCATGAAGGAAAGAGTTTATTCTATCGCAACCTGCTGTGGTTGACGGAG
 AAAGAGGGCAGCTACCTTAAGTTGAAGAACTCCTATGTGAACAAAAAGG
 CAAGGAGGTTCTGGTGCTGTGGGGCATACACCACCCCCCAATAGCAAGG

-continued

AGCAGCAGAATCTGTACCAAAACGAGAATGCCATATGTGAGCGTGGTCACT
 AGTAACTATAACCGTCGGTTCACTCCCCGAGATCGCCGAGCGTCCGAAGGT
 GAGGGACCAGGCAGGCCGGATGAACCTACTACTGGACCTATTGAAGCCAG
 GGGACACGATTATCTTCGAGGCAAACGGAAACCTCATAGCGCCGATGTAC
 GCCTTCGCGCTGAGCCGTGGCTTTGGATCGGGGATCATCACGTCTAACGC
 CTCGATGCACGAATGTAATACCAATGCCAGACCCCACTGGGTGCTATCA
 ACTCGTCCTTACCCTATCAAAATATACATCCGGTCACCATAGGCGAGTGT
 CCCAAATATGTCAGACACCATCATCACCACCATTAATAG
 SEQ ID NO:105 FOR PRIMER
 AGTCGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCAGGCCAAC
 GGAAAGCTGTGCCGTC

SEQ ID NO:106 REV PRIMER
 AGTCGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCAGGCCAAC
 CACTCGCCTATGGTGAC

SEQ ID NO:107 STF2.HA1-2
 ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTGCTGTCTCA
 CCCAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACCGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCCGCG
 TCAGGCCATCGCAAACCGCTTTACCGCCAATATCAAGGGACTGACGCAGG
 CTTTCGAGGAATGCTAACGATGGGATAAGCATCGCTCAAACCACGGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCGTCAGGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCAGCCAGCGCCTGAACGAGATTGACCGGGTGAGCGGTGAGACA
 CAGTTTAACGGCGTGAAGTACTTGCAGGATAACACACTTACGATACA
 GGTGGGCGCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAAGGCGTACGAC
 GTGAAAGACACGGCGGTCACTACCAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGCTACCGCGG
 GTACGAACGGAACCGCGTCCGTGACCGCGGTGCCGTGAAGTTGATGCT
 GACAACAATAAGTATTTCTCACCATTGGAGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGGTGCCACCAAGACCACCATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCCGCGGTGGTTAGCGCAGA
 TGCCAAAACGCGTTGATCGCGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAATGTCGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTGCTTACGATCGATGGCAAGACATACAACGCTTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGCGCGCGGA
 AAACAACGTGAAACCGGTTGCGAGAAGATTGATCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCGGTTCAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCGGCACTAGTGTCTGGCCCAGGCCAATCAAGT
 CCCGACGAATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCA
 GCCACAACGGAAAGCTGTGCCGTCTGAAAGGCATAGCGCCACTGCAGCTC
 GGCAATGTAAACATCGCAGGTTGGCTCCTTGGTAACCCGAGTGCGACCC
 CCTCCTCCCTGTACGATCCTGGAGTTATATCGTGGAGACCCCCAATAGCG
 AGAACGGAATTTGCTACCCAGCAGATTTTCATAGACTACGAGGAGTTGCGC
 GAGCAGCTTTTCGTCTGTGAGCAGCTTCGAAAGGTTGAGATATTCCCGAA
 GGAGAGCAGTGCGCGAATCATACACTAACGCTGTGACAGCCGCTGCA
 GTCATGAAGGAAAGAGTTTATTCTATCGCAACCTGCTGTGGTTGACGGAG
 AAAGAGGGCAGCTACCTTAAGTTGAAGAACTCCTATGTGAACAAAAAGG
 CAAGGAGGTTCTGGTGCTGTGGGGCATACACCACCCCCCAATAGCAAGG

-continued

CGCTGGCCACGATTTCAAAGCTCAACCCGAAGTGGCCGAGGCCGCGGCGA
 AAACAACCTGAGAACCCGTTGCAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGGCGCGTTCAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCCGCGACTAGTGTCTGGCCAGGCCAATCAAGT
 CCCGCAGAATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCA
 AAGGCATAGCGCCACTGCAGCTCGGCAAAATGTAACATCGCAGGTTGGCTC
 CTTGGTAACCCGAGTGCGACCCCTCTCTCCGTGACGATCCTGGAGTTA
 TATCGTGGAGACCCCAATAGCGAGAACGGAATTTGCTACCCAGGAGATT
 TCATAGACTACGAGGAGTTGCGCGAGCAGCTTTCGTCTGTGAGCAGCTTC
 GAGAGGTTTCAAATCTTCCCGAAGGAGAGCAGCTGGCCGAATCATAACAC
 TAACGGTGTGACAGCCGCTGCAGTCATGAAGGAAAGAGTTCAATCTATC
 GCAACCTGTGTGGTTGACGAGAGAAAGAGGCGAGCTACCCTAAGTTGAAG
 AACTCTATGTGAACAAAAAGGCAAGGAGGTTCTGTGTGTGGGGCAT
 ACACCACCCCAATAGCAAGGAGCAGCAGAATCTGTACCAAAACGAGA
 ATGCCTATGTGAGCGTGGTCACTAGTAACATATAACCGTCGGTTCACTCCC
 GAGATCGCCGAGCGTCCGAAGGTGAGGACAGGCAGGCCGATGAACATA
 CTACTGGACCTATTGAAGCCAGGGGACACGATTATCTTCGAGGCAAACG
 GAAACCTCATAGCGCCGATGTACGCGTTCCGCTGAGCCGTGGCTTTGGA
 TCGGGGATCATCAGCTCTACCATCATCACCACCATTAATAG

SEQ ID NO:108 FOR PRIMER
 AGTCGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCAAAGGCATA
 GCGCCACTGCAGCTCG

SEQ ID NO:109 REV PRIMER
 AGTCGCATGCCTATTAATGGTGGTGATGATGGTGAGACGTGATGATCCCC
 GATCCAAAGCCACGGCTCAG

SEQ ID NO:110 STF2.HA1-2mut
 ATGAAATTTCTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTT
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTCGTTGCTCA
 CCCAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACCGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCGCCGG
 TCAGGCCATCGAAACCGCTTTACCGCCAATATCAAGGACTGACGCAGG
 CTTTCGAGGAATGCTAACGATGGAATAAGCATCGCTCAAAACCGGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCGTCAGGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCACCCAGCGCTGAATGAGATTGACCGGTTGAGCGGTGAGACA
 CAGTTTAACGGCGTGAAGGTACTTGCACAGGATAACACACTTACGATACA
 GGTGGGCGCCAACGATGGTGAACCATAGACATTGATCTCAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCTGAATGTGCAAAAGGCGTACGAC

-continued

GTGAAAGACACGGCGGTCACTACCAAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGTACCGGCG
 GTACGAACGGAACCGCGTCCGTGACCGCGGTGCCCTGAAGTTGATGCT
 GACAACAATAAGTATTTCTGTCACCATTTGGAGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGGTGCCACCAAGACCACATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCGCGGTGGTTAGCGCAGA
 TGCCAAAAACGCGTTGATCGCCGCGGAGTGGATGCAACTGATGTAAATG
 GTGCGGAGCTGGTTAAATGTCTGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGCGCAGCGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTCGTTACGATCGATGGCAAGACATACAACGCCTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAAGTGGCCGAGGCCGCGCGA
 AAACAACCTGAGAACCCGTTGCAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCCGTTCAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCCGCGACTAGTGTCTTGGCCAGGCCAATCAAGT
 CCCGCGAATGTGTGAGCCTGCTGGCTTCTGGATCTGGATCTGGTTCTA
 AAGGCGCTGCCCTCTGCAACTCGGCAAGTGAATATTGCGGGGTGGCTG
 TTGGGCAACCCAGAATGTGACCCCTCTGCGCCGTCGTTCTTGGTCTGA
 CATCGCTGAAACACCTAACTCCGAGAACGGCATCTGTTACCCGGGCGACT
 TCATTGACTACGAGGAGCTCCGCGAACAGCTGTCTTCTGTCTCTTCTTC
 GAACGTTTCGAAATTTTCCCAAAGGAATCTCTGCGCAACCAATAACAC
 CAACGGAGTGACCGCGCTGTAGCCATGAGGGCAAGTCTTCTTTCTACC
 GTAATCTGCTGTGGCTGACTGAAAAAGAGGGTTCTTATCCAAACTGAAG
 AACTCTTATGTCAACAAGAAGGGCAAAGAGGTCCTGGTGTGTGGGGTAT
 CCACCACCCCCCAACTCCAAGGAGCAGAGAATCTGTACCAAAACGAAA
 ATGCTTACGTGTCTGTCTGACATCTAACTACAACCGCGCTTCACGCC
 GAAATCGCCGAGCGTCCCAAAGTGCAGATCAGGCCGGAAGGATGAACATA
 CTACTGGACCTGCTCAAAACCGGAGATACCATCATATTGGAAGCAATG
 GCAATCTCATCGCTCCCATGTATGCTTTGCTCTCTCTCGGGATTGCGT
 TCCGGCATAATCACTAGCCACCACCACCACCACCACTAATAG

SEQ ID NO:111 HA1-2mut
 AAGGCGCTGCCCTCTGCAACTCGGCAAGTGAATATTGCGGGGTGGCTG
 TTGGGCAACCCAGAATGTGACCCCTCTGCGCCGTCGTTCTTGGTCTGA
 CATCGCTGAAACACCTAACTCCGAGAACGGCATCTGTTACCCGGGCGACT
 TCATTGACTACGAGGAGCTCCGCGAACAGCTGTCTTCTGTCTCTTCTTC

-continued

GAACGTTTCGAAATTTTCCCAAAGGAATCCTCCTGGCCAAACCATAACAC
 CAACGGAGTGACCGCCGCTGTAGCCATGAGGGCAAGTCTTCTTTCTACC
 GTAATCTGCTGTGGCTGACTGAAAAAGAGGGTTCTTATCCCAAACCTGAAG
 AACTCTTATGTCAACAAGAAGGGCAAAGAGGTCTGGTGCTGTGGGGTAT
 CCACCACCCCCCAACTCCAAGGAGCAGCAGAATCTGTACAAAACGAAA
 ATGCTTACGTGTCTGTCTGTGACATCTAACTACAACCGGCGCTTCACGCCC
 GAAATCGCCGAGCGTCCCAAAGTGC GCGATCAGGCCGGAAGGATGAAC TA
 CTACTGGACCTGCTCAAACCCGAGATACCATCATATTCGAAGCCAATG
 GCAATCTCATCGTCCCATGTATGCTTTTCGCTCTCTCTCGCGGATTCGGT
 TCCGGCATAATCACTAGCCACCACCACCACCACCACCTAATAG
 SEQ ID NO:112 STF2.HA1-3
 ATGAAATTCCTAGTCAACGTTGCCCTTGTTTATGGTCGTATACATTC
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTCGTTGCTCA
 CCCAAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCGCGCG
 TCAGGCCATCGAAACCGCTTTACCGCCAATATCAAGGACTGACGCAAG
 CTTGAGGAATGCTAACGATGGGATAAGCATCGCTCAAACCACGGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCTCAGGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCACCAGCGCTGAACGAGATTGACCGGGTGAGCGGTGAGACA
 CAGTTTAACGGCGTGAAGGTACTTGCGCAGGATAACACACTTACGATACA
 GGTGGGCGCCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAAGGCGTACGAC
 GTGAAAGACACGCGGTCACTACCAAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGCTACCGCG
 GTACGAACGGAACCGCTCCGTGACCGCGGTGCCGTGAAGTTCGATGCT
 GACAACAATAAGTATTTCTGTCACCATTTGAGGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGTTGCCACCAAGACCACCATGCCAGCGCGGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCCGCGTGGTTAGCGCAGA
 TGCCAAAACGCGTTGATCGCCGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAATGTGCTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTGCTTACGATCGATGGCAAGACATACAACGCTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGGCCGCGCGA
 AAACAACTGAGAACCCTTGCGAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCCGTTCAAATCGATTCAATTCGCG

-continued

GATTACAAACCTGGGCAATACAGTAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCCGGCACTAGTGTCTGGCCAGGCCAATCAAGT
 CCCGCAAGATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCA
 ATAGCGAGAACGGAATTTGCTACCCAGGAGATTTCATAGACTACGAGGAG
 TTGCGCGAGCAGCTTTCGTCTGTGAGCAGCTTCGAGAGGTTGAAATCTT
 CCCGAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCCG
 CCTGCAGTCATGAAGGAAAGAGTTCATTCTATCGCAACCTGCTGTGGTTG
 ACGGAGAAGGAGGGCAGCTACCTAAGCTGAAGAACTCCTATGTGAACAA
 AAAAGGCAAGGAGGTTCTGGTGCTGTGGGCATACACACCCCCCAATA
 GCAAGGAGCAGCAGAATCTGTACAAAACGAGAATGCCTATGTGAGCGTG
 GTCCTAGTAACATAAACCGTCGGTTCCTCCGAGATCGCCGAGCGCTCC
 GAAGGTGAGGGACCGGAGGCGGATGAACACTACTACTGGACCTATTGA
 AGCCAGGGGACACGATTATCTTCGAGGCAACCGGAAACCTCATAGCGCG
 ATGTACGCCCTTCGCCCTGAGCCGTGGCCACCATCATCACCACCATTAATA
 G
 SEQ ID NO:113 FOR PRIMER
 AGTCGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCAATAGCGAG
 AACGGAATTTGCTACC
 SEQ ID NO:114 REV PRIMER
 AGTCGCATGCCATTAATGGTGGTGATGATGGTGCCACGGCTCAGGGCG
 AAGG
 SEQ ID NO:115 STF2.HA1-3mut
 ATGAAATTCCTAGTCAACGTTGCCCTTGTTTATGGTCGTATACATTC
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTCGTTGCTCA
 CCCAAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCCGCG
 TCAGGCCATCGCAAACCGCTTTACCGCCAATATCAAGGACTGACGAGG
 CTTGAGGAATGCTAACGATGGAATAAGCATCGCTCAAACCACGGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCTCAGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCACCAGCGCTGAATGAGATTGACCGGGTGAGCGGTGAGACA
 CAGTTTAACGGCGTGAAGGTACTTGACAGGATAACACACTTACGATACA
 GGTGGGCGCCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAAGGCGTACGAC
 GTGAAAGACACGCGGTCACTACCAAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGCTACCGCG
 GTACGAACGGAACCGCTCCGTGACCGCGGTGCCGTGAAGTTCGATGCT
 GACAACAATAAGTATTTCTGTCACCATTTGAGGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGTTGCCACCAAGACCACCATGCCAGCGCGGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCCGCGTGGTTAGCGCAGA
 TGCCAAAACGCGTTGATCGCCGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAATGTGCTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTGCTTACGATCGATGGCAAGACATACAACGCTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGGCCGCGCGA
 AAACAACTGAGAACCCTTGCGAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCCGTTCAAATCGATTCAATTCGCG

-continued

CGCTGGCCGCTGGTGCCACCAAGACCACCATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCGCGGTGGTTAGCGCAGA
 TGCCAAAAACGCGTTGATCGCCGGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAAATGTCGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGCTGATTA
 CGATGAGGCGACGGGAGCTATTAAGGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGGTTGACGGG
 AAGACAGAGGTGCTTACGATCGATGGCAAGACATACAACGCTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACTGGCCGAGGCCGCGCGA
 AAACAACTGAGAACCCTTGCAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCGGTTCAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGAGGCCGCGACTAGTGCTCTGGCCAGGCCAATCAAGT
 CCCGAGAAATGTGCTGAGCCTGCTGGCTTCTGGATCTGGATCTGGTTCTA
 ACTCCGAGAACGAAATCTGTTACCCGGCGGACTTCATTGACAAAGAGGAG
 CTCCGCGAACAGCTGTCTTCTGTCTCTTCTTCGAACGTTTCGAAATTTT
 CCCAAAGGAATCTCTCGGCCAAACCATAACACCAACGGAGTGACCGCCG
 CCTGTAGCCATGAGGGCAAGTCTTCTTCTACCGTAATCTGCTGTGGCTG
 ACTGAAAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAA
 GAAGGGCAAAGAGGTCTGTGTCTGTGGGGTATCCACCACCCCCCAACT
 CCAAGGAGCAGCAGAATCTGTACCAAACGAAATGCTTACGTGTCTGTC
 GTGACATCTAACTACAACCGGCGCTTACGCCCCGAAATCGCCGAGCGTCC
 CAAAGTGCAGGATCAGGCCGAAGGATGAACCTACTACTGGACCTGTCTCA
 AACCCGGAGATACCATCATATTCGAAGCCAATGGCAATCTCATCGCTCCC
 ATGTATGCTGCTGCTCTCTCTCGCGGACACCACCACCACCACCTAATA
 G

SEQ ID NO:116 HA1-3mut PR8

AACTCCGAGAACGAAATCTGTTACCCGGCGACTTCATTGACAAAGAGGA
 GCTCCGCGAACAGCTGTCTTCTGTCTCTTCTTTCGAACGTTTCGAAATTT
 TCCCAAAGGAATCCTCTGGCCAAACCATAACACCAACGGAGTGACCGCC
 GCCTGTAGCCATGAGGGCAAGTCTTCTTCTACCGTAATCTGCTGTGGCT
 GACTGAAAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACA
 AGAAGGGCAAAGAGGTCTGGTGTGTGGGGTATCCACCACCCCCCAAC
 TCCAAGGAGCAGCAGAATCTGTACCAAACGAAATGCTTACGTGTCTGT
 CGTGACATCTAACTACAACCGGCGCTTACGCCCCGAAATCGCCGAGCGTC
 CCAAAGTGCAGGATCAGGCCGAAGGATGAACCTACTACTGGACCTGTCTC
 AAACCCGGAGATACCATCATATTCGAAGCCAATGGCAATCTCATCGCTCC
 CATGTATGCTGCTGCTCTCTCTCGCGGACACCACCACCACCACCTAATA

-continued

AG

SEQ ID NO:117 ngSTF2.HA0s

ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGGCCCAAGTGATCAACACCAACTCCCTGTCCCTGCTCA
 CTCAAAACAACTCCAGAAATCTCAGTCTGCTCTCGGTACTGCTATCGAG
 CGTCTCTCTCTCCGGCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
 ACAAGCTATCGCTAACCCTTTACTGCCAACATTAAAGGCTGACCCAAG
 CCTCTAGGAACGCCAATGATGGCATTTCTATCGCTCAAACCACCGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCGAACTCGCCGT
 CCAGTCCGCCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTGTCTGGTCAAAC
 CAGTTTAAACGGCGTCAAAGTGCTGGCTCAAGACAACACACTCACCATCCA
 AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGGACTCGATGACGCCGCTATCAAGGCCGCTACTGGCG
 GCACCTCAGGGAAGTGCCTCTGTGACCGGAGCGCTGTGAAGTTGACGCC
 GATAACAACAAATACTTCGTCACTATTGGTGGTTTCACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTCGCCACTGACGGAACAGTGA
 CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGCTGGTGTCTACTACT
 AAGACAGAGGTGCAGGAATCAAGGACACCCCTGCCGTGGTGTCCGCCGA
 TGCTAAGAACGCTCTCATTGTGTGGTGGTGTGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAATGTCTTACACCGACAAGAACGGAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGCTGATTA
 TGACGAAGCTACAGGCGCCATTAAAGCTAAAACCACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGCTGCCAATCAGCTGGGTGGAGTCGATGGA
 AAAACCGAAGTGGTCACAATCGACGGAAGACCTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCCAACCCGAACTGGCCGAGGCTGCTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG
 GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAACCTCGGCAATACCGTCAATCAACTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATTCTCCAACAAGCCGGCACCTCCGTCCTCGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGCCTGCTGGCTTCTGGATCTGGATCTGGTTCTG
 ACACCATCTGCATTGGCTACCATGCCAGCAGTCTACAGATACAGTCGAT
 ACCGTCCTGGAGAAACAAGTCACCGTCACGCACTCCGTGAACCTCTGGAA
 GGACTCCCATAATGGCAAATTGTGTCGGCTGAAAGGCATCGCCCCCTGTC
 AACTCGGCAAGTGCAATATTGCGGGGTGGCTGTTGGGCAACCCAGAATGT
 GACCCCTCTCGCCGTCCGTTCTTGGTCTTACATCGTGAAACACCTAA

-continued

CTCCGAGAACGGCATCTGTTACCCGGGCGACTTCATTGACTACGAGGAGC
 TCCGCGAACAGCTGTCTTCTGTCTCTTCTTCGAACGTTTCGAATTTTCC
 CAAAGGAATCCTCCTGGCCAAACCATAACACCAACGGAGTGACCGCCGCC
 TGTAGCCATGAGGGCAAGTCTTCTTTCTACCGTAATCTGCTGTGGCTGAC
 TGAAAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAAGA
 AGGGCAAAGAGGTCTGGTGTGTGGGGTATCCACCACCCCCCAACTCC
 AAGGAGCAGCAGAATCTGTACCAAACGAAATGCTTACGTGTCTGTGCT
 GACATCTAACTACAACCGGCGCTTCACGCCGAAATCGCCGAGCGTCCCA
 AAGTGCGGATCAGGCCGGAAGGATGAACTACTACTGGACCCTGCTCAAA
 CCCGGAGATACCATCATATTGGAAGCCAATGGCAATCTCATCGCTCCCAT
 GTATGCTTTCGCTCTCTCTCGCGATTTCGGTTCGGGCATAATCACTAGCC
 AAGCCTCCATGCACGAGTGCAATACCAAGTGTCAAACCTCCACTGGGAGCC
 ATTCAAAGCTCCCTGCCCTACCAAACATCCATCCTGTGACCATTGGAGA
 GTGCCCTAAATACGTGCGCAGCGCGAAACTGCGCATGGTGACCGGTCTCC
 GCAATATACCCTCCATACAATCGCGCGCCTGTTTGGTGTATCGCTGGC
 TTTATCGAGGGAGGATGGACTGGAATGATCGACGGCTGGTATGGCTATCA
 TCATCAAAACGAACAGGGTTCCGGCTACGCCGTGACCAAAAGTCCACTC
 AAAACGCCATTACCGTATTACAACAAAGTAAACACCGTGATAGAGAAA
 ATGAATATCCAATTCACGTGCGTGGGCAAAGAGTTTAAACAGCTGGAGAA
 GCGCATGGAAAACTGTAACAAAAAGTCGATGATGGCTTCTCTGACATCT
 GGACTTACAACGCCGAACCTCTCGTGTGCTCGAAAACGAGAGGACTCTG
 GACTTCCACGACTCCAACGTGAAGAACTGTACGAGAAAGTCAAATCCCA
 ACTCAAGAACACGCCAAAGAAATCGGCAACGGCTGCTTCGAGTTCTACC
 ACAATGTGATAACGAGTGTATGGAATCCGTACGGCAAGGCATTACGAC
 TACCCCAATACAGCGAAGAGAGCAAATTGAACCGCGAGAAAGTGGACGG
 CGTGAAACTGGAGTCCATGGGCATCTATCAGCACCACCACCACCACCT
 AATAG

SEQ ID NO:118 ngHA0s

GACACCATCTGCATTGGCTACCATGCCAGCAGTCTACAGATACAGTCGA
 TACCGTCTTGAGAAACAAGTACCGTCACGCACTCCGTGAACCTCCTGG
 AGGACTCCCATAATGGCAAATTTGTGTCGGCTGAAAGGCATCGCCCTCTG
 CAACTCGGCAAGTGCAATATTGCGGGGTGGCTGTTGGGCAACCCAGAATG
 TGACCCCTCTCGCCGTCGGTTCTTGGTCTTACATCGTGAAACACCTA
 ACTCCGAGAACGGCATCTGTTACCCGGGCGACTTCATTGACTACGAGGAG
 CTCCGCGAACAGCTGTCTTCTGTCTCTTCTTCGAACGTTTCGAAATTTT
 CCCAAAGGAATCCTCCTGGCCAAACCATAACACCAACGGAGTGACCGCCG
 CCTGTAGCCATGAGGGCAAGTCTTCTTTCTACCGTAATCTGCTGTGGCTG
 ACTGAAAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAA
 GAAGGGCAAAGAGGTCTGTGTGTGTGGGGTATCCACCACCCCCCAACT

-continued

CCAAGGAGCAGCAGAATCTGTACCAAACGAAATGCTTACGTGTCTGTC
 GTGACATCTAACTACAACCGGCGCTTCACGCCGAAATCGCCGAGCGTCC
 CAAAGTGCGGATCAGGCCGGAAGGATGAACTACTACTGGACCCTGCTCA
 AACCCGGAGATACCATCATATTGGAAGCCAATGGCAATCTCATCGCTCCC
 ATGTATGCTTTCGCTCTCTCTCGCGATTTCGGTTCGGGCATAATCACTAG
 CCAAGCCTCCATGCACGAGTGCAATACCAAGTGTCAAACCTCCACTGGGAG
 CCATTCAAAGCTCCCTGCCCTACCAAACATCCATCCTGTGACCATTGGA
 GAGTGCCCTAAATACGTGCGCAGCGCGAAACTGCGCATGGTGACCGGTCT
 CCGCAATATACCCTCCATACAATCGCGCGCCTGTTTGGTGTATCGCTG
 GCTTTATCGAGGGAGGATGGACTGGAATGATCGACGGCTGGTATGGCTAT
 CATCATCAAAACGAACAGGGTTCCGGCTACGCCGTGACCAAAAGTCCAC
 TCAAAACGCCATTAAACGGTATTACAACAAAGTAAACACCGTGATAGAGA
 AAATGAATATCCAATTCACGTGCGGTGGGCAAAGAGTTTAAACAGCTGGAG
 AAGCGCATGGAAAACTGAACAAAAAGTCGATGATGGCTTCTCTGACAT
 CTGGACTTACAACGCCGAACCTCTCGTGTGCTCGAAAACGAGAGGACTC
 TGGACTTCCACGACTCCAACGTGAAGAACCTGTACGAGAAAGTCAAATCC
 CAACTCAAGAACAACGCCAAAGAAATCGGCAACGGCTGCTTCGAGTTCTA
 CCACAAATGTGATAACGAGTGTATGGAATCCGTACGGCAAGGCATTACG
 ACTACCCCAATACAGCGAAGAGAGCAAATTGAACCGCGAGAAAGTGGAC
 GGCGTGAAACTGGAGTCCATGGGCATCTATCAGCACCACCACCACCACCA
 CTAATAG

SEQ ID NO:119 ngSTF2.HA1-1

ATGAATTTCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGGCCAAGTGATCAACACCAACTCCCTGTCCCTGCTCA
 CTCAAAACAACCTCCAGAAATCTCAGTCTGTCTCGGTACTGCTATCGAG
 CGTCTCTCTCCCGCCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
 ACAAGCTATCGCTAACCGCTTTACTGCCAACATTAAGGCCTGACCCAAG
 CCTCTAGGAACGCCAATGATGGCATTCTATCGCTCAAACCACCGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCGAACTCGCCGT
 CCAGTCCGCCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTGTCTGGTCAAACCT
 CAGTTTAAACGGCGTCAAAGTGTGGCTCAAGACAACACACTCACCATCCA
 AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGGACTCGATGACGCCGTATCAAGGCCGCTACTGGCG
 GCACTCAGGGAACCTGCCTCTGTGACCGGAGGCGCTGTGAAGTTCGACGCC
 GATAACAACAATACTTCGTCACTATTGGTGGTTTCACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTGCGCACTGACGGAAACGTGA

-continued

CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGCTGGTGCTACTACT
 AAGACAGAGGTGCAGGAACCAAGGACACCCCTGCCGTGGTGTCGCCCGA
 TGCTAAGAACGCTCTCATTGCTGGTGGTQTCGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAAATGTCCTACACCGACAAGAACGGAAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGCTGATTA
 TGACGAAGCTACAGGCCCATTAAGCTAAAACACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGCTGCCAATCAGCTGGGTGGAGTCGATGGA
 AAAACCGAAGTGGTCAACAATCGACGGAAGACCTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCAACCCGAACCTGGCCGAGGCTGCTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG
 GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATTCTCCAACAAGCCGACCTCCGTCCTCGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGCCTGCTGGCTTCTGGATCTGGATCTGGTTCTT
 CCCATAATGGCAAATTGTGTCGGCTGAAAGGCATCGCCCCCTCTGCAACTC
 GGCAAGTGCAATATTGCGGGGTGGCTGTTGGGCAACCCAGAATGTGACCC
 CCTCTGCCCGCTCCGTTCTTGGTCTTACATCGTGGAACACCTAACTCCG
 AGAACGGCATCTGTTACCCGGGCGACTTCATTGACTACGAGGAGCTCCGC
 GAACAGCTGTCTTCTGTCTCTTTCGAACGTTTCGAAATTTTCCCAA
 GGAATCTCTCGGCCAAACATAACACCAACGGAGTGACCGCCGCTGTA
 GCCATGAGGGCAAGTCTTCTTTCTACCGTAATCTGCTGTGGCTGACTGAA
 AAAGAGGGTTCTTATCCAAACTGAAGAACTCTTATGTCAACAAGAAGGG
 CAAAGAGGTCTGGTGCTGTGGGGTATCCACCACCCCCCAACTCCAAGG
 AGCAGCAGAATCTGTACAAAACGAAATGCTTACGTGTCTGTCTGTGACA
 TCTAACTACAACCGGCGCTTCACGCCGAAATCGCCGAGCGTCCCAAAGT
 GCGCGATCAGGCCGAAGGATGAACCTACTACTGGACCTGTCTAAACCCG
 GAGATACCATCATATTGCAAGCCAATGGCAATCTCATCGCTCCCATGTAT
 GCTTTCGCTCTCTCTCGCGGATTTCGGTTCCGGCATAATCACTAGCCAAGC
 CTCCATGCACGAGTGCAATACCAAGTGCAAACTCCACTGGGAGCCATTCT
 AAAGCTCCCTGCCCTACAAAACATCCATCTGTGACCATTTGGAGAGTGC
 CCTAAATACGTGCGCCACCACCACCACCACCACCTAATAG
 SEQ ID NO:120 ng HA1-1
 TCCCATAAATGGCAAATGTGTCGGCTGAAAGGCATCGCCCCCTCTGCAACT
 CGGCAAGTGCAATATTGCGGGGTGGCTGTTGGGCAACCCAGAATGTGACC
 CCCTCTGCCCGCTCCGTTCTTGGTCTTACATCGTGGAACACCTAACTCC
 GAGAACGGCATCTGTTACCCGGGCGACTTCATTGACTACGAGGAGCTCCG
 CGAACAGCTGTCTTCTGTCTCTTCTTTCGAACGTTTCGAAATTTTCCCAA
 AGGAATCTCTCGGCCAAACATAACACCAACGGAGTGACCGCCGCTGT

-continued

AGCCATGAGGGCAAGTCTTCTTTCTACCGTAATCTGCTGTGGCTGACTGA
 AAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAAGAAGG
 GCAAAGAGGTCTGGTGCTGTGGGGTATCCACCACCCCCCAACTCCAAG
 GAGCAGCAGAATCTGTACAAAACGAAAATGCTTACGTGTCTGTCTGTGAC
 ATCTAACTACAACCGGCGCTTACGCCCCGAAATCGCCGAGCGTCCCAAAG
 TGCGCGATCAGGCCGAAGGATGAACCTACTACTGGACCTGTCTAAACCC
 GGAGATACCATCATATTGCAAGCCAATGGCAATCTCATCGCTCCCATGTA
 TGCTTTCGCTCTCTCTCGCGGATTTCGGTTCCGGCATAATCACTAGCCAAG
 CCTCCATGCACGAGTGCAATACCAAGTGCAAACTCCACTGGGAGCCATT
 CAAAGCTCCCTGCCCTACAAAACATCCATCTGTGACCATTTGGAGAGTG
 CCCTAAATACGTGCGCCACCACCACCACCACCACCTAATAG
 SEQ ID NO:121 ngSTF2.HA1-2
 ATGAAATTCCTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTT
 TTACATCTATGCGGCCCAAGTGATCAACACCAACTCCCTGTCCCTGCTCA
 CTCAAAACAACCTCCAGAAATCTCAGTCTGCTCTCGGTACTGCTATCGAG
 CGTCTCTCCTCCGGCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
 ACAAGCTATCGCTAACCCTTTTACTGCCAACATTAAAGGCTGACCCAAG
 CCTCTAGGAACGCCAATGATGGCATTTCTATCGCTCAAACCCGGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCACGTTCCGCGAACTCGCCGT
 CCAGTCCGCCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTGTCTGGTCAAAC
 CAGTTTAAACGGCTCAAGTGCTGGCTCAAGACAACACACTCACCATCCA
 AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGACTCGATGACGCCCTATCAAGGCCGCTACTGGCG
 GCACTCAGGGAAGTGCCTCTGTGACCGGAGGCGCTGTGAAGTTTCGACGCC
 GATAACAACAAATACTTCGTCACTATTTGGTGGTTTCACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTGCGCACTGACGGAACAGTGA
 CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGCTGGTGTACTACT
 AAGACAGAGGTGCAGGAACCTCAAGGACACCCCTGCCGTGGTGTCCGCCGA
 TGCTAAGAACGCTCTCATTGCTGGTGGTGTGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAAATGTCCTACACCGACAAGAACGGAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGCTGATTA
 TGACGAAGCTACAGCGCCATTAAAGCTAAAACACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGTGCCAATCAGCTGGGTGGAGTCGATGGA
 AAAACCGAAGTGGTCACAATCGACGGAAGACCTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCAACCCGAACCTGGCCGAGGCTGTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG

-continued

GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATTCTCCAACAAGCCGCGACCTCCGTCTCGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGCCTACTAGCATCAGTTCCGGCTCAGGTTCCA
 AAGGCATAGCGCCACTGCAGCTCGGCAAAATGTAACATCGCAGTTGGCTC
 CTTGGTAACCCGAGTGCGACCCCTCCTCCCTGTACGATCCTGGAGTTA
 TATCGTGGAGACCCCAATAGCGAGAACGGAATTTGCTACCAGGAGATT
 TCATAGACTACGAGGAGTTGCGCGAGCAGCTTTCTGTCTGTGAGCAGCTTC
 GAGAGGTTTGAATCTTCCGAAGGAGAGCAGCTGGCCGAATCATAACAC
 TAACGGTGTGACAGCCGCTGCAGTCATGAAGGAAAGAGTTTCATTCTATC
 GCAACCTGTGTGGTTGACGGAGAAAGAGGGCAGCTACCCTAAGTTGAAG
 AACTCCTATGTGAACAAAAAGGCAAGGAGGTTCTGGTGTGTGGGGCAT
 ACACCACCCCCCAATAGCAAGGAGCAGCAGAATCTGTACCAAAACGAGA
 ATGCTTATGTGAGCGTGGTCACTAGTAACATAAACCCTCGGTTCACTCCC
 GAGATCGCCGAGCGTCCGAAGGTGAGGGACCAGGCAGGCCGGATGAACATA
 CTACTGGACCTATTGAAGCCAGGGGACACGATTATCTTCGAGGCAACG
 GAAACCTCATAGCGCCGATGTACGCGTTGCGCCCTGAGCCGTGGCTTTGGA
 TCGGGGATCATCAGTCTCACCATCATCACCACCATTAATAG

SEQ ID NO:122 ng HA1-2

AAAGGCATAGCGCCACTGCAGCTCGGCAATGTAACATCGCAGGTTGGCT
 CCTTGGTAACCCGGAGTGCGACCCCTCCTCCCTGTACGATCCTGGAGTT
 ATATCGTGGAGACCCCAATAGCGAGAACGGAATTTGTACCCAGGAGAT
 TTCATAGACTACGAGGAGTTGCGCGAGCAGCTTTCGTCTGTGAGCAGCTT
 CGAAAGGTTTCGAGATATTTCCGAAGGAGAGCAGCTGGCCGAATCATAACA
 CTAACGGTGTGACAGCCGCTGCAGTCATGAAGGAAAGAGTTTCATTCTAT
 CGCAACCTGTGTGGTTGACGGAGAAAGAGGGCAGCTACCCTAAGTTGAA
 GAACTCCTATGTGAACAAAAAGGCAAGGAGGTTCTGGTGTGTGGGGCA
 TACACCACCCCCCAATAGCAAGGAGCAGCAGAATCTGTACCAAAACGAG
 AATGCCTATGTGAGCGTGGTCACTAGTAACATAAACCCTCGGTTCACTCC
 CGAGATCGCCGAGCGTCCGAAGGTGAGGGACCAGGCAGGCCGGATGAAC
 ACTACTGGACCTTATTGAAGCCAGGGACACGATTATCTTCGAGGCAAC
 GGAACCTCATAGCGCCGATGTACGCTTTCGCCCTGAGCCGTGGCTTTGG
 ATCGGGGATCATCAGTCTCACCATCATCACCACCATTAATAG

SEQ ID NO:123 ngSTF2.HA1-2mut

ATGAAATTTCTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTT
 TTACATCTATGCGGCCCAAGTGATCAACCAACTCCCTGTCCCTGTCTCA
 CTCAAAACAACCTCCAGAAATCTCAGTCTGCTCTCGGTACTGCTATCGAG
 CGTCTCTCTCCGGCCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
 ACAAGCTATCGCTAACCGCTTTACTGCCAACATTAAGGCCTGACCCAAG

-continued

CCTCTAGGAACGCCAATGATGGCATTTCATCGCTCAAACCACCGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCGAACTCGCCGT
 CCAGTCCGCCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTGTCTGGTCAAAC
 CAGTTTAACGGCGTCAAAGTGCTGGCTCAAGACAACACACTCACCATCCA
 AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCCTGGACTCCCTGAACGTCCAAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGGACTCGATGACGCCGCTATCAAGGCCGCTACTGGCG
 GCACCTCAGGGAACGCTCTGTGACCGGAGGCGCTGTGAAGTTGACGCC
 GATAACAACAAATACTTCGTCACTATTGGTGGTTTCACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTCCGCACTGACGGAACAGTGA
 CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGCTGGTGTCTACTACT
 AAGACAGAGGTGCAGGAACCAAGGACACCCCTGCCGTGGTGTCCGCCGA
 TGCTAAGAACGCTCTCATTGTCTGGTGGTGTGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAAATGTCTACACCGACAAGAACGAAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGTGATTA
 TGACGAAGCTACAGCGCCATTAAAGCTAAAACCACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGCTGCCAATCAGCTGGGTGGAGTCGATGGA
 AAAACCGAAGTGGTCACAATCGACGAAAGACTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCCAACCCGAAGTGGCCGAGGCTGTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG
 GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATTCTCCAACAAGCCGCGACCTCCGTCTCGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGCCTGCTGGCTTCTGGATCTGGATCTGGTTCTA
 AAGGCGCTGCCCTCTGCAACTCGGCAAGTGAATATTGCGGGGTGGCTG
 TTGGGCAACCCAGAATGTGACCCCTCCTGCCCGTCCGTCTTGGTCTGA
 CATCGCTGAAACACCTAACTCCGAGAACGGCATCTGTTACCCGGGCGACT
 TCATTGACTACGAGGAGCTCCGCGAACAGCTGTCTTCTGTCTCTTCTTTC
 GAACGTTTTGAAATTTTCCCAAAGGAATCCTCTGGCCAAACCAATAACAC
 CAACGGAGTGACCGCCGCTGTAGCCATGAGGGCAAGTCTTCTTTCTACC
 GTAATCTGCTGTGGCTGACTGAAAAAGAGGGTCTTATCCCAACTGAAG
 AACTCTTATGTCAACAAGAAGGGCAAGAGGTCCTGGTGTGTGGGGTAT
 CCACCACCCCCCAACTCCAAGGAGCAGCAGAATCTGTACCAAAACGAAA
 ATGCTTACGTGTCTGTCTGACATCTAACTACAACCGCGCTTCACGCC
 GAAATCGCCGAGCGTCCCAAAGTGCGCGATCAGGCCGGAAGGATGAACATA

-continued

CTACTGGACCCTGCTCAAACCCGGAGATACCATCATATTGGAAGCCAATG
GCAATCTCATCGCTCCCATGTATGCTTTCGCTCTCTCTCGCGGATTTCGGT
TCCGGCATAATCACTAGCCACCACCACCACCACCACCTAATAG

SEQ ID NO:124 ngHA1-2mut

AAAGGCGCTGCCCTCTGCAACTCGCAAGTGCAATATTGCGGGGTGGCT

GTTGGGCAACCCAGAATGTGACCCCTCCTGCCGTCCGTTCTTGGTCTG
ACATCGCTGAAACACCTAACTCCGAGAACGGCATCTGTTACCCGGGCGAC
TTCATTGACTACGAGGAGCTCCGCGAAGAGCTGTCTTCTGTCTCTTCTTT
CGAACGTTTTCGAAATTTTCCCAAAGGAATCCTCCTGGCCAAACCATAACA
CCAACGAGTGACCGCCGCTGTAGCCATGAGGCAAGTCTTCTTTCTAC
CGTAATCTGCTGTGGCTGACTGAAAAAGAGGGTTCTTATCCCAAAGTAA
GAACTCTTATGTCAACAAGAAGGGCAAGAGGTCTGGTGTGTGGGGTA
TCCACCACCCCCCAACTCCAAGGAGCAGCAGAATCTGTACCAAAACGAA
AATGCTTACGTGTCTGTCTGTGACATCTAACTACAACCGGCGCTTCACGCC
CGAAATCGCCGAGCGTCCCAAAGTGCGCGATCAGGCCGGAAGGATGAACT
ACTACTGGACCCTGCTCAAACCCGGAGATACCATCATATTGGAAGCCAAT
GGCAATCTCATCGCTCCCATGTATGCTTTCGCTCTCTCTCGCGGATTTCGG
TTCCGGCATAATCACTAGCCACCACCACCACCACCACCTAATAG

SEQ ID NO:125 ngSTF2.HA1-3

ATGAAATCTTAGTCAACGTTGCCCTTGTTTATGGTCGTATACATTTT

TTACATCTATGCGGCCCAAGTGATCAACACCAACTCCCTGTCCCTGTCTCA
CTCAAAACAACCTCCAGAAATCTCAGTCTGCTCTCGGTACTGCTATCGAG
CGTCTCTCCTCCGGCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
ACAAGCTATCGCTAACCGCTTTACTGCCAACATTAAAGGCCTGACCCAAAG
CCTCTAGGAACGCCAATGATGGCATTCTATCGCTCAAACCACCGAAGGC
GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCGAATCGCCGT
CCAGTCCGCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
CTGAGATCACCAAGGCTCAACGAGATCGATCGTGTGTCTGGTCAAAC
CAGTTTAAACGGCGTCAAAGTGCTGGCTCAAGACAACACACTCACCATCCA
AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAAGGCTTACGAC
GTGAAGGACACCGCTGTGACAACCAAAGCCTATGCTAATCAAGGAACAAC
CCTGGACGTGTCCGGACTCGATGACGCCGCTATCAAGGCCGCTACTGGCG
GCACTCAGGGAATGCCTCTGTGACCGGAGGCGCTGTGAAGTTCGACGCC
GATAACAACAATACTTCGTCACTATTGGTGGTTTCTACTGGCGCTGACGC
CGCTAAGAACGGTGACTACGAGGTCAACGTCGCCACTGACGGAACAGTGA
CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGTGTGGTGTACTACT
AAGACAGAGGTGACGGAATCAAGGACACCCCTGCCGTGGTGTCCGCCGA
TGCTAAGAACGCTCTCATTGCTGGTGGTGTGACGCTACGGACGCCAACG
GAGCCGAGCTTGTGAAATGTCTTACACCGACAAGAACGGAAGACTATC

-continued

GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGTGATTA
TGACGAAGCTACAGGCGCCATTAAAGCTAAAACCACATCTTATACCGCTG
CCGATGGTACTACCAAGACTGCTGCCAATCAGCTGGGTGGAGTCGATGGA
AAAACCAGAGTGGTCACAATCGACGGAAGACCTATCAAGCTTCCAAGGC
TGCTGGCCACGACTTCAAGGCCCAACCCGAACCTGGCCGAGGCTGCTGCTA
AAACCACTGAAAACCCCTGCGAGAAAATTGACGCTGCCCTGGCCCAAGTG
GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
TATTACAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
GTATTGAGGATTCCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
CAAATTTCTCCAAAGCCGGCACCTCCGTCCTCGCCCAAGCCAATCAGGT
GCCACAGAATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCA
ATAGCGAGAACGGAATTTGCTACCCAGGAGATTTATAGACTACGAGGAG
TTGCCGCGAGCAGCTTTCGTCTGTGAGCAGCTTCGAGAGGTTGCAAAATCTT
CCCGAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCCG
CCTGCAGTCATGAAGGAAAGAGTTCATTCTATCGCAACCTGCTGTGGTTG
ACGGAGAAGGAGGGCAGCTACCTAAGCTGAAGAACTCCTATGTGAACAA
AAAAGGCAAGGAGGTTCTGGTGTGTGGGGCATAACACCCCCCAATA
GCAAGGAGCAGCAGAATCTGTACCAAAACGAGAATGCCTATGTGAGCGTG
GTCACTAGTAACATAAACCGTCGGTTCCTCCGAGATCGCCGAGCGTCC
GAAGGTGAGGGACACGAGCAGGCGGATGAACCTACTACTGGACCCTATTGA
AGCCAGGGGACACGATTATCTTCGAGGCAACCGGAAACCTCATAGCGCCG
ATGTACGCCCTTCGCCCTGAGCCGTGGCCACCATCATCACCACCATTAAATA
G

SEQ ID NO:126 ngHA1-3

AATAGCGAGAACGGAATTTGCTACCCAGGAGATTTATAGACTACGAGGA

GTTCGCGAGCAGCTTTCGTCTGTGAGCAGCTTCGAAAGGTTTCGAGATAT
TCCCCAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCC
GCCTGCAGTCATGAAGGAAAGAGTTCATTCTATCGCAACCTGCTGTGGTT
GACGAGAAAGAGGGCAGCTACCTAAGTTGAAGAACTCCTATGTGAACA
AAAAAGGCAAGGAGGTTCTGGTGTGTGGGGCATAACACCCCCCAAT
AGCAAGGAGCAGCAGAATCTGTACCAAAACGAGAATGCCTATGTGAGCGT
GGTCACTAGTAACATAAACCGTCGGTTCCTCCGAGATCGCCGAGCGTC
CGAAGGTGAGGGACACGAGCAGGCGGATGAACCTACTACTGGACCCTATTG
AAGCCAGGGGACACGATTATCTTCGAGGCAACCGGAAACCTCATAGCGCC
GATGTACGCCCTTCGCCCTGAGCCGTGGCCACCATCATCACCACCATTAAAT
AG

SEQ ID NO:127 ngSTF2.HA1-3mut

ATGAAATCTTAGTCAACGTTGCCCTTGTTTATGGTCGTATACATTTT

TTACATCTATGCGGCCCAAGTGATCAACACCAACTCCCTGTCCCTGTCTCA

-continued

CTCAAAACAACCTCCAGAAATCTCAGTCTGCTCTCGGTACTGCTATCGAG
 CGTCTCTCTCCGGCCTGAGGATCAACTCCGCTAAAGATGACGCCGCCG
 ACAAGCTATCGCTAACCGCTTTACTGCCAACATTAAGGCCGTGACCCAAG
 CCTCTAGGAACGCCAATGATGGCATTCTATCGCTCAAACCACCGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCGAATCGCCGT
 CCAGTCCGCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTCTGGTCAAAC
 CAGTTTAAACGGCTCAAAAGTGTGGCTCAAGACAACACACTCACCATCCA
 AGTAGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGGACTCGATGACGCCGCTATCAAGGCCGCTACTGGCG
 GCACTCAGGGAATGCCTCTGTGACCGGAGCGCTGTGAAGTTCGACGCC
 GATAACAACAATACTTCGTCACATATTGGTGGTTTCACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTGCGCACTGACGGAACAGTGA
 CACTGGCCGCTGGCGCTACCAAGACCACAATGCCTGCTGGTGTCTACTACT
 AAGACAGAGGTGCAGGAATCAAGGACACCCCTGCCGTGGTGTCCGCCGA
 TGCTAAGAACGCTCTCATTGCTGGTGGTGTGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAAATGTCTACACCGACAAGAACGGAAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGCTGATTA
 TGACGAAGCTACAGGCCCATTAAGCTAAAACACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGCTGCCAATCAGCTGGGTGGAGTCGATGGA
 AAAACCGAAGTGGTCACAATCGACGGAAGACCTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCCAACCCGAACCTGGCCGAGGCTGCTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG
 GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATTCTCCAACAAGCCGGCACCTCCGTCTCGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGCCTGTGGCTTCTGGATCTGGATCTGGTTCTA
 ACTCCGAGAACGAAATCTGTTACCCGGCGACTTCATTGACAAAGAGGAG
 CTCCGCGAACAGCTGTCTTGTCTCTTCTTTTGAACGTTTCGAAATTTT
 CCCAAAGGAATCTCTCGCCAAACCATACACCAACGGAGTGACGCCCG
 CCTGTAGCCATGAGGGCAAGTCTTCTTTCTACCGTAATCTGCTGTGGCTG
 ACTGAAAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAA
 GAAGGGCAAAGAGTCTGTGTGTGTGGGTATCCACCACCCCCCAACT
 CCAAGGAGCAGCAGAATCTGTACCAAAACGAAAATGCTTACGTGTCTGTG
 GTGACATCTAACTACAACCGCGCTTCACGCCGAAATCGCCGAGCGTCC
 CAAAGTGCAGATCAGGCCGAAGGATGAACCTACTACTGGACCTGTCTCA

-continued

AACCCGGAGATACCATCATATTTCGAAGCCAATGGCAATCTCATCGCTCCC
 ATGTATGTGTGCTCTCTCTCGCGACACCACCACCACCACCACCTAATA
 G

SEQ ID NO:128 ngHA1-3mut

AACCTCCGAGAACGAAATCTGTTACCCGGCGACTTCATTGACAAAGAGGA
 GCTCCGCGAACAGCTGTCTTCTGTCTCTTCTTTTGAACGTTTCGAAATTT
 TCCCAAAGGAATCCTCCTGGCCAAACCATAACACCAACGAGTGACCGCC
 GCCTGTAGCCATGAGGGCAAGTCTTCTTTTACCCGTAATCTGCTGTGGCT
 GACTGAAAAAGAGGGTTCTTATCCCAAACCTGAAGAACTCTTATGTCAACA
 AGAAGGGCAAAGAGGTCTGTGTGTGTGGGTATCCACCACCCCCCAAC
 TCCAAGGAGCAGCAGAATCTGTACCAAAACGAAATGCTTACGTGTCTGT
 CGTGACATCTAACTACAACCGCGCTTCACGCCGAAATCGCCGAGCGTC
 CCAAAGTGCAGATCAGGCCGAAGGATGAACCTACTACTGGACCTGTCTC
 AAACCCGGAGATACCATCATATTTCGAAGCCAATGGCAATCTCATCGCTCC
 CATGTATGCTGCTGCTCTCTCTCGCGGACACCACCACCACCACCTAAT
 AG

SEQ ID NO:129 wtSTF2.HA1-1ng

ATGAAATTCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGGCCAGGTTATCAATACCAACTCCCTGTGCTGTCTCA
 CCCAAATAACCTTAATAAAAGCCAGAGCGCACTGGGAACCGCCATAGAA
 CGCCTCTCAAGCGGCTCCGGATCAATTCTGCAAAAGACGACGCCGCCG
 TCAGGCCATCGCAAACCGCTTTACCGCCAAATATCAAGGGACTGACGAGG
 CTTTCGAGGAATGCTAACGATGGAATAAGCATCGCTCAAACACGAGGGC
 GCCCTGAACGAGATCAACAACAACCTACAGCGCGTCAAGGAGCTCGCAGT
 GCAGTCCGCCAATTCGACCAACTCGCAGTCGGACCTGGACTCGATCCAAG
 CCGAAATCACCAGCGCCTGAATGAGATTGACCGGGTGAGCGGTGACACA
 CAGTTTAAACGGCGTGAAGTACTTGACAGGATAACACACTTACGATACA
 GGTGGGCGCCAACGATGGTGAAACCATAGACATTGATCTCAAACAGATTA
 ACAGCCAGACGCTCGGGTTGGATAGCCTGAATGTGCAAAAGCGGTACGAC
 GTGAAAGACACGGCGGTCACTACCAAGCCTACGCTAACAATGGCACTAC
 CTTGGATGTGAGCGGATTGGATGATGCAGCAATCAAGGCTGTACCGGCG
 GTACGAACGGAACCGCGTCCGTGACCGCGGTGCCGTGAAGTTCGATGCT
 GACAACAATAAGTATTTCTGTCACCATTTGAGGCTTTACTGGCGCCGACGC
 AGCAAAGAACGGCGACTATGAAGTGAACGTGGCAACCGATGGAACCGTGA
 CGCTGGCCGCTGGTGCCACCAAGACCACCATGCCAGCCGGCGCCACAAC
 AAGACCGAGGTGCAGGAGTTAAAGGACACCCCGCGGTGGTTAGCGCAGA
 TGCCAAAAACGCCTTGATCGCCGGCGGAGTGGATGCAACTGATGCTAATG
 GTGCGGAGCTGGTTAAATGTCGTATACAGACAAGAATGGTAAGACGATC
 GAGGGCGGTTATGCCCTTAAGGCAGGAGATAAGTATTACGCTGTGATTA

-continued

CGATGAGGCGACGGGAGCTATTAAAGCCAAGACAACGTCATACACGGCGG
 CGGACGGAACGACTAAGACGGCTGCCAATCAGTTGGGAGGGTTGACGGG
 AAGACAGAGGTCGTTACGATCGATGGCAAGACATACAACGCTTCCAAGGC
 CGCTGGCCACGATTTCAAAGCTCAACCCGAACCTGGCCGAGGCCGCGCGA
 AAACAACCTGAGAACCCGTTGCGAGAAGATTGATGCGGCCCTGGCGCAAGTA
 GATGCCCTGCGCTCAGACCTGGGCGCGGTTCAAAATCGATTCAATTCCGC
 GATTACAAACCTGGGCAATACAGTAACAATCTATCCGAGGCCAGATCCC
 GCATTGAAGACTCCGACTACGCGACAGAAGTAAGTAACATGAGTCGTGCC
 CAGATTCTGCAGCAGGCCGCGACTAGTGTCTTGGCCAGGCCAATCAAGT
 CCCGACAGATGTGCTGAGCCTACTAGCATCTGGATCTGGATCTGGTTCTT
 CCCATAATGGCAAATTGTGTGCGGTGAAAGCATCGCCCCCTCTGCAACTC
 GGCAAGTGCAATATTGCGGGGTGGCTGTTGGGCAACCCAGAATGTGACCC
 CCTCCTGCCGTCGCTTCTTGGTCTTACATCGTGGAAACACCTAACTCCG
 AGAAGGCATCTGTTACCCGGGCGACTTCATTGACTACGAGGAGCTCCGC
 GAACAGCTGTCTTCTGTCTCTTTCGAACGTTTCGAAATTTTCCCAA
 GGAATCCTCTGGCCAAACCATAACACCAACGGAGTGACCGCCGCTGTA
 GCCATGAGGGCAAGTCTTCTTTTACCGTAATCTGCTGTGGCTGACTGAA
 AAAGAGGGTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAAGAAGG
 CAAAGAGGTCCTGGTGTGTGGGTATCCACCACCCCCCAACTCCAAGG
 AGCAGCAGAATCTGTACCAAAACGAAATGCTTACGTGTCTGTCTGTGACA
 TCTAACTACAACCGCGCTTCACGCCCGAAATCGCCGAGCTCCCAAAGT
 GCGCGATCAGGCCGGAAGGATGAACTACTACTGGACCTGTCTAAACCCG
 GAGATACCATCATATTGGAAGCCAATGGCAATCTCATCGCTCCCATGTAT
 GCTTTCGCTCTCTCTCGCGGATTCCGGTTCCGGCATAATCACTAGCCAAGC
 CTCCATGCACGAGTGCAATACCAAGTGTCAAACTCCACTGGGAGCCATTTC
 AAAGCTCCCTGCCCTACCAAAACATCCATCCTGTGACCATTGGAGAGTGC
 CCTAAATACGTGCCACCACCACCACCACCACCTAATAG
 SEQ ID NO:130 ngHA1-1
 TCCCATATGGCAAATTGTGTGCGGTGAAAGGCATCGCCCCCTGCAACT
 CGGCAAGTGCAATATTGCGGGGTGGCTGTTGGGCAACCCAGAATGTGACC
 CCCTCCTGCCGTCGCTTCTTGGTCTTACATCGTGGAAACACCTAACTCC
 GAGAAGGCATCTGTTACCCGGGCGACTTCATTGACTACGAGGAGCTCCG
 CGAACAGCTGTCTTCTGTCTCTTCTTTCGAACGTTTCGAAATTTTCCCAA
 AGGAATCCTCCTGGCCAAACCATAACACCAACGGAGTGACCGCCGCTGT
 AGCCATGAGGGCAAGTCTTCTTTTACCGTAATCTGCTGTGGCTGACTGA
 AAAAGAGGGTCTTATCCCAAACCTGAAGAACTCTTATGTCAACAAGAAGG
 GCAAAGAGGTCCTGGTGTGTGGGTATCCACCACCCCCCAACTCCAAG
 GAGCAGCAGAATCTGTACCAAAACGAAATGCTTACGTGTCTGTCTGTGAC
 ATCTAACTACAACCGCGCTTCACGCCCGAAATCGCCGAGCGTCCCAAAG

-continued

TGCGCGATCAGGCCGGAAGGATGAACCTACTACTGGACCCCTGCTCAAACCC
 GGAGATACCATCATATTGGAAGCCAATGGCAATCTCATCGCTCCCATGTA
 TGCTTTCGCTCTCTCTCGCGGATTCCGGTTCCGGCATAATCACTAGCCAAG
 CCTCCATGCACGAGTGCAATACCAAGTGTCAAACTCCACTGGGAGCCATT
 CAAAGCTCCCTGCCCTACCAAAACATCCATCCTGTGACCATTGGAGAGTG
 CCCTAAATACGTGCCACCACCACCACCACCACCTAATAG
 SEQ ID NO:131 ngSTF2.HA1-1wt
 ATGAAATTCCTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGGCCCAAGTGATCAACACCAACTCCCTGTCCCTGCTCA
 CTCAAAACAACCTCCAGAAATCTCAGTCTGTCTCGGTACTGCTATCGAG
 CGTCTCTCCTCCGGCCTGAGGATCAACTCCGCTAAAGATGACGCCGCCGG
 ACAAGCTATCGCTAACCGCTTTACTGCCAACATTAAAGCCTGACCCAAG
 CCTCTAGGAACGCCAATGATGGCATTCTATCGCTCAAACCACCGAAGGC
 GCCCTGAACGAAATCAATAATAACCTGCAACGTGTCCGCGAACTCGCCGT
 CCAGTCCGCCCAATCTACCAACTCTCAGTCTGACCTGGACTCTATCCAAG
 CTGAGATCACCCAAAGGCTCAACGAGATCGATCGTGTGTCTGGTCAAACCT
 CAGTTTAAACGGCGTCAAAGTGCTGGCTCAAGACAACACACTCACCATCCA
 AGTAGGAGCCAATGACGGCGAGACAATCGATATCGACCTGAAGCAGATCA
 ATTCTCAAACCTCTGGGCTGGACTCCCTGAACGTCCAAAGGCTTACGAC
 GTGAAGGACACCGCTGTGACAACCAAGCCTATGCTAATCAAGGAACAAC
 CCTGGACGTGTCCGGACTCGATGACGCCGCTATCAAGGCCGCTACTGGCG
 GCACCTCAGGGAAGTGCCTCTGTGACCGGAGGCGCTGTGAAGTTTCAGCGC
 GATAACAACAATACTTCTGCTACTATTGGTGGTTTCTACTGGCGCTGACGC
 CGCTAAGAACGGTGACTACGAGGTCAACGTGCGCACTGACGGAACAGTGTA
 CACTGGCCGCTGGCGCTACCAAGACCACAAATGCCTGCTGGTGTCTACTACT
 AAGACAGAGGTGCAGGAACCTAAGGACACCCCTGCCGTGGTGTCCGCCGA
 TGCTAAGAACGCTCTCATTTGCTGGTGGTGTGACGCTACGGACGCCAACG
 GAGCCGAGCTTGTGAAAATGTCTACACCGACAAGAACGGAAGACTATC
 GAGGGAGGTTACGCTCTGAAGGCTGGCGATAAGTACTACGCCGCTGATTA
 TGACGAAGCTACAGGCGCCATTAAAGCTAAAACCACATCTTATACCGCTG
 CCGATGGTACTACCAAGACTGTGCGCAATCAGCTGGGTGGAGTCGATGGA
 AAAACCGAAGTGGTCACAATCGACGGAAGACCTATCAAGCCTCCAAGGC
 TGCTGGCCACGACTTCAAGGCCCAACCCGAACCTGGCCGAGGCTGTGCTA
 AAACCACTGAAAACCCCTGCAGAAAATTGACGCTGCCCTGGCCCAAGTG
 GATGCCCTCCGCTCCGACCTCGGCGCTGTGCAAAACCGCTTCAACTCCGC
 TATTACAAAACCTCGGCAATACCGTCAATCAGCTCTCTGAAGCTAGGTCTC
 GTATTGAGGATTCCGATTACGCTACCGAGGTCTCCAGATGTCCCGTGCC
 CAAATCTCCAACAAGCCGGCACCTCCGTCCTGCCCAAGCCAATCAGGT
 GCCACAGAATGTGCTGAGCCTACTAGCATCAGGTTCCGGCTCAGGTTCCA

-continued

GCCACAACGGAAAGCTGTGCCGTCTGAAAGGCATAGCGCCACTGCAGCTC
 GGCAAATGTAACATCGCAGGTTGGCTCCTTGGTAACCCGGAGTGCGACCC
 CCTCTCCCTGTACGATCCTGGAGTTATATCGTGGAGACCCCCAATAGCG
 AGAACGGAATTTGCTACCCAGGAGATTTCATAGACTACGAGGAGTTGCGC
 GAGCAGCTTTTCTGTGTGAGCAGCTTCGAGAGGTTTCGAAATCTTCCCGAA
 GGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCCGCTGCA
 GTCATGAAGGAAAGAGTTCATTCTATCGCAACCTGCTGTGGTTGACGGAG
 AAAGAGGGCAGCTACCCTAAGTTGAAGAACTCCTATGTGAACAAAAAGG
 CAAGGAGGTTCTGGTGTGTGGGGCATACACCACCCCCCAATAGCAAGG
 AGCAGCAGAATCTGTACAAAACGAGAATGCCCTATGTGAGCGTGGTCACT
 AGTAACATAACCGTCGGTTCACTCCCGAGATCGCCGAGCGTCCGAAGGT
 GAGGGACCAGGCAGGCCGGATGAACCTACTACTGGACCTATTGAAGCCAG
 GGGACACGATTATCTTCGAGGCAAACGGAACCTCATAGCGCCGATGTAC
 GCGTTCGCCCTGAGCCGCGCTTTGGATCGGGGATCATCAGCTTAACGC
 CTCGATGCACGAATGTAATACCAATGCCAGACCCCACTGGGTGCTATCA
 ACTCGTCCTTACCCTATCAAAATATACATCCGGTCACCATAGGCGAGTGT
 CCCAAATATGTACAGACCATCATCACCACCATTAAATAG

SEQ ID NO:132 HA1-1 PR8

AGCCACAACGGAAAGCTGTGCCGTCTGAAAGGCATAGCGCCACTGCAGCT
 CGGCAAATGTAACATCGCAGGTTGGCTCCTTGGTAACCCGGAGTGCGACC
 CCCTCTCCCTGTACGATCCTGGAGTTATATCGTGGAGACCCCCAATAGC
 GAGAACGGAATTTGCTACCCAGGAGATTTCATAGACTACGAGGAGTTGCG
 CGAGCAGCTTTCTGTGTGAGCAGCTTCGAAAGGTTTCGAGATATTCGGA
 AGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGTGACAGCCGCTGTC
 AGTCATGAAGGAAAGAGTTTCACTTATCGCAACCTGCTGTGGTTGACGGA
 GAAAGAGGGCAGCTACCCTAAGTTGAAGAACTCCTATGTGAACAAAAAG
 GCAAGGAGGTTCTGGTGTGTGGGGCATACACCACCCCCCAATAGCAAG
 GAGCAGCAGAATCTGTACAAAACGAGAATGCCCTATGTGAGCGTGGTCAC
 TAGTAACATAAACCGTCGGTTCACTCCCGAGATCGCCGAGCGTCCGAAGG
 TGAGGGACCAGGCAGGCCGGATGAACCTACTACTGGACCTATTGAAGCCA
 GGGGACACGATTATCTTCGAGGCAAACGGAACCTCATAGCGCCGATGTA
 CGCCTTCGCCCTGAGCCGTGGCTTTGGATCGGGGATCATCAGCTCTAACG
 CCTCGATGCACGAATGTAATACCAATGCCAGACCCCACTGGGTGCTATC
 AACTCGTCCTTACCCTATCAAAATATACATCCGGTCACCATAGGCGAGTG
 TCCCAAATATGTACAGACCATCATCACCACCATTAAATAG

SEQ ID NO:133 HA1-1 PR8

ATGAAATCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTC
 TTACATCTATGCGAGCCACAACGGAAGCTGTGCCGTCTGAAAGGCATAG
 CGCCACTGCAGCTCGGCAAATGTAACATCGCAGGTTGGCTCCTTGGTAAC
 CCGGAGTGCGACCCCTCCTCCTGTACGATCCTGGAGTTATATCGTGA

-continued

GACCCCCAATAGCGAGAACGGAATTTGCTACCCAGGAGATTTTCATAGACT
 ACGAGGAGTTGCGCGAGCAGCTTTCTGTCTGTGAGCAGCTTCGAAAGGTTTC
 GAGATATTTCCCGAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGT
 GACAGCCGCTGCAGTCATGAAGGAAAGAGTTTATTCTATCGCAACCTGC
 TGTGGTTGACGGAGAAAGAGGGCAGCTACCCTAAGTTGAAGAACTCCTAT
 GTGAACAAAAAGGCAAGGAGGTTCTGGTGTGTGGGGCATACACCACCC
 CCCCCAATAGCAAGGAGCAGCAGAATCTGTACAAAACGAGAATGCCTATG
 TGAGCGTGGTCACTAGTAACATAACCGTCGGTTCACTCCCGAGATCGCC
 GAGCGTCCGAAGGTGAGGACCAGGCAGGCCGGATGAACCTACTACTGGAC
 CCTATTGAAGCCAGGGGACACGATTATCTTCGAGGCAAACGGAACCTCA
 TAGCGCCGATGTACGCCCTTCGCCCTGAGCCGTGGCTTTGGATCGGGGATC
 ATCAGCTCTAACGCCCTCGATGCACGAATGTAATACCAATGCCAGACCCC
 ACTGGGTGCTATCAACTCGTCTTACCCTATCAAAATATACATCCGGTCA
 CCATAGGCGAGTGTCCCAAATATGTACAGACCATCATCACCACCATTAA
 TAG

SEQ ID NO:134 FOR PRIMER

AGGCAGATCTATGAAATCTTAGTCAACGTTGCCCTTGTTTTTATGGTCG
 TATACATTTCTTACATCTATGCGAGCCACAACGGAAGCTGTGCCGTCTG
 AAAG

SEQ ID NO:135 REV PRIMER

ACCTGCATGCCCTATTAATGGTGGTGATGATGGTGTCTGACATATTTGGGA
 CACTC

SEQ ID NO:136 HA0s PR8

ATGAAATCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTC
 TTACATCTATGCGGACACCATTTCGATTGGATACCATGCAACAACCTCAA
 CCGATACTGTTGATACCGTCTTGAAGAAGGTTACCGTCACGCACTCG
 GTCAACCTATTAGAGGATAGCCACAACGGAAGCTGTGCCGTCTGAAAGG
 CATAGCGCCACTGCAGCTCGGCAAATGTAACATCGCAGGTTGGCTCCTTG
 GTAACCCGGAGTGCGACCCCTCCTCCTGTACGATCCTGGAGTTATATC
 GTGGAGACCCCCAATAGCGAGAACGGAATTTGCTACCCAGGAGATTTTCAT
 AGACTACGAGGAGTTGCGCGAGCAGCTTTCTGTCTGTGAGCAGCTTCGAAA
 GGTTTCGAGATATTTCCGAAGGAGAGCAGCTGGCCGAATCATAACACTAAC
 GGTGTGACAGCCGCTGCAGTCATGAAGGAAAGAGTTTATTCTATCGCAA
 CCTGTGTGGTTGACGGAGAAAGAGGGCAGCTACCCTAAGTTGAAGAACT
 CCTATGTGAACAAAAAGGCAAGGAGGTTCTGGTGTGTGGGGCATACAC
 CACCCCCCAATAGCAAGGAGCAGCAGAATCTGTACAAAACGAGAATGC
 CTATGTGAGCGTGGTCACTAGTAACATAACCGTCGGTTCACTCCCGAGA
 TCGCCGAGCGTCCGAAGGTGAGGGACCAGGCAGGCCGGATGAACCTACTAC
 TGGACCTATTGAAGCCAGGGGACACGATTATCTTCGAGGCAAACGGA
 CCTCATAGCGCGATGTACGCCCTTCGCCCTGAGCCGTGGCTTTGGATCGG

-continued

GGATCATCACGTCTAACGCCTCGATGCACGAATGTAATACCAAATGCCAG
 ACCCCACTGGGTGCTATCAACTCGTCCTTACCCTATCAAAATATACATCC
 GGTCACCATAGGCAGTGTCCCAAATATGTGAGATCCGCCAAGTTGCCGA
 TGGTGACCGGCTCCGCAATATTCTAGTATTTCAGTCACGCGGCTTGTTT
 GGCGCCATCGCTGGTTTCATCGAAGCGGGTGGACAGGCATGATTGATGG
 CTGGTATGGCTATCACCACCAGAACGAGCAGGGCTCGGGCTACGCGGCTG
 ACCAGAAGTCGACTCAGAATGCCATCAATGGCATCAGCAATAAGGTGAAC
 ACGGTCATTGAAAAGATGAACATTCAATTTACAGCCGTAGGAAAAGAGTT
 TAATAAACTGGAAAAAGAATGGAGAATCTGAATAAGAAGTGGACGACG
 GATTTTTGGACATCTGGACGTACAACGCCGAGCTGCTGGTTCTGCTGGAA
 AATGAGCGAACACTGGATTTTCATGATTCTAACGTAAAGAATCTGTACGA
 GAAGGTGAAGTCCCAACTAAAGAATAATGCCAAGGAAATCGGAAATGGAT
 GCTTTGAGTTTACCACAAGTGCATAATGAGTGCATGGAATCCGTGCGA
 AATGGTACATACGATTACCCAAAGTACTCCGAAGAATCCAAGCTAAATCG
 CGAAAAGGTTGATGGTGTAAACTTGAATCCATGGGTATTTACCAACACC
 ATCATCACCACCATTAATAG

SEQ ID NO:137 HA1-1 PR8 (no his tag)
 ATGAAATCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGAGCCACAACGGAAAGCTGTGCCGTCTGAAAGGCATAG
 CGCCACTGCAGCTCGGCCAAATGTAACATCGCAGGTTGGCTCCTTGTTAAC
 CCGGAGTGCAGCCCCCTCCTCCCTGTACGATCCTGGAGTTATATCGTGA
 GACCCCCAATAGCGAGAACGGAATTTGCTACCCAGGAGATTTCATAGACT
 ACGAGGAGTTGCGCGAGCAGCTTTCGTCTGTGAGCAGCTTCGAAAGGTTT
 GAGATATTTCCGAAGGAGAGCAGCTGGCCGAATCATAACACTAACGGTGT
 GACAGCCGCCTGCAGTCATGAAGGAAAGAGTTTATTCTATCGCAACCTGC
 TGTGGTTGACGGAGAAAGAGGGCAGCTACCCCTAAGTTGAAGAACTCCTAT
 GTGAACAAAAAGGCAAGGAGGTTCTGGTGCTGTGGGGCATACACCACCC
 CCCCATAAGCAAGGAGCAGAGAATCTGTACCAAAAACGAGAATGCCTATG
 TGAGCGTGGTCACTAGTAATAACCGTCGGTTCACTCCCGAGATCGCC
 GAGCGTCCGAAGGTGAGGGACAGGCAGGCCGATGAACCTACTACTGGAC
 CCTATTGAAGCCAGGGGACACGATTATCTTCGAGGCAACGAAACCTCA
 TAGCGCCGATGTACGCCTTCGCCCTGAGCCGTGGCTTTGGATCGGGGATC
 ATCAGTCTAACGCCTCGATGCACGAATGTAATACCAAATGCCAGACCCC
 ACTGGGTGCTATCAACTCGTCCTTACCCTATCAAAATATACATCCGGTCA
 CCATAGGCGAGTGTCCCAAATATGTCAGATAATAG

SEQ ID NO:138 REV PRIMER
 ACCTGCATGCCTATTATCTGACATATTTGGGACACTCGCCTATGG

SEQ ID NO:139 HA1-1 NC
 ATGAAATCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTTCT
 TTACATCTATGCGAGCCACAATGGTAAGCTGTGTCTGTTGAAGGAATTG

-continued

CACCCCTCCAACCTGGGCAACTGTTCCGGTCGCTGGTTGGATCCTCGGAAAC
 CCAGAATGCGAGCTCCTTATCAGTAAGGAATCTTGGTCTTATATTGTCGA
 AACCCCGAACCCCGAGAACGGAACATGTACCCGGGTTACTTTGCTGATT
 ACGAAGAGCTTCGCGAGCAACTCAGCTCCGTATCTCTCTCGAGCGCTTC
 GAGATTTTTCCTAAAGAGTCCAGCTGGCCAAATCATACCGTCACCGCGGT
 GTCGGCCTCCTGTTCCTCACACGAAAGTCTAGCTTCTATAGAAATCTTC
 TCTGGCTGACGGGTAAGAATGGTCTTTACCCCAATTTGAGCAAGTCTTAC
 GTCAACAACAAAGAAAAGGAAGTTCTGGTATTGTGGGGTGTGCACCACCC
 TCCGAACATCGGCAATCAGCGCGCCCTGTATCACACAGAGAACCGGTATG
 TTTCCGTTGTCTCCTCACATTACTCGAGGCGCTTCACTCCTGAAATAGCT
 AAGCGTCCGAAAGTGCCTGACCAGGAGGACGTATCAACTATTATTGGAC
 GCTGTTGGAGCCAGGCGATACAATTATCTTCGAGGCTAACCGTAACCTTA
 TCGCTCCCTGGTACGCCCTTCGCCCTGTGCGGTGGTTTCGGTAGTGAATA
 ATCACTAGTAATGCTCCTATGGACGAGTGTGACGCTAAGTGCCAAACACC
 TCAGGGCGCTATCAATAGCTCCCTTCCATTCCAGAACGCTCCATCCGGTTA
 CCATTGGAGAGTGTCCAAAGTACGTGAGACACCATCATCACCATCACTAA
 TAG

SEQ ID NO:140 FOR PRIMER
 AGGCAGATCTATGAAATCTTAGTCAACGTTGCCCTTGTTTTTATGGTCG
 TATACATTTCTTACATCTATGCGAGCCACAATGGTAAGCTGTGTCTGTTG
 AAG

SEQ ID NO:141 REV PRIMER
 AGTCGCATGCCTATTAATGGTGGTATGATGGTGTCTCACGTACTTTGGA
 CACTCTCCAATGG

SEQ ID NO:142 HA0s NC
 AGATCTATGAAGTTTCTCGTGAACGTTGCACTGGTGTGTTATGGTAGTCTA
 TATATCGTACATTTATGCTGACACGATTTGCATCGGTTACCATGCGAACA
 ACTCCACGGATACCGTGGACACAGTGTGGAAAAGAACGTGACCGTCACG
 CACTCCGTAAACCTTCTGGAGGACAGCCACAATGGTAAGCTGTGTCTGTT
 GAAGGGAATTGCACCCCTCCAACCTGGGCAACTGTTCCGGTCGCTGGTTGGA
 TCCTCGGAAACCCAGAATGCGAGCTCCTTATCAGTAAGGAATCTTGGTCT
 TATATTGTGCAAAACCCGAACCCGAGAACGGAACATGTACCCGGGTTA
 CTTTGTGATTACGAAGAGCTTCGCGAGCAACTCAGCTCCGTATCTCTCT
 TCGAGCGCTTCGAGATTTTTCCTAAAGAGTCCAGCTGGCCAAATCATACC
 GTCACCGCGGTGTCGGCTCCTGTTCCTCACACGAAAGTCTAGCTTCTA
 TAGAAATCTTCTCTGGCTGACGGGTAAGAATGGTCTTTACCCCAATTTGA
 GCAAGTCTTACGTCAACAACAAAGAAAAGGAAGTTCTGGTATTGTGGGT
 GTGCACCAACCCCTCCGAACATCGGCAATCAGCGCGCCCTGTATCACACAGA
 GAACGCGTATGTTTCCGTTGTCTCCTCACATTACTCGAGGCGCTTCACTC
 CTGAAATAGCTAAGCGTCCGAAAGTGCCTGACCAGGAGGACGTATCAAC

-continued

TATTATTGGACGCTGTTGGAGCCAGGCGATACAATTATCTTCGAGGCTAA
 CGGTAACCTTATCGCTCCCTGGTACGCCTTCGCCCTGTCGCGTGTTTCG
 GTAGTGAATAATCACTAGTAATGCTCCTATGGACGAGTGTGACGCTAAG
 TGCCAAACACCTCAGGCGCTATCAATAGCTCCCTTCATTCCAGAACGT
 CCATCCGGTTACCATTGGAGAGTGTCCAAAGTACGTGAGATCGGCCAAAC
 TTCGCATGGTCACGGGTCTGCGCAACATCCCGTCAATCCAATCTAGGGGC
 CTCTTCGGCGCTATCGCGGTTTCATTGAGGGCGGTGGACTGGAATGGT
 TGACGGATGGTACGGCTATCATCACCAGAACAAGGTTCCGGTTACG
 CTGCTGACCAGAAATCTACTCAGAACGCGATCAATGGTATCAGGAACAAG
 GTGAACAGCGTCATTGAAAGATGAATACTCAGTTTACAGCCGTGGGCAA
 AGAGTTCAATAAACTCGAGAGACGTATGGAACCTCAATAAGAAGGTGG
 ATGACGGCTTCCTGGACATTTGGACTTACAACGCCGAGCTGCTGGTCTCG
 CTCGAGAACGAGAGAACCTTGACTTCCACGACAGCAACGTCAAGAACCT
 GTACGAGAAGGTGAAAAGTCAACTTAAAAACAATGCCAAGGAGATTGGTA
 ACGGCTGCTTCGAATTTACCACAAGTGTAATAATGAGTGCATGGAATCC
 GTTAAGAACGGCACCTACGATTACCTTAAATACTCAGAGGAGTCCAAGCT
 TAACCGCGAGAAGATCGACGGCGTAAACTGGAAGCATGGGCGTATACC
 AACACCATCATCACCATCACTAATAGGCATGC
 SEQ ID NO:143 HA1-1 VN
 ATGAAATTTCTTAGTCAACGTTGCCCTTGTTTTTATGGTCGTATACATTT
 TTACATCTATGCGGAGAAGAAACACAACGGTAAGCTTTGCGACTTGGATG
 GAGTCAAGCCCCCTACTACTTAGAGATTGAGTGTAGCGGTGGCTGTCTC
 GGTAACCCAATGTGCGATGAGTTTCATCAATGTTCCCGAATGGTCATATAT
 CGTCGAAAAAGCTAATCCTGTCAACGACCTGTGTACCCCGGTGATTTCA
 ATGACTATGAAGAACTGAAGCACCTGTCTCCCGCATCAACCATTTTCGAG
 AAAATCCAGATCATTTCCAAGAGTTCTGGTCTAGCCATGAGGCTAGTCT
 GGGTGTCTCATCCGCTGCCCATATCAGGGTAAAAGTTCTTTCTTTAGGA
 ACGTAGTATGGTTGATAAAGAAAACTCTACATACCCGACCATCAAGCGC
 TCTTACAACAATACGAACCAAGAGGATCTGCTTGTCTTTGGGAATCCA
 TCATCCTAATGATGCTGCCGAACAGACTAAGCTCTACCAAACCTTACCA
 CTTATATTTCCGTCGGCACCTCTACTCTGAACGAGCGCTTGTGCCAGG
 ATCGCTACGAGATCAAAAGTCAACGGCCAATCGGGCCGATGGAATTTCT
 CTGGACGATCCTGAAGCCTAATGACGTATCAACTTCGAGTCAATGGAA
 ACTTTATCGCTCCCGAGTACGCTTACAAGATCGTCAAGAAGGGCGACTCC
 ACGATTATGAAGTCAGAGTTGGAGTACGGCAACTGCAACACAAAGTGCCA
 AACTCCTATGGGCGCTATAAATTTCTCAATGCCGTTCCACAACATCCATC
 CGCTCAGCATCGGTGAGTGCCCGAAATATGTAAAGCACCATCACCACCAT
 CACTAATAG

-continued

SEQ ID NO:144 FOR PRIMER
 AGGCAGATCTATGAAATTTCTTAGTCAACGTTGCCCTTGTTTTTATGGTCG
 TATACATTCTTACATCTATGCGGAGAAGAAACACAACGTAAGCTTTG
 SEQ ID NO:145 REV PRIMER
 AGTCGCATGCCTATTATGGTGGTGATGATGGTGCTTTACATATTTCCGGC
 ACTCACCG
 SEQ ID NO:146 HA0s VN
 AGATCTATGAAATTTCTGGTTAATGTAGCCCTGGTGTATGGTAGTGTA
 CATTTTCATACATTTATGCTGATCAAATCTGCATTGGCTACCATGCCAACA
 ACAGCACCGAGCAAGTTGACACGATCATGGAGAAGAACGTAACCGTCACT
 CACGCTCAAGACATCCTGGAGAAGAAACACAACGGTAAGCTTTGCGCACTT
 GGATGGAGTCAAGCCCCCTACACTTAGAGATTGTAGTGTAGCCGGTTGGC
 TGCTCGGTAACCCAATGTGCGATGAGTTTCATCAATGTTCCCGAATGGTCA
 TATATCGTCGAAAAAGCTAATCCTGTCAACGACCTGTGTACCCCGGTGA
 TTTCAATGACTATGAAGAACTGAAGCACCTGTCTCCCGCATCAACCATT
 TCGAGAAAATCCAGATCATTTCCAAGAGTTCTGGTCTAGCCATGAGGCT
 AGTCTGGGTGTCTCATCCGCTGCCCATATCAGGGTAAAAGTTCTTTCTT
 TAGGAACGTAGTATGGTTGATAAAGAAAACTCTACATACCCGACCATCA
 AGCGCTCTTACAACAATACGAACCAAGAGGATCTGCTTGTCTTTGGGGA
 ATCCATCATCCTAATGATGCTGCCGAACAGACTAAGCTCTACCAAACCC
 TACCATTATATTTCCGTCGGCACCTCTACTCTGAACGAGCGCTTGTGTC
 CCAGGATCGCTACGAGATCAAAAGTCAACGGCCAATCGGGCCGATGGAA
 TTCTTCTGGACGATCCTGAAGCCTAATGACGCTATCAACTTCGAGTCAAA
 TGGAAACTTTATCGCTCCCGAGTACGCTTACAAGATCGTCAAGAAGGGCG
 ACTCCACGATTATGAAGTCAGAGTTGGAGTACGGCAACTGCAACACAAAG
 TGCCAAACTCCTATGGGCGCTATAAATTTCTCAATGCCGTTCCACAACAT
 CCATCCGCTCACGATCGGTGAGTGCCGAAATATGTAAAGTCGAATCGTC
 TCGTACTGGCGACAGGCGTGAGAAATAGTCCGCAACGTGAACGTCGTCGC
 AAGAAGAGAGGACTGTTTGGTGCCATGTCAGGCTTTATTGAGGGCGGCTG
 GCAGGCGATGGTTGACGGATGGTACGGCTACCACCATTCAAACGAGCAGG
 GATCTGGCTACGCCGTGACAAAGAAAGCACCCAAAAGGCCATTGATGGA
 GTGACGAATAAGGTGAATTCGATCATCGACAAAATGAACACGCAATTCGA
 AGCAGTGGGTGCGGAATTAATAACCTGGAGCGCGGTATCGAGAATCTGA
 ACAAGAAGATGGAAGACGGCTTTTGGATGTCTGGACATATAACGCTGAA
 TTGCTGGTCCCTCATGGAAAACGAGCGTACCCTTGATTTCCACGACAGCAA
 CGTTAAGAACCTCTACGACAAGGTCAGGCTCCAGCTCAGGGATAACGCCA
 AGGAATTGGGAAACGGATGCTTCGAGTTCTACCACAATGCGACAACGAG
 TGCATGGAGTCAGTCAGGAATGGTACCTACGACTACCCGCAATATTCGTA
 GGAGGCTCGCTTGAAGCGTGAGGAAATATCGGGTGTAAATGGAGAGTA
 TTGGAATCTACCAGCACCATCACCACCATCACTAATAGGCATGC

-continued

SEQ ID NO:147 HA1-1 IND
 ATGAAATTCCTTAGTCAACGTTGCCCTTGTTTTATGGTCGTATACATTTCTTACATCTATGCGGAGAAAACCCATAACGGTAAGTTGTGCGACCTTGACG
 GTGTAAAGCCCCCTGATCCTCCGTGACTGCAGTGTGTGCTGGTTGGCTTTTG
 GCGAACCCCATGTGTGACGAATTTATCAACGTGCCTGAATGGTCATACAT
 TGTAGAGAAGGCCAACCCACGAACGATCTCTGTTATCCCGGCAGCTTCA
 ATGACTATGAGGAACCTTAAGCACCTTCTGTACGTATCAACCACTTCGAA
 AAGATCCAGATCATCCCAGAGCTCCTGGAGCGACCACGAAGCCAGTTC
 GGGTGTGTCTTCCGCTTGCCCTACCTCGGTAGCCCTTCCTTCTCCGTA
 ACGTAGTGTGGCTGATCAAGAAGATAGCACTTACCCTACAATCAAAAAG
 TCGTATAACAATACTAACCAGAGGATCTGCTTGTACTCTGGGAATTCA
 TCATCCCAACGACGCGCGGAGCAGACCAGTTGTACCAGAACCCACCA
 CTTACATCTCCATCGGTACGTCCACACTGAATCAGCGTCTGGTCCCCAAG
 ATCGCAACCAGGTCCAAGGTTAACGGTCAGTCCCGTCGTATGGAGTTCTT
 CTGGACCATCCTGAAGCCCAACGACGCCATCAACTTCGAGTCCAACGGTA
 ATTTTCATTGCTCCGGAGTACGCCCTACAAGATAGTTAAGAAGGGTGATTCA
 GCGATCATGAAGTCGGAACCTTGAGTATGGCAACTGCAACACTAAATGCCA
 AACTCCAATGGGCGCTATCAACTCCGATATGCCATTCCATAACATCCACC
 CATTGACAATCGGTGAATGTCCCAAGTACGTGAAGCACCACCATCACCAT
 CACTAATAG

SEQ ID NO:148 FOR PRIMER
 AGGCAGATCTATGAAATTCCTTAGTCAACGTTGCCCTTGTTTTATGGTCG
 TATACATTTCTTACATCTATGCGGAGAAAACCCATAACGGTAAGTTGTG

SEQ ID NO:149 REV PRIMER
 AGTCGCATGCCTATTAATGGTGGTGATGATGGTCTTACGTACTTGGGA
 CATTACCGATTG

SEQ ID NO:150 HA0s IND
 AGATCTATGAAGTTCCTGGTCAATGTAGCCTTGGTATTATGGTAGTCTA
 TATCTCGTACATTTACGACAGACAGATTTGTATTGGATATACGCTAACA
 ACAGCACAGAGCAGGTAGATACTATTATGGAGAAAAATGTTACCGTCACT
 CACGCCCAGGACATCCTGGAGAAAACCCATAACGGTAAGTTGTGCGACCT
 TGACGGTGTAAAGCCCCCTGATCCTCCGTGACTGCAGTGTGCTGGTTGGC
 TTTTGGGCAACCCCATGTGTGACGAATTTATCAACGTGCCTGAATGGTCA
 TACATTGTAGAGAAGGCCAACCCACGAACGATCTCTGTTATCCCGGCAG
 CTTCAATGACTATGAGGAACCTTAAGCACCTTCTGTACGTATCAACCACT
 TCGAAAAGATCCAGATCATCCGAGAGCTCCTGGAGCGACCACGAAGCC
 AGTTCGGGTGTGTCTTCCGCTTGCCCTACCTCGGTAGCCCTTCTCTCTT
 CCGTAACGTAGTGTGGCTGATCAAGAAGATAGCACTTACCCTACAATCA
 AAAAGTCGTATAACAATACTAACCAGAGGATCTGCTTGTACTCTGGGGA
 ATTCATCATCCCAACGACGCGCGGAGCAGACCAGGTTGTACCAGAACCC
 CACCACCTTACATCTCCATCGGTACGTCCACACTGAATCAGCGTCTGGTCC

-continued

CCAAGATCGCAACCAGGTCCAAGGTTAACGGTCAGTCCGGTCGTATGGAG
 TTCTTCTGGACCATCCTGAAGCCCAACGACGCCATCAACTTCGAGTCCAA
 CGGTAATTTTCATTGCTCCGGAGTACGCCCTACAAGATAGTTAAGAAGGGTG
 ATTCAGCGATCATGAAGTCGGAACCTTGAGTATGGCAACTGCAACACTAAA
 TGCCAAACTCCAATGGGCGCTATCAACTCCAGTATGCCATTCCATAACAT
 CCACCCATTGACAATCGGTGAATGTCCCAGTACGTGAAGAGCAACAGGT
 TGGTATTGGCCACCGGTTTGAGAAACAGCCCCAGAGAGAGTCGCGTCGT
 AAAAGCGCGGCTTGTTTCGGAGCCATCGCTGGCTTCATAGAGGGTGGTTG
 GCAGGGAATGGTCGATGGTTGGTATGGTTATCATCATTTCCAACGAGCAGG
 GAAGTGGTTACGCCCGCCGACAAAGAATCGACCCAGAAGGCTATTGACGGC
 GTCACAAACAAAGTAAACTCTATCATTTGATAAGATGAACACCCAGTTCGA
 GGCTGTAGGTAGAGAATTCACAACCTCGAAAGACGTATTGAGAACCTGA
 ACAAGAAAATGGAGGATGGCTTCTCGGACGTGTGGACCTACAATGCTGAG
 CTGTTGGTCTTATGGAGAACGAGCGTACCCTCGATTTCATGACTCAAA
 CGTGAAAGAACCTGTATGACAAGGTGCGTTTGCAACTGAGGGACACGCAA
 AGGAGCTTGAAACCGTTGTTTTCGAATTTATCATAAGTCGCGACAATGAG
 TGTATGGAGTCGATTAGAAATGGCACGTACAACCTACCCTCAATACAGCGA
 AGAAGCTCGTCTCAAACGTGAGGAAATCAGCGGCGTCAAGCTCGAATCAA
 TCGGTACCTATCAGCACCACCATCACCATCACTAATAGGCATGC

SEQ ID NO:151 STF2.HA1-1His (PR8)
 MKFLVNVALVFMVVYISYIYAAQVINTNSLSLLTQNNLNKSQSALGTAIE
 RLSSGLRTNSAKDDAAGQAIA NRFTANIKGLTQASRNANDGISIAQTTEG
 ALNEINNQLQVRRELAVQSANSTNSQSDLSIQAEITQRLNEIDRVSGQT
 QFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
 VKDTAVTTKAYANNGTLLDVSGLDAAIKAAATGGTNGTASVTGGAVKFDA
 DNNKYFVTIGGFTGADAANKGDYEVNVATDGTVTLLAAGATKTTMPAGATT
 KTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYTDKNGKTI
 EGGYALKAGDKYYAADYDEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
 KTEVVTIDGKTYNASKAAGHDFKAQPELAEEAAKT TENPLQKIDAALAQV
 DALRSDLGAVQNRFN SAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
 QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSSHNGKLCRLKGIAPLQL
 GKCNIAGWLLGNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEELR
 EQLSSVSSFERFEIFPKESSWPNNHTNGVTAACSHGKSSFYRNLLWLTE
 KEGSYPKLN SYVNKKGKEVLVLWGIHHPNSKEQONLYQENAYVSVVT
 SNYNRRFTPEIAERP KVRDQAGRMNYWTLLKPGDTIIFEANGNL IAPMY
 AFALSRGFGSGIITSNASMHECNTKCQTPLGAINSSLPYQNIHPVTIGEC
 PKYVRHHHHHH

-continued

SEQ ID NO:152 ngSTF2.HA1-1His (PR8)
 MKFLVNVALVFMVVYISYIYAAQVINTNSLSLLTQNNLQKSQSALGTAIE
 RLSSGLRINSKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
 ALNEINNQLQRVRELAVQSAQSTNSQSDLDLSIQAEITQRLNEIDRVSGQT
 QFNGVKVLAQDNTLTITQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
 VKDTAVTTKAYANQGTTLDVSGLDAAIKAATGGTQGTASVTGGAVKFDA
 DNNKYFVTIGGFTGADAAKNGDYEYVNVATDGTVTLAAGATKTTMPAGATT
 KTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYTDKNGKTI
 EGGYALKAGDKYYAADYDEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
 KTEVVTIDGKTYQASKAAGHDFKAQPELAEEAAKTENPLQKIDAALAQV
 DALRSDLGAVQNRFNSAITNLGNTVNLSEARSRIEDSDYATEVSQMSRA
 QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSHNGKLCRLKGIAPLQL
 GKNCIAGWLLGNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEELR
 EQLSSVSSFERFEIFPKESSWPNNHTNGVTAACSHEGKSSFYRNLLWLTE
 KEGSYPKLKNSYVKKGKEVLVLWGIHPPNSKEQQONLYQENAYVSVVT
 SNYNRRFTPEIAERP KVRDQAGRMNYWTLLKPGDTIIFEANGNLIAPMY
 AFALSRGFGSGIITSQASMHECNTKCTPLGAIQSSLPYQNIHPVTIGEC
 PKYVRHHHHH

SEQ ID NO:153 STF2.HA1-2His (PR8)
 MKFLVNVALVFMVVYISYIYAAQVINTNSLSLLTQNNLQKSQSALGTAIE
 RLSSGLRINSKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
 ALNEINNQLQRVRELAVQSANSTNSQSDLDLSIQAEITQRLNEIDRVSGQT
 QFNGVKVLAQDNTLTITQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
 VKDTAVTTKAYANNGTTLDVSGLDAAATKAAIGGTNGTASVTGGAVKFDA
 DNNKYFVTIGGFTGADAAKNGDYEYVNVATDGTVTLAAGATKTTMPAGATT
 KTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYTDKNGKTI
 EGGYALKAGDKYYAADYDEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
 KTEVVTIDGKTYNASKAAGHDFKAQPELAEEAAKTENPLQKIDAALAQV
 DALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
 QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSGKIAPQLGKNCIAGWLL
 GNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEELREQLSSVSFE
 RFEIFPKESSWPNNHTNGVTAACSHEGKSSFYRNLLWLTEKESYPKLKN
 SYVKKGKEVLVLWGIHPPNSKEQQONLYQENAYVSVVTSNYNRRFTPE
 IAERP KVRDQAGRMNYWTLLKPGDTIIFEANGNLIAPMYAFALSRGFGS
 GIITSHHHHHH

SEQ ID NO:154 ngSTF2.HA1-2His (PR8)
 MKFLVNVALVFMVVYISYIYAAQVINTNSLSLLTQNNLQKSQSALGTAIE
 RLSSGLRINSKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
 ALNEINNQLQRVRELAVQSAQSTNSQSDLDLSIQAEITQRLNEIDRVSGQT
 QFNGVKVLAQDNTLTITQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD

-continued

VKDTAVTTKAYANQGTTLDVSGLDAAIKAATGGTQGTASVTGGAVKFDA
 DNNKYFVTIGGFTGADAAKNGDYEYVNVATDGTVTLAAGATKTTMPAGATT
 KTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYTDKNGKTI
 EGGYALKAGDKYYAADYDEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
 KTEVVTIDGKTYQASKAAGHDFKAQPELAEEAAKTENPLQKIDAALAQV
 DALRSDLGAVQNRFNSAITNLGNTVNLSEARSRIEDSDYATEVSQMSRA
 QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSGKIAPQLGKNCIAGWLL
 LGNPECDPLLPVRSWSYIVETPNSENGICYPGDFIDYEELREQLSSVSFF
 ERFEIFPKESSWPNNHTNGVTAACSHEGKSSFYRNLLWLTEKESYPKLKN
 NSYVKKGKEVLVLWGIHPPNSKEQQONLYQENAYVSVVTSNYNRRFTP
 EIAERP KVRDQAGRMNYWTLLKPGDTIIFEANGNLIAPMYAFALSRGFG
 SGITSHHHHHH

SEQ ID NO:155 STF2.HA1-2mutHis (PR8)
 MKFLVNVALVFMVVYISYIYAAQVINTNSLSLLTQNNLQKSQSALGTAIE
 RLSSGLRINSKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
 ALNEINNQLQRVRELAVQSANSTNSQSDLDLSIQAEITQRLNEIDRVSGQT
 QFNGVKVLAQDNTLTITQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
 VKDTAVTTKAYANNGTTLDVSGLDAAIKAATGGTNGTASVTGGAVKFDA
 DNNKYFVTIGGFTGADAAKNGDYEYVNVATDGTVTLAAGATKTTMPAGATT
 KTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYTDKNGKTI
 EGGYALKAGDKYYAADYDEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
 KTEVVTIDGKTYNASKAAGHDFKAQPELAEEAAKTENPLQKIDAALAQV
 DALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
 QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSGKAAPLQLGKNCIAGWLL
 LGNPECDPLLPVRSWSDIAETPNSENGICYPGDFIDYEELREQLSSVSFF
 ERFEIFPKESSWPNNHTNGVTAACSHEGKSSFYRNLLWLTEKESYPKLKN
 NSYVKKGKEVLVLWGIHPPNSKEQQONLYQENAYVSVVTSNYNRRFTP
 EIAERP KVRDQAGRMNYWTLLKPGDTIIFEANGNLIAPMYAFALSRGFG
 SGITSHHHHHH

SEQ ID NO:156 ngSTF2.HA1-2mutHis (PR8)
 MKFLVNVALVFMVVYISYIYAAQVINTNSLSLLTQNNLQKSQSALGTAIE
 RLSSGLRINSKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
 ALNEINNQLQRVRELAVQSAQSTNSQSDLDLSIQAEITQRLNEIDRVSGQT
 QFNGVKVLAQDNTLTITQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
 VKDTAVTTKAYANQGTTLDVSGLDAAIKAATGGTQGTASVTGGAVKFDA
 DNNKYFVTIGGFTGADAAKNGDYEYVNVATDGTVTLAAGATKTTMPAGATT
 KTEVQELKDTPAVVSADAKNALIAGGV DATDANGAELVKMSYTDKNGKTI
 EGGYALKAGDKYYAADYDEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
 KTEVVTIDGKTYQASKAAGHDFKAQPELAEEAAKTENPLQKIDAALAQV
 DALRSDLGAVQNRFNSAITNLGNTVNLSEARSRIEDSDYATEVSQMSRA

-continued

QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSKGAAPQLGKCNIAGWLL
GNPECDPLLPVRSWDIAETPNSENGICYPGDFIDYEELREQLSSVSSFE
RFEIFPKESSWPNNHNTNGVTAACSHGKSSFYRNLLWLTEKEGSYPKLKN
SYVNKKGKEVLVLWGIHHPNSKEQQNLYQENENAYVSVVTSNYNRRFTPE
IAERP KVRDQAGRMNYYWTLLKPGDTIIFEANGNLIAPMYAFALSRGPGS
GIITSHHHHHH

SEQ ID NO:157 STF2.HA1-3His (PR8)
MKFLVNVALVFMVYIISYIYAAQVINTNSLSLLTQNNLNKSQSALGTAIE
RLSSGLRINS AKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
ALNEINNQLQVRRELAVQSANSTNSQSDLDISQAEITQRLNEIDRVSGQT
QFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
VKDTAVTTKAYANNGTTLDSVGLDDAAIKAATGGTNGTASVTGGAVKFDA
DNNKYFVTIGGFTGADAAKNGDYEENVATDGTVTLAAGATKTTMPAGATT
KTEVQELKDTPAVVSADAKNALIAGGVDATDANGAELVKMSYTDKNGKTI
EGGYALKAGDKYYAADYEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
KTEVVTIDGKTYNASKAAGHDFKAQPELAEAAKTENPLQKIDAALAQV
DALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSNSENGICYPGDFIDYEEL
REQLSSVSSFERFEIFPKESSWPNNHNTNGVTAACSHGKSSFYRNLLWLT
EKEGSYPKLKNSYVNKKGKEVLVLWGIHHPNSKEQQNLYQENENAYVSVV
TSNYNRRFTPEIAERP KVRDQAGRMNYYWTLLKPGDTIIFEANGNLIAPM
YAFALSRGHHHHH

SEQ ID NO:158 STF2.HA1-3mutHis (PR8)
MKFLVNVALVFMVYIISYIYAAQVINTNSLSLLTQNNLNKSQSALGTAIE
RLSSGLRINS AKDDAAGQAIANRFTANIKGLTQASRNANDGISIAQTTEG
ALNEINNQLQVRRELAVQSANSTNSQSDLDISQAEITQRLNEIDRVSGQT
QFNGVKVLAQDNTLTIQVGANDGETIDIDLKQINSQTLGLDSLNVQKAYD
VKDTAVTTKAYANNGTTLDSVGLDDAAIKAATGGTNGTASVTGGAVKFDA
DNNKYFVTIGGFTGADAAKNGDYEENVATDGTVTLAAGATKTTMPAGATT
KTEVQELKDTPAVVSADAKNALIAGGVDATDANGAELVKMSYTDKNGKTI
EGGYALKAGDKYYAADYEATGAIAKAKTTSYTAADGTTKTAANQLGGVDG
KTEVVTIDGKTYNASKAAGHDFKAQPELAEAAKTENPLQKIDAALAQV
DALRSDLGAVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSRA
QILQQAGTSVLAQANQVPQNVLSLLASGSGSGSNSENEICYPGDFIDKEE
LREQLSSVSSFERFEIFPKESSWPNNHNTNGVTAACSHGKSSFYRNLLWLT
TEKEGSYPKLKNSYVNKKGKEVLVLWGIHHPNSKEQQNLYQENENAYVSVV
VTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLLKPGDTIIFEANGNLIAP
MYAAALSRGHHHHH

-continued

SEQ ID NO:159 STF2.HA1-2 (IND)
MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINS AKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRRELAVQSA
NSTNSQSDLDISQAEITQRLNEIDRVSGQTQPNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTTLDV
SGLDDAAIKAATGGTNGTASVTGGAVKFADANNKYFVTIGGFTGADAAKN
GDYEENVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
ALIAGGVDATDANGAELVKMSYTDKNGKTIIEGGYALKAGDKYYAADYEDEA
TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
DFKAQPELAEAAKTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
VLSLLAGVKPLILRDCSVAGWLLGNFMCDEFINVPESYIVEKANPTNDL
CYPGFSNDYEELKHLISRINHFEKIQIIPKSSWSDEASSGVSSACPYLG
SPSFERNVVLIKKNSTYPTIKKSYNNTNQEDLLVLWGIHHPNDAAEQTR
LYQNPTTYISIGTSTLNQRLVPKIAITRSKVNGQSGRMEFFWTILKPNDAI
NFESNGNFIAPYAYKIVKKGDSAIMKSE

Drosophila
SEQ ID NO:160 STF2A.HA0shis
ATGGCACAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGCGGATT
GCTAACCGTTTCACCGCGAACATCAAAGGCTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCAGTCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGGTATCCGCCAGACTCAGTTCAACG
GCGTGAAAGTCTTGGCGCAGGACACACCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCATGGAGCGCCGGTGGATCCTGCTA
GCCATGGACCGAAAACCGCTGCAGAAAATTGATGCCGCGCTGGCGCAG
GTGGATGCGCTGCGCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTC
TGCTATCACCAACCTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTA
GCCGTATCGAAGATTCGACTACGCGACCGAAGTTTCCAACATGTCTCGC
GCGCAGATTTTGCAGCAGCGCGGTACTTCCGTTCTGGCGCAGGCTAACCA
GGTCCCGCAGAACGTGCTGTCTCTGTACGTGAATTCTCTAGATATCCAG
CACAGTGGCGCGCGCTGCACACAATATGTATAGGCTACCATGCGAACAAT
TCAACCGACACTGTTGACACAGTACTCGAGAAGAATGTGACAGTGACACA
CTCTGTTAACTGCTCGAAGACAGCCACAACGGAAAACCTATGTAGATTAA
AAGGAATAGCCCCACTACAATTGGGGAAATGTAACTCGCCGGATGGCTT
TTGGGAAACCCAGAATGCGACCCACTGCTTCCAGTGAGATCATGGTCCTA

-continued

CATTGTAGAAACACCAAACCTCTGAGAAATGGAATATGTTATCCAGGAGATT
 TCATCGACTATGAGGAGCTGAGGGAGCAATGAGCTCAGTGTCATCATTC
 GAAAGATTGCAATATTTCCCAAAGAAAGCTCATGGCCCAACCACAACAC
 AAACGGAGTAACGGCAGCATGCTCCCATGAGGGGAAAAGCAGTTTTTACA
 GAAATTTGCTATGGCTGACGGAGAAGGAGGCTCATACCCAAAGCTGAAA
 AATTCTTATGTGAACAAAAAGGAAAGAGTCCTTGTACTGTGGGGTAT
 TCATCACCCGCCTAACAGTAAGGAACAACAGAATCTCTATCAGAATGAAA
 ATGCTTATGTCTCTGTAGTGACTTCAAATTATAACAGGAGATTTACCCCG
 GAAATAGCAGAAAGACCCAAAGTAAGAGATCAAGCTGGGAGGATGAACTA
 TTACTGGACCTTGCTAAAACCCGGAGACACAATAATATTTGAGGCAAATG
 GAAATCTAATAGCACCAATGTATGCTTTTCGCACTGAGTAGAGGCTTTGGG
 TCCGGCATCATCACCTCAAACGCATCAATGCATGAGTGAACACGAAGTG
 TCAAACACCCCTGGGAGCTATAAACAGCAGTCTCCCTTACCAGAATATAC
 ACCCAGTCACAATAGGAGAGTGCCCAAATACGTCAGGAGTGCCAAATTG
 AGGATGGTTACAGGACTAAGGAACATTCGGTCCATTCAATCCAGAGGTCT
 ATTTGGAGCCATTGCCGGTTTTATTGAAGGGGATGGACTGGAATGATAG
 ATGGATGGTATGGTTATCATCATCAGAATGAACAGGGATCAGGCTATGCA
 GCGGATCAAAAAAGCACAAAAATGCCATTAAACGGGATTACAAACAAGGT
 GAACACTGTTATCGAGAAAATGAACATTCATTCACAGCTGTGGGTAAAG
 AATTCAACAAATTAGAAAAAGGATGGAAAATTTAAATAAAAAAGTTGAT
 GATGGATTTCTGGACATTTGGACATATAATGCAGAATTGTTAGTTCTACT
 GGAAAATGAAAGGACTCTGGATTTCATGACTCAAATGTGAAGAATCTGT
 ATGAGAAAGTAAAAAGCAATTAAAGAATAATGCCAAAGAAATCGGAAAT
 GGATGTTTTGAGTTCTACCACAAGTGTGACAATGAATGCATGGAAGGTG
 AAGAAATGGGACTTATGATTATCCCAAATATTCAGAAGAGTCAAAGTTGA
 ACAGGGAAAAGGTAGATGGAGTGAATTTGGAATCAATGGGGATCTATCAG
 ACGCGTACCGGTATCATCATCACCATCACCATTGA

SEQ ID NO:161 FOR PRIMER
 ATTCTCCGGCGCGCTCGACACAATATGTATAGGCTACC

SEQ ID NO:162 REV PRIMER
 AGTCTTGCGCGCGCTATTAATGGTGATGGTGATGATGCTGATAGATCCC
 CATTGATTCC

SEQ ID NO:163 HA0s PR8 template
 GACACAATATGTATAGCTACCATGCGAACAATTCACCGACACTGTTGA
 CACAGTACTCGAGAAGATGTGACAGTGACACACTCTGTAACTGCTCG
 AAGACAGCCACAACGGAAAACATGTAGATTAAAGGAATAGCCCCACTA
 CAATTGGGGAAATGTAACATCGCCGGATGGCTTTTGGGAAACCCAGAATG
 CGACCCACTGCTTCCAGTGAGATCATGGTCCTACATTGTAGAAACACCAA
 ACTCTGAGAATGGAATATGTTATCCAGGAGATTTATCGACTATGAGGAG
 CTGAGGGAGCAATTGAGCTCAGTGTATCATTTGAAAGATTGCAAAATAT

-continued

TCCCAAAGAAAGCTCATGGCCCAACCACAACACAAACGGAGTAACGGCAG
 CATGCTCCCATGAGGGGAAAAGCAGTTTTTACAGAAATTTGCTATGGCTG
 ACGGAGAAGGAGGGCTCATACCCAAAGCTGAAAAATTTCTATGTGAACAA
 AAAAGGGAAAGAAGTCCCTTGTAAGTGTGGGTATTTCATCACCCGCCCTAACA
 GTAAGGAACAACAGAATCTCTATCAGAATGAAAATGCTTATGTCTCTGTA
 GTGACTTCAAATTATAACAGGAGATTTACCCCGAAATAGCAGAAAGACC
 CAAAGTAAGAGATCAAGCTGGGAGGATGAACATTACTGGACCTTGCTAA
 AACCCGGAGACACAATAATATTTGAGGCAATGGAATCTAATAGCACCA
 ATGTATGCTTTTCGCACTGAGTAGAGGCTTTGGGTCCGGCATCATCACCTC
 AAACGCATCAATGCATGAGTGTAAACACGAAGTGTCAAACACCCCTGGGAG
 CTATAAACAGCAGTCTCCCTTACCAGAATATACACCAGTCACAATAGGA
 GAGTGCCCAAAATACGTCAGGAGTGCCAAATTTGAGGATGGTTACAGGACT
 AAGGAACATTCGCTCCATTCAATCCAGAGGTCTATTGGAGCCATTGCCG
 GTTTTATTGAAGGGGGATGGACTGGAATGATAGATGGATGGTATGTTAT
 CATCATCAGAATGAACAGGGATCAGGCTATGCAGCGGATCAAAAAAGCAC
 ACAAATGCCATTAAACGGGATTACAAACAGGTGAACACTGTTATCGAGA
 AAATGAACATTCAATTCACAGCTGTGGGTAAAGAATTCAACAAATTAGAA
 AAAAGGATGGAATAATTAATAAAAAAGTTGATGATGGATTTCTGGACAT
 TTGGACATATAATGCAGAATTGTTAGTTCTACTGGAAAATGAAAGGACTC
 TGGATTTCCATGACTCAAATGTGAAGAATCTGTATGAGAAAGTAAAAAGC
 CAATTAAAGAATAATGCCAAAGAAATCGGAAATGGATGTTTTGAGTTCTA
 CCACAAGTGTGACAATGAATGCATGGAAAGTGAAGAAATGGGACTTATG
 ATTATCCCAATATTCAGAAGAGTCAAAGTTGAACAGGGAAAAGGTAGAT
 GGAGTGAAATTTGGAATCAATGGGGATCTATCAG

SEQ ID NO:164 STF2A.HA0s
 ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGCGATT
 GCTAACCGTTTACCAGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCTGAACGAAATCGACCGGTATCCGCGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCATGGAGCGCGGTGGATCTTGCTA
 GCCCATGGACCGAAAACCCGCTGCAGAAAATTTGATGCCGCGCTGGCGCAG
 GTGGATGCGCTGCGCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTC
 TGCTATCACCAACCTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTA

-continued

GCCGTATCGAAGATTCGACTACGCGACCGAAGTTTCCAACATGTCTCGC
 GCGCAGATTTTGCAGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACCA
 GTTCCCGCAGAACGTGCTGTCTCTGTTACGTGAATTCTCTAGATATCCAG
 CACAGTGGCGGCCGCTCGACACAATATGTATAGGCTACCATGCGAACAAT
 TCAACCGACACTGTTGACACAGTACTCGAGAAGAATGTGACAGTGACACA
 CTCTGTTAACCTGCTCGAAGACAGCCACAACGGAAAACATGTAGATTAA
 AAGGAATAGCCCCACTACAATTGGGGAAATGTAACATCGCCGGATGGCTT
 TTGGGAAACCCAGAATGCGACCCACTGCTTCCAGTGAGATCATGGTCTTA
 CATTGTAGAAACACCAAACTCTGAGAATGGAATATGTTATCCAGGAGATT
 TCATCGACTATGAGGAGCTGAGGGAGCAATTGAGCTCAGTGTCATCATT
 GAAAGATTTCGAAATATTTCCCAAAGAAAGCTCATGGCCCAACCAACAC
 AAACGGAGTAACGGCAGCATGCTCCCATGAGGGGAAAAGCAGTTTTCACA
 GAAATTTGCTATGGCTGACGGAGAAGGAGGCTCATACCCAAAGCTGAAA
 AATTCTTATGTGAACAAAAAGGAAAGAGTCCTTGTACTGTGGGGTAT
 TCATCACCCGCTTAACAGTAAGGAACAACAGAATCTCTATCAGAATGAAA
 ATGCTTATGTCTCTGTAGTGACTTCAATTATAACAGGAGATTTACCCCG
 GAAATAGCAGAAAGACCCAAAGTAAGAGATCAAGCTGGGAGGATGAACTA
 TTACTGGACCTTGCTAAACCCGGAGACACAATAATATTTGAGGCAATG
 GAAATCTAATAGCACCAATGTATGCTTTCGCACCTGAGTAGAGGCTTTGGG
 TCCGGCATCATCACCTCAAACGCATCAATGCATGAGTGAACACGAAGTG
 TCAAACACCCCTGGGAGCTATAAACAGCAGTCTCCCTTACCAGAATATAC
 ACCCAGTCACAATAGGAGAGTGCCCAAAATACGTGAGGAGTGCCAAATTG
 AGGATGGTTACAGGACTAAGGAACATTCCGTCATTCAATCCAGAGGTCT
 ATTTGGAGCCATTGCCGTTTATTTGAAGGGGATGGACTGGAATGATAG
 ATGGATGGTATGGTTATCATCATCAGAATGAACAGGGATCAGGCTATGCA
 GCGGATCAAAAAAGCACACAAAATGCCATTAAACGGGATTACAAACAAGGT
 GAACACTGTTATCGAGAAAATGAACATTCAATTACAGCTGTGGGTAAAG
 AATTCAACAAATTAGAAAAAGGATGGAAAATTTAAATAAAAAAGTTGAT
 GATGGATTTCTGGACATTTGGACATATAATGCAGAATTGTTAGTTCTACT
 GGAAAATGAAGGACTCTGGATTTCATGACTCAAATGTGAAGAATCTGT
 ATGAGAAAGTAAAAAGCAATTAAGAATAATGCCAAAGAAATCGGAAAT
 GGATGTTTTGAGTTCTACCACAAGTGTGACAATGAATGCATGGAAAGTGT
 AAGAAATGGGACTTATGATTATCCAAATATTAGAAGAGTCAAAGTTGA
 ACAGGGAAAAGGTAGATGGAGTGAAATTGGAATCAATGGGATCTATCAG
 TAG
 SEQ ID NO:165 FOR PRIMER
 ATTCTCCGGCGCGCTCGACACAATATGTATAGGCTACC
 SEQ ID NO:166 REV PRIMER
 AGTCTTGGCGCCGCTATTAATGGTGATGGTGATGCTGATAGATCCC
 CATTGATTCC

-continued

SEQ ID NO:167 HA0s.STF2A
 GACACAATATGTATAGGCTACCATGCGAACAATTCAACCGACACTGTTGA
 CACAGTACTCGAGAAGAATGTGACAGTGACACACTCTGTTAACCTGCTCG
 AAGACAGCCACAACGGAAAACATGTAGATTAAAAGGAATAGCCCCACTA
 CAATTGGGGAAATGTAACATCGCCGGATGGCTTTTGGGAAACCCAGAATG
 CGACCCACTGCTTCCAGTGAGATCATGGTCTCATATTGTAGAAACACCAA
 ACTCTGAGAATGGAATATGTTATCCAGGAGATTTTCATCGACTATGAGGAG
 CTGAGGGAGCAATTGAGCTCAGTGTCATCATTCGAAAGATTTCGAAATATT
 TCCCAAAGAAAGCTCATGGCCCAACCAACACAACCGAGTAACGGCAG
 CATGCTCCCATGAGGGGAAAAGCAGTTTTTACAGAAATTGCTATGGCTG
 ACGGAGAAGGAGGGCTCATACCCAAAGCTGAAAAATTCTTATGTGAACAA
 AAAAGGAAAAGAGTCCCTGTACTGTGGGTATTTCATCACCCGCTTAACA
 GTAAGGAACAACAGAATCTCTATCAGAATGAAAATGCTTATGTCTCTGTA
 GTGACTTCAAATTATAACAGGAGATTTACCCCGAAATAGCAGAAAGACC
 CAAAGTAAGAGATCAAGCTGGGAGGATGAACATTACTGGACCTTGCTAA
 AACCCGAGACACAATAATATTTGAGGCAATGGAATCTAATAGCACCA
 ATGTATGCTTTCGCACTGAGTAGAGGCTTTGGGTCCGGCATCATCACCTC
 AAACGCATCAATGCATGAGTGAACACGAAGTGTCAAACACCCCTGGGAG
 CTATAAACAGCAGTCTCCCTTACCAGAATATACACCCAGTCACAATAGGA
 GAGTGCCCAAAATACGTGAGGAGTGCCAAAGTTGAGGATGGTTACAGGACT
 AAGGAACATTCGCTCCATTCAATCCAGAGGTCTATTGGAGCCATTGCCG
 GTTTTATTGAAGGGGGATGGACTGGAATGATAGATGGATGGTATGGTTAT
 CATCATCAGAATGAACAGGGATCAGGCTATGCAGCGGATCAAAAAAGCAC
 ACAAATGCCATTAAACGGGATTACAAACAGGTGAACACTGTTATCGAGA
 AAATGAACATTCAATTCACAGCTGTGGGTAAAGAATTCAACAAATTAGAA
 AAAAGGATGAAAAATTTAAATAAAAAAGTTGATGATGGATTCTTGACAT
 TTGGACATATAATGCAGAATTGTTAGTTCTACTGAAAAATGAAGGACTC
 TGGATTTCATGACTCAAATGTGAAGAATCTGTATGAGAAAGTAAAAAGC
 CAATTAAAGAATAATGCCAAAGAAATCGGAAATGGATGTTTTGAGTTCTA
 CCACAAGTGTGACAATGAATGCATGGAAAGTGAAGAAATGGGACTTATG
 ATTATCCCAAATATTAGAAGAGTCAAAGTTGAACAGGGAAGGTAGAT
 GGAGTGAAATTGAATCAATGGGGATCTATCAGGAATTCTCTAGATATCC
 AGCACAGTGGCGGCCGCTCGCACAGTAATCAACACTAACAGTCTGTGCG
 TGCTGACCCAGAATAACCTGAACAAATCCAGTCCGCACTGGGCACCGCT
 ATCGAGCGTCTGTCTTCTGGTCTGCGTATCAACAGCGCGAAAGACGATGC
 GGCAGGTCAGGCGATTGCTAACCGTTTCACCGCAACATCAAAGGTCTGA
 CTCAGGCTTCCCGTAACGCTAACGACGGTATCTCCATTGCGCAGACCACT
 GAAGGCGCGCTGAACGAAATCAACAACAACCTGCAGCGTGTGCGTGAAC
 GCGGTTTCACTCTGCTAACAGCACCAACTCCAGTCTGACCTCGACTCCA

-continued

TCCAGGCTGAAATCACCCAGCGCCTGAACGAAATCGACCGTGTATCCGGC
 CAGACTCAGTTCAACGGCGTGAAAGTCCTGGCGCAGGACAACACCCCTGAC
 CATCCAGGTTGGCGCCAACGACGCTGAAACTATCGATATCGATCTGAAGC
 AGATCAACTCTCAGACCCCTGGGTCTGGACTCACTGAACGTGCATGGAGCG
 CCGGTGGATCCTGCTAGCCCATGGACCGAAAACCCGCTGCAGAAATTGA
 TGCCGCGCTGGCGCAGGTGGATGCGCTGCGCTCTGATCTGGGTGCGGTAC
 AAAACCGTTTCAACTCTGCTATCACCAACCTGGGCAATACCGTAAACAAT
 CTGTCTGAAGCGGTAGCCGTATCGAAGATTCCGACTACGCGACCGAAGT
 TTCCAACATGTCTCGCGCGCAGATTTTGCAGCAGGCGCGTACTTCCGTTT
 TGGCGCAGGCTAACCAGGTCCCGCAGAACGTGCTGTCTCTGTTACGTAA
 TAG

SEQ ID NO:168 FOR PRIMER
 AGGCAAGATCTGACACAATATGTATAGGCTACC

SEQ ID NO:169 REV PRIMER
 AGTCAGACGCGTCTATTAACTAACAGAGACAGCAC

SEQ ID NO:170 HA0s.his
 GACACAATATGTATAGGCTACCATGCGAACAATTCAACCGACACTGTTGA
 CACAGTACTCGAGAAGAAATGTGACAGTGACACACTCTGTTAACTTGCTCG
 AAGACAGCCACAACGGAACATATGTAGATTAAAAGGAATAGCCCCACTA
 CAATTGGGGAAATGTAACATCGCCGGATGGCTTTTGGGAAACCCAGAATG
 CGACCCACTGCTTCCAGTGAGATCATGGTCTACATTGTAGAAACACCAA
 ACTCTGAGAATGGAATATGTTATCCAGGAGATTTTCATCGACTATGAGGAG
 CTGAGGGAGCAATTGAGCTCAGTGTATCATATTGAAAGATTGCAAAATATT
 TCCCAAAGAAAGCTCATGGCCCAACCAACAACACGAGTAACGGCAG
 CATGCTCCCATGAGGGGAAAAGCAGTTTACAGAAATTTGCTATGGCTG
 ACGGAGAAGGAGGGCTCATACCCAAAGCTGAAAAATTCTTATGTGAACAA
 AAAAGGGAAAGAAGTCCTTGTACTGTGGGTATTTCATCACCCGCTAACA
 GTAAGGAACAACAGAATCTCTATCAGAATGAAATGCTTATGTCTCTGTA
 GTGACTTCAAATTATAACAGGAGATTTACCCCGGAAATAGCAGAAAGACC
 CAAAGTAAGAGATCAAGCTGGGAGGATGAACTATTACTGGACCTTGCTAA
 AACCCGGAGACACAATAATATTGAGGCAATGGAATCTAATAGCACCA
 ATGTATGCTTTTCGCACTGAGTAGAGGCTTTGGGTCCGGCATCATCACCTC
 AAACGCATCAATGCATGAGTGTAACACGAAGTGTAACACCCCTGGGAG
 CTATAAACAGCAGTCTCCCTTACCAGAATATACACCCAGTCACAATAGGA
 GAGTGCCCAAAATACGTCAGGAGTGCCAAATTGAGGATGGTTACAGGACT
 AAGGAACATTCCGTCCATTCAATCCAGAGGTCTATTGAGGCCATTGCGG
 GTTTTATTGAAGGGGGATGGACTGGAATGATAGATGGATGGTATGGTTAT
 CATCATCAGAATGAACAGGGATCAGGCTATGCAGCGGATCAAAAAGCAC
 ACAAATGCCATTAACGGGATTACAAACAAGGTGAACACTGTTATCGAGA
 AAATGAACATTCAATTACAGCTGTGGGTAAGAATTCAACAAATTAGAA

-continued

AAAAGGATGGAAAATTTAAATAAAAAAGTTGATGATGGATTCTTGGACAT
 TTGGACATATAATGCAGAATTGTTAGTTCTACTGAAAAATGAAAGGACTC
 TGGATTTCATGACTCAAATGTGAAGAATCTGTATGAGAAAGTAAAAAGC
 CAATTAAGAATAATGCCAAAGAAATCGGAAATGGATGTTTTGAGTTCTA
 CCACAAGTGTGACAATGAATGCATGGAAAGTGAAGAAATGGGACTTATG
 ATTATCCCAATATTTCAGAAGAGTCAAAGTTGAACAGGGAAAAGGTAGAT
 GGAGTGAAATTGGAATCAATGGGGATCTATCAGACGCGTACCGGTCATCA
 TCACCATCACCATTGA

SEQ ID NO:171 FOR PRIMER
 ATTCTCCGCGCGCCGCTCGACACAATATGTATAGGCTACC

SEQ ID NO:172 REV PRIMER
 AGTCTTGCGGCCGCTATTAAATGGTGTGGTGTATGCTGATAGATCCC
 CATTGATTCC

SEQ ID NO:173 HA0s
 GACACAATATGTATAGGCTACCATGCGAACAATTCAACCGACACTGTTGA
 CACAGTACTCGAGAAGAAATGTGACAGTGACACACTCTGTTAACTTGCTCG
 AAGACAGCCACAACGGAACATATGTAGATTAAAAGGAATAGCCCCACTA
 CAATTGGGGAAATGTAACATCGCCGGATGGCTTTTGGGAAACCCAGAATG
 CGACCCACTGCTTCCAGTGAGATCATGGTCTACATTGTAGAAACACCAA
 ACTCTGAGAATGGAATATGTTATCCAGGAGATTTTCATCGACTATGAGGAG
 CTGAGGGAGCAATTGAGCTCAGTGTATCATTCGAAAGATTGCAAAATATT
 TCCCAAAGAAAGCTCATGGCCCAACCAACAACACGAGTAACGGCAG
 CATGTCTCCCATGAGGGGAAAAGCAGTTTACAGAAATTTGCTATGGCTG
 ACGGAGAAGGAGGGCTCATACCCAAAGCTGAAAAATTTCTTATGTGAACAA
 AAAAGGGAAAGAAGTCTTGTACTGTGGGTATTTCATCACCCGCTAACA
 GTAAGGAACAACAGAATCTCTATCAGAATGAAATGCTTATGTCTCTGTA
 GTGACTTCAAATTATAACAGGAGATTTACCCCGGAAATAGCAGAAAGACC
 CAAAGTAAGAGATCAAGCTGGGAGGATGAACTATTACTGGACCTTGCTAA
 AACCCGGAGACACAATAATATTGAGGCAATGGAATCTAATAGCACCA
 ATGTATGCTTTTCGCACTGAGTAGAGGCTTTGGGTCCGGCATCATCACCTC
 AAACGCATCAATGCATGAGTGTAACACGAAGTGTAACACCCCTGGGAG
 CTATAAACAGCAGTCTCCCTTACCAGAATATACACCCAGTCACAATAGGA
 GAGTGCCCAAAATACGTCAGGAGTGCCAAATTGAGGATGGTTACAGGACT
 AAGGAACATTCCGTCCATTCAATCCAGAGGTCTATTGAGGCCATTGCGG
 GTTTTATTGAAGGGGGATGGACTGGAATGATAGATGGATGGTATGGTTAT
 CATCATCAGAATGAACAGGGATCAGGCTATGCAGCGGATCAAAAAGCAC
 ACAAATGCCATTAAACGGGATTACAAACAAGGTGAACACTGTTATCGAGA
 AAATGAACATTCAATTACAGCTGTGGGTAAGAATTCAACAAATTAGAA
 AAAAGGATGGAAAATTTAAATAAAAAAGTTGATGATGGATTCTTGGACAT
 TTGGACATATAATGCAGAATTGTTAGTTCTACTGAAAAATGAAAGGACTC

-continued

TGGATTTCATGACTCAAATGTGAAGAATCTGTATGAGAAAGTAAAAAGC
 CAATTAAAGAATAATGCCAAAGAAATCGGAAATGGATGTTTGGATTCTA
 CCACAAGTGTGACAATGAATGCATGGAAGTGTAAAGAAATGGGACTTATG
 ATTATCCCAAATATTGAGAAGAGTCAAAGTTGAACAGGGAAAAGGTAGAT
 GGAGTGAAATTGGAATCAATGGGGATCTATCAG
 SEQ ID No:174 FOR PRIMER
 AGGCAAGATCTGACACAATATGTATAGGCTACC
 SEQ ID No:175 REV PRIMER
 AGTCAGACGCGTCTATTACGTAACGAGACAGACAGCAC
 SEQ ID No:176 HA0sHis (PR8)
 DTICIGYHANNSTDVTDVLEKNVTVTHSVNLLSDSHNGKLCRLKGIAPL
 QLGKCNIAGWLLGNPECDPLLVPVRSWSYIVETPNSENGICYPGDFIDYEE
 LREQSSVSSFERFEIFPKESSWPNNHNTNGVTAACSHGKSSFYRNLLWL
 TEKEGSYPKLKNSVTNKKGKEVLVLWGIHHPNSKEQQNLYQENENAYVSV
 VTSNYNRRFTPEIAERP KVRDQAGRMNYYWTLKPGDTIIFEANGNLIAP
 MYAFALSRGFGSGIITSNASMHECNTKCQTPLGAINSSLPYQNIHPVTIG
 ECPKYVRS AKLRMVTGLRNIPSIQSRGLFGAIAGFIEGGWTGMIDGWYGY
 HHQNEQSGSYAADQKSTQNAINGITNKVNTVIEKMNIQFTAVGKEFNKLE
 KR MENLNKKVDDGFLDIWTYNAELLVLENERTLDFHDSNVKNLYEKVKS
 QLKNNAKEIGNGCFEFYHKCDNECMESVRNGTYDYPKYSEESKLNREKVD
 GVKLESMGIYQH HHHHH
 SEQ ID No:177 STF2A.HA0s (PR8)
 MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINS AKDDAAGQAI
 ANRFTANIKGLTQASRNDNGISIAQTTEGALNEINNNLQVRRELAVQSA
 NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVHGAPVDPASPWTENPLQKIDAALAQ
 VDALSRLDGVANQNRFSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSR
 AQILQQAGTSVLAQANQVPQNVLSLLREFSRYPQWRPLDTICIGYHANN
 STDVTDVTVLEKNVTVTHSVNLLSDSHNGKLCRLKGIAPLQLGKCNIAWL
 LGNPECDPLLVPVRSWSYIVETPNSENGICYPGDFIDYEELREQSSVSSF
 ERFEIFPKESSWPNNHNTNGVTAACSHGKSSFYRNLLWLTEKEGSYPKLK
 NSYVNKKGKEVLVLWGIHHPNSKEQQNLYQENENAYVSVVTSNYNRRFTP
 EIAERP KVRDQAGRMNYYWTLKPGDTIIFEANGNLIAPMYAFALSRGFG
 SGIITSNASMHECNTKCQTPLGAINSSLPYQNIHPVTIG ECPKYVRS AKL
 RMVTGLRNIPSIQSRGLFGAIAGFIEGGWTGMIDGWYGYHHQNEQSGGYA
 ADQKSTQNAINGITNKVNTVIEKMNIQFTAVGKEFNKLEKR MENLNKKVD
 DGFLDIWTYNAELLVLENERTLDFHDSNVKNLYEKVKS QLKNNAKEIGN
 GCFEFYHKCDNECMESVRNGTYDYPKYSEESKLNREKVDGVKLESMGIYQ
 SEQ ID No:178 STF2
 ATGGCACAAGTAATCAACACTAACAGTCTGTGCGTGCTGACCCAGAATAA
 CCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT

-continued

CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGCGGATT
 GCTAACCGGTTTCACCGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCGAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGCGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGCTGATGCCGCCAAAAAT
 GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTTGCGGC
 TGGCGCAACTAAAACCACAATGCCTGCTGGTGCGACAACATAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
 GCCTTAATGTCTGGCGGCGTTGACGCTACCGATGTCTAATGGCGCTGAGTT
 GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCCGCAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCTACAAGTTTACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCAT
 GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACCACCGA
 AAACCCGCTGCAGAAAAATTGATGCCGCGTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTACGT
 SEQ ID No:179 HA1-1his (PR8)
 MKFLVNVALVFMVVYISYIYASHNGKLCRLKGIAPLQLGKCNIAGWLLGN
 PECDDLVPVRSWSYIVETPNSENGICYPGDFIDYEELREQSSVSSFERF
 EIFPKESSWPNNHNTNGVTAACSHGKSSFYRNLLWLTEKEGSYPKLKNSY
 VNKKGKEVLVLWGIHHPNSKEQQNLYQENENAYVSVVTSNYNRRFTPEIA
 ERP KVRDQAGRMNYYWTLKPGDTIIFEANGNLIAPMYAFALSRGFGSGI
 ITSNASMHECNTKCQTPLGAINSSLPYQNIHPVTIG ECPKYVRHHHHHH
 SEQ ID No:180 HA1-1his (VN)
 MKFLVNVALVFMVVYISYIYAEKKHNGKLCDLGVKPLILRDCSVAGWLL
 GNP M CDEFIN VPEWSYIVEKANPVNDLCYPGDFNDYEELKHLISRINHFE

-continued

KIQIIPKSSWSSEASLGVSSACPYQKSSFFRNVVWLIIKKNSTYPTIKR
SYNNTNQEDLLVLWGIHHPNDAAEQTKLYQNPTTYISVGTSTLNQRLVPR
IATRISKVNGQSGRMEFFWTILKPDAINFESNGNFIAPYAYKIVKKGDS
TIMKSELEYGNCNTKCQTPMGAINSSMPFHNHPLTIGECPKYVKHHHHH
H

SEQ ID No:181 HA1-1his (IND)

MKFLVNVALVFMVVIISYIYAEKTHNGKLCDLGDKPLILRDCSVAGWLL
GNPMCEDEFINVPESWYIVEKANPTNDLCYPGSFNDYEELKHLISRINHFE
KIQIIPKSSWSDEASSGVSSACPYLGSPSFFRNVVWLIIKKNSTYPTIKK
SYNNTNQEDLLVLWGIHHPNDAAEQTRLYQNPTTYISIGTSTLNQRLVPR
IATRISKVNGQSGRMEFFWTILKPDAINFESNGNFIAPYAYKIVKKGDS
AIMKSELEYGNCNTKCQTPMGAINSSMPFHNHPLTIGECPKYVKHHHHH
H

SEQ ID No:182 HA1-1his (NC)

MKFLVNVALVFMVVIISYIYASHNGKLCCLLKIAPLQLGNCVAGWILGN
PECELLISKESWSYIVETPNPENGTCYPGYFADYEELREQLSSVSSFERF
EIFPKESSWPNHTVTGVSASCSHNGKSSFYRNLLWLTGKNGLYPNLSKSY
VNNKEKEVLVLWGVHHPNIGNQRLYHTENAYSVSVSSHYRRFTPEIA
KRPKVRDQEGRINYWTLLPEPGDTIIFEANGNLIAPWYAFALSRGFGSGT
ITSNAPMDECDACKQTPQGAINSSLPFQNVHPVTIGECPKYVRHHHHHH

SEQ ID No:183 HA1-1 (MAL)

ACCACTACCCCAACCAATCTCACTTCGCAAACTGAAAGGCACTGAAAC
CCGTGGCAAGCTGTGTCCGAAGTGTCTGAACTGCACCGATCTGGACGTCG
CACTGGGTCGTCCGAAATGTACTGGTAACATTCCTCCGCGCGTGTCTCC
ATCCTGCATGAAGTGCCTCCAGTGACCTCCGGCTGTTTCCGATTATGCA
TGATCGTACTAAATCCGTCAGCTGCCGAACCTGCTGCGTGGTTACGAAC
ACATTCGTCTGTCCACCCATAACGTTATCAACGCGGAAAACGCGCCGGGC
GGTAGCTATAAAATCGGTACCTCTGGTTCTTGCCCGAACGTGACTAACGG
TAACGGCTTCTTTGCAACCATGGCCTGGGCGGTCCCGAAAAACGACAACA
ACAAGACCGCGACCAATTCCTTGACCATCGAAGTCCCGTATATCTGCACC
GAAGGTGAAGATCAAATCACGGTTTGGGGCTTCCACTCCGACAACGAGGC
ACAAATGGCGAACTGTACGGTGACAGCAAACCGCAAAAATTCCTAGCT
CCGCTAACGGTGTACCACCCACTACGTTTCCAGATCGGTGGTTTCCCA
AACCAGACCGAAGATGGTGGTCTGCCGCGATCCGGTCGCATCGTTGTAGA
TTATATGGTGCAGAAAAGCGGTAAAACCGGTACCATCACCTACCAGCGTG
GCATCTGTCTGCCGAGAAAGTTTGGTGGCCTTCCGGTCGTAGCAAAGTA
ATCAAAGGTTCCCTGCCGCTGATCGGTGAAGCAGACTGCCTGCACGAGAA
ATACGGCGGTCTGAACAAAAGCAAGCCGTACTATACCGGCAACATGCGA
AAGCAATTGGTAACGTGCCAATTGGGTGAAATAGTAG

-continued

SEQ ID No:184 STF2.HA1-1 (MAL)

ATGGCACAAGTAATCAACACTAACAGTCTGTCTGCTGCTGACCCAGAATAA
CCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAAGCGATT
GCTAACCGTTTCACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTGTCT
AACAGCACCAACTCCCAGTCTGACCTGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGCCAGACTCAGTTCAACG
GCGTGAAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCGTGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACCAATGG
TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTGGCGGC
TGGCGCAACTAAAACCACAATGCCTGCTGGTGCAGCAACTAAAACAGAAG
TACAGGAGTTAAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
GCCTTAATGTCTGGCGCGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGTCTTATACCGATAAAAATGGTAAGACAATTGAAGGCGGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCCGCAGATTACGATGAAGCG
ACAGGAGCAATTAAAGCTAAAACCTACAAGTTATACTGCTGCTGACGGCAC
TACCAAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCAT
GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGGA
AAACCCGCTGCAGAAAATTGATGCCGCGTGGCGCAGGTGGATGCGCTGC
GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAC
CTGGGCAATACCGTAAACAATCTGTCTGAAGCGGTAGCCGTATCGAAGA
TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
GTGCTGTCTCTGTTAGCGACCACTACCCCAACCAATCTCACTTCGCAAA
CCTGAAAGGCACTGAAACCCGTGGCAAGCTGTCTCCGAAGTGTCTGAACT
GCACCGATCTGGACGTCGCACTGGGTGCTCCGAAATGTACTGGTAACATT
CCGTCCGCGCGTGTCTCCATCTGTCATGAAGTGGTCCAGTGACCTCCGG
CTGTTTTCCGATTATGCATGATCGTACTAAAATCCGTCAGTGCCGAAC
TGCTGCGTGGTTACGAACACATTCTGTCTGCCACCCATAACGTTATCAAC
GCGGAAAACGCGCCGGCGGTAGCTATAAAATCGGTACCTCTGTTCTTGC
CCCGAACGTGACTAACGGTAACGGCTTCTTTGCAACCATGGCCTGGGCGG

-continued

TCCCGAAAAACGACAACAACAAGACCGCGACCAATTCCCTGACCATCGAA
GTCCCGTATATCTGCACCGAAGGTGAAGATCAAATCACGGTTTGGGGCTT
CCACTCCGACAACGAGGCACAAATGGCGAAACTGTACGGTGACAGCAAAC
CGCAAAAATTCACTAGCTCCGCTAACGGTGTTACCACCCACTACGTTTCC
CAGATCGGTGGTTTCCCAAACCAGACCGAAGATGGTGGTCTGCCCGCAGTC
CGGTGCGATCGTTGTAGATTATATGGTGCAGAAAAGCGGTAAAACCGGTA
CCATCACCTACCAGCGTGGCATCCTGCTGCCGCAGAAAGTTTGGTGCCT
TCCGGTCGTAGCAAAGTAATCAAAGGTTCCCTGCCGCTGATCGGTGAAGC
AGACTGCCTGCACGAGAAATACGGCGGTCTGAACAAAAGCAAGCCGTACT
ATACCGCGCAACATGCGAAAGCAATTGGTAACGTCCCAATTGGGTGAAA
TAGTAG

SEQ ID No:185 HA1-2 (MAL)

AAAGGCACTGAAACCCGTGGCAAGCTGTGTCCGAAGTGTCTGAACTGCAC
CGATCTGGACGTGCGACTGGGTCTGCCGAATGTACTGGTAACATTCGCT
CCGCGCGTGTCTCCATCCTGCATGAAGTGCCTCCAGTGACCTCCGGCTGT
TTTCCGATTATGCATGATCTGACTAAAATCCGTGAGCTGCCGAACCTGCT
GCGTGGTTACGAACACATTCGTCTGTCCACCCATAACGTTATCAACGCGG
AAAACGCGCGGGCGGTAGCTATAAAATCGGTACTCTGGTTCTTGCCCG
AACGTGACTAACGGTAACGGCTTCTTTGCAACCATGGCCTGGGCGGTCCC
GAAAAACGACAACAACAAGACCGCGACCAATTCCCTGACCATCGAAGTCC
CGTATATCTGCACCGAAGGTGAAGATCAAATCACGGTTTGGGGCTTCCAC
TCCGACAACGAGGCACAAATGGCGAAACTGTACGGTGACAGCAAACCGCA
AAAATTCAGTCTCCGCTAACGGTGTACACCCACTACGTTTCCGAGA
TCGGTGGTTTCCCAAACCAGACCGAAGATGGTGGTCTGCCGCAGTCCGGT
CGCATCGTTGTAGATTATATGGTGCAGAAAAGCGGTAAAACCGGTACCAT
CACCTACCAGCGTGGCATCCTGCTGCCGCAGAAAGTTTGGTGCCTTCCG
GTCGTAGCAAAGTAATCAAAGGTTGATAG

SEQ ID No:186 STF2.HA1-2 (MAL)

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGGAAAGACGATGCGGCAGGTGACGGCGATT
GCTAACCGTTTACC CGCGAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAACGGCGGTTTCAGTCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
GCGTGAAAGTCTTGGCGCAGGACAACACCTTGACCATCCAGGTTGGCGCC
AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGCTACTACACTGGATGTA

-continued

TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAACCGGTGGTGCGGTTAAATTTGACGCAGATAATAACA
AGTACTTTGTTACTATTGGTGGCTTTACTGGTGCTGATGCCGCCAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
TGGCGCAACTAAAACCACAATGCCTGCTGGTGCAGCAACTAAAACAGAAG
TACAGGAGTTAAAAGATACACCGGCAGTTGTTTCAGCAGATGCTAAAAAT
GCCTTAATGTCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGTCTTATACCATAAAAATGGTAAGACAATTGAAGGCGGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCCGAGATTACGATGAAGCG
ACAGGAGCAATTAAAGCTAAAACCTACAAGTTTACTGCTGCTGACGGCAC
TACCAAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
TCGTTACTATCTGACGGTAAAACCTACAATGCCAGCAAGCCCGCTGGTCAT
GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA
AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCCGTGC
GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAC
CTGGGCAATACCGTAAACAATCTGTCTGAAGCGGTAGCCGTATCGAAGA
TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
GTGCTGTCTCTGTTAGCGAAAGGCACTGAAACCCGTGGCAAGCTGTGTCC
GAAGTGTCTGAACTGCACCGATCTGGACGTGCGACTGGGTCGTCCGAAAT
GTACTGGTAACATTCCGTCGCGCGTGTCTCCATCCTGCATGAAGTGCCT
CCAGTGACCTCCGGCTGTTTTCCGATTATGCATGATCTGACTAAAATCCG
TCAGCTGCCGAACCTGCTGCGTGGTTACGAACACATTCTGCTGTCCACCC
ATAACGTTATCAACGCGGAAAACGCGCCGGGCGGTAGCTATAAAATCGGT
ACCTCTGGTTCTTGCCCGAACGTGACTAACGGTAACGGCTTCTTTGCAAC
CATGGCCTGGGCGTCCCGAAAACGACAACAACAAGACCGCGACCAATT
CCCTGACCATCGAAGTCCCGTATATCTGCACCGAAGGTGAAGATCAAATC
ACGGTTTGQGGCTTCCACTCCGACAACGAGGCACAAATGGCGAAACTGTA
CGGTGACAGCAAAACCGCAAAAATTCACTAGCTCCGCTAACGGTGTACCA
CCCACTACGTTTCCAGATCGGTGGTTTCCCAAACCAGACCGAAGATGGT
GGTCTGCCGCGAGTCCGGTCGCATCGTTGTAGATTATATGGTGCAGAAAAG
CGGTAAAACCGGTACCATCACCTACCAGCGTGGCATCCTGCTGCCGCAGA
AAGTTTGGTGCCTTCCGGTCGTAGCAAAGTAATCAAAGGTTGATAG

SEQ ID No:187 HA1-2 (SH)

AAAGGCACTCGTACCCGCGGTAAGCTGTGCCCGGACTGCCTGAACTGTAC
CGATCTGGATGTTGCACTGGGTGCTCCGATGTGCGTGGTACCACCCCGT
CTGCCGAAAGCCAGCATCCTGCACGAAGTTCGCCCGGTTACTTCCGGTTGT
TTCCCGATTATGCATGATCGTACCAAAATTCGTGAGCTGCCAAACCTGCT
GCGTGGCTATGAAAACATTCGTCTGTCCACTCAAACGTAATCGATGCAG

-continued

AAAAAGCGCTGGGTGGCCGTATCGTCTGGGTACAGCGGCTCCTGCCCG
 AACGCGACGAGCAAAAGCGGCTTCTTCGCCACCATGGCTTGGGCGTTCC
 GAAAGACAACAACAAAACGCTACGAACCCGCTGACCGTCGAAGTCCCGT
 ACATCTGCACCGAAGGCGAAGATCAGATCACTGTGTGGGGCTTCCACAGC
 GATGATAAGACCCAGATGAAAAATCTGTACGGTGACTCCAACCCGAGAA
 ATTCACTCTTCTGCTAACGGTGTAAACGACCCACTACGTTTCTCAGATCG
 GTGGTTTCCCGGACCAGACGGAAGATGGCGGTCTGCCTCAGTCCGGCCGC
 ATCGTAGTTGATTACATGGTCCAGAAACCGGGTAAGACTGGTACCATTGT
 TTACGACGCTGGTGTACTGTGCCGAGAAAGTCTGGTGTGCTTCCGGCC
 GTTCCAAGGTCATTAAGGCTGATA

SEQ ID No:188 STF2.HA1-2 (SH)

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTTCAGCGATT
 GCTAACCGTTTACC CGCAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTTCAGTCTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGTGAAATCAC
 CCAGCGCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCCCTGGCGCAGGACAACACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAACCTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGACGCTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGCAGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTTCGGC
 TGGCGCAACTAAAACCAATGCCTGTGCTGCGACAATAAAACAGAAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTTCAGCAGATGTAAAAAT
 GCCTTAATTGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCTACAAGTTATACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCAT
 GATTTCAAAGCACACACGAGCTGGCGGAAGCAGCGCTAAAACCCGGA
 AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACC AAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCTAGCCGTATCGAAGA

-continued

TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTTAGCGAAAGGCACTCGTACCCCGGTAAGCTGTGCCC
 GGACTGCCTGAACGTACCGATCTGGATGTTGCACTGGGTCTCCGATGT
 GCGTTGGTACCACCCCGTCTGCGAAAGCCAGCATCTGCACGAAGTTTCGC
 CCGGTTACTTCCGTTGTTTCCCGATTATGCATGATCGTACCAAAATTCG
 TCAGTGCCTAAACCTGCTGCGTGGCTATGAAAAATTCTGCTGTCCACTC
 AAAACGTAATCGATGCAGAAAAAGCGCTGGGTGGCCCGTATCGTCTGGGT
 ACCAGCGGCTCCTGCCCGAACGCGACGAGCAAAAGCGGCTTCTTCGCCAC
 CATGGCTTGGGCGGTTCCGAAAGACAACAACAAAACGCTACGAACCCGC
 TGACCGTCAAGTCCCGTACATCTGCACCGAAGGCGAAGATCAGATCACT
 GTGTGGGGCTTCCACAGCGATGATAAGACCCAGATGAAAAATCTGTACGG
 TGACTCCAACCCGAGAAATTCACCTCTTCTGCTAACGGTGTAAACGACCC
 ACTACGTTTCTCAGATCGGTGGTTTCCCGACCGACGGAAGATGGCGGT
 CTGCCTCAGTCCGGCCGATCGTAGTTGATTACATGGTCCAGAAACCGGG
 TAAGACTGGTACCATTGTTTACCAGCGTGGTGTACTGCTGCCGAGAAAG
 TCTGTTGTGCTTCCGGCCGTTCCAAGGTCATTAAGGCTGATAG

SEQ ID No:189 HA1-2 (Lee)

AAAGGCACTCAGACCCGTGGCAAGCTGTGTCCGAACTGTTTCAACTGCAC
 CGATCTGGACGTTGCACTGGGTGCTCCGAAATGCATGGGTAAACATCCCGT
 CTGCGAAGGTAAGCATCTGCACGAAGTTAAACCGGTAACAGCGGCTGT
 TTCCCGATCATGCACGACAAAATAAAATTCTGTCAGCTGCCGAACTGCT
 GCGTGGTTATGAGAACATTCTGCTGTCTACCTCTAATGTTATCAACGCGG
 AGACTGCACCGAGGTGGCCATACAAAGTAGGTACCAGCGGTTCTGTCCG
 AACGTTGCGAATCGTAACGGCTTCTTCAACACTATGGCGTGGGTTATCCC
 GAAAGATAACAATAAAACTGCAATTAACCCGGTAACGTAGAAAGTTCCGT
 ACATCTGCTCCGAAGGCGAGGACAGATTACGGTATGGGGCTTTCACAGC
 GACGATAAAACCCAGATGGAGCGTCTGTACGGTGACTCTAACCCGAGAA
 ATTCACCTCCTCCGGAACGGCGTTACACCCACTATGTTTCTCAGATCG
 GCGGTTTCCCGAATCAGACCGAAGACGAAGCGCTGAAGCAGTCCGGCGGT
 ATTGTTGTAGACTACATGGTTGAGAAGCCGGGCAAAACTGGTACCATTGT
 ATACCAGCGCGCATCTGCTGCCGAGAAAGTTTGGTGGCTTCCGGTC
 GTAGCAAAGTAATCAAAGTTGATAG

SEQ ID No:190 STF2.HA1-2 (Lee)

ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTTCAGGCGATT
 GCTAACCGTTTACC CGCAACATCAAAGGCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG

-continued

AAATCAACAACAACCTGCAGCGTGTGCGTGAAC TGGCGGTT CAGTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCC TGGCGCAGGACAACACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGCTCTTGATGATGTCAGCTATTAAAGCGGCTACGGGTGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGT TACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTGCGGC
 TGGCGCAACTAAAACCACAATGCCTGTGTGGTGCAGACAATAAACAGAAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTT CAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCTACAAGTTATACTGCTGTGACGGCAC
 TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTTCAT
 GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGCA
 AAACCCGCTGCAGAAAATTGATGCCGCGTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACC AAC
 CTGGGCAATACCGTTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGCTCGCGCGCAGATTCTGC
 AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACCGGTCCCAGCAAG
 GTGCTGTCTCTGTTAGCGAAAGGCACTCAGACCCGTGGCAAGCTGTGTCC
 GAACTGTTTCAACTGCACCGATCTGGACGTTGCACTGGGTCTGCCGAAAT
 GCATGGGTAACATCCCGTCTGCGAAGGTAAGCATCCTGCACGAAGTTAAA
 CCGGTAACGAGCGGCTGTTTCCCGATCATGCACGACAAAACCTAAAATTCG
 TCAGCTGCCGAACCTGCTGCGTGGTTATGAGAACATTCGTCTGTCTACCT
 CTAATGTTATCAACGCGGAGACTGCACCGGTGGCCATACAAAGTAGGT
 ACCAGCGGTTCCGTGTCGGAACGTTGCGAATCGTAACGGCTTCTTCAACAC
 TATGGCGTGGGTTATCCCGAAAGATAACAATAAACTGCAATTAACCCGG
 TAACTGTAGAAGTTCCGTACATCTGCTCCGAAGCGAGGACCAGATTACG
 GTATGGGGCTTTTACAGCGACGATAAAACCCAGATGGAGCGTCTGTACGG
 TGACTCTAACCCGAGAAATTCACCTCCTCCGCGAACGGCGTTACCACCC
 ACTATGTTTCTCAGATCGGCGGTTTCCCGAATCAGACCGAAGACGAAGGC
 CTGAAGCAGTCCGGCGTATTGTTGTAGACTACATGGTT CAGAAGCCGGG

-continued

CAAAAC TGGTACCATTGTATACCAGCGCGGCATCCTGCTGCCGAGAAAG
 TTTGGTGCCTTCCGGTCGTAGCAAAGTAATCAAAGGTTGATAG
 SEQ ID No:191 HA1-2 (Ohio)
 AAAGGCAC TAAACCCGTGGCAAGCTGTGTCCGAAGTGTCTGAACTGCAC
 CGATCTGGACGTCGCACTGGGTCGTCCGAAATGTACTGGTAACATTCCGT
 CCGCGGAAGTCTCCATCCTGCATGAAGTGCCTCCAGTGACCTCCGGCTGT
 TTTCCGATTATGCATGATCGTACTAAAAATCCGTCAGCTGCCGAACCTGCT
 GCGTGGTTACGAACACATTCTGTCTGTCCACCATAACGTTATCAACGCGG
 AAAAGGCGCCGGGCGGTCCCTATAAAATCGGTACCTCTGGTTCTTGCCCCG
 AACGTGACTAACGGTAACGGCTTCTTTTGCAACCATGGCCTGGGCGGTCCC
 GAAAAACGACAACAACAAGACCGCGACCAATTCCTTGACCATCGAAGTCC
 CGTATATCTGCACCGAAGGTGAAGATCAAATCACGATTTGGGGCTTCCAC
 TCCGACAGCGAGACACAAATGGCGAAACTGTACGGTGACAGCAAAACCGCA
 AAAATTCACTAGCTCCGCTAACGGTGT TACCACCCACTACGTTTCCGAGA
 TCGGTGGTTTCCCAAACCAGACCGAAGATGGTGGTCTGCCGAGTCCGGT
 CGCATCGTTGTAGATTATATGGTGCAGAAAAGCGGTAAAACCGGTACCAT
 CACCTACCAGCGTGGCATCCTGCTGCCGAGAAAGTTGGTGCCTTCCG
 GTCGTAGCAAAGTAATCAAAGGTTGATAG
 SEQ ID No:192 STF2.HA1-2 (Ohio)
 ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
 CCTGAACAAATCCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGT CAGGCGATT
 GCTAACCGTTTACCAGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCACCGTGTGCGTGAAC TGGCGGTT CAGTCTGCT
 AACAGCACCAACTCCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGT TACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
 TGGCGCAACTAAAACCACAATGCCTGTGTGGTGCAGACAATAAACAGAAAG
 TACAGGAGTTAAAGATACACCGGCAGTTGTTT CAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG

-continued

ACAGGAGCAATTAAAGCTAAACCTACAAGTTATACTGCTGCTGACGGCAC
TACCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAAGCCGCTGGTCAT
GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACCCGA
AAACCCGCTGCAGAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
AGCAGGCCGGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCAGAGAAC
GTGCTGTCTCTGTAGCGAAAGGCACTAAAACCCGTGGCAAGCTGTGTCC
GAAGTGCTGAACTGCACCGATCTGGACGTGCGACTGGGTCTGCCGAAAT
GTACTGGTAACATTCCGTCGCGGAAGTCTCCATCCTGCATGAAGTGCCT
CCAGTGACCTCCGGCTGTTTTCCGATTATGCATGATCGTACTAAAATCCG
TCAGCTGCCGAACCTGCTGCGTGGTTACGAACACATTCTGTCTGCCACCC
ATAACGTTATCAACGCGGAAAGCGCGCGCGGCTCCCTATAAAATCCGT
ACCTCTGGTCTCTGCCCGAACGTGACTAACGGTAACGGCTCTTTTGCAC
CATGGCCTGGGCGGTCCCGAAAAACGACAACAACAAGACCGCGACCAATT
CCCTGACCATCGAAGTCCCGTATATCTGCACCGAAGGTGAAGATCAAATC
ACGATTTGGGGCTTCCACTCCGACAGCGAGACACAATGGCGAAACTGTA
CGGTGACAGCAAACCGCAAAAATCTACTAGCTCCGCTAACGGTGTTACCA
CCCACTACGTTTCCAGATCGGTGGTTTCCCAAACAGACCGAAGATGGT
GGTCTGCCGAGTCCGGTCGCATCGTTGTAGATTATATGGTGCAGAAAAAG
CGGTAAAACCGGTACCATCACCTACCAGCGTGGCATCCTGTGCCGCGAGA
AAGTTTGGTGCCTTCCGGTCGTAGCAAAGTAATCAAAGTTGATAG

SEQ ID No:193 FOR PRIMER
CGGATAACAATCCCCCTCTAG

SEQ ID No:194 REV PRIMER
CGAAGTGAGATTTGGTTGGAGTAGTGGTCGCTAACAGAGACAGCACGTTT

SEQ ID No:195 FOR PRIMER
CCCGCAGAACGTGCTGTCTCTGTTAGCGACCACTACCCCAACCAATCTC

SEQ ID No:196 REV PRIMER
CTATTTACCCCAATTTGGAC

SEQ ID No:197 REV PRIMER
CACAGCTTGCCACGGGTTTCAGTGCCTTTCGCTAACAGAGACAGCACGTT

C

SEQ ID No:198 FOR PRIMER
CCCGCAGAACGTGCTGTCTCTGTTAGCGAAAGGCACTGAAACCCGTGGC

SEQ ID No:199 REV PRIMER
GACTAGACGCTCAGCTATCAACCTTTGATTACTTTGCTACGACC

SEQ ID No:200 REV PRIMER
CACAGCTTACCGGGGTACGAGTGCCTTTCGCTAACAGAGACAGCACGTT

C

-continued

SEQ ID No:201 FOR PRIMER
CGGGATCCAAAGGCACTCGTACCCGCGGTAAG

SEQ ID No:202 REV PRIMER
GACTAGACGCTCAGCTATCAGCCCTTAATGACCTTGAACGGCC

SEQ ID No:203 REV PRIMER
GCATAGTTTTCTCTGTGCTGTGTTCTTTCGCTAACAGAGACAGCACGT

TC

SEQ ID No:204 FOR PRIMER
AAAGGAACACAGACCAGAGG

SEQ ID No:205 REV PRIMER
CTACTACCTTTTATTACCTTGCTCC

SEQ ID No:206 REV PRIMER
CACAGCTTGCCACGGGTTTGTAGTGCCTTTCGCTAACAGAGACAGCACGTT

C

SEQ ID No:207 FOR PRIMER
AAAGGCATAAAACCCGTGGC

SEQ ID No:208 REV PRIMER
GACTAGACGCTCAGCTATCAACCTTTGATTACTTTGCTACGACC

SEQ ID No:209 STF2.HA1-1 (MAL)
MAQVINTNSLSLLTQNNLNKQSALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQVRRELAVQSA
NSTNSQSDLDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV
SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAANK
GDYEVNVAATDGTVTTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
ALTAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
TGAIKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
DFKAQPELAEEAAKTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
VLSLLATTPTPKSHFANLKGTETRGKLPKCLNCTDLVALGRPKCTGNI
PSARVSILHEVRPVTSGCFPIMHDRTKIRQLPNLLRGYEHIRLSTHNVIN
AENAPGGSYKIGTSGSCPNTVNGNFFATMAWAVPKNDNNKTATNSLTIE
VPYICTEGEDQITVWGFHSDNEAQMAYLGDSPKQKFTSSANGVTTHYVS
QIGGFNPQTEDGGLPQSGRIVVDYVMVQKSGKTGITYQRGILLPQKVWCA
SGRSKVIKGSLLPLIGADCLHEKYGGLNKSHPYYTGEHAKAIGNCPIWVK

SEQ ID No:210 STF2.HA1-2 (MAL)
MAQVINTNSLSLLTQNNLNKQSALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNNLQVRRELAVQSA
NSTNSQSDLDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV
SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAANK
GDYEVNVAATDGTVTTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
ALIAAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYYAADYDEA
TGAIKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH

-continued

DFKAQPELAEEAAKTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLAKGTETRGKLCPKCLNCTDLVALGRPKCTGNIPSARVSVILHEVR
 PVTSGCFPTMHDRTKIRQLPNLLRGYEHIRLSTHNVINAEANAPGGSYKIG
 TSGSCPNTVNGNGFFATMAWAVPKNDNNKTATNSLTIEVPYICTEGEDQI
 TVWGFHSDNEAQMAYLYGDSKPQKFTSSANGVTTHYVSQIGGFPNQTEDG
 GLPQSGRIVVDYVMVQKSGTGTITYQRGILLPQKQVWCASGRSKVIKQ

SEQ ID NO:211 STF2.HA1-2 (SH)

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKADDAAGQAI
 ANRFTANIKGLTQASRNANDGTSIAQTTEGALNEINNQLRVRELAVQSA
 NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
 NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNNGTTLDV
 SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKN
 GDYEVNVATDGTVTILAAGATKTTMPAGATTKTEVQELKDTPAVVSADAKN
 ALIAGGVDATDANGAELVKMSYTDKNGKTIIEGGYALKAGDKYYAADYDEA
 TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
 DFKAQPELAEEAAKTENPLQKIDAALAQVDALRSDLGAVQNRFNSAITN
 LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
 VLSLLAKGTRTRGKLCPCDCLNCTDLVALGRPMC VGTTTPSAKASILHEVR
 PVTSGCFPIIMHDRTKIRQLPNLLRGYENIRLSTQNVIDAELKAGGYPYRLG
 TSGSCPNTATSKSGFFATMAWAVPKNDNNKATNPLTVEVPYICTEGEDQIT
 VWGFHSDDKTQMKNLYGDSNPQKFTSSANGVTTHYVSQIGGFPDQTEDGG
 LPQSGRIVVDYVMVQKPGKTGTIVYQRGVLLPQKQVWCASGRSKVIKQ

SEQ ID NO:212 STF2

ATGGCACAAGTAATCAACACTAACAGTCTGTGCGTGTGACCCAGAATAA
 CCTGAACAAATCCCGACTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTACAGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACGGCGGTTCAGTCTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGTGAAATCAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGCCAGACTCAGTTCAACG
 GCGTGAAAGTCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGCAGCTATTAAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAAACCGGTGGTGGGTTAAATTTGACGCAGATAATAACA
 AGTACTTTGTTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT

-continued

GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
 TGGCGCAACTAAAACCACAATGCCTGCTGGTGGCACAATAAACAGAAG
 TACAGGAGTTAAAAGATACACCGGCAGTTGTTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTCTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCCGCAGATTACGATGAAGCG
 ACAGGAGCAATTAAAGCTAAAACCTACAAGTTTACTGCTGCTGACGGCAC
 TACCAAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCCGCTGGTCAT
 GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCCGCTAAAACACCGA
 AAACCCGCTGCAGAAAAATTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
 CTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGCTCTCGCGCGCAGATTCTGC
 AGCAGGCCGCTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTACGT

SEQ ID NO:213 B/Yamagata/16/18 HA

MKAIIVLLMVVTSNADRICTGITSSNSPHVVKATQGEVNVTVIPLTTT
 PTKSHFANLKGTKTRGKLCPCNCLNCTDLVALGRPMCMTIPSASAKSILH
 EVRPVTS GCFPIIMHDRTKIRQLPNLLRGYENIRLSTHNVINAEAPGGPY
 RLGTSGSCPNTSRNGFFATMAWAVPRDNKTATNPLTVEVPYICTKGEDQ
 ITVWGFHSDDKTQMKNLYGDSNPQKFTSSANGVTTHYVSQIGGFPNQTED
 GGLPQSGRIVVDYVMVQKPGKTGTIVYQRGVLLPQKQVWCASGRSKVIKGL
 PLIGEADCLHEKYGGLNKS KPYYTGEHAKAIGNCPIWVKTPCLKANGTKY
 RPPAKLLKERGFFGAIAGFLEGGWEGMIAWHGYTSHGAHVAVAADLKS
 TQEAINKITKNLSLSELEVKNLQRLSGAMDELHNEILELDEKVDLDRAD
 TISSQIELAVLLSNEGIINSEDEHLLALERKCLKMLGPSAVDIGNGCFET
 KHKCNQTCLDRIAAGTFNAGEFSLPTFDSLNTAASLNDGLDNHTILLY
 YSTAASSLAVTLMIAFIVYMVSRDNVSCSICL

SEQ ID NO:787 B/Victoria/2/87 HA

MKAIIVLLMVVTSNADRICTGITSSNSPHVVKATQGEVNVTVIPLTTT
 PTKSHFANLKGTKTRGKLCPCNCLNCTDLVALGRPKCTGTIPSASAKSILH
 EVKPVTS GCFPIIMHDRTKIRQLPNLLRGYEHIRLSTHNVINAEAPGGPY
 KVGTS GSCPNTVNGNGFFATMAWAVPKNDNNKTATNPLTVEVPYICTEGE
 DQITVWGFHSDNEAQMVKLYGDSKPQKFTSSANGVTTHYVSQIGGFPNQA
 EDGGLPQSGRIVVDYVMVQKSGTGTITYQRGILLPQKQVWCASGRSKVIKQ
 SLPLIGEADCLHEKYGGLNKS KPYYTGEHAKAIGNCPIWVKTPCLKANGT
 KYRPPAKLLKEGFFGAIAGFLEGGWEGMIAWHGYTSHGAHVAVAADL
 KSTQEAINKITKNLSLSELEVKNLQRLSGAMDELHNKILELDEKVDLDR
 ADTISSQIELAVLLSNEGIINSEDEHLLALERKCLKMLGPSAVEIGNGCF

-continued

ETKHKCNQTCLDRIAAGTFNAGEFSLPTFDSLNTAASLNDDGLDNHTIL

LYYSTAASSLAVTLMIAIFIVYMVSRDNVSCSICL

Example 13 Flagellin Proteins with Engineered Cysteine Residues for Chemical Conjugation

[0561] The crystal structure for the bacterial flagellin *S. typhimurium* flagellin type 2 (STF2, SEQ ID NO: 312) has been reported (Yonekura, K., et al., *Nature* 424:643-650 (2003)). Complete atomic model of the bacterial flagellar filament by electron cryomicroscopy has been reported, and the most detailed structure-function study of the TLR5 activation was based on this structure (Smith, K. D., et al., *Nature Immunology* 4:1247-1253 (2003)). Toll-like Receptor 5 (TLR5) recognizes a conserved site on flagellin required for protofilament formation and bacterial motility (*Nature Immunology* 4(12):1247-1253). Mutational analysis demonstrated that the TLR5 activity of flagellin resides in two regions in the N- and C-terminal domains, stretches of 39 and 31 amino acids, respectively (Smith, et al.). These regions are highlighted in grey in FIG. 18. Alanine-scanning mutagenesis of these regions identified a number of mutations which reduced TLR5 activation, but no single mutation completely abrogated activity, suggesting that there is a degree of flexibility or redundancy in the TLR5 binding site. This makes sense from an evolutionary perspective, otherwise the bacteria could easily evolve to evade detection by TLR5. Deletion of the hypervariable hinge region and retention of the conserved N- and C-terminal domains results in a protein (STF2Δ) that retains full TLR5 activity, demonstrating that the N- and C-terminal domains (together) are necessary and sufficient for TLR5 activation.

[0562] Lysine residues are convenient substrates for chemical conjugation of diverse chemical structures to a protein carrier such as flagellin. Of the 30 lysine residues in STF2 (SEQ ID NO: 312), only one lies within the experimentally-defined TLR5 activation site, and this particular lysine is not conserved among all bacterial flagellins (Smith, et al., *Nature Immunology* 4:1247-1253 (2003)). Of the remaining 29 lysines, 24 are located in the hypervariable hinge region of flagellin, thus leaving only 5 lysine residues in STF2Δ (SEQ ID NO: 313). The structure in FIGS. 19 and 20 shows that most lysines are spatially distal to the TLR5 activation domain, thus it would seem that lysines could be randomly conjugated with peptide or carbohydrate antigens with a reasonable probability of not interfering with the TLR5 activity of the resulting conjugated STF2Δ (SEQ ID NO: 313) protein. Five lysines appear to border the TLR5 activation site quite closely; these residues would be the first to consider mutating if antigen conjugation to lysine affects TLR5 activity.

[0563] Possible additional sites for conjugation (such as cysteines or additional lysines) may be engineered into full-length flagellin (STF2 (SEQ ID NO: 312)) or STF2Δ (SEQ ID NO: 313) by placing such residues in the hypervariable region or in the tail of the conserved N- and C-terminal domains distal to the TLR5 binding site. Chemical methods for conjugation to other amino acids also exist. These include carboxy amino acids (glutamic acid, aspartic acid) and the carboxyl terminal amino acid, arginine, histi-

dine, tryptophan, tyrosine, and serine. Any of these strategies may be utilized to conjugate antigenic structures to flagellin without interfering with the TLR5 activation site.

[0564] Some types of antigens including polysaccharides are not amenable to recombinant fusion DNA technology nor to synthetic peptide chemistry, and thus cannot be genetically or synthetically linked to a ligand for a Toll-like Receptor (TLR). In addition, it is possible that genetic or synthetic fusion of a peptide to a TLR ligand may inhibit the proper folding of the TLR ligand. Chemical conjugation of the antigen to the TLR ligand, which is folded and purified, may be useful. The chemical conjugation of peptide antigens also permits the placement of antigen at specific sites on the TLR ligand, thus limiting interference with receptor binding of the TLR ligand or maximizing exposure to antigen receptors. The conjugation of many peptide antigens to a TLR ligand may also maximize immunogenicity by increasing the conformational heterogeneity with which antigen is presented to the immune system. To provide a site for chemical conjugation of such antigens to flagellin, a cysteine residue was engineered at different sites in the gene encoding STF2Δ (flagellin with the hypervariable hinge region deleted; SEQ ID NO: 313 and SEQ ID NO: 314).

Materials and Methods

[0565] Polymerase chain reaction (PCR): Platinum® PCR SuperMix High Fidelity kit (catalog number 12532-016, Invitrogen Corporation, Carlsbad, Calif.) was used for all PCR amplifications using the following protocol based on the manufacturer's instructions.

[0566] 1. The following components were added in any order to each reaction tube:

[0567] a. 45 μl Platinum® PCR SuperMix High Fidelity (PCR reaction buffer)

[0568] b. Primer solution (10 pMol final concentration of each primer)

[0569] c. Template DNA solution (10 ng plasmid DNA)

[0570] 2. Reaction volume was made up to 50 μl with water

[0571] 3. Tubes were capped and loaded in thermal cycler

[0572] 4. Tubes were incubated at 94° C. for 30 to 120 seconds to completely denature the template and activate the enzyme

[0573] 5. The following PCR amplification was performed for 25-35 cycles:

[0574] a. Denature at 94° C. for 15-30 seconds

[0575] b. Anneal at 50-55° C. for 15-30 seconds

[0576] c. Extend at 68-72° C. for 1 minute per kb of PCR product size

[0577] d. If necessary, cool to 4° C. and hold until ready for next process

[0578] Construction of STF2Δ gene (SEQ ID NO: 314): Full length flagellin of *Salmonella typhimurium* fljB (STF2, SEQ ID NO: 312) is encoded by a 1.5 kb gene (SEQ ID NO: 315). To generate STF2Δ (SEQ ID NO: 314), the sequence

corresponding to the hypervariable hinge region (amino acids 170 to 415 of SEQ ID NO: 312) was deleted and replaced with a short flexible linker (GAPVDPASPW (SEQ ID NO: 336)) designed to facilitate interactions of the NH₂ and COOH terminal sequences of STF2 (SEQ ID NO: 315) necessary for TLR5 signaling (Smith, K. D., et al., (2004). Toll-like Receptor 5 recognizes a conserved site on flagellin required for protofilament formation and bacterial motility (Smith, et al., *Nature Immunology* 4:1247-53 (2003)).

[0579] To generate the STF2Δ plasmid, a two-step PCR was used (FIG. 21). In the first reaction, a plasmid encoding STF2.OVA (a fusion of full-length STF2 with full-length ovalbumin, SEQ ID NO: 316) was mixed with primers STF28BGF-1 (SEQ ID NO: 317) and STF28MCR-1 (SEQ ID NO: 318) in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above. In a parallel reaction, STF2.OVA template plasmid was mixed with primers STF28.MCF-2 (SEQ ID NO: 319) and STF28ECR-2 (SEQ ID NO: 320) in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above.

[0580] The PCR amplification reactions generated ~500 bp and 270 bp fragments, respectively. These PCR products were combined with primers STF28BGF-1 (SEQ ID NO: 317) and STF28ECR-2 (SEQ ID NO: 320) in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above. The amplified DNA product from this reaction (770 bp) was digested with BglII and EcoRI restriction enzymes for 2 hours at 37° C., and ligated into pMTBiP/V5-His B (Invitrogen, Carlsbad, Calif.) that had previously been digested with BglII and EcoRI and treated with calf intestinal phosphatase (CIP). An aliquot of the ligation mix was used to transform *E. coli* TOP10 cells. The resultant construct pMT/STF2Δ was used to generate the cysteine modified STF2Δ constructs.

[0581] Construction of STF2Δ genes engineered to express a single cysteine residue: To introduce cysteine residues at defined positions in the STF2 polypeptide, pMT/STF2Δ was used as a template in PCR with primer pairs as defined below.

[0582] To construct the STF2Δ.3'Cys (SEQ ID NO: 325) plasmid, pMT/STF2Δ plasmid was mixed with 3'Forward1 (SEQ ID NO: 321) and 3'Reverse1 (SEQ ID NO: 322) primers in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above.

[0583] To construct the 5'Cys.STF2Δ (SEQ ID NO: 326) plasmid, pMT/STF2Δ plasmid was mixed with 5'Forward2 (SEQ ID NO: 323) and 5'Reverse2 (SEQ ID NO: 324) primers in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above.

[0584] For STF2Δ.3'Cys (SEQ ID NO: 325) and 5'Cys.STF2Δ (SEQ ID NO: 326) constructs, the PCR products generated were digested with NdeI and BlnI restriction enzymes and purified by agarose gel electrophoresis. The purified fragments were ligated by compatible ends to pET24a (Novagen, Madison, Wis.) plasmid DNA that had been previously digested with NdeI and BlnI restriction enzymes and CIP-treated. The resulting constructs are designated pET/STF2Δ.3'Cys and pET/5'Cys.STF2Δ, respectively.

[0585] To construct the STF2Δ.HingeCys (SEQ ID NO: 331) plasmid, a two step PCR similar to that described in

FIG. 18 was employed. One fragment was generated by mixing pMT/STF2Δ with paired HingeForward1 (SEQ ID NO: 327) and HingeReverse3 (SEQ ID NO: 328) primers in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above. Another fragment was generated by mixing pMT/STF2Δ with paired HingeForward2 (SEQ ID NO: 329) and HingeReverse4 (SEQ ID NO: 330) primers in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above. Aliquots from each of these PCR reactions were combined and mixed with HingeForward1 (SEQ ID NO: 327) and HingeReverse4 (SEQ ID NO: 330) primers in PCR reaction buffer, and the mixture was amplified in a PCR reaction as described above. The final PCR product was purified by agarose gel electrophoresis and digested with NdeI and BlnI, and the purified fragment was ligated into the pET24a vector as described above. The resulting construct is designated pET/STF2Δ.HingeCys.

[0586] Protein expression: Plasmids pET/STF2Δ.3'Cys, pET/5'Cys.STF2Δ, and pET/STF2Δ.HingeCys were transformed into competent *E. coli* BLR(DE3) cells as follows. To a 1.5-ml snap-cap polypropylene tubes pre-chilled on ice was added 20 μl aliquot of BLR (DE3) cells and 1 μl plasmid DNA (1 μg/μl), and the mixture was incubated for 30 minutes. The tubes were heated for exactly 30 seconds in a 42° C. water bath without shaking, and then placed on ice for an additional 2 minutes. To each tube of heat shocked cells were added 250 μl of room temperature SOC medium.

[0587] The cells were recovered at 37° C. (shaking at 250 rpm) for 60 minutes prior to plating on selective media containing kanamycin. Various aliquots (50-100 μl) of the transformation mix were plated and the plates were incubated at 37° C. for 15 to 18 hours. Colonies were picked and inoculated into 2 ml Luria-Bertani (LB) broth supplemented with 25 μg/ml kanamycin and 12.5 μg/ml tetracycline, and cultured overnight. Fresh LB cultures were inoculated by diluting an aliquot of the overnight cultures 1:100 and cultured at 37° C. with shaking. When the OD₆₀₀ of the culture reached 0.6 to 1.0, protein expression was induced by the addition of isopropyl thio-β-D-galactoside (IPTG) to a final concentration of 1 mM. Several hours after induction, the cells were harvested for analysis of protein expression by SDS-PAGE. Glycerol stocks were prepared from cultures grown in LB supplemented with 0.5% glucose, 25 μg/ml kanamycin and 12.5 μg/ml tetracycline and were frozen at -80° C. following the addition of glycerol (7% final).

[0588] Protein purification: Proteins STF2Δ.3'Cys (SEQ ID NO: 332) and STF2Δ.HingeCys (SEQ ID NO: 333) were expressed and purified as follows. Glycerol stocks of *E. coli* BLR(DE3) cells harboring the desired plasmids were inoculated into 10 L shake-flasks containing LB and incubated at 37° C. with constant shaking. When cultures reached an optical density of A₆₀₀=0.8, protein expression was induced by the addition of 1 mM IPTG and cultures were incubated at 37° C. with constant shaking for 4 hours before harvesting. Cells were collected from 10 L of culture by low-speed centrifugation at 5,000 rpm (SLA3000 rotor) for 10 minutes and suspended in 50 mM Tris-HCl pH 8.0, 1 mM EDTA, 1 mM DTT (100 ml/10 L).

[0589] The cells were disrupted by passing the cell suspension twice through a microfluidizer at 18,000 psi. The insoluble material was separated from soluble proteins by

centrifugation at 10,000 rpm (SS34 rotor) for 15 minutes. Under the culture conditions described, all STF2Δ proteins fractionated with the insoluble material and formed stable inclusion bodies that were collected as a solid pellet following centrifugation. The inclusion bodies were washed twice with 50 mM Tris-HCl, pH 8.0, 0.1 M NaCl and 0.5% Triton X-100, followed by two washes with the same buffer without detergent. Each wash was performed using a dounce homogenizer and the cleaned inclusion bodies (IBs) were collected by centrifugation at 10,000 rpm (SS34 rotor) for 15 minutes. The purified IBs were washed a final time with 50 mM Tris-HCl, pH 8.0 and stored as a cell pellet at -80° C. until needed.

[0590] IB material was thawed and solubilized in 8 M urea at pH 4.0. This step selectively solubilized the target protein while leaving a significant amount of debris and contaminating proteins as a solid precipitate which was removed by centrifugation. Because of the single cysteine present in these proteins, all subsequent purification procedures were carried out using buffers containing 1 mM DTT as a reducing agent. Solubilized protein was captured using SP fast flow sepharose (30 ml, XK16) and selectively eluted with 8 M urea in 25 mM Na Acetate (C₂H₃O₂Na), pH 4.0, 1 mM DTT, 1 mM EDTA and 0.2 M NaCl. To eliminate protein precipitation following this step, the pH of the SP elution was adjusted from 4.0 to 8.0 by dialysis against 50 mM Tris-HCl, pH 8.0, 1 mM DTT, 1 mM EDTA before protein refolding. The dialyzed material was refolded by direct dilution (10-fold) into 50 mM Tris-HCl pH 8.0, 1 mM DTT and 1 mM EDTA such that the final protein concentration was less than 0.1 mg/ml. The refolded SP pool was loaded directly onto Q high performance sepharose (30 ml, XK16) and bound protein was eluted with a 20 column volume linear gradient from 0 to 0.5 M NaCl in 50 mM Tris-HCl, pH 8.0, 1 mM DTT, 1 mM EDTA. This chromatography step yielded a single peak that eluted at a conductivity of approximately 15 ms/cm. The eluted material was pooled and stored at -80° C.

[0591] Protein characterization: Proteins were characterized for purity, identity, endotoxin content, and biological activity using the following assays.

[0592] SDS-PAGE: Proteins (typically 5 μg) were diluted in SDS-PAGE sample buffer (1% SDS, 30 mM Tris-HCl, pH 6.8, 4% glycerol, 0.1 mg/ml bromophenol blue) with and without 5 mM β-mercaptoethanol. The samples were boiled for 5 minutes and loaded onto a 4-20% SDS polyacrylamide gel. Following electrophoresis, gels were stained with coomassie blue to visualize protein bands.

[0593] Endotoxin assay: Endotoxin levels were measured using the QCL-1000 Quantitative Chromogenic LAL test kit (BioWhittaker #50-648U), following the manufacturer's instructions for the microplate method.

[0594] Protein Assay: Protein concentrations were determined by the MicroBCA Protein Assay Reagent Kit in a 96-well format using BSA as a standard (Pierce Biotechnology), following the manufacturer's instructions.

[0595] Flagellin ELISA: Protein integrity and concentration were examined by ELISA with antibodies specific for flagellin. 96-well ELISA plates were coated overnight at 4° C. with serial dilutions of each target protein, in PBS starting at 5 μg/ml. Plates were blocked with 200 μl/well of Assay

Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature then washed three times in phosphate-buffered saline containing Tween-20 (PBS-T, 12 mM NaPO₄, 137 mM NaCl, 2.7 mM KCl, 0.05% Tween 20). Rabbit polyclonal anti-flagellin antibody diluted in ADB (100 μl/well, 1:5000) was added to all wells and the plates were incubated for 1 hour at room temperature or overnight at 4° C., then washed three times with PBS-T. HRP-labeled goat anti-rabbit IgG antibody (Jackson Immunochemical) diluted in ADB was added (100 μl/well, 1:5000) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce) and monitoring color development, A₄₅₀ was measured on a Tecan Farcyte microplate spectrophotometer.

[0596] TLR5 bioactivity assay: HEK293 cells (ATCC, Cat#CRL-1573, Manassas, Va.) constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96 well microplates (50,000 cells/well), and recombinant test proteins were added. The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20° C. After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce; Rockford, Ill., #M801E and #M802B) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer

Results and Discussion

[0597] Protein yield and purity: The final yield and endotoxin levels of each protein are shown below.

Protein	SEQ ID NO:	Yield (mg)	Endotoxin (EU/μg)
STF2Δ.3'Cys	332	75.0	0.01
STF2Δ.HingeCys	333	117.0	0.004

[0598] Flagellin integrity: Engineering a single cysteine into STF2Δ does not diminish its recognition by flagellin-specific antibodies, as shown in FIG. 22.

[0599] TLR5 agonist activity in the HEK293 IL-8 assay: Engineering a single cysteine into STF2Δ does not diminish the TLR agonist activity of the protein, as shown in FIG. 23. Cells exposed to STF2Δ.3'Cys (SEQ ID NO: 332) or STF2Δ.HingeCys (SEQ ID NO: 333) secreted IL-8 at levels comparable to those induced by exposure to full-length flagellin (STF2, SEQ ID NO: 312) or STF2Δ without cysteine residues (SEQ ID NO: 313).

[0600] Fusion of full-length flagellin (STF2, SEQ ID NO: 312) or hinge region-deleted flagellin (STF2Δ, SEQ ID NO: 313) to a protein antigen such as West Nile virus envelope protein or influenza A hemagglutination significantly increases the immunogenicity of the fused antigen. This approach is useful for protein or peptide antigens which can be encoded in a genetic fusion construct expressing both flagellin and the antigen of interest in a single protein, but it cannot be applied to non-protein or non-peptide antigens, such as polysaccharide antigens. Since many bacteria and tumor cells express specific polysaccharide structures that

are antigenic but poorly immunogenic, it is important to devise scalable methods for increasing the immunogenicity and protective efficacy of such structures. Chemical conjugation to a TLR ligand, such as flagellin, is one potential strategy.

[0601] There are several methods of chemical linkage, but one of the simplest is to couple an antigen to a free thiol group such as that on a single, unmatched cysteine residue. Since flagellin does not natively contain any cysteines, cysteine residues were engineered into flagellin at one of three positions: the amino terminus of flagellin (5'Cys.STF2Δ, SEQ ID NO: 326 and SEQ ID NO: 334), the carboxy terminus of flagellin (STF2Δ.3'Cys, SEQ ID NO: 325 and SEQ ID NO: 332), and in the deleted hinge region of flagellin (STF2Δ.HingeCys, SEQ ID NO: 331 and SEQ ID NO: 333). All constructs were produced in the expression vector pET24a for *E. coli* expression. When purified from *E. coli* cells, STF2Δ.HingeCys (SEQ ID NO: 333) and STF2Δ.3'Cys (SEQ ID NO: 332) proteins retain all antigenic and TLR5 biological activity properties of the unmodified STF2Δ protein, thus confirming that the introduction of a single cysteine residue does not negatively impact the expression, purification, refolding, or biological activity of STF2Δ.

Conjugation of Influenza hemagglutinin Maturational Cleavage site peptide to STF2Δ.hingeCYS

[0602] A peptide representing the maturational cleavage site of influenza hemagglutinin was chemically conjugated a modified flagellin protein, STF2Δ.HingeCys (SEQ ID NO: 333).

Materials and Methods

[0603] Production of STF2Δ.HingeCys protein: STF2Δ.HingeCys protein (SEQ ID NO: 333) was expressed and purified as described in the previous Example.

[0604] Synthesis of H1C1 peptides The sequence NH₂-NIPSIQSRGLFFAIAAGFIE-COOH (SEQ ID NO: 337) represents the maturational cleavage site of influenza A/H1N1 hemagglutinin (Bianchi, E., et al. (2005). Universal influenza B vaccine based on the maturational cleavage site of the hemagglutinin precursor (Bianchi, et al., *J. Virol.* 79:7380-7388 (2005)). Two peptides were designed with an extra cysteine or an extra lysine on the N-terminus to facilitate chemical linkage to carrier proteins, resulting in the following peptide sequences:

CysH1C1	(SEQ ID NO: 338)
NH ₂ -CNIPSIQSRGLFFAIAAGFIE-COOH	(SEQ ID NO: 338)
LysH1C1	(SEQ ID NO: 339)
NH ₂ -KNIPSIQSRGLFFAIAAGFIE-COOH	(SEQ ID NO: 339)

[0605] The peptides were synthesized by Anaspec, Inc., (San Jose, Calif.) utilizing standard Fmoc chemistry, after which they were cleaved from the matrix with trifluoroacetic acid (TFA), purified by reversed-phase HPLC and lyophilized.

[0606] Conjugation of CysH1C1 peptide to STF2Δ.HingeCys: STF2Δ.HingeCys protein (SEQ ID NO:

333) at a concentration of 4.9 mg/mL was dialyzed overnight into Buffer A [1× phosphate-buffered saline (PBS), 5 mM EDTA, pH 7.2]. BM(PEO)₂ (Pierce Biotechnology, Rockford, Ill.), a homobifunctional maleimide crosslinker, was dissolved in DMSO (dimethyl sulfoxide) and added to a final concentration of 2.3 mM. After incubating for 1 hour at room temperature, the free crosslinker was removed from the protein using a 5 ml Superdex 25 HiTrap desalting column. CysH1C1 peptide (SEQ ID NO: 338) was dissolved in DMSO and added to the maleimide-derivatized STF2Δ.HingeCys protein at a final concentration of 0.32 mM. After incubating 3 hours at room temperature, the reaction was stopped by adding DTT (dithiothreitol) to a final concentration of 40 mM. The protein and protein-peptide conjugate were separated from free peptide by fractionation on a Superdex 200 10/300 size-exclusion chromatography (SEC) column (GE/Amersham; Piscataway, N.J.) equilibrated in 1× Tris-buffered saline (TBS), pH 8.0.

[0607] Protein characterization: Proteins were characterized for purity, identity, endotoxin content, and biological activity as described in the previous Example.

Results and Discussion

[0608] Characterization of the STF2Δ.HingeCys:CysH1C1 peptide conjugate: Conjugation of the CysH1C1 peptide (SEQ ID NO: 338) to STF2Δ.HingeCys (SEQ ID NO: 333) was assayed by SDS-PAGE with coomassie staining. Peptide conjugation to the protein resulted in a doublet band in which the faster-migrating band corresponded to the un-conjugated protein and the slower migrating band corresponded to the protein-peptide conjugate.

[0609] A cloudy precipitate was seen in the conjugation mixture due to the low solubility of the peptide. To ascertain if the protein-peptide conjugate was precipitating, the mixture was centrifuged (16,000×g for 15 minutes) and the resulting pellet and supernatant fractions analyzed by SDS-PAGE. The majority of the peptide-conjugated protein remains in the supernatant indicating that it is still soluble.

[0610] Size-exclusion chromatography (SEC) of the STF2Δ.HingeCys:CysH1C1 conjugate was performed to separate the conjugated protein from remaining crosslinker and unconjugated peptide. A single major protein peak eluted in the included range of the column at the expected volume for monomeric STF2Δ, while there is very little material eluting in the void (FIG. 24), indicating that most of the fractionating protein is monomeric.

[0611] The monomeric nature of the STF2Δ.HingeCys:CysH1C1 conjugate was confirmed by SDS-PAGE analysis of the S200 fractions (FIG. 25). The protein-peptide conjugate co-elutes with the unconjugated protein in the same ratio as in the load sample, indicating that the conjugate species is a monomer.

[0612] Bioactivity of the STF2Δ.HingeCys:CysH1C1 conjugate mixture was found to be equivalent to the unconjugated STF2Δ.HingeCys protein, indicating that the attachment of the BM(PEO)₂ crosslinker and the peptide conjugation reaction does not inhibit TLR5 stimulatory activity (FIG. 8).

Conjugation of a Pam₃CYS-Containing Lipopeptide to an Influenza Hemagglutinin Antigen

[0613] Several types of Toll-like Receptors (TLRs) are potentially activated by lipids or lipid-conjugates. These include TLR2/1 (diacyl lipoproteins and GPI-linked proteins), TLR2/6 (triacyl lipoproteins and GPI-linked proteins) and TLR4 (lipopolysaccharides). The utility of vaccines composed of antigens linked to such lipids is obvious: the lipids are strong activators of innate immune pathways and yet are themselves poorly immunogenic. However, the production of such conjugates is very complex. Biosynthesis of recombinant lipoproteins in bacteria is limited by low productivity and extreme difficulty in purifying the resulting protein-lipopeptide fusion. Chemical synthesis of lipopeptide antigens is more straight forward, but is limited to peptide antigens. Here we demonstrate the chemical conjugation of a Pam₃Cys lipopeptide, a TLR2/6 agonist, to influenza HA1-1His₆ in a process which bypasses the difficulties inherent in both biosynthesis and complete chemical synthesis of lipidated antigens.

Materials and Methods

[0614] Production of HA1-1His₆(PR8)Bv (SEQ ID NO: 179) HA1-1His₆ (PR8)Bv was prepared as described herein.

[0615] Synthesis of Pam₃CS(K)₄GC (SEQ ID NO: 335) Custom lipopeptide synthesis was performed by Anaspec, Inc. (San Jose, Calif.) using standard solid-phase Fmoc chemistry, after which the lipopeptide was cleaved from the matrix with trifluoroacetic acid (TFA), purified by reversed-phase HPLC and lyophilized.

[0616] Conjugation of Pam₃CS(K)₄GC (SEQ ID NO:335) to HA1-1His₆(PR8)Bv (SEQ ID NO: 179) HA1-1His₆(PR8)Bv (SEQ ID NO: 179) was dialyzed into Buffer A (1×PBS+5 mM EDTA, pH 7.2). Sulfo-SMCC (Pierce; Rockland, Ill.) a heterobifunctional maleimide/NHS-ester crosslinker was dissolved in DMSO and added to a final concentration of 0.432 mM. After incubating for 30 min. at room temperature the protein was desalted into Buffer A using a G-25 HiTrap desalting column (GE/Amersham; Piscataway, N.J.). Triton X-114 (TX-114) (Sigma; St Louis, Mo.) was added to a final concentration of 1% (w/v). Pam₃CS(K)₄GC was dissolved to 20 mg/ml in DMSO and added to a final concentration of 0.6 mg/ml. After incubating at room temperature for 3 hours, the reaction tube was placed in a 37° C. bath for 10 minutes to cause TX-114 droplet formation. The sample was then centrifuged at 16,000×g for 10 minutes to separate the detergent and aqueous phases. After drawing off the aqueous phase the detergent phase was resuspended to the original reaction volume with Buffer A. The total, detergent, and aqueous samples were then analyzed by SDS-PAGE.

[0617] Protein characterization: Proteins-lipopeptide conjugates were characterized by SDS-PAGE. Samples (typically 5 µg) were diluted in SDS-PAGE sample buffer (0.1M Tris, pH 8.0/4% SDS/25% glycerol/0.1M DTT). The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel. Following electrophoresis, gels were stained with coomassie blue to visualize protein bands.

Results and Discussion

[0618] Input HA1-1His₆(PR8)Bv (SEQ ID NO: 179) runs as a doublet band on SDS-PAGE. This has been seen for all

of the HA1-1His₆ proteins made in Baculovirus to date and is likely due to differences in glycosylation. HA1-1His₆(PR8)Bv (SEQ ID NO: 179) which has been derivatized with Sulfo-SMCC but not conjugated to lipopeptide stays in the aqueous phase during TX-114 separation, the expected behavior for a soluble, globular protein. HA1-1His₆(PR8)Bv (SEQ ID NO: 179) conjugated to Pam₃Cys shows 3 or 4 bands of higher molecular weight than the derivatized input protein, consistent with covalent modification by the lipopeptide. On TX-114 phase separation the higher mw species segregate primarily into the detergent phase while the lower species (corresponding to the unmodified protein) stay in the aqueous phase. This behavior is consistent with the commonly observed segregation of lipoproteins into detergent droplets and confirms that HA1-1His₆(PR8)Bv (SEQ ID NO: 179) has been covalently modified with the Pam₃CS(K)₄GC (SEQ ID NO: 335) lipopeptide. The higher molecular-weight bands of approximately 60, 120, and 180 kDa, which are seen in the SMCC-derivatized protein but not in the input, are covalent HA1-1 multimers formed by the crosslinker. This multimerization would be prevented, and the yield of singly lipidated HA1-1 improved, by engineering a single cysteine into HA1-1 and using a cysteine-specific rather than a lysine-specific crosslinker. Such a strategy would also give better control over the site of attachment of the lipopeptide on HA1-1, giving a more homogeneous product and possibly improving immunogenicity

SEQ ID NO: 312, amino acid sequence of STF2
MAQVINTNSLSLLTQNNLNKSQLGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQRVRELAVQSA
NSTNSQSDLDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDLSLVQKAYDVKDTAVTTKAYANNNGTLDV
SGLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIIGGFTGADAANK
GDYEVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKN
ALIAAGVDATDANGAELVKMSYTDKNGKTIIEGYALKAGDKYYAADYDEA
TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
DFKAQPELAEEAAKTTENPLQKIDAALAQVDALRSLDGAQVQNRFNSAITN
LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQN
VLSLLR

SEQ ID NO: 313, amino acid sequence of STF2A
MAQVINTNSLSLLTQNNLNKSQLGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQRVRELAVQSA
NSTNSQSDLDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDLSLVHGAQVDPASPTENPLQKIDAALAQ
VDALRSLDGAQVQNRFNSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSR
AQILQQAGTSVLAQANQVPQNVLSSLLR

SEQ ID NO: 314, nucleotide sequence of STF2A
ATGGCACAAAGTAATCAACACTAACAGTCTGTGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTCGGTATCAACAGCGCGAAAGACGATCGCGCAGGTACGGCGATT
GCTAACCGTTTACC CGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACCGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
AATCAACAACAACCTGCGAGCGTGTGCGTGAACCTGGCGGTTCACTGTGCT
AACAGCACCACCTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCCTGAACGAAATCGACCGTGTATCCGCGCCAGACTCAGTTCAACG
GCGTGAAAGTCTTGGCGCAGGACACACCTGACCATCCAGGTTGGCGCC
AACGACCGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGCTGGACTCACTGAACGTGCATGGAGCGCGGTTGGATCCTGCTA
GCCATGGAACGAAACCGCTGCGAGAAATTTGATCCCGCGCTGGCGCAG
GTGGATGCGCTGCGCTCTGATCTGGGTGCGGTACAAACCGTTTCAACTC
TGCTATACCAACCTGGGCAATACCGTAACCAATCTGTCTGAAGCGCGTA
GCCGTATCGAAGATTCCGACTACGCGACCGAAGTTTCCAACTGTCTCGC
GCGCAGATTTTGACGACGCGCGGTACTTCCGTTCTGGCGCAGGCTAACCA
GGTCCCGCAGAACGCTGCTCTCTGTTACGT

-continued

SEQ ID NO: 315, nucleotide sequence of STF2
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTACCGCGCAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCTGAACTGGCGTTTCACTCTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAATACAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAGTCCCTGGCGCAGGACAACACCTGACCATCCAGTTGGCGCC
 AACGACGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCAGTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCAATAATGGTACTACACTGGATGA
 TCGGGTCTTGATGATGACAGTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTAAATTTGACGCGAGATAATAACA
 AGTACTTTTGTACTTATTTGGTGGCTTTTACTGGTGTGATGCGCAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAACACCAATGCCTGCTGGTGGCAGCACTAAACAGAGAG
 TACAGGAGTTAAAGATACACCGGACGTTGTTTTCAGCAGATGCTAAAGT
 GCCTTAATTGCTGGCGGCTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGTTCAGGATACAGCTTACATGCGCAGGATAACGATGAAGC
 ACAGGAGCAATTAAAGCTAAACATAAGTTTACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCACTGGGTGGCGTAGACGGTAAACCGAAG
 TCGTTACTATTCGACGGTAAACCTTACATGCGCAGCAAGCGCTGATCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAACACCGCA
 AAACCGCGCTGCAGAAAATGATGCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGGGTACAAAACCGTTTCAACTCTGCTATCAACAA
 CTGGCAATACCGTAAACAATCTGCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGC
 AGCAGCGCGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
 GTGCTGTCTCTGTACTCT

SEQ ID NO: 316, nucleotide sequence of STF2.OVA
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTACCGCGCAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCTGAACTGGCGTTTCACTCTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAATACAC
 CCAGCGCCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAGTCCCTGGCGCAGGACAACACCTGACCATCCAGTTGGCGCC
 AACGACGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCAGTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGACAGTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTAAATTTGACGCGAGATAATAACA
 GTACTTTTGTACTTATTTGGTGGCTTTTACTGGTGTGATGCGCCAAAATG
 GCGATTATGAAGTTAAGCTTGTCTACTGACGGTACAGTAACCTTTCGCGCT
 GCGCGCACTAAACACCAATGCTGCTGGTGGCAGCACTAAACAGAGAT
 ACAGGAGTTAAAGATACACCGGACGAGTGTTCAGCAGATGCTAAAGATG
 CCTTAATTGCTGGCGGCTTGACGCTACCGATGCTAATGGCGCTGAGTTG
 GTCAAAATGCTTTATACCGATAAAATGGTAAGACAATTGAAGGCGGTTAT
 GCGCTTAAAGCTGGCGATAAGTATTACGCGCAGATTACGATGAAGCGAC
 AGGAGCAATTAAAGCTAAACACCAAGTTATAGCTGCTGCTGACGGCACTA
 CCAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAACCGAAGTC
 GTTACTATCGACGGTAAACCTTACAATGCCAGCAAGCGCGCTGGTCAATGA
 TTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAACACCGCAAA
 ACCCGCTGCAGAAAATGATGCCGCGCTGGCGCAGGTGGATGCGCTGCGC
 TCTGATCTGGGTGGGTGCTACAAAACCGTTTCAACTCTGCTATCAACCACT
 GGGCAATACCGTAAACAATCTGCTGAAGCGCTAGCCGTATCCGAAGATT
 CCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTTTGGCAG
 CAGGCGGGTACTTCCGTTCTGGCGCAGGCTAACAGGTTCCGCGACGCT
 GCTGTCTCTGTTAGCTCTCAGGCGCTCCATCGCGCAGCAAGCATGGAAAT
 TTTGTTTTGATGATTTCAAGGAGCTCAAAGTCCACCATGCCAATGAGAAAC
 ATCTCTACTGCGCCCATTTCCGTTCTGGCGCAGGCTAACAGGTTCCGCGACG
 GGTGCAAAAAGACAGCACCAGGACACAATAAATAAGGTTGTTGCTTTG
 ATAAACTTCCAGGATTCGGAGACAGTATTGAAGCTCAGTGTGGCACATCT
 GTAACCGTTCACTCTTCACTTTCAGGACATCTCAACCAATCAACCAAC
 AAATGATGTTTATTCGTTAGCCTTTCAGGATAGACTTTATGCTGAAGAGA
 GATACCAATCTTCCGAGAATACTTGCAGTGTGTGAAGAACTGTATAGA
 GGAGGCTTGAAGCACTTCACTTTCAGGACGCTGAGATCAAGCCAGAGA
 GCTCATCAATTCCTGGGTAGAAAGTCAGACAAATGAATATCAGAAATGTC
 CTTACGCAAGCTCCGTTGGATTTCAAACTGCAATGGTTCTGGTTAATGTC
 CATTGCTCTCAAGAGCTGTGGGAGAAAGCATTTAAGGATGAAGACACAC
 AAGCAATGCCCTTTCAGAGTGAAGTGAAGAAAGCAAACTGTGCGAGATG

-continued

ATGTACCAGATTGGTTTATTTAGAGTGGCATCAATGGCTCTGAGAAAAT
 GAAGATCCTGGAGCTTCCATTTGCCAGTGGGACAATGAGCATGTTGGTGC
 TGTTGCCCTGATGAAGTCTCAGGCCCTTGAGCAGCTTGAGAGTATAATCAAC
 TTTGAAAACCTGACTGAATGGACAGTTCTAATGTTATGGAAGAGAGAA
 GATCAAAGTGTACTTACCTCGCATGAAGATGGAGGAAAAATACAACCTCA
 CATCTGTCTTAATGGCTATGGGCATTACTGACGTGTTAGCTCTTCAGCC
 AATCTGTCTGGCATCTCCTCAGCAGAGAGCCTGAAGATATCTCAAGCTGT
 CCATGCAGCACATGCAGAAATCAATGAAGCAGGAGAGAGGTTGGTAGGGT
 CAGCAGAGGCTGGAGTGGATGCTGCAAGCGTCTCTGAAGAAATTTAGGGCT
 GACCATCCATTCTCTTCTGTATCAAGCACATCGCAACCAACGCGCTTCT
 CTCTTTTGGCAGATGTGTTTCCCTTTCGAAGCTTGAAGGTAAGCCTATCC
 CTAACCCCTCTCTCGGTCTCGATTCTACGCGTACCGGTCATCATCACCAT
 CACCATTGA

SEQ ID NO: 317, nucleotide sequence of STF28BGF-1
 CTCGGGAGATCTGCACAAGTAATCAACACTAACAGTCT

SEQ ID NO: 318, nucleotide sequence of STF28MCR-1
 CCATGGGCTAGCAGGATCCACCGCGCTCCCTGCAGCTTCA

SEQ ID NO: 319, nucleotide sequence of STF28MCF-2
 GGAGCGCGGTGGATCTCTGCTAGCCCATGGACCGAAAACCCG

SEQ ID NO: 320, nucleotide sequence of STF28ECR-2
 TCTGCAGAAATTCAGTAACAGAGACAGCAGCTTCTGCGGAGCGTCCCGCA
 GAACGTGCTGTCTCTGTTACGTGAATTCGCAGA

SEQ ID NO: 312, nucleotide sequence of 3'Forward1
 ACTGAGTGCATATGGCACAAGTAATCAACACTAACAG

SEQ ID NO: 322, nucleotide sequence of 3'Reverse1
 GACTGACTGCTCAGCCTATTAGCAGAGCGGCCACTGTGCTGGATATC
 TAGAG

SEQ ID NO: 323 nucleotide sequence of 5'Forward2
 AGTCAGGCGATATGTGCGCACAGTAATCAACACTAACAGTCTG

SEQ ID NO: 324, nucleotide sequence of 5'Reverse2
 GACTGACTGCTCAGCCTATTAAAGTAAACAGAGACAGCAGCTTCTGCGGGA
 CCTGGTTAG

SEQ ID NO: 325, nucleotide sequence of STF2A.3'Cys
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGGCGATT
 GCTAACCGTTTACCAGCAACATCAAAGGTTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCGAGGTGTGCGTGAACCTGGCGGTTCACTGCTGT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAG
 CCAGCGCTGAACGAAATCGACCGTGTATCCGCGCAGACTCAGTTCAACG
 GCGTGAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCGCG
 AACGACGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGACTACTGAACGTGATGGAGCGCGGTGGATCTTGGCTA
 GCCCATGGACCGAAAACCGCTGCAGAAAATGATGCCGCGCTGGCGCAG
 GTGGATGCGCTGCGCTCTGATCTGGGTGGGTACAAAACCGTTTCAACTC
 TGCTATCACCAACCTGGGCAATACCGTAAACAATCTGTCTGAAGCGGTA
 GCCGTATCGAAGATTCCGACTACGCGACGGAAGTTTCCAACATGTCTCGC
 GCGCAGATTTTGCAGCAGGCGGTTACTTCCGTTCTGGCGCAGGCTAACCA
 GGTCCCGCAGACGTGCTGTCTGTACGTGAATCTCTAGATATCCAG
 CACAGTGGCGCGGCTCTGCT

SEQ ID NO: 326, nucleotide sequence of 5'Cys.STF2A
 ATGTGCGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAA
 TAACCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGT
 CTCTTGTGCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGCGGA
 TTGCTAACCGTTTACCAGCAACATCAAAGGTTCTGACTCAGGCTTCCCGT
 AACGCTAACGACGGTATCTCCATTGCGCAGACCACTGATCGAGCGCTGTG
 CGAATCAACAACAACCTGACGCGTGTGCGTGAACCTGGCGGTTCACTCTG
 CTAACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATC
 ACCAGCGGCTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAAC
 CGGCTGGAAGTCTTGGCGCAGGACAACACCTGACCATCCAGGTTGGCG
 CCAACGACGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAG
 CACTTGGGCTTGGACTCACTGAACGTGATGATGAGCGCGGATGAGCTCTGC
 TAGCCCATGGACCGAAAACCGCTGCAGAAAATGATGCCGCGCTGGCGC
 AGGTGGATGCGCTGCGCTCTGATCTGGGTGGGTACAAAACCGTTTCAAC
 TCTGCTATCACCAACCTGGGCAATACCGTAACCGTGAACATCTGCTGAGCGG
 TAGCCGTATCGAAGATTCCGACTACGCGACCGAAGTTTCCAACATGTCTC

-continued

GCGCGCAGATTTTGCAGCAGCGCGGTACTTCCGTTCTGGCGCAGGCTAAC
CAGGTCCCGCAGAACGTGCTOTCTCTGTTACGT

SEQ ID NO: 327, nucleotide sequence of
HingeForward1
ACTGAGTGCATATGGCACAAGTAATCAACACTAACAG

SEQ ID NO: 328, nucleotide sequence of
HingeReverse3
GGTCCATGGGCAAGCAGGATCCACCGGCGCT

SEQ ID NO: 329, nucleotide sequence of
HingeForward2
AGCGCCGGTGGATCCTGCTTGCCCATGGACC

SEQ ID NO: 330, nucleotide sequence of
HingeReverse4
GACTGACTGCTCAGCCTATTAACGTAACAGAGACAGCACGTTCTGCGGGA
CCTGGTTAG

SEQ ID NO: 331, nucleotide sequence of
STF2A.HingeCys
ATGGCACAAGTAATCAACACTAACAGTCTGTGCTGTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAGACGATGCGGCAGGTGAGCGGATTG
CTAACCGTTTACCAGCAACATCAAAGGTCTGACTCAGGCTTCCCGTAAC
GCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACGA
AATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTTCAGTCTGCTA
ACAGACCAACTCCCACTCTGACCTCGACTCCATCCAGGCTGAAATCACC
CAGCGCCTGAACGAAATCGACCGTGTATCCGCGCAGACTCAGTTCAACGG
CGTGAAGTCTCTGGCGCAGGACAACACCCCTGACCATCCAGGTTGGCGCCA
ACGACGGTGAAACTATCGATATCGATCTGAAGCAGATCAACTCTCAGACC
CTGGGTCTGGACTCACTGAACGTGCATGGAGCGCGGTGGATCCTGCTTG
CCCATGGACCGAAACCCGCTGCAGAAATTTGATGCCGCGCTGGCGCAGG
TGGATGGCTGCGCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCT
GCTATACCAACCTGGGCAATACCGTAAACAATCTGTCTGAAGCGCGTAG
CCGTATCGAAGATTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCG
CGCAGATTTTGCAGCAGGCGCGTACTTCCGTTCTGGCGCAGGCTAACCA
GTCCCGCAGAACGTGCTGTCTCTGTTACGT

SEQ ID NO: 332, amino acid sequence of STF2A.3'Cys
MAQVINTNSLSLLTQNNLNKSSQALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRVRELAVQSA
NSTNSQSDLDLSIAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDLSNVHGAPVDPASPWTENPLQKIDAALAQ
VDALRSDLGAVQNRNFSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSR
AQILQQAGTSVLAQANQVPQNVLSLLREFSRYPQWRPL

SEQ ID NO: 333, amino acid sequence of
STF2A.HingeCys
MAQVINTNSLSLLTQNNLNKSSQALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRVRELAVQSA
NSTNSQSDLDLSIAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDLSNVHGAPVDPASPWTENPLQKIDAALAQ
VDALRSDLGAVQNRNFSAITNLGNTVNNLSEARSRIEDSDYATEVSNMSR
AQILQQAGTSVLAQANQVPQNVLSLLR

SEQ ID NO: 334, amino acid sequence of 5'Cys.STF2A
MAQVINTNSLSLLTQNNLNKSSQALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNQLQVRVRELAVQSA
NSTNSQSDLDLSIAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDLSNVHGAPVDPASPWTENPLQKIDAALAQ
VDALRSDLGAVQNRNFSAITNLGNTVNNLSEARSRIEDSDYATEVSNMS
RAQILQQAGTSVLAQANQVPQNVLSLLR

SEQ ID NO: 335,
CSKKKKGC

SEQ ID NO: 336,
GAPVDPASPW

Example 14 Cloning and Expression of STF2.4xH1C1, a Fusion Protein Comprising Flagellin (TLR5 Agonist) and the Maturation Cleavage Site Peptide of Influenza A Hemagglutinin

Materials and Methods

[0619] Construct design: To facilitate cloning of target genes in fusion with flagellin, a cassette plasmid containing a unique BspI site at the 3' end of the flagellin gene was used (STF2.blp, SEQ ID NO: 340). This cassette was made by introducing a silent mutation (5'-GTGCTGAGCCTGT-TACGT-3') at nucleotides 1501 to 1518 of STF2, creating the unique BspI site in the plasmid cassette, pET24/STF2.blp (SEQ ID NO: 340). Synthetic genes encoding one, two, three and four copies of the maturational cleavage site fragment (H1C1) of influenza A subtype H1 (NIPSIQSR-GLFGAIAGFIE; SEQ ID NO: 341) were codon-optimized for *E. coli* expression and obtained from a commercial vendor (DNA2.0 Inc., Menlo Park, Calif.). The synthetic genes were excised with BspI enzyme and ligated by compatible ends to pET24/STF2.blp which had been treated with BspI and bacterial alkaline phosphatase (BAP). These constructs were designated as STF2.1xH1C1, STF2.2xH1C1, STF2.3xH1C1 and STF2.4xH1C1 respectively (SEQ ID NO: 342; SEQ ID NO: 343; SEQ ID NO: 344; SEQ ID NO: 345).

[0620] Similarly, a fusion gene constructed to contain the H1C1 sequence and flanked by a pair of cysteines to facilitate loop formation of the concatemers was designated as STF2.4xH1C2 (SEQ ID NO: 346). A similar approach was employed to generate a construct harboring four tandem copies of the cleavage fragment of influenza A subtype H5 (RERRRKKRGLFGAIAGFIE; SEQ ID NO: 347) designated as construct STF2.4xH5C1 (SEQ ID NO: 348). The consensus cleavage fragments representing subtypes H3 (NVPEKQTRGIFGAIAGFIE; SEQ ID NO: 349), H2 (NVPQIESRGLFGAIAGFIE; SEQ ID NO: 350) and the B-strain, B/HA (PAKLLKERGFFFGAIAGFIE; SEQ ID NO: 351) are thus amenable to this experimental approach as outlined above. In each case, the constructed plasmids were used to transform competent *E. coli* TOP10 cells and putative recombinants were identified by PCR screening and restriction mapping analysis.

[0621] The integrity of the constructs was verified by DNA sequencing and they were used to transform the expression host, BLR3 (DE3) (Novagen, San Diego, Calif.; Cat #69053). Transformants were selected on plates containing kanamycin (50 µg/mL), tetracycline (5 µg/mL) and glucose (0.5%). Colonies were picked and inoculated into 2 ml of LB medium supplemented with 25 µg/ml kanamycin, 12.5 µg/ml tetracycline and 0.5% glucose and grown overnight. Aliquots of these cultures were used to inoculate fresh cultures in the same medium formulation, which were cultured until they reached an OD₆₀₀=0.6, at which time protein expression was induced by the addition of 1 mM IPTG and culturing for 3 hours at 37° C. The cells were harvested and analyzed for protein expression.

[0622] SDS-PAGE and Western blot: Protein expression and identity were determined by gel electrophoresis and immunoblot analysis. Cells were harvested by centrifugation and lysed in Laemmli buffer. An aliquot of 10 µl of each

lysate was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel and electrophoresed (SDS-PAGE). The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands. For western blot, 0.5 µl/lane cell lysate was electrophoresed and electrotransferred to a PVDF membrane and blocked with 5% (w/v) dry milk. The membrane was probed with anti-flagellin antibody 6H11 (Inotek; Beverly, Mass.) After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockland, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega, Madison, Wis.). Bacterial clones which yielded protein bands of the correct molecular weight and reactive with the appropriate antibodies were selected for production of protein for use in biological assays.

Results and Discussion

[0623] Cloning and expression of cleavage site constructs: The maturational cleavage site fragment of hemagglutinin precursor is well-conserved across subtypes of influenza strains A and B, and could therefore be a target for the development of a universal vaccine effective against most circulating strains of influenza. A series of plasmids encoding a fusion of the TLR5 ligand with cleavage fragment was generated and expressed in *E. coli* strain BLR(DE3). As assayed by Coomassie blue staining of the SDS-PAGE gel and confirmed by immunoblot assays, the *E. coli* strains harboring the constructs STF2.1xH1C1, STF2.2xH1C1, STF2.3xH1C1, and STF2.4xH1C1 displayed bands that correspond to the predicted molecular weights of 55, 58, 60 and 62 kDa respectively. Similarly, BLR(DE3) strains expressing the constructs STF2.4xH1C2 and STF2.4xH5C1 display recombinant fusion proteins migrating with the apparent molecular weight of 62 Kda and 66 Kda respectively. These data indicate that a fusion of flagellin and cleavage fragment of HA is abundantly expressed in *E. coli*.

Expression and Purification of STF2.4xH1C1, a
Fusion Protein Comprising Flagellin (TLR5
Agonist) and the Maturational Cleavage Site
Peptide of Influenza A Hemagglutinin

Materials and methods

[0624] Bacterial cell growth and cell lysis: The STF2.4xH1C1 (SEQ ID NO: 345) construct was expressed in the *E. coli* host strain BLR(DE3). The strain was retrieved from a glycerol stock and grown in shake flasks to a final volume of 12 L. Cells were grown in LB medium containing 50 µg/ml kanamycin, 12.5 µg/ml tetracycline, 0.5% dextrose to OD₆₀₀=0.6 and induced with 1 mM IPTG for 3 h at 37° C. The cells were harvested by centrifugation (7000 rpm×7 minutes in a Sorvall RC5C centrifuge) and resuspended in 20 mM Tris-HCl, pH 8.0, 1 µg/ml DNaseI, 1 mM PMSE, protease inhibitor cocktail and 1 mg/ml lysozyme. The cells were then lysed by two passes through a microfluidizer (Microfluidics; Newton, Mass.) at 15,000 psi. The lysate was centrifuged at 45,000 g for one hour in a Beckman Optima L ultracentrifuge (Beckman Coulter; Fullerton, Calif.) to separate the soluble and insoluble fractions.

[0625] Purification of STF2.4xH1C1 (SEQ ID NO: 345): After centrifugation, the supernatant fraction was collected

and passed through a Q sepharose Fast Flow column (GE/Amersham Biosciences; Piscataway, N.J.). The flow-through fraction from this step was supplemented with Triton X-100 (Sigma; St. Louis, Mo.) to a final concentration of 1% (w/v) and passed through the same Q sepharose column. The flow through fraction was collected again and supplemented with urea to a final concentration of 8 M and citric acid to a final concentration of 20 mM. After adjusting the pH to 3.5 with concentrated HCl, the solution was passed over a Source S column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated with 20 mM citric acid, pH 3.5.

[0626] The column was then washed with 10 column volumes of equilibration buffer supplemented with 1% (w/v) Triton X-100 to remove endotoxin. The protein was then eluted in a 5-column volume linear gradient of 0 to 1 M NaCl in equilibration buffer. The STF2.4xH1C1 was then re-folded by rapid dilution to a final concentration of 0.1 mg/ml protein in refolding buffer [0.1 M Tris-HCl, pH 8.0, 0.1 M NaCl, 1% (w/v) glycerol]. The refolded protein was concentrated using a pressurized ultrafiltration stir-cell (Millipore; Billerica, Mass.) and fractionated on a Superdex 200 size-exclusion column (GE/Amersham Biosciences; Piscataway, N.J.).

[0627] SDS-PAGE and Western Blot analysis: Protein identity was determined, and purity estimated, by SDS-PAGE. An aliquot of 5 µg of each sample was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel (LifeGels; French's Forrest, New South Wales, AUS) and electrophoresed. The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands. For western blot, 0.5 µg/lane total protein was electrophoresed as described above and then electro-transferred to a PVDF membrane and blocked with 5% (w/v) dry milk before probing with anti-flagellin antibody (Inotek; Beverly, Mass.) or serum from mice immunized with a synthetic, lipidated H1C1 peptide. After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockland, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega; Madison, Wis.).

[0628] Protein assay: Total protein concentration for all proteins was determined using the Micro BCA (bicinchoninic acid) Assay (Pierce; Rockford Ill.) in the microplate format, using bovine serum albumin as a standard, according to the manufacturer's instructions.

[0629] Endotoxin assay: Endotoxin levels for all proteins were determined using the QCL-1000 Quantitative Chromogenic LAL test kit (Cambrex; E. Rutherford, N.J.), following the manufacturer's instructions for the microplate method.

[0630] TLR5 bioactivity assay: HEK293 cells constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96-well microplates (50,000 cells/well), and the STF2.4xH1C1 test protein was added. The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20° C. After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce, Rockland, Ill.; #M801E and #M802B) following the manufacturer's instructions. Optical density

was measured using a microplate spectrophotometer (FAR-Cyte, GE/Amersham; Piscataway, N.J.).

Results and Discussion

[0631] The purified and refolded STF2.4×H1C1 protein (SEQ ID NO: 345) was found to be substantially aggregated as the majority of the protein fractionated in the void volume of the Superdex S200 column (FIG. 26). The protein also had poor in vitro TLR5 activity, with an EC_{50} value approximately two orders of magnitude higher than the standard flagellin fusion protein STF2.OVA. The purified protein reacted with serum from mice immunized with a synthetic, lipidated H1C1 peptide.

Expression and Purification of STF2.1×H1C1, a Fusion Protein Comprising Flagellin (TLR5 Agonist) and the Maturational Cleavage SITE Peptide of Influenza A Hemagglutinin

Materials and methods

[0632] Bacterial cell growth and cell lysis: The STF2.1×H1C1 (SEQ ID NO: 355) construct was expressed in the *E. coli* host strain BLR(DE3). The strain was retrieved from a glycerol stock and grown in shake flasks to a final volume of 12 L. Cells were grown in LB medium containing 50 µg/ml kanamycin/12.5 µg/ml tetracycline/0.5% dextrose to $OD_{600}=0.6$ and induced with 1 mM IPTG for 3 hours at 37° C. The cells were harvested by centrifugation (7000 rpm×7 minutes in a Sorvall RC5C centrifuge) and resuspended in 20 mM Tris-HCl, pH 8.0, 1 µg/ml DNaseI, 1 mM PMSF, protease inhibitor cocktail and 1 mg/ml lysozyme. The cells were then lysed by two passes through a microfluidizer (Microfluidics; Newton, Mass.) at 15,000 psi. The lysate was centrifuged at 45,000 g for one hour in a Beckman Optima L ultracentrifuge (Beckman Coulter; Fullerton, Calif.) to separate the soluble and insoluble fractions.

[0633] Purification of STF2.1×H1C1 (SEQ ID NO: 342): After centrifugation, the supernatant fraction was collected and passed through a Q sepharose Fast Flow column (GE/Amersham Biosciences; Piscataway, N.J.). The flow-through fraction from this step was supplemented with Triton X-100 (Sigma; St. Louis, Mo.) to a final concentration of 1% (w/v) and passed through the same Q sepharose column. The flow-through fraction was collected again and supplemented with urea to a final concentration of 8 M and citric acid to a final concentration of 20 mM. After adjusting the pH to 3.5 with concentrated HCl, the solution was passed over a Source S column (GE/Amersham Biosciences; Piscataway, N.J.) equilibrated with 20 mM citric acid, pH 3.5.

[0634] The column was then washed with 10 column volumes of equilibration buffer supplemented with 1% (w/v) Triton X-100 to remove endotoxin. The protein was then eluted in a 5-column volume linear gradient of 0 to 1 M NaCl in equilibration buffer. The STF2.1×H1C1 was refolded by rapid dilution to a final concentration of 0.1 mg/ml protein in refolding buffer [0.1M Tris-HCl, pH 8.0/0.1M NaCl/1% (w/v) glycerol]. The refolded protein was concentrated using a pressurized ultrafiltration stir-cell (Millipore; Billerica, Mass.) and fractionated on a Superdex 200 size-exclusion column (GE/Amersham Biosciences; Piscataway, N.J.).

[0635] SDS-PAGE and western blot analysis: Protein identity was determined, and purity estimated, by SDS-

PAGE. An aliquot of 5 µg of each sample was diluted in SDS-PAGE sample buffer with or without 100 mM DTT as a reductant. The samples were boiled for 5 minutes and loaded onto a 10% SDS polyacrylamide gel (LifeGels; French's Forrest, New South Wales, AUS) and electrophoresed. The gel was stained with Coomassie R-250 (Bio-Rad; Hercules, Calif.) to visualize protein bands. For western blot, 0.5 µg/lane total protein was electrophoresed as described above and the gels were then electro-transferred to a PVDF membrane and blocked with 5% (w/v) dry milk before probing with anti-flagellin antibody (Inotek; Beverly, Mass.) or serum from mice immunized with a Pam3Cys.H1C1 peptide (SEQ ID NO: 358). After probing with alkaline phosphatase-conjugated secondary antibodies (Pierce; Rockland, Ill.), protein bands were visualized with an alkaline phosphatase chromogenic substrate (Promega; Madison, Wis.).

Results and Discussion

[0636] The purified and refolded STF2.1×H1C1 protein (SEQ ID NO: 342) was found to be monomeric as judged by the elution profile on a Superdex 200 gel filtration column. The majority of the protein was found in the included volume with the major peak eluting at approximately 14 mls. This corresponds very closely with the known elution profile of purified, monomeric flagellin on this column. Almost no protein was seen eluting in the void volume, known to be approximately 7 mls for this column, demonstrating that virtually no aggregates are present.

Immunogenicity and Efficacy of PAM3Cys.H1C1, a Fusion Peptide Comprising PAM3CYS (TLR2 Agonist) and the Maturational Cleavage Site Peptide of Influenza A Hemagglutinin

Materials and methods

[0637] Peptide design and synthesis: Pam3 (tri-palmytyl) is the natural ligand for Toll-like Receptor 2 (TLR2), and it is natively expressed as the lipidation motif of bacterial lipoprotein (BLP, SEQ ID NO: 357). Pam3 can be made by chemical synthesis and conjugated to macromolecules such as proteins or coupled to the N-terminus of synthetic peptides using standard peptide synthesis chemistry. This strategy usually involves the synthesis of a peptide of interest concluded by the coupling of Pam3-modified cysteine (Pam3Cys) to the amino terminus to yield the lipopeptide of interest. This approach was used to synthesize a lipidated HA cleavage fragment peptide, Pam3Cys.H1C1 (Pam3Cys-SLWSEENIPSIQSRGLFGAAGFIEE, SEQ ID NO: 358).

[0638] Mice and immunization: Female BALB/c mice (Jackson Laboratory, Bar Harbor, Me.) were used at the age of 6-8 weeks. Mice were divided into groups of 10 and received inguinal subcutaneous (s.c) immunizations on days 0 and 14 as follows:

[0639] 8) PBS (Phosphate Buffered Saline).

[0640] 9) 20 µg of H1C1 native peptide (SEQ ID NO: 358) in saline buffer (10 mM Histidine, 10 mM Tris, 75 mM NaCl, 5% (vol/vol) sucrose, 0.02% (w/v) Polysorbate-80, 0.1 mM EDTA, 0.5% (v/v) ethanol, pH 7.2)

[0641] 10) 20 µg of Pam3Cys.H1C1 peptide (SEQ ID NO: 358) in saline buffer

[0642] An additional group of five mice received an experimentally determined sublethal challenge with 8×10^1 egg infectious dosages (EID) PR/8/34 and were allowed to convalesce for >21 days. These animals were then used as immune convalescent positive controls during the challenge studies. Mice were bled on days 10 (primary) and 21 (boost), and sera were clarified by clotting and centrifugation and stored at -20°C .

[0643] Serum antibody determination: H1C1-specific IgG levels were determined by ELISA. 96-well ELISA plates (Costar (Cat #9018) Corning, N.Y.) were coated overnight at 4°C with $100\ \mu\text{l/well}$ H1C1 peptide (SEQ ID NO: 358) in PBS (5 $\mu\text{g/ml}$). Plates were blocked with $200\ \mu\text{l/well}$ of Assay Diluent Buffer (ADB; BD Pharmingen, (Cat#: 555213) (San Diego, Calif.) for one hour at room temperature. The plates were washed three times in PBS+0.05% (v/v) Tween 20 (PBS-T). Dilutions of the sera in ADB were added ($100\ \mu\text{l/well}$) and the plates were incubated overnight at 4°C . The plates were washed three times with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical, West Grove, Pa. (Cat#: 115-035-146)) diluted in ADB were added ($100\ \mu\text{l/well}$) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce (Cat 34028), Rockford, Ill.) and monitoring color development, A_{450} was measured on a Tecan Farcyte (Durham, N.C.) microplate spectrophotometer.

[0644] Influenza virus challenge of mice. To assess efficacy, mice immunized as described above were challenged on day 28 by intranasal administration of an LD_{50} (dose lethal to 90% of mice) (8×10^5 EID) of influenza A isolate PR/8/34. Animals were monitored daily for 21 days following the challenge for survival, weight loss and clinical presentation. The % weight loss was calculated based on the mean of ((Daily weight (g)/Initial weight (g) day 28) $\times 100$) of each individual animal per group. Clinical scores were assigned as follows: 4 pts=healthy, 3 pts=reduced grooming, 2 pts=reduced physical activity and 1 pt=moribund. (Experimental results for clinical scores and weight loss reflect the results based on surviving animals on the day evaluated).

Results and Discussion

[0645] Induction of H1C1-specific antibody responses following immunization with Pam3Cys.H1C1: Mice were immunized with native H1C1 peptide (SEQ ID NO: 358) and the same peptide modified by the addition of an amino terminal Pam3Cys residue, to test the hypothesis that linkage of the peptide to a TLR2 ligand would increase its immunogenicity. Immunogenicity was determined by measuring levels of antibodies to native H1C1 (SEQ ID NO: 358) in the sera of the immunized mice. The results show clearly that mice immunized with the Pam3Cys-modified peptide generated higher antibody titers to the immunizing peptide backbone sequence than did mice immunized with the native peptide (SEQ ID NO: 358) (FIG. 27).

[0646] Protection from lethal influenza virus challenge following immunization with Pam3Cys.H1C1 The results in FIG. 27 demonstrated that immunization of mice with Pam3Cys.H1C1 generated an antibody response that recognized native H1C1 peptide. In order to evaluate efficacy, the same mice were challenged on day 28 with an LD_{50} (8×10^3 EID) of PR/8/34 virus administered intra-nasally. Mice were

monitored daily for 21 days following the challenge for survival, weight loss and clinical presentation. As shown in FIG. 28, PBS-immunized mice died between 6 and 10 days post-challenge, while convalescent mice survived beyond the 21-day observation period. Mice immunized with native H1C1 (SEQ ID NO: 358) began to die by day 6 post-infection at a rate similar to the PBS control mice, although 2 of the 10 mice survived beyond the 21-day observation period. In contrast, and similar to the convalescent mice, all mice immunized with Pam3Cys.H1C1 survived beyond the 21-day observation period. Thus, while H1C1 alone is not potentially immunogenic (FIG. 27) nor protective (FIG. 28), the same sequence coupled to a TLR2 ligand is both immunogenic and protective.

SEQ ID NO: 340 STF2.blp
 ATGGCACAAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAGACGATGCGGCAGGTCAAGCGGATT
 GCTAACCGTTTCACCGCAACATCAAGGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCGAGCCACTGAAGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTCAGTCTGCT
 AACAGCACCACTCCAGTCTGACCTCGACTCCATCCAGGCTGAATCAC
 CCAGCGCTGAACGAAATCGACCGTGTATCCGCGCAGACTCAGTTCAACG
 GCGTGAAGTCTTGGCGCAGGACAACACCTTACCATCCAGGTTCAGCGCC
 AACGACGGTGAACATATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAAGATA
 CAGCAGTAAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGCTCTTGATGATGCAGCTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGCGTTAAATTTGACGCGAGATAATAACA
 AGTACTTTTGTACTATTGGTGGCTTACTGGTGTCTGATCCGCCAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCTTTCGCGC
 TGGCGCACTAAAACCAATGCCTGCTGGTGGCGCACTAAACAGAGAAG
 TACAGGAGTTAAAAGATACACCGGCACTGTTTTCAGCAGATGCTAAAAGT
 GCCTTAATGCTGGCGGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTATACCGATAAAAATGGTAAGCAATTTGAAGGGCGTT
 ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGCGATGATGAAGCG
 ACAGGAGCAATTAAGCTAAAACCTACAAGTTATACTGCTGCTGACGGCAC
 TACCAAAACAGCGGTAACCAACTGGGTGGCGTAGACGGTAAACCGAAG
 TCGTTACTATCGACGGTAAAACCTACAATGCCAGCAAGCGCGTGGTCTAT
 GATTTCAAAGCACAAACAGAGCTGGCGGAAGCAGCGCTAAAACACCGA
 AAACCGCTGCAGAAAATTTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGGCGTACAAACCGTTTCAACTCTGATATCAACAAAC
 TGGGCAATACCGTAAACAATCTGTCTGAAGCGCTAGCGGTATCGAAGAT
 TCCGACTACGCGACCGAAGTTTCCCAACATGTCTCGCGCGCAGATTCTGCA
 GCAGGCGGTACTTCCGTTCTGGCGCAGGCTAACACAGGTCCCGCAGAAGC
 TGCTGAGCCTGTTACGT

SEQ ID NO: 341 HA maturation cleavage fragment
 (H1N1)
 NIPSTQSRGLFGAIAGFIE

SEQ ID NO: 342 STF2.1xH1C1
 MAQVINTNSLSLLTQNNLNKSQALGTAIERLSSGLRINSALDAAGQAIA
 NRFTANIKGLTQASRNANDGISIAQTTEGALNEINNLRVRELAVQSAN
 STNSQSDLDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGAN
 DGETIDIDLKQINSQTLGLDLSLNVQKAYDVKDTAVTTKAYANNNGTTLDVS
 GLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKNG
 DYEATDGTVTTLAAGATKTTMPAGATTKTEVQELKDTTPAVVSADAKNALI
 AGGVDATDANGAELVKMSYTDKNGKTIIEGGYALKAGDKYAAADYEATGA
 IKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTTIDGKTYNASKAAGHDFK
 AQPELAEEAAKTTENPLQKIDAALAQVDALRDLGAVQNRNFSAITNLGN
 TVNNLSSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLAQANQVPQNVLS
 LLAMEWENIPSIQSRGLFGAIAGFIE

SEQ ID NO: 343 STF2.2xH1C1
 MAQVINTNSLSLLTQNNLNKSQALGTAIERLSSGLRINSALDAAGQAIA
 NRFTANIKGLTQASRNANDGISIAQTTEGALNEINNLRVRELAVQSAN
 STNSQSDLDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGAN
 DGETIDIDLKQINSQTLGLDLSLNVQKAYDVKDTAVTTKAYANNNGTTLDVS
 GLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAAKNG
 DYEAVNATDGTVTTLAAGATKTTMPAGATTKTEVQELKDTTPAVVSADAKNA
 LIAGGVDATDANGAELVKMSYTDKNGKTIIEGGYALKAGDKYAAADYEATGA
 IKAIAKTTTSYTAADGTTKTAANQLGGVDGKTEVVTTIDGKTYNASKAAGHD

-continued

FKAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRNFSAITNL
GNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLIAQANQVPQN
VLSLLAMEWENIPSIQSRGLFGAIAGFIEEWENIPSIQSRGLFGAIAGFIE

SEQ ID NO: 344 STF2.3xH1C1

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNLRVRELAVQSA
NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
DGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTLLDVS
GLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAANKG
DYEYVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKNA
LIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYAAADYDEAT
GAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGHD
FKAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRNFSAITNL
GNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLIAQANQVPQN
VLSLLAMEWENIPSIQSRGLFGAIAGFIEEWENIPSIQSRGLFGAIAGFIE
EWENIPSIQSRGLFGAIAGFIE

SEQ ID NO: 345 STF2.4xH1C1

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNLRVRELAVQSA
NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTLLDVS
GLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAANKG
DYEYVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKNA
ALIAAGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYAAADYDEA
TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
DFAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRNFSAITNL
LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLIAQANQVPQN
VLSLLAMEWENIPSIQSRGLFGAIAGFIEEWENIPSIQSRGLFGAIAGFIE
EWENIPSIQSRGLFGAIAGFIEEWENIPSIQSRGLFGAIAGFIE

SEQ ID NO: 346 STF2.4xH1C2

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNLRVRELAVQSA
NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
DGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTLLDVS
GLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAANKG
DYEYVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKNA
LIAGGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYAAADYDEAT
GAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGHD
FKAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRNFSAITNL
LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLIAQANQVPQN
VLSLLAGCGSEWENIPSIQSRGLFGAIAGFIEEWENIPSIQSRGLFGAIA
GFIEEWENIPSIQSRGLFGAIAGFIEEWENIPSIQSRGLFGAIAGFIESG
C

SEQ ID NO: 347 HA maturation cleavage fragment
(H5N1)

RRRRRKRGLFGAIAGFIE

SEQ ID NO: 348 STF2.4xH5C1

MAQVINTNSLSLLTQNNLNKSQSALGTAIERLSSGLRINSKDDAAGQAI
ANRFTANIKGLTQASRNANDGISIAQTTEGALNEINNLRVRELAVQSA
NSTNSQSDLSIQAEITQRLNEIDRVSGQTQFNGVKVLAQDNTLTIQVGA
NDGETIDIDLKQINSQTLGLDSLNVQKAYDVKDTAVTTKAYANNGTLLDVS
GLDDAAIKAAATGGTNGTASVTGGAVKFDADNNKYFVTIGGFTGADAANKG
DYEYVNVATDGTVTLAAGATKTTMPAGATTKEVQELKDTPAVVSADAKNA
ALIAAGVDATDANGAELVKMSYTDKNGKTIEGGYALKAGDKYAAALYDEA
TGAIAKAKTTSYTAADGTTKTAANQLGGVDGKTEVVTIDGKTYNASKAAGH
DFAQPELAEAAAKTTENPLQKIDAALAQVDALRSDLGAVQNRNFSAITNL
LGNTVNNLSEARSRIEDSDYATEVSNMSRAQILQQAGTSVLIAQANQVPQN
VLSLLAEWERERKRGLFGAIAGFIEEWERERRRKRGLFGAIAGFIEEW
ERRRRKRGLFGAIAGFIEEWERERRRKRGLFGAIAGFIE

SEQ ID NO: 349 HA maturation cleavage fragment
(H3N2)

NVPEKQTRGIFGAIAGFIE

SEQ ID NO: 350 HA maturation cleavage fragment
(H2N1/H2N2/H2N3/H2N5/H2N8/H2N9)

NVPQIESRGLFGAIAGFIE

SEQ ID NO: 351 HA maturation cleavage fragment
(B strain)

PAKLLKERGFPGAIAGFLE

-continued

SEQ ID NO: 352 STF2.4xH1C1

ATGGCACAAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTGAGCGGATT
GCTAACCGTTTACCAGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTGCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCTGAACGAAATCGACCGTGTATCCGCGCAGACTCAGTTCAACG
GCGTGAAGTCTGCGCGCAGGACAACACCTGACCATCCAGGTTGCGCGCC
AACGACCGTGAACATTCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGACGCTATTAAAGCGCGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAAACCGGTGGTGGGTAAATTTGACGCGAGATAATAACA
AGTACTTTGTACTATTGGTGGCTTTACTGGTGTGATGCCGCCAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACCGGTACAGTAACCCCTTGGCGC
TGGCGCAACTAAACACCAATGCTGCTGGTGGCGCACTAAACGAAAG
TACAGGAGTTAAAGATACACCGCGAGTTGTTTACGAGATGCTAAAAAT
GCCTTAATTGCTGGCGCGGTTGACGCTACCGATGCTAATGGCGCTGAGTT
TACGCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTTGAAGCGGTT
ATGCGCTTAAAGCTGGCGATAAGTATTACGCCGAGATTACGATGAAGCG
ACAGGAGCAATTAAGCTAAACACTACAAGTTATCTGCTGCTGACGGCAC
TACCAAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGGTAAACCGAAG
TCGTTACTATCGAGCGTAAACCTACAATGCCAGCAAGCGCGTGGTCAAT
GATTTCAAGACACAACAGAGCTGGCGGAGCGAGCGCTTAAACACCGCA
AAACCGCTGCAGAAAAATGATGCGCGCTGGCGCGTGGGATGCGCTGCTG
GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
CTGGGCAATACCGTAAACAATCTGCTGTAAGCGCGTAGCGGATTCGAAGA
TTCGAGCTACGCGACCGAAGTTTCAACATGTTCTGCGCGAGGATTCTGTC
AGCAGCGCGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
GTGCTGTCTCTGTTAGCGATGGAATGGGAGAACATCCCTAGCATCCAATC
TCGCGCGCTGTTTGGCGCTATCGCGCGCTTTATCGAAGAAATGGAGAAC
TCCCAGCATCCAATCTCGCGGTCTGTTTGGTGGCATCGCTGGTTTCATC
GAGGAGTGGGAGAACATTCCTAGCATTCAAAGCGCTGGCGCTGTTGGCGC
TATTGCGAGGTTTATGTAAGAAATGGGAAAAATCCCGCTGATCAATCC
CGGCTGTGTTGGCGCGATCGCAGGTTTCAATTGAATAATAAGCTAAGC

SEQ ID NO: 353 STF2.3xH1C1

ATGGCACAAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
CTGGTCTGCGTATCAACAGCGCGAAGACGATGCGGCAGGTGAGCGGATT
GCTAACCGTTTACCAGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
CGCTAACGACGGTATCTCCATTGCGCAGACCACTGAAGCGCGCTGAACG
AAATCAACAACAACCTGCAGCGTGTGCGTGAAGTGGCGGTTCACTGCTGCT
AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAAATCAC
CCAGCGCTGAACGAAATCGACCGTGTATCCGCGCAGACTCAGTTCAACG
CGGTGAAGTCTGCGCGCAGGACAACACCTGACCATCCAGGTGCGCGCC
AACGACGGTGAACATTCGATATCGATCTGAAGCAGATCAACTCTCAGAC
CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
TCGGGTCTTGATGATGACGCTATTAAAGCGGCTACGGGTGGTACGAATGG
TACGGCTTCTGTAAACCGGTGGTGGCGTTAAATTTGACGCGAGATAATAACA
AGTACTTTGTTACTATTGGTGGCTTTACTGCTGATGCGCGCAAAAAAT
GGCGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCCTTGGCGC
TGGCGCAACTAAACACCAATGCTGCTGGTGGCACAACATAAACAGAAG
TACAGGAGTTAAAGATACACCGCGAGTTGTTTACGAGATGCTAAAAAT
GCCTTAATTGCTGGCGCGCTTACGCTACCGATGCTAATGGCGCTGAGTT
GGTCAAAATGTCTTATACCGATAAAAAATGGTAAGACAATTTGAAGCGGTT
ATTGCGCTTAAAGCTGGCGATAAGTATTACGCGCAGATGATACGATGAGG
ACAGGAGCAATTAAGCTAAACACTACAAGTTATCTGCTGCTGACGGCAC
TACCAAAAACAGCGGCTAACCAACTGGGTGGCGTAGACGTTAAACCGAAG
TGCTTACTATCGACGGTAAACCTACAATGCCAGCAAGCGCGTGGTCAAT
GATTTCAAGACACAACAGAGCTGGCGAAGCAGCGCTAAACACCGCA
AAACCGCTGCAGAAAAATGATGCGCGCTGGCGCAGGTGGATGCGCTGC
GCTCTGATCTGGGTGCGGTACAAAACCGTTTCAACTCTGCTATCACCAAC
CTGGGCAATACCGTAAACAATCTGCTGAAGCGCTGACCGTATCGAAGA
TTCGACTACGCGACCGAAGTTTCAACATGCTCGCGCGCAGATTCTGC
AGCAGCGCGTACTTCCGTTCTGGCGCAGGCTAACAGGTCCCGCAGAAC
GTGCTGTCTCTGTTAGCGATGGAATGGGAGAACATCCCTAGCATCCAATC
TCGCGCGCTGTTTGGCGCTATCGCGCGCTTTATCGAAGAAATGGAGAAC
TCCCAGCATCCAATCTCGCGGTCTGTTTGGTGGCATCGCTGGTTTCATC
GAGGAGTGGGAGAACATTCCTAGCATTCAAAGCGCTGGCGCTGTTGGCGC
TATTGCGAGGTTTATGTAAGAAATGGGAAAAATCCCGCTGATCAATCC
CGGCTGTGTTGGCGCGATCGCAGGTTTCAATTGAATAATAAGCTAAGC

-continued

SEQ ID NO: 354 STF2.2xH1C1
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCTGAAGTGGCGTTTCACTCTAAT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAATACAC
 CCAGCGCTTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAGTCCCTGGCGCAGGACAAACCCCTGACCATCCAGTTTGGCGCC
 AACGACGGTGAAGTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACACTGGATGTA
 TCGGGTCTTGATGATGACGCTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTTAAATTTGACGCGAGATAATAACA
 AGTACTTTGTGTTACTATTGGTGGCTTTTACTGGTGTGATGCGCGCAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAACACCAATGCTGCTGGTGGCAGCACTAAACAGAAAG
 TACAGGAGTTAAAGATACACCGCAGTTGTTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGTGGCGGATGAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAGCTAAACCTACAAGTTATACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCACTGGGTGGCGTAGACGGTAAACCGGAAG
 TCGTTACTATCGACGCTAAACCTTACAATGCCAGCAAGCGCTGGTCTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAACCCACCGA
 AAACCCGCTGCAGAAAATTTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGGCGGTACAAAACCGTTTCAACTCTGCTATCAACCAAC
 CTGGCAATACCGTAAACAATCTGCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGCG
 AGCAGCGCGGTACTTCCGTTCTGCGCGCAGGCTAACAGGTTCCCGCAGAAC
 GTGCTGAGCCTGTTAGCGATGGAATGGGAAAATATCCCTAGCATCCAATC
 TCGGGTCTGTTCCGTTGCTATTGCTGGCTTCATCGAGGAATGGGAGAAC
 TCCCATCTATTACGTTCTCGCGCGCTGTTTGGTGGCATCGCGGGTTTATT
 GAGTGATAAGCTTAGCCCAAGG

SEQ ID NO: 355 STF2.1xH1C1
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGTCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG
 AAATCAACAACAACCTGCAGCGTGTGCTGAAGTGGCGTTTCACTCTGCT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAATACAC
 CCAGCGCTTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAGTCCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAGTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACTAGGATGTA
 TCGGGTCTTGATGATGACGCTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTTAAATTTGACGCGAGATAATAACA
 AGTACTTTGTGTTACTATTGGTGGCTTTTACTGGTGTGATGCGCGCAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAACACCAATGCTGCTGGTGGCAGCACTAAACAGAAAG
 TACAGGAGTTAAAGATACACCGGCGAGTTGTTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGTGGCGGATGAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAGCTAAACCTACAAGTTATACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCACTGGGTGGCGTAGACGGTAAACCGGAAG
 TCGTTACTATCGACGCTAAACCTTACAATGCCAGCAAGCGCTGGTCTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAACCCACCGA
 AAACCCGCTGCAGAAAATTTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGGCGGTACAAAACCGTTTCAACTCTGCTATCAACCAAC
 CTGGCAATACCGTAAACAATCTGCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGCG
 AGCAGCGCGGTACTTCCGTTCTGCGCGCAGGCTAACAGGTTCCCGCAGAAC
 GTGCTGAGCCTGTTAGCGATGGAATGGGAAAATATCCCTAGCATCCAATC
 TCGGGTCTGTTCCGTTGCTATTGCTGGCTTCATCGAGGAATGGGAGAAC
 TCCCATCTATTACGTTCTCGCGCGCTGTTTGGTGGCATCGCGGGTTTATT
 GAGTGATAAGCTTAGCCCAAGG

SEQ ID NO: 356 STF2.4xH1C1
 ATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGCTGACCCAGAATAA
 CCTGAACAAATCCAGTCCGCACTGGGCACCGCTATCGAGCGTCTGTCTT
 CTGGTCTGCGTATCAACAGCGCGAAAGACGATGCGGCAGGTCAGGCGATT
 GCTAACCGTTTACCGCGAACATCAAAGGCTGACTCAGGCTTCCCGTAA
 CGCTAACGACGGTATCTCCATTGCGCAGACCCTGAAGGCGCGCTGAACG

-continued

AAATCAACAACAACCTGCAGCGTGTGCGTGAACCTGGCGGTTCACTGCTGT
 AACAGCACCAACTCCAGTCTGACCTCGACTCCATCCAGGCTGAATACAC
 CCAGCGCTTGAACGAAATCGACCGTGTATCCGGCCAGACTCAGTTCAACG
 GCGTGAAGTCCCTGGCGCAGGACAAACCCCTGACCATCCAGGTTGGCGCC
 AACGACGGTGAAGTATCGATATCGATCTGAAGCAGATCAACTCTCAGAC
 CCTGGGTCTGGACTCACTGAACGTGCAGAAAGCGTATGATGTGAAGATA
 CAGCAGTAACAACGAAAGCTTATGCCAATAATGGTACTACTAGGATGTA
 TCGGGTCTTGATGATGACGCTATTAAGCGGCTACGGGTGGTACGAATGG
 TACGGCTTCTGTAACCGGTGGTGGGTTAAATTTGACGCGAGATAATAACA
 AGTACTTTGTGTTACTATTGGTGGCTTTTACTGGTGTGATGCGCGCAAAAT
 GCGGATTATGAAGTTAACGTTGCTACTGACGGTACAGTAACCCTTGCGGC
 TGGCGCAACTAAACACCAATGCTGCTGGTGGCAGCACTAAACAGAAAG
 TACAGGAGTTAAAGATACACCGGCGAGTTGTTTTCAGCAGATGCTAAAAAT
 GCCTTAATTGCTGGCGCGGTTGACGCTACCGATGCTAATGGCGCTGAGTT
 GGTCAAAATGCTTTATACCGATAAAAAATGGTAAGACAATTGAAGCGGTT
 ATGCGCTTAAAGTGGCGGATGAAGTATTACGCCGAGATTACGATGAAGCG
 ACAGGAGCAATTAAGCTAAACCTACAAGTTATACTGCTGCTGACGGCAC
 TACCAAAACAGCGGCTAACCACTGGGTGGCGTAGACGGTAAACCGGAAG
 TCGTTACTATCGACGCTAAACCTTACAATGCCAGCAAGCGCTGGTCTCAT
 GATTTCAAAGCACAACAGAGCTGGCGGAAGCAGCGCTAAACCCACCGA
 AAACCCGCTGCAGAAAATTTGATGCCGCGCTGGCGCAGGTGGATGCGCTGC
 GCTCTGATCTGGGTGGCGGTACAAAACCGTTTCAACTCTGCTATCAACCAAC
 CTGGCAATACCGTAAACAATCTGCTGAAGCGCTAGCCGTATCGAAGA
 TTCCGACTACGCGACCGAAGTTTCCAACATGTCTCGCGCGCAGATTCTGCG
 AGCAGCGCGGTACTTCCGTTCTGCGCGCAGGCTAACAGGTTCCCGCAGAAC
 GTGCTGCTCTGTTAGCGGTTGTTGGTTCCGAGTGGGAAAATATTCGCTC
 TATCCAGAGCGGTGGTCTGTTTGGCGCAATTTGCTGGCTTCATTGAAGAAT
 GGGAAAACATCCCGTCCATCCAGGCGCGTGGCTGTTTGGCGCGCAATGCT
 GGTTCATCGAGGAATGGGAAAACATTCGCTCCATCCAGTCCCGCGGCTCT
 GTTTGGCGCTATCGCGGTTTTCATTGAGGAATGGGAAAATATCCCTTCCA
 TCCCATCTCGTGGTCTGTTTGGCGCGGATTGCGAGGCTTTATCGAATCTGGT
 TGCTAATAAGCTAAGC

SEQ ID NO: 357 Bacterial lipoprotein of *E. coli*
 MKATKLVLGAVILGSTLLAGCSSNAKIDQLSSDVQLTNAKVDQLSNDVNA
 MRSVDVQAAKDDAAPANQRLDNMATKYRK

SEQ ID NO: 358 H1C1 native peptide
 SLWSEENTPSIQSRFLGFAIAGFIE

Example 15

Flagellin-M2e Fusion Proteins

[0647] M2e is conserved across multiple influenza A subtypes (also referred to herein as “strain”). M2e is at least a portion of the M2 protein, in particular, a 24 amino-terminus (also referred to herein as an “ectodomain”) of the M2 protein. The M2 ectodomain is relatively small amino acid sequence (24 amino acids) compared to HA (about 566 amino acids) and NA (about 469 amino acids). The M2e sequence of exemplary avian influenza A isolates differs from that of human isolates, but is highly-conserved among the avian isolates (see, for example, SEQ ID NOS: 544-556, 570 and 573-578). Four tandem copies of M2e fused to the carboxy terminus of a flagellin STF2 (full-length or STF2 hinge region-deleted) were generated. The STF2 without the hinge region is also referred to herein as “STF2A.”

Construction of Fusion Protein

[0648] The carboxy-terminal fusion of the synthetic 4xM2e sequence (4 consecutive 24 amino acid sequences) with STF2 was constructed as follows. The pET24A vector was purchased from Novagen, San Diego, Calif. The strategy employed the Seamless Cloning Kit (Catalog number 214400) from Stratagene (La Jolla, Calif. www.stratagene.com) performed by DNA 2.0 Inc. (Menlo Park, Calif.). The gene encoding the fusion protein was in pDrive 4xM2e G00448 and was used as a PCR template for insert preparation for construction of the C-terminal fusion expres-

sion construct with STF2. The synthetic 4×M2e construct pDrive 4×M2e G00448 was used as a template for PCR as outlined in the Seamless Cloning Kit (Catalog number 214400) from Stratagene (La Jolla, Calif.). The expected product from this amplification includes the 318 bp and the restriction enzyme sites incorporated into the oligonucleotides used to amplify this insert. The procedure was as follows:

PCR Conditions

1 μL –20 ng of pDrive 4×M2e G00448
 5 μL of 10× cloned Pfu polymerase buffer
 1 μL of 40 mM dNTP mix
 1 μL –10 μmol of forward primer 4×M2eforbs1
 1 μL –10 μmol of reverse primer 4×M2erevwsto
 40 μL ddH₂O

[0649] Immediately before starting the thermal cycling 1 μL of PfuTurbo DNA Polymerase the following were added.

4×M2eforbs1 primer sequence: (SEQ ID NO: 566)
 5' –CGCTCTTCAMTGAGCTTGCTGACTGAGGTTGAGACCCGATTC
 4×M2erevwsto primer sequence: (SEQ ID NO: 567)
 5' –CGCTCTTCACGCTTATTATCTAGACGGGCTGAGCTATCGTTAGAGC
 GAG

[0650] This reaction was cycled as follows on a Thermo Hybaid Px thermal cycler (Waltham, Mass.).

Initial cycle	
Temperature	Duration
95°	3 minutes
65°	1 minute
72°	1 minute

[0651]

Subsequent nine cycles	
Temperature	Duration
95°	45 seconds
65°	35 seconds
72°	1 minute

[0652] At this point the following was added to each reaction.

5 μL of 10× cloned Pfu polymerase buffer
 1 μL of 5-methyl dNTP mix
 44 μL ddH₂O

[0653] Subsequently the following thermal cycling was repeated five times.

Temperature	Duration
95°	45 seconds
65°	35 seconds
72°	1 minute

[0654] The 100 μL product was brought to a volume of 300 μL by the addition of TE buffer. The resulting product was phenol chloroform (Invitrogen Carlsbad, Calif.-Catalog number 15593-031) extracted once and chloroform extracted once. The amplification product was then ethanol precipitated by addition of 30 μL of Sodium acetate buffer pH 5.2 and 750 μL of 100% Ethanol. The DNA pellet washed twice with 300 μL 70% Ethanol allowed to air dry for ten minutes and then resuspended in 50 μL TE buffer.

Amplification of Vector STF2 in pET24.

[0655] The previously constructed pET24a/STF2.M2e construct was used as a template for PCR as outlined in the Seamless Cloning Kit (Catalog number 214400) from Stratagene (La Jolla, Calif.). The expected product from this amplification includes the whole of the pET24 plasmid plus the STF2 sequences but does not include the single copy of M2E that exists in this construct. The procedure was as follow:

PCR Conditions

1 μL –40 ng of STF2.M2E pET22-2
 5 μL of 10× cloned Pfu polymerase buffer
 1 μL of 40 mM dNTP mix
 1 μL –10 μmol of primer 4×MECpET24
 1 μL –10 μmol of primer 4×M2eC-STF2
 40 μL ddH₂O

[0656] Immediately before starting the thermal cycling the following were added:

[0657] 1 μL of PfuTurbo DNA Polymerase

4×MECpET24 primer sequence: (SEQ ID NO: 568)
 5' –GCTCTTCAGCGGCTGAGCAATAACTAGCATAACCCCTTGGG
 4×M2eC-STF2 primer sequence: (SEQ ID NO: 569)
 5' –CGCTCTTCACAGACGTAACAGAGACAGCAGCTTCTGCGG

[0658] This reaction was cycled as follows on a Thermo Hybaid Px thermal cycler (Waltham, Mass.).

Initial cycle	
Temperature	Duration
95°	3 minutes
65°	1 minute
72°	18 minutes

[0659]

Subsequent nine cycles	
Temperature	Duration
95°	45 seconds
65°	35 seconds
72°	18 minutes

[0660] At this point the following was added to each reaction.

5 μ L of 10 $\times$ cloned Pfu polymerase buffer

1 μ L of 5-methyl dNTP mix

44 μ L ddH₂O

[0661] Subsequently the following thermal cycling was repeated five times.

Temperature	Duration
95°	45 seconds
65°	35 seconds
72°	18 minutes

[0662] The 100 μ L product was brought to a volume of 300 μ L by the addition of TE buffer. The resulting product was phenol chloroform (Invitrogen Carlsbad, Calif.-Catalog number 15593-031) extracted once and chloroform extracted once. The amplification product was then ethanol precipitated by addition of 30 μ L of Sodium acetate buffer pH 5.2 and 750 μ L of 100% Ethanol. The DNA pellet washed twice with 300 μ L 70% Ethanol allowed to air dry for ten minutes and then resuspended in 50 μ L TE buffer.

Digestion and Ligation of Vector and Insert Amplifications

[0663] Eam 1104 I digests were set up separately for vector and insert as follows:

[0664] 30 μ L of amplified product after ethanol precipitation

[0665] 5 μ L of 10 $\times$ Universal buffer (Supplied with Seamless Cloning Kit)

[0666] 4 μ L Eam 1104 I restriction enzyme (Supplied with Seamless Cloning Kit)

[0667] 11 μ L ddH₂O

[0668] Digests were mixed gently and incubated at 37° C. for one hour and ligation reactions of vector and insert products were prepared as above performed as follows (Reagents supplied with Seamless Cloning Kit):

[0669] Ingredients added in order listed:

[0670] 9 μ L ddH₂O

[0671] 5 μ L of Eam 1104 I digested 4 $\times$ M2e amplified insert

[0672] 5 μ L of Eam 1104 I digested STF2.M2E pET22-2 amplified vector

[0673] 2 μ L 10 $\times$ Ligase buffer

[0674] 2 μ L 10 mM rATP

[0675] 1 μ L T4 DNA Ligase (diluted from stock 1:16)

[0676] 1 μ L Eam 1104 I restriction enzyme

[0677] The ligation reactions were mixed gently and incubated for 30 minutes at 37° C. The ligations were then stored on ice until transformed into XL-10 competent cells (Stratagene Catalog number 200314) later than same day.

Transformation of Ligation into XL-10 Competent Cells

[0678] Eppendorf tubes were chilled for ten minutes while the XL-10 (Stratagene Catalog number 200314) competent cells thawed on ice.

[0679] 50 μ L of competent cells were aliquoted from the stock tube per ligation.

[0680] 2 μ L of β -mercaptoethanol stock which is provided with the XL-10 cells.

[0681] This mixture was incubated for ten minutes on ice gently mixing every 2 minutes. Seamless cloning ligation reaction (4 μ L) was added, swirled gently and then incubated on ice for 30 minutes. The tubes were heat shocked for 35 seconds at 42° C. in a water bath. The tubes were incubated on ice for at least two minutes. SOC medium (400 μ L) were added to the cells and incubated for one hour at 37° C. with agitation.

[0682] Two LB agar kanamycin (50 μ g/mL) plates are used to plate 200 μ L and 10 μ L of the transformed cells and allowed to grow overnight.

Screening of Kanamycin Resistant Clones

[0683] Recombinant candidates were grown up for mini-preps in Luria Broth containing Kanamycin (25 μ g/mL) and extracted using the QIAprep Spin Miniprep Kit (Qiagen Valencia, Calif. Catalog Number 27106). Candidate clones were screened by restriction enzymes (New England Biolabs Beverly, Mass.) and positive clones were grown up in 100 mL of Luria Broth containing kanamycin (25 μ g/mL) and extracted using the Qiagen HiSpeed Plasmid Midi Kit (Catalog number 12643). These clones were submitted to GENEWIZ (North Brunswick; NJ) for sequencing.

Production and Purification of STF2.4 $\times$ M2e Fusion Protein

[0684] STF2.4 $\times$ M2e in *E. coli* BLR(DE3)pLysS host (Novagen, San Diego, Calif., Catalog #69053) was retrieved from glycerol stock and scaled up to 5 L. Cells were grown in LB medium containing 15 μ g/ml Kanamycin and 12.5 μ g/ml Tetracycline to OD₆₀₀=0.4 and induced with 1 mM IPTG for 3 h at 37° C. The cells were harvested by centrifugation (7000 rpm $\times$ 7 minutes in a Sorvall RC5C centrifuge) and resuspended in 2 $\times$ PBS, 1% glycerol, DNase, 1 mM PMSF, protease inhibitor cocktail and 1 mg/ml lysozyme. The suspension was passed through a microfluidizer to lyse the cells. The lysate was centrifuged

(45,000 g for one hour in a Beckman Optima L ultracentrifuge) to separate the soluble fraction from inclusion bodies. Protein was detected by SDS-PAGE in the soluble and insoluble fractions.

[0685] The soluble fraction was applied to Sepharose Q resin in the presence of high salt via batch method to reduce DNA, endotoxin, and other contaminants. The flow through containing the protein of interest was loaded onto 30 ml Q Sepharose column (Amersham Biosciences). Bound protein was eluted using a linear gradient from Buffer A to B. (Buffer A: 100 mM Tris-Cl, pH 8.0. Buffer B: 100 mM Tris-Cl, 1 M NaCl, pH 8.0). Eluted protein was further purified using a 45 ml Source Q column that provided greater resolution needed to resolve contaminating proteins. Bound protein was eluted with a linear gradient from Buffer A to B (Buffer A: 100 mM Tris-Cl, pH 8.0 Buffer B: 100 mM Tris-Cl, 1 M NaCl, pH 8.0).

[0686] Final purification of protein was completed using Superdex-200 gel filtration chromatography. The column was developed with 100 mM Tris, 150 mM NaCl and 1% glycerol plus 1% Na-deoxycholate to remove the LPS. Buffer exchange was carried out using overnight dialysis against buffer containing 50 mM Tris, 100 mM NaCl and 1% glycerol was done to remove Na-deoxycholate. Protein concentration was determined by the MicroBCA Protein Assay Reagent Kit (Pierce Biotechnology). Purified preparations of STF2.4×M2e yielded a single band visible with Coomassie stain that migrated with an apparent molecular weight of about 64 kDa on 12% SDS polyacrylamide gels.

Example 16

Expression and Purification of Flagellin (STF2 and STF2Δ) Fusion Protein Constructs Encoding Influenza A M2 Ectodomain Sequences

[0687] The consensus M2e sequences from several influenza A strains of human and avian origin are depicted in SEQ ID NOS: 544-556, 570 and 573-578. To facilitate the cloning of the M2e sequence, two vector cassettes, pMT/STF2 and pMT/STF2Δ, each containing a multiple cloning site (MCS) were generated (See FIGS. 17A and 17B). To generate pMT/STF2, the 1.5 kb gene encoding full length flagellin of *Salmonella typhimurium* fljB type 2, or STF2, was fused to the Ig binding protein (BiP) secretion signal of pMTBiP/V5-His vector (Invitrogen Corporation, Carlsbad, Calif.) for expression in *Drosophila*. The BiP sequence is included at the 5' end of the construct as a secretion signal for expression in *Drosophila*. A chemically-synthesized 4×M2e gene representing the H1, H2 and H3 consensus sequence, SEQ ID NO: 544, was cloned into the MCS of pMT/STF2 to create pMT/STF2.4×M2e(H1).

[0688] A similar strategy prophetically is employed to clone two H5-associated M2e sequences, SLLTEVETP-

TRNEWECRCSDDSDP (SEQ ID NO: 553) (A/Viet Nam/1203/2004) and SLLTEVETLTRNGWGCRCSDSDP (SEQ ID NO: 552) (A/Hong Kong/156/97). Codon-optimized chemically synthesized genes containing four tandemly repeated copies of the indicated H5-associated M2e sequence prophetically are cloned into pMT/STF2 to generate STF2.4×M2e(HSVN) and STF2.4×M2e(H5HK), respectively. To generate a construct that contains multiple M2e forms, the heterologous 4×M2e sequence(s) prophetically are inserted into either of the primary constructs.

[0689] "Heterologous sequences," as used herein, means sequences from different species. For example, the H1 sequence is a human sequence and the H5 sequence is an avian sequence. Thus, the H1 and H5 sequences are heterologous sequences (e.g., SLLTEVETPTRNEWESRSDSDSDPLESLLTEVETPTRNEWESRSDSDSDPESSLLTEVETPTRNEWESRSDSDSDPGSSLLTEVETPTRNEWESRSDSDSDP (SEQ ID NO: 597), encoded by

(SEQ ID NO: 598)
 tctctgctgactgaagtagaaactccaacgcgtaatgaatgggaatcccg
 ttctagcgactcctctgatcctctcgagtcctctgctgacggagggtgaaa
 ccccgaccgcgaacgagtgaggaaagccgttcctccgattcctctgatccg
 gagagcagcctgctgaccgaggtagaaaccccgaccgcgtaatgagtgagg
 atctcgctcctctgattctcttgaccgggacccctctctgctgaccgaag
 tggagactccgactcgcaacgaatgggagagccgttctctgactcctct
 gaccgc.

[0690] Primary constructs comprise at least one pathogen-associated molecular pattern (e.g., STF2, STF2Δ) and at least a portion of at least one integral membrane protein (e.g., M2e, such as SEQ ID NOS: 510 and 544). If there is more than one integral membrane in a primary construct, the integral membrane proteins are from the same species.

[0691] A heterologous construct includes at least two integral membrane proteins such as H1 (human) and H5 (avian), for example, in SEQ ID NOS: 583 and 584.

[0692] To generate pMT/STF2Δ, the hyper-variable region that spans amino acids 170 to 415 of the full-length flagellin gene of SEQ ID NO: 499 was deleted and replaced with a short (10 amino acid) flexible linker (GAPVD-PASPW, SEQ ID NO: 594) designed to facilitate interactions of the amino and carboxy terminal sequences necessary for TLR5 signaling. The protein expressed from this construct retains potent TLR5 activity whether expressed alone or in fusion with test antigen. Thus, a second series of M2e constructs prophetically is generated based on pMT/STF2Δ. *Drosophila* Dmel-2 cells (Invitrogen Corporation, Carlsbad, Calif.) grown at room temperature in Schneider's medium supplemented with 10% FBS and antibiotics prophetically is

transfected with the constructs described above using Cellfectin reagent (Invitrogen) according to the manufacturer's instructions. Twenty-four hours post transfection, cells prophetically is induced with 0.5 mM CuSO₄ in medium lacking FBS and incubated for an additional 48 hours. Conditioned media (CM) prophetically is harvested from induced cultures and screened for protein expression by SDS-PAGE and Western blot analyses using anti-flagellin and anti-M2e specific antibodies. The identity, TLR bioactivity of the fusion protein, antigenicity assessed by ELISA and in vivo mouse studies for immunogenicity prophetically is performed.

Example 17

Construction and Expression of Flagellin-Hemagglutinin (HA) Constructs

[0693] The gene encoding HA from genomic DNA from the in-house laboratory strain PR8, an attenuated derivative of A/Puerto Rico/8/34 was isolated (SEQ ID NO: 565, encoding SEQ ID NO: 564). The gene was fused to the STF2A cassette that has been previously constructed in pPICZA generating STF2A.HAPR8 (SEQ ID NO: 560, encoding SEQ ID NO: 559) (See FIG. 42). Purified recombinant protein was tested for immunogenicity and efficacy in BALB/c mice. The gene encoding H5N1 of the A/Vietnam/1203/04 strain was custom synthesized and fused to STF2A cassette generating STF2A.HAH5 (SEQ ID NO: 558, encoding SEQ ID NO: 557). Both human and avian HA constructs were transformed into *Pichia pastoris* strains GS115 and X-33 (Invitrogen Corporation, Carlsbad, Calif.). Selected clones were screened for expression by fractionation on SDS-PAGE gel and staining by Coomassie Blue and Western blot analysis using anti-HA and anti-flagellin antibodies.

Example 18

Generation of a Pam3Cys Fusion Protein

[0694] M2e (SEQ ID NO: 544) was chemically coupled to a tri-palmitoylcysteine (Pam3Cys) moiety through the amino terminal serine residue of the peptide. The structure of the fusion protein (Pam3Cys.M2e) is shown in FIG. 39. The chemical name for Pam3Cys.M2e is [Palmitoyl-Cys((RS)-2,3-di(palmitoyloxy)-propyl)-Ser-Leu-Leu-Thr-Glu-Val-Glu-Thr-Pro-Ile-Arg-Asn-Glu-Trp-Gly-Ser-Arg-Ser-Asn-Asp-Ser-Ser-Asp-Pro-OH acetate salt]. The molecular mass of Pam3Cys.M2e is 3582.3 daltons.

[0695] Pam3Cys.M2e was synthesized using a solid phase peptide synthesis methodology based on a well established Fmoc-strategy (Houben-Weyl, 2004. Synthesis of peptides and peptidomimetics, Vol. 22, Georg Thieme Verlag Stuttgart, N.Y.). The synthetic scheme and manufacturing process for Pam3Cys.M2e is diagrammed in the flow chart

below. The Pam3Cys.M2e is a fusion protein (chemically linked) and is also referred to herein as a "lipidated peptide."

[0696] The first step in the synthesis included solid phase peptide synthesis. The amino acid sequence of Pam3Cys.M2e was assembled on an H-Pro-2-chlorotriethyl chloride resin by solid phase peptide synthesis. This resin is highly suitable for the formation of peptides with the Fmoc-strategy. The peptide chain was elongated by successive coupling of the amino acid derivatives. Each coupling step was preceded by an Fmoc-deprotection step and both steps were accompanied by repeated washing of the resin. After coupling of the last amino acid derivative, the final Fmoc-deprotection step was performed. Finally, the peptide resin washed and dried under reduced pressure. During solid phase peptide synthesis color indicator tests were performed for each step to monitor the completion of the Fmoc-cleavage and the subsequent coupling of the amino acid derivatives.


[0697] Stage 2 of the synthesis included coupling of Pam3Cys-OH. Pam3Cys-OH was pre-activated with N,N'-dicyclohexyl-carbodiimide (DCCI) in the presence of 1-hydroxybenzotriazole (HOBt). The resulting solution was filtered and added to the peptide resin. At the end of the reaction time the peptide resin washed and dried under reduced pressure. Color indicator tests were performed to control the coupling of Pam3Cys-OH.

[0698] Stage 3 of the synthesis included cleavage from the resin including cleavage of the side chain protecting groups. The peptide resin was treated with trifluoroacetic acid (TFA). The product was precipitated from the reaction mixture and lyophilized.

[0699] Stage 4 of the synthesis included purification by preparative reverse phase HPLC. The crude material obtained from Stage 3 was purified by preparative HPLC on a reverse phase column using a TFA system. The fractions were collected, checked by analytical HPLC and pooled accordingly. Pooled fractions from the TFA runs were lyophilized.

[0700] Stage 5 of the synthesis included precipitation in the presence of EDTA. The purified material from Stage 4 was precipitated from an aqueous solution of EDTA. The product was filtered off and dried under reduced pressure.

[0701] Stage 6 of the synthesis included ion exchange chromatography. The last stage of manufacturing Pam3Cys.M2e was the exchange from the trifluoroacetate salt into the acetate salt by ion exchange. The material from Stage 5 was loaded onto an ion exchange column and eluted with acetic acid. Fractions were checked by thin layer chromatography and the combined product-containing fractions were filtered and lyophilized to yield the final product.



Pam3-Cys-OH = Palmitoyl-Cys(RS)-2,3-di(palmitoyloxy)-propyl-OH

[0702] The purity specification for the Pam3Cys.M2e drug substance was >80% by RP-HPLC. The specification was based on the purity achieved with three non-GMP lots of Pam3Cys.M2e made from the same GMP batch of M2e-peptide intermediate resin. The purity of the three non-GMP lots of Pam3Cys.M2e was 80.2%, 80.3% and 80.8%, for lots D.001.Pam3Cys.M2e, D.002.Pam3Cys.M2e and D.003.Pam3Cys.M2e, respectively.

Example 19

Immunogenicity

Materials and methods

Synthesis and Purification of PAM3Cys.M2e

[0703] Pam3Cys.M2e was prepared by Genemed Synthesis and Bachem using solid phase synthesis methodologies

and Fmoc chemistry as described above. Mass spectroscopy analysis was used to verify the molecular weight of the final product.

Endotoxin Assay

[0704] Endotoxin levels of the STF2.4×M2e and the Pam3Cys.M2e were measured using the QCL-1000 Quantitative Chromogenic LAL test kit (BioWhittaker #50-648U), following the manufacturer's instructions for the microplate method.

TLR5 Bioactivity Assay

[0705] HEK293 cells constitutively express TLR5 and secrete several soluble factors, including IL-8, in response to TLR5 signaling. HEK293 cells were seeded in 96-well microplates (50,000 cells/well) and test proteins were added and incubated overnight. The next day, the conditioned

medium was harvested, transferred to a clean 96-well microplate and frozen at -20°C . After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce, #M801E and #M802B) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer (FARCyte, Amersham). Results are reported as pg of IL8 per ml as determined by inclusion of a standard curve for IL8 in the assay.

TLR2 Bioactivity Assay

[0706] RAW264.7 cells (ATCC) express TLR2 and secrete several soluble factors, including $\text{TNF}\alpha$, in response to TLR2 signaling. RAW264.7 cells were seeded in 96-well microplates (50,000 cells/well), test compounds were added and incubated overnight. The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate and frozen at -20°C . After thawing, the conditioned medium was assayed for the presence of $\text{TNF}\alpha$ in a sandwich ELISA using an anti-mouse $\text{TNF}\alpha$ matched antibody pair (Pierce) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer (FARCyte, Amersham). Results are reported as ng of TNF per ml as determined by reference to a standard curve for TNF included in the assay.

Mouse Immunogenicity

[0707] Female BALB/c mice (National Cancer Institute) were used at the age of about 6-8 weeks. Mice were divided into groups of 5 to 10 mice per group, and immunized subcutaneously on each side of the base of the tail on days 0 and 21 with the indicated concentrations of STF2.4xM2e or Pam3Cys.M2e fusion protein. On days 10 (primary) and 28 (boost), individual mice were bled by retro-orbital puncture. Sera were harvested by clotting and centrifugation of the heparin-free blood samples.

Mouse Serum Antibody Determination

[0708] M2e-specific IgG levels were determined by ELISA. 96-well ELISA plates were coated overnight at 4°C with 100 μl /well of a 5 $\mu\text{g}/\text{ml}$ solution of the M2e peptide in PBS. Plates were blocked with 200 μl /well of Assay Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature. The plates were washed three times in PBS containing 0.05% Tween-20 (PBS-T). Dilutions of the sera in ADB were added (100 μl /well) and the plates were incubated overnight at 4°C . The plates were washed three times with PBS-T. Horse radish peroxidase, or HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical) diluted in ADB were added (100 μl /well) and the plates were incubated at room temperature for 1 hour. The plates were washed three times with PBS-T. After adding TMB Ultra substrate (3,3',5,5'-tetramethylbenzidine; Pierce) and monitoring color development, the O.D. 450 was measured on a Tecan Farcyte microspectrophotometer.

Rabbit Immunogenicity

[0709] Female and male NZW rabbits (Covance Research Products) were used at the age of about 13-17 weeks. Rabbits were divided into groups of 3 male and 3 female per group, and immunized i.m. on alternating thighs on days 0 and 21 and 42 with the indicated concentrations of Pam3Cys.M2e peptide or STF2.4xM2e fusion protein. Ani-

mals were bled on day -1 (prebleed), 14 (primary) and 28 and 42 (boost). Sera were prepared by clotting and centrifugation of samples.

Rabbit Serum Antibody Determination

[0710] M2e-specific IgG levels were determined by ELISA. 96-well ELISA plates were coated overnight at about 4°C with 100 μl /well M2e peptide in PBS (5 $\mu\text{g}/\text{ml}$). Plates were blocked with 200 μl /well of Assay Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature. The plates were washed three times in PBS-T. Dilutions of the sera in ADB were added (100 μl /well) and the plates were incubated overnight at about 4°C . The plates were washed 3x with PBS-T. Bound IgG was detected using HRP-conjugated goat anti-rabbit IgG (Jackson Immunochemical). The plates were washed three times with PBS-T. After adding TMB Ultra substrate (Pierce) and monitoring color development, O.D. 450 was measured on a Molecular Devices Spectramax microspectrophotometer. Results are reported as the Delta O.D. which is determined by subtracting the O.D. 450 reading for the prebleed of each animal from the O.D. 450 for each animal post-immunization.

BALB/C Mouse Efficacy Model

[0711] In a typical experiment, about 5-6 week old female BALB/c mice (10-20 per group) were obtained and allowed to acclimate for one week. Fusion proteins formulated in PBS or other suitable formulation were administered by s.c. injection. Mice were immunized on days 0 and 14. On day 21, sera was harvested by retro-orbital puncture and evaluated for M2e specific IgG by ELISA. Mice were challenged by intranasal administration of 1xLD₅₀ of the well characterized mouse adapted Influenza A strain, A/Puerto Rico/8/34 (H1N1). Mice were monitored daily for 14 days for survival and weight loss. Mice that lost about 30% of their initial body weight were humanely sacrificed, and the day of sacrifice recorded as the day of death. Efficacy data were reported as survival times.

Results

In Vitro Bioactivity

[0712] These assays were based on cell lines expressing the relevant TLR and screened for the ability to produce either IL8 or $\text{TNF}\alpha$ in response to TLR triggering. In FIG. 52, the ability of STF2.4xM2e (■) or STF2.OVA (○) to stimulate TLR5 dependent IL8 production was evaluated following the stimulation of TLR5 positive, HEK293 cells. The results indicate that both fusion proteins stimulated IL8 production in a dose dependent manner and that the activity of the PAMP was retained in the context of the fusion.

[0713] TLR2 activity was similarly evaluated for Pam3Cys.M2e following stimulation of TLR2 positive RAW264.7 cells. In FIG. 51, the experimental groups are: the known endotoxin, LPS, as a positive control (◆), LPS plus the inhibitor of endotoxin polymyxin B (PMB) as a negative control (○), free Pam3Cys as a positive control for TLR2 signalling (■), free Pam3Cys plus PMB (□), Pam3Cys.M2e (◆) and Pam3Cys.M2e plus PMB (◇). The results showed similar activity profiles for Pam3Cys.M2e and the free TLR2 ligand Pam3Cys. The addition of polymyxin B (PMB) did not reduce its activity, indicating that there is no or low endotoxin contamination.

Physical Linkage of PAMP and Antigen Enhances Immunogenicity

[0714] Using mouse models of immunogenicity, chemical coupling of Pam3Cys to M2e enhances the immunogenicity of the M2e antigen as compared to either the M2e peptide delivered alone or the M2e peptide co-delivered with free Pam3Cys. In the experiment shown in FIG. 52, groups of mice were immunized on days 0 and 21 with PBS as a negative control (*), the free TLR2 ligand, Pam3CSK-4 ((), M2e peptide alone (○), free Pam3CSK-4 mixed with M2e peptide (□), or the fusion of Pam3Cys and M2e referred to as Pam3.M2e (◆). The relevant the molar ratio of M2e peptide delivered was held constant. On day 28, sera were harvested and analyzed for M2e-specific antibody titers by ELISA. The results show that chemical coupling of Pam3Cys to the M2e (Pam3Cys.M2e) generates a detectable serum antibody response to the M2e antigen.

[0715] Physical linkage between the TLR5 ligand STF2 and antigen was demonstrated using the model antigen ovalbumin (OVA). Mice received a single s.c. immunization with STF2, OVA, STF2.OVA fusion protein, STF2+OVA mixture or PBS alone. Dosages were calculated to deliver 12 µg equivalents of STF2 and OVA per group. Seven days later, sera were harvested and OVA-specific antibodies were examined by ELISA. Data shown in FIG. 53 depict IgG1 titers at a 1:100 dilution of the sera. These results demonstrate that physical linkage of the TLR5 ligand and antigen results in optimal immunogenicity in vivo.

PAMP Linked Antigens are More Immunogenic than Conventional Adjuvant

[0716] Groups of 5 BALB/c mice were immunized on day 0 and 14 with 30 µg of Pam3Cys.M2e (◆), 22.5 µg of M2e which is the molar equivalent of M2e in 30 µg of Pam3Cys.M2e (◇), 22.5 mg of M2e adsorbed to the conventional adjuvant Alum (□), or 25 mg of the recombinant protein STF2.4×M2e (■). A group receiving PBS was included as a negative control (○). Sera were harvested 7 days post the second dose and M2e specific IgG were evaluated by ELISA. The results shown in FIG. 54 indicate that M2e alone is poorly immunogenic in that it failed to elicit antibody titers above background. The conventional adjuvant Alum provided a modest enhancement in the immune response to M2e. The PAMP linked M2e constructs; however, provided the greatest enhancement in immunogenicity. These results indicate direct linkage of PAMPs with portions of an integral membrane protein of an influenza viral protein can elicit immune responses that are more potent than those elicited by the conventional adjuvant Alum.

Dose and Immunogenicity

[0717] Dose ranging studies were carried out to further assess the potency of Pam3Cys.M2e and STF2.4×M2e. For STF2.4×M2e, BALB/c mice were immunized on day 0 and 14 with dilutions of STF2.4×M2e that ranged from 0.25 to 25 µg of STF2.4×M2e per immunization. The prefix D002 refers to the specific batch of STF2.4×M2e used in this experiment, while R-028 refers to a historical reference batch of STF2.4×M2e used in this experiment. Seven days following the last immunization (Day 21) mice were bled and M2e-specific IgG responses were evaluated by ELISA. The results shown in FIG. 55 demonstrate that immunization

with doses as low as 0.25 µg per immunization of STF2.4×M2e induced detectable levels of M2e-specific IgG, with the optimal dose in mice falling in the range of about 2.5 to about 25 µg.

[0718] For Pam3Cys.M2e, BALB/c mice were immunized on day 0 and 14 with 0.05 to 30 µg of Pam3Cys.M2e per immunization. Seven days following the last immunization (Day 21) mice were bled and M2e-specific IgG responses were evaluated by ELISA. The results shown in FIG. 56 demonstrate that immunization with concentrations as low as 0.05 µg of Pam3Cys.M2e induced detectable levels of M2e-specific IgG, with the optimal dose for mice in this study of about 30 µg.

Immunogenicity In Multiple Mouse Strains

[0719] The immunogenicity of Pam3Cys.M2e was evaluated in multiple mouse strains including BALB/c (●), C57BL/6 (■), CB6/F1 (◆), DBA/2 (▲), Cr:NIH (Swiss) (X) and C3H/HeN (*). Groups of five for each strain were immunized on day 0 and 14 with 30 µg of Pam3Cys.M2e per immunization. Sera were harvested on day 21 and levels of M2e-specific IgG evaluated by ELISA. All strains exhibited significant levels of M2e-specific IgG indicating that the immunogenicity of Pam3Cys.M2e is not dependent on a particular MHC (FIG. 57).

Immunogenicity in Rabbits

[0720] Studies aimed at evaluating the immunogenicity of Pam3Cys.M2e and STF2.4×M2e in a second species, rabbit, were carried out. In the first study, rabbits (3 females and 3 males/group) were immunized with 500, 150, 50, 15 or 5 µg (i.m.) of Pam3Cys.M2e on day 0, 21 and 42. As a control, an additional group received the formulation buffer F111 (10 mM Tris, 10 mM histidine, 75 mM NaCl, 5% sucrose, 0.02% Polysorbate-80, 0.1 mM EDTA, 0.5% ethanol, 20 mg/mL hydroxypropyl-beta-cyclodextrin, pH 7.2). On day 7 post-boost 2, peripheral blood was obtained and the anti-M2e antibody titers were evaluated by ELISA. The results shown in FIG. 58 depict the individual rabbit antibody titers at a 1:125 dilution of the sera. The data suggest a dose-response relationship between the amount of Pam3Cys.M2e used for prime/boost vaccinations and the level of the antibody titer achieved.

[0721] In the second study, rabbits (3 females and 3 males/group) were immunized with 500, 150, 50, 15 or 5 µg (i.m.) of STF2.4×M2e. As a control, an additional group received saline alone. On day 14 post-immunization, peripheral blood was obtained and the anti-M2e antibody titers were evaluated by ELISA. Notably, significant M2e-specific IgG responses were detectable by day 14 post-prime in all animals immunized (FIG. 59). The results indicate that STF2.4×M2e elicits a rapid and consistent immune response in rabbits.

Efficacy in the Mouse Challenge Model

[0722] The efficacy of the Pam3Cys.M2e and STF2.4×M2e was evaluated in BALB/c mice using the well characterized mouse adapted strain, Influenza A/Puerto Rico/8/34 (PR/8) as the challenge virus. Groups of ten mice were immunized s.c. on day 0 and 14 with 30 µg of Pam3Cys.M2e in the formulation buffer F111 (■), 30 µg of Pam3Cys.M2e in the proprietary buffer F120 (10 mM Tris, 10 mM histidine, 10% sucrose, 0.02% Polysorbate-80, 0.1 mM EDTA,

0.5% ethanol, 0.075% docusate sodium, pH 7.2) ($\blacktriangle$), 30 μ g of Pam3Cys.M2e in the buffer F119 (10 mM Tris, 10 mM histidine, 75 mM NaCl, 5% sucrose, 0.02% Polysorbate-80, 0.1 mM EDTA, 0.5% ethanol, 0.1% docusate sodium, pH 7.2), 30 μ g of STF2.4xM2e in the buffer F105 (10 mM Tris, 10 mM histidine, 75 mM NaCl, 5% sucrose, 0.02% Polysorbate-80, 0.1 mM EDTA, 0.5% ethanol, pH 7.2), 3 μ g of STF2.4xM2e in buffer F105 (10 mM Tris, 10 mM histidine, 75 mM NaCl, 5% sucrose, 0.02% Polysorbate-80, 0.1 mM EDTA, 0.5% ethanol, pH 7.2) ($\bullet$) or 0.3 μ g of STF2.4xM2e in buffer F105 ($\square$). A group receiving PBS alone was included as a negative control ($\circ$), and a convalescent group with immunity to PR/8 following a sublethal challenge with the virus was included as a positive control ($\diamond$). On day 28, animals were challenge with an LD90 of the PR/8 challenge stock. Weight loss and survival was followed for 14 days post challenge (FIG. 60).

[0723] Animals in the convalescent group which had successfully cleared an earlier non-lethal infection with PR/8 demonstrated 100% protection to a subsequent viral challenge. Animals receiving the PBS buffer alone exhibited morbidity beginning on days 7 and 8, with 80% lethality occurring by day 10, while animals immunized with 30 μ g of Pam3Cys.M2e in F111 demonstrated enhanced survival, with 50% of mice surviving the challenge. Animals receiving Pam3Cys.M2e in F119 exhibited morbidity beginning on days 8 and 9 with 80% of the mice surviving. Animals receiving Pam3Cys.M2e in buffer F120 (10 mM Tris, 10 mM histidine, 10% sucrose, 0.02% Polysorbate-80, 0.1 mM EDTA, 0.5% ethanol, 0.075% docusate sodium, pH 7.2) or the STF2.4xM2e protein exhibited the mildest disease course with 90 to 100% of the mice in these groups surviving the lethal challenge. These results demonstrate that both Pam3Cys.M2e and STF2.4xM2e can confer protective immunity to a challenge with influenza A in vivo.

Discussion

[0724] *Salmonella typhimurium* flagellin (fljB) is a ligand for TLR5. A recombinant protein consisting of full-length fljB (STF2) fused to four tandem repeats of M2e was expressed in *E. coli* and purified to >95% purity with low endotoxin levels. In reporter cell lines, this protein (STF2.4xM2e) triggered IL8 production in a TLR5-dependent fashion. Mice immunized with dilutions of STF2.4xM2e that ranged from 0.25 μ g to 25 μ g, formulated in the buffer F105 which is without a conventional adjuvant or carrier, mounted a vigorous antibody response. The potency of the recombinant protein was further demonstrated in rabbit immunogenicity studies where animals receiving as little as 5 μ g of protein seroconverted after a single dose. The efficacy of the PAMP fusion protein was demonstrated in the mouse challenge model using Influenza A/Puerto Rico/8/34 as the challenge virus. Mice immunized with as little as about 0.3 μ g of the protein per dose exhibited mild morbidity with 100% of the mice surviving the challenge.

[0725] Synthetic tripalmitoylated peptides mimic the acylated amino terminus of lipidated bacterial proteins and are potent activators of TLR2. In these studies, a tripalmitoylated peptide consisting of three fatty acid chains linked to a cysteine residue and the amino terminus of the Influenza A M2 ectodomain (M2e) was synthesized using standard solid-phase peptide chemistries. This peptide (Pam3Cys.M2e) triggered TNF α production in a TLR2-

dependent fashion in reporter cell lines. When used to immunize mice without adjuvant, Pam3Cys.M2e generated an antibody response that was more potent than M2e when mixed with free Pam3CSK-4. Pam3Cys.M2e was also found to be immunogenic in rabbits where a dose response relationship was observed between the amount of Pam3Cys.M2e used for immunization and the antibody titer achieved. The efficacy of the Pam3Cys.M2e peptide in a number of different formulations was evaluated in the mouse challenge model using Influenza A/Puerto Rico/8/34 as the challenge virus. Pam3Cys.M2e formulated in F119 and F120 exhibited the mildest morbidity with about 80 to about 100% of the mice surviving the challenge.

Example 20

Materials and Methods

PCR Amplification and DNA Primers

[0726] All PCR amplifications were performed using Pfu Ultra Hotstart PCR Master Mix (Catalog number 600630) from Stratagene (La Jolla, Calif.) according to the manufacturer's recommendations. DNA primers were purchased from Sigma Genosys and are described below.

STF28BGF-1: (SEQ ID NO: 644)
CTCGGGAGATCTGCACAAGTAATCAACACTAACAGTCT

STF28MCR-1: (SEQ ID NO: 645)
CCATGGGCTAGCAGGATCCACGGCGCTCCCTGCACGTTCA

STF28MCF-2: (SEQ ID NO: 646)
GGAGCGCCGGTGGATCCTGCTAGCCCATGGACCGAAAACCCG

STF28ECR-2: (SEQ ID NO: 647)
TCTGCAGAATTCACGTAACAGAGACAGCAGCTTCTCGGGACGTCCTCCGCA
GAACGTGCTGTCTCTGTTACGTGAATTCCTGCAGA

pET24AR:5 (SEQ ID NO: 648)
TCCGGCGTAGAGGATCGAGA

STF2-E3R3: (SEQ ID NO: 649)
CAATTGACCTTCAAGCTTCGAATTGCCCTTACGTAAACAGAGACAGCAGCT
TCTG

AX-E3F3: (SEQ ID NO: 650)
AAGCTTGAAGTCAATTGGAATTCCTAGGACTAGTATGGAAAAATTGCA
GTTGAAG

pET24AF: (SEQ ID NO: 651)
GCTTAATGCGCCGCTACAGG

5'WNE28: (SEQ ID NO: 652)
GCGGCCGCTCATGGAAAAATTGCAGTTGAAGGGAACAACC

3'WNE28: (SEQ ID NO: 653)
CCGCGGTTTGCCAATGCTGCTTCCAGACTTGT

NdeI-STF2: (SEQ ID NO: 654)
CCGGCATGCCATATGGCACAAGTAATCAACACTAACAGTCTGTCGCTGC

-continued

BlpI-EdIII:
 (SEQ ID NO: 656)
 GCATGCTCAGCTTATTAAGGGTTTGCCAATGCTGCTTCCCAGACTTGTG
 JE EIII primer:
 (SEQ ID NO: 53)
 TACGTGAATTACAGCAGATATCCAGCAC

Cloning of pET/STF2Δ.EIII

[0727] Full length flagellin of *Salmonella typhimurium* fljB (flagellin phase 2) (also referred to herein as "STF2") is encoded by a 1.5 kb gene. A truncated version of the STF2 (STF2Δ, SEQ ID NO: 606, encoded by SEQ ID NO: 607) was generated by deleting the hyper-variable region that spans amino acids 170 to 415 of SEQ ID NO: 604. The deleted region was replaced with a short flexible linker (GAPVDPASPW, SEQ ID NO: 659) designed to facilitate interactions of the NH2 and COOH termini sequences necessary for TLR5 signaling. To generate this construct, a two-step PCR was used. In the first reaction, STF2.OVA (SEQ ID NO: 755 encoding amino acid sequence SEQ ID NO: 756) served as the DNA template and STF28BGF-1 and STF28MCR-1 were used as primer pairs. In a separate reaction, the same DNA template was combined with primers STF28MCF-2 and STF28ECR-2.

[0728] The PCR amplification reactions generated about 500 bp and about 270 bp fragments, respectively. These PCR products were combined in a final PCR reaction using STF28BGF-1 and STF28ECR-2 as primers. The amplified DNA product from this reaction (about 0.77 kb) was digested with BglII and EcoRI restriction enzymes and ligated into pMTBip/V5-His B (Invitrogen, Carlsbad, Calif.) that had previously been digested with BglII and EcoRI and treated with alkaline phosphatase. An aliquot of the ligation mix was used to transform TOP10 cells (Invitrogen, Carlsbad, Calif.). PCR screening was performed using vector specific primers, pMTFOR (methionine promoter) (CATCTCAGTGCAACTAAA, SEQ ID NO: 759) and BGHREV (bovine growth hormone poly A) (TAGAAGGCACAGTCGAGG, SEQ ID NO: 760), to identify several positive clones. All positive clones were further analyzed by restriction mapping analysis and confirmed by DNA sequencing. The resultant construct pMT/STF2Δ was used to generate pMT/STF2Δ.EIII+.

[0729] The domain III of the West Nile virus envelope protein of pET/STF2Δ.EIII+ (SEQ ID NOS: 673, 674) was derived from the *Drosophila* expression plasmid pMT/STF2.E. This plasmid contains full-length STF2 (amino acids 1-506, SEQ ID NO: 604) fused to the West Nile Virus envelope protein (amino acids 1-406, SEQ ID NO: 642). The pMT/STF2.E (SEQ ID NO: 761) clone AX-1 was used as a DNA template and 5'WNE28 (SEQ ID NO: 652) and 3'WNE28 (SEQ ID NO: 653) served as primers for PCR amplification. In order to facilitate restriction analysis and subsequent cloning steps, the 5' primer encoded a novel NotI site (New England Biolabs, Beverly, Mass.) and the 3' primer contained a unique SacII site. The amplified EIII+ DNA fragment (345 bp; SEQ ID NO: 781 that encodes amino acids 292-406 of SEQ ID NO: 642) was subcloned into pCR-Blunt II-TOPO cloning vector (Invitrogen, Carlsbad, Calif.) to generate plasmid TOPOEIII. A stop codon

was subsequently introduced downstream of the EIII+ sequence by blunting the SacII and SpeI restriction sites using T4 DNA polymerase.

[0730] To generate pMT/STF2Δ.EIII+ (SEQ ID NOS: 673, 674), the EIII+ fragment was isolated from TOPOEIII+ using NotI and BamHI restriction sites and ligated into the NotI and SacII restriction sites in pMT/STF2Δ. The BamHI site of the EIII+DNA fragment and the SacII site of pMT-STF2Δ were blunted with T4 DNA polymerase prior to ligation. The STF2Δ.EIII+ sequence (SEQ ID NOS: 673, 674) from pMT/STF2Δ.EII+ was isolated by PCR amplification using the primers NdeI-STF2 and BlpI-EdIII. To generate pET/STF2Δ.EIII+ (SEQ ID NO: 674), the PCR product was digested with NdeI and BlpI and ligated into pET24a plasmid that had been predigested with NdeI and BlpI. The ligation mix was transformed into Mach-1 cells (Invitrogen, Carlsbad, Calif.) and the cells were grown on LB supplemented with 50 μg/ml kanamycin. Several colonies were screened by restriction mapping and were verified by DNA sequencing.

Cloning of pET/STF2.EIII+

[0731] The West Nile virus EIII+ sequence of pET/STF2.EIII+ (SEQ ID NOS: 657, 658) was derived from pETSTF2.E (SEQ ID NOS: 761, 762). This *E. coli* expression plasmid contains full-length STF2 (amino acids 1-506) fused to the West Nile Virus envelope protein (amino acids 292-406 of SEQ ID NO: 642, which is SEQ ID NO: 610). In two independent PCR reactions, pET/STF2.E was used as the DNA template. One reaction used the primers pET24AR:5 (SEQ ID NO: 648) and STF2-E3R3: (SEQ ID NO: 649) and the other used AX-E3F3 (SEQ ID NO: 650) and pET24AF (SEQ ID NO: 651). These PCR reactions generated a 1.5 kb fragment that consisted of full-length STF2 and a 340 bp fragment that comprised the EIII domain plus additional amino acids that extended into domain I of the envelope protein. Aliquots of these PCR amplification reactions were combined, and the two products served as templates for a PCR reaction with the external primers pET24AR (SEQ ID NO: 648) and pET24AF (SEQ ID NO: 651). This resulted in the generation of about a 1.8 kb DNA fragment that fused EIII+ sequence (SEQ ID NO: 781, a nucleic acid sequence encoding amino acids 292-406 of SEQ ID NO: 642, which is SEQ ID NO: 610) to STF2. The PCR product was digested with NdeI and BlpI and gel purified and ligated by compatible ends to a pET24a vector that had previously been digested with compatible enzymes and de-phosphorylated. The ligation mix was transformed into Mach-1 cells (Invitrogen, Carlsbad, Calif.) as described for pET/STF2Δ.EIII+. Several colonies were screened by restriction mapping and two clones were verified by DNA sequencing.

Cloning of pET/STF2Δ.JEIII+

[0732] A portion of the envelope protein of a *Japanese encephalitis virus* (JEV) (strain SA-14-14-2 (Jai, L., et al., *Chin Med J (Eng)* 116:941-943 (2003)); currently employed in a JEV vaccine encoded by domain III was custom synthesized by DNA 2. Inc (Menlo Park, Calif.). The portion of domain III was excised from the pJ2:G01510 using NotI and Blp I site that flank the insert. The DNA insert was gel isolated and cloned by compatible ends to pET24A/STF2Δ.EIII+ (SEQ ID NOS: 673, 674) that had previously been digested with the appropriate enzymes to release the

West Nile virus EIII+ insert. The deleted vector was then gel purified and ligated to an aliquot of JE EIII+. The ligation mix was used to transform TOP-10 cells (Invitrogen, Carlsbad, Calif.) and the cells were grown on LB supplemented with 50 µg/ml kanamycin. Several colonies were screened by restriction mapping and were verified by DNA sequencing.

[0733] The resulting construct, pET24A/STF2Δ.EIII (SEQ ID NOS: 608, 609) was BLR (DE3) strain (Novagen) and expression was monitored in several clones using Coomassie Blue staining which was confirmed by Western blot using anti-flagellin antibodies. Using, pET24A/STF2Δ.EIII+ as the DNA template and the JE EIII+ oligonucleotide as primer (SEQ ID NO: 656) the cysteine residue in the linker region between STF2Δ and JEIII+ was changed to a serine residue using QuikChange Site Directed Mutagenesis Kit (Stratagene, LaJolla, Calif.) according to the manufacturer's instructions. The clone was verified by sequencing and assayed for expression as described for pET24A/STF2Δ.EIII+ above.

[0734] When a cysteine residue in a linker in change to a serine residue the fusion protein in also referred to herein by inclusion of an "s" in the designation of the fusion protein. For example, "STF2Δ.EIII+" includes a cysteine residue in the linker (SEQ ID NO: 674), whereas "STF2Δ.EIII+s" include a serine residue substituted for the cysteine residue in the linker (SEQ ID NO: 675).

Cloning the EIII Domain of Each Dengue Virus Fused to the C-Terminal End of Flagellin (STF2Δ)

[0735] Initially, obtaining biologically active material from the fusion of the entire envelope protein of West Nile virus was difficult, perhaps due to the presence of multiple cysteines residues (12 cysteines) in the envelope protein (see SEQ ID NO: 642). However, when the region encoding domain III (EIII) of the protein was sub-cloned, the fusion protein was abundantly expressed in *E. coli* and was highly efficacious in mice. Although there is an overall sequence dissimilarity among the 4 distinct DEN viruses (Den1, Den2, Den3, Den4, SEQ ID NOS: 763-770, the three-dimensional structures within domain III of the envelope protein are similar among the flaviviruses. This domain in DEN and other flaviviruses encodes the majority of the type-specific contiguous critical/dominant neutralizing epitopes. Domain III of the dengue viruses (Den1, Den2, Den3 and Den4) has been expressed, in bacteria and shown to be immunogenic, capable of inducing neutralizing antibodies in experimental animals (Simmons, M., et al., *Am. J. Trop. Med Hyg* 65:159 (2001)). Domain III corresponding to residues about 295 to about 399 (exact numbering depends on the particular DEN virus, for example, of SEQ ID NOS: 763, 765, 767, 769) of the four different DEN viruses have been codon-optimized for expression in *E. coli*. The synthetic gene was amplified by using PCR and sub-cloned into the NotI site of the vector pET/STF2Δ generating pET/STF2Δ.DEN1EIII, pET/STF2Δ.DEN2EIII, pET/STF2Δ.DEN3EIII and pET/STF2Δ.DEN4EIII (SEQ ID NOS: 683, 685, 687 and 689).

E. coli Production of STF2.EIII+, STF2Δ.EIII+, STF2Δ.EIII+s and STF2Δ.EIII+

[0736] Cell cultures (6 L) of BLR(DE3) pLysS that harbor pETSTF2.EIII+ (SEQ ID NOS: 657, 658), pETSTF2Δ.EIII+

(SEQ ID NOS: 673, 674), pETSTF2Δ.EIII+s (SEQ ID NOS: 675, 676) or pETSTF2Δ.EIII+ (SEQ ID NOS: 608, 609) were grown in LB medium containing 15 µg/ml kanamycin, 12.5 µg/ml tetracycline and 24 µg/ml chloramphenicol. At an OD₆₀₀ of about 0.6 protein expression was induced with 1 mM IPTG for about 3 h at about 37° C. Following induction, cells were harvested by centrifugation (7000 rpm×7 minutes in a Sorvall RC5C centrifuge) and resuspended in 2×PBS, 1% glycerol, DNase, 1 mM PMSF, protease inhibitor cocktail and 1 mg/ml lysozyme. The suspension was passed through a microfluidizer to lyse the cells and the lysate was centrifuged (45,000 g for one hour in a Beckman Optima L ultracentrifuge) to separate the soluble fraction from inclusion bodies. Under these growth and induction conditions, STF2.EIII+ was expressed as a soluble protein and STF2Δ.EIII+ (SEQ ID NOS: 673, 674), STF2Δ.EIII+s (SEQ ID NOS: 675, 676) and STF2Δ.EIII+ (SEQ ID NOS: 608, 609) formed inclusion bodies.

Purification of STF2.EIII+

[0737] Cell lysate containing soluble STF2.EIII+ (SEQ ID NOS: 657, 658) was applied to Sepharose Q resin (Amersham Biosciences, Piscataway, N.J.) in the presence of 0.5 M NaCl to reduce DNA, endotoxin, and other contaminants. The flow-through fraction was collected and the conductivity adjusted by a 10-fold dilution with buffer A (Buffer A: 100 mM Tris-Cl, pH 8.0). The diluted material was re-loaded onto Q Sepharose and bound protein was eluted with a linear gradient from 20% to 60% Buffer B (Buffer B: 100 mM Tris-Cl, 1 M NaCl, pH 8.0). Fractions containing STF2.EIII+ were pooled and further processed by Superdex-200 gel (SD200) filtration chromatography in the presence of Na-deoxycholate to remove residual endotoxin (running buffer: 1% Na-deoxycholate, 100 mM NaCl, 100 mM Tris-HCl, 1% glycerol, pH 8.0). Following SD200 chromatography, the eluted protein was loaded directly onto Q Sepharose and washed extensively with buffer A to remove detergent. Bound protein was again eluted with a linear gradient from 20% to 60% Buffer B. In one preparation (Batch 057), this step was substituted with a detergent removal procedure using Extract-D detergent removal gel (Pierce Biotechnology, Rockford, Ill.). The purified protein was dialyzed against buffer containing 50 mM Tris, 100 mM NaCl and 1% glycerol and stored at -80° C.

Purification of STF2Δ.EIII+

[0738] STF2Δ.EIII+ inclusion bodies were collected by low-speed centrifugation (7000 rpm×7 minutes in a Sorvall RC5C centrifuge) and solubilized with buffer containing 8 M urea, 100 mM Tris-HCl, 5 mM EDTA, pH 8.0. The urea concentration of the solubilized protein was adjusted to 1 M and the sample was loaded onto Q Sepharose. The bound protein was eluted using a linear gradient from 0% to 100% Buffer B. (Buffer A: 100 mM Tris-HCl, 5 mM EDTA, 1 M urea, pH 8.0. Buffer B: 100 mM Tris-Cl, 5 mM EDTA, 1 M NaCl, 1 M urea, pH 8.0). Due to the formation of protein aggregates following elution, the urea concentration of the Q Sepharose material was adjusted to 8 M. The protein was further purified by gel filtration chromatography using SD200. The column was pre-equilibrated with 100 mM Tris-HCl, pH 8.0, 100 mM NaCl, 1% glycerol, 8 M urea plus 1% Na-deoxycholate. The eluted protein was subjected to a second IEX chromatography step using Source Q to remove 1% Na-deoxycholate. Bound protein was eluted with a

linear gradient from 20% to 60% Buffer B. (Buffer A: 100 mM Tris-Cl, pH 8.0, 8 M urea, 5 mM EDTA. Buffer B: 100 mM Tris-HCl, pH 8.0, 5 mM EDTA, 8 M urea, 1 M NaCl). Final polishing of the protein was completed by gel filtration chromatography using SD200 (Running Buffer: 100 mM Tris-HCl, pH 8.0, 8 M urea, 100 mM NaCl and 1% glycerol). Reducing agent was added to the SD200 fraction (2.5 mM DTT) and the protein was refolded by step-wise dialysis against decreasing concentrations of urea. The urea concentration was reduced sequentially against buffers that contained 100 mM Tris-HCl, pH 8.0, 100 mM NaCl, 1% glycerol and 6 M, 4 M, 2 M or no urea.

Refolding and Purification of STF2Δ.EIII+ Trimer

[0739] STF2Δ.EIII+ (SEQ ID NOS: 673, 674) from urea-solubilized inclusion bodies was efficiently refolded to form trimer product by simple dialysis as described above the trimer (3 of these STFΔ.EIII fusion proteins) was deduced based on molecular weight in SDS-PAGE. Following dialysis, endotoxin was removed by multiple extractions with Triton X-114. The trimer was purified and separated from monomer and aggregates by S200 size exclusion chromatography. The final product migrated as a single band with an apparent molecular weight of about 130 kDa on SDS-PAGE.

Refolding and Purification of STF2Δ.EIII+ Monomer

[0740] The monomeric form of STF2Δ.EIII+ (SEQ ID NOS: 673, 674) was produced consistently and efficiently by refolding using rapid dilution, which prevented individual STF2Δ.EIII+ fusion proteins from interacting with one another to form multimers, such as trimers (supra). STF2Δ.EIII+ solubilized from inclusion bodies in 4M urea was raised to 8M urea without reductant. The protein was then rapidly diluted in Tris/NaCl/glycerol buffer, pH 8.0, to about 0.1 mg/ml and a final urea concentration of 0.1M at room temperature. The monomer was further purified and separated from aggregates by S200 size exclusion chromatography. The final product migrated as a single band with an apparent molecular weight of about 43 kDa on SDS-PAGE.

Purification of STF2Δ.EIII+ (Serine Substitution of the Linker between STF2Δ and EII+, SEQ ID NO: 675)

[0741] STF2Δ.EIII+ (SEQ ID NOS: 675, 676) from solubilized inclusion bodies was refolded using a rapid dilution method similar to that used to refold the STF2Δ.EIII+ monomer. The refolded protein was captured on a butyl sepharose column and eluted while removing most of the endotoxin contamination. Eluate from the butyl sepharose purification was concentrated and put through 4 cycles of Triton X-114 extractions to reduce endotoxin levels down to about <0.1 EU/μg before a final purification step over SD200 gel filtration. The final pooled product migrated as a single band with an apparent molecular weight of about 43 kDa on SDS-PAGE and contained a trace amount of Triton X-114 (about 0.000015%).

Purification of STF2Δ.JEIII+ (SEQ ID NOS: 608, 609)

[0742] Protein was isolated from inclusion bodies under denaturing conditions. Inclusion bodies were washed with detergent (0.5% Triton X 100) and solubilized in 8 M urea, resulting in partial purification of the target protein. For endotoxin removal, protein was applied on a Source S cation

exchange column at low pH (about 3.5) and eluted with a salt gradient (0 to about 1 M NaCl). The protein was refolded using rapid dilution as described for STF2Δ.EIII+ monomer. The protein was then concentrated and further purified using SD200 to separate the monomeric form of the protein from aggregates. The purified material migrated with an apparent molecular weight of about 43 kDa on SDS PAGE and contained acceptable levels of endotoxin (about 0.03 EU/ug).

Fed Batch Production of Fusion Proteins

[0743] STF2Δ.EIII+ was produced in an aerobic bioreactor using a fed batch process. Three control loops were placed to control pH by acid (2 N HCl) or base (3 N NH₄OH) addition, temperature by heating (heating blanket) or cooling (time cycled cooling loop), and dissolved oxygen by compressed air flow (manually controlled), agitation (mixing speed) and O₂ flow (timed cycled) in cascade. Cells [BLR(DE3) pLysS that harbor the STF2Δ.EIII+ were adapted to and banked in MRSF media (see infra), and frozen in 25% glycerol. Cells were scaled up for the bioreactor by adding 1 mL of banked cells to 1 L of MRSF media and agitating at about 37° C. for about 15.5 to about 16.5 hours. Cells from the scale up process were added in a about 1:10 ratio to MRSF or MRBR synthetic media at about 37° C. and about 0.5 vvm air flow.

[0744] The process, was run in batch mode at about 37° C. until the cells oxygen consumption was such that the compressed air flow is about 1.5 vvm and the agitation is at the maximum, about 6 hours, when the temperature is dropped to between about 25° C. and about 33° C. The feed can be started before the culture is induced, or up to about 1 and about ½ hours after. The feed rate can be kept constant, or adjusted based on process variables (dissolved oxygen, glucose concentration). The culture was induced with IPTG upon batch glucose exhaustion. The culture was maintained for a minimum of about 2 hours and a maximum of about 20 hours.

MRBR Media		Trace Metal Solution 1000x	
Composition	g/L	Component	g/L
Glucose	20	EDTA, disodium	5
KH ₂ PO ₄	2.2	FeSO ₄ (7H ₂ O)	10
(NH ₄) ₂ SO ₄	4.5	ZnSO ₄ (7H ₂ O)	2
Citric Acid	1.0	MnSO ₄ (H ₂ O)	2
MgSO ₄ (7H ₂ O)	1.0	CoCl ₂ (6H ₂ O)	0.2
CaCl ₂	0.04	CuSO ₄ (5H ₂ O)	0.1
Trace Metals	1 ml	Na ₂ MoO ₄ (2H ₂ O)	0.2
Thiamine HCl	0.01	H ₃ BO ₃	0.1
Antifoam	0.05		

MRSF Media		Feed Media	
Composition	g/L	Composition	g/L
Glucose	10 (20 in bioreactor)	NaCl	0.5
KH ₂ PO ₄	7.8	FeSO ₄ (7H ₂ O)	2
(NH ₄) ₂ SO ₄	2.33	CaCl ₂	3.5
Citric Acid	1.0	MgSO ₄ (7H ₂ O)	12
MgSO ₄ (7H ₂ O)	1.0	Thiamine HCl	1
CaCl ₂	0.04	Trace Metals	1 ml

-continued

Trace Metals	1 ml	Glucose	100
Thiamine HCl	0.01		
Kanamycin	0.0075 (shake flask only)		

[0745] STF2Δ.EIIs+ was produced as inclusion bodies. Upon harvest, the cells were separated from the conditioned media by centrifugation (Beckman Avanti J-20 XP, JLA 8.1000 rotor, 10 kxg for about 20 minutes at about 4° C.) and resuspended in equal volume of 50 mM Tris, 100 mM NaCl, 1 mM EDTA, pH 8.0. The centrifugation was repeated under the same conditions, with the cells resuspended in a minimum volume of the same buffer. The suspension was passed through a homogenizer (APV-1000) at >10,000 psi for at least two passes.

[0746] The solids can be separated and the STF2Δ.EIIs solubilized by one of three methods; centrifugation, filtration, or fluidized bed chromatography.

Method 1

[0747] Solids are separated by centrifugation (Beckman Avanti J-20 XP, JA 20 rotor, 20 kxg for 20 minutes at 4° C.) and resuspended in 50 mM tris, 1 M NaCl, 1 mM EDTA, 1% glycerol, 0.5% Triton X-100, pH 8.0. This process was repeated up to 6 times (total) at increasing speeds and times (to a maximum of about 40 kxg for about 20 minutes). After the final pellet recovery, the pellet was resuspended in 50 mM Tris, 0.1M NaCl, 1 mM EDTA, pH 8.0 and clarified by centrifugation (Beckman Avanti J-20 XP, JA 20 rotor, 40 kxg for about 20 minutes at about 4° C.) The pellet was resuspended and dissolved in 50 mM Tris, 0.1M NaCl, 1 mM EDTA, 4 M urea, pH 8.0. Insolubles were removed by centrifugation (Beckman Avanti J-20 XP, JA 20 rotor, 40 kxg for about 50 minutes at about 4° C.), the supernatant retained for further processing.

[0748] After the multiple washes described above, STF2Δ.EIIs can also be dissolved in 50 mM acetate, 10 mM NaCl, 8M urea, pH about 4.1 to about 5.3 and clarified by centrifugation (Beckman Avanti J-20 XP, JA 20 rotor, 20 kxg for about 20 minutes).

Method 2

[0749] After homogenization, the lysate was captured in body feed and STF2Δ.EIIs+ extracted with urea containing buffer. Body feed is a filter aid designed to trap particles in a cake above a depth filter. The body feed (Advanced Minerals Corporation CelPure 65) is a diatomite (silica powder) with a high surface area and low permeability, retaining <0.2 μm particles. The filter aid was pre-mixed with the lysate and pumped over a depth filter (Ertel 703), building up a cake containing both body feed and lysate particles. The suspension creates a depth filter as the particles settle on the filter pad. A 50 mM Tris, 100 mM NaCl pH 8.0 wash was performed to remove soluble proteins and nucleic acids. A subsequent wash with 50 mM Tris, 100 mM NaCl, 4 M urea, pH 8 solubilizes and removes the STF2Δ.EIIs from the body feed for further processing.

Method 3

[0750] After the cells were initially resuspended in buffer, they were resuspended in sodium chloride and urea contain-

ing buffer at pH about 6 to about 8 and homogenized. The lysate was applied on a Streamline CST fluidized bed column (GE Healthcare) where the STF2Δ.EIIs+ binds to the resin and the particulates flow through. STF2Δ.EIIs+ may be eluted in low salt conditions at a pH greater than the load pH, in the presence or absence of detergents such as Triton X-100 or polysorbate 80.

SDS-PAGE

[0751] Proteins (typically about 5 μg) were diluted in SDS-PAGE sample buffer with and without β-mercaptoethanol. The samples were boiled for 5 minutes and loaded onto a 4-20% SDS polyacrylamide gel. Following electrophoresis, gels were stained with coomassie blue to visualize protein bands.

Endotoxin Assay

[0752] Endotoxin levels were measured using the QCL-1000 Quantitative Chromogenic LAL test kit (BioWhittaker #50-648U, Walkersville, Md.), following the manufacturer's instructions for the microplate method.

Protein Assay

[0753] Protein concentrations were determined by the MicroBCA Protein Assay Reagent Kit in a 96-well format using BSA as a standard (Pierce Biotechnology, Rockford, Ill.).

TLR5 Bioactivity Assay

[0754] HEK293 cells (ATCC, Catalog No. CRL-1573 Manassas, Va.) constitutively express TLR5, and secrete several soluble factors, including IL-8, in response to TLR5 signaling. Cells were seeded in 96-well microplates (about 50,000 cells/well), fusion proteins added and incubated overnight. The next day, the conditioned medium was harvested, transferred to a clean 96-well microplate, and frozen at -20° C. After thawing, the conditioned medium was assayed for the presence of IL-8 in a sandwich ELISA using an anti-human IL-8 matched antibody pair (Pierce, #M801E and #M802B, Rockford, Ill.) following the manufacturer's instructions. Optical density was measured using a microplate spectrophotometer (FARCyte, Amersham Biosciences, Piscataway, N.J.).

Plaque Reduction Neutralization Test (PRNT)

[0755] PRNT was performed according to Wang, et al., *J. Immunol.* 167:5273-5277 (2001). Briefly, serum samples were heat inactivated by incubation in a 56° C. water bath for about 30 min and were serially diluted in PBS with 5% gelatin from 1/10 to 1/2560. West Nile virus was diluted in PBS with 5% gelatin so that the final concentration was about 100 PFU/well. Virus was mixed with about 75 μl serum in a 96-well plate at about 37° C. for about 1 h. Aliquots of serum-virus mixture were inoculated onto confluent monolayers of Vero cells in a six-well tissue culture plate. The cells were incubated at about 37° C. for 1 h, and the plates were shaken every 15 min. The agarose overlay was then added. The overlay was prepared by mixing equal volumes of a solution consisting of 100 ml 2xMEM (Life Technologies) with sterile 2% agarose. Both solutions were placed in a 40° C. water bath for 1 h before adding the overlay. The cells were incubated for 4 days at 37° C. in a humidified 5% CO₂-air mixture. A second overlay with an additional 4% neutral red was added on day 5. Virus plaques were counted about 12 h later.

Antigenicity of STF2Δ-Fusion Proteins

[0756] ELISA plates (96-well) were coated overnight at 4° C. with serial dilutions (100 µl/well) of purified STF2Δ-fusion proteins (SEQ ID NOS: 761, 762, 657, 658, 673, 674) in PBS (about 2 µg/ml). Plates were blocked with 200 µl/well of Assay Diluent Buffer (ADB; BD Pharmingen) for one hour at room temperature. The plates were washed 3× in PBS-Tween, and then incubated with antibodies reactive with flagellin or the E domain of the construct. The expression of flagellin was detected using the mAb 6H11 (Intotek), while the antigenicity of WNV-E was monitored using a panel of mAb (5C5, 7H2, 5H10, 3A3, and 3D9) (Beasley, D. W., et al., *J. Virol.* 76:13097-13100 (2002)) were purchased from Bioreliance (Road Rockville, Md.). Antibodies diluted in ADB (about 100 µl/well) were incubated overnight at 4° C. The plates were washed 3× with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical, West Grove, Pa.) diluted in ADB were added (100 µl/well) and the plates were incubated at room temperature for 1 hour. The plates were washed 3× with PBS-T. After adding TMB (3,3',5,5'-tetramethylbenzidine) Ultra substrate (Pierce Biotechnology, Rockford, Ill.) and monitoring color development, A₄₅₀ was measured on a Tecan Farcyte microspectrophotometer.

Immunization of Mice

[0757] C3H/HeN mice (10 per group) were immunized intraperitoneally or subcutaneously with the indicated concentrations of fusion proteins or synthetic peptides on days 0, 14 and 28. On days 21 and 35, immunized animals were bled by retro-orbital puncture. Sera were harvested by clotting and centrifugation of the heparin-free blood samples. On day 35, mice were challenged with a lethal dose of WNV strain 2741 (Wang, T., et al., *J. Immunol* 167:5273-5277 (2001)). Survival was monitored for 21 days post-challenge.

Serum Antibody Determination

[0758] West Nile envelope protein specific IgG levels were determined by ELISA. ELISA plates (96-wells) were coated overnight at about 4° C. with 100 µl/well of West Nile E protein mAb 5C5, 7H2, 5H10, 3A3, and 3D9 (Beasley, D. W., et al., *J. Virol.* 76:13097-13100 (2002)) (Bioreliance, Road Rockville, Md.) in PBS at a concentration of 2 µg/ml. Plates were blocked with 200 µl/well of Assay Diluent Buffer (ADB; BD Pharmingen, San Diego Calif.) for one hour at room temperature. The plates were washed 3× in PBS-T. Dilutions of the sera in ADB were added (100 µl/well) and the plates were incubated overnight at 4° C. The plates were washed 3× with PBS-T. HRP-labeled goat anti-mouse IgG antibodies (Jackson Immunochemical, West Grove, Pa.) diluted in ADB were added (100 µl/well) and the plates were incubated at room temperature for 1 hour. The plates were washed 3× with PBS-T. After adding TMB (3,3',5,5'-tetramethylbenzidine) Ultra substrate (Pierce Biotechnology, Rockford, Ill.) and monitoring color development, A₄₅₀ was measured on a Tecan Farcyte microspectrophotometer.

[0759] Production of Pam3Cys.WNV001 Peptide Synthesis

[0760] Pam3Cys.WNV001 was synthesized by Bachem Bioscience, Inc. (King of Prussia, Pa.). WNV001 is a 20 amino acid peptide (SEQ ID NO: 771) of the West Nile virus

envelope protein chemically coupled to a tri-palmitoylcysteine (Pam3Cys) moiety through the amino terminal serine residue of the peptide. The chemical name for Pam3Cys.WNV001 is [Palmitoyl-Cys((RS)-2,3-di(palmitoyloxy)-propyl)-LTSGHLKCRVKMEKLQLKGT (SEQ ID NO: 771) acetate salt]. The molecular mass of Pam3Cys.WNV001 is 3163.3 daltons. The peptide was synthesized by Bachem using solid phase synthesis methodologies and FMOC chemistry. The amino acid sequence of Pam3Cys.WNV001 was assembled on an H-Pro-2-chlorotrityl chloride resin by solid phase peptide synthesis. The peptide chain was elongated by successive coupling of the amino acid derivatives. Each coupling step was preceded by an Fmoc-deprotection step and were accompanied by repeated washing of the resin. After coupling of the last amino acid derivative, the final Fmoc-deprotection step was performed. Finally, the peptide resin washed and dried under reduced pressure. During solid phase peptide synthesis color indicator tests were performed for each step to monitor the completion of the Fmoc-cleavage and the subsequent coupling of the amino acid derivatives. To couple Pam3Cys-OH to the elongated peptide, the lipid moiety was pre-activated with N,N'-dicyclohexyl-carbodiimide (DCCI) in the presence of 1-hydroxybenzotriazole (HOBt). The resulting solution was filtered and added to the peptide resin. At the end of the reaction time the peptide resin washed and dried under reduced pressure. Color indicator tests were performed to control the coupling of Pam3Cys-OH. The completed peptide was cleaved from the resin by incubating with trifluoroacetic acid (TFA). The liberated product (crude peptide material) was precipitated from the reaction mixture and lyophilized. The crude product was used for initial immunogenicity studies.

Synthesis of WNV-E Peptide Arrays

[0761] Peptide arrays (SEQ ID NOS: 703-754) were synthesized by Sigma Genosys (Woodlands, Tex.).

Results

West Nile Fusion Protein

[0762] West Nile virus (WNV) has emerged in recent years in temperate regions of Europe and North America, presenting a threat to public and animal health. The most serious manifestation of WNV infection is fatal encephalitis (inflammation of the brain) in humans and horses, as well as mortality in certain domestic and wild birds. WNV has also been a significant cause of human illness in the United States. The envelope glycoprotein of West Nile (WNV-E) and other flaviviruses may generate neutralizing and protective antibodies. By linking this antigen to a Toll-like Receptor ligand, the compositions, fusion proteins and polypeptides described herein may target appropriate antigen presenting cells without the need for adjuvant or other immune modulator formulations.

[0763] As described herein, several strategies have been implemented to facilitate production of West Nile virus envelope (WNV-E) fusion proteins in *E. coli*. One approach is to engineer a smaller WNV-E antigen by fusing domain III (EIII) and, optionally, with amino acids of domain II of the WNV-E protein to full-length STF2 (e.g., STF2.E, STF2.EIII+). Domain III is responsible for virus-host interactions and retains many West Nile virus neutralizing antibody epitopes. It also contains only 2 of the 12 cysteine

residues present within the full length envelope protein, making expression in *E. coli* more feasible. A second approach has been to delete the hyper-variable hinge region of flagellin (e.g., STF2Δ) thereby creating a smaller fusion protein (STF2Δ.EIII+). The hyper-variable region of flagellin is not required for TLR5 signaling and its removal may also reduce the immunogenic potential of flagellin. Both STF2.EIII+ and STF2Δ.EIII+ have been expressed in *E. coli* and purified. The purified proteins have been characterized for TLR5 signaling activity in bioassays and for E epitope display in ELISA assays using a panel of WNV-E polyclonal and neutralizing monoclonal antibodies. Results from these studies indicate that STF2Δ.EIII+ has higher PAMP activity and more conformation-sensitive neutralizing WNV-E epitopes than STF2.EIII+.

Purity of STF2.EIII+ and STF2Δ.EIII+

[0764] Several lots of STF2.EIII+ and STF2Δ.EIII+ have been produced in *E. coli* and purified (Table 12). STF2.EIII+ was expressed as a soluble protein and purified under non-denaturing conditions using a 4-step process, as described above, that included anion exchange chromatography and gel filtration. Final yields from 6 L cultures ranged from about 0.9 mg to about 3.8 mg and all preparations contained low levels of endotoxin as measured by standard LAL procedures (about <0.1 EU/μg protein, see supra). In contrast, STF2Δ.EIII+ formed inclusion bodies in *E. coli*, and was purified under denaturing conditions. All chromatography steps used to purify STF2Δ.EIII+ required the use of 8M urea. Following purification, the denatured protein was refolded by step-wise dialysis to allow for gradual urea removal. Refolding was typically carried out at protein concentrations of about 0.3 mg/ml without any loss due to protein precipitation. Two preparations of STF2Δ.EIII+ from a single 6 L culture yielded about 1.2 and about 6.7 mg of protein, both of which had acceptable endotoxin levels. As expected, purified STF2.EIII+ and STF2Δ.EIII+ migrated on SDS PAGE under reducing conditions as about 65 kDa and about 43 kDa proteins, respectively. Notably, STF2Δ.EIII+ migrated slightly faster under non-reducing conditions. This altered migration may be due to disulfide bond formation involving the two cysteines residues in domain III of the envelope protein. As well, a larger species of STF2Δ.EIII+ was detected by Western blot analysis whose molecular weight is consistent with a trimer form of the protein (“(STF2Δ.EIII+)×3 or 3 units of STF2Δ.EIII+”).

TABLE 11

Endotoxin levels and TLR-5 activity for STF2.EIII+ (SEQ ID NO: 658) and STF2Δ.EIII+ (SEQ ID NO: 674) fusion proteins.				
Batch Number	Protein	Yield (mg)	Endotoxin Levels (EU/μg)	TLR-5 EC ₅₀
052	STF2.EIII+	3.8	0.03	>5000.00 ng/ml
054	STF2.EIII+	0.9	0.02	1195.00 ng/ml
057	STF2.EIII+	1.6	0.07	197.92 ng/ml
044	STF2Δ.EIII+	1.2	0.07	1.13 ng/ml
045	STF2Δ.EIII+	6.7	0.07	4.34 ng/ml

TLR5 Activity in the HEK293 IL-8 Assay

[0765] To compare the PAMP activity of both fusion proteins, a TLR5 bioassay was performed. HEK293 IL-8

cells were treated with serial dilutions of two independent protein batches (FIGS. 62A and 62B). Cultures were incubated for a 24 hour period and conditioned media were harvested and assayed for IL-8 production by ELISA. As shown in FIG. 62A, STF2Δ.EIII+ showed potent TLR-5 activity. Regression analysis of the titration curve determined the EC₅₀ of batches 2004-044 and 2004-045 to be 1.13 ng/ml and 4.34 ng/ml, respectively (Table 11, supra). In both cases, the TLR5 specific-activity was at least about 10-fold higher than the control protein STF2.OVA. In contrast, 2 preparations of STF2.EIII+ showed significantly weaker TLR5 activity than STF2.OVA. The EC₅₀ of STF2.EIII+ batches 054 and 057 were about 1195.00 ng/ml and about 197.92 ng/ml.

Antigenicity of STF2.EIII+ and STF2Δ.EIII+

[0766] The antigenicity of STF2.EIII+ and STF2Δ.EIII+ was examined by direct ELISA using a flagellin monoclonal antibody specific for the N-terminal region of STF2 (6H11, Inotek Pharmaceuticals, Beverly, Mass.) and a panel of WNV-E-specific antibodies (5C5, 5H10, 3A3, 7H2 and 3D9, Bioreliance, Road Rockville, Md.) previously shown to neutralize West Nile virus in vitro. As shown in FIG. 63, a comparison of the reactivity of full length West Nile virus envelope protein with STF2Δ.EIII+ revealed that West Nile virus monoclonal antibodies 5C5, 5H10, 3A3 and 7H2, but not 3D9 recognize the fusion protein. This pattern of reactivity is consistent with the proposed location of 5C5, 5H10, 3A3 and 7H2 epitopes within EIII. The epitope for 3D9 lies outside of domain III of the West Nile virus envelope protein. As expected, all West Nile virus monoclonal antibodies reacted with full length West Nile virus envelope protein and the flagellin monoclonal only reacted with STF2Δ.EIII+. Both proteins reacted with a polyclonal West Nile virus envelope antiserum, but STF2Δ.EIII+ reactivity was somewhat reduced, perhaps due to the reduced number of potential epitopes present in the smaller domain.

[0767] Using 5C5 and 7H10 WNV monoclonal antibodies, a direct antigenic comparison was made between STF2.EIII+ and STF2Δ.EIII+ (FIGS. 64A, 64B, 64C and 64D). In these studies, plates were coated with the indicated proteins and then detected with polyclonal rabbit anti-E, or mouse monoclonal antibodies as described. As shown in FIGS. 64A, 64B, 64C and 64D, both STF2.EIII+ and STF2Δ.EIII+ were readily detected with the flagellin monoclonal antibody with no significant differences in reactivity. However, distinct reactivity with the anti-envelope monoclonal antibodies was observed. The reactivity of STF2Δ.EIII+ with either 5C5 or 7H2 was significantly greater than that observed with STF2.EIII+. Collectively, these results indicate that the flagellin 6H11 epitope of STF2Δ.EIII+ is un compromised and is comparable to the flagellin sequence of STF2.EIII+. They also highlight distinct differences in the antigenicity of the EIII domains of these proteins and indicate that STF2Δ.EIII+ contains more of the critical conformation dependent neutralizing epitopes than STF2.EIII+.

Efficacy and Immunogenicity

[0768] Several efficacy studies designed to examine the protective efficacy our candidates in C3H/HeN mice following challenge with West Nile virus have been completed. Studies typically consisted of 5 groups of mice (10 mice per group) immunized intraperitoneally (i.p.) or subcutaneously

(s.c.) on days 0, 14 and 28. On days 21 and 35, sera were harvested and tested for West Nile virus envelope protein—IgG antibody (ELISA) and the ability to neutralize West Nile virus in vitro (PRNT assay). On day 35, mice were challenged with a lethal dose of West Nile virus strain 2741. Survival was monitored for 21 days post-challenge.

[0769] Mice were immunized with PBS, *Drosophila* conditioned medium containing STF2.E (CM, positive control), 25 µg of STF2Δ.EIII+ i.p., 25 µg STF2Δ.EIII+ s.c., 25 µg STF2.EIII+ i.p. and 25 µg STF2.EIII+ s.c. The West Nile virus envelope protein antibody responses and survival data are shown FIGS. 65 and 66. By day 35 all groups that received STF2Δ.EIII+ had significant levels of West Nile virus envelope protein IgG. In contrast, mice that received STF2.EIII+ had no measurable West Nile virus envelope protein antibody response. Administration of STF2Δ.EIII+ i.p. or s.c led to 100% survival following West Nile virus challenge. Consistent with the poor immunogenicity of STF2.EIII+, little to no protection was provided by this candidate when compared to the PBS control. The poor immunogenicity and efficacy of STF2.EIII+ in this study are attributed to the reduced TLR5 activity and/or the weak EIII epitope reactivity of this protein.

Plaque Reduction Neutralization Titers

[0770] To further evaluate the West Nile virus envelope protein antibody response elicited by STF2Δ.EIII+ and potentially correlate protective efficacy with neutralizing antibody titers, the plaque reduction neutralization test (PRNT) was performed. Day 35 serum samples from efficacy studies described above were tested for their ability to block West Nile virus infection in cultured Vero cells. Briefly, pooled mouse serum samples were heat-inactivated and serially diluted two-fold in PBS with 0.5% gelatin. Dilutions starting with 1:10 were incubated with about 100 pfu of the West Nile virus strain 2741. The virus/serum mixture was incubated at about 37° C. for 1 h and then inoculated onto confluent monolayers of Vero cells (ATCC, Catalog Number CCL-81, Manassas, Va.) in duplicate wells of 6-well tissue culture plates. The virus was allowed to adsorb to the cell monolayer prior to adding a 1% agarose overlay. Infected cell cultures were incubated for 4 days at 37° C. followed by a second agarose overlay containing 4% neutral red. Virus plaques were counted 12 h later. Serum titers that led to 80% reduction in viral plaque numbers (PRNT₈₀) were recorded.

[0771] A summary of the PRNT₈₀ data from efficacy studies concerning STF2.EIII+ and STF2Δ.EIII+ is presented in Table 12 below. In two independent studies involving STF2.EIII+ where survival of about 50% or less was reported, pooled sera failed to inhibit plaque formation. This finding is not surprising given the weak antibody response elicited by this construct. In three efficacy studies involving STF2Δ.EIII+ where survival was about 70% or greater, pooled sera had neutralization titers of 1:40 or better. Neutralization titers of 1:40 or greater typically correlate with protection in vivo.

TABLE 12

Survival and PRNT ₈₀ Results for STF2.EIII+ (SEQ ID NO: 658), STF2Δ.EIII+ (SEQ ID NO: 674) and STF2.E (SEQ ID NO: 762) CM (Control Media) Fusion Proteins					
Batch	Candidate	Route	Study #	Survival (%)	PRNT ₈₀ (dilution)
054	STF2.EIII+	i.p.	3	50	Negative
057	STF2.EIII+	i.p.	4	11	Negative
057	STF2.EIII+	s.c.	4	20	negative
044	STF2Δ.EIII+	i.p.	2	70	1:40
045	STF2Δ.EIII+	i.p.	3	90	1:40
045	STF2Δ.EIII+	s.c.	3	100	1:160
045	STF2Δ.EIII+	i.p.	4	100	1:80
045	STF2Δ.EIII+	s.c.	4	100	1:40
—	STF2.E CM	i.p.	3	90	1:640
—	STF2.E CM	i.p.	4	—	1:1280

STF2Δ.EIII+ a Modified Version of STF2Δ.EIII+

[0772] Protein preparations of STF2Δ.EIII+ tested in the mouse efficacy studies described above were purified by anion-exchange and size-exclusion chromatography steps carried out under denaturing conditions followed by refolding using step-wise dialysis. With this process, two predominant species that correspond to the monomeric and trimeric forms of STF2Δ.EIII+ were generated and present as a mixture in the final product. To minimize the heterogeneity of the final product, new refolding and purification methods have been developed that favor the production of either monomer or trimer. Because it is unclear which form of STF2Δ.EIII+ is the active component or if both are equally potent, both species have been produced in milligram quantities and tested for efficacy in mice.

[0773] It was initially unclear as to why STF2Δ.EIII+ refolding resulted in the formation of a trimeric species. However, when the sequence of the STF2Δ.EIII+ expression construct was re-examined, we identified a cysteine residue within the linker sequence that separates STF2Δ from EIII+. The presence of this cysteine would likely interfere with the formation of the appropriate disulfide bond during refolding and might account for the trimeric form of STF2Δ.EIII+. This unnecessary cysteine was changed to a serine using site-directed mutagenesis and the modified protein (STF2Δ.EIII_S+) was produced and purified. It should be noted that refolding the serine-substituted construct yielded only monomeric protein.

[0774] Protective efficacy of STF2Δ.EIII+ (monomer) and STF2Δ.EIII_S (trimer) were evaluated in C3H/HeN mice following challenge with West Nile virus. Five groups of mice (10 per group) were immunized with about 25 µg of protein s.c. on days 0, 14 and 28. On days 21 and 35, sera were harvested and tested for WNV-E IgG antibody (ELISA). On day 38, mice were challenged with a lethal dose of WNV strain 2741 and survival was monitored for 21 days. ELISA results from boost 2 (day 35, FIG. 67) and survival data (FIG. 68) indicate that all constructs elicited significant levels of WNV-E reactive IgG prior to viral challenge and provided about 90% to about 100% protection against the lethal infection. These findings indicate that monomeric or multimeric (e.g., trimers) forms of STFΔ.EIII+ are efficacious and removal of the additional cysteine from the construct does not appreciably impact

potency. Removal of the cysteine within the linker sequence may simplify purification of the protein by reducing heterogeneity following protein refolding.

Conclusion

[0775] Two recombinant fusion proteins containing the *Salmonella typhimurium* flagellin (STF2) fused to EIII+ domain of West Nile virus envelope protein have been generated. One includes the full length STF2 sequence (STF2.EIII+) and the other a modified version of STF2 that lacks the internal hypervariable region of STF2 (STF2Δ.EIII+). Both proteins have been expressed in *E. coli* and purified by conventional means using anion exchange chromatography and gel filtration. STF2.EIII+ was produced as a soluble protein and was purified under non-denaturing conditions. In contrast, STF2Δ.EIII+ was expressed as an insoluble protein and was purified under denaturing conditions and refolded by step-wise dialysis to remove urea. In HEK293 IL8 assays, preparations of STF2Δ.EIII+ showed greater TLR-5 activity than STF2.EIII+.

[0776] In envelope protein epitope display analysis using ELISA assays and West Nile virus envelope protein antibodies, STF2Δ.EIII+ displayed more of the critical conformation dependent neutralizing epitopes. Consistent with the potent TLR-5 activity and envelope protein epitope antigenicity observed with STF2Δ.EIII+, STF2Δ.EIII+ was highly immunogenic and efficacious in mice challenged with a lethal dose of West Nile virus. Because monomeric and trimeric species of STF2Δ.EIII+ were generated during the purification process of this protein, a cysteine within the linker sequence of the expression construct was changed to a serine. Removal of this cysteine eliminated the production of trimeric forms of the protein during refolding and resulted in the generation of monomeric product that displayed potent efficacy in vivo.

Japanese Encephalitis Fusion Protein

[0777] JE virus is localized in Asia and northern Australia (about 50,000 cases with about 10,000 deaths annually). An approved inactivated virus vaccine was recently associated with a case of acute disseminated encephalomyelitis, prompting the Japanese Ministry of Health, Labor and Welfare to recommend the nationwide suspension of the vaccine. Given the complexities of producing inactivated viruses in infected mouse brains or even in cell culture, and the potential for adverse events associated with inactivated viruses, the opportunity for recombinant-based JE vaccine is appealing.

[0778] A STF2Δ.JEIII+ fusion construct was constructed. The JE EIII+DNA fragment was generated synthetically and codon optimized for expression in *E. coli*. The sequence was ligated into pET24STF2Δ to generate pETSTF2Δ.JEIII+. Expression constructs have been screened by restriction analysis and for expression in *E. coli* BLR(DE3) by IPTG induction. The DNA sequence of each construct has been confirmed, and production of the protein has been scaled up. A batch of material has been generated. A total of about 24 mg of material was purified. This material has potent TLR5 activity, acceptable levels of endotoxin (about 0.03 EU/μg) and a A280/A260 ratio of about 1.3.

Flavivirus Peptides

Identification of WNV-E Specific Antibody Epitopes

[0779] To identify linear epitopes within the West Nile virus envelope protein that are recognized by antisera from STFΔ.EIII+ immunized mice, several synthetic peptide arrays were generated. One array consisted of overlapping peptides of 20 amino acids in length that spanned the entire West Nile virus domain III and parts of domain II (SEQ ID NOS: 728-754). ELISA results with this array identified a highly reactive 20 amino acid sequence that mapped to the N-terminal region of domain III and included part of the domain I domain CRVKMEKLQLKGT TYGVCSK (SEQ ID NO: 728). To fine map this epitope, additional arrays were generated that focused on the domain I and II junctions (SEQ ID NOS: 703-754). These arrays included an alanine substitution scan to identify amino acids critical for antibody binding (SEQ ID NOS: 728-754). As shown in FIGS. 69 and 70, antisera from STF2Δ.EIII (monomer and trimer) and STF2Δ.EIII+ immunized mice reacted with peptides that spanned the EI/EII junction (peptides E-30 to E-42) and included the E2-21 peptide CRVKMEKLQLKGT TYGVCSK (SEQ ID NO: 728). This reactivity was severely reduced when specific amino acids (E6, K7, L10 and K11) were changed to alanines (FIG. 71). Although it is not known if the antibodies that recognize this epitope are neutralizing, efforts are underway to design and test a peptide vaccine based on this region of WNV-E.

Immunogenicity of Pam3Cys.WNV001 Peptide Vaccine.

[0780] A lipidated West Nile virus envelope protein fused to Pam3Cys on the N-terminal end was synthesized using the 20 amino acid sequence LTSGHLKCRVKMEKLQLKGT (SEQ ID NO:772) (Putnak, R., et al, *Vaccine* 23:4442-4452 (2005)). The immunogenicity of this peptide was tested in C3H/HeN mice and compared to peptide without Pam3Cys (FIG. 72). The reactivity of antisera from immunized animals was tested by direct ELISA as described in the legend and the results indicate that the Pam3Cys.WNV001 peptide is significantly more immunogenic than the peptide without the TLR2 modification. The antisera from these studies will be tested in virus neutralization assays (PRNT) to determine if the antibodies elicited will neutralize West Nile virus in vitro. The lipidated peptide will also be tested in the West Nile virus challenge model to assess protective efficacy against a lethal virus challenge.

Assay Development

Competition ELISA Assay Development

[0781] To assess the neutralizing potential of antisera derived from immunized mice, a competition ELISA assay was developed using well-characterized monoclonal antibody (7H2) that neutralizes West Nile virus in culture and reacts with a conformation-sensitive epitope within the EIII domain of the West Nile virus envelope protein antigen. The assay was designed as a capture ELISA that measures the ability of sera from immunized animals to prevent 7H2 from binding West Nile virus envelope protein. Serial dilutions ranging from 1:10 to 1:5000 of day 35 mouse antisera from efficacy study 4 (FIGS. 65 and 66, Table 13) were incubated with biotinylated West Nile virus envelope protein and then added to ELISA plates pre-coated with 7H2 monoclonal antibody (Bioreliance, Road Rockville, Md.). Following several washes to remove unbound material, bound West Nile virus envelope protein was detected using avidin-HRP.

Results from a representative experiment are shown in FIG. 69. At dilutions of 1:25, a measurable loss of West Nile virus envelope protein binding to 7H2-coated plates was observed when antisera derived from animals immunized with STF2Δ.EIII were tested. No competition was detected with antisera derived from mock immunized animals that received PBS in place of antigen. These initial results demonstrate that antibodies elicited by STF2Δ.EIII+ compete with 7H2 for binding West Nile virus envelope protein. These findings are consistent with the protection from WNV

infection observed in animals immunized with STF2Δ.EIII+ and help establish a correlation between antibody epitope reactivity in vitro and efficacy in vivo.

[0782] While this invention has been particularly shown and described with references to example embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

SEQUENCE LISTING

The patent application contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<http://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US20070224205A1>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

What is claimed is:

1. A composition comprising a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

2. The composition of claim 1, further including at least a portion of at least one member selected from the group consisting of a Toll-like Receptor 1 agonist, a Toll-like Receptor 2 agonist, a Toll-like Receptor 3 agonist, a Toll-like Receptor 4 agonist, a Toll-like Receptor 6 agonist, a Toll-like Receptor 7 agonist, a Toll-like Receptor 8 agonist and a Toll-like Receptor 9 agonist.

3. The composition of claim 2, wherein the Toll-like Receptor 2 agonist is a Pam3Cys.

4. The composition of claim 3, wherein the Pam3Cys further includes at least one additional cysteine residue.

5. The composition of claim 1, wherein at least one cysteine residue substitutes for at least one amino acid residue in a naturally occurring flagellin amino acid sequence of the flagellin component.

6. The composition of claim 5, wherein the cysteine residue is remote to at least one amino acid of the Toll-like Receptor 5 recognition site of the flagellin component.

7. The composition of claim 1, wherein the flagellin component includes at least a portion of a naturally occurring flagellin amino acid sequence in combination with the cysteine residue.

8. The composition of claim 7, wherein the cysteine residue is at least one member selected from the group consisting of the amino-terminal amino acid of the flagellin component and the carboxy-terminal amino acid of the flagellin component.

9. The composition of claim 7, wherein the cysteine residue is remote to at least one amino acid of the Toll-like Receptor 5 recognition site of the flagellin component.

10. The composition of claim 1, wherein the flagellin component lacks at least a portion of a hypervariable region.

11. The composition of claim 1, further including at least a portion of at least one antigen.

12. The composition of claim 11, wherein the antigen is chemically conjugated to the flagellin component.

13. The composition of claim 12, wherein the antigen is essentially hydrophobic.

14. The composition of claim 13, wherein chemical conjugation results in increased aqueous solubility of the antigen as a component of the composition.

15. The composition of claim 1, wherein the cysteine residue is in the hypervariable region of the flagellin component.

16. The composition of claim 1, wherein the flagellin is at least a portion of at least one member selected from the group consisting of a *S. typhimurium* flagellin, an *E. coli* fliC flagellin, *S. muenchen* flagellin, *P. aeruginosa* flagellin and *Listeria monocytogenes* flagellin.

17. The composition of claim 16, wherein the flagellin is at least a portion of a *S. typhimurium* flagellin and the *S. typhimurium* flagellin is selected from the group consisting of SEQ ID NO: 812 and SEQ ID NO: 816.

18. The composition of claim 17, wherein at least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence of the flagellin component.

19. The composition of claim 18, wherein the amino acid is at least one amino acid selected from the group consisting of amino acid 1, 237, 238, 239, 240, 241 and 495 of SEQ ID NO: 812.

20. The composition of claim 17, wherein the flagellin is at least a portion of a *S. typhimurium* flagellin and the *S. typhimurium* flagellin includes at least a portion of SEQ ID NO: 816.

21. The composition of claim 20, wherein at least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component.

22. The composition of claim 21, wherein the amino acid is at least one amino acid selected from the group consisting of amino acid 1, 240, 241, 242, 243, 244 and 505 of SEQ ID NO: 816.

23. The composition of claim 16, wherein the flagellin is at least a portion of an *S. muenchen* flagellin and the *S. muenchen* flagellin includes at least a portion of SEQ ID NO: 504.

24. The composition of claim 23, wherein at least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component.

25. The composition of claim 24, wherein the amino acid is at least one amino acid selected from the group consisting of amino acid 1, 237, 238, 239, 240, 241 and 504 of SEQ ID NO: 504.

26. The composition of claim 16, wherein the flagellin is at least a portion of *P. aeruginosa* flagellin and the *P. aeruginosa* flagellin includes at least a portion of SEQ ID NO: 815.

27. The composition of claim 26, wherein at least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component.

28. The composition of claim 27, wherein the amino acid is at least one amino acid selected from the group consisting of amino acid 1, 211, 212, 213 and 393 of SEQ ID NO: 815.

29. The composition of claim 16, wherein the flagellin includes at least a portion of a *Listeria monocytogenes* flagellin and the *Listeria monocytogenes* flagellin includes at least a portion of SEQ ID NO: 820.

30. The composition of claim 29, wherein at least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component.

31. The composition of claim 30, wherein the amino acid is at least one amino acid selected from the group consisting of amino acid 1, 151, 152, 153, 154 and 287 of SEQ ID NO: 820.

32. The composition of claim 16, wherein the flagellin is at least a portion of *E. coli* flagellin and the *E. coli* flagellin includes at least a portion of SEQ ID NO: 502.

33. The composition of claim 32, wherein at least one cysteine residue substitutes for at least one amino acid in a naturally occurring flagellin amino acid sequence flagellin component.

34. The composition of claim 33, wherein the amino acid is at least one amino acid selected from the group consisting of amino acid 1, 238, 239, 240, 241, 242, 243 and 497 of SEQ ID NO: 502.

35. The composition of claim 11, wherein the antigen is at least a portion of an influenza antigen.

36. The composition of claim 35, wherein the influenza antigen is at least a portion of an integral membrane protein of influenza.

37. The composition of claim 35, wherein the influenza antigen is an influenza A antigen.

38. The composition of claim 35, wherein the influenza antigen is an influenza B antigen.

39. The composition of claim 35, wherein the influenza antigen is an influenza C antigen.

40. The composition of claim 36, wherein the integral membrane protein is at least a portion of a hemagglutinin protein.

41. The composition of claim 40, wherein the portion of the hemagglutinin protein includes at least a portion of a maturational cleavage site of the hemagglutinin protein.

42. The composition of claim 41, wherein the maturational cleavage site includes at least one member selected from the group consisting of SEQ ID NO: 530, SEQ ID NO: 532, SEQ ID NO: 533, SEQ ID NO: 534, SEQ ID NO: 535, SEQ ID NO: 600, SEQ ID NO: 601, SEQ ID NO: 602, SEQ ID NO: 603, SEQ ID NO: 821, SEQ ID NO: 796, SEQ ID NO: 797, SEQ ID NO: 798, SEQ ID NO: 799 and SEQ ID NO: 822.

43. The composition of claim 36, wherein the integral membrane protein includes at least a portion of a matrix 2 protein.

44. The composition of claim 11, wherein the antigen is at least a portion of at least one member selected from the group consisting of a West Nile viral protein, a Langat viral protein, a Kunjin viral protein, a Murray Valley encephalitis viral protein, a Japanese encephalitis viral protein, a Tick-borne encephalitis viral protein, Dengue 1 viral protein, Dengue 2 viral protein, Dengue 3 viral protein, Dengue 4 viral protein, hepatitis C viral protein and a Yellow fever viral protein.

45. The composition of claim 11, wherein the antigen includes a pathogen-related antigen.

46. The composition of claim 45, wherein the pathogen-related antigen is at least one member selected from the group consisting respiratory syncytial virus, *Plasmodium falciparum* and *Plasmodium vivax*.

47. The composition of claim 1, further including a carrier protein.

48. The composition of claim 47, wherein the carrier protein is at least one member selected from the group consisting of a tetanus toxoid, a *Vibrio cholerae* toxoid, a diphtheria toxoid, a cross-reactive mutant of diphtheria toxoid, a *E. coli* B subunit of a heat labile enterotoxin, a tobacco mosaic virus coat protein, a rabies virus envelope protein, a rabies virus envelope glycoprotein, a thyroglobulin, a heat shock protein 60, a keyhole limpet hemocyanin and an early secreted antigen tuberculosis-6.

49. The composition of claim 1, wherein at least one member selected from the group consisting of an arginine residue, a serine residue and a histidine residue substitutes for at least one lysine of at least one member selected from the group consisting of at least a portion of the flagellin component and at least a portion of the flagellin.

50. A composition comprising a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor.

51. The composition of claim 50, wherein the Toll-like Receptor agonist component includes at least a portion of a naturally occurring Toll-like Receptor agonist in combination with the cysteine residue.

52. The composition of claim 51, wherein the cysteine residue is remote to at least one amino acid of the Toll-like Receptor recognition site of the Toll-like Receptor agonist component.

53. The composition of claim 50, wherein the Toll-like Receptor agonist is at least a portion of at least one member selected from the group consisting of a Toll-like Receptor 1 agonist, a Toll-like Receptor 2 agonist, a Toll-like Receptor 3 agonist, a Toll-like Receptor 4 agonist, a Toll-like Receptor

5 agonist, a Toll-like Receptor 6 agonist, a Toll-like Receptor 7 agonist, a Toll-like Receptor 8 agonist and a Toll-like Receptor 9 agonist.

54. The composition of claim 53, further including at least a portion of at least one antigen.

55. A composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5.

56. The composition of claim 55, wherein the flagellin is at least a portion of at least one member selected from the group consisting of a *S. typhimurium* flagellin, an *E. coli* fliC flagellin, *S. muenchen* flagellin, *P. aeruginosa* flagellin and *Listeria monocytogenes* flagellin.

57. The composition of claim 56, wherein the flagellin is at least a portion of a *S. typhimurium* flagellin and the *S. typhimurium* flagellin is selected from the group consisting of SEQ ID NO: 812 and SEQ ID NO: 816.

58. The composition of claim 57, wherein the lysine residue that is substituted with the arginine is at least one lysine residue selected from the group consisting of lysine 19, 41, 58, 135, 160, 177, 179, 203, 215, 221, 228, 232, 241, 251, 279, 292, 308, 317, 326, 338, 348, 357, 362, 369, 378, 384, 391 and 410 of SEQ ID NO: 812.

59. The composition of claim 57, wherein the lysine residue that is substituted with the arginine is at least one lysine residue selected from the group consisting of lysine 20, 42, 59, 136, 161, 177, 182, 189, 209, 227, 234, 249, 271, 281, 288, 299, 319, 325, 328, 337, 341, 355, 357, 369, 381, 390, 396, 403, 414 and 422 of SEQ ID NO: 816.

60. The composition of claim 56, wherein the flagellin is at least a portion of an *E. coli* fliC flagellin and the *E. coli* flagellin includes at least a portion of SEQ ID NO: 814.

61. The composition of claim 56, wherein the flagellin is at least a portion of a *S. muenchen* flagellin and the *S. muenchen* flagellin includes at least a portion of SEQ ID NO: 813.

62. The composition of claim 56, wherein the flagellin is at least a portion of a *P. aeruginosa* flagellin and the *P. aeruginosa* flagellin includes at least a portion of SEQ ID NO: 815.

63. The composition of claim 56, wherein the flagellin is at least a portion of a *Listeria monocytogenes* flagellin and the *Listeria monocytogenes* flagellin includes SEQ ID NO: 500.

64. The composition of claim 55, further including at least a portion of at least one antigen.

65. The composition of claim 64, wherein the antigen is conjugated to the flagellin component after substitution of the lysine for arginine.

66. The composition of claim 64, wherein the antigen is at least a portion of an influenza antigen.

67. The composition of claim 66, wherein the influenza antigen is at least a portion of an integral membrane protein.

68. The composition of claim 67, wherein the integral membrane protein includes at least a portion of a hemagglutinin protein.

69. The composition of claim 67, wherein the integral membrane protein includes at least a portion of a matrix 2 protein.

70. The composition of claim 66, wherein the influenza antigen is an influenza A antigen.

71. The composition of claim 66, wherein the influenza antigen is an influenza B antigen.

72. The composition of claim 66, wherein the influenza antigen is an influenza C antigen.

73. A composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5.

74. A composition comprising a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, whereby the flagellin component activates a Toll-like Receptor 5.

75. A method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein the flagellin component includes at least one cysteine residue and whereby the flagellin component activates a Toll-like Receptor 5.

76. The method of claim 75, further including at least a portion of at least one antigen.

77. A method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one arginine, whereby the flagellin component activates a Toll-like Receptor 5.

78. The method of claim 77, further including at least a portion of at least one antigen.

79. A method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one serine residue, whereby the flagellin component activates a Toll-like Receptor 5.

80. The method of claim 79, further including at least a portion of at least one antigen.

81. A method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a flagellin component that is at least a portion of a flagellin, wherein at least one lysine of the flagellin component has been substituted with at least one histidine residue, and whereby the flagellin component activates a Toll-like Receptor 5.

82. The method of claim 81, further including at least a portion of at least one antigen.

83. A method of stimulating an immune response in a subject, comprising the step of administering to the subject a composition that includes a Toll-like Receptor agonist component that is at least a portion of a Toll-like Receptor agonist, wherein the Toll-like Receptor agonist component includes at least one cysteine residue in a position where a cysteine residue does not occur in the native Toll-like Receptor agonist, whereby the Toll-like Receptor agonist component activates a Toll-like Receptor agonist.

84. The method of claim 83, further including at least a portion of at least one antigen.