

(19) **DANMARK**

(10) **DK/EP 4132943 T3**



(12) **Oversættelse af  
europæisk patentskrift**

Patent- og  
Varemærkestyrelsen

- 
- (51) Int.Cl.: **C 07 K 1/22 (2006.01)** **C 07 K 14/31 (2006.01)** **C 07 K 14/315 (2006.01)**  
**C 07 K 16/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-07-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-05-29**
- (86) Europæisk ansøgning nr.: **21736946.1**
- (86) Europæisk indleveringsdag: **2021-07-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2023-02-15**
- (86) International ansøgning nr.: **EP2021069564**
- (87) Internationalt publikationsnr.: **WO2022013272**
- (30) Prioritet: **2020-07-15 EP 20186024** **2021-02-23 EP 21158714**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **NAVIGO PROTEINS GMBH, Heinrich-Damerow-Str. 1, 06120 Halle/Saale, Tyskland**
- (72) Opfinder: **KNICK, Paul, , 06120 Halle/Saale, Tyskland**  
**BOBOLOWSKI, Hanna, , 06120 Halle/Saale, Tyskland**  
**KAHL, Mathias, , 06120 Halle/Saale, Tyskland**  
**FIEDLER, Erik, , 06120 Halle/Saale, Tyskland**  
**Haupts, Ulrich, , 06120 Halle/Saale, Tyskland**
- (74) Fuldmægtig i Danmark: **Patrade A/S, Ceresbyen 75, 6., 8000 Århus C, Danmark**
- (54) Benævnelse: **Immunoglobulin-bindende proteiner til affinitetsoprensning**
- (56) Fremdragne publikationer:  
**EP-A1- 3 650 465**  
**WO-A1-2017/009421**  
**WO-A1-2019/093439**  
**WO-A1-2019/152318**  
**WO-A1-2021/122943**  
**US-A1- 2010 221 844**  
**US-A1- 2019 177 376**



# DESCRIPTION

Description

## FIELD OF THE INVENTION

[0001] The present invention relates to immunoglobulin (Ig) binding proteins comprising one or more domains having highly hydrophobic amino acids with branched side chains (Iso, Leu, Val), or aromatic amino acids (Tyr, Phe, or Trp), corresponding to positions 4 or 6 or 8 of the Ig binding protein of SEQ ID NO: 1 or functionally similar proteins. The novel proteins have superior properties for highly efficient purification methods for antibodies (immunoglobulins), for example, the proteins have high binding capacity and high chemical stability. The invention further relates to affinity matrices comprising the Ig binding proteins of the invention. The invention also relates to a use of these Ig binding proteins or affinity matrices for affinity purification of immunoglobulins and to methods of affinity purification using the Ig binding proteins of the invention.

## BACKGROUND OF THE INVENTION

[0002] Many biotechnological and pharmaceutical applications require the removal of contaminants from a sample containing antibodies. An established procedure for capturing and purifying antibodies is affinity chromatography using the bacterial cell surface Protein A from *Staphylococcus aureus* as selective ligand for immunoglobulins (see, for example, review by Huse et al., J. Biochem. Biophys. Methods 51, 2002: 217-231). Wild-type Protein A binds to the Fc region of IgG molecules with high affinity and selectivity. Variants of Protein A with improved properties such as alkaline stability are available for purifying antibodies and various chromatographic matrices comprising Protein A ligands are commercially available. However, currently available Protein A based chromatography matrices show a loss of binding capacity for immunoglobulins following exposure to alkaline conditions and require elution conditions at pH below 4.

[0003] EP 3 650 465 A1 relates to Ig binding proteins and discloses variants thereof for the design of polypeptide libraries. WO 2019/152318 A1 discloses an Ig binding protein having a D8E mutation, and further relates to alkaline stability in the context of cysteine residues in the C-terminal region.

## TECHNICAL PROBLEMS UNDERLYING THE INVENTION

**[0004]** Most large scale production processes for antibodies or Fc-containing fusion proteins use Protein A for affinity purification. However, due to limitations of Protein A applications in affinity chromatography there is a need in the art to provide novel Ig binding proteins with improved properties that specifically bind to immunoglobulins in order to facilitate affinity purification of immunoglobulins. To maximally exploit the value of the chromatographic matrices comprising Ig binding proteins it is desirable to use the affinity ligand matrices multiple times. Between chromatography cycles a thorough cleaning procedure is required for sanitization and removal of residual contaminants on the matrix. In this procedure, it is general practice to apply alkaline solutions with high concentrations of NaOH to the affinity ligand matrices. Wild-type Protein A domains cannot withstand such harsh alkaline conditions for an extended time and quickly lose binding capacity for immunoglobulin. Further, for a repeated use of affinity ligand matrices, a cleaning step under harsh acidic conditions is required.

**[0005]** Accordingly, there is an ongoing need in this field to obtain novel proteins capable of binding proteins comprising an Ig sequence, for example antibodies, and to withstand the harsh cleaning conditions applied in affinity purification of immunoglobulins.

**[0006]** The present invention relates to Ig binding proteins that are particularly well-suited for affinity purification of immunoglobulins. In particular, the Ig binding proteins of the invention have several advantages. One significant advantage of the Ig binding proteins of the invention is their improved stability at high pH for a prolonged time period (such as more than 2 days) without reducing the Ig binding capacities in combination with high dynamic binding capacities. Further, the novel proteins of the invention are particularly useful for affinity purification of antibodies where mild acidic elution conditions are required.

**[0007]** The above overview does not necessarily describe all problems solved by the present invention.

## **SUMMARY OF THE INVENTION**

**[0008]** An aspect of the present invention relates to an Ig binding protein suitable for affinity purification. The present invention is defined by the appended claims.

1. [1] Further disclosed herein is an Immunoglobulin (Ig) binding protein comprising one or more Ig binding domains, wherein at least one Ig binding domain corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to positions 4 or 6 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or an aromatic amino acid. The Ig binding protein is alkaline stable (at least 20 h at 0.5 M NaOH). The Ig binding protein of the present invention comprises one or more Ig binding domains, wherein at least one Ig binding domain corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 1, wherein the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I), leucine (L),

- valine (V) or an aromatic amino acid which is tryptophan (W) or phenylalanine (F), and wherein the Ig binding protein is stable under alkaline conditions of 0.5 M NaOH for at least 20 h.
2. [2] Further disclosed herein is an Ig binding protein according to item [1], wherein
    1. (a) the amino acid corresponding to position 4 of SEQ ID NO: 1 is tryptophan (W) or phenylalanine (F), or
    2. (b) the amino acid corresponding to position 6 of SEQ ID NO: 1 is isoleucine (I), tryptophan (W), or tyrosine (Y), or leucine (L).
  3. [3] Disclosed herein is an Ig binding protein according to item [1] or [2], wherein one or more amino acid(s) corresponding to position 10, 14, 16, 17, 18, or 28 of SEQ ID NO: 1 is/are selected from the group of histidine (H) or acidic amino acids selected from aspartate (D) or glutamate (E), in particular in position 14 or 28. In various embodiments, the Ig binding protein of the invention has at least 80 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or valine (V) or an aromatic amino acid which is tryptophan (W) or phenylalanine (F), and the amino acid corresponding to position 14 of SEQ ID NO: 1 is histidine. In various embodiments, the Ig binding protein of the invention has at least 80 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or valine (V) or an aromatic amino acid which is tryptophan (W) or phenylalanine (F), the amino acid corresponding to position 14 of SEQ ID NO: 1 is histidine (H), and the amino acid corresponding to position 29 of SEQ ID NO: 1 is lysine (K). Preferably, the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or an aromatic amino acid, which may be any of tryptophan (W), phenylalanine (F), or tyrosine (Y). More preferably, the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L), even more preferably the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I). In various embodiments, the Ig binding protein may have a cysteine (C) residue at the position corresponding to position 43 or 46 of SEQ ID NO: 1.
  4. [4] Disclosed herein is an Ig binding protein according to any one of items [1]-[3], wherein the amino acid corresponding to position 29 of SEQ ID NO: 1 is lysine (K).
  5. [5] Disclosed herein is an Ig binding protein according to any one of items [1]-[4], wherein at least one domain comprises or consists of an amino acid sequence of any one of SEQ ID NOs: 4-36 and 40-49. The Ig binding protein may comprise an amino acid sequence of any of SEQ ID NOs: 4-9, 20-36, and 40-49, or an amino acid sequence with at least 89.5 % identity thereto to any of SEQ ID NO: 4-36, 40-49.
  6. [6] Disclosed herein is an Ig binding protein according to any one of items [1]-[5], wherein said protein binds to one or more of IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>4</sub>, IgM, IgA, Ig fragments, Fc fragments, Fab fragments, fusion proteins comprising an Ig region, and conjugates comprising an Ig region. The Ig binding protein may bind to proteins comprising an Fc region or may bind to Fc fragments.
  7. [7] Disclosed herein is an Ig binding protein according to any one of items [1]-[6], wherein the protein comprises 2, 3, 4, 5, or 6 domains linked to each other.
  8. [8] Disclosed herein is an Ig binding protein according to item [7], wherein the protein is

- a homo-multimer or a hetero-multimer. The Ig binding protein may be a dimer comprising the sequence of SEQ ID NO: 37, SEQ ID NO: 50, or SEQ ID NO: 51.
9. [9] Disclosed herein is an Ig binding protein according to any one of items [1]-[8], wherein the protein is immobilized to a solid support. The Ig binding protein may be immobilized to a solid support by a cysteine (C) at the position corresponding to position 43 or 46 of SEQ ID NO: 1.
  10. [10] Disclosed herein is an Ig binding protein according to any one of items [1]-[9], wherein the Ig binding protein is stable under alkaline conditions, optionally for at least 20 h at 0.5 M NaOH.
  11. [11] Disclosed herein is an affinity separation matrix comprising the Ig binding protein of any one of items [1] to [10] coupled to said affinity separation matrix.
  12. [12] Disclosed herein is the use of the Ig binding protein of any one of items [1] to [10], or the affinity separation matrix of item [11] for affinity purification of any protein with affinity to the Ig binding protein.
  13. [13] A-Disclosed herein is a method of affinity purification of a protein comprising an Ig sequence, the method comprising:
    1. a) providing a liquid that contains protein comprising an Ig sequence;
    2. b) providing an affinity separation matrix according to item [11] comprising at least one Ig binding protein of any one of items [1] to [10] coupled to said affinity separation matrix of item [11];
    3. c) contacting said affinity separation matrix with the liquid under conditions that permit binding of the at least one Ig binding protein according to any one of items [1]-[10] to a protein comprising an Ig sequence; and
    4. d) eluting said protein comprising an Ig sequence from said affinity purification matrix, thereby obtaining an eluate containing said immunoglobulin.
  14. [14] Disclosed herein is a method according to item [13], wherein in step (d) more than 95 % of the protein comprising the Ig sequence is eluted at pH 3.7 or higher from the affinity separation matrix comprising the Ig binding protein according to any of items [1]-[10]. In step (d) more than 95 % of the protein comprising the Ig sequence may be eluted at pH 4.5 from the affinity separation matrix comprising the Ig binding protein according to any of items [1]-[10].
  15. [15] Disclosed herein is a method according to any of items [13]-[14], comprising the additional step (e) of cleaning the affinity purification matrix with an alkaline cleaning liquid, optionally wherein at least 90 % of the Ig binding protein retains Ig binding activity after incubation for at least 20 h at 0.5 M NaOH.
  16. [16] Further disclosed herein is an Immunoglobulin (Ig) binding domain having at least 80 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to positions 4, 6, or 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or an aromatic amino acid. The Immunoglobulin (Ig) binding domain is/corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to positions 4, 6, or 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or an aromatic amino acid, as described under item [1] above.

This summary of the invention does not necessarily describe all features of the present

invention as defined by the appended claims. Other embodiments will become apparent from a review of the ensuing detailed description.

## BRIEF DESCRIPTION OF THE FIGURES.

### [0009]

**Figure 1.** Amino acid sequences of novel Ig binding proteins. The numbers in the top row refer to the corresponding amino acid position in the Ig binding protein.

**Figure 2, 3, 4.** Caustic stability of Ig binding proteins coupled to Praesto Epoxy 85 resin (coupling 18 h at 35 °C) after incubation for 20 h with 0.5 M NaOH. SBC determination with 6 mg Gammanorm. Coupling of variants and parent to the resin via Cysteine located in the third helix (position 43C).

**Figure 2.** Variants 8I, 8V, 8F, 8W, 8L, and 8Y show significantly improved remaining activity after long term incubation at alkaline conditions, in comparison to the parent molecule (cs26).

**Figure 3.** Variants 4F and 4W show improved remaining activity after long term incubation at alkaline conditions, in comparison to the parent molecule (cs26).

**Figure 4.** Variants 6F, 6I, 6L, 6V, 6W, 6Y, 6R show improved remaining activity after long term incubation at alkaline conditions, in comparison to the parent molecule (cs26).

**Figure 5.** Elution profile of Belimumab from affinity ligand (SEQ ID NO: 51). The antibody was injected onto the column with immobilized affinity ligand (SEQ ID NO: 51) and eluted with linear pH gradient from pH 6.0 to 2.0. The pH of the elution peak maximum was used as readout.

**Figure 6.** DBC10% determination of affinity ligand SEQ ID NO: 51. Belimumab was loaded onto resin with immobilized affinity ligand SEQ ID NO: 51 ("ID51" in the figure) until 10 % target breakthrough. Elution of Belimumab was performed at pH 4.8 followed by CIP at pH 1.7. The chromatogram shows complete elution of bound Belimumab at pH 4.8.

## DETAILED DESCRIPTION OF THE INVENTION

**[0010]** Before the present invention is described in detail below, it is to be understood that this invention is not limited to the particular methodology, protocols and reagents described herein as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to limit the scope of the present invention which is limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs.

**[0011]** Preferably, the terms used herein are consistent with the definitions provided in "A multilingual glossary of biotechnological terms: (IUPAC Recommendations)", Leuenberger, H.G.W, Nagel, B. and Kölbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland).

**[0012]** Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated member, integer or step or group of members, integers or steps but not the exclusion of any other member, integer or step or group of members, integers or steps.

**[0013]** As used in the description of the invention and the appended claims, the singular forms "a", "an" and "the" are used interchangeably and intended to include the plural forms as well and fall within each meaning, unless the context clearly indicates otherwise. Also, as used herein, "and/or" refers to and encompasses any and all possible combinations of one or more of the listed items, as well as the lack of combinations when interpreted in the alternative ("or").

**[0014]** The term "about", as used herein, encompasses the explicitly recited amounts as well as deviations therefrom of  $\pm 10\%$ . More preferably, a deviation of  $5\%$  is encompassed by the term "about".

**[0015]** Several documents (for example: patents, patent applications, scientific publications, manufacturer's specifications etc.) are cited throughout the text of this specification. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention. In the event of a conflict between the definitions or teachings of such references and definitions or teachings recited in the present specification, the text of the present specification takes precedence.

**[0016]** All sequences referred to herein are disclosed in the attached sequence listing that, with its whole content and disclosure, is a part of this specification.

**[0017]** In the context of the present invention, the term "Ig binding protein" or "immunoglobulin-binding protein" is used to describe proteins that are capable to specifically bind to an immunoglobulin.

**[0018]** Further, in the context of the present invention, the term "Ig binding domain" or "immunoglobulin-binding domain" is used to describe proteins that are capable to specifically bind to an immunoglobulin. The Ig binding proteins or Ig binding domains of the present invention are sometimes referred to herein as ligands of the invention. The "immunoglobulin" or "Ig" as understood herein can include, but is not necessarily limited to, mammalian IgG, such as for example human IgG<sub>1</sub>, human IgG<sub>2</sub>, human IgG<sub>4</sub>, mouse IgG, rat IgG, goat IgG, bovine IgG, guinea pig IgG, rabbit IgG; human IgM, human IgA; and an immunoglobulin fragment comprising a Fc region (also referred to as "Fc fragment" or "Fc") and/or an

immunoglobulin fragment comprising a Fab region (also referred to as "Fab fragment" or "Fab"). The Ig binding proteins are capable of binding to entire immunoglobulins, and to Ig fragments comprising a Fc region and/or Ig fragments comprising a Fab region. The definition "immunoglobulin" as understood herein includes fusion proteins comprising an immunoglobulin, fragment of an immunoglobulin comprising a Fc region (Fc fragment), fragment of an immunoglobulin comprising a Fab region (Fab fragment), fusion proteins comprising a fragment of an immunoglobulin comprising a Fc region, fusion proteins comprising a fragment of an immunoglobulin comprising a Fab region, conjugates comprising an Ig or an Ig fragment comprising a Fc region (Fc fragment), and conjugates comprising an Ig fragment comprising a Fab region (Fab fragment).

**[0019]** As will be appreciated by a person of ordinary skill in the art, the terms "immunoglobulin" and "antibody" may be used interchangeably herein. Any definitions disclosed herein concerning the term "immunoglobulin" apply to the term "antibody" accordingly.

**[0020]** The term "binding" according to the invention preferably relates to a specific binding. "Specific binding" means that an Ig binding protein or an Ig binding domain binds stronger to an immunoglobulin for which it is specific compared to the binding to another non-immunoglobulin target.

**[0021]** The term "binding activity" refers to the ability of an Ig binding protein or Ig binding domain of the invention to bind to immunoglobulin. For example, the binding activity can be determined before and/or after alkaline treatment. The terms (immunoglobulin) "binding activity" and "binding capacity" may be used interchangeably herein. The binding activity can be determined for an Ig binding protein or for an Ig binding protein coupled to a matrix, i.e., for an immobilized Ig binding protein. Also, the binding activity can be determined for an Ig binding domain or for an Ig binding domain coupled to a matrix, i.e., for an immobilized Ig binding domain. The term "artificial" refers to an object that is not naturally occurring, i.e. the term refers to an object that has been produced or modified by man. For example, a polypeptide or polynucleotide sequence that has been generated by man (e.g. for example in a laboratory by genetic engineering, by shuffling methods, or by chemical reactions, etc.) or intentionally modified is artificial.

**[0022]** The term "dissociation constant" or " $K_D$ " defines the specific binding affinity. As used herein, the term " $K_D$ " (usually measured in "mol/L", sometimes abbreviated as "M") is intended to refer to the dissociation equilibrium constant of the particular interaction between a first protein and a second protein. In the context of the present invention, the term  $K_D$  is particularly used to describe the binding affinity between an Ig binding protein or an Ig binding domain and an immunoglobulin. An Ig binding protein or Ig binding domain of the invention is considered to bind to an immunoglobulin, if it has a dissociation constant  $K_D$  to immunoglobulin of at least 500 nM or less, or preferably 100 nM or less, more preferably 50 nM or less, even more preferably 10 nM or less.

**[0023]** The terms "protein" and "polypeptide" refer to any linear molecular chain of two or more amino acids linked by peptide bonds and does not refer to a specific length of the product. Thus, "peptides", "protein", "amino acid chain," or any other term used to refer to a chain of two or more amino acids, are included within the definition of "polypeptide," and the term "polypeptide" may be used instead of, or interchangeably with any of these terms. The term "polypeptide" is also intended to refer to the products of post-translational modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, proteolytic cleavage, modification by non-naturally occurring amino acids and similar modifications which are well-known in the art. Thus, Ig binding proteins comprising two or more protein domains also fall under the definition of the term "protein" or "polypeptides".

**[0024]** The terms "alkaline stable" or "alkaline stability" or "caustic stable" or "caustic stability" (also abbreviated as "cs" herein) may be used interchangeably herein and refer to the ability of the Ig binding protein or Ig binding domain of the invention to withstand alkaline conditions without significantly losing the ability to bind to immunoglobulins. The skilled person in this field can easily test alkaline stability by incubating an Ig binding protein or Ig binding domain with, for example, sodium hydroxide solutions, e.g., as described in the Examples, and subsequent testing of the binding capacity or binding activity to immunoglobulin by routine experiments known to someone skilled in the art, for example, by chromatographic approaches. The alkaline stability may be determined by coupling the Ig binding protein or Ig binding domain of the invention to a surface plasmon resonance (SPR) sensor chip, and assaying the binding capacity or binding activity for immunoglobulin before and after exposure to an alkaline solution. The alkaline treatment can be performed, for instance, in 0.5 M NaOH for an extended period of time, e.g., at least 20 h.

**[0025]** Ig binding proteins or Ig binding domains of the invention as well as matrices comprising Ig binding proteins or Ig binding domains of the invention exhibit an "increased" or "improved" alkaline stability, meaning that the molecules and matrices incorporating said Ig binding proteins or Ig binding domains are stable under alkaline conditions for an extended period of time relative to a reference. In various embodiments, the reference may be the parent molecule cs26 having the sequence of any one of SEQ ID NOs: 1-3, preferably the sequence of SEQ ID NO: 3 (cs26 43C). In various other embodiments, the reference may be the parent molecule cs26 of any of SEQ ID NOs: 1-3 having a substitution D8E (Asp8Glu), preferably the parent molecule cs26 of SEQ ID NO: 3 having a substitution D8E (Asp8Glu).

**[0026]** The term "variant" as used herein includes an amino acid sequence of an Ig binding protein or Ig binding domain that differs from another amino acid sequence by at least one amino acid substitution, deletion or insertion. These modifications may be generated by genetic engineering or by chemical synthesis or chemical reactions carried out by man.

**[0027]** The term "conjugate" as used herein relates to a molecule comprising or essentially consisting of at least a first protein attached chemically to other substances such as to a second protein or a non-proteinaceous moiety.

**[0028]** The term "modification" or "amino acid modification" refers to an exchange, a deletion, or an insertion of an amino acid at a particular position in a polypeptide sequence by another amino acid. Given the known genetic code, and recombinant and synthetic DNA techniques, the skilled scientist can readily construct DNAs encoding the amino acid variants.

**[0029]** The term "substitution" or "amino acid substitution" refers to an exchange of an amino acid at a particular position in a polypeptide sequence by another amino acid. The term "deletion" or "amino acid deletion" refers to the removal of an amino acid at a particular position in a polypeptide sequence.

**[0030]** The term "insertions" or "amino acid insertion" refers to the addition of amino acids to the polypeptide sequence.

**[0031]** Throughout this description, the amino acid residue position numbers are designated as corresponding to those for example in SEQ ID NO: 1.

**[0032]** The term "amino acid sequence identity" refers to a quantitative comparison of the identity (or differences) of the amino acid sequences of two or more proteins. "Percent (%) amino acid sequence identity" or "percent identical" or "percent identity" with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. In various embodiments, the term "sequence identity" means that two (nucleotide or) amino acid sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 70% sequence identity, or at least 80% sequence identity, or at least 85% sequence identity, or at least 90% sequence identity, or at least 95% sequence identity or more.

**[0033]** To determine the sequence identity, the sequence of a query protein is aligned and compared to the sequence of a reference protein. Methods for sequence alignment and sequence comparison algorithms are well known in the art. For example, for determining the extent of an amino acid sequence identity of an arbitrary polypeptide relative to the reference amino acid sequence, the SIM Local similarity program is preferably employed. For multiple alignment analysis, ClustalW as known to someone skilled in the art is preferably used.

**[0034]** The extent of sequence identity is generally calculated with respect to the total length of the unmodified sequence. As used herein, the phrases "percent identical" or "percent (%) amino acid sequence identity" or "percent identity", in the context of two polypeptide sequences, refer to two or more sequences or subsequences that have in some embodiments at least 89.5 %, in some embodiments at least 91 %, some embodiments at least 92 %, in some embodiments at least 93 %, in some embodiments at least 94 %, in some embodiments at least 95 %, in some embodiments at least 96 %, in some embodiments at least 97 %, in some embodiments at least 98 %, and in some embodiments 100 % amino acid residue identity, when compared and aligned for maximum correspondence, as measured using one of

the following sequence comparison algorithms or by visual inspection. For clarity reasons, for example a sequence with at least 89.5 % identity includes all sequences with identities higher than 89.5% identity, e.g. embodiments with at least 89.6 %, at least 90 %, at least 91 %, at least 92 %, at least 93 %, at least 94 %, at least 95 %, at least 96 %, at least 97 %, at least 98 %, at least 99 %, or 100 % amino acid identity.

**[0035]** The percent identity exists in some embodiments over a region of at least 52 residues, in some embodiments over a region of at least 53 residues, in some embodiments over a region of at least 54 residues, in some embodiments over a region of at least 55 residues, in some embodiments over a region of at least 56 residues, in some embodiments over a region of at least 57 residues, and in some embodiments over a region of at least 58 residues.

**[0036]** The term "fused" means that polypeptide components or units are linked by peptide bonds, either directly or via peptide linkers. In various embodiments, the term "fused" may mean that polypeptide components or units are linked by a non-peptide linker, e.g., through chemical conjugation.

**[0037]** The term "fusion protein" relates to a protein comprising at least a first protein joined genetically to at least a second protein. A fusion protein is created through joining of two or more genes that originally coded for separate proteins. Thus, a fusion protein may comprise a multimer of identical or different proteins which are expressed as a single, linear polypeptide. In various embodiments, a fusion protein is created through joining of two or more polypeptides via a non-peptide linker, e.g., through chemical conjugation. In various embodiments, a dimer of an Ig binding protein or Ig binding domain of the present invention may be considered as a "fusion protein".

**[0038]** As used herein, the term "linker" refers in its broadest meaning to a molecule that covalently joins at least two other molecules. In typical embodiments of the present invention, a "linker" is to be understood as a moiety that connects an Ig binding protein or Ig binding domain with at least one further Ig binding protein or Ig binding domain, i.e. a moiety linking two protein domains to each other to generate a dimer or a multimer. In preferred embodiments, the "linker" is a peptide linker, i.e. the moiety linking the two binding proteins or binding domains is one single amino acid or a peptide comprising two or more amino acids. In various embodiments, a dimer or multimer of the present invention may comprise a linker joining two or more Ig binding proteins or Ig binding domains with each other.

**[0039]** The term "chromatography" refers to separation technologies which employ a mobile phase and a stationary phase to separate one type of molecules (e.g., immunoglobulins) from other molecules (e.g. contaminants or other immunoglobulins) in the sample. The liquid mobile phase contains a mixture of molecules and transports these across or through a stationary phase (such as a solid matrix). Due to the differential interaction of the different molecules in the mobile phase with the stationary phase, molecules in the mobile phase can be separated. The term "affinity chromatography" refers to a specific mode of chromatography in which a ligand coupled to a stationary phase interacts with a molecule (i.e. immunoglobulin) in the

mobile phase (the sample) i.e. the ligand has a specific binding affinity or binding capacity for the molecule to be purified. As understood in the context of the invention, affinity chromatography involves the addition of a (liquid) sample containing an immunoglobulin to a stationary phase which comprises a chromatography ligand, such as an Ig binding protein or Ig binding domain of the invention.

**[0040]** The terms "solid support" or "solid matrix" are used interchangeably herein, and in various embodiments are used for the stationary phase.

**[0041]** The terms "affinity matrix" or "affinity separation matrix" or "affinity chromatography matrix", as used interchangeably herein, refer to a matrix, e.g. a chromatographic matrix, onto which an affinity ligand e.g., an Ig binding protein or Ig binding domain of the invention is attached. The ligand (e.g., Ig binding protein or Ig binding domain) is capable of specific binding to a molecule of interest (e.g., an immunoglobulin as defined above) which is to be purified or removed from a mixture (in a liquid sample). As will be appreciated by a person of ordinary skill in the art, the terms "affinity matrix" or "affinity separation matrix" or "affinity chromatography matrix" describe the separation of a molecule of interest (in particular an immunoglobulin) by using an Ig binding protein or Ig binding domain of the invention. Accordingly, the terms "affinity matrix" or "affinity separation matrix" or "affinity chromatography matrix" or "separation matrix" may be used interchangeably herein.

**[0042]** The term "affinity purification" as used herein refers to a method of purifying immunoglobulins of interest as defined above from a liquid (sample) by binding immunoglobulins of interest as defined above to an Ig binding protein or Ig binding domain that is immobilized to a matrix. Thereby, all other components of the mixture except immunoglobulins of interest are removed. In various embodiments, said other components of the mixture may include, e.g., other immunoglobulins that are not of interest. In a further step, immunoglobulins of interest are eluted in purified form. The terms "affinity purification" or "affinity chromatography purification" or "affinity separation" or "affinity chromatography separation" may be used interchangeably herein.

#### **EMBODIMENTS OF THE INVENTION**

**[0043]** The present invention will now be further described. In the following passages different embodiments of the invention are defined in more detail. Each embodiment defined below may be combined with any other embodiments unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous.

**[0044]** The present invention provides an Ig binding domain, which corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L) or valine (V) or an aromatic amino acid, which is tryptophan (W) or phenylalanine (F).

**[0045]** Preferably, the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or leucine (L), more preferably the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I). In various embodiments, the amino acid corresponding to position 14 of SEQ ID NO: 1 is histidine (H), and/or the amino acid corresponding to position 29 of SEQ ID NO: 1 is lysine (K). The Ig binding domain may further have a cysteine (C) residue at the position corresponding to position 43 or 46 of SEQ ID NO: 1. The Ig binding domain is stable under alkaline conditions of 0.5 M NaOH for at least 20 h. As described elsewhere herein, an Ig binding protein of the invention comprises one or more such Ig binding domains.

**[0046]** In one embodiment, the Ig binding protein comprises one or more domains, wherein at least one domain comprises or essentially consists of or consists of an amino acid substitution at one or more amino acids corresponding to positions 4, 6, or 8 of SEQ ID NO: 1, the substitution at the amino acid corresponding to position 8 of SEQ ID NO: 1 is an amino acid selected from the group of Iso (I), Leu (L), Phe (F), Val (V), or Trp (W), and wherein the amino acid sequence of the Ig binding protein is at least 80% identical to SEQ ID NO: 1.

**[0047]** In one embodiment, the Ig protein comprises one or more domains, wherein at least one domain comprises an amino acid substitution at an amino acid corresponding to position 8 of SEQ ID NO: 1 selected from the group of highly hydrophobic amino acids with branched side chains (Iso, Leu, Val), or aromatic amino acids (Phe, or Trp), and wherein the amino acid sequence of the Ig binding protein is at least 80 % identical to SEQ ID NO: 1. In some embodiments, the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I) or tyrosine (Y). In some preferred embodiments, the amino acid corresponding to position 8 of SEQ ID NO: 1 (or functionally equivalent proteins) is isoleucine (I). In various embodiments, Isoleucine (I) in a position corresponding to position 8 of SEQ ID NO: 1 improves the Ig binding capacity and caustic stability. Further, an isoleucine in position corresponding to position 8 of SEQ ID NO: 1 (or functionally equivalent proteins) might improve the expression of the protein. For an Ig binding ligand of the present invention with high binding capacity and caustic stability suitable for uses in affinity chromatography, it is important that the amino acid in position 8 is not selected from any one of arginine (R), serine (S), threonine (T), or alanine (A) (see **FIGURE 2**). The present invention demonstrates an improved caustic stability over these substitutions at position 8 of SEQ ID NO: 1 as well as over the substitution D8E (Asp8Glu). Accordingly, in various embodiments of the present invention the amino acid corresponding to position 8 of SEQ ID NO: 1 is not glutamic acid (E).

**[0048]** As further described herein, the Ig binding protein or Ig binding domain may be selected from the following (1) to (3): (1) a protein comprising an amino acid sequence corresponding to SEQ ID NO: 1 wherein an amino acid residue at the 8th position is substituted by Ile, Leu, Val, or an aromatic amino acid; (2) a protein comprising an amino acid sequence specified in the (1) further having deletion, substitution and/or addition of one or more amino acid residues in a position other than the 8th position; (3) a protein comprising an amino acid sequence having a sequence identity of at least 80% or more with the amino acid sequence specified in the (1), provided that the amino acid substitution specified in the (1) at the 8th

position is not further mutated in (3).

**[0049]** Further disclosed herein is an Ig protein comprising an amino acid substitution at a position corresponding to position 4 of SEQ ID NO: 1 selected from the group of from the group of highly hydrophobic amino acids with branched side chains (Iso, Leu), or aromatic amino acids (Tyr, Phe, or Trp), and wherein the amino acid sequence of the Ig binding protein is at least 80% identical to SEQ ID NO: 1. The amino acid corresponding to position 4 of SEQ ID NO: 1 may be tryptophan (W) or phenylalanine (F). Aromatic amino acids such as W or F in a position corresponding to position 4 of SEQ ID NO: 1 improve the binding capacity and caustic stability of the ligand in affinity chromatography. For an Ig binding ligand with high binding capacity and caustic stability suitable for uses in affinity chromatography, it is important that the amino acid in position 4 is not selected from arginine (R), serine (S), threonine (T), alanine (A), or valine (V) (see **FIGURE 3**).

**[0050]** Disclosed herein is a protein selected from the following (1) to (3): (1) a protein comprising an amino acid sequence corresponding to SEQ ID NO: 1 wherein an amino acid residue corresponding to the 4th position is substituted by Ile, Leu, or an aromatic amino acid; (2) a protein comprising an amino acid sequence specified in the (1) further having deletion, substitution and/or addition of one or more amino acid residues in a position other than the 4th position; (3) a protein comprising an amino acid sequence having a sequence identity of at least 80% or more with the amino acid sequence specified in the (1), provided that the amino acid substitution specified in the (1) at the 4th position is not further mutated in (3).

**[0051]** Further disclosed herein is an Ig protein comprising an amino acid substitution at a position corresponding to position 6 of SEQ ID NO: 1 selected from selected from the group of Iso, Leu, Tyr, Phe, or Trp, and wherein the amino acid sequence of the Ig binding protein is at least 80 % identical to SEQ ID NO: 1. The amino acid corresponding to position 6 of SEQ ID NO: 1 may be isoleucine (I), tryptophan (W), or tyrosine (Y), or leucine (L). Aromatic amino acids such as Trp (W) or Tyr (Y), or amino acids selected from Iso (I) or Leu (L) in a position corresponding to position 6 of SEQ ID NO: 1 improve the binding capacity and caustic stability of the ligand in affinity chromatography. For an Ig binding ligand with high binding capacity and caustic stability suitable for uses in affinity chromatography, it is important that the amino acid in position 6 is not selected from valine (V), serine (S) or alanine (A) (see **FIGURE 4**).

**[0052]** Disclosed herein is a protein selected from the following (1) to (3): (1) a protein comprising an amino acid sequence of SEQ ID NO: 1 wherein an amino acid residue at the 6th position is substituted by Ile, Leu, or an aromatic amino acid; (2) a protein comprising an amino acid sequence specified in the (1) further having deletion, substitution and/or addition of one or more amino acid residues in a position other than the 6th position; (3) a protein comprising an amino acid sequence having a sequence identity of at least 80% or more with the amino acid sequence specified in the (1), provided that the amino acid substitution specified in the (1) at the 6th position is not further mutated in (3).

**[0053]** The surprising advantage of the Ig binding proteins and Ig binding domains of the

invention is the stability under extreme conditions such as high pH (pH 13 and higher) without losing Ig binding properties. The Ig binding proteins and Ig binding domains as described herein demonstrate alkali stability for a prolonged period of time without impairing the Ig-binding properties (see **FIGURES 2, 3, 4** and **Examples**). Further, they are stable at low pH without significantly losing Ig binding properties. The alkali stability feature is particularly important for chromatography approaches with cleaning procedures using alkaline solutions with high NaOH concentrations to remove contaminants on the matrix so for example that the matrix can be used several times. In addition to high caustic stability, Ig binding proteins show high coupling efficiencies, as shown in the **Examples**.

**[0054]** Further, an important step in affinity chromatography is the elution of the protein of interest, particular an immunoglobulin of interest, that is bound to the Ig binding protein or Ig binding domain of the invention. This step is usually done at low pH. The affinity ligands of the invention do not lose binding properties to Ig after this treatment, while elution of the protein of interest is possible at low pH.

**[0055]** In some circumstances, it is important to have conditions for the elution of antibodies (immunoglobulins) from the affinity ligand at pH between 3.7 higher, such as pH 4.3 and above, for example, up to pH 5.5. In order to improve characteristics of the ligands of the invention, further modifications can be made to the ligands described above. In some embodiments, one or more amino acid(s) corresponding to position 10, 14, 16, 17, 18, or 28 of SEQ ID NO: 1 are selected from the group of histidine (H) or acidic amino acids selected from aspartate (D) or glutamate (E). In some embodiments, the amino acid corresponding to position 14 of SEQ ID NO: 1 is H. In some embodiments, the amino acid corresponding to position 16 of SEQ ID NO: 1 is H. In other embodiments, the amino acid corresponding to position 28 of SEQ ID NO: 1 is H. In some embodiments, the amino acid corresponding to position 28 of SEQ ID NO: 1 is E. In other embodiments, the amino acid corresponding to position 9 of SEQ ID NO: 1 is H. In some embodiments, the amino acid corresponding to position 10 of SEQ ID NO: 1 is H. Ligands of the invention with His (H), Asp (D), or Glu (E) in positions corresponding to positions 10, 14, 16, 17, 18, or 28 of SEQ ID NO: 1 weaken the Fc binding affinity and allow elution of the bound Ig protein of interest at pH higher than 4.0 or even pH 4.3, up to pH 5.5.

**[0056]** It has surprisingly been found that an Ig binding protein or Ig binding domain of the present invention comprising a histidine (H) at the position corresponding to position 14 of SEQ ID NO: 1 or a histidine (H) at the position corresponding to position 16 of SEQ ID NO: 1 is particularly suitable for elution of Ig molecules of interest from the immobilized ligand at mild pH conditions (up to pH 5.5) (see **Table 3**). This feature is particularly useful for isolating immunoglobulins, in particular Ig having an Fc region, using separation matrices, wherein the elution step has to be carried out at mild acidic conditions higher than pH 3.7, in particular in the range of pH 4.0 up to and including pH 5.5.

**[0057]** The Ig binding protein or Ig binding domain may be selected from the following (1) to (3): (1) a protein comprising an amino acid sequence corresponding to SEQ ID NO: 1 wherein

the amino acid residue corresponding to the 8th position is Ile, Leu, Val, or an aromatic amino acid; (2) a protein comprising an amino acid sequence specified in the (1) wherein the amino acid residue corresponding to the 10<sup>th</sup>, 14<sup>th</sup>, 16<sup>th</sup>, 17<sup>th</sup>, 18<sup>th</sup>, or 28<sup>th</sup> position is His, Asp, or Glu, preferably wherein the amino acid corresponding to position 14 is His, the amino acid corresponding to position 16 is His, the amino acid corresponding to position 10 is Asp or His, the amino acid corresponding to position 17 is His, the amino acid corresponding to position 18 is Glu, or the amino acid corresponding to position 28 is His or Glu, more preferably wherein the amino acid corresponding to position 14 is His, (3) a protein comprising an amino acid sequence having a sequence identity of at least 80% or more with the amino acid sequence specified in the (1), provided that the amino acid specified in (1) corresponding to the 8th position and the amino acid in the (2) corresponding to the 10<sup>th</sup>, 14<sup>th</sup>, 16<sup>th</sup>, 17<sup>th</sup>, 18<sup>th</sup>, or 28<sup>th</sup> position is not further mutated in (3).

**[0058]** Further modifications can be introduced to the protein to modify certain properties for affinity chromatography. For example, a cysteine can be added to the C-terminus. Alternatively, a cysteine can be introduced at a position within helix 3 of the protein, for example, in position 43 or position 46, to enable efficient coupling to the matrix.

**[0059]** The position corresponding to position 29 might be exchanged in order to lower the binding of Ig (by eliminating Fab-VH3 binding) and improve elution properties at higher pH values. The amino acid corresponding to position 29 of SEQ ID NO: 1 may be Lys (K). The resulting ligand of the invention has at least 80 % identity to SEQ ID NO: 1.

**[0060]** Disclosed herein is an Ig binding protein or Ig binding domain selected from the following (1) to (3):

1. (1) protein comprising an amino acid sequence corresponding to SEQ ID NO: 1 wherein the amino acid residue corresponding to the 8th position is Ile, Leu, Val, or an aromatic amino acid;
2. (2) a protein comprising an amino acid sequence specified in (1) wherein the amino acid residue corresponding to the 10<sup>th</sup>, 14<sup>th</sup>, 16<sup>th</sup>, 17<sup>th</sup>, 18<sup>th</sup>, or 28<sup>th</sup> position is His, Asp, or Glu, preferably wherein the amino acid corresponding to position 14 is His, the amino acid corresponding to position 16 is His, the amino acid corresponding to position 10 is Asp or His, the amino acid corresponding to position 17 is His, the amino acid corresponding to position 18 is Glu, or the amino acid corresponding to position 28 is His or Glu, in particular wherein the amino acid corresponding to position 14 is His ;
3. (3) a protein comprising an amino acid sequence specified in the (1) wherein the amino acid residue corresponding to the 29th position is Lys;
4. (4) a protein comprising an amino acid sequence having a sequence identity of at least 80 % or more with the amino acid sequence specified in the (1), provided that the amino acids specified in the (1), (2), and (3) are not further mutated in (4).

**[0061] Preferred Ig binding domains.** In various embodiments, the Ig binding domain comprises or consists of an amino acid sequence of any one of SEQ ID NOs: 4-36 and 40-49, or an amino acid with at least 89.5 %, at least 90 %, at least 91 %, at least 92 %, at least 93 %, at least 94 %, at least 95 %, at least 96 %, at least 97 %, at least 98 %, or at least 99 % identity thereto. In various embodiments, the Ig binding domain comprises or essentially consists of or consists of an amino acid sequence of any of SEQ ID NOs: 4-36 and 40-49, or an amino acid with at least 95 %, at least 96 %, at least 97 %, at least 98 %, or at least 99 % identity to any of SEQ ID NOs: 4-36 and 40-49.

**[0062] Preferred Ig binding proteins.** In some embodiments, the Ig binding protein comprises one or more binding domains wherein at least one domain comprises or consists of an amino acid sequence of any one of SEQ ID NOs: 4-36 and 40-49, or an amino acid with at least 89.5 %, at least 90 %, at least 91 %, at least 92 %, at least 93 %, at least 94 %, at least 95 %, at least 96 %, at least 97 %, at least 98 %, or at least 99 % identity thereto. In some embodiments, the Ig binding protein comprises one or more domains, wherein at least one domain comprises or essentially consists of or consists of an amino acid sequence of any of SEQ ID NOs: 4-36 and 40-49, or an amino acid with at least 95 %, at least 96 %, at least 97 %, at least 98 %, or at least 99 % identity to any of SEQ ID NOs: 4-36 and 40-49.

**[0063] Affinity to Immunoglobulin.** All Ig binding proteins or Ig binding domains as described herein bind to Immunoglobulin with a dissociation constant  $K_D$  preferably below 200 nM, or below 100 nM, even more preferably 10 nM or less. In some embodiments, the Ig binding protein or Ig binding domain binds to IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>4</sub>, IgM, IgA, Ig fragments, Fc fragments, Fab fragments, fusion proteins comprising an Ig region, and conjugates comprising an Ig region with a dissociation constant  $K_D$  preferably below 200 nM, or below 100 nM, even more preferably 10 nM or less. Methods for determining binding affinities or binding capacities of Ig binding proteins or domains, i.e. for determining the dissociation constant  $K_D$ , are known to a person of ordinary skill in the art and can be selected for instance from the following methods known in the art: Surface Plasmon Resonance (SPR) based technology, kinetic exclusion analysis (KinExA assay), Bio-layer interferometry (BLI), enzyme-linked immunosorbent assay (ELISA), flow cytometry, isothermal titration calorimetry (ITC), analytical ultracentrifugation, radioimmunoassay (RIA or IRMA) and enhanced chemiluminescence (ECL). Some of the methods are described further in the Examples. Typically, the dissociation constant  $K_D$  is determined at 20 °C, 25 °C, or 30 °C. If not specifically indicated otherwise, the  $K_D$  values recited herein are determined at 22 °C +/- 3 °C by surface plasmon resonance spectroscopy.

**[0064]** In one embodiment, the Ig binding protein has a dissociation constant  $K_D$  to human IgG<sub>1</sub> in the range between 0.1 nM and 100 nM, preferably between 0.1 nM and 50 nM.

**[0065] High alkaline stability of Ig binding proteins.** The Ig binding proteins and Ig binding domains of the invention provide surprisingly particularly good alkaline stability, as shown in the Examples and in the Figures, in addition to high dynamic binding capacities (DBC). The alkaline stability of the Ig binding protein or Ig binding domain is determined by comparing the

loss in Ig binding activity. In some embodiments, the alkaline liquid comprises 0.1 - 1.0 M NaOH or KOH, preferably 0.25 - 0.5 M NaOH or KOH. Due to the high alkaline stability of the Ig binding proteins and Ig binding domains of the invention, an alkaline liquid with pH higher than 13 can be used for cleaning affinity matrices with immobilized Ig binding proteins or immobilized Ig binding domains of the invention. In some embodiments, the alkaline stability of the Ig binding protein or Ig binding domain is determined by comparing the loss in Ig binding activity after at least 20 h incubation in 0.5 M NaOH (see **FIGURE 2**, **FIGURE 3**, **FIGURE 4** and **Examples**). In some embodiments, the alkaline stability of the Ig binding protein or Ig binding domain is determined by comparing the loss in Ig binding activity after very long incubation in alkaline solution, e.g. for at least 2 days (at least 48 h) incubation in 0.5 M NaOH (see **Examples**), reflecting an extraordinary stability of the Ig binding proteins as described herein.

**[0066]** The Ig binding proteins and Ig binding domains of the invention are stable under alkaline conditions, in particular under alkaline conditions of 0.5 M NaOH for at least 20 h. In various embodiments, the Ig binding protein or Ig binding domain of the invention is stable under alkaline conditions, in particular under alkaline conditions of 0.5 M NaOH for at least 24 h. In preferred embodiments, the Ig binding protein or Ig binding domain of the invention is stable under alkaline conditions, in particular under alkaline conditions of 0.5 M NaOH for at least 48 h, more preferably for at least 50 h.

**[0067]** The Ig binding proteins and Ig binding domains of the invention are alkaline-stable ligands for immunoglobulins. The Ig binding proteins and Ig binding domains of the invention retain binding capacity (or binding affinity) for immunoglobulin after exposure to 0.5 M NaOH for at least 20 h. As further described herein, the Ig binding proteins and Ig binding domains of the invention retain at least 85 % or at least 90 % binding capacity for immunoglobulin after exposure to alkaline conditions as described herein. In further preferred embodiments, the Ig binding proteins and Ig binding domains of the invention retain at least 94% binding capacity for immunoglobulin after exposure to alkaline conditions (0.5 M NaOH for at least 20 h or 24 h), preferably after exposure to 0.5 M NaOH for at least 48 h, more preferably after exposure to 0.5 M NaOH for at least 50 h. In various embodiments, the Ig binding proteins and Ig binding domains of the invention retain binding capacity for immunoglobulin as described above when immobilized to a solid support, preferably to a solid support of an affinity separation matrix.

**[0068]** As further described herein, the Ig binding proteins and Ig binding domains of the invention are typically stable under alkaline conditions at room temperature. The term room temperature may include temperatures between 15°C and 25°C, more specifically temperatures between 20°C and 25°C. In various embodiments, the Ig binding protein or Ig binding domain of the invention is stable under alkaline conditions at 22°C ± 3°C.

**[0069]** In various embodiments, the alkaline stability of the Ig binding protein or Ig binding domain as described above means alkaline stability of the Ig binding protein or Ig binding domain immobilized to a solid support, preferably to a solid support of an affinity separation matrix. Hence, in various embodiments, the alkaline stability of the Ig binding protein or Ig binding domain is determined by comparing the loss in Ig binding activity or Ig binding capacity

of the Ig binding protein or Ig binding domain when immobilized to a solid support, preferably to a solid support of an affinity separation matrix. Hence, in other embodiments, the alkaline stability of the Ig binding protein or Ig binding domain is determined by comparing the Ig binding activity of the Ig binding protein or Ig binding domain to a reference protein after alkaline treatment for a prolonged time when immobilized to a solid support (see **FIGURES 2, 3, 4**).

**[0070]** The binding capacity or binding affinity for immunoglobulin of the Ig binding protein or Ig binding domain of the present invention can be evaluated by a skilled person using methods well known in the art, in particular methods for determining the dissociation constant  $K_D$  as described elsewhere herein. In various embodiments, the binding capacity or binding affinity for immunoglobulin of the Ig binding protein or Ig binding domain of the present invention is determined using Surface Plasmon Resonance (SPR) spectroscopy, as also described elsewhere herein. In other embodiments, the binding capacity or binding affinity for immunoglobulin of the Ig binding protein or Ig binding domain of the present invention is determined using kinetic exclusion analysis (KinExA assay), or enzyme-linked immunosorbent assay (ELISA), as described elsewhere herein.

**[0071]** The binding capacity or binding affinity for immunoglobulin of the Ig binding protein or Ig binding domain of the present invention can be assessed for each candidate ligand before and after exposure to alkaline conditions as described herein.

**[0072] Multimers.** In one embodiment, the Ig binding protein comprises 1, 2, 3, 4, 5, or 6 Ig binding domains linked to each other, i.e. the Ig binding protein can be, for example, a monomer, a dimer, a trimer, a tetramer, a pentamer, or a hexamer. A multimer may comprise two, three, four, or even more binding domains. Multimers of the invention are fusion proteins generated artificially, generally by recombinant DNA technology well-known to a skilled person.

**[0073]** In some embodiments, the multimer is a homo-multimer, e.g. the amino acid sequences of all Ig binding domains of the Ig binding protein are identical. In some embodiments, the multimer is a hetero-multimer, e.g. at least one Ig binding domain has a different amino acid sequence than the other Ig binding domains within the Ig-binding protein.

**[0074]** A multimer may comprise two or more Ig binding domains, wherein said Ig binding domains preferably comprise or essentially consist of an amino acid sequence as described above. In some embodiments, the multimer is a dimer. The present invention provides dimers comprising monomers of any of SEQ ID NOs: 4-36 and 40-49. Further disclosed herein is an Ig binding protein, which is a dimer comprising two Ig binding domains, wherein each of the two Ig binding domains corresponds to an Ig binding protein having at least 80 % amino acid identity to any one of SEQ ID NOs: 4-36 and 40-49, wherein the dimeric Ig binding protein is stable under alkaline conditions. As further described herein, an Ig binding protein of the present disclosure may be a dimer comprising two Ig binding domains, wherein each of the two Ig binding domains corresponds to an Ig binding protein having at least 80 % amino acid identity to any one of SEQ ID NOs: 4-9, 20-36, and 40-49, wherein the amino acid

corresponding to position 8 of SEQ ID NOs: 4-9, 20-36, and 40-49 is isoleucine (I) or leucine (L) or valine (V) or an aromatic amino acid (Y, F, or W), and wherein the dimeric Ig binding protein is stable under alkaline conditions of 0.5 M NaOH for at least 20 h. As further described herein, a dimeric Ig binding protein of the present disclosure may comprise an Ig binding domain corresponding to an Ig binding protein having at least 80 % amino acid identity to any one of SEQ ID NOs: 4-5, 20-36, and 40-49, wherein the amino acid corresponding to position 8 of SEQ ID NOs: 4-5, 20-36 and 40-49 is isoleucine (I), and wherein the dimeric Ig binding protein is stable under alkaline conditions of 0.5 M NaOH for at least 20 h. As further described herein, a dimeric Ig binding protein of the present disclosure may comprise an Ig binding domain corresponding to an Ig binding protein having at least 80 % amino acid identity to any one of SEQ ID NOs: 4-5, 20-36, and 40-49, wherein the amino acid corresponding to position 8 of SEQ ID NOs: 4-5, 20-36 and 40-49 is isoleucine (I), and wherein the dimeric Ig binding protein is stable under alkaline conditions of 0.5 M NaOH for at least 20 h and wherein the dimeric Ig binding protein allows elution of the target at mild elution conditions of an pH of at least 4.0. As further described herein, the N-terminal Ig binding domain of a dimer comprising two Ig binding domains may have a cysteine (C) at the position corresponding to position 43 or 46 of SEQ ID NO: 1 or any of SEQ ID NOs: 4-36 and 40-49, respectively.

**[0075]** Disclosed herein is an Ig binding protein that comprises two Ig binding domains, wherein one Ig binding domain corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 4, and the other Ig binding domain corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 5. The domain based on SEQ ID NO: 5 may be located upstream of the domain based on SEQ ID NO: 4. As further described herein, an Ig binding protein of the present disclosure may comprise two Ig binding domains, wherein one Ig binding domain corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 43, and the other Ig binding domain corresponds to an Ig binding protein having at least 80 % amino acid identity to SEQ ID NO: 46. The domain based on SEQ ID NO: 46 may be located upstream of the domain based on SEQ ID NO: 43.

**[0076]** The Ig binding protein may be a dimer comprising the sequence of SEQ ID NO: 37, SEQ ID NO: 50, or SEQ ID NO: 51.

**[0077] Linker.** In various embodiments, the one or more Ig binding domains are directly linked to each other. In other embodiments, the one or more Ig binding domains are linked to each other with one or more linkers. Preferred in these typical embodiments are peptide linkers. This means that the peptide linker is one or more amino acids, e.g. an amino acid sequence, that connects a first Ig binding domain with a second Ig binding domain. The peptide linker is connected to the first Ig binding domain and to the second Ig binding domain by a peptide bond between the C-terminal and N-terminal ends of the domains, thereby generating a single, linear polypeptide chain. The length and composition of a linker may vary between at least one and up to about 30 amino acids. More specifically, a peptide linker has a length of between 1 and 30 amino acids; e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 amino acids. It is preferred that the amino acid sequence of the peptide linker is stable against caustic conditions and proteases. Linkers should not

destabilize the conformation of the domains in the Ig binding protein. Well-known are linkers that comprise or consist of small amino acids such as glycine and serine. The linkers can be glycine-rich (e.g., more than 50% of the residues in the linker can be glycine residues). Also preferred are linkers that comprise further amino acids. Other embodiments of the invention comprise linkers consisting of alanine, proline, and serine. Other linkers for the fusion of proteins are known in the art and can be used. In some embodiments, the multimer of Ig binding proteins comprises one or more linkers connecting the Ig binding domains wherein the linkers are identical or different.

**[0078] Non-Ig binding proteins.** The Ig binding protein as described above may further comprises at least one further polypeptide distinct from the Ig binding protein or Ig binding domain as disclosed. The further polypeptide distinct from the Ig binding protein or Ig binding domain as disclosed herein might be a non-Ig-binding protein, for example but not limited to, a protein that does not bind to the Fc part of immunoglobulin. Accordingly, disclosed herein are fusion proteins comprising one or two or more Ig binding protein(s) thereof as disclosed herein and one or two or more non-Ig-binding polypeptide(s). A fusion protein may comprise one (or more) Ig binding protein(s) and/or one (or more) Ig binding domain(s) as disclosed herein fused to one (or more) non-Ig binding protein(s).

**[0079]** As described herein, a non-Ig binding protein may have at least 89.5 % or at least 91 % or at least 93 % or at least 95 % or at least 96 % or at least 98 % or 100 % identity to SEQ ID NO: 38 or SEQ ID NO: 39. SEQ ID NO: 39 is a non-Ig binding protein. SEQ ID NO: 39 has the same basic scaffold as SEQ ID NO: 1 but with modifications D8I, F13D, Y14K, I31R, L42A in SEQ ID NO: 1 which lead to the non-Ig binding property of SEQ ID NO: 39. The modification in position 8 to isoleucine results in improved biochemical properties such as high stability in the non-Ig binding protein, for example, in high stability under alkaline conditions. Accordingly, an isoleucine in a position corresponding to position 8 results in higher stability of proteins having a similar triple-helical structure as present in SEQ ID NO: 1, regardless of the function (i.e. Ig binding protein or non-Ig binding protein). Accordingly, the further disclosed herein is a protein comprising one or more domains, wherein at least one domain corresponds to a protein having at least 70 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I). The protein comprising one or more domains, wherein at least one domain corresponds to a protein having at least 70 % amino acid identity to SEQ ID NO: 1 (cs26), wherein the amino acid corresponding to position 8 of SEQ ID NO: 1 is isoleucine (I), is stable under alkaline conditions.

**[0080]** In accordance with the above, disclosed herein is a non-Immunoglobulin (Ig) binding protein comprising an amino acid sequence having at least 70 %, preferably at least 80 %, more preferably at least 85 %, even more preferably at least 89.5 %, identity to SEQ ID NO: 1, wherein the non-Ig binding protein comprises:

1. (i) an amino acid selected from any one of isoleucine (I), leucine (L), valine (V), or an aromatic amino acid (Tyr, Phe, or Trp), preferably any one of isoleucine (I), leucine (L), or an aromatic amino acid (Y, F, W), more preferably isoleucine (I) or leucine (L), even

- more preferably isoleucine (I), at the position corresponding to position 8 of SEQ ID NO: 1;
2. (ii) an aspartate (D) at the position corresponding to position 13 of SEQ ID NO: 1;
  3. (iii) a lysine (K) or serine (S) at the position corresponding to position 14 of SEQ ID NO: 1;
  4. (iv) an arginine (R) at the position corresponding to position 31 of SEQ ID NO: 1; and (v) an alanine (A) or leucine (L) at the position corresponding to position 42 of SEQ ID NO: 1. In some embodiments, the non-Ig binding protein is stable under alkaline conditions of 0.5 M NaOH for at least 20 h.

**[0081]** Further disclosed herein is a non-Ig binding protein having at least 89.5 % or at least 91 % or at least 93 % or at least 95 % or at least 96 % or at least 98 % or 100 % sequence identity to the amino acid sequence of SEQ ID NO: 38 or SEQ ID NO: 39, wherein the non-Ig binding protein comprises:

1. (i) an amino acid selected from any one of isoleucine (I), leucine (L), valine (V), or an aromatic amino acid (Tyr, Phe, or Trp), preferably any one of isoleucine (I), leucine (L), or an aromatic amino acid (Y, F, W), more preferably isoleucine (I) or leucine (L), even more preferably isoleucine (I), at the position corresponding to position 8 of SEQ ID NO: 38 or 39;
2. (ii) an aspartate (D) at the position corresponding to position 13 of SEQ ID NO: 38 or 39;
3. (iii) a lysine (K) or a serine (S) at the position corresponding to position 14 of SEQ ID NO: 38 or 39;
4. (iv) an arginine (R) at the position corresponding to position 31 of SEQ ID NO: 38 or 39; and
5. (v) an alanine (A) or a leucine (L) at the position corresponding to position 42 of SEQ ID NO: 38 or 39. In some embodiments, the non-Ig binding protein is stable under alkaline conditions of 0.5 M NaOH for at least 20 h.

**[0082]** Affinity separation matrix. The present invention is further directed to an affinity separation matrix, comprising an Ig binding protein or Ig binding domain described herein.

**[0083]** In preferred embodiments, the affinity separation matrix is a solid support. The affinity separation matrix comprises at least one Ig binding protein or Ig binding domain as described above.

**[0084]** An affinity matrix is useful for separation of immunoglobulins and should retain the Ig binding property even after highly alkaline conditions as applied during cleaning processes. Such cleaning of matrices is essential for long-term repeated use of matrices.

**[0085]** Solid support matrices for affinity chromatography are known in the art and include for

example but are not limited to, agarose and stabilized derivatives of agarose (e.g. Praesto<sup>®</sup>Pure, Praesto<sup>®</sup> Jetted A50, Mabselect<sup>®</sup>, PrismA<sup>®</sup>, Sepharose 6B, CaptivA<sup>®</sup>, rPROTEIN A Sepharose Fast Flow, and other), cellulose or derivatives of cellulose, controlled pore glass (e.g. ProSep<sup>®</sup> vA resin), monolith (e.g. CIM<sup>®</sup> monoliths), silica, zirconium oxide (e.g. CM Zirconia or CPG<sup>®</sup>), titanium oxide, or synthetic polymers (e.g. polystyrene such as Poros 50A or Poros MabCapture<sup>®</sup> A resin, polyvinylether, polyvinyl alcohol, monodisperse polyacrylate resin (e.g. UniMab<sup>™</sup>, UniMab<sup>™</sup>Pro), polyhydroxyalkyl acrylates, polyhydroxyalkyl methacrylates, polyacrylamides, polymethacrylamides etc) and hydrogels of various compositions. In certain embodiments the support comprises a polyhydroxy polymer, such as a polysaccharide. Examples of polysaccharides suitable for supports include but are not limited to agar, agarose, dextran, starch, cellulose, pullulan, etc, and stabilized variants of these.

**[0086]** The formats for solid support matrices can be of any suitable well-known kind. Such solid support matrix for coupling the Ig binding protein or Ig binding domain as described herein might comprise for example, one of the following: columns, capillaries, particles, membranes, filters, monoliths, fibers, pads, gels, slides, plates, cassettes, or any other format commonly used in chromatography and known to someone skilled in the art.

**[0087]** In one embodiment, the matrix is comprised of substantially spherical particles, also known as beads, for example Sepharose or Agarose beads or monodisperse polyacrylate beads. Suitable particle sizes may be in the diameter range of 5-500  $\mu\text{m}$ , such as 10-100  $\mu\text{m}$ , such as 20-80  $\mu\text{m}$ , such as 40-70  $\mu\text{m}$ . Matrices in particle form can be used as a packed bed or in a suspended form including expanded beds.

**[0088]** In an alternative embodiment, the solid support matrix is a membrane, for example a hydrogel membrane. In some embodiments, the affinity purification involves a membrane as matrix to which the Ig binding protein or Ig binding domain of the one embodiment is covalently bound. The solid support can also be in the form of a membrane in a cartridge.

**[0089]** In some embodiments, the affinity purification involves a chromatography column containing a solid support matrix to which the Ig binding protein or Ig binding domain of the one embodiment is covalently bound.

**[0090] Immobilization to a solid support.** In embodiments of the invention, the Ig binding protein or Ig binding domain is conjugated to a solid support. In some embodiments of the invention, the Ig binding protein or Ig binding domain may comprise additional amino acid residues at the N-and/or C-terminal end. The Ig binding protein or Ig binding domain of the invention may be attached to a suitable solid support matrix via conventional coupling techniques. Methods for immobilization of protein ligands to solid supports are well-known in this field and easily performed by the skilled person in this field using standard techniques and equipment. In some embodiments, the coupling may be a multipoint coupling, for example via several lysines, or a single point coupling, for example via cysteine.

**[0091]** In some embodiments, the alkaline stable Ig binding protein or Ig binding domain comprises an attachment site for covalent attachment to a solid phase (matrix). Site-specific attachment sites comprise natural amino acids, such as cysteine or lysine, which enable specific chemical reactions with a reactive group of the solid phase or a linker between the solid phase and the protein.

**[0092]** In some embodiments, the attachment site may be directly at the C- or N-terminal end of the Ig binding protein or Ig binding domain. In some embodiments, a single cysteine is located at the C-terminal end for site-specific immobilization of the Ig binding protein or Ig binding domain. An advantage of having a C-terminal cysteine is that coupling of the Ig binding protein or Ig binding domain can be achieved through reaction of the cysteine thiol with an electrophilic group on a support resulting in a thioether bridge coupling. This provides excellent mobility of the coupled protein which provides increased binding capacity.

**[0093]** In other embodiments, the attachment site may be located in the third helix of the Ig binding protein or Ig binding domain, for example, in position corresponding to position 43 or position 46 of SEQ ID NO: 1.

**[0094]** In other embodiments, there may be a linker between the N- or C-terminus and the attachment site. In some embodiments of the invention, the Ig binding protein or Ig binding domain may comprise a N- or C-terminal amino acid sequence of 3 - 20 amino acids, preferably of 4 - 10 amino acids, with a terminal cysteine. Amino acids for a terminal attachment site may be selected from the group of proline, glycine, alanine, and serine, with a single cysteine at the C-terminal end for coupling.

**[0095]** In some embodiments of the invention, the Ig binding protein or Ig binding domain may also comprise additional amino acid residues at the N- and/or C-terminal end, such as for example a leader sequence at the N-terminal end and/or a coupling sequence with or without a tag at the N- or C-terminal end.

**[0096] Use of the Ig binding protein.** The present invention is further directed to the use of the Ig binding protein or Ig binding domain described herein, or an affinity matrix described herein, for affinity purification of immunoglobulins or variants thereof, i.e. the Ig binding protein or Ig binding domain of the invention is used for affinity chromatography. In some embodiments, the Ig binding protein or Ig binding domain of the invention is immobilized onto a solid support as described herein,

**[0097] Method of affinity purification of immunoglobulins.** The present invention is further directed to a method of affinity purification of immunoglobulins, the method comprising the following steps:

1. (a) providing a liquid (sample) that contains an Ig such as IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>4</sub>, IgM, IgA, Ig fragments, Fc fragments, or Fab fragments (including fusion proteins and conjugates, as defined above);

2. (b) providing an affinity separation matrix comprising an immobilized Ig binding protein or Ig binding domain as described above immobilized to said affinity separation matrix;
3. (c) contacting said liquid with said affinity separation matrix, under conditions that permit binding of the at least one Ig binding protein or Ig binding domain as described above to an Ig; and
4. (d) eluting said Ig from said matrix, thereby obtaining an eluate containing said Ig.

**[0098]** In some embodiments, the method of affinity purification may further comprise one or more washing steps carried out between steps (c) and (d) under conditions sufficient to remove from the affinity separation matrix some or all molecules that are non-specifically bound thereto. Non-specifically bound means any binding that does not involve an interaction between the at least one Ig binding protein or Ig binding domain and an Ig.

**[0099]** Affinity separation matrices suitable for the disclosed uses and methods are those matrices according to the embodiments described above and as known to someone skilled in the art.

**[0100]** The elution of the immunoglobulin from (the matrix comprising) the Ig binding protein or Ig binding domain in step (d) may be effected through a change in pH and/or a change in salt concentration. In general, suitable conditions for performing the method of affinity purification are well known to someone skilled in the art.

**[0101]** The disclosed uses or methods of affinity purification comprising the disclosed Ig binding proteins or Ig binding domains may provide elution of at least about 95 %, at least about 96 %, at least about 97 %, at least about 98 %, at least about 99 %, or 100 % of Ig containing proteins at a pH of greater than or equal to 3.7 (e.g., about pH 4.0, about pH 4.5, about pH 5.0, or about pH 5.5). Due to the high stability of the Ig binding proteins and Ig binding domains of the invention, solutions with greater than or equal to pH 3.7 can be used for the elution of Ig proteins (see **Examples**).

**[0102]** In step (d) of the method of affinity purification more than 95 % of the protein comprising the Ig sequence (e.g. antibody) is eluted at pH 3.7 or higher (up to about pH 5.5) from the matrix comprising the immobilized Ig binding protein or Ig binding domain as described above. A further step (e) for efficient cleaning of the affinity matrix may be added, in particular by using an alkaline liquid, for example, with pH of 13 - 14. As further described herein, the cleaning liquid may comprises 0.1 - 1.0 M NaOH or KOH, in particular 0.25 - 0.5 M NaOH or KOH. Due to the high alkaline stability of the Ig binding proteins or Ig binding domains of the invention, such strong alkaline solution can be used for cleaning purposes. After cleaning the affinity purification matrix with an alkaline cleaning liquid, at least 88 % of the Ig binding protein or Ig binding domain may have Ig binding activity if incubated for at least 48 h at 0.5 M NaOH. The Ig-binding capacity of the Ig binding protein or Ig binding domain may be at least about 80%, at least about 90 %, or 100 % of the Ig binding capacity before the

incubation under alkaline conditions, for example, as determined by the remaining Ig-binding capacity after at least 20 h incubation in 0.5 M NaOH.

**[0103]** The present invention further relates to a method of isolating an immunoglobulin, comprising the steps (a) contacting a liquid sample comprising an immunoglobulin with a separation matrix comprising a plurality of Ig binding proteins or Ig binding domains described herein (coupled to a solid support); (b) washing the separation matrix with a washing liquid at a pH above pH 3.7 (up to pH 5.5); (c) eluting the immunoglobulin from the separation matrix; and (d) obtaining the immunoglobulin.

**[0104] Nucleic acid molecule.** Further disclosed herein is a nucleic acid molecule, in particular an isolated nucleic acid molecule, encoding an Ig binding protein or Ig binding domain as disclosed above. Further disclosed herein is a vector comprising the nucleic acid molecule. A vector means any molecule or entity (e.g., nucleic acid, plasmid, bacteriophage or virus) that can be used to transfer protein coding information into a host cell. In one embodiment, the vector is an expression vector.

**[0105]** Further disclosed herein is an expression system which comprises a nucleic acid or a vector as disclosed above, for example a prokaryotic host cell, for example *E. coli*, or a eukaryotic host, for example yeast *Saccharomyces cerevisiae* or *Pichia pastoris* or mammalian cells such as CHO cells.

**Method for the production of a Ig binding protein.**

**[0106]** Further disclosed herein is a method for the production of an Ig binding protein or Ig binding domain of the invention, comprising the step(s): (a) culturing the host cell of the one embodiment under suitable conditions for the expression of the binding protein or Ig binding domain in order to obtain said Ig binding protein or Ig binding domain; and (b) optionally isolating said Ig binding protein or Ig binding domain. Suitable conditions for culturing a prokaryotic or eukaryotic host are well-known to the person skilled in the art.

**[0107]** Ig binding molecules of the invention may be prepared by any of the many conventional and well-known techniques such as plain organic synthetic strategies, solid phase-assisted synthesis techniques or by commercially available automated synthesizers. On the other hand, they may also be prepared by conventional recombinant techniques alone or in combination with conventional synthetic techniques.

**[0108] Further disclosed herein is** a method for the preparation of a Ig binding protein or Ig binding domain according to the invention as detailed above, said method comprising the following steps: (a) preparing a nucleic acid encoding an Ig binding protein or Ig binding domain as defined above; (b) introducing said nucleic acid into an expression vector; (c) introducing said expression vector into a host cell; (d) cultivating the host cell; (e) subjecting the host cell to culturing conditions under which an Ig binding protein or Ig binding domain is

expressed, thereby (e) producing an Ig binding protein or Ig binding domain as described above; optionally (f) isolating the Ig binding protein or Ig binding domain produced in step (e); and (g) optionally conjugating the Ig binding protein or Ig binding domain to solid matrices as described above. The production of the Ig binding protein or Ig binding domain may be performed by cell-free *in vitro* transcription / translation.

## EXAMPLES

[0109] The following Examples are provided for further illustration of the invention. The invention, however, is not limited thereto, and the following Examples merely show the practicability of the invention on the basis of the above description.

### Example 1. Generation of Ig binding proteins of the invention

[0110] Modifications were introduced in SEQ ID NO: 1 (cs26) by site-saturation mutagenesis of individual amino acid residues. The following exchanges were introduced and analyzed for properties in affinity chromatography: 8I, 8F, 8Y, 8W, 8L, 8V, 6I, 6Y, 6W, 6L, 4I, 4F, 4W, 4Y, 4L; position 43 was exchanged to C (cysteine) for coupling to the matrix.

### Example 2. Expression of Ig binding proteins

[0111] BL21 (DE3) competent cells were transformed with an expression plasmid encoding Ig binding proteins (for example, 203524 and 203550). Cells were spread onto selective agar plates (Kanamycin) and incubated either for 2 days at 21 °C or overnight at 37°C. Precultures were inoculated from single colony in 50 ml 2xYT medium supplemented with 50 µg/ml kanamycin and cultured for 17 hours at 37 °C at 210 rpm in a conventional orbital shaker in 250 mL Erlenmeyer flasks. The OD<sub>600</sub> readout should be in the range of 3.5-6. Main cultures were inoculated from previous overnight culture with an adjusted start-OD<sub>600</sub> of 0.3 in 300 ml superrich medium (modified H15 medium consisting of 2% glucose, 5% yeast extract, 0.89% glycerol, 0,76% lactose, 250 mM MOPS, 202 mM TRIS, 10 mM MgSO<sub>4</sub>, pH 7.4, antifoam SE15) that was supplemented with 50 µg/ml Kanamycin and trace elements (see Studier 2005) in 1 L thick-walled Erlenmeyer flasks. Cultures were transferred to a resonant acoustic mixer (RAM<sub>bio</sub>) and incubated at 37 °C with 20 x g. Aeration was facilitated by Oxy-Pump stoppers. Recombinant protein expression was induced by metabolizing glucose and subsequently allowing lactose to enter the cells. Cells were grown overnight for approx. 18 hours to reach a final OD<sub>600</sub> of about 35-55. Before the harvest, the OD<sub>600</sub> was measured, samples adjusted to 0.6/OD<sub>600</sub> were withdrawn, pelleted and frozen at -20 °C. To collect biomass cells were centrifuged at 12000 x g for 20 min at 20 °C. Pellets were weighed (wet weight). Cells were stored at -20 °C before processing.

**Example 3: SDS-PAGE Analysis of expression and solubility of Ig binding proteins**

[0112] Samples were resuspended in 90 µl extraction buffer (PBS supplemented with 0.2 mg/ml Lysozyme, 0.5x BugBuster, 6 mM MgSO<sub>4</sub>, 6 mM MgCl<sub>2</sub>, 15 U/mL Benzonase) and solubilized by agitation in a thermomixer at 850 rpm, rt for 15 min with a subsequent incubation at -80 °C for 15 min. After thawing, soluble proteins were separated from insoluble proteins by centrifugation (16000 x g, 2 min, rt). Supernatant was withdrawn (soluble fraction) and the pellet (insoluble fraction) was resuspended in equivalent amount of urea buffer (8 M urea, 0.2 M Tris, 20 mM EDTA, pH 7.0). 35 µl were taken both from the soluble and insoluble fraction, and 10 µl 5x sample buffer as well as 5 µl 0.5 M DTT were added. Samples were boiled at 95 °C for 5 min. Finally, 5 µl of those samples were applied to NuPage Novex 4-12 % Bis-Tris SDS gels which were run in accordance to the manufacturer's recommendations and stained with Coomassie. **Results:** High level expression was found under optimized conditions within the chosen period of time. All expressed Ig binding proteins were 100% soluble according to SDS-PAGE.

**Example 4: Purification of Ig binding proteins**

[0113] Ig binding proteins were expressed in the soluble fraction of *E. coli*. The cells were resuspended in cell disruption buffer and lysed by an ultrasonic cell disruption system (Sonopuls HD 2200, Bandelin). Purification step was performed with IEC Sepharose SP-HP (GE Healthcare) using an ÄKTAvant system (Ge Healthcare) according to the manufacturer's instructions using citric acid buffer at pH 3.0 (20 mM Citric acid, 1 mM EDTA, pH 3.0). Pure protein fractions were eluted by increasing sodium chloride concentration to 1 M with a linear gradient in 10 column volumes. Further purification was performed by size exclusion chromatography (Superdex 75) according to manufacturer's instructions using citric acid buffer pH 6.0 (20 mM Citric acid, 150 mM NaCl, 1 mM EDTA, pH 6.0). **Results:** The purity of variants 8I and 6L was > 95% after SE-HPLC and >90 % after RP HPLC.

**Example 5. The Ig binding proteins bind to IgG with high affinities (SPR)**

[0114] A CM5 sensor chip (GE Healthcare) was equilibrated with surface plasmon resonance (SPR) running buffer. Surface-exposed carboxylic groups were activated by passing a mixture of EDC and NHS to yield reactive ester groups. 700-1500 RU on-ligand were immobilized on a flow cell, off- ligand was immobilized on another flow cell. Injection of ethanolamine after ligand immobilization removes non-covalently bound Ig binding protein. Upon ligand binding, protein analyte was accumulated on the surface increasing the refractive index. This change in the refractive index was measured in real time and plotted as response or resonance units (RU) versus time. The analytes were applied to the chip in serial dilutions with a suitable flow rate

( $\mu\text{l}/\text{min}$ ). After each run, the chip surface was regenerated with regeneration buffer and equilibrated with running buffer. The control samples were applied to the matrix. Regeneration and re-equilibration were performed as previously mentioned. Binding studies were carried out by the use of the Biacore<sup>®</sup> 3000 (GE Healthcare) at 25 °C; data evaluation was operated via the BIAevaluation 3.0 software, provided by the manufacturer, by the use of the Langmuir 1:1 model ( $RI=0$ ). Evaluated dissociation constants ( $K_D$ ) were standardized against *off-target* and  $K_D$  values of Ig binding proteins for Cetuximab (IgG<sub>1</sub>), Natalizumab (IgG<sub>4</sub>), and Panitumab (IgG<sub>2</sub>) in Table 1.

**Table 1.  $K_D$  values of Ig binding proteins for IgG**

CID	Variant	vs. hlgG <sub>1</sub> (nM)	vs. hlgG <sub>2</sub> (nM)	vs. hlgG <sub>4</sub> (nM)
203524	D6L	4.2	110	4
203550	D8I	2.9	20.6	2.4
203704	N28H	17.2	107	20
203447	cs26	3.6	23.2	2.7

**Example 6. Ig binding proteins coupled to agarose-based chromatography beads Praesto™ Pure85 - coupling efficiencies, DBC10 %, elution**

**[0115] DBC10%:** Purified Ig binding proteins were coupled to agarose-based chromatography beads (Praesto™ Pure85, Purolite; Cat. No. PR01265-164) according to the manufacturer's instructions (coupling conditions: pH 9.5, 3 hours, 35 °C, 4.1 M NaSO<sub>4</sub>, blocking overnight with ethanolamine). Coupled Resin was packed into super compact 5/50 column (Götec GmbH). Polyclonal human IgG Gammanorm<sup>®</sup> (Ocatpharm) was used as IgG sample (conc. 2,2 mg/ml). Polyclonal hlgG sample was applied in saturated amounts to the matrix comprising immobilized Ig binding protein. **Results:** Variant 8I (203550) shows slightly increased DBC10% in comparison to the parent variant (203447)

**[0116] Elution of Immunoglobulin from matrix:** The matrix was washed with 100 mM acetic acid buffer, pH 3.7 and then with 0.1 M phosphoric acid pH 1.7 to elute hlgG (Load: 2.2 mg/mL Gammanorm, 6 min residence time) that was bound to the immobilized Ig binding protein. **Results:** For all variants tested, more than 99 % of the antibody was eluted (e.g. D8I, D6L), compared to 96% elution if the parent molecule was immobilized; see Table 2.

**Example 7. Alkaline stability of Ig binding proteins coupled to an epoxy-activated matrix**

**[0117]** Columns were incubated with 0.5 M NaOH for 0 h and 20 h at room temperature (22 °C

+/- 3 °C). The Ig binding activity of the immobilized proteins was analyzed after incubation with 0.5 M NaOH. Results are shown in **FIGURE 2, FIGURE 3, and FIGURE 4**. Praesto 85 epoxy resin with immobilized 25 mg/ml variants 6L (203524) and 8I (203550) and controls was incubated with 0.5 M NaOH for 50 h at room temperature (22 °C +/- 3 °C). *Results:* Even after more than 2 days in strong alkaline solution, variant 8I and variant 6L showed 94,4 % and 88,5 %, respectively, remaining binding capacity for Ig. The remaining IgG binding capacity after alkaline treatment for 50 h is improved compared to the parent (caustic stable Ig binding protein of SEQ ID NO: 3; CID203447) (83 % remaining binding capacity for Ig). Results see **Table 2**.

**Table 2. Caustic stability and elution**

CID	Affinity ligand	DBC10% at 6 min residence time (mg/ml)	DBC10 compared to cs26	Caustic stability 50 h, 0.5 M NaOH (%)	Elution recovery at 10 mM acetic acid at pH 3.7
203550	cs26 8I	56.6	103.1	94.4	99.6
203704	cs26 28H	56	102	n.d.	n.d.
203447	cs26	54.9	100	83	99.8

#### Example 8. Elution of hlgG from immobilized ligands

**[0118] Determination of elution pH with pH gradient.** 1 mg/ml hlgG (Gammanorm) in PBS pH 7.3 was injected onto the column; contact time: 6 min. The column was washed with 0.1 M citrate pH 6.0. hlgG that was bound to the immobilized ligand was eluted via pH gradient from pH 6.0 - 2.0. The pH of the eluted main fraction was determined (peak maximum). **Table 3** shows that all variants show a peak maximum at the range of pH 4.2 and pH 5.5, compared to cs26 a peak maximum of pH 3.7 for cs26.

**Table 3. Elution pH (gradient)**

SEQ ID NO:	CID	Substitution	Peak maximum (pH)
3	203447		3.7
60	203561	Q10D	4.6
59	203564	Q10H	5.3
58	203606	Y14H	5.5
57	203634	I16H	5.5
56	203648	L17H	4.9
55	203660	H18E	4.1
54	203702	N28E	4.3
53	203704	N28H	4.2

**[0119] Elution at high pH (pH 4.5).** The matrix was washed with 50 mM acetic acid buffer, pH 4.5 and then 100 mM acetic acid to elute hlgG (Load: 2.2 mg/mL Gammanorm, 6 min residence time) that was bound to the immobilized affinity ligand (variant). **Table 4** shows that all ligands show significantly higher percentage of recovered antibody at pH 4.5 compared to cs26.

**Table 4. Elution recovery (%)**

SEQ ID NO:	CID	ligand	50 mM acetic acid pH 4.5	100 mM citrate pH 4.5
2	184244	cs26	45	n.d.
60	203561	cs26 10D	85	n.d.
59	203564	cs26 10H	85	n.d.
58	203606	cs26 14H	98	n.d.
57	203634	cs26 16H	91	n.d.
56	203648	cs26 17H	82	n.d.
55	203660	cs26 18E	71	n.d.
54	203702	cs26 28E	70	n.d.
53	203704	cs26 28H	76	93.5

**Example 9. Characterization of SEQ ID NO: 51 as ligand for affinity purification of IgG**

**[0120]** Experiments were performed as described above, unless different procedures are mentioned here.

**[0121] Purity:** The purity of affinity ligand of SEQ ID NO: 51 was 100 % after RP HPLC. Protein was detected by 220 nm absorption.

**[0122] Affinity for hlgG<sub>1</sub>:** For the analysis of affinity ligand SEQ ID NO: 51, the monoclonal antibodies Cetuximab (IgG<sub>1</sub>) and Belimumab (IgG<sub>1</sub>) were used as target. The KD of SEQ ID NO: 51 for IgG Cetuximab was 40.8 nM and for IgG<sub>1</sub> Belimumab 47.4 nM.

**[0123] Binding capacity:** Binding capacity was determined with IgG<sub>1</sub> sample. The antibody was injected onto coupled resin with affinity ligand SEQ ID NO: 51 until 10 % target breakthrough at 6 min residence time. Loaded antibody was quantified and calculated as dynamic binding capacity DBC10 %, The DBC10 % at 6 min residence time with 2.2 mg/ml Belimumab compared to cs26 was 104.1% compared to cs26.

**[0124] Caustic stability:** Praesto 85 epoxy resin with immobilized 19.6 mg/ml SEQ ID NO: 51 (coupled at pH=10.5 and 2.05 M Na<sub>2</sub>SO<sub>4</sub>) was incubated with 0.5 M NaOH for 24 h at room

temperature (22 °C +/- 3 °C). Even after 24 h in strong alkaline solution, SEQ ID NO: 51 showed no reduction in binding capacity for Ig (99 %).

**[0125] Elution pH:** Elution pH of peak maximum was determined as described above. SEQ ID NO: 51 showed high elution pH (pH 5.0) for Gammanorm and Belimumab in comparison to cs26 (pH 3.5), see **Table 5**.

**Table 5. Elution of peak maximum.**

Affinity ligand	Elution pH peak maximum;	Elution pH peak maximum;
	Target: Gammanorm	Target: Belimumab
SEQ ID NO: 51	5.0	5.0
cs26	3.5	3.5

**[0126] Step Elution:** Step elution was analyzed for target elution at pH 4.8. Residual elution of the target (Gammanorm or Belimumab) from the ligand (SEQ ID NO: 51) was analyzed after 100 mM phosphoric acid CIP at pH 1.7 (recovery). The target protein was nearly completely eluted from the ligand of SEQ ID NO: 51 at pH 4.8, see **Table 6**.

**Table 6. Target elution at pH 4.8 compared to pH 1.7**

Affinity ligand	Elution-Recovery at pH 4.8 vs. 1.7;	Elution-Recovery at pH 4.8 vs. 1.7;
	Target: Gammanorm	Target: Belimumab
SEQ ID NO: 51	96.9 %	99.7 %
cs26	15.8 %	19.3 %

## REFERENCES CITED IN THE DESCRIPTION

### Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- [EP3650465A1](#) [0003]
- [WO2019152318A1](#) [0003]

**Non-patent literature cited in the description**

- **HUSE et al.**J. Biochem. Biophys. Methods, 2002, vol. 51, 217-231 [0002]
- A multilingual glossary of biotechnological terms: (IUPAC Recommendations)Helvetica Chimica Acta19950000 [0011]

## PATENTKRAV

- 5 1. Et Immunoglobulin (Ig)-bindende protein omfattende et eller flere Ig-bindende domæner, hvor mindst ét Ig-bindende domæne svarer til et Ig-bindende protein med mindst 80 % aminosyreidentitet med SEQ ID NO: 1, hvor aminosyren svarer til position 8 i SEQ ID NO: 1 er isoleucin (I), leucin (L), valin (V) eller en aromatisk aminosyre, som er tryptophan (W) eller phenylalanin (F), og hvor det Ig-bindende protein er stabil under alkaliske betingelser på 0,5 M NaOH i mindst 20 timer.
- 10 2. Ig-bindende protein ifølge krav 1, hvor aminosyren svarende til position 8 i SEQ ID NO: 1 er isoleucin (I).
- 15 3. Ig-bindende protein ifølge krav 1-2, hvor en eller flere aminosyre(r) svarende til position 10, 14, 16, 17, 18 eller 28 i SEQ ID NO: 1 er udvalgt blandt gruppen af histidin (H) eller aspartat (D) eller glutamat (E).
- 20 4. Ig-bindende protein ifølge et hvilket som helst af kravene 1-3, hvor mindst ét domæne omfatter en aminosyresekvens af en hvilken som helst af SEQ ID NOs: 4-9, 20-26 og 40-49, eller en amino. syresekvens med mindst 89,5 % identitet dertil med en hvilken som helst af SEQ ID NO: 4-9, 20-26 og 40-49.
- 25 5. Ig-bindende protein ifølge et hvilket som helst af kravene 1-4, hvor nævnte protein binder til IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>4</sub>, IgM, IgA, Ig-fragmenter, Fc-fragmenter, Fab-fragmenter, fusionsproteiner, der omfatter en Ig-region, og konjugater, der omfatter en Ig-region.
- 30 6. Ig-bindende protein ifølge et hvilket som helst af kravene 1-5, hvor proteinet omfatter 2, 3, 4, 5 eller 6 domæner bundet til hinanden.
- 35 7. Ig-bindende protein ifølge krav 6, hvor proteinet er en homo-multimer eller en hetero-multimer.
- 40 8. Ig-bindende protein ifølge et hvilket som helst af kravene 1-7, hvor proteinet er immobiliseret til en fast bærer.
- 45 9. Affinitetsseparationsmatrix omfattende det Ig-bindende protein ifølge et hvilket som helst af kravene 1 til 8 koblet til nævnte affinitetsseparationsmatrix.
- 50 10. Anvendelse af det Ig-bindende protein ifølge et hvilket som helst af kravene 1 til 8 eller af affinitetsseparationsmatrixen ifølge krav 9 til affinitetsoprensning af et hvilket som helst protein med affinitet til det Ig-bindende protein.
11. Fremgangsmåde til affinitetsoprensning af et protein omfattende en Ig-sekvens, fremgangsmåden omfatter:
  - a. tilvejebringelse af en væske, der indeholder protein omfattende en Ig-sekvens;
  - b. tilvejebringelse af en affinitetsseparationsmatrix ifølge krav 9 omfattende mindst ét Ig-bindende protein ifølge et hvilket som helst af kravene 1 til 8 koblet til affinitetsseparationsmatrixen ifølge krav 9;
  - c. at bringe nævnte affinitetsseparationsmatrix i kontakt med væsken under betingelser, der tillader binding af det mindst ene Ig-bindende protein ifølge et hvilket som helst af kravene 1-8 til et protein, der omfatter en Ig-sekvens; og

d. eluering af nævnte protein omfattende en Ig-sekvens fra affinitetsoprensningmatrixen, hvorved der opnås et eluat indeholdende immunoglobulinet.

- 5 12. Fremgangsmåde ifølge krav 11, hvor i trin (d), hvor mere end 95 % af proteinet omfattende Ig-sekvensen elueres ved pH 3,7 eller højere fra affinitetsoprensningmatrixen omfattende det Ig-bindende protein ifølge et hvilket som helst af kravene 1- 8.
- 10 13. Fremgangsmåde ifølge krav 12, hvor i trin (d) mere end 95 % af proteinet omfattende Ig-sekvensen elueres ved pH 4,5 eller højere fra affinitetsoprensningmatrixen omfattende det Ig-bindende protein ifølge et hvilket som helst af kravene 1- 8.
- 15 14. Fremgangsmåden ifølge et hvilket som helst af kravene 11-13, omfattende det yderligere trin (e) rensning af affinitetsoprensningmatrixen med en alkalisk renssevæske.

# DRAWINGS

Drawing

FIGURE 1. Selected novel Ig Binding proteins.

SEQ ID	remarks	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58
4	cs26 8I	I	A	A	Q	H	D	K	I	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
6	cs26 8F	I	A	A	Q	H	D	K	F	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
7	cs26 8Y	I	A	A	Q	H	D	K	Y	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
8	cs26 8W	I	A	A	Q	H	D	K	W	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
9	cs26 8L	I	A	A	Q	H	D	K	L	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
10	cs26 6I	I	A	A	Q	H	I	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
11	cs26 6Y	I	A	A	Q	H	Y	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
12	cs26 6W	I	A	A	Q	H	W	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
14	cs26 6L	I	A	A	Q	H	L	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	C	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
15	cs26 4I	I	A	A	I	H	D	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
16	cs26 4F	I	A	A	F	H	D	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
17	cs26 4W	I	A	A	W	H	D	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
18	cs26 4L	I	A	A	L	H	D	K	D	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
20	cs26 8I 14H	I	A	A	Q	H	D	K	I	Q	Q	A	A	F	H	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
25	cs27 8I	I	A	A	K	F	D	E	I	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	V	L	G	E	A	Q	K	L	N	D	S	Q	A	P	K
26	cs27 8I 14H	I	A	A	K	F	D	E	I	Q	Q	A	A	F	H	E	I	L	H	L	P	N	L	T	E	E	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	V	L	G	E	A	Q	K	L	N	D	S	Q	A	P	K
27	cs27 8I 28H	I	A	A	K	F	D	E	I	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	E	Q	R	H	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	V	L	G	E	A	Q	K	L	N	D	S	Q	A	P	K
28	cs59 8I	I	D	A	K	F	D	E	I	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	D	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	A	L	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
29	cs59 8I 14H	I	D	A	K	F	D	E	I	Q	Q	A	A	F	H	E	I	L	H	L	P	N	L	T	E	D	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	A	L	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
30	cs59 8I 28H	I	D	A	K	F	D	E	I	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	D	Q	R	H	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	A	L	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
31	cs60 8I	I	D	A	K	F	D	E	I	A	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	D	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	A	L	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
32	cs60 8I 14H	I	D	A	K	F	D	E	I	A	Q	A	A	F	H	E	I	L	H	L	P	N	L	T	E	D	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	A	L	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
33	cs60 8I 28H	I	D	A	K	F	D	E	I	A	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	D	Q	R	H	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	A	L	L	A	E	A	K	K	L	N	D	A	Q	A	P	K
34	cs_4 8I	I	A	A	K	H	D	K	I	Q	Q	A	A	F	H	E	I	L	H	L	P	N	L	T	E	D	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	G	E	A	K	K	L	N	D	A	Q	A	P	K
35	cs_4 8I 14H	I	A	A	K	H	D	K	I	Q	Q	A	A	F	H	E	I	L	H	L	P	N	L	T	E	D	Q	R	N	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	G	E	A	K	K	L	N	D	A	Q	A	P	K
36	cs_4 8I 28H	I	A	A	K	H	D	K	I	Q	Q	A	A	F	Y	E	I	L	H	L	P	N	L	T	E	D	Q	R	H	A	F	I	Q	S	L	R	D	D	P	S	V	S	L	E	I	L	G	E	A	K	K	L	N	D	A	Q	A	P	K

FIGURE 2. Remaining activity after NaOH incubation (pos. 8 modifications)




FIGURE 3. Remaining activity after NaOH incubation (pos. 4 modifications)




FIGURE 4. Remaining activity after NaOH incubation (pos. 6 modifications)




FIGURE 5. Elution profile of Belimumab from affinity ligand SEQ ID NO: 51 (pH gradient pH 6.0 to 2.0)





Figure 6. DBC10% determination of affinity ligand SEQ ID NO: 51; the antibody eluted completely at pH 4.8.



SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

