

(19)



(11)

EP 1 404 666 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

27.05.2009 Bulletin 2009/22

(21) Application number: **02747474.1**

(22) Date of filing: **05.07.2002**

(51) Int Cl.:

C07D 401/04 ^(2006.01) **C07D 213/81** ^(2006.01)
A01N 43/40 ^(2006.01) **C07D 405/12** ^(2006.01)
C07D 401/12 ^(2006.01) **C07D 401/14** ^(2006.01)
C07D 405/14 ^(2006.01) **C07D 405/04** ^(2006.01)

(86) International application number:

PCT/EP2002/008665

(87) International publication number:

WO 2003/006456 (23.01.2003 Gazette 2003/04)

(54) **4-SUBSTITUTED-PICOLINIC ACID AMIDE DERIVATIVES USEFUL AS FUNGICIDES**

4-SUBSTITUIERTE-PICOLINSÄUREAMID-DERIVATE VERWENDBAR ALS FUNGIZIDE

DERIVES D'AMIDE D'ACIDE PICOLINIQUE SUBSTITUES EN 4, UTILISES COMME FONGICIDES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
 IE IT LI LU MC NL PT SE SK TR**

(30) Priority: **11.07.2001 FR 0109195**

(43) Date of publication of application:
07.04.2004 Bulletin 2004/15

(73) Proprietor: **Bayer CropScience S.A.**
69009 Lyon (FR)

(72) Inventors:

- **HUTIN, Pierre**
F-69009 Lyon (FR)
- **MULLER, Benoît**
F-69002 Lyon (FR)
- **STEELE, Christopher, Richard**
F-69001 Lyon (FR)
- **PEREZ, Joseph**
F-69009 Lyon (FR)
- **GENIX, Pierre**
F-69004 Lyon (FR)

(74) Representative: **Gerhards, Frank et al**
Bayer CropScience SA
Département Brevets et Licences
14-20 Rue Pierre Baizet
BP 9163
69263 Lyon Cedex 09 (FR)

(56) References cited:

WO-A-01/05769 **WO-A-01/14339**
WO-A-01/49666

- **BRUNNER, HENRIET AL.:** "Asymmetric catalysis. Part 117. Novel chiral oxazoline ligands for potential charge-transfer effects in the Rh (I)-catalyzed enantioselective hydrosilylation" **EUROPEAN JOURNAL OF INORGANIC CHEMISTRY.**, no. 6, 1998, pages 771-781, XP002216185 WILEY-VCH VERLAG, WEINHEIM., DE ISSN: 1434-1948
- **MORKVED, EVA H. ET AL.:** "Potential acyl-transfer agents. Reactions of N-acyl-2-pyridinecarboxamides with nucleophiles" **ACTA CHEMICA SCANDINAVICA.**, vol. 36, no. 6, - 1982 pages 381-388, XP000974283 MUNKSGAARD, COPENHAGEN., DK ISSN: 0904-213X
- **MORKVED, EVA H.:** "Reactions of 4-(1-pyrrolidinyl)pyridine-2-carboxanilide s with acyl halide to give acyl imidate hydrohalides" **ACTA CHEMICA SCANDINAVICA. SERIES B-ORGANIC CHEMISTRY AND BIOCHEMISTRY.**, vol. 33, no. 6, - 1979 pages 433-436, XP002216197 COPENHAGEN DK
- **E.H. MORKVED ET AL.:** "Synthesis of some substituted picolinimidoyl chloride hydrochlorides" **ACTA CHEMICA SCANDINAVICA.**, vol. B32, no. 3, 1978, pages 231-234, XP002216089 MUNKSGAARD, COPENHAGEN., DK ISSN: 0904-213X

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 404 666 B1

Description

[0001] The present invention relates to novel picolinic acid derivatives, the method for preparing them, their use as fungicides, particularly in the form of fungicidal compositions, and a method for controlling phytopathogenic fungi of crops using these compounds or these compositions.

[0002] Picolinic acid derivatives with fungicidal action are known in the literature. Thus, actimycin and some of its derivatives, which are described particularly in patent application WO-A-99/11127 and by Kuzo SHIBATA et al. (The Journal of Antibiotics, 51 (12), (1998), 1113-1116), are presented as being effective against phytopathogenic fungi of plants, with a good efficacy. These compounds, as well as those described in patent US-A-3,228,950, do not have substituents in the 4-position of the pyridine nucleus.

[0003] Patent application WO-A-00/26191 presents picolinamide derivatives which are optionally substituted at the 4-position with the methoxy radical. Patent application WO-A-95/25723 proposes, for its part, 3-pyridylcarboxylic acid derivatives.

[0004] Patent application WO-A-01/14339 discloses heterocyclic aromatic amide derivatives and their use as fungicides. Compounds according to the present invention are not disclosed in that patent application.


[0005] Amide derivatives of picolinic acid are also known from the publication of patent application JP-11228542. These derivatives are presented as having potential antifungal activities and low toxicity to be used in pharmaceutical products.

[0006] Other picolinic acid derivatives are also known from patent application EP-A-0 690 061 in which such compounds are used as synthesis intermediates for the preparation of pyridothiadiazoles.

[0007] Nevertheless, these known compounds have the disadvantage of being toxic products, this precluding any use of these compounds in agriculture for the eradication of phytopathogenic diseases of crops. Furthermore, these compounds are obtained from fermentation liquors and possess relatively complex chemical structures. Thus, the preparation and purification of these compounds remain delicate and expensive operations, making any industrial synthesis and marketing scarcely profitable.




[0008] We have now found a new family of picolinic acid derivative which do not possess the above mentioned drawbacks and which have an improved fungicidal activity.

[0009] Accordingly, the present invention relates to picolinic acid derivative of general formula (I):



in which:




- Y is chosen in the group consisting of Y¹ to Y³ :





- k represents 0, 1 or 2;
- Het represents a five or six membered saturated or partially unsaturated or aromatic ring containing one to three heteroatoms of the group N, O, and S which can be identical or different and which can be substituted by one or two -R⁵;
- Q³ is -R¹ or -OR¹;

EP 1 404 666 B1



- **G** is chosen in the group consisting of $-(CH_2)_m-$, $-O-$, $-S-$ and $-NR^1$;
- **Z** is chosen in the group consisting of $-R^1$, C_1-C_4 alkylene, C_1-C_4 alkyne, $-Si(R^1)_3$, $-(CH_2)_p-O$ Me, $-(CH_2)_p-S$ Me,




- **X₁** and **X₂** are independently chosen in the group consisting of hydrogen, halogen, $-CF_3$, cyano group and nitro group;
- **Q¹** is chosen in the group consisting of $-(CH_2)_q-$,



- **Q²** is chosen in the group consisting of $-(O)_n-R^5$, cyano group,



and



- **R¹** is hydrogen or C₁-C₄ alkyl;
- **R²** is chosen in the group consisting of hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ halogenoalkyl and C₁-C₄ halogenoalkoxy;
- **R³** is chosen in the group consisting of hydrogen, C₁-C₄ alkyl and C₁-C₄ alkoxyalkyl;
- **R⁴** is chosen in the group consisting of halogen, C₁-C₄ alkyl and C₁-C₄ alkoxyalkyl;
- **R⁵ and R⁶** are independently of each other chosen in the group consisting of hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ halogenoalkyl;
- **R⁷ and R⁸** are independently of each other chosen in the group consisting of hydrogen and halogen;
- **n** is 0 or 1;
- **j, p, q and t** are independently chosen as being 0, 1, 2, 3 or 4;
- **m** is chosen as being 1, 2, 3 or 4;
- **r** is 0, 1, 2 or 3;

and the optional N-oxides, geometric and/or optical isomers, enantiomers and/or diastereoisomers, tautomeric forms, salts, metal and metalloid complexes of the compounds of formula (I) as they have just been defined; with the proviso that compound of formula (I) is different from :


- 3-methoxy-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-hydroxy-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-ethoxy-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- N-[4-(benzyloxy)phenyl]-3-(methylthio)-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-4-piperidin-1-yl-N-{4-[3-(trifluoromethyl)phenoxy]phenyl}pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-4-pyrrolidin-1-yl-N-{4-[3-(trifluoromethyl)phenoxy]phenyl}pyridine-2-carboxamide;
- 3-mercapto-4-piperidin-1-yl-N-{4-[3-(trifluoromethyl)-phenoxy]phenyl}pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-thiomorpholin-4-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(4-methylpiperidin-1-yl)pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(3,5-dimethylpiperidin-1-yl)pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(3-methylpiperidin-1-yl)pyridine-2-carboxamide;
- 3-mercapto-N-[4-(4-methylphenoxy)phenyl]-4-(4-methylpiperidin-1-yl)pyridine-2-carboxamide; and
- 3-mercapto-N-[4-(4-methylphenoxy)phenyl]-4-(3-methylpiperidin-1-yl)pyridine-2-carboxamide.

[0010] The tautomeric forms of the compound of general formula (I) are also included in the present invention. By tautomeric forms it is to be understood all of the isomeric forms well known in the art and as described in the work "The Tautomerism of Heterocycles, Advances in Heterocyclic Chemistry, Supplement 1, by J Elguero, C. Martin, A.R. Katritsky and P Linda, published by Academic Press, New York, 1976, pages 1-4.

[0011] The compound of general formula (I) can exist in one or more forms of geometrical isomers according to the number of double bonds in the compound. For example, compounds of general formula (I) can comprise 2 different geometrical isomers denoted (*E*) or (*Z*) depending on the configuration of the two double bonds. The *E* and *Z* notation can be replaced, respectively, by the term "syn" and "anti", or "cis" and "trans". Reference is made particularly to the work of E. Eliel and S. Wilen "Stereochemistry of Organic Compounds", published by Wiley (1994), for the description and use of these notations.


[0012] Preferably, the present invention relates to picolinic acid derivative of general formula (I) where the different substituents may be chosen independently from each other as being :

- as regards Y of the general formula (I), Y is Y². More preferably, Y is chosen in the group consisting of pyrroles, dimethylpyrroles and imidazole.
- as regards G of the general formula (I), G is O.
- as regards Q¹ of the general formula (I), Q¹ is chosen in the group consisting of



[0013] More preferably, the present invention relates to picolinic acid derivative of general formula (I) in which :


- **Y** is Y^2 ,
- **G** is O,
- and **Q¹** is



[0014] The compound of general formula (I) can exist in one or more optical isomeric or chiral forms according to the number of asymmetric centres in the compound. The present invention thus also includes all the optical isomers and their racemic or scalemic (scalemic designates a mixture of enantiomers in different proportions), as well as the mixtures of all possible stereoisomers in all proportions, including the racemic mixture. The separation of the diastereoisomers and/or optical isomers can be effected by known methods (E. Eliel *ibid.*).

[0015] The present invention also relates to a process of preparation of the compound of the general formula (I). Thus, according to a further aspect of the present invention there is provided a process for the preparation of compound of the general formula (I), **characterised in that** it comprises :

- as a first step, the reaction scheme S-I:




in which:

- W_1 is chosen from bromine, chlorine and iodine;
- W_2 is chosen from fluorine, chlorine, the group -O-DMG, the group -S-DMG and the group -NR¹-DMG; in which -DMG represents an *ortho*-directing metallation group such as those mentioned in the clarifications by V. Snieckus, Chemical Reviews (1990), pages 879-933, or by G. Quéguiner, Tetrahedron (2001), pages 4059-4090,
- k , R^1 , R^5 , R^6 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined above.

[0016] The *ortho*-lithiation of the pyridines of formula (IIa), according to methods such as those reported in the clarifications by V. Snieckus or G. Quéguiner cited above, makes it possible to prepare the compounds of formula (IIc), for which k represents 1 or 2, by reacting the intermediate lithium-containing pyridines with compounds of formula (IIj). The reaction of the same intermediates with formylation reagents such as dimethylformamide, ethyl formate, *N*-formylmorpholine or *N*-formylpiperidine, leads to the compounds of formula (IIb). The acetylation of the latter leads to the compounds of formula (IIc) for which k represents 0. The oxidation of the compounds of formula (IIc) with *meta*-chloroperbenzoic acid or hydrogen peroxide leads to the compounds of general formula (IIe). The latter are subjected to the Reissert-

Henze reaction or, more advantageously, to the modification of this reaction described by W.K. Fife, Journal of Organic Chemistry (1983), pages 1375-1377, to give the compounds of formula (IIf). The cleavage of the DMG group or, when W_2 represents a fluorine or chlorine atom, aromatic nucleophilic substitution reactions make it possible to convert the compounds of formula (IIf) to compounds of formula (IIg). Hydrolysis with potassium or sodium hydroxide in aqueous medium of the cyano radical of the compounds of general formula (IIg) makes it possible to prepare the compounds of general formula (IIh). Conversion of the carboxyl radical of the compounds of general formula (IIh) according to methods well known to persons skilled in the art leads to the compounds of general formula (IIi) for which a reaction for hydrolysing the acetal group makes it possible to prepare the compounds of general formula (Ia).

- as a second step, the preparation of the compound of general formula (Ic)




(Ic) ;

in which Y^3 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined above;

which compound may be prepared by reacting the compound of general formula (Ia) as defined above with a primary amine (R_1-NH_2), a hydrazine ($R_4R_5N-NH_2$) or a hydroxylamine (R_2O-NH_2), optionally in the presence of a Lewis acid and/or a water-capturing agent such as a molecular sieve.


[0017] The compound of general formula (Ib)



(Ib)

in which Y^2 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined above and the Het group is not directly linked to the rest of the structure (ie $k=0$) by a nitrogen atom may be prepared from the compounds of general formula (Ic) as defined above according to known heterocyclic methods of synthesis such as those reported in the book by A.R Katritzky "Handbook of heterocyclic chemistry", 2nd edition, Ed. Pergamon (2000).

[0018] The compound of general formula (Ib) in which Y^2 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined above and the Het group is directly linked to the rest of the structure (ie $k=0$) by a nitrogen atom may be prepared from the compound of general formula (IIIa) (prepared according to methods described in WO 01/49667) according to scheme S-II :



in which Y^2 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined above.

[0019] Aromatic nucleophilic substitution reactions with a base such as cesium or potassium carbonate in an aprotic solvent such as DMF or acetonitrile makes it possible to convert the compounds of formula (IIIa) to compounds of formula (IIIb). Hydrolysis with potassium or sodium hydroxide or with hydrochloric acid in aqueous medium of the cyano radical of the compounds of general formula (IIIb) makes it possible to prepare the compound of general formula (IIIc). Conversion of the carboxyl radical of the compound of general formula (IIIc) according to methods well known to persons skilled in the art leads to the compound of general formula (Ib).

[0020] The compound according to the present invention can be prepared according to the general method of preparation described above. It will nevertheless be understood that the skilled worker will be able to adapt this method according to the specifics of each of the compounds, which it is desired to synthesise.

[0021] The present invention also relates to fungicidal composition comprising an effective amount of an active material of general formula (I). Thus, according to the present invention, there is provided a fungicidal composition comprising, as an active ingredient, an effective amount of a compound as defined above and an agriculturally acceptable support.

[0022] In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid. Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

[0023] The composition may also comprise additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be between 5% and 40% by weight.

[0024] Additional but optional components also include protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.

[0025] In general, the composition according to the invention usually may contain from 0.05 to 99% (by weight) of active material.

[0026] Compositions according to the present invention can be used in quite diverse forms such as aerosol dispenser, bait (ready for use), bait concentrate, block bait, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, grain bait, granular bait, granule, hot fogging concen-

trate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil miscible liquid, paste, plant rodlet, plate bait, powder for dry seed treatment, scrap bait, seed coated with a pesticide, smoke candle, smoke cartridge, smoke generator, smoke pellet, smoke rodlet, smoke tablet, smoke tin, soluble concentrate, soluble powder, solution for seed treatment, suspension concentrate (flowable concentrate), tracking powder, ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, vapour releasing product, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

[0027] These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before they are applied to the crop.

[0028] The compounds of the invention can also be mixed with one or more insecticides, fungicides, bactericides, attractant acaricides or pheromones or other compounds with biological activity. The mixtures thus obtained have a broadened spectrum of activity. In particular the compounds of the present invention do not exhibit the problem of cross-resistance with strobilurin derivatives. In fact the compounds of the present invention are active on a different biochemical site to strobilurin derivatives.

[0029] The mixtures with other fungicides are particularly advantageous, especially the mixtures with acibenzolar-S-methyl, benalaxyl, benomyl, blasticidin-S, bromuconazole, captan, captan, carbendazim, carboxin, carpropamide, chlorothalonil, fungicidal compositions based on copper, derivatives of copper such as copper hydroxide and copper oxychloride, cyazofamide, cymoxanil, cyproconazole, cyprodinil, dichloran, diclocymet, diethofencarb, difenoconazole, diflumetorim, dimethomorph, diniconazole, dodemorph, dodine, edifenphos, epoxyconazole, ethaboxam, ethirimol, flumoxadone, fenamidone, fenarimol, fenbuconazole, fenhexamide, fenciclonil, fenpropidine, fenpropimorph, ferimzone, fluazinam, fludioxonil, flumetover, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpel, furalaxyl, furametpyr, guazatine, hexaconazole, hymexazol, imazalil, iprobenphos, iprodione, isoprothiolane, kasugamycin, mancozeb, maneb, mefenoxam, mepanipyrim, metalaxyl and their enantiomeric forms such as metalaxyl-M, metconazole, metiramzinc, oxadixyl, pefurazoate, penconazole, pencycuron, phosphorous acid and its derivatives such as fosetyl-Al, phthalide, probenazole, prochloraz, procymidone, propamocarb, propiconazole, pyrimethanil, pyroquilon, quinoxifen, silthiofam, simeconazole, spiroxamine, tebuconazole, tetraconazole, thiabendazole, thifluzamide, thiophanate, for example thiophanate-methyl, thiram, triadimefon, triadimenol, triazolopyrimidines e.g. cloransulam-methyl, flumetsulam, florasulam, metosulam, tricyclazole, tridemorph, trifloxystrobin, triticonazole, derivatives of valinamide such as for example, iprovalicarb and benthiavalicarb, vinclozolin, zineb and zoxamide, as well as fungicide of the strobilurin family, for example azoxystrobin, kresoxym-methyl, metominostrobin, discostrobin, dimoxystrobin, picoxystrobin, pyraclostrobin, and trifloxystrobin.

[0030] The fungicidal compositions of the present invention can be used to curatively or preventively combat the phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for curatively or preventively combating the phytopathogenic fungi of crops **characterised in that** a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to the fruit of the plant or to the soil in which the plant is growing or in which it is desired to grow.

[0031] The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

[0032] The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be combated, the type of crop, the climatic conditions and the compounds included in the fungicidal composition according to the invention.

[0033] This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

[0034] The method of treatment according to the present invention is useful to treat propagation material such as tubers and rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

[0035] Among the plants targeted by the method according to the invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae* sp. (for instance pip fruits such as apples and pears, but also stone fruits such as apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantains), *Rubiaceae* sp., *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruits); leguminous crops such as *Solanaceae* sp. (for instance tomatoes), *Liliaceae* sp., *Asteraceae* sp. (for instance lettuces), *Umbelliferae* sp., *Cruciferae* sp., *Chenopodiaceae* sp., *Cucurbitaceae* sp., *Papilionaceae* sp. (for instance peas), *Ro-*

saceae sp. (for instance strawberries); big crops such as *Graminae* sp. (for instance maize, cereals such as wheat, rice, barley and triticale), *Asteraceae* sp. (for instance sunflower), *Cruciferae* sp. (for instance colza), *Papilionaceae* sp. (for instance soja), *Solanaceae* sp. (for instance potatoes), *Chenopodiaceae* sp. (for instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

[0036] Among the plants and the possible diseases of these plants targeted by the method according to the present invention, mention may be made of :

- wheat, as regards controlling the following seed diseases: fusaria (*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*, *Tapesia acuiformis*), take-all (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts (*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and *Septoria nodorum*);
- wheat and barley, as regards controlling bacterial and viral diseases, for example barley yellow mosaic;
- barley, as regards controlling the following seed diseases: net blotch (*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- barley, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yallundae*), net blotch (*Pyrenophora teres* and *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis forma specie hordei*), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
- potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora infestans*) and certain viruses (virus Y);
- potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);
- cotton, as regards controlling the following diseases of young plants grown from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and black root rot (*Thielaviopsis basicola*);
- protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew (*Peronospora pisi*);
- oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;
- com, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);
- flax, as regards controlling the seed disease: *Alternaria linicola*;
- forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);
- rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);
- leguminous crops, as regards controlling the following diseases of seeds or of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium* sp.);
- leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis* sp.), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium* sp.), alternaria leaf spot (*Alternaria* sp.), anthracnose (*Colletotrichum* sp.), septoria leaf spot (*Septoria* sp.), black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora* sp., *Pseudoperonospora* sp., *Phytophthora* sp.);
- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);
- vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);
- beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

[0037] Cereals are preferably treated according to the method of the present invention. Wheat and rice are still preferred for carrying out the method according to the invention.

[0038] The fungicide composition according to the present invention may also be used against fungal diseases liable to grow on or inside lumber. The term "lumber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating lumber according to the invention consists in placing one or more compounds of the present invention, or a composition according to the invention, in contact. This placing in contact may cover the most diverse of forms such as,

for example, direct application, spraying, dipping, injection or any other suitable means.

[0039] The dose of active material applied in the treatment according to the present invention is generally and advantageously between 10 and 800 g/ha, preferably between 50 and 300 g/ha for applications in foliar treatment. The dose of active material applied is generally and advantageously between 2 and 200 g per 100 kg of seed, preferably between 3 and 150 g per 100 kg of seed in the case of seed treatments. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to tailor the application doses according to the nature of the crop to be treated.

[0040] The fungicidal composition according to the present invention may also be used in the treatment of genetically modified plants with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into whose genome a heterologous gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

[0041] Among the genes which give the transformed plants new agronomic properties, mention may be made of genes which impart a tolerance to certain herbicides, those which impart a resistance to certain insects, those which impart a tolerance to certain diseases, etc. Such genes are described in particular in patent applications WO 91/02071 and WO 95/06128.

[0042] Among the genes which impart a tolerance to certain herbicides, mention may be made of the *Bar* gene imparting tolerance to bialaphos, the gene encoding a suitable EPSPS imparting a resistance to herbicides having EPSPS as target, such as glyphosate and its salts (US 4,535,060, US 4,769,061, US 5,094,945, US 4,940,835, US 5,188,642, US 4,971,908, US 5,145,783, US 5,310,667, US 5,312,910, US 5,627,061, US 5,633,435 and FR 2 736 926), the gene encoding glyphosate oxidoreductase (US 5 463 175) or a gene encoding an HPPD imparting a tolerance to herbicides having HPPD as target, such as isoxazoles, in particular isoxafutol (FR 95/06800 and FR 95/13570), diketone nitriles (EP-A-0 496 630 and EP-A-0496631) or triketones, in particular sulcotrioline (EP-A-0 625 505, EP-A-0 625 508 and US 5 506 195). Such genes encoding an HPPD imparting a tolerance to herbicides having HPPD as target are disclosed in patent application WO 96/38567. In the case of genes encoding EPSPS or HPPD, and more particularly for the above genes, the sequence encoding these enzymes is advantageously preceded by a sequence encoding a transit peptide, in particular for the transit peptide known as optimized transit peptide, disclosed in patent US 5 510 471.

[0043] Among the genes imparting novel insect-resistance properties, mention will be made more particularly of the genes encoding the *Bt* proteins which are widely described in the literature and well known to those skilled in the art. Mention will also be made of the genes encoding proteins extracted from bacteria such as *Photobacterium* (WO 97/17432 and WO 98/08932).

[0044] Among the genes imparting novel disease-resistance properties, mention will be made in particular of the genes encoding chitinases, glucanases and oxalate oxidase, all these proteins and their coding sequences being widely described in the literature, or genes encoding antibacterial and/or antifungal peptides, in particular cysteine-rich peptides containing less than 100 amino acids, such as plant thionins or defensins, and more particularly lytic peptides of all origins comprising one or more disulphide bridges between the cysteines and regions comprising basic amino acids, in particular the following lytic peptides: androctonine (WO 97/30082 and PCT/FR98/01814, filed on 18 August 1998) or drosomicin (PCT/FR98/01462, filed on 8 July 1998). Mention will also be made of the genes encoding fungal elicitor peptides, in particular the elicitors (Kamoun et al., 1993; Panabières et al., 1995).

[0045] Among the genes which modify the constitution of modified plants, mention may be made in particular of genes which modify the content and quality of certain essential fatty acids (EP-A-0 666 918) or the content and quality of proteins, in particular in the leaves and/or seeds of the said plants. Mention will be made in particular of the genes encoding proteins that are rich in sulphur-containing amino acids (WO 98/20133; WO 97/41239; WO 95/31554; WO 94/20828 and WO 92/14822).

[0046] The fungicidal composition according to the present invention may, in particular, be used to the treatment of genetically modified plants comprising a heterologous gene, which gives the plant disease-resistance properties. The heterologous gene preferentially gives the genetically modified plant a spectrum of activity that is complementary to the spectrum of activity of the compounds according to the invention. According to the invention, the expression "complementary spectrum" means a spectrum of activity for the heterologous gene which is different from the spectrum of activity of the compounds according to the invention, or a spectrum of activity relating to identical infectious agents but allowing an identical or improved control for lower application doses of compounds according to the invention.

[0047] The compositions according to the present invention may also be used to curatively or preventively treat human and animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton diseases and candidiasis or diseases caused by *Aspergillus* spp., for example *Aspergillus fumigatus*.

[0048] The aspects of the present invention will now be illustrated with reference to the following tables of compounds and examples and reference examples. The following Tables I to III illustrate in a non-limiting manner examples of fungicidal compounds according to the present invention and reference examples. In the following Examples, "MP"

EP 1 404 666 B1

signifies "melting point" and is expressed in °Celsius (°C). M+1 (or M-1) means the molecular ion peak, plus or minus 1 a.m.u. (atomic mass units) respectively, as observed in mass spectroscopy.

5

10

15

20

25

30

35









40

45

50



















55

Table I :

















	N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
	1-1								M + 1 = 403	147
	1-2								M + 1 = 403	160
	1-3								M+1= 329	
	1-4								M + 1 = 415	112
















(continued)									
N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
1-5								M + 1 = 321	202
1-6								M + 1 = 430	

















Table II :

	N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
	2-1								M + 1 = 553	
	2-2								M + 1 = 432	
	2-3								M + 1 = 553	
	2-4								M + 1 = 432	
	2-5									















(continued)

5		melting point			131				
10		ion mol.							
15	Q2								
20	Q1								
25	Y								
30	Z								
35	G								
40	X2								
45	X1								
50	N°	2-6	2-7	2-8	2-9	2-10	2-11		
55									

















N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-12									
2-13									
2-14									
2-15									130
2-16									
2-17									

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-18									
2-19									
2-20									
2-21									155
2-22									
2-23									

(continued)
















5							
10							
15	Q2						
20							
25	Q1						
30	Y						
35							
40	Z						
45	G						
50	X2						
55	X1						
	N°	2-24	2-25	2-26	2-27	2-28	2-30

(continued)

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-31									
2-32								172	
2-33									
2-34									
2-35									
2-36									
2-37									















(continued)

5					113	
10						
15	Q2					
20						
25	Q1					
30	Y					
35						
40	Z					
45	G					
50	X2					
55	X1					
	N°	2-38	2-39	2-40	2-41	2-42

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-44									
2-45									
2-46									
2-47									
2-48									138
2-49									















N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-50									
2-51									
2-52									
2-53									
2-54									
2-55									

(continued)

5		melting point								
10		ion mol.								
15	Q2									
20	Q1									
25	Y									
30	Z									
35	G									
40	X2									
45	X1									
50	N°	2-56	2-57	2-58	2-59	2-60				
55										

(continued)

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-61									
2-62									
2-63									
2-64									
2-65									118

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-66									
2-67									
2-68									160
2-69									
2-70									
2-71									




















(continued)

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-72			-O-						156
2-73			-S-					M + 1 = 580	
2-74			-O-						140
2-75			-O-						
2-76			-O-						
2-77			-S-					M + 1 = 557	



















(continued)

N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
2-78									
2-79									
2-80									143
2-81									
2-82									
2-83									


















(continued)

5		melting point							
10		ion mol.					M + 1 = 592		
15	Q2								
20	Q1								
25	Y								
30	Z								
35	G								
40	X2								
45	X1								
50	N°	2-84	2-85	2-86	2-87	2-88	2-89		
55									

(continued)

5		melting point								
10		ion mol.		M + 1 = 440	M + 1 = 452	M + 1 = 438	M + 1 = 468	M + 1 = 400		
15	Q2									
20	Q1									
25	Y									
30	Z									
35	G	-S-	-O-	-O-	-O-	-O-	-O-	-O-		
40	X2									
45	X1									
50	N°	2-90	2-91	2-92	2-93	2-94	2-95			
55										

(continued)

5		melting point								
10		ion mol.	M + 1 = 468	M + 1 = 461	M + 1 = 373	M + 1 = 447	M + 1 = 454	M + 1 = 454		
15	Q2									
20	Q1									
25	Y									
30	Z									
35	G									
40	X2									
45	X1									
50	N°	2-96	2-97	2-98	2-99	2-100	2-101			

(continued)















	N°	X1	X2	G	Z	Y	Q1	Q2	ion mol.	melting point
5										
10										
15										
20										
25										
30										
35										
40										
45										
50										
55										
	2-102								M + 1 = 372	
	2-103								M + 1 = 482	
	2-104								M + 1 = 482	

Table III:


							
N°	X1	X2	G	Z	Y	Q1	Q2
3-1							
3-2							
ion mol.				melting point			
M+1= 336							
M+1= 418							

Examples of process for preparation of the compounds of general formula (I)

Example : Preparation of picolinic acid derivatives of formula (I) :

3-fluoro-4-(1,3-dioxolan-2-yl) pyridine-N-oxyde (example of compound of formula IIe according to scheme S-1)


[0049]



[0050] 11,1g (80 mmol) of 3-fluoro-4-formyl-pyridine (prepared according to the methods described in Queginer, Tetrahedron, (1983), pages 2021) are dissolved in 50ml of toluene. 6,2ml (112 mmol) of anhydride ethylene glycol, 17,0g (88 mmol) of paratoluene sulfonic acid and 250 ml of toluene are added successively. The resulting solution is heated at 130°C whilst the water formed is removed with a dean-stark. When the reaction is finished, the mixture is cooled down and the toluene removed in vacuum. The residue is taken up in a mixture of 200ml of chloroform and 100ml of a 1M sodium hydroxide solution. The phases are separated, and the aqueous phase is further extracted with chloroform. The organic phases are washed with brine, dried and solvent evaporated. The residue is taken up in 100ml of dichloromethane, and a solution of 22,0g (88 mmol) meta-chloro-perbenzoic acid in 200ml of dichloromethane is added. The resulting solution is stirred at room temperature overnight. 22g of potassium carbonate in 22ml of water were added, and the resulting mixture is stirred for 20 minutes, before being filtered off on celite. The filtrate was dried and solvent evaporated to give 11,8g (79% yield) of a yellow oil (M+1 = 186).

2-cyano-3-fluoro-4-(1,3-dioxolan-2-yl) pyridine (example of compound of formula IIe according to scheme S-1)


[0051]



[0052] 4g (21,6 mmol) of 3-fluoro-4-(1,3-dioxolan-2-yl)-pyridine-N-oxyde are dissolved in 34 ml of acetonitrile. 9,1ml (64,8 mmol) of triethylamine, followed by 8,6ml (64,8mmol) of trimethylsilyl cyanide are added and the resulting solution is heated at reflux for 12 hours. The solvent is evaporated and the residue is taken up in a mixture of 70ml of ethyl acetate, 70ml of water and 34g of potassium carbonate. The organic phases are washed with brine, dried filtrated on celite and solvent evaporated. Purification on silica give 3,7g (88% yield) of a brown oil (M+1 = 195)

3-fluoro-4-(1,3-dioxolan-2-yl) picolinic acid


[0053]



[0054] A mixture of 4,1g (21mmol) of 2-cyano-3-fluoro-4-(1,3-dioxolan-2-yl)-pyridine, 2,1g (53 mmol) sodium hydroxide in 100ml of water is heated at 90°C for 2 hours. After cooling, the reaction mixture is acidified to pH = 1 with aqueous hydrochloric acid, then extracted with ethyl acetate. The organic phases are washed with brine, dried filtrated on celite and solvent evaporated. There is obtained 2,4g (53% yield) of a white solid (M+1 = 214)

2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-fluoro-4-(1,3-dioxolan-2-yl) pyridine


[0055]



[0056] To a mixture of 0,95g (4,5 mmol) 3-fluoro-4-(1,3-dioxolan-2-yl) picolinic acid and 1,1g (4,5 mmol) of para-[4-(trifluoromethylphenoxy)] aniline in 30ml of dichloromethane are added 0,95g (4,5 mmol) of 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide chlorhydrate. The mixture is stirred at room temperature for 3 hours. After solvent evaporation, the residue is purified on silica. There is obtained 1,6g (80% yield) of a cream solid (M+1 = 449)

2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-(1,3-dioxolan-2-yl) pyridine (compounds n° 2-99)


[0057]



[0058] A mixture of 200mg (0,45 mmol) of 2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-fluoro-4-(1,3-dioxolan-2-yl) pyridine, few drops of water, 4 ml of DMSO and 125mg (2,2 mmol) of potassium hydroxide is heated at 100°C for 16 hours. After cooling, the reaction mixture is acidified with a 1M aqueous hydrochloric acid solution, and extracted with ethyl acetate. The organic phases are washed with brine, dried and solvent evaporated. There is obtained 139mg (69% yield) of a brown oil (M+1 = 447)

2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-formyl pyridine (compounds n° 1-2)


[0059]



[0060] A solution of 131 mg (0,29 mmol) of 2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-(1,3-dioxolan-2-yl) pyridine in 1ml of formic acid and 1ml of water is heated at 100°C for 3 hours. After cooling, the resulting precipitate is filtered off, rinsed with water, triturated with a mixture of diisopropyl ether and heptane, filtered off and dried. There is obtained 67mg (57% yield) of a cream solid (M+1 = 403)

2 -{N-(3-bromophenyl) aminocarbonyl}-3-hydroxy-4-carboxaldehyde oxime pyridine (compounds 3-1)


[0061]



[0062] A mixture of 50mg (0,160 mmol) of 2-{N-(3-bromophenyl) aminocarbonyl}-3-hydroxy-4-formyl pyridine, 11,4mg (0,164 mmol) of hydroxylamine hydrate, 32mg of Amberlyst A-21 and 1ml of ethanol is stirred at room temperature overnight. The resin is filtered off and rinsed with dichloromethane. The filtrate is evaporated to give 54mg (100% yield) of a brown solid. (M+1 = 336)

2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-(2,5 dimethyl pyrrol-1-yl) pyridine (compounds 2-94)

[0063]




[0064] A mixture of 0.5g (1.29 mmol) 2-{N-para-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-amino pyridine (prepared according to the method described in WO 0149666) and 293mg acetonylacetone (2.57 mmol) in 5ml of glacial acetic acid, is refluxed for 24 hours over a dean and stark containing 4A molecular sieves. The reaction

is cooled and solvent evaporated, the residue is dissolved in dichloromethane and washed with aqueous sodium bicarbonate, the organic phase was dried over MgSO_4 and solvent evaporated. The crude product was chromatographed on silica to yield 157mg (26% yield) of a brown oil. ($M+1 = 468$)

2-{N-meta-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-(pyrrol-1-yl) pyridine (compounds 2-93)


[0065]



[0066] A mixture of 0.2g (0.51 mmol) 2-{N-meta-[4-(trifluoromethylphenoxy) phenyl]} aminocarbonyl-3-hydroxy-4-amino pyridine (prepared according to the method described in WO 0149666) and 78mg (0.56 mmol) of 2,5-dimethoxytetrahydrofuran in 1ml of glacial acetic acid, is heated at 110°C for 1 hour. The reaction is cooled and solvent evaporated, the residue is dissolved in dichloromethane and washed with aqueous sodium bicarbonate, the organic phase was dried over MgSO_4 and solvent evaporated. The crude product was chromatographed on silica to yield 137mg (61% yield) of a yellow solid. ($M+1 = 440$)

2-N-(3-trifluoromethyl phenyl) aminocarbonyl-3-hydroxy-4-(imidazol-1-yl) pyridine (compound 2-24)

[0067]



[0068] A mixture of 1.36g (20 mmol) of imidazol and 6.52g (20 mmol) of cesium carbonate is stirred at room temperature in 20ml of DMF for 20 minutes. 3.58g (20 mmol) of 4-nitro-3-methoxy-2-cyano pyridine (prepared according to the methods described in WO 0149667) is added in one pot and the mixture is stirred overnight. DMF is removed in vacuo and the residue is partitioned between water and ethyl acetate. The aqueous phase is further extracted with ethyl acetate. The organic phases are washed with dilute brine, dried and solvent evaporated. Purification on silica gave 1.4 g of 4-(imidazol-1-yl)-3-methoxy-2-cyano pyridine.

A mixture of 1.4 g (7 mmol) of 4-(imidazol-1-yl)-3-methoxy-2-cyano pyridine in 12 mls of concentrated hydrochloric acid is heated at 85°C overnight. After cooling, the solvent is evaporated to dryness, more water is added and the solvent is evaporated. This is done 4 times. little water is added to the residue and the precipitate is filtered off and dried in an oven at 60°C over P_2O_5 under vacuum for 2 hours. There is obtained 0.8g of 4-(imidazol-1-yl)-3-methoxy nicotinic acid. To a stirred mixture of 144 mg (0.7 mmol) of 4-(imidazol-1-yl)-3-methoxy nicotinic acid, 95 mg (0.95 mmol) of HOBT and 113 mg (0.7mmol) of 3-trifluoromethyl aniline in 5 ml of pyridine is added 134mg (0.87 mmol) of EDCI. The mixture is stirred at reflux for 1.5 hours and at room temperature overnight. The solvent was evaporated and the residue is

partitioned between water and dichloromethane. The aqueous phase is further extracted with dichloromethane. The organic phases are washed with dilute brine, dried and solvent evaporated. Purification on silica yielded 97 mg (40% yield) of a yellow solid ($M+1 = 349$).

5 Examples of Biological Activity of the Compounds of the Invention

Example A: *in vivo* test on *Alternaria brassicae* (Leaf spot of crucifers):

[0069] An aqueous suspension, at a concentration of 1.5%, of the active material tested is obtained by grinding it finely in a formulation of concentrated suspension type. This aqueous suspension is then diluted in water so as to obtain the desired active material concentration. Radish plants (Pernot variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

[0070] After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Alternaria brassicae* spores (40,000 spores per cm³). The spores are collected from a 12-13 day-old culture. The contaminated radish plants are incubated for 6-7 days at about 18°C, under a humid atmosphere. Grading is carried out 6 to 7 days after the contamination, in comparison with the control plants. Under these conditions, good (at least 50%) or total protection is observed at a dose of 500 g/ha with the following compounds: 1-1, 1-2, 2-10, 2-11, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 2-20, 2-21, 2-22, 2-23, 2-24, 2-25, 2-29, 2-30, 2-31, 2-32, 2-34, 2-35, 2-362-38, 2-42, 2-522-47, 2-48, 2-49, 2-51, 2-52, 2-53, 2-56, 2-59, 2-60, 2-61, 2-67, 2-83, 2-85, 2-93, 2-94, 2-96, 2-97.

Example B: *in vivo* test on *Septoria nodorum* (wheat Glume blotch):

[0071] An aqueous suspension, at a concentration of 1.5%, of the active material tested is obtained by grinding it finely in a formulation of concentrated suspension type. This aqueous suspension is then diluted in water so as to obtain the desired active material concentration.

[0072] Wheat plants (Scipion variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying them with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material. After 24 hours, the plants are contaminated by spraying with an aqueous suspension of *Septoria nodorum* spores (500,000 spores per cm³). The spores are collected from a seven-day-old culture. The contaminated wheat plants are incubated for 72 hours at about 18°C, under a humid atmosphere, and then for 14 days at 90% relative humidity.

[0073] Grading is carried out 15 to 20 days after contamination, in comparison with the control plants. Under these conditions, good (at least 50%) or total protection is observed, at a dose of 500 g/ha, with the following compounds: 1-2, 2-5, 2-6, 2-7, 2-8, 2-9, 2-12, 2-13, 2-19, 2-20, 2-21, 2-23, 2-24, 2-25, 2-26, 2-28, 2-32, 2-33, 2-34, 2-35, 2-36, 2-37, 2-38, 2-40, 2-41, 2-42, 2-43, 2-46, 2-47, 2-70, 2-71, 2-72, 2-76, 2-78, 2-84, 2-85, 2-93, 2-96

Example C: *in vivo* test on *Erysiphe graminis* f. sp. *tritici* (powdery mildew of wheat):

[0074] An aqueous suspension, at a concentration of 1.5%, of the active material tested is obtained by grinding it finely in a formulation of concentrated suspension type. This aqueous suspension is then diluted in water so as to obtain the desired active material concentration. Wheat plants (Audace variety) in starter cups, sown on 50/50 peat soil-pozzulana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material. After 24 hours, the plants are contaminated by dusting them with *Erysiphe graminis* f. sp. *tritici* spores, the dusting being carried out using diseased plants.

Grading is carried out 7 to 14 days after the contamination, in comparison with the control plants. Under these conditions, good (at least 50%) or total protection is observed, at a dose of 500 g/ha, with the following compounds: 1-2, 1-3, 2-5, 2-6, 2-7, 2-8, 2-9, 2-12, 2-13, 2-14, 2-15, 2-17, 2-18; 2-19, 2-20, 2-21, 2-22, 2-23, 2-24, 2-25, 2-26, 2-42, 2-43, 2-44, 2-29, 2-31, 2-32, 2-33, 2-34, 2-35, 2-36, 2-37, 2-40, 2-41, 2-46, 2-47, 2-48, 2-49, 2-53, 2-68, 2-71, 2-76, 2-78, 2-79, 2-80, 2-83, 2-84, 2-85, 2-93, 2-96, 2-97.

Example D: *in vivo* test on *Septoria tritici* (Leaf spot of wheat):

[0075] An aqueous suspension, at a concentration of 1.5%, of the active material tested is obtained by grinding it finely in a formulation of concentrated suspension type. This aqueous suspension is then diluted in water so as to obtain the desired active material concentration. Wheat plants (Scipion variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous

suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

[0076] After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Septoria tritici* spores (500,000 spores per mL). The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of:

- 1.5 g/L of gelatine
- 0.5 g/L of sodium oleate
- 24 g/L of PDB

[0077] The contaminated wheat plants are incubated for 72 hours at about 20°C and at 100% relative humidity, and then for 15 days at 80% relative humidity.

[0078] Grading is carried out 15 to 20 days after the contamination, in comparison with the control plants. Under these conditions, good (at least 50%) protection is observed, at a dose of 500 g/ha, with the following compounds: 1-2, 1-3, 2-5, 2-6, 2-7, 2-8, 2-12, 2-18, 2-19, 2-20, 2-21, 2-22, 2-23, 2-30, 2-31, 2-34, 2-40, 2-46, 2-47, 2-70, 2-71, 2-72, 2-74, 2-75, 2-76, 2-80, 2-83, 2-84, 2-85, 2-91, 2-93, 2-96, 2-98.

Example E: *in vivo* test on *Puccinia recondita* (Wheat brown rust):

[0079] An aqueous suspension, at a concentration of 1.5%, of the active material tested is obtained by grinding it finely in a formulation of concentrated suspension type. This aqueous suspension is then diluted in water so as to obtain the desired active material concentration. Wheat plants (Scipion variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

[0080] After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Puccinia recondita* spores (150,000 spores per mL). The contaminated wheat plants are incubated for 24 hours at about 20°C and at 100% relative humidity, and then for 10 days at 70 % relative humidity.

Grading is carried out 10 days after the contamination, in comparison with the control plants. Under these conditions, good (at least 50%) protection is observed, at a dose of 500 g/ha, with the following compounds described as example: 1-2, 2-5, 2-7, 2-9, 2-12, 2-13, 2-18, 2-22, 2-34, 2-46, 2-47, 2-69, 2-71, 2-76, 2-79, 2-80, 2-84, 2-85, 2-94, 2-96, 2-98.

Example F: *in vivo* test on *Botrytis cinerea* (cucumber Grey mould):

[0081] An aqueous suspension, at a concentration of 1.5%, of the active material tested is obtained by grinding it finely in a formulation of concentrated suspension type. This aqueous suspension is then diluted in water so as to obtain the desired active material concentration. Cucumber plants (Marketer variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18- 20°C, are treated at the cotyledon Z11 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

[0082] After 24 hours, the plants are contaminated by depositing drops of an aqueous suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the leaves. The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of:


- 20 g/L of gelatine
- 50 g/L of cane sugar
- 2 g/L of NH₄NO₃
- 1 g/L of KH₂PO₄

[0083] The contaminated cucumber plants are settled for 5/7 days in a climatic room at 15-11 °c (day/night) and at 80% relative humidity.

[0084] Grading is carried out 5/7 days after the contamination, in comparison with the control plants. Under these conditions, good (at least 50%) protection is observed, at a dose of 500 g/ha, with the following compounds described as example: 2-11, 2-20, 2-70, 2-91, 2-92, 2-94, 2-96, 2-98.

Claims




1. Compound of general formula (I):



(I)

in which:

- Y is chosen in the group consisting of Y^1 to Y^3 :

 Y^1 , Y^2 and Y^3








- k represents 0, 1 or 2;

- Het represents a five or six membered saturated or partially unsaturated or aromatic ring containing one to three heteroatoms of the group N, O, and S which can be identical or different and which can be substituted by one or two $-R^5$;

- Q^3 is $-R^1$ or $-OR^1$;



- G is chosen in the group consisting of $-(CH_2)_m-$, $-O-$, $-S-$ and $-NR^1$;

- Z is chosen in the group consisting of $-R^1$, C_1-C_4 alkylene, C_1-C_4 alkyne, $-Si(R^1)_3$, $-(CH_2)_p-OMe$, $-(CH_2)_p-SMe$,





- X_1 and X_2 are independently chosen in the group consisting of hydrogen, halogen, $-CF_3$, cyano group and nitro group;


- Q^1 is chosen in the group consisting of $-(CH_2)_q-$,



- **Q²** is chosen in the group consisting of $-(\text{O})_n\text{---R}^5$, cyano group,



and



- **R¹** is hydrogen or C₁-C₄ alkyl;
- **R²** is chosen in the group consisting of hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ halogenoalkyl and C₁-C₄ halogenoalkoxy;
- **R³** is chosen in the group consisting of hydrogen, C₁-C₄ alkyl and C₁-C₄ alkoxyalkyl;
- **R⁴** is chosen in the group consisting of halogen, C₁-C₄ alkyl and C₁-C₄ alkoxyalkyl;
- **R⁵ and R⁶** are independently of each other chosen in the group consisting of hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ halogenoalkyl;
- **R⁷ and R⁸** are independently of each other chosen in the group consisting of hydrogen and halogen;
- **n** is 0 or 1;
- **j, p, q and t** are independently chosen as being 0, 1, 2, 3 or 4;
- **m** is chosen as being 1, 2, 3 or 4;
- **r** is 0, 1, 2 or 3

and the optional N-oxides, geometric and/or optical isomers, enantiomers and/or diastereoisomers, tautomeric forms, salts, metal and metalloid complexes of the compounds of formula (I) as they have just been defined; with the proviso that compound of formula (I) is different from :

- 3-methoxy-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-hydroxy-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-ethoxy-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- N-[4-(benzyloxy)phenyl]-3-(methylthio)-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-4-piperidin-1-yl-N-[4-[3-(trifluoromethyl)phenoxy]phenyl]pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-4-pyrrolidin-1-yl-N-[4-[3-(trifluoromethyl)phenoxy]phenyl]pyridine-2-carboxamide;
- 3-mercapto-4-piperidin-1-yl-N-[4-[3-(trifluoromethyl)phenoxy]phenyl]pyridine-2-carboxamide;


- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-thiomorpholin-4-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(4-methylpiperidin-1-yl)pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(3,5-dimethylpiperidin-1-yl)pyridine-2-carboxamide;
- 3-[(4-methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(3-methylpiperidin-1-yl)pyridine-2-carboxamide;
- 3-mercapto-N-[4-(4-methylphenoxy)phenyl]-4-(4-methylpiperidin-1-yl)pyridine-2-carboxamide; and
- 3-mercapto-N-[4-(4-methylphenoxy)phenyl]-4-(3-methylpiperidin-1-yl)pyridine-2-carboxamide.

2. Compound according to claim 1, **characterised in that** Y is Y².

3. Compound according to claim 2, **characterised in that** Y is chosen in the group consisting of pyrroles, dimethylpyrroles and imidazole.


4. Compound according to any of the preceding claims, **characterised in that** G is -O-.

5. Compound according to any of the preceding claims, **characterised in that** Q¹ is chosen in the group consisting of




6. Compound according to claim 1, **characterised in that:**

- Y is Y²,
- G is O,
- and Q¹ is




7. Process for the preparation of compound as defined in any of the claims 1 to 6, **characterised in that** it comprises :

- as a first step, the reaction scheme S- I:




in which:

- W_1 is chosen from bromine, chlorine and iodine;
- W_2 is chosen from fluorine, chlorine, the group $-O$ -DMG, the group $-S$ -DMG and the group $-NR^1$ -DMG; in which -DMG represents an *ortho*-directing metallation group,
- k , R^1 , R^5 , R^6 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as in any of the claims 1 to 6;
- as a second step, the preparation of the compound of general formula (Ic)




in which Y^3 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined in any of the claims 1 to 6; which compound may be prepared by reacting the compound of general formula (Ia) with a primary amine (R_1-NH_2), a hydrazine ($R_4R_5N-NH_2$) or a hydroxylamine (R_2O-NH_2), optionally in the presence of a Lewis acid and/or a water-capturing agent such as a molecular sieve.

8. Process to prepare the compound of general formula (Ib)



in which Y^2 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined in any of the claims 1 to 6 and the Het group is directly linked to the rest of the structure (ie $k=0$) by a nitrogen atom,

characterised in that a compound of general formula (III) is reacting according to scheme S-II :




in which Y^2 , X^1 , X^2 , G , Z , Q^1 and Q^2 are as defined in any of the claims 1 to 6.

9. Fungicidal composition comprising an effective amount of a compound according to any of the claims 1 to 6 and an agriculturally acceptable support.

10. Fungicidal composition according to claim 9 further comprising a surfactant.
11. Fungicidal composition according to either of claims 9 and 10, comprising from 0.05% to 99% by weight of active material.
12. Method for preventively or curatively combating the phytopathogenic fungi of crops, **characterised in that** an effective and non-phytotoxic amount of a composition according to any of the claims 9 to 11 is applied to the plant seeds or to the plant leaves and/or to the fruits of the plants or to the soil in which the plants are growing or in which it is desired to grow them.
13. Method according to claim 12, in which the dose of active material applied is between 10 g and 800 g of active material per hectare in the case of foliar treatments.
14. Method according to Claim 13, in which the dose of active material applied is between 50 g and 300 g of active material per hectare in the case of foliar treatments.
15. Method according to claim 12, in which the dose of active material applied is between 2 and 200 g of active material per 100 kg of seed, in the case of seed treatments.
16. Method according to claim 15, in which the dose of active material applied is between 3 g and 150 g per 100 kg of seed, in the case of seed treatments.




Patentansprüche

1. Verbindung der allgemeinen Formel (I)










in der:

- Y aus der Gruppe bestehend aus Y¹ bis Y³ ausgewählt ist:








- k 0, 1 oder 2 bedeutet;
- **Het** einen fünf- oder sechsgliedrigen gesättigten oder teilweise ungesättigten oder aromatischen Ring mit ein bis drei Heteroatomen aus der Gruppe N, O und S, die gleich oder verschieden sein können und die durch einen oder zwei Reste -R⁵ substituiert sein können, bedeutet;
- **Q³** -R¹ oder -OR¹ bedeutet;
- **G** aus der Gruppe bestehend aus -(CH₂)_m-, -O-, -S- und -NR¹ ausgewählt ist;
- **Z** aus der Gruppe bestehend aus -R¹, C₁-C₄-Alkylen, C₁-C₄-Alkylyn, -Si(R¹)₃, -(CH₂)_p-OMe, -(CH₂)_p-Sme



ausgewählt ist;







- X_1 und X_2 unabhängig aus der Gruppe bestehend aus Wasserstoff, Halogen, $-\text{CF}_3$, Cyanogruppe und Nitrogruppe ausgewählt sind,

- Q^1 aus der Gruppe bestehend aus $-(\text{CH}_2)_q-$,




ausgewählt ist;

- Q^2 aus der Gruppe bestehend aus $-(\text{O})_n-\text{R}^5$, Cyanogruppe,



und



ausgewählt ist;

- R^1 Wasserstoff oder C_1 - C_4 -Alkyl bedeutet;

- **R²** aus der Gruppe bestehend aus Wasserstoff, Halogen, C₁-C₄-Alkyl, C₁-C₄-Alkoxy, C₁-C₄-Halogenalkyl und C₁-C₄-Halogenalkoxy ausgewählt ist;
- **R³** aus der Gruppe bestehend aus Wasserstoff, C₁-C₄-Alkyl und C₁-C₄-Alkoxyalkyl ausgewählt ist;
- **R⁴** aus der Gruppe bestehend aus Wasserstoff, C₁-C₄-Alkyl und C₁-C₄-Alkoxyalkyl ausgewählt ist;
- **R⁵ und R⁶** unabhängig voneinander aus der Gruppe bestehend aus Wasserstoff, Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl ausgewählt sind;
- **R⁷ und R⁸** unabhängig voneinander aus der Gruppe bestehend aus Wasserstoff und Halogen ausgewählt sind;
- **n** 0 oder 1 bedeutet;
- j, p, q und t** unabhängig voneinander aus der Reihe 0, 1, 2, 3 oder 4 ausgewählt sind;
- m** aus der Reihe 1, 2, 3 oder 4 ausgewählt ist;
- r** 0, 1, 2 oder 3 bedeutet

und die gegebenenfalls existierenden N-Oxide, geometrischen und/oder optischen Isomere, Enantiomere und/oder Diastereomere, tautomeren Formen, Salze, Metall- und Metalloidkomplexe der soeben definierten Verbindungen der Formel (I);

mit der Maßgabe, daß sich die Verbindung der Formel (I) unterscheidet von:


- 3-Methoxy-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridin-2-carboxamid;
- 3-Hydroxy-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridin-2-carboxamid;
- 3-Ethoxy-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridin-2-carboxamid;
- N-[4-(Benzyloxy)phenyl]-3-(methylthio)-4-piperidin-1-ylpyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-4-piperidin-1-yl-N-[4-[3-(trifluormethyl)phenoxy]phenyl]pyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-4-pyrrolidin-1-yl-N-[4-[3-(trifluormethyl)phenoxy]phenyl]pyridin-2-carboxamid;
- 3-Mercapto-4-piperidin-1-yl-N-[4-[3-(trifluormethyl)phenoxy]phenyl]pyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-thiomorpholin-4-ylpyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-piperidin-1-ylpyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(4-methylpiperidin-1-yl)pyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(3,5-dimethylpiperidin-1-yl)pyridin-2-carboxamid;
- 3-[(4-Methoxybenzyl)thio]-N-[4-(4-methylphenoxy)phenyl]-4-(3-methylpiperidin-1-yl)pyridin-2-carboxamid;
- 3-Mercapto-N-[4-(4-methylphenoxy)phenyl]-4-(4-methylpiperidin-1-yl)pyridin-2-carboxamid; und
- 3-Mercapto-N-[4-(4-methylphenoxy)phenyl]-4-(3-methylpiperidin-1-yl)pyridin-2-carboxamid.

2. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, daß** Y Y² bedeutet.

3. Verbindung nach Anspruch 2, **dadurch gekennzeichnet, daß** Y aus der Gruppe bestehend aus Pyrrolen, Dimethylpyrrolen und Imidazol ausgewählt ist.

4. Verbindung nach einem der vorhergehenden Ansprüche, **dadurch gekennzeichnet, daß** G -O- bedeutet.


5. Verbindung nach einem der vorhergehenden Ansprüche, **dadurch gekennzeichnet, daß** Q¹ aus der Gruppe bestehend aus



ausgewählt ist.

6. Verbindung nach Anspruch 1, **dadurch gekennzeichnet, daß:**


- **Y** Y² bedeutet,
- **G** O bedeutet,
- und



bedeutet.

7. Verfahren zur Herstellung einer in einem der Ansprüche 1 bis 6 definierten Verbindung, **dadurch gekennzeichnet, daß** es folgendes umfaßt:


- als ersten Schritt das Reaktionsschema S-I:



in dem:

- W_1 aus der Reihe Brom, Chlor und Iod ausgewählt ist;
- W_2 aus der Reihe Fluor, Chlor, der Gruppe -O-DMG, der Gruppe -S-DMG und der Gruppe -NR¹-DMG ausgewählt ist, wobei -DMG eine ortho-dirigierende Metallierungsgruppe bedeutet,
- k , R^1 , R^5 , R^6 , X^1 , X^2 , G , Z , Q^1 und Q^2 wie in einem der Ansprüche 1 bis 6 beschrieben sind;


- als zweiten Schritt die Herstellung einer Verbindung der allgemeinen Formel (Ic)




in der Y³, X¹, X², G, Z, Q¹ und Q² wie in einem der Ansprüche 1 bis 6 definiert sind;

wobei die Verbindung **dadurch** hergestellt werden kann, daß man die Verbindung der allgemeinen Formel (Ia) mit einem primären Amin (R₁-NH₂), einem Hydrazin (R₄R₅N-NH₂) oder einem Hydroxylamin (R₂O-NH₂), gegebenenfalls in Gegenwart einer Lewis-Säure und/oder eines Wasserfängers wie einem Molekularsieb, umsetzt.

8. Verfahren zur Herstellung der Verbindung der allgemeinen Formel (Ib)



in der Y², X¹, X², G, Z, Q¹ und Q² wie in einem der Ansprüche 1 bis 6 definiert sind und die Het-Gruppe direkt durch ein Stickstoffatom mit dem Rest der Struktur verbunden ist (d.h. k=0), **dadurch gekennzeichnet, daß** man eine Verbindung der allgemeinen Formel (III) gemäß Schema S-II umsetzt:




Schema S-II

in dem Y², X¹, X², G, Z, Q¹ und Q² wie in einem der Ansprüche 1 bis 6 definiert sind.

9. Fungizide Zusammensetzung, die eine wirksame Menge einer Verbindung nach einem der Ansprüche 1 bis 6 und einen landwirtschaftlich annehmbaren Träger enthält.
10. Fungizide Zusammensetzung nach Anspruch 9, die weiterhin ein Tensid enthält.
11. Fungizide Zusammensetzung nach einem der Ansprüche 9 und 10, die 0,05 Gew.-% bis 99 Gew.-% Wirkstoff enthält.
12. Verfahren für die präventive oder kurative Bekämpfung von phytopathogenen Pilzen an Kulturpflanzen, **dadurch gekennzeichnet, daß** man eine wirksame und nichtphytotoxische Menge einer Verbindung nach einem der Ansprüche 9 bis 11 auf die Pflanzensamen oder auf die Pflanzenblätter und/oder auf die Früchte der Pflanzen oder auf den Boden, in dem die Pflanzen wachsen oder in dem sie wachsen sollen, ausbringt.
13. Verfahren nach Anspruch 12, bei dem die Dosis an ausgebrachtem Wirkstoff bei Blattbehandlungen zwischen 10 g und 800 g Wirkstoff pro Hektar liegt.
14. Verfahren nach Anspruch 13, bei dem die Dosis an ausgebrachtem Wirkstoff bei Blattbehandlungen zwischen 50 g und 300 g Wirkstoff pro Hektar liegt.
15. Verfahren nach Anspruch 12, bei dem die Dosis an ausgebrachtem Wirkstoff bei Saatgutbehandlungen zwischen 2 und 200 g Wirkstoff pro 100 kg Samen liegt.
16. Verfahren nach Anspruch 15, bei dem die Dosis an ausgebrachtem Wirkstoff bei Saatgutbehandlungen zwischen 3 und 150 g Wirkstoff pro 100 kg Samen liegt.




Revendications

1. Composé de formule générale (I) :



dans laquelle :

- **Y** est choisi dans le groupe constitué par Y^1 à Y^3 :






- **k** représente 0, 1 ou 2 ;

- **Het** représente un cycle saturé ou partiellement insaturé ou aromatique à cinq ou six chaînons contenant de un à trois hétéroatomes du groupe N, O et S qui peuvent être identiques ou différents et qui peuvent être substitués par un ou deux -R^5 ;

- Q^3 est -R^1 ou -OR^1 ;



- **G** est choisi dans le groupe constitué par $\text{-(CH}_2\text{)}_m\text{-}$, -O- , -S- et -NR^1 ;

- **Z** est choisi dans le groupe constitué par -R^1 , $\text{C}_1\text{-C}_4$ alkylène, $\text{C}_1\text{-C}_4$ alcylyne, $\text{-Si(R}^1\text{)}_3$, $\text{(CH}_2\text{)}_p\text{-OMe}$, $\text{-(CH}_2\text{)}_p\text{-SMe}$,





- **X₁** et **X₂** sont choisis indépendamment dans le groupe constitué par hydrogène, halogène, -CF_3 , un groupement cyano et un groupement nitro ;


- Q^1 est choisi dans le groupe constitué par $\text{-(CH}_2\text{)}_q\text{-}$



- **Q²** est choisi dans le groupe constitué par $\text{---}(\text{O})_n\text{---R}^5$, un groupement cyano,



et



- **R¹** est hydrogène ou C₁-C₄ alkyle ;
- **R²** est choisi dans le groupe constitué par hydrogène, halogène, C₁-C₄ alkyle, C₁-C₄ alcoxy, C₁-C₄ halogénoalkyle et C₁-C₄ halogénoalcoxy ;
- **R³** est choisi dans le groupe constitué par hydrogène, C₁-C₄ alkyle et C₁-C₄ alcoxyalkyle ;
- **R⁴** est choisi dans le groupe constitué par halogène, C₁-C₄ alkyle et C₁-C₄ alcoxyalkyle ;
- **R⁵ et R⁶** sont indépendamment l'un de l'autre choisis dans le groupe constitué par hydrogène, halogène, C₁-C₄ alkyle, C₁-C₄ halogénoalkyle ;
- **R⁷ et R⁸** sont indépendamment l'un de l'autre choisis dans le groupe constitué par hydrogène et halogène ;
- **n** est 0 ou 1 ;
- **j, p, q et t** sont choisis indépendamment comme étant 0, 1, 2, 3 ou 4 ;
- **m** est choisi comme étant 1, 2, 3 ou 4 ;
- **r** est 0, 1, 2 ou 3

et les N-oxydes, isomères géométriques et/ou optiques, énantiomères et/ou diastéréoisomères, formes tautomères, sels, complexes métalliques et métalloïdes éventuels des composés de formule (I) tels que l'on vient de les définir ; à condition que le composé de formule (I) soit différent de :

- 3-méthoxy-N-[4-(4-méthylphénoxy)phényl]-4-pipéridin-1-ylpyridine-2-carboxamide ;
- 3-hydroxy-N-[4-(4-méthylphénoxy)phényl]-4-pipéridin-1-ylpyridine-2-carboxamide ;
- 3-éthoxy-[4-(4-méthylphénoxy)phényl]-4-pipéridin-1-ylpyridine-2-carboxamide ;


- N-[4-(benzyloxy)phényl]-3-(méthylthio)-4-pipéridin-1-ylpyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-4-pipéridin-1-yl-N-[4-[3-(trifluorométhyl)phénoxy]phényl]pyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-4-pyrrolidin-1-yl-N-[4-[3-(trifluorométhyl)phénoxy]phényl]pyridine-2-carboxamide ;
- 3-mercapto-4-pipéridin-1-yl-N-[4-[3-(trifluorométhyl)-phénoxy]phényl]pyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-N-[4-(4-méthylphénoxy)phényl]-4-pipéridin-1-ylpyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-N-[4-(4-méthylphénoxy)phényl]-4-thiomorpholin-4-ylpyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-N-[4-(4-méthylphénoxy)phényl]-4-pipéridin-1-ylpyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-N-[4-(4-méthylphénoxy)phényl]-4-(4-méthylpipéridin-1-yl)pyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-N-[4-(4-méthylphénoxy)phényl]-4-(3,5-diméthylpipéridin-1-yl)pyridine-2-carboxamide ;
- 3-[(4-méthoxybenzyl)thio]-N-[4-(4-méthylphénoxy)phényl]-4-(3-méthylpipéridin-1-yl)pyridine-2-carboxamide ;
- 3-mercapto-N-[4-(4-méthylphénoxy)phényl]-4-(4-méthylpipéridin-1-yl)pyridine-2-carboxamide et
- 3-mercapto-N-[4-(4-méthylphénoxy)phényl]-4-(3-méthylpipéridin-1-yl)pyridine-2-carboxamide.

2. Composé selon la revendication 1, **caractérisé en ce que** Y est Y².

3. Composé selon la revendication 2, **caractérisé en ce que** Y est choisi dans le groupe constitué de pyrroles, de diméthylpyrroles et d'imidazole.

4. Composé selon l'une quelconque des revendications précédentes, **caractérisé en ce que** G est -O-.

5. Composé selon l'une quelconque des revendications précédentes, **caractérisé en ce que** Q¹ est choisi dans le groupe constitué par



6. Composé selon la revendication 1, **caractérisé en ce que** :

- Y est Y²,
- G est O,
- et Q¹ est



7. Procédé de préparation de composé tel que défini dans l'une quelconque des revendications 1 à 6, **caractérisé en ce qu'il** comprend :

- à titre de première étape, le schéma réactionnel S- I :

5

10

15

20





Schéma S-I

25

30

35

40




dans lequel :

45

- W_1 est choisi parmi le brome, le chlore et l'iode ;
- W_2 est choisi parmi le fluor, le chlore, le groupement -O-DMG, le groupement -S-DMG et le groupement -NR¹-DMG ; dans lesquels -DMG représente un groupe ortho-directeur de métallation,
- k , R^1 , R^5 , R^6 , X^1 , X^2 , G , Z , Q^1 et Q^2 sont tels que dans l'une quelconque des revendications 1 à 6 ;
- à titre de deuxième étape, la préparation du composé de formule générale (Ic)

50


55




(Ic)

dans laquelle Y^2 , X^1 , X^2 , G , Z , Q^1 et Q^2 sont tels que définis dans l'une quelconque des revendications 1 à 6 ;
lequel composé peut être préparé en faisant réagir le composé de formule générale (Ia) avec une amine primaire (R_1-NH_2), une hydrazine ($R_4R_5N-NH_2$) ou une hydroxylamine (R_2O-NH_2), éventuellement en présence d'un acide de Lewis et/ou d'un agent de captage d'eau tel qu'un tamis moléculaire.

8. Procédé de préparation du composé de formule générale (Ib)



dans laquelle Y^2 , X^1 , X^2 , G , Z , Q^1 et Q^2 sont tels que définis dans l'une quelconque des revendications 1 à 6 et le groupement Het est lié directement au reste de la structure (c'est-à-dire $k=0$) par un atome d'azote, **caractérisé en ce qu'un** composé de formule générale (III) est mis en réaction selon le schéma S-II :



dans lequel Y^2 , X^1 , X^2 , G , Z , Q^1 et Q^2 sont tels que définis dans l'une quelconque des revendications 1 à 6.

9. Composition fongicide comprenant une quantité efficace d'un composé selon l'une quelconque des revendications 1 à 6 et un support acceptable en agriculture.

10. Composition fongicide selon la revendication 9, comprenant en outre un tensioactif.

11. Composition fongicide selon l'une ou l'autre des revendications 9 et 10, comprenant de 0,05% à 99% en poids de matière active.

12. Méthode de lutte préventive ou curative contre les champignons phytopathogènes des cultures, **caractérisée en ce qu'une** quantité efficace et non phytotoxique d'une composition selon l'une quelconque des revendications 9 à 11 est appliquée aux semences de plantes ou aux feuilles de plantes et/ou aux fruits des plantes ou au sol dans lequel poussent les plantes ou dans lequel on souhaite les faire pousser.

13. Méthode selon la revendication 12, **caractérisée en ce que** la dose de matière active appliquée est comprise entre 10 g et 800 g de matière active par hectare dans le cas de traitements foliaires.

EP 1 404 666 B1

14. Méthode selon la revendication 13, **caractérisée en ce que** la dose de matière active appliquée est comprise entre 50 g et 300 g de matière active par hectare dans le cas de traitements foliaires.
15. Méthode selon la revendication 12, **caractérisée en ce que** la dose de matière active appliquée est comprise entre 2 et 200 g de matière active par 100 kg de semences, dans le cas de traitements de semences.
16. Méthode selon la revendication 15, **caractérisée en ce que** la dose de matière active appliquée est comprise entre 3 g et 150 g par 100 kg de semences, dans le cas de traitements de semences.

5

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9911127 A [0002]
- US 3228950 A [0002]
- WO 0026191 A [0003]
- WO 9525723 A [0003]
- WO 0114339 A [0004]
- JP 11228542 B [0005]
- EP 0690061 A [0006]
- WO 0149667 A [0018] [0068]
- WO 9102071 A [0041]
- WO 9506128 A [0041]
- US 4535060 A [0042]
- US 4769061 A [0042]
- US 5094945 A [0042]
- US 4940835 A [0042]
- US 5188642 A [0042]
- US 4971908 A [0042]
- US 5145783 A [0042]
- US 5310667 A [0042]
- US 5312910 A [0042]
- US 5627061 A [0042]
- US 5633435 A [0042]
- FR 2736926 [0042]
- US 5463175 A [0042]
- FR 9506800 [0042]
- FR 9513570 [0042]
- EP 0496630 A [0042]
- EP 0496631 A [0042]
- EP 0625505 A [0042]
- EP 0625508 A [0042]
- US 5506195 A [0042]
- WO 9638567 A [0042]
- US 5510471 A [0042]
- WO 9717432 A [0043]
- WO 9808932 A [0043]
- WO 9730082 A [0044]
- FR 9801814 W [0044]
- FR 9801462 W [0044]
- EP 0666918 A [0045]
- WO 9820133 A [0045]
- WO 9741239 A [0045]
- WO 9531554 A [0045]
- WO 9420828 A [0045]
- WO 9214822 A [0045]
- WO 0149666 A [0064] [0066]

Non-patent literature cited in the description

- **Kuzo SHIBATA et al.** *The Journal of Antibiotics*, 1998, vol. 51 (12), 1113-1116 [0002]
- **J Elguero ; C. Martin ; A.R. Katritzky ; P Linda.** *The Tautomerism of Heterocycles, Advances in Heterocyclic Chemistry.* Academic Press, 1976, vol. 1, 1-4 [0010]
- **E. Eliel ; S. Wilen.** *Stereochemistry of Organic Compounds.* Wiley, 1994 [0011]
- **V. Snieckus.** *Chemical Reviews*, 1990, 879-933 [0015]
- **G. Quéguiner.** *Tetrahedron*, 2001, 4059-4090 [0015]
- **W.K. Fife.** *Journal of Organic Chemistry*, 1983, 1375-1377 [0016]
- **A.R Katritzky.** *Handbook of heterocyclic chemistry.* Pergamon, 2000 [0017]
- **Queginer.** *Tetrahedron*, 1983, 2021 [0050]