

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 April 2008 (17.04.2008)

PCT

(10) International Publication Number
WO 2008/045214 A2

(51) International Patent Classification:
C12N 15/09 (2006.01)

(74) Agent: QUERTERMOUS, Elena E.; Danisco Us, Inc.,
Genencor Division, 925 Page Mill Road, Palo Alto, Cali-
fornia 94304 (US).

(21) International Application Number:
PCT/US2007/020851

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG,
ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,
PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(22) International Filing Date:
26 September 2007 (26.09.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/849,982 6 October 2006 (06.10.2006) US

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL,
PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US):
DANISCO US, INC., GENENCOR DIVISION
[US/US]; 925 Page Mill Road, Palo Alto, California
94304 (US).


(72) Inventor; and

(75) Inventor/Applicant (for US only): **CERVIN, Marguerite**
A. [CA/US]; Danisco Us, Inc., Genencor Division, 925
Page Mill Road, Palo Alto, California 94304 (US).

Published:

— without international search report and to be republished
upon receipt of that report

(54) Title: CELLULASE-FREE ENZYME COMPOSITIONS AND HOST CELLS FOR PRODUCING THE SAME



(57) Abstract: The present invention provides recombinant bacterial cells for producing a detergent-additive protein. In some embodiments, the cells are of the genus *Bacillus*. In additional embodiments, the cells comprise a genome comprising an inactivated *bglC* gene, as well as a recombinant nucleic acid for production of at least one secreted detergent-additive protein. In some preferred embodiments, the secreted detergent-additive protein is a protease. The present invention also provides methods of using the bacterial cells to produce at least one detergent-additive protein, as well as cellulase-free compositions containing at least one detergent-additive protein.

WO 2008/045214 A2

CELLULASE-FREE ENZYME COMPOSITIONS AND HOST CELLS
FOR PRODUCING THE SAME

5 FIELD OF THE INVENTION

[01] The present invention provides recombinant bacterial cells for producing a detergent-additive protein. In some embodiments, the cells are of the genus *Bacillus*. In additional embodiments, the cells comprise a genome comprising an inactivated *bglC* gene, as well as a recombinant nucleic acid for production of at least one secreted detergent-additive protein. In some preferred embodiments, the
10 secreted detergent-additive protein is a protease. The present invention also provides methods of using the bacterial cells to produce at least one detergent-additive protein, as well as cellulase-free compositions containing at least one detergent-additive protein.

BACKGROUND

15 [02] Expression and recombinant production of exogenous polypeptides is a widely used technique. It is well known that cells can be transformed with nucleic acids encoding exogenous polypeptides of interest for expression and production of large quantities of the desired polypeptides. In some applications, the methods are used to produce vast amounts of polypeptide over what would be produced naturally by the originating organism. Indeed, expression of exogenous nucleic acid sequences, as well as
20 over-expression of endogenous sequences have been extensively used in modern biotechnology.

[03] In some cases, undesirable products are produced along with the protein of interest. For example, in production of a recombinant enzyme (e.g., a protease, amylase or the like), other enzymes that are undesirable (e.g., cellulases) may also be produced.

[04] Despite advances in molecular biology and protein engineering, there remains a need for
25 methods and compositions that reduce, if not eliminate such undesirable activities.

SUMMARY OF THE INVENTION

[05] The present invention provides recombinant bacterial cells for producing a detergent-additive protein. In some embodiments, the cells are of the genus *Bacillus*. In additional embodiments, the cells
30 comprise a genome comprising an inactivated *bglC* gene, as well as a recombinant nucleic acid for production of at least one secreted detergent-additive protein. In some preferred embodiments, the secreted detergent-additive protein is a protease. The present invention also provides methods of using the bacterial cells to produce at least one detergent-additive protein, as well as cellulase-free compositions containing at least one detergent-additive protein.

35 [06] The present invention provides recombinant *Bacillus* sp. host cells comprising a genome comprising an inactivated *bglC* gene, wherein the cell further comprises a recombinant nucleic acid for production of a secreted detergent-additive protein. In some preferred embodiments, the inactivated *bglC*

gene contains a deletion, an insertion, a substitution or a rearrangement. In some alternative preferred embodiments, the *Bacillus* sp. cell is a *B. licheniformis*, *B. subtilis*, *B. clausii*, *B. alkalophilus* or *B. halodurans* cell. In still further preferred embodiments, the secreted detergent-additive protein is an enzyme selected from a protease, an amylase, a pectate lyase and a lipase. In some particularly preferred
5 embodiments, the enzyme is a protease. In some more particularly preferred embodiments, the protease is a subtilisin. In yet further preferred embodiments, the *bglC* gene encodes a polypeptide that is at least 80% identical to SEQ ID NO:2.

[07] The present invention also provides cultures of cells comprising culture medium and a recombinant *Bacillus* sp. host cell comprising a genome comprising an inactivated *bglC* gene, wherein
10 the cell further comprises a recombinant nucleic acid for production of a secreted detergent-additive protein. In some preferred embodiments, the inactivated *bglC* gene contains a deletion, an insertion, a substitution or a rearrangement. In some alternative preferred embodiments, the *Bacillus* sp. cell is a *B. licheniformis*, *B. subtilis*, *B. clausii*, *B. alkalophilus* or *B. halodurans* cell. In still further preferred
15 embodiments, the secreted detergent-additive protein is an enzyme selected from a protease, an amylase, a pectate lyase, an acyltransferase, an arylesterase and a lipase. In some particularly preferred
embodiments, the enzyme is a protease. In some more particularly preferred embodiments, the protease is a subtilisin. In yet further preferred embodiments, the *bglC* gene encodes a polypeptide that is at least 80% identical to SEQ ID NO:2. .

[08] The present invention further provides methods comprising maintaining the culture of cells under
20 conditions suitable to produce the secreted detergent-additive protein. In some embodiments, the methods further comprise recovering the secreted detergent-additive protein from the culture medium. In additional embodiments, the methods further comprise combining the secreted detergent-additive protein with a laundry detergent. In still further preferred embodiments, the detergent-additive protein is a subtilisin protease.

[09] The present invention also provides a cellulase-free protein or enzyme composition produced by the methods set forth herein. In some preferred embodiments, the composition comprises a secreted
25 detergent-additive protein produced by a *Bacillus* sp. and is characterized in that the composition is does not have detectable cellulase activity.

[010] The present invention further provides a cellulase-free laundry detergent comprising a secreted
30 detergent-additive protein produced by the methods set forth herein. In some embodiments, the cellulase-free laundry detergent comprises a cellulosic polymer.

[011] The present invention also provides methods for making a host cell, comprising introducing a first recombinant nucleic acid into a *Bacillus* sp. cell so that the recombinant nucleic acid recombines
35 with the *bglC* gene of the cell; and introducing a second recombinant nucleic acid into the cell that provides for expression of the secreted detergent-additive protein. In some preferred embodiments, the nucleic acid inserts into the *bglC* gene. In some alternative preferred embodiments, the nucleic acid

deletes at least a portion of the *bglC* gene. In still further preferred embodiments, the secreted detergent-additive protein is a subtilisin.

BRIEF DESCRIPTION OF THE DRAWINGS

- 5 [012] Certain aspects of the following detailed description are best understood when read in conjunction with the accompanying drawings. It is emphasized that, according to common practice, the various features of the drawings are not to-scale. On the contrary, the dimensions of the various features are arbitrarily expanded or reduced for clarity. Included in the drawings are the following figures:
- 10 [013] Figures 1A-B provide the strategy employed to construct a *Bacillus* host cell containing an inactivated *bglC* gene.
- [014] Figure 2 provides a graph showing the results of viscosity assays on the culture supernatants of two *Bacillus subtilis* strains having an inactivated *bglC* gene (i.e., "host A" and "host B") and two *Bacillus subtilis* strains having an inactivated *bglS* genes (i.e., "host C" and "host D"). Fluid viscosity is measured in centipoises (i.e., cP).
- 15 [015] Fig. 3 is a graph showing the results of viscosity assays on the culture supernatants of two *Bacillus subtilis* strains producing recombinant subtilisin. Strain MDT-05-28 has an inactivated *bglC* gene whereas strain MTD-04-250 contains wild type *bglC* gene.

DESCRIPTION OF THE INVENTION

- 20 [016] The present invention provides recombinant bacterial cells for producing a detergent-additive protein. In some embodiments, the cells are of the genus *Bacillus*. In additional embodiments, the cells comprise a genome comprising an inactivated *bglC* gene, as well as a recombinant nucleic acid for production of at least one secreted detergent-additive protein. In some preferred embodiments, the secreted detergent-additive protein is a protease. The present invention also provides methods of using
- 25 the bacterial cells to produce at least one detergent-additive protein, as well as cellulase-free compositions containing at least one detergent-additive protein.
- [017] Unless otherwise indicated, the practice of the present invention involves conventional techniques commonly used in molecular biology, microbiology, protein purification, protein engineering, protein and DNA sequencing, and recombinant DNA fields, which are within the skill of the art. Such
- 30 techniques are known to those of skill in the art and are described in numerous texts and reference works (See e.g., Sambrook et al., "Molecular Cloning: A Laboratory Manual", Second Edition (Cold Spring Harbor), [1989]); and Ausubel et al., "Current Protocols in Molecular Biology" [1987]). All patents, patent applications, articles and publications mentioned herein, both supra and infra, are hereby expressly incorporated herein by reference.
- 35 [018] Furthermore, the headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the

specification as a whole. Nonetheless, in order to facilitate understanding of the invention, a number of terms are defined below.

Definitions

5 [019] Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. For example, Singleton and Sainsbury, Dictionary of Microbiology and Molecular Biology, 2d Ed., John Wiley and Sons, NY (1994); and Hale and Marham, The Harper Collins Dictionary of Biology, Harper Perennial, NY (1991) provide those of skill in the art with a general dictionaries of many of the terms
10 used herein. Although any methods and materials similar or equivalent to those described herein find use in the practice of the present invention, the preferred methods and materials are described herein. Accordingly, the terms defined immediately below are more fully described by reference to the Specification as a whole. Also, as used herein, the singular terms "a," "an," and "the" include the plural reference unless the context clearly indicates otherwise. Thus, for example, reference to a "host cell"
15 includes a plurality of such host cells. Likewise, reference to "a gene" includes a plurality of such candidate agents and reference to "the cell" includes reference to one or more cells and equivalents thereof known to those skilled in the art, and so forth.

[020] Numeric ranges are inclusive of the numbers defining the range. Indeed, it is intended that every maximum numerical limitation given throughout this specification includes every lower numerical
20 limitation, as if such lower numerical limitations were expressly written herein. Every minimum numerical limitation given throughout this specification will include every higher numerical limitation, as if such higher numerical limitations were expressly written herein. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein. Where a range
25 of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limits of that range is also specifically disclosed. The upper and lower limits of these smaller ranges may independently be included or excluded in the range, and each range where either, neither or both limits are included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the
30 stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[021] All documents cited are, in relevant part, incorporated herein by reference; the citation of any document is not to be construed as an admission that it is prior art with respect to the present invention. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate
35 such publication by virtue of prior invention. All patents and publications, including all sequences disclosed within such patents and publications, referred to herein are expressly incorporated by reference.

[022] Although any suitable methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, exemplary and preferred methods and materials are now described.

5 [023] Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. The headings provided herein are not limitations of the various aspects or embodiments of the invention that can be had by reference to the specification as a whole. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary, depending upon the context they are used by those of skill in the art. Accordingly, the terms defined immediately below are
10 more fully defined by reference to the Specification as a whole.

[024] As used herein, the term "recombinant" refers to a polynucleotide or polypeptide that does not naturally occur in a host cell. A recombinant molecule may contain two or more naturally-occurring sequences that are linked together in a way that does not occur naturally. A recombinant cell contains a recombinant polynucleotide or polypeptide.

15 [025] As used herein, the term "heterologous" refers to elements that are not normally associated with each other. For example, if a host cell produces a heterologous protein, that protein that is not normally produced by that host cell. Likewise, a promoter that is operably linked to a heterologous coding sequence is a promoter that is operably linked to a coding sequence that it is not usually operably linked to in a wild-type host cell.

20 [026] As used herein, the term "homologous," when used in reference to a polynucleotide or protein, refers to a polynucleotide or protein that occurs naturally in a host cell.

[027] The terms "protein" and "polypeptide" are used interchangeably herein.

[028] As used herein, a "signal sequence" is a sequence of amino acids present at the N-terminal portion of a protein which facilitates the secretion of the mature form of the protein outside the cell. The
25 definition of a signal sequence is a functional one. The mature form of the extracellular protein lacks the signal sequence which is cleaved off during the secretion process.

[029] As used herein, a "coding sequence" is a DNA segment that encodes a polypeptide.

[030] As used herein, an "inactivated gene" is a locus of a genome that, prior to its inactivation, was capable of producing a protein (i.e., capable of being transcribed into an RNA that could be translated to
30 produce a full length polypeptide). A gene encoding an enzyme is inactivated when it not transcribed and translated into full length catalytically active protein. A gene may be inactivated by altering a sequence required for its transcription, for example by altering a sequence required for RNA processing (e.g., poly-A tail addition), or by altering a sequence required for translation. Examples of inactivated genes include but are not limited to a deleted gene, a gene containing a deleted region, a gene containing a rearranged
35 region, a gene having an inactivating point mutation or frameshift, and a gene containing an insertion. In addition, a gene may also be inactivated using antisense or any other method that abolishes expression of that gene.

[031] As used herein, the term "nucleic acid" encompasses DNA, RNA, whether single stranded or double stranded, and encompasses chemically modified DNA or RNA. The terms "nucleic acid" and "polynucleotide" are used interchangeably herein.

[032] As used herein, the term "vector" refers to a polynucleotide designed to introduce nucleic acids into one or more host cells. In preferred embodiments, vectors autonomously replicate in different host cells. The term is intended to encompass, but is not limited to cloning vectors, expression vectors, shuttle vectors, plasmids, phage particles, cassettes, and the like.

[033] An "expression vector" as used herein refers to a DNA construct comprising a protein-coding region that is operably linked to a suitable control sequence capable of effecting expression of the protein in a suitable host cell. In some embodiments, such control sequences include a promoter to effect transcription, an optional operator sequence to control transcription to produce mRNA, a sequence encoding suitable ribosome binding sites on the mRNA, and enhancers and sequences which control termination of transcription and translation.

[034] As used herein, the term "promoter" refers to a regulatory sequence that initiates transcription of a downstream nucleic acid.

[035] As used herein, the term "operably linked" refers to an arrangement of elements that allows them to be functionally related. For example, a promoter is operably linked to a coding sequence if it controls the transcription of the sequence.

[036] As used herein, the term "selective marker" refers to a protein capable of expression in a host that allows for ease of selection of those hosts containing an introduced nucleic acid or vector. Examples of selectable markers include but are not limited to antimicrobials (e.g., hygromycin, bleomycin, or chloramphenicol) and/or genes that confer a metabolic advantage, such as a nutritional advantage on the host cell.

[037] As used herein, the term "derived" encompasses the terms "originated from," "obtained," or "obtainable from," and "isolated from".

[038] As used herein, a "non-pathogenic" organism is an organism that is not pathogenic to humans and/or other animals.

[039] The terms "recovered," "isolated," and "separated," as used herein refer to a protein, cell, nucleic acid or amino acid that is removed from at least one component with which it is naturally associated.

[040] As used herein, the terms "transformed," "stably transformed," and "transgenic" when used in reference to a cell means the cell has a non-native (e.g., heterologous) nucleic acid sequence integrated into its genome or as an episomal plasmid that is maintained through multiple generations.

[041] As used herein, the term "expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

[042] As used herein, the term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection," "transformation," or "transduction," and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell wherein the nucleic acid

sequence may be incorporated into the genome of the cell (e.g., chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA).

[043] As used herein, the term “hybridization” refers to the process by which a strand of nucleic acid joins with a complementary strand through base pairing as known in the art. A nucleic acid is considered to be “selectively hybridizable” to a reference nucleic acid sequence if the two sequences specifically hybridize to one another under moderate to high stringency hybridization and wash conditions. Moderate and high stringency hybridization conditions are known to those of skill in the art (See e.g., Ausubel et al., Short Protocols in Molecular Biology, 3rd ed., Wiley & Sons, Hoboken, NJ [1995]; and Sambrook et al., Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor, NY [2001]). One example of high stringency conditions include hybridization at about 42°C in 50% formamide, 5X SSC, 5X Denhardt’s solution, 0.5% SDS and 100 ug/ml denatured carrier DNA followed by washing two times in 2X SSC and 0.5% SDS at room temperature and two additional times in 0.1 X SSC and 0.5% SDS at 42°C.

[044] As used herein, the term “cellulosic polymer” refers to cellulose, hemicellulose or modified cellulose or hemicellulose polymers that comprise at least one 1,4-beta-D glucosidic linkage.

[045] As used herein, the term “cellulose” refers to a polysaccharide polymer comprising glucose residues joined by beta-1,4 linkages.

[046] As used herein, the term “hemicellulose” refers to a polysaccharide polymer comprising at least one non-glucose saccharide residue (e.g., xylose, galactose, arabinose, rhamnose, mannose, uronic acid or galacturonic acid, or xylans), joined by a beta-(1-4) linkage. Hemicelluloses include xylan, glucuronoxylan, arabinoxylan, arabinogalactan, glucomannan, xyloglucan, and galactomannan.

[047] As used herein, “cellulase” refers to an enzyme that hydrolyzes the 1,4-beta-D-glucosidic linkages in cellulose, lichen and cereal beta-D-glucans. The cellulase described herein has an activity described as EC 3.2.1.4, according to IUBMB enzyme nomenclature. The systematic name for the cellulase described herein is 1,4-(1,3;1,4)-beta-D-glucan 4-glucanohydrolase.

[048] As used herein, the term “Bacillus sp.” (e.g., a Bacillus host cell) refers to any species of the genus Bacillus including but not limited to B. subtilis, B. licheniformis, B. lentus, B. brevis, B. stearothermophilus, B. alkalophilus, B. amyloliquefaciens, B. clausii, B. halodurans, B. megaterium, B. coagulans, B. circulans, B. lautus, and B. thuringiensis, as well as sub-species thereof.

[049] As used herein, “cellulase-free Bacillus sp.” refers to a genetically engineered Bacillus sp. host cell that does not secrete a detectable amount of cellulase. As will be discussed in greater detail below, in certain embodiments, a cellulase-free Bacillus strain may contain an inactivated bglC gene.

[050] As used herein, an “equivalent unaltered Bacillus sp. strain” refers to the host strain that is otherwise identical to a cellulase-free Bacillus sp. strain, except the bglC gene is not altered (i.e., is wild-type).

[051] As used herein, "culturing" refers to growing a population of microbial cells under suitable conditions in a liquid or solid medium. In some embodiments, culturing refers to fermentative recombinant production of an exogenous protein of interest or other desired end products (typically in a vessel or reactor).

5 [052] As used herein, a "detergent-additive protein" refers to a protein that is to be added to laundry detergent. A detergent-additive protein may be an enzyme (e.g., a protease, amylase, pectate lyase, lipase, acyltransferase, arylesterases or a protein that does not have enzymatic activity) In some particularly preferred embodiments, beta-glucan hydrolases (i.e., enzymes having an activity described as EC 3.2.1.8, EC 3.2.1.32, EC 3.2.1.72, EC 3.2.1.136, according to IUBMB enzyme nomenclature), are specifically
10 excluded from the term "detergent-additive protein." In further particularly preferred embodiments, xylanases (i.e., enzymes having an activity described as EC 3.2.1.75, according to IUBMB enzyme nomenclature) are also specifically excluded from the term "detergent-additive protein."

[053] Other definitions of terms may appear throughout the Specification.

[054] Before the exemplary embodiments are described in more detail, it is to be understood that the
15 present invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

Host Cells

[055] As noted above, a *Bacillus* sp. host cell that produces reduced (e.g., undetectable) levels of
20 cellulase is provided by the present invention. In general terms, the subject *Bacillus* sp. host cell typically produces less than about 50% (e.g., less than about 40%, less than about 30%, less than about 20%, or less than about 10%) of the cellulase of an equivalent wild-type *Bacillus* sp. host cell. In some embodiments, the subject cell produces less than about 5% of the cellulase of an equivalent *Bacillus* sp. host cell. In some particularly preferred embodiments, the cellulase is undetectable (i.e., the *Bacillus* sp.
25 host cell is a cellulase-free *Bacillus* sp. host cell). It is intended that cellulase activity be assessed by any suitable method known, for example, by staining cellulose-containing LP agar plates with Congo red (See e.g., Wolf et al., *Microbiol.*, 141:281-290 [1995]; and Carder, *Anal. Biochem.*, 153:75-9 [1986]), and/or by using a viscosity assay, as described in more detail below.

[056] In some embodiments, a *Bacillus* sp. host cell that produces a reduced amount of cellulase is
30 produced by reducing the expression of the *bglC* gene product by the cell. In such embodiments, *bglC* expression is reduced in a *Bacillus* sp. host cell using any suitable method, including but not limited to methods that employ antisense molecules, or ribozymes, for example. In some preferred embodiments, expression of *bglC* is reduced by inactivating the *bglC* gene in the cell.

[057] The DNA sequences of several *Bacillus* sp. *bglC* genes and the proteins encoded by those genes
35 have been determined and deposited in NCBI's Genbank database. The sequence is provided below:

ATGAAACGGTCAATCTCTATTTTTATTACGTGTTTATTGATTACGTTATTGACAATGG
 GCGGCATGATAGCTTCGCCGGCATCAGCAGCAGGGACAAAAACGCCAGTAGCCAAG
 AATGGCCAGCTTAATAAAAGGTACACAGCTCGTTAACCGAGACGGTAAAGCGGTAC
 AGCTGAAGGGGATCAGTTCACACGGATTGCAATGGTATGGAGAATATGTCAATAAA
 5 GACAGCTTAAATGGCTGAGAGATGATGGGTATCACCGTTTTCCGTGCAGCGATGTA
 TACGGCAGATGGCGGTTATATTGACAACCCGTCCGTGAAAAATAAAGTAAAAGAAG
 CGGTTGAAGCGGCAAAAGAGCTTGGGATATATGTCATCATGATGGCATATCTTAAAT
 GACGGTAATCCAAACCAAAATAAAGAGAAGGCCAAAAGAATTCTTCAAGGAAATGTC
 AAGCCTTTACGGAACACGCCAAACGTCATTTATGAAATTGCAAACGAACCAACGG
 10 GATGTGAACTGGAAGCGTGATATTAAACCATATGCGGAAGAAGTGATTTTCAGTTATC
 CGCAAAAATGATCCAGACAACATCATCATTGTCGGAACCGGTACATGGAGCCAGGA
 TGTGAATGATCTGCGATGACCAGCTAAAAGATGCAAACGTTATGTACGCACTTCATT
 TTTATGCCGGCACGCACGGCCAATTTTTACGGGATAAAGCAAACCTATGCACTCAGCA
 AAGGAGCACCTATTTTTGTGACGAGTGGGAACAAGCGACGCGTCTGGCAATGGCGG
 15 TGTATTCCTTGATCAATCGAGGGAATGGCTGAAATATCTCGACAGCAAGACCATTAG
 CTGGGTGAACTGGAATCTTTCTGATAAGCAGGAATATCCTCGCTTTAAAGCCGGGGG
 CATCTAAAACAGGCGGCTGGCGGTTGTCAGATTTATCTGCTTCAGGAACATTCTGTTA
 GAGAAAACATTCTCGGCACCAAAGATTTCGACGAAGGACATTCCTGAACGCCATCAA
 AGATAAACCCACACAGGAAAATGGTATTTCTGTACAGTACAGAGCAGGGGATGGGA
 20 GTATGAACAGCAACCAAATCCGTCCGCAGCTTCAAATAAAAAATAACGGCAATACC
 ACGGTGATTTAAAGATGTCACTGCCCGTTACTGGTATAAAGCGAAAAACAAAGGCC
 AAAACTTTGACTGTGACTACGCGCAGATTGGATGCGGCAATGTGACACACAAGTTTG
 TGACGTTGCATAAACCAAGCAAGGTGCGATACCTATCTGGAACCTTGATTAAAAAC
 GGAACGTTGGCACCGGGAGCAAGCACAGGGAATATTCAGCTCCGTCTTCACAATGA
 25 TGACTGGAGCAATTATGCACAAAGCGGCGATATTCCTTTTAAATCAAATACGTTTA
 AAACAACGAAAAAAATCACATTATATGATCAAGGAAAACCTGATTTGGGGAACAGAA
 CCAAATTAG (SEQ ID NO:1)

[058] The following is the amino acid sequence encoded by SEQ ID NO:1.

30 MKRSISIFITCLLITLLTMGGMIASPASAAGTKTPVAKNGQLSIKGTQLVNRDGVKAVQLKGISSHG
 LQWYGEYVNKDSLKWLRDDWGITVFRAAMYTADGGYIDNPSVKNKVKEAVEAAKELGIYVI
 WHILNDGNPNQNKEKAKEFFKEMSSLYGNTPNVIYEIANEPNGDVNWKRDIKPYAEEVISVIRK
 NDPDNIIIVGTGTWSQDVNDAADDQLKDANVMYALHFYAGTHGQFLRDKANYALSKGAPIFVE
 35 GTSDASGNNGGVFLDQSREWLYLDSKTISWVNWNLSDKQESSALKPGASKTG GWRLSDLSAS
 GTFVRENILGTKDSTKDIPETPSKDKPTQENGISVQYRAGDGSMSNSNQIRPQLQIKNNGNTTDL
 VTARYWYKAKNKGQNFDCDYAQIGCGNVTHKFVTLHKPKQGADTYLELGFKNGTLAPGASTG
 NIQLRLHNDDWSNYAQSGDYSFFKSNFTKTTKITLYDQGKLIWGTEPN (SEQ ID NO:2)

40 [059] Further, several conserved domains of cellulase enzymes have been identified, as well as a
 number of conserved amino acids, allowing the identification of further *Bacillus* sp. bglC genes and
 proteins by bioinformatic methods.

[060] In some embodiments, the *Bacillus* sp. bglC gene comprises at least 70% (e.g., at least 80%, at
 least 90%, at least 95%, at least 97% or at least 98% sequence identity) to a bglC sequence deposited in
 45 NCBI's Genbank database and provided above (SEQ ID NO:1). In some further embodiments, the
Bacillus sp. bglC gene hybridizes under stringent conditions to a bglC sequence deposited in NCBI's
 Genbank database or SEQ ID NO:1. In yet additional embodiments, , the *Bacillus* sp. bglC gene encodes
 a polypeptide that has at least 70% sequence identity (e.g., at least 80%, at least 90%, at least 93%, at
 least 95%, at least 97% or at least 98% sequence identity) to a bglC sequence deposited in NCBI's

Genbank database or SEQ ID NO:2. Exemplary bglC protein and nucleotide sequences deposited in NCBI's Genbank database include: GID:3100136 (*Bacillus licheniformis*), GID:52348343 (*Bacillus licheniformis*), GID:42491106 (*Bacillus amyloliquefaciens*) and GID:50812243 (*Bacillus subtilis*). The above Genbank accessions are incorporated by reference in their entirety, including the nucleic acid and protein sequences therein and the annotation of those sequences.

[061] In some preferred embodiments, the *Bacillus* sp. host cell is of any of the following species: *B. licheniformis*, *B. lentus*, *B. subtilis*, *B. amyloliquefaciens*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. coagulans*, *B. circulans*, *B. pumilus*, *B. thuringiensis*, *B. clausii*, or *B. megaterium*. *B. subtilis* host cells include, but not limited to those described in U.S. Patents 5,264,366 and 4,760,025 (RE 34,606), as well as 1A6 (ATCC 39085), 168 (1A01), SB19, W23, Ts85, B637, PB1753 through PB1758, PB3360, JH642, 1A243 (ATCC 39,087), ATCC 21332, ATCC 6051, MI113, DE100 (ATCC 39,094), GX4931, PBT 110, and PEP 211 strain (See e.g., Hoch et al., *Genetics* 73:215–228 [1973]; U.S. Patent No. 4,450,235; U.S. Patent No. 4,302,544; and EP 0134048 (See also, Palva et al., *Gene* 19:81–87 [1982]; and Fahnestock and Fischer, *J. Bacteriol.*, 165:796–804 [1986]; and Wang et al., *Gene* 69:39–47 [1988]).

[062] In some embodiments, the host cell comprises a recombinant nucleic acid comprising an expression cassette (i.e., a promoter, a polynucleotide encoding a detergent-additive protein, and a transcriptional terminator), wherein the expression cassette is sufficient for the production of the detergent-additive protein by the *Bacillus* sp. host cell. In some embodiments, the recombinant nucleic acid is integrated into the genome of the host cell, while in other embodiments, the recombinant nucleic acid is present in a vector that replicates autonomously from the genome. In some embodiments, the polynucleotide encoding the detergent-additive protein is codon optimized for expression of the protein in the *Bacillus* sp. host cell.

[063] In some particularly preferred embodiments, the *Bacillus* host cell is engineered to maximize protein expression. Thus, in some embodiments, the host cells contain an inactivating alteration in at least one of the following genes, *degU*, *degS*, *degR* and/or *degQ* (See, Msadek et al., *J. Bacteriol.*, 172:824–834 [1990]; and Olmos et al., *Mol. Gen. Genet.*, 253:562–567 [1997]). In some particularly preferred embodiments, the host cell is a *B. subtilis* that carries a *degU32(Hy)* mutation. In some additional embodiments, the *Bacillus* host cell comprises a mutation and/or deletion in *scoC4*, (See, Caldwell et al., *J. Bacteriol.*, 183:7329–7340 [2001]); *spoIIE* (See, Arigoni et al., *Mol. Microbiol.*, 31:1407–1415 [1999]); *oppA* or another gene in the *opp* operon (See, Perego et al., *Mol. Microbiol.*, 5:173–185 [1991]). Indeed, it is contemplated that any mutation in the *opp* operon that causes the same phenotype as a mutation in the *oppA* gene will find use in some embodiments of the altered *Bacillus* strain of the present invention. In some embodiments, these mutations occur alone, while in other embodiments, combinations of mutations are present. In some embodiments, an altered *Bacillus* of the invention is obtained from a *Bacillus* host strain that already includes a mutation to one or more of the above-mentioned genes. In

alternate embodiments, an altered *Bacillus* of the invention is further engineered to include mutation of one or more of the above-mentioned genes.

[064] *Bacillus* sp. host cells constructed using any convenient method find use in the present invention, including cells constructed by altering the sequence of the *bglC* gene of the cell by making an insertion, deletion, replacement, frameshift, point mutation, and/or rearrangement in the gene find use in the present invention. In some embodiments, the portion of the gene to be altered is within the coding region or a regulatory element required for expression of the coding region. In some embodiments, the regulatory or control sequence of a gene is a promoter sequence or a functional part thereof (i.e., a part which is necessary for expression of the gene). Such gene inactivation methods are well known in the art (See e.g., Wolf et al *Microbiol.*, 141:281-290 [1995]).

[065] In some embodiments, the host cell is constructed by: introducing a recombinant nucleic acid into a *Bacillus* sp. cell so that the recombinant nucleic acid recombines with the *bglC* gene of the cell's genome and introducing a second recombinant nucleic acid into the cell to provide for expression of a secreted detergent-additive protein. In some embodiments, the nucleic acid inserts into the *bglC* gene, while in other embodiments, it deletes at least a portion of the *bglC* gene.

[066] Suitable methods for introducing polynucleotide sequences into *Bacillus* cells are well known to those of skill in the art (See e.g., Ferrari et al., "Genetics," in Harwood et al. (ed.), *Bacillus*, Plenum Publishing Corp. [1989], pages 57-72; See also, Saunders et al., *J. Bacteriol.*, 157:718-726 [1984]; Hoch et al., *J. Bacteriol.*, 93:1925-1937 [1967]; Mann et al., *Curr. Microbiol.*, 13:131-135 [1986]; and Holubova, *Folia Microbiol.*, 30:97 [1985]; Chang et al., *Mol. Gen. Genet.*, 168:11-115; [1979]; Vorobjeva et al., *FEMS Microbiol. Lett.*, 7:261-263 [1980]; Smith et al., *Appl. Env. Microbiol.*, 51:634 [1986]; Fisher et al., *Arch. Microbiol.*, 139:213-217 [1981]; and McDonald, *J. Gen. Microbiol.*, 130:203 [1984]). Indeed, such methods as transformation including protoplast transformation and congression, transduction, and protoplast fusion are known and suited for use in the present invention.

[067] As also noted above, in addition to producing reduced (e.g., undetectable) levels of cellulase, the present invention provides a *Bacillus* sp. host cell that further contains a recombinant nucleic acid for production of a secreted detergent-additive protein, where a secreted detergent-additive protein is a protein (e.g., an enzyme) that is secreted from the cell and is added to laundry detergent. Exemplary detergent-additive proteins include, but are not limited to proteases (e.g., subtilisins), alpha-amylases, mannanases, cellulases, lyases, acyltransferases, arylesterases and lipases, etc.. In some embodiments, the detergent-additive protein may be expressed by a strain that is the same as the strain from which the detergent-additive protein is derived.

Enzymes

[068] Subtilisins (i.e., extracellular alkaline serine proteases), are of particular interest. Any suitable subtilisin finds use in the present invention (See e.g., Siezen, *Protein Sci.*, 6:501-523 [1997]; Bryan, *Biochim. Biophys. Acta*, 1543:203-222 [2000]; Maurer, *Curr. Op. Biotechnol.*, 2004 15:330-334 [2004];

and Gupta, *Appl. Microbiol. Biotechnol.*, 59:15-32 [2002]). In some embodiments, the subtilisin of interest has an activity described as EC 3.4.4.16, according to IUBMB enzyme nomenclature.

[069] In some embodiments, such a subtilisin has an amino acid sequences that is found in wild-type genomes (i.e., the subtilisin is a naturally-occurring subtilisin), while in other embodiments, the subtilisin is a variant of a naturally-occurring subtilisin. In some preferred embodiments, the variant subtilisin comprises an amino acid sequence that is at least 80%, at least 90%, at least 95% or at least 98% identical to a subtilisin encoded by a wild-type genome. Exemplary subtilisins include, but are not limited to: ALCANASE® (Novozymes), FNA™ (Genencor), SAVINASE® (Novozymes) PURAFECT™ (Genencor), KAP™ (Kao), EVERLASE™ (Novozymes), PURAFECT OxP™ (Genencor), FN4™ (Genencor), BLAP S™ (Henkel), BLAP X™ (Henkel), ESPERASE® (Novozymes), KANNASE™ (Novozymes) and PROPERASE™ (Genencor). In yet additional embodiments, the subtilisin includes, but is not limited to subtilisin 168, subtilisin BPN', subtilisin Carlsberg, subtilisin DY, subtilisin 147, or subtilisin 309 (See e.g., EP414279B; WO89/06279; and Stahl et al., *J. Bacteriol.*, 159:811-818 [1984]). Additional subtilisins and other proteases that find use in the present invention include but are not limited to those described in WO 99/20770; WO 99/20726; WO 99/20769; WO 89/06279; RE 34,606; U.S. Patent No. 4,914,031; U.S. Patent No. 4,980,288; U.S. Patent No. 5,208,158; U.S. Patent No. 5,310,675; U.S. Patent No. 5,336,611; U.S. Patent No. 5,399,283; U.S. Patent No. 5,441,882; U.S. Patent No. 5,482,849; U.S. Patent No. 5,631,217; U.S. Patent No. 5,665,587; U.S. Patent No. 5,700,676; U.S. Patent No. 5,741,694; U.S. Patent No. 5,858,757; U.S. Patent No. 5,880,080; U.S. Patent No. 6,197,567; and U.S. Patent No. 6,218,165.

Detergents and Detergent-Additive Proteins

[070] In some embodiments, the host cells are used to make protein compositions and laundry detergents, where the detergent, in some preferred embodiments, contains at least one cellulosic polymer.

[071] In some embodiments, the host cell is cultured to provide at least one secreted detergent-additive protein into the growth medium in which the cell is growing. In some particularly preferred embodiments, the secreted detergent-additive protein is recovered from the growth medium using any suitable method (e.g., precipitation, centrifugation, affinity, filtration or any other method known in the art). For example, affinity chromatography (Tilbeurgh et al., *FEBS Lett.*, 16:215 [1984]); ion-exchange chromatographic methods (Goyal et al., *Bio. Technol.*, 36:37 [1991]; Fliess et al., *Eur. J. Appl. Microbiol. Biotechnol.*, 17:314 [1983]; Bhikhabhai et al., *J. Appl. Biochem.*, 6:336 [1984]; and Ellouz et al., *Chromatogr.*, 396:307 [1987]), including ion-exchange using materials with high resolution power (See e.g., Medve et al., *J. Chromatogr. A* 808:153 [1998]); hydrophobic interaction chromatography (Tomaz and Queiroz, *J. Chromatogr. A* 865:123 [1999]; two-phase partitioning (Brumbauer et al., *Bioseparation* 7:287 [1999]); ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; and gel filtration (e.g., SEPHADEX® G-75), find use.

[072] In some preferred embodiments, the detergent-additive protein is used without purification from the other components the culture medium. In some of these embodiments, the culture medium is simply concentrated and then used without further purification of the protein from the components of the growth medium. In other embodiments, the culture medium is used without any further modification.

5 [073] The protein compositions produced using the host cells generally contain reduced cellulose, as compared to an equivalent *Bacillus* host cell that contains an unaltered (e.g., wild-type) *bgIC* gene.

[074] In one embodiment, the cellulase content of a composition is evaluated by measuring the change in viscosity of a solution of a cellulosic polymer (e.g., a solution containing 1% carboxymethyl cellulose (CMC)) upon addition of the composition to the solution. Such methods are well known (See e.g.,
10 Manning, *Biochem. Biophys. Meth.*, 5:189-202 [1981]; and Sheperd, *Biochem. J.*, 193:67-74 [1981]). One exemplary viscosity assay that finds use with the present invention is provided in the Examples below.

[075] In some embodiments comprising the protein composition of the present invention, the composition is capable of reducing the viscosity of a solution of cellulosic material by less than 80%
15 (e.g., less than 50%, less than 30%, less than 20% or less than 10%), as compared to the reduction by an equivalent protein composition produced using a *Bacillus* cell having an unaltered *bgIC* gene. In some embodiments, the protein composition does not produce a detectable reduction in the viscosity of a solution of cellulosic material, in which case the protein composition is considered to be a "cellulase-free" protein composition.

20 [076] Cellulase-free protein compositions find use in various settings, including but not limited to laundry detergents, particularly those detergents that contain cellulosic polymers. In some embodiments, laundry detergents comprising the cellulase-free protein composition of the present invention contain from about 1% to 80%, (e.g., 5% to 50%)(by weight) of surfactant. In some embodiments, the surfactant is a non-ionic surfactant, while in other embodiments, it is a cationic surfactant, and in still other
25 embodiments, it is an anionic surfactant, and in further embodiments it is a zwitterionic surfactant, and in still further embodiments, it comprises any mixture thereof (e.g., a mixture of anionic and nonionic surfactants). Exemplary surfactants include, but are not limited to alkyl benzene sulfonate (ABS), including linear alkyl benzene sulfonate and linear alkyl sodium sulfonate, alkyl phenoxy polyethoxy ethanol (e.g., nonyl phenoxy ethoxylate or nonyl phenol), diethanolamine,, triethanolamine and
30 monoethanolamine. Additional descriptions of surfactants that find use in laundry detergents are provided in U.S. Patent Nos. 3,664,961, 3,919,678, 4,222,905, and 4,239,659.

[077] The laundry detergent comprising the cellulase-free enzyme of the present invention finds use in any form (e.g., solid, liquid, gel, etc.). In some embodiments, the laundry detergents further contain a buffer such as sodium carbonate, sodium bicarbonate, or detergent builder, bleach, bleach activator, an
35 enzymes, an enzyme stabilizing agent, suds booster, suppresser, anti-tarnish agent, anti-corrosion agent, soil suspending agent, soil release agent, germicide, pH adjusting agent, non-builder alkalinity source, chelating agent, organic or inorganic filler, solvent, hydrotrope, optical brightener, dye or perfumes.

[078] As noted above, in some embodiments, the laundry detergents contain a cellulosic polymer (e.g., a cellulose polymer) or a modified cellulose polymer. Suitable cellulosic polymers include, but are not limited to anionically modified cellulose, nonionically modified cellulose, cationically modified cellulose, zwitterionically modified cellulose, and mixtures thereof, as well as cellulose, cellulose ethers, cellulose esters, cellulose amides, methyl cellulose, carboxy methyl cellulose, ethyl cellulose, hydroxyl ethyl cellulose, hydroxyl propyl methyl cellulose, ester carboxy methyl cellulose, and any mixture(s) thereof. In some embodiments, a modified cellulose ether polymer (See e.g., U.S. Patent No. 6,833,347), or other cellulosic polymer (See e.g., U.S. Patent Nos. 5,009,800 and 4,661,267) find use in the present invention. In some embodiments, the laundry detergent contains from about 0.1% to 8% by weight (e.g., about 0.5% to 4% or about 1% to 3%, of cellulosic polymer).

EXPERIMENTAL

[079] The following examples provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

[080] In the experimental disclosure which follows, the following abbreviations apply: °C (degrees Centigrade); rpm (revolutions per minute); H₂O (water); aa (amino acid); bp (base pair); kb (kilobase pair); kD (kilodaltons); gm (grams); µg and ug (micrograms); mg (milligrams); ng (nanograms); µl and ul (microliters); ml (milliliters); mm (millimeters); nm (nanometers); µm and um (micrometer); M (molar); mM (millimolar); µM and uM (micromolar); U (units); V (volts); MW (molecular weight); sec (seconds); min(s) (minute/minutes); h(s) and hr(s) (hour/hours); OD₂₈₀ (optical density at 280 nm); OD₄₀₅ (optical density at 405 nm); OD₆₀₀ (optical density at 600 nm); PAGE (polyacrylamide gel electrophoresis); LAS (lauryl sodium sulfonate); SDS (sodium dodecyl sulfate); and Tris (tris(hydroxymethyl)aminomethane).

EXAMPLE 1

Construction of bglC::Spectinomycin Strains

[081] The general strategy for the deletion of the bglC gene is depicted in the diagrams in Figs 1A-B. Briefly, upstream and downstream DNA fragments flanking the bglC gene were amplified by PCR and ligated together in a vector. The insert was opened by digestion with a restriction endonuclease and the spectinomycin cassette, flanked by loxP sites, was ligated in. The resultant plasmid was linearized by digestion with a restriction endonuclease and transformed into Bacillus. By selection with the introduced antimicrobial marker (spectinomycin), the replacement of the gene of interest was accomplished by a

double crossover event. In the case of bglC, the coding region was replaced with the spectinomycin resistance gene that was looped out using the Cre recombinase which recognizes the flanking loxP sites.

[082] DNA constructs were made using PCR technology using the strategy described in WO 03/083125 and the primers described in of Table 1.

5

Table 1. Primers			
Primer Name	Restr. Site in Primer	Primer Sequence	SEQ ID NO:
BglC SacI UF IUF	SacI	acaaatGAGCTCgctggagcattggatggcgcatcc	3
BglC BamHI UR	BamHI	TgatctGGATCCccttcgcatctttggctctacac	4
BglC BamHI DF	BamHI	AaaactGGATCCgggaacagaaccaaattagtaagc	5
bglC Sall DR	Sall	atggtaGTCGACgcaaacgcggctacaatatggctca	6
bglC Uout chk		ttccgcgaggggccggcctactata	7
bglC Dout chk		catattcacaatgcgatggttagagg	8
bglC DF chkdel		Ttatgcacaaagcggcgattattcc	9
SPECchkUR:		Atctcttgccagtcacgttacg	10
bglS xba UF	Xba	ggagtGTCTAGAactgaccagcttccgtttccctg	11
BglS BamHI UR	BamHI	ActaacGGATCCcctgtaactatcatcatcttcctc	12
BglS BamHI DF	BamHI	CaaaaaGGATCCgccaaatgtgaagagcctgctgca	13
bglS Sac DR	Sac	agaggtGAGCTCaccgctgattcccgtatgatgcc	14
bglS Uout chk		Caatatacacaatacagtgctgaaagc	15
bglS DROUT chk		Gcgggaatagcgaatgcttggttcgg	16
bglS DF chk del		Gatgaactgtggaatggcacgggt	17
PloxBamUF		TCGACGGTATCGATAAGCTGGATCCATAAC	18
ploxBamDR		GCCTAGGATGCATATGGGATCCGCATAACTTC	19

[083] The restriction sites are designated as follows: XbaI is TCTAGA (SEQ ID NO:20); BamHI is GGATCC (SEQ ID NO:21); SacI is GAGCTC (SEQ ID NO:22); Asp718 is GGTACC (SEQ ID NO:23); PstI is CTGCAG (SEQ ID NO:24) and HindIII is AAGCTT (SEQ ID NO:25).

10 [084] In this method, 100 μ L PCR reactions were carried out in 150 μ L Eppendorf tubes containing 84 μ L water, 10 μ L PCR buffer, 1 μ L of each primer (i.e., BglC SacI UF and BglC BamHI UR, or loxPBamHI F and loxPBamHI R), 2 μ L of dNTPs, 1 μ L of DNA template (e.g., wild type *Bacillus* chromosomal or control plasmid), and 1 μ L of DNA polymerase. DNA polymerases used included Taq Plus Precision polymerase and Herculanase (Stratagene). Reactions were carried out in a Hybaid

15 PCRExpress thermocycler using the following program. The samples were first heated at 94°C for 5 minutes, then cooled to a 50°C. hold. The DNA polymerase was added at this point. Twenty-five cycles of amplification consisted of 1 minute at 95°C, 1 minute at 50°C and 1 minute at 72°C. A final 10 minutes at 72°C ensured complete elongation. Samples were held at 4°C until analysis. The reactions yielded the flanking DNA cassettes (approximately 1 kb each) and the loxP-spectinomycin cassette with

20 BamHI restriction sites at the 5' and 3' ends.

[085] After completion of the PCR, 10 μ L of each reaction were electrophoresed on an Invitrogen 1.2% agarose E-gel at 60 volts for 30 minutes to check for the presence of a band at the correct size. All the gel electrophoresis methods described herein used these conditions. If a band was present, the remainder of the reaction tube was purified using the Qiagen QIAQUICK® PCR purification kit according to the manufacturer's instructions, then cut with the appropriate restriction enzyme pair. Digests were performed at 37°C for 1 hour as a 20 μ L reaction consisting of 9 μ L of water, 2 μ L of 10xBSA, 2 μ L of an appropriate NEB restriction buffer (according to the 2000-01 NEB Catalog and Technical Reference), 5 μ L of template, and 1 μ L of each restriction enzyme. For example, the bglC upstream fragment was cut with SacI and BamHI in NEB (New England BioLabs) restriction buffer B. The digested fragments were purified by gel electrophoresis and extraction using the Qiagen QIAQUICK® gel extraction kit following the manufacturer's instructions.

[086] Ligation of the fragments into a plasmid vector was done in two steps, using either the Takara ligation kit, following the manufacturer's instructions or T4 DNA ligase (Reaction contents: 5 μ L each insert fragment, 1 μ L cut pUC19 plasmid, 3 μ L T4 DNA ligase buffer, and 1 μ L T4 DNA ligase). First, the cut upstream and downstream fragments were ligated overnight at 15°C into unique restriction sites in the pUC19 plasmid (See, Yanisch-Raman et al., Gene 33:103-119 [1985]) polylinker that had been digested with SacI and gel purified, connecting at the common BamHI site to re-form a circular plasmid. This re-circularized plasmid was transformed into Invitrogen's competent "Top 10" E. coli cells, using the manufacturers One Shot transformation protocol.

[087] Transformants were selected on Luria-Bertani (LB) broth solidified with 1.5% agar (LA) plus 50 ug/ml carbenicillin containing X-gal (Sigma) for blue-white screening. Clones were picked and grown overnight at 37°C in 5mL of Luria Bertani broth (LB) plus 50 ug/ml carbenicillin and plasmids were isolated using Qiagen's QIAQUICK® Mini-Prep kit. Restriction analysis using SacI confirmed the presence of the 2 kb insert. Confirmed plasmids with the insert were cut with BamHI to linearize them in digestion reactions as described above (with an additional 1 μ L of water in place of a second restriction enzyme), treated with 1 μ L calf intestinal or shrimp phosphatases for 1 hour at 37°C to prevent re-circularization, and ligated to the BamHI digested spectinomycin resistance cassette flanked by loxP sites. This ligation mix was transformed into E. coli Top 10 cells, and colonies were selected on LA plates containing using 100 ug/ml spectinomycin. Confirmation of marker insertion in isolated plasmids was done as described above, by restriction analysis with BamHI. Prior to transformation into B. subtilis, the plasmid was linearized with ScaI to ensure a double crossover event.

EXAMPLE 2

Viscosity Assay

- 5 [088] Cells were grown in 250 ml baffled flasks containing 50 mL of growth media suitable for subtilisin production (See, Christianson et al., Anal. Biochem., 223:119–129 [1994]; and Hsia et al., Anal. Biochem., 242:221–227 [1996]). The media were inoculated with 50 μ L of an 8 hour 5mL culture, and grown for 40 hours at 37°C with shaking at 250 RPM. Cultures of Host A (-bglC) Host B (-bglC), Host C (-bglS) and Host D (-bglS) were grown for 48 hours.
- 10 [089] A 1 ml sample was taken from each shake flask at 37 hours. The cells were removed by centrifugation and 60 microliters of the supernatant were subjected to a viscosity assay. Viscosity was measured at t=0, t=22 hours and t=120 hours. Water was used as the control.
- [090] Viscosity was measured using the following method: a 2 ml volume of 1% cellulosic material (carboxymethyl cellulose) was dispensed into cryogenic vials for each experiment. A 60 microliter
- 15 sample was added to 2 ml volume of cellulosic material. After gently mixing, samples were taken for analysis at the appropriate preselected time (e.g., incubation for 20 hours). For analysis, 500 microliters of sample and substrate were pipetted into a sample cup of a viscometer (Brookfield LV DVIII Cone & Plate viscometer with a CP40 waterbath set at 25°C) and the viscometer was programmed so that the SSN (set speed) was at 16 RPM, and the WTI (wait time) was set at 30 seconds (Rheocalc software). After 30
- 20 seconds, a summary sheet was displayed.
- [091] The results of these assays are provided in Figure 2. No decrease in the viscosity was observed over the course of the experiment for the Control, Host A (-bglC) and Host B (-bglC) samples contacted with the 1% cellulosic material, indicating the absence of cellulase activity in the samples. Host strains with deleted bglS genomic sequences exhibited significant cellulase activity as indicated by the decrease
- 25 in viscosity.

EXAMPLE 3

Construction of bglC::Spectinomycin Strains Producing Protease

- [092] The following strains were tested in this experiment, a *Bacillus subtilis* strain making subtilisin protease and the same strain with an insertional inactivation of bglC.
- 30 [093] Strains to be tested were constructed as follows. SC6 strain (spoIIE, amyE, aprE Pxyl:comK) was transformed with chromosomal DNA from a subtilisin-producing strain. Transformants were selected on L agar plates containing chloramphenicol (5 micrograms/ml) and 1.6% skim milk. The presence of subtilisin was confirmed by halo production on plates containing skim milk. The resulting strain,
- 35 MDT04-250, was transformed with chromosomal DNA from a strain with an insertional inactivation of the bglC gene, and transformants were selected on L agar plates containing spectinomycin (100

micrograms/ml). The new strain, MDT05-28, was confirmed to be resistant to chloramphenicol (5 micrograms/ml) and spectinomycin (100 micrograms/ml). The two strains were amplified for protease by growing them sequentially on L agar plates containing chloramphenicol (10 micrograms/ml) and then chloramphenicol (25 micrograms/ml). These amplified strains were grown in shake flasks containing 25 ml of LB (Difco), glucose (0.1%) and chloramphenicol (25 micrograms/ml) in a 250 mL baffled flask. Shake flasks were incubated at 37°C with shaking at 280 rpm and at OD₅₅₀ of 0.8, 1 mL of culture was mixed with 0.5 ml 30 % glycerol and frozen at -70°C. Thirty microliters of the thawed vials were used to inoculate 40 ml of FNII media in 250 ml flasks. Two flasks per strain were used. The shake flasks were incubated at 37°C with shaking at 280 rpm for several days, and samples of the supernatant were collected for viscosity testing and subtilisin analysis.

[094] Supernatants from liquid cultures were harvested after different times during growth (e.g., 16, 48 and 68 h) and assayed for subtilisin as previously described (See, Estell et al., J. Biol. Chem., 260:6518-6521 [1985]) in a solution containing 0.3 mM N-succinyl-L-Ala-L-Ala-L-Pro-L-Phe-p-nitroanilide (Vega Biochemicals), 0.1 M Tris, pH 8.6 at 25°C. The assays measured the increase in absorbance at 410 nm/min due to hydrolysis and release of p-nitroaniline. These strains produced equivalent amounts of subtilisin at equivalent rates.

[095] Supernatant samples from shake flasks of either the parent strain, MDT04-250 or the bglC::spec strain (MDT05-28) were taken at 16, 48 or 68 h after inoculation and incubated with a cellulose substrate (4.5 ml of 5% CMC + 0.5 ml of supernatant) for 1, 24 or 48 h. The activity of cellulases on the cellulose substrate was measured by a drop in viscosity of the cellulose substrate as measured by a Rheometer using software as directed by the manufacturer, as discussed above. The data are presented as a percentage of the control (dH₂O) readings. Samples taken from the shake flasks of MDT05-28 did not show a reduction in viscosity of the cellulose substrate while all the samples from MDT04-250 quickly reduced the viscosity of the cellulose substrate. This indicates that the inactivation of bglC prevents activity on the cellulose substrate. The results of these assays are shown in Figure 3. These results show that the culture supernatant from host Bacillus strains expressing subtilisin that have a disrupted bglC gene exhibit no significant cellulase activity.

[096] While particular embodiments of the present invention have been illustrated and described, it will be apparent to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

[097] All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

[098] Having described the preferred embodiments of the present invention, it will appear to those ordinarily skilled in the art that various modifications may be made to the disclosed embodiments, and that such modifications are intended to be within the scope of the present invention.

[099] Those of skill in the art readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The compositions and methods described herein are representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. It is readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

[0100] The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

[0101] The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

CLAIMS

We claim:

- 5 1. A recombinant *Bacillus* sp. host cell comprising a genome comprising an inactivated *bglC* gene, wherein said cell further comprises a recombinant nucleic acid for production of a secreted detergent-additive protein.
2. The recombinant *Bacillus* sp. host cell of claim 1, wherein said inactivated *bglC* gene
10 contains a deletion, an insertion, a substitution or a rearrangement.
3. The recombinant *Bacillus* sp. host cell of claim 1, wherein said *Bacillus* sp. cell is a *B. licheniformis*, *B. subtilis*, *B. clausii*, *B. alkalophilus* or *B. halodurans* cell.
- 15 4. The recombinant *Bacillus* sp. host cell of claim 1, wherein said secreted detergent-additive protein is an enzyme selected from a protease, an amylase, a pectate lyase, an acyltransferase, an arylesterase and a lipase.
5. The recombinant *Bacillus* sp. host cell of claim 4, wherein said enzyme is a protease.
20 6. The recombinant *Bacillus* sp. host cell of claim 5, wherein said protease is a subtilisin.
7. The recombinant *Bacillus* sp. host cell of claim 1, wherein said *bglC* gene encodes a polypeptide that is at least 80% identical to SEQ ID NO:2.
25 8. A culture of cells comprising:
 culture medium; and
 a recombinant *Bacillus* sp. host cell of claim 1.
- 30 9. A method, comprising:
 maintaining the culture of cells of claim 8 under conditions suitable to produce said secreted detergent-additive protein.
10. The method of claim 9, further comprising:
35 recovering said secreted detergent-additive protein from said culture medium.
11. The method of claim 10, further comprising combining said secreted detergent-additive

protein with a laundry detergent.

12. The method of claim 9, wherein said secreted detergent-additive protein is a subtilisin protease.

13. A cellulase-free protein composition produced by the method of claim 9.

14. The cellulase-free protein composition of claim 13, wherein said composition comprises a secreted detergent-additive protein produced by a *Bacillus sp.* and is characterized in that the composition is does not have detectable cellulase activity.

15. A cellulase-free laundry detergent comprising a secreted detergent-additive protein produced by the method of claim 9.

16. The cellulase-free laundry detergent of claim 15, wherein said detergent comprises a cellulosic polymer.

17. A method of making a cell, comprising:

a) introducing a first recombinant nucleic acid into a *Bacillus sp.* cell so that said recombinant nucleic acid recombines with the *bglC* gene of said cell; and

b) introducing a second recombinant nucleic acid into said cell that provides for expression of said secreted detergent-additive protein.

18. The method of claim 17 wherein said nucleic acid inserts into said *bglC* gene.

19. The method of claim 17, wherein said nucleic acid deletes at least a portion of said *bglC* gene.

20. The method of claim 17, wherein said secreted detergent-additive protein is a subtilisin.




FIG. 1A

2 / 3






FIG. 1B

3 / 3

**FIG. 2****FIG. 3**