




(86) **Date de dépôt PCT/PCT Filing Date:** 2010/08/16
(87) **Date publication PCT/PCT Publication Date:** 2011/02/24
(45) **Date de délivrance/Issue Date:** 2015/02/24
(85) **Entrée phase nationale/National Entry:** 2012/02/16
(86) **N° demande PCT/PCT Application No.:** JP 2010/063804
(87) **N° publication PCT/PCT Publication No.:** 2011/021597
(30) **Priorité/Priority:** 2009/08/19 (JP2009-190145)

(51) **Cl.Int./Int.Cl.** **A61K 31/47** (2006.01),
A61K 47/32 (2006.01), **A61K 47/36** (2006.01),
A61K 47/38 (2006.01), **A61P 35/00** (2006.01),
A61P 35/04 (2006.01), **A61P 43/00** (2006.01),
C07D 215/48 (2006.01)
(72) **Inventeur/Inventor:**
BANDO, MASASHI, JP
(73) **Propriétaire/Owner:**
EISAI R&D MANAGEMENT CO. LTD., JP
(74) **Agent:** TORYS LLP

(54) **Titre : COMPOSITION PHARMACEUTIQUE CONTENANT UN DERIVE DE QUINOLEINE**
(54) **Title: QUINOLINE DERIVATIVE-CONTAINING PHARMACEUTICAL COMPOSITION**




(57) **Abrégé/Abstract:**

A. pharmaceutical composition comprising a compound represented by the formula (I) or pharmaceutically acceptable salt thereof or solvate thereof; and a basic substance is excellent in dissolution, is stable even after a long term storage, and is useful as a preventive or therapeutic agent against a tumor: (see formula I) wherein, R¹ is a hydrogen atom, a C₁₋₆ alkyl group or a C₃₋₈ cycloalkyl group; and R² is a hydrogen atom or a methoxy group.

ABSTRACT

A pharmaceutical composition comprising a compound represented by the formula (I) or pharmaceutically acceptable salt thereof or solvate thereof; and a basic substance is excellent in dissolution, is stable even after a long term storage, and is useful as a preventive or therapeutic agent against a tumor:

5



wherein, R^1 is a hydrogen atom, a C_{1-6} alkyl group or a C_{3-8} cycloalkyl group; and R^2 is a hydrogen atom or a methoxy group.

DESCRIPTION

Title of Invention


QUINOLINE DERIVATIVE-CONTAINING PHARMACEUTICAL COMPOSITION

Technical Field

5 [0001] The present invention relates to a pharmaceutical composition comprising a quinoline derivative, useful as a medicament. More specifically, the present invention relates to a pharmaceutical composition improved in dissolution of a quinoline derivative or a pharmaceutically acceptable salt thereof or a solvate thereof.

Background Art

10 [0002] A quinoline derivative represented by the formula (I) or a pharmaceutically acceptable salt thereof or a solvate thereof (hereinafter referred to as quinoline derivative (I)) has been known to have a potent angiogenesis inhibitory effect (Patent Literature 1) and a c-Kit kinase inhibitory effect (Patent Literature 2) and to be useful as a preventive or therapeutic agent against various tumors such as thyroid cancer, lung cancer, melanoma and
15 pancreatic cancer, and as an metastatic inhibitor against these tumors:



wherein, R^1 is a hydrogen atom, a C_{1-6} alkyl group or a C_{3-8} cycloalkyl group; and R^2 is a hydrogen atom or a methoxy group.

20 [0003] However, the quinoline derivative (I) has been found to degrade under humidifying and warming storage conditions when formulated into a pharmaceutical composition. In addition, when the pharmaceutical composition absorbs moisture, dissolution of the quinoline derivative (I) from the pharmaceutical composition that is an active ingredient may delay because of gelation on the surface of the composition. In order to overcome these problems, a pharmaceutical composition which includes the
25 quinoline derivative (I), (1) a compound, a 5% (w/w) aqueous solution or suspension of which has a pH of 8 or more, and/or (2) silicic acid, salt thereof or solvate thereof has been

developed (Patent Literature 3).

Citation List

Patent Literature

[0004] Patent Literature 1: WO 2002/32872

5 Patent Literature 2: WO 2004/080462

Patent Literature 3: WO 2006/030826

Summary of Invention

Technical Problem

10 [0005] However, development of a pharmaceutical composition further excellent in the dissolution of the quinoline derivative (I) has been desired. Thus, the present invention is aimed at providing a pharmaceutical composition that is excellent in dissolution of the quinoline derivative (I) that is maintained even after long term storage.


Solution to Problem

15 [0006] The present inventors have intensively studied in order to solve the problems above and surprisingly have discovered the configuration below could solve the problems and have completed the present invention.

Specifically, the present invention provides the following <1> to <12>.

[1] A pharmaceutical composition comprising:

20 (1) a compound represented by the formula (I) or pharmaceutically acceptable salt thereof or solvate thereof:



wherein R^1 is a hydrogen atom, a C_{1-6} alkyl group or a C_{3-8} cycloalkyl group; and R^2 represents a hydrogen atom or a methoxy group; and

(2) a basic substance.

25 [2] The composition according to [1], wherein the basic substance is a carbonate.

[3] The composition according to [2], wherein the salt is an alkaline earth metal salt.

[4] The composition according to [3], wherein the alkaline earth metal salt is a magnesium salt or a calcium salt.

[5] The composition according to any one of [1] to [4], further comprising a disintegrating agent.

5 [6] The composition according to [5], wherein the disintegrating agent is carmellose sodium, carmellose calcium, carboxymethyl starch sodium, croscarmellose sodium, low-substituted hydroxypropylcellulose or crospovidone.

[7] The composition according to any one of [1] to [6], wherein R^1 is a hydrogen atom, a methyl group, an ethyl group, an n-propyl group or a cyclopropyl group.

10 [8] The composition according to any one of [1] to [7], wherein R^1 is a cyclopropyl group.

[9] The composition according to any one of [1] to [8], wherein R^2 is a hydrogen atom, a methoxy group or an ethoxy group.

15 [10] The composition according to any one of [1] to [9], wherein R^2 is a hydrogen atom.

[11] The composition according to any one of [1] to [10], wherein the pharmaceutically acceptable salt is hydrochloride, hydrobromide, p-toluenesulfonate, sulfate, methanesulfonate or ethanesulfonate.

20 [12] The composition according to any one of [1] to [11], wherein the compound represented by the formula (I) is 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide methanesulfonate.

Advantageous Effects of Invention

25 [0007] The pharmaceutical composition of the present invention is excellent in dissolution of the quinoline derivative (I), which is a principal agent, and is also excellent in absorption into a living body. The pharmaceutical composition is also a pharmaceutical composition that is maintained even after long term storage.

Brief description of drawings

30 [0008] Fig. 1 shows the dissolution profiles of the compound A from the pharmaceutical compositions obtained in Examples 4 to 6 and Comparative Example 1.

Fig. 2 shows the dissolution profiles of the compound A from the pharmaceutical compositions obtained in Examples 7 to 9 and Comparative Example 2.

Fig. 3 shows the dissolution patterns of the compound A from the pharmaceutical

compositions obtained in Examples 10 to 12 and Comparative Example 3.

Fig. 4 shows the dissolution profiles of the compound A from the pharmaceutical compositions obtained in Examples 13 to 15 and Comparative Example 4.

5 Fig. 5 shows the dissolution profiles of the compound A from the pharmaceutical compositions obtained in Examples 16 to 17 and Comparative Example 5.

Fig. 6 shows the dissolution profiles of the compound A from the pharmaceutical compositions obtained in Example 18 and Comparative Examples 7 to 8.

Fig. 7 shows the dissolution profiles of the compound A from the pharmaceutical compositions obtained in Example 19 and Comparative Examples 9 to 10.

10 Description of Embodiments

[0009] The pharmaceutical composition of the present invention means a composition comprising the quinoline derivative (I) and a basic substance as essential ingredients. A mixing ratio of the quinoline derivative (I) and the basic substance is, but is not limited to, normally 1:0.5 to 50, preferably 1:1 to 25, further preferably 1:2 to 12.5.

15 [0010] In addition, a mixing rate of the quinoline derivative (I) with respect to the total weight of the pharmaceutical composition (excluding a capsule shell) is normally 0.25 to 50 weight %, preferably 0.5 to 25 weight %, further preferably 1 to 12.5 weight %.

[0011] A mixing rate of the basic substance with respect to the total weight of the pharmaceutical composition is normally 1 to 60 weight %, preferably 5 to 50 weight %, further preferably 10 to 40 weight %. At least one basic substance of the present invention may be included in the pharmaceutical composition, or two or more basic substances may also be included.

20 [0012] A dosage form of the pharmaceutical composition specifically means a solid preparation such as granules, fine granules, tablets or capsules and so on. It is preferably fine granules, granules or capsules filled with fine granules or granules.

[0013] The quinoline derivative (I) is a compound disclosed in WO 2002/32872. A preferable quinoline derivative (I) is a quinoline derivative or pharmacologically acceptable salt thereof or solvate thereof selected from the group consisting of

30 4-(3-fluoro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-(2-methoxyethoxy)-6-quinolinecarboxamide,

4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,

4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-(2-methoxyethoxy)-6-

- quinolinecarboxamide,
 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-(2-hydroxyethoxy)-6-quinolinecarboxamide,
 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-((2S)-2,3-
 5 dihydroxypropyl)oxy-6-quinolinecarboxamide,
 4-(3-chloro-4-(methylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 4-(3-chloro-4-(ethylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 N6-methoxy-4-(3-chloro-4-(((ethylamino)carbonyl)amino)phenoxy)-7-methoxy-6-quinolinecarboxamide,
 10 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-(2-ethoxyethoxy)-6-quinolinecarboxamide,
 4-(4-((cyclopropylamino)carbonyl)aminophenoxy)-7-(2-methoxyethoxy)-6-quinolinecarboxamide,
 N-(2-fluoro-4-[(6-carbamoyl-7-methoxy-4-quinolyl)oxy]phenyl)-N'-cyclopropylurea,
 15 N6-(2-hydroxyethyl)-4-(3-chloro-4-(((cyclopropylamino)carbonyl)amino)phenoxy)-7-methoxy-6-quinolinecarboxamide,
 4-(3-chloro-4-(1-propylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 4-(3-chloro-4-(cis-2-fluoro-cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 20 N6-methyl-4-(3-chloro-4-(((cyclopropylamino)carbonyl)amino)phenoxy)-7-(2-methoxyethoxy)-6-quinolinecarboxamide and
 N6-methyl-4-(3-chloro-4-(((ethylamino)carbonyl)amino)phenoxy)-7-methoxy-6-quinolinecarboxamide.
 25 [0014] A more preferable quinoline derivative (I) is a quinoline derivative or pharmacologically acceptable salt thereof or solvate thereof selected from the group consisting of
 4-(3-chloro-4-(methylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 4-(3-chloro-4-(ethylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 30 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide,
 N6-methoxy-4-(3-chloro-4-(((cyclopropylamino)carbonyl)amino)phenoxy)-7-methoxy-6-quinolinecarboxamide and

N6-methoxy-4-(3-chloro-4-(((ethylamino)carbonyl)amino)phenoxy)-7-methoxy-6-quinolinecarboxamide.

5 [0015] A particularly preferable quinoline derivative (I) is 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide or pharmacologically acceptable salt thereof or solvate thereof.

[0016] The pharmaceutically acceptable salt of the present invention means hydrochloride, hydrobromide, p-toluenesulfonate, sulfate, methanesulfonate or ethanesulfonate. It is preferably the methanesulfonate.

10 [0017] The solvate of the present invention means hydrate, dimethyl sulfoxide solvate or acetic acid solvate.

[0018] The quinoline derivative (I) is preferably a crystal of a salt of 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide, or a solvate thereof disclosed in WO 2005/063713. A particularly preferred quinoline derivative (I) is the C Form crystal of 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide
15 methanesulfonate.

[0019] The quinoline derivative (I) is useful as a preventive or therapeutic agent against various tumors and as a metastasis inhibitor against tumors. Examples of the tumors against which the quinoline derivative (I) is effective include thyroid cancer, non-small-cell
20 lung cancer, melanoma, laryngopharyngeal cancer, esophageal cancer, gastric cancer, colorectal cancer, hepatocellular carcinoma, renal cell carcinoma, pancreatic cancer, bladder cancer, breast cancer, uterine cancer, ovarian cancer, prostate cancer, testicular cancer, gastrointestinal stromal tumor, sarcoma, osteogenic sarcoma, angioma, malignant lymphoma, myeloid leukemia, neuroma and neuroglioma.

25 [0020] The basic substance of the present invention means a basic inorganic salt. Such basic inorganic salts include beryllium carbonate, magnesium carbonate, calcium carbonate, strontium carbonate, barium carbonate, potassium carbonate, calcium hydrogenphosphate and titanium oxide. It is preferably an alkaline earth metal salt of carbonic acid, further preferably magnesium carbonate or calcium carbonate.

30 [0021] It is also acceptable to further include a disintegrating agent in the pharmaceutical composition of the present invention. Such a disintegrating agent include corn starch, partially pregelatinized starch, hydroxypropyl starch, carmellose, carmellose sodium, carmellose calcium, carboxymethyl starch sodium, croscarmellose sodium, low-substituted

hydroxypropylcellulose and crospovidone. It is preferably the croscarmellose sodium, the low-substituted hydroxypropylcellulose or the crospovidone.

[0022] The pharmaceutical composition of the present invention may be prepared by a known method such as a method described in the General Rules for Preparations in the Japanese Pharmacopoeia Fifteenth Edition.

[0023] For example, in the case of the granule, it is possible to add an excipient, a binder, a disintegrating agent, a solvent, or the like to the quinoline derivative (I) as needed, to perform agitation granulation, extruding granulation, tumbling granulation, fluidized-bed granulation, spray granulation, or the like, and to prepare it. It is also acceptable to be coated with an atomizing agent containing the quinoline derivative (I) and an additive such as corn starch, microcrystalline cellulose, hydroxypropylcellulose, methylcellulose or polyvinylpyrrolidone while spraying water or a solution of a binder such as saccharose, hydroxypropylcellulose or hydroxypropylmethylcellulose on a core material such as a purified sucrose spherical granule, a lactose/crystalline cellulose spherical granule, a saccharose/starch spherical granule or a granular crystalline cellulose. It is also acceptable to perform sizing and milling as needed.

[0024] It is also possible to further, as needed, add an excipient, a binder, a disintegrating agent, a lubricant, an anti-oxidizing agent, a corrigent, a coloring agent, a flavoring agent, or the like to the granule prepared in this way and to compress it to be a tablet. A required excipient may be added to the quinoline derivative (I) to directly compress the mixture into a tablet. It is also possible to fill a capsule with the quinoline derivative (I) added/mixed with an excipient such as lactose, saccharose, glucose, starch, microcrystalline cellulose, powdered glycyrrhiza, mannitol, calcium phosphate or calcium sulfate, or with the granule.

[0025] Examples of the excipient include lactose, saccharose, glucose, fructose, starch, potato starch, corn starch, wheat starch, rice starch, crystalline cellulose, microcrystalline cellulose, powdered glycyrrhiza, mannitol, erythritol, maltitol, sorbitol, trehalose, silicic anhydride, calcium silicate, sodium hydrogencarbonate, calcium phosphate, anhydrous calcium phosphate and calcium sulfate.

[0026] Examples of the binder include gelatin, starch, gum arabic, tragacanth, carboxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, methylcellulose, partially pregelatinized starch, pregelatinized starch, polyvinyl alcohol, sodium arginine, pullulan and glycerin.

[0027] Examples of the disintegrating agent include corn starch, partially pregelatinized

starch, hydroxypropyl starch, carmellose, carmellose sodium, carmellose calcium, carboxymethyl starch sodium, croscarmellose sodium, low-substituted hydroxypropylcellulose and crospovidone.

[0028] Examples of the lubricant include magnesium stearate, stearic acid, calcium stearate, sodium stearyl fumarate, talc and macrogol.

[0029] Examples of the anti-oxidizing agent include sodium ascorbate, L-cysteine, sodium sulfite, tocopherol and soybean lecithin.

[0030] Examples of the corrigent include citric acid, ascorbic acid, tartaric acid, malic acid, aspartame, acesulfame potassium, thaumatin, saccharin sodium, dipotassium glycyrrhizinate, sodium glutamate, sodium 5'-inosinate and sodium 5'-guanylate.

[0031] Examples of the coloring agent include titanium oxide, iron sesquioxide, iron sesquioxide yellow, cochineal, carmine, riboflavin, food yellow No. 5 and food blue No. 2.

[0032] Examples of the flavoring agent include lemon oil, orange oil, menthol, peppermint oil, borneol and vanilla flavor.

Examples

[0033] The present invention will be described in more detail below with reference to Examples, but is not limited to the Examples.

[0034] Examples 1 to 3

Wet granulation was performed with purified water as a solvent using a high-shear granulator (apparatus name: FM-VG-10, manufactured by Powrex Corporation) with the C form crystal of 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide methanesulfonate (hereinafter referred to as compound A), D-mannitol (trade name: MannitolTM, Merck), precipitated calcium carbonate (trade name: Whiton F, Shiraishi CalciumTM), hydroxypropylcellulose (HPC-L, Nippon Soda), low-substituted hydroxypropylcellulose (trade name: L-HPC (LH-21), Shin-Etsu Chemical) and microcrystalline cellulose (trade name: Ceolus PH-101, Asahi Kasei Chemicals) according to the formulation proportions in Table 1. The granules of which a moisture content was reduced to be less than 2% by further drying were sized using a screen mill (apparatus name: Power Mill P-04STM, manufactured by Showa Giken KK) so that their granule diameters were less than 1 mm. Then, microcrystalline cellulose (trade name: Ceolus PH-102, Asahi Kasei Chemicals) and talc (trade name: Hi-Filler 17, Iwai Chemicals Company) were added to the sized granules according to the formulation proportions in Table 1, and the mixture was thoroughly mixed using a diffusion (tumbler-type) mixer

(trade name: 10L/20L Exchange-type Tumbler Mixer, manufactured by Toyo Packing Corporation). Hard capsules size #4 were filled with 100 mg of the resultant granules to prepare capsules containing the compound A.

[0035] [Table 1]

	Ex. 1	Ex. 2	Ex. 3
Compound A	1.25	5	12.5
Precipitated calcium carbonate	33	33	33
D-Mannitol	19.75	16	8.5
Hydroxypropylcellulose	3	3	3
Low-substituted hydroxypropylcellulose	25	25	25
Microcrystalline cellulose (PH-101)	10	10	10
Microcrystalline cellulose (PH-102)	5	5	5
Talc	3	3	3
Total	100	100	100

Unit: weight %

5

[0036] Examples 4 to 9, Comparative Examples 1 to 2

The compound A, precipitated calcium carbonate, low-substituted hydroxypropylcellulose, D-mannitol and talc were thoroughly mixed using a mortar and a pestle according to the formulation proportions in Table 2 and Table 3. Hard capsules size #3 were filled with 100 mg of the resultant mixtures to prepare capsules in Examples 4 to 9. Capsules in Comparative Examples 1 to 2, which contained no precipitated calcium carbonate, were also prepared by the same method.

10

[0037] [Table 2]

	Com.Ex. 1	Ex. 4	Ex. 5	Ex. 6
Compound A	5	5	5	5
Precipitated calcium carbonate	0	5	10	20
Low-substituted hydroxypropylcellulose	30	25	20	10
D-Mannitol	62	62	62	62
Talc	3	3	3	3
Total	100	100	100	100

Unit: weight %

15

[0038] [Table 3]

	Com.Ex. 2	Ex. 7	Ex. 8	Ex. 9
Compound A	20	20	20	20
Precipitated calcium carbonate	0	5	10	20
Low-substituted hydroxypropylcellulose	30	25	20	10
D-Mannitol	47	47	47	47
Talc	3	3	3	3
Total	100	100	100	100

Unit: weight %

[0039] Test Example 1

The dissolutions of the compound A in the capsules in Examples 4 to 9 and Comparative Examples 1 to 2 were examined according to the Dissolution Test (the Paddle method, test medium: JP1 solution) described in the Japanese Pharmacopoeia Fifteenth Edition. As a result, the dissolutions of the compound A in the capsules in Comparative Examples 1 to 2, in which no calcium carbonate was mixed, were insufficient. In contrast, the dissolutions of the compound A in the capsules in Examples 4 to 9, in which calcium carbonate was mixed, were good (Fig. 1 and Fig. 2).

10 [0040] Examples 10 to 15, Comparative Examples 3 to 4

The compound A, magnesium carbonate (Kyowa Chemical Industry), low-substituted hydroxypropylcellulose, D-mannitol and talc were thoroughly mixed using a mortar and a pestle according to the formulation proportions in Table 4 and Table 5. Hard capsules size #3 were filled with 100 mg of the resultant mixtures to prepare capsules in Examples 10 to 15. Capsules in Comparative Examples 3 to 4, which contained no magnesium carbonate, were also prepared by the same method.

15

[0041] [Table 4]

	Com.Ex. 3	Ex. 10	Ex. 11	Ex. 12
Compound A	5	5	5	5
Magnesium carbonate	0	5	10	20
Low-substituted hydroxypropylcellulose	30	25	20	10
D-Mannitol	62	62	62	62
Talc	3	3	3	3
Total	100	100	100	100

Unit: weight %

[0042] [Table 5]

	Com.Ex. 4	Ex. 13	Ex. 14	Ex. 15
Compound A	20	20	20	20
Magnesium carbonate	0	5	10	20
Low-substituted hydroxypropylcellulose	30	25	20	10
D-Mannitol	47	47	47	47
Talc	3	3	3	3
Total	100	100	100	100

Unit: weight %

[0043] Test Example 2

The dissolutions of the compound A in the capsules in Examples 10 to 15 and Comparative Examples 3 to 4 were examined by the same method as in Test Example 1. The dissolutions of the compound A in the capsules in Comparative Examples 3 to 4, in which no magnesium carbonate was mixed, were insufficient. In contrast, the dissolutions of the compound A in the capsules in Examples 10 to 15, in which the magnesium carbonate was mixed, were good (Fig. 3 and Fig. 4).

[0044] Examples 16 to 17, Comparative Examples 5 to 6

Purified water was added to the compound A, precipitated calcium carbonate or magnesium carbonate, hydroxypropylcellulose and croscarmellose sodium (trade name: Ac-Di-SolTM, Asahi Kasei Chemicals) to perform granulation using a mortar and a pestle, followed by sizing of the dried granules so that their granule diameters were less than 1 mm. Then, microcrystalline cellulose (trade name: Ceolus PH-102TM, Asahi Kasei Chemicals), low-substituted hydroxypropylcellulose and talc (trade name: Hi-Filler 17, Iwai Chemicals Company) were added to the sized granules according to the formulation proportions in Table 6, and the mixture was mixed thoroughly. Hard capsules size #4 were filled with 100 mg of the resultant mixtures to prepare capsules in Examples 16 to 17. Capsules in Comparative Examples 5 to 6, which contained neither precipitated calcium carbonate nor magnesium carbonate but contained mannitol or talc as a substitute, were also similarly prepared according to the formulation proportions in Table 7.

[0045] [Table 6]

	Ex. 16	Ex. 17
Compound A	10	10
Precipitated calcium carbonate	15	0
Magnesium carbonate	0	15
Hydroxypropylcellulose	2	2
Croscarmellose sodium	10	10
Low-substituted hydroxypropylcellulose	20	20
Microcrystalline cellulose (PH-102)	41	41
Talc	2	2
Total	100	100

Unit: weight %

[0046] [Table 7]

	Com.Ex. 5	Com.Ex. 6
Compound A	10	10
Mannitol	15	0
Talc	0	15
Hydroxypropylcellulose	2	2
Croscarmellose sodium	10	10
Low-substituted hydroxypropylcellulose	20	20
Microcrystalline cellulose (PH-102)	41	41
Talc	2	2
Total	100	100

Unit: weight %

[0047] Test Example 3

5 The dissolutions of the compound A in the capsules in Examples 16 to 17 and Comparative Example 5 were examined by the same method as in Test Example 1. The dissolution of the compound A in the capsule in Comparative Example 5, in which neither calcium carbonate nor magnesium carbonate was mixed, was insufficient. In contrast, the dissolutions of the compound A in the capsules in Examples 16 to 17, in which calcium carbonate or magnesium carbonate was mixed, were good (Fig. 5).

[0048] Test Example 4

10 The capsules in Examples 16 to 17 and Comparative Example 6 were stored for 1 week in an open system under an environment at a temperature of 60°C and a relative humidity of 75%, followed by determining the production of the degradants with high-performance liquid chromatography. In the capsule formulation in Comparative Example

15

6, in which neither calcium carbonate nor magnesium carbonate was mixed, an amount of the degradants was increased. In contrast, in the capsules in Examples 16 to 17, in which calcium carbonate or magnesium carbonate was mixed, no increase in amount of the degradants was observed (Table 8).

5 [0049] [Table 8]

	Degradants (%)	Quantitated compound A(%)
Compound A (Initial)	1.61%	98.38%
Com.Ex. 6	1.92%	98.08%
Ex. 16	1.50%	98.50%
Ex. 17	1.57%	98.44%

[0050] Examples 18 to 19, Comparative Examples 7 to 10

The respective ingredients were mixed according to the formulations of Tables 9 and 10 by the same method as in Examples 4 to 9 and Comparative Examples 1 to 2. Hard capsules size #3 were filled with 100 mg of the resultant mixtures to prepare capsules in Examples 18 to 19 and Comparative Examples 7 to 10.

10 [0051] [Table 9]

	Ex. 18	Com.Ex. 7	Com.Ex. 8
Compound A	20	20	20
Precipitated calcium carbonate	10	0	0
Calcium oxide	0	10	0
Calcium hydroxide	0	0	10
Low-substituted hydroxypropylcellulose	20	20	20
D-Mannitol	47	47	47
Talc	3	3	3
Total	100	100	100

Unit: weight %

[0052] [Table 10]

	Ex. 19	Com.Ex. 9	Com. Ex. 10
Compound A	20	20	20
Magnesium carbonate	10	0	0
Magnesium oxide	0	10	0
Magnesium hydroxide	0	0	10
Low-substituted hydroxypropylcellulose	20	20	20
D-Mannitol	47	47	47
Talc	3	3	3
Total	100	100	100

Unit: weight %

[0053] Test Example 5


The dissolutions of the compound A in the capsules in Examples 18 to 19 and Comparative Examples 7 to 10 were examined by the same method as in Test Example 1. As a result, the dissolutions of the compound A in the capsules in Comparative Examples 7 to 10, in which calcium oxide, calcium hydroxide, magnesium oxide or magnesium hydroxide was mixed, were insufficient. In contrast, the dissolutions of the compound A in the capsules in Examples 18 to 19, in which calcium carbonate or magnesium carbonate was mixed, were good (Fig. 6 and Fig. 7).

10 Industrial Applicability

[0054] The pharmaceutical composition of the present invention is excellent in dissolution of the quinoline derivative and also in stability, and is therefore useful as a medicament for prevention or treatment of a tumor.

CLAIMS

1. A pharmaceutical composition comprising
 - (1) a compound represented by the formula (I) or pharmaceutically acceptable salt thereof or solvate thereof:



wherein R^1 is a hydrogen atom, a C_{1-6} alkyl group or a C_{3-8} cycloalkyl group; and R^2 represents a hydrogen atom or a methoxy group; and

- (2) an alkaline earth metal carbonate.
2. The composition according to claim 1, wherein the alkaline earth metal carbonate is magnesium carbonate or calcium carbonate.
3. The composition according to claim 1 or 2, further comprising a disintegrating agent.
4. The composition according to claim 3, wherein the disintegrating agent is carmellose sodium, carmellose calcium, carboxymethyl starch sodium, croscarmellose sodium, low-substituted hydroxypropylcellulose or crospovidone.
5. The composition according to any one of claims 1 to 4, wherein R^1 is a hydrogen atom, a methyl group, an ethyl group, an n-propyl group or a cyclopropyl group.
6. The composition according to any one of claims 1 to 5, wherein R^1 is a cyclopropyl group.
7. The composition according to any one of claims 1 to 6, wherein R^2 is a hydrogen atom.
8. The composition according to any one of claims 1 to 7, wherein the pharmaceutically acceptable salt of the compound represented by the formula (I) is hydrochloride, hydrobromide, p-toluenesulfonate, sulfate, methanesulfonate or ethanesulfonate.
9. The composition according to any one of claims 1 to 8, wherein the compound represented by the formula (I) is 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide methanesulfonate.

10. A pharmaceutical composition comprising

(1) 4-(3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy)-7-methoxy-6-quinolinecarboxamide methanesulfonate,

(2) calcium carbonate,

(3) mannitol,

(4) hydroxypropylcellulose,


(5) low-substituted hydroxypropylcellulose,

(6) microcrystalline cellulose and

(7) talc.


FP10-0287-00

1/7

Fig.1


FP10-0287-00

2/7

Fig.2


FP10-0287-00

3/7

Fig.3

FP10-0287-00


4/7

Fig.4


FP10-0287-00

5/7

Fig.5



6/7

Fig.6

FP10-0287-00

7/7

Fig.7