




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : C07H 15/12, 17/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 86/ 02929 (43) International Publication Date: 22 May 1986 (22.05.86)</p>
<p>(21) International Application Number: PCT/US85/02198 (22) International Filing Date: 6 November 1985 (06.11.85) (31) Priority Application Number: 669,334 (32) Priority Date: 8 November 1984 (08.11.84) (33) Priority Country: US</p> <p>(71) Applicant: LIFE TECHNOLOGIES, INC. [US/US]; 8717 Grovemont Circle, Gaithersburg, MD 20760 (US). (72) Inventors: KLEVAN, Leonard ; 3650 39th Street, N.W., Washington, DC 20016 (US). GEBEYEHU, Gulilat ; 9130 Glen Avenue, #102, Silver Spring, MD 20910 (US). RAO, Prasad, Y. ; 9908 Durango Drive, Damas- cus, MD 20872 (US).</p>		<p>(74) Agent: HOCHBERG, D., Peter.; 1510 Ohio Savings Plaza, Cleveland, OH 44114 (US). (81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), LU (European patent), NL (European patent), SE (European patent).</p> <p>Published <i>With international search report.</i></p>


(54) Title: NUCLEOTIDE ANALOGS FOR NUCLEIC ACID LABELING AND DETECTION

(57) Abstract

Nucleotide analogs, modified by the attachment at hydrogen bonding positions of linker groups, that is, the 6-position of adenine, 4-position of cytosine, and 2-position of guanine, are prepared. Such analogs, alone or with reporter groups attached (Figure 1), may be incorporated into DNA probes which effectively hybridize adenine to target DNA. For example, Figure 1 shows adenine and cytosine analogs attached to a biotin reporter group.



Bio-3-dATP	L = (CH ₂) ₃
Bio-7-dATP	L = (CH ₂) ₇
Bio-10-dATP	L = (CH ₂) ₁₀ NHCO(CH ₂) ₃
Bio-14-dATP	L = (CH ₂) ₁₄ NHCO(CH ₂) ₃
Bio-10'-dATP	L = CH ₂ CONH(CH ₂) ₁₀
Bio-17-dATP	L = CH ₂ CONH(CH ₂) ₁₇ NHCO(CH ₂) ₃



Bio-3-dCTP	L = (CH ₂) ₃
Bio-7-dCTP	L = (CH ₂) ₇
Bio-10-dCTP	L = (CH ₂) ₁₀ NHCO(CH ₂) ₃
Bio-14-dCTP	L = (CH ₂) ₁₄ NHCO(CH ₂) ₃

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GA	Gabon	MR	Mauritania
AU	Australia	GB	United Kingdom	MW	Malawi
BB	Barbados	HU	Hungary	NL	Netherlands
BE	Belgium	IT	Italy	NO	Norway
BG	Bulgaria	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali		
FR	France				

NUCLEOTIDE ANALOGS FOR NUCLEIC ACID
LABELING AND DETECTION

This invention relates to techniques of detecting the hybridization of nucleic acid probes to target nucleic acids. In particular, it relates to a method of biotinylating such probes so that
5 biotin-avidin or biotin-streptavidin binding may be used to localize and detect the target.

BACKGROUND OF THE INVENTION

Nucleic acid hybridization has found widespread application in genetic research, biomedical
10 research and clinical diagnostics. The development of a technology for "mixed phase hybridization" (hybridizing probes to immobilized target DNA) and for cloning of unique sequence genetic probes has lead to major
15 advances in basic and applied areas of biochemistry and medicine (For a review see: Meinhoth, J. and Wahl, G., (1984) Analytical Biochemistry 138, 267-284). In the standard hybridization reaction, a radioisotope-labeled
20 probe is annealed to a DNA or RNA sample which has been immobilized on an inert solid support. The detection of a radioactive signal by autoradiography indicates the presence or absence of the complementary nucleic acid
25 sequence in the target sample. In the past, the potential health hazards, disposal problems and instability of radionucleotides have placed limitations on the use of nucleic acid probes. This has lead to
30 great interest in the development of alternative DNA labeling and detection systems which do not contain the drawbacks inherent in the use of radioisotopes.

The development of a nonradioactive DNA
35 detection system requires a method to label the nucleic acid probe and a means to recognize and detect the annealed probe-target hybrid. The high binding constant and specificity of biotin for avidin or streptavidin (reviewed in Green, M., (1975) Advances in Protein
Chemistry, 29, 85-133) have been used in different

systems to localize and detect a wide range of target molecules.

The principle of the biotin-avidin interaction as the basis of probe-ligand interactions in molecular biology was reviewed by Bayer and Wilchek in 1979: "Thus (in addition to biotin-requiring enzymes) biotin-derivatized hormones, phages, lectins, antibodies and other binding proteins can interact with avidin; and if the avidin is immobilized or covalently bound to a potentially perceptible probe, the avidin-biotin complex can be used for the localization or isolation of the compounds above and/or their receptors" (Bayer, E.A. and Wilchek, M. (1979), *Methods in Biochemical Analysis* 26, 1-45). This principle is further described in U.S. Patent 4,228,237 (Hevey, R.C. and Malmros, M.K., October 14, 1980) in which an avidin-coupled signaling enzyme is used to recognize a biotin-labeled reagent which will specifically bind to the ligand in question.

Other examples of biotin-avidin interactions as applied to probes are contained within an article by Langer et al. (Langer, P.R., Waldrop, A.A. and Ward, D.C. (1981) *Proc. Natl. Acad. Sci., USA* 78 6633-6637), who describe the synthesis of biotin-UTP and biotin-dUTP analogs which are substrates for a variety of RNA and DNA polymerases. The disclosures in this publication were further set forth in European Patent Application EP-A200063879, in which it is claimed that biotin may be covalently linked to the 8-position of a purine base or the 5-position of a pyrimidine base in a nucleoside triphosphate.

However, it is stated by Ward, et al., in that application that "...probe moieties should not be placed on ring positions that sterically, or otherwise, interfere with the normal Watson-Crick hydrogen bonding potential of the bases. Otherwise, the substituents will yield compounds that are inactive as polymerase substrates....Normally, such considerations limit substitution positions to the 5-position of a pyrimidine

and the 7-position of a purine or a 7-deazapurine."

(The Ward disclosure and claims show that the 8-position of a purine was intended.) This statement reflects the common supposition that if labels are attached to nucleotides at the hydrogen bonding positions the modified nucleotides cannot be useful in probes.

Notwithstanding the teachings of Ward and others, the inventors herein have discovered that nucleotide analogs in which biotin is attached at the hydrogen bonding position of the purine or pyrimidine base may be incorporated into probes. Moreover, probes labelled in this manner will still hybridize to "target" DNA in highly sensitive detection protocols. This result has occasioned the production of a new class of labeled nucleotides, the usefulness of which was hitherto unsuspected.

SUMMARY OF THE INVENTION

According to the present invention, dATP and dCTP are modified at the amino-nitrogen at the 6- and 4-positions, respectively, through a linker arm that varies in length from 3 to 17 atoms. dGTP may be similarly modified at the 2-position. Nucleotides labelled thereby with biotin are stable for at least one year, are inexpensive to prepare and are substrates for *E. coli* DNA polymerase I. Nucleic acid probes incorporating these analogs may be prepared by standard nick-translation protocols and employed for non-radioactive DNA detection when used with streptavidin-conjugated enzymes or enzyme polymers.

The same methods of synthesis may be used to prepare unlabelled nucleotide precursors which contain a linking group at the positions mentioned. The label or reporter group is added after incorporation. "Chain terminator" nucleotide analogs may also be synthesized.

Accordingly, it is an object of the present invention to provide improved methods for detecting the presence of specific polynucleotides or nucleic acids.

Another object of the present invention is to provide a novel means of labelling polynucleotides for detection of specific sequences, for use as probes, and the like.

5 It is another object of the present invention to provide a novel class of nucleotide analogs which may be incorporated into nucleic acids without interfering with hybridization to complementary nucleic acids.

10 A still further object of the invention is to provide nucleotide analogs which may be linked uniquely to biotin before or after incorporation into nucleic acids.

Other objects will be seen from the description of preferred embodiments and claims, as explained by the entire specification and the Figures, 15 in which

Figure 1 is a schematic molecular representation of some typical compounds synthesized according to the present invention;

20 Figure 2 is a schematic representation of syntheses of some of the dATP analogs of the present invention;

Figure 3 is a schematic representation of other synthetic methods for dATP analogs of the present 25 invention;

Figure 4 is a schematic representation of a synthetic method for the dCTP analogs of the present invention;

30 Figure 5 is a graph showing the incorporation into DNA of some dATP analogs of the present invention, as a function of time; and

Figure 6 is a graph showing nucleotide incorporation as measured by various radioactive labeling means.

GENERAL METHODS OF SYNTHESIS AND USE

We have used two approaches for the synthesis of dATP analogs. In the first, a modifiable amino group is introduced via reaction of chloropurine with diamino alkane. In the second approach, the amino group at the 6 position is alkylated by a modifiable group. In the case of dCTP, a modifiable amino group is introduced by transamination. dGTP analogs may also be prepared by the reaction of diaminoalkane with 2-chloro-2'-deoxyinosine following a similar procedure to that described for dATP. In all cases the amino group involved in Watson-Crick hydrogen bonding is preferably modified.

For convenience, the following abbreviations will be used:

B-NHS: N-hydroxysuccinimide biotin ester
CAB-NHS: N-hydroxysuccinimide
caproylamidobiotin ester
CDI: carbonyldiimidazole
EDC: ethyl dimethylaminopropyl carbodiimide
DAE: 1,2-diaminoethane
DAH: 1,6-diaminohexane
TEAB: triethylammonium bicarbonate

Synthesis of biotinylated dATP (Figures 2 and 3) and dGTP

In a first approach, the synthesis starts from the known compound 6-chloropurine-2'-deoxyriboside. This compound was prepared (Figure 2, (1)) in about 70% yield from 2'-deoxyinosine by using Robins' procedure. (M.J. Robins & G.L. Basom in "Nucleic Acid Chemistry" p. 602 (1978) edited by Townsend & Tipson.) It was phosphorylated (Figure 2, (2)) using $\text{POCl}_3/(\text{EtO})_3\text{PO}$ (M. Yoshikawa, T. Kato & T. Takenishi, Tetrahedron Lett. 5095 (1967)) in the presence of 4Å molecular sieves. The presence of molecular sieves is preferred since in their absence the reaction gives a mixture of products, the desired compound being only a minor component.

-6-

The resulting monophosphate was then treated (Figure 2, (3)) with diaminoalkane to give the desired N^6 -(n-aminoalkyl)dAMP. In our examples, DAE (n=2) and DAH (n=6) were employed, but n can range from 2 to 12, and probably beyond. N^6 -(6-aminohexyl)dAMP was obtained 60-70% yield, whereas the yield of N^6 -(2-aminoethyl)dAMP was very low. (The reaction condition was not maximized to increase the yield.)

The biotinylation (Ward et al., Proc. Natl. Acad. Sci. USA 78 6633 (1981)) was accomplished by treating the N^6 -(aminoalkyl)dAMP with B-NHS (Figure 2, (4a)), or CAB-NHS (Figure 2, (4b)). (S.M. Costello, R.T. Felix & R. W. Giese, Clin. Chem. 25 1572 (1979). Bio-7-dAMP and Bio-14-dAMP were obtained in 70-90% yield. Bio-3-dAMP and Bio-10-dAMP were also obtained in high yield, about 50-80%. In place of CAB-NHS one may employ the biotin ester of any acid of the form $H_2N(CH_2)_mCOOH$ for m such that the total linker length does not exceed 26 atoms.

Finally the triphosphates were prepared (Figure 2, (5a) and (5b)) by using the Hoard & Otts (D.E. Hoard & D.G. Otts, J.Am. Chem. Soc. 87, 1785 (1965)) method. That is, the monophosphates were treated with CDI followed with tributylammonium pyrophosphate to give Bio-7-dATP, Bio-3-dATP, Bio-14-dATP and Bio-10-dATP. The yield varied between 30 and 80%.

The Bio-7-dATP was also prepared (Figure 2, reactions (6)-(9)) from N^6 -(6-aminohexyl)dATP. This approach is particularly useful in the synthesis of radiolabelled biotinylated nucleotides or for attachment of other detectable groups. Where it is desired to fully phosphorylate the precursor before labelling, it is preferred to protect the amino group during the phosphorylation step, as Trayer et al. (Trayer et al., Biochem J. 139, 609 (1974)) have reported that the application of the Hoard-Ott method on the ribose analog N^6 -(6-aminohexyl)-AMP resulted in a complicated mixture and a low yield of the desired compound. Accordingly,

-7-

the amino group in N^6 -(6-aminohexyl)dAMP was protected with a trifluoro acetyl group by treatment (Figure 2,(6)) with ethyl trifluorothioacetate according to the procedure of Trayer, et al., to give N^6 -(6-Trifluoroacetamidohexyl)dAMP, which was then converted (Figure 2,(7)) to the triphosphate by the Hoard & Ott method. Basic hydrolysis (pH 11) of the protecting group (Figure 2,(8)) yielded N^6 -(6-Aminohexyl)dATP. This was then treated with B-NHS to yield Bio-7-dATP, (Figure 2, 9(a)), or, analogously, treatment with CAB-NHS will yield Bio-14-dATP (Figure 2, 9(b)). Tritium labelled Bio-7-dATP was synthesized analogously by using tritium labelled biotin-NHS ester. (Commercial product from Amersham).

In the second approach, (see Figure 3) dATP was alkylated (10) at the N-1 position with iodoacetic acid at pH 6.5 and subsequently rearranged (11) at pH 8.5, 90°C to the N-6 position to yield N^6 -carboxymethyl dATP. (M. Lindeberg and K. Mosback Eur. J. Biochem 53, 481 (1975)). That compound was then condensed (12) with diaminohexane using EDC, a water soluble coupling agent. Other diaminoalkanes with up to 12 carbon atoms may be employed. The aminohexyl adduct was then coupled with B-NHS (13a) or CAB-NHS (13b) to give the corresponding Bio-10'-dATP or Bio-17-dATP.

These compounds were also made by effecting the alkylation and the rearrangement on dAMP. The biotinylated compounds were then converted to Bio-10'-dATP and Bio-17-dATP respectively by using the Hoard-Ott method.

Generally the yield on the second approach is lower due to depurination on the first step.

The synthesis of dGTP analogs would start with 2-chloro-2'-deoxyinosine. Phosphorylation of this compound by the modified Yoshikawa method followed by treatment with diaminoalkane will give the corresponding N^2 (n-aminoalkyl)dGMP. This will then be subjected to the same treatment as in the case of

N^6 -(n-aminoalkyl)dAMP to obtain the corresponding biotinylated dGTP or the N^2 -(n-aminoalkyl)dGTP precursor.

Synthesis of biotinylated dCTP

5 The synthesis of biotinylated dCTP (figure 4) follows the procedure described by Draper (D.E. Draper, Nucleic Acid Res. 12 989 (1984)) where bisulfite-catalyzed transamination is effected at the N-4 position. Treatment of dCTP with DAE or DAH in the presence of bisulfite at a pH of about 5.5, followed by
10 adjustment of the pH to about 8.5, (Figure 4, (14)) gave N^4 -(2-aminoethyl)dCTP or N^4 -(6-aminohexyl)dCTP respectively. The yield in either case was less than 50%. The nucleotides are then treated (15(a),(b)) with B-NHS or CAB-NHS to give the desired compounds.

15 Use of these and analogous modified nucleotides

As will be seen from the experimental examples to follow, the modified nucleotides of the present invention may be incorporated into nucleic acids. Their presence is detectable with a high degree of
20 sensitivity, and labelled probes may be used to detect a single-copy gene sequence using existing methods.

Using reactions 1-3 plus 6-8, reaction 14, or reactions 10-12, nucleotide-analog precursors of these compounds, (such as N^6 -(6-aminohexyl)dATP) may also be
25 synthesized. They may be directly incorporated into nucleic acid probes and then linked to reporter molecules, such as biotin, or activated enzymes (e.g., enzymes activated with amino reactive, bifunctional crosslinking reagents such as
30 Bis(sulfosuccinimidyl)suberate or dimethyl suberimidate). Recently, it has been demonstrated that protein-linked single-stranded probes may be used in standard hybridization assays (Renz, M. and Kurz, C. (1984) Nucleic Acids Res. 12 3435-3444). It is also
35 clear to one skilled in the art that other reporter groups such as tetramethylrhodamine isothiocyanate, fluorescein isothiocyanate or dimethylaminoazobenzene

sulfonyl chloride may be condensed with the primary amino group of the nucleotide-analog precursors after incorporation into a nucleic acid (Richardson, R.W. and Gumpert, R.I., (1983) *Nucleic Acids Res.* 11 6167-6184).

5 The same synthetic procedures reported above may also be used to synthesize the corresponding chain terminator nucleotides such as 3'-deoxynucleoside-5'-triphosphate analogs or 2'-3'-dideoxynucleoside-5'-triphosphate analogs. The 3'-deoxy analogs may be
10 condensed, before or after the reporter group is added, with the 3'ends of a nucleic acid by chemical means or through the use of terminal deoxynucleotidyl transferase (TdT) (Tu, C.-P.D. and Cohen, S.P. (1980) *Gene* 10
177-183). This addition of a single biotin nucleotide
15 to the 3'end of each strand of DNA will permit the labelling of nucleic acids for detection of specific gene sequences in biomedical and research applications.

 The particular "linkers" we have described are presently preferred, as is their bonding to the label,
20 or reporter group, by an amido group. However, the linker can be any suitable compound which does not interfere either chemically or sterically with the desired application; further examples will be evident to one skilled in the art. For practical purposes, its
25 length (that is, the number of atoms between the amino group of the base and the reporter group) should not be more than about 26. A range of 3 to 17 is preferred. Our invention also contemplates the use of other bonding groups such as esters and thioesters. For example, the
30 chloronucleotides can be treated with aminoalkanol (e.g. 6-aminohexanol) to give N^6 -(n-hydroxyalkyl)dATP which can be esterified with biotin or longer chain analogs using standard coupling means such as dicyclohexyl carbodiimide. For the thioester, the chloronucleotide
35 can be treated with aminoalkanethiol to yield N^6 -(n-thioalkyl)dATP, which is esterified the same way.

EXAMPLES SHOWING PREFERRED EMBODIMENTSIncorporation into Plasmid DNA of Biotinylated dATP Derivatives

5 The time course and levels of incorporation were studied as a function of linker size.

Using the BRL Nick Translation System, dATP and various derivatives were incorporated into plasmid DNA. The 400 ul reaction mixture contained 50 mM Tris-HCl (pH 7.8), 5 mM MgCl₂, 10 mM 2-mercaptoethanol, 10 ug/ml BSA, 20 uM each of dGTP, dCTP, dTTP and dATP or a dATP derivative, 4 ug of a 5.4 Kb plasmid, 10uCi of ³H-dGTP (12 Ci/mmmole), 8 units of DNA polymerase I and 0.8 ng of DNase I. Reaction was carried out at 15°C.

15 At each time point of interest, 2 ul from the reaction mixture was withdrawn, spotted on glass fiber filters (GF/C), washed in 10% trichloroacetic acid (TCA) once and 5% TCA twice, and dried after washing in alcohol. The filters were counted in a liquid scintillation counter.

20 Figure 5 shows the time course of incorporation, given as labelled nucleotide incorporated per kilobase of DNA, when either dATP, a derivative precursor N⁶-(6-aminohexyl)dATP, biotinylated dATP with various linker lengths, or no dATP, was present in the reaction mixture.

25 The relative levels of incorporation of the modified nucleotides at the 90 minute time point are seen in Table 1 below, in which the data are reported as percent incorporation of the tracer radio-nucleotide (³H-dGTP) relative to incorporation resulting from a reaction mixture containing dNTP's and no derivatives. Duplicate figures indicate two determinations. Results for dCTP and analogs appear also.

TABLE 1

	<u>Nucleotide</u>	<u>% Incorporation (90 min.)</u>
	dATP	100
	N ⁶ -aminohexyldATP	67, 75
5	Biotin-7-dATP	53, 59
	Biotin-14-dATP	33, 39
	Biotin-3-dATP	29, 23
	Biotin-10-dATP	15, 17
	Biotin-10'-dATP	6.9, 7.5
10	Biotin-17-dATP	6.7, 5
	no ATP	6.5, 6
	dCTP	100
	N ⁴ -(6-aminohexyl)dCTP	74
	Biotin-7-dCTP	44
15	Biotin-14-dCTP	42
	Biotin-10-dCTP	22
	Biotin-3-dCTP	12

Using the same protocol, incorporation was then performed using a radiolabelled biotin-nucleotide. Thus, the level of biotin incorporation was directly determined. 200 ul of the reaction mixture contained 10uCi of α -³²PdCTP and either 20 uM dATP or 20 uM [³H-biotin]-7-dATP. The upper curve of Figure 6 shows the time course of incorporation of ³²P-dCMP when dATP was used in the reaction mixture. The lower curves show the time courses of incorporation of ³²P-dCMP and the tritium labelled and biotinylated derivative of dAMP. The close correspondence of these two curves indicates that the level of biotin incorporation is accurately reflected by ³²P-dNMP incorporation.

Studies on Sensitivity of Detection

To examine the detection sensitivity of probes made from the biotinylated nucleotides of the present invention, incorporation of various levels of Bio-7-dATP into plasmid DNA was carried out by nick translation as previously disclosed.

-12-

The reaction volume was 1 ml. At 10 minutes, 30 minutes and 90 minutes, 200 ul of the reaction mixture was removed and the reaction quenched with 30 mM EDTA. The DNA from each sample was purified on a Sephadex[®] G-50 column. Various picogram quantities of each sample, as seen in Table 2 below, were spotted on a nitrocellulose filter and checked for visualizability after use of BRL's DNA Detection System (a streptavidin-biotin alkaline phosphatase system) and dye development for 2 hours. In Table 2, "+" or "-" indicates the presence or absence, respectively of visible spots; it can be seen that a high degree of sensitivity exists even with a relatively low level of biotin incorporation.

TABLE 2

Time, min.	90	30	10
Bio-7-dAMP per Kb	98	35	7
50 pg of sample	+	+	+
20 pg of sample	+	+	+
10 pg of sample	+	+	+
5 pg of sample	+	+	+
2 pg of sample	+	+	+
1 pg of sample	+	+	-
0 pg of sample	-	-	-

Probes biotinylated with various linker lengths were then compared for sensitivity of detection. Plasmid DNA was nick translated according to the described protocol with Biotin-n-dATP, for n=7, 14, 3 or 10. The biotinylated DNA's were then purified from unincorporated nucleotides by gel filtration on Sephadex[®] G-50 in 1 x SSC (0.15 M NaCl, 0.015 M sodium citrate) containing 0.1% SDS. The biotinylated DNA was diluted into 6 x SSC containing 0.2 ug/ul sheared herring sperm DNA and spotted onto nitrocellulose paper in the range of 50 pg/5 ul to 1 pg/5 ul. After 30

minutes' incubation in the NBT/BCIP dye mixture of the BRL DNA Detection System, 2 pg spots of Bio-7, 14 or 10-dATP labeled probe and a 5 pg spot of Bio-3-dATP labeled probe were visible. After 1.5 hours, the 1 pg spot of each biotin-dATP-labeled probe was visible.

Likewise, probe labeled with Bio-10-dCTP exhibited 2 pg sensitivity after thirty minutes of detection, and the 1 pg spot was visible after 1.5 hours. Probe labeled with both Bio-7-dATP and Bio-10-dCTP was not detected with a higher sensitivity than probe labeled with either nucleotide individually.

Detection of Single-Copy Gene Sequences With the Nucleotides of the Present Invention

A plasmid containing the 1.1 Kb Mst II fragment of the human β -globin gene in the Eco RI site of pBR322 (obtained from Dr. Y.W. Kan, UCSF) was nick translated with either Biotin-n-dATP in the presence of dTTP, dCTP, and dGTP, or Biotin-n-dCTP in the presence of dATP, dTTP, and dGTP. All nucleotides were at 20uM and the reaction was allowed to proceed for 90 minutes. Conditions for nick translation of the DNA probe and purification of the biotin-labeled material are given in the manual of the BRL DNA Detection System.

The biotinylated probe was hybridized to a Southern blot of Eco RI digested human DNA or to an Eco RI digest of the plasmid DNA. Probe concentration was 100 ng/ml and hybridization was allowed to proceed for 16-24 hours under conditions described by Leary et al. (Leary, J.J., Brigati, D.J. and Ward, D.C. (1983) Proc. Natl. Acad. Sci. USA 80 4045-4049). The filter was developed with the BRL Detection System as described in the system's instruction manual. The single copy B-globin gene Fragment at 5.2 Kb was observed within 3 hours of staining with the NBT/BCIP dye system for DNA probes labeled with Biotin-n-dATP where n=3, 7, 10 or 14 and Biotin-n-dCTP where n=7 or 10.

-14-

Thus, Biotin-n-dATP where n=3,7,10 or 14 and Biotin-n-dCTP where n=7 or 10 may be used to label a nucleic acid probe which will then recognize a single copy gene sequence in a Southern blot of genomic DNA.

5 The annealed probe-target hybrid may then be detected by a commercially available DNA detection system.

Specific Example of Synthesis of Bio-7-dATP

Following is an example of the first described synthetic procedure, in which Bio-7-dATP was prepared:

1. Preparation of Biotin-N-hydroxysuccinimide

5 Biotin (1.0 g, 4.1 mmole) was dissolved in 10 ml DMF (dry) with heating at 80°C in oil bath. CDI (665 mg, 4.1 mmoles) was added and the mixture was heated at 80°C. The reaction mixture was stirred at 80°C for 30 minutes, then at room temperature for 2 hours; a white
10 precipitate formed. N-hydroxysuccinimide (475 mg, 4.1 mmoles) was added and the reaction mixture was stirred at room temperature overnight. DMF was removed under vacuum on rotary evaporator. The solid residue was dissolved in 250 ml of refluxing isopropanol, filtered,
15 and stored in the cold room overnight. The precipitate was filtered, washed one time with cold isopropanol and dried in vacuo at 45°C overnight to give 870 mg (61% yield) of the desired product.

2. Preparation of 6-Chloropurine-2'-deoxyriboside

20 2'-Deoxyinosine (5.0 g, .02 M) in methylene chloride (100 ml) was cooled to 0°C under nitrogen in a 3-neck 3 liter flask. Trifluoroacetic anhydride (45.0 g, 0.25 mole) was added and stirred for 2 hours. The
25 temperature was allowed to rise to 15°C and an aspirator was connected (a trap of drierite was used) and the major part of methylene chloride was removed at room temperature. A vacuum pump was attached and the residual trifluoroacetic anhydride and methylene
30 chloride were removed (about 30 minutes). A foam was obtained at this stage.

The flask was fitted with a dropping funnel, a condenser and a gas dispersion tube inlet. Methylene chloride (500 ml) was added and the solution heated to a
35 gentle reflux and a steady stream of nitrogen was bubbled through. A solution of freshly distilled

-16-

thionyl chloride (16 ml) and DMF (8 ml) in methylene chloride (200 ml) was added dropwise during a 2 hour period. The reaction was refluxed for 5 hours (to overnight) cooled and filtered. An offwhite solid of 0.5 g hypoxanthine was obtained. The solution was slowly poured into a vigorously stirred ice cold solution of sodium bicarbonate (30 g) in 500 ml H₂O. The two phases were separated and the aqueous layer was extracted with two 250 ml portions of methylene chloride. The organic phase was combined, dried over sodium sulfate and concentrated. The resulting semisolid was dissolved in 10 ml methanol, applied to a column of neutral alumina (2.5 cm X 40 cm, 75 g), and eluted with methanol. The first 75 ml was discarded and the column washed with 1 liter of methanol. The methanol was concentrated and the resulting solid crystallized from methanol/ethyl acetate (weight = 3.8 g yield = 70%, mp = 138°-140°C). HPLC analysis showed it to be a single compound.

20 3. 6-Chloropurine-2'-deoxyribose
 5'-monophosphate

Molecular Sieve (4 Å, 5.0 gm) was ground to give a coarse powder, added to triethyl phosphate (10 ml), and stirred vigorously for 10 minutes. The 6-chloropurine-2'-deoxyribose (135 mg) was added and the mixture cooled to 0°C. Phosphorous oxychloride (90 ul) was pretreated with water (9 ul) very carefully at 0°C and added to the reaction mixture in one portion with an oven dry glass pipet or glass syringe. The reaction mixture was monitored on HPLC for the appearance of product and disappearance of starting material. After the complete disappearance of starting material the reaction mixture was filtered and added to 20 ml of ice water and the pH was adjusted to about 7.5 with 1 M TEAB. The solution was extracted with ether (4 X 50 ml).

-17-

The aqueous layer was concentrated on the rotary evaporator to remove any residual ether and diluted to 200 ml with water. It was applied to a Sephadex column (HCO_3^- form, 70 ml, equilibrated with .01 M TEAB) and washed with 200 ml .01 M TEAB. It was then eluted with a gradient of .01 M to 0.5 M TEAB (400 ml each). 20 ml fractions were collected. Fractions 24-40 had UV active fractions; they were combined, concentrated, and coevaporated with ethanol (4 X 100 ml) to give a slightly brownish gum that weighed 245 mg (90%).

4. N^6 -(6-aminohexyl)-2'-deoxyadenosine
5'-monophosphate

The diaminohexane (1.12 g) was dissolved in 5 ml H_2O and the pH adjusted to about 9-9.5 with carbon dioxide. The 6-chloropurine-2'-deoxyriboside 5'-monophosphate (204 mg) in 5 ml H_2O was added and the reaction mixture was heated at 50°C and monitored on HPLC until the complete disappearance of the starting material (1.5 - 3 hours). It was cooled to room temperature, diluted to 200 ml with water, and loaded on Sephadex-A-25 column (70 ml, 2.5 cm X 45 cm, HCO_3^- form equilibrated with .01 M TEAB). The column was washed with 400 ml of .01 M TEAB followed with a gradient of .01 M to 0.5 M TEAB (350 ml each) and 20ml fractions were collected. Fractions 10-20 contained the desired compound, which was concentrated and coevaporated with ethanol (4 X 100 ml) to give an offwhite solid that weighed 184 mg (70%). HPLC analysis showed it to be a single peak.

5. N^6 -(6-hexylamidobiotin-2'-deoxyadenosine
5'-monophosphate

6-aminohexyl dAMP (80 mg) was dissolved in 0.1 M sodium borate (10 ml, pH 8.5). Biotin-NHS was dissolved in DMF (1.5 ml) and added to the amine. The

-18-

mixture was stirred vigorously until HPLC analysis showed the disappearance of the starting material (1-4 hours). The reaction mixture was concentrated to remove the DMF and then dissolved in 100 ml of water and loaded
5 onto a Sephadex-A-25 column (HCO_3^- form, 30 ml, 1.5 cm X 40 cm). The column was washed with 100 ml 0.01 M TEAB. It was then eluted with a gradient 0.1 M to 0.5 M TEAB (300 ml each); 18 ml fractions were collected. The
10 desired compound eluted in fractions 20-25. The fractions were combined and concentrated and coevaporated with ethanol (4 X 100 ml) to give a white solid that weighed 90 mg (81%). It was a single compound on HPLC analysis and was positive for biotin.

15 6. N^6 -(6-hexylamidobiotin)-2'-deoxyadenosine 5'-triphosphate (bio-7-dATP)

Bio-7-dAMP (45 mg) was coevaporated with anhydrous DMF (3 X 10 ml) and finally dissolved in 2 ml DMF. Carbonyldiimidazole (40 mg) was added and the flask was tightly stoppered and vigorously stirred for
20 about 1 hour. HPLC analysis revealed the starting material was consumed. (Reaction is stopped when the starting material is completely consumed. This may take 30 minutes to about 2 hours). 15 ul of methanol was added and stirring continued for 30 minutes.

25 Tributylammonium pyrophosphate (1.3 ml of 89 mg/ml DMF solution) was added and stirring continued overnight. The material was filtered. The filtrate was concentrated and then diluted to 100 ml with H_2O and loaded on Sephadex-A-25 (HCO_3^- form, 30 ml 1.5 X 30 cm).
30 It was washed with 100 ml 0.01 M TEAB followed with a gradient of 0.01 M to 0.5 M TEAB (200 ml each) and then 200 ml each of 0.5 M and 1.0 M TEAB. 20 ml fractions were collected. Fractions 30-38 contained the desired compound which weighed 40 mg (56%). It was a single
35 compound on HPLC.

-19-

The invention has been described in detail with particular emphasis on the preferred embodiments thereof, but it should be understood that variations and modifications within the spirit and scope of the
5 invention may occur to those skilled in the art to which the invention pertains.

We claim:

1. In the process of detecting the presence of a target polynucleotide, said process comprising

5 incorporating a deoxyribonucleotide, modified by the attachment thereto of a reporter group R, into a polynucleotide complementary to said target polynucleotide,

10 hybridizing said complementary polynucleotide to said target polynucleotide, and

detecting the presence of said reporter group R;

15 The improvement comprising using, for said modified deoxyribonucleotide, a compound of the form

(d)BTP-NH-M, where

20 (d)BTP represents dATP, dCTP, dGTP, 3'-deoxy ATP, 3'-deoxy CTP, 3'-deoxy GTP, 2'-3'-dideoxy ATP, 2'-3' dideoxy CTP or 2'-3' dideoxy GTP;

25 M is bonded through NH to the N⁶ position of B when B represents adenosine, to the N⁴ position of B when B represents cytidine, and to the N² position of B when B represents guanosine; and

30 M represents a moiety containing said reporter group R.

2. The improved process of claim 1 and in which (d)BTP represents dATP or dCTP.

3. The improved process of claim 1 and in which (d)BTP represents dGTP.

5 4. The improved process of claim 1 and in which M represents a compound of the form L-X-R, where X represents an amido, ester or thioester group, and L is an organic compound linking NH to a X by a chain of between one and 12 atoms.

10 5. The improved process of claim 4 and in which X represents NH and L represents either $(\text{CH}_2)_n$ for an n such that $1 \leq n \leq 12$, or $(\text{CH}_2)_n \text{NHCO}(\text{CH}_2)_m$ for an n and m such that $2 \leq n+m \leq 24$.

-22-

6. In the process of detecting the presence of a target polynucleotide, said process comprising

5 incorporating a deoxyribonucleotide, modifiable by the attachment thereto of a reporter group R, into a polynucleotide complementary to said target polynucleotide,

10 in either order, attaching said reporter group R to said modifiable deoxyribonucleotide and hybridizing said complementary polynucleotide to said target polynucleotide, and

15 detecting the presence of said reporter group R;

The improvement comprising using, for said modifiable deoxyribonucleotide, a compound of the form

20 (d)BTP-NH-M, where

25 (d)BTP represents dATP, dCTP, dGTP, 3'-deoxy ATP, 3'-deoxy CTP, 3'-deoxy GTP, 2'-3'-dideoxy ATP, 2'-3' dideoxy CTP or 2'-3' dideoxy GTP;

30 M is bonded through NH to the N⁶ position of B when B represents adenosine, to the N⁴ position of B when B represents cytidine, and to the N² position of B when B represents guanosine; and

M represents a moiety to which said reporter group R may be attached.

7. The improved process of claim 6 and in which (d)BTP represents dATP or dCTP.
8. The improved process of claim 6 and in which (d)BTP represents dGTP.
- 5 9. The improved process of claim 6 and in which M represents a compound of the form L-X, where X represents NH_2 , OH or SH, and L is an organic compound linking NH to a X by a chain of between one and 12 atoms.
- 10 10. The improved process of claim 9 and in which X represents NH_2 and L represents either $(\text{CH}_2)_n$ for an n such that $1 \leq n \leq 12$, or $(\text{CH}_2)_n \text{NHCO}(\text{CH}_2)_m$ for an n and m such that $2 \leq n+m \leq 24$.

11. A compound having the structure



5 wherein

(a) (d)BTP represents either dBTP (2'-deoxy BTP), 3'-deoxy BTP, or 2'-3'-dideoxy BTP, and B represents adenosine, cytidine or guanosine;

10 (b) when B represents adenosine or guanosine, L is bonded through NH to the N⁶ position of adenosine or the N² position of guanosine, and L represents either (CH₂)_n for an n such that 1 ≤ n ≤ 12,

15 (CH₂)_nNHCO(CH₂)_m for an n and m such that 2 ≤ n+m ≤ 24, CH₂CONH(CH₂)_n for 1 ≤ n ≤ 12, or CH₂CONH(CH₂)_nNHCO(CH₂)₅ for 1 ≤ n ≤ 12;

20 (c) when B represents cytidine, L is bonded through NH to the N⁴ position of B and L represents either (CH₂)_n for an n such that 1 ≤ n ≤ 12 or (CH₂)_nNHCO(CH₂)_m for an n and m such that 2 ≤ n+m ≤ 24, and

25 (d) X represents either H or a moiety capable of forming a detectable complex when the compound is incorporated into a nucleic acid.

12. The compound of claim 11 and wherein X is biotin.


30 13. The compound of claim 11 and wherein X is a protein.

14. The compound of claim 11 and wherein X is a fluorescent reporter group.

15. The compound of claim 11 and wherein B is either adenosine or cytidine.

35 16. The compound of claim 11 and wherein (d)BTP is either dATP or dCTP.


17. A compound having the structure



wherein

- 10
- (a) (d)R represents 2'-deoxyribose, 3'-deoxyribose or 2'-3'-dideoxyribose; and either
- (b) B represents either adenine or cytosine and L represents either $(\text{CH}_2)_n$ where $1 \leq n \leq 12$, or $(\text{CH}_2)_n \text{NHCO}(\text{CH}_2)_m$, where
- 15 $2 \leq n+m \leq 24$, or
- (c) B represents adenine and L represents either $\text{CH}_2\text{CONH}(\text{CH}_2)_n$ for $1 \leq n \leq 12$ or $\text{CH}_2\text{CONH}(\text{CH}_2)_n \text{NHCO}(\text{CH}_2)_5$ for $1 \leq n \leq 12$; and wherein, when B represents adenine, L is covalently
- 20 bonded through NH to the N^6 position of B, and when B represents cytosine, L is covalently bonded through NH to the N^4 position of B.

18. A method of preparing a compound having the structure



5

where

- (a) (d)A represents either 2'-deoxy adenosine, 3'-deoxyadenosine or 2'-3' dideoxy adenosine
- (b) R represents biotin, and
- (c) L is bonded through NH to the N⁶ position of adenosine, and L represents $(CH_2)_nNHCO(CH_2)_5$ for an n such that $1 \leq n \leq 12$;

10

15 said method comprising

reacting the corresponding 6-chloropurine-nucleoside with triethylphosphate and phosphorous oxychloride in the presence of molecular sieves to obtain 6-chloropurine-nucleoside-5'-monophosphate;

reacting said 6-chloropurine-nucleoside 5'-monophosphate with $H_2N(CH_2)_nNH_2$ to obtain N⁶-(n-aminoalkyl)(d)AMP;

20

reacting said N⁶-aminoalkyl (d)AMP with N-hydroxysuccinimide caproylamidobiotin ester to obtain N⁶-alkylcaproylamidobiotin-(d)AMP; and


25

reacting said N⁶-alkylcaproyl amidobiotin-(d)AMP with carbonyldiimidazole, followed by tributylammonium pyrophosphate, to obtain the desired said compound.

30

19. The method of claim 18 and in which (d)ATP represents 2'-deoxy ATP.

20. A method of preparing a compound having the structure



5

where

(a) (d)A represents either 2'-deoxy adenosine, 3'-deoxy adenosine, or 2'-3'dideoxy adenosine,

10

(b) R represents biotin, and

(c) L is bonded through NH to the N⁶ position of adenosine, and L represents

(CH₂)_n for an n such that
1 ≤ n ≤ 12;

15 said method comprising

reacting 6-chloropurine-nucleoside with triethylphosphate and phosphorous oxychloride in the presence of molecular sieves to obtain 6-chloropurine-nucleoside-5'-monophosphate;

20

reacting said 6-chloropurine-nucleoside-5'-monophosphate with H₂N(CH₂)_nNH₂ to obtain N⁶-(n-aminoalkyl)(d)AMP;

reacting said N⁶-(n-aminoalkyl)(d)AMP with N-hydroxysuccinimide biotin ester to obtain N⁶-alkylamidobiotin-(d)AMP; and

25

reacting said N⁶-alkylamidobiotin-(d)AMP with carbonyldiimidazole, followed by tributylammonium pyrophosphate, to obtain the desired said compound.

30

21. The method of claim 20 and in which (d)ATP represents 2'-deoxy ATP.

22. A method of preparing a compound having the structure



where

- 5 (a) (d)A represents either 2'-deoxy adenosine, 3'-deoxy adenosine or 2'-3' dideoxy adenosine
- (b) R represents biotin, and
- 10 (c) L is bonded through NH to the N⁶ position of adenosine, and L represents
- (1) $(CH_2)_n$ for an n such that $1 \leq n \leq 12$;
or
- (2) $(CH_2)_n NHCO(CH_2)_5$ for an n such that $1 \leq n \leq 12$;

15 said method comprising

reacting 6-chloropurine-nucleoside with triethylphosphate and phosphorous oxychloride in the presence of molecular sieves to obtain 6-chloropurine-nucleoside-5'-monophosphate;

20 reacting said 6-chloropurine-nucleoside-5'-monophosphate with $H_2N(CH_2)_nNH_2$ to obtain N⁶-(n-aminoalkyl)(d)AMP;

reacting said N⁶-(n-aminoalkyl)(d)AMP with ethyl trifluorothiolacetate to obtain N⁶-(n-trifluoroacetamidoalkyl)(d)AMP;

25 reacting said N⁶-(n-trifluoroacetamidoalkyl)(d)AMP with carbonyldiimidazole, followed by tributylammonium pyrophosphate, to obtain N⁶-(n-trifluoroacetamidoalkyl)(d)ATP;


30 hydrolyzing the trifluoroacetyl group at high pH to obtain N⁶-(n-aminoalkyl)(d)ATP; and

reacting said N⁶-(n-aminoalkyl)(d)ATP with N-hydroxysuccinimide biotin ester to obtain the desired said compound (1), or with N-hydroxysuccinimide caproylamidobiotin ester to obtain the desired said compound (2).

23. The method of claim 22 and in which (d)ATP represents 2'-deoxy ATP.

-29-

24. A method of preparing a compound having the structure



where

5

(a) (d)A represents either 2'-deoxy adenosine, 3'-deoxy adenosine or 2'-3' dideoxy adenosine;

(b) R represents biotin, and

10

(c) L is bonded through NH to the N⁶ position of adenosine, and L represents

(1) $CH_2CONH(CH_2)_n$ for an n such that $1 \leq n \leq 12$; or

(2) $CH_2CONH(CH_2)_nNHCO(CH_2)_5$ for an n such that $1 \leq n \leq 12$;

15 said method comprising

reacting said (d)ATP with iodoacetic acid at pH 6.5 to obtain N'-carboxymethyl (d)ATP, raising the pH to 8.5 and temperature to 90°C to obtain N⁶-carboxymethyl(d)ATP, reacting


20 said N⁶-carboxymethyl(d)ATP with $H_2N(CH_2)_nNH_2$ in the presence of ethyl dimethylaminopropyl carbodiimide to obtain N⁶-(n-alkylamido-carboxymethyl)(d)ATP, and

25


reacting said N⁶-(n-alkylamido-carboxymethyl)(d)ATP with N-hydroxysuccinimide biotin ester to obtain the desired said compound (1), or with N-hydroxysuccinimide caproylamidobiotin ester to obtain the desired said compound (2).

30

25. The method of claim 24 and in which (d)ATP represents 2'-deoxy ATP.



- | | |
|--------------|------------------------------------|
| Blo-3-dATP | $L = (CH_2)_2$ |
| Blo-7-dATP | $L = (CH_2)_6$ |
| Blo-10-dATP | $L = (CH_2)_2NHCO(CH_2)_6$ |
| Blo-14-dATP | $L = (CH_2)_6NHCO(CH_2)_6$ |
| Blo-10'-dATP | $L = CH_2CONH(CH_2)_6$ |
| Blo-17-dATP | $L = CH_2CONH(CH_2)_6NHCO(CH_2)_6$ |



- | | |
|-------------|----------------------------|
| Blo-3-dCTP | $L = (CH_2)_2$ |
| Blo-7-dCTP | $L = (CH_2)_6$ |
| Blo-10-dCTP | $L = (CH_2)_2NHCO(CH_2)_6$ |
| Blo-14-dCTP | $L = (CH_2)_6NHCO(CH_2)_6$ |

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 5

SUBSTITUTE SHEET





FIGURE 6

INTERNATIONAL SEARCH REPORT

International Application No PCT/US85/02198

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
National: 536/27; 536/28; 536/29		
IPC. CL ⁴ C07H 15/12; C07H 17/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
U.S.	536/27; 536/28; 536/29	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁵		
CAS ONLINE COMPUTER SEARCH		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
Y	N, <i>Chemica Scripta</i> (Sweden) Volume 2, No. 4, issued 1972, Guilford et al, "On Adenine Nucleotides for Affinity chromatography.	1-11 and 15-18
Y	US,A, 4,038,480, Published 26 July 1977.	11,13 and 15-18
A,Y	JP,A, 53-133283, Published 20 November 1978.	11-13 and 15-24
Y	US,A, 4,088,639, Published 09 May 1978.	1-11,13 and 15-18
<p>⁶ Special categories of cited documents: ¹⁵</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ¹	Date of Mailing of this International Search Report ²	
29 January 1986	24 FEB 1986	
International Searching Authority ¹	Signature of Authorized Officer ²¹	
ISA/US	 ETHEL G. LOVE PRIMARY EXAMINER	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	US,A, 4,213,893, Published 22 July 1980.	11,13 and 15-18
Y	US,A, 4,255,566, Published 10 March 1981.	1-11,13 and 15- 18
X	EP,A, 0063879, Published 11 March 1982.	1-25

V. OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹⁰

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. Claim numbers because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

VI. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ¹¹

This International Searching Authority found multiple inventions in this international application as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- The additional search fees were accompanied by applicant's protest.
- No protest accompanied the payment of additional search fees.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, ¹⁴ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No ¹⁸
Y	US,A, 4,443,594, Published 17 April 1984.	18-23
Y	US,A, 4,460,772, Published 17 July 1984.	11, 13 and 15-18
Y	JP,A, 57-11999, Published 21 January 1982.	1-11, 13 and 15-18
P,Y	JP,A, 60-96610, Published 30 May 1985.	11 and 15-18